

**Table 1. Estimated numbers of cases of HIV/AIDS, by year of diagnosis and selected characteristics of persons, 2000–2003—33 areas with confidential name-based HIV infection reporting**

	Year of diagnosis			
	2000	2001	2002	2003
<b>Age at diagnosis (yrs)</b>				
<13	196	238	210	161
13–14	36	43	36	50
15–24	3,502	3,568	3,601	3,847
25–34	9,340	8,920	9,001	8,722
35–44	11,312	11,072	11,382	11,299
45–54	4,924	5,179	5,504	5,821
55–64	1,412	1,496	1,614	1,681
≥65	439	485	457	466
<b>Race/ethnicity</b>				
White, not Hispanic	9,962	9,803	10,214	10,322
Black, not Hispanic	16,257	16,042	16,216	16,165
Hispanic	4,340	4,560	4,833	4,963
Asian/Pacific Islander	177	162	187	273
American Indian/Alaska Native	178	164	185	188
<b>Transmission category</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	13,112	13,483	14,355	14,532
Injection drug use	3,539	3,298	3,227	3,189
Male-to-male sexual contact and injection drug use	1,364	1,302	1,256	1,224
Heterosexual contact	3,876	3,879	3,949	4,041
Other <sup>a</sup>	172	172	175	168
Subtotal	22,064	22,134	22,963	23,153
<b>Female adult or adolescent</b>				
Injection drug use	1,884	1,722	1,621	1,628
Heterosexual contact	6,854	6,770	6,870	6,942
Other <sup>a</sup>	160	138	142	163
Subtotal	8,899	8,630	8,633	8,733
<b>Child(&lt;13 yrs at diagnosis)</b>				
Perinatal	175	218	193	147
Other <sup>b</sup>	22	20	17	15
Subtotal	197	238	210	162
<b>Subtotal for 32 states with confidential name-based HIV infection reporting</b>	<b>31,104</b>	<b>30,953</b>	<b>31,749</b>	<b>31,994</b>
<b>Subtotal for U.S. dependencies, possessions, and associated nations with confidential name-based HIV infection reporting</b>	<b>57</b>	<b>49</b>	<b>57</b>	<b>54</b>
<b>Total<sup>c</sup></b>	<b>31,161</b>	<b>31,002</b>	<b>31,805</b>	<b>32,048</b>

*Note.* These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1999, the following 33 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

<sup>a</sup> Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

<sup>b</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>c</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.