

Table 10. Estimated numbers of persons living with AIDS, by year and selected characteristics, 1999–2003—United States

	1999	2000	2001	2002	2003
Age as of end of year (yrs)					
<13	3,034	2,843	2,605	2,335	1,998
13–14	440	517	645	728	768
15–24	4,719	4,991	5,229	5,668	6,313
25–34	60,184	56,686	53,687	51,410	49,906
35–44	141,295	151,180	158,173	163,732	168,322
45–54	77,216	89,461	102,252	115,613	129,311
55–64	19,258	22,922	27,197	32,703	38,997
≥65	5,058	6,132	7,251	8,583	10,310
Race/ethnicity					
White, not Hispanic	119,674	126,162	132,258	139,089	146,544
Black, not Hispanic	126,044	137,524	148,469	160,022	172,278
Hispanic	61,194	66,266	71,034	75,782	80,623
Asian/Pacific Islander	2,484	2,755	3,056	3,414	3,826
American Indian/Alaska Native	1,047	1,166	1,262	1,380	1,498
Transmission category					
Male adult or adolescent					
Male-to-male sexual contact	140,216	150,172	160,076	171,035	182,989
Injection drug use	58,006	61,249	63,723	66,003	68,191
Male-to-male sexual contact and injection drug use	21,667	22,403	23,033	23,690	24,334
Heterosexual contact	20,595	23,478	26,471	29,835	33,324
Other ^a	3,807	3,922	4,062	4,204	4,345
Subtotal	244,291	261,223	277,366	294,767	313,183
Female adult or adolescent					
Injection drug use	25,744	27,317	28,602	29,670	30,710
Heterosexual contact	35,603	40,422	45,097	50,142	55,685
Other ^a	1,746	1,908	2,067	2,239	2,420
Subtotal	63,093	69,647	75,765	82,052	88,815
Child(<13 yrs at diagnosis)					
Perinatal	3,672	3,714	3,763	3,808	3,788
Other ^b	148	145	145	143	139
Subtotal	3,820	3,860	3,908	3,951	3,927
Region of residence					
Northeast	92,741	99,964	105,970	111,506	116,827
Midwest	31,016	33,470	35,725	38,513	41,668
South	115,991	125,396	135,465	146,421	158,962
West	62,300	66,280	69,931	74,253	78,333
U.S. dependencies, possessions, and associated nations	9,157	9,621	9,949	10,077	10,136
Total^c	311,205	334,731	357,040	380,771	405,926

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

^a Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^c Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.