

Table 13. Proportion of persons surviving for more than 12, 24, and 36 months after an AIDS diagnosis in 1999, by selected characteristics—United States

	No. of persons	Survival in months		
		>12	>24	>36
		Proportion		
Age at diagnosis (yrs)				
<13	172	0.92	0.89	0.89
13—14	55	0.96	0.95	0.93
15—24	1,498	0.95	0.92	0.89
25—34	10,940	0.93	0.90	0.87
35—44	16,419	0.91	0.87	0.84
45—54	7,618	0.86	0.82	0.78
55—64	2,043	0.79	0.74	0.71
≥65	658	0.69	0.62	0.58
Race/ethnicity				
White, not Hispanic	12,095	0.92	0.88	0.86
Black, not Hispanic	19,018	0.89	0.84	0.80
Hispanic	7,685	0.90	0.87	0.84
Asian/Pacific Islander	357	0.91	0.88	0.87
American Indian/Alaska Native	151	0.90	0.85	0.83
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	13,680	0.92	0.89	0.87
Injection drug use	6,048	0.88	0.82	0.78
Male-to-male sexual contact and injection drug use	1,984	0.92	0.87	0.83
Heterosexual contact	3,050	0.90	0.86	0.82
Other ^a	4,923	0.86	0.82	0.80
Subtotal	29,685	0.90	0.86	0.83
Female adult or adolescent				
Injection drug use	2,689	0.89	0.83	0.78
Heterosexual contact	4,464	0.93	0.89	0.86
Other ^a	2,393	0.88	0.83	0.80
Subtotal	9,546	0.90	0.86	0.82
Child(<13 yrs at diagnosis)				
Perinatal	162	0.93	0.90	0.90
Other ^b	10	0.80	0.80	0.80
Subtotal	172	0.92	0.89	0.89
Total^c	39,403	0.90	0.86	0.83

Note. Excludes persons whose date of death is before, or in the same month as, date of diagnosis. Excludes persons whose month of diagnosis or month of death is unknown.

^a Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^c Includes 97 persons of unknown race or multiple races.