

Table 19. Reported AIDS cases for male adults and adolescents, by transmission category and race/ethnicity, cumulative through 2003—United States

| Transmission category | White, not Hispanic | | | | Black, not Hispanic | | | | Hispanic | | | |
|--|---------------------|------------|--------------------------------------|------------|---------------------|------------|--------------------------------------|------------|--------------|------------|--------------------------------------|------------|
| | 2003 | | Cumulative through 2003 ^a | | 2003 | | Cumulative through 2003 ^a | | 2003 | | Cumulative through 2003 ^a | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sexual contact | 7,679 | 66 | 244,758 | 73 | 4,699 | 34 | 93,413 | 37 | 3,054 | 43 | 57,128 | 43 |
| Injection drug use | 1,051 | 9 | 31,164 | 9 | 2,454 | 18 | 80,282 | 32 | 1,290 | 18 | 44,277 | 33 |
| Male-to-male sexual contact and injection drug use | 793 | 7 | 28,795 | 9 | 548 | 4 | 19,182 | 8 | 311 | 4 | 9,313 | 7 |
| Hemophilia/coagulation disorder | 56 | 0 | 3,964 | 1 | 6 | 0 | 599 | 0 | 9 | 0 | 453 | 0 |
| Heterosexual contact: | 454 | 4 | 7,010 | 2 | 2,047 | 15 | 24,428 | 10 | 799 | 11 | 9,021 | 7 |
| <i>Sex with injection drug user</i> | 76 | 1 | 2,221 | 1 | 253 | 2 | 6,410 | 3 | 141 | 2 | 2,195 | 2 |
| <i>Sex with person with hemophilia</i> | 4 | 0 | 38 | 0 | 2 | 0 | 29 | 0 | 0 | 0 | 11 | 0 |
| <i>Sex with HIV-infected transfusion recipient</i> | 4 | 0 | 177 | 0 | 11 | 0 | 205 | 0 | 7 | 0 | 109 | 0 |
| <i>Sex with HIV-infected person, risk factor not specified</i> | 370 | 3 | 4,574 | 1 | 1,781 | 13 | 17,784 | 7 | 651 | 9 | 6,706 | 5 |
| Receipt of blood transfusion, blood components, or tissue | 30 | 0 | 3,227 | 1 | 49 | 0 | 1,205 | 0 | 28 | 0 | 646 | 0 |
| Other/risk factor not reported or identified | 1,640 | 14 | 14,519 | 4 | 3,932 | 29 | 33,905 | 13 | 1,544 | 22 | 12,659 | 9 |
| Total | 11,703 | 100 | 333,437 | 100 | 13,735 | 100 | 253,014 | 100 | 7,035 | 100 | 133,497 | 100 |

| Transmission category | Asian/Pacific Islander | | | | American Indian/Alaska Native | | | | Total | | | |
|--|------------------------|------------|--------------------------------------|------------|-------------------------------|------------|--------------------------------------|------------|---------------|------------|--------------------------------------|------------|
| | 2003 | | Cumulative through 2003 ^a | | 2003 | | Cumulative through 2003 ^a | | 2003 | | Cumulative through 2003 ^a | |
| | No. | % | No. | % | No. | % | No. | % | No. | % | No. | % |
| Male-to-male sexual contact | 254 | 56 | 4,084 | 69 | 93 | 58 | 1,299 | 56 | 15,859 | 48 | 401,392 | 55 |
| Injection drug use | 26 | 6 | 292 | 5 | 22 | 14 | 370 | 16 | 4,866 | 15 | 156,575 | 21 |
| Male-to-male sexual contact and injection drug use | 19 | 4 | 227 | 4 | 15 | 9 | 392 | 17 | 1,695 | 5 | 57,998 | 8 |
| Hemophilia/coagulation disorder | 2 | 0 | 72 | 1 | 1 | 1 | 32 | 1 | 74 | 0 | 5,130 | 1 |
| Heterosexual contact: | 42 | 9 | 305 | 5 | 11 | 7 | 92 | 4 | 3,371 | 10 | 40,947 | 6 |
| <i>Sex with injection drug user</i> | 3 | 1 | 55 | 1 | 2 | 1 | 28 | 1 | 477 | 1 | 10,930 | 1 |
| <i>Sex with person with hemophilia</i> | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 7 | 0 | 80 | 0 |
| <i>Sex with HIV-infected transfusion recipient</i> | 1 | 0 | 8 | 0 | 1 | 1 | 3 | 0 | 24 | 0 | 505 | 0 |
| <i>Sex with HIV-infected person, risk factor not specified</i> | 38 | 8 | 241 | 4 | 8 | 5 | 61 | 3 | 2,863 | 9 | 29,432 | 4 |
| Receipt of blood transfusion, blood components, or tissue | 3 | 1 | 118 | 2 | 0 | 0 | 9 | 0 | 111 | 0 | 5,219 | 1 |
| Other/risk factor not reported or identified | 110 | 24 | 792 | 13 | 19 | 12 | 130 | 6 | 7,274 | 22 | 62,217 | 9 |
| Total | 456 | 100 | 5,890 | 100 | 161 | 100 | 2,324 | 100 | 33,250 | 100 | 729,478 | 100 |

^a Includes persons with a diagnosis of AIDS, reported from the beginning of the epidemic through 2003. Cumulative total includes 1316 males of unknown race or multiple races.