

**Table 2. Estimated distribution of persons with and persons without a diagnosis of AIDS within 12 months of diagnosis of HIV infection, by selected characteristics, 2002—33 areas with confidential name-based HIV infection reporting**

	AIDS diagnoses					
	≥12 Months after diagnosis of HIV infection		<12 Months after diagnosis of HIV infection		Total	
	No.	(%) <sup>a</sup>	No.	(%) <sup>a</sup>	No.	(%) <sup>a</sup>
<b>Age at diagnosis (yrs)</b>						
<13	165	78	45	22	210	100
13—14	28	78	8	22	36	100
15—24	2,979	83	619	17	3,599	100
25—34	6,059	67	2,937	33	8,995	100
35—44	6,568	58	4,811	42	11,379	100
45—54	2,845	52	2,655	48	5,499	100
55—64	745	46	869	54	1,614	100
≥65	189	41	269	59	457	100
<b>Race/ethnicity</b>						
White, not Hispanic	6,377	62	3,831	38	10,207	100
Black, not Hispanic	10,119	62	6,092	38	16,211	100
Hispanic	2,751	57	2,079	43	4,830	100
Asian/Pacific Islander	119	64	67	36	186	100
American Indian/Alaska Native	118	64	67	36	185	100
<b>Transmission category</b>						
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	8,691	61	5,655	39	14,346	100
Injection drug use	1,728	54	1,497	46	3,225	100
Male-to-male sexual contact and injection drug use	752	60	504	40	1,256	100
Heterosexual contact	2,297	58	1,651	42	3,947	100
Other <sup>b</sup>	87	50	87	50	174	100
Subtotal	13,556	59	9,393	41	22,949	100
<b>Female adult or adolescent</b>						
Injection drug use	1,075	66	545	34	1,621	100
Heterosexual contact	4,712	69	2,157	31	6,869	100
Other <sup>b</sup>	69	49	72	51	142	100
Subtotal	5,857	68	2,774	32	8,631	100
<b>Child(&lt;13 yrs at diagnosis)</b>						
Perinatal	149	77	44	23	193	100
Other <sup>c</sup>	16	93	1	7	17	100
Subtotal	165	78	45	22	210	100
<b>Total<sup>d</sup></b>	<b>19,577</b>	<b>62</b>	<b>12,213</b>	<b>38</b>	<b>31,790</b>	<b>100</b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons in whom AIDS has developed and persons whose first diagnosis of HIV infection and the diagnosis of AIDS were made at the same time.

Data exclude 15 persons whose month of diagnosis of HIV infection is unknown.

Since 1999, the following 33 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

<sup>a</sup> Percentages represent proportions of the total number of diagnoses of HIV/AIDS made during 2002 for the corresponding group (see row entries).

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

<sup>c</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>d</sup> Includes 170 persons of unknown race or multiple races. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.