

Table 20. Reported cases of HIV infection (not AIDS) for male adults and adolescents, by transmission category and race/ethnicity, cumulative through 2003—41 areas with confidential name-based HIV infection reporting

Transmission category	White, not Hispanic				Black, not Hispanic				Hispanic			
	2003		Cumulative through 2003 ^a		2003		Cumulative through 2003 ^a		2003		Cumulative through 2003 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	5,464	65	41,048	65	2,944	32	21,472	33	1,853	37	8,941	42
Injection drug use	578	7	4,969	8	1,009	11	10,658	16	945	19	3,815	18
Male-to-male sexual contact and injection drug use	387	5	4,606	7	192	2	3,011	5	139	3	838	4
Hemophilia/coagulation disorder	33	0	381	1	9	0	107	0	6	0	27	0
Heterosexual contact	307	4	2,053	3	1,212	13	8,681	13	466	9	1,766	8
<i>Sex with injection drug user</i>	67	1	490	1	163	2	1,445	2	74	1	304	1
<i>Sex with person with hemophilia</i>	5	0	7	0	0	0	14	0	1	0	4	0
<i>Sex with HIV-infected transfusion recipient</i>	4	0	27	0	9	0	76	0	2	0	11	0
<i>Sex with HIV-infected person, risk factor not specified</i>	231	3	1,529	2	1,040	11	7,146	11	389	8	1,447	7
Receipt of blood transfusion, blood components, or tissue	13	0	216	0	8	0	204	0	3	0	46	0
Other/risk factor not reported or identified	1,572	19	9,676	15	3,708	41	21,183	32	1,618	32	5,968	28
Total	8,354	100	62,949	100	9,082	100	65,316	100	5,030	100	21,401	100

Transmission category	Asian/Pacific Islander				American Indian/Alaska Native				Total			
	2003		Cumulative through 2003 ^a		2003		Cumulative through 2003 ^a		2003		Cumulative through 2003 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	118	59	493	54	49	53	428	55	10,466	46	72,745	48
Injection drug use	9	5	46	5	8	9	89	11	2,551	11	19,652	13
Male-to-male sexual contact and injection drug use	5	3	20	2	6	7	103	13	732	3	8,623	6
Hemophilia/coagulation disorder	0	0	2	0	0	0	0	0	48	0	520	0
Heterosexual contact:	7	4	57	6	8	9	51	7	2,009	9	12,669	8
<i>Sex with injection drug user</i>	1	1	8	1	1	1	16	2	307	1	2,272	1
<i>Sex with person with hemophilia</i>	0	0	0	0	0	0	0	0	6	0	25	0
<i>Sex with HIV-infected transfusion recipient</i>	0	0	2	0	0	0	0	0	15	0	116	0
<i>Sex with HIV-infected person, risk factor not specified</i>	6	3	47	5	7	8	35	4	1,681	7	10,256	7
Receipt of blood transfusion, blood components, or tissue	1	1	3	0	0	0	2	0	26	0	477	0
Other/risk factor not reported or identified	60	30	290	32	21	23	111	14	7,003	31	38,053	25
Total	200	100	911	100	92	100	784	100	22,835	100	152,739	100

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2003, the following 41 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Georgia, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Puerto Rico, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

^a Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through December 2003. Cumulative total includes 1378 males of unknown race or multiple races.