

**Table 22. Reported cases of HIV infection (not AIDS) for female adults and adolescents, by transmission category and race/ethnicity, cumulative through 2003—41 areas with confidential name-based HIV infection reporting**

Transmission category	White, not Hispanic				Black, not Hispanic				Hispanic			
	2003		Cumulative through 2003 <sup>a</sup>		2003		Cumulative through 2003 <sup>a</sup>		2003		Cumulative through 2003 <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Injection drug use	410	22	3,545	26	565	9	6,255	15	359	18	1,512	18
Hemophilia/coagulation disorder	4	0	22	0	7	0	35	0	1	0	5	0
Heterosexual contact:	824	44	6,054	45	2,328	39	18,258	44	827	42	3,747	46
<i>Sex with injection drug user</i>	177	9	1,685	13	273	5	3,284	8	169	9	848	10
<i>Sex with bisexual male</i>	50	3	539	4	107	2	1,034	3	31	2	156	2
<i>Sex with person with hemophilia</i>	7	0	104	1	4	0	59	0	1	0	7	0
<i>Sex with HIV-infected transfusion recipient</i>	3	0	50	0	8	0	89	0	3	0	32	0
<i>Sex with HIV-infected person, risk factor not specified</i>	587	31	3,676	27	1,936	32	13,792	34	623	32	2,704	33
Receipt of blood transfusion, blood components, or tissue	12	1	152	1	39	1	325	1	9	0	53	1
Other/risk factor not reported or identified	624	33	3,632	27	3,077	51	16,194	39	774	39	2,877	35
<b>Total</b>	<b>1,874</b>	<b>100</b>	<b>13,405</b>	<b>100</b>	<b>6,016</b>	<b>100</b>	<b>41,067</b>	<b>100</b>	<b>1,970</b>	<b>100</b>	<b>8,194</b>	<b>100</b>

  

Transmission category	Asian/Pacific Islander				American Indian/Alaska Native				Total			
	2003		Cumulative through 2003 <sup>a</sup>		2003		Cumulative through 2003 <sup>a</sup>		2003		Cumulative through 2003 <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Injection drug use	6	10	20	7	11	25	99	32	1,355	14	11,480	18
Hemophilia/coagulation disorder	0	0	1	0	0	0	1	0	12	0	64	0
Heterosexual contact:	22	35	136	45	11	25	136	44	4,036	40	28,483	45
<i>Sex with injection drug user</i>	0	0	10	3	5	11	52	17	628	6	5,901	9
<i>Sex with bisexual male</i>	1	2	6	2	2	5	10	3	193	2	1,757	3
<i>Sex with person with hemophilia</i>	0	0	0	0	0	0	2	1	12	0	174	0
<i>Sex with HIV-infected transfusion recipient</i>	0	0	2	1	0	0	1	0	14	0	175	0
<i>Sex with HIV-infected person, risk factor not specified</i>	21	34	118	39	4	9	71	23	3,189	32	20,476	32
Receipt of blood transfusion, blood components, or tissue	0	0	6	2	1	2	1	0	61	1	539	1
Other/risk factor not reported or identified	34	55	141	46	21	48	74	24	4,543	45	23,174	36
<b>Total</b>	<b>62</b>	<b>100</b>	<b>304</b>	<b>100</b>	<b>44</b>	<b>100</b>	<b>311</b>	<b>100</b>	<b>10,007</b>	<b>100</b>	<b>63,740</b>	<b>100</b>

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2003, the following 41 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Georgia, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Puerto Rico, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

<sup>a</sup> Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through 2003. Cumulative total includes 459 females of unknown race or multiple races.