

Table 3. Estimated numbers of AIDS cases, by year of diagnosis and selected characteristics of persons, 1999–2003— United States

	Year of diagnosis					Cumulative through 2003 ^a
	1999	2000	2001	2002	2003	
Age at diagnosis (yrs)						
<13	187	117	119	105	59	9,419
13–14	57	56	76	68	59	891
15–24	1,541	1,642	1,625	1,810	1,991	37,599
25–34	11,349	10,385	9,947	9,504	9,605	311,137
35–44	17,165	17,295	16,890	17,008	17,633	365,432
45–54	8,099	8,566	8,929	9,310	10,051	148,347
55–64	2,218	2,422	2,468	2,724	2,888	43,451
≥65	739	783	779	759	886	13,711
Race/ethnicity						
White, not Hispanic	12,626	12,047	11,620	11,960	12,222	376,834
Black, not Hispanic	19,960	20,312	20,291	20,476	21,304	368,169
Hispanic	8,141	8,233	8,204	8,021	8,757	172,993
Asian/Pacific Islander	369	373	409	452	497	7,166
American Indian/Alaska Native	162	186	179	196	196	3,026
Transmission category						
Male adult or adolescent						
Male-to-male sexual contact	16,556	16,272	16,383	16,971	17,969	440,887
Injection drug use	7,710	7,425	6,772	6,406	6,353	175,988
Male-to-male sexual contact and injection drug use	2,323	2,071	2,026	1,942	1,877	62,418
Heterosexual contact	4,243	4,299	4,578	4,890	5,133	56,403
Other ^b	328	319	315	308	281	14,191
Subtotal	31,159	30,387	30,074	30,517	31,614	749,887
Female adult or adolescent						
Injection drug use	3,448	3,498	3,269	3,024	3,096	70,558
Heterosexual contact	6,350	7,011	7,119	7,380	8,127	93,586
Other ^b	212	254	251	261	276	6,535
Subtotal	10,010	10,763	10,639	10,666	11,498	170,679
Child(<13 yrs at diagnosis)						
Perinatal	185	115	116	103	58	8,749
Other ^c	3	2	3	3	1	670
Subtotal	187	117	119	105	59	9,419
Region of residence						
Northeast	11,885	12,516	11,350	10,551	11,461	285,040
Midwest	4,069	4,139	4,094	4,337	4,498	91,926
South	17,224	16,757	17,693	18,482	19,609	337,409
West	6,892	6,661	6,468	6,843	6,667	186,100
U.S. dependencies, possessions, and associated nations	1,286	1,194	1,228	1,075	935	29,511
Total^d	41,356	41,267	40,833	41,289	43,171	929,985

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

^a Includes persons with a diagnosis of AIDS from the beginning of the epidemic through 2003.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes persons of unknown race or multiple races and persons of unknown sex. Cumulative total includes 1796 persons of unknown race or multiple races and 1 person of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.