

**Table 4. Estimated numbers of AIDS cases in children <13 years of age, by year of diagnosis and transmission category, 1999–2003—United States**

	Year of diagnosis					Cumulative through 2003 <sup>a</sup>
	1999	2000	2001	2002	2003	
<b>Race/ethnicity</b>						
White, not Hispanic	18	12	15	13	9	1,620
Black, not Hispanic	133	86	78	68	40	5,562
Hispanic	34	17	24	22	7	2,128
Asian/Pacific Islander	1	2	2	1	0	56
American Indian/Alaska Native	0	0	0	0	0	30
<b>Transmission category</b>						
Hemophilia/coagulation disorder	0	0	0	0	0	234
Mother with, or at risk for, HIV infection	185	115	116	103	58	8,749
<i>Injection drug use</i>	<i>44</i>	<i>21</i>	<i>13</i>	<i>8</i>	<i>7</i>	<i>3,326</i>
<i>Sex with injection drug user</i>	<i>26</i>	<i>13</i>	<i>8</i>	<i>4</i>	<i>6</i>	<i>1,541</i>
<i>Sex with bisexual male</i>	<i>6</i>	<i>2</i>	<i>4</i>	<i>3</i>	<i>0</i>	<i>203</i>
<i>Sex with person with hemophilia</i>	<i>2</i>	<i>0</i>	<i>1</i>	<i>0</i>	<i>0</i>	<i>37</i>
<i>Sex with HIV-infected transfusion recipient</i>	<i>0</i>	<i>0</i>	<i>0</i>	<i>0</i>	<i>0</i>	<i>28</i>
<i>Sex with HIV-infected person, risk factor not specified</i>	<i>49</i>	<i>36</i>	<i>35</i>	<i>37</i>	<i>17</i>	<i>1,490</i>
<i>Receipt of blood transfusion, blood components, or tissue</i>	<i>1</i>	<i>2</i>	<i>2</i>	<i>2</i>	<i>0</i>	<i>156</i>
<i>Has HIV infection, risk factor not specified</i>	<i>57</i>	<i>41</i>	<i>53</i>	<i>48</i>	<i>28</i>	<i>1,968</i>
Receipt of blood transfusion, blood components, or tissue	0	1	0	1	0	391
Other/risk factor not reported or identified	3	1	3	2	1	45
<b>Total<sup>b</sup></b>	<b>187</b>	<b>117</b>	<b>119</b>	<b>105</b>	<b>59</b>	<b>9,419</b>


Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

<sup>a</sup> Includes children with a diagnosis of AIDS, from the beginning of the epidemic through 2003.

<sup>b</sup> Includes children of unknown race or multiple races. Cumulative total includes 24 children of unknown race or multiple races.

Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

**Figure 1. Estimated numbers of AIDS cases in children <13 years of age, by year of diagnosis, 1992–2003—United States**



Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays. The estimates do not include adjustments for incomplete reporting.