

Table 6. Estimated numbers of AIDS cases in adult or adolescent Hispanics, by transmission category and place of birth, 2003—United States

Transmission category	Place of birth											
	United States		Central/South America		Cuba		Mexico		Puerto Rico		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	1,503	40	375	49	108	62	888	61	238	16	3,609	41
Injection drug use	1,018	27	98	13	14	8	135	9	573	39	2,091	24
Male-to-male sexual contact and injection drug use	145	4	22	3	7	4	63	4	67	5	345	4
Heterosexual contact	1,024	27	259	34	44	25	351	24	569	39	2,593	30
Other ^b	53	1	10	1	1	0	22	2	10	1	111	1
Total^c	3,743	100	764	100	174	100	1,459	100	1,458	100	8,749	100

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

^a Includes 189 persons whose place of birth is not among those listed and 963 persons whose place of birth is unknown.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.