

**Table 8. Estimated numbers of persons living with HIV/AIDS by year and selected characteristics, 2000–2003—33 areas with confidential name-based HIV infection reporting**

	2000	2001	2002	2003
<b>Age as of end of year (yrs)</b>				
<13	2,898	2,867	2,796	2,614
13—14	336	442	519	618
15—24	11,297	11,931	12,409	13,134
25—34	67,688	66,711	66,416	66,446
35—44	124,116	132,137	139,133	145,288
45—54	60,616	70,407	80,757	91,567
55—64	14,579	17,314	20,988	25,237
≥65	4,002	4,840	5,687	6,710
<b>Race/ethnicity</b>				
White, not Hispanic	113,617	120,186	127,257	134,678
Black, not Hispanic	135,562	146,057	156,771	167,938
Hispanic	31,950	35,508	39,358	43,241
Asian/Pacific Islander	1,034	1,171	1,344	1,595
American Indian/Alaska Native	1,508	1,606	1,737	1,873
<b>Transmission category</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	128,956	138,629	149,336	160,433
Injection drug use	36,526	38,098	39,630	41,207
Male-to-male sexual contact and injection drug use	19,097	19,642	20,175	20,773
Heterosexual contact	25,262	28,115	31,042	34,124
Other <sup>a</sup>	2,817	2,891	2,972	3,071
Subtotal	212,658	227,375	243,154	259,609
<b>Female adult or adolescent</b>				
Injection drug use	19,789	20,650	21,381	22,173
Heterosexual contact	47,963	53,245	58,547	63,981
Other <sup>a</sup>	1,450	1,547	1,660	1,787
Subtotal	69,202	75,442	81,588	87,940
<b>Child(&lt;13 yrs at diagnosis)</b>				
Perinatal	3,260	3,434	3,593	3,720
Other <sup>b</sup>	406	393	367	342
Subtotal	3,666	3,827	3,960	4,062
<b>Total<sup>c</sup></b>	<b>285,531</b>	<b>306,649</b>	<b>328,705</b>	<b>351,614</b>

*Note.* These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1999, the following 33 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

<sup>a</sup> Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

<sup>b</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>c</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.