

Table 9. Estimated numbers of persons living with HIV/AIDS at the end of 2003, by race/ethnicity, sex, and transmission category—33 areas with confidential name-based HIV infection reporting

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male adult or adolescent												
Male-to-male sexual contact	86,674	76	50,675	47	20,498	60	809	67	840	61	160,433	62
Injection drug use	10,550	9	23,658	22	6,365	19	141	12	210	15	41,207	16
Male-to-male sexual contact and injection drug use	10,431	9	7,817	7	2,165	6	48	4	187	14	20,773	8
Heterosexual contact	5,178	5	23,513	22	4,814	14	190	16	122	9	34,124	13
Other ^b	1,524	1	1,198	1	276	1	28	2	19	1	3,071	1
Subtotal	114,358	100	106,861	100	34,118	100	1,216	100	1,379	100	259,609	100
Female adult or adolescent												
Injection drug use	6,625	34	13,244	23	1,957	23	51	14	164	34	22,173	25
Heterosexual contact	12,494	64	43,957	75	6,498	75	290	81	306	64	63,981	73
Other ^b	447	2	1,118	2	183	2	18	5	7	2	1,787	2
Subtotal	19,566	100	58,319	100	8,638	100	359	100	477	100	87,940	100
Child (<13 yrs at diagnosis)												
Perinatal	636	84	2,591	94	443	91	15	76	15	88	3,720	92
Other ^c	118	16	165	6	42	9	5	24	2	12	342	8
Subtotal	754	100	2,756	100	485	100	20	100	17	100	4,062	100
Total^d	134,678		167,938		43,241		1,595		1,873		351,614	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1999, the following 33 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Includes 2289 persons of unknown race or multiple races.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes 4 persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.