

Table 11. Annual number of deaths in adolescents and adults with AIDS, by sex and age at death — United States^a, 1994–2000

Age at death (yrs)	Number, by year of death						
	1994	1995	1996	1997	1998	1999	2000
Males							
13-24	484	447	264	165	129	106	101
25-29	3,092	2,823	1,794	971	642	546	387
30-34	8,412	8,080	5,413	2,708	1,909	1,574	1,345
35-39	9,825	9,926	7,005	3,617	2,955	2,602	2,381
40-44	8,462	8,612	6,279	3,532	2,953	2,926	2,484
45-49	5,163	5,575	4,286	2,600	2,389	2,347	2,284
50-54	2,816	2,939	2,225	1,379	1,360	1,513	1,522
55-59	1,512	1,590	1,208	778	750	814	745
60-64	859	871	696	470	441	410	457
65-69	427	481	355	282	226	267	242
≥70	270	277	272	181	192	203	224
Subtotal	41,322	41,621	29,797	16,683	13,946	13,307	12,171
Females							
13-24	247	257	182	129	109	108	73
25-29	784	835	635	376	332	289	284
30-34	1,610	1,667	1,316	801	683	632	580
35-39	1,728	1,824	1,572	992	857	874	821
40-44	1,310	1,511	1,323	895	803	920	825
45-49	688	853	788	568	543	600	630
50-54	353	382	344	285	268	342	321
55-59	215	246	192	143	169	159	178
60-64	118	136	142	84	92	102	99
65-69	73	89	75	42	53	60	60
≥70	83	82	63	66	55	45	68
Subtotal	7,209	7,882	6,632	4,381	3,964	4,131	3,939
Total	48,531	49,503	36,429	21,064	17,909	17,438	16,110

Note. These numbers do not represent the actual number of deaths in persons with an AIDS diagnosis. Rather, these numbers are point estimates of deaths, adjusted for reporting delays of death, but not for incomplete reporting of deaths. Subtotals and totals were calculated independently; values may not sum to subtotals or totals due to rounding.

^aThe 50 states and the District of Columbia (see Technical Notes).