

Table 14. Number of deaths in adolescents and adults with AIDS, by age at death, sex, and exposure category — United States^a, 2000

Age at death (yrs)	Number, by exposure category					Total
	Male-male sexual contact	Injection drug use	Male-male sexual contact and injection drug use	Heterosexual contact	Other	
Males						
13-24	51	13	4	12	21	101
25-29	245	53	35	44	10	387
30-34	790	248	152	138	16	1,345
35-39	1,301	549	285	214	32	2,381
40-44	1,123	850	278	201	31	2,484
45-49	922	950	208	173	32	2,284
50-54	594	639	103	161	26	1,522
55-59	341	238	47	105	13	745
60-64	207	135	20	86	9	457
65-69	105	59	11	58	8	242
≥70	109	46	4	53	12	224
Subtotal	5,788	3,781	1,147	1,244	211	12,171
Females						
13-24		18		51	4	73
25-29		73		206	5	284
30-34		232		336	11	580
35-39		402		405	13	821
40-44		463		351	12	825
45-49		372		245	13	630
50-54		166		145	10	321
55-59		69		100	10	178
60-64		21		72	6	99
65-69		17		38	4	60
≥70		14		45	9	68
Subtotal		1,847		1,995	97	3,939
Total	5,788	5,628	1,147	3,239	309	16,110

Note. These numbers do not represent the actual number of deaths in persons with an AIDS diagnosis. Rather, these numbers are point estimates of deaths, adjusted for reporting delays of death and for redistribution of cases initially reported with no identified risk, but not for incomplete reporting of deaths. Subtotals and totals were calculated independently; values may not sum to subtotals or totals due to rounding.

^aThe 50 states and the District of Columbia (see Technical Notes).