

Table 21. Number of adolescents and adults living with AIDS, by age as of December 31, 2002, sex, and exposure category — United States^a, 2000

Age as of 12/31/00 (yrs)	Number, by exposure category					Total
	Male-male sexual contact	Injection drug use	Male-male sexual contact and injection drug use	Heterosexual contact	Other	
Males						
13-24	1,309	252	121	288	350	2,320
25-29	5,837	1,143	689	1,094	325	9,087
30-34	19,737	4,627	2,537	2,926	517	30,344
35-39	37,448	10,405	5,469	4,799	719	58,840
40-44	33,413	14,876	5,306	4,666	723	58,985
45-49	23,399	14,412	3,444	3,380	638	45,273
50-54	14,353	8,636	1,728	2,135	368	27,220
55-59	7,335	3,199	696	1,292	237	12,759
60-64	3,564	1,312	248	850	142	6,116
65-69	1,670	562	90	494	110	2,925
≥70	1,108	346	56	353	121	1,985
Subtotal	149,172	59,770	20,384	22,278	4,250	255,854
Females						
13-24		333		1,446	176	1,955
25-29		1,114		3,668	103	4,884
30-34		3,324		6,870	237	10,431
35-39		6,154		8,648	372	15,174
40-44		6,810		7,346	402	14,558
45-49		5,101		4,622	294	10,017
50-54		2,306		2,626	201	5,133
55-59		779		1,494	117	2,390
60-64		292		811	85	1,188
65-69		130		467	60	657
≥70		98		372	84	553
Subtotal		26,441		38,369	2,132	66,941
Total	149,173	86,211	20,384	60,647	6,382	322,796

Note. These numbers do not represent the actual number of adolescents and adults living with AIDS. Rather, these numbers are point estimates derived by subtracting the cumulative number of deaths in persons with AIDS from the estimated cumulative number of persons with an AIDS diagnosis. The estimated number of new AIDS diagnoses and the estimated number of deaths are adjusted for reporting delays and for redistribution of cases initially reported with no identified risk, but not for incomplete reporting. Subtotals and totals were calculated independently; values may not sum to subtotals or totals due to rounding.

^aThe 50 states and the District of Columbia (see Technical Notes).