

Table 7. Number of new AIDS diagnoses in adolescents and adults, by age at AIDS diagnosis, sex, and exposure category — United States^a, 2000

Number, by exposure category						
Age at AIDS diagnosis (yrs)	Male-male sexual contact	Injection drug use	Male-male sexual contact and injection drug use	Heterosexual contact	Other	Total
Males						
13-24	581	109	52	144	34	919
25-29	1,490	302	131	314	27	2,265
30-34	2,958	811	312	627	41	4,748
35-39	4,028	1,454	487	852	79	6,900
40-44	2,779	1,714	375	744	55	5,668
45-49	1,773	1,422	227	516	55	3,993
50-54	1,034	810	126	357	23	2,351
55-59	555	326	52	193	20	1,145
60-64	294	145	17	144	18	617
65-69	142	61	10	88	9	310
≥70	115	52	9	57	12	246
Subtotal	15,749	7,206	1,799	4,035	372	29,162
Females						
13-24		128		542	36	705
25-29		243		892	25	1,160
30-34		569		1,213	45	1,827
35-39		777		1,286	50	2,113
40-44		763		1,093	48	1,903
45-49		500		672	29	1,201
50-54		219		379	28	626
55-59		86		228	15	328
60-64		49		140	11	199
65-69		25		79	9	112
≥70		16		52	6	74
Subtotal		3,373		6,575	301	10,250
Total	15,749	10,580	1,799	10,611	674	39,412

Note. These numbers do not represent actual cases in persons with an AIDS diagnosis. Rather, these numbers are point estimates of diagnoses in adolescents and adults, adjusted for reporting delays and for redistribution of cases initially reported with no identified risk, but not for incomplete reporting. Subtotals and totals were calculated independently; values may not sum to subtotals or totals due to rounding.

^aThe 50 states and the District of Columbia (see Technical Notes).