

**Table 11. Reported cases of HIV/AIDS for male adults and adolescents, by race/ethnicity and exposure category, 1999–2002—30 areas with confidential name-based HIV infection reporting**

Exposure category	Year of report			
	1999	2000	2001	2002
<b>White, not Hispanic</b>				
Male-to-male sexual contact	5,480	4,859	5,063	5,095
Injection drug use	695	549	572	475
Male-to-male sexual contact and injection drug use	553	469	369	358
Hemophilia/coagulation disorder	36	29	24	17
Heterosexual contact	368	296	308	242
<i>Sex with injection drug user</i>	84	82	56	59
<i>Sex with person with hemophilia</i>	0	2	0	0
<i>Sex with HIV-infected transfusion recipient</i>	2	1	3	4
<i>Sex with HIV-infected person, risk not specified</i>	282	211	249	179
Receipt of blood transfusion, blood components, or tissue	40	32	34	31
Other/risk not reported or identified <sup>a</sup>	1,297	1,263	1,264	1,776
<b>Total</b>	<b>8,469</b>	<b>7,497</b>	<b>7,634</b>	<b>7,994</b>
<b>Black, not Hispanic</b>				
Male-to-male sexual contact	3,052	2,810	2,877	2,716
Injection drug use	1,364	1,185	933	787
Male-to-male sexual contact and injection drug use	334	238	249	188
Hemophilia/coagulation disorder	14	5	7	4
Heterosexual contact	1,762	1,649	1,517	1,254
<i>Sex with injection drug user</i>	279	225	215	147
<i>Sex with person with hemophilia</i>	6	1	3	1
<i>Sex with HIV-infected transfusion recipient</i>	9	5	8	12
<i>Sex with HIV-infected person, risk not specified</i>	1,468	1,418	1,291	1,094
Receipt of blood transfusion, blood components, or tissue	43	38	24	29
Other/risk not reported or identified <sup>a</sup>	3,307	3,048	3,282	3,915
<b>Total</b>	<b>9,876</b>	<b>8,973</b>	<b>8,889</b>	<b>8,893</b>
<b>Hispanic</b>				
Male-to-male sexual contact	1,134	1,062	1,115	1,224
Injection drug use	335	314	324	257
Male-to-male sexual contact and injection drug use	96	78	68	57
Hemophilia/coagulation disorder	1	2	0	2
Heterosexual contact	241	209	242	207
<i>Sex with injection drug user</i>	42	42	32	34
<i>Sex with person with hemophilia</i>	0	0	0	1
<i>Sex with HIV-infected transfusion recipient</i>	0	3	2	2
<i>Sex with HIV-infected person, risk not specified</i>	199	164	208	170
Receipt of blood transfusion, blood components, or tissue	8	4	5	8
Other/risk not reported or identified <sup>a</sup>	604	654	677	787
<b>Total</b>	<b>2,419</b>	<b>2,323</b>	<b>2,431</b>	<b>2,542</b>
<b>Asian/Pacific Islander</b>				
Male-to-male sexual contact	60	42	58	64
Injection drug use	4	4	7	2
Male-to-male sexual contact and injection drug use	1	1	3	2
Hemophilia/coagulation disorder	0	0	0	1
Heterosexual contact	13	10	9	8
<i>Sex with injection drug user</i>	3	0	0	0
<i>Sex with person with hemophilia</i>	0	0	0	0
<i>Sex with HIV-infected transfusion recipient</i>	0	0	1	0
<i>Sex with HIV-infected person, risk not specified</i>	10	10	8	8
Receipt of blood transfusion, blood components, or tissue	1	0	1	0
Other/risk not reported or identified <sup>a</sup>	21	44	37	45
<b>Total</b>	<b>100</b>	<b>101</b>	<b>115</b>	<b>122</b>

**Table 11. (continued)**

Exposure category	Year of report			
	1999	2000	2001	2002
	<b>American Indian/Alaska Native</b>			
Male-to-male sexual contact	77	65	55	68
Injection drug use	17	15	17	15
Male-to-male sexual contact and injection drug use	19	10	21	8
Hemophilia/coagulation disorder	0	0	0	0
Heterosexual contact	13	10	11	8
<i>Sex with injection drug user</i>	3	3	5	3
<i>Sex with person with hemophilia</i>	0	0	0	0
<i>Sex with HIV-infected transfusion recipient</i>	1	0	0	0
<i>Sex with HIV-infected person, risk not specified</i>	9	7	6	5
Receipt of blood transfusion, blood components, or tissue	1	0	1	0
Other/risk not reported or identified <sup>a</sup>	22	13	17	18
<b>Total</b>	<b>149</b>	<b>113</b>	<b>122</b>	<b>117</b>
<b>Total cases<sup>b</sup></b>	<b>21,112</b>	<b>19,143</b>	<b>19,406</b>	<b>19,678</b>

Data include male adults and adolescents with a diagnosis of HIV infection. This includes male adults and adolescents with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1998, the following 30 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Arkansas, Arizona, Florida, Colorado, Iowa, Idaho, Indiana, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, New Jersey, New Mexico, Nevada, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Utah, Virginia, Wisconsin, West Virginia, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

<sup>a</sup> Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

<sup>b</sup> Includes male adults and adolescents of unknown race or multiple races.