

Table 15. Reported cases of AIDS for female adults and adolescents by race/ethnicity and exposure category, 1998–2002—United States

Exposure category	Year of report					Cumulative through 2002 ^a
	1998	1999	2000	2001	2002	
White, not Hispanic						
Injection drug use	762	672	641	615	583	13,035
Hemophilia/coagulation disorder	5	6	0	4	2	114
Heterosexual contact	847	833	752	770	787	12,995
<i>Sex with injection drug user</i>	280	275	258	218	225	5,063
<i>Sex with bisexual male</i>	77	87	55	55	49	1,642
<i>Sex with person with hemophilia</i>	19	8	8	5	10	302
<i>Sex with HIV-infected transfusion recipient</i>	9	9	12	3	12	330
<i>Sex with HIV-infected person, risk not specified</i>	462	454	419	489	491	5,658
Receipt of blood transfusion, blood components, or tissue	37	28	44	23	17	1,857
Other/risk not reported or identified ^b	346	358	422	582	542	4,011
Total	1,997	1,897	1,859	1,994	1,931	32,012
Black, not Hispanic						
Injection drug use	2,065	1,890	1,623	1,429	1,394	34,041
Hemophilia/coagulation disorder	13	8	4	4	8	122
Heterosexual contact	2,882	2,942	2,890	3,039	3,063	35,748
<i>Sex with injection drug user</i>	776	684	655	576	555	11,861
<i>Sex with bisexual male</i>	120	90	106	131	124	1,724
<i>Sex with person with hemophilia</i>	6	8	5	4	4	95
<i>Sex with HIV-infected transfusion recipient</i>	14	10	13	12	16	198
<i>Sex with HIV-infected person, risk not specified</i>	1,966	2,150	2,111	2,316	2,364	21,870
Receipt of blood transfusion, blood components, or tissue	64	81	82	71	76	1,434
Other/risk not reported or identified ^b	1,679	1,798	1,897	2,428	2,812	17,896
Total	6,703	6,719	6,496	6,971	7,353	89,241
Hispanic						
Injection drug use	616	577	540	437	386	11,104
Hemophilia/coagulation disorder	4	3	0	2	1	56
Heterosexual contact	1,050	980	879	914	837	14,001
<i>Sex with injection drug user</i>	311	290	206	206	193	5,827
<i>Sex with bisexual male</i>	48	22	35	26	30	617
<i>Sex with person with hemophilia</i>	6	2	3	1	1	43
<i>Sex with HIV-infected transfusion recipient</i>	1	5	5	3	4	107
<i>Sex with HIV-infected person, risk not specified</i>	684	661	630	678	609	7,407
Receipt of blood transfusion, blood components, or tissue	18	17	17	14	19	577
Other/risk not reported or identified ^b	347	356	409	504	629	3,634
Total	2,035	1,933	1,845	1,871	1,872	29,372

Table 15. (continued)

Exposure category	Year of report					Cumulative through 2002 ^a
	1998	1999	2000	2001	2002	
Asian/Pacific Islander						
Injection drug use	6	7	3	7	4	121
Hemophilia/coagulation disorder	0	0	0	1	0	7
Heterosexual contact	29	31	36	32	29	410
<i>Sex with injection drug user</i>	8	4	2	5	2	93
<i>Sex with bisexual male</i>	5	2	3	4	2	77
<i>Sex with person with hemophilia</i>	0	0	0	0	0	4
<i>Sex with HIV-infected transfusion recipient</i>	1	0	1	1	0	21
<i>Sex with HIV-infected person, risk not specified</i>	15	25	30	22	25	215
Receipt of blood transfusion, blood components, or tissue	3	3	3	1	3	103
Other/risk not reported or identified ^b	19	20	34	27	33	200
Total	57	61	76	68	69	841
American Indian/Alaska Native						
Injection drug use	17	12	29	16	13	221
Hemophilia/coagulation disorder	0	0	1	0	0	3
Heterosexual contact	10	15	29	16	20	199
<i>Sex with injection drug user</i>	4	5	9	9	9	89
<i>Sex with bisexual male</i>	1	1	6	1	0	26
<i>Sex with person with hemophilia</i>	0	1	0	0	0	2
<i>Sex with HIV-infected transfusion recipient</i>	1	2	0	0	0	3
<i>Sex with HIV-infected person, risk not specified</i>	4	6	14	6	11	79
Receipt of blood transfusion, blood components, or tissue	0	1	1	0	2	15
Other/risk not reported or identified ^b	4	12	10	10	7	71
Total	31	40	70	42	42	509
Total cases^c	10,832	10,656	10,350	10,955	11,279	152,060

^a Includes female adults and adolescents with a diagnosis AIDS, reported from the beginning of the epidemic through 2002.

^b Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

^c Includes female adults and adolescents of unknown race or multiple races.