

Table 5. Estimated proportions of adults and adolescents with HIV/AIDS for whom a CD4 test was performed within 12 months after HIV diagnosis, by year of diagnosis, disease category, and sex, 2001–2003—33 states with confidential name-based HIV infection reporting

	2001			2002			2003		
	CD4 count ^a	No CD4 count	Total	CD4 count ^a	No CD4 count	Total	CD4 count ^a	No CD4 count	Total
	No. (%)	No. (%)	No.	No. (%)	No. (%)	No.	No. (%)	No. (%)	No.
HIV without AIDS^b									
Males	5,380 (30.7)	12,168 (69.3)	17,548	5,349 (32.4)	11,168 (67.6)	16,516	5,343 (33.3)	10,688 (66.7)	16,030
Females	2,392 (28.7)	5,939 (71.3)	8,330	2,265 (29.8)	5,338 (70.2)	7,603	2,209 (32.0)	4,698 (68.0)	6,907
Subtotal	7,772 (30.0)	18,106 (70.0)	25,878	7,614 (31.6)	16,505 (68.4)	24,119	7,551 (32.9)	15,386 (67.1)	22,937
HIV to AIDS^c									
Males	4,224 (96.2)	165 (3.8)	4,390	4,446 (97.0)	135 (3.0)	4,582	4,815 (97.9)	104 (2.1)	4,919
Females	1,648 (96.5)	60 (3.5)	1,708	1,636 (97.6)	40 (2.4)	1,676	1,744 (98.4)	28 (1.6)	1,772
Subtotal	5,873 (96.3)	225 (3.7)	6,098	6,082 (97.2)	176 (2.8)	6,257	6,559 (98.0)	132 (2.0)	6,691
HIV with AIDS^d									
Males	6,130 (92.6)	488 (7.4)	6,618	6,100 (93.6)	419 (6.4)	6,519	5,925 (94.8)	326 (5.2)	6,251
Females	2,046 (92.6)	163 (7.4)	2,209	1,846 (93.1)	138 (6.9)	1,984	1,896 (94.8)	105 (5.2)	2,001
Subtotal	8,176 (92.6)	651 (7.4)	8,827	7,946 (93.5)	557 (6.5)	8,503	7,821 (94.8)	430 (5.2)	8,252

Note. These numbers do not represent actual cases in persons with a diagnosis of HIV infection or AIDS. Rather, these numbers are point estimates of case counts that have been adjusted for reporting delays. The estimates have not been adjusted for incomplete reporting.

Since 2000, the following states have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, and Wyoming. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

^a When only CD4 percentage was available, a CD4 count was interpreted.

^b For at least 12 months after HIV diagnosis.

^c AIDS diagnosed 1–12 months after HIV diagnosis.

^d HIV and AIDS diagnosed during the same calendar month.