

**Table 6. Median results of the first CD4 test performed within 12 months after HIV diagnosis among adults and adolescents, by year of diagnosis, disease category, and sex, 2001–2003—33 states with confidential name-based HIV infection reporting**

	2001			2002			2003		
	Median CD4 count <sup>a</sup>	25%–75%	Total No.	Median CD4 count <sup>a</sup>	25%–75%	Total No.	Median CD4 count <sup>a</sup>	25%–75%	Total No.
<b>HIV without AIDS<sup>b</sup></b>									
Males	457	341–640	5,380	470	350–644	5,349	445	336–610	5,343
Females	455	345–640	2,392	469	340–664	2,265	460	343–655	2,209
Subtotal	456	342–640	7,772	470	348–649	7,614	450	338–626	7,551
<b>HIV to AIDS<sup>c</sup></b>									
Males	117	38–184	4,224	113	38–182	4,446	117	40–182	4,815
Females	129	48–187	1,648	130	48–182	1,636	119	45–182	1,744
Subtotal	121	40–185	5,873	117	41–182	6,082	117	41–182	6,559
<b>HIV with AIDS<sup>d</sup></b>									
Males	62	20–146	6,130	58	20–140	6,100	59	20–137	5,925
Females	62	20–151	2,046	68	20–152	1,846	60	20–143	1,896
Subtotal	62	20–147	8,176	60	20–143	7,946	60	20–138	7,821

Note. These numbers do not represent actual cases in persons with a diagnosis of HIV infection or AIDS. Rather, these numbers are point estimates of case counts that have been adjusted for reporting delays. The estimates have not been adjusted for incomplete reporting.

Since 2000, the following states have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, and Wyoming. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

<sup>a</sup> When only CD4 percentage was available, a CD4 count was interpreted.

<sup>b</sup> For at least 12 months after HIV diagnosis.

<sup>c</sup> AIDS diagnosed 1–12 months after HIV diagnosis.

<sup>d</sup> HIV and AIDS diagnosed during the same calendar month.