

Table 9a. Estimated distribution of reported results of laboratory tests performed within 12 months after HIV diagnosis among adults and adolescents, by year of diagnosis, disease category, and sex, 2001–2003—33 states with confidential name-based HIV infection reporting

	CD4 only ^a No. (%)	CD4 & viral load No. (%)	Viral load only No. (%)	No lab test No. (%)	Total No ^b
2001					
HIV without AIDS^c					
Males	1,685 (9.6)	3,695 (21.1)	3,419 (19.5)	8,749 (49.9)	17,548
Females	736 (8.8)	1,655 (19.9)	1,746 (21.0)	4,193 (50.3)	8,330
HIV to AIDS^d					
Males	1,687 (38.4)	2,537 (57.8)	30 (0.7)	135 (3.1)	4,390
Females	607 (35.6)	1,041 (60.9)	10 (0.6)	50 (3.0)	1,708
HIV with AIDS^e					
Males	2,591 (39.2)	3,539 (53.5)	69 (1.0)	419 (6.3)	6,618
Females	867 (39.2)	1,179 (53.4)	21 (1.0)	142 (6.4)	2,209
Total^f	8,174 (20.0)	13,646 (33.4)	5,296 (13.0)	13,687 (33.5)	40,804
2002					
HIV without AIDS^c					
Males	1,414 (8.6)	3,935 (23.8)	2,849 (17.2)	8,319 (50.4)	16,516
Females	620 (8.1)	1,645 (21.6)	1,448 (19.0)	3,890 (51.2)	7,603
HIV to AIDS^d					
Males	1,729 (37.7)	2,717 (59.3)	34 (0.7)	102 (2.2)	4,582
Females	625 (37.3)	1,011 (60.3)	11 (0.6)	29 (1.7)	1,676
HIV with AIDS^e					
Males	2,476 (38.0)	3,624 (55.6)	62 (1.0)	357 (5.5)	6,519
Females	761 (38.3)	1,086 (54.7)	15 (0.8)	122 (6.2)	1,984
Total^f	7,624 (19.6)	14,018 (36.1)	4,419 (11.4)	12,819 (33.0)	38,880
2003					
HIV without AIDS^c					
Males	1,350 (8.4)	3,993 (24.9)	2,395 (14.9)	8,293 (51.7)	16,030
Females	515 (7.5)	1,693 (24.5)	1,159 (16.8)	3,539 (51.2)	6,907
HIV to AIDS^d					
Males	1,712 (34.8)	3,103 (63.1)	36 (0.7)	68 (1.4)	4,919
Females	593 (33.5)	1,151 (64.9)	3 (0.2)	26 (1.4)	1,772
HIV with AIDS^e					
Males	2,128 (34.0)	3,798 (60.8)	55 (0.9)	270 (4.3)	6,251
Females	733 (36.6)	1,164 (58.1)	16 (0.8)	89 (4.4)	2,001
Total^f	7,031 (18.6)	14,901 (39.3)	3,664 (9.7)	12,285 (32.4)	37,880

Note. These numbers do not represent actual cases in persons with a diagnosis of HIV infection or AIDS. Rather, these numbers are point estimates of case counts that have been adjusted for reporting delays. The estimates have not been adjusted for incomplete reporting.

Since 2000, the following states have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, and Wyoming. Since July 1997, Florida has had confidential name-based HIV infection reporting for new diagnoses only.

^a Count or percentage or both.

^b Because row totals were calculated independently of the row values, the values in each row may not sum to the row total.

^c For at least 12 months after HIV diagnosis.

^d AIDS diagnosed 1–12 months after HIV diagnosis.

^e HIV and AIDS diagnosed during the same calendar month.

^f Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.