

Table 4. Estimated numbers of AIDS cases, by year of diagnosis and selected characteristics, 2003–2007 and cumulative—United States and dependent areas

	Year of diagnosis					Cumulative ^a
	2003	2004	2005	2006	2007	
Data for 50 states and the District of Columbia						
Age at diagnosis (yr)						
<13	73	55	54	38	28	9,209
13–14	72	71	70	71	80	1,169
15–19	302	333	409	392	455	6,089
20–24	1,577	1,635	1,669	1,603	1,927	38,175
25–29	3,073	3,191	3,071	3,283	3,380	120,464
30–34	5,578	5,126	4,637	4,200	4,187	201,906
35–39	8,096	7,050	6,417	6,185	5,888	219,601
40–44	7,708	7,687	7,261	7,106	6,813	177,250
45–49	5,676	5,506	5,662	5,456	5,749	112,896
50–54	3,393	3,466	3,472	3,578	3,636	63,408
55–59	1,711	1,830	1,839	2,005	2,040	34,160
60–64	865	898	856	949	980	18,249
≥65	770	786	711	829	800	15,853
Race/ethnicity						
American Indian/Alaska Native	181	184	164	148	158	3,492
Asian ^b	394	389	378	425	475	7,511
Black/African American	19,580	18,719	17,690	17,257	17,507	426,003
Hispanic/Latino ^c	7,214	6,817	6,804	6,875	6,921	169,138
Native Hawaiian/Other Pacific Islander	52	51	58	61	76	721
White	11,061	11,064	10,580	10,521	10,407	404,465
Transmission category						
Male adult or adolescent						
Male-to-male sexual contact	16,782	16,627	16,172	16,235	16,749	487,695
Injection drug use	5,098	4,527	4,243	3,940	3,750	175,704
Male-to-male sexual contact and injection drug use	2,129	1,964	1,972	1,748	1,664	71,242
High-risk heterosexual contact ^d	4,140	4,204	3,909	4,054	4,011	63,927
Other ^e	220	222	230	209	181	12,108
Subtotal	28,370	27,545	26,525	26,185	26,355	810,676
Female adult or adolescent						
Injection drug use	3,002	2,884	2,604	2,331	2,260	80,155
High-risk heterosexual contact ^d	7,247	6,956	6,768	6,955	7,100	112,230
Other ^e	202	193	176	186	220	6,158
Subtotal	10,450	10,033	9,548	9,471	9,579	198,544
Child (<13 yrs at diagnosis)						
Perinatal	66	53	48	33	24	8,434
Other ^f	7	2	5	6	4	775
Subtotal	73	55	54	38	28	9,209
Region of residence						
Northeast	10,432	9,349	9,115	9,143	8,973	314,277
Midwest	4,264	4,074	4,328	4,082	4,074	105,573
South	17,643	18,089	16,641	16,271	16,683	390,479
West	6,555	6,122	6,043	6,199	6,232	208,099
Subtotal for 50 states and the District of Columbia	38,893	37,633	36,127	35,695	35,962	1,018,428
Data for U.S. dependent areas	1,085	917	955	832	812	32,051
Total^g	40,054	38,695	37,256	36,791	37,041	1,051,875^h

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and missing risk-factor information, but not for incomplete reporting.

^a From the beginning of the epidemic through 2007.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^h Includes 7,099 persons of unknown race or multiple races, 1,393 persons of unknown state of residence, and 3 persons who were residents of other areas.