

Table 8. Estimated numbers of deaths of persons with AIDS, by year of death and selected characteristics, 2003–2007 and cumulative—United States and dependent areas

	Year of death					Cumulative ^a
	2003	2004	2005	2006	2007	
Data for 50 states and the District of Columbia						
Age at death (yr)						
<13	24	14	6	16	5	4,891
13–14	7	15	10	4	17	292
15–19	36	34	38	44	41	1,143
20–24	162	173	143	164	155	8,880
25–29	522	489	452	437	440	44,219
30–34	1,298	1,147	1,062	841	766	96,379
35–39	2,821	2,443	2,078	1,815	1,600	118,886
40–44	3,582	3,434	3,316	2,857	2,660	107,417
45–49	3,307	3,283	3,287	3,103	2,865	76,249
50–54	2,442	2,543	2,622	2,494	2,411	47,022
55–59	1,368	1,425	1,558	1,532	1,476	26,767
60–64	721	738	798	813	817	15,244
≥65	792	830	879	868	858	15,404
Race/ethnicity						
American Indian/Alaska Native	75	85	70	79	70	1,792
Asian ^b	73	92	71	99	84	3,114
Black/African American	8,926	8,656	8,546	7,886	7,124	226,879
Hispanic/Latino ^c	2,627	2,601	2,450	2,256	2,312	82,894
Native Hawaiian/Other Pacific Islander	5	17	14	4	11	291
White	5,231	4,963	4,933	4,398	4,187	245,127
Transmission category						
Male adult or adolescent						
Male-to-male sexual contact	6,131	5,896	5,888	5,329	5,373	274,184
Injection drug use	3,655	3,356	3,245	2,820	2,397	112,068
Male-to-male sexual contact and injection drug use	1,343	1,262	1,318	1,154	1,054	42,551
High-risk heterosexual contact ^d	1,436	1,514	1,467	1,543	1,433	25,860
Other ^e	163	137	126	98	83	8,728
Subtotal	12,728	12,166	12,044	10,945	10,339	463,392
Female adult or adolescent						
Injection drug use	1,955	1,966	1,816	1,594	1,446	46,624
High-risk heterosexual contact ^d	2,257	2,312	2,283	2,321	2,211	43,432
Other ^e	86	71	70	75	57	3,928
Subtotal	4,298	4,350	4,169	3,991	3,714	93,984
Child (<13 yrs at diagnosis)						
Perinatal	50	52	33	46	50	4,842
Other ^f	7	2	3	7	7	575
Subtotal	57	54	37	53	57	5,417

Table 8. Estimated numbers of deaths of persons with AIDS, by year of death and selected characteristics, 2003–2007 and cumulative—United States and dependent areas (cont)

	Year of death					Cumulative ^a
	2003	2004	2005	2006	2007	
Region of residence						
Northeast	4,992	4,708	4,435	4,076	3,463	183,292
Midwest	1,700	1,491	1,387	1,478	1,368	56,137
South	7,735	7,678	7,812	7,282	7,080	206,654
West	2,655	2,693	2,615	2,152	2,200	116,710
Subtotal for 50 states and the District of Columbia	17,082	16,570	16,249	14,989	14,110	562,793
Data for U.S. dependent areas	574	568	535	541	403	20,178
Total^g	17,679	17,154	16,823	15,564	14,561	583,298^h

Note. These numbers do not represent reported death counts. Rather, these numbers are point estimates, which result from adjustments of reported death counts. The reported death counts have been adjusted for reporting delays and missing risk-factor information, but not for incomplete reporting.

^a From the beginning of the epidemic through 2007.

^b Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^c Hispanics/Latinos can be of any race.

^d Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^e Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^f Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^g Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

^h Includes 2,704 persons of unknown race or multiple races, 325 persons of unknown state of residence, and 2 persons who were residents of other areas.