

Table 8. Number and percentage of HIV-infected women with positive test results for selected conditions, by year of infant's birth, 2000–2003—24 areas conducting enhanced perinatal surveillance in the United States and dependent areas

Diagnosis	Year of infant's birth								Cumulative Total	
	2000		2001		2002		2003			
	No.	%	No.	%	No.	%	No.	%	No.	%
Sexually transmitted disease^{a,b}										
Yes	783	23	769	26	512	26	351	22	2,415	24
No	1,286	38	1,251	42	852	43	581	37	3,970	40
Unknown	1,353	40	943	32	623	31	629	40	3,548	36
Selected condition^c										
Yes	615	18	640	22	438	22	349	21	2,042	20
No	1,341	39	1,253	42	887	45	626	38	4,107	41
Unknown	1,466	43	1,070	36	662	33	661	40	3,859	39
Total	3,422		2,963		1,987		1,636		10,008	

Note. Because of rounding, column percentages may not total to 100 percent.

^a Includes presumptive or definitive diagnoses during pregnancy of the following sexually transmitted disease: Chlamydia, genital herpes (primary herpes and active lesions), gonorrhea, hepatitis B, syphilis, and *Trichomonas*.

^b Excludes 75 cases with missing information.

^c Includes presumptive or definitive diagnoses during pregnancy of the following conditions: bacterial vaginosis, group B strep, hepatitis C, or pelvic inflammatory disease.