

**Table 2. Prenatal care of HIV-infected women, by year of infant's birth, 2000–2003—24 areas conducting enhanced perinatal surveillance in the United States and dependent areas**

	Year of infant's birth								Cumulative Total	
	2000		2001		2002		2003			
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Prenatal care</b>										
Yes	3,002	88	2,606	88	1,778	89	1,449	89	8,835	88
No	270	8	235	8	144	7	115	7	764	8
Unknown	150	4	122	4	65	3	72	4	409	4
<b>Prenatal care visits<sup>a</sup></b>										
1–2 visits	119	4	110	4	70	4	87	6	386	4
≥3 visits	1,905	63	1,666	64	1,033	58	1,213	84	5,817	66
Unknown	978	33	830	32	675	38	149	10	2,632	30
Subtotal	3,002		2,606		1,778		1,449		8,835	
<b>Total</b>	<b>3,422</b>		<b>2,963</b>		<b>1,987</b>		<b>1,636</b>		<b>10,008</b>	

Note. Because of rounding, column percentages may not total to 100 percent.

<sup>a</sup> Includes only women who had documentation of prenatal care in their records.

**Table 3. Timing of HIV testing of HIV-infected women, by year of infant's birth, 2000–2003—24 areas conducting enhanced perinatal surveillance in the United States and dependent areas**

	Year of infant's birth								Cumulative Total	
	2000		2001		2002		2003			
	No.	%	No.	%	No.	%	No.	%	No.	%
<b>Timing of mother's HIV test</b>										
Before pregnancy	1,975	58	1,762	59	1,207	61	1,030	63	5,974	60
During labor	1,062	31	906	31	604	30	459	28	3,031	30
At delivery	154	5	124	4	60	3	46	3	384	4
After birth	94	3	72	2	26	1	36	2	228	2
Unknown	136	4	96	3	79	4	65	4	376	4
<b>Total<sup>a</sup></b>	<b>3,422</b>		<b>2,963</b>		<b>1,987</b>		<b>1,636</b>		<b>10,008</b>	

Note. Because of rounding, column percentages may not total to 100 percent.

<sup>a</sup> Includes 15 women who refused HIV testing.