



MMWRTM

Morbidity and Mortality Weekly Report

Surveillance Summaries

May 19, 2006 / Vol. 55 / No. SS-3

Youth Tobacco Surveillance — United States, 2001–2002



**DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION**

The *MMWR* series of publications is published by the Coordinating Center for Health Information and Service, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

General: Centers for Disease Control and Prevention. Youth Tobacco Surveillance — United States, 2001–2002. In: Surveillance Summaries, May 19, 2006. *MMWR* 2006;55(No. SS-3).

Specific: [Author(s)]. [Title of particular article]. In: Surveillance Summaries, May 19, 2006. *MMWR* 2006;55(No. SS-3):[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, MD, MPH
Director

Dixie E. Snider, MD, MPH
Chief Science Officer

Tanja Popovic, MD, PhD
Associate Director for Science

Coordinating Center for Health Information and Service

Steven L. Solomon, MD
Director

National Center for Health Marketing

Jay M. Bernhardt, PhD, MPH
Director

Division of Scientific Communications

Judith R. Aguilar
(Acting) Director

Mary Lou Lindegren, MD
Editor, MMWR Series

Suzanne M. Hewitt, MPA
Managing Editor, MMWR Series

Teresa F. Rutledge
Lead Technical Writer-Editor

Jeffrey D. Sokolow, MA
Project Editor

Beverly J. Holland
Lead Visual Information Specialist

Lynda G. Cupell
Malbea A. LaPete
Visual Information Specialists

Quang M. Doan, MBA
Erica R. Shaver
Information Technology Specialists

CONTENTS

Introduction	2
Methods	2
Results	4
Discussion	20
Conclusion	23
References	24

Youth Tobacco Surveillance — United States, 2001–2002

LaTisha Marshall, MPH¹
Michael Schooley, MPH¹
Heather Ryan, MPH¹
Patrick Cox¹
Alyssa Easton, PhD¹
Cheryl Heaton, DrPH²
Kat Jackson, MSPH³
Kevin C. Davis, MS³
Ghada Homsy, ME³

¹Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC

²American Legacy Foundation, District of Columbia

³Research Triangle Institute, Research Triangle Park, North Carolina

Abstract

Problem/Condition: Cigarette smoking is the leading preventable cause of death in the United States, accounting for approximately 440,000 deaths each year. The prevalence of cigarette smoking nationwide among high school students (grades 9–12) increased during the 1990s, peaking during 1996–1997, and then declined. Approximately 80% of tobacco users initiate use before age 18 years. An estimated 6.4 million children aged <18 years who are living today will die prematurely as adults because they began to smoke cigarettes during adolescence. The annual health-related economic cost associated with tobacco use exceeds \$167 billion. Because of these health and economic consequences, CDC has recommended that states establish and maintain comprehensive tobacco-control programs to reduce tobacco use among youth.

Reporting Period: This report covers data collected during January 2001–December 2002.

Description of the System: The National Youth Tobacco Survey (NYTS) and state youth tobacco surveys (YTS) were developed to provide states with data to support the design, implementation, and evaluation of comprehensive tobacco-control programs. NYTS is representative of middle and high school students in the 50 states and the District of Columbia. During spring 2002, a total of 26,149 students in 246 schools completed NYTS questionnaires. Weighted data for the YTS were achieved by 13 states in 2001 and by 20 states in 2002; state sample sizes varied (range: 982–38,934). This report summarizes data from the 2002 NYTS and the 2001 and 2002 YTS.

Results and Interpretation: Findings from the 2002 NYTS indicate that current use of any tobacco product ranged from 13.3% among middle school students to 28.2% among high school students. Cigarette smoking was the most prevalent form of tobacco use, with 9.8% of middle school students and 22.5% of high school students reporting that they currently smoke cigarettes. Cigar smoking was the second most prevalent form of tobacco use, with 6.0% of middle school students and 11.6% of high school students reporting that they currently smoke cigars. Among current cigarette smokers, 41.8% of middle school students and 52.0% of high school students reported that they usually smoke Marlboro[®] cigarettes. Black middle school and high school students who smoke were more likely to smoke Newport[®] cigarettes than any other brand (58.3% and 66.8%, respectively). Among middle school students aged <18 years, 75.9% were not asked to show proof of age when they bought or tried to buy cigarettes, and 63.4% were not refused purchase because of their age. Among high school students aged <18 years, 58.5% were not asked to show proof of age when they bought or tried to buy cigarettes, and 60.6% were not refused purchase because of their age.

Nearly half (49.6%) of middle school students and 62.1% of high school students who smoke reported a desire to stop smoking cigarettes, with 55.4% of middle school students and 53.1% of high school students reported having made at least one cessation attempt during the 12 months preceding the survey. Among students who have never smoked cigarettes, 21.3% of middle school students and 22.9% of high school students were susceptible to initiating cigarette smoking in the next year.

Exposure to secondhand smoke (i.e., environmental tobacco smoke) was high. During the week before the survey, 1) 88.3% of middle school students and 91.4% of high school students who currently smoke cigarettes and 47.1% of middle school students and 53.3% of high

Corresponding author: LaTisha Marshall, MPH, Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC, 1600 Clifton Rd., NE, MS K-50, Atlanta, GA 30333. Telephone: 770-488-5902; Fax: 770-488-5848; E-mail: ln19@cdc.gov.

school students who have never smoked cigarettes were in the same room with someone who was smoking cigarettes; 2) 81.7% of middle school students and 83.7% of high school students who currently smoke cigarettes and 31.5% of middle school students and 29.1% of high school students who have never smoked cigarettes rode in a car with someone who was smoking cigarettes; and 3) 71.5% of middle school students and 57.5% of high school students who currently smoke cigarettes and 33.3% of middle school students and 29.9% of high school students who have never smoked cigarettes lived in a home in which someone else smoked cigarettes. Media and advertising influence was also noted, with 58.1% of middle school students and 54.9% of high school students who currently use tobacco and 11.0% of middle school students and 13.7% of high school students who have never used tobacco reporting that they would wear or use an item with a tobacco company name or logo on it. Although 84.6% of middle school students and 91.2% of high school students had seen or heard antismoking commercials on television or radio, 89.9% of middle school students and 91.3% of high school students also had seen actors using tobacco on television or in the movies.

Public Health Actions: Health and education officials use YTS and NYTS data to plan, evaluate, and improve national and state programs to prevent and control youth tobacco use. States can use these data in presentations to their state legislators to demonstrate the need for funding comprehensive tobacco-control programs, including tobacco cessation and prevention programs for youth.

Introduction

Cigarette smoking is the leading preventable cause of death in the United States (1), accounting for approximately 440,000 deaths each year (2). The prevalence of cigarette smoking nationwide among high school (grades 9–12) students increased during the 1990s (3), peaking during 1996–1997, then declined (4,5). Approximately 80% of tobacco users initiate use before age 18 years (6). An estimated 6.4 million children aged <18 years who are living today will die prematurely as adults because they began to smoke cigarettes during adolescence (7). The annual health-related economic costs associated with tobacco use exceed \$167 billion (2).

The National Youth Tobacco Survey (NYTS) and state youth tobacco surveys (YTS) were developed to provide states with the data necessary to support the design, implementation, and evaluation of a comprehensive tobacco-control program (8,9). Certain states have data regarding the prevalence of selected tobacco use behaviors among high schools students from the Youth Risk Behavior Surveillance System (YRBSS). YTS supplements YRBSS by providing more comprehensive data regarding tobacco use (bidis^{*}, cigarettes, cigars, kreteks[†], pipes, and smokeless tobacco); exposure to secondhand smoke; smoking cessation; school curriculum; minors' ability to purchase or obtain tobacco products; knowledge and attitudes about tobacco and familiarity with protobacco and antitobacco media messages; and by providing information regarding both middle school (grades 6–8) and high school students.

* Bidis (or beedies) are small brown cigarettes from India consisting of tobacco wrapped in a leaf and tied with a thread.

† Kreteks (also called clove cigarettes) are flavored cigarettes containing tobacco and clove extract.

First conducted during fall 1999 (10) and then again during spring 2000 and 2002, NYTS is representative of all middle school and high school students in the 50 states and the District of Columbia. Funding for the first 3 years of NYTS was provided by the American Legacy Foundation (ALF) (District of Columbia), and all surveys were conducted during the spring semester of even years by ORC Macro (Calverton, Maryland). Implementation of the spring 2004 NYTS was supported and directed by CDC's Office on Smoking and Health. YTS was first conducted in 1998, with three states participating. Since then, the number of participating states has increased substantially; by the end of 2002, a total of 45 states and DC had conducted the YTS at least once, and 42 states and the District of Columbia had obtained weighted data that would provide representative estimates. Certain states conduct surveys annually, whereas others do so every other year or periodically. This report summarizes data from the 2002 NYTS and the 2001 and 2002 YTS.

Methods

Sampling

NYTS

The 2002 NYTS consisted of two components: 1) a newly drawn sample of 215 middle schools and high schools and 2) a panel sample of 83 middle schools and high schools drawn from schools that were selected to participate in the 2000 NYTS. The newly drawn sample employed a three-stage cluster sample design. The first-stage sampling frame contained 1,307 primary sampling units (PSUs) consisting of large counties or groups of smaller, adjacent counties. From these PSUs,

20 strata were formed by the United States being divided into four census regions, and then PSUs in each region were categorized into one of five “truth” levels, from lowest (1) to highest (5). These levels each represent and measure a different level of exposure to ALF’s national truth[®] antitobacco campaign (11). Of these, 100 PSUs were selected with a probability proportional to size (PPS) method in which school enrollment was the measure of size. At the second sampling stage, 200 public and private schools that contained any or all of the eligible grades were selected from the 100 PSUs with a probability proportional to weighted school enrollment. In addition, 15 small schools were selected from 15 PSUs that were selected randomly from the sample PSUs. Schools with substantial numbers of Asian, black, and Hispanic students were sampled at higher rates than all other schools. At the third sampling stage, approximately five intact classes of a required subject (e.g., English or social studies) were randomly selected from a class schedule provided by each participating school. Class schedules were constructed to ensure that all students in the eligible grades were accounted for once and were not duplicated. All students in the selected classes were eligible to participate in the survey.

The 83 panel sample schools were selected from the same 20 strata. Of these schools, 69 participated in the 2002 NYTS. The same data collection protocols were followed in panel schools as in the newly selected schools. Including both newly drawn and panel schools together, 298 schools were selected initially. Two schools were subsequently determined to be ineligible and were not replaced. One school had been part of the panel sample that was slated for participation in the 2000 NYTS sample but did not in fact participate and thus was ineligible for the 2002 survey. Students at the second school were predominantly Spanish speaking; because no Spanish version of the 2002 NYTS was available, and a substantial number of students were unable to read English at a level that would permit participation, the school was deemed ineligible. After exclusion of these two schools, 296 schools were determined to be eligible to participate in the survey.

A weighting factor was applied to each student record to adjust for nonresponse and for varying probabilities of selection, including taking into account the correct probabilities of selection for newly selected schools versus panel schools and those resulting from oversampling Asian, black, and Hispanic students. The numbers of students in other racial/ethnic populations were too low for meaningful analysis.[§] Weights were adjusted to ensure that the weighted proportions of students in each grade matched national population

proportions. Final adjusted weights were scaled to ensure that the weighted count of students was equal to the total sample size. SUDAAN (11) was used to compute 95% confidence intervals (CIs). Two-sided t-tests were conducted to test for statistically significant differences between prevalence estimates. If the p-value was <0.05, then results were considered statistically significant. In certain instances, although CIs overlapped, the differences between the estimates were statistically significant on the basis of the t-tests. Only those comparisons that were determined to be statistically significant by the t-test are discussed in this report.

NYTS produced a nationally representative sample of public and private middle school and high school students in all 50 states and the District of Columbia. Of the 296 eligible schools, 246 participated, and 26,149 questionnaires were completed by middle school and high school students in 246 schools. The school response rate was 83.1%, and the student response rate was 90.6%, resulting in an overall response rate (the school response rate multiplied by the student response rate) of 75.3%.

YTS

YTS employed a two-stage cluster sample design. The first-stage sampling frame included separate lists of public middle schools and high schools containing any or all of the eligible grades. Schools were selected with a probability proportional to school enrollment size. The number of schools selected varied by state. At the second sampling stage, classes were randomly selected from a class schedule provided by each participating school. Class schedules were constructed to ensure that all students in the eligible grades were counted once and were not duplicated. All students in the selected classes were eligible to participate in the survey. The number of classes selected varied by state. SAS (12) and SUDAAN (13) were used to compute 95% CIs.

A total of 40 states conducted YTS in either 2001 or 2002; one state, Florida, conducted the survey during both years. Of the 41 state surveys conducted during 2001–2002, a total of 11 (Florida, Idaho, Louisiana, Maine, Michigan, Nevada, Pennsylvania, Rhode Island, South Carolina, Texas, and Utah) were conducted in spring 2001; six (California, Georgia, New Hampshire, New Jersey, South Dakota, and Virginia) in fall 2001; 20 (Alabama, Arkansas, Connecticut, Delaware, Florida, Illinois, Iowa, Kentucky, Maryland, Massachusetts, Minnesota, Nebraska, New York, North Carolina, Ohio, Oklahoma, Tennessee, Vermont, West Virginia, and Wisconsin) in spring 2002; and four (Indiana, Kansas, Mississippi, and New Mexico) in fall 2002.

Of the 41 states that conducted the survey, 33 achieved an overall response rate of $\geq 60\%$ at either the middle or high

[§] Estimates are not reported if <35 cases are in the denominator.

school level or both (Table 1). These states' data were weighted, and YTS surveys produced state-representative samples of middle and high school students. Data are presented in this report only for states for which data were weighted. Data for 11 states (Arkansas, California, Indiana, Louisiana, Maine, Minnesota, Nevada, New Mexico, South Carolina, Tennessee, and Virginia) for which data were not weighted at either the middle or high school level or both are not included in this report. Data for Utah also were excluded from this report at the request of the Utah Department of Health. YTS student sample sizes varied (range: 982–38,934). School response rates varied (range: 71.4%–100.0%), as did student response rates (range: 72.0%–95.8%) and overall response rates (range: 60.2%–91.2%).

Data Collection

Survey procedures were designed to protect student privacy by ensuring that student participation was anonymous and voluntary. The survey was administered during one class period. Students completed a self-administered questionnaire in the classroom, recording their responses on an answer sheet. Questions varied by state. Both the YTS and NYTS questionnaires contained questions concerning tobacco use (bidis, cigarettes, cigars, kreteks, pipes, and smokeless tobacco), exposure to secondhand smoke, smoking cessation, school curriculum, minors' ability to purchase or obtain tobacco products, knowledge and attitudes about tobacco, and familiarity with protobacco and antitobacco media messages. Before the surveys were conducted, local parental permission procedures were followed, and state institutional review board criteria also were followed.

Results

Prevalence of Tobacco Use

Lifetime Tobacco Use

Lifetime use was defined by asking whether students have ever 1) tried cigarettes, even one or two puffs; 2) tried smoking cigars, cigarillos, or little cigars, even one or two puffs; 3) used chewing tobacco, snuff, or dip, such as Redman[®], Levi Garret[®], Beechnut[®], Skoal[®], Skoal Bandits[®], or Copenhagen[®]; 4) tried smoking bidis, even one or two puffs[‡]; 5) tried smoking kreteks, even one or two puffs[‡]; or 6) tried smoking bidis, kreteks, both, or neither.**


Middle School. Nationally, cigarettes were the most prevalent form of tobacco ever used (33.1%) among middle school

students (Figure 1), with non-Hispanic black (37.9%) and Hispanic (35.8%)^{††} students significantly more likely than white (31.3%) and Asian (24.5%) students to have ever smoked cigarettes. Cigars were the second most prevalent form of tobacco ever used (16.7%), with male students (22.0%) significantly more likely than female students (11.5%) and Hispanic students (19.1%) significantly more likely than white (16.1%)^{††} or Asian (12.7%)^{††} students to have ever smoked cigars. Smokeless tobacco was the third most prevalent form of tobacco ever used (8.2%), with male students (12.4%) significantly more likely than female students (3.9%) and white students (8.7%) significantly more likely than Hispanic (6.0%) or Asian (4.6%)^{††} students to have ever used smokeless tobacco. The percentage of students reporting they had ever smoked bidis was 4.3%, with male students (5.6%) significantly more likely than female students (3.1%) and black (5.5%) and Hispanic (6.2%) students significantly more likely than white students (3.2%) to have ever smoked bidis. The percentage of students who had ever smoked kreteks was 5.2%. No significant differences were identified among subgroups (Table 2).

Among the 31 states that asked students any or all of these questions, the percentage of students varied that had ever smoked cigarettes (range: 18.5% [New Hampshire]–50.0% [Louisiana]; median: 32.0%). Among 29 states, the percentage of students varied that had ever smoked cigars (range: 11.0% [New Hampshire]–28.5% [Louisiana]; median:

^{††} CIs overlap, but t-test is statistically significant.

FIGURE 1. Percentage of all middle school and high school students who ever* used tobacco, by type of tobacco product — National Youth Tobacco Survey, United States, 2002



* Ever use of tobacco products was determined by asking students if they had ever tried cigarettes, even one or two puffs; tried smoking cigars, cigarillos, or little cigars, even one or two puffs; used chewing tobacco, snuff, or dip, such as Redman[®], Levi Garret[®], Beechnut[®], Skoal[®], Skoal Bandits[®], or Copenhagen[®]; tried smoking bidis, even one or two puffs; or tried smoking kreteks, even one or two puffs.

[‡] Asked on NYTS only.

** Asked on state YTS only.

17.3%) (Table 3). Among 31 states, the percentage of students varied that had ever used smokeless tobacco (range: 5.5% [Maine]–22.3% [Kentucky]; median: 10.7%). Among 28 states, the percentage of students varied that had ever smoked bidis (range: 3.3% [Iowa]–10.7% [Louisiana]; median: 5.4%) or kreteks (range: 2.2% [New Hampshire]–7.7% [Texas]; median: 3.1%).

High School. Nationally, cigarettes were the most prevalent form of tobacco ever used (57.4%) among high school students (Figure 1), with male students (59.6%) significantly more likely than female students (55.3%)^{††}; white (57.2%), black (57.5%), and Hispanic (61.4%) students significantly more likely than Asian students (40.4%); and Hispanic students significantly more likely than white students^{††} to have ever smoked cigarettes. Cigars were the second most prevalent form of tobacco ever used (34.7%), with male students (44.4%) significantly more likely than female students (25.0%); white students (36.7%) significantly more likely than black (29.5%) or Asian (15.9%) students; and black and Hispanic (33.6%) students significantly more likely than Asian students to have ever smoked cigars. Smokeless tobacco was the third most prevalent form of tobacco ever used by students (16.2%), with male students (25.9%) significantly more likely than female students (6.4%); white students (19.6%) significantly more likely than black (7.0%), Hispanic (10.2%), or Asian (4.7%) students; and Hispanic students significantly more likely than black^{††} or Asian students to have ever used smokeless tobacco. The percentage of students reporting that they had ever smoked bidis was 8.5%, with male students (10.9%) significantly more likely than female students (6.0%); Hispanic (10.4%) students significantly more likely than white (7.7%)^{††} or Asian (6.3%) students to have ever smoked bidis; and black students (10.1%)^{††} significantly more likely than Asian students to have ever smoked bidis. The percentage of students reporting that they had ever smoked kreteks was 9.2%, with white (10.3%) and Hispanic (8.1%)^{††} students significantly more likely than black students (5.1%) to have ever smoked kreteks (Table 2).


Among the 25 states that asked students any or all of these questions, the percentage of students who had ever smoked cigarettes varied (range: 51.2% [Maryland]–69.7% [West Virginia]; median: 60.6%). Among 23 states, the percentage of students who had ever smoked cigars varied (range: 30.3% [Florida, Spring 2002]–48.3% [Kentucky]; median: 38.8%). The percentage of students varied that had ever used smokeless tobacco (range: 10.5% [Florida, Spring 2002]–30.6% [Kentucky]; median: 20.0%), smoked bidis (range: 6.6% [Iowa]–18.8% [Rhode Island]; median: 11.0%), and smoked kreteks (range: 4.6% [Kansas]–12.6% [Rhode Island]; median: 8.6%) (Table 3).

Current Tobacco Use

Students were asked on how many of the preceding 30 days they had used cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks[‡]. Current tobacco use was defined as use of any tobacco product on ≥ 1 of the preceding 30 days.

Middle School. Nationally, 13.3% of students were current users of any tobacco product (Figure 2), with male students (14.7%) significantly more likely than female students (11.7%)^{††} and white (13.2%),^{††} black (13.5%),^{††} and Hispanic (12.5%)^{††} students significantly more likely than Asian students (8.6%) to currently use any tobacco product. Cigarettes were the most prevalent form of tobacco currently used (9.8%), with no significant differences among subgroups. Cigars were the second most prevalent form of tobacco currently used (6.0%), with male students (7.9%) significantly more likely than female students (4.1%) to currently smoke cigars. Smokeless tobacco and pipe tobacco were the third most prevalent forms of tobacco currently used (3.5%). Male students (5.3%) were significantly more likely than female students (1.6%) and white students (3.8%) were significantly more likely than black (2.3%) students to currently use smokeless tobacco. Male students (5.1%) were significantly more likely than female students (1.9%) and Hispanic students (4.3%)^{††} were significantly more likely than white students (2.8%) to currently smoke pipe tobacco. The percentage of students reporting that they currently smoke bidis was 2.4%, with male students (3.1%) significantly more likely than female students (1.7%) to currently smoke bidis. Hispanic (2.9%)^{††} and black (3.1%)^{††} students were significantly more likely than white students (1.8%) to currently smoke bidis. The percentage of students reporting they currently smoke kreteks was 2.0%, with male students (2.7%) significantly

FIGURE 2. Percentage of all middle school and high school students who were current users of any tobacco product,* by type of tobacco product — National Youth Tobacco Survey, United States, 2002



* Use of cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥ 1 of the 30 days preceding the survey.

more likely than female students (1.1%) and Hispanic students (2.6%)^{††} significantly more likely than white students (1.5%) to currently smoke kreteks (Table 4).

Among the 31 states that asked students any or all of these questions, the percentage of students who were current users of any form of tobacco varied (range: 7.4% [New Hampshire]–26.3% [Louisiana]; median: 13.2%). The percentage of students who currently smoke cigarettes varied (range: 5.1% [New Hampshire]–17.1% [Louisiana]; median: 9.2%). Among 29 states, the percentage of students varied that currently smoke cigars (range: 1.9% [New Hampshire]–12.5% [Louisiana]; median: 5.0%) or use smokeless tobacco (range: 1.2% [Massachusetts]–10.9% [Kentucky]; median: 3.2%). Among 27 states, the percentage of students who currently smoke pipe tobacco varied (range: 1.9% [Delaware]–6.3% [Louisiana]; median: 3.1%). Among 28 states, the percentage of students who currently smoke bidis varied (range: 1.5% [Iowa]–7.1% [Louisiana]; median: 3.2%). The state YTS survey did not ask about current use of kreteks (Table 5).

High School. Nationally, 28.2% of students were current users of any tobacco product (Figure 2), with male students (32.6%) significantly more likely than female students (23.7%); white students (30.9%) significantly more likely than black (21.7%), Hispanic (24.1%), or Asian (14.6%) students; and black and Hispanic students significantly more likely than Asian students to currently use any tobacco product. Cigarettes were the most prevalent form of tobacco currently used (22.5%), with male students (23.9%)^{††} significantly more likely than female students (21.0%); white students (25.2%) significantly more likely than black (13.8%), Hispanic (19.8%), or Asian (12.2%) students; and Hispanic students significantly more likely than black or Asian students to currently smoke cigarettes. Cigars were the second most prevalent form of tobacco currently used (11.6%), with male students (16.9%) significantly more likely than female students (6.2%) and white (11.8%), black (12.0%), and Hispanic (10.8%) students significantly more likely than Asian students (5.4%) to currently smoke cigars. Smokeless tobacco was the third most prevalent form of tobacco used (5.9%), with male students (10.5%) significantly more likely than female students (1.2%); white students (7.3%) significantly more likely than black (1.8%), Hispanic (3.3%), or Asian (2.1%) students; and Hispanic students^{††} significantly more likely than black students to currently use smokeless tobacco. Pipe tobacco was the fourth most prevalent form of tobacco currently used (3.2%), with male students (5.0%) significantly more likely than female students (1.4%) and Hispanic students (4.6%) significantly more likely than white (2.8%) or Asian (2.7%)^{††} students to currently smoke pipe tobacco. The

prevalence of use of bidis (2.6%) was similar to that for kreteks (2.7%), with male students (3.7%) significantly more likely than female students (1.5%) and Hispanic students (3.5%)^{††} significantly more likely than white students (2.2%) to currently smoke bidis. Male students (3.5%) also were significantly more likely than female students (1.8%) to currently smoke kreteks (Table 4).

Among the 25 states that asked students any or all of these questions, the percentage of students that were current users of any form of tobacco varied (range: 24.3% [Florida, Spring 2002]–44.3% [Kentucky]; median: 32.0%); the percentage that currently smoke cigarettes also varied (range: 17.8% [Florida, Spring 2002]–34.2% [Kentucky]; median: 25.4%). Among 23 states, the percentage of students who currently smoke cigars varied (range: 9.1% [Connecticut]–17.1% [New Jersey]; median: 13.2%). Among 25 states, the percentage of students who currently use smokeless tobacco varied (range: 3.1% [Connecticut]–13.5% [Kentucky]; median: 6.9%). Among 21 states, the percentage of students who currently smoke pipe tobacco varied (range: 2.7% [Delaware and Wisconsin]–6.9% [Maryland]; median: 4.0%). Among 23 states, the percentage of students who currently smoke bidis varied (range: 2.7% [Iowa]–9.5% [Rhode Island]; median: 5.5%); YTS did not ask about current use of kreteks (Table 5).

Ever Smoked Cigarettes Daily

Students were asked if they had ever smoked cigarettes daily. Daily use was defined as ever having smoked at least one cigarette every day for 30 days.


Middle School. Nationally, 5.8% of students had ever smoked cigarettes daily (Figure 3), with white students (6.2%) significantly more likely than black (4.4%)^{††} or Hispanic students (4.2%)^{††} to have ever smoked cigarettes daily (Table 6). Among the 30 states that asked this question, the percentage of students who had ever smoked cigarettes daily varied (range: 2.4% [Kansas]–10.8% [Louisiana]; median: 4.9%) (Table 7).

High School. Nationally, 17.0% of students had ever smoked cigarettes daily (Figure 3), with white students (19.7%) significantly more likely than black (9.8%), Hispanic (12.1%), or Asian (10.1%) students to have ever smoked cigarettes daily (Table 6). Among the 24 states that asked this question, the percentage of students who had ever smoked cigarettes daily varied (range: 13.3% [Florida, Spring 2002]–29.1% [West Virginia]; median: 19.1%) (Table 7).

Number of Cigarettes Smoked Per Day Among Current Cigarette Smokers

Current cigarette smokers were asked how many cigarettes they smoked per day on the days they smoked during the preceding 30 days.

FIGURE 3. Percentage of all middle school and high school student who ever smoked cigarettes daily and current cigarette smokers who smoked six or more cigarettes per day on the days they smoked — National Youth Tobacco Survey, United States, 2002



* Students were asked, "Have you ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days?"

† Students were asked, "During the past 30 days, on the days you smoked, how many cigarettes did you smoke per day?"

Middle School. Nationally, 16.6% of current cigarette smokers had smoked six or more cigarettes per day on the days they smoked (Figure 3), with male students (19.8%)^{††} significantly more likely than female students (12.7%) to have smoked six or more cigarettes per day on the days they smoked (Table 6). Among the 31 states that asked this question, the percentage of current cigarette smokers that smoked six or more cigarettes per day on the days they smoked varied (range: 8.9% [Minnesota]–25.9% [Vermont]; median: 15.7%) (Table 7).


High School. Nationally, 31.3% of current cigarette smokers had smoked six or more cigarettes per day on the days they smoked (Figure 3), with male students (34.4%)^{††} significantly more likely than female students (27.9%) and white students (34.5%) significantly more likely than black (20.4%) or Hispanic (19.6%) students to have smoked six or more cigarettes per day on the days they smoked (Table 6). Among the 25 states that asked this question, the percentage of current cigarette smokers that smoked six or more cigarettes per day on the days they smoked varied (range: 21.7% [Kansas]–44.0% [Kentucky]; median: 30.7%) (Table 7).

Age of Initiation of Tobacco Use

Cigarettes

Middle School. Nationally, 8.1% of students first smoked a whole cigarette before age 11 years (Figure 4), with male students (9.8%) significantly more likely than female students (6.5%) to have done so (Table 8). Among the 30 states that asked this question, the percentage of students who first smoked

FIGURE 4. Percentage of all middle school and high school students who first smoked a cigarette before age 11* years — National Youth Tobacco Survey, United States, 2002



* Age of initiation was determined by asking, "How old were you when you smoked a whole cigarette for the first time?"

a whole cigarette before age 11 years varied (range: 4.9% [Maryland]–14.3% [Louisiana]; median: 8.9%) (Table 9).

High School. Nationally, 6.7% of students first smoked a whole cigarette before age 11 years (Figure 4), with male students (8.4%) significantly more likely than female students (5.0%) to have done so (Table 8). Among the 24 states that asked this question, the percentage of students who first smoked a whole cigarette before age 11 years varied (range: 5.7% [Connecticut]–11.9% [Florida]; median: 9.0%) (Table 9).

Cigars**

Middle Schools. Among the 27 states that asked a question concerning cigar use, the percentage of students who first smoked a cigar before age 11 years varied (range: 2.9% [Connecticut]–8.9% [Louisiana]; median: 5.1%) (Table 9).

High Schools. Among the 20 states that asked a question concerning cigar use, the percentage of students who first smoked a cigar before age 11 years varied (range: 2.6% [Delaware]–6.4% [West Virginia]; median: 4.3%) (Table 9).

Smokeless Tobacco**

Middle Schools. Among the 27 states that asked a question concerning smokeless tobacco use, the percentage of students who initiated use of smokeless tobacco before age 11 years varied (range: 1.2% [Massachusetts]–10.2% [Kentucky]; median: 3.5%) (Table 9).

High Schools. Among the 21 states that asked a question concerning smokeless tobacco use, the percentage of students who initiated use of smokeless tobacco before age 11 years varied (range: 1.3% [Massachusetts]–8.7% [Kentucky]; median: 3.2%) (Table 9).

Established Use of Tobacco Products

Smoked ≥ 100 Cigarettes in Lifetime


Middle School. Nationally, 10.6% of students who had ever smoked cigarettes had smoked ≥ 100 cigarettes during their lifetime (Figure 5), with male students (12.2%) significantly more likely than female students (8.6%)^{††} and white students (12.9%) significantly more likely than black (5.7%) or Hispanic (5.6%) students to have done so (Table 10). Among the 30 states that asked this question, the percentage of middle school students that had ever smoked cigarettes reported smoking ≥ 100 cigarettes varied (range: 5.9% [New Jersey]–17.5% [Kentucky]; median: 10.1%) (Table 11).

High School. Nationally, 28.0% of students who had ever smoked cigarettes had smoked ≥ 100 cigarettes in their lifetime (Figure 5), with male students (29.8%)^{††} significantly more likely than female students (26.0%); white students (33.5%) significantly more likely than black (12.1%), Hispanic (18.4%), or Asian (23.2%) students; and Hispanic and Asian students significantly more likely than black students to have done so (Table 10). Among the 24 states that asked this question, the percentage of high school students that had ever smoked cigarettes reported smoking ≥ 100 cigarettes varied (range: 20.3% [Texas]–39.2% [Kentucky]; median: 29.0%) (Table 11).

Frequent Use of Tobacco Products

Students were asked on how many of the preceding 30 days they had used a tobacco product. Frequent cigarette smoking was defined as having smoked on ≥ 20 of the preceding 30

FIGURE 5. Percentage of middle school and high school students who ever smoked cigarettes who smoked ≥ 100 cigarettes in their lifetime and percentage of all students who frequently smoked cigarettes — National Youth Tobacco Survey, United States, 2002



* Used cigarettes on ≥ 20 of the 30 days preceding the survey.

days. Questions also were asked regarding frequent use of cigars, smokeless tobacco, pipes, bidi,^{**} and kreteks.^{**}

Middle School. Nationally, 2.5% of students were frequent cigarette smokers (Figure 5), with white students (2.7%)^{††} significantly more likely than Hispanic students (1.5%) to smoke cigarettes frequently. Overall, 1.5% of students were frequent users of cigars, smokeless tobacco, pipes, bidis, or kreteks (Table 10). Among the 31 states that asked students any or all of these questions, the percentage of students who were frequent cigarette smokers varied (range: 1.1% [New Jersey]–5.4% [West Virginia]; median: 2.4%) (Table 11). Among 29 states, the percentage of students who were frequent cigar smokers varied (range: 0.1% [New Hampshire]–2.0% [Alabama]; median: 0.7%). Among the 31 states, the percentage of students who were frequent smokeless tobacco users varied (range: 0.2% [New Hampshire and Wisconsin]–3.1% [Kentucky]; median: 0.8%). Among 27 states, the percentage of students who were frequent smokers of pipe tobacco varied (range: 0.2% [Kansas]–1.1% [Kentucky]; median: 0.6%). Among 28 states, the percentage of students who were frequent bidi smokers varied (range: 0.1% [New Hampshire]–1.0% [Alabama]; median: 0.4%).

High School. Nationally, 11.1% of students were frequent cigarette smokers (Figure 5), with white students (13.4%) significantly more likely than black (5.4%), Hispanic (6.2%), or Asian (7.0%) students to frequently smoke cigarettes. Overall, students who were frequent users of cigars, smokeless tobacco, pipes, bidis, and kreteks, accounted for $<2.5\%$ of students (Table 10). Among the 25 states that asked any or all of these questions, the percentage of students who were frequent cigarette smokers varied (range: 7.4% [Florida, Spring 2002]–20.2% [Kentucky]; median: 12.0%). Among 23 states, the percentage of students who were frequent cigar smokers varied (range: 0.4% [Wisconsin]–2.2% [North Carolina]; median: 1.3%). Among 25 states, the percentage of students who were frequent smokeless tobacco users varied (range: 0.3% [Connecticut]–4.9% [Kentucky]; median: 1.9%). Among 21 states, the percentage of students who were frequent smokers of pipe tobacco varied (range: none [Wisconsin]–1.7% [New York]; median: 0.7%). Among 23 states, the percentage of students who were frequent bidi smokers varied (range: 0.2% [Ohio]–1.8% [New York]; median: 0.8%) (Table 11).

Brand of Cigarettes Usually Smoked


Current cigarette smokers were asked what brand of cigarettes they had usually smoked during the preceding 30 days. Options included American Spirit[®],[‡] Basic[®], Camel[®], Doral[®], GPC[®], Kool[®],[‡] Lucky Strike[®],[‡] Marlboro[®], Newport[®], Parliament[®],[‡] Virginia Slims[®], “some other brand,” and “no usual brand.” Because of low use of all brands

other than Camel, Marlboro, and Newport, all other specific brands were subsumed in “some other brand.”

Middle School. Nationally, 41.8% of current cigarette smokers identified Marlboro as the brand they usually smoked, followed by Newport (26.3%), other brands (17.4%), no usual brand (8.8%), and Camel (5.8%) (Figure 6). Among current smokers, white (47.6%), Hispanic (46.9%), and Asian (30.9%)^{††} students were significantly more likely than black students (9.0%) to smoke Marlboro; and black students (58.3%) were significantly more likely than white (22.3%), Hispanic (20.4%), or Asian (25.0%) students to smoke Newport (Table 12). Among the 29 states that asked this question, the percentage of current cigarette smokers who usually smoked one of these three specific brands varied: Marlboro (range: 24.4% [Delaware]–63.5% [New Hampshire]; median: 43.9%), Newport (range: 2.1% [Idaho]–52.7% [Delaware]; median: 16.7%), and Camel (range: 2.4% [Delaware]–26.0% [Idaho]; median: 5.5%) (Table 13).

High School. Nationally, 52.0% of current cigarette smokers identified Marlboro as the brand they usually smoked, followed by Newport (21.6%), other brands (11.7%), Camel (8.7%), and no usual brand (6.0%) (Figure 6). Among current smokers, white (56.6%), Hispanic (53.0%), and Asian (55.1%) students were significantly more likely than black students (12.8%) to smoke Marlboro; black students (66.8%) were significantly more likely than white (16.6%), Hispanic (21.5%), and Asian (16.4%) students to smoke Newport; and white students (9.5%) were significantly more likely than black (4.5%)^{††} or Hispanic (5.0%)^{††} students to smoke Camel

FIGURE 6. Brand of cigarettes smoked in the 30 days preceding the survey by current cigarette smokers* in middle school and high school — National Youth Tobacco Survey, United States, 2002



* Current cigarette smoking is defined as having smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

[†] Includes Kool[®], Lucky Strike[®], Virginia Slims[®], GPC[®], Basic[®], American Spirit[®], Parliament[®], and Doral[®].

(Table 12). Among the 23 states that asked this question, the percentage of current cigarette smokers varied who usually smoked one of these three brands varied: Marlboro (range: 31.9% [Delaware]–64.4% [Texas]; median: 52.4%), Newport (range: 6.9% [Oklahoma]–55.0% [Delaware]; median: 19.5%), and Camel (range: 1.6% [Delaware]–21.8% [Wisconsin]; median: 9.5%) (Table 13).

Susceptibility Among Students Who Have Never Smoked Cigarettes

Students who have never smoked cigarettes were defined as not susceptible to initiating cigarette smoking during the next year if they responded that 1) they would not smoke a cigarette soon, 2) they definitely would not smoke a cigarette during the next year, and 3) they definitely would not smoke a cigarette offered to them by one of their best friends. This definition has been published previously (13). All other students were classified as susceptible to initiating cigarette smoking during the next year.

Middle School. Nationally, 21.3% of students who have never smoked cigarettes were susceptible to initiating cigarette smoking during the next year. No significant differences were identified among subgroups (Table 14). Among the 29 states that asked this series of questions, the percentage of students who have never smoked cigarettes who were susceptible to initiating cigarette smoking during the next year varied (range: 17.4% [Idaho]–33.3% [New Jersey]; median: 23.9%) (Table 15).

High School. Nationally, 22.9% of students who have never smoked cigarettes were susceptible to initiating cigarette smoking in the next year. Among students who have never smoked cigarettes, Hispanic students (31.7%) were significantly more likely than white (21.6%) or black students (18.4%) and Asian students (30.4%) were significantly more likely than black^{††} students to be susceptible to initiating cigarette smoking in the next year (Table 14). Among the 23 states that asked this series of questions, the percentage of students who have never smoked cigarettes who were susceptible to initiating cigarette smoking in the next year varied (range: 18.4% [Florida, Spring 2000]–35.4% [New Jersey]; median: 23.8%) (Table 15).

Knowledge and Attitudes Regarding Tobacco Use

Social Influence

One or More Closest Friends Smoke Cigarettes

Students were asked how many of their four closest friends smoke cigarettes. Students were considered to have never smoked cigarettes if they answered “no” to whether they have

tried or experimented with cigarette smoking, even one or two puffs.

Middle School. Nationally, current cigarette smokers (89.7%) were significantly more likely than students who have never smoked cigarettes (15.4%) to report that one or more of their closest friends smoked cigarettes (Figure 7). No significant differences were observed among subgroups of students who have never smoked cigarettes. Among current cigarette smokers, female students (92.9%)^{††} were significantly more likely than male students (86.5%) to report that one or more of their best friends smoked cigarettes (Table 16). Among the 31 states that asked this question, the percentage of students who reported that one or more of their closest friends smoked cigarettes varied among students who have never smoked cigarettes (range: 13.1% [Massachusetts]–27.9% [West Virginia]; median: 18.6%) and among current cigarette smokers (range: 80.2% [Maryland and Wisconsin]–92.1% [Delaware]; median: 86.6%) (Table 17).

High School. Nationally, current cigarette smokers (91.2%) were significantly more likely than students who have never smoked cigarettes (27.8%) to report that one or more of their closest friends smoke cigarettes (Figure 7). Among students who have never smoked cigarettes, Hispanic students (31.5%) were significantly more likely than black (25.4%)^{††} or Asian (24.1%)^{††} students to report that one or more of their closest


friends smoke cigarettes. Among current cigarette smokers, white (92.6%) and Hispanic (90.2%)^{††} students were significantly more likely than black students (82.2%) to report that one or more of their closest friends smoke cigarettes (Table 16). Among the 25 states that asked this question, the percentage of students who reported that one or more of their closest friends smoked cigarettes varied among students who have never smoked cigarettes (range: 25.3% [Massachusetts]–47.5% [West Virginia]; median: 35.9%) and among current cigarette smokers (range: 81.2% [Connecticut]–95.1% [Iowa]; median: 90.6%) (Table 17).

One or More Closest Friends Use Smokeless Tobacco

Middle School. Nationally, current smokeless tobacco users (74.2%) were significantly more likely than students who have never used this product (6.4%) to report that one or more of their closest friends used smokeless tobacco (Figure 7). Among students who have never used smokeless tobacco, white (6.6%)^{††} and Hispanic (6.7%) students were significantly more likely than black students (4.2%) to report that one or more of their best friends use smokeless tobacco; no significant differences were identified among current smokeless tobacco users (Table 16). Among the 29 states that asked this question, the percentage of students who reported that one or more of their closest friends used smokeless tobacco varied among students who have never used smokeless tobacco (range: 6.2% [New York]–22.8% [Kentucky]; median: 12.8%) and among current smokeless tobacco users (range: 59.7% [Illinois]–90.1% [South Dakota]; median: 73.7%) (Table 17).

High School. Nationally, current smokeless tobacco users (83.0%) were significantly more likely than students who have never used this product (11.8%) to report that one or more of their closest friends use smokeless tobacco (Figure 7). Among students who have never used smokeless tobacco, male students (13.6%)^{††} were significantly more likely than female students (10.4%); white students (13.9%) significantly more likely than black (6.0%), Hispanic (9.8%), or Asian (7.0%) students; and Hispanic students significantly more likely than black students to report that one or more of their closest friends use smokeless tobacco. Among current smokeless tobacco users, male students (84.9%) were significantly more likely than female students (66.4%) to report that one or more of their closest friends used smokeless tobacco (Table 16). Among the 23 states that asked this question, the percentage of students who reported that one or more of their closest friends used smokeless tobacco varied among students who have never used smokeless tobacco (range: 9.1% [Massachusetts]–29.6% [West Virginia]; median: 19.6%) and among current users of smokeless tobacco (range: 67.5% [Connecticut]–90.8% [Ohio]; median: 82.6%) (Table 17).

FIGURE 7. Percentage of middle school and high school students with peers who use tobacco, by tobacco use status and type of tobacco — National Youth Tobacco Survey, United States, 2002



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

† Use of smokeless tobacco on ≥ 1 of the 30 days preceding the survey.


Social Perception Regarding Cigarette Use

Think that Cigarette Smokers Have More Friends

Middle School. Nationally, current cigarette smokers (42.9%) were significantly more likely than students who have never smoked cigarettes (14.2%) to think that cigarette smokers have more friends (Figure 8). Among students who have never smoked cigarettes, black (21.1%) and Hispanic (16.4%)^{††} students were significantly more likely than white students (12.3%), and black students were significantly more likely than Hispanic^{††} and Asian (13.6%) students to think that cigarette smokers have more friends. Among current cigarette smokers, black (51.7%) and Hispanic (51.3%) students were significantly more likely than white students (37.8%) to think that cigarette smokers have more friends (Table 18). Among the 31 states that asked this question, the percentage of students who think that cigarette smokers have more friends varied among students who have never smoked cigarettes (range: 5.8% [Idaho and Iowa]–22.4% [Mississippi]; median: 12.0%) and among current cigarette smokers (range: 31.0% [Iowa]–56.7% [Maryland]; median: 43.6) (Table 19).

High School. Nationally, current cigarette smokers (23.4%) were significantly more likely than students who have never smoked cigarettes (10.9%) to think that cigarette smokers have more friends (Figure 8). Among students who have never smoked cigarettes, male students (12.4%) were significantly more likely than female students (9.5%),^{††} and black (17.5%), Hispanic (15.7%), and Asian (14.7%)^{††} students were significantly more likely than white students (8.2%) to think

FIGURE 8. Percentage of middle school and high school students who think cigarette smokers have more friends, by cigarette smoking status — National Youth Tobacco Survey, United States, 2002



* Students were considered as never having smoked if they answered “no” to whether they have tried or experimented with cigarette smoking, even one or two puffs.

† Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

that cigarette smokers have more friends. Among current cigarette smokers, male students (28.8%) were significantly more likely than female students (17.3%); black (42.0%), Hispanic (29.5%), and Asian (41.7%) students significantly more likely than white students (19.4%); and black students were significantly more likely than Hispanic students to think that cigarette smokers have more friends (Table 18). Among the 25 states that asked this question, the percentage of students who think that cigarette smokers have more friends varied among students who have never smoked cigarettes (range: 8.1% [Iowa]–22.4% [Mississippi]; median: 13.6%) and among current cigarette smokers (range: 23.1% [Ohio]–42.1% [Mississippi]; median: 28.0%) (Table 19).

Access

How Current Cigarette Smokers Aged <18 Years Usually Acquired Cigarettes


Current cigarette smokers aged <18 years were asked how they usually had acquired cigarettes during the preceding 30 days. Options included 1) in a store, 2) from a vending machine, 3) someone else bought them, 4) borrowed them from someone else, 5) stole them (asked on NYTS), 6) took them from a store or family member (asked on YTS), 7) person aged ≥ 18 years gave them, and 8) obtained them some other way.

Middle School. Nationally, the most common means by which current cigarette smokers usually acquired their cigarettes were 1) obtained some other way (25.0%), 2) borrowed them from someone (24.9%), 3) had someone else buy them (21.7%), or 4) stole them (11.5%) (Figure 9). Among current cigarette smokers, white students (23.9%) were significantly more likely usually to have someone else buy their cigarettes for them than Hispanic students (14.7%) (Table 20). Hispanic (8.5%)^{††} students were significantly more likely than white (4.5%) students usually to buy their cigarettes in a store.

Among the 31 states that asked this question, for the majority of states, the most common ways by which middle school students aged <18 years who currently smoke usually acquired their cigarettes were similar nationwide: borrowed cigarettes (range: 9.3% [Maine]–36.5% [Nebraska]; median: 26.6%), had someone else buy cigarettes for them (range: 13.1% [New Jersey]–32.2% [South Dakota]; median: 23.6%), and acquired their cigarettes some other way (range: 9.3% [New Hampshire]–30.8% [Massachusetts]; median: 18.1%) (Table 21).

High School. Nationally, the most common means by which current cigarette smokers aged <18 years usually acquired their cigarettes was by having someone else buy them (28.7%), buying them in a store (24.7%), or borrowing them

FIGURE 9. Percentage of current cigarette smokers* aged <18 years in middle school and high school who obtained cigarettes, by usual methods of obtaining — National Youth Tobacco Survey, United States, 2002



* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

from someone (20.6%) (Figure 9). A total of 11.1% of current cigarette smokers reported that they were given cigarettes by a person aged ≥18 years. Among current cigarette smokers aged <18 years, female students (32.8%) were significantly more likely than male students (24.6%)^{††} and white students (30.4%) were significantly more likely than Hispanic (22.1%)^{††} or black (20.4%)^{††} students usually to have someone else buy their cigarettes for them. Male students (31.2%) were significantly more likely than female students (18.1%) to purchase their cigarettes in a store. Female students (14.1%) were more likely than male students (8.2%) to be given cigarettes by a person aged ≥18 years. Hispanic students (14.6%)^{††} were more likely than white students (8.5%) to obtain their cigarettes some other way. Male students (4.3%)^{††} were more likely than female students (2.2%) and black students (6.6%)^{††} were significantly more likely than white students (2.5%) usually to steal their cigarettes (Table 20). Among the 25 states that asked this question, ways by which current smokers aged <18 years usually obtained cigarettes varied: had someone else buy them (range: 19.8% [Connecticut]–43.2% [Iowa]; median: 31.9%), purchased them in a store (range: 6.9% [Nebraska]–33.7% [New Jersey]; median: 20.9%), and borrowed them (range: 18.3% [Delaware]–30.5% [Massachusetts]; median: 25.1%) (Table 21).

Where Current Cigarette Smokers Aged <18 Years Bought Last Pack of Cigarettes

Current cigarette smokers aged <18 years were asked where they bought their last pack of cigarettes during the preceding 30 days. Response categories differed between the national


and state surveys. NYTS response options were gas station, convenience store, discount store, grocery store, drug store, vending machine, and restaurant. YTS response options were gas station, convenience store, grocery store, drug store, vending machine, Internet, and other. The “other” category varied among the states.

Middle School. Nationally, the most common locations where current cigarette smokers bought their last pack of cigarettes were a gas station (44.7%), a convenience store (23.0%), and a vending machine (11.6%) (Figure 10). Among current cigarette smokers, white students (48.1%)^{††} were significantly more likely than black students (33.3%) to have bought their last pack of cigarettes at a gas station. White students (6.3%) also were significantly more likely than black (1.6%)^{††} or Hispanic students (2.0%)^{††} to have bought their last pack of cigarettes at a discount store. Hispanic (11.1%)^{††} and black students (16.2%)^{††} were significantly more likely than white students (4.0%) to have bought their last pack of cigarettes at a grocery store (Table 22). Among the 29 states that asked this question, percentages varied of current cigarette smokers who purchased their last pack of cigarettes at a gas station (range: 9.7% [Massachusetts]–38.6% [New York]; median: 22.2%), at a convenience store (range: 3.2% [Iowa]–31.4% [New York]; median: 13.0%), from a vending machine (range: 0.9% [Iowa]–15.6% [Maine]; median: 3.9%), and in another location (range: 2.9% [Texas^{§§}]–68.2% [Iowa]; median: 50.7%) (Table 23).

High School. Nationally, the most common locations where high school students aged <18 years who are current cigarette

§§ Only restaurants were included in this category.

FIGURE 10. Location in which current cigarette smokers* aged <18 years in middle school and high school bought their last pack of cigarettes — National Youth Tobacco Survey, United States, 2002



* Smoked cigarettes on ≥1 of the 30 days preceding the survey.

smokers bought their last pack of cigarettes were a gas station (57.0%) and a convenience store (27.8%) (Figure 10). Among current cigarette smokers aged <18 years, white students (61.1%) were significantly more likely than black (37.3%) or Hispanic (46.6%)^{††} students to have bought their last pack of cigarettes at a gas station, and Hispanic students (9.8%)^{††} were significantly more likely than white students (4.5%) to have bought their last pack of cigarettes at a grocery store (Table 22). Among the 23 states that asked this question, percentages varied of current smokers aged <18 years who purchased their last pack of cigarettes at a gas station (range: 20.2% [New Jersey]–63.2% [Illinois]; median: 44.4%), at a convenience store (range: 13.5% [Wisconsin]–52.7% [New Jersey]; median: 22.5%), and in another location (range: 2.1% [Texas^{§§}]–30.8% [Alabama]; median: 21.6%) (Table 23).

Enforcement

Current Cigarette Smokers Aged <18 Years Not Asked for Proof of Age


Middle School. Nationally, 75.9% of current cigarette smokers aged <18 years were not asked to show proof of age when purchasing or attempting to purchase cigarettes from a store (Figure 11). No significant differences were identified among current cigarette smoker subgroups (Table 24). Among the 26 states that asked this question, the percentage of current cigarette smokers who were not asked to show proof of age when purchasing or attempting to purchase cigarettes at a store varied (range: 57.8% [Delaware]–92.6% [Georgia]; median: 74.2%) (Table 25).

High School. Nationally, 58.5% of current cigarette smokers aged <18 years were not asked to show proof of age when purchasing or attempting to purchase cigarettes from a store (Figure 11). Among current smokers aged <18 years, black students (69.3%)^{††} were significantly more likely than white students (57.0%) to report that they were not asked to show proof of age (Table 24). Among the 24 states that asked this question, the percentage of current smokers aged <18 years who were not asked to show proof of age when purchasing or attempting to purchase cigarettes from a store varied (range: 54.3% [New York]–72.1% [Iowa]; median: 62.3%) (Table 25).

Current Cigarette Smokers Aged <18 Years Not Refused Purchase

Middle School. Nationally, 63.4% of current cigarette smokers were not refused purchase of cigarettes because of their age (Figure 11). No significant differences were identified among subgroups of current smokers (Table 24). Among the 26 states that asked this question, the percentage of cur-

FIGURE 11. Percentage of current cigarette smokers* aged <18 years in middle school and high school who purchased cigarettes in a store and were not asked to show proof of age or who were not refused purchase because of their age — National Youth Tobacco Survey, United States, 2002



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

rent smokers who were not refused purchase of cigarettes because of their age varied (range: 54.3% [Florida, Spring 2002]–78.2% [Minnesota]; median: 69.1%) (Table 25).

High School. Nationally, 60.6% of current cigarette smokers aged <18 years were not refused purchase of cigarettes because of their age (Figure 11). No significant differences were identified among subgroups of current smokers aged <18 years (Table 24). Among the 24 states that asked this question, the percentage of current smokers aged <18 years who were not refused purchase of cigarettes because of their age varied (range: 49.7% [Illinois]–68.0% [Wisconsin]; median: 60.9%) (Table 25).


Exposure to Media and Advertising Regarding Tobacco

Saw or Heard Antismoking Commercials

The following questions were used to ascertain students' exposure to anti-tobacco media messages during the preceding 30 days: 1) during the past 30 days, how often did you see anti-smoking commercials on TV[¶] and 2) during the past 30 days, how often did you hear anti-smoking commercials on the radio.[¶] Students were defined as exposed to these messages if they saw or heard such a message at least once.

Middle School. Nationally, 84.6% of middle school students saw or heard antismoking commercials (Figure 12). Female students (87.4%) were significantly more likely than male students (81.9%) and white (85.4%)^{††} and Hispanic (85.3%)^{††} students were significantly more likely than black students (81.8%) to have seen or heard such commercials (Table 26). Among the 29 states that asked this question, the

FIGURE 12. Percentage of all middle school and high school students who were exposed to tobacco-related media and advertising — National Youth Tobacco Survey, United States, 2002



percentage of students who saw or heard antismoking commercials varied (range: 69.1% [Maryland public schools]–91.3% [Iowa]; median: 80.3%) (Table 27).

High School. Nationally, 91.2% of high school students saw or heard antismoking commercials (Figure 12). Female students (92.3%)^{††} were significantly more likely than male students (90.1%) and white students (92.2%) significantly more likely than Hispanic (90.1%),^{††} black (88.7%), or Asian (88.3%)^{††} students to have seen or heard such commercials (Table 26). Among the 23 states that asked this question, the percentage of students who saw antismoking commercials on television or heard antismoking commercials on the radio varied (range: 78.1% [Maryland]–93.1% [Iowa]; median: 84.9%) (Table 27).

Saw Tobacco Use on Television or in Movies

Students were asked how often they see actors smoking[†] or using tobacco^{**} when they watch TV or go to the movies. For YTS, students who were exposed to these images are defined as those who responded to this question with “most of the time” or “some of the time.”

Middle School. Nationally, 89.9% of students had seen actors smoking on television or in the movies (Figure 12), with white (90.3%) and black (90.3%)^{††} students significantly more likely than Asian students (84.0%) and Hispanic (89.6%)^{††} students significantly more likely than Asian students to have to have done so (Table 26). Among the 28 states that asked this question, the percentage of students who had seen actors using tobacco on television or in the movies varied (range: 78.1% [New Hampshire]–90.2% [New Jersey]; median: 83.6%) (Table 27).

High School. Nationally, 91.3% of students had seen actors smoking on television or in the movies (Figure 12), with female students (92.4%) significantly more likely than male students (90.2%) to have done so (Table 26). Among the 22 states that asked this question, the percentage of students who had seen actors using tobacco on television or in the movies varied (range: 85.0% [Maryland public schools]–91.3% [New York]; median: 89.7%) (Table 27).

Saw Advertisements for Tobacco Products on Internet

Students were questioned regarding their exposure to advertisements for tobacco products on the Internet. For NYTS, students were asked: “When you are searching the Internet on a computer, how often do you see ads for cigarettes and other tobacco products?” and for YTS, students were asked: “When you are using the Internet, how often do you see ads for tobacco products?”


Middle School. Nationally, 42.7% of students had seen advertisements for tobacco products on the Internet (Figure 12), with Hispanic students (44.9%)^{††} significantly more likely than black students (41.3%) to have seen such advertisements (Table 26). Among the 27 states that asked this question, the percentage of students who had seen such advertisements varied (range: 25.3% [Texas]–43.9% [New York]; median: 34.1%) (Table 27).

High School. Nationally, 33.5% of students had seen advertisements for tobacco products on the Internet (Figure 12), with Hispanic (38.4%) and black (36.9%)^{††} students significantly more likely than white students (31.7%) to have seen such advertisements (Table 26). Among the 21 states that asked this question, the percentage of students who had seen such advertisements varied (range: 19.6% [Massachusetts]–37.9% [Maryland]; median: 28.4%) (Table 27).

Participated in Antitobacco Community Events

Middle School. Nationally, current tobacco users (17.9%) were significantly more likely than students who have never used tobacco (11.7%) to have participated in a community event to discourage persons from using tobacco products (Figure 13). Among students who have never used tobacco, female students (13.0%)^{††} were significantly more likely than male students (10.2%) and black students (15.6%) were significantly more likely than white (11.0%), Hispanic (10.1%) or Asian (9.0%)^{††} students to have participated in such events. Among current tobacco users, black students (23.9%)^{††} were significantly more likely than white students (16.4%) to have participated in such events (Table 28). Among the 29 states

FIGURE 13. Percentage of middle school and high school students who participated in antitobacco community events, by tobacco use status — National Youth Tobacco Survey, United States, 2002



* Students were considered as never having used tobacco if they did not use cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥1 of the 30 days preceding the survey.

† Used cigarettes, cigars, smokeless tobacco, tobacco in a pipe, bidis, or kreteks on ≥1 of the 30 days preceding the survey.

that asked this question, the percentage of students who had participated in an antitobacco community event varied among students who have never used tobacco (range: 11.6% [New York]–33.8% [Idaho]; median: 23.7%) and among current tobacco users (range: 14.6% [Florida, spring 2002]–27.8% [Maryland]; median: 21.2%) (Table 29).

High School. Nationally, 11.5% of students who have never used tobacco and 9.5% of current tobacco users participated in a community event to discourage persons from using tobacco products in the past year (Figure 13). Among students who have never used tobacco, female students (13.8%) were significantly more likely than male students (8.6%) and black students (14.4%)^{††} were significantly more likely than Hispanic students (9.7%) to have participated in such an event. Among current tobacco users, black (15.9%) and Hispanic (12.8%)^{††} students were significantly more likely than white students (7.8%) to have participated in such an event (Table 28). Among the 24 states that asked this question, the percentage of students who had participated in an antitobacco community event varied among students who have never used tobacco (range: 8.7% [Florida, spring 2002]–25.9% [Mississippi]; median: 15.4%) and among current tobacco users (range: 6.5% [Kentucky]–17.6% [Maryland public schools]; median: 11.4%) (Table 29).


Receptivity to Tobacco Company Merchandise

Bought or Received Item with Tobacco Company Name or Picture on It

Middle School. Nationally, current tobacco users (41.8%) were significantly more likely than students who have never used tobacco (8.7%) to have bought or received anything with a tobacco company name or picture on it (e.g., sports gear, T-shirt, cigarette lighter, hat, jacket, or sunglasses that they purchased or received for free) (Figure 14). Among students who have never used tobacco, Hispanic students (9.2%)^{††} were significantly more likely than Asian students (5.7%) to have bought or received any such items; no significant differences were identified among current tobacco user subgroups (Table 30). Among the 30 states that asked this question, the percentage of students who had bought or received anything with a tobacco company name or picture on it varied among students who have never used tobacco (range: 6.4% [New York]–16.1% [Iowa]; median: 11.2%) and among current tobacco users (range: 31.9% [Rhode Island]–52.7% [Maine]; median: 43.4%) (Table 31).

High School. Nationally, current tobacco users (31.3%) were significantly more likely than students who have never

FIGURE 14. Percentage of middle school and high school students receptive to tobacco company merchandise,* by tobacco status — National Youth Tobacco Survey, United States, 2002



* For example, a cigarette lighter or T-shirt.

† Students were considered as never having used tobacco if they did not use cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥1 of the 30 days preceding the survey.

§ Used cigarettes, cigars, smokeless tobacco, tobacco in a pipe, or bidis on ≥1 of the 30 days preceding the survey.

used tobacco (7.7%) to have bought or received anything with a tobacco company name or picture on it (Figure 14), with male students (34.6%) significantly more likely than female students (26.7%) to have bought or received such items (Table 30). Among the 25 states that asked this question, the percentage of students who had bought or received anything with a tobacco company name or picture on it varied among students who have never used tobacco (range: 5.7% [Kansas]–24.4% [Mississippi]; median: 11.9%) and among current tobacco users (range: 29.8% [Florida, spring 2001, and Rhode Island]–63.3% [Mississippi]; median: 39.4%) (Table 31).

Would Wear or Use an Item with Tobacco Company Name or Picture on It

Middle School. Nationally, current tobacco users (58.1%) were significantly more likely than students who have never used tobacco (11.0%) to report that they would wear or use items (e.g., sports gear, T-shirts, cigarette lighters, hats, jackets, or sunglasses) that had a tobacco company name or picture on it (Figure 14). Among students who have never used tobacco, male students (15.7%) were significantly more likely than female students (6.9%) and Hispanic students (12.7%)^{††} significantly more likely than black students (9.2%) to report that they would wear or use such items. Among current tobacco users, Hispanic students (60.9%)^{††} were significantly more likely than black students (49.9%) to report that they would wear or use such items (Table 30). Among the 31 states that asked this question, the percentage of students who would wear or use anything with a tobacco company name or picture on it varied among students who have never used tobacco (range: 7.9% [Rhode Island]–20.9% [Mississippi]; median: 11.6%) and among current tobacco users (range: 44.2% [Connecticut]–67.9% [Mississippi]; median: 54.7%) (Table 31).

High School. Nationally, current tobacco users (54.9%) were significantly more likely than students who have never used tobacco (13.7%) to report that they would wear or use an item with a tobacco company name or picture on it (Figure 14). Among students who have never used tobacco, male students (19.0%) were significantly more likely than female students (9.3%) and Hispanic students (17.4%) were significantly more likely than white (13.1%), black (12.9%)^{††} or Asian (12.0%)^{††} students to report that they would wear or use such items. Among current tobacco users, male students (57.8%) were significantly more likely than female students (50.9%);^{††} white students (57.9%) were significantly more likely than black (37.1%) or Hispanic (51.7%)^{††} students; and Hispanic students were significantly more likely than black students to report that they would wear or use

such items (Table 30). Among the 25 states that asked this question, the percentage of students who would wear or use anything with a tobacco company name or picture on it varied among students who have never used tobacco (range: 10.9% [Delaware]–24.4% [Mississippi]; median: 16.9%) and among current tobacco users (range: 48.2% [Delaware]–67.9% [Iowa]; median: 58.0%) (Table 31).

Cessation Attempts and Desire to Stop

Tried to Quit Smoking Cigarettes During Preceding 12 Months


Middle School. Nationally, 55.4% of current cigarette smokers had tried to quit smoking during the preceding 12 months (Figure 15). No significant differences were identified among subgroups of current cigarette smokers (Table 32). Among the 30 states that asked this question, the percentage of current cigarette smokers who had tried to quit smoking in the 12 months preceding the survey varied (range: 47.5% [Illinois]–69.7% [Iowa]; median: 56.3%) (Table 33).

High School. Nationally, 53.1% of current cigarette smokers had tried to quit smoking during the preceding 12 months (Figure 15). No significant differences were identified among subgroups of current cigarette smokers (Table 32). Among the 24 states that asked this question, the percentage of current cigarette smokers who had tried to quit smoking during the preceding 12 months varied (range: 41.8% [Texas]–64.4% [Wisconsin]; median: 57.5%) (Table 33).

Desire to Stop Smoking Cigarettes

Middle School. Nationally, 49.6% of current cigarette smokers reported that they wanted to stop smoking

FIGURE 15. Percentage of current cigarette smokers* in middle school and high school who tried to quit and who want to stop smoking — National Youth Tobacco Survey, United States, 2002



* Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

(Figure 15). No significant differences were identified among current cigarette smoker subgroups (Table 32). Among the 30 states that asked this question, the percentage of current smokers who wanted to stop smoking varied (range: 40.9% [Oklahoma]–65.9% [Iowa]; median: 52.2%) (Table 33).

High School. Nationally, 62.1% of current cigarette smokers reported that they wanted to stop smoking (Figure 15), with white (63.2%)^{††} and black (68.2%)^{††} students significantly more likely than Hispanic students (56.6%) to want to stop smoking cigarettes (Table 32). Among the 24 states that asked this question, the percentage of current cigarette smokers who wanted to stop smoking varied (range: 47.7% [Georgia]–67.8% [Massachusetts]; median: 58.6%) (Table 33).


Secondhand Cigarette Smoke

Was in Same Room with Someone Who Was Smoking Cigarettes on ≥ 1 of Preceding 7 Days

Middle School. Nationally, current cigarette smokers (88.3%) were significantly more likely than students who have never smoked cigarettes (47.1%) to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among students who have never smoked cigarettes, female students (49.4%) were significantly more likely than male students (44.8%) students; white students (51.6%) were significantly more likely than black (43.4%), Hispanic (35.1%), or Asian (27.6%) students; black students were significantly more likely than Hispanic or Asian students; and Hispanic students^{††} were significantly more likely than Asian students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Among current cigarette smokers, female students (91.5%) were significantly more likely than male students (84.8%)^{††} and white students (91.0%) were significantly more likely than black (77.7%) or Hispanic (84.1%)^{††} students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). Among the 30 states that asked this question, the percentage of students who had been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days varied among students who have never smoked cigarettes (range: 31.1% [Texas]–63.3% [Kentucky]; median: 45.3%) and among current smokers (range: 73.5% [Texas]–90.8% [Pennsylvania]; median: 86.1%) (Table 35).

High School. Nationally, current cigarette smokers (91.4%) were significantly more likely than students who have never smoked cigarettes (53.3%) to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the pre-

FIGURE 16. Percentage of middle school and high school students who were in a room or who rode in a car with someone who was smoking cigarettes on ≥ 1 of preceding 7 days or who think smoke from other persons' cigarettes is harmful, by smoking status — National Youth Tobacco Survey, United States, 2002



* Students were considered as never having smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

† Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

ceding 7 days (Figure 16). Among students who have never smoked cigarettes, white students (56.4%) were significantly more likely than black (50.7%),^{††} Hispanic (44.1%), or Asian (46.6%) students and black students^{††} were significantly more likely than Hispanic students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Among current cigarette smokers, female students (93.4%) were significantly more likely than male students (89.7%) and white students (93.4%) were significantly more likely than black (84.2%) or Hispanic (86.7%) students to have been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). Among the 24 states that asked this question, the percentage of students who had been in the same room with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days varied among students who have never smoked cigarettes (range: 40.4% [Texas]–73.6% [Kentucky]; median: 53.0%) and among current smokers (range: 81.6% [Maryland]–95.9% [Kentucky]; median: 90.5%) (Table 35).

Rode in Car with Someone Who Was Smoking Cigarettes on ≥ 1 of Preceding 7 Days

Middle School. Nationally, current cigarette smokers (81.7%) were significantly more likely than students who have never smoked cigarettes (31.5%) to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among students who have never smoked cigarettes, white (33.4%) and black (33.3%) students were significantly more likely than Hispanic (22.5%) or Asian (22.1%) students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Among current cigarette smokers, female students (85.8%) were significantly more likely than male students (77.3%) and white students (84.3%) were significantly more likely than black (71.3%) or Hispanic (76.2%)^{††} students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). Among the 30 states that asked this question, the percentage of students who had ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days varied among students who have never smoked cigarettes (range: 21.3% [Idaho]–49.8% [Kentucky]; median: 32.6%) and among current smokers (range: 70.0% [Massachusetts]–91.6% [Iowa]; median: 79.9%) (Table 35).

High School. Nationally, current smokers (83.7%) were significantly more likely than students who have never smoked cigarettes (29.1%) to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Figure 16). Among students who have never smoked cigarettes, female students (31.0%) were significantly more likely than male students (26.8%)^{††}; white (29.6%) students were significantly more likely than Hispanic (25.0%)^{††} or Asian (19.0%) students; black (34.2%) students were significantly more likely than Hispanic and Asian students; and Hispanic students were significantly more likely than Asian^{††} students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days. Among current cigarette smokers, female students (86.1%)^{††} were significantly more likely than male students (81.7%) and white (86.0%) and Asian (91.6%) students were significantly more likely than black (72.6%) or Hispanic (78.2%) students to have ridden in a car with someone who was smoking cigarettes on ≥ 1 of the preceding 7 days (Table 34). Among the 24 states that asked this question, the percentage of students who had ridden in a car with someone who was smoking cigarettes on ≥ 1 of the past 7 days varied among students who have never smoked cigarettes (range: 23.7% [Texas]–44.9% [Kentucky]; median: 30.7%) and among current cigarette smokers (range: 72.6% [Texas]–89.9% [Delaware]; median: 83.6%) (Table 35).

Think That Secondhand Cigarette Smoke is Harmful to Them

Middle School. Nationally, students who have never smoked cigarettes (91.1%) were significantly more likely than current cigarette smokers (78.6%) to think that secondhand cigarette smoke is harmful to them (Figure 16), with female students (93.4%) significantly more likely than male students (88.8%) and white (93.6%) and Asian (94.1%) students significantly more likely than black (84.6%) or Hispanic (87.1%) students to think that secondhand cigarette smoke is harmful to them. Among current smokers, female students (82.9%) were significantly more likely than male students (74.0%)^{††} and Asian students (91.0%) were significantly more likely than white (80.8%),^{††} black (72.2%), or Hispanic (75.6%) students to think that secondhand cigarette smoke is harmful to them (Table 34). Among the 31 states that asked this question, the percentage of students who think that secondhand cigarette smoke is harmful to them varied among students who have never smoked cigarettes (range: 87.0% [Louisiana]–96.0% [Maine]; median: 93.0%) and among current cigarette smokers (range: 70.6% [Texas]–89.7% [Nebraska]; median: 82.3%) (Table 35).

High School. Nationally, students who have never smoked cigarettes (95.1%) were significantly more likely than current cigarette smokers (90.2%) to think that secondhand cigarette smoke is harmful to them (Figure 16), with female students (96.7%) significantly more likely than male students (93.2%) and white (96.7%) and Asian (96.9%) students were significantly more likely than black (90.2%) or Hispanic (91.2%) students to think that secondhand cigarette smoke is harmful to them. Among current cigarette smokers, female students (93.3%) were significantly more likely than male students (87.5%) and white students (92.6%) were significantly more likely than black (81.2%) or Hispanic (84.5%) students to think that secondhand cigarette smoke is harmful to them (Table 34). Among the 25 states that asked this question, the percentage of students who think that secondhand cigarette smoke is harmful to them varied among students who have never smoked cigarettes (range: 89.1% [Florida]–97.4% [Nebraska]; median: 94.4%) and among current cigarette smokers (range: 80.9% [Maryland]–92.7% [Pennsylvania]; median: 88.8%) (Table 35).

Tobacco Use by Others in the Home Someone Else in Home Smokes Cigarettes

Middle School. Nationally, current cigarette smokers (71.5%) were significantly more likely than students who have never smoked cigarettes (33.3%) to live in a home in which

someone else smokes cigarettes (Figure 17). Among students who have never smoked cigarettes, black students (35.2%) were significantly more likely than Hispanic (29.7%)^{††} or Asian (26.6%)^{††} students to live in such a home. Among current cigarette smokers, white students (73.6%)^{††} were significantly more likely than Hispanic students (64.2%) students to live in such a home (Table 36). Among the 30 states that asked this question, the percentage of students who live in a home in which someone else smokes cigarettes varied among students who have never smoked cigarettes (range: 24.2% [Idaho]–45.0% [Kentucky]; median: 34.9%) and among current cigarette smokers (range: 58.0% [Illinois]–81.0% [Wisconsin]; median: 70.1%) (Table 37).

High School. Nationally, current cigarette smokers (57.5%) were significantly more likely than students who have never smoked cigarettes (29.9%) to live in a home in which someone else smokes cigarettes (Figure 17). Among students who have never smoked cigarettes, black students (35.3%) were significantly more likely than white (29.7%),^{††} Hispanic (27.8%), or Asian (24.2%) students to live in such a home. Among current cigarette smokers, female students (62.2%) were significantly more likely than male students (53.3%) and Asian students (70.9%)^{††} significantly more likely than Hispanic (52.8%) students to live in such a home (Table 36). Among the 24 states that asked this question, the percentage of students who live in a home in which someone else smokes cigarettes varied among students who have never smoked cigarettes (range: 24.7% [Illinois]–37.5% [Kentucky]; median: 29.6%) and among

current cigarette smokers (range: 48.6% [Connecticut]–66.3% [West Virginia]; median: 56.9%) (Table 37).

Someone Else in Home Uses Smokeless Tobacco

Students who have never used smokeless tobacco and current smokeless tobacco users were asked if anyone who lives with them now uses chewing tobacco, snuff, or dip.^{**}

Middle School. Among the 28 states that asked this question, the percentage of students who reported living in a home in which someone else uses smokeless tobacco varied among students who have never used smokeless tobacco (range: 3.2% [Massachusetts]–23.6% [West Virginia]; median: 10.3%) and among current users of smokeless tobacco (range: 24.3% [Delaware]–59.8% [Alabama]; median: 46.7%) (Table 37).

High School. Among the 22 states that asked this question, the percentage of students who reported living in a home in which someone else uses smokeless tobacco varied among students who have never used smokeless tobacco (range: 1.9% [Massachusetts]–20.7% [West Virginia]; median: 8.8%) and among current smokeless tobacco users (range: 22.7% [Ohio]–48.1% [Georgia]; median: 37.1%) (Table 37).


Schools

Practiced Tobacco Refusal Skills as Part of School Curriculum During School Year

Middle School. Nationally, 38.8% of students practiced (e.g., through role playing) ways to say “no” to tobacco as part of the school curriculum in any of their classes during the school year (Figure 18), with female students (41.9%)^{††} significantly more likely than male students (35.8%) and black (45.3%) students significantly more likely than white students (36.8%) to have done so (Table 38). Among the 29 states that asked this question, the percentage of students who practiced ways to say “no” to tobacco as part of the school curriculum varied (range: 23.5% [New Hampshire]–64.7% [Maryland]; median: 45.2%) (Table 39).

High School. Nationally, 16.4% of students practiced ways to say “no” to tobacco as part of the school curriculum (Figure 18), with female (17.8%)^{††} students significantly more likely than male (14.9%) students and black (22.5%), Hispanic (20.5%), and Asian (21.5%)^{††} students significantly more likely than white students (14.0%) to have done so (Table 38). Among the 23 states that asked this question, the percentage of students who practiced ways to say “no” to tobacco as part of the school curriculum varied (range: 9.4% [New Hampshire]–32.1% [Maryland]; median: 17.5%) (Table 39).


FIGURE 17. Percentage of middle school and high school students who were exposed to tobacco use at home, by tobacco status — National Youth Tobacco Survey, United States, 2002



* Students were considered as never having smoked if they answered “no” to whether they have tried or experimented with cigarette smoking, even one or two puffs.

† Smoked cigarettes on ≥ 1 of the 30 days preceding the survey.

FIGURE 18. Percentage of all middle school and high school students who practiced ways to say "no" to tobacco as part of school curriculum or who smoked cigarettes or used smokeless tobacco on school property during the 30 days preceding the survey — National Youth Tobacco Survey, United States, 2002



Tobacco Use on School Property

Smoked Cigarettes on School Property during Preceding 30 Days

Middle School. Nationally, 2.7% of students had smoked cigarettes on school property during the 30 days preceding the survey (Figure 18). No significant differences were identified among subgroups. Nationally, 26.7% of current smokers had smoked cigarettes on school property during the 30 days preceding the survey, with black (32.8%),^{††} Hispanic (32.1%),^{††} and Asian (48.2%)^{††} students significantly more likely than white (21.8%) students to have done so (Table 38). Among the 31 states that asked this question, the percentage of students who had smoked cigarettes on school property during the preceding 30 days varied among all students (range: 1.6% [Connecticut]–5.7% [Louisiana]; median: 3.1%) and among current cigarette smokers (range: 15.8% [Georgia]–41.3% [New Hampshire]; median: 27.6%) (Table 39).

High School. Nationally, 8.1% of students had smoked cigarettes on school property during the 30 days preceding the survey (Figure 18), with male students (9.4%)^{††} significantly more likely than female students (6.7%); white students (8.8%) significantly more likely than black (4.8%) or Asian (5.3%)^{††} students; and Hispanic students (7.9%) significantly more likely than black students to have done so. Nationally, 35.8% of current smokers had smoked cigarettes on school property during the 30 days preceding the survey, with male (39.4%) students significantly more likely than female (31.7%)^{††} students to have done so (Table 38). Among the 25 states that asked this question, the percentage of students who had smoked cigarettes on school property during

the preceding 30 days varied among all students (range: 4.9% [Florida, Spring 2002]–16.4% [Kentucky]; median: 10.1%) and among current cigarette smokers (range: 26.7% [Kansas]–57.0% [Rhode Island]; median: 36.0%) (Table 39).

Used Smokeless Tobacco on School Property during Preceding 30 Days

Middle School. Nationally, 1.9% of students had used smokeless tobacco on school property in the 30 days preceding the survey (Figure 18), with male students (2.7%) significantly more likely than female students (1.0%) to have done so. Nationally, 52.0% of users of smokeless tobacco reported using smokeless tobacco on school property in the 30 days preceding the survey, with black (65.4%)^{††} and Hispanic (64.8%)^{††} students significantly more likely than white (45.3%) students to have done so (Table 38). Among the 27 states that asked this question, the percentage of students who had used smokeless tobacco on school property during the preceding 30 days varied among all students (range: 0.8% [Iowa]–7.0% [Louisiana]; median: 2.0%) and among current users of smokeless tobacco (range: 20.9% [Kansas]–71.8% [Michigan]; median: 49.5%) (Table 39).

High School. Nationally, 3.5% of students had used smokeless tobacco on school property during the 30 days preceding the survey (Figure 18), with male students (6.5%) significantly more likely than female students (0.4%); white students (4.2%) significantly more likely than black (1.1%), Hispanic (2.5%),^{††} or Asian (1.0%) students; and Hispanic students significantly more likely than black^{††} or Asian^{††} students to have done so (Table 38). Nationally, 59.1% of smokeless tobacco users reported using smokeless tobacco on school property in the 30 days preceding the survey, with male students (62.0%) significantly more likely than female (34.4%) students and Hispanic students (74.5%)^{††} significantly more likely than white students (57.3%) to have done so. Among the 22 states that asked this question, the percentage of students who had used smokeless tobacco on school property during the preceding 30 days varied among all students (range: 2.8% [Connecticut]–8.1% [West Virginia]; median: 4.5%) and among current smokeless tobacco users (range: 40.7% [Ohio]–83.0% [Rhode Island]; median: 57.8%) (Table 39).

Discussion

CDC recommends that states establish comprehensive tobacco-control programs to reduce disease, disability, and death related to tobacco use by

- preventing young persons from ever starting to smoke,
- promoting quitting among young persons and adults,

- protecting persons from exposure to secondhand smoke, and
- identifying and eliminating disparities related to tobacco use and its effects on different population groups (8).

These goals can be addressed through comprehensive tobacco use prevention and control programs that include best practices components (e.g., statewide, school, counter marketing, community, cessation, and enforcement programs) (8). The Task Force on Community Preventive Services recently evaluated the evidence of effectiveness of 14 selected interventions across these program and goal areas. Recommended interventions (i.e., those that showed either sufficient or strong evidence of effectiveness) included 1) increasing the unit price for tobacco products, 2) mass media campaigns implemented with additional interventions (e.g., tobacco product excise tax increases, school-based education, or other community programs), 3) community mobilization combined with additional interventions around minor's access, 4) smoking bans and restrictions, 5) health-care systems-level interventions (e.g., provider reminders alone or in combination with provider education), 6) reducing patient costs for cessation treatments, and 7) multicomponent patient telephone support for cessation (15,16).

NYTS and YTS data can be used to monitor comprehensive tobacco-control programs and evaluate exposure to select intervention components and inform program planning, improvement, and effectiveness. Evaluation should include periodic and ongoing monitoring of tobacco related attitudes, behaviors, health outcomes, the prevalence of protobacco influences (e.g., advertising, promotions, and events that glamorize tobacco use), and measures of anti-tobacco influences. Ultimately, these data will help determine the effect of program elements on tobacco-related attitudes, behaviors, and policies. The following discussion provides examples in how youth surveillance and evaluation data are used for this purpose.

Prevalence and Initiation of Tobacco Use

Young persons are at the greatest risk for initiating tobacco use, becoming addicted to tobacco, and establishing use patterns that can become lifelong. However, if young persons remain tobacco-free, the majority will never start using tobacco (6). Major risk factors for initiation include perceptions that tobacco use is common and normative and the availability and accessibility of tobacco products (6). NYTS and YTS data provide measures that can be used to plan, monitor, and evaluate efforts to prevent initiation and reduce smoking prevalence among young persons.

Use of YTS as a Primary Source of Data for Evaluating Prevention Initiatives

In 1999, the Minnesota state legislature approved the Tobacco Prevention Endowment. The Minnesota Department of Health's Tobacco Prevention and Control Section (MN-TPCS) was charged with administering the Endowment, and in 2000 the Minnesota Youth Tobacco Prevention Initiative, Minnesota's only statewide coordinated youth tobacco prevention program, was established. In 1999, the state legislature set a goal of a 30% reduction in tobacco use among youth by 2005, and MN-TPCS was required to identify and establish measurable indicators and outcomes to assess progress toward this goal.

MN-TPCS uses the Minnesota Youth Tobacco Survey (MN YTS) as the primary data source for evaluation of the Minnesota Youth Tobacco Prevention Initiative. By monitoring trends from MN YTS, MN-TPCS was able to demonstrate movement toward the program's goal in its 2002 annual report to the legislature (17). MN YTS data indicated that, during 2000–2002, the percentage of middle school students who currently use some form of tobacco decreased 11%, from 12.6% to 11.2%, and the percentage of middle school students who currently smoke cigarettes decreased 21%, from 9.1% to 7.2%. The evaluation results from the MN YTS indicate that the Minnesota Youth Tobacco Prevention Initiative was on track to reach the 2005 goal. MN-TPCS estimates that achieving the goal would ultimately prevent approximately 1,700 premature deaths and save \$480 million in health-care costs every year in Minnesota. During the 2003 legislative session, the Minnesota state legislature approved an 80% reduction in MN-TPCS funding. MN YTS data will now be used to provide MN-TPCS with information on how this shift in programming affects these trends.

Quitting Tobacco Use among Young Persons

Young smokers become addicted to cigarettes and experience withdrawal symptoms similar to those reported by adults (6). Although evidence is limited regarding effective cessation programs among young persons, promising interventions and methods have emerged (18,19). To further this progress, state and national program providers and policy makers should continue to conduct scientifically sound youth cessation studies to produce quality data and address research gaps; develop and implement carefully considered youth cessation strategies; rigorously evaluate interventions using standardized methods; and uniformly report all results in the published literature (19,20). NYTS and YTS data provide measures for planning,

monitoring, and evaluating efforts to promote quitting tobacco use among young persons.

Use of YTS for Programmatic Decisions and Recommendations

New Jersey uses YTS data to guide programmatic decisions and recommendations. Since its inception in January 2000, the Comprehensive Tobacco Control Program in the New Jersey Department of Health and Senior Services (NJ-CTCP) has carried out a full range of statewide and local initiatives to reduce tobacco use. One of five NJ-CTCP program goals is increasing the number of youth tobacco users who initiate cessation treatment (21). Program progress is evaluated by monitoring indicators related to short and long term outcomes of the program, especially those related to tobacco use knowledge, attitudes, and behaviors. NJ-CTCP uses the New Jersey Youth Tobacco Survey (NJ YTS) as the primary evaluation system for youth tobacco use in the state to measure these and other types of youth indicators (22).

NJ-CTCP used NJ YTS data to demonstrate that youth cigarette smokers are interested in quitting. Nicotine dependence was demonstrated among current smokers, and a strong inverse association was identified between current smokers' level of confidence that they could successfully quit and how many cigarettes they smoked per day; these factors present challenges to cessation efforts. When 1999 and 2001 NJ YTS data were compared, no significant differences in interest in quitting, indicators of nicotine dependence, or quit attempts were identified. However, several important changes were noted with respect to physician tobacco counseling, a critical precursor to cessation treatment. During 1999–2001, the percentage of high school smokers who reported that they were asked about their smoking status while visiting their doctor increased 30%. NJ-CTCP continues to focus on developing options for improving overall youth smoking cessation. The program focus on making age-appropriate tobacco dependence treatment programs readily available and accessible to increase youth cessation in the state. To address this need, NJ-CTCP has implemented a youth cessation curriculum in several New Jersey high schools and established a new initiative to train pediatricians to identify and treat tobacco dependence.

Exposure to Secondhand Smoke

Exposure to secondhand smoke has direct health consequences for youth such as causing lower respiratory illness and exacerbating asthma (6). In addition, exposure to smoking by others contributes to perceptions that tobacco use is normative (6). Interventions to reduce secondhand smoke

exposure provide immediate protection for young persons in schools, public places, homes, and cars. NYTS and YTS data provide measures important to planning, monitoring, and evaluating efforts to reduce exposure to secondhand smoke among young persons.

Use of YTS as an Evaluation System

In New Hampshire, YTS is used as an evaluation system to inform program planning, assessment, and effectiveness. The New Hampshire Department of Health and Human Service's Tobacco Prevention and Control Program (NH-TPCP) has adopted all four of the National Tobacco Control Program's (NTCP) goal areas and is based on CDC/OSH's Best Practices for Comprehensive Tobacco Control Programs (8).

NH-TPCP's comprehensive program relies on the ongoing collection and detailed analysis of surveillance and evaluation data to determine program efficacy. NH-TPCP uses YTS data to measure overall program progress toward decreasing exposure to secondhand smoke among young persons aged 11–17 years and to guide program efforts in this area. NH-TPCP logic models and action plans connects the activity of providing community education on topics of secondhand smoke and promoting smoke-free homes and vehicles to outputs such as the recipients of the secondhand smoke education being exposed to pro-health/anti-tobacco messages. Outputs are further linked to Specific-Measurable-Achievable-Relevant and Time bound (SMART) outcome objectives such as decreasing the percentage of youth who report exposure to secondhand smoke and increasing the percentage of youth who report that they think that secondhand smoke is harmful. NH-TPCP uses NH YTS as the primary data source for measuring these objectives, with specific YTS questions directly providing the needed indicators.

Disparities in Tobacco Use

Tobacco-related measures often vary across different population groups (23). Analysis and review of data by different demographic characteristics (e.g., sex, race/ethnicity, or education level) can assist in targeting culturally appropriate programs that address varying levels of tobacco-related problems. YTS and NYTS data provide measures important to planning, monitoring, and evaluating efforts to reduce disparities in tobacco use among young persons.

Use of YTS to Identify Disparities in Tobacco Use

In Kansas, YTS is used as an integral component to identifying and eliminating disparities in tobacco use. The Kansas Department of Health and Environment's Tobacco Use Pre-

vention Program (KS-TUPP) has recognized the need for local tobacco control activities, programs, and monitoring to achieve lasting progress toward Kansas' and the Nation's goals and objectives. KS-TUPP's primary role is to provide resources and technical assistance to community coalitions for the development, enhancement, and evaluation of state and local tobacco-control initiatives. From this local-level focus, they have built a broad network of local, regional, and state coalitions. In keeping with their community focus, and goal of eliminating tobacco-related disparities among specific populations, KS-TUPP has issued a request for proposals for addressing tobacco use among minority youth for the last 3 years.

Using 2000 YTS data, KS-TUPP identified sub-state areas in which Hispanic youth showed substantially higher rates of current tobacco use and current cigarette smoking than Hispanic youth at the state level. In 2002, three Hispanic organizations were funded to address this disparity by implementing policy, media, and community mobilization prevention strategies to generate awareness and change in tobacco use knowledge, attitudes, and behavior among Hispanic youth aged 12–17 years. One of the funded organizations, Our Gang Inc., has created a Youth Advisory Board that makes tobacco control advocacy presentations and engages peers in youth empowerment. Board members reinforce their counter marketing messages by distributing pro-health/anti-tobacco cards and wearing pro-health/anti-tobacco T-shirts on assigned days. Over time, the YTS will provide measures to monitor the impact of these activities and identify other priority groups for KS-TUPP targeted strategies.

Limitations

The findings in this report are subject to at least four limitations. First, these data apply only to youth who attend middle school or high school and are not representative of all persons in this age group. In 2000, the most recent year for which data are available, only 4% of persons aged 16 years and 8% of those aged 17 years who had not completed high school were not enrolled in a high school program (24). Second, although NYTS and the majority of state surveys were conducted during the spring semester, a limited number of states conducted surveys during the fall semester. Conducting the survey at different times throughout the school year introduces variation which might influence survey estimates. These differences include cohort aging (i.e., the school population is older in the spring) and other potential contextual differences (e.g., level of tobacco-control program activity, tobacco-related policy, or price changes). Future analysis of data from fall and spring surveys will assess these effects. Third, all data are self reported. Because tobacco use by youth is considered

a socially undesirable behavior, young persons might not give an honest report of their behaviors and attitudes (6). This might possibly lead to under- or overreporting by respondents, though the direction and extent of this cannot be determined. However, those YTS questions that have been analyzed have demonstrated reliability (25), and youth self-reports have been determined to be generally accurate (26). Finally, in certain instances, question or response categories slightly differ either between different state YTS surveys or between NYTS and the core state YTS survey. For example, the definition of a current user of any tobacco product (Table 5) in certain states does not include use of cigars, pipes, or bidis because these core questions were removed from the states' survey. Also, the measure of how current smokers usually acquired cigarettes (Tables 20 and 21) differs for certain response categories, with wording varying slightly between NYTS and YTS. These differences limit comparability across the surveys on measures derived from those questions.

Conclusion

States can use NYTS and YTS data to plan, monitor, and evaluate aspects of their comprehensive tobacco-control programs that focus on youth. States can use these data as tobacco program indicators to 1) monitor trends, a step that could lead to better program planning and program improvement; and 2) compare their state data with those of other states and the nation to evaluate change.

The following conclusions can be drawn from NYTS and YTS data:

- Tobacco products are available and accessible to young persons.
- Tobacco refusal skills are not commonly practiced as part of middle and high school curricula.
- Current cigarette smokers have tried to or want to quit cigarette smoking.
- Young persons have high levels of exposure to second-hand smoke and pro-tobacco media.
- Tobacco-related disparities exist across racial/ethnic subgroups, sex, and middle/high school.

Three waves of NYTS have been completed: 1999, 2000, and 2002. Between the time of its inception in 1998 and completion of the 2002 surveys, 42 states and the District of Columbia have implemented YTS and obtained weighted data at least once. Nearly half of these states have weighted YTS data from ≥ 2 years. Mississippi and Texas now have 4 years of data, and Florida has 5 years of data. Alabama, Arkansas, Delaware, Iowa, Kansas, Kentucky, Maine, Maryland, Minnesota, New Jersey, New York, North Carolina, Ohio, Oklahoma,

Tennessee, West Virginia, and Wisconsin all have 2 years of data. Continued support for the NYTS and YTS will help inform tobacco-control program planning, improvement, and effectiveness.

References

- McGinnis JM, Foege WH. Actual causes of death in the United States. *JAMA* 1993;270:2207–12.
- CDC. Annual smoking-attributable mortality, years of potential life lost, and productivity losses—United States, 1997–2001. *MMWR* 2002;54:625–8.
- CDC. Tobacco use among high school students—United States, 1997. *MMWR* 1998;47:229–33.
- Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE. Monitoring the future: national survey results on drug use, 1975–2004. Volume 1: secondary school students (NIH publication no. 05-5727). Bethesda, MD: National Institute on Drug Abuse; 2004. Available at http://www.monitoringthefuture.org/pubs/monographs/vol1_2004.pdf.
- CDC. Trends in cigarette smoking among high schools students—United States, 1991–2001. *MMWR* 2002;51:409–12.
- US Department of Health and Human Services. Preventing tobacco use among young people: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1994.
- CDC. Sustaining state programs for tobacco control—data highlights 2004. Atlanta, GA: US Department of Health and Human Services, CDC; 2004. Available at <http://www.cdc.gov/tobacco/datahighlights/index.htm>.
- CDC. Best Practices for comprehensive tobacco control programs, August 1999. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1999.
- MacDonald G, Starr G, Schooley M, Yee SL, Klimowski K, Turner K. Introduction to program evaluation for comprehensive tobacco control programs. Atlanta, GA: US Department of Health and Human Services, CDC; 2001.
- CDC. Tobacco use among middle and high school students—United States, 1999. *MMWR* 2000;49:49–53.
- Farrelly M, Davis K, Yarsevich J, et al. Legacy first look report 9: getting to the truth: assessing youths' reactions to the truthsm and “think, don't smoke” tobacco countermarketing campaigns. Washington, DC: American Legacy Foundation; 2002.
- Shah BV, Barnwell BG, Bieler GS. SUDAAN user's manual, release 8.0. Research Triangle Park, NC: Research Triangle Institute; 2001.
- SAS Institute, Inc. SAS/STAT[®] user's guide, version 8.02. Cary, NC: SAS Institute; 2001.
- Pierce JP, Choi WS, Gilpin EA, Farkas AJ, Berry CC. Tobacco industry promotion of cigarettes and adolescent smoking. *JAMA* 1998;279:511–5.
- Hopkins DP, Fielding JE, Task Force on Community Preventive Services, eds. The Guide to Community Preventive Services: tobacco use prevention and control reviews, recommendations and expert commentary. *Am J Prev Med* 2001;20(Suppl 2):1–88.
- Zaza S, Briss PA, Harris KW, eds. The Guide to Community Preventive Services: what works to promote health? New York, NY: Oxford University Press; 2005.
- Minnesota Department of Health. Tobacco prevention and local public health endowment: annual report to the legislature, 2002 activities. St. Paul, MN: Minnesota Department of Health; 2003. Available at <http://www.health.state.mn.us/divs/opa/tobaclegprpt02.pdf>.
- Milton MH, Maule CO, Backinger C, Gregory DM. Recommendations and guidance for practice in youth tobacco cessation. *Am J Health Behav* 2003;27(Suppl 2):S159–69.
- McDonald P, Colwell B, Backinger CL, Huston C, Maule CO. Better practices for youth tobacco cessation: evidence of review panel. *Am J Health Behav* 2003;27(Suppl 2):S144–58.
- Backinger CL, McDonald P, Ossip-Klein DJ, et al. Improving the future of youth cessation. *Am J Health Behav* 2003;27(Suppl 2):170–84.
- Delnevo CD, Hrywna M, Lewis MJ, et al. Independent evaluation of the New Jersey Comprehensive Tobacco Control Program: annual update for the New Jersey Department of Health and Senior Services. New Brunswick, NJ: University of Medicine and Dentistry of New Jersey School of Public Health; 2003.
- Delnevo CD, Muthurajah S, Brown M, Hrywna M, Malka ES. The 2001 New Jersey Youth Tobacco Survey: a statewide report for the New Jersey Department of Health and Senior Services. New Brunswick, NJ: University of Medicine and Dentistry of New Jersey School of Public Health; 2002.
- US Department of Health and Human Services. Tobacco use among U.S. racial/ethnic minority groups: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Public Health Service, CDC; 1998.
- Kaufman P, Alt MN, Chapman CD. Dropout rates in the United States, 2000. Washington, DC: US Department of Education, National Center for Education Statistics, Office of Educational Research and Improvement; 2002; (publication no. NCES 02-114).
- Brener ND, Kann L, McManus T, Kinchen S, et al. Reliability of the 1999 Youth Risk Behavior Survey questionnaire. *J Adolesc Health* 31:336–42.
- US Department of Health and Human Services. Women and smoking: a report of the Surgeon General. Rockville, MD: US Department of Health and Human Services, Public Health Service, Office of the Surgeon General; 2001.

TABLE 1. Sample sizes and response rates for middle schools and high schools, by state — state youth tobacco surveys, United States, 2001–2002

State	Student completed sample size	Response rate (%)		
		School	Student	Overall
Middle school				
Alabama – Spring 02	1,594	81.6	89.2	72.8
Connecticut – Spring 02	2,677	79.7	85.7	68.3
Delaware – Spring 02	2,779	98.2	92.1	90.4
Florida – Spring 01	4,366	93.6	78.0	73.0
Florida – Spring 02	15,251	93.9	82.2	77.2
Georgia – Fall 01	2,848	100.0	91.2	91.2
Idaho – Spring 01*	1,878	86.8	87.8	76.3
Illinois – Spring 02	1,667	92.3	93.2	86.0
Iowa – Spring 02	1,633	83.3	88.4	73.6
Kansas – Fall 02	1,489	74.0	87.0	64.4
Kentucky – Spring 02	1,535	88.0	90.1	80.1
Louisiana – Spring 01	2,543	82.5	72.9	60.2
Maine – Spring 01	12,570	88.9	76.1	67.6
Maryland – Spring 02	27,338	100.0	89.9	89.9
Massachusetts – Spring 02	1,963	88.0	91.4	80.5
Michigan – Spring 01	1,820	80.0	90.6	72.5
Minnesota – Spring 02	4,751	85.0	86.2	73.3
Mississippi – Fall 02	3,287	86.5	78.2	67.7
Nebraska – Spring 02	2,812	75.4	95.5	72.0
New Hampshire – Fall 01	1,538	80.0	91.3	73.0
New Jersey – Fall 01*	5,413	88.2	83.6	73.3
New York – Spring 02†	8,124	85.7	89.6	76.8
North Carolina – Spring 02	5,747	86.7	84.5	73.3
Ohio – Spring 02	1,590	85.7	86.3	74.0
Oklahoma – Spring 02	2,043	86.0	84.4	72.6
Pennsylvania – Spring 01	8,419	72.3	87.9	63.6
Rhode Island – Spring 01	1,646	92.0	85.3	78.5
South Dakota – Fall 01	2,467	92.5	87.1	80.6
Texas – Spring 01	8,687	87.0	82.0	71.3
Utah – Spring 01*	1,303	100.0	83.3	83.3
Vermont – Spring 02	4,711	84.0	87.7	73.7
West Virginia – Spring 02	10,052	95.1	84.3	80.1
Wisconsin – Spring 02	1,298	75.0	83.3	62.5
High school				
Alabama – Spring 02	1,351	75.5	85.6	64.6
Connecticut – Spring 02	3,198	85.0	85.0	72.2
Delaware – Spring 02	2,517	100.0	82.9	82.9
Florida – Spring 01	4,327	93.6	78.6	73.6
Florida – Spring 02	8,895	93.3	78.1	72.9
Georgia – Fall 01	2,975	94.0	88.9	83.6
Illinois – Spring 02	1,730	88.5	90.9	80.4
Iowa – Spring 02	1,697	90.0	87.6	78.9
Kansas – Fall 02	1,517	83.7	83.8	70.1
Kentucky – Spring 02	1,530	86.0	84.1	72.3
Maryland – Spring 02	38,934	100.0	84.2	84.2
Massachusetts – Spring 02	1,675	88.0	83.5	73.5
Michigan – Spring 01	3,352	81.6	83.1	67.9
Mississippi – Fall 02	2,707	86.7	72.0	62.4
Nebraska – Spring 02	2,677	71.4	94.0	67.1
New Hampshire – Fall 01	1,446	74.0	84.9	62.8
New Jersey – Fall 01*	4,176	79.7	77.1	61.4
New York – Spring 02†	8,124	85.7	89.6	76.8
North Carolina – Spring 02	5,203	83.5	83.9	70.1
Ohio – Spring 02	1,271	86.0	87.4	75.2
Oklahoma – Spring 02	1,908	94.0	84.8	79.8
Pennsylvania – Spring 01	8,415	76.5	84.0	64.3
Rhode Island – Spring 01	1,574	92.0	74.3	68.4
Texas – Spring 01	8,696	73.4	82.2	60.3
Utah – Spring 01	982	97.1	77.0	74.8
West Virginia – Spring 02	1,640	83.7	79.7	66.7
Wisconsin – Spring 02	1,362	77.1	83.9	64.7

* New Jersey data combined public and private schools; middle school sample included only 7th and 8th graders.

† Because New York participated in the National Youth Tobacco Survey, which samples students in grades 6–12, New York response rates are for middle school and high school combined.

TABLE 2. Percentage of middle school and high school students who ever used* cigarettes, cigars, smokeless tobacco, bidis, or kreteks, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Cigarettes		Cigars		Smokeless tobacco		Bidis		Kreteks	
	%	(95% CI)†	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school										
Sex										
Male	34.7	(±2.3)	22.0	(±1.5)	12.4	(±2.0)	5.6	(±0.9)	6.5	(±3.1)
Female	31.4	(±2.5)	11.5	(±1.2)	3.9	(±0.8)	3.1	(±0.5)	3.8	(±2.1)
Race/Ethnicity										
White, non-Hispanic	31.3	(±2.9)	16.1	(±1.4)	8.7	(±1.7)	3.2	(±0.6)	4.9	(±3.3)
Black, non-Hispanic	37.9	(±3.2)	17.1	(±2.4)	6.9	(±1.5)	5.5	(±1.3)	4.9	(±2.0)
Hispanic	35.8	(±3.1)	19.1	(±2.1)	6.0	(±0.9)	6.2	(±0.9)	5.8	(±2.0)
Asian	24.5	(±6.6)	12.7	(±5.8)	4.6	(±2.9)	4.8	(±3.0)	6.2	(±3.9)
Total	33.1	(±2.3)	16.7	(±1.2)	8.2	(±1.3)	4.3	(±0.6)	5.2	(±2.5)
High school										
Sex										
Male	59.6	(±2.5)	44.4	(±2.4)	25.9	(±3.1)	10.9	(±1.6)	10.3	(±1.7)
Female	55.3	(±2.5)	25.0	(±1.7)	6.4	(±1.3)	6.0	(±0.9)	8.2	(±1.9)
Race/Ethnicity										
White, non-Hispanic	57.2	(±2.7)	36.7	(±2.1)	19.6	(±2.5)	7.7	(±1.2)	10.3	(±1.9)
Black, non-Hispanic	57.5	(±3.2)	29.5	(±3.2)	7.0	(±1.5)	10.1	(±2.1)	5.1	(±1.7)
Hispanic	61.4	(±3.2)	33.6	(±2.8)	10.2	(±2.1)	10.4	(±1.7)	8.1	(±1.8)
Asian	40.4	(±5.8)	15.9	(±3.8)	4.7	(±1.9)	6.3	(±2.1)	6.7	(±5.2)
Total	57.4	(±2.1)	34.7	(±1.7)	16.2	(±2.0)	8.5	(±1.0)	9.2	(±1.6)

* Ever having used tobacco products was determined by asking students whether they have ever tried cigarettes, even one or two puffs; tried smoking cigars, cigarillos, or little cigars, even one or two puffs; used chewing tobacco, snuff, or dip, such as Redman,[®] Levi Garrett,[®] Beechnut,[®] Skoal,[®] Skoal Bandits,[®] or Copenhagen[®]; tried smoking bidis, even one or two puffs; or tried smoking kreteks, even one or two puffs.

† Confidence interval.

TABLE 3. Percentage of middle school and high school students who ever used* cigarettes, cigars, smokeless tobacco, bidis, or kreteks, by state — state youth tobacco surveys, United States, 2001–2002

State	Cigarettes		Cigars		Smokeless tobacco		Bidis		Kreteks	
	%	(95% CI)†	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school										
Alabama – Spring 02	49.5	(±5.0)	26.7	(±5.0)	21.0	(±4.9)	9.0	(±2.2)	5.7	(±1.6)
Connecticut – Spring 02	23.4	(±3.8)	14.9	(±2.5)	6.0	(±1.2)	4.2	(±1.1)	3.0	(±0.9)
Delaware – Spring 02	34.3	(±3.6)	16.5	(±2.3)	5.7	(±1.0)	4.6	(±1.3)	2.3	(±1.0)
Florida – Spring 01	32.1	(±2.8)	16.6	(±1.6)	5.9	(±1.2)	5.0	(±0.9)	4.0	(±0.8)
Florida – Spring 02	30.6	(±1.6)	17.0	(±1.1)	7.4	(±0.8)	4.4	(±0.5)	3.9	(±0.4)
Georgia – Fall 01	32.3	(±4.5)	18.7	(±2.6)	12.6	(±2.6)	6.0	(±1.3)	2.5	(±0.8)
Idaho – Spring 01	33.7	(±3.6)	19.6	(±3.1)	10.7	(±1.6)	5.4	(±1.0)	3.1	(±0.9)
Illinois – Spring 02	28.1	(±6.0)	17.2	(±3.1)	7.1	(±1.6)	5.7	(±1.3)	2.9	(±1.2)
Iowa – Spring 02	28.7	(±4.5)	14.9	(±3.2)	10.7	(±2.3)	3.3	(±0.9)	2.7	(±0.9)
Kansas – Fall 02	25.0	(±4.3)	15.2	(±2.5)	10.4	(±2.4)	4.8	(±1.1)	2.3	(±0.8)
Kentucky – Spring 02	44.0	(±4.5)	24.0	(±2.7)	22.3	(±3.3)	6.6	(±1.7)	4.2	(±1.7)
Louisiana – Spring 01	50.0	(±3.4)	28.5	(±2.3)	20.4	(±3.0)	10.7	(±2.1)	6.4	(±1.5)
Maine – Spring 01	28.4	(±3.8)	12.1	(±1.4)	5.5	(±1.0)	NA§		NA	
Maryland – Spring 02	22.2	(±1.3)	14.8	(±0.8)	7.4	(±0.7)	8.2	(±0.8)	2.8	(±0.4)
Massachusetts – Spring 02	27.0	(±4.6)	15.0	(±2.3)	9.0	(±1.7)	5.8	(±1.7)	2.7	(±1.0)
Michigan – Spring 01	32.6	(±3.9)	17.3	(±2.5)	10.8	(±1.8)	7.2	(±1.7)	4.0	(±1.3)
Minnesota – Spring 02	27.4	(±3.4)	16.3	(±2.0)	11.2	(±1.6)	5.1	(±0.9)	2.4	(±0.5)
Mississippi – Fall 02	40.9	(±3.5)	NA		17.8	(±1.4)	NA		NA	
Nebraska – Spring 02	31.9	(±2.8)	NA		11.9	(±1.3)	NA		NA	
New Hampshire – Fall 01	18.5	(±3.5)	11.0	(±1.5)	5.6	(±1.3)	3.6	(±0.8)	2.2	(±0.5)
New Jersey – Fall 01	23.3	(±3.8)	15.8	(±2.2)	6.4	(±1.4)	6.3	(±1.6)	3.9	(±1.1)
New York – Spring 02	28.6	(±8.0)	14.1	(±2.9)	8.4	(±5.3)	3.7	(±1.0)	3.2	(±1.0)
North Carolina – Spring 02	35.1	(±4.7)	21.2	(±3.0)	13.5	(±1.8)	6.9	(±1.5)	3.7	(±0.9)
Ohio – Spring 02	31.7	(±4.7)	18.7	(±3.2)	11.8	(±1.9)	4.1	(±1.4)	2.9	(±1.2)
Oklahoma – Spring 02	37.3	(±4.9)	20.9	(±3.5)	15.4	(±2.0)	6.2	(±1.6)	3.5	(±1.0)
Pennsylvania – Spring 01	36.3	(±3.6)	19.5	(±1.8)	12.4	(±2.0)	5.4	(±0.8)	3.1	(±0.7)
Rhode Island – Spring 01	28.6	(±3.3)	17.1	(±2.8)	5.8	(±1.6)	6.5	(±1.4)	3.6	(±1.4)
South Dakota – Fall 01	33.9	(±6.7)	17.5	(±3.1)	19.9	(±4.0)	3.8	(±0.8)	2.4	(±0.7)
Texas – Spring 01	35.3	(±4.2)	22.0	(±3.2)	11.0	(±1.6)	7.2	(±1.5)	7.7	(±1.5)
Vermont – Spring 02	30.7	(±4.2)	17.3	(±2.4)	10.6	(±1.9)	4.4	(±0.6)	3.0	(±0.8)
West Virginia – Spring 02	44.5	(±3.3)	23.4	(±1.7)	20.2	(±2.0)	5.9	(±0.7)	4.3	(±0.5)
Wisconsin – Spring 02	34.0	(±4.7)	18.6	(±2.8)	10.0	(±2.6)	5.1	(±1.3)	2.5	(±1.1)
Median	32.0		17.3		10.7		5.4		3.1	
High school										
Alabama – Spring 02	66.8	(±3.5)	41.4	(±3.7)	25.7	(±3.7)	8.2	(±1.7)	5.2	(±1.3)
Connecticut – Spring 02	53.2	(±3.3)	34.7	(±2.3)	12.5	(±2.1)	12.6	(±2.6)	10.3	(±2.0)
Delaware – Spring 02	63.8	(±3.5)	38.3	(±3.8)	13.9	(±2.8)	14.0	(±2.1)	6.5	(±1.2)
Florida – Spring 01	53.7	(±3.0)	31.7	(±2.6)	10.7	(±1.9)	9.8	(±1.1)	9.7	(±1.4)
Florida – Spring 02	52.2	(±2.3)	30.3	(±2.2)	10.5	(±1.5)	8.4	(±1.8)	9.0	(±1.1)
Georgia – Fall 01	60.9	(±3.2)	41.0	(±3.0)	23.9	(±2.8)	12.3	(±1.8)	6.6	(±1.1)
Illinois – Spring 02	60.9	(±6.1)	41.5	(±3.9)	13.1	(±3.0)	7.7	(±2.1)	6.8	(±2.7)
Iowa – Spring 02	60.2	(±3.9)	41.9	(±3.2)	24.2	(±3.2)	6.6	(±1.6)	5.7	(±1.6)
Kansas – Fall 02	54.0	(±4.5)	37.8	(±3.6)	22.5	(±3.8)	7.4	(±1.5)	4.6	(±1.3)
Kentucky – Spring 02	69.2	(±3.2)	48.3	(±2.0)	30.6	(±2.7)	9.9	(±1.9)	6.3	(±1.5)
Maryland – Spring 02	51.2	(±1.1)	31.4	(±0.8)	13.0	(±0.6)	13.6	(±0.6)	5.4	(±0.4)
Massachusetts – Spring 02	55.9	(±4.3)	35.6	(±2.5)	12.8	(±3.1)	13.3	(±2.5)	10.6	(±2.6)
Michigan – Spring 01	62.2	(±3.0)	41.5	(±2.8)	20.4	(±2.6)	10.9	(±1.4)	11.1	(±1.4)
Mississippi – Fall 02	61.5	(±2.5)	NA§		20.7	(±2.6)	NA		NA	
Nebraska – Spring 02	60.3	(±3.7)	NA		24.9	(±2.7)	NA		NA	
New Hampshire – Fall 01	57.9	(±4.5)	36.9	(±3.5)	14.4	(±2.4)	12.0	(±2.4)	12.4	(±2.5)
New Jersey – Fall 01	59.6	(±4.2)	38.3	(±2.8)	13.2	(±2.5)	12.5	(±1.3)	8.8	(±1.4)
New York – Spring 02	56.9	(±3.1)	30.6	(±3.9)	13.6	(±6.1)	12.6	(±2.9)	12.2	(±3.9)
North Carolina – Spring 02	64.2	(±3.8)	40.0	(±3.2)	23.0	(±3.1)	12.0	(±2.2)	8.1	(±1.4)
Ohio – Spring 02	60.7	(±4.3)	42.3	(±2.9)	19.5	(±4.2)	10.8	(±2.7)	6.7	(±1.7)
Oklahoma – Spring 02	61.1	(±3.1)	31.6	(±4.2)	28.1	(±4.0)	8.0	(±1.9)	6.3	(±1.3)
Pennsylvania – Spring 01	64.4	(±2.2)	41.6	(±2.9)	20.7	(±2.7)	11.6	(±1.4)	10.9	(±1.3)
Rhode Island – Spring 01	59.9	(±3.0)	37.0	(±2.8)	11.2	(±1.7)	18.8	(±4.4)	12.6	(±2.6)
Texas – Spring 01	63.7	(±3.9)	39.3	(±2.1)	19.1	(±2.4)	11.1	(±2.3)	12.1	(±1.9)
West Virginia – Spring 02	69.7	(±3.6)	46.1	(±4.7)	28.6	(±2.9)	8.5	(±1.9)	8.3	(±1.7)
Wisconsin – Spring 02	60.4	(±3.7)	41.3	(±4.1)	23.2	(±4.0)	8.3	(±1.6)	8.6	(±2.3)
Median	60.6		38.8		20.0		11.0		8.6	

* Ever having used tobacco products was determined by asking students whether they have ever tried cigarettes, even one or two puffs; tried smoking cigars, cigarillos, or little cigars, even one or two puffs; used chewing tobacco, snuff, or dip, such as Redman,[®] Levi Garrett,[®] Beechnut,[®] Skoal,[®] Skoal Bandits,[®] or Copenhagen,[®] or tried smoking bidis or kreteks.

† Confidence interval.

§ Question not asked.

TABLE 4. Percentage of middle school and high school students who were current users* of any tobacco product, cigarettes, cigars, smokeless tobacco, pipes, bidis, or kreteks, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Any tobacco [†]		Cigarettes		Cigars		Smokeless tobacco		Pipes		Bidis		Kreteks	
	%	(95% CI) [§]	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school														
Sex														
Male	14.7	(±1.6)	9.8	(±1.3)	7.9	(±1.1)	5.3	(±1.3)	5.1	(±0.8)	3.1	(±0.6)	2.7	(±0.6)
Female	11.7	(±1.4)	9.7	(±1.4)	4.1	(±0.7)	1.6	(±0.5)	1.9	(±0.4)	1.7	(±0.4)	1.1	(±0.3)
Race/Ethnicity														
White, non-Hispanic	13.2	(±1.8)	10.1	(±1.6)	5.5	(±1.0)	3.8	(±1.1)	2.8	(±0.6)	1.8	(±0.4)	1.5	(±0.4)
Black, non-Hispanic	13.5	(±2.4)	9.0	(±2.3)	7.3	(±1.7)	2.3	(±0.9)	3.9	(±1.4)	3.1	(±1.0)	2.3	(±0.9)
Hispanic	12.5	(±1.9)	8.7	(±1.5)	6.3	(±1.1)	2.7	(±0.7)	4.3	(±0.9)	2.9	(±0.7)	2.6	(±0.7)
Asian	8.6	(±3.3)	7.4	(±3.3)	5.0	(±2.8)	3.5	(±2.7)	4.6	(±2.7)	3.1	(±2.2)	3.8	(±2.9)
Total	13.3	(±1.4)	9.8	(±1.2)	6.0	(±0.7)	3.5	(±0.7)	3.5	(±0.5)	2.4	(±0.3)	2.0	(±0.4)
High school														
Sex														
Male	32.6	(±2.3)	23.9	(±2.1)	16.9	(±1.4)	10.5	(±2.0)	5.0	(±0.9)	3.7	(±0.8)	3.5	(±0.7)
Female	23.7	(±1.8)	21.0	(±1.9)	6.2	(±0.9)	1.2	(±0.3)	1.4	(±0.4)	1.5	(±0.4)	1.8	(±0.5)
Race/Ethnicity														
White, non-Hispanic	30.9	(±2.0)	25.2	(±1.8)	11.8	(±1.0)	7.3	(±1.4)	2.8	(±0.6)	2.2	(±0.5)	2.7	(±0.6)
Black, non-Hispanic	21.7	(±2.9)	13.8	(±2.8)	12.0	(±1.8)	1.8	(±0.8)	3.7	(±1.2)	3.4	(±1.1)	1.9	(±0.8)
Hispanic	24.1	(±2.7)	19.8	(±2.5)	10.8	(±1.5)	3.3	(±1.1)	4.6	(±1.1)	3.5	(±0.9)	3.0	(±0.8)
Asian	14.6	(±3.8)	12.2	(±3.4)	5.4	(±2.3)	2.1	(±1.5)	2.7	(±1.5)	2.9	(±1.6)	2.1	(±1.7)
Total	28.2	(±1.7)	22.5	(±1.6)	11.6	(±0.9)	5.9	(±1.1)	3.2	(±0.6)	2.6	(±0.5)	2.7	(±0.5)

* Current use of tobacco products was determined by asking students on how many of the previous 30 days they had smoked cigarettes, cigars, cigarillos, or little cigars; used chewing tobacco, snuff, or dip; or smoked pipe tobacco, bidis, or kreteks.

[†] Use of cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥ 1 of the previous 30 days.

[§] Confidence interval.

TABLE 5. Percentage of middle school and high school students who were current users* of tobacco products, by state — state youth tobacco surveys, United States, 2001–2002

State	Any tobacco [†]		Cigarettes		Cigars		Smokeless tobacco		Pipes		Bidis	
	%	(95% CI) [§]	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school												
Alabama – Spring 02	25.3	(±3.7)	15.6	(±3.3)	11.3	(±2.8)	9.0	(±2.9)	5.0	(±1.6)	5.4	(±1.1)
Connecticut – Spring 02	10.0	(±2.1)	5.9	(±1.7)	3.5	(±1.2)	1.3	(±0.7)	2.0	(±0.6)	3.4	(±0.8)
Delaware – Spring 02	13.1	(±2.2)	10.3	(±1.9)	3.6	(±1.6)	2.3	(±0.6)	1.9	(±0.6)	2.1	(±0.9)
Florida – Spring 01	13.5	(±1.9) [¶]	9.8	(±1.6)	7.0	(±1.2)	3.0	(±0.7)	NA**		2.6	(±0.7)
Florida – Spring 02	13.0	(±1.0) [¶]	9.2	(±1.0)	7.0	(±0.7)	3.4	(±0.5)	NA		2.6	(±0.7)
Georgia – Fall 01	14.5	(±2.9)	8.9	(±2.1)	5.4	(±1.3)	4.5	(±1.3)	2.5	(±0.7)	2.8	(±0.9)
Idaho – Spring 01	13.4	(±1.8)	9.6	(±1.9)	5.2	(±1.1)	3.3	(±0.8)	3.4	(±0.9)	2.7	(±0.7)
Illinois – Spring 02	12.8	(±3.8)	7.6	(±2.9)	4.4	(±1.9)	2.4	(±0.9)	3.1	(±1.3)	4.3	(±1.2)
Iowa – Spring 02	10.9	(±2.5)	6.8	(±2.0)	3.1	(±1.1)	2.5	(±0.8)	2.1	(±0.7)	1.5	(±0.5)
Kansas – Fall 02	10.6	(±2.7)	5.7	(±1.8)	3.9	(±1.4)	2.5	(±1.0)	2.4	(±0.9)	2.7	(±0.9)
Kentucky – Spring 02	23.6	(±3.8)	15.3	(±3.1)	7.9	(±1.6)	10.9	(±2.5)	4.4	(±1.5)	4.7	(±1.7)
Louisiana – Spring 01	26.3	(±3.6)	17.1	(±2.5)	12.5	(±2.3)	9.9	(±2.8)	6.3	(±2.7)	7.1	(±2.7)
Maine – Spring 01	10.2	(±1.7)	8.6	(±1.6)	3.3	(±0.6)	2.3	(±0.6)	2.1	(±0.6)	NA [¶]	
Maryland – Spring 02	9.6	(±0.8)	5.3	(±0.5)	3.6	(±0.4)	2.1	(±0.3)	3.0	(±0.4)	3.5	(±0.5)
Massachusetts – Spring 02	10.4	(±2.5)	7.1	(±1.9)	3.6	(±1.0)	1.2	(±0.6)	2.4	(±0.9)	3.5	(±1.1)
Michigan – Spring 01	14.2	(±1.8)	9.3	(±1.8)	5.1	(±1.1)	3.0	(±1.1)	4.0	(±1.2)	4.1	(±1.6)
Minnesota – Spring 02	11.2	(±1.9)	7.2	(±1.7)	2.7	(±0.8)	2.2	(±0.6)	2.6	(±0.5)	2.8	(±0.6)
Mississippi – Fall 02	16.4	(±1.6) ^{††}	11.9	(±1.6)	NA		8.7	(±1.0)	NA		NA	
Nebraska – Spring 02	8.6	(±1.3) ^{††}	7.1	(±1.3)	NA		2.6	(±1.0)	NA		NA	
New Hampshire – Fall 01	7.4	(±1.9)	5.1	(±1.5)	1.9	(±0.7)	1.6	(±0.6)	2.2	(±0.6)	2.1	(±0.9)
New Jersey – Fall 01	11.8	(±2.5) [¶]	6.1	(±2.1)	6.0	(±1.7)	3.6	(±1.0)	NA		5.1	(±1.4)
New York – Spring 02	10.5	(±3.3) ^{§§}	6.7	(±2.4)	3.7	(±1.0)	4.0	(±2.2)	2.8	(±1.2)	2.6	(±1.0)
North Carolina – Spring 02	17.4	(±2.6)	11.3	(±1.9)	7.1	(±1.5)	4.1	(±1.0)	3.7	(±1.1)	4.5	(±0.9)
Ohio – Spring 02	15.0	(±2.6)	10.1	(±2.3)	6.3	(±1.9)	3.6	(±0.9)	3.3	(±1.1)	3.2	(±1.2)
Oklahoma – Spring 02	15.8	(±3.1)	10.3	(±2.8)	7.6	(±1.9)	4.9	(±1.1)	4.2	(±1.0)	4.0	(±1.1)
Pennsylvania – Spring 01	17.0	(±2.2)	13.1	(±2.0)	6.3	(±1.7)	4.4	(±1.3)	2.3	(±0.5)	3.0	(±0.5)
Rhode Island – Spring 01	13.2	(±1.7)	9.1	(±1.2)	4.2	(±1.1)	2.1	(±1.1)	2.8	(±1.0)	4.4	(±0.7)
South Dakota – Fall 01	16.8	(±4.5)	11.5	(±3.5)	4.2	(±1.0)	7.9	(±3.3)	3.5	(±1.2)	2.3	(±0.7)
Texas – Spring 01	16.6	(±2.8)	10.2	(±1.8)	8.7	(±1.5)	5.2	(±1.3)	4.6	(±1.0)	4.5	(±1.1)
Vermont – Spring 02	12.1	(±2.0)	8.6	(±1.7)	3.5	(±0.9)	2.7	(±0.6)	3.7	(±0.7)	3.1	(±0.6)
West Virginia – Spring 02	22.7	(±2.1)	16.3	(±2.0)	7.4	(±0.7)	7.8	(±1.2)	4.1	(±0.5)	4.2	(±0.4)
Wisconsin – Spring 02	13.2	(±3.3)	8.7	(±2.9)	4.9	(±1.9)	2.5	(±1.0)	3.2	(±1.5)	2.6	(±1.2)
Median	13.2		9.2		5.0		3.2		3.1		3.2	
High school												
Alabama – Spring 02	34.7	(±3.6)	25.5	(±4.0)	15.4	(±2.2)	9.5	(±2.1)	4.0	(±1.5)	4.4	(±1.3)
Connecticut – Spring 02	27.1	(±2.8)	22.0	(±2.6)	9.1	(±1.6)	3.1	(±0.7)	3.5	(±1.0)	5.7	(±1.4)
Delaware – Spring 02	26.0	(±2.2)	25.9	(±2.1)	10.1	(±1.2)	4.9	(±1.1)	2.7	(±0.7)	3.9	(±0.9)
Florida – Spring 01	25.8	(±2.5) [¶]	19.0	(±2.2)	13.4	(±1.4)	5.0	(±1.0)	NA		3.9	(±0.7)
Florida – Spring 02	24.3	(±2.0) [¶]	17.8	(±1.7)	11.6	(±1.1)	4.6	(±0.6)	NA		3.6	(±0.5)
Georgia – Fall 01	31.7	(±2.7)	23.7	(±2.3)	14.5	(±1.6)	9.5	(±1.9)	4.2	(±0.9)	5.5	(±1.1)
Illinois – Spring 02	35.7	(±5.2)	29.2	(±4.5)	13.9	(±2.9)	5.5	(±2.5)	4.0	(±1.8)	5.7	(±2.4)
Iowa – Spring 02	33.7	(±2.9)	26.7	(±3.0)	14.2	(±2.1)	8.4	(±1.6)	3.8	(±0.9)	2.7	(±0.9)
Kansas – Fall 02	29.3	(±3.9)	21.1	(±3.0)	11.6	(±1.9)	8.3	(±2.4)	3.3	(±1.0)	3.2	(±1.1)
Kentucky – Spring 02	44.3	(±3.1)	34.2	(±3.3)	17.0	(±2.4)	13.5	(±2.2)	4.6	(±1.8)	4.7	(±1.3)
Maryland – Spring 02	26.1	(±0.9)	19.3	(±0.8)	11.3	(±0.6)	5.4	(±0.4)	6.9	(±0.4)	7.3	(±0.5)
Massachusetts – Spring 02	28.5	(±2.7)	20.7	(±2.9)	9.3	(±1.7)	4.3	(±1.6)	4.5	(±1.2)	6.7	(±1.7)
Michigan – Spring 01	34.1	(±3.0)	27.6	(±2.6)	12.9	(±1.3)	6.7	(±1.4)	5.2	(±0.9)	5.5	(±0.9)
Mississippi – Fall 02	26.4	(±2.9) ^{††}	23.1	(±2.6)	NA		9.6	(±1.7)	NA		NA	
Nebraska – Spring 02	30.4	(±2.8) ^{††}	28.2	(±2.9)	NA		8.3	(±1.9)	NA		NA	
New Hampshire – Fall 01	30.7	(±3.9)	25.3	(±3.6)	11.8	(±2.3)	4.5	(±1.5)	3.6	(±1.1)	6.5	(±1.6)
New Jersey – Fall 01	33.6	(±3.1) [¶]	24.5	(±2.8)	17.1	(±2.2)	6.9	(±1.4)	NA		8.5	(±1.3)
New York – Spring 02	26.8	(±2.2) ^{§§}	21.3	(±2.5)	10.5	(±2.0)	6.5	(±3.7)	3.9	(±1.6)	6.1	(±2.0)
North Carolina – Spring 02	35.8	(±3.3)	27.8	(±3.5)	16.4	(±2.1)	8.9	(±1.4)	5.9	(±1.3)	7.4	(±1.8)
Ohio – Spring 02	32.6	(±4.4)	25.7	(±3.2)	15.1	(±3.7)	6.6	(±2.1)	3.3	(±1.2)	3.6	(±1.0)
Oklahoma – Spring 02	31.9	(±3.6)	24.0	(±3.1)	16.4	(±2.6)	10.1	(±2.4)	4.8	(±1.5)	3.4	(±1.0)
Pennsylvania – Spring 01	34.0	(±2.1)	27.6	(±2.1)	12.4	(±1.1)	6.9	(±1.4)	3.8	(±0.6)	5.7	(±1.0)
Rhode Island – Spring 01	32.1	(±3.2)	26.0	(±3.6)	11.5	(±2.2)	3.8	(±1.3)	4.9	(±1.5)	9.5	(±1.7)
Texas – Spring 01	33.4	(±2.2)	24.7	(±2.0)	16.9	(±1.6)	9.1	(±2.1)	4.7	(±1.0)	4.9	(±1.3)
West Virginia – Spring 02	40.9	(±3.3)	33.7	(±3.1)	14.9	(±2.7)	12.4	(±2.0)	4.1	(±1.7)	6.2	(±2.4)
Wisconsin – Spring 02	32.9	(±4.2)	27.1	(±4.1)	11.5	(±2.1)	7.3	(±1.9)	2.7	(±1.0)	3.0	(±1.1)
Median	32.0		25.4		13.2		6.9		4.0		5.5	

* Current use of tobacco products was determined by asking students on how many of the previous 30 days they had smoked cigarettes, cigars, cigarillos, or little cigars; used chewing tobacco, snuff, or dip; or smoked pipe tobacco, bidis, or kreteks.

[†] Use of cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥ 1 of the previous 30 days.

[§] Confidence interval.

[¶] Pipe tobacco use not included because question not asked.

^{**} Question not asked.

^{††} Cigars, pipe tobacco, and bidi use not included because questions not asked.

^{§§} New York survey also included kreteks.

TABLE 6. Percentage of middle school and high school students who ever smoked cigarettes daily* and percentage of current cigarette smokers† who smoked six or more cigarettes per day‡ on the days they smoked, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	All students who ever smoked daily		Current smokers who smoke six or more cigarettes daily§	
	%	(95% CI¶)	%	(95% CI)
Middle school				
Sex				
Male	6.1	(±0.9)	19.8	(±4.1)
Female	5.4	(±1.0)	12.7	(±3.9)
Race/Ethnicity				
White, non-Hispanic	6.2	(±1.1)	15.8	(±3.9)
Black, non-Hispanic	4.4	(±1.2)	18.6	(±7.2)
Hispanic	4.2	(±1.0)	11.7	(±5.1)
Asian	6.0	(±3.2)	38.1	(±28.1)
Total	5.8	(±0.8)	16.6	(±2.9)
High school				
Sex				
Male	18.0	(±2.0)	34.4	(±3.3)
Female	16.0	(±1.8)	27.9	(±3.8)
Race/Ethnicity				
White, non-Hispanic	19.7	(±1.9)	34.5	(±3.2)
Black, non-Hispanic	9.8	(±2.3)	20.4	(±5.8)
Hispanic	12.1	(±2.0)	19.6	(±4.9)
Asian	10.1	(±3.0)	26.3	(±12.9)
Total	17.0	(±1.6)	31.3	(±3.1)

* Students were asked, "Have you ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days?"

† Smoked cigarettes on ≥1 of the previous 30 days.

‡ Students were asked, "During the previous 30 days, on the days you smoked, how many cigarettes did you smoke per day?"

¶ Confidence interval.

TABLE 7. Percentage of all middle school and high school students who ever smoked cigarettes daily* and percentage of current cigarette smokers† who smoked six or more cigarettes per day‡ on the days they smoked, by state — state youth tobacco surveys, United States, 2001–2002

State	All students who ever smoked cigarettes daily		Current cigarette smokers who smoke six or more cigarettes daily	
	%	(95% CI¶)	%	(95% CI)
Middle school				
Alabama – Spring 02	8.7	(±3.0)	20.9	(±8.0)
Connecticut – Spring 02	3.3	(±1.2)	13.9	(±6.7)
Delaware – Spring 02	6.1	(±1.6)	13.8	(±5.4)
Florida – Spring 01	5.7	(±1.2)	13.1	(±3.9)
Florida – Spring 02	4.9	(±0.6)	15.9	(±2.4)
Georgia – Fall 01	4.6	(±1.3)	11.7	(±4.8)
Idaho – Spring 01	7.3	(±1.5)	22.1	(±7.3)
Illinois – Spring 02	4.8	(±1.4)	14.0	(±9.1)
Iowa – Spring 02	4.5	(±0.9)	13.1	(±3.6)
Kansas – Fall 02	2.4	(±1.3)	20.0	(±9.7)
Kentucky – Spring 02	9.6	(±1.6)	25.1	(±5.1)
Louisiana – Spring 01	10.8	(±1.8)	16.5	(±4.1)
Maine – Spring 01	3.9	(±1.0)	23.5	(±4.8)
Maryland – Spring 02	3.4	(±0.4)	16.3	(±3.1)
Massachusetts – Spring 02	4.3	(±1.4)	9.0	(±6.2)
Michigan – Spring 01	5.8	(±1.6)	19.5	(±6.8)
Minnesota – Spring 02	4.5	(±1.1)	8.9	(±3.6)
Mississippi – Fall 02	NA**		11.1	(±4.4)
Nebraska – Spring 02	4.4	(±0.9)	12.1	(±3.5)
New Hampshire – Fall 01	3.7	(±1.4)	14.0	(±10.0)
New Jersey – Fall 01	3.9	(±1.4)	11.4	(±7.3)
New York – Spring 02	4.7	(±2.2)	22.3	(±9.6)
North Carolina – Spring 02	6.6	(±1.3)	12.4	(±2.9)
Ohio – Spring 02	5.9	(±1.6)	18.1	(±7.2)
Oklahoma – Spring 02	6.4	(±2.1)	16.2	(±5.2)
Pennsylvania – Spring 01	7.6	(±1.5)	19.6	(±3.1)
Rhode Island – Spring 01	4.4	(±1.4)	21.4	(±10.5)
South Dakota – Fall 01	5.6	(±1.8)	12.3	(±5.4)
Texas – Spring 01	4.7	(±1.0)	15.5	(±0.5)
Vermont – Spring 02	7.0	(±1.4)	25.9	(±4.9)
West Virginia – Spring 02	10.5	(±1.5)	23.7	(±2.5)
Wisconsin – Spring 02	5.7	(±1.4)	13.0	(±7.1)
Median	4.9		15.7	
High school				
Alabama – Spring 02	19.1	(±3.6)	30.1	(±6.4)
Connecticut – Spring 02	14.8	(±1.7)	27.0	(±4.1)
Delaware – Spring 02	21.6	(±2.3)	38.0	(±6.8)
Florida – Spring 01	14.2	(±1.8)	25.4	(±3.4)
Florida – Spring 02	13.3	(±1.6)	25.6	(±2.5)
Georgia – Fall 01	17.3	(±2.2)	31.1	(±4.3)
Illinois – Spring 02	20.9	(±3.7)	31.0	(±6.3)
Iowa – Spring 02	20.7	(±3.3)	32.1	(±5.2)
Kansas – Fall 02	14.8	(±2.5)	21.7	(±3.8)
Kentucky – Spring 02	28.1	(±3.8)	44.0	(±5.5)
Maryland – Spring 02	14.5	(±0.6)	30.1	(±1.3)
Massachusetts – Spring 02	15.7	(±2.0)	28.2	(±5.3)
Michigan – Spring 01	22.2	(±2.0)	31.4	(±3.3)
Mississippi – Fall 02	NA**		23.0	(±4.7)
Nebraska – Spring 02	21.4	(±2.7)	26.8	(±4.3)
New Hampshire – Fall 01	18.2	(±3.9)	34.6	(±7.3)
New Jersey – Fall 01	16.2	(±2.7)	27.9	(±5.2)
New York – Spring 02	17.3	(±2.5)	36.3	(±4.5)
North Carolina – Spring 02	20.2	(±2.4)	30.3	(±4.4)
Ohio – Spring 02	19.8	(±3.3)	35.6	(±8.8)
Oklahoma – Spring 02	19.2	(±2.9)	32.3	(±7.3)
Pennsylvania – Spring 01	21.3	(±2.5)	35.5	(±2.4)
Rhode Island – Spring 01	19.1	(±4.0)	38.8	(±4.7)
Texas – Spring 01	15.8	(±2.4)	22.0	(±3.5)
West Virginia – Spring 02	29.1	(±3.3)	41.4	(±7.0)
Wisconsin – Spring 02	20.3	(±4.2)	29.4	(±4.5)
Median	19.1		30.7	

* Students were asked, "Have you ever smoked cigarettes daily, that is, at least one cigarette every day for 30 days?"

† Smoked cigarettes on ≥1 of the previous 30 days.

‡ Students were asked, "During the previous 30 days, on the days you smoked, how many cigarettes did you smoke per day?"

¶ Confidence interval.

** Question not asked.

TABLE 8. Percentage of middle school and high school students who first smoked a cigarette* before age 11 years, by sex and race/ethnicity — National Youth Tobacco Survey (NYTS), United States, 2002†

Sex and Race/Ethnicity	%	(95% CI)§
Middle school		
Sex		
Male	9.8	(±1.3)
Female	6.5	(±1.0)
Race/Ethnicity		
White, non-Hispanic	7.8	(±1.3)
Black, non-Hispanic	8.1	(±1.5)
Hispanic	7.5	(±1.2)
Asian	7.9	(±3.4)
Total	8.1	(±1.0)
High school		
Sex		
Male	8.4	(±1.2)
Female	5.0	(±0.8)
Race/Ethnicity		
White, non-Hispanic	6.6	(±1.2)
Black, non-Hispanic	6.1	(±1.5)
Hispanic	7.4	(±1.5)
Asian	5.0	(±2.0)
Total	6.7	(±0.9)

* Age of initiation was determined by asking students how old they were when they smoked a whole cigarette for the first time.

† NYTS did not include questions regarding initiation of cigars or smokeless tobacco.

§ Confidence interval.

TABLE 9. Percentage of middle school and high school students who first smoked a cigarette or cigar or used smokeless tobacco before age 11 years,* by state — state youth tobacco surveys, 2001–2002

State	Cigarette		Cigar		Smokeless tobacco	
	%	(95% CI)†	%	(95% CI)	%	(95% CI)
Middle school						
Alabama – Spring 02	13.7	(±3.2)	8.4	(±2.2)	10.1	(±2.8)
Connecticut – Spring 02	5.1	(±0.9)	2.9	(±1.0)	1.3	(±0.4)
Delaware – Spring 02	9.5	(±1.5)	3.1	(±1.0)	2.8	(±0.6)
Florida – Spring 01	12.6	(±1.7)	4.6	(±0.8)	2.4	(±0.7)
Florida – Spring 02	12.8	(±0.9)	5.1	(±0.5)	3.2	(±0.5)
Georgia – Fall 01	8.8	(±1.9)	5.7	(±1.2)	5.5	(±1.5)
Idaho – Spring 01	10.9	(±1.9)	4.4	(±0.9)	4.1	(±0.9)
Illinois – Spring 02	7.1	(±1.5)	4.6	(±1.2)	2.2	(±0.9)
Iowa – Spring 02	8.3	(±2.2)	5.2	(±1.8)	2.9	(±1.4)
Kansas – Fall 02	6.1	(±1.6)	4.6	(±1.1)	3.7	(±1.4)
Kentucky – Spring 02	13.4	(±2.5)	8.2	(±2.8)	10.2	(±2.6)
Louisiana – Spring 01	14.3	(±1.7)	8.9	(±1.3)	9.1	(±1.8)
Maine – Spring 01	8.7	(±1.4)	4.0	(±0.6)	2.1	(±0.4)
Maryland – Spring 02	4.9	(±0.4)	3.9	(±0.3)	2.4	(±0.3)
Massachusetts – Spring 02	5.2	(±1.3)	3.1	(±0.9)	1.2	(±0.5)
Michigan – Spring 01	9.4	(±1.5)	4.5	(±1.0)	3.4	(±1.1)
Minnesota – Spring 02	7.9	(±1.1)	5.3	(±0.6)	2.9	(±0.6)
Mississippi – Fall 02	11.3	(±1.3)	NA§		NA	
Nebraska – Spring 02	8.5	(±1.5)	NA		5.3	(±1.0)
New Hampshire – Fall 01	5.5	(±1.4)	4.0	(±0.9)	2.2	(±0.7)
New Jersey – Fall 01	5.4	(±1.3)	NA		NA	
New York – Spring 02	6.5	(±2.0)	NA		NA	
North Carolina – Spring 02	10.5	(±1.3)	7.2	(±1.1)	5.3	(±1.1)
Ohio – Spring 02	9.3	(±1.8)	5.8	(±1.4)	4.2	(±1.2)
Oklahoma – Spring 02	10.4	(±2.9)	6.0	(±1.7)	6.7	(±1.2)
Pennsylvania – Spring 01	8.9	(±1.1)	5.1	(±0.7)	3.6	(±0.9)
Rhode Island – Spring 01	7.2	(±0.9)	3.9	(±1.6)	1.7	(±0.7)
South Dakota – Fall 01	10.7	(±2.9)	6.0	(±1.4)	8.5	(±2.0)
Texas – Spring 01	5.7	(±1.2)¶	4.4	(±0.9)	3.3	(±0.6)
Vermont – Spring 02	7.3	(±1.6)	5.1	(±1.7)	3.6	(±0.7)
West Virginia – Spring 02	13.8	(±1.6)	7.9	(±0.9)	9.5	(±1.2)
Wisconsin – Spring 02	9.6	(±1.6)	4.9	(±1.3)	2.7	(±0.8)
Median	8.9**		5.1		3.5	
High School						
Alabama – Spring 02	9.5	(±1.8)	5.2	(±1.2)	5.8	(±1.2)
Connecticut – Spring 02	5.7	(±1.0)	4.1	(±1.0)	1.9	(±0.5)
Delaware – Spring 02	11.0	(±3.0)	2.6	(±0.5)	2.5	(±1.0)
Florida – Spring 01	11.6	(±1.3)	2.8	(±0.5)	2.9	(±0.8)
Florida – Spring 02	11.9	(±1.0)	3.6	(±0.5)	2.7	(±0.5)
Georgia – Fall 01	9.0	(±1.4)	5.0	(±1.0)	7.7	(±1.5)
Illinois – Spring 02	8.8	(±2.4)	4.8	(±1.3)	3.1	(±1.5)
Iowa – Spring 02	8.0	(±2.1)	4.0	(±1.1)	2.9	(±0.8)
Kansas – Fall 02	7.0	(±1.9)	3.6	(±1.0)	3.8	(±1.0)
Kentucky – Spring 02	10.6	(±2.0)	5.4	(±1.6)	8.7	(±2.1)
Maryland – Spring 02	6.6	(±0.4)	4.3	(±0.3)	3.9	(±0.3)
Massachusetts – Spring 02	6.3	(±1.8)	3.0	(±1.1)	1.3	(±0.7)
Michigan – Spring 01	9.4	(±1.6)	5.2	(±0.7)	4.5	(±0.9)
Mississippi – Fall 02	9.1	(±1.1)	NA§		NA	
Nebraska – Spring 02	7.8	(±1.2)	NA		4.9	(±1.0)
New Hampshire – Fall 01	9.7	(±2.0)	4.6	(±1.9)	2.8	(±1.1)
New Jersey – Fall 01	7.8	(±1.6)	NA		NA	
New York – Spring 02	6.2	(±1.8)	NA		NA	
North Carolina – Spring 02	10.4	(±1.5)	5.6	(±0.9)	5.1	(±0.8)
Ohio – Spring 02	7.2	(±1.6)	3.7	(±0.9)	2.3	(±0.7)
Oklahoma – Spring 02	10.0	(±1.6)	4.5	(±1.0)	6.2	(±2.2)
Pennsylvania – Spring 01	7.9	(±0.8)	4.1	(±0.6)	3.2	(±0.5)
Rhode Island – Spring 01	8.7	(±2.6)	3.5	(±0.9)	2.8	(±1.1)
Texas – Spring 01	6.5	(±2.5)¶	3.5	(±0.5)	3.3	(±0.6)
West Virginia – Spring 02	11.4	(±1.6)	6.4	(±1.5)	7.1	(±1.6)
Wisconsin – Spring 02	9.3	(±1.6)	5.4	(±1.2)	3.2	(±1.6)
Median	9.0**		4.3		3.2	

* Age of initiation was determined by asking students how old they were when they smoked a whole cigarette; smoked a cigar, cigarillo, or little cigar; or used chewing tobacco, snuff, or dip for the first time.

† Confidence interval.

§ Question not asked.

¶ Age of initiation was categorized as <10 years.

** Median does not include Texas because age of initiation was categorized as <10 years.

TABLE 10. Percentage of students who have ever smoked cigarettes* who smoked ≥ 100 cigarettes in their lifetime and percentage of all middle school and high school students who frequently used† cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Students who have smoked ≥ 100 cigarettes in lifetime		All students who reported frequent use											
	%	(95% CI)§	Cigarettes % (95% CI)	Cigars % (95% CI)	Smokeless tobacco % (95% CI)	Pipes % (95% CI)	Bidis % (95% CI)	Kreteks % (95% CI)						
Middle school														
Sex														
Male	12.2	(± 2.2)	2.7	(± 0.6)	1.6	(± 0.4)	1.5	(± 0.5)	1.4	(± 0.4)	1.1	(± 0.4)	0.8	(± 0.3)
Female	8.6	(± 2.5)	2.3	(± 0.9)	0.6	(± 0.2)	0.4	(± 0.2)	0.6	(± 0.3)	0.4	(± 0.2)	0.4	(± 0.2)
Race/Ethnicity														
White, non-Hispanic	12.9	(± 2.6)	2.7	(± 0.8)	0.8	(± 0.3)	0.8	(± 0.3)	0.8	(± 0.3)	0.4	(± 0.2)	0.3	(± 0.2)
Black, non-Hispanic	5.7	(± 2.2)	1.9	(± 0.9)	1.6	(± 0.8)	1.1	(± 0.7)	1.6	(± 0.8)	1.4	(± 0.7)	1.0	(± 0.7)
Hispanic	5.6	(± 1.7)	1.5	(± 0.7)	1.3	(± 0.4)	0.9	(± 0.5)	0.9	(± 0.3)	0.8	(± 0.4)	0.8	(± 0.4)
Asian	13.4	(± 9.9)	3.0	(± 2.7)	2.9	(± 2.6)	2.7	(± 2.6)	2.9	(± 2.6)	1.5	(± 1.8)	2.4	(± 2.6)
Total	10.6	(± 1.8)	2.5	(± 0.6)	1.1	(± 0.3)	1.0	(± 0.2)	1.0	(± 0.3)	0.7	(± 0.2)	0.6	(± 0.2)
High school														
Sex														
Male	29.8	(± 2.6)	11.9	(± 1.6)	1.8	(± 0.4)	3.9	(± 1.1)	1.5	(± 0.4)	1.0	(± 0.3)	0.8	(± 0.3)
Female	26.0	(± 2.7)	10.4	(± 1.5)	0.5	(± 0.2)	0.2	(± 0.1)	0.4	(± 0.2)	0.2	(± 0.1)	0.3	(± 0.1)
Race/Ethnicity														
White, non-Hispanic	33.5	(± 2.3)	13.4	(± 1.5)	0.8	(± 0.3)	2.6	(± 0.7)	0.7	(± 0.3)	0.4	(± 0.2)	0.5	(± 0.2)
Black, non-Hispanic	12.1	(± 2.9)	5.4	(± 1.6)	2.1	(± 0.6)	0.7	(± 0.4)	1.2	(± 0.6)	1.0	(± 0.5)	0.6	(± 0.4)
Hispanic	18.4	(± 2.8)	6.2	(± 1.4)	1.6	(± 0.6)	0.9	(± 0.5)	1.4	(± 0.7)	0.9	(± 0.5)	0.7	(± 0.4)
Asian	23.2	(± 5.9)	7.0	(± 2.7)	0.9	(± 0.7)	0.6	(± 0.7)	1.1	(± 0.8)	0.9	(± 0.7)	0.5	(± 0.5)
Total	28.0	(± 2.1)	11.1	(± 1.3)	1.2	(± 0.2)	2.1	(± 0.6)	0.9	(± 0.2)	0.6	(± 0.2)	0.6	(± 0.2)

* Students were asked, "Have you ever tried cigarettes, even one or two puffs?" and "About how many cigarettes have you smoked in your entire life?"

† Use of cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on >20 of the previous 30 days.

§ Confidence interval.

TABLE 11. Percentage of students who have ever smoked cigarettes* who smoked ≥ 100 cigarettes in their lifetime and percentage of middle school and high school students who frequently used† cigarettes, cigars, smokeless tobacco, pipe tobacco, or bidis, by state — state youth tobacco surveys, United States 2001–2002

States	Ever smokers who have smoked ≥ 100 cigarettes in lifetime		All students – frequent use†									
	%	(95% CI)§	Cigarettes		Cigars		Smokeless tobacco		Pipes		Bidis	
			%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school												
Alabama – Spring 02	11.5	(±5.4)	4.3	(±2.1)	2.0	(±1.1)	2.5	(±1.5)	1.0	(±0.7)	1.0	(±0.5)
Connecticut – Spring 02	8.4	(±2.9)	1.5	(±0.7)	0.6	(±0.4)	0.5	(±0.4)	0.4	(±0.3)	0.4	(±0.3)
Delaware – Spring 02	11.7	(±3.5)	2.6	(±1.0)	0.5	(±0.5)	0.9	(±0.5)	0.6	(±0.4)	0.4	(±0.3)
Florida – Spring 01	10.0	(±2.4)	2.1	(±0.8)	0.7	(±0.3)	0.8	(±0.3)	NA¶		0.6	(±0.3)
Florida – Spring 02	10.3	(±1.2)	2.2	(±0.4)	1.0	(±0.2)	0.9	(±0.2)	NA		0.8	(±0.2)
Georgia – Fall 01	8.0	(±2.5)	1.6	(±0.6)	0.3	(±0.2)	0.7	(±0.4)	0.4	(±0.3)	0.4	(±0.3)
Idaho – Spring 01	15.2	(±3.7)	3.3	(±1.2)	0.8	(±0.5)	1.0	(±0.6)	0.7	(±0.4)	0.5	(±0.4)
Illinois – Spring 02	7.9	(±2.0)	1.2	(±0.7)	0.7	(±0.4)	0.8	(±0.5)	0.8	(±0.6)	0.8	(±0.6)
Iowa – Spring 02	8.4	(±2.3)	1.6	(±0.9)	0.5	(±0.5)	0.4	(±0.4)	0.3	(±0.3)	0.2	(±0.2)
Kansas – Fall 02	6.9	(±3.5)	1.3	(±0.8)	0.8	(±0.4)	0.3	(±0.3)	0.2	(±0.3)	0.2	(±0.2)
Kentucky – Spring 02	17.5	(±3.9)	4.5	(±1.0)	1.0	(±0.6)	3.1	(±1.1)	1.1	(±0.7)	0.7	(±0.6)
Louisiana – Spring 01	15.7	(±2.1)	4.8	(±1.3)	1.4	(±0.5)	2.4	(±0.6)	0.8	(±0.6)	0.9	(±0.7)
Maine – Spring 01	17.1	(±4.6)	2.7	(±0.7)	0.5	(±0.2)	0.7	(±0.3)	0.6	(±0.2)	NA	
Maryland – Spring 02	6.2	(±1.0)	1.2	(±0.2)	0.5	(±0.1)	0.5	(±0.2)	0.5	(±0.1)	0.4	(±0.1)
Massachusetts – Spring 02	6.8	(±3.3)	1.4	(±0.7)	0.7	(±0.5)	0.3	(±0.2)	0.5	(±0.3)	0.2	(±0.2)
Michigan – Spring 01	14.0	(±3.3)	2.4	(±0.9)	0.5	(±0.4)	0.7	(±0.5)	0.7	(±0.4)	0.6	(±0.4)
Minnesota – Spring 02	8.7	(±2.5)	1.5	(±0.6)	0.4	(±0.1)	0.5	(±0.2)	0.4	(±0.2)	0.3	(±0.2)
Mississippi – Fall 02	NA		2.7	(±0.9)	NA		2.1	(±0.6)	NA		NA	
Nebraska – Spring 02	7.2	(±1.8)	1.8	(±0.6)	NA		0.4	(±0.4)	NA		NA	
New Hampshire – Fall 01	8.1	(±4.5)	1.5	(±0.8)	0.1	(±0.2)	0.2	(±0.2)	0.4	(±0.4)	0.1	(±0.2)
New Jersey – Fall 01	5.9	(±2.2)	1.1	(±0.6)	0.7	(±0.3)	0.5	(±0.4)	NA		0.4	(±0.3)
New York – Spring 02	10.0	(±3.5)	2.1	(±1.4)	1.1	(±0.8)	1.0	(±0.8)	0.8	(±0.8)	0.8	(±0.8)
North Carolina – Spring 02	8.5	(±1.8)	2.4	(±0.7)	0.7	(±0.2)	0.8	(±0.3)	0.6	(±0.3)	0.3	(±0.2)
Ohio – Spring 02	13.7	(±4.1)	3.2	(±0.9)	0.8	(±0.6)	0.8	(±0.7)	0.5	(±0.4)	0.5	(±0.4)
Oklahoma – Spring 02	10.1	(±3.5)	2.8	(±1.2)	0.6	(±0.4)	0.8	(±0.4)	0.7	(±0.4)	0.3	(±0.2)
Pennsylvania – Spring 01	16.7	(±3.8)	4.0	(±0.8)	0.7	(±0.4)	1.1	(±0.4)	0.4	(±0.2)	0.4	(±0.2)
Rhode Island – Spring 01	10.6	(±4.8)	2.4	(±1.1)	0.9	(±0.3)	0.6	(±0.4)	0.7	(±0.4)	0.9	(±0.4)
South Dakota – Fall 01	12.2	(±4.4)	2.5	(±1.2)	0.7	(±0.4)	1.6	(±0.8)	0.6	(±0.4)	0.3	(±0.2)
Texas – Spring 01	6.4	(±2.1)	1.9	(±0.8)	1.0	(±0.4)	1.0	(±0.3)	0.7	(±0.3)	0.6	(±0.3)
Vermont – Spring 02	13.2	(±2.6)	2.9	(±0.8)	0.7	(±0.2)	0.6	(±0.4)	0.9	(±0.4)	0.5	(±0.3)
West Virginia – Spring 02	17.3	(±2.0)	5.4	(±0.9)	0.9	(±0.2)	1.9	(±0.4)	0.6	(±0.1)	0.7	(±0.2)
Wisconsin – Spring 02	10.7	(±3.7)	2.5	(±1.1)	0.5	(±0.5)	0.2	(±0.4)	0.3	(±0.3)	0.2	(±0.3)
Median	10.1		2.4		0.7		0.8		0.6		0.4	
High school												
Alabama – Spring 02	27.4	(±5.1)	11.7	(±3.0)	1.9	(±0.9)	2.5	(±1.1)	0.6	(±0.5)	0.4	(±0.3)
Connecticut – Spring 02	27.3	(±3.0)	10.2	(±1.9)	1.2	(±0.5)	0.3	(±0.2)	0.6	(±0.4)	0.4	(±0.3)
Delaware – Spring 02	30.5	(±2.5)	14.2	(±2.6)	1.0	(±0.5)	1.0	(±0.6)	0.9	(±0.3)	1.0	(±0.3)
Florida – Spring 01	22.4	(±2.3)	8.5	(±1.2)	1.5	(±0.5)	1.7	(±0.6)	NA¶		0.8	(±0.3)
Florida – Spring 02	22.3	(±2.2)	7.4	(±1.1)	1.6	(±0.3)	1.4	(±0.4)	NA		1.0	(±0.3)
Georgia – Fall 01	26.8	(±2.8)	10.6	(±1.4)	1.3	(±0.4)	3.0	(±1.3)	0.9	(±0.3)	0.9	(±0.4)
Illinois – Spring 02	31.7	(±6.7)	12.3	(±3.2)	1.1	(±0.8)	1.1	(±0.7)	0.4	(±0.4)	0.6	(±0.4)
Iowa – Spring 02	33.6	(±5.4)	12.9	(±2.6)	1.0	(±0.5)	2.6	(±1.1)	0.7	(±0.5)	0.6	(±0.4)
Kansas – Fall 02	27.4	(±3.9)	8.7	(±1.6)	0.8	(±0.5)	3.3	(±1.2)	0.8	(±0.5)	0.7	(±0.4)
Kentucky – Spring 02	39.2	(±3.7)	20.2	(±2.7)	1.4	(±0.8)	4.9	(±1.3)	0.3	(±0.3)	0.3	(±0.2)
Maryland – Spring 02	22.9	(±0.9)	8.3	(±0.5)	1.5	(±0.2)	1.4	(±0.2)	1.4	(±0.1)	1.1	(±0.1)
Massachusetts – Spring 02	26.3	(±3.1)	9.7	(±2.0)	1.1	(±0.6)	1.1	(±0.6)	0.6	(±0.5)	0.7	(±0.4)
Michigan – Spring 01	33.7	(±3.2)	13.9	(±1.4)	1.4	(±0.5)	1.8	(±0.6)	1.1	(±0.4)	0.9	(±0.3)
Mississippi – Fall 02	NA		9.0	(±1.6)	NA		3.3	(±1.0)	NA		NA	
Nebraska – Spring 02	33.5	(±4.2)	13.8	(±2.3)	NA		2.5	(±0.7)	NA		NA	
New Hampshire – Fall 01	32.3	(±4.4)	11.7	(±2.7)	0.7	(±0.4)	0.6	(±0.4)	1.1	(±0.6)	0.6	(±0.4)
New Jersey – Fall 01	25.0	(±4.5)	10.6	(±2.0)	1.3	(±0.5)	1.2	(±0.6)	NA		1.4	(±0.7)
New York – Spring 02	27.8	(±3.5)	11.2	(±2.4)	1.8	(±0.8)	2.0	(±1.2)	1.7	(±0.8)	1.8	(±1.4)
North Carolina – Spring 02	29.0	(±3.5)	12.7	(±2.8)	2.2	(±0.5)	2.8	(±0.6)	1.3	(±0.5)	0.8	(±0.4)
Ohio – Spring 02	32.4	(±4.7)	13.4	(±3.2)	1.9	(±1.1)	1.5	(±1.2)	0.3	(±0.4)	0.2	(±0.3)
Oklahoma – Spring 02	30.8	(±4.8)	12.5	(±2.5)	1.5	(±0.6)	3.5	(±1.2)	0.6	(±0.4)	0.6	(±0.3)
Pennsylvania – Spring 01	32.8	(±3.1)	14.7	(±1.8)	1.0	(±0.3)	2.5	(±0.6)	0.7	(±0.2)	0.8	(±0.3)
Rhode Island – Spring 01	28.6	(±4.5)	13.2	(±2.6)	1.3	(±0.8)	1.1	(±0.7)	1.2	(±0.8)	1.1	(±0.7)
Texas – Spring 01	20.3	(±2.5)	8.1	(±1.3)	1.6	(±0.4)	2.8	(±0.7)	0.7	(±0.3)	0.8	(±0.4)
West Virginia – Spring 02	38.6	(±4.7)	18.7	(±3.3)	1.3	(±0.8)	4.6	(±1.3)	0.6	(±0.4)	0.8	(±0.6)
Wisconsin – Spring 02	34.0	(±6.4)	14.4	(±3.4)	0.4	(±0.5)	1.6	(±0.5)	0.0	(±0.0)	0.4	(±0.1)
Median	29.0		12.0		1.3		1.9		0.7		0.8	

* Students were asked, "Have you ever tried cigarettes, even one or two puffs?" and "About how many cigarettes have you smoked in your entire life?"

† Use of cigarettes or cigars or smokeless tobacco or pipes or bidis or kreteks on ≥ 20 of the previous 30 days.

§ Confidence interval.

¶ Question not asked.

TABLE 12. Brand of cigarettes usually smoked by current cigarette smokers* during the previous 30 days, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Marlboro®		Newport®		Camel®		Other†		No usual	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school										
Sex										
Male	41.7	(±6.0)	26.0	(±7.5)	6.4	(±2.1)	16.4	(±3.9)	9.5	(±3.0)
Female	42.2	(±5.9)	26.1	(±6.8)	5.1	(±2.8)	18.4	(±5.9)	8.2	(±2.6)
Race/Ethnicity										
White, non-Hispanic	47.6	(±6.2)	22.3	(±7.0)	5.4	(±2.8)	16.9	(±5.0)	7.8	(±2.4)
Black, non-Hispanic	9.0	(±7.3)	58.3	(±10.3)	7.0	(±6.7)	16.9	(±6.9)	8.8	(±5.1)
Hispanic	46.9	(±7.9)	20.4	(±7.5)	3.3	(±2.2)	16.0	(±5.0)	13.4	(±5.9)
Asian	30.9	(±18.1)	25.0	(±26.1)	8.4	(±10.0)	12.4	(±11.6)	23.4	(±24.0)
Total	41.8	(±4.7)	26.3	(±5.8)	5.8	(±2.0)	17.4	(±3.7)	8.8	(±2.1)
High school										
Sex										
Male	51.5	(±4.3)	18.6	(±4.8)	10.7	(±3.5)	13.0	(±2.9)	6.2	(±1.4)
Female	52.6	(±4.7)	24.9	(±5.3)	6.5	(±2.4)	10.1	(±2.6)	5.8	(±1.6)
Race/Ethnicity										
White, non-Hispanic	56.6	(±4.1)	16.6	(±4.8)	9.5	(±3.2)	10.9	(±2.7)	6.4	(±1.4)
Black, non-Hispanic	12.8	(±5.2)	66.8	(±7.8)	4.5	(±3.6)	11.9	(±5.0)	3.9	(±2.4)
Hispanic	53.0	(±6.5)	21.5	(±6.2)	5.0	(±2.5)	14.6	(±4.0)	6.0	(±2.4)
Asian	55.1	(±13.3)	16.4	(±14.0)	11.5	(±8.3)	15.3	(±10.6)	1.6	(±2.3)
Total	52.0	(±3.7)	21.6	(±4.6)	8.7	(±2.8)	11.7	(±2.3)	6.0	(±1.1)

* Smoked cigarettes on ≥1 of the previous 30 days.

† Includes Kool,® Lucky Strike,® Virginia Slims,® GPC,® Basic,® American Spirit,® Parliament,® and Doral.® The 2002 NYTS included two additional brands, American Spirit and Parliament, that were not included in the 2000 NYTS.

§ Confidence interval.

TABLE 13. Brand of cigarettes usually smoked by current cigarette smokers* during the previous 30 days, by state — state youth tobacco surveys, United States, 2001–2002

State	Marlboro®		Newport®		Camel®		Other†		No usual	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school										
Alabama – Spring 02	29.3	(±8.7)	39.6	(±9.0)	7.5	(±4.5)	14.8	(±7.0)	8.8	(±4.7)
Connecticut – Spring 02	29.2	(±6.1)	47.5	(±9.9)	2.7	(±3.0)	7.6	(±5.2)	13.0	(±7.1)
Delaware – Spring 02	24.4	(±6.5)	52.7	(±8.3)	2.4	(±2.1)	10.8	(±4.4)	9.7	(±4.5)
Florida – Spring 01	NA¶		NA		NA		NA		NA	
Florida – Spring 02	NA		NA		NA		NA		NA	
Georgia – Fall 01	39.3	(±9.4)	23.2	(±8.2)	3.5	(±2.3)	19.1	(±6.8)	15.0	(±5.6)
Idaho – Spring 01	43.9	(±7.1)	2.1	(±2.4)	26.0	(±7.7)	16.1	(±3.8)	11.8	(±4.6)
Illinois – Spring 02	40.3	(±12.4)	29.3	(±7.4)	7.1	(±3.1)	14.0	(±9.3)	9.2	(±4.8)
Iowa – Spring 02	58.7	(±10.7)	9.8	(±9.1)	6.1	(±5.6)	15.6	(±8.8)	9.8	(±6.4)
Kansas – Fall 02	41.2	(±17.1)	14.7	(±10.5)	6.7	(±4.5)	19.4	(±12.4)	17.9	(±12.0)
Kentucky – Spring 02	57.6	(±6.8)	8.6	(±3.7)	5.5	(±3.8)	15.9	(±4.5)	12.5	(±4.6)
Louisiana – Spring 01	52.5	(±6.7)	16.7	(±5.7)	4.9	(±1.8)	16.1	(±3.1)	9.9	(±3.9)
Maine – Spring 01	52.4	(±6.6)	5.0	(±2.2)	10.3	(±3.0)	18.4	(±5.7)	13.9	(±3.1)
Maryland – Spring 02	25.0	(±3.5)	46.3	(±4.3)	9.4	(±3.2)	75.2	(±3.7)	19.7	(±2.8)
Massachusetts – Spring 02	41.6	(±12.6)	32.8	(±11.3)	2.8	(±2.9)	12.6	(±6.5)	10.2	(±6.4)
Michigan – Spring 01	44.7	(±10.3)	21.9	(±8.1)	5.0	(±2.8)	12.1	(±5.3)	16.4	(±5.7)
Minnesota – Spring 02	50.2	(±6.7)	13.8	(±6.2)	10.5	(±3.6)	12.0	(±4.3)	13.6	(±4.1)
Mississippi – Fall 02	NA		NA		NA		NA		NA	
Nebraska – Spring 02	49.8	(±8.4)	10.8	(±4.5)	3.2	(±2.6)	17.0	(±6.2)	19.3	(±5.4)
New Hampshire – Fall 01	63.5	(±15.9)	11.1	(±7.9)	6.1	(±5.5)	8.2	(±6.5)	11.0	(±8.5)
New Jersey – Fall 01	25.6	(±8.8)	44.5	(±10.0)	3.1	(±3.0)	16.9	(±8.6)	9.9	(±4.4)
New York – Spring 02	28.7	(±13.9)	36.5	(±13.7)	3.1	(±3.3)	22.5	(±10.0)	9.2	(±4.9)
North Carolina – Spring 02	31.9	(±6.0)	40.9	(±7.8)	4.7	(±2.6)	12.4	(±4.0)	10.1	(±2.4)
Ohio – Spring 02	55.3	(±12.0)	14.2	(±8.0)	5.6	(±3.9)	12.6	(±4.0)	12.3	(±6.2)
Oklahoma – Spring 02	56.0	(±9.5)	7.4	(±4.8)	5.7	(±3.1)	18.3	(±6.5)	12.6	(±4.9)
Pennsylvania – Spring 01	40.7	(±12.6)	34.6	(±10.4)	4.4	(±2.3)	11.1	(±4.6)	9.2	(±2.5)
Rhode Island – Spring 01	30.3	(±14.3)	39.0	(±13.1)	4.6	(±4.2)	12.7	(±6.1)	13.4	(±8.3)
South Dakota – Fall 01	52.0	(±11.0)	5.7	(±3.3)	15.2	(±9.3)	15.8	(±8.2)	11.3	(±5.5)
Texas – Spring 01	52.7	(±6.3)	11.7	(±3.7)	5.2	(±2.0)	16.1	(±3.1)	14.4	(±4.5)
Vermont – Spring 02	52.9	(±4.6)	7.3	(±2.8)	14.2	(±4.5)	13.1	(±3.5)	12.5	(±3.4)
West Virginia – Spring 02	49.7	(±4.4)	10.3	(±3.6)	5.2	(±1.3)	44.5	(±3.3)	14.5	(±2.2)
Wisconsin – Spring 02	29.0	(±10.5)	35.5	(±15.7)	9.6	(±7.0)	16.1	(±6.1)	9.9	(±8.7)
Median	43.9		16.7		5.5		15.8		12.3	
High school										
Alabama – Spring 02	52.4	(±7.3)	27.3	(±7.9)	8.0	(±3.2)	6.4	(±1.4)	5.9	(±2.8)
Connecticut – Spring 02	36.7	(±4.6)	32.3	(±5.7)	12.3	(±5.5)	9.5	(±3.0)	9.2	(±2.5)
Delaware – Spring 02	31.9	(±4.9)	55.0	(±6.5)	1.6	(±1.2)	5.8	(±2.0)	5.6	(±3.5)
Florida – Spring 01	NA¶		NA		NA		NA		NA	
Florida – Spring 02	NA		NA		NA		NA		NA	
Georgia – Fall 01	52.5	(±5.0)	23.6	(±4.7)	11.9	(±2.9)	5.9	(±2.2)	6.1	(±2.0)
Illinois – Spring 02	44.8	(±7.6)	23.2	(±9.6)	16.8	(±4.9)	11.3	(±4.7)	4.0	(±2.0)
Iowa – Spring 02	59.9	(±6.7)	11.1	(±5.2)	12.7	(±3.3)	7.7	(±2.3)	8.5	(±3.5)
Kansas – Fall 02	53.0	(±6.5)	8.1	(±4.8)	19.3	(±7.1)	7.3	(±3.4)	12.3	(±4.6)
Kentucky – Spring 02	64.1	(±4.6)	10.1	(±2.9)	8.9	(±3.7)	9.7	(±2.6)	7.2	(±3.0)
Maryland – Spring 02	32.8	(±1.5)	41.1	(±1.6)	10.1	(±1.3)	81.5	(±1.1)	13.8	(±0.9)
Massachusetts – Spring 02	59.7	(±6.4)	17.4	(±5.4)	4.5	(±2.7)	6.8	(±3.3)	11.6	(±4.6)
Michigan – Spring 01	53.4	(±4.6)	19.5	(±6.1)	10.3	(±2.5)	8.6	(±2.2)	8.2	(±1.8)
Mississippi – Fall 02	NA		NA		NA		NA		NA	
Nebraska – Spring 02	58.3	(±3.4)	14.1	(±2.9)	9.6	(±2.7)	7.6	(±1.8)	10.4	(±3.2)
New Hampshire – Fall 01	47.6	(±6.0)	16.3	(±4.7)	13.6	(±5.5)	11.9	(±4.9)	10.5	(±4.6)
New Jersey – Fall 01	34.2	(±4.8)	41.1	(±7.1)	3.8	(±2.0)	10.2	(±2.9)	10.8	(±3.8)
New York – Spring 02	41.6	(±8.0)	29.2	(±9.4)	5.5	(±2.4)	15.1	(±6.3)	8.6	(±3.1)
North Carolina – Spring 02	39.5	(±4.0)	37.2	(±4.8)	9.2	(±2.5)	8.2	(±1.9)	5.9	(±1.6)
Ohio – Spring 02	57.2	(±6.0)	19.4	(±6.8)	10.4	(±5.3)	5.8	(±2.4)	7.1	(±3.3)
Oklahoma – Spring 02	56.5	(±5.6)	6.9	(±2.2)	9.5	(±4.3)	16.9	(±7.3)	10.2	(±3.6)
Pennsylvania – Spring 01	46.7	(±5.1)	31.3	(±5.3)	7.5	(±1.6)	7.4	(±1.8)	7.1	(±1.5)
Rhode Island – Spring 01	44.0	(±7.2)	35.0	(±7.7)	6.2	(±2.2)	7.3	(±3.2)	7.5	(±2.3)
Texas – Spring 01	64.4	(±5.4)	11.9	(±2.7)	7.8	(±2.1)	9.1	(±3.1)	6.8	(±2.2)
West Virginia – Spring 02	56.4	(±3.8)	14.3	(±4.3)	7.9	(±2.3)	69.7	(±3.6)	6.9	(±2.8)
Wisconsin – Spring 02	44.3	(±7.6)	16.8	(±5.3)	21.8	(±5.7)	8.6	(±2.9)	8.6	(±2.3)
Median	52.4		19.5		9.5		8.6		8.2	

* Smoked cigarettes on ≥1 of the previous 30 days.

† Includes Basic,® Doral,® GPC,® Virginia Slims,® and "some other brand."

§ Confidence interval.

¶ Question not asked.

TABLE 14. Cigarette smoking intentions and susceptibility* to initiate cigarette smoking among students who have never smoked cigarettes,† by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Will not try a cigarette soon		Will definitely not smoke in the next year		Would definitely not smoke if best friend offered a cigarette		Susceptible	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Sex								
Male	96.8	(±0.7)	84.9	(±1.5)	85.1	(±1.7)	21.8	(±1.8)
Female	96.6	(±0.6)	84.7	(±1.6)	85.6	(±1.6)	20.9	(±1.8)
Race/Ethnicity								
White, non-Hispanic	97.4	(±0.6)	84.6	(±1.7)	85.1	(±1.5)	20.9	(±1.9)
Black, non-Hispanic	96.3	(±1.2)	87.0	(±1.8)	88.1	(±2.4)	20.6	(±2.5)
Hispanic	94.2	(±1.5)	83.9	(±1.9)	84.1	(±2.3)	23.4	(±2.3)
Asian	96.1	(±2.9)	85.8	(±5.6)	85.9	(±5.8)	18.4	(±6.1)
Total	96.7	(±0.5)	84.8	(±1.3)	85.3	(±1.3)	21.3	(±1.5)
High school								
Sex								
Male	97.7	(±0.8)	84.1	(±1.9)	85.7	(±1.8)	21.5	(±2.7)
Female	97.7	(±0.7)	82.0	(±2.5)	84.7	(±2.2)	24.1	(±3.5)
Race/Ethnicity								
White, non-Hispanic	98.1	(±0.6)	82.8	(±2.2)	85.1	(±2.0)	21.6	(±2.8)
Black, non-Hispanic	98.5	(±1.2)	86.8	(±3.5)	89.8	(±2.9)	18.4	(±3.9)
Hispanic	94.9	(±1.8)	79.4	(±3.1)	80.4	(±3.0)	31.7	(±6.0)
Asian	97.9	(±1.6)	81.6	(±4.2)	82.9	(±4.6)	30.4	(±8.7)
Total	97.7	(±0.6)	83.0	(±1.8)	85.1	(±1.7)	22.9	(±2.8)

* Students were considered not susceptible if they answered, "no, will not try a cigarette soon"; "definitely no, will not smoke a cigarette in the next year"; and "definitely no, would not smoke a cigarette if best friend offered a cigarette." All other students were classified as susceptible.

† Students were considered never to have smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

§ Confidence interval.

TABLE 15. Cigarette smoking intentions and susceptibility* to initiate cigarette smoking among students who have never smoked cigarettes,† by state — state youth tobacco surveys, 2001–2002

State	Will not try a cigarette soon		Will definitely not smoke in the next year		Would definitely not smoke if best friend offered a cigarette		Susceptible	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Alabama – Spring 02	94.8	(±2.3)	84.7	(±2.9)	84.4	(±2.6)	25.0	(±4.1)
Connecticut – Spring 02	95.6	(±1.5)	84.5	(±3.4)	85.6	(±2.6)	23.4	(±3.7)
Delaware – Spring 02	97.2	(±1.1)	86.1	(±2.0)	85.7	(±2.1)	26.8	(±2.2)
Florida – Spring 01	94.7	(±1.0)	81.3	(±1.9)	83.0	(±1.7)	24.5	(±2.3)
Florida – Spring 02	95.7	(±0.6)	83.0	(±1.0)	84.4	(±0.9)	22.2	(±1.1)
Georgia – Fall 01	95.2	(±1.3)	85.5	(±2.1)	85.5	(±2.4)	24.3	(±3.4)
Idaho – Spring 01	95.6	(±1.3)	88.7	(±2.7)	89.3	(±2.1)	17.4	(±3.4)
Illinois – Spring 02	94.1	(±1.6)	82.6	(±3.7)	83.3	(±3.3)	26.7	(±3.5)
Iowa – Spring 02	96.4	(±1.0)	84.2	(±2.6)	85.5	(±2.3)	23.0	(±3.1)
Kansas – Fall 02	96.0	(±1.6)	82.9	(±2.5)	83.2	(±2.2)	26.9	(±3.4)
Kentucky – Spring 02	97.1	(±1.1)	87.6	(±2.4)	86.8	(±2.3)	19.6	(±3.3)
Louisiana – Spring 01	96.0	(±0.8)	84.3	(±2.1)	86.6	(±2.4)	22.3	(±2.6)
Maine – Spring 01	97.6	(±0.8)	84.1	(±1.7)	84.5	(±1.7)	23.7	(±1.9)
Maryland – Spring 02	94.0	(±0.6)	85.5	(±1.0)	86.0	(±1.1)	24.2	(±1.3)
Massachusetts – Spring 02	95.7	(±1.2)	85.9	(±2.4)	86.3	(±2.9)	21.6	(±2.7)
Michigan – Spring 01	96.4	(±1.1)	84.7	(±2.2)	83.3	(±2.4)	26.8	(±3.1)
Minnesota – Spring 02	95.8	(±1.0)	84.0	(±1.8)	83.5	(±1.9)	24.6	(±2.1)
Mississippi – Fall 02	NA¶		85.0	(±2.0)	84.4	(±2.6)	NA	
Nebraska – Spring 02	96.1	(±1.4)	84.5	(±2.0)	84.5	(±2.0)	23.8	(±2.5)
New Hampshire – Fall 01	96.3	(±1.3)	85.3	(±2.5)	85.4	(±2.5)	23.5	(±3.1)
New Jersey – Fall 01	92.8	(±1.7)	78.2	(±2.3)	81.2	(±1.6)	33.3	(±2.5)
New York – Spring 02	96.1	(±1.2)	84.2	(±3.5)	84.5	(±2.5)	22.1	(±3.9)
North Carolina – Spring 02	95.5	(±0.9)	86.0	(±1.9)	85.1	(±2.1)	23.3	(±2.6)
Ohio – Spring 02	95.8	(±1.1)	83.3	(±3.1)	84.4	(±2.7)	24.4	(±2.8)
Oklahoma – Spring 02	NA		87.4	(±2.8)	87.9	(±2.1)	NA	
Pennsylvania – Spring 01	95.0	(±1.1)	81.5	(±1.7)	81.6	(±2.5)	26.8	(±2.4)
Rhode Island – Spring 01	95.9	(±1.6)	83.7	(±2.8)	85.8	(±2.3)	23.6	(±2.7)
South Dakota – Fall 01	96.2	(±1.2)	85.6	(±2.3)	85.1	(±1.9)	22.5	(±2.5)
Texas – Spring 01	94.6	(±1.1)	96.4	(±1.1)	93.6	(±1.1)	25.1	(±2.3)
Vermont – Spring 02	96.4	(±0.9)	86.0	(±2.4)	85.1	(±1.6)	22.2	(±1.9)
West Virginia – Spring 02	95.8	(±0.8)	82.7	(±1.4)	83.6	(±0.9)	25.4	(±1.3)
Wisconsin – Spring 02	95.1	(±2.0)	84.5	(±2.7)	84.5	(±3.2)	24.0	(±4.2)
Median	95.8		84.5		84.8		23.9	
High school								
Alabama – Spring 02	97.1	(±1.8)	83.7	(±5.2)	85.9	(±4.4)	21.1	(±5.2)
Connecticut – Spring 02	95.0	(±1.4)	80.2	(±2.4)	81.9	(±2.6)	27.7	(±2.9)
Delaware – Spring 02	97.8	(±1.3)	85.5	(±2.5)	87.9	(±1.9)	23.8	(±2.9)
Florida – Spring 01	97.4	(±0.7)	83.4	(±1.9)	86.5	(±1.7)	19.7	(±2.0)
Florida – Spring 02	98.1	(±0.5)	85.2	(±1.5)	87.6	(±1.2)	18.4	(±1.7)
Georgia – Fall 01	96.0	(±1.0)	84.0	(±2.7)	85.4	(±2.7)	23.5	(±3.7)
Illinois – Spring 02	94.4	(±2.6)	80.6	(±2.9)	82.7	(±3.1)	26.6	(±3.5)
Iowa – Spring 02	97.6	(±0.8)	81.5	(±3.1)	85.3	(±2.3)	22.5	(±3.0)
Kansas – Fall 02	96.7	(±1.5)	81.3	(±3.7)	83.8	(±3.0)	25.2	(±3.8)
Kentucky – Spring 02	98.7	(±1.1)	83.4	(±3.4)	86.7	(±2.9)	22.9	(±3.2)
Maryland – Spring 02	93.8	(±0.5)	80.8	(±0.7)	82.5	(±0.7)	27.9	(±0.8)
Massachusetts – Spring 02	96.5	(±1.3)	81.2	(±3.6)	83.3	(±3.0)	24.4	(±4.0)
Michigan – Spring 01	96.7	(±1.3)	81.1	(±2.3)	85.9	(±1.8)	23.8	(±2.2)
Mississippi – Fall 02	NA		85.5	(±2.1)	85.8	(±2.3)	NA	
Nebraska	95.4	(±1.3)	81.1	(±2.6)	84.0	(±2.2)	25.9	(±3.3)
New Hampshire – Fall 01	96.4	(±1.3)	78.5	(±3.0)	84.9	(±3.2)	29.4	(±3.2)
New Jersey – Fall 01	92.5	(±1.9)	76.2	(±3.5)	81.0	(±3.4)	35.4	(±4.2)
New York – Spring 02	98.4	(±1.0)	84.0	(±2.2)	85.7	(±3.5)	20.2	(±1.3)
North Carolina – Spring 02	96.1	(±1.3)	80.7	(±1.4)	83.8	(±2.3)	25.8	(±2.5)
Ohio – Spring 02	95.4	(±1.9)	84.3	(±5.9)	85.6	(±4.8)	22.7	(±5.7)
Oklahoma – Spring 02	NA		85.8	(±3.2)	89.0	(±2.8)	NA	
Pennsylvania – Spring 01	96.8	(±0.7)	84.8	(±2.0)	86.9	(±1.5)	21.6	(±2.1)
Rhode Island – Spring 01	96.1	(±1.7)	83.2	(±3.7)	85.9	(±3.3)	23.1	(±3.7)
Texas – Spring 01	94.3	(±1.7)	95.7	(±1.7)	93.7	(±1.9)	26.7	(±3.4)
West Virginia – Spring 02	96.6	(±1.5)	82.9	(±4.2)	85.5	(±3.3)	22.9	(±5.1)
Wisconsin – Spring 02	95.5	(±1.7)	81.0	(±4.5)	83.5	(±3.4)	24.9	(±4.4)
Median	96.5		83.1		85.6		23.8	

* Students were considered not susceptible if they answered “no, will not try a cigarette soon”; “definitely no, will not smoke a cigarette in the next year”; and “definitely no, would not smoke a cigarette if best friend offered a cigarette.” All other students were classified as susceptible.

† Students were considered never to have smoked if they answered “no” to whether they have tried or experimented with cigarette smoking, even one or two puffs.

§ Confidence interval.

¶ Question not asked.

TABLE 16. Percentage of middle school and high school students with closest friends who use tobacco,* by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2001–2002

Sex and Race/Ethnicity	One or more closest friends smoke cigarettes				One or more closest friends use smokeless tobacco (SLT)			
	Never smoked [†]		Currently smoke [§]		Never used SLT [¶]		Currently use SLT ^{**}	
	%	(95% CI) ^{††}	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Sex								
Male	15.0	(±2.2)	86.5	(±3.9)	6.9	(±1.3)	77.3	(±8.4)
Female	15.7	(±2.2)	92.9	(±2.8)	5.9	(±1.1)	62.9	(±14.1)
Race/Ethnicity								
White, non-Hispanic	15.0	(±2.2)	91.1	(±3.0)	6.6	(±1.4)	74.0	(±8.7)
Black, non-Hispanic	15.3	(±2.6)	85.2	(±6.0)	4.2	(±1.2)	70.1	(±16.1)
Hispanic	16.2	(±2.5)	88.3	(±5.8)	6.7	(±1.2)	76.2	(±13.2)
Asian	11.9	(±4.6)		§§	7.9	(±3.6)		§§
Total	15.4	(±1.9)	89.7	(±2.6)	6.4	(±1.0)	74.2	(±7.3)
High school								
Sex								
Male	28.8	(±3.3)	90.7	(±1.5)	13.6	(±2.3)	84.9	(±3.8)
Female	26.9	(±2.7)	91.6	(±2.0)	10.4	(±1.5)	66.4	(±13.0)
Race/Ethnicity								
White, non-Hispanic	27.9	(±3.0)	92.6	(±1.3)	13.9	(±2.1)	83.9	(±4.1)
Black, non-Hispanic	25.4	(±4.5)	82.2	(±5.9)	6.0	(±1.7)		§§
Hispanic	31.5	(±3.4)	90.2	(±3.4)	9.8	(±1.7)	79.7	(±9.9)
Asian	24.1	(±6.5)	89.1	(±8.9)	7.0	(±3.1)		§§
Total	27.8	(±2.2)	91.2	(±1.4)	11.8	(±1.6)	83.0	(±3.8)

* Students were asked, "How many of your four closest friends smoke cigarettes?" and "How many of your four closest friends use chewing tobacco, snuff, or dip?"

[†] Students were considered to have never smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

[§] Smoked cigarettes on ≥ 1 of the previous 30 days.

[¶] Students were considered never to have used SLT if they answered "no" to whether they have ever used SLT.

^{**} Use of smokeless tobacco on ≥ 1 of the previous 30 days.

^{††} Confidence interval.

^{§§} Sample size <35.

TABLE 17. Percentage of middle school and high school students with closest friends who use tobacco,* by tobacco use status and state — state youth tobacco surveys, United States, 2001–2002

State	One or more closest friends smoke cigarettes				One or more closest friends use smokeless tobacco (SLT)			
	Never smoked cigarettes [†]		Currently smoke cigarettes [‡]		Never used SLT ^{**}		Currently use SLT ^{††}	
	%	(95% CI) [§]	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Alabama – Spring 02	20.5	(±4.7)	87.6	(±5.9)	17.7	(±3.9)	78.4	(±9.4)
Connecticut – Spring 02	15.2	(±2.9)	83.9	(±8.0)	9.8	(±1.9)	§§	
Delaware – Spring 02	21.6	(±2.1)	92.1	(±3.5)	12.9	(±1.6)	75.4	(±14.5)
Florida – Spring 01	24.2	(±2.8)	84.6	(±4.6)	13.7	(±1.4)	72.7	(±7.0)
Florida – Spring 02	22.1	(±1.2)	87.7	(±2.2)	13.6	(±0.8)	77.7	(±4.2)
Georgia – Fall 01	18.1	(±2.7)	85.1	(±4.0)	13.9	(±2.3)	69.9	(±10.3)
Idaho – Spring 01	16.2	(±3.0)	91.8	(±4.2)	10.1	(±1.8)	66.7	(±12.0)
Illinois – Spring 02	16.8	(±4.4)	83.5	(±5.4)	10.6	(±3.3)	59.7	(±9.5)
Iowa – Spring 02	18.7	(±4.3)	82.8	(±8.4)	11.1	(±1.7)	66.5	(±17.3)
Kansas – Fall 02	16.9	(±3.8)	80.6	(±10.4)	11.6	(±2.9)	§§	
Kentucky – Spring 02	22.4	(±3.6)	85.7	(±5.3)	22.8	(±3.0)	79.4	(±6.7)
Louisiana – Spring 01	26.1	(±2.6)	91.3	(±2.2)	18.1	(±3.5)	81.5	(±6.5)
Maine – Spring 01	22.2	(±2.7)	87.2	(±4.4)	15.8	(±2.0)	79.4	(±7.8)
Maryland – Spring 02	16.0	(±0.9)	80.2	(±4.6)	10.1	(±0.8)	60.3	(±7.0)
Massachusetts – Spring 02	13.1	(±3.0)	81.0	(±7.4)	7.0	(±1.9)	§§	
Michigan – Spring 01	21.6	(±3.3)	88.4	(±5.1)	12.6	(±2.6)	70.0	(±12.9)
Minnesota – Spring 02	16.8	(±2.2)	85.3	(±4.3)	12.2	(±1.6)	75.1	(±7.2)
Mississippi – Fall 02	24.8	(±3.0)	87.9	(±3.3)	NA ^{†††}		NA	
Nebraska – Spring 02	17.2	(±1.7)	84.7	(±5.7)	12.7	(±1.4)	70.1	(±16.8)
New Hampshire – Fall 01	17.4	(±2.8)	85.4	(±6.5)	11.1	(±2.5)	§§	
New Jersey – Fall 01	16.5	(±2.8)	82.3	(±6.0)	NA		NA	
New York – Spring 02	14.3	(±4.5)	89.3	(±8.4)	6.2	(±3.5)	73.7	(±14.7)
North Carolina – Spring 02	20.8	(±2.8)	89.1	(±4.3)	14.1	(±1.5)	71.8	(±8.0)
Ohio – Spring 02	19.5	(±3.6)	87.4	(±5.4)	13.3	(±2.2)	72.8	(±14.4)
Oklahoma – Spring 02	17.5	(±2.8)	89.3	(±4.5)	15.6	(±3.2)	70.4	(±9.0)
Pennsylvania – Spring 01	24.3	(±3.1)	91.3	(±2.6)	16.6	(±1.4)	80.6	(±8.7)
Rhode Island – Spring 01	18.5	(±3.6)	85.1	(±5.4)	11.4	(±2.0)	78.0	(±10.1)
South Dakota – Fall 01	17.7	(±3.4)	88.3	(±5.4)	14.4	(±3.1)	90.1	(±4.0)
Texas – Spring 01	25.2	(±3.6)	89.1	(±2.5)	15.7	(±1.6)	76.0	(±7.2)
Vermont – Spring 02	20.6	(±3.4)	85.9	(±6.3)	12.5	(±2.6)	72.2	(±8.3)
West Virginia – Spring 02	27.9	(±2.7)	89.9	(±1.6)	21.2	(±2.2)	77.4	(±4.0)
Wisconsin – Spring 02	17.8	(±3.0)	80.2	(±7.5)	12.0	(±2.2)	§§	
Median	18.6		86.6		12.8		73.7	
High school								
Alabama – Spring 02	38.9	(±5.0)	87.4	(±4.5)	22.3	(±3.8)	76.9	(±11.4)
Connecticut – Spring 02	27.6	(±4.5)	81.2	(±3.3)	12.3	(±2.0)	67.5	(±14.8)
Delaware – Spring 02	33.7	(±3.5)	87.5	(±3.8)	14.9	(±2.8)	84.1	(±10.7)
Florida – Spring 01	34.1	(±2.2)	93.0	(±1.9)	16.6	(±1.8)	81.1	(±6.5)
Florida – Spring 02	31.4	(±2.0)	90.2	(±1.7)	15.5	(±1.6)	85.1	(±4.7)
Georgia – Fall 01	36.9	(±3.7)	90.5	(±2.8)	21.7	(±2.7)	81.4	(±6.5)
Illinois – Spring 02	36.5	(±7.9)	91.5	(±3.8)	16.0	(±3.0)	82.1	(±9.8)
Iowa – Spring 02	37.2	(±4.5)	95.1	(±2.6)	22.2	(±3.3)	78.0	(±7.0)
Kansas – Fall 02	29.7	(±4.8)	91.1	(±3.3)	19.3	(±3.8)	84.2	(±4.9)
Kentucky – Spring 02	43.9	(±6.2)	91.2	(±2.5)	28.5	(±2.8)	84.8	(±4.5)
Maryland – Spring 02	32.0	(±1.1)	85.6	(±1.0)	15.9	(±0.6)	70.2	(±2.5)
Massachusetts – Spring 02	25.3	(±4.2)	88.0	(±4.0)	9.1	(±1.9)	70.8	(±14.5)
Michigan – Spring 01	36.6	(±2.9)	94.0	(±1.5)	18.1	(±2.2)	84.1	(±4.4)
Mississippi – Fall 02	36.0	(±2.8)	91.1	(±3.0)	NA ^{†††}		NA	
Nebraska – Spring 02	33.0	(±5.0)	90.7	(±2.3)	22.8	(±2.8)	85.6	(±5.1)
New Hampshire – Fall 01	35.5	(±3.6)	93.6	(±3.2)	20.3	(±3.9)	73.6	(±11.2)
New Jersey – Fall 01	38.1	(±4.4)	87.3	(±3.1)	NA		NA	
New York – Spring 02	28.7	(±4.3)	90.8	(±2.5)	10.4	(±3.7)	83.7	(±7.3)
North Carolina – Spring 02	39.2	(±4.0)	89.4	(±2.5)	21.9	(±2.3)	75.7	(±5.4)
Ohio – Spring 02	32.6	(±4.1)	92.4	(±3.0)	21.3	(±5.2)	90.8	(±8.2)
Oklahoma – Spring 02	35.9	(±4.6)	87.8	(±3.2)	29.2	(±3.7)	87.2	(±4.1)
Pennsylvania – Spring 01	35.9	(±2.1)	90.4	(±1.3)	19.9	(±2.4)	84.5	(±3.2)
Rhode Island – Spring 01	37.0	(±5.8)	88.8	(±5.3)	14.1	(±2.8)	71.2	(±11.2)
Texas – Spring 01	40.9	(±4.2)	92.9	(±2.0)	20.8	(±1.2)	90.6	(±4.8)
West Virginia – Spring 02	47.5	(±5.8)	92.0	(±3.5)	29.6	(±3.5)	83.1	(±9.0)
Wisconsin – Spring 02	28.2	(±5.0)	90.0	(±3.5)	18.2	(±3.9)	77.5	(±9.2)
Median	35.9		90.6		19.6		82.6	

* Students were asked, "How many your four closest friends smoke cigarettes?" and "How many of your four closest friends use chewing tobacco, snuff, or dip?"

† Students were considered to have never smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

§ Confidence interval.

‡ Smoked cigarettes on ≥1 of the previous 30 days.

** Students were considered never to have used SLT if they answered "no" to whether they have ever used SLT.

†† Use of SLT on ≥1 of the previous 30 days.

§§ Sample size <35.

††† Question not asked.

TABLE 18. Percentage of middle school and high school students who think that cigarette smokers have more friends, by cigarette smoking status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Think cigarette smokers have more friends			
	Never smoked cigarettes*		Currently smoke cigarettes†	
	%	(95% CI)§	%	(95% CI)
Middle school				
Sex				
Male	15.2	(±1.9)	44.6	(±4.3)
Female	13.3	(±2.0)	40.8	(±7.0)
Race/Ethnicity				
White, non-Hispanic	12.3	(±2.1)	37.8	(±5.0)
Black, non-Hispanic	21.1	(±2.9)	51.7	(±7.5)
Hispanic	16.4	(±2.6)	51.3	(±7.7)
Asian	13.6	(±4.4)	49.5	(±23.7)
Total	14.2	(±1.7)	42.9	(±3.9)
High school				
Sex				
Male	12.4	(±2.1)	28.8	(±3.7)
Female	9.5	(±1.3)	17.3	(±2.5)
Race/Ethnicity				
White, non-Hispanic	8.2	(±1.6)	19.4	(±2.8)
Black, non-Hispanic	17.5	(±3.3)	42.0	(±7.1)
Hispanic	15.7	(±3.0)	29.5	(±4.9)
Asian	14.7	(±5.2)	41.7	(±14.1)
Total	10.9	(±1.4)	23.4	(±2.3)

* Students were considered to have never smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

† Smoked cigarettes on ≥ 1 of the previous 30 days.

§ Confidence interval.

TABLE 19. Percentage of middle school and high school students who think that cigarette smokers have more friends, by smoking status and state — state youth tobacco surveys, United States, 2001–2002

State	Think cigarette smokers have more friends			
	Never smoked cigarettes*		Currently smoke cigarettes†	
	%	(95% CI)§	%	(95% CI)
Middle school				
Alabama – Spring 02	14.8	(±3.6)	51.1	(±9.2)
Connecticut – Spring 02	13.6	(±2.8)	38.0	(±10.9)
Delaware – Spring 02	11.3	(±1.9)	37.4	(±8.4)
Florida – Spring 01	15.7	(±2.3)	43.4	(±4.6)
Florida – Spring 02	14.6	(±1.1)	40.1	(±2.9)
Georgia – Fall 01	15.1	(±2.5)	44.9	(±6.9)
Idaho – Spring 01	5.8	(±1.7)	41.3	(±7.9)
Illinois – Spring 02	9.0	(±3.2)	36.5	(±12.8)
Iowa – Spring 02	5.8	(±1.4)	31.0	(±10.0)
Kansas – Fall 02	8.9	(±2.1)	46.0	(±12.1)
Kentucky – Spring 02	9.4	(±2.8)	44.0	(±7.4)
Louisiana – Spring 01	18.9	(±3.0)	49.1	(±5.8)
Maine – Spring 01	8.7	(±1.5)	43.2	(±3.9)
Maryland – Spring 02	17.2	(±1.2)	56.7	(±3.6)
Massachusetts – Spring 02	12.4	(±2.7)	40.6	(±7.5)
Michigan – Spring 01	9.2	(±2.1)	38.5	(±9.1)
Minnesota – Spring 02	10.3	(±1.5)	45.6	(±5.6)
Mississippi – Fall 02	22.4	(±2.1)	54.7	(±5.8)
Nebraska – Spring 02	9.3	(±1.8)	41.1	(±5.9)
New Hampshire – Fall 01	7.4	(±1.6)	32.5	(±10.5)
New Jersey – Fall 01	12.5	(±2.2)	38.4	(±9.8)
New York – Spring 02	12.1	(±2.9)	51.0	(±9.4)
North Carolina – Spring 02	14.9	(±2.3)	48.0	(±5.9)
Ohio – Spring 02	10.0	(±2.6)	45.9	(±7.6)
Oklahoma – Spring 02	11.0	(±2.9)	42.1	(±7.5)
Pennsylvania – Spring 01	13.1	(±2.9)	46.1	(±3.8)
Rhode Island – Spring 01	13.6	(±3.3)	36.5	(±7.1)
South Dakota – Fall 01	9.4	(±1.8)	48.0	(±6.6)
Texas – Spring 01	12.3	(±2.0)	51.1	(±6.2)
Vermont – Spring 02	9.7	(±2.4)	38.6	(±5.0)
West Virginia – Spring 02	11.9	(±1.9)	44.0	(±2.5)
Wisconsin – Spring 02	12.5	(±3.4)	43.7	(±12.9)
Median	12.0		43.6	
High school				
Alabama – Spring 02	19.4	(±5.4)	36.7	(±6.3)
Connecticut – Spring 02	15.6	(±2.3)	26.6	(±4.2)
Delaware – Spring 02	12.3	(±2.1)	26.1	(±4.2)
Florida – Spring 01	16.5	(±2.1)	29.0	(±3.4)
Florida – Spring 02	14.8	(±1.3)	27.7	(±2.5)
Georgia – Fall 01	16.9	(±2.1)	27.8	(±5.0)
Illinois – Spring 02	14.8	(±3.9)	25.7	(±5.2)
Iowa – Spring 02	8.1	(±2.3)	25.9	(±5.4)
Kansas – Fall 02	8.2	(±2.7)	24.8	(±4.8)
Kentucky – Spring 02	10.4	(±2.6)	32.0	(±5.9)
Maryland – Spring 02	19.8	(±0.8)	41.4	(±1.4)
Massachusetts – Spring 02	13.0	(±4.1)	29.3	(±6.1)
Michigan – Spring 01	9.4	(±2.2)	26.8	(±3.8)
Mississippi – Fall 02	22.4	(±4.1)	42.1	(±6.3)
Nebraska – Spring 02	10.4	(±2.5)	29.0	(±3.4)
New Hampshire – Fall 01	9.8	(±2.1)	25.8	(±5.5)
New Jersey – Fall 01	20.5	(±5.1)	31.2	(±4.5)
New York – Spring 02	12.1	(±4.3)	30.5	(±4.1)
North Carolina – Spring 02	17.9	(±2.8)	41.5	(±4.6)
Ohio – Spring 02	12.9	(±3.9)	23.1	(±4.7)
Oklahoma – Spring 02	10.6	(±2.5)	30.0	(±4.6)
Pennsylvania – Spring 01	13.6	(±1.7)	25.4	(±3.1)
Rhode Island – Spring 01	15.6	(±4.2)	28.2	(±6.8)
Texas – Spring 01	14.3	(±2.5)	31.5	(±6.9)
West Virginia – Spring 02	13.6	(±4.0)	27.1	(±5.2)
Wisconsin – Spring 02	12.3	(±3.4)	23.5	(±4.1)
Median	13.6		28.0	

* Students were considered to have never smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

† Smoked cigarettes on ≥ 1 of the previous 30 days.

§ Confidence interval.

TABLE 20. How current cigarette smokers* aged <18 years usually obtained cigarettes, by sex and race/ethnicity — National Youth Tobacco Survey, United States 2002

Sex and Race/Ethnicity	Bought in store		Bought from vending machine		Someone else bought them		Borrowed from someone else		Stole them†		Received from person aged >18 years		Some other way	
	%	(95% CI§)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school														
Sex														
Male	6.7	(±2.5)	2.6	(±1.4)	22.2	(±4.0)	22.4	(±5.2)	13.3	(±3.7)	8.3	(±3.0)	24.6	(±4.8)
Female	5.2	(±2.2)	2.0	(±1.5)	21.1	(±3.6)	27.4	(±3.9)	9.7	(±3.6)	9.2	(±3.1)	25.3	(±4.0)
Race/Ethnicity														
White, non-Hispanic	4.5	(±2.2)	1.5	(±1.1)	23.9	(±2.7)	26.8	(±3.9)	9.9	(±2.6)	8.1	(±2.9)	25.3	(±3.7)
Black, non-Hispanic	8.2	(±3.9)	2.3	(±1.9)	20.9	(±7.3)	20.2	(±7.0)	18.9	(±9.4)	9.8	(±5.4)	19.8	(±6.6)
Hispanic	8.5	(±3.1)	4.2	(±4.3)	14.7	(±4.7)	21.0	(±6.1)	15.2	(±5.7)	9.2	(±4.8)	27.3	(±6.0)
Asian	¶		¶		¶		¶		¶		¶		¶	
Total	5.9	(±1.7)	2.3	(±1.1)	21.7	(±2.2)	24.9	(±3.2)	11.5	(±2.5)	8.7	(±2.4)	25.0	(±2.7)
High school														
Sex														
Male	31.2	(±4.5)	1.8	(±1.2)	24.6	(±4.8)	19.9	(±2.7)	4.3	(±1.3)	8.2	(±1.9)	10.0	(±2.0)
Female	18.1	(±3.7)	1.8	(±1.0)	32.8	(±4.1)	21.5	(±3.6)	2.2	(±1.0)	14.1	(±2.7)	9.6	(±1.9)
Race/Ethnicity														
White, non-Hispanic	25.2	(±4.4)	1.9	(±1.1)	30.4	(±4.2)	20.8	(±2.6)	2.5	(±1.0)	10.7	(±2.2)	8.5	(±1.7)
Black, non-Hispanic	27.7	(±6.5)	2.2	(±2.7)	20.4	(±6.8)	17.9	(±6.7)	6.6	(±3.9)	12.4	(±4.7)	12.7	(±4.7)
Hispanic	21.5	(±5.8)	1.1	(±1.1)	22.1	(±4.7)	23.7	(±4.8)	3.8	(±2.0)	13.2	(±4.0)	14.6	(±4.7)
Asian	24.4	(±20.4)	0.5	(±1.0)	27.2	(±15.1)	9.1	(±6.9)	5.7	(±7.0)	15.4	(±13.5)	17.7	(±14.9)
Total	24.7	(±3.7)	1.8	(±0.8)	28.7	(±3.3)	20.6	(±2.0)	3.3	(±0.8)	11.1	(±1.9)	9.8	(±1.5)

* Smoked cigarettes on ≥1 of the previous 30 days.

† Not a response option on the state survey.

§ Confidence interval.

¶ Sample size <35.

TABLE 21. How current cigarette smokers* aged <18 years usually obtained their cigarettes, by state — state youth tobacco surveys, 2001–2002

Sex and Race/Ethnicity	Bought in store		Bought from vending machine		Someone else bought them		Borrowed from someone else		Stole from store or family member [§]		Received from person aged >18 years		Some other way	
	%	(95% CI) [†]	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school														
Alabama – Spring 02	6.8	(±3.2)	3.4	(±3.2)	31.3	(±10.6)	20.3	(±6.2)	13.2	(±3.1)	7.4	(±3.8)	17.6	(±6.1)
Connecticut – Spring 02	13.3	(±8.4)	2.1	(±2.1)	18.6	(±9.5)	24.0	(±10.0)	19.1	(±7.2)	11.7	(±5.6)	11.2	(±5.7)
Delaware – Spring 02	6.1	(±4.2)	1.4	(±2.1)	23.6	(±5.1)	25.9	(±7.5)	15.1	(±6.6)	9.1	(±4.9)	18.8	(±6.4)
Florida – Spring 01	4.1	(±2.9)	2.3	(±1.1)	21.6	(±4.1)	21.1	(±3.7)	14.8	(±3.4)	10.6	(±2.9)	21.0	(±4.2)
Florida – Spring 02	4.9	(±1.4)	1.3	(±0.7)	17.9	(±2.6)	23.6	(±2.6)	14.0	(±2.1)	13.9	(±2.1)	19.9	(±2.5)
Georgia – Fall 01	7.8	(±5.4)	0.8	(±1.3)	17.5	(±6.9)	25.2	(±5.5)	15.5	(±4.3)	12.0	(±5.7)	21.1	(±7.2)
Idaho – Spring 01	8.6	(±3.3)	1.6	(±1.8)	28.0	(±7.4)	27.5	(±6.9)	11.3	(±5.8)	4.1	(±3.0)	19.0	(±5.1)
Illinois – Spring 02	12.4	(±6.8)	3.3	(±2.7)	23.1	(±7.6)	24.7	(±10.6)	10.9	(±4.6)	4.4	(±4.1)	21.2	(±4.3)
Iowa – Spring 02	2.0	(±1.0)	0.6	(±1.2)	26.4	(±8.7)	34.3	(±7.3)	9.7	(±5.6)	8.9	(±7.8)	18.1	(±3.2)
Kansas – Fall 02	3.9	(±4.5)	2.4	(±3.1)	17.3	(±8.4)	31.4	(±11.7)	10.7	(±6.9)	11.0	(±7.3)	23.3	(±8.8)
Kentucky – Spring 02	7.0	(±2.0)	3.7	(±4.4)	24.1	(±5.2)	26.6	(±5.0)	8.3	(±2.9)	13.4	(±4.9)	16.8	(±5.2)
Louisiana – Spring 01	7.2	(±2.0)	3.6	(±2.3)	30.7	(±4.2)	20.8	(±4.4)	14.8	(±4.0)	8.9	(±2.7)	13.9	(±4.5)
Maine – Spring 01	3.4	(±1.6)	2.0	(±1.2)	17.4	(±4.4)	9.3	(±2.1)	25.9	(±5.3)	7.4	(±2.7)	9.8	(±4.3)
Maryland – Spring 02	8.3	(±2.2)	5.5	(±2.0)	21.3	(±3.1)	21.6	(±3.3)	15.2	(±3.6)	10.4	(±2.3)	17.7	(±2.9)
Massachusetts – Spring 02	5.5	(±4.6)	3.1	(±4.0)	19.9	(±9.3)	23.3	(±8.8)	12.8	(±6.1)	4.6	(±4.4)	30.8	(±10.6)
Michigan – Spring 01	12.6	(±8.2)	0.0	(±0.0)	NA [¶]		29.1	(±7.4)	16.3	(±5.0)	23.5	(±6.9)	18.5	(±5.7)
Minnesota – Spring 02	4.9	(±2.9)	0.9	(±1.3)	22.3	(±7.7)	28.2	(±7.4)	17.9	(±5.1)	7.5	(±3.1)	18.4	(±5.1)
Mississippi – Fall 02	8.7	(±4.6)	3.5	(±2.1)	20.4	(±4.7)	32.7	(±6.7)	15.4	(±4.8)	19.2	(±4.6)	NA	
Nebraska – Spring 02	2.9	(±3.4)	0.9	(±1.3)	19.1	(±5.7)	36.5	(±9.0)	18.8	(±5.1)	7.4	(±3.3)	14.4	(±5.9)
New Hampshire – Fall 01	7.7	(±6.2)	7.1	(±5.6)	24.1	(±12.0)	22.9	(±10.9)	19.5	(±8.9)	9.4	(±9.3)	9.3	(±5.5)
New Jersey – Fall 01	11.2	(±5.2)	7.5	(±5.4)	13.1	(±5.3)	29.1	(±5.7)	10.5	(±3.8)	11.9	(±5.4)	16.5	(±4.7)
New York – Spring 02	16.2	(±13.7)	2.2	(±3.7)	24.5	(±9.4)	26.7	(±5.3)	10.3	(±4.9)	4.6	(±4.5)	15.4	(±6.5)
North Carolina – Spring 02	4.2	(±2.8)	2.4	(±1.7)	24.1	(±5.1)	24.6	(±4.2)	12.1	(±4.7)	11.9	(±3.0)	20.6	(±4.9)
Ohio – Spring 02	5.8	(±3.9)	1.4	(±1.9)	28.8	(±6.9)	27.8	(±8.0)	15.3	(±7.3)	6.9	(±4.7)	14.0	(±8.4)
Oklahoma – Spring 02	5.2	(±3.5)	2.7	(±2.8)	30.5	(±5.4)	28.7	(±8.1)	11.1	(±4.5)	9.4	(±4.9)	12.4	(±3.8)
Pennsylvania – Spring 01	11.4	(±3.5)	3.7	(±1.8)	23.7	(±3.3)	28.5	(±4.3)	9.3	(±2.4)	7.6	(±3.4)	15.9	(±5.0)
Rhode Island – Spring 01	10.5	(±4.0)	4.9	(±3.5)	14.7	(±6.9)	31.1	(±5.6)	15.6	(±4.7)	6.6	(±4.1)	16.6	(±6.9)
South Dakota – Fall 01	2.1	(±1.8)	1.6	(±1.9)	32.2	(±4.8)	29.3	(±6.4)	9.7	(±3.6)	6.0	(±3.3)	19.1	(±3.9)
Texas – Spring 01	7.6	(±4.5)	3.0	(±1.6)	20.4	(±3.9)	25.3	(±5.1)	10.6	(±2.7)	10.2	(±3.7)	22.9	(±5.3)
Vermont – Spring 02	2.4	(±1.2)	1.4	(±1.1)	27.9	(±6.8)	26.6	(±4.6)	11.5	(±5.2)	7.6	(±2.9)	22.5	(±5.7)
West Virginia – Spring 02	4.8	(±1.1)	3.2	(±0.9)	23.9	(±3.6)	22.7	(±2.2)	14.1	(±1.8)	9.6	(±1.9)	21.8	(±2.4)
Wisconsin – Spring 02	7.3	(±5.9)	0.5	(±0.9)	31.3	(±11.2)	27.7	(±11.3)	12.6	(±7.9)	4.2	(±3.7)	16.5	(±7.6)
Median	6.9		2.4		23.6		26.6		13.6		9.0		18.1	
High School														
Alabama – Spring 02	21.3	(±6.5)	1.1	(±1.6)	35.0	(±6.3)	19.3	(±5.2)	4.4	(±2.7)	7.8	(±3.3)	11.0	(±4.5)
Connecticut – Spring 02	30.6	(±4.6)	2.6	(±1.9)	19.8	(±4.2)	25.7	(±4.3)	4.0	(±2.0)	9.2	(±2.9)	8.1	(±3.7)
Delaware – Spring 02	26.8	(±3.9)	3.1	(±2.6)	29.3	(±7.0)	18.3	(±3.6)	3.5	(±2.3)	7.0	(±1.7)	11.9	(±6.6)
Florida – Spring 01	18.5	(±3.3)	1.3	(±1.0)	23.9	(±3.6)	25.7	(±4.5)	5.5	(±2.2)	15.9	(±3.2)	6.5	(±2.3)
Florida – Spring 02	17.7	(±2.7)	1.2	(±0.7)	25.0	(±2.6)	28.0	(±2.3)	3.7	(±1.3)	12.7	(±1.8)	9.7	(±1.6)
Georgia – Fall 01	21.6	(±4.6)	1.8	(±1.2)	28.3	(±3.7)	23.8	(±2.3)	3.8	(±1.8)	11.2	(±3.1)	9.5	(±2.2)
Illinois – Spring 02	24.1	(±4.4)	2.2	(±2.7)	32.2	(±5.8)	23.1	(±4.8)	2.8	(±2.1)	9.6	(±4.2)	6.1	(±3.0)
Iowa – Spring 02	9.2	(±2.7)	1.7	(±1.6)	43.2	(±7.5)	26.2	(±6.2)	2.3	(±1.9)	8.6	(±3.1)	8.8	(±4.0)
Kansas – Fall 02	17.3	(±5.6)	1.6	(±1.9)	32.0	(±5.1)	30.0	(±5.2)	3.1	(±1.9)	6.9	(±2.8)	9.2	(±3.4)
Kentucky – Spring 02	19.9	(±5.0)	2.5	(±1.7)	31.9	(±4.7)	23.5	(±5.4)	4.1	(±2.4)	9.7	(±3.0)	8.3	(±3.0)
Maryland – Spring 02	24.0	(±1.6)	3.4	(±0.6)	26.8	(±1.3)	22.9	(±1.3)	5.2	(±0.6)	8.7	(±0.8)	9.1	(±0.8)
Massachusetts – Spring 02	20.0	(±7.0)	0.3	(±0.7)	29.3	(±7.4)	30.5	(±7.7)	4.7	(±2.6)	7.0	(±3.5)	8.1	(±3.4)
Michigan – Spring 01	21.5	(±5.9)	1.5	(±1.0)	NA [¶]		28.8	(±3.2)	4.9	(±1.6)	34.2	(±5.5)	9.1	(±1.7)
Mississippi – Fall 02	19.4	(±4.3)	3.0	(±2.6)	32.5	(±4.3)	30.3	(±4.4)	5.1	(±2.1)	9.6	(±3.6)	NA	
Nebraska – Spring 02	6.9	(±1.6)	1.2	(±1.0)	41.4	(±4.5)	29.5	(±4.7)	3.2	(±1.5)	9.6	(±2.6)	8.1	(±2.5)
New Hampshire – Fall 01	14.8	(±5.7)	2.5	(±2.1)	36.2	(±6.9)	26.8	(±5.4)	8.6	(±4.1)	6.4	(±3.1)	4.7	(±2.4)
New Jersey – Fall 01	33.7	(±6.9)	1.8	(±1.5)	24.2	(±5.4)	25.7	(±1.6)	3.7	(±2.0)	4.8	(±1.6)	6.0	(±2.6)
New York – Spring 02	22.7	(±5.9)	1.5	(±1.2)	27.3	(±9.2)	20.8	(±3.9)	5.5	(±5.1)	13.4	(±3.9)	8.9	(±1.8)
North Carolina – Spring 02	19.7	(±2.8)	1.8	(±0.7)	33.3	(±3.7)	22.9	(±3.6)	5.4	(±1.3)	8.5	(±2.5)	8.4	(±1.4)
Ohio – Spring 02	20.5	(±5.1)	1.3	(±1.4)	36.1	(±5.2)	22.8	(±5.4)	1.7	(±1.7)	8.6	(±4.1)	9.0	(±2.8)
Oklahoma – Spring 02	22.1	(±5.8)	1.5	(±1.4)	32.8	(±5.5)	21.1	(±4.3)	3.6	(±2.1)	10.2	(±3.7)	8.7	(±3.4)
Pennsylvania – Spring 01	29.0	(±2.7)	2.8	(±0.9)	26.5	(±3.1)	24.5	(±2.3)	2.8	(±1.0)	7.0	(±1.8)	7.4	(±2.1)
Rhode Island – Spring 01	32.0	(±6.7)	0.7	(±1.1)	26.9	(±5.2)	26.8	(±4.6)	4.1	(±2.9)	3.2	(±1.3)	6.2	(±2.5)
Texas – Spring 01	19.7	(±4.9)	1.5	(±1.0)	26.6	(±3.8)	25.9	(±3.0)	2.7	(±1.8)	10.5	(±3.4)	13.1	(±1.9)
West Virginia – Spring 02	19.0	(±6.9)	1.7	(±1.1)	36.4	(±4.6)	19.8	(±4.5)	4.1	(±2.2)	10.4	(±3.3)	8.7	(±2.9)
Wisconsin – Spring 02	21.7	(±5.4)	0.0	(±0.0)	39.1	(±5.0)	24.1	(±5.9)	3.8	(±2.0)	5.5	(±2.8)	5.8	(±2.6)
Median	20.9		1.7		31.9		25.1		3.9		9.0		8.7	

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Not a response option on the National Youth Tobacco Survey.

¶ Response option not given.

TABLE 22. Where current cigarette smokers* aged <18 years bought their last pack of cigarettes, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Gas station	Convenience store	Discount store	Grocery store	Drug store	Vending machine	Restaurant
	% (95% CI)†	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school							
Sex							
Male	43.3 (±7.1)	18.4 (±6.0)	7.1 (±3.6)	6.2 (±2.8)	6.6 (±3.9)	14.8 (±5.3)	3.5 (±2.6)
Female	46.1 (±9.7)	27.6 (±7.8)	3.1 (±2.4)	8.2 (±4.3)	4.6 (±2.9)	8.4 (±4.0)	2.1 (±1.9)
Race/Ethnicity							
White, non-Hispanic	48.1 (±8.1)	24.6 (±6.0)	6.3 (±3.5)	4.0 (±2.4)	4.7 (±2.7)	10.3 (±3.9)	2.0 (±2.1)
Black, non-Hispanic	33.3 (±11.2)	22.5 (±10.0)	1.6 (±2.4)	16.2 (±11.7)	5.1 (±3.6)	12.0 (±11.0)	9.3 (±8.1)
Hispanic	43.6 (±11.2)	22.5 (±11.0)	2.0 (±1.9)	11.1 (±5.4)	5.1 (±3.3)	15.0 (±7.6)	0.7 (±0.9)
Asian	§	§	§	§	§	§	§
Total	44.7 (±6.4)	23.0 (±4.7)	5.1 (±2.3)	7.2 (±2.5)	5.7 (±2.4)	11.6 (±3.3)	2.8 (±1.8)
High school							
Sex							
Male	56.6 (±6.3)	26.2 (±5.7)	3.2 (±1.5)	5.7 (±2.0)	3.5 (±1.3)	4.1 (±2.2)	0.6 (±0.6)
Female	57.5 (±5.6)	29.5 (±5.6)	2.0 (±1.4)	5.0 (±2.8)	2.1 (±1.4)	2.7 (±1.3)	1.1 (±1.0)
Race/Ethnicity							
White, non-Hispanic	61.1 (±6.0)	26.6 (±4.7)	2.0 (±1.1)	4.5 (±1.9)	2.1 (±1.0)	3.6 (±1.8)	0.1 (±0.2)
Black, non-Hispanic	37.3 (±10.8)	35.2 (±9.9)	7.4 (±5.6)	6.1 (±4.6)	8.4 (±5.9)	4.2 (±4.7)	1.4 (±2.2)
Hispanic	46.6 (±8.5)	29.5 (±7.3)	3.8 (±2.2)	9.8 (±4.5)	4.0 (±2.5)	2.4 (±2.0)	3.8 (±3.4)
Asian	§	§	§	§	§	§	§
Total	57.0 (±5.1)	27.8 (±4.1)	2.6 (±1.1)	5.4 (±1.9)	2.8 (±0.9)	3.5 (±1.4)	0.9 (±0.6)

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Sample size <35.

TABLE 23. Where current cigarette smokers* aged <18 years bought their last pack of cigarettes, by state — state youth tobacco surveys, United States, 2001–2002

State	Gas station	Convenience store	Grocery store	Drug store	Vending machine	Internet	Other
	% (95% CI)†	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school							
Alabama – Spring 02	25.9 (±8.7)	11.1 (±5.5)	2.8 (±2.7)	2.9 (±2.9)	2.2 (±2.4)	4.0 (±3.5)	51.2 (±13.3)
Connecticut – Spring 02	18.2 (±8.8)	18.3 (±8.2)	2.1 (±3.5)	3.8 (±3.7)	3.7 (±4.3)	3.5 (±4.3)	50.3 (±14.7)
Delaware – Spring 02	21.4 (±7.4)	13.6 (±9.4)	2.2 (±2.5)	4.5 (±5.4)	1.8 (±1.8)	1.5 (±2.3)	55.2 (±11.6)
Florida – Spring 01	NA§	NA	NA	NA	NA	NA	NA
Florida – Spring 02	NA	NA	NA	NA	NA	NA	NA
Georgia – Fall 01	18.3 (±5.7)	12.5 (±5.2)	3.3 (±3.4)	0.9 (±1.8)	6.2 (±5.0)	0.5 (±0.9)	58.3 (±7.2)
Idaho – Spring 01	26.7 (±6.6)	7.7 (±5.5)	8.0 (±4.6)	1.9 (±2.6)	2.9 (±3.5)	1.8 (±2.4)	51.0 (±9.3)
Illinois – Spring 02	31.8 (±11.6)	11.4 (±6.6)	7.2 (±5.7)	3.9 (±4.7)	2.8 (±2.3)	0.0 (±0.0)	42.8 (±12.5)
Iowa – Spring 02	22.7 (±7.0)	3.2 (±4.6)	5.0 (±2.7)	0.0 (±0.0)	0.9 (±1.7)	0.0 (±0.0)	68.2 (±9.5)
Kansas – Fall 02	17.8 (±2.0)	12.1 (±5.9)	5.7 (±5.9)	0.0 (±0.0)	1.2 (±2.7)	0.0 (±0.0)	63.1 (±12.4)
Kentucky – Spring 02	24.0 (±6.9)	9.1 (±3.5)	6.7 (±4.8)	3.0 (±3.0)	3.2 (±2.8)	0.5 (±1.0)	53.5 (±10.4)
Louisiana – Spring 01	22.3 (±5.8)	13.0 (±3.0)	5.1 (±2.4)	6.2 (±4.2)	3.6 (±2.1)	3.1 (±2.3)	46.8 (±6.5)
Maine – Spring 01	30.2 (±11.1)	27.4 (±6.4)	10.4 (±4.3)	6.8 (±3.7)	15.6 (±9.0)	NA	9.7 (±3.5)††
Maryland – Spring 02	23.0 (±3.1)	12.0 (±3.4)	4.6 (±1.6)	4.5 (±1.7)	6.9 (±2.8)	3.4 (±2.2)	45.6 (±4.1)
Massachusetts – Spring 02	9.7 (±7.8)	17.7 (±10.4)	3.9 (±4.1)	7.9 (±7.3)	8.3 (±6.2)	1.7 (±3.5)	50.7 (±15.0)
Michigan – Spring 01	22.8 (±6.2)	9.5 (±5.6)	2.0 (±3.8)	2.6 (±3.7)	1.4 (±2.5)	4.7 (±4.7)	56.9 (±9.8)
Minnesota – Spring 02	17.0 (±6.1)	8.7 (±4.3)	4.7 (±6.0)	1.4 (±2.3)	3.3 (±3.3)	0.0 (±0.0)	64.9 (±8.3)
Mississippi – Fall 02	NA	NA	NA	NA	NA	NA	NA
Nebraska – Spring 02	19.4 (±5.8)	3.6 (±4.0)	4.0 (±4.0)	4.8 (±3.8)	4.2 (±4.2)	0.0 (±0.0)	64.0 (±6.4)
New Hampshire – Fall 01	19.3 (±9.7)	20.3 (±10.2)	3.8 (±5.1)	0.0 (±0.0)	10.9 (±5.6)	2.9 (±4.0)	42.8 (±10.2)
New Jersey – Fall 01	16.5 (±9.7)	13.0 (±9.5)	4.0 (±1.9)	3.6 (±3.0)	10.0 (±6.6)	3.9 (±5.5)	49.0 (±6.1)
New York – Spring 02	38.6 (±19.8)	31.4 (±23.3)	8.3 (±5.3)	2.3 (±2.7)	7.9 (±9.0)	NA	11.4 (±13.1)††
North Carolina – Spring 02	17.6 (±6.3)	17.7 (±5.8)	4.8 (±2.6)	1.3 (±1.0)	3.6 (±2.3)	0.8 (±1.0)	54.3 (±9.4)
Ohio – Spring 02	22.8 (±7.7)	15.0 (±9.5)	3.1 (±3.6)	4.4 (±3.6)	3.4 (±3.2)	1.8 (±2.6)	49.4 (±13.6)
Oklahoma – Spring 02	22.2 (±8.9)	18.9 (±6.0)	4.2 (±3.6)	1.1 (±2.2)	3.9 (±2.6)	1.6 (±2.1)	48.2 (±7.9)
Pennsylvania – Spring 01	18.2 (±3.1)	18.3 (±4.5)	8.9 (±3.3)	2.1 (±1.4)	5.5 (±2.0)	1.1 (±1.3)	45.9 (±4.2)
Rhode Island – Spring 01	13.2 (±6.5)	21.8 (±8.8)	3.5 (±3.2)	2.9 (±3.3)	9.3 (±7.6)	2.4 (±4.8)	46.8 (±11.8)
South Dakota – Fall 01	20.8 (±5.7)	10.4 (±4.8)	6.2 (±4.4)	2.7 (±2.2)	4.5 (±2.7)	0.5 (±1.0)	54.9 (±8.2)
Texas – Spring 01	38.0 (±6.9)	27.9 (±5.9)	5.3 (±2.0)	12.3 (±6.4)	9.5 (±18.5)	4.1 (±2.9)	2.9 (±2.0)**
Vermont – Spring 02	19.4 (±6.3)	16.7 (±2.8)	4.2 (±1.9)	2.6 (±2.7)	3.9 (±3.3)	1.7 (±1.2)	51.5 (±7.7)
West Virginia – Spring 02	23.3 (±3.1)	12.2 (±2.1)	5.6 (±1.5)	2.0 (±1.0)	3.4 (±1.3)	0.8 (±0.7)	52.7 (±3.7)
Wisconsin – Spring 02	37.8 (±15.6)	13.4 (±9.2)	2.4 (±2.4)	0.8 (±1.5)	5.2 (±7.9)	0.0 (±0.0)	40.5 (±11.4)
Median	22.2	13.0	4.6	2.7	3.9	1.6	50.7
High school							
Alabama	47.1 (±8.8)	16.4 (±4.7)	2.4 (±2.5)	1.8 (±2.2)	0.0 (±0.0)	1.5 (±1.5)	30.8 (±8.4)
Connecticut – Spring 02	45.9 (±6.9)	18.9 (±6.3)	3.7 (±2.3)	3.2 (±2.0)	3.2 (±2.4)	3.1 (±1.5)	21.9 (±4.6)
Delaware – Spring 02	41.9 (±5.2)	28.6 (±4.6)	3.9 (±3.0)	2.7 (±2.1)	2.0 (±1.5)	0.9 (±0.9)	20.1 (±3.6)
Florida – Spring 01	NA§	NA	NA	NA	NA	NA	NA
Florida – Spring 02	NA	NA	NA	NA	NA	NA	NA
Georgia – Fall 01	47.1 (±6.8)	18.5 (±3.6)	6.3 (±2.6)	0.9 (±1.1)	2.2 (±1.2)	1.5 (±1.9)	23.5 (±5.7)
Illinois – Spring 02	63.2 (±8.6)	19.1 (±5.6)	1.0 (±1.3)	1.6 (±2.0)	0.9 (±1.0)	1.1 (±1.3)	13.1 (±4.5)
Iowa – Spring 02	51.1 (±11.0)	16.3 (±6.1)	8.6 (±6.1)	2.6 (±3.1)	0.6 (±0.9)	1.3 (±1.5)	19.5 (±5.0)
Kansas – Fall 02	52.4 (±8.4)	16.1 (±7.1)	4.4 (±2.4)	1.7 (±1.9)	2.2 (±2.1)	0.0 (±0.0)	23.2 (±5.6)
Kentucky – Spring 02	43.3 (±7.1)	19.0 (±5.0)	6.2 (±2.9)	2.3 (±2.0)	2.7 (±2.0)	1.7 (±1.4)	24.8 (±5.6)
Maryland – Spring 02	41.4 (±1.7)	19.7 (±1.3)	5.7 (±0.8)	3.9 (±0.7)	3.2 (±0.6)	3.0 (±0.5)	23.1 (±1.6)
Massachusetts – Spring 02	25.7 (±9.5)	46.3 (±11.1)	2.1 (±2.8)	6.5 (±4.8)	0.2 (±0.3)	2.5 (±2.0)	16.6 (±6.1)
Michigan – Spring 01	45.8 (±4.5)	18.3 (±4.1)	6.0 (±2.0)	4.1 (±2.1)	2.8 (±1.5)	2.0 (±1.2)	21.0 (±3.7)
Mississippi – Fall 02	NA	NA	NA	NA	NA	NA	NA
Nebraska – Spring 02	48.3 (±6.9)	14.3 (±4.1)	8.8 (±2.9)	1.4 (±1.3)	1.8 (±1.4)	1.0 (±1.0)	24.5 (±6.1)
New Hampshire – Fall 01	28.4 (±7.0)	39.6 (±8.6)	5.3 (±3.6)	0.8 (±1.2)	2.2 (±2.1)	0.5 (±0.9)	23.1 (±4.9)
New Jersey – Fall 01	20.2 (±5.0)	52.7 (±6.3)	3.5 (±3.3)	1.5 (±1.2)	2.7 (±1.9)	2.7 (±2.4)	16.6 (±3.3)
New York – Spring 02	44.4 (±11.2)	28.1 (±5.7)	10.3 (±5.3)	6.4 (±4.5)	4.0 (±3.3)	NA	6.8 (±2.9)††
North Carolina – Spring 02	39.5 (±4.2)	24.0 (±3.6)	7.1 (±1.7)	2.0 (±0.9)	1.7 (±1.1)	2.5 (±1.5)	23.2 (±3.5)
Ohio – Spring 02	40.1 (±6.6)	26.6 (±5.8)	6.3 (±4.4)	3.2 (±2.5)	1.4 (±1.6)	0.7 (±1.0)	21.6 (±7.2)
Oklahoma – Spring 02	31.2 (±5.8)	34.5 (±7.7)	2.7 (±2.9)	2.4 (±2.2)	3.7 (±2.5)	0.8 (±1.0)	24.8 (±5.7)
Pennsylvania – Spring 01	39.2 (±6.4)	28.6 (±5.1)	6.8 (±1.9)	3.4 (±1.4)	3.3 (±1.6)	1.0 (±0.8)	17.7 (±3.2)
Rhode Island – Spring 01	26.7 (±7.6)	49.2 (±6.5)	2.7 (±1.3)	3.5 (±2.3)	1.1 (±0.8)	2.7 (±2.2)	14.1 (±7.0)
Texas – Spring 01	45.1 (±4.2)	36.6 (±5.0)	7.6 (±2.8)	2.3 (±1.2)	4.4 (±3.0)	1.9 (±0.9)	2.1 (±1.4)**
West Virginia – Spring 02	45.1 (±6.9)	22.5 (±6.0)	6.6 (±3.6)	1.4 (±1.1)	0.7 (±0.8)	0.2 (±0.4)	23.4 (±5.3)
Wisconsin – Spring 02	61.9 (±6.3)	13.5 (±4.6)	3.7 (±2.5)	2.2 (±2.2)	1.1 (±1.6)	1.3 (±1.8)	16.4 (±4.6)
Median	44.4	22.5	5.7	2.3	2.2	1.4	21.6

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Question not asked.

†† Other category included discount store and restaurant.

** Included only restaurants.

TABLE 24. Percentage of current cigarette smokers* aged <18 years who bought or tried to buy cigarettes in a store and were not asked to show proof of age or who were not refused purchase because of their age, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Were not asked to show proof of age when purchasing cigarettes		Were not refused purchase because of age	
	%	(95% CI†)	%	(95% CI)
Middle school				
Sex				
Male	72.7	(±5.9)	59.0	(±7.4)
Female	79.9	(±6.0)	68.8	(±11.5)
Race/Ethnicity				
White, non-Hispanic	78.4	(±7.0)	63.4	(±9.2)
Black, non-Hispanic	73.3	(±12.6)	59.9	(±13.5)
Hispanic	71.5	(±9.1)	63.7	(±10.3)
Asian	§		§	
Total	75.9	(±4.5)	63.4	(±6.4)
High school				
Sex				
Male	58.3	(±6.1)	57.7	(±5.9)
Female	58.7	(±7.4)	64.8	(±5.5)
Race/Ethnicity				
White, non-Hispanic	57.0	(±5.4)	61.7	(±5.5)
Black, non-Hispanic	69.3	(±10.8)	57.0	(±12.9)
Hispanic	61.9	(±8.5)	60.7	(±8.8)
Asian	§		§	
Total	58.5	(±4.9)	60.6	(±4.4)

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Sample size <35.

TABLE 25. Percentage of current cigarette smokers* aged <18 years who bought or tried to buy cigarettes in a store or were not asked to show proof of age and who were not refused purchase because of their age, by state — state youth tobacco surveys, United States, 2001–2002

State	Were not asked to show proof of age		Were not refused purchase because of age	
	%	(95% CI†)	%	(95% CI)
Middle school				
Alabama – Spring 02	70.8	(±13.1)	60.4	(±9.1)
Connecticut – Spring 02	78.9	(±16.8)	66.1	(±17.4)
Delaware – Spring 02	57.8	(±12.0)	64.8	(±8.6)
Florida – Spring 01	65.7	(±12.5)	69.6	(±10.5)
Florida – Spring 02	64.4	(±6.7)	54.3	(±7.4)
Georgia – Fall 01	92.6	(±8.3)	77.0	(±9.5)
Idaho – Spring 01	83.0	(±12.1)	77.2	(±14.1)
Illinois – Spring 02	73.1	(±15.9)	68.7	(±18.4)
Iowa – Spring 02	78.7	(±22.4)	60.0	(±9.5)
Kansas – Fall 02	§		§	
Kentucky – Spring 02	74.5	(±12.7)	68.9	(±9.0)
Louisiana – Spring 01	70.5	(±7.8)	67.7	(±5.1)
Maine – Spring 01	72.6	(±9.1)	74.0	(±8.9)
Maryland – Spring 02	66.0	(±5.7)	63.8	(±5.3)
Massachusetts – Spring 02	76.2	(±12.9)	78.1	(±12.9)
Michigan – Spring 01	74.2	(±14.9)	77.4	(±14.4)
Minnesota – Spring 02	78.4	(±10.7)	78.2	(±6.9)
Mississippi – Fall 02	NA¶		NA	
Nebraska – Spring 02	§		75.9	(±10.2)
New Hampshire – Fall 01	§		§	
New Jersey – Fall 01	67.8	(±8.5)	58.1	(±8.2)
New York – Spring 02	82.0	(±13.9)	72.2	(±13.1)
North Carolina – Spring 02	69.7	(±9.8)	65.3	(±6.5)
Ohio – Spring 02	79.1	(±15.4)	61.0	(±11.4)
Oklahoma – Spring 02	78.2	(±15.5)	71.9	(±10.0)
Pennsylvania – Spring 01	77.7	(±5.8)	59.2	(±5.8)
Rhode Island – Spring 01	87.5	(±11.5)	71.5	(±10.8)
South Dakota – Fall 01	76.9	(±7.4)	75.0	(±9.7)
Texas – Spring 01	71.4	(±7.4)	71.2	(±6.5)
Vermont – Spring 02	72.2	(±8.8)	74.5	(±9.9)
West Virginia – Spring 02	70.0	(±4.3)	69.1	(±3.9)
Wisconsin – Spring 02	§		§	
Median	74.2		69.1	
High school				
Alabama – Spring 02	62.4	(±12.2)	61.6	(±9.9)
Connecticut – Spring 02	54.5	(±9.0)	55.2	(±7.8)
Delaware – Spring 02	62.6	(±5.0)	57.9	(±6.2)
Florida – Spring 01	62.5	(±7.1)	61.9	(±6.9)
Florida – Spring 02	54.6	(±5.1)	59.3	(±4.9)
Georgia – Fall 01	66.7	(±8.4)	67.5	(±7.3)
Illinois – Spring 02	54.7	(±8.7)	49.7	(±9.7)
Iowa – Spring 02	72.1	(±8.9)	63.5	(±7.1)
Kansas – Fall 02	59.0	(±10.3)	62.1	(±9.2)
Kentucky – Spring 02	61.9	(±6.2)	55.9	(±6.2)
Maryland – Spring 02	61.2	(±1.9)	62.2	(±2.0)
Massachusetts – Spring 02	61.7	(±11.4)	57.8	(±10.4)
Michigan – Spring 01	64.9	(±5.1)	64.2	(±6.3)
Mississippi – Fall 02	NA¶		NA	
Nebraska – Spring 02	62.4	(±9.6)	59.7	(±8.7)
New Hampshire – Fall 01	64.2	(±7.4)	58.6	(±12.1)
New Jersey – Fall 01	66.3	(±6.6)	65.4	(±4.9)
New York – Spring 02	54.3	(±10.2)	59.7	(±11.6)
North Carolina – Spring 02	62.2	(±3.7)	64.8	(±5.1)
Ohio – Spring 02	60.4	(±8.5)	55.9	(±11.3)
Oklahoma – Spring 02	64.2	(±7.8)	57.1	(±10.2)
Pennsylvania – Spring 01	64.4	(±4.0)	60.5	(±5.7)
Rhode Island – Spring 01	66.0	(±8.7)	60.9	(±6.9)
Texas – Spring 01	62.3	(±6.7)	63.4	(±7.1)
West Virginia – Spring 02	60.7	(±9.3)	63.2	(±7.5)
Wisconsin – Spring 02	55.7	(±10.6)	68.0	(±10.4)
Median	62.3		60.9	

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Sample size <35.

¶ Question not asked.

TABLE 26. Percentage of middle school and high school students who were exposed to tobacco-related media and advertising, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Saw or heard antismoking commercials on TV or radio		Saw actors on TV or in movies using tobacco		Saw ads for tobacco products on the Internet	
	%	(95% CI)*	%	(95% CI)	%	(95% CI)
Middle school						
Sex						
Male	81.9	(±1.6)	89.7	(±0.9)	42.8	(±2.1)
Female	87.4	(±1.5)	90.0	(±1.0)	42.5	(±2.2)
Race/Ethnicity						
White, non-Hispanic	85.4	(±1.5)	90.3	(±1.0)	42.1	(±2.1)
Black, non-Hispanic	81.8	(±2.5)	90.3	(±1.5)	41.3	(±2.6)
Hispanic	85.3	(±2.0)	89.6	(±1.7)	44.9	(±2.0)
Asian	86.7	(±4.3)	84.0	(±5.2)	39.8	(±8.4)
Total	84.6	(±1.2)	89.9	(±0.8)	42.7	(±1.6)
High school						
Sex						
Male	90.1	(±1.3)	90.2	(±0.9)	32.8	(±2.3)
Female	92.3	(±1.2)	92.4	(±0.8)	34.2	(±1.5)
Race/Ethnicity						
White, non-Hispanic	92.2	(±1.3)	91.4	(±0.7)	31.7	(±1.9)
Black, non-Hispanic	88.7	(±1.7)	91.5	(±1.6)	36.9	(±3.2)
Hispanic	90.1	(±1.5)	90.8	(±1.5)	38.4	(±2.6)
Asian	88.3	(±3.6)	92.4	(±1.9)	37.4	(±6.0)
Total	91.2	(±1.1)	91.3	(±0.6)	33.5	(±1.6)

* Confidence interval.

TABLE 27. Percentage of middle school and high school students who were exposed to tobacco-related media and advertising, by state — state youth tobacco surveys, United States 2001–2002

State	Saw or heard antismoking commercials on TV or radio		Saw actors on TV or in movies using tobacco		Saw ads for tobacco products on the Internet	
	%	(95% CI)*	%	(95% CI)	%	(95% CI)
Middle school						
Alabama – Spring 02	75.6	(±3.3)	85.6	(±1.9)	37.6	(±2.9)
Connecticut – Spring 02	78.9	(±1.9)	79.8	(±2.9)	30.8	(±3.7)
Delaware – Spring 02	83.5	(±1.8)	83.7	(±1.8)	34.1	(±2.5)
Florida – Spring 01	79.9	(±2.1)	NA†		NA	
Florida – Spring 02	80.7	(±1.2)	NA		NA	
Georgia – Fall 01	75.0	(±2.2)	83.5	(±1.6)	34.1	(±3.0)
Idaho – Spring 01	84.7	(±2.4)	83.8	(±2.9)	30.5	(±2.5)
Illinois – Spring 02	83.2	(±2.5)	83.4	(±1.9)	30.6	(±3.7)
Iowa – Spring 02	91.3	(±2.7)	85.9	(±2.5)	33.7	(±3.6)
Kansas – Fall 02	74.4	(±3.6)	81.5	(±3.1)	35.0	(±3.5)
Kentucky – Spring 02	75.8	(±2.7)	87.5	(±2.3)	43.0	(±3.2)
Louisiana – Spring 01	73.0	(±2.9)	82.7	(±2.0)	38.3	(±2.4)
Maine – Spring 01	81.5	(±1.8)	83.7	(±1.3)	29.9	(±3.8)
Maryland – Spring 02	69.1	(±1.1)	80.6	(±0.9)	40.6	(±1.4)
Massachusetts – Spring 02	84.4	(±2.6)	79.6	(±2.4)	28.9	(±3.9)
Michigan – Spring 01	78.3	(±3.7)	83.6	(±2.5)	NA	
Minnesota – Spring 02	81.2	(±1.7)	83.2	(±1.6)	35.3	(±1.7)
Mississippi – Fall 02	NA		NA		NA	
Nebraska – Spring 02	84.1	(±1.4)	83.5	(±1.4)	34.8	(±2.3)
New Hampshire – Fall 01	70.4	(±3.6)	78.1	(±2.2)	29.5	(±2.2)
New Jersey – Fall 01	88.4	(±1.4)	90.2	(±1.2)	NA	
New York – Spring 02	83.2	(±3.1)	89.3	(±2.0)	43.9	(±1.4)
North Carolina – Spring 02	73.1	(±1.9)	85.5	(±1.4)	38.7	(±3.5)
Ohio – Spring 02	83.4	(±2.9)	86.0	(±1.9)	34.2	(±4.2)
Oklahoma – Spring 02	74.1	(±2.5)	85.3	(±1.5)	33.7	(±2.9)
Pennsylvania – Spring 01	78.8	(±1.5)	83.3	(±2.0)	34.0	(±2.6)
Rhode Island – Spring 01	83.0	(±2.8)	81.8	(±3.3)	27.0	(±2.4)
South Dakota – Fall 01	82.7	(±1.7)	83.7	(±1.6)	35.4	(±2.5)
Texas – Spring 01	NA		NA		25.3	(±1.5)
Vermont – Spring 02	74.6	(±3.2)	80.0	(±2.8)	31.3	(±2.5)
West Virginia – Spring 02	76.9	(±1.4)	87.0	(±0.7)	40.1	(±2.2)
Wisconsin – Spring 02	83.2	(±2.3)	82.7	(±3.6)	35.8	(±5.1)
Median	80.3		83.6		34.1	
High school						
Alabama – Spring 02	82.2	(±2.6)	89.4	(±1.9)	30.7	(±3.5)
Connecticut – Spring 02	84.9	(±1.7)	89.2	(±1.3)	26.1	(±2.8)
Delaware – Spring 02	88.6	(±2.8)	90.3	(±2.1)	31.5	(±6.4)
Florida – Spring 01	84.4	(±1.7)	NA†		NA	
Florida – Spring 02	84.8	(±1.6)	NA		NA	
Georgia	81.2	(±2.3)	89.5	(±1.8)	29.8	(±2.2)
Illinois – Spring 02	88.5	(±2.2)	89.2	(±2.4)	32.2	(±4.4)
Iowa – Spring 02	93.1	(±1.7)	88.6	(±1.6)	23.5	(±3.8)
Kansas – Fall 02	83.6	(±2.6)	89.9	(±1.9)	33.3	(±2.9)
Kentucky – Spring 02	86.1	(±1.8)	91.1	(±1.4)	35.0	(±2.9)
Maryland – Spring 02	78.1	(±0.7)	85.0	(±0.5)	37.9	(±0.9)
Massachusetts – Spring 02	88.7	(±1.9)	85.3	(±3.3)	19.6	(±2.8)
Michigan – Spring 01	81.3	(±2.0)	86.2	(±1.2)	NA	
Mississippi – Fall 02	NA		NA		NA	
Nebraska – Spring 02	91.2	(±0.9)	90.0	(±1.3)	28.4	(±2.6)
New Hampshire – Fall 01	81.9	(±2.2)	87.3	(±1.9)	23.7	(±4.1)
New Jersey – Fall 01	89.2	(±1.5)	90.6	(±0.8)	NA	
New York – Spring 02	89.4	(±2.4)	91.3	(±1.8)	36.6	(±4.9)
North Carolina – Spring 02	80.2	(±1.7)	90.4	(±1.1)	34.1	(±2.0)
Ohio – Spring 02	89.7	(±1.9)	90.2	(±2.0)	26.2	(±3.3)
Oklahoma – Spring 02	84.5	(±2.3)	90.7	(±1.8)	28.3	(±3.4)
Pennsylvania – Spring 01	82.7	(±2.1)	89.0	(±1.1)	25.9	(±1.9)
Rhode Island – Spring 01	89.5	(±2.2)	87.8	(±1.9)	23.8	(±3.1)
Texas – Spring 01	NA		NA		23.7	(±1.6)
West Virginia – Spring 02	83.9	(±3.0)	89.8	(±2.4)	33.0	(±3.2)
Wisconsin – Spring 02	91.8	(±2.1)	90.3	(±1.6)	25.4	(±3.6)
Median	84.9		89.7		28.4	

* Confidence interval.

† Question not asked.

TABLE 28. Percentage of middle school and high school students who participated in any community events to discourage young persons from using tobacco products, by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Never used tobacco*		Currently use tobacco†	
	%	(95% CI)§	%	(95% CI)
Middle school				
Sex				
Male	10.2	(±1.1)	19.3	(±3.9)
Female	13.0	(±1.9)	15.7	(±3.1)
Race/Ethnicity				
White, non-Hispanic	11.0	(±1.6)	16.4	(±3.2)
Black, non-Hispanic	15.6	(±2.8)	23.9	(±6.3)
Hispanic	10.1	(±1.9)	16.4	(±4.1)
Asian	9.0	(±4.5)	27.3	(±20.7)
Total	11.7	(±1.3)	17.9	(±2.6)
High school				
Sex				
Male	8.6	(±1.5)	9.9	(±2.0)
Female	13.8	(±1.8)	9.0	(±1.9)
Race/Ethnicity				
White, non-Hispanic	11.3	(±1.7)	7.8	(±1.6)
Black, non-Hispanic	14.4	(±3.0)	15.9	(±4.0)
Hispanic	9.7	(±2.5)	12.8	(±3.8)
Asian	10.1	(±3.8)	11.4	(±5.6)
Total	11.5	(±1.4)	9.5	(±1.5)

* Students were considered to have never used tobacco if they answered "no" to whether they have tried or experimented with cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks.

† Used cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥1 of the previous 30 days.

§ Confidence interval.

TABLE 29. Percentage of middle school and high school students who participated in any community events to discourage young persons from using tobacco products, by tobacco use status and state — state youth tobacco surveys, United States, 2001–2002

State	Never used tobacco*		Current tobacco user§	
	%	(95% CI)†	%	(95% CI)
Middle school				
Alabama – Spring 02	22.4	(±5.5)	26.1	(±7.0)
Connecticut – Spring 02	19.7	(±9.6)	17.3	(±7.8)
Delaware – Spring 02	25.1	(±2.8)	24.6	(±6.8)
Florida – Spring 01	12.8	(±1.3)	17.2	(±3.7)
Florida – Spring 02	11.9	(±1.0)	14.6	(±2.0)
Georgia – Fall 01	24.2	(±3.9)	20.3	(±3.7)
Idaho – Spring 01	33.8	(±5.0)	22.8	(±5.7)
Illinois – Spring 02	26.2	(±5.9)	23.0	(±5.6)
Iowa – Spring 02	33.6	(±5.7)	21.8	(±10.5)
Kansas – Fall 02	19.4	(±4.7)	21.1	(±6.5)
Kentucky – Spring 02	23.3	(±5.1)	19.9	(±6.5)
Louisiana – Spring 01	29.0	(±4.8)	24.6	(±3.9)
Maine – Spring 01	NA††		NA	
Maryland – Spring 02	18.4	(±1.9)	27.8	(±3.1)
Massachusetts – Spring 02	16.9	(±5.3)	20.7	(±7.4)
Michigan – Spring 01	23.3	(±4.0)	23.1	(±7.2)
Minnesota – Spring 02	26.8	(±6.0)	23.9	(±4.3)
Mississippi – Fall 02	31.6	(±3.6)	23.6	(±4.9)
Nebraska – Spring 02	29.5	(±4.0)	23.0	(±8.1)
New Hampshire – Fall 01	15.9	(±3.3)	17.2	(±7.1)
New Jersey – Fall 01	NA		NA	
New York – Spring 02	11.6	(±3.1)	18.2	(±10.0)
North Carolina – Spring 02	21.7	(±2.7)	22.3	(±3.3)
Ohio – Spring 02	24.1	(±4.6)	18.1	(±8.1)
Oklahoma – Spring 02	20.5	(±3.4)	18.0	(±4.7)
Pennsylvania – Spring 01	28.5	(±3.5)	18.7	(±5.0)
Rhode Island – Spring 01	28.3	(±7.8)	16.8	(±6.9)
South Dakota – Fall 01	29.3	(±5.4)	21.2	(±6.1)
Texas – Spring 01	16.7	(±2.3)	19.1	(±2.3)
Vermont – Spring 02	21.8	(±3.9)	22.0	(±3.4)
West Virginia – Spring 02	25.3	(±2.4)	19.3	(±2.4)
Wisconsin – Spring 02	25.0	(±5.5)	24.3	(±8.3)
Median	23.7		21.2	
High school				
Alabama	18.2	(±4.7)	14.4	(±3.9)
Connecticut – Spring 02	10.6	(±2.1)	11.9	(±4.0)
Delaware – Spring 02	19.1	(±5.0)	11.3	(±2.4)
Florida – Spring 01	11.5	(±1.9)	10.9	(±2.2)
Florida – Spring 02	8.7	(±1.1)	9.8	(±1.8)
Georgia – Fall 01	18.3	(±2.9)	11.6	(±2.5)
Illinois – Spring 02	14.1	(±3.9)	12.0	(±3.4)
Iowa – Spring 02	20.5	(±3.7)	13.6	(±4.0)
Kansas – Fall 02	16.6	(±4.9)	11.1	(±2.7)
Kentucky – Spring 02	14.8	(±4.3)	6.5	(±2.4)
Maryland – Spring 02	13.7	(±1.0)	17.6	(±1.3)
Massachusetts – Spring 02	11.3	(±4.2)	10.6	(±3.1)
Michigan – Spring 01	12.5	(±3.2)	12.4	(±2.6)
Mississippi – Fall 02	25.9	(±3.3)	16.2	(±4.3)
Nebraska – Spring 02	22.4	(±3.1)	10.9	(±3.1)
New Hampshire – Fall 01	11.8	(±3.6)	8.2	(±3.0)
New Jersey – Fall 01	NA††		NA	
New York – Spring 02	12.8	(±4.3)	13.9	(±4.9)
North Carolina – Spring 02	13.4	(±2.3)	12.6	(±2.4)
Ohio – Spring 02	16.9	(±4.2)	9.5	(±4.0)
Oklahoma – Spring 02	15.4	(±4.4)	9.6	(±2.9)
Pennsylvania – Spring 01	17.2	(±2.4)	10.0	(±1.3)
Rhode Island – Spring 01	16.6	(±3.5)	11.4	(±2.4)
Texas – Spring 01	12.3	(±3.2)	11.4	(±2.1)
West Virginia – Spring 02	24.9	(±6.3)	16.4	(±6.1)
Wisconsin – Spring 02	18.6	(±6.6)	10.0	(±2.2)
Median	15.4		11.4	

* Students were considered to have never used tobacco if they answered "no" to whether they have tried or experimented with cigarettes, cigars, smokeless tobacco, pipe tobacco, or bidis. Use of kreteks was not asked.

† Confidence interval.

§ Used cigarettes, cigars, smokeless tobacco, pipe tobacco, or bidis on ≥1 of the previous 30 days.

†† Question not asked.

TABLE 30. Percentage of middle school and high school students who bought or received, or would wear or use, tobacco company merchandise*, by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Bought or received anything with tobacco company name or picture on it				Would wear or use something with tobacco company name or picture on it			
	Never used tobacco†		Currently use tobacco§		Never used tobacco		Currently use tobacco	
	%	(95% CI¶)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Sex								
Male	9.2	(±1.3)	43.5	(±4.1)	15.7	(±2.0)	60.4	(±4.6)
Female	8.3	(±1.2)	39.7	(±5.7)	6.9	(±1.1)	54.9	(±5.7)
Race/Ethnicity								
White, non-Hispanic	8.7	(±1.2)	42.3	(±4.1)	11.0	(±1.6)	58.1	(±4.5)
Black, non-Hispanic	8.4	(±1.8)	35.7	(±7.3)	9.2	(±1.7)	49.9	(±7.6)
Hispanic	9.2	(±1.4)	40.5	(±5.9)	12.7	(±1.9)	60.9	(±5.7)
Asian	5.7	(±3.1)	52.5	(±17.4)	10.3	(±3.9)	64.1	(±16.8)
Total	8.7	(±0.9)	41.8	(±3.1)	11.0	(±1.2)	58.1	(±3.2)
High school								
Sex								
Male	8.1	(±1.5)	34.6	(±3.0)	19.0	(±1.8)	57.8	(±3.5)
Female	7.2	(±1.2)	26.7	(±3.2)	9.3	(±1.4)	50.9	(±3.7)
Race/Ethnicity								
White, non-Hispanic	7.2	(±1.3)	31.6	(±2.7)	13.1	(±1.5)	57.9	(±3.3)
Black, non-Hispanic	8.2	(±2.8)	26.4	(±5.2)	12.9	(±2.4)	37.1	(±4.8)
Hispanic	9.5	(±2.2)	32.6	(±3.6)	17.4	(±2.7)	51.7	(±4.7)
Asian	6.0	(±3.1)	34.4	(±12.8)	12.0	(±3.4)	47.9	(±17.4)
Total	7.7	(±1.0)	31.3	(±2.4)	13.7	(±1.3)	54.9	(±2.9)

* For example, a cigarette lighter or T-shirt.

† Students were considered to have never used tobacco if they answered "no" to whether they have ever tried or experimented with cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks.

§ Used cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks on ≥1 of the previous 30 days.

¶ Confidence interval.

TABLE 31. Percentage of middle school and high school students who bought or received, or would wear or use, tobacco company merchandise,* by tobacco use status and state — state youth tobacco surveys, 2001–2002

State	Bought or received anything with tobacco company name or picture on it				Would wear or use something with tobacco company name or picture on it			
	Never used tobacco†		Currently use tobacco‡		Never used tobacco		Currently use tobacco	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Alabama – Spring 02	10.3	(±2.3)	42.5	(±10.1)	13.0	(±3.8)	56.8	(±6.4)
Connecticut – Spring 02	11.2	(±2.5)	39.5	(±8.2)	9.8	(±1.6)	44.2	(±6.9)
Delaware – Spring 02	7.8	(±1.2)	39.8	(±6.8)	10.4	(±2.1)	52.3	(±6.0)
Florida – Spring 01	9.9	(±1.5)	39.4	(±5.1)	12.8	(±1.4)	58.4	(±5.1)
Florida – Spring 02	8.3	(±0.8)	39.0	(±2.3)	13.0	(±0.9)	57.5	(±2.6)
Georgia – Fall 01	12.1	(±2.1)	38.9	(±6.3)	10.8	(±2.3)	54.4	(±5.1)
Idaho – Spring 01	9.6	(±1.7)	43.9	(±5.7)	9.9	(±2.0)	56.0	(±6.0)
Illinois – Spring 02	11.9	(±2.8)	39.0	(±8.4)	13.8	(±3.1)	54.4	(±7.7)
Iowa – Spring 02	16.1	(±1.3)	42.3	(±9.6)	12.6	(±2.6)	58.6	(±6.3)
Kansas – Fall 02	8.8	(±2.3)	39.2	(±7.8)	11.7	(±2.6)	57.6	(±8.5)
Kentucky – Spring 02	13.6	(±2.7)	50.2	(±6.2)	11.4	(±2.5)	59.4	(±7.9)
Louisiana – Spring 01	9.9	(±2.3)	45.1	(±3.9)	13.8	(±1.9)	56.9	(±3.9)
Maine – Spring 01	14.9	(±1.6)	52.7	(±3.9)	11.4	(±1.7)	65.3	(±5.6)
Maryland – Spring 02	13.0	(±1.0)	44.7	(±3.4)	11.3	(±1.0)	54.7	(±3.5)
Massachusetts – Spring 02	8.3	(±1.6)	33.0	(±7.6)	8.5	(±1.8)	46.6	(±9.7)
Michigan – Spring 01	12.1	(±2.3)	43.4	(±6.6)	13.3	(±2.5)	52.1	(±8.7)
Minnesota – Spring 02	10.9	(±1.1)	41.7	(±4.7)	8.0	(±1.3)	52.8	(±6.8)
Mississippi – Fall 02	NA**		NA		20.9	(±2.8)	67.9	(±5.6)
Nebraska – Spring 02	11.8	(±1.9)	47.1	(±7.7)	12.2	(±2.1)	54.6	(±6.6)
New Hampshire – Fall 01	12.4	(±2.0)	43.4	(±9.5)	11.6	(±1.8)	54.0	(±9.5)
New Jersey – Fall 01	13.7	(±1.7)	40.4	(±8.8)	12.2	(±1.8)	45.7	(±6.5)
New York – Spring 02	6.4	(±1.4)	46.0	(±5.7)	11.0	(±2.7)	54.5	(±7.4)
North Carolina – Spring 02	9.8	(±1.7)	45.6	(±4.0)	11.5	(±1.8)	53.7	(±5.1)
Ohio – Spring 02	12.7	(±2.6)	47.6	(±8.2)	11.1	(±2.5)	58.7	(±7.0)
Oklahoma – Spring 02	8.4	(±2.2)	48.3	(±8.5)	10.6	(±2.3)	65.2	(±7.5)
Pennsylvania – Spring 01	10.6	(±1.5)	46.3	(±2.6)	11.8	(±1.1)	62.8	(±3.2)
Rhode Island – Spring 01	9.6	(±2.3)	31.9	(±6.3)	7.9	(±1.9)	48.1	(±11.0)
South Dakota – Fall 01	12.6	(±2.3)	44.6	(±4.7)	11.0	(±2.3)	50.6	(±6.7)
Texas – Spring 01	14.8	(±1.6)	46.5	(±4.2)	10.2	(±1.7)	52.7	(±5.1)
Vermont – Spring 02	11.2	(±1.3)	50.7	(±6.9)	12.7	(±2.6)	62.2	(±3.8)
West Virginia – Spring 02	13.7	(±1.0)	49.2	(±2.7)	15.9	(±1.6)	67.5	(±2.3)
Wisconsin – Spring 02	10.1	(±2.4)	41.6	(±8.7)	11.6	(±1.8)	53.7	(±11.2)
Median	11.2		43.4		11.6		54.7	
High school								
Alabama – Spring 02	11.7	(±3.2)	39.1	(±6.1)	18.1	(±5.2)	61.1	(±6.8)
Connecticut – Spring 02	14.0	(±2.3)	43.5	(±4.7)	17.1	(±2.6)	50.9	(±4.3)
Delaware – Spring 02	6.1	(±1.4)	33.9	(±10.3)	10.9	(±4.0)	48.2	(±3.4)
Florida – Spring 01	9.4	(±1.4)	29.8	(±3.2)	18.5	(±1.5)	53.2	(±3.4)
Florida – Spring 02	7.5	(±0.9)	30.3	(±2.0)	16.7	(±1.5)	55.3	(±2.2)
Georgia – Fall 01	13.0	(±2.1)	43.7	(±3.9)	16.1	(±2.3)	60.4	(±4.0)
Illinois – Spring 02	15.5	(±4.4)	45.2	(±6.8)	18.0	(±3.8)	56.5	(±6.1)
Iowa – Spring 02	10.4	(±2.8)	40.7	(±3.4)	16.5	(±3.0)	67.9	(±4.4)
Kansas – Fall 02	5.7	(±2.2)	32.9	(±4.7)	13.7	(±3.6)	61.8	(±4.3)
Kentucky – Spring 02	16.3	(±3.3)	46.1	(±5.5)	19.2	(±5.0)	65.7	(±4.7)
Maryland – Spring 02	14.1	(±0.7)	42.9	(±1.3)	14.9	(±0.8)	50.2	(±1.3)
Massachusetts – Spring 02	10.8	(±2.7)	32.1	(±4.8)	15.7	(±3.4)	57.9	(±5.9)
Michigan – Spring 01	14.0	(±2.2)	35.5	(±4.4)	19.0	(±2.3)	55.8	(±2.8)
Mississippi – Fall 02	24.4	(±3.1)	63.3	(±4.9)	24.4	(±3.1)	63.3	(±4.9)
Nebraska – Spring 02	12.1	(±2.4)	41.4	(±2.4)	19.6	(±3.8)	62.1	(±3.6)
New Hampshire – Fall 01	12.5	(±3.8)	32.8	(±4.7)	15.0	(±3.1)	56.2	(±5.8)
New Jersey – Fall 01	15.2	(±1.9)	39.1	(±4.8)	18.3	(±2.6)	53.1	(±3.6)
New York – Spring 02	7.5	(±1.8)	34.2	(±6.3)	15.7	(±2.5)	55.9	(±8.6)
North Carolina – Spring 02	13.0	(±2.6)	39.6	(±3.7)	18.0	(±2.2)	58.0	(±3.6)
Ohio – Spring 02	12.5	(±3.1)	41.6	(±5.2)	16.5	(±3.4)	57.4	(±5.1)
Oklahoma – Spring 02	11.1	(±2.6)	40.9	(±4.1)	14.7	(±3.8)	64.9	(±4.9)
Pennsylvania – Spring 01	10.9	(±1.4)	35.8	(±3.4)	17.8	(±2.1)	58.4	(±4.1)
Rhode Island – Spring 01	7.9	(±1.5)	29.8	(±4.1)	15.3	(±3.8)	49.2	(±4.2)
Texas – Spring 01	17.7	(±1.8)	52.1	(±6.4)	16.1	(±2.7)	58.6	(±3.3)
West Virginia – Spring 02	9.8	(±3.1)	40.9	(±5.0)	19.8	(±3.7)	63.0	(±6.0)
Wisconsin – Spring 02	11.0	(±2.7)	30.2	(±5.6)	18.6	(±4.7)	59.0	(±5.2)
Median	11.9		39.4		16.9		58.0	

* For example, a cigarette lighter or T-shirt.

† Students were considered to have never used tobacco if they answered “no” to whether they have ever tried or experimented with cigarettes, cigars, smokeless tobacco, pipe tobacco, bidis, or kreteks.

§ Confidence interval.

¶ Used cigarettes, cigars, smokeless tobacco, pipe tobacco, or bidis on ≥1 of the previous 30 days.

** Question not asked.

TABLE 32. Percentage of current cigarette smokers* who tried to quit smoking during the previous 12 months and who want to stop smoking cigarettes, by sex and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Tried to quit in previous 12 months		Want to stop smoking cigarettes	
	%	(95% CI†)	%	(95% CI)
Middle school				
Sex				
Male	51.7	(±5.3)	47.5	(±6.0)
Female	59.6	(±6.2)	52.1	(±7.2)
Race/Ethnicity				
White, non-Hispanic	55.9	(±5.3)	49.9	(±6.7)
Black, non-Hispanic	53.5	(±9.5)	49.3	(±12.3)
Hispanic	55.2	(±7.6)	49.9	(±9.0)
Asian	§		§	
Total	55.4	(±4.5)	49.6	(±5.5)
High school				
Sex				
Male	51.1	(±3.2)	61.0	(±3.2)
Female	55.4	(±4.3)	63.4	(±3.5)
Race/Ethnicity				
White, non-Hispanic	52.5	(±3.3)	63.2	(±2.4)
Black, non-Hispanic	58.5	(±6.5)	68.2	(±7.4)
Hispanic	51.5	(±5.8)	56.6	(±5.7)
Asian	55.7	(±19.8)	54.2	(±21.6)
Total	53.1	(±2.8)	62.1	(±2.2)

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Sample size <35.

TABLE 33. Percentage of current cigarette smokers* who tried to quit smoking during the previous 12 months and who want to stop smoking cigarettes, by state — state youth tobacco surveys, United States, 2001–2002

State	Tried to quit smoking cigarettes during previous 12 months		Want to stop smoking cigarettes	
	%	(95% CI†)	%	(95% CI)
Middle school				
Alabama – Spring 02	54.5	(±9.3)	48.1	(±12.4)
Connecticut – Spring 02	56.2	(±10.3)	50.3	(±11.0)
Delaware – Spring 02	55.4	(±8.2)	55.1	(±12.0)
Florida – Spring 01	56.4	(±5.1)	52.2	(±3.4)
Florida – Spring 02	55.7	(±3.0)	52.2	(±3.6)
Georgia – Fall 01	60.6	(±6.1)	54.2	(±8.0)
Idaho – Spring 01	55.3	(±8.0)	54.9	(±9.2)
Illinois – Spring 02	47.5	(±12.6)	46.3	(±9.3)
Iowa – Spring 02	69.7	(±13.9)	65.9	(±8.3)
Kansas – Fall 02	48.4	(±14.5)	49.0	(±14.2)
Kentucky – Spring 02	61.2	(±5.9)	NA§	
Louisiana – Spring 01	52.6	(±4.8)	53.8	(±4.9)
Maine – Spring 01	54.3	(±5.4)	50.3	(±4.9)
Maryland – Spring 02	59.6	(±2.4)	51.6	(±5.4)
Massachusetts – Spring 02	51.8	(±7.4)	47.3	(±9.9)
Michigan – Spring 01	56.3	(±7.4)	41.2	(±8.5)
Minnesota – Spring 02	61.0	(±7.0)	61.5	(±6.3)
Mississippi – Fall 02	NA		49.1	(±7.3)
Nebraska – Spring 02	69.5	(±4.9)	58.7	(±10.8)
New Hampshire – Fall 01	59.6	(±12.1)	51.8	(±16.8)
New Jersey – Fall 01	53.5	(±9.8)	53.9	(±10.5)
New York – Spring 02	53.5	(±11.6)	52.6	(±13.5)
North Carolina – Spring 02	62.2	(±7.0)	53.2	(±7.2)
Ohio – Spring 02	59.8	(±9.5)	63.4	(±12.9)
Oklahoma – Spring 02	56.3	(±10.8)	40.9	(±9.0)
Pennsylvania – Spring 01	58.7	(±7.5)	61.0	(±10.6)
Rhode Island – Spring 01	53.5	(±10.2)	48.8	(±10.5)
South Dakota – Fall 01	61.2	(±9.7)	57.3	(±12.9)
Texas – Spring 01	54.7	(±5.3)	51.9	(±5.7)
Vermont – Spring 02	49.2	(±8.8)	47.4	(±9.0)
West Virginia – Spring 02	56.7	(±2.4)	50.8	(±2.9)
Wisconsin – Spring 02	61.6	(±10.6)	57.5	(±8.7)
Median	56.3		52.2	
High school				
Alabama – Spring 02	57.4	(±7.0)	61.4	(±7.6)
Connecticut – Spring 02	53.3	(±5.1)	62.7	(±5.9)
Delaware – Spring 02	60.3	(±3.8)	63.8	(±6.1)
Florida – Spring 01	56.6	(±3.4)	55.5	(±4.0)
Florida – Spring 02	57.7	(±2.8)	56.5	(±2.4)
Georgia – Fall 01	57.7	(±4.3)	47.7	(±5.6)
Illinois – Spring 02	55.9	(±7.6)	53.6	(±5.7)
Iowa – Spring 02	58.6	(±7.4)	59.0	(±5.7)
Kansas – Fall 02	50.0	(±6.5)	52.8	(±7.6)
Kentucky – Spring 02	54.3	(±3.5)	NA§	
Maryland – Spring 02	55.7	(±1.3)	51.3	(±1.7)
Massachusetts – Spring 02	63.3	(±7.4)	67.8	(±7.3)
Michigan – Spring 01	57.0	(±3.2)	57.8	(±4.2)
Mississippi – Fall 02	NA		60.4	(±3.6)
Nebraska – Spring 02	63.2	(±3.9)	63.7	(±6.5)
New Hampshire – Fall 01	59.9	(±5.6)	57.2	(±6.8)
New Jersey – Fall 01	51.6	(±6.4)	53.4	(±5.9)
New York – Spring 02	59.8	(±9.6)	61.5	(±7.8)
North Carolina – Spring 02	52.3	(±3.0)	49.5	(±3.7)
Ohio – Spring 02	61.8	(±5.5)	65.9	(±6.3)
Oklahoma – Spring 02	57.7	(±7.6)	54.6	(±5.1)
Pennsylvania – Spring 01	57.7	(±2.7)	59.9	(±2.9)
Rhode Island – Spring 01	57.5	(±5.6)	58.6	(±5.0)
Texas – Spring 01	41.8	(±7.0)	49.2	(±3.9)
West Virginia – Spring 02	57.9	(±4.5)	59.2	(±5.2)
Wisconsin – Spring 02	64.4	(±6.3)	60.8	(±6.2)
Median	57.5		58.6	

* Smoked cigarettes on ≥1 of the previous 30 days.

† Confidence interval.

§ Question not asked.

TABLE 34. Percentage of middle school and high school students who were in the same room as or who rode in a car with someone who was smoking cigarettes on ≥ 1 of the previous 7 days or who think smoke from other persons' cigarettes is harmful, by smoking status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	In the same room with someone who was smoking on ≥ 1 of previous 7 days				Rode in a car with someone who was smoking on ≥ 1 of previous 7 days				Think smoke from other persons' cigarettes is harmful			
	Never smoked*		Currently smoke [†]		Never smoked		Currently smoke		Never smoked		Currently smoke	
	%	(95% CI) [§]	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school												
Sex												
Male	44.8	(± 2.7)	84.8	(± 4.1)	30.2	(± 2.5)	77.3	(± 4.4)	88.8	(± 1.8)	74.0	(± 5.3)
Female	49.4	(± 2.5)	91.5	(± 2.7)	32.7	(± 2.9)	85.8	(± 3.6)	93.4	(± 1.2)	82.9	(± 4.4)
Race/Ethnicity												
White, non-Hispanic	51.6	(± 2.9)	91.0	(± 3.0)	33.4	(± 3.3)	84.3	(± 3.5)	93.6	(± 1.1)	80.8	(± 4.0)
Black, non-Hispanic	43.4	(± 3.2)	77.7	(± 6.4)	33.3	(± 3.7)	71.3	(± 7.6)	84.6	(± 2.5)	72.2	(± 7.8)
Hispanic	35.1	(± 3.6)	84.1	(± 5.1)	22.5	(± 2.4)	76.2	(± 6.6)	87.1	(± 2.7)	75.6	(± 6.1)
Asian	27.6	(± 5.6)	89.3	(± 10.3)	22.1	(± 6.4)	83.5	(± 12.3)	94.1	(± 2.8)	91.0	(± 8.3)
Total	47.1	(± 2.3)	88.3	(± 2.5)	31.5	(± 2.4)	81.7	(± 2.9)	91.1	(± 1.3)	78.6	(± 3.1)
High school												
Sex												
Male	51.5	(± 2.7)	89.7	(± 1.9)	26.8	(± 2.7)	81.7	(± 2.6)	93.2	(± 1.7)	87.5	(± 2.5)
Female	55.0	(± 2.8)	93.4	(± 1.8)	31.0	(± 2.8)	86.1	(± 2.2)	96.7	(± 0.8)	93.3	(± 1.8)
Race/Ethnicity												
White, non-Hispanic	56.4	(± 2.6)	93.4	(± 1.3)	29.6	(± 2.8)	86.0	(± 2.4)	96.7	(± 0.9)	92.6	(± 2.1)
Black, non-Hispanic	50.7	(± 3.8)	84.2	(± 5.1)	34.2	(± 5.3)	72.6	(± 6.2)	90.2	(± 2.5)	81.2	(± 5.2)
Hispanic	44.1	(± 4.6)	86.7	(± 3.2)	25.0	(± 3.2)	78.2	(± 4.5)	91.2	(± 2.5)	84.5	(± 4.3)
Asian	46.6	(± 6.0)	84.0	(± 12.1)	19.0	(± 4.8)	91.6	(± 6.7)	96.9	(± 2.0)	84.3	(± 12.5)
Total	53.3	(± 2.2)	91.4	(± 1.3)	29.1	(± 2.3)	83.7	(± 2.0)	95.1	(± 1.0)	90.2	(± 1.8)

* Students were considered to have never smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

[†] Smoked cigarettes on ≥ 1 of the previous 30 days.

[§] Confidence interval.

TABLE 35. Percentage of middle school and high school students who were in the same room as or who rode in a car with someone who was smoking cigarettes on ≥ 1 of the previous 7 days or who think smoke from other persons' cigarettes is harmful, by smoking status and state — State Youth Tobacco Surveys, 2001–2002

Sex and Race/Ethnicity	In the same room with someone who was smoking on ≥ 1 of previous 7 days				Rode in a car with someone who was smoking on ≥ 1 of previous 7 days				Think smoke from other persons' cigarettes is harmful			
	Never smoked*		Currently smoke [†]		Never smoked		Currently smoke		Never smoked		Currently smoke	
	%	(95% CI) [§]	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school												
Alabama – Spring 02	43.4	(± 4.9)	83.8	(± 6.0)	38.3	(± 5.0)	77.8	(± 7.0)	90.3	(± 2.5)	78.6	(± 6.4)
Connecticut – Spring 02	44.7	(± 4.8)	86.5	(± 6.7)	31.4	(± 5.6)	79.1	(± 9.5)	92.9	(± 1.5)	75.8	(± 8.8)
Delaware – Spring 02	45.0	(± 2.5)	85.7	(± 5.0)	33.7	(± 2.9)	81.8	(± 5.4)	95.4	(± 1.1)	80.9	(± 8.0)
Florida – Spring 01	47.8	(± 3.2)	87.1	(± 4.1)	33.6	(± 2.7)	78.9	(± 5.0)	87.1	(± 2.2)	79.0	(± 4.1)
Florida – Spring 02	46.9	(± 1.6)	84.2	(± 2.6)	32.6	(± 1.5)	78.6	(± 2.9)	87.6	(± 2.1)	79.3	(± 2.7)
Georgia – Fall 01	43.3	(± 2.7)	82.5	(± 5.9)	31.5	(± 3.0)	75.2	(± 5.3)	91.3	(± 2.6)	80.7	(± 5.8)
Idaho – Spring 01	34.5	(± 4.4)	85.8	(± 6.1)	21.3	(± 3.8)	80.9	(± 7.2)	94.1	(± 1.6)	84.5	(± 6.4)
Illinois – Spring 02	44.8	(± 3.4)	88.9	(± 2.7)	27.5	(± 4.7)	76.5	(± 6.3)	94.1	(± 1.7)	84.8	(± 13.6)
Iowa – Spring 02	50.5	(± 4.2)	89.4	(± 8.1)	37.1	(± 5.0)	91.6	(± 6.3)	95.7	(± 0.7)	89.1	(± 7.1)
Kansas – Fall 02	42.6	(± 3.4)	78.2	(± 8.5)	31.9	(± 4.9)	76.1	(± 8.4)	94.0	(± 2.1)	78.6	(± 8.0)
Kentucky – Spring 02	63.3	(± 5.0)	89.9	(± 3.3)	49.8	(± 4.5)	85.5	(± 4.4)	91.2	(± 2.6)	82.2	(± 6.2)
Louisiana – Spring 01	47.0	(± 2.8)	89.0	(± 2.9)	35.6	(± 3.1)	80.7	(± 3.8)	87.0	(± 3.3)	77.1	(± 5.0)
Maine – Spring 01	42.6	(± 3.5)	84.1	(± 4.5)	30.3	(± 3.3)	79.1	(± 4.5)	96.0	(± 0.7)	82.7	(± 3.8)
Maryland – Spring 02	37.7	(± 1.4)	76.9	(± 3.6)	27.1	(± 1.5)	73.2	(± 3.7)	90.6	(± 1.0)	79.0	(± 3.5)
Massachusetts – Spring 02	38.6	(± 4.2)	75.1	(± 8.5)	25.1	(± 4.0)	70.0	(± 8.7)	93.6	(± 1.3)	79.7	(± 8.7)
Michigan – Spring 01	45.3	(± 4.5)	84.5	(± 6.1)	35.2	(± 3.6)	79.9	(± 7.3)	93.0	(± 2.8)	85.8	(± 4.9)
Minnesota – Spring 02	41.8	(± 2.9)	86.6	(± 4.1)	27.4	(± 3.0)	80.5	(± 4.9)	94.3	(± 1.0)	84.7	(± 5.7)
Mississippi – Fall 02	NA [¶]		NA		NA		NA		93.1	(± 1.7)	79.8	(± 4.5)
Nebraska – Spring 02	48.4	(± 2.8)	86.2	(± 5.2)	35.6	(± 4.2)	79.7	(± 8.4)	95.6	(± 1.1)	89.7	(± 3.7)
New Hampshire – Fall 01	44.1	(± 4.1)	87.6	(± 8.1)	32.0	(± 3.9)	83.2	(± 9.1)	93.4	(± 1.4)	81.0	(± 7.7)
New Jersey – Fall 01	39.0	(± 3.5)	83.4	(± 5.0)	27.8	(± 4.3)	72.9	(± 6.5)	91.4	(± 1.6)	71.1	(± 5.9)
New York – Spring 02	47.8	(± 7.1)	86.1	(± 7.3)	33.0	(± 6.1)	80.1	(± 9.8)	91.8	(± 2.4)	77.3	(± 10.0)
North Carolina – Spring 02	45.9	(± 3.3)	85.8	(± 3.6)	35.7	(± 3.6)	76.5	(± 4.7)	89.3	(± 2.1)	82.3	(± 4.5)
Ohio – Spring 02	56.5	(± 4.3)	89.0	(± 4.7)	36.7	(± 5.1)	86.0	(± 5.0)	93.8	(± 2.2)	82.5	(± 5.4)
Oklahoma – Spring 02	46.0	(± 2.9)	85.2	(± 4.5)	33.4	(± 4.1)	81.4	(± 3.9)	92.8	(± 1.4)	83.7	(± 4.8)
Pennsylvania – Spring 01	51.7	(± 2.4)	90.8	(± 3.4)	34.4	(± 2.3)	80.3	(± 2.9)	93.9	(± 1.4)	83.3	(± 6.9)
Rhode Island – Spring 01	47.0	(± 4.6)	88.2	(± 4.2)	32.0	(± 3.3)	75.0	(± 4.1)	94.5	(± 1.4)	89.3	(± 4.2)
South Dakota – Fall 01	46.6	(± 3.7)	84.7	(± 7.2)	31.5	(± 3.8)	86.5	(± 5.5)	92.3	(± 1.2)	83.9	(± 6.0)
Texas – Spring 01	31.1	(± 2.2)	73.5	(± 3.6)	25.0	(± 2.8)	73.0	(± 4.8)	90.0	(± 1.9)	70.6	(± 3.6)
Vermont – Spring 02	41.6	(± 5.3)	90.5	(± 3.6)	31.9	(± 4.7)	81.4	(± 3.4)	91.8	(± 1.5)	82.5	(± 4.5)
West Virginia – Spring 02	57.3	(± 2.3)	90.7	(± 1.6)	39.0	(± 2.3)	84.4	(± 2.1)	94.2	(± 0.9)	82.3	(± 2.3)
Wisconsin – Spring 02	48.2	(± 5.3)	89.8	(± 4.6)	35.0	(± 5.5)	84.1	(± 7.1)	92.7	(± 2.3)	89.6	(± 6.6)
Median	45.3		86.1		32.6		79.9		93.0		82.3	
High school												
Alabama	50.3	(± 5.4)	86.5	(± 4.3)	31.6	(± 3.8)	88.6	(± 2.5)	93.3	(± 3.3)	85.5	(± 4.0)
Connecticut – Spring 02	51.4	(± 5.6)	87.9	(± 2.9)	28.2	(± 5.6)	81.7	(± 3.3)	94.4	(± 1.3)	87.1	(± 3.2)
Delaware – Spring 02	54.7	(± 2.8)	92.6	(± 2.0)	34.7	(± 4.7)	89.9	(± 4.4)	97.3	(± 1.1)	88.8	(± 6.2)
Florida – Spring 01	54.4	(± 3.4)	89.7	(± 2.1)	29.7	(± 2.2)	83.3	(± 2.9)	89.8	(± 2.2)	88.8	(± 2.5)
Florida – Spring 02	51.5	(± 2.9)	89.2	(± 1.7)	28.9	(± 2.1)	81.0	(± 2.3)	89.1	(± 1.7)	88.7	(± 2.1)
Georgia – Fall 01	53.0	(± 3.6)	90.6	(± 2.9)	30.7	(± 3.1)	79.8	(± 3.4)	95.2	(± 1.6)	90.7	(± 2.5)
Illinois – Spring 02	55.9	(± 5.0)	92.0	(± 3.5)	32.0	(± 6.7)	85.6	(± 4.8)	96.0	(± 2.0)	88.7	(± 4.5)
Iowa – Spring 02	54.1	(± 4.5)	92.2	(± 3.1)	29.6	(± 3.9)	86.2	(± 4.3)	96.1	(± 1.7)	90.0	(± 3.4)
Kansas – Fall 02	47.8	(± 4.4)	90.5	(± 3.5)	25.2	(± 5.3)	81.5	(± 5.5)	95.0	(± 1.5)	87.9	(± 4.8)
Kentucky – Spring 02	73.6	(± 4.1)	95.9	(± 2.3)	44.9	(± 5.8)	88.3	(± 3.1)	93.6	(± 2.3)	91.5	(± 2.8)
Maryland – Spring 02	45.6	(± 1.1)	81.6	(± 1.2)	27.3	(± 1.0)	78.9	(± 1.2)	92.0	(± 0.6)	80.9	(± 1.2)
Massachusetts – Spring 02	53.0	(± 3.9)	93.4	(± 3.6)	30.9	(± 4.8)	82.7	(± 3.7)	97.0	(± 1.4)	90.9	(± 3.5)
Michigan – Spring 01	48.1	(± 3.1)	87.8	(± 2.1)	28.7	(± 3.8)	82.4	(± 2.1)	94.0	(± 1.8)	88.4	(± 2.8)
Mississippi – Fall 02	NA [¶]		NA		NA		NA		93.9	(± 2.4)	81.8	(± 4.6)
Nebraska – Spring 02	55.3	(± 3.5)	92.3	(± 1.6)	28.0	(± 4.1)	86.2	(± 2.5)	97.4	(± 1.3)	90.1	(± 2.5)
New Hampshire – Fall 01	50.7	(± 3.5)	90.1	(± 3.2)	30.0	(± 3.5)	87.5	(± 3.0)	96.6	(± 1.5)	91.3	(± 4.3)
New Jersey – Fall 01	48.5	(± 3.8)	84.6	(± 3.0)	31.5	(± 4.5)	81.3	(± 4.3)	93.0	(± 1.5)	84.8	(± 2.5)
New York – Spring 02	51.1	(± 5.5)	90.3	(± 2.5)	26.8	(± 2.2)	83.2	(± 6.5)	93.7	(± 3.1)	88.9	(± 2.7)
North Carolina – Spring 02	61.0	(± 3.6)	90.5	(± 1.9)	38.4	(± 2.7)	82.8	(± 2.5)	92.9	(± 2.4)	86.9	(± 2.1)
Ohio – Spring 02	61.1	(± 5.6)	95.0	(± 3.4)	30.9	(± 5.3)	87.6	(± 5.4)	96.8	(± 1.0)	92.5	(± 3.4)
Oklahoma – Spring 02	57.3	(± 5.2)	93.6	(± 2.2)	30.7	(± 4.1)	87.7	(± 3.8)	95.9	(± 1.6)	87.4	(± 2.9)
Pennsylvania – Spring 01	60.4	(± 2.5)	91.6	(± 1.6)	32.9	(± 2.3)	83.9	(± 2.2)	94.1	(± 1.5)	92.7	(± 1.9)
Rhode Island – Spring 01	57.8	(± 4.3)	87.9	(± 5.1)	33.5	(± 5.0)	83.6	(± 6.1)	94.3	(± 2.0)	90.3	(± 4.0)
Texas – Spring 01	40.4	(± 4.0)	83.8	(± 3.6)	23.7	(± 3.1)	72.6	(± 4.0)	93.7	(± 1.5)	86.5	(± 3.5)
West Virginia – Spring 02	65.1	(± 4.3)	92.5	(± 2.7)	38.1	(± 4.2)	87.6	(± 4.4)	94.7	(± 2.5)	91.4	(± 3.4)
Wisconsin – Spring 02	51.4	(± 4.9)	90.6	(± 5.2)	25.2	(± 3.8)	83.7	(± 5.9)	94.4	(± 1.9)	91.9	(± 4.0)
Median	53.0		90.5		30.7		83.6		94.4		88.8	

* Students were considered to have never smoked if they answered "no" to whether they have tried or experimented with cigarette smoking, even one or two puffs.

[†] Smoked cigarettes on ≥ 1 of the previous 30 days.

[§] Confidence interval.

[¶] Question not asked.

TABLE 36. Percentage of middle school and high school students who live in a home in which someone else smokes cigarettes, by smoking status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Someone else in home smokes cigarettes			
	Never smoked*		Currently smoke†	
	%	(95% CI)§	%	(95% CI)
Middle school				
Sex				
Male	34.1	(±2.6)	69.8	(±4.9)
Female	32.7	(±2.4)	73.0	(±4.2)
Race/Ethnicity				
White, non-Hispanic	33.7	(±2.9)	73.6	(±4.0)
Black, non-Hispanic	35.2	(±3.6)	68.7	(±8.0)
Hispanic	29.7	(±3.8)	64.2	(±7.7)
Asian	26.6	(±6.8)	68.8	(±20.2)
Total	33.3	(±2.2)	71.5	(±3.2)
High school				
Sex				
Male	28.3	(±2.5)	53.3	(±3.1)
Female	31.3	(±2.3)	62.2	(±3.5)
Race/Ethnicity				
White, non-Hispanic	29.7	(±2.4)	58.1	(±3.6)
Black, non-Hispanic	35.3	(±4.2)	58.9	(±6.7)
Hispanic	27.8	(±2.7)	52.8	(±4.9)
Asian	24.2	(±5.4)	70.9	(±14.2)
Total	29.9	(±1.9)	57.5	(±2.9)

* Students were considered to have never smoked if they answered “no” to whether they have tried or experimented with cigarette smoking, even one or two puffs.

† Smoked cigarettes on ≥ 1 of the previous 30 days.

§ Confidence interval.

TABLE 37. Percentage of middle school and high school students who live in a home in which someone else uses tobacco, by tobacco use status and state — state youth tobacco surveys, United States, 2001–2002

States	Someone else in home smoke cigarettes				Someone else in home use smokeless tobacco (SLT)*			
	Never smoked†		Currently smoke§		Never used SLT¶		Currently use SLT**	
	%	(95% CI††)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school								
Alabama – Spring 02	36.9	(±3.1)	70.1	(±6.4)	15.3	(±2.7)	59.8	(±10.4)
Connecticut – Spring 02	36.8	(±5.8)	67.2	(±9.5)	3.5	(±0.8)	37.5	§§
Delaware – Spring 02	38.6	(±2.8)	73.4	(±6.9)	4.1	(±0.9)	24.3	(±14.6)
Florida – Spring 01	31.4	(±2.6)	69.7	(±4.7)	5.8	(±1.2)	37.5	(±8.2)
Florida – Spring 02	31.4	(±1.5)	65.8	(±3.0)	6.1	(±0.7)	39.1	(±5.4)
Georgia – Fall 01	33.9	(±2.6)	70.2	(±6.5)	12.9	(±2.4)	46.9	(±10.0)
Idaho – Spring 01	24.2	(±4.6)	73.7	(±6.9)	10.8	(±1.7)	56.2	(±14.8)
Illinois – Spring 02	30.3	(±4.7)	58.0	(±7.4)	8.8	(±2.5)	37.0	§§
Iowa – Spring 02	34.9	(±4.5)	72.2	(±10.2)	11.9	(±2.5)	37.0	(±17.4)
Kansas – Fall 02	34.9	(±4.6)	69.5	(±8.7)	14.6	(±2.5)	37.0	§§
Kentucky – Spring 02	45.0	(±4.8)	75.7	(±6.0)	17.4	(±3.1)	50.6	(±8.2)
Louisiana – Spring 01	37.7	(±4.6)	66.7	(±5.0)	15.7	(±1.9)	51.7	(±7.2)
Maine – Spring 01	36.0	(±3.4)	72.0	(±4.8)	4.9	(±0.8)	43.7	(±7.8)
Maryland – Spring 02	33.5	(±1.3)	63.9	(±3.1)	8.5	(±0.8)	42.2	(±7.3)
Massachusetts – Spring 02	29.6	(±3.9)	63.0	(±10.3)	3.2	(±0.9)	37.0	§§
Michigan – Spring 01	NA†††		NA		8.6	(±2.1)	49.2	(±20.2)
Minnesota – Spring 02	31.2	(±3.4)	72.9	(±8.5)	9.4	(±1.1)	51.2	(±14.1)
Mississippi – Fall 02	41.1	(±3.0)	71.4	(±5.1)	NA		NA	
Nebraska – Spring 02	33.5	(±3.7)	79.3	(±5.4)	12.8	(±1.5)	38.5	(±15.0)
New Hampshire – Fall 01	35.0	(±3.9)	65.8	(±9.7)	5.2	(±1.4)	37.0	§§
New Jersey – Fall 01	33.8	(±4.4)	62.8	(±6.8)	NA		NA	
New York – Spring 02	32.7	(±4.7)	72.6	(±7.4)	NA		NA	
North Carolina – Spring 02	39.1	(±2.6)	67.1	(±6.7)	12.3	(±1.8)	50.9	(±10.5)
Ohio – Spring 02	35.9	(±4.2)	80.2	(±7.8)	8.9	(±2.2)	46.4	(±16.7)
Oklahoma – Spring 02	34.3	(±3.3)	70.1	(±8.2)	20.0	(±2.1)	46.5	(±12.1)
Pennsylvania – Spring 01	36.1	(±3.2)	68.3	(±4.5)	11.5	(±1.5)	45.5	(±5.9)
Rhode Island – Spring 01	33.5	(±4.0)	67.5	(±7.5)	5.2	(±1.2)	37.0	§§
South Dakota – Fall 01	30.0	(±4.6)	77.3	(±6.4)	16.0	(±2.5)	52.9	(±12.1)
Texas – Spring 01	32.0	(±3.1)	59.2	(±5.2)	10.3	(±1.4)	25.6	(±3.7)
Vermont – Spring 02	35.1	(±4.4)	73.3	(±5.7)	8.5	(±1.2)	50.6	(±15.3)
West Virginia – Spring 02	38.5	(±2.6)	74.3	(±2.0)	23.6	(±2.6)	54.2	(±4.1)
Wisconsin – Spring 02	36.5	(±4.9)	81.0	(±7.4)	11.5	(±2.7)	37.0	§§
Median	34.9		70.1		10.3		46.7	
High school								
Alabama	33.4	(±5.1)	54.5	(±6.2)	14.4	(±3.1)	38.7	(±6.9)
Connecticut – Spring 02	26.9	(±4.2)	48.6	(±6.4)	4.9	(±1.0)	30.5	(±14.5)
Delaware – Spring 02	36.6	(±4.7)	65.5	(±9.4)	4.1	(±1.2)	30.7	(±6.1)
Florida – Spring 01	25.5	(±2.7)	58.3	(±4.2)	4.9	(±1.0)	43.4	(±8.1)
Florida – Spring 02	26.7	(±2.1)	57.8	(±2.9)	5.1	(±1.0)	32.6	(±5.3)
Georgia – Fall 01	31.4	(±3.4)	56.5	(±5.2)	11.7	(±1.5)	48.1	(±7.6)
Illinois – Spring 02	24.7	(±4.8)	59.4	(±4.3)	4.6	(±2.1)	37.1	(±13.2)
Iowa – Spring 02	29.5	(±5.3)	49.8	(±5.2)	9.7	(±2.0)	28.3	(±10.3)
Kansas – Fall 02	27.6	(±4.2)	53.8	(±7.7)	11.5	(±2.1)	40.3	(±9.3)
Kentucky – Spring 02	37.5	(±4.0)	60.7	(±4.9)	15.5	(±2.4)	31.7	(±7.8)
Maryland – Spring 02	31.4	(±1.0)	56.0	(±1.6)	9.1	(±0.5)	42.1	(±3.3)
Massachusetts – Spring 02	29.6	(±3.8)	56.5	(±6.4)	1.9	(±0.7)	26.2	(±12.6)
Michigan – Spring 01	NA†††		NA		8.5	(±1.3)	41.5	(±7.7)
Mississippi – Fall 02	28.7	(±3.3)	58.9	(±4.6)	NA		NA	
Nebraska – Spring 02	24.9	(±4.0)	52.5	(±4.7)	11.8	(±1.3)	40.1	(±10.2)
New Hampshire – Fall 01	27.2	(±3.1)	59.2	(±5.6)	5.5	(±2.0)	40.0	(±15.5)
New Jersey – Fall 01	35.4	(±4.9)	53.9	(±3.6)	NA		NA	
New York – Spring 02	27.2	(±4.7)	56.9	(±5.9)	NA		NA	
North Carolina – Spring 02	34.2	(±2.7)	61.0	(±3.2)	11.2	(±1.6)	40.2	(±6.8)
Ohio – Spring 02	32.7	(±5.2)	65.3	(±4.9)	8.3	(±2.0)	22.7	(±11.6)
Oklahoma – Spring 02	28.2	(±3.4)	61.2	(±4.7)	15.8	(±3.0)	38.9	(±8.7)
Pennsylvania – Spring 01	30.4	(±3.9)	59.2	(±3.2)	8.8	(±1.6)	36.7	(±5.8)
Rhode Island – Spring 01	29.8	(±2.3)	56.7	(±5.2)	4.1	(±1.2)	27.1	(±15.6)
Texas – Spring 01	30.4	(±3.7)	52.1	(±3.6)	8.9	(±2.4)	24.5	(±4.3)
West Virginia – Spring 02	35.1	(±5.6)	66.3	(±6.3)	20.7	(±3.0)	40.2	(±7.9)
Wisconsin – Spring 02	27.8	(±3.4)	54.7	(±5.4)	8.3	(±2.9)	35.0	(±14.2)
Median	29.6		56.9		8.8		37.1	

* Question not asked on National Youth Tobacco Survey.

† Students were considered to have never smoked if they answered “no” to whether they have tried or experimented with cigarette smoking, even one or two puffs.

§ Smoked cigarettes on ≥1 of the previous 30 days.

¶ Students were considered never having used SLT if they answered “no” to whether they have used SLT.

** Used SLT on ≥1 of the previous 30 days.

†† Confidence interval.

§§ Sample size <35.

††† Question not asked.

TABLE 38. Percentage of middle school and high school students who practiced ways to say “no” to tobacco as part of school curriculum or who smoked cigarettes or used smokeless tobacco (SLT) on school property during the previous 30 days, by tobacco use status, sex, and race/ethnicity — National Youth Tobacco Survey, United States, 2002

Sex and Race/Ethnicity	Practiced ways to say “no” to tobacco		Smoked cigarettes on school property				Used SLT on school property			
			All students		Currently smoke*		All students		Currently use SLT†	
	%	(95% CI)§	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Middle school										
Sex										
Male	35.8	(±2.5)	2.9	(±0.6)	28.9	(±5.2)	2.7	(±0.6)	50.4	(±8.1)
Female	41.9	(±3.7)	2.4	(±0.6)	24.0	(±5.8)	1.0	(±0.4)	55.4	(±14.6)
Race/Ethnicity										
White, non-Hispanic	36.8	(±3.5)	2.2	(±0.5)	21.8	(±3.9)	1.8	(±0.5)	45.3	(±6.9)
Black, non-Hispanic	45.3	(±3.5)	3.1	(±1.1)	32.8	(±10.0)	1.5	(±0.8)	65.4	(±16.3)
Hispanic	41.6	(±3.5)	2.8	(±0.8)	32.1	(±6.4)	1.9	(±0.7)	64.8	(±15.4)
Asian	38.8	(±6.0)	3.5	(±2.7)	48.2	(±25.0)	3.4	(±2.7)	¶	
Total	38.8	(±2.9)	2.7	(±0.5)	26.7	(±4.0)	1.9	(±0.4)	52.0	(±7.3)
High school										
Sex										
Male	14.9	(±2.0)	9.4	(±1.5)	39.4	(±4.5)	6.5	(±1.3)	62.0	(±4.9)
Female	17.8	(±2.1)	6.7	(±1.2)	31.7	(±4.8)	0.4	(±0.2)	34.4	(±13.1)
Race/Ethnicity										
White, non-Hispanic	14.0	(±2.2)	8.8	(±1.4)	35.0	(±4.8)	4.2	(±0.9)	57.3	(±5.1)
Black, non-Hispanic	22.5	(±3.0)	4.8	(±1.3)	33.6	(±6.8)	1.1	(±0.6)	¶	
Hispanic	20.5	(±3.8)	7.9	(±1.7)	39.2	(±6.4)	2.5	(±1.0)	74.5	(±13.5)
Asian	21.5	(±6.0)	5.3	(±2.2)	43.8	(±15.8)	1.0	(±0.8)	¶	
Total	16.4	(±1.9)	8.1	(±1.1)	35.8	(±3.9)	3.5	(±0.7)	59.1	(±4.7)

* Smoked cigarettes on ≥ 1 of the previous 30 days.

† Used SLT on ≥ 1 of the previous 30 days.

§ Confidence interval.

¶ Sample size <35.

TABLE 39. Percentage of middle school and high school students who practiced ways to say “no” to tobacco as part of school curriculum or who smoked cigarettes or used smokeless tobacco (SLT) on school property during the previous 30 days, by smoking status and state — state youth tobacco surveys, United States, 2001–2002

State	Practiced ways to say “no” to tobacco		Smoked cigarettes on school property		Used SLT on school property					
			All students	Currently smoke cigarettes*	All students	Currently use SLT†				
							% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
Middle school										
Alabama – Spring 02	43.1	(±6.2)	5.0	(±1.5)	26.7	(±7.3)	5.3	(±2.2)	46.7	(±11.2)
Connecticut – Spring 02	44.1	(±6.4)	1.6	(±0.6)	22.9	(±9.2)	1.0	(±0.4)	52.8	(±14.9)
Delaware – Spring 02	42.6	(±3.9)	3.3	(±0.9)	28.4	(±6.7)	1.5	(±0.6)	57.7	(±16.2)
Florida – Spring 01	45.8	(±3.6)	3.3	(±0.9)	27.5	(±5.5)	1.9	(±0.6)	61.7	(±8.8)
Florida – Spring 02	47.8	(±2.6)	3.0	(±0.7)	26.3	(±3.0)	2.0	(±0.4)	51.1	(±5.2)
Georgia – Fall 01	38.7	(±3.3)	2.2	(±0.9)	15.8	(±6.7)	2.3	(±0.9)	41.0	(±8.3)
Idaho – Spring 01	47.2	(±6.4)	2.8	(±0.7)	27.0	(±5.0)	2.0	(±0.6)	48.9	(±15.4)
Illinois – Spring 02	48.8	(±8.6)	2.8	(±1.3)	28.1	(±9.9)	2.1	(±1.2)	49.2	(±18.5)
Iowa – Spring 02	41.8	(±6.3)	2.3	(±0.7)	27.7	(±9.2)	0.8	(±0.4)	26.2	(±14.5)
Kansas – Fall 02	30.2	(±5.4)	2.1	(±0.7)	25.7	(±10.1)	0.9	(±0.7)	20.9	(±15.2)
Kentucky – Spring 02	52.2	(±4.3)	4.1	(±1.5)	24.5	(±6.1)	5.3	(±1.8)	39.7	(±7.6)
Louisiana – Spring 01	44.6	(±4.2)	5.7	(±1.7)	28.1	(±6.4)	7.0	(±2.0)	52.7	(±7.3)
Maine – Spring 01	62.9	(±6.1)	2.0	(±0.6)	23.7	(±5.3)	NA [¶]		NA	
Maryland – Spring 02	64.7	(±1.4)	2.6	(±0.3)	33.8	(±3.6)	2.0	(±0.4)	51.9	(±6.1)
Massachusetts – Spring 02	40.0	(±7.4)	2.7	(±1.1)	35.0	(±8.0)	2.5	(±0.9)	**	
Michigan – Spring 01	46.6	(±6.3)	3.5	(±1.1)	31.0	(±8.9)	3.0	(±1.1)	71.8	(±16.0)
Minnesota – Spring 02	51.5	(±4.6)	2.8	(±0.8)	31.5	(±7.3)	1.6	(±0.4)	52.3	(±12.2)
Mississippi – Fall 02	NA		3.9	(±1.8)	22.9	(±7.5)	NA		NA	
Nebraska – Spring 02	50.5	(±3.7)	2.6	(±0.6)	28.4	(±6.2)	1.2	(±0.5)	35.0	(±14.1)
New Hampshire – Fall 01	23.5	(±3.4)	2.2	(±0.7)	41.3	(±8.8)	1.2	(±0.4)	**	
New Jersey – Fall 01	NA		2.0	(±0.8)	26	(±9.3)	NA		NA	
New York – Spring 02	39.1	(±9.6)	3.1	(±1.4)	36.6	(±12.5)	3.2	(±1.2)	61.2	(±21.8)
North Carolina – Spring 02	43.9	(±2.6)	3.3	(±1.1)	23.2	(±6.3)	2.8	(±1.0)	45.2	(±8.9)
Ohio – Spring 02	48.8	(±5.0)	3.0	(±1.2)	27.2	(±8.5)	1.9	(±1.0)	33.8	(±16.5)
Oklahoma – Spring 02	37.4	(±5.7)	3.9	(±1.3)	30.1	(±7.5)	2.4	(±0.8)	31.2	(±8.9)
Pennsylvania – Spring 01	44.6	(±4.1)	3.5	(±0.9)	24.0	(±4.5)	2.8	(±1.3)	49.7	(±17.0)
Rhode Island – Spring 01	45.9	(±8.2)	3.7	(±1.2)	35.6	(±9.6)	2.0	(±0.9)	**	
South Dakota – Fall 01	34.4	(±4.1)	3.3	(±1.7)	24.9	(±7.6)	5.1	(±2.9)	52.5	(±13.0)
Texas – Spring 01	43.1	(±2.5)	4.4	(±0.8)	34.0	(±4.2)	NA		NA	
Vermont – Spring 02	47.3	(±8.0)	4.1	(±1.3)	40.5	(±8.8)	2.0	(±0.6)	50.5	(±13.2)
West Virginia – Spring 02	50.7	(±2.9)	4.9	(±1.1)	27.7	(±4.5)	3.6	(±0.6)	37.8	(±4.4)
Wisconsin – Spring 02	50.2	(±9.4)	2.6	(±1.3)	26.2	(±8.4)	1.6	(±0.8)	**	
Median	45.2		3.1		27.6		2.0		49.5	
High school										
Spring 02	21.1	(±3.7)	9.0	(±1.9)	32.9	(±5.94)	6.3	(±1.6)	53.4	(±9.8)
Connecticut – Spring 02	18.3	(±3.1)	10.8	(±1.6)	47.4	(±5.4)	2.8	(±0.7)	59.6	(±12.5)
Delaware – Spring 02	19.3	(±6.3)	11.2	(±1.4)	40.5	(±3.4)	3.5	(±0.9)	61.8	(±7.3)
Florida – Spring 01	22.2	(±2.3)	5.9	(±0.9)	31.7	(±4.2)	3.6	(±0.8)	68.0	(±7.0)
Florida – Spring 02	23.9	(±1.9)	4.9	(±0.7)	27.8	(±2.8)	3.1	(±0.5)	62.3	(±5.0)
Georgia – Fall 01	20.2	(±2.6)	8.2	(±1.1)	32.6	(±5.0)	6.9	(±1.8)	62.3	(±7.8)
Illinois – Spring 02	21.3	(±3.6)	10.0	(±2.9)	32.4	(±5.9)	4.5	(±2.3)	63.3	(±9.7)
Iowa – Spring 02	14.8	(±3.3)	10.1	(±2.0)	34.6	(±4.7)	3.9	(±1.2)	43.0	(±10.8)
Kansas – Fall 02	10.2	(±2.7)	6.5	(±1.8)	26.7	(±7.5)	4.5	(±1.6)	51.5	(±9.7)
Kentucky – Spring 02	22.2	(±5.4)	16.4	(±2.0)	47.2	(±6.5)	7.9	(±1.7)	54.3	(±8.8)
Maryland – Spring 02	32.1	(±1.0)	10.0	(±0.5)	44.2	(±1.5)	5.6	(±0.4)	65.9	(±2.3)
Massachusetts – Spring 02	11.7	(±2.3)	9.3	(±2.2)	42.9	(±6.2)	3.7	(±1.3)	57.8	(±17.8)
Michigan – Spring 01	15.5	(±2.4)	11.6	(±1.1)	39.9	(±3.6)	5.2	(±1.1)	60.5	(±9.8)
Mississippi – Fall 02	NA		8.4	(±1.5)	31.7	(±5.0)	NA		NA	
Nebraska – Spring 02	16.6	(±2.8)	10.2	(±1.6)	35.2	(±4.5)	4.4	(±1.0)	48.2	(±8.5)
New Hampshire – Fall 01	9.4	(±1.9)	9.9	(±1.6)	35.9	(±4.8)	3.0	(±1.2)	57.8	(±14.7)
New Jersey – Fall 01	NA		10.5	(±1.6)	41.2	(±5.7)	NA		NA	
New York – Spring 02	14.9	(±3.9)	10.1	(±2.4)	45.4	(±9.4)	4.5	(±2.4)	52.7	(±14.3)
North Carolina – Spring 02	16.6	(±2.0)	13.0	(±2.3)	44.0	(±4.1)	6.6	(±0.9)	61.1	(±5.7)
Ohio – Spring 02	18.3	(±3.3)	8.5	(±2.4)	32.3	(±8.2)	3.5	(±1.7)	40.7	(±12.9)
Oklahoma – Spring 02	10.1	(±1.9)	7.5	(±1.8)	30.4	(±5.4)	6.0	(±1.8)	52.5	(±10.6)
Pennsylvania – Spring 01	15.3	(±2.2)	10.4	(±1.1)	36.1	(±3.8)	5.3	(±1.0)	63.0	(±4.1)
Rhode Island – Spring 01	15.8	(±3.6)	14.9	(±2.4)	57.0	(±6.0)	4.4	(±1.3)	83.0	(±10.8)
Texas – Spring 01	20.7	(±1.8)	8.3	(±1.3)	30.4	(±4.0)	NA		NA	
West Virginia – Spring 02	24.4	(±4.3)	13.4	(±2.7)	37.9	(±7.6)	8.1	(±1.9)	55.5	(±9.0)
Wisconsin – Spring 02	13.7	(±3.8)	10.2	(±2.7)	36.5	(±7.7)	4.7	(±1.8)	53.2	(±7.1)
Median	17.5		10.1		36.0		4.5		57.8	

* Smoked cigarettes on ≥1 of the previous 30 days.

† Used SLT on ≥1 of the previous 30 days.

‡ Confidence interval.

¶ Question not asked.

** Sample size <35.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop E-90, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone 888-232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

All *MMWR* references are available on the Internet at <http://www.cdc.gov/mmwr>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.