

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	6,068	100.0	571.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,520	25.0	143.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	917	15.1	86.3
3	Sudden infant death syndrome (R95)	337	5.6	31.7
4	Newborn affected by maternal complications of pregnancy (P01)	307	5.1	28.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	240	4.0	22.6
6	Accidents (unintentional injuries) (V01-X59)	167	2.8	15.7
7	Bacterial sepsis of newborn (P36)	165	2.7	15.5
8	Respiratory distress of newborn (P22)	148	2.4	13.9
9	Neonatal hemorrhage (P50-P52,P54)	144	2.4	13.5
10	Diseases of the circulatory system (I00-I99)	139	2.3	13.1
11	Necrotizing enterocolitis of newborn (P77)	108	1.8	10.2
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	94	1.5	8.8
13	Atelectasis (P28.0-P28.1)	86	1.4	8.1
14	Assault (homicide) (*U01,X85-Y09)	81	1.3	7.6
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	80	1.3	7.5
...	All other causes (Residual)	1,535	25.3	144.4
Infants, Hispanic, male				
...	All causes	3,343	100.0	616.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	787	23.5	145.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	503	15.0	92.8
3	Sudden infant death syndrome (R95)	197	5.9	36.3
4	Newborn affected by maternal complications of pregnancy (P01)	178	5.3	32.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	122	3.6	22.5
6	Accidents (unintentional injuries) (V01-X59)	92	2.8	17.0
7	Bacterial sepsis of newborn (P36)	91	2.7	16.8
8	Neonatal hemorrhage (P50-P52,P54)	89	2.7	16.4
9	Respiratory distress of newborn (P22)	88	2.6	16.2
10	Diseases of the circulatory system (I00-I99)	73	2.2	13.5
11	Necrotizing enterocolitis of newborn (P77)	64	1.9	11.8
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	60	1.8	11.1
13	Atelectasis (P28.0-P28.1)	56	1.7	10.3
14	Assault (homicide) (*U01,X85-Y09)	44	1.3	8.1
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	43	1.3	7.9
...	All other causes (Residual)	856	25.6	157.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,725	100.0	523.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	733	26.9	140.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	414	15.2	79.5
3	Sudden infant death syndrome (R95)	140	5.1	26.9
4	Newborn affected by maternal complications of pregnancy (P01)	129	4.7	24.8
5	Newborn affected by complications of placenta, cord and membranes (P02)	118	4.3	22.7
6	Accidents (unintentional injuries) (V01-X59)	75	2.8	14.4
7	Bacterial sepsis of newborn (P36)	74	2.7	14.2
8	Diseases of the circulatory system (I00-I99)	66	2.4	12.7
9	Respiratory distress of newborn (P22)	60	2.2	11.5
10	Neonatal hemorrhage (P50-P52,P54)	55	2.0	10.6
11	Necrotizing enterocolitis of newborn (P77)	44	1.6	8.5
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	37	1.4	7.1
12	Assault (homicide) (*U01,X85-Y09)	37	1.4	7.1
14	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	34	1.2	6.5
15	Septicemia (A40-A41)	30	1.1	5.8
15	Atelectasis (P28.0-P28.1)	30	1.1	5.8
...	All other causes (Residual)	649	23.8	124.7
Infants, Non-Hispanic, white, both sexes				
...	All causes	12,998	100.0	562.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,871	22.1	124.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,829	14.1	79.2
3	Sudden infant death syndrome (R95)	1,302	10.0	56.4
4	Newborn affected by maternal complications of pregnancy (P01)	776	6.0	33.6
5	Accidents (unintentional injuries) (V01-X59)	641	4.9	27.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	508	3.9	22.0
7	Bacterial sepsis of newborn (P36)	353	2.7	15.3
8	Respiratory distress of newborn (P22)	337	2.6	14.6
9	Neonatal hemorrhage (P50-P52,P54)	284	2.2	12.3
10	Diseases of the circulatory system (I00-I99)	265	2.0	11.5
11	Necrotizing enterocolitis of newborn (P77)	207	1.6	9.0
12	Atelectasis (P28.0-P28.1)	173	1.3	7.5
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	164	1.3	7.1
14	Assault (homicide) (*U01,X85-Y09)	150	1.2	6.5
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	134	1.0	5.8
...	All other causes (Residual)	3,004	23.1	130.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, white, male				
...	All causes	7,340	100.0	619.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,559	21.2	131.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,003	13.7	84.7
3	Sudden infant death syndrome (R95)	767	10.4	64.7
4	Newborn affected by maternal complications of pregnancy (P01)	439	6.0	37.1
5	Accidents (unintentional injuries) (V01-X59)	365	5.0	30.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	279	3.8	23.6
7	Bacterial sepsis of newborn (P36)	211	2.9	17.8
8	Respiratory distress of newborn (P22)	198	2.7	16.7
9	Neonatal hemorrhage (P50-P52,P54)	171	2.3	14.4
10	Diseases of the circulatory system (I00-I99)	151	2.1	12.7
11	Necrotizing enterocolitis of newborn (P77)	115	1.6	9.7
12	Atelectasis (P28.0-P28.1)	99	1.3	8.4
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	97	1.3	8.2
14	Assault (homicide) (*U01,X85-Y09)	86	1.2	7.3
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	74	1.0	6.2
...	All other causes (Residual)	1,726	23.5	145.7
Infants, Non-Hispanic, white, female				
...	All causes	5,658	100.0	502.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,312	23.2	116.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	826	14.6	73.4
3	Sudden infant death syndrome (R95)	535	9.5	47.5
4	Newborn affected by maternal complications of pregnancy (P01)	337	6.0	29.9
5	Accidents (unintentional injuries) (V01-X59)	276	4.9	24.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	229	4.0	20.3
7	Bacterial sepsis of newborn (P36)	142	2.5	12.6
8	Respiratory distress of newborn (P22)	139	2.5	12.3
9	Diseases of the circulatory system (I00-I99)	114	2.0	10.1
10	Neonatal hemorrhage (P50-P52,P54)	113	2.0	10.0
11	Necrotizing enterocolitis of newborn (P77)	92	1.6	8.2
12	Atelectasis (P28.0-P28.1)	74	1.3	6.6
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	67	1.2	6.0
14	Assault (homicide) (*U01,X85-Y09)	64	1.1	5.7
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	60	1.1	5.3
...	All other causes (Residual)	1,278	22.6	113.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, both sexes				
...	All causes	8,629	100.0	1,375.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,861	21.6	296.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,096	12.7	174.7
3	Sudden infant death syndrome (R95)	714	8.3	113.8
4	Newborn affected by maternal complications of pregnancy (P01)	601	7.0	95.8
5	Accidents (unintentional injuries) (V01-X59)	422	4.9	67.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	333	3.9	53.1
7	Respiratory distress of newborn (P22)	270	3.1	43.0
8	Bacterial sepsis of newborn (P36)	259	3.0	41.3
9	Necrotizing enterocolitis of newborn (P77)	212	2.5	33.8
10	Diseases of the circulatory system (I00-I99)	183	2.1	29.2
11	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	165	1.9	26.3
12	Neonatal hemorrhage (P50-P52,P54)	138	1.6	22.0
13	Septicemia (A40-A41)	110	1.3	17.5
14	Assault (homicide) (*U01,X85-Y09)	106	1.2	16.9
15	Chronic respiratory disease originating in the perinatal period (P27)	101	1.2	16.1
...	All other causes (Residual)	2,058	23.8	328.1
Infants, Non-Hispanic, black, male				
...	All causes	4,794	100.0	1,504.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,035	21.6	324.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	564	11.8	177.0
3	Sudden infant death syndrome (R95)	398	8.3	124.9
4	Newborn affected by maternal complications of pregnancy (P01)	348	7.3	109.2
5	Accidents (unintentional injuries) (V01-X59)	219	4.6	68.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	190	4.0	59.6
7	Respiratory distress of newborn (P22)	165	3.4	51.8
8	Bacterial sepsis of newborn (P36)	136	2.8	42.7
9	Necrotizing enterocolitis of newborn (P77)	124	2.6	38.9
10	Diseases of the circulatory system (I00-I99)	115	2.4	36.1
11	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	101	2.1	31.7
12	Neonatal hemorrhage (P50-P52,P54)	88	1.8	27.6
13	Assault (homicide) (*U01,X85-Y09)	67	1.4	21.0
14	Atelectasis (P28.0-P28.1)	64	1.3	20.1
15	Septicemia (A40-A41)	60	1.3	18.8
...	All other causes (Residual)	1,120	23.4	351.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, female				
...	All causes	3,835	100.0	1,243.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	826	21.5	267.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	532	13.9	172.4
3	Sudden infant death syndrome (R95)	316	8.2	102.4
4	Newborn affected by maternal complications of pregnancy (P01)	253	6.6	82.0
5	Accidents (unintentional injuries) (V01-X59)	203	5.3	65.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	143	3.7	46.4
7	Bacterial sepsis of newborn (P36)	123	3.2	39.9
8	Respiratory distress of newborn (P22)	105	2.7	34.0
9	Necrotizing enterocolitis of newborn (P77)	88	2.3	28.5
10	Diseases of the circulatory system (I00-I99)	68	1.8	22.0
11	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	64	1.7	20.7
12	Septicemia (A40-A41)	50	1.3	16.2
12	Neonatal hemorrhage (P50-P52,P54)	50	1.3	16.2
14	Chronic respiratory disease originating in the perinatal period (P27)	47	1.2	15.2
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	46	1.2	14.9
...	All other causes (Residual)	921	24.0	298.5
Neonates, Hispanic, both sexes				
...	All causes	4,060	100.0	382.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,066	26.3	100.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	889	21.9	83.6
3	Newborn affected by maternal complications of pregnancy (P01)	305	7.5	28.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	237	5.8	22.3
5	Bacterial sepsis of newborn (P36)	155	3.8	14.6
6	Respiratory distress of newborn (P22)	145	3.6	13.6
7	Neonatal hemorrhage (P50-P52,P54)	143	3.5	13.5
8	Necrotizing enterocolitis of newborn (P77)	95	2.3	8.9
9	Atelectasis (P28.0-P28.1)	84	2.1	7.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	76	1.9	7.2
11	Hydrops fetalis not due to hemolytic disease (P83.2)	51	1.3	4.8
12	Pulmonary hemorrhage originating in the perinatal period (P26)	31	0.8	2.9
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	29	0.7	2.7
14	Newborn affected by other complications of labor and delivery (P03)	27	0.7	2.5
14	Hematological disorders (P60-P61)	27	0.7	2.5
...	All other causes (Residual)	700	17.2	65.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	2,233	100.0	411.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	558	25.0	102.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	486	21.8	89.6
3	Newborn affected by maternal complications of pregnancy (P01)	177	7.9	32.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	120	5.4	22.1
5	Neonatal hemorrhage (P50-P52,P54)	88	3.9	16.2
6	Respiratory distress of newborn (P22)	87	3.9	16.0
7	Bacterial sepsis of newborn (P36)	85	3.8	15.7
8	Necrotizing enterocolitis of newborn (P77)	57	2.6	10.5
9	Atelectasis (P28.0-P28.1)	54	2.4	10.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	40	1.8	7.4
11	Hydrops fetalis not due to hemolytic disease (P83.2)	23	1.0	4.2
12	Newborn affected by other complications of labor and delivery (P03)	20	0.9	3.7
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	18	0.8	*
13	Pulmonary hemorrhage originating in the perinatal period (P26)	18	0.8	*
15	Hematological disorders (P60-P61)	16	0.7	*
...	All other causes (Residual)	386	17.3	71.2
Neonates, Hispanic, female				
...	All causes	1,827	100.0	350.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	508	27.8	97.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	403	22.1	77.4
3	Newborn affected by maternal complications of pregnancy (P01)	128	7.0	24.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	117	6.4	22.5
5	Bacterial sepsis of newborn (P36)	70	3.8	13.4
6	Respiratory distress of newborn (P22)	58	3.2	11.1
7	Neonatal hemorrhage (P50-P52,P54)	55	3.0	10.6
8	Necrotizing enterocolitis of newborn (P77)	38	2.1	7.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	36	2.0	6.9
10	Atelectasis (P28.0-P28.1)	30	1.6	5.8
11	Hydrops fetalis not due to hemolytic disease (P83.2)	28	1.5	5.4
12	Pulmonary hemorrhage originating in the perinatal period (P26)	13	0.7	*
13	Neonatal aspiration syndromes (P24)	11	0.6	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	11	0.6	*
13	Hematological disorders (P60-P61)	11	0.6	*
...	All other causes (Residual)	310	17.0	59.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, both sexes				
...	All causes	8,417	100.0	364.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,074	24.6	89.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,798	21.4	77.8
3	Newborn affected by maternal complications of pregnancy (P01)	766	9.1	33.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	502	6.0	21.7
5	Bacterial sepsis of newborn (P36)	337	4.0	14.6
6	Respiratory distress of newborn (P22)	330	3.9	14.3
7	Neonatal hemorrhage (P50-P52,P54)	278	3.3	12.0
8	Necrotizing enterocolitis of newborn (P77)	182	2.2	7.9
9	Atelectasis (P28.0-P28.1)	163	1.9	7.1
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	153	1.8	6.6
11	Sudden infant death syndrome (R95)	104	1.2	4.5
12	Hydrops fetalis not due to hemolytic disease (P83.2)	84	1.0	3.6
13	Diseases of the circulatory system (I00-I99)	67	0.8	2.9
14	Accidents (unintentional injuries) (V01-X59)	66	0.8	2.9
15	Pulmonary hemorrhage originating in the perinatal period (P26)	60	0.7	2.6
...	All other causes (Residual)	1,453	17.3	62.9
Neonates, Non-Hispanic, white, male				
...	All causes	4,693	100.0	396.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,137	24.2	96.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	985	21.0	83.1
3	Newborn affected by maternal complications of pregnancy (P01)	434	9.2	36.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	275	5.9	23.2
5	Bacterial sepsis of newborn (P36)	200	4.3	16.9
6	Respiratory distress of newborn (P22)	195	4.2	16.5
7	Neonatal hemorrhage (P50-P52,P54)	169	3.6	14.3
8	Necrotizing enterocolitis of newborn (P77)	98	2.1	8.3
9	Atelectasis (P28.0-P28.1)	95	2.0	8.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	91	1.9	7.7
11	Sudden infant death syndrome (R95)	50	1.1	4.2
12	Diseases of the circulatory system (I00-I99)	41	0.9	3.5
13	Hydrops fetalis not due to hemolytic disease (P83.2)	40	0.9	3.4
14	Newborn affected by other complications of labor and delivery (P03)	38	0.8	3.2
14	Pulmonary hemorrhage originating in the perinatal period (P26)	38	0.8	3.2
...	All other causes (Residual)	807	17.2	68.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, female				
...	All causes	3,724	100.0	330.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	937	25.2	83.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	813	21.8	72.2
3	Newborn affected by maternal complications of pregnancy (P01)	332	8.9	29.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	227	6.1	20.2
5	Bacterial sepsis of newborn (P36)	137	3.7	12.2
6	Respiratory distress of newborn (P22)	135	3.6	12.0
7	Neonatal hemorrhage (P50-P52,P54)	109	2.9	9.7
8	Necrotizing enterocolitis of newborn (P77)	84	2.3	7.5
9	Atelectasis (P28.0-P28.1)	68	1.8	6.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	62	1.7	5.5
11	Sudden infant death syndrome (R95)	54	1.5	4.8
12	Hydrops fetalis not due to hemolytic disease (P83.2)	44	1.2	3.9
13	Accidents (unintentional injuries) (V01-X59)	32	0.9	2.8
14	Diseases of the circulatory system (I00-I99)	26	0.7	2.3
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	26	0.7	2.3
...	All other causes (Residual)	638	17.1	56.7
Neonates, Non-Hispanic, black, both sexes				
...	All causes	5,626	100.0	897.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,806	32.1	288.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	708	12.6	112.9
3	Newborn affected by maternal complications of pregnancy (P01)	592	10.5	94.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	329	5.8	52.5
5	Respiratory distress of newborn (P22)	262	4.7	41.8
6	Bacterial sepsis of newborn (P36)	244	4.3	38.9
7	Necrotizing enterocolitis of newborn (P77)	185	3.3	29.5
8	Neonatal hemorrhage (P50-P52,P54)	134	2.4	21.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	93	1.7	14.8
9	Atelectasis (P28.0-P28.1)	93	1.7	14.8
11	Sudden infant death syndrome (R95)	73	1.3	11.6
12	Pulmonary hemorrhage originating in the perinatal period (P26)	65	1.2	10.4
13	Accidents (unintentional injuries) (V01-X59)	41	0.7	6.5
14	Diseases of the circulatory system (I00-I99)	34	0.6	5.4
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	34	0.6	5.4
...	All other causes (Residual)	933	16.6	148.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, black, male				
...	All causes	3,131	100.0	982.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,002	32.0	314.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	386	12.3	121.1
3	Newborn affected by maternal complications of pregnancy (P01)	340	10.9	106.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	186	5.9	58.4
5	Respiratory distress of newborn (P22)	158	5.0	49.6
6	Bacterial sepsis of newborn (P36)	129	4.1	40.5
7	Necrotizing enterocolitis of newborn (P77)	111	3.5	34.8
8	Neonatal hemorrhage (P50-P52,P54)	85	2.7	26.7
9	Atelectasis (P28.0-P28.1)	62	2.0	19.5
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	48	1.5	15.1
11	Sudden infant death syndrome (R95)	35	1.1	11.0
12	Pulmonary hemorrhage originating in the perinatal period (P26)	31	1.0	9.7
13	Diseases of the circulatory system (I00-I99)	22	0.7	6.9
14	Newborn affected by other complications of labor and delivery (P03)	21	0.7	6.6
15	Accidents (unintentional injuries) (V01-X59)	18	0.6	*
...	All other causes (Residual)	497	15.9	155.9
Neonates, Non-Hispanic, black, female				
...	All causes	2,495	100.0	808.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	804	32.2	260.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	322	12.9	104.4
3	Newborn affected by maternal complications of pregnancy (P01)	252	10.1	81.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	143	5.7	46.4
5	Bacterial sepsis of newborn (P36)	115	4.6	37.3
6	Respiratory distress of newborn (P22)	104	4.2	33.7
7	Necrotizing enterocolitis of newborn (P77)	74	3.0	24.0
8	Neonatal hemorrhage (P50-P52,P54)	49	2.0	15.9
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	45	1.8	14.6
10	Sudden infant death syndrome (R95)	38	1.5	12.3
11	Pulmonary hemorrhage originating in the perinatal period (P26)	34	1.4	11.0
12	Atelectasis (P28.0-P28.1)	31	1.2	10.0
13	Accidents (unintentional injuries) (V01-X59)	23	0.9	7.5
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	18	0.7	*
15	Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	15	0.6	*
15	Slow fetal growth and fetal malnutrition (P05)	15	0.6	*
...	All other causes (Residual)	413	16.6	133.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	2,008	100.0	188.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	454	22.6	42.7
2	Sudden infant death syndrome (R95)	314	15.6	29.5
3	Accidents (unintentional injuries) (V01-X59)	154	7.7	14.5
4	Diseases of the circulatory system (I00-I99)	117	5.8	11.0
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	92	4.6	8.7
6	Assault (homicide) (*U01,X85-Y09)	74	3.7	7.0
7	Septicemia (A40-A41)	65	3.2	6.1
8	Influenza and pneumonia (J09-J18)	46	2.3	4.3
9	Chronic respiratory disease originating in the perinatal period (P27)	38	1.9	3.6
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	28	1.4	2.6
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	26	1.3	2.4
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.9	*
13	Malignant neoplasms (C00-C97)	16	0.8	*
14	Meningitis (G00,G03)	14	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	14	0.7	*
...	All other causes (Residual)	537	26.7	50.5
Postneonates, Hispanic, male				
...	All causes	1,110	100.0	204.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	20.6	42.2
2	Sudden infant death syndrome (R95)	183	16.5	33.8
3	Accidents (unintentional injuries) (V01-X59)	86	7.7	15.9
4	Diseases of the circulatory system (I00-I99)	61	5.5	11.3
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	58	5.2	10.7
6	Assault (homicide) (*U01,X85-Y09)	39	3.5	7.2
7	Septicemia (A40-A41)	35	3.2	6.5
8	Influenza and pneumonia (J09-J18)	28	2.5	5.2
9	Chronic respiratory disease originating in the perinatal period (P27)	18	1.6	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.5	*
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.8	*
11	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.8	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	0.8	*
14	Necrotizing enterocolitis of newborn (P77)	7	0.6	*
15	Malignant neoplasms (C00-C97)	6	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.5	*
15	Bacterial sepsis of newborn (P36)	6	0.5	*
...	All other causes (Residual)	298	26.8	55.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	898	100.0	172.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	25.1	43.2
2	Sudden infant death syndrome (R95)	131	14.6	25.2
3	Accidents (unintentional injuries) (V01-X59)	68	7.6	13.1
4	Diseases of the circulatory system (I00-I99)	56	6.2	10.8
5	Assault (homicide) (*U01,X85-Y09)	35	3.9	6.7
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	34	3.8	6.5
7	Septicemia (A40-A41)	30	3.3	5.8
8	Chronic respiratory disease originating in the perinatal period (P27)	20	2.2	3.8
9	Influenza and pneumonia (J09-J18)	18	2.0	*
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	17	1.9	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	11	1.2	*
12	Malignant neoplasms (C00-C97)	10	1.1	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	1.1	*
14	Meningitis (G00,G03)	9	1.0	*
15	Necrotizing enterocolitis of newborn (P77)	6	0.7	*
...	All other causes (Residual)	218	24.3	41.9
Postneonates, Non-Hispanic, white, both sexes				
...	All causes	4,581	100.0	198.3
1	Sudden infant death syndrome (R95)	1,198	26.2	51.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	797	17.4	34.5
3	Accidents (unintentional injuries) (V01-X59)	575	12.6	24.9
4	Diseases of the circulatory system (I00-I99)	198	4.3	8.6
5	Assault (homicide) (*U01,X85-Y09)	127	2.8	5.5
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	115	2.5	5.0
7	Septicemia (A40-A41)	98	2.1	4.2
8	Influenza and pneumonia (J09-J18)	88	1.9	3.8
9	Chronic respiratory disease originating in the perinatal period (P27)	82	1.8	3.5
10	Malignant neoplasms (C00-C97)	35	0.8	1.5
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	33	0.7	1.4
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	31	0.7	1.3
13	Meningitis (G00,G03)	28	0.6	1.2
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	27	0.6	1.2
15	Necrotizing enterocolitis of newborn (P77)	25	0.5	1.1
...	All other causes (Residual)	1,124	24.5	48.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, white, male				
...	All causes	2,647	100.0	223.4
1	Sudden infant death syndrome (R95)	717	27.1	60.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	422	15.9	35.6
3	Accidents (unintentional injuries) (V01-X59)	331	12.5	27.9
4	Diseases of the circulatory system (I00-I99)	110	4.2	9.3
5	Assault (homicide) (*U01,X85-Y09)	73	2.8	6.2
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	63	2.4	5.3
7	Influenza and pneumonia (J09-J18)	57	2.2	4.8
8	Chronic respiratory disease originating in the perinatal period (P27)	54	2.0	4.6
9	Septicemia (A40-A41)	52	2.0	4.4
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	21	0.8	1.8
11	Malignant neoplasms (C00-C97)	20	0.8	1.7
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	18	0.7	*
13	Necrotizing enterocolitis of newborn (P77)	17	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.5	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	12	0.5	*
...	All other causes (Residual)	667	25.2	56.3
Postneonates, Non-Hispanic, white, female				
...	All causes	1,934	100.0	171.8
1	Sudden infant death syndrome (R95)	481	24.9	42.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	375	19.4	33.3
3	Accidents (unintentional injuries) (V01-X59)	244	12.6	21.7
4	Diseases of the circulatory system (I00-I99)	88	4.6	7.8
5	Assault (homicide) (*U01,X85-Y09)	54	2.8	4.8
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	52	2.7	4.6
7	Septicemia (A40-A41)	46	2.4	4.1
8	Influenza and pneumonia (J09-J18)	31	1.6	2.8
9	Chronic respiratory disease originating in the perinatal period (P27)	28	1.4	2.5
10	Meningitis (G00,G03)	18	0.9	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	17	0.9	*
12	Malignant neoplasms (C00-C97)	15	0.8	*
13	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	13	0.7	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	12	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
...	All other causes (Residual)	449	23.2	39.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, both sexes				
...	All causes	3,003	100.0	478.8
1	Sudden infant death syndrome (R95)	641	21.3	102.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	388	12.9	61.9
3	Accidents (unintentional injuries) (V01-X59)	381	12.7	60.7
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	156	5.2	24.9
5	Diseases of the circulatory system (I00-I99)	149	5.0	23.8
6	Septicemia (A40-A41)	110	3.7	17.5
7	Assault (homicide) (*U01,X85-Y09)	96	3.2	15.3
8	Chronic respiratory disease originating in the perinatal period (P27)	88	2.9	14.0
9	Influenza and pneumonia (J09-J18)	74	2.5	11.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	55	1.8	8.8
11	Anoxic brain damage, not elsewhere classified (G93.1)	30	1.0	4.8
12	Necrotizing enterocolitis of newborn (P77)	27	0.9	4.3
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.6	*
14	Meningitis (G00,G03)	16	0.5	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	15	0.5	*
15	Bacterial sepsis of newborn (P36)	15	0.5	*
...	All other causes (Residual)	743	24.7	118.5
Postneonates, Non-Hispanic, black, male				
...	All causes	1,663	100.0	521.8
1	Sudden infant death syndrome (R95)	363	21.8	113.9
2	Accidents (unintentional injuries) (V01-X59)	201	12.1	63.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	178	10.7	55.9
4	Diseases of the circulatory system (I00-I99)	93	5.6	29.2
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	93	5.6	29.2
6	Assault (homicide) (*U01,X85-Y09)	62	3.7	19.5
7	Septicemia (A40-A41)	60	3.6	18.8
8	Chronic respiratory disease originating in the perinatal period (P27)	47	2.8	14.7
9	Influenza and pneumonia (J09-J18)	42	2.5	13.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	33	2.0	10.4
11	Anoxic brain damage, not elsewhere classified (G93.1)	17	1.0	*
12	Necrotizing enterocolitis of newborn (P77)	13	0.8	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	12	0.7	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	9	0.5	*
15	Meningitis (G00,G03)	8	0.5	*
15	Newborn affected by maternal complications of pregnancy (P01)	8	0.5	*
...	All other causes (Residual)	424	25.5	133.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2007

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, female				
...	All causes	1,340	100.0	434.4
1	Sudden infant death syndrome (R95)	278	20.7	90.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	15.7	68.1
3	Accidents (unintentional injuries) (V01-X59)	180	13.4	58.3
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	63	4.7	20.4
5	Diseases of the circulatory system (I00-I99)	56	4.2	18.2
6	Septicemia (A40-A41)	50	3.7	16.2
7	Chronic respiratory disease originating in the perinatal period (P27)	41	3.1	13.3
8	Assault (homicide) (*U01,X85-Y09)	34	2.5	11.0
9	Influenza and pneumonia (J09-J18)	32	2.4	10.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	1.6	7.1
11	Necrotizing enterocolitis of newborn (P77)	14	1.0	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	13	1.0	*
13	Meningitis (G00,G03)	8	0.6	*
13	Bacterial sepsis of newborn (P36)	8	0.6	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	7	0.5	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	7	0.5	*
...	All other causes (Residual)	317	23.7	102.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable