Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All races 3/, both sexes										
2002	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2
1999	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8
All races 3/, male										
2002	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4
1999	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6
All races 3/, female										
2002	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2
1999	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4
White, both sexes										
2002	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000	900.2	610.6	29.2	14.8	19.2	64.5	85.1	81.0	99.0	144.0
1999	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5
White, male										
2002	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000	887.8	667.6	32.6	16.5	23.1	88.1	124.6	113.9	133.6	188.2
1999	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
White, female										
2002	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000	912.3	550.5	25.5	13.0	15.2	39.3	43.0	46.5	63.0	99.2
1999	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5
Black, both sexes										
2002	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000	781.1	1,426.1	49.9	21.8	26.6	87.4	155.0	168.4	206.9	283.1
1999	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5
Black, male										
2002	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000	834.1	1,567.6	54.5	24.5	32.1	130.1	240.6	246.8	275.0	361.6
1999	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8
Black, female										
2002	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000	733.0	1,279.8	45.3	19.0	21.0	43.7	73.9	97.3	145.6	213.4
1999	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Salmonella infections (A01-A02)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Salmonella infections (A01-A02)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Shigellosis and amebiasis (A03,A06)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Shigellosis and amebiasis (A03,A06)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Certain other intestinal infections (A04,A07-A09)										
All races 3/, both sexes										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
White, male										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Certain other intestinal infections (A04,A07-A09)										
White, female										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Tuberculosis (A16-A19)										
All races 3/, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Tuberculosis (A16-A19)										
White, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.9	*	*	*	*	*	*	*	*	*
1999	0.9	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Respiratory tuberculosis (A16)										
All races 3/, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Respiratory tuberculosis (A16)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.6	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*
2000	0.8	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other tuberculosis (A17-A19)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other tuberculosis (A17-A19)										
White, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Whooping cough (A37)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Whooping cough (A37)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Scarlet fever and erysipelas (A38,A46)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Scarlet fever and erysipelas (A38,A46)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningococcal infection (A39)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	*	0.2	*	*	*	*
2000	0.1	0.6	*	*	*	0.2	0.1	*	*	*
1999	0.1	0.8	0.2	*	*	0.1	*	*	*	*
All races 3/, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	0.2	*	*	*	*
2000	0.1	*	*	*	*	0.2	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	0.2	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	0.2	*	*	*	*
2000	0.1	*	*	*	*	0.2	*	*	*	*
1999	0.1	1.0	0.2	*	*	0.1	*	*	*	*
White, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	0.2	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningococcal infection (A39)										
White, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
All races 3/, both sexes										
2002	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4
1999	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4
All races 3/, male										
2002	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4
1999	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5
All races 3/, female										
2002	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4
1999	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3
White, both sexes										
2002	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000	10.8	5.7	0.6	0.2	0.1	0.2	0.2	0.4	0.6	1.1
1999	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0
White, male										
2002	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000	9.4	5.8	0.6	*	*	*	0.3	0.4	0.7	1.2
1999	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
White, female										
2002	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000	12.1	5.6	0.5	*	*	*	*	0.4	0.5	1.0
1999	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9
Black, both sexes										
2002	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000	16.2	16.4	1.0	*	*	*	0.8	1.5	2.5	3.7
1999	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7
Black, male										
2002	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000	14.9	16.0	*	*	*	*	*	2.1	2.5	3.5
1999	15.3	19.8	*	*	*	*	*	*	2.1	4.0
Black, female										
2002	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000	17.4	16.9	*	*	*	*	*	*	2.5	3.9
1999	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Syphilis (A50-A53)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Syphilis (A50-A53)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute poliomyelitis (A80)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute poliomyelitis (A80)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Measles (B05)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Measles (B05)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2002	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001	2.0	*	*	*	*	*	*	*	0.3	0.9
2000	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999	1.7	*	*	*	*	*	*	*	0.3	0.9
All races 3/, male										
2002	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001	2.5	*	*	*	*	*	*	*	0.3	1.2
2000	2.4	*	*	*	*	*	*	*	0.3	1.3
1999	2.2	*	*	*	*	*	*	*	0.4	1.2
All races 3/, female										
2002	1.4	*	*	*	*	*	*	*	*	0.6
2001	1.4	*	*	*	*	*	*	*	0.2	0.6
2000	1.4	*	*	*	*	*	*	*	0.2	0.6
1999	1.3	*	*	*	*	*	*	*	0.2	0.6
White, both sexes										
2002	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001	1.9	*	*	*	*	*	*	*	0.2	0.9
2000	1.8	*	*	*	*	*	*	*	0.2	1.0
1999	1.7	*	*	*	*	*	*	*	0.3	0.9
White, male										
2002	2.6	*	*	*	*	*	*	*	0.4	1.5
2001	2.5	*	*	*	*	*	*	*	0.3	1.2
2000	2.4	*	*	*	*	*	*	*	0.3	1.3
1999	2.2	*	*	*	*	*	*	*	0.4	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
White, female										
2002	1.3	*	*	*	*	*	*	*	*	0.5
2001	1.3	*	*	*	*	*	*	*	*	0.6
2000	1.3	*	*	*	*	*	*	*	*	0.6
1999	1.2	*	*	*	*	*	*	*	0.2	0.5
Black, both sexes										
2002	2.3	*	*	*	*	*	*	*	*	1.1
2001	2.2	*	*	*	*	*	*	*	*	0.9
2000	2.2	*	*	*	*	*	*	*	*	1.0
1999	2.0	*	*	*	*	*	*	*	*	0.8
Black, male										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	2.7	*	*	*	*	*	*	*	*	*
2000	2.9	*	*	*	*	*	*	*	*	1.6
1999	2.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.6	*	*	*	*	*	*	*	*	*
2001	1.8	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	*
1999	1.5	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
All races 3/, both sexes										
2002	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8
1999	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0
All races 3/, male										
2002	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2
1999	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5
All races 3/, female										
2002	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4
1999	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6
White, both sexes										
2002	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000	2.8	*	*	*	*	*	0.3	1.5	4.5	7.6
1999	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9
White, male										
2002	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000	4.7	*	*	*	*	*	0.4	2.0	7.1	12.5
1999	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
White, female										
2002	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000	1.0	*	*	*	*	*	*	1.0	1.9	2.6
1999	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7
Black, both sexes										
2002	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000	21.4	*	*	*	*	0.7	3.7	15.2	37.1	49.8
1999	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9
Black, male										
2002	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000	31.0	*	*	*	*	*	3.6	16.4	47.9	71.2
1999	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9
Black, female										
2002	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000	12.8	*	*	*	*	*	3.8	14.1	27.4	30.8
1999	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malaria (B50-B54)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malaria (B50-B54)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)										
All races 3/, both sexes										
2002	2.3	5.6	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001	2.0	4.7	0.4	0.2	0.1	0.2	0.2	0.3	0.4	0.7
2000	2.0	4.4	0.5	0.1	0.2	0.2	0.3	0.3	0.4	0.7
1999	2.6	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0
All races 3/, male										
2002	2.6	6.1	0.5	*	0.2	0.2	0.3	0.4	0.5	0.9
2001	2.1	4.9	0.5	*	*	0.2	0.3	0.3	0.5	0.8
2000	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.4	0.9
1999	2.7	6.1	0.6	0.2	0.2	0.2	0.3	0.5	0.8	1.3
All races 3/, female										
2002	2.1	5.0	0.3	0.3	*	0.2	0.2	0.3	0.4	0.6
2001	1.9	4.6	0.4	0.2	*	*	*	0.3	0.4	0.6
2000	1.9	3.9	0.4	*	*	*	0.2	0.3	0.4	0.5
1999	2.5	4.5	0.4	0.2	*	0.2	0.2	0.5	0.6	0.7
White, both sexes										
2002	2.4	5.0	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.6
2001	2.0	3.8	0.4	0.2	0.1	0.2	0.2	0.3	0.3	0.6
2000	2.0	3.4	0.4	0.2	0.2	0.2	0.3	0.2	0.3	0.6
1999	2.5	4.5	0.5	0.2	0.2	0.2	0.2	0.4	0.5	0.8
White, male										
2002	2.6	5.3	0.5	*	0.2	*	*	0.3	0.5	0.7
2001	2.1	3.9	0.4	*	*	*	*	0.3	0.4	0.7
2000	2.1	3.9	0.4	*	*	*	0.3	*	0.3	0.7
1999	2.6	5.6	0.6	*	*	*	*	0.3	0.6	1.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)										
White, female										
2002	2.2	4.6	*	*	*	*	*	0.3	0.4	0.4
2001	1.9	3.8	*	*	*	*	*	*	0.2	0.5
2000	1.9	2.9	0.4	*	*	*	*	*	0.3	0.4
1999	2.5	3.4	*	*	*	*	*	0.4	0.5	0.5
Black, both sexes										
2002	2.5	8.7	*	*	*	*	*	*	1.0	1.8
2001	2.2	9.5	*	*	*	*	*	*	1.0	1.4
2000	2.3	9.9	*	*	*	*	*	*	1.2	1.6
1999	3.5	9.9	0.9	*	*	*	*	1.1	2.1	2.5
Black, male										
2002	2.9	9.6	*	*	*	*	*	*	*	2.1
2001	2.4	10.8	*	*	*	*	*	*	*	1.7
2000	2.5	9.9	*	*	*	*	*	*	*	2.2
1999	3.6	10.1	*	*	*	*	*	*	2.6	2.7
Black, female										
2002	2.1	7.9	*	*	*	*	*	*	*	1.6
2001	2.0	8.2	*	*	*	*	*	*	*	*
2000	2.1	9.9	*	*	*	*	*	*	*	*
1999	3.4	9.7	*	*	*	*	*	*	1.6	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2002	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
1999	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2
All races 3/, male										
2002	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
1999	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6
All races 3/, female										
2002	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
1999	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7
White, both sexes										
2002	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
1999	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0
White, male										
2002	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
1999	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
White, female										
2002	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
1999	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1
Black, both sexes										
2002	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
1999	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8
Black, male										
2002	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
1999	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8
Black, female										
2002	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5
1999	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes										
2002	2.7	*	*	*	*	*	*	0.1	0.2	0.4
2001	2.7	*	*	*	*	*	*	*	0.2	0.4
2000	2.7	*	*	*	*	*	*	*	0.2	0.4
1999	2.7	*	*	*	*	*	*	*	0.1	0.4
All races 3/, male										
2002	3.7	*	*	*	*	*	*	*	0.2	0.4
2001	3.7	*	*	*	*	*	*	*	0.2	0.5
2000	3.6	*	*	*	*	*	*	*	*	0.5
1999	3.6	*	*	*	*	*	*	*	*	0.5
All races 3/, female										
2002	1.7	*	*	*	*	*	*	*	*	0.3
2001	1.8	*	*	*	*	*	*	*	*	0.3
2000	1.8	*	*	*	*	*	*	*	*	0.2
1999	1.8	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	2.7	*	*	*	*	*	*	*	0.1	0.3
2001	2.7	*	*	*	*	*	*	*	0.1	0.3
2000	2.7	*	*	*	*	*	*	*	0.1	0.3
1999	2.7	*	*	*	*	*	*	*	0.1	0.3
White, male										
2002	3.7	*	*	*	*	*	*	*	*	0.4
2001	3.6	*	*	*	*	*	*	*	*	0.4
2000	3.5	*	*	*	*	*	*	*	*	0.4
1999	3.5	*	*	*	*	*	*	*	*	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
White, female										
2002	1.8	*	*	*	*	*	*	*	*	*
2001	1.9	*	*	*	*	*	*	*	*	*
2000	1.9	*	*	*	*	*	*	*	*	0.2
1999	1.9	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002	2.8	*	*	*	*	*	*	*	*	0.8
2001	2.9	*	*	*	*	*	*	*	*	0.8
2000	3.0	*	*	*	*	*	*	*	*	*
1999	3.2	*	*	*	*	*	*	*	*	0.8
Black, male										
2002	4.3	*	*	*	*	*	*	*	*	*
2001	4.4	*	*	*	*	*	*	*	*	*
2000	4.8	*	*	*	*	*	*	*	*	*
1999	4.8	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.4	*	*	*	*	*	*	*	*	*
2001	1.5	*	*	*	*	*	*	*	*	*
2000	1.4	*	*	*	*	*	*	*	*	*
1999	1.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of esophagus (C15)										
All races 3/, both sexes										
2002	4.4	*	*	*	*	*	*	*	*	0.4
2001	4.4	*	*	*	*	*	*	*	0.1	0.3
2000	4.3	*	*	*	*	*	*	*	0.1	0.3
1999	4.3	*	*	*	*	*	*	*	0.1	0.3
All races 3/, male										
2002	6.9	*	*	*	*	*	*	*	*	0.6
2001	6.9	*	*	*	*	*	*	*	*	0.5
2000	6.7	*	*	*	*	*	*	*	0.2	0.6
1999	6.6	*	*	*	*	*	*	*	0.2	0.6
All races 3/, female										
2002	2.0	*	*	*	*	*	*	*	*	*
2001	2.0	*	*	*	*	*	*	*	*	*
2000	2.1	*	*	*	*	*	*	*	*	*
1999	2.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	4.6	*	*	*	*	*	*	*	*	0.4
2001	4.6	*	*	*	*	*	*	*	*	0.3
2000	4.5	*	*	*	*	*	*	*	0.2	0.4
1999	4.4	*	*	*	*	*	*	*	0.1	0.4
White, male										
2002	7.3	*	*	*	*	*	*	*	*	0.7
2001	7.2	*	*	*	*	*	*	*	*	0.5
2000	7.0	*	*	*	*	*	*	*	0.3	0.7
1999	6.9	*	*	*	*	*	*	*	*	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of esophagus (C15)										
White, female										
2002	2.0	*	*	*	*	*	*	*	*	*
2001	2.1	*	*	*	*	*	*	*	*	*
2000	2.1	*	*	*	*	*	*	*	*	*
1999	2.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	4.2	*	*	*	*	*	*	*	*	*
2001	4.5	*	*	*	*	*	*	*	*	*
2000	4.5	*	*	*	*	*	*	*	*	*
1999	4.6	*	*	*	*	*	*	*	*	*
Black, male										
2002	6.3	*	*	*	*	*	*	*	*	*
2001	7.0	*	*	*	*	*	*	*	*	*
2000	6.7	*	*	*	*	*	*	*	*	*
1999	7.0	*	*	*	*	*	*	*	*	*
Black, female										
2002	2.3	*	*	*	*	*	*	*	*	*
2001	2.3	*	*	*	*	*	*	*	*	*
2000	2.5	*	*	*	*	*	*	*	*	*
1999	2.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of stomach (C16)										
All races 3/, both sexes										
2002	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000	4.5	*	*	*	*	*	*	0.2	0.4	0.7
1999	4.6	*	*	*	*	*	*	0.2	0.4	0.8
All races 3/, male										
2002	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000	5.3	*	*	*	*	*	*	0.2	0.5	0.9
1999	5.5	*	*	*	*	*	*	0.2	0.5	0.9
All races 3/, female										
2002	3.5	*	*	*	*	*	*	*	0.4	0.7
2001	3.5	*	*	*	*	*	*	*	0.4	0.7
2000	3.7	*	*	*	*	*	*	*	0.4	0.6
1999	3.7	*	*	*	*	*	*	0.3	0.3	0.8
White, both sexes										
2002	4.0	*	*	*	*	*	*	0.2	0.4	0.7
2001	4.1	*	*	*	*	*	*	0.2	0.4	0.6
2000	4.2	*	*	*	*	*	*	0.1	0.3	0.6
1999	4.3	*	*	*	*	*	*	0.2	0.4	0.7
White, male										
2002	4.8	*	*	*	*	*	*	0.3	0.4	0.7
2001	4.9	*	*	*	*	*	*	*	0.4	0.7
2000	5.0	*	*	*	*	*	*	*	0.4	0.7
1999	5.1	*	*	*	*	*	*	*	0.5	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of stomach (C16)										
White, female										
2002	3.2	*	*	*	*	*	*	*	0.3	0.7
2001	3.3	*	*	*	*	*	*	*	0.3	0.5
2000	3.5	*	*	*	*	*	*	*	0.3	0.5
1999	3.4	*	*	*	*	*	*	*	*	0.6
Black, both sexes										
2002	5.6	*	*	*	*	*	*	*	0.7	0.9
2001	5.6	*	*	*	*	*	*	*	0.8	1.7
2000	5.9	*	*	*	*	*	*	*	0.9	1.3
1999	6.1	*	*	*	*	*	*	*	0.8	1.4
Black, male										
2002	6.6	*	*	*	*	*	*	*	*	*
2001	6.7	*	*	*	*	*	*	*	*	1.8
2000	7.0	*	*	*	*	*	*	*	*	1.9
1999	7.3	*	*	*	*	*	*	*	*	1.4
Black, female										
2002	4.7	*	*	*	*	*	*	*	*	*
2001	4.5	*	*	*	*	*	*	*	*	1.5
2000	5.0	*	*	*	*	*	*	*	*	*
1999	5.0	*	*	*	*	*	*	*	*	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2002	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
1999	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9
All races 3/, male										
2002	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000	20.6	*	*	*	*	*	*	0.4	1.1	2.4
1999	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0
All races 3/, female										
2002	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000	20.2	*	*	*	*	*	*	0.4	0.8	1.8
1999	20.3	*	*	*	*	*	*	0.3	0.9	1.9
White, both sexes										
2002	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000	21.5	*	*	*	*	*	*	0.4	0.9	1.9
1999	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8
White, male										
2002	21.1	*	*	*	*	*	0.2	0.5	1.0	2.0
2001	21.2	*	*	*	*	*	*	0.4	1.1	2.5
2000	21.6	*	*	*	*	*	*	0.4	1.0	2.2
1999	21.8	*	*	*	*	*	*	0.5	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of colon, rectum and anus (C18-C21)										
White, female										
2002	20.3	*	*	*	*	*	*	0.4	0.9	1.9
2001	20.7	*	*	*	*	*	*	0.4	0.9	1.8
2000	21.4	*	*	*	*	*	*	0.3	0.7	1.6
1999	21.4	*	*	*	*	*	*	0.3	0.9	1.7
Black, both sexes										
2002	18.2	*	*	*	*	*	*	*	1.0	2.6
2001	18.6	*	*	*	*	*	*	*	1.6	3.4
2000	18.8	*	*	*	*	*	*	*	1.5	3.7
1999	18.7	*	*	*	*	*	*	0.8	1.3	3.1
Black, male										
2002	18.6	*	*	*	*	*	*	*	1.5	3.2
2001	18.4	*	*	*	*	*	*	*	1.8	3.8
2000	19.1	*	*	*	*	*	*	*	1.7	4.0
1999	18.6	*	*	*	*	*	*	*	*	3.1
Black, female										
2002	17.8	*	*	*	*	*	*	*	*	2.1
2001	18.8	*	*	*	*	*	*	*	*	3.0
2000	18.5	*	*	*	*	*	*	*	1.4	3.4
1999	18.9	*	*	*	*	*	*	*	*	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes										
2002	4.9	*	0.2	*	*	*	*	0.2	0.3	0.6
2001	4.7	*	*	*	*	*	*	0.2	0.2	0.6
2000	4.6	*	0.1	*	*	*	*	0.1	0.3	0.5
1999	4.4	*	0.2	*	*	*	0.1	0.2	0.2	0.6
All races 3/, male										
2002	6.4	*	*	*	*	*	*	0.2	0.4	0.8
2001	6.1	*	*	*	*	*	*	*	0.3	0.9
2000	5.9	*	*	*	*	*	*	*	0.4	0.7
1999	5.6	*	0.3	*	*	*	*	0.2	0.3	0.8
All races 3/, female										
2002	3.4	*	*	*	*	*	*	*	0.3	0.3
2001	3.4	*	*	*	*	*	*	*	*	0.3
2000	3.3	*	*	*	*	*	*	*	*	0.3
1999	3.3	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	4.8	*	0.2	*	*	*	*	*	0.2	0.4
2001	4.6	*	*	*	*	*	*	*	0.2	0.5
2000	4.5	*	*	*	*	*	*	*	0.2	0.4
1999	4.4	*	0.2	*	*	*	*	*	*	0.4
White, male										
2002	6.1	*	*	*	*	*	*	*	*	0.5
2001	5.9	*	*	*	*	*	*	*	*	0.6
2000	5.7	*	*	*	*	*	*	*	*	0.4
1999	5.5	*	*	*	*	*	*	*	*	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
White, female										
2002	3.5	*	*	*	*	*	*	*	0.2	0.3
2001	3.3	*	*	*	*	*	*	*	*	0.3
2000	3.3	*	*	*	*	*	*	*	*	0.3
1999	3.3	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002	4.6	*	*	*	*	*	*	*	*	0.8
2001	4.5	*	*	*	*	*	*	*	*	1.0
2000	4.4	*	*	*	*	*	*	*	*	0.8
1999	4.2	*	*	*	*	*	*	*	*	1.1
Black, male										
2002	6.5	*	*	*	*	*	*	*	*	*
2001	5.9	*	*	*	*	*	*	*	*	1.7
2000	6.1	*	*	*	*	*	*	*	*	*
1999	5.5	*	*	*	*	*	*	*	*	1.5
Black, female										
2002	2.9	*	*	*	*	*	*	*	*	*
2001	3.2	*	*	*	*	*	*	*	*	*
2000	2.8	*	*	*	*	*	*	*	*	*
1999	3.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of pancreas (C25)										
All races 3/, both sexes										
2002	10.5	*	*	*	*	*	*	*	0.2	0.7
2001	10.5	*	*	*	*	*	*	*	0.2	0.8
2000	10.4	*	*	*	*	*	*	*	0.2	0.6
1999	10.4	*	*	*	*	*	*	*	0.2	0.8
All races 3/, male										
2002	10.5	*	*	*	*	*	*	*	0.2	0.8
2001	10.3	*	*	*	*	*	*	*	0.2	0.9
2000	10.3	*	*	*	*	*	*	*	0.2	0.8
1999	10.4	*	*	*	*	*	*	*	0.3	0.9
All races 3/, female										
2002	10.5	*	*	*	*	*	*	*	*	0.6
2001	10.6	*	*	*	*	*	*	*	*	0.6
2000	10.5	*	*	*	*	*	*	*	*	0.5
1999	10.5	*	*	*	*	*	*	*	0.2	0.6
White, both sexes										
2002	11.1	*	*	*	*	*	*	*	0.1	0.7
2001	11.1	*	*	*	*	*	*	*	0.2	0.7
2000	11.0	*	*	*	*	*	*	*	0.2	0.6
1999	11.0	*	*	*	*	*	*	*	0.2	0.7
White, male										
2002	11.1	*	*	*	*	*	*	*	*	0.7
2001	11.0	*	*	*	*	*	*	*	0.2	0.8
2000	10.9	*	*	*	*	*	*	*	*	0.7
1999	10.9	*	*	*	*	*	*	*	*	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of pancreas (C25)										
White, female										
2002	11.0	*	*	*	*	*	*	*	*	0.6
2001	11.2	*	*	*	*	*	*	*	*	0.6
2000	11.1	*	*	*	*	*	*	*	*	0.4
1999	11.0	*	*	*	*	*	*	*	*	0.6
Black, both sexes										
2002	9.3	*	*	*	*	*	*	*	*	1.0
2001	9.1	*	*	*	*	*	*	*	*	0.9
2000	9.3	*	*	*	*	*	*	*	*	0.8
1999	9.4	*	*	*	*	*	*	*	*	1.2
Black, male										
2002	9.0	*	*	*	*	*	*	*	*	*
2001	8.6	*	*	*	*	*	*	*	*	*
2000	9.0	*	*	*	*	*	*	*	*	*
1999	9.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	9.5	*	*	*	*	*	*	*	*	*
2001	9.6	*	*	*	*	*	*	*	*	*
2000	9.7	*	*	*	*	*	*	*	*	*
1999	9.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of larynx (C32)										
All races 3/, both sexes										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.4	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	2.1	*	*	*	*	*	*	*	*	*
2001	2.2	*	*	*	*	*	*	*	*	*
2000	2.2	*	*	*	*	*	*	*	*	*
1999	2.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.3	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	*
White, male										
2002	2.0	*	*	*	*	*	*	*	*	*
2001	2.1	*	*	*	*	*	*	*	*	*
2000	2.1	*	*	*	*	*	*	*	*	*
1999	2.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of larynx (C32)										
White, female										
2002	0.6	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	1.8	*	*	*	*	*	*	*	*	*
2001	1.8	*	*	*	*	*	*	*	*	*
2000	2.0	*	*	*	*	*	*	*	*	*
1999	1.8	*	*	*	*	*	*	*	*	*
Black, male										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	3.1	*	*	*	*	*	*	*	*	*
2000	3.3	*	*	*	*	*	*	*	*	*
1999	3.1	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2002	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000	55.3	*	*	*	*	*	*	0.2	0.8	2.9
1999	54.5	*	*	*	*	*	*	0.2	0.7	2.9
All races 3/, male										
2002	63.7	*	*	*	*	*	*	*	0.7	2.6
2001	64.7	*	*	*	*	*	*	*	0.6	2.8
2000	65.5	*	*	*	*	*	*	0.3	0.7	3.3
1999	65.4	*	*	*	*	*	*	0.2	0.7	2.8
All races 3/, female										
2002	46.0	*	*	*	*	*	*	*	0.6	2.7
2001	45.3	*	*	*	*	*	*	*	0.6	2.9
2000	45.4	*	*	*	*	*	*	*	0.8	2.6
1999	44.1	*	*	*	*	*	*	*	0.6	3.0
White, both sexes										
2002	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000	59.5	*	*	*	*	*	*	0.2	0.8	2.7
1999	58.5	*	*	*	*	*	*	0.2	0.7	2.7
White, male										
2002	67.7	*	*	*	*	*	*	*	0.7	2.5
2001	68.6	*	*	*	*	*	*	*	0.6	2.8
2000	69.4	*	*	*	*	*	*	0.3	0.7	3.0
1999	69.1	*	*	*	*	*	*	*	0.8	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
White, female										
2002	50.7	*	*	*	*	*	*	*	0.5	2.8
2001	49.8	*	*	*	*	*	*	*	0.6	3.0
2000	49.9	*	*	*	*	*	*	*	0.8	2.4
1999	48.2	*	*	*	*	*	*	*	0.6	2.8
Black, both sexes										
2002	43.2	*	*	*	*	*	*	*	*	3.3
2001	43.1	*	*	*	*	*	*	*	*	3.5
2000	43.9	*	*	*	*	*	*	*	0.9	4.8
1999	44.1	*	*	*	*	*	*	*	*	4.5
Black, male										
2002	56.0	*	*	*	*	*	*	*	*	4.2
2001	57.2	*	*	*	*	*	*	*	*	3.9
2000	58.3	*	*	*	*	*	*	*	*	5.8
1999	58.9	*	*	*	*	*	*	*	*	4.5
Black, female										
2002	31.6	*	*	*	*	*	*	*	*	2.5
2001	30.2	*	*	*	*	*	*	*	*	3.2
2000	30.8	*	*	*	*	*	*	*	*	4.0
1999	30.7	*	*	*	*	*	*	*	*	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2002	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
1999	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
All races 3/, male										
2002	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000	3.3	*	*	*	*	*	*	0.3	0.7	1.0
1999	3.3	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2002	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999	1.9	*	*	*	*	*	*	0.2	0.6	0.8
White, both sexes										
2002	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
1999	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3
White, male										
2002	4.0	*	*	*	*	*	*	0.5	1.0	1.3
2001	4.1	*	*	*	*	*	*	0.5	0.7	1.4
2000	4.0	*	*	*	*	*	*	0.4	0.9	1.2
1999	4.0	*	*	*	*	*	*	0.5	0.8	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
White, female										
2002	2.3	*	*	*	*	*	*	0.4	0.5	1.1
2001	2.3	*	*	*	*	*	*	0.4	0.6	0.9
2000	2.3	*	*	*	*	*	*	0.3	0.6	1.2
1999	2.3	*	*	*	*	*	*	0.3	0.7	0.9
Black, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2002	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000	15.0	*	*	*	*	*	*	0.4	1.9	4.3
1999	14.9	*	*	*	*	*	*	0.4	1.7	4.3
All races 3/, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000	29.2	*	*	*	*	*	*	0.8	3.8	8.5
1999	28.9	*	*	*	*	*	*	0.9	3.4	8.5
White, both sexes										
2002	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000	15.7	*	*	*	*	*	*	0.4	1.6	3.9
1999	15.5	*	*	*	*	*	*	0.4	1.5	3.7
White, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
White, female										
2002	29.5	*	*	*	*	*	*	0.5	3.0	7.2
2001	29.8	*	*	*	*	*	*	0.8	3.0	7.8
2000	30.7	*	*	*	*	*	*	0.8	3.3	7.7
1999	30.2	*	*	*	*	*	*	0.8	3.0	7.5
Black, both sexes										
2002	14.9	*	*	*	*	*	*	0.8	3.7	8.1
2001	15.0	*	*	*	*	*	*	1.1	4.0	7.5
2000	14.8	*	*	*	*	*	*	0.7	3.7	7.7
1999	15.0	*	*	*	*	*	*	0.9	3.7	8.8
Black, male										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000	27.9	*	*	*	*	*	*	*	7.1	14.3
1999	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of cervix uteri (C53)										
All races 3/, both sexes										
2002	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000	1.5	*	*	*	*	*	*	0.4	0.7	1.3
1999	1.5	*	*	*	*	*	*	0.3	0.8	1.3
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000	2.9	*	*	*	*	*	*	0.8	1.5	2.6
1999	3.0	*	*	*	*	*	*	0.6	1.5	2.5
White, both sexes										
2002	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999	1.4	*	*	*	*	*	*	0.3	0.7	1.2
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of cervix uteri (C53)										
White, female										
2002	2.5	*	*	*	*	*	*	0.5	1.5	2.4
2001	2.7	*	*	*	*	*	*	0.6	1.5	2.5
2000	2.7	*	*	*	*	*	*	0.8	1.4	2.4
1999	2.8	*	*	*	*	*	*	0.7	1.5	2.4
Black, both sexes										
2002	2.2	*	*	*	*	*	*	*	1.1	2.1
2001	2.1	*	*	*	*	*	*	*	1.1	1.8
2000	2.4	*	*	*	*	*	*	*	1.4	2.1
1999	2.3	*	*	*	*	*	*	*	1.0	2.1
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	4.2	*	*	*	*	*	*	*	2.0	3.9
2001	4.0	*	*	*	*	*	*	*	2.1	3.5
2000	4.5	*	*	*	*	*	*	*	2.7	4.0
1999	4.4	*	*	*	*	*	*	*	1.9	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
All races 3/, both sexes										
2002	2.4	*	*	*	*	*	*	*	0.1	0.2
2001	2.4	*	*	*	*	*	*	*	0.1	0.2
2000	2.3	*	*	*	*	*	*	*	0.1	0.2
1999	2.3	*	*	*	*	*	*	*	*	0.2
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	4.7	*	*	*	*	*	*	*	0.2	0.4
2001	4.7	*	*	*	*	*	*	*	0.3	0.4
2000	4.6	*	*	*	*	*	*	*	0.2	0.4
1999	4.5	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	2.4	*	*	*	*	*	*	*	*	0.2
2001	2.4	*	*	*	*	*	*	*	0.1	0.2
2000	2.3	*	*	*	*	*	*	*	*	0.2
1999	2.4	*	*	*	*	*	*	*	*	0.2
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
White, female										
2002	4.7	*	*	*	*	*	*	*	*	0.4
2001	4.8	*	*	*	*	*	*	*	0.3	0.4
2000	4.6	*	*	*	*	*	*	*	*	0.4
1999	4.6	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	2.8	*	*	*	*	*	*	*	*	*
2000	2.9	*	*	*	*	*	*	*	*	*
1999	2.7	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	5.7	*	*	*	*	*	*	*	*	*
2001	5.4	*	*	*	*	*	*	*	*	*
2000	5.5	*	*	*	*	*	*	*	*	*
1999	5.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of ovary (C56)										
All races 3/, both sexes										
2002	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000	5.0	*	*	*	*	*	*	0.2	0.3	0.6
1999	4.9	*	*	*	*	*	*	0.1	0.2	0.7
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000	9.8	*	*	*	*	*	*	0.4	0.6	1.3
1999	9.6	*	*	*	*	*	*	0.3	0.5	1.4
White, both sexes										
2002	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000	5.5	*	*	*	*	*	*	0.2	0.3	0.6
1999	5.4	*	*	*	*	*	*	0.1	0.2	0.7
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of ovary (C56)										
White, female										
2002	11.1	*	*	*	*	*	0.3	0.4	0.6	1.4
2001	11.0	*	*	*	*	*	*	0.4	0.6	1.3
2000	10.9	*	*	*	*	*	*	0.4	0.6	1.3
1999	10.6	*	*	*	*	*	*	0.3	0.5	1.5
Black, both sexes										
2002	3.1	*	*	*	*	*	*	*	*	0.8
2001	3.1	*	*	*	*	*	*	*	*	0.9
2000	2.9	*	*	*	*	*	*	*	*	*
1999	3.0	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	5.9	*	*	*	*	*	*	*	*	1.6
2001	5.9	*	*	*	*	*	*	*	*	1.6
2000	5.6	*	*	*	*	*	*	*	*	*
1999	5.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of prostate (C61)										
All races 3/, both sexes										
2002	10.6	*	*	*	*	*	*	*	*	*
2001	10.8	*	*	*	*	*	*	*	*	*
2000	11.0	*	*	*	*	*	*	*	*	*
1999	11.4	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	21.5	*	*	*	*	*	*	*	*	*
2001	22.0	*	*	*	*	*	*	*	*	*
2000	22.5	*	*	*	*	*	*	*	*	*
1999	23.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
White, both sexes										
2002	10.6	*	*	*	*	*	*	*	*	*
2001	10.8	*	*	*	*	*	*	*	*	*
2000	11.0	*	*	*	*	*	*	*	*	*
1999	11.4	*	*	*	*	*	*	*	*	*
White, male										
2002	21.5	*	*	*	*	*	*	*	*	*
2001	21.9	*	*	*	*	*	*	*	*	*
2000	22.3	*	*	*	*	*	*	*	*	*
1999	23.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of prostate (C61)										
White, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, both sexes										
2002	13.6	*	*	*	*	*	*	*	*	*
2001	14.2	*	*	*	*	*	*	*	*	*
2000	14.6	*	*	*	*	*	*	*	*	*
1999	14.8	*	*	*	*	*	*	*	*	*
Black, male										
2002	28.6	*	*	*	*	*	*	*	*	*
2001	29.7	*	*	*	*	*	*	*	*	*
2000	30.7	*	*	*	*	*	*	*	*	*
1999	31.2	*	*	*	*	*	*	*	*	*
Black, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
All races 3/, both sexes										
2002	4.2	*	*	0.1	*	*	*	*	0.2	0.4
2001	4.2	*	0.1	0.1	*	*	*	*	0.2	0.3
2000	4.2	*	*	0.1	*	*	*	*	0.1	0.4
1999	4.0	*	*	0.1	*	*	*	0.1	0.1	0.5
All races 3/, male										
2002	5.4	*	*	*	*	*	*	*	0.2	0.6
2001	5.4	*	*	*	*	*	*	*	*	0.5
2000	5.3	*	*	*	*	*	*	*	*	0.5
1999	5.0	*	*	*	*	*	*	*	0.2	0.6
All races 3/, female										
2002	3.1	*	*	*	*	*	*	*	*	0.3
2001	3.1	*	*	0.2	*	*	*	*	*	*
2000	3.1	*	*	*	*	*	*	*	*	0.3
1999	3.0	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	4.6	*	*	*	*	*	*	*	0.1	0.4
2001	4.6	*	*	0.1	*	*	*	*	0.1	0.3
2000	4.6	*	*	*	*	*	*	*	0.1	0.3
1999	4.3	*	*	*	*	*	*	*	*	0.4
White, male										
2002	5.9	*	*	*	*	*	*	*	*	0.5
2001	5.8	*	*	*	*	*	*	*	*	0.5
2000	5.7	*	*	*	*	*	*	*	*	0.5
1999	5.4	*	*	*	*	*	*	*	*	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
White, female										
2002	3.4	*	*	*	*	*	*	*	*	0.2
2001	3.4	*	*	*	*	*	*	*	*	*
2000	3.4	*	*	*	*	*	*	*	*	*
1999	3.2	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002	2.8	*	*	*	*	*	*	*	*	0.7
2001	3.0	*	*	*	*	*	*	*	*	0.7
2000	2.9	*	*	*	*	*	*	*	*	0.7
1999	2.9	*	*	*	*	*	*	*	*	0.9
Black, male										
2002	3.5	*	*	*	*	*	*	*	*	*
2001	3.8	*	*	*	*	*	*	*	*	*
2000	3.6	*	*	*	*	*	*	*	*	*
1999	3.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	2.2	*	*	*	*	*	*	*	*	*
2001	2.2	*	*	*	*	*	*	*	*	*
2000	2.1	*	*	*	*	*	*	*	*	*
1999	2.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of bladder (C67)										
All races 3/, both sexes										
2002	4.4	*	*	*	*	*	*	*	*	0.1
2001	4.3	*	*	*	*	*	*	*	*	0.2
2000	4.3	*	*	*	*	*	*	*	*	0.2
1999	4.3	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	6.0	*	*	*	*	*	*	*	*	0.2
2001	6.0	*	*	*	*	*	*	*	*	*
2000	5.9	*	*	*	*	*	*	*	*	0.2
1999	5.9	*	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002	2.8	*	*	*	*	*	*	*	*	*
2001	2.7	*	*	*	*	*	*	*	*	0.2
2000	2.7	*	*	*	*	*	*	*	*	*
1999	2.7	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	4.9	*	*	*	*	*	*	*	*	0.2
2001	4.8	*	*	*	*	*	*	*	*	0.2
2000	4.8	*	*	*	*	*	*	*	*	0.2
1999	4.8	*	*	*	*	*	*	*	*	0.1
White, male										
2002	6.8	*	*	*	*	*	*	*	*	*
2001	6.8	*	*	*	*	*	*	*	*	*
2000	6.7	*	*	*	*	*	*	*	*	0.2
1999	6.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of bladder (C67)										
White, female										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	2.9	*	*	*	*	*	*	*	*	*
2000	2.9	*	*	*	*	*	*	*	*	*
1999	2.9	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	2.5	*	*	*	*	*	*	*	*	*
2001	2.4	*	*	*	*	*	*	*	*	*
2000	2.3	*	*	*	*	*	*	*	*	*
1999	2.5	*	*	*	*	*	*	*	*	*
Black, male										
2002	2.7	*	*	*	*	*	*	*	*	*
2001	2.5	*	*	*	*	*	*	*	*	*
2000	2.7	*	*	*	*	*	*	*	*	*
1999	2.9	*	*	*	*	*	*	*	*	*
Black, female										
2002	2.3	*	*	*	*	*	*	*	*	*
2001	2.2	*	*	*	*	*	*	*	*	*
2000	2.0	*	*	*	*	*	*	*	*	*
1999	2.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes										
2002	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9
All races 3/, male										
2002	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
1999	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3
All races 3/, female										
2002	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
1999	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5
White, both sexes										
2002	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
1999	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2
White, male										
2002	5.7	*	0.7	1.1	0.7	0.7	0.8	0.7	1.6	2.3
2001	5.7	*	0.8	0.8	0.6	0.7	0.7	1.1	1.9	2.6
2000	5.7	*	0.7	0.9	0.8	0.6	0.7	1.0	1.7	2.8
1999	5.8	*	0.8	0.9	0.6	0.6	0.7	1.2	1.8	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
White, female										
2002	4.4	*	0.6	1.0	0.6	0.4	0.3	0.7	1.2	1.6
2001	4.4	*	0.7	1.0	0.7	0.4	0.6	0.5	1.2	1.6
2000	4.4	*	0.6	0.8	0.7	0.4	0.6	0.8	0.9	1.5
1999	4.6	*	0.8	0.8	0.7	0.5	0.5	0.9	1.0	1.7
Black, both sexes										
2002	2.1	*	1.1	1.0	0.6	*	*	*	0.8	1.0
2001	2.0	*	*	*	0.7	*	*	*	0.7	*
2000	2.1	*	*	0.7	0.8	*	*	*	0.8	1.2
1999	2.1	*	*	1.1	*	0.7	*	*	0.9	0.9
Black, male										
2002	2.3	*	*	1.4	*	*	*	*	*	*
2001	2.2	*	*	*	*	*	*	*	*	*
2000	2.4	*	*	*	*	*	*	*	*	*
1999	2.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.9	*	*	*	*	*	*	*	*	*
2001	1.8	*	*	*	*	*	*	*	*	*
2000	1.9	*	*	*	*	*	*	*	*	*
1999	1.9	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All races 3/, both sexes										
2002	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
1999	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7
All races 3/, male										
2002	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
1999	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4
All races 3/, female										
2002	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
1999	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0
White, both sexes										
2002	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
1999	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6
White, male										
2002	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
1999	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, female										
2002	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
1999	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9
Black, both sexes										
2002	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
1999	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6
Black, male										
2002	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
1999	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3
Black, female										
2002	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1
1999	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
All races 3/, both sexes										
2002	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3
1999	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4
All races 3/, male										
2002	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5
All races 3/, female										
2002	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3
1999	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3
White, both sexes										
2002	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4
White, male										
2002	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2001	0.6	*	*	*	*	*	0.3	0.4	0.5	0.5
2000	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
White, female										
2002	0.5	*	*	*	*	*	0.4	0.3	0.5	0.3
2001	0.4	*	*	*	*	*	*	0.4	0.3	0.3
2000	0.5	*	*	*	*	*	*	0.3	0.4	0.3
1999	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3
Black, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	0.8	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2002	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4
All races 3/, male										
2002	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8
All races 3/, female										
2002	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0
White, both sexes										
2002	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4
White, male										
2002	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
White, female										
2002	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000	8.6	*	*	*	*	*	0.4	0.5	0.6	0.8
1999	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0
Black, both sexes										
2002	3.9	*	*	*	*	*	*	0.8	1.5	1.4
2001	4.0	*	*	*	*	*	*	1.1	1.2	2.0
2000	3.7	*	*	*	*	*	1.0	1.1	0.9	1.9
1999	4.0	*	*	*	*	*	*	0.9	1.2	2.1
Black, male										
2002	4.3	*	*	*	*	*	*	*	2.2	2.0
2001	4.5	*	*	*	*	*	*	*	*	3.0
2000	4.2	*	*	*	*	*	*	*	*	2.7
1999	4.6	*	*	*	*	*	*	*	1.7	2.6
Black, female										
2002	3.5	*	*	*	*	*	*	*	*	*
2001	3.6	*	*	*	*	*	*	*	*	*
2000	3.3	*	*	*	*	*	*	*	*	*
1999	3.5	*	*	*	*	*	*	*	*	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
All races 3/, both sexes										
2002	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6
1999	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6
All races 3/, male										
2002	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6
1999	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8
All races 3/, female										
2002	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6
1999	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5
White, both sexes										
2002	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000	8.3	1.2	1.0	0.7	0.9	1.2	1.2	1.1	1.1	1.6
1999	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6
White, male										
2002	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000	9.4	*	0.9	0.8	1.1	1.4	1.5	1.3	1.4	1.7
1999	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
White, female										
2002	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000	7.4	*	1.0	0.7	0.7	0.9	0.9	0.9	0.9	1.6
1999	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5
Black, both sexes										
2002	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000	4.8	*	*	*	0.6	1.5	1.3	1.4	1.8	1.7
1999	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8
Black, male										
2002	5.1	*	*	*	*	*	*	*	1.6	1.9
2001	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000	5.3	*	*	*	*	2.0	1.8	*	2.5	1.4
1999	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8
Black, female										
2002	4.3	*	*	*	*	*	*	*	*	*
2001	4.3	*	*	*	*	*	*	*	*	1.4
2000	4.2	*	*	*	*	*	*	1.5	*	1.9
1999	4.3	*	*	*	*	*	1.8	*	*	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
All races 3/, both sexes										
2002	4.0	*	*	*	*	*	*	*	*	0.2
2001	3.9	*	*	*	*	*	*	*	*	0.2
2000	3.9	*	*	*	*	*	*	*	*	0.2
1999	3.9	*	*	*	*	*	*	*	*	0.2
All races 3/, male										
2002	4.2	*	*	*	*	*	*	*	*	0.2
2001	4.1	*	*	*	*	*	*	*	*	0.2
2000	4.0	*	*	*	*	*	*	*	*	*
1999	4.1	*	*	*	*	*	*	*	*	0.3
All races 3/, female										
2002	3.7	*	*	*	*	*	*	*	*	*
2001	3.8	*	*	*	*	*	*	*	*	*
2000	3.8	*	*	*	*	*	*	*	*	0.2
1999	3.8	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	4.0	*	*	*	*	*	*	*	*	0.1
2001	3.9	*	*	*	*	*	*	*	*	0.2
2000	3.9	*	*	*	*	*	*	*	*	0.1
1999	3.9	*	*	*	*	*	*	*	*	0.2
White, male										
2002	4.3	*	*	*	*	*	*	*	*	*
2001	4.2	*	*	*	*	*	*	*	*	*
2000	4.1	*	*	*	*	*	*	*	*	*
1999	4.1	*	*	*	*	*	*	*	*	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
White, female										
2002	3.7	*	*	*	*	*	*	*	*	*
2001	3.7	*	*	*	*	*	*	*	*	*
2000	3.8	*	*	*	*	*	*	*	*	*
1999	3.8	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	4.7	*	*	*	*	*	*	*	*	*
2001	4.9	*	*	*	*	*	*	*	*	*
2000	5.0	*	*	*	*	*	*	*	*	*
1999	5.0	*	*	*	*	*	*	*	*	*
Black, male										
2002	4.7	*	*	*	*	*	*	*	*	*
2001	4.9	*	*	*	*	*	*	*	*	*
2000	4.8	*	*	*	*	*	*	*	*	*
1999	4.9	*	*	*	*	*	*	*	*	*
Black, female										
2002	4.7	*	*	*	*	*	*	*	*	*
2001	4.9	*	*	*	*	*	*	*	*	*
2000	5.2	*	*	*	*	*	*	*	*	*
1999	5.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
White, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)										
All races 3/, both sexes										
2002	21.8	0.6	0.8	0.7	0.7	1.1	1.5	1.5	2.1	3.3
2001	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2
2000	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4
1999	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6
All races 3/, male										
2002	22.7	*	0.8	0.8	0.7	1.3	1.9	2.0	2.5	3.7
2001	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5
2000	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7
1999	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1
All races 3/, female										
2002	21.0	*	0.8	0.6	0.8	0.9	1.0	1.0	1.7	2.8
2001	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0
2000	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0
1999	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2
White, both sexes										
2002	23.3	0.7	0.7	0.7	0.8	1.2	1.5	1.5	2.1	3.2
2001	23.5	*	0.8	0.6	0.8	1.2	1.5	1.7	1.9	3.1
2000	23.9	*	0.8	0.7	0.7	1.2	1.5	1.6	2.0	3.2
1999	25.3	*	0.8	0.7	0.7	1.2	1.4	1.6	2.2	3.5
White, male										
2002	24.2	*	0.8	0.7	0.7	1.4	1.9	2.0	2.5	3.6
2001	24.0	*	1.0	0.6	0.8	1.5	2.0	2.0	2.3	3.4
2000	24.4	*	1.0	0.7	0.7	1.3	1.9	2.1	2.5	3.6
1999	25.9	*	0.9	0.8	0.8	1.5	1.7	2.2	2.7	4.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)										
White, female										
2002	22.5	*	0.7	0.6	0.8	1.0	1.0	1.0	1.7	2.8
2001	22.9	*	0.7	0.5	0.7	1.0	0.8	1.4	1.5	2.8
2000	23.5	*	0.6	0.6	0.6	1.0	1.1	1.1	1.5	2.8
1999	24.6	*	0.7	0.7	0.6	1.0	1.1	1.0	1.7	3.0
Black, both sexes										
2002	18.1	*	1.1	0.8	0.7	1.1	1.6	1.9	2.6	4.5
2001	18.3	*	*	*	0.8	1.2	1.3	2.0	2.6	4.3
2000	18.3	*	1.0	*	0.7	1.5	1.3	1.3	2.0	5.2
1999	19.5	*	*	0.6	0.9	1.3	1.6	1.5	2.4	4.9
Black, male										
2002	18.8	*	*	*	*	1.4	2.2	2.5	2.7	5.1
2001	19.4	*	*	*	*	1.5	1.9	2.5	3.0	4.4
2000	19.4	*	*	*	*	1.7	2.1	1.6	2.0	5.4
1999	20.6	*	*	*	*	1.9	2.0	1.5	3.1	5.1
Black, female										
2002	17.4	*	*	*	*	*	*	1.4	2.6	4.0
2001	17.3	*	*	*	*	*	*	1.5	2.3	4.2
2000	17.3	*	*	*	*	1.4	*	*	1.9	5.1
1999	18.5	*	*	*	*	*	*	1.5	1.8	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All races 3/, both sexes										
2002	4.6	1.7	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001	4.8	1.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.7
2000	4.8	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7
1999	4.7	1.5	0.4	0.3	0.2	0.3	0.2	0.3	0.6	0.6
All races 3/, male										
2002	4.6	1.5	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001	4.8	1.3	0.4	0.3	0.2	0.2	0.3	0.3	0.4	0.7
2000	4.8	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8
1999	4.8	1.5	0.4	0.3	*	0.3	0.3	0.3	0.6	0.7
All races 3/, female										
2002	4.6	2.0	0.4	0.2	*	0.2	*	0.3	0.5	0.5
2001	4.8	1.3	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.7
2000	4.9	1.5	0.3	0.3	0.2	*	0.2	0.4	0.4	0.6
1999	4.7	1.5	0.4	0.3	*	0.3	*	0.2	0.5	0.5
White, both sexes										
2002	5.1	1.9	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001	5.3	1.2	0.4	0.2	0.2	0.2	0.3	0.4	0.4	0.7
2000	5.3	1.6	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7
1999	5.2	1.4	0.4	0.3	0.2	0.2	0.2	0.3	0.5	0.6
White, male										
2002	5.1	1.7	0.3	*	0.2	*	0.2	0.3	0.5	0.7
2001	5.3	1.2	0.4	*	*	*	0.3	0.4	0.4	0.7
2000	5.2	1.6	0.4	0.3	*	*	0.4	0.3	0.5	0.7
1999	5.2	1.4	0.4	0.3	*	*	0.3	0.3	0.5	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
White, female										
2002	5.1	2.1	0.4	0.3	*	*	*	*	0.4	0.5
2001	5.2	*	0.4	*	0.3	0.3	*	0.3	0.5	0.7
2000	5.4	1.6	*	0.3	*	*	*	0.3	0.4	0.6
1999	5.1	1.4	0.4	0.4	*	0.3	*	*	0.5	0.5
Black, both sexes										
2002	3.0	*	*	*	*	*	*	*	*	0.7
2001	3.1	*	*	*	*	*	*	*	*	0.9
2000	3.1	*	*	*	*	*	*	*	*	0.8
1999	3.2	*	*	*	*	*	*	*	0.8	1.0
Black, male										
2002	2.8	*	*	*	*	*	*	*	*	*
2001	3.0	*	*	*	*	*	*	*	*	*
2000	3.0	*	*	*	*	*	*	*	*	*
1999	3.0	*	*	*	*	*	*	*	*	*
Black, female										
2002	3.3	*	*	*	*	*	*	*	*	*
2001	3.2	*	*	*	*	*	*	*	*	*
2000	3.2	*	*	*	*	*	*	*	*	*
1999	3.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
All races 3/, both sexes										
2002	1.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.5
2001	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4
2000	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4
1999	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4
All races 3/, male										
2002	1.3	*	*	*	*	0.2	0.4	0.5	0.5	0.5
2001	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4
2000	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3
1999	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4
All races 3/, female										
2002	1.9	*	*	*	*	*	0.3	0.3	0.5	0.4
2001	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2000	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4
1999	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4
White, both sexes										
2002	1.5	*	*	*	*	*	*	*	0.2	0.1
2001	1.5	*	*	*	*	*	*	*	0.1	*
2000	1.5	*	*	*	*	*	*	*	*	*
1999	1.5	*	*	*	*	*	*	*	*	0.1
White, male										
2002	1.1	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.1	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
White, female										
2002	1.8	*	*	*	*	*	*	*	*	*
2001	1.9	*	*	*	*	*	*	*	*	*
2000	1.9	*	*	*	*	*	*	*	*	*
1999	1.9	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
1999	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2
Black, male										
2002	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
1999	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5
Black, female										
2002	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5
1999	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2002	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0
1999	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2
All races 3/, male										
2002	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5
1999	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8
All races 3/, female										
2002	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5
1999	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6
White, both sexes										
2002	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000	24.1	*	*	*	*	0.2	0.5	0.8	1.8	2.7
1999	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8
White, male										
2002	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000	22.9	*	*	*	*	0.3	0.5	1.0	2.3	3.1
1999	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
White, female										
2002	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000	25.3	*	*	*	*	*	0.4	0.7	1.2	2.3
1999	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1
Black, both sexes										
2002	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000	32.8	*	*	*	*	*	1.5	2.8	4.1	6.0
1999	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6
Black, male										
2002	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000	27.4	*	*	*	*	*	1.5	3.1	4.5	7.1
1999	27.7	*	*	*	*	*	*	1.6	5.0	7.7
Black, female										
2002	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000	37.8	*	*	*	*	*	1.5	2.5	3.8	5.0
1999	37.8	*	*	*	*	*	*	2.0	3.3	5.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nutritional deficiencies (E40-E64)										
All races 3/, both sexes										
2002	1.3	*	*	*	*	*	*	*	*	0.1
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	0.1
1999	1.5	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	0.9	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	1.7	*	*	*	*	*	*	*	*	*
2001	1.7	*	*	*	*	*	*	*	*	*
2000	2.0	*	*	*	*	*	*	*	*	0.2
1999	2.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.4	*	*	*	*	*	*	*	*	*
2001	1.4	*	*	*	*	*	*	*	*	*
2000	1.6	*	*	*	*	*	*	*	*	0.1
1999	1.6	*	*	*	*	*	*	*	*	*
White, male										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	0.9	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nutritional deficiencies (E40-E64)										
White, female										
2002	1.8	*	*	*	*	*	*	*	*	*
2001	1.8	*	*	*	*	*	*	*	*	*
2000	2.1	*	*	*	*	*	*	*	*	*
1999	2.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	*
1999	1.6	*	*	*	*	*	*	*	*	*
Black, male										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.5	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.6	*	*	*	*	*	*	*	*	*
2001	1.4	*	*	*	*	*	*	*	*	*
2000	1.7	*	*	*	*	*	*	*	*	*
1999	1.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malnutrition (E40-E46)										
All races 3/, both sexes										
2002	1.2	*	*	*	*	*	*	*	*	0.1
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.4	*	*	*	*	*	*	*	*	0.1
1999	1.4	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.9	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	1.6	*	*	*	*	*	*	*	*	*
2001	1.6	*	*	*	*	*	*	*	*	*
2000	1.8	*	*	*	*	*	*	*	*	0.2
1999	1.8	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	0.1
1999	1.5	*	*	*	*	*	*	*	*	*
White, male										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malnutrition (E40-E46)										
White, female										
2002	1.6	*	*	*	*	*	*	*	*	*
2001	1.7	*	*	*	*	*	*	*	*	*
2000	1.9	*	*	*	*	*	*	*	*	*
1999	2.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.4	*	*	*	*	*	*	*	*	*
1999	1.5	*	*	*	*	*	*	*	*	*
Black, male										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.5	*	*	*	*	*	*	*	*	*
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.6	*	*	*	*	*	*	*	*	*
1999	1.6	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other nutritional deficiencies (E50-E64)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other nutritional deficiencies (E50-E64)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningitis (G00,G03)										
All races 3/, both sexes										
2002	0.2	1.8	*	*	0.1	*	*	*	0.1	0.1
2001	0.3	1.9	0.2	*	*	*	0.1	*	0.1	0.1
2000	0.3	2.4	0.2	*	*	*	*	*	0.1	0.2
1999	0.3	3.1	0.3	*	*	*	0.1	0.1	0.1	0.1
All races 3/, male										
2002	0.3	2.0	*	*	*	*	*	*	*	*
2001	0.3	1.9	*	*	*	*	*	*	*	0.2
2000	0.3	2.5	0.3	*	*	*	*	*	*	0.2
1999	0.3	3.1	0.3	*	*	*	*	*	*	*
All races 3/, female										
2002	0.2	1.7	*	*	*	*	*	*	*	*
2001	0.2	2.0	*	*	*	*	*	*	*	*
2000	0.2	2.3	*	*	*	*	*	*	*	*
1999	0.3	3.0	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.2	1.7	*	*	*	*	*	*	*	*
2001	0.2	1.5	*	*	*	*	*	*	0.1	*
2000	0.2	1.8	0.2	*	*	*	*	*	*	0.1
1999	0.3	2.3	0.3	*	*	*	*	*	*	0.1
White, male										
2002	0.2	1.8	*	*	*	*	*	*	*	*
2001	0.2	1.4	*	*	*	*	*	*	*	*
2000	0.3	2.0	*	*	*	*	*	*	*	*
1999	0.3	3.0	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Meningitis (G00,G03)										
White, female										
2002	0.2	1.5	*	*	*	*	*	*	*	*
2001	0.2	1.6	*	*	*	*	*	*	*	*
2000	0.2	1.6	*	*	*	*	*	*	*	*
1999	0.3	1.5	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.5	3.0	*	*	*	*	*	*	*	*
2001	0.5	4.5	*	*	*	*	*	*	*	*
2000	0.6	5.7	*	*	*	*	*	*	*	*
1999	0.6	6.9	*	*	*	*	*	*	*	*
Black, male										
2002	0.6	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	9.7	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Parkinson's disease (G20-G21)										
All races 3/, both sexes										
2002	5.9	*	*	*	*	*	*	*	*	*
2001	5.8	*	*	*	*	*	*	*	*	*
2000	5.6	*	*	*	*	*	*	*	*	*
1999	5.2	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	6.8	*	*	*	*	*	*	*	*	*
2001	6.7	*	*	*	*	*	*	*	*	*
2000	6.4	*	*	*	*	*	*	*	*	*
1999	6.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	5.0	*	*	*	*	*	*	*	*	*
2001	4.9	*	*	*	*	*	*	*	*	*
2000	4.8	*	*	*	*	*	*	*	*	*
1999	4.5	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	6.9	*	*	*	*	*	*	*	*	*
2001	6.8	*	*	*	*	*	*	*	*	*
2000	6.5	*	*	*	*	*	*	*	*	*
1999	6.1	*	*	*	*	*	*	*	*	*
White, male										
2002	7.9	*	*	*	*	*	*	*	*	*
2001	7.8	*	*	*	*	*	*	*	*	*
2000	7.4	*	*	*	*	*	*	*	*	*
1999	7.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Parkinson's disease (G20-G21)										
White, female										
2002	5.9	*	*	*	*	*	*	*	*	*
2001	5.8	*	*	*	*	*	*	*	*	*
2000	5.6	*	*	*	*	*	*	*	*	*
1999	5.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	1.5	*	*	*	*	*	*	*	*	*
2001	1.5	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	*
Black, male										
2002	1.7	*	*	*	*	*	*	*	*	*
2001	1.7	*	*	*	*	*	*	*	*	*
2000	1.8	*	*	*	*	*	*	*	*	*
1999	1.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.3	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alzheimer's disease (G30)										
All races 3/, both sexes										
2002	20.4	*	*	*	*	*	*	*	*	*
2001	18.9	*	*	*	*	*	*	*	*	*
2000	17.6	*	*	*	*	*	*	*	*	*
1999	16.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	12.0	*	*	*	*	*	*	*	*	*
2001	11.3	*	*	*	*	*	*	*	*	*
2000	10.5	*	*	*	*	*	*	*	*	*
1999	9.8	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	28.5	*	*	*	*	*	*	*	*	*
2001	26.3	*	*	*	*	*	*	*	*	*
2000	24.5	*	*	*	*	*	*	*	*	*
1999	21.9	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	23.5	*	*	*	*	*	*	*	*	*
2001	21.7	*	*	*	*	*	*	*	*	*
2000	20.2	*	*	*	*	*	*	*	*	*
1999	18.3	*	*	*	*	*	*	*	*	*
White, male										
2002	13.7	*	*	*	*	*	*	*	*	*
2001	12.8	*	*	*	*	*	*	*	*	*
2000	11.9	*	*	*	*	*	*	*	*	*
1999	11.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alzheimer's disease (G30)										
White, female										
2002	33.0	*	*	*	*	*	*	*	*	*
2001	30.3	*	*	*	*	*	*	*	*	*
2000	28.2	*	*	*	*	*	*	*	*	*
1999	25.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	8.9	*	*	*	*	*	*	*	*	*
2001	8.4	*	*	*	*	*	*	*	*	*
2000	7.5	*	*	*	*	*	*	*	*	*
1999	6.5	*	*	*	*	*	*	*	*	*
Black, male										
2002	5.3	*	*	*	*	*	*	*	*	*
2001	5.1	*	*	*	*	*	*	*	*	*
2000	4.5	*	*	*	*	*	*	*	*	*
1999	4.0	*	*	*	*	*	*	*	*	*
Black, female										
2002	12.0	*	*	*	*	*	*	*	*	*
2001	11.3	*	*	*	*	*	*	*	*	*
2000	10.2	*	*	*	*	*	*	*	*	*
1999	8.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
All races 3/, both sexes										
2002	318.6	15.9	1.4	0.6	1.1	2.4	3.9	7.1	12.6	25.3
2001	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0
2000	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2
1999	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9
All races 3/, male										
2002	303.3	16.6	1.5	0.6	1.3	3.1	5.0	9.0	16.1	32.9
2001	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1
2000	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3
1999	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2
All races 3/, female										
2002	333.3	15.1	1.3	0.7	0.9	1.7	2.8	5.1	9.0	17.7
2001	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0
2000	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1
1999	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6
White, both sexes										
2002	339.4	13.4	1.3	0.6	0.9	2.1	3.3	5.7	10.2	21.6
2001	344.8	12.2	1.5	0.6	1.0	1.9	3.4	5.9	10.9	21.4
2000	354.7	14.6	1.3	0.6	0.9	2.1	3.5	5.5	10.1	20.6
1999	362.7	14.6	1.3	0.7	0.9	2.4	3.4	5.4	10.6	22.0
White, male										
2002	320.9	14.3	1.5	0.5	1.1	2.7	4.3	7.5	13.3	28.5
2001	323.5	13.2	1.6	0.6	1.0	2.3	4.3	7.3	14.0	28.8
2000	334.2	15.3	1.6	0.7	1.1	2.4	4.2	6.9	13.2	27.6
1999	341.2	15.3	1.5	0.8	0.9	2.9	3.9	6.5	14.0	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
White, female										
2002	357.5	12.5	1.1	0.7	0.8	1.5	2.2	3.9	7.1	14.5
2001	365.5	11.1	1.5	0.7	0.9	1.5	2.5	4.3	7.6	13.9
2000	374.8	13.9	1.1	0.5	0.7	1.7	2.8	4.1	6.9	13.5
1999	383.6	13.8	1.2	0.6	0.9	1.9	2.9	4.2	7.2	14.1
Black, both sexes										
2002	276.1	27.6	2.0	0.8	2.0	4.0	7.8	16.6	29.6	52.9
2001	280.4	28.6	3.5	1.0	1.6	4.0	9.4	16.4	28.5	51.5
2000	285.0	27.8	2.5	1.0	1.9	4.6	9.1	15.1	27.7	51.2
1999	289.4	29.2	2.4	1.0	1.8	4.8	9.0	14.4	26.8	55.2
Black, male										
2002	268.5	28.2	1.9	*	2.3	5.1	9.4	20.6	36.9	66.8
2001	272.4	28.1	3.6	*	1.5	5.1	11.3	20.3	35.2	65.6
2000	275.6	29.1	2.1	*	1.8	5.1	10.2	18.0	32.7	61.6
1999	282.5	30.6	2.5	*	2.1	6.3	10.9	17.7	33.0	69.6
Black, female										
2002	283.0	26.9	2.1	*	1.6	2.8	6.3	13.0	23.0	40.6
2001	287.8	29.0	3.4	1.3	1.6	2.8	7.5	12.9	22.4	38.9
2000	293.4	26.5	2.9	*	2.0	4.2	8.1	12.4	23.3	42.0
1999	295.7	27.9	2.4	*	1.5	3.3	7.1	11.5	21.3	42.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All races 3/, both sexes										
2002	241.7	12.4	1.1	0.5	0.8	2.0	3.1	5.7	10.0	20.3
2001	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9
2000	252.6	13.0	1.2	0.5	0.8	2.0	3.3	5.2	9.5	19.1
1999	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6
All races 3/, male										
2002	240.7	12.9	1.1	0.4	0.9	2.6	4.0	7.4	13.4	27.8
2001	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4
2000	249.8	13.3	1.4	0.6	0.9	2.4	4.0	6.5	12.4	26.0
1999	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5
All races 3/, female										
2002	242.7	11.8	1.0	0.5	0.6	1.3	2.1	3.9	6.5	12.8
2001	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4
2000	255.3	12.5	1.0	0.4	0.7	1.6	2.6	3.9	6.5	12.2
1999	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6
White, both sexes										
2002	258.5	10.6	0.9	0.4	0.7	1.7	2.5	4.6	8.1	17.7
2001	262.8	9.9	1.2	0.4	0.7	1.5	2.6	4.5	8.5	17.3
2000	270.2	11.5	1.1	0.4	0.7	1.7	2.6	4.2	7.9	16.7
1999	277.7	11.8	1.0	0.5	0.7	2.0	2.6	4.1	8.5	17.8
White, male										
2002	256.0	11.3	1.0	0.4	0.8	2.2	3.3	6.1	11.2	24.6
2001	257.8	10.6	1.2	0.4	0.8	1.8	3.4	5.9	11.3	24.4
2000	265.8	11.9	1.3	0.5	0.8	2.0	3.3	5.3	10.7	23.5
1999	272.9	12.0	1.2	0.6	0.7	2.4	3.0	5.2	11.4	25.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
White, female										
2002	261.0	9.9	0.8	0.5	0.5	1.1	1.6	3.0	5.0	10.7
2001	267.7	9.2	1.1	0.5	0.7	1.1	1.8	3.1	5.6	10.0
2000	274.5	11.1	0.8	0.4	0.5	1.3	1.9	2.9	5.0	9.8
1999	282.4	11.6	0.9	0.5	0.6	1.6	2.1	3.0	5.4	10.2
Black, both sexes										
2002	205.6	21.3	1.6	*	1.3	3.4	6.5	13.4	23.2	40.8
2001	208.8	21.6	3.0	0.8	1.3	3.3	7.5	12.9	22.6	39.8
2000	211.8	19.8	2.0	0.9	1.4	4.0	7.4	12.8	21.5	38.2
1999	217.2	23.8	2.0	0.9	1.4	4.2	7.7	11.7	21.5	42.7
Black, male										
2002	206.3	22.1	*	*	1.5	4.5	8.0	17.5	30.0	54.4
2001	209.0	19.5	3.2	*	1.4	4.4	9.7	16.0	29.3	52.1
2000	211.1	19.8	1.9	*	1.4	4.8	8.7	15.7	26.5	48.6
1999	218.5	23.4	2.1	*	1.6	5.8	9.3	14.8	27.1	56.3
Black, female										
2002	205.0	20.6	1.8	*	*	2.4	5.0	9.7	17.0	28.6
2001	208.6	23.9	2.7	*	1.2	2.3	5.5	10.1	16.7	28.9
2000	212.6	19.8	2.0	*	1.3	3.3	6.3	10.2	16.9	29.0
1999	216.0	24.2	1.9	*	*	2.5	6.1	9.0	16.5	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
All races 3/, both sexes										
2002	1.2	*	*	*	*	*	*	*	0.1	0.2
2001	1.2	*	*	*	*	*	*	*	0.1	0.2
2000	1.3	*	*	*	*	*	*	*	0.1	0.2
1999	1.3	*	*	*	*	*	*	*	0.1	0.2
All races 3/, male										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.8	*	*	*	*	*	*	*	*	0.2
1999	0.8	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	1.7	*	*	*	*	*	*	*	*	*
2001	1.7	*	*	*	*	*	*	*	*	0.2
2000	1.8	*	*	*	*	*	*	*	*	0.2
1999	1.9	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002	1.4	*	*	*	*	*	*	*	*	0.2
2001	1.3	*	*	*	*	*	*	*	*	0.2
2000	1.4	*	*	*	*	*	*	*	*	0.2
1999	1.4	*	*	*	*	*	*	*	*	0.2
White, male										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.9	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
White, female										
2002	1.9	*	*	*	*	*	*	*	*	*
2001	1.8	*	*	*	*	*	*	*	*	*
2000	1.9	*	*	*	*	*	*	*	*	*
1999	2.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2002	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4
1999	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5
All races 3/, male										
2002	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9
1999	6.8	*	*	*	*	*	*	0.4	1.0	2.0
All races 3/, female										
2002	10.2	*	*	*	*	*	*	*	0.6	1.0
2001	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000	9.7	*	*	*	*	*	*	0.3	0.5	1.0
1999	9.4	*	*	*	*	*	*	*	0.5	1.0
White, both sexes										
2002	8.3	*	*	*	*	*	*	0.3	0.6	1.3
2001	7.8	*	*	*	*	*	*	0.2	0.5	1.1
2000	7.6	*	*	*	*	*	*	0.2	0.4	0.9
1999	7.3	*	*	*	*	*	*	0.2	0.5	0.9
White, male										
2002	7.1	*	*	*	*	*	*	0.4	0.9	1.8
2001	6.4	*	*	*	*	*	*	0.3	0.9	1.6
2000	6.1	*	*	*	*	*	*	0.3	0.6	1.4
1999	5.8	*	*	*	*	*	*	0.3	0.8	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
White, female										
2002	9.5	*	*	*	*	*	*	*	0.3	0.7
2001	9.2	*	*	*	*	*	*	*	*	0.5
2000	9.0	*	*	*	*	*	*	*	*	0.5
1999	8.7	*	*	*	*	*	*	*	*	0.5
Black, both sexes										
2002	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000	16.1	*	*	*	*	*	*	1.4	2.8	4.9
1999	15.5	*	*	*	*	*	*	1.0	2.6	5.7
Black, male										
2002	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001	15.9	*	*	*	*	*	*	*	4.0	5.9
2000	15.6	*	*	*	*	*	*	1.6	3.5	5.8
1999	15.2	*	*	*	*	*	*	*	2.8	7.3
Black, female										
2002	16.9	*	*	*	*	*	*	*	2.6	3.4
2001	16.2	*	*	*	*	*	*	*	2.3	4.4
2000	16.5	*	*	*	*	*	*	*	2.1	4.1
1999	15.9	*	*	*	*	*	*	*	2.5	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart and renal disease (I13)										
All races 3/, both sexes										
2002	1.0	*	*	*	*	*	*	*	*	0.1
2001	1.0	*	*	*	*	*	*	*	*	0.2
2000	1.0	*	*	*	*	*	*	*	*	0.1
1999	1.2	*	*	*	*	*	*	*	*	0.2
All races 3/, male										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	0.2
2000	0.8	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	0.2
2000	1.1	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.8	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	*
White, male										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart and renal disease (I13)										
White, female										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.0	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	2.4	*	*	*	*	*	*	*	*	0.8
2001	2.2	*	*	*	*	*	*	*	*	1.0
2000	2.2	*	*	*	*	*	*	*	*	*
1999	2.8	*	*	*	*	*	*	*	*	0.9
Black, male										
2002	2.3	*	*	*	*	*	*	*	*	*
2001	2.1	*	*	*	*	*	*	*	*	*
2000	2.0	*	*	*	*	*	*	*	*	*
1999	2.5	*	*	*	*	*	*	*	*	*
Black, female										
2002	2.5	*	*	*	*	*	*	*	*	*
2001	2.3	*	*	*	*	*	*	*	*	*
2000	2.4	*	*	*	*	*	*	*	*	*
1999	3.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2002	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
1999	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4
All races 3/, male										
2002	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
1999	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8
All races 3/, female										
2002	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
1999	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1
White, both sexes										
2002	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
1999	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9
White, male										
2002	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
1999	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
White, female										
2002	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
1999	200.0	*	*	*	*	*	*	0.7	1.7	4.6
Black, both sexes										
2002	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
1999	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1
Black, male										
2002	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000	141.5	*	*	*	*	*	*	3.7	8.0	20.3
1999	146.0	*	*	*	*	*	*	2.9	8.4	23.3
Black, female										
2002	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000	138.8	*	*	*	*	*	*	*	4.3	9.7
1999	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2002	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
1999	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1
All races 3/, male										
2002	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
1999	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2
All races 3/, female										
2002	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000	64.6	*	*	*	*	*	*	0.3	0.7	2.2
1999	67.0	*	*	*	*	*	*	0.3	0.9	2.0
White, both sexes										
2002	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
1999	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0
White, male										
2002	71.8	*	*	*	*	*	0.3	0.6	2.1	5.9
2001	74.4	*	*	*	*	*	0.3	0.6	2.5	6.1
2000	78.8	*	*	*	*	*	*	0.6	1.8	6.0
1999	82.7	*	*	*	*	*	*	0.6	2.1	6.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
White, female										
2002	62.8	*	*	*	*	*	*	*	0.7	1.9
2001	66.0	*	*	*	*	*	*	0.3	0.9	2.0
2000	69.6	*	*	*	*	*	*	0.3	0.6	2.0
1999	72.3	*	*	*	*	*	*	0.3	0.9	1.9
Black, both sexes										
2002	48.5	*	*	*	*	*	*	0.9	2.3	5.5
2001	50.0	*	*	*	*	*	*	1.1	2.5	5.8
2000	52.2	*	*	*	*	*	*	1.2	2.3	4.9
1999	53.3	*	*	*	*	*	*	0.8	2.4	5.5
Black, male										
2002	48.3	*	*	*	*	*	*	*	3.0	7.6
2001	49.2	*	*	*	*	*	*	*	3.4	8.0
2000	52.0	*	*	*	*	*	*	1.6	2.7	6.5
1999	53.4	*	*	*	*	*	*	*	3.4	7.7
Black, female										
2002	48.8	*	*	*	*	*	*	*	1.6	3.6
2001	50.6	*	*	*	*	*	*	*	1.7	3.8
2000	52.5	*	*	*	*	*	*	*	1.9	3.6
1999	53.2	*	*	*	*	*	*	*	1.4	3.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	0.1
2000	1.2	*	*	*	*	*	*	*	*	0.1
1999	1.2	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.3	*	*	*	*	*	*	*	*	0.2
1999	1.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	1.1	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.1	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	0.1
2000	1.2	*	*	*	*	*	*	*	*	0.1
1999	1.2	*	*	*	*	*	*	*	*	*
White, male										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.3	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other acute ischemic heart diseases (I24)										
White, female										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	1.4	*	*	*	*	*	*	*	*	*
2001	1.4	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2002	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
1999	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2
All races 3/, male										
2002	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
1999	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4
All races 3/, female										
2002	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000	111.9	*	*	*	*	*	*	0.2	1.0	2.8
1999	116.4	*	*	*	*	*	*	0.4	1.0	3.1
White, both sexes										
2002	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
1999	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8
White, male										
2002	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
1999	126.5	*	*	*	*	*	*	0.9	3.0	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
White, female										
2002	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001	118.1	*	*	*	*	*	*	*	0.9	2.4
2000	121.6	*	*	*	*	*	*	*	0.8	2.4
1999	126.5	*	*	*	*	*	*	0.4	0.9	2.7
Black, both sexes										
2002	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000	86.6	*	*	*	*	*	*	1.3	3.6	9.5
1999	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1
Black, male										
2002	85.7	*	*	*	*	*	*	*	4.1	14.4
2001	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000	88.0	*	*	*	*	*	*	2.1	5.0	13.4
1999	91.1	*	*	*	*	*	*	1.7	5.0	14.9
Black, female										
2002	81.0	*	*	*	*	*	*	*	2.3	5.3
2001	83.5	*	*	*	*	*	*	*	1.9	4.9
2000	85.3	*	*	*	*	*	*	*	2.4	5.9
1999	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2002	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000	24.7	*	*	*	*	*	*	0.2	0.5	1.7
1999	25.8	*	*	*	*	*	*	0.2	0.6	2.0
All races 3/, male										
2002	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000	26.0	*	*	*	*	*	*	0.3	0.9	2.8
1999	26.8	*	*	*	*	*	*	0.2	0.9	3.1
All races 3/, female										
2002	21.9	*	*	*	*	*	*	*	0.3	0.8
2001	22.5	*	*	*	*	*	*	*	0.4	0.8
2000	23.5	*	*	*	*	*	*	*	0.2	0.7
1999	24.8	*	*	*	*	*	*	*	0.3	0.9
White, both sexes										
2002	24.1	*	*	*	*	*	*	0.2	0.6	1.7
2001	24.3	*	*	*	*	*	*	0.2	0.6	1.7
2000	25.1	*	*	*	*	*	*	*	0.5	1.6
1999	26.1	*	*	*	*	*	*	0.1	0.5	1.8
White, male										
2002	25.5	*	*	*	*	*	*	0.3	1.0	2.7
2001	25.3	*	*	*	*	*	*	0.3	0.9	2.7
2000	26.0	*	*	*	*	*	*	*	0.8	2.5
1999	26.6	*	*	*	*	*	*	*	0.8	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
White, female										
2002	22.7	*	*	*	*	*	*	*	*	0.7
2001	23.3	*	*	*	*	*	*	*	0.3	0.7
2000	24.3	*	*	*	*	*	*	*	*	0.6
1999	25.6	*	*	*	*	*	*	*	*	0.7
Black, both sexes										
2002	27.0	*	*	*	*	*	*	*	1.3	3.5
2001	27.7	*	*	*	*	*	*	*	1.0	3.0
2000	28.5	*	*	*	*	*	*	*	0.9	2.9
1999	29.9	*	*	*	*	*	*	*	1.2	3.7
Black, male										
2002	31.1	*	*	*	*	*	*	*	1.9	5.4
2001	31.5	*	*	*	*	*	*	*	*	4.6
2000	32.3	*	*	*	*	*	*	*	*	4.7
1999	33.8	*	*	*	*	*	*	*	1.7	5.8
Black, female										
2002	23.3	*	*	*	*	*	*	*	*	1.8
2001	24.3	*	*	*	*	*	*	*	*	1.6
2000	25.0	*	*	*	*	*	*	*	*	1.4
1999	26.4	*	*	*	*	*	*	*	*	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2002	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
1999	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3
All races 3/, male										
2002	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
1999	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3
All races 3/, female										
2002	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000	88.4	*	*	*	*	*	*	*	0.8	2.1
1999	91.6	*	*	*	*	*	*	0.3	0.7	2.2
White, both sexes										
2002	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
1999	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0
White, male										
2002	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000	97.5	*	*	*	*	*	*	0.7	2.1	5.6
1999	99.9	*	*	*	*	*	*	0.7	2.1	6.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, female										
2002	91.7	*	*	*	*	*	*	*	0.6	1.9
2001	94.9	*	*	*	*	*	*	*	0.6	1.7
2000	97.3	*	*	*	*	*	*	*	0.6	1.8
1999	100.8	*	*	*	*	*	*	0.3	0.6	2.0
Black, both sexes										
2002	56.2	*	*	*	*	*	*	0.9	1.9	6.1
2001	58.0	*	*	*	*	*	*	1.2	2.8	5.8
2000	58.1	*	*	*	*	*	*	0.8	2.8	6.5
1999	59.3	*	*	*	*	*	*	0.8	2.2	6.4
Black, male										
2002	54.5	*	*	*	*	*	*	*	2.3	9.0
2001	56.7	*	*	*	*	*	*	1.8	4.6	8.7
2000	55.8	*	*	*	*	*	*	*	3.8	8.7
1999	57.3	*	*	*	*	*	*	*	3.3	9.2
Black, female										
2002	57.7	*	*	*	*	*	*	*	1.6	3.6
2001	59.2	*	*	*	*	*	*	*	*	3.3
2000	60.3	*	*	*	*	*	*	*	1.9	4.5
1999	61.2	*	*	*	*	*	*	*	*	4.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2002	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5
1999	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2
All races 3/, male										
2002	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1
1999	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3
All races 3/, female										
2002	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9
1999	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1
White, both sexes										
2002	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000	62.5	10.6	1.0	0.4	0.6	1.4	2.1	2.8	4.3	6.1
1999	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7
White, male										
2002	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000	54.6	10.8	1.2	0.5	0.8	1.7	2.5	3.4	5.2	7.5
1999	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
White, female										
2002	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000	70.2	10.4	0.7	0.3	0.4	1.2	1.6	2.2	3.3	4.6
1999	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8
Black, both sexes										
2002	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000	52.7	19.0	1.9	0.8	1.4	3.6	6.0	8.8	11.9	17.8
1999	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5
Black, male										
2002	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000	51.5	18.9	1.9	*	1.4	4.3	6.7	10.1	14.3	21.6
1999	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3
Black, female										
2002	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000	53.9	19.2	2.0	*	1.3	2.9	5.3	7.6	9.8	14.5
1999	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute and subacute endocarditis (I33)										
All races 3/, both sexes										
2002	0.4	*	*	*	*	*	*	*	0.1	0.1
2001	0.4	*	*	*	*	*	*	*	0.1	0.2
2000	0.4	*	*	*	*	*	*	*	0.1	0.2
1999	0.4	*	*	*	*	*	*	*	*	0.2
All races 3/, male										
2002	0.5	*	*	*	*	*	*	*	0.2	*
2001	0.5	*	*	*	*	*	*	*	*	0.2
2000	0.5	*	*	*	*	*	*	*	*	0.2
1999	0.5	*	*	*	*	*	*	*	*	0.3
All races 3/, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	0.2
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002	0.4	*	*	*	*	*	*	*	0.1	0.1
2001	0.4	*	*	*	*	*	*	*	*	0.2
2000	0.4	*	*	*	*	*	*	*	*	0.1
1999	0.4	*	*	*	*	*	*	*	*	0.2
White, male										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	0.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute and subacute endocarditis (I33)										
White, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	0.7
Black, male										
2002	0.6	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All races 3/, both sexes										
2002	0.3	0.6	0.1	*	*	*	*	*	0.1	0.2
2001	0.3	0.8	0.2	*	*	*	0.1	0.1	0.1	0.2
2000	0.3	0.5	0.2	*	*	*	0.1	0.1	0.1	0.2
1999	0.3	0.7	0.2	*	*	*	0.1	*	0.2	0.2
All races 3/, male										
2002	0.3	*	*	*	*	*	*	*	0.2	*
2001	0.3	*	*	*	*	*	*	*	*	0.2
2000	0.3	*	*	*	*	*	*	*	*	0.3
1999	0.3	*	*	*	*	*	*	*	0.2	0.3
All races 3/, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	0.1
2001	0.3	*	*	*	*	*	*	*	0.1	0.2
2000	0.3	*	*	*	*	*	*	0.1	*	0.1
1999	0.3	*	*	*	*	*	*	*	0.1	0.2
White, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
White, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Heart failure (I50)										
All races 3/, both sexes										
2002	19.6	0.6	*	*	*	*	*	0.2	0.3	0.5
2001	20.0	0.6	*	*	*	*	*	0.2	0.2	0.5
2000	19.8	0.8	*	*	*	0.1	*	0.1	0.2	0.3
1999	19.7	*	*	*	*	*	*	0.1	0.2	0.4
All races 3/, male										
2002	15.3	*	*	*	*	*	*	0.2	0.4	0.5
2001	15.5	*	*	*	*	*	*	0.3	0.3	0.7
2000	15.3	1.1	*	*	*	*	*	*	0.2	0.4
1999	15.2	*	*	*	*	*	*	*	*	0.5
All races 3/, female										
2002	23.7	*	*	*	*	*	*	*	0.3	0.4
2001	24.3	*	*	*	*	*	*	*	0.2	0.4
2000	24.1	*	*	*	*	*	*	*	*	0.3
1999	24.0	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	21.6	*	*	*	*	*	*	*	0.2	0.3
2001	22.1	*	*	*	*	*	*	0.2	0.2	0.4
2000	21.8	*	*	*	*	*	*	*	0.1	0.2
1999	21.6	*	*	*	*	*	*	*	0.1	0.3
White, male										
2002	16.7	*	*	*	*	*	*	*	*	0.4
2001	16.9	*	*	*	*	*	*	*	*	0.5
2000	16.7	*	*	*	*	*	*	*	*	0.3
1999	16.4	*	*	*	*	*	*	*	*	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Heart failure (I50)										
White, female										
2002	26.4	*	*	*	*	*	*	*	*	0.3
2001	27.1	*	*	*	*	*	*	*	*	0.3
2000	26.7	*	*	*	*	*	*	*	*	*
1999	26.6	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002	13.7	*	*	*	*	*	*	*	1.0	1.4
2001	13.8	*	*	*	*	*	*	*	*	1.2
2000	13.6	*	*	*	*	*	*	*	*	1.1
1999	13.8	*	*	*	*	*	*	*	*	0.9
Black, male										
2002	11.5	*	*	*	*	*	*	*	*	1.6
2001	11.6	*	*	*	*	*	*	*	*	1.9
2000	11.2	*	*	*	*	*	*	*	*	*
1999	11.8	*	*	*	*	*	*	*	*	*
Black, female										
2002	15.8	*	*	*	*	*	*	*	*	*
2001	15.8	*	*	*	*	*	*	*	*	*
2000	15.7	*	*	*	*	*	*	*	*	*
1999	15.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All races 3/, both sexes										
2002	38.5	10.5	0.8	0.3	0.6	1.6	2.1	3.5	4.6	7.3
2001	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0
2000	38.3	10.6	0.9	0.4	0.6	1.5	2.4	3.2	4.8	6.8
1999	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4
All races 3/, male										
2002	36.3	10.9	0.8	0.3	0.7	2.2	2.7	4.2	5.8	8.8
2001	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8
2000	36.1	10.5	1.0	0.5	0.8	1.9	2.8	3.8	5.7	8.3
1999	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3
All races 3/, female										
2002	40.6	10.0	0.8	0.4	0.4	1.0	1.5	2.7	3.4	5.9
2001	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3
2000	40.4	10.7	0.7	0.3	0.5	1.2	2.0	2.6	3.8	5.3
1999	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4
White, both sexes										
2002	40.4	9.1	0.7	0.3	0.5	1.4	1.7	2.7	3.6	6.1
2001	39.6	8.2	0.9	0.3	0.6	1.1	1.8	2.6	3.7	5.7
2000	40.0	9.6	0.8	0.3	0.5	1.3	1.9	2.5	3.9	5.5
1999	40.6	10.3	0.8	0.4	0.6	1.6	1.9	2.5	4.0	6.0
White, male										
2002	37.5	9.5	0.8	0.3	0.6	1.9	2.3	3.3	4.6	7.3
2001	36.5	8.2	0.9	0.3	0.6	1.3	2.3	3.1	4.4	7.2
2000	37.1	9.4	0.9	0.4	0.7	1.5	2.3	3.1	4.8	6.9
1999	37.9	10.7	1.0	0.4	0.6	1.9	2.2	3.0	4.9	7.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
White, female										
2002	43.2	8.7	0.6	0.3	0.4	0.9	1.1	2.1	2.6	4.8
2001	42.6	8.2	0.8	0.3	0.5	0.9	1.3	2.1	3.0	4.2
2000	42.8	9.7	0.7	*	0.3	1.0	1.5	1.9	3.0	4.2
1999	43.2	10.0	0.7	0.4	0.5	1.4	1.7	2.0	3.1	4.3
Black, both sexes										
2002	37.8	17.2	1.1	*	0.9	2.9	4.8	8.7	11.4	17.0
2001	38.0	17.2	2.3	0.6	1.1	2.6	4.9	7.9	11.2	16.3
2000	38.2	15.6	1.4	0.7	1.1	3.2	5.4	8.0	10.8	15.9
1999	39.6	20.2	1.4	0.6	1.1	3.3	5.2	7.8	11.4	17.5
Black, male										
2002	38.3	18.9	*	*	1.2	3.7	5.8	10.9	14.5	20.5
2001	38.9	15.0	2.6	*	1.1	3.4	6.3	9.3	13.5	20.3
2000	39.2	14.7	1.6	*	1.2	3.9	6.1	9.1	13.2	19.3
1999	41.5	19.8	*	*	1.5	4.6	6.2	9.2	13.8	21.8
Black, female										
2002	37.4	15.4	*	*	*	2.0	3.8	6.7	8.7	13.9
2001	37.2	19.5	2.0	*	*	1.7	3.5	6.6	9.2	12.8
2000	37.3	16.5	*	*	*	2.6	4.8	6.9	8.7	12.9
1999	37.9	20.5	*	*	*	2.0	4.2	6.5	9.2	13.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
All races 3/, both sexes										
2002	7.0	*	*	*	*	*	*	0.2	0.2	0.6
2001	6.8	*	*	*	*	*	*	0.2	0.3	0.5
2000	6.4	*	*	*	*	*	*	0.1	0.2	0.5
1999	6.1	*	*	*	*	*	*	0.1	0.3	0.5
All races 3/, male										
2002	5.4	*	*	*	*	*	*	0.2	0.2	0.7
2001	5.2	*	*	*	*	*	*	0.2	0.4	0.5
2000	5.0	*	*	*	*	*	*	*	0.3	0.5
1999	4.7	*	*	*	*	*	*	*	0.3	0.6
All races 3/, female										
2002	8.6	*	*	*	*	*	*	*	0.2	0.5
2001	8.3	*	*	*	*	*	*	*	0.3	0.4
2000	7.8	*	*	*	*	*	*	*	0.2	0.4
1999	7.4	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	6.6	*	*	*	*	*	*	*	0.1	0.2
2001	6.3	*	*	*	*	*	*	*	0.2	0.2
2000	6.0	*	*	*	*	*	*	*	0.1	0.2
1999	5.6	*	*	*	*	*	*	*	0.1	0.2
White, male										
2002	4.9	*	*	*	*	*	*	*	*	0.3
2001	4.7	*	*	*	*	*	*	*	0.2	0.3
2000	4.5	*	*	*	*	*	*	*	*	0.3
1999	4.2	*	*	*	*	*	*	*	0.2	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
White, female										
2002	8.3	*	*	*	*	*	*	*	*	0.2
2001	8.0	*	*	*	*	*	*	*	*	*
2000	7.4	*	*	*	*	*	*	*	*	*
1999	7.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	11.1	*	*	*	*	*	*	*	1.0	2.8
2001	10.9	*	*	*	*	*	*	*	1.3	1.8
2000	10.6	*	*	*	*	*	*	*	1.0	2.1
1999	10.3	*	*	*	*	*	*	*	1.0	2.2
Black, male										
2002	9.6	*	*	*	*	*	*	*	*	3.6
2001	9.7	*	*	*	*	*	*	*	*	2.2
2000	9.1	*	*	*	*	*	*	*	*	2.2
1999	9.1	*	*	*	*	*	*	*	*	2.7
Black, female										
2002	12.5	*	*	*	*	*	*	*	*	2.0
2001	11.9	*	*	*	*	*	*	*	1.4	1.5
2000	12.0	*	*	*	*	*	*	*	*	2.0
1999	11.4	*	*	*	*	*	*	*	*	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2002	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9
1999	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0
All races 3/, male										
2002	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8
1999	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9
All races 3/, female										
2002	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0
1999	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1
White, both sexes										
2002	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000	62.8	2.5	0.3	0.1	0.2	0.3	0.6	1.0	1.6	3.0
1999	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2
White, male										
2002	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000	48.4	3.0	*	*	*	0.4	0.7	1.1	1.6	3.0
1999	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
White, female										
2002	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000	76.9	2.0	*	*	0.3	*	0.6	0.9	1.5	3.0
1999	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3
Black, both sexes										
2002	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000	52.5	7.5	*	*	*	*	1.1	1.5	4.3	9.6
1999	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1
Black, male										
2002	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000	46.1	8.6	*	*	*	*	*	*	4.1	9.2
1999	45.9	*	*	*	*	*	*	1.9	3.7	8.9
Black, female										
2002	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000	58.3	*	*	*	*	*	*	1.6	4.6	10.0
1999	57.9	*	*	*	*	*	*	1.8	3.6	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerosis (I70)										
All races 3/, both sexes										
2002	4.8	*	*	*	*	*	*	*	*	*
2001	4.9	*	*	*	*	*	*	*	*	*
2000	5.1	*	*	*	*	*	*	*	*	*
1999	5.4	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	3.7	*	*	*	*	*	*	*	*	*
2001	3.7	*	*	*	*	*	*	*	*	*
2000	3.9	*	*	*	*	*	*	*	*	*
1999	4.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	5.9	*	*	*	*	*	*	*	*	*
2001	6.2	*	*	*	*	*	*	*	*	*
2000	6.3	*	*	*	*	*	*	*	*	*
1999	6.7	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	5.4	*	*	*	*	*	*	*	*	*
2001	5.6	*	*	*	*	*	*	*	*	*
2000	5.7	*	*	*	*	*	*	*	*	*
1999	6.0	*	*	*	*	*	*	*	*	*
White, male										
2002	4.0	*	*	*	*	*	*	*	*	*
2001	4.1	*	*	*	*	*	*	*	*	*
2000	4.3	*	*	*	*	*	*	*	*	*
1999	4.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerosis (I70)										
White, female										
2002	6.7	*	*	*	*	*	*	*	*	*
2001	7.0	*	*	*	*	*	*	*	*	*
2000	7.1	*	*	*	*	*	*	*	*	*
1999	7.6	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	2.7	*	*	*	*	*	*	*	*	*
2001	2.6	*	*	*	*	*	*	*	*	*
2000	2.9	*	*	*	*	*	*	*	*	*
1999	3.0	*	*	*	*	*	*	*	*	*
Black, male										
2002	2.3	*	*	*	*	*	*	*	*	*
2001	2.1	*	*	*	*	*	*	*	*	*
2000	2.4	*	*	*	*	*	*	*	*	*
1999	2.5	*	*	*	*	*	*	*	*	*
Black, female										
2002	3.1	*	*	*	*	*	*	*	*	*
2001	3.1	*	*	*	*	*	*	*	*	*
2000	3.3	*	*	*	*	*	*	*	*	*
1999	3.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of circulatory system (I71-I78)										
All races 3/, both sexes										
2002	8.6	*	*	*	*	0.2	0.2	0.3	0.5	0.7
2001	8.9	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000	9.3	*	*	*	*	*	0.2	0.3	0.5	0.7
1999	9.2	*	*	*	*	*	0.2	0.3	0.4	0.8
All races 3/, male										
2002	9.4	*	*	*	*	*	0.2	0.3	0.7	0.7
2001	9.6	*	*	*	*	*	0.2	0.4	0.7	1.2
2000	10.2	*	*	*	*	*	0.2	0.4	0.6	0.8
1999	10.1	*	*	*	*	*	0.2	0.4	0.6	1.0
All races 3/, female										
2002	7.9	*	*	*	*	*	*	0.2	0.4	0.7
2001	8.2	*	*	*	*	*	*	0.3	0.3	0.6
2000	8.3	*	*	*	*	*	*	*	0.4	0.6
1999	8.4	*	*	*	*	*	*	*	0.2	0.5
White, both sexes										
2002	9.3	*	*	*	*	0.2	0.1	0.2	0.4	0.7
2001	9.6	*	*	*	*	*	0.2	0.3	0.5	0.8
2000	10.0	*	*	*	*	*	0.2	0.2	0.5	0.6
1999	10.0	*	*	*	*	*	0.2	0.2	0.4	0.7
White, male										
2002	10.2	*	*	*	*	*	*	0.3	0.5	0.7
2001	10.4	*	*	*	*	*	*	0.5	0.6	1.0
2000	11.1	*	*	*	*	*	*	0.3	0.6	0.7
1999	11.0	*	*	*	*	*	*	0.3	0.6	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of circulatory system (I71-I78)										
White, female										
2002	8.5	*	*	*	*	*	*	*	0.3	0.6
2001	8.8	*	*	*	*	*	*	*	0.3	0.5
2000	8.9	*	*	*	*	*	*	*	0.3	0.6
1999	9.1	*	*	*	*	*	*	*	*	0.5
Black, both sexes										
2002	6.7	*	*	*	*	*	*	*	1.1	1.1
2001	7.1	*	*	*	*	*	*	*	0.9	1.7
2000	7.1	*	*	*	*	*	*	*	0.9	1.2
1999	6.7	*	*	*	*	*	*	*	*	1.2
Black, male										
2002	6.7	*	*	*	*	*	*	*	1.7	*
2001	6.9	*	*	*	*	*	*	*	*	2.4
2000	6.9	*	*	*	*	*	*	*	*	1.6
1999	6.5	*	*	*	*	*	*	*	*	1.5
Black, female										
2002	6.6	*	*	*	*	*	*	*	*	*
2001	7.2	*	*	*	*	*	*	*	*	*
2000	7.2	*	*	*	*	*	*	*	*	*
1999	7.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes										
2002	5.1	*	*	*	*	0.1	*	0.2	0.4	0.5
2001	5.3	*	*	*	*	*	0.1	0.2	0.3	0.6
2000	5.6	*	*	*	*	*	0.1	0.2	0.3	0.5
1999	5.7	*	*	*	*	*	0.1	0.2	0.3	0.6
All races 3/, male										
2002	6.4	*	*	*	*	*	*	0.3	0.5	0.6
2001	6.5	*	*	*	*	*	*	0.4	0.5	0.9
2000	7.0	*	*	*	*	*	*	0.3	0.5	0.6
1999	7.0	*	*	*	*	*	*	0.3	0.5	0.9
All races 3/, female										
2002	4.0	*	*	*	*	*	*	*	0.2	0.3
2001	4.2	*	*	*	*	*	*	*	*	0.3
2000	4.3	*	*	*	*	*	*	*	0.2	0.3
1999	4.4	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	5.7	*	*	*	*	*	*	0.1	0.3	0.5
2001	5.9	*	*	*	*	*	0.1	0.2	0.3	0.5
2000	6.2	*	*	*	*	*	*	0.2	0.3	0.4
1999	6.3	*	*	*	*	*	*	0.2	0.3	0.6
White, male										
2002	7.0	*	*	*	*	*	*	*	0.4	0.6
2001	7.2	*	*	*	*	*	*	0.4	0.5	0.8
2000	7.8	*	*	*	*	*	*	*	0.4	0.6
1999	7.8	*	*	*	*	*	*	*	0.5	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Aortic aneurysm and dissection (I71)										
White, female										
2002	4.3	*	*	*	*	*	*	*	*	0.3
2001	4.6	*	*	*	*	*	*	*	*	*
2000	4.6	*	*	*	*	*	*	*	*	0.3
1999	4.8	*	*	*	*	*	*	*	*	0.3
Black, both sexes										
2002	3.2	*	*	*	*	*	*	*	0.9	*
2001	3.4	*	*	*	*	*	*	*	*	1.1
2000	3.5	*	*	*	*	*	*	*	*	0.8
1999	3.3	*	*	*	*	*	*	*	*	0.8
Black, male										
2002	3.8	*	*	*	*	*	*	*	*	*
2001	3.8	*	*	*	*	*	*	*	*	1.7
2000	3.9	*	*	*	*	*	*	*	*	*
1999	3.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	2.7	*	*	*	*	*	*	*	*	*
2001	3.0	*	*	*	*	*	*	*	*	*
2000	3.2	*	*	*	*	*	*	*	*	*
1999	3.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes										
2002	3.5	*	*	*	*	*	*	0.1	0.1	0.2
2001	3.5	*	*	*	*	*	*	0.1	0.2	0.3
2000	3.6	*	*	*	*	*	*	*	0.2	0.2
1999	3.6	*	*	*	*	*	*	*	0.1	0.2
All races 3/, male										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	3.1	*	*	*	*	*	*	*	*	0.2
2000	3.2	*	*	*	*	*	*	*	*	0.2
1999	3.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	4.0	*	*	*	*	*	*	*	*	0.3
2001	4.0	*	*	*	*	*	*	*	*	0.3
2000	4.0	*	*	*	*	*	*	*	*	0.3
1999	4.1	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002	3.7	*	*	*	*	*	*	*	0.1	0.2
2001	3.7	*	*	*	*	*	*	*	0.1	0.2
2000	3.8	*	*	*	*	*	*	*	0.2	0.2
1999	3.8	*	*	*	*	*	*	*	0.1	0.2
White, male										
2002	3.2	*	*	*	*	*	*	*	*	*
2001	3.2	*	*	*	*	*	*	*	*	*
2000	3.4	*	*	*	*	*	*	*	*	*
1999	3.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of arteries, arterioles and capillaries (I72-I78)										
White, female										
2002	4.2	*	*	*	*	*	*	*	*	0.3
2001	4.2	*	*	*	*	*	*	*	*	0.3
2000	4.2	*	*	*	*	*	*	*	*	0.3
1999	4.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	3.5	*	*	*	*	*	*	*	*	*
2001	3.7	*	*	*	*	*	*	*	*	*
2000	3.6	*	*	*	*	*	*	*	*	*
1999	3.5	*	*	*	*	*	*	*	*	*
Black, male										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	3.2	*	*	*	*	*	*	*	*	*
2000	3.1	*	*	*	*	*	*	*	*	*
1999	2.9	*	*	*	*	*	*	*	*	*
Black, female										
2002	3.9	*	*	*	*	*	*	*	*	*
2001	4.2	*	*	*	*	*	*	*	*	*
2000	4.1	*	*	*	*	*	*	*	*	*
1999	4.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2002	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4
All races 3/, male										
2002	1.4	*	*	*	*	*	*	*	0.5	0.6
2001	1.4	*	*	*	*	*	*	*	0.3	0.5
2000	1.4	*	*	*	*	*	*	*	0.4	0.6
1999	1.3	*	*	*	*	*	*	*	0.3	0.5
All races 3/, female										
2002	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000	1.8	*	*	*	*	*	*	0.4	0.4	0.5
1999	1.6	*	*	*	*	*	*	*	0.3	0.4
White, both sexes										
2002	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
1999	1.5	*	*	*	*	*	*	0.1	0.3	0.4
White, male										
2002	1.3	*	*	*	*	*	*	*	0.4	0.5
2001	1.4	*	*	*	*	*	*	*	0.3	0.4
2000	1.4	*	*	*	*	*	*	*	0.3	0.5
1999	1.2	*	*	*	*	*	*	*	0.2	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of circulatory system (I80-I99)										
White, female										
2002	1.9	*	*	*	*	*	*	*	0.3	0.4
2001	1.9	*	*	*	*	*	*	*	0.3	0.3
2000	1.9	*	*	*	*	*	*	0.3	0.3	0.4
1999	1.7	*	*	*	*	*	*	*	0.3	0.3
Black, both sexes										
2002	2.1	*	*	*	*	*	*	*	1.4	1.5
2001	2.0	*	*	*	*	*	*	*	1.0	1.2
2000	2.1	*	*	*	*	*	*	0.9	0.9	1.2
1999	1.7	*	*	*	*	*	*	*	*	1.0
Black, male										
2002	2.0	*	*	*	*	*	*	*	1.6	1.5
2001	1.9	*	*	*	*	*	*	*	*	*
2000	1.9	*	*	*	*	*	*	*	*	1.6
1999	1.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	2.2	*	*	*	*	*	*	*	*	1.4
2001	2.1	*	*	*	*	*	*	*	*	*
2000	2.2	*	*	*	*	*	*	*	*	*
1999	1.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Influenza and pneumonia (J10-J18)										
All races 3/, both sexes										
2002	22.8	6.5	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.7
2001	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8
2000	23.2	7.6	0.7	0.2	0.2	0.3	0.7	0.7	1.1	1.8
1999	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0
All races 3/, male										
2002	20.4	7.8	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0
2000	20.8	8.4	0.7	0.3	0.2	0.3	0.7	0.8	1.2	1.9
1999	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2
All races 3/, female										
2002	25.1	5.1	0.6	0.2	0.3	0.3	0.4	0.7	0.8	1.6
2001	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6
2000	25.6	6.8	0.7	0.2	*	0.3	0.6	0.6	1.0	1.7
1999	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8
White, both sexes										
2002	24.9	5.2	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.5
2001	23.6	5.3	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2000	25.2	5.8	0.6	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999	24.8	6.2	0.6	0.2	0.2	0.3	0.5	0.5	0.8	1.7
White, male										
2002	21.9	6.6	0.6	*	*	0.4	0.5	0.7	1.0	1.5
2001	20.7	6.2	0.7	*	*	0.3	0.7	0.5	1.0	1.7
2000	22.0	6.4	0.5	0.3	*	0.3	0.7	0.6	1.0	1.7
1999	21.5	6.7	0.5	0.3	0.3	0.3	0.5	0.6	1.0	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Influenza and pneumonia (J10-J18)										
White, female										
2002	27.8	3.7	0.5	*	0.3	0.3	0.4	0.7	0.7	1.4
2001	26.4	4.4	0.6	*	*	0.3	0.3	0.6	0.8	1.4
2000	28.2	5.1	0.7	*	*	0.3	0.5	0.6	0.9	1.6
1999	27.9	5.6	0.7	*	*	0.4	0.4	0.5	0.6	1.6
Black, both sexes										
2002	15.6	12.9	1.6	*	*	*	0.7	1.0	2.0	3.5
2001	15.5	16.3	1.0	*	*	*	1.0	1.2	2.1	3.7
2000	16.4	16.7	1.2	*	*	*	1.1	1.4	2.0	2.9
1999	16.2	19.0	1.7	*	*	*	1.3	1.7	2.2	4.1
Black, male										
2002	15.4	13.9	1.7	*	*	*	*	*	2.4	4.0
2001	15.9	15.9	*	*	*	*	*	*	2.7	4.5
2000	16.7	17.9	1.7	*	*	*	*	2.1	2.5	3.0
1999	16.4	18.9	2.0	*	*	*	*	1.9	2.5	4.5
Black, female										
2002	15.7	11.8	*	*	*	*	*	*	1.6	3.1
2001	15.2	16.7	*	*	*	*	*	*	1.4	3.0
2000	16.0	15.5	*	*	*	*	*	*	1.6	2.7
1999	16.1	19.1	*	*	*	*	*	1.5	1.9	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Influenza (J10-J11)										
All races 3/, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.8	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
White, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Influenza (J10-J11)										
White, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.9	*	*	*	*	*	*	*	*	*
1999	0.9	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
All races 3/, both sexes										
2002	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7
1999	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0
All races 3/, male										
2002	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8
1999	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2
All races 3/, female										
2002	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7
1999	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8
White, both sexes										
2002	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000	24.4	5.6	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7
White, male										
2002	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000	21.5	6.1	0.4	0.3	*	0.3	0.6	0.6	1.0	1.7
1999	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
White, female										
2002	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000	27.3	5.0	0.6	*	*	*	0.5	0.6	0.9	1.5
1999	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6
Black, both sexes										
2002	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000	16.1	16.4	1.2	*	*	*	1.0	1.4	2.0	2.9
1999	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0
Black, male										
2002	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000	16.5	17.9	1.7	*	*	*	*	2.0	2.5	3.0
1999	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4
Black, female										
2002	15.6	11.8	*	*	*	*	*	*	1.6	3.1
2001	15.1	16.7	*	*	*	*	*	*	1.4	3.0
2000	15.8	14.9	*	*	*	*	*	*	1.6	2.7
1999	15.9	18.8	*	*	*	*	*	1.5	1.9	3.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other acute lower respiratory infections (J20-J22)										
All races 3/, both sexes										
2002	0.1	1.3	0.1	*	*	*	*	*	*	*
2001	0.1	1.2	*	*	*	*	*	*	*	*
2000	0.2	1.1	*	*	*	*	*	*	*	*
1999	0.2	1.7	0.2	*	*	*	*	*	*	*
All races 3/, male										
2002	0.1	1.9	*	*	*	*	*	*	*	*
2001	0.1	1.3	*	*	*	*	*	*	*	*
2000	0.1	1.3	*	*	*	*	*	*	*	*
1999	0.1	1.9	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	1.2	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	1.5	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	1.1	0.2	*	*	*	*	*	*	*
2001	0.1	1.0	*	*	*	*	*	*	*	*
2000	0.2	0.8	*	*	*	*	*	*	*	*
1999	0.2	1.2	*	*	*	*	*	*	*	*
White, male										
2002	0.1	1.6	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other acute lower respiratory infections (J20-J22)										
White, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	4.0	*	*	*	*	*	*	*	*
Black, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute bronchitis and bronchiolitis (J20-J21)										
All races 3/, both sexes										
2002	0.1	1.2	0.1	*	*	*	*	*	*	*
2001	0.1	1.2	*	*	*	*	*	*	*	*
2000	0.1	1.1	*	*	*	*	*	*	*	*
1999	0.1	1.6	0.2	*	*	*	*	*	*	*
All races 3/, male										
2002	0.1	1.8	*	*	*	*	*	*	*	*
2001	0.1	1.2	*	*	*	*	*	*	*	*
2000	0.1	1.2	*	*	*	*	*	*	*	*
1999	0.1	1.9	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	1.2	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	1.4	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	1.0	0.2	*	*	*	*	*	*	*
2001	0.1	1.0	*	*	*	*	*	*	*	*
2000	0.1	0.8	*	*	*	*	*	*	*	*
1999	0.1	1.2	*	*	*	*	*	*	*	*
White, male										
2002	0.1	1.5	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute bronchitis and bronchiolitis (J20-J21)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	4.0	*	*	*	*	*	*	*	*
Black, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified acute lower respiratory infection (J22)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified acute lower respiratory infection (J22)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2002	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
1999	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3
All races 3/, male										
2002	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3
1999	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2
All races 3/, female										
2002	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4
1999	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4
White, both sexes										
2002	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000	49.0	0.8	0.2	0.1	0.2	0.3	0.4	0.4	0.6	1.0
1999	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1
White, male										
2002	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001	47.6	*	*	*	*	*	0.3	0.4	0.5	1.2
2000	48.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
White, female										
2002	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000	49.7	*	*	*	*	*	0.3	0.4	0.6	1.2
1999	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2
Black, both sexes										
2002	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000	20.8	*	0.8	0.7	1.4	1.4	1.3	1.9	2.4	3.5
1999	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2
Black, male										
2002	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000	24.3	*	*	*	1.7	2.0	1.5	2.4	3.1	3.7
1999	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3
Black, female										
2002	17.7	*	*	*	1.4	*	*	*	2.1	3.2
2001	17.5	*	*	*	*	*	1.4	1.5	2.6	4.7
2000	17.6	*	*	*	*	*	*	1.4	1.7	3.3
1999	18.0	*	*	*	*	*	*	2.4	2.8	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified (J40-J42)										
All races 3/, both sexes										
2002	0.3	0.6	*	*	*	*	*	*	*	*
2001	0.3	0.5	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	0.5	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
White, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified (J40-J42)										
White, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Emphysema (J43)										
All races 3/, both sexes										
2002	5.4	*	*	*	*	*	*	*	*	0.1
2001	5.7	*	*	*	*	*	*	*	*	0.1
2000	5.9	*	*	*	*	*	*	*	*	0.1
1999	6.4	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	5.7	*	*	*	*	*	*	*	*	*
2001	6.0	*	*	*	*	*	*	*	*	*
2000	6.3	*	*	*	*	*	*	*	*	*
1999	6.9	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	5.0	*	*	*	*	*	*	*	*	*
2001	5.4	*	*	*	*	*	*	*	*	*
2000	5.6	*	*	*	*	*	*	*	*	*
1999	5.8	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	6.2	*	*	*	*	*	*	*	*	*
2001	6.6	*	*	*	*	*	*	*	*	0.1
2000	6.8	*	*	*	*	*	*	*	*	*
1999	7.3	*	*	*	*	*	*	*	*	*
White, male										
2002	6.4	*	*	*	*	*	*	*	*	*
2001	6.8	*	*	*	*	*	*	*	*	*
2000	7.0	*	*	*	*	*	*	*	*	*
1999	7.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Emphysema (J43)										
White, female										
2002	5.9	*	*	*	*	*	*	*	*	*
2001	6.3	*	*	*	*	*	*	*	*	*
2000	6.5	*	*	*	*	*	*	*	*	*
1999	6.8	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	2.3	*	*	*	*	*	*	*	*	*
2001	2.2	*	*	*	*	*	*	*	*	*
2000	2.5	*	*	*	*	*	*	*	*	*
1999	2.6	*	*	*	*	*	*	*	*	*
Black, male										
2002	3.0	*	*	*	*	*	*	*	*	*
2001	2.7	*	*	*	*	*	*	*	*	*
2000	3.2	*	*	*	*	*	*	*	*	*
1999	3.5	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.6	*	*	*	*	*	*	*	*	*
2001	1.7	*	*	*	*	*	*	*	*	*
2000	1.8	*	*	*	*	*	*	*	*	*
1999	1.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
All races 3/, both sexes										
2002	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
1999	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9
All races 3/, male										
2002	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8
All races 3/, female										
2002	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
1999	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0
White, both sexes										
2002	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7
White, male										
2002	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
White, female										
2002	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001	1.7	*	*	*	*	*	*	*	0.6	0.6
2000	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8
Black, both sexes										
2002	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
1999	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6
Black, male										
2002	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
1999	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6
Black, female										
2002	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000	3.5	*	*	*	*	*	*	*	1.4	2.6
1999	3.5	*	*	*	*	*	*	2.0	2.3	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic lower respiratory diseases (J44,J47)										
All races 3/, both sexes										
2002	36.1	*	*	*	*	*	*	*	0.2	0.5
2001	35.7	*	*	*	*	*	*	*	0.2	0.5
2000	35.4	*	*	*	*	*	*	*	*	0.4
1999	36.0	*	*	*	*	*	*	0.1	0.1	0.3
All races 3/, male										
2002	35.7	*	*	*	*	*	*	*	0.2	0.4
2001	35.3	*	*	*	*	*	*	*	0.2	0.5
2000	35.6	*	*	*	*	*	*	*	*	0.4
1999	37.1	*	*	*	*	*	*	*	*	0.3
All races 3/, female										
2002	36.5	*	*	*	*	*	*	*	0.2	0.5
2001	36.0	*	*	*	*	*	*	*	*	0.5
2000	35.2	*	*	*	*	*	*	*	*	0.5
1999	35.0	*	*	*	*	*	*	*	*	0.3
White, both sexes										
2002	41.3	*	*	*	*	*	*	*	0.1	0.4
2001	40.8	*	*	*	*	*	*	*	0.2	0.5
2000	40.4	*	*	*	*	*	*	*	*	0.4
1999	41.0	*	*	*	*	*	*	*	*	0.3
White, male										
2002	40.2	*	*	*	*	*	*	*	*	0.4
2001	39.7	*	*	*	*	*	*	*	*	0.5
2000	39.9	*	*	*	*	*	*	*	*	0.4
1999	41.5	*	*	*	*	*	*	*	*	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic lower respiratory diseases (J44,J47)										
White, female										
2002	42.5	*	*	*	*	*	*	*	*	0.5
2001	41.8	*	*	*	*	*	*	*	*	0.4
2000	40.9	*	*	*	*	*	*	*	*	0.4
1999	40.4	*	*	*	*	*	*	*	*	0.2
Black, both sexes										
2002	15.4	*	*	*	*	*	*	*	*	0.8
2001	15.1	*	*	*	*	*	*	*	*	1.1
2000	14.9	*	*	*	*	*	*	*	*	0.7
1999	15.9	*	*	*	*	*	*	*	*	*
Black, male										
2002	18.2	*	*	*	*	*	*	*	*	*
2001	18.1	*	*	*	*	*	*	*	*	*
2000	18.2	*	*	*	*	*	*	*	*	*
1999	19.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	12.9	*	*	*	*	*	*	*	*	*
2001	12.3	*	*	*	*	*	*	*	*	*
2000	12.0	*	*	*	*	*	*	*	*	*
1999	12.5	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumoconioses and chemical effects (J60-J66,J68)										
All races 3/, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.8	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*
White, male										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	0.9	*	*	*	*	*	*	*	*	*
2000	0.9	*	*	*	*	*	*	*	*	*
1999	0.9	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumoconioses and chemical effects (J60-J66,J68)										
White, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes										
2002	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2001	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2000	5.9	*	*	*	*	*	0.1	0.1	0.2	0.3
1999	5.5	*	*	*	*	*	0.1	0.2	0.2	0.3
All races 3/, male										
2002	6.3	*	*	*	*	*	*	*	0.2	0.4
2001	6.1	*	*	*	*	*	*	*	0.2	0.4
2000	6.1	*	*	*	*	*	*	*	0.2	0.4
1999	5.7	*	*	*	*	*	*	0.2	0.3	0.3
All races 3/, female										
2002	5.9	*	*	*	*	*	*	*	*	0.3
2001	6.0	*	*	*	*	*	*	*	0.2	*
2000	5.8	*	*	*	*	*	*	*	*	0.2
1999	5.3	*	*	*	*	*	*	*	*	0.2
White, both sexes										
2002	6.7	*	*	*	*	*	0.1	0.2	0.2	0.3
2001	6.6	*	*	*	*	*	*	0.2	0.2	0.2
2000	6.5	*	*	*	*	*	*	*	0.2	0.2
1999	5.9	*	*	*	*	*	*	0.1	0.2	0.2
White, male										
2002	6.9	*	*	*	*	*	*	*	*	0.3
2001	6.7	*	*	*	*	*	*	*	0.2	0.3
2000	6.6	*	*	*	*	*	*	*	*	0.3
1999	6.1	*	*	*	*	*	*	*	0.2	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonitis due to solids and liquids (J69)										
White, female										
2002	6.6	*	*	*	*	*	*	*	*	0.3
2001	6.6	*	*	*	*	*	*	*	*	*
2000	6.3	*	*	*	*	*	*	*	*	*
1999	5.8	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	4.1	*	*	*	*	*	*	*	*	*
2001	4.3	*	*	*	*	*	*	*	*	*
2000	4.2	*	*	*	*	*	*	*	*	0.7
1999	4.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	4.2	*	*	*	*	*	*	*	*	*
2001	4.2	*	*	*	*	*	*	*	*	*
2000	4.4	*	*	*	*	*	*	*	*	*
1999	4.5	*	*	*	*	*	*	*	*	*
Black, female										
2002	3.9	*	*	*	*	*	*	*	*	*
2001	4.4	*	*	*	*	*	*	*	*	*
2000	3.9	*	*	*	*	*	*	*	*	*
1999	3.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
All races 3/, both sexes										
2002	8.7	6.9	0.5	0.2	0.2	0.3	0.4	0.4	0.5	0.9
2001	9.2	7.3	0.6	0.1	0.2	0.3	0.4	0.5	0.6	1.0
2000	9.1	7.2	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0
1999	9.0	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0
All races 3/, male										
2002	8.5	7.9	0.7	0.2	*	0.3	0.4	0.3	0.5	0.9
2001	8.9	9.0	0.6	*	0.2	0.3	0.5	0.5	0.6	0.9
2000	9.0	7.9	0.6	0.2	*	0.3	0.4	0.5	0.8	1.2
1999	8.9	7.5	0.7	0.2	*	0.4	0.3	0.5	0.8	1.1
All races 3/, female										
2002	8.8	5.8	0.4	0.2	0.2	0.2	0.4	0.4	0.6	0.8
2001	9.4	5.5	0.6	*	0.2	0.3	0.4	0.6	0.6	1.0
2000	9.2	6.4	0.6	*	0.2	0.3	0.3	0.5	0.7	0.9
1999	9.1	5.8	0.5	*	0.2	0.2	0.3	0.5	0.8	0.9
White, both sexes										
2002	9.5	4.8	0.5	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001	10.0	5.0	0.5	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000	9.9	5.6	0.5	0.2	0.2	0.2	0.3	0.5	0.7	0.8
1999	9.7	4.9	0.5	0.2	0.2	0.3	0.3	0.4	0.7	0.8
White, male										
2002	9.2	5.4	0.6	*	*	*	0.4	0.3	0.5	0.8
2001	9.7	5.9	0.5	*	0.2	0.3	0.4	0.3	0.5	0.8
2000	9.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9
1999	9.6	5.6	0.6	*	*	0.4	0.4	0.5	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
White, female										
2002	9.7	4.2	*	*	*	*	*	0.4	0.5	0.7
2001	10.2	4.0	0.6	*	*	*	0.4	0.5	0.5	0.7
2000	10.0	4.6	0.6	*	*	0.3	*	0.5	0.6	0.7
1999	9.8	4.3	0.5	*	*	*	*	0.3	0.7	0.7
Black, both sexes										
2002	6.0	16.6	1.1	*	*	*	0.9	*	1.0	1.8
2001	6.7	18.0	1.0	*	*	*	0.8	1.1	1.4	2.5
2000	6.5	16.7	0.8	*	*	*	0.8	0.8	1.3	2.4
1999	6.5	15.4	1.0	*	*	*	*	1.1	1.6	2.3
Black, male										
2002	6.0	19.5	*	*	*	*	*	*	*	1.9
2001	6.4	23.1	*	*	*	*	*	*	1.5	2.0
2000	6.5	16.3	*	*	*	*	*	*	*	3.0
1999	6.4	17.6	*	*	*	*	*	*	1.7	2.5
Black, female										
2002	6.0	13.6	*	*	*	*	*	*	*	1.6
2001	7.0	12.6	*	*	*	*	*	*	*	2.9
2000	6.4	17.2	*	*	*	*	*	*	*	1.9
1999	6.7	13.1	*	*	*	*	*	1.7	1.6	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Peptic ulcer (K25-K28)										
All races 3/, both sexes										
2002	1.4	*	*	*	*	*	*	*	*	0.2
2001	1.6	*	*	*	*	*	*	*	*	0.2
2000	1.6	*	*	*	*	*	*	*	*	0.2
1999	1.6	*	*	*	*	*	*	*	*	0.2
All races 3/, male										
2002	1.4	*	*	*	*	*	*	*	*	0.3
2001	1.5	*	*	*	*	*	*	*	*	0.2
2000	1.6	*	*	*	*	*	*	*	*	0.3
1999	1.6	*	*	*	*	*	*	*	*	0.2
All races 3/, female										
2002	1.5	*	*	*	*	*	*	*	*	*
2001	1.7	*	*	*	*	*	*	*	*	*
2000	1.6	*	*	*	*	*	*	*	*	*
1999	1.7	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.5	*	*	*	*	*	*	*	*	0.2
2001	1.7	*	*	*	*	*	*	*	*	*
2000	1.7	*	*	*	*	*	*	*	*	0.2
1999	1.8	*	*	*	*	*	*	*	*	0.1
White, male										
2002	1.5	*	*	*	*	*	*	*	*	0.2
2001	1.5	*	*	*	*	*	*	*	*	*
2000	1.7	*	*	*	*	*	*	*	*	0.2
1999	1.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Peptic ulcer (K25-K28)										
White, female										
2002	1.6	*	*	*	*	*	*	*	*	*
2001	1.9	*	*	*	*	*	*	*	*	*
2000	1.8	*	*	*	*	*	*	*	*	*
1999	1.9	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.4	*	*	*	*	*	*	*	*	*
2000	1.3	*	*	*	*	*	*	*	*	*
1999	1.4	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.9	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of appendix (K35-K38)										
All races 3/, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of appendix (K35-K38)										
White, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hernia (K40-K46)										
All races 3/, both sexes										
2002	0.6	0.7	*	*	*	*	*	*	*	*
2001	0.5	1.1	*	*	*	*	*	*	*	*
2000	0.5	1.2	*	*	*	*	*	*	*	*
1999	0.5	1.2	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	1.4	*	*	*	*	*	*	*	*
2000	0.4	1.4	*	*	*	*	*	*	*	*
1999	0.4	1.3	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.6	0.8	*	*	*	*	*	*	*	*
2001	0.6	1.0	*	*	*	*	*	*	*	*
2000	0.6	1.2	*	*	*	*	*	*	*	*
1999	0.6	1.1	*	*	*	*	*	*	*	*
White, male										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	1.5	*	*	*	*	*	*	*	*
1999	0.5	1.3	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hernia (K40-K46)										
White, female										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic liver disease and cirrhosis (K70,K73-K74)										
All races 3/, both sexes										
2002	9.5	*	*	*	*	*	0.1	0.4	1.4	4.2
2001	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0
2000	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8
1999	9.4	*	*	*	*	*	*	0.4	1.6	4.8
All races 3/, male										
2002	12.3	*	*	*	*	*	*	0.5	1.8	5.5
2001	12.4	*	*	*	*	*	*	0.6	2.0	6.4
2000	12.5	*	*	*	*	*	*	0.6	2.1	6.3
1999	12.5	*	*	*	*	*	*	0.4	2.1	6.3
All races 3/, female										
2002	6.7	*	*	*	*	*	*	0.3	1.0	3.0
2001	6.7	*	*	*	*	*	*	0.3	1.0	3.5
2000	6.5	*	*	*	*	*	*	0.3	1.1	3.2
1999	6.4	*	*	*	*	*	*	0.3	1.2	3.4
White, both sexes										
2002	10.1	*	*	*	*	*	*	0.4	1.4	4.3
2001	10.1	*	*	*	*	*	*	0.3	1.5	5.1
2000	10.0	*	*	*	*	*	*	0.4	1.5	4.7
1999	9.9	*	*	*	*	*	*	0.3	1.6	4.8
White, male										
2002	13.2	*	*	*	*	*	*	0.4	1.9	5.6
2001	13.1	*	*	*	*	*	*	0.5	2.1	6.5
2000	13.2	*	*	*	*	*	*	0.6	2.1	6.3
1999	13.1	*	*	*	*	*	*	0.4	2.1	6.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic liver disease and cirrhosis (K70,K73-K74)										
White, female										
2002	7.2	*	*	*	*	*	*	0.3	0.9	3.0
2001	7.1	*	*	*	*	*	*	*	1.0	3.7
2000	6.8	*	*	*	*	*	*	*	1.0	3.1
1999	6.8	*	*	*	*	*	*	*	1.1	3.3
Black, both sexes										
2002	6.9	*	*	*	*	*	*	*	1.3	3.0
2001	7.5	*	*	*	*	*	*	*	1.0	4.0
2000	7.5	*	*	*	*	*	*	0.8	1.3	4.9
1999	7.8	*	*	*	*	*	*	*	1.4	4.4
Black, male										
2002	9.0	*	*	*	*	*	*	*	*	3.7
2001	10.4	*	*	*	*	*	*	*	*	5.7
2000	9.9	*	*	*	*	*	*	*	*	6.6
1999	10.7	*	*	*	*	*	*	*	1.8	5.4
Black, female										
2002	4.9	*	*	*	*	*	*	*	*	2.4
2001	4.8	*	*	*	*	*	*	*	*	2.6
2000	5.2	*	*	*	*	*	*	*	1.4	3.3
1999	5.2	*	*	*	*	*	*	*	*	3.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2002	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000	4.3	*	*	*	*	*	*	0.3	1.1	3.3
1999	4.3	*	*	*	*	*	*	0.2	1.1	3.3
All races 3/, male										
2002	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000	6.5	*	*	*	*	*	*	0.5	1.5	4.3
1999	6.5	*	*	*	*	*	*	0.2	1.4	4.3
All races 3/, female										
2002	2.3	*	*	*	*	*	*	*	0.7	2.0
2001	2.3	*	*	*	*	*	*	*	0.7	2.3
2000	2.2	*	*	*	*	*	*	*	0.6	2.2
1999	2.1	*	*	*	*	*	*	*	0.8	2.4
White, both sexes										
2002	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000	4.5	*	*	*	*	*	*	0.3	1.1	3.2
1999	4.4	*	*	*	*	*	*	0.2	1.0	3.3
White, male										
2002	6.6	*	*	*	*	*	*	0.3	1.3	3.7
2001	6.7	*	*	*	*	*	*	0.3	1.5	4.5
2000	6.8	*	*	*	*	*	*	0.5	1.6	4.3
1999	6.7	*	*	*	*	*	*	0.3	1.4	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
White, female										
2002	2.3	*	*	*	*	*	*	*	0.6	2.0
2001	2.3	*	*	*	*	*	*	*	0.7	2.3
2000	2.1	*	*	*	*	*	*	*	0.5	2.2
1999	2.1	*	*	*	*	*	*	*	0.7	2.3
Black, both sexes										
2002	3.2	*	*	*	*	*	*	*	0.7	1.6
2001	3.6	*	*	*	*	*	*	*	0.7	2.6
2000	3.8	*	*	*	*	*	*	*	1.0	3.2
1999	4.1	*	*	*	*	*	*	*	0.9	2.8
Black, male										
2002	4.4	*	*	*	*	*	*	*	*	2.0
2001	5.3	*	*	*	*	*	*	*	*	3.7
2000	5.3	*	*	*	*	*	*	*	*	4.4
1999	5.9	*	*	*	*	*	*	*	*	3.5
Black, female										
2002	2.0	*	*	*	*	*	*	*	*	1.3
2001	2.1	*	*	*	*	*	*	*	*	1.5
2000	2.4	*	*	*	*	*	*	*	*	2.1
1999	2.4	*	*	*	*	*	*	*	*	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic liver disease and cirrhosis (K73-K74)										
All races 3/, both sexes										
2002	5.2	*	*	*	*	*	*	0.2	0.5	1.5
2001	5.2	*	*	*	*	*	*	0.2	0.4	1.6
2000	5.1	*	*	*	*	*	0.1	0.2	0.5	1.5
1999	5.1	*	*	*	*	*	*	0.2	0.6	1.5
All races 3/, male										
2002	6.1	*	*	*	*	*	*	*	0.6	1.9
2001	6.1	*	*	*	*	*	*	0.2	0.6	2.0
2000	6.0	*	*	*	*	*	*	*	0.6	2.0
1999	6.0	*	*	*	*	*	*	*	0.7	2.0
All races 3/, female										
2002	4.5	*	*	*	*	*	*	0.2	0.4	1.0
2001	4.4	*	*	*	*	*	*	*	0.3	1.3
2000	4.3	*	*	*	*	*	*	*	0.4	1.0
1999	4.3	*	*	*	*	*	*	*	0.4	1.0
White, both sexes										
2002	5.7	*	*	*	*	*	*	0.2	0.4	1.5
2001	5.6	*	*	*	*	*	*	*	0.5	1.7
2000	5.5	*	*	*	*	*	*	*	0.5	1.5
1999	5.5	*	*	*	*	*	*	0.1	0.6	1.5
White, male										
2002	6.5	*	*	*	*	*	*	*	0.6	2.0
2001	6.4	*	*	*	*	*	*	*	0.6	2.0
2000	6.4	*	*	*	*	*	*	*	0.5	2.1
1999	6.4	*	*	*	*	*	*	*	0.7	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic liver disease and cirrhosis (K73-K74)										
White, female										
2002	4.9	*	*	*	*	*	*	*	0.3	1.0
2001	4.8	*	*	*	*	*	*	*	0.4	1.4
2000	4.7	*	*	*	*	*	*	*	0.4	0.9
1999	4.6	*	*	*	*	*	*	*	0.4	1.0
Black, both sexes										
2002	3.7	*	*	*	*	*	*	*	*	1.4
2001	3.8	*	*	*	*	*	*	*	*	1.5
2000	3.7	*	*	*	*	*	*	*	*	1.7
1999	3.8	*	*	*	*	*	*	*	*	1.5
Black, male										
2002	4.6	*	*	*	*	*	*	*	*	1.7
2001	5.1	*	*	*	*	*	*	*	*	2.0
2000	4.7	*	*	*	*	*	*	*	*	2.2
1999	4.8	*	*	*	*	*	*	*	*	1.9
Black, female										
2002	2.9	*	*	*	*	*	*	*	*	*
2001	2.7	*	*	*	*	*	*	*	*	*
2000	2.8	*	*	*	*	*	*	*	*	*
1999	2.9	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All races 3/, both sexes										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.0	*	*	*	*	*	*	*	*	0.1
2000	1.0	*	*	*	*	*	*	*	*	0.1
1999	1.0	*	*	*	*	*	*	*	*	0.1
All races 3/, male										
2002	0.9	*	*	*	*	*	*	*	*	*
2001	1.0	*	*	*	*	*	*	*	*	*
2000	0.9	*	*	*	*	*	*	*	*	*
1999	0.9	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	1.1	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.1	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	1.1	*	*	*	*	*	*	*	*	*
2001	1.1	*	*	*	*	*	*	*	*	*
2000	1.1	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*
White, male										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.0	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cholelithiasis and other disorders of gallbladder (K80-K82)										
White, female										
2002	1.2	*	*	*	*	*	*	*	*	*
2001	1.2	*	*	*	*	*	*	*	*	*
2000	1.2	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.7	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.8	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.6	*	*	*	*	*	*	*	*	*
2001	0.7	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.9	*	*	*	*	*	*	*	*	*
2000	1.0	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes										
2002	14.2	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3
2001	13.9	3.3	*	*	*	0.1	0.4	0.4	0.7	1.3
2000	13.2	4.3	*	*	*	0.2	0.3	0.4	0.8	1.2
1999	12.7	4.4	*	*	*	0.1	0.3	0.5	0.8	1.2
All races 3/, male										
2002	13.9	5.3	*	*	*	*	0.3	0.5	0.8	1.4
2001	13.5	3.7	*	*	*	*	0.3	0.5	0.6	1.4
2000	12.9	5.3	*	*	*	*	0.3	0.4	0.8	1.3
1999	12.4	4.6	*	*	*	*	0.2	0.4	0.8	1.3
All races 3/, female										
2002	14.5	3.2	*	*	*	*	0.3	0.7	0.7	1.1
2001	14.2	2.9	*	*	*	*	0.4	0.4	0.7	1.3
2000	13.6	3.2	*	*	*	*	0.3	0.4	0.8	1.1
1999	13.0	4.1	*	*	*	*	0.3	0.6	0.7	1.0
White, both sexes										
2002	13.9	3.6	*	*	*	*	0.2	0.4	0.6	0.8
2001	13.5	2.4	*	*	*	*	0.3	0.3	0.5	0.9
2000	12.9	2.9	*	*	*	0.1	0.3	0.3	0.5	0.8
1999	12.3	3.4	*	*	*	*	0.2	0.3	0.5	0.8
White, male										
2002	13.7	4.4	*	*	*	*	*	0.3	0.6	0.9
2001	13.3	2.8	*	*	*	*	*	0.3	0.4	1.1
2000	12.7	3.8	*	*	*	*	0.3	0.4	0.5	0.9
1999	12.1	4.1	*	*	*	*	*	0.3	0.5	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
White, female										
2002	14.1	2.7	*	*	*	*	*	0.5	0.5	0.8
2001	13.7	2.1	*	*	*	*	0.4	0.3	0.6	0.7
2000	13.0	2.0	*	*	*	*	0.3	0.3	0.5	0.7
1999	12.4	2.8	*	*	*	*	*	0.4	0.5	0.7
Black, both sexes										
2002	19.8	8.2	*	*	*	*	0.9	1.9	2.1	4.0
2001	19.6	8.1	*	*	*	*	0.7	1.3	2.2	4.0
2000	18.9	11.5	*	*	*	*	*	1.1	2.6	4.4
1999	18.6	9.6	*	*	*	*	0.8	1.5	2.3	4.0
Black, male										
2002	19.1	10.2	*	*	*	*	*	1.6	2.2	4.4
2001	18.0	8.7	*	*	*	*	*	1.7	2.6	3.4
2000	17.7	13.4	*	*	*	*	*	*	2.8	5.1
1999	17.5	8.1	*	*	*	*	*	*	2.3	4.8
Black, female										
2002	20.5	6.1	*	*	*	*	*	2.1	2.1	3.6
2001	21.0	7.6	*	*	*	*	*	*	1.8	4.5
2000	20.0	9.6	*	*	*	*	*	*	2.3	3.8
1999	19.5	11.1	*	*	*	*	*	1.6	2.3	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)										
All races 3/, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)										
White, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
All races 3/, both sexes										
2002	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2
1999	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1
All races 3/, male										
2002	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3
1999	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3
All races 3/, female										
2002	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0
1999	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9
White, both sexes										
2002	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000	12.6	2.7	*	*	*	*	0.2	0.3	0.5	0.7
1999	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7
White, male										
2002	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000	12.4	3.6	*	*	*	*	0.3	0.3	0.5	0.8
1999	11.8	3.9	*	*	*	*	*	*	0.5	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
White, female										
2002	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000	12.8	1.7	*	*	*	*	*	*	0.5	0.7
1999	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6
Black, both sexes										
2002	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000	18.5	11.1	*	*	*	*	*	1.0	2.5	4.1
1999	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8
Black, male										
2002	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000	17.3	13.4	*	*	*	*	*	*	2.7	4.9
1999	17.1	8.1	*	*	*	*	*	*	2.2	4.5
Black, female										
2002	20.2	*	*	*	*	*	*	1.9	2.0	3.5
2001	20.7	7.6	*	*	*	*	*	*	1.7	4.4
2000	19.6	8.6	*	*	*	*	*	*	2.3	3.4
1999	19.1	11.1	*	*	*	*	*	1.6	2.2	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of kidney (N25,N27)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of kidney (N25,N27)										
White, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Infections of kidney (N10-N12,N13.6,N15.1)										
All races 3/, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
White, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Infections of kidney (N10-N12,N13.6,N15.1)										
White, female										
2002	0.4	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hyperplasia of prostate (N40)										
All races 3/, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
White, both sexes										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
White, male										
2002	0.3	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hyperplasia of prostate (N40)										
White, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
Black, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Inflammatory diseases of female pelvic organs (N70-N76)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Inflammatory diseases of female pelvic organs (N70-N76)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy, childbirth and the puerperium (O00-O99)										
All races 3/, both sexes										
2002	0.1	...	...	...	*	0.1	0.3	0.4	0.4	0.4
2001	0.1	...	...	...	*	0.2	0.4	0.5	0.5	0.3
2000	0.1	...	...	...	*	*	0.4	0.4	0.5	0.4
1999	0.1	...	...	...	*	0.2	0.3	0.5	0.4	0.4
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.3	...	...	...	*	0.3	0.6	0.9	0.9	0.7
2001	0.3	...	...	...	*	0.4	0.8	1.0	0.9	0.7
2000	0.3	...	...	...	*	*	0.8	0.9	0.9	0.8
1999	0.3	...	...	...	*	0.3	0.7	0.9	0.9	0.8
White, both sexes										
2002	0.1	...	...	...	*	*	0.2	0.3	0.3	0.2
2001	0.1	...	...	...	*	0.1	0.3	0.3	0.4	0.2
2000	0.1	...	...	...	*	*	0.3	0.3	0.3	0.3
1999	0.1	...	...	...	*	*	0.2	0.3	0.3	0.3
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy, childbirth and the puerperium (O00-O99)										
White, female										
2002	0.2	...	...	...	*	*	0.4	0.6	0.7	0.5
2001	0.2	...	...	...	*	0.3	0.6	0.7	0.8	0.4
2000	0.2	...	...	...	*	*	0.6	0.7	0.7	0.6
1999	0.2	...	...	...	*	*	0.4	0.6	0.6	0.6
Black, both sexes										
2002	0.4	...	...	...	*	*	1.0	1.4	1.0	1.2
2001	0.4	...	...	...	*	*	1.1	1.4	1.1	1.0
2000	0.4	...	...	...	*	*	1.1	1.1	1.2	0.9
1999	0.4	...	...	...	*	*	1.1	1.4	1.2	0.9
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	0.8	...	...	...	*	*	2.0	2.8	1.9	2.3
2001	0.8	...	...	...	*	*	2.1	2.6	2.1	1.9
2000	0.7	...	...	...	*	*	2.2	2.2	2.3	1.7
1999	0.9	...	...	...	*	*	2.1	2.7	2.3	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (O00-O07)										
All races 3/, both sexes										
2002	0.0	...	...	...	*	*	*	*	*	*
2001	0.0	...	...	...	*	*	*	*	*	*
2000	0.0	...	...	...	*	*	*	*	*	*
1999	0.0	...	...	...	*	*	*	*	*	*
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.0	...	...	...	*	*	*	*	*	*
2001	0.0	...	...	...	*	*	*	*	*	*
2000	0.0	...	...	...	*	*	*	*	*	*
1999	0.0	...	...	...	*	*	*	*	*	*
White, both sexes										
2002	*	...	...	...	*	*	*	*	*	*
2001	0.0	...	...	...	*	*	*	*	*	*
2000	*	...	...	...	*	*	*	*	*	*
1999	*	...	...	...	*	*	*	*	*	*
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (O00-O07)										
White, female										
2002	*	...	...	...	*	*	*	*	*	*
2001	0.0	...	...	...	*	*	*	*	*	*
2000	*	...	...	...	*	*	*	*	*	*
1999	*	...	...	...	*	*	*	*	*	*
Black, both sexes										
2002	*	...	...	...	*	*	*	*	*	*
2001	*	...	...	...	*	*	*	*	*	*
2000	*	...	...	...	*	*	*	*	*	*
1999	*	...	...	...	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	*	...	...	...	*	*	*	*	*	*
2001	*	...	...	...	*	*	*	*	*	*
2000	*	...	...	...	*	*	*	*	*	*
1999	*	...	...	...	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
All races 3/, both sexes										
2002	0.1	...	...	...	*	0.1	0.3	0.4	0.4	0.3
2001	0.1	...	...	...	*	0.2	0.3	0.5	0.4	0.3
2000	0.1	...	...	...	*	*	0.4	0.4	0.4	0.3
1999	0.1	...	...	...	*	0.1	0.3	0.4	0.4	0.4
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.2	...	...	...	*	0.3	0.6	0.8	0.8	0.7
2001	0.3	...	...	...	*	0.3	0.7	0.9	0.9	0.6
2000	0.3	...	...	...	*	*	0.8	0.8	0.9	0.7
1999	0.3	...	...	...	*	0.3	0.6	0.9	0.8	0.7
White, both sexes										
2002	0.1	...	...	...	*	*	0.2	0.3	0.3	0.2
2001	0.1	...	...	...	*	*	0.2	0.3	0.4	0.2
2000	0.1	...	...	...	*	*	0.3	0.3	0.3	0.3
1999	0.1	...	...	...	*	*	0.2	0.3	0.3	0.3
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
White, female										
2002	0.2	...	...	...	*	*	0.4	0.5	0.6	0.5
2001	0.2	...	...	...	*	*	0.5	0.6	0.7	0.3
2000	0.2	...	...	...	*	*	0.6	0.6	0.7	0.5
1999	0.2	...	...	...	*	*	0.4	0.6	0.6	0.6
Black, both sexes										
2002	0.4	...	...	...	*	*	0.9	1.4	0.9	1.0
2001	0.4	...	...	...	*	*	1.0	1.3	1.0	0.9
2000	0.3	...	...	...	*	*	1.0	1.0	1.1	0.7
1999	0.4	...	...	...	*	*	1.0	1.3	1.0	0.8
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	0.7	...	...	...	*	*	1.8	2.6	1.8	2.0
2001	0.7	...	...	...	*	*	1.9	2.5	1.8	1.7
2000	0.6	...	...	...	*	*	1.9	1.8	2.1	1.4
1999	0.8	...	...	...	*	*	1.9	2.4	2.0	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Certain conditions originating in the perinatal period (P00-P96)										
All races 3/, both sexes										
2002	4.9	349.7	0.4	*	*	*	*	*	*	*
2001	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000	5.0	365.7	0.5	0.1	*	*	*	*	*	*
1999	5.1	371.0	0.6	0.1	*	*	*	*	*	*
All races 3/, male										
2002	5.7	385.7	0.5	*	*	*	*	*	*	*
2001	5.6	376.6	0.6	*	*	*	*	*	*	*
2000	5.8	403.9	0.7	*	*	*	*	*	*	*
1999	5.9	408.6	0.7	*	*	*	*	*	*	*
All races 3/, female										
2002	4.2	312.0	0.3	*	*	*	*	*	*	*
2001	4.2	302.6	0.4	*	*	*	*	*	*	*
2000	4.3	325.6	0.4	*	*	*	*	*	*	*
1999	4.4	331.7	0.5	*	*	*	*	*	*	*
White, both sexes										
2002	3.7	277.2	0.3	*	*	*	*	*	*	*
2001	3.7	268.5	0.4	*	*	*	*	*	*	*
2000	3.7	283.7	0.4	*	*	*	*	*	*	*
1999	3.8	285.9	0.5	*	*	*	*	*	*	*
White, male										
2002	4.3	309.4	*	*	*	*	*	*	*	*
2001	4.2	297.1	0.5	*	*	*	*	*	*	*
2000	4.3	312.4	0.5	*	*	*	*	*	*	*
1999	4.3	315.0	0.5	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Certain conditions originating in the perinatal period (P00-P96)										
White, female										
2002	3.2	243.5	*	*	*	*	*	*	*	*
2001	3.1	238.6	*	*	*	*	*	*	*	*
2000	3.2	253.5	*	*	*	*	*	*	*	*
1999	3.2	255.3	0.4	*	*	*	*	*	*	*
Black, both sexes										
2002	13.1	724.8	1.0	*	*	*	*	*	*	*
2001	13.0	738.1	0.9	*	*	*	*	*	*	*
2000	13.7	809.5	1.1	*	*	*	*	*	*	*
1999	14.2	844.6	1.2	*	*	*	*	*	*	*
Black, male										
2002	15.2	783.2	*	*	*	*	*	*	*	*
2001	15.5	815.9	*	*	*	*	*	*	*	*
2000	16.3	901.3	*	*	*	*	*	*	*	*
1999	16.9	937.7	*	*	*	*	*	*	*	*
Black, female										
2002	11.2	663.8	*	*	*	*	*	*	*	*
2001	10.8	656.2	*	*	*	*	*	*	*	*
2000	11.3	714.6	*	*	*	*	*	*	*	*
1999	11.8	748.5	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All races 3/, both sexes										
2002	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
1999	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2
All races 3/, male										
2002	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
1999	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3
All races 3/, female										
2002	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
1999	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1
White, both sexes										
2002	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
1999	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2
White, male										
2002	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
1999	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
White, female										
2002	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
1999	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1
Black, both sexes										
2002	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
1999	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4
Black, male										
2002	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
1999	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*
Black, female										
2002	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5
1999	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All races 3/, both sexes										
2002	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
1999	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1
All races 3/, male										
2002	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
1999	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3
All races 3/, female										
2002	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
1999	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0
White, both sexes										
2002	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
1999	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7
White, male										
2002	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
1999	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
White, female										
2002	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
1999	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5
Black, both sexes										
2002	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
1999	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8
Black, male										
2002	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
1999	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4
Black, female										
2002	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3
1999	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9, N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
All races 3/, both sexes										
2002	67.5	28.4	4.2	1.9	2.3	3.5	5.6	6.6	9.0	14.1
2001	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1
2000	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9
1999	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0
All races 3/, male										
2002	55.8	33.4	4.6	2.2	2.5	4.0	6.3	7.4	10.6	16.4
2001	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8
2000	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5
1999	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8
All races 3/, female										
2002	78.8	23.0	3.9	1.7	2.1	3.0	4.9	5.8	7.3	11.9
2001	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5
2000	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3
1999	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3
White, both sexes										
2002	71.8	25.2	3.9	1.9	2.2	3.4	5.0	6.0	8.1	12.8
2001	69.4	23.7	3.9	2.0	2.2	3.6	4.9	5.8	7.9	12.6
2000	67.0	24.9	3.5	1.7	2.3	3.5	5.1	5.7	8.0	12.3
1999	65.0	25.2	4.0	2.0	2.3	3.8	4.8	5.9	8.1	12.1
White, male										
2002	58.2	30.8	4.1	2.2	2.3	4.1	5.9	6.8	9.9	15.5
2001	55.8	25.9	4.4	1.9	2.4	4.4	5.8	6.6	8.9	15.5
2000	54.1	26.8	3.9	1.9	2.6	4.1	6.2	7.0	9.6	15.0
1999	52.7	29.2	4.3	2.0	2.2	4.8	5.5	7.0	9.8	15.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9, N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
White, female										
2002	85.2	19.3	3.6	1.6	2.1	2.8	4.1	5.0	6.1	9.9
2001	82.7	21.4	3.4	2.0	2.0	2.8	3.9	4.8	6.8	9.7
2000	79.6	22.9	3.2	1.6	2.0	2.8	3.9	4.4	6.4	9.5
1999	77.0	20.9	3.7	1.9	2.3	2.8	4.1	4.7	6.3	9.1
Black, both sexes										
2002	59.7	45.2	5.9	2.2	2.9	4.3	9.4	12.2	16.6	25.7
2001	59.1	51.3	5.4	2.0	2.9	5.3	7.7	11.5	16.6	26.6
2000	58.1	49.6	5.9	2.3	3.3	4.8	8.3	12.8	17.1	26.9
1999	56.2	56.7	5.3	2.9	3.1	4.7	8.9	10.4	15.4	28.7
Black, male										
2002	54.8	48.8	6.4	2.3	3.4	4.2	9.4	12.7	17.9	26.4
2001	54.6	58.4	5.1	1.5	3.3	5.6	7.7	12.3	17.5	29.7
2000	54.3	59.8	6.1	2.0	3.1	5.1	9.1	12.9	18.3	29.9
1999	52.8	64.7	5.4	3.2	3.8	4.8	9.7	10.5	16.2	31.5
Black, female										
2002	64.1	41.5	5.4	2.2	2.5	4.3	9.5	11.8	15.5	25.0
2001	63.1	43.8	5.6	2.5	2.5	5.1	7.6	10.8	15.7	24.0
2000	61.6	39.0	5.7	2.6	3.5	4.6	7.6	12.6	16.1	24.2
1999	59.3	48.3	5.1	2.6	2.4	4.5	8.1	10.3	14.6	26.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2002	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
1999	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2
All races 3/, male										
2002	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
1999	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4
All races 3/, female										
2002	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
1999	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0
White, both sexes										
2002	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
1999	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4
White, male										
2002	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
1999	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
White, female										
2002	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
1999	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0
Black, both sexes										
2002	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
1999	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6
Black, male										
2002	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
1999	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4
Black, female										
2002	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
1999	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2002	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
1999	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9
All races 3/, male										
2002	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
1999	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0
All races 3/, female										
2002	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
1999	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8
White, both sexes										
2002	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
1999	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8
White, male										
2002	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
1999	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
White, female										
2002	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
1999	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8
Black, both sexes										
2002	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
1999	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9
Black, male										
2002	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
1999	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7
Black, female										
2002	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4
1999	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2002	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6
1999	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1
All races 3/, male										
2002	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1
1999	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2
All races 3/, female										
2002	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1
1999	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0
White, both sexes										
2002	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000	15.7	3.9	4.0	3.7	5.0	28.4	29.1	19.0	15.4	15.8
1999	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9
White, male										
2002	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000	21.6	4.2	4.5	4.0	5.9	36.8	42.6	27.9	22.4	22.1
1999	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
White, female										
2002	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000	10.0	3.5	3.6	3.4	4.0	19.5	14.6	9.6	8.2	9.3
1999	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9
Black, both sexes										
2002	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000	15.0	6.5	5.4	4.3	4.6	16.5	25.8	22.8	18.0	16.4
1999	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2
Black, male										
2002	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000	22.5	6.7	5.8	4.5	6.2	22.8	38.8	36.7	28.5	26.0
1999	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0
Black, female										
2002	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000	8.2	*	4.9	4.0	2.9	10.1	13.5	10.2	8.6	7.9
1999	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)										
All races 3/, both sexes										
2002	0.4	*	*	0.1	0.2	0.3	0.5	0.3	0.4	0.5
2001	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6
2000	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7
1999	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9
All races 3/, male										
2002	0.6	*	*	*	0.3	0.5	0.8	0.5	0.6	0.8
2001	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1
2000	0.8	*	*	0.2	0.4	1.0	1.0	0.8	0.9	1.1
1999	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	0.2	*	*	*	0.2
2000	0.3	*	*	*	*	0.3	0.2	*	0.3	0.2
1999	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4
White, both sexes										
2002	0.4	*	*	*	0.2	0.3	0.5	0.3	0.4	0.5
2001	0.5	*	0.2	*	0.2	0.5	0.6	0.5	0.4	0.5
2000	0.5	*	0.2	0.2	0.3	0.7	0.6	0.5	0.6	0.6
1999	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9
White, male										
2002	0.6	*	*	*	0.3	0.5	0.9	0.5	0.6	0.8
2001	0.7	*	*	*	0.3	0.7	0.9	0.8	0.7	0.9
2000	0.8	*	*	*	0.4	1.0	1.0	0.9	0.9	1.1
1999	1.1	*	*	0.4	0.5	1.2	1.7	1.3	1.1	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	0.3	*	*	0.3	0.2
1999	0.4	*	*	*	*	0.6	0.4	0.3	0.5	0.4
Black, both sexes										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	1.0
2000	0.6	*	*	*	*	*	*	*	0.8	0.9
1999	0.7	*	*	*	*	*	*	*	0.9	1.0
Black, male										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.9	*	*	*	*	*	*	*	*	1.6
2000	0.9	*	*	*	*	*	*	*	*	1.5
1999	1.1	*	*	*	*	*	*	*	*	1.6
Black, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
All races 3/, both sexes										
2002	0.7	*	*	0.1	0.2	0.4	0.6	0.5	0.7	0.9
2001	0.7	*	0.2	0.1	0.1	0.4	0.6	0.8	0.9	0.9
2000	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9
1999	0.7	*	*	0.2	0.2	0.4	0.6	0.8	0.8	0.9
All races 3/, male										
2002	1.1	*	*	*	0.2	0.6	0.9	0.9	1.2	1.5
2001	1.2	*	*	*	0.2	0.7	1.0	1.2	1.5	1.4
2000	1.2	*	*	*	0.2	0.6	1.2	1.3	1.3	1.5
1999	1.1	*	*	0.2	0.3	0.6	0.9	1.4	1.3	1.4
All races 3/, female										
2002	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2001	0.3	*	*	*	*	*	0.3	0.4	0.3	0.4
2000	0.2	*	*	*	*	0.3	*	*	0.2	0.2
1999	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.4
White, both sexes										
2002	0.7	*	*	*	0.2	0.5	0.6	0.6	0.8	0.9
2001	0.8	*	0.2	0.1	0.2	0.5	0.7	0.9	1.0	1.0
2000	0.7	*	*	*	0.2	0.5	0.8	0.9	0.8	0.9
1999	0.7	*	*	0.2	0.2	0.4	0.7	1.0	0.9	1.0
White, male										
2002	1.2	*	*	*	*	0.6	1.1	1.0	1.4	1.6
2001	1.3	*	*	*	*	0.8	1.1	1.4	1.6	1.5
2000	1.3	*	*	*	*	0.6	1.4	1.5	1.4	1.6
1999	1.2	*	*	*	0.3	0.6	1.1	1.5	1.4	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
White, female										
2002	0.2	*	*	*	*	0.3	*	*	*	0.3
2001	0.3	*	*	*	*	*	0.4	0.5	0.3	0.4
2000	0.2	*	*	*	*	0.3	*	*	*	*
1999	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4
Black, both sexes										
2002	0.4	*	*	*	*	*	*	*	*	0.7
2001	0.5	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	0.7
Black, male										
2002	0.8	*	*	*	*	*	*	*	*	*
2001	0.8	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	*	*	*
Black, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nontransport accidents (W00-X59,Y86)										
All races 3/, both sexes										
2002	20.2	20.3	6.4	2.4	2.4	6.6	11.1	11.8	13.6	17.8
2001	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7
2000	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8
1999	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3
All races 3/, male										
2002	25.2	23.3	8.1	3.1	3.3	10.6	18.0	18.4	20.6	25.9
2001	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8
2000	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2
1999	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4
All races 3/, female										
2002	15.4	17.2	4.7	1.6	1.5	2.5	3.9	4.9	6.5	9.7
2001	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5
2000	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4
1999	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2
White, both sexes										
2002	21.4	15.9	5.9	1.8	2.0	6.9	12.3	12.7	14.6	18.3
2001	20.0	16.9	6.7	2.2	2.1	6.3	11.3	11.0	12.2	17.2
2000	18.9	15.1	6.9	2.2	2.2	6.4	10.2	10.5	12.2	16.0
1999	19.0	14.3	7.3	2.2	2.4	6.2	9.7	10.4	12.1	15.6
White, male										
2002	26.2	19.2	7.4	2.4	2.8	10.9	19.7	19.7	22.0	26.4
2001	24.5	18.9	8.4	2.7	2.8	10.0	18.4	17.3	18.6	25.5
2000	23.4	17.3	8.4	2.8	3.1	10.3	16.7	16.9	19.1	24.6
1999	23.5	16.5	8.8	2.7	3.4	10.0	15.8	16.9	19.2	23.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nontransport accidents (W00-X59,Y86)										
White, female										
2002	16.8	12.5	4.4	1.2	1.2	2.7	4.3	5.3	6.9	10.1
2001	15.6	14.8	4.8	1.7	1.3	2.3	3.8	4.4	5.6	8.6
2000	14.6	12.8	5.2	1.6	1.2	2.2	3.3	3.8	5.1	7.3
1999	14.6	11.8	5.8	1.6	1.3	2.2	3.2	3.7	4.8	7.2
Black, both sexes										
2002	17.8	43.6	9.2	4.7	4.4	6.0	7.4	9.8	12.1	18.9
2001	17.6	40.2	8.4	4.1	4.1	5.9	7.2	9.4	11.1	17.7
2000	17.6	38.0	10.6	5.2	4.0	6.4	7.7	9.6	12.5	18.0
1999	18.9	34.4	10.6	5.6	4.6	6.3	8.8	10.2	14.4	16.7
Black, male										
2002	24.8	45.9	12.1	6.1	6.5	10.1	12.4	16.2	18.8	29.1
2001	24.4	46.4	9.7	6.0	5.7	9.6	11.8	15.1	16.5	26.1
2000	24.9	40.9	12.6	6.9	5.6	10.3	12.6	15.9	18.9	27.1
1999	26.8	40.0	13.0	7.4	6.6	10.8	15.5	16.7	22.3	25.7
Black, female										
2002	11.4	41.2	6.2	3.3	2.3	1.7	2.4	4.0	6.2	9.8
2001	11.5	33.7	7.1	2.2	2.5	2.1	2.7	4.3	6.2	10.2
2000	11.0	35.1	8.5	3.5	2.3	2.3	3.0	3.8	6.7	9.9
1999	11.7	28.5	8.1	3.7	2.5	1.6	2.5	4.3	7.3	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
All races 3/, both sexes										
2002	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0
1999	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2
All races 3/, male										
2002	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8
1999	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0
All races 3/, female										
2002	5.3	*	*	*	*	*	*	*	0.3	0.6
2001	4.8	*	*	*	*	*	*	*	0.3	0.4
2000	4.3	*	*	*	*	*	0.2	*	*	0.3
1999	4.3	*	*	*	*	*	*	*	0.3	0.4
White, both sexes										
2002	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000	5.3	*	0.2	*	*	0.6	0.8	0.8	0.9	1.0
1999	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3
White, male										
2002	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000	5.7	*	*	*	*	1.0	1.4	1.4	1.5	1.7
1999	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
White, female										
2002	6.2	*	*	*	*	*	*	*	0.3	0.7
2001	5.5	*	*	*	*	*	*	*	0.3	0.4
2000	5.0	*	*	*	*	*	0.3	*	*	0.3
1999	4.9	*	*	*	*	*	*	*	0.3	0.5
Black, both sexes										
2002	2.2	*	*	*	*	*	*	*	*	1.1
2001	2.3	*	*	*	*	*	*	*	*	1.1
2000	2.1	*	*	*	*	*	*	*	*	1.0
1999	2.4	*	*	*	*	*	*	*	1.1	0.9
Black, male										
2002	3.0	*	*	*	*	*	*	*	*	1.8
2001	3.1	*	*	*	*	*	*	*	*	2.0
2000	2.8	*	*	*	*	*	*	*	*	1.7
1999	3.3	*	*	*	*	*	*	*	1.8	1.7
Black, female										
2002	1.5	*	*	*	*	*	*	*	*	*
2001	1.6	*	*	*	*	*	*	*	*	*
2000	1.4	*	*	*	*	*	*	*	*	*
1999	1.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental discharge of firearms (W32-W34)										
All races 3/, both sexes										
2002	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.2
2001	0.3	*	*	*	0.2	0.5	0.5	0.4	0.2	0.3
2000	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.3
1999	0.3	*	*	*	0.3	0.6	0.7	0.4	0.3	0.3
All races 3/, male										
2002	0.5	*	*	*	0.3	1.0	0.9	0.7	0.6	0.4
2001	0.5	*	*	*	0.3	1.0	0.9	0.7	0.4	0.5
2000	0.5	*	*	*	0.4	0.9	0.9	0.7	0.4	0.5
1999	0.5	*	*	*	0.5	1.1	1.2	0.6	0.6	0.5
All races 3/, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.3	*	*	*	0.2	0.4	0.5	0.3	0.3	0.2
2001	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.4
2000	0.3	*	*	*	0.3	0.5	0.4	0.3	0.3	0.3
1999	0.3	*	*	*	0.3	0.6	0.6	0.4	0.3	0.3
White, male										
2002	0.4	*	*	*	0.3	0.8	0.8	0.6	0.5	0.4
2001	0.5	*	*	*	0.4	0.9	0.9	0.6	0.5	0.6
2000	0.5	*	*	*	0.5	0.9	0.8	0.6	0.4	0.5
1999	0.5	*	*	*	0.5	1.0	1.0	0.6	0.6	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental discharge of firearms (W32-W34)										
White, female										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.4	*	*	*	*	1.1	0.8	0.9	*	*
2001	0.3	*	*	*	*	0.8	0.8	*	*	*
2000	0.4	*	*	*	*	0.9	0.9	*	*	*
1999	0.4	*	*	*	*	1.2	1.1	*	*	*
Black, male										
2002	0.8	*	*	*	*	2.2	1.5	1.8	*	*
2001	0.6	*	*	*	*	1.6	1.5	*	*	*
2000	0.7	*	*	*	*	1.6	1.5	*	*	*
1999	0.8	*	*	*	*	2.1	2.1	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2002	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
1999	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1
All races 3/, male										
2002	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
1999	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9
All races 3/, female										
2002	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
1999	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3
White, both sexes										
2002	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1
White, male										
2002	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
1999	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
White, female										
2002	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4
Black, both sexes										
2002	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
1999	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1
Black, male										
2002	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
1999	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2
Black, female										
2002	0.5	*	*	*	*	*	*	*	*	*
2001	0.5	*	1.6	*	1.3	*	*	*	*	*
2000	0.6	*	1.7	*	*	*	*	*	*	*
1999	0.5	*	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2002	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
1999	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9
All races 3/, male										
2002	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
1999	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2
All races 3/, female										
2002	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
1999	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5
White, both sexes										
2002	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
1999	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8
White, male										
2002	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
1999	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
White, female										
2002	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5
Black, both sexes										
2002	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
1999	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9
Black, male										
2002	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
1999	3.0	*	5.0	2.4	*	*	*	*	*	2.8
Black, female										
2002	1.6	*	2.8	1.7	*	*	*	*	*	*
2001	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000	2.0	*	4.1	1.6	*	*	*	*	*	*
1999	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2002	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
1999	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1
All races 3/, male										
2002	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
1999	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3
All races 3/, female										
2002	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
1999	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9
White, both sexes										
2002	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
1999	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3
White, male										
2002	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
1999	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, female										
2002	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
1999	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8
Black, both sexes										
2002	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
1999	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3
Black, male										
2002	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
1999	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9
Black, female										
2002	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1
1999	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
All races 3/, both sexes										
2002	6.0	16.8	1.6	0.6	0.8	1.3	1.8	2.0	2.1	2.6
2001	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9
2000	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0
1999	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7
All races 3/, male										
2002	6.9	18.8	2.0	0.7	1.1	2.1	3.1	3.2	3.6	4.3
2001	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8
2000	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1
1999	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5
All races 3/, female										
2002	5.0	14.6	1.2	0.4	0.4	0.5	0.5	0.7	0.6	1.0
2001	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1
2000	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9
White, both sexes										
2002	6.3	12.9	1.4	0.5	0.8	1.4	2.0	2.2	2.3	2.7
2001	6.6	13.4	1.7	0.6	0.7	1.4	2.2	1.9	2.2	3.0
2000	6.6	12.2	1.6	0.6	0.8	1.4	2.1	2.3	2.4	3.0
1999	6.9	11.2	1.8	0.7	0.9	1.6	2.2	2.2	2.3	2.9
White, male										
2002	7.3	15.3	1.8	0.7	1.0	2.4	3.4	3.6	3.8	4.3
2001	7.5	15.2	2.0	0.7	1.0	2.1	3.7	3.2	3.7	5.0
2000	7.4	14.0	2.0	0.7	1.2	2.2	3.7	3.8	4.1	5.2
1999	7.8	12.9	2.1	1.0	1.3	2.5	3.7	3.9	3.8	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
White, female										
2002	5.4	10.5	1.1	0.4	0.5	0.5	0.5	0.7	0.6	1.0
2001	5.8	11.5	1.4	0.5	0.3	0.6	0.6	0.6	0.6	1.0
2000	5.7	10.3	1.2	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999	6.1	9.5	1.4	0.4	0.5	0.6	0.5	0.6	0.7	0.9
Black, both sexes										
2002	5.2	37.5	2.4	0.7	0.9	1.0	1.2	1.5	1.9	3.0
2001	5.4	34.2	2.5	*	0.7	0.8	1.3	1.8	1.9	2.6
2000	5.4	30.6	2.3	0.7	0.7	1.3	1.8	2.0	2.4	3.1
1999	6.1	27.9	2.5	0.7	0.7	1.0	1.6	2.2	2.6	2.2
Black, male										
2002	6.6	38.9	3.3	*	1.5	1.5	1.9	2.3	3.4	5.2
2001	6.8	40.4	2.8	*	*	*	2.0	2.8	3.6	3.7
2000	7.3	32.0	2.5	*	1.2	1.8	3.2	3.5	3.8	5.5
1999	8.0	33.2	3.4	*	1.2	1.8	3.0	3.5	4.4	3.5
Black, female										
2002	4.0	36.0	*	*	*	*	*	*	*	*
2001	4.1	27.7	2.2	*	*	*	*	*	*	1.5
2000	3.8	29.1	2.0	*	*	*	*	*	*	*
1999	4.4	22.5	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2002	11.0	...	...	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 4/	10.8	...	...	*	1.3	7.9	12.0	12.6	13.0	14.3
2000	10.4	...	...	*	1.5	8.0	12.5	12.1	12.0	13.7
1999	10.5	...	...	*	1.2	8.0	12.3	12.3	13.1	14.4
All races 3/, male										
2002	17.9	...	...	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 4/	17.6	...	...	*	1.9	12.9	20.5	21.0	21.0	22.7
2000	17.1	...	...	*	2.3	13.0	21.4	20.0	19.2	21.7
1999	17.1	...	...	*	1.9	13.0	20.8	20.3	21.1	22.7
All races 3/, female										
2002	4.3	...	...	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 4/	4.1	...	...	*	0.6	2.7	3.1	4.0	4.8	5.9
2000	4.0	...	...	*	0.6	2.7	3.2	4.0	4.6	5.8
1999	4.0	...	...	*	0.5	2.7	3.4	4.1	5.1	6.1
White, both sexes										
2002	12.2	...	...	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 4/	11.9	...	...	*	1.3	8.6	12.6	13.5	14.2	15.8
2000	11.5	...	...	*	1.5	8.5	13.1	12.7	13.0	15.1
1999	11.5	...	...	*	1.3	8.5	12.7	12.8	14.2	15.7
White, male										
2002	19.9	...	...	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 4/	19.5	...	...	*	2.0	14.0	21.3	22.2	22.6	24.7
2000	18.8	...	...	*	2.4	13.7	22.3	20.8	20.7	23.6
1999	18.7	...	...	*	2.0	13.7	21.5	21.0	22.5	24.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, female										
2002	4.8	...	...	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 4/	4.6	...	...	*	0.6	2.9	3.3	4.3	5.4	6.7
2000	4.4	...	...	*	0.7	2.9	3.3	4.1	5.1	6.5
1999	4.5	...	...	*	0.5	2.9	3.5	4.3	5.5	6.9
Black, both sexes										
2002	5.1	...	...	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 4/	5.3	...	...	*	1.1	4.3	10.1	9.6	8.6	7.8
2000	5.4	...	...	*	1.2	5.5	11.0	9.9	7.6	7.3
1999	5.4	...	...	*	0.9	5.8	10.7	9.6	8.7	8.4
Black, male										
2002	9.1	...	...	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 4/	9.2	...	...	*	1.7	7.3	19.4	17.7	15.4	13.8
2000	9.4	...	...	*	2.1	9.5	19.5	17.5	13.6	13.0
1999	9.6	...	...	*	1.3	9.9	19.5	17.9	15.7	15.6
Black, female										
2002	1.5	...	...	*	*	*	2.3	2.0	2.0	2.5
2001 4/	1.7	...	...	*	*	1.3	*	2.2	2.5	2.5
2000	1.7	...	...	*	*	1.4	3.0	3.0	2.2	2.2
1999	1.6	...	...	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2002	5.9	...	...	*	0.4	3.6	6.7	6.4	5.6	6.6
2001	5.9	...	...	*	0.4	4.1	6.6	6.7	6.3	6.4
2000	5.9	...	...	*	0.5	4.4	7.2	6.3	6.0	6.7
1999	5.9	...	...	*	0.5	4.9	7.2	6.6	6.5	7.1
All races 3/, male										
2002	10.6	...	...	*	0.6	6.4	11.9	11.1	9.6	11.3
2001	10.6	...	...	*	0.6	7.1	11.8	11.7	10.8	10.8
2000	10.5	...	...	*	0.9	7.6	12.9	11.1	10.1	11.1
1999	10.6	...	...	*	0.8	8.4	12.8	11.5	10.9	11.8
All races 3/, female										
2002	1.4	...	...	*	*	0.7	1.2	1.6	1.7	2.0
2001	1.5	...	...	*	0.2	1.0	1.0	1.5	1.7	2.1
2000	1.5	...	...	*	0.2	1.1	1.3	1.5	1.9	2.3
1999	1.5	...	...	*	0.2	1.1	1.4	1.6	1.9	2.3
White, both sexes										
2002	6.7	...	...	*	0.4	3.9	7.0	6.8	6.2	7.4
2001	6.7	...	...	*	0.4	4.5	6.7	7.2	7.0	7.2
2000	6.6	...	...	*	0.6	4.6	7.6	6.6	6.6	7.5
1999	6.6	...	...	*	0.6	5.2	7.4	7.0	7.1	7.8
White, male										
2002	11.9	...	...	*	0.7	6.9	12.3	11.3	10.4	12.4
2001	11.8	...	...	*	0.7	7.6	11.9	12.3	11.7	11.9
2000	11.6	...	...	*	1.0	7.9	13.5	11.4	11.0	12.2
1999	11.7	...	...	*	0.9	8.9	13.0	11.9	11.8	12.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, female										
2002	1.6	...	...	*	*	0.8	1.3	2.0	1.9	2.3
2001	1.7	...	...	*	*	1.1	1.1	1.7	2.0	2.4
2000	1.7	...	...	*	*	1.1	1.3	1.6	2.0	2.7
1999	1.7	...	...	*	0.3	1.2	1.4	1.8	2.2	2.7
Black, both sexes										
2002	2.8	...	...	*	*	2.5	5.8	6.1	4.0	4.0
2001	2.9	...	...	*	*	2.8	7.1	5.8	4.1	3.7
2000	3.0	...	...	*	*	3.8	6.8	6.0	4.0	3.4
1999	3.1	...	...	*	*	3.9	7.6	5.7	4.2	4.2
Black, male										
2002	5.4	...	...	*	*	4.3	11.0	12.4	7.7	7.4
2001	5.5	...	...	*	*	5.0	13.9	10.9	8.0	7.1
2000	5.5	...	...	*	*	6.9	12.6	11.3	7.3	6.4
1999	5.9	...	...	*	*	7.0	14.2	11.2	8.1	8.1
Black, female										
2002	0.5	...	...	*	*	*	*	*	*	*
2001	0.6	...	...	*	*	*	*	*	*	*
2000	0.7	...	...	*	*	*	*	*	*	*
1999	0.6	...	...	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
All races 3/, both sexes										
2002	5.0	...	...	*	0.8	3.8	5.7	6.4	6.9	7.7
2001 4/	4.8	...	...	*	0.9	3.8	5.4	5.9	6.7	7.9
2000	4.5	...	...	*	0.9	3.6	5.3	5.8	6.0	7.0
1999	4.5	...	...	*	0.7	3.2	5.1	5.7	6.7	7.3
All races 3/, male										
2002	7.3	...	...	*	1.2	5.8	8.9	9.7	10.6	11.4
2001 4/	7.1	...	...	*	1.3	5.8	8.7	9.3	10.2	11.9
2000	6.6	...	...	*	1.4	5.4	8.5	8.9	9.1	10.6
1999	6.6	...	...	*	1.1	4.6	8.0	8.8	10.2	10.9
All races 3/, female										
2002	2.9	...	...	*	0.4	1.6	2.3	2.9	3.1	4.0
2001 4/	2.6	...	...	*	0.4	1.7	2.0	2.5	3.1	3.9
2000	2.5	...	...	*	0.4	1.7	1.9	2.6	2.7	3.4
1999	2.5	...	...	*	0.3	1.6	2.0	2.5	3.2	3.7
White, both sexes										
2002	5.5	...	...	*	0.8	4.2	6.2	6.8	7.4	8.5
2001 4/	5.3	...	...	*	0.9	4.2	5.9	6.3	7.2	8.6
2000	4.9	...	...	*	0.9	3.8	5.5	6.1	6.4	7.6
1999	4.9	...	...	*	0.7	3.3	5.4	5.9	7.1	7.9
White, male										
2002	8.0	...	...	*	1.2	6.5	9.7	10.2	11.2	12.3
2001 4/	7.7	...	...	*	1.3	6.4	9.4	9.8	10.8	12.8
2000	7.1	...	...	*	1.3	5.8	8.8	9.4	9.7	11.4
1999	7.0	...	...	*	1.1	4.9	8.5	9.1	10.7	11.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
White, female										
2002	3.2	...	...	*	0.5	1.8	2.5	3.1	3.4	4.5
2001 4/	2.9	...	...	*	0.4	1.8	2.2	2.6	3.4	4.3
2000	2.8	...	...	*	0.5	1.8	1.9	2.5	3.1	3.8
1999	2.8	...	...	*	0.3	1.7	2.1	2.5	3.3	4.2
Black, both sexes										
2002	2.3	...	...	*	0.7	1.6	3.2	4.2	4.2	4.1
2001 4/	2.3	...	...	*	0.8	1.5	3.0	3.8	4.5	4.1
2000	2.4	...	...	*	1.1	1.8	4.2	3.9	3.6	3.8
1999	2.3	...	...	*	0.7	1.9	3.1	3.9	4.6	4.2
Black, male										
2002	3.7	...	...	*	1.1	2.6	5.0	7.0	7.5	7.0
2001 4/	3.7	...	...	*	1.1	2.3	5.5	6.8	7.4	6.7
2000	3.9	...	...	*	1.8	2.6	6.8	6.2	6.4	6.6
1999	3.7	...	...	*	*	2.9	5.3	6.7	7.6	7.5
Black, female										
2002	1.1	...	...	*	*	*	1.5	1.6	1.4	1.6
2001 4/	1.1	...	...	*	*	*	*	*	1.8	1.8
2000	1.0	...	...	*	*	*	1.7	1.8	*	1.4
1999	1.0	...	...	*	*	*	*	*	1.9	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All races 3/, both sexes										
2002	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 4/	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
1999	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7
All races 3/, male										
2002	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 4/	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
1999	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0
All races 3/, female										
2002	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 4/	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
1999	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3
White, both sexes										
2002	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 4/	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
1999	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1
White, male										
2002	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 4/	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
1999	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
White, female										
2002	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 4/	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
1999	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1
Black, both sexes										
2002	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 4/	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
1999	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1
Black, male										
2002	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 4/	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
1999	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1
Black, female										
2002	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 4/	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1
1999	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2002	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
1999	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6
All races 3/, male										
2002	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
1999	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1
All races 3/, female										
2002	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
1999	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2
White, both sexes										
2002	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
1999	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0
White, male										
2002	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
1999	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
White, female										
2002	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
1999	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6
Black, both sexes										
2002	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
1999	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9
Black, male										
2002	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
1999	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3
Black, female										
2002	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5
1999	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)										
All races 3/, both sexes										
2002	2.0	7.3	2.4	0.4	0.3	1.6	2.9	2.6	2.5	2.7
2001 4/	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2
2000	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9
1999	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0
All races 3/, male										
2002	2.6	7.8	2.6	0.4	0.3	2.1	3.9	3.4	3.0	3.4
2001 4/	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3
2000	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5
1999	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9
All races 3/, female										
2002	1.4	6.8	2.2	0.5	0.3	1.1	1.8	1.8	2.0	2.1
2001 4/	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2
2000	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4
1999	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2
White, both sexes										
2002	1.5	5.3	1.7	0.3	0.2	1.1	2.2	1.9	1.7	2.0
2001 4/	2.6	6.0	1.6	0.4	0.3	1.4	3.0	3.9	4.3	4.6
2000	1.5	6.5	1.5	0.4	0.4	1.3	2.3	2.0	2.0	2.1
1999	1.6	6.6	1.5	0.5	0.4	1.4	2.4	2.1	2.2	2.1
White, male										
2002	1.9	6.2	1.8	*	*	1.5	2.9	2.5	2.0	2.5
2001 4/	3.7	7.0	1.9	0.3	*	1.8	3.9	5.3	5.9	6.8
2000	2.0	8.1	1.7	0.5	0.4	1.6	3.0	2.5	2.5	2.5
1999	2.1	7.7	1.5	0.4	0.3	1.7	3.0	2.7	2.6	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)										
White, female										
2002	1.1	4.5	1.6	0.4	0.3	0.7	1.5	1.2	1.4	1.5
2001 4/	1.6	4.9	1.4	0.4	0.3	0.9	2.0	2.5	2.6	2.3
2000	1.1	4.8	1.3	0.3	0.5	1.0	1.5	1.4	1.4	1.7
1999	1.2	5.4	1.5	0.5	0.4	1.1	1.8	1.5	1.7	1.5
Black, both sexes										
2002	5.3	16.5	5.8	1.0	0.6	3.7	6.4	7.3	7.0	7.4
2001 4/	6.3	18.6	6.1	*	0.6	4.0	7.1	9.0	8.4	9.8
2000	5.9	21.6	5.5	0.8	0.8	4.5	8.8	7.6	7.6	8.6
1999	5.7	19.2	5.4	1.3	0.6	4.5	7.0	7.7	8.1	9.2
Black, male										
2002	7.3	15.7	6.4	*	*	4.8	9.4	9.7	9.1	9.5
2001 4/	8.5	20.7	7.3	*	*	5.4	9.8	12.8	11.0	11.3
2000	8.0	21.1	5.5	*	*	5.5	12.6	10.7	10.5	10.9
1999	7.7	19.2	5.4	1.2	*	6.1	9.8	9.5	10.7	12.8
Black, female										
2002	3.4	17.3	5.1	*	*	2.6	3.5	5.0	5.0	5.6
2001 4/	4.3	16.4	4.8	*	*	2.5	4.5	5.5	6.1	8.4
2000	4.0	22.2	5.5	*	*	3.4	5.2	4.7	4.9	6.7
1999	3.9	19.1	5.4	1.3	*	2.9	4.3	6.1	5.7	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Legal intervention (Y35,Y89.0)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	0.1	0.3	0.3	0.2	0.3
2001	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.3
2000	0.1	*	*	*	*	0.1	0.2	0.3	0.2	0.3
1999	0.1	*	*	*	*	0.1	0.3	0.3	0.3	0.3
All races 3/, male										
2002	0.3	*	*	*	*	0.2	0.6	0.5	0.5	0.5
2001	0.3	*	*	*	*	0.2	0.6	0.5	0.6	0.6
2000	0.3	*	*	*	*	0.3	0.5	0.6	0.5	0.5
1999	0.3	*	*	*	*	0.2	0.5	0.7	0.7	0.6
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3
2001	0.1	*	*	*	*	*	0.3	0.2	0.3	0.3
2000	0.1	*	*	*	*	*	0.2	0.2	0.2	0.2
1999	0.1	*	*	*	*	*	0.1	0.3	0.3	0.3
White, male										
2002	0.2	*	*	*	*	*	0.4	0.4	0.5	0.5
2001	0.2	*	*	*	*	*	0.5	0.4	0.5	0.5
2000	0.2	*	*	*	*	*	0.4	0.4	0.4	0.4
1999	0.2	*	*	*	*	*	0.3	0.5	0.6	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Legal intervention (Y35,Y89.0)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.3	*	*	*	*	*	0.9	0.8	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	0.7
1999	0.3	*	*	*	*	*	0.8	0.9	*	0.7
Black, male										
2002	0.6	*	*	*	*	*	1.7	1.6	*	*
2001	0.4	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	1.4
1999	0.7	*	*	*	*	*	1.7	1.9	*	1.5
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6
1999	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8
All races 3/, male										
2002	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7
1999	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9
All races 3/, female										
2002	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7
White, both sexes										
2002	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000	1.3	1.4	0.2	*	0.2	0.6	1.3	1.6	1.8	2.7
1999	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8
White, male										
2002	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000	1.8	*	*	*	0.3	1.0	1.9	2.3	2.5	3.7
1999	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, female										
2002	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000	0.8	1.6	*	*	*	0.3	0.7	0.9	1.1	1.6
1999	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8
Black, both sexes										
2002	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000	1.8	*	*	*	*	*	1.4	1.7	2.2	3.3
1999	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9
Black, male										
2002	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000	2.8	*	*	*	*	*	2.5	2.7	3.4	4.8
1999	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3
Black, female										
2002	1.2	*	*	*	*	*	*	*	1.5	1.9
2001	0.9	*	*	*	*	*	*	*	*	2.0
2000	0.9	*	*	*	*	*	*	*	*	2.0
1999	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Discharge of firearms, undetermined intent (Y22-Y24)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	0.2	0.2	0.1	*	0.1
2001	0.1	*	*	*	*	0.1	0.2	0.2	*	0.1
2000	0.1	*	*	*	*	0.1	0.2	0.2	0.1	*
1999	0.1	*	*	*	*	0.3	0.3	0.2	*	0.1
All races 3/, male										
2002	0.1	*	*	*	*	0.3	0.4	0.2	*	0.2
2001	0.1	*	*	*	*	0.2	0.4	0.3	*	0.2
2000	0.1	*	*	*	*	0.2	0.4	0.3	*	*
1999	0.2	*	*	*	*	0.6	0.5	0.3	*	0.2
All races 3/, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.1	*	*	*	*	0.1	0.2	0.1	*	0.1
2001	0.1	*	*	*	*	*	0.2	0.2	*	0.2
2000	0.1	*	*	*	*	0.1	0.2	0.1	*	*
1999	0.1	*	*	*	*	0.3	0.3	0.1	*	0.1
White, male										
2002	0.1	*	*	*	*	0.3	0.3	*	*	*
2001	0.1	*	*	*	*	*	0.3	0.3	*	*
2000	0.1	*	*	*	*	*	0.3	*	*	*
1999	0.2	*	*	*	*	0.5	0.5	*	*	0.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Discharge of firearms, undetermined intent (Y22-Y24)										
White, female										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	0.6	*	*	*	*
Black, male										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002	1.6	2.5	0.4	0.1	0.2	0.5	1.5	1.7	2.2	3.2
2001	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9
2000	1.3	1.7	0.3	*	0.2	0.5	1.1	1.4	1.7	2.6
1999	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7
All races 3/, male										
2002	2.1	2.5	0.3	*	0.2	0.7	2.1	2.4	3.0	4.3
2001	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8
2000	1.8	1.4	0.3	*	0.3	0.7	1.6	2.0	2.4	3.6
1999	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7
All races 3/, female										
2002	1.1	2.5	0.4	*	*	0.3	0.9	1.0	1.4	2.0
2001	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0
2000	0.8	2.0	*	*	*	0.3	0.5	0.8	1.0	1.6
1999	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7
White, both sexes										
2002	1.6	1.8	0.4	*	0.2	0.6	1.7	1.8	2.2	3.3
2001	1.4	1.4	0.3	*	0.2	0.6	1.3	1.6	2.0	2.9
2000	1.3	1.4	0.2	*	0.1	0.5	1.1	1.5	1.8	2.6
1999	1.3	1.2	0.2	*	0.1	0.5	1.2	1.5	2.0	2.6
White, male										
2002	2.0	1.8	0.4	*	*	0.8	2.3	2.5	3.0	4.3
2001	1.9	1.7	*	*	0.3	0.9	1.9	2.2	2.7	3.6
2000	1.7	*	*	*	0.3	0.7	1.6	2.1	2.4	3.6
1999	1.7	1.4	*	*	*	0.8	1.8	2.0	2.7	3.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
White, female										
2002	1.2	1.8	0.4	*	*	0.4	1.0	1.1	1.5	2.2
2001	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.1
2000	0.8	1.6	*	*	*	0.3	0.6	0.9	1.1	1.6
1999	0.8	*	*	*	*	*	0.5	0.9	1.4	1.8
Black, both sexes										
2002	1.9	5.5	*	*	*	*	1.1	1.8	2.4	3.9
2001	1.7	5.5	*	*	*	*	0.8	1.3	1.9	3.8
2000	1.7	*	*	*	*	*	0.9	1.4	1.9	3.2
1999	1.7	*	*	*	*	*	1.1	1.5	2.4	3.8
Black, male										
2002	2.7	*	*	*	*	*	1.4	2.7	3.4	6.3
2001	2.5	6.0	*	*	*	*	1.5	2.2	2.7	5.9
2000	2.5	*	*	*	*	*	1.5	2.2	3.1	4.5
1999	2.7	*	*	*	*	*	1.8	2.7	3.5	6.1
Black, female										
2002	1.2	*	*	*	*	*	*	*	1.5	1.8
2001	0.8	*	*	*	*	*	*	*	*	2.0
2000	0.9	*	*	*	*	*	*	*	*	2.0
1999	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Operations of war and their sequelae (Y36,Y89.1)										
All races 3/, both sexes										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	0.0	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Operations of war and their sequelae (Y36,Y89.1)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Complications of medical and surgical care (Y40-Y84,Y88)										
All races 3/, both sexes										
2002	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3
All races 3/, male										
2002	0.9	*	*	*	*	*	*	*	0.3	0.3
2001	1.0	*	*	*	*	*	*	*	0.3	0.4
2000	1.0	*	*	*	*	*	*	0.2	0.2	0.3
1999	1.0	1.6	*	*	*	*	*	*	0.3	0.3
All races 3/, female										
2002	1.1	*	*	*	*	*	*	*	0.2	0.3
2001	1.1	*	*	*	*	*	*	*	0.2	0.3
2000	1.1	*	*	*	*	*	*	*	0.2	0.3
1999	1.1	1.2	*	*	*	*	*	0.2	*	0.3
White, both sexes										
2002	1.0	*	*	*	*	*	*	0.1	0.2	0.3
2001	1.1	*	*	*	*	*	0.1	*	0.2	0.3
2000	1.1	*	*	*	*	*	*	0.1	0.2	0.3
1999	1.0	1.2	0.2	*	*	*	*	*	0.2	0.3
White, male										
2002	0.9	*	*	*	*	*	*	*	0.3	0.3
2001	0.9	*	*	*	*	*	*	*	0.3	0.3
2000	1.1	*	*	*	*	*	*	*	0.2	0.3
1999	1.0	1.4	*	*	*	*	*	*	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Complications of medical and surgical care (Y40-Y84,Y88)										
White, female										
2002	1.1	*	*	*	*	*	*	*	*	0.3
2001	1.2	*	*	*	*	*	*	*	*	0.3
2000	1.1	*	*	*	*	*	*	*	*	0.3
1999	1.1	*	*	*	*	*	*	*	*	0.2
Black, both sexes										
2002	1.2	*	*	*	*	*	*	*	*	0.7
2001	1.4	*	*	*	*	*	*	*	*	0.8
2000	1.3	*	*	*	*	*	*	*	*	*
1999	1.2	*	*	*	*	*	*	*	*	*
Black, male										
2002	1.0	*	*	*	*	*	*	*	*	*
2001	1.4	*	*	*	*	*	*	*	*	*
2000	1.1	*	*	*	*	*	*	*	*	*
1999	1.1	*	*	*	*	*	*	*	*	*
Black, female										
2002	1.3	*	*	*	*	*	*	*	*	*
2001	1.4	*	*	*	*	*	*	*	*	*
2000	1.5	*	*	*	*	*	*	*	*	*
1999	1.3	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70,R78.0,X45,X65,Y15) 5/										
All races 3/, both sexes										
2002	6.9	*	*	*	*	0.2	0.3	0.8	2.0	5.3
2001	7.0	*	*	*	*	0.2	0.3	0.9	2.2	5.9
2000	6.9	*	*	*	*	0.2	0.3	0.9	2.3	5.8
1999	6.9	*	*	*	*	0.1	0.4	0.7	2.3	6.0
All races 3/, male										
2002	10.6	*	*	*	*	0.3	0.5	1.3	2.9	7.4
2001	10.7	*	*	*	*	0.3	0.6	1.2	3.2	8.3
2000	10.7	*	*	*	*	0.3	0.5	1.4	3.4	8.5
1999	10.7	*	*	*	*	0.2	0.6	1.2	3.1	8.5
All races 3/, female										
2002	3.3	*	*	*	*	*	*	0.3	1.1	3.3
2001	3.4	*	*	*	*	*	*	0.5	1.3	3.5
2000	3.2	*	*	*	*	*	*	0.3	1.1	3.2
1999	3.2	*	*	*	*	*	*	0.3	1.5	3.6
White, both sexes										
2002	7.1	*	*	*	*	0.2	0.3	0.8	2.1	5.4
2001	7.1	*	*	*	*	0.2	0.3	0.8	2.2	5.9
2000	7.0	*	*	*	*	0.2	0.3	0.8	2.2	5.8
1999	6.9	*	*	*	*	*	0.4	0.8	2.3	5.9
White, male										
2002	11.0	*	*	*	*	0.3	0.5	1.3	3.0	7.5
2001	10.8	*	*	*	*	0.3	0.5	1.2	3.2	8.2
2000	10.9	*	*	*	*	0.3	0.5	1.4	3.3	8.5
1999	10.7	*	*	*	*	*	0.6	1.2	3.1	8.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70,R78.0,X45,X65,Y15) 5/										
White, female										
2002	3.4	*	*	*	*	*	*	*	1.0	3.2
2001	3.4	*	*	*	*	*	*	0.4	1.2	3.5
2000	3.1	*	*	*	*	*	*	*	1.1	3.1
1999	3.1	*	*	*	*	*	*	*	1.4	3.4
Black, both sexes										
2002	6.3	*	*	*	*	*	*	*	1.7	4.3
2001	7.1	*	*	*	*	*	*	*	1.8	5.0
2000	7.2	*	*	*	*	*	*	0.9	2.5	5.3
1999	7.6	*	*	*	*	*	*	*	2.0	6.7
Black, male										
2002	9.7	*	*	*	*	*	*	*	2.5	6.0
2001	11.2	*	*	*	*	*	*	*	2.3	8.0
2000	11.1	*	*	*	*	*	*	*	3.3	7.6
1999	11.9	*	*	*	*	*	*	*	2.6	9.5
Black, female										
2002	3.2	*	*	*	*	*	*	*	*	2.9
2001	3.4	*	*	*	*	*	*	*	*	2.3
2000	3.7	*	*	*	*	*	*	*	1.7	3.2
1999	3.8	*	*	*	*	*	*	*	1.5	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Drug-induced causes (F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5, F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) 5/										
All races 3/, both sexes										
2002	9.0	0.7	0.2	*	0.2	3.1	7.9	9.8	12.4	17.5
2001	7.6	0.5	0.2	*	0.2	2.7	6.5	8.2	10.6	15.5
2000	7.0	*	*	*	0.2	2.2	5.9	7.5	10.1	14.4
1999	6.8	0.6	0.2	*	0.1	1.8	5.3	7.4	10.3	14.2
All races 3/, male										
2002	11.8	1.0	*	*	*	4.3	11.4	14.0	17.0	23.1
2001	10.2	*	0.3	*	*	3.7	9.6	11.9	14.4	20.7
2000	9.5	*	*	*	*	3.1	8.8	10.8	14.1	19.8
1999	9.4	*	*	*	*	2.4	7.7	10.8	14.6	19.6
All races 3/, female										
2002	6.3	*	0.3	*	*	1.8	4.2	5.4	7.8	11.9
2001	5.1	*	*	*	*	1.5	3.2	4.4	6.7	10.3
2000	4.6	*	*	*	*	1.4	2.9	4.1	6.1	9.0
1999	4.4	*	*	*	*	1.2	2.8	4.0	6.0	8.7
White, both sexes										
2002	9.4	*	*	*	0.2	3.6	9.2	10.9	13.5	18.5
2001	7.8	*	*	*	0.2	3.1	7.5	9.1	11.4	16.1
2000	7.1	*	*	*	0.2	2.6	6.7	8.2	10.7	14.8
1999	6.9	*	*	*	0.1	2.1	5.8	7.9	10.7	14.4
White, male										
2002	12.2	*	*	*	*	5.0	13.2	15.5	18.5	24.2
2001	10.4	*	*	*	*	4.4	10.9	13.1	15.6	21.6
2000	9.6	*	*	*	*	3.6	10.0	11.6	14.9	20.4
1999	9.3	*	*	*	*	2.8	8.5	11.3	15.2	19.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Drug-induced causes (F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5, F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) 5/										
White, female										
2002	6.7	*	*	*	*	2.1	4.9	6.1	8.3	12.6
2001	5.4	*	*	*	*	1.6	3.8	5.0	7.0	10.6
2000	4.7	*	*	*	*	1.5	3.2	4.6	6.3	9.1
1999	4.5	*	*	*	*	1.4	3.0	4.3	6.1	8.9
Black, both sexes										
2002	9.2	*	*	*	*	1.0	3.2	6.6	10.2	17.3
2001	8.5	*	*	*	*	0.9	3.0	5.8	9.5	16.7
2000	8.3	*	*	*	*	0.9	3.2	6.2	10.3	16.5
1999	8.6	*	*	*	*	0.7	3.7	6.8	11.2	16.4
Black, male										
2002	12.8	*	*	*	*	1.4	4.6	9.9	13.9	24.2
2001	12.2	*	*	*	*	*	5.1	9.3	12.3	22.4
2000	12.0	*	*	*	*	*	4.6	9.8	14.3	22.8
1999	12.7	*	*	*	*	*	5.6	11.0	15.7	23.8
Black, female										
2002	5.8	*	*	*	*	*	1.9	3.6	6.8	11.2
2001	5.1	*	*	*	*	*	*	2.6	7.0	11.6
2000	4.9	*	*	*	*	*	1.8	2.9	6.8	11.0
1999	4.8	*	*	*	*	*	1.9	3.1	7.3	9.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
All races 3/, both sexes										
2002	10.5	*	0.4	0.4	1.3	12.1	21.3	17.8	13.2	12.4
2001	10.4	*	0.5	0.4	1.2	12.4	21.2	17.7	13.5	12.2
2000	10.2	*	0.3	0.3	1.5	12.9	20.9	16.3	12.8	12.0
1999	10.3	*	0.4	0.4	1.7	14.4	21.0	16.6	13.2	12.3
All races 3/, male										
2002	18.4	*	0.5	0.4	1.9	21.1	37.6	31.1	22.3	20.6
2001	18.2	*	0.5	0.5	1.9	21.8	37.7	30.9	22.9	20.0
2000	17.8	*	0.4	0.4	2.4	22.4	37.0	28.5	21.4	19.5
1999	18.1	*	0.5	0.5	2.5	24.8	36.8	28.8	22.3	20.0
All races 3/, female										
2002	2.8	*	0.3	0.3	0.7	2.7	4.2	4.1	3.9	4.2
2001	2.8	*	0.4	0.3	0.5	2.6	3.8	4.1	3.9	4.6
2000	2.8	*	*	0.2	0.5	2.9	4.2	3.9	4.1	4.5
1999	2.9	*	0.4	0.3	0.8	3.4	4.4	4.1	4.1	4.6
White, both sexes										
2002	9.3	*	0.3	0.3	1.1	8.7	13.9	12.3	10.2	10.9
2001	9.4	*	0.4	0.4	1.1	8.9	14.0	12.9	11.1	11.2
2000	9.1	*	0.2	0.3	1.4	9.2	13.9	11.5	10.4	10.9
1999	9.2	*	0.3	0.3	1.6	10.2	14.0	12.1	11.1	11.3
White, male										
2002	16.1	*	0.4	0.3	1.5	14.9	23.9	20.5	16.7	17.7
2001	16.2	*	0.4	0.5	1.7	15.0	24.1	21.8	18.2	17.9
2000	15.6	*	*	0.4	2.3	15.6	23.9	19.3	16.9	17.2
1999	15.9	*	*	0.3	2.3	17.3	23.8	20.3	18.1	18.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
White, female										
2002	2.7	*	*	0.3	0.6	2.1	3.1	3.7	3.4	4.0
2001	2.7	*	*	0.3	0.5	2.3	3.1	3.4	3.7	4.3
2000	2.7	*	*	*	0.5	2.3	3.2	3.3	3.7	4.5
1999	2.8	*	*	*	0.7	2.6	3.5	3.4	3.8	4.5
Black, both sexes										
2002	20.2	*	*	*	2.4	31.4	64.7	54.8	35.1	24.4
2001	19.3	*	0.9	*	1.9	32.9	64.4	50.4	31.5	22.6
2000	19.3	*	*	*	2.2	33.9	64.0	48.1	30.9	21.7
1999	19.4	*	1.1	0.9	2.3	38.0	63.5	46.5	29.0	20.9
Black, male										
2002	37.8	*	*	*	3.5	56.0	120.6	106.3	65.9	44.8
2001	36.4	*	*	*	3.0	60.5	123.4	96.0	60.1	39.5
2000	36.1	*	*	*	3.7	61.5	121.3	92.9	57.0	39.9
1999	36.0	*	*	1.3	3.4	67.1	119.9	88.6	54.3	37.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
Black, female										
2002	4.2	*	*	*	1.3	6.0	10.2	7.7	7.4	6.3
2001	3.8	*	*	*	*	4.5	7.8	8.9	5.8	7.5
2000	4.0	*	*	*	*	5.7	9.6	7.5	7.4	5.5
1999	4.4	*	*	*	*	8.2	9.9	8.4	6.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
All races 3/, both sexes										
2002	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000	239.1	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
All races 3/, male										
2002	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000	305.9	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
All races 3/, female										
2002	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000	173.4	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3
White, both sexes										
2002	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000	215.9	317.9	467.4	743.9	1,184.9	1,844.5	2,866.8	4,496.6	7,333.2	15,707.0
1999	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0
White, male										
2002	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000	279.4	416.1	588.3	925.1	1,465.1	2,285.1	3,593.1	5,620.3	9,092.5	17,716.4
1999	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
White, female										
2002	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000	152.2	220.7	349.3	571.5	926.5	1,459.3	2,284.0	3,706.2	6,286.6	14,890.7
1999	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6
Black, both sexes										
2002	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000	434.5	666.6	938.7	1,347.0	1,929.0	2,769.0	4,019.7	5,782.6	8,161.0	14,714.3
1999	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9
Black, male										
2002	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000	549.1	853.0	1,227.4	1,776.2	2,462.1	3,513.8	5,203.9	7,410.0	10,405.7	16,791.0
1999	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5
Black, female										
2002	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000	333.3	505.0	692.6	995.2	1,508.3	2,219.3	3,233.3	4,801.6	7,030.5	13,941.3
1999	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Salmonella infections (A01-A02)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Salmonella infections (A01-A02)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)										
All races 3/, both sexes										
2002	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.7	21.0
2001	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.8
2000	*	*	*	0.2	0.3	0.9	1.5	2.7	5.3	11.8
1999	*	*	*	*	0.3	0.7	1.2	2.3	4.5	10.3
All races 3/, male										
2002	*	*	*	0.4	0.8	1.5	3.1	5.1	10.1	20.4
2001	*	*	*	*	0.4	0.7	2.0	3.5	6.2	12.1
2000	*	*	*	0.3	*	1.0	1.6	2.9	5.8	11.4
1999	*	*	*	*	*	0.6	1.2	2.7	4.9	9.5
All races 3/, female										
2002	*	*	*	0.4	0.6	1.3	2.3	5.0	9.5	21.3
2001	*	*	0.2	*	0.5	0.8	1.8	3.0	6.5	13.1
2000	*	*	*	*	*	0.8	1.5	2.6	4.9	11.9
1999	*	*	*	*	*	0.8	1.3	2.0	4.3	10.7
White, both sexes										
2002	*	0.1	0.2	0.4	0.6	1.5	2.7	5.3	10.3	22.0
2001	*	*	0.2	0.2	0.4	0.7	2.0	3.3	6.6	13.5
2000	*	*	*	0.3	0.3	0.9	1.6	2.9	5.6	12.2
1999	*	*	*	*	0.3	0.7	1.3	2.3	4.7	10.9
White, male										
2002	*	*	*	0.4	0.7	1.6	3.0	5.3	10.7	21.4
2001	*	*	*	*	*	0.7	2.0	3.6	6.2	12.7
2000	*	*	*	*	*	1.0	1.6	3.0	6.0	11.9
1999	*	*	*	*	*	0.7	1.3	2.8	5.2	9.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)										
White, female										
2002	*	*	*	0.4	0.6	1.4	2.4	5.3	10.1	22.3
2001	*	*	0.3	*	0.5	0.8	1.9	3.2	6.8	13.9
2000	*	*	*	*	*	0.8	1.6	2.7	5.3	12.3
1999	*	*	*	*	*	0.8	1.3	2.1	4.4	11.3
Black, both sexes										
2002	*	*	*	*	*	*	2.9	*	*	13.1
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	8.1
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	13.1
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
All races 3/, both sexes										
2002	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2
All races 3/, male										
2002	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000	*	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1
All races 3/, female										
2002	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000	*	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0
White, both sexes										
2002	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2
White, male										
2002	*	*	*	*	0.5	0.8	1.0	1.3	2.5	3.2
2001	*	*	*	0.5	0.5	0.6	0.8	1.4	1.8	3.1
2000	*	*	0.3	*	*	0.7	0.8	1.5	2.3	3.8
1999	*	0.3	0.4	0.5	0.6	0.9	1.1	1.5	3.0	5.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
White, female										
2002	*	*	*	*	*	*	*	1.1	1.3	1.6
2001	*	*	*	*	*	*	0.5	0.9	1.2	1.8
2000	*	*	*	*	*	*	0.5	1.0	1.4	1.6
1999	*	*	*	*	*	*	0.5	1.1	1.4	2.4
Black, both sexes										
2002	*	*	*	*	*	2.7	*	*	5.2	*
2001	*	*	1.1	1.6	*	*	3.4	*	*	6.8
2000	*	0.9	1.1	*	*	2.8	3.6	5.2	8.8	*
1999	*	0.9	1.2	1.7	*	*	3.8	5.5	*	8.6
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	19.4	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
All races 3/, both sexes										
2002	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000	0.1	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7
All races 3/, male										
2002	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4
All races 3/, female										
2002	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000	*	*	*	*	*	*	0.6	1.1	1.4	1.6
1999	*	*	*	*	*	*	0.5	1.2	1.4	2.6
White, both sexes										
2002	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0
White, male										
2002	*	*	*	*	*	0.7	0.8	0.9	2.0	2.9
2001	*	*	*	0.4	0.5	*	0.7	1.2	1.4	2.6
2000	*	*	0.3	*	*	0.5	0.7	1.3	1.9	3.4
1999	*	*	*	0.4	0.5	0.7	1.0	1.3	2.6	5.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
White, female										
2002	*	*	*	*	*	*	*	0.9	1.1	1.3
2001	*	*	*	*	*	*	*	0.6	1.0	1.4
2000	*	*	*	*	*	*	*	0.7	1.2	1.2
1999	*	*	*	*	*	*	*	0.9	1.1	2.1
Black, both sexes										
2002	*	*	*	*	*	*	*	*	5.2	*
2001	*	*	*	*	*	*	*	*	*	6.1
2000	*	*	*	*	*	2.3	2.9	4.3	6.8	*
1999	*	*	*	1.5	*	*	2.7	4.6	*	7.3
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other tuberculosis (A17-A19)										
All races 3/, both sexes										
2002	*	*	*	*	*	0.2	0.2	0.4	*	*
2001	*	*	*	*	*	*	0.3	0.3	0.4	*
2000	*	*	*	*	*	*	*	0.3	*	*
1999	*	*	*	*	*	*	0.2	0.3	*	0.5
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other tuberculosis (A17-A19)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Whooping cough (A37)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Whooping cough (A37)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningococcal infection (A39)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningococcal infection (A39)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
All races 3/, both sexes										
2002	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000	2.5	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7
All races 3/, male										
2002	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000	2.6	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5
All races 3/, female										
2002	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000	2.3	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5
White, both sexes										
2002	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000	2.0	3.0	4.9	7.8	12.6	20.7	34.5	58.0	97.4	203.0
1999	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9
White, male										
2002	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000	2.2	3.5	5.5	8.4	14.0	23.1	39.4	66.7	114.0	209.8
1999	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
White, female										
2002	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000	1.9	2.5	4.3	7.2	11.3	18.7	30.6	51.9	87.6	200.2
1999	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4
Black, both sexes										
2002	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000	5.8	11.0	16.9	24.3	40.2	59.7	86.6	146.4	216.3	392.6
1999	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3
Black, male										
2002	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000	5.8	12.4	21.3	27.8	45.1	68.9	97.0	182.9	250.4	427.8
1999	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5
Black, female										
2002	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000	5.8	9.7	13.2	21.5	36.3	52.8	79.7	124.5	199.1	379.5
1999	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Syphilis (A50-A53)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Syphilis (A50-A53)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute poliomyelitis (A80)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute poliomyelitis (A80)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Measles (B05)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Measles (B05)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2002	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7
All races 3/, male										
2002	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9
All races 3/, female										
2002	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9
White, both sexes										
2002	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
1999	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5
White, male										
2002	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
1999	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
White, female										
2002	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
1999	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7
Black, both sexes										
2002	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
1999	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4
Black, male										
2002	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
1999	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*
Black, female										
2002	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*
1999	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)										
All races 3/, both sexes										
2002	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000	13.5	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*
All races 3/, male										
2002	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000	20.4	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*
All races 3/, female										
2002	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000	6.7	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*
White, both sexes										
2002	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000	7.5	6.9	4.7	3.4	2.1	1.3	0.7	0.5	*	*
1999	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*
White, male										
2002	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000	12.3	11.7	8.3	6.0	3.7	2.5	1.3	1.0	*	*
1999	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)										
White, female										
2002	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999	2.8	2.0	1.2	0.7	*	*	*	*	*	*
Black, both sexes										
2002	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000	57.3	58.7	46.6	29.3	22.0	15.1	11.6	6.2	*	*
1999	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*
Black, male										
2002	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000	84.1	92.7	77.4	51.0	39.5	26.3	22.4	11.8	*	*
1999	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*
Black, female										
2002	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000	33.7	29.2	20.4	11.5	8.2	6.9	4.4	*	*	*
1999	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malaria (B50-B54)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malaria (B50-B54)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)										
All races 3/, both sexes										
2002	1.3	1.9	2.7	3.0	3.8	5.4	8.1	11.6	13.8	21.7
2001	0.9	1.5	1.9	2.4	3.9	5.6	6.9	9.4	12.3	20.1
2000	0.8	1.4	1.8	2.4	3.6	4.7	7.0	10.0	12.9	21.2
1999	1.4	2.0	2.6	3.3	4.4	6.8	8.9	12.6	16.6	27.0
All races 3/, male										
2002	1.7	2.6	3.8	3.8	4.7	6.5	9.7	14.0	17.7	26.8
2001	1.2	2.1	2.4	3.0	4.5	6.2	7.5	11.4	15.9	23.6
2000	1.0	1.7	2.3	2.8	4.0	5.7	8.8	12.1	16.2	27.6
1999	1.7	2.4	3.0	3.5	5.2	8.0	10.5	15.3	20.4	31.9
All races 3/, female										
2002	0.8	1.2	1.7	2.2	3.0	4.5	6.8	9.9	11.5	19.5
2001	0.6	0.9	1.5	1.8	3.3	5.1	6.5	8.0	10.1	18.7
2000	0.6	1.1	1.3	2.1	3.2	3.9	5.6	8.5	10.9	18.6
1999	1.1	1.6	2.3	3.1	3.7	5.9	7.7	10.8	14.4	25.1
White, both sexes										
2002	1.1	1.8	2.5	2.6	3.6	5.4	7.9	11.3	13.8	22.0
2001	0.8	1.4	1.7	2.2	3.6	5.4	6.7	9.3	12.1	20.1
2000	0.7	1.3	1.6	2.3	3.2	4.6	6.8	9.8	12.7	21.0
1999	1.1	1.6	2.2	2.9	3.9	6.4	8.5	12.1	16.1	26.5
White, male										
2002	1.5	2.5	3.5	3.2	4.3	6.6	9.4	13.6	17.6	27.2
2001	1.0	1.9	2.2	2.8	4.3	6.0	7.2	11.3	15.9	23.8
2000	0.9	1.5	2.0	2.7	3.6	5.5	8.6	11.6	16.1	26.9
1999	1.4	2.0	2.6	3.3	4.5	7.5	10.2	14.8	18.6	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)										
White, female										
2002	0.7	1.1	1.5	2.0	2.9	4.3	6.7	9.7	11.5	19.8
2001	0.6	0.8	1.3	1.6	3.1	5.0	6.2	7.8	9.9	18.5
2000	0.6	1.0	1.1	1.9	2.8	3.8	5.4	8.6	10.8	18.7
1999	0.7	1.2	1.9	2.6	3.4	5.6	7.2	10.2	14.6	24.8
Black, both sexes										
2002	2.5	3.2	4.8	6.0	5.4	5.3	9.7	15.3	11.5	21.7
2001	1.8	2.5	3.4	4.4	6.0	7.1	9.8	9.7	13.3	20.6
2000	1.7	2.6	3.5	4.0	5.9	6.1	9.9	11.6	16.1	24.1
1999	3.6	5.1	5.9	6.3	8.2	10.7	13.1	18.7	23.4	34.1
Black, male										
2002	3.2	4.5	6.7	9.2	7.3	6.5	12.0	18.4	*	28.9
2001	2.7	3.6	4.3	5.5	7.3	9.7	10.4	10.2	16.1	*
2000	2.1	3.2	4.8	4.4	7.2	8.6	12.4	17.0	20.2	32.3
1999	3.6	5.8	6.5	5.8	10.4	12.5	15.9	21.8	41.9	45.7
Black, female										
2002	1.9	2.0	3.1	3.4	3.9	4.4	8.1	13.4	9.9	18.9
2001	*	1.6	2.7	3.6	5.0	5.3	9.4	9.4	11.9	20.2
2000	*	2.1	2.4	3.6	4.9	4.2	8.2	8.3	14.0	21.0
1999	3.5	4.5	5.3	6.8	6.5	9.4	11.3	16.9	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2002	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8
All races 3/, male										
2002	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2
All races 3/, female										
2002	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5
White, both sexes										
2002	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
1999	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8
White, male										
2002	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
1999	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
White, female										
2002	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
1999	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2
Black, both sexes										
2002	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
1999	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4
Black, male										
2002	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
1999	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7
Black, female										
2002	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7
1999	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes										
2002	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3
All races 3/, male										
2002	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1
All races 3/, female										
2002	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1
White, both sexes										
2002	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
1999	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8
White, male										
2002	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
1999	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
White, female										
2002	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
1999	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7
Black, both sexes										
2002	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
1999	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3
Black, male										
2002	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
1999	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6
Black, female										
2002	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
1999	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)										
All races 3/, both sexes										
2002	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8
All races 3/, male										
2002	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0
All races 3/, female										
2002	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3
White, both sexes										
2002	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
1999	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2
White, male										
2002	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
1999	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of esophagus (C15)										
White, female										
2002	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
1999	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9
Black, both sexes										
2002	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
1999	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9
Black, male										
2002	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
1999	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6
Black, female										
2002	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3
1999	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)										
All races 3/, both sexes										
2002	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1
All races 3/, male										
2002	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3
All races 3/, female										
2002	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6
White, both sexes										
2002	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
1999	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8
White, male										
2002	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
1999	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of stomach (C16)										
White, female										
2002	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
1999	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3
Black, both sexes										
2002	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
1999	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4
Black, male										
2002	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
1999	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6
Black, female										
2002	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
1999	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2002	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7
All races 3/, male										
2002	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0
All races 3/, female										
2002	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6
White, both sexes										
2002	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
1999	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9
White, male										
2002	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
1999	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)										
White, female										
2002	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
1999	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3
Black, both sexes										
2002	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
1999	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1
Black, male										
2002	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
1999	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4
Black, female										
2002	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0
1999	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes										
2002	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6
All races 3/, male										
2002	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1
All races 3/, female										
2002	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8
White, both sexes										
2002	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
1999	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9
White, male										
2002	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
1999	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
White, female										
2002	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
1999	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6
Black, both sexes										
2002	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
1999	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4
Black, male										
2002	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
1999	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5
Black, female										
2002	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
1999	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)										
All races 3/, both sexes										
2002	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7
All races 3/, male										
2002	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5
All races 3/, female										
2002	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6
White, both sexes										
2002	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
1999	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6
White, male										
2002	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
1999	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of pancreas (C25)										
White, female										
2002	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
1999	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0
Black, both sexes										
2002	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
1999	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6
Black, male										
2002	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
1999	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8
Black, female										
2002	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6
1999	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)										
All races 3/, both sexes										
2002	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0
All races 3/, male										
2002	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8
All races 3/, female										
2002	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6
White, both sexes										
2002	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
1999	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7
White, male										
2002	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
1999	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of larynx (C32)										
White, female										
2002	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
1999	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7
Black, both sexes										
2002	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
1999	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3
Black, male										
2002	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
1999	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6
Black, female										
2002	*	*	*	*	*	*	4.7	*	*	*
2001	*	*	*	*	3.4	*	5.6	*	*	*
2000	*	*	*	*	*	3.8	5.5	7.4	*	*
1999	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2002	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0
All races 3/, male										
2002	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2
All races 3/, female										
2002	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5
White, both sexes										
2002	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
1999	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8
White, male										
2002	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
1999	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
White, female										
2002	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
1999	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3
Black, both sexes										
2002	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
1999	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5
Black, male										
2002	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
1999	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2
Black, female										
2002	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9
1999	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2002	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4
All races 3/, male										
2002	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5
All races 3/, female										
2002	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9
White, both sexes										
2002	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
1999	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6
White, male										
2002	2.5	3.8	5.1	6.9	9.7	12.7	16.4	21.8	25.6	31.7
2001	2.6	3.8	5.5	7.2	10.0	13.3	18.0	21.5	26.1	32.7
2000	2.5	3.8	6.0	7.3	9.9	12.5	16.2	20.1	26.6	34.7
1999	2.7	4.1	5.3	7.7	9.5	13.4	16.2	21.5	24.5	29.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
White, female										
2002	1.5	1.7	2.9	3.2	4.2	5.5	6.7	8.8	10.7	13.8
2001	1.7	2.3	3.1	3.5	4.2	4.8	7.0	8.0	9.8	13.7
2000	1.6	2.5	2.8	3.4	4.6	5.7	6.9	8.2	9.7	14.5
1999	1.5	2.2	3.0	3.3	4.6	5.8	6.4	8.8	9.9	12.8
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	6.6
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2002	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4
All races 3/, male										
2002	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5
All races 3/, female										
2002	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8
White, both sexes										
2002	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
1999	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0
White, male										
2002	*	*	0.3	0.4	0.6	0.6	1.3	1.7	2.7	3.3
2001	*	*	0.3	0.5	0.7	1.2	1.6	2.1	2.3	3.3
2000	*	*	0.3	0.6	0.7	1.1	1.7	2.2	2.3	3.9
1999	*	*	0.3	0.5	0.6	1.0	1.3	1.8	2.2	4.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
White, female										
2002	13.9	23.2	36.3	48.6	63.1	76.2	93.3	114.7	142.0	192.6
2001	14.7	24.5	38.1	50.1	63.4	75.7	95.8	117.0	140.0	189.9
2000	15.0	24.4	38.7	51.8	65.4	80.2	98.6	119.7	144.9	205.5
1999	14.2	23.9	38.9	51.7	64.4	79.2	98.6	117.4	146.3	202.9
Black, both sexes										
2002	15.3	22.9	32.1	40.5	46.3	54.5	67.2	78.5	117.5	152.6
2001	14.8	22.6	33.1	42.4	47.4	58.0	70.4	83.5	105.9	152.0
2000	14.8	23.9	32.8	43.9	47.3	52.7	65.6	84.2	100.5	176.4
1999	14.6	24.2	34.7	44.4	48.6	57.8	69.9	88.0	104.9	147.3
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7
1999	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)										
All races 3/, both sexes										
2002	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1
White, both sexes										
2002	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
1999	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)										
White, female										
2002	3.1	3.7	4.3	4.9	4.5	5.2	5.4	6.2	5.7	7.3
2001	3.5	4.3	4.6	4.6	5.5	4.4	5.5	5.8	7.6	8.8
2000	3.1	4.0	4.7	5.3	5.0	5.5	5.9	6.5	8.0	8.8
1999	3.4	4.5	5.0	5.2	5.7	5.9	5.5	6.8	6.5	9.5
Black, both sexes										
2002	3.0	3.7	4.6	4.2	6.0	8.2	10.1	9.6	12.6	17.5
2001	3.0	3.2	4.4	4.6	5.9	7.9	7.7	9.7	15.2	19.0
2000	3.1	3.7	4.9	6.2	6.4	9.4	9.7	10.8	13.6	19.7
1999	3.0	3.0	4.1	6.1	8.3	8.7	13.1	12.3	9.5	21.7
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
1999	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
All races 3/, both sexes										
2002	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7
White, both sexes										
2002	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
1999	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
White, female										
2002	0.9	1.7	3.7	6.3	10.1	13.6	19.6	22.4	26.3	34.2
2001	0.8	1.7	3.8	6.1	9.9	13.6	18.9	23.5	29.8	34.7
2000	0.8	1.9	3.5	6.7	10.3	13.6	17.8	22.2	28.0	34.3
1999	0.9	1.8	3.2	5.9	9.9	14.2	18.4	23.3	30.0	32.8
Black, both sexes										
2002	*	1.1	2.9	6.6	10.8	20.0	25.2	27.8	29.0	38.9
2001	*	1.4	2.9	4.6	11.6	18.8	24.2	26.6	31.1	32.2
2000	*	1.0	2.4	5.3	12.0	19.6	26.4	27.7	35.0	33.2
1999	0.8	*	2.2	5.8	10.5	20.9	21.8	25.7	29.8	35.4
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5
1999	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)										
All races 3/, both sexes										
2002	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8
White, both sexes										
2002	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
1999	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of ovary (C56)										
White, female										
2002	2.9	6.6	11.4	16.8	25.9	33.9	41.6	53.4	62.5	55.9
2001	2.9	6.2	11.4	16.6	25.3	33.9	42.2	53.1	61.9	57.3
2000	3.0	6.4	11.1	17.0	24.5	33.7	40.3	54.5	60.0	59.0
1999	3.2	6.5	11.2	16.8	25.1	32.8	41.0	50.5	57.9	54.3
Black, both sexes										
2002	1.2	2.9	4.1	7.7	10.9	14.9	20.0	26.5	32.4	34.7
2001	1.4	1.9	4.4	7.9	12.6	16.3	20.7	26.2	27.9	36.5
2000	1.5	2.2	4.8	6.4	10.9	16.8	19.6	26.1	27.4	34.4
1999	1.1	2.4	5.0	7.2	11.4	15.0	22.7	28.9	28.3	28.7
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
1999	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of prostate (C61)										
All races 3/, both sexes										
2002	0.1	0.4	1.6	4.6	11.4	24.2	45.8	79.7	128.7	195.0
2001	0.1	0.5	1.7	4.6	11.3	25.0	47.4	82.4	133.2	200.9
2000	*	0.5	1.5	4.6	12.1	25.9	49.0	85.1	136.6	209.3
1999	0.1	0.5	1.6	4.9	12.3	28.4	51.9	89.0	141.4	207.5
All races 3/, male										
2002	0.2	0.8	3.3	9.6	24.0	52.2	103.2	192.6	340.3	644.5
2001	0.2	1.0	3.5	9.4	23.8	53.9	107.1	200.1	355.4	681.2
2000	*	1.1	3.1	9.6	25.4	56.0	111.3	207.4	368.2	723.1
1999	0.3	1.0	3.3	10.1	25.8	61.6	118.3	216.9	383.0	722.4
All races 3/, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
White, both sexes										
2002	*	0.3	1.3	3.9	9.6	21.2	40.7	72.6	120.3	186.7
2001	*	0.4	1.4	3.9	9.7	22.1	42.2	75.3	123.3	189.4
2000	*	0.4	1.3	3.8	10.4	22.4	43.2	77.6	125.8	198.6
1999	0.1	0.4	1.4	4.2	10.2	24.9	46.4	81.1	132.9	195.2
White, male										
2002	*	0.5	2.7	8.1	20.1	45.2	90.7	174.1	316.4	616.3
2001	*	0.8	2.8	8.0	20.2	47.3	94.3	181.5	327.2	641.8
2000	*	0.8	2.6	7.9	21.6	48.0	97.0	188.0	337.4	687.5
1999	0.2	0.8	2.8	8.6	21.2	53.4	104.7	196.5	358.0	681.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of prostate (C61)										
White, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, both sexes										
2002	*	1.2	4.2	11.7	29.4	57.9	110.6	176.8	261.9	328.4
2001	*	1.1	4.8	11.8	28.7	57.4	114.7	180.8	280.3	362.9
2000	*	1.5	3.6	12.9	29.5	64.7	120.7	190.1	292.8	356.6
1999	*	1.4	4.2	12.0	33.4	68.7	123.4	196.6	273.0	366.4
Black, male										
2002	*	2.6	9.1	25.9	66.9	135.9	275.4	472.6	765.0	1,185.7
2001	*	2.4	10.4	26.1	65.1	135.2	286.8	483.7	830.7	1,341.2
2000	*	3.2	7.8	28.7	66.9	152.5	302.6	505.6	874.2	1,314.7
1999	*	2.9	9.2	26.6	76.0	161.5	309.3	522.9	824.0	1,346.3
Black, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
All races 3/, both sexes										
2002	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4
All races 3/, male										
2002	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4
All races 3/, female										
2002	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1
White, both sexes										
2002	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
1999	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5
White, male										
2002	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
1999	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
White, female										
2002	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
1999	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2
Black, both sexes										
2002	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
1999	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2
Black, male										
2002	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
1999	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2
Black, female										
2002	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
1999	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)										
All races 3/, both sexes										
2002	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5
All races 3/, male										
2002	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8
All races 3/, female										
2002	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8
White, both sexes										
2002	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
1999	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8
White, male										
2002	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
1999	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of bladder (C67)										
White, female										
2002	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
1999	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4
Black, both sexes										
2002	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
1999	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3
Black, male										
2002	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
1999	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7
Black, female										
2002	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2
1999	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes										
2002	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0
All races 3/, male										
2002	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7
All races 3/, female										
2002	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1
White, both sexes										
2002	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
1999	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6
White, male										
2002	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
1999	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
White, female										
2002	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
1999	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5
Black, both sexes										
2002	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
1999	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9
Black, male										
2002	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
1999	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*
Black, female										
2002	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
1999	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All races 3/, both sexes										
2002	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4
All races 3/, male										
2002	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0
All races 3/, female										
2002	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3
White, both sexes										
2002	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
1999	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1
White, male										
2002	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
1999	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, female										
2002	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
1999	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3
Black, both sexes										
2002	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
1999	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0
Black, male										
2002	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
1999	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3
Black, female										
2002	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1
1999	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
All races 3/, both sexes										
2002	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4
All races 3/, male										
2002	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8
All races 3/, female										
2002	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8
White, both sexes										
2002	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
1999	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4
White, male										
2002	0.4	0.5	0.5	0.7	0.9	1.5	2.3	2.1	3.5	3.9
2001	0.5	0.5	0.7	0.6	0.9	1.6	1.8	2.5	3.0	2.7
2000	0.4	0.5	0.6	0.5	1.1	1.4	1.9	2.6	2.8	3.9
1999	0.5	0.5	0.6	0.9	1.3	1.7	2.4	2.5	3.0	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
White, female										
2002	0.3	0.3	0.5	*	0.7	0.6	1.2	1.6	1.7	2.5
2001	0.3	0.4	0.4	*	0.7	0.8	0.9	1.8	2.4	2.2
2000	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.6	2.4	2.1
1999	0.3	0.4	0.4	0.5	0.6	0.9	1.6	2.0	2.2	1.8
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2002	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7
All races 3/, male										
2002	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4
All races 3/, female										
2002	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9
White, both sexes										
2002	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
1999	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2
White, male										
2002	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
1999	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
White, female										
2002	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
1999	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4
Black, both sexes										
2002	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
1999	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5
Black, male										
2002	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
1999	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8
Black, female										
2002	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5
1999	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
All races 3/, both sexes										
2002	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4
All races 3/, male										
2002	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2
All races 3/, female										
2002	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6
White, both sexes										
2002	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
1999	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4
White, male										
2002	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
1999	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
White, female										
2002	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
1999	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9
Black, both sexes										
2002	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
1999	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8
Black, male										
2002	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
1999	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3
Black, female										
2002	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
1999	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
All races 3/, both sexes										
2002	0.4	1.3	2.5	4.7	8.3	14.1	21.0	27.9	35.8	36.7
2001	0.5	1.2	2.3	5.1	8.5	14.0	21.1	28.3	34.7	36.3
2000	0.5	1.2	2.6	4.5	8.5	14.4	20.4	28.3	36.4	37.8
1999	0.5	1.2	2.5	5.6	9.0	14.1	20.6	27.6	36.2	36.6
All races 3/, male										
2002	0.5	1.5	3.0	5.6	10.2	16.8	25.8	35.5	46.2	50.8
2001	0.7	1.5	2.8	6.0	10.1	16.8	25.3	35.6	43.4	48.0
2000	0.6	1.4	2.9	4.9	9.5	17.1	24.2	35.7	46.1	51.8
1999	0.6	1.4	3.0	6.5	11.0	16.6	24.1	34.1	45.5	51.3
All races 3/, female										
2002	0.4	1.0	2.0	4.0	6.5	11.7	17.2	22.5	29.5	30.6
2001	0.4	0.9	1.9	4.2	7.1	11.5	17.7	23.2	29.4	31.3
2000	0.4	1.0	2.3	4.2	7.6	12.1	17.3	23.1	30.7	32.2
1999	0.4	1.1	2.1	4.7	7.3	12.0	17.8	23.0	30.7	30.6
White, both sexes										
2002	0.4	1.1	2.1	4.2	7.5	13.0	19.9	26.9	34.8	35.7
2001	0.4	1.0	2.1	4.4	7.5	12.9	19.9	27.1	33.0	35.0
2000	0.4	1.0	2.2	3.8	7.8	13.4	19.1	26.5	35.2	35.2
1999	0.4	1.1	2.1	4.7	7.9	13.2	19.2	25.7	34.9	34.8
White, male										
2002	0.4	1.3	2.7	4.9	9.4	15.8	24.9	34.8	45.3	50.3
2001	0.6	1.3	2.5	5.2	9.0	15.7	24.1	34.6	41.7	47.1
2000	0.5	1.2	2.5	4.3	8.8	16.1	22.8	33.9	45.4	50.2
1999	0.4	1.3	2.6	5.6	10.0	15.8	22.5	32.4	44.3	49.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
White, female										
2002	0.3	0.8	1.6	3.4	5.7	10.5	15.8	21.2	28.4	29.4
2001	*	0.7	1.6	3.7	6.1	10.4	16.5	21.7	27.8	29.9
2000	0.2	0.8	2.0	3.4	6.8	11.1	16.2	21.3	29.2	29.1
1999	0.3	1.0	1.5	3.8	5.9	10.9	16.6	21.0	29.4	28.9
Black, both sexes										
2002	1.0	2.9	5.6	10.7	16.3	26.7	37.7	44.7	54.9	55.5
2001	1.5	2.6	4.8	11.8	19.4	25.8	38.2	47.9	61.2	58.3
2000	1.1	3.1	6.3	11.7	16.5	27.3	37.9	53.3	57.9	71.6
1999	1.4	2.4	6.6	13.8	20.0	24.2	37.6	51.1	56.3	63.1
Black, male										
2002	*	3.5	5.9	12.4	20.2	31.4	44.7	52.4	70.3	71.8
2001	1.6	3.4	5.3	14.5	23.4	30.2	45.1	57.1	80.3	71.5
2000	*	3.3	7.2	12.3	18.4	31.0	48.1	67.1	69.1	79.6
1999	1.9	2.6	6.7	16.1	21.6	26.2	46.8	59.4	74.1	80.9
Black, female										
2002	*	2.4	5.3	9.3	13.3	23.1	32.9	40.1	46.9	49.3
2001	1.5	2.0	4.3	9.5	16.3	22.5	33.6	42.4	51.5	53.5
2000	*	2.8	5.5	11.2	15.0	24.6	31.1	45.0	52.2	68.7
1999	*	2.3	6.6	11.9	18.8	22.7	31.5	46.1	47.5	56.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	0.5
2000	*	*	*	*	*	*	*	*	*	0.5
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)										
All races 3/, both sexes										
2002	5.8	10.4	17.5	28.8	46.5	66.0	97.0	131.0	168.7	230.1
2001	5.9	10.5	18.0	30.1	46.4	67.2	98.1	133.4	170.9	234.5
2000	5.9	10.5	18.2	30.4	48.2	71.5	100.4	136.0	177.4	239.8
1999	6.7	11.4	19.1	32.5	52.5	76.1	107.8	142.3	188.5	250.4
All races 3/, male										
2002	6.2	11.7	20.8	33.0	56.5	79.9	118.9	161.8	207.8	290.8
2001	6.4	12.0	21.5	34.9	55.4	79.4	120.3	162.3	211.0	295.2
2000	6.4	12.0	20.6	35.1	57.2	86.2	122.5	167.5	221.4	300.7
1999	7.6	13.2	22.0	38.2	62.6	90.6	132.8	174.0	233.0	318.9
All races 3/, female										
2002	5.4	9.1	14.3	24.9	37.5	53.9	79.5	109.2	144.9	203.8
2001	5.3	9.1	14.7	25.7	38.3	56.7	80.5	113.1	146.8	209.1
2000	5.3	9.1	15.8	25.9	40.0	59.0	83.1	114.1	151.4	215.0
1999	5.8	9.7	16.3	27.1	43.3	63.8	88.2	120.3	162.5	222.8
White, both sexes										
2002	5.5	9.9	16.8	28.1	45.7	65.6	96.9	131.7	170.0	233.0
2001	5.6	9.9	17.2	29.0	45.3	66.5	97.4	134.1	171.4	238.1
2000	5.4	9.9	17.2	29.7	47.1	71.4	99.9	135.6	179.5	243.7
1999	6.3	10.8	18.3	31.1	51.2	76.3	107.1	142.3	189.4	252.4
White, male										
2002	5.7	11.1	20.0	32.0	55.1	80.2	118.8	162.2	210.9	296.5
2001	6.2	11.3	20.1	33.1	53.5	78.0	119.7	163.1	212.0	300.2
2000	5.8	11.2	19.5	34.1	54.7	86.2	121.8	165.7	225.2	307.0
1999	7.2	12.2	20.8	36.6	60.9	90.1	131.8	174.6	235.3	323.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)										
White, female										
2002	5.4	8.7	13.7	24.3	36.9	52.7	79.1	109.9	144.9	205.5
2001	5.1	8.5	14.3	25.1	37.7	56.3	79.3	113.7	146.8	212.1
2000	5.0	8.6	14.8	25.6	40.1	58.4	82.4	114.5	152.3	218.0
1999	5.3	9.5	15.8	26.0	42.3	64.3	87.5	119.6	162.3	223.8
Black, both sexes										
2002	8.0	15.8	25.3	41.5	63.3	82.5	113.5	146.5	174.5	222.1
2001	7.7	16.7	27.8	44.1	64.1	85.2	120.3	140.7	182.3	217.1
2000	10.0	16.8	28.8	41.3	64.8	85.6	117.7	155.7	172.0	210.8
1999	10.1	18.3	29.0	48.3	73.5	85.4	128.3	157.3	193.0	242.6
Black, male										
2002	9.6	18.4	30.9	50.2	82.0	97.2	142.9	187.4	209.4	267.0
2001	8.2	19.0	35.9	57.1	80.9	110.0	151.5	178.9	219.1	283.7
2000	11.8	20.8	33.1	51.4	89.0	102.0	149.1	211.7	208.2	253.7
1999	11.7	23.2	36.5	61.9	94.0	109.8	163.8	191.4	233.7	292.9
Black, female										
2002	6.5	13.5	20.5	34.5	48.6	71.6	93.9	122.1	156.4	205.0
2001	7.3	14.8	20.9	33.5	50.9	66.8	99.5	117.9	163.5	192.4
2000	8.3	13.4	25.1	33.1	45.8	73.5	96.8	121.9	153.7	194.9
1999	8.8	14.0	22.7	37.2	57.4	67.3	104.6	136.7	172.9	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All races 3/, both sexes										
2002	1.0	1.5	2.1	3.7	5.4	9.7	16.3	28.8	46.3	78.7
2001	0.9	1.5	2.3	3.9	6.3	10.5	17.9	29.5	47.8	82.7
2000	0.9	1.5	2.3	3.8	6.6	10.6	17.4	28.9	50.6	84.9
1999	0.9	1.4	2.3	3.5	6.3	10.8	18.6	29.3	47.5	82.9
All races 3/, male										
2002	1.0	1.6	2.1	4.5	6.3	12.2	20.0	37.3	59.1	101.1
2001	1.0	1.7	2.4	4.1	7.5	12.8	22.5	39.2	62.4	107.6
2000	0.9	1.6	2.6	4.1	7.5	13.3	21.2	36.8	65.3	108.9
1999	1.1	1.5	2.7	4.0	7.9	13.4	23.9	37.2	61.0	109.2
All races 3/, female										
2002	0.9	1.5	2.0	3.0	4.6	7.5	13.4	22.8	38.5	68.9
2001	0.8	1.2	2.1	3.7	5.2	8.5	14.3	22.7	39.1	72.3
2000	1.0	1.3	1.9	3.5	5.7	8.2	14.4	23.4	41.9	75.2
1999	0.8	1.3	1.9	3.0	4.8	8.5	14.5	23.8	39.7	72.3
White, both sexes										
2002	0.9	1.6	2.0	3.7	5.6	9.9	16.7	29.7	47.2	80.7
2001	0.9	1.5	2.3	3.9	6.3	10.7	18.1	30.4	48.9	83.9
2000	0.9	1.3	2.2	3.9	6.6	10.7	17.3	29.5	52.2	87.3
1999	0.9	1.3	2.2	3.5	6.2	10.7	19.0	29.7	48.7	85.3
White, male										
2002	1.1	1.7	2.1	4.4	6.4	12.4	20.7	38.4	60.1	104.2
2001	1.0	1.7	2.5	4.1	7.4	13.2	23.0	40.4	63.7	109.9
2000	0.9	1.5	2.6	4.3	7.3	13.4	21.0	37.6	68.0	112.6
1999	1.1	1.4	2.6	3.9	8.0	13.3	24.6	37.7	62.9	114.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
White, female										
2002	0.8	1.4	1.8	3.0	4.8	7.6	13.5	23.4	39.2	70.5
2001	0.8	1.2	2.0	3.8	5.2	8.6	14.2	23.3	40.0	73.0
2000	1.0	1.2	1.9	3.5	6.0	8.4	14.3	23.8	42.9	77.1
1999	0.7	1.2	1.8	3.0	4.6	8.3	14.7	24.0	40.4	73.7
Black, both sexes										
2002	1.7	2.0	3.0	5.2	5.8	10.6	15.0	24.4	41.9	63.6
2001	1.2	1.4	2.7	4.5	7.6	10.9	18.6	24.5	42.5	73.4
2000	1.4	2.4	3.1	4.2	7.1	10.6	20.3	26.7	37.8	64.1
1999	1.3	2.2	3.2	5.0	7.8	13.7	18.1	29.5	38.7	61.9
Black, male										
2002	*	1.7	2.8	7.4	6.5	12.7	16.0	29.9	53.5	79.3
2001	*	*	2.4	4.5	9.2	13.0	22.9	31.6	58.6	88.5
2000	*	2.2	3.6	4.4	10.1	15.6	23.7	34.5	38.8	76.1
1999	*	*	3.3	6.1	8.3	16.9	21.7	36.1	42.7	63.3
Black, female										
2002	2.1	2.2	3.3	3.3	5.3	9.1	14.3	21.1	35.8	57.5
2001	*	*	3.0	4.5	6.4	9.4	15.7	20.2	34.3	67.8
2000	1.3	2.5	2.7	4.0	4.7	6.9	18.0	21.9	37.4	59.7
1999	1.4	2.5	3.0	4.0	7.5	11.4	15.8	25.5	36.7	61.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
All races 3/, both sexes										
2002	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8
All races 3/, male										
2002	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4
All races 3/, female										
2002	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0
White, both sexes										
2002	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
1999	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0
White, male										
2002	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
1999	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
White, female										
2002	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
1999	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9
Black, both sexes										
2002	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
1999	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9
Black, male										
2002	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
1999	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5
Black, female										
2002	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4
1999	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2002	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2
All races 3/, male										
2002	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7
All races 3/, female										
2002	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8
White, both sexes										
2002	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
1999	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9
White, male										
2002	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
1999	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
White, female										
2002	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
1999	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1
Black, both sexes										
2002	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
1999	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0
Black, male										
2002	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
1999	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2
Black, female										
2002	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
1999	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional deficiencies (E40-E64)										
All races 3/, both sexes										
2002	0.1	0.2	0.3	0.5	0.8	1.6	2.7	6.1	12.8	41.5
2001	0.1	0.2	0.3	0.5	0.8	1.4	2.8	5.9	12.8	42.9
2000	0.1	0.2	0.3	0.5	1.0	2.0	3.0	6.8	15.8	50.0
1999	0.2	0.2	0.4	0.6	0.9	1.6	3.2	6.4	15.9	53.9
All races 3/, male										
2002	*	0.2	0.3	0.6	0.9	1.6	3.0	6.3	12.0	37.0
2001	*	0.3	0.3	0.5	0.9	1.4	2.9	5.6	13.2	35.2
2000	*	0.3	0.2	0.7	1.0	2.0	3.4	6.8	14.4	43.7
1999	0.2	0.2	0.4	0.7	1.1	1.6	3.8	6.8	16.0	48.4
All races 3/, female										
2002	*	*	0.4	0.4	0.7	1.6	2.4	6.0	13.3	43.5
2001	*	0.2	0.4	0.4	0.7	1.3	2.7	6.1	12.6	46.1
2000	*	0.2	0.4	0.4	0.9	2.0	2.7	6.8	16.6	52.6
1999	0.2	0.2	0.4	0.5	0.7	1.5	2.6	6.2	15.8	56.2
White, both sexes										
2002	0.1	0.2	0.3	0.4	0.7	1.4	2.5	5.9	12.3	40.2
2001	*	0.2	0.3	0.4	0.8	1.2	2.6	5.5	12.0	42.2
2000	*	0.2	0.2	0.5	0.8	1.9	2.7	6.4	15.3	48.8
1999	0.2	0.2	0.3	0.4	0.8	1.4	2.7	5.9	14.8	52.8
White, male										
2002	*	*	0.3	0.4	0.7	1.4	2.8	6.1	11.0	36.3
2001	*	*	*	0.4	0.7	1.1	2.6	5.3	12.2	33.8
2000	*	0.2	*	0.5	0.8	1.8	2.9	6.2	13.9	42.8
1999	*	*	0.3	0.4	1.0	1.3	3.0	6.1	14.7	46.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional deficiencies (E40-E64)										
White, female										
2002	*	*	0.4	0.4	0.7	1.3	2.2	5.7	13.1	41.9
2001	*	0.2	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.7
2000	*	*	0.3	0.4	0.9	2.0	2.5	6.5	16.2	51.3
1999	*	*	0.3	0.4	0.6	1.4	2.4	5.9	14.9	55.4
Black, both sexes										
2002	*	*	*	1.5	2.0	3.5	5.1	10.5	20.7	60.6
2001	*	*	1.0	*	*	3.0	4.4	10.8	23.8	54.7
2000	*	*	*	*	2.1	3.2	6.5	12.1	22.6	66.9
1999	*	*	*	2.0	2.0	3.2	8.3	12.3	30.9	70.8
Black, male										
2002	*	*	*	*	*	*	*	9.6	27.5	47.2
2001	*	*	*	*	*	*	6.7	11.6	24.9	57.9
2000	*	*	*	*	*	*	9.7	15.1	22.8	63.4
1999	*	*	*	*	*	*	12.9	16.1	34.9	75.0
Black, female										
2002	*	*	*	*	*	*	4.7	11.0	17.1	65.7
2001	*	*	*	*	*	*	*	10.3	23.3	53.5
2000	*	*	*	*	*	*	4.4	10.3	22.5	68.3
1999	*	*	*	*	*	*	5.2	10.0	29.0	69.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
All races 3/, both sexes										
2002	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0
All races 3/, male										
2002	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000	*	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4
All races 3/, female										
2002	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000	*	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3
White, both sexes										
2002	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	45.2
1999	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9
White, male										
2002	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000	*	*	*	0.5	0.7	1.6	2.8	5.9	12.5	39.7
1999	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
White, female										
2002	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000	*	*	0.3	*	0.9	1.9	2.1	6.2	15.1	47.4
1999	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6
Black, both sexes										
2002	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000	*	*	*	*	1.9	3.1	6.5	11.2	20.9	63.8
1999	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0
Black, male										
2002	*	*	*	*	*	*	*	9.2	25.2	46.1
2001	*	*	*	*	*	*	*	11.2	24.1	55.6
2000	*	*	*	*	*	*	9.7	14.2	22.8	61.1
1999	*	*	*	*	*	*	12.2	13.3	33.1	73.8
Black, female										
2002	*	*	*	*	*	*	*	10.7	17.1	62.0
2001	*	*	*	*	*	*	*	9.4	21.3	50.1
2000	*	*	*	*	*	*	4.4	9.4	20.0	64.8
1999	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other nutritional deficiencies (E50-E64)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	0.4	0.6	3.1
2001	*	*	*	*	*	*	*	0.4	0.8	2.7
2000	*	*	*	*	*	*	0.3	0.4	1.2	3.7
1999	*	*	*	*	*	*	*	0.4	1.1	3.9
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	1.9
2001	*	*	*	*	*	*	*	*	*	1.6
2000	*	*	*	*	*	*	*	*	1.3	3.0
1999	*	*	*	*	*	*	*	*	1.3	3.9
All races 3/, female										
2002	*	*	*	*	*	*	*	*	0.6	3.6
2001	*	*	*	*	*	*	*	*	1.0	3.1
2000	*	*	*	*	*	*	*	*	1.1	4.0
1999	*	*	*	*	*	*	*	*	1.1	3.9
White, both sexes										
2002	*	*	*	*	*	*	*	0.5	0.6	3.1
2001	*	*	*	*	*	*	*	0.4	0.8	2.7
2000	*	*	*	*	*	*	0.3	0.3	1.2	3.7
1999	*	*	*	*	*	*	*	0.4	1.1	3.9
White, male										
2002	*	*	*	*	*	*	*	*	*	2.0
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	1.4	3.2
1999	*	*	*	*	*	*	*	*	1.3	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other nutritional deficiencies (E50-E64)										
White, female										
2002	*	*	*	*	*	*	*	*	0.7	3.6
2001	*	*	*	*	*	*	*	*	0.9	3.1
2000	*	*	*	*	*	*	*	*	1.0	3.9
1999	*	*	*	*	*	*	*	*	1.0	3.8
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
All races 3/, both sexes										
2002	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5
All races 3/, male										
2002	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7
All races 3/, female										
2002	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4
White, both sexes										
2002	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3
White, male										
2002	*	0.3	*	0.3	*	0.6	0.8	*	*	*
2001	*	0.2	0.3	0.4	0.5	*	0.7	0.8	*	*
2000	*	0.3	0.3	0.4	0.5	*	0.8	1.2	*	*
1999	*	0.3	*	0.4	*	0.5	0.7	*	1.3	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
White, female										
2002	*	*	*	*	*	*	0.5	0.5	*	1.2
2001	*	*	*	*	0.4	*	*	*	0.8	0.8
2000	*	*	*	*	*	*	*	0.6	1.0	1.0
1999	*	*	*	*	0.5	*	0.6	0.8	1.0	1.3
Black, both sexes										
2002	*	0.8	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	1.0	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
All races 3/, both sexes										
2002	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4
All races 3/, male										
2002	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6
All races 3/, female										
2002	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7
White, both sexes										
2002	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
1999	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2
White, male										
2002	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
1999	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
White, female										
2002	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
1999	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8
Black, both sexes										
2002	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
1999	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0
Black, male										
2002	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000	*	*	*	*	*	*	13.7	37.8	65.8	86.5
1999	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7
Black, female										
2002	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000	*	*	*	*	*	*	5.5	10.8	23.4	38.6
1999	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
All races 3/, both sexes										
2002	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3
All races 3/, male										
2002	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7
All races 3/, female										
2002	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0
White, both sexes										
2002	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
1999	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9
White, male										
2002	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
1999	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
White, female										
2002	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
1999	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5
Black, both sexes										
2002	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
1999	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4
Black, male										
2002	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
1999	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5
Black, female										
2002	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
1999	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Major cardiovascular diseases (I00-I78)										
All races 3/, both sexes										
2002	49.7	87.9	144.2	233.5	374.0	602.6	993.2	1,700.2	3,034.4	7,389.6
2001	48.8	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000	49.6	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
1999	50.0	88.3	148.7	254.8	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1
All races 3/, male										
2002	67.9	123.9	206.1	326.7	512.6	802.0	1,287.4	2,140.7	3,633.4	7,638.8
2001	66.5	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000	68.0	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
1999	68.9	125.7	212.5	358.4	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2
All races 3/, female										
2002	31.6	52.9	85.0	146.0	248.2	430.4	758.0	1,389.0	2,670.1	7,281.5
2001	31.5	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000	31.5	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
1999	31.4	52.1	87.5	158.0	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5
White, both sexes										
2002	43.6	76.6	127.4	211.8	344.7	567.9	957.1	1,663.0	3,023.3	7,482.7
2001	41.6	74.2	127.7	212.4	355.6	580.3	980.2	1,721.1	3,122.8	7,770.6
2000	42.6	76.5	130.6	225.6	376.3	612.7	1,027.2	1,783.3	3,243.6	8,133.9
1999	42.4	76.6	132.4	230.2	385.3	644.7	1,068.3	1,848.2	3,345.3	8,299.2
White, male										
2002	62.0	111.9	186.7	301.4	479.1	765.2	1,253.1	2,112.9	3,642.0	7,766.6
2001	59.1	109.0	185.6	301.7	493.8	781.8	1,286.2	2,192.9	3,777.7	8,060.7
2000	60.7	114.5	190.5	323.5	523.7	822.7	1,366.4	2,273.2	3,963.6	8,585.0
1999	60.8	113.0	193.5	329.9	533.4	874.1	1,420.2	2,361.9	4,089.2	8,717.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Major cardiovascular diseases (I00-I78)										
White, female										
2002	25.2	41.6	69.3	126.1	220.5	394.1	715.8	1,341.0	2,644.0	7,359.3
2001	24.0	39.8	71.0	127.1	228.1	403.3	732.3	1,386.9	2,726.9	7,649.2
2000	24.4	39.0	72.0	132.4	240.4	429.2	755.0	1,438.7	2,815.1	7,950.6
1999	24.0	40.6	72.8	135.3	248.8	444.5	788.7	1,487.4	2,906.2	8,131.1
Black, both sexes										
2002	99.1	184.1	302.3	468.4	697.7	1,001.3	1,510.2	2,343.9	3,555.3	7,081.7
2001	105.7	181.9	307.2	471.1	713.7	1,057.5	1,548.8	2,422.8	3,623.2	7,316.7
2000	104.9	192.0	303.5	489.3	736.9	1,061.9	1,623.2	2,496.2	3,719.8	7,518.9
1999	109.8	188.5	311.0	509.4	743.3	1,093.2	1,704.6	2,553.3	3,787.7	7,582.5
Black, male										
2002	122.7	237.2	406.5	633.2	934.4	1,292.7	1,908.8	2,803.5	4,093.1	6,929.0
2001	129.8	237.2	405.7	627.8	934.4	1,342.7	1,988.0	2,943.9	4,284.2	7,363.9
2000	131.7	247.2	407.0	670.8	955.1	1,328.0	2,040.7	2,999.5	4,397.8	7,687.4
1999	138.3	246.5	423.5	686.4	993.1	1,400.0	2,146.5	3,046.5	4,407.7	7,921.1
Black, female										
2002	78.2	138.0	213.9	333.0	512.5	785.0	1,242.7	2,069.3	3,275.3	7,140.2
2001	84.3	134.0	223.4	342.6	540.2	846.9	1,256.2	2,111.8	3,286.5	7,299.2
2000	81.2	144.0	215.2	340.5	564.6	865.5	1,346.0	2,192.8	3,378.3	7,456.2
1999	84.7	138.5	215.6	364.7	547.4	865.8	1,411.4	2,256.2	3,480.4	7,456.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)										
All races 3/, both sexes										
2002	40.2	71.3	119.2	192.8	304.4	479.6	766.1	1,278.2	2,234.3	5,446.8
2001	39.1	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000	39.4	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
1999	40.1	71.7	123.5	211.0	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0
All races 3/, male										
2002	57.7	104.7	177.0	278.2	429.1	654.3	1,025.8	1,667.3	2,786.9	5,823.5
2001	55.8	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000	57.0	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
1999	58.1	107.0	182.9	306.5	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6
All races 3/, female										
2002	22.9	38.8	63.9	112.5	191.2	328.8	558.5	1,003.3	1,898.3	5,283.3
2001	22.6	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000	22.2	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
1999	22.4	37.5	66.6	121.7	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1
White, both sexes										
2002	36.3	64.0	108.1	177.9	284.5	457.0	744.1	1,255.7	2,234.0	5,529.1
2001	34.2	61.8	108.6	178.5	294.7	468.4	765.0	1,297.7	2,308.3	5,752.1
2000	34.9	63.7	109.9	190.0	309.5	492.6	801.0	1,347.0	2,393.0	6,020.8
1999	34.9	64.1	112.7	193.7	321.3	524.0	839.7	1,403.0	2,484.1	6,152.7
White, male										
2002	53.9	96.8	164.4	261.0	406.6	631.3	1,006.4	1,654.3	2,805.0	5,939.8
2001	50.9	94.4	163.4	261.7	420.5	647.2	1,031.2	1,716.7	2,896.4	6,157.2
2000	52.3	99.4	166.1	280.7	442.0	680.1	1,096.8	1,771.0	3,030.0	6,560.8
1999	52.4	98.8	170.3	286.1	454.8	730.0	1,148.4	1,851.7	3,144.4	6,669.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)										
White, female										
2002	18.6	31.4	53.1	98.3	171.5	303.6	530.3	970.4	1,884.1	5,350.6
2001	17.5	29.4	54.9	99.1	178.6	311.5	549.5	1,001.0	1,952.9	5,582.5
2000	17.5	28.3	54.9	103.7	187.4	328.6	563.7	1,048.7	2,014.1	5,801.4
1999	17.4	29.7	56.4	105.8	198.3	344.3	594.5	1,088.0	2,094.3	5,945.3
Black, both sexes										
2002	74.8	138.7	232.1	365.8	542.7	764.2	1,126.8	1,715.1	2,557.1	5,115.2
2001	79.5	137.2	232.2	369.7	546.5	809.7	1,160.0	1,780.6	2,619.7	5,275.7
2000	77.5	142.2	231.2	380.7	571.0	813.9	1,230.4	1,836.2	2,673.7	5,389.5
1999	82.9	143.0	240.2	400.6	579.9	852.1	1,289.1	1,878.6	2,749.2	5,487.4
Black, male										
2002	96.7	186.2	319.6	503.6	740.6	1,001.5	1,448.8	2,095.3	3,039.3	5,125.7
2001	101.9	184.4	314.7	503.1	731.0	1,039.8	1,518.8	2,214.9	3,181.5	5,459.9
2000	101.3	190.8	319.1	533.0	754.7	1,033.7	1,570.9	2,245.4	3,226.1	5,633.5
1999	109.3	193.3	335.5	556.7	789.2	1,111.8	1,652.9	2,277.2	3,259.4	5,858.8
Black, female										
2002	55.5	97.5	157.9	252.6	388.0	588.1	910.7	1,488.0	2,306.1	5,111.2
2001	59.7	96.3	162.0	260.3	401.4	639.8	921.0	1,521.3	2,333.5	5,207.3
2000	56.5	100.1	156.3	255.9	426.1	651.7	1,004.3	1,589.5	2,395.4	5,298.7
1999	59.6	99.6	159.4	273.1	415.8	659.7	1,047.8	1,638.6	2,496.4	5,348.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
All races 3/, both sexes										
2002	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0
All races 3/, male										
2002	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3
All races 3/, female										
2002	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1
White, both sexes										
2002	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
1999	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9
White, male										
2002	*	0.3	0.5	0.9	1.5	1.8	4.4	5.9	8.8	12.6
2001	0.3	0.4	0.4	1.0	1.2	2.3	3.3	6.3	8.0	13.3
2000	0.3	0.5	0.7	1.1	1.9	2.3	3.5	6.1	8.4	12.7
1999	0.2	0.4	0.6	1.0	1.3	2.6	4.9	5.5	9.4	11.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
White, female										
2002	0.2	0.5	0.9	1.4	2.5	3.7	5.9	10.6	14.2	21.6
2001	0.3	0.4	0.7	1.1	2.3	3.9	6.4	9.9	14.7	21.0
2000	0.4	0.6	0.8	1.5	2.4	4.6	7.3	10.4	14.7	19.6
1999	0.3	0.6	0.9	1.8	2.7	4.6	6.9	11.0	15.4	22.0
Black, both sexes										
2002	*	0.8	1.1	2.0	2.9	3.2	*	4.8	*	*
2001	0.9	1.1	1.6	2.2	2.0	3.3	5.1	4.9	*	6.8
2000	0.9	1.0	1.1	1.9	2.2	3.4	3.6	3.7	*	8.4
1999	0.7	0.9	1.6	2.2	2.1	2.8	4.2	4.3	*	8.3
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	3.6	*	*	5.5	*	*
2001	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000	*	*	*	*	*	4.4	*	*	*	*
1999	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2002	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9
All races 3/, male										
2002	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6
All races 3/, female										
2002	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5
White, both sexes										
2002	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
1999	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3
White, male										
2002	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
1999	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
White, female										
2002	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
1999	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0
Black, both sexes										
2002	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
1999	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1
Black, male										
2002	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
1999	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6
Black, female										
2002	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
1999	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
All races 3/, both sexes										
2002	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1
All races 3/, male										
2002	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3
All races 3/, female										
2002	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2
White, both sexes										
2002	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
1999	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1
White, male										
2002	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
1999	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
White, female										
2002	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
1999	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0
Black, both sexes										
2002	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
1999	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4
Black, male										
2002	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
1999	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8
Black, female										
2002	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5
1999	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2002	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
All races 3/, male										
2002	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
All races 3/, female										
2002	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8
White, both sexes										
2002	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
1999	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2
White, male										
2002	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
1999	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
White, female										
2002	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
1999	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3
Black, both sexes										
2002	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
1999	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8
Black, male										
2002	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
1999	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1
Black, female										
2002	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
1999	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2002	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3
All races 3/, male										
2002	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1
All races 3/, female										
2002	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5
White, both sexes										
2002	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
1999	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2
White, male										
2002	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
1999	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
White, female										
2002	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
1999	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4
Black, both sexes										
2002	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
1999	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8
Black, male										
2002	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
1999	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7
Black, female										
2002	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5
1999	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes										
2002	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9
All races 3/, male										
2002	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3
All races 3/, female										
2002	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7
White, both sexes										
2002	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
1999	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4
White, male										
2002	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
1999	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
White, female										
2002	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
1999	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1
Black, both sexes										
2002	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
1999	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2
Black, male										
2002	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
1999	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*
Black, female										
2002	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001	*	*	*	*	*	*	5.2	*	11.9	24.0
2000	*	*	*	*	3.3	*	*	8.0	10.6	22.3
1999	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2002	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0
All races 3/, male										
2002	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1
All races 3/, female										
2002	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7
White, both sexes										
2002	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
1999	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6
White, male										
2002	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
1999	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
White, female										
2002	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
1999	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7
Black, both sexes										
2002	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
1999	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8
Black, male										
2002	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
1999	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5
Black, female										
2002	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2
1999	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2002	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4
All races 3/, male										
2002	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6
All races 3/, female										
2002	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2
White, both sexes										
2002	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
1999	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5
White, male										
2002	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
1999	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
White, female										
2002	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
1999	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0
Black, both sexes										
2002	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
1999	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6
Black, male										
2002	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
1999	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9
Black, female										
2002	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
1999	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2002	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6
All races 3/, male										
2002	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4
All races 3/, female										
2002	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4
White, both sexes										
2002	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
1999	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1
White, male										
2002	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
1999	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, female										
2002	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
1999	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8
Black, both sexes										
2002	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
1999	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2
Black, male										
2002	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
1999	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5
Black, female										
2002	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9
1999	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2002	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000	12.3	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9
All races 3/, male										
2002	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000	15.4	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9
All races 3/, female										
2002	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000	9.2	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5
White, both sexes										
2002	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000	10.2	14.6	22.1	35.9	59.0	97.3	160.6	284.5	545.7	1,541.6
1999	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3
White, male										
2002	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000	13.0	19.7	28.6	46.9	74.8	119.7	199.2	339.4	630.8	1,580.5
1999	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
White, female										
2002	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000	7.4	9.6	15.8	25.4	44.3	77.7	129.5	245.9	495.2	1,525.8
1999	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9
Black, both sexes										
2002	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000	28.5	43.3	62.0	86.2	122.8	181.5	266.8	406.9	590.7	1,306.0
1999	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8
Black, male										
2002	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000	34.8	54.3	82.1	112.1	153.6	224.8	318.9	480.5	696.3	1,402.3
1999	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1
Black, female										
2002	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000	22.9	33.9	44.8	64.9	98.4	149.6	232.2	362.6	537.5	1,270.2
1999	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
All races 3/, both sexes										
2002	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000	0.3	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9
All races 3/, male										
2002	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000	0.4	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7
All races 3/, female										
2002	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1
White, both sexes										
2002	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0
White, male										
2002	0.3	0.4	0.6	0.4	0.9	1.3	1.7	2.7	3.9	3.6
2001	0.3	0.5	0.5	0.5	0.9	0.9	1.8	3.0	4.3	3.9
2000	0.4	0.5	0.6	0.7	0.9	1.2	1.6	3.2	4.0	4.2
1999	0.3	0.4	0.5	0.8	0.7	1.2	1.6	3.0	4.0	5.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
White, female										
2002	*	*	*	0.3	0.7	0.7	1.2	1.8	2.3	2.3
2001	*	*	*	0.4	0.4	0.9	1.3	1.3	2.2	2.1
2000	*	*	0.3	0.4	0.5	1.0	1.0	1.6	2.0	2.1
1999	*	*	*	*	0.6	0.8	1.3	1.6	2.0	2.1
Black, both sexes										
2002	*	1.1	1.3	1.7	*	2.4	*	*	*	*
2001	0.9	1.1	1.5	*	2.0	2.3	3.0	4.7	*	*
2000	0.7	1.3	1.4	1.9	*	*	*	*	*	*
1999	0.8	1.1	1.3	2.3	*	2.3	2.8	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	1.8	*	*	*	*	*	*	*	*
2000	*	1.8	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All races 3/, both sexes										
2002	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0
All races 3/, male										
2002	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5
All races 3/, female										
2002	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9
White, both sexes										
2002	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0
White, male										
2002	0.3	0.3	0.3	0.4	0.6	0.6	1.1	1.1	1.2	2.5
2001	*	0.2	0.3	0.4	0.6	0.5	0.9	1.3	1.1	2.0
2000	0.2	*	*	0.5	*	*	1.1	1.2	1.9	2.1
1999	0.3	0.3	*	0.6	0.6	0.7	0.9	1.3	1.5	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
White, female										
2002	*	*	*	0.4	*	0.5	0.9	1.0	1.4	2.0
2001	*	0.3	*	0.3	*	*	0.9	1.2	1.4	1.9
2000	*	*	*	*	*	0.6	0.7	0.9	1.1	1.6
1999	*	0.3	*	*	0.5	0.7	0.8	1.1	1.4	1.8
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
All races 3/, both sexes										
2002	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9
All races 3/, male										
2002	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0
All races 3/, female										
2002	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9
White, both sexes										
2002	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
1999	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2
White, male										
2002	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
1999	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
White, female										
2002	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
1999	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0
Black, both sexes										
2002	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
1999	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9
Black, male										
2002	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
1999	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5
Black, female										
2002	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9
1999	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All races 3/, both sexes										
2002	11.2	15.7	21.7	32.5	46.9	73.3	116.3	194.4	338.2	793.7
2001	10.5	15.4	21.9	32.3	47.8	75.3	116.6	191.2	333.6	796.5
2000	10.9	15.5	22.1	33.1	50.4	78.0	118.4	193.4	338.9	820.0
1999	11.1	15.7	22.3	34.1	53.5	79.9	124.4	197.7	347.6	826.1
All races 3/, male										
2002	14.2	20.6	28.1	41.1	59.4	89.7	143.8	233.1	389.4	821.3
2001	13.4	19.9	27.7	40.3	61.1	91.7	143.7	232.4	390.0	821.3
2000	13.6	20.2	28.9	43.4	64.2	96.8	147.6	232.6	391.5	870.7
1999	13.9	20.2	28.7	44.5	68.7	103.5	155.5	238.5	411.1	874.7
All races 3/, female										
2002	8.2	11.0	15.6	24.4	35.6	59.2	94.3	167.1	307.1	781.7
2001	7.6	11.1	16.3	24.8	35.7	61.2	95.0	162.4	299.7	786.1
2000	8.2	11.0	15.6	23.4	37.8	61.9	95.4	166.0	307.9	799.3
1999	8.3	11.2	16.2	24.3	39.8	59.7	100.1	169.4	310.5	806.5
White, both sexes										
2002	9.7	13.3	18.8	28.9	42.7	69.1	111.9	191.2	339.3	806.1
2001	8.7	12.9	18.9	28.3	44.0	70.3	112.3	186.3	333.4	809.9
2000	9.1	13.0	18.9	29.6	46.2	73.1	113.5	188.4	339.9	833.4
1999	9.2	12.9	19.4	29.8	48.2	74.9	118.6	192.7	346.9	839.0
White, male										
2002	12.6	17.7	24.8	36.9	54.7	85.0	138.8	229.4	391.0	834.2
2001	11.5	17.0	24.7	35.4	56.5	86.9	138.2	226.4	388.3	831.0
2000	11.5	17.5	24.8	38.9	59.5	90.9	142.2	227.3	393.2	884.2
1999	11.9	17.3	25.1	39.0	62.4	97.5	148.4	232.5	409.5	883.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
White, female										
2002	6.7	9.0	13.0	21.4	31.5	55.1	90.0	163.8	307.6	793.9
2001	5.8	8.8	13.4	21.5	32.5	55.8	91.3	158.0	300.1	801.1
2000	6.7	8.6	13.2	20.7	33.9	57.6	90.4	161.1	308.1	812.7
1999	6.5	8.6	13.7	21.0	35.2	55.1	94.9	164.7	310.0	821.0
Black, both sexes										
2002	23.6	36.5	49.5	70.6	94.5	127.7	180.3	263.4	376.5	738.5
2001	25.1	36.9	50.0	73.6	91.5	135.2	179.9	276.1	383.5	730.2
2000	25.2	37.3	52.5	69.7	96.5	137.5	191.0	282.9	377.8	742.0
1999	26.3	38.5	51.8	77.4	109.1	140.1	206.0	283.3	400.8	746.1
Black, male										
2002	28.2	47.7	62.8	90.7	117.2	157.1	226.3	314.2	434.1	753.7
2001	30.0	45.9	60.5	95.7	121.2	158.8	227.2	347.1	479.1	805.6
2000	30.9	46.3	71.0	95.0	122.0	174.3	233.4	340.7	449.3	816.5
1999	31.8	47.7	67.7	103.6	142.8	183.3	265.3	341.0	503.1	849.5
Black, female										
2002	19.5	26.8	38.3	54.1	76.8	105.8	149.4	233.0	346.5	732.8
2001	20.8	29.1	41.0	55.4	68.2	117.7	148.4	233.8	334.8	702.2
2000	20.3	29.5	36.8	49.0	76.5	110.4	162.9	248.1	341.8	714.3
1999	21.5	30.6	38.3	56.1	82.7	108.1	166.8	248.6	350.0	707.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
All races 3/, both sexes										
2002	1.0	1.8	2.9	4.5	7.3	11.8	20.7	34.9	66.7	180.4
2001	1.0	1.9	3.0	4.4	7.5	12.2	19.0	35.5	65.2	171.9
2000	1.1	1.8	3.0	4.4	7.8	11.5	19.1	33.8	63.0	162.9
1999	0.9	1.7	2.8	4.1	7.3	12.0	18.6	32.7	59.9	152.1
All races 3/, male										
2002	1.3	2.3	3.5	5.5	8.2	12.7	22.2	35.9	64.8	153.4
2001	1.3	2.3	3.7	5.2	8.1	13.7	20.0	36.0	61.3	150.8
2000	1.4	2.3	3.8	5.4	9.1	11.5	20.6	34.5	62.9	142.4
1999	1.1	2.0	3.5	5.0	8.2	13.4	20.1	33.1	59.4	132.4
All races 3/, female										
2002	0.7	1.3	2.3	3.7	6.5	11.0	19.5	34.2	67.9	192.1
2001	0.7	1.4	2.3	3.8	6.9	11.0	18.2	35.2	67.5	180.8
2000	0.8	1.3	2.2	3.5	6.7	11.5	17.9	33.4	63.0	171.3
1999	0.7	1.3	2.1	3.3	6.5	10.8	17.5	32.4	60.2	160.0
White, both sexes										
2002	0.6	1.0	1.9	3.0	5.4	9.0	16.8	29.8	60.4	170.4
2001	0.5	1.1	1.8	2.9	5.3	9.2	15.3	30.9	58.6	163.3
2000	0.6	1.0	1.8	3.0	5.5	8.5	15.3	28.9	57.4	153.2
1999	0.5	1.0	1.6	2.7	5.0	9.1	15.2	27.6	53.3	142.0
White, male										
2002	0.8	1.4	2.4	3.8	6.3	9.8	18.3	30.8	57.7	145.2
2001	0.7	1.4	2.2	3.5	5.9	10.3	16.4	31.4	54.5	142.2
2000	0.9	1.4	2.5	3.8	6.6	8.8	16.8	29.9	57.5	132.5
1999	0.7	1.3	2.1	3.5	5.8	10.4	16.6	28.1	52.5	121.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
White, female										
2002	0.4	0.7	1.4	2.3	4.5	8.3	15.6	29.0	62.0	181.3
2001	0.3	0.7	1.4	2.4	4.8	8.2	14.5	30.5	61.1	172.2
2000	0.4	0.6	1.1	2.2	4.5	8.3	14.1	28.2	57.4	161.6
1999	0.3	0.8	1.2	1.9	4.3	7.9	14.0	27.2	53.8	150.5
Black, both sexes										
2002	3.8	7.4	10.9	18.4	24.3	36.5	61.5	96.4	151.0	308.8
2001	4.4	7.8	12.5	18.0	27.0	40.4	56.3	89.8	148.2	288.0
2000	4.2	7.2	12.7	17.9	29.0	40.3	58.8	92.4	130.4	283.4
1999	4.0	6.3	12.8	17.0	27.0	38.0	57.2	95.0	138.7	269.4
Black, male										
2002	4.7	9.2	13.3	21.7	25.7	40.6	67.1	105.2	164.3	274.5
2001	5.5	10.1	16.3	21.7	30.3	48.8	61.9	95.2	158.1	275.7
2000	5.3	9.0	15.6	22.6	34.4	40.1	63.9	93.1	137.4	269.9
1999	4.6	7.8	16.6	19.5	31.0	43.1	62.1	98.8	155.2	266.0
Black, female										
2002	3.0	5.9	8.8	15.6	23.2	33.5	57.8	91.1	144.0	322.0
2001	3.4	5.8	9.3	15.0	24.5	34.2	52.5	86.5	143.1	292.6
2000	3.2	5.7	10.3	14.2	24.7	40.5	55.5	92.0	126.9	288.5
1999	3.5	5.0	9.5	15.0	24.0	34.2	54.0	92.7	130.5	270.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2002	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8
All races 3/, male										
2002	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8
All races 3/, female										
2002	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0
White, both sexes										
2002	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
1999	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6
White, male										
2002	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
1999	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
White, female										
2002	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
1999	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4
Black, both sexes										
2002	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
1999	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5
Black, male										
2002	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
1999	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3
Black, female										
2002	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
1999	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
All races 3/, both sexes										
2002	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0
All races 3/, male										
2002	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0
All races 3/, female										
2002	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2
White, both sexes										
2002	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
1999	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7
White, male										
2002	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
1999	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
White, female										
2002	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
1999	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3
Black, both sexes										
2002	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
1999	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3
Black, male										
2002	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
1999	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0
Black, female										
2002	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8
1999	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of circulatory system (I71-I78)										
All races 3/, both sexes										
2002	1.3	2.2	3.1	6.2	11.7	22.3	36.5	57.8	83.8	148.4
2001	1.3	2.0	3.3	6.0	12.1	22.3	36.6	60.6	88.1	158.2
2000	1.3	2.1	3.5	6.4	12.7	24.5	40.6	62.9	92.2	163.2
1999	1.4	2.1	3.3	6.7	12.5	23.5	40.1	63.5	94.3	165.2
All races 3/, male										
2002	1.8	3.2	4.4	8.6	16.3	31.7	49.2	78.3	113.7	187.4
2001	1.8	2.8	4.4	8.2	16.8	30.8	51.2	81.0	122.5	197.3
2000	1.7	2.8	4.9	9.3	18.0	34.6	57.6	88.6	126.6	214.9
1999	1.6	3.0	4.4	9.4	18.2	33.5	56.7	89.1	129.3	207.5
All races 3/, female										
2002	0.7	1.4	2.0	3.9	7.5	14.2	26.4	43.4	65.6	131.5
2001	0.9	1.2	2.2	3.9	7.8	15.0	25.1	46.3	67.4	141.8
2000	0.9	1.5	2.1	3.7	7.9	15.7	27.2	45.0	71.9	142.2
1999	1.1	1.3	2.1	4.1	7.3	15.0	27.2	45.7	73.7	148.2
White, both sexes										
2002	1.1	1.9	2.7	5.8	11.4	22.4	37.0	58.7	84.8	148.9
2001	1.1	1.8	2.9	5.6	11.7	22.6	36.8	60.6	88.4	158.2
2000	1.0	1.8	3.0	6.0	12.3	24.5	41.4	63.0	93.0	163.0
1999	1.2	1.9	3.0	6.3	12.6	23.7	40.8	64.1	94.8	165.1
White, male										
2002	1.6	2.7	3.8	7.9	16.0	32.5	49.8	80.4	116.1	190.9
2001	1.4	2.5	3.8	7.9	16.4	31.5	51.7	82.0	124.0	201.3
2000	1.3	2.5	4.4	8.9	17.8	35.5	59.1	89.4	129.7	219.6
1999	1.5	2.7	4.0	9.2	18.6	34.4	58.4	91.1	131.1	211.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of circulatory system (I71-I78)										
White, female										
2002	0.6	1.0	1.7	3.7	7.2	13.5	26.5	43.1	65.6	130.7
2001	0.7	1.1	2.0	3.4	7.4	14.7	24.7	45.5	67.0	140.1
2000	0.7	1.1	1.8	3.3	7.2	15.0	27.1	44.5	71.2	140.1
1999	0.9	1.0	1.9	3.6	7.1	14.5	26.9	45.1	73.4	146.6
Black, both sexes										
2002	2.8	5.3	6.7	11.2	16.5	25.7	37.8	58.4	83.5	156.8
2001	3.2	4.0	6.5	10.7	18.0	22.9	42.8	69.3	96.2	174.7
2000	3.2	4.5	6.8	11.2	18.3	27.3	39.9	68.9	92.3	177.7
1999	2.8	4.5	6.1	11.4	13.5	26.1	38.6	64.3	96.5	178.9
Black, male										
2002	4.1	6.7	9.8	16.5	23.0	30.7	48.7	65.7	100.1	166.2
2001	4.5	5.5	9.4	13.1	23.8	28.0	55.9	79.0	127.6	165.7
2000	4.2	5.5	10.0	15.1	22.6	32.8	51.8	90.7	104.5	177.6
1999	3.1	6.1	8.6	14.1	17.8	31.9	49.2	74.6	117.7	184.0
Black, female										
2002	1.7	4.0	4.0	6.8	11.5	22.0	30.5	54.1	74.8	153.2
2001	2.0	2.7	4.1	8.7	13.4	19.1	34.1	63.5	80.1	178.1
2000	2.3	3.7	4.1	8.0	14.9	23.2	32.0	55.8	86.2	177.7
1999	2.5	3.2	4.0	9.1	10.1	21.7	31.5	58.1	86.0	177.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes										
2002	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3
All races 3/, male										
2002	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3
All races 3/, female										
2002	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2
White, both sexes										
2002	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
1999	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2
White, male										
2002	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
1999	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
White, female										
2002	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
1999	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1
Black, both sexes										
2002	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
1999	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3
Black, male										
2002	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
1999	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2
Black, female										
2002	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7
1999	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes										
2002	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9
All races 3/, male										
2002	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2
All races 3/, female										
2002	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0
White, both sexes										
2002	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
1999	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9
White, male										
2002	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
1999	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
White, female										
2002	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
1999	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5
Black, both sexes										
2002	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
1999	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6
Black, male										
2002	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
1999	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8
Black, female										
2002	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
1999	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2002	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4
All races 3/, male										
2002	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8
All races 3/, female										
2002	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5
White, both sexes										
2002	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
1999	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2
White, male										
2002	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
1999	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
White, female										
2002	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
1999	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2
Black, both sexes										
2002	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
1999	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4
Black, male										
2002	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
1999	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*
Black, female										
2002	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0
1999	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)										
All races 3/, both sexes										
2002	2.6	4.2	5.5	8.5	14.8	25.3	50.9	106.5	227.3	696.6
2001	2.6	3.9	5.4	8.1	13.9	24.4	49.3	101.1	216.9	685.6
2000	3.0	3.9	5.6	8.9	15.6	26.6	52.6	106.7	240.8	744.1
1999	2.7	4.1	5.3	8.4	14.3	25.0	50.5	105.9	233.8	751.8
All races 3/, male										
2002	3.1	4.9	6.7	10.2	18.0	31.2	63.9	133.9	287.7	805.8
2001	3.0	4.9	6.7	10.0	17.0	30.3	61.6	130.7	274.7	808.4
2000	3.8	4.7	7.0	10.9	18.9	32.9	65.7	134.1	308.8	892.0
1999	3.1	5.2	6.8	9.7	16.8	31.7	64.2	133.5	301.7	894.7
All races 3/, female										
2002	2.2	3.4	4.3	6.8	11.8	20.1	40.6	87.2	190.6	649.2
2001	2.2	3.0	4.1	6.4	11.1	19.2	39.5	80.5	182.3	634.2
2000	2.2	3.1	4.3	7.0	12.5	21.2	42.2	87.6	200.7	683.9
1999	2.3	3.0	3.9	7.2	12.0	19.2	39.7	86.7	194.1	694.2
White, both sexes										
2002	2.3	3.8	5.0	7.7	13.8	24.5	49.9	105.4	228.0	709.1
2001	2.2	3.3	4.8	7.5	13.0	23.3	48.1	98.6	216.6	695.0
2000	2.5	3.4	4.9	8.1	14.3	24.5	50.3	105.6	240.7	758.3
1999	2.1	3.4	4.6	7.7	13.1	24.0	48.8	105.5	235.5	764.3
White, male										
2002	2.7	4.4	6.0	9.1	16.5	29.8	62.4	131.4	286.5	815.9
2001	2.5	4.2	5.7	9.0	15.4	28.3	59.3	125.5	272.3	814.4
2000	3.2	3.9	5.9	9.5	17.2	29.7	62.2	131.8	306.4	906.1
1999	2.3	4.2	5.7	8.6	15.1	30.1	60.9	132.2	301.9	905.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza and pneumonia (J10-J18)										
White, female										
2002	1.9	3.2	3.9	6.3	11.3	19.8	39.7	86.8	192.1	662.6
2001	1.8	2.5	4.0	6.0	10.7	19.0	39.0	79.6	182.9	645.0
2000	1.8	2.8	4.0	6.7	11.6	20.0	40.8	87.1	201.7	698.3
1999	1.9	2.6	3.6	6.7	11.2	18.7	39.1	86.8	196.4	707.5
Black, both sexes										
2002	5.2	7.5	10.1	17.2	26.3	35.5	68.9	123.3	231.6	575.2
2001	5.8	9.0	10.7	15.5	23.3	34.3	65.7	133.7	224.0	581.3
2000	6.1	8.3	12.2	17.8	27.2	46.0	77.5	124.4	245.4	600.9
1999	6.9	9.7	11.8	16.2	26.6	36.8	75.2	118.7	219.9	634.8
Black, male										
2002	6.2	9.9	12.9	23.7	36.0	48.1	89.8	173.2	321.7	689.4
2001	7.0	11.4	15.7	20.8	33.6	50.1	94.9	203.5	309.0	714.8
2000	8.3	11.3	17.2	25.9	35.2	64.8	107.6	172.5	343.9	755.4
1999	9.0	13.4	17.2	21.7	34.2	51.9	109.6	160.0	312.2	803.8
Black, female										
2002	4.3	5.5	7.7	12.0	18.7	26.1	54.9	93.6	184.6	531.5
2001	4.8	6.9	6.5	11.2	15.2	22.7	46.2	92.1	180.7	531.7
2000	4.2	5.8	7.9	11.2	20.9	32.1	57.5	95.4	195.7	543.4
1999	5.1	6.6	7.1	11.8	20.6	25.6	52.4	93.8	174.1	571.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001	*	*	*	*	*	*	*	*	0.7	2.2
2000	*	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2
All races 3/, male										
2002	*	*	*	*	*	*	*	0.9	2.6	8.9
2001	*	*	*	*	*	*	*	*	*	2.2
2000	*	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9
All races 3/, female										
2002	*	*	*	*	*	*	*	0.7	2.0	8.7
2001	*	*	*	*	*	*	*	*	0.7	2.2
2000	*	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7
White, both sexes										
2002	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001	*	*	*	*	*	*	*	*	0.7	2.3
2000	*	*	0.1	0.3	0.5	0.8	1.4	2.7	6.5	21.4
1999	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8
White, male										
2002	*	*	*	*	*	*	*	1.0	2.8	9.7
2001	*	*	*	*	*	*	*	*	*	2.3
2000	*	*	*	*	0.5	1.0	1.7	2.7	6.7	21.6
1999	*	*	*	*	0.5	0.6	1.2	2.6	7.5	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
White, female										
2002	*	*	*	*	*	*	*	0.8	2.2	9.3
2001	*	*	*	*	*	*	*	*	0.7	2.3
2000	*	*	*	*	0.6	0.7	1.2	2.7	6.3	21.3
1999	*	*	*	*	*	0.7	1.1	2.0	5.7	23.2
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	9.4
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
All races 3/, both sexes										
2002	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000	2.9	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6
All races 3/, male										
2002	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000	3.7	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7
All races 3/, female										
2002	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000	2.1	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5
White, both sexes										
2002	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000	2.4	3.2	4.8	7.8	13.8	23.7	48.9	102.9	234.3	737.0
1999	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6
White, male										
2002	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000	3.1	3.8	5.7	9.2	16.7	28.6	60.6	129.1	299.7	884.4
1999	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
White, female										
2002	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000	1.8	2.7	3.9	6.4	11.1	19.4	39.6	84.5	195.4	677.1
1999	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4
Black, both sexes										
2002	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000	6.0	8.1	12.1	17.7	26.9	45.4	76.0	123.5	243.1	591.5
1999	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1
Black, male										
2002	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000	8.1	11.0	17.0	25.5	35.0	64.3	106.6	170.6	343.1	742.7
1999	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3
Black, female										
2002	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000	4.2	5.6	7.9	11.2	20.4	31.5	55.7	95.1	192.8	535.3
1999	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001	*	*	*	*	*	*	*	0.3	0.7	3.3
2000	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999	*	*	*	*	*	*	0.2	0.6	1.3	6.0
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	2.8
2001	*	*	*	*	*	*	*	*	*	2.3
2000	*	*	*	*	*	*	*	0.7	*	3.8
1999	*	*	*	*	*	*	*	0.7	1.6	4.0
All races 3/, female										
2002	*	*	*	*	*	*	*	*	0.9	3.5
2001	*	*	*	*	*	*	*	*	0.7	3.7
2000	*	*	*	*	*	*	*	0.5	0.9	4.8
1999	*	*	*	*	*	*	*	0.5	1.1	6.8
White, both sexes										
2002	*	*	*	*	*	*	*	*	0.8	3.5
2001	*	*	*	*	*	*	*	0.3	0.7	3.5
2000	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999	*	*	*	*	*	*	*	0.6	1.3	6.3
White, male										
2002	*	*	*	*	*	*	*	*	*	2.9
2001	*	*	*	*	*	*	*	*	*	2.4
2000	*	*	*	*	*	*	*	0.8	*	3.8
1999	*	*	*	*	*	*	*	0.8	1.6	4.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)										
White, female										
2002	*	*	*	*	*	*	*	*	0.9	3.7
2001	*	*	*	*	*	*	*	*	0.8	3.9
2000	*	*	*	*	*	*	*	*	1.0	5.1
1999	*	*	*	*	*	*	*	0.5	1.1	7.2
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	0.5	2.0
2001	*	*	*	*	*	*	*	*	0.4	2.1
2000	*	*	*	*	*	*	*	0.4	0.5	2.6
1999	*	*	*	*	*	*	*	0.4	0.8	3.6
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	1.7
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	2.4
1999	*	*	*	*	*	*	*	*	*	2.6
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	2.2
2001	*	*	*	*	*	*	*	*	*	2.4
2000	*	*	*	*	*	*	*	*	*	2.7
1999	*	*	*	*	*	*	*	*	0.7	4.0
White, both sexes										
2002	*	*	*	*	*	*	*	*	0.5	2.1
2001	*	*	*	*	*	*	*	*	0.5	2.1
2000	*	*	*	*	*	*	*	0.4	0.5	2.7
1999	*	*	*	*	*	*	*	0.4	0.9	3.8
White, male										
2002	*	*	*	*	*	*	*	*	*	1.7
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	2.4
1999	*	*	*	*	*	*	*	*	*	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)										
White, female										
2002	*	*	*	*	*	*	*	*	*	2.3
2001	*	*	*	*	*	*	*	*	*	2.5
2000	*	*	*	*	*	*	*	*	*	2.9
1999	*	*	*	*	*	*	*	*	0.7	4.3
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	0.4	1.3
2001	*	*	*	*	*	*	*	*	*	1.2
2000	*	*	*	*	*	*	*	*	*	1.9
1999	*	*	*	*	*	*	*	*	0.5	2.4
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	1.3
2001	*	*	*	*	*	*	*	*	*	1.3
2000	*	*	*	*	*	*	*	*	*	2.1
1999	*	*	*	*	*	*	*	*	*	2.8
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	1.4
2001	*	*	*	*	*	*	*	*	*	1.3
2000	*	*	*	*	*	*	*	*	*	2.0
1999	*	*	*	*	*	*	*	*	0.5	2.5
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)										
White, female										
2002	*	*	*	*	*	*	*	*	*	1.4
2001	*	*	*	*	*	*	*	*	*	1.5
2000	*	*	*	*	*	*	*	*	*	2.2
1999	*	*	*	*	*	*	*	*	*	2.9
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2002	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000	2.8	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0
All races 3/, male										
2002	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000	2.6	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7
All races 3/, female										
2002	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000	2.9	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6
White, both sexes										
2002	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000	2.3	5.3	11.4	30.1	65.5	130.5	229.6	343.1	495.5	675.6
1999	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9
White, male										
2002	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000	2.2	5.7	11.5	31.8	69.8	143.1	265.8	415.5	648.5	997.4
1999	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
White, female										
2002	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000	2.4	4.9	11.3	28.4	61.5	119.4	200.5	292.1	404.4	544.8
1999	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7
Black, both sexes										
2002	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000	6.1	11.0	18.2	33.0	57.4	97.3	151.2	211.6	271.3	373.8
1999	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2
Black, male										
2002	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000	5.9	11.4	19.6	39.5	73.5	138.4	226.3	326.5	448.5	693.1
1999	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9
Black, female										
2002	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000	6.2	10.6	16.9	27.7	44.6	67.0	101.2	142.4	182.1	255.0
1999	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)										
All races 3/, both sexes										
2002	*	*	0.1	0.1	0.3	0.5	1.0	1.8	2.9	8.1
2001	*	0.1	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000	*	*	*	0.1	0.4	0.7	1.0	2.1	3.6	11.8
1999	*	*	0.1	0.3	0.6	0.7	1.2	1.9	4.1	10.7
All races 3/, male										
2002	*	*	*	*	*	0.7	1.1	2.2	2.9	8.8
2001	*	*	*	*	0.4	0.8	1.1	2.2	2.5	9.2
2000	*	*	*	*	0.4	0.9	1.0	2.8	4.9	12.9
1999	*	*	*	0.4	0.5	0.8	1.5	2.4	4.7	11.9
All races 3/, female										
2002	*	*	*	*	*	*	0.9	1.4	2.9	7.8
2001	*	*	*	*	*	0.4	0.7	1.4	2.6	9.1
2000	*	*	*	*	0.4	0.5	1.1	1.6	2.8	11.4
1999	*	*	*	*	0.6	0.5	1.0	1.5	3.8	10.3
White, both sexes										
2002	*	*	*	*	0.3	0.6	1.1	1.9	3.0	8.6
2001	*	*	0.1	0.2	0.3	0.6	0.9	1.8	2.7	9.6
2000	*	*	*	*	0.5	0.7	1.0	2.2	3.7	12.7
1999	*	*	*	0.3	0.6	0.7	1.2	1.9	4.1	11.2
White, male										
2002	*	*	*	*	*	0.7	1.2	2.4	2.9	9.3
2001	*	*	*	*	*	0.8	1.0	2.2	2.5	9.4
2000	*	*	*	*	0.5	1.0	1.0	2.8	5.2	13.8
1999	*	*	*	0.4	0.5	0.9	1.5	2.4	4.4	12.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)										
White, female										
2002	*	*	*	*	*	*	1.0	1.6	3.0	8.3
2001	*	*	*	*	*	*	0.7	1.5	2.8	9.7
2000	*	*	*	*	0.5	0.5	1.1	1.7	2.8	12.2
1999	*	*	*	*	0.7	0.5	1.0	1.5	3.9	10.8
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
All races 3/, both sexes										
2002	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1
All races 3/, male										
2002	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4
All races 3/, female										
2002	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7
White, both sexes										
2002	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
1999	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7
White, male										
2002	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
1999	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
White, female										
2002	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
1999	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9
Black, both sexes										
2002	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
1999	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6
Black, male										
2002	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
1999	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6
Black, female										
2002	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
1999	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
All races 3/, both sexes										
2002	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3
All races 3/, male										
2002	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7
All races 3/, female										
2002	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2
White, both sexes										
2002	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
1999	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6
White, male										
2002	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
1999	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
White, female										
2002	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
1999	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9
Black, both sexes										
2002	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
1999	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4
Black, male										
2002	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
1999	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*
Black, female										
2002	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0
1999	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)										
All races 3/, both sexes										
2002	1.4	3.5	8.7	21.8	49.3	98.4	175.4	279.8	400.3	555.5
2001	1.3	3.3	8.4	22.4	49.8	100.7	177.4	275.4	386.8	557.4
2000	1.2	3.4	8.2	22.4	49.1	99.2	177.1	272.5	400.7	556.7
1999	1.2	3.1	8.2	23.5	51.6	102.5	183.2	280.8	405.5	555.8
All races 3/, male										
2002	1.6	3.8	9.5	23.3	53.2	109.1	204.3	343.0	511.9	785.1
2001	1.3	3.8	9.0	24.2	54.0	111.6	208.6	341.5	500.4	794.5
2000	1.2	3.8	8.7	24.4	54.5	111.6	210.9	337.1	533.3	838.0
1999	1.2	3.7	8.6	25.8	58.7	119.6	223.9	357.9	544.9	866.7
All races 3/, female										
2002	1.3	3.3	7.9	20.4	45.7	89.3	152.3	235.2	332.4	455.8
2001	1.4	2.8	7.8	20.7	46.1	91.4	152.7	229.1	318.6	458.2
2000	1.2	2.9	7.8	20.5	44.3	88.7	150.4	227.6	322.5	442.2
1999	1.1	2.6	7.8	21.4	45.2	87.9	151.4	227.2	324.0	430.5
White, both sexes										
2002	1.4	3.6	8.8	22.9	51.9	105.4	185.7	294.0	418.6	580.7
2001	1.4	3.3	8.6	23.4	53.1	106.6	187.6	289.8	404.4	581.3
2000	1.2	3.4	8.4	23.2	51.8	105.0	186.7	285.5	419.7	581.2
1999	1.1	3.0	8.4	24.4	54.2	108.1	191.8	292.0	422.6	580.3
White, male										
2002	1.6	3.8	9.5	23.7	54.7	114.7	211.4	353.0	526.8	810.4
2001	1.3	3.9	9.0	24.8	56.2	115.4	215.7	351.7	514.4	817.8
2000	1.2	3.8	8.5	24.9	56.2	115.1	217.3	345.8	550.1	866.7
1999	1.1	3.5	8.6	26.2	60.3	123.4	230.5	365.1	558.5	897.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)										
White, female										
2002	1.3	3.5	8.2	22.1	49.3	97.1	164.8	251.8	352.3	480.9
2001	1.4	2.8	8.2	22.1	50.2	98.9	164.8	246.0	337.9	482.3
2000	1.1	3.0	8.2	21.6	47.7	96.1	162.1	243.0	342.1	465.2
1999	1.1	2.6	8.3	22.7	48.7	94.7	161.1	240.6	342.4	453.1
Black, both sexes										
2002	1.7	3.9	10.1	20.3	41.8	67.8	116.9	184.3	252.5	320.1
2001	1.6	4.1	9.4	20.7	37.6	75.7	119.1	173.3	238.4	320.9
2000	1.9	3.9	10.1	21.7	42.2	72.8	120.7	174.7	227.9	310.6
1999	2.0	4.8	8.5	23.9	43.1	77.1	129.9	199.1	248.8	313.7
Black, male										
2002	2.0	4.8	12.7	27.5	56.2	88.8	174.3	291.6	407.3	565.0
2001	1.6	4.4	11.3	26.6	50.4	105.7	179.4	285.3	408.5	583.2
2000	1.7	5.0	12.5	27.3	56.1	104.1	187.9	284.9	383.6	576.6
1999	2.5	6.0	11.0	31.2	61.3	109.6	192.3	325.8	441.2	607.0
Black, female										
2002	1.4	3.1	8.0	14.4	30.5	52.2	78.4	120.2	171.9	226.3
2001	1.5	3.8	7.7	15.8	27.5	53.5	78.9	106.5	151.7	223.5
2000	1.9	2.8	8.0	17.0	31.2	49.7	76.1	108.2	149.4	211.6
1999	1.5	3.7	6.5	17.9	28.8	53.1	88.4	122.7	153.4	204.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)										
All races 3/, both sexes										
2002	*	*	*	0.2	0.4	0.8	1.6	3.4	4.2	6.8
2001	*	*	*	0.2	0.3	1.0	2.0	3.4	5.1	6.2
2000	*	*	*	0.2	0.3	0.9	2.2	3.4	4.8	7.5
1999	*	*	*	*	0.4	1.1	1.9	3.4	5.2	6.1
All races 3/, male										
2002	*	*	*	0.3	0.8	1.7	3.5	8.0	10.5	21.6
2001	*	*	*	0.4	0.7	2.0	4.4	7.8	13.2	20.2
2000	*	*	*	*	0.7	1.8	4.8	7.9	12.3	24.9
1999	*	*	*	*	0.7	2.3	4.4	8.1	13.5	20.4
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	0.2	0.4	0.9	1.8	3.6	4.4	7.2
2001	*	*	*	0.2	0.3	1.1	2.1	3.7	5.4	6.5
2000	*	*	*	0.2	0.4	1.0	2.4	3.5	5.0	7.9
1999	*	*	*	*	0.3	1.1	2.0	3.6	5.5	6.3
White, male										
2002	*	*	*	0.4	0.8	1.9	3.8	8.3	11.0	22.8
2001	*	*	*	0.4	0.7	2.2	4.6	8.4	13.7	21.2
2000	*	*	*	*	0.7	1.9	5.1	8.3	12.8	26.4
1999	*	*	*	*	0.6	2.4	4.6	8.5	14.3	21.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes										
2002	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6
All races 3/, male										
2002	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All races 3/, female										
2002	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2
White, both sexes										
2002	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
1999	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0
White, male										
2002	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
White, female										
2002	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6
Black, both sexes										
2002	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1
Black, male										
2002	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5
Black, female										
2002	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
All races 3/, both sexes										
2002	1.5	2.5	4.1	7.0	11.9	20.6	37.6	56.8	83.7	133.5
2001	1.5	2.6	4.4	7.3	12.6	22.8	38.0	61.8	89.6	142.7
2000	1.7	2.6	4.6	7.8	12.7	22.9	36.9	60.5	89.2	141.7
1999	1.6	2.7	4.1	6.9	12.7	22.5	36.8	59.5	87.6	147.6
All races 3/, male										
2002	1.6	2.5	4.3	7.9	13.9	24.4	47.8	70.5	104.2	156.6
2001	1.8	2.9	4.9	7.7	15.0	27.1	47.0	76.2	110.0	169.1
2000	1.8	2.8	5.0	8.8	14.6	26.8	45.7	77.3	112.4	179.7
1999	1.7	2.8	4.4	7.4	14.7	27.3	46.4	74.9	115.0	182.3
All races 3/, female										
2002	1.4	2.4	3.9	6.2	10.2	17.3	29.4	47.0	71.1	123.4
2001	1.3	2.3	4.0	6.8	10.5	19.1	30.8	51.7	77.3	131.6
2000	1.6	2.4	4.2	6.8	10.8	19.6	29.9	48.8	75.6	126.2
1999	1.4	2.6	3.8	6.5	10.9	18.4	29.3	48.8	71.6	133.6
White, both sexes										
2002	1.3	2.1	3.8	6.7	11.6	21.0	38.2	58.7	85.5	137.9
2001	1.3	2.3	3.9	7.0	12.4	23.0	38.7	63.5	92.3	146.0
2000	1.6	2.2	4.3	7.5	12.4	22.8	37.7	62.0	91.2	145.9
1999	1.3	2.3	3.8	6.4	12.6	22.6	37.0	60.6	89.5	150.7
White, male										
2002	1.4	2.1	4.0	7.6	13.4	25.3	48.3	72.5	106.8	162.0
2001	1.7	2.5	4.4	7.6	14.5	27.5	48.3	78.4	113.3	173.5
2000	1.7	2.4	4.6	8.5	14.4	26.8	46.3	78.9	115.0	184.4
1999	1.5	2.4	4.2	6.7	14.7	27.7	47.0	76.5	117.6	185.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
White, female										
2002	1.2	2.0	3.6	5.9	10.0	17.2	30.0	48.7	72.5	127.4
2001	1.0	2.0	3.4	6.4	10.4	19.1	31.0	52.9	79.6	134.4
2000	1.5	2.1	3.9	6.6	10.6	19.3	30.7	50.1	77.1	130.3
1999	1.1	2.2	3.5	6.2	10.7	18.2	29.1	49.5	72.9	136.6
Black, both sexes										
2002	2.9	5.5	6.9	11.1	16.8	20.7	33.9	40.5	68.3	90.3
2001	3.3	5.7	9.3	10.9	17.0	24.2	36.1	50.2	66.9	109.6
2000	3.0	6.0	7.8	11.6	15.5	28.1	33.4	49.8	71.2	98.2
1999	3.4	5.8	7.1	11.7	15.4	25.0	38.0	52.8	66.2	111.6
Black, male										
2002	3.4	5.7	7.6	12.4	21.2	22.2	44.4	53.3	86.4	100.8
2001	3.2	6.2	9.4	10.5	21.6	28.0	39.4	60.4	85.1	129.3
2000	3.3	7.2	8.6	13.3	17.6	32.5	40.5	67.6	82.6	132.6
1999	3.3	6.6	7.8	13.7	17.8	28.3	44.1	66.5	83.7	145.3
Black, female										
2002	2.3	5.3	6.2	10.0	13.3	19.6	26.9	32.9	58.9	86.3
2001	3.3	5.3	9.1	11.3	13.4	21.4	33.9	44.1	57.6	102.3
2000	2.6	5.0	7.1	10.3	13.9	24.8	28.6	39.0	65.4	85.4
1999	3.5	5.1	6.6	10.1	13.5	22.5	34.0	44.6	57.5	99.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
All races 3/, both sexes										
2002	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000	0.3	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6
All races 3/, male										
2002	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000	0.4	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5
All races 3/, female										
2002	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000	0.2	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5
White, both sexes										
2002	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000	0.3	0.6	0.9	1.4	2.2	3.0	5.1	8.5	16.0	36.3
1999	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3
White, male										
2002	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000	0.4	0.8	1.3	2.0	2.9	3.8	6.7	10.9	17.8	40.7
1999	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
White, female										
2002	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000	*	0.3	0.5	0.9	1.6	2.4	3.8	6.9	14.9	34.5
1999	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1
Black, both sexes										
2002	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000	*	1.4	2.0	2.6	2.7	4.1	5.7	9.4	11.0	26.6
1999	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5
Black, male										
2002	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000	*	*	3.3	3.5	*	*	10.0	12.8	*	31.1
1999	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1
Black, female										
2002	*	*	*	*	*	*	*	6.9	*	18.5
2001	*	*	*	*	*	*	*	6.1	9.0	19.8
2000	*	*	*	*	*	*	*	7.4	8.5	24.9
1999	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)										
All races 3/, both sexes										
2002	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5
All races 3/, male										
2002	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000	*	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999	*	*	*	*	*	*	0.5	0.9	*	2.9
All races 3/, female										
2002	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001	*	*	*	*	*	*	0.4	*	0.7	1.8
2000	*	*	*	*	*	*	0.4	0.5	1.2	1.5
1999	*	*	*	*	*	*	*	*	1.1	0.9
White, both sexes										
2002	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5
White, male										
2002	*	*	*	*	0.5	*	0.7	1.2	1.6	2.0
2001	*	*	*	*	*	*	1.0	0.7	1.5	2.5
2000	*	*	*	*	0.5	*	*	0.9	1.5	2.0
1999	*	*	*	*	*	*	*	0.9	*	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)										
White, female										
2002	*	*	*	*	*	*	0.5	0.6	0.9	2.1
2001	*	*	*	*	*	*	*	*	*	1.9
2000	*	*	*	*	*	*	*	*	1.2	1.6
1999	*	*	*	*	*	*	*	*	1.1	0.9
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
All races 3/, both sexes										
2002	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7
All races 3/, male										
2002	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2
All races 3/, female										
2002	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6
White, both sexes										
2002	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
1999	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0
White, male										
2002	*	*	*	0.4	0.5	0.8	1.6	2.9	5.4	11.9
2001	*	*	0.3	*	0.7	0.9	1.4	2.3	5.2	11.8
2000	*	*	0.3	*	0.6	0.8	1.5	3.1	5.7	12.7
1999	*	*	*	0.4	0.8	0.8	1.6	2.5	6.6	12.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
White, female										
2002	*	*	*	0.4	0.6	0.9	1.6	3.0	5.4	14.0
2001	*	*	0.3	0.5	0.6	0.8	1.6	2.6	5.0	13.2
2000	*	*	0.3	0.5	0.6	0.8	1.5	2.9	4.6	13.6
1999	*	*	*	*	0.5	0.7	1.5	2.5	4.9	13.1
Black, both sexes										
2002	*	*	*	*	*	*	*	*	5.2	7.7
2001	*	*	*	*	*	*	*	*	*	11.1
2000	*	*	*	*	*	*	*	3.9	*	12.5
1999	*	*	*	*	*	*	*	3.6	*	9.9
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	8.8
2000	*	*	*	*	*	*	*	*	*	12.4
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)										
All races 3/, both sexes										
2002	9.7	16.0	20.3	21.7	24.5	27.9	31.2	32.5	29.9	21.4
2001	9.8	17.1	20.1	21.1	24.7	28.9	31.1	31.3	28.7	22.2
2000	10.2	16.8	18.7	21.9	26.2	28.9	30.8	31.9	29.6	23.1
1999	9.9	16.4	18.6	21.4	26.5	29.7	31.6	32.6	30.8	23.2
All races 3/, male										
2002	12.9	23.3	31.0	32.0	34.5	37.2	40.0	39.4	36.9	28.4
2001	13.2	25.2	30.1	31.5	35.3	39.7	40.8	37.6	35.3	32.0
2000	14.4	25.4	28.6	31.7	37.7	39.3	40.7	40.7	34.6	34.3
1999	14.5	25.3	28.8	32.5	37.2	41.8	41.5	40.1	38.7	33.1
All races 3/, female										
2002	6.5	8.9	10.1	12.1	15.3	19.8	24.1	27.6	25.6	18.4
2001	6.4	9.3	10.5	11.4	15.2	19.6	23.5	26.9	24.7	18.1
2000	6.1	8.4	9.1	12.7	15.7	19.9	23.1	25.8	26.6	18.5
1999	5.4	7.8	8.8	11.1	16.8	19.4	23.9	27.4	26.2	19.3
White, both sexes										
2002	10.0	16.1	20.2	21.8	24.8	28.9	32.7	34.2	30.9	22.3
2001	9.8	17.2	19.6	20.7	25.3	29.7	32.3	32.6	30.0	23.2
2000	10.1	16.9	18.1	21.6	26.5	29.4	31.8	33.4	30.5	23.9
1999	9.7	15.8	18.3	20.8	26.3	30.2	32.6	33.7	32.0	23.8
White, male										
2002	13.5	23.8	30.9	32.1	34.7	38.2	41.6	41.6	38.1	29.6
2001	13.2	25.5	29.3	30.6	35.6	40.2	42.1	39.1	37.2	33.5
2000	14.4	26.0	27.8	31.5	38.3	40.2	41.7	42.3	35.9	35.5
1999	14.2	24.6	28.4	31.5	37.4	42.2	42.5	41.1	40.3	34.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)										
White, female										
2002	6.5	8.5	9.8	11.9	15.6	20.7	25.4	28.9	26.5	19.1
2001	6.3	8.9	10.0	11.3	15.8	20.4	24.3	28.0	25.7	18.9
2000	5.9	7.8	8.7	12.2	15.6	20.0	23.9	27.2	27.3	19.2
1999	5.1	7.2	8.5	10.7	16.2	19.8	24.7	28.4	27.2	19.6
Black, both sexes										
2002	7.9	17.4	25.0	24.6	25.5	22.9	20.0	20.1	16.0	12.2
2001	10.6	19.2	26.8	27.7	23.6	26.9	22.6	21.0	14.6	10.4
2000	10.9	19.2	25.9	25.9	25.9	27.5	24.7	18.3	17.5	13.8
1999	11.9	22.5	22.9	28.2	30.7	27.7	25.3	23.2	18.8	17.2
Black, male										
2002	9.8	23.4	38.2	36.9	39.0	34.9	29.0	23.0	21.4	*
2001	14.5	28.2	42.2	43.8	38.4	43.2	33.7	26.9	*	*
2000	14.6	26.4	40.7	38.5	37.3	38.0	37.4	27.4	19.4	*
1999	17.1	34.5	36.6	44.1	42.4	43.4	37.3	31.8	25.3	*
Black, female										
2002	6.3	12.2	13.7	14.5	14.9	14.1	13.9	18.4	13.1	10.7
2001	7.1	11.3	13.7	14.5	12.0	14.8	15.2	17.5	14.7	9.7
2000	7.7	13.0	13.3	15.5	17.0	19.8	16.2	12.8	16.6	11.6
1999	7.3	12.2	11.4	15.2	21.5	16.2	17.3	18.0	15.6	15.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2002	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8
All races 3/, male										
2002	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1
All races 3/, female										
2002	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5
White, both sexes										
2002	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
1999	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7
White, male										
2002	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
1999	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
White, female										
2002	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
1999	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5
Black, both sexes										
2002	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
1999	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*
Black, male										
2002	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
1999	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*
Black, female										
2002	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
1999	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)										
All races 3/, both sexes										
2002	3.7	6.6	8.9	10.3	12.6	17.2	21.7	25.8	25.5	19.3
2001	3.6	6.9	8.8	10.0	12.8	17.7	22.2	24.4	24.7	19.9
2000	3.6	6.8	7.9	10.2	13.3	17.5	21.7	24.7	25.2	20.9
1999	3.5	6.4	7.7	9.9	13.9	18.1	22.3	25.4	25.6	20.4
All races 3/, male										
2002	4.9	9.5	13.3	14.4	15.3	19.8	24.6	28.6	28.9	23.7
2001	4.8	10.0	13.1	13.5	16.3	21.8	25.6	25.5	28.4	26.5
2000	5.0	10.2	11.6	13.7	17.4	20.6	25.0	27.0	26.8	29.0
1999	5.2	9.8	11.6	13.8	17.3	22.0	25.3	28.0	29.0	27.0
All races 3/, female										
2002	2.6	3.8	4.7	6.5	10.2	14.9	19.4	23.9	23.4	17.4
2001	2.5	4.0	4.7	6.7	9.6	14.2	19.5	23.6	22.6	17.1
2000	2.2	3.6	4.3	6.9	9.5	14.8	19.2	23.2	24.2	17.5
1999	1.8	3.2	4.0	6.2	10.8	14.8	19.9	23.5	23.6	17.7
White, both sexes										
2002	3.9	6.5	8.7	10.4	12.8	18.0	22.8	27.4	26.4	20.2
2001	3.6	6.6	8.5	9.7	13.1	18.5	23.1	25.5	26.0	20.9
2000	3.6	6.9	7.5	10.1	13.5	17.8	22.5	26.1	26.2	21.6
1999	3.4	6.0	7.6	9.6	13.9	18.7	23.2	26.5	26.8	21.1
White, male										
2002	5.2	9.5	13.0	14.3	15.3	20.5	25.8	30.5	30.0	25.1
2001	4.7	9.6	12.5	13.0	16.2	22.4	26.7	26.6	30.0	28.1
2000	5.0	10.3	11.0	13.5	17.6	21.2	25.8	28.2	28.1	30.1
1999	5.1	9.2	11.3	13.4	17.3	22.8	26.3	29.1	30.5	28.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)										
White, female										
2002	2.5	3.5	4.6	6.6	10.5	15.7	20.4	25.1	24.3	18.1
2001	2.4	3.8	4.6	6.6	10.2	15.1	20.2	24.7	23.5	17.9
2000	2.1	3.4	4.2	6.9	9.6	14.8	19.9	24.6	25.1	18.2
1999	1.8	2.9	3.9	6.0	10.7	15.1	20.7	24.6	24.6	18.2
Black, both sexes										
2002	3.7	9.2	12.8	12.1	13.5	13.4	12.6	13.9	12.0	9.8
2001	4.7	10.2	12.1	13.6	12.2	13.9	14.8	15.3	11.6	7.7
2000	4.1	8.2	12.6	12.0	13.3	15.7	16.3	12.1	13.3	12.2
1999	4.7	9.8	10.1	13.4	15.1	14.8	14.9	15.2	13.3	12.8
Black, male										
2002	4.2	11.9	19.5	19.5	18.6	18.4	16.0	12.4	*	*
2001	6.4	15.3	19.9	20.3	19.7	20.3	18.5	18.1	*	*
2000	5.0	11.4	20.1	18.1	19.3	18.7	20.7	16.1	*	*
1999	7.0	15.3	16.4	19.2	19.1	19.2	18.7	17.6	*	*
Black, female										
2002	3.3	6.9	7.1	6.1	9.4	9.6	10.3	14.8	10.7	9.9
2001	3.2	5.8	5.5	8.1	6.2	9.2	12.3	13.6	13.1	*
2000	3.2	5.5	6.1	7.0	8.7	13.4	13.3	9.7	14.0	11.2
1999	2.6	5.0	4.8	8.7	12.0	11.5	12.4	13.7	12.1	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All races 3/, both sexes										
2002	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9
All races 3/, male										
2002	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7
All races 3/, female										
2002	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4
White, both sexes										
2002	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
1999	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4
White, male										
2002	*	0.3	0.5	0.8	1.1	2.3	4.0	7.4	13.3	24.7
2001	*	0.3	0.4	0.8	1.4	1.9	4.4	7.6	12.0	26.7
2000	*	*	0.3	0.7	1.1	2.3	3.4	7.4	13.2	28.9
1999	*	0.3	0.3	0.7	1.2	1.9	4.2	7.0	13.8	28.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)										
White, female										
2002	*	*	0.3	0.4	0.9	1.7	3.1	4.1	9.0	23.0
2001	*	0.3	*	0.4	1.1	1.8	3.1	4.4	8.4	22.8
2000	*	*	0.3	0.4	0.7	1.4	2.2	4.4	8.3	24.3
1999	*	0.3	0.4	0.6	0.6	1.4	2.7	4.8	9.3	22.9
Black, both sexes										
2002	*	*	*	*	*	2.3	4.6	7.7	12.8	14.8
2001	*	*	*	1.4	2.0	3.5	6.2	6.1	11.6	19.7
2000	*	*	*	*	*	3.1	4.9	6.6	11.0	17.8
1999	*	*	*	*	2.0	3.2	4.1	4.6	11.3	15.3
Black, male										
2002	*	*	*	*	*	*	*	*	16.0	*
2001	*	*	*	*	*	*	*	*	*	27.2
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000	*	*	*	*	*	*	4.9	6.3	11.5	19.3
1999	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes										
2002	2.1	3.7	5.9	10.0	16.8	29.4	50.0	85.4	142.2	288.6
2001	2.1	3.6	5.8	9.9	16.9	31.7	49.3	82.2	135.8	287.7
2000	2.1	3.4	5.5	9.4	16.9	29.5	47.2	79.5	132.8	277.8
1999	2.0	3.1	5.2	9.0	15.8	28.2	46.7	77.7	127.4	268.9
All races 3/, male										
2002	2.3	4.3	6.9	11.0	18.6	33.6	59.8	105.7	182.7	385.5
2001	2.3	4.4	6.7	10.9	18.2	36.5	57.0	100.5	176.8	393.7
2000	2.5	4.4	6.4	10.7	18.6	32.8	54.3	99.8	173.0	380.7
1999	2.3	3.8	5.8	9.9	17.8	31.0	54.9	99.2	168.9	371.1
All races 3/, female										
2002	1.9	3.1	4.9	9.2	15.2	25.8	42.1	71.1	117.6	246.6
2001	1.9	2.9	4.9	8.9	15.7	27.6	43.2	69.4	111.2	243.3
2000	1.6	2.5	4.6	8.2	15.5	26.6	41.6	65.3	109.0	235.9
1999	1.7	2.4	4.5	8.1	14.0	25.8	40.3	62.8	103.1	227.7
White, both sexes										
2002	1.4	2.6	3.9	7.4	13.1	24.4	43.2	78.1	135.5	278.8
2001	1.4	2.4	4.1	7.3	12.8	25.8	43.1	75.1	127.9	278.2
2000	1.5	2.2	3.8	6.8	13.2	23.8	40.6	73.0	125.3	268.7
1999	1.3	2.0	3.5	6.6	12.2	23.0	40.4	70.9	118.5	255.7
White, male										
2002	1.6	3.0	4.6	8.0	14.3	28.4	53.0	99.0	177.9	377.4
2001	1.6	2.8	4.9	8.2	14.4	30.5	51.3	94.6	169.0	388.9
2000	1.8	2.9	4.4	7.9	14.9	27.8	48.7	93.4	166.7	376.2
1999	1.5	2.4	4.0	7.8	14.0	26.0	48.9	91.6	159.6	360.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
White, female										
2002	1.3	2.3	3.3	6.7	12.0	20.8	35.1	63.2	109.4	235.9
2001	1.2	1.9	3.3	6.3	11.4	21.7	36.4	61.3	103.1	231.9
2000	1.1	1.7	3.2	5.8	11.6	20.3	34.1	58.6	100.7	225.0
1999	1.1	1.6	3.1	5.5	10.6	20.4	33.6	56.4	94.3	213.5
Black, both sexes										
2002	6.7	11.7	21.6	33.7	50.4	79.2	126.6	178.0	241.5	441.6
2001	6.7	12.5	19.8	32.5	53.1	88.6	119.8	172.3	248.6	429.6
2000	6.3	12.7	19.7	32.3	50.7	83.5	119.7	165.4	242.3	406.4
1999	7.0	11.8	19.6	29.1	47.1	77.8	117.0	163.7	246.2	433.3
Black, male										
2002	7.6	14.7	26.8	40.0	61.2	89.3	149.6	207.6	281.2	561.8
2001	7.0	16.5	22.8	37.7	55.5	101.1	130.9	189.1	309.8	511.7
2000	7.9	17.4	24.1	37.4	54.2	86.9	127.7	198.9	290.0	490.1
1999	8.2	15.5	23.4	29.2	54.5	81.8	130.6	208.0	327.0	526.1
Black, female										
2002	5.9	9.1	17.2	28.5	42.0	71.8	111.1	160.3	220.8	395.5
2001	6.4	9.1	17.1	28.2	51.2	79.4	112.3	162.3	217.5	399.1
2000	4.8	8.6	16.1	28.2	47.9	81.0	114.3	145.3	218.2	375.2
1999	5.9	8.6	16.4	29.1	41.3	74.8	108.0	137.0	206.1	398.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	0.6	0.9
2001	*	*	*	*	*	*	*	0.4	0.4	0.9
2000	*	*	*	*	*	*	*	*	0.7	1.2
1999	*	*	*	*	*	*	0.3	0.5	0.5	0.8
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	1.8
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	0.6	0.9
2001	*	*	*	*	*	*	*	*	*	0.7
2000	*	*	*	*	*	*	*	*	0.7	0.9
1999	*	*	*	*	*	*	*	0.5	*	0.7
White, both sexes										
2002	*	*	*	*	*	*	*	*	0.5	0.8
2001	*	*	*	*	*	*	*	0.4	*	0.8
2000	*	*	*	*	*	*	*	*	0.8	1.1
1999	*	*	*	*	*	*	0.3	0.5	0.5	0.8
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)										
White, female										
2002	*	*	*	*	*	*	*	*	*	0.8
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	0.7	1.0
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)										
All races 3/, both sexes										
2002	*	0.1	0.1	0.2	*	0.3	0.6	1.2	1.7	2.8
2001	*	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000	0.1	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	2.8
1999	*	*	0.2	0.2	0.3	0.4	0.7	1.4	2.0	3.2
All races 3/, male										
2002	*	*	*	0.3	*	*	0.9	1.6	2.1	3.8
2001	*	*	*	*	*	0.5	1.0	1.3	2.3	3.7
2000	*	*	*	*	*	0.5	0.8	1.5	2.1	4.5
1999	*	*	*	*	*	0.5	0.6	1.9	2.7	5.0
All races 3/, female										
2002	*	*	*	*	*	*	0.4	0.8	1.5	2.3
2001	*	*	*	*	*	0.5	0.5	0.7	1.0	2.4
2000	*	*	*	*	*	*	0.6	0.8	1.8	2.1
1999	*	*	*	*	*	0.4	0.7	1.1	1.5	2.4
White, both sexes										
2002	*	*	*	0.2	*	0.3	0.7	1.0	1.7	2.6
2001	*	*	*	*	*	0.4	0.7	0.9	1.5	2.9
2000	*	*	*	0.2	*	0.4	0.7	1.0	1.9	2.4
1999	*	*	*	*	0.3	0.3	0.6	1.4	1.8	2.9
White, male										
2002	*	*	*	*	*	*	0.9	1.5	2.1	3.8
2001	*	*	*	*	*	*	1.1	1.2	2.3	4.0
2000	*	*	*	*	*	*	0.8	1.4	2.2	4.0
1999	*	*	*	*	*	*	0.6	1.9	2.6	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)										
White, female										
2002	*	*	*	*	*	*	0.5	0.7	1.4	2.0
2001	*	*	*	*	*	*	0.5	0.7	0.9	2.4
2000	*	*	*	*	*	*	0.6	0.7	1.7	1.8
1999	*	*	*	*	*	*	0.7	1.0	1.4	2.2
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
All races 3/, both sexes										
2002	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000	2.0	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8
All races 3/, male										
2002	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000	2.4	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6
All races 3/, female										
2002	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000	1.6	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5
White, both sexes										
2002	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000	1.4	2.1	3.7	6.6	12.8	23.3	39.6	71.7	122.6	265.0
1999	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9
White, male										
2002	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000	1.7	2.7	4.2	7.6	14.5	27.2	47.7	91.6	163.8	370.4
1999	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
White, female										
2002	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000	1.1	1.6	3.1	5.6	11.3	19.9	33.2	57.8	98.1	222.1
1999	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6
Black, both sexes										
2002	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000	6.1	12.3	18.9	32.0	50.1	82.1	118.6	163.1	238.9	400.4
1999	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3
Black, male										
2002	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000	7.7	17.1	22.7	37.0	53.2	84.8	127.0	196.6	287.5	480.9
1999	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2
Black, female										
2002	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000	4.7	8.2	15.7	27.8	47.6	80.1	113.0	143.0	214.4	370.4
1999	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of kidney (N25,N27)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of kidney (N25,N27)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)										
All races 3/, both sexes										
2002	*	*	0.1	0.2	0.3	0.5	0.7	1.6	2.3	6.3
2001	*	0.1	0.1	0.2	0.2	0.4	0.6	1.7	2.4	6.0
2000	*	0.1	0.1	0.2	0.3	0.4	0.7	1.5	2.7	7.5
1999	*	0.1	0.2	0.2	0.3	0.5	0.9	1.5	2.7	7.5
All races 3/, male										
2002	*	*	*	*	*	*	0.7	1.4	1.4	5.3
2001	*	*	*	*	*	*	0.5	1.2	2.1	4.3
2000	*	*	*	*	*	*	*	1.1	2.2	5.6
1999	*	*	*	*	*	*	0.6	1.2	2.6	6.8
All races 3/, female										
2002	*	*	*	*	0.4	0.6	0.7	1.7	2.8	6.7
2001	*	*	0.2	*	*	*	0.6	2.1	2.6	6.7
2000	*	*	*	0.3	0.4	0.5	1.1	1.8	3.0	8.3
1999	*	*	0.3	*	*	0.6	1.1	1.7	2.8	7.8
White, both sexes										
2002	*	*	*	0.2	0.2	0.5	0.8	1.6	2.3	6.4
2001	*	*	*	*	*	0.3	0.6	1.7	2.2	6.1
2000	*	*	*	0.2	0.3	0.4	0.7	1.4	2.5	7.6
1999	*	*	0.2	0.2	0.2	0.5	0.9	1.5	2.8	7.7
White, male										
2002	*	*	*	*	*	*	0.8	1.4	1.4	5.4
2001	*	*	*	*	*	*	*	1.1	2.0	4.3
2000	*	*	*	*	*	*	*	1.1	2.1	5.9
1999	*	*	*	*	*	*	*	1.3	2.6	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)										
White, female										
2002	*	*	*	*	*	0.6	0.8	1.7	2.9	6.8
2001	*	*	*	*	*	*	0.7	2.2	2.4	6.9
2000	*	*	*	*	*	0.5	1.0	1.7	2.8	8.3
1999	*	*	*	*	*	0.6	1.1	1.7	2.9	7.9
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	6.2
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	7.8
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	9.0
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hyperplasia of prostate (N40)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	0.4	0.6	2.2	4.8
2001	*	*	*	*	*	*	0.4	0.8	1.6	5.5
2000	*	*	*	*	*	*	0.3	0.9	1.9	5.2
1999	*	*	*	*	*	*	0.4	0.8	2.1	5.0
All races 3/, male										
2002	*	*	*	*	*	*	0.8	1.6	5.8	15.8
2001	*	*	*	*	*	*	0.8	2.0	4.2	18.6
2000	*	*	*	*	*	*	0.7	2.1	5.2	18.1
1999	*	*	*	*	*	*	0.8	2.0	5.6	17.5
All races 3/, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
White, both sexes										
2002	*	*	*	*	*	*	0.3	0.6	2.2	4.9
2001	*	*	*	*	*	*	0.4	0.8	1.6	5.7
2000	*	*	*	*	*	*	0.3	0.9	2.0	5.2
1999	*	*	*	*	*	*	0.4	0.8	2.1	5.1
White, male										
2002	*	*	*	*	*	*	0.8	1.4	5.9	16.1
2001	*	*	*	*	*	*	0.8	2.0	4.2	19.4
2000	*	*	*	*	*	*	0.7	2.1	5.3	18.1
1999	*	*	*	*	*	*	0.8	2.1	5.5	17.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hyperplasia of prostate (N40)										
White, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	0.8
2001	*	*	*	*	*	*	*	*	*	0.9
2000	*	*	*	*	*	*	*	*	*	0.6
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	1.1
2001	*	*	*	*	*	*	*	*	*	1.2
2000	*	*	*	*	*	*	*	*	*	0.9
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	0.8
2001	*	*	*	*	*	*	*	*	*	0.9
2000	*	*	*	*	*	*	*	*	*	0.6
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)										
White, female										
2002	*	*	*	*	*	*	*	*	*	1.1
2001	*	*	*	*	*	*	*	*	*	1.2
2000	*	*	*	*	*	*	*	*	*	0.9
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
All races 3/, both sexes										
2002	0.1	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All races 3/, both sexes										
2002	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8
All races 3/, male										
2002	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5
All races 3/, female										
2002	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6
White, both sexes										
2002	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
1999	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0
White, male										
2002	1.4	1.8	2.0	2.3	2.8	2.6	2.7	3.4	4.6	7.3
2001	1.6	1.5	1.9	2.2	2.6	2.7	3.2	3.9	4.6	7.5
2000	1.4	1.6	2.3	2.0	2.8	2.6	2.6	3.4	5.6	11.1
1999	1.3	1.9	2.0	2.6	2.6	2.5	2.6	3.0	5.8	8.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
White, female										
2002	1.2	1.5	2.0	2.4	2.2	2.4	2.4	3.0	5.0	6.7
2001	1.3	1.4	1.9	2.2	2.8	2.1	2.3	3.1	4.2	7.3
2000	1.1	1.3	1.7	2.4	2.4	2.7	2.3	3.4	4.9	6.5
1999	1.2	1.7	1.7	2.3	2.3	2.9	2.3	3.6	4.6	7.7
Black, both sexes										
2002	1.8	1.8	1.9	2.8	2.4	2.1	2.7	*	*	*
2001	1.6	1.8	1.6	1.9	1.9	*	2.8	*	*	6.1
2000	1.6	1.7	1.5	2.4	2.5	2.8	*	*	*	*
1999	1.3	1.9	1.9	1.9	2.0	2.3	3.4	3.9	*	7.7
Black, male										
2002	2.1	2.3	2.3	3.3	*	*	*	*	*	*
2001	2.1	*	*	*	*	*	*	*	*	*
2000	2.0	2.0	*	*	*	*	*	*	*	*
1999	1.8	2.5	*	*	*	*	*	*	*	*
Black, female										
2002	1.6	1.5	*	*	*	*	*	*	*	*
2001	*	2.0	*	*	*	*	*	*	*	*
2000	*	*	*	2.8	3.3	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All races 3/, both sexes										
2002	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7
All races 3/, male										
2002	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3
All races 3/, female										
2002	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4
White, both sexes										
2002	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
1999	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2
White, male										
2002	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
1999	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
White, female										
2002	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
1999	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4
Black, both sexes										
2002	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
1999	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4
Black, male										
2002	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
1999	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0
Black, female										
2002	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7
1999	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9, N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
All races 3/, both sexes										
2002	23.5	33.6	42.6	56.2	77.1	110.4	173.8	297.8	555.5	1,493.0
2001	21.7	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
2000	22.1	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
1999	21.6	29.9	38.4	51.0	72.1	107.7	167.8	281.3	510.8	1,410.7
All races 3/, male										
2002	28.2	41.0	52.4	64.4	87.9	122.8	186.8	309.6	555.2	1,262.9
2001	25.7	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
2000	26.3	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
1999	26.6	36.3	45.4	59.5	80.9	117.1	184.3	298.1	528.1	1,225.4
All races 3/, female										
2002	18.9	26.4	33.2	48.6	67.3	99.6	163.5	289.5	555.6	1,592.8
2001	17.7	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
2000	17.9	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
1999	16.6	23.7	31.7	43.1	64.2	99.5	154.8	269.7	500.7	1,485.3
White, both sexes										
2002	21.6	30.9	39.2	53.1	73.4	107.7	170.2	295.6	556.2	1,517.6
2001	19.3	28.7	37.3	49.5	72.6	108.9	169.2	292.6	541.2	1,498.0
2000	19.8	27.4	37.1	49.3	70.1	102.5	164.5	285.7	537.2	1,464.6
1999	19.1	26.3	34.8	47.4	68.6	103.2	163.6	277.1	512.1	1,434.0
White, male										
2002	26.8	38.2	48.7	60.6	83.4	118.6	181.2	304.6	553.0	1,274.7
2001	23.4	35.3	45.1	55.3	81.8	119.9	182.1	302.0	540.6	1,282.9
2000	24.0	35.0	44.1	56.8	77.5	112.7	177.5	298.2	533.4	1,274.8
1999	24.3	32.5	41.6	55.4	76.8	111.8	179.1	290.9	525.5	1,231.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9, N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
White, female										
2002	16.3	23.6	29.9	46.0	64.2	98.1	161.2	289.2	558.2	1,623.1
2001	15.3	22.3	29.7	44.0	64.2	99.2	158.8	285.9	541.6	1,588.0
2000	15.5	19.8	30.3	42.2	63.2	93.6	154.2	277.0	539.4	1,541.7
1999	13.8	20.2	28.3	39.9	61.0	95.7	151.2	267.4	504.2	1,515.3
Black, both sexes										
2002	41.6	60.3	78.4	96.1	123.7	155.7	242.8	369.7	642.1	1,425.4
2001	41.2	63.0	79.4	95.6	122.1	163.6	237.0	380.4	612.1	1,418.0
2000	41.9	64.3	75.6	95.5	122.2	157.2	233.9	372.5	618.9	1,388.9
1999	42.5	61.8	75.4	92.1	117.2	167.2	239.7	369.5	556.5	1,286.2
Black, male										
2002	44.2	70.9	94.6	116.3	146.1	185.2	288.4	419.3	693.1	1,319.8
2001	47.0	74.8	95.9	114.8	141.6	196.9	273.3	445.6	659.7	1,313.9
2000	48.0	74.7	89.1	114.4	150.1	188.4	279.8	428.6	692.9	1,325.1
1999	47.6	73.3	89.5	111.0	134.9	189.5	274.1	434.1	635.7	1,291.3
Black, female										
2002	39.3	51.0	64.6	79.6	106.1	133.8	212.2	340.0	615.5	1,465.9
2001	36.2	52.9	65.3	79.8	106.7	138.9	212.8	341.4	587.9	1,456.6
2000	36.5	55.2	64.0	79.9	100.2	134.2	203.3	338.6	581.7	1,412.7
1999	37.9	52.0	63.5	76.6	103.3	150.6	216.9	330.7	517.3	1,284.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2002	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4
All races 3/, male										
2002	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4
All races 3/, female										
2002	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8
White, both sexes										
2002	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
1999	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4
White, male										
2002	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
1999	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
White, female										
2002	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
1999	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0
Black, both sexes										
2002	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
1999	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6
Black, male										
2002	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
1999	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4
Black, female										
2002	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
1999	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2002	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0
All races 3/, male										
2002	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5
All races 3/, female										
2002	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3
White, both sexes										
2002	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
1999	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3
White, male										
2002	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
1999	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
White, female										
2002	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
1999	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7
Black, both sexes										
2002	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
1999	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7
Black, male										
2002	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
1999	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0
Black, female										
2002	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9
1999	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2002	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000	15.1	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4
All races 3/, male										
2002	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000	21.9	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8
All races 3/, female										
2002	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000	8.4	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5
White, both sexes										
2002	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000	15.0	14.5	13.4	14.1	14.7	14.9	17.9	24.0	29.2	31.1
1999	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7
White, male										
2002	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000	21.6	20.9	18.9	19.3	19.2	19.1	23.4	32.0	41.7	59.2
1999	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
White, female										
2002	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000	8.4	8.1	7.9	9.1	10.6	11.3	13.5	18.4	21.8	19.7
1999	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8
Black, both sexes										
2002	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000	17.9	17.6	18.3	16.7	15.4	16.6	20.1	18.1	20.0	20.3
1999	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5
Black, male										
2002	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000	28.4	26.3	28.1	28.2	25.7	27.8	31.4	32.1	38.8	43.8
1999	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9
Black, female										
2002	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000	8.6	10.0	9.9	7.3	7.4	8.3	12.7	9.7	10.6	11.6
1999	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)										
All races 3/, both sexes										
2002	0.5	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.6	*
2001	0.6	0.7	0.6	0.5	0.6	0.7	0.6	0.8	0.8	0.7
2000	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.7	0.8	0.8
1999	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.3	1.4	1.3
All races 3/, male										
2002	0.9	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.2	*
2001	1.0	1.0	0.9	0.8	0.9	1.1	0.9	1.2	*	*
2000	1.2	0.9	0.9	0.7	1.0	0.9	0.9	1.1	1.3	*
1999	1.5	1.4	1.2	1.2	1.1	1.2	1.2	1.7	2.1	2.9
All races 3/, female										
2002	0.2	*	*	*	*	*	*	*	*	*
2001	0.2	0.4	0.2	*	*	*	*	0.5	0.7	*
2000	0.3	0.3	0.4	*	*	*	*	*	*	*
1999	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.9	1.0	*
White, both sexes										
2002	0.5	0.5	0.5	0.4	0.4	0.3	0.4	0.4	0.6	*
2001	0.6	0.6	0.6	0.5	0.5	0.6	0.6	0.8	0.7	0.6
2000	0.7	0.6	0.6	0.4	0.6	0.5	0.5	0.7	0.8	0.8
1999	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.2	1.4	1.3
White, male										
2002	0.8	0.9	0.8	0.6	0.7	0.5	0.7	*	1.3	*
2001	0.9	0.9	0.9	0.8	0.9	1.0	0.9	1.3	*	*
2000	1.2	0.8	0.8	0.6	1.0	0.8	0.9	1.1	1.3	*
1999	1.5	1.3	1.2	1.3	1.1	1.2	1.1	1.7	2.0	2.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	0.2	0.3	*	*	*	*	*	*	*	*
2000	0.3	0.3	0.4	*	*	*	*	*	*	*
1999	0.4	0.5	0.3	0.4	0.4	0.5	*	0.9	1.0	*
Black, both sexes										
2002	0.8	1.1	*	*	*	*	*	*	*	*
2001	1.1	1.2	*	*	*	*	*	*	*	*
2000	1.0	1.1	*	*	*	*	*	*	*	*
1999	0.8	1.0	*	*	*	*	*	*	*	*
Black, male										
2002	1.5	2.4	*	*	*	*	*	*	*	*
2001	2.0	1.9	*	*	*	*	*	*	*	*
2000	1.7	1.8	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
All races 3/, both sexes										
2002	0.9	1.1	1.0	1.1	1.1	1.0	0.8	0.8	0.7	0.4
2001	1.2	1.1	1.2	1.0	1.1	1.0	0.9	0.8	0.7	0.5
2000	1.0	0.9	1.0	1.0	1.1	1.2	0.7	0.6	0.7	*
1999	0.9	1.0	1.0	1.1	1.1	0.9	0.8	0.5	0.6	*
All races 3/, male										
2002	1.5	1.9	1.6	1.9	1.9	2.0	1.7	1.4	1.5	*
2001	1.8	1.8	1.9	1.7	1.9	1.8	1.7	1.3	1.4	*
2000	1.7	1.6	1.8	1.8	1.9	2.0	1.4	1.2	1.4	*
1999	1.5	1.7	1.8	2.0	1.9	1.6	1.6	0.9	*	*
All races 3/, female										
2002	0.3	0.3	0.3	0.3	0.4	*	*	*	*	*
2001	0.5	0.4	0.4	0.4	*	*	*	0.5	*	*
2000	0.3	0.3	0.3	*	0.4	0.4	*	*	*	*
1999	0.4	0.3	0.3	*	*	*	*	*	*	*
White, both sexes										
2002	1.0	1.2	1.0	1.1	1.2	1.0	0.9	0.8	0.7	*
2001	1.3	1.1	1.2	1.0	1.2	1.0	1.0	0.8	0.7	*
2000	1.1	1.0	1.1	1.1	1.2	1.2	0.8	0.6	0.7	*
1999	1.0	1.1	1.1	1.1	1.1	0.9	0.9	0.5	0.5	*
White, male										
2002	1.6	2.1	1.7	1.9	2.1	2.0	1.8	1.3	1.5	*
2001	2.0	1.8	2.0	1.6	2.1	1.8	1.8	1.3	1.5	*
2000	1.9	1.6	2.0	1.9	2.0	2.0	1.5	1.2	1.5	*
1999	1.5	1.8	1.8	2.0	1.9	1.6	1.6	0.9	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
White, female										
2002	0.3	0.2	0.4	0.4	0.4	*	*	*	*	*
2001	0.6	0.4	0.5	0.5	*	*	*	*	*	*
2000	0.4	0.3	0.3	*	0.4	*	*	*	*	*
1999	0.4	0.3	0.3	*	*	*	*	*	*	*
Black, both sexes										
2002	*	0.8	*	*	*	*	*	*	*	*
2001	*	1.1	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	2.0	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)										
All races 3/, both sexes										
2002	22.3	21.8	18.4	15.5	16.0	21.7	30.6	55.0	101.4	246.7
2001	19.4	19.7	16.3	14.1	16.1	20.3	29.8	54.3	100.1	247.5
2000	18.6	18.0	15.3	14.2	15.9	18.6	29.4	50.4	94.6	242.0
1999	18.8	18.2	14.3	13.9	15.5	20.6	30.0	52.7	101.3	250.3
All races 3/, male										
2002	32.3	31.5	26.8	22.6	22.9	30.0	41.4	71.9	127.5	293.6
2001	28.7	30.0	24.3	20.5	22.5	27.8	38.8	72.1	127.8	305.8
2000	28.3	28.0	23.7	20.7	22.7	25.5	38.4	65.3	120.8	299.8
1999	29.1	28.6	22.0	20.6	22.7	28.3	40.7	66.6	133.5	311.9
All races 3/, female										
2002	12.5	12.5	10.4	8.9	9.7	14.5	21.9	43.1	85.5	226.3
2001	10.2	9.7	8.6	8.2	10.3	13.9	22.6	41.8	83.5	223.1
2000	9.0	8.4	7.3	8.0	9.7	12.8	22.4	40.0	79.2	218.4
1999	8.7	8.1	6.9	7.6	9.0	14.0	21.7	43.0	82.4	225.5
White, both sexes										
2002	23.1	21.6	17.7	14.7	15.5	20.8	30.5	55.6	104.3	256.5
2001	19.3	19.0	15.4	13.4	15.3	20.0	29.4	54.6	102.1	255.7
2000	18.3	17.2	14.1	13.3	15.3	17.9	28.9	50.4	96.5	250.6
1999	18.5	17.1	13.2	12.9	14.2	19.7	28.7	51.9	102.6	258.1
White, male										
2002	33.3	31.1	25.2	20.9	21.7	28.2	41.1	72.1	130.1	302.8
2001	28.7	28.6	22.4	19.0	21.1	27.1	38.1	72.0	128.6	314.0
2000	27.9	26.4	21.5	19.0	21.6	24.0	37.5	64.4	122.0	308.2
1999	28.5	26.6	20.0	18.6	20.9	26.5	38.6	64.8	135.0	319.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)										
White, female										
2002	12.8	12.1	10.4	8.8	9.7	14.3	21.9	43.8	88.5	236.3
2001	9.7	9.4	8.5	8.1	10.0	13.8	22.3	42.2	86.1	231.2
2000	8.7	8.1	6.9	7.9	9.5	12.6	21.9	40.6	81.4	227.2
1999	8.5	7.8	6.5	7.4	8.0	13.7	20.8	42.9	83.5	233.3
Black, both sexes										
2002	22.9	29.4	28.6	25.2	23.2	29.9	35.0	54.8	72.7	155.0
2001	24.9	29.6	28.0	22.8	25.1	26.9	39.0	56.1	84.8	166.1
2000	24.5	27.6	27.7	23.9	22.9	27.8	38.8	56.0	79.1	159.5
1999	25.6	30.7	26.2	24.6	28.6	32.7	45.6	65.7	95.9	179.8
Black, male										
2002	32.7	42.9	46.3	42.1	38.0	47.1	50.1	79.5	105.5	214.4
2001	36.0	47.8	46.1	37.2	38.8	40.9	56.2	80.4	133.2	220.1
2000	37.8	45.3	46.5	40.5	37.9	42.9	52.8	85.1	116.3	230.6
1999	40.7	51.5	44.1	42.2	42.2	50.6	70.2	95.0	128.2	248.4
Black, female										
2002	14.2	17.7	13.6	11.2	11.6	17.2	24.9	40.1	55.7	132.3
2001	15.2	13.8	12.5	11.0	14.2	16.5	27.6	41.6	60.1	146.1
2000	12.7	12.3	11.7	10.3	11.1	16.7	29.5	38.4	60.3	133.1
1999	12.3	12.9	11.0	10.2	18.0	19.4	29.3	48.1	79.9	154.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
All races 3/, both sexes										
2002	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000	1.6	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2
All races 3/, male										
2002	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000	2.7	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5
All races 3/, female										
2002	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000	0.6	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2
White, both sexes										
2002	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000	1.7	2.1	2.5	3.2	4.6	6.2	12.3	23.1	46.6	119.1
1999	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9
White, male										
2002	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000	2.7	3.5	3.9	4.8	6.9	8.4	16.2	30.3	59.7	152.8
1999	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
White, female										
2002	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000	0.6	0.7	1.2	1.8	2.6	4.3	9.3	18.1	38.8	105.4
1999	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6
Black, both sexes										
2002	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000	1.6	2.0	3.6	4.5	5.1	7.1	9.7	11.6	20.3	38.8
1999	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3
Black, male										
2002	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000	2.6	3.4	6.3	7.7	8.7	10.7	12.4	18.4	26.1	61.1
1999	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6
Black, female										
2002	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000	*	*	*	*	*	4.4	8.0	7.4	17.4	30.5
1999	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental discharge of firearms (W32-W34)										
All races 3/, both sexes										
2002	0.3	0.2	0.2	0.2	0.2	*	*	*	*	*
2001	0.3	0.3	0.2	0.3	0.2	*	*	0.3	*	*
2000	0.3	0.2	0.2	0.3	*	0.3	*	*	*	*
1999	0.3	0.3	0.2	0.2	0.2	*	*	*	*	*
All races 3/, male										
2002	0.5	0.4	0.4	0.4	*	*	*	*	*	*
2001	0.6	0.5	0.4	0.5	0.4	*	*	0.7	*	*
2000	0.6	0.3	0.4	0.5	*	0.5	*	*	*	*
1999	0.5	0.5	0.4	0.3	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.3	0.2	0.2	0.2	*	*	*	*	*	*
2001	0.3	0.3	0.3	0.3	0.3	*	*	0.3	*	*
2000	0.4	0.2	0.2	0.3	*	0.3	*	*	*	*
1999	0.3	0.3	0.3	0.2	0.2	*	*	*	*	*
White, male										
2002	0.5	0.4	0.4	0.5	*	*	*	*	*	*
2001	0.6	0.5	0.5	0.5	0.4	*	*	0.8	*	*
2000	0.6	0.3	0.5	0.5	*	*	*	*	*	*
1999	0.5	0.5	0.4	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental discharge of firearms (W32-W34)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2002	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6
All races 3/, male										
2002	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3
All races 3/, female										
2002	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4
White, both sexes										
2002	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
1999	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6
White, male										
2002	1.9	1.4	1.4	1.3	1.2	1.4	1.6	1.8	2.4	3.7
2001	1.6	1.7	1.3	1.1	1.3	1.4	1.6	1.8	2.3	3.5
2000	1.6	1.5	1.5	1.6	1.6	1.4	1.7	1.1	3.0	2.7
1999	1.7	1.8	1.7	1.4	1.4	1.4	1.5	1.9	2.7	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
White, female										
2002	0.4	0.4	0.3	0.3	0.4	*	*	0.6	0.8	0.7
2001	0.4	0.3	0.3	0.4	0.5	*	*	0.7	0.7	1.0
2000	0.3	0.3	0.3	*	*	*	0.6	0.7	1.0	1.0
1999	0.4	0.4	0.3	0.4	*	*	0.7	0.5	0.7	1.3
Black, both sexes										
2002	1.1	1.3	1.1	*	*	*	*	*	*	*
2001	1.3	1.4	*	1.5	*	*	*	*	*	*
2000	1.5	1.0	1.6	*	*	*	*	*	*	*
1999	1.0	*	1.1	*	*	*	*	*	*	*
Black, male										
2002	2.1	2.4	*	*	*	*	*	*	*	*
2001	2.2	2.4	*	*	*	*	*	*	*	*
2000	2.8	1.9	2.8	*	*	*	*	*	*	*
1999	1.9	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2002	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9
All races 3/, male										
2002	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3
All races 3/, female										
2002	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5
White, both sexes										
2002	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7
White, male										
2002	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
1999	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
White, female										
2002	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3
Black, both sexes										
2002	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
1999	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7
Black, male										
2002	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
1999	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3
Black, female										
2002	*	*	*	*	*	*	*	6.0	9.2	11.9
2001	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6
1999	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2002	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4
All races 3/, male										
2002	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4
All races 3/, female										
2002	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0
White, both sexes										
2002	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
1999	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3
White, male										
2002	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
1999	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, female										
2002	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
1999	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8
Black, both sexes										
2002	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
1999	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*
Black, male										
2002	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
1999	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*
Black, female										
2002	10.5	12.5	8.1	4.3	*	*	*	*	*	*
2001	10.7	8.9	7.5	3.1	*	*	*	*	*	*
2000	8.6	8.0	6.6	3.0	*	*	*	*	*	*
1999	9.1	7.8	4.9	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
All races 3/, both sexes										
2002	3.3	3.5	4.0	4.8	5.7	8.6	11.1	21.6	38.8	105.6
2001	3.2	3.9	4.5	4.6	6.3	7.8	12.3	22.5	42.2	115.0
2000	3.4	3.9	4.1	5.0	6.3	7.8	12.2	22.0	41.4	117.1
1999	3.7	3.8	4.4	5.1	6.3	8.9	13.3	23.6	46.9	128.9
All races 3/, male										
2002	5.4	5.8	6.2	7.4	8.7	12.0	15.0	28.6	47.2	118.6
2001	5.3	6.3	6.8	6.7	8.9	10.9	15.9	29.3	52.7	134.0
2000	5.6	6.5	6.7	7.7	9.1	10.8	15.6	28.1	51.6	135.2
1999	6.1	6.0	7.2	7.8	9.1	12.0	18.0	29.9	59.3	152.7
All races 3/, female										
2002	1.3	1.3	1.9	2.4	3.0	5.7	8.1	16.6	33.8	100.0
2001	1.2	1.6	2.2	2.5	3.8	5.2	9.5	17.7	35.9	107.0
2000	1.3	1.4	1.7	2.4	3.8	5.2	9.6	17.8	35.4	109.8
1999	1.4	1.7	1.7	2.5	3.7	6.3	9.6	19.3	39.6	119.3
White, both sexes										
2002	3.4	3.5	4.0	4.6	5.6	7.9	11.0	21.4	39.1	108.8
2001	3.2	3.8	4.4	4.5	5.9	7.7	11.9	22.3	42.5	117.7
2000	3.4	3.7	4.0	4.7	6.2	7.4	11.8	21.7	41.9	120.5
1999	3.7	3.6	4.2	4.7	5.8	8.6	12.7	22.8	46.8	132.4
White, male										
2002	5.6	5.9	6.1	6.9	8.5	10.9	14.9	28.1	47.2	121.2
2001	5.2	6.2	6.6	6.5	8.5	10.7	15.3	29.1	51.9	137.0
2000	5.6	6.1	6.4	7.3	8.7	10.1	15.1	27.1	51.2	137.6
1999	6.0	5.5	6.8	7.1	8.5	11.3	17.0	28.7	59.0	155.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
White, female										
2002	1.2	1.1	1.9	2.4	2.9	5.3	7.8	16.5	34.2	103.4
2001	1.1	1.5	2.3	2.6	3.5	5.1	9.1	17.4	36.8	109.7
2000	1.3	1.4	1.6	2.3	3.8	5.1	9.1	17.9	36.4	113.6
1999	1.5	1.6	1.7	2.4	3.3	6.2	9.3	18.7	39.6	123.2
Black, both sexes										
2002	3.3	4.3	5.1	7.3	7.3	15.9	14.3	27.8	39.0	83.1
2001	3.9	5.4	5.8	5.9	10.4	10.4	18.7	27.3	43.9	96.1
2000	3.9	5.7	5.6	7.9	8.6	12.3	18.1	30.2	39.2	90.1
1999	4.7	6.6	6.6	9.3	10.9	13.9	22.9	36.2	53.7	105.5
Black, male										
2002	4.3	6.4	8.6	12.7	11.3	23.7	18.7	40.9	53.5	106.1
2001	6.6	8.6	9.5	9.9	14.6	15.8	25.2	35.3	72.2	114.6
2000	6.5	10.1	10.0	13.8	13.9	18.2	23.1	46.8	64.1	124.5
1999	8.1	11.1	11.0	15.4	16.1	20.8	34.9	50.3	69.8	153.5
Black, female										
2002	2.4	2.5	2.1	2.9	4.2	10.2	11.4	20.0	31.4	74.3
2001	1.6	2.7	2.7	2.7	7.0	6.4	14.3	22.5	29.4	89.3
2000	1.5	1.8	*	3.0	4.4	7.9	14.9	20.2	26.7	77.3
1999	1.6	2.7	2.8	4.3	6.8	8.9	14.9	27.7	45.8	87.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2002	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 4/	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000	15.4	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3
All races 3/, male										
2002	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 4/	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000	23.9	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1
All races 3/, female										
2002	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 4/	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000	7.0	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1
White, both sexes										
2002	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 4/	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000	17.0	16.5	15.3	13.9	12.6	12.3	14.7	17.6	20.4	21.0
1999	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7
White, male										
2002	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 4/	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000	26.2	25.6	23.2	21.8	20.4	21.3	27.6	37.1	47.7	61.6
1999	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, female										
2002	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 4/	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000	7.9	7.5	7.5	6.4	5.5	4.4	4.4	3.9	4.2	4.5
1999	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3
Black, both sexes										
2002	7.1	5.9	5.9	4.6	5.0	3.5	6.6	6.7	*	5.9
2001 4/	6.8	6.5	5.9	5.8	4.4	4.6	6.5	5.2	6.5	*
2000	8.0	6.6	6.4	4.6	3.7	4.3	6.1	5.3	6.5	*
1999	6.8	6.4	4.7	5.2	4.7	6.4	5.1	6.2	*	*
Black, male										
2002	11.5	11.0	9.2	8.6	8.1	7.7	12.4	15.6	*	*
2001 4/	10.8	10.1	9.6	10.2	8.3	8.1	14.1	12.1	16.1	*
2000	13.4	11.8	10.5	8.4	6.4	9.4	13.4	11.8	*	*
1999	10.8	10.9	8.7	9.3	8.9	12.2	11.2	14.2	*	*
Black, female										
2002	3.2	1.5	3.0	*	*	*	*	*	*	*
2001 4/	3.3	3.3	2.7	*	*	*	*	*	*	*
2000	3.2	2.1	2.9	*	*	*	*	*	*	*
1999	3.3	2.4	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2002	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000	7.4	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6
All races 3/, male										
2002	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000	12.6	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6
All races 3/, female										
2002	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000	2.4	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9
White, both sexes										
2002	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000	8.4	8.3	8.7	8.2	8.5	9.2	11.1	13.4	15.2	14.3
1999	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6
White, male										
2002	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000	14.0	14.2	14.7	14.2	15.3	17.3	22.7	30.4	38.1	46.5
1999	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, female										
2002	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0
Black, both sexes										
2002	3.0	2.9	2.9	2.7	2.9	2.3	4.6	5.3	*	*
2001	3.2	2.6	3.0	3.0	2.4	2.9	4.7	4.5	6.2	*
2000	3.3	3.3	3.3	2.9	3.0	3.1	4.5	*	*	*
1999	2.7	3.4	2.4	3.5	3.4	4.0	3.9	5.0	*	*
Black, male										
2002	5.2	5.7	5.3	5.5	5.5	5.0	10.0	13.3	*	*
2001	5.4	4.4	5.4	6.1	5.1	5.6	10.4	10.7	*	*
2000	6.1	6.0	5.6	5.8	5.6	7.0	11.0	*	*	*
1999	5.1	6.0	4.9	7.1	6.8	7.8	9.8	11.9	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
All races 3/, both sexes										
2002	8.7	8.5	7.2	5.9	4.2	3.7	3.9	4.2	5.0	6.0
2001 4/	8.2	8.2	6.6	5.6	4.0	3.7	3.6	4.0	4.8	5.3
2000	7.9	7.4	6.2	5.2	3.7	3.0	3.5	4.2	5.3	6.5
1999	7.7	6.9	6.2	4.7	3.9	3.8	3.6	4.5	5.7	6.7
All races 3/, male										
2002	12.1	11.6	9.9	8.2	5.7	5.0	5.5	6.5	8.1	12.9
2001 4/	11.6	11.5	8.6	7.5	5.3	5.4	5.4	6.1	8.7	12.2
2000	11.3	10.6	8.1	6.9	4.8	3.9	4.8	6.8	9.4	14.6
1999	10.8	9.6	8.1	6.4	5.1	5.2	5.3	6.6	9.5	15.5
All races 3/, female										
2002	5.4	5.6	4.6	3.9	2.9	2.5	2.6	2.6	3.1	3.0
2001 4/	4.8	5.1	4.6	3.8	2.8	2.2	2.3	2.6	2.4	2.4
2000	4.6	4.4	4.3	3.7	2.8	2.1	2.5	2.4	2.8	3.2
1999	4.5	4.2	4.4	3.2	2.8	2.5	2.3	3.1	3.4	3.1
White, both sexes										
2002	9.7	9.6	7.9	6.4	4.6	3.8	4.0	4.3	5.3	6.2
2001 4/	9.1	9.1	7.2	5.9	4.2	3.8	3.7	4.2	5.0	5.5
2000	8.7	8.2	6.6	5.7	4.1	3.1	3.6	4.2	5.3	6.7
1999	8.4	7.6	6.7	5.1	4.1	3.8	3.8	4.7	5.9	7.1
White, male										
2002	13.4	12.6	10.9	8.7	6.2	5.0	5.8	6.7	8.4	13.4
2001 4/	12.7	12.6	9.3	7.9	5.3	5.5	5.3	6.3	9.0	12.9
2000	12.2	11.4	8.5	7.6	5.2	4.0	4.9	6.7	9.6	15.2
1999	11.9	10.4	8.6	6.7	5.2	5.3	5.5	6.7	10.0	16.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
White, female										
2002	6.0	6.5	5.0	4.2	3.0	2.7	2.5	2.6	3.4	3.1
2001 4/	5.4	5.6	5.1	4.1	3.1	2.3	2.5	2.8	2.5	2.3
2000	5.1	5.0	4.8	4.0	3.1	2.3	2.6	2.5	2.7	3.3
1999	4.9	4.7	4.9	3.5	3.0	2.6	2.4	3.3	3.5	3.3
Black, both sexes										
2002	4.2	3.0	3.0	1.9	2.2	*	*	*	*	*
2001 4/	3.6	3.9	2.9	2.7	2.0	*	*	*	*	*
2000	4.7	3.3	3.1	1.7	*	*	*	*	*	*
1999	4.1	3.0	2.3	1.7	*	2.4	*	*	*	*
Black, male										
2002	6.3	5.3	3.9	3.1	*	*	*	*	*	*
2001 4/	5.3	5.7	4.2	4.1	*	*	*	*	*	*
2000	7.3	5.7	4.9	*	*	*	*	*	*	*
1999	5.7	4.9	3.8	*	*	*	*	*	*	*
Black, female										
2002	2.3	*	2.2	*	*	*	*	*	*	*
2001 4/	2.1	2.3	*	*	*	*	*	*	*	*
2000	2.3	*	*	*	*	*	*	*	*	*
1999	2.7	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All races 3/, both sexes										
2002	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 4/	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4
All races 3/, male										
2002	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 4/	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4
All races 3/, female										
2002	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 4/	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0
White, both sexes										
2002	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 4/	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
1999	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1
White, male										
2002	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 4/	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
1999	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
White, female										
2002	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 4/	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
1999	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9
Black, both sexes										
2002	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 4/	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
1999	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*
Black, male										
2002	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 4/	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
1999	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*
Black, female										
2002	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 4/	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2002	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6
All races 3/, male										
2002	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*
All races 3/, female										
2002	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*
White, both sexes										
2002	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
1999	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6
White, male										
2002	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
1999	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
White, female										
2002	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*
Black, both sexes										
2002	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
1999	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*
Black, male										
2002	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
1999	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*
Black, female										
2002	4.6	2.8	2.8	*	*	*	*	*	*	*
2001	3.9	2.4	1.9	*	*	*	*	*	*	*
2000	4.7	3.2	*	*	*	*	*	*	*	*
1999	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)										
All races 3/, both sexes										
2002	2.6	2.6	1.8	1.7	1.5	1.4	1.4	1.3	1.6	1.7
2001 4/	4.9	4.4	3.4	2.9	2.1	2.1	1.6	1.6	1.8	2.0
2000	2.9	2.4	1.8	1.6	1.4	1.4	1.4	1.6	2.1	1.7
1999	2.9	2.3	1.9	1.6	1.3	1.6	1.4	1.7	1.7	1.8
All races 3/, male										
2002	3.4	3.8	2.6	2.6	2.3	2.0	2.2	1.8	1.9	2.3
2001 4/	7.0	6.4	5.0	4.4	3.0	2.8	2.2	2.0	2.1	2.7
2000	3.9	3.3	2.6	2.6	2.0	2.0	2.1	2.0	3.0	2.5
1999	4.0	3.3	2.8	2.5	1.8	2.6	1.7	2.2	2.6	2.6
All races 3/, female										
2002	1.8	1.4	1.1	1.0	0.7	0.9	0.8	0.9	1.4	1.4
2001 4/	2.9	2.5	1.8	1.5	1.4	1.4	1.2	1.4	1.6	1.7
2000	1.8	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
1999	1.8	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.2	1.5
White, both sexes										
2002	1.9	1.8	1.3	1.4	1.3	1.2	1.2	1.0	1.3	1.4
2001 4/	4.2	3.7	2.9	2.6	1.8	1.7	1.4	1.4	1.5	1.8
2000	1.9	1.7	1.4	1.2	1.0	1.0	1.1	1.4	1.6	1.5
1999	2.0	1.7	1.5	1.3	1.0	1.4	1.2	1.4	1.4	1.5
White, male										
2002	2.6	2.5	1.6	1.9	1.9	1.7	1.8	1.2	1.5	2.0
2001 4/	6.0	5.3	4.2	3.9	2.5	2.4	1.8	1.7	2.0	2.3
2000	2.7	2.1	1.8	1.9	1.5	1.3	1.6	1.7	2.2	2.1
1999	2.8	2.4	2.1	1.9	1.5	2.1	1.3	1.8	2.2	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)										
White, female										
2002	1.2	1.2	0.9	0.9	0.7	0.7	0.7	0.8	1.2	1.2
2001 4/	2.3	2.0	1.6	1.4	1.1	1.1	1.1	1.2	1.2	1.6
2000	1.2	1.2	0.9	0.6	0.6	0.7	0.7	1.1	1.2	1.3
1999	1.2	0.9	1.0	0.8	0.6	0.8	1.0	1.0	1.0	1.4
Black, both sexes										
2002	7.5	8.0	5.9	4.6	3.5	3.1	3.8	5.7	*	*
2001 4/	10.1	9.7	7.2	5.6	4.6	4.8	4.0	4.7	*	*
2000	9.2	7.4	5.3	5.0	4.7	5.4	4.1	4.1	6.8	*
1999	9.0	6.5	5.3	4.3	3.2	4.4	3.4	5.2	*	*
Black, male										
2002	9.7	13.4	10.3	8.5	6.9	*	7.3	10.1	*	*
2001 4/	13.9	14.7	11.3	8.8	6.9	6.9	7.1	*	*	*
2000	12.6	12.0	9.3	8.9	7.7	9.9	6.7	*	*	*
1999	13.6	10.0	9.0	7.6	4.5	8.6	*	*	*	*
Black, female										
2002	5.5	3.4	2.1	*	*	*	*	*	*	*
2001 4/	6.8	5.3	3.7	3.1	*	*	*	*	*	*
2000	6.2	3.3	2.0	*	*	*	*	*	*	*
1999	5.0	3.5	2.2	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Legal intervention (Y35,Y89.0)										
All races 3/, both sexes										
2002	0.2	0.2	*	*	*	*	*	*	*	*
2001	0.2	0.2	*	*	*	*	*	*	*	*
2000	0.2	0.2	*	*	*	*	*	*	*	*
1999	0.2	*	0.2	*	*	*	*	*	*	*
All races 3/, male										
2002	0.4	0.4	*	*	*	*	*	*	*	*
2001	0.4	0.3	*	*	*	*	*	*	*	*
2000	0.3	0.4	*	*	*	*	*	*	*	*
1999	0.4	*	0.4	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	0.2	0.2	*	*	*	*	*	*	*	*
2001	0.2	0.1	*	*	*	*	*	*	*	*
2000	0.1	0.2	*	*	*	*	*	*	*	*
1999	0.2	*	0.2	*	*	*	*	*	*	*
White, male										
2002	0.3	0.3	*	*	*	*	*	*	*	*
2001	0.4	0.3	*	*	*	*	*	*	*	*
2000	0.3	0.4	*	*	*	*	*	*	*	*
1999	0.3	*	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Legal intervention (Y35,Y89.0)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000	3.1	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4
All races 3/, male										
2002	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000	4.2	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1
All races 3/, female										
2002	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1
White, both sexes										
2002	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000	2.9	2.6	1.6	1.0	0.7	0.5	0.5	0.6	0.8	1.3
1999	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4
White, male										
2002	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000	4.0	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, female										
2002	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.1
1999	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1
Black, both sexes										
2002	4.3	4.5	3.8	1.5	*	*	*	*	*	*
2001	4.0	4.1	3.3	1.5	*	*	*	*	*	*
2000	4.8	4.4	2.8	2.3	*	*	*	*	*	*
1999	4.6	4.8	3.3	1.5	*	*	*	*	*	*
Black, male										
2002	5.5	6.2	6.5	*	*	*	*	*	*	*
2001	6.6	7.0	5.4	*	*	*	*	*	*	*
2000	7.2	7.0	4.4	4.0	*	*	*	*	*	*
1999	6.5	8.3	5.7	*	*	*	*	*	*	*
Black, female										
2002	3.3	3.1	*	*	*	*	*	*	*	*
2001	1.7	1.6	*	*	*	*	*	*	*	*
2000	2.7	2.1	*	*	*	*	*	*	*	*
1999	2.9	1.7	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2002	3.3	3.2	2.2	1.3	0.9	0.8	0.6	0.6	0.9	1.4
2001	3.1	2.9	1.9	1.0	0.6	0.5	0.4	0.6	1.0	1.0
2000	3.0	2.7	1.6	1.1	0.7	0.5	0.5	0.5	0.7	1.3
1999	2.8	2.6	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3
All races 3/, male										
2002	4.1	4.0	2.8	1.7	1.3	0.9	0.8	0.7	1.0	2.2
2001	4.2	4.1	2.4	1.4	0.7	0.8	0.6	0.7	1.3	*
2000	4.1	3.7	2.1	1.5	0.9	0.7	0.7	0.8	*	*
1999	3.8	3.5	2.0	1.4	0.9	0.9	*	0.9	*	1.8
All races 3/, female										
2002	2.6	2.5	1.7	0.9	0.7	0.6	0.4	0.5	0.8	1.1
2001	2.0	1.7	1.5	0.6	0.5	*	*	0.6	0.7	0.8
2000	1.9	1.6	1.1	0.7	0.4	*	*	*	*	1.2
1999	1.8	1.7	1.0	0.6	0.5	0.5	*	*	0.7	1.1
White, both sexes										
2002	3.4	3.2	2.1	1.3	0.9	0.7	0.6	0.5	0.8	1.4
2001	3.1	2.9	1.8	1.0	0.6	0.5	0.4	0.6	0.9	1.0
2000	2.8	2.5	1.5	0.9	0.6	0.5	0.5	0.5	0.7	1.2
1999	2.6	2.4	1.3	0.9	0.7	0.7	0.4	0.6	0.8	1.3
White, male										
2002	4.2	3.9	2.4	1.7	1.2	0.8	0.7	*	*	2.3
2001	4.1	3.9	2.2	1.3	0.6	0.7	*	0.7	1.4	*
2000	3.8	3.5	2.0	1.2	0.8	0.6	0.6	0.8	*	*
1999	3.6	3.1	1.7	1.2	0.9	0.9	*	0.8	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
White, female										
2002	2.6	2.5	1.9	1.0	0.7	0.6	0.5	*	0.7	1.1
2001	2.1	1.8	1.5	0.7	0.6	*	*	0.6	*	0.8
2000	1.8	1.6	1.1	0.7	0.4	*	*	*	*	1.1
1999	1.7	1.7	1.0	0.7	0.6	0.5	*	*	0.8	1.1
Black, both sexes										
2002	4.3	4.5	3.8	1.5	*	*	*	*	*	*
2001	3.9	4.0	3.2	*	*	*	*	*	*	*
2000	4.7	4.3	2.8	2.1	*	*	*	*	*	*
1999	4.5	4.6	3.3	*	*	*	*	*	*	*
Black, male										
2002	5.4	6.2	6.4	*	*	*	*	*	*	*
2001	6.5	6.8	5.2	*	*	*	*	*	*	*
2000	7.1	6.9	4.3	3.5	*	*	*	*	*	*
1999	6.4	7.9	5.7	*	*	*	*	*	*	*
Black, female										
2002	3.3	3.1	*	*	*	*	*	*	*	*
2001	1.6	1.6	*	*	*	*	*	*	*	*
2000	2.6	2.1	*	*	*	*	*	*	*	*
1999	2.7	1.7	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)										
All races 3/, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
White, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)										
White, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, both sexes										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, male										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*
Black, female										
2002	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
All races 3/, both sexes										
2002	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7
All races 3/, male										
2002	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5
All races 3/, female										
2002	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2
White, both sexes										
2002	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
1999	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1
White, male										
2002	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
1999	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
White, female										
2002	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
1999	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7
Black, both sexes										
2002	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
1999	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1
Black, male										
2002	*	*	*	*	*	5.0	*	*	*	*
2001	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000	*	*	*	*	4.6	6.2	9.4	10.4	*	*
1999	*	*	*	*	*	*	9.2	10.4	*	23.4
Black, female										
2002	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9
1999	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70,R78.0,X45,X65,Y15) 5/										
All races 3/, both sexes										
2002	10.5	15.3	18.0	18.0	18.3	16.4	14.3	10.8	7.2	4.3
2001	10.5	16.2	17.7	17.3	18.6	17.0	13.6	10.9	7.5	5.0
2000	10.8	15.7	16.5	18.0	19.1	17.1	14.1	11.0	7.9	5.3
1999	10.7	15.8	16.6	17.9	19.1	17.4	13.9	11.8	8.5	5.5
All races 3/, male										
2002	15.0	23.2	28.9	28.8	30.5	27.7	23.5	18.6	13.4	9.6
2001	15.1	24.9	27.6	28.3	30.5	28.1	23.6	19.2	13.8	11.5
2000	16.0	24.7	26.6	28.4	31.2	28.7	24.6	20.8	14.8	11.7
1999	16.3	24.9	27.0	29.3	31.1	29.7	24.2	20.8	16.5	11.8
All races 3/, female										
2002	6.0	7.6	7.6	7.8	7.2	6.6	6.9	5.3	3.4	2.0
2001	5.9	7.8	8.2	7.1	7.8	7.4	5.6	5.1	3.8	2.3
2000	5.7	7.0	6.9	8.2	8.2	7.1	5.8	4.2	3.8	2.7
1999	5.2	7.0	6.7	7.2	8.2	6.8	5.9	5.5	3.7	3.0
White, both sexes										
2002	10.8	15.6	17.9	17.5	17.7	16.4	14.1	10.8	7.1	4.3
2001	10.3	16.3	16.8	16.5	18.4	16.5	13.6	10.9	7.6	4.8
2000	10.6	15.3	16.0	17.2	18.5	16.8	14.0	11.0	7.8	5.2
1999	10.3	15.1	15.9	16.8	18.3	16.8	13.5	11.5	8.3	5.3
White, male										
2002	15.5	23.6	28.8	28.0	29.4	27.2	22.9	18.6	13.2	9.2
2001	14.8	25.0	26.4	26.7	29.8	26.8	23.3	19.0	14.0	11.0
2000	15.7	24.3	25.8	27.3	30.1	27.9	24.2	20.6	14.8	11.4
1999	15.8	24.0	25.9	27.3	30.0	28.2	23.1	20.1	16.4	11.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcohol-induced causes (F10,G31.2,G62.1,I42.6,K29.2,K70,R78.0,X45,X65,Y15) 5/										
White, female										
2002	6.0	7.6	7.4	7.5	6.9	6.8	7.0	5.3	3.4	2.1
2001	5.8	7.6	7.5	6.9	7.8	7.5	5.8	5.1	3.8	2.2
2000	5.5	6.4	6.4	7.6	7.9	7.0	5.9	4.3	3.7	2.7
1999	4.8	6.4	6.2	6.9	7.6	6.9	5.8	5.4	3.5	2.9
Black, both sexes										
2002	8.8	15.2	21.3	24.9	24.6	18.5	17.2	12.4	9.2	*
2001	11.8	17.9	26.5	26.2	24.1	24.4	14.5	13.0	7.3	*
2000	12.3	20.9	23.7	26.0	25.6	23.8	17.1	12.8	7.9	6.3
1999	12.6	22.1	23.7	28.6	28.4	24.4	20.8	16.1	11.3	7.3
Black, male										
2002	12.3	23.4	34.7	41.2	44.3	35.9	33.7	23.0	19.1	*
2001	17.9	28.0	42.5	46.3	43.3	46.2	29.6	25.6	*	*
2000	18.1	31.8	39.2	43.0	43.3	42.7	34.8	27.4	*	*
1999	18.5	34.7	39.3	51.4	47.5	47.8	40.7	30.4	21.8	*
Black, female										
2002	5.8	8.1	10.0	11.5	9.1	5.6	6.0	6.0	*	*
2001	6.5	9.1	12.9	9.6	9.0	8.3	4.5	5.5	*	*
2000	7.2	11.5	10.5	12.1	11.6	9.8	5.3	*	*	*
1999	7.5	11.1	10.5	10.0	13.5	7.1	7.7	7.4	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Drug-induced causes (F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5, F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) 5/										
All races 3/, both sexes										
2002	21.8	20.9	14.5	8.2	5.1	3.8	3.3	3.5	4.1	5.6
2001	18.5	17.6	11.6	6.3	4.1	3.1	2.9	3.4	3.5	5.1
2000	17.5	15.8	10.3	5.9	3.6	2.5	2.7	3.1	3.9	5.5
1999	17.3	15.5	9.3	5.8	3.7	3.2	2.8	3.5	4.1	4.8
All races 3/, male										
2002	28.0	26.3	19.1	10.0	5.7	4.1	3.3	3.8	4.6	7.3
2001	24.6	23.9	15.0	7.7	4.3	3.7	3.3	3.8	3.8	5.6
2000	23.8	21.9	13.8	6.7	3.9	2.9	2.7	3.5	4.3	7.3
1999	24.2	21.9	12.2	7.1	4.6	3.7	3.1	3.4	4.4	6.0
All races 3/, female										
2002	15.8	15.6	10.2	6.6	4.5	3.5	3.2	3.3	3.7	4.8
2001	12.6	11.6	8.3	5.1	3.9	2.6	2.6	3.1	3.4	4.9
2000	11.4	9.8	7.0	5.2	3.4	2.3	2.7	2.8	3.7	4.8
1999	10.6	9.2	6.6	4.6	2.9	2.7	2.6	3.6	3.9	4.3
White, both sexes										
2002	22.6	20.8	13.8	7.9	4.9	3.8	3.4	3.6	4.3	5.7
2001	18.7	16.9	10.8	6.1	3.9	3.1	3.0	3.5	3.7	5.0
2000	17.3	15.0	9.4	5.7	3.6	2.5	2.8	3.3	4.1	5.5
1999	16.8	14.5	8.6	5.5	3.5	3.1	2.9	3.7	4.3	4.7
White, male										
2002	29.0	25.9	17.3	9.0	4.9	3.8	3.4	3.9	4.5	7.6
2001	24.7	22.3	13.3	6.9	3.6	3.6	3.2	3.6	4.0	5.5
2000	23.3	20.4	12.1	5.9	3.6	2.7	2.6	3.7	4.4	7.0
1999	23.5	20.0	10.6	6.2	4.0	3.5	3.1	3.4	4.6	6.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Drug-induced causes (F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5, F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,X40-X44,X60-X64,X85,Y10-Y14) 5/										
White, female										
2002	16.3	15.8	10.4	6.9	4.9	3.9	3.5	3.4	4.1	4.8
2001	12.7	11.6	8.4	5.3	4.1	2.6	2.8	3.5	3.4	4.8
2000	11.3	9.7	6.9	5.4	3.7	2.4	3.0	3.0	3.9	4.8
1999	10.1	9.1	6.6	4.8	3.1	2.8	2.7	3.8	4.2	4.1
Black, both sexes										
2002	24.1	28.2	25.0	13.6	8.8	3.7	*	*	*	*
2001	23.3	28.0	21.7	10.2	7.0	3.3	3.0	*	*	6.1
2000	23.9	26.0	21.0	9.6	4.8	3.6	*	*	*	7.5
1999	25.7	27.8	18.4	9.8	6.4	3.9	2.8	*	*	*
Black, male										
2002	31.1	38.7	39.9	22.8	16.0	7.0	*	*	*	*
2001	31.7	43.3	34.4	17.1	12.6	5.3	*	*	*	*
2000	33.8	40.7	33.6	16.0	7.9	6.0	*	*	*	*
1999	36.8	44.3	30.2	17.6	11.9	6.5	*	*	*	*
Black, female										
2002	17.9	19.1	12.4	6.1	3.1	*	*	*	*	*
2001	15.9	14.8	10.8	4.6	*	*	*	*	*	*
2000	15.1	13.3	10.2	4.5	*	*	*	*	*	*
1999	15.9	13.5	8.4	3.3	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
All races 3/, both sexes										
2002	11.9	11.1	10.5	10.7	9.6	10.1	11.8	13.6	15.5	12.5
2001	11.3	10.6	10.4	10.4	9.7	10.5	11.4	13.7	15.3	12.8
2000	11.8	10.7	10.3	9.6	9.1	9.7	11.5	13.4	14.7	14.2
1999	10.9	10.7	9.6	9.9	9.4	10.0	12.2	14.0	14.5	13.5
All races 3/, male										
2002	19.7	18.4	18.0	18.6	17.1	18.8	24.1	29.8	37.6	38.9
2001	18.6	17.9	17.6	17.9	16.7	19.6	22.9	29.9	37.3	40.2
2000	19.4	17.9	17.3	16.5	16.1	17.8	23.1	29.6	36.6	44.7
1999	17.7	17.8	16.1	17.1	17.1	18.9	24.5	30.8	36.2	43.3
All races 3/, female										
2002	4.2	4.0	3.2	3.3	2.8	2.5	2.0	2.2	2.0	1.1
2001	4.1	3.6	3.5	3.3	3.2	2.6	2.2	2.3	2.2	1.3
2000	4.3	3.7	3.5	3.1	2.8	2.7	2.4	2.2	1.9	1.7
1999	4.2	3.7	3.4	3.2	2.4	2.4	2.5	2.3	1.9	1.5
White, both sexes										
2002	11.6	11.3	10.8	11.5	10.2	11.0	12.8	14.4	16.7	13.4
2001	11.0	10.9	10.9	10.9	10.5	11.3	12.1	14.6	16.1	13.8
2000	11.5	10.7	10.6	10.0	9.8	10.4	12.1	14.5	15.7	15.2
1999	10.6	10.6	10.0	10.4	9.8	10.7	12.9	14.9	15.4	14.5
White, male										
2002	19.1	18.4	18.5	19.7	18.1	20.3	25.8	31.2	40.3	41.7
2001	18.0	18.2	18.3	18.6	18.0	20.9	24.1	31.8	39.2	43.3
2000	18.7	17.8	17.6	17.0	17.0	18.9	24.1	31.9	38.8	48.4
1999	17.1	17.6	16.7	17.5	17.7	20.1	25.8	32.6	38.2	46.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
White, female										
2002	4.1	4.2	3.3	3.6	2.9	2.8	2.1	2.3	2.2	1.2
2001	4.0	3.7	3.7	3.6	3.6	2.9	2.3	2.4	2.2	1.4
2000	4.1	3.7	3.7	3.3	3.1	2.9	2.6	2.2	1.9	1.8
1999	4.1	3.8	3.4	3.6	2.5	2.5	2.7	2.4	1.9	1.6
Black, both sexes										
2002	16.6	13.0	10.3	7.3	7.0	5.1	6.7	8.8	*	*
2001	15.4	11.3	9.6	8.1	5.5	5.9	7.4	6.9	8.4	*
2000	16.5	13.2	10.6	7.4	5.8	6.2	7.7	4.6	6.8	*
1999	15.6	13.6	8.6	8.5	7.9	6.4	6.9	8.4	6.1	*
Black, male										
2002	29.1	24.1	18.1	14.4	12.3	10.2	14.7	20.7	*	*
2001	26.8	20.4	17.4	16.3	10.4	12.2	15.8	16.3	20.9	*
2000	28.5	23.4	19.4	13.7	11.2	12.7	18.1	9.5	16.9	*
1999	26.9	24.4	15.2	17.3	15.5	12.7	15.6	19.5	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget guidelines. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/										
Black, female										
2002.....	5.6	3.3	3.7	*	*	*	*	*	*	*
2001.....	5.4	3.5	2.9	*	*	*	*	*	*	*
2000.....	5.9	4.4	3.1	*	*	*	*	*	*	*
1999.....	5.6	4.3	3.1	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
... Category not applicable.
1/ Figures for age not stated included in "All ages" but not distributed among age groups.
2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
3/ Includes races other than white and black.
4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
5/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.