Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All origins 3/, both sexes												
2002	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2314.7	5556.9	14,828.3
2001	848.5	683.5	33.3	17.3	80.7	105.3	203.7	428.9	964.8	2353.4	5582.7	15,113.3
2000	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2399.1	5666.5	15,524.4
1999	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1005.0	2457.3	5714.5	15,554.6
All origins 3/, male												
2002	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1184.0	2855.3	6760.5	16,254.5
2001	846.5	749.8	37.0	19.8	117.0	143.7	259.7	545.1	1192.8	2911.7	6833.4	16,745.5
2000	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1230.7	2979.6	6972.6	17,501.4
1999	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1252.0	3073.7	7083.3	17,597.2
All origins 3/, female												
2002	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1864.7	4757.9	14,209.6
2001	850.5	614.1	29.5	14.6	42.6	66.0	148.2	316.9	754.1	1890.9	4760.7	14,430.3
2000	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1921.2	4814.7	14,719.2
1999	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1952.3	4825.4	14,731.3
Hispanic, both sexes												
2002	302.2	592.7	29.8	15.5	77.2	79.3	145.8	321.3	719.8	1734.1	4143.1	10,679.0
2001	307.2	573.2	30.7	14.8	75.6	84.7	155.0	334.7	723.4	1778.1	4269.4	11,834.3
2000	303.8	596.3	29.6	15.7	72.8	84.6	157.6	330.2	744.4	1803.8	4335.9	11,822.9
1999	305.7	583.8	30.9	15.6	71.3	85.5	158.2	333.2	752.5	1853.0	4356.3	12,177.9
Hispanic, male												
2002	328.7	644.0	34.2	17.4	114.4	112.5	192.5	423.4	937.4	2193.4	5043.5	11,674.1
2001	333.4	624.4	33.9	16.8	111.5	118.3	208.7	444.4	925.3	2245.2	5265.6	12,903.5
2000	331.3	637.1	31.5	17.9	107.7	120.2	211.0	439.0	965.7	2287.9	5395.3	13,086.2
1999	332.6	623.4	32.9	17.8	104.1	120.6	215.1	444.4	974.8	2368.9	5379.2	13,485.9
Hispanic, female												
2002	274.0	539.1	25.3	13.5	34.1	40.0	94.9	219.8	524.3	1368.7	3526.4	10,186.0
2001	279.4	519.9	27.3	12.7	33.7	45.3	97.2	226.9	544.2	1409.7	3596.1	11,311.7
2000	274.6	553.6	27.5	13.4	31.7	43.4	100.5	223.8	548.4	1423.2	3624.5	11,202.8
1999	277.2	542.3	28.7	13.3	32.9	44.9	97.4	224.9	555.8	1448.8	3675.7	11,547.3
Non-Hispanic 4/, both sexes												
2002	928.8	713.0	31.4	17.8	81.9	108.4	210.3	438.8	967.2	2347.0	5610.8	14,958.7
2001	926.2	703.5	33.8	17.7	81.4	108.9	209.2	435.6	979.5	2383.9	5628.5	15,199.0
2000	929.6	760.4	32.9	18.4	80.9	104.1	202.9	431.8	1006.5	2427.9	5707.6	15,619.8
1999	929.9	760.7	34.8	19.1	80.5	104.7	201.6	423.3	1019.2	2485.0	5753.4	15,631.0
Non-Hispanic 4/, male												
2002	928.0	782.6	35.4	20.4	117.3	148.6	266.0	557.0	1198.1	2889.1	6825.9	16,418.0
2001	923.6	773.2	37.6	20.4	117.5	148.5	265.2	551.6	1207.0	2944.1	6888.0	16,866.4
2000	928.1	836.8	36.8	21.5	115.9	141.6	259.2	548.7	1244.2	3009.8	7020.7	17,639.8
1999	932.2	834.9	38.8	22.1	114.6	142.7	258.1	537.3	1265.6	3102.8	7133.0	17,710.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Non-Hispanic 4/, female												
2002	929.5	640.1	27.2	15.0	45.3	68.5	155.7	325.2	753.0	1894.4	4805.4	14,329.1
2001	928.6	630.4	29.9	14.9	44.1	69.6	154.3	324.0	768.7	1918.4	4801.8	14,505.6
2000	931.0	680.1	28.8	15.2	45.0	66.8	147.7	319.2	786.8	1947.5	4852.4	14,802.9
1999	927.8	682.9	30.6	16.0	45.4	67.1	146.3	313.7	791.7	1977.5	4858.2	14,798.4
Non-Hispanic white, both sexes												
2002	997.5	575.9	27.1	16.0	75.8	95.9	191.6	400.1	912.8	2296.1	5596.0	15,149.1
2001	991.1	555.4	30.1	16.3	74.1	96.4	188.3	396.0	924.1	2325.5	5604.4	15,371.2
2000	993.2	596.5	28.5	17.1	73.6	90.1	181.0	390.5	950.2	2366.2	5675.4	15,799.7
1999	990.7	593.1	30.1	17.3	72.4	91.3	177.8	381.5	957.8	2420.0	5716.0	15,795.3
Non-Hispanic white, male												
2002	983.9	643.5	30.3	18.3	106.7	130.9	244.9	509.9	1126.5	2824.1	6801.7	16,641.9
2001	975.6	611.6	33.7	18.6	105.1	131.5	241.6	502.7	1136.3	2869.4	6851.5	17,055.4
2000	978.5	658.7	32.4	20.0	103.5	123.0	233.9	497.7	1170.9	2930.5	6977.8	17,853.2
1999	979.6	658.1	33.4	19.9	100.8	124.5	230.0	483.7	1187.4	3023.2	7088.0	17,871.2
Non-Hispanic white, female												
2002	1,010.6	504.8	23.8	13.6	43.8	60.3	138.3	292.1	710.5	1846.0	4787.9	14,504.3
2001	1,006.1	496.4	26.3	13.9	41.9	60.9	134.9	291.0	723.5	1864.1	4777.3	14,670.6
2000	1,007.3	530.9	24.4	13.9	42.6	56.8	128.1	285.0	742.1	1891.0	4819.3	14,971.7
1999	1,001.3	524.6	26.7	14.7	42.9	57.8	125.6	281.0	741.6	1915.1	4817.7	14,967.5
Non-Hispanic black, both sexes												
2002	792.8	1280.3	48.4	25.3	116.9	193.9	357.8	782.0	1592.9	3204.8	6584.8	14,386.7
2001	798.1	1354.4	49.5	24.0	121.4	191.5	365.8	786.9	1602.5	3300.7	6645.3	14,660.8
2000	805.5	1477.8	51.7	25.0	123.2	194.6	365.1	802.2	1633.9	3378.8	6764.3	14,837.5
1999	812.1	1510.2	54.6	27.5	125.5	191.1	376.9	796.7	1681.7	3454.5	6805.4	14,839.2
Non-Hispanic black, male												
2002	842.3	1369.3	55.8	29.8	177.4	274.3	445.7	1002.9	2074.5	4078.9	8246.2	15,776.9
2001	849.7	1484.2	55.1	28.6	186.2	267.9	456.3	1014.6	2055.4	4218.9	8426.8	16,576.2
2000	859.5	1623.2	56.1	29.1	187.2	270.2	462.7	1036.2	2111.4	4303.9	8551.0	16,944.8
1999	872.8	1650.8	61.5	33.1	191.4	267.9	479.5	1048.1	2176.7	4397.8	8615.3	17,385.3
Non-Hispanic black, female												
2002	748.0	1187.5	40.8	20.7	56.1	121.3	280.1	592.7	1206.3	2584.8	5644.9	13,855.4
2001	751.2	1217.8	43.6	19.3	56.8	122.4	285.7	591.5	1238.2	2653.0	5645.1	13,951.4
2000	756.7	1327.7	47.1	20.7	60.1	126.5	278.8	601.2	1249.4	2728.0	5758.6	14,054.9
1999	757.3	1365.4	47.4	21.8	60.5	122.1	286.4	581.7	1285.1	2789.2	5792.0	13,887.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)												
All origins 3/, both sexes												
2002	22.2	14.4	1.1	0.6	1.2	6.2	17.5	24.9	26.1	50.7	113.4	252.8
2001	21.2	14.0	1.5	0.5	1.3	6.7	17.6	22.9	25.4	48.0	104.9	246.3
2000	21.0	13.4	1.4	0.5	1.2	7.5	17.7	22.9	24.4	45.8	103.7	256.0
1999	21.5	14.8	1.5	0.6	1.2	8.4	18.7	22.1	24.1	48.0	105.1	267.9
All origins 3/, male												
2002	24.2	14.8	1.2	0.6	1.2	7.6	24.6	36.1	33.0	58.0	130.8	270.9
2001	23.3	14.7	1.6	0.5	1.4	8.5	25.2	33.2	31.9	54.5	122.3	269.7
2000	23.0	14.1	1.5	0.5	1.3	9.5	25.4	33.7	30.8	52.7	120.5	273.0
1999	23.7	17.0	1.8	0.7	1.3	11.2	27.1	32.2	30.4	55.8	123.3	290.1
All origins 3/, female												
2002	20.2	14.0	0.9	0.5	1.1	4.7	10.4	14.1	19.7	44.6	101.9	245.0
2001	19.2	13.2	1.3	0.5	1.3	4.8	10.2	13.0	19.4	42.5	93.5	236.5
2000	19.0	12.7	1.4	0.5	1.0	5.4	10.1	12.5	18.4	40.1	92.7	249.1
1999	19.3	12.5	1.2	0.6	1.2	5.6	10.3	12.4	18.4	41.7	93.3	258.9
Hispanic, both sexes												
2002	12.7	15.0	1.0	0.5	1.1	5.1	17.7	30.3	31.7	50.1	95.6	197.9
2001	12.6	13.2	1.1	0.5	1.1	5.5	18.7	29.7	32.2	48.1	93.4	213.2
2000	12.5	14.3	1.4	0.5	1.0	6.9	19.9	30.1	31.0	44.6	93.6	185.1
1999	13.3	13.0	1.3	0.6	1.0	7.9	22.7	28.5	29.5	53.6	97.7	226.6
Hispanic, male												
2002	15.7	16.4	*	0.6	1.1	6.9	26.0	45.9	42.3	56.2	107.5	214.2
2001	15.7	13.4	*	0.7	1.4	7.4	26.9	45.4	45.4	56.1	111.3	205.7
2000	16.0	14.4	1.6	*	1.3	9.5	29.7	45.6	42.8	53.6	110.1	209.6
1999	16.7	14.7	1.7	*	1.1	10.8	33.8	42.5	40.9	63.7	110.0	230.8
Hispanic, female												
2002	9.5	13.5	*	*	1.1	3.0	8.7	14.7	22.2	45.2	87.4	189.8
2001	9.4	13.0	*	*	0.7	3.4	9.8	14.2	20.6	41.7	81.3	216.9
2000	8.9	14.1	*	*	0.8	3.8	9.4	14.9	20.5	37.5	82.5	173.1
1999	9.8	11.1	*	*	1.0	4.5	10.9	14.8	19.5	45.6	89.5	224.6
Non-Hispanic 4/, both sexes												
2002	23.5	14.1	1.1	0.6	1.1	6.3	17.2	24.1	25.5	50.5	114.0	254.3
2001	22.3	14.1	1.6	0.5	1.4	6.9	17.3	22.0	24.7	47.7	105.2	246.9
2000	22.0	13.0	1.5	0.5	1.2	7.5	17.2	22.0	23.7	45.6	103.7	257.7
1999	22.5	15.2	1.6	0.6	1.3	8.5	17.9	21.2	23.5	47.5	105.0	268.3
Non-Hispanic 4/, male												
2002	25.5	14.1	1.2	0.6	1.2	7.7	24.1	34.7	32.0	57.8	131.5	272.8
2001	24.3	15.0	1.8	0.5	1.4	8.7	24.6	31.5	30.6	54.1	122.4	272.0
2000	23.9	13.9	1.5	0.5	1.3	9.4	24.5	32.2	29.7	52.2	120.2	274.5
1999	24.5	17.5	1.8	0.7	1.3	11.2	25.9	30.7	29.3	55.1	123.3	291.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)												
Non-Hispanic 4/, female												
2002	21.7	14.1	1.0	0.6	1.1	5.0	10.5	13.9	19.4	44.5	102.4	246.3
2001	20.5	13.1	1.3	0.5	1.4	5.0	10.1	12.8	19.2	42.5	93.9	236.5
2000	20.3	12.2	1.4	0.5	1.1	5.7	10.1	12.2	18.1	40.1	92.9	250.9
1999	20.5	12.8	1.3	0.6	1.3	5.7	10.1	12.1	18.2	41.3	93.2	259.1
Non-Hispanic white, both sexes												
2002	20.8	11.4	0.9	0.5	0.7	2.9	9.9	16.4	19.6	44.4	107.1	243.2
2001	19.5	9.5	1.4	0.4	0.8	3.1	9.9	14.6	18.9	41.9	98.7	235.4
2000	19.0	9.2	1.1	0.4	0.7	3.3	10.0	14.4	18.0	39.7	95.7	244.1
1999	19.2	11.3	1.3	0.5	0.8	4.3	10.2	13.7	17.8	40.9	96.6	251.8
Non-Hispanic white, male												
2002	21.6	11.2	1.1	0.5	0.8	4.0	14.8	23.8	24.3	50.4	122.3	261.4
2001	20.3	10.2	1.6	0.4	0.8	4.4	15.3	21.2	22.9	46.9	114.8	258.2
2000	19.6	9.8	1.1	0.4	0.9	4.7	15.0	21.2	22.0	45.3	110.4	259.6
1999	20.0	13.8	1.5	0.5	0.9	6.2	15.7	19.8	21.3	47.1	113.6	271.2
Non-Hispanic white, female												
2002	20.0	11.6	0.7	0.4	0.7	1.9	5.0	9.1	15.1	39.2	97.0	235.3
2001	18.6	8.8	1.2	0.4	0.8	1.8	4.5	8.1	15.1	37.6	88.0	225.9
2000	18.3	8.5	1.2	0.4	0.5	2.0	5.0	7.6	14.2	35.0	86.0	237.8
1999	18.4	8.7	1.1	0.4	0.6	2.3	4.6	7.7	14.4	35.8	85.4	244.1
Non-Hispanic black, both sexes												
2002	44.2	24.6	1.7	0.9	3.1	25.1	63.3	81.2	75.8	108.8	206.1	417.4
2001	43.6	32.7	2.3	0.9	4.2	27.3	64.0	77.7	74.8	103.8	191.7	409.2
2000	44.3	30.7	2.5	0.8	3.5	30.9	63.9	80.4	70.5	103.9	208.1	439.8
1999	45.7	32.2	2.6	1.3	3.6	32.3	68.2	79.1	71.6	111.2	209.5	470.8
Non-Hispanic black, male												
2002	53.1	24.8	1.6	0.7	3.0	29.2	86.5	120.9	104.3	137.5	259.1	454.0
2001	52.7	35.4	2.8	0.9	4.1	33.9	88.4	114.4	103.3	129.5	233.5	478.2
2000	54.0	32.8	2.9	0.8	3.6	37.5	89.8	122.4	99.3	128.5	259.2	488.5
1999	56.0	34.3	3.3	1.4	3.2	41.4	96.5	121.0	101.0	140.5	254.2	536.0
Non-Hispanic black, female												
2002	36.1	24.3	1.8	1.0	3.3	21.3	42.7	47.2	52.8	88.4	176.1	403.4
2001	35.3	29.9	1.8	0.9	4.4	21.3	42.4	46.3	51.9	85.7	168.3	383.7
2000	35.5	28.5	2.2	0.8	3.5	24.9	40.9	44.3	47.3	86.7	179.4	421.7
1999	36.4	30.0	2.0	1.1	4.1	24.1	43.1	43.4	47.9	90.4	184.5	446.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intestinal infectious diseases (A00-A09)												
All origins 3/, both sexes												
2002	0.9	*	*	*	*	*	0.0	0.2	0.5	2.0	7.1	21.0
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.6	13.1
2000	0.5	*	*	*	*	*	0.0	0.1	0.3	1.3	3.9	11.9
1999	0.4	0.6	*	*	*	*	*	0.1	0.2	1.0	3.3	10.6
All origins 3/, male												
2002	0.7	*	*	*	*	*	*	0.2	0.6	2.3	7.1	20.4
2001	0.4	*	*	*	*	*	*	0.1	0.3	1.4	4.7	12.2
2000	0.4	*	*	*	*	*	*	0.1	0.4	1.3	4.1	11.5
1999	0.3	*	*	*	*	*	*	*	0.3	1.0	3.6	9.7
All origins 3/, female												
2002	1.0	*	*	*	*	*	*	0.1	0.5	1.8	7.0	21.3
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.6	13.4
2000	0.6	*	*	*	*	*	*	*	0.2	1.2	3.7	12.0
1999	0.5	*	*	*	*	*	*	*	0.2	1.0	3.1	10.9
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	5.2	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.0	*	*	*	*	*	0.1	0.2	0.6	2.1	7.1	21.5
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.7	13.1
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.3	3.9	12.0
1999	0.5	*	*	*	*	*	*	0.1	0.2	1.0	3.4	10.7
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	*	*	0.2	0.6	2.4	7.3	21.1
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.4	4.8	12.1
2000	0.4	*	*	*	*	*	*	0.1	0.3	1.3	4.2	11.6
1999	0.4	*	*	*	*	*	*	*	0.2	1.0	3.7	9.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intestinal infectious diseases (A00-A09)												
Non-Hispanic 4/, female												
2002	1.2	*	*	*	*	*	*	0.2	0.5	1.8	7.1	21.7
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.3	4.6	13.5
2000	0.6	*	*	*	*	*	*	*	0.2	1.2	3.7	12.2
1999	0.5	*	*	*	*	*	*	*	0.2	1.0	3.2	11.1
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	*	*	*	0.2	0.5	2.2	7.6	22.6
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.4	4.9	13.8
2000	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.2	12.5
1999	0.5	*	*	*	*	*	*	*	0.2	1.1	3.5	11.4
Non-Hispanic white, male												
2002	0.9	*	*	*	*	*	*	0.2	0.5	2.4	7.7	22.2
2001	0.5	*	*	*	*	*	*	*	0.3	1.4	4.9	12.7
2000	0.5	*	*	*	*	*	*	*	0.3	1.4	4.4	12.1
1999	0.4	*	*	*	*	*	*	*	0.2	1.1	3.8	10.3
Non-Hispanic white, female												
2002	1.4	*	*	*	*	*	*	0.2	0.5	2.0	7.5	22.7
2001	0.9	*	*	*	*	*	*	0.2	0.3	1.3	4.8	14.2
2000	0.7	*	*	*	*	*	*	*	0.3	1.3	4.0	12.6
1999	0.6	*	*	*	*	*	*	*	*	1.1	3.3	11.8
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	1.1	1.9	3.8	13.3
2001	0.3	*	*	*	*	*	*	*	*	1.4	3.2	6.6
2000	0.2	*	*	*	*	*	*	*	*	*	*	8.3
1999	0.2	*	*	*	*	*	*	*	*	*	3.0	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	3.1	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.8	13.4
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholera (A00)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholera (A00)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other intestinal infectious diseases (A01-A08)												
All origins 3/, both sexes												
2002	0.8	*	*	*	*	*	0.0	0.1	0.5	1.9	6.8	20.2
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.6
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.3
1999	0.4	0.6	*	*	*	*	*	0.1	0.2	0.9	3.0	9.9
All origins 3/, male												
2002	0.7	*	*	*	*	*	*	0.2	0.5	2.2	6.9	19.7
2001	0.4	*	*	*	*	*	*	0.1	0.3	1.3	4.5	11.9
2000	0.4	*	*	*	*	*	*	0.1	0.4	1.3	4.0	10.6
1999	0.3	*	*	*	*	*	*	*	0.2	0.9	3.4	9.3
All origins 3/, female												
2002	1.0	*	*	*	*	*	*	0.1	0.5	1.7	6.8	20.4
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.2	4.4	13.0
2000	0.6	*	*	*	*	*	*	*	0.2	1.1	3.6	11.5
1999	0.5	*	*	*	*	*	*	*	0.2	0.9	2.8	10.2
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	5.0	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	*	*	*	*	*	0.1	0.1	0.5	2.0	6.9	20.6
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.7
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.8	11.4
1999	0.4	*	*	*	*	*	*	0.1	0.2	0.9	3.1	10.0
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	*	0.2	0.5	2.2	7.0	20.3
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.4	4.6	11.8
2000	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	10.6
1999	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other intestinal infectious diseases (A01-A08)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	*	*	0.1	0.5	1.8	6.8	20.8
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.2	4.4	13.0
2000	0.6	*	*	*	*	*	*	*	0.2	1.1	3.6	11.7
1999	0.5	*	*	*	*	*	*	*	*	0.9	2.8	10.3
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	*	*	*	0.2	0.5	2.1	7.4	21.6
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.7	13.3
2000	0.6	*	*	*	*	*	*	0.1	0.3	1.2	4.0	11.8
1999	0.5	*	*	*	*	*	*	*	0.2	0.9	3.2	10.6
Non-Hispanic white, male												
2002	0.8	*	*	*	*	*	*	0.2	0.5	2.3	7.5	21.3
2001	0.5	*	*	*	*	*	*	*	0.3	1.4	4.7	12.3
2000	0.5	*	*	*	*	*	*	*	0.3	1.3	4.2	11.1
1999	0.4	*	*	*	*	*	*	*	0.2	0.9	3.7	9.9
Non-Hispanic white, female												
2002	1.3	*	*	*	*	*	*	0.1	0.5	1.9	7.3	21.7
2001	0.8	*	*	*	*	*	*	0.1	0.3	1.3	4.7	13.7
2000	0.7	*	*	*	*	*	*	*	0.2	1.2	3.9	12.1
1999	0.6	*	*	*	*	*	*	*	*	1.0	2.9	10.9
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	1.1	1.8	3.5	13.3
2001	0.3	*	*	*	*	*	*	*	*	1.3	3.2	6.6
2000	0.2	*	*	*	*	*	*	*	*	*	*	7.6
1999	0.2	*	*	*	*	*	*	*	*	*	2.7	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.5	13.4
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Typhoid fever (A01.0)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Typhoid fever (A01.0)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis (A03)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis (A03)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other bacterial food-borne intoxications (A05)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other bacterial food-borne intoxications (A05)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Amebiasis (A06)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Amebiasis (A06)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intestinal infections due to other specified organisms (A04,A07-A08)												
All origins 3/, both sexes												
2002	0.8	*	*	*	*	*	*	0.1	0.5	1.9	6.8	20.2
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.2	4.3	12.4
2000	0.4	*	*	*	*	*	*	0.1	0.3	1.1	3.6	11.1
1999	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.9	9.7
All origins 3/, male												
2002	0.6	*	*	*	*	*	*	0.1	0.5	2.1	6.8	19.6
2001	0.4	*	*	*	*	*	*	*	0.3	1.3	4.4	11.8
2000	0.4	*	*	*	*	*	*	*	0.3	1.2	3.9	10.5
1999	0.3	*	*	*	*	*	*	*	0.2	0.8	3.3	9.1
All origins 3/, female												
2002	1.0	*	*	*	*	*	*	0.1	0.5	1.7	6.7	20.4
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.2	4.3	12.7
2000	0.5	*	*	*	*	*	*	*	0.2	1.1	3.4	11.4
1999	0.4	*	*	*	*	*	*	*	0.2	0.9	2.7	9.9
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	5.0	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	*	*	*	*	*	*	0.1	0.5	2.0	6.8	20.6
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.4	12.4
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.1	3.7	11.3
1999	0.4	*	*	*	*	*	*	*	0.2	0.9	3.0	9.7
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	*	0.1	0.5	2.2	6.9	20.2
2001	0.4	*	*	*	*	*	*	*	0.3	1.3	4.5	11.7
2000	0.4	*	*	*	*	*	*	*	0.3	1.2	3.9	10.5
1999	0.3	*	*	*	*	*	*	*	0.2	0.8	3.4	9.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intestinal infections due to other specified organisms (A04,A07-A08)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	*	*	0.1	0.5	1.8	6.8	20.7
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.2	4.3	12.7
2000	0.6	*	*	*	*	*	*	*	0.2	1.1	3.5	11.5
1999	0.5	*	*	*	*	*	*	*	*	0.9	2.7	10.0
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	*	*	*	0.1	0.5	2.1	7.3	21.6
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000	0.6	*	*	*	*	*	*	0.1	0.3	1.2	3.9	11.7
1999	0.5	*	*	*	*	*	*	*	0.2	0.9	3.1	10.3
Non-Hispanic white, male												
2002	0.8	*	*	*	*	*	*	0.1	0.5	2.3	7.3	21.2
2001	0.5	*	*	*	*	*	*	*	0.3	1.3	4.5	12.3
2000	0.4	*	*	*	*	*	*	*	0.3	1.2	4.1	11.0
1999	0.4	*	*	*	*	*	*	*	*	0.9	3.6	9.6
Non-Hispanic white, female												
2002	1.3	*	*	*	*	*	*	*	0.5	1.9	7.3	21.7
2001	0.8	*	*	*	*	*	*	0.1	0.3	1.3	4.6	13.4
2000	0.7	*	*	*	*	*	*	*	0.2	1.1	3.8	11.9
1999	0.6	*	*	*	*	*	*	*	*	0.9	2.8	10.6
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	1.1	1.8	3.4	13.3
2001	0.3	*	*	*	*	*	*	*	*	1.3	3.0	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	7.6
1999	0.2	*	*	*	*	*	*	*	*	*	2.5	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.3	13.4
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diarrhea and gastroenteritis of infectious origin (A09)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.6
1999	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.7
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.9
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	0.8
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.7
1999	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.7
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diarrhea and gastroenteritis of infectious origin (A09)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.9
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.0
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.7
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.0
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
All origins 3/, male												
2002	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	1.8	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	3.6	*
2000	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999	0.4	*	*	*	*	*	*	*	1.2	2.3	5.6	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.8	2.5
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.7
2000	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.9	2.5
1999	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9	3.9
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	0.1	0.3	0.6	1.2	2.3	3.8
2001	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	1.9	4.3
2000	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.6	4.3
1999	0.4	*	*	*	*	*	0.2	0.4	0.8	1.3	2.6	6.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	1.9
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.1
2000	0.2	*	*	*	*	*	*	*	*	0.7	1.4	1.8
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.8
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.0
2000	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.8	1.7	2.9
2001	0.2	*	*	*	*	*	*	0.1	0.4	0.6	1.4	2.9
2000	0.2	*	*	*	*	*	*	0.2	*	0.6	1.5	3.6
1999	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.8	4.9
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	1.1	1.4
2001	0.2	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000	0.1	*	*	*	*	*	*	*	*	0.3	1.1	1.4
1999	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.2
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	0.4	0.8	1.2	2.6	3.9	*
2001	0.6	*	*	*	*	*	*	0.8	1.5	2.7	3.8	6.9
2000	0.7	*	*	*	*	*	0.5	1.0	1.4	3.2	6.4	*
1999	0.7	*	*	*	*	*	0.4	1.0	1.7	2.6	5.6	8.4
Non-Hispanic black, male												
2002	0.7	*	*	*	*	*	*	1.1	2.0	3.7	6.4	*
2001	0.7	*	*	*	*	*	*	1.4	2.4	4.4	*	*
2000	0.9	*	*	*	*	*	*	1.9	2.5	4.2	11.7	*
1999	0.9	*	*	*	*	*	*	1.8	3.0	4.2	7.8	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	3.5	*
2000	0.4	*	*	*	*	*	*	*	*	2.5	3.5	*
1999	0.5	*	*	*	*	*	*	*	*	*	4.4	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7
All origins 3/, male												
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	5.0	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.5	2.2
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.4
2000	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.6	2.2
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.5
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.3
2001	0.3	*	*	*	*	*	*	0.2	0.6	0.8	1.6	3.8
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.2	3.8
1999	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	2.3	6.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.7
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.2	1.5
1999	0.2	*	*	*	*	*	*	*	*	0.4	1.2	2.4
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.6
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
2000	0.1	*	*	*	*	*	*	0.1	*	0.3	1.1	1.8
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.2	2.7
2001	0.2	*	*	*	*	*	*	*	0.4	0.4	1.1	2.4
2000	0.2	*	*	*	*	*	*	0.2	*	0.5	1.3	3.2
1999	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.6	4.8
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.9
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.6	1.0	1.9	3.6	*
2001	0.4	*	*	*	*	*	*	0.6	1.2	2.0	3.0	6.2
2000	0.5	*	*	*	*	*	0.4	0.8	0.9	2.7	5.1	*
1999	0.6	*	*	*	*	*	*	0.7	1.5	2.0	4.8	7.1
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	1.0	1.8	*	6.2	*
2001	0.6	*	*	*	*	*	*	1.2	2.1	3.4	*	*
2000	0.8	*	*	*	*	*	*	1.5	2.1	3.9	9.3	*
1999	0.7	*	*	*	*	*	*	1.3	2.7	3.2	6.9	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	3.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	*
2000	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	0.1	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	0.1	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)												
All origins 3/, both sexes												
2002	12.3	8.6	0.6	0.3	0.4	0.9	2.1	5.5	13.3	36.2	90.2	210.5
2001	11.9	8.9	0.9	0.3	0.5	0.8	2.0	5.4	13.0	34.3	85.9	213.7
2000	11.6	8.6	0.8	0.2	0.4	0.8	2.1	5.3	12.6	32.3	83.8	222.5
1999	11.7	9.1	0.8	0.3	0.5	0.8	2.0	5.0	12.1	33.2	83.9	231.2
All origins 3/, male												
2002	11.1	8.6	0.7	0.3	0.4	0.9	2.2	6.1	14.4	39.7	103.7	222.5
2001	10.8	9.5	0.9	0.3	0.5	0.8	2.2	5.9	14.1	38.4	99.9	233.5
2000	10.4	8.9	0.9	0.2	0.5	0.9	2.2	6.1	13.7	36.1	96.0	233.1
1999	10.5	10.4	0.9	0.3	0.5	0.9	2.3	5.6	13.3	37.6	97.5	246.7
All origins 3/, female												
2002	13.5	8.5	0.6	0.3	0.4	0.8	2.0	4.9	12.3	33.3	81.3	205.3
2001	12.9	8.3	0.8	0.2	0.5	0.8	1.8	4.9	12.0	30.8	76.7	205.4
2000	12.9	8.3	0.7	0.2	0.4	0.7	2.0	4.5	11.5	29.2	75.9	218.2
1999	12.9	7.7	0.7	0.3	0.4	0.8	1.8	4.5	11.1	29.7	75.0	225.0
Hispanic, both sexes												
2002	4.2	9.4	*	*	0.3	0.6	1.3	4.1	8.8	29.7	66.1	163.5
2001	4.0	8.3	0.8	0.3	*	0.5	1.4	4.2	8.8	26.1	66.0	178.8
2000	3.8	10.2	0.7	*	0.3	0.6	1.6	4.2	9.2	22.6	62.6	148.6
1999	4.0	7.1	*	*	*	0.6	1.7	3.6	9.1	27.6	65.0	192.2
Hispanic, male												
2002	4.0	9.6	*	*	*	0.7	1.7	5.5	9.5	31.5	74.7	172.3
2001	3.9	8.6	*	*	*	*	1.6	5.3	10.7	30.8	80.5	165.7
2000	3.6	9.9	*	*	*	0.7	1.8	5.2	10.5	27.0	72.8	159.2
1999	3.8	8.3	*	*	*	0.6	2.1	4.4	11.5	30.8	76.6	194.1
Hispanic, female												
2002	4.4	9.1	*	*	*	*	0.8	2.7	8.2	28.3	60.1	159.2
2001	4.2	8.0	*	*	*	*	1.3	3.2	7.1	22.5	56.2	185.2
2000	3.9	10.6	*	*	*	*	1.4	3.2	8.0	19.1	55.8	143.4
1999	4.2	6.0	*	*	*	*	1.2	2.8	7.0	25.1	57.2	191.3
Non-Hispanic 4/, both sexes												
2002	13.5	8.2	0.6	0.3	0.4	0.9	2.2	5.7	13.6	36.5	91.1	211.7
2001	13.0	9.0	0.9	0.2	0.5	0.9	2.1	5.5	13.2	34.7	86.6	214.5
2000	12.7	8.1	0.8	0.2	0.4	0.8	2.1	5.3	12.8	32.8	84.4	224.4
1999	12.7	9.5	0.8	0.3	0.5	0.9	2.1	5.1	12.3	33.5	84.3	231.8
Non-Hispanic 4/, male												
2002	12.3	8.1	0.6	0.4	0.4	1.0	2.3	6.2	14.7	40.0	104.7	224.2
2001	11.9	9.7	1.0	0.2	0.6	0.9	2.2	5.9	14.2	38.7	100.5	236.2
2000	11.3	8.6	0.9	0.2	0.5	0.9	2.2	6.1	13.8	36.5	96.5	235.2
1999	11.4	10.9	0.9	0.3	0.5	0.9	2.3	5.6	13.3	37.9	97.9	247.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)												
Non-Hispanic 4/, female												
2002	14.8	8.3	0.6	0.3	0.4	0.9	2.1	5.2	12.6	33.5	82.1	206.4
2001	14.1	8.4	0.7	0.2	0.5	0.9	1.9	5.1	12.3	31.3	77.4	205.6
2000	14.0	7.6	0.8	0.2	0.4	0.8	2.1	4.6	11.7	29.8	76.5	220.0
1999	14.0	8.1	0.8	0.3	0.5	0.8	1.9	4.6	11.4	29.8	75.6	225.4
Non-Hispanic white, both sexes												
2002	13.5	6.6	0.6	0.3	0.4	0.7	1.8	4.6	11.3	32.5	85.0	200.1
2001	12.9	6.1	0.8	0.2	0.4	0.6	1.6	4.3	11.0	30.9	81.0	203.3
2000	12.5	5.9	0.7	0.2	0.4	0.6	1.7	4.1	10.6	28.9	77.5	211.0
1999	12.4	7.1	0.8	0.3	0.4	0.7	1.6	4.0	10.1	29.2	77.1	216.2
Non-Hispanic white, male												
2002	12.1	6.4	0.7	0.4	0.4	0.8	1.8	4.9	12.5	35.7	96.8	212.8
2001	11.6	6.8	1.0	0.2	0.5	0.6	1.8	4.6	11.8	34.2	94.1	223.0
2000	11.0	6.1	0.8	0.2	0.5	0.6	1.8	4.7	11.6	32.4	88.8	221.2
1999	11.0	8.4	0.8	0.2	0.5	0.7	1.7	4.3	10.6	32.8	89.6	230.9
Non-Hispanic white, female												
2002	14.8	6.8	0.5	0.2	0.4	0.7	1.8	4.2	10.2	29.8	77.1	194.6
2001	14.2	5.3	0.6	0.2	0.4	0.7	1.4	4.0	10.3	28.1	72.4	195.1
2000	13.9	5.6	0.7	0.2	0.3	0.6	1.6	3.6	9.6	26.0	70.0	206.9
1999	13.9	5.8	0.7	0.3	0.4	0.6	1.4	3.6	9.7	26.1	68.8	210.3
Non-Hispanic black, both sexes												
2002	17.5	14.5	0.9	0.5	0.6	2.1	5.1	13.9	35.1	78.5	177.5	380.2
2001	17.1	21.3	1.2	*	1.0	2.1	5.0	14.4	34.3	75.3	166.8	377.1
2000	17.4	18.7	1.3	0.3	0.8	2.2	5.2	14.7	32.9	74.9	179.7	405.4
1999	17.6	20.0	0.9	0.3	0.8	2.0	5.5	14.0	32.2	78.9	178.9	430.0
Non-Hispanic black, male												
2002	16.2	14.8	*	*	*	2.0	5.5	15.9	38.2	90.6	220.4	408.0
2001	16.1	22.0	*	*	1.1	2.1	5.2	16.0	39.4	90.2	202.2	442.4
2000	16.1	20.5	*	*	0.9	2.6	5.3	18.0	37.0	84.7	214.4	445.0
1999	16.8	21.5	*	*	0.8	2.2	6.5	16.5	39.7	96.2	216.4	470.6
Non-Hispanic black, female												
2002	18.6	14.2	*	0.7	*	2.2	4.7	12.1	32.6	69.9	153.2	369.6
2001	17.9	20.5	*	*	0.9	2.1	4.9	13.0	30.3	64.8	146.9	352.9
2000	18.5	16.9	*	*	*	1.9	5.2	11.9	29.5	67.9	160.1	390.7
1999	18.3	18.6	*	*	0.8	1.9	4.6	12.0	26.2	66.7	157.9	414.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Plague (A20)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Plague (A20)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Brucellosis (A23)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Brucellosis (A23)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tetanus (A33-A35)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tetanus (A33-A35)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diphtheria (A36)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diphtheria (A36)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
All origins 3/, both sexes												
2002	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.1	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	0.1	*	*	*	*
1999	0.1	0.8	0.2	0.1	0.1	*	0.1	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.1	1.0	0.3	*	0.1	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
All origins 3/, both sexes												
2002	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
All origins 3/, male												
2002	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
All origins 3/, female												
2002	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5
Hispanic, both sexes												
2002	3.9	7.8	*	*	*	0.5	1.2	3.7	8.4	28.9	63.7	156.3
2001	3.8	6.1	*	*	*	0.3	1.3	3.9	8.2	25.4	62.6	171.3
2000	3.5	8.0	*	*	*	0.5	1.5	3.9	8.5	21.8	59.3	144.7
1999	3.6	5.7	*	*	*	0.4	1.5	3.1	8.0	25.6	61.7	180.3
Hispanic, male												
2002	3.7	8.0	*	*	*	0.6	1.6	5.2	9.0	30.8	72.2	164.0
2001	3.6	7.0	*	*	*	*	1.5	4.7	9.9	29.8	76.5	160.0
2000	3.4	8.4	*	*	*	0.6	1.7	5.0	10.0	26.6	70.3	157.2
1999	3.5	7.0	*	*	*	*	2.0	3.8	10.2	29.4	75.1	181.2
Hispanic, female												
2002	4.1	7.6	*	*	*	*	0.7	2.3	7.8	27.4	57.9	152.5
2001	3.9	5.2	*	*	*	*	1.1	3.1	6.6	21.9	53.2	176.9
2000	3.5	7.7	*	*	*	*	1.3	2.8	7.2	18.1	51.8	138.5
1999	3.7	*	*	*	*	*	1.0	2.4	6.1	22.5	52.7	179.9
Non-Hispanic 4/, both sexes												
2002	12.9	7.1	0.5	0.2	0.3	0.8	2.0	5.3	12.9	34.9	87.3	204.2
2001	12.4	8.1	0.7	0.1	0.4	0.8	1.9	5.1	12.5	33.1	83.0	206.8
2000	12.1	6.9	0.7	0.2	0.3	0.7	2.0	5.0	12.1	31.4	81.0	217.5
1999	12.0	7.9	0.6	0.2	0.3	0.7	1.8	4.7	11.6	31.4	79.8	221.2
Non-Hispanic 4/, male												
2002	11.6	7.3	0.5	0.3	0.3	0.9	2.0	5.7	14.0	38.2	99.8	215.6
2001	11.2	8.8	0.9	0.1	0.4	0.8	2.1	5.5	13.4	37.0	96.1	226.8
2000	10.7	6.7	0.7	0.2	0.3	0.8	2.0	5.6	13.1	34.8	92.3	224.8
1999	10.7	9.4	0.6	0.2	0.4	0.8	2.1	5.2	12.5	35.4	92.2	234.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Non-Hispanic 4/, female												
2002	14.1	6.8	0.5	0.2	0.3	0.8	2.0	4.9	11.9	32.1	79.0	199.3
2001	13.5	7.4	0.6	0.2	0.3	0.8	1.7	4.8	11.8	29.9	74.4	198.5
2000	13.5	7.0	0.6	0.1	0.2	0.7	1.9	4.3	11.2	28.7	73.6	214.5
1999	13.2	6.2	0.5	0.2	0.3	0.7	1.6	4.2	10.8	28.2	71.8	215.9
Non-Hispanic white, both sexes												
2002	12.9	5.5	0.5	0.2	0.3	0.7	1.6	4.3	10.7	31.0	81.3	192.6
2001	12.3	5.1	0.7	0.1	0.3	0.6	1.4	4.0	10.4	29.4	77.6	195.6
2000	11.9	4.8	0.5	0.2	0.2	0.5	1.5	3.8	10.0	27.7	74.2	204.3
1999	11.7	5.3	0.5	0.2	0.3	0.6	1.4	3.6	9.5	27.3	73.0	206.0
Non-Hispanic white, male												
2002	11.5	5.6	0.5	0.3	0.3	0.7	1.6	4.6	11.8	34.0	92.1	203.9
2001	11.0	5.8	0.9	*	0.3	0.5	1.7	4.2	11.1	32.5	89.7	213.6
2000	10.4	4.7	0.5	0.2	0.3	0.6	1.6	4.3	10.9	30.9	84.7	211.2
1999	10.2	6.7	0.5	0.2	0.3	0.7	1.5	4.0	9.9	30.5	84.6	218.1
Non-Hispanic white, female												
2002	14.2	5.4	*	0.2	0.3	0.6	1.7	3.9	9.7	28.5	74.0	187.7
2001	13.6	4.4	0.5	*	0.3	0.6	1.2	3.8	9.9	26.8	69.6	188.1
2000	13.4	4.8	0.6	*	*	0.5	1.5	3.3	9.2	24.9	67.2	201.6
1999	13.1	3.8	0.5	0.2	0.2	0.5	1.2	3.3	9.1	24.5	65.4	201.2
Non-Hispanic black, both sexes												
2002	16.8	13.1	*	0.3	0.4	2.0	4.6	13.2	33.8	76.0	172.9	370.8
2001	16.4	20.3	0.9	*	0.7	1.9	4.8	13.7	32.6	72.8	161.9	368.6
2000	16.7	16.6	1.1	*	0.6	2.1	4.8	13.9	31.8	72.5	175.1	395.9
1999	16.6	18.6	*	*	0.6	1.6	5.1	13.2	30.8	75.3	169.8	414.2
Non-Hispanic black, male												
2002	15.5	13.9	*	*	*	1.9	5.0	15.0	36.6	87.6	213.7	399.3
2001	15.4	21.4	*	*	0.9	1.9	4.9	15.2	37.4	86.9	196.8	430.8
2000	15.3	16.4	*	*	*	2.4	4.8	16.8	35.9	81.6	208.9	432.1
1999	15.7	20.8	*	*	*	1.7	6.0	15.4	37.9	91.5	202.6	451.6
Non-Hispanic black, female												
2002	18.0	12.3	*	*	*	2.0	4.2	11.6	31.6	67.8	149.9	359.9
2001	17.2	19.1	*	*	*	1.9	4.7	12.5	28.8	62.8	142.4	345.6
2000	17.9	16.9	*	*	*	1.8	4.9	11.5	28.5	66.1	156.1	382.5
1999	17.4	16.4	*	*	*	1.6	4.3	11.3	25.1	63.9	151.4	400.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	*	0.2	0.3	0.6	1.5	3.7	7.4
2001	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.4	3.5	7.5
2000	0.5	0.5	*	*	*	*	0.1	0.3	0.6	1.3	3.3	6.7
1999	0.6	0.6	*	*	*	0.1	0.2	0.4	0.7	2.0	4.3	10.3
All origins 3/, male												
2002	0.5	*	*	*	*	*	0.2	0.4	0.7	1.7	4.6	8.5
2001	0.5	*	*	*	*	*	0.1	0.4	0.8	1.7	4.3	8.9
2000	0.5	*	*	*	*	*	0.1	0.4	0.7	1.6	4.0	9.9
1999	0.6	*	*	*	*	0.1	0.2	0.4	0.8	2.4	5.4	13.1
All origins 3/, female												
2002	0.5	*	*	*	*	*	0.1	0.2	0.5	1.3	3.0	6.9
2001	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	2.9	6.9
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.8	5.3
1999	0.6	*	*	*	*	*	0.2	0.3	0.6	1.6	3.6	9.2
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	2.0	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	*	*	*	*	*	0.2	0.3	0.6	1.5	3.7	7.4
2001	0.5	*	*	*	*	0.1	0.1	0.3	0.6	1.5	3.5	7.5
2000	0.5	0.7	*	*	*	*	0.1	0.3	0.6	1.3	3.3	6.7
1999	0.7	0.7	*	*	*	0.1	0.2	0.3	0.7	2.0	4.4	10.3
Non-Hispanic 4/, male												
2002	0.6	*	*	*	*	*	0.2	0.4	0.7	1.8	4.7	8.5
2001	0.6	*	*	*	*	*	0.1	0.4	0.8	1.7	4.4	9.1
2000	0.5	*	*	*	*	*	0.1	0.4	0.7	1.6	4.0	10.2
1999	0.7	*	*	*	*	*	0.2	0.4	0.8	2.4	5.6	13.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	*	*	0.1	0.2	0.5	1.3	3.1	6.9
2001	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	2.9	6.8
2000	0.5	*	*	*	*	*	0.1	0.2	0.5	1.0	2.8	5.3
1999	0.7	*	*	*	*	*	0.2	0.3	0.5	1.6	3.6	9.1
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	3.6	7.3
2001	0.6	*	*	*	*	*	0.1	0.3	0.5	1.3	3.4	7.4
2000	0.5	*	*	*	*	*	0.1	0.3	0.5	1.2	3.2	6.5
1999	0.7	*	*	*	*	*	0.2	0.3	0.6	1.8	4.0	9.8
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	0.2	0.3	0.6	1.7	4.6	8.7
2001	0.6	*	*	*	*	*	*	0.4	0.7	1.6	4.4	9.1
2000	0.5	*	*	*	*	*	*	0.4	0.7	1.5	4.0	9.8
1999	0.7	*	*	*	*	*	0.2	0.3	0.8	2.3	5.0	12.6
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.2	2.7	6.7
2000	0.5	*	*	*	*	*	*	0.2	0.4	1.0	2.7	5.1
1999	0.7	*	*	*	*	*	0.2	0.3	0.5	1.5	3.3	8.7
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	0.4	0.6	1.2	2.5	4.6	9.4
2001	0.6	*	*	*	*	*	*	0.7	1.5	2.5	4.7	8.1
2000	0.5	*	*	*	*	*	*	0.5	1.1	2.2	4.6	9.5
1999	0.8	*	*	*	*	*	*	0.8	1.4	3.5	9.0	15.8
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	*	*	3.1	6.7	*
2001	0.6	*	*	*	*	*	*	*	1.8	3.3	*	*
2000	0.6	*	*	*	*	*	*	*	*	3.1	*	*
1999	0.9	*	*	*	*	*	*	*	*	4.7	13.4	*
Non-Hispanic black, female												
2002	0.6	*	*	*	*	*	*	*	*	2.1	3.3	9.6
2001	0.6	*	*	*	*	*	*	*	*	*	4.5	*
2000	0.5	*	*	*	*	*	*	*	*	*	4.0	*
1999	0.8	*	*	*	*	*	*	*	*	2.7	6.5	14.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections with a predominately sexual mode of transmission (A50-A64)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections with a predominately sexual mode of transmission (A50-A64)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cardiovascular syphilis (A52.0)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cardiovascular syphilis (A52.0)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neurosyphilis (A52.1-A52.3)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neurosyphilis (A52.1-A52.3)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gonococcal infection (A54)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gonococcal infection (A54)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other infections with a predominately sexual mode of transmission (A55-A64)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other infections with a predominately sexual mode of transmission (A55-A64)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lyme disease (A69.2)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lyme disease (A69.2)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Typhus fever (A75)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Typhus fever (A75)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral diseases (A80-B34)												
All origins 3/, both sexes												
2002	7.4	4.1	0.3	0.2	0.6	5.0	14.6	17.3	9.6	7.8	7.8	10.1
2001	7.4	3.0	0.4	0.2	0.7	5.6	15.0	16.0	9.8	7.9	7.2	8.3
2000	7.5	2.9	0.4	0.2	0.6	6.4	15.0	16.3	9.4	8.0	8.1	9.6
1999	7.6	3.5	0.5	0.3	0.6	7.1	15.6	15.1	8.8	8.1	8.3	11.6
All origins 3/, male												
2002	10.6	4.6	0.4	0.1	0.6	6.3	21.3	26.9	14.5	10.0	9.6	11.7
2001	10.5	3.1	0.5	0.2	0.7	7.4	22.1	25.2	14.5	9.6	8.5	9.3
2000	10.8	2.9	0.4	0.2	0.7	8.3	22.3	25.8	14.3	9.9	10.4	10.8
1999	10.9	4.0	0.6	0.3	0.6	9.8	23.5	24.1	13.3	10.3	9.9	13.3
All origins 3/, female												
2002	4.4	3.6	*	0.2	0.6	3.6	7.9	7.9	5.1	6.0	6.5	9.3
2001	4.3	2.8	0.3	0.2	0.8	3.7	7.9	7.1	5.3	6.4	6.4	7.9
2000	4.4	2.9	0.4	0.2	0.6	4.5	7.7	7.1	5.0	6.4	6.6	9.1
1999	4.3	3.1	0.4	0.2	0.6	4.4	7.8	6.5	4.7	6.4	7.2	10.9
Hispanic, both sexes												
2002	7.0	4.0	*	*	0.6	4.0	15.3	22.9	18.1	13.0	13.2	*
2001	7.4	3.2	*	*	0.6	4.6	16.3	23.5	20.3	15.1	13.7	13.1
2000	7.7	*	*	*	0.5	5.9	17.5	24.2	18.5	16.3	17.4	*
1999	8.0	3.2	*	*	0.7	6.8	19.7	22.6	16.6	18.0	15.6	*
Hispanic, male												
2002	10.0	5.6	*	*	0.7	5.6	22.7	35.3	26.7	15.5	14.0	*
2001	10.4	*	*	*	0.7	6.2	24.0	37.3	29.7	16.8	15.7	*
2000	11.0	*	*	*	0.7	8.3	26.7	38.6	27.5	19.0	19.7	*
1999	11.4	*	*	*	0.7	9.6	29.8	35.2	25.5	23.1	13.0	*
Hispanic, female												
2002	3.9	*	*	*	*	2.1	7.3	10.5	10.4	10.9	12.6	*
2001	4.2	*	*	*	*	2.8	8.0	9.8	11.9	13.7	12.4	*
2000	4.1	*	*	*	*	3.1	7.6	10.2	10.6	14.3	15.8	*
1999	4.4	*	*	*	*	3.5	8.9	10.3	8.6	14.1	17.3	*
Non-Hispanic 4/, both sexes												
2002	7.4	4.0	0.4	0.2	0.6	5.1	14.3	16.5	8.8	7.4	7.5	10.1
2001	7.3	2.9	0.5	0.2	0.7	5.7	14.6	15.1	8.8	7.3	6.9	8.1
2000	7.4	3.1	0.4	0.2	0.6	6.5	14.5	15.4	8.6	7.4	7.7	9.6
1999	7.4	3.6	0.5	0.3	0.6	7.1	14.9	14.2	8.1	7.5	8.0	11.7
Non-Hispanic 4/, male												
2002	10.6	4.2	0.5	0.1	0.6	6.4	20.8	25.7	13.4	9.6	9.4	11.7
2001	10.4	3.0	0.6	0.2	0.7	7.6	21.6	23.7	13.1	9.1	8.2	9.1
2000	10.6	3.2	0.4	0.2	0.7	8.2	21.5	24.3	13.1	9.1	10.0	10.6
1999	10.7	4.1	0.6	0.3	0.6	9.8	22.4	22.7	12.2	9.4	9.7	13.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral diseases (A80-B34)												
Non-Hispanic 4/, female												
2002	4.4	3.8	*	0.2	0.6	3.9	7.9	7.6	4.6	5.6	6.2	9.3
2001	4.3	2.8	0.4	0.2	0.8	3.8	7.8	6.8	4.8	5.9	6.1	7.7
2000	4.4	3.0	0.4	0.2	0.6	4.7	7.6	6.8	4.5	5.9	6.2	9.1
1999	4.3	3.1	0.4	0.2	0.6	4.5	7.5	6.1	4.4	5.9	6.8	11.0
Non-Hispanic white, both sexes												
2002	4.7	3.5	0.3	0.1	0.2	2.0	7.5	10.1	5.7	5.6	6.8	10.3
2001	4.5	1.9	0.4	0.1	0.3	2.3	7.8	9.0	5.6	5.7	6.3	8.2
2000	4.6	2.1	0.2	0.2	0.2	2.6	7.9	9.2	5.5	5.8	7.2	9.6
1999	4.6	2.7	0.4	0.1	0.2	3.3	8.0	8.3	5.1	5.8	7.5	11.7
Non-Hispanic white, male												
2002	6.9	3.4	*	*	0.3	3.1	12.2	16.4	8.6	7.1	8.5	11.7
2001	6.7	1.9	*	*	0.3	3.6	12.9	15.0	8.4	7.1	7.5	9.4
2000	6.8	2.1	*	*	0.3	3.9	12.7	15.1	8.2	6.8	9.3	10.7
1999	6.8	3.4	0.5	0.2	0.3	5.2	13.2	13.6	7.6	7.2	9.3	12.9
Non-Hispanic white, female												
2002	2.5	3.6	*	*	0.2	1.0	2.8	4.0	3.0	4.4	5.7	9.7
2001	2.3	2.0	*	*	0.3	1.0	2.8	3.2	3.0	4.5	5.5	7.7
2000	2.5	2.0	*	0.2	0.2	1.3	3.1	3.3	3.0	4.9	5.8	9.1
1999	2.5	2.0	*	*	0.2	1.4	2.8	3.1	2.8	4.6	6.3	11.2
Non-Hispanic black, both sexes												
2002	24.2	5.9	*	0.4	2.3	22.1	56.4	63.4	34.0	21.7	13.7	7.6
2001	24.3	6.9	0.9	0.5	3.1	24.4	57.5	60.2	35.0	19.1	11.8	6.2
2000	24.6	7.5	1.0	0.4	2.6	27.9	56.7	62.3	32.9	20.0	12.7	8.6
1999	24.8	7.2	1.2	0.8	2.5	28.8	59.6	59.8	32.2	20.9	11.2	11.3
Non-Hispanic black, male												
2002	33.9	6.7	*	*	2.3	26.1	78.6	99.4	56.2	34.8	20.5	*
2001	34.0	8.0	*	*	2.7	31.2	81.0	93.6	57.0	27.1	15.9	*
2000	35.1	7.5	*	*	2.5	34.0	82.1	99.3	55.3	31.9	19.1	*
1999	35.7	7.6	*	0.9	2.1	37.6	86.8	97.9	52.9	30.5	13.1	*
Non-Hispanic black, female												
2002	15.3	*	*	*	2.3	18.5	36.9	32.7	16.2	12.4	9.8	*
2001	15.5	*	*	*	3.4	18.2	36.6	31.6	17.3	13.5	9.4	*
2000	15.2	7.4	*	*	2.7	22.4	34.3	30.6	14.8	11.7	9.0	8.7
1999	15.0	*	*	0.7	2.9	20.8	35.7	27.2	15.7	14.1	10.2	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rabies (A82)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rabies (A82)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Yellow fever (A95)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Yellow fever (A95)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified arthropod-borne viral and hemorrhagic fevers (A90-A94,A96-A99)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.8	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified arthropod-borne viral and hemorrhagic fevers (A90-A94,A96-A99)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.9
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Herpes-viral (herpes simplex) infections (B00)												
All origins 3/, both sexes												
2002	0.1	0.6	*	*	*	*	*	*	*	0.2	0.4	0.5
2001	0.1	0.6	*	*	*	*	*	*	0.1	0.2	0.4	0.5
2000	0.1	0.6	*	*	*	*	*	*	*	0.1	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.5
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	0.7	*	*	*	*	*	*	*	0.2	0.4	0.6
2001	0.1	0.7	*	*	*	*	*	*	0.1	0.2	0.4	0.5
2000	0.1	0.7	*	*	*	*	*	*	*	0.1	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.5
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Herpes-viral (herpes simplex) infections (B00)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2001	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	0.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Zoster (herpes zoster)(B02)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.7
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	1.6
2000	0.0	*	*	*	*	*	*	*	*	*	0.3	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.0
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	1.8
2000	0.1	*	*	*	*	*	*	*	*	*	*	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.2	1.7
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	1.7
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.1
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Zoster (herpes zoster)(B02)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	1.9
2000	0.1	*	*	*	*	*	*	*	*	*	*	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.2	1.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	1.8
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
2000	0.1	*	*	*	*	*	*	*	*	*	*	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.6
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Smallpox (B03)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Smallpox (B03)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rubella (German measles) (B06)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rubella (German measles) (B06)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
All origins 3/, both sexes												
2002	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7
All origins 3/, male												
2002	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9
All origins 3/, female												
2002	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9
Hispanic, both sexes												
2002	2.0	*	*	*	*	*	1.7	8.0	8.3	8.6	10.5	*
2001	2.1	*	*	*	*	*	2.3	7.8	10.4	9.9	11.0	*
2000	2.0	*	*	*	*	*	2.5	7.0	8.6	11.1	13.0	*
1999	1.9	*	*	*	*	*	2.1	6.8	7.8	12.4	13.5	*
Hispanic, male												
2002	2.5	*	*	*	*	*	2.5	12.0	11.6	7.6	10.3	*
2001	2.6	*	*	*	*	*	3.1	12.2	14.2	7.0	10.3	*
2000	2.5	*	*	*	*	*	3.8	10.6	11.7	11.0	12.3	*
1999	2.3	*	*	*	*	*	3.3	10.5	9.8	13.0	*	*
Hispanic, female												
2002	1.4	*	*	*	*	*	0.9	4.0	5.4	9.4	10.7	*
2001	1.7	*	*	*	*	*	1.4	3.5	7.1	12.3	11.5	*
2000	1.5	*	*	*	*	*	1.0	3.3	5.8	11.3	13.5	*
1999	1.5	*	*	*	*	*	0.8	3.3	6.0	12.0	16.3	*
Non-Hispanic 4/, both sexes												
2002	2.0	*	*	*	*	0.2	1.6	5.6	3.6	3.8	4.3	3.7
2001	1.9	*	*	*	*	0.2	1.7	5.0	3.5	4.2	4.6	3.6
2000	1.9	*	*	*	*	0.2	1.6	4.8	3.3	4.2	5.0	3.8
1999	1.7	*	*	*	*	0.2	1.5	3.8	3.0	4.3	5.0	4.7
Non-Hispanic 4/, male												
2002	2.7	*	*	*	*	0.3	2.3	8.6	4.9	4.3	5.1	4.8
2001	2.5	*	*	*	*	0.2	2.4	7.7	4.6	4.6	5.0	4.3
2000	2.4	*	*	*	*	0.2	2.2	7.4	4.4	4.4	6.1	5.4
1999	2.2	*	*	*	*	0.2	2.2	5.9	3.9	4.7	6.4	6.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Non-Hispanic 4/, female												
2002	1.4	*	*	*	*	0.2	1.0	2.7	2.4	3.5	3.8	3.2
2001	1.4	*	*	*	*	0.2	1.0	2.3	2.5	3.8	4.3	3.3
2000	1.4	*	*	*	*	0.2	1.0	2.4	2.3	4.0	4.3	3.2
1999	1.2	*	*	*	*	0.2	0.9	1.8	2.1	4.0	4.0	3.9
Non-Hispanic white, both sexes												
2002	1.9	*	*	*	*	0.2	1.6	5.3	3.0	3.3	4.1	3.5
2001	1.9	*	*	*	*	0.2	1.7	4.6	2.9	3.7	4.3	3.3
2000	1.8	*	*	*	*	0.1	1.6	4.3	2.7	3.7	4.7	3.7
1999	1.6	*	*	*	*	0.2	1.4	3.4	2.5	3.7	4.7	4.5
Non-Hispanic white, male												
2002	2.6	*	*	*	*	0.3	2.3	8.2	4.1	3.9	4.9	4.6
2001	2.5	*	*	*	*	0.2	2.3	7.2	4.0	4.3	5.1	4.4
2000	2.3	*	*	*	*	*	2.1	6.7	3.6	3.8	5.9	5.4
1999	2.1	*	*	*	*	0.2	2.1	5.2	3.4	4.3	6.4	6.5
Non-Hispanic white, female												
2002	1.3	*	*	*	*	*	0.9	2.5	2.0	2.8	3.5	3.1
2001	1.2	*	*	*	*	*	1.0	2.0	2.0	3.1	3.8	2.9
2000	1.3	*	*	*	*	*	1.0	2.0	1.8	3.6	4.0	3.1
1999	1.1	*	*	*	*	0.2	0.8	1.7	1.6	3.3	3.6	3.7
Non-Hispanic black, both sexes												
2002	2.3	*	*	*	*	0.4	2.0	8.2	7.2	6.0	5.2	*
2001	2.3	*	*	*	*	*	2.2	8.1	6.8	5.6	5.7	*
2000	2.2	*	*	*	*	0.4	2.1	8.0	5.8	5.8	6.3	*
1999	2.1	*	*	*	*	*	2.2	7.0	5.1	6.8	5.9	*
Non-Hispanic black, male												
2002	3.1	*	*	*	*	*	2.6	12.6	10.7	7.0	*	*
2001	2.8	*	*	*	*	*	3.0	12.0	9.0	5.0	*	*
2000	2.9	*	*	*	*	*	3.0	12.3	8.1	7.2	7.7	*
1999	2.6	*	*	*	*	*	3.1	11.8	6.3	5.7	*	*
Non-Hispanic black, female												
2002	1.7	*	*	*	*	*	1.5	4.4	4.4	5.2	5.3	*
2001	1.8	*	*	*	*	*	1.4	4.7	5.0	6.0	6.7	*
2000	1.6	*	*	*	*	*	1.4	4.3	3.9	4.7	5.6	*
1999	1.6	*	*	*	*	*	1.4	2.9	4.1	7.5	6.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
All origins 3/, both sexes												
2002	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*
All origins 3/, male												
2002	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001	7.6	*	*	0.1	0.5	7.1	19.5	16.9	8.6	3.5	1.5	*
2000	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*
All origins 3/, female												
2002	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*
Hispanic, both sexes												
2002	4.8	*	*	*	0.4	3.6	13.4	14.7	9.5	3.4	*	*
2001	5.0	*	*	*	0.5	4.4	13.9	15.5	9.2	4.5	*	*
2000	5.4	*	*	*	0.3	5.6	14.9	16.9	9.5	3.8	*	*
1999	5.8	*	*	*	0.5	6.4	17.3	15.4	8.4	4.7	*	*
Hispanic, male												
2002	7.2	*	*	*	*	5.1	20.1	23.1	15.0	6.9	*	*
2001	7.6	*	*	*	0.6	6.1	20.8	24.9	14.8	8.6	*	*
2000	8.2	*	*	*	*	7.8	22.7	27.5	15.3	7.2	*	*
1999	8.7	*	*	*	0.6	9.3	26.2	24.2	15.4	9.0	*	*
Hispanic, female												
2002	2.2	*	*	*	*	1.8	6.2	6.3	4.6	*	*	*
2001	2.3	*	*	*	*	2.5	6.5	6.2	4.1	*	*	*
2000	2.4	*	*	*	*	3.0	6.4	6.6	4.4	*	*	*
1999	2.6	*	*	*	*	3.2	7.9	6.7	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	4.8	*	*	0.1	0.4	4.8	12.5	10.6	4.7	2.1	0.7	*
2001	4.9	*	*	0.1	0.6	5.4	12.7	9.9	4.8	1.9	0.6	*
2000	5.0	*	*	0.1	0.5	6.2	12.7	10.3	4.7	2.0	0.6	*
1999	5.2	*	*	0.2	0.5	6.8	13.2	10.0	4.5	2.0	0.6	*
Non-Hispanic 4/, male												
2002	7.3	*	*	*	0.4	6.0	18.4	16.9	7.9	3.7	1.3	*
2001	7.5	*	*	0.1	0.5	7.3	19.0	15.8	8.0	3.1	1.3	*
2000	7.7	*	*	*	0.5	7.9	19.1	16.7	8.0	3.4	1.2	*
1999	8.0	*	*	0.2	0.5	9.4	20.1	16.5	7.6	3.4	1.0	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic 4/, female												
2002	2.5	*	*	*	0.5	3.6	6.7	4.7	1.7	0.8	0.3	*
2001	2.5	*	*	*	0.7	3.6	6.6	4.2	1.8	0.9	*	*
2000	2.5	*	*	0.1	0.5	4.4	6.4	4.2	1.6	0.8	*	*
1999	2.5	*	*	0.1	0.6	4.2	6.5	3.9	1.6	0.7	0.3	*
Non-Hispanic white, both sexes												
2002	2.1	*	*	*	0.1	1.7	5.8	4.6	2.2	0.8	0.3	*
2001	2.2	*	*	*	0.1	2.1	6.0	4.2	2.1	0.7	0.3	*
2000	2.3	*	*	*	0.1	2.4	6.2	4.6	2.2	0.8	0.3	*
1999	2.4	*	*	*	0.1	3.0	6.4	4.6	2.0	0.8	0.3	*
Non-Hispanic white, male												
2002	3.7	*	*	*	*	2.7	9.8	8.0	4.0	1.5	0.5	*
2001	3.8	*	*	*	*	3.2	10.4	7.5	3.8	1.4	0.6	*
2000	4.0	*	*	*	0.2	3.7	10.4	8.2	4.0	1.5	0.6	*
1999	4.2	*	*	*	0.2	4.9	10.9	8.2	3.6	1.5	0.6	*
Non-Hispanic white, female												
2002	0.6	*	*	*	*	0.8	1.8	1.2	0.5	*	*	*
2001	0.6	*	*	*	*	0.9	1.7	1.0	0.5	*	*	*
2000	0.7	*	*	*	*	1.1	2.0	1.1	0.5	*	*	*
1999	0.7	*	*	*	*	1.2	1.9	1.1	0.4	*	*	*
Non-Hispanic black, both sexes												
2002	21.3	*	*	*	2.1	21.5	54.0	54.8	26.3	14.6	5.7	*
2001	21.6	*	*	0.4	2.8	24.0	55.0	51.9	28.0	12.8	5.3	*
2000	22.0	*	*	0.4	2.2	27.2	54.5	53.9	26.3	13.2	4.6	*
1999	22.3	*	*	0.6	2.3	28.3	57.2	52.2	26.4	13.2	3.7	*
Non-Hispanic black, male												
2002	30.3	*	*	*	2.0	25.5	75.6	86.2	44.8	26.6	12.3	*
2001	30.8	*	*	*	2.4	30.8	77.8	81.4	47.7	21.4	11.4	*
2000	31.7	*	*	*	2.2	33.4	79.0	86.6	46.4	23.8	8.9	*
1999	32.6	*	*	0.7	1.9	37.1	83.5	85.5	45.6	24.0	6.6	*
Non-Hispanic black, female												
2002	13.2	*	*	*	2.2	17.9	34.9	28.0	11.5	6.0	*	*
2001	13.3	*	*	*	3.2	17.8	34.8	26.6	12.1	6.7	*	*
2000	13.1	*	*	*	2.3	21.7	32.8	25.8	10.1	5.8	*	*
1999	13.0	*	*	*	2.8	20.3	34.1	23.8	11.0	5.6	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)												
All origins 3/, both sexes												
2002	1.9	*	*	*	0.2	1.9	5.1	4.5	1.9	0.7	0.2	*
2001	2.0	*	*	*	0.2	2.1	5.1	4.2	1.9	0.7	0.2	*
2000	2.0	*	*	*	0.2	2.4	5.3	4.4	1.8	0.8	0.3	*
1999	2.1	*	*	0.1	0.2	2.8	5.6	4.4	1.6	0.7	0.3	*
All origins 3/, male												
2002	2.9	*	*	*	0.2	2.3	7.5	7.2	3.0	1.3	0.5	*
2001	2.9	*	*	*	0.3	2.8	7.6	6.7	3.1	1.2	0.5	*
2000	3.1	*	*	*	0.2	3.2	7.8	7.1	2.9	1.4	0.5	*
1999	3.3	*	*	*	0.2	3.8	8.4	7.2	2.7	1.3	0.6	*
All origins 3/, female												
2002	1.0	*	*	*	0.2	1.4	2.7	1.9	0.8	0.3	*	*
2001	1.0	*	*	*	0.2	1.5	2.6	1.8	0.8	0.4	*	*
2000	1.0	*	*	*	0.2	1.6	2.8	1.7	0.7	0.2	*	*
1999	1.0	*	*	*	0.3	1.8	2.7	1.8	0.5	0.3	*	*
Hispanic, both sexes												
2002	2.2	*	*	*	*	1.7	6.1	6.4	4.6	*	*	*
2001	2.3	*	*	*	*	2.1	6.5	7.0	4.2	*	*	*
2000	2.5	*	*	*	*	2.7	7.0	7.6	3.8	*	*	*
1999	2.6	*	*	*	*	3.1	7.6	7.1	3.0	*	*	*
Hispanic, male												
2002	3.2	*	*	*	*	2.2	8.9	10.1	7.3	*	*	*
2001	3.4	*	*	*	*	2.9	9.5	11.2	6.1	*	*	*
2000	3.7	*	*	*	*	3.8	10.6	12.3	5.5	*	*	*
1999	3.8	*	*	*	*	4.2	11.1	11.0	5.6	*	*	*
Hispanic, female												
2002	1.0	*	*	*	*	1.0	2.9	2.8	2.1	*	*	*
2001	1.1	*	*	*	*	1.0	3.2	2.9	2.5	*	*	*
2000	1.2	*	*	*	*	1.4	3.2	3.0	2.3	*	*	*
1999	1.3	*	*	*	*	1.8	3.8	3.3	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.9	*	*	*	0.2	1.9	4.9	4.2	1.7	0.7	0.2	*
2001	1.9	*	*	*	0.2	2.1	4.8	3.9	1.7	0.7	0.2	*
2000	1.9	*	*	*	0.2	2.4	5.0	4.0	1.6	0.7	0.2	*
1999	2.0	*	*	0.1	0.2	2.7	5.2	4.1	1.4	0.7	0.2	*
Non-Hispanic 4/, male												
2002	2.8	*	*	*	0.2	2.3	7.1	6.7	2.7	1.2	0.5	*
2001	2.8	*	*	*	0.2	2.8	7.2	6.1	2.8	1.0	0.4	*
2000	2.9	*	*	*	0.2	3.1	7.4	6.5	2.6	1.3	0.4	*
1999	3.1	*	*	*	0.2	3.6	8.0	6.7	2.4	1.2	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)												
Non-Hispanic 4/, female												
2002	1.0	*	*	*	0.2	1.5	2.7	1.8	0.7	0.3	*	*
2001	1.0	*	*	*	0.2	1.5	2.5	1.7	0.7	0.4	*	*
2000	1.0	*	*	*	0.2	1.7	2.7	1.6	0.6	0.2	*	*
1999	1.0	*	*	*	0.3	1.8	2.6	1.6	0.5	0.3	*	*
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	0.7	2.4	1.8	0.8	0.3	*	*
2001	0.8	*	*	*	*	0.9	2.3	1.7	0.8	0.3	*	*
2000	0.9	*	*	*	0.1	0.9	2.4	1.8	0.7	0.3	*	*
1999	0.9	*	*	*	*	1.2	2.5	1.8	0.7	0.3	*	*
Non-Hispanic white, male												
2002	1.4	*	*	*	*	1.0	3.8	3.1	1.4	0.5	*	*
2001	1.4	*	*	*	*	1.3	3.9	3.0	1.4	0.4	*	*
2000	1.5	*	*	*	*	1.4	3.9	3.2	1.3	0.6	*	*
1999	1.6	*	*	*	*	1.8	4.2	3.2	1.2	0.5	*	*
Non-Hispanic white, female												
2002	0.3	*	*	*	*	0.4	0.9	0.5	0.3	*	*	*
2001	0.3	*	*	*	*	0.4	0.8	0.4	0.2	*	*	*
2000	0.3	*	*	*	*	0.4	0.9	0.5	0.2	*	*	*
1999	0.3	*	*	*	*	0.5	0.8	0.5	*	*	*	*
Non-Hispanic black, both sexes												
2002	8.2	*	*	*	0.9	8.4	20.5	22.2	9.2	4.6	*	*
2001	8.2	*	*	*	1.1	9.2	20.5	20.3	9.9	4.5	*	*
2000	8.5	*	*	*	1.0	10.6	21.7	20.6	8.8	4.6	*	*
1999	8.8	*	*	*	1.2	11.5	22.6	21.6	8.2	4.4	*	*
Non-Hispanic black, male												
2002	11.7	*	*	*	0.9	9.9	29.1	35.5	15.1	7.8	*	*
2001	11.6	*	*	*	1.2	11.2	29.5	31.3	16.6	7.4	*	*
2000	12.2	*	*	*	0.9	13.4	31.0	33.2	15.1	8.8	*	*
1999	12.9	*	*	*	0.8	14.5	33.5	35.2	14.2	8.1	*	*
Non-Hispanic black, female												
2002	5.0	*	*	*	0.8	7.0	13.0	10.7	4.4	2.3	*	*
2001	5.1	*	*	*	1.1	7.4	12.5	10.9	4.5	2.4	*	*
2000	5.1	*	*	*	1.1	8.1	13.4	9.7	3.8	*	*	*
1999	5.2	*	*	*	1.5	8.7	13.0	10.0	3.4	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	0.2	0.8	0.8	0.4	0.2	*	*
2001	0.3	*	*	*	*	0.3	0.9	0.7	0.5	0.2	*	*
2000	0.4	*	*	*	*	0.4	0.9	0.8	0.4	0.2	*	*
1999	0.4	*	*	*	*	0.5	1.0	0.8	0.5	0.2	*	*
All origins 3/, male												
2002	0.5	*	*	*	*	0.3	1.3	1.4	0.6	0.4	*	*
2001	0.6	*	*	*	*	0.5	1.5	1.2	0.9	0.3	*	*
2000	0.6	*	*	*	*	0.5	1.6	1.4	0.8	0.3	*	*
1999	0.7	*	*	*	*	0.9	1.8	1.4	0.9	0.3	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	0.3	0.2	*	*	*	*
2001	0.1	*	*	*	*	0.2	0.2	0.2	0.2	*	*	*
2000	0.1	*	*	*	*	0.2	0.3	0.2	*	*	*	*
1999	0.1	*	*	*	*	0.2	0.3	0.2	*	*	*	*
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	0.8	0.9	*	*	*	*
2001	0.3	*	*	*	*	0.4	0.9	0.9	*	*	*	*
2000	0.4	*	*	*	*	0.4	1.0	1.3	*	*	*	*
1999	0.5	*	*	*	*	0.6	1.7	1.0	*	*	*	*
Hispanic, male												
2002	0.5	*	*	*	*	*	1.3	1.3	*	*	*	*
2001	0.5	*	*	*	*	0.5	1.4	1.5	*	*	*	*
2000	0.6	*	*	*	*	0.6	1.7	2.3	*	*	*	*
1999	0.8	*	*	*	*	0.9	2.9	1.6	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	0.2	0.8	0.8	0.4	0.2	*	*
2001	0.3	*	*	*	*	0.3	0.9	0.7	0.5	0.1	*	*
2000	0.4	*	*	*	*	0.3	0.9	0.8	0.4	0.2	*	*
1999	0.4	*	*	*	*	0.5	0.9	0.8	0.4	0.2	*	*
Non-Hispanic 4/, male												
2002	0.6	*	*	*	*	0.3	1.3	1.3	0.6	0.4	*	*
2001	0.6	*	*	*	*	0.5	1.5	1.1	0.8	*	*	*
2000	0.6	*	*	*	*	0.5	1.6	1.3	0.8	0.3	*	*
1999	0.7	*	*	*	*	0.8	1.6	1.3	0.8	0.3	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	0.3	0.2	*	*	*	*
2001	0.1	*	*	*	*	0.2	0.2	0.2	0.2	*	*	*
2000	0.1	*	*	*	*	0.2	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	*	0.2	0.3	0.2	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	0.1	0.5	0.5	0.2	*	*	*
2001	0.2	*	*	*	*	0.1	0.6	0.4	0.3	*	*	*
2000	0.2	*	*	*	*	0.2	0.6	0.5	0.3	*	*	*
1999	0.3	*	*	*	*	0.3	0.6	0.5	0.3	*	*	*
Non-Hispanic white, male												
2002	0.4	*	*	*	*	0.3	0.9	0.9	0.4	*	*	*
2001	0.4	*	*	*	*	0.2	1.1	0.8	0.4	*	*	*
2000	0.4	*	*	*	*	0.3	1.1	0.9	0.5	*	*	*
1999	0.5	*	*	*	*	0.6	1.1	0.9	0.6	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	1.1	*	*	*	*	0.8	2.7	3.1	1.4	*	*	*
2001	1.1	*	*	*	*	1.2	2.7	2.5	2.5	*	*	*
2000	1.2	*	*	*	*	1.1	3.1	2.9	1.6	*	*	*
1999	1.2	*	*	*	*	1.5	3.0	2.8	1.9	*	*	*
Non-Hispanic black, male												
2002	1.7	*	*	*	*	0.9	4.1	5.0	2.5	*	*	*
2001	1.8	*	*	*	*	1.8	4.4	4.0	4.5	*	*	*
2000	1.9	*	*	*	*	1.6	5.1	4.9	3.2	*	*	*
1999	2.0	*	*	*	*	2.3	5.1	4.9	3.2	*	*	*
Non-Hispanic black, female												
2002	0.6	*	*	*	*	*	1.4	1.5	*	*	*	*
2001	0.6	*	*	*	*	*	1.1	1.3	*	*	*	*
2000	0.5	*	*	*	*	0.7	1.3	1.2	*	*	*	*
1999	0.5	*	*	*	*	0.8	1.2	1.0	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	0.5	1.2	1.0	0.6	0.2	*	*
2001	0.5	*	*	*	*	0.6	1.3	1.0	0.6	0.3	*	*
2000	0.6	*	*	*	*	0.7	1.5	1.2	0.6	0.3	*	*
1999	0.6	*	*	*	*	0.7	1.6	1.1	0.5	0.2	*	*
All origins 3/, male												
2002	0.8	*	*	*	*	0.7	1.9	1.6	1.0	0.4	*	*
2001	0.8	*	*	*	*	0.8	2.0	1.6	1.0	0.4	*	*
2000	0.9	*	*	*	*	1.0	2.4	2.0	1.1	0.4	*	*
1999	0.9	*	*	*	*	1.1	2.5	1.8	0.9	0.4	*	*
All origins 3/, female												
2002	0.2	*	*	*	*	0.3	0.5	0.4	*	*	*	*
2001	0.2	*	*	*	*	0.3	0.5	0.3	0.2	*	*	*
2000	0.2	*	*	*	*	0.4	0.6	0.4	*	*	*	*
1999	0.2	*	*	*	*	0.4	0.6	0.3	*	*	*	*
Hispanic, both sexes												
2002	0.5	*	*	*	*	0.5	1.5	1.4	*	*	*	*
2001	0.5	*	*	*	*	0.5	1.5	1.4	1.2	*	*	*
2000	0.6	*	*	*	*	0.7	1.6	1.6	*	*	*	*
1999	0.6	*	*	*	*	0.6	2.0	1.6	*	*	*	*
Hispanic, male												
2002	0.9	*	*	*	*	1.0	2.4	2.5	*	*	*	*
2001	0.9	*	*	*	*	0.9	2.5	2.5	*	*	*	*
2000	1.0	*	*	*	*	1.1	2.6	2.7	*	*	*	*
1999	1.0	*	*	*	*	0.9	3.3	2.8	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	0.5	1.1	0.9	0.5	0.2	*	*
2001	0.5	*	*	*	*	0.6	1.2	0.9	0.6	0.3	*	*
2000	0.6	*	*	*	*	0.7	1.4	1.1	0.6	0.2	*	*
1999	0.6	*	*	*	*	0.7	1.5	1.0	0.5	0.2	*	*
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	0.7	1.8	1.5	1.0	0.4	*	*
2001	0.8	*	*	*	*	0.8	2.0	1.5	0.9	0.4	*	*
2000	0.9	*	*	*	*	1.0	2.3	1.9	1.0	0.3	*	*
1999	0.9	*	*	*	*	1.1	2.4	1.7	0.9	0.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	0.4	0.5	0.4	*	*	*	*
2001	0.2	*	*	*	*	0.3	0.5	0.3	0.2	*	*	*
2000	0.2	*	*	*	*	0.4	0.6	0.4	*	*	*	*
1999	0.2	*	*	*	*	0.4	0.6	0.3	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	0.2	0.6	0.5	0.3	*	*	*
2001	0.2	*	*	*	*	0.3	0.6	0.5	0.3	*	*	*
2000	0.3	*	*	*	*	0.3	0.9	0.6	0.3	*	*	*
1999	0.3	*	*	*	*	0.4	0.9	0.6	0.2	*	*	*
Non-Hispanic white, male												
2002	0.4	*	*	*	*	0.4	1.1	1.0	0.6	*	*	*
2001	0.4	*	*	*	*	0.4	1.1	0.9	0.5	*	*	*
2000	0.5	*	*	*	*	0.6	1.5	1.0	0.5	*	*	*
1999	0.6	*	*	*	*	0.7	1.5	1.2	0.4	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	*	0.2	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	1.7	*	*	*	*	2.1	4.4	3.8	2.7	1.5	*	*
2001	2.0	*	*	*	*	2.3	5.0	4.0	3.0	1.8	*	*
2000	2.2	*	*	*	*	2.9	5.3	5.3	3.0	1.4	*	*
1999	2.2	*	*	*	*	2.8	6.0	4.1	3.1	1.4	*	*
Non-Hispanic black, male												
2002	2.5	*	*	*	*	2.4	6.5	5.9	5.1	2.9	*	*
2001	2.9	*	*	*	*	3.0	7.7	6.3	4.6	*	*	*
2000	3.3	*	*	*	*	3.7	7.8	8.9	5.7	*	*	*
1999	3.3	*	*	*	*	4.0	8.9	6.7	5.7	*	*	*
Non-Hispanic black, female												
2002	1.0	*	*	*	*	1.8	2.5	1.9	*	*	*	*
2001	1.2	*	*	*	*	1.6	2.7	2.1	1.6	*	*	*
2000	1.2	*	*	*	*	2.2	3.1	2.3	*	*	*	*
1999	1.2	*	*	*	*	1.6	3.3	1.8	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)												
All origins 3/, both sexes												
2002	1.2	*	*	*	0.1	1.0	3.0	2.8	1.4	0.6	0.3	*
2001	1.2	*	*	*	0.2	1.2	3.1	2.6	1.4	0.5	0.2	*
2000	1.2	*	*	*	0.1	1.3	2.9	2.6	1.5	0.6	0.2	*
1999	1.2	*	*	0.1	0.1	1.4	3.0	2.6	1.4	0.7	*	*
All origins 3/, male												
2002	1.7	*	*	*	*	1.3	4.2	4.2	2.3	1.0	0.4	*
2001	1.8	*	*	*	0.1	1.6	4.4	4.1	2.3	0.9	0.4	*
2000	1.8	*	*	*	*	1.6	4.2	4.0	2.4	0.9	0.5	*
1999	1.8	*	*	*	0.1	1.9	4.3	4.1	2.3	1.2	*	*
All origins 3/, female												
2002	0.7	*	*	*	*	0.7	1.8	1.4	0.5	0.3	*	*
2001	0.7	*	*	*	0.2	0.9	1.9	1.3	0.5	*	*	*
2000	0.7	*	*	*	0.1	1.1	1.6	1.2	0.6	0.3	*	*
1999	0.7	*	*	*	0.1	0.9	1.8	1.1	0.5	0.2	*	*
Hispanic, both sexes												
2002	1.1	*	*	*	*	0.6	3.3	3.6	2.5	*	*	*
2001	1.2	*	*	*	*	0.9	3.2	3.9	2.3	*	*	*
2000	1.2	*	*	*	*	1.0	2.9	3.9	2.9	*	*	*
1999	1.2	*	*	*	*	1.2	3.1	3.1	2.7	*	*	*
Hispanic, male												
2002	1.6	*	*	*	*	0.7	4.8	5.6	3.6	*	*	*
2001	1.7	*	*	*	*	1.1	4.6	5.8	3.9	*	*	*
2000	1.7	*	*	*	*	1.3	4.3	6.3	4.5	*	*	*
1999	1.7	*	*	*	*	1.8	4.3	4.4	4.9	*	*	*
Hispanic, female												
2002	0.6	*	*	*	*	*	1.7	1.6	*	*	*	*
2001	0.7	*	*	*	*	0.7	1.8	1.9	*	*	*	*
2000	0.6	*	*	*	*	*	1.5	1.5	*	*	*	*
1999	0.6	*	*	*	*	*	1.8	1.9	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	*	0.1	1.1	2.9	2.6	1.3	0.6	0.2	*
2001	1.2	*	*	*	0.2	1.3	3.1	2.5	1.2	0.4	0.2	*
2000	1.2	*	*	*	0.1	1.4	2.9	2.4	1.3	0.5	0.2	*
1999	1.2	*	*	*	0.1	1.4	3.0	2.4	1.3	0.6	*	*
Non-Hispanic 4/, male												
2002	1.7	*	*	*	*	1.4	4.0	4.0	2.2	1.0	*	*
2001	1.8	*	*	*	*	1.6	4.3	3.8	2.0	0.8	0.4	*
2000	1.7	*	*	*	*	1.6	4.1	3.7	2.1	0.9	*	*
1999	1.8	*	*	*	*	1.9	4.2	3.9	2.1	1.0	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)												
Non-Hispanic 4/, female												
2002	0.7	*	*	*	*	0.8	1.8	1.4	0.5	0.3	*	*
2001	0.7	*	*	*	0.2	0.9	1.9	1.2	0.5	*	*	*
2000	0.7	*	*	*	*	1.2	1.6	1.2	0.6	0.3	*	*
1999	0.7	*	*	*	0.2	1.0	1.8	1.0	0.5	*	*	*
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	0.3	1.1	0.9	0.4	0.2	*	*
2001	0.4	*	*	*	*	0.4	1.2	0.8	0.5	*	*	*
2000	0.5	*	*	*	*	0.4	1.2	0.9	0.5	0.2	*	*
1999	0.5	*	*	*	*	0.5	1.2	0.8	0.4	0.2	*	*
Non-Hispanic white, male												
2002	0.7	*	*	*	*	0.5	1.8	1.4	0.9	0.3	*	*
2001	0.7	*	*	*	*	0.6	1.9	1.4	0.9	*	*	*
2000	0.7	*	*	*	*	0.6	2.0	1.4	0.9	0.3	*	*
1999	0.8	*	*	*	*	0.7	2.0	1.5	0.8	0.3	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	0.4	0.3	*	*	*	*
2001	0.1	*	*	*	*	0.2	0.5	0.2	*	*	*	*
2000	0.2	*	*	*	*	0.3	0.5	0.3	*	*	*	*
1999	0.2	*	*	*	*	0.2	0.5	0.2	*	*	*	*
Non-Hispanic black, both sexes												
2002	5.9	*	*	*	0.5	5.3	14.1	15.5	8.6	4.7	*	*
2001	5.9	*	*	*	0.7	6.1	14.6	14.6	7.8	3.4	2.2	*
2000	5.6	*	*	*	0.5	6.7	13.3	14.2	8.3	3.9	*	*
1999	5.9	*	*	*	0.6	6.9	14.6	14.6	8.3	4.4	*	*
Non-Hispanic black, male												
2002	8.2	*	*	*	*	6.7	19.0	23.7	14.5	8.6	*	*
2001	8.2	*	*	*	*	7.7	19.7	22.7	13.4	6.2	*	*
2000	7.9	*	*	*	*	7.8	18.7	22.3	13.7	6.9	*	*
1999	8.5	*	*	*	*	8.9	20.1	23.6	13.9	8.4	*	*
Non-Hispanic black, female												
2002	3.8	*	*	*	*	4.1	9.7	8.5	3.9	*	*	*
2001	3.7	*	*	*	1.0	4.6	10.1	7.7	3.4	*	*	*
2000	3.6	*	*	*	*	5.7	8.5	7.3	3.9	*	*	*
1999	3.6	*	*	*	0.7	5.1	9.7	6.9	3.8	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)												
All origins 3/, both sexes												
2002	1.0	*	*	*	0.1	1.0	2.7	2.1	0.9	0.4	*	*
2001	1.0	*	*	*	0.1	1.1	2.7	2.0	0.8	0.4	*	*
2000	1.0	*	*	*	0.1	1.3	2.5	2.1	0.9	0.4	*	*
1999	1.0	*	*	*	0.1	1.3	2.6	1.8	0.8	0.4	*	*
All origins 3/, male												
2002	1.5	*	*	*	*	1.2	3.9	3.3	1.5	0.7	*	*
2001	1.5	*	*	*	*	1.5	4.0	3.4	1.4	0.7	*	*
2000	1.5	*	*	*	0.1	1.6	3.8	3.3	1.6	0.7	*	*
1999	1.5	*	*	*	*	1.8	4.0	3.0	1.5	0.7	*	*
All origins 3/, female												
2002	0.5	*	*	*	0.1	0.8	1.4	0.9	0.3	*	*	*
2001	0.5	*	*	*	0.1	0.7	1.4	0.7	0.3	*	*	*
2000	0.5	*	*	*	*	0.9	1.2	0.8	0.2	*	*	*
1999	0.5	*	*	*	*	0.8	1.3	0.7	0.3	*	*	*
Hispanic, both sexes												
2002	0.7	*	*	*	*	0.6	1.8	2.4	1.3	*	*	*
2001	0.6	*	*	*	*	0.5	1.8	2.3	*	*	*	*
2000	0.8	*	*	*	*	0.9	2.3	2.5	1.4	*	*	*
1999	1.0	*	*	*	*	1.0	3.0	2.5	1.3	*	*	*
Hispanic, male												
2002	1.0	*	*	*	*	0.9	2.7	3.6	*	*	*	*
2001	1.0	*	*	*	*	0.6	2.7	3.9	*	*	*	*
2000	1.2	*	*	*	*	1.1	3.6	3.9	*	*	*	*
1999	1.5	*	*	*	*	1.4	4.6	4.3	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	0.9	1.1	*	*	*	*
2001	0.3	*	*	*	*	*	0.8	*	*	*	*	*
2000	0.4	*	*	*	*	*	1.0	*	*	*	*	*
1999	0.4	*	*	*	*	*	1.3	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.0	*	*	*	0.1	1.1	2.8	2.1	0.8	0.4	*	*
2001	1.0	*	*	*	0.1	1.2	2.8	1.9	0.8	0.4	*	*
2000	1.0	*	*	*	0.1	1.3	2.5	2.0	0.8	0.4	*	*
1999	1.0	*	*	*	0.1	1.4	2.5	1.7	0.8	0.3	*	*
Non-Hispanic 4/, male												
2002	1.5	*	*	*	*	1.3	4.0	3.3	1.4	0.7	*	*
2001	1.5	*	*	*	*	1.6	4.1	3.2	1.4	0.6	*	*
2000	1.5	*	*	*	0.1	1.7	3.7	3.2	1.5	0.7	*	*
1999	1.5	*	*	*	*	1.9	3.8	2.8	1.4	0.6	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	0.1	0.9	1.5	0.9	0.3	*	*	*
2001	0.5	*	*	*	0.2	0.7	1.5	0.7	0.3	*	*	*
2000	0.5	*	*	*	*	1.0	1.3	0.8	0.2	*	*	*
1999	0.5	*	*	*	*	0.9	1.3	0.7	0.3	*	*	*
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	0.4	1.2	0.9	0.4	0.1	*	*
2001	0.4	*	*	*	*	0.4	1.3	0.8	0.3	0.2	*	*
2000	0.4	*	*	*	*	0.5	1.1	0.8	0.4	0.1	*	*
1999	0.4	*	*	*	*	0.7	1.2	0.8	0.3	0.1	*	*
Non-Hispanic white, male												
2002	0.8	*	*	*	*	0.6	2.1	1.6	0.7	*	*	*
2001	0.8	*	*	*	*	0.7	2.3	1.5	0.6	0.4	*	*
2000	0.7	*	*	*	*	0.8	1.9	1.5	0.8	0.3	*	*
1999	0.8	*	*	*	*	1.0	2.1	1.4	0.6	0.3	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	0.2	0.3	0.2	*	*	*	*
2001	0.1	*	*	*	*	*	0.3	0.2	*	*	*	*
2000	0.1	*	*	*	*	0.2	0.4	0.2	*	*	*	*
1999	0.1	*	*	*	*	0.3	0.4	0.2	*	*	*	*
Non-Hispanic black, both sexes												
2002	4.4	*	*	*	0.5	4.9	12.3	10.2	4.4	2.9	*	*
2001	4.5	*	*	*	0.6	5.2	12.2	10.4	4.8	2.2	*	*
2000	4.5	*	*	*	0.5	5.9	11.2	10.9	4.5	2.5	*	*
1999	4.2	*	*	*	0.4	5.7	11.1	9.2	4.9	2.0	*	*
Non-Hispanic black, male												
2002	6.2	*	*	*	*	5.5	16.9	16.1	7.7	5.3	*	*
2001	6.4	*	*	*	*	7.0	16.5	17.1	8.5	3.7	*	*
2000	6.5	*	*	*	*	6.9	16.5	17.4	8.8	4.5	*	*
1999	6.0	*	*	*	*	7.4	15.9	15.1	8.5	3.3	*	*
Non-Hispanic black, female												
2002	2.9	*	*	*	*	4.4	8.2	5.2	1.7	*	*	*
2001	2.8	*	*	*	0.8	3.6	8.4	4.6	1.8	*	*	*
2000	2.6	*	*	*	*	5.0	6.5	5.3	*	*	*	*
1999	2.5	*	*	*	*	4.1	6.8	4.1	2.0	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)												
All origins 3/, both sexes												
2002	0.4	3.2	0.3	0.1	0.1	0.1	0.2	0.2	0.4	0.9	1.3	2.9
2001	0.3	2.1	0.3	0.1	0.1	0.1	0.1	0.2	0.4	0.9	1.0	2.2
2000	0.4	1.8	0.3	0.1	0.1	0.1	0.1	0.2	0.5	1.0	1.4	3.3
1999	0.4	2.7	0.3	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.7	4.0
All origins 3/, male												
2002	0.4	3.6	0.4	*	0.1	0.1	0.2	0.3	0.4	0.9	1.6	2.8
2001	0.3	2.1	0.4	*	0.1	*	0.2	0.2	0.5	0.9	1.2	2.4
2000	0.4	1.9	0.3	*	0.1	0.1	0.1	0.2	0.5	1.2	1.8	3.1
1999	0.4	2.8	0.4	*	*	*	0.2	0.3	0.6	1.1	1.6	3.9
All origins 3/, female												
2002	0.4	2.8	*	0.1	*	0.1	0.2	0.2	0.4	0.9	1.2	2.9
2001	0.3	2.0	0.3	0.1	*	*	0.1	0.2	0.4	0.9	0.9	2.1
2000	0.4	1.7	*	*	*	*	0.1	0.2	0.5	0.9	1.2	3.4
1999	0.4	2.5	*	*	*	0.1	0.1	0.3	0.6	0.8	1.8	4.0
Hispanic, both sexes												
2002	0.2	3.4	*	*	*	*	*	*	*	*	*	*
2001	0.2	2.8	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	4.7	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	3.2	0.3	0.1	0.1	0.1	0.2	0.2	0.4	1.0	1.4	3.0
2001	0.3	1.9	0.4	0.1	0.1	0.1	0.2	0.2	0.4	0.9	1.0	2.2
2000	0.4	1.9	0.3	0.1	0.1	0.1	0.1	0.2	0.5	1.0	1.4	3.4
1999	0.4	2.7	0.3	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.7	4.0
Non-Hispanic 4/, male												
2002	0.4	3.3	0.4	*	0.1	*	0.2	0.3	0.4	1.0	1.6	2.9
2001	0.3	1.8	0.4	*	*	*	0.2	0.2	0.5	0.9	1.2	2.4
2000	0.4	2.1	*	*	0.1	*	0.1	0.2	0.6	1.2	1.8	3.1
1999	0.4	2.8	0.4	*	*	*	0.2	0.3	0.6	1.1	1.6	3.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)												
Non-Hispanic 4/, female												
2002	0.4	3.0	*	*	*	*	0.2	0.2	0.4	0.9	1.2	3.0
2001	0.3	2.0	*	*	*	*	0.1	0.2	0.4	0.9	0.9	2.1
2000	0.4	1.8	*	*	*	*	0.1	0.2	0.5	0.9	1.2	3.4
1999	0.5	2.6	*	*	*	*	0.1	0.3	0.6	0.8	1.8	4.1
Non-Hispanic white, both sexes												
2002	0.4	2.9	0.2	*	0.1	0.1	0.2	0.2	0.4	1.0	1.4	3.2
2001	0.3	1.2	0.3	0.1	0.1	*	0.1	0.2	0.5	1.0	1.1	2.4
2000	0.4	1.5	0.2	0.1	0.1	*	0.1	0.2	0.5	1.1	1.4	3.5
1999	0.4	2.3	0.3	*	*	0.1	0.1	0.3	0.6	1.0	1.8	4.2
Non-Hispanic white, male												
2002	0.4	2.6	*	*	*	*	0.2	0.2	0.4	1.1	1.7	3.1
2001	0.3	*	*	*	*	*	0.2	0.2	0.5	1.0	1.3	2.6
2000	0.4	*	*	*	*	*	0.1	0.2	0.6	1.3	1.8	3.4
1999	0.4	2.6	*	*	*	*	0.2	0.3	0.6	1.1	1.7	4.0
Non-Hispanic white, female												
2002	0.4	3.2	*	*	*	*	0.1	0.2	0.4	1.0	1.2	3.3
2001	0.3	*	*	*	*	*	*	0.2	0.4	1.0	0.9	2.3
2000	0.4	*	*	*	*	*	*	0.2	0.5	0.9	1.2	3.6
1999	0.5	2.0	*	*	*	*	*	0.3	0.7	0.9	1.9	4.4
Non-Hispanic black, both sexes												
2002	0.3	4.0	*	*	*	*	*	*	*	*	*	*
2001	0.3	4.3	*	*	*	*	*	*	*	*	*	*
2000	0.4	3.6	*	*	*	*	*	*	*	*	*	*
1999	0.4	4.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
All origins 3/, both sexes												
2002	1.3	1.3	*	0.1	0.1	0.3	0.6	1.7	2.2	3.7	6.4	8.3
2001	1.1	1.6	*	0.0	0.1	0.2	0.5	1.2	1.9	3.6	5.4	8.2
2000	1.0	1.3	*	0.1	0.1	0.2	0.5	1.0	1.7	3.3	5.7	9.0
1999	1.4	1.6	0.1	0.1	0.1	0.4	0.8	1.6	2.4	4.7	7.4	10.0
All origins 3/, male												
2002	1.5	1.5	*	*	*	0.3	0.9	2.5	2.9	4.7	7.9	11.9
2001	1.2	1.6	*	*	0.1	0.3	0.7	1.6	2.3	3.8	7.1	10.1
2000	1.1	1.5	*	*	*	0.2	0.7	1.3	2.0	4.3	6.9	12.6
1999	1.6	1.9	*	*	0.1	0.4	1.0	1.9	2.7	5.4	9.4	13.1
All origins 3/, female												
2002	1.0	1.2	*	*	*	0.2	0.4	0.9	1.6	2.9	5.5	6.8
2001	1.0	1.6	*	*	*	0.2	0.4	0.8	1.5	3.5	4.4	7.4
2000	0.9	1.1	*	*	*	0.2	0.3	0.7	1.5	2.5	4.9	7.6
1999	1.3	1.2	*	*	*	0.4	0.6	1.3	2.1	4.0	6.2	8.8
Hispanic, both sexes												
2002	1.0	*	*	*	*	0.5	0.9	2.9	3.6	4.5	7.8	*
2001	0.7	*	*	*	*	0.3	0.7	1.6	1.8	4.4	6.3	*
2000	0.7	*	*	*	*	*	0.8	1.3	2.0	3.6	6.7	13.3
1999	0.9	*	*	*	*	0.4	1.1	1.5	2.3	4.8	9.8	*
Hispanic, male												
2002	1.3	*	*	*	*	0.5	1.3	4.6	4.9	5.7	9.0	*
2001	0.9	*	*	*	*	*	1.0	2.1	2.9	5.2	*	*
2000	0.9	*	*	*	*	*	1.2	1.4	3.2	4.6	*	*
1999	0.9	*	*	*	*	*	1.5	1.6	*	5.5	12.0	*
Hispanic, female												
2002	0.7	*	*	*	*	*	*	1.2	2.4	3.5	7.0	*
2001	0.6	*	*	*	*	*	*	*	*	3.8	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	*	*	1.3	2.4	4.3	8.3	*
Non-Hispanic 4/, both sexes												
2002	1.3	1.4	*	0.1	0.1	0.2	0.6	1.6	2.1	3.7	6.3	8.3
2001	1.1	1.6	*	*	0.1	0.2	0.5	1.1	1.9	3.6	5.4	8.3
2000	1.1	1.4	*	*	0.1	0.2	0.5	1.0	1.7	3.3	5.6	8.9
1999	1.5	1.5	0.2	0.1	0.1	0.4	0.8	1.6	2.4	4.6	7.3	9.9
Non-Hispanic 4/, male												
2002	1.5	1.6	*	*	*	0.2	0.8	2.3	2.8	4.6	7.8	11.7
2001	1.2	1.8	*	*	*	0.2	0.7	1.5	2.2	3.7	7.0	10.3
2000	1.2	1.5	*	*	*	0.2	0.6	1.2	1.9	4.2	6.7	12.3
1999	1.6	1.7	*	*	*	0.4	1.0	1.9	2.7	5.4	9.3	13.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	0.2	0.4	0.9	1.5	2.9	5.4	6.8
2001	1.0	1.5	*	*	*	0.2	0.4	0.7	1.5	3.4	4.3	7.5
2000	1.0	1.4	*	*	*	0.2	0.3	0.7	1.5	2.5	4.9	7.5
1999	1.4	*	*	*	*	0.4	0.6	1.3	2.1	4.0	6.0	8.6
Non-Hispanic white, both sexes												
2002	1.3	1.1	*	*	*	0.1	0.5	1.4	1.8	3.6	6.2	8.2
2001	1.1	1.1	*	*	*	0.1	0.4	1.0	1.6	3.4	5.3	8.0
2000	1.1	1.0	*	*	*	0.1	0.4	0.8	1.5	3.1	5.5	8.7
1999	1.4	1.1	*	*	0.1	0.2	0.5	1.2	2.0	4.3	7.0	9.5
Non-Hispanic white, male												
2002	1.5	*	*	*	*	*	0.7	2.1	2.4	4.5	7.6	11.7
2001	1.2	*	*	*	*	*	0.5	1.3	2.0	3.6	6.9	10.1
2000	1.1	*	*	*	*	*	0.5	1.1	1.7	4.0	6.3	11.7
1999	1.5	*	*	*	*	0.3	0.8	1.6	2.3	5.1	8.8	12.2
Non-Hispanic white, female												
2002	1.1	*	*	*	*	*	0.3	0.7	1.3	2.8	5.3	6.7
2001	1.0	*	*	*	*	*	0.3	0.7	1.3	3.2	4.2	7.1
2000	1.0	*	*	*	*	*	0.2	0.6	1.3	2.4	4.9	7.4
1999	1.3	*	*	*	*	0.2	0.3	0.9	1.6	3.7	5.8	8.4
Non-Hispanic black, both sexes												
2002	1.5	*	*	*	*	0.7	1.3	3.0	4.2	3.8	6.5	9.4
2001	1.3	3.9	*	*	*	0.6	1.1	2.1	3.4	5.0	5.8	11.5
2000	1.3	3.5	*	*	*	0.6	1.3	2.0	3.0	4.6	7.0	11.1
1999	2.3	3.9	*	*	*	1.3	2.5	4.2	5.1	7.6	10.2	14.5
Non-Hispanic black, male												
2002	1.8	*	*	*	*	0.9	1.7	4.3	6.5	5.0	7.0	*
2001	1.5	*	*	*	*	*	1.6	3.0	4.2	5.8	6.9	*
2000	1.5	*	*	*	*	*	1.7	2.7	3.9	6.4	9.9	*
1999	2.3	*	*	*	*	1.5	2.5	4.7	5.1	8.7	13.8	*
Non-Hispanic black, female												
2002	1.2	*	*	*	*	*	0.9	1.9	2.4	3.0	6.1	*
2001	1.2	*	*	*	*	*	0.7	1.3	2.8	4.5	5.2	12.0
2000	1.1	*	*	*	*	*	0.9	1.5	2.4	3.4	5.4	9.2
1999	2.3	*	*	*	*	1.2	2.5	3.8	5.1	6.9	8.2	11.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mycoses (B35-B49)												
All origins 3/, both sexes												
2002	0.4	1.2	*	*	*	0.1	0.2	0.4	0.7	1.4	2.5	2.5
2001	0.4	1.4	*	*	0.1	0.1	0.2	0.3	0.7	1.3	2.2	3.0
2000	0.4	1.3	*	*	*	0.1	0.2	0.4	0.8	1.5	2.2	3.2
1999	0.7	1.4	*	*	0.1	0.2	0.3	0.7	1.2	2.3	3.3	3.9
All origins 3/, male												
2002	0.5	1.2	*	*	*	0.2	0.3	0.5	1.0	2.0	3.5	4.2
2001	0.4	1.4	*	*	*	*	0.2	0.4	0.8	1.4	3.0	3.8
2000	0.5	1.5	*	*	*	0.1	0.3	0.4	0.9	2.0	2.9	4.9
1999	0.8	1.6	*	*	*	0.2	0.4	0.7	1.4	2.9	4.6	6.6
All origins 3/, female												
2002	0.3	1.1	*	*	*	*	0.1	0.2	0.5	0.9	1.7	1.7
2001	0.4	1.5	*	*	*	*	0.1	0.2	0.5	1.2	1.6	2.6
2000	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	1.8	2.6
1999	0.6	1.1	*	*	*	0.2	0.2	0.6	1.0	1.9	2.4	2.9
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	0.4	*	*	2.8	4.6	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	1.2	*	*	*	0.1	0.2	0.4	0.7	1.4	2.4	2.5
2001	0.4	1.5	*	*	*	0.1	0.2	0.3	0.7	1.3	2.2	3.0
2000	0.5	1.4	*	*	*	0.1	0.2	0.4	0.8	1.5	2.2	3.2
1999	0.7	1.3	*	*	0.1	0.2	0.3	0.7	1.2	2.3	3.2	3.9
Non-Hispanic 4/, male												
2002	0.6	1.3	*	*	*	0.2	0.3	0.5	0.9	2.0	3.5	4.1
2001	0.5	1.6	*	*	*	*	0.2	0.4	0.8	1.4	3.0	3.9
2000	0.5	1.5	*	*	*	*	0.2	0.4	0.9	2.0	2.9	4.9
1999	0.8	1.5	*	*	*	0.2	0.4	0.7	1.4	2.8	4.5	6.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mycoses (B35-B49)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	1.7	1.8
2001	0.4	1.3	*	*	*	*	0.1	0.3	0.6	1.2	1.6	2.6
2000	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	1.7	2.4
1999	0.6	*	*	*	*	0.2	0.2	0.6	1.0	1.8	2.3	2.7
Non-Hispanic white, both sexes												
2002	0.4	0.9	*	*	*	*	0.1	0.3	0.6	1.3	2.3	2.4
2001	0.4	1.0	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.6
2000	0.4	1.0	*	*	*	*	0.1	0.3	0.7	1.4	2.1	2.9
1999	0.6	0.9	*	*	*	0.1	0.2	0.6	1.0	2.1	3.0	3.4
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	0.2	0.3	0.8	1.9	3.4	4.1
2001	0.4	*	*	*	*	*	0.1	0.4	0.7	1.1	2.8	3.5
2000	0.5	*	*	*	*	*	0.2	0.4	0.8	1.8	2.7	4.4
1999	0.7	*	*	*	*	*	0.3	0.6	1.3	2.7	4.3	5.8
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.2	0.4	0.9	1.6	1.7
2001	0.4	*	*	*	*	*	*	0.2	0.5	1.1	1.5	2.3
2000	0.4	*	*	*	*	*	*	0.3	0.6	1.0	1.8	2.2
1999	0.5	*	*	*	*	*	0.1	0.5	0.8	1.6	2.1	2.4
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	0.4	0.5	1.0	1.6	2.2	3.0	*
2001	0.7	3.8	*	*	*	*	0.4	0.7	1.6	2.8	3.6	7.2
2000	0.7	*	*	*	*	*	0.6	0.8	1.5	2.6	3.0	6.7
1999	1.0	3.5	*	*	*	0.6	0.8	1.4	2.2	4.2	5.2	9.1
Non-Hispanic black, male												
2002	0.9	*	*	*	*	*	*	1.7	2.5	3.5	*	*
2001	0.8	*	*	*	*	*	*	*	2.0	3.9	*	*
2000	0.7	*	*	*	*	*	*	1.1	*	3.3	*	*
1999	1.1	*	*	*	*	*	1.0	1.8	2.2	4.8	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	2.1	*	*
1999	1.0	*	*	*	*	*	*	1.1	2.2	3.7	4.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Protozoal diseases (B50-B64)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.2	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.3	0.2	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
1999	0.1	*	*	*	*	0.1	0.2	0.1	0.2	0.3	0.3	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	0.3	0.4	*	*
2000	0.1	*	*	*	*	*	0.2	0.1	0.3	0.4	*	*
1999	0.1	*	*	*	*	*	0.2	0.2	0.2	0.3	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	0.2	0.3	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.3	0.2	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
1999	0.1	*	*	*	*	0.1	0.2	0.1	0.1	0.2	0.3	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	0.3	0.4	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	*	*
1999	0.1	*	*	*	*	*	0.2	0.2	*	0.3	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Protozoal diseases (B50-B64)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	0.2	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.2	*
2001	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.2	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.2	0.2	*
1999	0.1	*	*	*	*	*	0.1	0.1	0.1	0.3	0.3	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.4	*	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	*	*
1999	0.1	*	*	*	*	*	0.2	*	*	0.3	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	0.4	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	0.4	*	*	*	*	*
1999	0.2	*	*	*	*	*	0.5	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leishmaniasis (B55)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leishmaniasis (B55)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Trypanosomiasis (B56-B57)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Trypanosomiasis (B56-B57)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumocystosis (B59)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.3	0.2	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.2	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	*	*
1999	0.1	*	*	*	*	*	0.1	0.2	*	0.3	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	0.2	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.2	0.3	0.2	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.2	0.2	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.2	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	*	*
1999	0.1	*	*	*	*	*	0.2	0.2	*	0.3	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumocystosis (B59)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	0.2	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	*	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.3	0.2	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.2	0.2	*
1999	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.2	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.3	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	0.3	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	0.4	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified protozoal diseases (B58,B60-B64)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified protozoal diseases (B58,B60-B64)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Helminthiases (B65-B83)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Helminthiases (B65-B83)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Schistosomiasis (bilharziasis) (B65)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Schistosomiasis (bilharziasis) (B65)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified helminthiases (B66-B83)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified helminthiases (B66-B83)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sequelae of tuberculosis (B90)												
All origins 3/, both sexes												
2002	0.0	...	*	*	*	*	*	*	*	*	0.3	0.8
2001	0.0	...	*	*	*	*	*	*	*	0.1	0.3	0.6
2000	0.0	...	*	*	*	*	*	*	*	0.1	0.4	0.7
1999	0.1	...	*	*	*	*	*	*	*	0.2	0.6	0.8
All origins 3/, male												
2002	0.0	...	*	*	*	*	*	*	*	*	*	*
2001	0.0	...	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	...	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	...	*	*	*	*	*	*	*	*	0.6	*
All origins 3/, female												
2002	0.0	...	*	*	*	*	*	*	*	*	*	0.9
2001	0.0	...	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	...	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	...	*	*	*	*	*	*	*	*	0.6	0.8
Hispanic, both sexes												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	0.1	...	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	...	*	*	*	*	*	*	*	*	0.3	0.9
2001	0.0	...	*	*	*	*	*	*	*	*	0.3	0.7
2000	0.0	...	*	*	*	*	*	*	*	0.1	0.4	0.5
1999	0.1	...	*	*	*	*	*	*	*	0.2	0.6	0.8
Non-Hispanic 4/, male												
2002	0.0	...	*	*	*	*	*	*	*	*	*	*
2001	0.0	...	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	...	*	*	*	*	*	*	*	*	*	*
1999	0.1	...	*	*	*	*	*	*	*	*	0.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sequelae of tuberculosis (B90)												
Non-Hispanic 4/, female												
2002	0.0	...	*	*	*	*	*	*	*	*	*	0.9
2001	0.0	...	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	...	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	...	*	*	*	*	*	*	*	*	0.6	0.7
Non-Hispanic white, both sexes												
2002	0.0	...	*	*	*	*	*	*	*	*	0.2	0.9
2001	0.0	...	*	*	*	*	*	*	*	*	0.3	0.6
2000	0.0	...	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	...	*	*	*	*	*	*	*	*	0.6	0.8
Non-Hispanic white, male												
2002	0.0	...	*	*	*	*	*	*	*	*	*	*
2001	0.0	...	*	*	*	*	*	*	*	*	*	*
2000	0.0	...	*	*	*	*	*	*	*	*	*	*
1999	0.0	...	*	*	*	*	*	*	*	*	0.5	*
Non-Hispanic white, female												
2002	0.0	...	*	*	*	*	*	*	*	*	*	1.0
2001	0.0	...	*	*	*	*	*	*	*	*	*	*
2000	0.0	...	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	...	*	*	*	*	*	*	*	*	0.6	0.8
Non-Hispanic black, both sexes												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	0.1	...	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sequelae of poliomyelitis (B91)												
All origins 3/, both sexes												
2002	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.3
2001	0.1	...	*	*	*	*	*	0.1	0.2	0.5	0.9	1.2
2000	0.1	...	*	*	*	*	*	0.1	0.2	0.4	0.8	1.2
1999	0.1	...	*	*	*	*	*	0.1	0.1	0.4	0.6	1.3
All origins 3/, male												
2002	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.9
2001	0.1	...	*	*	*	*	*	*	0.2	0.6	1.1	1.8
2000	0.1	...	*	*	*	*	*	*	0.2	0.5	0.8	*
1999	0.1	...	*	*	*	*	*	*	*	0.5	0.7	*
All origins 3/, female												
2002	0.1	...	*	*	*	*	*	*	*	0.5	0.9	1.0
2001	0.1	...	*	*	*	*	*	*	0.2	0.5	0.7	1.0
2000	0.1	...	*	*	*	*	*	*	0.2	0.4	0.8	1.1
1999	0.1	...	*	*	*	*	*	*	*	0.3	0.6	1.2
Hispanic, both sexes												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.3
2001	0.1	...	*	*	*	*	*	0.1	0.2	0.6	0.9	1.3
2000	0.1	...	*	*	*	*	*	*	0.2	0.4	0.8	1.2
1999	0.1	...	*	*	*	*	*	*	0.2	0.4	0.6	1.3
Non-Hispanic 4/, male												
2002	0.1	...	*	*	*	*	*	*	0.2	0.5	1.0	2.0
2001	0.1	...	*	*	*	*	*	*	0.2	0.6	1.2	1.8
2000	0.1	...	*	*	*	*	*	*	*	0.5	0.8	*
1999	0.1	...	*	*	*	*	*	*	*	0.5	0.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sequelae of poliomyelitis (B91)												
Non-Hispanic 4/, female												
2002	0.1	...	*	*	*	*	*	*	*	0.5	0.9	1.1
2001	0.1	...	*	*	*	*	*	*	0.2	0.5	0.7	1.0
2000	0.1	...	*	*	*	*	*	*	0.2	0.4	0.8	1.1
1999	0.1	...	*	*	*	*	*	*	*	0.4	0.6	1.2
Non-Hispanic white, both sexes												
2002	0.2	...	*	*	*	*	*	*	0.2	0.6	1.0	1.5
2001	0.2	...	*	*	*	*	*	*	0.2	0.6	1.0	1.4
2000	0.1	...	*	*	*	*	*	*	0.2	0.5	0.9	1.4
1999	0.1	...	*	*	*	*	*	*	0.2	0.5	0.7	1.5
Non-Hispanic white, male												
2002	0.1	...	*	*	*	*	*	*	0.2	0.6	1.1	2.1
2001	0.2	...	*	*	*	*	*	*	0.2	0.6	1.3	2.0
2000	0.1	...	*	*	*	*	*	*	*	0.6	0.9	*
1999	0.1	...	*	*	*	*	*	*	*	0.5	0.8	*
Non-Hispanic white, female												
2002	0.2	...	*	*	*	*	*	*	*	0.5	1.0	1.2
2001	0.2	...	*	*	*	*	*	*	0.3	0.6	0.8	1.1
2000	0.2	...	*	*	*	*	*	*	0.2	0.4	0.9	1.3
1999	0.1	...	*	*	*	*	*	*	*	0.4	0.7	1.3
Non-Hispanic black, both sexes												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	...	*	*	*	*	*	*	*	*	*	*
2001	*	...	*	*	*	*	*	*	*	*	*	*
2000	*	...	*	*	*	*	*	*	*	*	*	*
1999	*	...	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)												
All origins 3/, both sexes												
2002	0.6	*	*	*	*	0.1	0.4	1.2	1.2	1.6	2.6	3.6
2001	0.4	*	*	*	*	0.1	0.3	0.6	0.7	1.3	1.8	3.2
2000	0.3	*	*	*	*	*	0.2	0.4	0.5	1.0	2.0	3.7
1999	0.5	*	*	*	*	0.1	0.4	0.7	0.8	1.5	2.6	3.9
All origins 3/, male												
2002	0.7	*	*	*	*	*	0.5	1.8	1.6	1.9	3.0	4.7
2001	0.5	*	*	*	*	*	0.4	0.9	0.9	1.3	2.2	3.4
2000	0.4	*	*	*	*	*	0.2	0.6	0.5	1.2	2.4	4.3
1999	0.5	*	*	*	*	0.1	0.4	0.9	0.9	1.6	3.2	3.8
All origins 3/, female												
2002	0.5	*	*	*	*	*	0.2	0.6	0.8	1.4	2.4	3.1
2001	0.4	*	*	*	*	*	0.2	0.4	0.5	1.3	1.6	3.2
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	1.7	3.4
1999	0.5	*	*	*	*	0.1	0.3	0.5	0.8	1.4	2.3	3.9
Hispanic, both sexes												
2002	0.7	*	*	*	*	*	0.6	2.3	2.5	3.0	5.0	*
2001	0.3	*	*	*	*	*	0.4	1.0	*	*	*	*
2000	0.3	*	*	*	*	*	*	0.8	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.8	*	*	*	*	*	0.9	3.5	3.4	3.8	*	*
2001	0.4	*	*	*	*	*	*	1.4	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	*	*	*	*	*	0.3	1.1	1.1	1.5	2.5	3.5
2001	0.4	*	*	*	*	*	0.2	0.6	0.7	1.3	1.8	3.2
2000	0.4	*	*	*	*	*	0.2	0.4	0.5	0.9	1.9	3.7
1999	0.5	*	*	*	*	0.1	0.4	0.7	0.8	1.5	2.6	3.8
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	0.4	1.6	1.4	1.7	2.8	4.7
2001	0.5	*	*	*	*	*	0.3	0.9	0.9	1.3	2.2	3.4
2000	0.4	*	*	*	*	*	0.2	0.5	0.5	1.2	2.2	4.4
1999	0.6	*	*	*	*	*	0.4	0.9	0.9	1.6	3.2	3.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	0.2	0.5	0.7	1.3	2.3	3.0
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.3	1.5	3.1
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.7	1.7	3.5
1999	0.5	*	*	*	*	*	0.3	0.5	0.8	1.4	2.2	3.9
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	0.3	1.0	0.9	1.4	2.5	3.4
2001	0.4	*	*	*	*	*	0.2	0.6	0.6	1.3	1.8	3.2
2000	0.4	*	*	*	*	*	0.2	0.4	0.4	0.9	1.9	3.7
1999	0.5	*	*	*	*	*	0.2	0.5	0.6	1.4	2.5	3.7
Non-Hispanic white, male												
2002	0.7	*	*	*	*	*	0.4	1.6	1.2	1.7	2.8	4.6
2001	0.5	*	*	*	*	*	0.3	0.8	0.8	1.3	2.1	3.5
2000	0.4	*	*	*	*	*	0.2	0.5	0.4	1.1	2.2	4.2
1999	0.5	*	*	*	*	*	0.3	0.7	0.7	1.5	2.9	3.7
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	0.2	0.5	0.6	1.2	2.2	2.8
2001	0.4	*	*	*	*	*	*	0.3	0.5	1.2	1.5	3.1
2000	0.3	*	*	*	*	*	*	0.2	0.3	0.8	1.6	3.5
1999	0.5	*	*	*	*	*	0.1	0.2	0.6	1.3	2.2	3.8
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	0.4	1.4	2.0	*	2.9	*
2001	0.4	*	*	*	*	*	0.4	0.9	1.0	1.9	*	*
2000	0.4	*	*	*	*	*	*	0.6	1.2	1.2	2.9	*
1999	1.0	*	*	*	*	0.4	1.1	2.1	2.5	2.9	4.4	*
Non-Hispanic black, male												
2002	0.7	*	*	*	*	*	*	1.9	3.2	*	*	*
2001	0.5	*	*	*	*	*	*	1.3	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	1.0	2.0	2.5	3.3	7.2	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	0.9	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	2.3	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	1.0	*	*	*	*	*	1.2	2.2	2.4	2.5	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
II. Neoplasms (C00-D48)												
All origins 3/, both sexes												
2002	197.9	3.5	3.0	2.8	4.5	10.1	36.6	125.6	355.6	805.0	1348.0	1,802.6
2001	199.3	2.9	3.1	2.7	4.5	10.5	37.6	128.3	361.5	816.9	1352.8	1,848.4
2000	201.3	4.0	3.1	2.7	4.6	10.2	37.4	129.3	371.7	830.2	1373.2	1,904.3
1999	201.8	3.3	3.1	2.7	4.7	10.4	37.9	129.4	379.3	841.6	1368.0	1,888.7
All origins 3/, male												
2002	208.5	3.5	3.1	3.1	5.1	9.7	32.4	129.9	405.1	980.6	1757.2	2,592.2
2001	210.2	2.8	3.3	2.8	5.3	9.7	33.4	132.4	410.7	1002.0	1775.4	2,721.5
2000	212.0	4.2	3.4	2.9	5.4	9.6	33.6	133.0	421.5	1018.9	1808.1	2,819.6
1999	213.7	3.4	3.3	2.9	5.4	10.2	34.2	133.7	434.1	1037.8	1813.1	2,831.4
All origins 3/, female												
2002	187.6	3.6	2.8	2.6	3.8	10.6	40.8	121.5	309.7	658.8	1076.3	1,460.0
2001	188.7	3.1	2.9	2.6	3.7	11.3	41.8	124.4	316.0	663.5	1075.1	1,483.1
2000	191.1	3.8	2.8	2.5	3.8	10.8	41.2	125.8	325.8	674.8	1089.5	1,531.6
1999	190.3	3.2	2.9	2.6	4.0	10.6	41.5	125.3	328.9	681.0	1079.0	1,508.8
Hispanic, both sexes												
2002	61.2	4.2	2.5	3.2	4.6	7.1	23.8	79.7	218.1	504.9	919.6	1,357.5
2001	62.1	2.4	3.2	2.8	4.4	7.6	26.3	84.4	227.5	513.6	938.1	1,424.3
2000	61.4	3.0	2.8	3.1	4.3	7.7	26.0	83.5	224.1	525.3	977.8	1,453.8
1999	61.1	3.2	2.9	3.0	4.3	7.8	25.0	86.2	228.8	531.1	949.9	1,451.6
Hispanic, male												
2002	62.6	*	2.4	3.3	5.4	6.5	18.9	79.9	258.0	632.6	1222.4	1,925.9
2001	63.7	*	3.3	2.9	5.7	6.5	22.6	83.2	257.9	653.6	1289.1	2,043.6
2000	62.6	*	3.0	3.7	4.8	7.3	20.7	80.6	256.2	660.9	1333.0	2,136.4
1999	62.5	*	2.6	3.0	4.9	7.0	20.6	86.4	272.0	679.3	1277.4	1,995.0
Hispanic, female												
2002	59.6	*	2.6	3.2	3.7	7.7	29.0	79.5	182.2	403.3	712.2	1,076.0
2001	60.3	*	3.1	2.8	3.0	8.8	30.3	85.6	200.5	403.1	700.8	1,121.7
2000	60.1	*	2.6	2.4	3.8	8.1	31.7	86.3	195.6	418.6	739.2	1,118.8
1999	59.6	*	3.3	2.9	3.6	8.7	29.7	86.0	190.6	414.9	732.0	1,189.6
Non-Hispanic 4/, both sexes												
2002	218.5	3.3	3.1	2.7	4.4	10.7	38.4	129.8	365.5	823.6	1365.8	1,816.6
2001	219.2	3.0	3.1	2.7	4.5	11.0	39.0	132.1	370.9	834.9	1368.9	1,860.1
2000	220.8	4.2	3.1	2.6	4.7	10.7	38.7	133.0	381.8	847.0	1386.9	1,916.2
1999	220.7	3.4	3.2	2.7	4.8	10.8	39.3	132.9	389.4	858.0	1381.5	1,899.9
Non-Hispanic 4/, male												
2002	231.8	3.2	3.3	3.0	5.0	10.4	34.4	134.5	415.5	1001.3	1779.6	2,616.1
2001	232.6	3.1	3.3	2.8	5.2	10.4	34.8	136.7	421.0	1021.6	1794.0	2,743.9
2000	234.0	4.4	3.4	2.8	5.5	10.1	35.2	137.3	432.4	1037.7	1824.3	2,841.8
1999	235.2	3.2	3.4	2.9	5.5	10.8	35.9	137.5	444.6	1055.7	1830.6	2,858.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
II. Neoplasms (C00-D48)												
Non-Hispanic 4/, female												
2002	205.9	3.5	2.9	2.4	3.8	11.1	42.3	125.2	319.2	675.3	1091.5	1,471.6
2001	206.4	3.0	2.9	2.6	3.9	11.7	43.2	127.7	324.5	679.7	1089.9	1,492.5
2000	208.4	4.1	2.8	2.5	3.8	11.3	42.2	128.9	334.9	689.5	1102.0	1,542.0
1999	207.0	3.5	2.9	2.5	4.0	10.9	42.7	128.4	338.4	695.7	1090.0	1,515.9
Non-Hispanic white, both sexes												
2002	236.6	3.4	2.9	2.7	4.5	10.4	37.0	124.0	357.6	822.7	1365.9	1,819.8
2001	236.5	3.1	3.3	2.6	4.5	10.8	37.3	126.5	363.4	830.6	1365.1	1,860.0
2000	238.1	4.5	3.2	2.7	4.5	10.5	36.4	126.9	374.5	842.0	1382.1	1,914.2
1999	237.2	3.1	3.4	2.6	4.6	10.6	37.3	126.6	379.8	852.1	1374.2	1,894.6
Non-Hispanic white, male												
2002	249.6	3.5	3.2	2.9	5.1	10.3	33.4	127.8	400.8	992.0	1764.1	2,613.7
2001	249.4	3.2	3.4	2.6	5.1	10.4	33.5	129.5	407.3	1006.5	1771.9	2,728.2
2000	250.2	4.7	3.5	2.8	5.4	10.1	33.1	129.3	417.8	1019.5	1800.1	2,827.2
1999	250.9	3.2	3.7	2.8	5.2	10.5	34.3	128.7	428.2	1038.5	1805.8	2,838.1
Non-Hispanic white, female												
2002	224.1	3.2	2.7	2.5	3.9	10.6	40.5	120.3	316.7	678.4	1099.0	1,476.8
2001	224.2	3.0	3.2	2.6	3.9	11.1	41.2	123.5	321.8	681.4	1095.3	1,498.8
2000	226.5	4.2	2.7	2.6	3.7	10.9	39.7	124.5	333.6	692.5	1107.4	1,546.1
1999	224.0	3.1	3.1	2.4	3.9	10.6	40.2	124.5	334.2	696.1	1091.6	1,518.3
Non-Hispanic black, both sexes												
2002	174.7	3.2	3.5	2.9	4.3	13.4	52.7	188.5	500.2	990.0	1569.1	1,968.7
2001	176.0	*	2.2	3.1	4.8	13.8	53.9	190.7	500.5	1020.1	1591.4	2,020.2
2000	178.1	3.5	3.2	2.5	5.6	13.0	56.4	199.4	510.8	1034.4	1614.1	2,079.4
1999	180.2	4.4	2.6	3.0	6.1	13.2	56.5	199.4	536.9	1049.5	1626.0	2,086.8
Non-Hispanic black, male												
2002	190.7	*	3.4	3.4	4.9	12.0	45.2	204.5	629.1	1307.9	2286.9	3,089.3
2001	193.9	*	2.7	3.4	5.6	11.4	46.9	211.5	620.7	1372.3	2352.8	3,326.3
2000	198.0	*	3.3	2.9	6.7	11.0	50.8	221.8	644.4	1402.2	2408.7	3,366.4
1999	200.5	*	2.9	3.2	7.1	12.7	49.8	228.2	678.2	1406.2	2416.2	3,425.2
Non-Hispanic black, female												
2002	160.1	*	3.7	2.3	3.8	14.7	59.2	174.8	396.8	764.6	1163.1	1,540.5
2001	159.8	*	*	2.7	4.0	15.9	60.1	172.8	403.7	771.6	1163.9	1,536.5
2000	160.2	*	3.1	2.2	4.4	14.8	61.4	180.1	403.3	775.7	1166.9	1,601.3
1999	161.8	*	2.2	2.9	5.1	13.7	62.5	174.8	423.6	797.8	1183.5	1,586.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
All origins 3/, both sexes												
2002	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1311.9	1,723.9
2001	194.5	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1315.8	1,765.7
2000	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1335.6	1,819.4
1999	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1331.5	1,805.8
All origins 3/, male												
2002	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1711.3	2,491.1
2001	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.7	1727.2	2,613.9
2000	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1001.9	1760.6	2,710.7
1999	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1019.4	1767.0	2,722.2
All origins 3/, female												
2002	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1046.7	1,391.1
2001	184.0	1.8	2.5	2.4	3.5	10.9	41.0	122.8	311.6	652.2	1045.5	1,410.8
2000	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1058.5	1,456.4
1999	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1048.5	1,436.5
Hispanic, both sexes												
2002	59.7	*	2.3	2.9	4.5	6.8	23.2	78.2	215.1	496.0	894.9	1,308.8
2001	60.6	*	2.8	2.5	4.3	7.3	25.6	82.9	223.9	504.7	915.0	1,366.8
2000	59.9	*	2.5	2.9	4.2	7.3	25.2	82.1	221.0	516.2	953.5	1,383.5
1999	59.6	*	2.5	2.7	4.1	7.4	24.4	84.9	225.1	520.5	926.8	1,399.6
Hispanic, male												
2002	61.2	*	2.2	2.9	5.2	6.3	18.4	78.4	254.3	622.3	1190.8	1,869.0
2001	62.3	*	2.9	2.6	5.5	6.4	21.7	81.5	253.8	643.6	1259.2	1,973.1
2000	61.3	*	2.9	3.5	4.5	6.9	20.1	79.4	253.1	651.2	1306.4	2,049.7
1999	61.2	*	2.3	2.8	4.6	6.6	19.8	84.6	268.1	668.9	1252.9	1,945.3
Hispanic, female												
2002	58.1	*	2.4	2.9	3.6	7.5	28.4	78.0	179.8	395.6	692.2	1,031.2
2001	58.8	*	2.7	2.5	2.9	8.4	29.8	84.4	197.2	395.3	682.4	1,070.5
2000	58.5	*	2.2	2.3	3.7	7.8	30.7	84.7	192.5	410.0	716.5	1,056.5
1999	58.0	*	2.7	2.6	3.5	8.3	29.4	85.1	187.2	404.3	709.8	1,136.5
Non-Hispanic 4/, both sexes												
2002	213.4	1.7	2.6	2.5	4.2	10.3	37.6	128.0	360.9	810.5	1329.2	1,736.8
2001	213.9	1.6	2.7	2.4	4.2	10.6	38.2	130.3	365.9	820.5	1331.4	1,776.6
2000	215.6	2.5	2.7	2.4	4.4	10.3	37.9	131.2	376.6	832.8	1348.9	1,831.1
1999	215.5	1.7	2.7	2.4	4.5	10.4	38.6	131.0	384.6	843.2	1344.5	1,816.1
Non-Hispanic 4/, male												
2002	226.7	1.9	2.8	2.8	4.8	9.9	33.5	132.6	410.0	985.1	1733.2	2,513.2
2001	227.2	1.6	2.9	2.5	4.9	9.9	34.0	134.6	415.3	1003.8	1745.1	2,634.8
2000	228.7	2.6	2.9	2.5	5.2	9.7	34.3	135.1	426.6	1020.2	1776.0	2,732.4
1999	229.9	1.6	3.0	2.6	5.3	10.3	35.0	135.4	438.8	1036.8	1783.7	2,747.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Non-Hispanic 4/, female												
2002	200.8	1.5	2.4	2.2	3.6	10.6	41.6	123.5	315.4	664.7	1061.5	1,401.8
2001	201.2	1.6	2.5	2.3	3.6	11.3	42.4	126.0	320.0	668.1	1059.8	1,419.6
2000	203.1	2.4	2.6	2.2	3.6	10.9	41.4	127.3	330.3	678.1	1070.6	1,466.6
1999	201.9	1.8	2.5	2.2	3.7	10.5	42.1	126.8	334.5	684.2	1059.4	1,443.2
Non-Hispanic white, both sexes												
2002	230.9	1.6	2.6	2.5	4.3	10.0	36.2	122.3	353.0	809.2	1328.4	1,737.9
2001	230.6	1.8	2.9	2.4	4.3	10.3	36.5	124.6	358.3	815.9	1326.5	1,775.2
2000	232.2	2.6	2.8	2.4	4.3	10.1	35.6	125.1	369.3	827.7	1342.9	1,826.6
1999	231.5	1.6	3.0	2.3	4.3	10.2	36.6	124.8	375.0	837.1	1336.4	1,808.3
Non-Hispanic white, male												
2002	243.8	2.0	2.8	2.7	4.9	9.8	32.5	125.9	395.5	975.3	1716.5	2,507.7
2001	243.4	1.7	3.1	2.4	4.9	10.0	32.6	127.3	401.7	988.2	1721.8	2,616.8
2000	244.4	2.9	3.1	2.6	5.0	9.7	32.3	127.2	412.0	1002.1	1750.2	2,714.1
1999	245.1	1.7	3.3	2.6	5.0	10.1	33.5	126.7	422.3	1019.3	1757.7	2,721.3
Non-Hispanic white, female												
2002	218.5	*	2.3	2.3	3.6	10.3	39.9	118.7	312.8	667.7	1068.3	1,405.4
2001	218.4	1.8	2.8	2.4	3.6	10.7	40.4	121.9	317.3	669.7	1064.4	1,425.1
2000	220.6	2.4	2.4	2.3	3.5	10.5	38.9	123.0	328.9	681.0	1075.3	1,468.7
1999	218.4	*	2.7	2.1	3.6	10.3	39.7	123.0	330.5	684.6	1060.5	1,444.2
Non-Hispanic black, both sexes												
2002	171.5	*	3.0	2.6	4.1	12.8	51.5	186.0	494.6	977.2	1537.4	1,904.3
2001	172.8	*	2.0	2.7	4.4	13.4	52.8	188.7	494.5	1005.6	1559.5	1,947.5
2000	174.9	*	2.8	2.3	5.3	12.3	55.3	196.7	505.2	1019.3	1582.8	2,015.1
1999	176.9	*	2.0	2.8	5.6	12.6	55.3	196.8	530.5	1033.6	1592.9	2,024.0
Non-Hispanic black, male												
2002	187.8	*	2.7	3.1	4.6	11.7	44.3	202.3	621.8	1293.5	2247.6	3,008.4
2001	190.8	*	2.4	3.0	5.1	10.9	45.8	209.6	614.1	1354.7	2310.5	3,238.5
2000	194.9	*	2.6	2.6	6.4	10.5	49.6	219.0	637.4	1382.7	2372.3	3,290.1
1999	197.4	*	2.0	2.9	6.6	11.9	48.6	225.8	670.9	1387.0	2377.4	3,361.1
Non-Hispanic black, female												
2002	156.7	*	3.2	2.0	3.6	13.9	57.8	172.1	392.5	752.8	1135.6	1,482.4
2001	156.5	*	*	2.3	3.8	15.7	59.0	170.7	398.3	759.4	1137.8	1,469.3
2000	156.9	*	3.0	1.9	4.2	14.0	60.4	177.5	398.8	763.6	1138.4	1,541.6
1999	158.3	*	2.0	2.7	4.8	13.3	61.3	172.0	417.9	784.2	1153.6	1,524.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All origins 3/, both sexes												
2002	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3
All origins 3/, male												
2002	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1
All origins 3/, female												
2002	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1
Hispanic, both sexes												
2002	0.7	*	*	*	*	*	*	1.3	3.3	5.3	9.0	16.1
2001	0.8	*	*	*	*	*	*	1.6	3.8	6.9	10.5	19.4
2000	0.8	*	*	*	*	*	*	1.3	4.4	6.8	9.5	18.6
1999	0.8	*	*	*	*	*	*	1.4	4.0	6.9	13.1	14.0
Hispanic, male												
2002	1.1	*	*	*	*	*	*	2.1	5.7	9.7	14.8	*
2001	1.2	*	*	*	*	*	*	2.4	6.4	11.6	17.4	*
2000	1.2	*	*	*	*	*	*	2.2	7.7	12.2	13.8	40.3
1999	1.2	*	*	*	*	*	*	2.3	7.2	13.6	18.2	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	3.1	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	6.6	*
1999	0.5	*	*	*	*	*	*	*	*	*	9.7	*
Non-Hispanic 4/, both sexes												
2002	3.0	*	*	*	0.1	0.2	0.6	2.9	7.1	10.7	13.7	18.7
2001	3.0	*	*	*	0.1	0.2	0.7	2.8	7.1	10.6	14.3	18.4
2000	2.9	*	*	*	0.1	0.1	0.7	3.0	6.9	10.8	13.3	19.5
1999	2.9	*	*	*	0.1	0.1	0.7	3.0	7.0	10.5	14.3	19.4
Non-Hispanic 4/, male												
2002	4.1	*	*	*	*	0.2	0.8	4.6	11.3	16.2	19.8	26.7
2001	4.1	*	*	*	*	0.2	1.0	4.6	11.6	16.2	20.6	25.8
2000	3.9	*	*	*	*	0.1	1.0	4.6	11.1	16.1	19.4	26.5
1999	4.0	*	*	*	0.1	0.2	1.0	4.8	11.0	15.8	21.0	27.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic 4/, female												
2002	1.9	*	*	*	*	*	0.4	1.2	3.2	6.1	9.7	15.3
2001	1.9	*	*	*	*	0.1	0.4	1.2	2.9	6.0	10.2	15.3
2000	2.0	*	*	*	*	0.1	0.4	1.3	3.0	6.5	9.3	16.6
1999	2.0	*	*	*	*	*	0.4	1.4	3.2	6.0	9.9	16.2
Non-Hispanic white, both sexes												
2002	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.6	13.9	18.9
2001	3.1	*	*	*	*	0.1	0.5	2.5	6.5	10.2	14.5	18.9
2000	3.0	*	*	*	*	0.1	0.6	2.5	6.1	10.4	13.3	19.8
1999	3.0	*	*	*	*	0.1	0.5	2.4	6.3	10.1	14.2	19.9
Non-Hispanic white, male												
2002	4.1	*	*	*	*	0.2	0.7	4.0	10.2	15.7	19.9	26.8
2001	4.1	*	*	*	*	0.2	0.8	4.0	10.4	15.3	20.5	26.4
2000	3.9	*	*	*	*	*	0.8	3.8	9.7	15.0	18.9	27.4
1999	3.9	*	*	*	*	*	0.8	3.8	9.7	15.1	20.3	27.5
Non-Hispanic white, female												
2002	2.1	*	*	*	*	*	0.3	1.0	2.9	6.2	9.8	15.5
2001	2.1	*	*	*	*	*	0.3	1.0	2.8	5.9	10.5	15.7
2000	2.1	*	*	*	*	*	0.4	1.2	2.8	6.5	9.6	16.8
1999	2.1	*	*	*	*	*	0.3	1.1	3.0	5.9	10.1	16.8
Non-Hispanic black, both sexes												
2002	2.9	*	*	*	*	*	1.1	5.6	12.4	12.9	12.1	16.0
2001	3.0	*	*	*	*	*	1.3	5.3	12.9	15.7	13.9	13.1
2000	3.1	*	*	*	*	*	1.1	6.1	13.7	15.7	15.1	14.9
1999	3.3	*	*	*	*	*	1.2	7.3	13.6	15.4	16.8	14.5
Non-Hispanic black, male												
2002	4.4	*	*	*	*	*	1.6	9.8	22.5	23.0	18.8	26.3
2001	4.6	*	*	*	*	*	1.7	8.9	23.9	27.3	23.4	*
2000	4.9	*	*	*	*	*	1.8	10.7	24.7	28.6	29.0	*
1999	5.0	*	*	*	*	*	1.8	12.3	23.5	25.4	31.9	25.0
Non-Hispanic black, female												
2002	1.5	*	*	*	*	*	0.7	2.1	4.3	5.8	8.3	12.1
2001	1.5	*	*	*	*	*	0.9	2.1	4.0	7.5	8.6	10.7
2000	1.5	*	*	*	*	*	*	2.1	4.9	6.6	7.3	14.4
1999	1.7	*	*	*	*	*	0.7	3.0	5.7	8.4	8.4	10.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of lip (C00)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of lip (C00)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of tongue (C01-C02)												
All origins 3/, both sexes												
2002	0.7	*	*	*	*	0.1	0.2	0.7	1.7	2.4	3.3	4.0
2001	0.6	*	*	*	*	0.1	0.2	0.7	1.7	2.3	3.3	3.8
2000	0.6	*	*	*	*	0.1	0.2	0.7	1.7	2.3	3.0	4.0
1999	0.6	*	*	*	*	*	0.2	0.7	1.5	2.4	3.1	4.0
All origins 3/, male												
2002	0.9	*	*	*	*	*	0.3	1.0	2.7	3.6	4.4	5.2
2001	0.9	*	*	*	*	*	0.2	1.1	2.8	3.5	4.4	4.1
2000	0.8	*	*	*	*	*	0.2	1.1	2.7	3.4	3.6	5.1
1999	0.8	*	*	*	*	*	0.3	1.1	2.4	3.7	4.5	5.4
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.3	0.9	1.5	2.5	3.5
2001	0.4	*	*	*	*	*	0.1	0.3	0.7	1.2	2.5	3.6
2000	0.5	*	*	*	*	*	0.1	0.4	0.8	1.4	2.5	3.6
1999	0.4	*	*	*	*	*	0.2	0.3	0.8	1.3	2.2	3.5
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	0.5	1.0	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	1.4	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	*	*	0.1	0.2	0.7	1.8	2.5	3.3	4.0
2001	0.7	*	*	*	*	0.1	0.2	0.7	1.7	2.4	3.3	3.7
2000	0.7	*	*	*	*	0.1	0.2	0.7	1.7	2.4	3.0	4.0
1999	0.7	*	*	*	*	*	0.2	0.8	1.6	2.4	3.1	4.1
Non-Hispanic 4/, male												
2002	0.9	*	*	*	*	*	0.3	1.1	2.7	3.7	4.5	5.0
2001	0.9	*	*	*	*	*	0.3	1.2	2.9	3.7	4.3	4.1
2000	0.9	*	*	*	*	*	0.3	1.1	2.8	3.4	3.7	4.8
1999	0.9	*	*	*	*	*	0.3	1.2	2.4	3.8	4.5	5.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of tongue (C01-C02)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.3	0.9	1.5	2.5	3.6
2001	0.5	*	*	*	*	*	0.1	0.3	0.7	1.3	2.5	3.5
2000	0.5	*	*	*	*	*	0.2	0.4	0.8	1.5	2.5	3.6
1999	0.5	*	*	*	*	*	0.1	0.4	0.8	1.3	2.2	3.6
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	*	0.2	0.6	1.7	2.5	3.5	4.0
2001	0.8	*	*	*	*	0.1	0.2	0.7	1.7	2.3	3.3	3.9
2000	0.7	*	*	*	*	*	0.2	0.7	1.6	2.4	3.1	4.2
1999	0.7	*	*	*	*	*	0.2	0.7	1.5	2.4	3.2	4.1
Non-Hispanic white, male												
2002	1.0	*	*	*	*	*	0.3	1.0	2.6	3.5	4.6	5.0
2001	1.0	*	*	*	*	*	0.2	1.1	2.7	3.6	4.4	4.4
2000	0.9	*	*	*	*	*	0.3	1.0	2.7	3.3	3.6	5.1
1999	0.9	*	*	*	*	*	0.3	1.0	2.2	3.7	4.6	5.5
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	0.3	0.9	1.6	2.7	3.6
2001	0.5	*	*	*	*	*	*	0.3	0.7	1.3	2.6	3.7
2000	0.6	*	*	*	*	*	0.2	0.4	0.7	1.5	2.7	3.8
1999	0.5	*	*	*	*	*	0.1	0.3	0.8	1.2	2.3	3.6
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	*	1.2	2.6	2.9	*	*
2001	0.6	*	*	*	*	*	*	1.1	2.4	2.6	2.5	*
2000	0.6	*	*	*	*	*	*	1.2	2.6	2.8	2.3	*
1999	0.6	*	*	*	*	*	*	1.7	2.7	2.9	2.5	*
Non-Hispanic black, male												
2002	0.9	*	*	*	*	*	*	1.8	4.6	5.4	*	*
2001	0.9	*	*	*	*	*	*	2.0	4.8	4.3	*	*
2000	0.9	*	*	*	*	*	*	2.3	4.5	5.1	*	*
1999	0.9	*	*	*	*	*	*	2.8	4.7	4.8	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pharynx (C10-C13,C14.0)												
All origins 3/, both sexes												
2002	1.1	*	*	*	*	*	0.2	1.2	3.0	4.5	5.1	5.4
2001	1.1	*	*	*	*	*	0.3	1.2	3.1	4.5	5.5	5.9
2000	1.1	*	*	*	*	*	0.2	1.2	3.1	4.7	5.3	5.6
1999	1.1	*	*	*	*	*	0.3	1.3	3.1	4.6	5.6	5.2
All origins 3/, male												
2002	1.6	*	*	*	*	*	0.3	1.9	4.8	7.1	8.2	9.8
2001	1.6	*	*	*	*	*	0.4	2.0	5.1	7.2	8.9	10.0
2000	1.6	*	*	*	*	*	0.4	1.9	5.2	7.5	8.6	9.9
1999	1.6	*	*	*	*	*	0.4	2.1	5.1	7.2	9.0	9.9
All origins 3/, female												
2002	0.6	*	*	*	*	*	0.2	0.5	1.3	2.3	3.0	3.5
2001	0.6	*	*	*	*	*	0.1	0.5	1.2	2.3	3.3	4.2
2000	0.6	*	*	*	*	*	*	0.5	1.3	2.5	3.1	3.9
1999	0.6	*	*	*	*	*	0.1	0.5	1.3	2.5	3.3	3.4
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	1.5	2.5	3.7	*
2001	0.4	*	*	*	*	*	*	0.7	2.1	3.2	3.8	*
2000	0.3	*	*	*	*	*	*	*	1.9	2.5	*	*
1999	0.4	*	*	*	*	*	*	*	1.8	3.2	4.8	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	2.5	4.8	*	*
2001	0.5	*	*	*	*	*	*	1.2	3.6	5.0	*	*
2000	0.5	*	*	*	*	*	*	*	3.6	*	*	*
1999	0.5	*	*	*	*	*	*	*	3.1	6.6	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	*	*	*	0.2	1.3	3.1	4.6	5.1	5.4
2001	1.2	*	*	*	*	*	0.3	1.3	3.2	4.6	5.6	5.9
2000	1.2	*	*	*	*	*	0.3	1.2	3.2	4.8	5.3	5.5
1999	1.2	*	*	*	*	*	0.3	1.3	3.2	4.6	5.6	5.3
Non-Hispanic 4/, male												
2002	1.7	*	*	*	*	*	0.3	2.1	5.0	7.2	8.2	9.5
2001	1.8	*	*	*	*	*	0.4	2.1	5.2	7.3	9.0	10.0
2000	1.8	*	*	*	*	*	0.4	2.0	5.3	7.6	8.7	9.5
1999	1.8	*	*	*	*	*	0.4	2.2	5.2	7.2	9.1	10.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pharynx (C10-C13,C14.0)												
Non-Hispanic 4/, female												
2002	0.7	*	*	*	*	*	0.2	0.5	1.4	2.4	3.1	3.6
2001	0.7	*	*	*	*	*	0.1	0.5	1.3	2.3	3.4	4.2
2000	0.7	*	*	*	*	*	*	0.5	1.3	2.6	3.1	3.8
1999	0.7	*	*	*	*	*	0.1	0.5	1.4	2.6	3.3	3.4
Non-Hispanic white, both sexes												
2002	1.2	*	*	*	*	*	0.2	1.0	2.6	4.4	4.9	5.2
2001	1.2	*	*	*	*	*	0.2	0.9	2.7	4.2	5.4	5.9
2000	1.1	*	*	*	*	*	0.1	0.9	2.6	4.4	5.1	5.2
1999	1.1	*	*	*	*	*	0.2	0.9	2.7	4.3	5.2	5.4
Non-Hispanic white, male												
2002	1.7	*	*	*	*	*	0.2	1.6	4.2	6.8	7.8	9.2
2001	1.7	*	*	*	*	*	0.3	1.6	4.4	6.5	8.6	9.9
2000	1.6	*	*	*	*	*	0.2	1.4	4.2	6.8	8.2	9.2
1999	1.6	*	*	*	*	*	0.3	1.5	4.4	6.5	8.3	10.0
Non-Hispanic white, female												
2002	0.7	*	*	*	*	*	*	0.4	1.1	2.4	2.9	3.5
2001	0.7	*	*	*	*	*	*	0.3	1.1	2.3	3.3	4.3
2000	0.7	*	*	*	*	*	*	0.4	1.1	2.4	3.1	3.6
1999	0.7	*	*	*	*	*	*	0.4	1.2	2.4	3.2	3.5
Non-Hispanic black, both sexes												
2002	1.5	*	*	*	*	*	0.6	2.9	6.5	6.1	6.4	7.3
2001	1.5	*	*	*	*	*	0.7	2.9	6.6	7.7	7.4	*
2000	1.7	*	*	*	*	*	0.6	2.9	7.9	8.7	8.0	7.3
1999	1.7	*	*	*	*	*	0.6	3.7	7.0	8.2	9.8	*
Non-Hispanic black, male												
2002	2.3	*	*	*	*	*	0.8	5.2	12.1	10.7	10.6	*
2001	2.4	*	*	*	*	*	1.0	4.9	12.0	14.5	12.9	*
2000	2.7	*	*	*	*	*	1.1	5.2	14.3	15.9	15.7	*
1999	2.7	*	*	*	*	*	1.0	6.7	12.1	13.7	19.4	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	0.9	2.1	2.9	4.0	*
2001	0.7	*	*	*	*	*	*	1.2	2.3	2.8	4.4	*
2000	0.7	*	*	*	*	*	*	1.0	2.8	3.7	3.6	*
1999	0.8	*	*	*	*	*	*	1.2	2.9	4.3	4.4	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)												
All origins 3/, both sexes												
2002	0.9	*	*	*	*	*	0.2	0.9	2.1	3.4	5.0	8.7
2001	0.9	*	*	*	*	*	0.2	0.8	2.0	3.6	5.3	8.3
2000	0.9	*	*	*	*	*	0.2	0.9	1.9	3.5	4.7	9.5
1999	0.9	*	*	*	*	*	0.2	0.9	2.1	3.3	5.4	9.7
All origins 3/, male												
2002	1.2	*	*	*	*	*	0.2	1.4	3.4	5.1	6.7	11.1
2001	1.2	*	*	*	*	*	0.2	1.2	3.3	5.1	7.0	10.6
2000	1.1	*	*	*	*	*	0.3	1.4	2.9	5.0	6.7	11.4
1999	1.1	*	*	*	*	*	0.3	1.4	3.2	4.7	7.2	11.4
All origins 3/, female												
2002	0.7	*	*	*	*	*	0.1	0.3	0.9	2.0	3.9	7.7
2001	0.7	*	*	*	*	*	0.1	0.4	0.9	2.3	4.1	7.3
2000	0.7	*	*	*	*	*	0.1	0.4	0.9	2.3	3.4	8.8
1999	0.7	*	*	*	*	*	0.1	0.4	1.0	2.1	4.3	9.0
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	2.6	*	*
2000	0.2	*	*	*	*	*	*	*	*	2.4	*	*
1999	0.3	*	*	*	*	*	*	*	1.2	2.1	4.6	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	5.0	*	*
2000	0.4	*	*	*	*	*	*	*	*	4.2	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.0	*	*	*	*	*	0.2	0.9	2.2	3.5	5.1	8.9
2001	1.0	*	*	*	*	*	0.2	0.8	2.1	3.6	5.3	8.4
2000	1.0	*	*	*	*	*	0.2	1.0	1.9	3.6	4.8	9.7
1999	1.0	*	*	*	*	*	0.2	0.9	2.1	3.3	5.5	9.7
Non-Hispanic 4/, male												
2002	1.3	*	*	*	*	*	0.2	1.5	3.6	5.3	6.8	11.3
2001	1.3	*	*	*	*	*	0.3	1.3	3.4	5.1	7.0	10.7
2000	1.2	*	*	*	*	*	0.3	1.5	2.9	5.0	6.7	11.5
1999	1.3	*	*	*	*	*	0.3	1.4	3.3	4.7	7.2	11.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)												
Non-Hispanic 4/, female												
2002	0.7	*	*	*	*	*	0.1	0.4	0.9	2.1	4.0	7.8
2001	0.8	*	*	*	*	*	0.1	0.4	0.9	2.4	4.2	7.4
2000	0.8	*	*	*	*	*	0.1	0.4	0.9	2.4	3.5	9.0
1999	0.8	*	*	*	*	*	0.1	0.5	1.1	2.2	4.3	9.0
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	*	*	0.2	0.8	2.1	3.6	5.3	9.2
2001	1.1	*	*	*	*	*	0.2	0.8	2.0	3.6	5.6	8.6
2000	1.1	*	*	*	*	*	0.2	0.9	1.8	3.6	4.9	10.1
1999	1.1	*	*	*	*	*	0.2	0.8	2.0	3.3	5.6	10.1
Non-Hispanic white, male												
2002	1.4	*	*	*	*	*	0.2	1.4	3.4	5.3	7.1	11.7
2001	1.3	*	*	*	*	*	0.3	1.3	3.1	5.0	7.2	11.1
2000	1.3	*	*	*	*	*	0.3	1.3	2.7	4.8	6.7	12.3
1999	1.3	*	*	*	*	*	0.2	1.3	3.0	4.6	7.3	11.5
Non-Hispanic white, female												
2002	0.8	*	*	*	*	*	0.1	0.3	0.9	2.2	4.1	8.1
2001	0.9	*	*	*	*	*	*	0.4	0.9	2.4	4.5	7.5
2000	0.9	*	*	*	*	*	0.1	0.4	0.9	2.6	3.7	9.2
1999	0.9	*	*	*	*	*	*	0.4	1.0	2.2	4.5	9.5
Non-Hispanic black, both sexes												
2002	0.8	*	*	*	*	*	*	1.6	3.2	3.8	4.0	6.4
2001	0.9	*	*	*	*	*	*	1.2	3.8	5.5	3.9	*
2000	0.8	*	*	*	*	*	*	1.9	3.2	4.1	4.7	*
1999	0.9	*	*	*	*	*	0.4	1.9	3.9	4.3	4.6	6.5
Non-Hispanic black, male												
2002	1.2	*	*	*	*	*	*	2.7	5.9	6.7	*	*
2001	1.3	*	*	*	*	*	*	1.9	7.0	8.5	6.6	*
2000	1.3	*	*	*	*	*	*	3.2	5.8	7.6	8.3	*
1999	1.3	*	*	*	*	*	*	2.9	6.7	6.9	7.8	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	3.5	*
2001	0.5	*	*	*	*	*	*	*	*	3.3	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	1.0	1.6	2.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)												
All origins 3/, both sexes												
2002	45.7	*	0.2	0.1	0.3	1.5	7.4	29.7	80.6	181.0	311.9	467.4
2001	46.0	*	*	*	0.3	1.5	7.4	30.1	80.4	183.8	315.0	488.0
2000	46.4	*	0.2	0.1	0.2	1.5	7.2	29.6	83.1	184.4	323.9	504.9
1999	46.4	*	0.2	*	0.3	1.5	7.4	29.3	84.0	187.0	320.2	507.4
All origins 3/, male												
2002	50.7	*	0.3	*	0.3	1.8	9.1	39.7	108.1	237.5	397.9	557.9
2001	50.5	*	*	*	0.3	1.7	8.8	39.7	106.9	238.5	402.1	593.3
2000	50.8	*	*	*	0.3	1.8	8.9	39.1	109.6	239.8	413.7	614.1
1999	50.7	*	0.3	*	0.3	1.8	8.9	38.9	110.0	243.0	411.2	620.8
All origins 3/, female												
2002	40.8	*	*	*	0.3	1.3	5.7	20.1	55.2	133.9	254.9	428.1
2001	41.7	*	*	*	0.2	1.3	6.0	20.9	56.0	138.5	257.7	443.9
2000	42.3	*	*	*	0.2	1.2	5.6	20.5	58.6	138.9	265.3	460.4
1999	42.3	*	*	*	0.3	1.2	5.9	20.1	60.0	141.1	261.1	461.7
Hispanic, both sexes												
2002	18.2	*	*	*	*	1.1	6.2	24.6	70.3	162.4	273.7	420.2
2001	18.5	*	*	*	*	1.2	6.4	26.1	72.0	163.3	291.2	446.4
2000	18.2	*	*	*	*	1.2	6.6	25.4	69.6	166.3	301.4	436.6
1999	17.9	*	*	*	*	1.4	5.9	25.9	69.4	166.6	295.0	443.4
Hispanic, male												
2002	20.0	*	*	*	*	1.3	6.5	32.7	94.8	217.7	339.4	480.2
2001	20.0	*	*	*	*	1.1	7.3	33.7	93.5	212.5	363.5	554.2
2000	19.8	*	*	*	*	1.3	6.9	31.4	90.1	224.3	394.3	560.3
1999	19.3	*	*	*	*	1.4	6.2	33.7	90.8	214.9	377.0	528.4
Hispanic, female												
2002	16.4	*	*	*	*	1.0	5.8	16.5	48.3	118.5	228.8	390.4
2001	17.0	*	*	*	*	1.5	5.4	18.6	52.9	124.6	242.4	393.7
2000	16.4	*	*	*	*	1.1	6.3	19.5	51.5	120.6	239.0	375.9
1999	16.5	*	*	*	*	1.3	5.6	18.4	50.5	128.7	240.4	402.4
Non-Hispanic 4/, both sexes												
2002	49.8	*	0.2	*	0.3	1.6	7.5	30.1	81.2	181.8	313.1	468.0
2001	50.0	*	*	*	0.3	1.6	7.5	30.4	80.9	184.6	315.5	488.5
2000	50.4	*	0.2	*	0.2	1.6	7.3	29.9	83.8	185.1	324.1	506.3
1999	50.2	*	0.2	*	0.3	1.5	7.5	29.5	84.8	187.6	320.4	508.3
Non-Hispanic 4/, male												
2002	55.6	*	*	*	0.4	1.8	9.4	40.3	108.7	238.3	399.8	559.8
2001	55.1	*	*	*	0.3	1.8	9.0	40.1	107.5	239.3	403.2	593.9
2000	55.3	*	*	*	0.3	1.9	9.1	39.7	110.8	240.0	413.6	615.2
1999	55.1	*	*	*	0.3	1.8	9.2	39.2	111.0	243.8	411.5	622.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)												
Non-Hispanic 4/, female												
2002	44.3	*	*	*	0.3	1.3	5.7	20.4	55.7	134.7	255.6	428.4
2001	45.1	*	*	*	0.3	1.3	6.0	21.1	56.1	139.2	257.9	444.6
2000	45.7	*	*	*	0.2	1.2	5.4	20.6	58.9	139.7	265.9	462.2
1999	45.5	*	*	*	0.3	1.2	5.9	20.2	60.6	141.5	261.3	462.5
Non-Hispanic white, both sexes												
2002	52.3	*	0.2	*	0.3	1.4	6.8	27.4	75.9	174.7	305.3	460.5
2001	52.3	*	*	*	0.2	1.4	6.7	27.6	75.6	176.9	307.1	481.5
2000	52.7	*	*	*	0.2	1.3	6.3	27.1	78.3	177.8	315.1	497.6
1999	52.5	*	0.2	*	0.2	1.3	6.6	26.8	78.8	180.7	311.5	500.1
Non-Hispanic white, male												
2002	58.2	*	*	*	0.3	1.6	8.5	36.7	101.7	229.8	390.8	551.6
2001	57.6	*	*	*	0.3	1.6	8.0	36.4	100.8	230.1	394.6	587.7
2000	57.6	*	*	*	0.3	1.7	8.0	35.9	103.2	231.0	403.5	605.1
1999	57.4	*	*	*	0.3	1.5	8.2	35.3	102.9	235.9	400.7	616.1
Non-Hispanic white, female												
2002	46.6	*	*	*	0.3	1.3	5.1	18.2	51.5	127.7	248.0	421.1
2001	47.2	*	*	*	0.2	1.2	5.4	18.9	51.8	131.9	249.2	437.3
2000	48.0	*	*	*	0.2	1.0	4.7	18.4	54.9	132.9	257.1	454.3
1999	47.8	*	*	*	0.2	1.0	5.0	18.4	56.2	134.5	253.2	453.9
Non-Hispanic black, both sexes												
2002	45.1	*	*	*	0.4	2.2	11.4	49.3	129.2	255.8	412.2	571.4
2001	45.6	*	*	*	*	2.3	11.9	51.0	128.7	260.9	421.1	590.9
2000	46.1	*	*	*	*	2.4	12.1	52.4	131.8	261.4	434.5	612.8
1999	46.1	*	*	*	0.4	2.5	12.2	50.2	136.7	263.0	433.6	617.2
Non-Hispanic black, male												
2002	50.2	*	*	*	*	2.8	14.4	66.7	176.9	338.4	535.7	678.2
2001	49.7	*	*	*	*	2.7	14.6	68.2	172.5	345.0	534.1	686.1
2000	50.9	*	*	*	*	2.8	15.3	70.8	182.1	343.5	557.5	736.2
1999	51.0	*	*	*	*	3.1	14.7	70.2	188.0	341.3	573.5	717.9
Non-Hispanic black, female												
2002	40.5	*	*	*	*	1.6	8.8	34.4	90.9	197.3	342.4	530.5
2001	41.8	*	*	*	*	1.8	9.6	36.2	93.4	201.5	357.7	555.6
2000	41.7	*	*	*	*	2.1	9.2	36.6	91.4	203.6	365.2	566.9
1999	41.7	*	*	*	*	2.0	10.0	33.2	95.6	207.7	355.2	579.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of esophagus (C15)												
All origins 3/, both sexes												
2002	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
All origins 3/, male												
2002	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.2
2000	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
All origins 3/, female												
2002	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3
Hispanic, both sexes												
2002	1.1	*	*	*	*	*	0.4	1.5	4.1	10.2	15.2	23.3
2001	1.1	*	*	*	*	*	*	1.9	5.2	8.5	18.4	20.6
2000	1.1	*	*	*	*	*	*	1.2	5.0	11.1	19.4	19.9
1999	1.1	*	*	*	*	*	*	1.5	5.2	9.2	18.1	26.7
Hispanic, male												
2002	1.6	*	*	*	*	*	*	2.7	8.0	18.3	27.1	36.8
2001	1.7	*	*	*	*	*	*	3.6	8.5	16.2	30.4	43.8
2000	1.6	*	*	*	*	*	*	2.2	8.0	19.4	35.4	40.3
1999	1.7	*	*	*	*	*	*	2.7	10.2	16.3	32.8	49.6
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	3.6	7.0	16.6
2001	0.5	*	*	*	*	*	*	*	2.2	*	10.3	*
2000	0.5	*	*	*	*	*	*	*	2.4	4.6	8.6	*
1999	0.4	*	*	*	*	*	*	*	*	3.6	8.3	*
Non-Hispanic 4/, both sexes												
2002	4.9	*	*	*	*	*	0.7	3.7	10.8	20.8	27.2	24.5
2001	4.9	*	*	*	*	0.1	0.7	3.7	10.4	21.3	26.6	26.0
2000	4.8	*	*	*	*	0.1	0.7	3.8	11.2	20.4	25.8	26.1
1999	4.7	*	*	*	*	0.1	0.7	3.6	11.0	20.6	24.6	26.8
Non-Hispanic 4/, male												
2002	7.8	*	*	*	*	*	1.2	6.3	18.9	36.6	49.3	43.5
2001	7.7	*	*	*	*	0.1	1.3	6.5	18.2	37.1	48.1	48.4
2000	7.5	*	*	*	*	0.2	1.3	6.6	19.4	34.8	46.1	47.8
1999	7.4	*	*	*	*	0.2	1.2	6.4	18.8	36.2	44.7	48.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of esophagus (C15)												
Non-Hispanic 4/, female												
2002	2.2	*	*	*	*	*	0.3	1.1	3.4	7.6	12.5	16.3
2001	2.2	*	*	*	*	*	0.3	1.1	3.2	8.2	12.5	16.7
2000	2.3	*	*	*	*	*	0.1	1.0	3.6	8.5	12.6	17.3
1999	2.2	*	*	*	*	*	0.2	0.9	3.8	7.8	11.6	18.3
Non-Hispanic white, both sexes												
2002	5.3	*	*	*	*	*	0.7	3.6	10.4	20.5	27.3	25.1
2001	5.1	*	*	*	*	0.1	0.8	3.5	9.7	20.5	26.8	26.5
2000	5.1	*	*	*	*	0.1	0.7	3.5	10.4	20.0	25.5	26.7
1999	4.9	*	*	*	*	0.1	0.7	3.3	10.2	20.0	24.4	27.2
Non-Hispanic white, male												
2002	8.4	*	*	*	*	*	1.3	6.3	18.2	36.3	49.9	44.7
2001	8.1	*	*	*	*	*	1.3	6.1	17.0	36.0	48.6	49.0
2000	8.0	*	*	*	*	0.2	1.4	6.3	18.3	34.3	45.9	49.4
1999	7.7	*	*	*	*	0.2	1.2	6.0	17.5	35.4	44.4	48.0
Non-Hispanic white, female												
2002	2.3	*	*	*	*	*	0.2	0.9	3.0	7.1	12.2	16.6
2001	2.3	*	*	*	*	*	0.2	1.0	2.7	7.5	12.4	17.1
2000	2.3	*	*	*	*	*	*	0.8	3.0	8.0	12.0	17.5
1999	2.2	*	*	*	*	*	0.2	0.8	3.3	7.1	11.3	18.9
Non-Hispanic black, both sexes												
2002	4.3	*	*	*	*	*	1.0	5.2	17.7	28.9	29.9	20.6
2001	4.7	*	*	*	*	*	1.0	6.1	18.9	32.8	30.0	24.0
2000	4.7	*	*	*	*	*	0.8	6.8	20.2	28.6	34.1	22.6
1999	4.7	*	*	*	*	*	1.0	6.6	20.3	30.2	31.4	23.6
Non-Hispanic black, male												
2002	6.5	*	*	*	*	*	1.5	8.4	30.7	49.8	52.2	38.3
2001	7.2	*	*	*	*	*	1.6	10.5	33.0	56.2	55.5	47.4
2000	7.0	*	*	*	*	*	1.2	11.5	34.8	47.6	58.9	38.7
1999	7.2	*	*	*	*	*	1.7	11.5	35.3	51.1	57.2	49.9
Non-Hispanic black, female												
2002	2.4	*	*	*	*	*	*	2.4	7.2	14.0	17.2	13.8
2001	2.4	*	*	*	*	*	*	2.3	7.5	16.2	15.7	15.4
2000	2.6	*	*	*	*	*	*	2.7	8.5	15.2	20.1	16.6
1999	2.4	*	*	*	*	*	*	2.4	8.3	15.5	17.0	13.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of stomach (C16)												
All origins 3/, both sexes												
2002	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1
All origins 3/, male												
2002	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3
All origins 3/, female												
2002	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6
Hispanic, both sexes												
2002	3.1	*	*	*	*	0.5	1.7	4.2	11.0	21.9	47.2	72.1
2001	3.2	*	*	*	*	0.6	1.8	5.4	11.4	26.5	45.6	63.8
2000	3.1	*	*	*	*	0.4	2.0	4.8	9.8	28.4	50.2	65.7
1999	3.1	*	*	*	*	0.6	1.7	4.1	11.7	26.2	49.8	73.7
Hispanic, male												
2002	3.4	*	*	*	*	0.7	1.5	4.7	13.7	32.9	61.6	107.1
2001	3.6	*	*	*	*	*	1.9	6.2	15.0	37.1	60.4	95.2
2000	3.6	*	*	*	*	*	2.0	5.0	13.7	43.3	70.8	98.8
1999	3.5	*	*	*	*	0.7	1.6	5.2	16.4	35.6	73.5	75.5
Hispanic, female												
2002	2.7	*	*	*	*	*	1.9	3.8	8.7	13.2	37.4	54.7
2001	2.8	*	*	*	*	0.7	1.7	4.6	8.2	18.1	35.7	48.4
2000	2.7	*	*	*	*	*	2.1	4.6	6.3	16.8	36.3	49.5
1999	2.7	*	*	*	*	*	1.8	3.0	7.6	18.8	34.0	72.8
Non-Hispanic 4/, both sexes												
2002	4.4	*	*	*	0.1	0.3	0.9	2.6	6.5	14.9	27.7	46.3
2001	4.5	*	*	*	*	0.2	0.9	2.6	6.6	15.8	29.4	44.7
2000	4.7	*	*	*	*	0.3	0.9	2.7	7.2	15.8	30.1	52.0
1999	4.7	*	*	*	*	0.3	1.0	2.9	7.5	16.5	30.8	49.2
Non-Hispanic 4/, male												
2002	5.3	*	*	*	*	0.3	1.0	3.6	9.1	21.7	39.4	65.0
2001	5.4	*	*	*	*	0.3	1.0	3.6	9.8	22.3	42.1	61.0
2000	5.5	*	*	*	*	0.3	1.2	3.7	10.3	22.4	42.6	74.5
1999	5.7	*	*	*	*	0.3	1.2	4.0	10.6	23.4	45.3	71.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of stomach (C16)												
Non-Hispanic 4/, female												
2002	3.5	*	*	*	*	0.3	0.7	1.7	4.1	9.2	19.9	38.3
2001	3.6	*	*	*	*	0.2	0.8	1.7	3.7	10.4	21.0	37.9
2000	3.8	*	*	*	*	0.3	0.7	1.8	4.4	10.4	21.9	42.9
1999	3.8	*	*	*	*	0.3	0.9	1.8	4.7	10.8	21.3	40.5
Non-Hispanic white, both sexes												
2002	4.1	*	*	*	*	0.2	0.7	2.1	5.4	12.6	24.1	40.7
2001	4.2	*	*	*	*	0.2	0.7	2.1	5.6	13.5	25.8	40.0
2000	4.4	*	*	*	*	0.2	0.7	2.2	6.1	13.5	26.1	46.3
1999	4.4	*	*	*	*	0.2	0.7	2.3	6.1	14.4	27.0	43.5
Non-Hispanic white, male												
2002	5.0	*	*	*	*	0.2	0.8	2.9	7.8	18.6	35.0	58.0
2001	5.0	*	*	*	*	0.2	0.8	2.8	8.2	18.8	37.4	55.7
2000	5.2	*	*	*	*	0.2	0.8	3.0	8.9	19.3	37.4	67.3
1999	5.4	*	*	*	*	0.2	0.9	3.3	8.7	20.6	39.9	62.6
Non-Hispanic white, female												
2002	3.2	*	*	*	*	0.2	0.5	1.2	3.2	7.5	16.8	33.3
2001	3.4	*	*	*	*	*	0.6	1.3	3.1	9.0	18.1	33.5
2000	3.5	*	*	*	*	0.2	0.5	1.4	3.6	8.6	18.7	37.9
1999	3.5	*	*	*	*	0.2	0.6	1.4	3.7	9.1	18.6	35.9
Non-Hispanic black, both sexes												
2002	5.8	*	*	*	*	0.6	1.6	5.5	13.2	31.5	57.7	100.1
2001	5.8	*	*	*	*	0.5	2.2	5.5	13.1	31.0	59.1	89.3
2000	6.1	*	*	*	*	0.7	2.1	5.8	14.1	32.3	64.3	102.7
1999	6.3	*	*	*	*	0.7	2.3	6.0	16.1	31.5	65.1	103.1
Non-Hispanic black, male												
2002	6.9	*	*	*	*	0.8	1.9	7.9	19.5	47.3	80.9	136.7
2001	6.9	*	*	*	*	*	2.4	8.0	20.0	49.2	87.9	106.3
2000	7.2	*	*	*	*	0.8	3.0	8.1	20.9	47.0	91.0	138.6
1999	7.6	*	*	*	*	*	2.8	8.7	23.3	46.4	101.6	149.8
Non-Hispanic black, female												
2002	4.9	*	*	*	*	*	1.4	3.4	8.1	20.3	44.6	86.1
2001	4.7	*	*	*	*	*	2.0	3.3	7.6	18.1	43.0	83.0
2000	5.2	*	*	*	*	*	1.4	3.9	8.6	22.0	49.3	89.4
1999	5.1	*	*	*	*	*	1.9	3.7	10.2	21.0	44.6	85.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of small intestine (C17)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.3	2.8
2001	0.4	*	*	*	*	*	0.1	0.3	0.8	1.4	2.5	3.2
2000	0.4	*	*	*	*	*	0.1	0.3	0.8	1.5	2.4	3.0
1999	0.4	*	*	*	*	*	0.1	0.4	0.6	1.4	2.4	3.2
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.1	0.3	0.8	1.6	2.8	3.7
2001	0.4	*	*	*	*	*	0.1	0.4	0.9	1.9	3.0	4.1
2000	0.4	*	*	*	*	*	0.1	0.4	0.9	1.8	3.0	4.2
1999	0.4	*	*	*	*	*	0.1	0.5	0.9	1.7	3.2	4.4
All origins 3/, female												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.0	2.4
2001	0.4	*	*	*	*	*	0.1	0.2	0.7	1.1	2.1	2.8
2000	0.4	*	*	*	*	*	0.1	0.3	0.6	1.2	2.1	2.5
1999	0.3	*	*	*	*	*	*	0.3	0.4	1.2	1.8	2.7
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.4	2.8
2001	0.4	*	*	*	*	*	0.1	0.3	0.8	1.5	2.5	3.2
2000	0.4	*	*	*	*	*	0.1	0.3	0.8	1.5	2.5	3.0
1999	0.4	*	*	*	*	*	0.1	0.4	0.6	1.4	2.4	3.2
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	0.1	0.4	0.9	1.6	2.9	3.8
2001	0.4	*	*	*	*	*	0.1	0.4	0.9	1.9	3.0	4.1
2000	0.4	*	*	*	*	*	0.1	0.4	0.9	1.8	3.0	4.2
1999	0.5	*	*	*	*	*	0.1	0.5	0.9	1.7	3.3	4.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of small intestine (C17)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	*	0.2	0.5	1.3	2.1	2.4
2001	0.4	*	*	*	*	*	*	0.2	0.6	1.1	2.1	2.8
2000	0.4	*	*	*	*	*	*	0.3	0.7	1.2	2.1	2.5
1999	0.4	*	*	*	*	*	*	0.3	0.4	1.2	1.9	2.7
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.6	1.4	2.3	2.7
2001	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.4	3.0
2000	0.4	*	*	*	*	*	0.1	0.3	0.8	1.4	2.4	3.1
1999	0.4	*	*	*	*	*	0.1	0.4	0.6	1.4	2.3	3.0
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	*	0.3	0.7	1.6	2.8	3.6
2001	0.4	*	*	*	*	*	0.1	0.3	0.8	1.8	2.9	3.7
2000	0.5	*	*	*	*	*	*	0.3	0.9	1.8	2.8	4.3
1999	0.5	*	*	*	*	*	*	0.5	0.8	1.7	3.1	4.3
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	0.2	0.5	1.3	2.0	2.3
2001	0.4	*	*	*	*	*	*	0.2	0.6	1.1	2.1	2.7
2000	0.4	*	*	*	*	*	*	0.2	0.7	1.1	2.1	2.5
1999	0.4	*	*	*	*	*	*	0.3	0.4	1.2	1.7	2.5
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.6	1.3	1.8	3.5	*
2001	0.5	*	*	*	*	*	*	0.6	1.5	2.3	3.1	*
2000	0.4	*	*	*	*	*	*	0.6	1.1	2.0	2.8	*
1999	0.4	*	*	*	*	*	*	0.7	1.0	1.7	4.5	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	1.9	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	6.3	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	2.1	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	3.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of colon, rectum and anus (C18-C21)												
All origins 3/, both sexes												
2002	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
All origins 3/, male												
2002	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
All origins 3/, female												
2002	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.9
2000	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6
Hispanic, both sexes												
2002	6.1	*	*	*	*	0.4	2.1	7.7	23.0	51.2	95.6	165.7
2001	6.1	*	*	*	*	0.3	2.3	7.2	22.5	52.2	96.5	194.5
2000	6.0	*	*	*	*	0.5	2.3	8.4	23.1	49.2	98.2	181.1
1999	6.0	*	*	*	*	0.5	1.7	8.3	21.0	56.1	101.0	176.8
Hispanic, male												
2002	6.4	*	*	*	*	*	2.0	9.3	31.1	67.1	121.1	174.0
2001	6.2	*	*	*	*	*	2.7	7.4	28.2	66.5	116.7	257.1
2000	6.2	*	*	*	*	*	2.0	9.0	28.2	63.5	131.3	235.8
1999	6.1	*	*	*	*	*	1.5	9.1	24.1	72.7	134.0	211.4
Hispanic, female												
2002	5.7	*	*	*	*	*	2.1	6.1	15.7	38.6	78.1	161.6
2001	5.9	*	*	*	*	*	1.9	7.1	17.4	40.9	82.8	163.8
2000	5.8	*	*	*	*	*	2.7	7.8	18.5	38.0	75.9	154.3
1999	5.9	*	*	*	*	*	2.0	7.5	18.3	43.0	79.1	160.1
Non-Hispanic 4/, both sexes												
2002	21.7	*	*	*	0.1	0.8	3.3	11.8	32.3	74.6	139.1	244.0
2001	22.0	*	*	*	0.1	0.9	3.3	12.0	32.2	76.7	141.1	257.4
2000	22.4	*	*	*	0.1	0.7	3.1	11.5	33.3	78.2	147.8	273.1
1999	22.4	*	*	*	0.1	0.7	3.1	11.3	34.3	78.5	148.1	274.3
Non-Hispanic 4/, male												
2002	22.3	*	*	*	0.2	0.9	3.8	13.4	39.9	92.7	169.3	279.2
2001	22.3	*	*	*	0.1	0.9	3.6	13.6	38.9	95.2	171.8	299.6
2000	22.8	*	*	*	0.1	0.8	3.6	13.1	40.3	97.2	181.3	318.6
1999	22.7	*	*	*	0.1	0.9	3.2	12.8	41.6	96.2	183.2	327.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of colon, rectum and anus (C18-C21)												
Non-Hispanic 4/, female												
2002	21.2	*	*	*	*	0.7	2.9	10.2	25.2	59.6	119.1	228.8
2001	21.7	*	*	*	*	0.8	3.0	10.5	25.9	61.3	121.0	239.9
2000	22.1	*	*	*	*	0.6	2.7	9.9	26.9	62.6	126.1	254.7
1999	22.2	*	*	*	*	0.6	3.0	9.9	27.6	63.9	125.3	253.0
Non-Hispanic white, both sexes												
2002	23.2	*	*	*	0.1	0.8	3.0	10.9	30.6	73.0	136.8	243.2
2001	23.5	*	*	*	0.1	0.8	3.1	11.1	30.6	74.7	138.5	256.9
2000	24.0	*	*	*	0.1	0.6	2.8	10.6	31.7	76.5	145.7	271.9
1999	24.0	*	*	*	0.1	0.7	2.9	10.6	32.8	76.7	145.9	273.4
Non-Hispanic white, male												
2002	23.8	*	*	*	0.2	0.9	3.4	12.5	38.0	91.1	165.8	277.6
2001	23.9	*	*	*	*	0.9	3.3	12.8	37.4	93.2	169.3	298.2
2000	24.2	*	*	*	*	0.8	3.2	12.2	38.2	95.1	178.4	315.8
1999	24.3	*	*	*	*	0.8	3.1	12.1	39.7	94.8	180.6	328.2
Non-Hispanic white, female												
2002	22.7	*	*	*	*	0.7	2.7	9.4	23.7	57.6	117.3	228.3
2001	23.1	*	*	*	*	0.7	2.8	9.5	24.2	59.0	118.1	239.8
2000	23.8	*	*	*	*	0.5	2.4	9.0	25.5	60.9	124.1	254.1
1999	23.7	*	*	*	*	0.6	2.7	9.0	26.2	61.5	123.2	251.5
Non-Hispanic black, both sexes												
2002	18.9	*	*	*	*	0.7	5.1	19.0	50.5	100.6	185.9	278.3
2001	19.2	*	*	*	*	1.1	5.0	19.5	50.6	105.9	189.7	289.0
2000	19.4	*	*	*	*	1.0	5.2	20.0	51.1	106.3	192.5	312.9
1999	19.3	*	*	*	*	1.1	4.9	18.1	53.7	107.2	194.5	308.8
Non-Hispanic black, male												
2002	19.3	*	*	*	*	0.9	6.0	21.8	62.7	123.0	240.9	332.6
2001	19.0	*	*	*	*	1.2	5.8	21.4	59.3	130.7	229.9	352.3
2000	19.8	*	*	*	*	1.1	6.1	22.7	63.7	133.7	244.4	381.6
1999	19.2	*	*	*	*	1.2	4.8	20.6	67.0	127.8	245.8	351.8
Non-Hispanic black, female												
2002	18.4	*	*	*	*	*	4.3	16.7	40.7	84.7	154.9	257.5
2001	19.4	*	*	*	*	1.0	4.3	17.9	43.6	88.4	167.1	265.6
2000	19.1	*	*	*	*	1.0	4.4	17.6	40.9	87.1	163.2	287.4
1999	19.5	*	*	*	*	0.9	5.0	15.9	43.1	92.7	165.8	292.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Colon (C18)												
All origins 3/, both sexes												
2002	16.5	*	*	*	0.1	0.5	2.4	8.9	25.3	61.0	118.5	211.3
2001	16.8	*	*	*	0.1	0.6	2.4	8.9	25.7	63.5	120.1	221.6
2000	17.3	*	*	*	0.1	0.5	2.3	8.9	26.5	64.3	126.5	236.6
1999	17.4	*	*	*	0.1	0.5	2.3	8.8	27.4	65.1	127.0	238.1
All origins 3/, male												
2002	16.6	*	*	*	0.1	0.6	2.6	9.9	30.9	75.5	142.6	240.2
2001	16.7	*	*	*	*	0.6	2.6	9.9	30.6	77.7	144.4	256.7
2000	17.1	*	*	*	*	0.6	2.6	9.9	31.4	78.6	154.0	275.8
1999	17.2	*	*	*	0.1	0.6	2.3	9.8	32.7	79.2	156.1	282.2
All origins 3/, female												
2002	16.4	*	*	*	*	0.5	2.1	7.9	20.2	49.0	102.6	198.7
2001	16.9	*	*	*	0.1	0.6	2.3	8.0	21.2	51.8	104.2	206.9
2000	17.4	*	*	*	*	0.5	2.1	7.9	22.0	52.5	108.5	220.6
1999	17.5	*	*	*	*	0.5	2.3	7.9	22.5	53.5	108.2	220.4
Hispanic, both sexes												
2002	5.1	*	*	*	*	*	1.7	6.0	18.1	43.1	84.4	145.8
2001	5.0	*	*	*	*	*	1.7	5.4	18.4	43.4	81.7	173.2
2000	5.0	*	*	*	*	0.4	1.8	6.7	18.1	42.0	83.6	160.6
1999	5.0	*	*	*	*	0.4	1.4	6.9	16.6	46.8	88.3	154.3
Hispanic, male												
2002	5.3	*	*	*	*	*	1.6	7.1	23.3	56.6	108.3	153.9
2001	5.0	*	*	*	*	*	1.8	5.2	22.2	54.9	96.6	220.9
2000	5.1	*	*	*	*	*	1.6	7.2	20.9	52.3	116.0	211.6
1999	5.0	*	*	*	*	*	1.2	7.4	19.4	58.9	118.4	185.5
Hispanic, female												
2002	4.9	*	*	*	*	*	1.8	4.9	13.5	32.4	68.0	141.7
2001	5.1	*	*	*	*	*	1.6	5.7	14.9	34.3	71.6	149.9
2000	4.8	*	*	*	*	*	2.1	6.2	15.5	33.8	61.7	135.5
1999	5.0	*	*	*	*	*	1.7	6.5	14.2	37.3	68.3	139.3
Non-Hispanic 4/, both sexes												
2002	18.2	*	*	*	0.1	0.6	2.5	9.2	25.8	62.1	119.9	213.5
2001	18.5	*	*	*	0.1	0.7	2.5	9.2	26.2	64.7	121.7	222.9
2000	19.0	*	*	*	0.1	0.6	2.4	9.1	27.1	65.6	128.1	238.9
1999	19.0	*	*	*	0.1	0.6	2.4	9.0	28.1	66.0	128.2	240.4
Non-Hispanic 4/, male												
2002	18.4	*	*	*	0.2	0.6	2.8	10.2	31.4	76.6	144.0	243.7
2001	18.5	*	*	*	*	0.7	2.7	10.3	31.1	78.9	146.5	257.8
2000	18.9	*	*	*	*	0.6	2.7	10.1	32.1	80.1	155.3	278.2
1999	19.0	*	*	*	*	0.6	2.5	10.0	33.5	80.2	157.1	285.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Colon (C18)												
Non-Hispanic 4/, female												
2002	18.1	*	*	*	*	0.5	2.2	8.2	20.7	50.1	104.0	200.5
2001	18.6	*	*	*	*	0.6	2.4	8.2	21.6	52.9	105.4	208.4
2000	19.1	*	*	*	*	0.5	2.1	8.0	22.3	53.6	110.4	223.0
1999	19.0	*	*	*	*	0.5	2.3	8.0	23.0	54.3	109.5	222.4
Non-Hispanic white, both sexes												
2002	19.4	*	*	*	0.1	0.6	2.3	8.4	24.3	60.4	117.7	212.6
2001	19.7	*	*	*	0.1	0.6	2.3	8.4	24.6	62.6	119.0	222.4
2000	20.2	*	*	*	*	0.5	2.1	8.2	25.5	64.0	125.8	237.3
1999	20.3	*	*	*	0.1	0.5	2.2	8.3	26.6	64.3	126.0	239.3
Non-Hispanic white, male												
2002	19.5	*	*	*	*	0.7	2.6	9.4	29.6	74.9	140.8	241.9
2001	19.7	*	*	*	*	0.7	2.5	9.5	29.7	77.0	143.6	256.3
2000	20.1	*	*	*	*	0.6	2.4	9.4	30.3	78.4	152.3	274.5
1999	20.3	*	*	*	*	0.5	2.4	9.5	31.9	79.0	154.6	285.9
Non-Hispanic white, female												
2002	19.3	*	*	*	*	0.5	2.0	7.4	19.3	48.1	102.3	200.0
2001	19.7	*	*	*	*	0.5	2.1	7.4	19.8	50.5	102.7	208.3
2000	20.4	*	*	*	*	0.4	1.8	7.1	21.1	51.9	108.4	222.2
1999	20.2	*	*	*	*	0.5	2.0	7.1	21.6	52.0	107.4	220.7
Non-Hispanic black, both sexes												
2002	16.2	*	*	*	*	0.6	3.9	15.9	42.5	87.3	163.8	247.4
2001	16.7	*	*	*	*	0.9	4.1	16.1	43.5	93.3	167.8	252.2
2000	16.8	*	*	*	*	0.8	4.2	16.7	43.3	91.1	173.2	282.4
1999	16.8	*	*	*	*	0.9	4.0	15.3	45.9	93.0	173.0	276.4
Non-Hispanic black, male												
2002	16.4	*	*	*	*	*	4.5	17.5	52.2	106.1	209.8	296.5
2001	16.2	*	*	*	*	0.9	4.3	17.7	49.0	111.5	204.0	311.8
2000	16.8	*	*	*	*	*	4.7	18.4	52.8	110.3	218.8	351.1
1999	16.3	*	*	*	*	0.9	3.7	16.7	55.6	109.6	215.8	306.6
Non-Hispanic black, female												
2002	16.1	*	*	*	*	*	3.5	14.6	34.7	73.9	137.8	228.7
2001	17.1	*	*	*	*	0.9	3.9	14.7	39.1	80.5	147.4	230.1
2000	16.9	*	*	*	*	0.8	3.7	15.2	35.7	77.5	147.6	256.9
1999	17.3	*	*	*	*	0.9	4.3	14.0	38.0	81.3	149.0	265.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)												
All origins 3/, both sexes												
2002	3.0	*	*	*	*	0.2	0.7	2.2	5.9	11.7	18.0	29.0
2001	3.0	*	*	*	*	0.2	0.7	2.5	5.5	11.2	18.3	32.6
2000	3.0	*	*	*	*	0.1	0.6	2.1	5.8	11.8	18.8	32.6
1999	3.0	*	*	*	*	0.2	0.6	2.0	5.8	11.7	18.8	32.1
All origins 3/, male												
2002	3.4	*	*	*	*	0.2	0.8	2.9	8.1	15.3	24.2	33.6
2001	3.4	*	*	*	*	0.2	0.8	3.0	7.4	15.5	24.4	40.7
2000	3.4	*	*	*	*	0.2	0.7	2.6	7.8	16.3	24.8	38.9
1999	3.3	*	*	*	*	0.2	0.6	2.5	7.6	15.4	25.1	40.7
All origins 3/, female												
2002	2.6	*	*	*	*	0.2	0.5	1.7	3.8	8.7	13.9	27.0
2001	2.6	*	*	*	*	0.1	0.5	1.9	3.7	7.6	14.4	29.2
2000	2.6	*	*	*	*	0.1	0.5	1.6	4.0	8.1	14.8	30.1
1999	2.7	*	*	*	*	0.1	0.5	1.6	4.1	8.7	14.7	28.7
Hispanic, both sexes												
2002	0.9	*	*	*	*	*	*	1.6	4.5	7.8	10.5	18.3
2001	1.0	*	*	*	*	*	0.6	1.8	4.0	8.1	14.1	21.3
2000	1.0	*	*	*	*	*	0.4	1.7	4.7	6.9	14.0	19.9
1999	0.9	*	*	*	*	*	*	1.2	4.3	9.1	12.5	21.0
Hispanic, male												
2002	1.1	*	*	*	*	*	*	2.0	7.2	10.1	12.3	*
2001	1.2	*	*	*	*	*	0.8	2.2	6.0	11.2	20.1	*
2000	1.0	*	*	*	*	*	*	1.8	6.8	10.6	14.8	*
1999	1.0	*	*	*	*	*	*	1.7	4.7	13.8	15.6	*
Hispanic, female												
2002	0.8	*	*	*	*	*	*	1.1	2.0	5.9	9.3	17.4
2001	0.8	*	*	*	*	*	*	1.3	2.2	5.7	10.0	*
2000	0.9	*	*	*	*	*	*	1.5	2.9	4.0	13.5	*
1999	0.8	*	*	*	*	*	*	*	4.0	5.3	10.4	*
Non-Hispanic 4/, both sexes												
2002	3.3	*	*	*	*	0.2	0.7	2.3	6.0	11.9	18.3	29.3
2001	3.3	*	*	*	*	0.2	0.7	2.5	5.6	11.4	18.5	33.0
2000	3.3	*	*	*	*	0.1	0.6	2.2	5.9	12.1	18.9	33.0
1999	3.2	*	*	*	*	0.2	0.6	2.1	5.9	11.9	19.0	32.5
Non-Hispanic 4/, male												
2002	3.8	*	*	*	*	0.3	0.9	2.9	8.2	15.6	24.7	34.2
2001	3.7	*	*	*	*	0.2	0.8	3.1	7.5	15.7	24.6	40.9
2000	3.7	*	*	*	*	0.2	0.8	2.7	7.8	16.6	25.1	39.3
1999	3.6	*	*	*	*	0.2	0.7	2.6	7.8	15.5	25.4	41.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)												
Non-Hispanic 4/, female												
2002	2.9	*	*	*	*	0.2	0.6	1.7	4.0	8.8	14.1	27.2
2001	2.9	*	*	*	*	0.2	0.5	2.0	3.8	7.8	14.5	29.7
2000	2.9	*	*	*	*	0.1	0.5	1.6	4.1	8.3	14.8	30.5
1999	2.9	*	*	*	*	*	0.6	1.7	4.1	8.9	14.8	29.0
Non-Hispanic white, both sexes												
2002	3.6	*	*	*	*	0.2	0.7	2.3	5.9	12.0	18.1	29.3
2001	3.6	*	*	*	*	0.2	0.7	2.5	5.6	11.4	18.5	33.0
2000	3.5	*	*	*	*	0.1	0.6	2.1	5.7	11.9	19.0	33.5
1999	3.5	*	*	*	*	0.2	0.6	2.1	5.8	11.9	19.0	32.6
Non-Hispanic white, male												
2002	4.0	*	*	*	*	0.2	0.8	2.8	8.0	15.8	24.4	34.3
2001	4.0	*	*	*	*	0.2	0.8	3.1	7.3	15.6	25.0	41.1
2000	4.0	*	*	*	*	*	0.7	2.6	7.6	16.2	25.4	40.3
1999	3.9	*	*	*	*	0.2	0.6	2.4	7.6	15.4	25.3	41.2
Non-Hispanic white, female												
2002	3.1	*	*	*	*	0.2	0.5	1.7	3.8	8.8	13.9	27.2
2001	3.1	*	*	*	*	0.2	0.6	1.8	3.9	7.9	14.3	29.6
2000	3.1	*	*	*	*	*	0.4	1.6	4.0	8.3	14.9	30.7
1999	3.2	*	*	*	*	*	0.6	1.7	4.1	8.9	14.9	29.2
Non-Hispanic black, both sexes												
2002	2.5	*	*	*	*	*	1.0	2.9	7.6	12.8	21.5	30.2
2001	2.4	*	*	*	*	*	0.8	3.1	6.8	12.1	20.9	35.9
2000	2.5	*	*	*	*	*	0.9	3.0	7.4	14.7	18.3	28.9
1999	2.4	*	*	*	*	*	0.7	2.5	7.7	13.4	20.8	32.0
Non-Hispanic black, male												
2002	2.8	*	*	*	*	*	1.2	4.0	10.2	16.4	30.2	35.0
2001	2.8	*	*	*	*	*	1.3	3.5	9.9	18.7	24.9	39.3
2000	2.9	*	*	*	*	*	1.3	3.9	10.5	22.8	24.4	29.4
1999	2.8	*	*	*	*	*	0.8	3.5	11.1	17.7	29.7	45.2
Non-Hispanic black, female												
2002	2.2	*	*	*	*	*	0.7	2.0	5.5	10.2	16.6	28.4
2001	2.1	*	*	*	*	*	*	2.8	4.3	7.5	18.7	34.6
2000	2.0	*	*	*	*	*	*	2.2	4.9	9.1	14.9	28.8
1999	2.0	*	*	*	*	*	*	1.6	4.9	10.3	15.8	27.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anus and anus canal (C21)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.2
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.4
2000	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.8	1.2
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.4
All origins 3/, male												
2002	0.2	*	*	*	*	*	0.1	0.3	0.4	0.5	0.6	*
2001	0.1	*	*	*	*	*	0.1	0.2	0.3	0.6	0.7	*
2000	0.1	*	*	*	*	*	0.1	0.2	0.4	0.5	0.8	*
1999	0.1	*	*	*	*	*	*	0.2	0.3	0.4	0.6	*
All origins 3/, female												
2002	0.2	*	*	*	*	*	0.1	0.3	0.5	0.6	1.0	1.2
2001	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.1	1.7
2000	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.3
1999	0.2	*	*	*	*	*	*	0.2	0.4	0.7	0.9	1.6
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.2
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.5
2000	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.8	1.2
1999	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.8	1.4
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	0.1	0.3	0.3	0.5	0.6	*
2001	0.1	*	*	*	*	*	*	0.2	0.3	0.6	0.7	*
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.5	0.8	*
1999	0.1	*	*	*	*	*	*	0.2	0.3	0.4	0.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anus and anus canal (C21)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	0.3	0.5	0.7	1.0	1.1
2001	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.1	1.7
2000	0.2	*	*	*	*	*	0.1	0.3	0.4	0.7	0.9	1.3
1999	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.0	1.6
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.9	1.2
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.5
2000	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	0.8	1.2
1999	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.5
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.3	0.3	0.5	0.6	*
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.6	0.7	*
2000	0.2	*	*	*	*	*	*	0.2	0.3	0.5	0.8	*
1999	0.1	*	*	*	*	*	*	0.2	0.3	0.5	0.7	*
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.3	0.5	0.7	1.1	1.2
2001	0.3	*	*	*	*	*	0.1	0.3	0.5	0.7	1.1	1.8
2000	0.3	*	*	*	*	*	*	0.3	0.5	0.7	0.9	1.2
1999	0.2	*	*	*	*	*	*	0.2	0.5	0.6	1.0	1.7
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)												
All origins 3/, both sexes												
2002	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6
All origins 3/, male												
2002	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1
All origins 3/, female												
2002	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8
Hispanic, both sexes												
2002	3.5	*	*	*	*	*	0.8	5.9	15.7	34.1	44.2	59.9
2001	3.4	*	*	*	*	*	0.8	6.7	14.2	28.9	52.1	52.5
2000	3.3	*	*	*	*	*	1.2	5.6	13.7	31.1	51.8	58.4
1999	3.2	*	*	*	*	*	0.9	5.7	13.1	31.3	49.8	53.3
Hispanic, male												
2002	4.5	*	*	*	*	*	1.1	9.8	24.1	48.9	59.9	75.3
2001	4.3	*	*	*	*	*	1.3	11.1	21.5	41.3	68.0	64.8
2000	4.2	*	*	*	*	*	1.7	9.1	20.5	43.9	74.7	72.6
1999	3.9	*	*	*	*	*	1.1	9.6	20.7	42.8	63.1	66.9
Hispanic, female												
2002	2.5	*	*	*	*	*	*	2.1	8.2	22.4	33.4	52.2
2001	2.5	*	*	*	*	*	*	2.3	7.7	19.1	41.4	46.5
2000	2.4	*	*	*	*	*	*	2.2	7.6	21.1	36.3	51.4
1999	2.4	*	*	*	*	*	*	1.9	6.3	22.2	40.9	46.8
Non-Hispanic 4/, both sexes												
2002	5.1	*	0.2	*	0.1	0.3	0.9	4.8	9.1	18.6	28.4	33.9
2001	4.9	*	*	*	0.1	0.2	0.9	4.6	8.9	18.1	27.9	33.2
2000	4.8	*	0.2	*	0.1	0.2	0.9	4.3	8.9	17.9	27.7	31.7
1999	4.6	*	0.2	*	0.1	0.2	1.0	3.9	8.4	18.2	27.1	29.7
Non-Hispanic 4/, male												
2002	6.6	*	*	*	*	0.4	1.4	8.0	13.7	27.1	40.5	44.8
2001	6.3	*	*	*	*	0.2	1.3	7.4	13.2	25.8	39.7	49.7
2000	6.2	*	*	*	*	0.3	1.3	6.9	13.3	25.9	40.5	43.4
1999	5.8	*	*	*	0.1	0.3	1.5	6.2	12.2	26.5	37.9	38.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)												
Non-Hispanic 4/, female												
2002	3.6	*	*	*	*	0.2	0.5	1.7	4.8	11.4	20.5	29.2
2001	3.5	*	*	*	*	0.2	0.4	1.9	4.8	11.7	20.2	26.4
2000	3.4	*	*	*	*	0.1	0.6	1.9	4.9	11.2	19.4	27.0
1999	3.4	*	*	*	*	0.2	0.5	1.8	4.9	11.3	20.0	26.1
Non-Hispanic white, both sexes												
2002	5.0	*	*	*	*	0.2	0.7	3.9	7.7	17.0	27.4	32.6
2001	4.8	*	*	*	0.1	0.1	0.6	3.7	7.6	16.6	26.5	32.4
2000	4.6	*	*	*	*	0.1	0.6	3.4	7.5	16.2	26.5	30.8
1999	4.5	*	0.2	*	0.1	0.1	0.7	3.1	7.1	16.8	26.1	28.9
Non-Hispanic white, male												
2002	6.4	*	*	*	*	0.2	1.1	6.4	11.5	24.6	38.7	43.6
2001	6.1	*	*	*	*	*	0.9	5.9	11.2	24.2	38.0	48.9
2000	5.9	*	*	*	*	0.2	0.8	5.3	11.0	23.6	39.2	41.6
1999	5.7	*	*	*	*	*	1.1	4.8	10.2	24.6	36.7	39.0
Non-Hispanic white, female												
2002	3.6	*	*	*	*	0.2	0.4	1.4	4.2	10.4	19.9	27.8
2001	3.4	*	*	*	*	*	0.3	1.5	4.1	10.2	18.9	25.5
2000	3.4	*	*	*	*	*	0.4	1.6	4.3	10.1	18.2	26.5
1999	3.4	*	*	*	*	*	0.4	1.6	4.2	10.2	19.1	24.8
Non-Hispanic black, both sexes												
2002	4.7	*	*	*	*	0.5	1.5	8.9	15.0	23.3	30.9	38.1
2001	4.6	*	*	*	*	*	1.7	8.9	15.1	21.9	32.3	35.3
2000	4.5	*	*	*	*	0.4	1.8	8.8	14.5	23.7	29.1	31.8
1999	4.3	*	*	*	*	0.4	1.7	7.4	13.4	24.6	29.7	34.9
Non-Hispanic black, male												
2002	6.6	*	*	*	*	*	2.3	15.4	24.4	38.2	46.3	41.6
2001	6.1	*	*	*	*	*	2.7	15.1	24.2	30.3	42.6	42.7
2000	6.3	*	*	*	*	*	2.5	15.0	24.1	37.3	40.4	48.1
1999	5.7	*	*	*	*	*	2.5	13.1	20.3	36.8	41.6	30.9
Non-Hispanic black, female												
2002	3.0	*	*	*	*	*	0.8	3.3	7.4	12.8	22.2	36.8
2001	3.3	*	*	*	*	*	0.8	3.6	7.8	16.0	26.4	32.5
2000	2.9	*	*	*	*	*	1.2	3.4	6.7	14.1	22.8	25.7
1999	3.1	*	*	*	*	*	0.9	2.5	7.9	15.9	23.1	36.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)												
All origins 3/, both sexes												
2002	3.8	*	0.2	*	0.1	0.2	0.7	4.3	7.6	15.7	22.2	24.7
2001	3.7	*	*	*	0.1	0.2	0.7	4.2	7.5	14.7	21.9	24.2
2000	3.6	*	0.1	0.0	0.1	0.2	0.8	3.8	7.5	14.8	22.1	23.4
1999	3.5	*	0.2	*	0.1	0.2	0.8	3.5	7.0	15.2	21.5	22.5
All origins 3/, male												
2002	5.3	*	*	*	*	0.3	1.2	7.4	12.3	24.0	32.7	34.6
2001	5.1	*	*	*	*	0.2	1.1	7.1	12.0	22.0	32.2	38.5
2000	5.0	*	*	*	*	0.2	1.2	6.4	11.9	22.4	33.7	33.9
1999	4.7	*	0.3	*	0.1	0.2	1.2	5.9	10.9	23.1	31.5	31.2
All origins 3/, female												
2002	2.4	*	*	*	*	0.1	0.3	1.3	3.2	8.7	15.2	20.4
2001	2.4	*	*	*	*	0.1	0.3	1.4	3.4	8.6	15.1	18.2
2000	2.3	*	*	*	*	0.1	0.4	1.3	3.5	8.5	14.4	19.1
1999	2.4	*	*	*	*	0.1	0.4	1.2	3.5	8.6	15.0	18.9
Hispanic, both sexes												
2002	3.0	*	*	*	*	*	0.7	5.6	13.2	29.1	37.4	49.9
2001	2.9	*	*	*	*	*	0.7	6.2	12.6	24.6	42.2	40.6
2000	2.8	*	*	*	*	*	1.0	4.9	11.6	25.7	43.9	51.1
1999	2.7	*	*	*	*	*	0.6	5.1	11.6	26.5	40.8	45.6
Hispanic, male												
2002	4.0	*	*	*	*	*	1.0	9.5	21.3	43.6	52.9	63.6
2001	3.9	*	*	*	*	*	1.1	10.6	20.4	36.1	57.2	59.0
2000	3.8	*	*	*	*	*	1.6	8.4	18.7	38.0	65.9	66.5
1999	3.6	*	*	*	*	*	0.9	9.3	19.5	38.7	53.7	58.2
Hispanic, female												
2002	2.0	*	*	*	*	*	*	1.8	5.9	17.6	26.7	43.1
2001	1.9	*	*	*	*	*	*	1.9	5.7	15.4	32.0	31.6
2000	1.8	*	*	*	*	*	*	1.5	5.3	16.1	29.1	43.5
1999	1.8	*	*	*	*	*	*	*	4.6	17.0	32.3	39.5
Non-Hispanic 4/, both sexes												
2002	3.9	*	0.2	*	0.1	0.2	0.7	4.1	7.1	14.7	21.4	23.6
2001	3.8	*	*	*	0.1	0.2	0.7	4.0	7.1	14.0	20.9	23.5
2000	3.7	*	0.2	*	0.1	0.2	0.8	3.7	7.2	14.1	21.1	22.3
1999	3.6	*	0.2	*	0.1	0.2	0.8	3.4	6.7	14.4	20.7	21.6
Non-Hispanic 4/, male												
2002	5.5	*	*	*	*	0.3	1.2	7.2	11.5	22.6	31.6	33.2
2001	5.2	*	*	*	*	0.2	1.1	6.7	11.3	21.0	31.0	37.6
2000	5.1	*	*	*	*	0.2	1.1	6.2	11.4	21.4	32.2	32.4
1999	4.9	*	*	*	0.1	0.2	1.3	5.6	10.2	22.1	30.5	29.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)												
Non-Hispanic 4/, female												
2002	2.5	*	*	*	*	0.2	0.3	1.2	3.0	8.1	14.6	19.4
2001	2.4	*	*	*	*	0.1	0.3	1.3	3.2	8.1	14.3	17.6
2000	2.4	*	*	*	*	0.1	0.4	1.3	3.3	8.0	13.8	18.2
1999	2.4	*	*	*	*	0.1	0.4	1.2	3.4	8.1	14.3	18.3
Non-Hispanic white, both sexes												
2002	3.8	*	*	*	*	0.1	0.5	3.3	5.8	13.1	20.4	22.4
2001	3.6	*	*	*	*	0.1	0.4	3.1	5.8	12.6	19.7	22.7
2000	3.5	*	*	*	*	0.1	0.5	2.8	5.9	12.4	19.8	21.3
1999	3.4	*	0.2	*	0.1	*	0.6	2.6	5.4	13.0	19.7	20.6
Non-Hispanic white, male												
2002	5.2	*	*	*	*	*	0.8	5.6	9.4	20.3	30.1	32.0
2001	5.0	*	*	*	*	*	0.7	5.2	9.4	19.5	29.5	36.8
2000	4.8	*	*	*	*	*	0.7	4.6	9.2	19.0	30.8	30.1
1999	4.6	*	*	*	*	*	0.9	4.1	8.2	20.1	29.4	30.0
Non-Hispanic white, female												
2002	2.4	*	*	*	*	*	0.2	0.9	2.5	7.1	14.0	18.3
2001	2.3	*	*	*	*	*	0.2	1.0	2.5	6.7	13.2	16.8
2000	2.3	*	*	*	*	*	0.3	1.0	2.8	6.8	12.6	17.7
1999	2.3	*	*	*	*	*	0.3	1.0	2.7	7.0	13.3	16.9
Non-Hispanic black, both sexes												
2002	4.1	*	*	*	*	0.4	1.3	8.3	13.2	20.0	24.4	28.7
2001	4.0	*	*	*	*	*	1.6	8.1	13.1	18.5	26.3	27.2
2000	3.9	*	*	*	*	*	1.6	8.1	12.4	20.4	24.3	24.8
1999	3.8	*	*	*	*	*	1.4	7.0	11.8	21.3	24.6	28.8
Non-Hispanic black, male												
2002	6.0	*	*	*	*	*	2.1	14.8	22.7	33.2	37.2	35.0
2001	5.5	*	*	*	*	*	2.6	14.4	21.5	26.0	35.7	34.6
2000	5.7	*	*	*	*	*	2.2	14.3	22.1	33.2	34.9	42.3
1999	5.2	*	*	*	*	*	2.3	12.8	19.1	32.8	35.0	25.0
Non-Hispanic black, female												
2002	2.3	*	*	*	*	*	*	2.7	5.5	10.6	17.1	26.3
2001	2.6	*	*	*	*	*	*	2.7	6.4	13.3	21.1	24.4
2000	2.3	*	*	*	*	*	1.1	2.7	4.7	11.4	18.4	18.3
1999	2.5	*	*	*	*	*	*	2.1	6.0	13.1	18.7	30.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)												
All origins 3/, both sexes												
2002	1.0	*	*	*	*	0.1	0.2	0.6	2.0	3.9	7.1	10.3
2001	1.0	*	*	*	*	*	0.2	0.6	1.8	4.1	7.1	9.9
2000	1.0	*	*	*	*	*	0.2	0.7	1.8	3.9	6.7	9.3
1999	0.9	*	*	*	*	*	0.2	0.6	1.7	3.8	6.5	8.1
All origins 3/, male												
2002	1.0	*	*	*	*	*	0.2	0.8	2.2	4.5	8.8	11.6
2001	1.0	*	*	*	*	*	0.2	0.7	1.9	4.8	8.8	11.9
2000	1.0	*	*	*	*	*	0.2	0.7	1.9	4.6	8.3	10.8
1999	0.9	*	*	*	*	*	0.2	0.6	1.9	4.4	7.5	8.9
All origins 3/, female												
2002	1.0	*	*	*	*	*	0.2	0.5	1.8	3.4	5.9	9.8
2001	1.0	*	*	*	*	*	0.1	0.5	1.6	3.6	6.0	9.0
2000	1.0	*	*	*	*	*	0.1	0.6	1.7	3.3	5.7	8.8
1999	0.9	*	*	*	*	*	0.2	0.6	1.6	3.3	5.8	7.8
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	*	2.5	5.0	6.8	*
2001	0.5	*	*	*	*	*	*	*	1.6	4.3	9.9	*
2000	0.5	*	*	*	*	*	*	0.7	2.1	5.4	7.9	*
1999	0.5	*	*	*	*	*	*	*	1.5	4.7	9.0	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	2.8	5.4	*	*
2001	0.4	*	*	*	*	*	*	*	*	5.2	10.7	*
2000	0.4	*	*	*	*	*	*	*	*	5.9	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	2.3	4.7	6.7	*
2001	0.6	*	*	*	*	*	*	*	*	3.6	9.4	*
2000	0.6	*	*	*	*	*	*	*	2.3	5.0	7.3	*
1999	0.6	*	*	*	*	*	*	*	*	5.2	8.7	*
Non-Hispanic 4/, both sexes												
2002	1.1	*	*	*	*	0.1	0.2	0.7	2.0	3.9	7.1	10.3
2001	1.1	*	*	*	*	*	0.2	0.6	1.8	4.1	7.0	9.8
2000	1.0	*	*	*	*	*	0.2	0.7	1.8	3.8	6.7	9.4
1999	1.0	*	*	*	*	*	0.2	0.6	1.8	3.7	6.4	8.1
Non-Hispanic 4/, male												
2002	1.1	*	*	*	*	*	0.2	0.8	2.2	4.5	8.8	11.6
2001	1.1	*	*	*	*	*	0.2	0.7	2.0	4.7	8.6	12.1
2000	1.0	*	*	*	*	*	0.2	0.7	1.9	4.5	8.3	11.0
1999	1.0	*	*	*	*	*	0.2	0.6	2.0	4.4	7.4	8.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	*	0.1	0.5	1.8	3.3	5.9	9.8
2001	1.1	*	*	*	*	*	0.1	0.6	1.6	3.5	5.9	8.8
2000	1.0	*	*	*	*	*	0.1	0.6	1.6	3.2	5.6	8.8
1999	1.0	*	*	*	*	*	0.1	0.5	1.6	3.2	5.7	7.8
Non-Hispanic white, both sexes												
2002	1.2	*	*	*	*	*	0.2	0.6	1.9	3.8	7.0	10.1
2001	1.2	*	*	*	*	*	0.2	0.6	1.7	4.0	6.9	9.7
2000	1.1	*	*	*	*	*	0.1	0.7	1.7	3.8	6.7	9.6
1999	1.1	*	*	*	*	*	0.2	0.6	1.7	3.8	6.4	8.2
Non-Hispanic white, male												
2002	1.2	*	*	*	*	*	0.2	0.8	2.1	4.4	8.6	11.6
2001	1.2	*	*	*	*	*	0.2	0.7	1.8	4.6	8.6	12.1
2000	1.1	*	*	*	*	*	0.2	0.7	1.8	4.6	8.4	11.5
1999	1.0	*	*	*	*	*	0.2	0.6	2.0	4.5	7.2	9.1
Non-Hispanic white, female												
2002	1.2	*	*	*	*	*	0.1	0.5	1.7	3.4	5.9	9.5
2001	1.1	*	*	*	*	*	*	0.5	1.6	3.5	5.8	8.7
2000	1.1	*	*	*	*	*	0.1	0.6	1.6	3.2	5.6	8.8
1999	1.1	*	*	*	*	*	0.1	0.5	1.4	3.2	5.8	7.9
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	0.6	1.8	3.3	6.6	9.4
2001	0.6	*	*	*	*	*	*	0.7	2.0	3.4	5.9	8.1
2000	0.6	*	*	*	*	*	*	0.7	2.0	3.3	4.8	7.0
1999	0.5	*	*	*	*	*	*	*	1.6	3.3	5.2	*
Non-Hispanic black, male												
2002	0.7	*	*	*	*	*	*	*	1.8	5.0	9.1	*
2001	0.6	*	*	*	*	*	*	*	2.7	4.3	6.9	*
2000	0.6	*	*	*	*	*	*	*	2.1	4.0	*	*
1999	0.5	*	*	*	*	*	*	*	*	4.1	6.6	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	*	1.9	2.2	5.1	10.5
2001	0.7	*	*	*	*	*	*	*	*	2.7	5.4	*
2000	0.6	*	*	*	*	*	*	*	2.0	2.7	4.3	*
1999	0.6	*	*	*	*	*	*	*	1.9	2.8	4.4	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of gallbladder and extrahepatic bile ducts (C23-C24)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.6	1.7	4.6	8.9	14.5
2001	1.3	*	*	*	*	*	0.2	0.6	1.8	4.8	9.9	15.8
2000	1.3	*	*	*	*	*	0.1	0.6	2.0	5.3	10.0	16.0
1999	1.3	*	*	*	*	*	0.1	0.6	2.0	5.1	10.0	16.2
All origins 3/, male												
2002	0.9	*	*	*	*	*	0.1	0.4	1.4	4.3	8.5	15.0
2001	1.0	*	*	*	*	*	0.1	0.5	1.4	4.0	9.9	15.8
2000	1.0	*	*	*	*	*	0.1	0.4	1.6	5.0	9.5	15.8
1999	1.0	*	*	*	*	*	0.1	0.5	1.6	4.5	9.3	15.2
All origins 3/, female												
2002	1.4	*	*	*	*	*	0.2	0.8	1.9	4.9	9.2	14.3
2001	1.6	*	*	*	*	*	0.2	0.7	2.1	5.5	9.9	15.8
2000	1.6	*	*	*	*	*	0.1	0.8	2.3	5.5	10.4	16.0
1999	1.6	*	*	*	*	*	0.2	0.7	2.4	5.5	10.5	16.7
Hispanic, both sexes												
2002	0.7	*	*	*	*	*	*	1.0	2.0	8.0	12.5	13.3
2001	0.8	*	*	*	*	*	*	0.9	3.1	8.5	12.3	20.6
2000	0.8	*	*	*	*	*	*	0.9	3.4	6.5	16.4	18.6
1999	0.9	*	*	*	*	*	*	1.1	3.7	8.3	14.2	21.7
Hispanic, male												
2002	0.5	*	*	*	*	*	*	*	*	6.5	10.7	*
2001	0.6	*	*	*	*	*	*	*	2.3	7.0	13.4	*
2000	0.5	*	*	*	*	*	*	*	*	5.5	12.8	*
1999	0.4	*	*	*	*	*	*	*	*	*	10.9	*
Hispanic, female												
2002	1.0	*	*	*	*	*	*	1.4	2.8	9.1	13.8	*
2001	1.1	*	*	*	*	*	*	1.3	3.9	9.8	11.5	25.1
2000	1.1	*	*	*	*	*	*	1.4	4.9	7.3	18.8	*
1999	1.3	*	*	*	*	*	*	1.7	5.1	11.7	16.3	22.9
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.6	1.6	4.4	8.7	14.5
2001	1.3	*	*	*	*	*	0.1	0.6	1.7	4.6	9.8	15.6
2000	1.4	*	*	*	*	*	0.1	0.6	1.8	5.2	9.7	15.8
1999	1.3	*	*	*	*	*	0.1	0.5	1.9	4.9	9.8	16.0
Non-Hispanic 4/, male												
2002	1.0	*	*	*	*	*	0.1	0.4	1.4	4.2	8.4	14.7
2001	1.0	*	*	*	*	*	0.1	0.5	1.4	3.8	9.7	16.0
2000	1.1	*	*	*	*	*	0.1	0.5	1.6	4.9	9.3	15.6
1999	1.0	*	*	*	*	*	0.1	0.5	1.6	4.6	9.2	15.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of gallbladder and extrahepatic bile ducts (C23-C24)												
Non-Hispanic 4/, female												
2002	1.5	*	*	*	*	*	0.1	0.7	1.8	4.6	9.0	14.4
2001	1.6	*	*	*	*	*	0.2	0.7	2.0	5.2	9.8	15.4
2000	1.7	*	*	*	*	*	0.1	0.8	2.1	5.4	10.0	15.9
1999	1.6	*	*	*	*	*	0.1	0.6	2.2	5.1	10.2	16.4
Non-Hispanic white, both sexes												
2002	1.3	*	*	*	*	*	0.1	0.5	1.6	4.3	8.6	14.5
2001	1.4	*	*	*	*	*	0.1	0.6	1.6	4.4	9.8	15.5
2000	1.5	*	*	*	*	*	0.1	0.6	1.8	5.1	9.8	16.2
1999	1.5	*	*	*	*	*	0.1	0.5	1.9	4.8	9.6	16.1
Non-Hispanic white, male												
2002	1.1	*	*	*	*	*	*	0.3	1.4	4.1	8.3	15.0
2001	1.1	*	*	*	*	*	*	0.5	1.4	3.7	9.6	15.6
2000	1.2	*	*	*	*	*	0.1	0.5	1.6	4.9	9.5	16.1
1999	1.1	*	*	*	*	*	0.1	0.4	1.6	4.4	9.3	15.8
Non-Hispanic white, female												
2002	1.6	*	*	*	*	*	0.1	0.7	1.8	4.5	8.8	14.4
2001	1.7	*	*	*	*	*	0.1	0.7	1.9	4.9	9.9	15.5
2000	1.8	*	*	*	*	*	0.1	0.7	2.0	5.3	10.1	16.2
1999	1.8	*	*	*	*	*	*	0.6	2.1	5.1	9.9	16.3
Non-Hispanic black, both sexes												
2002	0.9	*	*	*	*	*	*	0.9	2.3	5.2	9.5	13.9
2001	0.9	*	*	*	*	*	*	0.7	2.1	5.8	8.8	15.9
2000	0.8	*	*	*	*	*	*	0.8	2.0	5.9	7.8	11.4
1999	0.9	*	*	*	*	*	*	0.7	2.2	5.3	10.1	12.0
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	*	1.9	4.5	7.9	*
2001	0.6	*	*	*	*	*	*	*	*	4.6	9.3	*
2000	0.5	*	*	*	*	*	*	*	*	4.8	6.8	*
1999	0.6	*	*	*	*	*	*	*	*	5.3	9.1	*
Non-Hispanic black, female												
2002	1.2	*	*	*	*	*	*	1.2	2.7	5.7	10.4	15.9
2001	1.2	*	*	*	*	*	*	1.0	2.6	6.6	8.6	16.3
2000	1.1	*	*	*	*	*	*	1.3	2.5	6.6	8.3	12.2
1999	1.1	*	*	*	*	*	*	*	2.8	5.3	10.7	15.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pancreas (C25)												
All origins 3/, both sexes												
2002	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.6	74.1	100.2
2000	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
All origins 3/, male												
2002	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
All origins 3/, female												
2002	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6
Hispanic, both sexes												
2002	3.5	*	*	*	*	*	0.9	4.0	13.8	34.7	55.5	79.3
2001	3.6	*	*	*	*	*	0.8	3.7	14.2	37.0	61.5	84.4
2000	3.5	*	*	*	*	*	0.7	4.1	13.7	36.9	61.8	82.9
1999	3.3	*	*	*	*	*	0.9	4.6	13.3	32.6	56.7	84.2
Hispanic, male												
2002	3.3	*	*	*	*	*	1.2	5.3	15.7	40.9	55.8	60.2
2001	3.4	*	*	*	*	*	0.9	4.3	16.2	42.5	68.9	64.8
2000	3.4	*	*	*	*	*	0.8	5.2	16.6	45.2	62.9	80.6
1999	3.2	*	*	*	*	*	1.2	5.7	15.2	40.0	56.3	86.3
Hispanic, female												
2002	3.7	*	*	*	*	*	*	2.8	12.1	29.8	55.4	88.7
2001	3.9	*	*	*	*	*	0.8	3.2	12.5	32.6	56.5	94.0
2000	3.6	*	*	*	*	*	*	3.0	11.1	30.4	61.1	84.1
1999	3.5	*	*	*	*	*	*	3.5	11.6	26.8	56.9	83.2
Non-Hispanic 4/, both sexes												
2002	11.6	*	*	*	*	0.1	1.4	6.3	19.6	45.4	75.8	94.9
2001	11.5	*	*	*	*	0.1	1.3	6.4	19.6	44.9	74.6	100.6
2000	11.4	*	*	*	*	0.1	1.3	6.5	19.8	44.4	77.0	96.6
1999	11.4	*	*	*	*	0.1	1.4	6.6	20.2	45.6	73.5	100.1
Non-Hispanic 4/, male												
2002	11.7	*	*	*	*	0.1	1.7	8.0	24.1	52.1	85.3	102.4
2001	11.4	*	*	*	*	0.2	1.5	7.9	24.2	51.2	84.6	106.4
2000	11.3	*	*	*	*	0.2	1.5	8.3	24.1	50.7	86.7	103.5
1999	11.4	*	*	*	*	0.2	1.7	8.4	24.4	52.9	82.7	108.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of pancreas (C25)												
Non-Hispanic 4/, female												
2002	11.5	*	*	*	*	*	1.1	4.6	15.4	39.8	69.5	91.7
2001	11.5	*	*	*	*	0.1	1.2	4.9	15.4	39.8	68.0	98.2
2000	11.4	*	*	*	*	*	1.0	4.7	15.8	39.2	70.6	93.8
1999	11.4	*	*	*	*	*	1.0	4.8	16.4	39.6	67.6	96.8
Non-Hispanic white, both sexes												
2002	12.4	*	*	*	*	0.1	1.4	6.0	19.0	44.3	75.1	94.6
2001	12.3	*	*	*	*	0.1	1.3	6.1	19.3	44.1	73.7	99.4
2000	12.2	*	*	*	*	0.2	1.2	6.3	19.2	43.4	75.7	94.7
1999	12.1	*	*	*	*	0.1	1.3	6.3	19.5	44.7	72.1	98.9
Non-Hispanic white, male												
2002	12.6	*	*	*	*	*	1.7	7.7	23.5	51.2	85.4	102.9
2001	12.4	*	*	*	*	0.2	1.5	7.7	24.0	50.3	84.5	108.0
2000	12.2	*	*	*	*	0.2	1.5	8.0	23.4	50.0	86.1	103.0
1999	12.1	*	*	*	*	0.2	1.6	8.0	23.4	52.1	81.6	108.1
Non-Hispanic white, female												
2002	12.2	*	*	*	*	*	1.1	4.3	14.7	38.3	68.1	91.0
2001	12.3	*	*	*	*	*	1.2	4.6	14.9	38.8	66.6	95.8
2000	12.2	*	*	*	*	*	1.0	4.5	15.3	37.9	68.9	91.4
1999	12.1	*	*	*	*	*	1.0	4.7	15.7	38.5	65.9	95.2
Non-Hispanic black, both sexes												
2002	9.6	*	*	*	*	*	1.8	8.9	28.3	61.7	90.9	108.3
2001	9.4	*	*	*	*	*	1.6	9.2	25.9	59.0	93.3	122.0
2000	9.7	*	*	*	*	*	1.7	9.2	27.9	60.0	98.9	121.2
1999	9.7	*	*	*	*	*	2.0	9.9	28.8	60.4	95.0	119.0
Non-Hispanic black, male												
2002	9.3	*	*	*	*	*	2.2	11.6	34.9	69.5	97.3	102.8
2001	8.9	*	*	*	*	*	1.8	11.4	30.8	68.2	100.5	100.5
2000	9.3	*	*	*	*	*	2.1	12.1	34.6	66.9	108.0	110.4
1999	9.7	*	*	*	*	*	2.4	13.6	37.4	68.7	107.6	115.3
Non-Hispanic black, female												
2002	9.8	*	*	*	*	*	1.5	6.5	23.0	56.2	87.2	110.4
2001	9.9	*	*	*	*	*	1.4	7.3	21.9	52.4	89.3	130.0
2000	10.0	*	*	*	*	*	1.5	6.8	22.5	55.1	93.8	125.2
1999	9.7	*	*	*	*	*	1.7	6.7	21.9	54.6	87.9	120.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and ill-defined digestive organs (C26)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.6	1.7	3.7	6.9
2001	0.5	*	*	*	*	*	0.1	0.2	0.7	1.7	3.6	7.8
2000	0.5	*	*	*	*	*	0.1	0.2	0.7	1.7	3.5	7.8
1999	0.6	*	*	*	*	*	0.1	0.3	0.8	1.9	4.1	9.1
All origins 3/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.7	2.3	4.7	6.4
2001	0.5	*	*	*	*	*	*	0.3	1.0	2.0	4.2	8.9
2000	0.5	*	*	*	*	*	*	0.2	1.0	2.3	4.1	7.5
1999	0.6	*	*	*	*	*	0.1	0.4	1.0	2.2	5.2	10.4
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.1	0.5	1.1	3.0	7.2
2001	0.5	*	*	*	*	*	*	0.1	0.5	1.5	3.2	7.3
2000	0.5	*	*	*	*	*	*	0.2	0.6	1.2	3.1	8.0
1999	0.6	*	*	*	*	*	*	0.2	0.6	1.7	3.4	8.5
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	4.8	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.6	1.7	3.7	7.0
2001	0.5	*	*	*	*	*	0.1	0.2	0.7	1.8	3.7	7.8
2000	0.5	*	*	*	*	*	0.1	0.2	0.8	1.7	3.5	7.9
1999	0.6	*	*	*	*	*	0.1	0.3	0.8	2.0	4.0	9.1
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.7	2.3	4.8	6.5
2001	0.5	*	*	*	*	*	*	0.3	0.9	2.1	4.2	8.7
2000	0.5	*	*	*	*	*	*	0.2	1.0	2.3	4.1	7.5
1999	0.6	*	*	*	*	*	0.1	0.4	1.0	2.3	5.1	10.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and ill-defined digestive organs (C26)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.1	0.5	1.1	3.0	7.3
2001	0.6	*	*	*	*	*	*	0.1	0.5	1.5	3.3	7.4
2000	0.5	*	*	*	*	*	*	0.2	0.6	1.2	3.1	8.0
1999	0.6	*	*	*	*	*	*	0.2	0.5	1.7	3.3	8.7
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	*	0.2	0.6	1.6	3.8	7.1
2001	0.6	*	*	*	*	*	*	0.2	0.6	1.7	3.6	7.7
2000	0.6	*	*	*	*	*	*	0.2	0.8	1.6	3.4	7.9
1999	0.7	*	*	*	*	*	0.1	0.2	0.7	2.0	4.1	9.1
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	*	0.2	0.7	2.2	4.8	6.3
2001	0.6	*	*	*	*	*	*	0.3	0.9	2.1	4.2	8.5
2000	0.6	*	*	*	*	*	*	0.3	1.0	2.1	4.2	7.6
1999	0.7	*	*	*	*	*	0.1	0.3	0.9	2.2	5.2	10.0
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	0.5	1.0	3.0	7.4
2001	0.6	*	*	*	*	*	*	*	0.4	1.5	3.2	7.4
2000	0.6	*	*	*	*	*	*	0.2	0.5	1.1	3.0	8.0
1999	0.7	*	*	*	*	*	*	0.1	0.5	1.8	3.4	8.7
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.8	2.8	3.9	7.3
2001	0.5	*	*	*	*	*	*	*	1.3	2.3	4.7	10.0
2000	0.5	*	*	*	*	*	*	*	1.0	2.6	5.0	8.3
1999	0.5	*	*	*	*	*	*	0.8	1.2	2.1	3.3	10.3
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	4.0	5.9	*
2001	0.5	*	*	*	*	*	*	*	*	3.3	*	*
2000	0.4	*	*	*	*	*	*	*	*	3.9	*	*
1999	0.5	*	*	*	*	*	*	1.3	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	4.9	8.6
2000	0.5	*	*	*	*	*	*	*	*	*	5.4	8.7
1999	0.4	*	*	*	*	*	*	*	*	*	*	9.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
All origins 3/, both sexes												
2002	56.3	*	*	*	0.1	0.5	6.3	31.7	119.3	282.1	386.9	305.1
2001	56.5	*	*	*	0.1	0.5	6.6	32.2	121.9	287.3	380.6	312.2
2000	57.0	*	*	*	0.1	0.6	6.4	33.0	126.7	291.9	380.7	310.8
1999	56.2	*	*	*	0.1	0.5	6.4	32.7	127.9	289.4	371.2	306.6
All origins 3/, male												
2002	66.1	*	*	*	0.2	0.6	6.6	38.8	150.7	367.5	544.9	498.8
2001	67.2	*	*	*	0.1	0.4	7.2	39.7	154.1	382.6	546.2	534.1
2000	68.1	*	*	*	0.1	0.6	7.3	40.7	161.1	390.8	550.4	541.7
1999	68.0	*	*	*	*	0.6	7.1	40.6	165.3	390.0	544.0	550.6
All origins 3/, female												
2002	46.8	*	*	*	*	0.4	6.0	24.9	90.3	211.1	282.0	221.1
2001	46.1	*	*	*	*	0.5	5.9	25.0	92.3	208.4	271.8	219.3
2000	46.2	*	*	*	*	0.5	5.5	25.6	94.9	210.4	270.0	216.8
1999	45.0	*	*	*	*	0.5	5.7	25.1	93.4	207.0	258.9	208.3
Hispanic, both sexes												
2002	11.0	*	*	*	*	0.4	1.8	10.8	43.3	115.7	191.0	199.0
2001	11.0	*	*	*	*	*	2.1	11.6	43.7	115.0	190.3	210.7
2000	10.9	*	*	*	*	0.3	1.9	11.6	41.7	116.0	206.6	222.3
1999	11.1	*	*	*	*	0.4	2.2	11.5	46.5	118.8	197.7	240.6
Hispanic, male												
2002	14.0	*	*	*	*	*	1.9	14.0	65.3	170.1	312.7	321.3
2001	14.1	*	*	*	*	*	2.7	15.7	64.1	176.1	308.1	356.2
2000	14.1	*	*	*	*	*	2.3	16.1	63.1	176.4	343.2	391.0
1999	14.6	*	*	*	*	*	2.6	16.2	72.2	186.5	333.2	396.8
Hispanic, female												
2002	7.9	*	*	*	*	*	1.8	7.6	23.7	72.4	107.6	138.4
2001	7.6	*	*	*	*	*	1.6	7.6	25.6	66.8	110.6	139.6
2000	7.4	*	*	*	*	*	1.4	7.3	22.7	68.5	114.9	139.5
1999	7.4	*	*	*	*	*	1.8	7.0	23.7	65.7	107.5	165.3
Non-Hispanic 4/, both sexes												
2002	63.2	*	*	*	0.1	0.5	7.0	33.7	125.1	292.9	395.6	308.8
2001	63.1	*	*	*	0.1	0.5	7.1	34.0	127.7	298.1	388.6	315.2
2000	63.4	*	*	*	0.1	0.6	7.0	34.8	132.7	302.0	387.4	313.2
1999	62.3	*	*	*	0.1	0.5	6.9	34.4	133.5	298.7	377.3	308.2
Non-Hispanic 4/, male												
2002	74.5	*	*	*	0.2	0.6	7.4	41.2	157.0	379.8	555.2	505.7
2001	75.4	*	*	*	*	0.5	7.8	41.8	160.3	395.1	556.1	540.1
2000	76.1	*	*	*	0.1	0.7	8.0	42.8	167.8	402.7	558.2	546.6
1999	75.5	*	*	*	*	0.6	7.6	42.6	171.6	400.6	551.2	555.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)												
Non-Hispanic 4/, female												
2002	52.4	*	*	*	*	0.4	6.6	26.5	95.5	220.4	289.8	223.9
2001	51.5	*	*	*	*	0.5	6.5	26.5	97.5	217.4	278.7	221.7
2000	51.3	*	*	*	*	0.5	6.0	27.2	100.3	218.9	276.1	218.8
1999	49.8	*	*	*	*	0.5	6.1	26.6	98.4	215.1	264.4	209.2
Non-Hispanic white, both sexes												
2002	69.5	*	*	*	0.1	0.5	6.8	32.4	124.3	298.9	401.4	312.8
2001	69.2	*	*	*	*	0.5	7.0	32.7	128.0	302.4	391.9	317.9
2000	69.3	*	*	*	0.1	0.6	6.5	33.2	132.9	305.4	391.3	315.7
1999	67.8	*	*	*	*	0.6	6.5	32.8	133.0	301.2	379.3	309.6
Non-Hispanic white, male												
2002	80.6	*	*	*	*	0.6	6.9	38.4	151.7	380.6	554.8	510.5
2001	81.2	*	*	*	*	0.5	7.4	38.9	156.7	392.7	552.1	541.6
2000	81.8	*	*	*	*	0.7	7.1	39.5	163.8	399.3	555.4	548.0
1999	80.9	*	*	*	*	0.6	7.1	39.2	166.8	396.5	547.1	556.6
Non-Hispanic white, female												
2002	58.8	*	*	*	*	0.4	6.7	26.5	98.4	229.3	298.6	227.5
2001	57.6	*	*	*	*	0.5	6.5	26.7	100.9	225.7	285.7	224.8
2000	57.3	*	*	*	*	0.5	5.9	27.1	103.7	226.2	283.5	222.1
1999	55.2	*	*	*	*	0.5	6.0	26.5	101.2	221.5	269.5	211.0
Non-Hispanic black, both sexes												
2002	46.9	*	*	*	*	0.6	9.7	51.3	164.0	311.6	400.6	280.7
2001	46.8	*	*	*	*	*	9.9	50.9	158.3	324.7	404.8	291.2
2000	47.8	*	*	*	*	0.7	11.1	54.9	165.3	334.4	395.7	298.9
1999	47.8	*	*	*	*	0.5	10.5	55.0	170.9	330.9	402.3	303.3
Non-Hispanic black, male												
2002	61.6	*	*	*	*	0.8	12.0	71.6	244.1	473.3	665.6	504.3
2001	62.7	*	*	*	*	*	12.1	73.2	234.1	509.3	698.0	574.0
2000	64.2	*	*	*	*	*	14.8	78.2	246.3	521.9	680.3	589.4
1999	64.5	*	*	*	*	*	13.0	79.8	259.6	514.6	689.2	606.1
Non-Hispanic black, female												
2002	33.5	*	*	*	*	*	7.7	33.9	99.7	196.8	250.7	195.2
2001	32.3	*	*	*	*	*	7.9	31.7	97.4	194.5	240.2	186.5
2000	32.9	*	*	*	*	*	7.9	35.0	100.2	202.4	235.5	191.0
1999	32.7	*	*	*	*	*	8.2	33.7	99.8	201.2	241.6	190.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of nasal cavity, middle ear and accessory sinuses (C30-C31)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.2
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.1	1.7
2000	0.1	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.2
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.5
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.3	0.4	0.8	1.0	1.7
2001	0.2	*	*	*	*	*	*	0.2	0.5	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	0.1	0.2	0.5	0.8	1.0	*
1999	0.2	*	*	*	*	*	0.1	0.3	0.4	0.7	1.2	1.8
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	0.2	0.4	0.7	1.0
2001	0.2	*	*	*	*	*	*	*	0.2	0.4	0.9	1.6
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	1.1
1999	0.1	*	*	*	*	*	*	*	0.2	0.5	0.7	1.4
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.2
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.1	1.6
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.2
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.5
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.5	0.9	1.0	1.7
2001	0.2	*	*	*	*	*	*	0.2	0.5	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	0.1	0.2	0.5	0.8	1.1	*
1999	0.2	*	*	*	*	*	*	0.3	0.5	0.7	1.2	1.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of nasal cavity, middle ear and accessory sinuses (C30-C31)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	1.0
2001	0.2	*	*	*	*	*	*	*	0.2	0.4	0.9	1.6
2000	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	1.1
1999	0.2	*	*	*	*	*	*	*	*	0.5	0.7	1.4
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.6	0.9	1.3
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.1	1.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.3
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.5
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.3	0.5	0.8	1.1	1.8
2001	0.2	*	*	*	*	*	*	0.2	0.5	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.8	1.2	*
1999	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.2	1.9
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.0
2001	0.2	*	*	*	*	*	*	*	0.2	0.4	0.9	1.7
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.6	1.2
1999	0.2	*	*	*	*	*	*	*	*	0.5	0.8	1.4
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of larynx (C32)												
All origins 3/, both sexes												
2002	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
All origins 3/, male												
2002	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	*	2.2	4.1	5.5	*
2001	0.4	*	*	*	*	*	*	0.6	2.1	5.3	5.2	*
2000	0.4	*	*	*	*	*	*	*	1.7	4.2	6.9	*
1999	0.5	*	*	*	*	*	*	0.8	1.8	5.3	8.3	*
Hispanic, male												
2002	0.7	*	*	*	*	*	*	*	4.3	8.0	11.1	*
2001	0.7	*	*	*	*	*	*	*	3.9	10.6	10.7	*
2000	0.7	*	*	*	*	*	*	*	3.5	7.8	14.8	*
1999	0.9	*	*	*	*	*	*	1.6	3.5	10.5	16.2	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.4	*	*	*	*	*	0.2	1.1	3.5	6.0	7.7	5.4
2001	1.5	*	*	*	*	*	0.2	1.1	3.7	6.4	7.3	6.7
2000	1.5	*	*	*	*	*	0.1	1.2	3.7	6.7	8.1	6.5
1999	1.5	*	*	*	*	*	0.2	1.1	3.9	6.8	7.2	6.8
Non-Hispanic 4/, male												
2002	2.3	*	*	*	*	*	0.3	1.9	5.9	10.1	15.0	13.2
2001	2.4	*	*	*	*	*	0.3	1.9	6.1	11.1	14.3	18.4
2000	2.4	*	*	*	*	*	0.2	1.9	6.4	11.5	15.7	17.3
1999	2.4	*	*	*	*	*	0.3	1.9	6.6	11.4	14.0	17.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of larynx (C32)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	*	*	*	0.4	1.3	2.6	2.9	2.1
2001	0.6	*	*	*	*	*	0.1	0.4	1.4	2.6	2.7	1.8
2000	0.6	*	*	*	*	*	*	0.5	1.2	2.8	3.1	2.1
1999	0.6	*	*	*	*	*	*	0.3	1.4	3.0	2.8	2.6
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	*	0.1	0.9	3.1	5.8	7.3	5.5
2001	1.5	*	*	*	*	*	0.2	0.9	3.3	6.0	6.9	6.5
2000	1.5	*	*	*	*	*	0.1	1.0	3.2	6.2	7.7	6.5
1999	1.5	*	*	*	*	*	0.1	0.9	3.5	6.4	7.1	6.5
Non-Hispanic white, male												
2002	2.3	*	*	*	*	*	0.2	1.5	5.1	9.5	14.2	13.3
2001	2.3	*	*	*	*	*	0.3	1.5	5.5	10.2	13.3	18.3
2000	2.4	*	*	*	*	*	0.2	1.5	5.4	10.5	15.0	17.4
1999	2.3	*	*	*	*	*	0.2	1.6	5.8	10.7	13.6	16.0
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	0.3	1.3	2.6	2.7	2.2
2001	0.6	*	*	*	*	*	*	0.3	1.2	2.4	2.7	1.6
2000	0.6	*	*	*	*	*	*	0.4	1.1	2.6	3.0	2.1
1999	0.7	*	*	*	*	*	*	0.3	1.3	2.9	2.9	2.7
Non-Hispanic black, both sexes												
2002	1.9	*	*	*	*	*	0.5	2.7	7.7	10.6	13.9	*
2001	1.9	*	*	*	*	*	0.4	2.9	7.9	11.5	12.0	8.4
2000	2.0	*	*	*	*	*	*	2.9	9.2	12.9	14.3	7.0
1999	1.9	*	*	*	*	*	0.6	3.0	8.7	11.7	10.0	8.4
Non-Hispanic black, male												
2002	3.1	*	*	*	*	*	0.8	4.6	15.1	19.8	29.6	*
2001	3.1	*	*	*	*	*	*	5.2	14.2	21.4	28.2	*
2000	3.4	*	*	*	*	*	*	5.2	17.5	24.6	29.0	*
1999	3.2	*	*	*	*	*	1.0	5.3	16.3	22.1	22.5	25.0
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	1.1	1.8	4.1	5.0	*
2001	0.7	*	*	*	*	*	*	*	2.9	4.6	*	*
2000	0.8	*	*	*	*	*	*	1.0	2.4	4.7	6.1	*
1999	0.7	*	*	*	*	*	*	0.9	2.7	4.3	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)												
All origins 3/, both sexes												
2002	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
All origins 3/, male												
2002	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.6	512.4
2000	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
All origins 3/, female												
2002	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001	45.3	*	*	*	*	0.4	5.7	24.3	90.5	205.0	267.5	215.0
2000	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5
Hispanic, both sexes												
2002	10.4	*	*	*	*	0.4	1.5	9.9	40.9	110.4	183.7	189.0
2001	10.3	*	*	*	*	*	1.9	10.7	41.1	108.6	182.5	200.7
2000	10.3	*	*	*	*	*	1.8	10.9	39.3	110.6	198.7	213.7
1999	10.4	*	*	*	*	*	2.0	10.4	44.2	112.9	187.7	227.3
Hispanic, male												
2002	13.1	*	*	*	*	*	1.5	12.6	60.7	161.7	299.1	307.9
2001	13.1	*	*	*	*	*	2.4	14.2	59.6	163.9	294.2	335.2
2000	13.3	*	*	*	*	*	2.1	14.8	58.6	167.3	327.5	368.8
1999	13.6	*	*	*	*	*	2.3	14.2	68.1	175.3	316.0	362.3
Hispanic, female												
2002	7.6	*	*	*	*	*	1.6	7.3	23.2	69.5	104.6	130.1
2001	7.4	*	*	*	*	*	1.4	7.3	24.6	64.9	107.0	135.0
2000	7.2	*	*	*	*	*	1.4	7.1	22.2	66.0	112.3	137.5
1999	7.1	*	*	*	*	*	1.7	6.7	23.1	64.0	102.3	162.2
Non-Hispanic 4/, both sexes												
2002	61.4	*	*	*	*	0.4	6.7	32.3	121.0	285.7	386.2	301.0
2001	61.3	*	*	*	*	0.4	6.8	32.5	123.3	290.4	379.4	305.8
2000	61.5	*	*	*	*	0.5	6.6	33.4	128.4	294.2	377.4	304.5
1999	60.5	*	*	*	*	0.5	6.5	33.0	128.9	290.7	368.2	298.8
Non-Hispanic 4/, male												
2002	71.8	*	*	*	*	0.5	6.9	38.9	150.3	368.1	538.0	488.9
2001	72.6	*	*	*	*	0.4	7.3	39.5	153.2	382.4	539.5	518.4
2000	73.2	*	*	*	*	0.6	7.5	40.5	160.6	389.7	539.9	526.2
1999	72.7	*	*	*	*	0.5	7.1	40.1	164.0	387.8	534.9	534.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)												
Non-Hispanic 4/, female												
2002	51.6	*	*	*	*	0.4	6.4	25.9	93.9	216.9	285.5	220.0
2001	50.5	*	*	*	*	0.5	6.2	25.8	95.6	213.9	274.3	217.4
2000	50.4	*	*	*	*	0.5	5.8	26.5	98.6	215.4	271.6	214.8
1999	48.8	*	*	*	*	0.4	6.0	26.1	96.5	211.0	260.0	204.4
Non-Hispanic white, both sexes												
2002	67.7	*	*	*	*	0.4	6.5	31.2	120.6	292.0	392.2	304.8
2001	67.3	*	*	*	*	0.5	6.6	31.5	124.0	295.2	383.1	308.6
2000	67.4	*	*	*	*	0.5	6.2	32.0	129.1	298.0	381.7	306.8
1999	65.9	*	*	*	*	0.5	6.2	31.5	128.9	293.5	370.3	300.5
Non-Hispanic white, male												
2002	77.8	*	*	*	*	0.5	6.5	36.5	145.7	369.5	538.3	493.3
2001	78.4	*	*	*	*	0.4	6.9	37.0	150.3	381.1	536.5	520.0
2000	78.9	*	*	*	*	0.6	6.7	37.7	157.7	387.3	537.7	527.3
1999	78.1	*	*	*	*	0.5	6.6	37.1	160.1	384.4	531.1	536.9
Non-Hispanic white, female												
2002	57.9	*	*	*	*	0.3	6.5	26.0	96.8	225.9	294.3	223.4
2001	56.6	*	*	*	*	0.5	6.4	26.0	99.2	222.3	281.3	220.6
2000	56.4	*	*	*	*	0.5	5.7	26.4	102.2	222.9	279.2	218.0
1999	54.3	*	*	*	*	0.5	5.8	26.1	99.5	217.5	265.0	206.2
Non-Hispanic black, both sexes												
2002	44.7	*	*	*	*	0.4	9.0	48.1	155.6	299.5	385.6	274.9
2001	44.6	*	*	*	*	*	9.2	47.5	149.4	311.4	390.9	281.6
2000	45.4	*	*	*	*	0.6	10.4	51.5	155.0	320.2	379.4	291.0
1999	45.6	*	*	*	*	0.4	9.6	51.5	161.1	317.7	390.9	293.3
Non-Hispanic black, male												
2002	58.1	*	*	*	*	*	11.0	66.3	228.0	451.4	633.9	490.1
2001	59.2	*	*	*	*	*	11.4	67.5	218.3	484.4	668.0	550.9
2000	60.3	*	*	*	*	*	13.8	72.4	227.1	495.4	649.5	567.1
1999	60.9	*	*	*	*	*	11.7	73.7	241.9	491.0	665.1	580.0
Non-Hispanic black, female												
2002	32.6	*	*	*	*	*	7.3	32.5	97.5	191.8	245.1	192.7
2001	31.3	*	*	*	*	*	7.3	30.4	94.0	189.4	235.3	181.8
2000	31.9	*	*	*	*	*	7.4	33.6	97.0	196.9	227.3	188.4
1999	31.8	*	*	*	*	*	7.9	32.5	96.4	195.5	237.4	186.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of thymus, heart, mediastinum and pleura (C37-C38)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.6	0.8	1.1
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.0
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.0	0.9
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9	1.0
All origins 3/, male												
2002	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.7
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.8	0.9	1.5
2000	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.4	*
1999	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.0	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	0.1	*	0.5	0.7	0.9
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.7	0.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.7	0.7
1999	0.1	*	*	*	*	*	*	*	0.3	0.5	0.8	0.8
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.2	0.2	0.6	0.8	1.1
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.0
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.0	0.9
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	0.9	1.0
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.0	1.7
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.8	0.9	*
2000	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.4	*
1999	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.0	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of thymus, heart, mediastinum and pleura (C37-C38)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	0.1	*	0.5	0.7	0.8
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.5	0.7	0.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.7	*
1999	0.2	*	*	*	*	*	*	*	0.3	0.5	0.8	0.8
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.8	1.2
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.0
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	0.9
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9	1.0
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.0	1.9
2001	0.2	*	*	*	*	*	*	0.2	0.4	0.7	0.9	*
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4	*
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.1	*
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	0.1	*	0.5	0.7	0.9
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.5	0.6	0.9
2000	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	0.8	0.8
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and ill-defined sites in respiratory system and intrathoracic organs (C39)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of bone and articular cartilage (C40-C41)												
All origins 3/, both sexes												
2002	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.6	0.8	1.6	2.9
2001	0.5	*	*	0.2	0.5	0.2	0.2	0.3	0.5	1.1	1.9	3.5
2000	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.6	1.0	1.9	3.4
1999	0.4	*	*	0.2	0.5	0.2	0.2	0.4	0.4	1.0	1.8	3.0
All origins 3/, male												
2002	0.5	*	*	0.2	0.5	0.3	0.3	0.4	0.7	0.9	2.0	3.7
2001	0.5	*	*	0.2	0.7	0.2	0.3	0.4	0.6	1.4	2.5	3.6
2000	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.7	1.2	2.4	4.2
1999	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.6	1.2	2.0	4.5
All origins 3/, female												
2002	0.4	*	*	0.2	0.3	0.2	0.1	0.2	0.5	0.8	1.3	2.5
2001	0.4	*	*	0.2	0.3	0.2	0.2	0.2	0.5	0.8	1.5	3.4
2000	0.4	*	*	0.2	0.3	0.1	0.1	0.2	0.4	0.8	1.6	3.1
1999	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.3	0.8	1.7	2.4
Hispanic, both sexes												
2002	0.3	*	*	0.3	0.5	*	*	*	*	*	*	*
2001	0.3	*	*	*	0.4	*	*	*	*	*	*	*
2000	0.3	*	*	*	0.5	*	*	*	*	*	*	*
1999	0.3	*	*	*	0.5	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	0.7	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	0.2	0.4	0.3	0.2	0.3	0.6	0.8	1.6	2.9
2001	0.5	*	*	0.2	0.5	0.3	0.2	0.3	0.5	1.1	1.9	3.5
2000	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.6	0.9	1.9	3.4
1999	0.5	*	*	0.2	0.5	0.2	0.2	0.4	0.4	1.0	1.8	2.9
Non-Hispanic 4/, male												
2002	0.5	*	*	0.2	0.5	0.3	0.3	0.4	0.6	0.9	2.1	3.6
2001	0.5	*	*	0.2	0.7	0.3	0.3	0.4	0.6	1.4	2.4	3.7
2000	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.7	1.2	2.4	4.2
1999	0.5	*	*	0.2	0.6	0.2	0.2	0.4	0.6	1.3	1.9	4.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of bone and articular cartilage (C40-C41)												
Non-Hispanic 4/, female												
2002	0.4	*	*	0.2	0.3	0.2	0.1	0.2	0.5	0.8	1.3	2.6
2001	0.4	*	*	0.2	0.3	0.2	0.2	0.2	0.5	0.8	1.5	3.4
2000	0.4	*	*	0.2	0.3	*	0.1	0.2	0.4	0.7	1.6	3.0
1999	0.4	*	*	0.2	0.4	0.2	0.2	0.3	0.3	0.8	1.7	2.4
Non-Hispanic white, both sexes												
2002	0.5	*	*	0.2	0.5	0.3	0.2	0.3	0.6	0.8	1.7	3.0
2001	0.5	*	*	0.2	0.6	0.3	0.2	0.3	0.5	1.0	2.0	3.6
2000	0.5	*	*	0.2	0.4	0.2	0.2	0.3	0.5	1.0	1.9	3.5
1999	0.5	*	*	0.2	0.5	0.2	0.2	0.4	0.4	1.0	1.9	3.1
Non-Hispanic white, male												
2002	0.5	*	*	0.2	0.5	0.4	0.3	0.4	0.6	0.9	2.1	3.8
2001	0.6	*	*	0.2	0.8	0.3	0.3	0.4	0.6	1.4	2.5	3.6
2000	0.6	*	*	0.2	0.5	0.4	0.2	0.4	0.7	1.2	2.4	4.5
1999	0.5	*	*	0.2	0.6	0.3	0.2	0.4	0.5	1.3	2.0	4.6
Non-Hispanic white, female												
2002	0.4	*	*	0.2	0.4	0.2	*	0.1	0.5	0.8	1.4	2.7
2001	0.4	*	*	0.2	0.3	0.3	0.1	0.2	0.4	0.7	1.6	3.5
2000	0.4	*	*	0.2	0.3	*	0.1	0.2	0.4	0.8	1.6	3.1
1999	0.4	*	*	*	0.5	0.2	0.2	0.3	0.3	0.8	1.8	2.4
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.8	1.2	*	*
2001	0.4	*	*	*	0.4	*	*	*	0.9	1.6	*	*
2000	0.4	*	*	*	0.5	*	*	*	*	*	*	*
1999	0.3	*	*	*	0.4	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	0.8	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)												
All origins 3/, both sexes												
2002	3.4	*	*	*	0.1	0.5	1.5	3.3	6.4	11.5	19.5	29.8
2001	3.5	*	*	*	0.1	0.5	1.5	3.6	6.8	12.1	18.7	31.2
2000	3.4	*	*	*	0.1	0.5	1.5	3.7	6.7	11.9	18.0	31.8
1999	3.4	*	*	*	0.1	0.5	1.5	3.6	6.7	11.8	18.8	29.2
All origins 3/, male												
2002	4.5	*	*	*	0.1	0.6	1.8	4.6	9.3	17.4	31.1	50.2
2001	4.6	*	*	*	0.1	0.6	1.8	4.8	9.9	18.7	30.0	54.0
2000	4.5	*	*	*	0.1	0.6	1.7	4.9	9.5	17.8	29.6	55.4
1999	4.5	*	*	*	0.1	0.6	2.0	4.9	9.8	17.9	30.0	50.6
All origins 3/, female												
2002	2.4	*	*	*	0.1	0.4	1.1	2.1	3.7	6.6	11.8	21.0
2001	2.5	*	*	*	*	0.4	1.1	2.4	4.0	6.7	11.3	21.7
2000	2.5	*	*	*	0.1	0.4	1.2	2.5	4.1	7.0	10.5	22.2
1999	2.4	*	*	*	*	0.4	1.1	2.4	3.9	6.7	11.5	20.6
Hispanic, both sexes												
2002	0.6	*	*	*	*	*	0.4	0.9	2.0	2.8	8.3	14.4
2001	0.6	*	*	*	*	*	0.5	0.9	2.1	3.6	6.9	13.8
2000	0.6	*	*	*	*	*	0.6	1.1	2.5	4.3	6.5	*
1999	0.6	*	*	*	*	*	*	1.0	2.7	4.1	6.2	15.4
Hispanic, male												
2002	0.6	*	*	*	*	*	*	1.1	3.2	4.4	10.3	*
2001	0.7	*	*	*	*	*	0.7	*	2.9	5.6	8.9	*
2000	0.8	*	*	*	*	*	0.8	1.5	4.1	5.9	11.8	*
1999	0.7	*	*	*	*	*	*	1.6	3.9	5.7	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	7.0	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	3.9	*	*	*	0.1	0.6	1.6	3.6	6.7	12.1	20.0	30.4
2001	3.9	*	*	*	0.1	0.6	1.6	3.9	7.2	12.6	19.2	31.8
2000	3.8	*	*	*	0.1	0.6	1.6	3.9	7.0	12.3	18.5	32.5
1999	3.8	*	*	*	0.1	0.6	1.7	3.8	7.0	12.2	19.2	29.6
Non-Hispanic 4/, male												
2002	5.1	*	*	*	*	0.7	2.0	5.0	9.8	18.3	32.0	51.8
2001	5.2	*	*	*	0.1	0.7	2.0	5.2	10.4	19.5	30.9	55.4
2000	5.0	*	*	*	0.1	0.7	1.9	5.2	9.9	18.5	30.3	57.0
1999	5.0	*	*	*	0.1	0.7	2.2	5.2	10.2	18.5	30.7	51.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)												
Non-Hispanic 4/, female												
2002	2.7	*	*	*	*	0.5	1.2	2.2	3.9	7.0	12.0	21.2
2001	2.7	*	*	*	*	0.5	1.2	2.6	4.2	7.0	11.5	22.0
2000	2.7	*	*	*	*	0.5	1.3	2.6	4.3	7.2	10.8	22.6
1999	2.6	*	*	*	*	0.5	1.1	2.5	4.0	7.0	11.8	20.7
Non-Hispanic white, both sexes												
2002	4.7	*	*	*	0.1	0.8	2.0	4.1	7.6	13.4	21.9	32.9
2001	4.8	*	*	*	0.2	0.7	2.0	4.5	8.1	14.0	20.9	34.4
2000	4.6	*	*	*	0.2	0.7	1.9	4.5	7.8	13.6	20.0	35.0
1999	4.6	*	*	*	0.1	0.7	2.0	4.4	7.9	13.5	20.8	32.0
Non-Hispanic white, male												
2002	6.2	*	*	*	*	0.9	2.4	5.7	10.8	20.0	35.0	56.3
2001	6.3	*	*	*	0.2	0.8	2.4	5.9	11.6	21.4	33.7	60.4
2000	6.0	*	*	*	0.2	0.8	2.2	6.0	11.0	20.3	32.7	62.5
1999	6.0	*	*	*	*	0.8	2.6	5.8	11.4	20.5	33.1	56.5
Non-Hispanic white, female												
2002	3.3	*	*	*	*	0.6	1.5	2.6	4.5	7.8	13.1	22.8
2001	3.3	*	*	*	*	0.6	1.5	3.1	4.8	7.7	12.4	23.6
2000	3.3	*	*	*	*	0.6	1.6	3.1	4.8	7.9	11.7	24.0
1999	3.2	*	*	*	*	0.6	1.4	3.0	4.6	7.8	12.7	22.2
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	1.1	2.5	3.8	4.0	7.9
2001	0.7	*	*	*	*	*	*	0.9	2.3	4.5	4.6	8.1
2000	0.8	*	*	*	*	*	0.4	1.0	2.9	4.3	5.7	8.3
1999	0.7	*	*	*	*	*	*	1.3	2.2	3.1	5.1	6.8
Non-Hispanic black, male												
2002	1.0	*	*	*	*	*	*	1.7	4.2	6.4	*	*
2001	0.9	*	*	*	*	*	*	1.6	3.5	7.3	*	*
2000	0.9	*	*	*	*	*	*	1.5	4.2	5.5	8.6	*
1999	1.0	*	*	*	*	*	*	2.2	3.8	4.8	9.1	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	2.5	4.7	8.6
2000	0.7	*	*	*	*	*	*	*	1.8	3.4	4.0	10.0
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All origins 3/, both sexes												
2002	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
All origins 3/, male												
2002	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
All origins 3/, female												
2002	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	0.4	0.7	1.2	*	4.8	*
2001	0.4	*	*	*	*	*	0.4	0.7	1.4	2.5	5.0	*
2000	0.4	*	*	*	*	*	0.5	0.9	1.6	2.4	*	*
1999	0.4	*	*	*	*	*	*	0.7	1.8	2.6	4.2	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	2.9	*	*	*	0.1	0.6	1.5	3.0	5.4	9.3	14.1	18.2
2001	3.0	*	*	*	0.1	0.5	1.5	3.3	5.6	9.7	13.7	18.3
2000	3.0	*	*	*	0.1	0.5	1.5	3.4	5.7	9.3	13.5	19.3
1999	2.9	*	*	*	0.1	0.6	1.5	3.3	5.6	9.3	13.7	16.7
Non-Hispanic 4/, male												
2002	3.8	*	*	*	*	0.7	1.8	4.1	7.6	13.7	22.0	30.2
2001	3.9	*	*	*	0.1	0.6	1.9	4.2	7.8	14.6	22.0	31.0
2000	3.7	*	*	*	0.1	0.6	1.7	4.4	7.8	13.4	21.3	32.6
1999	3.7	*	*	*	0.1	0.6	2.0	4.2	7.8	13.7	21.5	28.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Non-Hispanic 4/, female												
2002	2.1	*	*	*	*	0.4	1.2	2.0	3.3	5.7	8.9	12.9
2001	2.1	*	*	*	*	0.5	1.2	2.4	3.5	5.6	8.3	12.9
2000	2.2	*	*	*	*	0.4	1.3	2.4	3.7	5.9	8.4	13.9
1999	2.1	*	*	*	*	0.5	1.1	2.3	3.5	5.7	8.7	12.0
Non-Hispanic white, both sexes												
2002	3.6	*	*	*	0.1	0.7	1.8	3.6	6.2	10.5	15.6	19.6
2001	3.7	*	*	*	0.2	0.7	1.9	4.0	6.5	10.8	15.0	19.9
2000	3.6	*	*	*	0.2	0.7	1.8	4.0	6.5	10.4	14.7	20.8
1999	3.5	*	*	*	0.1	0.7	1.9	3.9	6.4	10.5	14.9	17.9
Non-Hispanic white, male												
2002	4.7	*	*	*	*	0.9	2.2	4.8	8.6	15.3	24.1	32.9
2001	4.8	*	*	*	0.2	0.8	2.3	5.0	9.0	16.3	24.1	34.1
2000	4.6	*	*	*	0.2	0.8	2.0	5.2	8.9	15.0	23.2	35.7
1999	4.5	*	*	*	*	0.8	2.4	5.0	8.9	15.4	23.3	30.5
Non-Hispanic white, female												
2002	2.6	*	*	*	*	0.6	1.4	2.4	3.9	6.5	9.8	13.8
2001	2.6	*	*	*	*	0.6	1.5	2.9	4.1	6.2	9.0	13.9
2000	2.7	*	*	*	*	0.6	1.6	2.8	4.2	6.6	9.1	14.7
1999	2.5	*	*	*	*	0.6	1.4	2.8	4.1	6.4	9.4	12.9
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	0.8	1.5	*	*
2001	0.3	*	*	*	*	*	*	*	*	2.2	2.6	*
2000	0.4	*	*	*	*	*	*	*	*	1.9	3.6	6.7
1999	0.3	*	*	*	*	*	*	*	*	1.4	2.8	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	2.1	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other malignant neoplasms of skin (C44)												
All origins 3/, both sexes												
2002	0.8	*	*	*	*	*	0.1	0.5	1.3	2.7	5.8	12.0
2001	0.9	*	*	*	*	*	0.1	0.5	1.5	2.9	5.4	13.4
2000	0.8	*	*	*	*	*	0.1	0.5	1.3	3.0	4.9	12.9
1999	0.8	*	*	*	*	*	0.1	0.6	1.4	2.8	5.4	12.8
All origins 3/, male												
2002	1.2	*	*	*	*	*	0.2	0.8	2.2	4.4	9.9	21.0
2001	1.2	*	*	*	*	*	0.1	0.9	2.5	4.8	8.7	24.1
2000	1.1	*	*	*	*	*	0.2	0.8	2.1	4.9	8.8	23.6
1999	1.2	*	*	*	*	*	0.2	0.9	2.4	4.7	9.1	23.1
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.6	1.2	3.1	8.1
2001	0.5	*	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9
2000	0.5	*	*	*	*	*	*	0.2	0.6	1.3	2.4	8.5
1999	0.5	*	*	*	*	*	*	0.2	0.5	1.3	3.0	8.6
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	3.5	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	1.9	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	*	*	*	*	*	0.1	0.5	1.4	2.8	5.9	12.3
2001	1.0	*	*	*	*	*	0.1	0.5	1.6	3.0	5.5	13.5
2000	0.9	*	*	*	*	*	0.1	0.5	1.3	3.0	5.0	13.2
1999	0.9	*	*	*	*	*	0.2	0.6	1.5	2.9	5.5	12.9
Non-Hispanic 4/, male												
2002	1.3	*	*	*	*	*	0.2	0.9	2.2	4.6	10.0	21.6
2001	1.3	*	*	*	*	*	0.1	0.9	2.6	4.9	8.9	24.4
2000	1.3	*	*	*	*	*	0.2	0.8	2.1	5.0	8.9	24.4
1999	1.3	*	*	*	*	*	0.3	0.9	2.4	4.8	9.2	23.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other malignant neoplasms of skin (C44)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.1	8.2
2001	0.6	*	*	*	*	*	*	0.2	0.7	1.4	3.3	9.0
2000	0.5	*	*	*	*	*	*	0.2	0.6	1.3	2.4	8.6
1999	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.1	8.6
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	*	*	0.1	0.5	1.4	2.9	6.4	13.3
2001	1.1	*	*	*	*	*	0.1	0.5	1.6	3.1	5.9	14.5
2000	1.0	*	*	*	*	*	0.1	0.5	1.3	3.1	5.3	14.3
1999	1.1	*	*	*	*	*	0.2	0.5	1.5	3.1	5.9	14.1
Non-Hispanic white, male												
2002	1.5	*	*	*	*	*	0.2	0.9	2.2	4.7	10.9	23.4
2001	1.5	*	*	*	*	*	0.1	0.9	2.7	5.0	9.6	26.3
2000	1.4	*	*	*	*	*	0.1	0.8	2.1	5.3	9.5	26.8
1999	1.5	*	*	*	*	*	0.3	0.9	2.5	5.1	9.8	26.0
Non-Hispanic white, female												
2002	0.7	*	*	*	*	*	*	0.2	0.6	1.4	3.3	8.9
2001	0.7	*	*	*	*	*	*	0.2	0.7	1.5	3.5	9.6
2000	0.6	*	*	*	*	*	*	0.2	0.6	1.3	2.6	9.3
1999	0.7	*	*	*	*	*	*	0.2	0.6	1.4	3.3	9.3
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.8	1.7	2.3	2.1	*
2001	0.4	*	*	*	*	*	*	0.7	1.7	2.3	*	*
2000	0.4	*	*	*	*	*	*	0.7	2.1	2.4	*	*
1999	0.4	*	*	*	*	*	*	1.0	1.6	1.7	2.2	*
Non-Hispanic black, male												
2002	0.7	*	*	*	*	*	*	1.3	3.1	4.7	*	*
2001	0.7	*	*	*	*	*	*	1.4	3.1	4.9	*	*
2000	0.6	*	*	*	*	*	*	1.2	3.3	3.4	*	*
1999	0.7	*	*	*	*	*	*	1.9	2.8	3.5	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
All origins 3/, both sexes												
2002	2.4	*	0.1	0.2	0.3	0.4	0.8	1.6	4.3	9.1	14.9	15.7
2001	2.4	*	*	0.1	0.3	0.5	0.8	1.8	4.2	9.1	15.1	14.9
2000	2.4	*	0.1	0.1	0.4	0.5	0.8	1.8	4.4	9.6	15.1	15.7
1999	2.4	*	*	0.2	0.4	0.5	0.9	1.8	4.4	9.3	14.8	14.8
All origins 3/, male												
2002	2.8	*	*	0.2	0.3	0.5	0.9	1.6	5.0	12.3	22.5	28.1
2001	2.8	*	*	0.1	0.4	0.6	0.8	2.0	4.8	12.5	22.7	25.9
2000	2.9	*	*	0.1	0.4	0.6	0.8	1.8	5.3	12.9	23.5	28.4
1999	2.8	*	*	0.2	0.4	0.6	1.1	1.9	5.5	12.8	22.6	24.6
All origins 3/, female												
2002	1.9	*	*	0.2	0.3	0.3	0.7	1.5	3.7	6.4	9.8	10.4
2001	2.0	*	*	0.1	0.3	0.4	0.7	1.6	3.7	6.3	10.1	10.3
2000	2.0	*	*	0.1	0.3	0.5	0.7	1.7	3.6	6.9	9.6	10.5
1999	1.9	*	*	0.2	0.3	0.4	0.7	1.7	3.5	6.4	9.6	10.8
Hispanic, both sexes												
2002	0.9	*	*	*	0.3	0.3	0.7	1.1	3.0	6.1	11.2	*
2001	0.9	*	*	*	0.3	*	0.6	1.5	3.0	6.0	9.2	*
2000	0.8	*	*	*	*	0.4	*	1.3	2.1	6.3	11.3	*
1999	0.9	*	*	*	0.4	0.4	0.7	1.4	3.6	5.9	8.1	*
Hispanic, male												
2002	1.0	*	*	*	*	*	0.8	*	3.9	7.1	19.3	*
2001	0.9	*	*	*	*	*	*	1.2	3.6	7.0	14.3	*
2000	0.9	*	*	*	*	*	*	*	3.0	7.6	18.2	*
1999	1.0	*	*	*	*	*	0.8	1.4	4.0	8.3	*	*
Hispanic, female												
2002	0.8	*	*	*	*	*	*	1.5	2.2	5.3	5.6	*
2001	0.8	*	*	*	*	*	*	1.8	2.6	5.2	*	*
2000	0.7	*	*	*	*	*	*	1.5	*	5.3	6.6	*
1999	0.9	*	*	*	*	*	*	1.5	3.2	4.0	7.6	*
Non-Hispanic 4/, both sexes												
2002	2.6	*	0.2	0.2	0.3	0.5	0.8	1.6	4.4	9.2	15.0	15.9
2001	2.6	*	*	0.1	0.3	0.5	0.8	1.8	4.3	9.3	15.3	15.1
2000	2.6	*	*	0.1	0.4	0.5	0.8	1.8	4.6	9.8	15.2	16.1
1999	2.6	*	*	0.2	0.3	0.5	0.9	1.9	4.5	9.5	15.0	15.0
Non-Hispanic 4/, male												
2002	3.1	*	*	0.2	0.3	0.6	1.0	1.7	5.1	12.6	22.6	28.7
2001	3.1	*	*	0.1	0.4	0.6	0.8	2.1	4.8	12.8	23.1	26.2
2000	3.1	*	*	*	0.4	0.6	0.9	1.9	5.5	13.1	23.7	29.3
1999	3.1	*	*	0.2	0.4	0.6	1.1	2.0	5.5	13.1	23.1	25.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of mesothelial and soft tissue (C45-C49)												
Non-Hispanic 4/, female												
2002	2.1	*	*	0.2	0.3	0.4	0.7	1.5	3.8	6.5	9.9	10.4
2001	2.1	*	*	0.1	0.3	0.5	0.7	1.6	3.8	6.3	10.3	10.4
2000	2.2	*	*	*	0.3	0.5	0.8	1.7	3.7	7.0	9.7	10.8
1999	2.1	*	*	0.1	0.3	0.4	0.7	1.7	3.5	6.5	9.7	10.9
Non-Hispanic white, both sexes												
2002	2.9	*	*	0.2	0.3	0.4	0.8	1.6	4.5	9.7	15.7	16.7
2001	2.9	*	*	0.1	0.3	0.5	0.8	1.8	4.5	9.9	16.1	15.6
2000	3.0	*	*	0.1	0.4	0.6	0.8	1.8	4.7	10.2	16.1	16.6
1999	2.8	*	*	0.2	0.3	0.5	0.8	1.9	4.6	9.9	15.5	15.5
Non-Hispanic white, male												
2002	3.5	*	*	0.2	0.4	0.5	0.9	1.8	5.3	13.4	23.9	30.5
2001	3.5	*	*	*	0.4	0.6	0.8	2.1	5.1	13.9	24.3	27.4
2000	3.6	*	*	*	0.4	0.6	0.8	2.0	5.7	13.8	25.0	30.5
1999	3.5	*	*	0.2	0.3	0.6	1.0	2.0	5.8	13.7	24.2	26.9
Non-Hispanic white, female												
2002	2.3	*	*	*	0.3	0.4	0.7	1.5	3.8	6.6	10.3	10.7
2001	2.3	*	*	*	0.3	0.5	0.7	1.5	3.9	6.6	10.7	10.7
2000	2.4	*	*	*	0.3	0.5	0.8	1.6	3.8	7.2	10.2	11.0
1999	2.2	*	*	*	0.3	0.4	0.6	1.7	3.5	6.7	9.8	10.9
Non-Hispanic black, both sexes												
2002	1.6	*	*	*	0.4	0.7	1.1	1.8	4.2	7.0	10.4	7.6
2001	1.5	*	*	*	0.4	0.6	0.9	2.3	3.8	6.0	8.5	10.6
2000	1.6	*	*	*	0.5	0.6	1.2	2.2	4.3	8.3	8.4	11.1
1999	1.7	*	*	*	0.6	0.7	1.4	1.9	4.3	6.7	11.7	10.7
Non-Hispanic black, male												
2002	1.6	*	*	*	*	0.9	1.4	2.1	4.4	7.9	13.2	*
2001	1.5	*	*	*	0.7	0.8	1.1	2.2	3.9	6.4	11.1	*
2000	1.7	*	*	*	*	*	1.5	1.7	5.1	10.2	12.6	*
1999	1.8	*	*	*	0.8	0.8	1.7	1.9	5.2	8.1	14.7	*
Non-Hispanic black, female												
2002	1.5	*	*	*	*	*	0.9	1.6	4.1	6.4	8.8	*
2001	1.4	*	*	*	*	*	0.7	2.4	3.7	5.6	7.1	8.6
2000	1.6	*	*	*	*	*	1.1	2.7	3.7	7.0	6.1	8.7
1999	1.6	*	*	*	*	*	1.1	2.0	3.6	5.7	10.0	10.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mesothelioma (C45)												
All origins 3/, both sexes												
2002	0.8	*	*	*	*	*	0.1	0.3	1.4	4.0	7.2	5.9
2001	0.8	*	*	*	*	*	0.1	0.4	1.3	3.9	7.1	5.5
2000	0.8	*	*	*	*	*	0.1	0.3	1.5	4.2	6.9	5.7
1999	0.8	*	*	*	*	*	0.1	0.4	1.6	4.2	6.8	4.8
All origins 3/, male												
2002	1.4	*	*	*	*	*	0.1	0.4	2.4	7.6	14.7	15.9
2001	1.4	*	*	*	*	*	*	0.5	2.2	7.3	14.4	14.5
2000	1.4	*	*	*	*	*	*	0.5	2.5	7.8	14.2	15.5
1999	1.4	*	*	*	*	*	0.1	0.6	2.5	7.7	14.0	12.7
All origins 3/, female												
2002	0.3	*	*	*	*	*	*	0.1	0.5	1.0	2.2	1.6
2001	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.3	1.7
2000	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	1.8
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.3	2.1	1.6
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	2.5	5.0	*
2001	0.2	*	*	*	*	*	*	*	*	1.8	4.3	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	2.3	*	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	4.2	9.8	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	*	*	*	*	*	0.1	0.3	1.4	4.1	7.2	6.1
2001	0.9	*	*	*	*	*	0.1	0.4	1.4	4.0	7.2	5.6
2000	0.9	*	*	*	*	*	0.1	0.3	1.5	4.4	7.0	5.9
1999	0.9	*	*	*	*	*	0.1	0.4	1.6	4.3	7.0	4.8
Non-Hispanic 4/, male												
2002	1.6	*	*	*	*	*	0.1	0.4	2.4	7.8	14.9	16.4
2001	1.5	*	*	*	*	*	*	0.6	2.3	7.5	14.6	14.8
2000	1.6	*	*	*	*	*	*	0.5	2.6	8.0	14.4	16.1
1999	1.5	*	*	*	*	*	*	0.6	2.6	7.9	14.4	13.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mesothelioma (C45)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	*	0.1	0.5	1.0	2.2	1.6
2001	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.3	1.8
2000	0.4	*	*	*	*	*	*	0.2	0.5	1.4	2.1	1.8
1999	0.3	*	*	*	*	*	*	0.2	0.7	1.3	2.1	1.5
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	*	*	0.1	0.3	1.6	4.5	7.8	6.6
2001	1.1	*	*	*	*	*	0.1	0.4	1.6	4.5	7.8	6.1
2000	1.1	*	*	*	*	*	0.1	0.3	1.6	4.7	7.5	6.3
1999	1.1	*	*	*	*	*	0.1	0.4	1.8	4.7	7.3	5.1
Non-Hispanic white, male												
2002	1.9	*	*	*	*	*	0.1	0.4	2.7	8.5	15.8	17.8
2001	1.8	*	*	*	*	*	*	0.6	2.6	8.4	15.7	16.0
2000	1.8	*	*	*	*	*	*	0.5	2.8	8.5	15.5	17.3
1999	1.8	*	*	*	*	*	*	0.6	2.8	8.7	15.2	14.1
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	0.2	0.5	1.1	2.4	1.7
2001	0.4	*	*	*	*	*	*	0.2	0.6	1.2	2.5	2.0
2000	0.4	*	*	*	*	*	*	0.2	0.5	1.5	2.3	1.9
1999	0.4	*	*	*	*	*	*	0.2	0.7	1.4	2.2	1.5
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	1.6	3.3	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	0.9	2.6	2.2	*
1999	0.3	*	*	*	*	*	*	*	0.9	1.4	3.7	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	3.4	7.3	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	5.1	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	6.9	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Kaposi's sarcoma (C46)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Kaposi's sarcoma (C46)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)												
All origins 3/, both sexes												
2002	1.5	*	0.1	0.2	0.3	0.4	0.7	1.3	2.9	5.1	7.6	9.3
2001	1.5	*	*	0.1	0.3	0.5	0.7	1.4	2.8	5.2	8.0	9.1
2000	1.6	*	0.1	0.1	0.4	0.5	0.7	1.4	2.9	5.3	8.1	9.6
1999	1.5	*	*	0.2	0.4	0.5	0.8	1.5	2.9	5.1	7.9	9.6
All origins 3/, male												
2002	1.4	*	*	0.2	0.3	0.5	0.8	1.3	2.6	4.7	7.7	11.7
2001	1.4	*	*	0.1	0.4	0.5	0.7	1.4	2.5	5.2	8.2	10.7
2000	1.4	*	*	0.1	0.4	0.5	0.7	1.4	2.7	5.0	9.2	12.5
1999	1.5	*	*	0.2	0.4	0.5	0.9	1.4	2.9	5.1	8.5	11.3
All origins 3/, female												
2002	1.6	*	*	0.2	0.3	0.3	0.6	1.3	3.2	5.4	7.6	8.3
2001	1.6	*	*	0.1	0.2	0.4	0.7	1.4	3.2	5.1	7.8	8.4
2000	1.7	*	*	0.1	0.3	0.5	0.7	1.5	3.1	5.6	7.4	8.4
1999	1.6	*	*	0.2	0.3	0.4	0.6	1.5	2.8	5.1	7.5	8.9
Hispanic, both sexes												
2002	0.7	*	*	*	0.3	0.3	0.6	1.0	2.2	3.6	5.5	*
2001	0.7	*	*	*	0.3	*	0.5	1.3	2.5	4.1	4.7	*
2000	0.6	*	*	*	*	0.4	*	1.0	1.4	4.6	7.7	*
1999	0.7	*	*	*	0.4	0.3	0.6	1.1	2.8	3.6	5.6	*
Hispanic, male												
2002	0.6	*	*	*	*	*	*	*	2.5	*	8.2	*
2001	0.6	*	*	*	*	*	*	*	2.8	4.0	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	10.8	*
1999	0.7	*	*	*	*	*	*	*	2.7	4.4	*	*
Hispanic, female												
2002	0.7	*	*	*	*	*	*	1.4	2.0	4.3	*	*
2001	0.7	*	*	*	*	*	*	1.6	2.2	4.3	*	*
2000	0.7	*	*	*	*	*	*	*	*	5.1	*	*
1999	0.7	*	*	*	*	*	*	*	2.9	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.6	*	0.2	0.2	0.3	0.4	0.7	1.3	3.0	5.1	7.7	9.5
2001	1.6	*	*	0.1	0.3	0.5	0.7	1.4	2.9	5.2	8.1	9.1
2000	1.7	*	*	0.1	0.4	0.5	0.8	1.5	3.0	5.4	8.1	9.9
1999	1.6	*	*	0.2	0.3	0.5	0.8	1.5	2.9	5.2	8.0	9.7
Non-Hispanic 4/, male												
2002	1.5	*	*	0.2	0.3	0.5	0.8	1.3	2.6	4.8	7.6	11.8
2001	1.5	*	*	0.1	0.4	0.6	0.8	1.5	2.5	5.3	8.4	10.7
2000	1.6	*	*	*	0.4	0.5	0.8	1.4	2.8	5.1	9.1	12.8
1999	1.6	*	*	0.2	0.4	0.6	1.0	1.4	2.9	5.1	8.6	11.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)												
Non-Hispanic 4/, female												
2002	1.8	*	*	0.1	0.3	0.3	0.6	1.3	3.3	5.4	7.7	8.4
2001	1.8	*	*	0.1	0.2	0.4	0.7	1.4	3.2	5.2	7.9	8.4
2000	1.8	*	*	*	0.3	0.5	0.7	1.5	3.2	5.6	7.5	8.7
1999	1.7	*	*	0.1	0.3	0.4	0.6	1.6	2.8	5.2	7.6	9.0
Non-Hispanic white, both sexes												
2002	1.8	*	*	0.2	0.3	0.4	0.7	1.3	2.9	5.2	7.9	9.7
2001	1.8	*	*	0.1	0.3	0.5	0.7	1.4	2.9	5.4	8.3	9.2
2000	1.8	*	*	0.1	0.4	0.5	0.7	1.5	3.1	5.5	8.4	10.0
1999	1.7	*	*	0.2	0.3	0.4	0.7	1.5	2.9	5.2	8.0	9.9
Non-Hispanic white, male												
2002	1.6	*	*	0.2	0.4	0.5	0.8	1.3	2.6	4.9	8.0	12.1
2001	1.6	*	*	*	0.4	0.6	0.7	1.5	2.5	5.5	8.5	10.7
2000	1.7	*	*	*	0.4	0.6	0.7	1.5	2.9	5.2	9.3	12.8
1999	1.7	*	*	0.2	0.3	0.5	0.9	1.4	3.0	5.1	8.7	12.2
Non-Hispanic white, female												
2002	1.9	*	*	*	0.3	0.3	0.6	1.3	3.2	5.4	7.9	8.7
2001	1.9	*	*	*	0.3	0.5	0.7	1.3	3.3	5.3	8.1	8.5
2000	1.9	*	*	*	0.3	0.5	0.7	1.4	3.3	5.7	7.8	8.8
1999	1.8	*	*	*	0.3	0.4	0.5	1.5	2.8	5.3	7.6	9.0
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	0.4	0.6	1.0	1.7	3.6	5.4	7.1	6.0
2001	1.3	*	*	*	0.4	0.6	0.8	2.0	3.3	4.8	6.8	9.4
2000	1.3	*	*	*	0.5	0.4	1.1	2.0	3.3	5.5	5.9	9.2
1999	1.4	*	*	*	0.6	0.7	1.3	1.6	3.4	5.3	8.0	8.4
Non-Hispanic black, male												
2002	1.2	*	*	*	*	*	1.1	1.8	3.3	4.5	5.9	*
2001	1.2	*	*	*	0.7	*	1.0	1.7	2.9	4.3	7.2	*
2000	1.1	*	*	*	*	*	1.3	1.2	3.2	4.6	6.8	*
1999	1.3	*	*	*	0.8	*	1.5	1.4	3.4	5.7	7.8	*
Non-Hispanic black, female												
2002	1.4	*	*	*	*	*	0.8	1.6	3.9	6.0	7.8	*
2001	1.4	*	*	*	*	*	0.7	2.2	3.6	5.2	6.6	8.6
2000	1.5	*	*	*	*	*	1.0	2.6	3.4	6.1	5.4	*
1999	1.4	*	*	*	*	*	1.0	1.8	3.4	5.0	8.1	9.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
All origins 3/, both sexes												
2002	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
All origins 3/, female												
2002	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8
Hispanic, both sexes												
2002	4.5	*	*	*	*	0.7	3.8	11.0	17.6	27.4	43.7	71.0
2001	4.8	*	*	*	*	0.8	4.5	12.0	22.1	28.0	39.5	68.8
2000	4.8	*	*	*	*	0.8	4.3	12.1	21.0	31.3	45.6	71.7
1999	4.6	*	*	*	*	0.7	4.0	12.2	19.6	28.0	43.5	86.3
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	9.2	*	*	*	*	1.6	7.8	21.6	33.5	48.7	73.1	105.3
2001	9.7	*	*	*	*	1.7	9.2	23.8	41.7	49.3	64.7	102.4
2000	9.7	*	*	*	*	1.8	8.7	23.9	39.1	54.9	74.9	105.8
1999	9.3	*	*	*	*	1.6	8.3	23.9	36.6	49.1	71.8	127.9
Non-Hispanic 4/, both sexes												
2002	16.0	*	*	*	*	1.1	6.4	16.6	30.3	47.7	78.0	136.8
2001	16.1	*	*	*	*	1.2	6.5	17.2	30.7	48.8	78.4	136.4
2000	16.5	*	*	*	*	1.2	6.5	17.3	31.9	50.1	80.1	149.7
1999	16.3	*	*	*	*	1.2	6.3	17.3	31.9	50.6	80.2	146.2
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.3	3.5
2001	0.3	*	*	*	*	*	*	0.3	0.6	1.5	2.3	3.7
2000	0.3	*	*	*	*	*	*	0.3	0.7	1.4	2.2	4.2
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.1	4.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Non-Hispanic 4/, female												
2002	31.0	*	*	*	*	2.2	12.6	32.3	57.9	86.7	128.2	194.4
2001	31.1	*	*	*	*	2.5	12.8	33.6	58.6	88.1	128.3	191.6
2000	31.8	*	*	*	*	2.4	12.8	33.8	60.7	90.3	130.8	208.6
1999	31.4	*	*	*	*	2.3	12.4	33.6	60.7	91.2	130.9	202.8
Non-Hispanic white, both sexes												
2002	16.9	*	*	*	*	0.9	5.6	15.2	29.4	47.4	78.0	137.7
2001	17.0	*	*	*	*	1.0	5.8	16.0	29.6	48.2	78.7	137.1
2000	17.4	*	*	*	*	1.1	5.8	16.1	30.8	50.3	80.6	149.6
1999	17.1	*	*	*	*	1.0	5.6	16.0	30.6	50.2	80.3	147.8
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.5
2000	0.4	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.0
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.1	2.0	4.5
Non-Hispanic white, female												
2002	32.9	*	*	*	*	1.9	11.1	30.0	56.7	87.0	128.9	195.8
2001	32.9	*	*	*	*	2.0	11.5	31.6	57.0	88.0	129.4	192.6
2000	33.8	*	*	*	*	2.1	11.6	31.7	59.2	91.4	132.2	208.3
1999	33.3	*	*	*	*	2.0	11.1	31.5	58.9	91.2	131.5	204.9
Non-Hispanic black, both sexes												
2002	15.4	*	*	*	*	2.4	12.0	27.7	43.7	61.1	95.3	154.0
2001	15.5	*	*	*	*	2.7	11.5	27.9	45.6	64.6	93.0	153.9
2000	15.3	*	*	*	*	2.3	11.5	28.4	46.4	59.4	91.7	178.7
1999	15.5	*	*	*	*	2.4	12.0	29.2	47.1	64.5	95.6	149.1
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	29.1	*	*	*	*	4.5	22.5	51.0	77.8	102.7	147.1	210.7
2001	29.2	*	*	*	*	5.1	21.5	51.3	81.6	108.4	143.0	208.3
2000	28.9	*	*	*	*	4.3	21.5	52.5	82.5	100.1	141.5	241.6
1999	29.2	*	*	*	*	4.5	22.3	53.7	83.7	108.2	147.2	201.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of female genital organs (C51-C58)												
All origins 3/, both sexes												
2002	9.4	*	*	*	0.1	0.9	3.0	8.2	18.3	34.9	56.9	77.5
2001	9.4	*	*	*	0.1	0.9	3.0	8.4	18.2	34.7	58.2	79.9
2000	9.4	*	*	*	0.1	0.9	3.0	8.3	18.5	34.8	57.7	82.3
1999	9.3	*	*	*	0.1	0.8	3.1	8.3	18.5	35.0	56.9	79.1
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	18.5	*	*	*	0.3	1.7	5.9	16.2	35.3	63.9	94.6	111.2
2001	18.5	*	*	*	0.2	1.8	6.0	16.4	35.1	63.4	96.4	113.4
2000	18.4	*	*	*	0.2	1.9	6.0	16.3	35.6	63.5	95.3	115.8
1999	18.2	*	*	*	0.2	1.6	6.1	16.4	35.6	63.6	93.8	110.9
Hispanic, both sexes												
2002	3.6	*	*	*	*	0.6	2.7	6.8	12.9	26.8	44.5	61.0
2001	3.8	*	*	*	*	0.8	2.9	7.3	15.6	26.6	41.5	66.9
2000	3.8	*	*	*	*	0.7	3.2	7.0	15.0	28.6	47.4	75.6
1999	3.5	*	*	*	*	0.7	2.7	7.3	14.1	26.6	42.3	59.6
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	7.5	*	*	*	*	1.4	5.6	13.6	24.5	48.2	75.0	91.2
2001	7.8	*	*	*	*	1.8	6.1	14.4	29.4	47.6	69.5	99.6
2000	7.9	*	*	*	*	1.6	6.6	13.8	28.2	51.1	79.2	112.8
1999	7.2	*	*	*	*	1.5	5.5	14.3	26.6	47.5	70.4	88.4
Non-Hispanic 4/, both sexes												
2002	10.3	*	*	*	0.1	0.9	3.0	8.4	18.7	35.3	57.4	78.1
2001	10.2	*	*	*	0.1	0.9	3.0	8.4	18.4	35.1	58.9	80.2
2000	10.2	*	*	*	0.1	1.0	3.0	8.4	18.7	35.1	58.0	82.4
1999	10.0	*	*	*	0.1	0.8	3.1	8.4	18.8	35.4	57.2	79.6
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of female genital organs (C51-C58)												
Non-Hispanic 4/, female												
2002	20.0	*	*	*	0.3	1.8	6.0	16.4	36.0	64.8	95.4	111.8
2001	20.0	*	*	*	0.2	1.8	6.0	16.6	35.4	64.3	97.5	113.5
2000	19.8	*	*	*	0.2	1.9	5.9	16.4	36.1	64.1	95.8	115.7
1999	19.6	*	*	*	0.2	1.7	6.1	16.5	36.1	64.4	94.4	111.5
Non-Hispanic white, both sexes												
2002	10.9	*	*	*	0.1	0.9	2.9	8.2	18.3	34.2	56.8	77.3
2001	10.9	*	*	*	0.1	0.9	2.9	8.3	17.9	34.2	58.6	79.8
2000	10.8	*	*	*	0.1	0.9	2.8	8.2	18.4	33.9	57.5	82.1
1999	10.7	*	*	*	0.1	0.8	3.0	8.4	18.4	34.2	56.6	79.4
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	21.4	*	*	*	0.3	1.8	5.8	16.2	35.7	63.4	94.9	110.7
2001	21.4	*	*	*	0.2	1.7	5.8	16.5	34.7	63.2	97.4	113.1
2000	21.1	*	*	*	0.2	1.9	5.5	16.3	35.7	62.4	95.2	115.2
1999	21.0	*	*	*	0.2	1.7	6.0	16.7	35.7	62.8	93.7	111.0
Non-Hispanic black, both sexes												
2002	8.9	*	*	*	*	1.3	4.2	10.0	23.9	50.7	71.6	99.8
2001	8.8	*	*	*	*	1.4	4.2	9.6	24.4	50.0	72.1	95.5
2000	8.9	*	*	*	*	1.3	4.4	9.8	24.3	52.6	74.3	97.0
1999	8.7	*	*	*	*	1.0	4.0	9.1	25.4	53.2	72.3	92.5
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	17.0	*	*	*	*	2.4	7.9	18.6	43.1	86.7	112.1	138.0
2001	16.7	*	*	*	*	2.6	7.9	17.8	44.0	85.3	112.5	130.9
2000	17.0	*	*	*	*	2.5	8.2	18.1	43.9	89.5	116.2	133.0
1999	16.5	*	*	*	*	2.0	7.5	16.8	45.8	90.8	112.8	127.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of vulva and vagina (C51-C52)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.9	7.7
2001	0.4	*	*	*	*	*	0.1	0.2	0.4	1.1	3.1	7.3
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	1.3	3.0	7.7
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.2	3.1	8.6
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.8	*	*	*	*	*	0.2	0.4	0.9	2.0	4.9	11.0
2001	0.8	*	*	*	*	*	0.2	0.4	0.8	2.1	5.1	10.3
2000	0.8	*	*	*	*	*	0.2	0.4	0.8	2.3	4.9	10.9
1999	0.8	*	*	*	*	*	0.1	0.4	0.9	2.1	5.0	12.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	3.0	7.7
2001	0.4	*	*	*	*	*	0.1	0.2	0.4	1.1	3.1	7.3
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.3	2.9	7.7
1999	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	3.0	8.5
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of vulva and vagina (C51-C52)												
Non-Hispanic 4/, female												
2002	0.9	*	*	*	*	*	0.2	0.4	0.9	2.0	4.9	11.0
2001	0.9	*	*	*	*	*	0.2	0.5	0.8	2.1	5.2	10.3
2000	0.9	*	*	*	*	*	0.2	0.4	0.9	2.4	4.8	10.8
1999	0.9	*	*	*	*	*	0.2	0.4	0.9	2.2	5.0	11.9
Non-Hispanic white, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.5	1.1	3.0	8.0
2001	0.5	*	*	*	*	*	0.1	0.2	0.4	1.1	3.2	7.3
2000	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.0	7.9
1999	0.5	*	*	*	*	*	0.1	0.2	0.4	1.2	3.0	9.0
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	1.0	*	*	*	*	*	0.2	0.4	1.0	2.1	5.1	11.5
2001	0.9	*	*	*	*	*	0.2	0.4	0.7	2.1	5.3	10.4
2000	1.0	*	*	*	*	*	0.1	0.4	0.9	2.5	5.0	11.1
1999	1.0	*	*	*	*	*	0.2	0.4	0.9	2.2	5.0	12.5
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2001	0.4	*	*	*	*	*	*	*	0.9	1.5	3.5	6.6
2000	0.3	*	*	*	*	*	*	*	*	*	2.7	7.0
1999	0.3	*	*	*	*	*	*	*	*	1.4	3.1	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.8	*
2001	0.7	*	*	*	*	*	*	*	1.6	2.6	5.4	9.0
2000	0.5	*	*	*	*	*	*	*	*	*	4.2	9.6
1999	0.5	*	*	*	*	*	*	*	*	2.3	4.9	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of cervix uteri (C53)												
All origins 3/, both sexes												
2002	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1
Hispanic, both sexes												
2002	1.0	*	*	*	*	0.4	1.5	2.4	3.6	4.7	6.7	*
2001	1.1	*	*	*	*	0.5	1.9	3.0	4.1	3.4	7.0	*
2000	1.2	*	*	*	*	0.5	2.1	2.8	4.6	5.6	8.5	13.9
1999	1.1	*	*	*	*	0.5	1.8	3.3	3.9	4.9	6.5	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	2.1	*	*	*	*	0.9	3.1	4.7	6.8	8.4	11.2	*
2001	2.3	*	*	*	*	1.1	3.9	6.0	7.7	6.1	11.8	*
2000	2.5	*	*	*	*	1.1	4.4	5.5	8.6	10.0	14.2	20.8
1999	2.3	*	*	*	*	1.1	3.6	6.6	7.4	8.8	10.8	*
Non-Hispanic 4/, both sexes												
2002	1.4	*	*	*	*	0.5	1.5	2.2	2.6	3.4	4.0	6.1
2001	1.5	*	*	*	*	0.5	1.5	2.4	2.8	3.2	4.5	7.0
2000	1.5	*	*	*	*	0.6	1.5	2.4	3.0	3.6	4.7	7.1
1999	1.6	*	*	*	*	0.5	1.5	2.4	3.2	3.7	4.6	7.9
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of cervix uteri (C53)												
Non-Hispanic 4/, female												
2002	2.8	*	*	*	*	1.1	2.9	4.3	5.1	6.2	6.7	8.7
2001	2.9	*	*	*	*	1.0	3.0	4.6	5.3	5.8	7.5	10.0
2000	3.0	*	*	*	*	1.2	2.9	4.7	5.7	6.6	7.7	9.9
1999	3.0	*	*	*	*	1.1	3.0	4.7	6.1	6.7	7.5	11.0
Non-Hispanic white, both sexes												
2002	1.3	*	*	*	*	0.5	1.3	1.9	2.3	2.7	3.4	4.9
2001	1.4	*	*	*	*	0.5	1.4	2.1	2.4	2.6	3.8	6.1
2000	1.4	*	*	*	*	0.5	1.3	2.1	2.5	2.9	4.1	6.0
1999	1.4	*	*	*	*	0.6	1.4	2.3	2.7	3.0	3.9	6.7
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	2.5	*	*	*	*	1.0	2.7	3.8	4.5	5.0	5.7	7.0
2001	2.7	*	*	*	*	1.0	2.8	4.2	4.7	4.9	6.3	8.6
2000	2.7	*	*	*	*	1.1	2.5	4.2	4.8	5.3	6.8	8.4
1999	2.8	*	*	*	*	1.1	2.8	4.5	5.2	5.5	6.5	9.3
Non-Hispanic black, both sexes												
2002	2.3	*	*	*	*	0.8	2.6	4.2	5.1	9.2	10.7	17.5
2001	2.2	*	*	*	*	0.8	2.5	3.8	5.3	7.9	12.0	19.0
2000	2.5	*	*	*	*	1.0	2.6	4.3	6.4	9.8	12.0	19.7
1999	2.4	*	*	*	*	0.7	2.5	3.6	7.2	10.8	11.3	21.7
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	4.3	*	*	*	*	1.6	4.9	7.8	9.2	15.8	16.7	24.2
2001	4.2	*	*	*	*	1.6	4.6	7.1	9.6	13.5	18.7	26.1
2000	4.7	*	*	*	*	1.9	5.0	8.0	11.5	16.6	18.8	27.0
1999	4.5	*	*	*	*	1.2	4.8	6.6	13.0	18.3	17.7	29.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of corpus uteri and uterus, part unspecified (C54-C55)												
All origins 3/, both sexes												
2002	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7
Hispanic, both sexes												
2002	0.9	*	*	*	*	*	0.3	1.1	3.2	9.0	11.0	20.5
2001	0.9	*	*	*	*	*	*	1.4	3.8	7.5	11.7	18.8
2000	0.8	*	*	*	*	*	*	1.1	3.6	7.0	10.7	19.9
1999	0.8	*	*	*	*	*	*	1.1	3.7	8.3	10.6	15.4
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	1.8	*	*	*	*	*	0.7	2.1	6.2	16.1	18.5	30.7
2001	1.8	*	*	*	*	*	*	2.8	7.3	13.4	19.6	27.9
2000	1.6	*	*	*	*	*	*	2.2	6.8	12.4	17.8	29.7
1999	1.6	*	*	*	*	*	*	2.2	7.0	14.8	17.7	22.9
Non-Hispanic 4/, both sexes												
2002	2.6	*	*	*	*	*	0.3	1.4	4.6	10.1	15.5	24.6
2001	2.6	*	*	*	*	0.1	0.3	1.4	4.4	9.9	16.6	24.8
2000	2.6	*	*	*	*	0.1	0.3	1.4	4.7	9.8	16.0	24.9
1999	2.5	*	*	*	*	0.1	0.3	1.3	4.3	9.9	16.6	24.3
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic 4/, female												
2002	5.1	*	*	*	*	*	0.6	2.8	8.8	18.5	25.8	35.3
2001	5.1	*	*	*	*	0.2	0.6	2.8	8.5	18.1	27.5	35.1
2000	5.0	*	*	*	*	0.1	0.6	2.7	9.0	17.9	26.5	34.9
1999	4.9	*	*	*	*	0.1	0.6	2.5	8.3	18.0	27.4	34.1
Non-Hispanic white, both sexes												
2002	2.6	*	*	*	*	*	0.3	1.3	4.1	8.9	14.6	23.8
2001	2.7	*	*	*	*	*	0.3	1.3	4.0	8.9	15.9	24.5
2000	2.6	*	*	*	*	*	0.3	1.3	4.3	8.6	15.0	24.5
1999	2.6	*	*	*	*	*	0.3	1.3	4.0	8.9	16.0	23.6
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	5.1	*	*	*	*	*	0.6	2.7	8.0	16.5	24.4	34.1
2001	5.2	*	*	*	*	*	0.6	2.7	7.8	16.4	26.5	34.7
2000	5.1	*	*	*	*	*	0.5	2.7	8.4	15.9	24.8	34.3
1999	5.1	*	*	*	*	*	0.6	2.5	7.7	16.3	26.4	33.1
Non-Hispanic black, both sexes												
2002	3.1	*	*	*	*	*	0.4	2.0	8.6	22.7	28.4	39.6
2001	2.9	*	*	*	*	*	0.4	2.1	7.7	21.6	28.6	32.8
2000	3.0	*	*	*	*	*	0.4	1.7	8.4	22.9	31.1	33.7
1999	2.8	*	*	*	*	*	0.5	1.4	8.1	21.5	27.5	35.9
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	5.8	*	*	*	*	*	0.7	3.6	15.5	38.7	44.4	54.8
2001	5.5	*	*	*	*	*	0.7	4.0	13.9	36.8	44.6	44.9
2000	5.7	*	*	*	*	*	0.7	3.1	15.3	39.1	48.6	46.2
1999	5.4	*	*	*	*	*	0.9	2.7	14.7	36.7	42.9	49.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of ovary (C56)												
All origins 3/, both sexes												
2002	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001	5.1	*	*	*	0.1	0.3	1.0	4.2	10.3	20.0	33.2	39.6
2000	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001	9.9	*	*	*	0.1	0.5	2.1	8.2	19.9	36.6	55.0	56.1
2000	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8
Hispanic, both sexes												
2002	1.6	*	*	*	*	*	0.8	3.2	5.6	12.3	24.2	23.8
2001	1.6	*	*	*	*	*	0.8	2.7	7.1	14.4	20.0	30.6
2000	1.6	*	*	*	*	*	0.6	3.0	6.5	14.3	24.9	32.5
1999	1.4	*	*	*	*	*	0.6	2.7	6.1	12.2	20.8	23.9
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	3.3	*	*	*	*	*	1.7	6.3	10.7	22.1	40.8	35.6
2001	3.4	*	*	*	*	*	1.6	5.4	13.4	25.8	33.5	45.6
2000	3.4	*	*	*	*	*	1.2	5.9	12.2	25.5	41.6	48.5
1999	2.9	*	*	*	*	*	1.2	5.3	11.5	21.7	34.7	35.4
Non-Hispanic 4/, both sexes												
2002	5.6	*	*	*	0.1	0.3	1.1	4.4	10.7	20.2	33.8	38.7
2001	5.6	*	*	*	0.1	0.3	1.1	4.3	10.5	20.3	33.7	39.8
2000	5.5	*	*	*	*	0.3	1.1	4.3	10.3	19.9	33.4	41.2
1999	5.3	*	*	*	0.1	0.2	1.2	4.4	10.5	20.0	32.0	38.1
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of ovary (C56)												
Non-Hispanic 4/, female												
2002	11.0	*	*	*	0.2	0.5	2.1	8.6	20.6	37.0	56.3	55.3
2001	10.9	*	*	*	0.1	0.5	2.1	8.4	20.3	37.3	55.8	56.4
2000	10.7	*	*	*	*	0.5	2.2	8.4	19.9	36.3	55.2	57.8
1999	10.4	*	*	*	0.2	0.4	2.3	8.6	20.1	36.4	52.7	53.3
Non-Hispanic white, both sexes												
2002	6.3	*	*	*	0.1	0.3	1.1	4.6	11.1	20.9	34.8	39.6
2001	6.2	*	*	*	*	0.3	1.1	4.5	10.8	21.0	34.8	40.6
2000	6.2	*	*	*	*	0.3	1.1	4.5	10.8	20.5	34.5	42.2
1999	6.0	*	*	*	0.1	0.2	1.2	4.5	10.9	20.5	32.7	39.2
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	12.4	*	*	*	0.2	0.5	2.2	9.1	21.7	38.7	58.0	56.7
2001	12.2	*	*	*	*	0.5	2.2	9.0	21.0	38.8	57.8	57.5
2000	12.0	*	*	*	*	0.6	2.3	8.9	21.0	37.7	57.2	59.2
1999	11.7	*	*	*	0.2	0.4	2.4	9.0	21.2	37.7	54.1	54.9
Non-Hispanic black, both sexes												
2002	3.2	*	*	*	*	*	1.1	3.5	9.2	17.4	29.3	35.4
2001	3.2	*	*	*	*	*	1.2	3.1	10.2	18.5	27.2	36.5
2000	3.0	*	*	*	*	*	1.1	3.4	8.6	18.2	27.1	34.7
1999	3.1	*	*	*	*	*	0.9	3.6	9.1	18.7	28.7	29.1
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	6.1	*	*	*	*	*	2.0	6.5	16.6	29.8	45.9	48.9
2001	6.1	*	*	*	*	*	2.2	5.7	18.4	31.5	42.5	50.0
2000	5.8	*	*	*	*	*	2.0	6.3	15.6	31.1	42.4	47.5
1999	5.8	*	*	*	*	*	1.6	6.7	16.4	31.9	44.8	40.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified female genital organs (C57)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.3	0.6	1.0	1.0
2001	0.1	*	*	*	*	*	*	0.1	0.3	0.5	0.9	1.2
2000	0.1	*	*	*	*	*	*	0.1	0.3	0.5	0.9	1.6
1999	0.1	*	*	*	*	*	*	0.1	0.3	0.6	1.0	0.9
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.1	1.7	1.5
2001	0.3	*	*	*	*	*	*	0.2	0.5	1.0	1.5	1.7
2000	0.3	*	*	*	*	*	*	0.2	0.6	1.0	1.5	2.2
1999	0.3	*	*	*	*	*	*	0.3	0.6	1.1	1.7	1.3
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.0	1.0
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.6	0.9	1.3
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.5	0.9	1.6
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.0	0.9
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified female genital organs (C57)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	*	0.2	0.6	1.1	1.7	1.5
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.0	1.5	1.8
2000	0.3	*	*	*	*	*	*	0.2	0.6	0.9	1.5	2.2
1999	0.3	*	*	*	*	*	*	0.3	0.7	1.1	1.7	1.2
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.0	1.0
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.6	0.9	1.3
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.5	0.9	1.6
1999	0.2	*	*	*	*	*	*	0.1	0.4	0.6	1.0	0.9
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.1	1.7	1.5
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.1	1.6	1.9
2000	0.3	*	*	*	*	*	*	0.2	0.6	1.0	1.4	2.2
1999	0.3	*	*	*	*	*	*	0.3	0.7	1.1	1.7	1.2
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of placenta (C58)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of placenta (C58)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of male genital organs (C60-C63)												
All origins 3/, both sexes												
2002	10.8	*	*	*	0.1	0.2	0.3	1.2	7.9	35.0	100.8	195.9
2001	11.0	*	*	*	0.1	0.2	0.3	1.3	7.8	36.2	103.8	201.9
2000	11.3	*	*	*	0.1	0.2	0.3	1.2	8.2	37.4	106.5	210.1
1999	11.6	*	*	*	0.1	0.3	0.4	1.2	8.5	40.2	110.6	208.7
All origins 3/, male												
2002	21.9	*	*	*	0.2	0.5	0.6	2.4	16.4	77.0	252.6	647.3
2001	22.4	*	*	*	0.2	0.4	0.6	2.7	16.2	79.8	261.7	684.3
2000	22.9	*	*	*	0.2	0.4	0.6	2.4	17.1	82.9	269.8	726.1
1999	23.6	*	*	*	0.2	0.5	0.7	2.5	17.6	89.2	280.8	726.5
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	3.3	*	*	*	*	0.3	*	1.1	5.9	25.4	79.6	172.4
2001	3.5	*	*	*	*	*	*	1.1	6.3	29.8	88.5	171.3
2000	3.1	*	*	*	*	0.4	*	1.0	5.1	26.1	80.4	166.5
1999	3.3	*	*	*	*	*	*	0.9	5.9	29.0	81.5	179.6
Hispanic, male												
2002	6.5	*	*	*	*	0.5	*	2.2	12.4	57.3	195.7	520.4
2001	6.8	*	*	*	*	*	*	2.3	13.4	67.7	219.6	521.9
2000	6.1	*	*	*	*	0.7	*	2.0	11.0	59.3	200.1	505.9
1999	6.4	*	*	*	*	*	*	1.8	12.6	65.9	203.9	552.1
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	11.9	*	*	*	0.1	0.2	0.3	1.2	8.0	35.5	101.6	196.3
2001	12.1	*	*	*	0.1	0.2	0.3	1.3	7.9	36.5	104.2	202.5
2000	12.4	*	*	*	0.1	0.2	0.3	1.2	8.4	38.0	107.3	211.3
1999	12.7	*	*	*	0.1	0.3	0.4	1.3	8.6	40.7	111.6	209.3
Non-Hispanic 4/, male												
2002	24.4	*	*	*	0.2	0.5	0.7	2.4	16.7	78.0	254.8	651.4
2001	24.8	*	*	*	0.1	0.4	0.6	2.7	16.4	80.4	262.9	689.4
2000	25.4	*	*	*	0.2	0.4	0.7	2.4	17.6	84.1	272.1	733.7
1999	26.1	*	*	*	0.1	0.5	0.7	2.6	17.9	90.3	283.5	732.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of male genital organs (C60-C63)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	12.2	*	*	*	0.1	0.3	0.4	1.0	6.8	31.4	93.8	187.4
2001	12.2	*	*	*	*	0.2	0.3	1.1	6.7	32.3	95.6	190.4
2000	12.5	*	*	*	0.1	0.2	0.3	1.0	7.2	33.2	98.3	200.1
1999	12.9	*	*	*	*	0.3	0.4	1.1	7.2	36.1	103.3	196.5
Non-Hispanic white, male												
2002	24.8	*	*	*	0.2	0.5	0.7	2.0	14.0	68.2	233.7	621.4
2001	25.0	*	*	*	*	0.5	0.7	2.2	13.7	70.4	239.8	648.3
2000	25.6	*	*	*	0.2	0.4	0.7	2.1	14.8	72.7	247.9	696.6
1999	26.4	*	*	*	*	0.6	0.8	2.2	14.8	79.2	261.1	689.2
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	14.2	*	*	*	*	*	*	2.9	19.8	82.6	213.0	333.6
2001	14.8	*	*	*	*	*	*	3.2	19.8	84.8	222.6	366.1
2000	15.2	*	*	*	*	*	0.4	2.6	20.9	91.8	232.0	360.9
1999	15.5	*	*	*	*	*	*	2.8	22.5	95.2	228.1	369.9
Non-Hispanic black, male												
2002	29.9	*	*	*	*	*	*	6.3	44.5	199.1	589.4	1,206.6
2001	31.1	*	*	*	*	*	*	6.8	44.5	205.1	619.0	1,354.8
2000	32.0	*	*	*	*	*	0.8	5.7	46.7	222.2	644.2	1,332.7
1999	32.5	*	*	*	*	*	*	6.0	50.5	230.1	635.4	1,359.6
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of prostate (C61)												
All origins 3/, both sexes												
2002	10.6	*	*	*	*	*	0.1	1.0	7.6	34.5	100.1	195.0
2001	10.8	*	*	*	*	*	0.1	1.1	7.5	35.7	103.2	201.0
2000	11.0	*	*	*	*	*	0.1	1.0	7.9	37.0	105.7	209.3
1999	11.4	*	*	*	*	*	0.1	1.0	8.2	39.7	109.9	207.5
All origins 3/, male												
2002	21.5	*	*	*	*	*	0.1	1.9	15.8	75.9	251.0	644.5
2001	22.0	*	*	*	*	*	0.1	2.1	15.7	78.9	260.2	681.2
2000	22.5	*	*	*	*	*	0.1	2.0	16.6	82.0	267.9	723.1
1999	23.2	*	*	*	*	*	0.1	2.1	17.1	88.2	279.1	722.4
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	3.1	*	*	*	*	*	*	0.8	5.4	24.7	78.7	170.7
2001	3.3	*	*	*	*	*	*	0.8	5.8	29.2	87.8	170.7
2000	2.9	*	*	*	*	*	*	0.8	4.6	25.7	79.2	165.9
1999	3.1	*	*	*	*	*	*	0.7	5.6	28.5	80.4	176.1
Hispanic, male												
2002	6.0	*	*	*	*	*	*	1.6	11.3	55.8	193.7	515.4
2001	6.5	*	*	*	*	*	*	1.6	12.4	66.3	217.8	520.0
2000	5.7	*	*	*	*	*	*	1.7	9.7	58.5	197.2	503.9
1999	6.0	*	*	*	*	*	*	1.5	12.0	64.8	201.3	541.3
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	11.7	*	*	*	*	*	0.1	1.0	7.7	35.0	101.0	195.5
2001	11.9	*	*	*	*	*	0.1	1.1	7.6	36.1	103.6	201.6
2000	12.2	*	*	*	*	*	0.1	1.0	8.2	37.6	106.6	210.4
1999	12.5	*	*	*	*	*	0.1	1.0	8.3	40.3	110.9	208.2
Non-Hispanic 4/, male												
2002	24.0	*	*	*	*	*	0.2	2.0	16.1	76.9	253.3	648.7
2001	24.3	*	*	*	*	*	0.1	2.2	15.9	79.5	261.4	686.3
2000	25.0	*	*	*	*	*	0.1	2.0	17.0	83.2	270.2	730.7
1999	25.6	*	*	*	*	*	0.2	2.1	17.4	89.4	281.8	728.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of prostate (C61)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	11.9	*	*	*	*	*	0.1	0.8	6.5	30.9	93.1	186.6
2001	12.0	*	*	*	*	*	*	0.9	6.4	31.9	95.1	189.5
2000	12.3	*	*	*	*	*	*	0.8	6.9	32.8	97.6	199.3
1999	12.6	*	*	*	*	*	0.1	0.9	6.9	35.6	102.6	195.4
Non-Hispanic white, male												
2002	24.3	*	*	*	*	*	0.2	1.5	13.4	67.1	232.1	618.5
2001	24.6	*	*	*	*	*	*	1.8	13.3	69.6	238.4	645.0
2000	25.2	*	*	*	*	*	*	1.7	14.2	71.7	246.1	693.6
1999	25.9	*	*	*	*	*	0.1	1.8	14.3	78.2	259.4	685.4
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	14.1	*	*	*	*	*	*	2.6	19.5	82.1	212.3	333.3
2001	14.6	*	*	*	*	*	*	2.9	19.5	84.2	221.6	365.8
2000	15.1	*	*	*	*	*	*	2.4	20.8	91.6	230.8	359.6
1999	15.3	*	*	*	*	*	*	2.7	22.0	94.7	227.5	368.6
Non-Hispanic black, male												
2002	29.6	*	*	*	*	*	*	5.7	43.8	197.9	587.6	1,205.5
2001	30.8	*	*	*	*	*	*	6.2	43.7	203.5	616.3	1,353.6
2000	31.7	*	*	*	*	*	*	5.3	46.6	221.7	640.8	1,328.0
1999	32.2	*	*	*	*	*	*	5.8	49.5	229.1	633.8	1,354.9
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of testis (C62)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	0.1	*	*
2001	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	0.1	*	*
2000	0.1	*	*	*	0.1	0.2	0.2	0.1	0.1	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.3	0.2	0.1	0.1	*	*
All origins 3/, male												
2002	0.3	*	*	*	0.2	0.5	0.4	0.3	0.3	0.3	*	*
2001	0.2	*	*	*	0.2	0.4	0.4	0.3	0.2	0.3	*	*
2000	0.2	*	*	*	0.2	0.4	0.5	0.3	0.2	*	*	*
1999	0.3	*	*	*	0.2	0.5	0.5	0.3	0.2	0.3	*	*
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	0.4	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	0.7	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	0.1	*	*
2001	0.1	*	*	*	0.1	0.2	0.2	0.2	0.1	0.1	*	*
2000	0.1	*	*	*	0.1	0.2	0.2	0.1	0.1	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.3	0.2	0.1	*	*	*
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.2	0.5	0.5	0.3	0.3	0.3	*	*
2001	0.2	*	*	*	0.1	0.4	0.4	0.3	0.2	0.3	*	*
2000	0.2	*	*	*	0.2	0.4	0.5	0.3	0.2	*	*	*
1999	0.3	*	*	*	0.1	0.5	0.5	0.4	0.2	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of testis (C62)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.1	0.3	0.3	0.2	0.2	0.1	*	*
2001	0.1	*	*	*	*	0.2	0.2	0.2	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.3	0.1	0.1	*	*	*
1999	0.2	*	*	*	*	0.3	0.3	0.2	0.1	*	*	*
Non-Hispanic white, male												
2002	0.3	*	*	*	0.2	0.5	0.5	0.4	0.3	0.3	*	*
2001	0.3	*	*	*	*	0.5	0.5	0.3	*	*	*	*
2000	0.3	*	*	*	0.2	0.4	0.5	0.3	0.2	*	*	*
1999	0.3	*	*	*	*	0.6	0.6	0.4	0.3	*	*	*
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of penis and other and unspecified male genital organs (C60,C63)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.5	0.6
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.7	0.7
1999	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	1.1
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.4	0.8	1.3	2.0
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.8
2000	0.2	*	*	*	*	*	*	0.1	0.4	0.7	1.7	2.4
1999	0.2	*	*	*	*	*	*	*	0.3	0.7	1.4	3.7
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.6
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	0.7
1999	0.1	*	*	*	*	*	*	*	0.2	0.3	0.5	1.0
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.3	1.9
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.8
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.6	2.4
1999	0.2	*	*	*	*	*	*	*	0.3	0.7	1.4	3.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of penis and other and unspecified male genital organs (C60,C63)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.3	0.5	0.6
2001	0.1	*	*	*	*	*	*	*	0.1	0.3	0.4	0.9
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	0.6
1999	0.1	*	*	*	*	*	*	*	0.1	0.3	0.5	0.9
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	0.3	0.7	1.3	2.0
2001	0.2	*	*	*	*	*	*	*	0.3	0.6	1.1	2.9
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.5	2.3
1999	0.2	*	*	*	*	*	*	*	0.3	0.7	1.4	3.3
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)												
All origins 3/, both sexes												
2002	8.8	*	*	0.1	0.1	0.2	1.0	4.6	14.0	34.6	65.1	102.0
2001	8.7	*	0.1	0.1	0.1	0.2	1.1	4.6	14.4	33.8	63.8	103.9
2000	8.7	*	*	0.1	0.1	0.1	1.0	4.5	14.3	33.5	64.0	106.0
1999	8.5	*	*	0.1	*	0.2	1.0	4.5	14.4	33.5	61.8	103.0
All origins 3/, male												
2002	11.6	*	*	0.1	*	0.2	1.3	6.7	21.1	51.4	103.3	183.6
2001	11.6	*	*	*	*	0.2	1.4	6.8	21.6	51.5	102.5	192.4
2000	11.4	*	*	*	*	0.1	1.3	6.6	21.3	50.7	103.3	198.9
1999	11.2	*	*	*	*	0.2	1.4	6.6	21.5	50.6	100.8	188.4
All origins 3/, female												
2002	6.1	*	*	*	*	0.1	0.6	2.6	7.5	20.5	39.7	66.6
2001	6.0	*	*	0.1	*	0.2	0.8	2.6	7.9	19.2	38.3	66.9
2000	6.0	*	*	*	*	0.1	0.7	2.6	7.9	19.4	38.5	68.1
1999	5.9	*	*	*	*	0.1	0.7	2.5	7.9	19.5	36.5	68.6
Hispanic, both sexes												
2002	2.7	*	*	*	*	*	0.7	3.5	11.0	20.9	43.9	82.0
2001	2.7	*	*	*	*	*	0.8	3.9	9.3	22.5	43.1	81.9
2000	2.5	*	*	*	*	*	0.8	3.4	10.9	19.1	43.5	73.7
1999	2.5	*	*	*	*	*	0.7	3.6	11.4	22.6	39.8	68.1
Hispanic, male												
2002	3.4	*	*	*	*	*	0.9	4.9	15.5	31.9	68.5	142.2
2001	3.5	*	*	*	*	*	1.2	5.5	14.4	33.3	67.5	150.5
2000	3.3	*	*	*	*	*	1.2	4.6	16.2	28.9	73.8	125.0
1999	3.3	*	*	*	*	*	1.1	5.4	17.8	36.7	58.4	118.6
Hispanic, female												
2002	2.0	*	*	*	*	*	*	2.1	6.9	12.2	27.0	52.2
2001	1.9	*	*	*	*	*	*	2.5	4.7	13.9	26.6	48.4
2000	1.7	*	*	*	*	*	*	2.2	6.3	11.4	23.1	48.5
1999	1.7	*	*	*	*	*	*	1.9	5.6	11.5	27.4	43.7
Non-Hispanic 4/, both sexes												
2002	9.7	*	*	0.1	0.1	0.2	1.0	4.7	14.2	35.4	66.0	102.6
2001	9.6	*	*	0.1	0.1	0.2	1.1	4.7	14.8	34.5	64.6	104.5
2000	9.5	*	*	0.1	0.1	0.1	1.0	4.6	14.5	34.3	64.7	106.8
1999	9.3	*	*	0.1	*	0.2	1.0	4.5	14.6	34.1	62.6	104.0
Non-Hispanic 4/, male												
2002	12.9	*	*	*	*	0.2	1.4	6.9	21.4	52.6	104.9	185.0
2001	12.8	*	*	*	*	0.2	1.4	6.9	22.1	52.5	103.9	193.6
2000	12.6	*	*	*	*	0.1	1.3	6.7	21.6	51.8	104.2	201.3
1999	12.3	*	*	*	*	0.3	1.4	6.6	21.7	51.3	102.3	190.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)												
Non-Hispanic 4/, female												
2002	6.7	*	*	*	*	0.1	0.6	2.7	7.5	21.1	40.2	67.1
2001	6.6	*	*	0.2	*	0.2	0.9	2.5	8.1	19.5	38.8	67.4
2000	6.6	*	*	*	*	0.1	0.7	2.6	8.0	19.9	39.0	68.6
1999	6.4	*	*	*	*	0.1	0.7	2.5	8.1	19.9	36.8	69.2
Non-Hispanic white, both sexes												
2002	11.0	*	*	0.1	*	0.1	1.0	4.8	14.7	36.8	67.7	104.5
2001	10.8	*	*	*	*	0.1	1.1	4.8	15.2	36.0	66.1	106.9
2000	10.7	*	*	*	*	0.1	1.0	4.8	15.1	35.5	66.1	110.2
1999	10.4	*	*	0.1	*	0.1	1.0	4.5	15.0	35.0	64.0	105.6
Non-Hispanic white, male												
2002	14.7	*	*	*	*	0.2	1.4	7.1	22.3	54.9	108.1	192.4
2001	14.5	*	*	*	*	*	1.4	7.1	22.7	55.0	106.9	202.1
2000	14.3	*	*	*	*	*	1.4	7.0	22.4	53.5	107.0	209.7
1999	13.8	*	*	*	*	0.2	1.3	6.6	22.4	53.0	105.5	197.6
Non-Hispanic white, female												
2002	7.4	*	*	*	*	*	0.6	2.6	7.5	21.4	40.6	66.5
2001	7.2	*	*	*	*	*	0.8	2.5	8.1	19.8	38.9	67.4
2000	7.3	*	*	*	*	*	0.7	2.7	8.2	20.2	39.1	70.1
1999	7.0	*	*	*	*	*	0.7	2.6	8.1	20.0	36.8	68.9
Non-Hispanic black, both sexes												
2002	5.7	*	*	*	*	0.4	1.0	5.0	14.1	31.0	59.0	94.1
2001	5.7	*	*	*	*	0.4	1.7	5.1	15.3	28.3	59.1	87.4
2000	5.5	*	*	*	*	*	1.2	4.6	13.4	31.2	61.9	80.1
1999	5.7	*	*	*	*	0.5	1.5	5.6	14.8	32.3	54.3	94.1
Non-Hispanic black, male												
2002	6.5	*	*	*	*	*	1.5	6.4	20.3	41.0	86.2	135.6
2001	6.7	*	*	*	*	*	2.0	7.4	22.2	40.0	86.4	127.0
2000	6.6	*	*	*	*	*	1.2	6.4	19.5	45.5	93.2	139.7
1999	6.8	*	*	*	*	*	2.1	8.6	21.1	43.6	80.0	139.1
Non-Hispanic black, female												
2002	4.9	*	*	*	*	*	0.7	3.8	9.1	24.0	43.6	78.2
2001	4.8	*	*	*	*	*	1.5	3.2	9.8	20.0	43.8	72.7
2000	4.4	*	*	*	*	*	1.1	3.1	8.5	21.1	44.3	58.0
1999	4.7	*	*	*	*	*	0.9	3.1	9.7	24.4	39.9	77.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)												
All origins 3/, both sexes												
2002	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4
All origins 3/, male												
2002	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4
All origins 3/, female												
2002	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1
Hispanic, both sexes												
2002	1.7	*	*	*	*	*	0.6	2.8	8.1	12.9	24.4	26.1
2001	1.7	*	*	*	*	*	0.7	3.3	6.9	14.4	23.3	31.9
2000	1.6	*	*	*	*	*	0.6	2.8	7.9	12.3	24.7	25.9
1999	1.6	*	*	*	*	*	0.6	2.8	7.9	13.9	22.5	28.8
Hispanic, male												
2002	2.1	*	*	*	*	*	0.9	3.9	11.3	18.5	39.0	40.2
2001	2.2	*	*	*	*	*	1.2	4.7	9.9	21.4	35.3	57.1
2000	2.0	*	*	*	*	*	1.0	3.9	11.4	17.9	38.4	40.3
1999	2.0	*	*	*	*	*	0.9	4.3	11.8	21.1	29.7	*
Hispanic, female												
2002	1.2	*	*	*	*	*	*	1.7	5.2	8.4	14.3	19.1
2001	1.2	*	*	*	*	*	*	1.9	4.2	8.8	15.1	19.5
2000	1.1	*	*	*	*	*	*	1.8	4.7	7.8	15.5	*
1999	1.2	*	*	*	*	*	*	1.4	4.4	8.3	17.7	23.9
Non-Hispanic 4/, both sexes												
2002	4.6	*	*	0.1	*	0.1	0.7	3.3	8.8	18.0	26.4	32.7
2001	4.6	*	*	0.1	0.1	0.1	0.8	3.3	9.4	17.8	26.4	32.3
2000	4.5	*	*	0.1	*	0.1	0.8	3.3	9.1	17.3	26.6	33.9
1999	4.3	*	*	0.1	*	0.1	0.8	3.2	8.9	16.8	24.6	31.4
Non-Hispanic 4/, male												
2002	5.9	*	*	*	*	0.2	1.0	4.9	13.0	25.5	37.7	50.4
2001	5.9	*	*	*	*	0.1	1.0	4.9	13.7	25.4	37.5	52.6
2000	5.7	*	*	*	*	0.1	1.0	4.8	13.2	24.4	38.1	55.4
1999	5.4	*	*	*	*	0.2	1.1	4.6	13.0	23.7	35.9	49.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)												
Non-Hispanic 4/, female												
2002	3.4	*	*	*	*	*	0.4	1.8	4.9	11.8	18.8	25.1
2001	3.4	*	*	0.1	*	0.1	0.6	1.8	5.4	11.4	19.1	23.8
2000	3.4	*	*	*	*	*	0.5	1.9	5.3	11.4	19.1	25.2
1999	3.2	*	*	*	*	*	0.5	1.8	5.2	11.2	17.3	24.1
Non-Hispanic white, both sexes												
2002	5.1	*	*	0.1	*	0.1	0.7	3.4	9.1	18.7	26.8	33.5
2001	5.1	*	*	*	*	0.1	0.7	3.4	9.6	18.4	26.8	32.8
2000	5.0	*	*	*	*	0.1	0.7	3.4	9.3	17.7	26.9	34.6
1999	4.7	*	*	0.1	*	0.1	0.7	3.1	9.2	17.2	24.9	31.5
Non-Hispanic white, male												
2002	6.5	*	*	*	*	*	1.0	5.0	13.3	26.3	38.4	52.4
2001	6.5	*	*	*	*	*	1.0	4.9	13.8	26.2	37.7	54.3
2000	6.3	*	*	*	*	*	1.0	4.9	13.5	24.6	38.5	56.9
1999	6.0	*	*	*	*	*	1.0	4.5	13.3	24.0	36.5	50.2
Non-Hispanic white, female												
2002	3.8	*	*	*	*	*	0.4	1.8	5.1	12.2	19.0	25.3
2001	3.8	*	*	*	*	*	0.5	1.8	5.5	11.8	19.5	23.8
2000	3.8	*	*	*	*	*	0.5	1.9	5.4	11.8	19.3	25.7
1999	3.5	*	*	*	*	*	0.5	1.8	5.3	11.5	17.3	24.1
Non-Hispanic black, both sexes												
2002	2.9	*	*	*	*	*	0.8	3.6	8.8	15.8	26.5	28.1
2001	3.1	*	*	*	*	*	1.3	3.8	9.7	15.3	26.0	30.0
2000	2.9	*	*	*	*	*	0.9	3.5	9.0	16.8	25.9	29.6
1999	3.0	*	*	*	*	0.4	1.1	3.9	8.7	16.4	23.9	31.7
Non-Hispanic black, male												
2002	3.6	*	*	*	*	*	1.2	5.0	14.4	22.1	36.6	39.4
2001	4.0	*	*	*	*	*	1.5	5.5	15.3	22.7	40.8	46.2
2000	3.7	*	*	*	*	*	1.0	5.0	13.4	26.8	38.3	49.3
1999	3.7	*	*	*	*	*	1.6	6.0	13.3	23.9	32.5	49.9
Non-Hispanic black, female												
2002	2.3	*	*	*	*	*	*	2.5	4.4	11.3	20.7	23.8
2001	2.3	*	*	*	*	*	1.1	2.2	5.3	10.0	17.7	24.0
2000	2.2	*	*	*	*	*	0.8	2.2	5.4	9.7	18.9	22.2
1999	2.2	*	*	*	*	*	*	2.1	5.0	11.0	19.1	24.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of bladder (C67)												
All origins 3/, both sexes												
2002	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5
All origins 3/, male												
2002	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8
All origins 3/, female												
2002	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8
Hispanic, both sexes												
2002	1.0	*	*	*	*	*	*	0.7	2.9	7.7	18.3	54.9
2001	0.9	*	*	*	*	*	*	0.6	2.3	7.9	18.8	46.3
2000	0.9	*	*	*	*	*	*	*	2.9	6.6	17.6	47.1
1999	0.9	*	*	*	*	*	*	0.7	3.1	8.5	16.5	38.6
Hispanic, male												
2002	1.3	*	*	*	*	*	*	*	4.2	12.8	27.9	100.4
2001	1.2	*	*	*	*	*	*	*	4.4	11.8	30.0	87.6
2000	1.2	*	*	*	*	*	*	*	4.6	10.6	33.9	84.6
1999	1.3	*	*	*	*	*	*	*	5.6	15.2	27.1	77.6
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	*	3.6	11.8	32.3
2001	0.6	*	*	*	*	*	*	*	*	4.9	11.2	26.1
2000	0.5	*	*	*	*	*	*	*	*	3.5	6.6	28.7
1999	0.5	*	*	*	*	*	*	*	*	*	9.4	*
Non-Hispanic 4/, both sexes												
2002	4.9	*	*	*	*	*	0.3	1.3	5.1	16.4	37.8	66.9
2001	4.8	*	*	*	*	*	0.3	1.3	5.2	15.8	36.4	69.6
2000	4.7	*	*	*	*	*	0.3	1.2	5.2	16.1	36.3	70.3
1999	4.7	*	*	*	*	*	0.2	1.3	5.4	16.3	36.0	69.3
Non-Hispanic 4/, male												
2002	6.8	*	*	*	*	*	0.4	1.9	7.9	26.0	64.7	129.8
2001	6.7	*	*	*	*	*	0.4	1.9	8.1	26.0	63.8	136.4
2000	6.6	*	*	*	*	*	0.3	1.8	8.1	26.1	63.6	141.3
1999	6.6	*	*	*	*	*	0.3	2.0	8.4	26.3	63.8	136.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of bladder (C67)												
Non-Hispanic 4/, female												
2002	3.1	*	*	*	*	*	0.2	0.8	2.4	8.5	20.0	39.8
2001	3.0	*	*	*	*	*	0.3	0.6	2.5	7.4	18.3	41.9
2000	3.0	*	*	*	*	*	0.2	0.6	2.5	7.8	18.5	41.6
1999	3.0	*	*	*	*	*	0.2	0.6	2.6	8.1	17.9	42.3
Non-Hispanic white, both sexes												
2002	5.6	*	*	*	*	*	0.3	1.4	5.3	17.1	39.1	68.0
2001	5.5	*	*	*	*	*	0.3	1.3	5.4	16.6	37.4	71.5
2000	5.4	*	*	*	*	*	0.3	1.3	5.5	16.8	37.1	72.8
1999	5.4	*	*	*	*	*	0.2	1.3	5.5	16.8	37.1	70.7
Non-Hispanic white, male												
2002	7.9	*	*	*	*	*	0.4	2.0	8.5	27.4	67.2	135.2
2001	7.8	*	*	*	*	*	0.4	2.0	8.6	27.6	66.6	143.0
2000	7.7	*	*	*	*	*	0.3	2.0	8.6	27.5	65.9	148.0
1999	7.6	*	*	*	*	*	0.3	2.0	8.7	27.6	66.2	143.0
Non-Hispanic white, female												
2002	3.4	*	*	*	*	*	0.2	0.8	2.3	8.4	20.2	39.0
2001	3.2	*	*	*	*	*	0.3	0.6	2.4	7.4	18.1	41.7
2000	3.3	*	*	*	*	*	0.2	0.7	2.6	7.8	18.2	42.5
1999	3.2	*	*	*	*	*	0.2	0.7	2.5	7.8	18.0	41.9
Non-Hispanic black, both sexes												
2002	2.6	*	*	*	*	*	*	1.1	4.5	14.7	31.2	64.1
2001	2.4	*	*	*	*	*	0.4	1.3	5.1	12.1	31.4	55.6
2000	2.4	*	*	*	*	*	*	1.1	4.2	13.5	34.9	49.6
1999	2.6	*	*	*	*	*	*	1.5	5.6	15.2	29.4	60.8
Non-Hispanic black, male												
2002	2.8	*	*	*	*	*	*	1.3	5.5	18.7	48.1	91.9
2001	2.6	*	*	*	*	*	*	1.8	6.5	16.5	43.2	78.5
2000	2.8	*	*	*	*	*	*	1.4	5.9	17.8	52.8	88.1
1999	3.0	*	*	*	*	*	*	2.6	7.4	19.2	46.6	86.8
Non-Hispanic black, female												
2002	2.4	*	*	*	*	*	*	0.9	3.7	11.8	21.7	53.5
2001	2.3	*	*	*	*	*	*	*	3.9	9.0	24.8	47.1
2000	2.1	*	*	*	*	*	*	*	2.7	10.4	24.8	35.3
1999	2.3	*	*	*	*	*	*	*	4.2	12.3	19.8	51.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified sites within the urinary tract (C66,C68)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.8	2.9
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.9	1.8	2.7
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.8	2.6
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.9	3.1
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	0.1	0.4	1.1	2.4	4.7
2001	0.2	*	*	*	*	*	*	0.1	0.3	1.1	2.6	4.7
2000	0.2	*	*	*	*	*	*	*	0.3	1.3	2.5	4.4
1999	0.2	*	*	*	*	*	*	*	0.4	1.3	2.6	4.2
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	*	0.2	0.8	1.3	2.2
2001	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	1.8
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.4	2.7
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	*	0.1	0.3	1.0	1.8	3.0
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.9	1.8	2.6
2000	0.2	*	*	*	*	*	*	0.1	0.3	1.0	1.9	2.6
1999	0.3	*	*	*	*	*	*	0.1	0.3	0.9	1.9	3.2
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	0.1	0.5	1.1	2.5	4.8
2001	0.3	*	*	*	*	*	*	*	0.3	1.2	2.6	4.7
2000	0.3	*	*	*	*	*	*	*	0.3	1.3	2.5	4.6
1999	0.3	*	*	*	*	*	*	*	0.3	1.3	2.6	4.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified sites within the urinary tract (C66,C68)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	0.2	0.8	1.3	2.2
2001	0.2	*	*	*	*	*	*	*	0.3	0.7	1.3	1.8
2000	0.2	*	*	*	*	*	*	*	0.2	0.7	1.4	1.9
1999	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.8
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	*	*	*	0.1	0.3	1.0	1.8	3.0
2001	0.3	*	*	*	*	*	*	0.1	0.2	0.9	1.9	2.7
2000	0.3	*	*	*	*	*	*	0.1	0.3	1.0	2.0	2.8
1999	0.3	*	*	*	*	*	*	*	0.3	1.0	2.0	3.4
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	*	0.5	1.2	2.6	4.8
2001	0.3	*	*	*	*	*	*	*	0.2	1.2	2.7	4.9
2000	0.3	*	*	*	*	*	*	*	0.3	1.4	2.6	4.8
1999	0.3	*	*	*	*	*	*	*	0.3	1.4	2.8	4.4
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	0.9	1.4	2.3
2001	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	*	*	0.2	0.6	1.6	2.0
1999	0.3	*	*	*	*	*	*	*	0.2	0.6	1.5	3.0
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of eye and adnexa (C69)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.5	0.7
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.6
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.8
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.4	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	0.2	0.2	0.4	0.7
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.5	0.7
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.6
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.8
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	0.3	*	0.6	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of eye and adnexa (C69)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	0.2	*	0.4	0.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	0.8
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.5	0.8
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.6	0.7
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8
1999	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	1.0
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	0.3	*	0.6	*
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.5	0.5	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	0.9
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All origins 3/, both sexes												
2002	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
All origins 3/, male												
2002	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
All origins 3/, female												
2002	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1
Hispanic, both sexes												
2002	1.7	*	0.7	0.7	0.4	0.5	0.9	3.4	5.8	10.0	13.7	13.3
2001	1.7	*	0.7	0.5	0.4	0.6	1.4	2.6	5.9	10.3	14.8	*
2000	1.8	*	*	0.9	0.4	0.7	1.5	2.9	6.2	9.8	14.2	15.9
1999	1.7	*	0.9	0.7	*	0.5	1.3	3.0	6.0	10.0	14.0	*
Hispanic, male												
2002	1.8	*	*	0.7	*	*	1.0	4.4	6.6	12.4	16.0	*
2001	1.8	*	*	*	*	0.6	1.4	3.0	6.8	14.4	17.9	*
2000	1.9	*	*	1.2	*	0.8	1.5	2.6	7.7	12.5	18.7	*
1999	1.7	*	*	0.7	*	*	1.8	2.9	7.8	10.5	16.2	*
Hispanic, female												
2002	1.5	*	*	0.7	*	*	0.9	2.3	5.1	8.1	12.1	*
2001	1.6	*	*	0.6	*	*	1.4	2.2	5.1	7.1	12.7	*
2000	1.7	*	*	0.7	*	*	1.5	3.2	4.9	7.6	11.2	*
1999	1.7	*	*	0.8	*	*	0.8	3.1	4.5	9.6	12.5	*
Non-Hispanic 4/, both sexes												
2002	4.9	*	0.7	0.8	0.5	1.1	2.4	5.3	9.9	16.6	19.4	16.5
2001	4.8	*	0.6	0.8	0.6	1.2	2.4	5.2	10.0	15.7	20.6	16.5
2000	4.9	*	0.7	0.7	0.6	1.1	2.5	5.3	9.8	16.9	20.2	16.9
1999	5.0	*	0.7	0.8	0.6	1.2	2.5	5.3	10.6	17.3	20.6	16.1
Non-Hispanic 4/, male												
2002	5.5	*	0.8	0.9	0.7	1.3	3.0	6.4	12.0	20.7	24.1	20.2
2001	5.5	*	0.6	0.7	0.7	1.5	3.0	6.4	12.5	19.4	26.1	20.5
2000	5.5	*	0.7	0.8	0.7	1.4	3.3	6.5	12.1	20.6	24.5	22.3
1999	5.6	*	0.7	0.8	0.7	1.5	3.0	6.5	12.7	21.4	25.8	21.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic 4/, female												
2002	4.2	*	0.7	0.7	0.4	0.9	1.8	4.2	7.9	13.1	16.2	14.9
2001	4.2	*	0.6	0.8	0.5	0.8	1.8	4.1	7.7	12.7	17.0	14.9
2000	4.2	*	0.7	0.7	0.5	0.7	1.8	4.1	7.8	13.8	17.5	14.7
1999	4.4	*	0.7	0.8	0.5	0.9	2.0	4.2	8.6	14.0	17.2	14.0
Non-Hispanic white, both sexes												
2002	5.6	*	0.6	0.8	0.6	1.2	2.8	5.9	10.7	17.9	20.4	17.3
2001	5.6	*	0.7	0.8	0.6	1.3	2.7	5.7	10.8	16.8	21.9	17.2
2000	5.6	*	0.7	0.8	0.6	1.2	2.8	5.9	10.7	18.3	21.2	17.5
1999	5.7	*	0.7	0.8	0.6	1.4	2.8	5.8	11.5	18.7	21.7	16.7
Non-Hispanic white, male												
2002	6.4	*	0.7	0.9	0.8	1.4	3.5	7.1	12.8	22.3	25.2	21.4
2001	6.3	*	0.7	0.7	0.7	1.7	3.5	7.0	13.4	20.9	27.5	20.9
2000	6.3	*	0.7	0.8	0.7	1.5	3.7	7.1	13.1	22.2	25.7	23.5
1999	6.4	*	0.8	0.8	0.7	1.8	3.3	7.1	13.7	23.0	27.2	22.4
Non-Hispanic white, female												
2002	4.9	*	0.6	0.8	0.4	1.1	2.1	4.7	8.7	14.2	17.1	15.5
2001	4.8	*	0.6	0.9	0.5	0.9	2.0	4.5	8.4	13.4	18.2	15.7
2000	4.9	*	0.7	0.7	0.5	0.9	1.9	4.6	8.4	15.0	18.3	15.1
1999	5.0	*	0.7	0.7	0.5	1.1	2.3	4.6	9.5	15.1	18.1	14.4
Non-Hispanic black, both sexes												
2002	2.1	*	1.1	0.8	0.4	0.6	1.2	2.8	5.6	8.6	11.9	9.7
2001	2.1	*	*	0.7	0.5	0.5	1.1	3.1	5.9	9.4	9.9	11.9
2000	2.2	*	*	0.8	0.6	0.7	1.6	2.9	5.8	8.2	12.1	10.5
1999	2.2	*	*	0.8	0.7	0.7	1.4	3.3	6.1	8.2	11.7	10.0
Non-Hispanic black, male												
2002	2.3	*	*	1.1	*	0.9	1.3	2.7	8.1	10.2	16.1	*
2001	2.2	*	*	0.8	*	*	1.3	3.7	7.9	10.4	12.0	*
2000	2.4	*	*	0.9	0.8	1.1	2.0	3.7	7.2	9.9	13.3	*
1999	2.4	*	*	0.7	0.9	0.9	1.8	4.5	7.4	10.2	12.5	*
Non-Hispanic black, female												
2002	1.9	*	*	*	*	*	1.1	2.8	3.7	7.5	9.5	10.5
2001	1.9	*	*	*	*	*	1.0	2.5	4.2	8.7	8.8	9.8
2000	1.9	*	*	0.7	*	*	1.2	2.2	4.7	7.1	11.5	10.5
1999	1.9	*	*	0.9	*	*	1.0	2.2	5.1	6.8	11.2	9.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of brain (C71)												
All origins 3/, both sexes												
2002	4.4	*	0.7	0.8	0.5	0.9	2.2	5.0	9.5	15.8	18.7	16.0
2001	4.3	*	0.6	0.7	0.5	1.0	2.3	4.9	9.6	15.1	20.0	16.0
2000	4.4	0.5	0.7	0.8	0.6	1.0	2.4	5.0	9.5	16.2	19.7	16.3
1999	4.5	*	0.7	0.7	0.5	1.1	2.3	5.0	10.1	16.6	19.9	15.5
All origins 3/, male												
2002	4.9	*	0.8	0.9	0.6	1.1	2.7	6.1	11.5	19.7	23.1	19.7
2001	4.9	*	0.6	0.7	0.6	1.3	2.8	6.0	11.9	18.8	25.3	19.7
2000	5.0	*	0.7	0.8	0.6	1.3	3.1	6.1	11.7	19.9	23.9	21.8
1999	5.0	*	0.7	0.7	0.6	1.3	2.7	6.1	12.2	20.5	25.0	20.1
All origins 3/, female												
2002	3.8	*	0.6	0.7	0.4	0.8	1.7	4.0	7.6	12.5	15.8	14.4
2001	3.8	*	0.6	0.8	0.5	0.8	1.7	3.8	7.5	12.1	16.4	14.4
2000	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.4	13.2	16.9	14.1
1999	4.0	*	0.7	0.7	0.4	0.8	1.9	4.0	8.2	13.5	16.6	13.6
Hispanic, both sexes												
2002	1.6	*	0.7	0.7	0.3	0.5	0.9	3.3	5.8	9.7	13.5	12.2
2001	1.6	*	*	0.5	0.4	0.5	1.4	2.5	5.9	10.3	14.4	*
2000	1.8	*	*	0.9	0.4	0.7	1.5	2.8	6.1	9.8	14.2	15.9
1999	1.6	*	0.9	0.7	*	0.5	1.3	2.9	5.8	10.0	13.7	*
Hispanic, male												
2002	1.8	*	*	0.7	*	*	1.0	4.3	6.6	12.2	16.0	*
2001	1.7	*	*	*	*	0.5	1.4	3.0	6.8	14.4	17.9	*
2000	1.9	*	*	1.2	*	0.7	1.5	2.5	7.7	12.5	18.7	*
1999	1.7	*	*	0.7	*	*	1.7	2.9	7.7	10.5	16.2	*
Hispanic, female												
2002	1.5	*	*	0.7	*	*	0.9	2.3	5.0	7.8	11.8	*
2001	1.5	*	*	0.6	*	*	1.4	2.1	5.1	7.1	12.1	*
2000	1.6	*	*	0.7	*	*	1.5	3.0	4.6	7.6	11.2	*
1999	1.6	*	*	0.8	*	*	0.8	3.0	4.1	9.6	12.1	*
Non-Hispanic 4/, both sexes												
2002	4.8	*	0.7	0.8	0.5	1.1	2.4	5.2	9.7	16.2	18.9	16.1
2001	4.7	*	0.6	0.8	0.6	1.1	2.4	5.1	9.8	15.4	20.2	16.2
2000	4.8	*	0.7	0.7	0.6	1.0	2.5	5.2	9.7	16.6	19.9	16.3
1999	4.9	*	0.7	0.7	0.6	1.2	2.4	5.2	10.4	17.0	20.1	15.6
Non-Hispanic 4/, male												
2002	5.4	*	0.8	0.9	0.7	1.2	3.0	6.3	11.8	20.2	23.4	19.8
2001	5.4	*	0.6	0.7	0.6	1.5	3.0	6.3	12.2	19.0	25.6	20.2
2000	5.4	*	0.7	0.8	0.7	1.4	3.3	6.4	12.0	20.4	24.1	21.7
1999	5.5	*	0.7	0.7	0.7	1.5	2.9	6.4	12.5	21.0	25.2	20.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of brain (C71)												
Non-Hispanic 4/, female												
2002	4.1	*	0.6	0.7	0.4	0.9	1.8	4.1	7.8	12.8	15.9	14.5
2001	4.1	*	0.6	0.8	0.5	0.8	1.8	4.0	7.6	12.4	16.6	14.5
2000	4.1	*	0.7	0.7	0.5	0.7	1.7	4.0	7.6	13.5	17.1	14.1
1999	4.3	*	0.7	0.7	0.5	0.9	2.0	4.0	8.5	13.7	16.8	13.5
Non-Hispanic white, both sexes												
2002	5.5	*	0.6	0.8	0.6	1.2	2.7	5.8	10.6	17.5	20.0	16.9
2001	5.5	*	0.6	0.8	0.6	1.3	2.7	5.7	10.6	16.5	21.4	16.8
2000	5.5	*	0.7	0.7	0.6	1.2	2.7	5.7	10.5	18.0	20.9	16.9
1999	5.6	*	0.7	0.7	0.6	1.4	2.7	5.7	11.3	18.4	21.2	16.3
Non-Hispanic white, male												
2002	6.2	*	0.7	0.9	0.7	1.4	3.4	7.0	12.7	21.7	24.5	21.1
2001	6.2	*	0.7	0.7	0.6	1.7	3.4	6.9	13.1	20.4	27.0	20.6
2000	6.2	*	0.7	0.7	0.7	1.5	3.6	7.0	13.0	21.9	25.3	22.9
1999	6.3	*	0.8	0.8	0.7	1.7	3.2	6.9	13.5	22.6	26.6	21.8
Non-Hispanic white, female												
2002	4.8	*	0.5	0.8	0.4	1.0	2.1	4.6	8.6	13.9	16.9	15.0
2001	4.7	*	0.6	0.9	0.5	0.9	2.0	4.4	8.3	13.2	17.7	15.2
2000	4.8	*	0.7	0.7	0.5	0.9	1.9	4.5	8.2	14.7	17.9	14.5
1999	4.9	*	0.7	0.7	0.5	1.0	2.2	4.5	9.3	14.8	17.7	14.0
Non-Hispanic black, both sexes												
2002	2.0	*	1.1	0.8	0.4	0.6	1.2	2.7	5.3	8.1	11.3	9.4
2001	2.0	*	*	0.7	0.5	0.5	1.1	3.0	5.7	9.3	9.7	11.9
2000	2.1	*	*	0.8	0.6	0.7	1.5	2.8	5.6	8.1	11.7	9.9
1999	2.1	*	*	0.8	0.6	0.6	1.3	3.1	5.8	8.0	11.0	9.4
Non-Hispanic black, male												
2002	2.2	*	*	1.0	*	0.9	1.3	2.6	7.6	9.5	15.2	*
2001	2.2	*	*	0.8	*	*	1.2	3.6	7.6	10.4	12.0	*
2000	2.4	*	*	0.9	0.8	1.1	2.0	3.7	7.0	9.9	12.3	*
1999	2.3	*	*	0.7	0.8	0.8	1.7	4.3	7.2	9.8	11.9	*
Non-Hispanic black, female												
2002	1.9	*	*	*	*	*	1.1	2.8	3.4	7.2	9.1	10.5
2001	1.9	*	*	*	*	*	0.9	2.5	4.2	8.6	8.4	9.8
2000	1.9	*	*	0.7	*	*	1.1	2.1	4.6	6.8	11.3	10.0
1999	1.9	*	*	0.9	*	*	1.0	2.2	4.7	6.8	10.5	8.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other parts of central nervous system (C70,C72)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.4	*
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	*
2000	0.1	*	*	*	*	*	0.0	0.1	0.2	0.3	0.3	0.6
1999	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	0.2	0.5	0.7	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.4	0.5	*
2000	0.1	*	*	*	*	*	*	0.1	*	0.3	*	*
1999	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.5	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	0.1	0.1	0.3	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
2000	0.1	*	*	*	*	*	*	*	0.2	0.3	0.3	*
1999	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.5	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.4	*
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	*
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.4	0.6
1999	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	0.2	0.6	0.7	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.4	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	*	*
1999	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other parts of central nervous system (C70,C72)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	0.1	*	0.3	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.3	*
1999	0.1	*	*	*	*	*	*	0.1	*	0.2	0.5	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.4	*
2001	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.5	*
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.3	0.6
1999	0.1	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	0.6	0.7	*
2001	0.1	*	*	*	*	*	*	*	0.2	0.5	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	*	*
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.5	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)												
All origins 3/, both sexes												
2002	0.7	*	0.5	0.2	0.1	0.1	0.2	0.5	1.2	2.3	4.3	5.3
2001	0.7	*	0.5	0.2	0.1	0.1	0.2	0.6	1.3	2.6	4.1	5.3
2000	0.7	*	0.5	0.2	0.1	0.1	0.2	0.5	1.3	2.6	4.1	6.2
1999	0.7	*	0.5	0.2	0.1	0.1	0.2	0.5	1.3	2.3	4.1	5.8
All origins 3/, male												
2002	0.6	*	0.4	0.3	0.1	*	0.2	0.6	1.3	2.3	4.3	4.2
2001	0.7	*	0.5	0.2	*	0.1	0.2	0.6	1.4	2.9	3.9	4.5
2000	0.7	*	0.5	0.2	*	*	0.2	0.5	1.3	2.7	4.3	5.3
1999	0.7	*	0.5	0.3	*	*	0.2	0.6	1.3	2.6	4.2	5.1
All origins 3/, female												
2002	0.8	*	0.5	0.2	*	0.1	0.2	0.5	1.1	2.3	4.2	5.8
2001	0.8	*	0.4	0.2	*	0.1	0.2	0.5	1.2	2.4	4.3	5.7
2000	0.8	*	0.5	0.2	*	*	0.2	0.4	1.3	2.4	4.0	6.5
1999	0.8	*	0.4	0.2	*	*	0.2	0.5	1.3	2.1	4.0	6.1
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	*	1.5	2.9	5.0	*
2001	0.4	*	*	*	*	*	*	*	1.3	2.7	4.7	*
2000	0.4	*	*	*	*	*	*	*	1.6	2.7	4.0	*
1999	0.4	*	*	*	*	*	*	*	1.5	3.0	6.5	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	3.3	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	6.0	*
2000	0.5	*	*	*	*	*	*	*	2.3	3.3	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	6.9	*
Non-Hispanic 4/, both sexes												
2002	0.8	*	0.5	0.3	0.1	0.1	0.2	0.5	1.1	2.3	4.2	5.3
2001	0.8	*	0.5	0.2	0.1	0.1	0.2	0.6	1.3	2.6	4.1	5.3
2000	0.8	*	0.5	0.2	0.1	0.1	0.2	0.5	1.3	2.6	4.1	6.1
1999	0.8	*	0.6	0.3	0.1	0.1	0.2	0.5	1.3	2.2	4.0	5.7
Non-Hispanic 4/, male												
2002	0.7	*	0.5	0.3	*	*	0.2	0.6	1.2	2.2	4.2	4.2
2001	0.7	*	0.6	0.2	*	0.1	0.2	0.6	1.4	2.9	4.0	4.5
2000	0.7	*	0.5	0.2	*	*	0.2	0.5	1.4	2.8	4.3	5.0
1999	0.7	*	0.7	0.3	*	*	0.3	0.6	1.3	2.5	4.1	5.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)												
Non-Hispanic 4/, female												
2002	0.8	*	0.6	0.3	*	0.1	0.2	0.5	1.0	2.3	4.2	5.8
2001	0.8	*	0.4	0.2	*	*	0.2	0.6	1.2	2.4	4.2	5.6
2000	0.8	*	0.5	0.2	*	*	0.2	0.4	1.2	2.4	4.0	6.5
1999	0.8	*	0.5	0.3	*	*	0.2	0.5	1.3	2.0	3.9	5.9
Non-Hispanic white, both sexes												
2002	0.8	*	0.5	0.3	*	0.1	0.2	0.5	1.2	2.2	4.2	5.4
2001	0.9	*	0.6	0.2	*	0.1	0.2	0.6	1.3	2.8	4.0	5.3
2000	0.8	*	0.5	0.2	0.1	0.1	0.2	0.4	1.3	2.5	4.2	6.1
1999	0.8	*	0.7	0.3	*	0.1	0.2	0.5	1.3	2.2	4.0	5.7
Non-Hispanic white, male												
2002	0.7	*	0.5	0.3	*	*	0.2	0.6	1.3	2.3	4.4	4.2
2001	0.8	*	0.7	0.2	*	*	0.2	0.6	1.3	3.2	4.1	4.4
2000	0.8	*	0.5	0.3	*	*	0.2	0.5	1.4	2.9	4.5	5.3
1999	0.8	*	0.8	0.3	*	*	0.3	0.6	1.3	2.6	4.3	5.0
Non-Hispanic white, female												
2002	0.9	*	0.5	0.3	*	*	0.2	0.5	1.1	2.2	4.1	5.9
2001	0.9	*	0.6	0.2	*	*	0.2	0.5	1.2	2.4	4.0	5.7
2000	0.9	*	*	0.2	*	*	0.2	0.4	1.1	2.2	3.9	6.5
1999	0.9	*	0.5	0.3	*	*	0.2	0.5	1.3	2.0	3.8	6.0
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	*	0.5	1.1	1.9	4.7	*
2001	0.6	*	*	*	*	*	*	0.7	1.4	1.8	5.1	*
2000	0.6	*	*	*	*	*	*	0.8	1.4	2.7	3.7	*
1999	0.6	*	*	0.3	*	*	0.4	*	1.7	2.5	3.3	*
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.6	*	*	*	*	*	*	*	*	*	6.0	*
2001	0.7	*	*	*	*	*	*	1.0	*	2.2	6.1	*
2000	0.7	*	*	*	*	*	*	*	*	2.9	4.5	*
1999	0.6	*	*	*	*	*	*	*	1.6	2.3	3.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)												
All origins 3/, both sexes												
2002	14.1	*	0.1	0.1	0.3	0.6	2.4	8.7	24.1	53.7	96.7	154.5
2001	14.1	*	*	0.1	0.2	0.6	2.4	8.6	24.5	53.9	97.4	155.2
2000	14.5	*	0.1	0.1	0.2	0.6	2.5	9.0	25.1	56.4	101.9	159.0
1999	15.7	*	*	0.1	0.3	0.7	2.8	9.4	28.3	63.0	109.3	168.4
All origins 3/, male												
2002	14.7	*	*	*	0.4	0.7	2.4	10.4	29.0	66.0	118.5	196.9
2001	14.5	*	*	0.1	0.3	0.7	2.5	10.3	29.4	65.1	119.1	196.5
2000	14.8	*	*	0.1	0.2	0.6	2.6	10.5	30.0	68.4	124.6	200.1
1999	16.2	*	*	0.1	0.3	0.9	2.8	11.1	33.9	76.6	133.5	218.9
All origins 3/, female												
2002	13.5	*	*	*	0.2	0.6	2.4	7.1	19.6	43.4	82.3	136.0
2001	13.6	*	*	*	0.2	0.6	2.3	7.1	19.9	44.6	83.1	137.9
2000	14.1	*	*	*	0.2	0.5	2.4	7.5	20.6	46.6	87.0	142.3
1999	15.3	*	*	*	0.2	0.6	2.7	7.9	23.1	51.8	93.6	148.1
Hispanic, both sexes												
2002	4.4	*	*	*	0.3	0.5	1.3	5.3	15.6	35.5	69.2	109.8
2001	4.3	*	*	*	*	0.4	1.7	5.3	14.2	36.3	69.4	122.6
2000	4.4	*	*	*	*	0.3	1.4	5.2	15.4	39.1	77.8	132.0
1999	4.5	*	*	*	*	0.5	1.6	5.8	15.7	40.0	75.6	127.7
Hispanic, male												
2002	4.4	*	*	*	*	0.5	1.2	6.2	18.0	43.6	88.2	137.2
2001	4.3	*	*	*	*	*	1.6	6.2	17.8	43.1	90.3	131.4
2000	4.4	*	*	*	*	*	1.3	5.7	17.8	48.3	98.3	167.3
1999	4.7	*	*	*	*	*	1.5	6.4	19.1	53.6	100.1	153.1
Hispanic, female												
2002	4.3	*	*	*	*	0.6	1.4	4.5	13.6	29.0	56.2	96.2
2001	4.2	*	*	*	*	*	1.8	4.5	10.9	30.9	55.3	118.2
2000	4.4	*	*	*	*	*	1.5	4.7	13.2	31.9	64.1	114.7
1999	4.3	*	*	*	*	*	1.8	5.1	12.7	29.3	59.3	115.4
Non-Hispanic 4/, both sexes												
2002	15.6	*	*	0.1	0.3	0.7	2.6	9.0	24.8	54.8	97.9	155.8
2001	15.5	*	*	0.1	0.3	0.7	2.5	8.9	25.2	54.8	98.4	156.0
2000	15.9	*	*	0.1	0.2	0.6	2.6	9.2	25.8	57.3	102.5	159.7
1999	17.2	*	*	0.1	0.3	0.8	2.9	9.7	29.1	64.2	110.4	169.6
Non-Hispanic 4/, male												
2002	16.3	*	*	*	0.4	0.7	2.6	10.8	29.8	67.3	119.7	198.9
2001	16.0	*	*	*	0.3	0.7	2.7	10.6	30.2	66.2	120.2	198.7
2000	16.3	*	*	*	0.2	0.7	2.7	10.9	30.8	69.4	125.3	201.0
1999	17.8	*	*	0.1	0.3	0.9	3.0	11.4	34.9	77.7	134.4	221.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)												
Non-Hispanic 4/, female												
2002	14.9	*	*	*	0.2	0.6	2.5	7.4	20.1	44.3	83.4	137.1
2001	14.9	*	*	*	0.2	0.6	2.4	7.3	20.6	45.4	84.1	138.2
2000	15.4	*	*	*	0.2	0.6	2.5	7.6	21.2	47.4	87.7	143.1
1999	16.7	*	*	*	0.2	0.6	2.8	8.1	23.8	53.1	94.8	148.9
Non-Hispanic white, both sexes												
2002	16.8	*	*	*	0.3	0.6	2.4	8.5	24.0	54.1	97.6	156.2
2001	16.6	*	*	0.1	0.2	0.6	2.3	8.3	24.1	53.7	98.0	157.8
2000	17.0	*	*	*	0.3	0.6	2.4	8.6	25.0	56.6	101.8	161.0
1999	18.5	*	*	0.1	0.3	0.8	2.7	9.1	27.8	63.5	109.9	169.2
Non-Hispanic white, male												
2002	17.5	*	*	*	0.4	0.7	2.3	10.1	28.5	66.6	118.9	201.0
2001	17.0	*	*	*	0.3	0.7	2.4	9.7	28.3	64.3	119.2	201.1
2000	17.3	*	*	*	0.3	0.7	2.4	9.9	29.0	68.3	123.5	202.1
1999	18.9	*	*	*	0.3	0.9	2.8	10.4	33.1	76.3	133.8	221.3
Non-Hispanic white, female												
2002	16.1	*	*	*	0.2	0.5	2.5	7.0	19.7	43.4	83.3	136.9
2001	16.2	*	*	*	0.2	0.6	2.3	7.0	20.2	44.6	84.0	139.8
2000	16.8	*	*	*	0.3	0.6	2.4	7.2	21.1	46.8	87.6	144.4
1999	18.0	*	*	*	0.2	0.6	2.6	7.8	22.9	52.7	94.3	148.4
Non-Hispanic black, both sexes												
2002	13.0	*	*	*	*	0.9	3.8	13.8	36.7	71.7	117.5	169.7
2001	13.0	*	*	*	0.3	1.0	3.7	14.9	38.2	74.4	115.4	154.2
2000	13.1	*	*	*	*	0.6	4.4	15.6	37.5	72.5	123.1	155.8
1999	14.4	*	*	*	*	0.8	4.4	16.2	44.7	80.1	129.0	185.9
Non-Hispanic black, male												
2002	13.9	*	*	*	*	0.8	4.5	17.0	48.0	89.0	153.9	210.0
2001	14.5	*	*	*	*	0.9	4.0	19.7	51.9	97.6	151.2	212.5
2000	14.6	*	*	*	*	*	5.1	20.4	52.0	92.4	167.2	204.3
1999	15.5	*	*	*	*	0.9	4.4	21.2	58.8	105.4	160.7	242.5
Non-Hispanic black, female												
2002	12.3	*	*	*	*	1.1	3.2	11.1	27.6	59.4	97.0	154.3
2001	11.7	*	*	*	*	1.0	3.4	10.8	27.1	58.0	95.4	132.6
2000	11.8	*	*	*	*	*	3.7	11.4	25.9	58.5	98.3	137.8
1999	13.3	*	*	*	*	*	4.4	11.9	33.3	62.2	111.2	164.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All origins 3/, both sexes												
2002	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
All origins 3/, male												
2002	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.1	275.3
2000	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
All origins 3/, female												
2002	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3
Hispanic, both sexes												
2002	7.3	*	0.9	1.3	2.1	1.9	3.9	7.7	22.1	53.7	100.8	132.5
2001	7.4	*	1.3	1.2	2.2	2.3	3.9	8.2	23.9	52.6	103.7	134.4
2000	7.5	*	1.0	1.1	2.0	1.9	3.9	8.9	25.0	58.8	103.3	143.3
1999	7.5	*	1.0	1.2	2.0	2.1	3.9	10.1	24.2	58.5	101.0	131.9
Hispanic, male												
2002	7.6	*	*	1.3	2.7	2.3	4.4	8.9	26.2	63.3	119.0	180.7
2001	8.3	*	1.4	1.3	2.7	2.7	5.0	9.7	28.9	67.5	142.6	180.9
2000	8.0	*	*	1.4	2.3	2.4	4.3	11.1	30.7	71.3	126.9	197.5
1999	7.6	*	*	1.3	2.1	2.2	4.2	12.0	29.4	68.5	117.8	135.9
Hispanic, female												
2002	7.0	*	*	1.3	1.5	1.4	3.3	6.5	18.5	46.1	88.3	108.6
2001	6.5	*	*	1.1	1.5	1.7	2.6	6.8	19.5	40.9	77.4	111.7
2000	6.9	*	*	0.8	1.5	1.4	3.4	6.8	19.8	48.9	87.5	116.7
1999	7.4	*	*	1.0	1.9	2.1	3.4	8.3	19.6	50.6	89.9	130.0
Non-Hispanic 4/, both sexes												
2002	21.3	0.8	0.7	0.8	1.6	2.4	3.9	10.1	28.6	75.1	145.3	199.8
2001	21.6	0.6	0.9	0.8	1.7	2.5	4.1	10.5	29.8	75.8	146.8	202.1
2000	21.8	1.0	0.9	0.9	1.8	2.4	4.1	10.8	30.3	77.3	150.9	206.5
1999	21.9	0.7	0.9	0.8	1.8	2.4	4.3	10.8	32.3	78.8	149.5	201.4
Non-Hispanic 4/, male												
2002	23.6	*	0.8	1.0	1.8	2.8	4.7	12.1	35.6	97.0	190.9	272.8
2001	23.5	*	1.0	1.0	1.9	2.9	5.0	12.9	36.8	96.3	188.6	278.8
2000	23.7	*	1.0	1.0	2.2	2.7	4.8	13.1	36.4	98.1	195.2	285.0
1999	23.7	*	0.9	0.9	2.2	2.9	5.2	13.3	39.5	98.7	191.5	284.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic 4/, female												
2002	19.2	*	0.7	0.7	1.4	2.0	3.1	8.0	22.2	56.8	115.1	168.3
2001	19.7	*	0.9	0.7	1.4	2.2	3.3	8.2	23.3	58.9	119.3	170.2
2000	20.1	*	0.9	0.7	1.3	2.0	3.4	8.6	24.7	60.1	122.1	174.8
1999	20.1	*	0.8	0.7	1.4	2.0	3.4	8.5	25.6	62.5	122.3	168.2
Non-Hispanic white, both sexes												
2002	23.7	*	0.8	0.8	1.7	2.3	3.8	9.7	28.4	76.7	149.5	206.5
2001	23.8	*	1.1	0.8	1.7	2.4	4.0	10.3	29.5	77.2	150.5	208.2
2000	24.2	1.3	1.0	0.9	1.7	2.3	4.0	10.5	30.3	79.0	155.1	210.7
1999	24.1	*	1.0	0.8	1.8	2.4	4.2	10.6	32.0	80.4	153.0	206.4
Non-Hispanic white, male												
2002	26.2	*	0.9	1.0	1.9	2.7	4.5	11.8	35.4	99.3	196.8	283.5
2001	25.9	*	1.0	0.8	2.0	2.8	4.7	12.8	36.4	97.9	193.6	288.7
2000	26.2	*	1.1	1.0	2.1	2.6	4.8	12.7	36.3	100.1	201.0	293.8
1999	26.1	*	1.0	0.9	2.2	2.7	5.0	13.1	39.2	100.7	196.1	291.8
Non-Hispanic white, female												
2002	21.2	*	0.7	0.7	1.4	1.9	3.1	7.6	21.7	57.5	117.8	173.2
2001	21.8	*	1.1	0.7	1.4	2.0	3.2	7.9	23.0	59.5	122.0	174.6
2000	22.3	*	0.9	0.8	1.2	1.9	3.2	8.4	24.6	61.2	125.0	177.2
1999	22.2	*	0.9	0.6	1.3	2.0	3.3	8.3	25.2	63.4	124.7	172.4
Non-Hispanic black, both sexes												
2002	14.1	*	*	0.8	1.5	3.0	4.8	13.9	36.4	77.0	123.5	153.7
2001	14.5	*	*	1.0	1.8	3.5	5.8	13.5	37.1	79.0	127.6	157.3
2000	14.3	*	*	0.7	2.3	3.0	5.4	15.0	36.5	76.2	122.3	177.1
1999	14.5	*	*	0.9	2.4	3.0	5.6	14.1	39.5	77.5	128.2	162.3
Non-Hispanic black, male												
2002	15.1	*	*	1.0	1.5	3.8	6.2	16.5	45.2	99.4	155.3	203.5
2001	15.6	*	*	1.3	2.0	4.0	7.6	16.3	46.6	100.7	161.4	206.7
2000	15.2	*	*	0.8	2.8	3.7	5.7	18.2	45.3	97.8	157.4	226.6
1999	15.6	*	*	1.2	2.6	3.9	7.0	17.5	48.4	96.9	162.9	221.1
Non-Hispanic black, female												
2002	13.1	*	*	*	1.5	2.3	3.5	11.6	29.4	61.2	105.4	134.6
2001	13.5	*	*	*	1.7	3.1	4.2	11.1	29.5	63.7	108.7	139.0
2000	13.5	*	*	*	1.8	2.5	5.2	12.2	29.4	61.0	102.6	158.7
1999	13.6	*	*	0.7	2.1	2.2	4.3	11.1	32.3	63.8	108.7	140.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All origins 3/, both sexes												
2002	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
All origins 3/, male												
2002	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001	0.5	*	*	*	0.2	0.5	0.5	0.6	0.7	1.6	2.6	2.6
2000	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
All origins 3/, female												
2002	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	1.8	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	4.5	*
1999	0.3	*	*	*	*	*	*	*	*	2.5	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.8
2001	0.5	*	*	*	0.2	0.5	0.4	0.5	0.5	1.1	2.2	2.2
2000	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.4
1999	0.5	*	*	*	0.3	0.4	0.4	0.4	0.7	1.5	2.1	2.3
Non-Hispanic 4/, male												
2002	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.4	3.7
2001	0.5	*	*	*	0.2	0.5	0.5	0.6	0.7	1.5	2.6	2.5
2000	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.3	3.7
1999	0.6	*	*	*	0.3	0.5	0.5	0.5	1.0	1.9	2.3	3.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	1.5	2.4
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.5	0.8	2.0	2.0
2000	0.4	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.7	2.0
1999	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.7
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	0.2	0.5	0.4	0.5	0.6	1.4	2.0	2.9
2001	0.5	*	*	*	0.2	0.5	0.4	0.5	0.6	1.2	2.3	2.3
2000	0.5	*	*	*	0.2	0.5	0.4	0.4	0.6	1.3	2.1	2.6
1999	0.6	*	*	*	0.3	0.4	0.4	0.5	0.8	1.5	2.2	2.3
Non-Hispanic white, male												
2002	0.6	*	*	*	0.2	0.5	0.4	0.5	0.8	1.9	2.6	4.0
2001	0.6	*	*	*	0.3	0.5	0.5	0.6	0.7	1.6	2.7	2.6
2000	0.6	*	*	*	0.2	0.5	0.4	0.5	0.7	1.6	2.5	3.9
1999	0.7	*	*	*	0.3	0.5	0.5	0.5	1.1	2.0	2.5	3.8
Non-Hispanic white, female												
2002	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.6	2.5
2001	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	2.0	2.3
2000	0.5	*	*	*	0.2	0.4	0.3	0.4	0.6	1.1	1.9	2.1
1999	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	1.2	2.0	1.7
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	0.4	0.4	0.4	0.8	*	*	*
2001	0.4	*	*	*	*	0.8	0.4	*	*	*	*	*
2000	0.3	*	*	*	*	*	0.4	*	*	*	*	*
1999	0.4	*	*	*	*	0.5	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	0.8	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)												
All origins 3/, both sexes												
2002	1.2	*	*	0.1	0.1	0.2	0.3	0.8	1.9	4.8	8.5	10.0
2001	1.2	*	*	*	0.1	0.2	0.4	0.8	2.1	4.6	8.3	10.1
2000	1.2	*	*	0.1	0.1	0.2	0.3	0.8	2.0	4.6	8.9	10.7
1999	1.3	*	*	*	0.1	0.1	0.3	0.9	2.3	5.3	8.8	10.0
All origins 3/, male												
2002	1.4	*	*	0.1	0.2	0.2	0.4	1.0	2.5	6.6	11.6	13.9
2001	1.4	*	*	*	0.2	0.2	0.6	1.0	2.8	6.2	11.1	14.2
2000	1.4	*	*	*	0.2	0.2	0.4	1.0	2.7	6.0	11.5	17.0
1999	1.4	*	*	*	0.2	0.2	0.4	1.2	2.9	6.7	11.0	13.1
All origins 3/, female												
2002	1.0	*	*	*	*	0.2	0.2	0.6	1.4	3.3	6.5	8.3
2001	1.0	*	*	*	*	0.1	0.2	0.5	1.4	3.3	6.4	8.3
2000	1.1	*	*	*	*	0.1	0.2	0.6	1.4	3.5	7.2	8.2
1999	1.1	*	*	*	*	*	0.2	0.6	1.7	4.1	7.4	8.8
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	0.6	1.4	3.6	5.8	*
2001	0.4	*	*	*	*	*	*	0.7	1.7	3.2	4.3	*
2000	0.4	*	*	*	*	*	*	*	1.7	2.7	4.3	*
1999	0.4	*	*	*	*	*	*	*	1.8	3.5	5.4	*
Hispanic, male												
2002	0.5	*	*	*	*	*	*	*	*	5.0	*	*
2001	0.6	*	*	*	*	*	*	*	*	4.8	*	*
2000	0.5	*	*	*	*	*	*	*	2.5	*	*	*
1999	0.5	*	*	*	*	*	*	*	2.6	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	6.5	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	3.8	*	*
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	0.1	0.1	0.2	0.3	0.8	2.0	4.9	8.7	10.1
2001	1.3	*	*	*	0.1	0.2	0.4	0.8	2.1	4.7	8.4	10.2
2000	1.4	*	*	0.1	0.1	0.2	0.3	0.8	2.0	4.7	9.1	10.8
1999	1.4	*	*	*	0.1	0.1	0.3	0.9	2.3	5.4	8.9	10.1
Non-Hispanic 4/, male												
2002	1.6	*	*	0.1	0.2	0.2	0.4	1.0	2.6	6.7	11.9	14.1
2001	1.6	*	*	*	0.2	0.2	0.6	1.0	2.9	6.3	11.2	14.5
2000	1.6	*	*	*	0.2	0.2	0.4	1.1	2.7	6.2	11.7	17.0
1999	1.6	*	*	*	0.2	0.2	0.5	1.2	2.9	6.9	11.2	13.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	0.1	0.2	0.6	1.4	3.3	6.5	8.4
2001	1.1	*	*	*	*	0.1	0.2	0.5	1.4	3.3	6.6	8.4
2000	1.2	*	*	*	*	0.1	0.2	0.6	1.5	3.5	7.4	8.3
1999	1.2	*	*	*	*	*	0.2	0.6	1.8	4.1	7.4	8.9
Non-Hispanic white, both sexes												
2002	1.5	*	*	0.1	0.1	0.1	0.3	0.7	1.9	5.1	9.1	10.4
2001	1.5	*	*	*	0.1	0.2	0.4	0.8	2.2	4.8	8.7	10.6
2000	1.5	*	*	0.1	0.1	0.1	0.3	0.8	2.1	4.9	9.5	11.4
1999	1.6	*	*	*	0.1	0.1	0.3	0.9	2.3	5.6	9.4	10.5
Non-Hispanic white, male												
2002	1.8	*	*	0.2	0.2	0.2	0.4	1.0	2.6	7.0	12.5	14.2
2001	1.7	*	*	*	0.2	0.2	0.5	1.0	3.0	6.5	11.5	14.9
2000	1.7	*	*	*	0.2	0.2	0.4	1.0	2.8	6.3	12.1	18.0
1999	1.8	*	*	*	0.2	0.2	0.4	1.3	2.9	7.1	11.6	13.6
Non-Hispanic white, female												
2002	1.2	*	*	*	*	*	*	0.5	1.4	3.4	6.8	8.7
2001	1.2	*	*	*	*	*	0.2	0.5	1.4	3.4	6.8	8.8
2000	1.3	*	*	*	*	*	0.2	0.6	1.5	3.7	7.8	8.8
1999	1.4	*	*	*	*	*	0.2	0.6	1.8	4.3	7.9	9.2
Non-Hispanic black, both sexes												
2002	0.9	*	*	*	*	*	0.4	1.4	2.2	4.7	4.3	7.6
2001	0.8	*	*	*	*	*	0.6	1.0	2.2	3.7	5.6	7.5
2000	0.8	*	*	*	*	*	0.4	1.2	1.8	4.0	5.2	*
1999	0.8	*	*	*	*	*	0.5	1.0	2.5	4.2	4.5	*
Non-Hispanic black, male												
2002	1.0	*	*	*	*	*	*	1.4	2.7	6.3	5.9	*
2001	1.0	*	*	*	*	*	0.9	1.6	2.9	4.6	8.4	*
2000	0.9	*	*	*	*	*	*	1.8	2.5	5.2	7.4	*
1999	1.0	*	*	*	*	*	*	1.1	3.2	6.6	6.9	*
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	1.3	1.8	3.5	3.5	*
2001	0.6	*	*	*	*	*	*	*	1.6	3.0	4.0	*
2000	0.6	*	*	*	*	*	*	*	*	3.2	4.0	*
1999	0.6	*	*	*	*	*	*	*	1.9	2.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All origins 3/, both sexes												
2002	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
All origins 3/, male												
2002	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
All origins 3/, female												
2002	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9
Hispanic, both sexes												
2002	2.8	*	*	*	0.4	0.6	1.3	3.0	9.4	22.7	43.7	50.4
2001	2.7	*	*	*	0.3	0.6	1.4	3.6	10.9	18.3	43.5	59.4
2000	2.9	*	*	*	0.3	0.6	1.6	3.6	11.1	25.7	41.5	59.7
1999	2.8	*	*	*	*	0.6	1.4	4.4	9.3	25.2	42.7	56.1
Hispanic, male												
2002	2.7	*	*	*	*	0.6	1.5	3.6	11.7	26.0	45.5	60.2
2001	3.0	*	*	*	*	0.7	2.0	4.4	13.2	24.4	58.6	68.6
2000	3.1	*	*	*	*	0.7	1.9	4.6	13.9	30.2	48.2	82.6
1999	2.8	*	*	*	*	*	1.7	5.7	11.8	27.5	48.0	60.4
Hispanic, female												
2002	2.8	*	*	*	*	*	1.2	2.3	7.4	20.1	42.4	45.6
2001	2.4	*	*	*	*	*	0.8	2.8	8.8	13.5	33.2	54.9
2000	2.7	*	*	*	*	*	1.3	2.6	8.6	22.2	37.0	48.5
1999	2.8	*	*	*	*	*	1.0	3.1	7.1	23.4	39.2	54.1
Non-Hispanic 4/, both sexes												
2002	8.3	*	*	0.1	0.3	0.7	1.5	4.0	11.6	29.2	57.7	78.2
2001	8.6	*	*	0.1	0.4	0.8	1.7	4.5	12.3	29.9	59.2	80.3
2000	8.8	*	*	0.1	0.5	0.8	1.7	4.7	13.2	30.7	61.7	80.9
1999	8.9	*	*	0.1	0.4	0.7	1.8	4.8	13.5	32.1	62.4	78.3
Non-Hispanic 4/, male												
2002	9.0	*	*	0.2	0.4	0.9	2.0	5.1	14.6	36.5	71.8	101.4
2001	9.2	*	*	0.1	0.5	0.9	2.3	5.9	15.1	37.2	71.7	108.1
2000	9.4	*	*	0.2	0.7	0.9	2.2	6.1	15.9	37.7	75.6	106.7
1999	9.4	*	*	0.2	0.5	0.9	2.3	6.2	16.5	38.8	75.5	106.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic 4/, female												
2002	7.7	*	*	*	0.3	0.5	1.1	3.0	8.9	23.0	48.3	68.2
2001	8.0	*	*	*	0.3	0.6	1.2	3.1	9.7	23.8	50.9	68.7
2000	8.3	*	*	*	0.3	0.6	1.2	3.4	10.6	24.8	52.6	70.4
1999	8.4	*	*	*	0.3	0.5	1.3	3.4	10.8	26.5	53.9	67.2
Non-Hispanic white, both sexes												
2002	9.5	*	*	0.2	0.3	0.7	1.5	4.0	12.0	30.6	60.9	83.1
2001	9.7	*	*	0.1	0.3	0.7	1.7	4.5	12.7	31.4	62.3	84.1
2000	10.1	*	*	0.1	0.4	0.7	1.7	4.8	13.8	32.3	65.1	85.1
1999	10.1	*	*	0.1	0.4	0.7	1.7	4.9	14.1	33.9	65.6	82.0
Non-Hispanic white, male												
2002	10.2	*	*	0.2	0.4	0.8	1.9	5.1	15.0	38.3	75.6	107.6
2001	10.4	*	*	*	0.4	0.9	2.1	6.0	15.5	38.8	75.2	113.1
2000	10.6	*	*	0.2	0.6	0.8	2.2	6.1	16.5	39.5	79.6	113.0
1999	10.6	*	*	0.2	0.5	0.8	2.2	6.3	17.0	40.8	78.9	110.0
Non-Hispanic white, female												
2002	8.8	*	*	*	0.2	0.5	1.1	2.9	9.2	24.1	51.0	72.6
2001	9.2	*	*	*	0.3	0.5	1.2	3.1	10.1	25.2	53.7	72.0
2000	9.5	*	*	*	0.3	0.6	1.1	3.4	11.3	26.3	55.6	73.8
1999	9.6	*	*	*	0.2	0.5	1.2	3.5	11.4	28.1	56.9	70.8
Non-Hispanic black, both sexes												
2002	4.0	*	*	*	0.4	1.2	2.0	4.8	10.8	21.6	30.1	31.2
2001	4.1	*	*	*	0.6	1.2	2.3	5.1	10.9	21.3	30.6	40.0
2000	3.9	*	*	*	0.6	1.1	2.1	5.0	10.8	19.7	27.1	36.6
1999	4.1	*	*	*	0.6	1.1	2.4	4.8	11.2	20.5	32.1	40.4
Non-Hispanic black, male												
2002	4.5	*	*	*	*	1.6	2.9	5.7	13.9	28.3	35.8	43.8
2001	4.7	*	*	*	0.8	1.3	3.4	6.7	13.7	28.2	37.2	55.4
2000	4.3	*	*	*	*	1.2	2.6	7.0	14.8	25.3	31.8	43.4
1999	4.7	*	*	*	*	1.6	3.3	6.6	14.8	26.2	41.0	59.4
Non-Hispanic black, female												
2002	3.6	*	*	*	*	0.8	1.2	4.0	8.3	16.9	26.9	26.3
2001	3.7	*	*	*	*	1.1	1.4	3.7	8.6	16.5	27.0	34.2
2000	3.5	*	*	*	*	0.9	1.7	3.3	7.5	15.7	24.5	34.0
1999	3.6	*	*	*	*	*	1.7	3.2	8.2	16.5	27.1	33.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant immunoproliferative diseases (C88)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.2	0.4	1.4	2.0
2001	0.1	*	*	*	*	*	*	*	0.2	0.5	1.1	2.0
2000	0.1	*	*	*	*	*	*	*	0.1	0.4	1.2	2.1
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.7
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	2.2	2.8
2001	0.1	*	*	*	*	*	*	*	0.2	0.6	1.6	2.9
2000	0.1	*	*	*	*	*	*	*	0.2	0.6	1.7	3.4
1999	0.2	*	*	*	*	*	*	*	0.3	0.8	1.9	3.9
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.6
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.9	1.6
1999	0.1	*	*	*	*	*	*	*	0.2	0.3	1.1	2.2
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	2.0
2001	0.1	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	1.2	2.2
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	2.8
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.7	2.2	2.9
2001	0.2	*	*	*	*	*	*	*	0.2	0.7	1.6	3.0
2000	0.2	*	*	*	*	*	*	*	*	0.6	1.8	3.5
1999	0.2	*	*	*	*	*	*	*	0.3	0.8	1.9	4.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant immunoproliferative diseases (C88)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.9	1.7
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.9	1.7
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.9	1.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.3	1.1	2.3
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	2.1
2001	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.2
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.3	2.2
1999	0.2	*	*	*	*	*	*	*	0.2	0.6	1.5	3.0
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	0.3	0.7	2.3	3.0
2001	0.2	*	*	*	*	*	*	*	0.2	0.7	1.8	3.2
2000	0.2	*	*	*	*	*	*	*	*	0.7	1.9	3.6
1999	0.2	*	*	*	*	*	*	*	0.3	0.9	2.0	4.4
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.9	1.7
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.9	1.7
1999	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.4
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)												
All origins 3/, both sexes												
2002	3.8	*	*	*	*	*	0.3	1.8	6.1	17.0	29.8	34.7
2001	3.8	*	*	*	*	*	0.3	1.7	6.4	16.9	29.8	34.2
2000	3.8	*	*	*	*	*	0.3	1.8	6.2	16.9	30.3	35.7
1999	3.8	*	*	*	*	0.1	0.3	1.8	6.9	16.7	29.6	33.9
All origins 3/, male												
2002	4.0	*	*	*	*	*	0.3	2.2	7.3	20.4	37.5	48.0
2001	4.0	*	*	*	*	*	0.4	2.1	7.6	20.2	37.1	45.1
2000	3.9	*	*	*	*	*	0.4	2.1	6.8	19.8	37.9	48.3
1999	3.9	*	*	*	*	*	0.4	2.1	8.2	19.4	36.5	47.4
All origins 3/, female												
2002	3.6	*	*	*	*	*	0.3	1.4	5.0	14.1	24.7	29.0
2001	3.6	*	*	*	*	*	0.3	1.3	5.4	14.2	25.0	29.7
2000	3.7	*	*	*	*	*	0.3	1.6	5.6	14.4	25.4	30.6
1999	3.7	*	*	*	*	*	0.3	1.5	5.7	14.5	25.2	28.4
Hispanic, both sexes												
2002	1.4	*	*	*	*	*	*	1.8	4.9	14.1	26.2	32.2
2001	1.4	*	*	*	*	*	*	1.3	5.8	14.7	24.5	29.4
2000	1.3	*	*	*	*	*	*	1.4	5.1	14.9	24.1	30.5
1999	1.4	*	*	*	*	*	*	1.7	6.8	13.2	22.5	27.4
Hispanic, male												
2002	1.5	*	*	*	*	*	*	2.0	6.5	17.2	30.8	51.9
2001	1.5	*	*	*	*	*	*	1.4	6.0	20.0	35.8	38.1
2000	1.3	*	*	*	*	*	*	1.5	5.2	18.1	30.0	*
1999	1.3	*	*	*	*	*	*	1.8	7.8	14.3	24.0	*
Hispanic, female												
2002	1.4	*	*	*	*	*	*	1.5	3.5	11.6	23.0	22.4
2001	1.3	*	*	*	*	*	*	1.2	5.6	10.6	16.9	25.1
2000	1.4	*	*	*	*	*	*	*	5.1	12.3	20.1	29.7
1999	1.5	*	*	*	*	*	*	1.5	5.9	12.4	21.5	30.2
Non-Hispanic 4/, both sexes												
2002	4.2	*	*	*	*	*	0.3	1.8	6.2	17.1	29.9	34.7
2001	4.1	*	*	*	*	*	0.3	1.7	6.5	17.0	29.9	34.4
2000	4.1	*	*	*	*	*	0.3	1.9	6.2	16.9	30.5	35.9
1999	4.1	*	*	*	*	*	0.4	1.8	6.9	16.9	29.8	34.0
Non-Hispanic 4/, male												
2002	4.4	*	*	*	*	*	0.3	2.2	7.4	20.5	37.8	47.7
2001	4.3	*	*	*	*	*	0.4	2.1	7.7	20.1	37.0	45.3
2000	4.3	*	*	*	*	*	0.4	2.1	6.8	19.9	38.1	49.0
1999	4.3	*	*	*	*	*	0.4	2.1	8.2	19.6	36.9	48.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)												
Non-Hispanic 4/, female												
2002	3.9	*	*	*	*	*	0.3	1.4	5.1	14.3	24.7	29.1
2001	4.0	*	*	*	*	*	0.3	1.3	5.4	14.4	25.2	29.8
2000	4.0	*	*	*	*	*	0.3	1.6	5.6	14.5	25.6	30.5
1999	4.0	*	*	*	*	*	0.3	1.5	5.7	14.6	25.2	28.3
Non-Hispanic white, both sexes												
2002	4.3	*	*	*	*	*	0.2	1.5	5.5	15.9	28.8	33.6
2001	4.2	*	*	*	*	*	0.3	1.5	5.6	15.7	28.4	32.9
2000	4.2	*	*	*	*	*	0.3	1.6	5.4	15.7	28.9	33.0
1999	4.1	*	*	*	*	*	0.3	1.5	5.8	15.6	28.2	32.0
Non-Hispanic white, male												
2002	4.6	*	*	*	*	*	0.3	1.9	6.6	19.5	36.8	47.1
2001	4.4	*	*	*	*	*	0.3	1.9	6.7	18.8	35.6	44.1
2000	4.4	*	*	*	*	*	0.4	1.8	6.1	18.5	36.6	47.3
1999	4.4	*	*	*	*	*	0.3	1.9	7.2	18.4	35.4	46.2
Non-Hispanic white, female												
2002	3.9	*	*	*	*	*	0.2	1.1	4.4	12.9	23.4	27.7
2001	4.0	*	*	*	*	*	0.2	1.1	4.5	13.1	23.6	28.2
2000	4.0	*	*	*	*	*	0.2	1.4	4.8	13.3	23.8	27.2
1999	3.9	*	*	*	*	*	0.2	1.2	4.5	13.3	23.5	26.3
Non-Hispanic black, both sexes												
2002	4.8	*	*	*	*	*	0.7	4.2	13.2	31.7	48.6	54.4
2001	5.0	*	*	*	*	*	0.9	3.6	15.1	31.6	52.9	58.1
2000	5.1	*	*	*	*	*	0.8	4.5	14.1	32.3	54.9	71.2
1999	5.1	*	*	*	*	*	0.9	4.2	16.5	30.6	53.0	62.7
Non-Hispanic black, male												
2002	4.8	*	*	*	*	*	0.8	4.8	15.8	37.2	59.2	70.0
2001	5.0	*	*	*	*	*	1.1	4.3	18.3	36.9	65.7	70.5
2000	4.9	*	*	*	*	*	*	4.9	15.3	39.1	67.3	77.5
1999	4.9	*	*	*	*	*	1.1	4.3	18.5	35.3	64.1	80.8
Non-Hispanic black, female												
2002	4.8	*	*	*	*	*	*	3.8	11.2	27.7	42.6	48.5
2001	5.0	*	*	*	*	*	0.8	3.0	12.5	27.8	45.7	53.5
2000	5.4	*	*	*	*	*	0.9	4.1	13.1	27.6	47.9	68.9
1999	5.2	*	*	*	*	*	*	4.2	15.0	27.2	46.7	56.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All origins 3/, both sexes												
2002	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
All origins 3/, male												
2002	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
All origins 3/, female												
2002	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6
Hispanic, both sexes												
2002	2.8	*	0.8	1.1	1.6	1.1	2.1	2.5	7.2	15.5	27.4	47.1
2001	3.0	*	1.3	1.1	1.7	1.5	1.9	3.0	6.5	17.4	33.4	42.5
2000	2.9	*	0.9	1.1	1.5	1.1	1.9	3.5	7.8	16.9	32.6	49.1
1999	2.9	*	0.9	1.1	1.6	1.4	2.0	3.4	7.0	17.3	31.0	44.2
Hispanic, male												
2002	3.0	*	*	1.1	2.0	1.4	2.3	2.8	7.2	18.5	38.6	65.3
2001	3.4	*	1.4	1.1	2.2	1.8	2.3	3.3	8.4	19.6	45.6	68.6
2000	3.3	*	*	1.3	1.7	1.5	2.0	4.1	10.2	20.7	42.8	76.6
1999	3.1	*	*	1.1	1.7	1.4	1.9	3.6	8.6	23.5	39.1	49.6
Hispanic, female												
2002	2.6	*	*	1.2	1.1	0.7	1.9	2.3	7.3	13.1	19.7	38.1
2001	2.7	*	*	1.1	1.1	1.1	1.4	2.6	4.8	15.7	25.1	29.8
2000	2.6	*	*	0.8	1.3	*	1.8	2.9	5.6	13.9	25.8	35.6
1999	2.8	*	*	1.0	1.5	1.3	2.1	3.2	5.6	12.4	25.7	41.6
Non-Hispanic 4/, both sexes												
2002	8.2	*	0.6	0.7	1.1	1.2	1.6	3.7	10.0	27.0	54.3	81.7
2001	8.2	*	0.9	0.7	1.1	1.2	1.7	3.8	10.3	27.2	54.1	82.7
2000	8.2	1.0	0.9	0.7	1.1	1.2	1.7	3.8	10.1	27.8	55.3	84.7
1999	8.2	*	0.7	0.7	1.2	1.2	1.7	3.8	10.8	27.8	53.6	83.7
Non-Hispanic 4/, male												
2002	9.4	*	0.7	0.8	1.2	1.4	1.9	4.3	12.6	37.3	76.5	116.5
2001	9.3	*	0.9	0.8	1.2	1.4	1.9	4.3	13.1	36.6	75.4	119.3
2000	9.3	*	0.9	0.8	1.4	1.4	1.9	4.3	12.6	38.2	77.0	121.5
1999	9.2	*	0.8	0.8	1.4	1.3	1.9	4.4	13.4	37.5	74.6	122.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Non-Hispanic 4/, female												
2002	7.0	*	0.5	0.6	0.9	1.0	1.4	3.2	7.6	18.4	39.5	66.6
2001	7.2	*	0.9	0.6	0.9	1.1	1.5	3.4	7.7	19.4	40.1	67.5
2000	7.2	*	0.9	0.6	0.8	1.0	1.6	3.3	7.7	19.3	41.1	69.8
1999	7.2	*	0.7	0.6	0.9	1.1	1.5	3.2	8.5	19.8	40.0	68.3
Non-Hispanic white, both sexes												
2002	9.1	*	0.7	0.7	1.1	1.2	1.7	3.7	10.0	28.2	56.2	84.4
2001	9.1	*	1.0	0.7	1.1	1.2	1.7	3.8	10.5	28.1	56.0	86.1
2000	9.2	1.3	0.9	0.7	1.1	1.1	1.7	3.7	10.2	29.0	57.5	87.3
1999	9.1	*	0.8	0.7	1.1	1.2	1.7	3.7	11.0	28.6	55.3	86.8
Non-Hispanic white, male												
2002	10.5	*	0.8	0.7	1.3	1.3	1.9	4.2	12.7	38.7	79.2	121.3
2001	10.3	*	1.0	0.7	1.3	1.4	1.8	4.3	13.3	37.7	78.0	125.3
2000	10.4	*	1.0	0.8	1.3	1.3	1.9	4.2	12.7	39.5	79.9	125.5
1999	10.2	*	0.8	0.7	1.4	1.3	1.9	4.3	13.6	38.6	77.0	127.2
Non-Hispanic white, female												
2002	7.8	*	0.5	0.6	0.9	1.0	1.5	3.2	7.5	19.3	40.7	68.4
2001	8.0	*	1.1	0.6	0.9	1.0	1.6	3.3	7.8	19.9	41.4	69.8
2000	8.1	*	0.9	0.6	0.8	0.9	1.5	3.2	7.8	20.1	42.7	72.0
1999	8.0	*	0.9	0.6	0.8	1.1	1.5	3.2	8.5	20.3	41.1	70.8
Non-Hispanic black, both sexes												
2002	4.8	*	*	0.6	0.9	1.3	1.6	4.4	11.5	22.6	42.9	65.3
2001	4.9	*	*	0.9	1.0	1.5	2.1	4.3	10.1	25.0	42.1	57.1
2000	4.9	*	*	0.6	1.4	1.7	2.1	5.0	10.8	23.1	38.8	66.1
1999	4.9	*	*	0.8	1.5	1.3	1.9	4.4	10.8	25.2	41.6	55.9
Non-Hispanic black, male												
2002	5.3	*	*	0.7	1.0	1.5	1.8	5.2	14.1	32.2	58.0	87.5
2001	5.4	*	*	1.2	1.1	1.8	2.6	4.6	13.4	34.4	56.1	75.1
2000	5.5	*	*	0.7	1.9	2.0	2.0	5.6	14.0	32.3	55.8	98.6
1999	5.4	*	*	1.0	1.6	1.5	2.1	5.7	13.7	33.4	55.7	78.4
Non-Hispanic black, female												
2002	4.4	*	*	*	0.8	1.1	1.4	3.6	9.4	15.8	34.3	56.9
2001	4.4	*	*	*	1.0	1.1	1.7	4.0	7.4	18.4	34.2	50.5
2000	4.4	*	*	*	0.9	1.4	2.2	4.4	8.3	16.5	29.2	54.1
1999	4.4	*	*	0.6	1.3	1.1	1.7	3.3	8.5	19.4	33.8	47.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lymphoid leukemia (C91)												
All origins 3/, both sexes												
2002	2.2	*	0.2	0.4	0.5	0.3	0.4	0.9	2.5	6.7	14.9	30.1
2001	2.2	*	0.3	0.4	0.5	0.4	0.4	0.8	2.5	7.0	15.1	31.2
2000	2.2	*	0.3	0.4	0.5	0.3	0.3	0.8	2.4	7.1	15.7	32.0
1999	2.2	*	0.3	0.4	0.5	0.3	0.3	0.8	2.8	7.6	15.9	30.8
All origins 3/, male												
2002	2.6	*	*	0.5	0.7	0.4	0.6	1.1	3.3	10.0	21.3	43.0
2001	2.5	*	0.3	0.5	0.7	0.4	0.5	1.1	3.6	10.0	21.2	43.8
2000	2.6	*	0.3	0.5	0.7	0.4	0.4	1.0	3.5	10.4	22.4	43.8
1999	2.6	*	0.3	0.4	0.7	0.4	0.4	1.1	3.9	10.7	22.7	45.6
All origins 3/, female												
2002	1.8	*	*	0.4	0.4	0.3	0.3	0.6	1.7	4.0	10.6	24.5
2001	1.8	*	0.3	0.3	0.3	0.3	0.3	0.6	1.5	4.4	11.0	25.9
2000	1.8	*	0.3	0.3	0.3	0.2	0.2	0.6	1.4	4.5	11.2	27.3
1999	1.9	*	0.3	0.3	0.3	0.2	0.3	0.5	1.7	5.0	11.5	24.8
Hispanic, both sexes												
2002	0.8	*	*	0.7	0.8	0.3	0.7	0.7	1.6	2.4	5.7	*
2001	0.9	*	*	0.7	1.0	0.5	0.5	*	1.5	4.0	6.0	*
2000	0.9	*	*	0.6	0.9	0.4	*	1.1	1.2	3.5	6.5	13.3
1999	0.9	*	*	0.5	0.9	0.5	0.5	0.7	1.8	2.8	7.5	15.4
Hispanic, male												
2002	0.9	*	*	0.6	0.9	*	0.9	*	*	*	*	*
2001	1.0	*	*	0.7	1.3	0.6	*	*	*	4.2	*	*
2000	1.0	*	*	0.9	1.0	0.6	*	*	*	4.9	*	*
1999	1.0	*	*	*	0.9	0.6	*	*	*	*	*	*
Hispanic, female												
2002	0.8	*	*	0.8	*	*	*	*	*	*	*	*
2001	0.7	*	*	0.7	0.6	*	*	*	*	3.8	*	*
2000	0.7	*	*	*	0.7	*	*	*	*	*	*	*
1999	0.8	*	*	*	0.9	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	2.4	*	0.2	0.4	0.5	0.3	0.4	0.9	2.5	7.0	15.3	30.8
2001	2.4	*	0.2	0.3	0.4	0.3	0.4	0.8	2.6	7.1	15.5	31.9
2000	2.4	*	0.3	0.4	0.4	0.2	0.3	0.8	2.5	7.4	16.0	32.6
1999	2.4	*	0.3	0.3	0.4	0.3	0.3	0.8	2.8	7.8	16.2	31.3
Non-Hispanic 4/, male												
2002	2.8	*	*	0.4	0.6	0.4	0.5	1.1	3.4	10.4	22.0	44.3
2001	2.8	*	*	0.4	0.6	0.4	0.4	1.1	3.7	10.4	21.8	44.8
2000	2.8	*	*	0.5	0.6	0.3	0.4	1.0	3.6	10.7	23.0	44.8
1999	2.9	*	*	0.4	0.6	0.4	0.4	1.1	4.0	11.1	23.3	46.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lymphoid leukemia (C91)												
Non-Hispanic 4/, female												
2002	2.0	*	*	0.3	0.4	0.3	0.2	0.6	1.7	4.1	10.9	25.0
2001	2.0	*	*	0.3	0.3	0.3	0.3	0.6	1.5	4.5	11.3	26.6
2000	2.0	*	*	0.3	0.2	0.2	0.2	0.6	1.4	4.6	11.4	27.8
1999	2.0	*	*	0.3	0.2	0.2	0.2	0.5	1.7	5.1	11.7	25.2
Non-Hispanic white, both sexes												
2002	2.7	*	*	0.4	0.5	0.3	0.4	0.9	2.5	7.3	15.8	32.5
2001	2.7	*	0.2	0.3	0.5	0.3	0.4	0.8	2.6	7.3	16.0	33.7
2000	2.7	*	0.3	0.4	0.4	0.3	0.3	0.7	2.6	7.6	16.7	33.8
1999	2.7	*	0.3	0.3	0.4	0.3	0.3	0.8	2.8	8.1	16.8	32.6
Non-Hispanic white, male												
2002	3.2	*	*	0.4	0.6	0.4	0.6	1.2	3.4	10.9	22.7	46.5
2001	3.1	*	*	0.4	0.6	0.4	0.4	1.1	3.7	10.6	22.6	47.5
2000	3.1	*	*	0.5	0.6	0.3	0.5	0.9	3.7	11.0	23.9	46.2
1999	3.2	*	*	0.4	0.6	0.4	0.4	1.1	4.0	11.4	24.2	48.9
Non-Hispanic white, female												
2002	2.2	*	*	0.4	0.4	0.3	0.3	0.6	1.6	4.2	11.2	26.4
2001	2.2	*	*	0.3	0.3	0.3	0.3	0.5	1.5	4.5	11.7	28.0
2000	2.3	*	*	0.3	0.2	0.2	0.2	0.5	1.5	4.8	11.9	28.9
1999	2.3	*	*	0.3	0.2	0.2	0.2	0.5	1.7	5.4	12.0	26.2
Non-Hispanic black, both sexes												
2002	1.4	*	*	*	0.4	*	*	0.8	3.4	6.4	14.1	18.5
2001	1.5	*	*	0.4	0.4	*	*	1.3	3.2	7.7	13.4	18.4
2000	1.3	*	*	*	0.4	*	*	1.1	2.4	6.7	11.9	25.4
1999	1.4	*	*	0.3	0.5	*	0.4	0.9	3.4	7.5	12.6	22.0
Non-Hispanic black, male												
2002	1.6	*	*	*	*	*	*	1.0	4.5	9.1	20.8	31.7
2001	1.7	*	*	*	*	*	*	1.5	4.7	11.1	19.2	25.4
2000	1.6	*	*	*	*	*	*	1.5	4.1	10.3	18.5	41.1
1999	1.8	*	*	*	*	*	*	1.3	4.7	11.6	17.8	33.3
Non-Hispanic black, female												
2002	1.2	*	*	*	*	*	*	*	2.6	4.6	10.3	13.4
2001	1.3	*	*	*	*	*	*	1.1	2.0	5.3	10.1	15.8
2000	1.1	*	*	*	*	*	*	*	*	4.2	8.2	19.6
1999	1.2	*	*	*	*	*	*	*	2.4	4.7	9.6	17.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myeloid leukemia (C92)												
All origins 3/, both sexes												
2002	3.4	*	0.3	0.2	0.4	0.6	1.0	2.0	5.4	13.4	23.8	25.7
2001	3.5	*	0.5	0.2	0.5	0.7	1.1	2.3	5.4	13.6	23.6	25.1
2000	3.5	*	0.4	0.2	0.5	0.7	1.1	2.2	5.4	13.4	23.8	25.3
1999	3.3	*	0.4	0.2	0.5	0.7	1.1	2.2	5.5	12.9	21.5	25.6
All origins 3/, male												
2002	3.9	*	0.4	0.2	0.4	0.7	1.0	2.2	6.6	17.8	33.7	37.6
2001	3.9	*	0.5	0.2	0.5	0.8	1.1	2.4	6.7	17.5	32.6	37.2
2000	3.9	*	0.4	0.2	0.6	0.8	1.1	2.5	6.4	17.8	33.2	39.1
1999	3.7	*	0.4	0.2	0.5	0.7	1.1	2.4	6.4	17.0	29.4	37.7
All origins 3/, female												
2002	3.0	*	*	0.2	0.4	0.5	0.9	1.8	4.3	9.7	17.3	20.5
2001	3.1	*	0.5	0.2	0.5	0.6	1.0	2.2	4.3	10.4	17.6	20.1
2000	3.1	*	0.3	0.2	0.4	0.6	1.2	1.9	4.5	9.9	17.7	19.7
1999	3.0	*	0.3	0.2	0.5	0.6	1.0	2.0	4.6	9.5	16.3	20.7
Hispanic, both sexes												
2002	1.2	*	*	*	0.5	0.5	1.0	1.4	3.5	8.6	12.0	15.5
2001	1.4	*	*	*	0.4	0.7	1.1	1.8	3.6	8.9	15.0	16.3
2000	1.4	*	*	*	0.5	0.5	1.2	1.8	4.9	9.2	15.6	16.6
1999	1.4	*	*	0.4	0.4	0.6	1.1	2.1	3.7	9.7	15.0	14.0
Hispanic, male												
2002	1.3	*	*	*	0.6	0.6	1.1	1.4	3.5	10.9	16.4	*
2001	1.5	*	*	*	0.6	0.9	1.2	2.0	5.3	10.0	17.0	*
2000	1.5	*	*	*	*	0.7	1.1	2.4	6.0	12.0	20.2	*
1999	1.4	*	*	*	0.6	0.7	0.9	2.0	4.0	13.6	18.8	*
Hispanic, female												
2002	1.1	*	*	*	*	*	0.9	1.5	3.5	6.7	9.0	*
2001	1.2	*	*	*	*	*	0.9	1.7	2.1	8.0	13.6	*
2000	1.2	*	*	*	*	*	1.2	*	3.9	7.0	12.5	*
1999	1.4	*	*	*	*	*	1.4	2.2	3.3	6.7	12.5	*
Non-Hispanic 4/, both sexes												
2002	3.8	*	0.3	0.2	0.4	0.7	1.0	2.1	5.5	13.7	24.3	26.0
2001	3.8	*	0.5	0.2	0.5	0.7	1.1	2.3	5.6	13.9	23.9	25.4
2000	3.8	*	0.4	0.2	0.5	0.7	1.1	2.3	5.4	13.7	24.2	25.5
1999	3.6	*	0.3	0.2	0.5	0.7	1.1	2.2	5.6	13.1	21.7	26.0
Non-Hispanic 4/, male												
2002	4.3	*	0.4	0.2	0.4	0.8	1.0	2.3	6.8	18.3	34.4	37.9
2001	4.2	*	0.4	0.2	0.4	0.8	1.1	2.4	6.8	18.0	33.3	37.5
2000	4.2	*	0.4	0.2	0.6	0.8	1.1	2.5	6.4	18.1	33.7	39.4
1999	4.0	*	0.3	0.2	0.5	0.7	1.1	2.5	6.6	17.2	29.8	38.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myeloid leukemia (C92)												
Non-Hispanic 4/, female												
2002	3.2	*	*	0.2	0.4	0.5	0.9	1.9	4.3	9.9	17.7	20.9
2001	3.4	*	0.5	0.2	0.5	0.6	1.0	2.2	4.4	10.5	17.8	20.4
2000	3.3	*	0.3	0.2	0.3	0.7	1.2	2.0	4.6	10.1	17.9	19.9
1999	3.2	*	0.4	0.2	0.5	0.6	1.0	2.0	4.7	9.7	16.4	20.8
Non-Hispanic white, both sexes												
2002	4.2	*	0.3	0.2	0.4	0.6	1.0	2.1	5.6	14.4	25.3	26.9
2001	4.2	*	0.5	0.2	0.5	0.7	1.0	2.3	5.8	14.5	24.8	26.3
2000	4.2	*	0.4	0.2	0.4	0.6	1.1	2.2	5.4	14.4	25.2	26.3
1999	4.0	*	0.4	0.2	0.4	0.6	1.1	2.2	5.8	13.5	22.4	27.1
Non-Hispanic white, male												
2002	4.8	*	*	0.2	0.4	0.7	1.0	2.3	6.9	19.1	35.8	39.0
2001	4.7	*	0.4	0.2	0.4	0.7	1.0	2.4	7.0	18.5	34.7	39.4
2000	4.7	*	0.4	0.2	0.5	0.7	1.0	2.5	6.4	18.9	35.4	41.4
1999	4.4	*	*	0.2	0.5	0.6	1.1	2.4	6.7	17.7	30.8	40.7
Non-Hispanic white, female												
2002	3.6	*	*	0.2	0.4	0.5	0.9	1.9	4.3	10.4	18.3	21.7
2001	3.8	*	0.6	0.2	0.5	0.6	1.0	2.2	4.6	11.0	18.3	20.8
2000	3.6	*	*	0.2	0.3	0.6	1.1	2.0	4.5	10.6	18.6	20.2
1999	3.5	*	0.4	0.2	0.4	0.7	1.0	2.0	4.8	10.0	16.9	21.7
Non-Hispanic black, both sexes												
2002	2.2	*	*	*	0.4	0.8	0.9	2.2	5.7	10.3	16.6	19.7
2001	2.2	*	*	*	0.4	0.8	1.4	2.2	4.5	11.5	16.8	17.5
2000	2.4	*	*	*	0.8	1.1	1.5	2.9	6.1	10.0	14.6	19.7
1999	2.2	*	*	0.3	0.8	0.7	1.2	2.4	5.0	11.1	15.8	14.5
Non-Hispanic black, male												
2002	2.4	*	*	*	*	0.9	1.1	2.6	7.0	14.2	21.4	32.8
2001	2.4	*	*	*	*	1.0	1.6	2.2	5.9	15.9	20.7	*
2000	2.5	*	*	*	1.0	1.4	1.4	3.2	7.2	13.2	18.5	23.5
1999	2.4	*	*	*	0.8	0.9	1.2	3.1	6.5	14.4	20.6	*
Non-Hispanic black, female												
2002	2.0	*	*	*	*	0.7	0.8	1.9	4.7	7.6	13.9	14.6
2001	2.0	*	*	*	*	*	1.2	2.2	3.3	8.4	14.7	16.7
2000	2.2	*	*	*	*	0.9	1.5	2.7	5.3	7.7	12.3	18.3
1999	2.0	*	*	*	0.8	*	1.1	1.8	3.8	8.7	13.1	12.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Monocytic leukemia (C93)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.7
2000	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	0.7
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.9
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.8	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.7
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.5	0.9
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.8	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Monocytic leukemia (C93)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.0
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.8	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified leukemia (C94-C95)												
All origins 3/, both sexes												
2002	1.8	*	*	0.1	0.2	0.2	0.3	0.7	1.9	6.1	14.0	24.1
2001	1.8	*	0.2	0.2	0.2	0.2	0.3	0.7	2.0	5.9	14.2	24.4
2000	1.9	*	0.2	0.1	0.2	0.2	0.3	0.7	2.0	6.5	14.6	25.6
1999	1.9	*	*	0.2	0.3	0.2	0.3	0.7	2.3	6.5	14.9	25.2
All origins 3/, male												
2002	2.0	*	*	0.2	0.3	0.3	0.3	0.8	2.3	8.2	19.3	32.6
2001	2.0	*	0.3	0.2	0.2	0.3	0.3	0.8	2.4	8.0	19.8	35.3
2000	2.0	*	*	0.1	0.2	0.3	0.4	0.8	2.5	8.8	19.5	35.7
1999	2.1	*	*	0.2	0.3	0.2	0.4	0.8	2.7	8.8	20.3	34.6
All origins 3/, female												
2002	1.6	*	*	0.1	0.2	0.2	0.3	0.6	1.6	4.4	10.5	20.4
2001	1.6	*	*	0.1	0.2	0.2	0.2	0.5	1.7	4.3	10.5	19.8
2000	1.7	*	*	0.1	0.2	0.2	0.3	0.7	1.6	4.5	11.4	21.4
1999	1.8	*	*	0.1	0.2	0.2	0.3	0.6	1.9	4.6	11.5	21.3
Hispanic, both sexes												
2002	0.7	*	*	*	0.4	0.3	*	*	2.1	4.3	9.3	21.1
2001	0.7	*	*	*	*	*	*	0.7	1.4	4.5	12.3	16.9
2000	0.7	*	*	*	*	*	0.4	0.6	1.6	4.0	10.3	19.2
1999	0.6	*	*	*	*	*	*	*	1.6	4.6	8.1	14.0
Hispanic, male												
2002	0.8	*	*	*	*	*	*	*	2.1	4.6	14.0	*
2001	0.8	*	*	*	*	*	*	*	*	5.2	20.1	*
2000	0.7	*	*	*	*	*	*	*	*	*	13.8	*
1999	0.7	*	*	*	*	*	*	*	*	6.2	*	*
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	2.0	4.1	6.2	19.9
2001	0.7	*	*	*	*	*	*	*	*	3.9	7.0	*
2000	0.6	*	*	*	*	*	*	*	*	4.3	7.9	*
1999	0.6	*	*	*	*	*	*	*	*	3.4	6.9	*
Non-Hispanic 4/, both sexes												
2002	2.0	*	*	0.1	0.2	0.2	0.3	0.7	1.9	6.2	14.3	24.2
2001	2.0	*	0.2	0.1	0.2	0.2	0.3	0.7	2.1	6.0	14.2	24.6
2000	2.0	*	0.2	0.1	0.2	0.2	0.3	0.7	2.1	6.6	14.8	25.8
1999	2.1	*	*	0.2	0.3	0.2	0.3	0.7	2.3	6.6	15.2	25.5
Non-Hispanic 4/, male												
2002	2.2	*	*	0.2	0.2	0.2	0.3	0.8	2.3	8.4	19.6	33.0
2001	2.2	*	*	0.2	0.2	0.3	0.3	0.8	2.5	8.1	19.6	35.8
2000	2.2	*	*	0.1	0.2	0.3	0.3	0.7	2.5	9.1	19.7	36.0
1999	2.3	*	*	0.2	0.3	0.3	0.4	0.8	2.7	8.9	20.7	35.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified leukemia (C94-C95)												
Non-Hispanic 4/, female												
2002	1.8	*	*	*	0.1	0.2	0.3	0.7	1.6	4.3	10.7	20.4
2001	1.7	*	*	*	0.2	0.2	0.2	0.5	1.7	4.3	10.7	20.0
2000	1.9	*	*	*	0.2	0.2	0.3	0.7	1.7	4.5	11.6	21.7
1999	1.9	*	*	0.1	0.2	0.2	0.3	0.6	2.0	4.7	11.6	21.6
Non-Hispanic white, both sexes												
2002	2.2	*	*	0.1	0.2	0.2	0.3	0.7	1.9	6.4	14.7	24.4
2001	2.2	*	0.3	0.1	0.2	0.2	0.2	0.7	2.0	6.1	14.7	25.3
2000	2.3	*	0.3	0.1	0.2	0.2	0.3	0.7	2.1	6.8	15.2	26.5
1999	2.3	*	*	0.1	0.3	0.2	0.3	0.7	2.4	6.7	15.5	26.1
Non-Hispanic white, male												
2002	2.4	*	*	*	0.3	0.2	0.3	0.7	2.3	8.5	20.2	34.5
2001	2.4	*	*	0.2	0.2	0.3	0.3	0.8	2.5	8.3	20.1	36.9
2000	2.4	*	*	*	0.2	0.3	0.3	0.7	2.5	9.4	20.1	36.6
1999	2.5	*	*	*	0.3	0.2	0.4	0.7	2.8	9.2	21.2	36.1
Non-Hispanic white, female												
2002	2.0	*	*	*	0.2	0.2	0.3	0.6	1.5	4.6	10.9	20.0
2001	1.9	*	*	*	*	*	0.2	0.5	1.6	4.3	11.1	20.5
2000	2.1	*	*	*	0.3	0.2	0.3	0.7	1.7	4.6	11.9	22.4
1999	2.1	*	*	*	0.2	0.2	0.3	0.6	2.0	4.7	11.8	22.1
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	*	*	*	1.3	2.3	5.8	12.0	26.3
2001	1.2	*	*	*	*	*	0.4	0.8	2.4	5.8	11.7	20.9
2000	1.2	*	*	*	*	*	0.4	0.9	2.2	6.2	12.2	20.4
1999	1.2	*	*	*	*	*	*	1.1	2.4	6.3	12.8	19.4
Non-Hispanic black, male												
2002	1.3	*	*	*	*	*	*	1.5	2.6	8.9	15.5	21.9
2001	1.3	*	*	*	*	*	*	*	2.7	7.4	15.9	30.0
2000	1.4	*	*	*	*	*	*	*	2.7	8.8	18.5	32.9
1999	1.2	*	*	*	*	*	*	1.2	2.5	7.2	16.6	25.0
Non-Hispanic black, female												
2002	1.3	*	*	*	*	*	*	1.1	2.1	3.5	9.9	28.0
2001	1.1	*	*	*	*	*	*	*	2.0	4.7	9.3	17.5
2000	1.0	*	*	*	*	*	*	*	1.8	4.4	8.7	15.7
1999	1.3	*	*	*	*	*	*	1.0	2.2	5.7	10.7	17.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms (D00-D09)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms (D00-D09)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Carcinoma in situ of breast and genitourinary system (D05-D07)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Carcinoma in situ of breast and genitourinary system (D05-D07)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Carcinoma in situ of other and unspecified sites (D00-D04,D09)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Carcinoma in situ of other and unspecified sites (D00-D04,D09)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Benign neoplasms (D10-D36)												
All origins 3/, both sexes												
2002	0.6	*	*	*	*	0.1	0.2	0.4	0.6	1.6	3.6	7.8
2001	0.5	*	*	*	0.1	0.1	0.1	0.3	0.7	1.5	3.6	8.7
2000	0.6	0.6	*	*	*	0.1	0.2	0.3	0.6	1.5	4.2	9.0
1999	0.6	*	*	*	*	0.1	0.1	0.3	0.7	1.6	4.0	8.4
All origins 3/, male												
2002	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.6	3.4	6.3
2001	0.5	*	*	*	*	*	0.1	0.3	0.8	1.4	3.9	8.9
2000	0.4	*	*	*	*	*	0.1	0.3	0.4	1.5	3.8	8.7
1999	0.4	*	*	*	*	*	0.2	0.3	0.7	1.6	3.6	9.2
All origins 3/, female												
2002	0.7	*	*	*	*	*	0.2	0.4	0.6	1.6	3.8	8.4
2001	0.6	*	*	*	*	0.1	0.2	0.3	0.7	1.5	3.4	8.6
2000	0.7	*	*	*	*	0.1	0.3	0.3	0.7	1.5	4.4	9.1
1999	0.7	*	*	*	*	*	0.1	0.3	0.8	1.6	4.2	8.1
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	4.0	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	*	*	*	*	0.1	0.2	0.4	0.6	1.6	3.6	7.8
2001	0.6	*	*	*	0.1	0.1	0.1	0.3	0.7	1.5	3.6	8.7
2000	0.6	*	*	*	*	0.1	0.2	0.3	0.6	1.5	4.2	8.9
1999	0.6	*	*	*	*	0.1	0.1	0.3	0.7	1.6	4.1	8.6
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	0.2	0.3	0.6	1.6	3.3	6.2
2001	0.5	*	*	*	*	*	0.1	0.3	0.8	1.4	4.0	9.1
2000	0.5	*	*	*	*	*	0.1	0.3	0.5	1.5	3.8	8.8
1999	0.5	*	*	*	*	*	0.1	0.3	0.7	1.6	3.7	9.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Benign neoplasms (D10-D36)												
Non-Hispanic 4/, female												
2002	0.7	*	*	*	*	*	0.2	0.4	0.6	1.6	3.9	8.6
2001	0.7	*	*	*	*	0.1	0.2	0.3	0.7	1.5	3.4	8.5
2000	0.7	*	*	*	*	*	0.3	0.3	0.7	1.5	4.4	8.9
1999	0.7	*	*	*	*	*	0.1	0.3	0.7	1.6	4.3	8.2
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	0.1	0.2	0.3	0.6	1.5	3.5	7.4
2001	0.6	*	*	*	*	0.1	0.1	0.3	0.6	1.4	3.5	8.2
2000	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	4.1	8.8
1999	0.6	*	*	*	*	*	0.1	0.2	0.6	1.5	3.9	8.6
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	0.2	0.3	0.6	1.6	3.2	5.6
2001	0.5	*	*	*	*	*	*	0.3	0.6	1.3	3.8	8.6
2000	0.5	*	*	*	*	*	*	0.3	0.4	1.3	3.8	8.8
1999	0.5	*	*	*	*	*	*	0.2	0.6	1.5	3.5	9.7
Non-Hispanic white, female												
2002	0.7	*	*	*	*	*	0.2	0.4	0.6	1.5	3.7	8.2
2001	0.7	*	*	*	*	*	0.1	0.3	0.6	1.4	3.3	8.0
2000	0.8	*	*	*	*	*	0.2	0.2	0.7	1.5	4.3	8.8
1999	0.7	*	*	*	*	*	*	0.2	0.7	1.4	4.2	8.2
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	*	0.7	1.1	2.7	5.5	11.5
2001	0.7	*	*	*	*	*	*	0.6	1.8	2.5	5.6	15.3
2000	0.6	*	*	*	*	*	0.5	0.6	0.8	2.7	5.7	10.2
1999	0.7	*	*	*	*	*	*	0.7	1.8	2.8	6.6	8.7
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	2.1	*	6.3	*
2000	0.4	*	*	*	*	*	*	*	*	3.0	*	*
1999	0.6	*	*	*	*	*	*	*	2.1	*	7.5	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	0.9	*	2.8	5.5	11.3
2001	0.8	*	*	*	*	*	*	*	1.5	2.3	5.2	14.5
2000	0.8	*	*	*	*	*	0.7	*	*	2.5	6.4	10.0
1999	0.8	*	*	*	*	*	*	*	*	3.2	6.1	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of female genital organs (D25-D28)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of female genital organs (D25-D28)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of eye, brain and other parts of central nervous system (D31-D33)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.2	0.3	1.0	2.4	5.1
2001	0.3	*	*	*	*	*	0.0	0.1	0.4	1.0	2.3	5.9
2000	0.3	*	*	*	*	*	0.1	0.1	0.3	1.0	2.9	6.2
1999	0.3	*	*	*	*	*	0.1	0.1	0.5	1.0	2.7	5.5
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.3	1.0	2.0	3.2
2001	0.2	*	*	*	*	*	*	*	0.4	0.9	2.2	5.4
2000	0.3	*	*	*	*	*	*	0.2	0.2	0.9	2.5	5.9
1999	0.2	*	*	*	*	*	*	0.1	0.4	1.0	2.4	5.5
All origins 3/, female												
2002	0.4	*	*	*	*	*	0.1	0.2	0.4	1.1	2.7	5.9
2001	0.4	*	*	*	*	*	*	0.2	0.4	1.0	2.4	6.2
2000	0.4	*	*	*	*	*	*	0.1	0.5	1.0	3.1	6.4
1999	0.4	*	*	*	*	*	*	0.1	0.5	1.1	3.0	5.5
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.3	1.1	2.5	5.1
2001	0.4	*	*	*	*	*	*	0.1	0.4	1.0	2.3	6.0
2000	0.4	*	*	*	*	*	0.1	0.1	0.4	1.0	2.9	6.2
1999	0.4	*	*	*	*	*	0.1	0.1	0.5	1.1	2.8	5.7
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	0.2	0.3	1.0	2.0	3.2
2001	0.3	*	*	*	*	*	*	*	0.3	0.9	2.2	5.6
2000	0.3	*	*	*	*	*	*	0.2	0.2	0.9	2.5	5.9
1999	0.3	*	*	*	*	*	*	0.1	0.4	1.0	2.4	5.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of eye, brain and other parts of central nervous system (D31-D33)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.1	2.8	6.0
2001	0.4	*	*	*	*	*	*	0.2	0.4	1.0	2.4	6.1
2000	0.5	*	*	*	*	*	*	0.1	0.5	1.0	3.1	6.4
1999	0.5	*	*	*	*	*	*	0.1	0.5	1.1	3.0	5.6
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.3	1.0	2.3	5.0
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.2	5.6
2000	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.8	6.3
1999	0.4	*	*	*	*	*	*	0.1	0.4	1.0	2.7	5.7
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	0.2	0.3	1.0	1.8	2.9
2001	0.3	*	*	*	*	*	*	*	0.2	0.9	2.1	5.1
2000	0.3	*	*	*	*	*	*	0.2	*	0.8	2.5	6.0
1999	0.3	*	*	*	*	*	*	*	0.3	1.0	2.2	5.7
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.0	2.7	5.9
2001	0.4	*	*	*	*	*	*	0.1	0.4	0.9	2.3	5.8
2000	0.5	*	*	*	*	*	*	*	0.5	1.0	3.0	6.4
1999	0.5	*	*	*	*	*	*	*	0.5	1.0	2.9	5.7
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	1.6	3.9	6.0
2001	0.4	*	*	*	*	*	*	*	1.2	1.8	3.7	10.9
2000	0.3	*	*	*	*	*	*	*	*	1.3	4.0	*
1999	0.4	*	*	*	*	*	*	*	1.1	1.9	4.5	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	4.1	*
2001	0.5	*	*	*	*	*	*	*	*	*	3.9	10.7
2000	0.4	*	*	*	*	*	*	*	*	*	4.9	*
1999	0.5	*	*	*	*	*	*	*	*	2.7	4.2	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.6	1.2	2.6
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.2	2.5
2000	0.2	0.5	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.2	2.8
All origins 3/, male												
2002	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.4	3.1
2001	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.7	3.5
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.3	2.9
1999	0.2	*	*	*	*	*	0.1	0.1	0.4	0.6	1.3	3.7
All origins 3/, female												
2002	0.2	*	*	*	*	*	0.1	0.2	0.2	0.5	1.0	2.4
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.4	0.9	2.1
2000	0.2	*	*	*	*	*	0.1	*	0.2	0.5	1.2	2.6
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	2.4
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.5	1.2	2.6
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.3	2.5
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.5
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.2	2.8
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4	3.0
2001	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.8	3.5
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.3	2.9
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.3	3.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Of other and unspecified sites (D10-D24,D29-D30,D34-D36)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.0	2.4
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.0
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	2.4
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	2.3
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.2	2.4
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	2.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.8
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.4	2.7
2001	0.2	*	*	*	*	*	*	0.1	0.4	0.4	1.7	3.5
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.7
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	4.0
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.2
2001	0.2	*	*	*	*	*	*	*	0.2	0.4	0.9	2.0
2000	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	2.3
1999	0.2	*	*	*	*	*	*	*	*	0.4	1.2	2.4
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	1.4	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)												
All origins 3/, both sexes												
2002	3.8	0.9	0.3	0.2	0.2	0.3	0.5	1.3	3.6	10.8	31.1	67.0
2001	4.0	0.6	0.3	0.2	0.2	0.2	0.6	1.4	4.0	12.0	32.2	70.1
2000	4.0	0.7	0.3	0.2	0.2	0.3	0.6	1.4	4.2	11.8	32.0	72.0
1999	4.0	0.8	0.4	0.2	0.2	0.3	0.6	1.4	3.8	12.5	31.4	70.8
All origins 3/, male												
2002	4.0	*	0.3	0.2	0.2	0.3	0.6	1.4	4.5	13.7	41.3	91.9
2001	4.2	*	0.3	0.2	0.1	0.3	0.7	1.6	4.6	15.3	43.1	96.6
2000	4.2	*	0.4	0.2	0.2	0.3	0.7	1.6	4.9	15.1	42.4	97.6
1999	4.2	*	0.4	0.2	0.2	0.4	0.7	1.6	4.8	16.2	41.4	97.8
All origins 3/, female												
2002	3.7	*	0.3	0.2	0.2	0.3	0.5	1.1	2.8	8.3	24.3	56.2
2001	3.9	*	0.3	0.2	0.2	0.2	0.6	1.2	3.4	9.3	25.0	59.0
2000	3.9	*	*	0.2	0.1	0.3	0.5	1.2	3.5	9.2	25.2	61.5
1999	3.8	*	0.3	0.2	0.2	0.3	0.5	1.2	2.9	9.5	24.9	60.0
Hispanic, both sexes												
2002	1.1	*	*	0.3	*	*	0.4	1.0	2.2	7.4	21.2	40.5
2001	1.2	*	*	*	*	*	0.6	1.1	2.5	7.2	20.4	47.5
2000	1.1	*	*	*	*	*	0.5	0.8	2.7	7.2	19.2	59.1
1999	1.2	*	*	*	*	0.3	0.4	1.0	2.6	9.0	20.2	46.3
Hispanic, male												
2002	1.1	*	*	*	*	*	*	*	2.9	9.0	26.7	50.2
2001	1.2	*	*	*	*	*	*	1.4	2.9	8.6	27.3	62.9
2000	1.1	*	*	*	*	*	*	*	2.7	7.8	21.6	82.6
1999	1.1	*	*	*	*	*	*	1.5	2.7	9.0	22.9	49.6
Hispanic, female												
2002	1.2	*	*	*	*	*	*	*	*	6.1	17.4	35.6
2001	1.2	*	*	*	*	*	*	*	2.2	6.1	15.7	40.0
2000	1.2	*	*	*	*	*	*	*	2.8	6.6	17.5	47.5
1999	1.3	*	*	*	*	*	*	*	2.5	8.9	18.4	44.7
Non-Hispanic 4/, both sexes												
2002	4.2	0.8	0.3	0.1	0.2	0.3	0.5	1.3	3.7	11.0	31.5	68.0
2001	4.4	*	0.3	0.2	0.2	0.2	0.6	1.4	4.1	12.3	32.6	70.8
2000	4.4	0.8	0.3	0.2	0.2	0.3	0.6	1.5	4.3	12.1	32.5	72.2
1999	4.4	0.9	0.4	0.2	0.2	0.3	0.6	1.4	3.9	12.7	31.8	71.5
Non-Hispanic 4/, male												
2002	4.5	*	*	0.2	0.2	0.3	0.6	1.5	4.6	14.0	41.9	93.6
2001	4.6	*	*	0.2	0.2	0.3	0.7	1.6	4.7	15.7	43.7	97.9
2000	4.6	*	0.4	0.2	0.2	0.3	0.7	1.7	5.1	15.5	43.2	97.8
1999	4.6	*	0.4	0.2	0.2	0.4	0.7	1.7	4.9	16.6	42.1	99.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)												
Non-Hispanic 4/, female												
2002	4.0	*	0.3	0.1	0.2	0.3	0.5	1.1	2.9	8.4	24.6	57.0
2001	4.3	*	*	0.2	0.2	0.2	0.6	1.2	3.5	9.5	25.3	59.6
2000	4.3	*	*	0.2	0.1	0.3	0.5	1.2	3.5	9.4	25.5	61.9
1999	4.1	*	0.3	0.2	0.2	0.2	0.5	1.3	2.9	9.5	25.1	60.3
Non-Hispanic white, both sexes												
2002	4.8	0.9	0.3	0.1	0.2	0.3	0.5	1.3	3.8	11.4	32.5	70.3
2001	5.0	*	0.3	0.2	0.1	0.3	0.6	1.5	4.2	12.8	33.7	72.5
2000	5.0	*	0.4	0.2	0.2	0.3	0.6	1.4	4.4	12.4	33.7	74.5
1999	4.9	*	0.4	0.2	0.2	0.3	0.6	1.4	3.9	13.1	32.8	74.1
Non-Hispanic white, male												
2002	5.1	*	*	*	0.2	0.3	0.7	1.5	4.6	14.6	43.1	97.2
2001	5.3	*	*	0.2	*	0.3	0.7	1.7	4.8	16.4	45.0	100.7
2000	5.2	*	0.4	0.2	0.2	0.3	0.7	1.7	5.1	15.7	44.8	101.4
1999	5.2	*	*	0.2	0.2	0.3	0.7	1.7	5.0	17.1	43.6	104.8
Non-Hispanic white, female												
2002	4.6	*	*	*	0.2	0.2	0.4	1.1	3.0	8.7	25.4	58.7
2001	4.8	*	*	0.2	0.2	0.2	0.6	1.2	3.6	9.7	26.2	60.8
2000	4.8	*	*	0.2	*	0.3	0.5	1.2	3.6	9.6	26.4	63.7
1999	4.7	*	*	0.2	0.2	0.3	0.5	1.2	2.9	9.7	25.7	61.9
Non-Hispanic black, both sexes												
2002	2.3	*	*	*	*	0.4	0.7	1.5	4.4	9.5	25.2	50.5
2001	2.4	*	*	0.4	*	*	0.8	1.3	3.9	10.6	25.4	54.9
2000	2.4	*	*	*	*	0.4	0.5	2.0	4.3	11.6	23.6	51.8
1999	2.5	*	*	*	*	*	0.8	1.7	4.2	12.4	25.1	49.5
Non-Hispanic black, male												
2002	2.3	*	*	*	*	*	*	1.7	5.5	11.3	33.1	67.8
2001	2.3	*	*	*	*	*	0.9	1.3	4.4	13.2	35.1	70.5
2000	2.5	*	*	*	*	*	0.8	2.2	5.5	15.1	29.9	64.6
1999	2.4	*	*	*	*	*	0.8	1.7	4.7	16.1	29.4	53.5
Non-Hispanic black, female												
2002	2.4	*	*	*	*	*	0.8	1.4	3.4	8.3	20.7	43.9
2001	2.4	*	*	*	*	*	0.8	1.3	3.5	8.8	19.9	49.2
2000	2.3	*	*	*	*	*	*	1.8	3.3	9.1	20.0	47.1
1999	2.5	*	*	*	*	*	0.8	1.7	3.8	9.8	22.8	48.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.4	1.4	3.8
2001	0.2	*	*	*	*	*	0.0	0.1	0.2	0.5	1.2	3.8
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	3.9
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.5
All origins 3/, male												
2002	0.2	*	*	*	*	*	0.1	*	0.2	0.5	1.1	3.0
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.1	1.8
2000	0.2	*	*	*	*	*	*	*	0.3	0.4	1.3	2.4
1999	0.1	*	*	*	*	*	*	*	0.2	0.5	1.0	2.1
All origins 3/, female												
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.4	1.5	4.2
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.5	1.2	4.6
2000	0.3	*	*	*	*	*	*	0.1	0.3	0.5	1.5	4.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	4.1
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.1	0.3	0.5	1.4	3.8
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.2	3.8
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	4.0
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.6
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	0.1	*	0.2	0.5	1.1	3.1
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.1	1.8
2000	0.2	*	*	*	*	*	*	*	0.3	0.4	1.3	2.5
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.0	2.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neoplasms of uncertain or unknown behavior of unspecified sites (D48)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.4	1.5	4.1
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.6	1.3	4.6
2000	0.3	*	*	*	*	*	*	*	0.3	0.5	1.5	4.5
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.2	4.2
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	3.9
2001	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.2	4.0
2000	0.3	*	*	*	*	*	*	0.1	0.3	0.4	1.4	4.2
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.5
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	3.2
2001	0.2	*	*	*	*	*	*	0.2	*	0.5	1.2	1.9
2000	0.2	*	*	*	*	*	*	*	0.2	0.3	1.3	2.7
1999	0.2	*	*	*	*	*	*	*	0.3	0.5	1.0	2.3
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.4	1.6	4.2
2001	0.3	*	*	*	*	*	*	0.2	0.3	0.5	1.3	4.9
2000	0.3	*	*	*	*	*	*	*	0.3	0.5	1.4	4.7
1999	0.3	*	*	*	*	*	*	*	*	0.4	1.2	3.9
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)												
All origins 3/, both sexes												
2002	3.4	2.0	0.5	0.2	0.5	0.8	1.5	2.4	3.6	7.6	18.4	56.8
2001	3.3	2.6	0.5	0.2	0.4	0.8	1.5	2.3	3.6	7.3	18.9	58.5
2000	3.3	1.8	0.5	0.2	0.4	0.8	1.4	2.3	3.7	7.6	18.8	58.9
1999	3.2	2.5	0.5	0.2	0.5	0.8	1.4	2.1	3.3	7.3	18.2	60.3
All origins 3/, male												
2002	2.9	2.5	0.5	0.2	0.5	0.9	1.4	2.3	3.7	7.7	18.9	57.2
2001	2.8	2.9	0.6	0.2	0.4	0.9	1.5	2.2	3.3	7.2	20.0	58.1
2000	2.8	2.1	0.5	0.3	0.4	0.8	1.4	2.3	3.5	8.1	19.1	60.4
1999	2.8	3.0	0.6	0.2	0.5	0.9	1.5	2.1	3.1	7.5	19.3	63.2
All origins 3/, female												
2002	3.8	1.5	0.5	0.2	0.4	0.7	1.5	2.5	3.5	7.5	18.1	56.6
2001	3.9	2.4	0.3	0.2	0.4	0.7	1.5	2.3	3.8	7.5	18.2	58.7
2000	3.8	1.6	0.6	0.2	0.4	0.8	1.5	2.2	3.8	7.2	18.6	58.3
1999	3.7	1.9	0.4	0.2	0.5	0.8	1.4	2.1	3.5	7.1	17.4	59.2
Hispanic, both sexes												
2002	1.3	2.5	*	*	0.4	0.4	0.6	1.5	2.7	8.3	15.0	46.6
2001	1.3	2.4	*	*	*	0.5	0.8	1.2	3.1	5.9	18.9	41.9
2000	1.2	*	*	*	*	0.4	0.9	1.6	3.0	6.4	15.0	27.9
1999	1.2	*	*	*	*	0.4	0.9	1.5	2.8	6.8	13.7	47.7
Hispanic, male												
2002	1.3	*	*	*	*	*	0.7	1.6	3.1	8.0	16.4	73.6
2001	1.2	*	*	*	*	*	1.1	*	3.5	5.6	23.3	*
2000	1.0	*	*	*	*	*	0.8	1.9	*	7.6	16.2	*
1999	1.2	*	*	*	*	*	1.2	1.4	*	9.0	15.1	56.1
Hispanic, female												
2002	1.3	*	*	*	*	*	*	1.3	2.4	8.5	14.1	33.2
2001	1.3	*	*	*	*	*	*	1.4	2.9	6.1	16.0	44.7
2000	1.3	*	*	*	*	*	1.0	1.3	4.1	5.5	14.2	28.7
1999	1.2	*	*	*	*	*	*	1.5	3.1	5.0	12.8	43.7
Non-Hispanic 4/, both sexes												
2002	3.7	1.8	0.5	0.2	0.5	0.9	1.6	2.5	3.7	7.5	18.5	57.1
2001	3.6	2.6	0.5	0.2	0.4	0.9	1.6	2.4	3.6	7.4	18.8	59.0
2000	3.6	1.9	0.6	0.2	0.4	0.9	1.5	2.3	3.7	7.6	18.9	59.9
1999	3.5	2.5	0.5	0.3	0.6	0.9	1.5	2.1	3.4	7.3	18.3	60.6
Non-Hispanic 4/, male												
2002	3.1	2.3	0.5	0.2	0.6	1.1	1.5	2.4	3.7	7.6	18.9	56.5
2001	3.1	2.8	0.6	0.2	0.5	1.0	1.5	2.4	3.3	7.3	19.8	58.9
2000	3.1	2.2	0.6	0.3	0.4	0.8	1.5	2.3	3.6	8.0	19.2	61.8
1999	3.0	3.1	0.6	0.3	0.6	0.9	1.5	2.2	3.2	7.4	19.5	63.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)												
Non-Hispanic 4/, female												
2002	4.2	1.4	0.5	0.2	0.4	0.8	1.7	2.6	3.6	7.4	18.2	57.4
2001	4.2	2.4	0.3	0.2	0.4	0.8	1.7	2.4	3.9	7.5	18.2	59.0
2000	4.1	1.6	0.6	0.2	0.4	0.9	1.5	2.3	3.8	7.3	18.7	59.2
1999	4.0	1.9	0.4	0.2	0.5	0.9	1.4	2.1	3.5	7.2	17.5	59.5
Non-Hispanic white, both sexes												
2002	3.4	1.5	0.4	0.1	0.2	0.4	0.7	1.4	2.8	6.8	17.9	56.4
2001	3.4	2.1	0.4	0.2	0.2	0.4	0.7	1.3	2.8	6.8	18.2	58.7
2000	3.4	1.7	0.4	0.1	0.2	0.3	0.7	1.3	2.9	7.0	18.5	59.8
1999	3.2	1.7	0.3	0.2	0.3	0.3	0.6	1.3	2.5	6.6	18.0	59.8
Non-Hispanic white, male												
2002	2.8	2.2	*	*	0.3	0.5	0.8	1.4	3.0	7.0	18.4	55.3
2001	2.8	2.1	0.5	0.2	0.2	0.4	0.7	1.4	2.7	6.7	19.3	59.7
2000	2.8	1.8	0.5	0.2	0.2	0.3	0.8	1.4	3.0	7.6	18.8	61.2
1999	2.7	2.2	*	0.2	0.3	0.3	0.7	1.4	2.4	6.8	19.0	63.8
Non-Hispanic white, female												
2002	3.9	*	*	0.2	0.2	0.3	0.7	1.4	2.6	6.7	17.6	56.9
2001	3.9	2.1	*	*	0.2	0.4	0.7	1.2	3.0	6.9	17.5	58.3
2000	3.9	*	*	*	0.2	0.2	0.6	1.2	2.8	6.5	18.3	59.3
1999	3.7	*	*	0.2	0.3	0.3	0.6	1.1	2.6	6.5	17.3	58.3
Non-Hispanic black, both sexes												
2002	6.1	*	*	0.5	1.6	3.6	6.7	9.9	11.5	15.1	28.3	72.9
2001	6.1	3.8	*	0.5	1.6	3.7	7.2	10.3	10.3	14.4	28.3	65.6
2000	6.0	*	1.2	0.6	1.4	4.1	6.5	10.0	11.2	14.0	26.9	67.7
1999	5.9	5.5	1.4	0.6	1.9	4.2	6.8	8.7	10.9	13.9	24.9	71.5
Non-Hispanic black, male												
2002	5.6	*	*	*	2.0	4.3	6.3	9.6	11.5	15.5	28.7	79.9
2001	5.4	*	*	*	1.8	4.5	6.9	9.8	9.2	14.5	29.4	53.1
2000	5.2	*	*	0.7	1.3	3.9	6.2	10.1	10.0	13.2	25.6	75.1
1999	5.5	*	*	0.7	2.1	4.5	6.9	8.5	10.5	14.0	27.2	63.0
Non-Hispanic black, female												
2002	6.6	*	*	*	1.2	2.9	7.1	10.2	11.5	14.9	28.0	70.2
2001	6.6	*	*	*	1.4	3.0	7.4	10.8	11.1	14.3	27.6	70.2
2000	6.6	*	*	*	1.5	4.2	6.8	10.0	12.1	14.5	27.6	65.0
1999	6.3	*	*	*	1.6	3.8	6.7	8.9	11.2	13.8	23.6	74.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All origins 3/, both sexes												
2002	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
All origins 3/, male												
2002	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
All origins 3/, female												
2002	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	*	*	3.1	7.3	34.4
2001	0.5	*	*	*	*	*	*	*	*	2.2	7.8	28.8
2000	0.4	*	*	*	*	*	*	*	*	2.2	7.1	17.3
1999	0.5	*	*	*	*	*	*	*	*	2.6	7.3	35.1
Hispanic, male												
2002	0.5	*	*	*	*	*	*	*	*	*	*	53.5
2001	0.4	*	*	*	*	*	*	*	*	*	9.4	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	3.0	7.3	24.9
2001	0.6	*	*	*	*	*	*	*	*	*	6.6	32.6
2000	0.5	*	*	*	*	*	*	*	*	*	6.9	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	32.2
Non-Hispanic 4/, both sexes												
2002	1.8	*	*	0.1	0.3	0.5	0.5	0.7	1.0	2.7	9.2	40.5
2001	1.8	*	*	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.5
2000	1.8	*	0.3	0.1	0.2	0.4	0.5	0.6	1.1	2.8	9.5	43.2
1999	1.8	*	0.2	0.1	0.4	0.4	0.5	0.7	1.0	2.6	9.6	44.0
Non-Hispanic 4/, male												
2002	1.4	*	*	0.1	0.4	0.6	0.4	0.6	1.0	2.9	9.0	37.6
2001	1.4	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	10.0	40.8
2000	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	2.8	9.6	41.0
1999	1.4	*	*	0.1	0.4	0.5	0.5	0.7	1.0	2.6	10.1	43.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Non-Hispanic 4/, female												
2002	2.1	*	*	0.2	0.2	0.4	0.5	0.7	1.0	2.6	9.3	41.8
2001	2.2	*	*	0.2	0.3	0.4	0.6	0.6	1.0	2.6	9.7	43.1
2000	2.2	*	*	*	0.3	0.4	0.5	0.6	1.0	2.8	9.5	44.1
1999	2.1	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.3	44.2
Non-Hispanic white, both sexes												
2002	1.6	*	*	*	0.1	0.1	0.1	0.3	0.7	2.4	8.7	39.9
2001	1.7	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	42.1
2000	1.7	*	*	*	*	*	0.1	0.3	0.8	2.5	9.3	42.9
1999	1.6	*	*	*	0.1	*	0.1	0.3	0.6	2.4	9.4	43.2
Non-Hispanic white, male												
2002	1.2	*	*	*	*	0.2	*	0.3	0.7	2.5	8.5	36.3
2001	1.3	*	*	*	*	*	*	0.3	0.7	2.6	9.8	41.2
2000	1.2	*	*	*	*	*	0.1	0.3	0.9	2.7	9.3	39.9
1999	1.2	*	*	*	*	*	*	0.3	0.5	2.4	9.9	43.4
Non-Hispanic white, female												
2002	2.1	*	*	*	*	*	*	0.3	0.7	2.4	8.8	41.4
2001	2.1	*	*	*	*	*	0.2	0.3	0.8	2.4	9.2	42.4
2000	2.1	*	*	*	*	*	*	0.3	0.7	2.4	9.4	44.1
1999	2.0	*	*	*	*	*	*	0.3	0.7	2.4	9.1	43.1
Non-Hispanic black, both sexes												
2002	2.9	*	*	0.4	1.3	2.5	2.5	3.1	3.7	6.1	16.2	52.9
2001	2.8	*	*	0.4	1.2	2.3	2.8	3.1	3.0	5.5	16.1	49.3
2000	2.7	*	*	0.5	1.1	2.4	2.5	3.1	3.5	5.5	13.4	51.8
1999	2.9	*	0.9	0.5	1.5	2.7	2.6	3.2	4.3	5.0	13.4	53.0
Non-Hispanic black, male												
2002	2.7	*	*	*	1.7	2.9	2.5	2.6	4.0	7.5	17.0	58.0
2001	2.6	*	*	*	1.4	3.0	2.5	3.6	3.2	6.8	15.3	42.7
2000	2.4	*	*	*	1.0	2.2	2.1	3.4	3.5	4.6	13.9	56.4
1999	2.7	*	*	*	1.8	2.9	2.8	3.4	5.1	5.6	12.8	44.0
Non-Hispanic black, female												
2002	3.0	*	*	*	1.0	2.0	2.6	3.5	3.4	5.1	15.8	51.0
2001	2.9	*	*	*	1.1	1.6	3.0	2.7	2.9	4.6	16.5	51.8
2000	3.1	*	*	*	1.1	2.5	2.8	2.9	3.5	6.1	13.2	50.2
1999	3.1	*	*	*	1.2	2.4	2.4	3.0	3.6	4.7	13.7	56.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional anemias (D50-D53)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.4	3.5
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	4.3
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	4.0
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	2.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	5.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	3.9
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.8
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.4	3.5
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	4.4
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	4.0
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	2.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000	0.1	*	*	*	*	*	*	*	*	*	*	5.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	3.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional anemias (D50-D53)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.8
2000	0.2	*	*	*	*	*	*	*	*	*	0.7	4.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	4.1
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	3.6
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.7
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	4.4
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.9
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	2.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.4
2000	0.1	*	*	*	*	*	*	*	*	*	*	5.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	4.1
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2001	0.2	*	*	*	*	*	*	*	*	*	0.6	3.8
2000	0.2	*	*	*	*	*	*	*	*	*	0.7	4.2
1999	0.2	*	*	*	*	*	*	*	*	*	0.7	3.9
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hemolytic anemias (D55-D59)												
All origins 3/, both sexes												
2002	0.3	*	*	0.1	0.2	0.4	0.3	0.4	0.2	0.4	0.8	2.2
2001	0.3	*	*	0.1	0.2	0.3	0.3	0.3	0.3	0.3	0.9	2.2
2000	0.3	*	0.2	0.1	0.2	0.3	0.3	0.3	0.2	0.4	0.9	2.4
1999	0.3	*	0.1	0.1	0.2	0.4	0.3	0.3	0.3	0.3	0.9	2.2
All origins 3/, male												
2002	0.3	*	*	*	0.2	0.4	0.3	0.3	0.2	0.3	0.6	2.2
2001	0.3	*	*	*	0.2	0.4	0.3	0.3	0.2	0.3	1.0	2.2
2000	0.3	*	*	*	0.1	0.3	0.3	0.3	0.2	0.3	0.8	2.5
1999	0.3	*	*	*	0.3	0.4	0.3	0.3	0.3	0.3	1.1	1.8
All origins 3/, female												
2002	0.3	*	*	*	0.2	0.3	0.3	0.4	0.2	0.4	0.9	2.2
2001	0.3	*	*	*	0.1	0.3	0.4	0.3	0.3	0.3	0.8	2.2
2000	0.3	*	*	*	0.2	0.3	0.3	0.3	0.3	0.4	0.9	2.3
1999	0.3	*	*	*	0.2	0.3	0.3	0.3	0.3	0.4	0.8	2.4
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	0.1	0.2	0.4	0.4	0.4	0.2	0.4	0.8	2.2
2001	0.3	*	*	0.1	0.2	0.4	0.4	0.3	0.3	0.3	0.9	2.3
2000	0.3	*	0.2	0.1	0.2	0.3	0.3	0.3	0.3	0.4	0.9	2.4
1999	0.3	*	0.2	0.1	0.2	0.4	0.3	0.3	0.3	0.4	0.9	2.3
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.3	0.5	0.3	0.3	0.2	0.4	0.5	2.3
2001	0.3	*	*	*	0.3	0.4	0.3	0.4	0.2	0.3	1.0	2.2
2000	0.3	*	*	*	0.2	0.3	0.3	0.3	0.2	0.3	0.9	2.6
1999	0.3	*	*	*	0.3	0.4	0.4	0.3	0.3	0.3	1.1	1.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hemolytic anemias (D55-D59)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	0.2	0.4	0.4	0.4	0.2	0.4	1.0	2.2
2001	0.3	*	*	*	0.2	0.3	0.4	0.3	0.3	0.4	0.9	2.3
2000	0.4	*	*	*	0.2	0.4	0.4	0.3	0.3	0.4	0.9	2.3
1999	0.4	*	*	*	0.2	0.4	0.3	0.3	0.3	0.4	0.8	2.5
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	*	0.3	0.8	2.3
2001	0.2	*	*	*	*	*	*	*	0.1	0.3	0.9	2.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.9	2.6
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.9	2.5
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.5
2001	0.1	*	*	*	*	*	*	*	*	*	1.1	2.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.9	2.9
1999	0.1	*	*	*	*	*	*	*	*	*	1.1	*
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.2
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	2.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.4
1999	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.7
Non-Hispanic black, both sexes												
2002	1.5	*	*	0.3	1.2	2.3	2.3	2.5	1.4	*	*	*
2001	1.4	*	*	*	1.1	2.0	2.4	2.4	1.5	*	*	*
2000	1.4	*	*	0.4	1.0	2.1	2.2	2.4	1.7	*	*	*
1999	1.5	*	*	0.4	1.3	2.5	2.3	2.3	2.0	*	*	*
Non-Hispanic black, male												
2002	1.5	*	*	*	1.5	2.7	2.3	2.1	*	*	*	*
2001	1.5	*	*	*	1.3	2.7	2.2	2.7	*	*	*	*
2000	1.3	*	*	*	0.9	1.9	2.0	2.5	*	*	*	*
1999	1.7	*	*	*	1.7	2.7	2.6	2.3	2.0	*	*	*
Non-Hispanic black, female												
2002	1.4	*	*	*	0.9	2.0	2.3	2.8	1.5	*	*	*
2001	1.3	*	*	*	0.8	1.4	2.5	2.2	1.5	*	*	*
2000	1.5	*	*	*	1.0	2.2	2.4	2.4	2.0	*	*	*
1999	1.4	*	*	*	0.9	2.3	2.0	2.3	1.9	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aplastic anemias (D60-D61)												
All origins 3/, both sexes												
2002	0.3	*	*	0.1	0.1	0.1	0.1	0.1	0.3	0.8	2.4	5.3
2001	0.4	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.9	2.7	6.8
2000	0.4	*	*	*	0.1	0.1	0.1	0.2	0.4	1.0	2.5	7.4
1999	0.4	*	*	0.1	0.1	*	0.1	0.2	0.3	1.0	2.6	7.7
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.8	7.0
2001	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.9	7.8
2000	0.3	*	*	*	*	*	*	0.1	0.4	1.1	2.8	7.6
1999	0.4	*	*	*	*	*	*	0.2	0.3	1.1	3.1	10.1
All origins 3/, female												
2002	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.1	4.5
2001	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.5	6.3
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.4	7.3
1999	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.3	6.8
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	0.1	0.1	*	0.1	0.1	0.3	0.8	2.4	5.3
2001	0.4	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.9	2.6	6.8
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.6	7.6
1999	0.4	*	*	*	0.1	*	0.1	0.2	0.3	1.0	2.7	7.9
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.8	6.8
2001	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.9	7.9
2000	0.3	*	*	*	*	*	*	0.1	0.4	1.0	2.9	7.8
1999	0.4	*	*	*	*	*	*	0.2	0.3	1.1	3.2	10.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aplastic anemias (D60-D61)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.1	4.6
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.0	2.5	6.3
2000	0.5	*	*	*	*	*	0.1	0.2	0.4	0.9	2.4	7.5
1999	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.3	6.9
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.5	5.2
2001	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.7	7.1
2000	0.4	*	*	*	*	*	0.1	0.1	0.4	1.0	2.6	7.6
1999	0.4	*	*	*	*	*	0.1	0.1	0.3	0.9	2.7	8.0
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	*	0.3	0.8	2.9	6.7
2001	0.4	*	*	*	*	*	*	0.2	0.3	0.8	3.0	8.6
2000	0.4	*	*	*	*	*	*	*	0.4	1.1	2.7	7.3
1999	0.4	*	*	*	*	*	*	*	0.3	1.0	3.2	10.4
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	*	0.3	0.8	2.1	4.6
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.0	2.5	6.4
2000	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.5	7.7
1999	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.3	7.0
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	1.2	*	6.4
2001	0.3	*	*	*	*	*	*	*	*	1.3	2.2	*
2000	0.3	*	*	*	*	*	*	*	*	*	2.4	8.3
1999	0.3	*	*	*	*	*	*	*	1.0	1.3	*	6.8
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute posthemorrhagic and other anemias (D62,D64)												
All origins 3/, both sexes												
2002	0.9	*	*	*	*	*	*	0.1	0.4	1.4	5.5	29.4
2001	0.8	*	*	*	*	*	0.1	0.1	0.3	1.3	5.6	29.5
2000	0.8	*	*	*	*	*	*	0.1	0.4	1.4	5.5	28.2
1999	0.8	*	*	*	*	*	0.0	0.2	0.3	1.2	5.3	29.9
All origins 3/, male												
2002	0.6	*	*	*	*	*	*	0.1	0.5	1.6	5.3	26.3
2001	0.6	*	*	*	*	*	*	*	0.3	1.5	5.4	26.9
2000	0.6	*	*	*	*	*	*	0.1	0.5	1.4	5.5	24.9
1999	0.6	*	*	*	*	*	*	0.2	0.3	1.2	5.2	27.7
All origins 3/, female												
2002	1.1	*	*	*	*	*	*	0.1	0.4	1.2	5.7	30.7
2001	1.1	*	*	*	*	*	*	0.1	0.3	1.1	5.7	30.5
2000	1.1	*	*	*	*	*	*	0.1	0.3	1.4	5.5	29.6
1999	1.1	*	*	*	*	*	*	0.2	0.3	1.2	5.4	30.8
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	4.3	23.8
2001	0.2	*	*	*	*	*	*	*	*	*	4.1	19.4
2000	0.2	*	*	*	*	*	*	*	*	*	5.5	*
1999	0.3	*	*	*	*	*	*	*	*	*	4.6	28.8
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	38.5
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	16.6
2001	0.3	*	*	*	*	*	*	*	*	*	*	22.3
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	29.1
Non-Hispanic 4/, both sexes												
2002	1.0	*	*	*	*	*	*	0.1	0.4	1.4	5.6	29.6
2001	0.9	*	*	*	*	*	0.1	0.1	0.3	1.3	5.6	29.8
2000	0.9	*	*	*	*	*	*	0.1	0.4	1.4	5.5	28.8
1999	0.9	*	*	*	*	*	0.0	0.2	0.3	1.2	5.3	29.8
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	*	0.1	0.5	1.6	5.4	25.7
2001	0.6	*	*	*	*	*	*	*	0.3	1.5	5.3	27.5
2000	0.6	*	*	*	*	*	*	0.1	0.5	1.4	5.4	25.5
1999	0.6	*	*	*	*	*	*	0.2	0.3	1.1	5.1	27.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute posthemorrhagic and other anemias (D62,D64)												
Non-Hispanic 4/, female												
2002	1.2	*	*	*	*	*	*	0.2	0.4	1.2	5.7	31.2
2001	1.2	*	*	*	*	*	*	*	0.3	1.1	5.7	30.8
2000	1.2	*	*	*	*	*	*	*	0.3	1.4	5.5	30.2
1999	1.2	*	*	*	*	*	*	0.2	0.3	1.2	5.4	30.7
Non-Hispanic white, both sexes												
2002	1.0	*	*	*	*	*	*	0.1	0.3	1.2	5.0	28.8
2001	1.0	*	*	*	*	*	*	*	0.2	1.2	5.1	29.0
2000	1.0	*	*	*	*	*	*	0.1	0.3	1.2	5.2	28.3
1999	0.9	*	*	*	*	*	*	0.1	0.2	1.0	5.1	28.8
Non-Hispanic white, male												
2002	0.7	*	*	*	*	*	*	*	0.3	1.3	4.8	24.3
2001	0.7	*	*	*	*	*	*	*	0.2	1.3	5.0	26.9
2000	0.6	*	*	*	*	*	*	*	0.4	1.3	5.2	24.8
1999	0.6	*	*	*	*	*	*	*	*	1.0	5.0	27.1
Non-Hispanic white, female												
2002	1.3	*	*	*	*	*	*	*	0.3	1.1	5.2	30.7
2001	1.3	*	*	*	*	*	*	*	0.2	1.0	5.2	29.9
2000	1.3	*	*	*	*	*	*	*	0.3	1.2	5.2	29.7
1999	1.2	*	*	*	*	*	*	*	0.3	1.0	5.2	29.5
Non-Hispanic black, both sexes												
2002	1.1	*	*	*	*	*	*	*	1.5	3.3	13.0	42.0
2001	1.0	*	*	*	*	*	*	0.5	0.8	2.9	12.0	41.8
2000	0.9	*	*	*	*	*	*	*	1.3	3.4	9.6	37.8
1999	0.9	*	*	*	*	*	*	0.7	1.2	2.8	8.9	39.8
Non-Hispanic black, male												
2002	0.9	*	*	*	*	*	*	*	1.9	4.5	13.8	45.9
2001	0.7	*	*	*	*	*	*	*	*	3.7	11.1	38.1
2000	0.7	*	*	*	*	*	*	*	*	*	8.9	34.1
1999	0.7	*	*	*	*	*	*	*	*	*	8.4	30.9
Non-Hispanic black, female												
2002	1.2	*	*	*	*	*	*	*	*	2.4	12.6	40.6
2001	1.2	*	*	*	*	*	*	*	*	2.4	12.5	43.2
2000	1.1	*	*	*	*	*	*	*	*	3.8	9.9	39.2
1999	1.2	*	*	*	*	*	*	*	*	3.0	9.1	43.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
All origins 3/, both sexes												
2002	0.9	0.6	*	*	0.1	0.2	0.3	0.7	1.1	2.8	6.0	11.3
2001	0.9	0.8	*	*	0.1	0.2	0.4	0.7	1.2	2.7	6.0	11.1
2000	0.9	0.6	*	*	0.1	0.1	0.3	0.7	1.4	2.9	6.1	11.6
1999	0.9	1.1	*	0.1	0.1	0.1	0.4	0.6	1.2	2.9	6.0	11.5
All origins 3/, male												
2002	0.8	*	*	*	*	0.2	0.3	0.6	1.2	2.7	6.7	13.9
2001	0.8	*	*	*	*	0.1	0.3	0.7	1.2	2.8	6.7	12.3
2000	0.8	*	*	*	*	0.1	0.3	0.7	1.2	3.2	6.2	14.6
1999	0.8	1.4	*	*	*	0.1	0.4	0.7	1.1	3.1	6.2	14.3
All origins 3/, female												
2002	1.0	*	*	*	0.1	0.1	0.3	0.7	1.1	2.9	5.6	10.1
2001	1.0	*	*	*	*	0.2	0.4	0.7	1.3	2.6	5.6	10.5
2000	1.0	*	*	*	0.1	0.2	0.3	0.6	1.5	2.7	6.0	10.4
1999	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.8	5.8	10.3
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	0.6	1.1	3.8	4.8	*
2001	0.4	*	*	*	*	*	*	*	1.2	2.4	8.5	*
2000	0.4	*	*	*	*	*	*	0.8	1.3	2.4	6.1	*
1999	0.4	*	*	*	*	*	*	*	1.3	3.3	5.2	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	10.3	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	4.1	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	7.3	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.0	*	*	*	0.1	0.2	0.3	0.7	1.1	2.7	6.0	11.3
2001	1.0	0.9	*	*	0.1	0.2	0.4	0.7	1.2	2.7	5.9	11.0
2000	1.0	*	*	*	0.1	0.1	0.3	0.6	1.4	2.9	6.0	11.7
1999	1.0	1.1	*	*	0.1	0.2	0.4	0.6	1.2	2.9	6.0	11.5
Non-Hispanic 4/, male												
2002	0.9	*	*	*	*	0.2	0.3	0.7	1.2	2.7	6.6	13.8
2001	0.9	*	*	*	*	0.1	0.4	0.7	1.2	2.8	6.5	12.3
2000	0.9	*	*	*	*	*	0.3	0.7	1.3	3.2	6.2	14.9
1999	0.9	1.3	*	*	*	0.1	0.4	0.7	1.1	3.0	6.2	14.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	0.1	0.3	0.7	1.1	2.8	5.6	10.2
2001	1.1	*	*	*	*	0.3	0.4	0.7	1.3	2.6	5.5	10.5
2000	1.1	*	*	*	*	0.2	0.3	0.6	1.4	2.7	5.9	10.4
1999	1.1	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.8	10.3
Non-Hispanic white, both sexes												
2002	1.0	*	*	*	*	0.2	0.3	0.6	1.0	2.6	6.0	11.4
2001	1.0	*	*	*	*	0.2	0.3	0.5	1.1	2.6	5.8	11.0
2000	1.0	*	*	*	*	0.1	0.3	0.5	1.2	2.8	5.9	11.9
1999	1.0	0.9	*	*	0.1	0.1	0.3	0.5	1.1	2.6	5.8	11.5
Non-Hispanic white, male												
2002	0.9	*	*	*	*	0.2	0.3	0.5	1.1	2.6	6.5	14.0
2001	0.9	*	*	*	*	*	0.3	0.5	1.1	2.7	6.3	12.4
2000	0.9	*	*	*	*	*	0.3	0.6	1.2	3.0	6.2	15.3
1999	0.9	*	*	*	*	*	0.3	0.6	0.9	2.8	5.8	14.8
Non-Hispanic white, female												
2002	1.1	*	*	*	*	*	0.3	0.6	0.9	2.6	5.6	10.2
2001	1.1	*	*	*	*	0.2	0.3	0.5	1.2	2.6	5.4	10.4
2000	1.1	*	*	*	*	*	0.3	0.5	1.3	2.6	5.7	10.6
1999	1.1	*	*	*	*	*	0.2	0.4	1.2	2.5	5.8	10.1
Non-Hispanic black, both sexes												
2002	1.0	*	*	*	*	*	0.4	1.6	2.1	4.6	7.5	13.3
2001	1.1	*	*	*	*	0.5	0.9	1.9	2.1	3.6	8.1	12.2
2000	1.1	*	*	*	*	0.4	0.6	1.7	2.6	4.5	8.2	11.4
1999	1.2	*	*	*	*	*	1.0	1.5	2.7	5.0	8.3	13.9
Non-Hispanic black, male												
2002	0.9	*	*	*	*	*	*	1.6	2.0	4.2	8.2	*
2001	1.0	*	*	*	*	*	0.8	1.9	2.0	4.0	9.0	*
2000	1.0	*	*	*	*	*	*	1.8	2.4	5.4	6.8	*
1999	1.1	*	*	*	*	*	0.8	1.4	2.5	5.4	10.9	*
Non-Hispanic black, female												
2002	1.2	*	*	*	*	*	*	1.6	2.2	4.9	7.1	12.5
2001	1.3	*	*	*	*	*	1.0	1.8	2.3	3.2	7.6	13.7
2000	1.3	*	*	*	*	*	*	1.6	2.8	3.9	9.0	10.5
1999	1.3	*	*	*	*	*	1.1	1.5	2.8	4.8	6.8	13.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of blood and blood-forming organs (D70-D76)												
All origins 3/, both sexes												
2002	0.3	*	0.2	*	*	*	0.1	0.2	0.4	0.9	2.0	3.4
2001	0.3	0.6	0.1	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000	0.3	*	*	*	0.1	0.1	0.1	0.2	0.4	0.9	1.8	3.4
1999	0.3	0.6	*	*	0.1	0.1	0.1	0.2	0.4	0.9	1.6	3.6
All origins 3/, male												
2002	0.3	*	0.3	*	*	*	0.1	0.2	0.5	1.1	2.2	3.5
2001	0.3	*	*	*	*	*	*	0.2	0.4	0.8	2.2	3.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	2.2	4.1
1999	0.3	*	*	*	*	*	0.1	0.1	0.5	1.0	2.1	4.0
All origins 3/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.9	3.3
2001	0.3	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6	3.8
2000	0.3	*	*	*	*	*	*	0.2	0.4	0.7	1.6	3.1
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.3	3.4
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.8	3.4
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.7	3.7
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.2	3.5
2001	0.3	*	*	*	*	*	*	0.2	0.4	0.8	2.2	3.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	2.2	4.2
1999	0.3	*	*	*	*	*	0.1	0.1	0.5	1.0	2.2	4.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of blood and blood-forming organs (D70-D76)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.8	3.4
2001	0.3	*	*	*	*	*	0.1	0.1	0.4	0.8	1.7	3.9
2000	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.6	3.1
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.4	3.5
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.9	3.5
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	3.7
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.3	3.4
2001	0.3	*	*	*	*	*	*	0.2	0.4	0.8	2.2	4.0
2000	0.3	*	*	*	*	*	*	0.2	0.3	1.1	2.1	4.3
1999	0.3	*	*	*	*	*	*	*	0.5	1.0	2.2	4.0
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	0.2	0.4	0.8	1.8	3.4
2001	0.4	*	*	*	*	*	*	*	0.4	0.8	1.7	3.9
2000	0.4	*	*	*	*	*	*	0.2	0.4	0.7	1.7	3.2
1999	0.3	*	*	*	*	*	*	0.2	0.3	0.8	1.4	3.7
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	0.9	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain disorders involving the immune mechanism (D80-D89)												
All origins 3/, both sexes												
2002	0.5	0.5	*	*	*	0.2	0.6	0.9	1.0	1.1	1.3	1.8
2001	0.5	0.8	*	*	*	0.2	0.6	0.9	0.9	1.1	1.2	1.5
2000	0.5	*	*	*	*	0.2	0.6	0.8	0.9	0.9	1.5	1.6
1999	0.4	*	0.2	*	*	0.2	0.5	0.7	0.7	0.9	1.0	1.4
All origins 3/, male												
2002	0.5	*	*	*	*	0.2	0.6	0.9	1.0	0.9	1.1	1.5
2001	0.4	*	*	*	*	0.3	0.6	0.8	0.7	0.7	1.1	1.8
2000	0.5	*	*	*	*	0.2	0.6	0.8	0.8	0.9	1.3	1.6
1999	0.4	*	*	*	*	0.3	0.5	0.7	0.6	0.7	1.0	*
All origins 3/, female												
2002	0.6	*	*	*	*	0.1	0.6	0.9	1.1	1.2	1.4	1.9
2001	0.6	*	*	*	*	0.1	0.5	1.0	1.1	1.5	1.3	1.4
2000	0.5	*	*	*	*	0.2	0.6	0.8	0.9	1.0	1.6	1.5
1999	0.4	*	*	*	*	0.2	0.5	0.6	0.8	1.0	1.1	1.5
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	*	*	*	*	0.2	0.7	0.9	1.1	1.1	1.3	1.9
2001	0.5	0.7	*	*	*	0.2	0.6	0.9	1.0	1.2	1.3	1.6
2000	0.5	*	*	*	*	0.2	0.6	0.8	0.9	0.9	1.5	1.5
1999	0.4	*	0.2	*	*	0.2	0.5	0.7	0.7	0.9	1.0	1.5
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	0.2	0.7	0.9	1.0	0.9	1.1	1.6
2001	0.5	*	*	*	*	0.3	0.7	0.8	0.7	0.8	1.2	1.9
2000	0.5	*	*	*	*	0.3	0.7	0.8	0.8	0.9	1.3	*
1999	0.4	*	*	*	*	0.3	0.6	0.7	0.6	0.7	1.0	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain disorders involving the immune mechanism (D80-D89)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	*	0.1	0.7	0.9	1.1	1.3	1.4	2.0
2001	0.6	*	*	*	*	0.1	0.6	1.0	1.2	1.5	1.3	1.5
2000	0.5	*	*	*	*	0.2	0.6	0.9	0.9	0.9	1.7	1.5
1999	0.5	*	*	*	*	0.2	0.5	0.7	0.8	1.1	1.1	1.5
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	*	*	0.2	0.4	0.6	0.9	1.2	1.8
2001	0.3	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.2	1.7
2000	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.4	1.5
1999	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.0	1.5
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	0.3	0.4	0.7	0.8	1.1	*
2001	0.3	*	*	*	*	0.2	0.2	0.4	0.4	0.6	1.0	2.0
2000	0.3	*	*	*	*	*	0.3	0.3	0.6	0.8	1.2	*
1999	0.3	*	*	*	*	*	0.2	0.4	0.5	0.6	1.0	*
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	0.2	0.3	0.6	0.9	1.3	1.9
2001	0.4	*	*	*	*	*	0.1	0.3	0.7	1.1	1.3	1.6
2000	0.3	*	*	*	*	*	0.2	0.3	0.4	0.7	1.5	1.4
1999	0.3	*	*	*	*	*	*	0.2	0.4	0.8	1.0	1.4
Non-Hispanic black, both sexes												
2002	1.9	*	*	*	*	0.7	3.6	4.9	5.1	3.7	2.5	*
2001	1.9	*	*	*	*	0.9	3.4	5.1	4.6	4.3	2.6	*
2000	1.8	*	*	*	*	1.1	3.2	4.9	4.5	2.8	3.6	*
1999	1.5	*	*	*	*	1.1	3.0	3.8	3.0	2.8	*	*
Non-Hispanic black, male												
2002	1.7	*	*	*	*	1.0	3.2	5.0	4.6	*	*	*
2001	1.6	*	*	*	*	1.2	3.5	3.9	3.5	*	*	*
2000	1.6	*	*	*	*	1.1	3.3	4.5	3.4	*	*	*
1999	1.3	*	*	*	*	1.1	2.9	3.4	2.1	*	*	*
Non-Hispanic black, female												
2002	2.1	*	*	*	*	*	4.0	4.9	5.6	4.6	3.3	*
2001	2.2	*	*	*	*	*	3.3	6.1	5.5	5.5	*	*
2000	2.0	*	*	*	*	1.0	3.1	5.1	5.4	3.5	4.2	*
1999	1.7	*	*	*	*	1.0	3.2	4.1	3.7	3.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
All origins 3/, both sexes												
2002	34.5	6.6	0.9	0.5	1.3	3.1	7.4	18.7	47.9	111.1	235.3	504.8
2001	34.0	7.3	1.2	0.5	1.2	2.8	6.7	18.6	46.9	110.8	232.6	519.2
2000	33.5	6.8	1.0	0.5	1.3	2.9	6.6	17.8	46.9	109.9	232.0	521.0
1999	33.3	6.9	1.0	0.5	1.4	2.6	6.5	17.2	46.5	110.5	229.3	529.8
All origins 3/, male												
2002	31.9	7.4	0.8	0.6	1.4	3.6	8.9	22.1	56.1	123.3	255.6	504.5
2001	30.9	8.0	1.2	0.6	1.2	3.0	7.8	22.0	53.6	124.1	251.9	507.7
2000	30.2	7.4	1.1	0.6	1.4	3.4	7.7	20.8	53.1	120.8	251.5	517.4
1999	29.8	7.3	1.1	0.5	1.3	2.9	7.6	20.3	52.3	120.9	250.5	521.0
All origins 3/, female												
2002	37.0	5.8	1.0	0.4	1.2	2.6	5.9	15.5	40.2	100.9	221.9	504.9
2001	37.0	6.4	1.2	0.5	1.3	2.5	5.5	15.3	40.7	99.7	219.8	524.1
2000	36.7	6.2	1.0	0.5	1.2	2.3	5.6	14.8	41.2	100.8	219.4	522.5
1999	36.6	6.5	0.9	0.5	1.4	2.4	5.4	14.2	41.2	102.0	215.4	533.3
Hispanic, both sexes												
2002	18.0	5.4	0.8	0.4	0.7	1.4	4.9	16.8	56.7	147.9	324.3	554.3
2001	18.0	5.6	1.0	0.4	0.7	1.5	4.4	17.3	58.6	153.7	316.3	620.9
2000	17.6	5.2	0.8	0.4	0.6	1.6	4.3	16.7	60.3	161.0	313.3	595.9
1999	17.8	5.7	0.7	0.3	0.6	1.2	4.3	16.3	61.6	161.8	321.2	648.3
Hispanic, male												
2002	16.3	4.9	*	0.6	0.8	1.6	5.7	19.3	62.8	158.6	346.3	543.8
2001	16.1	5.3	*	*	0.7	1.8	5.4	20.9	64.8	164.5	325.1	558.0
2000	16.0	*	*	*	0.6	2.2	4.9	18.4	70.1	176.2	330.4	560.3
1999	15.6	7.0	*	*	*	1.2	4.8	19.9	67.8	168.9	334.7	590.9
Hispanic, female												
2002	19.7	5.9	*	*	*	1.3	4.1	14.3	51.1	139.3	309.2	559.5
2001	20.1	6.0	*	*	0.7	1.2	3.3	13.8	53.2	145.2	310.4	651.6
2000	19.4	6.1	*	*	*	0.9	3.6	15.0	51.7	149.0	301.8	613.3
1999	20.1	*	*	*	*	1.1	3.8	12.7	56.2	156.3	312.2	675.9
Non-Hispanic 4/, both sexes												
2002	36.9	6.9	0.9	0.6	1.4	3.5	7.8	18.9	47.0	108.2	230.1	501.4
2001	36.3	7.6	1.2	0.6	1.3	3.0	7.0	18.6	45.8	107.5	228.1	514.2
2000	35.7	7.2	1.1	0.5	1.4	3.1	6.9	17.8	45.7	106.3	227.9	516.7
1999	35.3	7.1	1.0	0.6	1.5	2.9	6.7	17.2	45.2	107.2	224.9	524.2
Non-Hispanic 4/, male												
2002	34.3	8.0	0.8	0.6	1.5	4.1	9.4	22.3	55.4	120.5	250.1	501.0
2001	33.1	8.7	1.2	0.6	1.3	3.3	8.2	22.0	52.6	121.0	247.8	504.3
2000	32.3	8.0	1.1	0.6	1.5	3.6	8.0	21.0	51.7	117.1	247.1	514.0
1999	31.8	7.2	1.3	0.6	1.4	3.2	7.9	20.2	51.0	117.8	246.4	516.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IV. Endocrine, nutritional and metabolic diseases (E00-E88)												
Non-Hispanic 4/, female												
2002	39.4	5.8	1.1	0.5	1.4	2.9	6.2	15.6	39.2	97.9	216.9	501.6
2001	39.3	6.5	1.3	0.5	1.3	2.7	5.8	15.4	39.6	96.3	215.1	518.4
2000	38.9	6.2	1.1	0.5	1.3	2.6	5.9	14.7	40.2	97.5	215.4	517.7
1999	38.7	6.9	0.8	0.6	1.6	2.6	5.6	14.3	39.9	98.5	210.9	527.2
Non-Hispanic white, both sexes												
2002	37.1	6.5	1.0	0.6	1.4	3.1	6.9	16.3	40.8	96.0	215.1	482.1
2001	36.5	6.9	1.3	0.6	1.3	2.6	6.2	16.1	39.7	96.2	213.0	495.7
2000	35.9	6.3	1.2	0.6	1.4	2.7	6.2	15.3	39.8	95.0	212.0	497.5
1999	35.3	6.0	1.1	0.6	1.5	2.6	5.9	14.6	38.1	95.0	209.5	504.3
Non-Hispanic white, male												
2002	34.9	8.4	0.8	0.7	1.4	3.6	8.5	19.8	49.3	110.6	239.8	493.9
2001	33.7	7.6	1.3	0.6	1.3	2.8	7.4	19.3	46.8	112.1	235.5	495.6
2000	32.9	7.2	1.3	0.7	1.5	3.2	7.3	18.7	46.0	108.0	235.9	505.5
1999	32.1	5.9	1.4	0.6	1.5	2.9	7.0	17.7	44.6	108.0	235.2	499.1
Non-Hispanic white, female												
2002	39.2	4.4	1.2	0.4	1.4	2.6	5.3	12.9	32.6	83.5	198.6	477.0
2001	39.2	6.2	1.3	0.6	1.3	2.5	5.0	12.9	33.0	82.7	198.1	495.7
2000	38.7	5.4	1.1	0.5	1.3	2.1	5.1	11.9	34.0	83.9	196.3	494.3
1999	38.3	6.0	0.9	0.6	1.6	2.4	4.7	11.5	32.0	84.0	192.7	506.4
Non-Hispanic black, both sexes												
2002	44.1	8.3	0.9	0.6	2.0	6.5	14.4	39.7	103.7	223.2	414.2	765.8
2001	43.0	11.0	0.9	0.5	1.8	5.7	13.2	39.1	100.6	219.9	413.8	770.4
2000	42.8	9.5	0.8	0.4	1.8	6.3	12.7	38.7	98.3	220.2	429.6	772.4
1999	42.8	12.0	0.9	0.6	1.6	5.2	13.4	38.7	106.0	225.6	409.7	766.6
Non-Hispanic black, male												
2002	37.9	7.3	*	*	2.4	7.8	16.8	44.3	116.8	226.4	408.5	618.1
2001	37.1	12.4	*	*	1.7	6.8	14.9	44.1	110.0	224.8	428.5	664.1
2000	35.7	10.6	*	*	2.0	7.1	14.0	42.0	106.6	219.8	417.1	662.2
1999	36.4	13.2	*	0.6	1.6	5.9	15.2	42.7	111.5	227.0	400.2	748.8
Non-Hispanic black, female												
2002	49.7	9.5	*	0.7	1.6	5.4	12.4	35.7	93.1	220.9	417.3	822.3
2001	48.4	9.4	*	*	1.9	4.7	11.7	34.9	93.1	216.5	405.5	809.7
2000	49.2	8.5	*	*	1.7	5.6	11.5	35.9	91.6	220.5	436.6	813.3
1999	48.7	10.7	*	*	1.6	4.7	11.9	35.4	101.6	224.7	414.9	773.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Endocrine diseases (E00-E34)												
All origins 3/, both sexes												
2002	26.4	1.0	*	0.1	0.5	1.7	5.0	14.1	38.4	93.2	188.8	347.6
2001	26.1	1.1	*	0.1	0.4	1.6	4.6	14.0	38.5	93.0	187.7	350.3
2000	25.6	1.2	0.1	0.1	0.5	1.7	4.5	13.5	38.6	92.3	185.5	345.2
1999	25.5	1.0	0.1	0.1	0.5	1.6	4.5	13.4	39.0	93.7	183.9	343.6
All origins 3/, male												
2002	24.8	1.1	*	0.1	0.5	2.0	6.1	16.7	45.1	102.8	206.6	365.2
2001	24.1	1.3	*	0.1	0.5	1.8	5.5	16.7	43.7	103.2	203.6	362.2
2000	23.4	1.3	*	*	0.5	2.0	5.3	15.5	43.0	100.4	203.5	360.9
1999	23.3	*	*	*	0.5	1.7	5.3	15.9	43.2	101.3	203.1	361.8
All origins 3/, female												
2002	28.0	*	*	*	0.5	1.4	3.9	11.6	32.2	85.3	177.1	339.9
2001	28.0	*	*	*	0.4	1.4	3.7	11.4	33.7	84.6	177.3	345.4
2000	27.7	1.1	*	0.1	0.4	1.4	3.7	11.5	34.5	85.6	173.8	338.8
1999	27.5	1.3	*	0.1	0.5	1.5	3.8	11.0	35.0	87.4	171.3	336.3
Hispanic, both sexes												
2002	15.6	*	*	*	*	0.8	3.7	14.3	52.6	135.8	289.2	459.0
2001	15.6	*	*	*	0.3	0.8	3.1	15.2	53.7	143.0	284.2	511.5
2000	15.4	*	*	*	*	1.1	3.1	14.3	56.1	149.0	282.3	481.1
1999	15.6	*	*	*	*	0.7	3.2	14.2	57.2	151.6	290.6	515.7
Hispanic, male												
2002	14.1	*	*	*	*	1.0	4.2	16.9	59.2	144.5	309.0	441.7
2001	13.8	*	*	*	*	0.9	3.9	18.8	58.8	153.0	289.3	436.1
2000	14.0	*	*	*	*	1.4	3.7	15.7	65.2	161.4	301.4	463.6
1999	13.6	*	*	*	*	0.7	3.3	17.4	62.7	157.3	299.3	478.8
Hispanic, female												
2002	17.1	*	*	*	*	0.7	3.1	11.7	46.7	128.8	275.7	467.5
2001	17.6	*	*	*	*	0.7	2.3	11.7	49.1	135.1	280.8	548.3
2000	16.8	*	*	*	*	*	2.5	12.9	48.1	139.2	269.4	489.7
1999	17.7	*	*	*	*	0.8	3.1	11.1	52.4	147.2	284.8	533.4
Non-Hispanic 4/, both sexes												
2002	28.0	1.1	*	0.1	0.5	1.9	5.2	14.0	37.1	90.0	183.2	342.2
2001	27.6	1.1	*	0.1	0.4	1.8	4.8	13.8	37.2	89.4	182.8	343.4
2000	27.0	1.2	0.2	0.1	0.5	1.8	4.7	13.4	37.1	88.5	180.9	339.1
1999	26.8	1.1	0.2	0.1	0.5	1.7	4.7	13.3	37.5	90.0	178.9	336.5
Non-Hispanic 4/, male												
2002	26.4	*	*	0.1	0.6	2.3	6.3	16.6	43.8	99.7	200.7	360.7
2001	25.6	1.5	*	0.2	0.5	2.0	5.7	16.5	42.4	99.5	199.0	358.3
2000	24.8	*	*	*	0.6	2.1	5.5	15.5	41.2	96.3	198.6	355.4
1999	24.7	*	*	*	0.5	1.9	5.5	15.7	41.6	97.8	198.6	355.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Endocrine diseases (E00-E34)												
Non-Hispanic 4/, female												
2002	29.5	*	*	*	0.5	1.6	4.0	11.5	31.0	82.0	171.7	334.3
2001	29.4	*	*	*	0.4	1.5	3.9	11.3	32.4	80.9	172.1	337.2
2000	29.1	*	*	*	0.4	1.5	3.9	11.3	33.3	82.0	169.4	332.4
1999	28.8	1.4	*	0.1	0.5	1.6	3.8	10.9	33.6	83.6	166.2	328.8
Non-Hispanic white, both sexes												
2002	27.6	1.2	*	0.1	0.4	1.7	4.5	11.6	31.3	78.5	169.4	324.5
2001	27.2	1.1	*	0.1	0.3	1.4	4.1	11.6	31.4	78.5	168.9	326.7
2000	26.5	0.9	*	*	0.4	1.5	4.1	11.1	31.6	77.6	165.9	321.5
1999	26.1	1.0	*	0.1	0.5	1.5	4.0	10.9	30.6	78.5	164.8	317.3
Non-Hispanic white, male												
2002	26.6	*	*	*	0.5	2.0	5.6	14.1	38.1	90.2	191.5	355.7
2001	25.8	*	*	*	0.4	1.5	5.0	14.0	37.0	91.1	187.9	351.8
2000	24.9	*	*	*	0.5	1.8	4.9	13.2	36.0	87.7	188.4	346.0
1999	24.6	*	*	*	0.5	1.6	4.8	13.3	35.5	88.8	189.1	342.3
Non-Hispanic white, female												
2002	28.6	*	*	*	0.4	1.3	3.4	9.2	24.8	68.6	154.6	311.1
2001	28.6	*	*	*	0.2	1.3	3.2	9.2	26.1	67.8	156.3	316.3
2000	28.1	*	*	*	0.3	1.2	3.3	8.9	27.5	69.0	151.1	311.7
1999	27.6	*	*	*	0.4	1.3	3.2	8.5	26.1	69.9	148.8	307.3
Non-Hispanic black, both sexes												
2002	35.7	*	*	*	1.1	4.0	10.4	32.0	87.6	194.7	346.4	566.8
2001	35.1	*	*	*	1.0	4.0	9.6	30.9	87.9	193.9	347.0	560.0
2000	35.0	*	*	*	1.0	4.1	8.9	31.5	85.0	194.5	364.4	556.2
1999	35.1	*	*	*	0.9	3.4	9.7	32.1	94.0	197.7	345.0	554.5
Non-Hispanic black, male												
2002	30.6	*	*	*	1.2	4.9	12.5	36.8	98.2	196.5	336.7	434.3
2001	30.2	*	*	*	0.9	5.2	11.1	35.8	94.9	193.1	354.1	472.4
2000	29.0	*	*	*	1.0	4.4	10.1	34.7	90.9	190.3	345.9	479.1
1999	29.5	*	*	*	0.8	3.6	11.2	36.2	97.6	194.3	326.8	524.1
Non-Hispanic black, female												
2002	40.4	*	*	*	1.0	3.3	8.6	27.9	79.2	193.4	351.8	617.5
2001	39.6	*	*	*	1.1	3.0	8.2	26.8	82.3	194.4	343.0	592.4
2000	40.4	*	*	*	1.1	3.8	7.9	28.8	80.2	197.4	374.8	584.8
1999	40.3	*	*	*	1.0	3.2	8.3	28.6	91.0	200.1	355.2	565.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of thyroid gland (E00-E07)												
All origins 3/, both sexes												
2002	0.7	*	*	*	*	0.1	0.1	0.2	0.3	0.9	3.9	22.7
2001	0.7	*	*	*	*	*	0.1	0.2	0.3	0.8	4.2	23.2
2000	0.6	*	*	*	*	0.1	0.1	0.1	0.4	0.9	3.9	21.1
1999	0.6	*	*	*	*	0.1	0.1	0.2	0.3	1.0	4.0	22.4
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.1	12.7
2001	0.3	*	*	*	*	*	*	0.1	0.2	0.5	2.5	14.1
2000	0.3	*	*	*	*	*	*	*	0.3	0.6	1.9	12.1
1999	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.6	12.9
All origins 3/, female												
2002	1.0	*	*	*	*	*	0.1	0.2	0.4	1.2	5.1	27.0
2001	1.0	*	*	*	*	*	0.1	0.2	0.4	1.1	5.3	27.1
2000	1.0	*	*	*	*	*	0.2	0.2	0.5	1.1	5.3	24.8
1999	1.0	*	*	*	*	*	0.1	0.2	0.4	1.2	4.9	26.3
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	18.3
2001	0.2	*	*	*	*	*	*	*	*	*	*	16.9
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	15.4
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	19.9
2001	0.2	*	*	*	*	*	*	*	*	*	*	19.5
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	*	*	0.1	0.1	0.2	0.3	1.0	4.0	22.8
2001	0.7	*	*	*	*	*	0.1	0.2	0.3	0.8	4.2	23.4
2000	0.7	*	*	*	*	0.1	0.1	0.1	0.4	0.9	4.0	21.4
1999	0.7	*	*	*	*	0.1	0.1	0.2	0.3	1.0	4.0	22.7
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	*	0.2	0.6	2.2	12.6
2001	0.3	*	*	*	*	*	*	0.1	0.2	0.5	2.4	14.2
2000	0.3	*	*	*	*	*	*	*	0.3	0.7	2.0	12.1
1999	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.6	13.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of thyroid gland (E00-E07)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	*	0.1	0.2	0.4	1.2	5.2	27.2
2001	1.1	*	*	*	*	*	0.1	0.2	0.4	1.1	5.4	27.2
2000	1.1	*	*	*	*	*	0.2	0.2	0.5	1.1	5.3	25.1
1999	1.1	*	*	*	*	*	0.1	0.3	0.4	1.2	4.9	26.5
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	*	*	0.1	0.3	0.9	4.1	23.6
2001	0.8	*	*	*	*	*	0.1	0.1	0.3	0.8	4.1	24.0
2000	0.8	*	*	*	*	*	0.1	0.1	0.4	0.8	4.0	21.9
1999	0.8	*	*	*	*	*	0.1	0.1	0.3	0.9	3.9	23.3
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	*	*	0.6	2.2	13.0
2001	0.4	*	*	*	*	*	*	*	0.2	0.5	2.3	14.7
2000	0.3	*	*	*	*	*	*	*	0.3	0.5	2.1	12.0
1999	0.3	*	*	*	*	*	*	*	*	0.5	2.4	13.5
Non-Hispanic white, female												
2002	1.3	*	*	*	*	*	*	0.2	0.4	1.2	5.4	28.2
2001	1.3	*	*	*	*	*	*	0.2	0.4	1.1	5.3	27.9
2000	1.2	*	*	*	*	*	*	0.2	0.5	1.1	5.2	25.9
1999	1.2	*	*	*	*	*	*	0.2	0.3	1.2	4.9	27.2
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	*	0.5	0.8	1.3	3.2	16.9
2001	0.6	*	*	*	*	*	*	*	*	1.4	5.8	19.7
2000	0.6	*	*	*	*	*	0.5	*	*	1.9	5.0	19.1
1999	0.6	*	*	*	*	*	*	0.6	0.9	1.5	5.3	18.8
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	*	*	*	4.0	20.1
2001	0.8	*	*	*	*	*	*	*	*	*	6.2	22.2
2000	0.8	*	*	*	*	*	*	*	*	*	6.9	20.1
1999	0.8	*	*	*	*	*	*	*	*	*	5.8	21.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All origins 3/, both sexes												
2002	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
All origins 3/, male												
2002	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
All origins 3/, female												
2002	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8
Hispanic, both sexes												
2002	15.3	*	*	*	*	0.8	3.5	14.0	52.1	134.7	285.6	437.9
2001	15.3	*	*	*	*	0.7	3.1	14.9	53.2	141.4	280.1	490.2
2000	15.1	*	*	*	*	1.0	3.0	14.0	55.5	147.9	278.5	465.1
1999	15.3	*	*	*	*	0.7	3.2	13.9	56.6	149.7	285.2	493.9
Hispanic, male												
2002	13.9	*	*	*	*	0.9	4.1	16.7	58.9	143.7	306.9	425.0
2001	13.6	*	*	*	*	0.8	3.9	18.5	58.7	151.6	286.2	422.8
2000	13.8	*	*	*	*	1.4	3.6	15.5	64.7	160.8	299.4	449.4
1999	13.4	*	*	*	*	0.6	3.3	17.2	62.5	155.8	293.0	468.0
Hispanic, female												
2002	16.7	*	*	*	*	0.7	2.9	11.4	46.0	127.6	270.9	444.3
2001	17.1	*	*	*	*	0.7	2.2	11.4	48.4	133.4	275.9	523.1
2000	16.5	*	*	*	*	*	2.4	12.6	47.4	137.7	264.5	472.8
1999	17.3	*	*	*	*	0.8	3.1	10.7	51.4	144.9	280.0	506.4
Non-Hispanic 4/, both sexes												
2002	26.9	*	*	0.1	0.5	1.8	5.0	13.6	36.4	88.2	177.2	315.0
2001	26.4	*	*	0.1	0.4	1.7	4.5	13.4	36.5	87.7	176.4	314.8
2000	25.9	*	*	0.1	0.5	1.7	4.4	13.0	36.4	86.8	174.8	313.4
1999	25.7	*	*	0.1	0.4	1.6	4.4	12.8	36.8	88.1	173.1	310.0
Non-Hispanic 4/, male												
2002	25.8	*	*	0.1	0.5	2.2	6.2	16.3	43.1	98.2	196.6	344.0
2001	25.0	*	*	0.1	0.4	1.9	5.5	16.1	41.8	98.0	194.3	338.9
2000	24.2	*	*	*	0.5	2.0	5.3	15.2	40.7	95.0	194.6	339.7
1999	24.1	*	*	*	0.4	1.8	5.3	15.3	41.0	96.3	193.8	339.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Non-Hispanic 4/, female												
2002	27.9	*	*	*	0.4	1.4	3.8	11.1	30.1	79.8	164.3	302.5
2001	27.8	*	*	*	0.3	1.4	3.6	10.9	31.6	79.1	164.6	304.7
2000	27.6	*	*	*	0.4	1.4	3.5	11.0	32.4	80.1	161.8	302.8
1999	27.3	*	*	*	0.4	1.4	3.6	10.4	32.8	81.4	159.6	298.2
Non-Hispanic white, both sexes												
2002	26.4	*	*	*	0.4	1.5	4.3	11.3	30.6	76.8	163.3	296.4
2001	26.0	*	*	*	0.3	1.3	3.9	11.3	30.7	77.0	162.7	297.4
2000	25.4	*	*	*	0.4	1.4	3.9	10.7	30.9	76.1	159.9	295.5
1999	25.0	*	*	*	0.4	1.4	3.8	10.5	30.1	76.8	159.1	290.2
Non-Hispanic white, male												
2002	26.0	*	*	*	0.4	1.9	5.5	13.8	37.5	88.9	187.4	338.4
2001	25.1	*	*	*	0.3	1.5	4.8	13.7	36.4	89.8	183.5	332.1
2000	24.3	*	*	*	0.5	1.7	4.8	13.0	35.4	86.6	184.5	330.4
1999	24.0	*	*	*	0.4	1.5	4.6	13.0	35.0	87.7	184.6	326.0
Non-Hispanic white, female												
2002	26.8	*	*	*	0.4	1.2	3.2	8.8	24.1	66.5	147.0	278.3
2001	26.9	*	*	*	0.2	1.2	2.9	8.8	25.3	66.0	148.9	283.0
2000	26.5	*	*	*	0.3	1.1	3.1	8.6	26.6	67.3	143.8	281.4
1999	26.0	*	*	*	0.3	1.2	3.0	8.1	25.4	67.7	142.4	275.9
Non-Hispanic black, both sexes												
2002	34.8	*	*	*	1.0	3.8	9.9	30.9	85.9	191.5	340.2	544.7
2001	34.1	*	*	*	1.0	3.8	9.1	30.1	86.5	190.8	337.9	533.8
2000	34.0	*	*	*	1.0	3.6	8.2	31.0	83.9	190.6	354.8	530.1
1999	34.1	*	*	*	0.8	3.1	9.1	31.0	92.0	194.0	336.1	530.9
Non-Hispanic black, male												
2002	30.0	*	*	*	1.1	4.6	12.1	35.8	96.6	193.3	332.6	423.4
2001	29.4	*	*	*	0.9	5.0	10.5	35.0	93.8	189.5	345.1	452.7
2000	28.3	*	*	*	1.0	4.0	9.7	34.2	89.7	186.9	339.4	456.8
1999	28.6	*	*	*	*	3.4	10.7	35.3	95.4	189.7	317.7	506.3
Non-Hispanic black, female												
2002	39.2	*	*	*	0.8	3.0	8.0	26.8	77.3	190.3	344.5	591.1
2001	38.4	*	*	*	1.0	2.8	7.8	25.9	80.6	191.7	333.9	563.8
2000	39.1	*	*	*	1.0	3.3	7.0	28.2	79.1	193.3	363.5	557.3
1999	39.0	*	*	*	0.9	2.7	7.7	27.2	89.2	197.0	346.5	540.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of pituitary gland (E22-E23)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	1.1
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.1	0.5	1.0
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.1
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.1
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.1
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.6	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.3	1.2
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.1
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.1	0.5	1.0
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.1
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.0
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.8	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of pituitary gland (E22-E23)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.4	1.1
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	1.1
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.5	1.0
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.5	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.1
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	1.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.3	1.2
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of adrenal glands (E24-E27)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.6	1.1
2001	0.1	*	*	*	*	*	0.0	0.1	0.1	0.3	0.8	1.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.7	1.0
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.5	1.0
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.9	2.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.7	1.1
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.7
2000	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.4	1.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.6	1.1
2001	0.1	*	*	*	*	*	0.1	0.1	0.1	0.3	0.8	1.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.7	1.1
1999	0.1	*	*	*	*	*	*	0.1	*	0.3	0.5	0.9
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.9	2.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of adrenal glands (E24-E27)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.7	1.1
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.7
2000	0.1	*	*	*	*	*	*	*	0.2	0.3	0.7	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.5	1.0
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.3	0.6	1.1
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.8	1.8
2000	0.1	*	*	*	*	*	*	*	0.2	0.2	0.7	1.1
1999	0.1	*	*	*	*	*	*	0.1	*	0.3	0.5	0.9
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.7	1.1
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	1.7
2000	0.1	*	*	*	*	*	*	*	0.2	*	0.7	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.5	1.0
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of thymus (E32)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of thymus (E32)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other endocrine diseases (E15-E21,E28-E31,E34)												
All origins 3/, both sexes												
2002	0.2	0.7	*	*	*	*	*	0.1	0.2	0.4	1.0	2.2
2001	0.2	0.7	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.5
2000	0.1	0.7	*	*	*	*	0.1	0.1	0.1	0.3	0.9	2.2
1999	0.2	0.7	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8
2001	0.1	*	*	*	*	*	*	*	0.2	0.4	0.7	2.2
2000	0.1	*	*	*	*	*	*	0.1	*	*	0.9	1.7
1999	0.1	*	*	*	*	*	*	0.1	0.2	0.5	1.0	*
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.3
2001	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	2.6
2000	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	2.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.8	2.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	0.8	*	*	*	*	*	0.1	0.2	0.4	1.0	2.2
2001	0.2	0.7	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.5
2000	0.2	0.7	*	*	*	*	0.1	0.1	0.1	0.3	0.9	2.2
1999	0.2	0.8	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.9	1.9
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.8	2.3
2000	0.1	*	*	*	*	*	*	0.1	*	*	0.9	1.7
1999	0.1	*	*	*	*	*	*	0.1	0.2	0.5	1.0	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other endocrine diseases (E15-E21,E28-E31,E34)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.1	2.4
2001	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	2.5
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.9	2.4
1999	0.2	*	*	*	*	*	*	0.1	*	0.3	0.8	2.0
Non-Hispanic white, both sexes												
2002	0.2	0.9	*	*	*	*	*	0.1	0.2	0.4	1.0	2.2
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.2	0.7	2.4
2000	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.7	2.0
1999	0.2	*	*	*	*	*	*	0.1	0.1	0.3	0.8	1.8
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	0.2	0.3	0.9	2.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	2.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	1.0	*
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	2.2
2001	0.2	*	*	*	*	*	*	*	0.2	*	0.8	2.4
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.0
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	1.4	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	3.2	*
1999	0.3	*	*	*	*	*	*	*	*	1.2	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional diseases (E40-E68)												
All origins 3/, both sexes												
2002	2.6	*	*	*	0.2	0.8	1.6	2.6	3.9	4.9	11.3	42.9
2001	2.4	*	*	*	0.2	0.6	1.3	2.4	3.3	4.6	10.8	44.5
2000	2.6	*	*	*	0.1	0.6	1.3	2.3	3.2	4.9	12.6	51.7
1999	2.5	*	*	*	0.2	0.5	1.2	2.0	2.9	4.7	12.3	55.3
All origins 3/, male												
2002	2.2	*	*	*	0.2	0.9	1.8	2.6	3.7	4.5	10.3	37.8
2001	1.9	*	*	*	0.2	0.7	1.4	2.4	3.1	4.3	9.8	35.8
2000	2.0	*	*	*	0.2	0.7	1.5	2.4	3.2	4.6	11.2	44.6
1999	2.0	*	*	*	0.2	0.6	1.4	2.0	3.1	4.6	11.6	48.7
All origins 3/, female												
2002	3.0	*	*	*	0.2	0.6	1.4	2.6	4.0	5.1	12.0	45.1
2001	2.9	*	*	*	0.2	0.5	1.2	2.5	3.5	4.9	11.4	48.1
2000	3.1	*	*	*	0.1	0.4	1.2	2.2	3.2	5.2	13.6	54.6
1999	2.9	*	*	*	0.2	0.4	1.0	2.0	2.6	4.7	12.8	58.0
Hispanic, both sexes												
2002	0.9	*	*	*	*	0.4	0.9	1.4	1.7	3.5	9.0	29.9
2001	0.8	*	*	*	*	0.5	0.9	1.2	1.6	2.4	9.6	41.9
2000	0.8	*	*	*	*	0.4	0.9	1.4	1.3	4.0	9.7	41.8
1999	0.8	*	*	*	*	0.3	0.8	1.0	1.9	2.5	7.5	52.6
Hispanic, male												
2002	0.7	*	*	*	*	*	1.0	1.1	*	*	*	*
2001	0.7	*	*	*	*	0.6	1.0	*	*	*	10.3	45.7
2000	0.7	*	*	*	*	0.6	1.0	1.4	*	*	*	*
1999	0.7	*	*	*	*	*	1.2	*	*	*	*	51.8
Hispanic, female												
2002	1.0	*	*	*	*	*	0.8	1.7	2.0	3.5	9.8	29.8
2001	0.9	*	*	*	*	*	0.8	1.3	*	*	9.1	40.0
2000	1.0	*	*	*	*	*	*	1.4	*	4.3	10.6	44.5
1999	0.8	*	*	*	*	*	*	*	*	*	7.6	53.0
Non-Hispanic 4/, both sexes												
2002	2.9	*	*	*	0.2	0.8	1.7	2.7	4.0	4.9	11.4	43.3
2001	2.6	*	*	*	0.2	0.6	1.3	2.5	3.4	4.8	10.8	44.5
2000	2.8	*	*	*	0.1	0.6	1.4	2.4	3.4	5.0	12.7	51.9
1999	2.7	*	*	*	0.2	0.5	1.3	2.1	2.9	4.8	12.5	55.3
Non-Hispanic 4/, male												
2002	2.4	*	*	*	0.2	0.9	1.9	2.8	3.9	4.6	10.3	38.0
2001	2.0	*	*	*	0.2	0.7	1.4	2.5	3.2	4.4	9.8	35.4
2000	2.2	*	*	*	0.2	0.7	1.5	2.5	3.4	4.7	11.2	44.8
1999	2.2	*	*	*	0.2	0.7	1.4	2.1	3.1	4.7	11.8	48.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional diseases (E40-E68)												
Non-Hispanic 4/, female												
2002	3.3	*	*	*	0.2	0.7	1.4	2.7	4.2	5.2	12.1	45.5
2001	3.2	*	*	*	0.2	0.5	1.3	2.6	3.6	5.0	11.5	48.3
2000	3.3	*	*	*	*	0.4	1.3	2.3	3.3	5.2	13.7	54.8
1999	3.2	*	*	*	0.2	0.4	1.1	2.1	2.7	4.8	13.0	58.1
Non-Hispanic white, both sexes												
2002	2.9	*	*	*	0.1	0.6	1.5	2.6	3.8	4.6	10.8	42.0
2001	2.7	*	*	*	0.1	0.5	1.1	2.3	3.3	4.4	10.1	43.7
2000	2.9	*	*	*	0.1	0.4	1.3	2.2	3.1	4.6	12.1	50.7
1999	2.7	*	*	*	0.1	0.4	1.1	1.9	2.7	4.3	11.7	54.2
Non-Hispanic white, male												
2002	2.4	*	*	*	*	0.7	1.8	2.8	3.8	4.4	9.8	37.4
2001	2.0	*	*	*	*	0.6	1.2	2.3	3.1	4.1	9.2	33.9
2000	2.3	*	*	*	0.2	0.6	1.4	2.5	3.2	4.3	10.6	43.8
1999	2.2	*	*	*	*	0.5	1.3	2.1	2.9	4.2	10.8	46.6
Non-Hispanic white, female												
2002	3.3	*	*	*	*	0.6	1.2	2.3	3.8	4.8	11.5	44.0
2001	3.3	*	*	*	*	0.4	1.0	2.3	3.5	4.7	10.7	47.8
2000	3.4	*	*	*	*	0.3	1.1	1.9	3.1	4.9	13.1	53.5
1999	3.3	*	*	*	*	0.3	0.9	1.8	2.5	4.4	12.3	57.2
Non-Hispanic black, both sexes												
2002	3.4	*	*	*	0.5	2.0	2.9	4.5	7.5	8.9	20.0	61.7
2001	3.2	*	*	*	0.5	1.3	2.9	4.9	5.6	8.7	20.1	56.8
2000	3.2	*	*	*	0.4	1.6	2.6	4.4	6.0	9.0	20.8	69.0
1999	3.1	*	*	*	0.4	1.3	2.4	3.6	5.5	9.9	23.3	72.4
Non-Hispanic black, male												
2002	2.7	*	*	*	*	2.4	3.1	3.3	6.5	7.8	19.0	47.0
2001	2.6	*	*	*	*	1.3	3.0	4.2	5.1	8.3	18.3	58.9
2000	2.6	*	*	*	*	1.8	2.6	3.4	6.2	9.1	20.7	66.9
1999	2.7	*	*	*	*	1.6	2.4	2.9	5.9	10.2	24.7	76.1
Non-Hispanic black, female												
2002	4.1	*	*	*	*	1.7	2.8	5.6	8.4	9.7	20.6	67.3
2001	3.7	*	*	*	*	1.3	2.9	5.5	6.1	9.0	21.1	56.0
2000	3.7	*	*	*	*	1.4	2.6	5.2	5.8	9.0	20.8	69.8
1999	3.6	*	*	*	*	1.1	2.4	4.2	5.1	9.6	22.6	71.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
All origins 3/, male												
2002	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
All origins 3/, female												
2002	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	*	*	1.8	8.2	28.3
2001	0.4	*	*	*	*	*	*	*	*	*	7.9	38.1
2000	0.4	*	*	*	*	*	*	*	*	2.0	7.7	38.5
1999	0.4	*	*	*	*	*	*	*	*	*	6.2	47.7
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	8.9	41.9
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	45.3
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	8.4	28.2
2001	0.4	*	*	*	*	*	*	*	*	*	7.3	36.3
2000	0.5	*	*	*	*	*	*	*	*	*	7.6	41.5
1999	0.5	*	*	*	*	*	*	*	*	*	*	48.9
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	*	*	0.1	0.2	0.6	1.9	8.3	38.7
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.2	40.2
2000	1.5	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.5
1999	1.6	*	*	*	*	*	0.1	0.3	0.7	2.2	9.6	50.0
Non-Hispanic 4/, male												
2002	1.0	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.3
2001	0.9	*	*	*	*	*	*	0.3	0.6	2.0	8.0	33.2
2000	1.0	*	*	*	*	*	*	0.2	0.8	2.5	8.8	40.9
1999	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	9.5	44.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Non-Hispanic 4/, female												
2002	1.7	*	*	*	*	*	0.1	0.3	0.5	1.8	8.6	40.1
2001	1.7	*	*	*	*	*	*	0.3	0.6	1.9	8.3	43.2
2000	2.0	*	*	*	*	*	0.2	0.3	0.6	2.2	10.3	48.7
1999	2.0	*	*	*	*	*	0.2	0.3	0.5	2.0	9.7	52.2
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.4
2001	1.4	*	*	*	*	*	*	0.2	0.5	1.8	7.6	39.4
2000	1.6	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.3
1999	1.6	*	*	*	*	*	0.1	0.2	0.5	1.9	8.9	48.8
Non-Hispanic white, male												
2002	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.5	34.5
2001	0.9	*	*	*	*	*	*	0.2	0.5	1.7	7.4	31.7
2000	1.1	*	*	*	*	*	*	0.2	0.6	2.1	8.3	39.8
1999	1.1	*	*	*	*	*	*	0.2	0.6	2.0	8.6	42.0
Non-Hispanic white, female												
2002	1.8	*	*	*	*	*	*	0.3	0.5	1.7	8.3	38.6
2001	1.9	*	*	*	*	*	*	0.3	0.6	1.9	7.7	42.7
2000	2.2	*	*	*	*	*	0.1	0.2	0.6	2.0	10.0	47.4
1999	2.2	*	*	*	*	*	0.2	0.3	0.4	1.8	9.2	51.6
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	*	*	*	*	1.7	3.6	14.2	57.2
2001	1.2	*	*	*	*	*	*	0.6	1.1	3.4	15.0	52.1
2000	1.4	*	*	*	*	*	*	0.6	1.6	4.8	15.1	64.2
1999	1.6	*	*	*	*	*	*	0.6	1.8	5.5	18.1	66.0
Non-Hispanic black, male												
2002	1.0	*	*	*	*	*	*	*	2.7	4.0	15.2	44.9
2001	1.1	*	*	*	*	*	*	1.0	*	5.0	15.9	56.6
2000	1.2	*	*	*	*	*	*	*	2.5	6.4	17.3	62.2
1999	1.4	*	*	*	*	*	*	*	2.6	8.0	20.6	74.9
Non-Hispanic black, female												
2002	1.5	*	*	*	*	*	*	*	*	3.4	13.6	61.9
2001	1.3	*	*	*	*	*	*	*	*	2.3	14.5	50.5
2000	1.6	*	*	*	*	*	*	*	*	3.6	13.9	65.0
1999	1.7	*	*	*	*	*	*	*	*	3.7	16.6	62.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional marasmus (E41)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.7
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.5	2.5
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	1.7
2001	0.0	*	*	*	*	*	*	*	*	*	0.5	2.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	3.0
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.5	2.5
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.1
2000	0.1	*	*	*	*	*	*	*	*	*	*	2.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	3.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional marasmus (E41)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2000	0.1	*	*	*	*	*	*	*	*	*	*	2.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	2.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.0	36.0
2001	1.1	*	*	*	*	*	0.1	0.2	0.5	1.8	7.7	37.7
2000	1.3	*	*	*	*	*	0.1	0.2	0.6	2.2	9.2	43.6
1999	1.4	*	*	*	*	*	0.1	0.3	0.6	2.1	9.0	47.5
All origins 3/, male												
2002	0.8	*	*	*	*	*	0.1	0.2	0.6	2.1	7.7	33.5
2001	0.8	*	*	*	*	*	*	0.2	0.6	1.8	7.5	31.3
2000	0.9	*	*	*	*	*	*	0.2	0.7	2.4	8.4	37.8
1999	0.9	*	*	*	*	*	0.1	0.3	0.8	2.3	8.9	41.4
All origins 3/, female												
2002	1.5	*	*	*	*	*	0.1	0.2	0.5	1.7	8.2	37.1
2001	1.5	*	*	*	*	*	*	0.3	0.5	1.8	7.9	40.4
2000	1.7	*	*	*	*	*	0.2	0.2	0.5	2.1	9.7	45.9
1999	1.8	*	*	*	*	*	0.1	0.3	0.5	1.9	9.1	50.0
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	7.8	24.9
2001	0.3	*	*	*	*	*	*	*	*	*	7.6	33.8
2000	0.4	*	*	*	*	*	*	*	*	2.0	7.3	37.8
1999	0.4	*	*	*	*	*	*	*	*	*	6.2	46.3
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	43.1
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	8.2	26.5
2001	0.4	*	*	*	*	*	*	*	*	*	7.0	32.6
2000	0.5	*	*	*	*	*	*	*	*	*	7.6	41.5
1999	0.5	*	*	*	*	*	*	*	*	*	*	47.8
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	*	*	0.1	0.2	0.6	1.9	8.0	36.3
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	1.9	7.7	37.8
2000	1.5	*	*	*	*	*	0.1	0.2	0.6	2.2	9.2	43.7
1999	1.5	*	*	*	*	*	0.1	0.3	0.6	2.1	9.1	47.5
Non-Hispanic 4/, male												
2002	0.9	*	*	*	*	*	0.1	0.2	0.7	2.0	7.6	33.8
2001	0.9	*	*	*	*	*	*	0.2	0.6	1.9	7.5	31.1
2000	1.0	*	*	*	*	*	*	0.2	0.7	2.4	8.4	38.1
1999	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	9.0	41.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)												
Non-Hispanic 4/, female												
2002	1.6	*	*	*	*	*	0.1	0.2	0.5	1.7	8.2	37.4
2001	1.7	*	*	*	*	*	*	0.3	0.6	1.8	7.9	40.6
2000	1.9	*	*	*	*	*	0.2	0.3	0.6	2.1	9.8	46.0
1999	1.9	*	*	*	*	*	0.2	0.3	0.5	1.9	9.2	50.0
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	*	0.1	0.2	0.5	1.7	7.6	35.0
2001	1.3	*	*	*	*	*	*	0.2	0.5	1.7	7.2	37.0
2000	1.5	*	*	*	*	*	0.1	0.2	0.5	2.0	8.8	42.5
1999	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.5	46.4
Non-Hispanic white, male												
2002	1.0	*	*	*	*	*	*	0.2	0.5	1.9	7.2	33.0
2001	0.9	*	*	*	*	*	*	0.1	0.5	1.7	6.9	29.7
2000	1.0	*	*	*	*	*	*	0.2	0.5	2.1	7.9	37.1
1999	1.0	*	*	*	*	*	*	0.2	0.6	1.9	8.1	39.2
Non-Hispanic white, female												
2002	1.7	*	*	*	*	*	*	0.2	0.5	1.6	7.9	35.9
2001	1.8	*	*	*	*	*	*	0.3	0.6	1.8	7.4	40.0
2000	2.0	*	*	*	*	*	0.1	0.2	0.5	1.9	9.5	44.7
1999	2.1	*	*	*	*	*	0.2	0.2	0.4	1.7	8.7	49.3
Non-Hispanic black, both sexes												
2002	1.2	*	*	*	*	*	*	*	1.6	3.5	13.2	55.4
2001	1.2	*	*	*	*	*	*	0.6	0.9	3.3	14.5	49.6
2000	1.4	*	*	*	*	*	*	0.5	1.5	4.6	14.8	60.1
1999	1.5	*	*	*	*	*	*	0.6	1.7	5.2	17.1	62.7
Non-Hispanic black, male												
2002	0.9	*	*	*	*	*	*	*	2.5	4.0	14.1	43.8
2001	1.1	*	*	*	*	*	*	*	*	4.9	15.6	54.3
2000	1.1	*	*	*	*	*	*	*	2.4	6.1	16.4	56.4
1999	1.4	*	*	*	*	*	*	*	2.4	7.7	20.0	70.1
Non-Hispanic black, female												
2002	1.5	*	*	*	*	*	*	*	*	3.2	12.8	59.8
2001	1.3	*	*	*	*	*	*	*	*	2.2	13.8	47.9
2000	1.6	*	*	*	*	*	*	*	*	3.5	13.9	61.5
1999	1.6	*	*	*	*	*	*	*	*	3.4	15.4	60.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.1
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.2
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.6	2.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.8
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	4.0
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.2
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.6	3.8
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.8	4.2
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.6	4.0
1999	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)												
All origins 3/, both sexes												
2002	1.3	*	*	*	0.2	0.7	1.5	2.3	3.2	2.8	2.4	1.4
2001	1.1	*	*	*	0.2	0.6	1.2	2.2	2.7	2.6	2.0	1.6
2000	1.1	*	*	*	0.1	0.5	1.2	2.1	2.5	2.4	2.2	1.7
1999	0.9	*	*	*	0.1	0.5	1.1	1.7	2.1	2.3	2.1	1.4
All origins 3/, male												
2002	1.2	*	*	*	0.2	0.8	1.7	2.4	3.0	2.3	1.7	*
2001	1.0	*	*	*	0.1	0.6	1.3	2.1	2.4	2.2	1.3	*
2000	1.0	*	*	*	0.2	0.7	1.4	2.2	2.4	1.9	1.6	*
1999	0.9	*	*	*	0.1	0.6	1.3	1.7	2.2	1.9	1.4	*
All origins 3/, female												
2002	1.3	*	*	*	0.1	0.6	1.2	2.3	3.5	3.2	2.9	1.6
2001	1.2	*	*	*	0.2	0.5	1.1	2.2	2.9	2.9	2.5	2.0
2000	1.1	*	*	*	*	0.4	1.0	1.9	2.6	2.8	2.7	2.0
1999	1.0	*	*	*	0.1	0.4	0.9	1.7	2.1	2.7	2.6	1.9
Hispanic, both sexes												
2002	0.5	*	*	*	*	0.4	0.8	1.3	1.5	1.7	*	*
2001	0.5	*	*	*	*	0.5	0.8	1.1	1.3	*	*	*
2000	0.4	*	*	*	*	0.4	0.8	1.2	*	*	*	*
1999	0.4	*	*	*	*	0.3	0.8	0.8	1.5	*	*	*
Hispanic, male												
2002	0.4	*	*	*	*	*	0.9	*	*	*	*	*
2001	0.5	*	*	*	*	0.6	0.9	*	*	*	*	*
2000	0.4	*	*	*	*	*	0.9	*	*	*	*	*
1999	0.4	*	*	*	*	*	1.1	*	*	*	*	*
Hispanic, female												
2002	0.6	*	*	*	*	*	0.8	1.6	*	*	*	*
2001	0.5	*	*	*	*	*	*	1.2	*	*	*	*
2000	0.5	*	*	*	*	*	*	1.3	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.4	*	*	*	0.2	0.8	1.5	2.4	3.4	2.8	2.5	1.4
2001	1.2	*	*	*	0.2	0.6	1.2	2.2	2.8	2.7	2.0	1.5
2000	1.1	*	*	*	0.1	0.5	1.3	2.1	2.6	2.5	2.3	1.7
1999	1.0	*	*	*	0.1	0.5	1.1	1.8	2.2	2.4	2.2	1.4
Non-Hispanic 4/, male												
2002	1.3	*	*	*	0.2	0.9	1.8	2.5	3.1	2.4	1.8	*
2001	1.1	*	*	*	0.1	0.7	1.3	2.2	2.5	2.2	1.3	*
2000	1.1	*	*	*	0.2	0.7	1.4	2.3	2.5	2.0	1.6	*
1999	1.0	*	*	*	0.1	0.6	1.3	1.8	2.2	2.0	1.4	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)												
Non-Hispanic 4/, female												
2002	1.5	*	*	*	0.2	0.7	1.3	2.4	3.6	3.2	3.0	1.7
2001	1.3	*	*	*	0.2	0.5	1.2	2.3	3.0	3.0	2.5	1.9
2000	1.2	*	*	*	*	0.4	1.1	2.0	2.7	2.8	2.7	2.0
1999	1.1	*	*	*	0.2	0.4	0.9	1.7	2.1	2.7	2.6	1.9
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	0.1	0.6	1.4	2.3	3.2	2.8	2.3	1.5
2001	1.1	*	*	*	0.1	0.5	1.1	2.1	2.7	2.5	1.9	1.5
2000	1.1	*	*	*	0.1	0.4	1.1	2.0	2.5	2.4	2.2	1.7
1999	1.0	*	*	*	0.1	0.4	1.0	1.7	2.1	2.3	2.1	1.4
Non-Hispanic white, male												
2002	1.4	*	*	*	*	0.7	1.7	2.6	3.2	2.4	1.8	*
2001	1.1	*	*	*	*	0.6	1.2	2.1	2.6	2.2	1.3	*
2000	1.1	*	*	*	*	0.6	1.4	2.3	2.5	2.0	1.6	*
1999	1.0	*	*	*	*	0.5	1.2	1.8	2.2	2.1	1.4	*
Non-Hispanic white, female												
2002	1.3	*	*	*	*	0.5	1.1	2.0	3.2	3.0	2.6	1.7
2001	1.2	*	*	*	*	0.4	1.0	2.0	2.9	2.7	2.4	1.9
2000	1.1	*	*	*	*	0.3	0.9	1.7	2.5	2.6	2.6	2.1
1999	1.0	*	*	*	*	0.3	0.7	1.6	2.0	2.5	2.5	1.8
Non-Hispanic black, both sexes												
2002	2.0	*	*	*	0.5	2.0	2.6	4.1	5.7	4.6	5.3	*
2001	1.8	*	*	*	0.5	1.2	2.7	4.2	4.5	5.0	4.0	*
2000	1.7	*	*	*	*	1.4	2.3	3.8	4.2	4.2	4.4	*
1999	1.5	*	*	*	0.4	1.3	2.1	2.9	3.4	4.3	3.7	*
Non-Hispanic black, male												
2002	1.6	*	*	*	*	2.3	2.8	2.6	3.6	3.2	*	*
2001	1.4	*	*	*	*	1.3	2.6	3.0	3.2	*	*	*
2000	1.4	*	*	*	*	1.6	2.4	2.7	3.4	*	*	*
1999	1.2	*	*	*	*	1.5	2.1	2.2	3.0	*	*	*
Non-Hispanic black, female												
2002	2.5	*	*	*	*	1.6	2.5	5.3	7.4	5.6	6.8	*
2001	2.3	*	*	*	*	1.1	2.7	5.2	5.5	6.6	5.2	*
2000	2.0	*	*	*	*	1.2	2.3	4.7	4.9	5.4	5.4	*
1999	1.7	*	*	*	*	1.1	2.2	3.5	3.8	5.9	4.9	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Metabolic disorders (E70-E88)												
All origins 3/, both sexes												
2002	5.5	5.5	0.8	0.4	0.6	0.6	0.8	2.0	5.6	13.0	35.2	114.3
2001	5.5	6.1	1.1	0.4	0.6	0.6	0.8	2.1	5.0	13.1	34.1	124.4
2000	5.4	5.3	0.9	0.4	0.7	0.6	0.8	2.0	5.1	12.6	33.9	124.1
1999	5.3	5.6	0.8	0.4	0.7	0.6	0.8	1.8	4.7	12.2	33.1	130.8
All origins 3/, male												
2002	4.9	6.1	0.8	0.5	0.6	0.7	1.0	2.8	7.3	16.0	38.8	101.5
2001	4.9	6.5	1.1	0.4	0.6	0.6	1.0	2.8	6.8	16.7	38.6	109.8
2000	4.8	5.5	0.9	0.5	0.7	0.7	0.9	2.9	6.8	15.8	36.7	112.0
1999	4.5	6.3	0.9	0.4	0.7	0.6	0.9	2.4	6.1	15.1	35.8	110.5
All origins 3/, female												
2002	6.0	4.9	0.9	0.4	0.6	0.5	0.7	1.3	4.0	10.5	32.8	119.9
2001	6.1	5.6	1.1	0.4	0.7	0.5	0.6	1.5	3.4	10.2	31.1	130.6
2000	6.0	5.0	0.8	0.3	0.7	0.5	0.7	1.1	3.5	10.0	32.0	129.1
1999	6.1	4.9	0.7	0.4	0.8	0.6	0.6	1.2	3.5	9.8	31.3	139.0
Hispanic, both sexes												
2002	1.6	4.4	0.7	0.4	*	*	0.3	1.1	2.4	8.6	26.0	65.4
2001	1.5	4.6	0.8	0.3	*	*	0.4	0.9	3.4	8.4	22.5	67.5
2000	1.4	4.2	0.7	0.4	*	*	*	1.1	2.9	8.0	21.3	73.0
1999	1.4	4.9	*	*	*	*	*	1.0	2.5	7.7	23.1	80.0
Hispanic, male												
2002	1.5	*	*	0.6	*	*	*	1.3	2.4	10.7	29.5	71.9
2001	1.5	*	*	*	*	*	*	*	4.6	10.0	25.5	76.2
2000	1.3	*	*	*	*	*	*	1.4	3.7	11.2	20.7	60.5
1999	1.3	6.2	*	*	*	*	*	*	3.1	9.4	28.2	60.4
Hispanic, female												
2002	1.6	5.2	*	*	*	*	*	*	2.4	7.0	23.6	62.2
2001	1.6	*	*	*	*	*	*	*	2.4	7.1	20.6	63.3
2000	1.6	5.3	*	*	*	*	*	*	2.2	5.5	21.8	79.1
1999	1.6	*	*	*	*	*	*	*	*	6.4	19.8	89.4
Non-Hispanic 4/, both sexes												
2002	6.1	5.8	0.9	0.4	0.7	0.7	0.9	2.1	5.8	13.2	35.5	115.9
2001	6.1	6.4	1.1	0.4	0.7	0.6	0.9	2.3	5.1	13.4	34.5	126.3
2000	5.9	5.5	0.9	0.4	0.8	0.7	0.9	2.0	5.3	12.9	34.3	125.7
1999	5.9	5.7	0.9	0.5	0.8	0.7	0.8	1.9	4.8	12.4	33.4	132.3
Non-Hispanic 4/, male												
2002	5.5	6.7	0.8	0.5	0.7	0.8	1.1	2.9	7.7	16.3	39.1	102.3
2001	5.4	7.0	1.1	0.4	0.7	0.6	1.1	3.0	6.9	17.0	39.1	110.6
2000	5.3	6.1	0.9	0.5	0.8	0.8	1.0	3.0	7.0	16.1	37.3	113.7
1999	5.0	6.2	1.0	0.5	0.8	0.7	1.0	2.5	6.2	15.4	36.0	112.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Metabolic disorders (E70-E88)												
Non-Hispanic 4/, female												
2002	6.6	4.8	1.0	0.4	0.7	0.6	0.7	1.4	4.1	10.7	33.2	121.8
2001	6.7	5.7	1.1	0.4	0.7	0.6	0.7	1.5	3.5	10.3	31.5	132.9
2000	6.5	4.8	0.9	0.3	0.8	0.6	0.7	1.1	3.6	10.2	32.4	130.5
1999	6.7	5.2	0.7	0.4	0.8	0.6	0.7	1.3	3.6	10.0	31.7	140.3
Non-Hispanic white, both sexes												
2002	6.6	5.3	0.9	0.5	0.8	0.8	0.9	2.1	5.7	12.8	34.9	115.5
2001	6.6	5.7	1.2	0.5	0.9	0.7	0.9	2.2	5.0	13.2	34.0	125.3
2000	6.5	5.1	1.0	0.5	0.9	0.7	0.9	2.0	5.1	12.7	34.0	125.3
1999	6.4	4.7	1.0	0.5	0.9	0.8	0.8	1.8	4.8	12.1	33.1	132.8
Non-Hispanic white, male												
2002	5.9	6.9	0.7	0.5	0.8	1.0	1.2	2.9	7.4	15.9	38.5	100.9
2001	5.9	6.1	1.3	0.4	0.8	0.7	1.1	3.0	6.7	16.8	38.5	109.9
2000	5.7	5.8	1.1	0.6	0.8	0.8	1.0	3.0	6.9	16.0	36.9	115.8
1999	5.3	5.2	1.1	0.5	0.9	0.7	0.9	2.4	6.2	15.0	35.2	110.3
Non-Hispanic white, female												
2002	7.3	3.5	1.0	0.4	0.9	0.7	0.7	1.3	4.0	10.1	32.5	121.9
2001	7.4	5.3	1.1	0.5	0.9	0.7	0.7	1.5	3.5	10.2	31.1	131.6
2000	7.2	4.4	0.9	0.4	1.0	0.7	0.7	1.0	3.4	10.0	32.1	129.1
1999	7.4	4.2	0.7	0.5	1.0	0.8	0.6	1.1	3.4	9.8	31.7	141.8
Non-Hispanic black, both sexes												
2002	5.0	7.6	0.8	0.3	0.4	0.4	1.1	3.2	8.5	19.6	47.8	137.3
2001	4.7	9.3	*	*	*	0.4	0.7	3.3	7.1	17.4	46.7	153.6
2000	4.6	6.6	*	*	0.4	0.7	1.1	2.8	7.4	16.7	44.4	147.2
1999	4.5	9.7	*	0.4	*	0.5	1.4	3.0	6.5	18.0	41.3	139.7
Non-Hispanic black, male												
2002	4.6	7.0	*	*	0.7	*	1.3	4.3	12.2	22.1	52.8	136.7
2001	4.4	9.6	*	*	*	*	0.8	4.1	10.0	23.4	56.1	132.8
2000	4.1	7.8	*	*	*	0.9	1.3	4.0	9.5	20.4	50.6	116.2
1999	4.2	10.7	*	*	*	*	1.6	3.6	7.9	22.4	48.8	148.6
Non-Hispanic black, female												
2002	5.3	8.2	*	*	*	*	0.9	2.3	5.5	17.9	44.9	137.5
2001	5.1	9.1	*	*	*	*	*	2.6	4.8	13.1	41.4	161.3
2000	5.0	*	*	*	*	*	0.9	1.9	5.6	14.1	41.0	158.7
1999	4.8	8.6	*	*	*	*	1.2	2.5	5.4	14.9	37.1	136.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cystic fibrosis (E84)												
All origins 3/, both sexes												
2002	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
2001	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
2000	0.2	*	*	0.1	0.4	0.3	0.2	0.1	*	*	*	*
1999	0.2	*	*	0.1	0.5	0.3	0.1	0.1	*	*	*	*
All origins 3/, male												
2002	0.2	*	*	*	0.3	0.4	0.2	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.3	0.2	0.1	*	*	*	*
2000	0.2	*	*	0.1	0.4	0.3	0.2	*	*	*	*	*
1999	0.2	*	*	0.1	0.4	0.4	0.2	*	*	*	*	*
All origins 3/, female												
2002	0.2	*	*	0.1	0.4	0.3	0.2	*	*	*	*	*
2001	0.2	*	*	0.1	0.5	0.3	0.1	0.1	*	*	*	*
2000	0.2	*	*	0.1	0.5	0.4	0.2	*	*	*	*	*
1999	0.2	*	*	0.1	0.5	0.3	0.1	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	0.1	0.4	0.4	0.2	0.1	*	*	*	*
2001	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
2000	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
1999	0.2	*	*	0.1	0.5	0.4	0.2	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.2	*	*	*	0.4	0.5	0.2	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.4	0.2	0.1	*	*	*	*
2000	0.2	*	*	*	0.4	0.4	0.2	*	*	*	*	*
1999	0.2	*	*	*	0.5	0.4	0.2	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cystic fibrosis (E84)												
Non-Hispanic 4/, female												
2002	0.2	*	*	0.1	0.5	0.3	0.2	*	*	*	*	*
2001	0.2	*	*	*	0.6	0.4	0.1	*	*	*	*	*
2000	0.2	*	*	0.1	0.5	0.4	0.3	*	*	*	*	*
1999	0.2	*	*	0.2	0.6	0.4	0.1	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	0.1	0.6	0.5	0.2	0.1	*	*	*	*
2001	0.2	*	*	0.1	0.6	0.5	0.2	0.1	*	*	*	*
2000	0.2	*	*	0.1	0.6	0.5	0.3	0.1	*	*	*	*
1999	0.2	*	*	0.2	0.7	0.5	0.2	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.2	*	*	*	0.5	0.6	0.3	*	*	*	*	*
2001	0.2	*	*	*	0.5	0.5	0.3	0.1	*	*	*	*
2000	0.2	*	*	*	0.6	0.5	0.3	*	*	*	*	*
1999	0.2	*	*	*	0.6	0.5	0.2	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	*	*	*	0.7	0.4	0.2	*	*	*	*	*
2001	0.2	*	*	*	0.7	0.4	0.2	*	*	*	*	*
2000	0.2	*	*	0.2	0.7	0.5	0.3	*	*	*	*	*
1999	0.2	*	*	0.2	0.8	0.5	0.2	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)												
All origins 3/, both sexes												
2002	2.5	1.8	*	*	0.1	0.1	0.2	0.5	1.2	3.8	16.0	83.2
2001	2.7	2.2	0.2	*	*	0.1	0.2	0.5	1.2	4.1	16.8	93.9
2000	2.8	1.9	*	*	0.1	0.1	0.3	0.5	1.3	4.0	17.4	96.8
1999	2.9	1.9	0.1	*	0.1	0.1	0.3	0.5	1.1	4.2	18.2	106.4
All origins 3/, male												
2002	1.7	2.1	*	*	*	*	0.2	0.6	1.3	3.8	15.3	67.9
2001	1.8	2.2	*	*	*	*	0.2	0.6	1.3	4.1	16.8	76.5
2000	2.0	1.8	*	*	*	*	0.3	0.6	1.5	4.5	17.7	84.3
1999	2.0	2.3	*	*	*	*	0.3	0.7	1.2	4.5	18.4	85.0
All origins 3/, female												
2002	3.3	1.5	*	*	*	0.1	0.2	0.5	1.2	3.7	16.5	89.8
2001	3.6	2.2	*	*	*	0.1	0.2	0.5	1.1	4.1	16.8	101.2
2000	3.5	2.0	*	*	*	*	0.2	0.4	1.2	3.6	17.2	101.9
1999	3.8	1.5	*	*	*	*	0.2	0.4	1.0	3.9	18.1	115.1
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	*	*	2.5	9.7	33.3
2001	0.5	*	*	*	*	*	*	*	*	3.1	9.2	38.8
2000	0.5	*	*	*	*	*	*	*	*	2.1	9.7	46.4
1999	0.6	*	*	*	*	*	*	*	*	2.1	12.3	54.7
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	10.7	33.5
2001	0.4	*	*	*	*	*	*	*	*	*	9.8	45.7
2000	0.4	*	*	*	*	*	*	*	*	*	11.3	*
1999	0.5	*	*	*	*	*	*	*	*	*	16.7	*
Hispanic, female												
2002	0.6	*	*	*	*	*	*	*	*	*	9.0	33.2
2001	0.6	*	*	*	*	*	*	*	*	4.3	8.8	35.4
2000	0.7	*	*	*	*	*	*	*	*	*	8.6	51.4
1999	0.7	*	*	*	*	*	*	*	*	*	9.4	63.4
Non-Hispanic 4/, both sexes												
2002	2.9	2.0	*	*	*	0.1	0.2	0.5	1.3	3.8	16.3	84.9
2001	3.1	2.4	0.2	*	*	0.1	0.3	0.6	1.2	4.1	17.1	95.8
2000	3.1	2.1	*	*	0.1	0.1	0.3	0.5	1.4	4.1	17.7	98.4
1999	3.2	2.0	*	*	0.1	0.1	0.3	0.5	1.1	4.3	18.4	108.0
Non-Hispanic 4/, male												
2002	1.9	2.6	*	*	*	*	0.2	0.6	1.4	3.9	15.5	69.0
2001	2.1	2.4	*	*	*	*	0.3	0.6	1.3	4.2	17.1	77.4
2000	2.2	2.2	*	*	*	*	0.3	0.6	1.5	4.6	18.0	86.0
1999	2.2	2.3	*	*	*	*	0.3	0.7	1.2	4.7	18.5	86.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)												
Non-Hispanic 4/, female												
2002	3.7	1.4	*	*	*	*	0.2	0.5	1.3	3.8	16.8	91.8
2001	4.0	2.4	*	*	*	0.1	0.3	0.5	1.1	4.0	17.1	103.5
2000	3.9	2.0	*	*	*	*	0.2	0.4	1.2	3.7	17.6	103.3
1999	4.2	1.8	*	*	*	*	0.2	0.4	1.1	4.0	18.4	116.5
Non-Hispanic white, both sexes												
2002	3.0	1.5	*	*	*	0.1	0.2	0.4	1.2	3.5	15.4	84.2
2001	3.3	1.7	0.2	*	*	0.1	0.2	0.4	1.0	3.7	16.3	94.9
2000	3.3	1.5	*	*	*	*	0.2	0.4	1.2	3.8	17.2	97.7
1999	3.5	1.3	*	*	*	0.1	0.2	0.4	0.9	3.9	18.0	108.3
Non-Hispanic white, male												
2002	2.0	2.2	*	*	*	*	0.2	0.5	1.2	3.4	14.5	67.2
2001	2.1	*	*	*	*	*	0.2	0.4	1.1	3.7	16.0	76.7
2000	2.3	*	*	*	*	*	0.2	0.5	1.3	4.2	17.1	86.8
1999	2.2	*	*	*	*	*	0.2	0.6	1.0	3.9	17.4	84.3
Non-Hispanic white, female												
2002	4.1	*	*	*	*	*	0.2	0.4	1.2	3.5	16.0	91.5
2001	4.4	1.9	*	*	*	*	0.3	0.4	1.0	3.7	16.6	102.5
2000	4.3	*	*	*	*	*	0.2	0.3	1.1	3.4	17.3	102.1
1999	4.7	*	*	*	*	*	0.2	0.3	0.9	3.8	18.3	117.9
Non-Hispanic black, both sexes												
2002	2.6	3.6	*	*	*	*	0.6	1.3	2.6	8.1	28.8	106.2
2001	2.9	5.4	*	*	*	*	0.4	1.5	2.9	8.5	29.6	121.1
2000	2.8	3.5	*	*	*	*	0.6	1.4	2.8	8.4	27.6	119.9
1999	2.8	5.1	*	*	*	*	0.8	1.4	2.9	9.4	26.5	115.8
Non-Hispanic black, male												
2002	2.2	*	*	*	*	*	*	1.8	3.4	8.8	30.5	101.7
2001	2.3	*	*	*	*	*	*	1.7	3.8	10.4	35.4	99.3
2000	2.3	*	*	*	*	*	0.8	1.8	3.8	10.0	34.2	98.6
1999	2.6	*	*	*	*	*	1.2	1.7	3.5	13.2	33.1	126.0
Non-Hispanic black, female												
2002	3.1	*	*	*	*	*	*	0.9	1.9	7.7	27.9	107.9
2001	3.3	*	*	*	*	*	*	1.4	2.2	7.1	26.3	129.2
2000	3.2	*	*	*	*	*	*	1.1	2.0	7.3	23.8	127.8
1999	3.0	*	*	*	*	*	*	1.2	2.5	6.8	22.8	111.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)												
All origins 3/, both sexes												
2002	2.8	3.4	0.7	0.3	0.2	0.2	0.5	1.5	4.3	9.2	19.1	31.1
2001	2.6	3.7	0.8	0.3	0.2	0.1	0.4	1.5	3.8	9.0	17.2	30.5
2000	2.4	3.2	0.8	0.3	0.2	0.2	0.3	1.4	3.8	8.6	16.4	27.3
1999	2.2	3.4	0.7	0.3	0.2	0.1	0.4	1.2	3.6	8.0	14.8	24.3
All origins 3/, male												
2002	3.0	3.6	0.7	0.4	0.3	0.2	0.6	2.2	6.0	12.1	23.4	33.6
2001	2.9	4.1	0.9	0.3	0.2	0.2	0.5	2.2	5.5	12.6	21.7	33.3
2000	2.6	3.6	0.8	0.3	0.2	0.3	0.4	2.2	5.3	11.3	19.0	27.7
1999	2.4	3.7	0.8	0.3	0.2	0.1	0.5	1.7	4.8	10.5	17.3	25.5
All origins 3/, female												
2002	2.5	3.2	0.7	0.2	0.1	0.1	0.3	0.8	2.8	6.7	16.3	30.0
2001	2.3	3.2	0.8	0.3	0.1	0.1	0.3	0.9	2.3	6.1	14.3	29.3
2000	2.3	2.8	0.7	0.2	0.2	*	0.2	0.6	2.3	6.3	14.8	27.1
1999	2.1	3.1	0.6	0.3	0.2	0.1	0.3	0.8	2.4	5.9	13.2	23.9
Hispanic, both sexes												
2002	1.0	2.9	0.7	*	*	*	*	0.7	2.0	6.2	16.3	32.2
2001	0.9	3.1	0.7	*	*	*	*	*	2.8	5.3	13.2	28.8
2000	0.9	3.0	0.7	*	*	*	*	0.8	2.0	5.9	11.7	26.5
1999	0.8	3.4	*	*	*	*	*	*	2.0	5.6	10.8	25.3
Hispanic, male												
2002	1.0	*	*	*	*	*	*	*	2.1	8.0	18.9	38.5
2001	1.0	*	*	*	*	*	*	*	4.0	8.4	15.2	*
2000	0.9	*	*	*	*	*	*	*	2.9	8.9	*	*
1999	0.8	*	*	*	*	*	*	*	2.6	7.0	11.5	*
Hispanic, female												
2002	0.9	*	*	*	*	*	*	*	*	4.7	14.6	29.0
2001	0.8	*	*	*	*	*	*	*	*	*	11.8	27.9
2000	0.8	*	*	*	*	*	*	*	*	3.5	13.2	27.7
1999	0.8	*	*	*	*	*	*	*	*	4.5	10.4	26.0
Non-Hispanic 4/, both sexes												
2002	3.0	3.6	0.7	0.3	0.2	0.2	0.5	1.6	4.5	9.4	19.2	30.9
2001	2.9	3.8	0.8	0.3	0.2	0.1	0.4	1.6	3.9	9.3	17.3	30.4
2000	2.7	3.3	0.8	0.3	0.2	0.2	0.4	1.5	3.9	8.7	16.6	27.2
1999	2.4	3.3	0.7	0.3	0.2	0.2	0.4	1.3	3.7	8.1	15.0	24.3
Non-Hispanic 4/, male												
2002	3.3	3.8	0.6	0.4	0.3	0.2	0.7	2.3	6.3	12.3	23.5	33.2
2001	3.2	4.4	0.9	0.3	0.2	0.2	0.6	2.3	5.6	12.8	21.9	33.2
2000	2.9	3.9	0.8	0.3	0.3	0.3	0.5	2.3	5.5	11.4	19.4	27.7
1999	2.6	3.5	0.9	0.3	0.2	0.2	0.5	1.8	5.0	10.7	17.5	25.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)												
Non-Hispanic 4/, female												
2002	2.7	3.3	0.7	0.2	0.1	0.2	0.3	0.9	2.8	6.9	16.3	29.9
2001	2.5	3.3	0.7	0.3	0.1	0.1	0.3	0.9	2.4	6.3	14.3	29.3
2000	2.4	2.6	0.7	0.2	0.2	*	0.2	0.7	2.4	6.5	14.8	27.0
1999	2.3	3.0	0.5	0.3	0.2	0.2	0.3	0.8	2.5	6.0	13.3	23.8
Non-Hispanic white, both sexes												
2002	3.3	3.5	0.7	0.3	0.2	0.2	0.5	1.6	4.5	9.3	19.4	31.3
2001	3.2	3.8	0.9	0.3	0.2	0.2	0.5	1.7	4.0	9.5	17.6	30.3
2000	2.9	3.4	0.8	0.3	0.2	0.2	0.4	1.5	3.9	8.9	16.8	27.4
1999	2.7	2.9	0.8	0.3	0.2	0.2	0.4	1.3	3.8	8.3	15.1	24.5
Non-Hispanic white, male												
2002	3.7	4.3	0.6	0.4	0.3	0.2	0.7	2.3	6.2	12.5	23.8	33.6
2001	3.5	4.2	1.0	0.3	0.3	0.2	0.6	2.4	5.6	13.1	22.4	33.2
2000	3.2	4.0	0.9	0.4	0.2	0.2	0.5	2.4	5.6	11.7	19.8	29.0
1999	2.9	3.2	1.0	0.3	0.2	*	0.5	1.8	5.2	11.0	17.7	25.9
Non-Hispanic white, female												
2002	3.0	2.6	0.8	0.2	*	0.2	0.3	0.8	2.8	6.6	16.4	30.3
2001	2.8	3.3	0.7	0.3	*	*	0.3	0.9	2.4	6.5	14.4	29.1
2000	2.7	2.7	0.7	0.2	0.2	*	0.2	0.7	2.3	6.6	14.8	26.8
1999	2.5	2.5	0.6	0.3	0.2	0.2	0.2	0.8	2.5	5.9	13.3	23.9
Non-Hispanic black, both sexes												
2002	2.2	3.9	*	*	*	*	0.5	1.9	5.8	11.4	19.0	31.2
2001	1.9	3.9	*	*	*	*	*	1.7	4.2	8.9	17.2	32.5
2000	1.8	*	*	*	*	*	0.4	1.3	4.5	8.3	16.9	27.3
1999	1.7	4.6	*	0.3	*	*	0.5	1.5	3.6	8.6	14.8	23.9
Non-Hispanic black, male												
2002	2.3	*	*	*	*	*	*	2.5	8.7	13.2	22.3	35.0
2001	2.0	*	*	*	*	*	*	2.3	6.2	13.1	20.7	33.5
2000	1.8	*	*	*	*	*	*	2.1	5.7	10.3	16.4	*
1999	1.6	*	*	*	*	*	*	1.9	4.5	9.2	15.6	*
Non-Hispanic black, female												
2002	2.1	*	*	*	*	*	*	1.4	3.6	10.2	17.1	29.7
2001	1.7	*	*	*	*	*	*	1.2	2.6	6.0	15.2	32.1
2000	1.8	*	*	*	*	*	*	*	3.6	6.8	17.2	31.0
1999	1.8	*	*	*	*	*	0.7	1.2	2.9	8.2	14.4	24.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)												
All origins 3/, both sexes												
2002	19.3	*	*	0.1	0.6	1.7	5.3	8.1	8.8	19.0	107.6	646.7
2001	17.8	*	*	0.1	0.6	1.8	4.9	7.8	8.4	18.2	99.0	604.9
2000	16.4	*	*	0.1	0.6	1.8	4.9	7.3	8.1	16.9	89.9	561.6
1999	15.1	*	*	0.1	0.6	1.8	5.1	7.1	8.0	16.0	81.3	515.3
All origins 3/, male												
2002	14.8	*	*	*	0.8	2.3	7.5	12.7	13.5	23.9	106.9	506.0
2001	13.8	*	*	*	0.8	2.6	7.0	12.2	12.8	23.3	97.3	488.0
2000	12.8	*	*	*	0.9	2.6	7.2	11.5	12.4	21.9	89.9	451.0
1999	12.3	*	*	0.1	0.8	2.6	7.7	11.3	12.6	21.2	85.5	415.8
All origins 3/, female												
2002	23.7	*	*	*	0.4	1.0	3.0	3.7	4.5	14.9	108.0	707.8
2001	21.7	*	*	*	0.3	1.0	2.8	3.6	4.3	13.9	100.1	653.8
2000	19.8	*	*	*	0.4	1.0	2.5	3.3	4.2	12.9	90.0	606.6
1999	17.8	*	*	*	0.3	1.0	2.5	3.1	3.8	11.8	78.6	555.4
Hispanic, both sexes												
2002	5.1	*	*	*	0.6	1.6	4.6	7.1	8.2	13.2	59.2	332.6
2001	4.9	*	*	*	0.7	1.5	4.2	6.9	8.4	12.6	54.6	372.7
2000	4.6	*	*	*	0.7	1.9	4.8	6.6	8.1	11.1	50.0	307.2
1999	4.8	*	*	*	0.7	2.3	4.7	7.8	8.3	11.3	47.7	343.1
Hispanic, male												
2002	6.0	*	*	*	1.0	2.5	7.3	12.3	13.9	21.6	68.1	331.3
2001	5.4	*	*	*	1.0	2.1	6.6	12.0	15.1	19.6	57.2	327.6
2000	5.5	*	*	*	1.1	2.9	8.0	11.1	14.4	20.0	48.2	298.3
1999	5.8	*	*	*	1.2	3.7	8.0	13.3	15.4	17.8	51.6	325.7
Hispanic, female												
2002	4.2	*	*	*	*	0.7	1.7	1.9	3.0	6.5	53.1	333.2
2001	4.4	*	*	*	*	0.8	1.6	2.0	2.5	7.1	52.9	394.7
2000	3.6	*	*	*	*	0.8	1.3	2.1	2.4	4.0	51.2	311.6
1999	3.8	*	*	*	*	0.8	1.2	2.5	*	6.2	45.1	351.5
Non-Hispanic 4/, both sexes												
2002	21.5	*	*	0.1	0.6	1.6	5.3	8.1	8.8	19.3	109.5	658.0
2001	19.7	*	*	0.1	0.6	1.9	4.9	7.8	8.3	18.5	100.9	612.1
2000	18.0	*	*	0.1	0.6	1.8	4.8	7.3	8.1	17.2	91.4	569.7
1999	16.4	*	*	0.1	0.5	1.7	5.0	7.0	7.9	16.2	82.5	520.1
Non-Hispanic 4/, male												
2002	16.2	*	*	*	0.8	2.2	7.4	12.5	13.4	23.9	108.5	512.8
2001	15.1	*	*	*	0.8	2.8	6.9	12.1	12.5	23.5	99.0	493.5
2000	13.8	*	*	*	0.8	2.5	7.0	11.3	12.1	21.8	91.4	456.4
1999	13.1	*	*	*	0.7	2.4	7.5	11.0	12.2	21.2	86.7	418.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)												
Non-Hispanic 4/, female												
2002	26.5	*	*	*	0.4	1.1	3.2	3.8	4.6	15.5	110.2	720.6
2001	24.1	*	*	*	0.3	1.0	2.9	3.7	4.4	14.3	102.1	661.4
2000	21.9	*	*	*	0.4	1.0	2.6	3.4	4.3	13.4	91.4	615.5
1999	19.6	*	*	*	0.3	1.0	2.6	3.2	3.9	12.1	79.8	560.8
Non-Hispanic white, both sexes												
2002	24.1	*	*	*	0.6	1.7	5.1	7.5	8.1	18.9	111.0	675.7
2001	22.0	*	*	*	0.6	1.9	4.5	6.8	7.6	18.1	102.8	629.4
2000	19.9	*	*	*	0.6	1.7	4.5	6.5	7.3	16.7	92.3	586.2
1999	18.2	*	*	0.1	0.5	1.7	4.5	6.0	7.1	15.7	83.6	540.2
Non-Hispanic white, male												
2002	17.4	*	*	*	0.8	2.2	7.2	11.5	11.9	22.4	108.2	522.2
2001	16.1	*	*	*	0.8	2.7	6.4	10.5	11.0	22.3	99.7	504.2
2000	14.6	*	*	*	0.8	2.4	6.5	10.0	10.7	20.5	91.2	464.8
1999	13.8	*	*	*	0.7	2.5	6.8	9.5	10.7	20.0	86.1	432.2
Non-Hispanic white, female												
2002	30.5	*	*	*	0.4	1.1	2.9	3.6	4.5	16.0	112.8	742.0
2001	27.6	*	*	*	0.3	1.0	2.6	3.2	4.4	14.5	104.8	681.5
2000	25.0	*	*	*	0.5	1.0	2.4	3.0	4.2	13.5	93.0	635.1
1999	22.5	*	*	*	0.3	1.0	2.2	2.6	3.7	12.1	81.9	583.2
Non-Hispanic black, both sexes												
2002	13.7	*	*	*	0.5	1.9	7.4	13.9	16.4	26.7	117.0	548.7
2001	12.9	*	*	*	0.6	2.0	7.8	15.9	15.7	26.6	102.1	492.3
2000	12.2	*	*	*	0.7	2.4	7.4	14.6	15.4	25.6	99.2	448.1
1999	11.0	*	*	*	0.6	1.9	8.8	14.8	16.1	24.4	85.6	350.5
Non-Hispanic black, male												
2002	13.5	*	*	*	*	2.5	10.0	22.7	28.8	42.6	137.5	461.6
2001	13.4	*	*	*	0.9	3.1	11.1	26.1	28.3	40.6	118.2	427.3
2000	12.9	*	*	*	1.1	3.5	10.9	24.2	26.5	40.1	114.8	421.5
1999	12.2	*	*	*	0.8	2.6	12.7	23.4	28.2	38.0	112.3	312.6
Non-Hispanic black, female												
2002	13.8	*	*	*	*	1.3	5.0	6.3	6.5	15.4	105.4	581.9
2001	12.5	*	*	*	*	1.0	4.8	7.1	5.6	16.6	93.0	516.3
2000	11.6	*	*	*	*	1.5	4.3	6.4	6.5	15.4	90.5	457.9
1999	9.9	*	*	*	*	1.3	5.3	7.4	6.4	14.8	70.7	364.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Organic dementia (F01,F03)												
All origins 3/, both sexes												
2002	14.5	*	*	*	*	*	*	0.1	1.1	10.9	94.3	598.3
2001	13.1	*	*	*	*	*	*	0.1	1.1	10.5	84.9	554.1
2000	11.7	*	*	*	*	*	*	0.1	1.0	9.3	76.0	505.6
1999	10.3	*	*	*	*	*	*	0.1	0.9	8.3	65.8	454.4
All origins 3/, male												
2002	8.7	*	*	*	*	*	*	0.1	1.1	12.0	91.4	465.7
2001	7.8	*	*	*	*	*	*	0.2	1.2	11.5	81.1	442.4
2000	6.9	*	*	*	*	*	*	0.1	1.1	10.4	73.7	398.9
1999	6.2	*	*	*	*	*	*	0.1	1.1	9.8	66.4	361.7
All origins 3/, female												
2002	20.1	*	*	*	*	*	*	0.1	1.0	10.0	96.3	655.9
2001	18.2	*	*	*	*	*	*	0.1	1.0	9.6	87.3	600.8
2000	16.3	*	*	*	*	*	*	0.1	0.9	8.3	77.5	549.1
1999	14.2	*	*	*	*	*	*	*	0.7	7.1	65.5	491.7
Hispanic, both sexes												
2002	2.3	*	*	*	*	*	*	*	*	5.8	48.5	293.8
2001	2.4	*	*	*	*	*	*	*	*	6.0	44.7	338.9
2000	1.9	*	*	*	*	*	*	*	*	4.6	40.9	268.1
1999	1.9	*	*	*	*	*	*	*	*	4.5	35.2	295.4
Hispanic, male												
2002	1.7	*	*	*	*	*	*	*	*	7.6	50.5	291.1
2001	1.6	*	*	*	*	*	*	*	*	7.8	41.1	300.9
2000	1.4	*	*	*	*	*	*	*	*	7.2	38.4	266.0
1999	1.3	*	*	*	*	*	*	*	*	4.8	34.4	280.4
Hispanic, female												
2002	3.0	*	*	*	*	*	*	*	*	4.4	47.2	295.1
2001	3.2	*	*	*	*	*	*	*	*	4.6	47.1	357.4
2000	2.5	*	*	*	*	*	*	*	*	*	42.6	269.1
1999	2.6	*	*	*	*	*	*	*	*	4.3	35.7	302.6
Non-Hispanic 4/, both sexes												
2002	16.4	*	*	*	*	*	*	0.1	1.1	11.2	96.3	609.3
2001	14.7	*	*	*	*	*	*	0.1	1.1	10.7	86.6	560.9
2000	13.0	*	*	*	*	*	*	0.1	1.0	9.5	77.3	513.2
1999	11.4	*	*	*	*	*	*	0.1	0.9	8.5	66.9	458.9
Non-Hispanic 4/, male												
2002	9.8	*	*	*	*	*	*	0.1	1.1	12.3	93.2	472.6
2001	8.8	*	*	*	*	*	*	0.1	1.2	11.7	82.8	447.3
2000	7.7	*	*	*	*	*	*	0.1	1.1	10.6	75.0	403.5
1999	6.9	*	*	*	*	*	*	0.1	1.1	10.0	67.6	364.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Organic dementia (F01,F03)												
Non-Hispanic 4/, female												
2002	22.6	*	*	*	*	*	*	0.1	1.1	10.4	98.3	668.3
2001	20.3	*	*	*	*	*	*	0.1	1.0	9.9	89.0	608.1
2000	18.1	*	*	*	*	*	*	0.1	0.9	8.7	78.8	557.6
1999	15.7	*	*	*	*	*	*	*	0.7	7.3	66.5	496.8
Non-Hispanic white, both sexes												
2002	18.8	*	*	*	*	*	*	0.1	1.1	11.2	97.5	625.6
2001	16.9	*	*	*	*	*	*	0.1	1.1	10.6	88.2	576.4
2000	14.9	*	*	*	*	*	*	0.1	0.9	9.3	78.0	527.7
1999	13.2	*	*	*	*	*	*	0.1	0.9	8.4	67.9	476.4
Non-Hispanic white, male												
2002	11.1	*	*	*	*	*	*	*	1.2	11.9	93.0	481.1
2001	9.9	*	*	*	*	*	*	*	1.2	11.3	83.6	457.4
2000	8.6	*	*	*	*	*	*	*	1.0	10.0	74.9	410.6
1999	7.8	*	*	*	*	*	*	*	1.1	9.8	67.5	376.4
Non-Hispanic white, female												
2002	26.3	*	*	*	*	*	*	0.2	1.1	10.6	100.4	688.0
2001	23.5	*	*	*	*	*	*	0.1	1.1	10.0	91.2	626.0
2000	20.9	*	*	*	*	*	*	*	0.9	8.6	79.9	574.9
1999	18.3	*	*	*	*	*	*	*	0.7	7.3	68.1	516.2
Non-Hispanic black, both sexes												
2002	8.2	*	*	*	*	*	*	*	1.3	14.6	103.6	510.3
2001	7.2	*	*	*	*	*	*	*	1.5	15.0	87.9	455.7
2000	6.6	*	*	*	*	*	*	*	1.6	14.0	84.3	408.0
1999	5.1	*	*	*	*	*	*	*	1.1	11.4	68.1	310.7
Non-Hispanic black, male												
2002	5.5	*	*	*	*	*	*	*	*	19.0	117.5	428.8
2001	4.8	*	*	*	*	*	*	*	2.0	19.4	96.3	385.8
2000	4.6	*	*	*	*	*	*	*	2.0	18.4	92.9	378.1
1999	3.7	*	*	*	*	*	*	*	*	14.3	81.9	268.6
Non-Hispanic black, female												
2002	10.6	*	*	*	*	*	*	*	1.4	11.4	95.7	541.4
2001	9.4	*	*	*	*	*	*	*	*	11.8	83.2	481.6
2000	8.4	*	*	*	*	*	*	*	*	10.8	79.5	419.1
1999	6.5	*	*	*	*	*	*	*	*	9.3	60.4	326.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified organic mental disorders (F04-F09)												
All origins 3/, both sexes												
2002	0.7	*	*	*	*	*	*	0.1	0.2	0.8	4.1	26.7
2001	0.8	*	*	*	*	*	*	*	0.2	0.8	4.9	29.3
2000	0.8	*	*	*	*	*	*	*	0.2	0.8	5.0	33.7
1999	0.9	*	*	*	*	*	*	*	0.2	0.8	5.5	37.7
All origins 3/, male												
2002	0.4	*	*	*	*	*	*	*	0.2	1.0	4.2	20.1
2001	0.5	*	*	*	*	*	*	*	0.2	1.0	4.8	23.8
2000	0.5	*	*	*	*	*	*	*	0.2	1.0	4.5	28.4
1999	0.6	*	*	*	*	*	*	*	0.2	0.9	5.6	29.8
All origins 3/, female												
2002	0.9	*	*	*	*	*	*	*	0.1	0.7	4.1	29.6
2001	1.0	*	*	*	*	*	*	*	*	0.7	5.0	31.6
2000	1.1	*	*	*	*	*	*	*	0.2	0.7	5.2	35.8
1999	1.2	*	*	*	*	*	*	*	*	0.7	5.4	40.9
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	5.0	26.6
2001	0.2	*	*	*	*	*	*	*	*	*	3.8	25.0
2000	0.2	*	*	*	*	*	*	*	*	*	*	29.2
1999	0.3	*	*	*	*	*	*	*	*	*	5.4	35.8
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	28.2
2001	0.3	*	*	*	*	*	*	*	*	*	*	28.9
2000	0.3	*	*	*	*	*	*	*	*	*	*	33.6
1999	0.3	*	*	*	*	*	*	*	*	*	*	37.4
Non-Hispanic 4/, both sexes												
2002	0.8	*	*	*	*	*	*	0.1	0.2	0.8	4.0	26.7
2001	0.8	*	*	*	*	*	*	*	0.1	0.8	5.0	29.4
2000	0.9	*	*	*	*	*	*	*	0.2	0.8	5.0	33.7
1999	1.0	*	*	*	*	*	*	*	0.2	0.8	5.5	37.7
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	*	*	0.2	1.0	4.1	19.9
2001	0.5	*	*	*	*	*	*	*	0.2	1.0	4.8	23.9
2000	0.6	*	*	*	*	*	*	*	0.2	1.0	4.7	28.7
1999	0.6	*	*	*	*	*	*	*	0.2	0.9	5.6	29.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified organic mental disorders (F04-F09)												
Non-Hispanic 4/, female												
2002	1.0	*	*	*	*	*	*	*	*	0.7	4.0	29.6
2001	1.1	*	*	*	*	*	*	*	*	0.7	5.1	31.7
2000	1.2	*	*	*	*	*	*	*	0.2	0.8	5.2	35.7
1999	1.3	*	*	*	*	*	*	*	*	0.7	5.4	40.9
Non-Hispanic white, both sexes												
2002	0.9	*	*	*	*	*	*	*	0.2	0.8	4.2	27.5
2001	1.0	*	*	*	*	*	*	*	0.2	0.8	5.1	30.3
2000	1.0	*	*	*	*	*	*	*	0.2	0.9	5.1	34.7
1999	1.1	*	*	*	*	*	*	*	0.2	0.8	5.5	38.9
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	*	*	*	0.9	4.2	20.4
2001	0.6	*	*	*	*	*	*	*	*	1.0	4.8	24.3
2000	0.6	*	*	*	*	*	*	*	0.2	0.9	4.8	28.8
1999	0.7	*	*	*	*	*	*	*	0.2	0.8	5.4	30.0
Non-Hispanic white, female												
2002	1.2	*	*	*	*	*	*	*	*	0.6	4.1	30.5
2001	1.3	*	*	*	*	*	*	*	*	0.7	5.2	32.7
2000	1.4	*	*	*	*	*	*	*	*	0.8	5.3	37.1
1999	1.5	*	*	*	*	*	*	*	*	0.7	5.5	42.4
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	*	*	3.4	23.3
2001	0.5	*	*	*	*	*	*	*	*	1.2	5.5	24.7
2000	0.5	*	*	*	*	*	*	*	*	*	5.6	25.8
1999	0.5	*	*	*	*	*	*	*	*	1.4	6.7	28.8
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	6.3	23.1
2000	0.4	*	*	*	*	*	*	*	*	*	*	25.8
1999	0.5	*	*	*	*	*	*	*	*	*	9.7	29.7
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.6	25.5
2001	0.6	*	*	*	*	*	*	*	*	*	5.1	25.2
2000	0.6	*	*	*	*	*	*	*	*	*	5.7	25.7
1999	0.6	*	*	*	*	*	*	*	*	*	5.1	28.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to psychoactive substance use (F10-F19)												
All origins 3/, both sexes												
2002	3.3	*	*	*	0.5	1.5	4.9	7.4	6.8	5.7	4.4	3.7
2001	3.2	*	*	*	0.5	1.6	4.5	7.3	6.5	5.4	4.6	4.0
2000	3.1	*	*	*	0.5	1.6	4.6	6.8	6.2	5.5	4.7	4.1
1999	3.1	*	*	*	0.4	1.6	4.7	6.6	6.4	5.4	4.9	3.8
All origins 3/, male												
2002	5.1	*	*	*	0.7	2.1	7.1	11.9	11.4	9.3	7.2	7.0
2001	4.9	*	*	*	0.7	2.4	6.6	11.6	10.7	9.2	7.2	7.5
2000	4.8	*	*	*	0.7	2.4	6.9	10.9	10.3	9.1	8.0	7.3
1999	4.8	*	*	*	0.7	2.4	7.3	10.6	10.6	8.9	8.8	7.4
All origins 3/, female												
2002	1.6	*	*	*	0.2	0.8	2.7	3.1	2.6	2.7	2.5	2.2
2001	1.5	*	*	*	0.2	0.8	2.4	3.1	2.6	2.3	2.9	2.5
2000	1.5	*	*	*	0.2	0.8	2.2	2.8	2.5	2.5	2.5	2.8
1999	1.4	*	*	*	0.2	0.8	2.2	2.7	2.5	2.6	2.4	2.4
Hispanic, both sexes												
2002	2.3	*	*	*	0.6	1.6	4.5	6.8	6.8	6.0	4.3	*
2001	2.2	*	*	*	0.6	1.4	4.1	6.7	7.2	4.9	3.8	*
2000	2.3	*	*	*	0.7	1.8	4.7	6.3	6.6	5.5	4.0	*
1999	2.4	*	*	*	0.7	2.1	4.6	7.5	7.0	5.5	4.2	*
Hispanic, male												
2002	3.9	*	*	*	0.9	2.4	7.2	12.0	12.6	11.7	9.0	*
2001	3.5	*	*	*	0.9	2.0	6.5	11.6	13.6	9.8	*	*
2000	3.8	*	*	*	1.0	2.7	7.8	10.8	12.7	11.0	*	*
1999	4.1	*	*	*	1.2	3.5	7.8	13.0	13.4	11.2	*	*
Hispanic, female												
2002	0.7	*	*	*	*	*	1.6	1.7	*	*	*	*
2001	0.7	*	*	*	*	0.7	1.5	1.8	*	*	*	*
2000	0.7	*	*	*	*	0.8	1.3	2.0	*	*	*	*
1999	0.6	*	*	*	*	*	1.2	2.1	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	3.4	*	*	*	0.4	1.4	4.9	7.4	6.8	5.6	4.3	3.6
2001	3.3	*	*	*	0.4	1.6	4.5	7.2	6.3	5.4	4.6	4.1
2000	3.2	*	*	*	0.4	1.6	4.5	6.7	6.2	5.4	4.6	4.1
1999	3.1	*	*	*	0.4	1.5	4.7	6.4	6.2	5.3	4.9	3.9
Non-Hispanic 4/, male												
2002	5.2	*	*	*	0.6	2.0	7.0	11.7	11.2	9.0	7.1	6.8
2001	5.0	*	*	*	0.7	2.4	6.5	11.3	10.3	9.1	7.1	7.7
2000	4.9	*	*	*	0.7	2.3	6.7	10.7	10.0	8.8	7.8	7.4
1999	4.8	*	*	*	0.6	2.1	7.1	10.2	10.1	8.6	8.7	7.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to psychoactive substance use (F10-F19)												
Non-Hispanic 4/, female												
2002	1.7	*	*	*	0.2	0.8	2.8	3.2	2.7	2.8	2.5	2.3
2001	1.6	*	*	*	0.1	0.8	2.6	3.2	2.7	2.4	2.9	2.5
2000	1.5	*	*	*	0.2	0.8	2.4	2.8	2.6	2.5	2.6	2.8
1999	1.5	*	*	*	0.2	0.8	2.3	2.7	2.6	2.7	2.4	2.4
Non-Hispanic white, both sexes												
2002	3.3	*	*	*	0.5	1.5	4.7	6.8	6.0	5.3	4.3	3.5
2001	3.1	*	*	*	0.5	1.6	4.1	6.3	5.7	5.1	4.6	4.0
2000	3.0	*	*	*	0.5	1.5	4.1	5.8	5.4	5.0	4.6	4.1
1999	2.9	*	*	*	0.4	1.5	4.1	5.4	5.4	4.9	4.8	3.9
Non-Hispanic white, male												
2002	5.0	*	*	*	0.7	2.1	6.8	10.7	9.6	8.1	6.8	6.7
2001	4.7	*	*	*	0.7	2.4	5.9	9.9	8.9	8.3	6.9	7.5
2000	4.6	*	*	*	0.7	2.2	6.2	9.4	8.7	8.0	7.6	7.3
1999	4.5	*	*	*	0.6	2.1	6.3	8.8	8.7	7.6	8.3	7.4
Non-Hispanic white, female												
2002	1.7	*	*	*	0.2	0.8	2.5	2.9	2.5	3.0	2.6	2.2
2001	1.6	*	*	*	0.2	0.8	2.3	2.7	2.6	2.4	3.0	2.6
2000	1.5	*	*	*	0.2	0.8	2.0	2.4	2.4	2.6	2.7	2.9
1999	1.4	*	*	*	0.2	0.8	1.9	2.1	2.4	2.7	2.5	2.5
Non-Hispanic black, both sexes												
2002	4.6	*	*	*	*	1.6	6.8	13.1	14.1	9.1	5.5	*
2001	4.7	*	*	*	0.4	1.7	7.2	14.9	13.0	8.8	5.8	*
2000	4.6	*	*	*	0.5	2.2	6.9	14.0	12.9	9.4	5.9	*
1999	4.8	*	*	*	0.4	1.7	8.3	13.9	13.8	10.3	7.4	*
Non-Hispanic black, male												
2002	7.2	*	*	*	*	2.3	9.2	21.7	26.3	19.0	11.7	*
2001	7.6	*	*	*	0.7	2.6	10.4	24.5	24.6	18.2	12.0	*
2000	7.4	*	*	*	0.8	3.2	10.0	23.4	23.5	18.7	13.3	*
1999	7.6	*	*	*	*	2.3	12.2	22.1	25.5	20.6	15.6	*
Non-Hispanic black, female												
2002	2.2	*	*	*	*	1.1	4.7	5.8	4.2	*	*	*
2001	2.1	*	*	*	*	0.9	4.3	6.7	3.7	2.2	*	*
2000	2.2	*	*	*	*	1.3	4.2	5.9	4.5	2.8	*	*
1999	2.4	*	*	*	*	1.1	4.9	6.9	4.5	3.1	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to use of alcohol (F10)												
All origins 3/, both sexes												
2002	2.4	*	*	*	0.2	0.6	3.1	5.6	5.8	4.6	3.0	2.0
2001	2.3	*	*	*	0.2	0.8	2.9	5.5	5.7	4.6	3.2	2.1
2000	2.2	*	*	*	0.1	0.7	2.9	5.1	5.4	4.6	3.2	2.5
1999	2.2	*	*	*	0.2	0.7	2.9	5.0	5.5	4.4	3.5	2.3
All origins 3/, male												
2002	3.8	*	*	*	0.2	1.0	4.7	9.1	10.0	8.0	5.7	4.4
2001	3.7	*	*	*	0.3	1.1	4.5	8.9	9.5	8.1	5.6	4.7
2000	3.6	*	*	*	0.2	1.1	4.6	8.4	9.2	8.1	6.0	5.2
1999	3.6	*	*	*	0.3	1.1	4.6	8.3	9.2	7.6	6.8	4.9
All origins 3/, female												
2002	1.0	*	*	*	*	0.3	1.5	2.2	2.0	1.8	1.3	1.0
2001	1.0	*	*	*	*	0.4	1.4	2.2	2.2	1.7	1.6	1.1
2000	0.9	*	*	*	*	0.3	1.2	1.9	1.9	1.7	1.3	1.5
1999	0.9	*	*	*	*	0.4	1.3	1.9	2.1	1.8	1.3	1.3
Hispanic, both sexes												
2002	1.5	*	*	*	0.3	0.8	2.6	4.4	5.7	5.3	3.5	*
2001	1.5	*	*	*	0.4	0.9	2.3	4.7	6.1	4.3	*	*
2000	1.5	*	*	*	0.3	0.8	2.8	4.6	5.7	5.1	*	*
1999	1.5	*	*	*	0.3	1.0	2.6	4.8	5.8	4.9	*	*
Hispanic, male												
2002	2.6	*	*	*	*	1.4	4.2	8.0	10.7	10.7	*	*
2001	2.5	*	*	*	0.6	1.3	3.9	8.2	11.4	9.0	*	*
2000	2.7	*	*	*	0.6	1.4	4.9	8.2	11.2	10.3	*	*
1999	2.7	*	*	*	*	1.7	4.7	8.5	10.8	10.1	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	0.8	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	1.2	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	2.5	*	*	*	0.1	0.6	3.1	5.6	5.8	4.5	3.0	1.9
2001	2.4	*	*	*	0.1	0.7	3.0	5.5	5.6	4.6	3.2	2.2
2000	2.3	*	*	*	0.1	0.7	2.9	5.1	5.3	4.5	3.1	2.5
1999	2.3	*	*	*	0.1	0.7	2.9	5.0	5.4	4.3	3.4	2.3
Non-Hispanic 4/, male												
2002	4.0	*	*	*	0.2	0.9	4.7	9.1	9.8	7.7	5.6	4.1
2001	3.8	*	*	*	0.3	1.1	4.5	8.8	9.2	8.0	5.5	4.9
2000	3.7	*	*	*	0.2	1.1	4.5	8.3	9.0	7.8	5.8	5.3
1999	3.6	*	*	*	0.2	0.9	4.5	8.1	9.0	7.3	6.7	4.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to use of alcohol (F10)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	0.3	1.6	2.3	2.0	1.8	1.3	1.0
2001	1.1	*	*	*	*	0.4	1.5	2.3	2.2	1.7	1.6	1.1
2000	1.0	*	*	*	*	0.3	1.3	2.0	2.0	1.7	1.4	1.4
1999	1.0	*	*	*	*	0.4	1.4	1.9	2.1	1.8	1.3	1.3
Non-Hispanic white, both sexes												
2002	2.5	*	*	*	0.1	0.6	3.2	5.6	5.2	4.1	2.9	1.9
2001	2.4	*	*	*	0.1	0.7	2.9	5.1	5.0	4.2	3.1	2.1
2000	2.2	*	*	*	0.1	0.7	2.8	4.7	4.7	4.1	3.0	2.5
1999	2.2	*	*	*	0.1	0.7	2.7	4.5	4.8	3.9	3.2	2.3
Non-Hispanic white, male												
2002	3.9	*	*	*	0.2	0.9	4.8	8.9	8.6	6.8	5.3	3.8
2001	3.7	*	*	*	0.2	1.0	4.2	8.1	8.1	7.2	5.3	4.6
2000	3.6	*	*	*	*	1.0	4.4	7.6	7.8	6.9	5.4	5.2
1999	3.4	*	*	*	0.2	1.0	4.2	7.4	7.7	6.3	6.3	4.7
Non-Hispanic white, female												
2002	1.1	*	*	*	*	0.3	1.5	2.3	1.9	1.9	1.3	1.0
2001	1.1	*	*	*	*	0.3	1.5	2.1	2.1	1.7	1.7	1.1
2000	1.0	*	*	*	*	0.4	1.2	1.8	1.9	1.7	1.5	1.4
1999	0.9	*	*	*	*	0.4	1.2	1.7	2.0	1.8	1.2	1.4
Non-Hispanic black, both sexes												
2002	2.8	*	*	*	*	0.6	3.2	7.3	11.1	8.1	4.8	*
2001	3.1	*	*	*	*	0.6	3.7	9.0	11.1	8.1	4.7	*
2000	3.0	*	*	*	*	0.9	3.2	8.8	10.9	8.6	5.0	*
1999	3.1	*	*	*	*	0.6	4.1	8.5	11.5	9.1	6.6	*
Non-Hispanic black, male												
2002	4.8	*	*	*	*	1.1	4.8	12.5	21.1	17.3	10.0	*
2001	5.3	*	*	*	*	1.0	6.1	15.3	21.2	16.9	10.2	*
2000	5.0	*	*	*	*	1.5	4.8	15.0	20.5	17.7	12.3	*
1999	5.0	*	*	*	*	*	6.2	14.0	21.6	18.6	13.4	*
Non-Hispanic black, female												
2002	1.1	*	*	*	*	*	1.8	2.8	3.2	*	*	*
2001	1.1	*	*	*	*	*	1.6	3.6	2.9	*	*	*
2000	1.1	*	*	*	*	*	1.8	3.4	3.2	2.2	*	*
1999	1.3	*	*	*	*	*	2.3	3.7	3.5	2.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)												
All origins 3/, both sexes												
2002	0.9	*	*	*	0.3	0.8	1.8	1.8	1.0	1.1	1.3	1.7
2001	0.9	*	*	*	0.3	0.8	1.6	1.8	0.8	0.8	1.4	1.8
2000	0.9	*	*	*	0.3	0.9	1.6	1.6	0.8	0.9	1.5	1.6
1999	0.9	*	*	*	0.3	0.9	1.8	1.6	0.9	1.0	1.4	1.5
All origins 3/, male												
2002	1.3	*	*	*	0.4	1.1	2.4	2.8	1.4	1.3	1.5	2.6
2001	1.2	*	*	*	0.4	1.2	2.1	2.7	1.2	1.1	1.6	2.8
2000	1.2	*	*	*	0.5	1.2	2.3	2.5	1.1	1.0	1.9	2.1
1999	1.3	*	*	*	0.4	1.3	2.7	2.3	1.3	1.3	2.0	2.5
All origins 3/, female												
2002	0.6	*	*	*	0.2	0.5	1.2	0.9	0.6	1.0	1.2	1.2
2001	0.5	*	*	*	0.1	0.4	1.0	0.9	0.5	0.6	1.3	1.4
2000	0.6	*	*	*	0.2	0.6	1.0	0.8	0.6	0.8	1.2	1.3
1999	0.5	*	*	*	0.1	0.4	0.9	0.8	0.4	0.8	1.1	1.1
Hispanic, both sexes												
2002	0.8	*	*	*	*	0.7	1.9	2.4	1.1	*	*	*
2001	0.7	*	*	*	*	0.5	1.8	2.0	1.1	*	*	*
2000	0.7	*	*	*	0.3	1.0	1.9	1.7	*	*	*	*
1999	0.9	*	*	*	0.4	1.1	2.0	2.6	1.2	*	*	*
Hispanic, male												
2002	1.2	*	*	*	*	1.1	3.0	4.0	*	*	*	*
2001	1.0	*	*	*	*	0.7	2.6	3.4	*	*	*	*
2000	1.1	*	*	*	*	1.3	2.9	2.6	*	*	*	*
1999	1.5	*	*	*	0.6	1.8	3.1	4.5	2.6	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	0.8	*	*	*	*	*
2001	0.3	*	*	*	*	*	0.9	*	*	*	*	*
2000	0.4	*	*	*	*	0.7	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	0.8	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	*	*	*	0.3	0.8	1.7	1.7	1.0	1.1	1.3	1.7
2001	0.9	*	*	*	0.3	0.9	1.5	1.7	0.8	0.8	1.4	1.8
2000	0.9	*	*	*	0.3	0.9	1.6	1.6	0.8	0.9	1.5	1.6
1999	0.9	*	*	*	0.3	0.8	1.7	1.4	0.8	1.0	1.5	1.5
Non-Hispanic 4/, male												
2002	1.3	*	*	*	0.4	1.1	2.2	2.6	1.4	1.3	1.5	2.7
2001	1.2	*	*	*	0.5	1.4	2.0	2.6	1.1	1.1	1.6	2.8
2000	1.2	*	*	*	0.5	1.2	2.2	2.4	1.1	1.0	2.0	2.1
1999	1.2	*	*	*	0.4	1.2	2.5	2.1	1.2	1.3	2.0	2.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental and behavioral disorders due to other psychoactive substance use (F11-F19)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	0.2	0.5	1.2	0.9	0.6	1.0	1.2	1.3
2001	0.6	*	*	*	*	0.5	1.0	0.9	0.5	0.7	1.3	1.4
2000	0.6	*	*	*	0.2	0.5	1.0	0.8	0.6	0.8	1.2	1.4
1999	0.5	*	*	*	0.1	0.5	0.9	0.8	0.5	0.8	1.1	1.1
Non-Hispanic white, both sexes												
2002	0.9	*	*	*	0.3	0.9	1.5	1.2	0.8	1.2	1.4	1.7
2001	0.8	*	*	*	0.3	0.9	1.2	1.2	0.7	0.9	1.5	1.9
2000	0.8	*	*	*	0.4	0.8	1.3	1.1	0.7	1.0	1.6	1.7
1999	0.8	*	*	*	0.3	0.8	1.4	0.9	0.6	1.1	1.6	1.6
Non-Hispanic white, male												
2002	1.1	*	*	*	0.5	1.2	2.0	1.8	1.0	1.3	1.5	2.8
2001	1.1	*	*	*	0.5	1.4	1.7	1.8	0.9	1.1	1.6	2.8
2000	1.1	*	*	*	0.5	1.2	1.8	1.7	0.9	1.0	2.1	2.1
1999	1.1	*	*	*	0.4	1.2	2.1	1.4	0.9	1.3	2.0	2.7
Non-Hispanic white, female												
2002	0.6	*	*	*	0.2	0.5	1.0	0.6	0.6	1.1	1.3	1.2
2001	0.5	*	*	*	*	0.4	0.8	0.6	0.5	0.7	1.4	1.5
2000	0.5	*	*	*	0.2	0.5	0.8	0.5	0.5	0.9	1.2	1.5
1999	0.5	*	*	*	*	0.4	0.6	0.5	0.4	0.9	1.3	1.2
Non-Hispanic black, both sexes												
2002	1.7	*	*	*	*	1.0	3.6	5.8	2.9	*	*	*
2001	1.7	*	*	*	*	1.1	3.5	5.9	1.9	*	*	*
2000	1.7	*	*	*	*	1.3	3.7	5.2	2.1	*	*	*
1999	1.8	*	*	*	*	1.1	4.2	5.5	2.3	*	*	*
Non-Hispanic black, male												
2002	2.5	*	*	*	*	1.1	4.4	9.2	5.3	*	*	*
2001	2.4	*	*	*	*	1.6	4.4	9.1	3.4	*	*	*
2000	2.4	*	*	*	*	1.7	5.2	8.3	3.0	*	*	*
1999	2.5	*	*	*	*	1.6	6.0	8.1	3.9	*	*	*
Non-Hispanic black, female												
2002	1.1	*	*	*	*	0.9	2.9	2.9	*	*	*	*
2001	1.0	*	*	*	*	*	2.7	3.1	*	*	*	*
2000	1.0	*	*	*	*	1.0	2.5	2.6	*	*	*	*
1999	1.1	*	*	*	*	0.8	2.7	3.2	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Schizophrenia, schizotypal and delusional disorders (F20-F29)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.1	3.3
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.7
2000	0.2	*	*	*	*	*	0.0	0.1	0.2	0.5	1.2	3.3
1999	0.2	*	*	*	*	*	0.0	0.1	0.2	0.4	1.4	4.9
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.5	0.9	1.9
2001	0.1	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	1.5
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.4
1999	0.2	*	*	*	*	*	*	0.1	*	0.4	1.2	4.6
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.3	3.9
2001	0.2	*	*	*	*	*	*	*	0.2	0.3	1.3	3.1
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.3	3.8
1999	0.2	*	*	*	*	*	*	*	*	0.4	1.6	5.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	3.4
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.5	5.0
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.2	0.5	0.9	2.0
2001	0.1	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	*
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	2.5
1999	0.2	*	*	*	*	*	*	0.1	*	0.4	1.2	4.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Schizophrenia, schizotypal and delusional disorders (F20-F29)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.4	4.0
2001	0.2	*	*	*	*	*	*	*	0.2	0.3	1.3	3.2
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.4	3.8
1999	0.3	*	*	*	*	*	*	*	*	0.4	1.6	5.1
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	3.4
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.2	2.8
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	3.5
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.1
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.2	0.2	0.5	0.8	2.0
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.0	*
2000	0.2	*	*	*	*	*	*	*	0.2	0.6	1.1	2.5
1999	0.2	*	*	*	*	*	*	*	*	0.5	1.1	4.8
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	*	*	0.5	1.4	4.1
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.4	3.4
2000	0.3	*	*	*	*	*	*	*	*	0.5	1.4	3.9
1999	0.3	*	*	*	*	*	*	*	*	0.4	1.7	5.2
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mood (affective) disorders (F30-F39)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	*	0.1	0.1	0.4	2.2	9.1
2001	0.3	*	*	*	*	*	0.1	0.1	0.1	0.5	2.2	9.6
2000	0.3	*	*	*	*	*	*	0.1	0.1	0.4	1.9	9.8
1999	0.3	*	*	*	*	*	*	0.1	0.1	0.4	2.4	9.4
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.6	6.6
2001	0.2	*	*	*	*	*	*	0.1	*	0.5	1.7	8.0
2000	0.2	*	*	*	*	*	*	*	*	0.3	1.4	9.1
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	2.0	8.8
All origins 3/, female												
2002	0.4	*	*	*	*	*	*	*	0.2	0.5	2.7	10.1
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.5	10.3
2000	0.4	*	*	*	*	*	*	*	*	0.4	2.2	10.1
1999	0.4	*	*	*	*	*	*	*	*	0.5	2.6	9.7
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	*	0.1	0.1	0.4	2.3	9.3
2001	0.3	*	*	*	*	*	0.1	0.1	0.1	0.5	2.2	9.8
2000	0.3	*	*	*	*	*	*	0.1	0.1	0.4	1.9	10.1
1999	0.3	*	*	*	*	*	*	*	0.1	0.4	2.4	9.6
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	*	0.3	1.6	6.8
2001	0.2	*	*	*	*	*	*	0.1	*	0.5	1.8	8.3
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.5	9.4
1999	0.2	*	*	*	*	*	*	*	*	0.4	2.1	9.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mood (affective) disorders (F30-F39)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	*	0.2	0.5	2.8	10.4
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.5	10.4
2000	0.4	*	*	*	*	*	*	*	*	0.4	2.3	10.3
1999	0.4	*	*	*	*	*	*	*	*	0.5	2.6	9.8
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.4	10.0
2001	0.4	*	*	*	*	*	0.1	*	0.1	0.5	2.4	10.3
2000	0.4	*	*	*	*	*	*	0.1	0.2	0.4	2.1	10.9
1999	0.4	*	*	*	*	*	*	*	0.1	0.5	2.6	10.4
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.7	7.4
2001	0.3	*	*	*	*	*	*	*	*	0.6	1.9	8.7
2000	0.3	*	*	*	*	*	*	*	*	0.4	1.5	10.5
1999	0.3	*	*	*	*	*	*	*	*	0.4	2.2	9.8
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	*	0.6	2.9	11.2
2001	0.5	*	*	*	*	*	*	*	*	0.5	2.7	11.0
2000	0.5	*	*	*	*	*	*	*	*	0.5	2.4	11.0
1999	0.5	*	*	*	*	*	*	*	*	0.5	2.9	10.6
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neurotic, stress-related and somatoform disorders (F40-F48)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.6
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	0.8
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.7
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.7
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	0.8
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neurotic, stress-related and somatoform disorders (F40-F48)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	0.9
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.7
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.7
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	0.8
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	0.9
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	0.9
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental retardation (F70-F79)												
All origins 3/, both sexes												
2002	0.2	*	*	*	0.1	0.1	0.2	0.2	0.4	0.4	0.4	0.5
2001	0.1	*	*	*	0.1	0.1	0.1	0.2	0.3	0.4	0.4	*
2000	0.1	*	*	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.5
1999	0.2	*	*	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.4	*
All origins 3/, male												
2002	0.2	*	*	*	*	*	0.2	0.3	0.4	0.4	0.5	*
2001	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.4	0.6	*
2000	0.2	*	*	*	0.1	0.1	0.2	0.2	0.3	0.3	0.5	*
1999	0.2	*	*	*	*	0.1	0.2	0.3	0.3	0.5	0.6	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	0.1	0.1	0.3	0.3	0.3	*
2001	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.3	*
2000	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2	0.3	*
1999	0.1	*	*	*	*	0.1	0.2	0.1	0.2	0.2	0.3	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	0.1	0.1	0.2	0.2	0.4	0.4	0.4	0.5
2001	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.4	0.4	*
2000	0.2	*	*	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.5
1999	0.2	*	*	0.1	0.1	0.1	0.2	0.2	0.3	0.3	0.4	*
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	0.2	0.3	0.5	0.4	0.5	*
2001	0.2	*	*	*	*	0.1	0.2	0.2	0.4	0.4	0.7	*
2000	0.2	*	*	*	0.1	*	0.2	0.2	0.3	0.3	0.5	*
1999	0.2	*	*	*	*	0.1	0.2	0.3	0.3	0.5	0.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Mental retardation (F70-F79)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	0.1	0.1	0.3	0.3	0.4	*
2001	0.1	*	*	*	*	*	0.1	0.2	0.3	0.3	0.3	*
2000	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2	0.3	*
1999	0.1	*	*	*	*	*	0.2	0.1	0.3	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.2	0.2	0.4	0.4	0.4	0.6
2001	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.4	0.4	*
2000	0.2	*	*	*	0.1	0.1	0.2	0.2	0.3	0.3	0.4	0.6
1999	0.2	*	*	*	0.1	0.1	0.2	0.2	0.3	0.3	0.4	*
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	0.2	0.3	0.5	0.5	0.5	*
2001	0.2	*	*	*	*	*	0.2	0.2	0.3	0.4	0.7	*
2000	0.2	*	*	*	*	*	0.2	0.2	0.4	0.3	0.5	*
1999	0.2	*	*	*	*	*	0.2	0.3	0.3	0.5	0.7	*
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	0.1	0.2	0.4	0.4	0.4	*
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.3	0.3	*
2000	0.1	*	*	*	*	*	0.1	0.2	0.3	0.2	0.3	*
1999	0.1	*	*	*	*	*	0.2	*	0.2	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.7	4.4
2001	0.1	*	*	*	*	0.1	*	*	*	0.2	0.7	4.1
2000	0.1	*	*	*	*	*	*	*	0.1	0.1	0.7	3.9
1999	0.1	*	*	*	*	*	0.1	0.1	*	0.2	0.7	4.0
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.9	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.5
All origins 3/, female												
2002	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.6	4.7
2001	0.2	*	*	*	*	*	*	*	*	*	0.7	4.4
2000	0.2	*	*	*	*	*	*	*	*	*	0.8	4.1
1999	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.7
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.8	4.4
2001	0.2	*	*	*	*	0.1	*	*	*	0.2	0.7	4.1
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.7	3.9
1999	0.2	*	*	*	*	*	0.1	0.1	*	0.3	0.6	4.0
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.9	3.5
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	3.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	3.2
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.5	2.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	0.2	0.1	0.1	*	0.2	0.7	4.7
2001	0.2	*	*	*	*	*	*	*	*	*	0.7	4.4
2000	0.2	*	*	*	*	*	*	*	*	*	0.8	4.2
1999	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.6
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	*	0.2	0.8	4.3
2001	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.4
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999	0.2	*	*	*	*	*	0.1	*	*	0.3	0.7	4.3
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.9	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	3.2
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.7
Non-Hispanic white, female												
2002	0.3	*	*	*	*	0.2	0.1	*	*	*	0.7	4.7
2001	0.2	*	*	*	*	*	*	*	*	*	0.8	4.7
2000	0.2	*	*	*	*	*	*	*	*	*	0.8	4.3
1999	0.3	*	*	*	*	*	*	*	*	*	0.8	5.0
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VI. Diseases of the nervous system (G00-G98)												
All origins 3/, both sexes												
2002	35.7	9.5	2.1	1.2	2.0	2.2	4.2	9.1	19.7	62.0	270.2	951.0
2001	33.9	8.9	2.1	1.2	2.1	2.2	4.1	8.7	19.4	60.2	258.9	906.2
2000	32.3	10.2	2.1	1.3	1.9	2.1	4.1	9.1	19.2	59.1	248.5	864.9
1999	30.1	11.6	2.4	1.3	2.0	2.1	3.9	8.5	17.9	57.1	234.2	790.8
All origins 3/, male												
2002	28.5	11.0	2.4	1.4	2.6	2.7	4.9	9.8	21.2	70.3	293.9	867.4
2001	27.4	10.0	2.2	1.3	2.8	2.4	4.7	9.2	20.3	68.7	289.2	861.6
2000	26.2	11.5	2.1	1.3	2.4	2.5	4.6	9.9	20.6	68.3	277.0	826.2
1999	25.0	12.2	2.6	1.4	2.8	2.4	4.4	9.0	19.7	65.6	268.8	772.7
All origins 3/, female												
2002	42.8	7.8	1.7	1.1	1.4	1.8	3.5	8.4	18.3	55.0	254.4	987.3
2001	40.2	7.8	1.9	1.1	1.4	1.9	3.5	8.2	18.4	53.2	239.0	924.9
2000	38.2	8.8	2.0	1.2	1.3	1.8	3.6	8.2	18.0	51.6	229.9	880.7
1999	35.1	11.0	2.1	1.1	1.3	1.7	3.4	8.1	16.2	50.2	211.8	798.2
Hispanic, both sexes												
2002	8.6	6.6	1.6	1.1	1.5	1.4	1.9	5.0	12.1	40.3	154.0	564.9
2001	7.8	5.6	1.7	0.9	1.4	1.1	2.2	4.4	10.6	35.4	154.2	527.1
2000	7.8	8.6	1.9	0.9	1.2	1.3	2.3	4.9	11.2	38.3	150.7	510.3
1999	7.3	8.1	2.1	0.9	1.4	1.2	2.3	4.1	9.8	36.0	142.7	480.6
Hispanic, male												
2002	7.4	7.7	1.8	1.3	1.9	1.4	2.3	5.7	13.5	47.2	162.5	507.0
2001	7.1	6.7	2.2	1.0	2.1	1.3	2.7	5.1	11.4	42.1	176.2	493.3
2000	7.2	9.6	1.9	1.0	1.6	1.5	2.5	6.3	13.7	47.5	179.0	483.7
1999	6.5	9.6	2.3	1.0	1.6	1.3	2.4	4.9	11.6	43.1	162.7	440.0
Hispanic, female												
2002	9.9	5.4	1.4	0.9	1.0	1.3	1.5	4.3	10.9	34.8	148.1	593.5
2001	8.7	*	*	0.9	0.7	0.9	1.6	3.7	9.8	30.1	139.3	543.6
2000	8.4	7.4	1.9	0.8	0.7	1.0	2.1	3.6	8.9	31.0	131.7	523.3
1999	8.2	6.5	2.0	0.8	1.2	1.0	2.1	3.3	8.3	30.5	129.4	500.2
Non-Hispanic 4/, both sexes												
2002	39.8	10.2	2.2	1.2	2.1	2.4	4.5	9.4	20.2	63.3	275.2	964.4
2001	37.7	9.7	2.1	1.2	2.2	2.4	4.4	9.1	20.0	61.7	263.2	918.3
2000	35.8	10.6	2.1	1.3	2.0	2.3	4.3	9.4	19.8	60.3	252.2	875.9
1999	33.2	12.3	2.4	1.3	2.2	2.2	4.1	8.9	18.4	58.3	237.5	800.0
Non-Hispanic 4/, male												
2002	31.9	11.9	2.6	1.4	2.7	3.0	5.2	10.1	21.7	71.7	299.8	881.9
2001	30.5	10.7	2.2	1.3	2.9	2.6	4.9	9.6	20.9	70.3	293.9	874.7
2000	29.1	12.0	2.2	1.4	2.5	2.6	4.8	10.2	21.1	69.3	280.7	838.8
1999	27.6	12.8	2.6	1.4	3.0	2.6	4.7	9.3	20.2	66.8	272.8	784.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VI. Diseases of the nervous system (G00-G98)												
Non-Hispanic 4/, female												
2002	47.4	8.4	1.8	1.1	1.5	1.8	3.8	8.7	18.9	56.3	258.9	1,000.0
2001	44.5	8.6	2.0	1.2	1.5	2.2	3.8	8.6	19.1	54.6	243.0	936.5
2000	42.1	9.1	2.0	1.2	1.5	1.9	3.8	8.6	18.6	52.8	233.6	891.0
1999	38.5	11.9	2.2	1.2	1.3	1.8	3.5	8.5	16.8	51.3	214.6	806.3
Non-Hispanic white, both sexes												
2002	45.8	10.1	1.9	1.1	1.9	2.3	4.4	9.2	20.5	65.6	287.5	1,004.6
2001	43.1	8.6	1.9	1.2	2.1	2.2	4.2	8.9	20.1	64.1	273.5	955.6
2000	40.8	9.5	1.7	1.2	2.0	2.2	4.1	9.2	20.0	62.1	262.3	912.9
1999	37.8	11.0	2.1	1.2	2.0	2.1	3.9	8.6	18.4	59.9	246.9	835.7
Non-Hispanic white, male												
2002	36.2	11.9	2.2	1.2	2.6	2.9	5.1	9.8	22.0	73.8	312.5	921.0
2001	34.6	9.3	2.1	1.3	2.8	2.3	4.7	9.3	21.1	72.6	305.4	911.0
2000	32.8	10.6	1.9	1.3	2.6	2.6	4.5	10.1	21.0	71.5	291.5	873.6
1999	31.2	12.2	2.5	1.2	2.9	2.5	4.4	8.9	20.2	68.7	284.2	819.5
Non-Hispanic white, female												
2002	55.0	8.1	1.5	1.0	1.3	1.6	3.6	8.6	19.1	58.6	270.7	1,040.8
2001	51.2	7.8	1.6	1.1	1.4	2.0	3.6	8.5	19.1	56.9	252.3	974.2
2000	48.5	8.4	1.5	1.1	1.3	1.7	3.6	8.2	19.0	54.2	243.1	928.7
1999	44.1	9.7	1.7	1.2	1.1	1.7	3.4	8.2	16.7	52.4	222.5	842.2
Non-Hispanic black, both sexes												
2002	20.2	12.1	3.4	1.7	3.0	4.0	6.8	14.0	23.2	58.5	199.3	660.9
2001	19.6	15.1	3.3	1.7	3.0	4.2	6.6	12.9	24.4	55.2	202.6	631.8
2000	18.5	14.7	3.5	1.9	2.6	3.6	6.8	13.8	23.2	56.8	185.8	567.3
1999	17.2	18.3	3.5	1.9	3.1	3.2	6.3	13.8	22.5	55.8	168.9	487.6
Non-Hispanic black, male												
2002	17.1	12.4	3.9	1.9	3.7	4.7	7.9	15.5	25.1	70.4	219.8	571.0
2001	16.7	17.6	3.0	1.5	3.8	4.9	7.7	14.0	25.7	64.1	223.0	599.4
2000	15.9	18.1	3.3	2.0	2.7	3.8	7.9	13.9	27.4	65.3	206.7	544.8
1999	15.1	16.6	3.4	2.4	4.1	3.7	7.3	15.3	24.2	61.8	181.7	486.1
Non-Hispanic black, female												
2002	22.9	11.7	2.8	1.5	2.3	3.3	5.8	12.7	21.7	50.1	187.7	695.2
2001	22.3	12.4	3.6	1.9	2.2	3.6	5.6	11.8	23.4	48.9	191.2	643.8
2000	20.9	11.3	3.6	1.8	2.5	3.5	5.8	13.8	19.9	50.9	174.0	575.6
1999	19.1	20.0	3.6	1.5	2.1	2.8	5.5	12.5	21.1	51.6	161.8	488.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
All origins 3/, both sexes												
2002	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5
All origins 3/, male												
2002	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7
All origins 3/, female												
2002	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	2.1	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.6	1.2
2001	0.3	2.2	0.2	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.7	0.9
2000	0.3	2.7	0.3	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999	0.3	3.2	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.4
Non-Hispanic 4/, male												
2002	0.3	2.1	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001	0.3	2.1	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000	0.3	2.7	*	*	*	*	0.2	0.3	0.5	0.8	1.2	*
1999	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.1	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Non-Hispanic 4/, female												
2002	0.2	2.0	*	*	*	*	*	0.2	0.3	0.5	0.5	1.2
2001	0.2	2.3	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.7
2000	0.3	2.7	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.1
1999	0.3	3.3	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4
Non-Hispanic white, both sexes												
2002	0.2	1.9	*	0.1	*	*	0.1	0.2	0.2	0.5	0.5	1.2
2001	0.2	1.6	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.7
2000	0.2	2.1	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.1
1999	0.3	2.2	0.3	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3
Non-Hispanic white, male												
2002	0.2	1.9	*	*	*	*	0.1	0.2	0.3	0.7	0.7	*
2001	0.3	*	*	*	*	*	*	0.3	0.4	0.5	0.8	*
2000	0.3	2.3	*	*	*	*	0.1	0.3	0.4	0.7	1.1	*
1999	0.3	2.8	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
Non-Hispanic white, female												
2002	0.2	1.8	*	*	*	*	*	0.2	0.2	0.4	0.4	1.3
2001	0.2	1.9	*	*	*	*	*	0.2	0.3	0.4	0.6	*
2000	0.2	1.8	*	*	*	*	*	0.2	0.2	0.3	0.8	1.0
1999	0.3	*	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.2
Non-Hispanic black, both sexes												
2002	0.5	3.1	*	*	*	*	*	0.9	0.9	1.3	*	*
2001	0.5	4.7	*	*	*	*	0.5	0.7	1.0	*	*	*
2000	0.6	5.9	*	*	*	*	0.5	0.6	0.9	1.7	*	*
1999	0.6	7.0	*	*	*	*	0.4	0.9	*	1.3	2.7	*
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	1.3	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	1.5	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	9.6	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other inflammatory diseases of central nervous system (G04-G09)												
All origins 3/, both sexes												
2002	0.3	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.6	1.2	1.2
2001	0.2	*	*	*	0.1	0.1	0.1	0.2	0.3	0.7	1.0	1.0
2000	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	1.1	1.0
1999	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.8	1.1
All origins 3/, male												
2002	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.9	1.6	*
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.2	1.8
2000	0.2	*	*	*	*	*	0.2	0.2	0.4	0.8	1.6	*
1999	0.2	*	*	0.1	*	*	0.2	0.2	0.4	0.6	1.1	*
All origins 3/, female												
2002	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.0	1.1
2001	0.2	*	*	*	*	0.2	0.1	0.2	0.3	0.6	0.9	0.7
2000	0.2	*	*	*	*	0.1	0.1	0.1	0.3	0.5	0.8	0.9
1999	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	0.6	0.9
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.6	1.3	1.2
2001	0.2	*	*	*	0.1	0.1	0.1	0.2	0.4	0.7	1.0	1.0
2000	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	1.1	1.0
1999	0.2	*	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.8	1.0
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	0.1	0.2	0.3	0.4	0.8	1.7	*
2001	0.2	*	*	*	*	*	0.1	0.3	0.4	0.7	1.2	1.8
2000	0.2	*	*	*	*	*	0.2	0.2	0.3	0.7	1.5	*
1999	0.2	*	*	*	*	*	0.2	0.2	0.4	0.6	1.2	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other inflammatory diseases of central nervous system (G04-G09)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.0	1.2
2001	0.2	*	*	*	*	0.2	0.1	0.2	0.3	0.6	0.9	0.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.9	0.9
1999	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	0.6	0.9
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	0.1	0.1	0.1	0.2	0.4	0.6	1.2	1.4
2001	0.2	*	*	*	*	0.1	0.1	0.2	0.4	0.6	1.0	1.0
2000	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.1	1.0
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	1.0
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.6	*
2001	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.1	1.9
2000	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	*
1999	0.2	*	*	*	*	*	*	0.1	0.4	0.5	1.2	*
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	1.3
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.5	0.9	*
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.4	0.8	0.9
1999	0.2	*	*	*	*	*	*	0.2	*	0.6	0.6	0.9
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	0.4	0.6	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	1.3	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All origins 3/, both sexes												
2002	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
All origins 3/, male												
2002	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
All origins 3/, female												
2002	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7
Hispanic, both sexes												
2002	1.3	*	*	*	*	*	*	*	1.1	8.6	37.2	76.5
2001	1.3	*	*	*	*	*	*	*	*	8.6	41.5	86.3
2000	1.2	*	*	*	*	*	*	*	*	7.8	37.7	86.3
1999	1.1	*	*	*	*	*	*	*	*	8.1	38.3	77.2
Hispanic, male												
2002	1.4	*	*	*	*	*	*	*	*	13.6	50.1	113.8
2001	1.4	*	*	*	*	*	*	*	*	11.2	63.0	127.6
2000	1.3	*	*	*	*	*	*	*	*	9.7	55.1	143.1
1999	1.3	*	*	*	*	*	*	*	*	11.4	58.9	105.7
Hispanic, female												
2002	1.1	*	*	*	*	*	*	*	*	4.7	28.4	58.0
2001	1.2	*	*	*	*	*	*	*	*	6.6	26.9	66.1
2000	1.1	*	*	*	*	*	*	*	*	6.3	26.1	58.4
1999	1.0	*	*	*	*	*	*	*	*	5.5	24.6	63.4
Non-Hispanic 4/, both sexes												
2002	6.6	*	*	*	*	*	*	0.1	1.2	12.4	65.1	137.4
2001	6.5	*	*	*	*	*	*	0.1	1.2	11.9	65.5	135.6
2000	6.2	*	*	*	*	*	*	0.1	1.1	11.7	62.8	133.3
1999	5.8	*	*	*	*	*	*	0.1	1.1	11.1	58.9	125.8
Non-Hispanic 4/, male												
2002	7.6	*	*	*	*	*	*	0.2	1.5	18.2	96.7	224.4
2001	7.5	*	*	*	*	*	*	0.2	1.8	17.1	98.3	227.8
2000	7.1	*	*	*	*	*	*	0.2	1.5	16.5	94.2	225.2
1999	6.7	*	*	*	*	*	*	0.2	1.5	15.9	90.6	209.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Non-Hispanic 4/, female												
2002	5.6	*	*	*	*	*	*	*	0.9	7.6	44.2	99.8
2001	5.4	*	*	*	*	*	*	*	0.7	7.6	44.0	97.3
2000	5.3	*	*	*	*	*	*	*	0.8	7.7	42.3	96.1
1999	4.9	*	*	*	*	*	*	*	0.7	7.1	38.3	92.3
Non-Hispanic white, both sexes												
2002	7.9	*	*	*	*	*	*	0.2	1.3	13.3	69.6	145.8
2001	7.7	*	*	*	*	*	*	0.1	1.3	12.8	69.9	143.5
2000	7.3	*	*	*	*	*	*	0.1	1.2	12.6	66.7	141.2
1999	6.8	*	*	*	*	*	*	0.1	1.1	11.9	62.4	132.8
Non-Hispanic white, male												
2002	9.1	*	*	*	*	*	*	0.2	1.6	19.3	102.9	237.8
2001	9.0	*	*	*	*	*	*	0.2	2.0	18.3	104.5	241.7
2000	8.4	*	*	*	*	*	*	0.2	1.6	17.6	99.2	238.6
1999	7.9	*	*	*	*	*	*	0.2	1.6	17.1	95.7	221.0
Non-Hispanic white, female												
2002	6.7	*	*	*	*	*	*	*	1.0	8.1	47.3	106.0
2001	6.5	*	*	*	*	*	*	*	0.7	8.2	46.9	102.7
2000	6.3	*	*	*	*	*	*	*	0.9	8.3	45.3	101.9
1999	5.8	*	*	*	*	*	*	*	0.7	7.6	40.5	97.7
Non-Hispanic black, both sexes												
2002	1.5	*	*	*	*	*	*	*	*	6.7	25.7	51.7
2001	1.5	*	*	*	*	*	*	*	0.8	5.6	25.7	59.0
2000	1.5	*	*	*	*	*	*	*	*	5.8	27.3	52.5
1999	1.4	*	*	*	*	*	*	*	0.9	5.9	24.8	51.7
Non-Hispanic black, male												
2002	1.8	*	*	*	*	*	*	*	*	11.1	40.4	87.5
2001	1.7	*	*	*	*	*	*	*	*	8.8	38.1	99.3
2000	1.8	*	*	*	*	*	*	*	*	9.0	47.5	88.1
1999	1.6	*	*	*	*	*	*	*	*	8.7	38.5	87.9
Non-Hispanic black, female												
2002	1.3	*	*	*	*	*	*	*	*	3.6	17.4	38.0
2001	1.4	*	*	*	*	*	*	*	*	3.3	18.7	44.1
2000	1.2	*	*	*	*	*	*	*	*	3.5	16.0	39.2
1999	1.3	*	*	*	*	*	*	*	*	3.9	17.2	38.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All origins 3/, both sexes												
2002	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
All origins 3/, male												
2002	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
All origins 3/, female												
2002	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0
Hispanic, both sexes												
2002	3.8	*	*	*	*	*	*	*	1.0	10.8	89.2	438.5
2001	3.4	*	*	*	*	*	*	*	*	9.9	83.9	399.5
2000	3.2	*	*	*	*	*	*	*	*	11.9	79.2	380.2
1999	2.9	*	*	*	*	*	*	*	*	9.7	75.4	351.5
Hispanic, male												
2002	2.3	*	*	*	*	*	*	*	*	10.5	80.4	341.3
2001	2.1	*	*	*	*	*	*	*	*	8.8	83.2	306.6
2000	2.1	*	*	*	*	*	*	*	*	13.3	81.6	284.2
1999	1.8	*	*	*	*	*	*	*	*	11.2	66.7	278.2
Hispanic, female												
2002	5.4	*	*	*	*	*	*	*	*	10.9	95.3	486.6
2001	4.7	*	*	*	*	*	*	*	*	10.7	84.3	444.9
2000	4.3	*	*	*	*	*	*	*	*	10.8	77.6	427.3
1999	4.0	*	*	*	*	*	*	*	*	8.4	81.2	386.8
Non-Hispanic 4/, both sexes												
2002	22.9	*	*	*	*	*	*	0.1	2.0	20.3	161.1	763.2
2001	21.2	*	*	*	*	*	*	0.2	2.2	19.2	150.2	720.5
2000	19.6	*	*	*	*	*	*	0.2	2.1	19.0	142.0	676.7
1999	17.7	*	*	*	*	*	*	0.2	2.0	17.8	131.5	608.9
Non-Hispanic 4/, male												
2002	13.6	*	*	*	*	*	*	0.2	1.9	19.1	145.0	581.4
2001	12.7	*	*	*	*	*	*	0.2	2.0	18.3	138.6	565.2
2000	11.7	*	*	*	*	*	*	0.2	1.8	19.0	129.6	531.0
1999	10.9	*	*	*	*	*	*	0.2	2.0	17.7	124.8	492.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Non-Hispanic 4/, female												
2002	31.9	*	*	*	*	*	*	0.1	2.1	21.2	171.8	841.6
2001	29.3	*	*	*	*	*	*	0.2	2.4	20.0	157.8	785.1
2000	27.2	*	*	*	*	*	*	0.2	2.3	19.1	150.0	735.6
1999	24.2	*	*	*	*	*	*	0.2	1.9	18.0	135.9	655.4
Non-Hispanic white, both sexes												
2002	26.9	*	*	*	*	*	*	0.2	2.1	21.3	168.2	793.9
2001	24.8	*	*	*	*	*	*	0.2	2.4	20.3	155.6	749.2
2000	22.9	*	*	*	*	*	*	0.2	2.2	19.9	147.4	705.1
1999	20.7	*	*	*	*	*	*	0.2	2.1	18.6	136.7	636.6
Non-Hispanic white, male												
2002	15.8	*	*	*	*	*	*	0.2	2.0	19.9	150.5	605.4
2001	14.7	*	*	*	*	*	*	0.2	2.1	19.1	143.0	587.2
2000	13.6	*	*	*	*	*	*	0.2	2.0	19.8	134.6	551.7
1999	12.7	*	*	*	*	*	*	0.2	2.2	18.3	129.8	515.1
Non-Hispanic white, female												
2002	37.6	*	*	*	*	*	*	0.2	2.3	22.5	180.1	875.3
2001	34.4	*	*	*	*	*	*	0.2	2.7	21.3	163.9	816.6
2000	31.9	*	*	*	*	*	*	0.2	2.5	20.0	155.8	767.0
1999	28.4	*	*	*	*	*	*	0.2	2.0	18.8	141.2	685.0
Non-Hispanic black, both sexes												
2002	9.1	*	*	*	*	*	*	*	1.2	16.8	124.2	548.7
2001	8.7	*	*	*	*	*	*	*	1.1	15.5	126.9	513.2
2000	7.7	*	*	*	*	*	*	*	1.3	16.1	111.6	450.9
1999	6.7	*	*	*	*	*	*	*	1.2	15.5	99.1	374.7
Non-Hispanic black, male												
2002	5.5	*	*	*	*	*	*	*	*	17.9	122.5	414.6
2001	5.3	*	*	*	*	*	*	*	*	16.3	123.6	420.4
2000	4.6	*	*	*	*	*	*	*	*	16.9	98.7	378.1
1999	4.2	*	*	*	*	*	*	*	*	16.8	90.7	320.9
Non-Hispanic black, female												
2002	12.5	*	*	*	*	*	*	*	*	16.1	125.2	599.9
2001	11.8	*	*	*	*	*	*	*	*	15.0	128.7	547.5
2000	10.6	*	*	*	*	*	*	*	*	15.6	118.8	478.0
1999	9.0	*	*	*	*	*	*	*	*	14.5	103.8	394.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple sclerosis (G35)												
All origins 3/, both sexes												
2002	1.1	*	*	*	*	0.1	0.6	2.0	3.2	3.4	3.4	2.2
2001	1.1	*	*	*	*	0.1	0.7	1.9	3.1	3.4	3.5	2.1
2000	1.0	*	*	*	*	0.1	0.7	1.9	3.1	3.1	3.1	1.7
1999	1.0	*	*	*	*	0.1	0.7	1.9	2.6	2.9	3.0	2.1
All origins 3/, male												
2002	0.8	*	*	*	*	*	0.4	1.3	2.6	2.5	2.6	1.9
2001	0.7	*	*	*	*	*	0.5	1.4	2.2	2.7	2.7	1.5
2000	0.7	*	*	*	*	*	0.5	1.4	2.3	2.4	2.3	1.8
1999	0.6	*	*	*	*	*	0.5	1.3	1.9	2.4	2.3	*
All origins 3/, female												
2002	1.4	*	*	*	*	0.1	0.8	2.6	3.8	4.2	3.9	2.4
2001	1.4	*	*	*	*	0.2	0.9	2.5	3.9	3.9	4.0	2.3
2000	1.3	*	*	*	*	0.2	0.8	2.4	3.8	3.6	3.6	1.7
1999	1.2	*	*	*	*	0.2	0.9	2.5	3.2	3.3	3.5	2.4
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	0.6	*	*	*	*
2001	0.2	*	*	*	*	*	*	0.6	*	*	*	*
2000	0.2	*	*	*	*	*	*	0.6	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	*	*	0.1	0.7	2.1	3.4	3.6	3.5	2.3
2001	1.2	*	*	*	*	0.1	0.7	2.1	3.3	3.5	3.6	2.1
2000	1.1	*	*	*	*	0.2	0.7	2.0	3.2	3.2	3.2	1.8
1999	1.1	*	*	*	*	0.1	0.7	2.0	2.7	3.0	3.1	2.1
Non-Hispanic 4/, male												
2002	0.9	*	*	*	*	*	0.5	1.4	2.7	2.6	2.7	2.0
2001	0.8	*	*	*	*	*	0.5	1.5	2.4	2.8	2.8	1.6
2000	0.8	*	*	*	*	*	0.6	1.5	2.3	2.4	2.4	1.9
1999	0.7	*	*	*	*	*	0.5	1.3	1.9	2.5	2.3	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple sclerosis (G35)												
Non-Hispanic 4/, female												
2002	1.6	*	*	*	*	*	0.9	2.8	4.0	4.4	4.0	2.4
2001	1.5	*	*	*	*	0.2	1.0	2.6	4.2	4.1	4.1	2.4
2000	1.4	*	*	*	*	0.2	0.8	2.5	4.0	3.8	3.7	1.8
1999	1.4	*	*	*	*	0.2	0.9	2.7	3.4	3.4	3.6	2.4
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	*	0.7	2.2	3.6	3.9	3.7	2.5
2001	1.3	*	*	*	*	0.1	0.7	2.2	3.5	3.8	3.9	2.3
2000	1.2	*	*	*	*	0.1	0.7	2.1	3.3	3.4	3.4	2.0
1999	1.2	*	*	*	*	0.1	0.7	2.1	2.9	3.2	3.4	2.3
Non-Hispanic white, male												
2002	1.0	*	*	*	*	*	0.5	1.5	3.0	2.8	2.9	2.2
2001	1.0	*	*	*	*	*	0.5	1.6	2.6	3.1	3.0	1.8
2000	0.9	*	*	*	*	*	0.5	1.6	2.4	2.7	2.6	2.1
1999	0.8	*	*	*	*	*	0.5	1.4	2.1	2.7	2.5	*
Non-Hispanic white, female												
2002	1.7	*	*	*	*	*	0.8	2.8	4.2	4.8	4.3	2.6
2001	1.7	*	*	*	*	0.2	0.9	2.7	4.4	4.4	4.4	2.5
2000	1.6	*	*	*	*	0.2	0.8	2.6	4.2	4.1	4.0	1.9
1999	1.5	*	*	*	*	*	0.9	2.8	3.7	3.6	4.0	2.6
Non-Hispanic black, both sexes												
2002	0.9	*	*	*	*	*	1.1	2.7	2.9	2.5	*	*
2001	0.8	*	*	*	*	*	1.1	2.2	2.8	2.2	*	*
2000	0.9	*	*	*	*	*	1.2	2.3	3.4	2.0	*	*
1999	0.7	*	*	*	*	*	1.0	2.0	1.9	2.4	*	*
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	1.6	1.8	*	*	*
2001	0.5	*	*	*	*	*	*	1.4	*	*	*	*
2000	0.6	*	*	*	*	*	1.0	1.3	2.9	*	*	*
1999	0.5	*	*	*	*	*	*	1.5	*	*	*	*
Non-Hispanic black, female												
2002	1.3	*	*	*	*	*	1.6	3.6	3.9	3.4	*	*
2001	1.2	*	*	*	*	*	1.6	2.9	3.8	2.7	*	*
2000	1.1	*	*	*	*	*	1.4	3.2	3.7	2.7	*	*
1999	0.9	*	*	*	*	*	1.4	2.3	2.2	3.2	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Epilepsy (G40-G41)												
All origins 3/, both sexes												
2002	0.5	*	0.1	0.1	0.3	0.4	0.6	0.6	0.5	0.8	1.3	1.8
2001	0.5	*	*	0.1	0.2	0.4	0.6	0.5	0.6	0.7	1.2	1.6
2000	0.5	*	0.2	0.1	0.2	0.4	0.5	0.6	0.6	0.8	1.2	2.0
1999	0.5	*	0.2	0.1	0.3	0.4	0.5	0.6	0.6	0.9	1.2	1.8
All origins 3/, male												
2002	0.5	*	*	0.1	0.3	0.5	0.7	0.8	0.6	0.8	1.3	2.3
2001	0.5	*	*	0.1	0.3	0.5	0.7	0.6	0.7	0.8	0.9	2.0
2000	0.5	*	*	0.1	0.2	0.5	0.6	0.7	0.8	0.9	1.0	2.4
1999	0.5	*	0.3	0.1	0.3	0.5	0.7	0.7	0.6	1.0	1.1	2.3
All origins 3/, female												
2002	0.4	*	*	*	0.3	0.3	0.4	0.4	0.5	0.8	1.3	1.5
2001	0.4	*	*	*	0.2	0.3	0.4	0.4	0.5	0.6	1.3	1.4
2000	0.4	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8	1.4	1.9
1999	0.4	*	*	0.1	0.2	0.3	0.4	0.5	0.5	0.8	1.4	1.7
Hispanic, both sexes												
2002	0.3	*	*	*	*	0.3	0.4	*	*	*	*	*
2001	0.3	*	*	*	*	*	0.4	*	*	*	*	*
2000	0.3	*	*	*	*	0.4	0.4	0.6	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	0.1	0.3	0.4	0.6	0.6	0.5	0.8	1.3	1.7
2001	0.5	*	*	0.1	0.2	0.4	0.6	0.5	0.6	0.7	1.2	1.6
2000	0.5	*	*	0.1	0.3	0.4	0.5	0.6	0.6	0.8	1.2	2.0
1999	0.5	*	0.2	0.1	0.3	0.5	0.6	0.6	0.6	0.9	1.2	1.7
Non-Hispanic 4/, male												
2002	0.6	*	*	0.1	0.3	0.5	0.7	0.8	0.6	0.8	1.3	2.3
2001	0.5	*	*	0.1	0.3	0.5	0.8	0.6	0.7	0.8	0.9	2.0
2000	0.5	*	*	*	0.3	0.5	0.6	0.7	0.8	0.9	0.9	2.4
1999	0.6	*	*	0.1	0.4	0.6	0.8	0.7	0.6	1.0	1.0	2.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Epilepsy (G40-G41)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	0.3	0.3	0.4	0.5	0.5	0.8	1.4	1.5
2001	0.4	*	*	*	0.2	0.4	0.4	0.4	0.5	0.7	1.4	1.5
2000	0.5	*	*	*	0.2	0.3	0.5	0.4	0.5	0.8	1.4	1.9
1999	0.4	*	*	0.1	0.2	0.3	0.4	0.5	0.5	0.8	1.3	1.6
Non-Hispanic white, both sexes												
2002	0.5	*	*	0.1	0.2	0.4	0.6	0.5	0.5	0.6	1.2	1.6
2001	0.4	*	*	0.1	0.2	0.4	0.5	0.5	0.5	0.6	0.9	1.6
2000	0.4	*	*	0.1	0.2	0.4	0.5	0.5	0.5	0.7	1.0	1.7
1999	0.4	*	0.2	0.1	0.2	0.4	0.5	0.5	0.4	0.7	1.1	1.6
Non-Hispanic white, male												
2002	0.5	*	*	*	0.2	0.5	0.7	0.6	0.5	0.7	1.0	2.2
2001	0.5	*	*	*	0.3	0.4	0.6	0.6	0.6	0.6	0.7	2.0
2000	0.5	*	*	*	0.2	0.4	0.6	0.7	0.6	0.7	0.6	*
1999	0.5	*	*	*	0.3	0.5	0.6	0.6	0.4	0.8	0.9	2.2
Non-Hispanic white, female												
2002	0.4	*	*	*	0.3	0.3	0.4	0.4	0.5	0.6	1.3	1.3
2001	0.4	*	*	*	0.2	0.4	0.4	0.4	0.4	0.6	1.1	1.4
2000	0.4	*	*	*	0.2	0.3	0.5	0.3	0.4	0.7	1.3	1.7
1999	0.4	*	*	*	0.2	0.3	0.4	0.4	0.4	0.6	1.2	1.4
Non-Hispanic black, both sexes												
2002	0.9	*	*	*	0.5	0.7	0.9	1.5	1.3	2.2	3.6	*
2001	0.8	*	*	*	0.4	0.8	1.2	0.9	1.7	1.6	3.9	*
2000	0.9	*	*	*	0.5	0.6	0.8	1.3	1.7	2.2	4.0	*
1999	1.0	*	*	*	0.4	0.8	1.1	1.4	1.7	2.7	3.5	*
Non-Hispanic black, male												
2002	1.0	*	*	*	*	1.0	1.2	2.1	*	*	*	*
2001	1.0	*	*	*	*	1.0	1.7	1.2	2.0	*	*	*
2000	0.9	*	*	*	*	*	1.0	1.5	2.3	*	*	*
1999	1.1	*	*	*	*	1.1	1.7	1.9	2.3	*	*	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	1.1	*	2.1	*	*
2001	0.7	*	*	*	*	*	0.7	*	*	*	4.0	*
2000	0.8	*	*	*	*	*	*	1.2	*	*	3.5	*
1999	0.8	*	*	*	*	*	*	1.0	*	2.7	3.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transient cerebral ischemic attacks and related syndromes (G45)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.5	2.9
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.5	3.8
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.6
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	1.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	4.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	2.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.8
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.4
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	3.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	4.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	4.9
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.0
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.5	3.9
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.7
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	4.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transient cerebral ischemic attacks and related syndromes (G45)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.4
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	3.7
2000	0.2	*	*	*	*	*	*	*	*	*	0.7	4.2
1999	0.2	*	*	*	*	*	*	*	*	*	0.5	5.0
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.6	3.1
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	4.0
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	4.0
1999	0.2	*	*	*	*	*	*	*	*	0.2	0.7	5.0
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	4.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.9	3.1
1999	0.1	*	*	*	*	*	*	*	*	*	1.0	4.1
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	3.8
2000	0.2	*	*	*	*	*	*	*	*	*	0.7	4.4
1999	0.2	*	*	*	*	*	*	*	*	*	0.5	5.4
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infantile cerebral palsy (G80)												
All origins 3/, both sexes												
2002	0.4	*	0.6	0.5	0.7	0.4	0.2	0.3	0.4	0.4	0.5	*
2001	0.4	*	0.5	0.5	0.7	0.3	0.2	0.3	0.4	0.4	0.5	*
2000	0.4	*	0.4	0.6	0.6	0.3	0.2	0.3	0.3	0.5	0.5	*
1999	0.4	*	0.5	0.5	0.5	0.3	0.2	0.2	0.3	0.4	0.4	0.5
All origins 3/, male												
2002	0.5	*	0.7	0.5	0.8	0.4	0.3	0.3	0.5	0.5	0.6	*
2001	0.5	*	0.5	0.5	0.8	0.4	0.3	0.3	0.4	0.6	0.7	*
2000	0.4	*	0.3	0.6	0.7	0.4	0.2	0.3	0.3	0.7	0.5	*
1999	0.4	*	0.6	0.5	0.6	0.4	0.2	0.3	0.4	0.5	0.5	*
All origins 3/, female												
2002	0.4	*	0.5	0.5	0.5	0.3	0.2	0.3	0.3	0.4	0.4	*
2001	0.3	*	0.4	0.5	0.5	0.3	0.2	0.2	0.4	0.3	0.4	*
2000	0.3	*	0.4	0.5	0.5	0.3	0.2	0.2	0.3	0.3	0.5	*
1999	0.3	*	0.4	0.4	0.4	0.3	0.2	0.2	0.2	0.3	0.4	*
Hispanic, both sexes												
2002	0.3	*	*	0.5	0.4	*	*	*	*	*	*	*
2001	0.3	*	*	0.4	0.5	*	*	*	*	*	*	*
2000	0.2	*	*	*	0.4	*	*	*	*	*	*	*
1999	0.3	*	0.7	0.4	0.3	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	0.6	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	0.7	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	0.6	0.5	0.7	0.4	0.2	0.3	0.4	0.5	0.5	*
2001	0.4	*	0.5	0.5	0.7	0.4	0.2	0.3	0.4	0.5	0.5	*
2000	0.4	*	0.4	0.6	0.6	0.4	0.2	0.3	0.3	0.5	0.5	*
1999	0.4	*	0.5	0.5	0.5	0.4	0.2	0.2	0.3	0.4	0.4	0.5
Non-Hispanic 4/, male												
2002	0.5	*	0.8	0.5	0.9	0.4	0.3	0.3	0.5	0.5	0.7	*
2001	0.5	*	0.5	0.5	0.8	0.4	0.3	0.3	0.4	0.7	0.7	*
2000	0.5	*	*	0.7	0.7	0.4	0.2	0.3	0.3	0.7	0.5	*
1999	0.4	*	0.6	0.5	0.6	0.4	0.2	0.3	0.4	0.5	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infantile cerebral palsy (G80)												
Non-Hispanic 4/, female												
2002	0.4	*	0.4	0.5	0.5	0.3	0.2	0.3	0.3	0.4	0.4	*
2001	0.4	*	0.4	0.5	0.6	0.3	0.2	0.3	0.4	0.3	0.4	*
2000	0.4	*	0.4	0.6	0.5	0.3	0.2	0.2	0.3	0.3	0.5	*
1999	0.3	*	0.4	0.4	0.4	0.3	0.2	0.2	0.2	0.3	0.4	*
Non-Hispanic white, both sexes												
2002	0.4	*	0.5	0.4	0.6	0.3	0.2	0.3	0.4	0.5	0.5	*
2001	0.4	*	0.4	0.4	0.6	0.3	0.3	0.3	0.4	0.5	0.6	*
2000	0.4	*	*	0.5	0.6	0.4	0.2	0.3	0.4	0.5	0.5	*
1999	0.4	*	0.4	0.4	0.5	0.4	0.3	0.3	0.3	0.4	0.5	0.6
Non-Hispanic white, male												
2002	0.5	*	0.7	0.4	0.8	0.4	0.3	0.3	0.5	0.6	0.7	*
2001	0.5	*	0.5	0.4	0.7	0.4	0.3	0.3	0.4	0.7	0.8	*
2000	0.4	*	*	0.6	0.7	0.4	0.2	0.3	0.4	0.7	*	*
1999	0.4	*	0.5	0.4	0.6	0.4	0.2	0.3	0.4	0.5	0.6	*
Non-Hispanic white, female												
2002	0.3	*	*	0.4	0.4	0.3	0.2	0.3	0.4	0.4	0.4	*
2001	0.3	*	*	0.4	0.5	0.3	0.2	0.2	0.4	0.3	0.4	*
2000	0.3	*	*	0.5	0.5	0.3	0.2	0.2	0.4	0.3	0.5	*
1999	0.3	*	*	0.4	0.3	0.4	0.3	0.2	*	0.3	0.4	*
Non-Hispanic black, both sexes												
2002	0.7	*	1.0	0.9	1.2	0.8	0.4	*	*	*	*	*
2001	0.6	*	*	0.8	1.1	0.7	*	*	*	*	*	*
2000	0.6	*	1.1	1.0	0.9	0.5	*	*	*	*	*	*
1999	0.5	*	1.2	0.7	0.9	0.4	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.8	*	*	0.9	1.5	0.9	*	*	*	*	*	*
2001	0.7	*	*	0.8	1.2	0.9	*	*	*	*	*	*
2000	0.7	*	*	1.0	1.0	*	*	*	*	*	*	*
1999	0.7	*	*	0.9	1.2	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.6	*	*	0.8	1.0	0.7	*	*	*	*	*	*
2001	0.5	*	*	0.8	0.9	*	*	*	*	*	*	*
2000	0.6	*	*	1.0	0.8	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)												
All origins 3/, both sexes												
2002	6.9	6.9	1.1	0.5	0.9	1.2	2.5	5.4	11.7	24.1	40.5	53.8
2001	6.7	6.1	1.2	0.5	1.0	1.1	2.3	5.2	11.2	24.0	39.4	51.9
2000	6.7	7.1	1.2	0.5	0.9	1.1	2.4	5.5	11.4	23.3	39.4	55.4
1999	6.5	7.7	1.2	0.6	1.1	1.0	2.1	5.0	10.9	23.3	39.3	53.6
All origins 3/, male												
2002	7.3	8.4	1.3	0.6	1.3	1.5	3.0	6.4	13.4	28.2	49.3	66.1
2001	7.1	7.1	1.2	0.6	1.5	1.2	2.9	5.9	12.5	28.5	49.0	69.7
2000	7.2	8.1	1.3	0.5	1.3	1.4	2.8	6.5	13.2	27.9	49.1	72.0
1999	7.0	8.6	1.2	0.6	1.6	1.2	2.6	5.8	12.6	27.4	49.7	70.3
All origins 3/, female												
2002	6.5	5.4	0.9	0.3	0.5	0.8	1.9	4.4	10.1	20.6	34.6	48.5
2001	6.3	5.0	1.2	0.5	0.5	0.9	1.8	4.4	10.0	20.2	33.1	44.5
2000	6.3	6.0	1.1	0.5	0.5	0.8	1.9	4.6	9.7	19.5	33.0	48.7
1999	6.1	6.8	1.2	0.5	0.5	0.7	1.7	4.3	9.2	20.0	32.6	46.8
Hispanic, both sexes												
2002	2.4	5.2	0.7	0.4	0.7	0.6	1.0	3.1	7.3	17.4	23.5	44.9
2001	2.1	3.5	0.9	0.4	0.6	0.5	1.2	2.6	7.0	13.6	24.7	38.1
2000	2.4	6.9	0.9	0.4	0.6	0.6	1.3	2.8	6.7	15.2	30.4	39.1
1999	2.2	5.0	0.9	0.4	0.6	0.5	1.2	2.4	6.8	15.4	24.6	42.8
Hispanic, male												
2002	2.5	6.1	*	*	1.0	0.8	1.2	3.9	8.0	18.5	27.5	46.9
2001	2.4	*	*	*	1.0	0.6	1.4	3.3	8.1	18.0	25.9	55.2
2000	2.6	7.3	*	*	0.8	0.8	1.4	3.9	8.3	20.3	36.4	52.4
1999	2.4	5.7	*	*	0.9	0.6	1.3	3.0	8.6	17.8	32.8	49.6
Hispanic, female												
2002	2.3	*	*	*	*	*	0.8	2.2	6.6	16.6	20.8	43.9
2001	1.9	*	*	*	*	*	0.9	2.0	6.1	10.1	23.9	29.8
2000	2.1	6.4	*	*	*	*	1.1	1.7	5.2	11.3	26.4	32.6
1999	2.0	*	*	*	*	*	1.1	1.8	5.2	13.6	19.1	39.5
Non-Hispanic 4/, both sexes												
2002	7.6	7.4	1.2	0.5	1.0	1.3	2.7	5.6	12.0	24.5	41.2	53.9
2001	7.3	6.7	1.3	0.6	1.1	1.2	2.5	5.4	11.5	24.6	40.0	52.2
2000	7.3	7.1	1.3	0.5	1.0	1.2	2.5	5.7	11.7	23.7	39.7	55.9
1999	7.1	8.3	1.2	0.6	1.1	1.1	2.2	5.3	11.1	23.7	39.8	53.8
Non-Hispanic 4/, male												
2002	8.1	9.0	1.4	0.6	1.4	1.7	3.3	6.6	13.8	28.8	50.3	66.7
2001	7.8	7.7	1.3	0.6	1.6	1.4	3.1	6.1	12.8	29.1	49.9	69.9
2000	7.8	8.3	1.4	0.5	1.4	1.5	3.0	6.7	13.5	28.3	49.6	72.5
1999	7.6	9.2	1.3	0.6	1.7	1.3	2.8	6.1	12.9	27.8	50.3	70.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)												
Non-Hispanic 4/, female												
2002	7.0	5.8	1.0	0.3	0.5	0.8	2.0	4.5	10.4	20.8	35.2	48.4
2001	6.9	5.6	1.3	0.5	0.6	1.0	1.9	4.6	10.3	20.9	33.4	44.9
2000	6.8	5.9	1.2	0.4	0.6	0.8	2.0	4.8	10.0	20.0	33.2	49.1
1999	6.6	7.3	1.2	0.5	0.5	0.8	1.7	4.5	9.5	20.3	33.0	46.9
Non-Hispanic white, both sexes												
2002	8.2	7.4	1.1	0.5	1.0	1.3	2.7	5.4	12.0	24.8	41.9	54.9
2001	7.9	6.4	1.1	0.6	1.1	1.1	2.4	5.2	11.2	24.9	40.4	52.8
2000	7.9	6.8	1.2	0.5	1.1	1.2	2.4	5.6	11.6	23.8	40.5	56.7
1999	7.6	8.1	1.1	0.5	1.2	1.1	2.2	5.0	10.9	23.8	40.5	54.5
Non-Hispanic white, male												
2002	8.8	9.2	1.2	0.6	1.4	1.8	3.3	6.5	13.8	28.9	51.5	68.2
2001	8.4	7.2	1.2	0.6	1.7	1.2	3.0	5.9	12.6	29.5	50.7	70.4
2000	8.4	7.4	1.3	0.5	1.6	1.5	2.8	6.7	13.3	28.5	50.6	73.8
1999	8.2	9.0	1.2	0.5	1.9	1.4	2.7	5.9	12.8	28.1	51.6	72.2
Non-Hispanic white, female												
2002	7.6	5.5	0.8	0.3	0.5	0.8	2.0	4.4	10.3	21.2	35.5	49.1
2001	7.3	5.6	1.0	0.5	0.5	0.9	1.9	4.5	9.8	21.0	33.6	45.5
2000	7.3	6.1	1.1	0.4	0.5	0.8	1.9	4.5	10.1	19.8	33.8	49.8
1999	7.1	7.2	1.1	0.5	0.5	0.8	1.6	4.1	9.2	20.3	33.3	47.4
Non-Hispanic black, both sexes												
2002	6.1	8.2	1.7	0.5	1.1	1.7	3.6	7.9	15.4	27.5	40.0	50.8
2001	6.3	8.3	2.1	0.7	1.2	2.0	3.3	8.0	16.3	27.5	41.5	49.0
2000	6.0	7.6	1.7	0.5	1.0	1.7	3.7	8.7	14.6	27.4	37.4	50.9
1999	5.9	9.3	1.4	0.9	1.2	1.3	3.1	8.6	14.9	26.6	36.8	50.8
Non-Hispanic black, male												
2002	6.5	8.2	2.0	0.6	1.4	2.0	4.5	9.3	17.8	34.3	45.1	58.0
2001	6.5	10.5	1.9	*	1.6	2.5	4.1	9.4	17.4	31.9	51.9	62.4
2000	6.4	10.9	1.7	*	1.0	2.1	4.4	9.3	18.1	32.2	49.7	62.2
1999	5.9	10.7	*	1.0	1.6	1.5	3.6	9.4	15.9	28.6	43.5	65.4
Non-Hispanic black, female												
2002	5.8	8.2	*	*	0.7	1.4	2.7	6.7	13.6	22.6	37.1	48.1
2001	6.1	*	2.2	0.8	0.9	1.6	2.6	6.9	15.5	24.5	35.7	44.1
2000	5.7	*	*	*	1.0	1.4	3.1	8.2	11.8	24.0	30.6	46.7
1999	5.9	7.9	*	0.7	*	1.2	2.7	8.0	14.2	25.2	33.1	45.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VII. Diseases of the eye and adnexa (H00-H57)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	*	0.6
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	*	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	*	0.5
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VII. Diseases of the eye and adnexa (H00-H57)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VIII. Diseases of the ear and mastoid process (H60-H93)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
VIII. Diseases of the ear and mastoid process (H60-H93)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)												
All origins 3/, both sexes												
2002	320.2	16.5	1.5	0.9	3.2	10.3	38.4	115.6	297.1	792.2	2266.8	7,414.0
2001	325.5	15.4	1.9	0.9	3.3	10.5	37.7	114.6	303.5	815.0	2331.8	7,700.0
2000	334.6	17.4	1.5	0.9	3.4	9.8	37.5	117.0	321.3	853.6	2405.6	8,055.8
1999	342.0	17.6	1.5	0.9	3.5	9.9	38.3	117.3	329.2	891.0	2484.0	8,220.5
All origins 3/, male												
2002	304.7	17.3	1.5	0.9	4.2	13.1	51.5	163.7	409.5	1031.7	2740.6	7,659.3
2001	307.6	15.8	2.0	0.9	4.0	13.0	50.6	161.8	416.6	1062.6	2833.2	7,958.3
2000	317.2	18.3	1.7	1.0	4.0	12.1	50.3	166.7	443.8	1118.8	2936.2	8,470.3
1999	324.2	18.5	1.6	1.0	4.1	12.3	51.9	166.9	454.0	1172.7	3033.6	8,624.0
All origins 3/, female												
2002	335.2	15.7	1.4	0.8	2.3	7.5	25.5	69.2	193.1	592.8	1952.2	7,307.6
2001	342.8	15.0	1.8	0.9	2.5	7.9	25.0	69.1	198.9	610.0	2002.2	7,591.9
2000	351.3	16.5	1.4	0.8	2.9	7.5	24.9	69.2	208.3	635.2	2059.6	7,887.0
1999	359.1	16.6	1.4	0.8	2.8	7.4	24.9	69.5	214.3	660.4	2127.0	8,057.9
Hispanic, both sexes												
2002	94.8	13.9	1.2	0.7	2.7	6.3	20.7	74.6	227.2	624.9	1734.8	5,388.0
2001	96.9	12.6	1.5	0.8	2.5	6.5	20.2	76.3	217.8	649.9	1838.9	6,075.6
2000	97.1	14.5	1.3	0.8	2.6	6.6	22.0	76.0	237.6	664.0	1861.5	6,192.8
1999	99.7	15.5	1.4	0.7	2.6	6.6	21.7	81.6	244.1	694.1	1940.9	6,360.5
Hispanic, male												
2002	94.6	13.4	*	0.6	3.4	8.3	27.7	103.8	318.2	825.7	2074.3	5,503.2
2001	96.0	13.7	1.4	0.8	3.1	8.4	27.1	108.5	298.9	857.4	2232.6	6,193.7
2000	96.7	14.4	1.4	0.9	3.1	8.3	28.6	109.0	332.3	873.4	2283.4	6,330.5
1999	99.7	12.7	1.4	0.9	3.1	8.7	30.0	117.3	334.6	929.0	2362.9	6,621.1
Hispanic, female												
2002	95.1	14.5	1.4	0.8	1.8	3.9	13.2	45.4	145.5	465.2	1502.2	5,330.9
2001	97.8	11.5	1.6	0.8	1.8	4.3	12.7	44.6	145.7	486.3	1572.8	6,017.9
2000	97.6	14.6	*	*	2.0	4.6	14.8	43.8	153.7	499.4	1578.2	6,125.2
1999	99.8	18.4	1.4	*	2.0	4.0	12.8	46.9	164.1	510.0	1660.0	6,234.8
Non-Hispanic 4/, both sexes												
2002	354.0	17.1	1.5	0.9	3.4	11.2	40.9	119.2	301.4	800.5	2286.3	7,476.1
2001	358.5	16.0	2.0	0.9	3.4	11.3	39.9	117.7	308.7	823.2	2347.6	7,741.0
2000	367.4	18.0	1.6	0.9	3.6	10.4	39.3	120.1	326.0	862.1	2421.6	8,102.0
1999	374.3	18.1	1.5	0.9	3.6	10.5	40.2	119.8	333.7	899.1	2498.5	8,263.1
Non-Hispanic 4/, male												
2002	338.0	18.3	1.7	1.0	4.3	14.1	54.9	169.0	414.7	1041.1	2765.3	7,734.9
2001	339.7	16.2	2.1	0.9	4.2	14.0	53.7	166.1	423.2	1071.8	2852.4	8,012.2
2000	349.2	19.0	1.7	1.0	4.2	12.9	52.8	171.0	449.6	1128.8	2955.6	8,535.2
1999	355.6	20.0	1.7	1.0	4.3	13.0	54.4	170.4	459.9	1181.4	3051.5	8,678.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)												
Non-Hispanic 4/, female												
2002	369.3	15.9	1.4	0.8	2.4	8.3	27.2	71.3	196.2	599.7	1968.8	7,364.4
2001	376.3	15.9	1.8	0.9	2.7	8.6	26.5	71.0	202.5	616.6	2016.2	7,628.2
2000	384.6	16.9	1.5	0.8	3.0	8.0	26.0	71.2	211.6	642.0	2073.7	7,926.7
1999	392.0	16.0	1.4	0.8	3.0	8.0	26.3	71.2	217.3	667.3	2139.6	8,096.9
Non-Hispanic white, both sexes												
2002	383.7	14.0	1.4	0.8	2.8	8.8	35.5	104.1	274.0	764.3	2261.8	7,570.6
2001	387.6	12.6	1.6	0.8	2.7	9.2	33.8	102.4	280.5	783.7	2322.1	7,837.2
2000	397.3	15.1	1.4	0.8	2.9	8.4	33.4	105.1	297.2	824.2	2394.0	8,204.4
1999	404.3	14.9	1.3	0.8	3.0	8.5	33.9	104.9	304.3	860.2	2471.9	8,365.9
Non-Hispanic white, male												
2002	363.7	15.4	1.6	0.9	3.6	11.4	49.3	152.4	383.2	1004.9	2753.4	7,866.2
2001	364.6	13.4	1.7	0.8	3.4	11.6	47.2	149.3	392.0	1031.7	2836.9	8,137.5
2000	375.2	16.3	1.6	0.9	3.4	10.7	46.9	154.6	417.4	1092.7	2940.1	8,672.2
1999	381.1	16.9	1.5	0.8	3.5	10.8	47.5	153.6	426.4	1143.7	3039.5	8,796.3
Non-Hispanic white, female												
2002	402.9	12.5	1.1	0.7	1.9	6.3	21.7	56.5	170.7	559.2	1932.4	7,442.9
2001	409.7	11.7	1.5	0.8	2.1	6.8	20.4	56.2	175.2	573.4	1980.8	7,712.2
2000	418.5	14.0	1.2	0.6	2.3	6.0	20.0	56.4	183.9	598.2	2035.1	8,015.8
1999	426.5	12.9	1.1	0.7	2.5	6.2	20.2	57.0	189.3	623.1	2100.2	8,194.2
Non-Hispanic black, both sexes												
2002	287.3	29.4	2.1	1.5	6.2	25.3	79.9	245.2	583.0	1250.0	2868.8	7,166.9
2001	291.7	30.9	3.8	1.3	7.1	24.3	82.2	245.5	592.2	1301.4	2930.1	7,396.3
2000	296.3	29.6	2.6	1.5	7.3	23.3	80.7	249.2	614.2	1342.1	3009.6	7,606.4
1999	300.2	30.8	2.6	1.4	7.2	22.2	85.1	249.2	626.7	1389.5	3058.9	7,652.9
Non-Hispanic black, male												
2002	279.3	29.7	2.0	1.5	7.6	31.2	99.3	322.4	781.1	1584.6	3336.9	7,014.4
2001	283.2	30.6	4.0	1.2	8.5	29.6	101.8	321.5	778.6	1648.8	3477.5	7,442.7
2000	286.3	30.7	2.3	1.5	8.1	27.4	99.4	327.5	812.8	1667.2	3541.0	7,786.1
1999	292.8	32.6	2.8	1.6	8.8	27.2	107.1	332.5	835.8	1747.0	3559.7	7,974.8
Non-Hispanic black, female												
2002	294.5	29.0	2.2	1.4	4.8	19.9	62.8	179.2	424.0	1012.7	2603.9	7,225.2
2001	299.5	31.2	3.6	1.4	5.7	19.5	64.9	180.3	442.2	1056.3	2622.8	7,379.1
2000	305.4	28.5	2.9	1.6	6.5	19.6	64.1	182.0	454.3	1113.4	2710.4	7,539.6
1999	307.0	28.9	2.4	1.3	5.6	17.7	65.8	178.0	459.0	1137.4	2778.5	7,532.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
All origins 3/, male												
2002	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
All origins 3/, female												
2002	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	0.7	1.8	4.6	7.0	*
2001	0.4	*	*	*	*	*	*	*	1.5	4.2	5.4	*
2000	0.4	*	*	*	*	*	*	*	1.8	4.4	5.3	*
1999	0.5	*	*	*	*	*	*	0.8	1.6	4.2	7.3	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.8	*	*	*	*	*	*	*	2.1	5.9	10.7	*
2001	0.6	*	*	*	*	*	*	*	2.0	5.2	*	*
2000	0.6	*	*	*	*	*	*	*	*	5.6	6.9	*
1999	0.8	*	*	*	*	*	*	*	*	5.9	10.4	*
Non-Hispanic 4/, both sexes												
2002	1.4	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.7	18.1
2001	1.3	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.6	18.0
2000	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.7	17.1
1999	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.1	18.3
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	7.0	12.0
2001	0.8	*	*	*	*	*	0.3	0.5	1.1	2.8	6.7	13.1
2000	0.9	*	*	*	*	*	0.3	0.6	1.4	2.9	6.7	12.5
1999	0.8	*	*	*	*	*	0.2	0.5	1.2	3.6	6.9	10.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic 4/, female												
2002	1.8	*	*	*	*	*	0.2	0.7	2.0	4.5	11.6	20.8
2001	1.8	*	*	*	*	*	0.3	0.7	1.7	5.1	11.5	20.0
2000	1.9	*	*	*	*	*	0.4	0.8	2.0	5.8	11.6	19.0
1999	2.0	*	*	*	*	0.1	0.3	0.8	2.3	5.5	12.2	21.3
Non-Hispanic white, both sexes												
2002	1.5	*	*	*	*	*	0.2	0.5	1.5	3.9	10.2	19.3
2001	1.5	*	*	*	*	*	0.2	0.5	1.3	4.0	10.2	19.0
2000	1.6	*	*	*	*	*	0.3	0.6	1.7	4.6	10.3	18.0
1999	1.6	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.6	19.2
Non-Hispanic white, male												
2002	0.9	*	*	*	*	*	0.2	0.4	1.1	3.0	7.3	12.9
2001	0.9	*	*	*	*	*	0.2	0.4	1.1	2.7	7.0	13.7
2000	0.9	*	*	*	*	*	0.3	0.5	1.5	2.9	7.2	13.0
1999	0.9	*	*	*	*	*	0.2	0.5	1.1	3.8	7.2	11.6
Non-Hispanic white, female												
2002	2.1	*	*	*	*	*	0.2	0.6	1.9	4.6	12.2	22.0
2001	2.0	*	*	*	*	*	0.2	0.5	1.5	5.1	12.2	21.2
2000	2.1	*	*	*	*	*	0.3	0.7	1.9	6.0	12.4	20.0
1999	2.2	*	*	*	*	*	0.2	0.7	2.2	5.7	12.9	22.3
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	0.4	1.0	2.4	2.9	4.4	*
2001	0.8	*	*	*	*	*	0.7	1.4	2.1	4.2	4.3	6.9
2000	0.7	*	*	*	*	*	0.6	1.0	2.0	3.6	4.0	8.6
1999	0.8	*	*	*	*	*	0.5	1.2	2.2	3.4	4.4	8.4
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	1.8	2.9	*	*
2001	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	1.1	2.9	2.9	5.1	*
2001	1.2	*	*	*	*	*	0.9	1.7	2.6	5.4	5.4	*
2000	1.0	*	*	*	*	*	0.8	1.4	2.7	4.4	4.9	*
1999	1.0	*	*	*	*	*	*	1.5	2.7	4.1	4.4	11.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever (I00-I02)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever (I00-I02)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.5	17.5
2001	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.0	9.3	17.2
2000	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.4	9.5	16.7
1999	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	17.9
All origins 3/, male												
2002	0.7	*	*	*	*	*	0.2	0.5	1.2	2.9	6.6	11.4
2001	0.7	*	*	*	*	*	0.2	0.4	1.1	2.7	6.5	12.3
2000	0.8	*	*	*	*	*	0.3	0.6	1.4	2.8	6.5	12.1
1999	0.8	*	*	*	*	*	0.2	0.5	1.1	3.5	6.7	10.3
All origins 3/, female												
2002	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.4	20.2
2001	1.7	*	*	*	*	*	0.3	0.7	1.7	5.0	11.2	19.3
2000	1.7	*	*	*	*	0.1	0.3	0.7	2.0	5.7	11.4	18.5
1999	1.8	*	*	*	*	0.1	0.3	0.8	2.2	5.5	12.1	21.0
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	0.7	1.8	4.5	7.0	*
2001	0.4	*	*	*	*	*	*	*	1.5	4.1	5.4	*
2000	0.4	*	*	*	*	*	*	*	1.8	4.3	5.3	*
1999	0.5	*	*	*	*	*	*	0.8	1.6	4.2	7.3	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	2.1	5.8	10.7	*
2001	0.6	*	*	*	*	*	*	*	2.0	5.2	*	*
2000	0.6	*	*	*	*	*	*	*	*	5.6	6.9	*
1999	0.8	*	*	*	*	*	*	*	*	5.9	10.4	*
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.9
2001	1.3	*	*	*	*	0.1	0.3	0.6	1.4	4.0	9.5	17.5
2000	1.4	*	*	*	*	0.1	0.3	0.7	1.7	4.4	9.6	17.0
1999	1.4	*	*	*	*	0.1	0.3	0.6	1.7	4.6	10.1	18.2
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	*	0.2	0.5	1.2	2.9	6.9	11.7
2001	0.8	*	*	*	*	*	0.2	0.4	1.1	2.7	6.6	12.6
2000	0.8	*	*	*	*	*	0.3	0.6	1.4	2.8	6.7	12.4
1999	0.8	*	*	*	*	*	0.2	0.5	1.1	3.5	6.8	10.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)												
Non-Hispanic 4/, female												
2002	1.8	*	*	*	*	*	0.2	0.7	2.0	4.5	11.4	20.6
2001	1.8	*	*	*	*	*	0.3	0.7	1.7	5.0	11.4	19.5
2000	1.9	*	*	*	*	*	0.3	0.8	2.0	5.7	11.6	18.8
1999	2.0	*	*	*	*	0.1	0.3	0.8	2.3	5.5	12.2	21.2
Non-Hispanic white, both sexes												
2002	1.5	*	*	*	*	*	0.2	0.5	1.5	3.9	10.1	19.0
2001	1.5	*	*	*	*	*	0.2	0.5	1.3	4.0	10.0	18.6
2000	1.5	*	*	*	*	*	0.3	0.6	1.6	4.5	10.3	17.9
1999	1.6	*	*	*	*	0.1	0.2	0.6	1.6	4.8	10.6	19.2
Non-Hispanic white, male												
2002	0.9	*	*	*	*	*	0.2	0.4	1.1	3.0	7.2	12.5
2001	0.9	*	*	*	*	*	0.2	0.4	1.1	2.7	6.9	13.2
2000	0.9	*	*	*	*	*	0.3	0.5	1.4	2.8	7.1	12.9
1999	0.9	*	*	*	*	*	0.2	0.4	1.1	3.7	7.1	11.6
Non-Hispanic white, female												
2002	2.1	*	*	*	*	*	0.2	0.6	1.8	4.6	12.0	21.8
2001	2.0	*	*	*	*	*	0.2	0.5	1.5	5.1	12.1	20.8
2000	2.1	*	*	*	*	*	0.2	0.7	1.9	6.0	12.4	19.9
1999	2.2	*	*	*	*	*	0.2	0.7	2.2	5.6	12.8	22.2
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	1.0	2.4	2.9	4.2	*
2001	0.8	*	*	*	*	*	0.7	1.4	2.1	4.0	4.3	*
2000	0.7	*	*	*	*	*	0.6	1.0	2.0	3.5	4.0	8.3
1999	0.8	*	*	*	*	*	0.5	1.2	2.1	3.4	4.3	7.8
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	1.8	2.9	*	*
2001	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	1.1	2.9	2.9	4.8	*
2001	1.1	*	*	*	*	*	0.9	1.7	2.6	5.1	5.4	*
2000	1.0	*	*	*	*	*	0.8	1.3	2.7	4.3	4.9	*
1999	1.0	*	*	*	*	*	*	1.5	2.6	4.1	4.4	10.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatic mitral valve diseases (I05)												
All origins 3/, both sexes												
2002	0.6	*	*	*	*	0.1	0.1	0.4	1.0	2.0	5.0	7.4
2001	0.7	*	*	*	*	0.1	0.2	0.4	0.8	2.3	5.1	8.5
2000	0.7	*	*	*	*	0.1	0.2	0.4	1.0	2.7	5.4	8.5
1999	0.7	*	*	*	*	*	0.1	0.4	1.0	2.6	5.7	9.0
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.1	0.3	0.7	1.6	3.0	4.2
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.6	3.2	6.2
2000	0.4	*	*	*	*	*	0.2	0.3	0.9	1.7	3.8	6.1
1999	0.4	*	*	*	*	*	0.1	0.3	0.6	1.9	3.7	4.9
All origins 3/, female												
2002	0.9	*	*	*	*	*	0.1	0.4	1.2	2.3	6.2	8.7
2001	0.9	*	*	*	*	*	0.2	0.5	1.1	2.9	6.3	9.4
2000	1.0	*	*	*	*	*	0.2	0.5	1.2	3.5	6.4	9.5
1999	1.0	*	*	*	*	*	0.2	0.5	1.3	3.3	7.0	10.7
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	1.0	2.1	*	*
2001	0.2	*	*	*	*	*	*	*	*	2.1	*	*
2000	0.2	*	*	*	*	*	*	*	*	2.4	*	*
1999	0.3	*	*	*	*	*	*	*	*	2.4	4.8	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	3.3	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	6.9	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	*	*	*	0.1	0.4	1.0	2.0	5.0	7.5
2001	0.7	*	*	*	*	0.1	0.2	0.4	0.8	2.3	5.2	8.6
2000	0.8	*	*	*	*	0.1	0.2	0.4	1.0	2.7	5.5	8.7
1999	0.8	*	*	*	*	*	0.1	0.4	1.0	2.6	5.7	9.1
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	0.1	0.3	0.7	1.6	3.1	4.3
2001	0.4	*	*	*	*	*	0.1	0.3	0.6	1.6	3.3	6.4
2000	0.5	*	*	*	*	*	0.2	0.3	0.8	1.7	3.9	6.2
1999	0.5	*	*	*	*	*	0.1	0.3	0.6	1.9	3.7	5.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatic mitral valve diseases (I05)												
Non-Hispanic 4/, female												
2002	1.0	*	*	*	*	*	0.1	0.4	1.2	2.3	6.3	8.9
2001	1.0	*	*	*	*	*	0.2	0.5	1.1	2.9	6.4	9.5
2000	1.1	*	*	*	*	*	0.2	0.5	1.2	3.5	6.5	9.7
1999	1.1	*	*	*	*	*	0.2	0.6	1.3	3.2	7.0	10.7
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	*	0.1	0.3	0.9	2.0	5.3	8.0
2001	0.8	*	*	*	*	*	0.2	0.3	0.8	2.3	5.5	9.2
2000	0.9	*	*	*	*	*	0.2	0.4	1.0	2.8	5.8	9.3
1999	0.9	*	*	*	*	*	0.1	0.4	1.0	2.7	6.0	9.6
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	*	0.2	0.6	1.6	3.3	4.6
2001	0.5	*	*	*	*	*	0.2	0.2	0.6	1.5	3.5	6.7
2000	0.5	*	*	*	*	*	0.2	0.4	0.9	1.7	4.1	6.5
1999	0.5	*	*	*	*	*	*	0.3	0.6	2.0	3.8	5.5
Non-Hispanic white, female												
2002	1.1	*	*	*	*	*	0.1	0.4	1.1	2.3	6.7	9.5
2001	1.1	*	*	*	*	*	0.2	0.4	1.0	2.9	6.8	10.2
2000	1.2	*	*	*	*	*	0.2	0.5	1.1	3.7	7.0	10.5
1999	1.3	*	*	*	*	*	0.1	0.4	1.3	3.3	7.5	11.2
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.6	1.6	1.3	*	*
2001	0.5	*	*	*	*	*	0.4	0.9	1.2	2.4	2.3	*
2000	0.4	*	*	*	*	*	*	*	1.1	1.8	2.2	*
1999	0.5	*	*	*	*	*	*	0.9	1.2	2.5	2.5	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	1.9	*	*	*
2001	0.7	*	*	*	*	*	*	1.1	1.7	2.9	*	*
2000	0.5	*	*	*	*	*	*	*	1.5	2.3	*	*
1999	0.6	*	*	*	*	*	*	1.2	1.6	3.1	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatic aortic valve diseases (I06)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.4	1.0
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatic aortic valve diseases (I06)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	1.2
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of both mitral and aortic valves (I08.0)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.5	3.5
2001	0.1	*	*	*	*	*	*	*	0.1	0.5	1.0	2.5
2000	0.1	*	*	*	*	*	*	*	0.1	0.4	0.9	2.0
1999	0.1	*	*	*	*	*	*	*	0.1	0.4	0.8	2.0
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.5	1.4	2.4
2001	0.1	*	*	*	*	*	*	*	*	0.4	1.0	2.3
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.9
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.0
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.1	2.5
2000	0.1	*	*	*	*	*	*	*	*	0.4	1.0	2.1
1999	0.1	*	*	*	*	*	*	*	0.2	0.4	0.8	2.2
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.5	3.6
2001	0.1	*	*	*	*	*	*	*	0.1	0.5	1.0	2.5
2000	0.1	*	*	*	*	*	*	*	0.1	0.3	0.9	2.1
1999	0.1	*	*	*	*	*	*	*	0.1	0.4	0.8	2.0
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.5	1.4	2.5
2001	0.1	*	*	*	*	*	*	*	*	0.5	1.0	2.3
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
1999	0.1	*	*	*	*	*	*	*	*	0.4	0.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of both mitral and aortic valves (I08.0)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.6	4.1
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.1	2.6
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.1
1999	0.2	*	*	*	*	*	*	*	*	0.4	0.8	2.2
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.7	3.8
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.1	2.6
2000	0.1	*	*	*	*	*	*	*	0.1	0.3	1.0	2.1
1999	0.1	*	*	*	*	*	*	*	0.1	0.4	0.8	2.1
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.5	2.7
2001	0.1	*	*	*	*	*	*	*	*	0.5	1.0	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	2.0
1999	0.1	*	*	*	*	*	*	*	*	0.5	0.7	*
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	*	*	0.5	1.8	4.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.7
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.1
1999	0.2	*	*	*	*	*	*	*	*	0.4	0.9	2.2
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.1	0.5	1.3	2.7	5.6
2001	0.4	*	*	*	*	*	0.1	0.1	0.5	1.2	3.0	5.9
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.3	3.2	5.8
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.5	3.4	6.6
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.4	0.8	1.9	3.8
2001	0.2	*	*	*	*	*	*	0.1	0.4	0.7	2.0	3.6
2000	0.2	*	*	*	*	*	*	0.1	0.4	0.8	2.0	4.0
1999	0.2	*	*	*	*	*	*	0.1	0.3	1.1	2.2	3.7
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.6	1.6	3.2	6.3
2001	0.5	*	*	*	*	*	0.1	0.2	0.5	1.6	3.7	6.8
2000	0.6	*	*	*	*	*	0.1	0.2	0.7	1.8	4.0	6.6
1999	0.6	*	*	*	*	*	0.1	0.2	0.7	1.8	4.2	7.8
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.1	0.5	1.2	2.7	5.7
2001	0.4	*	*	*	*	*	0.1	0.1	0.4	1.2	3.1	6.0
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.4	3.2	5.9
1999	0.5	*	*	*	*	*	0.1	0.1	0.6	1.5	3.4	6.8
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.4	0.8	2.0	3.9
2001	0.2	*	*	*	*	*	*	0.1	0.4	0.7	2.0	3.7
2000	0.3	*	*	*	*	*	*	0.1	0.4	0.8	2.0	4.1
1999	0.3	*	*	*	*	*	*	*	0.3	1.2	2.3	3.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.6	1.6	3.1	6.4
2001	0.6	*	*	*	*	*	*	0.2	0.5	1.6	3.7	6.9
2000	0.6	*	*	*	*	*	0.1	0.2	0.7	1.8	4.0	6.7
1999	0.7	*	*	*	*	*	0.1	0.2	0.7	1.7	4.2	8.0
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	*	0.1	0.5	1.3	2.8	6.0
2001	0.4	*	*	*	*	*	*	0.1	0.4	1.2	3.2	6.4
2000	0.5	*	*	*	*	*	0.1	0.1	0.5	1.4	3.4	6.1
1999	0.5	*	*	*	*	*	*	0.1	0.5	1.5	3.6	7.1
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	*	0.3	0.8	2.1	4.1
2001	0.2	*	*	*	*	*	*	*	0.4	0.7	2.1	3.8
2000	0.3	*	*	*	*	*	*	*	0.4	0.8	2.2	4.2
1999	0.3	*	*	*	*	*	*	*	0.3	1.2	2.4	4.1
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	0.6	1.7	3.2	6.8
2001	0.6	*	*	*	*	*	*	*	0.4	1.6	3.9	7.4
2000	0.7	*	*	*	*	*	*	0.2	0.6	1.9	4.2	6.9
1999	0.8	*	*	*	*	*	*	0.2	0.7	1.8	4.4	8.3
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	0.5	*	1.3	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive diseases (I10-I13)												
All origins 3/, both sexes												
2002	17.2	*	*	*	0.1	1.0	3.7	9.9	19.2	39.5	104.6	390.8
2001	16.4	*	*	*	0.1	1.0	3.4	9.0	17.7	37.5	103.1	384.7
2000	15.9	*	*	*	0.1	0.8	3.1	8.3	17.8	37.0	100.7	380.3
1999	15.4	*	*	*	0.1	0.8	3.1	7.9	17.3	36.8	100.1	366.0
All origins 3/, male												
2002	14.5	*	*	*	0.2	1.4	5.0	13.2	24.8	44.6	103.7	311.1
2001	13.5	*	*	*	0.2	1.2	4.5	12.0	21.9	42.7	101.0	309.4
2000	13.0	*	*	*	0.2	1.0	4.0	11.2	22.4	42.0	98.8	309.3
1999	12.6	*	*	*	0.2	1.1	4.1	10.3	21.4	41.1	99.3	297.3
All origins 3/, female												
2002	19.9	*	*	*	*	0.7	2.4	6.7	14.1	35.2	105.2	425.4
2001	19.2	*	*	*	*	0.7	2.3	6.1	13.8	33.2	104.5	416.3
2000	18.6	*	*	*	0.1	0.6	2.2	5.5	13.5	32.9	101.9	409.3
1999	18.1	*	*	*	*	0.5	2.1	5.6	13.6	33.3	100.7	393.7
Hispanic, both sexes												
2002	6.5	*	*	*	*	0.6	2.3	6.3	19.7	40.7	107.1	337.0
2001	6.1	*	*	*	*	0.5	2.2	6.9	15.7	40.6	103.2	338.3
2000	5.5	*	*	*	*	0.5	1.9	6.0	15.5	35.9	98.2	316.5
1999	5.4	*	*	*	*	0.4	1.8	5.9	15.1	35.1	101.2	305.2
Hispanic, male												
2002	6.2	*	*	*	*	0.8	3.2	8.9	26.0	47.2	115.7	294.5
2001	5.7	*	*	*	*	0.6	3.4	9.7	19.3	46.5	106.9	304.7
2000	5.2	*	*	*	*	0.7	2.5	8.7	19.4	41.6	112.1	276.1
1999	4.9	*	*	*	*	0.6	2.6	7.5	19.3	42.4	101.7	267.4
Hispanic, female												
2002	6.9	*	*	*	*	*	1.2	3.7	14.1	35.6	101.2	358.1
2001	6.5	*	*	*	*	*	1.0	4.2	12.5	35.9	100.6	354.7
2000	5.8	*	*	*	*	*	1.3	3.3	12.0	31.5	88.8	336.3
1999	5.9	*	*	*	*	*	1.0	4.3	11.4	29.4	101.0	323.4
Non-Hispanic 4/, both sexes												
2002	18.8	*	*	*	0.1	1.1	3.8	10.2	19.1	39.1	104.1	391.7
2001	17.9	*	*	*	0.1	1.0	3.5	9.1	17.7	37.1	102.8	385.4
2000	17.3	*	*	*	0.2	0.8	3.2	8.4	17.9	36.8	100.4	381.4
1999	16.7	*	*	*	0.1	0.8	3.2	8.0	17.3	36.7	99.7	366.9
Non-Hispanic 4/, male												
2002	15.7	*	*	*	0.2	1.5	5.2	13.5	24.5	44.0	102.6	310.5
2001	14.6	*	*	*	0.2	1.3	4.6	12.1	21.8	42.2	100.4	308.4
2000	14.1	*	*	*	0.2	1.0	4.2	11.3	22.5	41.6	97.7	309.5
1999	13.6	*	*	*	0.2	1.1	4.3	10.4	21.3	40.8	98.7	297.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive diseases (I10-I13)												
Non-Hispanic 4/, female												
2002	21.7	*	*	*	*	0.7	2.5	7.0	14.1	35.0	105.1	426.7
2001	21.0	*	*	*	*	0.7	2.4	6.3	13.9	32.9	104.4	417.5
2000	20.3	*	*	*	0.1	0.6	2.3	5.6	13.6	32.9	102.2	410.5
1999	19.7	*	*	*	*	0.5	2.2	5.7	13.6	33.4	100.4	394.7
Non-Hispanic white, both sexes												
2002	17.3	*	*	*	*	0.6	2.3	6.7	13.0	29.8	90.0	368.4
2001	16.5	*	*	*	*	0.6	2.0	5.6	12.0	27.9	89.6	363.9
2000	15.7	*	*	*	*	0.4	2.0	5.1	11.7	27.6	86.6	357.0
1999	15.1	*	*	*	*	0.5	1.7	4.8	11.4	27.3	85.1	343.1
Non-Hispanic white, male												
2002	13.8	*	*	*	*	0.8	3.4	9.4	17.3	34.0	88.1	289.1
2001	12.8	*	*	*	*	0.8	2.9	7.8	15.5	32.3	87.0	286.3
2000	12.2	*	*	*	*	0.6	2.8	7.4	15.4	32.0	84.4	285.6
1999	11.7	*	*	*	*	0.8	2.5	6.6	14.6	30.7	84.1	272.3
Non-Hispanic white, female												
2002	20.7	*	*	*	*	0.3	1.3	4.0	8.9	26.2	91.2	402.6
2001	20.0	*	*	*	*	0.3	1.1	3.4	8.7	24.0	91.3	396.2
2000	19.1	*	*	*	*	0.2	1.2	2.9	8.3	23.9	88.1	385.8
1999	18.4	*	*	*	*	0.2	0.9	3.0	8.3	24.5	85.8	371.3
Non-Hispanic black, both sexes												
2002	31.7	*	*	*	0.4	4.1	13.2	36.7	72.8	126.6	273.9	691.7
2001	30.1	*	*	*	0.4	3.7	12.8	35.5	66.9	124.6	259.7	670.8
2000	29.9	*	*	*	0.6	3.2	11.4	33.7	71.1	127.1	264.1	680.8
1999	29.5	*	*	*	0.6	2.8	12.8	33.0	68.1	127.6	272.8	646.7
Non-Hispanic black, male												
2002	30.2	*	*	*	*	5.6	17.0	47.0	93.5	146.9	290.7	603.9
2001	28.5	*	*	*	*	4.5	15.9	46.8	81.9	147.3	275.8	611.0
2000	27.6	*	*	*	*	3.8	14.0	43.4	89.7	144.5	270.0	609.4
1999	27.6	*	*	*	0.8	3.3	15.9	42.1	83.7	146.7	288.3	595.4
Non-Hispanic black, female												
2002	33.0	*	*	*	*	2.7	9.7	27.9	56.1	112.3	264.5	725.3
2001	31.5	*	*	*	*	3.0	10.1	25.8	54.8	108.6	250.7	693.0
2000	31.9	*	*	*	*	2.7	9.2	25.4	56.2	114.8	260.8	707.3
1999	31.3	*	*	*	*	2.4	10.0	25.2	55.5	114.2	264.2	665.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension (I10)												
All origins 3/, both sexes												
2002	4.0	*	*	*	*	0.1	0.4	1.3	3.2	8.6	26.3	112.0
2001	3.9	*	*	*	*	0.1	0.3	1.3	3.4	8.4	25.9	107.4
2000	3.7	*	*	*	*	0.1	0.3	1.4	3.4	8.2	24.1	101.1
1999	3.6	*	*	*	*	0.1	0.3	1.2	3.3	8.1	24.0	98.1
All origins 3/, male												
2002	2.8	*	*	*	*	*	0.5	1.7	4.0	9.4	23.3	74.2
2001	2.8	*	*	*	*	0.1	0.4	1.7	4.2	9.3	23.2	74.8
2000	2.6	*	*	*	*	*	0.4	1.9	4.2	8.9	22.0	70.3
1999	2.5	*	*	*	*	*	0.4	1.5	4.1	9.2	21.9	68.1
All origins 3/, female												
2002	5.2	*	*	*	*	*	0.3	1.0	2.5	8.0	28.2	128.3
2001	5.0	*	*	*	*	*	0.3	0.9	2.7	7.7	27.6	121.1
2000	4.6	*	*	*	*	*	0.3	0.9	2.6	7.5	25.5	113.6
1999	4.5	*	*	*	*	*	0.2	1.0	2.7	7.3	25.4	110.2
Hispanic, both sexes												
2002	1.1	*	*	*	*	*	*	0.5	2.2	5.6	20.4	80.9
2001	1.1	*	*	*	*	*	*	0.8	1.5	6.2	24.2	90.0
2000	1.1	*	*	*	*	*	*	0.8	2.9	6.1	21.3	80.3
1999	1.0	*	*	*	*	*	*	0.8	2.2	5.6	21.2	72.3
Hispanic, male												
2002	0.7	*	*	*	*	*	*	*	2.6	5.2	14.8	58.6
2001	0.8	*	*	*	*	*	*	*	*	6.2	24.6	53.3
2000	1.0	*	*	*	*	*	*	1.3	3.7	6.3	25.1	72.6
1999	0.8	*	*	*	*	*	*	*	*	7.0	21.9	51.8
Hispanic, female												
2002	1.4	*	*	*	*	*	*	*	1.9	5.9	24.2	92.0
2001	1.4	*	*	*	*	*	*	*	*	6.1	23.9	108.0
2000	1.2	*	*	*	*	*	*	*	*	6.0	18.8	84.1
1999	1.2	*	*	*	*	*	*	*	*	4.5	20.8	82.1
Non-Hispanic 4/, both sexes												
2002	4.5	*	*	*	*	0.1	0.4	1.4	3.3	8.8	26.5	112.7
2001	4.3	*	*	*	*	0.1	0.4	1.3	3.5	8.5	25.9	107.9
2000	4.0	*	*	*	*	0.1	0.4	1.4	3.4	8.3	24.2	101.4
1999	3.9	*	*	*	*	0.1	0.3	1.3	3.4	8.3	24.0	98.8
Non-Hispanic 4/, male												
2002	3.1	*	*	*	*	*	0.5	1.8	4.1	9.6	23.6	74.5
2001	3.0	*	*	*	*	*	0.4	1.7	4.3	9.5	23.0	75.4
2000	2.9	*	*	*	*	*	0.4	1.9	4.2	9.0	21.7	70.0
1999	2.8	*	*	*	*	*	0.4	1.5	4.2	9.3	21.7	68.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension (I10)												
Non-Hispanic 4/, female												
2002	5.8	*	*	*	*	*	0.3	1.0	2.6	8.1	28.3	129.2
2001	5.5	*	*	*	*	*	0.3	0.9	2.8	7.8	27.7	121.3
2000	5.1	*	*	*	*	*	0.3	0.9	2.6	7.6	25.8	114.2
1999	5.0	*	*	*	*	*	0.2	1.0	2.7	7.4	25.5	111.0
Non-Hispanic white, both sexes												
2002	4.6	*	*	*	*	*	0.3	1.0	2.8	7.6	24.3	109.1
2001	4.3	*	*	*	*	*	0.2	0.9	2.8	7.1	23.9	104.8
2000	4.0	*	*	*	*	*	0.3	1.0	2.7	7.2	22.2	97.7
1999	3.9	*	*	*	*	*	0.2	0.9	2.7	7.2	21.9	94.7
Non-Hispanic white, male												
2002	3.1	*	*	*	*	*	0.4	1.4	3.6	8.4	21.2	71.3
2001	2.9	*	*	*	*	*	0.3	1.3	3.5	8.0	20.5	71.6
2000	2.7	*	*	*	*	*	0.3	1.4	3.3	8.0	19.5	65.7
1999	2.7	*	*	*	*	*	0.3	1.2	3.5	8.1	19.5	62.7
Non-Hispanic white, female												
2002	6.0	*	*	*	*	*	0.2	0.7	2.1	6.9	26.4	125.5
2001	5.7	*	*	*	*	*	0.2	0.6	2.2	6.4	26.1	118.7
2000	5.3	*	*	*	*	*	0.2	0.6	2.1	6.6	23.9	110.5
1999	5.1	*	*	*	*	*	*	0.6	2.1	6.5	23.5	107.5
Non-Hispanic black, both sexes												
2002	5.1	*	*	*	*	*	1.2	4.1	8.0	21.0	53.8	160.3
2001	5.1	*	*	*	*	*	1.1	4.1	10.1	22.0	50.8	152.3
2000	4.9	*	*	*	*	*	1.0	4.5	9.9	19.1	48.3	150.7
1999	4.8	*	*	*	*	*	1.1	4.1	9.5	19.3	49.3	145.2
Non-Hispanic black, male												
2002	4.2	*	*	*	*	*	1.5	4.9	9.6	23.0	56.3	119.2
2001	4.5	*	*	*	*	*	1.3	5.3	12.6	25.5	56.4	132.8
2000	4.2	*	*	*	*	*	1.0	5.9	13.3	21.1	49.7	124.5
1999	4.0	*	*	*	*	*	1.2	4.4	11.3	22.7	50.7	126.0
Non-Hispanic black, female												
2002	6.0	*	*	*	*	*	0.9	3.4	6.7	19.6	52.4	176.0
2001	5.7	*	*	*	*	*	1.0	3.1	8.1	19.6	47.7	159.5
2000	5.5	*	*	*	*	*	1.0	3.3	7.1	17.7	47.6	160.5
1999	5.5	*	*	*	*	*	1.0	3.8	8.1	16.9	48.5	152.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All origins 3/, both sexes												
2002	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
All origins 3/, male												
2002	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
All origins 3/, female												
2002	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5
Hispanic, both sexes												
2002	3.7	*	*	*	*	0.4	1.7	4.9	12.2	22.4	51.4	165.7
2001	3.3	*	*	*	*	0.3	1.7	4.9	10.1	21.1	46.9	151.9
2000	2.8	*	*	*	*	0.3	1.2	3.8	8.2	18.4	46.0	143.3
1999	2.7	*	*	*	*	*	1.2	3.3	8.9	16.8	44.8	152.2
Hispanic, male												
2002	3.9	*	*	*	*	0.6	2.6	7.4	17.0	28.5	61.1	137.2
2001	3.4	*	*	*	*	*	2.7	7.0	13.8	26.4	48.3	144.7
2000	2.7	*	*	*	*	*	1.6	5.4	10.7	22.4	50.6	112.9
1999	2.6	*	*	*	*	*	1.8	4.7	12.5	20.9	42.8	107.8
Hispanic, female												
2002	3.4	*	*	*	*	*	*	2.5	7.9	17.5	44.7	179.9
2001	3.2	*	*	*	*	*	*	2.7	6.7	17.0	45.9	155.5
2000	2.9	*	*	*	*	*	0.8	2.1	6.1	15.3	42.9	158.3
1999	2.9	*	*	*	*	*	*	2.0	5.8	13.6	46.1	173.7
Non-Hispanic 4/, both sexes												
2002	10.0	*	*	*	0.1	0.8	2.8	7.4	12.4	21.0	49.3	187.1
2001	9.4	*	*	*	*	0.7	2.5	6.3	10.9	19.6	48.6	190.3
2000	9.2	*	*	*	0.1	0.6	2.3	5.7	11.2	19.5	48.1	194.3
1999	8.8	*	*	*	0.1	0.6	2.2	5.4	10.7	19.1	47.6	185.3
Non-Hispanic 4/, male												
2002	8.9	*	*	*	0.1	1.2	4.0	10.0	16.8	24.6	48.5	133.7
2001	8.1	*	*	*	*	0.9	3.4	8.6	14.2	23.3	48.3	133.4
2000	7.8	*	*	*	0.1	0.7	3.0	7.9	14.6	23.5	46.0	139.6
1999	7.4	*	*	*	0.1	0.8	3.1	7.2	13.6	21.7	46.7	134.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Non-Hispanic 4/, female												
2002	11.1	*	*	*	*	0.5	1.7	4.8	8.4	18.0	49.9	210.2
2001	10.7	*	*	*	*	0.4	1.7	4.2	7.8	16.5	48.8	214.0
2000	10.6	*	*	*	*	0.4	1.6	3.6	8.0	16.2	49.5	216.5
1999	10.2	*	*	*	*	0.3	1.4	3.7	8.0	16.9	48.2	205.8
Non-Hispanic white, both sexes												
2002	9.1	*	*	*	*	0.4	1.9	5.1	8.8	16.1	42.2	177.0
2001	8.6	*	*	*	*	0.4	1.6	4.1	7.6	14.8	42.2	180.4
2000	8.3	*	*	*	*	0.3	1.5	3.6	7.5	14.5	41.2	182.5
1999	7.9	*	*	*	*	0.3	1.3	3.4	7.3	14.1	40.6	175.2
Non-Hispanic white, male												
2002	7.6	*	*	*	*	0.7	2.8	7.3	12.2	19.0	41.2	123.5
2001	6.9	*	*	*	*	0.6	2.3	5.9	10.5	18.1	41.5	122.0
2000	6.6	*	*	*	*	0.4	2.1	5.4	10.3	18.0	38.8	129.5
1999	6.2	*	*	*	*	0.6	2.0	4.9	9.6	16.2	39.4	123.8
Non-Hispanic white, female												
2002	10.5	*	*	*	*	0.2	1.0	2.9	5.5	13.6	42.9	200.1
2001	10.2	*	*	*	*	0.2	0.8	2.3	4.9	11.9	42.6	204.7
2000	9.9	*	*	*	*	0.2	0.8	1.9	4.9	11.6	42.7	203.9
1999	9.6	*	*	*	*	*	0.6	2.0	5.0	12.4	41.4	195.7
Non-Hispanic black, both sexes												
2002	17.7	*	*	*	*	2.9	8.8	24.8	45.1	68.3	134.2	322.1
2001	16.5	*	*	*	*	2.2	8.5	23.2	39.4	66.1	125.6	328.7
2000	16.6	*	*	*	*	2.2	7.5	21.6	43.1	67.6	132.7	344.1
1999	16.0	*	*	*	0.4	1.9	8.1	21.3	40.1	67.4	132.7	308.5
Non-Hispanic black, male												
2002	18.0	*	*	*	*	4.1	11.7	31.9	61.9	84.0	141.8	264.7
2001	16.3	*	*	*	*	2.7	10.5	30.4	50.1	79.8	137.4	288.7
2000	16.1	*	*	*	*	2.7	9.2	28.4	56.0	81.9	139.5	275.9
1999	15.6	*	*	*	*	2.2	10.3	26.8	51.1	80.6	143.2	266.2
Non-Hispanic black, female												
2002	17.4	*	*	*	*	1.8	6.2	18.7	31.7	57.1	129.8	344.1
2001	16.7	*	*	*	*	1.7	6.7	17.1	30.8	56.5	118.9	343.5
2000	17.1	*	*	*	*	1.8	5.9	15.8	32.7	57.5	128.9	369.4
1999	16.4	*	*	*	*	1.6	6.1	16.5	31.3	58.1	126.8	324.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive renal disease (I12)												
All origins 3/, both sexes												
2002	3.0	*	*	*	*	0.2	0.4	1.0	2.5	7.3	21.9	68.4
2001	2.9	*	*	*	*	0.2	0.4	1.1	2.4	7.0	21.8	64.5
2000	2.8	*	*	*	*	0.1	0.4	1.0	2.6	7.0	21.4	61.8
1999	2.5	*	*	*	*	0.1	0.4	1.0	2.2	7.1	19.6	53.9
All origins 3/, male												
2002	2.6	*	*	*	*	0.2	0.5	1.2	2.7	7.7	24.1	79.2
2001	2.4	*	*	*	*	0.2	0.5	1.3	2.3	7.3	22.6	76.0
2000	2.4	*	*	*	*	0.1	0.5	1.1	2.8	6.8	23.2	72.0
1999	2.2	*	*	*	*	0.2	0.5	1.2	2.4	7.4	21.1	64.3
All origins 3/, female												
2002	3.4	*	*	*	*	0.1	0.3	0.8	2.3	7.1	20.6	63.7
2001	3.3	*	*	*	*	0.2	0.3	0.9	2.5	6.8	21.3	59.7
2000	3.1	*	*	*	*	0.1	0.3	0.8	2.4	7.1	20.2	57.7
1999	2.8	*	*	*	*	*	0.3	0.7	2.1	6.8	18.6	49.8
Hispanic, both sexes												
2002	1.4	*	*	*	*	*	*	*	4.0	9.9	26.9	67.6
2001	1.3	*	*	*	*	*	*	1.0	3.0	10.3	24.7	70.0
2000	1.2	*	*	*	*	*	*	0.9	3.5	8.9	24.5	67.7
1999	1.2	*	*	*	*	*	*	1.3	2.8	9.7	27.1	57.5
Hispanic, male												
2002	1.2	*	*	*	*	*	*	*	4.5	10.1	31.6	73.6
2001	1.1	*	*	*	*	*	*	1.2	*	10.6	26.4	76.2
2000	1.1	*	*	*	*	*	*	1.5	4.4	8.9	27.5	68.5
1999	1.2	*	*	*	*	*	*	1.4	3.0	11.4	29.7	79.8
Hispanic, female												
2002	1.5	*	*	*	*	*	*	*	3.6	9.7	23.6	64.7
2001	1.5	*	*	*	*	*	*	*	3.8	10.1	23.6	67.0
2000	1.3	*	*	*	*	*	*	*	2.8	9.0	22.5	67.3
1999	1.3	*	*	*	*	*	*	*	2.6	8.3	25.3	46.8
Non-Hispanic 4/, both sexes												
2002	3.2	*	*	*	*	0.2	0.4	1.0	2.4	7.1	21.7	68.4
2001	3.1	*	*	*	*	0.2	0.4	1.1	2.3	6.8	21.6	64.2
2000	3.0	*	*	*	*	0.1	0.4	1.0	2.5	6.8	21.2	61.4
1999	2.7	*	*	*	*	0.1	0.4	0.9	2.2	6.9	19.2	53.5
Non-Hispanic 4/, male												
2002	2.8	*	*	*	*	0.2	0.5	1.2	2.5	7.5	23.6	79.4
2001	2.6	*	*	*	*	0.2	0.5	1.3	2.2	7.1	22.3	75.7
2000	2.5	*	*	*	*	*	0.6	1.1	2.7	6.7	22.9	72.0
1999	2.4	*	*	*	*	0.1	0.5	1.2	2.3	7.1	20.6	63.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive renal disease (I12)												
Non-Hispanic 4/, female												
2002	3.6	*	*	*	*	0.1	0.3	0.9	2.2	6.8	20.4	63.6
2001	3.5	*	*	*	*	0.2	0.3	0.9	2.3	6.6	21.1	59.4
2000	3.4	*	*	*	*	0.1	0.3	0.8	2.3	7.0	20.0	57.2
1999	3.0	*	*	*	*	*	0.3	0.7	2.0	6.7	18.3	49.5
Non-Hispanic white, both sexes												
2002	2.8	*	*	*	*	*	0.1	0.5	1.0	4.9	17.8	61.4
2001	2.7	*	*	*	*	*	0.1	0.4	1.1	4.6	17.9	58.0
2000	2.5	*	*	*	*	*	0.1	0.3	1.2	4.3	17.7	55.0
1999	2.2	*	*	*	*	*	0.2	0.3	0.9	4.6	15.4	47.0
Non-Hispanic white, male												
2002	2.4	*	*	*	*	*	0.2	0.5	1.1	5.2	19.8	74.1
2001	2.2	*	*	*	*	*	0.2	0.4	1.0	4.8	19.1	70.5
2000	2.2	*	*	*	*	*	0.2	0.4	1.3	4.4	20.1	66.0
1999	1.9	*	*	*	*	*	0.2	0.4	1.0	4.9	17.0	57.1
Non-Hispanic white, female												
2002	3.1	*	*	*	*	*	*	0.4	0.9	4.6	16.5	56.0
2001	3.1	*	*	*	*	*	*	0.4	1.1	4.5	17.1	52.8
2000	2.9	*	*	*	*	*	*	0.3	1.0	4.3	16.2	50.6
1999	2.5	*	*	*	*	*	*	0.3	0.8	4.3	14.3	43.0
Non-Hispanic black, both sexes												
2002	6.4	*	*	*	*	0.6	2.2	5.1	13.4	27.3	66.2	152.1
2001	6.1	*	*	*	*	0.8	2.0	6.0	12.3	26.4	63.1	139.2
2000	6.1	*	*	*	*	0.5	2.2	5.4	13.5	30.4	60.6	135.1
1999	5.9	*	*	*	*	0.4	2.1	5.3	12.3	28.2	63.6	127.1
Non-Hispanic black, male												
2002	5.7	*	*	*	*	*	2.8	6.4	14.4	29.6	73.3	157.5
2001	5.5	*	*	*	*	0.9	2.6	7.8	13.3	30.0	63.3	144.4
2000	5.3	*	*	*	*	*	2.8	6.2	15.1	30.4	60.8	144.4
1999	5.4	*	*	*	*	*	2.5	7.7	13.5	29.6	68.5	142.6
Non-Hispanic black, female												
2002	7.0	*	*	*	*	*	1.7	4.0	12.7	25.7	62.2	150.1
2001	6.7	*	*	*	*	0.7	1.6	4.5	11.6	23.9	63.0	137.3
2000	6.9	*	*	*	*	*	1.7	4.7	12.1	30.4	60.4	131.7
1999	6.3	*	*	*	*	*	1.6	3.3	11.3	27.2	60.9	121.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All origins 3/, both sexes												
2002	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
All origins 3/, male												
2002	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
All origins 3/, female												
2002	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	*	1.3	2.9	8.5	22.7
2001	0.4	*	*	*	*	*	*	*	1.1	3.0	7.4	26.3
2000	0.4	*	*	*	*	*	*	*	*	2.5	6.3	25.2
1999	0.4	*	*	*	*	*	*	*	*	3.1	8.1	23.2
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	8.2	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	8.7	21.6
2001	0.5	*	*	*	*	*	*	*	*	*	7.3	24.2
2000	0.4	*	*	*	*	*	*	*	*	*	*	26.7
1999	0.5	*	*	*	*	*	*	*	*	*	8.7	20.8
Non-Hispanic 4/, both sexes												
2002	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.1	6.7	23.5
2001	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.7	23.1
2000	1.1	*	*	*	*	0.1	0.2	0.4	0.9	2.2	7.0	24.2
1999	1.3	*	*	*	*	0.1	0.3	0.4	1.1	2.5	8.9	29.3
Non-Hispanic 4/, male												
2002	0.9	*	*	*	*	*	0.2	0.6	1.2	2.3	6.9	22.9
2001	0.9	*	*	*	*	*	0.3	0.5	1.0	2.4	6.7	23.8
2000	0.9	*	*	*	*	*	0.2	0.5	0.9	2.4	7.0	27.9
1999	1.1	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Non-Hispanic 4/, female												
2002	1.2	*	*	*	*	*	0.2	0.3	0.8	2.0	6.5	23.7
2001	1.2	*	*	*	*	*	0.2	0.2	0.9	2.0	6.7	22.8
2000	1.2	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.8
1999	1.5	*	*	*	*	*	0.2	0.3	0.9	2.4	8.4	28.4
Non-Hispanic white, both sexes												
2002	0.9	*	*	*	*	*	*	0.1	0.4	1.2	5.6	20.8
2001	0.9	*	*	*	*	*	*	0.1	0.5	1.4	5.6	20.7
2000	0.9	*	*	*	*	*	0.1	0.1	0.4	1.5	5.6	21.8
1999	1.1	*	*	*	*	*	*	0.2	0.5	1.4	7.3	26.1
Non-Hispanic white, male												
2002	0.7	*	*	*	*	*	*	*	0.5	1.4	5.9	20.2
2001	0.7	*	*	*	*	*	*	0.2	0.4	1.5	5.8	22.2
2000	0.7	*	*	*	*	*	*	0.2	0.4	1.6	6.1	24.3
1999	0.9	*	*	*	*	*	*	0.2	0.5	1.6	8.3	28.7
Non-Hispanic white, female												
2002	1.1	*	*	*	*	*	*	*	0.3	1.1	5.4	21.0
2001	1.1	*	*	*	*	*	*	*	0.5	1.3	5.4	20.1
2000	1.1	*	*	*	*	*	*	*	0.3	1.4	5.2	20.8
1999	1.2	*	*	*	*	*	*	*	0.4	1.3	6.6	25.1
Non-Hispanic black, both sexes												
2002	2.5	*	*	*	*	*	1.0	2.8	6.2	10.0	19.8	57.2
2001	2.3	*	*	*	*	0.4	1.2	2.1	5.1	10.1	20.2	50.6
2000	2.3	*	*	*	*	*	0.8	2.2	4.7	10.0	22.6	50.9
1999	2.8	*	*	*	*	*	1.5	2.3	6.2	12.7	27.3	66.0
Non-Hispanic black, male												
2002	2.3	*	*	*	*	*	1.0	3.9	7.7	10.2	19.3	62.4
2001	2.2	*	*	*	*	*	1.6	3.2	6.0	12.0	18.6	45.0
2000	2.1	*	*	*	*	*	1.0	2.9	5.4	11.1	20.1	64.6
1999	2.6	*	*	*	*	*	1.8	3.3	7.8	13.8	26.0	60.6
Non-Hispanic black, female												
2002	2.6	*	*	*	*	*	0.9	1.8	5.0	9.9	20.1	55.2
2001	2.4	*	*	*	*	*	0.8	1.2	4.3	8.7	21.1	52.6
2000	2.4	*	*	*	*	*	*	1.6	4.2	9.2	24.0	45.8
1999	3.1	*	*	*	*	*	1.2	1.5	4.8	12.0	28.0	68.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart disease (I20-I25)												
All origins 3/, both sexes												
2002	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1220.2	3,775.0
2001	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1271.4	3,958.3
2000	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1324.2	4,182.5
1999	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1384.8	4,318.1
All origins 3/, male												
2002	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1598.9	4,205.3
2001	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.8	1663.6	4,396.0
2000	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1736.8	4,701.7
1999	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1813.7	4,837.5
All origins 3/, female												
2002	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1013.6	3,775.2
2000	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1055.2	3,971.0
1999	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1106.3	4,108.8
Hispanic, both sexes												
2002	54.0	*	*	*	*	1.2	7.6	38.0	135.9	382.0	1010.8	3,182.9
2001	56.0	*	*	*	*	1.6	6.8	38.7	130.4	401.3	1100.8	3,624.0
2000	55.9	*	*	*	*	1.1	7.6	39.5	143.6	405.3	1111.3	3,697.2
1999	58.7	*	*	*	*	1.4	7.4	44.8	150.4	433.5	1178.0	3,876.2
Hispanic, male												
2002	56.5	*	*	*	*	1.9	11.4	58.0	204.0	530.4	1271.7	3,374.9
2001	58.0	*	*	*	*	2.4	10.5	60.3	192.4	560.7	1395.6	3,833.9
2000	58.2	*	*	*	*	1.6	12.1	62.7	214.8	561.5	1420.0	3,865.6
1999	60.8	*	*	*	*	2.1	12.2	71.3	218.3	608.5	1495.9	3,987.7
Hispanic, female												
2002	51.4	*	*	*	*	*	3.5	18.1	74.6	263.9	832.2	3,087.8
2001	53.8	*	*	*	*	*	2.8	17.4	75.4	275.6	901.5	3,521.4
2000	53.6	*	*	*	*	*	2.8	16.8	80.6	282.5	904.0	3,614.6
1999	56.6	*	*	*	*	*	2.3	19.0	90.5	296.4	966.5	3,822.4
Non-Hispanic 4/, both sexes												
2002	189.0	*	*	*	0.3	2.8	18.4	66.1	180.1	462.1	1226.5	3,787.0
2001	193.6	0.7	*	*	0.4	3.1	18.0	66.1	187.3	480.9	1274.9	3,959.4
2000	200.5	0.8	*	*	0.3	2.6	17.7	68.1	199.5	509.0	1328.8	4,187.0
1999	207.1	0.8	*	0.1	0.4	2.7	18.3	69.4	206.3	540.3	1388.1	4,319.6
Non-Hispanic 4/, male												
2002	197.6	*	*	*	0.5	4.1	28.3	103.1	266.5	644.3	1609.7	4,228.4
2001	200.4	*	*	*	0.6	4.5	27.5	102.8	276.1	667.7	1670.1	4,408.5
2000	207.6	*	*	*	0.4	3.9	27.4	106.7	294.0	710.4	1744.6	4,721.1
1999	214.1	*	*	*	0.5	4.0	28.3	108.8	303.4	754.2	1820.2	4,854.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart disease (I20-I25)												
Non-Hispanic 4/, female												
2002	180.7	*	*	*	0.2	1.5	8.8	30.5	100.0	309.9	972.5	3,596.5
2001	187.2	*	*	*	0.2	1.6	8.7	30.8	105.0	325.6	1015.5	3,772.7
2000	193.8	*	*	*	0.2	1.3	8.2	31.0	112.1	342.8	1057.9	3,971.0
1999	200.6	*	*	*	0.3	1.5	8.5	31.5	116.6	364.6	1107.8	4,105.4
Non-Hispanic white, both sexes												
2002	208.0	*	*	*	0.3	2.6	17.9	62.8	171.2	450.8	1223.1	3,848.2
2001	212.6	*	*	*	0.3	2.9	17.3	62.6	177.8	467.8	1271.3	4,022.8
2000	220.2	*	*	*	0.3	2.4	16.9	64.6	189.5	496.4	1324.5	4,261.3
1999	227.2	*	*	*	0.3	2.5	17.3	65.8	196.4	527.7	1386.5	4,390.2
Non-Hispanic white, male												
2002	217.2	*	*	*	0.4	4.0	28.0	100.0	256.8	636.0	1618.3	4,324.5
2001	219.6	*	*	*	0.5	4.3	26.9	99.5	266.3	656.5	1678.7	4,506.9
2000	227.6	*	*	*	0.4	3.7	26.8	103.5	283.5	701.7	1752.3	4,833.4
1999	234.2	*	*	*	0.5	3.6	27.2	105.1	292.3	745.2	1834.9	4,954.1
Non-Hispanic white, female												
2002	199.1	*	*	*	0.2	1.3	7.8	26.1	90.2	293.0	958.3	3,642.5
2001	205.8	*	*	*	0.2	1.4	7.6	26.4	94.2	307.7	1001.1	3,821.4
2000	213.1	*	*	*	0.2	1.0	6.9	26.4	100.9	323.6	1043.3	4,030.7
1999	220.5	*	*	*	*	1.4	7.4	27.2	106.1	345.6	1092.9	4,165.3
Non-Hispanic black, both sexes												
2002	137.0	*	*	*	0.5	4.1	25.8	103.6	292.4	643.9	1433.9	3,481.1
2001	141.2	*	*	*	0.9	4.8	26.4	104.6	302.1	681.4	1487.0	3,628.7
2000	144.4	*	*	*	0.4	4.4	26.1	107.7	318.9	710.1	1534.5	3,701.7
1999	147.9	*	*	*	0.8	4.1	28.2	110.8	324.7	741.6	1559.0	3,817.9
Non-Hispanic black, male												
2002	139.5	*	*	*	*	5.3	36.2	146.9	413.2	850.8	1762.3	3,572.8
2001	143.2	*	*	*	1.2	6.8	37.1	149.0	418.0	903.1	1834.9	3,791.8
2000	145.7	*	*	*	*	6.1	36.3	152.2	444.2	916.5	1887.6	3,911.2
1999	150.0	*	*	*	0.9	6.0	40.3	160.0	458.4	967.5	1873.3	4,082.5
Non-Hispanic black, female												
2002	134.8	*	*	*	*	2.9	16.7	66.5	195.4	497.1	1248.1	3,446.0
2001	139.4	*	*	*	*	2.9	16.9	66.5	208.9	525.0	1291.6	3,568.3
2000	143.2	*	*	*	*	2.9	17.1	69.4	218.1	565.0	1335.7	3,623.9
1999	146.0	*	*	*	0.7	2.4	17.4	68.7	217.5	582.3	1383.0	3,719.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All origins 3/, both sexes												
2002	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.3	1,265.5
2000	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
All origins 3/, male												
2002	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.2	1,461.5
2000	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
All origins 3/, female												
2002	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5
Hispanic, both sexes												
2002	19.7	*	*	*	*	0.5	3.0	14.2	55.1	146.2	364.3	1,048.2
2001	20.1	*	*	*	*	0.7	2.6	14.8	49.6	149.9	395.7	1,188.0
2000	20.1	*	*	*	*	0.5	3.0	15.5	55.0	154.3	404.7	1,177.1
1999	21.1	*	*	*	*	0.5	2.9	17.8	56.1	164.0	433.7	1,227.8
Hispanic, male												
2002	20.6	*	*	*	*	0.7	4.2	21.0	80.8	199.3	460.0	1,107.7
2001	20.9	*	*	*	*	1.0	3.9	22.7	69.6	207.1	490.1	1,337.0
2000	21.1	*	*	*	*	0.7	4.7	23.4	80.0	207.6	514.8	1,273.8
1999	22.0	*	*	*	*	0.8	4.6	26.7	79.0	225.8	558.9	1,276.8
Hispanic, female												
2002	18.7	*	*	*	*	*	1.8	7.4	32.1	103.9	298.8	1,018.8
2001	19.1	*	*	*	*	*	1.1	7.0	31.8	104.9	331.8	1,115.2
2000	19.1	*	*	*	*	*	1.2	7.8	32.9	112.3	330.8	1,129.7
1999	20.1	*	*	*	*	*	1.1	9.1	35.9	115.5	350.4	1,204.1
Non-Hispanic 4/, both sexes												
2002	68.7	*	*	*	0.2	1.1	7.6	27.7	76.3	188.2	447.3	1,194.0
2001	71.4	*	*	*	0.2	1.3	7.6	28.8	80.3	195.6	473.6	1,265.3
2000	75.3	*	*	*	0.1	1.0	7.4	30.0	88.6	211.9	506.3	1,340.8
1999	78.2	*	*	*	0.2	1.2	7.7	30.5	92.2	227.9	531.9	1,381.0
Non-Hispanic 4/, male												
2002	73.5	*	*	*	0.2	1.6	11.5	42.6	111.3	259.2	573.4	1,374.2
2001	75.9	*	*	*	0.2	1.8	11.6	44.2	117.0	267.7	607.9	1,463.9
2000	80.3	*	*	*	0.2	1.5	11.4	46.5	128.7	290.9	652.6	1,576.2
1999	83.8	*	*	*	0.2	1.7	11.8	47.8	135.1	313.8	687.7	1,631.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Non-Hispanic 4/, female												
2002	64.1	*	*	*	*	0.6	3.8	13.3	44.0	128.9	363.8	1,116.3
2001	67.1	*	*	*	*	0.8	3.8	13.9	46.3	135.7	385.4	1,182.7
2000	70.6	*	*	*	0.1	0.6	3.5	14.1	51.6	146.8	411.0	1,245.6
1999	72.9	*	*	*	*	0.7	3.7	13.9	52.5	157.3	430.7	1,280.8
Non-Hispanic white, both sexes												
2002	75.5	*	*	*	0.1	1.1	7.5	27.0	74.7	186.1	445.9	1,202.8
2001	78.4	*	*	*	0.1	1.2	7.5	28.1	78.7	193.0	473.0	1,276.8
2000	82.7	*	*	*	0.1	0.9	7.3	29.3	86.9	209.3	505.3	1,355.3
1999	86.0	*	*	*	0.1	1.1	7.5	30.0	90.7	226.1	532.4	1,394.2
Non-Hispanic white, male												
2002	81.2	*	*	*	0.2	1.6	11.5	42.5	110.8	260.2	575.8	1,393.0
2001	83.7	*	*	*	0.2	1.7	11.6	44.3	116.8	267.7	612.5	1,483.2
2000	88.5	*	*	*	0.2	1.4	11.4	46.7	128.0	291.0	656.7	1,600.7
1999	92.4	*	*	*	0.2	1.5	11.8	48.1	134.6	315.0	695.7	1,653.8
Non-Hispanic white, female												
2002	70.1	*	*	*	*	0.5	3.6	11.8	40.6	122.9	358.8	1,120.7
2001	73.4	*	*	*	*	0.8	3.4	12.2	42.7	129.6	380.6	1,190.9
2000	77.2	*	*	*	*	0.5	3.1	12.3	48.2	140.5	405.9	1,256.3
1999	79.8	*	*	*	*	0.7	3.3	12.3	49.3	151.8	425.5	1,290.7
Non-Hispanic black, both sexes												
2002	50.2	*	*	*	*	1.7	10.0	39.0	109.1	242.6	523.0	1,212.9
2001	51.6	*	*	*	0.4	1.9	10.2	40.2	111.6	256.0	543.7	1,247.0
2000	54.0	*	*	*	*	1.8	10.1	41.9	122.3	273.0	576.6	1,281.8
1999	55.1	*	*	*	*	1.7	10.5	42.0	123.2	282.4	589.9	1,328.6
Non-Hispanic black, male												
2002	50.1	*	*	*	*	2.2	13.7	53.4	144.5	305.4	631.3	1,296.3
2001	50.9	*	*	*	*	2.5	13.5	53.4	146.1	320.8	646.6	1,389.4
2000	53.7	*	*	*	*	2.2	13.6	56.3	163.1	341.2	693.3	1,426.6
1999	55.2	*	*	*	*	2.5	14.5	58.1	168.7	357.0	691.4	1,509.4
Non-Hispanic black, female												
2002	50.4	*	*	*	*	1.3	6.7	26.7	80.7	198.0	461.8	1,181.0
2001	52.3	*	*	*	*	1.3	7.4	28.8	83.8	210.3	485.9	1,194.3
2000	54.2	*	*	*	*	1.4	7.0	29.6	89.5	225.0	510.9	1,228.0
1999	54.9	*	*	*	*	1.0	7.0	28.2	86.8	229.7	533.1	1,261.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
All origins 3/, male												
2002	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
All origins 3/, female												
2002	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	*	*	0.2	0.9	1.8	3.6	7.3	21.8
2001	1.3	*	*	*	*	*	0.2	0.8	1.9	3.6	7.1	22.0
2000	1.3	*	*	*	*	*	0.2	0.8	2.1	3.6	7.5	22.2
1999	1.3	*	*	*	*	*	0.2	0.7	2.0	3.8	7.2	22.2
Non-Hispanic 4/, male												
2002	1.4	*	*	*	*	*	0.3	1.4	2.6	4.9	9.0	22.1
2001	1.4	*	*	*	*	*	0.3	1.3	2.7	4.8	8.7	20.9
2000	1.4	*	*	*	*	*	0.3	1.3	3.1	4.9	9.3	21.0
1999	1.4	*	*	*	*	*	0.3	1.1	2.9	5.2	9.0	22.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Non-Hispanic 4/, female												
2002	1.3	*	*	*	*	*	0.1	0.5	1.1	2.5	6.1	21.6
2001	1.3	*	*	*	*	*	0.1	0.4	1.1	2.6	6.0	22.4
2000	1.2	*	*	*	*	*	0.1	0.3	1.2	2.5	6.3	22.7
1999	1.2	*	*	*	*	*	*	0.4	1.1	2.7	6.0	21.9
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	*	0.2	0.8	1.7	3.5	7.2	22.0
2001	1.4	*	*	*	*	*	0.2	0.8	1.7	3.4	7.0	22.0
2000	1.4	*	*	*	*	*	0.2	0.7	1.9	3.4	7.3	22.5
1999	1.4	*	*	*	*	*	0.2	0.6	1.8	3.6	7.0	22.7
Non-Hispanic white, male												
2002	1.5	*	*	*	*	*	0.3	1.2	2.5	4.6	9.0	22.7
2001	1.4	*	*	*	*	*	0.3	1.2	2.5	4.5	8.7	20.9
2000	1.5	*	*	*	*	*	0.3	1.2	2.8	4.6	9.0	21.3
1999	1.4	*	*	*	*	*	0.2	0.9	2.6	4.9	8.7	23.6
Non-Hispanic white, female												
2002	1.4	*	*	*	*	*	*	0.5	1.0	2.5	6.0	21.8
2001	1.4	*	*	*	*	*	*	0.3	1.0	2.5	5.9	22.4
2000	1.3	*	*	*	*	*	*	0.3	1.0	2.4	6.1	23.0
1999	1.3	*	*	*	*	*	*	0.3	1.1	2.6	5.8	22.4
Non-Hispanic black, both sexes												
2002	1.2	*	*	*	*	*	*	1.7	3.0	6.1	9.4	21.5
2001	1.3	*	*	*	*	*	0.5	1.6	4.0	5.8	8.5	24.3
2000	1.3	*	*	*	*	*	0.4	1.5	4.3	5.6	10.8	21.9
1999	1.2	*	*	*	*	*	0.5	1.9	3.6	5.8	10.9	17.5
Non-Hispanic black, male												
2002	1.4	*	*	*	*	*	*	2.8	4.4	9.4	11.7	21.9
2001	1.5	*	*	*	*	*	*	2.5	5.6	8.8	10.5	24.3
2000	1.5	*	*	*	*	*	*	2.3	6.3	8.7	14.2	*
1999	1.5	*	*	*	*	*	0.8	2.7	6.1	8.4	12.8	*
Non-Hispanic black, female												
2002	1.1	*	*	*	*	*	*	0.8	2.0	3.7	8.1	21.3
2001	1.1	*	*	*	*	*	*	*	2.6	3.8	7.4	24.4
2000	1.1	*	*	*	*	*	*	*	2.7	3.5	8.9	22.2
1999	1.0	*	*	*	*	*	*	1.1	1.6	3.9	9.8	16.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
All origins 3/, both sexes												
2002	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001	110.3	*	*	*	0.2	1.6	9.5	35.6	103.9	280.9	793.2	2,671.3
2000	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0
All origins 3/, male												
2002	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1021.0	2,817.4
2001	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1050.9	2,913.8
2000	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1079.4	3,112.7
1999	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1120.5	3,193.1
All origins 3/, female												
2002	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.8	2,569.9
2000	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7
Hispanic, both sexes												
2002	34.2	*	*	*	*	0.7	4.5	23.6	80.1	234.6	644.0	2,129.7
2001	35.7	*	*	*	*	0.9	4.2	23.8	80.1	249.7	702.6	2,427.9
2000	35.6	*	*	*	*	0.6	4.6	23.9	88.0	249.7	703.2	2,508.8
1999	37.5	*	*	*	*	0.9	4.5	26.8	93.5	268.8	741.8	2,640.7
Hispanic, male												
2002	35.7	*	*	*	*	1.2	7.2	36.6	122.3	329.1	808.8	2,262.2
2001	36.9	*	*	*	*	1.4	6.5	37.5	121.7	350.8	902.3	2,479.8
2000	36.9	*	*	*	*	0.9	7.4	39.1	134.2	351.6	900.8	2,577.7
1999	38.7	*	*	*	*	1.3	7.6	44.3	138.5	381.8	932.8	2,706.7
Hispanic, female												
2002	32.6	*	*	*	*	*	1.6	10.5	42.2	159.4	531.2	2,064.1
2001	34.5	*	*	*	*	*	1.7	10.3	43.3	169.9	567.6	2,402.5
2000	34.2	*	*	*	*	*	1.7	8.9	47.2	169.6	570.5	2,475.0
1999	36.3	*	*	*	*	*	1.3	9.7	53.8	180.2	614.8	2,608.9
Non-Hispanic 4/, both sexes												
2002	118.9	*	*	*	0.2	1.6	10.6	37.5	102.0	270.3	771.9	2,571.2
2001	120.9	*	*	*	0.2	1.7	10.2	36.5	105.1	281.6	794.2	2,672.2
2000	123.9	*	*	*	0.2	1.5	10.1	37.3	108.8	293.5	815.0	2,824.0
1999	127.6	*	*	*	0.2	1.5	10.4	38.1	112.1	308.6	849.1	2,916.3
Non-Hispanic 4/, male												
2002	122.7	*	*	*	0.3	2.4	16.6	59.2	152.6	380.2	1027.3	2,832.2
2001	123.1	*	*	*	0.3	2.6	15.7	57.3	156.4	395.1	1053.5	2,923.7
2000	126.0	*	*	*	0.2	2.3	15.7	58.9	162.3	414.7	1082.7	3,123.9
1999	128.9	*	*	*	0.3	2.3	16.2	59.9	165.4	435.2	1123.5	3,200.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic 4/, female												
2002	115.4	*	*	*	*	0.8	4.8	16.7	55.0	178.4	602.6	2,458.6
2001	118.8	*	*	*	0.1	0.8	4.8	16.5	57.5	187.3	624.1	2,567.6
2000	122.0	*	*	*	*	0.7	4.6	16.5	59.3	193.5	640.6	2,702.7
1999	126.4	*	*	*	0.1	0.8	4.7	17.3	62.9	204.6	671.1	2,802.7
Non-Hispanic white, both sexes												
2002	131.0	*	*	*	0.2	1.5	10.2	34.9	94.8	261.3	770.0	2,623.4
2001	132.7	*	*	*	0.2	1.6	9.6	33.8	97.4	271.4	791.2	2,724.1
2000	136.1	*	*	*	0.2	1.4	9.4	34.6	100.7	283.7	811.9	2,883.6
1999	139.9	*	*	*	0.2	1.4	9.6	35.2	103.9	297.9	847.2	2,973.2
Non-Hispanic white, male												
2002	134.5	*	*	*	0.2	2.3	16.2	56.4	143.5	371.2	1033.6	2,908.8
2001	134.4	*	*	*	0.3	2.5	15.0	54.1	147.0	384.3	1057.5	3,002.8
2000	137.7	*	*	*	0.2	2.2	15.1	55.7	152.7	406.1	1086.5	3,211.3
1999	140.4	*	*	*	0.2	2.1	15.2	56.1	155.1	425.3	1130.5	3,276.7
Non-Hispanic white, female												
2002	127.7	*	*	*	*	0.7	4.2	13.8	48.7	167.6	593.4	2,500.1
2001	131.1	*	*	*	*	0.7	4.1	13.9	50.5	175.6	614.6	2,608.1
2000	134.6	*	*	*	*	0.5	3.8	13.8	51.8	180.6	631.3	2,751.4
1999	139.3	*	*	*	*	0.7	4.0	14.6	55.7	191.3	661.7	2,852.2
Non-Hispanic black, both sexes												
2002	85.5	*	*	*	*	2.2	15.5	62.9	180.3	395.2	901.5	2,246.7
2001	88.2	*	*	*	0.5	2.9	15.6	62.9	186.6	419.6	934.8	2,357.3
2000	89.1	*	*	*	*	2.5	15.6	64.3	192.3	431.5	947.1	2,398.0
1999	91.6	*	*	*	0.4	2.4	17.2	66.9	197.9	453.5	958.2	2,471.9
Non-Hispanic black, male												
2002	88.0	*	*	*	*	3.0	22.2	90.8	264.4	536.0	1119.3	2,254.6
2001	90.8	*	*	*	*	4.3	22.9	93.1	266.3	573.5	1177.8	2,378.1
2000	90.4	*	*	*	*	3.7	22.1	93.6	274.7	566.6	1180.1	2,463.5
1999	93.3	*	*	*	*	3.5	25.1	99.2	283.6	602.1	1169.2	2,552.9
Non-Hispanic black, female												
2002	83.3	*	*	*	*	1.6	9.6	39.0	112.8	295.4	778.3	2,243.7
2001	86.0	*	*	*	*	1.6	9.2	36.9	122.4	310.9	798.4	2,349.6
2000	88.0	*	*	*	*	1.5	10.0	39.1	125.9	336.5	815.9	2,373.7
1999	90.1	*	*	*	*	1.3	10.2	39.4	129.2	348.6	840.0	2,441.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All origins 3/, both sexes												
2002	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
All origins 3/, male												
2002	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001	25.4	*	*	*	*	0.7	5.0	21.0	51.4	92.4	194.0	502.0
2000	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
All origins 3/, female												
2002	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.6	490.1
2000	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2
Hispanic, both sexes												
2002	8.2	*	*	*	*	0.4	1.9	8.9	29.7	61.1	124.1	358.6
2001	8.5	*	*	*	*	0.4	1.5	10.1	29.7	65.9	132.9	417.0
2000	8.4	*	*	*	*	*	1.6	9.5	30.5	65.7	128.2	452.5
1999	9.0	*	*	*	*	*	1.5	10.4	33.0	72.1	147.5	469.4
Hispanic, male												
2002	9.7	*	*	*	*	0.6	2.9	14.8	49.3	90.8	153.9	415.0
2001	10.1	*	*	*	*	0.6	2.2	17.1	47.9	100.4	181.5	440.0
2000	9.9	*	*	*	*	*	2.7	16.6	50.5	92.4	180.9	493.8
1999	10.6	*	*	*	*	*	2.5	17.8	50.0	115.8	194.0	485.3
Hispanic, female												
2002	6.5	*	*	*	*	*	*	3.1	12.0	37.6	103.7	330.7
2001	6.8	*	*	*	*	*	*	3.2	13.5	38.7	100.0	405.8
2000	6.8	*	*	*	*	*	*	2.6	12.8	44.6	92.8	432.3
1999	7.4	*	*	*	*	*	*	3.2	18.0	37.9	116.6	461.7
Non-Hispanic 4/, both sexes												
2002	25.9	*	*	*	*	0.5	3.6	13.7	32.9	64.3	146.4	477.8
2001	26.0	*	*	*	*	0.5	3.5	13.0	33.3	65.0	152.6	494.8
2000	26.9	*	*	*	*	0.4	3.2	13.4	33.8	67.9	158.3	535.9
1999	27.9	*	*	*	*	0.4	3.5	13.1	35.0	72.0	165.7	564.9
Non-Hispanic 4/, male												
2002	27.8	*	*	*	*	0.7	5.8	22.0	50.7	91.0	185.6	493.3
2001	27.5	*	*	*	*	0.7	5.4	21.2	51.1	91.0	193.2	503.0
2000	28.2	*	*	*	*	0.7	5.1	21.7	52.1	96.4	198.5	551.0
1999	28.9	*	*	*	*	0.6	5.6	21.0	53.3	101.1	205.6	580.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic 4/, female												
2002	24.0	*	*	*	*	0.2	1.4	5.7	16.5	42.0	120.5	471.1
2001	24.6	*	*	*	*	0.3	1.5	5.2	16.8	43.3	126.0	491.4
2000	25.6	*	*	*	*	0.2	1.3	5.4	16.9	44.4	132.2	529.8
1999	27.0	*	*	*	*	0.2	1.4	5.4	18.2	48.2	139.8	558.8
Non-Hispanic white, both sexes												
2002	26.8	*	*	*	*	0.4	3.3	12.4	28.9	57.2	138.2	473.5
2001	26.8	*	*	*	*	0.4	3.2	11.6	28.4	57.1	144.2	490.7
2000	27.6	*	*	*	*	0.4	2.9	11.9	28.9	60.2	149.0	530.2
1999	28.6	*	*	*	*	0.4	3.1	11.3	29.8	63.6	156.2	558.8
Non-Hispanic white, male												
2002	28.2	*	*	*	*	0.7	5.4	20.3	44.8	81.3	174.4	489.2
2001	27.8	*	*	*	*	0.6	5.0	19.1	44.3	80.4	182.9	499.4
2000	28.4	*	*	*	*	0.6	4.8	19.6	45.2	86.3	186.1	544.7
1999	29.0	*	*	*	*	0.6	5.2	18.4	46.0	90.0	193.5	572.8
Non-Hispanic white, female												
2002	25.3	*	*	*	*	0.2	1.1	4.6	13.8	36.6	113.9	466.8
2001	25.8	*	*	*	*	0.2	1.3	4.1	13.3	37.3	118.6	487.0
2000	26.9	*	*	*	*	*	1.1	4.3	13.4	38.2	124.6	524.4
1999	28.2	*	*	*	*	0.2	1.1	4.2	14.6	41.6	131.8	553.2
Non-Hispanic black, both sexes												
2002	27.7	*	*	*	*	0.8	6.0	25.7	71.9	137.2	261.1	580.4
2001	28.5	*	*	*	*	0.7	5.8	25.7	78.1	144.8	266.7	595.9
2000	29.3	*	*	*	*	0.7	5.3	27.3	78.9	147.2	280.2	652.2
1999	30.7	*	*	*	*	0.8	6.5	28.1	82.0	158.7	291.0	670.3
Non-Hispanic black, male												
2002	31.9	*	*	*	*	1.1	8.9	38.8	111.8	198.7	358.4	603.9
2001	32.3	*	*	*	*	1.0	8.8	39.9	118.1	209.4	352.3	609.8
2000	33.2	*	*	*	*	1.1	7.9	42.3	120.8	209.3	371.5	711.6
1999	34.5	*	*	*	*	1.1	9.8	43.6	123.7	228.2	378.4	714.3
Non-Hispanic black, female												
2002	24.0	*	*	*	*	*	3.4	14.4	39.8	93.6	206.1	571.5
2001	25.0	*	*	*	*	*	3.2	13.6	45.9	99.3	218.7	590.7
2000	25.8	*	*	*	*	*	2.9	14.4	45.1	103.4	228.9	630.2
1999	27.1	*	*	*	*	*	3.6	14.7	48.5	109.7	242.1	653.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All origins 3/, both sexes												
2002	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.8
2000	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
All origins 3/, male												
2002	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	857.0	2,411.8
2000	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
All origins 3/, female												
2002	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4
Hispanic, both sexes												
2002	26.0	*	*	*	*	0.4	2.7	14.6	50.4	173.4	519.9	1,771.1
2001	27.2	*	*	*	*	0.6	2.7	13.7	50.5	183.7	569.7	2,010.8
2000	27.2	*	*	*	*	0.5	3.0	14.3	57.5	184.0	575.0	2,056.3
1999	28.5	*	*	*	*	0.6	3.0	16.4	60.5	196.7	594.3	2,171.4
Hispanic, male												
2002	25.9	*	*	*	*	0.6	4.3	21.8	73.0	238.3	654.9	1,847.2
2001	26.7	*	*	*	*	0.9	4.3	20.4	73.8	250.4	720.8	2,039.8
2000	27.0	*	*	*	*	0.7	4.7	22.6	83.6	259.1	719.8	2,084.0
1999	28.1	*	*	*	*	1.0	5.1	26.5	88.5	266.0	738.8	2,221.4
Hispanic, female												
2002	26.1	*	*	*	*	*	0.9	7.4	30.2	121.8	427.5	1,733.3
2001	27.7	*	*	*	*	*	1.0	7.1	29.8	131.2	467.5	1,996.7
2000	27.5	*	*	*	*	*	1.2	6.2	34.4	124.9	477.7	2,042.7
1999	28.9	*	*	*	*	*	0.8	6.5	35.8	142.3	498.2	2,147.2
Non-Hispanic 4/, both sexes												
2002	93.1	*	*	*	0.1	1.1	7.1	23.8	69.0	206.0	625.4	2,093.4
2001	94.9	*	*	*	0.2	1.2	6.7	23.5	71.8	216.7	641.6	2,177.4
2000	97.0	*	*	*	0.1	1.1	6.9	23.9	75.0	225.6	656.7	2,288.1
1999	99.7	*	*	*	0.2	1.1	6.9	25.1	77.1	236.5	683.4	2,351.4
Non-Hispanic 4/, male												
2002	94.9	*	*	*	0.2	1.6	10.8	37.2	102.0	289.3	841.7	2,338.9
2001	95.6	*	*	*	0.2	2.0	10.3	36.1	105.3	304.1	860.3	2,420.7
2000	97.8	*	*	*	0.2	1.7	10.5	37.2	110.2	318.2	884.2	2,572.9
1999	100.0	*	*	*	0.2	1.6	10.6	38.9	112.1	334.1	917.9	2,619.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Non-Hispanic 4/, female												
2002	91.4	*	*	*	*	0.6	3.4	11.0	38.5	136.4	482.1	1,987.5
2001	94.2	*	*	*	*	0.5	3.2	11.3	40.7	143.9	498.0	2,076.2
2000	96.3	*	*	*	*	0.5	3.3	11.1	42.4	149.1	508.5	2,172.9
1999	99.4	*	*	*	*	0.6	3.3	11.8	44.8	156.4	531.2	2,243.9
Non-Hispanic white, both sexes												
2002	104.3	*	*	*	0.1	1.1	6.9	22.5	65.9	204.1	631.9	2,149.8
2001	106.0	*	*	*	0.1	1.1	6.4	22.2	69.0	214.3	647.0	2,233.4
2000	108.4	*	*	*	0.1	1.0	6.5	22.7	71.9	223.5	662.9	2,353.3
1999	111.3	*	*	*	0.1	1.0	6.5	23.9	74.1	234.2	690.9	2,414.4
Non-Hispanic white, male												
2002	106.2	*	*	*	0.2	1.6	10.7	36.1	98.7	289.9	859.2	2,419.7
2001	106.6	*	*	*	0.2	1.8	10.0	34.9	102.7	303.9	874.6	2,503.4
2000	109.2	*	*	*	0.2	1.6	10.3	36.1	107.5	319.8	900.4	2,666.6
1999	111.4	*	*	*	0.2	1.5	10.1	37.7	109.1	335.3	936.9	2,703.9
Non-Hispanic white, female												
2002	102.4	*	*	*	*	0.5	3.1	9.2	34.9	131.0	479.6	2,033.3
2001	105.3	*	*	*	*	0.5	2.9	9.8	37.2	138.2	496.0	2,121.0
2000	107.7	*	*	*	*	0.4	2.7	9.5	38.3	142.4	506.8	2,227.0
1999	111.1	*	*	*	*	0.5	2.8	10.4	41.1	149.7	529.9	2,299.0
Non-Hispanic black, both sexes												
2002	57.8	*	*	*	*	1.5	9.6	37.2	108.4	258.0	640.3	1,666.3
2001	59.7	*	*	*	0.4	2.1	9.8	37.1	108.4	274.7	668.1	1,761.4
2000	59.8	*	*	*	*	1.8	10.4	37.0	113.4	284.4	666.8	1,745.8
1999	61.0	*	*	*	*	1.6	10.7	38.9	115.9	294.7	667.1	1,801.6
Non-Hispanic black, male												
2002	56.1	*	*	*	*	1.9	13.3	52.0	152.5	337.3	760.8	1,650.8
2001	58.4	*	*	*	*	3.3	14.1	53.2	148.2	364.1	825.5	1,768.3
2000	57.3	*	*	*	*	2.6	14.1	51.3	153.9	357.2	808.7	1,751.9
1999	58.8	*	*	*	*	2.4	15.3	55.5	160.0	374.0	790.8	1,838.6
Non-Hispanic black, female												
2002	59.3	*	*	*	*	1.1	6.2	24.5	73.0	201.8	572.2	1,672.2
2001	60.9	*	*	*	*	1.0	6.0	23.3	76.5	211.7	579.7	1,758.9
2000	62.1	*	*	*	*	1.1	7.1	24.7	80.7	233.1	587.0	1,743.5
1999	63.0	*	*	*	*	0.8	6.6	24.6	80.7	238.9	597.9	1,787.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All origins 3/, both sexes												
2002	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.6
2000	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9
All origins 3/, male												
2002	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9
All origins 3/, female												
2002	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.5
2000	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5
Hispanic, both sexes												
2002	13.3	9.7	0.7	0.4	1.5	2.6	5.1	11.4	25.7	72.6	219.8	740.0
2001	13.3	9.2	1.0	0.5	1.3	2.6	4.9	10.9	28.0	75.5	217.8	799.7
2000	13.6	10.2	1.0	0.5	1.7	2.8	5.5	11.6	28.8	75.9	227.0	829.4
1999	13.8	12.2	1.2	0.5	1.6	2.8	6.3	11.6	30.6	78.7	229.6	827.9
Hispanic, male												
2002	12.9	9.1	*	*	2.1	3.4	6.4	14.3	31.7	92.5	256.9	758.0
2001	12.6	9.6	*	*	1.6	3.2	6.5	15.0	34.1	91.1	249.1	775.2
2000	13.2	9.9	1.3	0.7	2.1	3.5	6.7	15.1	37.2	94.1	250.8	890.8
1999	13.7	10.3	1.4	0.6	1.8	3.8	8.2	15.4	38.8	102.4	267.5	888.6
Hispanic, female												
2002	13.7	10.3	*	*	0.8	1.6	3.7	8.5	20.3	56.7	194.5	731.1
2001	14.0	8.7	*	*	0.9	1.9	3.2	6.9	22.7	63.3	196.7	811.7
2000	14.0	10.6	*	*	1.1	1.9	4.3	8.1	21.4	61.6	211.0	799.3
1999	13.9	14.1	*	*	1.3	1.6	4.2	8.0	23.4	60.1	204.3	798.6
Non-Hispanic 4/, both sexes												
2002	65.7	12.2	1.1	0.6	2.2	4.9	11.0	22.7	50.4	133.2	398.4	1,469.2
2001	65.2	11.6	1.5	0.6	2.1	4.7	10.6	22.6	51.1	134.7	396.5	1,497.8
2000	65.1	12.4	1.1	0.6	2.3	4.7	10.4	22.3	52.4	137.5	397.5	1,530.9
1999	65.6	13.1	1.1	0.6	2.4	4.7	10.8	22.4	54.3	142.0	404.4	1,533.9
Non-Hispanic 4/, male												
2002	58.7	12.8	1.1	0.7	2.8	6.2	13.7	29.5	62.9	160.4	456.6	1,477.3
2001	57.8	11.0	1.6	0.6	2.6	5.6	13.2	28.7	63.6	162.5	460.6	1,499.0
2000	58.0	12.7	1.2	0.7	2.7	5.5	12.9	28.9	66.4	167.8	458.9	1,573.6
1999	58.6	13.8	1.3	0.7	2.9	5.5	13.5	28.7	69.3	176.2	466.4	1,571.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Non-Hispanic 4/, female												
2002	72.3	11.6	1.0	0.5	1.5	3.7	8.4	16.1	38.7	110.4	359.7	1,465.7
2001	72.3	12.2	1.3	0.6	1.6	3.9	8.0	16.7	39.4	111.6	354.4	1,497.3
2000	71.8	12.0	1.0	0.5	1.9	3.8	7.9	15.9	39.4	112.5	357.5	1,513.7
1999	72.2	12.3	1.0	0.6	1.9	3.9	8.1	16.2	40.5	113.9	364.2	1,519.0
Non-Hispanic white, both sexes												
2002	71.3	10.0	0.9	0.5	1.8	3.8	9.3	19.0	45.3	127.6	398.1	1,503.1
2001	70.6	9.1	1.1	0.5	1.7	3.7	8.5	19.0	45.8	128.7	394.8	1,533.2
2000	70.3	10.5	0.9	0.5	1.7	3.7	8.4	18.7	47.4	131.2	395.8	1,564.2
1999	70.4	10.5	0.9	0.5	2.0	3.7	8.7	18.7	48.3	135.2	401.1	1,565.9
Non-Hispanic white, male												
2002	62.7	10.7	1.1	0.6	2.3	4.7	11.8	25.2	57.1	154.1	457.1	1,512.9
2001	61.4	9.4	1.2	0.6	2.0	4.4	10.9	24.8	57.5	155.8	457.4	1,530.7
2000	61.6	10.9	1.1	0.6	2.0	4.4	10.7	24.6	60.7	161.0	457.6	1,606.4
1999	61.7	11.6	1.1	0.5	2.3	4.4	11.2	24.5	62.1	168.5	462.9	1,595.5
Non-Hispanic white, female												
2002	79.6	9.3	0.8	0.5	1.2	2.8	6.8	13.0	34.1	105.0	358.5	1,498.8
2001	79.4	8.8	1.0	0.5	1.3	3.1	6.1	13.4	34.9	105.7	353.3	1,534.2
2000	78.7	10.1	0.8	0.4	1.4	2.9	6.2	12.9	34.8	106.0	355.3	1,547.2
1999	78.8	9.4	0.8	0.5	1.7	3.0	6.2	13.1	35.3	107.3	360.6	1,554.1
Non-Hispanic black, both sexes												
2002	54.3	20.9	1.6	0.9	4.2	11.8	23.4	52.6	104.1	211.2	474.7	1,285.5
2001	54.7	21.3	3.0	1.0	4.4	11.0	24.5	52.2	105.7	216.1	486.4	1,292.9
2000	54.6	19.6	2.0	1.1	4.9	10.8	23.7	52.8	104.5	223.6	484.3	1,319.4
1999	56.2	23.2	2.0	1.0	4.7	11.0	25.3	53.5	115.3	232.9	499.9	1,314.0
Non-Hispanic black, male												
2002	52.6	21.8	*	0.9	5.1	14.8	28.2	67.5	130.4	259.4	543.4	1,258.0
2001	53.5	18.8	3.3	0.9	5.5	13.4	29.3	63.9	134.9	261.9	593.5	1,362.9
2000	53.3	19.1	2.0	1.2	5.7	12.7	28.7	68.3	132.9	269.8	565.5	1,416.1
1999	56.2	22.9	2.3	1.2	6.0	13.4	30.9	68.7	149.5	291.0	584.7	1,466.6
Non-Hispanic black, female												
2002	55.9	19.9	1.9	0.8	3.2	9.0	19.2	39.9	83.0	177.0	435.9	1,296.0
2001	55.8	23.8	2.7	1.0	3.2	8.9	20.3	42.2	82.3	183.7	426.2	1,267.0
2000	55.8	20.1	1.9	1.0	4.2	9.0	19.2	39.6	81.6	191.1	438.5	1,283.4
1999	56.3	23.6	1.8	0.9	3.4	8.9	20.4	40.5	87.9	192.0	452.4	1,257.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
All origins 3/, both sexes												
2002	4.8	4.6	0.2	0.1	0.3	0.8	1.7	3.4	6.9	14.9	29.8	56.3
2001	4.7	4.2	0.1	0.1	0.3	0.9	1.7	3.4	6.9	14.6	28.5	57.5
2000	4.6	4.9	0.2	0.1	0.3	0.9	1.7	3.5	6.7	14.5	28.3	57.4
1999	4.7	5.7	*	0.1	0.4	0.9	1.8	3.3	6.7	14.8	30.0	59.6
All origins 3/, male												
2002	3.8	4.7	*	*	0.2	0.7	1.4	3.1	6.5	14.1	27.7	52.1
2001	3.8	3.9	*	*	0.2	0.6	1.5	3.0	6.5	14.5	27.9	55.6
2000	3.8	5.1	*	*	0.2	0.7	1.4	3.3	6.4	13.9	28.3	54.8
1999	3.9	5.9	*	*	0.3	0.6	1.5	3.0	6.8	15.1	29.1	60.3
All origins 3/, female												
2002	5.7	4.5	*	*	0.4	1.0	1.9	3.6	7.2	15.6	31.2	58.1
2001	5.5	4.6	*	*	0.4	1.1	2.0	3.7	7.3	14.6	28.9	58.2
2000	5.5	4.6	*	*	0.5	1.0	2.1	3.7	7.0	15.0	28.4	58.5
1999	5.5	5.6	*	*	0.5	1.1	2.0	3.6	6.6	14.6	30.5	59.2
Hispanic, both sexes												
2002	1.3	3.1	*	*	*	0.4	0.9	1.8	3.1	7.5	16.8	30.5
2001	1.3	3.7	*	*	*	0.5	0.9	1.4	3.5	6.9	14.6	32.5
2000	1.2	4.3	*	*	*	0.5	1.1	2.0	3.0	6.2	14.4	27.9
1999	1.4	5.4	*	*	0.3	0.5	1.0	1.6	3.4	7.1	16.2	37.2
Hispanic, male												
2002	0.9	*	*	*	*	*	0.7	1.6	2.8	8.6	11.9	*
2001	1.1	*	*	*	*	*	0.9	1.4	2.9	7.2	15.2	43.8
2000	0.9	*	*	*	*	*	0.8	2.0	2.5	5.7	12.3	*
1999	1.1	5.2	*	*	*	*	0.8	*	3.3	6.8	15.1	*
Hispanic, female												
2002	1.6	*	*	*	*	*	1.3	2.0	3.5	6.5	20.2	35.6
2001	1.5	*	*	*	*	0.8	1.0	1.5	4.1	6.6	14.2	27.0
2000	1.5	*	*	*	*	0.7	1.4	2.0	3.4	6.6	15.8	32.6
1999	1.7	5.7	*	*	*	0.7	1.3	2.0	3.5	7.4	17.0	35.4
Non-Hispanic 4/, both sexes												
2002	5.3	5.0	0.2	0.1	0.3	0.9	1.8	3.5	7.1	15.4	30.4	57.3
2001	5.2	4.3	*	0.1	0.3	1.0	1.8	3.5	7.1	15.0	29.1	58.3
2000	5.1	5.0	0.2	0.1	0.4	0.9	1.8	3.6	6.9	14.9	28.9	58.4
1999	5.2	5.8	*	0.1	0.4	0.9	1.9	3.5	6.9	15.2	30.4	60.2
Non-Hispanic 4/, male												
2002	4.2	5.4	*	*	0.2	0.8	1.6	3.3	6.7	14.5	28.4	53.5
2001	4.2	4.0	*	*	0.3	0.7	1.6	3.1	6.7	14.9	28.3	55.8
2000	4.2	5.3	*	*	0.2	0.8	1.5	3.4	6.7	14.3	28.9	56.3
1999	4.3	6.0	*	*	0.3	0.7	1.6	3.2	7.0	15.5	29.6	60.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Non-Hispanic 4/, female												
2002	6.3	4.5	*	*	0.4	1.1	2.0	3.7	7.5	16.2	31.7	58.9
2001	6.1	4.7	*	*	0.4	1.2	2.1	3.9	7.6	15.2	29.6	59.3
2000	6.0	4.7	*	*	0.5	1.1	2.1	3.8	7.2	15.4	28.9	59.2
1999	6.0	5.5	*	*	0.6	1.1	2.1	3.7	6.8	15.0	31.0	59.9
Non-Hispanic white, both sexes												
2002	5.4	4.1	*	0.1	0.2	0.7	1.6	3.1	6.6	14.7	29.7	57.1
2001	5.3	3.5	*	*	0.3	0.8	1.5	3.2	6.6	14.3	28.4	57.6
2000	5.2	4.0	*	*	0.3	0.7	1.5	3.2	6.5	14.3	28.2	57.4
1999	5.3	4.9	*	0.1	0.4	0.7	1.5	3.0	6.1	14.1	29.4	59.9
Non-Hispanic white, male												
2002	4.4	4.6	*	*	0.2	0.6	1.4	3.1	6.4	14.0	27.9	53.4
2001	4.3	3.3	*	*	0.2	0.6	1.4	2.9	6.1	14.4	27.5	53.8
2000	4.3	4.4	*	*	0.2	0.7	1.3	3.1	6.3	14.1	28.4	54.4
1999	4.4	5.5	*	*	0.2	0.5	1.4	2.9	6.4	14.5	28.5	60.9
Non-Hispanic white, female												
2002	6.4	3.6	*	*	0.3	0.7	1.7	3.2	6.8	15.3	30.9	58.7
2001	6.3	3.8	*	*	0.3	0.9	1.7	3.4	7.0	14.3	29.0	59.2
2000	6.1	3.6	*	*	0.4	0.8	1.7	3.3	6.7	14.5	28.2	58.6
1999	6.1	4.4	*	*	0.5	0.8	1.6	3.2	5.9	13.8	29.9	59.5
Non-Hispanic black, both sexes												
2002	6.1	8.5	*	*	0.6	2.5	3.6	7.0	14.0	26.4	46.5	69.6
2001	6.0	7.0	*	*	0.6	2.2	4.0	7.0	13.8	26.1	42.6	78.0
2000	5.9	8.1	*	*	0.8	2.2	4.1	7.6	12.2	25.1	42.9	77.6
1999	6.4	10.0	*	*	0.7	2.3	4.1	7.4	15.0	29.2	49.7	72.1
Non-Hispanic black, male												
2002	4.7	9.4	*	*	*	2.1	3.0	5.7	12.0	23.4	43.1	66.7
2001	4.9	7.3	*	*	*	1.6	3.4	5.5	13.8	24.6	44.7	94.7
2000	4.8	7.8	*	*	*	1.7	3.3	6.6	12.2	21.6	43.8	93.9
1999	5.3	9.4	*	*	*	2.0	3.3	6.3	14.4	30.5	50.7	71.3
Non-Hispanic black, female												
2002	7.4	7.6	*	*	0.8	2.8	4.2	8.0	15.6	28.6	48.4	70.7
2001	7.0	6.7	*	*	*	2.8	4.5	8.2	13.8	27.2	41.4	71.9
2000	7.0	8.5	*	*	1.2	2.5	4.8	8.3	12.3	27.6	42.4	71.5
1999	7.4	10.7	*	*	0.8	2.7	4.9	8.2	15.5	28.3	49.2	72.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary embolism (I26)												
All origins 3/, both sexes												
2002	3.0	*	*	*	0.2	0.5	1.2	2.2	4.5	9.5	18.5	35.4
2001	3.0	*	*	*	0.2	0.6	1.2	2.3	4.7	9.5	18.4	37.8
2000	3.0	*	*	*	0.2	0.5	1.2	2.3	4.5	9.3	18.4	38.4
1999	3.2	*	*	*	0.2	0.6	1.2	2.3	4.7	10.1	21.1	42.0
All origins 3/, male												
2002	2.5	*	*	*	0.1	0.5	1.0	2.2	4.5	9.4	17.5	34.1
2001	2.6	*	*	*	0.1	0.5	1.1	2.1	4.5	10.0	19.0	38.0
2000	2.6	*	*	*	0.1	0.5	1.0	2.4	4.6	9.5	19.3	37.5
1999	2.7	*	*	*	0.2	0.5	1.1	2.2	5.0	10.9	20.4	42.0
All origins 3/, female												
2002	3.5	*	*	*	0.2	0.6	1.3	2.2	4.5	9.5	19.2	36.0
2001	3.5	*	*	*	0.2	0.7	1.3	2.5	4.9	9.1	18.0	37.7
2000	3.4	*	*	*	0.3	0.6	1.3	2.2	4.4	9.1	17.8	38.8
1999	3.7	*	*	*	0.3	0.7	1.3	2.4	4.4	9.6	21.5	42.1
Hispanic, both sexes												
2002	0.7	*	*	*	*	*	0.4	0.8	1.7	4.6	9.8	21.1
2001	0.7	*	*	*	*	*	0.5	0.6	2.2	4.4	9.2	22.5
2000	0.6	*	*	*	*	*	*	1.0	1.7	3.7	7.7	14.6
1999	0.7	*	*	*	*	*	*	1.0	2.3	5.0	10.2	28.1
Hispanic, male												
2002	0.5	*	*	*	*	*	*	*	*	5.9	*	*
2001	0.6	*	*	*	*	*	*	*	*	5.4	10.7	*
2000	0.5	*	*	*	*	*	*	*	*	4.2	*	*
1999	0.6	*	*	*	*	*	*	*	*	4.8	*	*
Hispanic, female												
2002	0.8	*	*	*	*	*	*	*	*	3.5	12.1	23.2
2001	0.7	*	*	*	*	*	*	*	2.5	3.6	8.2	19.5
2000	0.6	*	*	*	*	*	*	*	*	3.3	7.9	*
1999	0.9	*	*	*	*	*	*	*	2.3	5.2	11.1	26.0
Non-Hispanic 4/, both sexes												
2002	3.3	*	*	*	0.2	0.6	1.3	2.4	4.7	9.8	18.9	36.0
2001	3.4	*	*	*	0.2	0.7	1.2	2.4	4.9	9.8	18.8	38.3
2000	3.3	*	*	*	0.2	0.6	1.3	2.5	4.7	9.6	18.8	39.2
1999	3.6	*	*	*	0.3	0.6	1.3	2.4	4.9	10.4	21.4	42.4
Non-Hispanic 4/, male												
2002	2.8	*	*	*	0.1	0.5	1.1	2.3	4.8	9.6	18.0	34.9
2001	2.8	*	*	*	0.1	0.5	1.1	2.2	4.7	10.2	19.3	38.2
2000	2.9	*	*	*	0.1	0.6	1.1	2.6	4.8	9.8	19.8	38.6
1999	3.0	*	*	*	0.2	0.5	1.1	2.3	5.2	11.2	20.8	42.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pulmonary embolism (I26)												
Non-Hispanic 4/, female												
2002	3.9	*	*	*	0.3	0.7	1.4	2.4	4.7	9.9	19.5	36.4
2001	3.9	*	*	*	0.2	0.8	1.3	2.6	5.0	9.5	18.4	38.4
2000	3.8	*	*	*	0.3	0.7	1.4	2.3	4.6	9.5	18.2	39.5
1999	4.1	*	*	*	0.4	0.7	1.4	2.5	4.5	9.8	21.8	42.5
Non-Hispanic white, both sexes												
2002	3.4	*	*	*	0.2	0.5	1.2	2.2	4.4	9.3	18.3	35.3
2001	3.5	*	*	*	0.2	0.5	1.0	2.2	4.6	9.4	18.3	37.5
2000	3.4	*	*	*	0.2	0.5	1.0	2.2	4.4	9.2	18.2	38.2
1999	3.6	*	*	*	0.2	0.5	1.1	2.1	4.3	9.7	20.6	42.0
Non-Hispanic white, male												
2002	2.9	*	*	*	*	0.4	1.1	2.2	4.5	9.3	17.5	34.1
2001	2.9	*	*	*	*	0.4	1.0	2.1	4.3	9.9	18.6	36.3
2000	2.9	*	*	*	*	0.5	0.9	2.3	4.5	9.6	19.1	36.9
1999	3.0	*	*	*	*	0.4	1.1	2.1	4.7	10.4	19.7	41.6
Non-Hispanic white, female												
2002	4.0	*	*	*	0.2	0.5	1.2	2.1	4.3	9.3	18.8	35.9
2001	4.0	*	*	*	0.2	0.7	1.1	2.3	4.8	9.0	18.0	38.0
2000	3.9	*	*	*	0.2	0.6	1.2	2.0	4.4	8.8	17.6	38.8
1999	4.1	*	*	*	0.3	0.5	1.1	2.2	4.0	9.0	21.2	42.2
Non-Hispanic black, both sexes												
2002	3.9	*	*	*	0.4	1.6	2.3	4.6	9.0	17.0	30.8	49.9
2001	4.0	*	*	*	*	1.7	2.8	4.8	8.9	16.7	28.9	55.9
2000	4.0	*	*	*	0.6	1.3	2.9	5.3	8.2	16.8	30.9	57.2
1999	4.4	*	*	*	0.5	1.7	2.6	5.2	10.5	20.9	36.6	53.3
Non-Hispanic black, male												
2002	3.2	*	*	*	*	1.6	1.9	4.4	8.8	15.5	29.0	52.5
2001	3.4	*	*	*	*	1.4	2.6	3.8	10.1	16.2	33.3	71.6
2000	3.4	*	*	*	*	1.0	2.6	5.4	9.3	14.8	33.9	72.8
1999	3.9	*	*	*	*	1.6	2.1	4.8	11.0	23.6	41.0	60.6
Non-Hispanic black, female												
2002	4.6	*	*	*	*	1.6	2.7	4.8	9.1	18.0	31.8	48.9
2001	4.4	*	*	*	*	1.9	3.1	5.6	7.9	17.1	26.4	50.0
2000	4.5	*	*	*	0.9	1.6	3.1	5.1	7.4	18.1	29.2	51.5
1999	4.9	*	*	*	*	1.9	3.2	5.5	10.0	19.0	34.1	50.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
All origins 3/, both sexes												
2002	1.8	4.6	0.2	0.1	0.1	0.3	0.5	1.1	2.4	5.5	11.3	20.9
2001	1.6	4.2	0.1	0.1	0.1	0.3	0.6	1.1	2.2	5.1	10.1	19.6
2000	1.6	4.7	0.2	0.1	0.1	0.3	0.6	1.1	2.2	5.2	9.9	19.0
1999	1.5	5.7	*	0.1	0.2	0.3	0.6	1.0	2.0	4.6	8.9	17.5
All origins 3/, male												
2002	1.3	4.7	*	*	*	0.2	0.4	0.9	1.9	4.7	10.3	18.0
2001	1.2	3.9	*	*	0.1	0.2	0.4	0.9	2.0	4.5	8.9	17.6
2000	1.2	4.9	*	*	*	0.2	0.5	0.9	1.8	4.4	9.0	17.3
1999	1.2	5.9	*	*	0.1	0.2	0.5	0.8	1.7	4.2	8.7	18.4
All origins 3/, female												
2002	2.2	4.4	*	*	0.1	0.3	0.6	1.3	2.8	6.1	12.0	22.1
2001	2.0	4.5	*	*	0.1	0.4	0.7	1.3	2.5	5.5	11.0	20.5
2000	2.1	4.5	*	*	0.2	0.4	0.7	1.4	2.5	5.8	10.6	19.7
1999	1.8	5.5	*	*	0.2	0.4	0.7	1.2	2.2	5.0	9.0	17.2
Hispanic, both sexes												
2002	0.6	3.0	*	*	*	*	0.5	0.9	1.4	2.9	7.0	*
2001	0.6	3.7	*	*	*	0.4	0.4	0.8	1.4	2.5	5.4	*
2000	0.7	3.9	*	*	*	0.3	0.7	1.0	1.3	2.5	6.7	13.3
1999	0.6	5.3	*	*	*	*	0.7	*	*	2.1	6.0	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	5.2	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.8	*	*	*	*	*	*	1.3	*	3.0	8.2	*
2001	0.8	*	*	*	*	*	*	*	*	*	6.0	*
2000	0.9	*	*	*	*	*	1.0	*	*	3.3	7.9	*
1999	0.8	5.4	*	*	*	*	0.9	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.9	4.9	0.2	0.1	0.1	0.3	0.5	1.1	2.4	5.6	11.5	21.3
2001	1.8	4.3	*	0.1	0.1	0.3	0.6	1.1	2.3	5.2	10.3	20.0
2000	1.8	4.9	0.2	*	0.1	0.3	0.6	1.2	2.2	5.3	10.1	19.1
1999	1.6	5.8	*	0.1	0.2	0.3	0.6	1.1	2.1	4.8	9.0	17.8
Non-Hispanic 4/, male												
2002	1.5	5.3	*	*	*	0.2	0.4	0.9	2.0	4.8	10.5	18.6
2001	1.4	4.0	*	*	0.1	0.2	0.4	0.9	2.0	4.7	9.0	17.6
2000	1.3	5.2	*	*	*	0.2	0.5	0.9	1.9	4.5	9.1	17.7
1999	1.3	6.0	*	*	*	0.2	0.5	0.9	1.8	4.3	8.8	18.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Non-Hispanic 4/, female												
2002	2.4	4.5	*	*	0.1	0.4	0.6	1.3	2.8	6.3	12.2	22.5
2001	2.2	4.5	*	*	0.1	0.4	0.7	1.3	2.5	5.6	11.2	21.0
2000	2.2	4.6	*	*	0.2	0.4	0.7	1.4	2.6	6.0	10.7	19.7
1999	2.0	5.5	*	*	0.2	0.4	0.7	1.2	2.3	5.2	9.2	17.4
Non-Hispanic white, both sexes												
2002	2.0	4.1	*	*	*	0.2	0.4	1.0	2.2	5.4	11.4	21.8
2001	1.8	3.5	*	*	0.1	0.2	0.5	1.0	2.0	4.9	10.2	20.1
2000	1.8	4.0	*	*	0.1	0.2	0.5	1.0	2.1	5.1	10.1	19.1
1999	1.6	4.9	*	0.1	0.2	0.2	0.5	0.9	1.8	4.5	8.8	17.8
Non-Hispanic white, male												
2002	1.5	4.5	*	*	*	0.2	0.3	0.9	1.9	4.7	10.4	19.3
2001	1.4	3.3	*	*	*	0.2	0.4	0.8	1.9	4.4	8.9	17.5
2000	1.4	4.4	*	*	*	*	0.4	0.8	1.8	4.5	9.3	17.5
1999	1.3	5.5	*	*	*	0.2	0.4	0.8	1.6	4.1	8.8	19.3
Non-Hispanic white, female												
2002	2.5	3.6	*	*	*	0.2	0.5	1.1	2.5	6.0	12.1	22.9
2001	2.3	3.6	*	*	*	0.3	0.6	1.1	2.1	5.3	11.0	21.2
2000	2.2	3.6	*	*	*	0.3	0.5	1.2	2.3	5.7	10.6	19.8
1999	1.9	4.4	*	*	0.2	0.3	0.5	1.0	1.9	4.8	8.7	17.2
Non-Hispanic black, both sexes												
2002	2.2	8.5	*	*	*	0.9	1.3	2.3	5.0	9.5	15.7	19.7
2001	2.1	7.0	*	*	*	0.6	1.1	2.2	4.9	9.4	13.7	22.2
2000	1.9	7.8	*	*	*	0.9	1.2	2.3	4.0	8.3	12.0	20.4
1999	2.0	9.8	*	*	*	0.6	1.5	2.2	4.6	8.3	13.1	18.8
Non-Hispanic black, male												
2002	1.5	9.4	*	*	*	*	1.1	1.3	3.2	7.9	14.1	*
2001	1.5	7.3	*	*	*	*	0.8	1.7	3.7	8.5	11.4	23.1
2000	1.3	7.5	*	*	*	*	0.8	1.2	2.8	6.7	9.9	*
1999	1.4	9.4	*	*	*	*	1.2	1.5	3.4	6.9	9.7	*
Non-Hispanic black, female												
2002	2.9	7.6	*	*	*	1.2	1.5	3.2	6.5	10.6	16.6	21.7
2001	2.6	6.7	*	*	*	0.9	1.5	2.6	5.9	10.1	15.0	21.8
2000	2.5	8.1	*	*	*	1.0	1.7	3.2	4.9	9.5	13.2	20.1
1999	2.5	10.4	*	*	*	0.8	1.7	2.8	5.5	9.3	15.1	21.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other non-pulmonary forms of heart disease (I30-I51)												
All origins 3/, both sexes												
2002	54.0	7.1	0.8	0.5	1.8	3.7	8.6	18.4	41.8	114.6	361.1	1,387.2
2001	53.9	6.9	1.3	0.5	1.7	3.5	8.2	18.3	42.6	116.8	361.0	1,418.1
2000	54.1	7.1	0.9	0.5	1.8	3.5	8.1	18.0	44.2	119.7	362.9	1,452.0
1999	54.7	7.2	1.0	0.5	1.9	3.6	8.5	18.2	46.2	124.0	368.5	1,453.4
All origins 3/, male												
2002	48.6	7.3	0.9	0.5	2.5	5.0	11.4	25.1	54.3	142.4	420.4	1,396.5
2001	48.0	6.9	1.3	0.5	2.2	4.5	10.9	24.6	55.3	144.2	424.1	1,417.8
2000	48.5	7.1	1.0	0.6	2.4	4.5	10.8	24.5	58.3	150.2	422.7	1,494.4
1999	49.1	7.3	1.1	0.6	2.4	4.6	11.4	24.7	60.8	157.6	430.5	1,488.5
All origins 3/, female												
2002	59.3	6.9	0.8	0.4	1.1	2.4	5.9	11.9	30.2	91.6	321.7	1,383.1
2001	59.7	7.0	1.2	0.6	1.1	2.4	5.5	12.2	31.0	94.1	319.5	1,418.3
2000	59.6	7.2	0.7	0.4	1.3	2.5	5.5	11.7	31.3	94.7	323.9	1,434.8
1999	60.0	7.1	0.9	0.5	1.3	2.5	5.7	12.0	32.8	96.6	328.2	1,439.2
Hispanic, both sexes												
2002	12.0	6.6	0.7	0.3	1.3	2.2	4.2	9.6	22.5	65.1	203.0	709.5
2001	12.0	5.5	0.9	0.4	1.1	2.1	4.0	9.4	24.5	68.7	203.2	767.2
2000	12.3	6.0	0.8	0.5	1.5	2.3	4.4	9.6	25.8	69.7	212.5	801.6
1999	12.5	6.7	1.0	0.4	1.3	2.3	5.2	10.1	27.2	71.5	213.3	790.7
Hispanic, male												
2002	12.0	6.8	*	*	1.9	3.1	5.8	12.7	28.9	83.9	245.0	737.9
2001	11.6	6.2	*	*	1.4	2.9	5.7	13.5	31.2	83.9	233.9	731.4
2000	12.2	5.6	*	0.7	2.0	3.2	5.9	13.1	34.7	88.4	238.5	872.7
1999	12.7	5.2	*	*	1.4	3.5	7.4	14.2	35.5	95.6	252.4	847.6
Hispanic, female												
2002	12.0	6.4	*	*	0.6	1.1	2.4	6.6	16.8	50.2	174.3	695.5
2001	12.5	*	*	*	0.6	1.1	2.2	5.4	18.6	56.7	182.5	784.7
2000	12.5	6.4	*	*	0.9	1.3	2.9	6.2	18.0	54.9	195.1	766.6
1999	12.3	8.4	*	*	1.0	0.9	2.9	6.0	19.9	52.7	187.3	763.2
Non-Hispanic 4/, both sexes												
2002	60.4	7.2	0.9	0.5	1.9	4.0	9.2	19.2	43.2	117.8	368.0	1,411.9
2001	60.0	7.2	1.3	0.6	1.8	3.8	8.7	19.0	43.9	119.7	367.5	1,439.5
2000	60.0	7.4	0.9	0.5	1.9	3.7	8.5	18.7	45.5	122.5	368.6	1,472.5
1999	60.4	7.3	1.0	0.5	2.0	3.8	8.9	18.9	47.4	126.8	374.0	1,473.7
Non-Hispanic 4/, male												
2002	54.5	7.4	0.9	0.6	2.6	5.4	12.1	26.2	56.2	145.9	428.2	1,423.8
2001	53.6	7.0	1.5	0.5	2.3	4.9	11.6	25.5	56.9	147.6	432.3	1,443.2
2000	53.8	7.4	1.0	0.6	2.4	4.8	11.4	25.5	59.8	153.5	430.0	1,517.3
1999	54.3	7.8	1.1	0.6	2.6	4.8	11.8	25.5	62.3	160.7	436.8	1,510.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other non-pulmonary forms of heart disease (I30-I51)												
Non-Hispanic 4/, female												
2002	66.0	7.0	0.8	0.4	1.1	2.7	6.4	12.4	31.2	94.3	328.1	1,406.8
2001	66.2	7.5	1.2	0.6	1.2	2.7	6.0	12.8	31.9	96.4	324.9	1,438.0
2000	65.9	7.4	0.8	0.4	1.3	2.7	5.8	12.1	32.2	97.0	328.6	1,454.4
1999	66.1	6.7	0.9	0.5	1.4	2.8	6.0	12.5	33.6	99.0	333.2	1,459.0
Non-Hispanic white, both sexes												
2002	65.9	5.9	0.8	0.4	1.5	3.1	7.7	15.9	38.7	112.9	368.4	1,446.0
2001	65.2	5.6	1.0	0.5	1.4	3.0	7.0	15.9	39.3	114.4	366.4	1,475.6
2000	65.1	6.5	0.7	0.4	1.4	2.9	6.9	15.5	40.8	116.8	367.6	1,506.8
1999	65.2	5.6	0.8	0.4	1.6	3.0	7.2	15.7	42.1	121.0	371.7	1,506.0
Non-Hispanic white, male												
2002	58.3	6.1	0.9	0.5	2.2	4.1	10.4	22.2	50.7	140.0	429.2	1,459.5
2001	57.1	6.1	1.1	0.5	1.8	3.8	9.6	21.9	51.3	141.5	429.8	1,476.9
2000	57.3	6.5	0.9	0.5	1.8	3.8	9.4	21.5	54.3	146.9	429.2	1,551.9
1999	57.4	6.1	0.9	0.5	2.1	3.9	9.7	21.6	55.7	154.0	434.3	1,534.6
Non-Hispanic white, female												
2002	73.1	5.7	0.6	0.4	0.9	2.1	5.0	9.7	27.3	89.8	327.6	1,440.1
2001	73.1	5.0	0.9	0.5	1.0	2.1	4.4	10.0	27.9	91.4	324.3	1,475.0
2000	72.6	6.4	0.6	0.3	1.0	2.1	4.5	9.6	28.1	91.5	327.1	1,488.7
1999	72.7	4.9	0.7	0.4	1.1	2.2	4.6	9.9	29.4	93.5	330.7	1,494.6
Non-Hispanic black, both sexes												
2002	48.2	12.4	1.2	0.8	3.6	9.3	19.8	45.7	90.1	184.8	428.2	1,215.9
2001	48.7	14.2	2.8	0.9	3.7	8.8	20.6	45.3	91.9	189.9	443.8	1,214.9
2000	48.7	11.4	1.6	1.0	4.2	8.6	19.6	45.3	92.2	198.5	441.4	1,241.8
1999	49.8	13.2	1.9	1.0	4.0	8.7	21.2	46.1	100.3	203.7	450.2	1,241.9
Non-Hispanic black, male												
2002	47.9	12.4	*	0.9	4.7	12.7	25.2	61.7	118.4	236.0	500.3	1,191.3
2001	48.6	11.5	3.1	0.8	4.9	11.8	25.9	58.4	121.1	237.3	548.8	1,268.2
2000	48.5	11.3	*	1.1	5.3	11.0	25.3	61.7	120.7	248.2	521.7	1,322.1
1999	50.9	13.5	2.1	1.1	5.4	11.4	27.6	62.4	135.1	260.5	534.1	1,395.3
Non-Hispanic black, female												
2002	48.5	12.3	*	0.7	2.4	6.2	15.0	31.9	67.4	148.4	387.5	1,225.3
2001	48.8	17.1	2.4	0.9	2.6	6.1	15.8	34.0	68.5	156.5	384.8	1,195.1
2000	48.8	11.6	*	0.9	3.0	6.5	14.5	31.2	69.3	163.5	396.1	1,211.9
1999	48.9	12.9	*	0.9	2.7	6.2	15.5	32.3	72.4	163.7	403.2	1,184.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
All origins 3/, male												
2002	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
All origins 3/, female												
2002	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.7
2001	0.5	*	*	*	*	0.1	0.3	0.5	0.6	1.4	2.5	2.6
2000	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.5	2.9
1999	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.6
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.6	3.9
2000	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.4	3.6	4.2
1999	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.3
2001	0.4	*	*	*	*	*	0.3	0.3	0.5	1.2	1.8	2.0
2000	0.4	*	*	*	*	*	0.2	0.3	0.6	1.0	1.8	2.3
1999	0.4	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.2
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001	0.4	*	*	*	*	*	0.2	0.4	0.5	1.2	2.5	2.7
2000	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.1	2.4	2.8
1999	0.4	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	3.0
Non-Hispanic white, male												
2002	0.5	*	*	*	*	0.2	0.3	0.5	0.6	1.5	3.2	3.7
2001	0.5	*	*	*	*	*	0.2	0.5	0.7	1.4	3.6	4.0
2000	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.6	4.3
1999	0.5	*	*	*	*	*	0.2	0.4	0.7	1.3	3.3	5.1
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001	0.4	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7	2.2
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.2
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	0.4	1.3	1.6	2.3	*	*
2001	0.7	*	*	*	*	*	0.7	1.3	1.7	2.6	4.0	*
2000	0.6	*	*	*	*	*	0.6	1.4	1.6	1.9	2.9	*
1999	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	1.6	1.9	*	*	*
2001	0.8	*	*	*	*	*	0.8	1.9	2.2	3.9	*	*
2000	0.7	*	*	*	*	*	0.8	1.8	*	*	*	*
1999	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001	0.5	*	*	*	*	*	0.7	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	1.0	1.6	*	*	*
1999	0.5	*	*	*	*	*	*	*	1.6	2.9	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All origins 3/, both sexes												
2002	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0
All origins 3/, male												
2002	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5
All origins 3/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	0.7	0.2	*	0.1	0.1	0.2	0.3	0.4	0.8	1.3	2.1
2001	0.3	0.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999	0.3	0.8	0.2	*	0.1	0.1	0.3	0.3	0.6	0.8	1.3	2.1
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.9	1.2	2.3
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000	0.3	*	*	*	0.1	0.2	0.3	0.3	0.4	0.9	1.4	2.1
1999	0.3	*	*	*	*	0.2	0.3	0.3	0.6	0.9	1.4	2.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.0
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	1.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.3	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.9
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	2.1
2001	0.3	*	*	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.1	1.8
1999	0.3	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.3	2.0
Non-Hispanic white, male												
2002	0.3	*	*	*	*	0.2	0.2	0.3	0.4	0.9	1.2	2.3
2001	0.3	*	*	*	*	*	0.2	0.3	0.5	0.7	1.2	2.0
2000	0.3	*	*	*	*	*	0.2	0.2	0.4	0.8	1.4	2.1
1999	0.3	*	*	*	*	*	0.2	0.3	0.7	0.8	1.4	2.6
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.1
2001	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.8
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	0.7	1.2	1.8
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.6	0.9	*	2.1	*
2001	0.4	*	*	*	*	*	0.4	0.7	*	*	*	*
2000	0.4	*	*	*	*	*	0.4	0.5	*	*	*	*
1999	0.5	*	*	*	*	*	0.5	0.5	*	1.2	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of endocardium (I34-I38)												
All origins 3/, both sexes												
2002	6.9	*	*	*	0.1	0.3	0.8	1.6	3.7	12.0	49.3	203.9
2001	6.9	*	*	*	0.1	0.3	0.7	1.6	3.7	13.0	48.8	204.2
2000	7.0	*	*	*	0.1	0.3	0.8	1.6	4.0	13.2	49.8	213.2
1999	7.0	*	*	*	0.2	0.4	0.8	1.6	3.9	13.2	51.7	209.8
All origins 3/, male												
2002	5.4	*	*	*	0.1	0.3	0.9	2.0	4.1	12.7	51.6	198.5
2001	5.4	*	*	*	*	0.3	0.8	2.1	4.2	14.0	52.8	200.8
2000	5.5	*	*	*	0.1	0.3	0.9	2.1	4.7	14.0	51.6	215.8
1999	5.5	*	*	*	0.2	0.3	1.0	1.9	4.5	14.7	56.4	206.4
All origins 3/, female												
2002	8.4	*	*	*	*	0.3	0.6	1.2	3.4	11.4	47.7	206.3
2001	8.2	*	*	*	0.1	0.3	0.5	1.2	3.2	12.1	46.2	205.6
2000	8.5	*	*	*	0.1	0.3	0.7	1.2	3.3	12.6	48.6	212.1
1999	8.5	*	*	*	0.2	0.4	0.7	1.4	3.3	12.1	48.7	211.2
Hispanic, both sexes												
2002	1.5	*	*	*	*	*	0.6	1.3	3.5	7.8	24.4	92.6
2001	1.5	*	*	*	*	*	*	1.2	3.1	10.0	27.6	95.0
2000	1.6	*	*	*	*	*	0.4	1.4	3.6	9.3	29.6	116.8
1999	1.6	*	*	*	*	*	0.6	0.9	3.1	9.4	34.4	98.2
Hispanic, male												
2002	1.4	*	*	*	*	*	0.8	1.6	3.7	8.4	26.7	103.7
2001	1.3	*	*	*	*	*	*	1.5	3.9	10.4	33.1	81.9
2000	1.4	*	*	*	*	*	*	1.9	4.4	9.3	32.0	120.9
1999	1.5	*	*	*	*	*	*	1.4	4.6	10.5	37.0	110.0
Hispanic, female												
2002	1.6	*	*	*	*	*	*	*	3.4	7.3	22.8	87.0
2001	1.7	*	*	*	*	*	*	*	2.5	9.8	23.9	101.5
2000	1.8	*	*	*	*	*	*	*	2.9	9.3	28.1	114.7
1999	1.7	*	*	*	*	*	*	*	*	8.4	32.6	92.5
Non-Hispanic 4/, both sexes												
2002	7.8	*	*	*	0.1	0.3	0.8	1.6	3.7	12.2	50.3	208.2
2001	7.7	*	*	*	0.1	0.3	0.7	1.7	3.7	13.2	49.7	207.9
2000	7.8	*	*	*	0.1	0.3	0.8	1.6	4.0	13.5	50.6	216.2
1999	7.8	*	*	*	0.2	0.4	0.9	1.7	4.0	13.4	52.3	213.4
Non-Hispanic 4/, male												
2002	6.1	*	*	*	0.1	0.4	0.9	2.0	4.1	12.9	52.8	202.4
2001	6.1	*	*	*	*	0.4	0.8	2.1	4.2	14.2	53.7	205.5
2000	6.1	*	*	*	0.1	0.3	0.9	2.1	4.7	14.2	52.4	219.2
1999	6.1	*	*	*	0.2	0.4	1.0	1.9	4.5	14.9	57.0	209.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of endocardium (I34-I38)												
Non-Hispanic 4/, female												
2002	9.4	*	*	*	*	0.3	0.7	1.3	3.4	11.7	48.7	210.7
2001	9.2	*	*	*	0.1	0.3	0.6	1.2	3.2	12.3	47.1	208.8
2000	9.4	*	*	*	*	0.3	0.7	1.2	3.3	12.8	49.4	214.9
1999	9.3	*	*	*	0.2	0.4	0.7	1.5	3.4	12.2	49.2	214.8
Non-Hispanic white, both sexes												
2002	9.0	*	*	*	0.1	0.3	0.7	1.3	3.6	12.4	53.2	221.1
2001	8.8	*	*	*	0.1	0.3	0.6	1.5	3.5	13.3	52.2	220.7
2000	8.9	*	*	*	0.1	0.3	0.7	1.5	3.7	13.6	52.8	229.3
1999	8.9	*	*	*	0.2	0.4	0.8	1.5	3.7	13.3	54.6	225.5
Non-Hispanic white, male												
2002	7.0	*	*	*	*	0.3	0.9	1.7	4.0	13.3	55.7	216.3
2001	6.9	*	*	*	*	0.3	0.7	1.9	4.0	14.4	56.4	217.3
2000	6.9	*	*	*	*	0.2	0.8	1.8	4.5	14.5	54.7	233.1
1999	6.9	*	*	*	0.2	0.3	0.9	1.7	4.3	14.9	59.6	222.3
Non-Hispanic white, female												
2002	11.0	*	*	*	*	0.3	0.5	1.0	3.2	11.7	51.5	223.2
2001	10.7	*	*	*	*	0.3	0.5	1.0	3.0	12.4	49.5	222.1
2000	10.9	*	*	*	*	0.3	0.6	1.1	2.9	12.8	51.6	227.8
1999	10.8	*	*	*	*	0.4	0.6	1.3	3.2	12.0	51.4	226.8
Non-Hispanic black, both sexes												
2002	3.3	*	*	*	*	0.5	1.5	3.8	5.8	13.1	28.6	87.4
2001	3.4	*	*	*	*	0.6	1.3	3.3	6.2	13.7	31.0	86.5
2000	3.6	*	*	*	*	0.6	1.5	3.2	6.6	14.9	33.9	92.5
1999	3.6	*	*	*	*	0.6	1.6	3.5	6.6	16.1	32.3	92.1
Non-Hispanic black, male												
2002	2.7	*	*	*	*	*	1.4	4.3	6.7	12.3	28.7	67.8
2001	3.0	*	*	*	*	*	1.5	4.3	7.5	14.5	31.2	87.8
2000	3.1	*	*	*	*	*	1.6	4.3	7.2	14.8	34.9	85.7
1999	3.1	*	*	*	*	*	1.7	4.0	7.4	17.0	34.1	77.3
Non-Hispanic black, female												
2002	3.9	*	*	*	*	*	1.5	3.3	5.1	13.6	28.5	94.9
2001	3.7	*	*	*	*	*	1.0	2.5	5.3	13.2	30.8	86.0
2000	4.1	*	*	*	*	*	1.5	2.3	6.1	15.0	33.3	95.1
1999	4.1	*	*	*	*	*	1.4	3.1	6.0	15.5	31.3	97.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)												
All origins 3/, both sexes												
2002	1.0	*	*	*	0.1	0.1	0.2	0.3	0.7	2.1	7.0	22.1
2001	1.0	*	*	*	0.1	0.1	0.2	0.3	0.6	2.2	7.3	22.7
2000	1.0	*	*	*	0.1	0.1	0.2	0.3	0.7	2.2	7.7	24.3
1999	1.0	*	*	*	0.1	0.1	0.2	0.4	0.7	2.3	7.7	25.1
All origins 3/, male												
2002	0.7	*	*	*	*	*	0.2	0.2	0.6	1.8	5.7	21.0
2001	0.7	*	*	*	*	0.1	0.2	0.3	0.6	2.2	6.1	19.0
2000	0.7	*	*	*	*	*	0.2	0.3	0.6	2.1	7.2	22.9
1999	0.7	*	*	*	*	*	0.2	0.3	0.6	2.1	7.5	22.2
All origins 3/, female												
2002	1.2	*	*	*	*	0.1	0.2	0.3	0.8	2.4	7.9	22.6
2001	1.3	*	*	*	*	0.2	0.2	0.3	0.6	2.2	8.1	24.3
2000	1.3	*	*	*	*	0.1	0.3	0.3	0.8	2.4	8.1	24.9
1999	1.3	*	*	*	*	0.2	0.2	0.4	0.8	2.4	7.8	26.2
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	3.7	*
2001	0.2	*	*	*	*	*	*	*	*	2.0	4.9	*
2000	0.2	*	*	*	*	*	*	*	*	*	4.3	*
1999	0.2	*	*	*	*	*	*	*	*	*	4.2	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.1	*	*	*	*	0.1	0.2	0.3	0.7	2.2	7.2	22.7
2001	1.1	*	*	*	0.1	0.2	0.2	0.4	0.6	2.2	7.4	23.2
2000	1.1	*	*	*	*	0.1	0.3	0.3	0.7	2.3	7.9	24.8
1999	1.2	*	*	*	0.1	0.2	0.3	0.4	0.7	2.3	7.8	25.5
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	0.2	0.2	0.6	1.9	5.9	21.7
2001	0.8	*	*	*	*	0.1	0.2	0.4	0.6	2.2	6.1	19.3
2000	0.8	*	*	*	*	*	0.2	0.4	0.6	2.0	7.2	23.4
1999	0.8	*	*	*	*	*	0.3	0.3	0.6	2.2	7.7	22.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)												
Non-Hispanic 4/, female												
2002	1.4	*	*	*	*	0.2	0.2	0.3	0.8	2.5	8.0	23.2
2001	1.4	*	*	*	*	0.2	0.3	0.4	0.6	2.2	8.2	24.8
2000	1.4	*	*	*	*	0.2	0.3	0.3	0.8	2.4	8.3	25.3
1999	1.5	*	*	*	*	0.2	0.3	0.4	0.8	2.5	7.8	26.7
Non-Hispanic white, both sexes												
2002	1.2	*	*	*	*	0.1	0.2	0.2	0.7	2.2	7.5	24.5
2001	1.2	*	*	*	*	0.2	0.2	0.3	0.6	2.2	7.8	24.8
2000	1.3	*	*	*	*	0.1	0.3	0.3	0.7	2.3	8.3	26.0
1999	1.3	*	*	*	*	0.2	0.3	0.3	0.7	2.3	8.1	27.0
Non-Hispanic white, male												
2002	0.8	*	*	*	*	*	0.2	0.2	0.6	1.9	6.2	23.3
2001	0.9	*	*	*	*	*	0.2	0.4	0.6	2.2	6.6	20.9
2000	0.9	*	*	*	*	*	0.2	0.3	0.6	2.1	7.7	24.8
1999	0.9	*	*	*	*	*	0.3	0.3	0.7	2.2	8.1	24.2
Non-Hispanic white, female												
2002	1.6	*	*	*	*	0.2	0.2	0.3	0.8	2.5	8.4	25.0
2001	1.6	*	*	*	*	0.2	0.3	0.3	0.6	2.2	8.6	26.4
2000	1.6	*	*	*	*	0.2	0.3	0.3	0.8	2.5	8.7	26.4
1999	1.7	*	*	*	*	0.2	0.3	0.4	0.8	2.5	8.1	28.1
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	*	0.6	0.9	2.2	4.0	7.0
2001	0.5	*	*	*	*	*	*	0.6	0.9	2.7	3.6	7.8
2000	0.5	*	*	*	*	*	*	0.6	1.0	2.2	4.4	12.1
1999	0.6	*	*	*	*	*	*	0.6	1.0	2.5	4.5	10.3
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.6	*	*	*	*	*	*	*	*	2.5	4.5	*
2001	0.7	*	*	*	*	*	*	*	*	2.8	4.5	9.8
2000	0.7	*	*	*	*	*	*	*	*	2.7	5.0	13.5
1999	0.7	*	*	*	*	*	*	*	*	2.4	5.4	12.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)												
All origins 3/, both sexes												
2002	4.3	*	*	*	*	0.1	0.2	0.6	1.6	6.3	30.2	144.8
2001	4.3	*	*	*	*	0.1	0.2	0.7	1.7	6.8	30.0	144.1
2000	4.4	*	*	*	*	0.1	0.2	0.6	1.8	7.1	30.9	151.3
1999	4.4	*	*	*	*	0.1	0.3	0.7	1.8	6.9	32.4	147.8
All origins 3/, male												
2002	3.5	*	*	*	*	0.1	0.3	0.8	2.0	7.4	34.4	144.9
2001	3.5	*	*	*	*	0.1	0.3	1.0	2.1	8.0	34.8	145.9
2000	3.5	*	*	*	*	0.1	0.3	1.0	2.5	8.2	33.3	158.2
1999	3.5	*	*	*	*	0.1	0.4	0.9	2.3	8.6	37.2	150.3
All origins 3/, female												
2002	5.2	*	*	*	*	*	0.2	0.4	1.3	5.3	27.4	144.8
2001	5.1	*	*	*	*	*	*	0.4	1.3	5.8	26.9	143.4
2000	5.3	*	*	*	*	*	0.1	0.3	1.2	6.1	29.2	148.5
1999	5.2	*	*	*	*	*	0.2	0.4	1.3	5.6	29.2	146.8
Hispanic, both sexes												
2002	0.9	*	*	*	*	*	*	0.6	1.5	4.8	15.5	69.3
2001	0.8	*	*	*	*	*	*	*	1.6	4.7	16.6	70.7
2000	1.0	*	*	*	*	*	*	*	1.6	4.6	19.8	88.9
1999	0.9	*	*	*	*	*	*	*	1.3	5.2	22.7	75.8
Hispanic, male												
2002	0.8	*	*	*	*	*	*	*	*	5.5	19.3	83.7
2001	0.8	*	*	*	*	*	*	*	2.4	5.2	21.9	59.0
2000	0.9	*	*	*	*	*	*	*	2.6	4.6	24.1	96.7
1999	0.9	*	*	*	*	*	*	*	*	6.6	27.6	82.0
Hispanic, female												
2002	0.9	*	*	*	*	*	*	*	*	4.3	12.9	62.2
2001	0.9	*	*	*	*	*	*	*	*	4.3	13.0	76.3
2000	1.0	*	*	*	*	*	*	*	*	4.6	16.8	85.1
1999	0.9	*	*	*	*	*	*	*	*	4.1	19.4	72.8
Non-Hispanic 4/, both sexes												
2002	4.9	*	*	*	*	0.1	0.2	0.6	1.6	6.3	30.8	147.7
2001	4.8	*	*	*	*	0.1	0.2	0.7	1.7	6.9	30.6	146.6
2000	4.9	*	*	*	*	0.1	0.2	0.6	1.8	7.2	31.3	153.2
1999	4.8	*	*	*	*	0.1	0.3	0.7	1.8	7.0	32.7	150.0
Non-Hispanic 4/, male												
2002	3.9	*	*	*	*	0.1	0.3	0.8	2.0	7.5	35.1	147.4
2001	3.9	*	*	*	*	0.1	0.3	1.0	2.1	8.1	35.3	149.3
2000	3.9	*	*	*	*	*	0.3	1.0	2.5	8.5	33.7	160.3
1999	3.9	*	*	*	*	0.1	0.4	0.9	2.3	8.7	37.5	152.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)												
Non-Hispanic 4/, female												
2002	5.8	*	*	*	*	*	0.2	0.4	1.3	5.3	28.0	147.8
2001	5.6	*	*	*	*	*	*	0.4	1.3	5.9	27.5	145.4
2000	5.8	*	*	*	*	*	0.1	0.3	1.2	6.2	29.7	150.4
1999	5.7	*	*	*	*	*	0.2	0.4	1.3	5.6	29.5	148.9
Non-Hispanic white, both sexes												
2002	5.7	*	*	*	*	*	0.2	0.6	1.7	6.5	32.7	156.5
2001	5.6	*	*	*	*	0.1	0.2	0.7	1.7	7.2	32.2	155.6
2000	5.7	*	*	*	*	0.1	0.2	0.6	1.8	7.4	32.9	162.9
1999	5.6	*	*	*	*	*	0.3	0.6	1.8	7.0	34.4	158.8
Non-Hispanic white, male												
2002	4.6	*	*	*	*	*	0.3	0.8	2.1	7.9	37.2	157.4
2001	4.5	*	*	*	*	*	0.3	1.0	2.1	8.4	37.0	157.3
2000	4.5	*	*	*	*	*	0.3	0.9	2.5	8.6	35.4	171.0
1999	4.5	*	*	*	*	*	0.4	0.8	2.3	8.8	39.3	162.1
Non-Hispanic white, female												
2002	6.8	*	*	*	*	*	*	0.3	1.4	5.4	29.7	156.2
2001	6.7	*	*	*	*	*	*	0.4	1.4	6.1	29.1	154.9
2000	6.9	*	*	*	*	*	*	0.3	1.1	6.3	31.3	159.7
1999	6.7	*	*	*	*	*	0.2	0.4	1.3	5.6	31.1	157.5
Non-Hispanic black, both sexes												
2002	1.6	*	*	*	*	*	0.4	1.0	1.6	5.9	16.5	63.5
2001	1.6	*	*	*	*	*	*	1.0	1.9	5.6	18.6	60.2
2000	1.7	*	*	*	*	*	0.4	0.9	2.4	7.3	18.6	63.0
1999	1.8	*	*	*	*	*	0.4	1.2	2.2	7.6	18.0	64.7
Non-Hispanic black, male												
2002	1.3	*	*	*	*	*	*	1.4	2.2	6.3	17.0	52.5
2001	1.5	*	*	*	*	*	*	1.6	2.3	6.4	21.9	69.3
2000	1.5	*	*	*	*	*	*	1.5	2.8	8.4	19.7	57.5
1999	1.6	*	*	*	*	*	*	1.8	2.8	9.3	21.9	58.2
Non-Hispanic black, female												
2002	1.9	*	*	*	*	*	*	*	*	5.6	16.2	67.7
2001	1.7	*	*	*	*	*	*	*	1.5	5.1	16.7	56.9
2000	2.0	*	*	*	*	*	*	*	2.0	6.5	17.9	65.0
1999	1.9	*	*	*	*	*	*	*	1.8	6.5	15.8	67.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of endocardium (I36-I38)												
All origins 3/, both sexes												
2002	1.7	*	*	*	*	0.1	0.3	0.7	1.4	3.6	12.1	37.0
2001	1.6	*	*	*	*	0.1	0.3	0.6	1.4	4.0	11.5	37.4
2000	1.6	*	*	*	*	0.1	0.3	0.7	1.4	3.9	11.2	37.6
1999	1.6	*	*	*	0.1	0.1	0.3	0.6	1.4	4.0	11.7	37.0
All origins 3/, male												
2002	1.3	*	*	*	*	0.1	0.4	0.9	1.5	3.5	11.5	32.6
2001	1.3	*	*	*	*	0.1	0.3	0.8	1.5	3.9	12.0	35.9
2000	1.2	*	*	*	*	*	0.4	0.8	1.6	3.6	11.1	34.7
1999	1.3	*	*	*	*	0.1	0.4	0.7	1.6	3.9	11.7	33.9
All origins 3/, female												
2002	2.0	*	*	*	*	0.1	0.3	0.5	1.3	3.8	12.4	38.9
2001	1.9	*	*	*	*	*	0.2	0.4	1.3	4.1	11.2	38.0
2000	1.9	*	*	*	*	0.1	0.2	0.6	1.3	4.1	11.3	38.7
1999	2.0	*	*	*	*	0.1	0.3	0.6	1.3	4.1	11.7	38.2
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	0.3	*	1.3	1.8	5.2	15.5
2001	0.4	*	*	*	*	*	*	*	1.1	3.3	6.1	15.0
2000	0.4	*	*	*	*	*	*	0.6	1.3	2.9	5.5	17.9
1999	0.4	*	*	*	*	*	*	*	1.5	2.9	7.5	*
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	5.6	16.6
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	3.3	7.6	20.8
1999	0.5	*	*	*	*	*	*	*	*	*	7.6	*
Non-Hispanic 4/, both sexes												
2002	1.8	*	*	*	*	0.1	0.3	0.7	1.4	3.7	12.4	37.8
2001	1.8	*	*	*	*	0.1	0.3	0.6	1.4	4.0	11.7	38.1
2000	1.8	*	*	*	*	0.1	0.3	0.7	1.4	4.0	11.5	38.2
1999	1.8	*	*	*	0.1	0.1	0.3	0.6	1.4	4.1	11.8	37.8
Non-Hispanic 4/, male												
2002	1.4	*	*	*	*	0.1	0.4	0.9	1.5	3.5	11.8	33.3
2001	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.9	12.2	36.9
2000	1.4	*	*	*	*	*	0.4	0.8	1.6	3.7	11.5	35.6
1999	1.4	*	*	*	*	0.1	0.4	0.7	1.6	4.0	11.8	34.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of endocardium (I36-I38)												
Non-Hispanic 4/, female												
2002	2.2	*	*	*	*	0.1	0.3	0.6	1.3	3.9	12.7	39.7
2001	2.1	*	*	*	*	*	0.3	0.5	1.3	4.2	11.4	38.6
2000	2.1	*	*	*	*	0.1	0.2	0.6	1.3	4.2	11.5	39.2
1999	2.1	*	*	*	*	0.1	0.3	0.6	1.3	4.1	11.8	39.2
Non-Hispanic white, both sexes												
2002	2.0	*	*	*	*	0.1	0.3	0.5	1.2	3.6	13.0	40.1
2001	2.0	*	*	*	*	*	0.2	0.4	1.1	4.0	12.2	40.3
2000	1.9	*	*	*	*	*	0.2	0.5	1.2	3.9	11.6	40.4
1999	1.9	*	*	*	*	0.1	0.2	0.5	1.2	4.0	12.2	39.8
Non-Hispanic white, male												
2002	1.5	*	*	*	*	*	0.3	0.7	1.2	3.5	12.3	35.6
2001	1.5	*	*	*	*	*	0.3	0.6	1.2	3.8	12.9	39.1
2000	1.5	*	*	*	*	*	0.3	0.6	1.4	3.7	11.6	37.4
1999	1.5	*	*	*	*	*	0.3	0.6	1.3	4.0	12.2	36.0
Non-Hispanic white, female												
2002	2.5	*	*	*	*	*	0.2	0.4	1.1	3.7	13.5	42.1
2001	2.4	*	*	*	*	*	0.2	0.3	1.1	4.1	11.8	40.8
2000	2.4	*	*	*	*	*	0.1	0.5	1.1	4.1	11.7	41.6
1999	2.4	*	*	*	*	*	0.2	0.5	1.1	4.0	12.2	41.3
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	*	*	0.8	2.2	3.2	5.0	8.0	16.9
2001	1.3	*	*	*	*	*	0.6	1.8	3.5	5.4	8.8	18.4
2000	1.3	*	*	*	*	*	0.8	1.8	3.2	5.4	10.9	17.5
1999	1.3	*	*	*	*	*	0.9	1.7	3.3	6.0	9.9	17.1
Non-Hispanic black, male												
2002	1.1	*	*	*	*	*	0.8	2.5	3.7	4.2	8.5	*
2001	1.2	*	*	*	*	*	*	2.2	4.5	5.6	7.5	*
2000	1.2	*	*	*	*	*	0.8	2.2	3.6	5.1	11.7	*
1999	1.1	*	*	*	*	*	0.9	1.8	4.0	5.1	9.4	*
Non-Hispanic black, female												
2002	1.4	*	*	*	*	*	0.9	1.9	2.8	5.5	7.8	19.6
2001	1.3	*	*	*	*	*	*	1.4	2.7	5.2	9.6	19.2
2000	1.4	*	*	*	*	*	0.8	1.4	3.0	5.7	10.4	16.6
1999	1.4	*	*	*	*	*	0.9	1.6	2.8	6.6	10.2	18.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cardiomyopathy (I42)												
All origins 3/, both sexes												
2002	9.6	2.9	0.2	0.2	0.7	1.6	3.5	6.5	11.9	27.0	61.1	142.0
2001	9.4	3.4	0.4	0.2	0.7	1.4	3.3	6.3	12.3	27.2	60.4	138.3
2000	9.7	3.1	0.3	0.2	0.7	1.5	3.3	6.7	13.0	28.7	61.4	142.3
1999	9.8	3.5	0.2	0.2	0.7	1.5	3.6	6.7	14.0	30.4	60.4	139.2
All origins 3/, male												
2002	11.3	2.7	*	0.2	1.0	2.3	4.8	9.4	16.7	37.8	87.2	189.7
2001	11.0	2.9	0.5	0.2	0.9	2.0	4.8	8.9	17.3	37.4	84.9	184.7
2000	11.4	2.8	0.3	0.3	0.9	2.1	4.8	9.6	18.2	40.0	87.6	195.3
1999	11.5	3.1	*	0.2	0.9	2.0	5.2	9.6	19.6	43.0	84.3	188.2
All origins 3/, female												
2002	8.0	3.1	*	0.1	0.4	0.9	2.2	3.6	7.5	18.1	43.7	121.3
2001	8.0	3.9	0.3	0.2	0.4	0.8	1.9	3.9	7.7	18.8	44.2	118.9
2000	8.0	3.4	*	0.1	0.4	1.0	1.9	4.0	8.2	19.4	44.3	120.7
1999	8.1	3.8	*	0.1	0.4	1.0	2.0	4.0	8.9	20.0	44.9	119.4
Hispanic, both sexes												
2002	3.4	3.7	*	*	0.5	1.1	1.8	4.4	7.6	19.5	51.2	100.9
2001	3.4	2.9	*	*	0.5	0.9	1.8	4.0	8.9	21.8	50.3	122.6
2000	3.5	2.6	*	*	0.6	1.1	2.1	4.0	9.2	21.8	52.7	122.8
1999	3.4	3.7	*	*	0.3	0.9	2.5	4.8	10.5	21.5	47.9	105.2
Hispanic, male												
2002	4.1	*	*	*	0.8	1.6	2.6	6.1	10.5	28.3	71.0	137.2
2001	3.9	*	*	*	0.7	1.4	2.7	6.0	12.7	30.6	64.8	129.5
2000	4.1	*	*	*	0.8	1.5	2.9	5.9	13.6	29.3	70.3	151.2
1999	4.2	*	*	*	*	1.5	3.7	7.0	14.6	33.4	65.2	125.1
Hispanic, female												
2002	2.6	*	*	*	*	*	1.0	2.7	5.1	12.5	37.7	82.9
2001	2.9	*	*	*	*	*	0.9	2.1	5.6	15.0	40.5	119.1
2000	2.9	*	*	*	*	*	1.2	2.3	5.3	15.9	40.9	108.8
1999	2.6	5.4	*	*	*	*	1.1	2.7	6.9	12.2	36.4	95.7
Non-Hispanic 4/, both sexes												
2002	10.6	2.7	0.2	0.2	0.7	1.7	3.7	6.7	12.2	27.5	61.4	143.5
2001	10.3	3.5	0.4	0.2	0.7	1.5	3.5	6.5	12.6	27.5	60.7	138.6
2000	10.5	3.2	0.2	0.2	0.7	1.6	3.5	6.9	13.3	29.0	61.7	142.7
1999	10.6	3.4	0.2	0.2	0.7	1.6	3.7	6.9	14.2	30.8	60.9	140.1
Non-Hispanic 4/, male												
2002	12.5	2.5	*	0.2	1.0	2.5	5.1	9.7	17.2	38.4	87.8	192.0
2001	12.0	2.9	0.5	0.2	0.9	2.1	5.1	9.1	17.6	37.7	85.7	186.3
2000	12.4	2.8	*	0.2	1.0	2.2	5.0	9.9	18.5	40.5	88.2	196.7
1999	12.5	3.4	*	0.2	1.0	2.1	5.3	9.8	19.8	43.4	84.9	190.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cardiomyopathy (I42)												
Non-Hispanic 4/, female												
2002	8.7	2.9	*	0.1	0.4	1.0	2.4	3.7	7.7	18.5	43.9	122.6
2001	8.7	4.1	0.3	0.1	0.4	0.9	2.0	4.1	7.9	19.0	44.3	118.7
2000	8.7	3.6	*	*	0.4	1.0	2.0	4.1	8.4	19.6	44.4	120.8
1999	8.8	3.4	*	0.1	0.4	1.1	2.1	4.1	9.0	20.5	45.3	120.0
Non-Hispanic white, both sexes												
2002	10.7	2.5	0.2	0.2	0.5	1.2	2.9	5.3	10.5	25.7	60.1	143.2
2001	10.3	2.7	0.3	0.1	0.5	1.1	2.6	5.1	10.6	25.6	58.8	139.3
2000	10.5	3.0	*	0.1	0.5	1.1	2.6	5.3	11.3	26.6	59.6	142.6
1999	10.5	2.8	0.2	0.1	0.5	1.2	2.7	5.3	11.7	28.5	58.9	139.5
Non-Hispanic white, male												
2002	12.5	2.0	*	0.2	0.7	1.7	4.0	7.8	14.9	35.9	86.9	193.1
2001	12.0	2.5	0.4	*	0.7	1.5	3.8	7.3	15.0	35.4	83.7	188.4
2000	12.3	2.7	*	0.2	0.6	1.5	3.7	7.8	15.9	37.4	85.7	198.7
1999	12.2	2.8	*	0.2	0.7	1.6	3.9	7.6	16.4	40.1	83.0	189.9
Non-Hispanic white, female												
2002	8.9	3.0	*	*	0.2	0.6	1.7	2.8	6.3	17.1	42.1	121.7
2001	8.7	2.8	*	*	0.3	0.6	1.3	2.9	6.4	17.3	42.4	118.8
2000	8.8	3.4	*	*	0.2	0.7	1.4	2.9	7.0	17.6	42.4	119.9
1999	8.9	2.9	*	*	0.3	0.8	1.4	3.0	7.3	18.7	43.2	119.3
Non-Hispanic black, both sexes												
2002	12.7	3.4	*	*	1.7	4.8	9.5	17.1	29.1	48.4	86.0	161.8
2001	12.9	7.2	0.9	0.3	1.7	4.1	9.8	17.8	31.0	49.6	91.4	143.6
2000	13.3	3.6	*	0.4	1.9	4.4	9.1	19.9	30.9	55.8	93.9	153.0
1999	13.6	5.1	*	0.4	1.9	3.9	10.3	19.7	35.9	56.9	89.5	154.9
Non-Hispanic black, male												
2002	15.4	*	*	*	2.2	7.0	13.0	25.3	40.8	69.7	114.9	207.9
2001	15.5	*	*	*	2.3	5.7	13.6	24.3	44.4	67.8	129.6	183.6
2000	16.3	*	*	*	2.5	6.1	13.2	27.6	44.6	78.5	133.6	196.1
1999	17.0	*	*	*	2.7	5.4	14.7	28.4	52.9	82.8	118.5	218.7
Non-Hispanic black, female												
2002	10.2	*	*	*	1.2	2.8	6.4	10.1	19.7	33.2	69.6	144.2
2001	10.5	9.4	*	*	1.1	2.7	6.4	12.2	20.2	36.7	69.9	128.8
2000	10.6	*	*	*	1.2	2.9	5.5	13.2	20.0	39.9	71.6	136.9
1999	10.6	*	*	*	1.1	2.6	6.4	12.2	22.2	38.6	73.2	131.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)												
All origins 3/, both sexes												
2002	13.6	1.4	0.1	0.1	0.5	0.8	2.0	4.9	12.6	31.5	93.4	323.2
2001	13.5	0.9	0.2	0.1	0.4	0.7	1.7	5.0	12.6	32.6	92.4	328.2
2000	13.6	1.0	0.2	0.1	0.5	0.7	1.9	4.8	13.1	33.3	92.2	340.2
1999	14.3	1.0	0.3	0.1	0.5	0.8	1.9	5.2	14.5	35.7	97.1	353.7
All origins 3/, male												
2002	12.2	1.6	*	0.1	0.6	0.9	2.5	6.6	16.8	39.0	103.8	310.5
2001	12.0	*	*	0.1	0.6	0.9	2.2	6.7	16.5	39.8	104.9	311.6
2000	12.1	1.1	*	0.1	0.5	0.8	2.3	6.6	17.5	42.9	101.9	333.2
1999	12.9	1.2	0.3	0.2	0.6	1.0	2.3	7.1	19.2	45.4	110.8	353.4
All origins 3/, female												
2002	15.1	1.2	*	*	0.3	0.6	1.4	3.3	8.6	25.3	86.6	328.8
2001	15.1	*	*	0.1	0.2	0.6	1.3	3.4	9.0	26.7	84.2	335.2
2000	15.1	*	*	*	0.4	0.6	1.5	3.0	8.9	25.4	85.8	343.0
1999	15.7	*	*	0.1	0.4	0.6	1.4	3.4	10.2	27.8	88.2	353.8
Hispanic, both sexes												
2002	2.0	*	*	*	*	0.4	0.6	1.3	3.9	10.7	32.9	134.7
2001	2.1	*	*	*	*	0.3	0.6	1.4	3.9	12.7	36.4	145.7
2000	2.2	*	*	*	0.4	0.4	0.8	1.7	4.6	13.1	39.3	142.7
1999	2.3	*	*	*	*	0.3	0.6	1.5	5.5	14.0	40.6	164.2
Hispanic, male												
2002	2.0	*	*	*	*	0.6	0.8	1.8	5.0	13.8	39.0	130.5
2001	1.9	*	*	*	*	*	0.8	1.9	4.8	14.8	38.9	137.1
2000	2.2	*	*	*	*	*	0.9	2.6	5.6	18.4	39.8	155.2
1999	2.3	*	*	*	*	*	0.8	2.3	7.2	17.1	45.9	198.4
Hispanic, female												
2002	2.1	*	*	*	*	*	*	*	3.0	8.2	28.7	136.8
2001	2.3	*	*	*	*	*	*	*	3.1	11.0	34.8	149.9
2000	2.3	*	*	*	*	*	*	*	3.6	9.0	39.0	136.5
1999	2.4	*	*	*	*	*	*	*	4.0	11.5	37.1	147.7
Non-Hispanic 4/, both sexes												
2002	15.4	1.6	*	0.1	0.5	0.9	2.2	5.3	13.2	32.8	96.2	330.2
2001	15.2	1.1	0.2	0.2	0.4	0.8	1.9	5.4	13.2	33.8	94.8	334.2
2000	15.2	1.2	0.2	0.1	0.5	0.8	2.0	5.0	13.7	34.5	94.3	346.6
1999	15.9	1.2	0.3	0.1	0.5	0.9	2.0	5.5	15.1	36.9	99.2	359.7
Non-Hispanic 4/, male												
2002	13.8	1.6	*	0.2	0.7	1.0	2.8	7.1	17.7	40.6	106.8	318.1
2001	13.5	*	*	0.1	0.6	0.9	2.4	7.1	17.4	41.2	107.7	318.1
2000	13.6	1.3	*	0.1	0.5	0.9	2.5	6.9	18.4	44.3	104.4	339.7
1999	14.4	1.4	0.3	0.2	0.6	1.0	2.5	7.5	20.0	46.9	113.2	359.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Non-Hispanic 4/, female												
2002	16.9	1.5	*	*	0.3	0.7	1.6	3.5	9.1	26.4	89.2	335.4
2001	16.8	*	*	0.2	0.2	0.6	1.4	3.7	9.4	27.7	86.2	340.9
2000	16.8	*	*	*	0.4	0.6	1.6	3.2	9.3	26.4	87.7	349.3
1999	17.4	*	*	*	0.4	0.7	1.5	3.6	10.6	28.7	90.2	360.0
Non-Hispanic white, both sexes												
2002	16.6	1.3	*	0.1	0.5	0.7	2.0	4.7	12.0	31.4	94.6	331.7
2001	16.3	1.0	*	0.1	0.4	0.7	1.6	4.8	12.1	32.1	92.7	336.0
2000	16.4	1.1	*	0.1	0.4	0.7	1.8	4.5	12.6	33.0	93.1	348.3
1999	17.2	0.9	0.3	0.1	0.5	0.7	1.8	5.0	14.0	35.5	97.5	362.4
Non-Hispanic white, male												
2002	14.7	1.7	*	0.2	0.7	0.9	2.7	6.5	16.2	38.9	104.7	316.8
2001	14.2	*	*	0.2	0.5	0.8	2.1	6.5	15.8	39.3	104.4	316.9
2000	14.5	*	*	*	0.5	0.8	2.3	6.3	17.1	42.6	103.7	339.9
1999	15.3	*	*	0.2	0.6	0.9	2.4	7.0	18.7	45.3	110.4	357.1
Non-Hispanic white, female												
2002	18.5	*	*	*	0.3	0.5	1.4	3.0	7.9	24.9	87.8	338.2
2001	18.3	*	*	*	0.2	0.6	1.1	3.1	8.5	26.0	84.9	343.9
2000	18.3	*	*	*	0.3	0.5	1.3	2.8	8.4	25.0	86.1	351.7
1999	19.0	*	*	*	0.4	0.6	1.3	3.1	9.7	27.3	89.0	364.5
Non-Hispanic black, both sexes												
2002	13.0	*	*	*	0.6	1.7	3.7	10.4	26.6	52.4	125.1	342.4
2001	13.1	*	*	*	0.8	1.5	3.8	10.8	25.5	54.9	129.1	343.0
2000	12.7	*	*	*	0.7	1.4	3.7	10.1	25.6	53.6	123.6	349.8
1999	13.4	*	*	*	0.8	1.8	3.7	10.8	27.4	56.8	134.0	362.1
Non-Hispanic black, male												
2002	12.4	*	*	*	0.8	2.0	4.3	13.6	34.7	66.0	142.1	341.3
2001	12.7	*	*	*	1.1	1.9	4.3	13.4	34.6	67.0	161.1	359.2
2000	12.1	*	*	*	*	1.6	4.3	13.9	34.2	68.4	134.5	362.8
1999	13.3	*	*	*	1.0	2.0	4.1	13.8	37.0	72.0	165.4	413.6
Non-Hispanic black, female												
2002	13.5	*	*	*	*	1.5	3.2	7.7	20.1	42.8	115.4	342.8
2001	13.4	*	*	*	*	1.2	3.3	8.5	18.3	46.3	111.2	337.1
2000	13.3	*	*	*	0.8	1.2	3.3	6.8	18.7	43.2	117.4	345.0
1999	13.6	*	*	*	*	1.6	3.4	8.2	19.8	46.1	116.4	342.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All origins 3/, both sexes												
2002	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
All origins 3/, male												
2002	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.9	646.4
2000	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
All origins 3/, female												
2002	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9
Hispanic, both sexes												
2002	3.6	*	*	*	*	*	*	1.2	4.6	19.9	75.2	316.0
2001	3.7	*	*	*	*	*	0.4	1.3	5.6	18.8	73.0	350.1
2000	3.6	*	*	*	*	*	*	0.7	5.8	18.9	72.1	369.6
1999	3.7	*	*	*	*	*	*	1.0	5.3	19.5	73.5	377.4
Hispanic, male												
2002	3.0	*	*	*	*	*	*	1.1	5.8	23.1	85.8	296.2
2001	3.0	*	*	*	*	*	*	1.8	6.0	21.8	76.0	333.3
2000	2.9	*	*	*	*	*	*	*	7.7	23.0	74.2	380.9
1999	3.0	*	*	*	*	*	*	*	6.1	24.6	82.9	358.0
Hispanic, female												
2002	4.4	*	*	*	*	*	*	1.3	3.6	17.3	68.0	325.8
2001	4.5	*	*	*	*	*	*	*	5.2	16.4	71.0	358.4
2000	4.3	*	*	*	*	*	*	*	4.2	15.6	70.7	364.0
1999	4.4	*	*	*	*	*	*	*	4.6	15.5	67.3	386.8
Non-Hispanic 4/, both sexes												
2002	22.0	*	*	*	0.1	0.3	0.7	2.6	9.1	34.6	135.3	656.8
2001	22.4	*	*	*	0.1	0.3	0.7	2.5	9.6	35.2	138.8	685.6
2000	22.1	0.8	*	*	0.1	0.2	0.6	2.3	9.3	35.7	138.5	694.8
1999	21.8	*	*	*	0.1	0.2	0.6	2.2	9.1	36.3	139.4	691.0
Non-Hispanic 4/, male												
2002	17.3	*	*	*	*	0.4	0.9	3.3	10.9	40.2	151.8	634.7
2001	17.4	*	*	*	*	0.3	0.9	3.2	11.4	41.4	157.2	658.0
2000	17.2	*	*	*	0.1	0.2	0.8	2.9	11.0	42.1	156.9	683.3
1999	16.9	*	*	*	0.1	0.2	0.8	2.8	11.0	43.5	154.5	677.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Non-Hispanic 4/, female												
2002	26.5	*	*	*	*	0.2	0.5	1.9	7.5	29.9	124.3	666.4
2001	27.1	*	*	*	*	0.2	0.6	1.9	8.0	30.1	126.8	697.0
2000	26.7	*	*	*	*	0.2	0.5	1.7	7.7	30.5	126.5	699.5
1999	26.5	*	*	*	*	0.2	0.5	1.6	7.2	30.4	129.7	696.5
Non-Hispanic white, both sexes												
2002	24.7	*	*	*	*	0.2	0.5	2.0	8.2	33.3	135.9	677.3
2001	25.1	*	*	*	*	0.2	0.5	2.1	8.7	33.9	139.7	706.8
2000	24.7	*	*	*	*	0.1	0.5	1.9	8.4	34.3	139.2	714.3
1999	24.3	*	*	*	*	0.1	0.5	1.7	8.1	34.9	139.1	709.1
Non-Hispanic white, male												
2002	19.2	*	*	*	*	0.2	0.7	2.6	9.8	38.6	153.2	656.2
2001	19.3	*	*	*	*	0.2	0.7	2.7	10.5	40.0	158.1	679.1
2000	19.0	*	*	*	*	0.2	0.6	2.4	10.1	41.0	157.9	701.6
1999	18.6	*	*	*	*	*	0.6	2.2	10.1	42.2	154.7	690.8
Non-Hispanic white, female												
2002	30.0	*	*	*	*	0.2	0.4	1.4	6.7	28.8	124.2	686.4
2001	30.7	*	*	*	*	0.2	0.4	1.4	6.9	28.8	127.5	718.4
2000	30.2	*	*	*	*	*	0.3	1.3	6.8	28.7	126.9	719.4
1999	29.9	*	*	*	*	*	0.3	1.2	6.3	28.8	128.8	716.4
Non-Hispanic black, both sexes												
2002	14.2	*	*	*	*	0.9	1.9	7.4	19.0	54.1	158.3	534.5
2001	14.3	*	*	*	*	0.6	2.0	6.6	20.0	55.5	159.1	551.2
2000	14.1	*	*	*	*	0.5	1.7	5.9	19.0	56.2	156.3	564.8
1999	14.3	*	*	*	*	0.6	1.7	6.5	19.2	57.3	164.8	557.4
Non-Hispanic black, male												
2002	12.0	*	*	*	*	1.1	2.3	9.2	23.3	65.9	176.4	489.0
2001	12.0	*	*	*	*	0.9	2.3	7.6	22.6	66.3	185.4	547.5
2000	11.6	*	*	*	*	*	1.8	7.3	21.8	62.9	176.5	589.4
1999	12.2	*	*	*	*	*	1.9	8.6	23.0	66.4	180.4	609.7
Non-Hispanic black, female												
2002	16.3	*	*	*	*	*	1.5	5.7	15.5	45.8	148.1	551.8
2001	16.4	*	*	*	*	*	1.7	5.6	18.0	47.9	144.4	552.6
2000	16.3	*	*	*	*	*	1.6	4.7	16.8	51.5	145.0	555.6
1999	16.2	*	*	*	*	*	1.5	4.8	16.1	50.9	156.0	537.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congestive heart failure (I50.0)												
All origins 3/, both sexes												
2002	18.2	*	*	*	*	0.2	0.5	2.0	7.6	30.3	123.7	609.0
2001	18.6	*	*	*	0.1	0.2	0.5	2.1	8.0	31.1	126.6	636.1
2000	18.3	*	*	*	0.1	0.1	0.5	1.8	7.9	31.0	126.4	644.4
1999	18.2	*	*	*	0.1	0.1	0.5	1.7	7.4	31.9	127.3	642.7
All origins 3/, male												
2002	14.0	*	*	*	*	0.3	0.6	2.5	8.9	35.2	138.1	584.6
2001	14.2	*	*	*	*	0.2	0.6	2.6	9.3	36.2	141.2	609.3
2000	14.0	*	*	*	*	0.1	0.5	2.2	9.3	36.4	142.4	632.2
1999	13.9	*	*	*	*	0.1	0.6	2.2	8.9	38.0	140.5	628.2
All origins 3/, female												
2002	22.2	*	*	*	*	0.2	0.4	1.5	6.3	26.3	114.1	619.6
2001	22.8	*	*	*	*	0.1	0.4	1.6	6.8	26.8	117.0	647.4
2000	22.4	*	*	*	*	0.1	0.4	1.4	6.6	26.6	116.1	649.4
1999	22.4	*	*	*	*	0.1	0.4	1.3	6.1	26.9	118.7	648.5
Hispanic, both sexes												
2002	3.3	*	*	*	*	*	*	0.9	4.3	17.2	69.2	296.0
2001	3.4	*	*	*	*	*	*	1.2	4.9	17.3	68.2	323.3
2000	3.3	*	*	*	*	*	*	*	5.4	16.9	67.0	343.7
1999	3.4	*	*	*	*	*	*	0.7	4.9	17.9	68.7	359.2
Hispanic, male												
2002	2.6	*	*	*	*	*	*	*	5.2	20.3	76.7	274.4
2001	2.7	*	*	*	*	*	*	1.7	5.3	20.6	68.9	295.2
2000	2.6	*	*	*	*	*	*	*	7.0	20.7	67.4	342.6
1999	2.7	*	*	*	*	*	*	*	5.6	22.2	77.7	338.6
Hispanic, female												
2002	4.0	*	*	*	*	*	*	*	3.4	14.8	64.1	306.7
2001	4.2	*	*	*	*	*	*	*	4.5	14.6	67.7	337.0
2000	4.0	*	*	*	*	*	*	*	4.0	13.9	66.7	344.2
1999	4.1	*	*	*	*	*	*	*	4.4	14.6	62.8	369.1
Non-Hispanic 4/, both sexes												
2002	20.4	*	*	*	*	0.2	0.6	2.1	7.8	31.1	126.1	620.2
2001	20.8	*	*	*	0.1	0.2	0.6	2.1	8.2	31.9	129.0	646.7
2000	20.4	*	*	*	0.1	0.1	0.5	1.9	8.1	31.8	128.7	653.9
1999	20.2	*	*	*	0.1	0.1	0.5	1.8	7.6	32.6	129.4	651.2
Non-Hispanic 4/, male												
2002	15.9	*	*	*	*	0.3	0.7	2.6	9.2	36.1	140.8	597.3
2001	15.9	*	*	*	*	0.2	0.7	2.6	9.6	37.1	144.4	621.0
2000	15.7	*	*	*	*	0.2	0.6	2.3	9.4	37.2	145.4	643.2
1999	15.5	*	*	*	*	0.1	0.6	2.3	9.0	38.8	142.7	637.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congestive heart failure (I50.0)												
Non-Hispanic 4/, female												
2002	24.8	*	*	*	*	0.2	0.4	1.6	6.6	27.0	116.3	630.1
2001	25.4	*	*	*	*	0.2	0.5	1.7	6.9	27.6	118.9	657.3
2000	24.9	*	*	*	*	0.1	0.4	1.4	6.8	27.3	117.8	658.3
1999	24.7	*	*	*	*	0.1	0.4	1.3	6.2	27.6	120.7	656.6
Non-Hispanic white, both sexes												
2002	23.1	*	*	*	*	0.1	0.4	1.6	7.0	30.1	127.0	641.1
2001	23.4	*	*	*	*	0.2	0.4	1.7	7.4	30.8	130.2	667.8
2000	23.0	*	*	*	*	0.1	0.4	1.5	7.3	30.7	129.7	673.7
1999	22.6	*	*	*	*	*	0.4	1.3	6.8	31.4	129.4	669.4
Non-Hispanic white, male												
2002	17.7	*	*	*	*	0.2	0.5	2.0	8.2	34.6	142.7	619.6
2001	17.8	*	*	*	*	0.2	0.5	2.2	8.9	36.0	145.9	641.8
2000	17.5	*	*	*	*	*	0.5	1.9	8.7	36.5	146.8	662.2
1999	17.0	*	*	*	*	*	0.5	1.7	8.2	37.8	143.1	652.0
Non-Hispanic white, female												
2002	28.2	*	*	*	*	*	0.3	1.1	5.8	26.2	116.5	650.4
2001	28.8	*	*	*	*	*	0.3	1.3	6.0	26.4	119.8	678.6
2000	28.2	*	*	*	*	*	0.3	1.1	6.0	25.7	118.4	678.4
1999	27.9	*	*	*	*	*	0.3	0.9	5.4	26.1	120.4	676.3
Non-Hispanic black, both sexes												
2002	12.8	*	*	*	*	0.8	1.6	6.2	16.6	48.2	143.5	488.2
2001	12.8	*	*	*	*	0.4	1.7	5.6	16.9	49.9	143.8	509.4
2000	12.5	*	*	*	*	*	1.3	5.1	16.4	49.3	141.8	515.8
1999	12.8	*	*	*	*	0.5	1.3	5.7	16.1	51.2	149.5	514.4
Non-Hispanic black, male												
2002	10.6	*	*	*	*	1.1	1.9	7.6	20.2	60.6	157.1	437.6
2001	10.6	*	*	*	*	*	2.0	6.6	18.6	58.6	162.0	507.0
2000	10.2	*	*	*	*	*	1.4	6.4	18.4	53.6	157.4	535.4
1999	10.8	*	*	*	*	*	1.6	7.3	19.0	58.9	165.1	561.0
Non-Hispanic black, female												
2002	14.7	*	*	*	*	*	1.4	4.9	13.7	39.5	135.8	507.5
2001	14.9	*	*	*	*	*	1.4	4.8	15.6	43.7	133.6	510.3
2000	14.7	*	*	*	*	*	1.3	4.0	14.9	46.2	133.0	508.5
1999	14.7	*	*	*	*	*	1.1	4.3	13.9	45.8	140.8	497.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)												
All origins 3/, both sexes												
2002	1.4	*	*	*	*	*	0.1	0.5	1.2	3.4	9.1	36.1
2001	1.4	*	*	*	*	0.1	0.1	0.4	1.4	3.2	9.7	38.5
2000	1.5	0.6	*	*	*	*	0.1	0.4	1.2	3.8	9.6	40.5
1999	1.5	*	*	*	*	*	0.1	0.4	1.4	3.6	9.9	39.2
All origins 3/, male												
2002	1.3	*	*	*	*	*	0.2	0.7	1.6	4.0	10.9	36.8
2001	1.3	*	*	*	*	*	0.2	0.5	1.8	4.1	12.7	37.2
2000	1.3	*	*	*	*	*	0.2	0.5	1.5	4.7	11.3	40.0
1999	1.3	*	*	*	*	*	0.2	0.5	1.9	4.5	11.6	38.8
All origins 3/, female												
2002	1.5	*	*	*	*	*	*	0.3	0.9	2.9	7.9	35.7
2001	1.6	*	*	*	*	*	0.1	0.2	1.0	2.4	7.7	39.1
2000	1.7	*	*	*	*	*	*	0.3	0.9	3.1	8.5	40.6
1999	1.6	*	*	*	*	*	0.1	0.3	1.0	2.7	8.8	39.4
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	2.7	6.0	20.0
2001	0.3	*	*	*	*	*	*	*	*	*	4.9	26.9
2000	0.3	*	*	*	*	*	*	*	*	2.0	5.1	25.9
1999	0.3	*	*	*	*	*	*	*	*	*	4.8	18.2
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	9.0	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	38.1
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	19.1
2001	0.3	*	*	*	*	*	*	*	*	*	*	21.4
2000	0.3	*	*	*	*	*	*	*	*	*	*	19.8
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.6	*	*	*	*	*	0.1	0.5	1.3	3.5	9.2	36.6
2001	1.6	*	*	*	*	*	0.1	0.4	1.4	3.3	9.9	38.9
2000	1.6	*	*	*	*	*	0.1	0.4	1.2	3.9	9.8	40.9
1999	1.6	*	*	*	*	*	0.1	0.4	1.5	3.6	10.1	39.7
Non-Hispanic 4/, male												
2002	1.4	*	*	*	*	*	0.2	0.7	1.7	4.1	11.0	37.4
2001	1.5	*	*	*	*	*	0.2	0.5	1.8	4.2	12.9	37.0
2000	1.4	*	*	*	*	*	0.2	0.5	1.6	4.9	11.5	40.1
1999	1.4	*	*	*	*	*	0.2	0.6	2.0	4.6	11.8	39.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)												
Non-Hispanic 4/, female												
2002	1.7	*	*	*	*	*	*	0.3	0.9	2.9	8.0	36.3
2001	1.7	*	*	*	*	*	0.1	0.2	1.1	2.5	7.9	39.7
2000	1.8	*	*	*	*	*	*	0.3	0.9	3.1	8.7	41.3
1999	1.8	*	*	*	*	*	0.1	0.3	1.0	2.8	9.0	39.9
Non-Hispanic white, both sexes												
2002	1.7	*	*	*	*	*	0.1	0.4	1.2	3.2	8.9	36.2
2001	1.7	*	*	*	*	*	0.1	0.3	1.3	3.1	9.5	39.0
2000	1.7	*	*	*	*	*	0.1	0.4	1.1	3.7	9.5	40.6
1999	1.7	*	*	*	*	*	0.1	0.4	1.4	3.5	9.7	39.7
Non-Hispanic white, male												
2002	1.5	*	*	*	*	*	0.1	0.6	1.6	4.0	10.6	36.6
2001	1.5	*	*	*	*	*	0.1	0.5	1.7	4.0	12.2	37.3
2000	1.5	*	*	*	*	*	0.2	0.5	1.4	4.5	11.0	39.5
1999	1.5	*	*	*	*	*	0.1	0.5	1.8	4.4	11.6	38.8
Non-Hispanic white, female												
2002	1.8	*	*	*	*	*	*	0.2	0.9	2.5	7.8	36.0
2001	1.9	*	*	*	*	*	*	0.2	0.9	2.3	7.7	39.8
2000	2.0	*	*	*	*	*	*	0.2	0.8	3.0	8.5	41.1
1999	1.9	*	*	*	*	*	*	0.3	0.9	2.7	8.5	40.1
Non-Hispanic black, both sexes												
2002	1.5	*	*	*	*	*	*	1.2	2.4	5.9	14.8	46.3
2001	1.5	*	*	*	*	*	*	0.9	3.1	5.6	15.3	41.8
2000	1.5	*	*	*	*	*	0.4	0.8	2.6	6.9	14.6	49.0
1999	1.5	*	*	*	*	*	0.4	0.8	3.1	6.1	15.3	43.0
Non-Hispanic black, male												
2002	1.4	*	*	*	*	*	*	1.7	3.1	5.3	19.3	51.4
2001	1.5	*	*	*	*	*	*	1.0	3.9	7.7	23.4	40.4
2000	1.5	*	*	*	*	*	*	*	3.4	9.3	19.1	54.0
1999	1.3	*	*	*	*	*	*	1.3	4.1	7.5	15.3	48.7
Non-Hispanic black, female												
2002	1.6	*	*	*	*	*	*	*	1.8	6.3	12.3	44.3
2001	1.5	*	*	*	*	*	*	0.9	2.4	4.2	10.8	42.3
2000	1.5	*	*	*	*	*	*	*	2.0	5.3	12.0	47.1
1999	1.6	*	*	*	*	*	*	*	2.2	5.1	15.2	40.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
All origins 3/, both sexes												
2002	0.3	0.8	0.1	*	0.2	0.2	0.3	0.3	0.3	0.5	0.9	3.2
2001	0.3	0.6	0.3	0.1	0.1	0.2	0.3	0.3	0.2	0.4	0.5	2.0
2000	0.2	0.6	0.1	0.1	0.1	0.2	0.3	0.2	0.3	0.3	0.6	2.4
1999	0.3	0.7	0.2	0.1	0.2	0.2	0.3	0.3	0.2	0.3	0.5	2.4
All origins 3/, male												
2002	0.3	*	*	*	0.1	0.2	0.3	0.3	0.3	0.6	0.9	3.8
2001	0.3	*	0.3	*	0.1	0.2	0.3	0.3	0.2	0.5	0.8	2.2
2000	0.2	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	0.7	2.6
1999	0.3	*	*	*	0.2	0.3	0.3	0.3	0.2	0.4	0.6	2.1
All origins 3/, female												
2002	0.3	*	*	*	0.2	0.1	0.3	0.3	0.3	0.4	0.9	3.0
2001	0.2	*	*	*	0.1	0.2	0.2	0.2	0.2	0.3	0.4	1.9
2000	0.2	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3	0.5	2.4
1999	0.2	*	*	*	0.2	0.1	0.2	0.3	0.2	0.2	0.4	2.5
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	0.8	*	*	0.2	0.1	0.3	0.3	0.3	0.5	0.9	3.2
2001	0.3	0.6	0.3	0.1	0.1	0.2	0.3	0.3	0.2	0.4	0.5	2.1
2000	0.3	*	0.2	*	0.1	0.2	0.3	0.2	0.3	0.3	0.6	2.5
1999	0.3	0.7	0.2	*	0.2	0.2	0.3	0.3	0.2	0.3	0.5	2.4
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.2	0.2	0.4	0.3	0.3	0.6	0.9	3.8
2001	0.3	*	0.3	*	0.1	0.2	0.3	0.3	0.3	0.5	0.8	2.2
2000	0.2	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	0.7	2.7
1999	0.3	*	*	*	0.2	0.2	0.3	0.3	0.2	0.4	0.6	2.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	0.2	*	0.3	0.3	0.4	0.4	0.9	3.0
2001	0.3	*	*	*	*	0.2	0.3	0.3	0.2	0.3	0.4	2.0
2000	0.3	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3	0.5	2.4
1999	0.3	*	*	*	0.2	0.1	0.2	0.3	0.2	0.2	0.4	2.5
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	0.1	0.2	0.3	0.3	0.3	0.5	0.9	3.2
2001	0.3	*	*	*	0.1	0.2	0.3	0.2	0.2	0.4	0.5	2.1
2000	0.3	*	*	*	0.1	0.2	0.2	0.2	0.3	0.3	0.5	2.6
1999	0.2	*	*	*	0.2	0.1	0.2	0.2	0.2	0.3	0.5	2.4
Non-Hispanic white, male												
2002	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.6	0.9	4.0
2001	0.3	*	*	*	*	*	0.3	0.3	0.2	0.5	0.7	2.2
2000	0.2	*	*	*	*	0.2	0.3	*	0.3	0.3	0.6	2.8
1999	0.2	*	*	*	0.2	0.2	0.3	0.3	*	0.4	0.6	2.0
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	0.2	0.3	0.4	0.4	0.9	2.8
2001	0.2	*	*	*	*	0.2	0.2	0.2	0.2	0.3	0.4	2.0
2000	0.3	*	*	*	*	0.2	0.2	0.2	0.3	0.3	0.5	2.5
1999	0.2	*	*	*	0.2	*	0.2	0.2	0.2	0.3	0.5	2.6
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	0.7	0.6	*	*	*	*
2001	0.4	*	*	*	*	0.6	0.5	0.6	*	*	*	*
2000	0.4	*	*	*	*	*	0.5	0.5	*	*	*	*
1999	0.4	*	*	*	*	0.5	0.6	*	*	*	*	*
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	0.8	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)												
All origins 3/, both sexes												
2002	3.3	0.6	*	*	0.2	0.4	1.1	1.9	3.4	7.8	20.0	65.0
2001	3.2	*	*	0.1	0.2	0.4	1.0	1.8	3.4	7.3	18.9	66.3
2000	3.1	0.8	*	*	0.2	0.4	0.9	1.8	3.7	7.5	19.3	64.6
1999	2.9	0.8	*	*	0.2	0.4	0.9	1.6	3.5	6.9	17.8	61.5
All origins 3/, male												
2002	3.3	*	*	*	0.4	0.6	1.5	2.7	4.6	10.6	23.6	66.7
2001	3.2	*	*	*	0.3	0.7	1.5	2.6	4.7	9.9	22.1	66.4
2000	3.1	*	*	*	0.4	0.6	1.3	2.5	5.3	9.6	22.4	69.0
1999	3.0	*	*	*	0.3	0.6	1.3	2.3	4.9	9.4	21.8	64.3
All origins 3/, female												
2002	3.2	*	*	*	*	0.2	0.6	1.2	2.2	5.4	17.5	64.3
2001	3.1	*	*	*	*	0.2	0.5	1.1	2.2	5.2	16.8	66.3
2000	3.1	*	*	*	*	0.2	0.5	1.0	2.2	5.7	17.3	62.7
1999	2.8	*	*	*	*	0.2	0.5	0.9	2.1	4.8	15.2	60.4
Hispanic, both sexes												
2002	1.0	*	*	*	*	*	0.6	0.7	1.3	4.9	15.5	58.2
2001	0.9	*	*	*	*	*	0.4	1.0	2.1	3.9	12.1	48.1
2000	0.9	*	*	*	*	*	0.5	0.8	1.4	4.3	15.8	46.4
1999	0.9	*	*	*	*	0.3	0.6	1.1	2.0	4.8	12.5	39.3
Hispanic, male												
2002	1.1	*	*	*	*	*	0.9	*	*	7.5	19.3	60.2
2001	1.0	*	*	*	*	*	*	1.6	3.0	4.4	17.0	43.8
2000	1.0	*	*	*	*	*	0.8	*	*	5.9	17.7	60.5
1999	1.1	*	*	*	*	0.6	0.9	1.8	*	7.2	16.2	49.6
Hispanic, female												
2002	0.9	*	*	*	*	*	*	*	*	*	12.9	57.2
2001	0.8	*	*	*	*	*	*	*	*	3.5	8.8	50.3
2000	0.8	*	*	*	*	*	*	*	*	*	14.5	39.6
1999	0.7	*	*	*	*	*	*	*	*	*	10.1	34.3
Non-Hispanic 4/, both sexes												
2002	3.6	0.6	*	*	0.2	0.4	1.1	2.0	3.5	8.0	20.1	65.2
2001	3.5	*	*	*	0.2	0.5	1.1	1.9	3.5	7.5	19.2	66.8
2000	3.4	0.9	*	*	0.2	0.4	0.9	1.8	3.9	7.7	19.4	65.1
1999	3.2	0.7	*	*	0.2	0.4	0.9	1.6	3.6	7.0	18.0	62.2
Non-Hispanic 4/, male												
2002	3.7	*	*	*	0.4	0.6	1.6	2.8	4.8	10.8	23.8	66.8
2001	3.5	*	*	*	0.3	0.7	1.6	2.7	4.8	10.2	22.3	67.4
2000	3.5	*	*	*	0.4	0.7	1.3	2.6	5.5	9.8	22.5	69.2
1999	3.2	*	*	*	0.3	0.6	1.3	2.3	5.1	9.5	22.0	64.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)												
Non-Hispanic 4/, female												
2002	3.5	*	*	*	*	0.2	0.7	1.3	2.3	5.6	17.7	64.4
2001	3.5	*	*	*	*	0.2	0.6	1.1	2.3	5.3	17.1	66.6
2000	3.4	*	*	*	*	0.2	0.5	1.0	2.3	5.9	17.4	63.4
1999	3.1	*	*	*	*	0.2	0.5	1.0	2.1	4.9	15.3	61.2
Non-Hispanic white, both sexes												
2002	3.8	*	*	*	0.2	0.3	1.0	1.7	3.2	7.6	20.1	64.6
2001	3.6	*	*	*	0.2	0.4	0.9	1.6	3.2	7.2	18.6	66.2
2000	3.6	*	*	*	0.1	0.4	0.8	1.6	3.6	7.1	18.8	65.1
1999	3.3	*	*	*	0.2	0.3	0.8	1.4	3.1	6.6	17.4	62.0
Non-Hispanic white, male												
2002	3.8	*	*	*	0.3	0.5	1.4	2.5	4.4	10.5	23.4	67.1
2001	3.6	*	*	*	0.3	0.6	1.5	2.4	4.6	9.8	21.6	67.0
2000	3.5	*	*	*	0.2	0.6	1.2	2.4	5.2	9.2	21.7	69.3
1999	3.3	*	*	*	0.3	0.5	1.1	2.1	4.7	8.9	21.4	64.8
Non-Hispanic white, female												
2002	3.8	*	*	*	*	0.2	0.6	1.0	2.1	5.3	17.8	63.5
2001	3.7	*	*	*	*	0.2	0.4	0.8	2.0	4.9	16.7	65.8
2000	3.6	*	*	*	*	*	0.4	0.8	2.1	5.4	17.0	63.5
1999	3.3	*	*	*	*	*	0.4	0.7	1.7	4.6	14.9	60.9
Non-Hispanic black, both sexes												
2002	3.6	*	*	*	0.5	0.8	1.9	4.6	6.8	12.8	25.1	81.4
2001	3.6	*	*	*	0.4	1.0	2.1	4.2	6.4	12.1	27.2	84.9
2000	3.6	*	*	*	0.7	0.9	2.0	3.8	7.3	14.8	28.7	74.7
1999	3.4	*	*	*	0.4	0.9	2.1	3.5	8.0	12.4	25.6	69.8
Non-Hispanic black, male												
2002	3.9	*	*	*	0.8	1.3	2.5	6.0	9.5	17.1	33.1	77.7
2001	3.7	*	*	*	*	1.5	2.7	5.2	8.5	15.6	33.0	85.5
2000	4.0	*	*	*	1.1	1.3	2.6	5.5	10.2	19.5	36.4	81.0
1999	3.7	*	*	*	*	1.5	2.9	4.5	10.8	17.7	32.2	73.7
Non-Hispanic black, female												
2002	3.3	*	*	*	*	*	1.3	3.3	4.6	9.7	20.6	82.8
2001	3.5	*	*	*	*	*	1.6	3.2	4.6	9.6	23.9	84.7
2000	3.3	*	*	*	*	*	1.5	2.3	5.0	11.6	24.3	72.4
1999	3.1	*	*	*	*	*	1.4	2.7	5.7	8.6	21.9	68.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All origins 3/, both sexes												
2002	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	444.0	1,500.3
2000	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
All origins 3/, male												
2002	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.3
2000	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
All origins 3/, female												
2002	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.7	1,550.5
2000	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0
Hispanic, both sexes												
2002	16.6	2.4	*	0.3	0.4	1.3	4.4	15.3	35.9	102.9	319.9	898.0
2001	17.4	*	*	*	0.6	1.4	5.1	15.9	35.5	106.2	342.1	1,044.8
2000	17.5	*	*	*	0.5	1.7	5.2	15.2	39.8	113.2	344.1	1,082.9
1999	17.4	*	*	*	0.5	1.6	4.9	15.7	37.7	116.9	348.3	1,077.6
Hispanic, male												
2002	15.0	*	*	*	*	1.5	5.1	18.6	45.0	124.6	338.5	856.7
2001	15.7	*	*	*	0.7	1.6	5.4	18.7	43.7	127.2	386.3	1,007.5
2000	15.8	*	*	*	*	1.9	5.5	18.1	48.8	136.1	392.9	1,029.9
1999	16.1	*	*	*	0.6	1.8	5.7	19.5	46.1	140.8	395.2	1,136.6
Hispanic, female												
2002	18.4	*	*	*	*	1.0	3.7	12.0	27.6	85.6	307.2	918.5
2001	19.2	*	*	*	*	1.2	4.7	13.1	28.2	89.6	312.2	1,063.0
2000	19.4	*	*	*	*	1.4	4.9	12.4	31.9	95.2	311.3	1,108.9
1999	18.8	*	*	*	*	1.4	4.0	11.9	30.3	98.1	317.1	1,049.2
Non-Hispanic 4/, both sexes												
2002	62.4	3.0	0.3	0.2	0.4	1.5	5.5	15.0	37.2	121.3	435.5	1,465.0
2001	63.2	2.8	0.3	0.2	0.5	1.5	5.6	14.9	38.0	124.3	447.7	1,514.0
2000	65.4	3.6	0.3	0.2	0.5	1.5	5.8	15.9	40.9	129.2	465.0	1,604.5
1999	65.7	2.9	0.3	0.2	0.5	1.4	5.8	15.1	40.6	131.2	473.8	1,630.4
Non-Hispanic 4/, male												
2002	48.9	3.5	0.4	0.2	0.5	1.4	5.3	16.4	42.3	135.4	450.1	1,336.0
2001	49.7	3.3	0.3	0.2	0.5	1.5	5.7	16.4	43.2	140.9	469.8	1,392.9
2000	51.5	4.2	*	0.2	0.5	1.5	5.8	17.3	46.8	145.1	493.7	1,499.9
1999	51.5	3.7	0.3	0.2	0.5	1.4	5.8	16.4	46.3	147.3	503.6	1,524.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Non-Hispanic 4/, female												
2002	75.3	2.5	*	0.2	0.3	1.5	5.8	13.7	32.4	109.5	425.8	1,520.7
2001	76.1	2.4	0.4	0.2	0.5	1.5	5.5	13.5	33.2	110.4	433.2	1,564.4
2000	78.7	2.8	0.4	0.2	0.5	1.5	5.8	14.6	35.5	116.1	446.4	1,646.8
1999	79.2	2.0	*	0.2	0.5	1.4	5.8	13.9	35.4	118.1	454.4	1,672.7
Non-Hispanic white, both sexes												
2002	67.0	2.2	0.3	0.2	0.4	1.2	4.1	11.1	30.8	110.8	427.5	1,479.2
2001	67.6	1.9	0.3	0.2	0.4	1.3	4.2	10.8	31.2	113.0	439.6	1,525.0
2000	70.0	2.6	0.3	0.2	0.5	1.2	4.3	12.0	34.2	119.3	455.7	1,616.8
1999	70.4	2.4	0.3	0.2	0.4	1.1	4.3	11.2	33.4	120.7	465.1	1,648.0
Non-Hispanic white, male												
2002	51.3	2.7	*	0.2	0.5	1.1	4.0	12.1	34.5	123.2	441.1	1,345.9
2001	52.0	2.2	*	0.2	0.4	1.3	4.3	11.9	35.1	128.0	460.5	1,399.0
2000	53.9	3.3	*	0.2	0.5	1.2	4.3	13.0	38.7	133.1	482.3	1,505.9
1999	53.9	3.1	*	0.2	0.4	1.1	4.4	12.0	38.0	135.0	492.8	1,535.1
Non-Hispanic white, female												
2002	82.1	*	*	0.2	0.3	1.2	4.3	10.1	27.2	100.2	418.4	1,536.7
2001	82.6	*	*	*	0.3	1.3	4.1	9.8	27.4	100.3	425.7	1,577.4
2000	85.5	1.9	*	0.2	0.4	1.2	4.3	10.9	29.9	107.6	438.3	1,661.6
1999	86.1	*	*	*	0.4	1.1	4.3	10.5	29.1	108.8	447.0	1,693.0
Non-Hispanic black, both sexes												
2002	51.7	5.6	*	0.4	0.6	3.2	13.3	41.6	89.7	218.7	558.8	1,398.0
2001	52.9	6.9	*	*	0.9	3.0	13.6	43.5	92.8	226.0	567.2	1,465.5
2000	54.3	8.0	*	*	0.7	3.0	14.9	44.3	95.2	226.3	593.8	1,545.8
1999	53.9	4.7	*	*	0.7	2.9	14.7	42.2	97.7	235.7	596.4	1,501.5
Non-Hispanic black, male												
2002	45.1	*	*	*	0.7	2.9	12.7	47.5	112.7	267.3	595.2	1,266.8
2001	46.2	8.3	*	*	0.7	3.1	13.5	49.9	113.9	273.5	620.5	1,359.4
2000	47.7	9.2	*	*	*	2.8	15.1	50.4	116.9	272.1	664.9	1,479.5
1999	47.5	*	*	*	*	2.9	15.0	49.6	119.5	279.1	656.7	1,455.9
Non-Hispanic black, female												
2002	57.7	*	*	*	*	3.5	13.8	36.6	71.3	184.2	538.2	1,448.1
2001	59.0	*	*	*	1.1	2.9	13.6	38.1	75.9	192.5	537.2	1,504.8
2000	60.3	*	*	*	0.8	3.2	14.8	39.2	77.8	194.1	553.8	1,570.4
1999	59.8	*	*	*	0.8	2.8	14.4	35.9	80.2	205.1	562.7	1,518.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Subarachnoid hemorrhage (I60)												
All origins 3/, both sexes												
2002	2.1	*	*	0.0	0.1	0.5	1.8	3.1	4.2	5.8	8.9	12.2
2001	2.2	*	*	*	0.2	0.5	1.8	3.2	4.2	5.8	8.8	13.2
2000	2.3	*	*	0.1	0.2	0.6	1.9	3.5	4.9	5.8	9.4	14.1
1999	2.3	*	*	*	0.2	0.6	2.1	3.5	4.7	6.2	9.4	13.1
All origins 3/, male												
2002	1.5	*	*	*	0.2	0.4	1.4	2.3	3.3	4.4	5.8	9.1
2001	1.5	*	*	*	0.2	0.5	1.4	2.6	3.2	4.3	6.4	9.5
2000	1.6	*	*	*	0.2	0.6	1.5	2.7	3.5	4.2	6.8	11.2
1999	1.6	*	*	*	0.2	0.6	1.8	2.8	3.4	4.3	6.7	9.0
All origins 3/, female												
2002	2.8	*	*	*	0.1	0.6	2.1	3.8	5.1	6.9	11.0	13.6
2001	2.8	*	*	*	0.2	0.5	2.1	3.7	5.2	7.0	10.3	14.8
2000	3.0	*	*	*	0.2	0.6	2.3	4.3	6.2	7.0	11.2	15.3
1999	3.0	*	*	*	0.2	0.6	2.4	4.1	5.9	7.7	11.1	14.8
Hispanic, both sexes												
2002	1.3	*	*	*	*	0.6	1.6	3.3	5.0	6.6	9.3	*
2001	1.4	*	*	*	*	0.6	1.6	3.6	5.4	7.0	9.7	16.3
2000	1.5	*	*	*	*	0.8	1.8	4.0	5.0	6.8	11.1	15.9
1999	1.6	*	*	*	*	0.6	2.0	4.1	5.1	7.3	12.5	23.2
Hispanic, male												
2002	1.1	*	*	*	*	0.6	1.7	3.0	4.4	4.8	*	*
2001	1.1	*	*	*	*	*	1.3	3.1	4.3	5.4	*	*
2000	1.1	*	*	*	*	0.9	1.5	2.9	4.1	*	10.8	*
1999	1.2	*	*	*	*	0.8	2.0	3.9	4.3	4.4	*	*
Hispanic, female												
2002	1.6	*	*	*	*	*	1.5	3.5	5.5	8.1	11.8	*
2001	1.8	*	*	*	*	0.6	1.8	4.1	6.3	8.3	11.5	20.5
2000	1.9	*	*	*	*	0.7	2.1	5.0	5.7	9.0	11.2	19.8
1999	1.9	*	*	*	*	*	2.0	4.3	5.9	9.6	15.6	32.2
Non-Hispanic 4/, both sexes												
2002	2.3	*	*	*	0.1	0.5	1.8	3.1	4.2	5.7	8.9	12.3
2001	2.3	*	*	*	0.1	0.5	1.8	3.1	4.1	5.7	8.7	13.1
2000	2.4	*	*	*	0.2	0.6	1.9	3.5	4.8	5.7	9.4	14.1
1999	2.4	*	*	*	0.1	0.6	2.1	3.4	4.7	6.1	9.3	12.7
Non-Hispanic 4/, male												
2002	1.6	*	*	*	0.2	0.4	1.4	2.3	3.2	4.4	5.8	9.2
2001	1.6	*	*	*	0.1	0.6	1.4	2.5	3.1	4.2	6.3	9.5
2000	1.7	*	*	*	0.2	0.5	1.5	2.7	3.4	4.2	6.7	11.4
1999	1.7	*	*	*	0.1	0.5	1.7	2.7	3.4	4.3	6.6	9.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Subarachnoid hemorrhage (I60)												
Non-Hispanic 4/, female												
2002	2.9	*	*	*	*	0.6	2.2	3.8	5.0	6.8	10.9	13.6
2001	2.9	*	*	*	0.2	0.5	2.1	3.7	5.1	6.9	10.3	14.5
2000	3.1	*	*	*	0.2	0.6	2.3	4.2	6.2	6.9	11.1	15.1
1999	3.1	*	*	*	0.2	0.6	2.5	4.1	5.9	7.6	11.0	14.1
Non-Hispanic white, both sexes												
2002	2.2	*	*	*	0.1	0.4	1.6	2.7	3.9	5.4	8.7	12.0
2001	2.2	*	*	*	0.1	0.5	1.5	2.7	3.8	5.3	8.5	12.6
2000	2.4	*	*	*	0.2	0.5	1.6	3.1	4.6	5.3	9.1	13.5
1999	2.4	*	*	*	0.1	0.5	1.8	3.0	4.3	5.9	9.1	12.2
Non-Hispanic white, male												
2002	1.6	*	*	*	0.2	0.3	1.2	2.1	3.0	4.2	5.7	9.2
2001	1.6	*	*	*	*	0.5	1.3	2.2	2.8	4.0	6.4	9.2
2000	1.7	*	*	*	0.2	0.5	1.3	2.5	3.2	3.9	6.7	12.1
1999	1.7	*	*	*	*	0.5	1.6	2.4	3.2	4.1	6.8	9.1
Non-Hispanic white, female												
2002	2.9	*	*	*	*	0.5	1.9	3.3	4.7	6.4	10.6	13.2
2001	2.8	*	*	*	*	0.4	1.7	3.2	4.8	6.3	10.0	14.0
2000	3.1	*	*	*	*	0.5	1.9	3.7	6.0	6.4	10.7	14.1
1999	3.1	*	*	*	0.2	0.5	2.0	3.5	5.4	7.5	10.6	13.4
Non-Hispanic black, both sexes												
2002	2.4	*	*	*	*	1.0	3.3	5.6	5.9	6.4	8.7	13.0
2001	2.6	*	*	*	*	0.9	3.5	5.9	6.2	7.4	8.6	14.0
2000	2.7	*	*	*	*	0.9	3.7	6.2	6.2	8.0	10.8	17.5
1999	2.7	*	*	*	*	1.0	3.9	6.2	6.8	7.0	9.3	14.5
Non-Hispanic black, male												
2002	1.6	*	*	*	*	*	2.2	3.3	5.3	6.1	5.9	*
2001	1.8	*	*	*	*	0.9	2.6	4.7	5.1	4.9	6.3	*
2000	1.7	*	*	*	*	*	2.7	4.4	4.2	6.3	7.1	*
1999	1.8	*	*	*	*	0.8	2.8	5.0	5.0	5.4	*	*
Non-Hispanic black, female												
2002	3.2	*	*	*	*	1.3	4.2	7.5	6.4	6.5	10.3	15.9
2001	3.3	*	*	*	*	0.9	4.3	6.9	7.2	9.2	9.9	16.7
2000	3.6	*	*	*	*	1.3	4.6	7.8	7.8	9.3	12.9	23.1
1999	3.4	*	*	*	*	1.1	4.8	7.3	8.2	8.1	11.6	18.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)												
All origins 3/, both sexes												
2002	9.4	1.0	*	0.1	0.2	0.5	2.1	6.0	11.7	28.1	72.3	127.2
2001	9.4	1.0	*	0.1	0.2	0.6	2.3	6.2	11.8	27.9	71.7	127.9
2000	9.7	1.1	*	0.1	0.2	0.5	2.3	6.5	12.9	29.3	74.1	137.9
1999	9.1	1.1	*	0.1	0.2	0.5	2.2	5.9	12.3	28.3	71.9	122.3
All origins 3/, male												
2002	8.7	1.1	*	*	0.2	0.5	2.3	7.3	13.7	32.3	77.2	135.0
2001	8.8	1.2	*	*	0.3	0.6	2.6	7.5	13.9	32.0	78.6	134.2
2000	9.1	1.3	*	*	0.2	0.6	2.7	7.8	15.9	33.1	79.0	149.6
1999	8.5	1.3	*	*	0.2	0.5	2.6	7.0	14.7	32.0	75.9	130.7
All origins 3/, female												
2002	10.0	1.0	*	*	0.1	0.4	1.8	4.8	9.8	24.6	69.0	123.8
2001	9.9	*	*	*	0.2	0.5	1.9	5.1	10.0	24.6	67.2	125.2
2000	10.4	*	*	*	0.2	0.4	2.0	5.3	10.2	26.1	70.9	133.1
1999	9.8	*	*	*	0.1	0.4	1.8	4.9	10.1	25.2	69.2	118.9
Hispanic, both sexes												
2002	3.9	*	*	*	*	0.4	1.8	6.9	12.5	24.8	60.6	98.7
2001	4.0	*	*	*	*	0.6	2.3	7.2	12.6	25.4	60.2	107.5
2000	4.0	*	*	*	*	0.6	2.3	6.5	15.6	26.8	61.8	106.8
1999	4.0	*	*	*	*	0.6	2.0	7.4	13.8	29.5	63.3	98.2
Hispanic, male												
2002	4.0	*	*	*	*	*	2.3	9.7	16.5	30.4	60.3	90.4
2001	4.2	*	*	*	*	0.9	2.9	9.5	16.7	29.0	65.7	112.4
2000	4.2	*	*	*	*	0.8	3.1	9.0	21.8	31.7	62.0	112.9
1999	4.4	*	*	*	*	0.7	2.7	10.3	19.0	37.3	67.8	107.8
Hispanic, female												
2002	3.8	*	*	*	*	*	1.3	4.0	8.9	20.4	60.7	102.8
2001	3.8	*	*	*	*	*	1.6	5.0	9.0	22.7	56.5	105.2
2000	3.7	*	*	*	*	*	1.5	4.0	10.0	22.9	61.7	103.9
1999	3.6	*	*	*	*	*	1.3	4.5	9.2	23.4	60.4	93.6
Non-Hispanic 4/, both sexes												
2002	10.2	0.9	*	0.1	0.1	0.5	2.1	5.9	11.6	28.3	72.7	128.1
2001	10.1	0.9	*	0.1	0.2	0.6	2.3	6.1	11.7	28.0	72.1	128.3
2000	10.5	1.2	*	0.1	0.2	0.5	2.3	6.5	12.7	29.4	74.4	138.9
1999	9.8	0.9	*	*	0.2	0.4	2.2	5.7	12.1	28.1	72.1	122.7
Non-Hispanic 4/, male												
2002	9.5	*	*	*	0.2	0.6	2.3	7.0	13.5	32.3	77.9	136.6
2001	9.5	*	*	*	0.2	0.6	2.5	7.2	13.6	32.2	79.2	134.9
2000	9.8	1.5	*	*	0.2	0.6	2.6	7.7	15.3	33.1	79.5	151.0
1999	9.0	*	*	*	0.2	0.5	2.5	6.6	14.3	31.6	76.3	131.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)												
Non-Hispanic 4/, female												
2002	10.8	*	*	*	*	0.5	1.9	4.9	9.9	24.9	69.2	124.4
2001	10.7	*	*	*	0.2	0.6	2.0	5.0	10.0	24.6	67.5	125.6
2000	11.2	*	*	*	0.2	0.4	2.1	5.4	10.2	26.3	71.1	134.0
1999	10.5	*	*	*	0.1	0.4	1.8	4.9	10.1	25.2	69.4	119.5
Non-Hispanic white, both sexes												
2002	10.4	*	*	*	0.1	0.4	1.3	3.8	9.6	26.6	72.8	130.3
2001	10.3	*	*	*	0.2	0.4	1.5	3.9	9.5	26.1	72.1	129.3
2000	10.7	*	*	*	0.2	0.4	1.5	4.4	10.3	28.0	74.2	140.4
1999	10.0	*	*	*	0.1	0.3	1.4	3.7	9.8	26.9	71.9	123.7
Non-Hispanic white, male												
2002	9.5	*	*	*	0.2	0.4	1.4	4.5	11.0	30.5	78.6	138.7
2001	9.4	*	*	*	0.2	0.4	1.7	4.6	11.2	29.9	79.2	135.6
2000	9.7	*	*	*	0.2	0.4	1.6	5.2	12.2	31.2	79.5	151.7
1999	8.9	*	*	*	*	0.3	1.6	4.3	11.6	30.4	75.5	132.1
Non-Hispanic white, female												
2002	11.3	*	*	*	*	0.3	1.2	3.2	8.2	23.3	68.9	126.7
2001	11.1	*	*	*	*	0.4	1.4	3.1	8.0	22.8	67.4	126.7
2000	11.7	*	*	*	*	0.3	1.3	3.5	8.5	25.3	70.8	135.8
1999	11.0	*	*	*	*	0.3	1.2	3.2	8.1	23.9	69.5	120.3
Non-Hispanic black, both sexes												
2002	10.0	*	*	*	*	1.2	6.2	19.3	26.3	39.7	69.4	97.4
2001	10.4	*	*	*	0.4	1.4	6.1	21.1	27.2	40.7	69.5	106.8
2000	10.6	*	*	*	*	1.2	7.3	21.5	29.9	39.4	69.4	118.0
1999	9.8	*	*	*	*	1.1	7.0	19.1	28.3	37.2	68.9	96.7
Non-Hispanic black, male												
2002	10.4	*	*	*	*	1.4	7.0	24.5	33.8	48.2	68.6	101.7
2001	10.7	*	*	*	*	1.5	6.9	25.9	32.8	50.3	72.3	113.2
2000	11.2	*	*	*	*	1.5	8.5	26.2	40.8	47.2	73.7	132.7
1999	10.1	*	*	*	*	1.3	8.4	23.6	35.5	40.7	74.7	108.2
Non-Hispanic black, female												
2002	9.7	*	*	*	*	1.1	5.5	14.9	20.4	33.8	69.8	95.7
2001	10.1	*	*	*	*	1.3	5.4	17.0	22.7	33.9	67.9	104.4
2000	10.1	*	*	*	*	1.0	6.2	17.4	21.2	33.9	67.0	112.5
1999	9.5	*	*	*	*	0.9	5.8	15.2	22.5	34.8	65.7	92.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebral infarction (I63)												
All origins 3/, both sexes												
2002	4.3	*	*	*	*	0.1	0.2	1.0	2.5	8.6	32.2	118.3
2001	4.4	*	*	*	*	0.1	0.3	0.9	2.4	9.4	33.9	124.8
2000	4.9	*	*	*	*	0.1	0.3	0.9	2.9	9.6	38.1	141.0
1999	5.3	*	*	*	*	0.1	0.3	0.9	3.1	11.3	41.2	153.1
All origins 3/, male												
2002	3.4	*	*	*	*	0.1	0.2	1.2	3.1	9.6	35.4	108.7
2001	3.5	*	*	*	*	0.1	0.3	1.1	3.1	11.1	37.8	111.3
2000	3.9	*	*	*	*	*	0.4	1.1	3.6	11.4	42.3	133.3
1999	4.2	*	*	*	*	0.1	0.4	1.2	3.8	13.5	45.0	144.6
All origins 3/, female												
2002	5.1	*	*	*	*	0.1	0.2	0.7	1.9	7.7	30.1	122.4
2001	5.3	*	*	*	*	*	0.3	0.8	1.8	8.0	31.3	130.4
2000	5.8	*	*	*	*	0.1	0.3	0.8	2.2	8.1	35.3	144.2
1999	6.3	*	*	*	*	*	0.3	0.7	2.5	9.6	38.8	156.6
Hispanic, both sexes												
2002	1.1	*	*	*	*	*	*	0.8	1.6	7.1	19.8	68.7
2001	1.2	*	*	*	*	*	*	0.8	1.8	8.2	23.4	89.4
2000	1.3	*	*	*	*	*	*	0.8	2.2	8.5	26.7	84.9
1999	1.3	*	*	*	*	*	*	0.7	2.0	9.6	26.2	100.3
Hispanic, male												
2002	0.9	*	*	*	*	*	*	*	2.8	7.5	20.1	71.9
2001	1.1	*	*	*	*	*	*	*	2.6	12.2	25.5	66.7
2000	1.2	*	*	*	*	*	*	*	3.4	10.6	32.9	92.7
1999	1.1	*	*	*	*	*	*	*	2.6	11.9	27.1	105.7
Hispanic, female												
2002	1.2	*	*	*	*	*	*	*	*	6.8	19.7	67.1
2001	1.4	*	*	*	*	*	*	*	*	5.0	22.1	100.5
2000	1.3	*	*	*	*	*	*	*	*	7.0	22.5	81.1
1999	1.5	*	*	*	*	*	*	*	*	7.7	25.7	97.7
Non-Hispanic 4/, both sexes												
2002	4.8	*	*	*	*	0.1	0.3	1.0	2.5	8.7	32.8	120.1
2001	4.9	*	*	*	*	0.1	0.3	0.9	2.5	9.4	34.2	126.0
2000	5.4	*	*	*	*	0.1	0.4	0.9	3.0	9.6	38.4	142.7
1999	5.8	*	*	*	*	0.1	0.4	1.0	3.2	11.4	41.8	154.5
Non-Hispanic 4/, male												
2002	3.8	*	*	*	*	0.1	0.2	1.3	3.1	9.7	36.1	110.1
2001	3.9	*	*	*	*	0.1	0.3	1.1	3.1	11.0	38.2	113.2
2000	4.3	*	*	*	*	*	0.4	1.1	3.7	11.3	42.6	134.6
1999	4.7	*	*	*	*	*	0.4	1.2	3.8	13.5	45.6	145.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebral infarction (I63)												
Non-Hispanic 4/, female												
2002	5.7	*	*	*	*	0.1	0.3	0.7	2.0	7.8	30.6	124.4
2001	5.9	*	*	*	*	*	0.3	0.8	1.9	8.1	31.7	131.3
2000	6.4	*	*	*	*	0.1	0.3	0.8	2.3	8.2	35.7	146.0
1999	7.0	*	*	*	*	*	0.3	0.7	2.6	9.7	39.3	158.0
Non-Hispanic white, both sexes												
2002	5.2	*	*	*	*	0.1	0.2	0.8	2.3	8.1	32.2	121.0
2001	5.3	*	*	*	*	0.1	0.2	0.8	2.1	8.8	33.8	127.2
2000	5.8	*	*	*	*	*	0.3	0.8	2.5	8.9	37.7	143.4
1999	6.3	*	*	*	*	0.1	0.3	0.8	2.7	10.5	41.1	155.8
Non-Hispanic white, male												
2002	4.1	*	*	*	*	*	0.2	1.1	2.7	9.0	35.5	110.0
2001	4.2	*	*	*	*	*	0.3	0.9	2.7	10.4	37.3	112.5
2000	4.5	*	*	*	*	*	0.3	0.9	3.1	10.5	41.4	134.2
1999	4.9	*	*	*	*	*	0.4	1.0	3.3	12.5	44.7	145.3
Non-Hispanic white, female												
2002	6.3	*	*	*	*	*	0.2	0.6	1.8	7.2	29.9	125.8
2001	6.5	*	*	*	*	*	0.2	0.6	1.6	7.4	31.4	133.3
2000	7.1	*	*	*	*	*	0.2	0.7	2.0	7.6	35.3	147.1
1999	7.7	*	*	*	*	*	0.2	0.6	2.2	8.9	38.8	160.0
Non-Hispanic black, both sexes												
2002	3.6	*	*	*	*	*	0.5	2.2	5.0	15.2	42.7	113.7
2001	3.8	*	*	*	*	*	0.7	2.2	5.8	16.7	42.8	123.0
2000	4.2	*	*	*	*	*	0.8	2.1	6.6	16.9	49.6	144.1
1999	4.5	*	*	*	*	*	0.7	2.2	6.7	20.8	53.6	152.0
Non-Hispanic black, male												
2002	3.2	*	*	*	*	*	*	2.8	6.7	18.4	46.9	114.9
2001	3.4	*	*	*	*	*	*	2.6	7.5	19.1	54.6	129.4
2000	3.8	*	*	*	*	*	0.8	2.9	8.5	21.1	60.2	139.7
1999	4.1	*	*	*	*	*	*	3.0	8.4	26.5	63.8	162.8
Non-Hispanic black, female												
2002	4.0	*	*	*	*	*	*	1.6	3.7	13.0	40.3	113.3
2001	4.1	*	*	*	*	*	0.8	1.8	4.5	15.1	36.2	120.6
2000	4.7	*	*	*	*	*	0.8	1.5	5.0	14.0	43.6	145.7
1999	4.9	*	*	*	*	*	0.8	1.6	5.4	16.8	48.0	147.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)												
All origins 3/, both sexes												
2002	30.1	*	*	*	*	0.2	0.9	4.0	15.0	60.5	237.6	859.4
2001	31.1	*	*	*	*	0.2	0.9	3.8	15.6	62.8	249.5	906.3
2000	32.1	*	*	*	*	0.2	0.9	3.8	16.2	66.4	257.6	954.7
1999	32.6	*	*	*	*	0.2	0.8	3.8	16.3	66.5	263.9	982.4
All origins 3/, male												
2002	22.6	*	*	*	*	0.2	1.0	4.6	18.0	68.8	241.0	762.8
2001	23.5	*	*	*	*	0.1	0.9	4.5	18.7	72.1	256.3	819.7
2000	24.2	*	*	*	*	0.2	0.9	4.7	19.4	75.1	270.6	863.2
1999	24.7	*	*	*	*	0.2	0.8	4.6	19.4	75.9	280.4	896.4
All origins 3/, female												
2002	37.4	*	*	*	*	0.2	0.9	3.3	12.1	53.6	235.4	901.3
2001	38.5	*	*	*	*	0.2	0.8	3.0	12.7	55.1	245.0	942.5
2000	39.7	*	*	*	*	0.2	0.8	3.0	13.1	59.2	249.0	992.0
1999	40.3	*	*	*	*	0.2	0.7	3.0	13.5	58.9	253.3	1,017.1
Hispanic, both sexes												
2002	7.6	*	*	*	*	*	0.5	3.3	13.0	49.2	171.5	508.9
2001	8.0	*	*	*	*	*	0.6	3.6	12.0	49.3	183.9	609.0
2000	8.1	*	*	*	*	*	0.6	2.8	13.5	56.9	179.6	650.9
1999	7.9	*	*	*	*	*	0.5	2.7	13.3	53.8	181.2	646.1
Hispanic, male												
2002	6.6	*	*	*	*	*	*	3.8	16.5	61.2	186.7	480.2
2001	6.9	*	*	*	*	*	*	4.5	14.6	59.1	210.2	622.8
2000	7.0	*	*	*	*	*	*	3.9	15.3	69.8	213.9	608.7
1999	7.0	*	*	*	*	*	*	3.6	16.1	65.2	212.2	683.7
Hispanic, female												
2002	8.7	*	*	*	*	*	*	2.8	9.9	39.7	161.0	523.1
2001	9.1	*	*	*	*	*	*	2.8	9.6	41.6	166.2	602.3
2000	9.3	*	*	*	*	*	*	1.8	11.9	46.8	156.5	671.7
1999	8.9	*	*	*	*	*	*	1.7	10.8	44.7	160.6	628.1
Non-Hispanic 4/, both sexes												
2002	33.5	*	*	*	*	0.2	1.0	4.0	15.1	61.1	240.4	871.6
2001	34.5	*	*	*	*	0.2	0.9	3.7	15.8	63.6	251.9	915.4
2000	35.5	*	*	*	*	0.2	0.9	3.9	16.3	66.8	260.2	964.0
1999	36.0	*	*	*	*	0.2	0.8	3.9	16.5	67.1	266.8	992.4
Non-Hispanic 4/, male												
2002	25.2	*	*	*	*	0.2	1.0	4.6	18.1	69.1	243.0	774.2
2001	26.0	*	*	*	*	*	1.0	4.5	18.9	72.8	257.7	826.2
2000	26.8	*	*	*	*	0.3	1.0	4.7	19.7	75.3	272.3	871.9
1999	27.2	*	*	*	*	0.2	0.9	4.7	19.5	76.3	282.5	903.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)												
Non-Hispanic 4/, female												
2002	41.5	*	*	*	*	0.3	0.9	3.4	12.3	54.4	238.6	913.6
2001	42.6	*	*	*	*	0.2	0.8	3.0	12.9	55.9	248.1	952.5
2000	43.8	*	*	*	*	0.2	0.8	3.1	13.2	59.8	252.3	1,001.2
1999	44.3	*	*	*	*	0.2	0.7	3.1	13.6	59.6	256.6	1,027.8
Non-Hispanic white, both sexes												
2002	36.4	*	*	*	*	0.2	0.7	2.9	11.9	54.9	235.0	879.7
2001	37.3	*	*	*	*	0.2	0.7	2.7	12.5	57.0	246.4	922.8
2000	38.4	*	*	*	*	0.2	0.7	2.9	13.2	60.7	254.2	972.8
1999	38.9	*	*	*	*	0.1	0.6	2.8	13.1	60.7	261.2	1,003.8
Non-Hispanic white, male												
2002	26.6	*	*	*	*	0.2	0.8	3.5	14.2	61.5	236.6	778.0
2001	27.4	*	*	*	*	*	0.8	3.3	14.8	64.8	251.1	829.8
2000	28.3	*	*	*	*	0.2	0.8	3.6	16.1	68.1	264.9	873.6
1999	28.8	*	*	*	*	*	0.6	3.4	15.7	68.9	276.0	910.5
Non-Hispanic white, female												
2002	45.7	*	*	*	*	0.2	0.6	2.3	9.8	49.3	233.9	923.6
2001	46.8	*	*	*	*	0.2	0.5	2.1	10.4	50.3	243.3	961.5
2000	48.1	*	*	*	*	0.2	0.6	2.2	10.5	54.5	247.3	1,012.8
1999	48.7	*	*	*	*	0.2	0.5	2.2	10.7	53.8	251.5	1,041.1
Non-Hispanic black, both sexes												
2002	26.9	*	*	*	*	0.4	2.6	11.7	42.5	122.7	329.5	851.1
2001	27.5	*	*	*	*	0.4	2.6	11.3	43.4	128.5	337.4	891.5
2000	28.2	*	*	*	*	0.5	2.3	11.5	42.7	131.5	354.7	929.5
1999	28.4	*	*	*	*	0.4	2.3	12.1	45.5	134.2	354.4	923.4
Non-Hispanic black, male												
2002	22.9	*	*	*	*	*	2.4	13.6	54.5	153.6	349.6	774.5
2001	23.5	*	*	*	*	*	2.7	13.4	56.5	158.5	367.3	824.7
2000	24.0	*	*	*	*	*	2.3	14.0	52.3	157.4	395.8	911.2
1999	24.1	*	*	*	*	*	2.4	14.7	56.2	158.3	389.6	885.4
Non-Hispanic black, female												
2002	30.6	*	*	*	*	*	2.8	10.0	32.8	100.7	318.2	880.4
2001	31.2	*	*	*	*	*	2.5	9.5	32.9	107.2	320.6	916.2
2000	32.0	*	*	*	*	*	2.4	9.4	34.9	113.2	331.5	936.3
1999	32.3	*	*	*	*	*	2.2	9.9	36.9	117.2	334.8	937.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)												
All origins 3/, both sexes												
2002	10.5	1.4	0.2	0.1	0.0	0.1	0.4	1.1	3.8	17.4	80.0	328.9
2001	10.3	1.2	0.2	0.1	*	0.1	0.3	1.0	3.9	17.5	80.2	328.2
2000	10.5	1.6	0.2	*	0.1	0.1	0.3	1.1	4.1	17.6	82.1	341.4
1999	10.6	1.1	0.1	0.1	0.1	0.1	0.3	1.1	4.1	18.5	83.3	343.9
All origins 3/, male												
2002	7.9	1.7	0.3	0.1	*	0.1	0.3	1.2	4.5	20.0	86.4	302.3
2001	7.9	1.5	*	*	*	0.1	0.3	1.1	4.6	20.8	88.2	305.5
2000	8.1	1.8	*	*	*	*	0.3	1.2	4.7	21.2	92.0	326.9
1999	8.1	1.4	*	*	*	*	0.3	1.1	5.2	21.6	92.4	332.1
All origins 3/, female												
2002	13.0	1.1	*	*	*	*	0.4	1.0	3.2	15.3	75.7	340.4
2001	12.7	*	*	*	*	0.1	0.3	1.0	3.2	14.7	74.9	337.6
2000	12.8	1.4	*	*	*	0.1	0.3	1.1	3.6	14.6	75.6	347.4
1999	12.9	*	*	*	*	0.1	0.4	1.0	3.2	15.9	77.4	348.7
Hispanic, both sexes												
2002	2.8	*	*	*	*	*	*	1.1	3.7	15.2	58.7	212.3
2001	2.8	*	*	*	*	*	*	0.6	3.8	16.2	64.7	222.6
2000	2.7	*	*	*	*	*	*	1.1	3.6	14.2	65.0	224.3
1999	2.6	*	*	*	*	*	*	0.8	3.4	16.7	65.0	209.8
Hispanic, male												
2002	2.5	*	*	*	*	*	*	1.4	4.9	20.8	65.7	210.8
2001	2.4	*	*	*	*	*	*	*	5.5	21.6	77.8	198.1
2000	2.3	*	*	*	*	*	*	1.4	4.2	20.0	73.3	207.6
1999	2.4	*	*	*	*	*	*	*	4.0	22.0	80.3	235.1
Hispanic, female												
2002	3.1	*	*	*	*	*	*	*	2.7	10.6	54.0	213.0
2001	3.1	*	*	*	*	*	*	*	2.2	12.0	55.9	234.6
2000	3.1	*	*	*	*	*	*	*	3.1	9.6	59.4	232.5
1999	2.8	*	*	*	*	*	*	*	2.9	12.6	54.8	197.6
Non-Hispanic 4/, both sexes												
2002	11.7	1.6	0.2	0.1	0.1	0.1	0.4	1.1	3.8	17.5	80.9	333.0
2001	11.4	1.4	0.2	*	*	0.1	0.3	1.0	3.9	17.5	80.7	331.2
2000	11.6	1.9	0.2	*	0.1	0.1	0.3	1.1	4.2	17.7	82.6	344.9
1999	11.7	1.2	*	0.1	0.1	0.1	0.4	1.1	4.2	18.5	83.8	348.1
Non-Hispanic 4/, male												
2002	8.8	2.0	*	*	*	0.1	0.3	1.2	4.4	19.9	87.3	305.9
2001	8.7	1.6	*	*	*	0.1	0.4	1.1	4.5	20.8	88.4	309.0
2000	9.0	2.2	*	*	*	*	0.4	1.1	4.8	21.2	92.7	331.0
1999	9.0	1.7	*	*	*	*	0.3	1.2	5.3	21.6	92.5	335.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)												
Non-Hispanic 4/, female												
2002	14.4	1.3	*	*	*	*	0.5	1.0	3.2	15.6	76.6	344.6
2001	14.0	*	*	*	*	0.1	0.3	1.0	3.3	14.9	75.7	340.5
2000	14.1	1.6	*	*	*	0.1	0.3	1.1	3.6	14.9	76.1	350.5
1999	14.2	*	*	*	*	0.2	0.4	1.1	3.2	16.0	78.2	353.3
Non-Hispanic white, both sexes												
2002	12.8	0.9	*	*	*	0.1	0.3	0.8	3.1	15.9	78.9	336.1
2001	12.5	1.0	*	*	*	0.1	0.3	0.8	3.2	15.9	78.7	333.1
2000	12.6	1.4	*	*	*	0.1	0.3	0.9	3.5	16.4	80.4	346.8
1999	12.7	1.1	*	*	*	0.1	0.3	0.9	3.4	16.7	81.8	352.5
Non-Hispanic white, male												
2002	9.5	*	*	*	*	*	0.3	0.9	3.6	18.1	84.7	310.0
2001	9.4	*	*	*	*	*	0.3	0.8	3.7	18.9	86.5	311.8
2000	9.6	2.0	*	*	*	*	0.3	0.9	4.1	19.3	89.9	334.4
1999	9.5	*	*	*	*	*	0.2	0.9	4.2	19.1	89.9	338.1
Non-Hispanic white, female												
2002	15.9	*	*	*	*	*	0.4	0.7	2.7	14.0	75.0	347.4
2001	15.4	*	*	*	*	*	0.3	0.7	2.6	13.4	73.6	341.9
2000	15.5	*	*	*	*	*	0.2	0.8	2.9	13.9	74.2	351.8
1999	15.8	*	*	*	*	*	0.4	0.9	2.7	14.7	76.5	358.2
Non-Hispanic black, both sexes												
2002	8.7	3.6	*	*	*	*	0.7	2.9	10.0	34.7	108.5	322.7
2001	8.6	3.3	*	*	*	*	0.7	3.0	10.2	32.7	108.9	330.2
2000	8.5	4.2	*	*	*	*	0.8	2.9	9.8	30.5	109.3	336.8
1999	8.5	*	*	*	*	*	0.8	2.6	10.4	36.6	110.1	314.9
Non-Hispanic black, male												
2002	7.0	*	*	*	*	*	*	3.2	12.4	41.0	124.3	270.2
2001	6.8	*	*	*	*	*	0.8	3.2	12.0	40.6	120.0	285.3
2000	6.8	*	*	*	*	*	0.8	2.8	11.0	40.1	128.0	293.5
1999	7.3	*	*	*	*	*	0.8	3.4	14.3	48.2	123.2	294.7
Non-Hispanic black, female												
2002	10.2	*	*	*	*	*	0.8	2.6	8.1	30.2	99.6	342.8
2001	10.2	*	*	*	*	*	*	2.9	8.7	27.1	102.6	346.9
2000	10.0	*	*	*	*	*	0.7	3.1	8.9	23.7	98.8	352.8
1999	9.7	*	*	*	*	*	0.8	1.9	7.2	28.4	102.8	322.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I72-I78)												
All origins 3/, both sexes												
2002	13.4	*	*	*	0.2	0.4	1.1	3.1	10.4	36.2	100.6	316.5
2001	13.8	*	*	*	0.2	0.4	1.2	3.1	10.5	36.5	104.9	337.3
2000	14.4	*	*	*	0.2	0.4	1.1	3.2	11.1	40.0	109.2	352.0
1999	14.6	*	*	*	0.1	0.4	1.2	3.2	11.3	39.5	111.6	367.2
All origins 3/, male												
2002	13.0	*	*	*	0.2	0.5	1.4	4.3	14.3	48.6	127.8	344.0
2001	13.3	*	*	*	0.2	0.6	1.6	4.2	14.3	49.3	133.0	364.5
2000	14.1	*	*	*	0.2	0.5	1.4	4.4	15.7	55.1	142.5	391.4
1999	14.1	*	*	*	0.2	0.5	1.5	4.3	16.0	54.0	144.2	396.5
All origins 3/, female												
2002	13.8	*	*	*	0.1	0.3	0.8	2.0	6.8	25.8	82.5	304.6
2001	14.4	*	*	*	0.1	0.3	0.8	2.0	7.1	25.8	86.4	325.9
2000	14.6	*	*	*	0.1	0.3	0.8	2.1	6.8	27.5	87.5	336.0
1999	15.2	*	*	*	*	0.2	0.9	2.0	7.0	27.7	90.5	355.4
Hispanic, both sexes												
2002	3.3	*	*	*	*	0.3	0.8	1.9	6.9	19.1	63.7	201.2
2001	3.2	*	*	*	*	*	0.7	2.6	5.8	17.9	62.2	225.7
2000	3.5	*	*	*	*	0.4	0.8	2.1	6.7	26.0	67.0	235.6
1999	3.4	*	*	*	*	*	0.7	1.8	7.3	21.6	71.0	239.9
Hispanic, male												
2002	3.2	*	*	*	*	*	1.1	2.6	8.3	25.0	83.3	204.1
2001	3.1	*	*	*	*	*	0.9	3.3	7.6	25.0	80.9	236.2
2000	3.4	*	*	*	*	*	0.9	2.6	9.0	34.0	93.9	235.8
1999	3.4	*	*	*	*	*	0.9	2.2	9.8	29.0	93.3	314.9
Hispanic, female												
2002	3.3	*	*	*	*	*	*	1.3	5.7	14.3	50.3	199.8
2001	3.2	*	*	*	*	*	*	1.9	4.2	12.3	49.6	220.6
2000	3.5	*	*	*	*	*	*	1.6	4.7	19.7	48.9	235.4
1999	3.3	*	*	*	*	*	*	1.3	5.1	15.8	56.2	203.8
Non-Hispanic 4/, both sexes												
2002	15.0	*	*	*	0.2	0.4	1.1	3.2	10.7	37.3	102.1	320.4
2001	15.4	*	*	*	0.2	0.5	1.3	3.1	10.9	37.6	106.5	340.5
2000	15.9	*	*	*	0.2	0.4	1.1	3.3	11.4	40.7	110.8	355.4
1999	16.1	*	*	*	0.1	0.4	1.2	3.3	11.6	40.5	113.1	370.7
Non-Hispanic 4/, male												
2002	14.6	*	*	*	0.2	0.6	1.5	4.5	14.7	50.1	129.7	349.5
2001	14.8	*	*	*	0.2	0.6	1.7	4.3	14.8	50.7	135.0	368.8
2000	15.6	*	*	*	0.2	0.6	1.4	4.6	16.1	56.2	144.3	396.6
1999	15.6	*	*	*	0.2	0.6	1.5	4.5	16.4	55.2	146.0	398.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic 4/, female												
2002	15.3	*	*	*	0.1	0.3	0.8	2.0	6.9	26.6	83.8	307.9
2001	15.9	*	*	*	*	0.3	0.8	2.0	7.3	26.7	87.9	328.8
2000	16.1	*	*	*	0.1	0.3	0.8	2.1	7.0	28.0	88.9	338.7
1999	16.7	*	*	*	*	0.2	0.9	2.1	7.2	28.4	91.7	359.5
Non-Hispanic white, both sexes												
2002	16.7	*	*	*	0.1	0.4	1.0	2.7	10.2	37.7	103.4	327.9
2001	17.1	*	*	*	0.2	0.4	1.1	2.8	10.4	38.0	107.6	347.6
2000	17.7	*	*	*	0.2	0.4	0.9	2.8	10.9	41.3	111.8	361.6
1999	18.0	*	*	*	0.1	0.4	1.1	2.9	11.4	41.1	114.4	376.6
Non-Hispanic white, male												
2002	16.3	*	*	*	0.2	0.5	1.3	3.9	14.1	51.0	132.2	360.3
2001	16.4	*	*	*	0.2	0.6	1.4	3.7	14.3	51.6	136.9	379.8
2000	17.4	*	*	*	0.2	0.5	1.2	4.0	15.7	57.6	146.9	406.3
1999	17.4	*	*	*	0.2	0.6	1.4	4.1	16.3	56.7	148.3	407.4
Non-Hispanic white, female												
2002	17.2	*	*	*	*	0.3	0.7	1.6	6.5	26.4	84.0	313.9
2001	17.8	*	*	*	*	0.3	0.7	1.8	6.6	26.4	88.2	334.2
2000	17.9	*	*	*	*	0.2	0.7	1.7	6.3	27.5	88.8	343.5
1999	18.6	*	*	*	*	0.2	0.8	1.8	6.7	28.1	92.1	364.3
Non-Hispanic black, both sexes												
2002	9.7	*	*	*	*	0.9	2.2	7.1	16.6	39.8	106.1	275.5
2001	10.0	*	*	*	*	0.8	2.6	6.0	17.5	41.2	111.2	301.8
2000	10.3	*	*	*	*	0.7	2.4	6.8	17.9	42.5	114.5	320.2
1999	10.0	*	*	*	*	0.6	2.2	6.1	15.8	41.5	112.3	335.9
Non-Hispanic black, male												
2002	9.3	*	*	*	*	1.3	2.9	9.7	23.8	49.5	123.7	277.9
2001	9.3	*	*	*	*	1.0	3.7	8.5	21.8	51.3	134.4	281.8
2000	9.6	*	*	*	*	0.9	3.1	9.1	22.7	52.9	137.3	329.9
1999	9.3	*	*	*	*	1.0	2.6	8.3	20.4	52.0	135.1	350.6
Non-Hispanic black, female												
2002	10.0	*	*	*	*	*	1.5	4.8	10.9	32.9	96.2	274.7
2001	10.6	*	*	*	*	*	1.7	3.8	14.0	34.0	98.2	309.3
2000	10.9	*	*	*	*	*	1.8	4.8	14.0	35.2	101.6	316.6
1999	10.7	*	*	*	*	*	1.8	4.3	12.0	34.0	99.6	330.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
All origins 3/, both sexes												
2002	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0
All origins 3/, male												
2002	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0
All origins 3/, female												
2002	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2
Hispanic, both sexes												
2002	1.1	*	*	*	*	*	*	*	1.0	4.2	22.7	108.6
2001	1.1	*	*	*	*	*	*	*	*	4.0	23.1	130.7
2000	1.2	*	*	*	*	*	*	*	*	5.9	24.5	132.7
1999	1.2	*	*	*	*	*	*	*	1.5	5.2	27.1	129.1
Hispanic, male												
2002	0.9	*	*	*	*	*	*	*	*	5.2	26.7	102.1
2001	0.9	*	*	*	*	*	*	*	*	4.8	27.3	127.6
2000	1.0	*	*	*	*	*	*	*	*	8.2	32.0	104.8
1999	1.1	*	*	*	*	*	*	*	*	6.8	31.3	151.0
Hispanic, female												
2002	1.3	*	*	*	*	*	*	*	*	3.5	20.0	111.9
2001	1.4	*	*	*	*	*	*	*	*	3.5	20.2	132.2
2000	1.4	*	*	*	*	*	*	*	*	4.0	19.5	146.4
1999	1.4	*	*	*	*	*	*	*	*	4.0	24.3	118.5
Non-Hispanic 4/, both sexes												
2002	5.4	*	*	*	*	*	0.1	0.5	1.9	7.3	32.2	170.2
2001	5.5	*	*	*	*	*	0.1	0.5	1.9	7.5	33.4	180.5
2000	5.7	*	*	*	*	*	0.1	0.5	1.9	7.8	35.0	190.3
1999	5.9	*	*	*	*	*	0.1	0.5	2.1	8.1	36.1	203.9
Non-Hispanic 4/, male												
2002	4.1	*	*	*	*	*	0.1	0.6	2.5	9.0	35.8	158.7
2001	4.1	*	*	*	*	*	0.1	0.6	2.4	9.2	36.1	168.2
2000	4.3	*	*	*	*	*	0.2	0.7	2.6	9.7	39.9	178.7
1999	4.4	*	*	*	*	*	0.2	0.7	2.7	9.7	40.3	189.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Non-Hispanic 4/, female												
2002	6.6	*	*	*	*	*	*	0.3	1.4	5.9	29.8	175.1
2001	6.8	*	*	*	*	*	*	0.3	1.5	6.0	31.6	185.5
2000	7.0	*	*	*	*	*	*	0.3	1.3	6.3	31.8	195.1
1999	7.4	*	*	*	*	*	*	0.3	1.5	6.9	33.4	209.6
Non-Hispanic white, both sexes												
2002	6.1	*	*	*	*	*	0.1	0.4	1.8	7.3	32.5	177.0
2001	6.3	*	*	*	*	*	0.1	0.5	1.8	7.5	34.1	187.4
2000	6.5	*	*	*	*	*	0.1	0.4	1.8	7.8	35.5	196.8
1999	6.7	*	*	*	*	*	0.1	0.5	2.0	8.2	36.7	209.9
Non-Hispanic white, male												
2002	4.6	*	*	*	*	*	0.1	0.5	2.3	9.0	35.9	165.7
2001	4.7	*	*	*	*	*	0.1	0.6	2.3	9.2	36.6	175.2
2000	4.9	*	*	*	*	*	0.2	0.6	2.6	9.8	40.6	183.6
1999	4.9	*	*	*	*	*	0.1	0.7	2.6	9.6	40.4	194.6
Non-Hispanic white, female												
2002	7.6	*	*	*	*	*	*	0.3	1.3	5.9	30.2	181.9
2001	7.9	*	*	*	*	*	*	0.3	1.3	6.1	32.5	192.6
2000	8.0	*	*	*	*	*	*	0.2	1.1	6.2	32.1	202.1
1999	8.5	*	*	*	*	*	*	0.3	1.4	6.9	34.3	215.9
Non-Hispanic black, both sexes												
2002	2.8	*	*	*	*	*	*	1.0	3.0	8.1	37.0	118.0
2001	2.7	*	*	*	*	*	*	0.7	3.4	8.9	30.7	126.1
2000	3.0	*	*	*	*	*	*	1.1	3.3	9.1	35.6	140.2
1999	3.1	*	*	*	*	*	*	0.7	3.3	9.3	35.0	155.8
Non-Hispanic black, male												
2002	2.4	*	*	*	*	*	*	1.4	4.3	10.2	44.8	109.4
2001	2.2	*	*	*	*	*	*	1.0	3.9	11.0	37.2	113.2
2000	2.5	*	*	*	*	*	*	1.5	4.1	11.1	40.4	150.3
1999	2.6	*	*	*	*	*	*	*	4.5	12.0	44.4	166.4
Non-Hispanic black, female												
2002	3.2	*	*	*	*	*	*	*	2.0	6.6	32.5	121.2
2001	3.2	*	*	*	*	*	*	*	3.0	7.4	27.1	130.9
2000	3.5	*	*	*	*	*	*	*	2.7	7.7	32.8	136.5
1999	3.5	*	*	*	*	*	*	*	2.3	7.4	29.8	151.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All origins 3/, both sexes												
2002	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
All origins 3/, male												
2002	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.5	114.8
2000	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
All origins 3/, female												
2002	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2
Hispanic, both sexes												
2002	1.1	*	*	*	*	*	0.4	1.0	3.8	7.5	21.2	31.0
2001	1.1	*	*	*	*	*	0.4	1.2	2.5	8.2	23.1	31.9
2000	1.3	*	*	*	*	*	0.4	1.3	3.4	11.2	27.1	41.8
1999	1.1	*	*	*	*	*	0.5	0.9	3.4	9.2	24.0	40.7
Hispanic, male												
2002	1.4	*	*	*	*	*	0.7	1.6	5.5	12.0	32.0	43.5
2001	1.5	*	*	*	*	*	0.7	1.8	3.8	14.2	35.3	53.3
2000	1.6	*	*	*	*	*	*	1.8	5.2	16.0	42.3	64.5
1999	1.5	*	*	*	*	*	*	1.5	5.3	14.7	33.9	79.8
Hispanic, female												
2002	0.8	*	*	*	*	*	*	*	2.3	4.0	13.8	24.9
2001	0.7	*	*	*	*	*	*	*	*	3.5	14.8	21.4
2000	1.0	*	*	*	*	*	*	*	*	7.5	16.8	30.7
1999	0.8	*	*	*	*	*	*	*	*	4.8	17.3	21.8
Non-Hispanic 4/, both sexes												
2002	5.7	*	*	*	0.1	0.3	0.7	1.8	5.6	19.8	44.3	70.8
2001	6.0	*	*	*	0.1	0.3	0.8	1.7	5.7	19.8	46.7	79.0
2000	6.2	*	*	*	0.1	0.3	0.7	1.8	6.3	22.4	47.9	80.3
1999	6.3	*	*	*	0.1	0.3	0.8	1.8	6.2	22.4	49.5	81.7
Non-Hispanic 4/, male												
2002	7.2	*	*	*	0.2	0.4	1.1	2.9	8.7	29.6	64.1	109.1
2001	7.3	*	*	*	0.2	0.5	1.2	2.7	8.8	29.5	67.7	117.0
2000	7.8	*	*	*	*	0.4	1.0	2.7	9.8	33.8	71.6	129.6
1999	7.8	*	*	*	*	0.4	1.1	2.8	9.8	33.6	74.3	127.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Non-Hispanic 4/, female												
2002	4.4	*	*	*	*	0.1	0.4	0.9	2.7	11.6	31.2	54.3
2001	4.7	*	*	*	*	0.2	0.3	0.8	2.9	11.8	32.9	63.2
2000	4.7	*	*	*	*	0.1	0.4	0.9	3.1	13.0	32.5	60.4
1999	4.8	*	*	*	*	*	0.4	0.9	2.9	13.2	33.4	63.5
Non-Hispanic white, both sexes												
2002	6.5	*	*	*	0.1	0.2	0.6	1.5	5.5	20.7	46.0	72.8
2001	6.7	*	*	*	0.1	0.3	0.6	1.5	5.6	20.7	47.9	81.3
2000	7.0	*	*	*	*	0.2	0.6	1.6	6.3	23.4	49.3	82.0
1999	7.1	*	*	*	0.1	0.2	0.7	1.6	6.3	23.4	51.1	83.7
Non-Hispanic white, male												
2002	8.0	*	*	*	*	0.3	0.9	2.4	8.6	31.0	67.1	114.7
2001	8.2	*	*	*	0.2	0.5	1.0	2.3	8.7	30.9	70.0	122.9
2000	8.8	*	*	*	*	0.3	0.8	2.5	9.8	35.6	74.2	135.3
1999	8.9	*	*	*	*	0.4	1.0	2.4	10.1	35.4	77.5	131.8
Non-Hispanic white, female												
2002	4.9	*	*	*	*	*	0.3	0.7	2.6	11.9	31.9	54.7
2001	5.2	*	*	*	*	*	0.3	0.6	2.7	12.0	33.3	64.0
2000	5.2	*	*	*	*	*	0.3	0.7	2.9	13.1	32.9	60.5
1999	5.3	*	*	*	*	*	0.4	0.7	2.7	13.3	33.9	64.5
Non-Hispanic black, both sexes												
2002	3.3	*	*	*	*	0.6	1.4	4.3	7.5	14.8	30.2	50.5
2001	3.5	*	*	*	*	0.4	1.6	3.6	7.7	14.7	38.0	55.9
2000	3.6	*	*	*	*	0.5	1.5	3.3	8.2	17.3	36.8	63.0
1999	3.4	*	*	*	*	0.4	1.5	3.9	6.5	16.8	34.0	61.1
Non-Hispanic black, male												
2002	3.9	*	*	*	*	1.0	2.3	6.6	12.1	19.3	34.9	49.2
2001	3.9	*	*	*	*	*	2.6	5.4	11.3	19.9	47.4	49.7
2000	4.0	*	*	*	*	*	2.3	4.7	11.6	22.6	47.2	70.5
1999	3.7	*	*	*	*	*	2.0	6.0	9.3	22.1	38.8	77.3
Non-Hispanic black, female												
2002	2.8	*	*	*	*	*	*	2.2	3.8	11.6	27.5	51.0
2001	3.1	*	*	*	*	*	0.7	2.0	4.8	11.1	32.7	58.2
2000	3.3	*	*	*	*	*	0.9	2.1	5.5	13.5	30.9	60.2
1999	3.1	*	*	*	*	*	1.0	2.0	4.3	13.1	31.3	55.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All origins 3/, both sexes												
2002	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
All origins 3/, male												
2002	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.7	30.6	82.5
2000	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
All origins 3/, female												
2002	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0
Hispanic, both sexes												
2002	1.1	*	*	*	*	*	0.3	0.7	2.1	7.3	19.8	61.5
2001	1.0	*	*	*	*	*	*	0.9	2.2	5.6	16.0	63.2
2000	1.0	*	*	*	*	*	*	*	2.4	8.9	15.4	61.0
1999	1.0	*	*	*	*	*	*	*	2.4	7.2	20.0	70.2
Hispanic, male												
2002	0.9	*	*	*	*	*	*	*	*	7.8	24.6	58.6
2001	0.8	*	*	*	*	*	*	*	2.5	6.0	18.3	55.2
2000	0.9	*	*	*	*	*	*	*	2.7	9.7	19.7	66.5
1999	0.9	*	*	*	*	*	*	*	*	7.5	28.2	84.1
Hispanic, female												
2002	1.3	*	*	*	*	*	*	*	2.6	6.8	16.6	63.0
2001	1.1	*	*	*	*	*	*	*	*	5.4	14.5	67.0
2000	1.1	*	*	*	*	*	*	*	*	8.3	12.5	58.4
1999	1.1	*	*	*	*	*	*	*	2.4	7.1	14.6	63.4
Non-Hispanic 4/, both sexes												
2002	3.9	*	*	*	0.1	0.1	0.3	0.9	3.2	10.2	25.6	79.5
2001	3.9	*	*	*	*	0.1	0.4	0.9	3.2	10.3	26.5	81.1
2000	4.0	*	*	*	0.1	0.1	0.3	1.0	3.1	10.5	27.9	84.7
1999	3.9	*	*	*	*	0.1	0.3	1.0	3.3	9.9	27.5	85.1
Non-Hispanic 4/, male												
2002	3.4	*	*	*	*	*	0.3	1.0	3.5	11.5	29.8	81.7
2001	3.4	*	*	*	*	0.1	0.3	0.9	3.5	12.0	31.1	83.5
2000	3.5	*	*	*	*	0.1	0.2	1.2	3.6	12.7	32.8	88.2
1999	3.4	*	*	*	*	0.1	0.3	1.0	3.9	11.9	31.5	82.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic 4/, female												
2002	4.3	*	*	*	*	0.2	0.3	0.8	2.8	9.0	22.8	78.5
2001	4.4	*	*	*	*	0.1	0.4	0.9	2.9	8.8	23.5	80.1
2000	4.4	*	*	*	*	*	0.4	0.9	2.6	8.7	24.6	83.3
1999	4.5	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.8	86.4
Non-Hispanic white, both sexes												
2002	4.1	*	*	*	*	0.1	0.3	0.8	2.9	9.7	24.9	78.2
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.9	9.8	25.6	78.8
2000	4.3	*	*	*	*	0.1	0.3	0.9	2.8	10.1	27.1	82.8
1999	4.2	*	*	*	*	0.1	0.3	0.9	3.1	9.6	26.5	83.0
Non-Hispanic white, male												
2002	3.6	*	*	*	*	*	0.2	0.9	3.2	11.0	29.3	79.9
2001	3.6	*	*	*	*	*	0.2	0.8	3.3	11.5	30.3	81.8
2000	3.8	*	*	*	*	*	0.2	1.0	3.3	12.3	32.1	87.4
1999	3.6	*	*	*	*	*	0.3	1.0	3.6	11.6	30.4	81.0
Non-Hispanic white, female												
2002	4.7	*	*	*	*	*	0.3	0.7	2.6	8.6	21.9	77.4
2001	4.7	*	*	*	*	*	0.4	0.9	2.6	8.3	22.4	77.6
2000	4.7	*	*	*	*	*	0.3	0.7	2.2	8.2	23.8	81.0
1999	4.8	*	*	*	*	*	0.4	0.8	2.5	7.9	24.0	83.9
Non-Hispanic black, both sexes												
2002	3.6	*	*	*	*	*	0.6	1.8	6.2	16.8	39.0	107.1
2001	3.8	*	*	*	*	*	0.9	1.7	6.4	17.5	42.5	119.9
2000	3.7	*	*	*	*	*	0.6	2.4	6.3	16.1	42.1	117.0
1999	3.6	*	*	*	*	*	0.6	1.5	5.9	15.3	43.3	119.0
Non-Hispanic black, male												
2002	3.1	*	*	*	*	*	*	1.6	7.4	19.9	44.0	119.2
2001	3.3	*	*	*	*	*	0.9	2.0	6.6	20.5	49.8	119.0
2000	3.2	*	*	*	*	*	*	2.9	7.1	19.2	49.7	109.2
1999	3.0	*	*	*	*	*	*	1.4	6.7	17.9	51.9	107.0
Non-Hispanic black, female												
2002	4.0	*	*	*	*	*	0.8	1.9	5.1	14.6	36.1	102.4
2001	4.4	*	*	*	*	*	0.9	1.4	6.3	15.5	38.4	120.2
2000	4.2	*	*	*	*	*	0.7	1.9	5.7	14.0	37.9	119.9
1999	4.2	*	*	*	*	*	0.8	1.6	5.4	13.5	38.5	123.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
All origins 3/, both sexes												
2002	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4
All origins 3/, male												
2002	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8
All origins 3/, female												
2002	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5
Hispanic, both sexes												
2002	0.6	*	*	*	*	*	0.4	1.0	1.4	3.1	6.3	18.3
2001	0.6	*	*	*	*	*	*	0.9	*	4.2	7.4	32.5
2000	0.6	*	*	*	*	*	0.6	1.1	1.3	3.3	8.7	24.6
1999	0.5	*	*	*	*	*	*	1.1	1.5	4.1	5.4	24.6
Hispanic, male												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	1.3	*	4.0	8.9	*
2000	0.6	*	*	*	*	*	0.8	*	*	*	10.8	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	*	3.2	6.2	23.2
2001	0.6	*	*	*	*	*	*	*	*	4.4	6.3	33.5
2000	0.7	*	*	*	*	*	*	*	*	*	7.3	23.7
1999	0.6	*	*	*	*	*	*	*	*	4.3	*	23.9
Non-Hispanic 4/, both sexes												
2002	1.8	0.6	*	*	0.1	0.4	0.7	1.4	2.3	3.9	10.0	24.7
2001	1.8	*	*	*	0.1	0.4	0.7	1.2	2.3	4.6	9.5	25.8
2000	1.8	0.7	*	*	0.1	0.4	0.7	1.4	2.2	4.4	9.4	25.6
1999	1.6	0.7	*	*	0.1	0.3	0.6	1.0	1.9	3.7	9.3	23.3
Non-Hispanic 4/, male												
2002	1.5	*	*	*	*	0.4	0.7	1.5	2.5	3.9	9.6	21.1
2001	1.5	*	*	*	*	0.3	0.7	1.4	2.6	4.9	9.8	21.6
2000	1.5	*	*	*	*	0.3	0.8	1.6	2.3	4.8	9.6	22.1
1999	1.4	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.7	20.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Non-Hispanic 4/, female												
2002	2.1	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.2	26.2
2001	2.0	*	*	*	0.1	0.4	0.6	1.0	2.1	4.3	9.2	27.6
2000	2.0	*	*	*	*	0.4	0.6	1.2	2.1	4.1	9.3	27.0
1999	1.8	*	*	*	*	0.3	0.5	0.9	1.8	3.3	9.0	24.4
Non-Hispanic white, both sexes												
2002	1.8	*	*	*	*	0.3	0.6	1.2	2.1	3.6	9.6	24.5
2001	1.8	*	*	*	*	0.3	0.6	1.0	2.1	4.3	9.2	25.7
2000	1.8	*	*	*	*	0.3	0.6	1.2	1.9	4.0	9.2	25.5
1999	1.6	*	*	*	*	0.2	0.5	0.9	1.8	3.5	9.0	23.0
Non-Hispanic white, male												
2002	1.5	*	*	*	*	0.3	0.6	1.3	2.2	3.7	9.1	20.5
2001	1.5	*	*	*	*	0.2	0.6	1.2	2.3	4.6	9.4	21.2
2000	1.5	*	*	*	*	0.2	0.7	1.5	2.0	4.5	9.4	21.8
1999	1.4	*	*	*	*	0.2	0.6	0.9	2.0	3.9	9.3	20.3
Non-Hispanic white, female												
2002	2.2	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.9	26.3
2001	2.1	*	*	*	*	0.3	0.5	0.8	1.9	4.1	9.0	27.6
2000	2.0	*	*	*	*	0.3	0.5	1.0	1.9	3.6	9.0	27.0
1999	1.8	*	*	*	*	0.2	0.5	0.8	1.7	3.1	8.8	24.1
Non-Hispanic black, both sexes												
2002	2.2	*	*	*	*	1.0	1.7	2.7	5.0	6.9	16.8	29.9
2001	2.0	*	*	*	*	0.8	1.6	2.3	4.9	7.9	14.3	29.7
2000	2.1	*	*	*	*	0.9	1.5	2.8	4.6	9.0	14.4	29.9
1999	1.8	*	*	*	*	0.5	1.5	2.4	3.0	6.8	14.0	28.5
Non-Hispanic black, male												
2002	2.1	*	*	*	*	1.1	1.8	3.0	5.6	7.8	18.5	31.7
2001	2.0	*	*	*	*	*	1.8	2.4	6.5	9.2	15.9	28.9
2000	2.0	*	*	*	*	0.9	1.8	3.5	5.3	9.1	13.3	30.5
1999	1.7	*	*	*	*	*	1.9	2.9	2.7	8.4	17.2	*
Non-Hispanic black, female												
2002	2.3	*	*	*	*	1.0	1.6	2.4	4.5	6.2	15.9	29.3
2001	2.1	*	*	*	*	0.9	1.4	2.1	3.7	7.0	13.5	29.9
2000	2.3	*	*	*	*	1.0	1.3	2.3	4.0	8.8	15.1	29.7
1999	1.8	*	*	*	*	*	1.1	2.0	3.2	5.6	12.3	30.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
All origins 3/, both sexes												
2002	1.2	*	*	*	0.1	0.3	0.6	1.0	1.7	2.8	7.3	18.6
2001	1.2	*	*	*	0.1	0.3	0.5	0.8	1.7	3.4	7.1	19.6
2000	1.2	*	*	*	0.1	0.3	0.6	1.0	1.5	3.2	7.0	18.7
1999	0.9	*	*	*	0.1	0.2	0.5	0.7	1.2	2.4	5.8	14.6
All origins 3/, male												
2002	1.0	*	*	*	0.1	0.3	0.5	1.0	1.7	2.8	6.6	14.8
2001	1.0	*	*	*	*	0.2	0.5	0.9	1.8	3.5	6.8	15.2
2000	1.0	*	*	*	*	0.3	0.6	1.0	1.4	3.3	6.9	15.0
1999	0.8	*	*	*	*	0.2	0.5	0.6	1.1	2.7	5.7	11.2
All origins 3/, female												
2002	1.5	*	*	*	*	0.3	0.6	1.0	1.7	2.9	7.8	20.2
2001	1.5	*	*	*	*	0.3	0.5	0.8	1.6	3.3	7.3	21.4
2000	1.4	*	*	*	0.1	0.3	0.5	1.0	1.6	3.1	7.1	20.2
1999	1.1	*	*	*	*	0.2	0.4	0.7	1.3	2.3	5.8	15.9
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	0.6	*	2.0	4.7	13.3
2001	0.4	*	*	*	*	*	*	0.6	*	3.4	5.4	26.9
2000	0.5	*	*	*	*	*	0.4	*	*	2.2	7.3	19.9
1999	0.4	*	*	*	*	*	*	0.7	*	2.9	*	16.1
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	*	18.2
2001	0.5	*	*	*	*	*	*	*	*	3.8	*	27.9
2000	0.5	*	*	*	*	*	*	*	*	*	6.6	21.8
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.4	*	*	*	0.1	0.3	0.6	1.1	1.8	2.9	7.5	18.8
2001	1.4	*	*	*	0.1	0.3	0.6	0.9	1.8	3.4	7.2	19.3
2000	1.3	*	*	*	0.1	0.3	0.6	1.0	1.5	3.3	7.0	18.6
1999	1.0	*	*	*	0.1	0.2	0.5	0.7	1.2	2.4	5.8	14.5
Non-Hispanic 4/, male												
2002	1.1	*	*	*	*	0.3	0.6	1.1	1.8	2.9	6.8	15.3
2001	1.1	*	*	*	*	0.3	0.6	0.9	1.9	3.5	6.9	14.8
2000	1.1	*	*	*	*	0.3	0.6	1.1	1.4	3.3	6.8	14.9
1999	0.8	*	*	*	*	0.2	0.5	0.6	1.1	2.6	5.7	10.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Non-Hispanic 4/, female												
2002	1.6	*	*	*	*	0.4	0.6	1.1	1.8	2.9	7.9	20.2
2001	1.6	*	*	*	*	0.4	0.6	0.8	1.7	3.3	7.4	21.2
2000	1.5	*	*	*	*	0.4	0.5	1.0	1.6	3.2	7.1	20.1
1999	1.2	*	*	*	*	0.2	0.5	0.7	1.3	2.2	5.9	15.9
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	0.3	0.5	0.9	1.6	2.7	7.2	18.6
2001	1.4	*	*	*	*	0.3	0.4	0.8	1.6	3.2	7.0	19.3
2000	1.3	*	*	*	*	0.2	0.5	0.9	1.4	3.0	6.8	18.7
1999	1.0	*	*	*	*	0.2	0.4	0.6	1.2	2.2	5.7	14.2
Non-Hispanic white, male												
2002	1.1	*	*	*	*	0.2	0.5	0.9	1.6	2.7	6.5	14.9
2001	1.1	*	*	*	*	0.2	0.4	0.8	1.6	3.3	6.7	14.7
2000	1.1	*	*	*	*	0.2	0.5	1.0	1.3	3.1	6.8	14.8
1999	0.8	*	*	*	*	0.2	0.4	0.6	1.1	2.4	5.5	11.0
Non-Hispanic white, female												
2002	1.7	*	*	*	*	0.3	0.5	1.0	1.6	2.8	7.6	20.2
2001	1.6	*	*	*	*	0.3	0.5	0.7	1.5	3.0	7.1	21.2
2000	1.6	*	*	*	*	0.3	0.4	0.9	1.5	2.9	6.8	20.3
1999	1.2	*	*	*	*	0.2	0.4	0.6	1.3	2.1	5.7	15.5
Non-Hispanic black, both sexes												
2002	1.8	*	*	*	*	1.0	1.5	2.2	3.8	5.3	13.0	23.9
2001	1.7	*	*	*	*	0.8	1.3	1.8	4.0	6.4	10.9	22.8
2000	1.7	*	*	*	*	0.9	1.3	2.2	3.4	7.0	10.9	21.6
1999	1.2	*	*	*	*	0.5	1.2	1.6	2.0	4.7	9.4	18.8
Non-Hispanic black, male												
2002	1.6	*	*	*	*	1.0	1.5	2.3	4.3	5.6	12.3	23.0
2001	1.5	*	*	*	*	*	1.4	1.8	5.0	6.8	11.1	*
2000	1.5	*	*	*	*	0.8	1.5	2.4	3.6	6.7	8.3	*
1999	1.1	*	*	*	*	*	1.4	1.5	1.9	5.9	10.3	*
Non-Hispanic black, female												
2002	1.9	*	*	*	*	1.0	1.5	2.1	3.4	5.1	13.4	24.2
2001	1.8	*	*	*	*	0.9	1.2	1.8	3.2	6.2	10.8	23.5
2000	1.9	*	*	*	*	1.0	1.1	2.0	3.3	7.2	12.3	21.8
1999	1.4	*	*	*	*	*	1.1	1.7	2.1	3.9	8.9	21.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified disorders of circulatory system (I83-I99)												
All origins 3/, both sexes												
2002	0.4	0.5	*	*	*	*	0.1	0.3	0.5	1.0	2.5	5.9
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	2.3	6.5
2000	0.4	0.5	*	*	*	*	0.2	0.4	0.6	1.1	2.4	6.8
1999	0.5	*	*	*	*	*	0.1	0.3	0.7	1.3	3.4	8.8
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.2	0.4	0.7	1.1	2.9	5.7
2001	0.4	*	*	*	*	*	0.2	0.5	0.7	1.4	2.9	6.8
2000	0.4	*	*	*	*	*	0.2	0.6	0.8	1.4	2.8	7.3
1999	0.5	*	*	*	*	*	0.2	0.5	0.9	1.6	3.9	9.6
All origins 3/, female												
2002	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.2	5.9
2001	0.4	*	*	*	*	*	*	0.2	0.4	1.0	1.8	6.4
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.1	6.7
1999	0.5	*	*	*	*	*	*	0.2	0.5	1.1	3.0	8.5
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.5	1.0	2.5	5.9
2001	0.4	*	*	*	*	*	0.1	0.3	0.6	1.2	2.3	6.5
2000	0.5	*	*	*	*	*	0.1	0.4	0.6	1.1	2.4	6.9
1999	0.5	*	*	*	*	*	0.1	0.3	0.7	1.3	3.4	8.8
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	0.2	0.4	0.7	1.1	2.9	5.7
2001	0.4	*	*	*	*	*	0.2	0.5	0.7	1.4	2.9	6.7
2000	0.5	*	*	*	*	*	0.2	0.5	0.9	1.4	2.8	7.1
1999	0.5	*	*	*	*	*	0.2	0.5	0.9	1.6	4.0	9.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified disorders of circulatory system (I83-I99)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.3	6.0
2001	0.4	*	*	*	*	*	*	0.2	0.4	1.0	1.9	6.4
2000	0.4	*	*	*	*	*	*	0.2	0.4	0.8	2.2	6.8
1999	0.5	*	*	*	*	*	*	0.2	0.5	1.1	3.1	8.5
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.5	0.9	2.5	5.9
2001	0.5	*	*	*	*	*	0.1	0.3	0.5	1.2	2.2	6.5
2000	0.5	*	*	*	*	*	0.1	0.3	0.6	1.0	2.4	6.8
1999	0.6	*	*	*	*	*	0.1	0.3	0.6	1.2	3.3	8.8
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	0.1	0.4	0.6	1.0	2.6	5.6
2001	0.5	*	*	*	*	*	0.2	0.4	0.7	1.2	2.7	6.6
2000	0.5	*	*	*	*	*	0.2	0.5	0.8	1.4	2.7	7.0
1999	0.5	*	*	*	*	*	0.2	0.3	0.9	1.5	3.8	9.3
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	*	0.2	0.3	0.9	2.3	6.1
2001	0.5	*	*	*	*	*	*	*	0.4	1.1	1.9	6.4
2000	0.5	*	*	*	*	*	*	0.2	0.4	0.7	2.2	6.7
1999	0.6	*	*	*	*	*	*	0.2	0.4	1.0	3.0	8.6
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.5	1.2	1.6	3.8	6.0
2001	0.4	*	*	*	*	*	*	0.5	0.9	1.5	3.5	6.9
2000	0.5	*	*	*	*	*	*	0.6	1.1	2.0	3.6	8.3
1999	0.5	*	*	*	*	*	*	0.8	0.9	2.1	4.6	9.7
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	6.2	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	1.1	*	*	*	*
1999	0.6	*	*	*	*	*	*	1.4	*	*	6.9	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	8.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)												
All origins 3/, both sexes												
2002	81.4	15.8	1.9	0.8	1.3	2.3	6.0	17.6	65.7	240.3	662.2	1,663.7
2001	80.8	17.3	1.8	0.7	1.3	2.3	6.0	17.6	67.3	245.9	651.9	1,671.9
2000	82.1	17.1	1.7	0.8	1.4	2.5	6.2	17.9	68.8	250.2	667.5	1,734.0
1999	82.4	18.1	2.1	0.8	1.4	2.5	6.0	17.3	70.8	255.0	671.7	1,733.0
All origins 3/, male												
2002	78.9	18.8	2.2	0.8	1.5	2.4	6.5	19.5	73.0	282.2	832.8	2,150.0
2001	78.2	19.7	1.9	0.8	1.5	2.4	6.5	19.7	74.6	289.4	824.1	2,187.3
2000	80.2	18.9	1.7	0.8	1.6	2.7	6.8	19.9	77.1	297.1	851.4	2,344.4
1999	81.3	19.6	2.2	0.9	1.5	2.6	6.4	19.1	79.5	310.4	870.5	2,356.4
All origins 3/, female												
2002	83.7	12.6	1.5	0.8	1.1	2.1	5.6	15.8	58.9	205.5	548.9	1,452.7
2001	83.2	14.7	1.8	0.7	1.2	2.3	5.6	15.5	60.6	209.9	538.8	1,456.2
2000	84.0	15.2	1.7	0.7	1.2	2.3	5.7	16.0	61.1	211.7	547.6	1,485.4
1999	83.4	16.6	1.9	0.7	1.2	2.4	5.7	15.6	62.9	209.7	542.6	1,481.7
Hispanic, both sexes												
2002	20.0	12.7	1.5	0.6	0.8	1.4	3.4	10.0	34.7	119.9	406.5	1,273.8
2001	20.1	13.8	2.1	0.4	0.7	1.0	3.2	11.0	33.9	123.3	401.6	1,442.5
2000	20.1	15.8	1.6	0.7	0.8	1.7	4.1	10.5	37.2	122.6	413.2	1,406.0
1999	19.8	15.9	1.4	0.6	0.9	1.3	4.1	10.9	35.3	126.3	402.3	1,420.7
Hispanic, male												
2002	19.9	14.3	1.9	0.6	1.0	1.6	3.7	11.7	44.4	155.8	525.7	1,551.1
2001	19.7	14.9	2.0	*	0.9	1.1	3.6	12.5	40.9	156.4	515.6	1,839.8
2000	20.1	16.0	*	0.7	0.9	1.8	4.6	12.8	47.5	156.4	562.0	1,751.4
1999	19.7	18.1	1.5	0.7	0.8	1.3	4.8	11.8	41.5	161.7	538.1	1,882.8
Hispanic, female												
2002	20.1	11.0	*	0.6	*	1.2	3.1	8.4	26.1	91.4	324.9	1,136.5
2001	20.5	12.7	2.2	*	*	0.8	2.8	9.5	27.7	97.3	324.6	1,248.3
2000	20.0	15.7	1.9	0.7	0.7	1.6	3.5	8.2	28.0	96.1	313.3	1,236.5
1999	19.8	13.6	*	0.6	1.0	1.3	3.4	10.0	29.8	98.6	311.9	1,197.9
Non-Hispanic 4/, both sexes												
2002	90.6	16.5	2.0	0.8	1.4	2.5	6.4	18.3	68.0	247.9	673.0	1,675.8
2001	89.5	18.0	1.8	0.8	1.4	2.6	6.4	18.1	69.7	253.1	661.8	1,676.5
2000	90.7	17.3	1.7	0.8	1.5	2.6	6.5	18.4	70.9	257.3	676.4	1,741.6
1999	90.7	18.5	2.2	0.8	1.5	2.7	6.2	17.8	73.1	261.9	680.7	1,738.9
Non-Hispanic 4/, male												
2002	88.3	19.8	2.3	0.8	1.6	2.6	6.9	20.2	75.0	289.6	845.6	2,172.0
2001	87.1	20.9	1.9	0.8	1.6	2.7	6.8	20.2	76.7	296.7	836.5	2,196.7
2000	89.0	19.6	1.8	0.9	1.7	2.8	7.0	20.4	78.9	304.4	861.1	2,363.3
1999	90.0	19.8	2.3	1.0	1.7	2.9	6.5	19.6	81.8	317.8	881.7	2,367.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)												
Non-Hispanic 4/, female												
2002	92.9	12.9	1.6	0.9	1.2	2.3	5.9	16.5	61.4	213.1	558.6	1,461.8
2001	91.9	15.0	1.7	0.7	1.3	2.5	6.0	16.0	63.1	216.9	547.2	1,460.1
2000	92.4	14.9	1.6	0.7	1.3	2.4	5.9	16.6	63.4	218.4	556.1	1,490.2
1999	91.5	17.2	2.0	0.7	1.3	2.6	5.9	16.0	65.1	216.0	550.3	1,487.1
Non-Hispanic white, both sexes												
2002	103.2	11.5	1.4	0.6	1.2	2.1	5.6	17.2	68.6	258.9	695.5	1,726.5
2001	101.3	11.8	1.4	0.6	1.1	2.1	5.4	16.7	70.7	263.4	682.9	1,718.1
2000	102.5	12.1	1.4	0.6	1.2	2.1	5.5	16.9	71.3	265.5	697.1	1,791.6
1999	102.2	12.2	1.6	0.7	1.2	2.1	5.1	16.2	73.7	270.4	700.2	1,785.3
Non-Hispanic white, male												
2002	99.0	14.6	1.6	0.6	1.3	2.2	5.9	18.7	74.0	297.2	862.1	2,224.6
2001	97.2	14.2	1.3	0.7	1.3	2.1	5.8	18.6	76.3	303.3	851.4	2,239.8
2000	99.0	14.4	1.3	0.7	1.4	2.2	6.0	18.2	77.8	307.5	877.2	2,421.7
1999	99.9	13.3	1.4	0.8	1.4	2.3	5.2	17.5	80.7	322.7	896.0	2,422.2
Non-Hispanic white, female												
2002	107.1	8.3	1.2	0.7	1.1	2.1	5.2	15.8	63.6	226.2	583.8	1,511.4
2001	105.3	9.2	1.4	0.6	1.0	2.1	5.0	14.9	65.4	229.6	571.1	1,501.0
2000	105.8	9.6	1.5	0.5	1.0	2.1	5.1	15.6	65.0	230.1	578.7	1,537.5
1999	104.4	11.1	1.8	0.6	1.1	1.9	5.0	14.9	67.0	226.7	572.0	1,531.4
Non-Hispanic black, both sexes												
2002	48.2	35.2	4.2	1.8	2.8	4.7	12.8	29.7	81.1	209.0	526.5	1,214.4
2001	48.6	41.2	3.3	1.4	2.9	5.8	13.3	31.9	78.7	215.1	518.4	1,266.4
2000	49.6	39.7	3.2	1.6	3.1	5.4	13.0	33.7	85.0	233.6	522.4	1,247.8
1999	50.7	43.6	4.1	1.6	2.9	6.5	13.9	33.2	86.2	233.8	537.4	1,301.4
Non-Hispanic black, male												
2002	51.9	40.3	4.8	1.8	3.4	5.3	14.6	34.8	104.5	282.9	774.6	1,705.5
2001	52.1	45.3	3.9	1.5	3.2	6.5	14.0	37.0	99.0	295.7	778.7	1,792.5
2000	54.1	41.3	4.1	1.9	3.5	6.9	14.9	40.2	106.7	330.3	763.9	1,822.3
1999	55.7	46.8	5.4	2.0	3.4	6.7	15.2	40.2	111.0	327.8	813.9	1,907.5
Non-Hispanic black, female												
2002	44.8	30.0	3.6	1.9	2.2	4.2	11.2	25.2	62.3	156.6	386.2	1,026.7
2001	45.5	36.9	2.6	1.3	2.6	5.2	12.7	27.6	62.5	158.3	372.3	1,071.5
2000	45.4	38.1	2.2	1.3	2.6	4.0	11.3	28.1	67.6	165.6	386.4	1,034.4
1999	46.2	40.3	2.9	1.2	2.4	6.3	12.8	27.1	66.4	167.5	382.5	1,074.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute upper respiratory infections (J00-J06)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.2	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.2	1.3
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.3	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	1.7
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.2	1.3
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.3	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	1.7
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute upper respiratory infections (J00-J06)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	0.2	1.4
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.3	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	1.8
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.7
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute pharyngitis and tonsillitis (J02-J03)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute pharyngitis and tonsillitis (J02-J03)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute laryngitis and tracheitis (J04)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute laryngitis and tracheitis (J04)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	1.0
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	1.6
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	*	1.9
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	1.1
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	1.6
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified acute upper respiratory infections (J00-J01,J05-J06)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	*	1.9
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.1
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	1.7
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999	0.1	*	*	*	*	*	*	*	*	*	*	2.1
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)												
All origins 3/, both sexes												
2002	81.3	15.5	1.8	0.8	1.3	2.3	6.0	17.6	65.6	240.2	662.0	1,662.6
2001	80.7	17.0	1.7	0.7	1.3	2.3	6.0	17.5	67.3	245.9	651.7	1,670.6
2000	82.0	16.9	1.7	0.8	1.4	2.4	6.2	17.9	68.7	250.1	667.2	1,732.2
1999	82.3	17.8	2.0	0.8	1.4	2.5	6.0	17.3	70.8	255.0	671.3	1,731.2
All origins 3/, male												
2002	78.9	18.6	2.1	0.8	1.5	2.4	6.5	19.4	72.9	282.1	832.7	2,149.3
2001	78.2	19.4	1.8	0.7	1.5	2.4	6.4	19.7	74.5	289.4	824.0	2,186.7
2000	80.1	18.7	1.7	0.8	1.6	2.6	6.8	19.9	77.1	296.9	851.2	2,343.3
1999	81.3	19.2	2.1	0.9	1.5	2.6	6.3	19.1	79.5	310.3	870.1	2,355.3
All origins 3/, female												
2002	83.7	12.4	1.4	0.8	1.1	2.1	5.6	15.8	58.9	205.4	548.8	1,451.5
2001	83.2	14.6	1.7	0.6	1.1	2.3	5.6	15.5	60.6	209.8	538.5	1,454.6
2000	83.9	15.1	1.6	0.7	1.2	2.2	5.6	16.0	61.1	211.5	547.3	1,483.4
1999	83.3	16.4	1.9	0.7	1.2	2.4	5.7	15.6	62.8	209.6	542.2	1,479.7
Hispanic, both sexes												
2002	20.0	12.5	1.3	0.6	0.8	1.4	3.4	10.0	34.7	119.9	406.0	1,273.3
2001	20.1	13.5	2.0	0.4	0.7	1.0	3.2	11.0	33.8	123.3	401.6	1,442.5
2000	20.0	15.8	1.6	0.7	0.8	1.7	4.1	10.4	37.1	122.4	413.0	1,406.0
1999	19.7	15.5	1.3	0.6	0.9	1.3	4.0	10.9	35.3	126.3	402.3	1,419.3
Hispanic, male												
2002	19.9	14.1	1.7	0.6	1.0	1.6	3.7	11.6	44.4	155.8	525.2	1,551.1
2001	19.7	14.4	2.0	*	0.9	1.1	3.6	12.5	40.9	156.4	515.6	1,839.8
2000	20.1	16.0	*	0.7	0.9	1.7	4.6	12.6	47.4	156.2	562.0	1,751.4
1999	19.7	17.3	1.4	0.7	0.8	1.2	4.6	11.8	41.5	161.7	538.1	1,880.6
Hispanic, female												
2002	20.1	10.8	*	0.6	*	1.2	3.1	8.4	26.0	91.4	324.3	1,135.6
2001	20.5	12.5	2.1	*	*	0.8	2.8	9.5	27.5	97.2	324.6	1,248.3
2000	20.0	15.7	1.9	0.7	0.7	1.6	3.5	8.1	28.0	95.9	313.0	1,236.5
1999	19.8	13.6	*	0.6	1.0	1.3	3.4	9.9	29.8	98.6	311.9	1,196.8
Non-Hispanic 4/, both sexes												
2002	90.6	16.3	1.9	0.8	1.4	2.4	6.4	18.2	67.9	247.8	672.9	1,674.7
2001	89.5	17.8	1.6	0.7	1.4	2.6	6.4	18.1	69.6	253.1	661.6	1,675.1
2000	90.6	17.1	1.7	0.8	1.5	2.6	6.4	18.4	70.8	257.1	676.1	1,739.8
1999	90.7	18.3	2.1	0.8	1.5	2.7	6.2	17.7	73.1	261.8	680.4	1,737.1
Non-Hispanic 4/, male												
2002	88.3	19.6	2.2	0.8	1.6	2.6	6.8	20.1	74.9	289.5	845.4	2,171.2
2001	87.1	20.6	1.7	0.8	1.6	2.7	6.8	20.2	76.7	296.7	836.3	2,196.0
2000	88.9	19.2	1.8	0.9	1.7	2.8	7.0	20.3	78.9	304.2	860.9	2,362.1
1999	89.9	19.6	2.2	0.9	1.7	2.8	6.5	19.6	81.8	317.7	881.3	2,366.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)												
Non-Hispanic 4/, female												
2002	92.8	12.7	1.5	0.9	1.2	2.3	5.9	16.4	61.4	213.0	558.5	1,460.5
2001	91.8	14.9	1.5	0.7	1.2	2.5	6.0	16.0	63.0	216.8	546.9	1,458.5
2000	92.3	14.9	1.5	0.7	1.2	2.3	5.9	16.6	63.4	218.3	555.7	1,488.2
1999	91.4	16.9	2.0	0.7	1.3	2.6	5.9	16.0	65.1	215.9	550.0	1,485.1
Non-Hispanic white, both sexes												
2002	103.1	11.5	1.3	0.6	1.1	2.1	5.5	17.2	68.6	258.8	695.4	1,725.4
2001	101.3	11.7	1.2	0.6	1.1	2.1	5.4	16.7	70.7	263.4	682.6	1,716.6
2000	102.4	11.8	1.3	0.6	1.2	2.1	5.5	16.8	71.2	265.4	696.8	1,789.7
1999	102.1	12.0	1.5	0.7	1.2	2.1	5.1	16.2	73.6	270.4	699.8	1,783.5
Non-Hispanic white, male												
2002	99.0	14.6	1.5	0.6	1.2	2.2	5.9	18.7	74.0	297.2	862.0	2,224.0
2001	97.2	14.1	1.2	0.6	1.3	2.0	5.8	18.5	76.3	303.2	851.2	2,239.1
2000	98.9	13.9	1.3	0.7	1.4	2.1	5.9	18.1	77.8	307.4	877.0	2,420.6
1999	99.8	13.1	1.3	0.8	1.4	2.3	5.2	17.4	80.7	322.6	895.6	2,421.4
Non-Hispanic white, female												
2002	107.0	8.2	1.2	0.6	1.0	2.1	5.2	15.7	63.5	226.1	583.7	1,510.0
2001	105.2	9.2	1.3	0.6	1.0	2.1	5.0	14.9	65.3	229.6	570.8	1,499.3
2000	105.7	9.6	1.4	0.5	1.0	2.1	5.0	15.5	65.0	230.0	578.3	1,535.3
1999	104.3	10.9	1.8	0.6	1.1	1.9	5.0	14.9	67.0	226.7	571.6	1,529.2
Non-Hispanic black, both sexes												
2002	48.1	34.6	4.1	1.8	2.8	4.7	12.8	29.6	81.1	208.9	526.3	1,214.4
2001	48.6	40.7	3.1	1.4	2.9	5.8	13.2	31.9	78.6	215.1	518.3	1,265.4
2000	49.5	39.5	3.1	1.6	3.0	5.4	13.0	33.6	85.0	233.5	522.3	1,247.2
1999	50.6	43.1	4.0	1.6	2.9	6.4	13.9	33.1	86.1	233.6	537.3	1,299.8
Non-Hispanic black, male												
2002	51.8	39.4	4.8	1.8	3.4	5.3	14.6	34.7	104.5	282.7	774.3	1,705.5
2001	52.0	44.4	3.6	1.5	3.2	6.5	13.9	37.0	98.8	295.7	778.7	1,792.5
2000	54.0	41.3	4.0	1.9	3.5	6.9	14.9	40.2	106.7	330.0	763.6	1,820.0
1999	55.6	46.4	5.1	1.9	3.3	6.5	15.2	40.2	110.8	327.7	813.9	1,902.8
Non-Hispanic black, female												
2002	44.8	29.7	3.4	1.9	2.1	4.2	11.2	25.2	62.2	156.5	386.0	1,026.7
2001	45.4	36.9	2.5	1.2	2.6	5.2	12.7	27.5	62.4	158.2	372.1	1,070.2
2000	45.4	37.7	2.2	1.3	2.5	4.0	11.3	27.9	67.6	165.6	386.4	1,034.4
1999	46.1	39.6	2.9	1.2	2.4	6.3	12.7	27.1	66.3	167.3	382.4	1,074.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
All origins 3/, female												
2002	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	9.0
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.3
2000	0.7	*	*	*	*	*	0.0	0.1	0.4	1.1	4.1	20.5
1999	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.6
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	9.2
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2000	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.1	21.1
1999	0.5	*	*	*	*	*	*	*	0.3	0.8	4.3	20.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000	0.9	*	*	*	*	*	*	0.1	0.4	1.0	4.1	20.3
1999	0.8	*	*	*	*	*	*	*	0.3	0.9	3.4	22.0
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.6	9.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.8	*	*	*	*	*	*	0.1	0.4	1.2	4.3	21.7
1999	0.8	*	*	*	*	*	*	0.1	0.3	0.9	3.9	23.3
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	*	*	0.4	1.7	10.0
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000	0.6	*	*	*	*	*	*	*	0.4	1.4	4.3	22.0
1999	0.6	*	*	*	*	*	*	*	0.3	0.9	4.6	22.5
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	*	*	*	1.4	9.5
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2000	1.0	*	*	*	*	*	*	0.1	0.5	1.0	4.3	21.6
1999	1.0	*	*	*	*	*	*	*	0.3	1.0	3.5	23.5
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	9.5
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All origins 3/, both sexes												
2002	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
All origins 3/, male												
2002	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
All origins 3/, female												
2002	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5
Hispanic, both sexes												
2002	7.2	5.5	*	*	*	0.6	1.3	3.4	10.5	33.6	142.1	563.2
2001	7.3	7.0	1.0	*	*	0.3	1.3	3.6	9.1	35.3	136.5	660.3
2000	7.3	7.5	*	*	0.3	0.7	1.6	3.8	11.0	37.2	134.3	633.0
1999	6.5	7.9	*	*	0.4	0.4	1.5	3.4	9.2	30.9	116.7	607.6
Hispanic, male												
2002	6.9	7.3	*	*	*	0.6	1.7	4.7	13.9	43.6	172.8	654.2
2001	6.9	7.4	*	*	*	*	1.7	5.0	11.4	40.7	173.0	765.6
2000	7.1	7.6	*	*	*	0.7	2.1	5.2	14.6	45.6	178.0	751.8
1999	6.1	10.3	*	*	*	*	2.0	4.7	9.9	39.5	147.0	718.2
Hispanic, female												
2002	7.5	*	*	*	*	*	0.9	2.1	7.4	25.7	121.1	518.1
2001	7.8	6.5	*	*	*	*	0.9	2.1	7.1	31.0	111.8	608.8
2000	7.6	7.4	*	*	*	*	1.0	2.3	7.9	30.7	105.0	574.7
1999	6.8	5.4	*	*	*	*	0.8	2.2	8.5	24.1	96.4	554.2
Non-Hispanic 4/, both sexes												
2002	24.8	6.5	0.7	0.2	0.4	0.9	2.3	4.9	11.1	37.3	155.6	691.2
2001	23.8	7.2	0.6	0.2	0.5	0.9	2.3	4.6	10.6	36.0	148.2	682.6
2000	24.7	7.3	0.6	0.2	0.5	0.9	2.4	4.6	11.4	37.9	156.6	725.2
1999	24.3	8.1	0.8	0.2	0.4	0.9	2.4	4.6	10.7	36.5	154.2	732.4
Non-Hispanic 4/, male												
2002	22.3	7.6	0.8	0.2	0.5	1.0	2.5	5.8	13.4	45.9	193.5	801.5
2001	21.4	7.8	0.6	0.2	0.5	1.0	2.6	5.7	13.0	44.8	186.1	806.0
2000	22.2	8.2	0.7	0.2	0.5	1.0	2.8	5.6	13.9	46.8	195.7	873.8
1999	21.7	8.0	0.8	0.2	0.5	1.1	2.6	5.8	12.5	46.2	193.4	877.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Non-Hispanic 4/, female												
2002	27.2	5.3	0.6	0.2	0.4	0.8	2.0	4.0	9.1	30.1	130.6	643.7
2001	26.0	6.6	0.6	0.2	0.4	0.8	2.0	3.6	8.5	28.8	123.3	631.3
2000	27.1	6.4	0.5	0.1	0.4	0.8	2.0	3.7	9.2	30.4	131.2	665.2
1999	26.8	8.2	0.9	0.2	0.4	0.8	2.2	3.5	9.1	28.5	128.8	674.4
Non-Hispanic white, both sexes												
2002	27.6	4.7	0.5	0.2	0.4	0.8	2.0	4.4	10.2	36.4	155.3	703.1
2001	26.1	4.6	0.5	0.2	0.4	0.8	1.9	4.0	9.8	34.9	146.7	691.6
2000	27.1	4.8	0.5	0.2	0.4	0.8	2.1	3.9	10.3	35.6	155.9	738.1
1999	26.8	5.2	0.6	0.2	0.4	0.7	1.9	3.9	9.7	35.2	154.8	743.7
Non-Hispanic white, male												
2002	24.3	6.0	0.5	0.2	0.5	0.9	2.2	5.1	11.9	44.5	191.5	810.6
2001	23.0	5.3	0.6	0.2	0.5	0.9	2.1	4.8	11.6	42.6	182.0	811.5
2000	23.8	5.4	0.4	0.2	0.5	0.8	2.4	4.6	12.2	43.3	193.2	886.1
1999	23.4	5.4	*	0.2	0.4	0.9	2.0	4.7	11.1	43.8	192.7	886.8
Non-Hispanic white, female												
2002	30.7	3.4	0.5	0.2	0.3	0.8	1.8	3.7	8.5	29.5	131.1	656.6
2001	29.1	3.8	0.4	0.2	0.3	0.7	1.7	3.2	8.0	28.3	123.3	641.7
2000	30.3	4.2	0.5	*	0.3	0.8	1.7	3.3	8.5	29.1	131.4	678.4
1999	30.0	4.9	0.7	0.2	0.3	0.6	1.8	3.0	8.4	28.0	130.0	686.6
Non-Hispanic black, both sexes												
2002	16.0	13.1	1.5	*	0.6	1.6	4.5	8.9	21.4	50.5	167.3	577.4
2001	15.9	16.9	1.0	*	0.8	1.7	4.9	9.9	19.0	48.7	169.1	581.5
2000	16.6	17.2	1.2	*	0.8	1.8	4.5	10.1	22.1	59.9	170.8	595.0
1999	16.5	19.3	1.7	*	0.9	2.0	5.4	10.7	20.7	54.0	156.6	628.9
Non-Hispanic black, male												
2002	15.8	13.9	1.6	*	*	1.8	5.2	11.5	29.3	66.0	230.6	693.6
2001	16.3	16.6	*	*	0.8	2.0	5.9	13.5	26.4	69.7	242.5	716.1
2000	17.1	18.8	1.7	*	0.7	2.3	5.7	14.0	30.1	83.4	233.9	750.3
1999	16.8	19.1	2.0	*	0.8	2.3	6.6	15.1	27.2	77.0	213.3	799.9
Non-Hispanic black, female												
2002	16.1	12.3	*	*	*	1.4	3.9	6.6	15.1	39.5	131.5	533.0
2001	15.6	17.1	*	*	0.8	1.3	4.0	6.9	13.0	33.9	127.9	531.7
2000	16.3	15.5	*	*	0.8	1.2	3.5	6.8	15.6	43.3	135.3	537.3
1999	16.3	19.6	*	*	0.9	1.7	4.4	6.9	15.5	37.7	124.8	565.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral pneumonia, not elsewhere classified (J12)												
All origins 3/, both sexes												
2002	0.1	1.4	*	*	*	*	*	*	*	0.2	0.5	1.7
2001	0.1	1.9	*	*	*	*	*	*	*	0.1	0.5	1.4
2000	0.1	1.4	*	*	*	*	*	*	0.1	0.3	0.7	2.9
1999	0.1	2.1	0.1	*	*	*	*	*	0.1	0.2	0.8	2.4
All origins 3/, male												
2002	0.1	1.8	*	*	*	*	*	*	*	*	0.7	1.9
2001	0.1	2.0	*	*	*	*	*	*	*	*	0.8	1.9
2000	0.1	1.7	*	*	*	*	*	*	*	0.2	0.8	2.4
1999	0.1	2.2	*	*	*	*	*	*	*	*	1.2	1.8
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2001	0.1	1.7	*	*	*	*	*	*	*	*	0.3	1.2
2000	0.2	1.2	*	*	*	*	*	*	*	0.3	0.7	3.0
1999	0.2	1.9	*	*	*	*	*	*	*	0.3	0.5	2.6
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	1.4	*	*	*	*	*	*	*	0.2	0.5	1.7
2001	0.1	1.9	*	*	*	*	*	*	*	0.1	0.5	1.5
2000	0.1	1.5	*	*	*	*	*	*	0.1	0.2	0.7	2.9
1999	0.2	2.0	*	*	*	*	*	*	0.1	0.2	0.8	2.4
Non-Hispanic 4/, male												
2002	0.1	1.8	*	*	*	*	*	*	*	*	0.7	2.0
2001	0.1	2.3	*	*	*	*	*	*	*	*	0.8	1.9
2000	0.1	1.7	*	*	*	*	*	*	*	*	0.8	2.5
1999	0.1	2.1	*	*	*	*	*	*	*	*	1.3	1.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral pneumonia, not elsewhere classified (J12)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.7
2001	0.1	1.5	*	*	*	*	*	*	*	*	0.3	1.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	3.0
1999	0.2	1.9	*	*	*	*	*	*	*	0.3	0.5	2.6
Non-Hispanic white, both sexes												
2002	0.1	1.1	*	*	*	*	*	*	*	0.2	0.6	1.8
2001	0.1	1.3	*	*	*	*	*	*	*	*	0.5	1.6
2000	0.2	1.1	*	*	*	*	*	*	0.1	0.2	0.8	3.1
1999	0.2	1.4	*	*	*	*	*	*	0.1	0.2	0.8	2.5
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	*	0.9	2.0
2000	0.1	*	*	*	*	*	*	*	*	*	0.9	2.8
1999	0.1	*	*	*	*	*	*	*	*	*	1.2	2.0
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	3.2
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.5	2.7
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	3.8	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bacterial pneumonia (J13-J15)												
All origins 3/, both sexes												
2002	1.6	1.1	*	*	*	0.1	0.3	0.7	1.3	3.6	11.3	35.7
2001	1.6	0.8	*	*	*	0.1	0.3	0.6	1.5	3.5	11.0	34.8
2000	1.6	0.8	*	*	*	0.1	0.4	0.7	1.6	3.7	11.0	35.7
1999	1.4	1.3	0.1	*	*	0.1	0.4	0.5	1.3	3.3	9.8	30.1
All origins 3/, male												
2002	1.6	1.2	*	*	*	0.1	0.3	0.7	1.6	4.4	15.1	46.8
2001	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	15.1	46.8
2000	1.7	1.1	*	*	*	0.1	0.4	0.8	1.8	4.8	15.6	52.3
1999	1.4	1.2	*	*	*	*	0.4	0.7	1.4	4.2	12.5	40.4
All origins 3/, female												
2002	1.6	*	*	*	*	0.1	0.3	0.6	1.1	2.9	8.8	31.0
2001	1.5	*	*	*	*	*	0.2	0.5	1.2	2.6	8.3	29.8
2000	1.5	*	*	*	*	0.1	0.3	0.6	1.3	2.9	8.0	29.0
1999	1.4	1.5	*	*	*	*	0.4	0.4	1.2	2.6	7.9	26.0
Hispanic, both sexes												
2002	0.7	*	*	*	*	*	*	0.9	1.4	3.0	10.0	41.0
2001	0.7	*	*	*	*	*	*	*	1.5	4.0	12.1	50.0
2000	0.8	*	*	*	*	*	*	0.7	2.0	4.1	11.7	45.8
1999	0.6	*	*	*	*	*	*	*	1.4	3.3	9.8	37.2
Hispanic, male												
2002	0.7	*	*	*	*	*	*	*	*	4.2	9.0	51.9
2001	0.8	*	*	*	*	*	*	*	*	6.4	15.2	49.5
2000	0.9	*	*	*	*	*	*	*	*	5.5	16.2	66.5
1999	0.6	*	*	*	*	*	*	*	*	*	13.6	*
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	*	*	10.7	35.6
2001	0.7	*	*	*	*	*	*	*	*	*	10.0	50.3
2000	0.7	*	*	*	*	*	*	*	*	*	8.6	35.6
1999	0.6	*	*	*	*	*	*	*	*	*	7.3	38.5
Non-Hispanic 4/, both sexes												
2002	1.7	1.0	*	*	*	0.1	0.3	0.6	1.3	3.7	11.3	35.3
2001	1.7	0.6	*	*	*	0.1	0.3	0.6	1.5	3.4	10.9	34.1
2000	1.7	0.9	*	*	*	0.1	0.4	0.7	1.5	3.7	10.9	35.3
1999	1.5	1.5	*	*	*	0.1	0.4	0.6	1.3	3.3	9.7	29.8
Non-Hispanic 4/, male												
2002	1.8	*	*	*	*	0.1	0.3	0.7	1.6	4.4	15.3	46.4
2001	1.7	*	*	*	*	0.1	0.3	0.7	1.7	4.4	15.1	46.5
2000	1.8	*	*	*	*	*	0.4	0.7	1.8	4.7	15.5	51.6
1999	1.5	*	*	*	*	*	0.4	0.7	1.3	4.2	12.5	40.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bacterial pneumonia (J13-J15)												
Non-Hispanic 4/, female												
2002	1.7	*	*	*	*	0.1	0.3	0.6	1.1	3.0	8.7	30.6
2001	1.6	*	*	*	*	*	0.2	0.6	1.2	2.6	8.2	29.0
2000	1.6	*	*	*	*	0.1	0.3	0.6	1.3	2.9	8.0	28.6
1999	1.5	1.8	*	*	*	*	0.4	0.4	1.2	2.5	8.0	25.5
Non-Hispanic white, both sexes												
2002	1.9	*	*	*	*	0.1	0.3	0.6	1.2	3.6	11.2	35.8
2001	1.8	*	*	*	*	0.1	0.2	0.6	1.4	3.3	10.8	34.2
2000	1.8	*	*	*	*	0.1	0.4	0.6	1.4	3.5	10.7	35.0
1999	1.6	1.0	*	*	*	0.1	0.3	0.5	1.1	3.2	9.7	29.9
Non-Hispanic white, male												
2002	1.9	*	*	*	*	*	0.3	0.6	1.4	4.3	14.9	47.5
2001	1.8	*	*	*	*	*	0.2	0.6	1.6	4.1	14.6	46.2
2000	1.9	*	*	*	*	*	0.4	0.6	1.6	4.4	15.2	50.2
1999	1.5	*	*	*	*	*	0.3	0.6	1.2	4.0	12.3	40.3
Non-Hispanic white, female												
2002	1.9	*	*	*	*	*	0.3	0.5	1.1	3.0	8.7	30.8
2001	1.8	*	*	*	*	*	0.2	0.6	1.2	2.6	8.3	29.2
2000	1.7	*	*	*	*	*	0.3	0.5	1.3	2.7	7.7	28.9
1999	1.6	*	*	*	*	*	0.4	0.4	1.1	2.5	8.0	25.8
Non-Hispanic black, both sexes												
2002	1.2	*	*	*	*	*	0.7	1.0	2.3	4.0	11.4	28.1
2001	1.3	*	*	*	*	*	0.6	1.0	2.2	4.7	12.5	33.1
2000	1.4	*	*	*	*	*	0.5	1.2	2.4	5.8	12.3	35.0
1999	1.2	*	*	*	*	*	0.8	1.0	2.4	4.7	9.7	24.9
Non-Hispanic black, male												
2002	1.3	*	*	*	*	*	*	1.2	3.3	5.0	17.9	31.7
2001	1.5	*	*	*	*	*	0.9	1.5	2.9	7.4	20.1	50.8
2000	1.5	*	*	*	*	*	*	1.3	3.4	7.9	17.9	61.1
1999	1.3	*	*	*	*	*	0.9	1.6	2.7	6.6	14.4	39.2
Non-Hispanic black, female												
2002	1.2	*	*	*	*	*	*	0.9	1.4	3.3	7.8	26.8
2001	1.0	*	*	*	*	*	*	*	1.7	2.7	8.3	26.5
2000	1.2	*	*	*	*	*	*	1.1	1.6	4.3	9.2	25.3
1999	1.1	*	*	*	*	*	0.8	*	2.2	3.4	7.0	19.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)												
All origins 3/, both sexes												
2002	20.8	3.9	0.5	0.2	0.3	0.7	1.8	4.1	9.7	33.4	143.7	650.3
2001	20.0	4.6	0.5	0.2	0.4	0.7	1.9	3.9	9.1	32.6	136.6	647.2
2000	20.8	5.1	0.4	0.1	0.4	0.7	1.9	3.9	9.8	34.0	144.6	685.3
1999	20.7	4.7	0.5	0.1	0.4	0.7	1.9	4.0	9.4	32.8	142.8	698.1
All origins 3/, male												
2002	18.5	4.6	0.5	0.1	0.4	0.8	2.1	5.0	11.8	41.4	177.4	748.2
2001	17.8	4.9	0.5	0.2	0.4	0.8	2.1	4.9	11.2	40.2	170.1	757.5
2000	18.5	5.3	0.4	0.2	0.4	0.9	2.3	4.8	12.1	42.0	179.5	816.5
1999	18.3	5.0	0.5	0.2	0.4	0.8	2.2	5.1	11.0	41.7	178.5	832.5
All origins 3/, female												
2002	23.0	3.1	0.4	0.2	0.3	0.6	1.5	3.2	7.8	26.7	121.3	607.9
2001	22.2	4.2	0.5	0.2	0.3	0.6	1.6	2.9	7.1	26.3	114.6	601.0
2000	23.1	4.9	0.4	0.1	0.4	0.6	1.6	3.0	7.7	27.4	121.9	631.9
1999	23.0	4.3	0.5	0.1	0.4	0.6	1.6	2.9	7.9	25.6	119.6	643.9
Hispanic, both sexes												
2002	6.5	2.9	*	*	*	0.4	1.1	2.4	9.0	30.5	131.3	521.6
2001	6.5	3.9	0.7	*	*	0.3	1.0	3.0	7.5	31.1	124.3	609.6
2000	6.4	5.6	*	*	*	0.6	1.2	2.9	9.0	32.4	121.9	584.6
1999	5.8	4.8	*	*	0.3	0.3	1.1	3.1	7.8	27.3	106.4	569.7
Hispanic, male												
2002	6.2	*	*	*	*	0.5	1.4	3.6	11.9	39.0	163.3	600.7
2001	6.0	4.8	*	*	*	*	1.3	4.1	9.2	34.1	157.4	714.2
2000	6.1	5.3	*	*	*	0.6	1.5	4.0	12.2	39.0	161.3	685.2
1999	5.4	6.2	*	*	*	*	1.6	4.1	7.9	35.6	133.5	683.7
Hispanic, female												
2002	6.8	*	*	*	*	*	0.8	1.3	6.4	23.7	109.3	482.4
2001	7.1	*	*	*	*	*	*	1.9	5.9	28.7	101.8	558.5
2000	6.8	5.8	*	*	*	*	0.8	1.9	6.2	27.2	95.4	535.2
1999	6.2	*	*	*	*	*	*	2.1	7.6	20.8	88.5	514.7
Non-Hispanic 4/, both sexes												
2002	23.0	4.1	0.5	0.2	0.4	0.8	1.9	4.2	9.7	33.5	143.8	654.2
2001	22.0	4.7	0.5	0.2	0.4	0.8	2.0	4.0	9.1	32.5	136.7	647.0
2000	22.8	4.9	0.4	0.1	0.4	0.8	2.0	3.9	9.8	33.9	145.0	687.1
1999	22.7	4.6	0.5	0.2	0.4	0.8	2.0	4.0	9.4	33.0	143.7	700.1
Non-Hispanic 4/, male												
2002	20.4	4.9	0.6	0.2	0.4	0.8	2.2	5.1	11.7	41.3	177.5	753.1
2001	19.6	4.9	0.5	0.2	0.5	0.9	2.2	4.9	11.2	40.2	170.2	757.5
2000	20.2	5.2	0.4	0.2	0.5	0.9	2.4	4.8	12.0	41.9	179.4	819.6
1999	20.1	4.8	0.5	0.2	0.4	0.9	2.2	5.1	11.1	41.8	179.6	834.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)												
Non-Hispanic 4/, female												
2002	25.4	3.3	0.5	0.2	0.3	0.7	1.6	3.4	7.9	26.9	121.4	611.5
2001	24.3	4.5	0.4	0.2	0.4	0.7	1.7	3.0	7.2	26.1	114.8	601.0
2000	25.3	4.7	0.4	0.1	0.4	0.7	1.6	3.1	7.8	27.3	122.6	633.5
1999	25.1	4.5	0.5	0.1	0.4	0.7	1.7	3.0	7.8	25.8	120.4	646.2
Non-Hispanic white, both sexes												
2002	25.6	2.9	0.4	0.2	0.3	0.7	1.7	3.8	8.9	32.6	143.6	665.4
2001	24.2	2.8	0.4	0.2	0.4	0.7	1.6	3.4	8.3	31.5	135.4	655.8
2000	25.2	3.3	0.3	0.1	0.4	0.7	1.7	3.3	8.7	31.9	144.4	700.0
1999	25.0	2.8	0.4	0.1	0.3	0.6	1.6	3.4	8.5	31.8	144.3	711.2
Non-Hispanic white, male												
2002	22.4	3.9	*	*	0.4	0.7	1.9	4.5	10.5	40.0	175.9	761.2
2001	21.1	3.2	*	0.2	0.4	0.7	1.8	4.1	9.9	38.3	166.6	763.4
2000	21.8	3.8	*	0.2	0.4	0.7	2.0	3.9	10.4	38.7	177.2	833.0
1999	21.7	3.0	*	0.2	0.3	0.7	1.7	4.2	9.8	39.5	179.2	844.5
Non-Hispanic white, female												
2002	28.6	1.8	*	0.2	0.3	0.6	1.4	3.1	7.4	26.3	121.9	624.1
2001	27.2	2.4	*	0.2	0.3	0.6	1.4	2.6	6.7	25.7	114.7	611.1
2000	28.3	2.7	*	*	0.3	0.6	1.4	2.7	7.1	26.1	122.9	646.3
1999	28.2	2.5	0.5	*	0.3	0.5	1.4	2.6	7.2	25.3	121.5	658.1
Non-Hispanic black, both sexes												
2002	14.6	8.5	1.0	*	0.5	1.3	3.8	7.8	19.2	46.4	155.7	548.4
2001	14.5	12.1	*	*	0.7	1.4	4.3	8.9	16.7	43.9	156.4	548.1
2000	15.2	11.8	0.8	*	0.7	1.6	4.0	8.9	19.5	53.9	158.3	559.7
1999	15.1	12.1	1.1	*	0.8	1.8	4.6	9.6	18.3	49.1	146.4	602.7
Non-Hispanic black, male												
2002	14.4	8.5	*	*	*	1.6	4.5	10.3	26.0	61.1	212.2	659.7
2001	14.7	11.5	*	*	0.7	1.9	5.0	11.9	23.5	62.0	222.4	664.1
2000	15.4	10.9	*	*	*	2.1	5.1	12.7	26.5	75.3	216.0	689.3
1999	15.2	12.8	*	*	0.8	2.1	5.7	13.5	24.5	70.3	197.3	759.4
Non-Hispanic black, female												
2002	14.9	8.5	*	*	*	1.1	3.2	5.7	13.7	36.0	123.7	505.8
2001	14.4	12.8	*	*	0.7	1.0	3.7	6.3	11.3	31.1	119.4	505.2
2000	15.0	12.7	*	*	*	1.2	3.0	5.7	14.0	38.8	125.9	511.5
1999	15.0	11.4	*	*	0.8	1.5	3.6	6.4	13.3	34.1	117.8	544.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
All origins 3/, both sexes												
2002	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
All origins 3/, male												
2002	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	1.3	0.2	*	*	*	*	*	*	0.2	0.5	3.4
2001	0.1	1.3	*	*	*	*	*	*	*	0.2	0.4	3.3
2000	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.1
Non-Hispanic 4/, male												
2002	0.1	2.1	*	*	*	*	*	*	*	*	0.6	2.9
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	2.4
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.7
1999	0.2	2.0	*	*	*	*	*	*	*	*	1.1	4.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.6
2001	0.2	1.4	*	*	*	*	*	*	*	*	0.5	3.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.7	4.8
1999	0.3	1.5	*	*	*	*	*	*	*	*	0.8	6.9
Non-Hispanic white, both sexes												
2002	0.2	1.1	*	*	*	*	*	*	*	0.2	0.5	3.5
2001	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	4.7
1999	0.2	1.1	*	*	*	*	*	*	*	0.2	0.9	6.4
Non-Hispanic white, male												
2002	0.1	1.8	*	*	*	*	*	*	*	*	0.6	3.0
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	3.7
1999	0.2	*	*	*	*	*	*	*	*	*	1.1	4.1
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	5.1
1999	0.3	*	*	*	*	*	*	*	*	*	0.8	7.3
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
All origins 3/, both sexes												
2002	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6
All origins 3/, male												
2002	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2001	0.1	1.3	*	*	*	*	*	*	*	0.1	0.3	2.1
2000	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999	0.1	1.7	0.2	*	*	*	*	*	*	0.2	0.6	3.6
Non-Hispanic 4/, male												
2002	0.1	2.0	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
1999	0.1	2.0	*	*	*	*	*	*	*	*	0.7	2.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
1999	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0
Non-Hispanic white, both sexes												
2002	0.1	1.0	*	*	*	*	*	*	*	0.1	0.3	2.1
2001	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.2
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	2.7
1999	0.2	1.1	*	*	*	*	*	*	*	0.1	0.6	3.9
Non-Hispanic white, male												
2002	0.1	1.7	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	2.3
1999	0.1	*	*	*	*	*	*	*	*	*	0.8	2.6
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	2.8
1999	0.2	*	*	*	*	*	*	*	*	*	0.5	4.4
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.2	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	3.0
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of upper respiratory tract (J30-J39)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	2.1
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	0.9	2.4
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	0.9	2.1
1999	0.2	*	*	*	*	*	0.0	0.1	0.2	0.4	1.0	2.4
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.7	2.5
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.2	3.4
2000	0.2	*	*	*	*	*	*	0.1	*	0.5	1.1	3.1
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	3.5
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	0.2	0.3	0.6	1.9
2001	0.2	*	*	*	*	*	*	0.1	*	0.3	0.8	1.9
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.7	1.7
1999	0.2	*	*	*	*	*	*	*	0.2	0.3	0.8	2.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.7	2.1
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	0.9	2.3
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	0.9	2.2
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	2.4
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	2.6
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.1	3.3
2000	0.2	*	*	*	*	*	*	0.1	*	0.5	1.2	3.1
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.1	3.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of upper respiratory tract (J30-J39)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	0.2	0.3	0.6	1.9
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	1.9
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.7	1.8
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.9	2.0
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.6	2.2
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	2.3
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.2
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.9	2.4
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.7	2.7
2001	0.2	*	*	*	*	*	*	*	*	*	1.0	3.1
2000	0.2	*	*	*	*	*	*	0.1	*	0.5	1.2	3.0
1999	0.2	*	*	*	*	*	*	*	*	0.5	1.2	3.2
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	0.2	0.3	0.6	2.0
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	1.9
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	1.9
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.8	2.1
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
All origins 3/, both sexes												
2002	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0
All origins 3/, male												
2002	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7
All origins 3/, female												
2002	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6
Hispanic, both sexes												
2002	7.9	*	*	*	*	0.4	1.2	3.5	13.9	53.8	176.0	457.3
2001	7.7	*	*	*	*	0.3	1.0	3.6	14.0	55.8	169.5	480.2
2000	7.6	*	*	*	*	0.3	1.1	3.3	14.3	53.1	181.9	481.1
1999	8.4	*	*	*	*	0.5	1.6	3.9	14.6	62.6	194.8	536.0
Hispanic, male												
2002	8.1	*	*	*	*	*	1.0	3.8	17.5	69.2	243.3	602.4
2001	7.8	*	*	*	*	*	0.7	3.2	16.1	75.5	224.0	676.1
2000	8.0	*	*	*	*	*	0.9	3.4	18.2	72.4	250.3	671.1
1999	8.9	*	*	*	*	*	1.4	3.6	17.6	81.9	272.7	836.8
Hispanic, female												
2002	7.6	*	*	*	*	*	1.4	3.1	10.6	41.5	129.8	385.5
2001	7.5	*	*	*	*	*	1.2	4.1	12.1	40.3	132.7	384.4
2000	7.2	*	*	*	*	*	1.3	3.3	10.8	38.0	136.0	387.8
1999	8.0	*	*	*	*	*	1.8	4.2	12.0	47.5	142.9	391.0
Non-Hispanic 4/, both sexes												
2002	48.6	1.1	0.4	0.4	0.5	0.8	2.4	9.2	44.6	170.1	396.0	643.5
2001	48.4	1.1	0.3	0.3	0.5	0.8	2.3	8.9	46.3	174.8	388.6	649.4
2000	48.3	1.0	0.3	0.4	0.5	0.8	2.2	9.0	46.3	176.1	393.8	653.2
1999	49.4	1.0	0.4	0.4	0.6	0.9	2.0	8.9	49.8	183.5	405.1	648.0
Non-Hispanic 4/, male												
2002	48.4	1.2	0.6	0.4	0.6	0.9	2.4	9.6	47.3	191.9	491.2	906.0
2001	48.0	*	0.4	0.3	0.6	0.8	2.2	9.3	49.0	198.1	485.7	924.8
2000	48.7	*	0.4	0.4	0.6	0.9	2.1	9.4	49.7	201.9	497.6	978.0
1999	50.8	*	0.4	0.5	0.7	0.8	1.9	9.0	54.5	217.4	521.9	1,000.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Non-Hispanic 4/, female												
2002	48.9	*	*	0.3	0.4	0.8	2.4	8.7	42.0	151.8	332.9	530.3
2001	48.7	*	*	0.2	0.4	0.8	2.4	8.5	43.8	155.4	324.9	534.8
2000	48.0	*	*	0.3	0.4	0.7	2.3	8.7	43.2	154.8	326.2	521.9
1999	48.0	*	*	0.2	0.5	0.9	2.1	8.7	45.4	155.7	329.3	507.0
Non-Hispanic white, both sexes												
2002	56.4	0.9	0.2	0.2	0.4	0.6	2.0	9.0	46.7	181.6	416.3	673.1
2001	55.9	*	*	0.2	0.3	0.5	1.9	8.7	48.6	185.9	408.6	676.9
2000	55.7	*	*	0.2	0.4	0.6	1.8	8.6	48.2	186.5	413.2	680.8
1999	56.6	*	0.2	0.2	0.4	0.6	1.6	8.5	51.8	193.6	422.4	675.6
Non-Hispanic white, male												
2002	55.1	*	*	0.2	0.4	0.6	2.0	9.3	48.3	200.4	506.7	935.4
2001	54.6	*	*	0.2	0.3	0.5	1.9	9.1	50.5	206.1	500.9	951.5
2000	55.1	*	*	0.2	0.4	0.5	1.7	8.9	50.8	208.8	513.6	1,008.6
1999	57.3	*	*	0.3	0.5	0.5	1.5	8.5	55.6	224.9	534.9	1,033.5
Non-Hispanic white, female												
2002	57.7	*	*	*	0.4	0.6	2.1	8.6	45.1	165.5	355.7	559.8
2001	57.2	*	*	*	0.3	0.6	2.0	8.3	46.8	168.8	347.3	562.7
2000	56.2	*	*	0.2	0.3	0.6	1.9	8.3	45.8	167.6	347.2	548.7
1999	55.9	*	*	*	0.4	0.6	1.8	8.5	48.3	167.4	348.8	532.8
Non-Hispanic black, both sexes												
2002	21.4	*	1.2	1.1	1.3	2.0	5.2	13.1	41.6	115.0	250.5	370.8
2001	21.1	*	0.9	0.8	1.3	2.5	4.9	12.7	41.3	120.4	236.3	379.3
2000	21.5	*	0.9	1.1	1.4	2.2	4.9	14.5	44.7	123.2	236.5	376.2
1999	22.6	*	*	1.0	1.4	2.7	4.7	14.1	46.7	129.9	265.3	377.0
Non-Hispanic black, male												
2002	24.9	*	1.6	1.2	2.0	2.3	5.7	14.6	53.1	160.0	395.7	653.1
2001	24.5	*	*	1.0	1.7	2.8	4.7	13.5	50.4	170.1	385.6	682.6
2000	25.1	*	*	1.3	1.8	2.8	4.9	15.2	55.3	178.9	372.4	696.3
1999	27.0	*	*	1.4	1.9	2.7	4.7	15.6	60.4	186.1	438.4	707.2
Non-Hispanic black, female												
2002	18.2	*	*	1.1	0.7	1.7	4.8	11.8	32.3	83.0	168.5	263.0
2001	18.0	*	*	0.7	1.0	2.2	5.0	11.9	34.0	85.3	152.5	266.9
2000	18.2	*	*	0.8	1.0	1.6	4.9	13.8	36.1	84.1	159.9	257.3
1999	18.6	*	*	0.6	0.9	2.7	4.7	12.8	35.8	90.3	168.4	253.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
All origins 3/, both sexes												
2002	42.9	1.0	0.4	0.3	0.5	0.7	2.2	8.6	42.1	162.1	383.3	631.8
2001	42.9	1.0	0.3	0.3	0.4	0.7	2.1	8.4	43.9	167.0	376.9	637.7
2000	43.0	0.9	0.3	0.3	0.5	0.7	2.0	8.6	44.0	168.5	382.8	642.4
1999	44.2	0.9	0.3	0.3	0.5	0.8	2.0	8.4	47.2	176.3	394.4	639.8
All origins 3/, male												
2002	42.7	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.0	184.2	478.6	890.1
2001	42.5	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.6	190.6	473.2	912.1
2000	43.3	1.2	0.4	0.4	0.6	0.8	1.9	8.9	47.5	194.5	486.4	963.6
1999	45.4	*	0.4	0.4	0.6	0.7	1.8	8.6	52.0	210.1	510.1	990.1
All origins 3/, female												
2002	43.2	*	0.3	0.3	0.4	0.7	2.2	8.2	39.4	143.7	320.0	519.7
2001	43.2	*	*	0.2	0.4	0.7	2.2	8.1	41.3	147.3	313.6	522.9
2000	42.8	*	0.3	0.3	0.4	0.6	2.1	8.2	40.6	147.1	315.2	511.6
1999	43.0	*	0.3	0.2	0.5	0.9	2.1	8.2	42.8	148.6	319.3	498.6
Hispanic, both sexes												
2002	7.8	*	*	*	*	0.4	1.2	3.4	13.5	53.1	174.0	454.0
2001	7.6	*	*	*	*	0.3	0.9	3.5	13.9	54.8	167.4	477.7
2000	7.5	*	*	*	*	*	1.0	3.3	14.0	52.5	178.6	476.4
1999	8.3	*	*	*	*	0.5	1.6	3.7	14.5	61.9	192.1	531.8
Hispanic, male												
2002	8.1	*	*	*	*	*	1.0	3.8	17.3	68.6	241.7	599.0
2001	7.7	*	*	*	*	*	0.7	3.1	16.1	74.1	223.1	672.3
2000	7.9	*	*	*	*	*	0.8	3.4	17.9	72.2	247.3	669.1
1999	8.8	*	*	*	*	*	1.4	3.6	17.6	81.7	269.0	836.8
Hispanic, female												
2002	7.5	*	*	*	*	*	1.4	3.1	10.1	40.8	127.6	382.1
2001	7.4	*	*	*	*	*	1.1	4.0	12.0	39.5	129.7	382.6
2000	7.0	*	*	*	*	*	1.2	3.2	10.5	37.0	132.4	381.8
1999	7.8	*	*	*	*	*	1.8	3.9	11.7	46.3	140.9	384.7
Non-Hispanic 4/, both sexes												
2002	48.3	1.1	0.4	0.4	0.5	0.8	2.4	9.1	44.3	169.1	392.6	637.6
2001	48.0	1.0	0.3	0.3	0.5	0.8	2.3	8.8	46.0	173.9	385.7	642.2
2000	48.0	1.0	0.3	0.4	0.5	0.8	2.2	9.0	46.0	175.1	390.6	647.0
1999	49.0	1.0	0.4	0.3	0.6	0.8	2.0	8.8	49.5	182.6	401.6	641.8
Non-Hispanic 4/, male												
2002	48.2	1.2	0.6	0.4	0.6	0.9	2.4	9.6	47.1	191.3	489.0	901.2
2001	47.8	*	0.4	0.3	0.5	0.8	2.2	9.3	48.7	197.5	483.8	919.9
2000	48.5	*	0.4	0.4	0.6	0.9	2.1	9.3	49.5	201.1	495.4	973.6
1999	50.6	*	0.4	0.5	0.7	0.8	1.9	9.0	54.3	216.7	518.9	993.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)												
Non-Hispanic 4/, female												
2002	48.3	*	*	0.3	0.4	0.7	2.4	8.6	41.7	150.7	328.7	523.9
2001	48.1	*	*	0.2	0.4	0.8	2.4	8.4	43.5	154.2	321.2	526.7
2000	47.5	*	*	0.3	0.4	0.7	2.2	8.6	42.8	153.7	322.3	515.0
1999	47.5	*	*	0.2	0.5	0.9	2.1	8.6	45.0	154.6	325.6	501.0
Non-Hispanic white, both sexes												
2002	56.0	0.9	0.2	0.2	0.4	0.6	2.0	8.9	46.4	180.7	412.6	666.9
2001	55.5	*	*	0.2	0.3	0.5	1.9	8.6	48.3	184.9	405.5	669.3
2000	55.3	*	*	0.2	0.3	0.5	1.7	8.5	47.9	185.5	409.8	674.2
1999	56.2	*	0.2	0.2	0.4	0.5	1.6	8.4	51.5	192.6	418.9	668.8
Non-Hispanic white, male												
2002	54.9	*	*	0.2	0.4	0.6	1.9	9.3	48.1	199.9	504.4	930.6
2001	54.4	*	*	0.2	0.3	0.5	1.8	9.0	50.2	205.5	499.2	946.6
2000	54.9	*	*	0.2	0.4	0.5	1.6	8.8	50.5	208.1	511.4	1,003.9
1999	57.1	*	*	0.3	0.5	0.5	1.5	8.4	55.4	224.2	531.8	1,026.2
Non-Hispanic white, female												
2002	57.0	*	*	*	0.4	0.6	2.0	8.6	44.8	164.3	351.2	553.0
2001	56.5	*	*	*	0.3	0.6	2.0	8.2	46.5	167.5	343.3	553.9
2000	55.6	*	*	0.2	0.3	0.6	1.9	8.3	45.5	166.5	343.0	541.3
1999	55.4	*	*	*	0.4	0.6	1.8	8.4	47.9	166.3	345.0	526.3
Non-Hispanic black, both sexes												
2002	21.3	*	1.2	1.1	1.3	2.0	5.2	12.9	41.5	114.4	249.6	369.6
2001	21.0	*	0.9	0.8	1.3	2.4	4.9	12.6	41.2	120.1	235.2	378.0
2000	21.4	*	0.9	1.1	1.4	2.2	4.9	14.4	44.3	122.4	235.9	375.5
1999	22.5	*	*	1.0	1.4	2.6	4.7	14.0	46.5	129.4	263.8	376.0
Non-Hispanic black, male												
2002	24.8	*	1.6	1.2	2.0	2.3	5.7	14.5	52.9	159.1	394.5	653.1
2001	24.4	*	*	1.0	1.7	2.8	4.7	13.5	50.2	169.5	383.5	680.3
2000	25.0	*	*	1.3	1.8	2.8	4.9	15.2	55.1	177.6	371.8	696.3
1999	26.9	*	*	1.4	1.9	2.7	4.6	15.5	60.0	185.8	436.8	706.0
Non-Hispanic black, female												
2002	18.1	*	*	1.1	0.7	1.7	4.8	11.6	32.2	82.7	167.6	261.3
2001	17.9	*	*	0.7	1.0	2.1	5.0	11.8	33.9	85.2	152.0	266.1
2000	18.1	*	*	0.8	1.0	1.6	4.8	13.7	35.7	83.6	159.4	256.4
1999	18.4	*	*	0.6	0.9	2.6	4.7	12.7	35.6	89.6	166.9	252.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, not specified as acute or chronic (J40)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.1	0.3	1.0	5.2
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.1	5.2
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.5	6.8
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	6.2
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	1.0	4.7
2001	0.1	*	*	*	*	*	*	*	*	0.4	1.1	5.1
2000	0.2	*	*	*	*	*	*	*	0.2	0.4	2.0	6.4
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	5.9
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.3	1.0	5.4
2001	0.2	*	*	*	*	*	*	*	*	0.3	1.1	5.3
2000	0.3	*	*	*	*	*	*	*	*	0.3	1.1	7.0
1999	0.3	*	*	*	*	*	*	*	0.2	0.5	1.3	6.3
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.1	0.3	1.0	5.1
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.1	5.3
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.4	1.5	7.0
1999	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.4	6.3
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	1.0	4.8
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.1	5.2
2000	0.2	*	*	*	*	*	*	*	0.2	0.4	2.0	6.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	5.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, not specified as acute or chronic (J40)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.3	1.0	5.2
2001	0.2	*	*	*	*	*	*	*	*	0.3	1.1	5.3
2000	0.3	*	*	*	*	*	*	*	*	0.3	1.2	7.1
1999	0.3	*	*	*	*	*	*	*	0.2	0.5	1.3	6.4
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.1	0.4	1.1	5.4
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.1	5.6
2000	0.3	*	*	*	*	*	*	*	0.2	0.4	1.6	7.5
1999	0.3	*	*	*	*	*	*	*	0.2	0.5	1.4	6.5
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.1	5.2
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.0	5.4
2000	0.2	*	*	*	*	*	*	*	*	0.4	2.1	7.2
1999	0.2	*	*	*	*	*	*	*	*	0.5	1.5	6.3
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	*	*	0.4	1.1	5.4
2001	0.3	*	*	*	*	*	*	*	*	0.3	1.2	5.7
2000	0.3	*	*	*	*	*	*	*	*	0.3	1.2	7.6
1999	0.3	*	*	*	*	*	*	*	*	0.5	1.3	6.6
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic bronchitis (J41-J42)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.1	0.4	1.2	2.9
2001	0.2	*	*	*	*	*	*	*	0.1	0.4	1.0	3.9
2000	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	5.0
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	4.5
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	*	0.6	1.5	4.0
2001	0.1	*	*	*	*	*	*	*	*	0.6	1.2	4.1
2000	0.2	*	*	*	*	*	*	*	*	0.6	1.6	6.4
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.7	5.9
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	1.0	2.4
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	3.8
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.0	4.4
1999	0.2	*	*	*	*	*	*	*	0.2	0.3	1.1	4.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.1	0.5	1.2	2.9
2001	0.2	*	*	*	*	*	*	*	0.1	0.4	1.0	3.9
2000	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	4.8
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	4.4
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	*	0.6	1.5	4.0
2001	0.1	*	*	*	*	*	*	*	*	0.6	1.2	4.1
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.6	6.2
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.7	5.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic bronchitis (J41-J42)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.4
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	3.8
2000	0.2	*	*	*	*	*	*	*	*	0.5	0.9	4.3
1999	0.2	*	*	*	*	*	*	*	0.2	0.3	1.1	3.8
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	3.1
2001	0.2	*	*	*	*	*	*	*	0.2	0.4	1.0	4.0
2000	0.2	*	*	*	*	*	*	*	0.1	0.5	1.2	5.1
1999	0.2	*	*	*	*	*	*	*	0.3	0.5	1.4	4.7
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	*	0.6	1.5	4.2
2001	0.2	*	*	*	*	*	*	*	*	0.6	1.2	4.2
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.7	6.6
1999	0.2	*	*	*	*	*	*	*	0.3	0.7	1.7	6.1
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.6
2001	0.2	*	*	*	*	*	*	*	*	*	0.9	4.0
2000	0.2	*	*	*	*	*	*	*	*	0.5	0.9	4.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.3	1.1	4.1
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All origins 3/, both sexes												
2002	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
All origins 3/, male												
2002	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
All origins 3/, female												
2002	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7
Hispanic, both sexes												
2002	0.8	*	*	*	*	*	*	*	1.5	6.7	18.8	36.6
2001	0.8	*	*	*	*	*	*	*	1.4	6.6	20.4	39.4
2000	0.8	*	*	*	*	*	*	*	1.5	6.0	22.3	31.2
1999	0.9	*	*	*	*	*	*	*	1.3	7.1	23.5	44.2
Hispanic, male												
2002	0.8	*	*	*	*	*	*	*	*	9.2	26.7	53.5
2001	0.9	*	*	*	*	*	*	*	*	9.2	34.0	60.9
2000	0.9	*	*	*	*	*	*	*	*	9.1	32.5	58.4
1999	1.0	*	*	*	*	*	*	*	*	11.2	36.0	77.6
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	*	4.7	13.5	28.2
2001	0.6	*	*	*	*	*	*	*	*	4.6	11.2	28.9
2000	0.6	*	*	*	*	*	*	*	*	3.6	15.5	*
1999	0.7	*	*	*	*	*	*	*	*	4.0	15.3	28.1
Non-Hispanic 4/, both sexes												
2002	6.1	*	*	*	*	*	0.2	1.2	6.8	25.1	49.3	60.1
2001	6.4	*	*	*	*	*	0.2	1.2	7.6	27.5	51.7	62.8
2000	6.7	*	*	*	*	*	0.2	1.3	7.9	29.1	53.8	64.3
1999	7.1	*	*	*	*	*	0.2	1.3	9.0	32.0	58.1	62.5
Non-Hispanic 4/, male												
2002	6.5	*	*	*	*	*	0.3	1.5	7.7	29.3	65.2	91.3
2001	6.8	*	*	*	*	*	0.3	1.5	8.4	32.1	66.6	102.7
2000	7.1	*	*	*	*	*	0.2	1.6	8.5	34.3	70.6	105.9
1999	7.8	*	*	*	*	*	0.2	1.5	10.0	39.2	77.8	106.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Non-Hispanic 4/, female												
2002	5.7	*	*	*	*	*	0.1	1.0	6.0	21.7	38.8	46.6
2001	6.1	*	*	*	*	*	0.1	1.0	7.0	23.6	41.9	46.3
2000	6.3	*	*	*	*	*	0.2	1.1	7.3	24.8	42.9	47.5
1999	6.5	*	*	*	*	*	0.2	1.1	8.1	26.1	45.3	45.0
Non-Hispanic white, both sexes												
2002	7.1	*	*	*	*	*	0.2	1.3	7.3	27.0	52.3	63.1
2001	7.5	*	*	*	*	*	0.2	1.3	8.2	29.6	54.9	65.9
2000	7.8	*	*	*	*	*	0.2	1.3	8.4	31.2	57.0	66.9
1999	8.3	*	*	*	*	*	0.2	1.3	9.7	34.5	61.2	65.2
Non-Hispanic white, male												
2002	7.4	*	*	*	*	*	0.3	1.5	8.0	30.6	67.7	94.3
2001	7.8	*	*	*	*	*	0.3	1.5	8.7	33.8	69.6	106.6
2000	8.1	*	*	*	*	*	0.2	1.4	8.9	35.8	73.9	107.3
1999	8.9	*	*	*	*	*	0.2	1.4	10.4	41.1	80.6	110.0
Non-Hispanic white, female												
2002	6.8	*	*	*	*	*	*	1.1	6.7	24.0	41.9	49.7
2001	7.2	*	*	*	*	*	0.1	1.0	7.7	26.1	45.2	48.9
2000	7.4	*	*	*	*	*	0.2	1.1	8.0	27.3	45.8	50.6
1999	7.7	*	*	*	*	*	0.2	1.2	9.0	29.0	48.5	47.3
Non-Hispanic black, both sexes												
2002	2.3	*	*	*	*	*	0.5	1.2	4.6	16.2	28.2	32.1
2001	2.2	*	*	*	*	*	*	1.4	5.6	15.6	23.6	35.9
2000	2.5	*	*	*	*	*	*	2.0	6.1	17.1	27.0	37.5
1999	2.7	*	*	*	*	*	0.4	1.9	6.3	17.3	31.8	37.8
Non-Hispanic black, male												
2002	3.1	*	*	*	*	*	*	1.8	6.7	25.3	49.5	65.6
2001	2.8	*	*	*	*	*	*	1.6	7.9	23.0	40.2	69.3
2000	3.3	*	*	*	*	*	*	3.0	8.2	27.4	41.3	89.2
1999	3.6	*	*	*	*	*	*	3.0	9.1	29.3	55.7	73.7
Non-Hispanic black, female												
2002	1.6	*	*	*	*	*	*	*	2.8	9.8	16.1	19.2
2001	1.7	*	*	*	*	*	*	1.1	3.7	10.4	14.3	23.5
2000	1.8	*	*	*	*	*	*	1.3	4.5	9.8	18.9	18.3
1999	1.8	*	*	*	*	*	*	1.0	4.0	8.9	18.4	24.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)												
All origins 3/, both sexes												
2002	35.8	*	*	*	*	0.1	0.9	5.9	33.5	134.2	326.7	549.7
2001	35.3	*	*	*	0.1	0.1	0.9	5.6	34.2	136.5	318.0	550.3
2000	35.1	*	*	*	*	0.1	0.8	5.6	34.0	135.8	320.6	550.6
1999	35.7	*	*	*	*	0.1	0.7	5.4	35.9	140.5	327.2	549.6
All origins 3/, male												
2002	35.5	*	*	*	*	0.1	1.0	6.4	36.0	152.8	407.6	780.4
2001	35.1	*	*	*	*	0.1	0.9	6.2	36.9	156.3	401.1	789.6
2000	35.4	*	*	*	*	*	0.8	6.0	37.5	157.5	408.7	833.7
1999	36.9	*	*	*	*	*	0.7	5.9	40.3	167.9	424.8	860.0
All origins 3/, female												
2002	36.0	*	*	*	*	0.1	0.9	5.4	31.2	118.7	273.0	449.5
2001	35.5	*	*	*	*	*	0.9	5.1	31.7	120.2	263.4	450.2
2000	34.7	*	*	*	*	*	0.8	5.2	30.8	117.9	263.1	435.3
1999	34.5	*	*	*	*	0.1	0.7	4.9	31.8	118.0	263.8	424.5
Hispanic, both sexes												
2002	6.2	*	*	*	*	*	0.4	2.0	10.0	42.4	145.8	394.1
2001	6.0	*	*	*	*	*	*	1.8	10.0	44.1	137.4	416.4
2000	5.8	*	*	*	*	*	*	1.6	9.6	40.0	145.8	418.0
1999	6.4	*	*	*	*	*	*	1.7	10.7	48.5	155.4	458.1
Hispanic, male												
2002	6.6	*	*	*	*	*	*	2.2	13.8	57.3	207.2	532.1
2001	6.2	*	*	*	*	*	*	2.0	12.2	61.1	180.6	601.8
2000	6.1	*	*	*	*	*	*	1.6	13.3	56.3	206.0	582.5
1999	7.0	*	*	*	*	*	*	1.6	13.9	65.7	225.2	735.4
Hispanic, female												
2002	5.7	*	*	*	*	*	*	1.8	6.6	30.6	103.7	325.8
2001	5.7	*	*	*	*	*	*	1.7	8.0	30.7	108.2	325.8
2000	5.4	*	*	*	*	*	*	1.6	6.3	27.2	105.3	337.3
1999	5.7	*	*	*	*	*	*	1.8	7.8	35.1	108.9	324.4
Non-Hispanic 4/, both sexes												
2002	40.3	*	*	*	*	0.1	1.0	6.2	35.3	140.1	334.7	554.9
2001	39.6	*	*	*	0.1	0.1	1.0	5.9	35.9	142.2	325.6	554.1
2000	39.2	*	*	*	*	0.1	0.8	5.9	35.7	141.3	327.2	554.3
1999	39.6	*	*	*	*	0.1	0.8	5.7	37.6	145.5	333.3	551.4
Non-Hispanic 4/, male												
2002	40.2	*	*	*	*	0.1	1.0	6.8	37.6	158.6	416.3	790.1
2001	39.6	*	*	*	*	0.1	1.0	6.5	38.6	161.9	410.5	795.7
2000	39.7	*	*	*	*	*	0.9	6.4	39.0	163.0	416.3	842.2
1999	41.1	*	*	*	*	*	0.8	6.2	42.0	173.2	432.1	862.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)												
Non-Hispanic 4/, female												
2002	40.3	*	*	*	*	*	0.9	5.7	33.1	124.6	280.6	453.4
2001	39.6	*	*	*	*	*	1.0	5.4	33.5	125.9	269.9	453.6
2000	38.6	*	*	*	*	*	0.8	5.4	32.6	123.4	269.2	437.9
1999	38.2	*	*	*	*	0.1	0.7	5.1	33.5	122.7	269.2	426.7
Non-Hispanic white, both sexes												
2002	47.1	*	*	*	*	0.1	1.0	6.5	37.4	150.2	352.0	580.7
2001	46.2	*	*	*	*	0.1	1.0	6.1	38.3	151.7	342.5	578.0
2000	45.6	*	*	*	*	*	0.8	6.0	37.7	150.3	343.6	578.9
1999	45.9	*	*	*	*	0.1	0.7	5.8	39.5	153.8	347.9	576.0
Non-Hispanic white, male												
2002	46.2	*	*	*	*	*	1.1	6.9	38.8	166.4	429.5	816.2
2001	45.3	*	*	*	*	*	1.0	6.6	40.2	168.7	423.4	819.0
2000	45.4	*	*	*	*	*	0.9	6.3	40.3	169.1	429.2	871.8
1999	46.8	*	*	*	*	*	0.7	6.2	43.1	179.5	442.8	893.3
Non-Hispanic white, female												
2002	48.0	*	*	*	*	*	1.0	6.1	36.0	136.4	300.0	478.9
2001	46.9	*	*	*	*	*	1.0	5.7	36.5	137.2	288.9	477.8
2000	45.7	*	*	*	*	*	0.8	5.8	35.2	134.4	287.4	460.8
1999	45.0	*	*	*	*	*	0.7	5.4	36.2	132.3	285.7	449.5
Non-Hispanic black, both sexes												
2002	15.8	*	*	*	*	*	1.2	6.6	30.1	89.8	212.1	321.5
2001	15.5	*	*	*	*	*	1.3	6.5	28.5	95.8	199.7	322.1
2000	15.3	*	*	*	*	*	1.2	6.7	31.1	95.0	196.2	312.0
1999	16.3	*	*	*	*	*	1.2	6.4	32.9	101.3	218.5	313.0
Non-Hispanic black, male												
2002	18.6	*	*	*	*	*	1.3	8.3	40.2	125.8	336.7	572.1
2001	18.7	*	*	*	*	*	1.4	7.5	37.4	138.5	332.8	590.2
2000	18.7	*	*	*	*	*	1.3	8.4	40.5	141.3	321.8	578.9
1999	20.2	*	*	*	*	*	1.4	8.2	45.0	146.2	367.7	609.7
Non-Hispanic black, female												
2002	13.2	*	*	*	*	*	1.1	5.1	22.1	64.3	141.6	225.7
2001	12.6	*	*	*	*	*	1.2	5.6	21.4	65.7	125.0	222.9
2000	12.3	*	*	*	*	*	1.2	5.3	23.5	62.4	125.6	212.8
1999	12.8	*	*	*	*	*	1.0	4.9	23.2	69.7	135.0	202.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
All origins 3/, both sexes												
2002	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
All origins 3/, male												
2002	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
All origins 3/, female												
2002	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2
Hispanic, both sexes												
2002	0.7	*	*	*	*	0.3	0.7	1.2	1.9	3.8	7.0	12.2
2001	0.7	*	*	*	*	*	0.6	1.4	2.4	3.8	7.9	13.8
2000	0.8	*	*	*	*	*	0.6	1.3	2.8	5.6	7.9	16.6
1999	0.9	*	*	*	*	0.3	0.9	1.6	2.4	5.8	10.4	17.5
Hispanic, male												
2002	0.6	*	*	*	*	*	*	1.2	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	*	1.3	*	5.3	*	*
1999	0.7	*	*	*	*	*	0.9	1.4	*	*	*	*
Hispanic, female												
2002	0.9	*	*	*	*	*	1.0	1.2	2.2	5.3	7.6	*
2001	1.0	*	*	*	*	*	*	2.0	2.9	3.9	9.4	18.6
2000	1.0	*	*	*	*	*	*	1.3	3.2	5.8	9.2	*
1999	1.2	*	*	*	*	*	1.0	1.9	3.1	7.1	13.2	*
Non-Hispanic 4/, both sexes												
2002	1.6	*	0.3	0.3	0.5	0.6	1.1	1.5	2.0	3.1	6.3	14.7
2001	1.6	*	0.2	0.3	0.4	0.7	1.1	1.5	2.2	3.4	6.3	16.1
2000	1.7	*	0.2	0.3	0.5	0.7	1.1	1.6	2.2	3.9	6.8	16.6
1999	1.8	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.1	7.4	17.3
Non-Hispanic 4/, male												
2002	1.2	*	0.4	0.3	0.6	0.7	0.9	1.2	1.6	2.4	4.9	11.1
2001	1.1	*	*	0.3	0.4	0.6	0.9	1.2	1.5	2.6	4.4	12.2
2000	1.2	*	*	0.4	0.6	0.8	0.9	1.3	1.6	2.9	4.9	12.7
1999	1.3	*	*	0.4	0.6	0.7	0.8	1.1	1.8	3.1	5.6	12.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Non-Hispanic 4/, female												
2002	2.0	*	*	0.3	0.3	0.6	1.2	1.9	2.4	3.8	7.2	16.3
2001	2.1	*	*	0.2	0.4	0.7	1.2	1.9	2.8	4.1	7.5	17.7
2000	2.1	*	*	0.3	0.4	0.6	1.2	2.0	2.6	4.7	8.0	18.2
1999	2.2	*	*	0.2	0.4	0.7	1.2	2.2	3.0	4.9	8.6	19.1
Non-Hispanic white, both sexes												
2002	1.4	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.6	6.1	14.7
2001	1.4	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.9	5.8	15.7
2000	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.5	3.2	6.4	15.8
1999	1.5	*	*	0.2	0.4	0.4	0.7	1.2	1.9	3.4	7.1	16.5
Non-Hispanic white, male												
2002	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.1	1.8	4.5	10.8
2001	0.9	*	*	0.2	0.2	0.3	0.6	0.8	1.0	2.1	3.9	11.4
2000	1.0	*	*	0.2	0.3	0.5	0.6	0.9	1.0	2.3	4.6	11.1
1999	1.0	*	*	0.2	0.4	0.4	0.5	0.7	1.4	2.4	5.2	10.6
Non-Hispanic white, female												
2002	1.8	*	*	*	0.3	0.5	0.9	1.3	1.8	3.2	7.1	16.4
2001	1.9	*	*	*	0.2	0.4	0.8	1.4	2.1	3.6	7.1	17.5
2000	1.9	*	*	0.2	0.3	0.4	0.9	1.3	2.0	4.0	7.6	17.7
1999	2.1	*	*	*	0.3	0.5	0.9	1.6	2.3	4.2	8.3	18.8
Non-Hispanic black, both sexes												
2002	3.0	*	0.9	1.1	1.3	1.6	3.5	4.9	6.6	8.0	7.9	11.8
2001	3.1	*	*	0.8	1.2	2.2	3.3	4.7	6.9	7.7	10.7	16.5
2000	3.3	*	*	1.1	1.3	2.0	3.2	5.5	6.9	9.4	10.4	22.3
1999	3.3	*	*	1.0	1.3	2.4	3.0	5.4	7.0	9.8	10.6	20.7
Non-Hispanic black, male												
2002	2.8	*	*	1.1	1.9	2.0	3.7	4.0	5.9	7.5	6.7	*
2001	2.7	*	*	0.9	1.4	2.5	3.0	4.1	4.7	6.8	8.7	*
2000	2.8	*	*	1.3	1.6	2.5	3.0	3.7	6.1	7.9	6.5	24.7
1999	2.9	*	*	1.4	1.7	2.7	2.8	4.0	5.5	9.2	8.4	*
Non-Hispanic black, female												
2002	3.1	*	*	1.0	*	1.3	3.4	5.7	7.2	8.4	8.6	12.5
2001	3.4	*	*	0.7	0.9	2.0	3.6	5.1	8.6	8.4	11.8	17.1
2000	3.7	*	*	0.8	1.0	1.5	3.4	7.0	7.6	10.4	12.7	21.4
1999	3.6	*	*	0.6	0.9	2.2	3.2	6.5	8.1	10.2	11.7	21.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchiectasis (J47)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	*	0.1	0.3	0.9	3.4	5.8
2001	0.3	*	*	*	*	*	*	0.1	0.3	1.0	3.0	7.0
2000	0.3	*	*	*	*	*	*	*	0.3	0.9	3.3	6.1
1999	0.3	*	*	*	*	*	*	0.1	0.3	0.9	3.4	6.2
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	2.2	4.7
2001	0.2	*	*	*	*	*	*	*	0.3	0.7	1.8	4.8
2000	0.2	*	*	*	*	*	*	*	0.2	0.7	2.2	4.3
1999	0.2	*	*	*	*	*	*	*	0.2	0.7	3.1	6.6
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	*	0.3	1.1	4.1	6.3
2001	0.5	*	*	*	*	*	*	*	0.3	1.2	3.7	7.9
2000	0.5	*	*	*	*	*	*	*	0.3	1.1	4.0	6.9
1999	0.5	*	*	*	*	*	*	0.1	0.4	1.2	3.6	6.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.9	3.4	5.9
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.9	3.0	7.2
2000	0.4	*	*	*	*	*	*	*	0.3	1.0	3.3	6.2
1999	0.4	*	*	*	*	*	*	0.1	0.3	0.9	3.4	6.2
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	2.2	4.7
2001	0.2	*	*	*	*	*	*	*	0.3	0.7	1.9	4.9
2000	0.2	*	*	*	*	*	*	*	0.2	0.8	2.1	4.4
1999	0.3	*	*	*	*	*	*	*	0.2	0.7	3.1	6.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchiectasis (J47)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	*	0.3	1.2	4.2	6.4
2001	0.5	*	*	*	*	*	*	*	0.3	1.2	3.7	8.1
2000	0.5	*	*	*	*	*	*	*	0.3	1.1	4.0	6.9
1999	0.5	*	*	*	*	*	*	*	0.4	1.2	3.7	6.0
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.3	0.9	3.7	6.2
2001	0.4	*	*	*	*	*	*	*	0.3	1.0	3.1	7.6
2000	0.4	*	*	*	*	*	*	*	0.3	0.9	3.4	6.6
1999	0.4	*	*	*	*	*	*	0.1	0.3	1.0	3.5	6.7
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	2.4	4.7
2001	0.2	*	*	*	*	*	*	*	0.3	0.6	1.8	4.9
2000	0.2	*	*	*	*	*	*	*	0.2	0.7	2.2	4.7
1999	0.3	*	*	*	*	*	*	*	0.2	0.7	3.0	7.2
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	0.4	1.3	4.6	6.8
2001	0.6	*	*	*	*	*	*	*	0.3	1.3	4.0	8.8
2000	0.6	*	*	*	*	*	*	*	0.3	1.1	4.2	7.4
1999	0.6	*	*	*	*	*	*	*	0.4	1.2	3.8	6.5
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lung diseases due to external agents (J60-J70)												
All origins 3/, both sexes												
2002	6.5	*	*	*	0.1	0.2	0.4	0.9	2.9	11.2	50.3	192.9
2001	6.5	*	*	*	0.1	0.2	0.4	1.0	2.9	11.6	50.1	195.9
2000	6.3	*	*	*	0.1	0.2	0.4	1.0	2.7	12.0	48.6	195.3
1999	5.9	*	*	*	0.1	0.2	0.4	0.8	2.8	11.1	45.5	181.8
All origins 3/, male												
2002	7.0	*	*	*	0.1	0.2	0.5	1.2	3.7	15.8	73.1	290.1
2001	7.0	*	*	*	0.1	0.2	0.6	1.3	3.7	16.6	73.3	291.3
2000	6.9	*	*	*	0.1	0.2	0.5	1.3	3.5	17.7	72.1	301.5
1999	6.5	*	*	*	0.1	0.2	0.5	1.0	3.8	16.4	70.2	279.1
All origins 3/, female												
2002	6.0	*	*	*	*	0.2	0.3	0.6	2.1	7.4	35.2	150.7
2001	6.1	*	*	*	*	0.2	0.3	0.8	2.1	7.4	34.9	156.0
2000	5.8	*	*	*	*	0.1	0.3	0.8	2.0	7.2	33.3	152.1
1999	5.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.5	142.6
Hispanic, both sexes												
2002	1.5	*	*	*	*	*	*	*	2.1	6.5	30.2	136.4
2001	1.6	*	*	*	*	*	*	0.6	1.9	6.7	28.7	165.1
2000	1.5	*	*	*	*	*	*	0.7	2.0	6.4	30.4	140.7
1999	1.3	*	*	*	*	*	*	*	1.7	6.2	25.2	132.6
Hispanic, male												
2002	1.6	*	*	*	*	*	*	*	3.0	8.2	45.5	182.4
2001	1.6	*	*	*	*	*	*	*	2.8	9.8	39.3	230.5
2000	1.5	*	*	*	*	*	*	*	2.6	8.4	49.2	177.4
1999	1.2	*	*	*	*	*	*	*	*	8.1	35.5	161.8
Hispanic, female												
2002	1.5	*	*	*	*	*	*	*	*	5.2	19.7	113.6
2001	1.5	*	*	*	*	*	*	*	*	4.3	21.5	133.1
2000	1.4	*	*	*	*	*	*	*	*	4.8	17.8	122.7
1999	1.4	*	*	*	*	*	*	*	*	4.6	18.4	118.5
Non-Hispanic 4/, both sexes												
2002	7.3	*	*	*	0.1	0.2	0.5	1.0	2.9	11.5	51.2	194.7
2001	7.2	*	*	*	0.1	0.2	0.4	1.0	3.0	11.9	51.0	196.5
2000	7.0	*	*	*	0.1	0.2	0.4	1.0	2.8	12.3	49.2	196.9
1999	6.5	*	*	*	0.1	0.2	0.4	0.9	2.8	11.4	46.2	183.1
Non-Hispanic 4/, male												
2002	7.9	*	*	*	0.1	0.2	0.6	1.3	3.7	16.2	74.2	294.1
2001	7.8	*	*	*	0.1	0.2	0.6	1.3	3.8	17.0	74.7	293.0
2000	7.7	*	*	*	0.1	0.2	0.6	1.3	3.6	18.3	72.8	305.9
1999	7.2	*	*	*	0.1	0.3	0.5	1.1	3.9	16.8	71.5	283.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Lung diseases due to external agents (J60-J70)												
Non-Hispanic 4/, female												
2002	6.7	*	*	*	*	0.2	0.4	0.7	2.2	7.5	35.9	151.9
2001	6.7	*	*	*	*	0.2	0.3	0.8	2.2	7.6	35.4	156.4
2000	6.4	*	*	*	*	0.1	0.3	0.8	2.0	7.4	33.9	152.8
1999	5.8	*	*	*	*	0.2	0.3	0.7	1.9	6.9	29.9	143.2
Non-Hispanic white, both sexes												
2002	8.2	*	*	*	0.1	0.2	0.5	0.9	2.8	11.3	51.6	198.7
2001	8.1	*	*	*	0.1	0.2	0.4	1.0	2.9	11.7	51.3	198.1
2000	7.8	*	*	*	0.1	0.2	0.4	1.0	2.6	11.9	49.3	200.8
1999	7.2	*	*	*	0.1	0.2	0.4	0.8	2.7	11.0	46.1	185.9
Non-Hispanic white, male												
2002	8.9	*	*	*	*	0.3	0.6	1.2	3.6	15.9	74.9	301.6
2001	8.7	*	*	*	*	0.3	0.6	1.3	3.7	16.9	75.6	299.5
2000	8.6	*	*	*	*	0.2	0.5	1.2	3.5	17.4	73.1	315.9
1999	8.0	*	*	*	*	0.2	0.5	1.0	3.6	16.1	71.0	289.6
Non-Hispanic white, female												
2002	7.5	*	*	*	*	0.2	0.4	0.6	2.1	7.3	36.1	154.2
2001	7.4	*	*	*	*	0.2	0.3	0.7	2.2	7.2	35.2	155.9
2000	7.1	*	*	*	*	0.2	0.3	0.8	1.9	7.2	33.7	154.4
1999	6.5	*	*	*	*	0.2	0.3	0.7	1.8	6.7	29.8	144.5
Non-Hispanic black, both sexes												
2002	4.4	*	*	*	*	*	0.7	1.5	4.1	15.9	55.2	171.5
2001	4.6	*	*	*	*	*	0.6	1.8	4.4	16.2	55.8	191.0
2000	4.5	*	*	*	*	*	0.7	1.9	4.4	19.0	54.8	164.4
1999	4.5	*	*	*	*	*	0.8	1.6	5.0	17.4	54.9	173.6
Non-Hispanic black, male												
2002	4.7	*	*	*	*	*	0.9	2.1	5.3	23.7	80.0	251.6
2001	4.6	*	*	*	*	*	0.8	2.0	6.2	22.5	81.6	257.6
2000	4.9	*	*	*	*	*	0.9	2.7	5.4	31.0	82.1	224.3
1999	5.1	*	*	*	*	*	0.9	2.0	7.5	27.5	86.9	247.2
Non-Hispanic black, female												
2002	4.1	*	*	*	*	*	*	0.9	3.1	10.3	41.1	140.9
2001	4.6	*	*	*	*	*	*	1.5	2.9	11.7	41.3	166.4
2000	4.1	*	*	*	*	*	*	1.2	3.6	10.5	39.4	142.2
1999	4.0	*	*	*	*	*	*	1.2	2.9	10.3	36.9	146.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.3	1.2	3.8	6.8
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.1	6.2
2000	0.4	*	*	*	*	*	*	*	0.2	1.5	3.9	7.5
1999	0.4	*	*	*	*	*	*	*	0.2	1.5	4.1	6.1
All origins 3/, male												
2002	0.7	*	*	*	*	*	*	*	0.5	2.6	9.0	21.6
2001	0.8	*	*	*	*	*	*	*	0.5	3.1	9.9	20.2
2000	0.8	*	*	*	*	*	*	*	0.5	3.2	9.6	24.9
1999	0.8	*	*	*	*	*	*	*	0.5	3.2	10.1	20.4
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.3	1.2	3.9	6.9
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.2	6.2
2000	0.5	*	*	*	*	*	*	*	0.2	1.6	4.0	7.6
1999	0.4	*	*	*	*	*	*	*	0.3	1.5	4.3	6.2
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	*	*	*	0.5	2.6	9.3	21.9
2001	0.9	*	*	*	*	*	*	*	0.5	3.3	10.2	20.5
2000	0.9	*	*	*	*	*	*	*	0.5	3.3	9.7	25.7
1999	0.9	*	*	*	*	*	*	*	0.5	3.4	10.5	21.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.5	*	*	*	*	*	*	*	0.3	1.4	4.1	7.3
2001	0.5	*	*	*	*	*	*	*	0.3	1.6	4.5	6.6
2000	0.5	*	*	*	*	*	*	*	0.3	1.7	4.1	8.1
1999	0.5	*	*	*	*	*	*	*	0.2	1.6	4.6	6.4
Non-Hispanic white, male												
2002	1.0	*	*	*	*	*	*	*	0.6	2.8	9.8	23.2
2001	1.0	*	*	*	*	*	*	*	0.5	3.5	10.8	21.6
2000	1.1	*	*	*	*	*	*	*	0.5	3.6	10.1	27.2
1999	1.0	*	*	*	*	*	*	*	0.5	3.6	11.1	21.8
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	2.3	*
2001	0.2	*	*	*	*	*	*	*	*	*	2.2	*
2000	0.2	*	*	*	*	*	*	*	*	*	2.6	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	5.9	*
2001	0.3	*	*	*	*	*	*	*	*	*	6.0	*
2000	0.3	*	*	*	*	*	*	*	*	*	6.5	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
All origins 3/, both sexes												
2002	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
All origins 3/, male												
2002	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4
All origins 3/, female												
2002	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2
Hispanic, both sexes												
2002	1.4	*	*	*	*	*	*	*	1.8	5.3	29.0	131.4
2001	1.5	*	*	*	*	*	*	*	1.8	6.1	27.4	161.3
2000	1.4	*	*	*	*	*	*	0.6	1.9	5.9	27.9	137.4
1999	1.2	*	*	*	*	*	*	*	1.6	5.5	24.6	130.5
Hispanic, male												
2002	1.4	*	*	*	*	*	*	*	2.6	6.1	43.1	167.3
2001	1.5	*	*	*	*	*	*	*	2.5	8.4	36.2	220.9
2000	1.4	*	*	*	*	*	*	*	2.5	7.4	42.8	169.3
1999	1.1	*	*	*	*	*	*	*	*	6.8	33.9	159.6
Hispanic, female												
2002	1.4	*	*	*	*	*	*	*	*	4.6	19.4	113.6
2001	1.5	*	*	*	*	*	*	*	*	4.3	21.5	132.2
2000	1.4	*	*	*	*	*	*	*	*	4.8	17.8	121.7
1999	1.3	*	*	*	*	*	*	*	*	4.5	18.4	116.5
Non-Hispanic 4/, both sexes												
2002	6.8	*	*	*	0.1	0.2	0.5	0.9	2.6	10.1	47.1	187.8
2001	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.2	46.5	190.0
2000	6.5	*	*	*	0.1	0.2	0.4	1.0	2.5	10.6	45.1	189.0
1999	6.0	*	*	*	0.1	0.2	0.4	0.8	2.5	9.7	41.7	176.8
Non-Hispanic 4/, male												
2002	7.0	*	*	*	0.1	0.2	0.6	1.2	3.1	13.4	64.7	272.1
2001	6.8	*	*	*	0.1	0.2	0.6	1.3	3.2	13.6	64.2	271.6
2000	6.8	*	*	*	0.1	0.2	0.5	1.2	3.1	14.8	62.8	279.7
1999	6.3	*	*	*	0.1	0.3	0.5	1.0	3.3	13.3	60.6	261.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Non-Hispanic 4/, female												
2002	6.6	*	*	*	*	0.2	0.3	0.6	2.1	7.3	35.4	151.4
2001	6.6	*	*	*	*	0.2	0.3	0.8	2.2	7.4	34.9	156.0
2000	6.3	*	*	*	*	0.1	0.3	0.8	2.0	7.1	33.6	152.3
1999	5.8	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.4	142.9
Non-Hispanic white, both sexes												
2002	7.7	*	*	*	0.1	0.2	0.4	0.9	2.5	9.7	47.3	191.3
2001	7.5	*	*	*	0.1	0.2	0.4	0.9	2.6	9.9	46.6	191.2
2000	7.3	*	*	*	0.1	0.2	0.4	0.9	2.3	10.0	45.0	192.5
1999	6.7	*	*	*	0.1	0.2	0.4	0.7	2.4	9.2	41.2	179.3
Non-Hispanic white, male												
2002	7.9	*	*	*	*	0.3	0.6	1.1	3.0	12.9	64.9	278.4
2001	7.6	*	*	*	*	0.3	0.6	1.2	3.1	13.2	64.4	276.9
2000	7.5	*	*	*	*	0.2	0.5	1.1	2.9	13.7	62.7	288.1
1999	6.9	*	*	*	*	0.2	0.4	0.9	3.0	12.4	59.6	267.4
Non-Hispanic white, female												
2002	7.4	*	*	*	*	0.2	0.3	0.6	2.0	7.1	35.6	153.7
2001	7.4	*	*	*	*	0.2	0.3	0.7	2.1	7.0	34.7	155.5
2000	7.1	*	*	*	*	*	0.3	0.7	1.8	7.0	33.4	153.9
1999	6.5	*	*	*	*	0.2	0.3	0.6	1.7	6.5	29.3	144.2
Non-Hispanic black, both sexes												
2002	4.2	*	*	*	*	*	0.7	1.4	3.8	15.2	52.8	167.3
2001	4.4	*	*	*	*	*	0.6	1.6	4.1	15.4	53.6	187.3
2000	4.3	*	*	*	*	*	0.7	1.9	4.3	17.9	52.0	160.6
1999	4.3	*	*	*	*	*	0.7	1.5	4.5	16.4	52.6	168.5
Non-Hispanic black, male												
2002	4.4	*	*	*	*	*	0.9	2.0	4.9	22.3	74.1	237.4
2001	4.3	*	*	*	*	*	0.8	1.9	5.5	20.8	75.6	243.7
2000	4.5	*	*	*	*	*	0.8	2.7	5.2	28.4	75.0	210.2
1999	4.7	*	*	*	*	*	0.9	1.8	6.6	25.3	81.6	229.4
Non-Hispanic black, female												
2002	4.1	*	*	*	*	*	*	0.9	3.0	10.2	40.8	140.5
2001	4.6	*	*	*	*	*	*	1.4	2.9	11.6	41.3	166.4
2000	4.1	*	*	*	*	*	*	1.2	3.6	10.4	39.1	142.2
1999	4.0	*	*	*	*	*	*	1.2	2.9	10.2	36.4	145.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)												
All origins 3/, both sexes												
2002	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
All origins 3/, male												
2002	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.0	270.2
2000	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.0	59.7	258.4
All origins 3/, female												
2002	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.5
2000	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.0	142.2
Hispanic, both sexes												
2002	1.4	*	*	*	*	*	*	*	1.8	5.3	29.0	131.4
2001	1.5	*	*	*	*	*	*	*	1.8	6.1	27.4	161.3
2000	1.4	*	*	*	*	*	*	0.6	1.9	5.9	27.9	137.4
1999	1.2	*	*	*	*	*	*	*	1.6	5.5	24.6	130.5
Hispanic, male												
2002	1.4	*	*	*	*	*	*	*	2.6	6.1	43.1	167.3
2001	1.5	*	*	*	*	*	*	*	2.5	8.4	36.2	220.9
2000	1.4	*	*	*	*	*	*	*	2.5	7.4	42.8	169.3
1999	1.1	*	*	*	*	*	*	*	*	6.8	33.9	159.6
Hispanic, female												
2002	1.4	*	*	*	*	*	*	*	*	4.6	19.4	113.6
2001	1.5	*	*	*	*	*	*	*	*	4.3	21.5	132.2
2000	1.4	*	*	*	*	*	*	*	*	4.8	17.8	121.7
1999	1.3	*	*	*	*	*	*	*	*	4.5	18.4	116.5
Non-Hispanic 4/, both sexes												
2002	6.8	*	*	*	0.1	0.2	0.5	0.9	2.6	10.1	47.1	187.8
2001	6.7	*	*	*	0.1	0.2	0.4	1.0	2.7	10.2	46.5	189.9
2000	6.5	*	*	*	0.1	0.2	0.4	1.0	2.5	10.6	45.1	189.0
1999	6.0	*	*	*	0.1	0.2	0.4	0.8	2.5	9.7	41.6	176.8
Non-Hispanic 4/, male												
2002	7.0	*	*	*	0.1	0.2	0.6	1.2	3.1	13.4	64.7	272.1
2001	6.8	*	*	*	0.1	0.2	0.6	1.3	3.2	13.6	64.1	271.5
2000	6.8	*	*	*	0.1	0.2	0.5	1.2	3.1	14.8	62.8	279.6
1999	6.3	*	*	*	0.1	0.3	0.5	1.0	3.3	13.3	60.6	261.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)												
Non-Hispanic 4/, female												
2002	6.6	*	*	*	*	0.2	0.3	0.6	2.1	7.3	35.4	151.4
2001	6.6	*	*	*	*	0.2	0.3	0.8	2.2	7.4	34.9	156.0
2000	6.3	*	*	*	*	0.1	0.3	0.8	1.9	7.1	33.5	152.3
1999	5.8	*	*	*	*	0.2	0.3	0.7	1.8	6.8	29.3	142.8
Non-Hispanic white, both sexes												
2002	7.6	*	*	*	0.1	0.2	0.4	0.9	2.5	9.7	47.3	191.3
2001	7.5	*	*	*	0.1	0.2	0.4	0.9	2.6	9.9	46.6	191.1
2000	7.3	*	*	*	0.1	0.2	0.4	0.9	2.3	10.0	45.0	192.4
1999	6.7	*	*	*	0.1	0.2	0.4	0.7	2.4	9.2	41.2	179.3
Non-Hispanic white, male												
2002	7.9	*	*	*	*	0.3	0.6	1.1	3.0	12.9	64.9	278.4
2001	7.6	*	*	*	*	0.3	0.6	1.2	3.1	13.2	64.4	276.9
2000	7.5	*	*	*	*	0.2	0.5	1.1	2.9	13.7	62.7	288.0
1999	6.9	*	*	*	*	0.2	0.4	0.9	3.0	12.4	59.6	267.4
Non-Hispanic white, female												
2002	7.4	*	*	*	*	0.2	0.3	0.6	2.0	7.1	35.5	153.7
2001	7.4	*	*	*	*	0.2	0.3	0.7	2.1	7.0	34.7	155.4
2000	7.1	*	*	*	*	*	0.3	0.7	1.8	7.0	33.4	153.9
1999	6.4	*	*	*	*	0.2	0.3	0.6	1.7	6.5	29.2	144.1
Non-Hispanic black, both sexes												
2002	4.2	*	*	*	*	*	0.7	1.4	3.8	15.2	52.7	167.3
2001	4.4	*	*	*	*	*	0.6	1.6	4.1	15.4	53.6	187.3
2000	4.3	*	*	*	*	*	0.7	1.9	4.3	17.9	52.0	160.6
1999	4.3	*	*	*	*	*	0.7	1.5	4.5	16.4	52.5	168.5
Non-Hispanic black, male												
2002	4.4	*	*	*	*	*	0.9	2.0	4.9	22.3	74.1	237.4
2001	4.3	*	*	*	*	*	0.8	1.9	5.5	20.8	75.6	243.7
2000	4.5	*	*	*	*	*	0.8	2.7	5.2	28.4	75.0	210.2
1999	4.7	*	*	*	*	*	0.9	1.8	6.6	25.1	81.6	229.4
Non-Hispanic black, female												
2002	4.1	*	*	*	*	*	*	0.9	3.0	10.2	40.6	140.5
2001	4.6	*	*	*	*	*	*	1.4	2.9	11.6	41.3	166.4
2000	4.1	*	*	*	*	*	*	1.2	3.6	10.4	39.1	142.2
1999	4.0	*	*	*	*	*	*	1.2	2.9	10.2	36.2	145.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other pneumonitis due to solids and liquids (J69.1-J69.8)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other lung diseases due to external agents (J67,J70)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	0.2	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	0.1	0.3	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	0.2	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other lung diseases due to external agents (J67,J70)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	0.2	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.2	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.3
2001	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	1.7	3.0
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.9
1999	0.3	*	*	*	*	0.1	0.1	0.3	0.4	1.0	1.5	2.8
All origins 3/, male												
2002	0.3	*	*	*	*	*	0.1	0.4	0.6	1.1	2.3	3.4
2001	0.3	*	*	*	*	*	0.2	0.4	0.6	1.0	2.3	4.7
2000	0.3	*	*	*	*	*	0.1	0.3	0.5	1.2	2.2	5.8
1999	0.4	*	*	*	*	*	0.2	0.4	0.6	1.5	2.1	4.0
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.0	1.8
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.3
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.1	1.7
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.1	2.3
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5	2.3
2001	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	1.7	3.1
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.5	2.9
1999	0.3	*	*	*	*	0.1	0.2	0.3	0.5	1.0	1.5	2.8
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	0.2	0.4	0.6	1.0	2.3	3.5
2001	0.4	*	*	*	*	*	0.2	0.4	0.6	1.1	2.3	4.9
2000	0.4	*	*	*	*	*	0.1	0.3	0.6	1.2	2.2	5.8
1999	0.4	*	*	*	*	*	0.2	0.4	0.6	1.5	2.2	3.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.0	1.8
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.2	2.4
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.1	1.7
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.1	2.3
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.4	2.2
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.6	3.1
2000	0.3	*	*	*	*	*	0.1	0.2	0.3	0.8	1.5	2.7
1999	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	1.5	2.5
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	0.3	0.5	1.0	2.1	3.2
2001	0.4	*	*	*	*	*	0.1	0.3	0.6	0.9	2.2	4.9
2000	0.4	*	*	*	*	*	0.1	0.3	0.5	1.1	2.1	5.5
1999	0.4	*	*	*	*	*	0.2	0.4	0.4	1.4	2.2	3.2
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	0.2	0.5	1.0	1.7
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.3	2.3
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.2	1.5
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.1	2.3
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	0.6	*	*	2.9	*
2001	0.3	*	*	*	*	*	*	0.5	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	1.5	*	*
1999	0.4	*	*	*	*	*	*	*	1.3	1.5	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	2.2	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pleural effusion and plaque (J90,J92)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.7	2.5	9.9
2001	0.3	*	*	*	*	*	*	0.1	0.2	0.7	2.5	10.0
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.4	8.9
1999	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.4	9.0
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	*	0.2	0.6	2.9	11.0
2001	0.3	*	*	*	*	*	*	*	0.2	0.8	2.8	12.2
2000	0.3	*	*	*	*	*	*	*	0.2	0.8	2.8	10.8
1999	0.3	*	*	*	*	*	*	*	0.2	0.8	3.0	11.5
All origins 3/, female												
2002	0.4	*	*	*	*	*	*	*	*	0.7	2.2	9.4
2001	0.4	*	*	*	*	*	*	*	0.2	0.5	2.4	9.1
2000	0.3	*	*	*	*	*	*	*	*	0.5	2.1	8.1
1999	0.3	*	*	*	*	*	*	*	0.2	0.5	2.0	8.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	*	0.1	0.2	0.7	2.5	10.1
2001	0.4	*	*	*	*	*	*	0.1	0.2	0.7	2.6	10.2
2000	0.3	*	*	*	*	*	*	0.1	0.1	0.6	2.4	8.9
1999	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	9.0
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	*	0.2	0.7	3.0	11.2
2001	0.3	*	*	*	*	*	*	*	*	0.9	2.9	12.4
2000	0.3	*	*	*	*	*	*	*	*	0.7	2.8	11.1
1999	0.3	*	*	*	*	*	*	*	0.2	0.8	3.0	11.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pleural effusion and plaque (J90,J92)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	*	*	*	0.7	2.2	9.6
2001	0.4	*	*	*	*	*	*	*	0.2	0.5	2.4	9.3
2000	0.4	*	*	*	*	*	*	*	*	0.5	2.1	8.0
1999	0.4	*	*	*	*	*	*	*	0.2	0.5	2.0	8.0
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.2	0.7	2.5	10.4
2001	0.4	*	*	*	*	*	*	*	0.2	0.7	2.6	10.1
2000	0.4	*	*	*	*	*	*	*	0.1	0.6	2.4	9.0
1999	0.4	*	*	*	*	*	*	*	0.2	0.7	2.4	9.0
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	*	*	0.2	0.6	3.0	11.8
2001	0.4	*	*	*	*	*	*	*	*	0.9	2.8	12.7
2000	0.3	*	*	*	*	*	*	*	*	0.7	2.8	11.2
1999	0.3	*	*	*	*	*	*	*	*	0.9	3.0	11.7
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	*	*	*	0.7	2.2	9.8
2001	0.5	*	*	*	*	*	*	*	*	0.5	2.4	9.1
2000	0.4	*	*	*	*	*	*	*	*	0.5	2.1	8.2
1999	0.4	*	*	*	*	*	*	*	0.2	0.5	2.1	8.0
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	2.8	8.5
2001	0.3	*	*	*	*	*	*	*	*	*	3.0	12.2
2000	0.2	*	*	*	*	*	*	*	*	*	2.7	7.9
1999	0.2	*	*	*	*	*	*	*	*	*	2.4	8.7
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	8.8
2001	0.3	*	*	*	*	*	*	*	*	*	*	12.4
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)												
All origins 3/, both sexes												
2002	7.9	6.1	0.4	0.1	0.2	0.4	1.0	2.7	8.3	26.6	63.0	118.1
2001	8.3	6.5	0.4	0.1	0.3	0.4	1.0	3.0	8.8	28.1	67.6	125.7
2000	8.3	6.4	0.4	0.2	0.3	0.5	1.1	3.1	9.2	27.5	66.8	125.8
1999	8.1	5.8	0.4	0.1	0.2	0.5	1.0	2.8	8.6	27.1	65.3	131.6
All origins 3/, male												
2002	7.7	7.3	0.5	0.1	0.2	0.4	1.0	2.7	9.4	32.9	77.5	138.9
2001	8.1	8.1	0.4	0.2	0.3	0.4	1.1	3.2	9.9	34.1	82.5	147.4
2000	8.1	7.1	0.4	0.2	0.3	0.5	1.2	3.2	10.4	33.0	83.8	158.3
1999	7.9	6.6	0.4	0.1	0.3	0.5	1.1	2.9	9.5	33.1	82.9	161.8
All origins 3/, female												
2002	8.0	4.9	0.3	0.2	0.3	0.4	1.0	2.8	7.3	21.4	53.4	109.0
2001	8.6	4.9	0.4	0.1	0.3	0.5	1.0	2.7	7.7	23.2	57.9	116.6
2000	8.4	5.7	0.3	0.1	0.3	0.5	1.1	2.9	8.0	23.0	55.7	112.6
1999	8.3	4.9	0.3	0.2	0.2	0.6	1.0	2.8	7.7	22.2	53.8	119.4
Hispanic, both sexes												
2002	3.0	4.2	*	*	*	0.3	0.6	2.3	7.6	24.5	54.2	104.2
2001	3.2	4.4	*	*	*	*	0.7	2.6	8.2	24.0	61.0	125.7
2000	3.2	4.9	*	*	*	0.4	1.0	2.4	8.9	24.0	60.4	124.1
1999	3.1	3.7	*	*	*	*	0.8	2.7	8.7	23.9	61.2	122.1
Hispanic, male												
2002	2.9	*	*	*	*	*	0.7	2.2	9.2	32.3	58.7	98.7
2001	3.1	*	*	*	*	*	0.8	2.8	9.7	28.6	70.6	154.3
2000	3.1	*	*	*	*	*	1.1	2.6	11.0	27.4	74.7	120.9
1999	2.9	*	*	*	*	*	0.9	2.4	10.3	27.7	77.2	140.2
Hispanic, female												
2002	3.1	*	*	*	*	*	*	2.4	6.2	18.2	51.2	106.9
2001	3.3	*	*	*	*	*	*	2.5	6.9	20.5	54.4	111.7
2000	3.4	*	*	*	*	*	0.8	2.1	7.2	21.2	50.8	125.6
1999	3.2	*	*	*	*	*	*	3.0	7.4	21.0	50.7	113.3
Non-Hispanic 4/, both sexes												
2002	8.6	6.6	0.4	0.1	0.3	0.4	1.1	2.8	8.4	26.7	63.3	118.4
2001	9.1	7.0	0.4	0.2	0.3	0.5	1.1	3.0	8.8	28.3	67.7	125.4
2000	8.9	6.7	0.4	0.2	0.3	0.5	1.2	3.1	9.2	27.6	66.8	125.6
1999	8.8	6.2	0.4	0.2	0.2	0.6	1.1	2.8	8.5	27.3	65.3	131.6
Non-Hispanic 4/, male												
2002	8.5	8.0	0.5	0.1	0.3	0.4	1.1	2.7	9.4	32.9	78.3	140.4
2001	8.8	9.0	0.4	0.2	0.3	0.4	1.1	3.3	9.9	34.3	82.8	147.0
2000	8.9	7.6	0.4	0.2	0.3	0.5	1.2	3.3	10.4	33.2	84.0	159.5
1999	8.6	7.3	0.5	0.2	0.3	0.5	1.1	2.9	9.4	33.3	82.8	162.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)												
Non-Hispanic 4/, female												
2002	8.7	5.1	*	0.2	0.3	0.5	1.0	2.8	7.4	21.5	53.4	108.9
2001	9.3	4.9	0.5	0.1	0.3	0.6	1.1	2.8	7.8	23.3	57.9	116.4
2000	9.0	5.8	0.3	0.1	0.3	0.6	1.1	2.9	8.0	23.0	55.7	111.9
1999	8.9	5.2	0.4	0.2	0.2	0.6	1.1	2.7	7.7	22.3	53.9	119.3
Non-Hispanic white, both sexes												
2002	9.6	4.2	0.3	0.1	0.2	0.4	0.9	2.4	8.0	27.3	65.4	122.5
2001	10.0	4.4	0.3	0.2	0.2	0.4	0.9	2.6	8.5	28.9	69.9	128.7
2000	9.9	4.8	0.3	0.2	0.2	0.5	1.0	2.8	8.9	28.2	68.5	129.7
1999	9.7	4.2	0.4	0.1	0.3	0.5	0.9	2.5	8.3	27.5	66.7	134.8
Non-Hispanic white, male												
2002	9.4	5.2	*	*	0.2	0.3	0.9	2.5	9.0	33.7	80.8	145.5
2001	9.8	5.7	*	0.2	0.2	0.3	1.0	2.9	9.6	35.4	85.8	151.0
2000	9.8	5.9	*	0.2	0.3	0.4	1.0	2.9	10.1	33.8	86.0	164.6
1999	9.5	4.9	*	*	0.3	0.5	0.9	2.6	9.2	34.0	84.8	166.9
Non-Hispanic white, female												
2002	9.7	3.2	*	*	0.2	0.4	0.9	2.4	7.1	21.8	55.1	112.6
2001	10.3	3.0	*	*	0.2	0.5	0.8	2.3	7.5	23.5	59.5	119.4
2000	10.0	3.5	*	*	0.2	0.5	1.0	2.7	7.8	23.5	57.1	115.6
1999	9.8	3.3	*	*	0.2	0.5	0.8	2.4	7.5	22.2	54.9	122.0
Non-Hispanic black, both sexes												
2002	5.5	15.5	0.9	0.3	0.6	0.9	2.0	5.3	12.2	24.4	45.6	77.4
2001	6.0	17.5	*	*	0.5	1.1	2.4	6.4	12.9	26.6	50.5	90.8
2000	5.8	15.9	*	*	0.5	0.9	2.4	6.2	12.4	26.8	53.0	83.6
1999	5.8	15.1	*	*	*	1.1	2.4	5.7	11.3	28.1	53.0	95.1
Non-Hispanic black, male												
2002	5.4	18.5	*	*	*	0.8	2.2	5.2	14.1	29.6	56.9	88.6
2001	5.6	22.3	*	*	*	1.1	1.9	6.5	14.2	29.5	58.2	108.6
2000	5.8	15.7	*	*	*	0.9	2.6	7.0	13.9	31.0	67.0	106.9
1999	5.5	17.7	*	*	*	0.8	2.2	6.1	12.2	31.6	66.9	115.3
Non-Hispanic black, female												
2002	5.6	12.3	*	*	*	0.9	1.8	5.4	10.7	20.7	39.3	73.2
2001	6.4	12.4	*	*	*	1.1	2.9	6.3	11.8	24.6	46.2	84.3
2000	5.9	16.2	*	*	*	0.9	2.1	5.5	11.3	23.9	45.2	75.0
1999	6.1	12.5	*	*	*	1.4	2.6	5.4	10.5	25.7	45.2	87.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)												
All origins 3/, both sexes												
2002	30.1	12.6	0.8	0.3	0.6	2.3	11.7	29.8	44.7	81.7	170.0	415.3
2001	30.1	11.9	0.8	0.3	0.5	2.4	11.7	29.7	45.2	83.8	168.8	431.4
2000	29.9	13.1	0.7	0.3	0.7	2.4	11.9	28.6	45.6	82.1	170.4	440.7
1999	29.7	13.2	0.7	0.3	0.6	2.4	11.8	28.0	45.8	85.2	169.0	436.4
All origins 3/, male												
2002	30.7	15.1	0.9	0.3	0.7	2.9	15.0	42.5	59.0	96.9	184.1	404.4
2001	30.6	13.4	0.9	0.3	0.6	2.9	15.2	41.9	59.8	99.5	181.8	428.2
2000	30.3	14.8	0.9	0.3	0.7	3.0	15.8	40.7	60.1	97.7	186.4	441.9
1999	30.4	16.5	0.7	0.3	0.6	2.9	15.8	40.4	60.7	102.3	185.8	438.5
All origins 3/, female												
2002	29.5	10.0	0.7	0.3	0.6	1.8	8.5	17.7	31.5	69.0	160.6	420.0
2001	29.7	10.3	0.6	0.3	0.5	1.9	8.3	18.0	31.8	70.8	160.2	432.7
2000	29.4	11.4	0.5	0.3	0.5	1.8	8.1	16.9	32.3	69.3	159.9	440.2
1999	29.1	9.7	0.8	0.4	0.6	1.9	7.8	15.9	32.2	71.1	158.1	435.6
Hispanic, both sexes												
2002	16.2	11.8	0.9	0.3	0.7	2.2	12.1	36.4	58.1	102.6	168.1	290.5
2001	16.4	10.8	0.8	*	0.5	2.1	12.3	36.9	58.4	102.0	178.9	364.5
2000	16.4	11.5	0.7	*	0.6	2.4	13.4	37.3	59.5	102.9	183.3	366.3
1999	16.7	12.0	0.7	*	0.7	2.7	14.3	37.1	60.5	106.5	172.3	411.1
Hispanic, male												
2002	19.9	12.9	*	*	0.7	3.1	18.1	57.5	85.3	133.2	194.9	284.4
2001	19.8	12.7	*	*	*	2.8	18.2	58.6	87.9	128.2	204.8	373.3
2000	20.1	12.4	*	*	0.7	3.5	20.3	58.7	90.2	134.8	208.5	387.0
1999	20.3	15.8	*	*	0.7	3.6	21.9	58.9	89.8	138.0	196.0	427.0
Hispanic, female												
2002	12.3	10.6	*	*	*	1.2	5.5	15.3	33.7	78.3	149.8	293.4
2001	12.7	8.7	*	*	*	1.4	5.9	15.5	32.1	81.4	161.4	360.2
2000	12.5	10.6	*	*	*	1.0	6.1	16.3	32.2	77.8	166.4	356.1
1999	12.9	8.1	*	*	*	1.6	6.2	15.9	34.6	81.9	156.5	403.5
Non-Hispanic 4/, both sexes												
2002	32.1	12.6	0.8	0.3	0.6	2.4	11.6	29.1	43.4	79.9	169.6	419.6
2001	32.1	11.9	0.8	0.3	0.5	2.4	11.6	28.9	44.0	82.3	167.9	433.0
2000	31.7	13.3	0.7	0.3	0.7	2.4	11.7	27.7	44.4	80.5	169.4	442.5
1999	31.4	13.3	0.7	0.3	0.6	2.4	11.4	27.0	44.6	83.6	168.4	436.0
Non-Hispanic 4/, male												
2002	32.3	15.6	0.9	0.3	0.7	2.8	14.4	40.7	56.5	94.0	183.0	408.8
2001	32.2	13.4	0.9	0.4	0.6	2.9	14.8	40.1	57.2	97.2	180.3	430.0
2000	31.7	15.1	1.0	0.3	0.8	2.8	15.1	38.8	57.5	94.9	185.0	443.4
1999	31.7	16.4	0.7	0.3	0.6	2.8	15.0	38.5	58.3	99.9	184.9	437.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)												
Non-Hispanic 4/, female												
2002	31.9	9.5	0.7	0.3	0.6	1.9	8.9	17.8	31.1	68.2	160.8	424.2
2001	32.0	10.4	0.6	0.3	0.5	1.9	8.5	18.2	31.7	70.0	159.8	434.3
2000	31.6	11.4	0.4	0.3	0.5	1.9	8.3	16.9	32.2	68.5	159.3	442.1
1999	31.1	10.0	0.8	0.4	0.6	2.0	8.0	15.9	31.9	70.3	157.7	435.4
Non-Hispanic white, both sexes												
2002	34.9	9.2	0.6	0.2	0.5	2.1	11.5	27.9	42.0	80.0	171.8	430.9
2001	34.5	7.7	0.6	0.3	0.4	2.2	11.1	27.4	42.0	82.1	169.3	443.2
2000	33.9	9.2	0.4	0.3	0.6	2.0	11.0	25.9	42.4	79.8	170.5	451.0
1999	33.6	9.1	0.6	0.3	0.5	2.2	10.7	25.2	42.5	82.6	169.3	444.7
Non-Hispanic white, male												
2002	34.5	11.5	0.7	0.3	0.6	2.5	14.4	39.1	54.4	93.5	184.2	417.2
2001	33.9	7.9	0.8	0.3	0.5	2.6	14.0	38.0	54.2	95.9	180.8	436.1
2000	33.4	10.5	0.6	0.3	0.6	2.5	14.3	36.5	54.9	93.6	185.2	449.5
1999	33.3	11.0	0.5	0.3	0.5	2.5	14.1	36.0	55.6	98.8	185.6	441.9
Non-Hispanic white, female												
2002	35.2	6.8	0.6	0.2	0.5	1.6	8.6	16.8	30.2	68.6	163.5	436.9
2001	35.2	7.4	0.5	0.3	0.3	1.8	8.2	17.0	30.5	70.3	161.7	446.1
2000	34.4	7.8	*	0.3	0.5	1.5	7.7	15.3	30.6	68.2	160.9	451.6
1999	33.8	7.2	0.8	0.3	0.4	1.8	7.2	14.6	30.2	69.1	158.7	445.8
Non-Hispanic black, both sexes												
2002	24.4	26.3	1.5	0.5	1.1	3.5	13.2	41.9	59.9	90.6	166.4	336.6
2001	25.9	29.8	1.3	0.5	1.1	3.6	15.5	43.8	66.1	98.1	172.0	354.3
2000	26.1	30.5	1.6	0.4	1.1	4.0	16.4	44.2	65.7	97.9	171.8	385.7
1999	25.8	30.6	1.1	0.5	1.3	3.5	16.5	43.5	66.9	101.3	173.3	366.7
Non-Hispanic black, male												
2002	26.8	32.4	1.7	*	0.9	3.9	15.4	58.7	83.2	114.9	189.9	359.9
2001	29.3	35.1	*	*	1.0	4.5	20.5	63.1	92.1	128.9	196.8	391.5
2000	28.6	34.5	2.2	*	1.4	4.2	21.1	61.6	89.7	123.7	203.3	396.9
1999	28.6	37.8	*	*	1.2	4.2	21.3	64.1	90.3	123.1	197.0	402.9
Non-Hispanic black, female												
2002	22.2	19.9	*	*	1.3	3.1	11.1	27.5	41.2	73.3	153.0	327.8
2001	22.8	24.2	*	*	1.2	2.8	11.0	27.3	45.1	76.4	158.0	340.5
2000	23.7	26.4	*	*	0.9	3.7	12.2	29.3	46.4	79.8	154.0	381.6
1999	23.4	23.2	*	*	1.4	2.9	12.4	26.0	48.2	85.9	160.0	353.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.3	1.5
2001	0.0	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
1999	0.0	*	*	*	*	*	*	*	*	0.1	0.2	1.0
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	1.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.3	1.2
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	1.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	1.5
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	1.1
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of oral cavity, salivary glands and jaws (K00-K14)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	1.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	*	1.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	1.0
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	1.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.2	1.1
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	*	1.2
1999	0.1	*	*	*	*	*	*	*	*	*	*	1.0
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)												
All origins 3/, both sexes												
2002	30.0	12.5	0.8	0.3	0.6	2.3	11.7	29.8	44.6	81.6	169.7	413.8
2001	30.1	11.8	0.8	0.3	0.5	2.4	11.7	29.7	45.2	83.8	168.4	430.2
2000	29.8	13.0	0.7	0.3	0.7	2.4	11.9	28.6	45.6	82.1	170.1	439.5
1999	29.7	13.1	0.7	0.3	0.6	2.4	11.8	27.9	45.8	85.0	168.8	435.4
All origins 3/, male												
2002	30.6	15.0	0.9	0.3	0.7	2.9	15.0	42.4	58.9	96.7	183.8	403.1
2001	30.5	13.3	0.9	0.3	0.6	2.8	15.2	41.8	59.7	99.5	181.4	427.0
2000	30.3	14.8	0.9	0.3	0.7	3.0	15.8	40.6	60.1	97.6	186.2	440.8
1999	30.3	16.4	0.7	0.3	0.6	2.9	15.8	40.4	60.6	102.2	185.6	437.4
All origins 3/, female												
2002	29.4	9.9	0.7	0.3	0.6	1.8	8.5	17.6	31.4	68.9	160.3	418.4
2001	29.7	10.2	0.6	0.3	0.5	1.9	8.3	18.0	31.8	70.8	159.8	431.5
2000	29.4	11.2	0.5	0.3	0.5	1.8	8.1	16.9	32.3	69.3	159.7	439.0
1999	29.0	9.7	0.8	0.4	0.6	1.9	7.8	15.9	32.2	71.0	157.8	434.6
Hispanic, both sexes												
2002	16.2	11.8	0.9	0.3	0.7	2.2	12.0	36.3	57.9	102.6	168.1	289.9
2001	16.3	10.8	0.8	*	0.5	2.1	12.3	36.9	58.3	102.0	178.7	363.3
2000	16.4	11.4	0.7	*	0.6	2.4	13.4	37.3	59.5	102.9	183.1	365.6
1999	16.7	12.0	0.7	*	0.7	2.7	14.3	37.1	60.5	106.3	172.3	411.1
Hispanic, male												
2002	19.8	12.9	*	*	0.7	3.1	18.0	57.5	85.1	133.2	194.9	284.4
2001	19.8	12.7	*	*	*	2.8	18.2	58.6	87.8	128.2	204.8	369.5
2000	20.1	12.4	*	*	0.7	3.5	20.3	58.7	90.2	134.8	208.5	387.0
1999	20.2	15.8	*	*	0.7	3.5	21.9	58.8	89.8	138.0	196.0	427.0
Hispanic, female												
2002	12.3	10.6	*	*	*	1.2	5.5	15.3	33.6	78.2	149.8	292.6
2001	12.7	8.7	*	*	*	1.4	5.9	15.5	32.1	81.4	161.1	360.2
2000	12.5	10.4	*	*	*	1.0	6.1	16.3	32.2	77.8	166.1	355.1
1999	12.9	8.1	*	*	*	1.6	6.2	15.9	34.6	81.6	156.5	403.5
Non-Hispanic 4/, both sexes												
2002	32.1	12.5	0.8	0.3	0.6	2.3	11.6	29.0	43.3	79.8	169.3	418.0
2001	32.0	11.9	0.8	0.3	0.5	2.4	11.6	28.9	44.0	82.3	167.5	431.9
2000	31.6	13.2	0.7	0.3	0.7	2.4	11.6	27.6	44.4	80.4	169.2	441.3
1999	31.3	13.2	0.7	0.3	0.6	2.4	11.4	27.0	44.6	83.5	168.2	435.0
Non-Hispanic 4/, male												
2002	32.2	15.5	0.9	0.3	0.6	2.8	14.4	40.7	56.4	93.9	182.7	407.5
2001	32.1	13.4	0.9	0.3	0.6	2.9	14.8	40.1	57.2	97.2	179.9	428.8
2000	31.7	15.1	1.0	0.3	0.8	2.8	15.1	38.8	57.5	94.8	184.8	442.3
1999	31.7	16.3	0.7	0.3	0.6	2.8	14.9	38.5	58.3	99.8	184.7	436.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)												
Non-Hispanic 4/, female												
2002	31.9	9.5	0.7	0.3	0.6	1.9	8.9	17.8	31.1	68.1	160.5	422.6
2001	32.0	10.3	0.6	0.3	0.5	1.9	8.5	18.2	31.7	69.9	159.3	433.1
2000	31.6	11.3	0.4	0.3	0.5	1.9	8.3	16.9	32.2	68.5	159.0	440.9
1999	31.0	10.0	0.8	0.4	0.6	1.9	7.9	15.8	31.9	70.2	157.5	434.4
Non-Hispanic white, both sexes												
2002	34.8	9.1	0.6	0.2	0.5	2.1	11.5	27.8	42.0	79.9	171.5	429.4
2001	34.5	7.6	0.6	0.3	0.4	2.2	11.1	27.4	42.0	82.0	168.9	441.9
2000	33.9	9.1	0.4	0.3	0.6	2.0	11.0	25.8	42.4	79.7	170.3	449.8
1999	33.5	9.1	0.6	0.3	0.5	2.1	10.6	25.2	42.5	82.6	169.1	443.6
Non-Hispanic white, male												
2002	34.4	11.4	0.6	0.3	0.6	2.5	14.4	39.1	54.4	93.3	183.9	415.9
2001	33.8	7.8	0.7	0.3	0.5	2.6	13.9	38.0	54.2	95.9	180.4	434.9
2000	33.4	10.5	0.6	0.3	0.6	2.5	14.3	36.5	54.9	93.5	185.0	448.2
1999	33.3	10.8	0.5	0.3	0.5	2.5	14.1	36.0	55.6	98.6	185.5	440.6
Non-Hispanic white, female												
2002	35.2	6.7	0.6	0.2	0.5	1.6	8.6	16.7	30.2	68.5	163.2	435.2
2001	35.1	7.4	0.5	0.3	0.3	1.8	8.2	17.0	30.5	70.3	161.3	444.9
2000	34.3	7.7	*	0.3	0.5	1.5	7.7	15.3	30.6	68.1	160.6	450.4
1999	33.7	7.2	0.8	0.3	0.4	1.8	7.2	14.5	30.2	69.1	158.4	444.7
Non-Hispanic black, both sexes												
2002	24.4	26.3	1.5	0.5	1.1	3.4	13.1	41.9	59.9	90.5	166.1	335.7
2001	25.8	29.8	1.3	0.5	1.1	3.6	15.5	43.8	66.1	98.1	171.1	353.3
2000	26.0	30.3	1.6	0.4	1.1	3.9	16.3	44.2	65.7	97.8	171.5	385.4
1999	25.8	30.6	1.1	0.5	1.3	3.5	16.5	43.5	66.9	101.2	172.8	366.0
Non-Hispanic black, male												
2002	26.7	32.4	1.7	*	0.9	3.9	15.4	58.6	83.2	114.8	189.6	359.9
2001	29.2	35.1	*	*	1.0	4.5	20.5	63.1	92.1	128.9	195.6	390.4
2000	28.6	34.5	2.2	*	1.4	4.2	21.1	61.6	89.7	123.5	203.3	396.9
1999	28.5	37.8	*	*	1.2	4.2	21.2	64.1	90.2	123.1	196.7	402.9
Non-Hispanic black, female												
2002	22.2	19.9	*	*	1.3	3.1	11.1	27.5	41.2	73.3	152.9	326.5
2001	22.7	24.2	*	*	1.2	2.8	11.0	27.3	45.1	76.4	157.3	339.6
2000	23.7	26.1	*	*	0.9	3.7	12.2	29.3	46.4	79.7	153.5	381.1
1999	23.3	23.2	*	*	1.4	2.9	12.3	26.0	48.2	85.8	159.5	352.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of esophagus (K20-K22)												
All origins 3/, both sexes												
2002	0.7	*	*	*	*	0.1	0.1	0.3	0.4	1.3	4.9	20.7
2001	0.7	*	*	*	*	*	0.1	0.2	0.5	1.4	5.1	21.2
2000	0.8	0.5	*	*	*	*	0.1	0.2	0.5	1.4	5.2	23.3
1999	0.8	*	*	*	*	0.1	0.1	0.2	0.5	1.5	5.0	23.5
All origins 3/, male												
2002	0.7	*	*	*	*	*	0.2	0.4	0.4	1.6	5.8	25.4
2001	0.7	*	*	*	*	*	0.1	0.3	0.7	1.7	5.9	23.6
2000	0.7	*	*	*	*	*	0.1	0.3	0.7	1.8	6.4	28.6
1999	0.7	*	*	*	*	*	0.1	0.3	0.6	2.0	6.5	28.3
All origins 3/, female												
2002	0.8	*	*	*	*	*	*	0.2	0.3	1.1	4.3	18.7
2001	0.8	*	*	*	*	*	*	*	0.3	1.1	4.6	20.2
2000	0.8	*	*	*	*	*	*	0.1	0.3	1.1	4.5	21.1
1999	0.8	*	*	*	*	*	*	0.2	0.4	1.1	4.1	21.5
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	3.7	11.1
2001	0.2	*	*	*	*	*	*	*	*	*	*	13.1
2000	0.2	*	*	*	*	*	*	*	*	*	4.7	*
1999	0.3	*	*	*	*	*	*	*	*	2.1	*	24.6
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	23.9
Non-Hispanic 4/, both sexes												
2002	0.8	*	*	*	*	0.1	0.1	0.3	0.4	1.3	4.9	21.1
2001	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.2	21.5
2000	0.9	*	*	*	*	*	0.1	0.2	0.5	1.5	5.2	23.7
1999	0.8	*	*	*	*	0.1	0.1	0.2	0.5	1.5	5.1	23.4
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	*	0.1	0.4	0.4	1.6	5.8	25.7
2001	0.7	*	*	*	*	*	0.1	0.3	0.7	1.7	6.0	24.1
2000	0.8	*	*	*	*	*	0.2	0.4	0.7	1.8	6.4	28.7
1999	0.8	*	*	*	*	*	0.1	0.3	0.6	2.0	6.6	28.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of esophagus (K20-K22)												
Non-Hispanic 4/, female												
2002	0.9	*	*	*	*	*	*	0.2	0.3	1.1	4.4	19.1
2001	0.9	*	*	*	*	*	*	*	0.3	1.2	4.7	20.5
2000	0.9	*	*	*	*	*	*	0.1	0.3	1.2	4.5	21.6
1999	0.9	*	*	*	*	*	*	0.2	0.4	1.1	4.1	21.4
Non-Hispanic white, both sexes												
2002	0.9	*	*	*	*	*	0.1	0.3	0.4	1.3	5.0	22.1
2001	0.9	*	*	*	*	*	0.1	0.2	0.5	1.4	5.3	22.6
2000	1.0	*	*	*	*	*	0.1	0.2	0.5	1.5	5.4	24.8
1999	1.0	*	*	*	*	*	0.1	0.2	0.5	1.5	5.3	24.4
Non-Hispanic white, male												
2002	0.8	*	*	*	*	*	0.2	0.4	0.4	1.6	5.8	27.1
2001	0.8	*	*	*	*	*	0.1	0.2	0.7	1.7	6.0	25.3
2000	0.9	*	*	*	*	*	0.1	0.3	0.7	1.8	6.6	30.1
1999	0.9	*	*	*	*	*	0.1	0.2	0.6	2.0	6.7	29.9
Non-Hispanic white, female												
2002	1.0	*	*	*	*	*	*	0.2	0.4	1.1	4.5	20.0
2001	1.1	*	*	*	*	*	*	*	0.4	1.2	4.9	21.4
2000	1.1	*	*	*	*	*	*	*	0.3	1.2	4.7	22.7
1999	1.0	*	*	*	*	*	*	0.2	0.4	1.1	4.4	22.2
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	*	2.0	4.6	12.1
2001	0.4	*	*	*	*	*	*	*	*	1.5	4.2	13.1
2000	0.5	*	*	*	*	*	*	*	*	1.8	4.1	14.3
1999	0.5	*	*	*	*	*	*	*	*	1.7	3.6	15.5
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	2.9	5.9	*
2001	0.5	*	*	*	*	*	*	*	*	*	6.6	*
2000	0.6	*	*	*	*	*	*	*	*	*	6.5	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	3.8	11.3
2001	0.4	*	*	*	*	*	*	*	*	*	*	12.4
2000	0.4	*	*	*	*	*	*	*	*	*	*	13.5
1999	0.4	*	*	*	*	*	*	*	*	*	*	15.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All origins 3/, both sexes												
2002	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.5	33.4
2000	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
All origins 3/, male												
2002	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
All origins 3/, female												
2002	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	0.5	1.6	3.3	8.8	20.5
2001	0.5	*	*	*	*	*	*	*	1.2	2.9	10.1	23.8
2000	0.5	*	*	*	*	*	*	*	1.6	3.3	9.9	30.5
1999	0.5	*	*	*	*	*	*	*	1.5	3.6	9.2	26.7
Hispanic, male												
2002	0.6	*	*	*	*	*	*	*	2.5	5.4	11.1	*
2001	0.5	*	*	*	*	*	*	*	*	*	17.0	*
2000	0.5	*	*	*	*	*	*	*	2.9	4.4	*	*
1999	0.6	*	*	*	*	*	*	*	*	4.8	14.1	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	7.3	17.4
2001	0.4	*	*	*	*	*	*	*	*	*	*	25.1
2000	0.5	*	*	*	*	*	*	*	*	*	10.9	29.7
1999	0.4	*	*	*	*	*	*	*	*	*	*	22.9
Non-Hispanic 4/, both sexes												
2002	1.6	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.2	28.6
2001	1.7	*	*	*	*	0.1	0.3	0.7	1.7	4.0	11.5	33.6
2000	1.8	*	*	*	*	*	0.3	0.8	1.8	4.1	11.5	35.7
1999	1.8	*	*	*	*	*	0.3	0.7	1.8	4.6	12.4	34.8
Non-Hispanic 4/, male												
2002	1.5	*	*	*	*	*	0.4	0.9	1.9	4.5	12.3	31.7
2001	1.6	*	*	*	*	0.1	0.4	1.0	2.2	4.8	12.8	35.5
2000	1.8	*	*	*	*	*	0.4	1.2	2.4	5.4	13.9	41.0
1999	1.7	*	*	*	*	*	0.4	1.0	2.3	6.1	14.8	37.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Non-Hispanic 4/, female												
2002	1.6	*	*	*	*	*	0.2	0.5	1.2	3.0	8.8	27.3
2001	1.8	*	*	*	*	*	0.2	0.5	1.2	3.3	10.6	32.8
2000	1.8	*	*	*	*	*	0.1	0.5	1.3	3.1	9.9	33.5
1999	1.9	*	*	*	*	*	0.2	0.5	1.3	3.3	10.8	33.7
Non-Hispanic white, both sexes												
2002	1.7	*	*	*	*	*	0.2	0.6	1.4	3.6	10.4	29.4
2001	1.9	*	*	*	*	*	0.2	0.7	1.5	3.9	11.6	34.7
2000	1.9	*	*	*	*	*	0.2	0.7	1.8	4.0	11.6	36.3
1999	2.0	*	*	*	*	*	0.2	0.6	1.7	4.5	12.4	35.6
Non-Hispanic white, male												
2002	1.6	*	*	*	*	*	0.3	0.8	1.7	4.4	12.4	32.6
2001	1.7	*	*	*	*	*	0.3	0.8	2.0	4.5	12.7	36.3
2000	1.9	*	*	*	*	*	0.3	1.1	2.3	5.2	13.7	41.0
1999	1.8	*	*	*	*	*	0.3	0.8	2.1	5.9	14.9	38.6
Non-Hispanic white, female												
2002	1.8	*	*	*	*	*	0.2	0.5	1.1	3.0	9.0	27.9
2001	2.1	*	*	*	*	*	0.1	0.5	1.1	3.4	10.9	34.0
2000	2.0	*	*	*	*	*	0.1	0.4	1.3	3.1	10.2	34.4
1999	2.1	*	*	*	*	*	0.2	0.4	1.3	3.2	10.8	34.4
Non-Hispanic black, both sexes												
2002	1.0	*	*	*	*	*	0.6	1.2	2.6	4.2	7.9	18.8
2001	1.1	*	*	*	*	*	0.6	1.4	2.8	4.5	9.0	20.9
2000	1.2	*	*	*	*	*	0.5	1.7	2.6	4.9	10.1	26.7
1999	1.3	*	*	*	*	*	0.8	1.5	2.8	5.7	11.0	25.5
Non-Hispanic black, male												
2002	1.2	*	*	*	*	*	0.8	2.0	3.9	6.0	11.1	*
2001	1.4	*	*	*	*	*	1.0	2.3	4.3	7.0	11.7	24.3
2000	1.4	*	*	*	*	*	0.8	2.4	3.8	7.2	13.9	31.7
1999	1.5	*	*	*	*	*	1.2	2.4	4.5	8.3	12.2	28.5
Non-Hispanic black, female												
2002	0.9	*	*	*	*	*	*	*	1.6	2.9	6.1	18.4
2001	0.9	*	*	*	*	*	*	*	1.5	2.7	7.4	19.7
2000	1.0	*	*	*	*	*	*	1.1	1.7	3.3	8.0	24.9
1999	1.1	*	*	*	*	*	*	*	*	3.9	10.3	24.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastric ulcer (K25)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.8	7.7
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.3	3.3	8.0
2000	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0	9.5
1999	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.5	8.8
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.1	0.3	0.7	1.4	2.9	8.1
2001	0.4	*	*	*	*	*	0.1	0.3	0.6	1.4	3.4	6.0
2000	0.4	*	*	*	*	*	0.1	0.4	0.7	1.6	3.3	9.9
1999	0.4	*	*	*	*	*	0.1	0.4	0.7	1.5	3.7	8.7
All origins 3/, female												
2002	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.8	7.5
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.1	3.2	8.8
2000	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.9	9.4
1999	0.5	*	*	*	*	*	0.1	0.2	0.5	1.1	3.4	8.8
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	3.8	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.5	1.1	2.9	7.7
2001	0.5	*	*	*	*	*	0.1	0.3	0.5	1.3	3.3	8.2
2000	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.0	9.4
1999	0.5	*	*	*	*	*	0.1	0.3	0.5	1.3	3.5	8.8
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	0.1	0.3	0.6	1.3	2.9	7.9
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.4	3.3	6.3
2000	0.5	*	*	*	*	*	0.1	0.4	0.7	1.6	3.3	9.6
1999	0.5	*	*	*	*	*	0.1	0.4	0.6	1.5	3.7	8.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastric ulcer (K25)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.8	7.6
2001	0.6	*	*	*	*	*	*	0.2	0.4	1.1	3.3	9.0
2000	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.8	9.3
1999	0.6	*	*	*	*	*	0.1	0.2	0.5	1.1	3.4	8.8
Non-Hispanic white, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.4	1.1	2.8	7.7
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.3	8.2
2000	0.5	*	*	*	*	*	0.1	0.3	0.5	1.2	3.0	9.3
1999	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	3.5	8.7
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	*	0.2	0.5	1.2	2.8	8.0
2001	0.4	*	*	*	*	*	*	0.2	0.5	1.2	3.3	6.1
2000	0.5	*	*	*	*	*	*	0.4	0.5	1.4	3.2	8.9
1999	0.5	*	*	*	*	*	*	0.3	0.5	1.4	3.6	8.1
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.0	2.7	7.6
2001	0.6	*	*	*	*	*	*	0.2	0.4	1.1	3.2	9.0
2000	0.6	*	*	*	*	*	*	0.1	0.5	1.1	2.9	9.4
1999	0.6	*	*	*	*	*	*	0.2	0.5	1.1	3.5	8.9
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	0.5	0.9	*	3.5	6.4
2001	0.4	*	*	*	*	*	*	0.5	0.9	2.0	2.7	7.5
2000	0.5	*	*	*	*	*	*	0.8	1.2	1.9	3.2	9.2
1999	0.5	*	*	*	*	*	0.4	0.7	1.2	2.2	3.4	9.7
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	1.8	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	3.4	*	*
2000	0.6	*	*	*	*	*	*	*	1.9	3.6	*	*
1999	0.6	*	*	*	*	*	*	1.2	*	3.0	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	3.6	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Duodenal ulcer (K26)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.6	1.3	3.6	9.0
2001	0.5	*	*	*	*	*	0.1	0.3	0.6	1.3	3.9	11.0
2000	0.5	*	*	*	*	*	0.1	0.3	0.7	1.4	4.0	10.1
1999	0.5	*	*	*	*	*	0.1	0.3	0.7	1.5	3.7	10.0
All origins 3/, male												
2002	0.5	*	*	*	*	*	0.1	0.3	0.8	1.7	4.5	10.1
2001	0.6	*	*	*	*	*	0.2	0.4	0.8	1.7	4.7	13.8
2000	0.6	*	*	*	*	*	0.2	0.5	1.0	1.8	4.8	12.3
1999	0.6	*	*	*	*	*	0.1	0.4	0.9	2.4	5.1	11.9
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.0	8.5
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.4	9.8
2000	0.5	*	*	*	*	*	*	0.1	0.5	1.0	3.6	9.2
1999	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.9	9.3
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.6	1.3	3.7	9.1
2001	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	4.0	11.0
2000	0.6	*	*	*	*	*	0.1	0.3	0.7	1.4	4.2	10.2
1999	0.6	*	*	*	*	*	0.1	0.3	0.7	1.6	3.8	10.1
Non-Hispanic 4/, male												
2002	0.6	*	*	*	*	*	0.2	0.3	0.8	1.7	4.7	10.3
2001	0.6	*	*	*	*	*	0.2	0.5	0.8	1.8	4.7	13.8
2000	0.6	*	*	*	*	*	0.2	0.5	0.9	1.8	4.9	12.5
1999	0.6	*	*	*	*	*	0.1	0.4	0.9	2.4	5.1	12.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Duodenal ulcer (K26)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.0	8.6
2001	0.6	*	*	*	*	*	*	0.2	0.4	1.0	3.5	9.8
2000	0.6	*	*	*	*	*	*	0.1	0.5	1.0	3.7	9.2
1999	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.9	9.3
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	0.1	0.2	0.6	1.3	3.8	9.7
2001	0.7	*	*	*	*	*	0.1	0.3	0.6	1.4	4.1	11.7
2000	0.7	*	*	*	*	*	0.1	0.3	0.7	1.4	4.3	10.6
1999	0.6	*	*	*	*	*	0.1	0.2	0.7	1.6	3.9	10.3
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	0.1	0.3	0.8	1.7	4.8	11.1
2001	0.7	*	*	*	*	*	0.2	0.4	0.8	1.8	4.8	14.6
2000	0.7	*	*	*	*	*	0.1	0.4	1.0	1.8	5.0	12.9
1999	0.7	*	*	*	*	*	*	0.3	1.0	2.4	5.3	12.8
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	0.1	0.4	1.1	3.2	9.1
2001	0.7	*	*	*	*	*	*	0.2	0.4	1.1	3.7	10.5
2000	0.7	*	*	*	*	*	*	0.1	0.5	1.0	3.8	9.6
1999	0.6	*	*	*	*	*	*	0.2	0.5	0.9	3.0	9.3
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	0.4	1.0	1.5	*	*
2001	0.3	*	*	*	*	*	*	0.6	0.9	*	2.4	*
2000	0.3	*	*	*	*	*	*	0.6	*	1.5	2.8	*
1999	0.4	*	*	*	*	*	*	0.6	*	1.3	3.0	6.5
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	1.1	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer, site unspecified (K27)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.6	11.3
2001	0.6	*	*	*	*	*	0.1	0.2	0.6	1.3	4.1	14.2
2000	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.2	15.5
1999	0.6	*	*	*	*	*	0.1	0.1	0.5	1.6	4.9	15.6
All origins 3/, male												
2002	0.5	*	*	*	*	*	*	0.3	0.4	1.4	4.6	12.8
2001	0.5	*	*	*	*	*	*	0.2	0.8	1.6	4.8	14.9
2000	0.6	*	*	*	*	*	*	0.3	0.8	1.9	5.4	18.2
1999	0.6	*	*	*	*	*	*	0.2	0.7	2.1	5.8	16.4
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.9	10.7
2001	0.6	*	*	*	*	*	*	0.1	0.4	1.0	3.7	13.9
2000	0.6	*	*	*	*	*	*	0.1	0.3	1.0	3.5	14.5
1999	0.7	*	*	*	*	*	*	0.1	0.4	1.2	4.3	15.2
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	3.8	*
2001	0.2	*	*	*	*	*	*	*	*	*	4.1	12.5
2000	0.2	*	*	*	*	*	*	*	*	*	4.7	*
1999	0.2	*	*	*	*	*	*	*	*	*	4.4	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.6	11.5
2001	0.6	*	*	*	*	*	0.1	0.2	0.6	1.3	4.1	14.2
2000	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.2	15.7
1999	0.7	*	*	*	*	*	0.1	0.1	0.5	1.7	4.9	15.7
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	*	0.3	0.4	1.4	4.5	13.0
2001	0.6	*	*	*	*	*	*	0.2	0.8	1.6	4.6	15.2
2000	0.6	*	*	*	*	*	*	0.3	0.8	1.8	5.4	18.6
1999	0.6	*	*	*	*	*	*	0.2	0.7	2.2	5.8	16.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer, site unspecified (K27)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	*	*	*	0.1	0.4	0.9	2.9	10.8
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.1	3.8	13.8
2000	0.7	*	*	*	*	*	*	0.1	0.3	1.0	3.4	14.5
1999	0.7	*	*	*	*	*	*	*	0.4	1.2	4.3	15.4
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	*	0.2	0.4	1.2	3.6	11.7
2001	0.7	*	*	*	*	*	*	0.2	0.5	1.3	4.1	14.6
2000	0.7	*	*	*	*	*	0.1	0.2	0.5	1.4	4.2	16.0
1999	0.8	*	*	*	*	*	*	0.1	0.5	1.6	4.9	16.3
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	*	0.3	0.4	1.4	4.6	13.1
2001	0.6	*	*	*	*	*	*	0.2	0.7	1.5	4.5	15.4
2000	0.7	*	*	*	*	*	*	0.2	0.8	1.8	5.4	18.8
1999	0.7	*	*	*	*	*	*	0.2	0.7	2.1	5.8	17.3
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	0.2	0.4	1.0	3.0	11.0
2001	0.8	*	*	*	*	*	*	0.1	0.4	1.1	3.8	14.3
2000	0.7	*	*	*	*	*	*	*	0.2	1.0	3.4	14.9
1999	0.8	*	*	*	*	*	*	*	0.4	1.2	4.3	16.0
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	0.8	1.5	2.4	9.1
2001	0.4	*	*	*	*	*	*	*	1.1	1.3	3.9	9.7
2000	0.4	*	*	*	*	*	*	*	*	1.2	3.8	12.1
1999	0.4	*	*	*	*	*	*	*	0.9	2.2	4.5	9.4
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	1.9	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	3.3	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	8.4
2001	0.3	*	*	*	*	*	*	*	*	*	*	8.6
2000	0.4	*	*	*	*	*	*	*	*	*	3.6	11.3
1999	0.4	*	*	*	*	*	*	*	*	*	4.2	10.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastrojejunal ulcer (K28)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastrojejunal ulcer (K28)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastritis and duodenitis (K29)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.8	2.9
2001	0.1	*	*	*	*	*	*	*	0.1	0.3	0.7	3.8
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.8	3.5
1999	0.1	*	*	*	*	*	0.0	0.1	0.1	0.3	0.9	3.8
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	0.2	0.3	0.9	2.9
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	4.8
2000	0.1	*	*	*	*	*	*	*	0.3	0.4	0.8	3.3
1999	0.1	*	*	*	*	*	*	0.2	*	0.3	1.1	3.7
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	2.9
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.3
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.9	3.6
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.8	3.9
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.8	3.0
2001	0.1	*	*	*	*	*	*	*	0.1	0.3	0.7	3.7
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.8	3.4
1999	0.2	*	*	*	*	*	*	0.1	0.1	0.3	0.9	3.8
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	0.2	0.3	0.9	2.9
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	4.6
2000	0.1	*	*	*	*	*	*	*	0.3	0.4	0.8	3.3
1999	0.1	*	*	*	*	*	*	0.2	*	0.3	1.1	3.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Gastritis and duodenitis (K29)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	3.0
2001	0.2	*	*	*	*	*	*	*	*	0.3	0.8	3.4
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.8	3.4
1999	0.2	*	*	*	*	*	*	*	*	0.3	0.7	3.9
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.1	0.2	0.8	3.0
2001	0.2	*	*	*	*	*	*	*	0.1	0.3	0.7	4.0
2000	0.2	*	*	*	*	*	*	*	0.2	0.3	0.9	3.4
1999	0.2	*	*	*	*	*	*	0.1	0.1	0.2	0.7	3.9
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	1.0	3.0
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	4.8
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	0.8	3.3
1999	0.1	*	*	*	*	*	*	0.2	*	*	0.9	3.8
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	*	0.6	2.9
2001	0.2	*	*	*	*	*	*	*	*	*	0.7	3.7
2000	0.2	*	*	*	*	*	*	*	*	*	0.9	3.4
1999	0.2	*	*	*	*	*	*	*	*	*	0.6	4.0
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	3.2
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.6
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.3	4.0
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.2
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.9
2001	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	3.3
2000	0.2	*	*	*	*	*	*	*	0.2	0.8	1.4	4.3
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	4.8
All origins 3/, female												
2002	0.2	*	*	*	*	*	0.1	0.1	0.3	0.4	1.4	3.4
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	3.7
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.3	3.9
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	3.3
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.6
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.4	4.1
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.1
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.7	1.3	2.8
2001	0.2	*	*	*	*	*	*	*	*	0.6	1.4	3.2
2000	0.2	*	*	*	*	*	*	*	*	0.8	1.5	4.3
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.4	4.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Dyspepsia and other diseases of stomach and duodenum (K30-K31)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	0.1	0.1	0.3	0.5	1.4	3.5
2001	0.2	*	*	*	*	*	*	0.1	*	0.5	1.2	3.8
2000	0.3	*	*	*	*	*	*	0.1	0.3	0.4	1.3	3.9
1999	0.3	*	*	*	*	*	*	*	0.2	0.4	1.4	4.0
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.3	3.3
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	3.8
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	4.2
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.4	4.2
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.7
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.4	3.4
2000	0.2	*	*	*	*	*	*	*	*	0.7	1.5	4.4
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.4	4.5
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.1	0.3	0.5	1.3	3.6
2001	0.3	*	*	*	*	*	*	0.2	*	0.5	1.2	3.9
2000	0.3	*	*	*	*	*	*	0.1	0.3	0.4	1.3	4.1
1999	0.3	*	*	*	*	*	*	*	0.2	0.4	1.4	4.0
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	2.1
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.2
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.7
1999	0.1	*	*	*	*	*	0.0	0.1	0.2	0.4	0.8	1.5
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.3	2.3
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.0	3.0
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.4	1.0	3.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.0	2.1
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.8
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.6
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.4	2.1
2001	0.2	*	*	*	*	*	*	*	0.2	0.7	1.0	2.6
2000	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.1
1999	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.4	2.0
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.0
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
All origins 3/, both sexes												
2002	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7
All origins 3/, male												
2002	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2
All origins 3/, female												
2002	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	4.3	12.5
2000	0.2	*	*	*	*	*	*	*	*	*	4.5	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.2	3.8	13.0
2001	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.5
2000	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.1
1999	0.6	1.3	*	*	*	*	0.1	0.1	0.5	1.1	3.7	12.7
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.3	3.7	11.9
2001	0.5	1.2	*	*	*	*	*	0.2	0.5	1.2	3.5	12.1
2000	0.5	1.4	*	*	*	*	*	0.2	0.4	1.1	3.9	12.8
1999	0.5	1.5	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Non-Hispanic 4/, female												
2002	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.4
2001	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.5	12.7
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.2
1999	0.7	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.4
Non-Hispanic white, both sexes												
2002	0.7	0.9	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.6
2001	0.6	0.9	*	*	*	*	*	0.2	0.5	1.2	3.5	12.8
2000	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.7	13.3
1999	0.6	1.4	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.8	12.2
2001	0.5	*	*	*	*	*	*	0.2	0.5	1.2	3.5	11.9
2000	0.5	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.0
1999	0.5	*	*	*	*	*	*	0.2	0.6	1.1	4.1	13.0
Non-Hispanic white, female												
2002	0.8	*	*	*	*	*	*	0.2	0.4	1.2	4.1	14.2
2001	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.5	13.1
2000	0.8	*	*	*	*	*	*	0.2	0.5	1.1	3.6	13.4
1999	0.7	*	*	*	*	*	*	*	0.4	1.1	3.4	12.9
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.8	1.6	3.2	7.9
2001	0.4	*	*	*	*	*	*	*	0.9	1.5	3.6	11.2
2000	0.4	*	*	*	*	*	*	*	*	1.3	4.0	12.7
1999	0.4	*	*	*	*	*	*	*	*	1.5	4.2	10.0
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	9.0
2000	0.5	*	*	*	*	*	*	*	*	*	4.3	12.6
1999	0.5	*	*	*	*	*	*	*	*	*	4.2	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Crohn's disease and ulcerative colitis (K50-K51)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	0.2	0.2	0.3	0.9	2.0	3.1
2001	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.9	1.9	3.8
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9	3.3
1999	0.3	*	*	*	*	0.1	0.1	0.3	0.4	0.8	2.0	3.3
All origins 3/, male												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.9	2.7
2001	0.3	*	*	*	*	*	*	0.2	0.5	1.0	1.9	3.5
2000	0.2	*	*	*	*	*	*	0.2	0.3	0.8	1.6	2.9
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.9	2.7
All origins 3/, female												
2002	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	2.0	3.2
2001	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	1.9	3.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	2.0	3.4
1999	0.3	*	*	*	*	*	0.1	0.3	0.4	0.8	2.0	3.5
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	*	*	0.2	0.3	0.3	0.9	2.0	3.2
2001	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0	3.8
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.9	3.4
1999	0.3	*	*	*	*	0.1	0.1	0.3	0.4	0.8	2.0	3.4
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	0.1	0.2	0.3	0.9	2.0	2.9
2001	0.3	*	*	*	*	*	*	0.2	0.5	1.0	1.9	3.6
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.8	1.7	3.0
1999	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.0	2.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Crohn's disease and ulcerative colitis (K50-K51)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	0.2	0.3	0.4	0.8	2.0	3.3
2001	0.4	*	*	*	*	*	0.1	0.2	0.5	0.8	2.0	3.9
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	2.0	3.5
1999	0.4	*	*	*	*	*	0.1	0.3	0.4	0.8	2.1	3.6
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.2	0.3	0.4	1.0	2.2	3.5
2001	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.1	3.9
2000	0.3	*	*	*	*	*	0.1	0.2	0.4	1.0	2.0	3.6
1999	0.4	*	*	*	*	*	0.1	0.3	0.4	0.9	2.1	3.6
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	0.1	0.2	0.3	1.0	2.1	3.1
2001	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.1	3.6
2000	0.3	*	*	*	*	*	*	0.2	0.4	0.9	1.8	3.2
1999	0.3	*	*	*	*	*	*	0.2	0.4	1.0	2.1	3.1
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	0.2	0.3	0.4	1.0	2.2	3.7
2001	0.4	*	*	*	*	*	*	0.2	0.5	0.9	2.1	4.1
2000	0.4	*	*	*	*	*	*	0.2	0.4	1.1	2.2	3.7
1999	0.4	*	*	*	*	*	0.1	0.4	0.4	0.9	2.2	3.8
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
All origins 3/, both sexes												
2002	5.0	5.8	0.1	0.1	0.1	0.1	0.4	1.1	3.7	12.8	39.4	113.3
2001	5.0	5.8	0.1	0.1	*	0.1	0.3	1.2	3.9	13.8	38.9	113.5
2000	5.1	6.0	0.1	0.1	0.1	0.1	0.4	1.3	3.9	13.4	40.1	116.2
1999	5.0	6.1	0.1	0.1	0.1	0.1	0.4	1.1	4.0	13.5	38.9	116.2
All origins 3/, male												
2002	3.6	6.5	*	*	0.1	0.1	0.4	1.1	3.4	11.6	35.8	96.7
2001	3.6	6.5	*	*	*	0.1	0.3	1.1	3.5	13.1	35.2	98.5
2000	3.6	6.3	*	0.1	0.1	0.1	0.4	1.2	3.8	12.5	36.8	98.5
1999	3.5	7.6	*	*	*	0.1	0.4	1.1	3.8	12.5	35.6	95.6
All origins 3/, female												
2002	6.4	5.0	*	*	*	0.1	0.3	1.1	3.8	13.8	41.7	120.5
2001	6.4	5.1	*	*	*	0.1	0.3	1.2	4.1	14.4	41.3	119.8
2000	6.5	5.8	*	*	*	0.1	0.4	1.3	4.0	14.0	42.1	123.4
1999	6.4	4.4	*	*	*	0.2	0.4	1.1	4.2	14.3	41.1	124.5
Hispanic, both sexes												
2002	1.5	5.3	*	*	*	*	*	0.8	2.1	10.8	31.4	75.9
2001	1.5	5.1	*	*	*	*	*	0.6	3.0	10.0	29.8	95.7
2000	1.5	5.1	*	*	*	*	*	0.8	2.3	9.8	30.4	88.3
1999	1.5	6.1	*	*	*	*	*	0.8	2.9	9.7	29.8	92.6
Hispanic, male												
2002	1.2	5.2	*	*	*	*	*	*	*	12.4	32.0	60.2
2001	1.2	5.5	*	*	*	*	*	*	2.5	8.2	28.2	104.8
2000	1.2	*	*	*	*	*	*	*	*	10.8	33.9	70.5
1999	1.2	8.5	*	*	*	*	*	*	*	8.6	31.3	69.0
Hispanic, female												
2002	1.9	5.4	*	*	*	*	*	*	2.5	9.6	30.9	83.7
2001	1.9	*	*	*	*	*	*	*	3.4	11.5	30.8	91.2
2000	1.8	5.6	*	*	*	*	*	*	2.6	9.1	28.1	96.9
1999	1.9	*	*	*	*	*	*	*	3.5	10.7	28.8	104.0
Non-Hispanic 4/, both sexes												
2002	5.6	5.9	0.2	0.1	0.1	0.1	0.4	1.2	3.8	12.9	39.7	114.6
2001	5.5	5.8	*	0.1	*	0.1	0.3	1.2	3.9	14.0	39.2	113.9
2000	5.6	6.1	*	0.1	0.1	0.1	0.4	1.3	4.0	13.5	40.4	117.0
1999	5.5	5.9	*	0.1	0.1	0.2	0.4	1.1	4.1	13.7	39.2	116.9
Non-Hispanic 4/, male												
2002	4.0	6.8	*	*	*	0.2	0.5	1.2	3.6	11.6	35.9	98.0
2001	4.0	6.7	*	*	*	0.1	0.4	1.2	3.6	13.3	35.4	98.2
2000	4.0	6.4	*	0.1	0.1	0.1	0.4	1.2	3.9	12.6	36.9	99.4
1999	3.9	7.2	*	*	*	0.1	0.4	1.1	3.9	12.7	35.7	96.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Vascular disorders and obstruction of intestine without hernia (K55-K56)												
Non-Hispanic 4/, female												
2002	7.1	4.9	*	*	*	0.1	0.3	1.2	3.9	14.1	42.2	121.8
2001	7.0	4.9	*	*	*	0.1	0.3	1.3	4.2	14.6	41.7	120.4
2000	7.1	5.9	*	*	*	0.1	0.4	1.3	4.1	14.3	42.6	124.1
1999	7.0	4.6	*	*	*	0.2	0.4	1.2	4.3	14.5	41.4	125.0
Non-Hispanic white, both sexes												
2002	6.1	3.7	*	*	0.1	0.1	0.4	1.0	3.6	12.5	39.8	117.3
2001	6.1	3.5	*	*	*	0.1	0.3	1.1	3.6	13.8	39.2	116.5
2000	6.0	3.3	*	0.1	0.1	0.1	0.3	1.1	3.6	13.3	40.1	118.8
1999	6.0	3.5	*	*	0.1	0.1	0.3	1.0	4.0	13.1	39.0	118.7
Non-Hispanic white, male												
2002	4.2	4.3	*	*	*	*	0.4	1.0	3.2	11.0	35.5	98.9
2001	4.2	3.7	*	*	*	*	0.3	1.0	3.3	12.8	35.0	98.3
2000	4.1	3.4	*	*	*	*	0.3	1.0	3.3	12.0	35.9	99.7
1999	4.1	4.3	*	*	*	*	0.3	0.9	3.7	12.1	35.1	95.5
Non-Hispanic white, female												
2002	7.9	3.2	*	*	*	*	0.3	1.1	3.9	13.8	42.7	125.3
2001	7.8	3.2	*	*	*	*	0.3	1.1	3.8	14.6	42.0	124.1
2000	7.9	3.2	*	*	*	*	0.4	1.1	3.9	14.3	43.0	126.5
1999	7.8	2.7	*	*	*	0.2	0.3	1.0	4.3	14.0	41.5	127.9
Non-Hispanic black, both sexes												
2002	4.2	14.5	*	*	*	*	0.7	2.4	5.9	18.8	45.1	101.0
2001	4.3	16.5	*	*	*	*	0.6	2.8	7.5	19.6	45.7	97.7
2000	4.5	17.2	*	*	*	*	0.8	3.2	7.8	18.1	48.1	112.9
1999	4.3	16.0	*	*	*	*	1.0	2.4	5.8	20.8	47.6	110.3
Non-Hispanic black, male												
2002	3.6	17.3	*	*	*	*	0.9	2.6	7.0	19.0	46.0	106.1
2001	3.9	19.8	*	*	*	*	0.8	2.8	7.2	22.7	48.3	122.4
2000	4.1	18.4	*	*	*	*	1.0	3.6	9.9	19.9	55.2	111.5
1999	3.6	18.7	*	*	*	*	1.1	2.8	6.3	19.8	50.0	111.7
Non-Hispanic black, female												
2002	4.7	11.7	*	*	*	*	*	2.2	5.1	18.6	44.6	99.1
2001	4.7	13.1	*	*	*	*	*	2.8	7.8	17.5	44.3	88.5
2000	4.8	15.9	*	*	*	*	0.7	2.9	6.1	16.7	44.1	113.4
1999	5.0	13.2	*	*	*	*	0.9	2.1	5.4	21.5	46.2	109.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diverticular disease of intestine (K57)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.2	0.7	2.7	9.9	27.7
2001	1.2	*	*	*	*	*	0.0	0.2	0.9	2.9	9.4	31.4
2000	1.2	*	*	*	*	*	0.1	0.2	0.7	3.0	9.2	30.9
1999	1.2	*	*	*	*	*	0.1	0.2	0.9	3.1	9.3	33.4
All origins 3/, male												
2002	0.8	*	*	*	*	*	0.1	0.3	0.7	2.4	8.0	19.9
2001	0.8	*	*	*	*	*	*	0.3	0.8	2.4	8.2	23.4
2000	0.7	*	*	*	*	*	*	0.3	0.7	2.7	7.3	22.3
1999	0.8	*	*	*	*	*	*	0.2	0.8	2.7	7.4	25.8
All origins 3/, female												
2002	1.5	*	*	*	*	*	*	0.2	0.7	2.9	11.1	31.0
2001	1.6	*	*	*	*	*	*	0.2	1.0	3.3	10.2	34.7
2000	1.6	*	*	*	*	*	*	0.2	0.7	3.1	10.5	34.4
1999	1.7	*	*	*	*	*	*	0.2	0.9	3.4	10.5	36.4
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	2.1	9.2	16.6
2001	0.4	*	*	*	*	*	*	*	*	2.3	7.6	22.5
2000	0.3	*	*	*	*	*	*	*	*	2.8	7.1	21.2
1999	0.4	*	*	*	*	*	*	*	*	2.4	8.1	33.0
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	10.4	17.4
2001	0.4	*	*	*	*	*	*	*	*	*	8.8	27.0
2000	0.5	*	*	*	*	*	*	*	*	*	9.2	22.8
1999	0.5	*	*	*	*	*	*	*	*	*	8.0	35.4
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	*	*	0.1	0.3	0.7	2.7	9.9	28.0
2001	1.3	*	*	*	*	*	*	0.2	0.9	2.9	9.5	31.6
2000	1.3	*	*	*	*	*	0.1	0.2	0.7	3.0	9.3	31.2
1999	1.3	*	*	*	*	*	0.1	0.2	0.9	3.2	9.3	33.3
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	*	*	0.4	0.7	2.4	8.0	20.1
2001	0.9	*	*	*	*	*	*	0.3	0.8	2.3	8.3	23.8
2000	0.8	*	*	*	*	*	*	0.2	0.7	2.7	7.4	22.5
1999	0.8	*	*	*	*	*	*	0.2	0.9	2.7	7.3	25.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diverticular disease of intestine (K57)												
Non-Hispanic 4/, female												
2002	1.7	*	*	*	*	*	*	0.2	0.7	2.9	11.1	31.4
2001	1.8	*	*	*	*	*	*	0.2	1.0	3.4	10.2	34.9
2000	1.7	*	*	*	*	*	*	0.2	0.7	3.1	10.5	34.7
1999	1.8	*	*	*	*	*	*	0.2	0.9	3.5	10.6	36.4
Non-Hispanic white, both sexes												
2002	1.5	*	*	*	*	*	0.1	0.3	0.7	2.8	10.0	28.9
2001	1.5	*	*	*	*	*	*	0.2	0.9	2.9	9.9	32.7
2000	1.5	*	*	*	*	*	*	0.2	0.7	3.0	9.6	32.6
1999	1.5	*	*	*	*	*	*	0.2	0.9	3.2	9.6	34.5
Non-Hispanic white, male												
2002	0.9	*	*	*	*	*	*	0.4	0.7	2.5	7.8	20.8
2001	1.0	*	*	*	*	*	*	0.3	0.8	2.3	8.5	24.8
2000	0.9	*	*	*	*	*	*	0.2	0.7	2.7	7.8	23.9
1999	0.9	*	*	*	*	*	*	0.2	0.8	2.8	7.5	26.6
Non-Hispanic white, female												
2002	2.0	*	*	*	*	*	*	0.2	0.7	3.0	11.5	32.4
2001	2.1	*	*	*	*	*	*	0.2	1.0	3.3	10.8	36.0
2000	2.0	*	*	*	*	*	*	0.2	0.7	3.2	10.8	36.1
1999	2.1	*	*	*	*	*	*	0.2	0.9	3.5	11.0	37.7
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	*	0.9	2.6	10.0	22.1
2001	0.7	*	*	*	*	*	*	*	1.1	4.0	6.9	24.3
2000	0.7	*	*	*	*	*	*	*	1.3	3.8	7.2	20.4
1999	0.7	*	*	*	*	*	*	*	1.5	3.5	7.3	23.9
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	11.4	*
2001	0.4	*	*	*	*	*	*	*	*	3.0	7.2	*
2000	0.4	*	*	*	*	*	*	*	*	3.7	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	6.6	*
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	*	*	3.1	9.1	25.1
2001	0.9	*	*	*	*	*	*	*	*	4.7	6.7	27.8
2000	0.9	*	*	*	*	*	*	*	*	3.8	8.7	24.9
1999	1.0	*	*	*	*	*	*	*	1.6	4.2	7.7	27.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)												
All origins 3/, both sexes												
2002	1.7	2.0	0.2	0.0	0.0	0.1	0.2	0.6	1.6	4.3	12.5	32.8
2001	1.7	1.4	0.2	*	*	0.1	0.2	0.6	1.4	4.5	12.8	34.1
2000	1.6	1.7	*	*	*	0.1	0.2	0.5	1.6	4.1	12.4	33.3
1999	1.6	1.5	0.2	*	*	0.1	0.2	0.5	1.5	4.4	12.1	32.0
All origins 3/, male												
2002	1.3	2.4	*	*	*	*	0.2	0.6	1.5	4.4	11.8	29.4
2001	1.2	1.6	*	*	*	*	0.2	0.5	1.4	4.2	12.1	29.9
2000	1.2	1.8	*	*	*	*	0.2	0.4	1.7	4.0	11.4	28.7
1999	1.2	2.0	*	*	*	*	0.2	0.6	1.5	4.2	11.9	28.9
All origins 3/, female												
2002	2.0	1.5	*	*	*	*	0.2	0.5	1.7	4.3	13.0	34.3
2001	2.1	1.1	*	*	*	*	0.2	0.6	1.5	4.7	13.3	35.8
2000	2.0	1.6	*	*	*	*	0.2	0.6	1.4	4.2	13.1	35.2
1999	1.9	*	*	*	*	*	0.2	0.5	1.6	4.6	12.2	33.2
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	*	1.3	2.3	8.3	14.4
2001	0.5	*	*	*	*	*	*	*	*	3.2	11.0	25.6
2000	0.5	*	*	*	*	*	*	*	*	2.1	10.1	21.2
1999	0.6	*	*	*	*	*	*	0.7	1.4	3.3	8.1	21.7
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	8.2	*
2001	0.5	*	*	*	*	*	*	*	*	*	13.4	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	8.4	*
2001	0.6	*	*	*	*	*	*	*	*	3.1	9.4	28.9
2000	0.5	*	*	*	*	*	*	*	*	*	11.2	20.8
1999	0.6	*	*	*	*	*	*	*	*	*	9.4	27.0
Non-Hispanic 4/, both sexes												
2002	1.9	1.9	0.2	*	0.1	0.1	0.3	0.6	1.6	4.5	12.7	33.5
2001	1.8	1.4	*	*	*	*	0.2	0.6	1.5	4.5	12.9	34.4
2000	1.8	1.6	*	*	*	0.1	0.2	0.6	1.6	4.2	12.5	33.7
1999	1.7	1.3	0.2	*	*	0.1	0.2	0.5	1.5	4.5	12.3	32.2
Non-Hispanic 4/, male												
2002	1.4	2.4	*	*	*	*	0.2	0.6	1.5	4.6	12.0	29.9
2001	1.4	1.7	*	*	*	*	0.2	0.6	1.4	4.2	12.1	30.4
2000	1.3	1.8	*	*	*	*	0.2	0.5	1.7	4.1	11.4	28.9
1999	1.3	1.7	*	*	*	*	0.2	0.6	1.4	4.2	12.1	29.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)												
Non-Hispanic 4/, female												
2002	2.3	1.4	*	*	*	*	0.3	0.6	1.7	4.4	13.2	35.1
2001	2.3	*	*	*	*	*	0.3	0.7	1.5	4.8	13.5	36.0
2000	2.2	1.5	*	*	*	*	0.2	0.6	1.5	4.3	13.2	35.6
1999	2.1	*	*	*	*	*	0.2	0.4	1.7	4.7	12.3	33.3
Non-Hispanic white, both sexes												
2002	2.1	1.4	*	*	*	*	0.2	0.6	1.6	4.5	13.1	34.9
2001	2.0	*	*	*	*	*	0.2	0.6	1.4	4.6	12.9	35.5
2000	1.9	1.2	*	*	*	0.1	0.2	0.5	1.5	4.3	12.7	34.1
1999	1.9	0.9	*	*	*	0.1	0.2	0.5	1.5	4.5	12.4	33.4
Non-Hispanic white, male												
2002	1.6	2.2	*	*	*	*	0.2	0.6	1.5	4.4	12.1	30.9
2001	1.5	*	*	*	*	*	0.2	0.6	1.2	4.1	12.0	31.7
2000	1.4	*	*	*	*	*	0.2	0.4	1.5	4.2	11.4	28.3
1999	1.4	*	*	*	*	*	0.2	0.6	1.3	4.3	12.1	28.9
Non-Hispanic white, female												
2002	2.6	*	*	*	*	*	0.3	0.6	1.7	4.6	13.8	36.6
2001	2.6	*	*	*	*	*	0.2	0.6	1.5	5.1	13.4	37.0
2000	2.5	*	*	*	*	*	0.2	0.6	1.5	4.3	13.6	36.5
1999	2.4	*	*	*	*	*	0.2	0.4	1.7	4.7	12.6	35.1
Non-Hispanic black, both sexes												
2002	1.2	3.7	*	*	*	*	0.4	0.7	2.4	5.4	10.8	23.6
2001	1.4	4.1	*	*	*	*	0.4	1.0	2.5	5.0	16.0	27.5
2000	1.2	3.6	*	*	*	*	0.4	0.8	2.6	4.8	10.9	29.9
1999	1.2	*	*	*	*	*	0.4	1.0	1.9	5.2	11.8	23.3
Non-Hispanic black, male												
2002	1.1	*	*	*	*	*	*	*	2.5	7.6	12.3	25.2
2001	1.2	*	*	*	*	*	*	*	3.1	7.0	16.5	*
2000	1.1	*	*	*	*	*	*	*	3.2	4.0	13.6	29.4
1999	1.1	*	*	*	*	*	*	1.3	1.9	4.7	12.8	40.4
Non-Hispanic black, female												
2002	1.3	*	*	*	*	*	*	*	2.3	3.8	9.9	23.0
2001	1.5	*	*	*	*	*	*	1.2	2.0	3.6	15.7	29.9
2000	1.4	*	*	*	*	*	*	1.0	2.1	5.3	9.4	30.1
1999	1.2	*	*	*	*	*	*	*	1.9	5.5	11.2	16.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of peritoneum (K65-K66)												
All origins 3/, both sexes												
2002	0.7	*	*	*	*	0.1	0.2	0.5	0.8	1.8	4.8	10.3
2001	0.7	*	*	*	*	*	0.2	0.5	0.8	1.8	4.4	10.4
2000	0.7	*	*	*	*	0.1	0.2	0.4	1.0	2.0	4.4	10.6
1999	0.7	0.5	*	*	*	0.1	0.2	0.5	0.9	2.2	4.4	9.7
All origins 3/, male												
2002	0.5	*	*	*	*	*	0.2	0.5	0.9	1.6	4.3	9.1
2001	0.6	*	*	*	*	*	0.2	0.5	1.0	2.0	4.6	10.1
2000	0.6	*	*	*	*	*	0.2	0.5	1.1	2.0	4.6	11.0
1999	0.6	*	*	*	*	*	0.2	0.6	1.0	2.4	4.9	10.1
All origins 3/, female												
2002	0.8	*	*	*	*	*	0.2	0.4	0.8	1.9	5.1	10.8
2001	0.7	*	*	*	*	*	0.1	0.4	0.7	1.7	4.3	10.5
2000	0.8	*	*	*	*	*	0.2	0.4	0.9	2.0	4.3	10.5
1999	0.7	*	*	*	*	*	0.2	0.3	0.8	2.0	4.1	9.6
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	1.1	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	3.6	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	*	*	0.1	0.2	0.5	0.8	1.8	4.9	10.5
2001	0.7	*	*	*	*	*	0.2	0.5	0.9	1.8	4.5	10.4
2000	0.8	*	*	*	*	0.1	0.2	0.4	1.0	2.1	4.5	10.8
1999	0.7	*	*	*	*	0.1	0.2	0.5	0.9	2.2	4.5	9.6
Non-Hispanic 4/, male												
2002	0.6	*	*	*	*	*	0.1	0.5	0.9	1.7	4.4	9.2
2001	0.6	*	*	*	*	*	0.2	0.6	1.0	1.9	4.6	10.3
2000	0.7	*	*	*	*	*	0.2	0.4	1.2	2.1	4.6	11.0
1999	0.7	*	*	*	*	*	0.2	0.6	1.0	2.4	5.0	9.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of peritoneum (K65-K66)												
Non-Hispanic 4/, female												
2002	0.9	*	*	*	*	*	0.2	0.4	0.8	1.9	5.2	11.1
2001	0.8	*	*	*	*	*	0.2	0.4	0.8	1.7	4.3	10.4
2000	0.9	*	*	*	*	*	0.2	0.4	0.9	2.1	4.3	10.7
1999	0.8	*	*	*	*	*	0.2	0.3	0.8	2.0	4.2	9.5
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	*	0.2	0.4	0.7	1.7	4.9	10.8
2001	0.7	*	*	*	*	*	0.1	0.4	0.7	1.7	4.3	10.4
2000	0.8	*	*	*	*	*	0.2	0.3	0.9	2.0	4.4	10.8
1999	0.7	*	*	*	*	*	0.2	0.4	0.8	2.1	4.4	9.4
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	*	0.5	0.8	1.6	4.3	9.5
2001	0.6	*	*	*	*	*	0.2	0.5	0.8	1.8	4.6	9.9
2000	0.7	*	*	*	*	*	0.2	0.3	1.0	1.9	4.5	10.8
1999	0.7	*	*	*	*	*	0.2	0.5	0.9	2.3	4.9	10.0
Non-Hispanic white, female												
2002	1.0	*	*	*	*	*	0.2	0.3	0.7	1.8	5.3	11.3
2001	0.8	*	*	*	*	*	*	0.3	0.6	1.6	4.2	10.6
2000	0.9	*	*	*	*	*	0.2	0.3	0.8	2.1	4.3	10.7
1999	0.8	*	*	*	*	*	0.2	0.3	0.7	1.9	4.1	9.2
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	1.1	1.5	2.9	5.5	10.6
2001	0.8	*	*	*	*	*	0.5	1.2	2.1	2.9	6.1	11.9
2000	0.8	*	*	*	*	*	0.5	1.1	2.2	3.2	6.3	11.1
1999	0.8	*	*	*	*	*	*	1.0	2.1	3.9	5.9	12.0
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	1.3	*	*	6.2	*
2001	0.7	*	*	*	*	*	*	1.1	2.5	3.4	*	*
2000	0.8	*	*	*	*	*	*	1.3	2.5	3.9	7.4	*
1999	0.7	*	*	*	*	*	*	1.3	2.6	4.2	7.2	*
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	1.0	1.4	3.3	5.1	11.3
2001	0.9	*	*	*	*	*	*	1.2	1.8	2.6	6.4	10.7
2000	0.9	*	*	*	*	*	*	1.0	2.0	2.6	5.7	10.0
1999	0.8	*	*	*	*	*	*	*	1.7	3.6	5.3	12.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of liver (K70-K76)												
All origins 3/, both sexes												
2002	12.4	2.2	0.2	0.1	0.2	1.3	8.9	23.0	29.1	38.1	41.9	35.7
2001	12.2	1.9	0.2	0.1	0.2	1.3	9.0	22.9	28.8	38.1	41.4	36.1
2000	12.0	2.0	0.2	0.1	0.2	1.4	9.2	21.8	29.3	37.9	41.2	36.7
1999	12.0	2.3	0.1	0.1	0.2	1.4	9.0	21.4	29.4	39.4	42.2	37.2
All origins 3/, male												
2002	15.8	3.1	*	*	0.2	1.6	11.6	33.9	41.2	49.5	51.7	47.3
2001	15.7	2.3	*	*	0.2	1.6	11.9	33.7	40.9	50.2	50.9	49.4
2000	15.5	2.7	*	*	0.2	1.8	12.5	32.5	41.3	49.7	51.0	52.2
1999	15.6	2.9	*	*	0.2	1.8	12.5	32.4	42.1	52.5	52.0	52.3
All origins 3/, female												
2002	9.0	1.3	*	*	0.2	1.1	6.2	12.5	17.9	28.5	35.4	30.7
2001	8.9	1.6	*	*	0.2	1.0	6.2	12.5	17.5	28.1	35.1	30.5
2000	8.7	1.3	*	*	0.2	1.0	5.9	11.4	18.3	28.3	34.7	30.4
1999	8.6	1.6	*	*	0.2	1.1	5.6	10.7	17.7	28.8	35.8	31.1
Hispanic, both sexes												
2002	10.7	2.5	*	*	*	1.6	10.2	31.8	46.1	67.9	67.6	57.1
2001	10.7	*	*	*	*	1.5	10.8	32.5	46.5	66.5	77.7	55.6
2000	10.7	*	*	*	*	1.6	11.5	32.0	47.4	69.6	77.6	67.0
1999	10.7	*	*	*	*	1.7	12.1	31.8	47.2	68.2	73.3	73.0
Hispanic, male												
2002	14.5	*	*	*	*	2.1	15.7	51.6	70.0	92.7	82.9	61.9
2001	14.7	*	*	*	*	2.0	16.3	53.3	73.0	91.7	97.0	68.6
2000	14.7	*	*	*	*	2.5	17.6	52.0	74.7	95.8	96.4	88.7
1999	14.6	*	*	*	*	2.4	18.9	52.1	74.3	94.5	86.6	99.2
Hispanic, female												
2002	6.6	*	*	*	*	0.9	4.2	12.2	24.7	48.2	57.1	54.7
2001	6.5	*	*	*	*	0.8	4.8	12.0	23.0	46.6	64.7	49.3
2000	6.5	*	*	*	*	*	5.0	12.4	23.2	48.9	65.0	56.4
1999	6.5	*	*	*	*	0.9	4.8	12.0	23.3	47.7	64.5	60.3
Non-Hispanic 4/, both sexes												
2002	12.6	2.1	*	0.1	0.2	1.3	8.7	22.0	27.5	35.8	40.5	34.8
2001	12.4	1.8	0.2	0.1	0.2	1.3	8.8	21.9	27.2	36.0	39.6	35.3
2000	12.2	2.2	0.2	0.1	0.2	1.3	8.8	20.7	27.8	35.8	39.5	35.5
1999	12.2	2.2	0.2	0.1	0.2	1.4	8.6	20.3	28.0	37.6	40.8	35.8
Non-Hispanic 4/, male												
2002	15.9	3.0	*	*	0.2	1.5	10.9	32.0	38.6	46.4	50.0	46.6
2001	15.8	2.1	*	*	0.2	1.5	11.3	31.7	38.2	47.3	48.6	48.5
2000	15.5	3.0	*	*	0.2	1.6	11.7	30.5	38.5	46.6	49.0	50.5
1999	15.7	3.1	*	*	0.2	1.6	11.6	30.4	39.6	49.8	50.5	50.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of liver (K70-K76)												
Non-Hispanic 4/, female												
2002	9.4	*	*	*	0.2	1.1	6.5	12.5	17.3	27.0	34.3	29.7
2001	9.2	1.5	*	*	0.2	1.1	6.3	12.5	17.0	26.7	33.6	29.8
2000	9.0	*	*	*	0.2	1.1	6.0	11.3	17.9	26.9	33.3	29.4
1999	8.9	1.3	*	*	0.2	1.1	5.7	10.6	17.3	27.6	34.5	30.1
Non-Hispanic white, both sexes												
2002	13.4	1.6	*	*	0.2	1.2	8.8	21.5	27.1	36.8	41.7	35.7
2001	13.1	1.3	*	*	0.1	1.2	8.6	21.2	26.6	36.6	40.6	36.1
2000	12.7	1.7	*	*	0.1	1.2	8.5	19.8	27.1	36.2	40.5	36.1
1999	12.7	1.4	*	*	0.1	1.2	8.1	19.3	26.8	37.9	41.7	36.2
Non-Hispanic white, male												
2002	17.0	2.4	*	*	*	1.4	11.2	31.3	37.9	47.4	51.4	47.9
2001	16.5	*	*	*	*	1.4	10.9	30.7	36.9	47.7	49.6	49.7
2000	16.2	2.1	*	*	*	1.4	11.4	29.4	37.7	47.1	50.3	51.3
1999	16.3	1.9	*	*	*	1.5	11.1	29.0	38.2	50.0	51.8	51.7
Non-Hispanic white, female												
2002	10.0	*	*	*	0.2	0.9	6.5	11.8	16.9	27.8	35.3	30.4
2001	9.8	*	*	*	*	1.0	6.3	11.8	16.8	27.2	34.6	30.4
2000	9.4	*	*	*	*	0.9	5.6	10.4	17.1	27.1	34.1	30.0
1999	9.2	*	*	*	*	0.9	5.2	9.7	16.2	27.7	35.2	30.0
Non-Hispanic black, both sexes												
2002	10.2	3.9	*	*	0.4	1.5	8.1	29.0	34.4	31.8	29.4	27.2
2001	10.7	3.6	*	*	0.5	1.5	10.0	29.9	36.1	35.5	31.1	22.2
2000	10.8	4.5	*	*	0.5	1.8	10.7	30.1	35.6	36.6	29.1	26.4
1999	11.1	5.1	*	*	0.4	1.8	11.1	30.2	39.6	37.7	32.1	30.4
Non-Hispanic black, male												
2002	13.0	*	*	*	*	1.6	9.4	41.4	51.4	44.8	37.2	36.1
2001	14.4	*	*	*	*	1.7	13.6	44.9	54.8	51.9	41.4	27.7
2000	13.9	6.8	*	*	*	1.9	13.8	43.9	51.3	51.2	37.6	36.4
1999	14.5	7.3	*	*	*	1.7	14.5	45.8	56.7	54.0	39.7	34.5
Non-Hispanic black, female												
2002	7.6	*	*	*	*	1.4	7.0	18.4	20.8	22.5	25.0	23.8
2001	7.4	*	*	*	*	1.3	6.8	17.0	21.0	24.0	25.3	20.1
2000	7.9	*	*	*	*	1.8	7.9	18.3	22.9	26.3	24.3	22.7
1999	8.1	*	*	*	*	1.8	8.0	16.8	25.8	26.3	27.8	28.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All origins 3/, both sexes												
2002	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
All origins 3/, male												
2002	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
All origins 3/, female												
2002	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5
Hispanic, both sexes												
2002	4.3	*	*	*	*	0.7	5.5	16.1	19.9	20.8	10.5	*
2001	4.7	*	*	*	*	0.9	6.3	17.2	21.4	21.1	17.1	*
2000	4.8	*	*	*	*	1.0	6.9	16.8	23.4	22.9	15.4	*
1999	4.6	*	*	*	*	0.8	6.9	16.8	21.7	20.6	14.4	*
Hispanic, male												
2002	7.1	*	*	*	*	1.2	9.0	27.0	36.3	40.3	18.1	*
2001	7.7	*	*	*	*	1.5	10.0	29.4	38.9	41.9	30.9	*
2000	7.9	*	*	*	*	1.8	10.6	29.3	42.8	46.6	33.9	*
1999	7.4	*	*	*	*	1.2	10.8	29.1	40.0	39.5	27.1	*
Hispanic, female												
2002	1.4	*	*	*	*	*	1.8	5.1	5.1	5.3	*	*
2001	1.5	*	*	*	*	*	2.2	5.1	5.9	4.7	7.9	*
2000	1.4	*	*	*	*	*	2.9	4.5	6.2	4.3	*	*
1999	1.5	*	*	*	*	*	2.8	4.9	5.6	5.7	*	*
Non-Hispanic 4/, both sexes												
2002	4.2	*	*	*	*	0.6	4.2	9.7	10.8	9.3	5.5	1.9
2001	4.2	*	*	*	*	0.6	4.6	10.1	10.7	9.3	5.2	2.2
2000	4.2	*	*	*	*	0.6	4.7	9.7	11.3	9.5	5.7	2.2
1999	4.2	*	*	*	*	0.6	4.6	9.8	11.3	9.9	6.1	2.5
Non-Hispanic 4/, male												
2002	6.0	*	*	*	*	0.7	5.3	14.4	16.7	14.8	9.2	4.3
2001	6.2	*	*	*	*	0.8	5.9	14.8	16.7	14.9	9.2	5.1
2000	6.2	*	*	*	*	0.8	6.3	14.7	17.1	15.4	10.5	5.0
1999	6.4	*	*	*	*	0.7	6.2	15.1	17.7	16.7	10.5	5.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Non-Hispanic 4/, female												
2002	2.4	*	*	*	*	0.5	3.1	5.2	5.4	4.8	3.0	0.9
2001	2.4	*	*	*	*	0.5	3.2	5.5	5.0	4.7	2.5	1.0
2000	2.3	*	*	*	*	0.4	3.1	4.8	5.9	4.5	2.5	1.0
1999	2.2	*	*	*	*	0.5	2.9	4.7	5.3	4.2	3.3	1.4
Non-Hispanic white, both sexes												
2002	4.4	*	*	*	*	0.5	4.3	9.7	10.8	9.5	5.5	1.9
2001	4.4	*	*	*	*	0.6	4.5	9.9	10.5	9.3	5.3	2.1
2000	4.3	*	*	*	*	0.6	4.5	9.4	11.1	9.5	5.6	2.2
1999	4.3	*	*	*	*	0.6	4.4	9.5	10.8	9.7	6.0	2.4
Non-Hispanic white, male												
2002	6.4	*	*	*	*	0.7	5.4	14.4	16.7	14.9	9.4	4.1
2001	6.4	*	*	*	*	0.8	5.8	14.7	16.4	14.6	9.3	4.8
2000	6.5	*	*	*	*	0.8	6.1	14.5	17.0	15.4	10.6	5.1
1999	6.5	*	*	*	*	0.7	6.0	14.7	17.1	16.3	10.3	5.1
Non-Hispanic white, female												
2002	2.4	*	*	*	*	0.4	3.2	5.1	5.2	4.9	3.0	1.0
2001	2.4	*	*	*	*	0.4	3.2	5.3	5.0	4.8	2.5	1.0
2000	2.2	*	*	*	*	0.4	3.0	4.4	5.5	4.6	2.4	1.0
1999	2.2	*	*	*	*	0.5	2.8	4.3	4.9	4.2	3.2	1.3
Non-Hispanic black, both sexes												
2002	3.2	*	*	*	*	0.5	3.0	10.2	12.3	8.7	5.1	*
2001	3.7	*	*	*	*	0.5	4.4	11.7	13.1	10.8	4.6	*
2000	3.9	*	*	*	*	0.7	5.1	12.3	13.5	10.4	5.6	*
1999	4.2	*	*	*	*	0.5	5.1	12.9	15.4	11.9	7.2	*
Non-Hispanic black, male												
2002	4.6	*	*	*	*	*	3.9	15.1	18.6	15.1	8.8	*
2001	5.5	*	*	*	*	*	6.1	17.4	21.7	20.0	7.8	*
2000	5.5	*	*	*	*	*	7.1	17.9	19.6	18.4	10.2	*
1999	6.1	*	*	*	*	*	6.9	19.7	24.6	21.9	12.8	*
Non-Hispanic black, female												
2002	2.1	*	*	*	*	*	2.2	6.1	7.2	4.2	*	*
2001	2.2	*	*	*	*	*	2.8	6.8	6.2	4.3	*	*
2000	2.5	*	*	*	*	*	3.4	7.5	8.6	4.8	*	*
1999	2.4	*	*	*	*	*	3.5	7.0	8.1	4.8	4.0	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic hepatitis, not elsewhere classified (K73)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.0	0.8
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	0.8	0.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	0.8	0.8
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	0.6
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	0.2	*	0.4	0.6	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.3	0.4	*
2000	0.1	*	*	*	*	*	*	0.1	*	*	*	*
1999	0.1	*	*	*	*	*	*	0.2	0.2	0.3	0.5	*
All origins 3/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.3	0.9
2001	0.2	*	*	*	*	*	*	0.2	0.5	0.9	1.1	0.7
2000	0.2	*	*	*	*	*	*	0.2	0.5	1.0	1.1	0.9
1999	0.2	*	*	*	*	*	*	0.2	0.5	0.8	1.1	0.7
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	2.0	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	3.3	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.9	0.7
2001	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	0.8	0.8
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	0.8	0.8
1999	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	0.9	0.6
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	0.2	*	0.4	0.6	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.3	0.4	*
2000	0.1	*	*	*	*	*	*	0.1	*	*	*	*
1999	0.1	*	*	*	*	*	*	0.2	*	0.3	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic hepatitis, not elsewhere classified (K73)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.2	0.8
2001	0.2	*	*	*	*	*	*	0.2	0.4	0.9	1.0	0.7
2000	0.2	*	*	*	*	*	*	0.1	0.4	0.9	1.1	0.9
1999	0.2	*	*	*	*	*	*	0.2	0.5	0.7	1.1	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.0	0.8
2001	0.2	*	*	*	*	*	*	0.2	0.3	0.6	0.8	0.8
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.6	0.8	0.8
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	0.8	0.6
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.6	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.4	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	0.2	*	*	*	*
Non-Hispanic white, female												
2002	0.3	*	*	*	*	*	*	0.2	0.4	0.9	1.2	0.8
2001	0.2	*	*	*	*	*	*	0.2	0.5	0.8	1.0	0.8
2000	0.2	*	*	*	*	*	*	*	0.5	0.9	1.1	0.9
1999	0.2	*	*	*	*	*	*	0.1	0.4	0.7	1.1	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)												
All origins 3/, both sexes												
2002	5.1	*	*	*	*	0.3	2.5	7.5	11.0	18.6	24.7	18.5
2001	5.0	*	*	*	*	0.3	2.6	7.6	10.9	19.2	23.7	19.1
2000	5.0	*	*	*	0.1	0.3	2.5	7.2	11.3	18.9	24.1	20.0
1999	5.0	*	*	*	*	0.3	2.4	6.8	11.3	19.6	24.6	19.8
All origins 3/, male												
2002	6.0	*	*	*	*	0.3	3.4	11.1	14.7	21.7	28.1	23.2
2001	6.0	*	*	*	*	0.4	3.3	11.3	14.6	23.2	26.2	25.7
2000	5.9	*	*	*	*	0.4	3.4	10.7	15.3	22.5	26.5	28.4
1999	5.9	*	*	*	*	0.4	3.5	10.4	15.2	23.3	27.9	26.6
All origins 3/, female												
2002	4.2	*	*	*	*	0.2	1.7	4.1	7.6	16.0	22.4	16.5
2001	4.2	*	*	*	*	0.2	1.8	4.1	7.5	15.9	22.1	16.4
2000	4.1	*	*	*	*	0.2	1.5	3.8	7.6	15.9	22.5	16.6
1999	4.1	*	*	*	*	0.2	1.3	3.4	7.8	16.5	22.5	17.0
Hispanic, both sexes												
2002	4.3	*	*	*	*	0.3	3.1	10.6	17.8	33.3	40.7	37.1
2001	4.1	*	*	*	*	*	3.4	10.5	16.9	32.1	42.0	30.0
2000	4.1	*	*	*	*	0.4	2.8	10.3	17.4	33.8	46.6	41.8
1999	4.1	*	*	*	*	0.4	3.5	9.5	16.7	34.5	44.2	38.6
Hispanic, male												
2002	5.0	*	*	*	*	*	4.5	16.8	23.6	37.1	46.0	*
2001	5.0	*	*	*	*	*	5.0	16.9	24.7	34.7	45.2	*
2000	4.7	*	*	*	*	*	4.3	16.2	23.0	36.5	46.2	46.4
1999	4.8	*	*	*	*	0.6	5.7	15.0	22.4	39.3	43.3	45.3
Hispanic, female												
2002	3.4	*	*	*	*	*	1.6	4.4	12.5	30.3	37.1	39.8
2001	3.2	*	*	*	*	*	1.7	4.3	10.0	30.1	39.9	32.6
2000	3.5	*	*	*	*	*	1.1	4.5	12.4	31.7	46.9	39.6
1999	3.3	*	*	*	*	*	1.2	4.1	11.6	30.8	44.8	35.4
Non-Hispanic 4/, both sexes												
2002	5.2	*	*	*	*	0.3	2.5	7.2	10.4	17.5	23.8	17.7
2001	5.2	*	*	*	*	0.3	2.5	7.3	10.4	18.3	22.8	18.6
2000	5.1	*	*	*	0.1	0.3	2.4	6.9	10.8	17.8	23.1	19.2
1999	5.1	*	*	*	*	0.3	2.3	6.6	10.9	18.6	23.7	19.0
Non-Hispanic 4/, male												
2002	6.1	*	*	*	*	0.3	3.2	10.4	13.9	20.5	27.2	22.8
2001	6.1	*	*	*	*	0.4	3.1	10.7	13.7	22.4	25.3	25.6
2000	6.0	*	*	*	*	0.3	3.3	10.2	14.6	21.5	25.7	27.5
1999	6.0	*	*	*	*	0.4	3.2	9.9	14.6	22.3	27.2	25.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)												
Non-Hispanic 4/, female												
2002	4.3	*	*	*	*	0.2	1.7	4.0	7.2	15.0	21.6	15.6
2001	4.3	*	*	*	*	0.2	1.8	4.0	7.3	14.8	21.2	15.7
2000	4.2	*	*	*	*	0.3	1.6	3.7	7.2	14.8	21.4	15.8
1999	4.2	*	*	*	*	0.2	1.4	3.3	7.5	15.6	21.5	16.3
Non-Hispanic white, both sexes												
2002	5.7	*	*	*	*	0.3	2.5	6.9	10.4	18.3	25.0	18.5
2001	5.6	*	*	*	*	0.3	2.4	6.9	10.2	19.0	23.9	19.6
2000	5.5	*	*	*	*	0.2	2.4	6.6	10.7	18.3	24.2	19.8
1999	5.5	*	*	*	*	0.3	2.2	6.2	10.7	19.3	24.8	19.6
Non-Hispanic white, male												
2002	6.6	*	*	*	*	0.3	3.4	10.2	13.6	21.3	28.7	24.0
2001	6.5	*	*	*	*	0.3	3.0	10.0	13.2	23.2	26.5	27.2
2000	6.5	*	*	*	*	0.3	3.3	9.8	14.5	22.1	26.8	28.4
1999	6.5	*	*	*	*	0.4	3.1	9.3	14.2	23.1	28.5	27.1
Non-Hispanic white, female												
2002	4.8	*	*	*	*	0.2	1.7	3.7	7.3	15.8	22.5	16.2
2001	4.7	*	*	*	*	0.2	1.8	3.9	7.4	15.5	22.2	16.4
2000	4.6	*	*	*	*	0.2	1.5	3.5	7.2	15.1	22.4	16.3
1999	4.6	*	*	*	*	0.2	1.3	3.1	7.4	16.1	22.4	16.7
Non-Hispanic black, both sexes												
2002	3.6	*	*	*	*	*	2.4	10.6	12.7	12.5	12.8	9.1
2001	3.8	*	*	*	*	*	3.0	11.0	13.0	13.9	12.7	7.5
2000	3.7	*	*	*	*	*	2.8	10.1	12.5	15.4	12.2	11.8
1999	3.7	*	*	*	*	0.4	3.0	9.8	13.9	14.2	13.8	11.6
Non-Hispanic black, male												
2002	4.6	*	*	*	*	*	2.9	15.2	19.6	17.1	12.9	*
2001	5.2	*	*	*	*	*	4.2	17.5	20.2	19.4	14.7	*
2000	4.7	*	*	*	*	*	3.5	15.3	18.8	19.6	14.8	*
1999	4.8	*	*	*	*	*	4.4	15.8	19.3	18.8	16.6	*
Non-Hispanic black, female												
2002	2.7	*	*	*	*	*	1.9	6.7	7.2	9.2	12.8	8.8
2001	2.5	*	*	*	*	*	2.0	5.4	7.2	10.0	11.6	*
2000	2.7	*	*	*	*	*	2.2	5.6	7.4	12.4	10.8	10.5
1999	2.7	*	*	*	*	*	1.7	4.8	9.5	11.0	12.3	12.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)												
All origins 3/, both sexes												
2002	2.9	2.1	0.2	0.1	0.1	0.4	1.9	5.0	6.2	8.6	10.5	14.3
2001	2.7	1.9	0.2	0.1	0.1	0.4	1.6	4.4	6.1	8.1	11.1	13.8
2000	2.6	1.9	0.2	0.1	0.1	0.4	1.7	4.1	5.5	8.1	10.2	13.6
1999	2.6	2.1	0.1	0.1	0.1	0.4	1.7	4.0	5.7	8.8	10.3	14.0
All origins 3/, male												
2002	3.5	2.8	*	*	0.1	0.5	2.3	7.1	8.1	11.0	13.3	18.9
2001	3.2	2.2	*	*	0.1	0.3	2.1	6.2	7.8	10.0	14.1	17.4
2000	3.0	2.5	*	*	0.1	0.4	2.2	5.6	6.9	9.7	12.6	17.8
1999	3.1	2.7	*	*	0.1	0.5	2.1	5.5	7.5	10.8	12.5	19.2
All origins 3/, female												
2002	2.3	1.3	*	*	0.1	0.4	1.5	3.0	4.4	6.6	8.6	12.3
2001	2.3	1.5	*	*	0.1	0.4	1.2	2.7	4.4	6.6	9.2	12.3
2000	2.2	1.3	*	*	*	0.3	1.2	2.7	4.3	6.8	8.6	11.9
1999	2.2	1.5	*	*	0.2	0.3	1.3	2.5	4.0	7.2	8.8	11.9
Hispanic, both sexes												
2002	1.9	2.4	*	*	*	0.5	1.4	5.0	7.7	11.8	14.0	11.6
2001	1.8	*	*	*	*	*	1.0	4.4	7.5	12.1	17.1	18.8
2000	1.7	*	*	*	*	*	1.7	4.7	6.0	11.2	13.8	18.6
1999	1.9	*	*	*	*	0.5	1.6	5.2	8.2	12.0	14.0	21.7
Hispanic, male												
2002	2.3	*	*	*	*	0.6	2.1	7.5	9.9	14.9	18.5	*
2001	2.1	*	*	*	*	*	1.3	6.7	9.1	14.6	20.6	*
2000	2.0	*	*	*	*	*	2.6	6.4	8.6	12.5	16.2	*
1999	2.4	*	*	*	*	0.6	2.3	7.8	11.7	15.4	15.6	*
Hispanic, female												
2002	1.4	*	*	*	*	*	*	2.5	5.8	9.3	11.0	*
2001	1.5	*	*	*	*	*	0.8	2.2	6.0	10.1	14.8	*
2000	1.4	*	*	*	*	*	0.8	3.1	3.6	10.3	12.2	*
1999	1.5	*	*	*	*	*	*	2.7	5.2	9.3	12.8	*
Non-Hispanic 4/, both sexes												
2002	3.1	1.9	*	0.1	0.1	0.4	2.0	5.0	6.0	8.3	10.3	14.4
2001	2.9	1.7	0.2	0.1	0.1	0.4	1.7	4.4	5.9	7.8	10.8	13.6
2000	2.7	2.1	0.2	*	0.1	0.4	1.7	4.1	5.5	7.9	10.0	13.4
1999	2.7	2.1	*	*	0.1	0.4	1.7	3.8	5.5	8.6	10.1	13.7
Non-Hispanic 4/, male												
2002	3.7	2.7	*	*	*	0.4	2.4	7.0	7.9	10.7	13.0	18.9
2001	3.4	2.1	*	*	0.1	0.4	2.2	6.1	7.6	9.6	13.8	17.0
2000	3.1	2.8	*	*	0.1	0.4	2.1	5.5	6.7	9.5	12.5	17.4
1999	3.2	2.8	*	*	*	0.5	2.1	5.2	7.2	10.5	12.3	18.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)												
Non-Hispanic 4/, female												
2002	2.5	*	*	*	0.1	0.4	1.6	3.1	4.2	6.4	8.5	12.4
2001	2.4	1.4	*	*	0.1	0.4	1.2	2.7	4.3	6.3	8.9	12.2
2000	2.3	*	*	*	*	0.4	1.3	2.6	4.3	6.6	8.4	11.8
1999	2.3	*	*	*	0.1	0.3	1.3	2.5	4.0	7.0	8.6	11.7
Non-Hispanic white, both sexes												
2002	3.1	1.5	*	*	0.1	0.4	1.9	4.7	5.6	8.3	10.2	14.5
2001	2.9	1.2	*	*	0.1	0.3	1.6	4.1	5.5	7.7	10.7	13.6
2000	2.7	1.7	*	*	0.1	0.3	1.6	3.6	5.0	7.8	9.9	13.3
1999	2.7	1.3	*	*	0.1	0.3	1.5	3.4	5.0	8.3	10.1	13.6
Non-Hispanic white, male												
2002	3.8	2.1	*	*	*	0.4	2.3	6.6	7.4	10.7	12.7	19.2
2001	3.5	*	*	*	*	0.3	2.0	5.8	7.2	9.6	13.3	16.9
2000	3.1	2.1	*	*	*	0.3	2.0	4.9	6.1	9.3	12.5	17.2
1999	3.2	*	*	*	*	0.4	1.9	4.7	6.6	10.3	12.5	19.0
Non-Hispanic white, female												
2002	2.5	*	*	*	*	0.3	1.5	2.8	4.0	6.2	8.5	12.4
2001	2.4	*	*	*	*	0.4	1.2	2.5	4.0	6.1	8.9	12.2
2000	2.3	*	*	*	*	0.3	1.1	2.3	4.0	6.5	8.2	11.8
1999	2.3	*	*	*	*	0.2	1.1	2.1	3.5	6.7	8.5	11.4
Non-Hispanic black, both sexes												
2002	3.1	3.7	*	*	*	0.6	2.5	7.8	9.1	10.0	11.1	15.1
2001	3.0	3.6	*	*	*	0.6	2.5	6.9	9.8	10.2	12.8	11.5
2000	3.1	4.2	*	*	*	0.8	2.7	7.5	9.4	10.1	11.0	12.4
1999	3.1	5.1	*	*	*	0.7	2.8	7.2	9.8	10.9	10.4	13.6
Non-Hispanic black, male												
2002	3.7	*	*	*	*	*	2.5	10.8	13.2	12.4	15.2	*
2001	3.7	*	*	*	*	*	3.2	9.8	12.8	12.2	18.9	*
2000	3.7	*	*	*	*	0.8	3.1	10.5	12.8	13.0	12.6	*
1999	3.5	7.3	*	*	*	*	3.2	10.0	12.9	12.9	9.7	*
Non-Hispanic black, female												
2002	2.6	*	*	*	*	*	2.5	5.3	5.8	8.3	8.8	13.4
2001	2.5	*	*	*	*	*	1.8	4.5	7.3	8.9	9.4	10.3
2000	2.5	*	*	*	*	*	2.3	4.8	6.7	8.0	10.1	10.9
1999	2.7	*	*	*	*	0.8	2.6	4.8	7.4	9.4	10.9	13.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All origins 3/, both sexes												
2002	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9
All origins 3/, male												
2002	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7
All origins 3/, female												
2002	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4
Hispanic, both sexes												
2002	0.5	*	*	*	*	*	*	*	1.3	3.6	9.2	24.4
2001	0.5	*	*	*	*	*	*	*	1.2	4.5	8.5	31.3
2000	0.5	*	*	*	*	*	*	*	*	3.8	10.1	29.2
1999	0.5	*	*	*	*	*	*	*	1.2	3.3	11.9	32.3
Hispanic, male												
2002	0.5	*	*	*	*	*	*	*	*	4.0	9.4	*
2001	0.5	*	*	*	*	*	*	*	*	5.0	11.2	*
2000	0.5	*	*	*	*	*	*	*	*	5.3	15.7	*
1999	0.5	*	*	*	*	*	*	*	*	*	15.1	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	3.2	9.0	23.2
2001	0.6	*	*	*	*	*	*	*	*	4.1	6.6	30.7
2000	0.5	*	*	*	*	*	*	*	*	*	*	31.7
1999	0.6	*	*	*	*	*	*	*	*	*	9.7	32.2
Non-Hispanic 4/, both sexes												
2002	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.7	22.8
2001	1.1	*	*	*	*	*	0.1	0.3	0.9	2.8	7.5	23.4
2000	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.4	24.6
1999	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.6	23.6
Non-Hispanic 4/, male												
2002	1.0	*	*	*	*	*	0.1	0.4	0.9	3.0	9.8	23.9
2001	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.2	26.6
2000	1.0	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	27.7
1999	1.0	*	*	*	*	*	0.1	0.3	1.0	3.0	9.0	27.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Non-Hispanic 4/, female												
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	2.4	6.3	22.4
2001	1.2	*	*	*	*	*	*	0.3	0.8	2.5	6.3	22.1
2000	1.2	*	*	*	*	*	0.1	0.3	0.6	2.0	6.2	23.4
1999	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.6	22.0
Non-Hispanic white, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.3	0.7	2.6	7.5	23.4
2001	1.2	*	*	*	*	*	0.1	0.3	0.8	2.6	7.3	23.6
2000	1.2	*	*	*	*	*	0.1	0.2	0.7	2.2	7.3	25.3
1999	1.2	*	*	*	*	*	0.1	0.3	0.7	2.4	7.6	24.0
Non-Hispanic white, male												
2002	1.1	*	*	*	*	*	*	0.3	0.9	3.0	9.8	24.5
2001	1.1	*	*	*	*	*	*	0.4	1.0	2.9	9.2	26.4
2000	1.0	*	*	*	*	*	*	0.2	0.8	2.6	9.3	28.9
1999	1.1	*	*	*	*	*	*	0.3	0.9	2.9	9.3	27.8
Non-Hispanic white, female												
2002	1.3	*	*	*	*	*	*	0.2	0.6	2.3	6.1	22.9
2001	1.3	*	*	*	*	*	*	0.2	0.7	2.2	6.1	22.4
2000	1.3	*	*	*	*	*	*	0.2	0.6	1.8	6.0	23.9
1999	1.3	*	*	*	*	*	*	0.3	0.6	1.9	6.5	22.5
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.6	15.1
2001	0.9	*	*	*	*	*	*	0.7	1.7	4.8	8.3	20.0
2000	0.8	*	*	*	*	*	*	0.7	1.2	3.8	8.3	18.1
1999	0.7	*	*	*	*	*	0.4	0.5	1.5	3.7	7.3	15.5
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	*	*	*	10.8	*
2001	0.8	*	*	*	*	*	*	*	2.2	5.0	6.3	27.7
2000	0.6	*	*	*	*	*	*	*	*	3.4	8.0	*
1999	0.6	*	*	*	*	*	*	*	*	3.8	7.2	*
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	*	*	3.8	9.0	16.3
2001	1.0	*	*	*	*	*	*	*	*	4.6	9.4	17.1
2000	1.0	*	*	*	*	*	*	*	*	4.0	8.5	19.6
1999	0.8	*	*	*	*	*	*	*	*	3.6	7.4	15.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)												
All origins 3/, both sexes												
2002	1.7	*	*	*	0.1	0.3	0.8	1.4	2.5	4.8	9.7	21.7
2001	1.7	*	*	*	0.1	0.4	0.9	1.6	2.5	5.1	9.4	22.9
2000	1.7	*	*	*	0.1	0.3	0.8	1.4	2.5	4.6	9.4	22.0
1999	1.6	*	*	*	0.1	0.4	0.8	1.4	2.4	4.5	8.5	22.2
All origins 3/, male												
2002	1.9	*	*	*	0.2	0.5	1.1	2.0	3.4	6.4	11.9	21.2
2001	1.8	*	*	*	0.1	0.5	1.1	2.0	3.3	6.5	10.2	22.4
2000	1.8	*	*	*	0.1	0.5	1.2	1.9	3.0	5.9	11.4	22.9
1999	1.7	*	*	*	0.1	0.5	1.1	1.9	3.1	5.6	10.1	23.1
All origins 3/, female												
2002	1.5	*	*	*	*	0.2	0.5	0.8	1.6	3.5	8.3	21.9
2001	1.7	*	*	*	*	0.3	0.6	1.1	1.8	3.9	8.8	23.2
2000	1.6	*	*	*	*	0.2	0.5	1.0	2.0	3.5	8.0	21.6
1999	1.5	*	*	*	0.1	0.2	0.5	0.9	1.7	3.6	7.4	21.8
Hispanic, both sexes												
2002	0.8	*	*	*	*	0.3	0.6	1.2	1.5	4.7	8.5	22.7
2001	0.7	*	*	*	*	0.3	0.6	0.9	1.9	4.8	10.8	18.8
2000	0.8	*	*	*	*	*	0.8	1.1	1.9	4.0	10.7	27.2
1999	0.8	*	*	*	*	0.4	0.7	1.0	2.3	5.1	8.3	30.9
Hispanic, male												
2002	0.9	*	*	*	*	0.5	0.7	1.4	2.6	5.9	13.1	*
2001	0.7	*	*	*	*	*	0.8	*	2.4	5.4	9.4	*
2000	0.9	*	*	*	*	*	1.3	1.5	*	4.6	14.3	*
1999	0.9	*	*	*	*	*	0.8	1.7	*	6.8	10.9	*
Hispanic, female												
2002	0.6	*	*	*	*	*	*	*	*	3.6	*	25.7
2001	0.8	*	*	*	*	*	*	*	*	4.3	11.8	24.2
2000	0.7	*	*	*	*	*	*	*	*	3.5	8.3	23.7
1999	0.7	*	*	*	*	*	*	*	2.3	3.8	*	30.2
Non-Hispanic 4/, both sexes												
2002	1.8	*	*	*	0.1	0.3	0.8	1.4	2.6	4.8	9.8	21.6
2001	1.9	*	*	*	0.1	0.4	0.9	1.6	2.6	5.1	9.3	23.0
2000	1.8	*	*	*	0.1	0.4	0.8	1.4	2.5	4.6	9.3	21.8
1999	1.7	*	*	*	0.1	0.3	0.8	1.5	2.4	4.4	8.4	21.8
Non-Hispanic 4/, male												
2002	2.0	*	*	*	0.1	0.5	1.1	2.1	3.5	6.4	11.8	21.3
2001	2.0	*	*	*	*	0.5	1.2	2.1	3.3	6.5	10.2	22.9
2000	1.9	*	*	*	*	0.5	1.2	1.9	3.1	5.9	11.3	22.4
1999	1.8	*	*	*	*	0.5	1.1	1.9	3.2	5.5	10.0	22.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)												
Non-Hispanic 4/, female												
2002	1.7	*	*	*	*	0.2	0.6	0.8	1.7	3.5	8.4	21.7
2001	1.8	*	*	*	*	0.3	0.6	1.1	1.9	3.9	8.7	23.1
2000	1.7	*	*	*	*	0.2	0.5	1.0	2.0	3.5	8.0	21.5
1999	1.6	*	*	*	*	0.2	0.5	1.0	1.7	3.6	7.4	21.4
Non-Hispanic white, both sexes												
2002	1.9	*	*	*	0.1	0.3	0.8	1.2	2.4	4.8	9.9	21.8
2001	2.0	*	*	*	*	0.3	0.8	1.4	2.4	5.0	9.3	23.1
2000	1.8	*	*	*	*	0.3	0.8	1.3	2.3	4.3	9.2	21.7
1999	1.7	*	*	*	0.1	0.3	0.7	1.2	2.2	4.4	8.3	21.5
Non-Hispanic white, male												
2002	2.1	*	*	*	*	0.4	1.0	1.8	3.2	6.3	12.0	21.2
2001	2.0	*	*	*	*	0.4	1.1	1.8	3.1	6.5	10.3	22.8
2000	1.9	*	*	*	*	0.4	1.0	1.7	2.8	5.6	11.2	21.5
1999	1.8	*	*	*	*	0.4	1.0	1.6	2.8	5.6	9.9	21.4
Non-Hispanic white, female												
2002	1.8	*	*	*	*	0.2	0.5	0.7	1.6	3.5	8.5	22.0
2001	1.9	*	*	*	*	0.2	0.5	1.0	1.8	3.7	8.7	23.2
2000	1.8	*	*	*	*	0.2	0.5	0.9	1.8	3.3	7.9	21.7
1999	1.7	*	*	*	*	0.2	0.5	0.9	1.5	3.4	7.2	21.5
Non-Hispanic black, both sexes												
2002	1.8	*	*	*	*	0.6	1.5	3.1	4.6	6.1	8.8	21.8
2001	2.0	*	*	*	*	0.8	2.0	3.8	4.2	6.7	9.5	22.2
2000	1.9	*	*	*	*	0.8	1.5	3.0	4.6	7.4	10.1	23.8
1999	1.8	*	*	*	*	0.7	1.5	3.4	4.8	5.5	10.0	23.3
Non-Hispanic black, male												
2002	2.2	*	*	*	*	0.9	2.0	4.7	6.6	8.6	10.0	26.3
2001	2.2	*	*	*	*	1.1	2.4	5.2	6.0	8.8	10.2	*
2000	2.2	*	*	*	*	1.1	2.2	4.2	6.1	10.3	12.0	32.9
1999	2.2	*	*	*	*	1.2	2.2	5.3	6.6	6.2	10.9	33.3
Non-Hispanic black, female												
2002	1.4	*	*	*	*	*	1.0	1.7	3.0	4.3	8.1	20.1
2001	1.7	*	*	*	*	*	1.6	2.5	2.7	5.3	9.1	22.7
2000	1.6	*	*	*	*	*	0.8	2.0	3.4	5.3	9.0	20.5
1999	1.5	*	*	*	*	*	0.9	1.7	3.3	5.0	9.5	19.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)												
All origins 3/, both sexes												
2002	1.2	*	*	*	0.1	0.3	0.7	1.2	2.0	3.4	6.5	11.9
2001	1.3	*	*	*	0.1	0.3	0.8	1.3	2.1	3.8	6.4	13.2
2000	1.2	*	*	*	0.1	0.3	0.8	1.2	2.0	3.4	6.3	12.1
1999	1.2	*	*	*	0.1	0.3	0.7	1.2	1.9	3.5	5.8	12.8
All origins 3/, male												
2002	1.4	*	*	*	0.1	0.4	1.0	1.8	2.8	4.6	7.8	11.2
2001	1.4	*	*	*	*	0.4	1.0	1.8	2.8	4.9	6.8	12.9
2000	1.3	*	*	*	0.1	0.4	1.1	1.6	2.5	4.5	7.4	11.9
1999	1.3	*	*	*	*	0.5	1.0	1.7	2.5	4.4	6.8	12.7
All origins 3/, female												
2002	1.1	*	*	*	*	0.1	0.5	0.7	1.3	2.4	5.7	12.2
2001	1.2	*	*	*	*	0.2	0.5	0.9	1.5	3.0	6.1	13.4
2000	1.1	*	*	*	*	0.2	0.4	0.8	1.5	2.5	5.6	12.2
1999	1.1	*	*	*	0.1	0.2	0.5	0.8	1.3	2.7	5.2	12.8
Hispanic, both sexes												
2002	0.6	*	*	*	*	0.3	0.5	1.0	1.2	3.6	4.7	13.3
2001	0.6	*	*	*	*	0.3	0.6	0.6	1.4	4.0	6.7	*
2000	0.6	*	*	*	*	*	0.7	0.9	1.3	2.3	6.5	16.6
1999	0.6	*	*	*	*	0.4	0.6	0.9	1.6	4.0	5.2	14.0
Hispanic, male												
2002	0.7	*	*	*	*	0.5	0.7	1.3	*	4.8	8.2	*
2001	0.6	*	*	*	*	*	0.8	*	*	4.6	*	*
2000	0.7	*	*	*	*	*	1.3	1.3	*	*	*	*
1999	0.7	*	*	*	*	*	0.8	1.5	*	5.3	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	3.5	8.5	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	0.1	0.3	0.7	1.2	2.1	3.4	6.6	11.8
2001	1.4	*	*	*	0.1	0.3	0.8	1.4	2.2	3.8	6.3	13.4
2000	1.3	*	*	*	0.1	0.3	0.8	1.2	2.0	3.4	6.3	12.0
1999	1.3	*	*	*	0.1	0.3	0.7	1.3	1.9	3.4	5.8	12.7
Non-Hispanic 4/, male												
2002	1.5	*	*	*	0.1	0.4	1.0	1.9	2.9	4.6	7.7	11.1
2001	1.5	*	*	*	*	0.4	1.1	1.8	2.8	4.9	6.9	13.3
2000	1.4	*	*	*	*	0.4	1.1	1.6	2.5	4.6	7.3	11.4
1999	1.4	*	*	*	*	0.5	1.0	1.7	2.5	4.4	6.8	12.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)												
Non-Hispanic 4/, female												
2002	1.1	*	*	*	*	0.2	0.5	0.7	1.4	2.4	5.9	12.1
2001	1.3	*	*	*	*	0.2	0.6	0.9	1.5	2.9	6.0	13.4
2000	1.2	*	*	*	*	0.2	0.4	0.9	1.6	2.5	5.6	12.2
1999	1.1	*	*	*	*	0.2	0.5	0.9	1.3	2.7	5.2	12.8
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	0.2	0.7	1.1	2.0	3.4	6.8	12.0
2001	1.4	*	*	*	*	0.2	0.7	1.1	2.0	3.8	6.4	13.4
2000	1.3	*	*	*	*	0.3	0.7	1.1	1.8	3.3	6.3	12.0
1999	1.3	*	*	*	*	0.3	0.7	1.0	1.7	3.4	5.7	12.7
Non-Hispanic white, male												
2002	1.5	*	*	*	*	0.3	0.9	1.6	2.6	4.6	7.8	10.8
2001	1.5	*	*	*	*	0.3	0.9	1.5	2.6	5.0	6.9	13.4
2000	1.4	*	*	*	*	0.4	1.0	1.4	2.3	4.3	7.2	10.9
1999	1.3	*	*	*	*	0.4	0.9	1.3	2.3	4.3	6.8	11.7
Non-Hispanic white, female												
2002	1.2	*	*	*	*	*	0.5	0.5	1.3	2.4	6.1	12.4
2001	1.3	*	*	*	*	*	0.5	0.8	1.5	2.8	6.1	13.4
2000	1.2	*	*	*	*	0.2	0.4	0.8	1.4	2.4	5.6	12.5
1999	1.2	*	*	*	*	0.2	0.4	0.8	1.1	2.6	5.0	13.1
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	*	0.5	1.3	2.7	3.9	3.8	5.3	11.5
2001	1.5	*	*	*	*	0.7	1.7	3.4	3.4	4.7	6.5	13.4
2000	1.5	*	*	*	*	0.7	1.4	2.6	4.0	5.5	6.2	12.1
1999	1.5	*	*	*	*	0.6	1.4	3.1	3.8	4.4	7.2	12.9
Non-Hispanic black, male												
2002	1.7	*	*	*	*	*	1.8	4.1	5.7	5.3	6.7	*
2001	1.8	*	*	*	*	0.9	2.2	4.6	5.3	5.8	7.5	*
2000	1.8	*	*	*	*	1.0	2.2	3.5	5.3	8.2	8.0	*
1999	1.9	*	*	*	*	1.1	2.2	4.9	5.3	5.1	7.8	*
Non-Hispanic black, female												
2002	1.0	*	*	*	*	*	0.8	1.5	2.4	2.7	4.5	9.6
2001	1.3	*	*	*	*	*	1.3	2.3	2.0	4.0	5.9	14.1
2000	1.1	*	*	*	*	*	0.7	1.8	2.9	3.6	5.2	10.0
1999	1.1	*	*	*	*	*	0.8	1.6	2.6	3.8	6.8	10.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.2	9.8
2001	0.5	*	*	*	*	0.1	0.1	0.3	0.4	1.2	3.0	9.7
2000	0.5	*	*	*	*	*	0.1	0.2	0.5	1.2	3.1	9.9
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.6	9.4
All origins 3/, male												
2002	0.5	*	*	*	*	*	0.1	0.2	0.6	1.8	4.2	10.0
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.6	3.4	9.5
2000	0.4	*	*	*	*	*	*	0.3	0.5	1.3	4.0	11.0
1999	0.4	*	*	*	*	*	*	0.3	0.6	1.2	3.2	10.5
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.1	0.3	1.1	2.5	9.8
2001	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.7	9.8
2000	0.5	*	*	*	*	*	0.1	0.1	0.4	1.0	2.5	9.4
1999	0.4	*	*	*	*	*	*	0.1	0.5	0.9	2.2	9.0
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	3.8	*
2001	0.2	*	*	*	*	*	*	*	*	*	4.1	*
2000	0.2	*	*	*	*	*	*	*	*	*	4.1	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	16.8
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.2	0.5	1.4	3.2	9.8
2001	0.5	*	*	*	*	*	0.1	0.3	0.4	1.3	2.9	9.7
2000	0.5	*	*	*	*	*	0.1	0.2	0.5	1.1	3.0	9.8
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.6	9.0
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	0.1	0.2	0.6	1.8	4.1	10.2
2001	0.5	*	*	*	*	*	0.1	0.3	0.5	1.6	3.4	9.6
2000	0.5	*	*	*	*	*	*	0.4	0.5	1.3	4.0	11.0
1999	0.4	*	*	*	*	*	*	0.3	0.6	1.2	3.2	10.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.1	0.3	1.1	2.5	9.7
2001	0.5	*	*	*	*	*	*	0.2	0.3	0.9	2.7	9.7
2000	0.5	*	*	*	*	*	0.1	0.1	0.4	1.0	2.4	9.3
1999	0.5	*	*	*	*	*	*	*	0.4	0.9	2.2	8.6
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	0.1	0.2	0.4	1.3	3.1	9.8
2001	0.5	*	*	*	*	*	0.1	0.2	0.4	1.2	2.9	9.7
2000	0.5	*	*	*	*	*	0.1	0.2	0.4	1.1	3.0	9.6
1999	0.5	*	*	*	*	*	0.1	0.2	0.5	1.0	2.6	8.8
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	*	0.2	0.6	1.6	4.2	10.4
2001	0.5	*	*	*	*	*	*	0.3	0.5	1.5	3.4	9.4
2000	0.5	*	*	*	*	*	*	0.3	0.5	1.3	4.0	10.5
1999	0.4	*	*	*	*	*	*	0.3	0.6	1.2	3.2	9.7
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	0.3	1.1	2.5	9.6
2001	0.6	*	*	*	*	*	*	0.2	0.3	0.9	2.6	9.8
2000	0.5	*	*	*	*	*	*	*	0.4	0.9	2.3	9.3
1999	0.5	*	*	*	*	*	*	*	0.4	0.8	2.2	8.4
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	*	2.3	3.5	10.3
2001	0.4	*	*	*	*	*	*	*	*	2.0	3.0	8.7
2000	0.4	*	*	*	*	*	*	*	*	1.9	3.9	11.8
1999	0.4	*	*	*	*	*	*	*	0.9	*	2.8	10.3
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	3.4	*	*
2001	0.4	*	*	*	*	*	*	*	*	3.0	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.6	10.5
2001	0.4	*	*	*	*	*	*	*	*	*	*	8.6
2000	0.4	*	*	*	*	*	*	*	*	*	3.8	10.5
1999	0.4	*	*	*	*	*	*	*	*	*	*	8.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of digestive system (K90,K92)												
All origins 3/, both sexes												
2002	2.9	0.6	0.1	*	*	0.1	0.3	1.0	2.1	6.1	19.7	76.2
2001	2.9	0.6	*	*	*	0.1	0.3	0.8	2.4	6.2	19.9	77.8
2000	2.9	0.6	*	*	*	0.1	0.3	0.9	2.1	5.9	20.6	80.6
1999	2.8	0.7	0.1	*	*	0.1	0.3	0.8	2.2	6.2	19.6	77.2
All origins 3/, male												
2002	2.6	*	*	*	*	0.1	0.4	1.4	2.9	7.8	23.0	76.2
2001	2.6	*	*	*	*	0.1	0.4	1.3	3.1	7.9	23.5	81.4
2000	2.6	*	*	*	*	0.1	0.4	1.3	2.8	7.1	25.1	83.1
1999	2.5	*	*	*	*	*	0.4	1.2	2.9	7.9	23.7	80.8
All origins 3/, female												
2002	3.1	*	*	*	*	*	0.1	0.5	1.4	4.7	17.6	76.2
2001	3.1	*	*	*	*	*	0.2	0.4	1.7	4.8	17.5	76.3
2000	3.2	*	*	*	*	*	0.2	0.5	1.4	4.9	17.7	79.6
1999	3.0	*	*	*	*	*	0.2	0.5	1.6	4.8	16.9	75.8
Hispanic, both sexes												
2002	0.6	*	*	*	*	*	*	*	1.2	4.0	11.8	29.4
2001	0.6	*	*	*	*	*	*	*	1.5	3.3	9.7	46.9
2000	0.6	*	*	*	*	*	*	*	*	3.5	11.5	40.5
1999	0.6	*	*	*	*	*	*	*	*	4.1	8.5	44.2
Hispanic, male												
2002	0.7	*	*	*	*	*	*	*	2.1	5.7	14.4	*
2001	0.6	*	*	*	*	*	*	*	*	*	10.3	66.7
2000	0.6	*	*	*	*	*	*	*	*	4.2	11.8	42.3
1999	0.6	*	*	*	*	*	*	*	*	5.9	*	56.1
Hispanic, female												
2002	0.6	*	*	*	*	*	*	*	*	*	10.1	31.5
2001	0.6	*	*	*	*	*	*	*	*	3.5	9.4	37.2
2000	0.6	*	*	*	*	*	*	*	*	*	11.2	39.6
1999	0.6	*	*	*	*	*	*	*	*	*	8.7	38.5
Non-Hispanic 4/, both sexes												
2002	3.2	*	*	*	*	0.1	0.3	1.0	2.2	6.2	20.1	77.9
2001	3.2	0.7	*	*	*	0.1	0.3	0.9	2.4	6.4	20.3	78.8
2000	3.2	*	*	*	*	0.1	0.3	0.9	2.1	6.0	20.9	81.9
1999	3.1	0.8	*	*	*	*	0.3	0.8	2.3	6.3	20.0	78.1
Non-Hispanic 4/, male												
2002	2.9	*	*	*	*	*	0.4	1.4	3.0	7.9	23.4	78.3
2001	2.9	*	*	*	*	0.1	0.5	1.3	3.1	8.2	24.1	82.0
2000	2.9	*	*	*	*	0.1	0.4	1.3	2.9	7.3	25.6	84.8
1999	2.8	*	*	*	*	*	0.4	1.2	3.0	8.0	24.3	81.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases of digestive system (K90,K92)												
Non-Hispanic 4/, female												
2002	3.5	*	*	*	*	*	0.2	0.6	1.4	4.8	17.8	77.8
2001	3.5	*	*	*	*	*	0.2	0.5	1.8	4.9	17.9	77.5
2000	3.5	*	*	*	*	*	0.2	0.5	1.5	5.0	17.9	80.7
1999	3.4	*	*	*	*	*	0.2	0.5	1.7	4.9	17.2	76.7
Non-Hispanic white, both sexes												
2002	3.5	*	*	*	*	*	0.2	0.9	2.0	5.9	19.8	79.8
2001	3.5	*	*	*	*	0.1	0.3	0.8	2.1	6.1	20.2	80.2
2000	3.5	*	*	*	*	*	0.2	0.8	1.8	5.7	20.5	83.1
1999	3.4	*	*	*	*	*	0.2	0.7	2.1	6.0	19.6	79.7
Non-Hispanic white, male												
2002	3.1	*	*	*	*	*	0.3	1.3	2.8	7.4	23.2	79.5
2001	3.1	*	*	*	*	*	0.4	1.1	2.8	7.8	23.8	83.4
2000	3.0	*	*	*	*	*	0.3	1.1	2.5	6.8	25.1	86.7
1999	3.0	*	*	*	*	*	0.4	1.0	2.7	7.6	23.9	82.8
Non-Hispanic white, female												
2002	3.9	*	*	*	*	*	0.1	0.5	1.2	4.6	17.6	79.9
2001	3.9	*	*	*	*	*	0.2	0.4	1.5	4.8	17.8	78.8
2000	3.8	*	*	*	*	*	0.1	0.4	1.2	4.7	17.5	81.6
1999	3.7	*	*	*	*	*	0.1	0.4	1.5	4.7	16.8	78.5
Non-Hispanic black, both sexes												
2002	2.5	*	*	*	*	*	0.6	1.8	4.4	10.0	26.7	65.9
2001	2.5	*	*	*	*	*	0.6	1.7	5.2	9.9	26.0	73.4
2000	2.6	*	*	*	*	*	0.7	2.0	5.0	10.0	28.2	79.2
1999	2.3	*	*	*	*	*	0.4	1.7	4.3	10.0	26.7	66.9
Non-Hispanic black, male												
2002	2.5	*	*	*	*	*	0.8	2.6	5.7	14.1	30.8	72.2
2001	2.6	*	*	*	*	*	0.8	2.8	6.9	13.9	32.7	71.6
2000	2.6	*	*	*	*	*	1.0	2.8	6.7	12.1	35.5	79.8
1999	2.3	*	*	*	*	*	*	2.5	5.7	13.1	32.8	72.5
Non-Hispanic black, female												
2002	2.4	*	*	*	*	*	*	1.1	3.3	7.1	24.4	63.5
2001	2.5	*	*	*	*	*	*	*	3.8	7.0	22.2	74.0
2000	2.7	*	*	*	*	*	*	1.3	3.7	8.4	24.1	78.9
1999	2.4	*	*	*	*	*	*	1.0	3.1	7.8	23.3	64.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
All origins 3/, both sexes												
2002	1.4	*	*	*	0.1	0.1	0.3	0.6	1.2	3.1	8.7	32.8
2001	1.3	*	*	*	*	0.1	0.3	0.6	1.2	2.8	8.7	32.2
2000	1.3	*	*	*	*	0.1	0.2	0.5	1.2	3.0	8.9	34.7
1999	1.3	*	*	*	*	0.1	0.2	0.5	1.0	2.9	9.0	35.0
All origins 3/, male												
2002	1.0	*	*	*	*	*	0.2	0.6	1.1	3.2	8.6	28.6
2001	0.9	*	*	*	*	*	0.2	0.6	1.2	3.0	8.3	24.9
2000	0.9	*	*	*	*	*	0.2	0.4	1.3	2.8	8.4	29.7
1999	0.9	*	*	*	*	*	0.2	0.5	1.0	2.8	8.4	31.7
All origins 3/, female												
2002	1.7	*	*	*	*	0.2	0.3	0.6	1.3	3.0	8.8	34.7
2001	1.7	*	*	*	*	0.2	0.3	0.6	1.2	2.6	9.1	35.2
2000	1.7	*	*	*	*	0.2	0.3	0.5	1.1	3.1	9.2	36.7
1999	1.7	*	*	*	*	0.1	0.3	0.5	1.1	2.9	9.4	36.3
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	*	*	1.9	7.0	24.4
2001	0.5	*	*	*	*	*	*	0.6	1.8	2.3	6.5	24.4
2000	0.4	*	*	*	*	*	*	*	1.2	2.0	6.5	29.9
1999	0.4	*	*	*	*	*	*	*	*	2.0	7.5	25.3
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	6.7	24.0
2001	0.5	*	*	*	*	*	*	*	*	*	8.5	24.2
2000	0.6	*	*	*	*	*	*	*	*	*	7.3	30.7
1999	0.5	*	*	*	*	*	*	*	*	*	9.0	29.1
Non-Hispanic 4/, both sexes												
2002	1.5	*	*	*	*	0.1	0.3	0.6	1.2	3.1	8.8	33.1
2001	1.4	*	*	*	*	0.1	0.3	0.6	1.1	2.8	8.8	32.4
2000	1.5	*	*	*	*	0.1	0.2	0.4	1.2	3.0	9.0	34.7
1999	1.4	*	*	*	*	0.1	0.2	0.5	1.0	2.9	9.1	35.2
Non-Hispanic 4/, male												
2002	1.1	*	*	*	*	*	0.2	0.6	1.1	3.2	8.6	28.6
2001	1.0	*	*	*	*	*	0.2	0.6	1.1	3.0	8.4	24.9
2000	1.0	*	*	*	*	*	0.2	0.4	1.3	2.8	8.5	29.6
1999	1.0	*	*	*	*	*	0.2	0.5	1.0	2.8	8.5	32.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XII. Diseases of the skin and subcutaneous tissue (L00-L98)												
Non-Hispanic 4/, female												
2002	1.9	*	*	*	*	0.2	0.3	0.6	1.4	3.0	8.8	35.0
2001	1.8	*	*	*	*	0.2	0.3	0.6	1.2	2.7	9.1	35.5
2000	1.9	*	*	*	*	0.1	0.3	0.5	1.1	3.2	9.3	36.8
1999	1.8	*	*	*	*	*	0.3	0.5	1.1	3.0	9.4	36.4
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	*	*	0.2	0.5	1.0	2.5	7.4	28.6
2001	1.3	*	*	*	*	*	0.2	0.5	0.9	2.4	7.5	28.2
2000	1.3	*	*	*	*	*	0.2	0.3	0.9	2.4	7.6	29.4
1999	1.3	*	*	*	*	0.1	0.2	0.4	0.8	2.3	7.4	29.3
Non-Hispanic white, male												
2002	1.1	*	*	*	*	*	0.2	0.5	0.9	2.6	7.7	26.3
2001	0.9	*	*	*	*	*	0.1	0.5	1.0	2.6	7.0	21.6
2000	0.9	*	*	*	*	*	0.2	0.3	1.1	2.1	7.1	25.4
1999	0.9	*	*	*	*	*	0.1	0.3	0.7	2.2	6.9	27.9
Non-Hispanic white, female												
2002	1.7	*	*	*	*	*	0.2	0.4	1.0	2.5	7.3	29.5
2001	1.7	*	*	*	*	*	0.2	0.5	0.8	2.2	7.9	30.9
2000	1.7	*	*	*	*	*	0.2	0.3	0.8	2.6	7.9	31.0
1999	1.6	*	*	*	*	*	0.2	0.4	0.8	2.4	7.8	29.9
Non-Hispanic black, both sexes												
2002	2.6	*	*	*	*	0.5	1.0	1.5	3.7	8.6	25.9	93.2
2001	2.4	*	*	*	*	0.4	0.8	1.4	3.2	7.4	25.2	89.6
2000	2.5	*	*	*	*	0.4	0.7	1.3	3.1	9.2	26.6	98.9
1999	2.8	*	*	*	*	*	0.8	1.9	3.6	9.0	29.5	106.4
Non-Hispanic black, male												
2002	1.7	*	*	*	*	*	*	1.3	3.0	10.2	24.0	61.3
2001	1.8	*	*	*	*	*	*	1.6	2.6	7.4	28.8	72.8
2000	1.8	*	*	*	*	*	*	*	3.0	10.3	28.4	83.4
1999	2.0	*	*	*	*	*	*	2.1	3.3	9.3	30.0	86.8
Non-Hispanic black, female												
2002	3.4	*	*	*	*	*	1.2	1.6	4.4	7.4	27.0	105.3
2001	3.0	*	*	*	*	*	1.0	1.2	3.6	7.4	23.2	95.8
2000	3.2	*	*	*	*	*	0.7	1.7	3.2	8.3	25.5	104.7
1999	3.4	*	*	*	*	*	0.9	1.6	3.8	8.8	29.2	113.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of skin and subcutaneous tissue (L00-L08)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.6	1.1	2.8	9.3
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.0	2.6	9.4
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.5	9.3
1999	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.1	8.3
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.1	0.4	0.6	1.1	3.0	9.1
2001	0.4	*	*	*	*	*	0.1	0.4	0.6	1.2	2.4	8.1
2000	0.3	*	*	*	*	*	*	0.2	0.7	1.1	2.8	8.9
1999	0.3	*	*	*	*	*	*	0.3	0.4	0.9	2.1	9.1
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.6	9.4
2001	0.5	*	*	*	*	*	*	0.2	0.4	0.9	2.7	10.0
2000	0.5	*	*	*	*	*	0.1	0.1	0.4	0.9	2.4	9.5
1999	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.1	8.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.3	0.6	1.1	2.8	9.5
2001	0.5	*	*	*	*	*	0.1	0.3	0.5	1.0	2.6	9.7
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.6	9.4
1999	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.1	8.4
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	0.1	0.4	0.6	1.2	3.0	9.3
2001	0.4	*	*	*	*	*	0.1	0.4	0.6	1.2	2.5	8.3
2000	0.4	*	*	*	*	*	*	0.2	0.7	1.1	2.9	9.2
1999	0.3	*	*	*	*	*	*	0.3	0.4	0.8	2.1	9.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of skin and subcutaneous tissue (L00-L08)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.7	9.6
2001	0.6	*	*	*	*	*	*	0.2	0.4	0.9	2.7	10.2
2000	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.4	9.5
1999	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.0	8.0
Non-Hispanic white, both sexes												
2002	0.5	*	*	*	*	*	*	0.2	0.5	1.0	2.8	9.9
2001	0.5	*	*	*	*	*	0.1	0.3	0.4	1.0	2.6	9.9
2000	0.5	*	*	*	*	*	0.1	0.1	0.5	1.0	2.5	9.4
1999	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.1	8.4
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	*	0.3	0.6	1.1	3.1	9.7
2001	0.4	*	*	*	*	*	*	0.3	0.6	1.2	2.5	8.6
2000	0.4	*	*	*	*	*	*	0.2	0.7	1.1	2.8	9.0
1999	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.0	9.2
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	0.2	0.4	0.9	2.6	10.0
2001	0.6	*	*	*	*	*	*	0.2	0.3	0.9	2.7	10.4
2000	0.5	*	*	*	*	*	*	*	0.4	0.9	2.4	9.5
1999	0.5	*	*	*	*	*	*	*	0.3	0.8	2.2	8.1
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	*	*	*	*	0.9	2.0	3.8	7.0
2001	0.4	*	*	*	*	*	*	0.5	1.0	*	3.1	9.4
2000	0.4	*	*	*	*	*	*	*	*	1.5	3.1	10.2
1999	0.4	*	*	*	*	*	*	0.7	1.1	*	*	8.7
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	4.1	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	10.7
2000	0.4	*	*	*	*	*	*	*	*	*	*	10.5
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
All origins 3/, both sexes												
2002	0.9	*	*	*	*	0.1	0.2	0.3	0.7	2.0	5.9	23.5
2001	0.9	*	*	*	*	0.1	0.1	0.3	0.7	1.8	6.1	22.7
2000	0.9	*	*	*	*	0.1	0.2	0.3	0.6	1.9	6.4	25.4
1999	1.0	*	*	*	*	0.1	0.2	0.3	0.6	2.0	7.0	26.6
All origins 3/, male												
2002	0.6	*	*	*	*	*	0.1	0.2	0.5	2.0	5.7	19.4
2001	0.6	*	*	*	*	*	0.1	0.2	0.5	1.8	5.8	16.8
2000	0.6	*	*	*	*	*	0.1	0.2	0.6	1.6	5.6	20.8
1999	0.6	*	*	*	*	*	0.1	0.2	0.5	2.0	6.3	22.6
All origins 3/, female												
2002	1.2	*	*	*	*	0.2	0.3	0.4	0.8	2.0	6.1	25.3
2001	1.2	*	*	*	*	0.1	0.2	0.3	0.8	1.8	6.4	25.2
2000	1.2	*	*	*	*	0.1	0.2	0.4	0.7	2.2	6.9	27.3
1999	1.3	*	*	*	*	*	0.2	0.4	0.7	2.0	7.4	28.3
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	5.2	20.5
2001	0.3	*	*	*	*	*	*	*	*	*	4.5	21.3
2000	0.3	*	*	*	*	*	*	*	*	*	4.7	23.9
1999	0.3	*	*	*	*	*	*	*	*	*	5.6	19.6
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	20.7
2001	0.4	*	*	*	*	*	*	*	*	*	6.0	21.4
2000	0.4	*	*	*	*	*	*	*	*	*	*	22.8
1999	0.4	*	*	*	*	*	*	*	*	*	6.9	21.8
Non-Hispanic 4/, both sexes												
2002	1.0	*	*	*	*	0.1	0.2	0.3	0.7	2.1	6.0	23.6
2001	1.0	*	*	*	*	0.1	0.2	0.3	0.6	1.8	6.2	22.7
2000	1.0	*	*	*	*	0.1	0.2	0.3	0.6	2.0	6.4	25.3
1999	1.1	*	*	*	*	0.1	0.2	0.3	0.6	2.0	7.0	26.8
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	0.1	0.2	0.5	2.1	5.6	19.3
2001	0.6	*	*	*	*	*	*	0.2	0.5	1.8	5.9	16.6
2000	0.6	*	*	*	*	*	0.1	0.2	0.6	1.6	5.7	20.4
1999	0.7	*	*	*	*	*	0.1	0.3	0.6	1.9	6.4	22.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)												
Non-Hispanic 4/, female												
2002	1.3	*	*	*	*	0.1	0.3	0.4	0.9	2.0	6.2	25.4
2001	1.3	*	*	*	*	0.2	0.2	0.3	0.7	1.8	6.4	25.3
2000	1.4	*	*	*	*	*	0.2	0.4	0.7	2.3	6.9	27.2
1999	1.4	*	*	*	*	*	0.2	0.4	0.7	2.1	7.4	28.5
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	*	0.1	0.2	0.5	1.6	4.7	18.6
2001	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	4.9	18.3
2000	0.8	*	*	*	*	*	0.1	0.2	0.4	1.4	5.0	20.0
1999	0.9	*	*	*	*	*	0.1	0.2	0.4	1.4	5.3	20.9
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	*	0.2	0.3	1.5	4.6	16.7
2001	0.5	*	*	*	*	*	*	0.1	0.4	1.4	4.5	13.0
2000	0.5	*	*	*	*	*	*	0.2	0.4	1.0	4.2	16.4
1999	0.6	*	*	*	*	*	*	0.2	0.4	1.3	4.9	18.7
Non-Hispanic white, female												
2002	1.1	*	*	*	*	*	0.1	0.3	0.6	1.6	4.7	19.5
2001	1.1	*	*	*	*	*	*	0.3	0.5	1.3	5.2	20.5
2000	1.2	*	*	*	*	*	*	0.2	0.5	1.8	5.6	21.4
1999	1.2	*	*	*	*	*	0.2	0.3	0.5	1.5	5.6	21.8
Non-Hispanic black, both sexes												
2002	2.2	*	*	*	*	0.4	0.7	1.1	2.8	6.6	22.1	86.2
2001	2.0	*	*	*	*	*	0.6	0.9	2.1	6.3	22.1	80.2
2000	2.2	*	*	*	*	*	0.5	1.0	2.4	7.7	23.4	88.7
1999	2.4	*	*	*	*	*	0.7	1.2	2.5	7.9	27.8	97.6
Non-Hispanic black, male												
2002	1.4	*	*	*	*	*	*	*	2.2	7.9	20.8	55.8
2001	1.4	*	*	*	*	*	*	*	1.8	6.4	25.8	67.0
2000	1.5	*	*	*	*	*	*	*	2.4	8.2	25.0	74.0
1999	1.7	*	*	*	*	*	*	1.1	2.2	8.9	26.9	74.9
Non-Hispanic black, female												
2002	3.0	*	*	*	*	*	1.1	1.3	3.3	5.6	22.9	97.8
2001	2.5	*	*	*	*	*	0.8	0.9	2.4	6.2	20.0	85.1
2000	2.8	*	*	*	*	*	*	1.5	2.4	7.3	22.6	94.2
1999	3.1	*	*	*	*	*	0.8	1.3	2.7	7.2	28.4	106.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
All origins 3/, both sexes												
2002	5.0	*	*	0.1	0.4	0.7	1.4	2.7	5.5	12.9	31.7	94.9
2001	4.9	*	*	0.1	0.5	0.7	1.5	2.6	5.3	12.8	32.2	93.4
2000	4.9	*	*	0.1	0.4	0.8	1.4	2.6	5.4	12.6	32.2	94.0
1999	4.8	*	*	0.1	0.4	0.7	1.4	2.7	5.4	12.7	31.1	91.2
All origins 3/, male												
2002	3.0	*	*	*	0.2	0.4	0.9	1.9	4.0	10.2	24.1	63.5
2001	2.9	*	*	*	0.2	0.4	1.0	1.9	3.9	9.6	24.9	63.2
2000	2.9	*	*	*	0.3	0.5	0.9	1.7	4.0	9.8	25.2	67.5
1999	2.8	*	*	*	0.2	0.4	0.9	1.8	3.8	9.5	24.2	63.7
All origins 3/, female												
2002	7.0	*	*	0.1	0.6	1.0	1.8	3.3	6.9	15.1	36.8	108.5
2001	6.9	*	*	*	0.7	1.1	2.0	3.2	6.6	15.6	37.0	106.0
2000	6.8	*	*	0.1	0.5	1.1	1.9	3.5	6.6	15.0	36.8	104.8
1999	6.7	*	*	0.1	0.6	1.1	1.9	3.7	6.9	15.3	35.5	102.3
Hispanic, both sexes												
2002	1.9	*	*	*	0.4	0.5	1.4	2.7	5.6	12.6	20.9	53.2
2001	2.0	*	*	*	0.6	0.6	1.6	3.0	5.0	12.9	22.4	55.6
2000	1.9	*	*	*	*	0.9	1.2	2.5	5.7	13.0	26.9	55.1
1999	2.1	*	*	*	0.4	0.8	1.5	3.0	6.9	12.8	22.7	73.7
Hispanic, male												
2002	1.0	*	*	*	*	*	1.1	1.6	3.1	6.7	13.5	36.8
2001	1.2	*	*	*	*	*	1.1	2.6	2.9	7.4	19.2	41.9
2000	1.0	*	*	*	*	*	0.8	1.5	4.5	7.4	14.3	40.3
1999	1.1	*	*	*	*	*	0.9	1.6	4.3	8.6	16.7	47.4
Hispanic, female												
2002	2.9	*	*	*	0.8	0.9	1.8	3.7	7.9	17.3	25.9	61.3
2001	2.8	*	*	*	1.1	0.9	2.1	3.5	7.0	17.3	24.5	62.4
2000	2.9	*	*	*	*	1.4	1.7	3.5	6.7	17.4	35.3	62.3
1999	3.1	*	*	*	0.7	1.3	2.2	4.3	9.2	16.2	26.7	86.3
Non-Hispanic 4/, both sexes												
2002	5.5	*	*	0.1	0.4	0.7	1.4	2.6	5.5	12.8	32.1	96.2
2001	5.4	*	*	0.1	0.4	0.8	1.5	2.5	5.3	12.8	32.6	94.6
2000	5.3	*	*	0.1	0.4	0.8	1.5	2.6	5.3	12.6	32.4	95.2
1999	5.1	*	*	0.1	0.4	0.7	1.4	2.7	5.3	12.7	31.3	91.5
Non-Hispanic 4/, male												
2002	3.3	*	*	*	0.2	0.5	0.8	2.0	4.1	10.4	24.6	64.7
2001	3.2	*	*	*	0.2	0.4	1.0	1.8	3.9	9.7	25.0	64.0
2000	3.2	*	*	*	0.3	0.5	0.9	1.7	3.9	9.9	25.6	68.4
1999	3.0	*	*	*	0.2	0.4	0.9	1.8	3.8	9.6	24.5	63.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)												
Non-Hispanic 4/, female												
2002	7.5	*	*	0.2	0.6	1.0	1.9	3.3	6.8	14.9	37.2	109.9
2001	7.4	*	*	*	0.6	1.1	1.9	3.2	6.5	15.4	37.6	107.3
2000	7.3	*	*	0.1	0.5	1.1	1.9	3.5	6.6	14.8	36.7	106.1
1999	7.1	*	*	0.1	0.6	1.0	1.9	3.6	6.6	15.3	35.8	102.5
Non-Hispanic white, both sexes												
2002	5.8	*	*	0.1	0.2	0.4	0.8	2.1	5.0	12.7	32.6	99.7
2001	5.7	*	*	0.1	0.2	0.4	1.0	2.0	4.8	12.5	33.3	97.1
2000	5.6	*	*	*	0.2	0.5	0.9	2.0	4.8	12.5	32.8	98.2
1999	5.4	*	*	*	0.2	0.4	0.8	2.1	4.7	12.5	32.0	94.7
Non-Hispanic white, male												
2002	3.4	*	*	*	*	0.3	0.6	1.7	3.7	10.0	24.5	65.5
2001	3.3	*	*	*	0.2	0.3	0.8	1.5	3.5	9.4	25.3	65.1
2000	3.3	*	*	*	*	0.3	0.6	1.4	3.7	9.4	25.4	68.2
1999	3.1	*	*	*	*	0.3	0.6	1.5	3.4	9.5	24.4	64.6
Non-Hispanic white, female												
2002	8.1	*	*	*	0.2	0.5	1.1	2.6	6.2	14.9	38.0	114.5
2001	7.9	*	*	*	0.3	0.6	1.2	2.5	6.0	15.3	38.6	110.4
2000	7.8	*	*	*	0.3	0.6	1.2	2.6	5.9	15.0	37.6	110.3
1999	7.6	*	*	*	0.2	0.6	1.0	2.6	6.0	15.1	37.0	106.7
Non-Hispanic black, both sexes												
2002	5.0	*	*	*	1.2	2.6	4.3	6.4	10.2	16.2	31.3	68.4
2001	4.9	*	*	*	1.2	2.5	4.4	6.5	9.6	16.7	29.7	76.2
2000	5.0	*	*	*	1.3	2.5	4.7	7.3	10.0	15.5	31.1	68.7
1999	4.8	*	*	*	1.2	2.2	4.9	7.2	9.8	16.4	27.3	60.5
Non-Hispanic black, male												
2002	3.2	*	*	*	*	1.5	2.3	4.1	8.0	15.5	28.4	61.3
2001	3.2	*	*	*	*	1.2	2.8	4.9	7.9	13.9	28.2	60.1
2000	3.5	*	*	*	1.0	1.4	3.1	4.1	7.2	16.6	30.9	78.7
1999	3.0	*	*	*	*	0.9	3.0	4.2	6.7	13.1	29.4	64.2
Non-Hispanic black, female												
2002	6.6	*	*	*	1.9	3.6	6.2	8.3	12.1	16.6	33.0	71.1
2001	6.5	*	*	*	2.0	3.7	5.7	7.8	11.0	18.7	30.5	82.1
2000	6.4	*	*	*	1.6	3.5	6.1	10.0	12.4	14.6	31.3	65.0
1999	6.4	*	*	*	1.9	3.4	6.6	9.8	12.3	18.7	26.1	59.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
All origins 3/, both sexes												
2002	1.0	*	*	*	*	*	0.0	0.2	1.0	3.3	9.1	14.2
2001	0.9	*	*	*	*	*	0.1	0.2	1.0	3.3	8.8	12.7
2000	1.0	*	*	*	*	*	0.1	0.2	1.0	3.5	9.2	12.9
1999	1.0	*	*	*	*	*	0.0	0.2	1.1	3.5	9.5	13.9
All origins 3/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.7	2.1	5.3	7.6
2001	0.4	*	*	*	*	*	*	0.1	0.7	2.0	5.1	5.5
2000	0.5	*	*	*	*	*	*	0.1	0.6	2.3	5.2	6.9
1999	0.5	*	*	*	*	*	*	0.2	0.8	2.3	5.2	7.2
All origins 3/, female												
2002	1.4	*	*	*	*	*	*	0.3	1.3	4.3	11.6	17.0
2001	1.4	*	*	*	*	*	0.1	0.3	1.3	4.4	11.2	15.7
2000	1.4	*	*	*	*	*	*	0.4	1.3	4.5	11.7	15.3
1999	1.5	*	*	*	*	*	*	0.3	1.5	4.4	12.2	16.7
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	3.0	6.7	*
2001	0.3	*	*	*	*	*	*	*	*	4.5	6.0	*
2000	0.3	*	*	*	*	*	*	*	*	3.7	8.3	*
1999	0.4	*	*	*	*	*	*	*	1.4	3.3	7.3	20.3
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	4.3	10.1	*
2001	0.6	*	*	*	*	*	*	*	*	7.2	7.3	*
2000	0.5	*	*	*	*	*	*	*	*	5.1	11.9	*
1999	0.6	*	*	*	*	*	*	*	*	5.2	10.1	25.0
Non-Hispanic 4/, both sexes												
2002	1.1	*	*	*	*	*	*	0.2	1.0	3.3	9.2	14.3
2001	1.0	*	*	*	*	*	0.1	0.2	1.0	3.2	8.9	12.8
2000	1.0	*	*	*	*	*	*	0.3	1.0	3.5	9.2	13.2
1999	1.1	*	*	*	*	*	*	0.2	1.1	3.5	9.5	13.7
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.7	2.2	5.5	7.7
2001	0.5	*	*	*	*	*	*	0.1	0.7	2.1	5.1	5.7
2000	0.5	*	*	*	*	*	*	*	0.6	2.4	5.3	7.1
1999	0.5	*	*	*	*	*	*	0.2	0.8	2.4	5.3	7.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)												
Non-Hispanic 4/, female												
2002	1.6	*	*	*	*	*	*	0.3	1.3	4.3	11.6	17.1
2001	1.5	*	*	*	*	*	0.1	0.3	1.3	4.2	11.4	15.8
2000	1.6	*	*	*	*	*	*	0.4	1.3	4.4	11.7	15.6
1999	1.6	*	*	*	*	*	*	0.3	1.4	4.4	12.3	16.4
Non-Hispanic white, both sexes												
2002	1.2	*	*	*	*	*	*	0.2	1.0	3.5	9.5	14.9
2001	1.2	*	*	*	*	*	*	0.2	1.1	3.4	9.3	13.0
2000	1.2	*	*	*	*	*	*	0.3	1.0	3.7	9.5	13.7
1999	1.2	*	*	*	*	*	*	0.3	1.2	3.7	9.9	14.6
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	*	0.2	0.7	2.2	5.7	8.1
2001	0.6	*	*	*	*	*	*	*	0.8	2.2	5.3	5.8
2000	0.6	*	*	*	*	*	*	*	0.7	2.5	5.4	7.5
1999	0.6	*	*	*	*	*	*	0.2	0.9	2.5	5.4	7.6
Non-Hispanic white, female												
2002	1.8	*	*	*	*	*	*	0.3	1.3	4.5	12.0	17.8
2001	1.8	*	*	*	*	*	*	0.3	1.3	4.5	12.0	16.0
2000	1.8	*	*	*	*	*	*	0.4	1.3	4.7	12.2	16.1
1999	1.9	*	*	*	*	*	*	0.3	1.5	4.7	12.8	17.4
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	*	*	0.9	2.5	6.7	8.2
2001	0.5	*	*	*	*	*	*	*	0.9	2.2	5.1	12.2
2000	0.4	*	*	*	*	*	*	*	0.9	2.0	5.2	7.9
1999	0.4	*	*	*	*	*	*	*	*	2.0	6.2	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	*	*	2.7	8.1	10.0
2001	0.7	*	*	*	*	*	*	*	*	2.7	5.4	15.0
2000	0.6	*	*	*	*	*	*	*	*	*	6.4	9.6
1999	0.6	*	*	*	*	*	*	*	*	2.2	6.8	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Systemic lupus erythematosus (M32)												
All origins 3/, both sexes												
2002	0.4	*	*	*	0.2	0.3	0.5	0.6	0.8	1.0	1.3	1.3
2001	0.5	*	*	*	0.3	0.4	0.5	0.6	0.8	1.0	1.5	1.2
2000	0.5	*	*	*	0.2	0.3	0.6	0.7	0.9	0.9	1.5	1.2
1999	0.5	*	*	*	0.2	0.3	0.5	0.7	0.8	1.1	1.3	1.4
All origins 3/, male												
2002	0.1	*	*	*	*	*	0.1	0.2	0.2	0.4	0.6	*
2001	0.1	*	*	*	*	0.1	0.1	0.2	0.2	0.3	0.6	*
2000	0.1	*	*	*	*	*	0.2	0.1	0.3	0.4	0.7	*
1999	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.6	*
All origins 3/, female												
2002	0.8	*	*	*	0.4	0.5	0.9	1.0	1.4	1.5	1.8	1.7
2001	0.8	*	*	*	0.5	0.7	0.8	1.1	1.3	1.6	2.1	1.3
2000	0.8	*	*	*	0.3	0.6	1.0	1.1	1.4	1.3	2.0	1.4
1999	0.8	*	*	*	0.4	0.6	0.9	1.2	1.3	1.8	1.8	1.6
Hispanic, both sexes												
2002	0.4	*	*	*	*	0.3	0.6	0.8	1.2	1.8	*	*
2001	0.4	*	*	*	0.4	*	0.5	0.7	*	*	*	*
2000	0.4	*	*	*	*	0.4	0.5	0.7	1.3	*	*	*
1999	0.5	*	*	*	*	0.4	0.7	1.1	1.4	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.7	*	*	*	*	*	1.0	1.4	2.1	*	*	*
2001	0.7	*	*	*	0.8	*	0.8	1.2	*	*	*	*
2000	0.7	*	*	*	*	0.8	0.9	1.5	2.3	*	*	*
1999	0.8	*	*	*	*	0.9	1.3	2.0	2.4	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	0.2	0.3	0.5	0.6	0.8	0.9	1.3	1.4
2001	0.5	*	*	*	0.2	0.4	0.5	0.6	0.8	1.0	1.5	1.3
2000	0.5	*	*	*	0.2	0.3	0.6	0.6	0.9	0.9	1.4	1.2
1999	0.5	*	*	*	0.2	0.3	0.5	0.6	0.7	1.1	1.3	1.5
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	0.2	0.2	0.3	0.6	*
2001	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.7	*
2000	0.2	*	*	*	*	*	0.2	0.2	0.3	0.4	0.6	*
1999	0.1	*	*	*	*	*	0.1	0.2	0.2	0.3	0.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Systemic lupus erythematosus (M32)												
Non-Hispanic 4/, female												
2002	0.8	*	*	*	0.4	0.5	0.8	1.0	1.3	1.4	1.8	1.8
2001	0.8	*	*	*	0.4	0.7	0.8	1.1	1.3	1.6	2.1	1.3
2000	0.8	*	*	*	0.3	0.6	1.0	1.1	1.3	1.3	2.0	1.5
1999	0.8	*	*	*	0.3	0.6	0.9	1.1	1.1	1.8	1.8	1.6
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	*	0.1	0.2	0.4	0.6	0.7	1.3	1.4
2001	0.4	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.5	1.2
2000	0.4	*	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4	1.4
1999	0.3	*	*	*	*	0.1	0.2	0.4	0.6	0.9	1.4	1.5
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	0.1	*	0.3	0.6	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	*
2000	0.1	*	*	*	*	*	*	*	0.3	0.4	0.6	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	*
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	0.3	0.6	0.9	1.0	1.8	1.8
2001	0.6	*	*	*	*	0.3	0.3	0.7	1.0	1.4	2.1	1.2
2000	0.6	*	*	*	*	0.2	0.5	0.7	1.0	1.2	1.9	1.6
1999	0.6	*	*	*	*	0.2	0.4	0.7	0.9	1.5	1.8	1.7
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	0.8	1.4	2.0	2.0	2.5	2.6	*	*
2001	1.2	*	*	*	0.8	1.5	2.0	2.0	2.1	1.7	*	*
2000	1.3	*	*	*	0.8	1.3	2.1	2.4	2.5	1.9	*	*
1999	1.2	*	*	*	0.8	1.2	2.2	2.2	1.9	2.9	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	0.8	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	2.2	*	*	*	1.4	2.3	3.6	3.5	4.4	3.8	*	*
2001	2.0	*	*	*	1.5	2.5	3.4	3.3	3.1	2.6	*	*
2000	2.1	*	*	*	1.1	2.3	3.3	4.1	3.8	2.8	*	*
1999	2.1	*	*	*	1.4	2.1	3.7	3.8	3.0	4.5	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
All origins 3/, both sexes												
2002	1.7	*	*	*	0.1	0.3	0.6	1.2	2.2	4.7	9.7	27.2
2001	1.7	*	*	*	0.1	0.2	0.6	1.1	2.1	5.0	10.1	27.7
2000	1.7	*	*	*	0.1	0.3	0.6	1.1	2.2	5.0	10.5	28.0
1999	1.7	*	*	*	0.1	0.2	0.6	1.3	2.3	5.0	9.5	27.6
All origins 3/, male												
2002	1.0	*	*	*	*	0.2	0.4	0.8	1.6	3.3	7.6	18.6
2001	1.0	*	*	*	*	0.2	0.4	0.9	1.5	3.4	8.1	20.5
2000	1.1	*	*	*	*	0.3	0.4	0.8	1.6	3.4	8.8	21.1
1999	1.0	*	*	*	*	0.2	0.5	0.8	1.6	3.5	7.8	22.5
All origins 3/, female												
2002	2.3	*	*	*	0.1	0.3	0.7	1.5	2.8	5.9	11.1	31.0
2001	2.3	*	*	*	0.2	0.3	0.8	1.3	2.7	6.3	11.5	30.7
2000	2.3	*	*	*	0.1	0.4	0.7	1.5	2.8	6.3	11.6	30.8
1999	2.3	*	*	*	0.2	0.3	0.7	1.7	3.0	6.3	10.6	29.7
Hispanic, both sexes												
2002	0.7	*	*	*	*	*	0.6	1.0	2.3	4.8	6.7	12.7
2001	0.8	*	*	*	*	*	0.7	1.2	2.7	5.2	7.4	18.8
2000	0.7	*	*	*	*	0.3	0.4	1.1	2.3	4.6	9.3	19.9
1999	0.7	*	*	*	*	*	0.5	1.2	2.6	5.5	6.9	21.0
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	4.4	*	*
Hispanic, female												
2002	1.0	*	*	*	*	*	*	1.4	3.3	7.5	7.9	*
2001	1.1	*	*	*	*	*	0.9	1.4	3.8	6.6	9.1	19.5
2000	1.0	*	*	*	*	*	*	1.3	2.5	6.8	11.9	*
1999	1.1	*	*	*	*	*	*	1.7	3.6	6.4	6.9	27.0
Non-Hispanic 4/, both sexes												
2002	1.8	*	*	*	0.1	0.3	0.6	1.2	2.2	4.7	9.8	27.7
2001	1.8	*	*	*	0.1	0.2	0.6	1.1	2.1	5.0	10.3	27.9
2000	1.8	*	*	*	0.1	0.3	0.6	1.1	2.2	5.0	10.5	28.2
1999	1.8	*	*	*	0.1	0.2	0.6	1.2	2.3	5.0	9.6	27.7
Non-Hispanic 4/, male												
2002	1.1	*	*	*	*	0.2	0.4	0.8	1.6	3.4	7.8	19.0
2001	1.1	*	*	*	*	0.2	0.4	0.9	1.5	3.4	8.2	20.6
2000	1.1	*	*	*	*	0.3	0.4	0.7	1.6	3.4	8.9	20.7
1999	1.1	*	*	*	*	0.1	0.5	0.8	1.6	3.5	7.8	22.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)												
Non-Hispanic 4/, female												
2002	2.5	*	*	*	0.1	0.4	0.7	1.5	2.8	5.8	11.2	31.4
2001	2.5	*	*	*	0.2	0.3	0.8	1.3	2.6	6.3	11.6	30.9
2000	2.5	*	*	*	0.1	0.4	0.8	1.5	2.8	6.3	11.5	31.3
1999	2.4	*	*	*	0.1	0.3	0.7	1.6	2.9	6.2	10.7	29.6
Non-Hispanic white, both sexes												
2002	1.9	*	*	*	*	0.2	0.4	0.9	2.1	4.7	10.0	28.5
2001	1.9	*	*	*	0.1	0.1	0.4	0.8	1.9	4.8	10.6	28.8
2000	1.9	*	*	*	*	0.2	0.3	0.8	2.0	4.9	10.7	29.0
1999	1.8	*	*	*	*	0.1	0.4	0.9	2.0	4.9	9.8	28.5
Non-Hispanic white, male												
2002	1.1	*	*	*	*	*	0.3	0.6	1.4	3.3	7.9	19.6
2001	1.2	*	*	*	*	*	0.3	0.6	1.4	3.3	8.4	21.2
2000	1.1	*	*	*	*	0.2	0.2	0.6	1.4	3.3	8.8	20.2
1999	1.1	*	*	*	*	*	0.3	0.6	1.4	3.4	7.9	23.8
Non-Hispanic white, female												
2002	2.7	*	*	*	*	0.2	0.5	1.2	2.7	5.8	11.5	32.4
2001	2.6	*	*	*	*	0.2	0.6	1.0	2.4	6.2	12.1	31.9
2000	2.6	*	*	*	*	0.2	0.5	1.0	2.6	6.3	11.9	32.6
1999	2.5	*	*	*	*	0.2	0.4	1.1	2.6	6.2	11.0	30.4
Non-Hispanic black, both sexes												
2002	1.8	*	*	*	*	0.9	1.8	2.8	3.9	5.7	8.9	22.1
2001	1.8	*	*	*	*	0.8	1.8	3.2	3.2	6.1	8.6	22.5
2000	1.8	*	*	*	*	0.9	2.0	3.2	4.0	5.8	9.3	21.9
1999	1.9	*	*	*	*	0.8	2.0	3.8	4.8	6.1	7.9	20.7
Non-Hispanic black, male												
2002	1.2	*	*	*	*	*	1.3	2.1	3.9	5.0	6.4	*
2001	1.2	*	*	*	*	*	1.4	2.7	2.4	4.3	7.5	*
2000	1.3	*	*	*	*	*	1.5	1.8	3.2	5.1	10.8	27.0
1999	1.3	*	*	*	*	*	1.8	2.4	3.6	4.7	7.8	*
Non-Hispanic black, female												
2002	2.3	*	*	*	*	1.1	2.3	3.5	3.9	6.1	10.3	25.1
2001	2.3	*	*	*	*	1.1	2.2	3.6	3.9	7.4	9.3	24.4
2000	2.3	*	*	*	*	1.0	2.4	4.4	4.6	6.2	8.5	20.1
1999	2.5	*	*	*	*	1.1	2.2	4.9	5.7	7.1	7.9	22.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Dorsopathies (M40-M54)												
All origins 3/, both sexes												
2002	0.3	*	*	*	*	*	0.1	0.1	0.4	0.9	2.2	6.0
2001	0.3	*	*	*	*	*	0.1	0.1	0.3	0.8	2.4	7.4
2000	0.3	*	*	*	*	*	0.0	0.1	0.3	0.8	2.4	6.1
1999	0.3	*	*	*	*	*	0.0	0.2	0.3	0.8	1.9	5.4
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.4	5.5
2001	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	2.5	6.4
2000	0.3	*	*	*	*	*	*	0.2	0.3	1.0	2.4	6.5
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.8	2.2	5.9
All origins 3/, female												
2002	0.4	*	*	*	*	*	*	*	0.3	0.9	2.1	6.2
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	7.8
2000	0.3	*	*	*	*	*	*	*	0.3	0.6	2.4	6.0
1999	0.3	*	*	*	*	*	*	0.2	0.2	0.7	1.7	5.2
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.3	6.2
2001	0.4	*	*	*	*	*	0.1	0.1	0.3	0.8	2.4	7.5
2000	0.3	*	*	*	*	*	0.0	0.1	0.3	0.8	2.4	6.1
1999	0.3	*	*	*	*	*	0.1	0.2	0.2	0.8	1.9	5.3
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	0.2	0.4	0.9	2.4	5.6
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.8	2.4	6.3
2000	0.3	*	*	*	*	*	*	0.2	0.3	1.0	2.4	6.5
1999	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	5.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Dorsopathies (M40-M54)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	*	*	0.4	0.9	2.2	6.4
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.4	7.9
2000	0.4	*	*	*	*	*	*	*	0.3	0.6	2.4	6.0
1999	0.3	*	*	*	*	*	*	0.2	0.2	0.7	1.8	5.2
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	*	0.2	0.3	0.9	2.3	6.4
2001	0.4	*	*	*	*	*	0.1	0.1	0.3	0.8	2.4	7.9
2000	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.4	6.4
1999	0.3	*	*	*	*	*	*	0.2	0.2	0.7	2.0	5.4
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.4	5.5
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.7	2.5	6.6
2000	0.3	*	*	*	*	*	*	0.2	0.3	0.9	2.4	7.0
1999	0.3	*	*	*	*	*	*	0.2	0.2	0.8	2.2	5.6
Non-Hispanic white, female												
2002	0.5	*	*	*	*	*	*	*	0.4	0.9	2.3	6.8
2001	0.5	*	*	*	*	*	*	*	0.3	0.8	2.3	8.4
2000	0.4	*	*	*	*	*	*	*	0.3	0.6	2.4	6.2
1999	0.4	*	*	*	*	*	*	0.2	0.2	0.7	1.8	5.4
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	2.8	*
2000	0.2	*	*	*	*	*	*	*	*	1.2	2.6	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Soft tissue disorders (M60-M79)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.1	2.4	5.3
2001	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.0	2.3	4.3
2000	0.4	*	*	*	*	0.1	0.1	0.4	0.5	1.0	2.1	4.2
1999	0.3	*	*	*	*	0.1	0.2	0.3	0.5	1.0	2.1	4.3
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.2	0.4	0.6	1.4	2.9	7.8
2001	0.4	*	*	*	*	*	0.2	0.4	0.6	1.4	3.2	6.2
2000	0.4	*	*	*	*	*	0.2	0.4	0.6	1.2	2.9	6.7
1999	0.4	*	*	*	*	*	0.2	0.3	0.6	1.3	3.1	6.8
All origins 3/, female												
2002	0.4	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
2001	0.3	*	*	*	*	*	0.1	0.3	0.6	0.8	1.7	3.5
2000	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	1.6	3.2
1999	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.5	3.3
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	0.7	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.1	2.5	5.3
2001	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.3	4.4
2000	0.4	*	*	*	*	*	0.1	0.4	0.5	1.0	2.2	4.3
1999	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	2.1	4.2
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	0.2	0.4	0.5	1.4	2.9	8.1
2001	0.4	*	*	*	*	*	0.2	0.4	0.6	1.4	3.2	6.3
2000	0.4	*	*	*	*	*	0.2	0.4	0.5	1.2	2.9	6.9
1999	0.4	*	*	*	*	*	0.2	0.3	0.6	1.3	3.1	6.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Soft tissue disorders (M60-M79)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	0.1	0.2	0.5	0.9	2.1	4.2
2001	0.4	*	*	*	*	*	0.1	0.2	0.6	0.8	1.7	3.6
2000	0.4	*	*	*	*	*	0.1	0.3	0.5	0.8	1.7	3.2
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.5	3.3
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.3	0.5	1.0	2.4	5.0
2001	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	2.1	4.3
2000	0.4	*	*	*	*	*	0.1	0.3	0.4	0.9	2.0	4.0
1999	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.0	3.9
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	0.2	0.3	0.5	1.3	2.9	7.9
2001	0.4	*	*	*	*	*	0.2	0.3	0.5	1.3	3.1	6.1
2000	0.4	*	*	*	*	*	*	0.3	0.5	1.1	2.8	6.5
1999	0.4	*	*	*	*	*	0.1	0.2	0.4	1.2	2.8	6.0
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	0.2	0.5	0.8	2.1	3.8
2001	0.3	*	*	*	*	*	*	0.2	0.5	0.7	1.5	3.5
2000	0.3	*	*	*	*	*	*	0.3	0.3	0.8	1.4	3.0
1999	0.3	*	*	*	*	*	*	0.2	0.3	0.6	1.5	3.0
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	*	0.8	1.1	1.9	3.2	9.4
2001	0.5	*	*	*	*	*	*	0.6	1.7	2.1	4.5	*
2000	0.6	*	*	*	*	*	0.4	0.9	1.3	2.0	4.9	7.3
1999	0.5	*	*	*	*	*	0.4	0.7	1.3	1.9	3.4	8.4
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	1.1	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	1.1	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*	6.6	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	*	8.8
2001	0.5	*	*	*	*	*	*	*	1.6	*	3.9	*
2000	0.6	*	*	*	*	*	*	*	1.7	*	4.9	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
All origins 3/, both sexes												
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	7.0	40.8
2001	1.1	*	*	*	*	*	0.1	0.2	0.5	1.7	7.1	40.1
2000	1.1	*	*	*	*	*	0.1	0.1	0.4	1.4	6.6	41.6
1999	1.0	*	*	*	*	*	*	0.2	0.4	1.4	6.7	38.6
All origins 3/, male												
2002	0.6	*	*	*	*	*	*	0.2	0.6	2.1	5.3	23.6
2001	0.6	*	*	*	*	*	0.1	0.2	0.6	1.7	5.4	23.6
2000	0.6	*	*	*	*	*	0.1	0.2	0.5	1.5	5.3	25.6
1999	0.5	*	*	*	*	*	*	0.2	0.4	1.3	5.3	20.4
All origins 3/, female												
2002	1.7	*	*	*	*	*	*	0.1	0.5	1.7	8.1	48.3
2001	1.6	*	*	*	*	*	*	0.1	0.4	1.7	8.2	47.0
2000	1.6	*	*	*	*	*	*	0.1	0.4	1.4	7.5	48.1
1999	1.5	*	*	*	*	*	*	0.1	0.5	1.5	7.7	45.9
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	3.8	23.8
2001	0.2	*	*	*	*	*	*	*	*	*	*	22.5
2000	0.2	*	*	*	*	*	*	*	*	1.9	4.1	22.6
1999	0.2	*	*	*	*	*	*	*	*	*	4.2	18.9
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	24.9
2001	0.3	*	*	*	*	*	*	*	*	*	*	25.1
2000	0.4	*	*	*	*	*	*	*	*	*	*	31.7
1999	0.3	*	*	*	*	*	*	*	*	*	*	23.9
Non-Hispanic 4/, both sexes												
2002	1.3	*	*	*	*	*	0.1	0.2	0.6	1.9	7.1	41.4
2001	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	7.2	40.7
2000	1.2	*	*	*	*	*	0.1	0.1	0.4	1.4	6.7	42.2
1999	1.1	*	*	*	*	*	*	0.1	0.4	1.4	6.8	39.1
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	*	0.2	0.6	2.1	5.4	23.7
2001	0.7	*	*	*	*	*	*	0.2	0.6	1.7	5.4	23.8
2000	0.7	*	*	*	*	*	0.1	0.2	0.5	1.5	5.5	26.4
1999	0.6	*	*	*	*	*	*	0.2	0.4	1.3	5.3	20.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)												
Non-Hispanic 4/, female												
2002	1.9	*	*	*	*	*	*	0.1	0.5	1.7	8.3	49.0
2001	1.8	*	*	*	*	*	*	0.1	0.5	1.8	8.4	47.7
2000	1.7	*	*	*	*	*	*	0.1	0.4	1.4	7.5	48.6
1999	1.7	*	*	*	*	*	*	0.1	0.5	1.5	7.8	46.4
Non-Hispanic white, both sexes												
2002	1.5	*	*	*	*	*	*	0.2	0.5	1.9	7.0	43.4
2001	1.4	*	*	*	*	*	0.1	0.2	0.5	1.6	7.3	42.0
2000	1.4	*	*	*	*	*	0.1	0.1	0.4	1.3	6.8	43.7
1999	1.3	*	*	*	*	*	*	0.1	0.4	1.3	6.9	40.8
Non-Hispanic white, male												
2002	0.8	*	*	*	*	*	*	0.2	0.6	2.0	5.0	23.9
2001	0.7	*	*	*	*	*	*	0.2	0.4	1.5	5.3	24.1
2000	0.7	*	*	*	*	*	*	0.1	0.4	1.3	5.3	26.3
1999	0.6	*	*	*	*	*	*	0.2	0.4	1.2	5.2	20.6
Non-Hispanic white, female												
2002	2.2	*	*	*	*	*	*	0.1	0.4	1.8	8.4	51.9
2001	2.1	*	*	*	*	*	*	0.1	0.5	1.7	8.7	49.4
2000	2.0	*	*	*	*	*	*	*	0.3	1.4	7.7	50.8
1999	2.0	*	*	*	*	*	*	0.1	0.5	1.4	8.0	48.8
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	*	1.2	2.9	8.8	24.2
2001	0.7	*	*	*	*	*	*	*	1.0	3.4	6.8	30.0
2000	0.7	*	*	*	*	*	*	*	1.1	2.5	7.4	27.7
1999	0.6	*	*	*	*	*	*	*	*	2.2	7.0	21.0
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	*	*	4.1	10.0	25.2
2001	0.6	*	*	*	*	*	*	*	*	3.9	7.5	26.6
2000	0.7	*	*	*	*	*	*	*	*	4.0	8.3	32.9
1999	0.5	*	*	*	*	*	*	*	*	*	7.2	25.0
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	*	*	*	8.1	23.8
2001	0.8	*	*	*	*	*	*	*	*	3.0	6.4	31.2
2000	0.7	*	*	*	*	*	*	*	*	*	6.9	25.7
1999	0.7	*	*	*	*	*	*	*	*	2.1	6.8	19.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Osteoporosis (M80-M81)												
All origins 3/, both sexes												
2002	0.6	*	*	*	*	*	*	*	*	0.4	3.4	29.0
2001	0.6	*	*	*	*	*	*	*	*	0.4	3.3	26.8
2000	0.6	*	*	*	*	*	*	*	*	0.4	3.4	28.5
1999	0.6	*	*	*	*	*	*	*	*	0.4	3.5	27.1
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	*	*	*	1.1	9.9
2001	0.1	*	*	*	*	*	*	*	*	*	1.0	9.3
2000	0.1	*	*	*	*	*	*	*	*	*	1.3	9.9
1999	0.1	*	*	*	*	*	*	*	*	*	1.3	7.0
All origins 3/, female												
2002	1.1	*	*	*	*	*	*	*	*	0.6	4.9	37.2
2001	1.0	*	*	*	*	*	*	*	*	0.5	4.8	34.1
2000	1.1	*	*	*	*	*	*	*	*	0.5	4.9	36.0
1999	1.0	*	*	*	*	*	*	*	*	0.6	5.0	35.1
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	11.1
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	13.3
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	19.8
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	*	*	*	*	*	*	0.4	3.5	29.6
2001	0.7	*	*	*	*	*	*	*	*	0.4	3.4	27.4
2000	0.7	*	*	*	*	*	*	*	*	0.4	3.5	29.0
1999	0.7	*	*	*	*	*	*	*	*	0.4	3.6	27.5
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	*	*	*	1.1	10.0
2001	0.2	*	*	*	*	*	*	*	*	*	1.0	9.6
2000	0.2	*	*	*	*	*	*	*	*	*	1.3	10.3
1999	0.1	*	*	*	*	*	*	*	*	*	1.3	7.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Osteoporosis (M80-M81)												
Non-Hispanic 4/, female												
2002	1.3	*	*	*	*	*	*	*	*	0.6	5.0	38.0
2001	1.2	*	*	*	*	*	*	*	*	0.6	4.9	34.8
2000	1.2	*	*	*	*	*	*	*	*	0.5	4.9	36.5
1999	1.2	*	*	*	*	*	*	*	*	0.6	5.0	35.6
Non-Hispanic white, both sexes												
2002	0.9	*	*	*	*	*	*	*	*	0.5	3.7	32.0
2001	0.8	*	*	*	*	*	*	*	*	0.4	3.6	29.5
2000	0.8	*	*	*	*	*	*	*	*	0.4	3.8	31.3
1999	0.8	*	*	*	*	*	*	*	*	0.4	3.9	29.6
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	*	*	1.2	10.7
2001	0.2	*	*	*	*	*	*	*	*	*	1.1	10.4
2000	0.2	*	*	*	*	*	*	*	*	*	1.4	10.7
1999	0.2	*	*	*	*	*	*	*	*	*	1.4	7.6
Non-Hispanic white, female												
2002	1.5	*	*	*	*	*	*	*	*	0.7	5.4	41.3
2001	1.4	*	*	*	*	*	*	*	*	0.6	5.3	37.4
2000	1.4	*	*	*	*	*	*	*	*	0.6	5.4	39.5
1999	1.4	*	*	*	*	*	*	*	*	0.7	5.5	38.3
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
All origins 3/, both sexes												
2002	0.5	*	*	*	*	*	0.0	0.2	0.5	1.5	3.6	11.8
2001	0.5	*	*	*	*	*	0.1	0.2	0.4	1.3	3.8	13.3
2000	0.5	*	*	*	*	*	0.1	0.1	0.4	1.1	3.2	13.1
1999	0.4	*	*	*	*	*	*	0.1	0.4	1.0	3.2	11.5
All origins 3/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.6	1.9	4.2	13.7
2001	0.5	*	*	*	*	*	*	0.2	0.5	1.5	4.4	14.2
2000	0.5	*	*	*	*	*	0.1	0.1	0.5	1.4	4.1	15.6
1999	0.4	*	*	*	*	*	*	0.2	0.4	1.1	4.0	13.3
All origins 3/, female												
2002	0.6	*	*	*	*	*	*	0.1	0.5	1.1	3.2	11.1
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.2	3.4	12.9
2000	0.5	*	*	*	*	*	*	*	0.3	0.9	2.6	12.0
1999	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.8	10.8
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	12.7
2001	0.2	*	*	*	*	*	*	*	*	*	*	12.5
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	*	*	*	*	*	0.1	0.2	0.5	1.5	3.6	11.8
2001	0.6	*	*	*	*	*	0.1	0.2	0.4	1.4	3.9	13.3
2000	0.5	*	*	*	*	*	0.1	0.1	0.4	1.1	3.2	13.2
1999	0.5	*	*	*	*	*	*	0.1	0.4	1.0	3.3	11.6
Non-Hispanic 4/, male												
2002	0.5	*	*	*	*	*	*	0.2	0.6	2.0	4.3	13.7
2001	0.5	*	*	*	*	*	*	0.2	0.5	1.5	4.4	14.2
2000	0.5	*	*	*	*	*	0.1	0.2	0.5	1.3	4.1	16.1
1999	0.4	*	*	*	*	*	*	0.2	0.4	1.2	4.0	13.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)												
Non-Hispanic 4/, female												
2002	0.6	*	*	*	*	*	*	0.1	0.4	1.1	3.2	11.0
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.3	3.5	12.9
2000	0.5	*	*	*	*	*	*	*	0.3	0.8	2.6	12.1
1999	0.5	*	*	*	*	*	*	0.1	0.4	0.9	2.8	10.8
Non-Hispanic white, both sexes												
2002	0.6	*	*	*	*	*	*	0.2	0.4	1.4	3.3	11.4
2001	0.6	*	*	*	*	*	*	0.2	0.4	1.2	3.7	12.5
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.9	3.0	12.5
1999	0.5	*	*	*	*	*	*	0.1	0.3	0.9	3.0	11.2
Non-Hispanic white, male												
2002	0.6	*	*	*	*	*	*	0.2	0.6	1.8	3.9	13.2
2001	0.5	*	*	*	*	*	*	0.2	0.4	1.3	4.2	13.7
2000	0.5	*	*	*	*	*	*	*	0.4	1.1	3.9	15.5
1999	0.4	*	*	*	*	*	*	0.1	0.3	1.1	3.8	12.9
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	0.3	1.1	2.9	10.6
2001	0.7	*	*	*	*	*	*	0.1	0.4	1.1	3.3	12.0
2000	0.6	*	*	*	*	*	*	*	0.3	0.8	2.4	11.2
1999	0.5	*	*	*	*	*	*	*	0.3	0.7	2.6	10.5
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	*	*	1.2	2.7	7.7	19.7
2001	0.7	*	*	*	*	*	*	*	1.0	3.3	6.4	24.3
2000	0.6	*	*	*	*	*	*	*	1.0	2.4	6.9	23.8
1999	0.5	*	*	*	*	*	*	*	*	2.2	6.2	17.1
Non-Hispanic black, male												
2002	0.6	*	*	*	*	*	*	*	*	4.1	9.7	24.1
2001	0.6	*	*	*	*	*	*	*	*	3.9	7.5	24.3
2000	0.7	*	*	*	*	*	*	*	*	3.9	7.7	28.2
1999	0.4	*	*	*	*	*	*	*	*	*	6.9	*
Non-Hispanic black, female												
2002	0.7	*	*	*	*	*	*	*	*	*	6.6	18.0
2001	0.7	*	*	*	*	*	*	*	*	2.8	5.7	24.4
2000	0.6	*	*	*	*	*	*	*	*	*	6.4	22.2
1999	0.6	*	*	*	*	*	*	*	*	2.1	5.8	15.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)												
All origins 3/, both sexes												
2002	20.2	4.6	*	0.1	0.3	0.9	2.1	5.7	15.8	49.7	153.5	469.4
2001	20.0	3.9	0.2	*	0.3	0.8	2.1	5.7	15.8	51.3	149.9	476.4
2000	19.4	5.0	0.1	0.1	0.3	0.8	2.0	5.4	15.6	48.0	148.2	471.4
1999	19.0	5.1	0.2	0.1	0.2	0.8	2.0	5.0	15.1	48.1	144.7	471.1
All origins 3/, male												
2002	18.1	5.6	*	*	0.2	0.8	2.1	6.5	17.0	55.9	178.6	553.0
2001	17.7	4.2	*	*	0.3	0.7	2.2	6.4	16.7	56.9	173.5	564.2
2000	17.2	6.1	*	*	0.3	0.8	2.2	6.3	16.8	52.6	173.2	567.5
1999	16.9	5.0	*	0.1	0.2	0.8	2.1	5.6	16.5	53.3	172.1	558.3
All origins 3/, female												
2002	22.3	3.6	*	*	0.3	0.9	2.0	5.0	14.8	44.5	136.8	433.2
2001	22.2	3.6	*	*	0.3	0.8	2.0	5.1	15.0	46.5	134.3	439.6
2000	21.5	3.9	*	*	0.3	0.8	1.8	4.6	14.5	44.3	132.0	432.3
1999	21.1	5.1	*	0.1	0.3	0.8	1.8	4.5	13.8	43.9	127.0	436.0
Hispanic, both sexes												
2002	6.6	4.7	*	*	*	0.6	1.4	4.9	15.7	45.9	120.8	327.6
2001	6.8	2.6	*	*	*	0.4	1.8	5.0	15.4	52.1	130.4	361.4
2000	6.5	3.8	*	*	*	0.5	1.2	4.5	16.5	47.0	125.8	372.9
1999	6.5	3.3	*	*	*	0.4	1.7	4.7	17.0	46.2	126.4	385.2
Hispanic, male												
2002	5.9	5.4	*	*	*	*	1.6	5.3	17.2	53.5	141.2	319.6
2001	5.9	*	*	*	*	*	2.0	4.9	17.6	54.3	140.4	400.0
2000	5.7	5.1	*	*	*	*	1.5	5.7	16.3	50.0	138.7	411.1
1999	6.0	*	*	*	*	*	1.6	5.2	21.2	50.1	147.6	455.1
Hispanic, female												
2002	7.3	*	*	*	*	0.8	1.3	4.4	14.4	39.8	106.8	331.6
2001	7.8	*	*	*	*	0.6	1.5	5.1	13.5	50.4	123.6	342.6
2000	7.3	*	*	*	*	0.7	1.0	3.3	16.8	44.6	117.2	354.1
1999	7.1	*	*	*	*	*	1.8	4.3	13.4	43.2	112.4	351.5
Non-Hispanic 4/, both sexes												
2002	22.3	4.6	*	0.1	0.3	0.9	2.2	5.8	15.8	49.8	154.7	474.0
2001	21.9	4.2	0.2	*	0.3	0.8	2.2	5.7	15.8	51.0	150.3	479.6
2000	21.2	5.2	*	0.1	0.3	0.8	2.1	5.5	15.4	48.0	148.8	474.3
1999	20.7	5.5	0.2	0.1	0.3	0.9	2.0	5.1	14.9	48.1	145.2	472.8
Non-Hispanic 4/, male												
2002	20.1	5.7	*	*	0.3	0.9	2.2	6.6	16.9	55.8	179.9	562.0
2001	19.5	4.3	*	*	0.3	0.8	2.3	6.4	16.6	56.9	174.6	569.8
2000	18.9	6.1	*	*	0.3	0.9	2.3	6.3	16.8	52.5	174.3	573.3
1999	18.4	5.1	*	*	0.2	0.8	2.2	5.6	16.1	53.3	172.8	560.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)												
Non-Hispanic 4/, female												
2002	24.4	3.5	*	*	0.3	1.0	2.1	5.1	14.8	44.8	138.0	436.0
2001	24.2	4.0	*	*	0.3	0.8	2.1	5.0	15.1	46.2	134.3	442.0
2000	23.4	4.3	*	*	0.3	0.8	1.9	4.7	14.2	44.2	132.2	434.2
1999	22.9	5.9	*	*	0.3	0.9	1.8	4.5	13.8	43.8	127.2	437.6
Non-Hispanic white, both sexes												
2002	22.7	3.6	*	*	0.2	0.6	1.5	4.0	12.0	42.9	146.3	462.7
2001	22.2	3.0	*	*	0.2	0.6	1.5	3.9	11.9	43.7	141.1	467.7
2000	21.4	3.4	*	*	0.3	0.6	1.4	3.8	11.7	40.6	140.0	462.6
1999	20.8	4.1	*	0.1	0.2	0.6	1.3	3.4	11.4	40.8	136.0	456.8
Non-Hispanic white, male												
2002	20.3	4.3	*	*	0.2	0.6	1.5	4.5	12.6	48.7	172.0	551.9
2001	19.8	3.2	*	*	0.2	0.5	1.7	4.4	12.7	49.7	166.6	561.9
2000	19.1	4.2	*	*	0.2	0.6	1.5	4.2	12.8	45.9	165.9	566.3
1999	18.3	4.3	*	*	0.2	0.6	1.4	3.7	12.5	45.9	162.5	543.9
Non-Hispanic white, female												
2002	25.0	2.8	*	*	0.2	0.6	1.4	3.5	11.5	37.9	129.0	424.2
2001	24.5	2.8	*	*	0.3	0.7	1.3	3.4	11.1	38.6	124.2	428.5
2000	23.7	2.6	*	*	0.3	0.6	1.3	3.4	10.6	36.1	123.0	420.8
1999	23.2	3.8	*	*	0.2	0.6	1.2	3.1	10.3	36.6	118.6	422.1
Non-Hispanic black, both sexes												
2002	26.0	8.7	*	*	0.8	2.6	6.4	18.7	47.9	119.2	270.9	667.5
2001	26.0	9.7	*	*	0.5	2.2	6.3	18.8	48.2	123.9	274.7	670.5
2000	25.2	13.2	*	*	0.5	2.2	6.4	18.2	47.0	120.4	270.1	645.2
1999	25.2	11.6	*	*	0.7	2.3	6.4	18.0	44.1	119.8	266.7	685.5
Non-Hispanic black, male												
2002	24.0	10.9	*	*	*	2.3	6.7	22.6	56.6	135.6	309.8	804.1
2001	23.0	9.6	*	*	*	2.7	5.9	22.3	52.7	137.9	304.3	764.6
2000	22.7	14.3	*	*	*	2.2	7.4	23.0	52.8	126.8	312.5	726.8
1999	23.2	8.7	*	*	*	2.0	7.4	21.8	48.5	134.1	330.2	812.9
Non-Hispanic black, female												
2002	27.9	6.3	*	*	1.0	2.8	6.2	15.4	40.9	107.6	248.9	615.4
2001	28.8	9.7	*	*	*	1.7	6.6	15.8	44.6	114.0	258.1	635.6
2000	27.6	12.0	*	*	*	2.1	5.4	14.1	42.3	116.0	246.3	614.9
1999	27.0	14.6	*	*	0.8	2.5	5.5	14.7	40.6	109.7	231.1	637.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of urinary system (N00-N39)												
All origins 3/, both sexes												
2002	19.9	4.6	*	0.1	0.3	0.8	2.0	5.6	15.7	49.1	151.2	460.8
2001	19.7	3.7	0.2	*	0.3	0.7	2.1	5.6	15.6	50.5	147.6	467.4
2000	19.1	4.8	0.1	0.1	0.3	0.7	2.0	5.3	15.4	47.4	145.9	462.5
1999	18.7	4.7	0.1	0.1	0.2	0.8	1.9	4.9	14.8	47.5	142.3	462.6
All origins 3/, male												
2002	17.7	5.6	*	*	0.2	0.8	2.1	6.4	16.8	55.0	174.6	533.8
2001	17.3	4.2	*	*	0.3	0.7	2.2	6.3	16.5	56.0	169.8	542.9
2000	16.8	6.0	*	*	0.3	0.8	2.2	6.2	16.5	51.7	169.2	546.9
1999	16.4	5.0	*	0.1	0.2	0.8	2.1	5.5	16.2	52.4	167.7	537.0
All origins 3/, female												
2002	22.0	3.5	*	*	0.3	0.9	1.9	4.9	14.6	44.2	135.7	429.1
2001	22.0	3.2	*	*	0.3	0.8	1.9	4.9	14.8	46.0	133.0	435.8
2000	21.2	3.6	*	*	0.3	0.7	1.7	4.4	14.3	43.8	130.6	428.1
1999	20.9	4.3	*	0.1	0.3	0.8	1.7	4.4	13.5	43.4	125.7	432.6
Hispanic, both sexes												
2002	6.4	4.6	*	*	*	0.6	1.4	4.8	15.5	45.1	118.3	323.7
2001	6.7	2.6	*	*	*	0.4	1.7	4.8	15.3	51.0	128.9	356.4
2000	6.3	3.8	*	*	*	0.5	1.2	4.4	16.2	45.9	124.0	364.3
1999	6.4	3.0	*	*	*	0.4	1.6	4.6	16.8	45.3	124.8	379.6
Hispanic, male												
2002	5.7	5.4	*	*	*	*	1.6	5.3	17.0	52.2	136.2	314.6
2001	5.8	*	*	*	*	*	2.0	4.8	17.5	52.9	138.2	394.2
2000	5.5	5.1	*	*	*	*	1.5	5.6	16.1	48.5	136.7	389.0
1999	5.9	*	*	*	*	*	1.6	5.2	20.8	48.6	145.5	448.6
Hispanic, female												
2002	7.2	*	*	*	*	0.7	1.2	4.3	14.2	39.5	106.0	328.3
2001	7.6	*	*	*	*	*	1.4	4.8	13.4	49.4	122.7	337.9
2000	7.2	*	*	*	*	*	0.8	3.2	16.3	43.8	115.6	352.2
1999	7.0	*	*	*	*	*	1.7	4.1	13.2	42.7	111.0	346.3
Non-Hispanic 4/, both sexes												
2002	21.9	4.5	*	0.1	0.3	0.9	2.1	5.7	15.6	49.2	152.4	465.2
2001	21.6	3.9	0.2	*	0.3	0.8	2.1	5.6	15.6	50.3	148.0	470.4
2000	20.8	5.0	*	0.1	0.3	0.8	2.0	5.4	15.3	47.3	146.4	465.4
1999	20.3	5.0	*	0.1	0.3	0.8	1.9	5.0	14.6	47.4	142.6	464.1
Non-Hispanic 4/, male												
2002	19.6	5.7	*	*	0.3	0.9	2.2	6.5	16.7	54.9	175.9	542.4
2001	19.0	4.2	*	*	0.3	0.8	2.2	6.4	16.4	56.0	170.9	547.9
2000	18.4	6.0	*	*	0.3	0.8	2.3	6.2	16.5	51.7	170.2	552.8
1999	17.9	5.1	*	*	0.2	0.8	2.1	5.5	15.8	52.4	168.3	538.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of urinary system (N00-N39)												
Non-Hispanic 4/, female												
2002	24.1	3.3	*	*	0.3	0.9	2.0	4.9	14.6	44.5	136.9	431.9
2001	23.9	3.6	*	*	0.3	0.8	2.0	4.9	14.9	45.6	133.0	438.2
2000	23.1	3.9	*	*	0.3	0.7	1.8	4.5	14.1	43.7	130.9	430.0
1999	22.7	5.0	*	*	0.3	0.8	1.7	4.4	13.5	43.4	126.0	434.3
Non-Hispanic white, both sexes												
2002	22.3	3.5	*	*	0.2	0.6	1.4	3.9	11.9	42.4	144.1	453.9
2001	21.8	2.8	*	*	0.2	0.6	1.5	3.8	11.7	43.0	138.8	458.5
2000	21.0	3.3	*	*	0.3	0.6	1.4	3.7	11.6	40.0	137.6	453.7
1999	20.4	3.8	*	0.1	0.2	0.6	1.3	3.3	11.2	40.3	133.5	448.1
Non-Hispanic white, male												
2002	19.8	4.3	*	*	0.2	0.6	1.5	4.4	12.5	48.0	168.3	532.0
2001	19.2	3.1	*	*	0.2	0.5	1.7	4.3	12.5	48.8	162.8	539.4
2000	18.6	4.1	*	*	0.2	0.6	1.5	4.2	12.6	45.2	161.9	545.9
1999	17.8	4.3	*	*	0.2	0.6	1.4	3.6	12.3	45.1	158.0	521.7
Non-Hispanic white, female												
2002	24.7	2.6	*	*	0.2	0.6	1.3	3.4	11.3	37.7	127.8	420.2
2001	24.2	2.5	*	*	0.2	0.7	1.3	3.3	10.9	38.1	122.9	424.8
2000	23.4	2.4	*	*	0.3	0.5	1.3	3.2	10.5	35.6	121.7	416.5
1999	23.0	3.2	*	*	0.2	0.6	1.1	3.0	10.1	36.3	117.4	418.8
Non-Hispanic black, both sexes												
2002	25.7	8.7	*	*	0.7	2.5	6.2	18.5	47.5	117.7	268.0	657.3
2001	25.7	9.3	*	*	0.5	2.2	6.0	18.4	47.5	123.1	271.1	661.1
2000	24.9	12.7	*	*	0.5	2.1	6.1	18.1	46.5	119.2	267.0	636.0
1999	24.8	10.4	*	*	0.6	2.2	6.3	17.6	43.4	118.1	262.7	675.4
Non-Hispanic black, male												
2002	23.6	10.9	*	*	*	2.3	6.6	22.5	56.1	133.2	303.0	785.5
2001	22.6	9.6	*	*	*	2.7	5.7	22.0	51.8	136.9	298.6	745.0
2000	22.3	14.3	*	*	*	2.2	7.4	22.9	51.8	125.2	306.7	705.7
1999	22.8	8.7	*	*	*	2.0	7.4	21.5	47.5	131.7	324.3	789.2
Non-Hispanic black, female												
2002	27.5	6.3	*	*	0.9	2.7	5.7	15.1	40.6	106.8	248.2	608.3
2001	28.4	9.1	*	*	*	1.7	6.3	15.2	44.0	113.3	255.7	630.1
2000	27.3	10.9	*	*	*	2.0	5.0	14.0	42.3	115.0	244.7	610.1
1999	26.6	12.1	*	*	*	2.3	5.2	14.3	40.2	108.5	228.3	632.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
All origins 3/, both sexes												
2002	0.7	*	*	*	*	0.1	0.1	0.3	0.5	1.5	4.9	13.9
2001	0.6	*	*	*	*	0.1	0.2	0.3	0.5	1.6	4.6	13.7
2000	0.7	0.5	*	*	*	0.1	0.1	0.3	0.6	1.5	4.8	15.7
1999	0.7	*	*	*	*	0.1	0.1	0.3	0.7	1.7	5.4	16.4
All origins 3/, male												
2002	0.6	*	*	*	*	*	0.1	0.3	0.6	1.7	5.5	18.1
2001	0.6	*	*	*	*	*	0.2	0.2	0.5	1.8	5.3	17.2
2000	0.6	*	*	*	*	*	0.1	0.3	0.6	1.4	5.4	20.0
1999	0.7	*	*	*	*	*	0.1	0.3	0.7	1.6	6.7	22.6
All origins 3/, female												
2002	0.7	*	*	*	*	*	0.1	0.3	0.5	1.4	4.5	12.0
2001	0.7	*	*	*	*	*	0.2	0.3	0.5	1.4	4.1	12.3
2000	0.8	*	*	*	*	*	0.1	0.3	0.6	1.7	4.5	13.9
1999	0.8	*	*	*	*	*	0.2	0.4	0.6	1.9	4.6	13.9
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	1.8	5.3	11.6
2001	0.3	*	*	*	*	*	*	*	*	*	4.1	*
2000	0.3	*	*	*	*	*	*	*	*	*	4.3	13.9
1999	0.3	*	*	*	*	*	*	*	*	2.0	5.0	14.7
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	*	*	0.1	0.1	0.3	0.6	1.5	4.8	13.9
2001	0.7	*	*	*	*	0.1	0.2	0.2	0.5	1.6	4.6	13.8
2000	0.7	*	*	*	*	0.1	0.1	0.3	0.6	1.5	4.8	15.7
1999	0.8	*	*	*	*	0.1	0.1	0.3	0.6	1.7	5.4	16.5
Non-Hispanic 4/, male												
2002	0.7	*	*	*	*	*	0.1	0.3	0.5	1.7	5.4	18.1
2001	0.6	*	*	*	*	*	0.1	0.2	0.4	1.8	5.4	17.2
2000	0.6	*	*	*	*	*	0.2	0.3	0.6	1.4	5.4	20.0
1999	0.7	*	*	*	*	*	0.1	0.3	0.6	1.6	6.7	22.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Glomerular and renal tubulo-interstitial diseases (N00-N15)												
Non-Hispanic 4/, female												
2002	0.8	*	*	*	*	*	0.1	0.3	0.6	1.3	4.5	12.1
2001	0.8	*	*	*	*	*	0.2	0.3	0.6	1.4	4.1	12.4
2000	0.8	*	*	*	*	*	0.1	0.3	0.6	1.6	4.5	14.0
1999	0.9	*	*	*	*	*	0.2	0.4	0.6	1.8	4.5	13.9
Non-Hispanic white, both sexes												
2002	0.8	*	*	*	*	*	0.1	0.2	0.5	1.6	4.8	13.9
2001	0.7	*	*	*	*	*	0.1	0.2	0.4	1.5	4.5	14.0
2000	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	4.6	15.3
1999	0.8	*	*	*	*	0.1	0.1	0.3	0.5	1.5	5.3	16.3
Non-Hispanic white, male												
2002	0.7	*	*	*	*	*	0.1	0.2	0.5	1.7	5.2	18.4
2001	0.7	*	*	*	*	*	0.1	0.1	0.4	1.7	5.2	17.5
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.3	5.3	19.4
1999	0.7	*	*	*	*	*	*	0.3	0.5	1.3	6.6	22.2
Non-Hispanic white, female												
2002	0.8	*	*	*	*	*	0.1	0.2	0.5	1.5	4.5	11.9
2001	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	4.1	12.5
2000	0.9	*	*	*	*	*	*	0.2	0.6	1.6	4.2	13.6
1999	0.9	*	*	*	*	*	*	0.3	0.5	1.7	4.5	14.0
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	*	0.6	0.9	1.3	5.5	15.7
2001	0.6	*	*	*	*	*	*	0.6	0.9	2.5	5.9	12.5
2000	0.8	*	*	*	*	*	0.4	0.9	1.0	2.7	7.2	18.4
1999	0.7	*	*	*	*	*	*	0.6	1.3	3.4	5.9	20.0
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	7.6	*
2001	0.6	*	*	*	*	*	*	*	*	3.9	6.6	*
2000	0.7	*	*	*	*	*	*	1.1	*	3.0	7.1	24.7
1999	0.8	*	*	*	*	*	*	*	*	4.1	7.5	35.7
Non-Hispanic black, female												
2002	0.6	*	*	*	*	*	*	*	*	*	4.3	14.6
2001	0.6	*	*	*	*	*	*	*	*	*	5.6	12.0
2000	0.8	*	*	*	*	*	*	*	*	2.4	7.3	16.1
1999	0.7	*	*	*	*	*	*	*	*	2.9	5.1	14.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Glomerular diseases (N00-N07)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.6	1.7	3.3
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.7	1.5	3.3
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	3.6
1999	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	2.0	3.7
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.9	4.3
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.9	2.0	4.6
2000	0.3	*	*	*	*	*	*	0.2	0.3	0.8	2.0	5.6
1999	0.3	*	*	*	*	*	*	0.2	0.3	0.7	2.6	5.6
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	2.8
2001	0.2	*	*	*	*	*	*	*	*	0.6	1.1	2.8
2000	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	2.8
1999	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.5	2.9
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.7	3.3
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.7	1.5	3.3
2000	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	3.6
1999	0.3	*	*	*	*	*	0.1	0.1	0.3	0.7	2.0	3.7
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.9	4.4
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.9	2.0	4.7
2000	0.3	*	*	*	*	*	*	0.2	0.3	0.8	2.1	5.4
1999	0.3	*	*	*	*	*	*	0.2	0.3	0.7	2.7	5.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Glomerular diseases (N00-N07)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	2.8
2001	0.2	*	*	*	*	*	*	*	*	0.6	1.2	2.7
2000	0.3	*	*	*	*	*	*	*	0.2	0.6	1.4	2.9
1999	0.3	*	*	*	*	*	*	0.1	0.3	0.7	1.5	2.9
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.6	2.9
2001	0.2	*	*	*	*	*	*	*	0.2	0.7	1.4	3.3
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.6	3.2
1999	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.9	3.4
Non-Hispanic white, male												
2002	0.3	*	*	*	*	*	*	*	0.3	0.7	1.7	4.2
2001	0.2	*	*	*	*	*	*	*	*	0.8	2.0	4.8
2000	0.3	*	*	*	*	*	*	*	0.3	0.7	2.1	4.5
1999	0.3	*	*	*	*	*	*	0.2	0.2	0.6	2.6	5.1
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.4	2.4
2001	0.2	*	*	*	*	*	*	*	*	0.6	1.1	2.6
2000	0.2	*	*	*	*	*	*	*	*	0.6	1.4	2.7
1999	0.3	*	*	*	*	*	*	*	0.2	0.6	1.4	2.7
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	0.4	*	*	2.5	6.7
2001	0.2	*	*	*	*	*	*	*	*	1.2	*	*
2000	0.3	*	*	*	*	*	*	0.5	*	*	2.4	*
1999	0.3	*	*	*	*	*	*	*	*	*	2.6	7.1
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic syndrome (N00-N01)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic syndrome (N00-N01)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephrotic syndrome (N04)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	0.1	0.4	0.7
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.7
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.0
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.7
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.0
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.5	0.7
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephrotic syndrome (N04)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	0.6
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.6
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified glomerular diseases (N02-N03,N05-N07)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.3	2.4
2001	0.2	*	*	*	*	*	0.1	0.1	0.1	0.6	1.1	2.5
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.5
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.5	2.9
All origins 3/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	3.3
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.6	3.5
2000	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.5	3.8
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.4	2.0	4.4
All origins 3/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.1	1.9
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.1
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	1.9
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.3
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.3	2.3
2001	0.2	*	*	*	*	*	0.1	0.1	0.1	0.6	1.1	2.5
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.4
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.5	2.9
Non-Hispanic 4/, male												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	3.3
2001	0.2	*	*	*	*	*	*	0.1	*	0.8	1.6	3.4
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	3.6
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.4	2.1	4.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified glomerular diseases (N02-N03,N05-N07)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.3	1.1	1.9
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.0
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.0
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.2
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	0.1	0.1	0.1	0.5	1.2	2.1
2001	0.2	*	*	*	*	*	*	*	0.1	0.6	1.1	2.5
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.1
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	2.6
Non-Hispanic white, male												
2002	0.2	*	*	*	*	*	*	*	*	0.6	1.4	3.2
2001	0.2	*	*	*	*	*	*	*	*	0.8	1.6	3.7
2000	0.2	*	*	*	*	*	*	*	*	0.6	1.5	2.9
1999	0.2	*	*	*	*	*	*	*	*	0.4	2.0	4.3
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.7
2001	0.2	*	*	*	*	*	*	*	*	0.4	0.7	2.0
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.7
1999	0.2	*	*	*	*	*	*	*	*	0.5	1.0	2.0
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal tubulo-interstitial diseases (N10-N15)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.1	0.3	1.0	3.2	10.6
2001	0.4	*	*	*	*	*	0.1	0.1	0.3	0.8	3.1	10.4
2000	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	3.2	12.0
1999	0.5	*	*	*	*	*	0.1	0.2	0.4	1.0	3.4	12.7
All origins 3/, male												
2002	0.4	*	*	*	*	*	*	0.1	0.3	1.1	3.5	13.8
2001	0.3	*	*	*	*	*	*	*	0.2	0.9	3.3	12.5
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.7	3.3	14.4
1999	0.4	*	*	*	*	*	*	0.2	0.4	0.9	4.0	17.0
All origins 3/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.3	0.9	3.0	9.2
2001	0.5	*	*	*	*	*	0.1	0.2	0.4	0.7	3.0	9.5
2000	0.5	*	*	*	*	*	*	0.2	0.4	1.0	3.0	11.1
1999	0.6	*	*	*	*	*	0.1	0.2	0.3	1.1	3.0	11.0
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	3.5	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	4.2	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.1	0.3	0.9	3.2	10.7
2001	0.5	*	*	*	*	*	0.1	0.1	0.3	0.8	3.1	10.5
2000	0.5	*	*	*	*	*	0.1	0.2	0.3	0.8	3.1	12.1
1999	0.5	*	*	*	*	*	0.1	0.2	0.3	1.0	3.4	12.7
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	*	0.1	0.3	1.0	3.5	13.8
2001	0.4	*	*	*	*	*	*	*	0.2	0.9	3.4	12.5
2000	0.4	*	*	*	*	*	*	*	0.2	0.6	3.3	14.7
1999	0.4	*	*	*	*	*	*	0.1	0.3	0.9	4.0	17.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal tubulo-interstitial diseases (N10-N15)												
Non-Hispanic 4/, female												
2002	0.5	*	*	*	*	*	*	0.2	0.3	0.9	3.0	9.3
2001	0.6	*	*	*	*	*	0.1	0.2	0.4	0.7	3.0	9.7
2000	0.6	*	*	*	*	*	*	0.2	0.4	1.0	3.0	11.1
1999	0.6	*	*	*	*	*	*	0.2	0.3	1.1	3.0	11.0
Non-Hispanic white, both sexes												
2002	0.5	*	*	*	*	*	0.1	0.1	0.3	1.0	3.2	11.0
2001	0.5	*	*	*	*	*	0.1	0.1	0.3	0.8	3.1	10.7
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.8	3.0	12.0
1999	0.6	*	*	*	*	*	*	0.2	0.3	0.9	3.4	12.9
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	*	*	0.3	1.0	3.4	14.2
2001	0.4	*	*	*	*	*	*	*	*	0.8	3.2	12.7
2000	0.4	*	*	*	*	*	*	*	0.2	0.5	3.2	14.9
1999	0.5	*	*	*	*	*	*	*	0.3	0.7	4.0	17.1
Non-Hispanic white, female												
2002	0.6	*	*	*	*	*	*	*	0.3	1.0	3.1	9.5
2001	0.6	*	*	*	*	*	*	0.2	0.4	0.8	3.0	9.9
2000	0.6	*	*	*	*	*	*	0.2	0.4	1.0	2.8	10.9
1999	0.7	*	*	*	*	*	*	0.2	0.3	1.1	3.1	11.2
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	*	3.0	9.1
2001	0.4	*	*	*	*	*	*	*	*	1.3	4.0	9.1
2000	0.4	*	*	*	*	*	*	*	*	1.5	4.8	13.0
1999	0.4	*	*	*	*	*	*	*	*	2.2	3.4	12.9
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	3.0	*	23.8
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	3.4	*
2000	0.5	*	*	*	*	*	*	*	*	*	4.7	12.2
1999	0.4	*	*	*	*	*	*	*	*	*	*	8.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
All origins 3/, both sexes												
2002	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.8	39.4	102.6	284.0
2000	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
All origins 3/, male												
2002	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
All origins 3/, female												
2002	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5
Hispanic, both sexes												
2002	4.8	4.2	*	*	*	0.4	1.1	4.2	13.2	35.7	86.9	198.4
2001	4.9	*	*	*	*	0.3	1.2	3.7	12.5	40.6	93.4	217.6
2000	4.5	2.9	*	*	*	0.4	1.0	3.3	13.3	36.9	83.0	225.6
1999	4.5	*	*	*	*	*	1.2	3.5	13.6	34.4	86.9	228.0
Hispanic, male												
2002	4.6	4.9	*	*	*	*	1.3	4.8	14.4	40.7	109.2	232.6
2001	4.6	*	*	*	*	*	1.4	3.7	14.7	42.7	110.9	278.1
2000	4.3	*	*	*	*	*	1.3	4.2	13.8	40.5	99.3	280.1
1999	4.5	*	*	*	*	*	1.1	4.3	17.2	38.2	109.0	330.0
Hispanic, female												
2002	5.0	*	*	*	*	0.6	0.8	3.6	12.0	31.8	71.7	181.5
2001	5.2	*	*	*	*	*	1.0	3.7	10.5	38.9	81.6	188.0
2000	4.8	*	*	*	*	*	*	2.4	12.8	34.0	72.0	198.8
1999	4.5	*	*	*	*	*	1.3	2.7	10.4	31.3	72.2	178.8
Non-Hispanic 4/, both sexes												
2002	15.3	4.1	*	*	0.2	0.7	1.7	4.6	12.6	38.6	107.9	287.4
2001	14.9	3.3	*	*	0.2	0.6	1.7	4.6	12.7	39.2	102.7	285.9
2000	14.1	4.2	*	*	0.2	0.6	1.6	4.3	12.3	37.1	99.3	274.9
1999	13.5	4.7	*	*	0.2	0.6	1.5	3.9	11.5	36.4	95.4	265.0
Non-Hispanic 4/, male												
2002	15.1	5.1	*	*	0.2	0.6	1.8	5.4	13.9	45.1	134.7	386.2
2001	14.6	3.5	*	*	0.2	0.6	1.8	5.4	13.7	45.3	128.5	392.7
2000	13.8	5.2	*	*	0.2	0.6	1.9	5.1	13.7	42.0	125.9	377.7
1999	13.2	4.6	*	*	0.1	0.6	1.8	4.5	12.7	41.5	122.6	364.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Non-Hispanic 4/, female												
2002	15.5	2.9	*	*	0.2	0.7	1.5	3.8	11.4	33.2	90.2	244.8
2001	15.2	3.1	*	*	0.2	0.5	1.6	3.7	11.8	34.1	85.8	241.5
2000	14.4	3.1	*	*	0.2	0.5	1.4	3.5	11.0	33.2	82.0	233.4
1999	13.8	4.8	*	*	0.2	0.6	1.3	3.3	10.4	32.1	77.8	225.2
Non-Hispanic white, both sexes												
2002	15.2	3.1	*	*	0.1	0.4	1.1	3.0	9.2	32.3	100.8	277.6
2001	14.6	2.2	*	*	0.2	0.4	1.1	3.0	9.2	32.7	95.1	276.1
2000	13.9	2.5	*	*	0.2	0.4	1.1	2.8	9.0	30.7	92.3	265.7
1999	13.1	3.5	*	*	0.1	0.4	0.9	2.4	8.4	30.4	87.8	251.6
Non-Hispanic white, male												
2002	15.0	3.9	*	*	*	0.4	1.2	3.5	10.0	38.8	128.8	377.8
2001	14.5	2.3	*	*	*	0.3	1.3	3.6	10.3	39.1	121.7	387.6
2000	13.7	3.3	*	*	0.2	0.4	1.2	3.3	10.3	36.5	119.4	373.4
1999	13.0	3.8	*	*	*	0.4	1.1	2.8	9.6	35.9	114.4	353.8
Non-Hispanic white, female												
2002	15.3	2.2	*	*	*	0.4	1.0	2.5	8.5	26.7	82.0	234.3
2001	14.7	2.1	*	*	0.2	0.4	0.9	2.4	8.1	27.3	77.5	229.7
2000	14.0	*	*	*	*	0.3	0.9	2.3	7.7	25.8	74.4	222.2
1999	13.2	3.1	*	*	*	0.4	0.8	2.1	7.3	25.8	70.3	210.8
Non-Hispanic black, both sexes												
2002	20.2	8.0	*	*	0.6	2.0	5.4	16.0	41.4	100.9	202.7	436.5
2001	20.0	8.5	*	*	0.4	1.7	5.3	15.8	42.1	102.9	201.9	428.9
2000	19.2	11.4	*	*	0.4	1.8	5.3	15.6	40.7	99.9	194.6	404.2
1999	18.8	10.2	*	*	0.5	1.9	5.4	15.2	37.0	95.3	193.0	428.7
Non-Hispanic black, male												
2002	19.4	10.0	*	*	*	1.8	6.1	20.1	49.8	115.4	233.3	562.3
2001	18.4	8.9	*	*	*	2.1	5.1	19.3	45.7	114.3	233.5	514.0
2000	17.9	13.6	*	*	*	1.9	6.5	19.9	45.0	104.2	231.7	487.3
1999	17.7	8.7	*	*	*	1.8	6.4	19.0	40.5	102.5	246.7	519.4
Non-Hispanic black, female												
2002	20.9	*	*	*	*	2.1	4.8	12.5	34.7	90.6	185.4	388.4
2001	21.4	8.1	*	*	*	1.4	5.5	12.8	39.2	94.9	184.1	397.4
2000	20.3	9.2	*	*	*	1.7	4.1	11.8	37.3	96.9	173.7	373.3
1999	19.8	11.8	*	*	*	2.0	4.5	12.0	34.3	90.2	163.0	394.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Urolithiasis (N20-N23)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.7	1.8
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.7	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.8
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.6	1.9
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.0
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.7
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.7	1.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.8
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	2.0
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.7	1.8
2001	0.1	*	*	*	*	*	*	*	0.1	0.2	0.7	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.8
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.8
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.0
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Urolithiasis (N20-N23)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.7	1.8
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.7	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.8
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	1.9
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.9
2001	0.1	*	*	*	*	*	*	*	0.1	0.2	0.7	1.9
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.8
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.6	1.7
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	2.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.7	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	1.9
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified disorders of kidney (N25-N27)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified disorders of kidney (N25-N27)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.6
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Urinary tract infection, site not specified (N39.0)												
All origins 3/, both sexes												
2002	4.8	*	*	*	*	0.1	0.2	0.5	1.9	7.7	35.6	152.2
2001	4.9	*	*	*	*	0.1	0.2	0.5	1.8	8.2	37.0	160.4
2000	4.9	*	*	*	*	0.1	0.1	0.5	1.7	7.3	38.4	163.4
1999	5.0	*	*	*	*	0.1	0.2	0.5	1.8	8.0	38.1	172.0
All origins 3/, male												
2002	3.0	*	*	*	*	*	0.2	0.4	1.6	6.7	31.0	123.5
2001	3.0	*	*	*	*	*	0.1	0.4	1.6	7.4	32.6	125.8
2000	3.1	*	*	*	*	*	0.1	0.5	1.5	6.7	34.3	139.8
1999	3.1	*	*	*	*	*	0.1	0.4	1.8	7.5	34.7	137.5
All origins 3/, female												
2002	6.5	*	*	*	*	*	0.2	0.6	2.1	8.5	38.7	164.7
2001	6.7	*	*	*	*	*	0.2	0.6	2.0	8.9	39.8	174.9
2000	6.6	*	*	*	*	*	0.1	0.6	1.9	7.8	41.1	173.0
1999	6.9	*	*	*	*	*	0.2	0.6	1.9	8.4	40.3	185.9
Hispanic, both sexes												
2002	1.2	*	*	*	*	*	*	*	1.5	6.3	24.0	104.2
2001	1.3	*	*	*	*	*	*	*	2.1	7.0	28.5	115.7
2000	1.3	*	*	*	*	*	*	*	1.8	6.5	32.4	120.1
1999	1.3	*	*	*	*	*	*	*	1.5	7.7	29.8	124.9
Hispanic, male												
2002	0.7	*	*	*	*	*	*	*	*	8.0	17.2	51.9
2001	0.8	*	*	*	*	*	*	*	*	6.2	20.1	83.8
2000	0.8	*	*	*	*	*	*	*	*	5.5	27.5	84.6
1999	0.9	*	*	*	*	*	*	*	*	7.2	28.2	92.7
Hispanic, female												
2002	1.7	*	*	*	*	*	*	*	*	4.9	28.7	130.1
2001	1.9	*	*	*	*	*	*	*	2.1	7.6	34.2	131.2
2000	1.9	*	*	*	*	*	*	*	2.5	7.3	35.7	137.5
1999	1.8	*	*	*	*	*	*	*	*	8.1	30.9	140.4
Non-Hispanic 4/, both sexes												
2002	5.3	*	*	*	*	*	0.2	0.5	1.9	7.8	36.1	154.0
2001	5.4	*	*	*	*	0.1	0.2	0.5	1.8	8.3	37.2	161.6
2000	5.4	*	*	*	*	0.1	0.1	0.5	1.7	7.3	38.6	164.8
1999	5.5	*	*	*	*	0.1	0.2	0.5	1.9	8.0	38.4	173.4
Non-Hispanic 4/, male												
2002	3.3	*	*	*	*	*	0.1	0.5	1.7	6.6	31.6	126.6
2001	3.3	*	*	*	*	*	0.1	0.4	1.5	7.5	33.0	127.1
2000	3.4	*	*	*	*	*	0.1	0.5	1.6	6.8	34.5	141.8
1999	3.4	*	*	*	*	*	0.1	0.4	1.8	7.5	34.8	139.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Urinary tract infection, site not specified (N39.0)												
Non-Hispanic 4/, female												
2002	7.2	*	*	*	*	*	0.2	0.6	2.1	8.7	39.1	165.8
2001	7.4	*	*	*	*	*	0.2	0.6	2.0	9.0	40.0	175.9
2000	7.2	*	*	*	*	*	0.2	0.6	1.9	7.8	41.2	174.1
1999	7.5	*	*	*	*	*	0.2	0.6	1.9	8.4	40.7	187.1
Non-Hispanic white, both sexes												
2002	5.8	*	*	*	*	*	0.1	0.5	1.7	7.2	35.0	152.5
2001	5.9	*	*	*	*	0.1	0.2	0.4	1.6	7.6	35.8	159.2
2000	5.8	*	*	*	*	0.1	0.1	0.5	1.5	6.7	37.2	163.0
1999	6.0	*	*	*	*	0.1	0.1	0.4	1.6	7.2	37.2	171.0
Non-Hispanic white, male												
2002	3.6	*	*	*	*	*	0.1	0.4	1.5	6.0	30.3	124.3
2001	3.5	*	*	*	*	*	0.2	0.3	1.3	6.7	32.0	123.2
2000	3.6	*	*	*	*	*	*	0.4	1.3	6.1	32.8	139.7
1999	3.6	*	*	*	*	*	*	0.4	1.6	6.3	33.1	133.4
Non-Hispanic white, female												
2002	7.9	*	*	*	*	*	0.1	0.5	1.9	8.3	38.3	164.7
2001	8.1	*	*	*	*	*	0.2	0.5	1.8	8.4	38.3	174.2
2000	7.9	*	*	*	*	*	0.1	0.5	1.7	7.2	40.0	172.5
1999	8.2	*	*	*	*	*	0.2	0.5	1.7	7.9	40.0	186.0
Non-Hispanic black, both sexes												
2002	4.3	*	*	*	*	*	0.4	1.0	3.8	13.5	54.5	193.9
2001	4.5	*	*	*	*	*	*	1.2	3.4	14.6	58.9	210.7
2000	4.4	*	*	*	*	*	*	1.1	3.8	13.8	60.3	200.0
1999	4.6	*	*	*	*	*	0.4	1.2	3.8	16.6	57.1	215.3
Non-Hispanic black, male												
2002	3.0	*	*	*	*	*	*	*	3.6	13.8	55.4	190.3
2001	3.1	*	*	*	*	*	*	1.0	3.7	15.0	53.7	206.7
2000	3.2	*	*	*	*	*	*	1.4	4.3	14.4	62.3	178.5
1999	3.6	*	*	*	*	*	*	*	4.1	20.7	62.5	218.7
Non-Hispanic black, female												
2002	5.4	*	*	*	*	*	*	1.2	3.9	13.3	54.1	195.2
2001	5.8	*	*	*	*	*	*	1.4	3.1	14.3	61.8	212.2
2000	5.5	*	*	*	*	*	*	*	3.3	13.4	59.2	208.0
1999	5.5	*	*	*	*	*	*	1.5	3.6	13.7	54.1	214.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of urinary system (N28-N36,N39.1-N39.9)												
All origins 3/, both sexes												
2002	0.4	*	*	*	*	0.1	0.1	0.2	0.4	1.0	2.6	7.5
2001	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.7	7.1
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.8	7.5
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.6	7.1
All origins 3/, male												
2002	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	3.2	9.0
2001	0.4	*	*	*	*	*	0.1	0.2	0.5	1.2	3.2	8.9
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.3	3.5	10.4
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.5	3.2	9.9
All origins 3/, female												
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.2	6.9
2001	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.3	6.3
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	0.8	2.2	6.3
1999	0.4	*	*	*	*	*	*	0.1	0.4	0.7	2.2	6.0
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.4	1.0	2.7	7.5
2001	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.7	6.9
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.0	2.7	7.7
1999	0.4	*	*	*	*	*	0.1	0.2	0.5	1.1	2.6	7.1
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	*	0.3	0.5	1.1	3.3	9.0
2001	0.4	*	*	*	*	*	0.1	0.3	0.5	1.2	3.2	8.7
2000	0.4	*	*	*	*	*	0.1	0.2	0.5	1.3	3.5	10.6
1999	0.4	*	*	*	*	*	*	0.2	0.5	1.5	3.2	10.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of urinary system (N28-N36,N39.1-N39.9)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.9	2.3	6.8
2001	0.4	*	*	*	*	*	*	0.2	0.4	0.9	2.3	6.2
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	0.8	2.2	6.5
1999	0.4	*	*	*	*	*	*	0.1	0.4	0.7	2.2	5.9
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	*	*	0.1	0.2	0.3	1.0	2.6	7.4
2001	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.6	7.0
2000	0.4	*	*	*	*	*	0.1	0.2	0.4	0.9	2.6	7.5
1999	0.4	*	*	*	*	*	0.1	0.1	0.4	0.9	2.4	7.2
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	*	0.3	0.4	1.1	3.2	8.8
2001	0.4	*	*	*	*	*	*	0.2	0.5	1.0	3.3	8.8
2000	0.5	*	*	*	*	*	0.1	0.2	0.4	1.1	3.4	10.5
1999	0.4	*	*	*	*	*	*	0.2	0.4	1.3	3.1	10.3
Non-Hispanic white, female												
2002	0.4	*	*	*	*	*	*	*	0.2	0.8	2.2	6.8
2001	0.4	*	*	*	*	*	*	0.1	0.4	0.7	2.2	6.2
2000	0.4	*	*	*	*	*	*	0.2	0.4	0.7	2.1	6.3
1999	0.4	*	*	*	*	*	*	*	0.4	0.7	2.0	5.9
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	*	0.7	1.3	1.8	4.2	8.8
2001	0.5	*	*	*	*	*	*	0.6	1.1	2.6	3.6	7.2
2000	0.5	*	*	*	*	*	*	*	0.9	2.5	4.0	11.4
1999	0.5	*	*	*	*	*	*	*	1.1	2.3	5.5	7.8
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	*	*	*	3.1	*	*
2000	0.5	*	*	*	*	*	*	*	*	3.0	*	*
1999	0.6	*	*	*	*	*	*	*	*	3.8	*	*
Non-Hispanic black, female												
2002	0.5	*	*	*	*	*	*	*	*	*	3.6	*
2001	0.5	*	*	*	*	*	*	*	*	2.2	*	*
2000	0.5	*	*	*	*	*	*	*	*	2.1	*	10.0
1999	0.4	*	*	*	*	*	*	*	*	*	5.3	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of male genital organs (N40-N50)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.8
2001	0.2	*	*	*	*	*	*	*	0.1	0.4	1.5	6.3
2000	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.9
1999	0.2	*	*	*	*	*	*	*	0.1	0.4	1.7	6.1
All origins 3/, male												
2002	0.4	*	*	*	*	*	*	*	0.2	0.9	4.0	19.2
2001	0.4	*	*	*	*	*	*	*	0.2	0.9	3.7	21.2
2000	0.4	*	*	*	*	*	*	*	0.3	0.9	3.9	20.5
1999	0.4	*	*	*	*	*	*	*	0.3	1.0	4.3	21.2
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.9
2001	0.2	*	*	*	*	*	*	*	0.1	0.4	1.5	6.4
2000	0.2	*	*	*	*	*	*	*	0.1	0.4	1.6	5.9
1999	0.2	*	*	*	*	*	*	*	0.1	0.4	1.7	6.2
Non-Hispanic 4/, male												
2002	0.4	*	*	*	*	*	*	*	0.2	0.8	3.9	19.6
2001	0.5	*	*	*	*	*	*	*	0.2	0.9	3.8	21.9
2000	0.4	*	*	*	*	*	*	*	0.3	0.8	4.0	20.5
1999	0.5	*	*	*	*	*	*	*	0.3	0.9	4.4	21.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of male genital organs (N40-N50)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	0.3	1.5	6.0
2001	0.3	*	*	*	*	*	*	*	*	0.4	1.5	6.6
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.6	5.8
1999	0.3	*	*	*	*	*	*	*	*	0.4	1.8	6.3
Non-Hispanic white, male												
2002	0.5	*	*	*	*	*	*	*	*	0.7	3.7	19.9
2001	0.5	*	*	*	*	*	*	*	*	0.9	3.7	22.5
2000	0.5	*	*	*	*	*	*	*	*	0.8	3.9	20.3
1999	0.5	*	*	*	*	*	*	*	*	0.8	4.4	22.0
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	6.5
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	6.7	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	23.8
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
All origins 3/, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.8
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.5
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.0
All origins 3/, male												
2002	0.3	*	*	*	*	*	*	*	*	0.5	3.2	15.8
2001	0.3	*	*	*	*	*	*	*	*	0.5	2.8	18.6
2000	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999	0.3	*	*	*	*	*	*	*	*	0.6	3.3	17.5
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.9
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.1
Non-Hispanic 4/, male												
2002	0.3	*	*	*	*	*	*	*	*	0.5	3.2	16.2
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.8	19.3
2000	0.3	*	*	*	*	*	*	*	*	0.4	3.3	18.0
1999	0.3	*	*	*	*	*	*	*	*	0.5	3.4	18.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.0
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.9
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.2	1.4	5.2
Non-Hispanic white, male												
2002	0.4	*	*	*	*	*	*	*	*	0.4	3.2	16.6
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.8	20.1
2000	0.4	*	*	*	*	*	*	*	*	0.4	3.4	18.0
1999	0.4	*	*	*	*	*	*	*	*	0.5	3.4	18.4
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of male genital organs (N41-N50)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	0.2	0.3	1.0
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.8
2000	0.0	*	*	*	*	*	*	*	0.1	0.2	0.3	0.7
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.1
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.4	0.7	3.4
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.6
2000	0.1	*	*	*	*	*	*	*	0.2	0.4	0.7	2.4
1999	0.1	*	*	*	*	*	*	*	*	0.4	1.0	3.7
All origins 3/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.3	1.0
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.8
2000	0.0	*	*	*	*	*	*	*	*	0.2	0.3	0.7
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.1
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.4
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.6
2000	0.1	*	*	*	*	*	*	*	*	0.4	0.7	2.5
1999	0.1	*	*	*	*	*	*	*	*	0.4	1.1	3.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of male genital organs (N41-N50)												
Non-Hispanic 4/, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.0
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
2000	0.0	*	*	*	*	*	*	*	*	0.2	0.2	0.7
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.0
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	3.3
2001	0.1	*	*	*	*	*	*	*	*	0.4	0.9	2.4
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.6	2.4
1999	0.1	*	*	*	*	*	*	*	*	0.3	1.0	3.7
Non-Hispanic white, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of breast (N60-N64)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.6
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.7
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.5
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.6
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of breast (N60-N64)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	0.7
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.6
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of female genital organs (N70-N98)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	*	0.1	0.6	2.4
2001	0.1	*	*	*	*	*	0.0	0.1	0.1	0.3	0.7	2.1
2000	0.1	*	*	*	*	*	0.1	0.1	*	0.3	0.7	2.5
1999	0.1	*	*	*	*	*	0.0	0.1	0.1	0.2	0.7	2.0
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.2	*	*	*	*	*	0.1	0.2	*	0.3	1.0	3.4
2001	0.2	*	*	*	*	*	0.1	0.2	0.2	0.5	1.1	3.0
2000	0.2	*	*	*	*	*	0.1	0.2	*	0.5	1.2	3.6
1999	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	2.9
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	*	0.1	0.6	2.4
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.7	2.1
2000	0.1	*	*	*	*	*	0.1	0.1	*	0.2	0.7	2.6
1999	0.1	*	*	*	*	*	0.0	*	0.1	0.2	0.7	2.0
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders of female genital organs (N70-N98)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	*	*	0.1	0.2	*	0.3	1.0	3.4
2001	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.1	2.9
2000	0.3	*	*	*	*	*	0.1	0.2	*	0.5	1.2	3.6
1999	0.2	*	*	*	*	*	0.1	*	0.2	0.4	1.1	2.8
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	2.3
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.7	2.0
2000	0.1	*	*	*	*	*	*	0.1	*	0.2	0.7	2.6
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.6	2.0
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.2	*	*	*	*	*	*	*	*	*	1.1	3.3
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.1	2.8
2000	0.3	*	*	*	*	*	*	0.2	*	0.4	1.2	3.7
1999	0.2	*	*	*	*	*	*	*	0.2	0.3	1.0	2.7
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Noninflammatory disorders of female genital tract (N80-N98)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	*	*	0.4	1.6
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.4	1.2
2000	0.1	*	*	*	*	*	*	0.1	*	0.1	0.5	1.9
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.6
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	0.1	*	*	0.7	2.3
2001	0.1	*	*	*	*	*	*	0.1	*	0.3	0.7	1.8
2000	0.2	*	*	*	*	*	*	0.1	*	0.3	0.8	2.7
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.7	2.3
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.6
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
2000	0.1	*	*	*	*	*	*	0.1	*	0.1	0.5	1.9
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.6
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Noninflammatory disorders of female genital tract (N80-N98)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	2.2
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.7
2000	0.2	*	*	*	*	*	*	0.1	*	0.3	0.8	2.7
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.7	2.2
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.5
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.1
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.5	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	1.5
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.7	2.1
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.6
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.8	2.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	2.2
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XV. Pregnancy, childbirth and the puerperium (O00-O99)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	*	*	*	0.3	0.5	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.5	0.3	*	*	*	*	*
1999	0.1	*	*	*	0.3	0.5	0.3	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.3	*	*	*	0.5	0.9	0.5	*	*	*	*	*
2001	0.3	*	*	*	0.6	1.0	0.5	*	*	*	*	*
2000	0.3	*	*	*	0.5	0.9	0.5	*	*	*	*	*
1999	0.3	*	*	*	0.5	0.9	0.5	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	*	0.4	0.4	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.6	*	*	*	*	*	*
2000	0.2	*	*	*	0.4	0.6	0.4	*	*	*	*	*
1999	0.2	*	*	*	*	0.4	0.5	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	0.4	*	*	*	*	0.8	0.8	*	*	*	*	*
2001	0.5	*	*	*	0.8	1.2	*	*	*	*	*	*
2000	0.5	*	*	*	0.8	1.2	0.9	*	*	*	*	*
1999	0.4	*	*	*	*	0.9	1.0	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	*	*	*	0.3	0.5	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.4	0.2	*	*	*	*	*
1999	0.1	*	*	*	0.2	0.5	0.2	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XV. Pregnancy, childbirth and the puerperium (O00-O99)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	0.4	0.9	0.4	*	*	*	*	*
2001	0.3	*	*	*	0.5	0.9	0.4	*	*	*	*	*
2000	0.3	*	*	*	0.4	0.9	0.5	*	*	*	*	*
1999	0.3	*	*	*	0.5	0.9	0.5	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.1	0.3	0.1	*	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.1	0.3	0.2	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.3	0.1	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.1	*	*	*	0.2	0.6	0.3	*	*	*	*	*
2001	0.2	*	*	*	0.3	0.6	0.2	*	*	*	*	*
2000	0.2	*	*	*	0.3	0.6	0.3	*	*	*	*	*
1999	0.2	*	*	*	0.2	0.6	0.3	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	0.8	1.2	0.7	*	*	*	*	*
2001	0.4	*	*	*	0.8	1.3	0.7	*	*	*	*	*
2000	0.4	*	*	*	0.6	1.3	0.7	*	*	*	*	*
1999	0.5	*	*	*	0.8	1.3	0.7	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.8	*	*	*	1.5	2.4	1.4	*	*	*	*	*
2001	0.8	*	*	*	1.6	2.4	1.4	*	*	*	*	*
2000	0.8	*	*	*	1.2	2.4	1.3	*	*	*	*	*
1999	0.9	*	*	*	1.6	2.5	1.4	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ectopic pregnancy (O00)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ectopic pregnancy (O00)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Spontaneous abortion (O03)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Spontaneous abortion (O03)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Medical abortion (O04)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Medical abortion (O04)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other abortion (O05)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other abortion (O05)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified pregnancy with abortive outcome (O01-O02,O06-O07)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified pregnancy with abortive outcome (O01-O02,O06-O07)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other direct obstetric deaths (O10-O92)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.2	0.3	0.2	*	*	*	*	*
2001	0.1	*	*	*	0.2	0.4	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	*	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.2	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.2	*	*	*	0.3	0.7	0.4	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.8	0.3	*	*	*	*	*
2000	0.2	*	*	*	0.4	0.7	0.4	*	*	*	*	*
1999	0.2	*	*	*	0.4	0.7	0.4	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.5	*	*	*	*	*	*
2000	0.2	*	*	*	*	0.4	*	*	*	*	*	*
1999	0.2	*	*	*	*	0.3	0.4	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	0.8	1.1	*	*	*	*	*	*
2000	0.4	*	*	*	*	0.9	*	*	*	*	*	*
1999	0.3	*	*	*	*	0.8	0.8	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.1	0.3	0.2	*	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	*	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.2	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other direct obstetric deaths (O10-O92)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	0.3	0.7	0.3	*	*	*	*	*
2001	0.2	*	*	*	0.3	0.7	0.3	*	*	*	*	*
2000	0.2	*	*	*	0.3	0.7	0.3	*	*	*	*	*
1999	0.2	*	*	*	0.4	0.7	0.4	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	0.2	0.1	*	*	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.1	*	*	*	*	0.4	0.2	*	*	*	*	*
2001	0.1	*	*	*	0.2	0.5	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.4	0.2	*	*	*	*	*
1999	0.1	*	*	*	0.2	0.4	0.2	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	0.6	0.9	0.5	*	*	*	*	*
2001	0.3	*	*	*	0.5	1.0	0.5	*	*	*	*	*
2000	0.3	*	*	*	0.5	0.9	0.4	*	*	*	*	*
1999	0.3	*	*	*	0.6	1.0	0.6	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.6	*	*	*	1.2	1.8	1.0	*	*	*	*	*
2001	0.6	*	*	*	1.1	1.8	0.9	*	*	*	*	*
2000	0.5	*	*	*	1.0	1.6	0.8	*	*	*	*	*
1999	0.6	*	*	*	1.2	1.9	1.1	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Eclampsia and pre-eclampsia (O11,O13-O16)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	0.1	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	0.1	0.2	*	*	*	*	*	*
2000	0.1	*	*	*	*	0.2	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Eclampsia and pre-eclampsia (O11,O13-O16)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.2	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,072)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,072)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications predominately related to the puerperium (O85-O92)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	0.1	0.1	*	*	*	*	*
2001	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2000	0.0	*	*	*	0.1	0.1	0.0	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	0.0	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.1	*	*	*	*	0.2	0.1	*	*	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
1999	0.1	*	*	*	*	0.2	0.1	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	0.1	0.1	*	*	*	*	*
2001	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2000	0.0	*	*	*	0.1	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications predominately related to the puerperium (O85-O92)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	0.2	0.1	*	*	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	*	*	*	*	*	*
1999	0.1	*	*	*	*	0.2	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.2	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.2	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.2	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obstetric embolism (O88)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	0.0	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	0.1	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obstetric embolism (O88)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.2	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications predominately related to the puerperium (O85-O87,O89-O92)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications predominately related to the puerperium (O85-O87,O89-O92)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)												
All origins 3/, both sexes												
2002	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2001	0.0	*	*	*	0.1	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
1999	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.1	*	*	*	0.2	0.3	0.2	*	*	*	*	*
2001	0.1	*	*	*	0.1	0.3	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.2	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	0.1	0.2	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	0.1	0.1	0.1	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	0.1	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	0.1	0.3	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	0.2	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.1	*	*	*	*	*
1999	0.1	*	*	*	*	0.2	0.1	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.0	*	*	*	*	0.2	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.2	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obstetric death of unspecified cause (O95)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Obstetric death of unspecified cause (O95)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Indirect obstetric deaths (O98-O99)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
All origins 3/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2001	0.0	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	0.1	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Indirect obstetric deaths (O98-O99)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XVI. Certain conditions originating in the perinatal period (P00-P96)												
All origins 3/, both sexes												
2002	4.9	349.7	0.4	0.1	0.1	*	*	*	*	*	*	*
2001	4.9	340.5	0.5	0.1	*	*	*	*	*	*	*	*
2000	5.0	365.7	0.5	0.1	*	*	*	*	*	*	*	*
1999	5.1	371.0	0.6	0.1	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	5.7	385.7	0.5	*	*	*	*	*	*	*	*	*
2001	5.6	376.6	0.6	0.1	*	*	*	*	*	*	*	*
2000	5.8	403.9	0.7	0.1	*	*	*	*	*	*	*	*
1999	5.9	408.6	0.7	0.1	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	4.2	312.0	0.3	*	*	*	*	*	*	*	*	*
2001	4.2	302.6	0.4	*	*	*	*	*	*	*	*	*
2000	4.3	325.6	0.4	*	*	*	*	*	*	*	*	*
1999	4.4	331.7	0.5	0.1	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	6.2	284.8	*	*	*	*	*	*	*	*	*	*
2001	6.0	269.6	*	*	*	*	*	*	*	*	*	*
2000	6.1	274.9	*	*	*	*	*	*	*	*	*	*
1999	6.3	281.4	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	6.8	313.1	*	*	*	*	*	*	*	*	*	*
2001	6.7	302.0	*	*	*	*	*	*	*	*	*	*
2000	6.6	299.0	*	*	*	*	*	*	*	*	*	*
1999	6.8	305.4	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	5.6	255.2	*	*	*	*	*	*	*	*	*	*
2001	5.3	235.9	*	*	*	*	*	*	*	*	*	*
2000	5.5	249.7	*	*	*	*	*	*	*	*	*	*
1999	5.8	256.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	4.7	361.1	0.4	0.1	*	*	*	*	*	*	*	*
2001	4.6	353.5	0.5	0.1	*	*	*	*	*	*	*	*
2000	4.7	380.5	0.5	0.1	*	*	*	*	*	*	*	*
1999	4.8	383.8	0.6	0.1	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	5.4	398.6	0.5	*	*	*	*	*	*	*	*	*
2001	5.4	390.1	0.6	0.1	*	*	*	*	*	*	*	*
2000	5.5	421.7	0.7	*	*	*	*	*	*	*	*	*
1999	5.6	423.2	0.8	0.1	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XVI. Certain conditions originating in the perinatal period (P00-P96)												
Non-Hispanic 4/, female												
2002	4.0	321.7	0.3	*	*	*	*	*	*	*	*	*
2001	3.9	315.1	0.3	*	*	*	*	*	*	*	*	*
2000	4.0	337.2	0.4	*	*	*	*	*	*	*	*	*
1999	4.1	342.4	0.4	0.1	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	3.2	270.0	0.2	*	*	*	*	*	*	*	*	*
2001	3.2	260.3	0.4	*	*	*	*	*	*	*	*	*
2000	3.2	275.5	0.4	0.1	*	*	*	*	*	*	*	*
1999	3.2	275.1	0.4	0.1	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	3.8	303.0	*	*	*	*	*	*	*	*	*	*
2001	3.7	286.6	0.5	*	*	*	*	*	*	*	*	*
2000	3.7	305.5	0.5	*	*	*	*	*	*	*	*	*
1999	3.8	304.4	0.5	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	2.7	235.5	*	*	*	*	*	*	*	*	*	*
2001	2.7	232.8	*	*	*	*	*	*	*	*	*	*
2000	2.7	243.9	*	*	*	*	*	*	*	*	*	*
1999	2.7	244.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	13.2	731.8	1.1	*	*	*	*	*	*	*	*	*
2001	13.2	765.0	0.9	*	*	*	*	*	*	*	*	*
2000	13.9	838.3	1.1	*	*	*	*	*	*	*	*	*
1999	14.4	872.7	1.2	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	15.4	791.0	*	*	*	*	*	*	*	*	*	*
2001	15.8	846.6	*	*	*	*	*	*	*	*	*	*
2000	16.6	932.9	*	*	*	*	*	*	*	*	*	*
1999	17.1	968.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	11.3	670.1	*	*	*	*	*	*	*	*	*	*
2001	10.9	679.1	*	*	*	*	*	*	*	*	*	*
2000	11.5	740.7	*	*	*	*	*	*	*	*	*	*
1999	12.0	774.4	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)												
All origins 3/, both sexes												
2002	1.1	75.9	*	*	*	*	*	*	*	*	*	*
2001	1.0	70.2	*	*	*	*	*	*	*	*	*	*
2000	1.0	72.7	*	*	*	*	*	*	*	*	*	*
1999	1.0	72.1	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	1.2	83.2	*	*	*	*	*	*	*	*	*	*
2001	1.2	77.8	*	*	*	*	*	*	*	*	*	*
2000	1.1	78.7	*	*	*	*	*	*	*	*	*	*
1999	1.1	76.7	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.9	68.3	*	*	*	*	*	*	*	*	*	*
2001	0.8	62.2	*	*	*	*	*	*	*	*	*	*
2000	0.9	66.5	*	*	*	*	*	*	*	*	*	*
1999	0.9	67.4	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	1.2	55.6	*	*	*	*	*	*	*	*	*	*
2001	1.1	49.2	*	*	*	*	*	*	*	*	*	*
2000	1.0	47.9	*	*	*	*	*	*	*	*	*	*
1999	1.0	46.6	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	1.3	62.6	*	*	*	*	*	*	*	*	*	*
2001	1.3	59.3	*	*	*	*	*	*	*	*	*	*
2000	1.1	49.2	*	*	*	*	*	*	*	*	*	*
1999	1.0	45.5	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	1.0	48.3	*	*	*	*	*	*	*	*	*	*
2001	0.9	38.7	*	*	*	*	*	*	*	*	*	*
2000	1.0	46.5	*	*	*	*	*	*	*	*	*	*
1999	1.1	47.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.0	80.3	*	*	*	*	*	*	*	*	*	*
2001	1.0	74.5	*	*	*	*	*	*	*	*	*	*
2000	1.0	77.8	*	*	*	*	*	*	*	*	*	*
1999	1.0	76.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	1.2	87.5	*	*	*	*	*	*	*	*	*	*
2001	1.1	81.3	*	*	*	*	*	*	*	*	*	*
2000	1.1	84.7	*	*	*	*	*	*	*	*	*	*
1999	1.1	82.7	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)												
Non-Hispanic 4/, female												
2002	0.9	72.7	*	*	*	*	*	*	*	*	*	*
2001	0.8	67.3	*	*	*	*	*	*	*	*	*	*
2000	0.8	70.5	*	*	*	*	*	*	*	*	*	*
1999	0.8	70.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.7	62.1	*	*	*	*	*	*	*	*	*	*
2001	0.7	58.4	*	*	*	*	*	*	*	*	*	*
2000	0.7	61.3	*	*	*	*	*	*	*	*	*	*
1999	0.7	57.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.8	68.6	*	*	*	*	*	*	*	*	*	*
2001	0.8	63.2	*	*	*	*	*	*	*	*	*	*
2000	0.8	67.0	*	*	*	*	*	*	*	*	*	*
1999	0.8	61.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.6	55.2	*	*	*	*	*	*	*	*	*	*
2001	0.6	53.4	*	*	*	*	*	*	*	*	*	*
2000	0.6	55.3	*	*	*	*	*	*	*	*	*	*
1999	0.6	52.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	2.8	153.5	*	*	*	*	*	*	*	*	*	*
2001	2.6	148.3	*	*	*	*	*	*	*	*	*	*
2000	2.6	154.3	*	*	*	*	*	*	*	*	*	*
1999	2.7	165.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	3.2	165.1	*	*	*	*	*	*	*	*	*	*
2001	3.0	161.5	*	*	*	*	*	*	*	*	*	*
2000	2.9	165.2	*	*	*	*	*	*	*	*	*	*
1999	3.2	180.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	2.4	141.3	*	*	*	*	*	*	*	*	*	*
2001	2.1	134.3	*	*	*	*	*	*	*	*	*	*
2000	2.2	143.1	*	*	*	*	*	*	*	*	*	*
1999	2.3	150.0	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders related to short gestation and low birth weight, not elsewhere classified (P07)												
All origins 3/, both sexes												
2002	1.6	115.0	*	*	*	*	*	*	*	*	*	*
2001	1.6	109.3	*	*	*	*	*	*	*	*	*	*
2000	1.6	115.5	*	*	*	*	*	*	*	*	*	*
1999	1.6	115.7	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	1.8	125.5	*	*	*	*	*	*	*	*	*	*
2001	1.7	117.5	*	*	*	*	*	*	*	*	*	*
2000	1.7	122.7	*	*	*	*	*	*	*	*	*	*
1999	1.8	126.3	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	1.4	103.9	*	*	*	*	*	*	*	*	*	*
2001	1.4	100.7	*	*	*	*	*	*	*	*	*	*
2000	1.4	108.0	*	*	*	*	*	*	*	*	*	*
1999	1.4	104.6	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	2.0	91.5	*	*	*	*	*	*	*	*	*	*
2001	1.8	82.2	*	*	*	*	*	*	*	*	*	*
2000	1.8	82.1	*	*	*	*	*	*	*	*	*	*
1999	1.9	85.0	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	2.1	99.0	*	*	*	*	*	*	*	*	*	*
2001	1.9	88.3	*	*	*	*	*	*	*	*	*	*
2000	1.9	85.4	*	*	*	*	*	*	*	*	*	*
1999	2.1	93.1	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	1.8	83.7	*	*	*	*	*	*	*	*	*	*
2001	1.7	75.9	*	*	*	*	*	*	*	*	*	*
2000	1.7	78.6	*	*	*	*	*	*	*	*	*	*
1999	1.7	76.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.5	118.9	*	*	*	*	*	*	*	*	*	*
2001	1.5	114.1	*	*	*	*	*	*	*	*	*	*
2000	1.5	120.7	*	*	*	*	*	*	*	*	*	*
1999	1.5	119.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	1.8	130.1	*	*	*	*	*	*	*	*	*	*
2001	1.7	122.7	*	*	*	*	*	*	*	*	*	*
2000	1.7	128.6	*	*	*	*	*	*	*	*	*	*
1999	1.7	130.7	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Disorders related to short gestation and low birth weight, not elsewhere classified (P07)												
Non-Hispanic 4/, female												
2002	1.3	107.2	*	*	*	*	*	*	*	*	*	*
2001	1.3	105.2	*	*	*	*	*	*	*	*	*	*
2000	1.3	112.4	*	*	*	*	*	*	*	*	*	*
1999	1.3	108.6	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.9	77.6	*	*	*	*	*	*	*	*	*	*
2001	0.9	74.0	*	*	*	*	*	*	*	*	*	*
2000	0.9	75.4	*	*	*	*	*	*	*	*	*	*
1999	0.9	74.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	1.0	84.3	*	*	*	*	*	*	*	*	*	*
2001	1.0	78.9	*	*	*	*	*	*	*	*	*	*
2000	1.0	79.5	*	*	*	*	*	*	*	*	*	*
1999	1.0	80.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.8	70.5	*	*	*	*	*	*	*	*	*	*
2001	0.8	68.8	*	*	*	*	*	*	*	*	*	*
2000	0.8	71.0	*	*	*	*	*	*	*	*	*	*
1999	0.7	67.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	5.0	281.1	*	*	*	*	*	*	*	*	*	*
2001	5.0	291.4	*	*	*	*	*	*	*	*	*	*
2000	5.2	315.0	*	*	*	*	*	*	*	*	*	*
1999	5.3	322.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	6.0	309.0	*	*	*	*	*	*	*	*	*	*
2001	5.8	315.3	*	*	*	*	*	*	*	*	*	*
2000	6.0	340.2	*	*	*	*	*	*	*	*	*	*
1999	6.3	357.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	4.2	252.1	*	*	*	*	*	*	*	*	*	*
2001	4.2	266.3	*	*	*	*	*	*	*	*	*	*
2000	4.5	288.9	*	*	*	*	*	*	*	*	*	*
1999	4.4	286.7	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Birth trauma (P10-P15)												
All origins 3/, both sexes												
2002	0.1	8.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	3.6	*	*	*	*	*	*	*	*	*	*
2000	0.0	0.8	*	*	*	*	*	*	*	*	*	*
1999	0.0	0.5	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.2	10.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	4.2	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	6.4	*	*	*	*	*	*	*	*	*	*
2001	0.0	3.0	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	9.1	*	*	*	*	*	*	*	*	*	*
2001	0.1	3.3	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	11.0	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	7.1	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	8.3	*	*	*	*	*	*	*	*	*	*
2001	0.1	3.6	*	*	*	*	*	*	*	*	*	*
2000	0.0	0.9	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	10.4	*	*	*	*	*	*	*	*	*	*
2001	0.1	4.2	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Birth trauma (P10-P15)												
Non-Hispanic 4/, female												
2002	0.1	6.1	*	*	*	*	*	*	*	*	*	*
2001	0.0	2.9	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	7.3	*	*	*	*	*	*	*	*	*	*
2001	0.0	2.8	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.1	9.1	*	*	*	*	*	*	*	*	*	*
2001	0.0	2.9	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	5.4	*	*	*	*	*	*	*	*	*	*
2001	0.0	2.7	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	13.0	*	*	*	*	*	*	*	*	*	*
2001	0.1	7.2	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	16.1	*	*	*	*	*	*	*	*	*	*
2001	0.2	9.6	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	9.8	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intrauterine hypoxia and birth asphyxia (P20-P21)												
All origins 3/, both sexes												
2002	0.2	14.5	*	*	*	*	*	*	*	*	*	*
2001	0.2	13.2	*	*	*	*	*	*	*	*	*	*
2000	0.2	16.6	*	*	*	*	*	*	*	*	*	*
1999	0.2	16.1	0.1	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.2	14.5	*	*	*	*	*	*	*	*	*	*
2001	0.2	14.3	*	*	*	*	*	*	*	*	*	*
2000	0.3	18.1	*	*	*	*	*	*	*	*	*	*
1999	0.3	17.2	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.2	14.4	*	*	*	*	*	*	*	*	*	*
2001	0.2	12.1	*	*	*	*	*	*	*	*	*	*
2000	0.2	15.0	*	*	*	*	*	*	*	*	*	*
1999	0.2	15.0	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.3	12.4	*	*	*	*	*	*	*	*	*	*
2001	0.3	11.4	*	*	*	*	*	*	*	*	*	*
2000	0.3	12.8	*	*	*	*	*	*	*	*	*	*
1999	0.3	11.8	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	12.9	*	*	*	*	*	*	*	*	*	*
2001	0.3	13.4	*	*	*	*	*	*	*	*	*	*
2000	0.4	15.5	*	*	*	*	*	*	*	*	*	*
1999	0.3	11.4	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	11.8	*	*	*	*	*	*	*	*	*	*
2001	0.2	9.2	*	*	*	*	*	*	*	*	*	*
2000	0.3	10.1	*	*	*	*	*	*	*	*	*	*
1999	0.3	12.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	14.8	*	*	*	*	*	*	*	*	*	*
2001	0.2	13.6	*	*	*	*	*	*	*	*	*	*
2000	0.2	17.3	*	*	*	*	*	*	*	*	*	*
1999	0.2	17.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.2	14.7	*	*	*	*	*	*	*	*	*	*
2001	0.2	14.4	*	*	*	*	*	*	*	*	*	*
2000	0.3	18.6	*	*	*	*	*	*	*	*	*	*
1999	0.3	18.4	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intrauterine hypoxia and birth asphyxia (P20-P21)												
Non-Hispanic 4/, female												
2002	0.2	14.8	*	*	*	*	*	*	*	*	*	*
2001	0.2	12.7	*	*	*	*	*	*	*	*	*	*
2000	0.2	15.9	*	*	*	*	*	*	*	*	*	*
1999	0.2	15.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	13.2	*	*	*	*	*	*	*	*	*	*
2001	0.2	12.2	*	*	*	*	*	*	*	*	*	*
2000	0.2	15.1	*	*	*	*	*	*	*	*	*	*
1999	0.2	14.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.2	13.4	*	*	*	*	*	*	*	*	*	*
2001	0.2	12.6	*	*	*	*	*	*	*	*	*	*
2000	0.2	15.5	*	*	*	*	*	*	*	*	*	*
1999	0.2	15.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	13.0	*	*	*	*	*	*	*	*	*	*
2001	0.1	11.8	*	*	*	*	*	*	*	*	*	*
2000	0.2	14.6	*	*	*	*	*	*	*	*	*	*
1999	0.2	13.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.4	22.4	*	*	*	*	*	*	*	*	*	*
2001	0.4	20.5	*	*	*	*	*	*	*	*	*	*
2000	0.5	27.0	*	*	*	*	*	*	*	*	*	*
1999	0.5	26.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	21.5	*	*	*	*	*	*	*	*	*	*
2001	0.4	22.7	*	*	*	*	*	*	*	*	*	*
2000	0.6	31.1	*	*	*	*	*	*	*	*	*	*
1999	0.6	30.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	23.3	*	*	*	*	*	*	*	*	*	*
2001	0.3	18.1	*	*	*	*	*	*	*	*	*	*
2000	0.4	22.9	*	*	*	*	*	*	*	*	*	*
1999	0.4	22.9	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory distress of newborn (P22)												
All origins 3/, both sexes												
2002	0.3	23.4	*	*	*	*	*	*	*	*	*	*
2001	0.4	25.1	*	*	*	*	*	*	*	*	*	*
2000	0.4	26.3	*	*	*	*	*	*	*	*	*	*
1999	0.4	29.2	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.4	26.2	*	*	*	*	*	*	*	*	*	*
2001	0.4	27.8	*	*	*	*	*	*	*	*	*	*
2000	0.5	33.0	*	*	*	*	*	*	*	*	*	*
1999	0.5	33.7	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.3	20.4	*	*	*	*	*	*	*	*	*	*
2001	0.3	22.2	*	*	*	*	*	*	*	*	*	*
2000	0.2	19.2	*	*	*	*	*	*	*	*	*	*
1999	0.3	24.6	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.4	18.8	*	*	*	*	*	*	*	*	*	*
2001	0.5	22.1	*	*	*	*	*	*	*	*	*	*
2000	0.5	21.5	*	*	*	*	*	*	*	*	*	*
1999	0.5	23.4	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.5	21.6	*	*	*	*	*	*	*	*	*	*
2001	0.5	23.5	*	*	*	*	*	*	*	*	*	*
2000	0.5	23.1	*	*	*	*	*	*	*	*	*	*
1999	0.6	26.4	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	15.9	*	*	*	*	*	*	*	*	*	*
2001	0.5	20.7	*	*	*	*	*	*	*	*	*	*
2000	0.4	19.9	*	*	*	*	*	*	*	*	*	*
1999	0.5	20.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	24.3	*	*	*	*	*	*	*	*	*	*
2001	0.3	25.6	*	*	*	*	*	*	*	*	*	*
2000	0.3	26.9	*	*	*	*	*	*	*	*	*	*
1999	0.4	29.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.4	27.1	*	*	*	*	*	*	*	*	*	*
2001	0.4	28.6	*	*	*	*	*	*	*	*	*	*
2000	0.5	34.8	*	*	*	*	*	*	*	*	*	*
1999	0.4	34.5	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory distress of newborn (P22)												
Non-Hispanic 4/, female												
2002	0.3	21.4	*	*	*	*	*	*	*	*	*	*
2001	0.3	22.3	*	*	*	*	*	*	*	*	*	*
2000	0.2	18.6	*	*	*	*	*	*	*	*	*	*
1999	0.3	24.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	18.5	*	*	*	*	*	*	*	*	*	*
2001	0.2	19.0	*	*	*	*	*	*	*	*	*	*
2000	0.2	19.9	*	*	*	*	*	*	*	*	*	*
1999	0.3	22.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.3	22.6	*	*	*	*	*	*	*	*	*	*
2001	0.3	20.5	*	*	*	*	*	*	*	*	*	*
2000	0.3	27.7	*	*	*	*	*	*	*	*	*	*
1999	0.3	25.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	14.1	*	*	*	*	*	*	*	*	*	*
2001	0.2	17.5	*	*	*	*	*	*	*	*	*	*
2000	0.1	11.7	*	*	*	*	*	*	*	*	*	*
1999	0.2	19.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.9	50.4	*	*	*	*	*	*	*	*	*	*
2001	1.0	55.5	*	*	*	*	*	*	*	*	*	*
2000	1.0	59.1	*	*	*	*	*	*	*	*	*	*
1999	1.0	63.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	1.0	49.7	*	*	*	*	*	*	*	*	*	*
2001	1.2	64.8	*	*	*	*	*	*	*	*	*	*
2000	1.2	70.0	*	*	*	*	*	*	*	*	*	*
1999	1.4	77.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.9	51.1	*	*	*	*	*	*	*	*	*	*
2001	0.7	45.7	*	*	*	*	*	*	*	*	*	*
2000	0.7	47.9	*	*	*	*	*	*	*	*	*	*
1999	0.8	48.9	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other respiratory conditions originating in the perinatal period (P23-P28)												
All origins 3/, both sexes												
2002	0.4	30.9	0.1	*	*	*	*	*	*	*	*	*
2001	0.5	37.3	0.2	*	*	*	*	*	*	*	*	*
2000	0.6	40.2	0.3	*	*	*	*	*	*	*	*	*
1999	0.6	45.3	0.3	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.5	35.5	*	*	*	*	*	*	*	*	*	*
2001	0.6	42.1	*	*	*	*	*	*	*	*	*	*
2000	0.7	45.2	0.3	*	*	*	*	*	*	*	*	*
1999	0.8	51.4	0.3	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.4	26.0	*	*	*	*	*	*	*	*	*	*
2001	0.5	32.2	*	*	*	*	*	*	*	*	*	*
2000	0.5	34.9	*	*	*	*	*	*	*	*	*	*
1999	0.5	38.8	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.5	23.5	*	*	*	*	*	*	*	*	*	*
2001	0.6	26.1	*	*	*	*	*	*	*	*	*	*
2000	0.7	32.4	*	*	*	*	*	*	*	*	*	*
1999	0.8	36.8	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.6	27.9	*	*	*	*	*	*	*	*	*	*
2001	0.6	28.8	*	*	*	*	*	*	*	*	*	*
2000	0.8	36.2	*	*	*	*	*	*	*	*	*	*
1999	1.0	43.2	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	18.9	*	*	*	*	*	*	*	*	*	*
2001	0.6	23.2	*	*	*	*	*	*	*	*	*	*
2000	0.6	28.4	*	*	*	*	*	*	*	*	*	*
1999	0.7	30.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	32.5	*	*	*	*	*	*	*	*	*	*
2001	0.5	39.7	*	*	*	*	*	*	*	*	*	*
2000	0.5	41.3	0.3	*	*	*	*	*	*	*	*	*
1999	0.6	46.6	0.3	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.5	37.0	*	*	*	*	*	*	*	*	*	*
2001	0.6	45.1	*	*	*	*	*	*	*	*	*	*
2000	0.6	46.8	0.3	*	*	*	*	*	*	*	*	*
1999	0.7	52.5	0.4	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other respiratory conditions originating in the perinatal period (P23-P28)												
Non-Hispanic 4/, female												
2002	0.3	27.7	*	*	*	*	*	*	*	*	*	*
2001	0.4	34.0	*	*	*	*	*	*	*	*	*	*
2000	0.4	35.5	*	*	*	*	*	*	*	*	*	*
1999	0.5	40.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.3	25.9	*	*	*	*	*	*	*	*	*	*
2001	0.4	30.5	*	*	*	*	*	*	*	*	*	*
2000	0.4	31.1	*	*	*	*	*	*	*	*	*	*
1999	0.4	37.6	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.4	30.0	*	*	*	*	*	*	*	*	*	*
2001	0.4	35.0	*	*	*	*	*	*	*	*	*	*
2000	0.4	34.7	*	*	*	*	*	*	*	*	*	*
1999	0.5	43.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	21.6	*	*	*	*	*	*	*	*	*	*
2001	0.3	25.7	*	*	*	*	*	*	*	*	*	*
2000	0.3	27.3	*	*	*	*	*	*	*	*	*	*
1999	0.3	31.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	1.1	61.0	*	*	*	*	*	*	*	*	*	*
2001	1.4	81.2	*	*	*	*	*	*	*	*	*	*
2000	1.5	86.3	*	*	*	*	*	*	*	*	*	*
1999	1.5	91.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	1.4	68.2	*	*	*	*	*	*	*	*	*	*
2001	1.7	92.2	*	*	*	*	*	*	*	*	*	*
2000	1.8	100.3	*	*	*	*	*	*	*	*	*	*
1999	1.8	98.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.9	53.6	*	*	*	*	*	*	*	*	*	*
2001	1.1	69.5	*	*	*	*	*	*	*	*	*	*
2000	1.1	71.9	*	*	*	*	*	*	*	*	*	*
1999	1.3	84.6	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections specific to the perinatal period (P35-P39)												
All origins 3/, both sexes												
2002	0.3	23.5	*	*	*	*	*	*	*	*	*	*
2001	0.3	21.5	*	*	*	*	*	*	*	*	*	*
2000	0.3	24.4	*	*	*	*	*	*	*	*	*	*
1999	0.3	22.9	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.4	25.1	*	*	*	*	*	*	*	*	*	*
2001	0.4	24.5	*	*	*	*	*	*	*	*	*	*
2000	0.4	27.6	*	*	*	*	*	*	*	*	*	*
1999	0.4	25.5	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.3	21.8	*	*	*	*	*	*	*	*	*	*
2001	0.3	18.5	*	*	*	*	*	*	*	*	*	*
2000	0.3	21.0	*	*	*	*	*	*	*	*	*	*
1999	0.3	20.3	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.4	20.4	*	*	*	*	*	*	*	*	*	*
2001	0.4	19.1	*	*	*	*	*	*	*	*	*	*
2000	0.4	20.0	*	*	*	*	*	*	*	*	*	*
1999	0.4	17.6	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.5	23.2	*	*	*	*	*	*	*	*	*	*
2001	0.5	22.8	*	*	*	*	*	*	*	*	*	*
2000	0.5	24.3	*	*	*	*	*	*	*	*	*	*
1999	0.5	20.2	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	17.4	*	*	*	*	*	*	*	*	*	*
2001	0.3	15.2	*	*	*	*	*	*	*	*	*	*
2000	0.3	15.4	*	*	*	*	*	*	*	*	*	*
1999	0.4	14.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	24.0	*	*	*	*	*	*	*	*	*	*
2001	0.3	21.9	*	*	*	*	*	*	*	*	*	*
2000	0.3	25.1	*	*	*	*	*	*	*	*	*	*
1999	0.3	23.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.3	25.3	*	*	*	*	*	*	*	*	*	*
2001	0.3	24.7	*	*	*	*	*	*	*	*	*	*
2000	0.4	28.2	*	*	*	*	*	*	*	*	*	*
1999	0.3	26.3	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections specific to the perinatal period (P35-P39)												
Non-Hispanic 4/, female												
2002	0.3	22.6	*	*	*	*	*	*	*	*	*	*
2001	0.2	19.1	*	*	*	*	*	*	*	*	*	*
2000	0.3	21.9	*	*	*	*	*	*	*	*	*	*
1999	0.3	21.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	19.4	*	*	*	*	*	*	*	*	*	*
2001	0.2	16.4	*	*	*	*	*	*	*	*	*	*
2000	0.2	18.7	*	*	*	*	*	*	*	*	*	*
1999	0.2	17.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.3	21.4	*	*	*	*	*	*	*	*	*	*
2001	0.3	19.6	*	*	*	*	*	*	*	*	*	*
2000	0.3	21.5	*	*	*	*	*	*	*	*	*	*
1999	0.2	18.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	17.4	*	*	*	*	*	*	*	*	*	*
2001	0.2	13.1	*	*	*	*	*	*	*	*	*	*
2000	0.2	15.8	*	*	*	*	*	*	*	*	*	*
1999	0.2	16.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.8	45.1	*	*	*	*	*	*	*	*	*	*
2001	0.8	45.8	*	*	*	*	*	*	*	*	*	*
2000	0.9	53.6	*	*	*	*	*	*	*	*	*	*
1999	0.9	53.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.9	44.8	*	*	*	*	*	*	*	*	*	*
2001	0.9	48.5	*	*	*	*	*	*	*	*	*	*
2000	1.1	60.1	*	*	*	*	*	*	*	*	*	*
1999	1.1	63.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.8	45.4	*	*	*	*	*	*	*	*	*	*
2001	0.7	43.0	*	*	*	*	*	*	*	*	*	*
2000	0.7	46.9	*	*	*	*	*	*	*	*	*	*
1999	0.7	43.6	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)												
All origins 3/, both sexes												
2002	0.8	58.1	*	*	*	*	*	*	*	*	*	*
2001	0.9	60.2	*	*	*	*	*	*	*	*	*	*
2000	0.9	69.3	*	*	*	*	*	*	*	*	*	*
1999	0.9	69.1	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	1.0	65.1	*	*	*	*	*	*	*	*	*	*
2001	1.0	68.5	*	*	*	*	*	*	*	*	*	*
2000	1.1	78.0	*	*	*	*	*	*	*	*	*	*
1999	1.1	77.0	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.7	50.7	*	*	*	*	*	*	*	*	*	*
2001	0.7	51.6	*	*	*	*	*	*	*	*	*	*
2000	0.8	60.1	*	*	*	*	*	*	*	*	*	*
1999	0.8	60.7	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	1.2	53.5	*	*	*	*	*	*	*	*	*	*
2001	1.3	56.2	*	*	*	*	*	*	*	*	*	*
2000	1.3	57.6	*	*	*	*	*	*	*	*	*	*
1999	1.3	60.1	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	1.2	54.9	*	*	*	*	*	*	*	*	*	*
2001	1.4	62.2	*	*	*	*	*	*	*	*	*	*
2000	1.4	64.9	*	*	*	*	*	*	*	*	*	*
1999	1.5	65.2	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	1.1	52.0	*	*	*	*	*	*	*	*	*	*
2001	1.1	49.9	*	*	*	*	*	*	*	*	*	*
2000	1.1	49.9	*	*	*	*	*	*	*	*	*	*
1999	1.2	54.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.8	58.1	*	*	*	*	*	*	*	*	*	*
2001	0.8	60.5	*	*	*	*	*	*	*	*	*	*
2000	0.9	70.6	*	*	*	*	*	*	*	*	*	*
1999	0.9	69.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.9	66.6	*	*	*	*	*	*	*	*	*	*
2001	1.0	69.0	*	*	*	*	*	*	*	*	*	*
2000	1.0	79.3	*	*	*	*	*	*	*	*	*	*
1999	1.0	77.3	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)												
Non-Hispanic 4/, female												
2002	0.6	49.1	*	*	*	*	*	*	*	*	*	*
2001	0.6	51.5	*	*	*	*	*	*	*	*	*	*
2000	0.7	61.3	*	*	*	*	*	*	*	*	*	*
1999	0.7	61.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.6	46.1	*	*	*	*	*	*	*	*	*	*
2001	0.6	47.0	*	*	*	*	*	*	*	*	*	*
2000	0.6	53.4	*	*	*	*	*	*	*	*	*	*
1999	0.6	51.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.7	53.6	*	*	*	*	*	*	*	*	*	*
2001	0.7	53.8	*	*	*	*	*	*	*	*	*	*
2000	0.7	58.8	*	*	*	*	*	*	*	*	*	*
1999	0.7	58.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.4	38.2	*	*	*	*	*	*	*	*	*	*
2001	0.5	39.8	*	*	*	*	*	*	*	*	*	*
2000	0.5	47.6	*	*	*	*	*	*	*	*	*	*
1999	0.5	43.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	1.9	105.2	*	*	*	*	*	*	*	*	*	*
2001	2.0	115.2	*	*	*	*	*	*	*	*	*	*
2000	2.3	141.6	*	*	*	*	*	*	*	*	*	*
1999	2.4	148.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	2.3	116.6	*	*	*	*	*	*	*	*	*	*
2001	2.5	132.1	*	*	*	*	*	*	*	*	*	*
2000	2.9	165.8	*	*	*	*	*	*	*	*	*	*
1999	2.8	159.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	1.6	93.4	*	*	*	*	*	*	*	*	*	*
2001	1.6	97.4	*	*	*	*	*	*	*	*	*	*
2000	1.8	116.6	*	*	*	*	*	*	*	*	*	*
1999	2.1	137.1	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins 3/, both sexes												
2002	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8
All origins 3/, male												
2002	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5
All origins 3/, female												
2002	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001	3.4	130.0	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6
Hispanic, both sexes												
2002	4.4	150.7	3.5	1.0	1.0	0.7	0.7	1.1	1.3	*	3.8	*
2001	4.4	146.5	3.4	0.8	1.1	0.7	0.8	1.1	1.3	1.9	4.3	*
2000	4.6	153.3	3.8	0.8	0.8	0.7	0.8	1.1	1.8	2.0	4.7	*
1999	4.4	141.8	3.5	0.9	0.9	0.9	0.7	1.4	1.6	2.9	*	*
Hispanic, male												
2002	4.5	153.1	3.8	0.9	1.1	0.9	*	1.3	*	*	*	*
2001	4.7	153.4	3.5	1.0	1.1	0.7	1.0	1.5	*	*	*	*
2000	4.7	153.3	4.0	0.8	1.1	0.7	1.0	1.4	*	*	*	*
1999	4.3	138.1	4.3	0.9	0.9	0.7	*	1.6	*	*	*	*
Hispanic, female												
2002	4.3	148.2	3.3	1.0	0.9	*	0.8	*	*	*	*	*
2001	4.1	139.3	3.3	0.6	1.1	*	*	*	*	*	*	*
2000	4.4	153.3	3.5	0.7	*	0.7	*	*	*	*	*	*
1999	4.5	145.8	2.7	0.9	0.9	1.0	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	3.6	134.9	3.3	1.0	1.2	1.3	1.4	1.8	2.5	2.5	3.8	6.7
2001	3.5	132.8	3.7	0.9	1.3	1.3	1.4	1.7	2.4	2.5	3.7	7.2
2000	3.6	148.8	3.1	1.0	1.2	1.3	1.3	1.7	2.4	2.6	3.9	7.5
1999	3.6	143.1	3.6	1.1	1.2	1.2	1.3	1.8	2.4	2.5	4.0	7.9
Non-Hispanic 4/, male												
2002	3.8	138.1	3.5	1.0	1.4	1.5	1.4	2.0	2.6	2.6	3.9	7.0
2001	3.7	139.4	3.6	1.0	1.6	1.4	1.5	1.7	2.3	2.7	4.0	7.6
2000	3.9	154.4	3.1	1.0	1.4	1.5	1.5	1.9	2.2	2.7	3.9	10.4
1999	3.8	148.4	3.9	1.1	1.4	1.3	1.4	1.9	2.6	2.4	4.2	8.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic 4/, female												
2002	3.4	131.5	3.2	1.0	1.1	1.1	1.3	1.7	2.4	2.3	3.8	6.6
2001	3.3	125.8	3.7	0.9	1.0	1.2	1.3	1.7	2.4	2.2	3.5	7.0
2000	3.4	142.8	3.0	1.0	1.0	1.1	1.2	1.4	2.5	2.4	3.9	6.4
1999	3.4	137.4	3.3	1.1	0.9	1.1	1.2	1.6	2.2	2.6	3.9	7.6
Non-Hispanic white, both sexes												
2002	3.4	131.2	3.0	1.0	1.2	1.3	1.4	1.9	2.5	2.5	3.9	6.9
2001	3.4	127.4	3.1	0.9	1.2	1.3	1.4	1.7	2.5	2.5	3.8	7.4
2000	3.4	141.2	2.6	0.9	1.1	1.3	1.3	1.7	2.4	2.6	4.0	7.8
1999	3.4	136.8	3.0	0.9	1.1	1.3	1.3	1.8	2.5	2.5	4.0	8.0
Non-Hispanic white, male												
2002	3.6	137.2	3.2	1.0	1.4	1.6	1.4	2.0	2.6	2.7	3.8	7.3
2001	3.6	135.5	3.1	1.0	1.4	1.5	1.5	1.7	2.4	2.9	4.1	7.4
2000	3.6	147.9	2.6	0.9	1.3	1.6	1.4	1.9	2.3	2.6	4.1	10.8
1999	3.6	144.4	3.2	0.9	1.4	1.3	1.4	2.0	2.7	2.4	4.1	8.6
Non-Hispanic white, female												
2002	3.3	125.0	2.9	1.0	1.1	1.1	1.3	1.8	2.5	2.4	3.9	6.7
2001	3.2	118.8	3.2	0.7	1.0	1.2	1.3	1.7	2.6	2.3	3.6	7.4
2000	3.2	134.2	2.6	1.0	1.0	1.1	1.2	1.6	2.5	2.6	4.0	6.6
1999	3.3	128.9	2.9	1.0	0.8	1.2	1.2	1.7	2.3	2.6	4.0	7.8
Non-Hispanic black, both sexes												
2002	4.7	156.9	4.7	1.3	1.7	1.5	1.6	1.8	2.7	2.4	3.4	*
2001	4.7	163.5	5.4	1.1	2.0	1.3	1.6	1.8	1.9	2.1	3.0	*
2000	5.0	184.1	5.2	1.4	1.6	1.4	1.6	1.6	2.4	2.7	2.8	*
1999	4.8	174.8	5.8	1.6	1.5	1.1	1.4	1.9	2.0	2.9	4.3	7.4
Non-Hispanic black, male												
2002	5.2	153.3	4.8	1.4	2.0	1.8	1.7	2.3	3.5	3.1	*	*
2001	5.1	167.5	5.5	1.0	2.5	1.2	1.8	1.7	2.1	*	*	*
2000	5.5	186.0	5.6	1.5	2.1	1.7	1.9	2.2	*	3.9	*	*
1999	5.3	179.5	6.6	1.6	1.8	1.3	1.6	2.3	2.1	*	*	*
Non-Hispanic black, female												
2002	4.4	160.6	4.7	1.1	1.3	1.2	1.6	1.5	2.0	*	*	*
2001	4.4	159.2	5.4	1.3	1.4	1.4	1.5	1.8	1.8	2.4	*	*
2000	4.5	182.2	4.8	1.2	1.2	1.1	1.4	1.2	3.1	*	*	*
1999	4.4	170.0	5.0	1.6	1.1	1.0	1.2	1.5	1.9	3.2	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anencephaly and similar malformations (Q00)												
All origins 3/, both sexes												
2002	0.1	7.4	*	*	*	*	*	*	*	*	*	*
2001	0.1	6.5	*	*	*	*	*	*	*	*	*	*
2000	0.1	8.5	*	*	*	*	*	*	*	*	*	*
1999	0.1	8.2	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.1	6.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	5.3	*	*	*	*	*	*	*	*	*	*
2000	0.1	8.0	*	*	*	*	*	*	*	*	*	*
1999	0.1	6.3	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	8.1	*	*	*	*	*	*	*	*	*	*
2001	0.1	7.8	*	*	*	*	*	*	*	*	*	*
2000	0.1	9.0	*	*	*	*	*	*	*	*	*	*
1999	0.1	10.2	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	10.0	*	*	*	*	*	*	*	*	*	*
2001	0.2	9.7	*	*	*	*	*	*	*	*	*	*
2000	0.2	10.4	*	*	*	*	*	*	*	*	*	*
1999	0.2	11.0	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	9.9	*	*	*	*	*	*	*	*	*	*
2001	0.2	8.6	*	*	*	*	*	*	*	*	*	*
2000	0.2	8.9	*	*	*	*	*	*	*	*	*	*
1999	0.2	8.0	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	10.1	*	*	*	*	*	*	*	*	*	*
2001	0.2	10.7	*	*	*	*	*	*	*	*	*	*
2000	0.3	12.0	*	*	*	*	*	*	*	*	*	*
1999	0.3	14.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	6.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	5.6	*	*	*	*	*	*	*	*	*	*
2000	0.1	8.0	*	*	*	*	*	*	*	*	*	*
1999	0.1	7.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	5.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	4.4	*	*	*	*	*	*	*	*	*	*
2000	0.1	7.7	*	*	*	*	*	*	*	*	*	*
1999	0.1	5.8	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anencephaly and similar malformations (Q00)												
Non-Hispanic 4/, female												
2002	0.1	7.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	6.8	*	*	*	*	*	*	*	*	*	*
2000	0.1	8.2	*	*	*	*	*	*	*	*	*	*
1999	0.1	8.9	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	6.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	5.8	*	*	*	*	*	*	*	*	*	*
2000	0.1	8.2	*	*	*	*	*	*	*	*	*	*
1999	0.1	7.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.1	5.9	*	*	*	*	*	*	*	*	*	*
2001	0.1	4.6	*	*	*	*	*	*	*	*	*	*
2000	0.1	7.8	*	*	*	*	*	*	*	*	*	*
1999	0.1	5.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	7.4	*	*	*	*	*	*	*	*	*	*
2001	0.1	6.9	*	*	*	*	*	*	*	*	*	*
2000	0.1	8.6	*	*	*	*	*	*	*	*	*	*
1999	0.1	9.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	6.0	*	*	*	*	*	*	*	*	*	*
2001	0.1	5.1	*	*	*	*	*	*	*	*	*	*
2000	0.1	6.9	*	*	*	*	*	*	*	*	*	*
1999	0.1	6.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	8.2	*	*	*	*	*	*	*	*	*	*
2001	0.1	7.1	*	*	*	*	*	*	*	*	*	*
2000	0.1	7.8	*	*	*	*	*	*	*	*	*	*
1999	0.1	7.9	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Spina bifida (Q05)												
All origins 3/, both sexes												
2002	0.1	0.5	*	*	0.1	0.1	0.1	*	*	*	*	*
2001	0.0	*	*	*	0.1	0.1	*	*	*	*	*	*
2000	0.0	0.7	*	*	0.1	0.1	*	*	*	*	*	*
1999	0.0	0.6	*	*	0.1	0.0	*	*	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2001	0.1	*	*	*	0.1	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	0.1	0.1	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Spina bifida (Q05)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	0.1	*	*	*	*	*	*
2000	0.0	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)												
All origins 3/, both sexes												
2002	0.3	9.3	0.8	0.3	0.1	0.1	0.1	0.1	*	*	*	*
2001	0.3	9.7	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.3	10.7	0.6	0.3	0.1	0.1	0.1	*	*	*	*	*
1999	0.3	9.4	0.7	0.3	0.1	0.1	0.1	0.1	*	*	*	*
All origins 3/, male												
2002	0.3	9.0	0.7	0.2	0.1	*	*	*	*	*	*	*
2001	0.3	9.4	0.8	0.2	0.1	0.1	0.1	*	*	*	*	*
2000	0.3	10.6	0.6	0.2	0.1	*	*	*	*	*	*	*
1999	0.3	8.6	0.7	0.2	0.1	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.3	9.6	0.9	0.3	0.2	*	0.1	*	*	*	*	*
2001	0.3	10.0	0.9	0.2	0.1	*	*	*	*	*	*	*
2000	0.3	10.9	0.7	0.3	*	*	*	*	*	*	*	*
1999	0.3	10.2	0.7	0.3	0.1	0.1	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.4	13.9	0.8	*	*	*	*	*	*	*	*	*
2001	0.4	12.5	0.9	*	*	*	*	*	*	*	*	*
2000	0.4	11.3	0.9	*	*	*	*	*	*	*	*	*
1999	0.3	8.1	0.9	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.4	12.7	*	*	*	*	*	*	*	*	*	*
2001	0.4	13.0	*	*	*	*	*	*	*	*	*	*
2000	0.4	9.6	*	*	*	*	*	*	*	*	*	*
1999	0.3	5.4	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	15.2	*	*	*	*	*	*	*	*	*	*
2001	0.4	12.0	*	*	*	*	*	*	*	*	*	*
2000	0.4	13.0	*	*	*	*	*	*	*	*	*	*
1999	0.4	10.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	8.1	0.8	0.3	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.3	8.9	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.2	10.4	0.6	0.3	0.1	0.1	0.1	*	*	*	*	*
1999	0.3	9.6	0.6	0.3	0.1	0.1	0.1	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.2	8.1	0.7	0.2	0.1	*	*	*	*	*	*	*
2001	0.3	8.3	0.8	0.2	0.1	*	*	*	*	*	*	*
2000	0.3	10.7	0.5	0.2	0.1	*	*	*	*	*	*	*
1999	0.3	9.4	0.6	0.2	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)												
Non-Hispanic 4/, female												
2002	0.3	8.1	1.0	0.3	0.2	*	0.1	*	*	*	*	*
2001	0.2	9.4	0.9	0.2	0.1	*	*	*	*	*	*	*
2000	0.2	10.2	0.7	0.3	*	*	*	*	*	*	*	*
1999	0.3	9.9	0.6	0.3	0.1	0.1	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	7.7	0.7	0.3	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.2	8.5	0.7	0.2	0.2	0.1	0.1	*	*	*	*	*
2000	0.2	9.8	0.5	0.2	0.1	*	0.1	*	*	*	*	*
1999	0.2	8.7	0.6	0.2	0.1	0.1	0.1	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.2	7.6	0.6	0.3	*	*	*	*	*	*	*	*
2001	0.3	8.2	0.7	0.2	0.2	*	*	*	*	*	*	*
2000	0.2	10.2	0.5	0.2	*	*	*	*	*	*	*	*
1999	0.2	8.7	0.6	0.2	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	7.7	0.8	0.3	0.2	*	*	*	*	*	*	*
2001	0.2	8.7	0.7	*	*	*	*	*	*	*	*	*
2000	0.2	9.3	0.6	0.2	*	*	*	*	*	*	*	*
1999	0.2	8.6	0.5	0.3	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.4	10.0	1.3	0.4	*	*	*	*	*	*	*	*
2001	0.4	11.3	1.5	*	*	*	*	*	*	*	*	*
2000	0.4	14.0	*	0.4	*	*	*	*	*	*	*	*
1999	0.4	13.5	1.0	0.4	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	10.6	*	*	*	*	*	*	*	*	*	*
2001	0.4	10.5	*	*	*	*	*	*	*	*	*	*
2000	0.4	14.0	*	*	*	*	*	*	*	*	*	*
1999	0.4	12.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	9.5	*	*	*	*	*	*	*	*	*	*
2001	0.4	12.1	*	*	*	*	*	*	*	*	*	*
2000	0.4	14.1	*	*	*	*	*	*	*	*	*	*
1999	0.4	15.0	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)												
All origins 3/, both sexes												
2002	1.1	37.4	1.2	0.4	0.5	0.6	0.5	0.5	0.5	0.7	1.2	2.3
2001	1.2	38.6	1.3	0.3	0.6	0.5	0.5	0.6	0.5	0.7	1.2	2.6
2000	1.2	42.9	1.2	0.3	0.6	0.6	0.5	0.5	0.6	0.7	1.4	2.5
1999	1.3	42.1	1.3	0.4	0.6	0.6	0.6	0.6	0.7	0.9	1.4	3.3
All origins 3/, male												
2002	1.2	40.4	1.5	0.4	0.7	0.7	0.6	0.5	0.5	0.6	1.1	1.4
2001	1.3	41.8	1.4	0.3	0.8	0.6	0.6	0.6	0.5	0.8	1.2	1.7
2000	1.3	44.5	1.3	0.4	0.7	0.7	0.6	0.6	0.6	0.8	1.5	2.7
1999	1.4	45.7	1.5	0.4	0.7	0.6	0.6	0.7	0.7	0.8	1.3	3.1
All origins 3/, female												
2002	1.0	34.4	1.0	0.3	0.4	0.5	0.5	0.6	0.5	0.7	1.3	2.6
2001	1.1	35.2	1.2	0.3	0.4	0.4	0.5	0.6	0.5	0.5	1.2	3.0
2000	1.1	41.3	1.2	0.3	0.4	0.5	0.5	0.5	0.6	0.7	1.4	2.4
1999	1.2	38.3	1.1	0.3	0.4	0.6	0.6	0.5	0.6	0.9	1.5	3.4
Hispanic, both sexes												
2002	1.3	41.4	1.2	0.3	0.4	0.4	*	*	*	*	*	*
2001	1.3	40.9	1.2	*	0.4	*	*	*	*	*	*	*
2000	1.4	41.4	1.5	*	0.3	0.3	0.4	*	*	*	*	*
1999	1.4	42.5	1.4	0.4	0.4	0.4	*	*	*	*	*	*
Hispanic, male												
2002	1.4	43.9	1.3	*	0.5	*	*	*	*	*	*	*
2001	1.4	43.7	1.3	*	*	*	*	*	*	*	*	*
2000	1.4	39.0	1.6	*	*	*	*	*	*	*	*	*
1999	1.5	42.4	1.8	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	1.2	38.8	*	*	*	*	*	*	*	*	*	*
2001	1.2	37.9	*	*	*	*	*	*	*	*	*	*
2000	1.3	43.8	1.4	*	*	*	*	*	*	*	*	*
1999	1.4	42.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.1	36.1	1.2	0.4	0.5	0.6	0.6	0.5	0.5	0.6	1.2	2.3
2001	1.1	37.6	1.3	0.3	0.7	0.6	0.6	0.6	0.5	0.7	1.2	2.6
2000	1.2	43.0	1.2	0.4	0.6	0.6	0.6	0.5	0.6	0.7	1.4	2.4
1999	1.2	41.7	1.3	0.4	0.6	0.6	0.6	0.6	0.7	0.9	1.4	3.3
Non-Hispanic 4/, male												
2002	1.2	39.0	1.5	0.4	0.7	0.8	0.6	0.5	0.5	0.6	1.1	*
2001	1.3	41.1	1.4	0.3	0.9	0.7	0.6	0.6	0.5	0.8	1.2	1.6
2000	1.3	45.5	1.2	0.4	0.7	0.8	0.6	0.6	0.6	0.8	1.4	2.5
1999	1.4	46.2	1.4	0.4	0.8	0.7	0.7	0.7	0.7	0.8	1.3	3.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)												
Non-Hispanic 4/, female												
2002	1.0	33.0	0.9	0.3	0.4	0.5	0.5	0.6	0.5	0.7	1.4	2.7
2001	1.0	34.0	1.2	0.3	0.4	0.5	0.5	0.6	0.5	0.6	1.3	3.0
2000	1.1	40.4	1.1	0.3	0.5	0.5	0.5	0.5	0.6	0.7	1.4	2.4
1999	1.1	37.0	1.2	0.3	0.4	0.6	0.6	0.5	0.7	0.9	1.5	3.4
Non-Hispanic white, both sexes												
2002	1.0	33.9	1.2	0.3	0.5	0.7	0.6	0.6	0.5	0.7	1.2	2.4
2001	1.1	35.1	1.1	0.3	0.6	0.6	0.6	0.6	0.5	0.7	1.3	2.7
2000	1.1	40.6	1.0	0.3	0.5	0.7	0.6	0.6	0.6	0.7	1.5	2.6
1999	1.2	40.7	1.0	0.3	0.6	0.7	0.6	0.6	0.7	0.9	1.5	3.3
Non-Hispanic white, male												
2002	1.1	37.7	1.6	0.4	0.7	0.8	0.6	0.5	0.5	0.6	1.0	*
2001	1.2	39.5	1.3	0.3	0.8	0.8	0.6	0.6	0.5	0.9	1.3	*
2000	1.2	44.1	1.0	0.3	0.6	0.8	0.6	0.6	0.6	0.7	1.5	2.7
1999	1.3	46.2	1.1	0.4	0.8	0.8	0.7	0.6	0.7	0.8	1.2	3.0
Non-Hispanic white, female												
2002	1.0	29.8	0.9	0.3	0.4	0.5	0.5	0.6	0.5	0.7	1.4	2.8
2001	1.0	30.5	1.0	0.2	0.4	0.5	0.6	0.6	0.4	0.6	1.3	3.2
2000	1.0	36.8	1.0	0.3	0.4	0.5	0.5	0.5	0.6	0.7	1.5	2.5
1999	1.1	35.0	1.0	0.3	0.4	0.6	0.6	0.6	0.7	0.9	1.6	3.5
Non-Hispanic black, both sexes												
2002	1.5	45.3	1.4	0.5	0.7	0.7	0.6	*	*	*	*	*
2001	1.6	49.7	2.0	0.4	1.2	0.6	0.7	0.6	*	*	*	*
2000	1.6	52.9	1.9	0.5	0.9	0.7	0.7	0.5	*	*	*	*
1999	1.6	50.1	2.3	0.6	0.8	0.5	0.7	0.5	*	*	*	*
Non-Hispanic black, male												
2002	1.6	43.6	*	0.7	1.0	1.0	*	*	*	*	*	*
2001	1.8	50.7	1.9	*	1.7	*	*	*	*	*	*	*
2000	1.8	50.8	2.2	*	1.3	0.9	0.8	*	*	*	*	*
1999	1.9	52.7	2.6	*	1.1	*	0.8	*	*	*	*	*
Non-Hispanic black, female												
2002	1.3	47.0	*	*	*	*	*	*	*	*	*	*
2001	1.5	48.7	2.1	*	0.7	*	*	*	*	*	*	*
2000	1.4	55.0	*	*	*	*	*	*	*	*	*	*
1999	1.4	47.5	2.1	0.7	*	*	0.7	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other congenital malformations of circulatory system (Q25-Q28)												
All origins 3/, both sexes												
2002	0.3	5.5	0.2	0.1	0.1	0.1	0.1	0.2	0.3	0.5	1.0	1.5
2001	0.3	4.8	*	0.1	0.1	0.1	0.1	0.2	0.3	0.6	0.7	1.7
2000	0.3	5.5	0.2	0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.9	2.1
1999	0.3	5.8	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.9	1.7
All origins 3/, male												
2002	0.3	5.8	*	*	*	0.2	0.1	0.3	0.4	0.6	1.0	1.8
2001	0.3	4.9	*	0.1	0.1	*	0.1	0.2	0.3	0.6	1.0	2.1
2000	0.3	5.5	*	*	0.1	0.1	0.2	0.3	0.3	0.5	1.0	3.0
1999	0.3	5.9	*	*	0.1	0.1	0.2	0.3	0.4	0.5	1.1	2.2
All origins 3/, female												
2002	0.3	5.2	*	*	*	*	0.1	0.2	0.2	0.4	1.1	1.4
2001	0.3	4.6	*	*	*	*	0.1	0.2	0.2	0.6	0.6	1.6
2000	0.3	5.4	*	0.1	*	*	0.2	0.2	0.3	0.4	0.9	1.7
1999	0.3	5.7	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.5
Hispanic, both sexes												
2002	0.3	5.5	*	*	*	*	*	*	*	*	*	*
2001	0.3	5.4	*	*	*	*	*	*	*	*	*	*
2000	0.3	6.2	*	*	*	*	*	*	*	*	*	*
1999	0.3	3.8	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	6.0	*	*	*	*	*	*	*	*	*	*
2000	0.3	7.1	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	6.6	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	5.3	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	5.5	*	0.1	0.1	0.1	0.1	0.2	0.3	0.5	1.0	1.6
2001	0.3	4.5	*	0.1	0.1	*	0.1	0.2	0.3	0.5	0.7	1.7
2000	0.3	5.2	0.2	0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.9	2.1
1999	0.3	6.2	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.9	1.7
Non-Hispanic 4/, male												
2002	0.3	6.2	*	*	*	0.2	0.1	0.3	0.4	0.6	1.0	1.8
2001	0.3	4.6	*	0.1	*	*	0.1	0.2	0.3	0.5	0.9	2.1
2000	0.3	5.1	*	*	*	0.2	0.2	0.3	0.3	0.5	1.0	3.1
1999	0.3	6.4	*	*	*	*	0.2	0.3	0.4	0.4	1.1	2.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other congenital malformations of circulatory system (Q25-Q28)												
Non-Hispanic 4/, female												
2002	0.3	4.9	*	*	*	*	0.1	0.2	0.2	0.5	1.0	1.5
2001	0.3	4.5	*	*	*	*	*	0.2	0.2	0.6	0.5	1.6
2000	0.3	5.3	*	*	*	*	0.2	0.2	0.3	0.4	0.8	1.7
1999	0.3	5.9	*	*	*	*	0.1	0.2	0.3	0.6	0.8	1.5
Non-Hispanic white, both sexes												
2002	0.3	4.8	*	*	*	0.1	0.1	0.2	0.3	0.5	1.0	1.7
2001	0.3	4.2	*	0.1	*	*	0.1	0.2	0.3	0.5	0.8	1.7
2000	0.3	4.6	*	0.1	0.1	0.1	0.2	0.2	0.3	0.4	0.9	2.2
1999	0.3	5.4	*	*	*	0.1	0.2	0.3	0.3	0.5	0.9	1.7
Non-Hispanic white, male												
2002	0.3	4.9	*	*	*	0.2	*	0.2	0.3	0.6	0.9	1.9
2001	0.3	4.4	*	*	*	*	0.1	0.2	0.3	0.5	1.0	2.0
2000	0.3	4.6	*	*	*	*	0.2	0.3	0.3	0.5	1.1	3.3
1999	0.3	5.9	*	*	*	*	0.2	0.3	0.3	0.4	1.0	2.2
Non-Hispanic white, female												
2002	0.3	4.7	*	*	*	*	*	0.2	0.2	0.5	1.1	1.5
2001	0.3	4.0	*	*	*	*	*	0.1	0.3	0.5	0.6	1.6
2000	0.3	4.7	*	*	*	*	0.2	0.1	0.3	0.4	0.8	1.7
1999	0.3	4.9	*	*	*	*	0.1	0.2	0.3	0.5	0.8	1.5
Non-Hispanic black, both sexes												
2002	0.5	7.6	*	*	*	*	*	0.5	*	*	*	*
2001	0.3	6.1	*	*	*	*	*	*	*	*	*	*
2000	0.4	8.1	*	*	*	*	*	0.6	*	*	*	*
1999	0.4	9.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.5	9.4	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	8.5	*	*	*	*	*	*	*	*	*	*
1999	0.4	9.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	7.8	*	*	*	*	*	*	*	*	*	*
1999	0.4	10.4	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations of respiratory system (Q30-Q34)												
All origins 3/, both sexes												
2002	0.2	15.8	*	*	*	*	*	*	*	*	*	*
2001	0.2	14.7	*	*	*	*	*	*	*	*	*	*
2000	0.3	18.0	*	*	*	*	*	*	*	*	*	*
1999	0.2	14.9	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.3	17.2	*	*	*	*	*	*	*	*	*	*
2001	0.3	16.7	*	*	*	*	*	*	*	*	*	*
2000	0.3	19.8	*	*	*	*	*	*	*	*	*	*
1999	0.3	17.1	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.2	14.4	*	*	*	*	*	*	*	*	*	*
2001	0.2	12.7	*	*	*	*	*	*	*	*	*	*
2000	0.2	16.2	*	*	*	*	*	*	*	*	*	*
1999	0.2	12.5	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.4	16.2	*	*	*	*	*	*	*	*	*	*
2001	0.3	14.3	*	*	*	*	*	*	*	*	*	*
2000	0.4	16.6	*	*	*	*	*	*	*	*	*	*
1999	0.3	13.5	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.4	18.5	*	*	*	*	*	*	*	*	*	*
2001	0.3	14.6	*	*	*	*	*	*	*	*	*	*
2000	0.4	18.8	*	*	*	*	*	*	*	*	*	*
1999	0.3	14.7	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	13.7	*	*	*	*	*	*	*	*	*	*
2001	0.3	14.0	*	*	*	*	*	*	*	*	*	*
2000	0.3	14.3	*	*	*	*	*	*	*	*	*	*
1999	0.3	12.2	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	15.6	*	*	*	*	*	*	*	*	*	*
2001	0.2	14.6	*	*	*	*	*	*	*	*	*	*
2000	0.2	18.1	*	*	*	*	*	*	*	*	*	*
1999	0.2	15.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.2	16.6	*	*	*	*	*	*	*	*	*	*
2001	0.3	17.1	*	*	*	*	*	*	*	*	*	*
2000	0.3	19.8	*	*	*	*	*	*	*	*	*	*
1999	0.2	17.5	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations of respiratory system (Q30-Q34)												
Non-Hispanic 4/, female												
2002	0.2	14.5	*	*	*	*	*	*	*	*	*	*
2001	0.2	12.0	*	*	*	*	*	*	*	*	*	*
2000	0.2	16.3	*	*	*	*	*	*	*	*	*	*
1999	0.2	12.5	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	13.9	*	*	*	*	*	*	*	*	*	*
2001	0.2	13.7	*	*	*	*	*	*	*	*	*	*
2000	0.2	16.5	*	*	*	*	*	*	*	*	*	*
1999	0.2	14.0	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.2	15.6	*	*	*	*	*	*	*	*	*	*
2001	0.2	16.0	*	*	*	*	*	*	*	*	*	*
2000	0.2	18.0	*	*	*	*	*	*	*	*	*	*
1999	0.2	15.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	12.1	*	*	*	*	*	*	*	*	*	*
2001	0.2	11.2	*	*	*	*	*	*	*	*	*	*
2000	0.2	14.9	*	*	*	*	*	*	*	*	*	*
1999	0.1	12.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.4	23.0	*	*	*	*	*	*	*	*	*	*
2001	0.4	20.3	*	*	*	*	*	*	*	*	*	*
2000	0.4	25.7	*	*	*	*	*	*	*	*	*	*
1999	0.4	20.6	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	22.1	*	*	*	*	*	*	*	*	*	*
2001	0.5	25.2	*	*	*	*	*	*	*	*	*	*
2000	0.5	27.6	*	*	*	*	*	*	*	*	*	*
1999	0.5	26.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	24.0	*	*	*	*	*	*	*	*	*	*
2001	0.2	15.1	*	*	*	*	*	*	*	*	*	*
2000	0.4	23.6	*	*	*	*	*	*	*	*	*	*
1999	0.2	14.3	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)												
All origins 3/, both sexes												
2002	0.9	36.2	0.7	0.2	0.3	0.2	0.2	0.4	0.5	0.8	1.3	2.7
2001	0.9	36.4	0.8	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.5	2.5
2000	0.9	37.9	0.7	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.4	2.8
1999	0.9	38.3	0.9	0.2	0.2	0.2	0.2	0.3	0.5	0.8	1.5	2.6
All origins 3/, male												
2002	1.0	39.2	0.6	0.2	0.3	0.2	0.3	0.5	0.5	0.9	1.5	3.6
2001	1.0	41.1	0.8	0.2	0.3	0.2	0.3	0.3	0.5	0.9	1.5	3.7
2000	1.0	42.1	0.7	0.2	0.2	0.3	0.3	0.4	0.5	0.8	1.4	4.6
1999	1.0	42.5	1.1	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.6	3.0
All origins 3/, female												
2002	0.8	33.1	0.8	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.2	2.3
2001	0.8	31.4	0.8	0.2	0.2	0.2	0.3	0.3	0.5	0.7	1.5	2.0
2000	0.8	33.6	0.6	0.2	0.2	0.2	0.2	0.2	0.6	0.8	1.4	2.1
1999	0.8	33.8	0.8	0.2	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.5
Hispanic, both sexes												
2002	1.1	38.1	0.7	*	*	*	*	*	*	*	*	*
2001	1.0	35.4	0.7	*	*	*	*	*	*	*	*	*
2000	1.1	40.3	0.7	*	*	*	*	*	*	*	*	*
1999	1.1	38.4	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	1.1	38.5	*	*	*	*	*	*	*	*	*	*
2001	1.2	39.4	*	*	*	*	*	*	*	*	*	*
2000	1.3	44.1	*	*	*	*	*	*	*	*	*	*
1999	1.2	41.9	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	1.1	37.8	*	*	*	*	*	*	*	*	*	*
2001	0.9	31.2	*	*	*	*	*	*	*	*	*	*
2000	1.0	36.4	*	*	*	*	*	*	*	*	*	*
1999	1.0	34.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	35.3	0.7	0.2	0.3	0.2	0.3	0.4	0.5	0.8	1.4	2.7
2001	0.9	36.3	0.8	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.5	2.5
2000	0.9	36.9	0.7	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.4	2.8
1999	0.9	37.6	1.0	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.5	2.6
Non-Hispanic 4/, male												
2002	1.0	39.1	0.6	0.2	0.3	0.2	0.3	0.5	0.6	0.9	1.6	3.7
2001	1.0	41.2	0.7	0.2	0.3	0.2	0.3	0.3	0.5	0.8	1.6	3.7
2000	1.0	41.2	0.7	0.2	0.2	0.3	0.3	0.4	0.5	0.8	1.3	4.4
1999	1.0	41.7	1.1	0.2	0.2	0.2	0.3	0.4	0.5	0.8	1.6	3.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)												
Non-Hispanic 4/, female												
2002	0.8	31.4	0.8	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.2	2.2
2001	0.8	31.0	0.8	0.2	0.2	0.2	0.3	0.3	0.5	0.7	1.5	2.0
2000	0.8	32.4	0.6	0.2	0.2	0.2	0.2	0.2	0.6	0.9	1.4	2.1
1999	0.8	33.2	0.8	0.2	0.1	0.2	0.2	0.3	0.4	0.7	1.4	2.5
Non-Hispanic white, both sexes												
2002	0.9	35.8	0.6	0.2	0.3	0.2	0.3	0.4	0.6	0.8	1.4	2.7
2001	0.9	35.1	0.7	0.2	0.2	0.2	0.3	0.3	0.5	0.8	1.5	2.6
2000	0.9	35.3	0.6	0.2	0.2	0.2	0.3	0.3	0.6	0.9	1.4	2.8
1999	0.9	35.1	0.9	0.2	0.2	0.2	0.3	0.4	0.5	0.8	1.5	2.8
Non-Hispanic white, male												
2002	1.0	40.4	0.5	0.2	0.3	0.3	0.3	0.5	0.6	0.9	1.5	3.7
2001	0.9	39.8	0.6	0.2	0.3	0.2	0.3	0.3	0.5	0.9	1.5	3.8
2000	0.9	39.2	0.6	0.2	0.3	0.3	0.3	0.4	0.5	0.8	1.4	4.4
1999	1.0	40.3	1.0	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.6	3.1
Non-Hispanic white, female												
2002	0.8	31.0	0.7	0.2	0.2	0.2	0.2	0.3	0.5	0.7	1.3	2.3
2001	0.8	30.2	0.7	*	*	0.2	0.3	0.3	0.5	0.7	1.5	2.1
2000	0.8	31.1	0.6	0.2	0.2	0.2	0.2	0.3	0.6	0.9	1.4	2.1
1999	0.8	29.5	0.8	0.2	*	0.2	0.2	0.3	0.4	0.8	1.4	2.6
Non-Hispanic black, both sexes												
2002	1.1	37.1	1.1	*	0.4	*	*	*	*	*	*	*
2001	1.2	43.7	0.9	*	*	*	0.4	*	*	*	*	*
2000	1.2	44.6	1.0	0.3	*	*	*	*	*	*	*	*
1999	1.2	47.5	1.4	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	1.3	40.3	*	*	*	*	*	*	*	*	*	*
2001	1.4	49.8	*	*	*	*	*	*	*	*	*	*
2000	1.4	50.5	*	*	*	*	*	*	*	*	*	*
1999	1.4	51.3	1.7	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.9	33.8	*	*	*	*	*	*	*	*	*	*
2001	1.0	37.3	*	*	*	*	*	*	*	*	*	*
2000	1.0	38.4	*	*	*	*	*	*	*	*	*	*
1999	1.0	43.6	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Down's syndrome (Q90)												
All origins 3/, both sexes												
2002	0.3	2.9	0.1	*	0.1	0.1	0.2	0.5	0.9	0.4	*	*
2001	0.3	3.0	0.1	*	0.1	0.2	0.2	0.5	0.9	0.3	*	*
2000	0.3	2.4	*	*	0.1	0.1	0.1	0.5	0.8	0.4	*	*
1999	0.2	2.5	*	*	0.1	0.1	0.1	0.4	0.8	0.2	*	*
All origins 3/, male												
2002	0.3	3.1	*	*	*	0.1	0.2	0.5	1.0	0.3	*	*
2001	0.3	3.4	*	*	*	0.2	0.2	0.5	0.9	0.4	*	*
2000	0.3	2.8	*	*	*	0.1	0.2	0.5	0.8	0.3	*	*
1999	0.2	2.4	*	*	*	0.1	0.1	0.4	0.8	*	*	*
All origins 3/, female												
2002	0.3	2.7	*	*	*	*	0.2	0.4	0.9	0.4	*	*
2001	0.3	2.4	*	*	*	0.2	0.2	0.5	1.0	0.3	*	*
2000	0.3	2.1	*	*	*	0.2	0.1	0.5	0.8	0.4	*	*
1999	0.2	2.5	*	*	*	*	0.1	0.4	0.7	0.3	*	*
Hispanic, both sexes												
2002	0.2	3.2	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	2.9	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	2.8	*	*	0.1	0.1	0.2	0.5	1.0	0.4	*	*
2001	0.3	3.1	*	*	0.1	0.2	0.2	0.5	1.0	0.3	*	*
2000	0.3	2.3	*	*	*	0.1	0.1	0.5	0.8	0.4	*	*
1999	0.2	2.5	*	*	*	0.1	0.1	0.4	0.8	0.2	*	*
Non-Hispanic 4/, male												
2002	0.3	2.7	*	*	*	*	0.2	0.6	1.0	0.4	*	*
2001	0.3	3.5	*	*	*	0.2	0.2	0.5	0.9	0.4	*	*
2000	0.3	2.6	*	*	*	*	0.2	0.5	0.8	0.3	*	*
1999	0.2	2.7	*	*	*	*	0.1	0.4	0.9	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Down's syndrome (Q90)												
Non-Hispanic 4/, female												
2002	0.3	2.8	*	*	*	*	0.2	0.5	0.9	0.4	*	*
2001	0.3	2.7	*	*	*	0.2	0.2	0.5	1.1	0.3	*	*
2000	0.3	2.0	*	*	*	0.1	0.1	0.5	0.8	0.4	*	*
1999	0.2	2.4	*	*	*	*	0.1	0.4	0.7	0.3	*	*
Non-Hispanic white, both sexes												
2002	0.3	3.0	*	*	*	0.1	0.2	0.6	1.1	0.4	*	*
2001	0.3	3.2	*	*	*	0.2	0.2	0.5	1.1	0.3	*	*
2000	0.3	2.2	*	*	*	0.1	0.2	0.5	0.9	0.4	*	*
1999	0.2	2.7	*	*	*	0.1	0.1	0.5	0.9	0.2	*	*
Non-Hispanic white, male												
2002	0.3	2.8	*	*	*	*	0.2	0.6	1.1	0.4	*	*
2001	0.3	3.7	*	*	*	0.2	0.2	0.5	1.0	0.4	*	*
2000	0.3	2.8	*	*	*	*	0.2	0.5	0.8	0.3	*	*
1999	0.3	2.9	*	*	*	*	0.1	0.5	1.0	*	*	*
Non-Hispanic white, female												
2002	0.3	3.2	*	*	*	*	0.3	0.5	1.1	0.4	*	*
2001	0.3	2.7	*	*	*	*	0.2	0.6	1.2	0.3	*	*
2000	0.3	*	*	*	*	0.2	0.1	0.6	0.9	0.5	*	*
1999	0.2	2.5	*	*	*	*	0.1	0.5	0.8	0.2	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	0.8	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)												
All origins 3/, both sexes												
2002	0.4	24.3	0.2	0.1	0.1	*	*	*	*	*	*	*
2001	0.4	22.7	0.3	0.1	*	*	*	*	*	*	*	*
2000	0.4	24.2	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	22.5	0.3	0.1	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.3	20.7	*	*	*	*	*	*	*	*	*	*
2001	0.3	20.1	*	*	*	*	*	*	*	*	*	*
2000	0.3	21.4	*	*	*	*	*	*	*	*	*	*
1999	0.3	18.7	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.4	28.0	*	*	*	*	*	*	*	*	*	*
2001	0.4	25.4	0.5	*	*	*	*	*	*	*	*	*
2000	0.4	27.2	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	26.5	0.3	0.1	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.5	22.2	*	*	*	*	*	*	*	*	*	*
2001	0.6	25.7	*	*	*	*	*	*	*	*	*	*
2000	0.5	23.3	*	*	*	*	*	*	*	*	*	*
1999	0.5	21.8	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.5	20.6	*	*	*	*	*	*	*	*	*	*
2001	0.6	25.0	*	*	*	*	*	*	*	*	*	*
2000	0.5	21.3	*	*	*	*	*	*	*	*	*	*
1999	0.5	20.4	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.6	23.8	*	*	*	*	*	*	*	*	*	*
2001	0.6	26.5	*	*	*	*	*	*	*	*	*	*
2000	0.6	25.5	*	*	*	*	*	*	*	*	*	*
1999	0.6	23.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	24.4	0.2	0.1	*	*	*	*	*	*	*	*
2001	0.3	21.7	0.3	0.1	*	*	*	*	*	*	*	*
2000	0.3	24.2	0.3	*	*	*	*	*	*	*	*	*
1999	0.3	22.4	0.3	0.1	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.3	20.3	*	*	*	*	*	*	*	*	*	*
2001	0.3	18.7	*	*	*	*	*	*	*	*	*	*
2000	0.3	21.1	*	*	*	*	*	*	*	*	*	*
1999	0.3	18.0	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)												
Non-Hispanic 4/, female												
2002	0.4	28.7	*	*	*	*	*	*	*	*	*	*
2001	0.4	24.9	0.5	*	*	*	*	*	*	*	*	*
2000	0.4	27.4	0.3	*	*	*	*	*	*	*	*	*
1999	0.4	27.0	0.3	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.3	25.0	0.2	*	*	*	*	*	*	*	*	*
2001	0.3	21.4	0.3	*	*	*	*	*	*	*	*	*
2000	0.3	23.4	*	*	*	*	*	*	*	*	*	*
1999	0.3	22.4	0.2	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.3	21.4	*	*	*	*	*	*	*	*	*	*
2001	0.3	18.8	*	*	*	*	*	*	*	*	*	*
2000	0.3	20.4	*	*	*	*	*	*	*	*	*	*
1999	0.2	18.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.4	28.8	*	*	*	*	*	*	*	*	*	*
2001	0.3	24.1	0.5	*	*	*	*	*	*	*	*	*
2000	0.3	26.5	*	*	*	*	*	*	*	*	*	*
1999	0.3	26.6	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.5	24.7	*	*	*	*	*	*	*	*	*	*
2001	0.5	24.4	*	*	*	*	*	*	*	*	*	*
2000	0.5	28.4	*	*	*	*	*	*	*	*	*	*
1999	0.5	24.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	20.0	*	*	*	*	*	*	*	*	*	*
2001	0.4	19.5	*	*	*	*	*	*	*	*	*	*
2000	0.5	25.9	*	*	*	*	*	*	*	*	*	*
1999	0.4	19.7	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.5	29.7	*	*	*	*	*	*	*	*	*	*
2001	0.5	29.6	*	*	*	*	*	*	*	*	*	*
2000	0.5	31.0	*	*	*	*	*	*	*	*	*	*
1999	0.5	29.3	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All origins 3/, both sexes												
2002	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7
All origins 3/, male												
2002	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.1
2000	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3
All origins 3/, female												
2002	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4
Hispanic, both sexes												
2002	4.0	56.6	1.3	*	1.3	2.0	3.0	3.9	4.8	7.0	18.7	91.5
2001	5.9	60.7	1.6	0.5	2.8	4.2	6.0	8.3	6.9	9.9	19.3	108.8
2000	5.9	65.6	1.7	0.5	2.1	4.0	6.7	8.0	6.7	9.9	20.5	102.2
1999	3.8	54.9	1.0	*	1.3	2.2	3.0	3.1	3.8	7.0	19.4	87.0
Hispanic, male												
2002	4.6	66.4	1.6	*	1.7	2.9	4.0	4.7	6.3	9.6	23.4	88.7
2001	7.4	67.2	1.8	*	3.9	6.0	8.5	12.6	10.6	14.2	25.0	89.5
2000	7.5	78.6	2.1	*	2.9	5.8	9.9	12.4	9.0	14.3	25.6	106.8
1999	4.4	61.3	*	*	1.9	2.9	4.3	4.7	5.5	8.3	24.0	77.6
Hispanic, female												
2002	3.3	46.4	*	*	0.9	1.0	2.0	3.2	3.5	5.0	15.5	92.8
2001	4.3	53.9	*	0.6	1.5	2.1	3.4	4.1	3.6	6.5	15.4	118.2
2000	4.1	52.1	1.4	*	1.3	2.0	3.2	3.7	4.6	6.5	17.2	99.9
1999	3.1	48.2	*	*	*	1.3	1.6	1.7	2.4	6.0	16.3	91.5
Non-Hispanic 4/, both sexes												
2002	11.3	92.8	1.7	0.4	1.9	3.6	5.6	6.9	7.7	13.5	39.6	211.9
2001	12.1	92.6	1.8	0.4	2.3	3.7	7.0	8.9	9.2	15.4	40.7	209.6
2000	12.0	102.4	1.6	0.4	2.2	3.8	6.5	8.2	9.3	16.8	41.4	211.6
1999	10.3	101.9	1.6	0.4	1.6	2.8	4.7	6.0	8.0	15.0	36.7	190.9
Non-Hispanic 4/, male												
2002	10.9	106.2	1.9	0.4	2.5	4.8	7.3	9.1	10.1	16.3	44.6	178.6
2001	11.8	106.2	2.1	0.5	3.0	4.9	9.0	11.7	12.0	18.2	44.6	178.2
2000	11.9	118.0	1.7	0.5	2.9	5.1	8.6	11.2	12.3	20.7	46.6	185.4
1999	10.0	116.7	1.9	0.4	2.2	3.7	6.1	8.2	10.6	18.7	40.6	169.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic 4/, female												
2002	11.7	78.7	1.4	0.3	1.3	2.3	4.0	4.8	5.4	11.2	36.3	226.3
2001	12.3	78.3	1.5	0.4	1.5	2.6	5.1	6.2	6.6	13.1	38.0	222.7
2000	12.1	86.1	1.5	0.4	1.5	2.6	4.5	5.4	6.6	13.6	38.0	222.3
1999	10.5	86.3	1.4	0.3	1.0	2.0	3.3	3.9	5.7	11.9	34.2	199.5
Non-Hispanic white, both sexes												
2002	11.6	76.0	1.3	0.3	1.8	3.4	5.4	6.1	7.1	13.1	39.4	215.8
2001	12.4	73.1	1.5	0.3	2.2	3.6	6.9	8.5	8.6	14.7	40.4	213.3
2000	12.2	79.7	1.4	0.3	2.1	3.6	6.3	7.6	8.6	15.9	40.9	215.8
1999	10.2	79.3	1.4	0.3	1.4	2.5	4.1	5.2	7.3	14.3	35.7	193.8
Non-Hispanic white, male												
2002	10.7	88.2	1.4	0.3	2.5	4.7	6.9	7.9	9.3	15.7	43.6	181.5
2001	11.8	85.2	1.7	0.4	2.9	4.9	8.8	11.1	10.9	17.3	44.7	180.4
2000	11.8	91.8	1.7	0.4	2.8	4.8	8.5	10.3	11.1	19.3	46.0	189.6
1999	9.6	92.3	1.7	0.3	1.9	3.3	5.5	6.9	9.5	17.7	39.2	171.1
Non-Hispanic white, female												
2002	12.5	63.2	1.3	0.2	1.1	2.2	3.8	4.3	5.1	10.9	36.6	230.6
2001	13.1	60.4	1.3	0.3	1.5	2.4	5.0	5.9	6.4	12.5	37.6	227.0
2000	12.6	66.9	1.2	0.3	1.3	2.3	4.1	4.9	6.2	13.0	37.5	226.4
1999	10.9	65.5	1.0	0.3	0.9	1.8	2.8	3.5	5.2	11.4	33.4	202.8
Non-Hispanic black, both sexes												
2002	12.6	165.4	2.7	0.8	2.6	5.3	8.8	13.9	13.9	20.4	49.9	198.1
2001	12.9	176.9	3.0	0.6	2.7	5.2	9.3	14.0	15.1	24.6	50.3	194.2
2000	13.9	205.6	2.3	0.7	3.2	5.9	9.3	14.8	17.3	29.2	56.1	189.2
1999	13.0	202.0	2.9	0.5	2.7	5.0	9.1	13.2	16.0	24.6	54.8	182.0
Non-Hispanic black, male												
2002	14.4	186.3	3.6	0.9	3.1	7.1	11.7	19.6	20.0	26.4	68.6	182.7
2001	14.2	195.9	3.7	0.8	3.8	6.1	11.4	18.2	23.8	30.7	52.2	186.0
2000	15.7	236.8	2.2	0.9	3.8	7.8	10.9	20.6	25.1	40.0	66.0	163.2
1999	14.9	228.0	2.9	0.6	3.6	6.9	11.6	19.4	23.0	32.8	66.9	180.7
Non-Hispanic black, female												
2002	10.9	143.6	1.9	0.7	2.0	3.7	6.2	9.0	8.9	16.1	39.3	204.0
2001	11.7	156.8	2.2	*	1.7	4.3	7.4	10.3	8.1	20.3	49.2	197.2
2000	12.4	173.4	2.4	*	2.6	4.3	7.8	9.9	11.0	21.6	50.5	198.9
1999	11.2	175.3	2.8	*	1.9	3.2	7.0	7.9	10.3	18.8	48.0	182.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Senility (R54)												
All origins 3/, both sexes												
2002	1.5	*	*	*	*	*	*	*	*	0.8	6.3	72.0
2001	1.5	*	*	*	*	*	*	*	*	0.8	6.4	73.0
2000	1.3	*	*	*	*	*	*	*	*	0.8	6.3	67.2
1999	1.1	*	*	*	*	*	*	*	*	0.7	5.3	56.5
All origins 3/, male												
2002	0.8	*	*	*	*	*	*	*	*	0.8	6.3	56.5
2001	0.8	*	*	*	*	*	*	*	*	0.9	6.4	57.3
2000	0.7	*	*	*	*	*	*	*	*	0.8	6.4	51.7
1999	0.7	*	*	*	*	*	*	*	*	0.9	5.2	45.8
All origins 3/, female												
2002	2.1	*	*	*	*	*	*	*	*	0.8	6.2	78.8
2001	2.1	*	*	*	*	*	*	*	*	0.8	6.3	79.6
2000	1.9	*	*	*	*	*	*	*	*	0.8	6.2	73.6
1999	1.6	*	*	*	*	*	*	*	*	0.5	5.4	60.8
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	27.2
2001	0.2	*	*	*	*	*	*	*	*	*	*	33.1
2000	0.2	*	*	*	*	*	*	*	*	*	*	33.8
1999	0.2	*	*	*	*	*	*	*	*	*	*	26.7
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	29.8
2001	0.3	*	*	*	*	*	*	*	*	*	*	34.4
2000	0.2	*	*	*	*	*	*	*	*	*	*	38.6
1999	0.2	*	*	*	*	*	*	*	*	*	*	32.2
Non-Hispanic 4/, both sexes												
2002	1.7	*	*	*	*	*	*	*	*	0.8	6.4	73.8
2001	1.7	*	*	*	*	*	*	*	*	0.9	6.5	74.3
2000	1.5	*	*	*	*	*	*	*	*	0.8	6.5	68.4
1999	1.3	*	*	*	*	*	*	*	*	0.7	5.4	57.5
Non-Hispanic 4/, male												
2002	0.9	*	*	*	*	*	*	*	*	0.8	6.4	57.9
2001	0.9	*	*	*	*	*	*	*	*	1.0	6.5	58.2
2000	0.8	*	*	*	*	*	*	*	*	0.9	6.5	52.7
1999	0.7	*	*	*	*	*	*	*	*	1.0	5.4	47.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Senility (R54)												
Non-Hispanic 4/, female												
2002	2.4	*	*	*	*	*	*	*	*	0.8	6.5	80.6
2001	2.3	*	*	*	*	*	*	*	*	0.8	6.5	81.0
2000	2.2	*	*	*	*	*	*	*	*	0.8	6.4	74.7
1999	1.8	*	*	*	*	*	*	*	*	0.5	5.5	61.7
Non-Hispanic white, both sexes												
2002	2.0	*	*	*	*	*	*	*	*	0.8	6.7	76.1
2001	1.9	*	*	*	*	*	*	*	*	0.9	6.6	76.6
2000	1.7	*	*	*	*	*	*	*	*	0.8	6.4	70.1
1999	1.5	*	*	*	*	*	*	*	*	0.8	5.3	59.9
Non-Hispanic white, male												
2002	1.1	*	*	*	*	*	*	*	*	0.8	6.6	59.5
2001	1.1	*	*	*	*	*	*	*	*	1.0	6.7	59.4
2000	1.0	*	*	*	*	*	*	*	*	0.9	6.5	54.9
1999	0.8	*	*	*	*	*	*	*	*	1.0	5.2	48.7
Non-Hispanic white, female												
2002	2.8	*	*	*	*	*	*	*	*	0.8	6.7	83.3
2001	2.7	*	*	*	*	*	*	*	*	0.7	6.6	83.7
2000	2.5	*	*	*	*	*	*	*	*	0.7	6.4	76.3
1999	2.1	*	*	*	*	*	*	*	*	0.5	5.4	64.4
Non-Hispanic black, both sexes												
2002	0.7	*	*	*	*	*	*	*	*	*	5.3	56.0
2001	0.8	*	*	*	*	*	*	*	*	*	6.8	57.7
2000	0.8	*	*	*	*	*	*	*	*	1.5	8.7	56.6
1999	0.6	*	*	*	*	*	*	*	*	*	8.0	39.8
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	50.3
2001	0.5	*	*	*	*	*	*	*	*	*	6.3	58.9
2000	0.4	*	*	*	*	*	*	*	*	*	8.9	35.2
1999	0.4	*	*	*	*	*	*	*	*	*	9.4	36.8
Non-Hispanic black, female												
2002	1.0	*	*	*	*	*	*	*	*	*	5.5	58.1
2001	1.0	*	*	*	*	*	*	*	*	*	7.1	57.3
2000	1.2	*	*	*	*	*	*	*	*	*	8.5	64.5
1999	0.8	*	*	*	*	*	*	*	*	*	7.2	40.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)												
All origins 3/, both sexes												
2002	6.0	83.8	1.3	0.2	1.5	2.6	4.4	5.4	5.6	7.6	16.5	58.3
2001	7.1	84.8	1.5	0.3	2.0	3.3	6.0	7.8	7.2	9.6	18.1	62.1
2000	7.3	93.9	1.2	0.3	1.8	3.2	5.7	7.1	7.2	11.0	20.5	73.0
1999	5.9	91.3	1.3	0.3	1.3	2.2	3.7	4.8	5.7	9.2	17.3	66.7
All origins 3/, male												
2002	6.8	96.5	1.5	0.3	2.0	3.5	5.8	7.3	7.7	9.8	20.0	55.0
2001	8.2	96.5	1.8	0.3	2.7	4.5	7.8	10.4	9.6	12.2	21.8	58.1
2000	8.5	108.9	1.4	0.4	2.4	4.4	7.6	10.0	9.8	14.0	24.3	72.1
1999	6.7	104.3	1.4	0.3	1.8	2.9	4.9	6.6	7.8	12.0	20.5	64.4
All origins 3/, female												
2002	5.1	70.4	1.1	0.2	0.9	1.6	3.0	3.5	3.8	5.8	14.1	59.8
2001	6.1	72.4	1.2	0.3	1.2	2.0	4.3	5.3	5.0	7.4	15.7	63.8
2000	6.2	78.0	1.1	0.3	1.2	2.0	3.8	4.4	4.9	8.4	17.9	73.3
1999	5.2	77.7	1.1	0.3	0.7	1.4	2.6	3.0	3.8	6.8	15.2	67.7
Hispanic, both sexes												
2002	3.1	55.2	0.9	*	1.1	1.6	2.3	3.1	3.8	5.0	9.8	33.8
2001	5.1	60.3	1.4	0.4	2.6	3.6	5.4	7.8	6.2	8.1	10.8	40.0
2000	5.1	63.8	1.5	0.4	1.9	3.6	6.1	7.4	6.1	7.7	13.6	39.1
1999	3.0	54.3	0.8	*	1.1	1.9	2.5	2.7	2.9	4.2	11.9	28.8
Hispanic, male												
2002	3.8	65.4	*	*	1.4	2.4	3.2	3.8	5.0	7.1	13.5	45.2
2001	6.7	66.7	1.6	*	3.6	5.4	7.9	11.9	9.9	12.4	16.1	*
2000	6.8	77.3	1.9	*	2.6	5.2	8.9	11.5	8.6	11.0	16.2	52.4
1999	3.8	60.2	*	*	1.6	2.6	3.7	4.1	4.2	5.5	16.7	*
Hispanic, female												
2002	2.3	44.4	*	*	0.7	0.7	1.4	2.3	2.8	3.3	7.3	28.2
2001	3.4	53.7	*	*	1.3	1.5	2.8	3.8	2.9	4.7	7.3	44.7
2000	3.3	49.7	*	*	1.1	1.6	3.0	3.3	3.9	5.1	11.9	32.6
1999	2.3	48.0	*	*	*	1.1	1.1	1.4	*	*	8.7	28.1
Non-Hispanic 4/, both sexes												
2002	6.3	90.8	1.4	0.3	1.5	2.8	4.7	5.5	5.7	7.8	16.7	58.7
2001	7.4	90.4	1.5	0.3	1.8	3.2	6.1	7.7	7.2	9.6	18.4	62.8
2000	7.6	100.7	1.2	0.3	1.8	3.1	5.5	7.0	7.2	11.0	20.7	74.2
1999	6.3	99.7	1.3	0.3	1.3	2.2	3.8	4.9	5.8	9.3	17.5	67.8
Non-Hispanic 4/, male												
2002	7.2	104.1	1.6	0.3	2.0	3.8	6.2	7.5	7.8	9.9	20.1	55.0
2001	8.4	103.4	1.8	0.3	2.4	4.2	7.7	10.2	9.5	12.1	22.0	59.0
2000	8.6	116.2	1.3	0.3	2.3	4.2	7.3	9.7	9.7	14.0	24.6	72.9
1999	7.0	114.1	1.6	0.3	1.8	2.9	5.0	6.6	7.9	12.3	20.5	65.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)												
Non-Hispanic 4/, female												
2002	5.5	76.8	1.2	0.2	1.0	1.8	3.2	3.6	3.8	5.9	14.4	60.3
2001	6.4	76.8	1.2	0.3	1.2	2.1	4.5	5.3	5.1	7.6	16.1	64.4
2000	6.6	84.4	1.0	0.2	1.2	2.0	3.8	4.4	4.9	8.6	18.2	74.7
1999	5.6	84.6	1.1	0.3	0.8	1.5	2.7	3.2	3.9	6.9	15.5	68.7
Non-Hispanic white, both sexes												
2002	6.2	74.5	1.1	0.2	1.5	2.7	4.6	5.0	5.4	7.6	16.7	59.6
2001	7.3	71.1	1.2	0.2	1.8	3.1	6.1	7.5	6.9	9.4	18.3	63.9
2000	7.4	78.2	1.1	0.2	1.7	3.0	5.4	6.6	6.8	10.6	20.7	75.7
1999	6.0	77.4	1.1	0.2	1.2	2.0	3.4	4.3	5.3	9.0	17.1	68.2
Non-Hispanic white, male												
2002	6.9	86.6	1.2	0.2	2.1	3.8	5.9	6.7	7.3	9.6	19.6	56.1
2001	8.2	82.5	1.4	0.2	2.4	4.2	7.7	9.8	8.9	11.7	21.8	60.4
2000	8.3	90.1	1.3	0.3	2.3	4.1	7.3	9.1	9.1	13.1	24.4	73.8
1999	6.5	90.1	1.4	0.2	1.7	2.7	4.5	5.7	7.2	11.7	19.9	64.6
Non-Hispanic white, female												
2002	5.5	61.8	1.0	*	0.9	1.7	3.2	3.3	3.7	5.8	14.7	61.1
2001	6.5	59.0	1.1	0.2	1.2	2.0	4.4	5.1	5.0	7.4	15.9	65.4
2000	6.6	65.7	0.8	0.2	1.1	1.9	3.6	4.1	4.7	8.5	18.2	76.4
1999	5.5	64.0	0.8	0.2	0.6	1.3	2.4	2.9	3.6	6.8	15.3	69.7
Non-Hispanic black, both sexes												
2002	8.6	161.0	2.4	0.6	1.6	3.6	6.7	10.7	9.7	10.9	20.3	57.2
2001	9.0	174.6	2.4	0.5	1.9	4.0	7.0	11.1	10.5	12.9	22.7	56.2
2000	10.0	202.3	1.7	0.5	2.4	4.3	7.5	11.8	12.0	17.1	25.1	67.1
1999	9.3	198.3	2.4	0.4	2.0	3.5	7.1	10.1	11.2	13.7	24.5	71.1
Non-Hispanic black, male												
2002	10.6	181.5	3.0	*	2.0	4.7	9.2	15.5	14.5	15.4	31.1	54.7
2001	10.6	193.1	3.1	*	2.6	4.9	8.3	14.2	16.2	17.6	27.6	49.7
2000	12.1	233.7	1.8	0.7	2.9	5.8	8.7	16.4	17.3	25.3	33.0	71.6
1999	11.4	223.1	2.4	*	2.7	5.0	9.0	15.0	16.6	20.1	31.6	86.8
Non-Hispanic black, female												
2002	6.7	139.8	1.8	*	1.3	2.7	4.4	6.6	5.8	7.8	14.3	58.1
2001	7.6	155.2	1.7	*	1.1	3.2	5.9	8.5	5.9	9.5	19.9	58.6
2000	8.1	169.8	1.7	*	2.0	2.9	6.4	7.8	7.7	11.3	20.7	65.4
1999	7.4	172.8	2.4	*	1.4	2.1	5.4	5.9	6.8	9.1	20.5	65.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sudden infant death syndrome (R95)												
All origins 3/, both sexes												
2002	0.8	56.9	...	...	...	...	...	...	...	...	...	...
2001	0.8	55.4	...	...	...	...	...	...	...	...	...	...
2000	0.9	66.3	...	...	...	...	...	...	...	...	...	...
1999	0.9	69.8	...	...	...	...	...	...	...	...	...	...
All origins 3/, male												
2002	1.0	65.5	...	...	...	...	...	...	...	...	...	...
2001	0.9	63.8	...	...	...	...	...	...	...	...	...	...
2000	1.1	77.4	...	...	...	...	...	...	...	...	...	...
1999	1.1	79.9	...	...	...	...	...	...	...	...	...	...
All origins 3/, female												
2002	0.6	47.9	...	...	...	...	...	...	...	...	...	...
2001	0.6	46.6	...	...	...	...	...	...	...	...	...	...
2000	0.7	54.6	...	...	...	...	...	...	...	...	...	...
1999	0.8	59.1	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2002	0.8	35.4	...	...	...	...	...	...	...	...	...	...
2001	0.7	30.6	...	...	...	...	...	...	...	...	...	...
2000	0.9	39.2	...	...	...	...	...	...	...	...	...	...
1999	0.9	41.3	...	...	...	...	...	...	...	...	...	...
Hispanic, male												
2002	0.9	42.7	...	...	...	...	...	...	...	...	...	...
2001	0.7	34.1	...	...	...	...	...	...	...	...	...	...
2000	1.0	46.4	...	...	...	...	...	...	...	...	...	...
1999	1.0	45.8	...	...	...	...	...	...	...	...	...	...
Hispanic, female												
2002	0.6	27.7	...	...	...	...	...	...	...	...	...	...
2001	0.6	27.0	...	...	...	...	...	...	...	...	...	...
2000	0.7	31.6	...	...	...	...	...	...	...	...	...	...
1999	0.8	36.6	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, both sexes												
2002	0.8	62.3	...	...	...	...	...	...	...	...	...	...
2001	0.8	61.5	...	...	...	...	...	...	...	...	...	...
2000	0.9	72.9	...	...	...	...	...	...	...	...	...	...
1999	0.9	76.4	...	...	...	...	...	...	...	...	...	...
Non-Hispanic 4/, male												
2002	1.0	71.1	...	...	...	...	...	...	...	...	...	...
2001	1.0	71.1	...	...	...	...	...	...	...	...	...	...
2000	1.1	85.1	...	...	...	...	...	...	...	...	...	...
1999	1.1	87.7	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sudden infant death syndrome (R95)												
Non-Hispanic 4/, female												
2002	0.6	53.1	...	...	...	...	...	...	...	...	...	...
2001	0.6	51.4	...	...	...	...	...	...	...	...	...	...
2000	0.7	60.1	...	...	...	...	...	...	...	...	...	...
1999	0.8	64.5	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, both sexes												
2002	0.6	52.1	...	...	...	...	...	...	...	...	...	...
2001	0.6	48.9	...	...	...	...	...	...	...	...	...	...
2000	0.7	58.6	...	...	...	...	...	...	...	...	...	...
1999	0.7	61.4	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, male												
2002	0.7	60.2	...	...	...	...	...	...	...	...	...	...
2001	0.7	57.6	...	...	...	...	...	...	...	...	...	...
2000	0.8	68.6	...	...	...	...	...	...	...	...	...	...
1999	0.9	71.4	...	...	...	...	...	...	...	...	...	...
Non-Hispanic white, female												
2002	0.5	43.6	...	...	...	...	...	...	...	...	...	...
2001	0.5	39.9	...	...	...	...	...	...	...	...	...	...
2000	0.5	48.0	...	...	...	...	...	...	...	...	...	...
1999	0.6	50.9	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, both sexes												
2002	1.9	106.5	...	...	...	...	...	...	...	...	...	...
2001	2.1	119.6	...	...	...	...	...	...	...	...	...	...
2000	2.3	140.2	...	...	...	...	...	...	...	...	...	...
1999	2.4	145.4	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, male												
2002	2.3	119.4	...	...	...	...	...	...	...	...	...	...
2001	2.4	132.1	...	...	...	...	...	...	...	...	...	...
2000	2.8	161.4	...	...	...	...	...	...	...	...	...	...
1999	2.9	162.8	...	...	...	...	...	...	...	...	...	...
Non-Hispanic black, female												
2002	1.6	93.1	...	...	...	...	...	...	...	...	...	...
2001	1.7	106.5	...	...	...	...	...	...	...	...	...	...
2000	1.8	118.4	...	...	...	...	...	...	...	...	...	...
1999	2.0	127.5	...	...	...	...	...	...	...	...	...	...

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other ill-defined and unknown causes of mortality (R96-R99)												
All origins 3/, both sexes												
2002	5.2	26.9	1.3	0.2	1.5	2.6	4.4	5.4	5.6	7.6	16.5	58.3
2001	6.3	29.4	1.5	0.3	2.0	3.3	6.0	7.8	7.2	9.6	18.1	62.1
2000	6.4	27.6	1.2	0.3	1.8	3.2	5.7	7.1	7.2	11.0	20.5	73.0
1999	5.0	21.6	1.3	0.3	1.3	2.2	3.7	4.8	5.7	9.2	17.3	66.7
All origins 3/, male												
2002	5.9	31.1	1.5	0.3	2.0	3.5	5.8	7.3	7.7	9.8	20.0	55.0
2001	7.3	32.7	1.8	0.3	2.7	4.5	7.8	10.4	9.6	12.2	21.8	58.1
2000	7.4	31.5	1.4	0.4	2.4	4.4	7.6	10.0	9.8	14.0	24.3	72.1
1999	5.5	24.4	1.4	0.3	1.8	2.9	4.9	6.6	7.8	12.0	20.5	64.4
All origins 3/, female												
2002	4.5	22.5	1.1	0.2	0.9	1.6	3.0	3.5	3.8	5.8	14.1	59.8
2001	5.5	25.8	1.2	0.3	1.2	2.0	4.3	5.3	5.0	7.4	15.7	63.8
2000	5.5	23.4	1.1	0.3	1.2	2.0	3.8	4.4	4.9	8.4	17.9	73.3
1999	4.5	18.6	1.1	0.3	0.7	1.4	2.6	3.0	3.8	6.8	15.2	67.7
Hispanic, both sexes												
2002	2.3	19.8	0.9	*	1.1	1.6	2.3	3.1	3.8	5.0	9.8	33.8
2001	4.4	29.7	1.4	0.4	2.6	3.6	5.4	7.8	6.2	8.1	10.8	40.0
2000	4.2	24.6	1.5	0.4	1.9	3.6	6.1	7.4	6.1	7.7	13.6	39.1
1999	2.1	13.0	0.8	*	1.1	1.9	2.5	2.7	2.9	4.2	11.9	28.8
Hispanic, male												
2002	2.9	22.7	*	*	1.4	2.4	3.2	3.8	5.0	7.1	13.5	45.2
2001	6.0	32.6	1.6	*	3.6	5.4	7.9	11.9	9.9	12.4	16.1	*
2000	5.8	30.9	1.9	*	2.6	5.2	8.9	11.5	8.6	11.0	16.2	52.4
1999	2.7	14.5	*	*	1.6	2.6	3.7	4.1	4.2	5.5	16.7	*
Hispanic, female												
2002	1.7	16.7	*	*	0.7	0.7	1.4	2.3	2.8	3.3	7.3	28.2
2001	2.8	26.7	*	*	1.3	1.5	2.8	3.8	2.9	4.7	7.3	44.7
2000	2.6	18.1	*	*	1.1	1.6	3.0	3.3	3.9	5.1	11.9	32.6
1999	1.5	11.4	*	*	*	1.1	1.1	1.4	*	*	8.7	28.1
Non-Hispanic 4/, both sexes												
2002	5.5	28.5	1.4	0.3	1.5	2.8	4.7	5.5	5.7	7.8	16.7	58.7
2001	6.6	28.9	1.5	0.3	1.8	3.2	6.1	7.7	7.2	9.6	18.4	62.8
2000	6.7	27.7	1.2	0.3	1.8	3.1	5.5	7.0	7.2	11.0	20.7	74.2
1999	5.3	23.3	1.3	0.3	1.3	2.2	3.8	4.9	5.8	9.3	17.5	67.8
Non-Hispanic 4/, male												
2002	6.3	33.0	1.6	0.3	2.0	3.8	6.2	7.5	7.8	9.9	20.1	55.0
2001	7.4	32.3	1.8	0.3	2.4	4.2	7.7	10.2	9.5	12.1	22.0	59.0
2000	7.5	31.1	1.3	0.3	2.3	4.2	7.3	9.7	9.7	14.0	24.6	72.9
1999	5.8	26.3	1.6	0.3	1.8	2.9	5.0	6.6	7.9	12.3	20.5	65.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other ill-defined and unknown causes of mortality (R96-R99)												
Non-Hispanic 4/, female												
2002	4.8	23.7	1.2	0.2	1.0	1.8	3.2	3.6	3.8	5.9	14.4	60.3
2001	5.8	25.4	1.2	0.3	1.2	2.1	4.5	5.3	5.1	7.6	16.1	64.4
2000	5.9	24.3	1.0	0.2	1.2	2.0	3.8	4.4	4.9	8.6	18.2	74.7
1999	4.8	20.1	1.1	0.3	0.8	1.5	2.7	3.2	3.9	6.9	15.5	68.7
Non-Hispanic white, both sexes												
2002	5.6	22.4	1.1	0.2	1.5	2.7	4.6	5.0	5.4	7.6	16.7	59.6
2001	6.7	22.1	1.2	0.2	1.8	3.1	6.1	7.5	6.9	9.4	18.3	63.9
2000	6.8	19.6	1.1	0.2	1.7	3.0	5.4	6.6	6.8	10.6	20.7	75.7
1999	5.2	16.0	1.1	0.2	1.2	2.0	3.4	4.3	5.3	9.0	17.1	68.2
Non-Hispanic white, male												
2002	6.2	26.4	1.2	0.2	2.1	3.8	5.9	6.7	7.3	9.6	19.6	56.1
2001	7.5	24.9	1.4	0.2	2.4	4.2	7.7	9.8	8.9	11.7	21.8	60.4
2000	7.5	21.5	1.3	0.3	2.3	4.1	7.3	9.1	9.1	13.1	24.4	73.8
1999	5.6	18.8	1.4	0.2	1.7	2.7	4.5	5.7	7.2	11.7	19.9	64.6
Non-Hispanic white, female												
2002	5.0	18.2	1.0	*	0.9	1.7	3.2	3.3	3.7	5.8	14.7	61.1
2001	6.0	19.2	1.1	0.2	1.2	2.0	4.4	5.1	5.0	7.4	15.9	65.4
2000	6.1	17.7	0.8	0.2	1.1	1.9	3.6	4.1	4.7	8.5	18.2	76.4
1999	4.9	13.1	0.8	0.2	0.6	1.3	2.4	2.9	3.6	6.8	15.3	69.7
Non-Hispanic black, both sexes												
2002	6.6	54.6	2.4	0.6	1.6	3.6	6.7	10.7	9.7	10.9	20.3	57.2
2001	7.0	55.0	2.4	0.5	1.9	4.0	7.0	11.1	10.5	12.9	22.7	56.2
2000	7.7	62.1	1.7	0.5	2.4	4.3	7.5	11.8	12.0	17.1	25.1	67.1
1999	6.9	52.9	2.4	0.4	2.0	3.5	7.1	10.1	11.2	13.7	24.5	71.1
Non-Hispanic black, male												
2002	8.3	62.1	3.0	*	2.0	4.7	9.2	15.5	14.5	15.4	31.1	54.7
2001	8.1	60.9	3.1	*	2.6	4.9	8.3	14.2	16.2	17.6	27.6	49.7
2000	9.3	72.3	1.8	0.7	2.9	5.8	8.7	16.4	17.3	25.3	33.0	71.6
1999	8.5	60.3	2.4	*	2.7	5.0	9.0	15.0	16.6	20.1	31.6	86.8
Non-Hispanic black, female												
2002	5.1	46.7	1.8	*	1.3	2.7	4.4	6.6	5.8	7.8	14.3	58.1
2001	5.9	48.7	1.7	*	1.1	3.2	5.9	8.5	5.9	9.5	19.9	58.6
2000	6.3	51.4	1.7	*	2.0	2.9	6.4	7.8	7.7	11.3	20.7	65.4
1999	5.4	45.3	2.4	*	1.4	2.1	5.4	5.9	6.8	9.1	20.5	65.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)												
All origins 3/, both sexes												
2002	2.9	1.9	0.3	0.1	0.4	0.7	0.9	1.3	1.8	4.7	16.0	77.7
2001	2.7	1.8	0.3	0.1	0.4	0.6	0.9	1.2	1.8	4.7	15.3	71.3
2000	2.6	1.8	0.4	0.1	0.4	0.7	0.9	1.2	2.0	4.8	13.9	67.9
1999	2.5	1.9	0.3	0.1	0.3	0.6	0.8	1.1	2.1	4.8	13.5	64.5
All origins 3/, male												
2002	2.4	1.9	0.4	0.1	0.5	0.9	1.1	1.5	2.2	5.3	17.6	64.0
2001	2.3	2.3	0.3	0.2	0.5	0.7	1.2	1.5	2.3	4.9	15.6	59.7
2000	2.3	1.7	0.3	0.1	0.5	0.8	1.3	1.5	2.4	5.7	15.3	58.9
1999	2.1	2.3	0.3	0.1	0.4	0.7	1.1	1.4	2.5	5.3	14.4	56.1
All origins 3/, female												
2002	3.5	1.9	*	*	0.3	0.5	0.8	1.1	1.4	4.2	15.0	83.6
2001	3.2	1.2	0.3	0.1	0.3	0.5	0.7	0.8	1.3	4.6	15.1	76.2
2000	3.0	1.8	0.4	0.1	0.3	0.5	0.6	0.9	1.6	4.0	13.1	71.6
1999	2.9	1.5	*	*	0.3	0.5	0.6	0.7	1.7	4.4	12.9	67.9
Hispanic, both sexes												
2002	0.7	*	*	*	*	0.4	0.7	0.9	*	1.9	6.7	30.5
2001	0.6	*	*	*	*	0.6	0.6	*	*	*	5.2	35.6
2000	0.6	*	*	*	*	0.4	0.6	0.6	*	2.2	5.5	29.2
1999	0.6	*	*	*	*	*	0.5	*	*	2.6	5.0	31.6
Hispanic, male												
2002	0.7	*	*	*	*	0.5	0.8	*	*	*	*	*
2001	0.6	*	*	*	*	0.5	*	*	*	*	*	*
2000	0.7	*	*	*	*	*	0.9	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.8	*	*	*	*	*	*	*	*	*	6.7	34.8
2001	0.7	*	*	*	*	*	*	*	*	*	*	39.1
2000	0.6	*	*	*	*	*	*	*	*	*	*	28.7
1999	0.6	*	*	*	*	*	*	*	*	*	*	31.2
Non-Hispanic 4/, both sexes												
2002	3.3	2.0	0.3	0.1	0.4	0.8	1.0	1.3	1.9	4.9	16.5	79.4
2001	3.1	2.1	0.3	0.1	0.5	0.6	1.0	1.2	1.9	4.9	15.7	72.5
2000	2.9	1.7	0.4	0.1	0.4	0.7	1.0	1.2	2.1	4.9	14.2	69.1
1999	2.8	2.2	0.3	0.1	0.3	0.6	0.9	1.1	2.1	4.9	13.8	65.6
Non-Hispanic 4/, male												
2002	2.7	2.1	*	0.1	0.5	1.1	1.1	1.5	2.3	5.5	18.1	65.8
2001	2.5	2.8	*	0.2	0.6	0.7	1.3	1.6	2.4	5.1	16.1	60.9
2000	2.5	1.8	0.4	0.1	0.5	0.9	1.3	1.6	2.6	5.8	15.5	59.9
1999	2.3	2.6	0.3	0.1	0.4	0.7	1.1	1.5	2.6	5.4	14.7	56.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)												
Non-Hispanic 4/, female												
2002	3.9	1.9	*	*	0.3	0.6	0.8	1.1	1.5	4.4	15.4	85.3
2001	3.6	1.5	0.3	*	0.3	0.5	0.7	0.9	1.4	4.7	15.5	77.3
2000	3.3	1.7	0.4	0.1	0.3	0.6	0.7	0.9	1.6	4.2	13.4	72.9
1999	3.2	1.8	*	*	0.3	0.5	0.6	0.8	1.7	4.5	13.2	69.1
Non-Hispanic white, both sexes												
2002	3.5	1.5	0.2	0.1	0.3	0.7	0.8	1.1	1.6	4.7	16.1	80.1
2001	3.2	2.0	0.3	0.1	0.4	0.5	0.8	1.0	1.6	4.4	15.5	72.8
2000	3.0	1.5	0.4	0.1	0.4	0.6	0.9	1.0	1.8	4.5	13.8	70.0
1999	2.8	1.9	0.3	0.1	0.3	0.5	0.7	0.9	1.9	4.5	13.3	65.7
Non-Hispanic white, male												
2002	2.7	*	*	*	0.4	0.9	1.0	1.2	2.0	5.2	17.4	66.0
2001	2.5	2.7	*	0.2	0.5	0.6	1.0	1.3	1.9	4.5	16.2	60.6
2000	2.5	*	*	*	0.5	0.7	1.2	1.2	2.0	5.3	15.1	60.8
1999	2.3	2.2	*	*	0.3	0.6	0.9	1.2	2.3	5.0	14.1	57.9
Non-Hispanic white, female												
2002	4.2	*	*	*	0.2	0.5	0.6	1.0	1.4	4.3	15.3	86.1
2001	3.9	*	*	*	0.3	0.4	0.6	0.8	1.3	4.4	15.0	77.9
2000	3.5	*	*	*	0.3	0.4	0.5	0.8	1.5	3.8	13.0	73.7
1999	3.3	*	*	*	0.2	0.4	0.5	0.6	1.6	4.1	12.7	68.8
Non-Hispanic black, both sexes												
2002	3.3	4.3	*	*	0.9	1.7	2.1	3.2	4.2	8.3	24.2	85.0
2001	3.2	*	*	*	0.9	1.2	2.2	2.7	4.4	10.6	20.8	80.2
2000	3.1	*	*	*	0.7	1.6	1.7	3.1	5.2	10.6	22.3	65.5
1999	3.1	3.7	*	*	0.7	1.5	2.1	3.1	4.6	10.1	22.3	71.1
Non-Hispanic black, male												
2002	3.4	*	*	*	1.1	2.4	2.5	4.0	5.5	9.7	32.5	77.7
2001	3.2	*	*	*	1.2	1.3	3.1	3.8	7.4	12.3	18.3	77.4
2000	3.1	*	*	*	0.9	2.0	2.1	4.2	7.6	13.5	24.1	56.4
1999	3.1	*	*	*	0.9	1.9	2.6	4.3	6.1	11.6	26.0	57.0
Non-Hispanic black, female												
2002	3.3	*	*	*	0.8	1.1	1.8	2.4	3.1	7.4	19.6	87.8
2001	3.1	*	*	*	*	1.1	1.5	1.8	2.0	9.3	22.2	81.3
2000	3.1	*	*	*	*	1.4	1.4	2.1	3.3	8.5	21.4	68.9
1999	3.1	*	*	*	*	1.1	1.6	2.0	3.4	9.1	20.3	76.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)												
All origins 3/, both sexes												
2002	56.9	33.8	13.7	8.4	62.2	57.9	63.7	60.8	50.7	63.9	128.4	308.3
2001 5/	56.2	35.1	14.3	8.5	60.6	58.2	63.4	58.9	49.9	62.7	128.7	310.1
2000	53.8	34.5	14.6	9.1	60.0	54.1	59.2	54.8	48.6	60.9	122.6	310.5
1999	54.2	33.9	15.3	9.4	59.7	55.2	58.7	53.2	48.2	64.4	128.6	318.2
All origins 3/, male												
2002	79.8	37.6	16.3	10.4	95.1	89.9	93.3	89.2	75.3	92.9	180.3	414.8
2001 5/	79.3	38.9	16.8	10.7	93.7	90.5	93.9	88.1	73.0	90.0	183.4	429.1
2000	75.8	38.1	17.0	11.4	91.9	84.5	88.8	82.5	71.1	86.9	175.5	438.2
1999	76.2	38.3	17.3	11.8	90.8	85.7	87.7	80.1	71.5	92.9	183.6	454.8
All origins 3/, female												
2002	34.8	29.8	11.1	6.3	27.5	25.1	34.3	33.3	28.0	39.8	94.0	262.0
2001 5/	34.0	31.2	11.8	6.2	25.8	25.1	33.1	30.8	28.5	40.1	92.7	260.3
2000	32.5	30.8	12.1	6.7	26.5	23.3	29.9	28.1	27.9	39.4	88.1	258.5
1999	33.0	29.2	13.3	6.9	27.0	24.3	30.1	27.3	26.8	41.0	92.8	263.2
Hispanic, both sexes												
2002	40.5	23.4	14.1	6.9	61.0	48.0	48.7	46.9	42.0	52.7	94.9	170.7
2001 5/	41.2	26.8	13.0	6.8	58.1	50.8	52.2	48.4	42.4	54.4	91.4	193.8
2000	39.5	23.3	12.6	7.3	56.6	47.4	49.4	46.2	40.6	53.8	92.4	208.3
1999	40.1	26.2	14.3	7.5	55.8	49.3	50.5	46.5	41.4	54.5	88.3	209.1
Hispanic, male												
2002	61.8	27.9	17.2	8.3	95.6	75.5	74.6	72.1	66.7	81.2	134.6	262.7
2001 5/	62.3	26.9	15.6	8.2	90.7	79.0	81.8	74.6	65.2	79.9	136.4	264.7
2000	60.2	24.6	13.8	8.5	89.0	75.3	77.5	72.9	63.6	81.5	132.3	298.3
1999	61.5	29.0	15.8	9.4	87.0	77.8	80.1	74.4	65.9	84.6	133.0	312.7
Hispanic, female												
2002	17.9	18.6	10.8	5.4	20.7	15.3	20.5	21.8	19.9	30.0	67.7	125.2
2001 5/	18.7	26.7	10.4	5.4	20.0	17.9	20.3	22.8	22.2	34.3	61.0	159.2
2000	17.5	22.0	11.4	6.1	18.4	15.1	19.3	20.2	20.3	32.0	65.7	164.2
1999	17.6	23.3	12.7	5.5	19.3	16.3	19.0	19.3	19.8	31.0	58.6	159.1
Non-Hispanic 4/, both sexes												
2002	59.1	36.3	13.6	8.7	62.2	59.7	65.5	61.8	51.1	64.3	129.6	313.0
2001 5/	58.1	36.9	14.6	8.9	60.8	59.3	64.5	59.5	50.1	62.9	130.0	313.6
2000	55.5	37.0	15.0	9.4	60.4	55.2	60.1	55.2	48.9	61.0	123.6	313.4
1999	55.7	35.5	15.5	9.7	60.2	56.0	59.3	53.3	48.3	64.7	129.7	321.0
Non-Hispanic 4/, male												
2002	82.2	39.8	16.0	10.7	94.5	92.9	95.7	90.3	75.3	93.0	181.8	420.1
2001 5/	81.4	41.5	17.0	11.2	93.9	92.5	94.9	88.6	73.0	90.1	185.0	434.9
2000	77.7	41.2	17.7	12.0	92.1	85.9	89.7	82.7	71.0	86.7	176.9	442.7
1999	77.8	40.5	17.5	12.2	91.2	86.7	88.1	79.8	71.2	92.9	184.9	458.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)												
Non-Hispanic 4/, female												
2002	37.1	32.7	11.1	6.5	28.7	26.9	36.0	34.3	28.6	40.4	95.0	266.8
2001 5/	35.9	32.1	12.1	6.4	26.8	26.3	34.7	31.4	28.8	40.3	93.9	263.1
2000	34.4	32.7	12.2	6.8	27.9	24.6	31.1	28.7	28.4	39.8	88.8	261.1
1999	34.8	30.2	13.3	7.1	28.3	25.5	31.2	27.8	27.2	41.5	93.9	265.8
Non-Hispanic white, both sexes												
2002	59.8	26.9	11.6	7.7	58.4	55.8	65.8	61.6	51.0	64.6	132.6	324.9
2001 5/	58.4	28.9	13.1	8.2	56.3	55.7	64.1	58.6	49.6	62.7	132.6	323.5
2000	55.2	28.4	13.0	8.9	55.6	50.6	58.9	53.7	48.3	60.3	125.5	323.5
1999	55.2	27.9	13.1	8.9	54.8	51.6	57.9	51.6	47.0	64.0	130.6	329.8
Non-Hispanic white, male												
2002	81.2	31.3	13.5	9.5	86.0	84.1	94.9	88.9	74.2	92.5	184.9	433.3
2001 5/	80.1	32.5	15.6	10.1	84.1	84.9	93.6	86.3	71.3	89.5	188.0	447.7
2000	75.5	33.3	15.6	11.2	82.0	76.8	87.1	79.3	69.1	84.9	178.8	456.7
1999	75.3	33.3	14.9	11.1	80.1	78.2	85.1	76.2	68.3	91.3	185.8	470.1
Non-Hispanic white, female												
2002	39.3	22.2	9.7	5.9	29.7	27.1	36.6	34.7	29.0	40.8	97.5	278.1
2001 5/	37.6	25.2	10.4	6.1	27.4	26.0	34.5	31.5	29.1	39.9	95.9	271.8
2000	35.8	23.2	10.2	6.4	28.2	23.9	30.7	28.5	28.7	39.6	90.4	269.7
1999	36.0	22.3	11.3	6.6	28.6	24.7	30.7	27.5	26.8	41.1	94.5	273.8
Non-Hispanic black, both sexes												
2002	65.4	74.5	21.6	12.5	85.0	92.2	77.7	75.6	59.5	66.1	107.6	201.4
2001 5/	65.8	71.3	22.1	12.0	86.9	88.1	79.1	76.6	61.1	71.2	111.8	217.3
2000	65.6	73.7	23.7	12.5	88.7	87.9	78.0	74.7	59.4	73.1	109.6	215.0
1999	67.0	70.5	25.6	13.8	90.2	86.8	77.9	74.0	62.5	76.9	128.3	240.9
Non-Hispanic black, male												
2002	102.4	74.8	26.6	16.1	142.6	160.4	120.5	119.5	99.0	105.3	168.2	308.5
2001 5/	102.7	80.4	25.1	16.4	148.4	151.1	121.4	124.3	99.4	107.8	171.0	307.2
2000	102.8	75.8	26.3	16.1	149.5	151.2	123.8	121.9	98.1	115.7	169.1	320.6
1999	104.5	74.8	28.7	17.9	152.5	147.8	123.2	122.5	102.4	120.7	190.4	370.8
Non-Hispanic black, female												
2002	31.8	74.1	16.3	8.7	27.2	30.6	39.9	37.9	27.9	38.3	73.3	160.5
2001 5/	32.3	61.8	19.1	7.4	25.7	31.2	41.7	35.6	30.3	45.3	78.5	183.9
2000	31.8	71.5	21.0	8.7	28.8	30.8	37.6	34.1	28.2	43.2	76.1	175.7
1999	33.0	66.1	22.5	9.5	28.7	31.9	37.8	32.5	30.5	46.0	93.5	192.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All origins 3/, both sexes												
2002	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4
All origins 3/, male												
2002	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001	47.3	26.2	13.3	8.6	52.9	45.5	51.8	50.7	43.0	57.3	133.4	358.5
2000	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4
All origins 3/, female												
2002	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8
Hispanic, both sexes												
2002	26.1	15.2	10.9	5.4	35.5	27.6	31.7	32.0	28.7	39.5	77.4	153.0
2001	25.8	18.8	10.3	5.4	32.6	28.3	32.3	31.8	30.5	40.1	73.9	166.3
2000	25.0	15.3	10.4	5.6	31.8	27.3	31.5	30.4	28.8	38.8	72.1	184.5
1999	25.5	17.7	11.9	5.6	30.8	28.4	32.6	31.2	28.5	42.5	69.6	182.4
Hispanic, male												
2002	38.5	19.5	13.5	6.6	53.5	43.0	48.5	48.7	43.5	58.7	105.9	224.2
2001	37.7	18.0	12.7	6.2	49.2	43.9	50.2	48.9	45.0	55.3	105.5	198.1
2000	36.9	17.0	11.6	6.1	48.2	42.9	49.6	48.4	43.2	57.2	93.9	251.9
1999	37.7	20.7	13.5	6.8	46.2	44.1	51.3	50.2	43.5	63.3	94.9	258.8
Hispanic, female												
2002	12.8	10.8	8.2	4.1	14.5	9.5	13.4	15.3	15.3	24.2	57.9	117.7
2001	13.2	19.7	7.8	4.5	13.3	10.1	13.0	14.9	17.7	28.2	52.6	150.8
2000	12.4	13.5	9.2	5.2	12.5	9.2	12.2	12.8	16.0	24.4	57.4	151.3
1999	12.6	14.6	10.3	4.2	12.8	10.1	12.6	12.8	15.2	26.2	52.7	145.6
Non-Hispanic 4/, both sexes												
2002	38.5	25.5	10.4	6.9	38.3	32.2	37.8	36.8	31.4	44.4	102.1	279.8
2001	36.9	25.4	11.3	7.2	36.6	30.1	35.6	34.0	30.0	42.8	101.9	279.8
2000	36.0	25.0	12.2	7.5	36.7	29.8	34.2	32.5	30.9	41.9	95.8	276.1
1999	36.2	23.4	12.4	7.9	36.0	29.6	33.7	31.6	30.5	44.5	101.3	285.0
Non-Hispanic 4/, male												
2002	50.3	28.6	12.7	8.4	55.0	48.6	54.2	53.4	44.9	60.3	132.3	351.2
2001	48.5	28.0	13.4	9.0	53.5	45.6	51.7	50.4	42.5	57.1	134.3	364.5
2000	47.3	27.7	14.5	9.4	52.7	45.2	50.8	48.6	43.8	55.5	125.2	363.1
1999	47.4	26.6	14.5	9.9	51.4	44.8	49.7	47.3	43.5	59.7	133.2	378.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Non-Hispanic 4/, female												
2002	27.3	22.3	8.1	5.2	21.1	15.9	21.8	20.9	18.8	31.0	82.2	249.0
2001	26.0	22.6	9.2	5.3	19.3	14.7	19.9	18.3	18.5	30.8	80.6	244.6
2000	25.2	22.2	9.7	5.6	20.3	14.5	18.1	17.0	18.9	30.6	76.7	240.9
1999	25.5	20.0	10.2	5.9	20.2	14.6	18.0	16.5	18.5	32.0	80.6	247.5
Non-Hispanic white, both sexes												
2002	40.6	19.8	9.4	6.3	41.6	33.7	38.7	36.3	30.6	43.9	104.4	290.9
2001	38.5	20.9	10.9	6.6	39.0	31.1	35.9	32.8	29.0	42.4	103.7	289.3
2000	37.3	20.1	11.1	7.2	39.2	30.0	34.2	31.0	30.0	41.0	97.2	285.6
1999	37.2	19.0	11.1	7.4	38.1	29.9	33.5	30.0	28.9	43.3	101.7	293.3
Non-Hispanic white, male												
2002	52.0	23.3	11.1	7.7	59.2	50.3	55.0	52.1	43.0	58.8	133.5	361.6
2001	49.7	22.9	13.2	8.1	56.6	46.8	51.8	48.0	40.3	56.1	135.2	374.7
2000	48.1	22.9	13.4	8.9	55.9	45.2	50.4	45.8	41.6	53.6	125.2	373.8
1999	47.8	22.0	12.9	9.1	53.6	44.9	49.0	44.2	40.7	57.3	132.7	387.1
Non-Hispanic white, female												
2002	29.6	16.0	7.5	4.8	23.2	16.8	22.4	20.6	18.8	31.2	84.9	260.3
2001	27.8	18.7	8.5	5.0	20.7	15.2	19.9	17.9	18.3	30.7	82.8	253.8
2000	26.9	17.1	8.7	5.3	21.9	14.6	18.0	16.3	19.1	30.4	78.8	250.0
1999	27.0	15.9	9.3	5.6	22.0	14.7	17.9	15.9	17.7	31.5	81.5	255.9
Non-Hispanic black, both sexes												
2002	34.0	50.1	14.6	9.4	28.4	32.5	39.9	47.6	41.9	49.7	84.6	175.4
2001	34.4	44.5	14.2	9.7	29.6	31.1	40.9	48.5	42.4	51.3	89.6	183.9
2000	34.4	46.5	16.8	9.6	29.6	33.5	40.6	47.8	41.3	53.0	85.3	183.2
1999	36.1	44.1	18.2	10.9	30.5	33.2	40.7	48.3	45.1	58.6	103.8	208.9
Non-Hispanic black, male												
2002	49.0	52.1	18.9	11.9	42.6	52.2	59.7	73.3	69.2	77.2	131.3	262.5
2001	49.5	50.4	15.8	13.1	44.5	48.8	61.3	78.6	68.2	74.9	138.3	256.4
2000	50.3	49.5	19.4	12.3	44.2	53.5	63.3	76.7	69.4	80.4	133.3	277.1
1999	52.3	48.5	21.8	14.3	46.4	53.0	63.0	79.4	71.8	89.6	151.0	318.5
Non-Hispanic black, female												
2002	20.3	48.0	10.1	6.8	14.2	14.7	22.5	25.6	20.0	30.2	58.2	142.1
2001	20.6	38.3	12.4	6.2	14.8	15.1	22.7	22.7	21.6	34.6	62.2	157.0
2000	20.1	43.3	14.2	6.7	15.1	15.5	20.5	23.0	18.7	33.7	58.4	148.3
1999	21.4	39.6	14.5	7.4	14.8	15.4	21.1	21.8	23.7	36.8	77.4	167.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
All origins 3/, both sexes												
2002	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
All origins 3/, male												
2002	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
All origins 3/, female												
2002	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3
Hispanic, both sexes												
2002	15.3	3.1	5.8	3.8	27.6	18.3	16.0	15.2	15.2	18.4	26.2	25.5
2001	15.6	5.3	5.3	3.8	26.0	18.9	17.3	16.4	17.9	20.3	30.3	23.8
2000	14.8	5.4	5.2	4.1	24.6	17.3	16.2	16.2	15.7	20.4	23.9	25.2
1999	14.4	5.3	5.5	3.8	23.1	17.8	15.4	15.0	15.5	22.0	25.0	25.3
Hispanic, male												
2002	22.4	*	7.1	4.6	40.4	27.5	23.9	21.6	22.1	28.5	37.3	45.2
2001	22.8	*	6.0	4.3	38.4	28.9	25.9	23.2	24.6	28.6	51.4	43.8
2000	21.6	6.1	5.6	4.5	36.1	26.4	24.1	24.1	23.5	29.5	36.4	44.3
1999	20.8	5.7	5.8	4.5	33.3	26.7	23.2	22.6	21.0	31.6	40.1	*
Hispanic, female												
2002	7.7	*	4.5	2.9	12.6	7.3	7.5	8.7	8.9	10.3	18.5	*
2001	8.0	6.7	4.6	3.2	11.6	7.1	8.0	9.7	12.0	13.9	16.0	*
2000	7.6	*	4.8	3.7	11.0	6.8	7.8	8.5	8.7	13.3	15.5	*
1999	7.6	*	5.2	3.1	11.0	7.6	7.0	7.7	10.6	14.5	14.9	*
Non-Hispanic 4/, both sexes												
2002	16.9	3.2	3.7	4.3	29.3	18.8	17.2	16.4	15.6	18.3	26.9	28.9
2001	16.7	3.4	4.2	4.6	28.2	18.8	17.3	15.8	15.0	17.7	27.7	29.0
2000	16.8	4.3	4.3	4.8	28.8	18.7	17.0	15.7	15.9	17.8	27.1	31.7
1999	16.9	4.9	4.4	5.0	28.4	18.5	16.9	15.4	16.0	19.2	28.3	32.2
Non-Hispanic 4/, male												
2002	23.8	3.4	4.2	5.0	40.6	28.1	24.6	24.1	22.4	24.9	37.5	53.2
2001	23.5	3.2	4.7	5.5	39.7	28.1	24.8	23.2	21.3	24.0	39.2	52.9
2000	23.4	4.5	4.8	5.6	39.8	27.8	24.7	22.8	22.2	23.9	38.0	60.4
1999	23.3	5.1	4.9	6.0	39.0	27.2	24.0	22.2	22.4	25.6	40.4	64.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Non-Hispanic 4/, female												
2002	10.4	2.9	3.1	3.6	17.7	9.5	10.0	9.0	9.3	12.7	19.8	18.4
2001	10.2	3.6	3.6	3.6	16.3	9.5	10.0	8.8	9.1	12.5	20.2	19.1
2000	10.5	4.2	3.8	3.9	17.4	9.7	9.4	8.9	10.1	12.8	20.0	20.1
1999	10.7	4.7	3.9	4.0	17.5	10.0	9.9	8.8	10.0	13.9	20.5	19.3
Non-Hispanic white, both sexes												
2002	17.5	2.5	3.3	4.3	31.8	19.0	17.3	16.4	15.5	18.3	27.4	29.7
2001	17.0	3.1	3.8	4.4	29.9	19.0	17.4	15.6	14.7	17.6	28.3	29.9
2000	17.1	3.5	3.8	4.8	30.9	18.5	17.0	15.4	15.8	17.6	27.6	32.4
1999	17.1	4.3	3.7	5.0	30.3	18.6	16.7	15.0	15.6	19.1	28.6	32.5
Non-Hispanic white, male												
2002	24.3	2.9	3.6	5.0	43.6	28.1	24.6	24.0	22.0	24.5	37.5	54.0
2001	23.8	2.9	4.4	5.2	41.9	28.2	24.6	22.7	20.8	23.7	39.6	54.3
2000	23.7	3.8	4.3	5.7	42.5	27.1	24.5	22.2	21.6	23.2	38.1	62.3
1999	23.5	4.9	4.0	5.9	41.2	26.9	23.6	21.6	21.7	25.2	40.3	64.6
Non-Hispanic white, female												
2002	10.9	2.1	2.9	3.5	19.4	9.8	10.1	8.9	9.4	13.0	20.7	19.2
2001	10.6	3.3	3.2	3.5	17.4	9.7	10.2	8.6	9.1	12.5	20.8	19.8
2000	10.9	3.2	3.4	4.0	18.9	9.7	9.5	8.6	10.4	12.8	20.7	20.4
1999	11.0	3.6	3.4	4.0	19.1	10.2	9.9	8.6	9.9	13.9	20.9	19.7
Non-Hispanic black, both sexes												
2002	15.7	5.6	5.0	4.7	21.6	21.1	18.5	18.0	17.3	17.3	21.8	18.8
2001	16.3	*	5.4	5.5	22.8	20.5	19.2	19.1	18.4	18.8	21.4	17.8
2000	16.4	6.9	5.9	4.8	22.4	22.2	19.1	19.7	17.5	19.9	19.9	21.9
1999	16.7	7.7	7.1	5.6	22.7	20.6	19.5	19.2	18.5	19.5	26.0	27.8
Non-Hispanic black, male												
2002	23.6	*	6.5	5.4	31.0	34.1	28.0	28.0	28.6	28.7	40.7	47.0
2001	24.5	*	5.8	7.1	33.4	32.6	29.7	30.7	30.2	27.4	36.9	35.8
2000	24.8	7.2	6.4	5.8	32.5	35.6	30.5	30.1	29.6	32.6	36.4	45.8
1999	24.9	*	8.0	7.0	32.9	33.0	29.8	30.7	29.3	29.5	43.5	68.9
Non-Hispanic black, female												
2002	8.5	*	3.6	3.9	12.1	9.3	10.1	9.4	8.2	9.1	11.1	*
2001	8.9	*	5.0	3.8	12.3	9.6	9.9	9.2	9.0	12.8	12.6	11.1
2000	8.8	*	5.4	3.7	12.5	10.1	8.9	10.8	7.8	11.0	10.6	13.1
1999	9.4	9.3	6.2	4.1	12.6	9.4	10.4	9.5	9.8	12.5	16.3	12.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
All origins 3/, both sexes												
2002	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
1999	0.2	*	*	0.1	0.2	0.2	0.3	0.2	0.1	*	*	*
All origins 3/, male												
2002	0.3	*	*	*	0.4	0.4	0.5	0.4	0.3	*	*	*
2001	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	*	*	*
2000	0.3	*	*	*	0.4	0.3	0.5	0.4	0.2	*	*	*
1999	0.3	*	*	0.1	0.4	0.4	0.5	0.3	0.2	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	0.3	0.4	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	0.3	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	0.6	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	0.6	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001	0.2	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
1999	0.2	*	*	0.1	0.2	0.2	0.3	0.2	0.1	*	*	*
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.3	0.4	0.5	0.4	0.3	*	*	*
2001	0.3	*	*	*	0.3	0.4	0.4	0.4	0.2	*	*	*
2000	0.3	*	*	*	0.4	0.3	0.5	0.3	0.2	*	*	*
1999	0.3	*	*	*	0.4	0.4	0.5	0.3	0.2	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Railway accidents (V05,V15,V80.6,V81.2-V81.9)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2000	0.2	*	*	*	0.3	0.2	0.3	0.2	0.1	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
Non-Hispanic white, male												
2002	0.3	*	*	*	0.4	0.4	0.4	0.3	0.2	*	*	*
2001	0.2	*	*	*	0.3	0.4	0.4	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.4	0.3	0.5	0.3	*	*	*	*
1999	0.3	*	*	*	0.4	0.3	0.5	0.3	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	*	*	0.4	0.6	*	*	*	*
2001	0.2	*	*	*	*	*	0.4	*	*	*	*	*
2000	0.3	*	*	*	*	*	0.5	*	*	*	*	*
1999	0.2	*	*	*	*	*	0.4	*	*	*	*	*
Non-Hispanic black, male												
2002	0.5	*	*	*	*	*	*	1.2	*	*	*	*
2001	0.4	*	*	*	*	*	0.8	*	*	*	*	*
2000	0.5	*	*	*	*	*	0.9	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All origins 3/, both sexes												
2002	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4
All origins 3/, male												
2002	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8
All origins 3/, female												
2002	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5
Hispanic, both sexes												
2002	14.6	3.1	5.6	3.6	26.6	17.6	15.2	13.9	14.3	17.9	24.4	24.9
2001	14.3	4.9	4.7	3.5	25.1	17.5	15.3	13.8	15.1	17.9	26.5	20.0
2000	13.9	5.2	4.9	3.8	23.7	16.4	14.7	15.1	14.4	18.4	22.3	22.6
1999	13.3	5.3	5.2	3.6	21.9	16.4	13.9	13.7	14.5	19.7	23.5	23.2
Hispanic, male												
2002	21.3	*	6.8	4.4	38.9	26.4	22.6	19.6	20.5	27.5	34.5	43.5
2001	21.0	*	5.3	4.0	36.9	26.9	23.1	20.2	21.2	25.0	44.7	*
2000	20.1	6.1	5.2	4.1	34.7	24.9	21.6	22.0	21.2	26.0	33.4	*
1999	19.1	5.7	5.6	4.1	31.6	24.4	20.8	20.1	19.0	27.2	37.0	*
Hispanic, female												
2002	7.4	*	4.4	2.7	12.4	7.2	7.3	8.2	8.7	10.2	17.4	*
2001	7.2	6.2	4.1	3.0	11.2	6.5	6.9	7.6	9.7	12.3	14.2	*
2000	7.2	*	4.6	3.6	10.6	6.5	7.3	8.3	8.4	12.4	14.9	*
1999	7.2	*	4.7	3.0	10.6	7.1	6.5	7.4	10.6	13.8	14.6	*
Non-Hispanic 4/, both sexes												
2002	15.9	3.0	3.5	4.0	28.5	17.7	15.8	14.8	14.1	16.9	25.7	28.1
2001	15.5	3.2	3.9	4.2	27.1	17.5	15.7	14.2	13.5	16.2	26.3	28.0
2000	15.6	4.2	4.1	4.4	27.5	17.4	15.4	14.1	14.4	16.3	25.8	30.6
1999	15.4	4.7	4.0	4.5	26.9	16.9	15.0	13.5	14.0	17.6	26.5	30.5
Non-Hispanic 4/, male												
2002	21.9	3.2	4.0	4.6	39.2	26.4	22.2	21.3	19.7	22.3	35.1	51.9
2001	21.5	3.1	4.3	5.1	38.1	25.9	22.2	20.4	18.8	21.3	36.9	51.6
2000	21.4	4.2	4.5	5.1	37.9	25.5	22.0	20.2	19.6	21.3	35.7	58.3
1999	21.0	4.8	4.4	5.2	36.7	24.6	21.1	19.2	19.1	22.9	37.7	60.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic 4/, female												
2002	10.1	2.9	3.0	3.4	17.4	9.1	9.5	8.6	8.9	12.4	19.4	17.8
2001	9.8	3.4	3.5	3.3	15.8	9.1	9.4	8.2	8.7	12.0	19.3	18.2
2000	10.0	4.2	3.6	3.6	16.9	9.2	8.9	8.2	9.6	12.2	19.3	19.4
1999	10.0	4.7	3.6	3.7	16.8	9.2	9.1	8.1	9.3	13.2	19.2	18.5
Non-Hispanic white, both sexes												
2002	16.3	2.4	3.0	4.0	30.8	17.8	15.8	14.8	14.0	16.9	26.2	28.9
2001	15.8	3.0	3.5	4.1	28.7	17.6	15.7	13.9	13.3	16.1	26.9	29.0
2000	15.8	3.3	3.7	4.4	29.5	17.0	15.3	13.7	14.2	16.1	26.2	31.3
1999	15.6	4.1	3.4	4.4	28.7	16.8	14.8	13.1	13.7	17.5	26.8	30.9
Non-Hispanic white, male												
2002	22.3	2.7	3.3	4.6	42.1	26.1	22.0	21.2	19.4	21.8	35.2	52.6
2001	21.7	2.9	3.9	4.8	40.1	25.9	21.9	19.9	18.2	21.0	37.3	53.1
2000	21.5	3.4	4.1	5.1	40.3	24.7	21.6	19.6	18.9	20.7	35.6	60.0
1999	21.1	4.7	3.6	5.1	38.8	24.2	20.6	18.5	18.5	22.6	37.6	61.2
Non-Hispanic white, female												
2002	10.5	2.1	2.8	3.3	19.0	9.4	9.7	8.4	8.8	12.7	20.2	18.6
2001	10.1	3.1	3.1	3.3	17.0	9.3	9.6	8.1	8.7	11.9	20.0	19.0
2000	10.3	3.2	3.2	3.6	18.4	9.3	9.0	7.9	9.8	12.3	20.0	19.8
1999	10.3	3.5	3.1	3.7	18.3	9.4	9.0	7.9	9.2	13.2	19.7	18.8
Non-Hispanic black, both sexes												
2002	14.8	5.4	4.9	4.3	21.0	20.3	17.2	16.3	15.7	16.0	20.2	18.5
2001	15.2	*	5.3	5.0	22.2	19.6	17.6	17.2	16.6	17.4	19.9	16.9
2000	15.5	6.9	5.6	4.5	21.6	21.1	17.6	18.3	16.3	18.4	18.8	20.7
1999	15.5	7.6	6.5	5.1	21.7	19.2	17.9	17.4	16.7	18.1	24.0	25.5
Non-Hispanic black, male												
2002	22.0	*	6.3	5.0	30.1	32.7	25.7	24.8	25.4	25.6	36.6	45.9
2001	22.7	*	5.6	6.5	32.4	31.0	26.9	27.5	26.8	24.2	34.8	35.8
2000	23.1	7.2	6.0	5.4	31.1	33.6	27.9	27.6	27.2	29.8	34.6	44.6
1999	22.8	*	7.0	6.2	31.5	30.7	26.9	27.5	25.8	26.8	41.3	61.8
Non-Hispanic black, female												
2002	8.3	*	3.5	3.6	11.9	9.1	9.7	9.0	8.0	9.1	10.9	*
2001	8.5	*	4.9	3.4	12.1	9.2	9.4	8.4	8.4	12.6	11.5	9.8
2000	8.5	*	5.2	3.6	12.3	9.8	8.5	10.2	7.5	10.4	9.9	11.8
1999	8.9	9.3	5.9	4.0	12.1	8.9	9.9	8.8	9.3	12.0	14.4	12.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)												
All origins 3/, both sexes												
2002	1.3	*	1.0	0.6	1.1	1.0	1.4	1.4	1.4	2.0	2.6	3.3
2001	1.2	*	1.1	0.7	1.0	1.0	1.4	1.4	1.4	1.6	3.0	3.0
2000	1.2	*	1.2	0.6	0.9	1.0	1.3	1.3	1.5	1.6	2.8	3.6
1999	1.2	*	1.1	0.7	0.9	1.0	1.3	1.3	1.5	1.7	2.5	3.7
All origins 3/, male												
2002	1.7	*	1.3	0.7	1.6	1.6	1.9	2.1	2.0	2.7	3.4	5.8
2001	1.8	*	1.4	0.9	1.5	1.5	2.0	2.0	2.0	2.1	4.1	5.9
2000	1.7	*	1.4	0.8	1.3	1.5	1.9	2.0	2.2	2.2	3.8	7.0
1999	1.7	*	1.2	0.8	1.4	1.5	1.9	2.0	2.2	2.3	3.7	6.5
All origins 3/, female												
2002	0.8	*	0.6	0.5	0.5	0.5	0.8	0.7	0.9	1.4	2.0	2.2
2001	0.8	*	0.8	0.4	0.5	0.4	0.8	0.8	0.7	1.1	2.2	1.8
2000	0.8	*	0.9	0.5	0.5	0.4	0.7	0.7	0.8	1.1	2.1	2.3
1999	0.7	*	1.0	0.5	0.5	0.5	0.6	0.6	0.9	1.2	1.7	2.6
Hispanic, both sexes												
2002	1.7	*	1.8	0.7	1.3	1.5	1.9	2.1	2.8	5.0	5.5	*
2001	1.9	*	1.8	0.6	1.3	1.8	2.5	2.4	3.1	4.1	7.9	*
2000	1.6	*	1.5	0.6	1.1	1.4	1.9	2.1	2.7	4.3	6.7	*
1999	1.5	*	1.9	0.5	1.2	1.5	1.5	2.2	3.0	3.7	5.6	*
Hispanic, male												
2002	2.6	*	2.4	0.9	2.1	2.2	3.0	3.7	4.4	8.6	9.8	*
2001	2.8	*	2.2	0.7	2.0	3.0	4.0	3.9	4.9	6.2	10.7	*
2000	2.3	*	1.5	0.6	1.7	2.3	2.8	3.3	4.4	6.3	10.3	*
1999	2.2	*	2.1	0.6	2.0	2.4	2.1	3.6	4.3	5.7	*	*
Hispanic, female												
2002	0.8	*	*	*	*	0.6	0.7	*	*	*	*	*
2001	0.9	*	1.4	*	*	*	1.0	*	*	*	6.0	*
2000	0.9	*	1.4	*	*	*	0.9	*	*	*	*	*
1999	0.8	*	1.6	*	*	*	0.8	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	0.8	0.5	1.0	0.9	1.3	1.3	1.3	1.8	2.4	3.1
2001	1.1	*	0.9	0.7	0.9	0.8	1.2	1.3	1.2	1.4	2.7	3.0
2000	1.2	*	1.1	0.6	0.8	0.9	1.2	1.3	1.4	1.4	2.6	3.4
1999	1.2	*	0.9	0.7	0.9	0.9	1.2	1.2	1.4	1.5	2.3	3.6
Non-Hispanic 4/, male												
2002	1.6	*	1.1	0.6	1.4	1.4	1.7	1.9	1.8	2.3	3.0	5.5
2001	1.6	*	1.2	0.9	1.4	1.2	1.7	1.8	1.8	1.8	3.8	5.7
2000	1.6	*	1.4	0.8	1.2	1.3	1.7	1.9	2.0	1.9	3.5	6.5
1999	1.6	*	1.0	0.8	1.3	1.3	1.9	1.8	2.0	2.0	3.5	6.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)												
Non-Hispanic 4/, female												
2002	0.8	*	0.5	0.5	0.5	0.5	0.8	0.7	0.8	1.3	2.0	2.0
2001	0.7	*	0.6	0.4	0.5	0.4	0.7	0.8	0.7	1.0	2.0	1.8
2000	0.7	*	0.8	0.5	0.5	0.5	0.7	0.6	0.8	1.0	2.0	2.1
1999	0.7	*	0.8	0.6	0.5	0.5	0.6	0.6	0.8	1.1	1.6	2.5
Non-Hispanic white, both sexes												
2002	1.1	*	0.6	0.5	0.9	0.8	1.0	1.1	1.2	1.6	2.2	2.9
2001	1.0	*	0.8	0.6	0.9	0.7	1.0	1.1	1.1	1.2	2.5	2.9
2000	1.0	*	0.9	0.6	0.9	0.8	1.0	1.1	1.2	1.3	2.5	3.1
1999	1.0	*	0.7	0.6	0.9	0.7	1.0	1.0	1.2	1.3	2.2	3.4
Non-Hispanic white, male												
2002	1.4	*	0.8	0.5	1.4	1.3	1.4	1.6	1.6	2.0	2.7	5.0
2001	1.4	*	1.1	0.8	1.3	1.1	1.4	1.5	1.5	1.6	3.5	5.3
2000	1.4	*	1.1	0.7	1.1	1.1	1.3	1.6	1.7	1.6	3.1	5.7
1999	1.4	*	0.7	0.7	1.2	1.1	1.4	1.5	1.7	1.8	3.1	5.8
Non-Hispanic white, female												
2002	0.7	*	0.4	0.4	0.5	0.4	0.7	0.6	0.7	1.2	1.9	1.9
2001	0.7	*	0.4	0.3	0.5	0.4	0.6	0.7	0.7	0.9	1.8	1.9
2000	0.7	*	0.6	0.4	0.6	0.4	0.6	0.5	0.7	1.0	2.1	2.1
1999	0.7	*	0.8	0.5	0.5	0.4	0.5	0.5	0.7	1.0	1.5	2.5
Non-Hispanic black, both sexes												
2002	1.6	*	1.2	0.8	1.2	1.4	2.2	2.4	2.0	2.4	3.0	*
2001	1.7	*	1.6	1.1	1.1	1.1	2.5	2.6	2.3	2.0	2.7	*
2000	1.7	*	1.9	1.0	0.8	1.3	2.5	2.5	2.5	2.2	2.6	*
1999	1.9	*	1.7	1.2	1.0	1.6	2.8	2.3	2.7	2.3	2.7	*
Non-Hispanic black, male												
2002	2.4	*	1.6	0.9	1.7	2.2	3.2	4.0	3.4	3.8	6.2	*
2001	2.5	*	1.8	1.7	1.5	1.6	3.7	3.8	4.3	3.1	*	*
2000	2.6	*	2.3	1.3	1.4	2.0	3.8	3.9	4.1	4.3	*	*
1999	2.9	*	2.1	1.4	1.4	2.6	4.6	4.2	4.7	3.9	*	*
Non-Hispanic black, female												
2002	0.9	*	*	0.7	0.7	0.7	1.3	1.1	*	*	*	*
2001	1.0	*	*	*	*	*	1.5	1.5	*	*	*	*
2000	0.9	*	*	0.7	*	*	1.4	1.2	*	*	*	*
1999	0.9	*	*	1.0	*	*	1.2	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)												
All origins 3/, both sexes												
2002	0.2	*	*	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.2	*
2001	0.2	*	*	0.3	0.2	0.2	0.3	0.3	0.2	0.2	*	*
2000	0.2	*	*	0.3	0.2	0.1	0.2	0.3	0.2	0.2	0.2	*
1999	0.2	*	*	0.4	0.3	0.2	0.3	0.2	0.1	0.2	*	*
All origins 3/, male												
2002	0.4	*	*	0.4	0.3	0.3	0.4	0.5	0.3	0.3	*	*
2001	0.4	*	*	0.4	0.4	0.3	0.5	0.5	0.4	0.4	*	*
2000	0.4	*	*	0.6	0.3	0.2	0.4	0.5	0.4	0.4	0.4	*
1999	0.4	*	*	0.5	0.5	0.4	0.5	0.4	0.2	0.4	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	0.1	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	0.2	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.3	*	*	*	*	0.3	0.4	*	*	*	*	*
2001	0.3	*	*	*	*	0.4	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	0.3	0.4	*	*	*	*	*
Hispanic, male												
2002	0.5	*	*	*	*	0.6	0.7	*	*	*	*	*
2001	0.5	*	*	*	*	0.8	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	0.6	0.9	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	0.3	0.2	0.1	0.2	0.2	0.2	0.1	*	*
2001	0.2	*	*	0.3	0.2	0.1	0.3	0.3	0.2	0.2	*	*
2000	0.2	*	*	0.4	0.2	0.1	0.2	0.2	0.2	0.2	*	*
1999	0.2	*	*	0.4	0.3	0.2	0.3	0.2	0.1	0.2	*	*
Non-Hispanic 4/, male												
2002	0.3	*	*	0.4	0.3	0.2	0.4	0.4	0.3	0.3	*	*
2001	0.4	*	*	0.5	0.4	0.2	0.5	0.5	0.4	0.4	*	*
2000	0.4	*	*	0.6	0.3	0.2	0.4	0.4	0.4	0.3	*	*
1999	0.4	*	*	0.6	0.5	0.3	0.5	0.4	0.2	0.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	0.1	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	0.2	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.2	*	*	*
2001	0.2	*	*	0.3	0.2	0.1	0.2	0.2	0.2	0.1	*	*
2000	0.2	*	*	0.3	0.2	0.1	0.2	0.2	0.2	0.2	*	*
1999	0.2	*	*	0.4	0.3	0.2	0.2	0.2	0.1	0.1	*	*
Non-Hispanic white, male												
2002	0.3	*	*	0.4	0.3	0.3	0.4	0.4	0.3	*	*	*
2001	0.4	*	*	0.4	0.4	0.2	0.5	0.5	0.4	0.3	*	*
2000	0.3	*	*	0.6	0.3	*	0.3	0.4	0.4	0.3	*	*
1999	0.4	*	*	0.6	0.5	0.3	0.4	0.4	*	0.3	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	0.2	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	0.4	*	*	0.4	*	*	*	*	*
2001	0.3	*	*	0.4	*	*	0.4	*	*	*	*	*
2000	0.3	*	*	0.5	*	*	0.5	*	*	*	*	*
1999	0.3	*	*	0.5	*	*	0.4	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	*	*	0.6	*	*	0.8	*	*	*	*	*
2001	0.6	*	*	0.8	*	*	0.9	*	*	*	*	*
2000	0.6	*	*	0.9	*	*	1.2	*	*	*	*	*
1999	0.6	*	*	0.8	*	*	0.9	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)												
All origins 3/, both sexes												
2002	1.1	*	*	0.1	1.5	1.9	1.7	1.6	1.0	0.4	0.2	*
2001	1.1	*	*	0.1	1.6	2.0	1.7	1.4	0.7	0.3	0.2	*
2000	1.0	*	*	0.1	1.4	1.8	1.6	1.3	0.8	0.4	*	*
1999	0.8	*	*	0.1	1.3	1.5	1.3	1.0	0.7	0.3	*	*
All origins 3/, male												
2002	2.1	*	*	0.1	2.8	3.5	3.1	3.0	1.9	0.8	0.4	*
2001	2.0	*	*	0.2	2.8	3.7	2.9	2.5	1.4	0.7	0.4	*
2000	1.8	*	*	0.1	2.6	3.4	2.8	2.3	1.4	0.8	*	*
1999	1.6	*	*	0.1	2.3	2.9	2.4	1.8	1.4	0.6	*	*
All origins 3/, female												
2002	0.2	*	*	*	0.2	0.3	0.4	0.4	0.2	*	*	*
2001	0.2	*	*	*	0.2	0.3	0.4	0.3	*	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.4	0.3	0.2	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.3	0.3	*	*	*	*
Hispanic, both sexes												
2002	0.6	*	*	*	0.9	1.0	0.8	0.5	*	*	*	*
2001	0.6	*	*	*	1.0	1.3	0.6	*	*	*	*	*
2000	0.5	*	*	*	0.8	0.9	0.9	*	*	*	*	*
1999	0.4	*	*	*	0.6	0.8	0.6	*	*	*	*	*
Hispanic, male												
2002	1.0	*	*	*	1.6	1.8	1.3	*	*	*	*	*
2001	1.1	*	*	*	1.7	2.3	1.2	*	*	*	*	*
2000	0.9	*	*	*	1.4	1.6	1.5	*	*	*	*	*
1999	0.7	*	*	*	1.1	1.4	1.1	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	0.1	1.6	2.1	1.9	1.8	1.1	0.4	0.2	*
2001	1.1	*	*	0.1	1.7	2.2	1.8	1.5	0.8	0.4	0.2	*
2000	1.1	*	*	0.1	1.6	2.0	1.7	1.3	0.8	0.4	*	*
1999	0.9	*	*	0.1	1.4	1.7	1.4	1.1	0.7	0.3	*	*
Non-Hispanic 4/, male												
2002	2.2	*	*	0.1	3.0	3.9	3.3	3.2	2.0	0.9	0.4	*
2001	2.1	*	*	0.2	3.1	4.0	3.2	2.7	1.5	0.7	*	*
2000	2.0	*	*	0.1	2.9	3.7	3.0	2.4	1.5	0.8	*	*
1999	1.7	*	*	0.1	2.5	3.1	2.5	1.9	1.4	0.6	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)												
Non-Hispanic 4/, female												
2002	0.2	*	*	*	0.2	0.3	0.4	0.4	0.2	*	*	*
2001	0.2	*	*	*	0.2	0.4	0.5	0.3	*	*	*	*
2000	0.2	*	*	*	0.2	0.3	0.4	0.3	0.2	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.3	0.3	*	*	*	*
Non-Hispanic white, both sexes												
2002	1.3	*	*	0.1	1.8	2.1	2.1	2.0	1.2	0.4	0.2	*
2001	1.2	*	*	0.1	1.8	2.2	2.1	1.7	0.9	0.4	*	*
2000	1.1	*	*	0.1	1.7	1.9	1.9	1.5	0.9	0.4	*	*
1999	1.0	*	*	0.1	1.4	1.6	1.6	1.3	0.8	0.3	*	*
Non-Hispanic white, male												
2002	2.4	*	*	*	3.3	3.8	3.7	3.7	2.2	0.9	*	*
2001	2.3	*	*	0.2	3.3	3.9	3.5	3.0	1.6	0.8	*	*
2000	2.1	*	*	0.2	3.1	3.5	3.2	2.7	1.6	0.9	*	*
1999	1.7	*	*	0.2	2.7	2.9	2.7	2.2	1.5	0.6	*	*
Non-Hispanic white, female												
2002	0.3	*	*	*	0.2	0.4	0.5	0.4	0.2	*	*	*
2001	0.3	*	*	*	0.2	0.4	0.6	0.4	*	*	*	*
2000	0.2	*	*	*	0.2	0.3	0.5	0.4	0.2	*	*	*
1999	0.2	*	*	*	0.2	0.3	0.4	0.3	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.9	*	*	*	1.2	2.7	1.3	0.5	*	*	*	*
2001	0.9	*	*	*	1.3	2.9	1.0	0.5	*	*	*	*
2000	0.9	*	*	*	1.4	3.1	1.1	*	*	*	*	*
1999	0.8	*	*	*	1.4	2.4	0.9	*	*	*	*	*
Non-Hispanic black, male												
2002	1.8	*	*	*	2.3	5.5	2.7	1.0	*	*	*	*
2001	1.8	*	*	*	2.4	5.9	2.2	1.1	*	*	*	*
2000	1.9	*	*	*	2.5	6.4	2.3	*	*	*	*	*
1999	1.6	*	*	*	2.6	5.2	1.9	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	0.1	0.2	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*
All origins 3/, male												
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.1	0.2	*	*	*	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.2	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	0.1	0.2	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	0.2	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.2	*	*	*	*	*
2000	0.2	*	*	*	0.3	0.2	0.1	0.2	*	*	*	*
1999	0.1	*	*	*	0.2	*	0.1	0.2	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accident involving collision with railway train (V25,V35,V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.2	*	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*
Non-Hispanic white, male												
2002	0.1	*	*	*	0.2	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*	*
2000	0.2	*	*	*	0.3	0.2	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	0.2	*	*	0.2	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motorcyclist involved in collision with railway train (V25,V35)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motorcyclist involved in collision with railway train (V25,V35)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	0.1	0.2	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*
All origins 3/, male												
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	0.1	0.2	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	0.2	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.1	*	*	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.1	0.2	*	*	*	*
1999	0.1	*	*	*	0.2	*	0.1	0.2	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other motor vehicle accident involving collision with railway train (V45,V55,V65,V75,V81.0-V81.1,V87.6,V88.6)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.2	*	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	0.2	*
Non-Hispanic white, male												
2002	0.1	*	*	*	0.2	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*	*
2000	0.2	*	*	*	0.3	0.2	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	0.2	*	*	0.2	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)												
All origins 3/, both sexes												
2002	1.8	*	0.2	0.3	4.5	2.3	1.7	1.4	1.3	1.2	1.6	1.5
2001	1.5	*	0.1	0.2	3.8	2.0	1.4	1.2	0.9	1.1	1.6	1.7
2000	1.4	*	0.2	0.2	3.5	1.7	1.4	1.0	1.0	1.2	1.4	1.7
1999	1.4	*	0.1	0.2	3.3	1.8	1.3	1.0	1.0	1.2	1.5	1.2
All origins 3/, male												
2002	2.6	*	*	0.3	6.7	3.6	2.5	1.9	2.0	1.7	2.2	3.2
2001	2.2	*	*	0.2	5.6	3.0	2.2	1.8	1.5	1.5	2.5	3.1
2000	2.0	*	*	0.2	5.2	2.7	2.0	1.5	1.4	1.6	2.1	3.0
1999	2.0	*	*	0.3	4.9	2.8	1.9	1.6	1.4	1.8	2.3	2.5
All origins 3/, female												
2002	0.9	*	*	0.3	2.3	0.9	0.9	0.8	0.6	0.8	1.2	0.8
2001	0.8	*	*	0.2	1.9	0.9	0.7	0.6	0.4	0.7	1.1	1.2
2000	0.8	*	*	0.2	1.6	0.8	0.8	0.6	0.6	0.8	0.9	1.2
1999	0.7	*	*	0.2	1.7	0.7	0.7	0.5	0.6	0.8	1.0	0.7
Hispanic, both sexes												
2002	1.8	*	*	*	4.4	2.6	1.6	1.0	*	*	*	*
2001	1.6	*	*	*	3.7	2.1	1.4	1.4	1.1	*	*	*
2000	1.3	*	*	*	3.2	1.6	1.1	1.0	*	*	*	*
1999	1.3	*	*	*	3.1	1.8	1.3	0.9	*	*	*	*
Hispanic, male												
2002	2.8	*	*	*	6.9	4.2	2.6	1.2	*	*	*	*
2001	2.4	*	*	*	5.7	3.2	2.2	2.1	*	*	*	*
2000	1.9	*	*	*	4.8	2.6	1.5	1.4	*	*	*	*
1999	2.1	*	*	*	4.8	2.9	2.1	1.4	*	*	*	*
Hispanic, female												
2002	0.7	*	*	*	1.5	0.8	*	*	*	*	*	*
2001	0.7	*	*	*	1.4	0.8	*	*	*	*	*	*
2000	0.6	*	*	*	1.3	*	*	*	*	*	*	*
1999	0.5	*	*	*	1.1	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.7	*	0.2	0.3	4.5	2.2	1.7	1.4	1.3	1.2	1.6	1.6
2001	1.5	*	*	0.2	3.8	1.9	1.4	1.2	0.9	1.1	1.6	1.7
2000	1.4	*	0.2	0.2	3.5	1.7	1.5	1.0	1.0	1.2	1.4	1.8
1999	1.4	*	*	0.2	3.4	1.8	1.3	1.1	1.0	1.2	1.5	1.2
Non-Hispanic 4/, male												
2002	2.6	*	*	0.3	6.6	3.4	2.5	2.0	2.0	1.8	2.3	3.3
2001	2.2	*	*	0.2	5.6	3.0	2.1	1.8	1.5	1.5	2.5	3.0
2000	2.1	*	*	0.2	5.3	2.7	2.1	1.5	1.4	1.7	2.1	3.1
1999	2.0	*	*	0.3	4.8	2.8	1.9	1.6	1.5	1.8	2.4	2.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57,V60-V61,V66-V67,V70-V71,V76-V77)												
Non-Hispanic 4/, female												
2002	1.0	*	*	0.3	2.4	1.0	1.0	0.8	0.6	0.8	1.1	0.8
2001	0.8	*	*	0.2	2.0	0.9	0.7	0.6	0.4	0.7	1.0	1.2
2000	0.8	*	*	0.2	1.6	0.8	0.9	0.6	0.6	0.8	0.9	1.3
1999	0.7	*	*	0.2	1.9	0.8	0.7	0.5	0.7	0.8	0.9	0.7
Non-Hispanic white, both sexes												
2002	1.8	*	*	0.3	4.8	2.2	1.8	1.4	1.2	1.2	1.7	1.7
2001	1.5	*	*	0.2	3.8	1.9	1.4	1.1	0.9	1.1	1.6	1.9
2000	1.4	*	*	0.2	3.6	1.7	1.5	1.0	1.0	1.1	1.4	1.9
1999	1.3	*	*	0.2	3.5	1.7	1.3	1.0	1.0	1.2	1.5	1.2
Non-Hispanic white, male												
2002	2.6	*	*	0.3	6.9	3.4	2.5	1.9	1.9	1.6	2.3	3.3
2001	2.2	*	*	0.2	5.6	2.9	2.1	1.7	1.5	1.5	2.5	3.2
2000	2.0	*	*	0.2	5.4	2.5	2.1	1.4	1.3	1.5	2.0	3.3
1999	2.0	*	*	0.3	4.9	2.7	1.9	1.5	1.4	1.7	2.4	2.5
Non-Hispanic white, female												
2002	1.0	*	*	0.3	2.7	1.0	1.0	0.8	0.6	0.9	1.2	0.9
2001	0.8	*	*	0.2	2.0	0.9	0.7	0.6	0.4	0.8	1.1	1.3
2000	0.8	*	*	0.2	1.8	0.9	0.9	0.5	0.6	0.8	0.9	1.3
1999	0.7	*	*	0.2	1.9	0.8	0.7	0.5	0.6	0.8	0.9	*
Non-Hispanic black, both sexes												
2002	1.9	*	*	0.5	3.9	2.7	1.8	1.8	2.3	1.8	*	*
2001	1.8	*	*	*	4.2	2.5	1.9	1.7	1.1	*	2.2	*
2000	1.6	*	*	*	3.2	2.1	1.7	1.7	1.5	2.0	*	*
1999	1.7	*	*	*	3.5	2.3	1.7	1.5	1.6	1.4	2.2	*
Non-Hispanic black, male												
2002	3.1	*	*	*	6.0	4.7	2.7	3.0	4.1	3.7	*	*
2001	2.8	*	*	*	6.4	4.1	3.0	2.7	2.0	*	*	*
2000	2.5	*	*	*	4.9	3.8	2.7	2.5	2.6	3.3	*	*
1999	2.6	*	*	*	5.2	4.0	2.6	2.5	2.6	*	*	*
Non-Hispanic black, female												
2002	0.9	*	*	*	1.7	0.9	1.1	0.9	*	*	*	*
2001	0.8	*	*	*	2.0	1.1	0.9	*	*	*	*	*
2000	0.8	*	*	*	1.5	*	0.8	1.0	*	*	*	*
1999	0.8	*	*	*	1.7	0.8	0.9	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
All origins 3/, both sexes												
2002	3.9	1.1	1.0	0.9	6.8	3.8	3.4	3.3	3.8	5.2	9.2	9.9
2001	3.7	1.6	1.0	0.9	6.1	3.7	3.3	3.1	3.5	5.1	8.4	10.1
2000	3.7	1.5	0.9	0.9	6.1	3.7	3.1	3.2	3.8	5.0	9.0	10.1
1999	3.7	1.6	0.9	0.9	5.8	3.7	3.1	3.2	3.8	5.3	8.8	9.2
All origins 3/, male												
2002	4.8	1.5	1.0	0.9	8.7	5.1	4.2	4.1	4.7	6.0	12.2	16.9
2001	4.5	1.6	0.9	0.9	7.9	5.0	4.2	3.8	4.1	6.1	10.9	18.1
2000	4.5	1.5	0.9	0.9	7.5	4.8	3.9	4.0	4.5	5.8	11.6	18.2
1999	4.4	1.7	1.0	0.9	7.3	4.9	3.8	3.9	4.3	6.2	11.6	17.8
All origins 3/, female												
2002	3.0	*	0.9	0.8	4.9	2.4	2.6	2.5	3.0	4.4	7.2	6.8
2001	2.9	1.6	1.1	0.9	4.3	2.3	2.5	2.4	3.0	4.3	6.7	6.8
2000	2.9	1.5	0.9	0.8	4.5	2.6	2.3	2.4	3.2	4.3	7.3	6.8
1999	2.9	1.4	0.7	1.0	4.3	2.5	2.4	2.5	3.3	4.6	7.0	5.7
Hispanic, both sexes												
2002	3.9	*	1.3	0.9	7.6	4.4	4.0	3.7	4.1	4.8	6.7	*
2001	3.6	*	1.0	0.9	6.7	4.1	3.6	3.4	3.5	5.1	6.3	*
2000	3.1	*	1.2	0.9	4.8	3.6	3.0	3.7	3.6	5.1	7.7	*
1999	2.9	*	0.9	0.9	4.8	3.8	2.4	3.0	3.8	5.2	6.5	*
Hispanic, male												
2002	5.4	*	1.6	1.2	10.5	6.4	5.4	4.7	4.8	7.1	11.9	*
2001	4.9	*	*	1.0	9.5	5.9	5.1	4.2	4.3	6.6	13.0	*
2000	4.2	*	*	0.9	6.8	4.8	4.3	5.2	5.0	6.5	12.8	*
1999	3.9	*	*	0.8	6.2	5.6	3.2	4.2	4.7	5.7	*	*
Hispanic, female												
2002	2.3	*	*	*	4.1	2.1	2.4	2.7	3.4	3.0	*	*
2001	2.2	*	*	0.8	3.5	2.1	2.1	2.7	2.8	3.9	*	*
2000	1.9	*	*	0.8	2.5	2.2	1.6	2.2	2.4	4.0	*	*
1999	1.9	*	*	0.9	3.1	1.8	1.5	1.8	3.0	4.8	*	*
Non-Hispanic 4/, both sexes												
2002	3.9	1.1	0.9	0.9	6.7	3.6	3.3	3.2	3.8	5.2	9.3	10.1
2001	3.7	1.4	1.0	0.9	6.0	3.6	3.2	3.1	3.6	5.1	8.4	10.2
2000	3.8	1.4	0.8	0.9	6.3	3.7	3.1	3.1	3.8	5.0	9.0	10.3
1999	3.8	1.7	0.8	1.0	6.0	3.7	3.2	3.2	3.8	5.3	8.9	9.3
Non-Hispanic 4/, male												
2002	4.7	1.3	0.9	0.8	8.3	4.8	4.0	4.0	4.6	5.9	12.2	17.2
2001	4.5	1.5	0.9	0.9	7.5	4.7	4.0	3.8	4.1	6.1	10.8	18.4
2000	4.5	1.5	0.9	0.9	7.7	4.7	3.9	3.9	4.4	5.8	11.5	18.8
1999	4.5	1.7	1.0	0.9	7.4	4.7	3.9	3.9	4.3	6.2	11.7	18.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)												
Non-Hispanic 4/, female												
2002	3.1	*	0.9	0.9	5.0	2.5	2.6	2.5	3.0	4.5	7.4	7.0
2001	3.0	1.3	1.1	1.0	4.4	2.4	2.5	2.4	3.0	4.3	6.8	6.7
2000	3.1	*	0.8	0.8	4.9	2.6	2.4	2.4	3.2	4.4	7.4	6.9
1999	3.0	1.6	0.7	1.0	4.5	2.7	2.5	2.5	3.3	4.5	7.1	5.8
Non-Hispanic white, both sexes												
2002	4.1	0.9	0.8	0.9	7.2	3.5	3.3	3.2	3.8	5.3	9.7	10.6
2001	3.8	1.3	1.0	0.9	6.3	3.6	3.2	3.0	3.6	5.1	8.8	10.7
2000	3.9	1.3	0.8	0.9	6.8	3.7	3.1	3.0	3.8	5.1	9.3	10.9
1999	3.9	1.6	0.8	1.0	6.5	3.7	3.1	3.2	3.7	5.3	9.2	9.5
Non-Hispanic white, male												
2002	4.9	*	0.7	0.9	8.8	4.6	4.0	3.9	4.6	5.9	12.5	18.2
2001	4.6	*	0.8	0.9	7.8	4.7	3.9	3.5	4.1	6.1	11.1	19.2
2000	4.6	*	0.8	0.9	8.1	4.6	3.8	3.7	4.2	5.9	11.7	20.0
1999	4.6	1.9	0.9	0.9	7.9	4.7	3.6	3.8	4.1	6.1	11.9	18.6
Non-Hispanic white, female												
2002	3.3	*	0.8	0.9	5.6	2.4	2.6	2.4	3.0	4.8	7.9	7.3
2001	3.1	*	1.1	1.0	4.8	2.5	2.6	2.4	3.0	4.3	7.3	7.2
2000	3.2	*	0.8	0.9	5.4	2.6	2.4	2.3	3.4	4.5	7.8	7.2
1999	3.2	*	0.6	1.1	5.0	2.7	2.6	2.6	3.4	4.7	7.4	5.9
Non-Hispanic black, both sexes												
2002	3.4	*	1.3	0.8	4.9	4.8	3.6	3.6	3.8	4.0	5.8	*
2001	3.4	*	1.2	0.9	4.8	4.0	3.8	4.0	3.9	6.0	5.2	*
2000	3.4	*	1.1	0.7	4.8	4.3	3.5	4.3	4.1	4.5	5.7	*
1999	3.5	*	1.1	1.0	4.4	4.3	4.1	3.7	4.1	5.1	6.3	7.8
Non-Hispanic black, male												
2002	4.5	*	1.7	*	6.4	6.8	4.8	4.9	5.3	5.4	10.0	*
2001	4.6	*	*	0.9	6.4	5.7	5.3	6.0	4.9	7.3	8.7	*
2000	4.5	*	*	0.9	6.4	6.0	4.8	5.9	6.8	5.7	8.6	*
1999	4.6	*	*	1.1	6.0	5.8	5.8	5.1	5.5	7.8	10.9	*
Non-Hispanic black, female												
2002	2.4	*	*	0.9	3.4	3.0	2.6	2.6	2.5	3.0	3.5	*
2001	2.4	*	*	0.9	3.2	2.5	2.5	2.3	3.1	5.1	*	*
2000	2.3	*	*	*	3.2	2.7	2.4	2.9	1.9	3.6	4.0	*
1999	2.4	*	*	0.9	2.9	2.9	2.6	2.5	2.9	3.2	3.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	0.2	0.2	0.3	0.2	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.2	0.2	*	*	*
2000	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.2	0.2	0.2	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.2	0.2	*	*	*
2000	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of heavy transport vehicle or bus involved in collision with other motor vehicle (V62-V64,V69,V72-V74,V79)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	0.1	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	0.1	*	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	0.1	0.1	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2001	0.1	*	*	*	*	*	*	0.2	*	*	*	*
2000	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
All origins 3/, both sexes												
2002	1.5	*	0.3	0.4	3.5	2.0	1.6	1.3	1.2	1.0	1.0	0.9
2001	1.4	*	0.2	0.4	3.0	1.8	1.5	1.3	1.1	1.0	1.0	0.7
2000	1.3	0.6	0.2	0.4	3.1	1.8	1.4	1.2	0.9	0.8	0.9	1.0
1999	1.3	*	0.3	0.3	3.1	1.7	1.5	1.1	1.1	0.9	0.8	1.0
All origins 3/, male												
2002	2.2	*	0.4	0.4	4.8	3.0	2.3	2.0	1.6	1.4	1.5	1.8
2001	2.0	*	*	0.4	4.2	2.7	2.1	1.9	1.6	1.4	1.5	*
2000	1.9	*	*	0.4	4.3	2.8	2.0	1.8	1.3	1.2	1.4	2.2
1999	1.9	*	0.3	0.3	4.2	2.6	2.1	1.6	1.6	1.3	1.2	2.0
All origins 3/, female												
2002	0.9	*	*	0.4	2.2	1.0	1.0	0.7	0.7	0.6	0.7	*
2001	0.8	*	*	0.4	1.8	0.9	0.9	0.6	0.6	0.6	0.8	*
2000	0.7	*	*	0.3	1.7	0.8	0.8	0.6	0.6	0.5	0.6	*
1999	0.8	*	0.3	0.3	1.9	0.9	0.8	0.6	0.6	0.7	0.5	*
Hispanic, both sexes												
2002	1.7	*	0.7	0.6	3.3	2.3	1.8	1.5	1.3	*	*	*
2001	1.6	*	*	0.5	3.2	1.8	1.9	1.7	1.1	*	*	*
2000	1.5	*	*	0.5	2.7	1.8	1.7	1.5	*	*	*	*
1999	1.3	*	*	0.3	2.6	1.6	1.4	1.2	*	*	*	*
Hispanic, male												
2002	2.4	*	*	0.6	4.5	3.2	2.5	2.1	2.5	*	*	*
2001	2.3	*	*	*	4.5	2.7	2.9	2.4	*	*	*	*
2000	2.2	*	*	*	4.2	2.8	2.6	1.9	*	*	*	*
1999	1.9	*	*	*	4.0	2.6	2.1	1.6	*	*	*	*
Hispanic, female												
2002	1.0	*	*	*	1.9	1.2	1.0	*	*	*	*	*
2001	0.8	*	*	*	1.6	0.7	0.8	*	*	*	*	*
2000	0.7	*	*	*	0.9	0.7	*	*	*	*	*	*
1999	0.6	*	*	*	1.1	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.5	*	0.2	0.3	3.5	2.0	1.6	1.3	1.1	1.0	1.0	0.9
2001	1.3	*	0.2	0.3	3.0	1.8	1.4	1.2	1.1	0.9	1.0	0.8
2000	1.3	*	0.2	0.3	3.1	1.8	1.4	1.1	0.9	0.8	0.9	1.0
1999	1.3	*	0.3	0.3	3.2	1.7	1.5	1.1	1.1	0.9	0.8	1.0
Non-Hispanic 4/, male												
2002	2.1	*	*	0.4	4.9	2.9	2.2	1.9	1.6	1.4	1.5	1.8
2001	1.9	*	*	0.3	4.1	2.7	2.0	1.9	1.5	1.4	1.4	*
2000	1.9	*	*	0.4	4.3	2.8	2.0	1.8	1.3	1.2	1.3	2.2
1999	1.9	*	*	0.3	4.3	2.5	2.1	1.6	1.6	1.3	1.2	2.0

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)												
Non-Hispanic 4/, female												
2002	0.9	*	*	0.3	2.2	1.0	0.9	0.7	0.8	0.6	0.6	*
2001	0.8	*	*	0.3	1.9	0.9	0.9	0.6	0.6	0.5	0.7	*
2000	0.8	*	*	0.3	1.9	0.9	0.8	0.5	0.6	0.5	0.6	*
1999	0.8	*	0.4	0.3	2.0	0.9	0.9	0.6	0.6	0.6	0.5	*
Non-Hispanic white, both sexes												
2002	1.5	*	*	0.4	3.8	2.1	1.6	1.3	1.2	1.0	1.0	0.9
2001	1.4	*	*	0.3	3.3	1.8	1.5	1.2	1.0	1.0	1.1	0.7
2000	1.3	*	*	0.3	3.4	1.9	1.4	1.1	0.9	0.8	0.9	1.0
1999	1.3	*	0.2	0.3	3.4	1.8	1.4	1.0	1.0	0.9	0.8	1.0
Non-Hispanic white, male												
2002	2.2	*	*	0.4	5.3	3.0	2.2	1.9	1.6	1.5	1.6	1.7
2001	1.9	*	*	0.3	4.6	2.6	2.0	1.8	1.4	1.5	1.5	*
2000	1.9	*	*	0.3	4.7	2.9	2.0	1.7	1.2	1.2	1.3	2.2
1999	1.9	*	*	0.3	4.6	2.6	2.1	1.6	1.5	1.2	1.2	1.9
Non-Hispanic white, female												
2002	0.9	*	*	0.3	2.3	1.2	0.9	0.7	0.8	0.7	0.6	*
2001	0.8	*	*	0.3	2.0	0.9	0.9	0.6	0.6	0.5	0.8	*
2000	0.7	*	*	0.2	2.1	0.9	0.8	0.5	0.5	0.4	0.6	*
1999	0.8	*	*	0.3	2.2	1.0	0.8	0.5	0.6	0.7	0.5	*
Non-Hispanic black, both sexes												
2002	1.2	*	*	0.4	2.0	1.5	1.6	1.2	1.1	*	*	*
2001	1.1	*	*	0.4	1.7	1.8	1.1	1.2	1.8	*	*	*
2000	1.1	*	*	0.4	2.0	1.4	1.4	1.1	1.6	*	*	*
1999	1.2	*	*	*	2.0	1.5	1.6	1.3	1.0	*	*	*
Non-Hispanic black, male												
2002	1.8	*	*	*	2.7	2.7	2.3	2.3	1.8	*	*	*
2001	1.7	*	*	*	2.1	3.0	1.8	2.2	3.2	*	*	*
2000	1.7	*	*	*	2.7	2.4	2.1	1.8	2.5	*	*	*
1999	1.7	*	*	*	2.7	2.5	2.3	2.2	*	*	*	*
Non-Hispanic black, female												
2002	0.6	*	*	*	1.4	*	0.9	*	*	*	*	*
2001	0.6	*	*	*	1.3	0.8	*	*	*	*	*	*
2000	0.6	*	*	*	1.3	*	*	*	*	*	*	*
1999	0.7	*	*	*	1.4	0.7	1.0	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of special-use motor vehicle involved in any accident (V83-V86)												
All origins 3/, both sexes												
2002	0.3	*	*	0.3	0.5	0.4	0.3	0.3	0.3	0.4	0.5	*
2001	0.4	*	*	0.3	0.6	0.4	0.3	0.2	0.4	0.4	0.6	*
2000	0.4	*	*	0.3	0.6	0.4	0.3	0.3	0.5	0.4	0.4	*
1999	0.4	*	*	0.3	0.4	0.4	0.3	0.3	0.4	0.6	0.6	0.7
All origins 3/, male												
2002	0.6	*	*	0.5	0.9	0.6	0.6	0.5	0.5	0.8	1.0	*
2001	0.7	*	*	0.5	1.0	0.7	0.6	0.4	0.7	0.9	1.4	*
2000	0.7	*	*	0.4	1.0	0.7	0.6	0.5	0.9	0.9	1.1	*
1999	0.7	*	*	0.5	0.8	0.7	0.6	0.6	0.8	1.3	1.4	2.2
All origins 3/, female												
2002	0.1	*	*	0.2	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.2	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	0.2	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	0.3	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	*	*	0.4	0.6	0.4	0.4	0.3	0.3	0.4	0.5	*
2001	0.4	*	*	0.4	0.6	0.4	0.4	0.2	0.4	0.4	0.6	*
2000	0.4	*	*	0.4	0.7	0.4	0.3	0.3	0.5	0.5	0.5	*
1999	0.4	*	*	0.3	0.5	0.4	0.3	0.3	0.4	0.6	0.6	0.7
Non-Hispanic 4/, male												
2002	0.7	*	*	0.6	1.0	0.7	0.6	0.6	0.6	0.8	1.1	*
2001	0.7	*	*	0.5	1.1	0.8	0.7	0.5	0.7	0.9	1.5	*
2000	0.7	*	*	0.5	1.2	0.7	0.6	0.5	1.0	0.9	1.2	*
1999	0.7	*	*	0.5	0.9	0.8	0.6	0.6	0.8	1.3	1.4	2.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Occupant of special-use motor vehicle involved in any accident (V83-V86)												
Non-Hispanic 4/, female												
2002	0.1	*	*	0.2	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.2	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	0.2	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.4	*	*	0.5	0.7	0.5	0.4	0.4	0.3	0.4	0.5	*
2001	0.5	*	*	0.4	0.8	0.6	0.4	0.3	0.4	0.5	0.7	*
2000	0.5	*	*	0.5	0.8	0.5	0.4	0.3	0.5	0.5	0.5	*
1999	0.5	*	*	0.4	0.6	0.5	0.4	0.4	0.4	0.7	0.6	0.7
Non-Hispanic white, male												
2002	0.8	*	*	0.7	1.2	0.9	0.7	0.7	0.6	0.9	1.2	*
2001	0.8	*	*	0.6	1.3	1.0	0.8	0.5	0.8	1.0	1.6	*
2000	0.8	*	*	0.6	1.4	0.9	0.7	0.6	1.0	1.0	1.2	*
1999	0.9	*	*	0.6	1.1	0.9	0.7	0.7	0.9	1.4	1.6	2.4
Non-Hispanic white, female												
2002	0.1	*	*	0.2	0.2	*	*	*	*	*	*	*
2001	0.1	*	*	0.2	0.2	*	*	*	*	*	*	*
2000	0.1	*	*	0.3	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)												
All origins 3/, both sexes												
2002	5.5	1.1	1.3	1.1	9.9	6.0	5.2	5.0	4.8	6.6	10.3	11.4
2001	5.8	1.3	1.5	1.2	10.4	6.3	5.6	5.1	5.3	6.5	11.3	11.7
2000	6.0	1.8	1.6	1.4	11.1	6.5	5.8	5.5	5.7	6.7	10.7	13.1
1999	6.0	2.3	1.6	1.5	10.8	6.4	5.7	5.2	5.3	7.4	11.8	13.9
All origins 3/, male												
2002	7.3	1.1	1.4	1.3	13.3	8.5	7.0	6.8	6.5	8.7	13.9	21.7
2001	7.7	1.1	1.6	1.4	14.4	8.9	7.6	7.1	6.9	8.1	15.8	21.2
2000	8.0	1.9	1.8	1.5	14.8	9.2	8.0	7.4	7.5	8.5	14.7	24.7
1999	7.9	2.1	1.8	1.6	14.4	8.8	7.6	7.0	7.1	9.3	16.7	27.5
All origins 3/, female												
2002	3.7	1.1	1.3	1.0	6.3	3.4	3.4	3.3	3.3	4.8	8.0	6.9
2001	3.9	1.5	1.3	1.1	6.2	3.6	3.7	3.2	3.8	5.1	8.3	7.7
2000	4.1	1.7	1.4	1.4	7.1	3.8	3.5	3.6	4.0	5.3	8.1	8.4
1999	4.2	2.5	1.5	1.3	7.0	3.9	3.8	3.5	3.7	5.9	8.7	8.5
Hispanic, both sexes												
2002	4.4	*	1.5	0.9	8.4	5.0	4.4	4.3	4.3	5.5	8.0	*
2001	4.6	*	1.5	1.0	8.5	5.6	4.6	3.9	5.4	5.5	8.7	*
2000	5.4	*	1.8	1.3	10.4	6.4	5.5	5.5	5.9	6.4	5.5	*
1999	5.4	3.8	1.8	1.3	9.1	6.2	6.0	5.3	5.6	7.7	10.0	*
Hispanic, male												
2002	6.1	*	1.4	0.9	12.1	7.3	6.4	5.5	5.6	7.8	8.6	*
2001	6.6	*	1.6	1.0	12.6	8.3	6.9	5.7	6.4	7.6	14.8	*
2000	7.8	*	1.9	1.4	14.7	9.7	7.8	7.9	8.0	9.3	*	*
1999	7.4	*	1.7	1.5	12.7	8.5	8.9	7.3	6.4	10.8	17.7	*
Hispanic, female												
2002	2.5	*	1.7	1.0	4.1	2.3	2.1	3.0	3.2	3.6	7.6	*
2001	2.4	*	1.4	1.0	3.9	2.4	2.2	2.0	4.5	3.8	*	*
2000	3.0	*	1.6	1.2	5.2	2.7	3.0	3.1	4.1	4.1	*	*
1999	3.2	*	1.9	1.1	4.9	3.5	2.9	3.3	4.8	5.3	*	*
Non-Hispanic 4/, both sexes												
2002	5.6	1.2	1.3	1.2	10.2	6.2	5.3	5.1	4.8	6.6	10.4	11.5
2001	5.9	1.3	1.5	1.3	10.7	6.4	5.7	5.2	5.2	6.5	11.3	11.8
2000	6.1	1.8	1.6	1.4	11.2	6.5	5.8	5.4	5.6	6.7	10.9	13.3
1999	6.1	1.9	1.6	1.5	11.1	6.4	5.6	5.1	5.2	7.4	11.8	14.0
Non-Hispanic 4/, male												
2002	7.5	1.3	1.4	1.4	13.6	8.7	7.0	6.9	6.5	8.7	14.0	21.9
2001	7.9	1.2	1.6	1.4	14.7	9.0	7.6	7.2	6.9	8.1	15.9	21.5
2000	8.1	1.9	1.8	1.5	14.8	9.1	8.0	7.3	7.4	8.4	15.1	25.2
1999	7.9	1.7	1.8	1.6	14.7	8.8	7.4	6.9	7.0	9.2	16.5	27.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)												
Non-Hispanic 4/, female												
2002	3.9	*	1.2	1.0	6.8	3.7	3.6	3.4	3.3	4.9	8.0	7.0
2001	4.1	1.5	1.3	1.1	6.6	3.9	3.9	3.3	3.8	5.2	8.4	7.8
2000	4.3	1.8	1.4	1.4	7.4	4.0	3.6	3.6	4.0	5.3	8.2	8.5
1999	4.3	2.2	1.4	1.4	7.4	4.0	3.9	3.5	3.6	5.9	8.8	8.5
Non-Hispanic white, both sexes												
2002	5.8	0.9	1.2	1.1	11.1	6.2	5.3	4.9	4.7	6.7	10.6	11.9
2001	6.1	1.1	1.3	1.2	11.4	6.5	5.7	5.1	5.1	6.5	11.7	12.3
2000	6.2	1.3	1.5	1.4	12.0	6.3	5.8	5.2	5.6	6.5	11.2	13.7
1999	6.2	1.7	1.3	1.5	11.9	6.4	5.6	4.9	5.1	7.4	12.0	14.4
Non-Hispanic white, male												
2002	7.6	*	1.2	1.3	14.6	8.7	6.9	6.7	6.3	8.7	14.0	22.2
2001	8.0	*	1.4	1.3	15.5	9.1	7.4	7.0	6.5	8.1	16.1	22.6
2000	8.1	*	1.7	1.5	15.9	8.7	7.9	6.9	7.2	8.0	15.3	26.5
1999	8.0	*	1.4	1.6	15.6	8.8	7.4	6.5	6.8	9.2	16.4	28.7
Non-Hispanic white, female												
2002	4.1	*	1.1	1.0	7.4	3.7	3.6	3.3	3.3	4.9	8.3	7.4
2001	4.2	*	1.2	1.1	7.2	3.9	4.0	3.3	3.7	5.2	8.7	8.1
2000	4.4	*	1.2	1.4	8.0	3.9	3.7	3.4	4.1	5.3	8.5	8.6
1999	4.5	1.8	1.3	1.3	8.1	4.1	3.9	3.3	3.5	5.9	9.0	8.7
Non-Hispanic black, both sexes												
2002	5.2	*	1.5	1.3	7.6	6.9	5.8	6.1	5.7	6.3	9.1	7.3
2001	5.7	*	1.8	1.8	8.5	6.9	6.6	6.3	6.8	6.8	9.3	7.2
2000	6.1	3.8	2.0	1.4	9.1	8.4	6.5	7.6	5.8	8.1	7.7	8.9
1999	6.0	*	2.6	1.8	8.9	6.7	6.1	7.4	6.4	7.7	11.9	10.0
Non-Hispanic black, male												
2002	7.5	*	1.9	1.6	10.5	10.3	8.3	8.4	9.0	10.0	15.8	21.9
2001	8.2	*	1.9	2.2	12.3	10.1	9.6	9.9	10.7	9.0	16.2	*
2000	8.7	*	2.2	1.5	12.5	12.1	10.3	11.1	9.3	13.0	13.3	*
1999	8.4	*	3.1	2.1	12.5	10.1	8.4	11.3	9.7	9.6	21.3	23.8
Non-Hispanic black, female												
2002	3.2	*	*	0.9	4.6	3.8	3.6	4.0	3.0	3.7	5.3	*
2001	3.5	*	*	1.3	4.8	4.0	3.8	3.3	3.6	5.2	5.4	*
2000	3.8	*	1.9	1.3	5.8	5.1	3.2	4.6	2.9	4.6	4.5	*
1999	3.8	*	2.2	1.5	5.3	3.7	4.1	4.1	3.7	6.4	6.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Streetcar accidents (V82.2-V82.9)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Streetcar accidents (V82.2-V82.9)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
All origins 3/, both sexes												
2002	0.2	*	*	0.1	0.2	0.1	0.2	0.3	0.3	0.3	0.4	*
2001	0.3	*	0.2	0.1	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.6
2000	0.4	*	0.2	0.2	0.4	0.3	0.4	0.4	0.4	0.5	0.6	0.7
1999	0.6	*	0.3	0.3	0.7	0.6	0.6	0.6	0.6	0.7	1.2	1.2
All origins 3/, male												
2002	0.3	*	*	0.2	0.3	0.2	0.4	0.5	0.4	0.5	0.9	*
2001	0.5	*	*	0.2	0.5	0.3	0.6	0.6	0.6	0.8	0.9	*
2000	0.5	*	*	0.2	0.6	0.5	0.6	0.5	0.6	0.7	1.0	*
1999	0.8	*	0.3	0.3	1.0	0.8	0.9	1.0	1.0	1.1	1.7	2.7
All origins 3/, female												
2002	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
2001	0.2	*	*	0.1	0.1	*	0.2	0.2	0.2	0.3	0.5	*
2000	0.2	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.4	*
1999	0.3	*	0.3	0.2	0.4	0.3	0.3	0.3	0.3	0.4	0.9	*
Hispanic, both sexes												
2002	0.2	*	*	*	0.4	*	*	*	*	*	*	*
2001	0.3	*	*	*	0.4	*	0.4	*	*	*	*	*
2000	0.4	*	*	*	0.4	0.4	0.5	*	*	*	*	*
1999	0.6	*	*	*	0.6	0.6	0.8	0.8	*	1.9	*	*
Hispanic, male												
2002	0.3	*	*	*	0.6	*	*	*	*	*	*	*
2001	0.5	*	*	*	0.7	*	0.7	*	*	*	*	*
2000	0.6	*	*	*	0.6	0.6	0.8	*	*	*	*	*
1999	0.9	*	*	*	0.9	0.9	1.2	1.4	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	0.1	0.1	0.1	0.2	0.3	0.3	0.3	0.4	*
2001	0.3	*	*	0.2	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.5
2000	0.3	*	*	0.2	0.4	0.3	0.4	0.4	0.3	0.4	0.6	0.7
1999	0.6	*	0.3	0.3	0.7	0.5	0.6	0.6	0.7	0.6	1.2	1.2
Non-Hispanic 4/, male												
2002	0.3	*	*	0.2	0.2	0.2	0.4	0.5	0.4	0.6	0.8	*
2001	0.5	*	*	0.2	0.4	0.3	0.6	0.6	0.6	0.9	0.8	*
2000	0.5	*	*	0.2	0.6	0.5	0.6	0.5	0.6	0.7	0.9	*
1999	0.8	*	0.4	0.3	1.0	0.7	0.8	1.0	1.0	0.9	1.6	2.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.2	*	*	0.1	0.1	*	0.2	0.2	0.2	0.3	0.5	*
2000	0.2	*	*	0.2	0.2	0.2	0.2	0.3	*	*	0.4	*
1999	0.4	*	*	0.2	0.4	0.3	0.3	0.3	0.4	0.4	0.9	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	0.1	0.1	0.1	0.2	0.3	0.3	0.3	0.4	*
2001	0.3	*	*	0.1	0.3	0.2	0.3	0.4	0.3	0.5	0.6	*
2000	0.4	*	*	0.2	0.4	0.3	0.4	0.4	0.3	0.4	0.6	0.7
1999	0.6	*	0.3	0.3	0.7	0.5	0.6	0.6	0.7	0.6	1.1	1.2
Non-Hispanic white, male												
2002	0.3	*	*	0.2	0.2	0.2	0.3	0.5	0.4	0.5	0.8	*
2001	0.4	*	*	0.2	0.4	0.3	0.5	0.6	0.5	0.8	0.8	*
2000	0.5	*	*	0.3	0.6	0.5	0.5	0.5	0.5	0.7	1.0	*
1999	0.8	*	*	0.3	1.0	0.7	0.8	0.9	1.0	0.9	1.6	2.6
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	0.1	0.2	*	0.3	0.4	*
2000	0.2	*	*	0.2	0.2	0.2	0.2	0.3	*	*	0.4	*
1999	0.4	*	*	0.2	0.5	0.3	0.3	0.3	0.4	0.4	0.9	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	0.7	0.5	*	*	*	*
2000	0.3	*	*	*	*	*	0.5	*	*	*	*	*
1999	0.5	*	*	*	0.4	0.5	0.6	0.6	*	*	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.5	*	*	*	*	*	1.0	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.8	*	*	*	*	0.8	0.9	1.2	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water transport accidents (V90-V94)												
All origins 3/, both sexes												
2002	0.2	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.3	0.2	*
2001	0.2	*	*	0.1	0.3	0.2	0.2	0.3	0.3	0.2	*	*
2000	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.2	0.3	*	*
1999	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.3	0.2	*	*
All origins 3/, male												
2002	0.4	*	*	*	0.4	0.5	0.6	0.5	0.5	0.7	0.4	*
2001	0.4	*	*	0.1	0.5	0.4	0.4	0.5	0.5	0.5	*	*
2000	0.4	*	*	0.1	0.5	0.5	0.6	0.5	0.5	0.5	*	*
1999	0.4	*	*	0.2	0.5	0.5	0.5	0.6	0.6	0.5	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*
2001	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.3	0.3	*	*
2000	0.2	*	*	0.1	0.3	0.3	0.3	0.3	0.2	0.3	*	*
1999	0.3	*	*	0.1	0.3	0.3	0.3	0.3	0.3	0.3	*	*
Non-Hispanic 4/, male												
2002	0.4	*	*	*	0.4	0.5	0.6	0.5	0.5	0.7	0.4	*
2001	0.4	*	*	*	0.6	0.5	0.5	0.5	0.5	0.6	*	*
2000	0.4	*	*	0.1	0.5	0.6	0.6	0.5	0.5	0.6	*	*
1999	0.5	*	*	0.2	0.5	0.6	0.5	0.6	0.6	0.5	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water transport accidents (V90-V94)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.3	0.3	0.3	0.3	0.3	0.3	0.2	*
2001	0.2	*	*	0.1	0.4	0.3	0.3	0.3	0.3	0.2	*	*
2000	0.3	*	*	0.1	0.4	0.3	0.4	0.3	0.3	0.3	*	*
1999	0.3	*	*	0.1	0.4	0.4	0.3	0.3	0.3	0.2	*	*
Non-Hispanic white, male												
2002	0.4	*	*	*	0.5	0.6	0.6	0.5	0.5	0.7	*	*
2001	0.4	*	*	*	0.6	0.5	0.5	0.5	0.5	0.5	*	*
2000	0.5	*	*	*	0.6	0.6	0.6	0.5	0.5	0.6	*	*
1999	0.5	*	*	0.2	0.6	0.6	0.5	0.6	0.6	0.5	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	0.4	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Air and space transport accidents (V95-V97)												
All origins 3/, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.4	0.5	0.4	0.2	*
2001	0.3	*	*	*	0.2	0.4	0.5	0.5	0.5	0.5	0.2	*
2000	0.3	*	*	*	0.2	0.3	0.3	0.5	0.6	0.4	0.2	*
1999	0.3	*	*	0.0	0.1	0.3	0.3	0.4	0.5	0.4	0.2	*
All origins 3/, male												
2002	0.4	*	*	*	0.2	0.3	0.5	0.7	0.9	0.8	*	*
2001	0.5	*	*	*	0.2	0.6	0.7	0.8	0.8	0.8	0.5	*
2000	0.5	*	*	*	0.3	0.5	0.6	0.8	1.0	0.7	0.5	*
1999	0.4	*	*	*	0.2	0.5	0.5	0.7	1.0	0.7	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	0.1	0.1	*	*	*
2001	0.2	*	*	*	0.1	0.2	0.2	0.3	0.2	*	*	*
2000	0.1	*	*	*	*	0.1	*	0.1	0.2	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	0.5	1.0	1.4	1.7	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	1.1	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.6	*	*	*	*	*	0.8	1.7	2.0	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	*	0.1	0.2	0.3	0.4	0.5	0.4	0.2	*
2001	0.3	*	*	*	0.1	0.4	0.4	0.5	0.4	0.4	0.2	*
2000	0.3	*	*	*	0.2	0.4	0.3	0.5	0.6	0.4	0.2	*
1999	0.3	*	*	0.1	0.1	0.4	0.4	0.4	0.5	0.4	0.2	*
Non-Hispanic 4/, male												
2002	0.5	*	*	*	0.2	0.4	0.6	0.8	1.0	0.8	*	*
2001	0.5	*	*	*	0.2	0.6	0.7	0.8	0.8	0.7	*	*
2000	0.5	*	*	*	0.3	0.6	0.6	0.9	1.0	0.7	0.5	*
1999	0.5	*	*	*	0.2	0.6	0.6	0.7	1.0	0.7	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Air and space transport accidents (V95-V97)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.1	*	0.2	0.2	*	*	*
1999	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.3	*	*	*	0.2	0.3	0.4	0.5	0.6	0.4	0.2	*
2001	0.3	*	*	*	0.2	0.5	0.5	0.5	0.5	0.4	0.2	*
2000	0.4	*	*	*	0.2	0.5	0.4	0.6	0.7	0.4	0.2	*
1999	0.3	*	*	*	0.2	0.4	0.5	0.5	0.6	0.5	0.2	*
Non-Hispanic white, male												
2002	0.6	*	*	*	0.3	0.4	0.7	0.9	1.1	0.9	*	*
2001	0.6	*	*	*	0.2	0.8	0.8	0.9	0.9	0.8	*	*
2000	0.6	*	*	*	0.4	0.8	0.8	1.0	1.2	0.7	0.5	*
1999	0.6	*	*	*	0.3	0.7	0.7	0.8	1.2	0.8	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.2	*	0.2	0.2	*	*	*
1999	0.1	*	*	*	*	0.2	0.2	0.2	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)												
All origins 3/, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.4	0.3	0.2	0.4	*
2001	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3	0.4	*
2000	0.2	*	*	*	0.1	0.2	0.3	0.2	0.3	0.3	0.3	*
1999	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.2	0.3	*
All origins 3/, male												
2002	0.3	*	*	*	0.2	0.3	0.4	0.6	0.5	0.4	0.7	*
2001	0.3	*	*	*	0.2	0.3	0.5	0.6	0.4	0.4	0.6	*
2000	0.3	*	*	*	0.1	0.3	0.4	0.4	0.4	0.5	0.5	*
1999	0.3	*	*	*	0.1	0.3	0.4	0.4	0.4	0.4	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*
2001	0.1	*	*	*	*	0.1	0.2	0.1	*	*	0.3	*
2000	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	0.2	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.4	0.3	0.2	0.4	*
2001	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.2	0.4	*
2000	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3	0.3	*
1999	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.2	0.3	*
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.2	0.3	0.5	0.6	0.5	0.4	0.7	*
2001	0.3	*	*	*	0.2	0.3	0.5	0.6	0.4	0.4	0.6	*
2000	0.3	*	*	*	0.1	0.3	0.4	0.4	0.4	0.5	0.5	*
1999	0.3	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified transport accidents and their sequelae (V98-V99,Y85)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	*	*	0.3	*
2000	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.2	0.2	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.4	0.3	0.2	0.4	*
2001	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3	0.4	*
2000	0.2	*	*	*	0.1	0.2	0.3	0.2	0.2	0.3	0.3	*
1999	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic white, male												
2002	0.3	*	*	*	0.2	0.3	0.5	0.6	0.5	0.4	0.6	*
2001	0.3	*	*	*	0.2	0.3	0.5	0.5	0.4	0.4	0.6	*
2000	0.3	*	*	*	*	0.3	0.4	0.4	0.4	0.4	0.5	*
1999	0.3	*	*	*	*	0.3	0.4	0.4	0.3	0.4	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	0.2	0.2	0.2	*	*	*
2001	0.1	*	*	*	*	*	0.2	*	*	*	*	*
2000	0.1	*	*	*	*	*	0.2	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	0.7	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	1.2	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All origins 3/, both sexes												
2002	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2
All origins 3/, male												
2002	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
All origins 3/, female												
2002	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2
Hispanic, both sexes												
2002	2.1	*	*	*	0.7	0.9	1.6	2.5	4.4	10.0	29.2	71.5
2001	2.0	*	*	*	0.7	1.1	1.8	2.6	4.2	9.1	23.6	70.0
2000	2.0	*	*	*	0.7	1.1	1.3	2.5	4.3	8.0	26.5	81.0
1999	1.9	*	*	*	0.7	1.1	1.4	2.2	3.8	8.1	24.2	86.3
Hispanic, male												
2002	2.8	*	*	*	1.3	1.6	2.7	3.8	7.0	13.8	36.9	97.0
2001	2.8	*	*	*	1.2	1.9	3.3	4.5	7.1	11.4	27.7	72.4
2000	2.7	*	*	*	1.2	2.0	2.3	4.4	6.6	12.7	30.5	104.8
1999	2.7	*	*	*	1.3	1.9	2.4	3.8	7.2	11.4	31.3	120.8
Hispanic, female												
2002	1.4	*	*	*	*	*	*	1.1	2.1	7.0	23.9	58.9
2001	1.3	*	*	*	*	*	*	*	*	7.2	20.9	68.9
2000	1.2	*	*	*	*	*	*	*	2.3	4.3	23.8	69.2
1999	1.1	*	*	*	*	*	*	*	*	5.5	19.4	69.7
Non-Hispanic 4/, both sexes												
2002	6.2	*	0.2	0.1	0.6	0.7	1.5	2.5	4.0	10.8	38.6	132.6
2001	5.7	*	0.2	0.1	0.6	0.8	1.4	2.6	3.9	10.0	35.8	123.1
2000	5.1	*	0.2	0.1	0.6	0.7	1.3	2.3	3.9	9.1	31.2	113.5
1999	5.1	*	0.4	0.1	0.6	0.8	1.4	2.2	3.7	9.0	31.7	110.7
Non-Hispanic 4/, male												
2002	6.5	*	0.3	0.1	0.9	1.3	2.3	3.9	5.9	14.4	48.4	161.3
2001	6.2	*	0.3	0.2	1.1	1.4	2.3	4.2	5.8	13.1	46.4	159.0
2000	5.5	*	*	0.2	1.0	1.2	2.2	3.6	5.7	11.9	40.2	146.5
1999	5.5	*	0.5	0.2	1.0	1.3	2.3	3.6	5.5	12.4	40.6	142.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Non-Hispanic 4/, female												
2002	5.9	*	*	*	0.2	0.2	0.7	1.2	2.3	7.7	32.1	120.2
2001	5.3	*	*	*	0.1	0.3	0.5	1.0	2.2	7.4	28.8	108.1
2000	4.7	*	*	*	0.2	0.2	0.5	1.0	2.1	6.8	25.3	100.1
1999	4.6	*	*	*	0.2	0.3	0.5	0.9	2.0	6.3	25.9	97.8
Non-Hispanic white, both sexes												
2002	7.1	*	*	0.1	0.7	0.8	1.5	2.5	4.1	11.1	40.8	140.4
2001	6.6	*	0.2	0.2	0.8	0.9	1.4	2.6	3.9	10.3	37.6	129.9
2000	5.8	*	*	0.1	0.7	0.7	1.4	2.3	3.8	9.2	32.7	120.2
1999	5.7	*	0.4	0.2	0.7	0.8	1.5	2.1	3.5	9.1	33.1	116.6
Non-Hispanic white, male												
2002	7.3	*	*	*	1.0	1.4	2.3	3.9	5.9	14.8	50.6	169.2
2001	7.0	*	*	0.2	1.3	1.5	2.3	4.1	5.7	13.5	48.2	166.0
2000	6.2	*	*	0.2	1.1	1.3	2.2	3.6	5.6	12.0	41.7	154.1
1999	6.1	*	0.5	0.2	1.1	1.4	2.3	3.4	5.1	12.2	42.0	149.7
Non-Hispanic white, female												
2002	7.0	*	*	*	0.2	0.2	0.7	1.2	2.4	8.1	34.2	128.0
2001	6.2	*	*	*	0.2	0.3	0.5	1.1	2.2	7.7	30.6	114.8
2000	5.5	*	*	*	0.2	0.2	0.5	1.0	2.1	6.9	26.8	106.5
1999	5.4	*	*	*	0.2	0.3	0.6	0.8	1.9	6.4	27.2	103.4
Non-Hispanic black, both sexes												
2002	2.3	*	*	*	*	0.4	1.3	2.8	4.3	8.2	16.3	49.3
2001	2.3	*	*	*	*	0.5	1.6	3.0	4.4	8.8	18.7	44.0
2000	2.1	*	*	*	*	0.4	1.3	2.7	4.8	8.3	15.2	39.1
1999	2.5	*	*	*	0.4	0.8	1.3	3.5	5.5	9.3	17.2	45.3
Non-Hispanic black, male												
2002	3.0	*	*	*	*	0.8	2.2	4.7	6.8	11.9	24.9	76.6
2001	3.2	*	*	*	*	*	2.7	5.4	6.9	12.6	29.1	68.1
2000	2.8	*	*	*	*	*	2.1	4.6	8.2	11.4	21.6	61.1
1999	3.4	*	*	*	*	1.4	2.5	5.9	8.7	15.0	24.1	58.2
Non-Hispanic black, female												
2002	1.6	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.9
2001	1.6	*	*	*	*	*	*	*	2.3	6.1	12.8	35.1
2000	1.5	*	*	*	*	*	*	1.0	2.0	6.1	11.6	31.0
1999	1.7	*	*	*	*	*	*	1.4	2.9	5.3	13.3	40.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fall on same level (W00-W09,W18)												
All origins 3/, both sexes												
2002	1.8	*	*	*	0.1	0.1	0.3	0.6	1.1	3.1	12.6	50.8
2001	1.4	*	*	0.1	0.1	0.1	0.2	0.6	1.0	2.6	9.9	37.3
2000	1.2	*	*	*	0.1	0.1	0.2	0.4	0.8	2.0	8.0	33.0
1999	0.8	*	*	*	0.1	0.1	0.1	0.3	0.4	1.5	5.8	21.4
All origins 3/, male												
2002	1.8	*	*	*	0.2	0.2	0.4	0.9	1.4	4.2	15.2	60.8
2001	1.4	*	*	*	0.2	0.2	0.3	0.8	1.3	3.0	12.7	45.3
2000	1.1	*	*	*	0.1	0.1	0.3	0.6	1.0	2.5	9.8	39.6
1999	0.8	*	*	*	0.1	*	0.2	0.5	0.5	2.0	7.3	26.6
All origins 3/, female												
2002	1.9	*	*	*	*	*	0.1	0.3	0.8	2.2	10.9	46.4
2001	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	8.0	33.9
2000	1.2	*	*	*	*	*	*	0.2	0.6	1.6	6.9	30.4
1999	0.8	*	*	*	*	*	*	0.2	0.3	1.1	4.9	19.4
Hispanic, both sexes												
2002	0.6	*	*	*	*	*	*	0.6	1.3	3.1	10.7	37.1
2001	0.5	*	*	*	*	*	*	*	*	3.0	8.1	26.9
2000	0.4	*	*	*	*	*	*	*	*	2.0	9.3	24.6
1999	0.3	*	*	*	*	*	*	*	*	*	6.9	21.0
Hispanic, male												
2002	0.7	*	*	*	*	*	*	*	*	4.4	12.3	55.2
2001	0.6	*	*	*	*	*	*	*	*	4.0	9.8	*
2000	0.4	*	*	*	*	*	*	*	*	*	10.8	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	9.6	28.2
2001	0.5	*	*	*	*	*	*	*	*	*	7.0	25.1
2000	0.4	*	*	*	*	*	*	*	*	*	8.3	22.8
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	2.0	*	*	*	0.1	0.1	0.3	0.6	1.0	3.1	12.6	51.3
2001	1.5	*	*	0.1	0.1	0.1	0.2	0.5	1.0	2.5	9.9	37.5
2000	1.3	*	*	*	0.1	0.1	0.2	0.4	0.8	2.0	8.0	33.3
1999	0.9	*	*	*	0.1	0.1	0.1	0.3	0.4	1.5	5.7	21.4
Non-Hispanic 4/, male												
2002	1.9	*	*	*	0.2	0.2	0.4	0.8	1.3	4.2	15.3	61.1
2001	1.5	*	*	*	0.2	0.2	0.3	0.8	1.3	2.9	12.8	45.9
2000	1.2	*	*	*	*	*	0.3	0.6	1.0	2.5	9.7	39.9
1999	0.8	*	*	*	0.1	*	0.2	0.5	0.5	2.0	7.2	26.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fall on same level (W00-W09,W18)												
Non-Hispanic 4/, female												
2002	2.1	*	*	*	*	*	0.1	0.3	0.8	2.2	10.9	47.1
2001	1.6	*	*	*	*	*	0.1	0.3	0.6	2.1	8.0	34.0
2000	1.3	*	*	*	*	*	*	0.2	0.6	1.6	6.8	30.6
1999	0.9	*	*	*	*	*	*	0.2	0.3	1.1	4.8	19.4
Non-Hispanic white, both sexes												
2002	2.4	*	*	*	0.1	0.1	0.3	0.6	1.1	3.3	13.3	54.4
2001	1.8	*	*	*	0.1	0.1	0.2	0.6	1.0	2.6	10.4	39.6
2000	1.5	*	*	*	0.1	*	0.2	0.4	0.7	2.1	8.4	35.3
1999	1.0	*	*	*	0.1	0.1	0.1	0.3	0.3	1.5	5.9	22.6
Non-Hispanic white, male												
2002	2.2	*	*	*	0.2	0.2	0.4	0.9	1.4	4.3	15.7	63.6
2001	1.7	*	*	*	0.2	0.2	0.3	0.8	1.3	3.1	13.2	48.3
2000	1.4	*	*	*	*	*	0.3	0.6	0.9	2.6	10.1	42.3
1999	0.9	*	*	*	*	*	0.2	0.4	0.4	2.0	7.3	27.8
Non-Hispanic white, female												
2002	2.5	*	*	*	*	*	0.1	0.3	0.8	2.4	11.7	50.4
2001	1.8	*	*	*	*	*	*	0.3	0.6	2.2	8.5	36.0
2000	1.6	*	*	*	*	*	*	0.2	0.6	1.7	7.3	32.5
1999	1.0	*	*	*	*	*	*	0.2	0.2	1.1	5.0	20.6
Non-Hispanic black, both sexes												
2002	0.5	*	*	*	*	*	*	0.5	*	1.9	4.6	16.3
2001	0.5	*	*	*	*	*	*	0.5	1.1	1.8	4.3	11.5
2000	0.4	*	*	*	*	*	*	*	1.1	1.3	2.9	9.2
1999	0.4	*	*	*	*	*	*	0.6	0.9	1.8	3.5	7.1
Non-Hispanic black, male												
2002	0.7	*	*	*	*	*	*	*	*	3.1	9.1	26.3
2001	0.6	*	*	*	*	*	*	*	*	*	6.6	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	12.5
2001	0.4	*	*	*	*	*	*	*	*	*	*	10.3
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fall from one level to another (W10-W17)												
All origins 3/, both sexes												
2002	1.2	*	0.1	*	0.4	0.5	0.8	0.9	1.4	2.8	6.1	10.9
2001	1.2	*	0.1	0.1	0.5	0.6	0.7	1.1	1.4	2.7	6.1	11.3
2000	1.0	*	0.2	*	0.4	0.5	0.7	1.0	1.3	2.4	5.1	8.8
1999	1.1	*	0.2	0.1	0.4	0.6	0.7	0.9	1.5	2.6	5.8	10.4
All origins 3/, male												
2002	1.7	*	*	*	0.7	0.9	1.3	1.6	2.4	4.3	9.7	17.4
2001	1.7	*	*	0.1	0.8	1.0	1.3	2.0	2.4	4.2	9.5	18.2
2000	1.6	*	*	*	0.8	1.0	1.3	1.8	2.2	3.8	8.4	16.4
1999	1.7	*	0.3	0.1	0.7	1.0	1.3	1.6	2.6	4.1	9.0	17.8
All origins 3/, female												
2002	0.6	*	*	*	0.1	0.1	0.2	0.3	0.5	1.5	3.7	8.1
2001	0.6	*	*	*	*	0.1	0.1	0.3	0.5	1.4	3.9	8.3
2000	0.5	*	*	*	0.1	*	0.2	0.3	0.5	1.2	2.9	5.7
1999	0.6	*	*	*	0.1	0.1	0.2	0.2	0.6	1.4	3.7	7.4
Hispanic, both sexes												
2002	0.7	*	*	*	0.6	0.7	0.8	0.9	1.4	2.3	*	*
2001	0.7	*	*	*	0.5	0.8	0.9	1.2	1.7	2.1	3.8	*
2000	0.7	*	*	*	0.6	0.8	0.7	1.3	1.7	2.1	*	*
1999	0.6	*	*	*	0.4	0.8	0.9	0.8	1.3	*	*	*
Hispanic, male												
2002	1.1	*	*	*	1.0	1.3	1.5	1.4	2.7	4.0	*	*
2001	1.3	*	*	*	0.9	1.4	1.6	2.4	3.2	*	*	*
2000	1.2	*	*	*	1.0	1.5	1.4	2.5	2.9	*	*	*
1999	1.1	*	*	*	0.8	1.4	1.5	1.5	2.6	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	*	0.4	0.5	0.7	0.9	1.4	2.8	6.3	11.2
2001	1.2	*	*	0.1	0.4	0.5	0.7	1.1	1.3	2.7	6.2	11.5
2000	1.1	*	*	*	0.4	0.5	0.7	1.0	1.3	2.4	5.2	8.8
1999	1.2	*	0.2	0.1	0.4	0.5	0.7	0.9	1.5	2.6	5.9	10.5
Non-Hispanic 4/, male												
2002	1.8	*	*	*	0.6	0.9	1.3	1.6	2.4	4.3	9.9	17.8
2001	1.8	*	*	*	0.8	0.9	1.3	1.9	2.3	4.2	9.6	19.0
2000	1.6	*	*	*	0.7	0.9	1.2	1.7	2.2	3.8	8.6	16.4
1999	1.7	*	*	*	0.7	1.0	1.3	1.6	2.6	4.2	9.3	18.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Fall from one level to another (W10-W17)												
Non-Hispanic 4/, female												
2002	0.7	*	*	*	0.1	0.1	0.2	0.3	0.5	1.6	3.8	8.3
2001	0.7	*	*	*	*	0.1	0.1	0.3	0.5	1.4	4.0	8.4
2000	0.6	*	*	*	0.1	*	0.2	0.3	0.5	1.2	3.0	5.8
1999	0.7	*	*	*	*	0.1	0.2	0.2	0.6	1.4	3.8	7.4
Non-Hispanic white, both sexes												
2002	1.4	*	*	*	0.4	0.6	0.8	1.0	1.5	3.0	6.6	11.5
2001	1.4	*	*	*	0.5	0.6	0.7	1.1	1.3	2.8	6.5	12.0
2000	1.2	*	*	*	0.5	0.5	0.8	1.0	1.3	2.5	5.5	9.2
1999	1.3	*	0.2	*	0.5	0.6	0.8	0.9	1.6	2.8	6.2	11.1
Non-Hispanic white, male												
2002	2.0	*	*	*	0.7	1.0	1.3	1.7	2.5	4.6	10.4	18.5
2001	2.0	*	*	*	1.0	1.0	1.3	1.8	2.3	4.3	10.0	19.8
2000	1.8	*	*	*	0.9	1.0	1.3	1.7	2.2	3.9	9.0	17.0
1999	1.9	*	*	*	0.8	1.0	1.3	1.6	2.6	4.2	9.5	19.6
Non-Hispanic white, female												
2002	0.8	*	*	*	0.2	*	0.2	0.3	0.5	1.6	4.0	8.5
2001	0.8	*	*	*	*	0.2	0.2	0.3	0.5	1.5	4.2	8.7
2000	0.6	*	*	*	*	*	0.2	0.3	0.5	1.3	3.2	6.1
1999	0.8	*	*	*	*	*	0.2	0.2	0.7	1.5	4.0	7.7
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	*	*	0.6	0.8	1.2	1.8	2.9	7.6
2001	0.7	*	*	*	*	*	0.7	1.3	1.2	2.1	3.3	6.2
2000	0.6	*	*	*	*	*	0.5	1.1	1.3	2.0	*	*
1999	0.6	*	*	*	*	0.5	0.6	1.0	1.0	1.9	2.7	*
Non-Hispanic black, male												
2002	0.8	*	*	*	*	*	1.1	1.3	2.0	*	*	*
2001	1.1	*	*	*	*	*	1.3	2.5	2.4	4.3	6.6	*
2000	1.0	*	*	*	*	*	0.8	2.2	2.5	3.6	*	*
1999	1.0	*	*	*	*	0.8	1.2	1.8	2.0	4.1	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified fall (W19)												
All origins 3/, both sexes												
2002	2.7	*	*	*	0.1	0.1	0.5	1.0	1.6	4.8	19.6	68.8
2001	2.7	*	*	*	0.1	0.2	0.5	1.0	1.6	4.8	19.3	73.0
2000	2.5	*	*	*	0.1	0.1	0.4	0.9	1.8	4.6	17.9	70.7
1999	2.8	*	*	*	0.1	0.2	0.6	1.0	1.8	4.9	19.9	78.4
All origins 3/, male												
2002	2.5	*	*	*	0.1	0.2	0.7	1.4	2.2	5.9	23.2	80.7
2001	2.6	*	*	*	0.1	0.3	0.8	1.5	2.2	5.9	23.4	92.1
2000	2.5	*	*	*	0.2	0.2	0.7	1.4	2.6	5.6	21.7	89.4
1999	2.8	*	*	*	0.2	0.3	0.9	1.6	2.6	6.3	24.2	98.1
All origins 3/, female												
2002	2.8	*	*	*	*	*	0.3	0.6	1.0	3.9	17.2	63.6
2001	2.7	*	*	*	*	*	0.2	0.4	1.1	3.9	16.6	64.9
2000	2.6	*	*	*	*	*	0.2	0.5	1.0	3.8	15.5	63.1
1999	2.8	*	*	*	*	0.1	0.3	0.5	1.0	3.8	17.1	70.4
Hispanic, both sexes												
2002	0.8	*	*	*	*	*	0.5	0.9	1.7	4.6	15.5	31.6
2001	0.8	*	*	*	*	*	0.6	0.9	1.5	4.0	11.7	40.0
2000	0.8	*	*	*	*	*	0.4	0.9	1.7	3.9	14.6	49.1
1999	1.0	*	*	*	*	*	0.5	1.1	2.0	5.1	14.6	58.9
Hispanic, male												
2002	0.9	*	*	*	*	*	0.9	1.4	2.2	5.4	20.5	35.1
2001	0.9	*	*	*	*	*	1.2	1.4	2.6	4.0	10.7	41.9
2000	1.0	*	*	*	*	*	0.8	1.5	2.7	6.3	15.2	60.5
1999	1.2	*	*	*	*	*	0.8	1.9	4.0	6.8	18.8	82.0
Hispanic, female												
2002	0.7	*	*	*	*	*	*	*	*	4.0	12.1	29.8
2001	0.7	*	*	*	*	*	*	*	*	3.9	12.4	39.1
2000	0.7	*	*	*	*	*	*	*	*	*	14.2	43.5
1999	0.7	*	*	*	*	*	*	*	*	3.8	11.8	47.8
Non-Hispanic 4/, both sexes												
2002	2.9	*	*	*	0.1	0.1	0.4	1.0	1.6	4.8	19.7	70.0
2001	3.0	*	*	*	0.1	0.2	0.5	1.0	1.6	4.8	19.7	74.1
2000	2.8	*	*	*	0.1	0.1	0.4	0.9	1.8	4.6	18.1	71.4
1999	3.0	*	*	*	0.1	0.2	0.6	1.0	1.7	4.9	20.0	78.8
Non-Hispanic 4/, male												
2002	2.8	*	*	*	*	0.2	0.6	1.4	2.2	6.0	23.2	82.4
2001	2.9	*	*	*	0.1	0.3	0.7	1.5	2.2	6.0	24.0	94.1
2000	2.7	*	*	*	0.1	0.2	0.6	1.4	2.6	5.6	21.9	90.2
1999	3.0	*	*	*	0.2	0.3	0.9	1.5	2.5	6.2	24.2	98.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified fall (W19)												
Non-Hispanic 4/, female												
2002	3.1	*	*	*	*	*	0.3	0.6	1.0	3.9	17.4	64.7
2001	3.0	*	*	*	*	*	0.2	0.5	1.1	3.9	16.8	65.7
2000	2.9	*	*	*	*	*	0.2	0.5	1.0	3.9	15.5	63.7
1999	3.1	*	*	*	*	*	0.3	0.5	1.0	3.8	17.3	70.9
Non-Hispanic white, both sexes												
2002	3.4	*	*	*	0.1	0.1	0.4	0.9	1.5	4.9	20.9	74.5
2001	3.4	*	*	*	0.1	0.2	0.4	1.0	1.6	4.9	20.6	78.3
2000	3.2	*	*	*	0.1	0.1	0.4	0.9	1.7	4.7	18.8	75.6
1999	3.4	*	*	*	0.1	0.2	0.6	0.9	1.5	4.8	21.0	82.8
Non-Hispanic white, male												
2002	3.1	*	*	*	*	*	0.6	1.3	2.0	5.9	24.4	87.1
2001	3.3	*	*	*	0.2	0.3	0.7	1.5	2.1	6.0	24.9	97.9
2000	3.0	*	*	*	*	0.2	0.6	1.3	2.4	5.6	22.6	94.9
1999	3.3	*	*	*	0.2	0.3	0.8	1.4	2.1	6.1	25.2	102.2
Non-Hispanic white, female												
2002	3.7	*	*	*	*	*	0.3	0.6	1.0	4.0	18.6	69.1
2001	3.6	*	*	*	*	*	0.2	0.5	1.1	4.0	17.8	70.1
2000	3.3	*	*	*	*	*	0.2	0.5	1.0	3.9	16.3	67.9
1999	3.6	*	*	*	*	*	0.3	0.4	0.9	3.8	18.2	75.1
Non-Hispanic black, both sexes												
2002	1.2	*	*	*	*	*	0.5	1.5	2.4	4.5	8.9	25.4
2001	1.2	*	*	*	*	*	0.7	1.2	2.1	4.8	11.0	26.2
2000	1.2	*	*	*	*	*	0.7	1.2	2.4	4.9	10.2	26.1
1999	1.5	*	*	*	*	*	0.6	1.8	3.5	5.7	11.0	33.3
Non-Hispanic black, male												
2002	1.5	*	*	*	*	*	0.8	2.7	4.0	6.4	10.8	38.3
2001	1.4	*	*	*	*	*	1.0	2.0	3.0	6.2	15.9	46.2
2000	1.5	*	*	*	*	*	1.1	1.9	4.2	6.1	14.5	38.7
1999	1.9	*	*	*	*	*	1.1	3.1	5.5	8.4	14.7	46.4
Non-Hispanic black, female												
2002	0.8	*	*	*	*	*	*	*	*	3.1	7.8	20.5
2001	0.9	*	*	*	*	*	*	*	*	3.8	8.3	18.8
2000	0.9	*	*	*	*	*	*	*	*	4.1	7.8	21.4
1999	1.1	*	*	*	*	*	*	*	1.9	3.7	8.9	28.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
All origins 3/, both sexes												
2002	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
All origins 3/, male												
2002	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	0.4	*	*	*	*	*	*	*
2001	0.2	*	*	*	0.5	0.3	*	*	*	*	*	*
2000	0.2	*	*	*	0.4	*	*	*	*	*	*	*
1999	0.2	*	*	*	0.5	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	0.6	*	*	*	*	*	*	*
2001	0.4	*	*	*	0.9	*	*	*	*	*	*	*
2000	0.3	*	*	*	0.7	*	*	*	*	*	*	*
1999	0.3	*	*	*	0.9	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	*	*	0.1	0.5	0.4	0.3	0.3	0.2	0.2	0.2	*
2001	0.3	*	*	0.2	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.4	0.4	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.7	0.4	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic 4/, male												
2002	0.5	*	*	0.2	1.0	0.7	0.5	0.5	0.4	0.4	0.5	*
2001	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	1.0	0.6	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.2	0.6	0.5	0.5	0.4	0.4	0.4	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.2	*
2001	0.3	*	*	0.2	0.5	0.3	0.4	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.3	0.4	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic white, male												
2002	0.5	*	*	0.2	0.8	0.6	0.5	0.5	0.4	0.5	0.5	*
2001	0.5	*	*	0.3	0.8	0.6	0.6	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.0	0.6	0.5	0.5	0.4	0.4	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.4	*	*	*	1.0	0.8	0.4	*	*	*	*	*
2001	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2000	0.4	*	*	*	0.9	0.6	*	*	*	*	*	*
1999	0.4	*	*	*	1.2	0.4	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.8	*	*	*	1.9	1.6	*	*	*	*	*	*
2001	0.6	*	*	*	1.6	*	*	*	*	*	*	*
2000	0.7	*	*	*	1.6	1.2	*	*	*	*	*	*
1999	0.8	*	*	*	2.2	0.9	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental explosion (W39-W40)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	0.1	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	0.1	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	0.2	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.2	0.2	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental explosion (W39-W40)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	0.2	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All origins 3/, both sexes												
2002	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6
All origins 3/, male												
2002	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
All origins 3/, female												
2002	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4
Hispanic, both sexes												
2002	1.2	*	2.1	0.4	2.1	1.1	0.9	0.8	*	*	*	*
2001	1.1	*	2.6	0.6	1.5	1.1	1.0	0.7	*	*	*	*
2000	1.2	*	2.5	0.6	2.2	1.0	0.8	0.6	*	*	*	*
1999	1.4	*	2.7	0.7	2.3	1.3	0.9	0.8	*	*	*	*
Hispanic, male												
2002	2.0	*	3.0	0.6	3.7	2.0	1.6	1.5	*	*	*	*
2001	1.8	*	3.5	0.8	2.7	1.8	1.7	1.2	*	*	*	*
2000	1.9	*	2.9	0.6	3.8	1.7	1.4	*	*	*	*	*
1999	2.3	*	2.9	1.1	4.1	2.2	1.5	1.6	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	1.7	*	*	*	*	*	*	*	*	*
2000	0.5	*	2.2	0.6	*	*	*	*	*	*	*	*
1999	0.5	*	2.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	1.4	3.1	0.9	1.4	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001	1.1	1.6	3.1	0.9	1.5	0.9	1.0	0.9	0.8	0.9	1.3	1.7
2000	1.2	2.1	3.4	1.0	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.5
1999	1.2	1.7	3.3	0.9	1.5	1.1	1.1	1.0	0.9	1.0	1.3	1.7
Non-Hispanic 4/, male												
2002	1.9	1.6	4.1	1.3	2.6	1.8	2.0	1.5	1.3	1.6	2.0	3.6
2001	1.8	1.3	3.8	1.2	2.6	1.6	1.7	1.6	1.3	1.5	2.1	3.4
2000	2.0	2.4	4.3	1.4	2.7	1.8	1.9	1.6	1.6	1.6	1.8	2.8
1999	2.0	1.7	4.2	1.4	2.7	1.9	1.8	1.7	1.5	1.6	2.3	2.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Non-Hispanic 4/, female												
2002	0.5	*	2.1	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001	0.5	1.9	2.4	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.7	1.0
2000	0.5	1.9	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.4	0.6	0.6	1.4
Non-Hispanic white, both sexes												
2002	1.1	1.4	3.3	0.5	1.2	1.0	1.2	0.9	0.8	0.9	1.2	1.6
2001	1.1	1.5	3.5	0.5	1.3	0.8	1.0	0.9	0.8	0.9	1.2	1.7
2000	1.1	1.7	3.7	0.7	1.3	1.0	1.1	0.9	0.9	1.0	1.3	1.4
1999	1.1	1.3	3.6	0.5	1.3	1.0	1.1	1.0	0.9	1.0	1.2	1.6
Non-Hispanic white, male												
2002	1.7	*	4.3	0.7	2.2	1.7	1.9	1.4	1.2	1.5	2.0	3.7
2001	1.6	*	4.3	0.7	2.2	1.4	1.5	1.5	1.2	1.5	2.1	3.5
2000	1.8	1.9	4.7	1.0	2.2	1.7	1.8	1.5	1.6	1.5	1.8	2.7
1999	1.8	*	4.5	0.7	2.2	1.6	1.8	1.7	1.5	1.4	2.2	2.4
Non-Hispanic white, female												
2002	0.5	*	2.2	0.3	0.3	0.3	0.4	0.4	0.4	0.3	0.7	0.7
2001	0.5	1.7	2.6	0.4	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000	0.5	*	2.6	0.4	0.3	0.3	0.4	0.3	0.3	0.5	0.9	0.9
1999	0.5	*	2.6	0.3	0.3	0.3	0.4	0.4	0.4	0.6	0.6	1.3
Non-Hispanic black, both sexes												
2002	1.7	*	2.6	2.1	2.1	1.4	1.3	1.3	0.8	*	*	*
2001	1.6	*	2.0	2.1	2.3	1.2	1.3	1.2	1.3	*	*	*
2000	1.7	4.3	2.3	2.1	2.3	1.1	1.4	1.3	*	*	*	*
1999	1.8	3.7	2.5	2.5	2.5	1.6	1.1	0.9	1.2	1.3	*	*
Non-Hispanic black, male												
2002	2.9	*	3.5	3.3	4.1	2.7	2.5	2.3	*	*	*	*
2001	2.8	*	2.4	3.1	4.4	2.5	2.2	2.1	2.4	*	*	*
2000	2.8	*	2.8	3.1	4.4	2.1	2.6	2.4	*	*	*	*
1999	3.2	*	3.4	3.8	4.6	3.2	2.1	1.9	2.1	*	*	*
Non-Hispanic black, female												
2002	0.5	*	*	0.8	*	*	*	*	*	*	*	*
2001	0.5	*	*	1.0	*	*	*	*	*	*	*	*
2000	0.6	*	1.8	1.0	*	*	*	*	*	*	*	*
1999	0.6	*	*	1.1	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)												
All origins 3/, both sexes												
2002	1.4	1.5	0.5	0.1	0.1	0.2	0.4	0.6	1.2	3.1	9.5	28.4
2001	1.5	1.5	0.5	0.1	0.1	0.1	0.4	0.9	1.3	2.9	9.5	30.0
2000	1.5	2.0	0.6	0.1	0.1	0.2	0.4	0.8	1.4	3.1	10.0	32.6
1999	1.4	1.7	0.6	0.1	0.1	0.2	0.4	0.7	1.1	3.1	8.8	30.3
All origins 3/, male												
2002	1.5	2.0	0.6	0.1	0.2	0.3	0.5	0.8	1.6	3.9	12.1	37.0
2001	1.5	2.0	0.6	*	0.2	0.2	0.5	1.1	1.6	3.5	12.4	39.8
2000	1.6	2.3	0.7	*	0.2	0.3	0.5	1.1	1.6	3.6	13.2	41.0
1999	1.4	1.9	0.8	0.1	0.1	0.3	0.5	0.8	1.3	3.8	10.9	39.2
All origins 3/, female												
2002	1.4	1.0	0.3	*	*	0.1	0.3	0.5	0.9	2.4	7.8	24.6
2001	1.4	1.0	0.4	*	0.1	0.1	0.3	0.7	1.0	2.4	7.6	26.0
2000	1.5	1.8	0.5	*	*	0.2	0.3	0.5	1.1	2.7	7.9	29.1
1999	1.4	1.4	0.4	0.1	*	0.1	0.2	0.5	1.0	2.5	7.4	26.7
Hispanic, both sexes												
2002	0.4	*	0.7	*	*	*	*	*	*	2.0	6.7	12.7
2001	0.5	*	0.8	*	*	*	*	*	*	2.1	6.1	15.6
2000	0.5	*	0.7	*	*	*	*	*	*	*	6.3	23.2
1999	0.5	*	*	*	*	*	*	*	*	2.3	5.2	22.5
Hispanic, male												
2002	0.4	*	*	*	*	*	*	*	*	*	9.0	*
2001	0.5	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	9.8	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	19.8
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.6	1.6	0.4	0.1	0.1	0.2	0.4	0.7	1.3	3.2	9.6	28.9
2001	1.6	1.7	0.4	0.1	0.2	0.2	0.4	0.9	1.3	3.0	9.6	30.5
2000	1.7	2.2	0.6	*	0.1	0.3	0.4	0.8	1.4	3.2	10.1	32.8
1999	1.5	1.8	0.6	0.1	0.1	0.2	0.4	0.7	1.2	3.1	8.9	30.4
Non-Hispanic 4/, male												
2002	1.7	2.1	0.5	*	0.2	0.3	0.5	0.9	1.7	4.0	12.2	37.6
2001	1.7	2.2	0.4	*	0.2	0.2	0.5	1.1	1.6	3.6	12.5	40.4
2000	1.7	2.5	0.7	*	0.2	0.3	0.5	1.1	1.7	3.7	13.3	41.2
1999	1.5	2.0	0.7	0.1	0.1	0.3	0.5	0.9	1.3	3.8	11.1	39.5

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)												
Non-Hispanic 4/, female												
2002	1.5	*	*	*	*	0.2	0.3	0.5	0.9	2.5	7.9	25.2
2001	1.6	*	0.4	*	0.1	0.1	0.3	0.7	1.1	2.5	7.7	26.4
2000	1.7	1.9	0.5	*	*	0.2	0.3	0.5	1.1	2.7	8.1	29.4
1999	1.5	1.6	0.5	*	*	0.2	0.3	0.5	1.0	2.6	7.4	26.8
Non-Hispanic white, both sexes												
2002	1.7	1.4	0.3	*	0.1	0.3	0.4	0.7	1.2	2.9	9.5	28.9
2001	1.7	1.4	0.3	0.1	0.2	0.2	0.4	0.9	1.2	2.8	9.2	30.2
2000	1.8	1.8	0.5	*	0.1	0.2	0.4	0.8	1.3	3.0	10.0	32.7
1999	1.6	1.5	0.4	0.1	0.1	0.2	0.4	0.7	1.0	2.9	8.5	30.5
Non-Hispanic white, male												
2002	1.7	2.0	*	*	0.2	0.3	0.5	0.8	1.6	3.6	11.9	37.8
2001	1.7	2.0	*	*	0.2	0.2	0.5	1.0	1.5	3.4	11.7	40.0
2000	1.8	2.1	0.6	*	*	0.3	0.6	1.0	1.5	3.4	13.0	40.4
1999	1.6	1.8	0.5	*	*	0.2	0.5	0.8	1.1	3.5	10.6	39.9
Non-Hispanic white, female												
2002	1.6	*	*	*	*	0.2	0.3	0.5	0.9	2.2	7.9	25.0
2001	1.7	*	*	*	*	*	0.3	0.8	1.0	2.3	7.5	26.1
2000	1.8	*	*	*	*	0.2	0.3	0.5	1.1	2.6	8.1	29.6
1999	1.6	*	*	*	*	*	0.3	0.5	0.9	2.4	7.1	26.8
Non-Hispanic black, both sexes												
2002	1.5	*	0.9	*	*	*	0.4	1.1	2.1	6.6	13.2	33.6
2001	1.6	*	*	*	*	*	0.6	1.2	2.6	5.5	14.6	37.1
2000	1.5	4.0	1.0	*	*	0.4	0.4	1.2	2.2	6.0	13.1	35.6
1999	1.5	*	1.1	*	*	0.4	0.4	1.1	2.5	5.5	14.7	33.6
Non-Hispanic black, male												
2002	1.6	*	*	*	*	*	*	1.4	2.7	8.6	19.6	42.7
2001	1.8	*	*	*	*	*	*	1.6	3.6	7.0	23.7	50.8
2000	1.7	*	*	*	*	*	*	1.8	2.9	7.5	20.1	51.7
1999	1.7	*	*	*	*	*	*	1.4	3.3	6.9	20.6	44.0
Non-Hispanic black, female												
2002	1.4	*	*	*	*	*	*	*	1.6	5.1	9.6	30.1
2001	1.3	*	*	*	*	*	*	*	1.7	4.4	9.4	32.1
2000	1.4	*	*	*	*	*	*	*	1.7	5.0	9.2	29.7
1999	1.4	*	*	*	*	*	*	*	1.9	4.6	11.4	29.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)												
All origins 3/, both sexes												
2002	0.5	14.3	0.4	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.4	1.0
2001	0.5	13.7	0.4	0.2	0.2	0.2	0.4	0.3	0.2	0.3	0.4	0.9
2000	0.5	11.8	0.4	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.5	1.2
1999	0.6	10.8	0.5	0.2	0.3	0.2	0.3	0.4	0.3	0.6	1.5	4.7
All origins 3/, male												
2002	0.7	15.7	0.5	0.3	0.4	0.5	0.5	0.5	0.4	0.3	0.5	*
2001	0.7	15.3	0.5	0.3	0.4	0.4	0.6	0.5	0.3	0.4	0.5	*
2000	0.7	13.2	0.5	0.3	0.5	0.5	0.6	0.5	0.5	0.4	0.6	*
1999	0.8	12.6	0.5	0.4	0.4	0.4	0.5	0.6	0.5	0.9	1.8	5.4
All origins 3/, female												
2002	0.3	12.8	0.4	*	*	*	0.1	*	*	*	*	1.1
2001	0.3	12.0	0.3	*	*	*	0.1	*	*	0.2	0.4	0.9
2000	0.3	10.3	0.3	0.1	*	*	0.1	0.1	*	0.2	0.4	1.1
1999	0.4	8.9	0.5	*	*	*	0.1	0.1	0.2	0.4	1.3	4.5
Hispanic, both sexes												
2002	0.3	6.5	*	*	*	*	*	*	*	*	*	*
2001	0.3	7.6	*	*	*	*	*	*	*	*	*	*
2000	0.3	5.6	*	*	*	*	*	*	*	*	*	*
1999	0.4	6.6	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.4	8.0	*	*	*	*	*	*	*	*	*	*
2001	0.4	8.2	*	*	*	*	*	*	*	*	*	*
2000	0.4	5.8	*	*	*	*	*	*	*	*	*	*
1999	0.4	6.5	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.2	4.9	*	*	*	*	*	*	*	*	*	*
2001	0.2	7.0	*	*	*	*	*	*	*	*	*	*
2000	0.2	5.3	*	*	*	*	*	*	*	*	*	*
1999	0.3	6.8	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.5	16.2	0.5	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.4	1.1
2001	0.5	15.2	0.5	0.2	0.2	0.3	0.4	0.3	0.2	0.3	0.4	0.9
2000	0.5	13.3	0.4	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.5	1.2
1999	0.6	11.8	0.5	0.2	0.3	0.3	0.3	0.4	0.3	0.6	1.6	4.7
Non-Hispanic 4/, male												
2002	0.7	17.5	0.6	0.3	0.5	0.5	0.5	0.5	0.4	0.4	0.5	*
2001	0.7	17.0	0.6	0.3	0.4	0.5	0.7	0.6	0.3	0.4	0.5	*
2000	0.7	15.1	0.5	0.4	0.5	0.5	0.7	0.5	0.5	0.4	0.6	*
1999	0.8	14.1	0.5	0.4	0.5	0.5	0.6	0.6	0.5	0.8	1.9	5.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)												
Non-Hispanic 4/, female												
2002	0.3	14.8	0.4	*	*	*	0.1	*	*	*	*	1.1
2001	0.3	13.3	0.4	*	*	*	0.1	*	*	0.2	0.4	0.9
2000	0.3	11.6	*	*	*	*	0.1	0.1	*	*	0.4	1.1
1999	0.4	9.4	0.5	*	*	*	0.1	0.1	0.2	0.4	1.4	4.5
Non-Hispanic white, both sexes												
2002	0.5	12.1	0.5	0.2	0.3	0.4	0.4	0.3	0.3	0.3	0.4	1.0
2001	0.5	11.8	0.4	0.3	0.3	0.3	0.5	0.3	0.2	0.3	0.5	0.9
2000	0.5	10.6	0.5	0.3	0.3	0.3	0.4	0.3	0.4	0.3	0.5	1.2
1999	0.6	9.3	0.5	0.3	0.3	0.3	0.4	0.4	0.3	0.6	1.5	4.9
Non-Hispanic white, male												
2002	0.7	13.7	0.6	0.4	0.6	0.6	0.6	0.6	0.4	0.4	0.5	*
2001	0.7	13.0	0.5	0.4	0.5	0.5	0.8	0.6	0.3	0.5	0.5	*
2000	0.7	12.4	0.6	0.4	0.5	0.6	0.7	0.5	0.6	0.4	0.6	*
1999	0.8	10.8	0.5	0.5	0.5	0.5	0.6	0.7	0.5	0.8	1.8	5.6
Non-Hispanic white, female												
2002	0.3	10.3	0.4	*	*	*	0.1	*	*	*	*	1.1
2001	0.3	10.5	*	*	*	*	0.1	*	*	*	0.4	0.9
2000	0.3	8.7	*	*	*	*	*	*	*	*	0.4	1.1
1999	0.4	7.6	0.5	*	*	*	*	0.1	*	0.5	1.3	4.6
Non-Hispanic black, both sexes												
2002	0.8	34.2	*	*	*	*	*	*	*	*	*	*
2001	0.7	30.4	*	*	*	*	*	*	*	*	*	*
2000	0.7	25.8	*	*	*	*	*	*	*	*	*	*
1999	0.7	23.6	*	*	*	*	*	*	*	*	2.6	*
Non-Hispanic black, male												
2002	0.9	34.5	*	*	*	*	*	*	*	*	*	*
2001	0.9	35.4	*	*	*	*	*	*	*	*	*	*
2000	0.9	27.3	*	*	*	*	*	*	*	*	*	*
1999	1.0	28.4	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.7	33.8	*	*	*	*	*	*	*	*	*	*
2001	0.5	25.2	*	*	*	*	*	*	*	*	*	*
2000	0.5	24.3	*	*	*	*	*	*	*	*	*	*
1999	0.5	18.6	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to electric current (W85-W87)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.3	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	0.1	*	*
All origins 3/, male												
2002	0.3	*	*	*	0.4	0.5	0.5	0.4	0.2	*	*	*
2001	0.3	*	*	*	0.3	0.4	0.5	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.3	0.4	0.4	0.4	0.3	*	*	*
1999	0.3	*	*	*	0.4	0.4	0.5	0.3	0.3	0.2	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	*	0.4	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	0.4	*	*	*	*	*
2000	0.2	*	*	*	0.3	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	0.7	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	0.8	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.2	0.2	0.3	0.2	0.1	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.2	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	*	*	*
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.4	0.4	0.5	0.4	0.2	*	*	*
2001	0.3	*	*	*	0.3	0.5	0.5	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.3	0.4	0.4	0.4	0.3	*	*	*
1999	0.3	*	*	*	0.4	0.4	0.5	0.3	0.3	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to electric current (W85-W87)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.2	0.2	0.3	0.2	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000	0.2	*	*	*	0.2	0.3	0.3	0.2	0.1	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	*	*	*
Non-Hispanic white, male												
2002	0.3	*	*	*	0.4	0.4	0.6	0.4	*	*	*	*
2001	0.3	*	*	*	0.3	0.5	0.5	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.4	0.5	0.5	0.4	0.3	*	*	*
1999	0.3	*	*	*	0.5	0.5	0.6	0.3	0.4	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to radiation (W88-W91)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to radiation (W88-W91)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins 3/, both sexes												
2002	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9
All origins 3/, male												
2002	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
All origins 3/, female												
2002	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5
Hispanic, both sexes												
2002	0.6	*	1.2	0.4	0.4	*	0.5	0.6	*	1.7	*	*
2001	0.5	*	*	0.3	*	0.3	0.5	*	*	*	*	*
2000	0.5	*	0.7	*	0.3	*	0.5	*	*	*	*	*
1999	0.6	*	1.4	0.4	0.4	*	0.4	*	*	*	*	*
Hispanic, male												
2002	0.7	*	*	*	*	*	0.7	*	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.7	*	1.6	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	1.0	1.5	0.7	0.5	0.8	0.9	1.2	1.3	2.1	3.4	4.8
2001	1.3	1.3	1.7	0.7	0.6	0.7	1.0	1.1	1.5	2.3	3.4	5.7
2000	1.3	1.0	2.2	0.7	0.5	0.7	0.9	1.2	1.5	2.2	3.8	6.3
1999	1.3	1.2	2.1	0.7	0.5	0.8	1.0	1.0	1.4	2.3	4.0	6.0
Non-Hispanic 4/, male												
2002	1.5	*	1.7	0.8	0.7	1.0	1.2	1.5	1.7	2.9	4.6	7.0
2001	1.6	1.5	2.0	0.8	0.8	0.9	1.5	1.6	2.0	2.8	4.2	8.3
2000	1.6	*	2.6	0.8	0.7	0.9	1.3	1.7	2.1	2.8	5.0	9.3
1999	1.6	*	2.3	0.8	0.7	1.1	1.4	1.4	1.7	3.0	5.1	9.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic 4/, female												
2002	0.9	*	1.2	0.6	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.8
2001	1.0	*	1.3	0.5	0.4	0.5	0.6	0.7	1.1	1.9	2.9	4.6
2000	1.0	*	1.7	0.7	0.4	0.5	0.6	0.7	1.0	1.7	3.1	5.0
1999	1.0	*	1.8	0.6	0.4	0.5	0.6	0.6	1.1	1.8	3.3	4.7
Non-Hispanic white, both sexes												
2002	1.0	*	1.0	0.4	0.4	0.7	0.8	1.1	1.1	1.8	3.0	3.9
2001	1.1	1.1	1.3	0.5	0.5	0.6	0.9	1.0	1.3	2.0	2.9	4.7
2000	1.1	*	1.5	0.5	0.5	0.6	0.8	1.1	1.3	1.8	3.1	5.1
1999	1.1	0.9	1.5	0.5	0.5	0.8	0.9	0.9	1.2	1.9	3.2	4.7
Non-Hispanic white, male												
2002	1.3	*	1.1	0.6	0.6	0.9	1.1	1.4	1.4	2.4	4.2	5.7
2001	1.4	*	1.7	0.6	0.7	0.8	1.3	1.3	1.7	2.3	3.5	6.9
2000	1.4	*	1.8	0.6	0.6	0.8	1.1	1.6	1.7	2.3	4.1	8.1
1999	1.4	*	1.7	0.6	0.7	1.1	1.3	1.2	1.5	2.3	4.2	7.8
Non-Hispanic white, female												
2002	0.8	*	0.8	0.3	0.3	0.5	0.5	0.7	0.8	1.3	2.2	3.2
2001	0.9	*	0.9	0.4	0.3	0.4	0.6	0.6	1.0	1.7	2.5	3.8
2000	0.9	*	1.1	0.5	0.3	0.4	0.5	0.7	0.8	1.5	2.4	3.9
1999	0.8	*	1.3	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.6	3.5
Non-Hispanic black, both sexes												
2002	2.1	*	3.7	1.5	0.8	1.2	1.1	2.2	3.3	5.0	8.7	16.0
2001	2.3	*	3.4	1.3	0.7	1.2	1.8	2.4	3.6	6.1	9.9	17.8
2000	2.6	*	5.1	1.6	0.7	1.2	1.7	2.4	4.2	6.1	13.0	20.7
1999	2.6	*	4.7	1.6	0.7	1.0	1.8	2.2	3.4	7.0	13.5	22.0
Non-Hispanic black, male												
2002	2.6	*	4.3	1.6	1.0	1.4	1.7	3.1	4.6	7.8	11.1	26.3
2001	3.0	*	3.7	1.6	0.8	1.4	2.8	3.9	5.1	9.0	13.2	26.6
2000	3.2	*	5.8	1.7	0.9	1.6	2.4	3.5	6.3	9.0	17.9	28.2
1999	3.1	*	5.3	1.8	0.9	1.2	2.7	3.3	4.3	10.4	18.4	29.7
Non-Hispanic black, female												
2002	1.6	*	3.0	1.3	*	1.0	*	1.4	2.2	3.1	7.3	12.1
2001	1.8	*	3.1	1.0	*	1.1	1.0	1.1	2.4	4.0	8.1	14.5
2000	2.0	*	4.3	1.4	*	0.9	1.2	1.4	2.5	4.0	10.2	17.9
1999	2.1	*	4.1	1.4	*	0.8	1.1	1.3	2.8	4.7	10.7	19.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental contact with heat and hot substances (X10-X19)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	0.8
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental contact with heat and hot substances (X10-X19)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	0.7
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	0.6
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Victim of lightning (X33)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Victim of lightning (X33)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Earthquake, volcanic eruption, avalanche, landslide and other earth movements (X34-X36)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cataclysmic storm and flood (X37-X38)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cataclysmic storm and flood (X37-X38)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins 3/, both sexes												
2002	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4
All origins 3/, male												
2002	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4
All origins 3/, female												
2002	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0
Hispanic, both sexes												
2002	4.3	*	*	*	2.9	5.0	10.1	10.1	4.0	2.2	*	*
2001	3.6	*	*	*	2.2	4.9	8.7	7.7	3.5	*	*	*
2000	3.7	*	*	*	2.0	5.3	9.6	8.0	3.1	*	*	*
1999	4.4	*	*	*	2.3	6.1	11.9	9.8	3.5	*	*	*
Hispanic, male												
2002	6.6	*	*	*	4.4	7.8	15.4	15.8	6.4	*	*	*
2001	5.6	*	*	*	3.1	7.5	13.7	12.9	5.8	*	*	*
2000	6.0	*	*	*	3.2	8.4	15.6	13.6	4.4	*	*	*
1999	7.3	*	*	*	3.6	9.8	19.4	16.4	6.6	*	*	*
Hispanic, female												
2002	1.7	*	*	*	1.1	1.6	4.3	4.4	*	*	*	*
2001	1.5	*	*	*	1.1	1.9	3.4	2.6	*	*	*	*
2000	1.3	*	*	*	*	1.6	3.1	2.5	*	*	*	*
1999	1.4	*	*	*	0.8	1.7	3.9	3.3	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	6.3	0.6	0.2	0.1	4.4	8.4	13.8	11.8	4.1	2.1	2.3	4.1
2001	5.1	*	0.2	0.1	3.6	6.6	11.4	9.1	3.1	1.6	2.3	3.6
2000	4.6	*	0.2	0.1	3.1	6.0	10.3	8.0	2.8	1.5	2.0	4.4
1999	4.3	*	0.2	0.1	2.5	5.8	9.7	7.5	2.7	1.6	2.2	3.5
Non-Hispanic 4/, male												
2002	8.8	*	*	0.1	6.6	12.3	18.7	15.9	5.2	2.4	2.6	5.3
2001	7.2	*	*	0.1	5.7	9.7	15.9	12.9	3.9	2.0	2.8	3.9
2000	6.6	*	*	0.1	4.9	9.1	14.8	11.7	3.3	1.7	2.1	6.2
1999	6.3	*	*	0.1	3.8	8.8	14.2	11.1	3.6	2.0	2.5	4.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Non-Hispanic 4/, female												
2002	4.0	*	*	*	2.0	4.5	8.9	7.9	3.0	1.8	2.1	3.5
2001	3.1	*	*	*	1.5	3.5	7.1	5.4	2.3	1.2	1.9	3.4
2000	2.7	*	*	*	1.4	3.0	5.9	4.4	2.3	1.3	2.0	3.6
1999	2.4	*	*	*	1.2	2.8	5.4	4.0	1.8	1.3	2.0	3.1
Non-Hispanic white, both sexes												
2002	6.6	*	*	0.1	5.3	9.6	14.5	11.4	3.7	2.0	2.3	4.1
2001	5.2	*	*	0.1	4.3	7.3	11.7	8.3	2.7	1.5	2.1	3.4
2000	4.6	*	*	0.1	3.7	6.5	10.4	7.2	2.5	1.4	2.1	4.1
1999	4.2	*	*	0.1	2.8	6.0	9.6	6.7	2.4	1.5	2.3	3.3
Non-Hispanic white, male												
2002	9.1	*	*	0.2	8.0	13.9	19.6	15.0	4.3	2.2	2.5	5.4
2001	7.3	*	*	*	6.7	10.8	16.3	11.5	3.1	1.9	2.5	3.8
2000	6.6	*	*	*	5.8	9.8	14.9	10.3	2.7	1.5	2.2	5.8
1999	6.1	*	*	*	4.3	9.2	14.0	9.6	2.9	1.8	2.5	4.6
Non-Hispanic white, female												
2002	4.3	*	*	*	2.5	5.2	9.3	7.8	3.1	1.9	2.2	3.5
2001	3.1	*	*	*	1.7	3.8	7.1	5.2	2.4	1.2	1.9	3.2
2000	2.7	*	*	*	1.5	3.2	5.8	4.2	2.3	1.4	2.0	3.4
1999	2.4	*	*	*	1.3	2.8	5.1	3.8	1.9	1.2	2.1	2.8
Non-Hispanic black, both sexes												
2002	6.4	*	*	*	1.5	5.8	14.1	18.3	8.6	2.5	2.8	*
2001	6.0	*	*	*	1.6	5.3	13.4	17.0	6.4	2.4	3.7	6.6
2000	5.7	*	*	*	1.4	5.7	13.2	15.7	5.5	2.5	*	8.3
1999	5.8	*	*	*	1.7	6.3	13.2	15.6	6.0	3.0	*	*
Non-Hispanic black, male												
2002	9.3	*	*	*	2.2	8.7	19.8	27.1	15.2	4.5	*	*
2001	8.6	*	*	*	2.3	7.6	18.2	27.0	11.0	3.7	6.0	*
2000	8.4	*	*	*	1.8	8.5	19.0	25.3	9.5	4.3	*	*
1999	8.8	*	*	*	2.8	9.3	19.5	25.9	11.0	4.4	*	*
Non-Hispanic black, female												
2002	3.8	*	*	*	0.7	3.2	9.1	10.8	3.3	*	*	*
2001	3.6	*	*	*	0.9	3.2	9.2	8.4	2.6	*	*	*
2000	3.2	*	*	*	1.0	3.1	8.0	7.5	2.4	*	*	*
1999	3.1	*	*	*	*	3.5	7.7	6.7	1.9	2.1	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
All origins 3/, both sexes												
2002	5.7	*	*	*	3.9	7.4	12.8	11.1	3.6	1.6	1.7	3.1
2001	4.6	*	*	0.1	3.1	6.0	10.6	8.5	2.7	1.3	1.6	2.5
2000	4.2	*	*	*	2.7	5.6	9.8	7.6	2.3	1.1	1.4	3.3
1999	4.0	*	*	*	2.2	5.6	9.5	7.2	2.3	1.2	1.6	2.8
All origins 3/, male												
2002	7.9	*	*	*	5.9	10.9	17.5	15.0	4.5	1.7	1.8	3.7
2001	6.5	*	*	*	4.8	8.9	14.8	12.2	3.5	1.5	1.7	1.8
2000	6.1	*	*	*	4.1	8.5	14.3	11.2	2.7	1.3	1.3	4.4
1999	5.9	*	*	*	3.3	8.5	14.0	10.9	3.2	1.5	1.5	3.3
All origins 3/, female												
2002	3.5	*	*	*	1.8	3.9	8.1	7.3	2.7	1.4	1.6	2.8
2001	2.7	*	*	*	1.3	3.1	6.4	5.0	2.0	1.1	1.4	2.7
2000	2.3	*	*	*	1.2	2.7	5.5	4.1	1.9	1.0	1.5	2.8
1999	2.1	*	*	*	1.0	2.5	5.1	3.7	1.5	1.0	1.6	2.6
Hispanic, both sexes												
2002	3.9	*	*	*	2.5	4.4	9.5	9.6	3.6	1.8	*	*
2001	3.3	*	*	*	1.7	4.5	8.3	7.4	3.1	*	*	*
2000	3.4	*	*	*	1.6	4.9	9.2	7.6	2.8	*	*	*
1999	4.1	*	*	*	1.9	5.8	11.2	9.2	3.2	*	*	*
Hispanic, male												
2002	6.0	*	*	*	3.9	6.9	14.5	15.0	5.7	*	*	*
2001	5.1	*	*	*	2.5	6.9	12.8	12.3	5.3	*	*	*
2000	5.6	*	*	*	2.6	7.9	14.9	13.0	4.0	*	*	*
1999	6.8	*	*	*	3.1	9.4	18.2	15.5	6.0	*	*	*
Hispanic, female												
2002	1.6	*	*	*	1.0	1.5	4.0	4.3	*	*	*	*
2001	1.3	*	*	*	0.8	1.7	3.4	2.5	*	*	*	*
2000	1.2	*	*	*	*	1.6	3.1	2.3	*	*	*	*
1999	1.3	*	*	*	*	1.7	3.8	3.2	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	5.9	*	*	*	4.1	8.1	13.2	11.2	3.6	1.5	1.6	3.2
2001	4.7	*	*	*	3.3	6.3	10.8	8.5	2.7	1.2	1.6	2.5
2000	4.2	*	*	*	2.9	5.7	9.8	7.4	2.2	1.1	1.5	3.3
1999	3.9	*	*	*	2.2	5.5	9.2	7.0	2.2	1.2	1.6	2.8
Non-Hispanic 4/, male												
2002	8.1	*	*	*	6.3	11.8	17.8	14.9	4.4	1.6	1.7	3.8
2001	6.7	*	*	*	5.2	9.3	15.0	12.0	3.3	1.4	1.7	1.9
2000	6.1	*	*	*	4.4	8.5	14.0	10.8	2.5	1.3	1.3	4.3
1999	5.7	*	*	*	3.4	8.3	13.3	10.3	2.9	1.4	1.5	3.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)												
Non-Hispanic 4/, female												
2002	3.8	*	*	*	1.9	4.4	8.7	7.6	2.8	1.4	1.6	2.9
2001	2.9	*	*	*	1.4	3.4	6.8	5.2	2.1	1.1	1.5	2.8
2000	2.5	*	*	*	1.3	2.9	5.7	4.2	1.9	1.0	1.5	2.9
1999	2.2	*	*	*	1.1	2.6	5.2	3.7	1.6	1.0	1.6	2.6
Non-Hispanic white, both sexes												
2002	6.2	*	*	*	5.0	9.3	13.9	10.8	3.2	1.5	1.6	3.2
2001	4.8	*	*	*	4.0	7.0	11.1	7.8	2.3	1.2	1.6	2.4
2000	4.2	*	*	*	3.4	6.2	9.9	6.7	2.0	1.1	1.5	3.1
1999	3.9	*	*	*	2.5	5.7	9.0	6.2	1.9	1.1	1.6	2.6
Non-Hispanic white, male												
2002	8.5	*	*	*	7.6	13.4	18.7	14.1	3.6	1.5	1.5	3.9
2001	6.8	*	*	*	6.2	10.4	15.4	10.7	2.6	1.3	1.7	1.8
2000	6.1	*	*	*	5.3	9.2	14.1	9.5	1.9	1.1	1.3	4.1
1999	5.6	*	*	*	3.8	8.7	13.1	8.9	2.2	1.2	1.6	3.3
Non-Hispanic white, female												
2002	4.0	*	*	*	2.4	5.0	9.2	7.5	2.9	1.5	1.7	2.9
2001	2.9	*	*	*	1.6	3.7	6.8	5.0	2.1	1.1	1.5	2.7
2000	2.5	*	*	*	1.4	3.1	5.6	3.9	2.0	1.0	1.6	2.7
1999	2.2	*	*	*	1.2	2.7	5.0	3.5	1.6	1.0	1.7	2.4
Non-Hispanic black, both sexes												
2002	6.0	*	*	*	1.4	5.5	13.4	17.5	7.6	1.8	*	*
2001	5.5	*	*	*	1.4	5.0	12.7	16.1	5.9	1.9	*	*
2000	5.3	*	*	*	1.3	5.4	12.6	15.0	4.8	1.9	*	6.4
1999	5.4	*	*	*	1.5	6.0	12.7	14.7	5.1	2.2	*	*
Non-Hispanic black, male												
2002	8.6	*	*	*	2.1	8.2	18.6	25.7	13.5	2.9	*	*
2001	8.0	*	*	*	2.1	7.3	17.2	25.7	10.2	3.0	*	*
2000	7.8	*	*	*	1.6	8.1	18.0	24.3	8.2	3.4	*	*
1999	8.2	*	*	*	2.4	9.0	18.6	24.5	9.5	3.3	*	*
Non-Hispanic black, female												
2002	3.6	*	*	*	*	3.1	8.7	10.4	3.0	*	*	*
2001	3.3	*	*	*	*	3.0	8.8	7.9	2.4	*	*	*
2000	3.0	*	*	*	0.9	2.9	7.9	7.1	2.0	*	*	*
1999	2.8	*	*	*	*	3.4	7.5	6.3	1.6	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)												
All origins 3/, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.1	0.2	*
2001	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	0.1	*	*
2000	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	0.1	*	*
1999	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.2	*
All origins 3/, male												
2002	0.3	*	*	*	0.2	0.3	0.5	0.5	0.5	*	*	*
2001	0.2	*	*	*	0.2	0.2	0.5	0.4	0.3	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.4	0.4	0.3	*	*	*
1999	0.3	*	*	*	0.2	0.2	0.5	0.4	0.3	0.3	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
1999	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	*	0.3	0.4	*	*	*	*	*
2001	0.2	*	*	*	*	*	0.4	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	0.4	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	0.6	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.2	0.2	*
2001	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	*	*	*
2000	0.2	*	*	*	0.1	0.1	0.3	0.3	0.3	0.1	*	*
1999	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.1	0.3	0.5	0.5	0.5	*	*	*
2001	0.2	*	*	*	0.2	0.1	0.5	0.4	0.3	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.4	0.4	0.4	*	*	*
1999	0.3	*	*	*	0.2	0.2	0.5	0.4	0.3	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to other and unspecified solid or liquids substances (X45-X46,X48-X49)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
2001	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	0.2	*	*	*
1999	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.1	0.2	0.3	0.3	0.2	0.1	*	*
2001	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	*	*	*
2000	0.2	*	*	*	0.1	0.1	0.3	0.3	0.3	*	*	*
1999	0.2	*	*	*	0.1	0.1	0.3	0.3	0.2	0.2	0.2	*
Non-Hispanic white, male												
2002	0.3	*	*	*	0.2	0.2	0.4	0.4	0.4	*	*	*
2001	0.2	*	*	*	0.2	0.2	0.5	0.4	0.2	*	*	*
2000	0.2	*	*	*	0.2	0.2	0.4	0.4	0.3	*	*	*
1999	0.2	*	*	*	0.2	0.2	0.5	0.4	0.2	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
2001	0.1	*	*	*	*	*	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	*	*	*	0.1	0.2	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	*	0.5	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to other gases and vapors (X47)												
All origins 3/, both sexes												
2002	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	0.4	0.5	0.6
2001	0.2	*	*	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.6	0.9
2000	0.2	*	*	0.1	0.2	0.2	0.2	0.2	0.3	0.3	0.5	0.8
1999	0.2	*	*	*	0.2	0.2	0.2	0.2	0.3	0.2	0.4	0.5
All origins 3/, male												
2002	0.3	*	*	*	0.2	0.3	0.5	0.4	0.4	0.6	0.7	*
2001	0.3	*	*	*	0.3	0.2	0.4	0.4	0.3	0.3	0.8	1.7
2000	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.3	0.7	*
1999	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.4	0.7	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	*	*	*	0.2	0.2	0.3	0.3	0.3	0.4	0.5	0.7
2001	0.2	*	*	*	0.2	0.2	0.3	0.3	0.2	0.2	0.6	0.9
2000	0.2	*	*	0.1	0.2	0.2	0.2	0.3	0.3	0.3	0.5	0.8
1999	0.2	*	*	*	0.1	0.2	0.3	0.3	0.3	0.3	0.4	0.5
Non-Hispanic 4/, male												
2002	0.3	*	*	*	0.2	0.3	0.5	0.4	0.4	0.6	0.7	*
2001	0.3	*	*	*	0.3	0.2	0.4	0.4	0.3	0.4	0.9	1.8
2000	0.3	*	*	*	0.3	0.4	0.4	0.4	0.5	0.3	0.7	*
1999	0.3	*	*	*	0.2	0.3	0.4	0.4	0.4	0.4	0.7	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning by and exposure to other gases and vapors (X47)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	0.1	0.2	0.2	0.3	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	0.1	*	*	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.2	0.2	0.3	0.3	0.3	0.4	0.5	0.7
2001	0.2	*	*	*	0.2	0.2	0.3	0.2	0.2	0.2	0.4	0.9
2000	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	0.5	0.7
1999	0.2	*	*	*	0.2	0.2	0.2	0.2	0.3	0.3	0.4	*
Non-Hispanic white, male												
2002	0.3	*	*	*	0.3	0.3	0.5	0.4	0.4	0.6	0.7	*
2001	0.3	*	*	*	0.3	0.2	0.4	0.4	0.3	0.4	0.7	1.8
2000	0.3	*	*	*	0.3	0.4	0.4	0.4	0.4	0.3	0.7	*
1999	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.4	0.7	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	0.2	*	0.3	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	*	*	*	*	*	0.4	*	*	*	*	*
2001	0.3	*	*	*	*	*	0.4	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)												
All origins 3/, both sexes												
2002	3.8	0.9	0.6	0.3	0.9	1.2	1.9	2.4	3.5	6.2	18.5	75.8
2001	4.0	1.3	0.8	0.3	1.0	1.2	1.9	2.7	3.5	6.4	20.1	83.4
2000	3.9	1.1	0.7	0.3	1.0	1.3	2.1	2.6	3.6	6.2	18.9	82.6
1999	4.3	1.2	0.8	0.4	1.0	1.4	2.0	2.7	3.7	7.0	22.1	92.8
All origins 3/, male												
2002	4.3	1.2	0.7	0.4	1.5	2.1	3.0	4.0	5.5	8.8	22.8	79.9
2001	4.5	1.6	0.9	0.4	1.6	2.1	3.1	4.4	5.3	8.7	24.8	92.4
2000	4.5	1.2	0.8	0.4	1.7	2.3	3.6	4.5	5.6	8.5	22.6	91.7
1999	4.8	1.4	1.0	0.5	1.7	2.3	3.4	4.4	5.8	9.6	27.3	107.0
All origins 3/, female												
2002	3.2	*	0.5	0.2	0.3	0.3	0.7	0.9	1.6	4.0	15.7	74.0
2001	3.5	1.1	0.7	0.2	0.3	0.3	0.7	1.1	1.8	4.4	17.1	79.6
2000	3.3	*	0.5	0.2	0.3	0.3	0.6	0.8	1.7	4.4	16.5	78.9
1999	3.7	*	0.5	0.3	0.3	0.4	0.7	1.0	1.8	4.8	18.7	87.1
Hispanic, both sexes												
2002	1.5	*	*	*	0.9	1.3	1.8	2.0	2.7	3.7	8.0	35.5
2001	1.6	*	*	*	1.0	1.2	1.9	2.5	2.6	4.9	9.0	47.5
2000	1.6	*	*	*	1.0	1.6	2.1	2.1	2.6	4.3	9.3	45.1
1999	1.5	*	0.8	*	0.9	1.0	1.7	2.0	2.9	5.1	8.7	37.2
Hispanic, male												
2002	2.3	*	*	*	1.6	2.3	3.1	3.5	4.9	5.5	11.1	48.5
2001	2.3	*	*	*	1.7	2.1	3.4	4.3	4.1	6.8	12.5	47.6
2000	2.4	*	*	*	1.8	2.7	3.8	3.4	4.4	6.3	*	54.4
1999	2.2	*	*	*	1.6	1.7	2.8	3.4	4.4	7.5	*	47.4
Hispanic, female												
2002	0.6	*	*	*	*	*	*	*	*	*	5.9	29.0
2001	0.9	*	*	*	*	*	*	*	*	3.5	6.6	47.5
2000	0.8	*	*	*	*	*	*	*	*	*	9.6	40.6
1999	0.8	*	*	*	*	*	*	*	*	*	8.3	32.2
Non-Hispanic 4/, both sexes												
2002	4.1	1.0	0.7	0.4	0.9	1.2	1.9	2.4	3.5	6.3	19.0	77.2
2001	4.3	1.4	0.9	0.3	1.0	1.3	1.9	2.7	3.5	6.5	20.6	84.5
2000	4.2	1.2	0.7	0.3	1.0	1.2	2.1	2.6	3.6	6.3	19.2	83.7
1999	4.6	1.3	0.7	0.4	1.0	1.4	2.1	2.7	3.7	7.1	22.6	94.6
Non-Hispanic 4/, male												
2002	4.6	1.3	0.8	0.5	1.5	2.0	3.0	4.0	5.5	9.0	23.3	81.0
2001	4.8	1.8	1.0	0.4	1.6	2.2	3.1	4.4	5.4	8.8	25.3	94.3
2000	4.8	1.4	0.9	0.4	1.7	2.2	3.5	4.5	5.7	8.6	23.2	93.3
1999	5.2	1.6	1.0	0.5	1.7	2.4	3.4	4.5	5.8	9.7	27.9	108.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)												
Non-Hispanic 4/, female												
2002	3.6	*	0.5	0.2	0.3	0.4	0.8	0.9	1.7	4.1	16.1	75.5
2001	3.8	*	0.8	0.2	0.4	0.4	0.7	1.1	1.8	4.5	17.6	80.5
2000	3.6	*	0.5	0.2	0.3	0.3	0.6	0.8	1.7	4.5	16.7	79.9
1999	4.1	*	0.4	0.3	0.4	0.4	0.8	1.0	1.8	4.9	19.1	88.8
Non-Hispanic white, both sexes												
2002	4.5	0.9	0.6	0.3	0.9	1.3	1.8	2.4	3.4	6.2	19.2	80.6
2001	4.8	1.3	0.9	0.3	1.1	1.2	1.8	2.6	3.5	6.4	21.1	87.7
2000	4.6	1.1	0.7	0.3	1.0	1.3	2.0	2.5	3.5	6.2	19.4	87.5
1999	5.0	1.1	0.7	0.4	1.1	1.4	2.1	2.5	3.5	6.8	22.6	98.1
Non-Hispanic white, male												
2002	4.9	*	0.7	0.4	1.6	2.1	3.0	4.0	5.3	8.7	23.4	83.8
2001	5.2	1.8	1.1	0.4	1.7	2.1	3.0	4.2	5.3	8.7	25.8	97.8
2000	5.0	*	0.8	0.4	1.7	2.2	3.3	4.1	5.2	8.3	22.9	96.9
1999	5.4	*	1.0	0.5	1.7	2.4	3.4	4.1	5.4	9.1	27.8	110.8
Non-Hispanic white, female												
2002	4.1	*	0.4	0.3	0.3	0.4	0.7	0.9	1.7	4.0	16.4	79.3
2001	4.3	*	0.7	0.2	0.4	0.4	0.6	1.1	1.8	4.4	18.1	83.5
2000	4.2	*	0.6	0.2	0.3	0.3	0.6	0.8	1.8	4.4	17.2	83.7
1999	4.6	*	0.5	0.3	0.4	0.4	0.8	0.9	1.6	4.8	19.2	93.0
Non-Hispanic black, both sexes												
2002	2.9	*	1.0	0.5	0.7	1.2	2.5	3.2	4.7	8.2	18.8	47.5
2001	3.0	*	0.9	0.3	0.7	1.3	2.3	3.9	4.6	7.8	18.3	58.4
2000	3.1	*	0.8	0.3	1.0	1.4	2.8	4.0	5.5	8.1	19.4	51.5
1999	3.7	*	0.9	0.3	1.0	1.6	2.5	4.7	6.4	11.3	25.3	67.9
Non-Hispanic black, male												
2002	3.8	*	*	0.7	1.1	2.1	3.7	5.1	8.3	11.9	25.8	58.0
2001	3.8	*	*	*	1.1	2.3	3.7	6.4	7.3	11.4	24.0	63.5
2000	4.6	*	*	*	1.7	2.4	4.9	7.7	10.2	11.5	30.2	69.3
1999	5.1	*	*	*	1.8	2.7	4.4	8.2	10.5	17.7	32.8	108.2
Non-Hispanic black, female												
2002	2.1	*	*	*	*	*	1.4	1.5	1.8	5.6	14.9	43.5
2001	2.2	*	*	*	*	*	1.0	1.8	2.5	5.2	15.2	56.5
2000	1.8	*	*	*	*	*	0.9	*	1.7	5.7	13.4	44.9
1999	2.4	*	*	*	*	*	0.9	1.7	3.2	6.7	21.0	52.9

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins 3/, both sexes												
2002	11.0	...	...	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 5/	10.8	...	...	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3
All origins 3/, male												
2002	17.9	...	...	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 5/	17.6	...	...	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000	17.1	...	...	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999	17.1	...	...	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
All origins 3/, female												
2002	4.3	...	...	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 5/	4.1	...	...	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000	4.0	...	...	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999	4.0	...	...	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1
Hispanic, both sexes												
2002	5.0	...	...	0.5	6.7	6.6	7.0	7.1	7.2	7.5	9.2	12.7
2001 5/	5.0	...	...	0.5	6.2	7.1	7.3	7.0	6.2	7.6	8.8	13.8
2000	5.1	...	...	0.4	6.8	6.5	7.5	7.8	6.0	7.8	11.7	*
1999	5.0	...	...	0.5	6.2	6.7	7.5	7.1	6.8	7.0	13.1	15.4
Hispanic, male												
2002	8.3	...	...	0.6	10.6	10.7	11.2	11.3	13.2	14.7	19.3	35.1
2001 5/	8.3	...	...	0.7	9.5	11.3	12.4	11.4	11.5	15.4	20.1	41.9
2000	8.4	...	...	0.6	10.9	10.6	12.1	12.5	11.1	15.6	26.1	*
1999	8.2	...	...	0.6	9.9	10.7	12.1	11.6	12.2	13.8	28.2	*
Hispanic, female												
2002	1.6	...	...	*	2.1	1.7	2.4	2.9	*	*	*	*
2001 5/	1.5	...	...	*	2.3	2.2	1.8	2.7	*	*	*	*
2000	1.5	...	...	*	2.0	1.7	2.6	3.2	*	*	*	*
1999	1.6	...	...	*	1.8	2.0	2.6	2.7	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	11.8	...	...	0.7	10.5	13.9	16.4	16.5	14.0	13.8	18.1	18.1
2001 5/	11.6	...	...	0.7	10.7	13.9	15.7	15.9	13.6	13.6	17.8	17.5
2000	11.1	...	...	0.8	10.8	13.0	15.4	15.0	12.5	12.7	17.9	19.9
1999	11.2	...	...	0.6	10.8	13.8	15.1	14.4	12.5	13.7	18.3	19.3
Non-Hispanic 4/, male												
2002	19.4	...	...	1.0	17.6	22.7	25.5	25.6	22.8	25.2	39.0	50.9
2001 5/	19.0	...	...	1.1	18.1	23.1	24.6	24.5	21.7	25.0	38.5	51.3
2000	18.3	...	...	1.3	18.3	21.4	24.1	23.2	19.9	23.1	39.1	58.4
1999	18.4	...	...	1.0	18.2	22.7	23.6	22.2	20.2	25.2	39.1	57.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic 4/, female												
2002	4.6	...	...	0.3	3.1	5.2	7.4	7.8	5.9	4.3	4.3	3.9
2001 5/	4.5	...	...	0.3	3.0	4.9	7.0	7.6	6.1	4.1	4.2	3.5
2000	4.3	...	...	0.3	3.1	4.8	6.8	7.0	5.7	4.1	4.1	4.3
1999	4.3	...	...	0.2	3.2	5.0	6.8	6.9	5.4	4.3	4.8	4.0
Non-Hispanic white, both sexes												
2002	13.4	...	...	0.7	11.5	15.4	18.7	18.7	15.6	15.1	19.4	19.2
2001 5/	13.0	...	...	0.7	11.6	15.4	17.7	17.9	15.0	14.7	19.0	18.7
2000	12.5	...	...	0.9	11.6	14.2	17.2	16.7	13.9	13.8	19.0	21.3
1999	12.4	...	...	0.7	11.5	15.0	16.9	16.0	13.7	14.8	19.5	20.8
Non-Hispanic white, male												
2002	21.9	...	...	1.1	19.3	24.6	28.7	28.6	25.0	27.5	41.7	54.3
2001 5/	21.4	...	...	1.1	19.6	25.2	27.4	27.4	23.6	26.9	41.1	55.2
2000	20.4	...	...	1.4	19.5	23.1	26.6	25.5	21.8	24.7	41.7	62.7
1999	20.3	...	...	1.1	19.2	24.3	26.0	24.4	21.9	27.0	41.6	62.2
Non-Hispanic white, female												
2002	5.3	...	...	0.3	3.4	6.1	8.6	9.0	6.6	4.6	4.5	4.1
2001 5/	5.0	...	...	0.3	3.3	5.5	8.1	8.6	6.8	4.4	4.4	3.5
2000	4.9	...	...	0.4	3.3	5.2	7.8	7.9	6.4	4.6	4.2	4.6
1999	4.9	...	...	0.3	3.4	5.5	7.8	7.8	6.0	4.6	5.0	4.2
Non-Hispanic black, both sexes												
2002	5.2	...	...	0.5	6.6	9.6	7.7	5.9	4.9	4.9	5.4	*
2001 5/	5.3	...	...	0.6	7.3	9.2	7.5	6.2	5.1	5.5	5.6	*
2000	5.5	...	...	0.6	8.4	9.0	7.7	6.5	4.2	5.3	5.9	*
1999	5.5	...	...	0.5	8.3	9.5	7.8	5.7	5.0	5.8	5.8	*
Non-Hispanic black, male												
2002	9.3	...	...	0.9	11.6	18.0	13.2	10.2	8.5	9.7	13.5	*
2001 5/	9.3	...	...	0.9	13.4	16.6	12.5	9.8	9.3	10.8	13.2	*
2000	9.6	...	...	1.0	14.6	16.1	13.3	11.1	7.5	11.4	12.3	*
1999	9.9	...	...	0.7	14.8	17.5	13.7	10.1	9.2	11.7	14.1	*
Non-Hispanic black, female												
2002	1.6	...	...	*	1.7	2.1	2.9	2.2	2.0	*	*	*
2001 5/	1.7	...	...	*	1.3	2.4	3.0	3.1	1.8	*	*	*
2000	1.7	...	...	*	2.3	2.6	2.7	2.5	1.6	*	*	*
1999	1.6	...	...	*	2.0	2.4	2.6	2.0	1.6	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)												
All origins 3/, both sexes												
2002	1.3	...	...	*	0.6	1.4	2.6	2.8	1.7	1.0	1.0	0.8
2001	1.3	...	...	*	0.6	1.3	2.4	2.7	1.5	1.0	0.8	0.9
2000	1.2	...	...	*	0.5	1.2	2.2	2.4	1.5	0.8	0.9	1.0
1999	1.1	...	...	*	0.5	1.2	2.2	2.3	1.4	1.0	1.1	0.8
All origins 3/, male												
2002	1.3	...	...	*	0.6	1.5	2.5	2.7	1.5	0.9	1.2	*
2001	1.2	...	...	*	0.5	1.3	2.6	2.4	1.2	1.0	0.7	*
2000	1.1	...	...	*	0.5	1.2	2.2	2.3	1.2	0.7	1.0	*
1999	1.1	...	...	*	0.5	1.1	2.3	2.2	1.2	1.0	0.9	*
All origins 3/, female												
2002	1.4	...	...	*	0.6	1.2	2.6	3.0	2.0	1.1	0.9	0.7
2001	1.3	...	...	*	0.6	1.2	2.3	2.9	1.7	0.9	0.8	0.6
2000	1.2	...	...	*	0.5	1.1	2.3	2.5	1.7	0.9	0.7	0.9
1999	1.2	...	...	*	0.5	1.2	2.2	2.4	1.6	0.9	1.2	*
Hispanic, both sexes												
2002	0.4	...	...	*	*	0.5	0.9	1.1	*	*	*	*
2001	0.4	...	...	*	*	0.4	0.8	0.9	*	*	*	*
2000	0.3	...	...	*	*	0.4	0.9	0.8	*	*	*	*
1999	0.4	...	...	*	*	0.5	0.9	1.1	*	*	*	*
Hispanic, male												
2002	0.4	...	...	*	*	0.6	1.1	*	*	*	*	*
2001	0.3	...	...	*	*	*	1.0	*	*	*	*	*
2000	0.4	...	...	*	*	*	0.9	*	*	*	*	*
1999	0.4	...	...	*	*	*	1.1	*	*	*	*	*
Hispanic, female												
2002	0.4	...	...	*	*	*	0.8	1.1	*	*	*	*
2001	0.4	...	...	*	*	*	*	*	*	*	*	*
2000	0.3	...	...	*	*	*	0.8	*	*	*	*	*
1999	0.4	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.5	...	...	*	0.6	1.5	2.8	3.0	1.8	1.0	1.0	0.9
2001	1.4	...	...	*	0.6	1.4	2.7	2.8	1.5	1.0	0.8	0.9
2000	1.3	...	...	*	0.6	1.3	2.4	2.5	1.5	0.8	0.9	1.0
1999	1.2	...	...	*	0.5	1.3	2.4	2.4	1.4	1.0	1.1	0.8
Non-Hispanic 4/, male												
2002	1.4	...	...	*	0.7	1.7	2.7	2.8	1.6	0.9	1.2	*
2001	1.4	...	...	*	0.6	1.5	2.9	2.5	1.2	1.0	0.7	*
2000	1.2	...	...	*	0.6	1.4	2.3	2.4	1.3	0.7	1.0	*
1999	1.2	...	...	*	0.5	1.3	2.4	2.3	1.3	1.0	1.0	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)												
Non-Hispanic 4/, female												
2002	1.5	...	...	*	0.6	1.4	2.8	3.2	2.1	1.1	1.0	0.7
2001	1.4	...	...	*	0.7	1.3	2.5	3.1	1.8	0.9	0.8	0.7
2000	1.3	...	...	*	0.6	1.3	2.5	2.6	1.7	0.9	0.8	0.9
1999	1.3	...	...	*	0.6	1.3	2.3	2.5	1.6	0.9	1.2	*
Non-Hispanic white, both sexes												
2002	1.7	...	...	*	0.8	1.8	3.2	3.5	2.1	1.1	1.1	1.0
2001	1.6	...	...	*	0.7	1.7	3.1	3.2	1.7	1.1	0.9	1.0
2000	1.5	...	...	*	0.7	1.5	2.7	2.8	1.7	0.9	0.9	1.1
1999	1.4	...	...	*	0.6	1.4	2.7	2.7	1.6	1.1	1.2	0.9
Non-Hispanic white, male												
2002	1.7	...	...	*	0.8	1.9	3.1	3.3	1.8	1.0	1.2	*
2001	1.6	...	...	*	0.7	1.9	3.3	2.9	1.4	1.1	0.8	*
2000	1.4	...	...	*	0.7	1.6	2.7	2.6	1.4	0.7	1.1	*
1999	1.4	...	...	*	0.6	1.4	2.7	2.6	1.4	1.1	1.0	*
Non-Hispanic white, female												
2002	1.8	...	...	*	0.8	1.6	3.3	3.7	2.3	1.2	1.0	0.8
2001	1.6	...	...	*	0.8	1.5	2.9	3.6	2.1	1.0	0.9	*
2000	1.5	...	...	*	0.7	1.5	2.8	3.0	2.0	1.1	0.8	1.0
1999	1.5	...	...	*	0.6	1.5	2.7	2.9	1.8	1.1	1.3	*
Non-Hispanic black, both sexes												
2002	0.5	...	...	*	*	0.8	1.0	0.8	*	*	*	*
2001	0.4	...	...	*	*	0.5	1.0	0.9	*	*	*	*
2000	0.4	...	...	*	*	0.6	1.0	1.1	*	*	*	*
1999	0.4	...	...	*	*	0.7	0.9	0.8	*	*	*	*
Non-Hispanic black, male												
2002	0.5	...	...	*	*	1.1	1.0	*	*	*	*	*
2001	0.4	...	...	*	*	*	1.0	*	*	*	*	*
2000	0.5	...	...	*	*	*	0.9	1.3	*	*	*	*
1999	0.4	...	...	*	*	0.9	0.9	*	*	*	*	*
Non-Hispanic black, female												
2002	0.4	...	...	*	*	*	0.9	0.9	*	*	*	*
2001	0.4	...	...	*	*	*	1.0	1.0	*	*	*	*
2000	0.4	...	...	*	*	*	1.0	0.9	*	*	*	*
1999	0.4	...	...	*	*	*	1.0	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)												
All origins 3/, both sexes												
2002	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
2001	0.1	...	...	*	*	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	...	...	*	*	0.1	0.1	0.1	0.1	*	*	*
All origins 3/, male												
2002	0.1	...	...	*	*	0.1	0.2	0.2	*	*	*	*
2001	0.1	...	...	*	*	0.1	0.2	0.1	*	*	*	*
2000	0.1	...	...	*	*	0.1	0.2	0.2	*	*	*	*
1999	0.1	...	...	*	*	0.1	0.2	0.2	0.2	*	*	*
All origins 3/, female												
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
2001	0.1	...	...	*	*	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	...	...	*	*	0.1	0.1	0.1	0.1	*	*	*
Non-Hispanic 4/, male												
2002	0.1	...	...	*	*	0.1	0.2	0.2	*	*	*	*
2001	0.1	...	...	*	*	0.1	0.2	0.1	*	*	*	*
2000	0.1	...	...	*	*	*	0.2	0.2	*	*	*	*
1999	0.1	...	...	*	*	0.1	0.2	0.2	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)												
Non-Hispanic 4/, female												
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
2001	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
2000	0.1	...	...	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	...	...	*	*	0.1	0.1	0.1	0.1	*	*	*
Non-Hispanic white, male												
2002	0.1	...	...	*	*	*	0.2	0.2	*	*	*	*
2001	0.1	...	...	*	*	*	0.2	0.1	*	*	*	*
2000	0.1	...	...	*	*	*	0.2	0.2	*	*	*	*
1999	0.1	...	...	*	*	0.2	0.2	0.2	*	*	*	*
Non-Hispanic white, female												
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)												
All origins 3/, both sexes												
2002	0.5	...	...	*	0.2	0.5	0.8	1.0	0.6	0.4	0.7	0.8
2001	0.5	...	...	*	0.2	0.6	0.9	0.9	0.7	0.4	0.6	0.8
2000	0.5	...	...	*	0.2	0.6	0.9	0.9	0.7	0.4	0.6	0.8
1999	0.6	...	...	*	0.3	0.7	0.9	0.9	0.6	0.4	0.8	1.2
All origins 3/, male												
2002	0.8	...	...	*	0.3	0.9	1.4	1.5	1.0	0.6	1.2	2.2
2001	0.8	...	...	*	0.4	0.9	1.3	1.4	1.1	0.7	1.2	1.9
2000	0.8	...	...	*	0.4	1.0	1.4	1.5	1.0	0.6	1.2	2.3
1999	0.9	...	...	*	0.6	1.1	1.5	1.4	0.9	0.6	1.5	3.9
All origins 3/, female												
2002	0.2	...	...	*	*	0.2	0.3	0.4	0.2	0.2	0.3	*
2001	0.2	...	...	*	*	0.2	0.4	0.4	0.4	0.2	*	*
2000	0.2	...	...	*	*	0.2	0.4	0.4	0.3	0.2	*	*
1999	0.2	...	...	*	*	0.3	0.4	0.5	0.4	*	0.4	*
Hispanic, both sexes												
2002	0.1	...	...	*	*	*	*	*	*	*	*	*
2001	0.1	...	...	*	*	*	*	*	*	*	*	*
2000	0.1	...	...	*	*	*	*	*	*	*	*	*
1999	0.1	...	...	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	...	...	*	*	*	*	*	*	*	*	*
2001	0.2	...	...	*	*	*	*	*	*	*	*	*
2000	0.2	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.6	...	...	*	0.2	0.6	0.9	1.0	0.6	0.4	0.7	0.8
2001	0.6	...	...	*	0.3	0.6	0.9	1.0	0.8	0.5	0.7	0.8
2000	0.6	...	...	*	0.3	0.7	1.0	1.0	0.7	0.4	0.6	0.8
1999	0.6	...	...	*	0.4	0.8	1.0	1.0	0.7	0.4	0.8	1.2
Non-Hispanic 4/, male												
2002	0.9	...	...	*	0.4	1.0	1.5	1.6	1.0	0.7	1.3	2.3
2001	0.9	...	...	*	0.5	1.1	1.5	1.5	1.1	0.8	1.3	2.0
2000	0.9	...	...	*	0.5	1.1	1.6	1.5	1.1	0.6	1.3	2.4
1999	1.0	...	...	*	0.7	1.3	1.6	1.5	1.0	0.6	1.5	4.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)												
Non-Hispanic 4/, female												
2002	0.2	...	...	*	*	0.3	0.4	0.4	0.3	0.2	0.3	*
2001	0.3	...	...	*	*	0.2	0.4	0.5	0.5	0.2	*	*
2000	0.2	...	...	*	*	0.3	0.4	0.5	0.3	0.2	*	*
1999	0.3	...	...	*	*	0.3	0.4	0.5	0.4	*	0.4	*
Non-Hispanic white, both sexes												
2002	0.7	...	...	*	0.3	0.8	1.1	1.2	0.7	0.5	0.7	0.9
2001	0.7	...	...	*	0.3	0.8	1.1	1.1	0.9	0.5	0.7	0.9
2000	0.7	...	...	*	0.3	0.8	1.2	1.2	0.8	0.4	0.7	0.9
1999	0.7	...	...	*	0.5	1.0	1.2	1.2	0.8	0.4	0.9	1.4
Non-Hispanic white, male												
2002	1.1	...	...	*	0.5	1.3	1.8	1.9	1.2	0.8	1.4	2.4
2001	1.1	...	...	*	0.6	1.3	1.7	1.8	1.2	0.8	1.4	2.2
2000	1.1	...	...	*	0.6	1.3	1.8	1.8	1.3	0.6	1.3	2.6
1999	1.2	...	...	*	0.8	1.5	1.9	1.7	1.1	0.7	1.6	4.5
Non-Hispanic white, female												
2002	0.3	...	...	*	*	0.3	0.5	0.5	0.3	0.3	0.3	*
2001	0.3	...	...	*	*	0.3	0.5	0.5	0.5	0.3	*	*
2000	0.3	...	...	*	*	0.4	0.5	0.6	0.4	0.3	*	*
1999	0.3	...	...	*	*	0.4	0.5	0.6	0.5	*	0.4	*
Non-Hispanic black, both sexes												
2002	0.1	...	...	*	*	*	*	*	*	*	*	*
2001	0.1	...	...	*	*	*	*	*	*	*	*	*
2000	0.1	...	...	*	*	*	*	*	*	*	*	*
1999	0.1	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	...	...	*	*	*	*	*	*	*	*	*
2001	0.2	...	...	*	*	*	*	*	*	*	*	*
2000	0.2	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)												
All origins 3/, both sexes												
2002	2.2	...	...	0.4	3.2	3.5	3.4	2.7	1.7	1.4	1.6	2.3
2001	2.2	...	...	0.4	3.1	3.5	3.4	2.4	1.5	1.3	1.7	2.1
2000	2.0	...	...	0.4	2.9	3.1	3.1	2.2	1.4	1.1	1.8	2.9
1999	1.9	...	...	0.3	2.5	3.3	3.0	2.0	1.4	1.4	1.9	2.8
All origins 3/, male												
2002	3.8	...	...	0.6	5.2	6.0	5.7	4.4	2.9	2.3	2.7	5.2
2001	3.7	...	...	0.6	5.1	5.9	5.8	4.2	2.5	2.3	3.1	5.2
2000	3.4	...	...	0.7	4.8	5.4	5.3	3.8	2.2	1.7	3.3	7.2
1999	3.3	...	...	0.5	4.1	5.6	5.2	3.3	2.3	2.2	3.1	6.6
All origins 3/, female												
2002	0.8	...	...	0.2	1.0	1.0	1.1	1.0	0.6	0.6	0.8	1.0
2001	0.7	...	...	0.2	0.9	0.9	1.0	0.7	0.6	0.5	0.8	0.8
2000	0.7	...	...	0.2	0.9	0.8	0.8	0.7	0.7	0.7	0.9	1.2
1999	0.7	...	...	0.1	0.9	0.9	0.8	0.7	0.5	0.7	1.0	1.3
Hispanic, both sexes												
2002	1.8	...	...	0.3	2.7	2.4	2.5	2.1	1.9	1.8	*	*
2001	1.8	...	...	0.4	2.5	2.6	2.9	2.0	1.4	2.2	*	*
2000	1.7	...	...	0.3	2.6	2.3	2.7	2.1	1.6	*	*	*
1999	1.6	...	...	*	1.9	2.4	2.5	2.0	2.3	*	*	*
Hispanic, male												
2002	2.9	...	...	*	3.9	4.1	4.1	3.5	3.7	*	*	*
2001	3.1	...	...	0.6	3.7	4.2	5.0	3.8	2.5	4.2	*	*
2000	2.9	...	...	*	4.0	3.7	4.7	3.6	2.9	*	*	*
1999	2.8	...	...	*	3.2	4.1	4.1	3.5	4.4	*	*	*
Hispanic, female												
2002	0.5	...	...	*	1.2	*	0.7	*	*	*	*	*
2001	0.5	...	...	*	1.0	0.8	*	*	*	*	*	*
2000	0.5	...	...	*	1.0	*	*	*	*	*	*	*
1999	0.4	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	2.3	...	...	0.4	3.2	3.8	3.5	2.7	1.6	1.3	1.6	2.2
2001	2.2	...	...	0.4	3.2	3.6	3.5	2.5	1.6	1.2	1.6	2.0
2000	2.0	...	...	0.4	2.9	3.3	3.1	2.2	1.4	1.1	1.7	2.9
1999	2.0	...	...	0.3	2.6	3.4	3.1	2.0	1.3	1.3	1.8	2.7
Non-Hispanic 4/, male												
2002	3.9	...	...	0.6	5.5	6.5	5.9	4.5	2.8	2.2	2.6	5.0
2001	3.8	...	...	0.7	5.4	6.3	5.9	4.2	2.5	2.2	2.9	4.6
2000	3.5	...	...	0.7	4.9	5.7	5.3	3.8	2.1	1.6	3.1	7.0
1999	3.3	...	...	0.5	4.2	5.9	5.3	3.3	2.2	2.0	3.0	6.3

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)												
Non-Hispanic 4/, female												
2002	0.8	...	...	0.2	0.9	1.1	1.1	1.0	0.6	0.6	0.9	1.0
2001	0.7	...	...	0.1	0.9	1.0	1.1	0.8	0.6	0.5	0.8	0.9
2000	0.7	...	...	0.2	0.9	0.9	0.9	0.7	0.7	0.7	0.9	1.2
1999	0.7	...	...	0.1	0.9	1.0	0.8	0.7	0.5	0.7	1.1	1.3
Non-Hispanic white, both sexes												
2002	2.5	...	...	0.4	3.6	4.1	3.9	3.0	1.7	1.3	1.6	2.2
2001	2.4	...	...	0.4	3.6	3.9	3.8	2.7	1.6	1.2	1.7	1.9
2000	2.2	...	...	0.4	3.1	3.5	3.3	2.4	1.5	1.1	1.7	2.9
1999	2.1	...	...	0.3	2.7	3.6	3.3	2.1	1.3	1.3	1.8	2.8
Non-Hispanic white, male												
2002	4.2	...	...	0.6	6.2	6.9	6.5	4.8	2.9	2.2	2.7	4.9
2001	4.1	...	...	0.6	6.1	6.6	6.4	4.6	2.6	2.1	3.0	4.6
2000	3.7	...	...	0.7	5.3	6.1	5.7	4.0	2.3	1.6	3.0	7.0
1999	3.5	...	...	0.5	4.4	6.2	5.7	3.4	2.2	2.1	3.0	6.5
Non-Hispanic white, female												
2002	0.8	...	...	0.2	1.0	1.2	1.3	1.2	0.6	0.5	0.8	1.0
2001	0.7	...	...	*	0.9	1.0	1.2	0.8	0.7	0.5	0.8	0.8
2000	0.7	...	...	0.2	0.9	0.8	0.9	0.8	0.7	0.7	0.8	1.3
1999	0.7	...	...	*	0.9	1.0	0.9	0.8	0.5	0.7	1.0	1.4
Non-Hispanic black, both sexes												
2002	1.1	...	...	0.4	1.5	2.4	1.8	1.0	*	*	*	*
2001	1.1	...	...	0.4	1.5	2.5	1.6	1.0	*	*	*	*
2000	1.2	...	...	0.5	1.8	2.2	1.8	1.1	*	*	*	*
1999	1.1	...	...	0.3	1.5	2.2	1.7	0.9	*	*	*	*
Non-Hispanic black, male												
2002	2.1	...	...	0.6	2.3	4.3	3.5	2.0	*	*	*	*
2001	2.1	...	...	0.6	2.5	4.7	2.9	1.9	*	*	*	*
2000	2.2	...	...	0.9	3.1	3.9	3.5	2.2	*	*	*	*
1999	2.0	...	...	*	2.6	4.2	3.3	1.8	*	*	*	*
Non-Hispanic black, female												
2002	0.3	...	...	*	*	*	*	*	*	*	*	*
2001	0.3	...	...	*	*	*	*	*	*	*	*	*
2000	0.3	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All origins 3/, both sexes												
2002	5.9	...	...	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001	5.9	...	...	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000	5.9	...	...	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999	5.9	...	...	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6
All origins 3/, male												
2002	10.6	...	...	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001	10.6	...	...	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000	10.5	...	...	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999	10.6	...	...	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6
All origins 3/, female												
2002	1.4	...	...	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001	1.5	...	...	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000	1.5	...	...	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999	1.5	...	...	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9
Hispanic, both sexes												
2002	2.2	...	...	*	3.3	2.7	2.5	2.7	3.4	4.4	4.7	*
2001	2.2	...	...	*	2.9	3.3	2.5	3.0	3.2	4.0	4.7	*
2000	2.3	...	...	*	3.4	2.8	2.8	4.0	2.7	4.6	5.3	*
1999	2.3	...	...	*	3.5	2.9	3.0	3.2	2.9	3.6	8.1	*
Hispanic, male												
2002	3.8	...	...	*	5.7	4.6	4.2	4.8	6.6	9.4	11.5	*
2001	3.8	...	...	*	4.8	5.5	4.5	5.3	6.6	8.8	11.6	*
2000	4.1	...	...	*	5.8	4.7	4.7	6.8	5.7	9.7	12.8	*
1999	4.0	...	...	*	5.8	4.8	5.0	5.5	5.9	7.7	19.8	*
Hispanic, female												
2002	0.4	...	...	*	*	*	*	*	*	*	*	*
2001	0.4	...	...	*	0.7	0.7	*	*	*	*	*	*
2000	0.4	...	...	*	*	*	*	*	*	*	*	*
1999	0.5	...	...	*	0.8	*	0.9	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	6.5	...	...	0.2	5.5	6.7	7.7	8.3	8.8	10.0	13.6	12.1
2001	6.5	...	...	0.2	5.8	7.1	7.3	8.1	8.6	9.9	13.5	12.4
2000	6.4	...	...	0.3	6.2	6.8	7.6	7.9	7.9	9.5	13.4	13.4
1999	6.4	...	...	0.3	6.5	7.2	7.3	7.7	8.2	10.1	13.3	12.7
Non-Hispanic 4/, male												
2002	11.7	...	...	0.4	9.8	11.6	13.1	14.4	15.7	20.1	31.9	38.4
2001	11.6	...	...	0.4	10.4	12.5	12.3	14.0	15.1	19.8	31.5	39.7
2000	11.4	...	...	0.5	11.1	11.7	12.7	13.5	14.0	18.9	31.6	43.8
1999	11.5	...	...	0.4	11.5	12.5	12.3	13.1	14.5	20.1	31.5	42.2

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic 4/, female												
2002	1.6	...	...	*	1.1	1.9	2.4	2.4	2.3	1.6	1.4	0.8
2001	1.6	...	...	0.1	1.1	1.8	2.4	2.5	2.5	1.8	1.6	1.0
2000	1.6	...	...	0.1	1.3	1.9	2.6	2.4	2.3	1.8	1.5	1.1
1999	1.6	...	...	*	1.3	1.9	2.4	2.4	2.3	1.8	1.5	0.9
Non-Hispanic white, both sexes												
2002	7.5	...	...	0.2	5.9	7.5	8.8	9.5	9.9	11.2	14.7	13.1
2001	7.4	...	...	0.3	6.2	8.0	8.3	9.3	9.6	10.9	14.5	13.4
2000	7.3	...	...	0.4	6.7	7.4	8.6	8.9	8.7	10.4	14.5	14.5
1999	7.3	...	...	0.3	6.8	7.9	8.3	8.6	9.0	11.0	14.3	13.7
Non-Hispanic white, male												
2002	13.4	...	...	0.4	10.5	12.6	14.9	16.3	17.4	22.2	34.3	41.3
2001	13.2	...	...	0.4	11.0	13.8	13.9	15.9	16.7	21.6	33.8	43.0
2000	12.9	...	...	0.6	11.8	12.7	14.3	15.1	15.3	20.4	34.2	47.6
1999	12.9	...	...	0.5	12.0	13.4	13.7	14.6	15.9	21.8	33.8	45.7
Non-Hispanic white, female												
2002	1.8	...	...	*	1.2	2.3	2.8	2.8	2.7	1.8	1.6	0.9
2001	1.9	...	...	*	1.2	2.1	2.7	2.9	2.9	1.9	1.7	1.1
2000	1.9	...	...	*	1.4	2.1	3.0	2.8	2.6	2.0	1.5	1.2
1999	1.9	...	...	*	1.4	2.3	2.9	2.7	2.6	2.0	1.6	1.0
Non-Hispanic black, both sexes												
2002	2.9	...	...	*	4.2	5.3	3.6	2.9	2.8	3.3	4.3	*
2001	3.0	...	...	*	5.1	5.0	3.5	2.8	2.7	3.8	5.1	*
2000	3.1	...	...	*	5.4	5.2	3.5	3.2	3.0	3.8	3.6	*
1999	3.2	...	...	*	5.8	5.1	3.6	3.0	3.5	4.0	4.8	*
Non-Hispanic black, male												
2002	5.5	...	...	*	7.7	10.5	6.5	5.5	5.5	7.0	11.4	*
2001	5.6	...	...	*	9.6	9.5	6.4	4.9	5.5	7.9	12.0	*
2000	5.7	...	...	*	9.9	9.6	6.4	5.6	5.7	9.0	8.3	*
1999	6.1	...	...	*	10.6	10.0	6.9	5.6	7.1	8.9	12.2	*
Non-Hispanic black, female												
2002	0.5	...	...	*	0.7	*	1.0	*	*	*	*	*
2001	0.6	...	...	*	*	0.9	1.0	1.0	*	*	*	*
2000	0.7	...	...	*	1.0	1.1	0.8	1.1	*	*	*	*
1999	0.6	...	...	*	1.0	0.8	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by jumping from a high place (X80)												
All origins 3/, both sexes												
2002	0.3	...	...	*	0.3	0.4	0.3	0.3	0.3	0.2	0.3	0.6
2001	0.2	...	...	*	0.2	0.3	0.3	0.3	0.3	0.3	0.4	*
2000	0.2	...	...	*	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.6
1999	0.2	...	...	*	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.5
All origins 3/, male												
2002	0.4	...	...	*	0.4	0.6	0.5	0.4	0.4	0.3	0.5	*
2001	0.3	...	...	*	0.3	0.4	0.4	0.4	0.4	0.3	0.6	*
2000	0.3	...	...	*	0.4	0.4	0.4	0.5	0.2	0.3	0.6	*
1999	0.4	...	...	*	0.4	0.5	0.6	0.5	0.2	0.4	0.7	*
All origins 3/, female												
2002	0.2	...	...	*	0.1	0.2	0.2	0.2	0.2	*	*	*
2001	0.1	...	...	*	0.1	0.1	0.2	0.2	0.2	0.2	*	*
2000	0.1	...	...	*	0.1	0.1	0.1	0.1	0.2	*	*	*
1999	0.1	...	...	*	0.1	0.1	0.2	0.2	0.2	*	*	*
Hispanic, both sexes												
2002	0.2	...	...	*	*	*	*	*	*	*	*	*
2001	0.2	...	...	*	*	*	*	*	*	*	*	*
2000	0.2	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	...	...	*	*	*	*	*	*	*	*	*
2001	0.3	...	...	*	*	*	*	*	*	*	*	*
2000	0.2	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.3	...	...	*	0.3	0.4	0.4	0.3	0.3	0.2	0.3	0.6
2001	0.2	...	...	*	0.2	0.3	0.3	0.3	0.3	0.3	0.4	*
2000	0.2	...	...	*	0.3	0.3	0.3	0.3	0.2	0.2	0.3	0.6
1999	0.3	...	...	*	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.5
Non-Hispanic 4/, male												
2002	0.4	...	...	*	0.5	0.6	0.5	0.4	0.4	0.3	0.5	*
2001	0.3	...	...	*	0.3	0.4	0.4	0.4	0.4	0.3	0.6	*
2000	0.3	...	...	*	0.4	0.4	0.4	0.5	0.2	*	0.5	*
1999	0.4	...	...	*	0.5	0.5	0.6	0.5	0.2	0.5	0.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by jumping from a high place (X80)												
Non-Hispanic 4/, female												
2002	0.2	...	...	*	0.1	0.2	0.2	0.2	*	*	*	*
2001	0.2	...	...	*	*	0.2	0.2	0.3	*	0.2	*	*
2000	0.1	...	...	*	0.1	0.1	0.1	0.1	0.2	*	*	*
1999	0.1	...	...	*	*	0.1	0.2	0.2	0.2	*	*	*
Non-Hispanic white, both sexes												
2002	0.3	...	...	*	0.3	0.4	0.4	0.3	0.3	0.2	0.3	0.6
2001	0.2	...	...	*	0.2	0.3	0.3	0.3	0.3	0.3	0.4	*
2000	0.2	...	...	*	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.6
1999	0.3	...	...	*	0.3	0.3	0.4	0.3	0.2	0.3	0.4	*
Non-Hispanic white, male												
2002	0.4	...	...	*	0.5	0.6	0.5	0.4	0.4	*	0.5	*
2001	0.3	...	...	*	0.4	0.4	0.4	0.4	0.4	0.3	0.6	*
2000	0.3	...	...	*	0.5	0.3	0.4	0.5	0.2	*	0.5	*
1999	0.4	...	...	*	0.5	0.5	0.6	0.5	*	0.5	0.6	*
Non-Hispanic white, female												
2002	0.2	...	...	*	*	0.2	0.2	0.3	*	*	*	*
2001	0.2	...	...	*	*	0.2	0.2	0.2	*	0.2	*	*
2000	0.1	...	...	*	*	*	*	0.2	*	*	*	*
1999	0.1	...	...	*	*	*	0.2	0.2	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	...	...	*	*	*	*	*	*	*	*	*
2001	0.2	...	...	*	*	*	*	*	*	*	*	*
2000	0.2	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	0.4	*	*	*	*	*
Non-Hispanic black, male												
2002	0.3	...	...	*	*	*	*	*	*	*	*	*
2001	0.3	...	...	*	*	*	*	*	*	*	*	*
2000	0.3	...	...	*	*	*	*	*	*	*	*	*
1999	0.4	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)												
All origins 3/, both sexes												
2002	0.6	...	...	*	0.5	0.7	1.0	1.0	0.8	0.7	0.8	1.4
2001 5/	0.6	...	...	*	0.5	0.7	0.9	1.0	0.8	0.6	0.8	1.0
2000	0.6	...	...	*	0.5	0.6	0.9	0.8	0.8	0.6	0.9	1.2
1999	0.6	...	...	*	0.5	0.7	0.9	0.9	0.6	0.7	0.8	1.3
All origins 3/, male												
2002	1.0	...	...	*	0.8	1.1	1.5	1.6	1.2	1.0	1.4	3.1
2001 5/	0.9	...	...	*	0.7	1.1	1.3	1.5	1.2	0.9	1.4	2.5
2000	0.9	...	...	*	0.7	0.9	1.4	1.2	1.2	1.0	1.6	2.4
1999	0.8	...	...	*	0.7	1.1	1.2	1.3	0.9	1.0	1.4	2.5
All origins 3/, female												
2002	0.3	...	...	*	0.2	0.3	0.4	0.4	0.5	0.4	0.5	0.7
2001 5/	0.3	...	...	*	0.2	0.3	0.4	0.5	0.4	0.4	0.4	*
2000	0.3	...	...	*	0.2	0.2	0.4	0.5	0.4	0.4	0.5	0.7
1999	0.3	...	...	*	0.2	0.2	0.5	0.5	0.3	0.5	0.4	0.8
Hispanic, both sexes												
2002	0.4	...	...	*	0.4	0.6	0.7	0.8	*	*	*	*
2001 5/	0.4	...	...	*	0.3	0.5	0.6	0.6	*	*	*	*
2000	0.4	...	...	*	0.4	0.5	0.6	*	*	*	*	*
1999	0.3	...	...	*	*	0.5	0.5	*	*	*	*	*
Hispanic, male												
2002	0.7	...	...	*	0.6	0.9	1.1	1.2	*	*	*	*
2001 5/	0.6	...	...	*	*	0.8	0.9	*	*	*	*	*
2000	0.6	...	...	*	0.6	0.8	1.0	*	*	*	*	*
1999	0.5	...	...	*	*	0.8	0.9	*	*	*	*	*
Hispanic, female												
2002	0.1	...	...	*	*	*	*	*	*	*	*	*
2001 5/	0.2	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	0.2	...	...	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	...	...	*	0.5	0.7	1.0	1.0	0.8	0.7	0.8	1.3
2001 5/	0.6	...	...	*	0.5	0.7	0.9	1.0	0.8	0.6	0.9	1.0
2000	0.6	...	...	*	0.5	0.6	0.9	0.9	0.8	0.6	0.9	1.1
1999	0.6	...	...	*	0.5	0.7	0.9	0.9	0.6	0.7	0.8	1.3
Non-Hispanic 4/, male												
2002	1.0	...	...	*	0.8	1.1	1.6	1.6	1.2	1.0	1.4	2.8
2001 5/	1.0	...	...	*	0.8	1.1	1.4	1.6	1.1	0.9	1.5	2.3
2000	0.9	...	...	*	0.7	0.9	1.5	1.3	1.2	1.0	1.5	2.4
1999	0.9	...	...	*	0.8	1.1	1.2	1.3	0.9	1.0	1.4	2.6

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,X87.0)												
Non-Hispanic 4/, female												
2002	0.3	...	...	*	0.2	0.3	0.5	0.4	0.5	0.4	0.5	0.7
2001 5/	0.3	...	...	*	0.2	0.3	0.5	0.5	0.4	0.4	0.5	*
2000	0.3	...	...	*	0.2	0.3	0.4	0.5	0.4	0.4	0.5	*
1999	0.3	...	...	*	0.2	0.2	0.6	0.5	0.4	0.5	0.4	0.7
Non-Hispanic white, both sexes												
2002	0.7	...	...	*	0.5	0.7	1.1	1.1	0.9	0.7	0.9	1.4
2001 5/	0.7	...	...	*	0.5	0.7	1.0	1.1	0.8	0.6	0.9	1.0
2000	0.6	...	...	*	0.4	0.6	0.9	0.9	0.8	0.7	0.9	1.2
1999	0.6	...	...	*	0.5	0.6	0.9	0.9	0.6	0.7	0.8	1.3
Non-Hispanic white, male												
2002	1.1	...	...	*	0.8	1.1	1.6	1.7	1.2	1.0	1.4	2.9
2001 5/	1.0	...	...	*	0.8	1.2	1.4	1.6	1.2	0.9	1.5	2.6
2000	0.9	...	...	*	0.7	0.9	1.5	1.3	1.2	1.0	1.5	2.6
1999	0.9	...	...	*	0.8	1.1	1.2	1.4	0.9	0.9	1.5	2.9
Non-Hispanic white, female												
2002	0.4	...	...	*	0.2	0.4	0.5	0.5	0.5	0.4	0.5	0.7
2001 5/	0.3	...	...	*	0.2	0.3	0.5	0.5	0.4	0.4	0.5	*
2000	0.3	...	...	*	0.2	0.3	0.4	0.5	0.5	0.4	0.5	*
1999	0.3	...	...	*	0.2	0.2	0.6	0.5	0.3	0.4	0.4	*
Non-Hispanic black, both sexes												
2002	0.5	...	...	*	0.4	0.6	0.9	0.8	*	*	*	*
2001 5/	0.4	...	...	*	0.4	0.6	0.7	0.9	*	*	*	*
2000	0.5	...	...	*	0.6	0.6	1.0	0.7	*	*	*	*
1999	0.5	...	...	*	0.4	0.8	0.9	0.6	*	*	*	*
Non-Hispanic black, male												
2002	0.8	...	...	*	0.7	1.2	1.4	1.4	*	*	*	*
2001 5/	0.7	...	...	*	0.7	1.1	1.1	1.2	*	*	*	*
2000	0.8	...	...	*	0.9	0.9	1.7	1.1	*	*	*	*
1999	0.7	...	...	*	0.7	1.3	1.4	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	...	...	*	*	*	*	*	*	*	*	*
2001 5/	0.2	...	...	*	*	*	*	*	*	*	*	*
2000	0.2	...	...	*	*	*	*	*	*	*	*	*
1999	0.3	...	...	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
All origins 3/, both sexes												
2002	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 5/	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
All origins 3/, male												
2002	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 5/	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
All origins 3/, female												
2002	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 5/	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0
Hispanic, both sexes												
2002	8.1	6.2	2.6	0.9	17.6	12.3	8.0	5.9	4.4	2.5	*	*
2001 5/	9.0	6.7	2.3	0.8	18.3	14.0	10.4	7.3	4.4	4.0	*	*
2000	8.3	7.0	1.9	1.1	17.1	12.4	8.7	6.7	4.5	3.3	*	*
1999	8.4	7.0	1.8	1.4	17.7	12.7	8.8	6.6	4.2	3.2	*	*
Hispanic, male												
2002	13.2	6.6	3.1	1.0	29.6	19.8	12.1	9.1	7.7	4.8	*	*
2001 5/	14.5	7.9	2.5	1.1	30.5	21.7	16.0	11.3	6.8	6.6	*	*
2000	13.4	6.6	2.0	1.6	28.5	19.9	13.5	10.2	7.1	4.6	*	*
1999	13.8	6.7	1.7	1.8	29.1	20.6	13.9	10.3	7.3	5.7	*	*
Hispanic, female												
2002	2.6	5.9	2.1	0.8	3.8	3.4	3.5	2.7	*	*	*	*
2001 5/	3.2	5.5	2.1	*	4.0	5.1	4.4	3.4	2.4	*	*	*
2000	2.8	7.4	1.9	0.6	3.7	3.7	3.6	3.3	2.2	*	*	*
1999	2.8	7.3	1.9	0.9	4.4	3.5	3.3	3.1	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	5.7	7.7	2.7	0.9	11.8	10.9	7.0	4.6	3.0	2.3	2.2	2.1
2001 5/	6.7	8.4	2.8	0.8	12.1	12.8	9.2	6.1	3.9	2.8	2.5	2.4
2000	5.6	9.6	2.4	0.9	11.6	10.0	6.9	4.4	2.9	2.3	2.4	2.3
1999	5.7	8.9	2.6	1.0	11.9	10.0	6.8	4.3	2.9	2.5	2.5	2.4
Non-Hispanic 4/, male												
2002	8.9	8.0	2.9	0.9	19.6	17.6	10.4	6.8	4.5	3.2	3.1	2.9
2001 5/	10.4	9.8	3.2	0.9	20.1	20.3	13.4	9.2	5.8	3.8	3.1	3.1
2000	8.6	11.2	2.6	1.0	19.1	15.9	9.8	6.6	4.4	3.2	3.1	3.1
1999	8.6	10.1	2.6	1.1	19.4	15.6	9.7	6.4	4.5	3.5	3.3	3.4

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,X87.1)												
Non-Hispanic 4/, female												
2002	2.7	7.4	2.6	0.8	3.7	4.3	3.8	2.5	1.7	1.5	1.7	1.7
2001 5/	3.2	6.9	2.4	0.7	3.8	5.3	5.1	3.1	2.2	2.0	2.1	2.1
2000	2.7	7.9	2.2	0.7	3.8	4.1	4.0	2.3	1.5	1.6	1.9	2.0
1999	2.9	7.6	2.5	0.9	4.2	4.6	4.0	2.3	1.4	1.7	2.0	2.0
Non-Hispanic white, both sexes												
2002	2.8	5.0	1.8	0.5	3.7	3.9	4.1	3.0	2.4	1.9	1.8	1.8
2001 5/	4.0	5.8	1.7	0.6	4.2	6.5	6.6	4.6	3.3	2.3	2.1	2.2
2000	2.8	6.2	1.6	0.6	3.5	4.0	4.0	2.9	2.2	1.8	2.0	2.1
1999	2.9	6.3	1.7	0.7	3.9	4.3	4.1	2.9	2.2	2.1	2.0	2.1
Non-Hispanic white, male												
2002	3.8	5.8	1.9	0.5	5.2	5.3	5.6	3.9	3.3	2.4	2.3	2.7
2001 5/	5.6	6.8	2.0	0.6	5.7	9.3	9.5	6.7	4.7	2.9	2.6	2.8
2000	3.6	8.3	1.8	0.7	4.7	5.2	5.2	3.9	3.1	2.2	2.4	2.6
1999	3.9	8.0	1.7	0.7	5.1	5.7	5.4	4.2	3.2	2.7	2.6	2.6
Non-Hispanic white, female												
2002	1.9	4.1	1.8	0.6	2.2	2.6	2.5	2.1	1.5	1.4	1.5	1.5
2001 5/	2.4	4.7	1.5	0.7	2.7	3.7	3.7	2.6	1.9	1.8	1.8	2.0
2000	1.9	4.1	1.3	0.6	2.3	2.7	2.7	1.9	1.3	1.4	1.7	1.9
1999	2.0	4.4	1.6	0.7	2.6	2.9	2.8	1.7	1.3	1.6	1.6	1.9
Non-Hispanic black, both sexes												
2002	22.5	17.6	6.5	2.0	48.3	47.0	24.3	15.8	8.4	5.8	7.3	*
2001 5/	22.7	19.5	7.1	1.4	48.7	45.3	25.5	16.2	8.9	7.2	6.8	*
2000	22.1	24.1	6.1	1.9	49.3	42.2	24.5	15.0	8.4	7.6	7.1	*
1999	21.7	20.0	6.4	2.3	49.5	40.6	23.7	14.1	8.4	6.4	7.5	*
Non-Hispanic black, male												
2002	39.4	16.4	7.3	2.6	85.8	85.3	39.8	27.5	15.8	10.5	13.8	*
2001 5/	39.3	21.7	8.4	2.0	88.3	81.7	39.5	27.2	15.0	12.6	10.2	*
2000	38.3	24.9	6.1	2.1	88.1	76.5	39.4	26.3	15.0	13.9	12.0	*
1999	37.2	20.1	6.2	2.7	88.1	71.5	37.9	23.7	15.4	11.9	12.2	*
Non-Hispanic black, female												
2002	7.2	18.9	5.7	1.5	10.6	12.3	10.6	5.7	2.5	2.4	3.6	*
2001 5/	7.6	17.1	5.7	0.8	9.2	12.5	13.2	6.9	4.0	3.3	4.9	*
2000	7.4	23.3	6.0	1.6	11.0	11.4	11.3	5.3	3.2	3.2	4.3	*
1999	7.7	20.0	6.7	1.8	11.4	12.8	11.1	5.9	2.7	2.5	4.9	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by drugs, biological substances and other and unspecified noxious substances (*U01.6-*U01.7,X85-X90)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by hanging, strangulation and suffocation (X91)												
All origins 3/, both sexes												
2002	0.2	0.8	0.2	0.1	0.3	0.3	0.3	0.3	0.1	0.2	*	*
2001	0.2	1.0	*	*	0.3	0.3	0.3	0.2	0.2	0.3	0.2	*
2000	0.2	0.7	0.2	0.1	0.3	0.3	0.3	0.2	0.2	0.2	0.2	*
1999	0.3	0.9	0.1	0.1	0.3	0.4	0.3	0.2	0.1	0.2	0.2	*
All origins 3/, male												
2002	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	*	*	*
2001	0.2	1.2	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2000	0.2	*	*	*	0.1	0.2	0.2	0.1	0.2	*	*	*
1999	0.2	1.1	*	*	0.1	0.2	0.2	0.1	*	0.3	*	*
All origins 3/, female												
2002	0.3	*	*	0.1	0.4	0.4	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	0.2	0.3	*
2000	0.3	*	*	*	0.5	0.5	0.4	0.3	*	*	*	*
1999	0.3	*	*	0.1	0.5	0.6	0.5	0.2	*	0.2	0.3	*
Hispanic, both sexes												
2002	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2001	0.3	*	*	*	0.3	0.3	*	*	*	*	*	*
2000	0.2	*	*	*	*	0.3	*	*	*	*	*	*
1999	0.3	*	*	*	*	0.4	0.4	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.2	0.9	0.2	0.1	0.3	0.3	0.3	0.3	0.1	0.1	*	*
2001	0.2	1.0	*	*	0.3	0.3	0.3	0.2	0.2	0.2	0.2	*
2000	0.2	0.7	0.2	0.1	0.3	0.3	0.3	0.2	0.1	0.2	0.2	*
1999	0.2	1.0	0.2	0.1	0.3	0.4	0.3	0.2	0.1	0.2	0.2	*
Non-Hispanic 4/, male												
2002	0.2	*	*	*	0.2	0.2	0.2	0.2	*	*	*	*
2001	0.2	1.3	*	*	0.1	0.2	0.2	0.2	0.2	0.3	*	*
2000	0.2	*	*	*	*	0.2	0.2	0.1	*	*	*	*
1999	0.2	1.4	*	*	*	0.2	0.2	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by hanging, strangulation and suffocation (X91)												
Non-Hispanic 4/, female												
2002	0.3	*	*	*	0.4	0.4	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.2	0.2	0.2	0.3	*
2000	0.3	*	*	*	0.5	0.5	0.4	0.2	*	*	*	*
1999	0.3	*	*	*	0.5	0.6	0.5	0.2	*	0.2	*	*
Non-Hispanic white, both sexes												
2002	0.2	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
2001	0.2	*	*	*	0.2	0.3	0.2	0.2	0.1	0.2	*	*
2000	0.2	*	*	*	0.2	0.2	0.2	0.2	0.1	*	*	*
1999	0.2	*	*	0.1	0.2	0.3	0.2	0.1	*	0.2	0.2	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
2001	0.1	*	*	*	*	*	*	0.2	*	*	*	*
2000	0.1	*	*	*	*	0.2	0.2	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic white, female												
2002	0.2	*	*	*	0.3	0.3	0.3	0.3	*	*	*	*
2001	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
2000	0.2	*	*	*	0.4	0.3	0.3	0.2	*	*	*	*
1999	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.6	*	*	*	0.6	0.9	1.0	0.7	*	*	*	*
2001	0.6	*	*	*	0.7	0.8	1.0	0.6	*	*	*	*
2000	0.6	*	*	*	0.7	1.0	0.9	*	*	*	*	*
1999	0.6	*	*	*	0.7	1.2	1.0	*	*	*	*	*
Non-Hispanic black, male												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.8	*	*	*	0.8	1.3	1.5	*	*	*	*	*
2001	0.7	*	*	*	0.9	1.1	1.4	*	*	*	*	*
2000	0.8	*	*	*	1.2	1.5	1.3	*	*	*	*	*
1999	0.8	*	*	*	1.0	1.7	1.4	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins 3/, both sexes												
2002	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6
All origins 3/, male												
2002	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*
All origins 3/, female												
2002	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*
Hispanic, both sexes												
2002	5.6	*	*	0.6	14.3	9.3	5.1	3.2	1.6	*	*	*
2001	5.7	*	*	0.4	14.6	9.5	5.6	3.0	1.1	*	*	*
2000	5.5	*	*	0.6	13.7	8.9	4.9	4.0	2.0	*	*	*
1999	5.7	*	*	0.8	14.4	8.7	5.2	3.8	2.1	*	*	*
Hispanic, male												
2002	9.7	*	*	0.8	24.9	15.6	8.1	5.0	2.8	*	*	*
2001	10.0	*	*	0.6	24.9	16.1	9.1	4.8	*	*	*	*
2000	9.6	*	*	1.0	23.5	15.0	7.8	6.4	3.2	*	*	*
1999	9.8	*	*	1.2	24.2	14.7	8.6	5.7	3.5	*	*	*
Hispanic, female												
2002	1.2	*	*	*	2.1	1.8	1.8	1.4	*	*	*	*
2001	1.3	*	*	*	2.5	1.8	1.9	1.2	*	*	*	*
2000	1.3	*	*	*	2.2	1.9	1.7	1.7	*	*	*	*
1999	1.4	*	*	*	2.9	1.7	1.7	1.9	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	3.8	*	0.3	0.5	9.8	8.5	4.4	2.5	1.5	0.9	0.9	*
2001	3.7	*	0.3	0.4	9.6	8.0	4.2	2.3	1.5	1.0	0.8	*
2000	3.6	*	0.2	0.4	9.3	7.4	4.1	2.4	1.5	1.0	0.7	0.7
1999	3.6	*	0.4	0.5	9.6	7.3	4.0	2.3	1.5	1.1	0.8	0.5
Non-Hispanic 4/, male												
2002	6.5	*	0.3	0.6	17.0	14.7	7.1	3.7	2.3	1.2	1.2	*
2001	6.3	*	0.3	0.6	17.0	13.8	6.5	3.7	2.2	1.4	1.1	*
2000	6.0	*	*	0.6	16.3	12.7	6.4	3.7	2.2	1.3	0.8	*
1999	6.0	*	0.4	0.7	16.7	12.4	6.0	3.6	2.5	1.5	0.9	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic 4/, female												
2002	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001	1.2	*	0.3	0.3	2.0	2.3	2.0	1.0	0.8	0.7	0.6	*
2000	1.2	*	*	0.2	2.2	2.2	1.9	1.1	0.8	0.7	0.6	*
1999	1.3	*	0.4	0.3	2.4	2.3	2.0	1.1	0.6	0.7	0.6	*
Non-Hispanic white, both sexes												
2002	1.5	*	0.2	0.3	2.5	2.6	2.4	1.6	1.2	0.7	0.8	*
2001	1.6	*	0.2	0.4	2.4	2.7	2.4	1.6	1.2	0.8	0.7	*
2000	1.4	*	*	0.3	2.2	2.4	2.3	1.6	1.2	0.8	0.6	0.6
1999	1.5	*	0.2	0.3	2.4	2.6	2.3	1.5	1.1	0.9	0.6	0.6
Non-Hispanic white, male												
2002	2.2	*	*	0.3	3.7	3.7	3.4	2.2	1.7	0.9	1.0	*
2001	2.2	*	*	0.4	3.6	3.9	3.4	2.3	1.6	1.0	0.9	*
2000	2.0	*	*	0.4	3.2	3.4	3.2	2.2	1.6	0.9	0.6	*
1999	2.2	*	*	0.4	3.5	3.9	3.2	2.2	1.7	1.2	0.6	*
Non-Hispanic white, female												
2002	0.9	*	*	0.3	1.2	1.4	1.3	1.1	0.7	0.6	0.7	*
2001	0.9	*	*	0.3	1.2	1.5	1.4	0.8	0.8	0.7	0.6	*
2000	0.9	*	*	0.2	1.2	1.4	1.4	0.9	0.8	0.7	0.6	*
1999	0.9	*	*	0.2	1.2	1.4	1.5	0.8	0.6	0.7	0.6	*
Non-Hispanic black, both sexes												
2002	17.1	*	*	1.2	43.1	39.7	16.7	8.6	4.2	2.4	2.1	*
2001	16.3	*	*	0.9	43.0	36.4	15.5	7.7	3.7	2.6	*	*
2000	16.0	*	*	1.0	42.6	34.5	15.4	8.5	3.6	2.8	*	*
1999	15.9	*	*	1.3	43.6	32.4	14.4	8.1	4.5	2.4	2.4	*
Non-Hispanic black, male												
2002	31.9	*	*	1.6	78.6	75.6	30.0	15.3	7.7	4.7	*	*
2001	30.6	*	*	1.3	80.6	69.4	26.7	14.0	7.1	5.5	*	*
2000	30.1	*	*	1.6	79.1	65.7	27.4	15.3	6.5	5.5	*	*
1999	29.3	*	*	1.8	80.0	61.1	24.6	14.1	9.1	4.7	*	*
Non-Hispanic black, female												
2002	3.7	*	*	0.8	7.5	7.2	5.0	2.8	*	*	*	*
2001	3.2	*	*	*	5.6	6.5	5.5	2.2	*	*	*	*
2000	3.3	*	*	*	6.7	6.4	4.7	2.6	*	*	*	*
1999	3.7	*	*	0.7	7.8	6.6	5.5	3.0	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by sharp object (X99)												
All origins 3/, both sexes												
2002	0.7	*	*	0.1	1.0	1.2	1.1	0.8	0.5	0.4	0.4	*
2001	0.7	*	*	0.1	1.2	1.2	1.0	0.7	0.4	0.4	0.4	*
2000	0.6	*	*	0.1	1.0	1.1	1.0	0.6	0.4	0.3	0.5	*
1999	0.7	*	*	0.1	1.1	1.2	1.0	0.7	0.4	0.3	0.3	*
All origins 3/, male												
2002	1.0	*	*	0.1	1.6	1.7	1.5	1.2	0.8	0.7	0.4	*
2001	1.0	*	*	*	1.8	1.7	1.3	1.0	0.6	0.5	0.4	*
2000	0.9	*	*	*	1.6	1.5	1.4	0.8	0.7	0.5	0.7	*
1999	1.0	*	*	*	1.6	1.7	1.6	1.0	0.7	0.5	0.5	*
All origins 3/, female												
2002	0.4	*	*	*	0.4	0.8	0.7	0.4	0.3	0.3	0.3	*
2001	0.4	*	*	*	0.5	0.7	0.7	0.3	0.3	0.4	0.4	*
2000	0.4	*	*	*	0.4	0.6	0.7	0.4	0.2	*	0.3	*
1999	0.4	*	*	*	0.5	0.6	0.5	0.4	0.3	0.2	*	*
Hispanic, both sexes												
2002	1.1	*	*	*	2.0	1.7	1.5	1.0	*	*	*	*
2001	1.1	*	*	*	2.0	1.7	1.4	1.0	1.2	*	*	*
2000	1.2	*	*	*	2.0	1.9	1.8	1.0	*	*	*	*
1999	1.0	*	*	*	1.7	1.9	1.3	1.1	*	*	*	*
Hispanic, male												
2002	1.7	*	*	*	3.2	2.4	2.1	1.4	*	*	*	*
2001	1.7	*	*	*	3.3	2.4	1.9	1.5	*	*	*	*
2000	1.9	*	*	*	3.3	2.9	2.8	1.4	*	*	*	*
1999	1.6	*	*	*	2.7	2.9	2.1	1.6	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	0.6	0.8	0.8	*	*	*	*	*
2001	0.5	*	*	*	*	0.8	0.8	*	*	*	*	*
2000	0.5	*	*	*	*	0.7	*	*	*	*	*	*
1999	0.4	*	*	*	*	0.7	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.7	*	*	0.1	0.9	1.1	1.1	0.7	0.5	0.5	0.3	*
2001	0.6	*	*	*	1.0	1.1	1.0	0.6	0.4	0.4	0.4	*
2000	0.6	*	*	0.1	0.8	0.9	0.9	0.5	0.4	0.3	0.5	*
1999	0.6	*	*	0.1	0.9	1.0	1.0	0.6	0.4	0.3	0.3	*
Non-Hispanic 4/, male												
2002	0.9	*	*	0.1	1.3	1.5	1.5	1.1	0.7	0.6	*	*
2001	0.9	*	*	*	1.5	1.5	1.2	1.0	0.5	0.5	*	*
2000	0.8	*	*	*	1.2	1.2	1.2	0.7	0.6	0.5	0.7	*
1999	0.9	*	*	*	1.3	1.4	1.5	0.9	0.6	0.4	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by sharp object (X99)												
Non-Hispanic 4/, female												
2002	0.4	*	*	*	0.4	0.8	0.7	0.3	0.3	0.3	0.3	*
2001	0.4	*	*	*	0.6	0.6	0.7	0.3	0.3	0.4	0.4	*
2000	0.3	*	*	*	0.4	0.5	0.7	0.3	0.2	*	*	*
1999	0.4	*	*	*	0.5	0.6	0.5	0.3	0.2	0.3	*	*
Non-Hispanic white, both sexes												
2002	0.4	*	*	*	0.4	0.5	0.6	0.4	0.3	0.4	0.2	*
2001	0.4	*	*	*	0.6	0.6	0.5	0.4	0.3	0.3	0.3	*
2000	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	0.2	0.3	*
1999	0.4	*	*	*	0.5	0.6	0.5	0.4	0.3	0.2	0.2	*
Non-Hispanic white, male												
2002	0.5	*	*	*	0.6	0.6	0.9	0.6	0.4	0.5	*	*
2001	0.5	*	*	*	0.8	0.8	0.6	0.5	0.4	0.4	*	*
2000	0.4	*	*	*	0.6	0.6	0.6	0.4	0.4	0.3	0.6	*
1999	0.5	*	*	*	0.7	0.8	0.7	0.6	0.4	0.3	*	*
Non-Hispanic white, female												
2002	0.2	*	*	*	0.2	0.4	0.4	0.2	0.3	0.3	*	*
2001	0.3	*	*	*	0.3	0.4	0.4	0.2	*	0.3	0.3	*
2000	0.2	*	*	*	0.2	0.4	0.4	0.2	*	*	*	*
1999	0.2	*	*	*	0.3	0.4	0.4	0.2	0.2	*	*	*
Non-Hispanic black, both sexes												
2002	2.2	*	*	*	2.7	3.9	3.6	3.0	1.8	1.3	*	*
2001	2.1	*	*	*	3.0	3.6	3.6	2.4	1.3	1.2	*	*
2000	1.9	*	*	*	2.5	2.8	3.7	2.3	1.6	1.4	*	*
1999	2.1	*	*	*	2.9	3.2	3.7	2.4	1.4	*	*	*
Non-Hispanic black, male												
2002	3.3	*	*	*	4.3	5.8	4.9	5.1	3.3	*	*	*
2001	3.1	*	*	*	4.5	5.4	5.1	4.5	2.0	*	*	*
2000	2.9	*	*	*	3.8	4.5	5.0	3.7	2.5	*	*	*
1999	3.2	*	*	*	4.3	4.7	6.4	3.6	2.7	*	*	*
Non-Hispanic black, female												
2002	1.2	*	*	*	1.2	2.2	2.5	1.1	*	*	*	*
2001	1.1	*	*	*	1.6	1.8	2.2	*	*	*	*	*
2000	1.1	*	*	*	1.3	1.3	2.6	1.1	*	*	*	*
1999	1.0	*	*	*	1.6	1.8	1.3	1.3	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by blunt object (Y00)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*
All origins 3/, male												
2002	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
2000	0.1	*	*	*	0.1	*	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.2	0.1	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	0.1	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.2	*	*	*	*
2000	0.1	*	*	*	*	*	0.2	0.2	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.2	0.1	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by blunt object (Y00)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	0.1	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	0.2	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by bodily force (Y04)												
All origins 3/, both sexes												
2002	0.0	*	*	*	0.0	0.1	*	0.1	*	*	*	*
2001	0.0	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.0	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.2	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.2	0.1	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.0	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.0	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	*	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.2	0.1	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by bodily force (Y04)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sexual assault (homicide) by bodily force (Y05)												
All origins 3/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Sexual assault (homicide) by bodily force (Y05)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)												
All origins 3/, both sexes												
2002	0.1	2.4	0.6	*	*	*	*	*	*	*	*	*
2001	0.1	2.9	0.5	*	*	*	*	*	*	*	*	*
2000	0.1	3.4	0.5	*	*	*	*	*	*	*	*	*
1999	0.1	3.3	0.4	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.1	2.5	0.6	*	*	*	*	*	*	*	*	*
2001	0.1	3.2	0.7	*	*	*	*	*	*	*	*	*
2000	0.1	4.0	0.5	*	*	*	*	*	*	*	*	*
1999	0.1	3.7	0.4	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.1	2.4	0.6	*	*	*	*	*	*	*	*	*
2001	0.1	2.5	0.4	*	*	*	*	*	*	*	*	*
2000	0.1	2.7	0.4	*	*	*	*	*	*	*	*	*
1999	0.1	2.9	0.4	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	2.6	*	*	*	*	*	*	*	*	*	*
2001	0.1	2.6	*	*	*	*	*	*	*	*	*	*
2000	0.1	2.6	*	*	*	*	*	*	*	*	*	*
1999	0.1	2.6	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	2.3	0.6	*	*	*	*	*	*	*	*	*
2001	0.1	3.0	0.5	*	*	*	*	*	*	*	*	*
2000	0.1	3.6	0.5	*	*	*	*	*	*	*	*	*
1999	0.1	3.4	0.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	2.3	0.6	*	*	*	*	*	*	*	*	*
2001	0.1	3.3	0.7	*	*	*	*	*	*	*	*	*
2000	0.1	4.4	0.6	*	*	*	*	*	*	*	*	*
1999	0.1	3.9	0.5	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Neglect, abandonment and other maltreatment syndromes (Y06-Y07)												
Non-Hispanic 4/, female												
2002	0.1	2.3	0.6	*	*	*	*	*	*	*	*	*
2001	0.1	2.5	0.4	*	*	*	*	*	*	*	*	*
2000	0.1	2.7	0.4	*	*	*	*	*	*	*	*	*
1999	0.1	2.8	0.4	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	1.5	0.4	*	*	*	*	*	*	*	*	*
2001	0.1	2.0	0.4	*	*	*	*	*	*	*	*	*
2000	0.1	2.6	0.4	*	*	*	*	*	*	*	*	*
1999	0.0	2.4	0.2	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	0.4	*	*	*	*	*	*	*	*	*
2001	0.1	2.4	0.5	*	*	*	*	*	*	*	*	*
2000	0.1	3.8	0.4	*	*	*	*	*	*	*	*	*
1999	0.1	3.1	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	0.5	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.2	5.7	1.2	*	*	*	*	*	*	*	*	*
2001	0.2	7.0	1.3	*	*	*	*	*	*	*	*	*
2000	0.2	8.3	1.1	*	*	*	*	*	*	*	*	*
1999	0.3	7.7	1.4	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	7.7	*	*	*	*	*	*	*	*	*	*
2000	0.3	8.2	*	*	*	*	*	*	*	*	*	*
1999	0.3	7.3	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	8.5	*	*	*	*	*	*	*	*	*	*
1999	0.2	8.2	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)												
All origins 3/, both sexes												
2002	0.9	3.7	1.4	0.1	0.7	0.9	1.0	1.0	0.8	0.7	0.7	0.9
2001 5/	2.0	3.5	1.6	0.2	1.1	3.1	3.5	2.8	1.8	1.0	0.9	1.1
2000	1.0	4.3	1.3	0.2	1.0	1.1	1.3	1.1	0.8	0.7	1.0	0.9
1999	1.0	3.8	1.3	0.3	0.9	1.1	1.3	1.1	0.8	0.7	1.0	1.1
All origins 3/, male												
2002	1.2	4.0	1.6	0.1	1.0	1.1	1.4	1.5	1.4	1.2	1.1	*
2001 5/	2.9	4.0	1.7	0.2	1.5	4.4	5.3	4.2	2.8	1.6	1.2	1.6
2000	1.4	4.7	1.5	0.2	1.3	1.5	1.8	1.7	1.3	1.2	1.4	*
1999	1.3	4.0	1.4	0.3	1.2	1.4	1.8	1.6	1.2	1.1	1.6	*
All origins 3/, female												
2002	0.5	3.3	1.2	0.2	0.5	0.6	0.6	0.5	0.3	0.4	0.5	0.8
2001 5/	1.1	3.1	1.4	0.2	0.8	1.8	1.7	1.4	1.0	0.5	0.7	0.9
2000	0.7	3.9	1.1	0.3	0.7	0.8	0.9	0.6	0.4	0.3	0.8	0.7
1999	0.7	3.6	1.2	0.3	0.7	0.8	0.9	0.5	0.4	0.4	0.6	0.9
Hispanic, both sexes												
2002	0.9	2.4	1.3	*	0.8	0.9	1.0	1.2	1.4	*	*	*
2001 5/	1.7	2.7	1.1	*	1.2	2.4	2.9	2.9	1.9	*	*	*
2000	1.1	3.1	1.1	*	1.0	1.2	1.5	1.3	1.3	*	*	*
1999	1.2	3.0	1.3	0.4	1.2	1.4	1.7	1.4	*	*	*	*
Hispanic, male												
2002	1.3	*	1.6	*	1.1	1.4	1.4	2.0	2.7	*	*	*
2001 5/	2.4	*	1.4	*	1.8	2.8	4.4	4.5	3.0	*	*	*
2000	1.5	*	*	*	1.4	1.5	2.3	2.0	*	*	*	*
1999	1.8	*	*	*	1.8	2.1	2.7	2.4	*	*	*	*
Hispanic, female												
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001 5/	0.9	*	*	*	*	1.9	1.3	1.2	*	*	*	*
2000	0.6	*	*	*	*	0.7	*	*	*	*	*	*
1999	0.6	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.8	3.9	1.5	0.1	0.7	0.8	1.0	1.0	0.8	0.7	0.7	0.9
2001 5/	2.0	3.6	1.7	0.2	1.1	3.2	3.5	2.7	1.8	0.9	0.9	1.2
2000	1.0	4.5	1.4	0.2	1.0	1.1	1.3	1.1	0.7	0.7	1.0	0.9
1999	0.9	3.8	1.3	0.3	0.9	1.1	1.3	1.0	0.7	0.7	1.0	1.1
Non-Hispanic 4/, male												
2002	1.1	4.3	1.6	0.1	0.9	1.1	1.4	1.4	1.3	1.0	1.2	*
2001 5/	2.8	3.9	1.8	0.2	1.4	4.6	5.3	4.0	2.7	1.4	1.2	1.6
2000	1.3	5.0	1.6	0.2	1.3	1.5	1.7	1.6	1.2	1.1	1.4	*
1999	1.2	4.1	1.5	0.2	1.1	1.2	1.7	1.5	1.1	1.0	1.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)												
Non-Hispanic 4/, female												
2002	0.5	3.6	1.3	0.2	0.5	0.6	0.6	0.5	0.3	0.4	0.4	0.8
2001 5/	1.1	3.3	1.5	0.2	0.8	1.7	1.8	1.4	0.9	0.5	0.7	1.0
2000	0.7	4.0	1.2	0.3	0.7	0.8	0.9	0.5	0.3	0.3	0.8	0.7
1999	0.7	3.6	1.2	0.3	0.7	0.9	0.9	0.5	0.4	0.4	0.6	0.9
Non-Hispanic white, both sexes												
2002	0.6	2.6	0.9	0.1	0.5	0.5	0.8	0.7	0.6	0.5	0.6	0.8
2001 5/	1.7	2.6	1.0	0.2	0.9	2.9	3.3	2.4	1.6	0.8	0.8	1.0
2000	0.6	2.7	0.7	0.2	0.6	0.7	0.9	0.7	0.5	0.5	0.8	0.8
1999	0.7	2.7	0.9	0.2	0.6	0.7	0.8	0.7	0.6	0.6	0.8	0.9
Non-Hispanic white, male												
2002	0.8	3.0	1.0	*	0.6	0.7	1.0	0.9	0.9	0.8	0.8	*
2001 5/	2.6	2.9	1.1	*	1.0	4.3	5.2	3.5	2.4	1.2	1.0	*
2000	0.8	3.4	0.9	0.2	0.6	0.8	1.1	1.0	0.8	0.8	1.0	*
1999	0.8	3.0	1.0	*	0.7	0.7	1.1	1.0	0.9	0.8	1.4	*
Non-Hispanic white, female												
2002	0.4	2.1	0.7	*	0.4	0.4	0.5	0.5	0.3	0.3	0.4	0.7
2001 5/	0.9	2.3	0.9	0.2	0.7	1.4	1.4	1.2	0.8	0.4	0.6	0.9
2000	0.5	2.0	0.5	0.2	0.5	0.5	0.6	0.5	0.3	0.3	0.7	*
1999	0.5	2.3	0.8	0.2	0.6	0.6	0.6	0.5	0.3	0.4	0.5	0.9
Non-Hispanic black, both sexes												
2002	2.2	9.3	3.9	0.4	1.6	2.2	2.6	3.1	2.0	1.6	2.6	*
2001 5/	3.2	8.2	4.3	0.3	1.8	4.3	5.1	5.0	2.9	2.5	2.3	*
2000	3.0	12.0	4.0	0.6	3.1	3.4	3.9	3.5	2.4	2.2	3.4	*
1999	2.5	9.0	3.4	0.6	1.9	3.2	3.9	2.9	1.6	2.1	2.4	*
Non-Hispanic black, male												
2002	3.2	9.1	4.3	*	2.3	2.9	3.8	5.5	4.1	2.9	6.2	*
2001 5/	4.4	8.3	5.1	*	2.5	6.0	6.5	7.4	4.8	4.3	*	*
2000	4.3	11.9	4.1	*	4.4	5.2	5.4	6.4	4.8	4.3	6.2	*
1999	3.5	9.4	3.7	0.6	3.0	4.2	5.4	5.0	2.5	4.2	*	*
Non-Hispanic black, female												
2002	1.3	9.5	3.4	*	0.9	1.5	1.6	1.1	*	*	*	*
2001 5/	2.1	8.1	3.5	*	1.0	2.7	3.8	3.0	*	*	*	*
2000	1.8	12.0	4.0	0.8	1.7	1.9	2.5	1.0	*	*	*	*
1999	1.6	8.6	3.0	0.6	0.9	2.2	2.6	1.0	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All origins 3/, both sexes												
2002	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
All origins 3/, male												
2002	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
All origins 3/, female												
2002	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1
Hispanic, both sexes												
2002	0.8	*	*	*	0.7	1.0	1.5	1.3	*	*	*	*
2001	0.8	*	*	*	0.5	0.9	1.7	1.6	*	*	*	*
2000	0.6	*	*	*	0.5	0.9	1.2	0.8	*	*	*	*
1999	0.7	*	*	*	0.8	1.1	1.2	0.9	*	*	*	*
Hispanic, male												
2002	1.2	*	*	*	1.2	1.5	2.1	2.2	*	*	*	*
2001	1.1	*	*	*	0.7	1.4	2.6	2.1	*	*	*	*
2000	0.9	*	*	*	0.8	1.4	1.8	1.3	*	*	*	*
1999	1.1	*	*	*	1.3	1.6	1.9	1.6	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	0.9	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.8	2.7	0.4	0.2	1.3	2.3	3.6	3.0	1.2	0.7	0.8	1.4
2001	1.6	2.4	0.3	0.1	1.1	2.0	3.3	2.6	0.9	0.5	0.9	0.9
2000	1.5	1.9	0.3	0.2	1.0	1.8	3.0	2.4	0.9	0.6	0.6	1.4
1999	1.5	1.6	0.3	0.1	1.2	2.0	3.1	2.3	0.9	0.6	0.8	1.4
Non-Hispanic 4/, male												
2002	2.3	2.8	0.3	0.2	1.8	3.2	4.7	3.6	1.6	1.0	0.9	2.1
2001	2.1	2.7	*	0.2	1.7	2.8	4.4	3.5	1.2	0.7	1.2	*
2000	2.0	1.6	0.4	0.3	1.5	2.6	4.2	3.2	1.3	0.8	1.0	1.7
1999	2.1	1.9	*	0.2	1.8	2.7	4.2	3.1	1.3	0.9	1.1	2.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic 4/, female												
2002	1.3	2.6	0.4	0.2	0.7	1.4	2.6	2.3	0.9	0.5	0.7	1.1
2001	1.0	2.0	0.4	*	0.5	1.2	2.2	1.7	0.6	0.4	0.7	0.8
2000	0.9	2.2	*	*	0.5	1.1	1.9	1.5	0.6	0.3	0.4	1.2
1999	0.9	*	*	*	0.5	1.2	2.0	1.5	0.6	0.4	0.6	1.1
Non-Hispanic white, both sexes												
2002	1.8	1.8	0.4	0.2	1.4	2.4	3.7	2.9	1.2	0.7	0.7	1.5
2001	1.6	1.7	0.3	0.1	1.3	2.2	3.3	2.5	0.9	0.5	0.9	1.0
2000	1.4	1.5	0.2	0.1	1.1	1.9	3.0	2.3	0.8	0.5	0.7	1.3
1999	1.4	1.3	*	0.1	1.2	2.0	3.0	2.1	0.9	0.6	0.8	1.4
Non-Hispanic white, male												
2002	2.3	1.9	*	0.2	2.0	3.4	4.8	3.4	1.5	0.8	0.9	2.2
2001	2.1	2.0	*	0.2	1.8	3.0	4.3	3.3	1.1	0.7	1.2	*
2000	2.0	*	*	0.2	1.6	2.6	4.1	3.0	1.1	0.7	1.0	*
1999	2.0	*	*	0.2	1.8	2.7	4.0	2.6	1.1	0.8	1.1	2.1
Non-Hispanic white, female												
2002	1.3	1.8	*	0.2	0.8	1.5	2.7	2.4	0.9	0.6	0.6	1.1
2001	1.1	*	*	*	0.6	1.4	2.4	1.8	0.7	0.4	0.6	0.8
2000	0.9	*	*	*	0.6	1.2	1.9	1.5	0.6	0.3	0.4	1.2
1999	1.0	*	*	*	0.5	1.3	1.9	1.5	0.7	0.4	0.6	1.1
Non-Hispanic black, both sexes												
2002	2.0	5.7	*	*	0.9	2.3	4.3	4.3	1.5	*	*	*
2001	1.8	5.9	*	*	0.7	1.7	4.0	3.8	1.2	*	*	*
2000	1.9	*	*	0.3	0.9	2.0	4.1	3.7	1.8	*	*	*
1999	2.0	*	*	*	1.3	2.3	4.3	4.2	1.2	*	*	*
Non-Hispanic black, male												
2002	2.9	*	*	*	1.3	3.4	6.2	6.5	2.6	*	*	*
2001	2.8	6.4	*	*	1.3	2.7	6.3	6.3	2.4	*	*	*
2000	2.9	*	*	*	1.6	3.2	6.1	6.0	3.0	*	*	*
1999	3.1	*	*	*	2.0	3.7	6.6	7.3	2.6	*	*	*
Non-Hispanic black, female												
2002	1.2	*	*	*	*	1.3	2.7	2.4	*	*	*	*
2001	0.9	*	*	*	*	0.9	1.9	1.5	*	*	*	*
2000	1.0	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999	0.9	*	*	*	*	1.0	2.4	1.6	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)												
All origins 3/, both sexes												
2002	1.1	*	*	*	0.7	1.5	2.6	2.2	0.7	0.2	*	*
2001	1.0	*	*	*	0.6	1.3	2.4	1.8	0.5	0.2	0.2	*
2000	0.9	*	*	*	0.5	1.1	2.2	1.6	0.4	*	*	*
1999	0.9	*	*	*	0.5	1.3	2.2	1.5	0.4	0.2	*	*
All origins 3/, male												
2002	1.4	*	*	*	0.9	2.0	3.2	2.5	0.8	*	*	*
2001	1.3	*	*	*	0.8	1.7	3.1	2.5	0.5	*	*	*
2000	1.2	*	*	*	0.7	1.6	3.1	2.2	0.5	*	*	*
1999	1.2	*	*	*	0.7	1.8	3.0	2.0	0.5	*	*	*
All origins 3/, female												
2002	0.8	*	*	*	0.4	1.0	2.0	1.8	0.6	0.3	*	*
2001	0.7	*	*	*	0.3	0.9	1.7	1.2	0.4	*	*	*
2000	0.5	*	*	*	0.3	0.7	1.4	1.1	0.3	*	*	*
1999	0.6	*	*	*	0.3	0.8	1.5	1.1	0.4	0.2	*	*
Hispanic, both sexes												
2002	0.4	*	*	*	*	0.6	1.0	0.7	*	*	*	*
2001	0.4	*	*	*	*	0.6	1.0	0.8	*	*	*	*
2000	0.3	*	*	*	*	0.5	0.7	*	*	*	*	*
1999	0.4	*	*	*	*	0.7	0.8	*	*	*	*	*
Hispanic, male												
2002	0.5	*	*	*	*	0.7	1.3	*	*	*	*	*
2001	0.5	*	*	*	*	0.9	1.5	*	*	*	*	*
2000	0.4	*	*	*	*	0.9	0.8	*	*	*	*	*
1999	0.5	*	*	*	*	1.1	1.3	*	*	*	*	*
Hispanic, female												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.2	*	*	*	0.8	1.7	2.8	2.3	0.7	0.2	*	*
2001	1.1	*	*	*	0.7	1.5	2.6	1.9	0.5	0.2	0.2	*
2000	0.9	*	*	*	0.5	1.2	2.4	1.7	0.4	*	*	*
1999	1.0	*	*	*	0.6	1.4	2.4	1.6	0.5	0.2	*	*
Non-Hispanic 4/, male												
2002	1.5	*	*	*	1.1	2.3	3.5	2.6	0.8	*	*	*
2001	1.4	*	*	*	1.0	1.9	3.4	2.6	0.6	*	*	*
2000	1.3	*	*	*	0.8	1.7	3.4	2.3	0.5	*	*	*
1999	1.3	*	*	*	0.8	1.9	3.2	2.1	0.6	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)												
Non-Hispanic 4/, female												
2002	0.9	*	*	*	0.5	1.1	2.1	2.0	0.6	0.2	*	*
2001	0.7	*	*	*	0.4	1.0	1.9	1.3	0.5	*	*	*
2000	0.6	*	*	*	0.3	0.8	1.5	1.2	0.3	*	*	*
1999	0.7	*	*	*	0.3	0.9	1.6	1.2	0.4	*	*	*
Non-Hispanic white, both sexes												
2002	1.3	*	*	*	1.0	1.9	3.0	2.3	0.7	0.2	*	*
2001	1.1	*	*	*	0.8	1.6	2.7	1.9	0.5	0.1	*	*
2000	1.0	*	*	*	0.7	1.4	2.4	1.7	0.4	*	*	*
1999	1.0	*	*	*	0.7	1.5	2.3	1.5	0.5	0.2	*	*
Non-Hispanic white, male												
2002	1.6	*	*	*	1.3	2.6	3.7	2.5	0.8	*	*	*
2001	1.4	*	*	*	1.2	2.1	3.3	2.5	0.5	*	*	*
2000	1.3	*	*	*	0.9	1.8	3.4	2.2	0.5	*	*	*
1999	1.2	*	*	*	1.0	2.0	3.1	1.8	0.5	*	*	*
Non-Hispanic white, female												
2002	1.0	*	*	*	0.6	1.3	2.3	2.1	0.7	0.3	*	*
2001	0.8	*	*	*	0.5	1.2	2.0	1.4	0.5	*	*	*
2000	0.6	*	*	*	0.4	0.9	1.5	1.2	0.4	*	*	*
1999	0.7	*	*	*	0.4	1.1	1.6	1.2	0.5	*	*	*
Non-Hispanic black, both sexes												
2002	1.3	*	*	*	0.3	1.4	3.2	3.3	0.9	*	*	*
2001	1.1	*	*	*	*	1.2	3.1	2.8	*	*	*	*
2000	1.1	*	*	*	*	1.1	3.2	2.7	*	*	*	*
1999	1.2	*	*	*	*	1.4	3.3	3.0	*	*	*	*
Non-Hispanic black, male												
2002	1.7	*	*	*	*	1.9	4.3	4.6	*	*	*	*
2001	1.7	*	*	*	*	1.7	4.9	4.6	*	*	*	*
2000	1.6	*	*	*	*	1.7	4.6	4.1	*	*	*	*
1999	1.8	*	*	*	*	2.3	4.9	5.1	*	*	*	*
Non-Hispanic black, female												
2002	0.9	*	*	*	*	1.0	2.3	2.1	*	*	*	*
2001	0.5	*	*	*	*	0.8	1.5	1.2	*	*	*	*
2000	0.6	*	*	*	*	*	2.0	1.5	*	*	*	*
1999	0.6	*	*	*	*	*	2.0	1.3	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Poisoning by and exposure to other and unspecified solid or liquid substance, undetermined intent (Y15-Y16,Y18-Y19)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	0.1	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	0.0	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	0.0	*	*	*	*	*
1999	0.0	*	*	*	*	*	0.0	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Poisoning by and exposure to other gases and vapors, undetermined intent (Y17)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.5	0.2	0.2	*	*	*	*	*
Non-Hispanic white, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	0.6	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	1.1	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falling, jumping or pushed from a high place, undetermined intent (Y30)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falling, jumping or pushed from a high place, undetermined intent (Y30)												
Non-Hispanic 4/, female												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)												
All origins 3/, both sexes												
2002	0.4	2.1	0.3	0.1	0.3	0.3	0.5	0.5	0.4	0.4	0.5	1.1
2001	0.3	1.7	0.2	0.1	0.2	0.3	0.5	0.5	0.3	0.3	0.5	0.7
2000	0.3	1.6	0.2	0.1	0.2	0.4	0.4	0.5	0.4	0.3	0.4	0.9
1999	0.3	1.2	0.2	0.1	0.2	0.3	0.4	0.5	0.4	0.3	0.4	1.0
All origins 3/, male												
2002	0.5	2.0	0.3	0.1	0.3	0.5	0.8	0.7	0.6	0.6	0.6	1.7
2001	0.5	2.0	*	0.1	0.4	0.5	0.7	0.7	0.5	0.4	0.7	*
2000	0.5	1.2	0.3	0.1	0.4	0.5	0.6	0.7	0.6	0.5	0.7	*
1999	0.5	1.4	*	0.1	0.4	0.5	0.6	0.7	0.6	0.4	0.6	*
All origins 3/, female												
2002	0.3	2.1	0.3	0.1	0.2	0.1	0.3	0.2	0.2	0.2	0.4	0.8
2001	0.2	1.5	0.3	*	0.1	0.1	0.2	0.3	*	*	0.4	*
2000	0.2	1.9	*	*	0.1	0.2	0.3	0.2	0.2	0.2	0.3	0.9
1999	0.2	*	*	*	*	0.2	0.2	0.2	*	*	0.3	0.8
Hispanic, both sexes												
2002	0.3	*	*	*	*	0.3	0.4	0.6	*	*	*	*
2001	0.3	*	*	*	*	*	0.6	*	*	*	*	*
2000	0.2	*	*	*	*	*	0.4	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.5	*	*	*	*	*	0.7	1.1	*	*	*	*
2001	0.4	*	*	*	*	*	1.0	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.4	2.2	0.3	0.1	0.3	0.3	0.5	0.5	0.4	0.4	0.5	1.1
2001	0.3	2.0	0.3	0.1	0.2	0.3	0.4	0.5	0.3	0.3	0.5	0.6
2000	0.4	1.7	0.2	0.1	0.3	0.4	0.4	0.5	0.4	0.3	0.4	1.0
1999	0.3	1.4	0.2	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	0.9
Non-Hispanic 4/, male												
2002	0.5	2.2	*	0.1	0.3	0.5	0.8	0.7	0.6	0.6	0.7	1.7
2001	0.5	2.2	*	0.1	0.4	0.5	0.6	0.6	0.4	0.4	0.7	*
2000	0.5	1.4	*	0.2	0.4	0.5	0.5	0.7	0.6	0.5	0.6	*
1999	0.5	1.6	*	0.1	0.4	0.5	0.6	0.7	0.5	0.4	0.6	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified events of undetermined intent and their sequelae (Y20-Y21,Y25-Y29,Y31-Y34,Y87.2,Y89.9)												
Non-Hispanic 4/, female												
2002	0.3	2.2	*	0.1	0.2	0.2	0.3	0.2	0.2	0.2	0.4	0.9
2001	0.2	1.7	*	*	0.1	0.1	0.2	0.3	*	*	0.4	*
2000	0.2	2.0	*	*	0.1	0.2	0.3	0.3	0.2	*	0.3	0.9
1999	0.2	*	*	*	*	0.2	0.2	0.2	*	*	0.3	0.8
Non-Hispanic white, both sexes												
2002	0.4	1.6	0.3	0.1	0.3	0.3	0.5	0.4	0.3	0.3	0.5	1.2
2001	0.3	1.3	0.2	0.1	0.2	0.3	0.4	0.4	0.2	0.2	0.5	0.7
2000	0.3	1.4	*	0.1	0.3	0.3	0.4	0.4	0.3	0.3	0.4	0.8
1999	0.3	1.2	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	0.9
Non-Hispanic white, male												
2002	0.5	1.7	*	*	0.3	0.4	0.7	0.5	0.5	0.4	0.6	1.8
2001	0.4	*	*	*	0.3	0.4	0.6	0.6	0.4	0.3	0.7	*
2000	0.4	*	*	0.2	0.4	0.4	0.4	0.5	0.5	0.4	0.6	*
1999	0.4	*	*	*	0.4	0.4	0.5	0.6	0.5	0.4	0.5	*
Non-Hispanic white, female												
2002	0.3	*	*	*	0.2	0.2	0.3	0.2	0.2	*	0.4	0.9
2001	0.2	*	*	*	*	*	0.2	0.3	*	*	0.4	*
2000	0.2	*	*	*	*	0.2	0.3	0.3	*	*	*	0.8
1999	0.2	*	*	*	*	0.2	0.2	0.2	*	*	0.3	*
Non-Hispanic black, both sexes												
2002	0.6	4.2	*	*	*	0.5	0.9	1.0	*	*	*	*
2001	0.5	5.2	*	*	*	*	0.7	0.8	*	*	*	*
2000	0.6	*	*	*	*	0.5	0.6	0.9	0.9	*	*	*
1999	0.5	*	*	*	0.4	0.5	0.8	1.0	*	*	*	*
Non-Hispanic black, male												
2002	1.0	*	*	*	*	0.9	1.6	1.8	*	*	*	*
2001	0.8	*	*	*	*	*	1.1	1.3	*	*	*	*
2000	0.9	*	*	*	*	0.9	1.0	1.6	*	*	*	*
1999	0.8	*	*	*	*	*	1.4	1.9	*	*	*	*
Non-Hispanic black, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
All origins 3/, male												
2002	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.4	0.5	0.4	0.3	*	*	*	*
1999	0.3	*	*	*	0.3	0.7	0.5	0.3	0.2	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2001	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	0.4	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	0.6	0.5	*	*	*	*	*	*
2001	0.3	*	*	*	0.6	0.6	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	0.7	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.2	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000	0.2	*	*	*	0.3	0.6	0.4	0.3	*	*	*	*
1999	0.3	*	*	*	0.3	0.7	0.5	0.3	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.2	*	*	*	0.2	0.4	0.4	0.2	*	*	*	*
2001	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2000	0.2	*	*	*	0.2	0.4	0.3	0.3	*	*	*	*
1999	0.2	*	*	*	0.2	0.5	0.4	0.2	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	0.6	0.5	0.5	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.4	0.4	*	*	*	*	*
2000	0.3	*	*	*	0.5	0.6	0.6	*	*	*	*	*
1999	0.4	*	*	*	0.6	0.8	0.7	*	*	*	*	*
Non-Hispanic black, male												
2002	0.6	*	*	*	1.2	1.1	1.0	*	*	*	*	*
2001	0.5	*	*	*	0.8	0.8	0.8	*	*	*	*	*
2000	0.6	*	*	*	1.0	1.3	1.3	*	*	*	*	*
1999	0.8	*	*	*	1.1	1.7	1.4	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal execution (Y35.5)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.0	*	*	*	*	*	0.1	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.0	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.0	*	*	*	*	*	0.1	*	*	*	*	*
Non-Hispanic 4/, male												
2002	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.0	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.2	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal execution (Y35.5)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)												
All origins 3/, both sexes												
2002	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
All origins 3/, male												
2002	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
2001	0.2	*	*	*	0.4	0.5	0.4	0.2	*	*	*	*
2000	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
1999	0.2	*	*	*	0.3	0.6	0.3	0.2	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2001	0.2	*	*	*	0.3	0.3	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	0.3	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	0.6	*	*	*	*	*	*	*
2001	0.3	*	*	*	0.6	0.6	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	0.6	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
Non-Hispanic 4/, male												
2002	0.2	*	*	*	0.4	0.4	0.3	0.2	*	*	*	*
2001	0.2	*	*	*	0.4	0.4	0.4	0.2	*	*	*	*
2000	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
1999	0.2	*	*	*	0.3	0.6	0.3	0.2	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other legal intervention and their sequelae (Y35.0-Y35.4,Y35.6-Y35.7,Y89.0)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	0.1	0.2	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	*	0.2	0.1	0.1	*	*	*	*
Non-Hispanic white, male												
2002	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
2001	0.2	*	*	*	0.3	0.4	0.3	0.2	*	*	*	*
2000	0.2	*	*	*	0.2	0.4	0.2	0.2	*	*	*	*
1999	0.2	*	*	*	*	0.4	0.3	0.2	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.3	*	*	*	0.6	0.4	0.4	*	*	*	*	*
2001	0.2	*	*	*	0.4	*	*	*	*	*	*	*
2000	0.2	*	*	*	0.5	*	0.4	*	*	*	*	*
1999	0.3	*	*	*	0.6	0.7	0.4	*	*	*	*	*
Non-Hispanic black, male												
2002	0.5	*	*	*	1.2	0.9	*	*	*	*	*	*
2001	0.4	*	*	*	0.8	*	*	*	*	*	*	*
2000	0.4	*	*	*	1.0	*	*	*	*	*	*	*
1999	0.6	*	*	*	1.1	1.4	0.8	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
All origins 3/, both sexes												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All origins 3/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Non-Hispanic 4/, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
All origins 3/, both sexes												
2002	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7
All origins 3/, male												
2002	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5
All origins 3/, female												
2002	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2
Hispanic, both sexes												
2002	0.4	*	*	*	*	*	*	*	*	2.4	4.8	*
2001	0.4	*	*	*	*	*	*	0.6	*	2.2	5.2	*
2000	0.4	*	*	*	*	*	*	*	*	3.3	5.1	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	1.1	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.6
2001	1.2	0.7	*	*	0.1	0.2	0.4	0.7	1.6	3.1	6.9	12.9
2000	1.2	*	*	0.1	0.1	0.2	0.3	0.8	1.6	3.5	6.7	13.7
1999	1.1	1.6	0.2	*	0.1	0.2	0.4	0.6	1.3	3.3	6.8	12.8
Non-Hispanic 4/, male												
2002	1.0	*	*	0.1	*	0.3	0.4	0.7	1.4	3.1	6.3	12.9
2001	1.1	*	*	*	*	0.2	0.4	0.8	1.7	3.3	7.7	14.5
2000	1.1	*	*	*	*	0.3	0.4	0.8	1.6	4.0	8.2	16.1
1999	1.0	1.8	*	*	*	0.2	0.4	0.6	1.5	3.5	7.9	16.8

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic 4/, female												
2002	1.2	*	*	*	*	0.2	0.4	0.7	1.3	3.1	6.2	11.1
2001	1.2	*	*	*	*	0.2	0.4	0.7	1.5	3.0	6.4	12.2
2000	1.2	*	*	*	*	0.2	0.3	0.8	1.6	3.1	5.7	12.7
1999	1.2	1.4	*	*	*	0.2	0.4	0.6	1.2	3.1	6.1	11.2
Non-Hispanic white, both sexes												
2002	1.1	*	*	*	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.5
2001	1.2	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.2
2000	1.2	*	*	*	*	0.2	0.3	0.8	1.4	3.3	6.5	13.2
1999	1.1	1.3	*	*	0.1	0.2	0.3	0.5	1.2	3.1	6.5	12.1
Non-Hispanic white, male												
2002	1.0	*	*	*	*	0.3	0.4	0.6	1.3	2.9	6.3	12.4
2001	1.0	*	*	*	*	0.2	0.3	0.6	1.4	2.9	7.8	13.5
2000	1.2	*	*	*	*	0.2	0.4	0.8	1.4	3.7	8.2	15.9
1999	1.1	*	*	*	*	0.2	0.3	0.5	1.3	3.3	7.7	15.8
Non-Hispanic white, female												
2002	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	6.0	11.1
2001	1.3	*	*	*	*	0.2	0.3	0.6	1.4	2.7	6.2	11.6
2000	1.2	*	*	*	*	0.2	0.3	0.7	1.4	3.0	5.4	12.0
1999	1.2	*	*	*	*	*	0.3	0.6	1.1	3.0	5.7	10.7
Non-Hispanic black, both sexes												
2002	1.2	*	*	*	*	*	0.9	1.7	2.7	4.7	8.5	14.2
2001	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.4	9.3	23.7
2000	1.4	*	*	*	*	0.4	0.6	1.4	3.6	6.2	10.6	21.3
1999	1.3	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.4
Non-Hispanic black, male												
2002	1.1	*	*	*	*	*	*	1.4	2.6	5.4	7.3	*
2001	1.5	*	*	*	*	*	1.0	1.9	4.3	7.9	9.0	30.0
2000	1.2	*	*	*	*	*	*	1.3	3.2	7.8	10.2	*
1999	1.2	*	*	*	*	*	*	1.4	2.9	6.0	11.9	23.8
Non-Hispanic black, female												
2002	1.4	*	*	*	*	*	1.1	1.9	2.8	4.2	9.1	11.7
2001	1.4	*	*	*	*	*	0.7	1.4	2.7	5.3	9.4	21.4
2000	1.6	*	*	*	*	*	*	1.5	3.9	5.2	10.8	22.2
1999	1.4	*	*	*	*	*	*	1.2	2.4	4.7	9.8	19.1

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.3	0.4	0.5
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.9
2000	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.5	0.8
1999	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.3	*
All origins 3/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2001	0.1	*	*	*	*	0.1	0.1	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	0.1	0.3	0.4	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.4	1.1
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.3	0.4	0.5
2001	0.1	*	*	*	*	0.1	0.1	0.1	0.1	0.2	0.4	0.9
2000	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.5	0.8
1999	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.3	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drugs, medicaments and biological substances causing adverse effects in therapeutic use, and their sequelae (Y40-Y59,Y88.0)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.4	1.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.4	0.5
2001	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.4	0.9
2000	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.5	0.8
1999	0.1	*	*	*	*	*	0.1	0.1	0.1	0.2	0.3	*
Non-Hispanic white, male												
2002	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)												
All origins 3/, both sexes												
2002	0.1	*	*	*	*	*	*	0.0	0.1	0.2	0.3	0.5
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.4	0.5
2000	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.6
1999	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	*
All origins 3/, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	*
All origins 3/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	*	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
Hispanic, both sexes												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	0.5
2001	0.1	*	*	*	*	*	*	0.1	*	0.2	0.4	0.5
2000	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	0.6
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
Non-Hispanic 4/, male												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.5	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Misadventures to patients during medical and surgical care and their sequelae (Y60-Y69,Y88.1)												
Non-Hispanic 4/, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	*	*
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.4	*
Non-Hispanic white, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.3	*
2001	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.5
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	0.5
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	*
Non-Hispanic white, male												
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic white, female												
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*
Non-Hispanic black, both sexes												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, male												
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic black, female												
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
All origins 3/, both sexes												
2002	0.8	*	*	0.1	0.1	0.1	0.3	0.5	1.1	2.6	5.5	10.4
2001	0.9	*	*	*	0.1	0.1	0.3	0.6	1.4	2.7	6.1	11.4
2000	0.9	*	*	0.1	0.1	0.1	0.2	0.6	1.3	3.1	5.8	12.2
1999	0.9	1.2	0.1	*	0.1	0.1	0.2	0.5	1.1	2.7	5.9	11.8
All origins 3/, male												
2002	0.7	*	*	*	*	0.1	0.3	0.5	1.1	2.7	5.7	11.7
2001	0.8	*	*	*	*	0.1	0.2	0.6	1.4	2.8	6.8	13.2
2000	0.9	*	*	*	*	0.1	0.2	0.6	1.3	3.6	7.1	14.3
1999	0.8	1.4	*	*	*	0.1	0.2	0.4	1.2	2.9	7.0	15.4
All origins 3/, female												
2002	0.9	*	*	*	*	0.1	0.3	0.6	1.1	2.6	5.4	9.9
2001	1.0	*	*	*	0.1	0.1	0.3	0.5	1.3	2.5	5.6	10.6
2000	1.0	*	*	*	*	0.1	0.2	0.6	1.3	2.7	5.0	11.4
1999	0.9	*	*	*	*	0.1	0.2	0.5	1.0	2.6	5.2	10.4
Hispanic, both sexes												
2002	0.3	*	*	*	*	*	*	*	*	2.0	4.2	*
2001	0.3	*	*	*	*	*	*	*	*	2.0	4.9	*
2000	0.3	*	*	*	*	*	*	*	*	2.8	4.7	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, male												
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Hispanic, female												
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 4/, both sexes												
2002	0.9	*	*	0.1	*	0.1	0.3	0.6	1.1	2.7	5.6	10.7
2001	1.0	*	*	*	0.1	0.1	0.3	0.6	1.4	2.7	6.1	11.5
2000	1.0	*	*	0.1	0.1	0.2	0.3	0.7	1.4	3.1	5.9	12.4
1999	0.9	1.3	*	*	0.1	0.1	0.2	0.5	1.1	2.8	6.1	11.9
Non-Hispanic 4/, male												
2002	0.8	*	*	*	*	0.2	0.3	0.5	1.1	2.7	5.7	12.2
2001	0.9	*	*	*	*	0.1	0.3	0.6	1.4	2.9	6.8	13.2
2000	1.0	*	*	*	*	0.2	0.3	0.7	1.4	3.6	7.1	14.4
1999	0.9	1.5	*	*	*	0.2	0.2	0.4	1.3	3.1	7.1	15.7

Worktable Orig12. Death rates for 358 selected causes, by 10-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)												
Non-Hispanic 4/, female												
2002	1.0	*	*	*	*	*	0.3	0.6	1.1	2.6	5.5	10.0
2001	1.1	*	*	*	*	0.2	0.3	0.5	1.4	2.5	5.6	10.7
2000	1.1	*	*	*	*	0.2	0.3	0.6	1.4	2.7	5.1	11.6
1999	1.0	*	*	*	*	0.1	0.2	0.5	1.0	2.6	5.4	10.4
Non-Hispanic white, both sexes												
2002	1.0	*	*	*	*	0.1	0.3	0.5	1.0	2.6	5.5	10.5
2001	1.0	*	*	*	*	0.1	0.2	0.5	1.2	2.4	6.0	10.7
2000	1.0	*	*	*	*	0.2	0.2	0.6	1.2	2.9	5.6	11.9
1999	1.0	1.1	*	*	*	0.1	0.2	0.4	1.0	2.7	5.8	11.2
Non-Hispanic white, male												
2002	0.9	*	*	*	*	0.2	0.3	0.5	1.1	2.5	5.7	11.7
2001	0.9	*	*	*	*	*	0.2	0.5	1.2	2.6	6.8	12.3
2000	1.0	*	*	*	*	0.2	0.2	0.6	1.2	3.4	7.0	14.0
1999	0.9	*	*	*	*	0.2	0.2	0.4	1.1	2.9	6.9	14.8
Non-Hispanic white, female												
2002	1.0	*	*	*	*	*	0.3	0.5	1.0	2.6	5.3	9.9
2001	1.1	*	*	*	*	0.2	0.3	0.5	1.3	2.3	5.5	10.0
2000	1.1	*	*	*	*	0.2	0.2	0.6	1.2	2.6	4.7	11.0
1999	1.0	*	*	*	*	*	0.2	0.4	0.9	2.6	5.1	9.8
Non-Hispanic black, both sexes												
2002	1.0	*	*	*	*	*	0.7	1.3	2.1	4.1	7.8	13.6
2001	1.2	*	*	*	*	*	0.7	1.2	3.0	5.3	8.0	22.5
2000	1.2	*	*	*	*	*	0.5	1.0	3.0	5.3	9.6	19.4
1999	1.1	*	*	*	*	*	0.5	1.0	2.4	4.2	9.3	19.4
Non-Hispanic black, male												
2002	0.9	*	*	*	*	*	*	1.3	2.2	4.8	7.0	*
2001	1.2	*	*	*	*	*	*	1.4	4.0	6.8	7.8	28.9
2000	1.0	*	*	*	*	*	*	1.1	2.7	6.4	9.6	*
1999	1.0	*	*	*	*	*	*	1.2	2.6	5.3	10.6	*
Non-Hispanic black, female												
2002	1.1	*	*	*	*	*	0.9	1.4	2.0	3.5	8.3	11.3
2001	1.2	*	*	*	*	*	0.7	0.9	2.3	4.3	8.1	20.1
2000	1.3	*	*	*	*	*	*	0.9	3.2	4.5	9.6	19.6
1999	1.1	*	*	*	*	*	*	*	2.2	3.5	8.6	18.2

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
... Category not applicable.
1/ Figures for age not stated included in "All ages" but not distributed among age groups.
2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.
4/ Includes races other than white and black.
5/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.