Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All causes														
All origins														
Both sexes														
2002	2,443,387	28,034	4,858	3,018	4,132	13,812	19,234	17,959	23,396	35,347	55,793	76,065	96,320	115,441
2001	2,416,425	27,568	5,107	3,093	4,002	13,555	18,697	18,211	23,472	36,895	54,779	73,684	94,381	109,523
2000	2,403,351	28,035	4,979	3,253	4,160	13,563	17,744	17,681	22,770	36,140	53,658	70,832	89,509	106,751
1999	2,391,399	27,937	5,249	3,474	4,121	13,778	16,878	17,703	23,363	36,804	52,452	68,120	84,854	104,884
Male														
2002	1,199,264	15,717	2,806	1,702	2,496	9,844	14,572	12,954	15,782	22,626	34,967	47,701	60,021	69,830
2001	1,183,421	15,477	2,899	1,727	2,441	9,766	14,197	13,006	15,751	23,727	34,437	46,567	58,281	65,801
2000	1,177,578	15,718	2,824	1,850	2,551	9,697	13,374	12,619	15,271	23,252	34,045	45,121	55,277	64,425
1999	1,175,460	15,646	2,975	1,964	2,528	9,780	12,634	12,542	15,734	23,681	33,437	43,303	52,356	63,441
Female														
2002	1,244,123	12,317	2,052	1,316	1,636	3,968	4,662	5,005	7,614	12,721	20,826	28,364	36,299	45,611
2001	1,233,004	12,091	2,208	1,366	1,561	3,789	4,500	5,205	7,721	13,168	20,342	27,117	36,100	43,722
2000	1,225,773	12,317	2,155	1,403	1,609	3,866	4,370	5,062	7,499	12,888	19,613	25,711	34,232	42,326
1999	1,215,939	12,291	2,274	1,510	1,593	3,998	4,244	5,161	7,629	13,123	19,015	24,817	32,498	41,443
Hispanic														
Both sexes														
2002	117,135	4,943	933	538	586	2,086	3,169	2,803	3,013	3,644	4,823	5,527	6,218	6,634
2001	113,413	4,680	917	480	558	1,998	3,060	2,863	2,934	3,859	4,582	5,397	5,965	6,114
2000	107,254	4,598	872	510	555	1,953	2,840	2,598	2,911	3,656	4,429	4,990	5,366	5,876
1999	103,740	4,412	882	464	550	1,871	2,634	2,527	2,870	3,624	4,174	4,767	5,109	5,495
Male														
2002	65,703	2,746	546	303	344	1,614	2,568	2,239	2,235	2,514	3,314	3,695	4,024	4,247
2001	63,317	2,601	518	281	319	1,530	2,489	2,223	2,143	2,739	3,156	3,655	3,822	3,760
2000	60,172	2,514	475	277	345	1,528	2,309	2,014	2,186	2,543	3,054	3,357	3,451	3,656
1999	57,991	2,411	481	266	326	1,432	2,115	1,956	2,128	2,601	2,875	3,213	3,284	3,409
Female														
2002	51,432	2,197	387	235	242	472	601	564	778	1,130	1,509	1,832	2,194	2,387
2001	50,096	2,079	399	199	239	468	571	640	791	1,120	1,426	1,742	2,143	2,354
2000	47,082	2,084	397	233	210	425	531	584	725	1,113	1,375	1,633	1,915	2,220
1999	45,749	2,001	401	198	224	439	519	571	742	1,023	1,299	1,554	1,825	2,086
Non-Hispanic[1]														
Both sexes														
2002	2,318,269	22,816	3,904	2,472	3,530	11,671	15,987	15,073	20,256	31,512	50,713	70,178	89,665	108,370
2001	2,295,244	22,629	4,171	2,599	3,434	11,508	15,551	15,252	20,404	32,857	49,906	67,927	87,989	102,952
2000	2,287,846	23,074	4,086	2,725	3,591	11,550	14,824	15,000	19,741	32,283	48,899	65,454	83,687	100,433
1999	2,279,325	23,126	4,343	3,001	3,559	11,861	14,182	15,091	20,380	32,997	47,952	62,981	79,321	98,934
Male														
2002	1,129,090	12,815	2,250	1,394	2,140	8,192	11,937	10,657	13,457	19,976	31,475	43,743	55,690	65,283
2001	1,115,683	12,739	2,371	1,438	2,116	8,197	11,639	10,715	13,510	20,859	31,059	42,656	54,159	61,716
2000	1,112,704	13,007	2,338	1,569	2,201	8,131	11,005	10,542	13,006	20,556	30,748	41,490	51,494	60,462
1999	1,112,718	12,998	2,479	1,692	2,193	8,323	10,474	10,525	13,522	20,947	30,325	39,813	48,754	59,720

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes										
All origins										
Both sexes										
2002	137,901	175,591	247,399	330,140	377,514	345,385	228,587	88,282	18,822	357
2001	134,616	178,232	252,728	333,677	368,252	339,775	221,314	85,954	18,488	422
2000	134,095	181,739	259,470	337,700	362,745	335,813	219,283	84,641	18,434	356
1999	134,095	186,695	265,905	338,349	360,241	330,903	213,991	83,334	17,913	356
Male										
2002	81,533	100,590	136,431	169,625	173,879	134,637	68,817	19,522	2,930	282
2001	79,157	102,110	139,471	171,311	169,431	130,405	65,527	18,732	2,869	331
2000	78,896	103,935	143,473	173,327	166,892	128,877	64,439	18,552	2,874	289
1999	79,283	107,529	147,391	174,853	166,117	126,531	62,511	18,111	2,836	277
Female										
2002	56,368	75,001	110,968	160,515	203,635	210,748	159,770	68,760	15,892	75
2001	55,459	76,122	113,257	162,366	198,821	209,370	155,787	67,222	15,619	91
2000	55,199	77,804	115,997	164,373	195,853	206,936	154,844	66,089	15,560	67
1999	54,812	79,166	118,514	163,496	194,124	204,372	151,480	65,223	15,077	79
Hispanic										
Both sexes										
2002	7,581	9,060	11,416	12,797	12,041	9,591	6,460	2,668	546	58
2001	7,223	8,910	11,262	12,560	11,077	9,601	6,177	2,582	547	67
2000	6,857	8,546	10,874	11,578	10,373	9,170	5,853	2,334	461	54
1999	6,824	8,843	10,358	11,081	9,831	8,953	5,662	2,285	458	66
Male										
2002	4,515	5,214	6,262	6,654	5,637	3,846	2,226	768	137	55
2001	4,266	5,129	6,102	6,535	5,224	3,821	2,051	754	141	58
2000	4,104	4,838	6,004	6,069	4,904	3,617	2,035	725	116	51
1999	4,081	5,026	5,757	5,860	4,457	3,483	1,960	703	107	60
Female										
2002	3,066	3,846	5,154	6,143	6,404	5,745	4,234	1,900	409	3
2001	2,957	3,781	5,160	6,025	5,853	5,780	4,126	1,828	406	9
2000	2,753	3,708	4,870	5,509	5,469	5,553	3,818	1,609	345	3
1999	2,743	3,817	4,601	5,221	5,374	5,470	3,702	1,582	351	6
Non-Hispanic[1]										
Both sexes										
2002	129,815	165,934	235,248	316,397	364,481	334,920	221,569	85,363	18,226	169
2001	126,860	168,796	240,704	320,213	356,299	329,357	214,629	83,144	17,890	173
2000	126,685	172,601	247,768	325,151	351,481	325,765	212,888	82,088	17,932	140
1999	126,725	177,240	254,712	326,279	349,451	321,050	207,750	80,833	17,402	155
Male										
2002	76,683	94,990	129,716	162,434	167,757	130,450	66,440	18,694	2,783	134
2001	74,498	96,644	132,895	164,283	163,796	126,292	63,325	17,924	2,715	137
2000	74,428	98,692	136,956	166,709	161,575	124,927	62,247	17,768	2,746	107
1999	74,820	102,119	141,091	168,453	161,229	122,676	60,373	17,354	2,725	113

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All causes														
Non-Hispanic[1]														
Female														
2002	1,189,179	10,001	1,654	1,078	1,390	3,479	4,050	4,416	6,799	11,536	19,238	26,435	33,975	43,087
2001	1,179,561	9,890	1,800	1,161	1,318	3,311	3,912	4,537	6,894	11,998	18,847	25,271	33,830	41,236
2000	1,175,142	10,067	1,748	1,156	1,390	3,419	3,819	4,458	6,735	11,727	18,151	23,964	32,193	39,971
1999	1,166,607	10,128	1,864	1,309	1,366	3,538	3,708	4,566	6,858	12,050	17,627	23,168	30,567	39,214
Non-Hispanic White														
Both sexes														
2002	1,981,973	13,463	2,502	1,643	2,388	8,552	10,906	9,900	13,858	22,230	37,299	52,044	67,768	85,916
2001	1,962,810	13,311	2,772	1,777	2,374	8,264	10,466	10,130	14,061	23,334	36,212	50,151	66,935	81,505
2000	1,959,919	13,513	2,641	1,850	2,547	8,327	9,934	9,822	13,369	22,912	35,319	48,005	63,959	79,771
1999	1,953,197	13,553	2,819	1,990	2,498	8,490	9,378	10,048	13,938	23,345	34,285	45,974	60,834	78,170
Male														
2002	957,645	7,706	1,431	923	1,442	5,884	8,073	6,988	9,351	14,299	23,750	33,102	42,622	52,169
2001	945,967	7,508	1,592	957	1,471	5,756	7,786	7,168	9,445	15,102	23,111	32,045	41,686	49,212
2000	944,781	7,660	1,541	1,077	1,575	5,726	7,340	6,920	9,014	14,922	22,700	31,133	39,655	48,255
1999	944,913	7,718	1,605	1,115	1,528	5,799	6,878	7,056	9,393	15,144	22,147	29,448	37,703	47,570
Female														
2002	1,024,328	5,757	1,071	720	946	2,668	2,833	2,912	4,507	7,931	13,549	18,942	25,146	33,747
2001	1,016,843	5,803	1,180	820	903	2,508	2,680	2,962	4,616	8,232	13,101	18,106	25,249	32,293
2000	1,015,138	5,853	1,100	773	972	2,601	2,594	2,902	4,355	7,990	12,619	16,872	24,304	31,516
1999	1,008,284	5,835	1,214	875	970	2,691	2,500	2,992	4,545	8,201	12,138	16,526	23,131	30,600
Non-Hispanic Black														
Both sexes														
2002	286,573	8,284	1,163	686	948	2,542	4,362	4,487	5,522	8,102	11,853	16,002	19,209	19,483
2001	284,343	8,277	1,156	685	872	2,662	4,362	4,388	5,471	8,287	12,149	15,657	18,441	18,646
2000	282,676	8,525	1,217	745	875	2,672	4,254	4,489	5,614	8,247	12,032	15,545	17,366	17,965
1999	281,979	8,589	1,283	869	902	2,813	4,167	4,390	5,602	8,563	12,207	15,131	16,246	18,176
Male														
2002	144,802	4,520	681	392	585	1,914	3,334	3,179	3,544	4,906	6,755	9,372	11,467	11,414
2001	143,971	4,651	655	400	541	2,041	3,336	3,056	3,492	4,959	7,012	9,354	10,953	10,845
2000	143,297	4,757	672	425	534	2,012	3,212	3,127	3,522	4,947	7,114	9,225	10,422	10,649
1999	143,883	4,765	734	508	573	2,133	3,153	3,038	3,595	5,104	7,280	9,266	9,760	10,627
Female														
2002	141,771	3,764	482	294	363	628	1,028	1,308	1,978	3,196	5,098	6,630	7,742	8,069
2001	140,372	3,626	501	285	331	621	1,026	1,332	1,979	3,328	5,137	6,303	7,488	7,801
2000	139,379	3,768	545	320	341	660	1,042	1,362	2,092	3,300	4,918	6,320	6,944	7,316
1999	138,096	3,824	549	361	329	680	1,014	1,352	2,007	3,459	4,927	5,865	6,486	7,549
Origin not stated[2]														
Both sexes														
2002	7,983	275	21	8	16	55	78	83	127	191	257	360	437	437
2001	7,768	259	19	14	10	49	86	96	134	179	291	360	427	457

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes										
Non-Hispanic[1]										
Female										
2002	53,132	70,944	105,532	153,963	196,724	204,470	155,129	66,669	15,443	35
2001	52,362	72,152	107,809	155,930	192,503	203,065	151,304	65,220	15,175	36
2000	52,257	73,909	110,812	158,442	189,906	200,838	150,641	64,320	15,186	33
1999	51,905	75,121	113,621	157,826	188,222	198,374	147,377	63,479	14,677	42
Non-Hispanic White										
Both sexes										
2002	104,927	137,191	201,770	278,144	328,287	306,287	203,219	77,698	15,859	122
2001	102,604	139,503	206,886	282,018	321,125	301,010	197,069	75,705	15,485	113
2000	102,577	143,765	213,104	287,117	317,541	297,730	195,687	74,671	15,669	89
1999	102,623	148,130	219,360	288,018	316,326	292,962	191,334	73,705	15,326	91
Male										
2002	62,366	79,363	112,481	143,909	151,854	119,953	60,743	16,825	2,313	98
2001	60,781	80,805	115,353	145,725	148,288	115,799	57,884	16,125	2,276	92
2000	60,776	83,126	118,912	148,070	146,772	114,484	56,864	15,857	2,332	70
1999	61,122	86,299	122,856	149,671	146,857	111,859	55,125	15,589	2,365	66
Female										
2002	42,561	57,828	89,289	134,235	176,433	186,334	142,476	60,873	13,546	24
2001	41,823	58,698	91,533	136,293	172,837	185,211	139,185	59,580	13,209	21
2000	41,801	60,639	94,192	139,047	170,769	183,246	138,823	58,814	13,337	19
1999	41,501	61,831	96,504	138,347	169,469	181,103	136,209	58,116	12,961	25
Non-Hispanic Black										
Both sexes										
2002	21,395	24,508	28,240	32,076	30,107	23,674	15,316	6,521	2,054	39
2001	20,877	25,103	28,700	32,243	29,349	23,592	14,846	6,423	2,107	50
2000	20,902	24,831	29,814	32,219	28,655	23,696	14,579	6,362	2,023	49
1999	20,887	25,217	30,350	32,618	28,016	23,831	13,993	6,229	1,842	58
Male										
2002	12,292	13,292	14,567	15,246	12,891	8,195	4,407	1,441	379	29
2001	11,755	13,617	14,829	15,429	12,654	8,313	4,249	1,415	375	40
2000	11,753	13,355	15,390	15,491	12,224	8,341	4,218	1,523	349	35
1999	11,866	13,720	15,539	15,706	11,846	8,736	4,134	1,451	307	42
Female										
2002	9,103	11,216	13,673	16,830	17,216	15,479	10,909	5,080	1,675	10
2001	9,122	11,486	13,871	16,814	16,695	15,279	10,597	5,008	1,732	10
2000	9,149	11,476	14,424	16,728	16,431	15,355	10,361	4,839	1,674	14
1999	9,021	11,497	14,811	16,912	16,170	15,095	9,859	4,778	1,535	16
Origin not stated[2]										
Both sexes										
2002	505	597	735	946	992	874	558	251	50	130
2001	533	526	762	904	876	817	508	228	51	182

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All causes														
Origin not stated[2]														
Both sexes														
2000	8,251	363	21	18	14	60	80	83	118	201	330	388	456	442
1999	8,334	399	24	9	12	46	62	85	113	183	326	372	424	455
Male														
2002	4,471	156	10	5	12	38	67	58	90	136	178	263	307	300
2001	4,421	137	10	8	6	39	69	68	98	129	222	256	300	325
2000	4,702	197	11	4	5	38	60	63	79	153	243	274	332	307
1999	4,751	237	15	6	9	25	45	61	84	133	237	277	318	312
Female														
2002	3,512	119	11	3	4	17	11	25	37	55	79	97	130	137
2001	3,347	122	9	6	4	10	17	28	36	50	69	104	127	132
2000	3,549	166	10	14	9	22	20	20	39	48	87	114	124	135
1999	3,583	162	9	3	3	21	17	24	29	50	89	95	106	143
Salmonella infections (A01-A02)														
All origins														
Both sexes														
2002	21	3	-	-	-	-	-	-	-	-	2	1	2	-
2001	41	2	-	-	-	-	-	-	2	1	-	1	2	2
2000	28	3	2	-	-	1	-	-	1	-	-	1	3	2
1999	38	3	-	-	-	-	2	-	-	-	2	-	2	1
Male														
2002	10	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	21	-	-	-	-	-	-	-	1	1	-	1	2	2
2000	15	2	-	-	-	1	-	-	1	-	-	1	1	2
1999	21	2	-	-	-	-	2	-	-	-	1	-	2	-
Female														
2002	11	2	-	-	-	-	-	-	-	-	2	1	1	-
2001	20	2	-	-	-	-	-	-	1	-	-	-	-	-
2000	13	1	2	-	-	-	-	-	-	-	-	-	2	-
1999	17	1	-	-	-	-	-	-	-	-	1	-	-	1
Hispanic														
Both sexes														
2002	3	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	2	2	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes										
Origin not stated[2]										
Both sexes										
2000	553	592	828	971	891	878	542	219	41	162
1999	546	612	835	989	959	900	579	216	53	135
Male										
2002	335	386	453	537	485	341	151	60	10	93
2001	393	337	474	493	411	292	151	54	13	136
2000	364	405	513	549	413	333	157	59	12	131
1999	382	384	543	540	431	372	178	54	4	104
Female										
2002	170	211	282	409	507	533	407	191	40	37
2001	140	189	288	411	465	525	357	174	38	46
2000	189	187	315	422	478	545	385	160	29	31
1999	164	228	292	449	528	528	401	162	49	31
Salmonella infections (A01-A02)										
All origins										
Both sexes										
2002	-	3	1	3	5	1	-	-	-	-
2001	1	4	3	5	9	6	3	-	-	-
2000	-	1	6	3	2	1	2	-	-	-
1999	2	2	5	8	4	4	2	1	-	-
Male										
2002	-	1	1	1	4	1	-	-	-	-
2001	-	3	3	3	4	1	-	-	-	-
2000	-	1	4	1	-	-	1	-	-	-
1999	2	2	3	2	3	1	1	-	-	-
Female										
2002	-	2	-	2	1	-	-	-	-	-
2001	1	1	-	2	5	5	3	-	-	-
2000	-	-	2	2	2	1	1	-	-	-
1999	-	-	2	6	1	3	1	1	-	-
Hispanic										
Both sexes										
2002	-	2	-	-	-	-	-	-	-	-
2001	-	1	-	-	2	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	1	1	-	-	-	-	-
Male										
2002	-	1	-	-	-	-	-	-	-	-
2001	-	1	-	-	2	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Salmonella infections (A01-A02)														
Hispanic														
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	18	3	-	-	-	-	-	-	-	-	2	1	1	-
2001	36	2	-	-	-	-	-	-	2	1	-	1	1	1
2000	26	1	2	-	-	1	-	-	1	-	-	1	3	2
1999	34	3	-	-	-	-	2	-	-	-	2	-	2	1
Male														
2002	9	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	16	-	-	-	-	-	-	-	1	1	-	1	1	1
2000	14	1	-	-	-	1	-	-	1	-	-	1	1	2
1999	18	2	-	-	-	-	2	-	-	-	1	-	2	-
Female														
2002	9	2	-	-	-	-	-	-	-	-	2	1	-	-
2001	20	2	-	-	-	-	-	-	1	-	-	-	-	-
2000	12	-	2	-	-	-	-	-	-	-	-	-	2	-
1999	16	1	-	-	-	-	-	-	-	-	1	-	-	1
Non-Hispanic White														
Both sexes														
2002	15	2	-	-	-	-	-	-	-	-	2	1	1	-
2001	28	1	-	-	-	-	-	-	2	-	-	1	1	1
2000	23	-	2	-	-	-	-	-	1	-	-	1	3	2
1999	28	3	-	-	-	-	1	-	-	-	2	-	2	1
Male														
2002	8	1	-	-	-	-	-	-	-	-	-	-	1	-
2001	13	-	-	-	-	-	-	-	1	-	-	1	1	1
2000	12	-	-	-	-	-	-	-	1	-	-	1	1	2
1999	13	2	-	-	-	-	1	-	-	-	1	-	2	-
Female														
2002	7	1	-	-	-	-	-	-	-	-	2	1	-	-
2001	15	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	11	-	2	-	-	-	-	-	-	-	-	-	2	-
1999	15	1	-	-	-	-	-	-	-	-	1	-	-	1
Non-Hispanic Black														
Both sexes														
2002	3	1	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Salmonella infections (A01-A02)										
Hispanic										
Female										
2002	-	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	1	1	3	5	1	-	-	-	-
2001	1	3	3	5	7	6	3	-	-	-
2000	-	1	6	3	2	1	2	-	-	-
1999	1	1	5	7	3	4	2	1	-	-
Male										
2002	-	-	1	1	4	1	-	-	-	-
2001	-	2	3	3	2	1	-	-	-	-
2000	-	1	4	1	-	-	1	-	-	-
1999	1	1	3	2	2	1	1	-	-	-
Female										
2002	-	1	-	2	1	-	-	-	-	-
2001	1	1	-	2	5	5	3	-	-	-
2000	-	-	2	2	2	1	1	-	-	-
1999	-	-	2	5	1	3	1	1	-	-
Non-Hispanic White										
Both sexes										
2002	-	1	-	2	5	1	-	-	-	-
2001	1	1	2	5	5	6	2	-	-	-
2000	-	1	5	3	2	1	2	-	-	-
1999	-	-	4	6	2	4	2	1	-	-
Male										
2002	-	-	-	1	4	1	-	-	-	-
2001	-	1	2	3	2	1	-	-	-	-
2000	-	1	4	1	-	-	1	-	-	-
1999	-	-	2	2	1	1	1	-	-	-
Female										
2002	-	1	-	1	1	-	-	-	-	-
2001	1	-	-	2	3	5	2	-	-	-
2000	-	-	1	2	2	1	1	-	-	-
1999	-	-	2	4	1	3	1	1	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	1	1	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Salmonella infections (A01-A02)														
Non-Hispanic Black														
Both sexes														
2001	5	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	1	-	-	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	1	-	-	-	1	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	1	-	-	-	-	-	-	-
Female														
2002	2	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Shigellosis and amebiasis (A03,A06)														
All origins														
Both sexes														
2002	8	-	1	-	-	1	-	-	-	-	1	2	-	1
2001	7	1	1	1	-	-	-	-	-	-	-	1	2	-
2000	12	-	-	-	-	-	-	-	-	-	1	-	1	-
1999	10	-	1	1	-	-	-	-	-	-	1	-	-	-
Male														
2002	4	-	-	-	-	1	-	-	-	-	1	2	-	-
2001	6	1	1	1	-	-	-	-	-	-	-	1	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Salmonella infections (A01-A02)										
Non-Hispanic Black										
Both sexes										
2001	-	-	1	-	2	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	1	1	-	-	-	-	-
Male										
2002	-	-	1	-	-	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	1	-	-	-	-	-
Female										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	2	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Shigellosis and amebiasis (A03,A06)										
All origins										
Both sexes										
2002	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	1	1	-	2	5	-	-	-	1	-
1999	1	-	1	1	-	1	1	2	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Shigellosis and amebiasis (A03,A06)														
All origins														
Male														
2000	5	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	6	-	1	1	-	-	-	-	-	-	1	-	-	-
Female														
2002	4	-	1	-	-	-	-	-	-	-	-	-	-	1
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	7	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	1	-	-
2001	3	-	1	1	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	1	1	-	-	-	-	-	-	1	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	1	-	-
2001	3	-	1	1	-	-	-	-	-	-	-	1	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	1	1	-	-	-	-	-	-	1	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	7	-	1	-	-	1	-	-	-	-	1	1	-	1
2001	4	1	-	-	-	-	-	-	-	-	-	-	2	-
2000	12	-	-	-	-	-	-	-	-	-	1	-	1	-
1999	6	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	3	-	-	-	-	1	-	-	-	-	1	1	-	-
2001	3	1	-	-	-	-	-	-	-	-	-	-	2	-
2000	5	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	4	-	1	-	-	-	-	-	-	-	-	-	-	1
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	7	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)										
All origins										
Male										
2000	1	-	-	1	2	-	-	-	-	-
1999	1	-	1	-	-	-	1	-	-	-
Female										
2002	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	1	-	1	3	-	-	-	1	-
1999	-	-	-	1	-	1	-	2	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	1	1	-	2	5	-	-	-	1	-
1999	1	-	1	-	-	1	1	2	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	1	2	-	-	-	-	-
1999	1	-	1	-	-	-	1	-	-	-
Female										
2002	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	1	-	1	3	-	-	-	1	-
1999	-	-	-	-	-	1	-	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Shigellosis and amebiasis (A03,A06)														
Non-Hispanic White														
Both sexes														
2002	6	-	-	-	-	1	-	-	-	-	1	1	-	1
2001	4	1	-	-	-	-	-	-	-	-	-	-	2	-
2000	10	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	6	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	3	-	-	-	-	1	-	-	-	-	1	1	-	-
2001	3	1	-	-	-	-	-	-	-	-	-	-	2	-
2000	4	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	3	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	6	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)										
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	1	-	-	2	5	-	-	-	1	-
1999	1	-	1	-	-	1	1	2	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	1	2	-	-	-	-	-
1999	1	-	1	-	-	-	1	-	-	-
Female										
2002	-	-	-	-	1	-	1	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	1	3	-	-	-	1	-
1999	-	-	-	-	-	1	-	2	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Shigellosis and amebiasis (A03,A06)														
Origin not stated[2]														
Male														
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Certain other intestinal infections (A04,A07-A09)														
All origins														
Both sexes														
2002	2,465	13	8	1	2	4	2	3	1	5	13	25	31	63
2001	1,537	10	18	1	3	-	-	1	3	4	8	9	33	28
2000	1,328	13	10	-	-	-	1	2	1	7	12	11	18	33
1999	1,108	12	8	2	1	1	4	3	3	2	5	11	12	15
Male														
2002	947	3	5	1	-	3	-	1	-	2	6	12	17	29
2001	572	7	11	1	1	-	-	-	2	4	4	4	11	14
2000	524	10	6	-	-	-	-	2	1	4	6	7	9	20
1999	420	8	4	2	1	1	2	2	3	1	3	5	6	10
Female														
2002	1,518	10	3	-	2	1	2	2	1	3	7	13	14	34
2001	965	3	7	-	2	-	-	1	1	-	4	5	22	14
2000	804	3	4	-	-	-	1	-	-	3	6	4	9	13
1999	688	4	4	-	-	-	2	1	-	1	2	6	6	5
Hispanic														
Both sexes														
2002	78	5	5	-	-	-	1	-	-	-	1	-	3	-
2001	53	1	1	-	-	-	-	-	-	1	3	-	2	2
2000	46	3	2	-	-	-	-	-	-	1	1	-	-	5
1999	34	3	1	-	-	1	1	1	-	-	-	2	2	2
Male														
2002	31	1	3	-	-	-	-	-	-	-	-	-	2	-
2001	19	1	1	-	-	-	-	-	-	1	2	-	-	1
2000	23	2	1	-	-	-	-	-	-	1	-	-	-	3
1999	17	2	-	-	-	1	1	1	-	-	-	1	1	-
Female														
2002	47	4	2	-	-	-	1	-	-	-	1	-	1	-
2001	34	-	-	-	-	-	-	-	-	-	1	-	2	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Shigellosis and amebiasis (A03,A06)										
Origin not stated[2]										
Male										
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Certain other intestinal infections (A04,A07-A09)										
All origins										
Both sexes										
2002	79	133	229	374	515	509	318	114	23	-
2001	53	73	162	237	329	293	187	75	10	-
2000	37	85	137	202	260	267	168	53	11	-
1999	34	67	109	168	222	211	146	58	14	-
Male										
2002	42	67	118	156	202	160	91	27	5	-
2001	21	32	76	107	120	93	50	14	-	-
2000	19	44	61	88	107	81	48	9	2	-
1999	16	28	46	80	89	54	44	13	2	-
Female										
2002	37	66	111	218	313	349	227	87	18	-
2001	32	41	86	130	209	200	137	61	10	-
2000	18	41	76	114	153	186	120	44	9	-
1999	18	39	63	88	133	157	102	45	12	-
Hispanic										
Both sexes										
2002	6	4	6	15	16	7	6	2	1	-
2001	2	3	5	8	8	8	7	2	-	-
2000	3	5	4	6	5	6	3	1	1	-
1999	3	4	3	3	1	1	4	2	-	-
Male										
2002	4	2	3	6	6	3	-	-	1	-
2001	1	-	1	4	1	3	3	-	-	-
2000	2	3	3	3	1	2	1	-	1	-
1999	2	2	1	2	-	1	1	1	-	-
Female										
2002	2	2	3	9	10	4	6	2	-	-
2001	1	3	4	4	7	5	4	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain other intestinal infections (A04,A07-A09)														
Hispanic														
Female														
2000	23	1	1	-	-	-	-	-	-	-	1	-	-	2
1999	17	1	1	-	-	-	-	-	-	-	-	1	1	2
Non-Hispanic[1]														
Both sexes														
2002	2,381	8	3	1	2	4	1	3	1	5	12	25	27	63
2001	1,474	9	16	1	3	-	-	1	3	3	5	9	31	26
2000	1,277	10	8	-	-	-	1	2	1	6	11	11	18	28
1999	1,071	9	7	2	1	-	3	2	3	2	5	9	10	13
Male														
2002	915	2	2	1	-	3	-	1	-	2	6	12	14	29
2001	550	6	10	1	1	-	-	-	2	3	2	4	11	13
2000	497	8	5	-	-	-	-	2	1	3	6	7	9	17
1999	403	6	4	2	1	-	1	1	3	1	3	4	5	10
Female														
2002	1,466	6	1	-	2	1	1	2	1	3	6	13	13	34
2001	924	3	6	-	2	-	-	1	1	-	3	5	20	13
2000	780	2	3	-	-	-	1	-	-	3	5	4	9	11
1999	668	3	3	-	-	-	2	1	-	1	2	5	5	3
Non-Hispanic White														
Both sexes														
2002	2,205	2	1	1	2	4	1	2	-	4	10	21	24	47
2001	1,357	5	13	1	2	-	-	-	2	2	4	6	29	22
2000	1,175	5	5	-	-	-	1	1	1	2	7	8	14	25
1999	990	5	6	-	1	-	2	2	3	2	2	6	9	11
Male														
2002	840	-	1	1	-	3	-	1	-	1	5	11	13	22
2001	497	3	8	1	-	-	-	-	2	2	1	2	10	11
2000	453	4	4	-	-	-	-	1	1	1	5	5	7	14
1999	373	4	3	-	1	-	1	1	3	1	2	2	4	9
Female														
2002	1,365	2	-	-	2	1	1	1	-	3	5	10	11	25
2001	860	2	5	-	2	-	-	-	-	-	3	4	19	11
2000	722	1	1	-	-	-	1	-	-	1	2	3	7	11
1999	617	1	3	-	-	-	1	1	-	1	-	4	5	2
Non-Hispanic Black														
Both sexes														
2002	155	5	1	-	-	-	-	1	1	1	2	3	2	15
2001	98	3	3	-	1	-	-	1	1	1	1	3	2	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)										
Hispanic										
Female										
2000	1	2	1	3	4	4	2	1	-	-
1999	1	2	2	1	1	-	3	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	72	129	222	358	499	500	312	112	22	-
2001	51	69	156	228	318	284	179	72	10	-
2000	34	80	131	196	253	260	165	52	10	-
1999	31	63	106	165	221	209	142	54	14	-
Male										
2002	38	65	115	150	196	157	91	27	4	-
2001	20	31	74	103	119	90	46	14	-	-
2000	17	41	57	85	104	78	47	9	1	-
1999	14	26	45	78	89	53	43	12	2	-
Female										
2002	34	64	107	208	303	343	221	85	18	-
2001	31	38	82	125	199	194	133	58	10	-
2000	17	39	74	111	149	182	118	43	9	-
1999	17	37	61	87	132	156	99	42	12	-
Non-Hispanic White										
Both sexes										
2002	57	120	199	336	478	473	297	107	19	-
2001	40	56	146	214	295	275	170	65	10	-
2000	29	71	120	184	246	240	159	48	9	-
1999	25	58	97	152	208	202	135	51	13	-
Male										
2002	30	59	100	139	189	149	85	27	4	-
2001	18	26	68	95	108	85	45	12	-	-
2000	15	37	52	80	100	73	45	8	1	-
1999	11	25	44	73	85	52	40	11	1	-
Female										
2002	27	61	99	197	289	324	212	80	15	-
2001	22	30	78	119	187	190	125	53	10	-
2000	14	34	68	104	146	167	114	40	8	-
1999	14	33	53	79	123	150	95	40	12	-
Non-Hispanic Black										
Both sexes										
2002	14	9	22	18	17	25	12	4	3	-
2001	9	12	10	10	18	7	7	5	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain other intestinal infections (A04,A07-A09)														
Non-Hispanic Black														
Both sexes														
2000	83	4	2	-	-	-	-	1	-	3	3	3	4	3
1999	74	4	-	2	-	-	1	-	-	-	3	2	1	2
Male														
2002	66	1	1	-	-	-	-	-	-	1	1	1	1	7
2001	42	2	2	-	1	-	-	-	-	1	1	2	1	2
2000	36	3	1	-	-	-	-	1	-	1	1	2	2	3
1999	25	2	-	2	-	-	-	-	-	-	1	1	1	1
Female														
2002	89	4	-	-	-	-	-	1	1	-	1	2	1	8
2001	56	1	1	-	-	-	-	1	1	-	-	1	1	2
2000	47	1	1	-	-	-	-	-	-	2	2	1	2	-
1999	49	2	-	-	-	-	1	-	-	-	2	1	-	1
Origin not stated[2]														
Both sexes														
2002	6	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	10	-	1	-	-	-	-	-	-	-	-	-	-	-
2000	5	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	4	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	5	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	7	-	1	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Tuberculosis (A16-A19)														
All origins														
Both sexes														
2002	784	1	-	-	-	-	9	7	11	18	34	35	46	44
2001	764	2	1	-	-	6	6	11	14	18	22	37	47	63
2000	776	1	6	-	1	2	9	7	13	17	25	38	53	31
1999	930	2	2	1	2	1	10	3	16	25	32	51	55	67
Male														
2002	462	-	-	-	-	-	5	4	7	11	27	20	35	36
2001	453	-	-	-	-	5	3	7	7	12	15	27	36	47

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Certain other intestinal infections (A04,A07-A09)										
Non-Hispanic Black										
Both sexes										
2000	3	8	7	10	6	17	5	4	-	-
1999	4	5	9	13	12	5	7	3	1	-
Male										
2002	8	6	14	8	5	7	5	-	-	-
2001	2	4	6	4	8	3	1	2	-	-
2000	1	3	3	4	4	4	2	1	-	-
1999	1	1	1	5	4	-	3	1	1	-
Female										
2002	6	3	8	10	12	18	7	4	3	-
2001	7	8	4	6	10	4	6	3	-	-
2000	2	5	4	6	2	13	3	3	-	-
1999	3	4	8	8	8	5	4	2	-	-
Origin not stated[2]										
Both sexes										
2002	1	-	1	1	-	2	-	-	-	-
2001	-	1	1	1	3	1	1	1	-	-
2000	-	-	2	-	2	1	-	-	-	-
1999	-	-	-	-	-	1	-	2	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	1	-	-	-	1	-	-	-
2000	-	-	1	-	2	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	1	-	1	1	-	2	-	-	-	-
2001	-	-	-	1	3	1	-	1	-	-
2000	-	-	1	-	-	-	-	-	-	-
1999	-	-	-	-	-	1	-	2	-	-
Tuberculosis (A16-A19)										
All origins										
Both sexes										
2002	63	82	79	114	120	68	42	9	2	-
2001	53	60	92	102	104	85	30	10	-	1
2000	50	70	85	132	120	74	31	10	-	1
1999	63	70	103	134	119	115	41	16	1	1
Male										
2002	39	54	48	56	65	36	16	2	1	-
2001	38	40	58	53	47	34	18	5	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Tuberculosis (A16-A19)														
All origins														
Male														
2000	468	-	1	-	-	2	5	3	11	11	17	35	41	23
1999	584	-	2	-	2	1	3	1	10	20	23	42	45	48
Female														
2002	322	1	-	-	-	-	4	3	4	7	7	15	11	8
2001	311	2	1	-	-	1	3	4	7	6	7	10	11	16
2000	308	1	5	-	1	-	4	4	2	6	8	3	12	8
1999	346	2	-	1	-	-	7	2	6	5	9	9	10	19
Hispanic														
Both sexes														
2002	95	-	-	-	-	-	2	2	2	5	6	3	7	4
2001	97	-	-	-	-	4	2	6	3	4	5	3	6	9
2000	83	-	2	-	1	-	3	4	3	1	2	2	7	4
1999	125	1	-	-	1	1	3	1	2	7	6	10	9	9
Male														
2002	52	-	-	-	-	-	1	2	1	4	5	1	5	2
2001	67	-	-	-	-	4	2	5	2	2	3	3	6	8
2000	46	-	-	-	-	-	1	2	2	1	1	2	3	1
1999	78	-	-	-	1	1	2	1	2	5	4	8	8	3
Female														
2002	43	-	-	-	-	-	1	-	1	1	1	2	2	2
2001	30	-	-	-	-	-	-	1	1	2	2	-	-	1
2000	37	-	2	-	1	-	2	2	1	-	1	-	4	3
1999	47	1	-	-	-	-	1	-	-	2	2	2	1	6
Non-Hispanic[1]														
Both sexes														
2002	682	1	-	-	-	-	6	5	9	11	28	32	38	40
2001	660	2	1	-	-	2	4	5	11	13	17	34	39	52
2000	683	1	4	-	-	2	6	3	10	16	23	34	46	27
1999	788	1	2	1	1	-	6	2	14	18	24	39	43	57
Male														
2002	404	-	-	-	-	-	4	2	6	5	22	19	29	34
2001	379	-	-	-	-	1	1	2	5	9	12	24	28	37
2000	414	-	1	-	-	2	4	1	9	10	16	31	38	22
1999	492	-	2	-	1	-	-	-	8	15	17	32	34	44
Female														
2002	278	1	-	-	-	-	2	3	3	6	6	13	9	6
2001	281	2	1	-	-	1	3	3	6	4	5	10	11	15
2000	269	1	3	-	-	-	2	2	1	6	7	3	8	5
1999	296	1	-	1	-	-	6	2	6	3	7	7	9	13

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)										
All origins										
Male										
2000	36	46	41	69	70	38	13	5	-	1
1999	48	49	67	71	66	58	16	11	-	1
Female										
2002	24	28	31	58	55	32	26	7	1	-
2001	15	20	34	49	57	51	12	5	-	-
2000	14	24	44	63	50	36	18	5	-	-
1999	15	21	36	63	53	57	25	5	1	-
Hispanic										
Both sexes										
2002	13	11	10	6	13	5	5	1	-	-
2001	9	8	10	10	10	4	3	1	-	-
2000	8	5	8	14	9	8	1	1	-	-
1999	11	11	13	17	10	9	2	2	-	-
Male										
2002	4	6	5	4	7	3	2	-	-	-
2001	5	6	8	4	6	3	-	-	-	-
2000	6	4	4	8	6	3	1	1	-	-
1999	8	8	8	9	4	4	2	-	-	-
Female										
2002	9	5	5	2	6	2	3	1	-	-
2001	4	2	2	6	4	1	3	1	-	-
2000	2	1	4	6	3	5	-	-	-	-
1999	3	3	5	8	6	5	-	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	48	70	69	108	107	63	37	8	2	-
2001	44	52	82	92	94	80	27	9	-	-
2000	41	64	76	115	111	66	29	9	-	-
1999	52	59	89	116	106	105	39	14	-	-
Male										
2002	33	47	43	52	58	33	14	2	1	-
2001	33	34	50	49	41	30	18	5	-	-
2000	29	42	36	58	64	35	12	4	-	-
1999	40	41	58	61	61	53	14	11	-	-
Female										
2002	15	23	26	56	49	30	23	6	1	-
2001	11	18	32	43	53	50	9	4	-	-
2000	12	22	40	57	47	31	17	5	-	-
1999	12	18	31	55	45	52	25	3	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Tuberculosis (A16-A19)														
Non-Hispanic White														
Both sexes														
2002	366	-	-	-	-	-	1	1	1	2	8	10	16	16
2001	368	1	1	-	-	-	2	1	1	2	10	17	15	24
2000	353	-	2	-	-	1	2	1	3	7	3	10	24	5
1999	430	-	-	-	-	-	-	1	1	5	6	14	16	27
Male														
2002	218	-	-	-	-	-	1	-	-	-	7	6	14	13
2001	200	-	-	-	-	-	-	1	1	2	8	11	9	19
2000	205	-	-	-	-	1	2	-	3	7	2	9	19	5
1999	263	-	-	-	-	-	-	-	1	4	5	12	15	23
Female														
2002	148	-	-	-	-	-	-	1	1	2	1	4	2	3
2001	168	1	1	-	-	-	2	-	-	-	2	6	6	5
2000	148	-	2	-	-	-	-	1	-	-	1	1	5	-
1999	167	-	-	-	-	-	-	1	-	1	1	2	1	4
Non-Hispanic Black														
Both sexes														
2002	198	-	-	-	-	-	3	4	4	8	15	17	17	18
2001	200	-	-	-	-	-	1	3	6	9	7	14	22	23
2000	238	-	2	-	-	-	2	2	5	8	17	20	21	17
1999	239	1	2	1	-	-	4	1	8	10	14	20	21	22
Male														
2002	123	-	-	-	-	-	2	2	3	4	12	11	12	15
2001	124	-	-	-	-	-	-	1	2	6	4	11	18	16
2000	156	-	1	-	-	-	1	1	4	2	12	18	18	13
1999	154	-	2	-	-	-	-	-	4	8	10	17	16	15
Female														
2002	75	-	-	-	-	-	1	2	1	4	3	6	5	3
2001	76	-	-	-	-	-	1	2	4	3	3	3	4	7
2000	82	-	1	-	-	-	1	1	1	6	5	2	3	4
1999	85	1	-	1	-	-	4	1	4	2	4	3	5	7
Origin not stated[2]														
Both sexes														
2002	7	-	-	-	-	-	1	-	-	2	-	-	1	-
2001	7	-	-	-	-	-	-	-	-	1	-	-	2	2
2000	10	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	17	-	-	-	-	-	1	-	-	-	2	2	3	1
Male														
2002	6	-	-	-	-	-	-	-	-	2	-	-	1	-
2001	7	-	-	-	-	-	-	-	-	1	-	-	2	2
2000	8	-	-	-	-	-	-	-	-	-	-	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)										
Non-Hispanic White										
Both sexes										
2002	25	29	37	72	73	43	25	6	1	-
2001	24	30	42	63	57	54	18	6	-	-
2000	17	32	40	64	67	45	24	6	-	-
1999	26	37	47	68	74	77	22	9	-	-
Male										
2002	18	24	28	33	39	24	9	1	1	-
2001	18	18	21	34	25	18	11	4	-	-
2000	12	22	22	31	32	24	11	3	-	-
1999	18	25	32	33	44	38	7	6	-	-
Female										
2002	7	5	9	39	34	19	16	5	-	-
2001	6	12	21	29	32	36	7	2	-	-
2000	5	10	18	33	35	21	13	3	-	-
1999	8	12	15	35	30	39	15	3	-	-
Non-Hispanic Black										
Both sexes										
2002	13	25	18	17	20	10	9	-	-	-
2001	14	19	25	16	19	11	8	3	-	-
2000	16	25	27	27	31	13	2	3	-	-
1999	17	15	27	31	19	14	8	4	-	-
Male										
2002	8	16	9	11	11	3	4	-	-	-
2001	10	13	17	7	7	5	6	1	-	-
2000	13	17	11	15	23	6	-	1	-	-
1999	16	11	17	17	8	6	3	4	-	-
Female										
2002	5	9	9	6	9	7	5	-	-	-
2001	4	6	8	9	12	6	2	2	-	-
2000	3	8	16	12	8	7	2	2	-	-
1999	1	4	10	14	11	8	5	-	-	-
Origin not stated[2]										
Both sexes										
2002	2	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	1
2000	1	1	1	3	-	-	1	-	-	1
1999	-	-	1	1	3	1	-	-	1	1
Male										
2002	2	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	1
2000	1	-	1	3	-	-	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Tuberculosis (A16-A19)														
Origin not stated[2]														
Male														
1999	14	-	-	-	-	-	1	-	-	-	2	2	3	1
Female														
2002	1	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Respiratory tuberculosis (A16)														
All origins														
Both sexes														
2002	626	-	-	-	-	-	6	6	8	13	27	28	36	37
2001	600	2	1	-	-	3	4	8	9	14	14	27	34	51
2000	615	-	3	-	-	2	5	5	6	15	20	30	42	24
1999	764	1	1	-	1	-	8	2	12	19	26	38	38	53
Male														
2002	372	-	-	-	-	-	4	4	5	9	22	16	27	30
2001	371	-	-	-	-	3	2	5	5	9	9	20	29	41
2000	393	-	1	-	-	2	2	3	5	11	15	29	35	21
1999	486	-	1	-	1	-	2	-	8	15	18	32	32	39
Female														
2002	254	-	-	-	-	-	2	2	3	4	5	12	9	7
2001	229	2	1	-	-	-	2	3	4	5	5	7	5	10
2000	222	-	2	-	-	-	3	2	1	4	5	1	7	3
1999	278	1	-	-	-	-	6	2	4	4	8	6	6	14
Hispanic														
Both sexes														
2002	78	-	-	-	-	-	1	2	2	5	5	3	5	4
2001	77	-	-	-	-	2	1	5	3	2	3	2	5	8
2000	60	-	1	-	-	-	2	2	1	1	1	-	5	3
1999	93	-	-	-	-	-	2	-	1	5	5	6	4	6
Male														
2002	44	-	-	-	-	-	1	2	1	4	4	1	3	2
2001	56	-	-	-	-	2	1	4	2	1	2	2	5	7
2000	40	-	-	-	-	-	1	2	1	1	1	-	3	1
1999	55	-	-	-	-	-	1	-	1	3	3	5	4	2
Female														
2002	34	-	-	-	-	-	-	-	1	1	1	2	2	2
2001	21	-	-	-	-	-	-	1	1	1	1	-	-	1
2000	20	-	1	-	-	-	1	-	-	-	-	-	2	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)										
Origin not stated[2]										
Male										
1999	-	-	1	1	1	1	-	-	-	1
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	-	1	-	-	-
1999	-	-	-	-	2	-	-	-	1	-
Respiratory tuberculosis (A16)										
All origins										
Both sexes										
2002	47	61	59	87	105	60	37	7	2	-
2001	45	47	68	82	84	71	28	7	-	1
2000	36	52	67	107	102	64	26	8	-	1
1999	56	58	81	112	104	100	36	16	1	1
Male										
2002	30	44	36	40	56	32	15	1	1	-
2001	37	31	45	46	38	29	17	4	-	1
2000	29	38	35	58	58	33	13	4	-	1
1999	44	41	56	60	60	52	13	11	-	1
Female										
2002	17	17	23	47	49	28	22	6	1	-
2001	8	16	23	36	46	42	11	3	-	-
2000	7	14	32	49	44	31	13	4	-	-
1999	12	17	25	52	44	48	23	5	1	-
Hispanic										
Both sexes										
2002	8	9	7	3	13	5	5	1	-	-
2001	7	7	10	7	10	3	2	-	-	-
2000	6	5	7	10	8	6	1	1	-	-
1999	10	7	11	16	8	8	2	2	-	-
Male										
2002	3	6	3	2	7	3	2	-	-	-
2001	5	5	8	4	6	2	-	-	-	-
2000	5	4	4	7	5	3	1	1	-	-
1999	7	5	8	8	3	3	2	-	-	-
Female										
2002	5	3	4	1	6	2	3	1	-	-
2001	2	2	2	3	4	1	2	-	-	-
2000	1	1	3	3	3	3	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Respiratory tuberculosis (A16)														
Hispanic														
Female														
1999	38	-	-	-	-	-	1	-	-	2	2	1	-	4
Non-Hispanic[1]														
Both sexes														
2002	543	-	-	-	-	-	5	4	6	6	22	25	31	33
2001	517	2	1	-	-	1	3	3	6	11	11	25	28	41
2000	546	-	2	-	-	2	3	3	5	14	19	28	37	21
1999	657	1	1	-	1	-	5	2	11	14	19	31	32	46
Male														
2002	323	-	-	-	-	-	3	2	4	3	18	15	24	28
2001	309	-	-	-	-	1	1	1	3	7	7	18	23	32
2000	345	-	1	-	-	2	1	1	4	10	14	27	32	20
1999	419	-	1	-	1	-	-	-	7	12	13	26	26	36
Female														
2002	220	-	-	-	-	-	2	2	2	3	4	10	7	5
2001	208	2	1	-	-	-	2	2	3	4	4	7	5	9
2000	201	-	1	-	-	-	2	2	1	4	5	1	5	1
1999	238	1	-	-	-	-	5	2	4	2	6	5	6	10
Non-Hispanic White														
Both sexes														
2002	296	-	-	-	-	-	1	-	1	2	7	9	12	13
2001	288	1	1	-	-	-	1	-	-	1	7	12	11	20
2000	288	-	-	-	-	1	1	1	1	7	2	8	21	4
1999	371	-	-	-	-	-	-	1	1	3	6	13	14	20
Male														
2002	177	-	-	-	-	-	1	-	-	-	7	5	11	10
2001	162	-	-	-	-	-	-	-	-	1	5	7	9	17
2000	174	-	-	-	-	1	1	-	1	7	2	8	18	4
1999	234	-	-	-	-	-	-	-	1	3	5	11	13	18
Female														
2002	119	-	-	-	-	-	-	-	1	2	-	4	1	3
2001	126	1	1	-	-	-	1	-	-	-	2	5	2	3
2000	114	-	-	-	-	-	-	1	-	-	-	-	3	-
1999	137	-	-	-	-	-	-	1	-	-	1	2	1	2
Non-Hispanic Black														
Both sexes														
2002	160	-	-	-	-	-	2	4	4	3	11	13	16	15
2001	157	-	-	-	-	-	1	3	3	8	4	12	16	18
2000	187	-	2	-	-	-	1	2	4	6	14	16	15	12
1999	194	1	1	-	-	-	4	1	7	9	10	15	14	20

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)										
Hispanic										
Female										
1999	3	2	3	8	5	5	-	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	37	51	52	84	92	55	32	6	2	-
2001	38	40	58	75	74	67	26	7	-	-
2000	29	46	59	94	94	58	25	7	-	-
1999	46	51	69	95	94	91	34	14	-	-
Male										
2002	25	37	33	38	49	29	13	1	1	-
2001	32	26	37	42	32	26	17	4	-	-
2000	23	34	30	48	53	30	12	3	-	-
1999	37	36	47	51	56	48	11	11	-	-
Female										
2002	12	14	19	46	43	26	19	5	1	-
2001	6	14	21	33	42	41	9	3	-	-
2000	6	12	29	46	41	28	13	4	-	-
1999	9	15	22	44	38	43	23	3	-	-
Non-Hispanic White										
Both sexes										
2002	19	24	29	56	58	37	22	5	1	-
2001	21	22	30	52	45	42	17	5	-	-
2000	13	21	31	54	58	39	21	5	-	-
1999	25	31	38	54	65	70	21	9	-	-
Male										
2002	14	21	23	23	30	21	9	1	1	-
2001	18	13	16	30	19	14	10	3	-	-
2000	10	16	18	28	26	21	11	2	-	-
1999	18	22	26	28	39	37	7	6	-	-
Female										
2002	5	3	6	33	28	16	13	4	-	-
2001	3	9	14	22	26	28	7	2	-	-
2000	3	5	13	26	32	18	10	3	-	-
1999	7	9	12	26	26	33	14	3	-	-
Non-Hispanic Black										
Both sexes										
2002	11	19	13	14	20	8	7	-	-	-
2001	12	15	17	13	15	10	8	2	-	-
2000	10	21	22	22	24	12	2	2	-	-
1999	15	13	19	26	17	11	7	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Respiratory tuberculosis (A16)														
Non-Hispanic Black														
Male														
2002	101	-	-	-	-	-	1	2	3	2	8	8	12	13
2001	103	-	-	-	-	-	-	1	1	5	2	10	14	14
2000	129	-	1	-	-	-	-	1	3	2	10	15	13	12
1999	122	-	1	-	-	-	-	-	3	7	6	13	11	13
Female														
2002	59	-	-	-	-	-	1	2	1	1	3	5	4	2
2001	54	-	-	-	-	-	1	2	2	3	2	2	2	4
2000	58	-	1	-	-	-	1	1	1	4	4	1	2	-
1999	72	1	-	-	-	-	4	1	4	2	4	2	3	7
Origin not stated[2]														
Both sexes														
2002	5	-	-	-	-	-	-	-	-	2	-	-	-	-
2001	6	-	-	-	-	-	-	-	-	1	-	-	1	2
2000	9	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	14	-	-	-	-	-	1	-	-	-	2	1	2	1
Male														
2002	5	-	-	-	-	-	-	-	-	2	-	-	-	-
2001	6	-	-	-	-	-	-	-	-	1	-	-	1	2
2000	8	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	12	-	-	-	-	-	1	-	-	-	2	1	2	1
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Other tuberculosis (A17-A19)														
All origins														
Both sexes														
2002	158	1	-	-	-	-	3	1	3	5	7	7	10	7
2001	164	-	-	-	-	3	2	3	5	4	8	10	13	12
2000	161	1	3	-	1	-	4	2	7	2	5	8	11	7
1999	166	1	1	1	1	1	2	1	4	6	6	13	17	14
Male														
2002	90	-	-	-	-	-	1	-	2	2	5	4	8	6
2001	82	-	-	-	-	2	1	2	2	3	6	7	7	6
2000	75	-	-	-	-	-	3	-	6	-	2	6	6	2
1999	98	-	1	-	1	1	1	1	2	5	5	10	13	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Respiratory tuberculosis (A16)										
Non-Hispanic Black										
Male										
2002	7	12	7	10	11	2	3	-	-	-
2001	9	10	13	6	6	5	6	1	-	-
2000	10	16	10	11	19	5	-	1	-	-
1999	15	9	12	14	8	4	2	4	-	-
Female										
2002	4	7	6	4	9	6	4	-	-	-
2001	3	5	4	7	9	5	2	1	-	-
2000	-	5	12	11	5	7	2	1	-	-
1999	-	4	7	12	9	7	5	-	-	-
Origin not stated[2]										
Both sexes										
2002	2	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	1
2000	1	1	1	3	-	-	-	-	-	1
1999	-	-	1	1	2	1	-	-	1	1
Male										
2002	2	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	1
2000	1	-	1	3	-	-	-	-	-	1
1999	-	-	1	1	1	1	-	-	-	1
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	1	-
Other tuberculosis (A17-A19)										
All origins										
Both sexes										
2002	16	21	20	27	15	8	5	2	-	-
2001	8	13	24	20	20	14	2	3	-	-
2000	14	18	18	25	18	10	5	2	-	-
1999	7	12	22	22	15	15	5	-	-	-
Male										
2002	9	10	12	16	9	4	1	1	-	-
2001	1	9	13	7	9	5	1	1	-	-
2000	7	8	6	11	12	5	-	1	-	-
1999	4	8	11	11	6	6	3	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other tuberculosis (A17-A19)														
All origins														
Female														
2002	68	1	-	-	-	-	2	1	1	3	2	3	2	1
2001	82	-	-	-	-	1	1	1	3	1	2	3	6	6
2000	86	1	3	-	1	-	1	2	1	2	3	2	5	5
1999	68	1	-	1	-	-	1	-	2	1	1	3	4	5
Hispanic														
Both sexes														
2002	17	-	-	-	-	-	1	-	-	-	1	-	2	-
2001	20	-	-	-	-	2	1	1	-	2	2	1	1	1
2000	23	-	1	-	1	-	1	2	2	-	1	2	2	1
1999	32	1	-	-	1	1	1	1	1	2	1	4	5	3
Male														
2002	8	-	-	-	-	-	-	-	-	-	1	-	2	-
2001	11	-	-	-	-	2	1	1	-	1	1	1	1	1
2000	6	-	-	-	-	-	-	-	1	-	-	2	-	-
1999	23	-	-	-	1	1	1	1	1	2	1	3	4	1
Female														
2002	9	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	9	-	-	-	-	-	-	-	-	1	1	-	-	-
2000	17	-	1	-	1	-	1	2	1	-	1	-	2	1
1999	9	1	-	-	-	-	-	-	-	-	-	1	1	2
Non-Hispanic[1]														
Both sexes														
2002	139	1	-	-	-	-	1	1	3	5	6	7	7	7
2001	143	-	-	-	-	1	1	2	5	2	6	9	11	11
2000	137	1	2	-	-	-	3	-	5	2	4	6	9	6
1999	131	-	1	1	-	-	1	-	3	4	5	8	11	11
Male														
2002	81	-	-	-	-	-	1	-	2	2	4	4	5	6
2001	70	-	-	-	-	-	-	1	2	2	5	6	5	5
2000	69	-	-	-	-	-	3	-	5	-	2	4	6	2
1999	73	-	1	-	-	-	-	-	1	3	4	6	8	8
Female														
2002	58	1	-	-	-	-	-	1	1	3	2	3	2	1
2001	73	-	-	-	-	1	1	1	3	-	1	3	6	6
2000	68	1	2	-	-	-	-	-	-	2	2	2	3	4
1999	58	-	-	1	-	-	1	-	2	1	1	2	3	3
Non-Hispanic White														
Both sexes														
2002	70	-	-	-	-	-	-	1	-	-	1	1	4	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)										
All origins										
Female										
2002	7	11	8	11	6	4	4	1	-	-
2001	7	4	11	13	11	9	1	2	-	-
2000	7	10	12	14	6	5	5	1	-	-
1999	3	4	11	11	9	9	2	-	-	-
Hispanic										
Both sexes										
2002	5	2	3	3	-	-	-	-	-	-
2001	2	1	-	3	-	1	1	1	-	-
2000	2	-	1	4	1	2	-	-	-	-
1999	1	4	2	1	2	1	-	-	-	-
Male										
2002	1	-	2	2	-	-	-	-	-	-
2001	-	1	-	-	-	1	-	-	-	-
2000	1	-	-	1	1	-	-	-	-	-
1999	1	3	-	1	1	1	-	-	-	-
Female										
2002	4	2	1	1	-	-	-	-	-	-
2001	2	-	-	3	-	-	1	1	-	-
2000	1	-	1	3	-	2	-	-	-	-
1999	-	1	2	-	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	11	19	17	24	15	8	5	2	-	-
2001	6	12	24	17	20	13	1	2	-	-
2000	12	18	17	21	17	8	4	2	-	-
1999	6	8	20	21	12	14	5	-	-	-
Male										
2002	8	10	10	14	9	4	1	1	-	-
2001	1	8	13	7	9	4	1	1	-	-
2000	6	8	6	10	11	5	-	1	-	-
1999	3	5	11	10	5	5	3	-	-	-
Female										
2002	3	9	7	10	6	4	4	1	-	-
2001	5	4	11	10	11	9	-	1	-	-
2000	6	10	11	11	6	3	4	1	-	-
1999	3	3	9	11	7	9	2	-	-	-
Non-Hispanic White										
Both sexes										
2002	6	5	8	16	15	6	3	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other tuberculosis (A17-A19)														
Non-Hispanic White														
Both sexes														
2001	80	-	-	-	-	-	1	1	1	1	3	5	4	4
2000	65	-	2	-	-	-	1	-	2	-	1	2	3	1
1999	59	-	-	-	-	-	-	-	-	2	-	1	2	7
Male														
2002	41	-	-	-	-	-	-	-	-	-	-	1	3	3
2001	38	-	-	-	-	-	-	1	1	1	3	4	-	2
2000	31	-	-	-	-	-	1	-	2	-	-	1	1	1
1999	29	-	-	-	-	-	-	-	-	1	-	1	2	5
Female														
2002	29	-	-	-	-	-	-	1	-	-	1	-	1	-
2001	42	-	-	-	-	-	1	-	-	-	-	1	4	2
2000	34	-	2	-	-	-	-	-	-	-	1	1	2	-
1999	30	-	-	-	-	-	-	-	-	1	-	-	-	2
Non-Hispanic Black														
Both sexes														
2002	38	-	-	-	-	-	1	-	-	5	4	4	1	3
2001	43	-	-	-	-	-	-	-	3	1	3	2	6	5
2000	51	-	-	-	-	-	1	-	1	2	3	4	6	5
1999	45	-	1	1	-	-	-	-	1	1	4	5	7	2
Male														
2002	22	-	-	-	-	-	1	-	-	2	4	3	-	2
2001	21	-	-	-	-	-	-	-	1	1	2	1	4	2
2000	27	-	-	-	-	-	1	-	1	-	2	3	5	1
1999	32	-	1	-	-	-	-	-	1	1	4	4	5	2
Female														
2002	16	-	-	-	-	-	-	-	-	3	-	1	1	1
2001	22	-	-	-	-	-	-	-	2	-	1	1	2	3
2000	24	-	-	-	-	-	-	-	-	2	1	1	1	4
1999	13	-	-	1	-	-	-	-	-	-	-	1	2	-
Origin not stated[2]														
Both sexes														
2002	2	-	-	-	-	-	1	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	1	1	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)										
Non-Hispanic White										
Both sexes										
2001	3	8	12	11	12	12	1	1	-	-
2000	4	11	9	10	9	6	3	1	-	-
1999	1	6	9	14	9	7	1	-	-	-
Male										
2002	4	3	5	10	9	3	-	-	-	-
2001	-	5	5	4	6	4	1	1	-	-
2000	2	6	4	3	6	3	-	1	-	-
1999	-	3	6	5	5	1	-	-	-	-
Female										
2002	2	2	3	6	6	3	3	1	-	-
2001	3	3	7	7	6	8	-	-	-	-
2000	2	5	5	7	3	3	3	-	-	-
1999	1	3	3	9	4	6	1	-	-	-
Non-Hispanic Black										
Both sexes										
2002	2	6	5	3	-	2	2	-	-	-
2001	2	4	8	3	4	1	-	1	-	-
2000	6	4	5	5	7	1	-	1	-	-
1999	2	2	8	5	2	3	1	-	-	-
Male										
2002	1	4	2	1	-	1	1	-	-	-
2001	1	3	4	1	1	-	-	-	-	-
2000	3	1	1	4	4	1	-	-	-	-
1999	1	2	5	3	-	2	1	-	-	-
Female										
2002	1	2	3	2	-	1	1	-	-	-
2001	1	1	4	2	3	1	-	1	-	-
2000	3	3	4	1	3	-	-	1	-	-
1999	1	-	3	2	2	1	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	1	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other tuberculosis (A17-A19)														
Origin not stated[2]														
Female														
2002	1	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Whooping cough (A37)														
All origins														
Both sexes														
2002	18	18	-	-	-	-	-	-	-	-	-	-	-	-
2001	17	16	1	-	-	-	-	-	-	-	-	-	-	-
2000	12	12	-	-	-	-	-	-	-	-	-	-	-	-
1999	7	7	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	6	6	-	-	-	-	-	-	-	-	-	-	-	-
2001	7	6	1	-	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	12	12	-	-	-	-	-	-	-	-	-	-	-	-
2001	10	10	-	-	-	-	-	-	-	-	-	-	-	-
2000	7	7	-	-	-	-	-	-	-	-	-	-	-	-
1999	6	6	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic														
Both sexes														
2002	3	3	-	-	-	-	-	-	-	-	-	-	-	-
2001	10	10	-	-	-	-	-	-	-	-	-	-	-	-
2000	7	7	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	2	2	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	4	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	1	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	6	6	-	-	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other tuberculosis (A17-A19)										
Origin not stated[2]										
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	1	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Whooping cough (A37)										
All origins										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Whooping cough (A37)														
Non-Hispanic[1]														
Both sexes														
2002	15	15	-	-	-	-	-	-	-	-	-	-	-	-
2001	7	6	1	-	-	-	-	-	-	-	-	-	-	-
2000	5	5	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	4	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	4	4	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	2	1	-	-	-	-	-	-	-	-	-	-	-
2000	3	3	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	11	11	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	4	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White														
Both sexes														
2002	12	12	-	-	-	-	-	-	-	-	-	-	-	-
2001	6	5	1	-	-	-	-	-	-	-	-	-	-	-
2000	4	4	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	3	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	3	3	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	2	1	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	9	9	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	3	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	2	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	2	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black														
Both sexes														
2002	2	2	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	1	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)										
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Whooping cough (A37)														
Non-Hispanic Black														
Male														
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	1	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Scarlet fever and erysipelas (A38,A46)														
All origins														
Both sexes														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Whooping cough (A37)										
Non-Hispanic Black										
Male										
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Scarlet fever and erysipelas (A38,A46)										
All origins										
Both sexes										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	1	-	-	1	-	1	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Scarlet fever and erysipelas (A38,A46)														
All origins														
Female														
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White														
Both sexes														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)										
All origins										
Female										
1999	-	1	1	1	-	-	1	-	1	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	1	-	-	1	-	1	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	1	1	-	-	1	-	1	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	-	1	-	-	1	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Scarlet fever and erysipelas (A38,A46)														
Non-Hispanic White														
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)										
Non-Hispanic White										
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	1	-	-	1	-	1	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Scarlet fever and erysipelas (A38,A46)														
Origin not stated[2]														
Female														
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Meningococcal infection (A39)														
All origins														
Both sexes														
2002	161	17	12	6	14	21	12	12	4	4	5	10	9	9
2001	199	16	19	12	14	43	12	6	4	8	5	9	4	11
2000	211	22	19	12	6	34	22	9	6	7	6	13	16	6
1999	227	32	30	13	12	28	11	11	12	12	15	11	6	3
Male														
2002	83	8	9	5	8	11	6	8	2	2	3	5	5	2
2001	94	6	9	6	9	20	7	4	2	3	3	6	2	5
2000	106	14	12	5	2	24	11	3	3	6	3	5	6	1
1999	101	14	19	7	6	13	6	8	4	3	6	1	3	-
Female														
2002	78	9	3	1	6	10	6	4	2	2	2	5	4	7
2001	105	10	10	6	5	23	5	2	2	5	2	3	2	6
2000	105	8	7	7	4	10	11	6	3	1	3	8	10	5
1999	126	18	11	6	6	15	5	3	8	9	9	10	3	3
Hispanic														
Both sexes														
2002	19	7	-	1	-	3	2	2	1	-	1	1	-	-
2001	21	4	5	2	-	2	-	1	1	1	1	1	-	2
2000	34	9	3	2	-	3	6	3	2	-	-	2	1	-
1999	27	7	1	1	2	-	2	2	4	3	2	2	-	-
Male														
2002	9	3	-	1	-	1	1	1	1	-	-	-	-	-
2001	11	-	3	2	-	2	-	1	-	-	1	1	-	-
2000	16	4	2	-	-	2	4	2	-	-	-	1	-	-
1999	11	5	1	1	1	-	1	1	1	-	-	-	-	-
Female														
2002	10	4	-	-	-	2	1	1	-	-	1	1	-	-
2001	10	4	2	-	-	-	-	-	1	1	-	-	-	2
2000	18	5	1	2	-	1	2	1	2	-	-	1	1	-
1999	16	2	-	-	1	-	1	1	3	3	2	2	-	-
Non-Hispanic[1]														
Both sexes														
2002	142	10	12	5	14	18	10	10	3	4	4	9	9	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Scarlet fever and erysipelas (A38,A46)										
Origin not stated[2]										
Female										
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Meningococcal infection (A39)										
All origins										
Both sexes										
2002	4	3	4	3	7	2	-	3	-	-
2001	4	7	8	7	1	6	1	2	-	-
2000	3	3	5	7	8	2	4	1	-	-
1999	1	9	-	7	6	3	4	1	-	-
Male										
2002	2	1	1	1	2	1	-	1	-	-
2001	2	2	3	2	1	1	-	1	-	-
2000	1	-	2	4	3	-	-	1	-	-
1999	-	5	-	2	2	-	1	1	-	-
Female										
2002	2	2	3	2	5	1	-	2	-	-
2001	2	5	5	5	-	5	1	1	-	-
2000	2	3	3	3	5	2	4	-	-	-
1999	1	4	-	5	4	3	3	-	-	-
Hispanic										
Both sexes										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	1	-	-	1	1	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Male										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	3	3	4	3	7	2	-	3	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningococcal infection (A39)														
Non-Hispanic[1]														
Both sexes														
2001	178	12	14	10	14	41	12	5	3	7	4	8	4	9
2000	177	13	16	10	6	31	16	6	4	7	6	11	15	6
1999	198	25	28	12	10	28	9	9	8	9	13	8	6	3
Male														
2002	74	5	9	4	8	10	5	7	1	2	3	5	5	2
2001	83	6	6	4	9	18	7	3	2	3	2	5	2	5
2000	90	10	10	5	2	22	7	1	3	6	3	4	6	1
1999	89	9	17	6	5	13	5	7	3	3	6	1	3	-
Female														
2002	68	5	3	1	6	8	5	3	2	2	1	4	4	7
2001	95	6	8	6	5	23	5	2	1	4	2	3	2	4
2000	87	3	6	5	4	9	9	5	1	1	3	7	9	5
1999	109	16	11	6	5	15	4	2	5	6	7	7	3	3
Non-Hispanic White														
Both sexes														
2002	106	7	8	4	5	14	8	7	3	1	3	8	7	7
2001	138	10	10	6	12	29	9	4	1	5	4	7	4	8
2000	129	10	11	3	5	27	10	5	1	3	4	5	11	5
1999	155	23	24	9	6	20	7	4	4	5	9	7	5	3
Male														
2002	58	5	6	4	4	7	5	6	1	1	2	5	3	2
2001	62	5	5	2	8	13	5	2	1	1	2	4	2	4
2000	60	7	6	2	2	18	4	1	-	2	3	1	3	1
1999	67	9	15	4	3	8	4	3	1	2	4	1	2	-
Female														
2002	48	2	2	-	1	7	3	1	2	-	1	3	4	5
2001	76	5	5	4	4	16	4	2	-	4	2	3	2	4
2000	69	3	5	1	3	9	6	4	1	1	1	4	8	4
1999	88	14	9	5	3	12	3	1	3	3	5	6	3	3
Non-Hispanic Black														
Both sexes														
2002	29	3	3	-	8	3	2	2	-	3	1	1	2	1
2001	36	2	4	4	2	10	3	-	2	1	-	1	-	1
2000	44	3	5	7	1	3	4	1	3	4	2	5	4	1
1999	39	2	4	2	4	7	2	4	4	4	4	1	1	-
Male														
2002	12	-	2	-	4	2	-	-	-	1	1	-	2	-
2001	18	1	1	2	1	4	2	-	1	1	-	1	-	1
2000	27	3	4	3	-	3	2	-	3	4	-	2	3	-
1999	20	-	2	1	2	5	1	3	2	1	2	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)										
Non-Hispanic[1]										
Both sexes										
2001	4	7	8	6	1	6	1	2	-	-
2000	2	3	5	6	7	2	4	1	-	-
1999	1	9	-	6	6	3	4	1	-	-
Male										
2002	1	1	1	1	2	1	-	1	-	-
2001	2	2	3	1	1	1	-	1	-	-
2000	1	-	2	3	3	-	-	1	-	-
1999	-	5	-	2	2	-	1	1	-	-
Female										
2002	2	2	3	2	5	1	-	2	-	-
2001	2	5	5	5	-	5	1	1	-	-
2000	1	3	3	3	4	2	4	-	-	-
1999	1	4	-	4	4	3	3	-	-	-
Non-Hispanic White										
Both sexes										
2002	3	2	4	3	7	2	-	3	-	-
2001	1	6	8	5	1	5	1	2	-	-
2000	2	2	5	6	7	2	4	1	-	-
1999	1	9	-	6	5	3	4	1	-	-
Male										
2002	1	-	1	1	2	1	-	1	-	-
2001	1	2	3	-	1	-	-	1	-	-
2000	1	-	2	3	3	-	-	1	-	-
1999	-	5	-	2	2	-	1	1	-	-
Female										
2002	2	2	3	2	5	1	-	2	-	-
2001	-	4	5	5	-	5	1	1	-	-
2000	1	2	3	3	4	2	4	-	-	-
1999	1	4	-	4	3	3	3	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	3	1	-	1	-	1	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	1	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningococcal infection (A39)														
Non-Hispanic Black														
Female														
2002	17	3	1	-	4	1	2	2	-	2	-	1	-	1
2001	18	1	3	2	1	6	1	-	1	-	-	-	-	-
2000	17	-	1	4	1	-	2	1	-	-	2	3	1	1
1999	19	2	2	1	2	2	1	1	2	3	2	1	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	1	-	-	-	-	-	-	-	-	1	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	1	-	-
Septicemia (A40-A41)														
All origins														
Both sexes														
2002	33,865	296	79	42	53	49	69	119	195	318	538	904	1,170	1,499
2001	32,238	312	108	29	40	57	76	96	185	309	520	837	1,140	1,392
2000	31,224	274	99	38	25	40	60	104	176	324	553	781	1,064	1,260
1999	30,680	283	87	47	30	32	89	106	169	314	498	731	954	1,178
Male														
2002	14,947	158	41	26	26	22	37	63	103	159	282	495	629	767
2001	14,307	174	60	17	22	31	44	50	89	170	281	439	604	728
2000	13,537	139	53	24	14	23	37	62	96	162	289	439	600	657
1999	13,395	175	48	22	14	17	50	53	94	170	290	406	513	613
Female														
2002	18,918	138	38	16	27	27	32	56	92	159	256	409	541	732
2001	17,931	138	48	12	18	26	32	46	96	139	239	398	536	664
2000	17,687	135	46	14	11	17	23	42	80	162	264	342	464	603
1999	17,285	108	39	25	16	15	39	53	75	144	208	325	441	565

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningococcal infection (A39)										
Non-Hispanic Black										
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	2	1	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Septicemia (A40-A41)										
All origins										
Both sexes										
2002	1,861	2,589	3,747	5,195	5,815	5,026	2,988	1,111	199	3
2001	1,719	2,409	3,589	5,000	5,351	4,856	3,001	1,020	192	-
2000	1,639	2,303	3,401	4,745	5,193	4,900	2,949	1,079	216	1
1999	1,536	2,352	3,398	4,686	5,024	4,854	2,984	1,133	195	-
Male										
2002	981	1,292	1,862	2,476	2,552	1,857	834	247	35	3
2001	878	1,227	1,817	2,402	2,360	1,788	869	225	32	-
2000	856	1,182	1,685	2,249	2,233	1,690	812	199	35	1
1999	799	1,218	1,701	2,252	2,174	1,712	817	236	21	-
Female										
2002	880	1,297	1,885	2,719	3,263	3,169	2,154	864	164	-
2001	841	1,182	1,772	2,598	2,991	3,068	2,132	795	160	-
2000	783	1,121	1,716	2,496	2,960	3,210	2,137	880	181	-
1999	737	1,134	1,697	2,434	2,850	3,142	2,167	897	174	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Septicemia (A40-A41)														
Hispanic														
Both sexes														
2002	1,521	65	16	7	7	7	8	11	27	27	40	56	80	71
2001	1,386	50	18	8	10	5	8	9	14	28	44	70	63	57
2000	1,226	62	15	3	4	3	9	17	14	28	51	51	71	61
1999	1,215	43	14	6	2	5	4	7	19	30	44	35	57	63
Male														
2002	744	34	9	6	2	5	3	8	17	17	30	40	54	35
2001	684	29	5	7	7	4	5	5	10	20	23	37	43	32
2000	625	33	9	3	2	2	4	9	12	16	30	36	42	37
1999	605	27	8	3	1	3	2	3	11	20	31	22	34	37
Female														
2002	777	31	7	1	5	2	5	3	10	10	10	16	26	36
2001	702	21	13	1	3	1	3	4	4	8	21	33	20	25
2000	601	29	6	-	2	1	5	8	2	12	21	15	29	24
1999	610	16	6	3	1	2	2	4	8	10	13	13	23	26
Non-Hispanic[1]														
Both sexes														
2002	32,225	227	63	35	46	42	60	107	168	291	496	843	1,088	1,420
2001	30,754	261	90	21	29	52	68	87	170	280	475	766	1,071	1,327
2000	29,864	208	83	35	21	37	51	86	162	294	499	728	985	1,194
1999	29,351	239	73	41	28	27	85	98	150	283	453	695	890	1,108
Male														
2002	14,136	120	32	20	24	17	33	55	86	142	250	451	574	727
2001	13,573	145	55	10	15	27	39	45	78	150	258	402	555	694
2000	12,842	104	44	21	12	21	33	53	84	144	257	402	555	615
1999	12,737	147	40	19	13	14	48	49	83	149	259	384	474	573
Female														
2002	18,089	107	31	15	22	25	27	52	82	149	246	392	514	693
2001	17,181	116	35	11	14	25	29	42	92	130	217	364	516	633
2000	17,022	104	39	14	9	16	18	33	78	150	242	326	430	579
1999	16,614	92	33	22	15	13	37	49	67	134	194	311	416	535
Non-Hispanic White														
Both sexes														
2002	25,545	129	42	24	31	31	41	63	102	184	326	572	702	996
2001	24,389	122	63	17	18	32	37	52	93	162	296	478	703	919
2000	23,517	108	51	25	17	21	27	44	90	175	323	445	647	836
1999	23,023	121	47	29	18	21	50	58	97	165	274	421	594	749
Male														
2002	11,148	67	25	15	19	11	26	37	49	87	162	312	368	524
2001	10,668	71	41	7	11	18	19	28	41	95	171	250	365	470
2000	10,056	55	26	16	9	13	20	25	49	93	169	254	359	437

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)										
Hispanic										
Both sexes										
2002	94	132	209	206	176	144	96	41	1	-
2001	94	129	159	191	155	136	93	35	10	-
2000	84	97	138	157	143	119	70	24	5	-
1999	68	119	146	162	134	131	83	37	6	-
Male										
2002	49	61	100	96	80	52	32	13	1	-
2001	54	69	80	101	69	49	25	9	1	-
2000	43	60	66	79	64	41	24	11	2	-
1999	41	67	67	90	54	46	28	10	-	-
Female										
2002	45	71	109	110	96	92	64	28	-	-
2001	40	60	79	90	86	87	68	26	9	-
2000	41	37	72	78	79	78	46	13	3	-
1999	27	52	79	72	80	85	55	27	6	-
Non-Hispanic[1]										
Both sexes										
2002	1,762	2,442	3,527	4,971	5,622	4,865	2,884	1,066	197	3
2001	1,614	2,275	3,415	4,798	5,179	4,712	2,901	983	180	-
2000	1,545	2,194	3,248	4,568	5,033	4,765	2,868	1,049	210	1
1999	1,462	2,224	3,242	4,505	4,873	4,708	2,887	1,092	188	-
Male										
2002	930	1,221	1,754	2,367	2,463	1,799	801	233	34	3
2001	816	1,154	1,728	2,294	2,281	1,738	843	216	30	-
2000	809	1,113	1,611	2,160	2,156	1,641	787	187	32	1
1999	757	1,147	1,630	2,149	2,112	1,659	785	225	21	-
Female										
2002	832	1,221	1,773	2,604	3,159	3,066	2,083	833	163	-
2001	798	1,121	1,687	2,504	2,898	2,974	2,058	767	150	-
2000	736	1,081	1,637	2,408	2,877	3,124	2,081	862	178	-
1999	705	1,077	1,612	2,356	2,761	3,049	2,102	867	167	-
Non-Hispanic White										
Both sexes										
2002	1,245	1,805	2,778	4,043	4,763	4,168	2,466	883	149	2
2001	1,162	1,704	2,681	3,928	4,424	4,040	2,499	825	134	-
2000	1,083	1,610	2,563	3,690	4,211	4,075	2,468	848	159	1
1999	1,043	1,646	2,494	3,628	4,089	3,974	2,452	917	136	-
Male										
2002	680	903	1,407	1,921	2,085	1,557	676	192	23	2
2001	601	868	1,355	1,892	1,959	1,492	712	184	18	-
2000	578	826	1,305	1,748	1,831	1,401	668	150	23	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Septicemia (A40-A41)														
Non-Hispanic White														
Male														
1999	9,863	79	25	15	9	11	33	33	54	89	154	234	317	381
Female														
2002	14,397	62	17	9	12	20	15	26	53	97	164	260	334	472
2001	13,721	51	22	10	7	14	18	24	52	67	125	228	338	449
2000	13,461	53	25	9	8	8	7	19	41	82	154	191	288	399
1999	13,160	42	22	14	9	10	17	25	43	76	120	187	277	368
Non-Hispanic Black														
Both sexes														
2002	6,074	85	17	7	13	9	15	41	61	96	158	238	355	400
2001	5,829	124	21	2	8	19	22	28	70	109	159	261	334	381
2000	5,863	96	25	8	3	12	21	39	68	109	160	258	314	324
1999	5,772	106	16	7	7	5	27	37	48	109	170	251	267	329
Male														
2002	2,668	46	5	3	4	5	5	15	32	50	80	120	192	191
2001	2,611	67	12	1	2	9	16	13	33	51	77	133	171	211
2000	2,559	48	13	4	2	6	12	27	32	48	77	136	183	165
1999	2,594	60	10	4	3	3	11	14	27	55	99	136	143	173
Female														
2002	3,406	39	12	4	9	4	10	26	29	46	78	118	163	209
2001	3,218	57	9	1	6	10	6	15	37	58	82	128	163	170
2000	3,304	48	12	4	1	6	9	12	36	61	83	122	131	159
1999	3,178	46	6	3	4	2	16	23	21	54	71	115	124	156
Origin not stated[2]														
Both sexes														
2002	119	4	-	-	-	-	1	1	-	-	2	5	2	8
2001	98	1	-	-	1	-	-	-	1	1	1	1	6	8
2000	134	4	1	-	-	-	-	1	-	2	3	2	8	5
1999	114	1	-	-	-	-	-	1	-	1	1	1	7	7
Male														
2002	67	4	-	-	-	-	1	-	-	-	2	4	1	5
2001	50	-	-	-	-	-	-	-	1	-	-	-	6	2
2000	70	2	-	-	-	-	-	-	-	2	2	1	3	5
1999	53	1	-	-	-	-	-	1	-	1	-	-	5	3
Female														
2002	52	-	-	-	-	-	-	1	-	-	-	1	1	3
2001	48	1	-	-	1	-	-	-	-	1	1	1	-	6
2000	64	2	1	-	-	-	-	1	-	-	1	1	5	-
1999	61	-	-	-	-	-	-	-	-	-	1	1	2	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)										
Non-Hispanic White										
Male										
1999	523	847	1,264	1,762	1,776	1,396	659	184	18	-
Female										
2002	565	902	1,371	2,122	2,678	2,611	1,790	691	126	-
2001	561	836	1,326	2,036	2,465	2,548	1,787	641	116	-
2000	505	784	1,258	1,942	2,380	2,674	1,800	698	136	-
1999	520	799	1,230	1,866	2,313	2,578	1,793	733	118	-
Non-Hispanic Black										
Both sexes										
2002	468	581	670	845	788	632	381	166	47	1
2001	423	526	660	797	704	626	364	148	43	-
2000	432	537	636	817	759	631	378	185	51	-
1999	387	520	692	792	721	673	394	162	52	-
Male										
2002	227	286	312	395	334	212	108	35	10	1
2001	200	259	327	365	291	225	110	27	11	-
2000	216	262	283	384	293	216	109	34	9	-
1999	219	274	335	343	305	234	106	37	3	-
Female										
2002	241	295	358	450	454	420	273	131	37	-
2001	223	267	333	432	413	401	254	121	32	-
2000	216	275	353	433	466	415	269	151	42	-
1999	168	246	357	449	416	439	288	125	49	-
Origin not stated[2]										
Both sexes										
2002	5	15	11	18	17	17	8	4	1	-
2001	11	5	15	11	17	8	7	2	2	-
2000	10	12	15	20	17	16	11	6	1	-
1999	6	9	10	19	17	15	14	4	1	-
Male										
2002	2	10	8	13	9	6	1	1	-	-
2001	8	4	9	7	10	1	1	-	1	-
2000	4	9	8	10	13	8	1	1	1	-
1999	1	4	4	13	8	7	4	1	-	-
Female										
2002	3	5	3	5	8	11	7	3	1	-
2001	3	1	6	4	7	7	6	2	1	-
2000	6	3	7	10	4	8	10	5	-	-
1999	5	5	6	6	9	8	10	3	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Syphilis (A50-A53)														
All origins														
Both sexes														
2002	41	1	1	-	-	-	-	-	-	1	1	1	4	2
2001	36	1	-	-	1	-	-	-	1	3	-	2	1	3
2000	41	1	-	-	-	-	-	-	1	-	-	2	4	2
1999	33	2	-	-	-	-	-	-	1	-	1	3	2	1
Male														
2002	21	-	1	-	-	-	-	-	-	1	1	-	4	1
2001	19	1	-	-	1	-	-	-	-	2	-	2	1	2
2000	24	1	-	-	-	-	-	-	1	-	-	1	4	2
1999	13	-	-	-	-	-	-	-	1	-	-	2	1	1
Female														
2002	20	1	-	-	-	-	-	-	-	-	-	1	-	1
2001	17	-	-	-	-	-	-	-	1	1	-	-	-	1
2000	17	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	20	2	-	-	-	-	-	-	-	-	1	1	1	-
Hispanic														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	7	1	-	-	-	-	-	-	-	-	-	-	-	2
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	2	-	-	-	-	-	-	-	-	-	2	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	5	1	-	-	-	-	-	-	-	-	-	-	-	2
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	1	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	2	-	-	-	-	-	-	-	-	-	1	-	-
Non-Hispanic[1]														
Both sexes														
2002	39	1	1	-	-	-	-	-	-	1	1	1	4	2
2001	28	-	-	-	1	-	-	-	1	3	-	2	1	-
2000	39	1	-	-	-	-	-	-	1	-	-	2	4	2
1999	27	-	-	-	-	-	-	-	1	-	1	1	1	1
Male														
2002	20	-	1	-	-	-	-	-	-	1	1	-	4	1
2001	14	-	-	-	1	-	-	-	-	2	-	2	1	-
2000	24	1	-	-	-	-	-	-	1	-	-	1	4	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)										
All origins										
Both sexes										
2002	3	4	2	5	7	6	3	-	-	-
2001	1	3	2	4	9	2	2	1	-	-
2000	1	1	6	7	6	6	3	1	-	-
1999	3	2	4	6	4	2	1	1	-	-
Male										
2002	1	4	1	3	1	2	1	-	-	-
2001	1	1	2	2	3	1	-	-	-	-
2000	1	1	1	3	4	3	2	-	-	-
1999	2	2	3	-	-	1	-	-	-	-
Female										
2002	2	-	1	2	6	4	2	-	-	-
2001	-	2	-	2	6	1	2	1	-	-
2000	-	-	5	4	2	3	1	1	-	-
1999	1	-	1	6	4	1	1	1	-	-
Hispanic										
Both sexes										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	1	2	1	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	1	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	2	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	3	4	2	4	7	5	3	-	-	-
2001	1	3	2	3	7	1	2	1	-	-
2000	1	1	6	5	6	6	3	1	-	-
1999	3	2	4	6	3	2	1	1	-	-
Male										
2002	1	4	1	2	1	2	1	-	-	-
2001	1	1	2	1	3	-	-	-	-	-
2000	1	1	1	3	4	3	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Syphilis (A50-A53)														
Non-Hispanic[1]														
Male														
1999	11	-	-	-	-	-	-	-	1	-	-	1	-	1
Female														
2002	19	1	-	-	-	-	-	-	-	-	-	1	-	1
2001	14	-	-	-	-	-	-	-	1	1	-	-	-	-
2000	15	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	16	-	-	-	-	-	-	-	-	-	1	-	1	-
Non-Hispanic White														
Both sexes														
2002	16	-	-	-	-	-	-	-	-	-	1	-	2	-
2001	14	-	-	-	-	-	-	-	-	1	-	-	1	-
2000	20	-	-	-	-	-	-	-	1	-	-	1	1	-
1999	14	-	-	-	-	-	-	-	1	-	1	1	1	-
Male														
2002	8	-	-	-	-	-	-	-	-	-	1	-	2	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	7	-	-	-	-	-	-	-	1	-	-	-	1	-
1999	6	-	-	-	-	-	-	-	1	-	-	1	-	-
Female														
2002	8	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	9	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	13	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	8	-	-	-	-	-	-	-	-	-	1	-	1	-
Non-Hispanic Black														
Both sexes														
2002	23	1	1	-	-	-	-	-	-	1	-	1	2	2
2001	14	-	-	-	1	-	-	-	1	2	-	2	-	-
2000	19	1	-	-	-	-	-	-	-	-	-	1	3	2
1999	12	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	12	-	1	-	-	-	-	-	-	1	-	-	2	1
2001	9	-	-	-	1	-	-	-	-	2	-	2	-	-
2000	17	1	-	-	-	-	-	-	-	-	-	1	3	2
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	11	1	-	-	-	-	-	-	-	-	-	1	-	1
2001	5	-	-	-	-	-	-	-	1	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	7	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)										
Non-Hispanic[1]										
Male										
1999	2	2	3	-	-	1	-	-	-	-
Female										
2002	2	-	1	2	6	3	2	-	-	-
2001	-	2	-	2	4	1	2	1	-	-
2000	-	-	5	2	2	3	1	1	-	-
1999	1	-	1	6	3	1	1	1	-	-
Non-Hispanic White										
Both sexes										
2002	1	2	1	-	4	3	2	-	-	-
2001	1	1	1	3	4	1	-	1	-	-
2000	-	-	3	2	4	4	3	1	-	-
1999	3	2	1	2	1	-	-	1	-	-
Male										
2002	-	2	1	-	-	2	-	-	-	-
2001	1	-	1	1	1	-	-	-	-	-
2000	-	-	-	-	2	1	2	-	-	-
1999	2	2	-	-	-	-	-	-	-	-
Female										
2002	1	-	-	-	4	1	2	-	-	-
2001	-	1	-	2	3	1	-	1	-	-
2000	-	-	3	2	2	3	1	1	-	-
1999	1	-	1	2	1	-	-	1	-	-
Non-Hispanic Black										
Both sexes										
2002	2	2	1	4	3	2	1	-	-	-
2001	-	2	1	-	3	-	2	-	-	-
2000	1	1	3	3	2	2	-	-	-	-
1999	-	-	3	3	2	2	1	-	-	-
Male										
2002	1	2	-	2	1	-	1	-	-	-
2001	-	1	1	-	2	-	-	-	-	-
2000	1	1	1	3	2	2	-	-	-	-
1999	-	-	3	-	-	1	-	-	-	-
Female										
2002	1	-	1	2	2	2	-	-	-	-
2001	-	1	-	-	1	-	2	-	-	-
2000	-	-	2	-	-	-	-	-	-	-
1999	-	-	-	3	2	1	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Syphilis (A50-A53)														
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	1	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	1	-
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Acute poliomyelitis (A80)														
All origins														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Syphilis (A50-A53)										
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Acute poliomyelitis (A80)										
All origins										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute poliomyelitis (A80)														
Hispanic														
Male														
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute poliomyelitis (A80)										
Hispanic										
Male										
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute poliomyelitis (A80)														
Non-Hispanic White														
Female														
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
All origins														
Both sexes														
2002	9	-	-	-	-	-	1	1	-	-	-	-	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute poliomyelitis (A80)										
Non-Hispanic White										
Female										
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
All origins										
Both sexes										
2002	1	-	1	1	1	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
All origins														
Both sexes														
2001	9	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	4	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	6	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	5	-	-	-	-	-	1	1	-	-	-	-	1	-
2001	3	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	9	-	-	-	-	-	1	1	-	-	-	-	2	-
2001	9	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	4	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	6	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
All origins										
Both sexes										
2001	-	-	5	1	1	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	1	-	-	-	-
Male										
2002	1	-	1	1	-	-	-	-	-	-
2001	-	-	4	1	1	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	1	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1	-	1	1	1	1	-	-	-	-
2001	-	-	5	1	1	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	1	-	-	-	-
Male										
2002	1	-	1	1	-	-	-	-	-	-
2001	-	-	4	1	1	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
Non-Hispanic[1]														
Male														
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	5	-	-	-	-	-	1	1	-	-	-	-	1	-
2001	3	1	-	-	-	-	-	-	-	-	-	-	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White														
Both sexes														
2002	6	-	-	-	-	-	1	1	-	-	-	-	1	-
2001	6	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	4	-	-	-	-	-	1	1	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black														
Both sexes														
2002	3	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	2	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	2	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
Non-Hispanic[1]										
Male										
1999	-	1	1	-	-	1	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	1	1	1	-	-	-	-
2001	-	-	4	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	1	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	4	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	1	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	1	-	1	-	-	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	1	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Arthropod-borne viral encephalitis (A83-A84,A85.2)														
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Measles (B05)														
All origins														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	-	-	-	-	-	1
Hispanic														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Arthropod-borne viral encephalitis (A83-A84,A85.2)										
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Measles (B05)										
All origins										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Measles (B05)														
Hispanic														
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	-	-	-	-	-	1
Non-Hispanic White														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Measles (B05)										
Hispanic										
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Measles (B05)														
Non-Hispanic White														
Female														
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	1
Non-Hispanic Black														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Viral hepatitis (B15-B19)														
All origins														
Both sexes														
2002	5,793	2	3	-	2	12	11	37	57	216	524	1,192	1,139	644

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Measles (B05)										
Non-Hispanic White										
Female										
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Viral hepatitis (B15-B19)										
All origins										
Both sexes										
2002	428	369	393	352	235	136	34	7	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Viral hepatitis (B15-B19)														
All origins														
Both sexes														
2001	5,585	1	-	3	-	9	11	19	55	202	595	1,101	959	586
2000	5,357	6	1	2	-	4	14	25	54	219	560	1,044	850	492
1999	4,853	2	3	1	2	3	11	19	69	204	518	871	635	438
Male														
2002	3,770	1	3	-	1	4	9	20	43	154	365	887	868	448
2001	3,539	1	-	2	-	6	5	7	35	133	420	833	737	406
2000	3,381	2	1	1	-	2	9	13	33	151	389	787	638	338
1999	3,043	2	2	-	2	3	7	9	44	139	381	651	487	298
Female														
2002	2,023	1	-	-	1	8	2	17	14	62	159	305	271	196
2001	2,046	-	-	1	-	3	6	12	20	69	175	268	222	180
2000	1,976	4	-	1	-	2	5	12	21	68	171	257	212	154
1999	1,810	-	1	1	-	-	4	10	25	65	137	220	148	140
Hispanic														
Both sexes														
2002	758	-	-	-	-	2	3	4	13	30	71	142	149	92
2001	791	-	-	2	-	3	3	2	8	35	91	155	111	117
2000	708	3	-	-	-	1	2	2	12	39	87	119	99	67
1999	656	-	1	-	-	-	3	3	13	38	66	99	104	69
Male														
2002	491	-	-	-	-	2	3	4	10	22	53	113	105	65
2001	487	-	-	2	-	1	1	2	1	23	66	119	87	85
2000	456	1	-	-	-	1	1	2	9	30	70	91	74	43
1999	406	-	1	-	-	-	1	-	11	30	54	76	77	43
Female														
2002	267	-	-	-	-	-	-	-	3	8	18	29	44	27
2001	304	-	-	-	-	2	2	-	7	12	25	36	24	32
2000	252	2	-	-	-	-	1	-	3	9	17	28	25	24
1999	250	-	-	-	-	-	2	3	2	8	12	23	27	26
Non-Hispanic[1]														
Both sexes														
2002	5,004	2	3	-	2	10	8	33	44	184	450	1,044	986	543
2001	4,776	1	-	1	-	6	8	17	47	165	504	940	845	465
2000	4,629	3	1	2	-	3	12	23	42	180	472	920	745	423
1999	4,176	2	2	1	2	3	8	16	55	166	447	765	529	366
Male														
2002	3,256	1	3	-	1	2	6	16	33	130	309	768	761	376
2001	3,038	1	-	-	-	5	4	5	34	108	354	709	648	317
2000	2,909	1	1	1	-	1	8	11	24	121	319	692	558	293
1999	2,620	2	1	-	2	3	6	9	32	109	323	570	408	253

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)										
All origins										
Both sexes										
2001	428	407	429	375	238	120	41	5	-	1
2000	412	410	435	401	259	123	35	10	-	1
1999	348	427	454	389	262	145	45	5	2	-
Male										
2002	254	193	180	174	96	56	11	3	-	-
2001	237	207	189	156	107	42	13	2	-	1
2000	239	202	194	198	113	55	11	4	-	1
1999	193	217	209	185	132	61	20	1	-	-
Female										
2002	174	176	213	178	139	80	23	4	-	-
2001	191	200	240	219	131	78	28	3	-	-
2000	173	208	241	203	146	68	24	6	-	-
1999	155	210	245	204	130	84	25	4	2	-
Hispanic										
Both sexes										
2002	72	58	44	43	20	12	1	2	-	-
2001	75	52	61	43	18	10	5	-	-	-
2000	80	62	58	42	24	9	1	1	-	-
1999	58	68	61	42	23	6	2	-	-	-
Male										
2002	43	22	18	20	5	6	-	-	-	-
2001	38	17	18	15	8	3	1	-	-	-
2000	51	28	24	17	8	5	-	1	-	-
1999	32	32	27	14	4	4	-	-	-	-
Female										
2002	29	36	26	23	15	6	1	2	-	-
2001	37	35	43	28	10	7	4	-	-	-
2000	29	34	34	25	16	4	1	-	-	-
1999	26	36	34	28	19	2	2	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	355	309	346	309	214	124	33	5	-	-
2001	352	354	367	332	220	110	36	5	-	1
2000	328	348	376	359	235	113	34	9	-	1
1999	290	358	392	346	239	139	43	5	2	-
Male										
2002	210	169	162	154	91	50	11	3	-	-
2001	198	190	171	141	99	39	12	2	-	1
2000	186	174	169	181	105	49	11	3	-	1
1999	161	184	181	170	128	57	20	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Viral hepatitis (B15-B19)														
Non-Hispanic[1]														
Female														
2002	1,748	1	-	-	1	8	2	17	11	54	141	276	225	167
2001	1,738	-	-	1	-	1	4	12	13	57	150	231	197	148
2000	1,720	2	-	1	-	2	4	12	18	59	153	228	187	130
1999	1,556	-	1	1	-	-	2	7	23	57	124	195	121	113
Non-Hispanic White														
Both sexes														
2002	3,857	1	-	-	2	7	8	17	32	147	349	820	770	388
2001	3,663	1	-	1	-	3	4	10	33	131	391	742	622	330
2000	3,513	1	-	2	-	1	4	15	22	142	359	707	536	286
1999	3,176	1	2	1	1	3	3	8	43	131	333	580	378	254
Male														
2002	2,551	1	-	-	1	-	6	8	25	110	241	613	599	271
2001	2,410	1	-	-	-	3	1	3	25	88	280	570	492	233
2000	2,234	-	-	1	-	1	3	7	12	94	248	545	413	205
1999	2,026	1	1	-	1	3	3	4	25	88	245	423	294	180
Female														
2002	1,306	-	-	-	1	7	2	9	7	37	108	207	171	117
2001	1,253	-	-	1	-	-	3	7	8	43	111	172	130	97
2000	1,279	1	-	1	-	-	1	8	10	48	111	162	123	81
1999	1,150	-	1	1	-	-	-	4	18	43	88	157	84	74
Non-Hispanic Black														
Both sexes														
2002	849	1	1	-	-	2	-	15	7	31	80	180	188	113
2001	819	-	-	-	-	2	4	7	8	27	94	162	187	100
2000	777	2	-	-	-	2	8	7	14	29	88	170	159	90
1999	718	1	-	-	1	-	4	6	11	25	94	160	117	73
Male														
2002	532	-	1	-	-	1	-	7	4	16	51	121	141	82
2001	479	-	-	-	-	1	3	2	5	15	64	112	128	63
2000	490	1	-	-	-	-	5	3	8	21	56	119	115	61
1999	433	1	-	-	1	-	2	3	6	15	64	125	90	44
Female														
2002	317	1	-	-	-	1	-	8	3	15	29	59	47	31
2001	340	-	-	-	-	1	1	5	3	12	30	50	59	37
2000	287	1	-	-	-	2	3	4	6	8	32	51	44	29
1999	285	-	-	-	-	-	2	3	5	10	30	35	27	29
Origin not stated[2]														
Both sexes														
2002	31	-	-	-	-	-	-	-	-	2	3	6	4	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)										
Non-Hispanic[1]										
Female										
2002	145	140	184	155	123	74	22	2	-	-
2001	154	164	196	191	121	71	24	3	-	-
2000	142	174	207	178	130	64	23	6	-	-
1999	129	174	211	176	111	82	23	4	2	-
Non-Hispanic White										
Both sexes										
2002	247	219	270	256	184	109	29	2	-	-
2001	256	263	281	279	188	94	29	5	-	-
2000	234	258	301	308	198	101	29	8	-	1
1999	211	262	305	285	212	119	38	4	2	-
Male										
2002	144	131	132	132	82	45	10	-	-	-
2001	151	153	143	128	90	38	9	2	-	-
2000	134	128	137	158	90	44	11	2	-	1
1999	128	144	151	146	122	49	17	1	-	-
Female										
2002	103	88	138	124	102	64	19	2	-	-
2001	105	110	138	151	98	56	20	3	-	-
2000	100	130	164	150	108	57	18	6	-	-
1999	83	118	154	139	90	70	21	3	2	-
Non-Hispanic Black										
Both sexes										
2002	71	56	42	34	15	7	4	2	-	-
2001	67	46	45	35	18	12	4	-	-	1
2000	48	47	46	32	25	8	2	-	-	-
1999	45	59	50	36	17	16	2	1	-	-
Male										
2002	40	31	17	14	3	-	1	2	-	-
2001	36	18	16	9	4	-	2	-	-	1
2000	25	26	22	13	12	3	-	-	-	-
1999	21	21	17	13	2	7	1	-	-	-
Female										
2002	31	25	25	20	12	7	3	-	-	-
2001	31	28	29	26	14	12	2	-	-	-
2000	23	21	24	19	13	5	2	-	-	-
1999	24	38	33	23	15	9	1	1	-	-
Origin not stated[2]										
Both sexes										
2002	1	2	3	-	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Viral hepatitis (B15-B19)														
Origin not stated[2]														
Both sexes														
2001	18	-	-	-	-	-	-	-	-	2	-	6	3	4
2000	20	-	-	-	-	-	-	-	-	-	1	5	6	2
1999	21	-	-	-	-	-	-	-	1	-	5	7	2	3
Male														
2002	23	-	-	-	-	-	-	-	-	2	3	6	2	7
2001	14	-	-	-	-	-	-	-	-	2	-	5	2	4
2000	16	-	-	-	-	-	-	-	-	-	-	4	6	2
1999	17	-	-	-	-	-	-	-	1	-	4	5	2	2
Female														
2002	8	-	-	-	-	-	-	-	-	-	-	-	2	2
2001	4	-	-	-	-	-	-	-	-	-	-	1	1	-
2000	4	-	-	-	-	-	-	-	-	-	1	1	-	-
1999	4	-	-	-	-	-	-	-	-	-	1	2	-	1
Human immunodeficiency virus (HIV) disease (B20-B24)														
All origins														
Both sexes														
2002	14,095	7	5	4	21	38	140	500	1,339	2,618	3,089	2,680	1,794	885
2001	14,175	10	13	17	21	42	183	584	1,517	2,741	3,126	2,511	1,609	821
2000	14,478	10	14	17	29	35	144	647	1,790	2,896	3,023	2,551	1,591	797
1999	14,802	14	24	38	28	31	167	763	1,966	2,981	3,251	2,440	1,467	777
Male														
2002	10,468	4	1	4	10	17	76	288	899	1,943	2,255	2,039	1,446	710
2001	10,590	7	8	11	12	21	84	354	1,067	2,042	2,318	1,942	1,299	673
2000	10,868	8	6	9	13	19	80	371	1,231	2,172	2,269	2,008	1,288	655
1999	11,234	9	16	20	16	15	80	504	1,410	2,224	2,482	1,936	1,198	646
Female														
2002	3,627	3	4	-	11	21	64	212	440	675	834	641	348	175
2001	3,585	3	5	6	9	21	99	230	450	699	808	569	310	148
2000	3,610	2	8	8	16	16	64	276	559	724	754	543	303	142
1999	3,568	5	8	18	12	16	87	259	556	757	769	504	269	131
Hispanic														
Both sexes														
2002	1,855	4	-	-	3	6	23	80	182	353	426	320	217	123
2001	1,853	1	-	1	2	4	28	80	220	355	404	316	210	111
2000	1,908	-	-	5	4	7	16	98	267	382	380	322	209	96
1999	1,958	3	5	5	4	4	29	115	292	422	433	287	168	89
Male														
2002	1,440	3	-	-	3	1	18	59	142	281	326	251	170	95

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Viral hepatitis (B15-B19)										
Origin not stated[2]										
Both sexes										
2001	1	1	1	-	-	-	-	-	-	-
2000	4	-	1	-	-	1	-	-	-	-
1999	-	1	1	1	-	-	-	-	-	-
Male										
2002	1	2	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	2	-	1	-	-	1	-	-	-	-
1999	-	1	1	1	-	-	-	-	-	-
Female										
2002	-	-	3	-	1	-	-	-	-	-
2001	-	1	1	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Human immunodeficiency virus (HIV) disease (B20-B24)										
All origins										
Both sexes										
2002	462	275	126	71	25	8	3	1	-	4
2001	496	258	126	68	20	5	3	2	-	2
2000	442	250	147	69	16	4	3	2	-	1
1999	372	261	137	52	18	7	5	1	1	1
Male										
2002	373	223	100	53	18	4	-	1	-	4
2001	378	187	105	55	18	4	1	2	-	2
2000	358	199	114	54	11	1	1	-	-	1
1999	302	211	106	39	11	6	1	1	-	1
Female										
2002	89	52	26	18	7	4	3	-	-	-
2001	118	71	21	13	2	1	2	-	-	-
2000	84	51	33	15	5	3	2	2	-	-
1999	70	50	31	13	7	1	4	-	1	-
Hispanic										
Both sexes										
2002	65	28	12	6	3	-	1	-	-	3
2001	58	38	13	9	1	1	-	1	-	-
2000	67	28	13	11	3	-	-	-	-	-
1999	48	35	14	2	2	-	1	-	-	-
Male										
2002	45	25	11	4	3	-	-	-	-	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Human immunodeficiency virus (HIV) disease (B20-B24)														
Hispanic														
Male														
2001	1,437	-	-	1	1	2	19	56	166	273	316	241	179	88
2000	1,493	-	-	3	2	4	14	64	210	309	294	252	174	79
1999	1,525	1	4	2	1	3	17	87	227	338	328	224	130	76
Female														
2002	415	1	-	-	-	5	5	21	40	72	100	69	47	28
2001	416	1	-	-	1	2	9	24	54	82	88	75	31	23
2000	415	-	-	2	2	3	2	34	57	73	86	70	35	17
1999	433	2	1	3	3	1	12	28	65	84	105	63	38	13
Non-Hispanic[1]														
Both sexes														
2002	12,094	3	5	4	18	32	117	417	1,147	2,237	2,634	2,324	1,550	755
2001	12,141	9	13	16	19	38	155	499	1,280	2,363	2,681	2,163	1,370	702
2000	12,389	10	14	12	25	28	128	545	1,508	2,476	2,601	2,200	1,362	688
1999	12,638	11	19	33	23	27	138	640	1,661	2,535	2,770	2,108	1,269	675
Male														
2002	8,911	1	1	4	7	16	58	226	751	1,640	1,909	1,756	1,253	609
2001	9,011	7	8	10	11	19	65	295	892	1,754	1,971	1,671	1,095	579
2000	9,224	8	6	6	11	15	66	304	1,009	1,834	1,944	1,728	1,097	563
1999	9,538	8	12	18	15	12	63	412	1,172	1,867	2,117	1,673	1,042	557
Female														
2002	3,183	2	4	-	11	16	59	191	396	597	725	568	297	146
2001	3,130	2	5	6	8	19	90	204	388	609	710	492	275	123
2000	3,165	2	8	6	14	13	62	241	499	642	657	472	265	125
1999	3,100	3	7	15	8	15	75	228	489	668	653	435	227	118
Non-Hispanic White														
Both sexes														
2002	4,224	-	-	-	3	6	17	105	324	842	948	864	502	302
2001	4,286	2	3	6	2	2	28	110	409	914	997	771	477	295
2000	4,562	1	1	3	10	6	24	139	480	1,007	993	826	499	291
1999	4,744	2	2	7	8	6	28	188	606	1,044	1,038	766	520	252
Male														
2002	3,582	-	-	-	1	4	8	77	258	710	807	739	444	269
2001	3,669	2	-	4	1	-	15	79	331	781	858	674	432	262
2000	3,837	1	-	1	4	4	16	95	380	844	834	720	441	253
1999	4,043	2	1	4	6	4	18	139	504	880	888	666	470	224
Female														
2002	642	-	-	-	2	2	9	28	66	132	141	125	58	33
2001	617	-	3	2	1	2	13	31	78	133	139	97	45	33

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)										
Hispanic										
Male										
2001	41	32	11	8	1	1	-	1	-	-
2000	44	22	12	8	2	-	-	-	-	-
1999	42	28	13	2	2	-	-	-	-	-
Female										
2002	20	3	1	2	-	-	1	-	-	-
2001	17	6	2	1	-	-	-	-	-	-
2000	23	6	1	3	1	-	-	-	-	-
1999	6	7	1	-	-	-	1	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	393	245	114	65	22	8	2	1	-	1
2001	421	215	110	58	19	4	3	1	-	2
2000	370	213	130	57	13	4	3	2	-	-
1999	312	218	121	49	16	7	4	1	1	-
Male										
2002	324	197	89	49	15	4	-	1	-	1
2001	323	150	91	46	17	3	1	1	-	2
2000	309	169	98	46	9	1	1	-	-	-
1999	249	177	91	36	9	6	1	1	-	-
Female										
2002	69	48	25	16	7	4	2	-	-	-
2001	98	65	19	12	2	1	2	-	-	-
2000	61	44	32	11	4	3	2	2	-	-
1999	63	41	30	13	7	1	3	-	1	-
Non-Hispanic White										
Both sexes										
2002	158	78	34	24	9	5	2	1	-	-
2001	128	64	46	22	5	2	2	1	-	-
2000	129	80	42	23	5	2	1	-	-	-
1999	117	83	38	22	10	3	3	-	1	-
Male										
2002	140	68	31	17	5	3	-	1	-	-
2001	108	50	45	19	5	1	1	1	-	-
2000	116	72	31	21	4	-	-	-	-	-
1999	107	75	29	19	5	2	-	-	-	-
Female										
2002	18	10	3	7	4	2	2	-	-	-
2001	20	14	1	3	-	1	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Human immunodeficiency virus (HIV) disease (B20-B24)														
Non-Hispanic White														
Female														
2000	725	-	1	2	6	2	8	44	100	163	159	106	58	38
1999	701	-	1	3	2	2	10	49	102	164	150	100	50	28
Non-Hispanic Black														
Both sexes														
2002	7,714	3	5	4	15	25	99	306	804	1,368	1,643	1,441	1,028	445
2001	7,710	7	10	9	17	34	127	381	853	1,417	1,655	1,372	877	404
2000	7,708	9	12	9	14	22	103	405	1,009	1,447	1,581	1,360	851	388
1999	7,751	9	17	26	15	21	109	446	1,033	1,458	1,696	1,322	735	421
Male														
2002	5,203	1	1	4	6	11	49	145	479	911	1,067	999	792	332
2001	5,227	5	8	5	10	18	50	210	543	947	1,093	981	648	315
2000	5,291	7	6	5	7	11	49	208	614	973	1,087	995	647	303
1999	5,377	6	11	14	9	8	44	268	651	957	1,200	992	560	331
Female														
2002	2,511	2	4	-	9	14	50	161	325	457	576	442	236	113
2001	2,483	2	2	4	7	16	77	171	310	470	562	391	229	89
2000	2,417	2	6	4	7	11	54	197	395	474	494	365	204	85
1999	2,374	3	6	12	6	13	65	178	382	501	496	330	175	90
Origin not stated[2]														
Both sexes														
2002	146	-	-	-	-	-	-	3	10	28	29	36	27	7
2001	181	-	-	-	-	-	-	5	17	23	41	32	29	8
2000	181	-	-	-	-	-	-	4	15	38	42	29	20	13
1999	206	-	-	-	1	-	-	8	13	24	48	45	30	13
Male														
2002	117	-	-	-	-	-	-	3	6	22	20	32	23	6
2001	142	-	-	-	-	-	-	3	9	15	31	30	25	6
2000	151	-	-	-	-	-	-	3	12	29	31	28	17	13
1999	171	-	-	-	-	-	-	5	11	19	37	39	26	13
Female														
2002	29	-	-	-	-	-	-	-	4	6	9	4	4	1
2001	39	-	-	-	-	-	-	2	8	8	10	2	4	2
2000	30	-	-	-	-	-	-	1	3	9	11	1	3	-
1999	35	-	-	-	1	-	-	3	2	5	11	6	4	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Human immunodeficiency virus (HIV) disease (B20-B24)										
Non-Hispanic White										
Female										
2000	13	8	11	2	1	2	1	-	-	-
1999	10	8	9	3	5	1	3	-	1	-
Non-Hispanic Black										
Both sexes										
2002	230	163	77	41	13	3	-	-	-	1
2001	286	146	62	36	13	1	1	-	-	2
2000	237	130	84	33	8	2	2	2	-	-
1999	192	132	81	27	6	3	1	1	-	-
Male										
2002	180	125	57	32	10	1	-	-	-	1
2001	209	99	45	27	11	1	-	-	-	2
2000	189	95	64	24	5	1	1	-	-	-
1999	140	99	61	17	4	3	1	1	-	-
Female										
2002	50	38	20	9	3	2	-	-	-	-
2001	77	47	17	9	2	-	1	-	-	-
2000	48	35	20	9	3	1	1	2	-	-
1999	52	33	20	10	2	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	4	2	-	-	-	-	-	-	-	-
2001	17	5	3	1	-	-	-	-	-	-
2000	5	9	4	1	-	-	-	-	-	1
1999	12	8	2	1	-	-	-	-	-	1
Male										
2002	4	1	-	-	-	-	-	-	-	-
2001	14	5	3	1	-	-	-	-	-	-
2000	5	8	4	-	-	-	-	-	-	1
1999	11	6	2	1	-	-	-	-	-	1
Female										
2002	-	1	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-
2000	-	1	-	1	-	-	-	-	-	-
1999	1	2	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malaria (B50-B54)														
All origins														
Both sexes														
2002	12	-	-	-	-	-	1	-	1	-	1	1	4	2
2001	9	-	-	-	1	-	-	-	-	2	-	2	-	-
2000	3	-	-	-	-	-	1	-	-	-	-	-	1	-
1999	7	1	-	-	-	-	-	2	-	-	1	-	-	1
Male														
2002	8	-	-	-	-	-	-	-	-	-	1	1	3	1
2001	5	-	-	-	1	-	-	-	-	-	-	1	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	4	1	-	-	-	-	-	1	-	-	-	-	-	1
Female														
2002	4	-	-	-	-	-	1	-	1	-	-	-	1	1
2001	4	-	-	-	-	-	-	-	-	2	-	1	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	1	-	-	1	-	-	-
Hispanic														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	12	-	-	-	-	-	1	-	1	-	1	1	4	2
2001	9	-	-	-	1	-	-	-	-	2	-	2	-	-
2000	3	-	-	-	-	-	1	-	-	-	-	-	1	-
1999	7	1	-	-	-	-	-	2	-	-	1	-	-	1
Male														
2002	8	-	-	-	-	-	-	-	-	-	1	1	3	1
2001	5	-	-	-	1	-	-	-	-	-	-	1	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malaria (B50-B54)										
All origins										
Both sexes										
2002	-	1	1	-	-	-	-	-	-	-
2001	2	1	-	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	-	-	-	-	-
Male										
2002	-	1	1	-	-	-	-	-	-	-
2001	1	1	-	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	1	1	-	-	-	-	-	-	-
2001	2	1	-	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	1	-	-	-	-	-	-	-
Male										
2002	-	1	1	-	-	-	-	-	-	-
2001	1	1	-	1	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malaria (B50-B54)														
Non-Hispanic[1]														
Male														
1999	4	1	-	-	-	-	-	1	-	-	-	-	-	1
Female														
2002	4	-	-	-	-	-	1	-	1	-	-	-	1	1
2001	4	-	-	-	-	-	-	-	-	2	-	1	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	1	-	-	1	-	-	-
Non-Hispanic White														
Both sexes														
2002	7	-	-	-	-	-	-	-	-	-	1	-	3	1
2001	5	-	-	-	-	-	-	-	-	2	-	1	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	1	-	-	1	-	-	1
Male														
2002	7	-	-	-	-	-	-	-	-	-	1	-	3	1
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	1	-	-	-	-	-	1
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	2	-	1	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	1	-	-	-
Non-Hispanic Black														
Both sexes														
2002	4	-	-	-	-	-	1	-	1	-	-	1	1	-
2001	3	-	-	-	1	-	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	3	1	-	-	-	-	-	1	-	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	1	-	-
2001	2	-	-	-	1	-	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	1	1	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	3	-	-	-	-	-	1	-	1	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	1	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malaria (B50-B54)										
Non-Hispanic[1]										
Male										
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	1	1	-	-	-	-	-	-	-
2001	-	1	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Male										
2002	-	1	1	-	-	-	-	-	-	-
2001	-	1	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malaria (B50-B54)														
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
All origins														
Both sexes														
2002	6,707	224	61	41	43	42	55	67	102	163	290	403	513	445
2001	5,715	191	64	34	30	40	46	58	89	162	207	307	356	338
2000	5,532	167	71	30	37	44	52	56	85	162	188	277	308	323
1999	7,279	203	81	40	39	43	47	92	150	234	311	392	444	427
Male														
2002	3,622	125	39	16	24	20	32	38	58	97	194	278	348	273
2001	2,931	101	37	14	15	23	28	31	52	93	134	211	215	209
2000	2,861	94	41	16	19	25	29	29	46	105	115	164	194	179
1999	3,655	119	51	20	24	23	26	46	85	149	184	231	247	220
Female														
2002	3,085	99	22	25	19	22	23	29	44	66	96	125	165	172
2001	2,784	90	27	20	15	17	18	27	37	69	73	96	141	129
2000	2,671	73	30	14	18	19	23	27	39	57	73	113	114	144
1999	3,624	84	30	20	15	20	21	46	65	85	127	161	197	207
Hispanic														
Both sexes														
2002	593	41	11	8	9	7	12	18	33	22	45	49	78	52
2001	423	41	7	6	5	6	6	14	19	24	22	40	29	34

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malaria (B50-B54)										
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)										
All origins										
Both sexes										
2002	444	520	703	859	735	580	289	112	16	-
2001	433	537	606	698	632	483	288	98	18	-
2000	384	451	623	739	636	513	265	93	28	-
1999	471	652	791	927	812	633	330	138	22	-
Male										
2002	259	290	373	429	356	230	110	30	3	-
2001	240	275	290	350	306	185	101	21	-	-
2000	204	253	344	367	298	216	97	22	4	-
1999	262	351	409	460	367	253	99	28	1	-
Female										
2002	185	230	330	430	379	350	179	82	13	-
2001	193	262	316	348	326	298	187	77	18	-
2000	180	198	279	372	338	297	168	71	24	-
1999	209	301	382	467	445	380	231	110	21	-
Hispanic										
Both sexes										
2002	33	41	34	33	35	22	7	2	1	-
2001	22	31	36	37	20	11	6	6	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
Hispanic														
Both sexes														
2000	412	23	18	6	7	5	14	12	14	22	28	30	31	30
1999	497	36	13	8	6	5	8	17	22	37	30	33	35	27
Male														
2002	363	27	7	5	6	3	4	9	22	13	35	36	59	34
2001	250	21	4	1	4	4	5	9	13	16	15	26	21	25
2000	237	14	12	4	4	2	10	7	13	21	16	16	17	21
1999	260	22	10	4	3	3	3	10	12	29	20	17	23	9
Female														
2002	230	14	4	3	3	4	8	9	11	9	10	13	19	18
2001	173	20	3	5	1	2	1	5	6	8	7	14	8	9
2000	175	9	6	2	3	3	4	5	1	1	12	14	14	9
1999	237	14	3	4	3	2	5	7	10	8	10	16	12	18
Non-Hispanic[1]														
Both sexes														
2002	6,097	181	49	33	34	35	42	48	69	141	243	354	433	392
2001	5,271	149	57	28	25	34	39	44	70	137	182	264	325	302
2000	5,100	144	53	24	30	37	38	44	71	139	158	247	274	290
1999	6,753	166	68	32	31	38	39	75	127	197	278	356	407	397
Male														
2002	3,250	97	31	11	18	17	27	29	36	84	158	242	288	239
2001	2,666	79	33	13	11	19	23	22	39	76	117	182	192	182
2000	2,613	80	29	12	15	23	19	22	33	84	98	148	174	157
1999	3,379	96	41	16	20	20	23	36	72	120	162	213	222	208
Female														
2002	2,847	84	18	22	16	18	15	19	33	57	85	112	145	153
2001	2,605	70	24	15	14	15	16	22	31	61	65	82	133	120
2000	2,487	64	24	12	15	14	19	22	38	55	60	99	100	133
1999	3,374	70	27	16	11	18	16	39	55	77	116	143	185	189
Non-Hispanic White														
Both sexes														
2002	4,939	113	34	24	25	24	23	28	38	82	161	267	315	285
2001	4,257	80	37	22	16	21	24	25	38	87	124	194	241	227
2000	4,131	79	33	18	22	26	25	22	36	81	108	183	200	232
1999	5,288	101	43	24	18	27	24	39	65	121	166	227	282	300

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)										
Hispanic										
Both sexes										
2000	24	27	35	31	26	19	6	4	-	-
1999	35	47	33	43	25	23	9	5	-	-
Male										
2002	18	24	16	15	15	12	3	-	-	-
2001	13	19	19	17	10	3	2	3	-	-
2000	14	10	18	12	15	10	-	1	-	-
1999	20	24	12	21	8	7	2	1	-	-
Female										
2002	15	17	18	18	20	10	4	2	1	-
2001	9	12	17	20	10	8	4	3	1	-
2000	10	17	17	19	11	9	6	3	-	-
1999	15	23	21	22	17	16	7	4	-	-
Non-Hispanic[1]										
Both sexes										
2002	409	479	669	823	699	557	282	110	15	-
2001	409	505	568	660	612	471	281	92	17	-
2000	359	424	588	704	608	494	258	88	28	-
1999	434	603	755	880	786	610	319	133	22	-
Male										
2002	240	266	357	412	341	217	107	30	3	-
2001	226	256	269	332	296	182	99	18	-	-
2000	190	243	326	352	281	206	96	21	4	-
1999	241	326	396	438	359	246	96	27	1	-
Female										
2002	169	213	312	411	358	340	175	80	12	-
2001	183	249	299	328	316	289	182	74	17	-
2000	169	181	262	352	327	288	162	67	24	-
1999	193	277	359	442	427	364	223	106	21	-
Non-Hispanic White										
Both sexes										
2002	321	403	570	716	628	514	253	102	13	-
2001	324	417	482	577	546	426	254	82	13	-
2000	273	353	498	621	544	451	229	72	25	-
1999	328	488	638	751	687	542	278	121	18	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
Non-Hispanic White														
Male														
2002	2,608	57	21	9	15	13	14	15	18	52	109	185	213	169
2001	2,148	40	21	10	7	10	14	15	24	49	80	137	144	139
2000	2,092	45	16	8	10	14	15	12	16	47	68	111	128	128
1999	2,644	64	27	14	10	13	15	17	38	82	108	145	158	166
Female														
2002	2,331	56	13	15	10	11	9	13	20	30	52	82	102	116
2001	2,109	40	16	12	9	11	10	10	14	38	44	57	97	88
2000	2,039	34	17	10	12	12	10	10	20	34	40	72	72	104
1999	2,644	37	16	10	8	14	9	22	27	39	58	82	124	134
Non-Hispanic Black														
Both sexes														
2002	927	58	13	5	5	9	16	18	28	52	69	79	100	89
2001	815	61	16	6	7	9	14	17	28	41	50	62	68	62
2000	815	60	14	5	5	8	13	16	34	48	47	61	63	52
1999	1,247	57	23	7	10	7	13	31	58	73	96	116	103	83
Male														
2002	516	33	9	1	2	3	10	13	15	27	41	52	65	62
2001	419	36	10	3	3	7	8	7	12	22	34	41	39	34
2000	428	31	10	3	2	6	4	6	16	30	28	35	40	27
1999	617	29	14	2	8	5	7	14	34	36	45	62	52	34
Female														
2002	411	25	4	4	3	6	6	5	13	25	28	27	35	27
2001	396	25	6	3	4	2	6	10	16	19	16	21	29	28
2000	387	29	4	2	3	2	9	10	18	18	19	26	23	25
1999	630	28	9	5	2	2	6	17	24	37	51	54	51	49
Origin not stated[2]														
Both sexes														
2002	17	2	1	-	-	-	1	1	-	-	2	-	2	1
2001	21	1	-	-	-	-	1	-	-	1	3	3	2	2
2000	20	-	-	-	-	2	-	-	-	1	2	-	3	3
1999	29	1	-	-	2	-	-	-	1	-	3	3	2	3
Male														
2002	9	1	1	-	-	-	1	-	-	-	1	-	1	-
2001	15	1	-	-	-	-	-	-	-	1	2	3	2	2
2000	11	-	-	-	-	-	-	-	-	-	1	-	3	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)										
Non-Hispanic White										
Male										
2002	189	230	307	360	305	202	96	27	2	-
2001	181	211	232	294	268	166	90	16	-	-
2000	147	201	282	304	253	187	81	16	3	-
1999	179	266	342	382	299	214	80	25	-	-
Female										
2002	132	173	263	356	323	312	157	75	11	-
2001	143	206	250	283	278	260	164	66	13	-
2000	126	152	216	317	291	264	148	56	22	-
1999	149	222	296	369	388	328	198	96	18	-
Non-Hispanic Black										
Both sexes										
2002	62	50	72	87	43	35	27	8	2	-
2001	67	64	71	56	49	35	20	8	4	-
2000	64	55	74	64	56	33	25	15	3	-
1999	86	96	96	105	81	54	36	12	4	-
Male										
2002	37	26	36	39	19	12	10	3	1	-
2001	36	37	29	22	20	10	7	2	-	-
2000	35	33	37	35	23	10	11	5	1	-
1999	48	47	47	46	48	24	12	2	1	-
Female										
2002	25	24	36	48	24	23	17	5	1	-
2001	31	27	42	34	29	25	13	6	4	-
2000	29	22	37	29	33	23	14	10	2	-
1999	38	49	49	59	33	30	24	10	3	-
Origin not stated[2]										
Both sexes										
2002	2	-	-	3	1	1	-	-	-	-
2001	2	1	2	1	-	1	1	-	-	-
2000	1	-	-	4	2	-	1	1	-	-
1999	2	2	3	4	1	-	2	-	-	-
Male										
2002	1	-	-	2	-	1	-	-	-	-
2001	1	-	2	1	-	-	-	-	-	-
2000	-	-	-	3	2	-	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)														
Origin not stated[2]														
Male														
1999	16	1	-	-	1	-	-	-	1	-	2	1	2	3
Female														
2002	8	1	-	-	-	-	-	1	-	-	1	-	1	1
2001	6	-	-	-	-	-	1	-	-	-	1	-	-	-
2000	9	-	-	-	-	2	-	-	-	1	1	-	-	2
1999	13	-	-	-	1	-	-	-	-	-	1	2	-	-
Malignant neoplasms (C00-C97)														
All origins														
Both sexes														
2002	557,271	74	402	537	535	723	1,007	1,258	2,614	5,150	10,935	19,295	30,342	41,487
2001	553,768	66	420	493	515	732	972	1,389	2,605	5,466	11,103	19,199	30,363	39,827
2000	553,091	92	420	489	525	745	968	1,359	2,557	5,565	10,955	18,803	29,231	38,930
1999	549,838	68	418	509	503	745	979	1,429	2,576	5,764	10,968	18,412	28,269	38,637
Male														
2002	288,768	42	216	297	303	424	592	678	1,190	2,166	4,890	9,418	15,774	22,407
2001	287,075	31	229	253	280	447	577	694	1,166	2,331	4,966	9,377	15,694	21,390
2000	286,082	50	233	269	300	446	576	692	1,162	2,402	4,942	9,243	14,972	20,912
1999	285,832	36	227	279	276	448	575	755	1,204	2,467	5,004	9,151	14,483	20,891
Female														
2002	268,503	32	186	240	232	299	415	580	1,424	2,984	6,045	9,877	14,568	19,080
2001	266,693	35	191	240	235	285	395	695	1,439	3,135	6,137	9,822	14,669	18,437
2000	267,009	42	187	220	225	299	392	667	1,395	3,163	6,013	9,560	14,259	18,018
1999	264,006	32	191	230	227	297	404	674	1,372	3,297	5,964	9,261	13,786	17,746
Hispanic														
Both sexes														
2002	23,141	18	72	103	107	121	183	177	325	500	846	1,190	1,669	1,971
2001	22,371	14	84	87	90	143	146	204	293	545	851	1,186	1,629	1,814
2000	21,160	17	75	91	107	122	153	196	280	515	780	1,141	1,434	1,735
1999	20,233	16	72	87	89	94	165	189	277	471	733	1,093	1,423	1,635
Male														
2002	12,235	10	35	51	57	81	110	103	149	198	359	566	864	1,116
2001	11,825	4	44	49	41	95	104	100	133	236	379	579	793	948
2000	11,138	10	43	55	67	72	90	105	136	204	330	540	692	911
1999	10,668	11	34	45	47	55	103	98	124	198	306	539	698	901

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49, B55-B99)										
Origin not stated[2]										
Male										
1999	1	1	1	1	-	-	1	-	-	-
Female										
2002	1	-	-	1	1	-	-	-	-	-
2001	1	1	-	-	-	1	1	-	-	-
2000	1	-	-	1	-	-	-	1	-	-
1999	1	1	2	3	1	-	1	-	-	-
Malignant neoplasms (C00-C97)										
All origins										
Both sexes										
2002	51,904	63,598	81,159	89,565	77,497	50,298	22,323	5,772	789	7
2001	50,396	64,172	82,846	89,967	75,478	49,680	21,660	5,684	727	8
2000	50,075	65,843	84,288	90,655	74,444	49,155	21,564	5,653	764	11
1999	50,430	66,814	85,524	89,874	72,896	48,041	20,702	5,492	779	9
Male										
2002	28,708	35,374	44,711	47,734	39,218	23,345	9,106	1,952	218	5
2001	27,854	35,900	45,799	47,951	38,175	23,140	8,628	1,951	235	7
2000	27,515	36,730	46,461	48,284	37,623	22,714	8,506	1,850	190	10
1999	27,942	37,315	47,229	48,184	36,876	22,147	8,287	1,826	224	6
Female										
2002	23,196	28,224	36,448	41,831	38,279	26,953	13,217	3,820	571	2
2001	22,542	28,272	37,047	42,016	37,303	26,540	13,032	3,733	492	1
2000	22,560	29,113	37,827	42,371	36,821	26,441	13,058	3,803	574	1
1999	22,488	29,499	38,295	41,690	36,020	25,894	12,415	3,666	555	3
Hispanic										
Both sexes										
2002	2,276	2,712	3,145	3,120	2,245	1,446	707	188	20	-
2001	2,312	2,608	3,116	2,988	2,080	1,322	615	222	22	-
2000	2,045	2,532	3,025	2,783	2,044	1,292	586	183	24	-
1999	2,051	2,598	2,796	2,586	1,863	1,231	588	150	26	-
Male										
2002	1,261	1,532	1,724	1,737	1,165	700	319	92	6	-
2001	1,254	1,465	1,754	1,718	1,096	642	271	109	11	-
2000	1,123	1,398	1,688	1,545	1,112	632	292	88	5	-
1999	1,159	1,469	1,576	1,436	967	564	271	59	8	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms (C00-C97)														
Hispanic														
Female														
2002	10,906	8	37	52	50	40	73	74	176	302	487	624	805	855
2001	10,546	10	40	38	49	48	42	104	160	309	472	607	836	866
2000	10,022	7	32	36	40	50	63	91	144	311	450	601	742	824
1999	9,565	5	38	42	42	39	62	91	153	273	427	554	725	734
Non-Hispanic[1]														
Both sexes														
2002	532,712	55	329	433	428	598	820	1,077	2,273	4,636	10,057	18,043	28,585	39,406
2001	530,058	52	333	404	424	588	825	1,179	2,306	4,901	10,220	17,961	28,656	37,907
2000	530,531	75	341	395	417	619	813	1,159	2,272	5,036	10,134	17,602	27,703	37,088
1999	528,218	52	343	422	413	650	812	1,236	2,289	5,281	10,201	17,265	26,771	36,902
Male														
2002	275,770	31	181	246	246	342	479	573	1,037	1,962	4,513	8,817	14,865	21,232
2001	274,492	27	183	203	239	352	472	593	1,030	2,084	4,568	8,775	14,856	20,368
2000	274,185	40	186	214	232	372	486	584	1,023	2,192	4,595	8,675	14,227	19,938
1999	274,419	25	190	234	229	393	471	655	1,077	2,266	4,680	8,581	13,742	19,936
Female														
2002	256,942	24	148	187	182	256	341	504	1,236	2,674	5,544	9,226	13,720	18,174
2001	255,566	25	150	201	185	236	353	586	1,276	2,817	5,652	9,186	13,800	17,539
2000	256,346	35	155	181	185	247	327	575	1,249	2,844	5,539	8,927	13,476	17,150
1999	253,799	27	153	188	184	257	341	581	1,212	3,015	5,521	8,684	13,029	16,966
Non-Hispanic White														
Both sexes														
2002	458,754	37	236	331	311	454	643	795	1,688	3,509	7,725	13,970	22,639	32,418
2001	456,709	42	268	302	305	448	628	865	1,729	3,728	7,823	13,929	22,907	31,194
2000	458,200	60	256	307	320	452	607	884	1,715	3,802	7,656	13,598	22,263	30,664
1999	456,315	37	281	311	297	476	594	953	1,729	4,025	7,823	13,404	21,540	30,407
Male														
2002	237,323	24	134	182	173	264	375	427	791	1,511	3,530	6,912	11,787	17,454
2001	236,004	21	144	146	164	270	362	449	811	1,621	3,539	6,840	11,839	16,696
2000	235,932	34	146	161	177	270	366	452	807	1,705	3,496	6,745	11,347	16,345
1999	236,384	20	160	180	165	283	346	510	831	1,789	3,636	6,622	10,966	16,371
Female														
2002	221,431	13	102	149	138	190	268	368	897	1,998	4,195	7,058	10,852	14,964
2001	220,705	21	124	156	141	178	266	416	918	2,107	4,284	7,089	11,068	14,498
2000	222,268	26	110	146	143	182	241	432	908	2,097	4,160	6,853	10,916	14,319
1999	219,931	17	121	131	132	193	248	443	898	2,236	4,187	6,782	10,574	14,036
Non-Hispanic Black														
Both sexes														
2002	61,996	12	71	81	87	110	132	211	452	907	1,963	3,420	4,956	5,899

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms (C00-C97)										
Hispanic										
Female										
2002	1,015	1,180	1,421	1,383	1,080	746	388	96	14	-
2001	1,058	1,143	1,362	1,270	984	680	344	113	11	-
2000	922	1,134	1,337	1,238	932	660	294	95	19	-
1999	892	1,129	1,220	1,150	896	667	317	91	18	-
Non-Hispanic[1]										
Both sexes										
2002	49,480	60,718	77,827	86,232	75,071	48,738	21,564	5,569	768	5
2001	47,935	61,421	79,522	86,761	73,262	48,251	20,990	5,451	702	7
2000	47,889	63,135	81,065	87,661	72,245	47,750	20,926	5,456	739	11
1999	48,244	64,056	82,516	87,052	70,852	46,695	20,070	5,337	750	9
Male										
2002	27,355	33,742	42,879	45,877	37,962	22,595	8,763	1,857	212	4
2001	26,503	34,350	43,916	46,112	37,009	22,450	8,335	1,839	222	6
2000	26,315	35,223	44,655	46,620	36,427	22,037	8,194	1,755	185	10
1999	26,708	35,752	45,519	46,616	35,827	21,532	7,999	1,765	216	6
Female										
2002	22,125	26,976	34,948	40,355	37,109	26,143	12,801	3,712	556	1
2001	21,432	27,071	35,606	40,649	36,253	25,801	12,655	3,612	480	1
2000	21,574	27,912	36,410	41,041	35,818	25,713	12,732	3,701	554	1
1999	21,536	28,304	36,997	40,436	35,025	25,163	12,071	3,572	534	3
Non-Hispanic White										
Both sexes										
2002	41,391	51,562	67,902	76,462	67,495	44,205	19,422	4,915	642	2
2001	40,195	52,117	69,412	77,026	65,734	43,739	18,926	4,811	577	4
2000	40,201	54,052	70,790	77,899	65,180	43,210	18,871	4,800	606	7
1999	40,381	54,833	72,286	77,288	64,007	42,207	18,092	4,716	621	7
Male										
2002	22,756	28,750	37,500	40,610	34,029	20,477	7,837	1,629	169	2
2001	22,188	29,183	38,371	40,816	33,069	20,248	7,441	1,599	183	4
2000	22,023	30,179	38,906	41,247	32,705	19,854	7,309	1,507	144	7
1999	22,287	30,626	39,892	41,239	32,296	19,315	7,148	1,519	179	4
Female										
2002	18,635	22,812	30,402	35,852	33,466	23,728	11,585	3,286	473	-
2001	18,007	22,934	31,041	36,210	32,665	23,491	11,485	3,212	394	-
2000	18,178	23,873	31,884	36,652	32,475	23,356	11,562	3,293	462	-
1999	18,094	24,207	32,394	36,049	31,711	22,892	10,944	3,197	442	3
Non-Hispanic Black										
Both sexes										
2002	6,794	7,759	8,325	8,221	6,297	3,818	1,819	550	109	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms (C00-C97)														
Non-Hispanic Black														
Both sexes														
2001	61,580	5	46	78	94	110	147	241	450	949	2,001	3,370	4,805	5,694
2000	61,390	11	66	71	76	135	162	213	428	1,006	2,066	3,398	4,670	5,447
1999	61,409	11	47	84	95	136	178	227	433	1,049	2,000	3,278	4,471	5,587
Male														
2002	32,287	4	33	52	51	64	73	102	185	345	813	1,612	2,591	3,223
2001	32,335	2	29	43	57	65	83	106	161	365	836	1,628	2,568	3,133
2000	32,493	4	31	42	44	82	96	93	166	377	915	1,648	2,504	3,079
1999	32,542	3	24	40	54	84	97	113	181	382	873	1,687	2,411	3,117
Female														
2002	29,709	8	38	29	36	46	59	109	267	562	1,150	1,808	2,365	2,676
2001	29,245	3	17	35	37	45	64	135	289	584	1,165	1,742	2,237	2,561
2000	28,897	7	35	29	32	53	66	120	262	629	1,151	1,750	2,166	2,368
1999	28,867	8	23	44	41	52	81	114	252	667	1,127	1,591	2,060	2,470
Origin not stated[2]														
Both sexes														
2002	1,418	1	1	1	-	4	4	4	16	14	32	62	88	110
2001	1,339	-	3	2	1	1	1	6	6	20	32	52	78	106
2000	1,400	-	4	3	1	4	2	4	5	14	41	60	94	107
1999	1,387	-	3	-	1	1	2	4	10	12	34	54	75	100
Male														
2002	763	1	-	-	-	1	3	2	4	6	18	35	45	59
2001	758	-	2	1	-	-	1	1	3	11	19	23	45	74
2000	759	-	4	-	1	2	-	3	3	6	17	28	53	63
1999	745	-	3	-	-	-	1	2	3	3	18	31	43	54
Female														
2002	655	-	1	1	-	3	1	2	12	8	14	27	43	51
2001	581	-	1	1	1	1	-	5	3	9	13	29	33	32
2000	641	-	-	3	-	2	2	1	2	8	24	32	41	44
1999	642	-	-	-	1	1	1	2	7	9	16	23	32	46
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
All origins														
Both sexes														
2002	7,737	-	3	1	7	12	11	22	35	77	181	431	670	875
2001	7,702	-	-	2	2	10	19	19	37	93	190	427	649	861
2000	7,492	-	-	-	4	9	17	18	32	88	205	408	660	770
1999	7,486	-	2	1	5	9	18	14	29	96	195	449	612	753
Male														
2002	5,184	-	2	1	2	7	4	14	25	47	132	336	534	681

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms (C00-C97)										
Non-Hispanic Black										
Both sexes										
2001	6,502	7,877	8,515	8,163	6,291	3,776	1,780	567	116	3
2000	6,572	7,733	8,752	8,264	5,980	3,860	1,769	591	117	3
1999	6,735	7,888	8,737	8,347	5,845	3,905	1,703	531	121	1
Male										
2002	3,883	4,276	4,559	4,413	3,256	1,755	770	187	38	2
2001	3,619	4,430	4,704	4,432	3,268	1,830	734	203	37	2
2000	3,684	4,332	4,903	4,532	3,157	1,811	734	219	38	2
1999	3,816	4,423	4,805	4,603	3,000	1,887	710	198	33	1
Female										
2002	2,911	3,483	3,766	3,808	3,041	2,063	1,049	363	71	1
2001	2,883	3,447	3,811	3,731	3,023	1,946	1,046	364	79	1
2000	2,888	3,401	3,849	3,732	2,823	2,049	1,035	372	79	1
1999	2,919	3,465	3,932	3,744	2,845	2,018	993	333	88	-
Origin not stated[2]										
Both sexes										
2002	148	168	187	213	181	114	52	15	1	2
2001	149	143	208	218	136	107	55	11	3	1
2000	141	176	198	211	155	113	52	14	1	-
1999	135	160	212	236	181	115	44	5	3	-
Male										
2002	92	100	108	120	91	50	24	3	-	1
2001	97	85	129	121	70	48	22	3	2	1
2000	77	109	118	119	84	45	20	7	-	-
1999	75	94	134	132	82	51	17	2	-	-
Female										
2002	56	68	79	93	90	64	28	12	1	1
2001	52	58	79	97	66	59	33	8	1	-
2000	64	67	80	92	71	68	32	7	1	-
1999	60	66	78	104	99	64	27	3	3	-
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All origins										
Both sexes										
2002	936	846	1,052	973	748	505	262	77	13	-
2001	879	893	1,019	1,004	783	502	240	67	5	1
2000	872	942	1,017	933	693	488	249	74	13	-
1999	863	889	1,007	993	751	501	220	64	15	-
Male										
2002	711	619	698	590	407	245	101	24	4	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
All origins														
Male														
2001	5,156	-	-	1	-	3	9	12	23	61	141	325	521	687
2000	4,952	-	-	-	4	6	7	12	16	60	155	309	516	618
1999	4,944	-	1	-	3	6	14	10	17	58	146	346	474	568
Female														
2002	2,553	-	1	-	5	5	7	8	10	30	49	95	136	194
2001	2,546	-	-	1	2	7	10	7	14	32	49	102	128	174
2000	2,540	-	-	-	-	3	10	6	16	28	50	99	144	152
1999	2,542	-	1	1	2	3	4	4	12	38	49	103	138	185
Hispanic														
Both sexes														
2002	284	-	-	-	2	2	1	1	3	4	13	16	31	38
2001	311	-	-	-	-	1	3	-	1	6	8	25	31	29
2000	287	-	-	-	1	-	1	2	4	5	8	15	27	40
1999	282	-	-	-	1	-	-	-	1	9	9	19	22	29
Male														
2002	215	-	-	-	-	1	-	-	3	4	10	14	25	32
2001	219	-	-	-	-	1	1	-	1	3	8	17	24	23
2000	218	-	-	-	1	-	-	1	3	3	8	13	21	34
1999	204	-	-	-	1	-	-	-	-	4	5	16	17	27
Female														
2002	69	-	-	-	2	1	1	1	-	-	3	2	6	6
2001	92	-	-	-	-	-	2	-	-	3	-	8	7	6
2000	69	-	-	-	-	-	1	1	1	2	-	2	6	6
1999	78	-	-	-	-	-	-	-	1	5	4	3	5	2
Non-Hispanic[1]														
Both sexes														
2002	7,431	-	3	1	5	10	10	21	32	73	165	413	636	837
2001	7,370	-	-	2	2	9	16	19	36	87	181	402	616	828
2000	7,181	-	-	-	3	9	15	16	28	83	197	390	632	727
1999	7,181	-	2	1	4	9	18	14	28	86	186	429	589	720
Male														
2002	4,950	-	2	1	2	6	4	14	22	43	119	320	506	649
2001	4,920	-	-	1	-	2	8	12	22	58	132	308	496	661
2000	4,719	-	-	-	3	6	7	11	13	57	147	294	494	582
1999	4,725	-	1	-	2	6	14	10	17	53	141	329	456	538
Female														
2002	2,481	-	1	-	3	4	6	7	10	30	46	93	130	188
2001	2,450	-	-	1	2	7	8	7	14	29	49	94	120	167

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All origins										
Male										
2001	685	631	696	598	427	232	83	18	2	1
2000	652	667	659	569	370	217	96	19	-	-
1999	663	629	679	605	402	229	79	14	1	-
Female										
2002	225	227	354	383	341	260	161	53	9	-
2001	194	262	323	406	356	270	157	49	3	-
2000	220	275	358	364	323	271	153	55	13	-
1999	200	260	328	388	349	272	141	50	14	-
Hispanic										
Both sexes										
2002	28	30	32	34	20	14	11	3	1	-
2001	40	41	37	39	19	21	6	4	-	-
2000	35	40	33	27	21	16	10	2	-	-
1999	37	37	35	36	27	13	6	-	1	-
Male										
2002	21	23	28	25	11	11	5	2	-	-
2001	32	31	27	26	13	8	2	2	-	-
2000	28	32	26	15	13	9	9	2	-	-
1999	28	33	29	27	8	7	2	-	-	-
Female										
2002	7	7	4	9	9	3	6	1	1	-
2001	8	10	10	13	6	13	4	2	-	-
2000	7	8	7	12	8	7	1	-	-	-
1999	9	4	6	9	19	6	4	-	1	-
Non-Hispanic[1]										
Both sexes										
2002	902	815	1,015	938	728	490	251	74	12	-
2001	838	849	980	960	764	479	234	63	5	-
2000	833	896	981	905	670	472	239	72	13	-
1999	822	849	969	954	724	487	214	64	12	-
Male										
2002	686	595	666	564	396	233	96	22	4	-
2001	652	597	667	568	414	223	81	16	2	-
2000	622	631	632	553	355	208	87	17	-	-
1999	634	593	648	576	394	221	77	14	1	-
Female										
2002	216	220	349	374	332	257	155	52	8	-
2001	186	252	313	392	350	256	153	47	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
Non-Hispanic[1]														
Female														
2000	2,462	-	-	-	-	3	8	5	15	26	50	96	138	145
1999	2,456	-	1	1	2	3	4	4	11	33	45	100	133	182
Non-Hispanic White														
Both sexes														
2002	6,141	-	2	1	2	6	6	13	22	48	116	286	458	646
2001	6,052	-	-	1	1	7	11	14	22	49	116	272	464	632
2000	5,843	-	-	-	1	6	10	10	19	55	131	275	448	541
1999	5,826	-	2	1	2	5	9	10	19	51	125	269	408	530
Male														
2002	4,035	-	2	1	1	4	3	9	16	31	83	224	363	506
2001	3,961	-	-	1	-	1	5	10	12	34	86	211	375	500
2000	3,726	-	-	-	1	3	4	7	9	35	93	209	337	434
1999	3,746	-	1	-	1	3	6	6	11	30	95	203	320	390
Female														
2002	2,106	-	-	-	1	2	3	4	6	17	33	62	95	140
2001	2,091	-	-	-	1	6	6	4	10	15	30	61	89	132
2000	2,117	-	-	-	-	3	6	3	10	20	38	66	111	107
1999	2,080	-	1	1	1	2	3	4	8	21	30	66	88	140
Non-Hispanic Black														
Both sexes														
2002	1,034	-	-	-	2	4	2	3	8	22	41	106	148	150
2001	1,056	-	-	1	-	1	3	2	4	24	47	105	124	172
2000	1,089	-	-	-	2	3	4	5	2	15	45	94	155	161
1999	1,131	-	-	-	2	4	4	2	6	25	42	133	154	162
Male														
2002	757	-	-	-	1	2	1	2	5	10	31	80	123	122
2001	773	-	-	-	-	1	1	1	4	14	31	79	100	141
2000	821	-	-	-	2	3	2	3	1	11	35	67	136	130
1999	819	-	-	-	1	3	3	2	4	17	30	107	117	125
Female														
2002	277	-	-	-	1	2	1	1	3	12	10	26	25	28
2001	283	-	-	1	-	-	2	1	-	10	16	26	24	31
2000	268	-	-	-	-	-	2	2	1	4	10	27	19	31
1999	312	-	-	-	1	1	1	-	2	8	12	26	37	37
Origin not stated[2]														
Both sexes														
2002	22	-	-	-	-	-	-	-	-	-	3	2	3	-
2001	21	-	-	-	-	-	-	-	-	-	1	-	2	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
Non-Hispanic[1]										
Female										
2000	211	265	349	352	315	264	152	55	13	-
1999	188	256	321	378	330	266	137	50	11	-
Non-Hispanic White										
Both sexes										
2002	712	659	906	843	661	444	236	64	10	-
2001	656	679	845	859	701	440	216	62	5	-
2000	636	733	833	806	606	436	221	65	11	-
1999	650	699	828	837	660	443	207	60	11	-
Male										
2002	536	476	591	510	357	210	92	17	3	-
2001	504	478	567	502	378	204	75	16	2	-
2000	470	507	529	480	317	196	79	16	-	-
1999	502	487	556	496	354	197	74	13	1	-
Female										
2002	176	183	315	333	304	234	144	47	7	-
2001	152	201	278	357	323	236	141	46	3	-
2000	166	226	304	326	289	240	142	49	11	-
1999	148	212	272	341	306	246	133	47	10	-
Non-Hispanic Black										
Both sexes										
2002	168	124	89	66	48	32	12	7	2	-
2001	146	140	116	80	49	25	16	1	-	-
2000	166	133	121	80	56	25	14	6	2	-
1999	154	128	120	96	54	37	5	2	1	-
Male										
2002	135	96	61	36	28	17	3	3	1	-
2001	122	98	86	52	26	12	5	-	-	-
2000	132	104	87	60	34	7	7	-	-	-
1999	118	91	78	69	33	19	1	1	-	-
Female										
2002	33	28	28	30	20	15	9	4	1	-
2001	24	42	30	28	23	13	11	1	-	-
2000	34	29	34	20	22	18	7	6	2	-
1999	36	37	42	27	21	18	4	1	1	-
Origin not stated[2]										
Both sexes										
2002	6	1	5	1	-	1	-	-	-	-
2001	1	3	2	5	-	2	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)														
Origin not stated[2]														
Both sexes														
2000	24	-	-	-	-	-	1	-	-	-	-	3	1	3
1999	23	-	-	-	-	-	-	-	-	1	-	1	1	4
Male														
2002	19	-	-	-	-	-	-	-	-	-	3	2	3	-
2001	17	-	-	-	-	-	-	-	-	-	1	-	1	3
2000	15	-	-	-	-	-	-	-	-	-	-	2	1	2
1999	15	-	-	-	-	-	-	-	-	1	-	1	1	3
Female														
2002	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	9	-	-	-	-	-	1	-	-	-	-	1	-	1
1999	8	-	-	-	-	-	-	-	-	-	-	-	-	1
Malignant neoplasm of esophagus (C15)														
All origins														
Both sexes														
2002	12,701	-	-	-	1	-	6	4	15	84	227	503	887	1,286
2001	12,530	-	-	-	-	2	4	9	20	68	241	492	916	1,114
2000	12,232	1	-	-	-	1	2	9	30	78	219	478	870	1,180
1999	11,918	-	-	-	-	-	-	10	27	75	218	469	793	1,139
Male														
2002	9,789	-	-	-	-	-	3	4	10	71	186	426	759	1,099
2001	9,607	-	-	-	-	1	4	8	18	52	203	417	787	940
2000	9,279	1	-	-	-	-	2	7	24	69	196	415	746	988
1999	9,094	-	-	-	-	-	-	7	22	65	190	410	684	969
Female														
2002	2,912	-	-	-	1	-	3	-	5	13	41	77	128	187
2001	2,923	-	-	-	-	1	-	1	2	16	38	75	129	174
2000	2,953	-	-	-	-	1	-	2	6	9	23	63	124	192
1999	2,824	-	-	-	-	-	-	3	5	10	28	59	109	170
Hispanic														
Both sexes														
2002	411	-	-	-	-	-	1	-	1	5	18	22	31	36
2001	401	-	-	-	-	-	-	2	-	3	9	21	42	34
2000	384	-	-	-	-	-	-	-	2	3	7	15	23	37
1999	367	-	-	-	-	-	-	-	2	6	10	22	22	36
Male														
2002	329	-	-	-	-	-	1	-	1	5	14	22	27	35
2001	316	-	-	-	-	-	-	2	-	2	7	19	42	25

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
Origin not stated[2]										
Both sexes										
2000	4	6	3	1	2	-	-	-	-	-
1999	4	3	3	3	-	1	-	-	2	-
Male										
2002	4	1	4	1	-	1	-	-	-	-
2001	1	3	2	4	-	1	-	-	-	1
2000	2	4	1	1	2	-	-	-	-	-
1999	1	3	2	2	-	1	-	-	-	-
Female										
2002	2	-	1	-	-	-	-	-	-	-
2001	-	-	-	1	-	1	-	-	-	-
2000	2	2	2	-	-	-	-	-	-	-
1999	3	-	1	1	-	-	-	-	2	-
Malignant neoplasm of esophagus (C15)										
All origins										
Both sexes										
2002	1,477	1,736	1,955	1,909	1,480	766	295	65	4	1
2001	1,440	1,765	2,014	1,941	1,366	757	317	56	7	1
2000	1,430	1,711	1,960	1,817	1,346	764	274	55	7	-
1999	1,392	1,739	1,955	1,784	1,202	755	279	72	8	1
Male										
2002	1,222	1,394	1,558	1,437	1,017	452	127	23	1	-
2001	1,197	1,436	1,553	1,423	942	452	153	18	2	1
2000	1,181	1,349	1,482	1,337	896	437	122	23	4	-
1999	1,112	1,396	1,529	1,328	808	421	127	22	3	1
Female										
2002	255	342	397	472	463	314	168	42	3	1
2001	243	329	461	518	424	305	164	38	5	-
2000	249	362	478	480	450	327	152	32	3	-
1999	280	343	426	456	394	334	152	50	5	-
Hispanic										
Both sexes										
2002	44	58	62	54	37	23	14	5	-	-
2001	62	37	58	49	52	12	18	2	-	-
2000	49	56	64	62	36	20	6	3	1	-
1999	49	56	39	54	33	22	14	2	-	-
Male										
2002	40	47	49	41	25	13	8	1	-	-
2001	49	33	47	33	35	7	15	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of esophagus (C15)														
Hispanic														
Male														
2000	292	-	-	-	-	-	-	-	2	3	5	13	21	26
1999	294	-	-	-	-	-	-	-	2	5	10	20	19	34
Female														
2002	82	-	-	-	-	-	-	-	-	-	4	-	4	1
2001	85	-	-	-	-	-	-	-	-	1	2	2	-	9
2000	92	-	-	-	-	-	-	-	-	-	2	2	2	11
1999	73	-	-	-	-	-	-	-	-	1	-	2	3	2
Non-Hispanic[1]														
Both sexes														
2002	12,254	-	-	-	1	-	5	3	14	79	208	480	854	1,243
2001	12,085	-	-	-	-	2	4	7	20	65	231	468	871	1,077
2000	11,819	1	-	-	-	1	2	9	28	75	212	463	845	1,143
1999	11,520	-	-	-	-	-	-	10	25	69	207	447	768	1,100
Male														
2002	9,431	-	-	-	-	-	2	3	9	66	171	403	730	1,058
2001	9,254	-	-	-	-	1	4	6	18	50	195	395	743	912
2000	8,964	1	-	-	-	-	2	7	22	66	191	402	724	962
1999	8,775	-	-	-	-	-	-	7	20	60	179	390	663	932
Female														
2002	2,823	-	-	-	1	-	3	-	5	13	37	77	124	185
2001	2,831	-	-	-	-	1	-	1	2	15	36	73	128	165
2000	2,855	-	-	-	-	1	-	2	6	9	21	61	121	181
1999	2,745	-	-	-	-	-	-	3	5	9	28	57	105	168
Non-Hispanic White														
Both sexes														
2002	10,484	-	-	-	1	-	5	2	13	68	159	382	694	1,002
2001	10,193	-	-	-	-	2	4	5	17	56	182	365	674	823
2000	10,005	-	-	-	-	1	2	9	27	71	168	362	651	897
1999	9,703	-	-	-	-	-	-	9	21	61	152	340	596	860
Male														
2002	8,169	-	-	-	-	-	2	2	9	60	136	332	606	866
2001	7,863	-	-	-	-	1	4	4	15	43	157	316	581	713
2000	7,680	-	-	-	-	-	2	7	22	64	157	328	567	766
1999	7,440	-	-	-	-	-	-	6	16	55	133	303	524	739
Female														
2002	2,315	-	-	-	1	-	3	-	4	8	23	50	88	136
2001	2,330	-	-	-	-	1	-	1	2	13	25	49	93	110
2000	2,325	-	-	-	-	1	-	2	5	7	11	34	84	131
1999	2,263	-	-	-	-	-	-	3	5	6	19	37	72	121

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of esophagus (C15)										
Hispanic										
Male										
2000	38	44	48	48	24	14	3	2	1	-
1999	44	42	32	42	21	16	7	-	-	-
Female										
2002	4	11	13	13	12	10	6	4	-	-
2001	13	4	11	16	17	5	3	2	-	-
2000	11	12	16	14	12	6	3	1	-	-
1999	5	14	7	12	12	6	7	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,428	1,675	1,885	1,853	1,443	740	278	60	4	1
2001	1,370	1,721	1,943	1,888	1,313	745	298	54	7	1
2000	1,378	1,649	1,889	1,751	1,307	742	266	52	6	-
1999	1,339	1,677	1,910	1,725	1,168	732	264	70	8	1
Male										
2002	1,177	1,344	1,504	1,394	992	437	118	22	1	-
2001	1,140	1,396	1,498	1,386	906	445	138	18	2	1
2000	1,140	1,300	1,428	1,285	871	421	118	21	3	-
1999	1,064	1,349	1,492	1,281	787	405	120	22	3	1
Female										
2002	251	331	381	459	451	303	160	38	3	1
2001	230	325	445	502	407	300	160	36	5	-
2000	238	349	461	466	436	321	148	31	3	-
1999	275	328	418	444	381	327	144	48	5	-
Non-Hispanic White										
Both sexes										
2002	1,171	1,384	1,643	1,645	1,316	679	261	56	3	-
2001	1,102	1,413	1,647	1,690	1,198	688	268	51	7	1
2000	1,100	1,380	1,637	1,535	1,179	687	245	48	6	-
1999	1,063	1,390	1,648	1,508	1,068	673	244	65	5	-
Male										
2002	983	1,137	1,328	1,261	910	403	113	21	-	-
2001	932	1,162	1,296	1,255	831	413	120	17	2	1
2000	936	1,110	1,256	1,136	805	396	108	17	3	-
1999	859	1,148	1,303	1,127	730	369	108	18	2	-
Female										
2002	188	247	315	384	406	276	148	35	3	-
2001	170	251	351	435	367	275	148	34	5	-
2000	164	270	381	399	374	291	137	31	3	-
1999	204	242	345	381	338	304	136	47	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of esophagus (C15)														
Non-Hispanic Black														
Both sexes														
2002	1,572	-	-	-	-	-	-	1	1	9	47	91	143	221
2001	1,676	-	-	-	-	-	-	2	1	9	45	92	172	225
2000	1,645	1	-	-	-	-	-	-	1	3	41	96	182	228
1999	1,629	-	-	-	-	-	-	-	3	6	47	99	162	222
Male														
2002	1,118	-	-	-	-	-	-	1	-	4	34	66	109	179
2001	1,222	-	-	-	-	-	-	2	1	7	34	69	141	176
2000	1,162	1	-	-	-	-	-	-	-	1	31	69	149	185
1999	1,187	-	-	-	-	-	-	-	3	4	40	79	130	177
Female														
2002	454	-	-	-	-	-	-	-	1	5	13	25	34	42
2001	454	-	-	-	-	-	-	-	-	2	11	23	31	49
2000	483	-	-	-	-	-	-	-	1	2	10	27	33	43
1999	442	-	-	-	-	-	-	-	-	2	7	20	32	45
Origin not stated[2]														
Both sexes														
2002	36	-	-	-	-	-	-	1	-	-	1	1	2	7
2001	44	-	-	-	-	-	-	-	-	-	1	3	3	3
2000	29	-	-	-	-	-	-	-	-	-	-	-	2	-
1999	31	-	-	-	-	-	-	-	-	-	1	-	3	3
Male														
2002	29	-	-	-	-	-	-	1	-	-	1	1	2	6
2001	37	-	-	-	-	-	-	-	-	-	1	3	2	3
2000	23	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	25	-	-	-	-	-	-	-	-	-	1	-	2	3
Female														
2002	7	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	7	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	6	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	6	-	-	-	-	-	-	-	-	-	-	-	1	-
Malignant neoplasm of stomach (C16)														
All origins														
Both sexes														
2002	12,198	-	1	-	-	7	22	48	90	160	276	443	663	841
2001	12,319	-	-	-	-	5	15	33	89	178	286	452	675	751
2000	12,645	-	-	-	1	3	11	36	88	167	317	468	629	753
1999	12,711	-	-	-	1	5	8	47	89	189	316	450	651	808
Male														
2002	7,133	-	1	-	-	-	8	31	49	82	157	285	440	556

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of esophagus (C15)										
Non-Hispanic Black										
Both sexes										
2002	233	258	217	182	100	52	12	3	1	1
2001	241	272	262	176	102	51	23	3	-	-
2000	253	243	219	201	106	48	19	4	-	-
1999	250	257	229	198	82	49	17	5	2	1
Male										
2002	172	181	159	115	63	30	3	1	1	-
2001	187	202	177	118	67	27	13	1	-	-
2000	184	171	147	137	54	21	8	4	-	-
1999	188	176	164	141	42	28	10	4	-	1
Female										
2002	61	77	58	67	37	22	9	2	-	1
2001	54	70	85	58	35	24	10	2	-	-
2000	69	72	72	64	52	27	11	-	-	-
1999	62	81	65	57	40	21	7	1	2	-
Origin not stated[2]										
Both sexes										
2002	5	3	8	2	-	3	3	-	-	-
2001	8	7	13	4	1	-	1	-	-	-
2000	3	6	7	4	3	2	2	-	-	-
1999	4	6	6	5	1	1	1	-	-	-
Male										
2002	5	3	5	2	-	2	1	-	-	-
2001	8	7	8	4	1	-	-	-	-	-
2000	3	5	6	4	1	2	1	-	-	-
1999	4	5	5	5	-	-	-	-	-	-
Female										
2002	-	-	3	-	-	1	2	-	-	-
2001	-	-	5	-	-	-	1	-	-	-
2000	-	1	1	-	2	-	1	-	-	-
1999	-	1	1	-	1	1	1	-	-	-
Malignant neoplasm of stomach (C16)										
All origins										
Both sexes										
2002	990	1,179	1,637	1,903	1,753	1,344	664	153	24	-
2001	1,020	1,308	1,713	2,004	1,789	1,216	620	149	16	-
2000	1,051	1,317	1,737	2,009	1,824	1,379	641	188	26	-
1999	1,052	1,359	1,793	2,064	1,796	1,277	607	170	29	-
Male										
2002	659	804	1,063	1,118	946	628	258	38	10	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of stomach (C16)														
All origins														
Male														
2001	7,191	-	-	-	-	4	5	20	51	98	155	288	443	514
2000	7,328	-	-	-	-	-	6	22	47	100	184	290	413	506
1999	7,459	-	-	-	1	2	4	22	54	101	182	303	443	558
Female														
2002	5,065	-	-	-	-	7	14	17	41	78	119	158	223	285
2001	5,128	-	-	-	-	1	10	13	38	80	131	164	232	237
2000	5,317	-	-	-	1	3	5	14	41	67	133	178	216	247
1999	5,252	-	-	-	-	3	4	25	35	88	134	147	208	250
Hispanic														
Both sexes														
2002	1,185	-	-	-	-	1	5	11	27	45	52	71	83	97
2001	1,189	-	-	-	-	1	3	16	25	40	58	78	104	93
2000	1,111	-	-	-	1	-	4	10	14	48	57	73	78	68
1999	1,055	-	-	-	-	1	3	14	24	34	51	50	71	83
Male														
2002	671	-	-	-	-	-	2	8	18	15	29	42	43	59
2001	680	-	-	-	-	1	1	7	12	23	31	43	61	51
2000	655	-	-	-	-	-	1	7	9	26	26	35	43	43
1999	607	-	-	-	-	-	1	8	17	14	27	35	41	55
Female														
2002	514	-	-	-	-	1	3	3	9	30	23	29	40	38
2001	509	-	-	-	-	-	2	9	13	17	27	35	43	42
2000	456	-	-	-	1	-	3	3	5	22	31	38	35	25
1999	448	-	-	-	-	1	2	6	7	20	24	15	30	28
Non-Hispanic[1]														
Both sexes														
2002	10,974	-	1	-	-	6	17	36	63	114	223	372	578	742
2001	11,099	-	-	-	-	4	12	17	64	138	227	374	569	656
2000	11,491	-	-	-	-	3	7	25	74	119	259	394	549	681
1999	11,615	-	-	-	1	4	5	33	65	155	265	395	575	724
Male														
2002	6,438	-	1	-	-	-	6	23	31	66	128	243	396	495
2001	6,492	-	-	-	-	3	4	13	39	75	124	245	380	461
2000	6,648	-	-	-	-	-	5	15	38	74	157	254	368	461
1999	6,827	-	-	-	1	2	3	14	37	87	155	263	399	502
Female														
2002	4,536	-	-	-	-	6	11	13	32	48	95	129	182	247
2001	4,607	-	-	-	-	1	8	4	25	63	103	129	189	195
2000	4,843	-	-	-	-	3	2	10	36	45	102	140	181	220

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of stomach (C16)										
All origins										
Male										
2001	724	858	1,070	1,194	954	550	222	36	5	-
2000	722	860	1,106	1,201	943	617	247	56	8	-
1999	694	893	1,112	1,250	989	568	223	52	8	-
Female										
2002	331	375	574	785	807	716	406	115	14	-
2001	296	450	643	810	835	666	398	113	11	-
2000	329	457	631	808	881	762	394	132	18	-
1999	358	466	681	814	807	709	384	118	21	-
Hispanic										
Both sexes										
2002	121	111	148	162	121	73	42	15	-	-
2001	116	135	165	142	111	60	30	11	1	-
2000	99	139	167	134	120	65	26	8	-	-
1999	109	134	137	137	102	63	32	6	4	-
Male										
2002	69	71	101	87	63	39	20	5	-	-
2001	79	86	100	76	59	32	10	7	1	-
2000	67	93	112	73	71	34	12	3	-	-
1999	71	86	76	87	54	22	9	2	2	-
Female										
2002	52	40	47	75	58	34	22	10	-	-
2001	37	49	65	66	52	28	20	4	-	-
2000	32	46	55	61	49	31	14	5	-	-
1999	38	48	61	50	48	41	23	4	2	-
Non-Hispanic[1]										
Both sexes										
2002	867	1,064	1,484	1,736	1,627	1,264	619	137	24	-
2001	902	1,171	1,541	1,853	1,676	1,155	588	137	15	-
2000	946	1,173	1,566	1,869	1,699	1,309	613	180	25	-
1999	938	1,221	1,646	1,924	1,691	1,212	572	164	25	-
Male										
2002	589	729	959	1,028	880	585	236	33	10	-
2001	644	771	966	1,114	893	517	211	28	4	-
2000	652	762	991	1,123	871	582	234	53	8	-
1999	620	805	1,029	1,162	934	545	213	50	6	-
Female										
2002	278	335	525	708	747	679	383	104	14	-
2001	258	400	575	739	783	638	377	109	11	-
2000	294	411	575	746	828	727	379	127	17	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of stomach (C16)														
Non-Hispanic[1]														
Female														
1999	4,788	-	-	-	-	2	2	19	28	68	110	132	176	222
Non-Hispanic White														
Both sexes														
2002	8,124	-	1	-	-	4	11	21	34	75	130	246	370	525
2001	8,308	-	-	-	-	1	5	9	34	72	139	244	371	474
2000	8,600	-	-	-	-	2	3	10	43	65	148	242	383	491
1999	8,712	-	-	-	-	1	3	15	35	95	142	267	379	502
Male														
2002	4,864	-	1	-	-	-	3	13	18	44	79	167	264	366
2001	4,884	-	-	-	-	1	1	7	22	44	78	164	252	335
2000	5,033	-	-	-	-	-	2	6	23	40	89	158	266	336
1999	5,171	-	-	-	-	1	2	7	23	57	85	184	270	362
Female														
2002	3,260	-	-	-	-	4	8	8	16	31	51	79	106	159
2001	3,424	-	-	-	-	-	4	2	12	28	61	80	119	139
2000	3,567	-	-	-	-	2	1	4	20	25	59	84	117	155
1999	3,541	-	-	-	-	-	1	8	12	38	57	83	109	140
Non-Hispanic Black														
Both sexes														
2002	2,112	-	-	-	-	2	6	13	20	26	64	92	156	155
2001	2,051	-	-	-	-	1	4	4	21	49	72	90	147	128
2000	2,156	-	-	-	-	1	3	11	25	37	80	109	130	147
1999	2,183	-	-	-	1	1	2	14	22	40	87	89	148	170
Male														
2002	1,184	-	-	-	-	-	3	10	10	16	33	60	105	102
2001	1,177	-	-	-	-	1	1	3	11	25	38	57	104	86
2000	1,202	-	-	-	-	-	2	5	15	26	51	71	83	98
1999	1,252	-	-	-	1	-	1	7	12	20	52	60	98	109
Female														
2002	928	-	-	-	-	2	3	3	10	10	31	32	51	53
2001	874	-	-	-	-	-	3	1	10	24	34	33	43	42
2000	954	-	-	-	-	1	1	6	10	11	29	38	47	49
1999	931	-	-	-	-	1	1	7	10	20	35	29	50	61
Origin not stated[2]														
Both sexes														
2002	39	-	-	-	-	-	-	1	-	1	1	-	2	2
2001	31	-	-	-	-	-	-	-	-	-	1	-	2	2
2000	43	-	-	-	-	-	-	1	-	-	1	1	2	4
1999	41	-	-	-	-	-	-	-	-	-	-	5	5	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of stomach (C16)										
Non-Hispanic[1]										
Female										
1999	318	416	617	762	757	667	359	114	19	-
Non-Hispanic White										
Both sexes										
2002	612	753	1,108	1,331	1,282	1,011	496	100	14	-
2001	636	827	1,183	1,448	1,331	948	471	105	10	-
2000	689	850	1,181	1,436	1,345	1,061	497	141	13	-
1999	654	930	1,251	1,499	1,360	976	463	125	15	-
Male										
2002	425	534	731	815	708	481	187	24	4	-
2001	459	540	748	894	712	431	172	22	2	-
2000	489	567	763	877	703	476	192	42	4	-
1999	436	626	800	916	754	430	174	40	4	-
Female										
2002	187	219	377	516	574	530	309	76	10	-
2001	177	287	435	554	619	517	299	83	8	-
2000	200	283	418	559	642	585	305	99	9	-
1999	218	304	451	583	606	546	289	85	11	-
Non-Hispanic Black										
Both sexes										
2002	184	229	289	306	239	196	96	30	9	-
2001	196	251	254	292	256	161	94	26	5	-
2000	188	239	284	316	263	196	86	30	11	-
1999	203	212	295	326	254	193	85	32	9	-
Male										
2002	121	145	178	159	117	73	40	7	5	-
2001	134	172	160	162	131	59	25	6	2	-
2000	124	145	169	173	122	80	27	8	3	-
1999	132	134	175	188	137	89	28	7	2	-
Female										
2002	63	84	111	147	122	123	56	23	4	-
2001	62	79	94	130	125	102	69	20	3	-
2000	64	94	115	143	141	116	59	22	8	-
1999	71	78	120	138	117	104	57	25	7	-
Origin not stated[2]										
Both sexes										
2002	2	4	5	5	5	7	3	1	-	-
2001	2	2	7	9	2	1	2	1	-	-
2000	6	5	4	6	5	5	2	-	1	-
1999	5	4	10	3	3	2	3	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of stomach (C16)														
Origin not stated[2]														
Male														
2002	24	-	-	-	-	-	-	-	-	1	-	-	1	2
2001	19	-	-	-	-	-	-	-	-	-	-	-	2	2
2000	25	-	-	-	-	-	-	-	-	-	1	1	2	2
1999	25	-	-	-	-	-	-	-	-	-	-	5	3	1
Female														
2002	15	-	-	-	-	-	-	1	-	-	1	-	1	-
2001	12	-	-	-	-	-	-	-	-	-	1	-	-	-
2000	18	-	-	-	-	-	-	1	-	-	-	-	-	2
1999	16	-	-	-	-	-	-	-	-	-	-	-	2	-
Malignant neoplasms of colon, rectum and anus (C18-C21)														
All origins														
Both sexes														
2002	56,741	1	-	-	1	9	41	88	202	439	986	1,755	2,818	3,694
2001	56,887	-	-	-	-	11	35	81	218	505	928	1,744	2,804	3,610
2000	57,477	-	-	-	3	13	27	79	193	480	889	1,656	2,581	3,397
1999	57,155	1	-	-	1	11	38	87	196	439	894	1,623	2,434	3,309
Male														
2002	28,501	1	-	-	1	6	28	48	113	233	564	958	1,606	2,185
2001	28,225	-	-	-	-	6	18	38	119	290	491	967	1,545	2,116
2000	28,462	-	-	-	2	8	16	42	111	271	488	893	1,466	1,989
1999	28,240	-	-	-	1	5	22	57	103	226	452	876	1,374	1,932
Female														
2002	28,240	-	-	-	-	3	13	40	89	206	422	797	1,212	1,509
2001	28,662	-	-	-	-	5	17	43	99	215	437	777	1,259	1,494
2000	29,015	-	-	-	1	5	11	37	82	209	401	763	1,115	1,408
1999	28,915	1	-	-	-	6	16	30	93	213	442	747	1,060	1,377
Hispanic														
Both sexes														
2002	2,364	-	-	-	1	1	1	12	15	44	77	104	178	197
2001	2,250	-	-	-	-	-	6	5	15	44	83	88	157	176
2000	2,117	-	-	-	-	3	5	8	25	47	71	111	152	184
1999	2,031	-	-	-	-	1	6	7	24	33	53	113	132	132
Male														
2002	1,287	-	-	-	1	-	-	7	6	23	39	66	104	135
2001	1,185	-	-	-	-	-	3	4	4	28	48	44	80	99
2000	1,124	-	-	-	-	1	2	6	11	20	32	53	87	102
1999	1,062	-	-	-	-	1	4	3	11	15	24	59	74	72

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of stomach (C16)										
Origin not stated[2]										
Male										
2002	1	4	3	3	3	4	2	-	-	-
2001	1	1	4	4	2	1	1	1	-	-
2000	3	5	3	5	1	1	1	-	-	-
1999	3	2	7	1	1	1	1	-	-	-
Female										
2002	1	-	2	2	2	3	1	1	-	-
2001	1	1	3	5	-	-	1	-	-	-
2000	3	-	1	1	4	4	1	-	1	-
1999	2	2	3	2	2	1	2	-	-	-
Malignant neoplasms of colon, rectum and anus (C18-C21)										
All origins										
Both sexes										
2002	4,727	5,901	7,491	8,826	8,672	6,453	3,433	1,075	128	1
2001	4,374	5,997	7,794	8,988	8,539	6,618	3,482	1,028	131	-
2000	4,543	5,913	8,188	9,410	8,643	6,735	3,518	1,061	148	-
1999	4,658	6,095	8,154	9,473	8,456	6,672	3,503	973	138	-
Male										
2002	2,850	3,455	4,125	4,536	3,968	2,508	1,024	267	24	1
2001	2,536	3,479	4,294	4,568	3,883	2,520	1,068	251	36	-
2000	2,619	3,407	4,517	4,852	3,908	2,571	1,012	260	30	-
1999	2,689	3,431	4,452	4,857	3,897	2,525	1,081	236	24	-
Female										
2002	1,877	2,446	3,366	4,290	4,704	3,945	2,409	808	104	-
2001	1,838	2,518	3,500	4,420	4,656	4,098	2,414	777	95	-
2000	1,924	2,506	3,671	4,558	4,735	4,164	2,506	801	118	-
1999	1,969	2,664	3,702	4,616	4,559	4,147	2,422	737	114	-
Hispanic										
Both sexes										
2002	257	275	330	338	235	161	108	28	2	-
2001	239	255	337	317	218	169	96	41	4	-
2000	211	242	288	279	218	162	73	32	6	-
1999	212	271	310	274	211	153	72	25	2	-
Male										
2002	156	168	183	185	110	65	29	10	-	-
2001	146	150	183	167	94	76	39	19	1	-
2000	125	147	154	148	119	78	28	11	-	-
1999	113	152	179	147	110	58	31	8	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of colon, rectum and anus (C18-C21)														
Hispanic														
Female														
2002	1,077	-	-	-	-	1	1	5	9	21	38	38	74	62
2001	1,065	-	-	-	-	-	3	1	11	16	35	44	77	77
2000	993	-	-	-	-	2	3	2	14	27	39	58	65	82
1999	969	-	-	-	-	-	2	4	13	18	29	54	58	60
Non-Hispanic[1]														
Both sexes														
2002	54,244	1	-	-	-	8	40	75	184	394	905	1,647	2,634	3,488
2001	54,513	-	-	-	-	11	29	76	203	458	843	1,650	2,639	3,421
2000	55,244	-	-	-	3	10	22	71	167	432	815	1,538	2,425	3,201
1999	54,976	1	-	-	1	10	31	80	172	404	837	1,506	2,298	3,168
Male														
2002	27,140	1	-	-	-	6	28	41	105	209	521	888	1,500	2,045
2001	26,978	-	-	-	-	6	15	34	115	260	441	920	1,462	2,008
2000	27,287	-	-	-	2	7	14	36	99	250	455	837	1,378	1,885
1999	27,103	-	-	-	1	4	18	54	92	211	425	816	1,297	1,854
Female														
2002	27,104	-	-	-	-	2	12	34	79	185	384	759	1,134	1,443
2001	27,535	-	-	-	-	5	14	42	88	198	402	730	1,177	1,413
2000	27,957	-	-	-	1	3	8	35	68	182	360	701	1,047	1,316
1999	27,873	1	-	-	-	6	13	26	80	193	412	690	1,001	1,314
Non-Hispanic White														
Both sexes														
2002	46,147	1	-	-	-	6	30	57	149	292	655	1,211	2,060	2,788
2001	46,450	-	-	-	-	7	21	55	148	339	627	1,235	2,049	2,721
2000	47,317	-	-	-	3	9	14	49	118	303	595	1,137	1,897	2,546
1999	47,275	-	-	-	1	6	20	56	126	303	630	1,152	1,804	2,540
Male														
2002	23,119	1	-	-	-	4	20	32	82	151	383	670	1,183	1,661
2001	23,128	-	-	-	-	5	8	24	87	196	327	701	1,173	1,630
2000	23,369	-	-	-	2	6	7	26	75	184	331	637	1,095	1,507
1999	23,446	-	-	-	1	2	10	38	67	166	334	634	1,045	1,500
Female														
2002	23,028	-	-	-	-	2	10	25	67	141	272	541	877	1,127
2001	23,322	-	-	-	-	2	13	31	61	143	300	534	876	1,091
2000	23,948	-	-	-	1	3	7	23	43	119	264	500	802	1,039
1999	23,829	-	-	-	-	4	10	18	59	137	296	518	759	1,040

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)										
Hispanic										
Female										
2002	101	107	147	153	125	96	79	18	2	-
2001	93	105	154	150	124	93	57	22	3	-
2000	86	95	134	131	99	84	45	21	6	-
1999	99	119	131	127	101	95	41	17	1	-
Non-Hispanic[1]										
Both sexes										
2002	4,460	5,615	7,143	8,466	8,416	6,284	3,312	1,045	126	1
2001	4,124	5,733	7,439	8,655	8,308	6,434	3,379	985	126	-
2000	4,322	5,656	7,888	9,114	8,413	6,562	3,436	1,027	142	-
1999	4,436	5,811	7,829	9,180	8,210	6,494	3,424	948	136	-
Male										
2002	2,686	3,279	3,929	4,338	3,850	2,441	991	257	24	1
2001	2,382	3,322	4,097	4,396	3,785	2,441	1,028	232	34	-
2000	2,488	3,250	4,359	4,694	3,782	2,492	980	249	30	-
1999	2,570	3,271	4,267	4,699	3,768	2,458	1,047	228	23	-
Female										
2002	1,774	2,336	3,214	4,128	4,566	3,843	2,321	788	102	-
2001	1,742	2,411	3,342	4,259	4,523	3,993	2,351	753	92	-
2000	1,834	2,406	3,529	4,420	4,631	4,070	2,456	778	112	-
1999	1,866	2,540	3,562	4,481	4,442	4,036	2,377	720	113	-
Non-Hispanic White										
Both sexes										
2002	3,613	4,658	6,124	7,384	7,437	5,694	2,954	925	108	1
2001	3,371	4,747	6,377	7,509	7,394	5,839	3,044	866	101	-
2000	3,539	4,739	6,804	8,034	7,485	5,923	3,100	899	123	-
1999	3,644	4,874	6,769	8,085	7,342	5,856	3,121	839	107	-
Male										
2002	2,198	2,763	3,426	3,805	3,406	2,205	883	227	18	1
2001	1,986	2,793	3,576	3,871	3,392	2,211	922	198	28	-
2000	2,054	2,752	3,801	4,178	3,361	2,249	874	207	23	-
1999	2,138	2,791	3,768	4,150	3,406	2,215	957	203	21	-
Female										
2002	1,415	1,895	2,698	3,579	4,031	3,489	2,071	698	90	-
2001	1,385	1,954	2,801	3,638	4,002	3,628	2,122	668	73	-
2000	1,485	1,987	3,003	3,856	4,124	3,674	2,226	692	100	-
1999	1,506	2,083	3,001	3,935	3,936	3,641	2,164	636	86	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of colon, rectum and anus (C18-C21)														
Non-Hispanic Black														
Both sexes														
2002	6,817	-	-	-	-	2	8	11	27	75	208	367	490	591
2001	6,847	-	-	-	-	2	6	13	43	97	183	353	492	607
2000	6,820	-	-	-	-	-	6	12	42	109	181	357	462	534
1999	6,709	1	-	-	-	4	10	20	35	90	178	301	410	547
Male														
2002	3,322	-	-	-	-	2	6	5	18	43	113	184	268	317
2001	3,221	-	-	-	-	1	6	6	24	53	99	186	242	325
2000	3,307	-	-	-	-	-	5	4	22	56	104	180	250	309
1999	3,160	-	-	-	-	2	7	13	17	42	81	158	216	315
Female														
2002	3,495	-	-	-	-	-	2	6	9	32	95	183	222	274
2001	3,626	-	-	-	-	1	-	7	19	44	84	167	250	282
2000	3,513	-	-	-	-	-	1	8	20	53	77	177	212	225
1999	3,549	1	-	-	-	2	3	7	18	48	97	143	194	232
Origin not stated[2]														
Both sexes														
2002	133	-	-	-	-	-	-	1	3	1	4	4	6	9
2001	124	-	-	-	-	-	-	-	-	3	2	6	8	13
2000	116	-	-	-	-	-	-	-	1	1	3	7	4	12
1999	148	-	-	-	-	-	1	-	-	2	4	4	4	9
Male														
2002	74	-	-	-	-	-	-	-	2	1	4	4	2	5
2001	62	-	-	-	-	-	-	-	-	2	2	3	3	9
2000	51	-	-	-	-	-	-	-	1	1	1	3	1	2
1999	75	-	-	-	-	-	-	-	-	-	3	1	3	6
Female														
2002	59	-	-	-	-	-	-	1	1	-	-	-	4	4
2001	62	-	-	-	-	-	-	-	-	1	-	3	5	4
2000	65	-	-	-	-	-	-	-	-	-	2	4	3	10
1999	73	-	-	-	-	-	1	-	-	2	1	3	1	3
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
All origins														
Both sexes														
2002	14,047	3	28	7	11	10	17	33	72	123	291	731	1,245	1,208
2001	13,351	5	18	7	8	14	15	29	46	137	264	795	1,073	1,063
2000	12,916	2	23	9	11	15	15	28	58	113	316	742	944	1,040
1999	12,382	8	29	5	6	19	24	33	45	127	317	666	826	928

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of colon, rectum and anus (C18-C21)										
Non-Hispanic Black										
Both sexes										
2002	706	809	847	927	829	505	308	94	13	-
2001	641	839	887	985	773	503	296	104	23	-
2000	681	782	938	936	796	556	293	117	18	-
1999	701	808	916	958	775	573	266	92	24	-
Male										
2002	400	433	407	452	370	195	86	19	4	-
2001	327	449	432	442	324	187	86	27	5	-
2000	367	417	476	440	352	198	81	39	7	-
1999	377	420	430	475	311	205	76	15	-	-
Female										
2002	306	376	440	475	459	310	222	75	9	-
2001	314	390	455	543	449	316	210	77	18	-
2000	314	365	462	496	444	358	212	78	11	-
1999	324	388	486	483	464	368	190	77	24	-
Origin not stated[2]										
Both sexes										
2002	10	11	18	22	21	8	13	2	-	-
2001	11	9	18	16	13	15	7	2	1	-
2000	10	15	12	17	12	11	9	2	-	-
1999	10	13	15	19	35	25	7	-	-	-
Male										
2002	8	8	13	13	8	2	4	-	-	-
2001	8	7	14	5	4	3	1	-	1	-
2000	6	10	4	10	7	1	4	-	-	-
1999	6	8	6	11	19	9	3	-	-	-
Female										
2002	2	3	5	9	13	6	9	2	-	-
2001	3	2	4	11	9	12	6	2	-	-
2000	4	5	8	7	5	10	5	2	-	-
1999	4	5	9	8	16	16	4	-	-	-
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
All origins										
Both sexes										
2002	1,350	1,618	1,963	2,049	1,679	1,094	419	80	16	-
2001	1,288	1,556	1,884	2,003	1,646	1,009	386	97	8	-
2000	1,216	1,526	1,913	2,011	1,547	943	359	81	4	-
1999	1,163	1,558	1,933	1,914	1,511	863	325	75	7	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
All origins														
Male														
2002	9,011	2	19	4	8	9	9	20	42	85	223	596	1,015	899
2001	8,483	3	7	4	3	6	9	16	27	100	202	618	872	780
2000	8,176	1	14	6	2	11	9	18	41	74	223	595	729	761
1999	7,689	5	20	5	4	12	12	22	26	88	230	533	628	671
Female														
2002	5,036	1	9	3	3	1	8	13	30	38	68	135	230	309
2001	4,868	2	11	3	5	8	6	13	19	37	62	177	201	283
2000	4,740	1	9	3	9	4	6	10	17	39	93	147	215	279
1999	4,693	3	9	-	2	7	12	11	19	39	87	133	198	257
Hispanic														
Both sexes														
2002	1,363	-	3	-	-	-	1	1	7	11	37	95	122	146
2001	1,256	1	3	1	2	1	1	5	9	19	27	107	119	127
2000	1,176	1	3	3	4	2	2	2	5	14	45	92	84	111
1999	1,084	1	4	1	1	2	2	3	4	13	30	76	94	89
Male														
2002	891	-	2	-	-	-	1	-	4	7	26	74	105	114
2001	812	1	1	-	-	1	-	2	5	12	24	96	91	96
2000	762	1	2	3	1	2	1	2	3	10	35	72	69	82
1999	688	1	3	1	1	1	2	3	2	6	21	58	83	69
Female														
2002	472	-	1	-	-	-	-	1	3	4	11	21	17	32
2001	444	-	2	1	2	-	1	3	4	7	3	11	28	31
2000	414	-	1	-	3	-	1	-	2	4	10	20	15	29
1999	396	-	1	-	-	1	-	-	2	7	9	18	11	20
Non-Hispanic[1]														
Both sexes														
2002	12,638	3	25	7	11	10	16	31	65	112	253	632	1,119	1,058
2001	12,053	4	15	6	6	13	14	24	37	116	236	686	951	929
2000	11,702	1	20	6	7	13	13	25	53	98	270	646	856	929
1999	11,265	7	23	4	5	17	22	30	41	114	284	588	732	833
Male														
2002	8,085	2	17	4	8	9	8	19	38	78	196	518	908	782
2001	7,640	2	6	4	3	5	9	14	22	86	177	520	779	677
2000	7,384	-	12	3	1	9	8	15	38	63	188	519	656	679
1999	6,976	4	15	4	3	11	10	19	24	82	208	474	545	598
Female														
2002	4,553	1	8	3	3	1	8	12	27	34	57	114	211	276

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
All origins										
Male										
2002	959	1,100	1,267	1,212	900	463	155	22	2	-
2001	911	1,036	1,188	1,200	847	481	140	31	2	-
2000	846	1,039	1,207	1,196	856	407	114	26	1	-
1999	794	1,041	1,243	1,103	774	352	109	17	-	-
Female										
2002	391	518	696	837	779	631	264	58	14	-
2001	377	520	696	803	799	528	246	66	6	-
2000	370	487	706	815	691	536	245	55	3	-
1999	369	517	690	811	737	511	216	58	7	-
Hispanic										
Both sexes										
2002	164	198	205	165	100	68	38	2	-	-
2001	134	157	170	179	110	59	21	4	-	-
2000	123	145	190	150	112	61	18	9	-	-
1999	125	169	155	135	104	49	20	5	2	-
Male										
2002	111	130	126	97	49	31	13	1	-	-
2001	91	100	106	100	52	24	9	1	-	-
2000	83	95	113	88	64	23	10	3	-	-
1999	90	94	101	72	49	18	12	1	-	-
Female										
2002	53	68	79	68	51	37	25	1	-	-
2001	43	57	64	79	58	35	12	3	-	-
2000	40	50	77	62	48	38	8	6	-	-
1999	35	75	54	63	55	31	8	4	2	-
Non-Hispanic[1]										
Both sexes										
2002	1,176	1,416	1,755	1,878	1,574	1,022	381	78	16	-
2001	1,150	1,395	1,708	1,817	1,535	947	364	93	7	-
2000	1,089	1,375	1,716	1,857	1,430	881	341	72	4	-
1999	1,036	1,385	1,773	1,777	1,403	812	304	70	5	-
Male										
2002	839	967	1,139	1,111	846	431	142	21	2	-
2001	818	932	1,077	1,095	794	457	131	30	2	-
2000	761	938	1,090	1,105	788	383	104	23	1	-
1999	702	944	1,137	1,029	723	332	96	16	-	-
Female										
2002	337	449	616	767	728	591	239	57	14	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
Non-Hispanic[1]														
Female														
2001	4,413	2	9	2	3	8	5	10	15	30	59	166	172	252
2000	4,318	1	8	3	6	4	5	10	15	35	82	127	200	250
1999	4,289	3	8	-	2	6	12	11	17	32	76	114	187	235
Non-Hispanic White														
Both sexes														
2002	9,878	2	19	7	6	6	12	17	32	63	164	427	740	729
2001	9,433	4	11	1	4	9	11	14	20	62	127	445	644	640
2000	9,121	-	14	6	2	8	8	10	27	51	150	420	567	647
1999	8,890	5	20	2	2	12	13	14	15	60	178	371	509	571
Male														
2002	6,215	2	12	4	3	5	4	8	15	41	127	353	598	537
2001	5,956	2	4	1	3	4	7	7	10	43	94	334	525	460
2000	5,679	-	9	3	1	5	5	6	16	27	107	331	428	462
1999	5,455	2	13	2	1	9	5	8	8	39	134	293	367	411
Female														
2002	3,663	-	7	3	3	1	8	9	17	22	37	74	142	192
2001	3,477	2	7	-	1	5	4	7	10	19	33	111	119	180
2000	3,442	-	5	3	1	3	3	4	11	24	43	89	139	185
1999	3,435	3	7	-	1	3	8	6	7	21	44	78	142	160
Non-Hispanic Black														
Both sexes														
2002	1,708	1	4	-	3	3	1	8	18	23	60	144	256	227
2001	1,649	-	3	3	2	2	2	6	9	29	66	173	212	215
2000	1,581	1	2	-	2	4	2	9	11	24	77	162	198	171
1999	1,498	1	3	1	3	2	5	9	12	31	60	149	142	156
Male														
2002	1,143	-	3	-	3	3	1	5	11	16	44	113	208	167
2001	1,036	-	2	2	-	-	1	4	6	22	48	131	171	162
2000	1,047	-	1	-	-	3	2	4	10	15	50	134	151	134
1999	939	1	2	1	2	-	2	7	7	21	44	124	114	109
Female														
2002	565	1	1	-	-	-	-	3	7	7	16	31	48	60
2001	613	-	1	1	2	2	1	2	3	7	18	42	41	53
2000	534	1	1	-	2	1	-	5	1	9	27	28	47	37
1999	559	-	1	-	1	2	3	2	5	10	16	25	28	47
Origin not stated[2]														
Both sexes														
2002	46	-	-	-	-	-	-	1	-	-	1	4	4	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
Non-Hispanic[1]										
Female										
2001	332	463	631	722	741	490	233	63	5	-
2000	328	437	626	752	642	498	237	49	3	-
1999	334	441	636	748	680	480	208	54	5	-
Non-Hispanic White										
Both sexes										
2002	884	1,100	1,404	1,603	1,367	904	316	65	11	-
2001	865	1,084	1,393	1,552	1,305	835	323	78	6	-
2000	800	1,061	1,387	1,565	1,259	769	307	61	2	-
1999	769	1,089	1,457	1,524	1,231	715	270	58	5	-
Male										
2002	630	759	913	953	728	387	118	17	1	-
2001	628	752	899	938	694	403	120	26	2	-
2000	558	733	891	957	698	329	95	18	-	-
1999	523	753	949	887	647	301	90	13	-	-
Female										
2002	254	341	491	650	639	517	198	48	10	-
2001	237	332	494	614	611	432	203	52	4	-
2000	242	328	496	608	561	440	212	43	2	-
1999	246	336	508	637	584	414	180	45	5	-
Non-Hispanic Black										
Both sexes										
2002	158	180	204	171	121	75	43	6	2	-
2001	158	182	175	148	151	71	31	10	1	-
2000	173	183	200	161	101	70	21	7	2	-
1999	156	189	206	148	117	69	29	10	-	-
Male										
2002	112	126	135	92	66	23	12	3	-	-
2001	104	108	96	81	61	31	6	-	-	-
2000	122	126	123	77	54	33	4	3	1	-
1999	101	122	123	84	49	18	6	2	-	-
Female										
2002	46	54	69	79	55	52	31	3	2	-
2001	54	74	79	67	90	40	25	10	1	-
2000	51	57	77	84	47	37	17	4	1	-
1999	55	67	83	64	68	51	23	8	-	-
Origin not stated[2]										
Both sexes										
2002	10	4	3	6	5	4	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)														
Origin not stated[2]														
Both sexes														
2001	42	-	-	-	-	-	-	-	-	2	1	2	3	7
2000	38	-	-	-	-	-	-	1	-	1	1	4	4	-
1999	33	-	2	-	-	-	-	-	-	-	3	2	-	6
Male														
2002	35	-	-	-	-	-	-	1	-	-	1	4	2	3
2001	31	-	-	-	-	-	-	-	-	2	1	2	2	7
2000	30	-	-	-	-	-	-	1	-	1	-	4	4	-
1999	25	-	2	-	-	-	-	-	-	-	1	1	-	4
Female														
2002	11	-	-	-	-	-	-	-	-	-	-	-	2	1
2001	11	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	8	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	8	-	-	-	-	-	-	-	-	-	2	1	-	2
Malignant neoplasm of pancreas (C25)														
All origins														
Both sexes														
2002	30,264	1	-	-	-	2	5	7	32	156	440	889	1,544	2,321
2001	29,803	-	1	1	1	1	1	11	44	167	407	833	1,577	2,111
2000	29,332	2	1	-	1	1	4	17	38	140	408	834	1,528	2,030
1999	29,082	1	-	-	-	-	3	9	46	178	416	887	1,466	2,039
Male														
2002	14,877	-	-	-	-	2	3	3	23	89	275	555	971	1,394
2001	14,467	-	1	-	-	1	1	5	25	99	222	503	964	1,269
2000	14,238	1	1	-	-	1	2	11	22	87	237	531	955	1,206
1999	14,176	-	-	-	-	-	1	7	26	107	262	575	907	1,204
Female														
2002	15,387	1	-	-	-	-	2	4	9	67	165	334	573	927
2001	15,336	-	-	1	1	-	-	6	19	68	185	330	613	842
2000	15,094	1	-	-	1	-	2	6	16	53	171	303	573	824
1999	14,906	1	-	-	-	-	2	2	20	71	154	312	559	835
Hispanic														
Both sexes														
2002	1,363	-	-	-	-	-	1	-	4	17	34	59	89	127
2001	1,336	-	-	-	-	1	-	2	4	16	30	47	80	122
2000	1,243	1	-	-	-	-	2	2	4	13	24	53	75	92
1999	1,135	-	-	-	-	-	-	1	5	18	26	63	74	95
Male														
2002	667	-	-	-	-	-	1	-	2	14	21	39	57	71

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of liver and intrahepatic bile ducts (C22)										
Origin not stated[2]										
Both sexes										
2001	4	4	6	7	1	3	1	-	1	-
2000	4	6	7	4	5	1	-	-	-	-
1999	2	4	5	2	4	2	1	-	-	-
Male										
2002	9	3	2	4	5	1	-	-	-	-
2001	2	4	5	5	1	-	-	-	-	-
2000	2	6	4	3	4	1	-	-	-	-
1999	2	3	5	2	2	2	1	-	-	-
Female										
2002	1	1	1	2	-	3	-	-	-	-
2001	2	-	1	2	-	3	1	-	1	-
2000	2	-	3	1	1	-	-	-	-	-
1999	-	1	-	-	2	-	-	-	-	-
Malignant neoplasm of pancreas (C25)										
All origins										
Both sexes										
2002	2,784	3,564	4,623	5,210	4,345	2,824	1,217	263	36	1
2001	2,757	3,525	4,633	4,987	4,335	2,861	1,214	293	42	1
2000	2,678	3,546	4,557	5,223	4,236	2,571	1,196	286	34	1
1999	2,668	3,632	4,661	4,931	4,002	2,673	1,120	309	40	1
Male										
2002	1,618	1,924	2,346	2,424	1,846	985	351	60	7	1
2001	1,605	1,921	2,294	2,360	1,831	984	310	64	7	1
2000	1,543	1,965	2,224	2,447	1,742	891	310	59	2	1
1999	1,517	1,976	2,371	2,281	1,659	910	311	59	3	-
Female										
2002	1,166	1,640	2,277	2,786	2,499	1,839	866	203	29	-
2001	1,152	1,604	2,339	2,627	2,504	1,877	904	229	35	-
2000	1,135	1,581	2,333	2,776	2,494	1,680	886	227	32	-
1999	1,151	1,656	2,290	2,650	2,343	1,763	809	250	37	1
Hispanic										
Both sexes										
2002	146	176	234	202	131	97	35	10	1	-
2001	141	184	235	196	144	85	39	10	-	-
2000	142	176	221	185	128	74	44	7	-	-
1999	123	155	183	172	100	73	33	14	-	-
Male										
2002	76	97	117	85	51	27	7	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of pancreas (C25)														
Hispanic														
Male														
2001	643	-	-	-	-	1	-	1	2	9	17	28	45	68
2000	620	1	-	-	-	-	1	1	1	8	13	37	43	58
1999	563	-	-	-	-	-	-	-	2	12	18	43	41	58
Female														
2002	696	-	-	-	-	-	-	-	2	3	13	20	32	56
2001	693	-	-	-	-	-	-	1	2	7	13	19	35	54
2000	623	-	-	-	-	-	1	1	3	5	11	16	32	34
1999	572	-	-	-	-	-	-	1	3	6	8	20	33	37
Non-Hispanic[1]														
Both sexes														
2002	28,839	1	-	-	-	2	4	7	28	139	406	827	1,451	2,187
2001	28,407	-	1	1	1	-	1	9	39	151	375	786	1,493	1,987
2000	28,026	1	1	-	1	1	2	15	34	127	383	778	1,450	1,934
1999	27,869	1	-	-	-	-	3	8	41	160	387	820	1,387	1,942
Male														
2002	14,181	-	-	-	-	2	2	3	21	75	254	516	912	1,317
2001	13,792	-	1	-	-	-	1	4	23	90	204	475	918	1,200
2000	13,590	-	1	-	-	1	1	10	21	79	223	492	909	1,145
1999	13,572	-	-	-	-	-	1	7	24	95	243	528	863	1,145
Female														
2002	14,658	1	-	-	-	-	2	4	7	64	152	311	539	870
2001	14,615	-	-	1	1	-	-	5	16	61	171	311	575	787
2000	14,436	1	-	-	1	-	1	5	13	48	160	286	541	789
1999	14,297	1	-	-	-	-	2	1	17	65	144	292	524	797
Non-Hispanic White														
Both sexes														
2002	24,647	-	-	-	-	2	4	4	21	103	321	656	1,133	1,795
2001	24,424	-	1	1	1	-	1	5	31	119	296	606	1,210	1,632
2000	24,034	1	1	-	1	1	1	13	26	97	294	625	1,172	1,584
1999	23,903	-	-	-	-	-	2	6	28	119	302	663	1,108	1,564
Male														
2002	12,242	-	-	-	-	2	2	1	16	52	205	413	728	1,091
2001	11,994	-	1	-	-	-	1	3	19	69	164	376	755	1,014
2000	11,764	-	1	-	-	1	1	8	17	61	175	399	742	946
1999	11,684	-	-	-	-	-	1	6	17	73	193	430	684	920
Female														
2002	12,405	-	-	-	-	-	2	3	5	51	116	243	405	704
2001	12,430	-	-	1	1	-	-	2	12	50	132	230	455	618
2000	12,270	1	-	-	1	-	-	5	9	36	119	226	430	638

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of pancreas (C25)										
Hispanic										
Male										
2001	73	102	110	94	60	22	7	4	-	-
2000	75	106	108	78	50	21	15	4	-	-
1999	59	85	97	68	40	28	7	5	-	-
Female										
2002	70	79	117	117	80	70	28	8	1	-
2001	68	82	125	102	84	63	32	6	-	-
2000	67	70	113	107	78	53	29	3	-	-
1999	64	70	86	104	60	45	26	9	-	-
Non-Hispanic[1]										
Both sexes										
2002	2,634	3,380	4,383	4,996	4,204	2,721	1,180	253	35	1
2001	2,611	3,331	4,388	4,779	4,184	2,771	1,174	282	42	1
2000	2,531	3,365	4,327	5,029	4,095	2,490	1,149	278	34	1
1999	2,538	3,467	4,461	4,747	3,891	2,595	1,086	294	40	1
Male										
2002	1,539	1,825	2,226	2,334	1,792	954	343	58	7	1
2001	1,528	1,813	2,176	2,262	1,769	959	302	59	7	1
2000	1,464	1,857	2,113	2,367	1,687	869	293	55	2	1
1999	1,453	1,887	2,263	2,206	1,616	880	304	54	3	-
Female										
2002	1,095	1,555	2,157	2,662	2,412	1,767	837	195	28	-
2001	1,083	1,518	2,212	2,517	2,415	1,812	872	223	35	-
2000	1,067	1,508	2,214	2,662	2,408	1,621	856	223	32	-
1999	1,085	1,580	2,198	2,541	2,275	1,715	782	240	37	1
Non-Hispanic White										
Both sexes										
2002	2,176	2,798	3,735	4,369	3,765	2,450	1,067	219	29	-
2001	2,215	2,802	3,762	4,199	3,732	2,493	1,034	250	33	1
2000	2,106	2,824	3,721	4,416	3,650	2,218	1,013	242	27	1
1999	2,111	2,926	3,858	4,160	3,465	2,322	971	267	30	1
Male										
2002	1,301	1,549	1,931	2,088	1,627	869	312	50	5	-
2001	1,310	1,544	1,897	2,026	1,598	883	273	55	5	1
2000	1,237	1,603	1,841	2,120	1,518	790	256	45	2	1
1999	1,224	1,613	1,992	1,958	1,454	792	273	51	3	-
Female										
2002	875	1,249	1,804	2,281	2,138	1,581	755	169	24	-
2001	905	1,258	1,865	2,173	2,134	1,610	761	195	28	-
2000	869	1,221	1,880	2,296	2,132	1,428	757	197	25	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of pancreas (C25)														
Non-Hispanic White														
Female														
1999	12,219	-	-	-	-	-	1	-	11	46	109	233	424	644
Non-Hispanic Black														
Both sexes														
2002	3,466	1	-	-	-	-	-	2	4	30	71	142	257	332
2001	3,352	-	-	-	-	-	-	4	5	26	63	156	243	291
2000	3,391	-	-	-	-	-	1	1	8	24	73	134	245	302
1999	3,368	1	-	-	-	-	-	2	10	36	76	141	247	327
Male														
2002	1,606	-	-	-	-	-	-	1	4	19	39	85	157	193
2001	1,501	-	-	-	-	-	-	1	3	16	31	89	140	153
2000	1,546	-	-	-	-	-	-	1	4	15	39	79	150	174
1999	1,601	-	-	-	-	-	-	1	7	19	44	88	158	200
Female														
2002	1,860	1	-	-	-	-	-	1	-	11	32	57	100	139
2001	1,851	-	-	-	-	-	-	3	2	10	32	67	103	138
2000	1,845	-	-	-	-	-	1	-	4	9	34	55	95	128
1999	1,767	1	-	-	-	-	-	1	3	17	32	53	89	127
Origin not stated[2]														
Both sexes														
2002	62	-	-	-	-	-	-	-	-	-	-	3	4	7
2001	60	-	-	-	-	-	-	-	1	-	2	-	4	2
2000	63	-	-	-	-	-	-	-	-	-	1	3	3	4
1999	78	-	-	-	-	-	-	-	-	-	3	4	5	2
Male														
2002	29	-	-	-	-	-	-	-	-	-	-	-	2	6
2001	32	-	-	-	-	-	-	-	-	-	1	-	1	1
2000	28	-	-	-	-	-	-	-	-	-	1	2	3	3
1999	41	-	-	-	-	-	-	-	-	-	1	4	3	1
Female														
2002	33	-	-	-	-	-	-	-	-	-	-	3	2	1
2001	28	-	-	-	-	-	-	-	1	-	1	-	3	1
2000	35	-	-	-	-	-	-	-	-	-	-	1	-	1
1999	37	-	-	-	-	-	-	-	-	-	2	-	2	1
Malignant neoplasm of larynx (C32)														
All origins														
Both sexes														
2002	3,723	-	-	-	-	-	-	3	8	19	57	156	269	412
2001	3,797	-	-	-	-	-	-	1	2	18	68	149	270	414

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of pancreas (C25)										
Non-Hispanic White										
Female										
1999	887	1,313	1,866	2,202	2,011	1,530	698	216	27	1
Non-Hispanic Black										
Both sexes										
2002	394	471	545	509	349	225	95	32	6	1
2001	347	442	519	481	384	231	122	29	9	-
2000	362	452	518	524	366	227	117	32	5	-
1999	342	459	513	486	360	234	99	25	10	-
Male										
2002	206	225	250	196	136	62	22	8	2	1
2001	186	231	229	200	135	59	22	4	2	-
2000	193	213	234	213	137	60	26	8	-	-
1999	186	230	227	210	134	72	22	3	-	-
Female										
2002	188	246	295	313	213	163	73	24	4	-
2001	161	211	290	281	249	172	100	25	7	-
2000	169	239	284	311	229	167	91	24	5	-
1999	156	229	286	276	226	162	77	22	10	-
Origin not stated[2]										
Both sexes										
2002	4	8	6	12	10	6	2	-	-	-
2001	5	10	10	12	7	5	1	1	-	-
2000	5	5	9	9	13	7	3	1	-	-
1999	7	10	17	12	11	5	1	1	-	-
Male										
2002	3	2	3	5	3	4	1	-	-	-
2001	4	6	8	4	2	3	1	1	-	-
2000	4	2	3	2	5	1	2	-	-	-
1999	5	4	11	7	3	2	-	-	-	-
Female										
2002	1	6	3	7	7	2	1	-	-	-
2001	1	4	2	8	5	2	-	-	-	-
2000	1	3	6	7	8	6	1	1	-	-
1999	2	6	6	5	8	3	1	1	-	-
Malignant neoplasm of larynx (C32)										
All origins										
Both sexes										
2002	492	508	576	588	381	172	76	5	-	1
2001	499	531	639	553	357	208	74	13	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of larynx (C32)														
All origins														
Both sexes														
2000	3,861	-	-	-	2	1	-	-	4	9	55	156	270	384
1999	3,815	-	-	-	-	-	-	-	6	20	58	151	254	410
Male														
2002	2,922	-	-	-	-	-	-	3	3	16	42	130	219	339
2001	3,018	-	-	-	-	-	-	1	1	12	54	123	223	339
2000	3,041	-	-	-	-	-	-	-	2	6	43	122	218	319
1999	2,992	-	-	-	-	-	-	-	5	18	46	130	216	339
Female														
2002	801	-	-	-	-	-	-	-	5	3	15	26	50	73
2001	779	-	-	-	-	-	-	-	1	6	14	26	47	75
2000	820	-	-	-	2	1	-	-	2	3	12	34	52	65
1999	823	-	-	-	-	-	-	-	1	2	12	21	38	71
Hispanic														
Both sexes														
2002	165	-	-	-	-	-	-	-	1	3	5	7	12	21
2001	163	-	-	-	-	-	-	-	-	-	3	14	6	18
2000	143	-	-	-	1	-	-	-	1	1	3	5	12	13
1999	171	-	-	-	-	-	-	-	-	-	5	15	10	17
Male														
2002	139	-	-	-	-	-	-	-	-	3	4	6	10	21
2001	141	-	-	-	-	-	-	-	-	-	3	13	5	16
2000	124	-	-	-	-	-	-	-	-	1	2	5	11	13
1999	149	-	-	-	-	-	-	-	-	-	4	15	9	16
Female														
2002	26	-	-	-	-	-	-	-	1	-	1	1	2	-
2001	22	-	-	-	-	-	-	-	-	-	-	1	1	2
2000	19	-	-	-	1	-	-	-	1	-	1	-	1	-
1999	22	-	-	-	-	-	-	-	-	-	1	-	1	1
Non-Hispanic[1]														
Both sexes														
2002	3,548	-	-	-	-	-	-	3	7	16	52	149	256	388
2001	3,614	-	-	-	-	-	-	1	2	18	64	135	263	390
2000	3,699	-	-	-	1	1	-	-	3	8	51	150	256	370
1999	3,627	-	-	-	-	-	-	-	6	20	53	135	242	393
Male														
2002	2,775	-	-	-	-	-	-	3	3	13	38	124	208	315
2001	2,858	-	-	-	-	-	-	1	1	12	50	110	217	318
2000	2,901	-	-	-	-	-	-	-	2	5	40	116	206	305
1999	2,832	-	-	-	-	-	-	-	5	18	42	114	206	323

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of larynx (C32)										
All origins										
Both sexes										
2000	489	574	645	595	401	191	73	11	1	-
1999	494	584	655	529	363	200	72	16	3	-
Male										
2002	399	395	437	455	300	126	56	1	-	1
2001	398	417	502	442	268	171	58	8	1	-
2000	407	441	504	451	314	155	51	8	-	-
1999	396	458	490	397	284	154	46	10	3	-
Female										
2002	93	113	139	133	81	46	20	4	-	-
2001	101	114	137	111	89	37	16	5	-	-
2000	82	133	141	144	87	36	22	3	1	-
1999	98	126	165	132	79	46	26	6	-	-
Hispanic										
Both sexes										
2002	22	21	27	24	9	7	6	-	-	-
2001	20	35	25	20	9	9	4	-	-	-
2000	16	27	18	15	20	8	3	-	-	-
1999	12	30	25	24	16	11	4	2	-	-
Male										
2002	19	19	23	19	8	2	5	-	-	-
2001	18	29	24	16	8	5	4	-	-	-
2000	15	23	14	12	18	7	3	-	-	-
1999	11	26	22	20	11	10	4	1	-	-
Female										
2002	3	2	4	5	1	5	1	-	-	-
2001	2	6	1	4	1	4	-	-	-	-
2000	1	4	4	3	2	1	-	-	-	-
1999	1	4	3	4	5	1	-	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	469	486	547	563	371	165	70	5	-	1
2001	475	494	612	530	347	199	70	13	1	-
2000	470	543	623	579	380	182	70	11	1	-
1999	478	552	628	503	344	189	67	14	3	-
Male										
2002	380	376	412	435	291	124	51	1	-	1
2001	376	386	476	423	259	166	54	8	1	-
2000	389	415	486	438	295	148	48	8	-	-
1999	383	430	467	375	271	144	42	9	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of larynx (C32)														
Non-Hispanic[1]														
Female														
2002	773	-	-	-	-	-	-	-	4	3	14	25	48	73
2001	756	-	-	-	-	-	-	-	1	6	14	25	46	72
2000	798	-	-	-	1	1	-	-	1	3	11	34	50	65
1999	795	-	-	-	-	-	-	-	1	2	11	21	36	70
Non-Hispanic White														
Both sexes														
2002	2,838	-	-	-	-	-	-	2	4	7	33	97	179	293
2001	2,876	-	-	-	-	-	-	1	2	11	47	84	184	287
2000	2,940	-	-	-	1	1	-	-	3	7	37	102	179	266
1999	2,924	-	-	-	-	-	-	-	4	14	26	85	171	291
Male														
2002	2,205	-	-	-	-	-	-	2	1	5	25	82	148	234
2001	2,267	-	-	-	-	-	-	1	1	10	37	66	150	233
2000	2,291	-	-	-	-	-	-	-	2	4	27	80	140	215
1999	2,263	-	-	-	-	-	-	-	3	14	20	71	148	236
Female														
2002	633	-	-	-	-	-	-	-	3	2	8	15	31	59
2001	609	-	-	-	-	-	-	-	1	1	10	18	34	54
2000	649	-	-	-	1	1	-	-	1	3	10	22	39	51
1999	661	-	-	-	-	-	-	-	1	-	6	14	23	55
Non-Hispanic Black														
Both sexes														
2002	674	-	-	-	-	-	-	1	3	8	19	48	75	91
2001	668	-	-	-	-	-	-	-	-	7	16	48	76	98
2000	714	-	-	-	-	-	-	-	-	1	14	47	74	101
1999	657	-	-	-	-	-	-	-	2	6	26	48	69	98
Male														
2002	541	-	-	-	-	-	-	1	2	7	13	38	58	77
2001	531	-	-	-	-	-	-	-	-	2	12	41	64	80
2000	574	-	-	-	-	-	-	-	-	1	13	35	64	87
1999	532	-	-	-	-	-	-	-	2	4	21	41	56	85
Female														
2002	133	-	-	-	-	-	-	-	1	1	6	10	17	14
2001	137	-	-	-	-	-	-	-	-	5	4	7	12	18
2000	140	-	-	-	-	-	-	-	-	-	1	12	10	14
1999	125	-	-	-	-	-	-	-	-	2	5	7	13	13
Origin not stated[2]														
Both sexes														
2002	10	-	-	-	-	-	-	-	-	-	-	-	1	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of larynx (C32)										
Non-Hispanic[1]										
Female										
2002	89	110	135	128	80	41	19	4	-	-
2001	99	108	136	107	88	33	16	5	-	-
2000	81	128	137	141	85	34	22	3	1	-
1999	95	122	161	128	73	45	25	5	-	-
Non-Hispanic White										
Both sexes										
2002	360	394	456	469	324	151	65	4	-	-
2001	370	396	497	442	305	174	64	11	1	-
2000	346	436	500	485	337	162	67	10	1	-
1999	369	444	534	443	308	165	54	13	3	-
Male										
2002	284	303	342	361	258	113	47	-	-	-
2001	296	309	388	347	223	147	51	7	1	-
2000	285	331	390	370	262	133	45	7	-	-
1999	298	345	394	330	238	123	32	8	3	-
Female										
2002	76	91	114	108	66	38	18	4	-	-
2001	74	87	109	95	82	27	13	4	-	-
2000	61	105	110	115	75	29	22	3	1	-
1999	71	99	140	113	70	42	22	5	-	-
Non-Hispanic Black										
Both sexes										
2002	107	86	88	89	42	12	3	1	-	1
2001	97	87	101	77	34	20	6	1	-	-
2000	117	95	114	88	41	19	2	1	-	-
1999	105	102	86	57	32	15	10	1	-	-
Male										
2002	95	68	67	72	29	10	2	1	-	1
2001	76	68	76	66	28	15	3	-	-	-
2000	99	75	89	62	32	14	2	1	-	-
1999	83	80	67	42	30	13	7	1	-	-
Female										
2002	12	18	21	17	13	2	1	-	-	-
2001	21	19	25	11	6	5	3	1	-	-
2000	18	20	25	26	9	5	-	-	-	-
1999	22	22	19	15	2	2	3	-	-	-
Origin not stated[2]										
Both sexes										
2002	1	1	2	1	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of larynx (C32)														
Origin not stated[2]														
Both sexes														
2001	20	-	-	-	-	-	-	-	-	-	1	-	1	6
2000	19	-	-	-	-	-	-	-	-	-	1	1	2	1
1999	17	-	-	-	-	-	-	-	-	-	-	1	2	-
Male														
2002	8	-	-	-	-	-	-	-	-	-	-	-	1	3
2001	19	-	-	-	-	-	-	-	-	-	1	-	1	5
2000	16	-	-	-	-	-	-	-	-	-	1	1	1	1
1999	11	-	-	-	-	-	-	-	-	-	-	1	1	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	3	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	6	-	-	-	-	-	-	-	-	-	-	-	1	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)														
All origins														
Both sexes														
2002	157,713	-	1	2	2	4	17	35	130	583	2,115	4,341	7,809	12,930
2001	156,058	-	5	1	1	6	14	36	121	640	2,156	4,255	7,786	12,595
2000	155,521	2	2	-	1	3	19	37	160	664	2,092	4,220	7,695	12,453
1999	152,156	-	-	3	2	6	14	43	136	668	2,056	3,916	7,518	12,165
Male														
2002	90,171	-	1	1	2	2	14	19	71	290	1,096	2,471	4,723	7,769
2001	90,426	-	4	1	-	2	10	17	60	315	1,189	2,487	4,703	7,523
2000	90,469	2	1	-	-	3	11	28	74	370	1,172	2,540	4,583	7,546
1999	89,453	-	-	2	2	5	8	24	73	323	1,151	2,346	4,505	7,465
Female														
2002	67,542	-	-	1	-	2	3	16	59	293	1,019	1,870	3,086	5,161
2001	65,632	-	1	-	1	4	4	19	61	325	967	1,768	3,083	5,072
2000	65,052	-	1	-	1	-	8	9	86	294	920	1,680	3,112	4,907
1999	62,703	-	-	1	-	1	6	19	63	345	905	1,570	3,013	4,700
Hispanic														
Both sexes														
2002	4,038	-	-	2	-	-	4	13	14	23	67	124	238	364
2001	3,816	-	2	-	1	1	6	6	10	27	79	127	237	325
2000	3,643	-	-	-	-	1	2	4	13	28	62	124	218	303
1999	3,545	-	-	-	-	-	3	3	14	32	66	111	197	303
Male														
2002	2,617	-	-	1	-	-	2	5	9	8	37	71	158	256

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of larynx (C32)										
Origin not stated[2]										
Both sexes										
2001	4	2	2	3	1	-	-	-	-	-
2000	3	4	4	1	1	1	-	-	-	-
1999	4	2	2	2	3	-	1	-	-	-
Male										
2002	-	-	2	1	1	-	-	-	-	-
2001	4	2	2	3	1	-	-	-	-	-
2000	3	3	4	1	1	-	-	-	-	-
1999	2	2	1	2	2	-	-	-	-	-
Female										
2002	1	1	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	1	-	-	-	-
1999	2	-	1	-	1	-	1	-	-	-
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All origins										
Both sexes										
2002	17,754	22,358	27,898	27,969	20,113	9,781	3,214	576	81	-
2001	17,182	22,850	28,375	27,723	18,977	9,623	3,064	587	57	4
2000	17,268	23,438	28,820	27,305	18,531	9,302	2,902	542	60	5
1999	17,179	23,239	28,609	26,556	17,705	8,945	2,766	549	79	2
Male										
2002	10,646	13,209	16,334	15,833	10,988	5,022	1,439	216	25	-
2001	10,350	13,798	16,901	15,968	10,438	5,026	1,370	237	23	4
2000	10,391	14,189	17,193	15,726	10,240	4,840	1,335	204	16	5
1999	10,548	14,186	17,101	15,461	9,936	4,739	1,314	234	28	2
Female										
2002	7,108	9,149	11,564	12,136	9,125	4,759	1,775	360	56	-
2001	6,832	9,052	11,474	11,755	8,539	4,597	1,694	350	34	-
2000	6,877	9,249	11,627	11,579	8,291	4,462	1,567	338	44	-
1999	6,631	9,053	11,508	11,095	7,769	4,206	1,452	315	51	-
Hispanic										
Both sexes										
2002	444	574	729	681	420	224	97	16	4	-
2001	433	564	666	626	385	202	90	27	2	-
2000	369	545	646	595	411	204	91	21	6	-
1999	421	556	614	553	348	215	92	15	2	-
Male										
2002	311	393	453	463	266	121	54	6	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)														
Hispanic														
Male														
2001	2,495	-	2	-	-	-	6	3	7	22	46	83	156	219
2000	2,411	-	-	-	-	1	2	2	8	20	36	81	148	210
1999	2,373	-	-	-	-	-	2	3	7	17	41	74	134	219
Female														
2002	1,421	-	-	1	-	-	2	8	5	15	30	53	80	108
2001	1,321	-	-	-	1	1	-	3	3	5	33	44	81	106
2000	1,232	-	-	-	-	-	-	2	5	8	26	43	70	93
1999	1,172	-	-	-	-	-	1	-	7	15	25	37	63	84
Non-Hispanic[1]														
Both sexes														
2002	153,292	-	1	-	2	4	13	22	113	559	2,042	4,204	7,548	12,541
2001	151,882	-	3	1	-	5	8	30	111	611	2,067	4,119	7,523	12,239
2000	151,483	2	2	-	1	2	17	33	147	635	2,018	4,081	7,444	12,110
1999	148,206	-	-	3	2	6	11	40	121	635	1,983	3,795	7,292	11,829
Male														
2002	87,335	-	1	-	2	2	12	14	62	281	1,056	2,392	4,550	7,498
2001	87,706	-	2	1	-	2	4	14	53	293	1,138	2,400	4,529	7,278
2000	87,816	2	1	-	-	2	9	26	66	350	1,129	2,453	4,413	7,307
1999	86,822	-	-	2	2	5	6	21	66	306	1,105	2,266	4,350	7,224
Female														
2002	65,957	-	-	-	-	2	1	8	51	278	986	1,812	2,998	5,043
2001	64,176	-	1	-	-	3	4	16	58	318	929	1,719	2,994	4,961
2000	63,667	-	1	-	1	-	8	7	81	285	889	1,628	3,031	4,803
1999	61,384	-	-	1	-	1	5	19	55	329	878	1,529	2,942	4,605
Non-Hispanic White														
Both sexes														
2002	134,435	-	1	-	1	3	10	12	85	441	1,585	3,294	6,041	10,506
2001	133,266	-	2	1	-	4	5	23	91	494	1,606	3,201	6,100	10,333
2000	133,061	1	2	-	1	-	11	23	116	470	1,526	3,134	6,042	10,244
1999	129,967	-	-	2	1	3	6	30	101	479	1,546	2,910	5,922	10,004
Male														
2002	75,754	-	1	-	1	2	9	9	49	213	803	1,835	3,580	6,189
2001	76,066	-	1	1	-	2	2	12	43	232	863	1,827	3,594	6,032
2000	76,218	1	1	-	-	-	7	18	55	257	819	1,869	3,486	6,053
1999	75,338	-	-	1	1	3	3	14	57	231	847	1,708	3,447	6,023
Female														
2002	58,681	-	-	-	-	1	1	3	36	228	782	1,459	2,461	4,317
2001	57,200	-	1	-	-	2	3	11	48	262	743	1,374	2,506	4,301

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
Hispanic										
Male										
2001	298	369	451	413	244	119	43	13	1	-
2000	261	351	442	412	254	117	55	10	1	-
1999	304	394	404	377	229	107	51	9	1	-
Female										
2002	133	181	276	218	154	103	43	10	1	-
2001	135	195	215	213	141	83	47	14	1	-
2000	108	194	204	183	157	87	36	11	5	-
1999	117	162	210	176	119	108	41	6	1	-
Non-Hispanic[1]										
Both sexes										
2002	17,261	21,725	27,111	27,217	19,645	9,534	3,114	559	77	-
2001	16,697	22,236	27,651	27,041	18,558	9,396	2,968	560	54	4
2000	16,855	22,831	28,109	26,655	18,087	9,076	2,801	518	54	5
1999	16,720	22,620	27,916	25,928	17,315	8,710	2,667	534	77	2
Male										
2002	10,308	12,776	15,852	15,326	10,700	4,889	1,383	209	22	-
2001	10,013	13,402	16,416	15,518	10,177	4,892	1,325	224	21	4
2000	10,102	13,802	16,709	15,279	9,966	4,712	1,277	191	15	5
1999	10,214	13,750	16,647	15,037	9,687	4,620	1,260	225	27	2
Female										
2002	6,953	8,949	11,259	11,891	8,945	4,645	1,731	350	55	-
2001	6,684	8,834	11,235	11,523	8,381	4,504	1,643	336	33	-
2000	6,753	9,029	11,400	11,376	8,121	4,364	1,524	327	39	-
1999	6,506	8,870	11,269	10,891	7,628	4,090	1,407	309	50	-
Non-Hispanic White										
Both sexes										
2002	14,712	18,892	24,212	24,544	17,962	8,732	2,840	502	60	-
2001	14,380	19,332	24,640	24,359	16,864	8,593	2,707	487	42	2
2000	14,533	19,953	24,999	24,044	16,622	8,287	2,552	455	43	3
1999	14,325	19,694	24,883	23,335	15,818	7,934	2,427	483	63	1
Male										
2002	8,629	11,013	14,089	13,707	9,701	4,458	1,253	193	20	-
2001	8,520	11,536	14,520	13,853	9,169	4,432	1,205	202	18	2
2000	8,629	11,945	14,759	13,629	9,089	4,275	1,149	164	10	3
1999	8,631	11,872	14,724	13,441	8,778	4,188	1,143	206	19	1
Female										
2002	6,083	7,879	10,123	10,837	8,261	4,274	1,587	309	40	-
2001	5,860	7,796	10,120	10,506	7,695	4,161	1,502	285	24	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)														
Non-Hispanic White														
Female														
2000	56,843	-	1	-	1	-	4	5	61	213	707	1,265	2,556	4,191
1999	54,629	-	-	1	-	-	3	16	44	248	699	1,202	2,475	3,981
Non-Hispanic Black														
Both sexes														
2002	16,170	-	-	-	1	1	3	4	19	96	407	822	1,343	1,784
2001	15,883	-	1	-	-	1	2	3	13	102	413	800	1,260	1,674
2000	15,924	1	-	-	-	2	3	8	23	142	436	833	1,280	1,638
1999	15,834	-	-	1	1	3	5	5	15	134	397	795	1,234	1,628
Male														
2002	9,983	-	-	-	1	-	3	2	12	57	230	504	874	1,145
2001	10,030	-	1	-	-	-	1	1	6	53	245	511	841	1,099
2000	10,056	1	-	-	-	2	-	6	8	80	279	518	854	1,109
1999	10,039	-	-	1	1	2	3	4	7	63	238	512	826	1,088
Female														
2002	6,187	-	-	-	-	1	-	2	7	39	177	318	469	639
2001	5,853	-	-	-	-	1	1	2	7	49	168	289	419	575
2000	5,868	-	-	-	-	-	3	2	15	62	157	315	426	529
1999	5,795	-	-	-	-	1	2	1	8	71	159	283	408	540
Origin not stated[2]														
Both sexes														
2002	383	-	-	-	-	-	-	-	3	1	6	13	23	25
2001	360	-	-	-	-	-	-	-	-	2	10	9	26	31
2000	395	-	-	-	-	-	-	-	-	1	12	15	33	40
1999	405	-	-	-	-	-	-	-	1	1	7	10	29	33
Male														
2002	219	-	-	-	-	-	-	-	-	1	3	8	15	15
2001	225	-	-	-	-	-	-	-	-	-	5	4	18	26
2000	242	-	-	-	-	-	-	-	-	-	7	6	22	29
1999	258	-	-	-	-	-	-	-	-	-	5	6	21	22
Female														
2002	164	-	-	-	-	-	-	-	3	-	3	5	8	10
2001	135	-	-	-	-	-	-	-	-	2	5	5	8	5
2000	153	-	-	-	-	-	-	-	-	1	5	9	11	11
1999	147	-	-	-	-	-	-	-	1	1	2	4	8	11

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
Non-Hispanic White										
Female										
2000	5,904	8,008	10,240	10,415	7,533	4,012	1,403	291	33	-
1999	5,694	7,822	10,159	9,894	7,040	3,746	1,284	277	44	-
Non-Hispanic Black										
Both sexes										
2002	2,210	2,483	2,447	2,254	1,387	630	220	44	15	-
2001	2,011	2,509	2,567	2,264	1,359	635	195	61	11	2
2000	2,050	2,512	2,666	2,206	1,208	647	206	53	9	1
1999	2,115	2,532	2,579	2,237	1,246	660	196	38	13	-
Male										
2002	1,461	1,575	1,508	1,360	803	334	102	10	2	-
2001	1,301	1,635	1,631	1,410	816	373	86	15	3	2
2000	1,301	1,646	1,663	1,379	726	351	104	23	5	1
1999	1,412	1,637	1,630	1,371	756	376	92	13	7	-
Female										
2002	749	908	939	894	584	296	118	34	13	-
2001	710	874	936	854	543	262	109	46	8	-
2000	749	866	1,003	827	482	296	102	30	4	-
1999	703	895	949	866	490	284	104	25	6	-
Origin not stated[2]										
Both sexes										
2002	49	59	58	71	48	23	3	1	-	-
2001	52	50	58	56	34	25	6	-	1	-
2000	44	62	65	55	33	22	10	3	-	-
1999	38	63	79	75	42	20	7	-	-	-
Male										
2002	27	40	29	44	22	12	2	1	-	-
2001	39	27	34	37	17	15	2	-	1	-
2000	28	36	42	35	20	11	3	3	-	-
1999	30	42	50	47	20	12	3	-	-	-
Female										
2002	22	19	29	27	26	11	1	-	-	-
2001	13	23	24	19	17	10	4	-	-	-
2000	16	26	23	20	13	11	7	-	-	-
1999	8	21	29	28	22	8	4	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant melanoma of skin (C43)														
All origins														
Both sexes														
2002	7,514	-	3	-	1	7	34	70	124	219	385	495	640	659
2001	7,542	-	-	-	-	11	30	72	114	213	409	536	672	653
2000	7,420	-	-	-	1	7	34	58	128	234	392	534	665	627
1999	7,215	1	-	1	-	9	30	62	132	239	386	518	602	615
Male														
2002	4,725	-	2	-	1	3	17	38	81	120	243	343	403	445
2001	4,763	-	-	-	-	9	15	40	63	130	250	331	425	430
2000	4,592	-	-	-	1	4	17	33	77	118	237	321	448	420
1999	4,529	1	-	-	-	7	16	38	72	151	246	337	380	425
Female														
2002	2,789	-	1	-	-	4	17	32	43	99	142	152	237	214
2001	2,779	-	-	-	-	2	15	32	51	83	159	205	247	223
2000	2,828	-	-	-	-	3	17	25	51	116	155	213	217	207
1999	2,686	-	-	1	-	2	14	24	60	88	140	181	222	190
Hispanic														
Both sexes														
2002	147	-	-	-	-	-	2	2	5	7	14	13	14	10
2001	145	-	-	-	-	1	1	3	3	8	16	11	12	14
2000	144	-	-	-	-	-	2	2	3	14	12	15	13	17
1999	133	1	-	1	-	1	1	1	4	6	8	9	13	17
Male														
2002	77	-	-	-	-	-	1	1	5	4	9	8	5	8
2001	78	-	-	-	-	-	-	1	1	7	10	8	3	9
2000	92	-	-	-	-	-	1	1	3	9	8	11	8	13
1999	79	1	-	-	-	1	-	1	3	4	5	7	9	11
Female														
2002	70	-	-	-	-	-	1	1	-	3	5	5	9	2
2001	67	-	-	-	-	1	1	2	2	1	6	3	9	5
2000	52	-	-	-	-	-	1	1	-	5	4	4	5	4
1999	54	-	-	1	-	-	1	-	1	2	3	2	4	6
Non-Hispanic[1]														
Both sexes														
2002	7,349	-	3	-	1	7	31	68	118	211	370	480	626	648
2001	7,387	-	-	-	-	10	29	69	110	204	393	525	659	638
2000	7,263	-	-	-	1	7	32	56	125	219	380	519	648	610
1999	7,062	-	-	-	-	8	29	61	128	233	378	507	587	597
Male														
2002	4,640	-	2	-	1	3	16	37	76	116	234	334	398	437
2001	4,678	-	-	-	-	9	15	39	61	123	240	323	421	420
2000	4,490	-	-	-	1	4	16	32	74	108	229	310	436	407

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)										
All origins										
Both sexes										
2002	692	748	868	954	794	512	233	58	17	1
2001	680	746	943	919	759	489	213	72	11	-
2000	676	753	884	887	737	495	252	52	4	-
1999	648	791	861	945	694	435	181	53	12	-
Male										
2002	470	500	575	609	469	264	115	21	6	-
2001	463	524	632	600	463	256	99	31	2	-
2000	446	495	572	559	454	277	100	13	-	-
1999	426	526	570	596	410	223	90	11	4	-
Female										
2002	222	248	293	345	325	248	118	37	11	1
2001	217	222	311	319	296	233	114	41	9	-
2000	230	258	312	328	283	218	152	39	4	-
1999	222	265	291	349	284	212	91	42	8	-
Hispanic										
Both sexes										
2002	14	9	10	16	13	7	7	3	1	-
2001	11	13	15	15	13	4	2	2	1	-
2000	10	9	17	11	8	7	2	2	-	-
1999	13	15	12	15	5	4	5	2	-	-
Male										
2002	8	4	7	7	6	3	1	-	-	-
2001	6	10	9	9	4	-	-	1	-	-
2000	6	3	11	6	7	4	1	-	-	-
1999	8	10	7	5	4	1	2	-	-	-
Female										
2002	6	5	3	9	7	4	6	3	1	-
2001	5	3	6	6	9	4	2	1	1	-
2000	4	6	6	5	1	3	1	2	-	-
1999	5	5	5	10	1	3	3	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	675	739	857	935	779	504	226	55	16	-
2001	669	733	927	901	745	484	211	70	10	-
2000	666	742	866	875	727	487	249	50	4	-
1999	634	772	848	926	686	431	174	51	12	-
Male										
2002	459	496	567	601	461	261	114	21	6	-
2001	457	514	622	588	459	256	99	30	2	-
2000	440	490	561	552	446	273	98	13	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant melanoma of skin (C43)														
Non-Hispanic[1]														
Male														
1999	4,438	-	-	-	-	6	16	37	69	147	241	328	370	414
Female														
2002	2,709	-	1	-	-	4	15	31	42	95	136	146	228	211
2001	2,709	-	-	-	-	1	14	30	49	81	153	202	238	218
2000	2,773	-	-	-	-	3	16	24	51	111	151	209	212	203
1999	2,624	-	-	-	-	2	13	24	59	86	137	179	217	183
Non-Hispanic White														
Both sexes														
2002	7,190	-	3	-	1	6	31	66	118	202	366	471	614	631
2001	7,250	-	-	-	-	9	29	67	110	202	388	518	650	632
2000	7,090	-	-	-	1	6	32	53	122	214	365	510	637	594
1999	6,936	-	-	-	-	7	27	60	123	230	374	498	583	591
Male														
2002	4,555	-	2	-	1	2	16	35	76	114	230	329	390	425
2001	4,618	-	-	-	-	8	15	37	61	122	237	320	417	417
2000	4,414	-	-	-	1	4	16	31	73	106	221	308	429	398
1999	4,370	-	-	-	-	5	14	36	66	145	240	322	367	410
Female														
2002	2,635	-	1	-	-	4	15	31	42	88	136	142	224	206
2001	2,632	-	-	-	-	1	14	30	49	80	151	198	233	215
2000	2,676	-	-	-	-	2	16	22	49	108	144	202	208	196
1999	2,566	-	-	-	-	2	13	24	57	85	134	176	216	181
Non-Hispanic Black														
Both sexes														
2002	102	-	-	-	-	1	-	1	-	7	2	3	9	13
2001	102	-	-	-	-	1	-	1	-	2	5	3	6	5
2000	134	-	-	-	-	1	-	2	1	4	11	6	7	11
1999	94	-	-	-	-	1	-	1	3	1	3	6	3	5
Male														
2002	52	-	-	-	-	1	-	1	-	2	2	2	5	10
2001	36	-	-	-	-	1	-	1	-	1	3	2	1	3
2000	59	-	-	-	-	-	-	1	1	2	7	1	5	6
1999	47	-	-	-	-	1	-	1	1	1	-	4	2	3
Female														
2002	50	-	-	-	-	-	-	-	-	5	-	1	4	3
2001	66	-	-	-	-	-	-	-	-	1	2	1	5	2
2000	75	-	-	-	-	1	-	1	-	2	4	5	2	5
1999	47	-	-	-	-	-	-	-	2	-	3	2	1	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)										
Non-Hispanic[1]										
Male										
1999	417	514	562	589	405	222	86	11	4	-
Female										
2002	216	243	290	334	318	243	112	34	10	-
2001	212	219	305	313	286	228	112	40	8	-
2000	226	252	305	323	281	214	151	37	4	-
1999	217	258	286	337	281	209	88	40	8	-
Non-Hispanic White										
Both sexes										
2002	660	722	832	919	768	493	220	52	15	-
2001	654	711	905	885	729	476	207	69	9	-
2000	653	726	844	859	707	475	240	48	4	-
1999	622	760	833	905	672	420	169	50	12	-
Male										
2002	451	486	553	592	458	256	112	21	6	-
2001	450	503	613	580	454	253	99	30	2	-
2000	435	480	552	545	436	272	94	13	-	-
1999	409	508	556	577	399	218	84	10	4	-
Female										
2002	209	236	279	327	310	237	108	31	9	-
2001	204	208	292	305	275	223	108	39	7	-
2000	218	246	292	314	271	203	146	35	4	-
1999	213	252	277	328	273	202	85	40	8	-
Non-Hispanic Black										
Both sexes										
2002	8	11	14	10	8	10	5	-	-	-
2001	8	18	18	9	15	5	4	1	1	-
2000	8	11	19	14	18	12	7	2	-	-
1999	9	9	14	14	11	8	5	1	-	-
Male										
2002	3	5	7	6	2	4	2	-	-	-
2001	1	9	7	2	4	1	-	-	-	-
2000	4	7	7	6	9	1	2	-	-	-
1999	7	3	6	8	4	3	2	1	-	-
Female										
2002	5	6	7	4	6	6	3	-	-	-
2001	7	9	11	7	11	4	4	1	1	-
2000	4	4	12	8	9	11	5	2	-	-
1999	2	6	8	6	7	5	3	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant melanoma of skin (C43)														
Origin not stated[2]														
Both sexes														
2002	18	-	-	-	-	-	1	-	1	1	1	2	-	1
2001	10	-	-	-	-	-	-	-	1	1	-	-	1	1
2000	13	-	-	-	-	-	-	-	-	1	-	-	4	-
1999	20	-	-	-	-	-	-	-	-	-	-	2	2	1
Male														
2002	8	-	-	-	-	-	-	-	-	-	-	1	-	-
2001	7	-	-	-	-	-	-	-	1	-	-	-	1	1
2000	10	-	-	-	-	-	-	-	-	1	-	-	4	-
1999	12	-	-	-	-	-	-	-	-	-	-	2	1	-
Female														
2002	10	-	-	-	-	-	1	-	1	1	1	1	-	1
2001	3	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	8	-	-	-	-	-	-	-	-	-	-	-	1	1
Malignant neoplasm of breast (C50)														
All origins														
Both sexes														
2002	41,883	-	-	-	-	1	9	63	362	904	1,818	2,747	3,707	3,924
2001	41,809	-	-	-	-	1	8	88	375	960	1,863	2,790	3,803	3,843
2000	42,300	-	-	-	-	1	10	83	384	982	1,859	2,730	3,655	3,790
1999	41,528	-	1	-	1	3	10	86	353	986	1,755	2,619	3,555	3,682
Male														
2002	369	-	-	-	-	-	-	2	1	4	8	16	28	29
2001	415	-	-	-	-	-	-	-	2	5	9	13	33	32
2000	428	-	-	-	-	-	-	2	1	12	7	18	28	44
1999	384	-	-	-	-	-	-	-	3	6	9	11	30	36
Female														
2002	41,514	-	-	-	-	1	9	61	361	900	1,810	2,731	3,679	3,895
2001	41,394	-	-	-	-	1	8	88	373	955	1,854	2,777	3,770	3,811
2000	41,872	-	-	-	-	1	10	81	383	970	1,852	2,712	3,627	3,746
1999	41,144	-	1	-	1	3	10	86	350	980	1,746	2,608	3,525	3,646
Hispanic														
Both sexes														
2002	1,736	-	-	-	-	-	2	9	45	76	142	185	216	181
2001	1,760	-	-	-	-	-	1	15	38	104	140	184	224	216
2000	1,689	-	-	-	-	-	1	16	38	90	128	184	197	204
1999	1,551	-	-	-	-	1	1	13	34	74	123	160	202	164
Male														
2002	12	-	-	-	-	-	-	1	-	-	-	1	4	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant melanoma of skin (C43)										
Origin not stated[2]										
Both sexes										
2002	3	-	1	3	2	1	-	-	-	1
2001	-	-	1	3	1	1	-	-	-	-
2000	-	2	1	1	2	1	1	-	-	-
1999	1	4	1	4	3	-	2	-	-	-
Male										
2002	3	-	1	1	2	-	-	-	-	-
2001	-	-	1	3	-	-	-	-	-	-
2000	-	2	-	1	1	-	1	-	-	-
1999	1	2	1	2	1	-	2	-	-	-
Female										
2002	-	-	-	2	-	1	-	-	-	1
2001	-	-	-	-	1	1	-	-	-	-
2000	-	-	1	-	1	1	-	-	-	-
1999	-	2	-	2	2	-	-	-	-	-
Malignant neoplasm of breast (C50)										
All origins										
Both sexes										
2002	3,914	3,954	4,551	4,978	4,770	3,528	1,869	658	126	-
2001	3,789	3,972	4,743	5,130	4,533	3,330	1,848	644	89	-
2000	3,783	4,135	4,889	5,256	4,494	3,596	1,880	658	115	-
1999	3,717	4,147	4,962	5,140	4,511	3,428	1,837	611	124	-
Male										
2002	38	35	49	57	55	31	13	3	-	-
2001	40	54	66	69	46	31	9	6	-	-
2000	41	47	68	65	44	36	13	1	1	-
1999	34	47	56	56	42	33	13	7	1	-
Female										
2002	3,876	3,919	4,502	4,921	4,715	3,497	1,856	655	126	-
2001	3,749	3,918	4,677	5,061	4,487	3,299	1,839	638	89	-
2000	3,742	4,088	4,821	5,191	4,450	3,560	1,867	657	114	-
1999	3,683	4,100	4,906	5,084	4,469	3,395	1,824	604	123	-
Hispanic										
Both sexes										
2002	167	173	150	138	124	79	35	11	3	-
2001	191	158	160	137	82	55	36	17	2	-
2000	155	183	154	135	96	64	34	7	3	-
1999	157	146	144	115	94	61	45	14	3	-
Male										
2002	-	1	2	-	2	-	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of breast (C50)														
Hispanic														
Male														
2001	14	-	-	-	-	-	-	-	-	1	1	-	1	-
2000	19	-	-	-	-	-	-	-	-	1	1	-	2	3
1999	13	-	-	-	-	-	-	-	-	-	-	1	2	3
Female														
2002	1,724	-	-	-	-	-	2	8	45	76	142	184	212	181
2001	1,746	-	-	-	-	-	1	15	38	103	139	184	223	216
2000	1,670	-	-	-	-	-	1	16	38	89	127	184	195	201
1999	1,538	-	-	-	-	1	1	13	34	74	123	159	200	161
Non-Hispanic[1]														
Both sexes														
2002	40,037	-	-	-	-	1	7	54	314	825	1,670	2,554	3,479	3,729
2001	39,957	-	-	-	-	1	7	73	336	855	1,721	2,597	3,573	3,621
2000	40,506	-	-	-	-	1	9	67	346	891	1,724	2,541	3,448	3,580
1999	39,881	-	1	-	1	2	9	72	318	911	1,626	2,453	3,347	3,507
Male														
2002	355	-	-	-	-	-	-	1	1	4	8	14	24	28
2001	399	-	-	-	-	-	-	-	2	4	8	13	32	32
2000	409	-	-	-	-	-	-	2	1	11	6	18	26	41
1999	370	-	-	-	-	-	-	-	3	6	9	10	28	33
Female														
2002	39,682	-	-	-	-	1	7	53	313	821	1,662	2,540	3,455	3,701
2001	39,558	-	-	-	-	1	7	73	334	851	1,713	2,584	3,541	3,589
2000	40,097	-	-	-	-	1	9	65	345	880	1,718	2,523	3,422	3,539
1999	39,511	-	1	-	1	2	9	72	315	905	1,617	2,443	3,319	3,474
Non-Hispanic White														
Both sexes														
2002	33,611	-	-	-	-	-	4	31	199	558	1,175	1,876	2,683	3,021
2001	33,629	-	-	-	-	-	3	43	212	596	1,232	1,947	2,790	2,922
2000	34,386	-	-	-	-	1	3	45	230	628	1,249	1,887	2,733	2,895
1999	33,790	-	-	-	1	2	4	46	211	627	1,175	1,820	2,650	2,838
Male														
2002	288	-	-	-	-	-	-	1	-	2	7	10	17	23
2001	329	-	-	-	-	-	-	-	1	3	3	10	23	27
2000	341	-	-	-	-	-	-	-	1	7	2	13	22	31
1999	299	-	-	-	-	-	-	-	2	-	8	8	20	26
Female														
2002	33,323	-	-	-	-	-	4	30	199	556	1,168	1,866	2,666	2,998
2001	33,300	-	-	-	-	-	3	43	211	593	1,229	1,937	2,767	2,895
2000	34,045	-	-	-	-	1	3	45	229	621	1,247	1,874	2,711	2,864

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of breast (C50)										
Hispanic										
Male										
2001	1	4	1	3	2	-	-	-	-	-
2000	1	2	4	2	2	-	1	-	-	-
1999	-	2	3	1	1	-	-	-	-	-
Female										
2002	167	172	148	138	122	79	34	11	3	-
2001	190	154	159	134	80	55	36	17	2	-
2000	154	181	150	133	94	64	33	7	3	-
1999	157	144	141	114	93	61	45	14	3	-
Non-Hispanic[1]										
Both sexes										
2002	3,741	3,770	4,391	4,830	4,634	3,442	1,829	645	122	-
2001	3,587	3,805	4,574	4,976	4,442	3,267	1,808	627	87	-
2000	3,612	3,941	4,729	5,105	4,389	3,521	1,842	648	112	-
1999	3,549	3,991	4,812	5,011	4,408	3,355	1,790	597	121	-
Male										
2002	38	34	47	57	53	31	12	3	-	-
2001	38	49	65	66	44	31	9	6	-	-
2000	40	45	64	63	42	36	12	1	1	-
1999	34	45	52	55	41	33	13	7	1	-
Female										
2002	3,703	3,736	4,344	4,773	4,581	3,411	1,817	642	122	-
2001	3,549	3,756	4,509	4,910	4,398	3,236	1,799	621	87	-
2000	3,572	3,896	4,665	5,042	4,347	3,485	1,830	647	111	-
1999	3,515	3,946	4,760	4,956	4,367	3,322	1,777	590	120	-
Non-Hispanic White										
Both sexes										
2002	3,124	3,188	3,813	4,320	4,136	3,135	1,665	578	105	-
2001	2,973	3,201	3,984	4,458	4,014	2,959	1,660	568	67	-
2000	3,016	3,394	4,187	4,586	4,005	3,194	1,658	576	99	-
1999	2,941	3,396	4,227	4,474	4,014	3,084	1,634	541	105	-
Male										
2002	28	23	42	47	47	29	9	3	-	-
2001	32	41	54	56	40	26	9	4	-	-
2000	32	41	56	58	36	29	11	1	1	-
1999	25	38	41	48	36	30	11	5	1	-
Female										
2002	3,096	3,165	3,771	4,273	4,089	3,106	1,656	575	105	-
2001	2,941	3,160	3,930	4,402	3,974	2,933	1,651	564	67	-
2000	2,984	3,353	4,131	4,528	3,969	3,165	1,647	575	98	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of breast (C50)														
Non-Hispanic White														
Female														
1999	33,491	-	-	-	1	2	4	46	209	627	1,167	1,812	2,630	2,812
Non-Hispanic Black														
Both sexes														
2002	5,579	-	-	-	-	1	2	22	101	232	438	578	669	596
2001	5,528	-	-	-	-	1	4	30	110	219	422	552	657	598
2000	5,384	-	-	-	-	-	5	20	100	227	414	556	611	589
1999	5,386	-	1	-	-	-	5	24	100	259	402	545	605	577
Male														
2002	62	-	-	-	-	-	-	-	1	2	1	4	6	5
2001	64	-	-	-	-	-	-	-	1	-	5	3	9	4
2000	63	-	-	-	-	-	-	2	-	4	4	4	4	8
1999	67	-	-	-	-	-	-	-	1	6	1	1	8	7
Female														
2002	5,517	-	-	-	-	1	2	22	100	230	437	574	663	591
2001	5,464	-	-	-	-	1	4	30	109	219	417	549	648	594
2000	5,321	-	-	-	-	-	5	18	100	223	410	552	607	581
1999	5,319	-	1	-	-	-	5	24	99	253	401	544	597	570
Origin not stated[2]														
Both sexes														
2002	110	-	-	-	-	-	-	-	3	3	6	8	12	14
2001	92	-	-	-	-	-	-	-	1	1	2	9	6	6
2000	105	-	-	-	-	-	-	-	-	1	7	5	10	6
1999	96	-	-	-	-	-	-	1	1	1	6	6	6	11
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	1	-	1
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	108	-	-	-	-	-	-	-	3	3	6	7	12	13
2001	90	-	-	-	-	-	-	-	1	1	2	9	6	6
2000	105	-	-	-	-	-	-	-	-	1	7	5	10	6
1999	95	-	-	-	-	-	-	1	1	1	6	6	6	11
Malignant neoplasm of cervix uteri (C53)														
All origins														
Both sexes														
2002	3,952	-	-	-	-	1	16	47	161	275	391	439	455	401
2001	4,092	-	-	-	1	-	12	51	158	291	420	475	478	371

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of breast (C50)										
Non-Hispanic White										
Female										
1999	2,916	3,358	4,186	4,426	3,978	3,054	1,623	536	104	-
Non-Hispanic Black										
Both sexes										
2002	526	508	497	455	445	285	146	63	15	-
2001	527	532	521	473	389	274	141	59	19	-
2000	514	473	488	472	353	305	173	71	13	-
1999	518	522	515	490	362	251	140	54	16	-
Male										
2002	9	11	5	8	5	2	3	-	-	-
2001	6	7	10	9	4	4	-	2	-	-
2000	8	3	8	5	5	7	1	-	-	-
1999	9	6	10	6	5	3	2	2	-	-
Female										
2002	517	497	492	447	440	283	143	63	15	-
2001	521	525	511	464	385	270	141	57	19	-
2000	506	470	480	467	348	298	172	71	13	-
1999	509	516	505	484	357	248	138	52	16	-
Origin not stated[2]										
Both sexes										
2002	6	11	10	10	12	7	5	2	1	-
2001	11	9	9	17	9	8	4	-	-	-
2000	16	11	6	16	9	11	4	3	-	-
1999	11	10	6	14	9	12	2	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	6	11	10	10	12	7	5	2	1	-
2001	10	8	9	17	9	8	4	-	-	-
2000	16	11	6	16	9	11	4	3	-	-
1999	11	10	5	14	9	12	2	-	-	-
Malignant neoplasm of cervix uteri (C53)										
All origins										
Both sexes										
2002	324	315	315	305	223	155	99	24	6	-
2001	352	279	306	298	285	193	90	30	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of cervix uteri (C53)														
All origins														
Both sexes														
2000	4,200	-	-	-	1	3	14	77	153	294	400	448	467	416
1999	4,205	-	-	-	-	2	10	62	155	292	407	462	452	393
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	3,952	-	-	-	-	1	16	47	161	275	391	439	455	401
2001	4,092	-	-	-	1	-	12	51	158	291	420	475	478	371
2000	4,200	-	-	-	1	3	14	77	153	294	400	448	467	416
1999	4,205	-	-	-	-	2	10	62	155	292	407	462	452	393
Hispanic														
Both sexes														
2002	388	-	-	-	-	-	2	8	22	42	44	42	45	40
2001	412	-	-	-	-	-	3	10	26	42	60	60	42	41
2000	434	-	-	-	-	-	4	13	20	54	54	47	40	40
1999	378	-	-	-	-	-	2	7	24	34	53	55	44	34
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	388	-	-	-	-	-	2	8	22	42	44	42	45	40
2001	412	-	-	-	-	-	3	10	26	42	60	60	42	41
2000	434	-	-	-	-	-	4	13	20	54	54	47	40	40
1999	378	-	-	-	-	-	2	7	24	34	53	55	44	34
Non-Hispanic[1]														
Both sexes														
2002	3,551	-	-	-	-	1	14	39	138	231	346	396	409	359
2001	3,669	-	-	-	1	-	9	40	132	247	359	412	434	329
2000	3,755	-	-	-	1	3	10	64	133	239	344	399	426	376
1999	3,817	-	-	-	-	2	8	55	131	258	353	407	406	358
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)										
All origins										
Both sexes										
2000	332	342	344	320	279	180	92	35	3	-
1999	376	350	346	345	223	199	98	26	6	1
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	324	315	315	305	223	155	99	24	6	-
2001	352	279	306	298	285	193	90	30	2	-
2000	332	342	344	320	279	180	92	35	3	-
1999	376	350	346	345	223	199	98	26	6	1
Hispanic										
Both sexes										
2002	31	24	31	24	16	8	6	2	1	-
2001	34	16	23	21	18	6	8	1	1	-
2000	38	30	30	24	19	12	8	1	-	-
1999	30	27	24	20	11	8	4	1	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	31	24	31	24	16	8	6	2	1	-
2001	34	16	23	21	18	6	8	1	1	-
2000	38	30	30	24	19	12	8	1	-	-
1999	30	27	24	20	11	8	4	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	291	291	282	281	206	147	93	22	5	-
2001	317	263	283	277	267	187	82	29	1	-
2000	292	312	313	294	260	168	84	34	3	-
1999	344	322	322	323	212	191	93	25	6	1
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of cervix uteri (C53)														
Non-Hispanic[1]														
Female														
2002	3,551	-	-	-	-	1	14	39	138	231	346	396	409	359
2001	3,669	-	-	-	1	-	9	40	132	247	359	412	434	329
2000	3,755	-	-	-	1	3	10	64	133	239	344	399	426	376
1999	3,817	-	-	-	-	2	8	55	131	258	353	407	406	358
Non-Hispanic White														
Both sexes														
2002	2,580	-	-	-	-	-	10	26	103	168	249	281	300	277
2001	2,725	-	-	-	-	-	8	29	96	182	266	311	321	241
2000	2,725	-	-	-	-	3	6	50	91	170	233	288	312	275
1999	2,854	-	-	-	-	-	7	47	100	188	263	316	314	263
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	2,580	-	-	-	-	-	10	26	103	168	249	281	300	277
2001	2,725	-	-	-	-	-	8	29	96	182	266	311	321	241
2000	2,725	-	-	-	-	3	6	50	91	170	233	288	312	275
1999	2,854	-	-	-	-	-	7	47	100	188	263	316	314	263
Non-Hispanic Black														
Both sexes														
2002	820	-	-	-	-	1	2	13	30	59	85	95	93	62
2001	778	-	-	-	1	-	1	11	31	54	83	78	87	65
2000	860	-	-	-	1	-	4	12	39	60	87	86	91	82
1999	825	-	-	-	-	2	1	7	27	60	80	68	72	80
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	820	-	-	-	-	1	2	13	30	59	85	95	93	62
2001	778	-	-	-	1	-	1	11	31	54	83	78	87	65
2000	860	-	-	-	1	-	4	12	39	60	87	86	91	82
1999	825	-	-	-	-	2	1	7	27	60	80	68	72	80
Origin not stated[2]														
Both sexes														
2002	13	-	-	-	-	-	-	-	1	2	1	1	1	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)										
Non-Hispanic[1]										
Female										
2002	291	291	282	281	206	147	93	22	5	-
2001	317	263	283	277	267	187	82	29	1	-
2000	292	312	313	294	260	168	84	34	3	-
1999	344	322	322	323	212	191	93	25	6	1
Non-Hispanic White										
Both sexes										
2002	203	203	196	216	154	109	68	13	4	-
2001	237	180	212	207	202	144	63	25	1	-
2000	204	213	225	228	206	135	60	24	2	-
1999	245	235	219	244	170	153	72	13	4	1
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	203	203	196	216	154	109	68	13	4	-
2001	237	180	212	207	202	144	63	25	1	-
2000	204	213	225	228	206	135	60	24	2	-
1999	245	235	219	244	170	153	72	13	4	1
Non-Hispanic Black										
Both sexes										
2002	69	77	75	54	47	29	19	9	1	-
2001	66	72	57	55	56	39	18	4	-	-
2000	70	85	73	60	48	30	21	10	1	-
1999	87	77	96	68	33	35	21	9	2	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	69	77	75	54	47	29	19	9	1	-
2001	66	72	57	55	56	39	18	4	-	-
2000	70	85	73	60	48	30	21	10	1	-
1999	87	77	96	68	33	35	21	9	2	-
Origin not stated[2]										
Both sexes										
2002	2	-	2	-	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of cervix uteri (C53)														
Origin not stated[2]														
Both sexes														
2001	11	-	-	-	-	-	-	1	-	2	1	3	2	1
2000	11	-	-	-	-	-	-	-	-	1	2	2	1	-
1999	10	-	-	-	-	-	-	-	-	-	1	-	2	1
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	13	-	-	-	-	-	-	-	1	2	1	1	1	2
2001	11	-	-	-	-	-	-	1	-	2	1	3	2	1
2000	11	-	-	-	-	-	-	-	-	1	2	2	1	-
1999	10	-	-	-	-	-	-	-	-	-	1	-	2	1
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)														
All origins														
Both sexes														
2002	6,853	-	-	-	-	-	-	2	21	46	100	189	376	531
2001	6,784	-	-	-	-	-	-	10	28	47	92	192	374	457
2000	6,586	-	-	-	-	-	3	2	25	41	97	190	317	478
1999	6,468	-	-	-	-	-	2	11	12	38	104	182	286	427
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	6,853	-	-	-	-	-	-	2	21	46	100	189	376	531
2001	6,784	-	-	-	-	-	-	10	28	47	92	192	374	457
2000	6,586	-	-	-	-	-	3	2	25	41	97	190	317	478
1999	6,468	-	-	-	-	-	2	11	12	38	104	182	286	427
Hispanic														
Both sexes														
2002	337	-	-	-	-	-	-	-	5	7	13	18	21	38
2001	319	-	-	-	-	-	-	-	7	3	10	16	32	32
2000	277	-	-	-	-	-	-	-	2	4	15	11	24	31
1999	272	-	-	-	-	-	1	2	1	4	11	10	23	28
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of cervix uteri (C53)										
Origin not stated[2]										
Both sexes										
2001	1	-	-	-	-	-	-	-	-	-
2000	2	-	1	2	-	-	-	-	-	-
1999	2	1	-	2	-	-	1	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	2	-	2	-	1	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	2	-	1	2	-	-	-	-	-	-
1999	2	1	-	2	-	-	1	-	-	-
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
All origins										
Both sexes										
2002	668	809	1,026	1,051	906	694	314	109	11	-
2001	648	785	999	1,075	988	662	312	104	10	1
2000	644	800	981	1,039	919	617	333	84	15	1
1999	592	835	972	1,061	947	593	290	100	16	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	668	809	1,026	1,051	906	694	314	109	11	-
2001	648	785	999	1,075	988	662	312	104	10	1
2000	644	800	981	1,039	919	617	333	84	15	1
1999	592	835	972	1,061	947	593	290	100	16	-
Hispanic										
Both sexes										
2002	26	54	52	40	26	26	7	4	-	-
2001	39	39	46	34	31	20	6	4	-	-
2000	31	35	40	35	19	20	7	2	1	-
1999	33	45	41	30	21	16	4	2	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)														
Hispanic														
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	337	-	-	-	-	-	-	-	5	7	13	18	21	38
2001	319	-	-	-	-	-	-	-	7	3	10	16	32	32
2000	277	-	-	-	-	-	-	-	2	4	15	11	24	31
1999	272	-	-	-	-	-	1	2	1	4	11	10	23	28
Non-Hispanic[1]														
Both sexes														
2002	6,487	-	-	-	-	-	-	2	15	39	87	171	353	488
2001	6,449	-	-	-	-	-	-	10	21	44	82	176	341	423
2000	6,294	-	-	-	-	-	3	2	23	37	82	178	291	446
1999	6,188	-	-	-	-	-	1	9	11	34	93	172	263	398
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	6,487	-	-	-	-	-	-	2	15	39	87	171	353	488
2001	6,449	-	-	-	-	-	-	10	21	44	82	176	341	423
2000	6,294	-	-	-	-	-	3	2	23	37	82	178	291	446
1999	6,188	-	-	-	-	-	1	9	11	34	93	172	263	398
Non-Hispanic White														
Both sexes														
2002	5,213	-	-	-	-	-	-	1	10	27	70	130	273	370
2001	5,283	-	-	-	-	-	-	3	16	30	64	134	264	349
2000	5,113	-	-	-	-	-	1	2	16	29	59	147	236	363
1999	5,114	-	-	-	-	-	1	6	9	28	68	140	213	313
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	5,213	-	-	-	-	-	-	1	10	27	70	130	273	370
2001	5,283	-	-	-	-	-	-	3	16	30	64	134	264	349

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
Hispanic										
Male										
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	26	54	52	40	26	26	7	4	-	-
2001	39	39	46	34	31	20	6	4	-	-
2000	31	35	40	35	19	20	7	2	1	-
1999	33	45	41	30	21	16	4	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	636	753	972	1,007	877	665	306	105	11	-
2001	606	746	951	1,039	957	637	305	100	10	1
2000	611	765	937	1,004	897	595	326	82	14	1
1999	558	790	929	1,029	925	577	286	97	16	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	636	753	972	1,007	877	665	306	105	11	-
2001	606	746	951	1,039	957	637	305	100	10	1
2000	611	765	937	1,004	897	595	326	82	14	1
1999	558	790	929	1,029	925	577	286	97	16	-
Non-Hispanic White										
Both sexes										
2002	491	540	772	823	757	580	273	87	9	-
2001	453	557	762	878	834	576	268	88	7	-
2000	464	567	731	831	763	535	287	72	10	-
1999	440	587	764	874	813	507	249	88	14	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	491	540	772	823	757	580	273	87	9	-
2001	453	557	762	878	834	576	268	88	7	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)														
Non-Hispanic White														
Female														
2000	5,113	-	-	-	-	-	1	2	16	29	59	147	236	363
1999	5,114	-	-	-	-	-	1	6	9	28	68	140	213	313
Non-Hispanic Black														
Both sexes														
2002	1,107	-	-	-	-	-	-	1	4	8	13	27	61	98
2001	1,037	-	-	-	-	-	-	6	3	8	14	34	59	63
2000	1,055	-	-	-	-	-	1	-	6	6	15	24	44	70
1999	980	-	-	-	-	-	-	3	2	5	22	18	39	76
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	1,107	-	-	-	-	-	-	1	4	8	13	27	61	98
2001	1,037	-	-	-	-	-	-	6	3	8	14	34	59	63
2000	1,055	-	-	-	-	-	1	-	6	6	15	24	44	70
1999	980	-	-	-	-	-	-	3	2	5	22	18	39	76
Origin not stated[2]														
Both sexes														
2002	29	-	-	-	-	-	-	-	1	-	-	-	2	5
2001	16	-	-	-	-	-	-	-	-	-	-	-	1	2
2000	15	-	-	-	-	-	-	-	-	-	-	1	2	1
1999	8	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	29	-	-	-	-	-	-	-	1	-	-	-	2	5
2001	16	-	-	-	-	-	-	-	-	-	-	-	1	2
2000	15	-	-	-	-	-	-	-	-	-	-	1	2	1
1999	8	-	-	-	-	-	-	-	-	-	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)										
Non-Hispanic White										
Female										
2000	464	567	731	831	763	535	287	72	10	-
1999	440	587	764	874	813	507	249	88	14	-
Non-Hispanic Black										
Both sexes										
2002	123	187	186	160	108	81	31	17	2	-
2001	127	173	179	150	115	55	36	11	3	1
2000	131	177	194	156	124	56	36	10	4	1
1999	113	187	159	142	103	65	35	9	2	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	123	187	186	160	108	81	31	17	2	-
2001	127	173	179	150	115	55	36	11	3	1
2000	131	177	194	156	124	56	36	10	4	1
1999	113	187	159	142	103	65	35	9	2	-
Origin not stated[2]										
Both sexes										
2002	6	2	2	4	3	3	1	-	-	-
2001	3	-	2	2	-	5	1	-	-	-
2000	2	-	4	-	3	2	-	-	-	-
1999	1	-	2	2	1	-	-	1	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	6	2	2	4	3	3	1	-	-	-
2001	3	-	2	2	-	5	1	-	-	-
2000	2	-	4	-	3	2	-	-	-	-
1999	1	-	2	2	1	-	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of ovary (C56)														
All origins														
Both sexes														
2002	14,682	1	1	1	1	12	25	33	64	150	319	688	1,029	1,249
2001	14,414	-	-	-	2	5	16	37	65	152	320	610	1,016	1,171
2000	14,060	-	-	2	-	7	16	36	65	145	327	617	966	1,123
1999	13,627	-	1	-	2	9	18	26	51	162	334	616	933	1,081
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	14,682	1	1	1	1	12	25	33	64	150	319	688	1,029	1,249
2001	14,414	-	-	-	2	5	16	37	65	152	320	610	1,016	1,171
2000	14,060	-	-	2	-	7	16	36	65	145	327	617	966	1,123
1999	13,627	-	1	-	2	9	18	26	51	162	334	616	933	1,081
Hispanic														
Both sexes														
2002	622	-	-	1	1	2	1	1	9	10	37	45	71	47
2001	603	-	-	-	1	-	-	9	4	13	30	39	52	57
2000	578	-	-	1	-	-	4	3	6	11	19	42	52	55
1999	478	-	1	-	-	1	1	5	1	11	18	29	51	41
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	622	-	-	1	1	2	1	1	9	10	37	45	71	47
2001	603	-	-	-	1	-	-	9	4	13	30	39	52	57
2000	578	-	-	1	-	-	4	3	6	11	19	42	52	55
1999	478	-	1	-	-	1	1	5	1	11	18	29	51	41
Non-Hispanic[1]														
Both sexes														
2002	14,031	1	1	-	-	10	24	32	55	140	281	640	956	1,199
2001	13,783	-	-	-	1	5	16	28	61	139	289	569	964	1,112
2000	13,454	-	-	1	-	7	12	33	59	133	308	571	913	1,067
1999	13,103	-	-	-	2	8	17	21	50	150	316	585	881	1,035
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of ovary (C56)										
All origins										
Both sexes										
2002	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-
Hispanic										
Both sexes										
2002	64	71	74	90	55	31	12	-	-	-
2001	74	89	75	58	53	36	12	1	-	-
2000	56	75	79	80	46	33	15	1	-	-
1999	59	51	75	59	41	24	9	1	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	64	71	74	90	55	31	12	-	-	-
2001	74	89	75	58	53	36	12	1	-	-
2000	56	75	79	80	46	33	15	1	-	-
1999	59	51	75	59	41	24	9	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,434	1,583	1,862	2,155	1,952	1,142	435	115	14	-
2001	1,363	1,589	1,906	2,182	1,870	1,142	438	94	15	-
2000	1,262	1,590	1,850	2,212	1,752	1,142	436	97	9	-
1999	1,281	1,570	1,904	2,073	1,683	1,027	403	90	7	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of ovary (C56)														
Non-Hispanic[1]														
Male														
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	14,031	1	1	-	-	10	24	32	55	140	281	640	956	1,199
2001	13,783	-	-	-	1	5	16	28	61	139	289	569	964	1,112
2000	13,454	-	-	1	-	7	12	33	59	133	308	571	913	1,067
1999	13,103	-	-	-	2	8	17	21	50	150	316	585	881	1,035
Non-Hispanic White														
Both sexes														
2002	12,560	1	1	-	-	7	20	28	39	112	231	535	834	1,053
2001	12,338	-	-	-	-	3	11	23	45	107	240	496	839	972
2000	12,138	-	-	-	-	5	8	29	46	109	258	498	786	952
1999	11,795	-	-	-	2	7	13	15	36	124	270	506	762	920
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	12,560	1	1	-	-	7	20	28	39	112	231	535	834	1,053
2001	12,338	-	-	-	-	3	11	23	45	107	240	496	839	972
2000	12,138	-	-	-	-	5	8	29	46	109	258	498	786	952
1999	11,795	-	-	-	2	7	13	15	36	124	270	506	762	920
Non-Hispanic Black														
Both sexes														
2002	1,155	-	-	-	-	3	4	3	10	24	35	72	86	113
2001	1,140	-	-	-	1	1	3	5	10	25	40	47	87	112
2000	1,065	-	-	1	-	-	2	4	8	16	43	50	88	86
1999	1,065	-	-	-	-	1	3	6	7	16	31	54	89	94
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	1,155	-	-	-	-	3	4	3	10	24	35	72	86	113
2001	1,140	-	-	-	1	1	3	5	10	25	40	47	87	112
2000	1,065	-	-	1	-	-	2	4	8	16	43	50	88	86
1999	1,065	-	-	-	-	1	3	6	7	16	31	54	89	94

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of ovary (C56)										
Non-Hispanic[1]										
Male										
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	1,434	1,583	1,862	2,155	1,952	1,142	435	115	14	-
2001	1,363	1,589	1,906	2,182	1,870	1,142	438	94	15	-
2000	1,262	1,590	1,850	2,212	1,752	1,142	436	97	9	-
1999	1,281	1,570	1,904	2,073	1,683	1,027	403	90	7	-
Non-Hispanic White										
Both sexes										
2002	1,276	1,407	1,675	1,965	1,801	1,055	404	105	11	-
2001	1,179	1,396	1,729	2,001	1,740	1,050	413	84	10	-
2000	1,120	1,413	1,674	2,047	1,633	1,066	401	85	8	-
1999	1,141	1,400	1,719	1,890	1,566	962	372	83	7	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	1,276	1,407	1,675	1,965	1,801	1,055	404	105	11	-
2001	1,179	1,396	1,729	2,001	1,740	1,050	413	84	10	-
2000	1,120	1,413	1,674	2,047	1,633	1,066	401	85	8	-
1999	1,141	1,400	1,719	1,890	1,566	962	372	83	7	-
Non-Hispanic Black										
Both sexes										
2002	124	139	148	153	124	78	27	9	3	-
2001	139	150	151	149	103	79	24	9	5	-
2000	119	149	146	147	97	65	31	12	1	-
1999	117	135	166	159	97	54	30	6	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	124	139	148	153	124	78	27	9	3	-
2001	139	150	151	149	103	79	24	9	5	-
2000	119	149	146	147	97	65	31	12	1	-
1999	117	135	166	159	97	54	30	6	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of ovary (C56)														
Origin not stated[2]														
Both sexes														
2002	29	-	-	-	-	-	-	-	-	-	1	3	2	3
2001	28	-	-	-	-	-	-	-	-	-	1	2	-	2
2000	28	-	-	-	-	-	-	-	-	1	-	4	1	1
1999	46	-	-	-	-	-	-	-	-	1	-	2	1	5
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	29	-	-	-	-	-	-	-	-	-	1	3	2	3
2001	28	-	-	-	-	-	-	-	-	-	1	2	-	2
2000	28	-	-	-	-	-	-	-	-	1	-	4	1	1
1999	46	-	-	-	-	-	-	-	-	1	-	2	1	5
Malignant neoplasm of prostate (C61)														
All origins														
Both sexes														
2002	30,446	-	-	-	-	-	1	-	1	7	25	80	302	696
2001	30,719	-	-	-	-	-	1	1	-	3	27	99	313	647
2000	31,078	-	-	-	-	-	-	1	-	6	19	108	263	623
1999	31,729	1	-	-	-	-	1	1	1	5	28	100	274	636
Male														
2002	30,446	-	-	-	-	-	1	-	1	7	25	80	302	696
2001	30,719	-	-	-	-	-	1	1	-	3	27	99	313	647
2000	31,078	-	-	-	-	-	-	1	-	6	19	108	263	623
1999	31,729	1	-	-	-	-	1	1	1	5	28	100	274	636
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic														
Both sexes														
2002	1,209	-	-	-	-	-	-	-	1	-	1	7	22	36
2001	1,226	-	-	-	-	-	-	-	-	-	-	5	22	40
2000	1,033	-	-	-	-	-	-	-	-	-	1	7	19	21
1999	1,050	-	-	-	-	-	1	1	-	-	2	2	20	40
Male														
2002	1,209	-	-	-	-	-	-	-	1	-	1	7	22	36

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of ovary (C56)										
Origin not stated[2]										
Both sexes										
2002	1	8	3	4	3	1	-	-	-	-
2001	4	4	3	6	1	3	1	1	-	-
2000	3	4	3	3	5	1	2	-	-	-
1999	5	5	7	11	8	1	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	1	8	3	4	3	1	-	-	-	-
2001	4	4	3	6	1	3	1	1	-	-
2000	3	4	3	3	5	1	2	-	-	-
1999	5	5	7	11	8	1	-	-	-	-
Malignant neoplasm of prostate (C61)										
All origins										
Both sexes										
2002	1,324	2,316	3,984	5,915	6,838	5,474	2,708	703	72	-
2001	1,260	2,379	4,164	6,119	6,857	5,500	2,575	689	85	-
2000	1,306	2,466	4,342	6,314	6,757	5,532	2,638	628	75	-
1999	1,310	2,711	4,605	6,532	6,903	5,418	2,467	644	91	1
Male										
2002	1,324	2,316	3,984	5,915	6,838	5,474	2,708	703	72	-
2001	1,260	2,379	4,164	6,119	6,857	5,500	2,575	689	85	-
2000	1,306	2,466	4,342	6,314	6,757	5,532	2,638	628	75	-
1999	1,310	2,711	4,605	6,532	6,903	5,418	2,467	644	91	1
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic										
Both sexes										
2002	70	131	161	235	237	171	89	47	1	-
2001	68	142	190	278	209	155	78	37	2	-
2000	57	95	182	208	193	150	75	25	-	-
1999	52	135	160	209	177	142	82	23	4	-
Male										
2002	70	131	161	235	237	171	89	47	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of prostate (C61)														
Hispanic														
Male														
2001	1,226	-	-	-	-	-	-	-	-	-	-	5	22	40
2000	1,033	-	-	-	-	-	-	-	-	-	1	7	19	21
1999	1,050	-	-	-	-	-	1	1	-	-	2	2	20	40
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic[1]														
Both sexes														
2002	29,151	-	-	-	-	-	1	-	-	7	24	72	279	658
2001	29,407	-	-	-	-	-	1	1	-	3	26	94	291	603
2000	29,964	-	-	-	-	-	-	1	-	6	18	101	243	601
1999	30,614	1	-	-	-	-	-	-	1	5	26	97	254	595
Male														
2002	29,151	-	-	-	-	-	1	-	-	7	24	72	279	658
2001	29,407	-	-	-	-	-	1	1	-	3	26	94	291	603
2000	29,964	-	-	-	-	-	-	1	-	6	18	101	243	601
1999	30,614	1	-	-	-	-	-	-	1	5	26	97	254	595
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White														
Both sexes														
2002	23,690	-	-	-	-	-	-	-	-	6	18	40	186	472
2001	23,811	-	-	-	-	-	1	-	-	3	15	65	193	434
2000	24,287	-	-	-	-	-	-	1	-	4	11	63	174	422
1999	24,936	1	-	-	-	-	-	-	1	4	19	65	179	430
Male														
2002	23,690	-	-	-	-	-	-	-	-	6	18	40	186	472
2001	23,811	-	-	-	-	-	1	-	-	3	15	65	193	434
2000	24,287	-	-	-	-	-	-	1	-	4	11	63	174	422
1999	24,936	1	-	-	-	-	-	-	1	4	19	65	179	430
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of prostate (C61)										
Hispanic										
Male										
2001	68	142	190	278	209	155	78	37	2	-
2000	57	95	182	208	193	150	75	25	-	-
1999	52	135	160	209	177	142	82	23	4	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic[1]										
Both sexes										
2002	1,248	2,177	3,806	5,667	6,584	5,290	2,612	655	71	-
2001	1,187	2,231	3,966	5,820	6,629	5,335	2,486	651	83	-
2000	1,245	2,368	4,144	6,085	6,549	5,370	2,558	600	75	-
1999	1,253	2,573	4,431	6,308	6,717	5,263	2,383	619	87	1
Male										
2002	1,248	2,177	3,806	5,667	6,584	5,290	2,612	655	71	-
2001	1,187	2,231	3,966	5,820	6,629	5,335	2,486	651	83	-
2000	1,245	2,368	4,144	6,085	6,549	5,370	2,558	600	75	-
1999	1,253	2,573	4,431	6,308	6,717	5,263	2,383	619	87	1
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White										
Both sexes										
2002	889	1,614	2,947	4,568	5,523	4,600	2,234	541	52	-
2001	851	1,679	3,076	4,713	5,517	4,569	2,111	519	65	-
2000	903	1,755	3,191	4,958	5,441	4,630	2,181	497	56	-
1999	880	1,930	3,483	5,134	5,716	4,488	2,038	499	68	1
Male										
2002	889	1,614	2,947	4,568	5,523	4,600	2,234	541	52	-
2001	851	1,679	3,076	4,713	5,517	4,569	2,111	519	65	-
2000	903	1,755	3,191	4,958	5,441	4,630	2,181	497	56	-
1999	880	1,930	3,483	5,134	5,716	4,488	2,038	499	68	1
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of prostate (C61)														
Non-Hispanic White														
Female														
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black														
Both sexes														
2002	5,085	-	-	-	-	-	1	-	-	1	6	31	87	169
2001	5,215	-	-	-	-	-	-	1	-	-	11	28	96	163
2000	5,292	-	-	-	-	-	-	-	-	2	7	35	65	174
1999	5,315	-	-	-	-	-	-	-	-	1	6	31	74	157
Male														
2002	5,085	-	-	-	-	-	1	-	-	1	6	31	87	169
2001	5,215	-	-	-	-	-	-	1	-	-	11	28	96	163
2000	5,292	-	-	-	-	-	-	-	-	2	7	35	65	174
1999	5,315	-	-	-	-	-	-	-	-	1	6	31	74	157
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Origin not stated[2]														
Both sexes														
2002	86	-	-	-	-	-	-	-	-	-	-	1	1	2
2001	86	-	-	-	-	-	-	-	-	-	1	-	-	4
2000	81	-	-	-	-	-	-	-	-	-	-	-	1	1
1999	65	-	-	-	-	-	-	-	-	-	-	1	-	1
Male														
2002	86	-	-	-	-	-	-	-	-	-	-	1	1	2
2001	86	-	-	-	-	-	-	-	-	-	1	-	-	4
2000	81	-	-	-	-	-	-	-	-	-	-	-	1	1
1999	65	-	-	-	-	-	-	-	-	-	-	1	-	1
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Malignant neoplasms of kidney and renal pelvis (C64-C65)														
All origins														
Both sexes														
2002	12,165	4	8	23	7	7	13	14	34	91	215	514	808	1,075

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of prostate (C61)										
Non-Hispanic White										
Female										
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black										
Both sexes										
2002	331	539	813	1,016	989	637	344	103	18	-
2001	318	526	846	1,027	1,027	699	338	118	17	-
2000	320	584	897	1,050	1,027	672	345	96	18	-
1999	355	619	905	1,092	935	709	310	102	19	-
Male										
2002	331	539	813	1,016	989	637	344	103	18	-
2001	318	526	846	1,027	1,027	699	338	118	17	-
2000	320	584	897	1,050	1,027	672	345	96	18	-
1999	355	619	905	1,092	935	709	310	102	19	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Origin not stated[2]										
Both sexes										
2002	6	8	17	13	17	13	7	1	-	-
2001	5	6	8	21	19	10	11	1	-	-
2000	4	3	16	21	15	12	5	3	-	-
1999	5	3	14	15	9	13	2	2	-	-
Male										
2002	6	8	17	13	17	13	7	1	-	-
2001	5	6	8	21	19	10	11	1	-	-
2000	4	3	16	21	15	12	5	3	-	-
1999	5	3	14	15	9	13	2	2	-	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
All origins										
Both sexes										
2002	1,265	1,533	1,706	1,829	1,526	967	429	90	7	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)														
All origins														
Both sexes														
2001	12,078	1	22	30	5	8	12	19	33	74	284	498	813	1,006
2000	11,736	2	15	21	8	8	12	14	28	83	251	450	789	1,026
1999	11,116	2	19	23	10	8	9	25	29	105	237	427	720	959
Male														
2002	7,587	1	-	14	5	2	5	8	23	63	154	355	592	771
2001	7,524	1	10	7	2	7	9	11	16	58	182	355	586	722
2000	7,249	1	8	12	3	4	6	7	19	53	175	309	571	727
1999	6,867	1	11	12	4	4	7	15	22	70	166	310	506	671
Female														
2002	4,578	3	8	9	2	5	8	6	11	28	61	159	216	304
2001	4,554	-	12	23	3	1	3	8	17	16	102	143	227	284
2000	4,487	1	7	9	5	4	6	7	9	30	76	141	218	299
1999	4,249	1	8	11	6	4	2	10	7	35	71	117	214	288
Hispanic														
Both sexes														
2002	659	2	3	1	1	-	1	2	4	9	28	43	60	75
2001	640	-	6	5	1	-	-	4	4	7	31	49	63	48
2000	567	-	3	2	1	-	3	3	4	7	25	42	46	60
1999	547	-	1	4	-	1	-	1	3	9	22	32	52	61
Male														
2002	426	-	-	1	1	-	-	1	3	5	21	27	45	54
2001	426	-	1	3	1	-	-	3	2	7	27	37	43	33
2000	372	-	1	2	1	-	2	1	3	6	21	30	30	46
1999	353	-	1	2	-	-	-	-	2	7	16	28	35	38
Female														
2002	233	2	3	-	-	-	1	1	1	4	7	16	15	21
2001	214	-	5	2	-	-	-	1	2	-	4	12	20	15
2000	195	-	2	-	-	-	1	2	1	1	4	12	16	14
1999	194	-	-	2	-	1	-	1	1	2	6	4	17	23
Non-Hispanic[1]														
Both sexes														
2002	11,477	2	5	22	6	7	12	12	30	82	186	471	745	996
2001	11,415	1	16	25	4	8	12	15	29	66	253	448	748	958
2000	11,124	2	12	19	7	8	9	11	24	76	226	408	739	961
1999	10,542	2	18	19	10	7	9	23	26	94	215	393	667	897
Male														
2002	7,141	1	-	13	4	2	5	7	20	58	132	328	545	713
2001	7,084	1	9	4	1	7	9	8	14	50	155	317	542	689
2000	6,847	1	7	10	2	4	4	6	16	47	154	279	539	678

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
All origins										
Both sexes										
2001	1,322	1,395	1,826	1,810	1,496	947	390	82	5	-
2000	1,169	1,432	1,714	1,819	1,463	945	397	83	7	-
1999	1,152	1,374	1,702	1,696	1,314	867	341	91	6	-
Male										
2002	887	1,020	1,067	1,123	801	457	201	34	4	-
2001	912	928	1,165	1,052	814	500	164	23	-	-
2000	803	942	1,068	1,054	811	477	165	31	3	-
1999	803	894	1,064	1,016	701	421	140	29	-	-
Female										
2002	378	513	639	706	725	510	228	56	3	-
2001	410	467	661	758	682	447	226	59	5	-
2000	366	490	646	765	652	468	232	52	4	-
1999	349	480	638	680	613	446	201	62	6	-
Hispanic										
Both sexes										
2002	85	78	74	88	58	30	14	3	-	-
2001	79	77	86	86	43	37	9	4	1	-
2000	75	65	67	70	55	28	8	3	-	-
1999	68	77	67	65	43	28	9	3	1	-
Male										
2002	52	50	47	64	31	16	7	1	-	-
2001	53	50	57	52	27	23	6	1	-	-
2000	46	40	45	47	31	16	3	1	-	-
1999	53	49	47	38	19	13	4	1	-	-
Female										
2002	33	28	27	24	27	14	7	2	-	-
2001	26	27	29	34	16	14	3	3	1	-
2000	29	25	22	23	24	12	5	2	-	-
1999	15	28	20	27	24	15	5	2	1	-
Non-Hispanic[1]										
Both sexes										
2002	1,177	1,451	1,628	1,735	1,465	937	414	87	7	-
2001	1,240	1,314	1,737	1,722	1,450	910	379	76	4	-
2000	1,092	1,357	1,637	1,744	1,406	911	388	80	7	-
1999	1,082	1,295	1,632	1,624	1,268	838	330	88	5	-
Male										
2002	833	967	1,017	1,057	768	441	193	33	4	-
2001	857	875	1,106	999	785	477	157	22	-	-
2000	755	895	1,016	1,003	779	457	162	30	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)														
Non-Hispanic[1]														
Male														
1999	6,501	1	10	10	4	4	7	15	20	62	150	281	470	633
Female														
2002	4,336	1	5	9	2	5	7	5	10	24	54	143	200	283
2001	4,331	-	7	21	3	1	3	7	15	16	98	131	206	269
2000	4,277	1	5	9	5	4	5	5	8	29	72	129	200	283
1999	4,041	1	8	9	6	3	2	8	6	32	65	112	197	264
Non-Hispanic White														
Both sexes														
2002	10,179	2	5	18	5	5	10	8	16	57	154	385	626	872
2001	10,073	-	11	16	1	2	6	6	17	43	192	375	623	833
2000	9,884	-	7	10	6	6	5	5	19	52	187	339	638	840
1999	9,319	2	11	15	6	4	5	13	15	65	173	313	564	782
Male														
2002	6,363	1	-	11	3	1	4	4	11	40	110	277	464	624
2001	6,263	-	6	1	-	1	4	4	7	33	124	275	451	608
2000	6,082	-	4	4	2	3	3	1	12	36	129	236	465	590
1999	5,754	1	7	8	1	4	3	7	11	43	122	226	392	550
Female														
2002	3,816	1	5	7	2	4	6	4	5	17	44	108	162	248
2001	3,810	-	5	15	1	1	2	2	10	10	68	100	172	225
2000	3,802	-	3	6	4	3	2	4	7	16	58	103	173	250
1999	3,565	1	4	7	5	-	2	6	4	22	51	87	172	232
Non-Hispanic Black														
Both sexes														
2002	1,062	-	-	4	1	2	1	3	14	21	24	72	91	104
2001	1,105	-	3	9	3	5	5	9	10	21	51	63	100	104
2000	1,033	1	5	7	1	2	4	6	4	21	30	58	84	99
1999	1,028	-	4	3	3	3	4	10	10	27	34	67	87	99
Male														
2002	626	-	-	2	1	1	1	2	9	15	16	41	62	79
2001	673	-	1	3	1	5	5	4	6	15	25	39	72	67
2000	625	-	3	4	-	1	1	5	3	9	18	35	59	68
1999	618	-	2	1	2	-	4	8	8	19	23	44	65	72
Female														
2002	436	-	-	2	-	1	-	1	5	6	8	31	29	25
2001	432	-	2	6	2	-	-	5	4	6	26	24	28	37
2000	408	1	2	3	1	1	3	1	1	12	12	23	25	31
1999	410	-	2	2	1	3	-	2	2	8	11	23	22	27

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
Non-Hispanic[1]										
Male										
1999	748	843	1,015	975	682	408	135	28	-	-
Female										
2002	344	484	611	678	697	496	221	54	3	-
2001	383	439	631	723	665	433	222	54	4	-
2000	337	462	621	741	627	454	226	50	4	-
1999	334	452	617	649	586	430	195	60	5	-
Non-Hispanic White										
Both sexes										
2002	1,026	1,287	1,467	1,561	1,342	860	388	78	7	-
2001	1,071	1,178	1,563	1,554	1,326	836	345	71	4	-
2000	954	1,198	1,467	1,582	1,289	848	353	73	6	-
1999	958	1,149	1,464	1,470	1,166	773	293	74	4	-
Male										
2002	731	865	920	967	701	408	187	30	4	-
2001	732	788	1,002	895	721	444	146	21	-	-
2000	668	782	915	914	714	425	149	28	2	-
1999	670	748	914	887	641	381	119	19	-	-
Female										
2002	295	422	547	594	641	452	201	48	3	-
2001	339	390	561	659	605	392	199	50	4	-
2000	286	416	552	668	575	423	204	45	4	-
1999	288	401	550	583	525	392	174	55	4	-
Non-Hispanic Black										
Both sexes										
2002	122	135	125	149	101	65	21	7	-	-
2001	136	108	141	137	104	63	28	5	-	-
2000	114	134	137	132	101	56	29	7	1	-
1999	103	122	141	129	84	55	31	11	1	-
Male										
2002	85	80	71	75	50	28	6	2	-	-
2001	101	67	86	83	53	30	9	1	-	-
2000	74	96	83	68	56	29	10	2	1	-
1999	65	77	82	72	32	21	13	8	-	-
Female										
2002	37	55	54	74	51	37	15	5	-	-
2001	35	41	55	54	51	33	19	4	-	-
2000	40	38	54	64	45	27	19	5	-	-
1999	38	45	59	57	52	34	18	3	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)														
Origin not stated[2]														
Both sexes														
2002	29	-	-	-	-	-	-	-	-	-	1	-	3	4
2001	23	-	-	-	-	-	-	-	-	1	-	1	2	-
2000	45	-	-	-	-	-	-	-	-	-	-	-	4	5
1999	27	-	-	-	-	-	-	1	-	2	-	2	1	1
Male														
2002	20	-	-	-	-	-	-	-	-	-	1	-	2	4
2001	14	-	-	-	-	-	-	-	-	1	-	1	1	-
2000	30	-	-	-	-	-	-	-	-	-	-	-	2	3
1999	13	-	-	-	-	-	-	-	-	1	-	1	1	-
Female														
2002	9	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	9	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	15	-	-	-	-	-	-	-	-	-	-	-	2	2
1999	14	-	-	-	-	-	-	1	-	1	-	1	-	1
Malignant neoplasm of bladder (C67)														
All origins														
Both sexes														
2002	12,628	-	-	2	1	-	1	3	11	31	92	182	321	572
2001	12,225	-	-	1	-	1	2	6	9	35	93	161	314	484
2000	12,002	-	-	-	1	2	1	1	9	40	71	160	280	486
1999	11,911	-	-	-	1	-	1	6	10	32	68	140	323	492
Male														
2002	8,504	-	-	1	-	-	-	2	7	21	57	129	223	425
2001	8,343	-	-	-	-	1	2	1	7	14	56	118	233	359
2000	8,163	-	-	-	-	2	1	-	2	21	41	110	212	369
1999	8,105	-	-	-	1	-	-	3	6	20	43	107	243	374
Female														
2002	4,124	-	-	1	1	-	1	1	4	10	35	53	98	147
2001	3,882	-	-	1	-	-	-	5	2	21	37	43	81	125
2000	3,839	-	-	-	1	-	-	1	7	19	30	50	68	117
1999	3,806	-	-	-	-	-	1	3	4	12	25	33	80	118
Hispanic														
Both sexes														
2002	387	-	-	1	-	-	-	-	-	-	5	11	13	27
2001	336	-	-	-	-	1	1	-	-	2	2	12	9	16
2000	307	-	-	-	-	-	-	-	1	3	5	8	9	25
1999	301	-	-	-	-	-	-	-	2	1	3	5	17	21

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of kidney and renal pelvis (C64-C65)										
Origin not stated[2]										
Both sexes										
2002	3	4	4	6	3	-	1	-	-	-
2001	3	4	3	2	3	-	2	2	-	-
2000	2	10	10	5	2	6	1	-	-	-
1999	2	2	3	7	3	1	2	-	-	-
Male										
2002	2	3	3	2	2	-	1	-	-	-
2001	2	3	2	1	2	-	1	-	-	-
2000	2	7	7	4	1	4	-	-	-	-
1999	2	2	2	3	-	-	1	-	-	-
Female										
2002	1	1	1	4	1	-	-	-	-	-
2001	1	1	1	1	1	-	1	2	-	-
2000	-	3	3	1	1	2	1	-	-	-
1999	-	-	1	4	3	1	1	-	-	-
Malignant neoplasm of bladder (C67)										
All origins										
Both sexes										
2002	738	1,151	1,756	2,366	2,341	1,864	908	255	32	1
2001	781	1,147	1,667	2,179	2,309	1,865	864	265	42	-
2000	737	1,119	1,736	2,155	2,252	1,790	875	256	31	-
1999	752	1,126	1,797	2,178	2,139	1,714	852	249	30	1
Male										
2002	559	845	1,244	1,659	1,540	1,163	498	118	12	1
2001	597	859	1,229	1,554	1,562	1,129	482	123	17	-
2000	547	838	1,256	1,542	1,512	1,110	478	110	12	-
1999	559	849	1,283	1,575	1,433	1,039	438	116	15	1
Female										
2002	179	306	512	707	801	701	410	137	20	-
2001	184	288	438	625	747	736	382	142	25	-
2000	190	281	480	613	740	680	397	146	19	-
1999	193	277	514	603	706	675	414	133	15	-
Hispanic										
Both sexes										
2002	30	40	51	53	57	55	39	4	1	-
2001	25	40	50	51	53	41	20	11	2	-
2000	25	21	50	46	43	39	21	8	3	-
1999	30	35	53	41	38	37	14	2	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of bladder (C67)														
Hispanic														
Male														
2002	252	-	-	-	-	-	-	-	-	-	2	7	9	15
2001	223	-	-	-	-	1	1	-	-	-	1	7	4	15
2000	216	-	-	-	-	-	-	-	-	3	3	5	7	21
1999	222	-	-	-	-	-	-	-	2	1	3	4	12	18
Female														
2002	135	-	-	1	-	-	-	-	-	-	3	4	4	12
2001	113	-	-	-	-	-	-	-	-	2	1	5	5	1
2000	91	-	-	-	-	-	-	-	1	-	2	3	2	4
1999	79	-	-	-	-	-	-	-	-	-	-	1	5	3
Non-Hispanic[1]														
Both sexes														
2002	12,211	-	-	1	1	-	1	3	11	31	87	170	306	543
2001	11,858	-	-	1	-	-	1	6	9	33	91	147	305	467
2000	11,665	-	-	-	1	2	1	1	8	36	66	152	271	458
1999	11,582	-	-	-	1	-	1	6	8	31	65	135	304	470
Male														
2002	8,235	-	-	1	-	-	-	2	7	21	55	122	213	409
2001	8,100	-	-	-	-	-	1	1	7	14	55	110	229	344
2000	7,925	-	-	-	-	2	1	-	2	18	38	105	205	346
1999	7,865	-	-	-	1	-	-	3	4	19	40	103	229	355
Female														
2002	3,976	-	-	-	1	-	1	1	4	10	32	48	93	134
2001	3,758	-	-	1	-	-	-	5	2	19	36	37	76	123
2000	3,740	-	-	-	1	-	-	1	6	18	28	47	66	112
1999	3,717	-	-	-	-	-	1	3	4	12	25	32	75	115
Non-Hispanic White														
Both sexes														
2002	11,109	-	-	1	1	-	-	2	8	29	71	151	265	474
2001	10,823	-	-	1	-	-	1	6	8	28	72	121	265	409
2000	10,712	-	-	-	1	1	1	1	7	33	58	129	245	412
1999	10,548	-	-	-	1	-	-	3	5	23	54	109	266	407
Male														
2002	7,649	-	-	1	-	-	-	1	5	19	47	109	193	375
2001	7,541	-	-	-	-	-	1	1	7	13	43	95	201	308
2000	7,390	-	-	-	-	1	1	-	2	18	34	91	188	312
1999	7,290	-	-	-	1	-	-	1	2	13	33	83	198	321
Female														
2002	3,460	-	-	-	1	-	-	1	3	10	24	42	72	99
2001	3,282	-	-	1	-	-	-	5	1	15	29	26	64	101

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of bladder (C67)										
Hispanic										
Male										
2002	24	31	36	34	34	33	24	3	-	-
2001	22	27	32	31	36	21	15	8	2	-
2000	16	16	34	39	30	21	15	4	2	-
1999	25	30	39	28	24	26	9	1	-	-
Female										
2002	6	9	15	19	23	22	15	1	1	-
2001	3	13	18	20	17	20	5	3	-	-
2000	9	5	16	7	13	18	6	4	1	-
1999	5	5	14	13	14	11	5	1	2	-
Non-Hispanic[1]										
Both sexes										
2002	705	1,108	1,701	2,312	2,279	1,803	867	251	31	-
2001	753	1,103	1,615	2,120	2,252	1,819	843	253	40	-
2000	711	1,097	1,683	2,103	2,201	1,747	852	247	28	-
1999	720	1,088	1,744	2,129	2,099	1,671	834	247	28	1
Male										
2002	532	812	1,207	1,625	1,503	1,127	472	115	12	-
2001	573	829	1,196	1,518	1,523	1,104	466	115	15	-
2000	530	821	1,219	1,499	1,475	1,087	462	105	10	-
1999	533	816	1,244	1,542	1,408	1,010	427	115	15	1
Female										
2002	173	296	494	687	776	676	395	136	19	-
2001	180	274	419	602	729	715	377	138	25	-
2000	181	276	464	604	726	660	390	142	18	-
1999	187	272	500	587	691	661	407	132	13	-
Non-Hispanic White										
Both sexes										
2002	638	989	1,542	2,126	2,106	1,663	790	225	28	-
2001	667	995	1,484	1,956	2,070	1,697	772	235	36	-
2000	644	998	1,535	1,926	2,031	1,624	809	233	24	-
1999	635	969	1,587	1,964	1,957	1,550	770	219	28	1
Male										
2002	491	742	1,122	1,511	1,410	1,061	441	109	12	-
2001	522	768	1,116	1,428	1,428	1,044	440	113	13	-
2000	491	762	1,135	1,405	1,379	1,015	445	101	10	-
1999	479	746	1,161	1,445	1,326	957	402	106	15	1
Female										
2002	147	247	420	615	696	602	349	116	16	-
2001	145	227	368	528	642	653	332	122	23	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasm of bladder (C67)														
Non-Hispanic White														
Female														
2000	3,322	-	-	-	1	-	-	1	5	15	24	38	57	100
1999	3,258	-	-	-	-	-	-	2	3	10	21	26	68	86
Non-Hispanic Black														
Both sexes														
2002	931	-	-	-	-	-	-	1	3	1	12	16	33	59
2001	870	-	-	-	-	-	-	-	1	4	19	20	35	48
2000	842	-	-	-	-	1	-	-	-	3	7	20	24	40
1999	906	-	-	-	-	-	1	2	2	8	9	23	37	59
Male														
2002	475	-	-	-	-	-	-	1	2	1	6	10	16	29
2001	444	-	-	-	-	-	-	-	-	1	12	10	26	27
2000	460	-	-	-	-	1	-	-	-	-	4	11	16	28
1999	487	-	-	-	-	-	-	1	1	6	6	17	30	32
Female														
2002	456	-	-	-	-	-	-	-	1	-	6	6	17	30
2001	426	-	-	-	-	-	-	-	1	3	7	10	9	21
2000	382	-	-	-	-	-	-	-	-	3	3	9	8	12
1999	419	-	-	-	-	-	1	1	1	2	3	6	7	27
Origin not stated[2]														
Both sexes														
2002	30	-	-	-	-	-	-	-	-	-	-	1	2	2
2001	31	-	-	-	-	-	-	-	-	-	-	2	-	1
2000	30	-	-	-	-	-	-	-	-	1	-	-	-	3
1999	28	-	-	-	-	-	-	-	-	-	-	-	2	1
Male														
2002	17	-	-	-	-	-	-	-	-	-	-	-	1	1
2001	20	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	22	-	-	-	-	-	-	-	-	-	-	-	-	2
1999	18	-	-	-	-	-	-	-	-	-	-	-	2	1
Female														
2002	13	-	-	-	-	-	-	-	-	-	-	1	1	1
2001	11	-	-	-	-	-	-	-	-	-	-	1	-	1
2000	8	-	-	-	-	-	-	-	-	1	-	-	-	1
1999	10	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasm of bladder (C67)										
Non-Hispanic White										
Female										
2000	153	236	400	521	652	609	364	132	14	-
1999	156	223	426	519	631	593	368	113	13	-
Non-Hispanic Black										
Both sexes										
2002	57	103	139	154	141	120	68	21	3	-
2001	77	93	104	140	151	97	61	17	3	-
2000	59	86	132	159	155	102	39	12	3	-
1999	71	105	139	140	122	109	54	25	-	-
Male										
2002	34	57	71	90	74	55	26	3	-	-
2001	45	50	61	72	72	44	20	2	2	-
2000	35	48	71	86	85	58	14	3	-	-
1999	44	59	69	81	68	48	18	7	-	-
Female										
2002	23	46	68	64	67	65	42	18	3	-
2001	32	43	43	68	79	53	41	15	1	-
2000	24	38	61	73	70	44	25	9	3	-
1999	27	46	70	59	54	61	36	18	-	-
Origin not stated[2]										
Both sexes										
2002	3	3	4	1	5	6	2	-	-	1
2001	3	4	2	8	4	5	1	1	-	-
2000	1	1	3	6	8	4	2	1	-	-
1999	2	3	-	8	2	6	4	-	-	-
Male										
2002	3	2	1	-	3	3	2	-	-	1
2001	2	3	1	5	3	4	1	-	-	-
2000	1	1	3	4	7	2	1	1	-	-
1999	1	3	-	5	1	3	2	-	-	-
Female										
2002	-	1	3	1	2	3	-	-	-	-
2001	1	1	1	3	1	1	-	1	-	-
2000	-	-	-	2	1	2	1	-	-	-
1999	1	-	-	3	1	3	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)														
All origins														
Both sexes														
2002	12,830	12	114	202	135	108	102	129	259	375	634	900	1,158	1,276
2001	12,609	16	100	170	130	104	120	134	282	399	629	830	1,121	1,174
2000	12,655	20	104	166	150	105	120	159	247	454	642	815	1,109	1,110
1999	12,765	10	113	184	124	115	103	177	267	443	622	842	1,038	1,206
Male														
2002	7,125	8	63	112	79	65	71	67	156	222	396	560	665	773
2001	7,015	8	52	81	64	63	66	88	174	245	393	500	673	734
2000	7,011	10	58	89	87	63	69	94	170	293	411	481	667	660
1999	7,001	8	56	96	58	69	62	104	172	268	370	507	613	670
Female														
2002	5,705	4	51	90	56	43	31	62	103	153	238	340	493	503
2001	5,594	8	48	89	66	41	54	46	108	154	236	330	448	440
2000	5,644	10	46	77	63	42	51	65	77	161	231	334	442	450
1999	5,764	2	57	88	66	46	41	73	95	175	252	335	425	536
Hispanic														
Both sexes														
2002	649	2	21	29	23	15	9	14	19	21	34	50	73	53
2001	617	4	20	22	15	18	10	16	22	39	38	42	45	47
2000	631	4	16	36	27	17	11	20	25	30	46	41	51	36
1999	568	4	27	30	18	2	16	20	11	31	34	37	51	47
Male														
2002	359	2	12	14	13	11	5	6	10	9	21	30	50	27
2001	335	1	10	12	4	13	6	11	10	21	20	29	21	31
2000	346	1	10	24	17	9	7	11	16	18	22	23	18	20
1999	293	4	11	13	9	2	8	12	5	20	25	18	24	24
Female														
2002	290	-	9	15	10	4	4	8	9	12	13	20	23	26
2001	282	3	10	10	11	5	4	5	12	18	18	13	24	16
2000	285	3	6	12	10	8	4	9	9	12	24	18	33	16
1999	275	-	16	17	9	-	8	8	6	11	9	19	27	23
Non-Hispanic[1]														
Both sexes														
2002	12,152	10	93	172	112	91	93	115	240	352	599	848	1,081	1,219
2001	11,967	12	79	147	114	86	110	117	260	359	591	787	1,075	1,124
2000	11,995	16	87	130	123	88	109	139	222	423	596	773	1,055	1,070
1999	12,173	6	86	154	106	112	87	157	256	412	588	803	986	1,157
Male														
2002	6,752	6	51	98	66	54	66	61	146	212	374	530	612	744

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All origins										
Both sexes										
2002	1,284	1,407	1,547	1,454	981	536	174	39	4	-
2001	1,290	1,354	1,469	1,506	1,060	528	157	31	5	-
2000	1,227	1,409	1,623	1,437	1,041	521	166	28	2	-
1999	1,247	1,498	1,621	1,543	949	496	144	22	1	-
Male										
2002	721	819	859	750	460	213	54	10	2	-
2001	736	763	826	777	511	202	48	9	2	-
2000	719	789	889	705	482	212	53	9	1	-
1999	747	858	870	785	441	192	47	7	1	-
Female										
2002	563	588	688	704	521	323	120	29	2	-
2001	554	591	643	729	549	326	109	22	3	-
2000	508	620	734	732	559	309	113	19	1	-
1999	500	640	751	758	508	304	97	15	-	-
Hispanic										
Both sexes										
2002	62	60	58	54	28	17	6	-	1	-
2001	62	60	57	47	35	14	4	-	-	-
2000	70	58	47	53	19	19	4	1	-	-
1999	52	47	57	38	29	12	3	2	-	-
Male										
2002	35	36	29	27	12	8	1	-	1	-
2001	28	32	40	24	16	6	-	-	-	-
2000	42	32	27	27	11	10	1	-	-	-
1999	36	25	23	19	12	1	1	1	-	-
Female										
2002	27	24	29	27	16	9	5	-	-	-
2001	34	28	17	23	19	8	4	-	-	-
2000	28	26	20	26	8	9	3	1	-	-
1999	16	22	34	19	17	11	2	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,219	1,345	1,486	1,396	953	518	168	39	3	-
2001	1,222	1,292	1,411	1,456	1,023	513	153	31	5	-
2000	1,152	1,348	1,575	1,381	1,017	500	162	27	2	-
1999	1,191	1,449	1,561	1,501	917	482	141	20	1	-
Male										
2002	683	782	830	720	448	205	53	10	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)														
Non-Hispanic[1]														
Male														
2001	6,667	7	41	68	60	50	60	77	164	223	373	471	651	700
2000	6,646	9	47	65	70	54	62	83	154	274	389	457	646	639
1999	6,696	4	45	83	49	67	54	92	167	248	345	488	588	644
Female														
2002	5,400	4	42	74	46	37	27	54	94	140	225	318	469	475
2001	5,300	5	38	79	54	36	50	40	96	136	218	316	424	424
2000	5,349	7	40	65	53	34	47	56	68	149	207	316	409	431
1999	5,477	2	41	71	57	45	33	65	89	164	243	315	398	513
Non-Hispanic White														
Both sexes														
2002	11,139	5	59	131	83	73	77	95	210	318	547	774	982	1,112
2001	11,004	8	63	121	84	70	85	103	232	334	530	711	984	1,023
2000	11,026	12	65	99	95	68	88	120	195	376	520	704	974	980
1999	11,238	3	70	110	85	83	69	145	227	377	525	736	899	1,074
Male														
2002	6,195	3	34	70	46	44	57	50	125	193	347	484	567	673
2001	6,154	5	35	53	43	41	47	70	147	210	337	432	598	642
2000	6,128	8	35	48	53	43	50	71	129	247	344	426	589	591
1999	6,193	2	39	63	41	48	43	85	148	229	304	444	536	599
Female														
2002	4,944	2	25	61	37	29	20	45	85	125	200	290	415	439
2001	4,850	3	28	68	41	29	38	33	85	124	193	279	386	381
2000	4,898	4	30	51	42	25	38	49	66	129	176	278	385	389
1999	5,045	1	31	47	44	35	26	60	79	148	221	292	363	475
Non-Hispanic Black														
Both sexes														
2002	766	3	27	34	20	14	12	11	22	30	38	49	76	82
2001	736	1	10	19	25	12	17	8	20	18	45	64	69	76
2000	762	2	18	26	25	16	19	15	21	32	56	51	68	72
1999	754	3	11	36	18	23	14	10	26	26	50	59	69	70
Male														
2002	400	2	12	23	12	9	5	5	17	16	19	25	31	54
2001	378	-	3	12	13	6	10	4	10	8	25	34	41	44
2000	407	1	10	15	15	9	12	9	19	17	35	23	48	39
1999	399	2	4	17	7	15	9	6	17	15	32	39	42	35
Female														
2002	366	1	15	11	8	5	7	6	5	14	19	24	45	28

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
Non-Hispanic[1]										
Male										
2001	706	731	785	751	494	196	48	9	2	-
2000	674	754	861	677	468	201	52	9	1	-
1999	709	832	846	764	427	191	46	6	1	-
Female										
2002	536	563	656	676	505	313	115	29	2	-
2001	516	561	626	705	529	317	105	22	3	-
2000	478	594	714	704	549	299	110	18	1	-
1999	482	617	715	737	490	291	95	14	-	-
Non-Hispanic White										
Both sexes										
2002	1,131	1,251	1,395	1,310	899	490	158	36	3	-
2001	1,132	1,188	1,321	1,385	971	485	144	26	4	-
2000	1,067	1,261	1,495	1,306	954	466	154	26	1	-
1999	1,097	1,351	1,488	1,423	870	459	128	18	1	-
Male										
2002	633	733	782	677	420	197	50	9	1	-
2001	651	685	742	710	471	184	43	7	1	-
2000	628	713	818	641	445	190	50	9	-	-
1999	653	781	810	725	411	182	44	5	1	-
Female										
2002	498	518	613	633	479	293	108	27	2	-
2001	481	503	579	675	500	301	101	19	3	-
2000	439	548	677	665	509	276	104	17	1	-
1999	444	570	678	698	459	277	84	13	-	-
Non-Hispanic Black										
Both sexes										
2002	62	72	70	69	43	20	9	3	-	-
2001	69	83	70	52	40	24	8	5	1	-
2000	66	67	66	58	51	25	6	1	1	-
1999	72	77	55	66	38	21	9	1	-	-
Male										
2002	38	35	35	32	23	4	2	1	-	-
2001	43	36	34	25	15	8	4	2	1	-
2000	37	29	37	25	18	6	2	-	1	-
1999	41	42	26	30	10	8	2	-	-	-
Female										
2002	24	37	35	37	20	16	7	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)														
Non-Hispanic Black														
Female														
2001	358	1	7	7	12	6	7	4	10	10	20	30	28	32
2000	355	1	8	11	10	7	7	6	2	15	21	28	20	33
1999	355	1	7	19	11	8	5	4	9	11	18	20	27	35
Origin not stated[2]														
Both sexes														
2002	29	-	-	1	-	2	-	-	-	2	1	2	4	4
2001	25	-	1	1	1	-	-	1	-	1	-	1	1	3
2000	29	-	1	-	-	-	-	-	-	1	-	1	3	4
1999	24	-	-	-	-	1	-	-	-	-	-	2	1	2
Male														
2002	14	-	-	-	-	-	-	-	-	1	1	-	3	2
2001	13	-	1	1	-	-	-	-	-	1	-	-	1	3
2000	19	-	1	-	-	-	-	-	-	1	-	1	3	1
1999	12	-	-	-	-	-	-	-	-	-	-	1	1	2
Female														
2002	15	-	-	1	-	2	-	-	-	1	-	2	1	2
2001	12	-	-	-	1	-	-	1	-	-	-	1	-	-
2000	10	-	-	-	-	-	-	-	-	-	-	-	-	3
1999	12	-	-	-	-	1	-	-	-	-	-	1	-	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
All origins														
Both sexes														
2002	56,225	28	119	166	213	310	393	368	548	695	1,056	1,599	2,361	3,217
2001	56,341	27	157	165	202	311	394	436	552	768	1,089	1,674	2,390	3,224
2000	56,518	39	146	168	204	323	395	412	506	776	1,077	1,675	2,350	3,156
1999	56,318	30	137	142	209	303	417	423	537	837	1,076	1,648	2,302	3,346
Male														
2002	30,302	15	63	87	130	188	229	225	320	409	629	953	1,388	1,947
2001	30,060	12	82	95	122	192	234	246	328	464	669	1,002	1,440	1,900
2000	29,888	21	76	98	125	202	253	226	313	438	637	1,015	1,384	1,850
1999	29,721	12	75	84	122	180	257	241	316	497	647	986	1,382	1,957
Female														
2002	25,923	13	56	79	83	122	164	143	228	286	427	646	973	1,270
2001	26,281	15	75	70	80	119	160	190	224	304	420	672	950	1,324
2000	26,630	18	70	70	79	121	142	186	193	338	440	660	966	1,306

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
Non-Hispanic Black										
Female										
2001	26	47	36	27	25	16	4	3	-	-
2000	29	38	29	33	33	19	4	1	-	-
1999	31	35	29	36	28	13	7	1	-	-
Origin not stated[2]										
Both sexes										
2002	3	2	3	4	-	1	-	-	-	-
2001	6	2	1	3	2	1	-	-	-	-
2000	5	3	1	3	5	2	-	-	-	-
1999	4	2	3	4	3	2	-	-	-	-
Male										
2002	3	1	-	3	-	-	-	-	-	-
2001	2	-	1	2	1	-	-	-	-	-
2000	3	3	1	1	3	1	-	-	-	-
1999	2	1	1	2	2	-	-	-	-	-
Female										
2002	-	1	3	1	-	1	-	-	-	-
2001	4	2	-	1	1	1	-	-	-	-
2000	2	-	-	2	2	1	-	-	-	-
1999	2	1	2	2	1	2	-	-	-	-
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All origins										
Both sexes										
2002	4,283	5,672	7,836	9,319	8,970	6,003	2,433	574	62	-
2001	4,231	5,602	8,057	9,580	8,677	5,848	2,344	536	77	-
2000	4,128	5,932	8,109	9,743	8,705	5,716	2,329	559	70	-
1999	4,208	5,996	8,336	9,596	8,493	5,468	2,233	526	55	-
Male										
2002	2,523	3,391	4,512	5,047	4,506	2,624	906	186	24	-
2001	2,512	3,316	4,543	5,065	4,262	2,537	846	168	25	-
2000	2,355	3,445	4,593	5,168	4,231	2,475	783	185	15	-
1999	2,469	3,436	4,631	5,102	3,998	2,337	816	155	21	-
Female										
2002	1,760	2,281	3,324	4,272	4,464	3,379	1,527	388	38	-
2001	1,719	2,286	3,514	4,515	4,415	3,311	1,498	368	52	-
2000	1,773	2,487	3,516	4,575	4,474	3,241	1,546	374	55	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
All origins														
Female														
1999	26,597	18	62	58	87	123	160	182	221	340	429	662	920	1,389
Hispanic														
Both sexes														
2002	2,829	3	28	46	47	62	83	62	77	105	120	135	147	206
2001	2,747	7	38	46	41	77	67	75	79	90	121	118	161	179
2000	2,643	8	29	30	44	55	74	65	59	85	115	138	142	172
1999	2,538	8	28	28	47	45	83	63	72	88	102	141	158	182
Male														
2002	1,518	1	12	21	26	44	53	43	49	62	70	76	87	116
2001	1,577	2	21	28	20	52	47	45	54	59	83	72	92	108
2000	1,460	5	14	15	32	34	49	44	39	43	72	95	77	106
1999	1,319	3	16	19	23	23	50	34	41	57	51	84	91	102
Female														
2002	1,311	2	16	25	21	18	30	19	28	43	50	59	60	90
2001	1,170	5	17	18	21	25	20	30	25	31	38	46	69	71
2000	1,183	3	15	15	12	21	25	21	20	42	43	43	65	66
1999	1,219	5	12	9	24	22	33	29	31	31	51	57	67	80
Non-Hispanic[1]														
Both sexes														
2002	53,249	24	91	120	166	247	307	306	469	587	931	1,459	2,204	3,005
2001	53,458	20	117	118	161	233	326	357	471	675	965	1,549	2,223	3,040
2000	53,749	31	116	136	159	266	320	347	446	688	957	1,532	2,203	2,973
1999	53,669	22	108	114	162	258	334	360	463	749	970	1,503	2,139	3,158
Male														
2002	28,698	13	51	66	104	144	173	182	270	345	555	873	1,295	1,826
2001	28,410	10	60	67	102	140	186	200	273	404	583	925	1,344	1,790
2000	28,358	16	61	83	92	167	204	182	273	393	563	917	1,302	1,736
1999	28,347	9	58	65	99	157	207	207	274	440	592	900	1,290	1,851
Female														
2002	24,551	11	40	54	62	103	134	124	199	242	376	586	909	1,179
2001	25,048	10	57	51	59	93	140	157	198	271	382	624	879	1,250
2000	25,391	15	55	53	67	99	116	165	173	295	394	615	901	1,237
1999	25,322	13	50	49	63	101	127	153	189	309	378	603	849	1,307
Non-Hispanic White														
Both sexes														
2002	47,058	15	72	94	119	183	247	225	345	441	742	1,138	1,759	2,485
2001	47,190	17	97	81	117	178	244	240	355	519	736	1,241	1,816	2,523

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
All origins										
Female										
1999	1,739	2,560	3,705	4,494	4,495	3,131	1,417	371	34	-
Hispanic										
Both sexes										
2002	231	311	323	337	267	164	58	15	2	-
2001	263	271	326	346	227	148	53	11	3	-
2000	255	274	359	317	206	143	53	19	1	-
1999	214	287	319	273	212	117	60	11	-	-
Male										
2002	129	167	164	169	121	79	22	6	1	-
2001	143	159	179	205	113	65	23	4	3	-
2000	141	156	182	155	103	64	25	9	-	-
1999	124	145	167	137	89	42	18	3	-	-
Female										
2002	102	144	159	168	146	85	36	9	1	-
2001	120	112	147	141	114	83	30	7	-	-
2000	114	118	177	162	103	79	28	10	1	-
1999	90	142	152	136	123	75	42	8	-	-
Non-Hispanic[1]										
Both sexes										
2002	4,043	5,344	7,495	8,959	8,677	5,829	2,372	554	60	-
2001	3,954	5,317	7,710	9,211	8,434	5,691	2,287	525	74	-
2000	3,860	5,643	7,736	9,406	8,487	5,564	2,270	540	69	-
1999	3,985	5,698	8,006	9,299	8,262	5,343	2,167	514	55	-
Male										
2002	2,391	3,213	4,333	4,863	4,373	2,542	884	179	23	-
2001	2,361	3,151	4,354	4,843	4,142	2,467	822	164	22	-
2000	2,205	3,279	4,403	5,005	4,122	2,408	756	176	15	-
1999	2,344	3,284	4,456	4,953	3,900	2,291	797	152	21	-
Female										
2002	1,652	2,131	3,162	4,096	4,304	3,287	1,488	375	37	-
2001	1,593	2,166	3,356	4,368	4,292	3,224	1,465	361	52	-
2000	1,655	2,364	3,333	4,401	4,365	3,156	1,514	364	54	-
1999	1,641	2,414	3,550	4,346	4,362	3,052	1,370	362	34	-
Non-Hispanic White										
Both sexes										
2002	3,445	4,651	6,676	8,160	8,042	5,450	2,202	515	52	-
2001	3,355	4,581	6,911	8,395	7,804	5,296	2,131	484	69	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
Non-Hispanic White														
Both sexes														
2000	47,695	30	92	105	127	192	228	251	334	531	752	1,204	1,813	2,514
1999	47,608	16	91	86	113	188	249	271	347	590	762	1,220	1,760	2,654
Male														
2002	25,523	9	42	50	75	112	142	137	196	256	445	695	1,060	1,541
2001	25,162	8	47	45	66	110	142	137	215	300	448	758	1,115	1,498
2000	25,272	16	50	62	72	120	148	136	202	311	464	733	1,070	1,478
1999	25,219	9	48	51	69	114	163	153	204	351	465	733	1,079	1,569
Female														
2002	21,535	6	30	44	44	71	105	88	149	185	297	443	699	944
2001	22,028	9	50	36	51	68	102	103	140	219	288	483	701	1,025
2000	22,423	14	42	43	55	72	80	115	132	220	288	471	743	1,036
1999	22,389	7	43	35	44	74	86	118	143	239	297	487	681	1,085
Non-Hispanic Black														
Both sexes														
2002	5,080	5	14	17	34	43	44	62	94	104	162	267	357	439
2001	5,162	3	13	27	35	46	61	84	97	133	189	255	329	427
2000	5,026	1	19	21	22	58	72	74	84	128	174	288	327	390
1999	5,038	4	14	21	40	53	80	71	86	134	173	236	318	431
Male														
2002	2,593	2	6	12	20	24	20	35	58	66	96	155	187	240
2001	2,642	2	9	14	29	27	30	47	51	90	109	134	192	239
2000	2,535	-	8	14	12	36	43	33	57	64	85	159	187	222
1999	2,566	-	8	11	27	32	41	43	54	74	106	138	180	242
Female														
2002	2,487	3	8	5	14	19	24	27	36	38	66	112	170	199
2001	2,520	1	4	13	6	19	31	37	46	43	80	121	137	188
2000	2,491	1	11	7	10	22	29	41	27	64	89	129	140	168
1999	2,472	4	6	10	13	21	39	28	32	60	67	98	138	189
Origin not stated[2]														
Both sexes														
2002	147	1	-	-	-	1	3	-	2	3	5	5	10	6
2001	136	-	2	1	-	1	1	4	2	3	3	7	6	5
2000	126	-	1	2	1	2	1	-	1	3	5	5	5	11
1999	111	-	1	-	-	-	-	-	2	-	4	4	5	6
Male														
2002	86	1	-	-	-	-	3	-	1	2	4	4	6	5
2001	73	-	1	-	-	-	1	1	1	1	3	5	4	2
2000	70	-	1	-	1	1	-	-	1	2	2	3	5	8

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Non-Hispanic White										
Both sexes										
2000	3,298	4,968	6,943	8,611	7,917	5,146	2,088	493	58	-
1999	3,388	5,004	7,204	8,487	7,685	4,971	2,004	471	47	-
Male										
2002	2,056	2,833	3,913	4,480	4,077	2,398	818	169	19	-
2001	2,021	2,748	3,945	4,456	3,851	2,312	768	151	21	-
2000	1,901	2,939	3,960	4,623	3,868	2,246	698	161	14	-
1999	2,022	2,919	4,048	4,538	3,664	2,131	733	139	17	-
Female										
2002	1,389	1,818	2,763	3,680	3,965	3,052	1,384	346	33	-
2001	1,334	1,833	2,966	3,939	3,953	2,984	1,363	333	48	-
2000	1,397	2,029	2,983	3,988	4,049	2,900	1,390	332	44	-
1999	1,366	2,085	3,156	3,949	4,021	2,840	1,271	332	30	-
Non-Hispanic Black										
Both sexes										
2002	496	579	689	647	519	322	145	34	7	-
2001	488	618	670	661	522	329	134	37	4	-
2000	478	561	671	639	462	353	158	38	8	-
1999	486	582	665	665	477	317	138	39	8	-
Male										
2002	277	313	366	299	231	120	52	10	4	-
2001	273	343	336	300	238	123	44	11	1	-
2000	259	281	372	311	199	130	49	13	1	-
1999	258	309	336	333	188	122	50	10	4	-
Female										
2002	219	266	323	348	288	202	93	24	3	-
2001	215	275	334	361	284	206	90	26	3	-
2000	219	280	299	328	263	223	109	25	7	-
1999	228	273	329	332	289	195	88	29	4	-
Origin not stated[2]										
Both sexes										
2002	9	17	18	23	26	10	3	5	-	-
2001	14	14	21	23	16	9	4	-	-	-
2000	13	15	14	20	12	9	6	-	-	-
1999	9	11	11	24	19	8	6	1	-	-
Male										
2002	3	11	15	15	12	3	-	1	-	-
2001	8	6	10	17	7	5	1	-	-	-
2000	9	10	8	8	6	3	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)														
Origin not stated[2]														
Male														
1999	55	-	1	-	-	-	-	-	1	-	4	2	1	4
Female														
2002	61	-	-	-	-	1	-	-	1	1	1	1	4	1
2001	63	-	1	1	-	1	-	3	1	2	-	2	2	3
2000	56	-	-	2	-	1	1	-	-	1	3	2	-	3
1999	56	-	-	-	-	-	-	-	1	-	-	2	4	2
Hodgkin's disease (C81)														
All origins														
Both sexes														
2002	1,352	-	-	3	3	27	59	64	88	81	78	83	94	74
2001	1,323	-	-	1	5	25	53	76	91	81	80	91	91	61
2000	1,287	-	-	3	5	21	52	60	85	78	73	73	88	64
1999	1,403	-	1	-	9	31	63	77	71	90	75	82	83	87
Male														
2002	751	-	-	2	3	15	26	37	44	46	46	53	52	53
2001	725	-	-	1	2	15	28	39	51	50	51	54	56	40
2000	686	-	-	1	3	6	29	31	48	42	40	46	53	30
1999	785	-	1	-	3	20	35	42	44	56	47	48	50	54
Female														
2002	601	-	-	1	-	12	33	27	44	35	32	30	42	21
2001	598	-	-	-	3	10	25	37	40	31	29	37	35	21
2000	601	-	-	2	2	15	23	29	37	36	33	27	35	34
1999	618	-	-	-	6	11	28	35	27	34	28	34	33	33
Hispanic														
Both sexes														
2002	99	-	-	3	1	2	7	5	10	6	6	8	7	7
2001	99	-	-	1	-	4	3	7	6	6	7	3	11	8
2000	95	-	-	-	-	3	6	9	5	3	5	6	7	1
1999	113	-	-	-	3	2	6	2	5	5	7	6	11	7
Male														
2002	53	-	-	2	1	2	2	4	3	4	4	4	5	4
2001	66	-	-	1	-	3	1	3	5	3	5	1	10	8
2000	63	-	-	-	-	1	6	3	4	2	4	6	5	-
1999	65	-	-	-	2	2	4	1	2	4	4	3	8	4
Female														
2002	46	-	-	1	-	-	5	1	7	2	2	4	2	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Origin not stated[2]										
Male										
1999	1	7	8	12	9	4	1	-	-	-
Female										
2002	6	6	3	8	14	7	3	4	-	-
2001	6	8	11	6	9	4	3	-	-	-
2000	4	5	6	12	6	6	4	-	-	-
1999	8	4	3	12	10	4	5	1	-	-
Hodgkin's disease (C81)										
All origins										
Both sexes										
2002	94	94	138	126	121	83	35	7	-	-
2001	82	103	108	151	127	66	24	7	-	-
2000	94	92	140	134	121	70	24	10	-	-
1999	91	120	163	149	113	70	22	5	1	-
Male										
2002	53	62	84	60	65	37	11	2	-	-
2001	45	65	64	75	55	26	7	1	-	-
2000	57	58	76	71	50	30	10	5	-	-
1999	61	72	88	69	50	28	16	1	-	-
Female										
2002	41	32	54	66	56	46	24	5	-	-
2001	37	38	44	76	72	40	17	6	-	-
2000	37	34	64	63	71	40	14	5	-	-
1999	30	48	75	80	63	42	6	4	1	-
Hispanic										
Both sexes										
2002	3	6	7	7	11	2	1	-	-	-
2001	6	11	10	6	5	5	-	-	-	-
2000	12	4	7	12	11	2	1	1	-	-
1999	10	14	12	9	9	5	-	-	-	-
Male										
2002	3	2	4	2	6	1	-	-	-	-
2001	3	8	7	3	2	3	-	-	-	-
2000	8	4	6	9	3	-	1	1	-	-
1999	5	7	6	5	6	2	-	-	-	-
Female										
2002	-	4	3	5	5	1	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hodgkin's disease (C81)														
Hispanic														
Female														
2001	33	-	-	-	-	1	2	4	1	3	2	2	1	-
2000	32	-	-	-	-	2	-	6	1	1	1	-	2	1
1999	48	-	-	-	1	-	2	1	3	1	3	3	3	3
Non-Hispanic[1]														
Both sexes														
2002	1,248	-	-	-	2	25	51	59	77	74	72	75	86	67
2001	1,221	-	-	-	5	21	50	69	85	75	72	87	80	53
2000	1,189	-	-	3	5	18	46	51	80	75	68	67	81	62
1999	1,286	-	1	-	6	29	57	75	66	85	68	75	72	79
Male														
2002	693	-	-	-	2	13	23	33	40	41	42	49	46	49
2001	657	-	-	-	2	12	27	36	46	47	45	52	46	32
2000	621	-	-	1	3	5	23	28	44	40	36	40	48	29
1999	719	-	1	-	1	18	31	41	42	52	43	45	42	49
Female														
2002	555	-	-	-	-	12	28	26	37	33	30	26	40	18
2001	564	-	-	-	3	9	23	33	39	28	27	35	34	21
2000	568	-	-	2	2	13	23	23	36	35	32	27	33	33
1999	567	-	-	-	5	11	26	34	24	33	25	30	30	30
Non-Hispanic White														
Both sexes														
2002	1,093	-	-	-	1	19	44	47	65	58	60	64	71	54
2001	1,069	-	-	-	4	17	38	50	63	63	61	73	73	44
2000	1,059	-	-	2	3	14	34	46	71	63	56	60	68	54
1999	1,149	-	1	-	5	24	43	58	58	75	59	66	63	70
Male														
2002	599	-	-	-	1	11	21	24	33	32	33	40	35	39
2001	577	-	-	-	2	10	24	25	37	38	38	44	41	27
2000	551	-	-	1	1	3	20	25	38	34	30	36	40	25
1999	643	-	1	-	1	15	24	31	38	45	39	39	34	44
Female														
2002	494	-	-	-	-	8	23	23	32	26	27	24	36	15
2001	492	-	-	-	2	7	14	25	26	25	23	29	32	17
2000	508	-	-	1	2	11	14	21	33	29	26	24	28	29
1999	506	-	-	-	4	9	19	27	20	30	20	27	29	26
Non-Hispanic Black														
Both sexes														
2002	126	-	-	-	1	4	6	11	12	14	11	10	10	10
2001	136	-	-	-	1	4	11	18	21	11	10	12	5	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)										
Hispanic										
Female										
2001	3	3	3	3	3	2	-	-	-	-
2000	4	-	1	3	8	2	-	-	-	-
1999	5	7	6	4	3	3	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	91	88	131	118	110	81	34	7	-	-
2001	76	92	97	145	122	61	24	7	-	-
2000	81	88	133	122	109	68	23	9	-	-
1999	80	106	151	140	103	65	22	5	1	-
Male										
2002	50	60	80	57	59	36	11	2	-	-
2001	42	57	57	72	53	23	7	1	-	-
2000	49	54	70	62	46	30	9	4	-	-
1999	56	65	82	64	44	26	16	1	-	-
Female										
2002	41	28	51	61	51	45	23	5	-	-
2001	34	35	40	73	69	38	17	6	-	-
2000	32	34	63	60	63	38	14	5	-	-
1999	24	41	69	76	59	39	6	4	1	-
Non-Hispanic White										
Both sexes										
2002	78	79	121	111	104	79	32	6	-	-
2001	66	88	90	132	117	59	24	7	-	-
2000	69	82	118	122	102	66	21	8	-	-
1999	74	92	142	135	100	57	21	5	1	-
Male										
2002	41	55	75	54	57	36	10	2	-	-
2001	36	55	54	67	50	21	7	1	-	-
2000	40	51	61	62	43	29	8	4	-	-
1999	53	57	78	62	43	23	15	1	-	-
Female										
2002	37	24	46	57	47	43	22	4	-	-
2001	30	33	36	65	67	38	17	6	-	-
2000	29	31	57	60	59	37	13	4	-	-
1999	21	35	64	73	57	34	6	4	1	-
Non-Hispanic Black										
Both sexes										
2002	10	6	9	5	4	1	1	1	-	-
2001	9	4	7	9	3	2	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hodgkin's disease (C81)														
Non-Hispanic Black														
Both sexes														
2000	109	-	-	1	1	4	10	5	7	12	12	7	10	8
1999	125	-	-	-	-	5	14	17	8	10	8	9	9	9
Male														
2002	80	-	-	-	1	2	2	8	7	8	9	8	7	8
2001	72	-	-	-	-	2	2	11	9	8	7	7	4	5
2000	57	-	-	-	1	2	3	3	5	6	6	4	5	4
1999	74	-	-	-	-	3	7	10	4	7	4	6	8	5
Female														
2002	46	-	-	-	-	2	4	3	5	6	2	2	3	2
2001	64	-	-	-	1	2	9	7	12	3	3	5	1	4
2000	52	-	-	1	-	2	7	2	2	6	6	3	5	4
1999	51	-	-	-	-	2	7	7	4	3	4	3	1	4
Origin not stated[2]														
Both sexes														
2002	5	-	-	-	-	-	1	-	1	1	-	-	1	-
2001	3	-	-	-	-	-	-	-	-	-	1	1	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	4	-	-	-	-	-	-	-	-	-	-	1	-	1
Male														
2002	5	-	-	-	-	-	1	-	1	1	-	-	1	-
2001	2	-	-	-	-	-	-	-	-	-	1	1	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	1	-	-
Non-Hodgkin's lymphoma (C82-C85)														
All origins														
Both sexes														
2002	21,910	2	15	15	42	44	98	111	170	253	434	638	943	1,319
2001	22,305	1	1	11	27	61	87	129	169	301	455	711	1,021	1,367
2000	22,729	1	7	23	24	60	120	124	169	290	461	730	1,015	1,434
1999	22,802	2	16	12	25	51	93	109	180	324	465	696	1,045	1,414
Male														
2002	11,538	1	6	10	30	31	62	71	107	161	278	404	574	823
2001	11,710	1	1	8	21	43	51	72	107	195	305	450	663	817
2000	11,812	-	5	16	12	46	80	64	108	192	290	466	646	865

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)										
Non-Hispanic Black										
Both sexes										
2000	9	4	11	-	5	1	2	-	-	-
1999	5	11	7	5	1	6	1	-	-	-
Male										
2002	7	3	5	3	1	-	1	-	-	-
2001	5	2	3	3	2	2	-	-	-	-
2000	7	2	5	-	3	-	1	-	-	-
1999	3	8	4	2	1	1	1	-	-	-
Female										
2002	3	3	4	2	3	1	-	1	-	-
2001	4	2	4	6	1	-	-	-	-	-
2000	2	2	6	-	2	1	1	-	-	-
1999	2	3	3	3	-	5	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	1	-	-	-	1	-	-	-	-	-
1999	1	-	-	-	1	-	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	1	-	-	-	-	-
Non-Hodgkin's lymphoma (C82-C85)										
All origins										
Both sexes										
2002	1,730	2,225	3,039	3,722	3,561	2,446	881	200	22	-
2001	1,731	2,193	3,165	3,776	3,593	2,392	904	186	24	-
2000	1,736	2,386	3,212	3,984	3,551	2,330	856	193	23	-
1999	1,740	2,441	3,407	4,009	3,545	2,198	834	178	18	-
Male										
2002	1,015	1,334	1,652	1,898	1,696	1,013	315	51	6	-
2001	1,008	1,286	1,739	1,873	1,685	1,020	305	55	5	-
2000	984	1,342	1,763	2,018	1,616	959	274	62	4	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Non-Hodgkin's lymphoma (C82-C85)														
All origins														
Male														
1999	11,794	-	9	9	21	30	61	69	109	204	306	459	652	851
Female														
2002	10,372	1	9	5	12	13	36	40	63	92	156	234	369	496
2001	10,595	-	-	3	6	18	36	57	62	106	150	261	358	550
2000	10,917	1	2	7	12	14	40	60	61	98	171	264	369	569
1999	11,008	2	7	3	4	21	32	40	71	120	159	237	393	563
Hispanic														
Both sexes														
2002	1,068	-	2	3	4	5	20	19	22	32	46	52	56	76
2001	1,008	-	-	2	4	11	11	20	17	33	44	55	67	83
2000	1,026	-	2	2	-	4	17	19	19	26	57	51	62	78
1999	958	-	1	1	2	5	14	10	27	25	43	64	67	62
Male														
2002	549	-	-	1	2	5	14	14	11	19	26	34	31	46
2001	579	-	-	2	4	7	8	11	13	22	34	35	39	50
2000	560	-	2	1	-	4	13	11	12	16	35	34	38	48
1999	493	-	-	1	1	4	6	5	13	19	24	42	42	38
Female														
2002	519	-	2	2	2	-	6	5	11	13	20	18	25	30
2001	429	-	-	-	-	4	3	9	4	11	10	20	28	33
2000	466	-	-	1	-	-	4	8	7	10	22	17	24	30
1999	465	-	1	-	1	1	8	5	14	6	19	22	25	24
Non-Hispanic[1]														
Both sexes														
2002	20,780	2	13	12	38	38	77	92	148	219	384	583	883	1,240
2001	21,246	1	1	9	23	49	76	108	150	268	410	653	952	1,282
2000	21,657	1	5	21	24	56	103	105	149	264	404	677	949	1,353
1999	21,792	2	15	11	23	46	79	99	153	299	419	630	976	1,350
Male														
2002	10,953	1	6	9	28	26	47	57	96	141	248	368	542	774
2001	11,105	1	1	6	17	36	43	60	93	173	270	413	623	767
2000	11,225	-	3	15	12	42	67	53	95	176	255	432	604	814
1999	11,271	-	9	8	20	26	55	64	96	185	279	416	609	812
Female														
2002	9,827	1	7	3	10	12	30	35	52	78	136	215	341	466
2001	10,141	-	-	3	6	13	33	48	57	95	140	240	329	515
2000	10,432	1	2	6	12	14	36	52	54	88	149	245	345	539
1999	10,521	2	6	3	3	20	24	35	57	114	140	214	367	538

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)										
All origins										
Male										
1999	996	1,355	1,821	2,052	1,544	892	292	53	9	-
Female										
2002	715	891	1,387	1,824	1,865	1,433	566	149	16	-
2001	723	907	1,426	1,903	1,908	1,372	599	131	19	-
2000	752	1,044	1,449	1,966	1,935	1,371	582	131	19	-
1999	744	1,086	1,586	1,957	2,001	1,306	542	125	9	-
Hispanic										
Both sexes										
2002	110	122	146	155	107	65	19	6	1	-
2001	118	91	117	147	93	68	21	5	1	-
2000	112	111	166	128	82	63	18	9	-	-
1999	91	117	144	117	88	51	25	4	-	-
Male										
2002	63	69	67	66	45	26	8	2	-	-
2001	65	55	67	82	49	28	5	2	1	-
2000	64	60	83	60	38	28	9	4	-	-
1999	53	56	69	59	33	20	8	-	-	-
Female										
2002	47	53	79	89	62	39	11	4	1	-
2001	53	36	50	65	44	40	16	3	-	-
2000	48	51	83	68	44	35	9	5	-	-
1999	38	61	75	58	55	31	17	4	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,617	2,095	2,888	3,561	3,438	2,377	861	193	21	-
2001	1,610	2,093	3,041	3,619	3,494	2,322	881	181	23	-
2000	1,617	2,271	3,040	3,848	3,464	2,263	836	184	23	-
1999	1,646	2,318	3,257	3,880	3,447	2,144	807	173	18	-
Male										
2002	951	1,258	1,580	1,828	1,644	987	307	49	6	-
2001	941	1,227	1,670	1,783	1,633	991	300	53	4	-
2000	914	1,279	1,675	1,955	1,578	929	265	58	4	-
1999	942	1,296	1,748	1,986	1,504	870	284	53	9	-
Female										
2002	666	837	1,308	1,733	1,794	1,390	554	144	15	-
2001	669	866	1,371	1,836	1,861	1,331	581	128	19	-
2000	703	992	1,365	1,893	1,886	1,334	571	126	19	-
1999	704	1,022	1,509	1,894	1,943	1,274	523	120	9	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Non-Hodgkin's lymphoma (C82-C85)														
Non-Hispanic White														
Both sexes														
2002	18,901	1	12	12	26	23	57	65	97	164	299	479	719	1,074
2001	19,300	1	1	8	18	33	54	69	113	201	322	528	809	1,105
2000	19,843	-	2	16	22	42	68	70	111	202	329	553	814	1,209
1999	19,956	-	12	7	16	31	60	69	108	228	331	517	855	1,188
Male														
2002	9,961	-	6	9	20	18	37	44	62	104	193	314	449	678
2001	10,052	1	1	5	12	25	29	40	75	127	212	342	531	661
2000	10,264	-	2	11	12	34	45	39	70	136	217	355	515	720
1999	10,259	-	8	6	15	18	43	43	69	144	218	344	535	713
Female														
2002	8,940	1	6	3	6	5	20	21	35	60	106	165	270	396
2001	9,248	-	-	3	6	8	25	29	38	74	110	186	278	444
2000	9,579	-	-	5	10	8	23	31	41	66	112	198	299	489
1999	9,697	-	4	1	1	13	17	26	39	84	113	173	320	475
Non-Hispanic Black														
Both sexes														
2002	1,446	1	1	-	10	10	14	22	41	36	76	89	126	136
2001	1,478	-	-	1	4	16	16	29	33	59	70	103	118	137
2000	1,355	1	2	4	1	8	27	30	25	54	64	104	102	119
1999	1,427	2	2	3	6	13	19	24	34	62	73	92	97	138
Male														
2002	767	1	-	-	7	4	6	11	29	26	50	48	71	76
2001	795	-	-	1	4	11	11	15	17	42	46	56	79	77
2000	719	-	-	3	-	3	16	12	18	35	33	66	67	82
1999	778	-	1	2	4	6	12	17	22	36	50	58	62	81
Female														
2002	679	-	1	-	3	6	8	11	12	10	26	41	55	60
2001	683	-	-	-	-	5	5	14	16	17	24	47	39	60
2000	636	1	2	1	1	5	11	18	7	19	31	38	35	37
1999	649	2	1	1	2	7	7	7	12	26	23	34	35	57
Origin not stated[2]														
Both sexes														
2002	62	-	-	-	-	1	1	-	-	2	4	3	4	3
2001	51	-	-	-	-	1	-	1	2	-	1	3	2	2
2000	46	-	-	-	-	-	-	-	1	-	-	2	4	3
1999	52	-	-	-	-	-	-	-	-	-	3	2	2	2
Male														
2002	36	-	-	-	-	-	1	-	-	1	4	2	1	3
2001	26	-	-	-	-	-	-	1	1	-	1	2	1	-
2000	27	-	-	-	-	-	-	-	1	-	-	-	4	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)										
Non-Hispanic White										
Both sexes										
2002	1,443	1,886	2,637	3,339	3,258	2,278	827	185	20	-
2001	1,430	1,870	2,813	3,400	3,302	2,197	832	173	21	-
2000	1,446	2,065	2,813	3,626	3,312	2,149	797	176	21	-
1999	1,479	2,129	3,016	3,632	3,303	2,037	762	160	16	-
Male										
2002	848	1,143	1,457	1,725	1,562	950	291	45	6	-
2001	836	1,108	1,547	1,678	1,548	936	284	50	4	-
2000	818	1,181	1,544	1,854	1,511	890	251	55	4	-
1999	841	1,190	1,632	1,847	1,455	818	266	47	7	-
Female										
2002	595	743	1,180	1,614	1,696	1,328	536	140	14	-
2001	594	762	1,266	1,722	1,754	1,261	548	123	17	-
2000	628	884	1,269	1,772	1,801	1,259	546	121	17	-
1999	638	939	1,384	1,785	1,848	1,219	496	113	9	-
Non-Hispanic Black										
Both sexes										
2002	141	165	191	153	131	74	24	5	-	-
2001	132	174	174	146	138	83	36	7	2	-
2000	137	150	168	145	99	79	30	6	-	-
1999	121	153	177	179	107	78	34	11	2	-
Male										
2002	83	91	102	68	54	27	9	4	-	-
2001	74	96	94	65	59	35	11	2	-	-
2000	75	74	95	65	38	22	12	3	-	-
1999	72	90	84	99	32	32	11	5	2	-
Female										
2002	58	74	89	85	77	47	15	1	-	-
2001	58	78	80	81	79	48	25	5	2	-
2000	62	76	73	80	61	57	18	3	-	-
1999	49	63	93	80	75	46	23	6	-	-
Origin not stated[2]										
Both sexes										
2002	3	8	5	6	16	4	1	1	-	-
2001	3	9	7	10	6	2	2	-	-	-
2000	7	4	6	8	5	4	2	-	-	-
1999	3	6	6	12	10	3	2	1	-	-
Male										
2002	1	7	5	4	7	-	-	-	-	-
2001	2	4	2	8	3	1	-	-	-	-
2000	6	3	5	3	-	2	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Non-Hodgkin's lymphoma (C82-C85)														
Origin not stated[2]														
Male														
1999	30	-	-	-	-	-	-	-	-	-	3	1	1	1
Female														
2002	26	-	-	-	-	1	-	-	-	1	-	1	3	-
2001	25	-	-	-	-	1	-	-	1	-	-	1	1	2
2000	19	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	22	-	-	-	-	-	-	-	-	-	-	1	1	1
Leukemia (C91-C95)														
All origins														
Both sexes														
2002	21,498	21	104	148	168	237	235	190	281	325	442	606	853	1,111
2001	21,451	25	155	153	170	225	254	230	283	346	433	630	845	1,073
2000	21,339	38	137	142	173	242	222	228	245	368	433	622	789	1,041
1999	21,014	27	119	130	175	221	261	234	266	374	425	626	744	1,112
Male														
2002	12,016	11	57	75	97	142	140	117	163	178	251	335	486	667
2001	11,850	10	81	86	99	134	155	135	165	196	236	346	467	632
2000	11,777	21	69	81	109	150	143	131	152	185	238	356	435	629
1999	11,519	11	65	75	98	130	161	128	148	204	231	346	431	636
Female														
2002	9,482	10	47	73	71	95	95	73	118	147	191	271	367	444
2001	9,601	15	74	67	71	91	99	95	118	150	197	284	378	441
2000	9,562	17	68	61	64	92	79	97	93	183	195	266	354	412
1999	9,495	16	54	55	77	91	100	106	118	170	194	280	313	476
Hispanic														
Both sexes														
2002	1,088	2	26	40	42	54	55	38	42	63	58	51	42	74
2001	1,114	7	38	43	37	62	53	47	54	46	57	49	51	47
2000	1,038	8	27	28	44	48	51	37	34	52	45	58	52	61
1999	994	8	26	27	42	38	63	49	37	52	46	50	50	62
Male														
2002	601	1	12	18	23	37	36	25	32	35	34	22	29	36
2001	640	2	21	25	16	42	38	31	35	31	34	30	26	26
2000	595	5	12	14	32	29	30	30	22	24	28	38	26	41
1999	534	3	16	18	20	17	40	26	23	28	20	30	22	31
Female														
2002	487	1	14	22	19	17	19	13	10	28	24	29	13	38
2001	474	5	17	18	21	20	15	16	19	15	23	19	25	21
2000	443	3	15	14	12	19	21	7	12	28	17	20	26	20

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)										
Origin not stated[2]										
Male										
1999	1	3	4	7	7	2	-	-	-	-
Female										
2002	2	1	-	2	9	4	1	1	-	-
2001	1	5	5	2	3	1	2	-	-	-
2000	1	1	1	5	5	2	2	-	-	-
1999	2	3	2	5	3	1	2	1	-	-
Leukemia (C91-C95)										
All origins										
Both sexes										
2002	1,500	1,998	2,820	3,392	3,371	2,294	1,086	281	35	-
2001	1,462	1,963	2,923	3,541	3,155	2,256	1,008	271	50	-
2000	1,373	2,073	2,937	3,515	3,218	2,194	1,041	271	37	-
1999	1,404	2,078	2,934	3,401	3,059	2,151	984	258	31	-
Male										
2002	891	1,242	1,768	1,995	1,809	1,062	420	95	15	-
2001	920	1,216	1,746	2,023	1,676	1,041	380	89	17	-
2000	823	1,287	1,803	1,982	1,713	1,018	359	85	8	-
1999	853	1,275	1,779	1,946	1,579	964	375	75	9	-
Female										
2002	609	756	1,052	1,397	1,562	1,232	666	186	20	-
2001	542	747	1,177	1,518	1,479	1,215	628	182	33	-
2000	550	786	1,134	1,533	1,505	1,176	682	186	29	-
1999	551	803	1,155	1,455	1,480	1,187	609	183	22	-
Hispanic										
Both sexes										
2002	69	88	95	81	83	53	25	7	-	-
2001	73	90	107	113	72	44	17	5	2	-
2000	72	81	101	95	70	44	21	8	1	-
1999	53	86	93	85	64	36	22	5	-	-
Male										
2002	31	45	52	50	44	26	9	4	-	-
2001	47	46	52	72	30	22	10	2	2	-
2000	41	45	53	48	39	25	10	3	-	-
1999	35	45	62	45	30	17	5	1	-	-
Female										
2002	38	43	43	31	39	27	16	3	-	-
2001	26	44	55	41	42	22	7	3	-	-
2000	31	36	48	47	31	19	11	5	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Leukemia (C91-C95)														
Hispanic														
Female														
1999	460	5	10	9	22	21	23	23	14	24	26	20	28	31
Non-Hispanic[1]														
Both sexes														
2002	20,363	18	78	108	126	183	179	152	238	262	383	554	806	1,036
2001	20,285	18	115	109	133	163	200	180	229	298	375	578	793	1,023
2000	20,248	30	109	112	128	192	170	191	211	313	384	563	736	974
1999	19,981	19	92	103	133	183	198	185	227	322	378	575	691	1,048
Male														
2002	11,386	9	45	57	74	105	103	92	131	143	217	312	453	630
2001	11,180	8	59	61	83	92	116	104	130	164	201	314	440	604
2000	11,155	16	56	67	76	120	113	101	130	159	208	317	408	584
1999	10,968	8	48	57	78	113	121	102	124	176	210	315	409	603
Female														
2002	8,977	9	33	51	52	78	76	60	107	119	166	242	353	406
2001	9,105	10	56	48	50	71	84	76	99	134	174	264	353	419
2000	9,093	14	53	45	52	72	57	90	81	154	176	246	328	390
1999	9,013	11	44	46	55	70	77	83	103	146	168	260	282	445
Non-Hispanic White														
Both sexes														
2002	18,162	13	60	82	92	140	146	111	178	200	324	429	673	869
2001	18,105	15	95	73	95	128	152	121	177	231	293	479	644	879
2000	18,128	30	88	87	100	136	126	135	149	245	301	441	618	835
1999	17,885	15	78	79	92	133	146	144	171	256	308	473	567	911
Male														
2002	10,218	8	36	41	54	83	84	69	99	108	186	240	386	543
2001	10,018	6	46	40	52	75	89	72	101	123	154	267	365	527
2000	10,009	16	46	50	58	83	83	72	92	128	170	253	340	506
1999	9,853	8	39	45	53	81	96	79	91	142	171	255	338	530
Female														
2002	7,944	5	24	41	38	57	62	42	79	92	138	189	287	326
2001	8,087	9	49	33	43	53	63	49	76	108	139	212	279	352
2000	8,119	14	42	37	42	53	43	63	57	117	131	188	278	329
1999	8,032	7	39	34	39	52	50	65	80	114	137	218	229	381
Non-Hispanic Black														
Both sexes														
2002	1,752	4	13	17	23	29	24	28	40	42	47	93	104	136
2001	1,741	3	13	26	30	26	34	37	38	53	65	75	112	112
2000	1,729	-	17	16	20	46	34	39	49	49	67	107	97	106
1999	1,694	2	12	18	34	35	47	29	38	52	52	79	95	106

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)										
Hispanic										
Female										
1999	18	41	31	40	34	19	17	4	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,428	1,906	2,716	3,305	3,281	2,238	1,060	271	35	-
2001	1,383	1,869	2,808	3,423	3,078	2,207	989	266	48	-
2000	1,300	1,984	2,833	3,411	3,144	2,147	1,017	263	36	-
1999	1,347	1,989	2,837	3,308	2,990	2,113	959	253	31	-
Male										
2002	859	1,195	1,709	1,940	1,762	1,034	411	90	15	-
2001	870	1,168	1,689	1,947	1,643	1,016	369	87	15	-
2000	782	1,238	1,749	1,931	1,670	992	348	82	8	-
1999	818	1,227	1,713	1,898	1,548	947	370	74	9	-
Female										
2002	569	711	1,007	1,365	1,519	1,204	649	181	20	-
2001	513	701	1,119	1,476	1,435	1,191	620	179	33	-
2000	518	746	1,084	1,480	1,474	1,155	669	181	28	-
1999	529	762	1,124	1,410	1,442	1,166	589	179	22	-
Non-Hispanic White										
Both sexes										
2002	1,230	1,704	2,465	3,024	3,063	2,085	993	251	30	-
2001	1,205	1,630	2,555	3,139	2,892	2,084	928	244	46	-
2000	1,117	1,786	2,586	3,168	2,953	2,008	945	243	31	-
1999	1,164	1,751	2,597	3,073	2,775	1,999	892	233	28	-
Male										
2002	747	1,072	1,559	1,793	1,653	973	387	85	12	-
2001	762	1,024	1,555	1,803	1,546	968	348	80	15	-
2000	677	1,129	1,597	1,804	1,573	924	327	74	7	-
1999	716	1,094	1,579	1,767	1,453	894	345	68	9	-
Female										
2002	483	632	906	1,231	1,410	1,112	606	166	18	-
2001	443	606	1,000	1,336	1,346	1,116	580	164	31	-
2000	440	657	989	1,364	1,380	1,084	618	169	24	-
1999	448	657	1,018	1,306	1,322	1,105	547	165	19	-
Non-Hispanic Black										
Both sexes										
2002	159	160	212	228	177	131	61	19	5	-
2001	136	202	206	236	154	108	54	20	1	-
2000	152	160	213	195	154	122	63	18	5	-
1999	146	200	205	196	175	96	56	18	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Leukemia (C91-C95)														
Non-Hispanic Black														
Male														
2002	914	1	6	12	12	18	12	16	21	25	23	57	52	74
2001	915	2	9	13	25	14	17	21	23	32	35	32	61	63
2000	923	-	8	11	11	31	23	18	32	20	32	54	53	60
1999	887	-	7	9	23	23	22	16	22	25	29	47	56	61
Female														
2002	838	3	7	5	11	11	12	12	19	17	24	36	52	62
2001	826	1	4	13	5	12	17	16	15	21	30	43	51	49
2000	806	-	9	5	9	15	11	21	17	29	35	53	44	46
1999	807	2	5	9	11	12	25	13	16	27	23	32	39	45
Origin not stated[2]														
Both sexes														
2002	47	1	-	-	-	-	1	-	1	-	1	1	5	1
2001	52	-	2	1	-	-	1	3	-	2	1	3	1	3
2000	53	-	1	2	1	2	1	-	-	3	4	1	1	6
1999	39	-	1	-	-	-	-	-	2	-	1	1	3	2
Male														
2002	29	1	-	-	-	-	1	-	-	-	-	1	4	1
2001	30	-	1	-	-	-	1	-	-	1	1	2	1	2
2000	27	-	1	-	1	1	-	-	-	2	2	1	1	4
1999	17	-	1	-	-	-	-	-	1	-	1	1	-	2
Female														
2002	18	-	-	-	-	-	-	-	1	-	1	-	1	-
2001	22	-	1	1	-	-	-	3	-	1	-	1	-	1
2000	26	-	-	2	-	1	1	-	-	1	2	-	-	2
1999	22	-	-	-	-	-	-	-	1	-	-	-	3	-
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
All origins														
Both sexes														
2002	11,392	2	-	-	-	2	-	3	9	36	102	270	466	711
2001	11,172	-	-	-	-	-	-	1	8	39	118	241	430	721
2000	11,063	-	-	-	1	-	-	-	7	38	108	245	455	611
1999	11,016	-	-	-	-	-	-	3	19	47	109	241	424	727
Male														
2002	5,953	-	-	-	-	-	-	-	6	24	54	159	274	403
2001	5,724	-	-	-	-	-	-	-	4	22	75	151	252	410
2000	5,553	-	-	-	-	-	-	-	5	18	67	143	248	321
1999	5,584	-	-	-	-	-	-	2	14	33	62	132	246	411

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)										
Non-Hispanic Black										
Male										
2002	87	93	127	114	84	51	21	5	3	-
2001	84	125	107	111	76	42	17	6	-	-
2000	89	86	130	105	76	60	16	7	1	-
1999	81	111	111	108	70	42	20	4	-	-
Female										
2002	72	67	85	114	93	80	40	14	2	-
2001	52	77	99	125	78	66	37	14	1	-
2000	63	74	83	90	78	62	47	11	4	-
1999	65	89	94	88	105	54	36	14	3	-
Origin not stated[2]										
Both sexes										
2002	3	4	9	6	7	3	1	3	-	-
2001	6	4	8	5	5	5	2	-	-	-
2000	1	8	3	9	4	3	3	-	-	-
1999	4	3	4	8	5	2	3	-	-	-
Male										
2002	1	2	7	5	3	2	-	1	-	-
2001	3	2	5	4	3	3	1	-	-	-
2000	-	4	1	3	4	1	1	-	-	-
1999	-	3	4	3	1	-	-	-	-	-
Female										
2002	2	2	2	1	4	1	1	2	-	-
2001	3	2	3	1	2	2	1	-	-	-
2000	1	4	2	6	-	2	2	-	-	-
1999	4	-	-	5	4	2	3	-	-	-
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
All origins										
Both sexes										
2002	958	1,348	1,828	2,067	1,904	1,171	426	84	5	-
2001	950	1,332	1,849	2,101	1,785	1,123	400	71	3	-
2000	919	1,374	1,805	2,097	1,799	1,109	401	84	10	-
1999	967	1,347	1,825	2,023	1,765	1,040	389	85	5	-
Male										
2002	563	748	998	1,090	928	507	159	37	3	-
2001	535	741	984	1,089	837	446	152	23	3	-
2000	487	751	946	1,087	845	462	137	33	3	-
1999	557	729	940	1,026	820	450	133	26	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
All origins														
Female														
2002	5,439	2	-	-	-	2	-	3	3	12	48	111	192	308
2001	5,448	-	-	-	-	-	-	1	4	17	43	90	178	311
2000	5,510	-	-	-	1	-	-	-	2	20	41	102	207	290
1999	5,432	-	-	-	-	-	-	1	5	14	47	109	178	316
Hispanic														
Both sexes														
2002	572	1	-	-	-	1	-	-	3	4	10	24	42	49
2001	524	-	-	-	-	-	-	1	2	5	13	11	32	41
2000	478	-	-	-	-	-	-	-	1	4	8	22	21	31
1999	468	-	-	-	-	-	-	2	2	6	6	21	30	51
Male														
2002	314	-	-	-	-	-	-	-	3	4	6	16	22	30
2001	291	-	-	-	-	-	-	-	1	3	10	6	17	24
2000	239	-	-	-	-	-	-	-	1	1	5	16	8	16
1999	225	-	-	-	-	-	-	2	2	6	3	9	19	29
Female														
2002	258	1	-	-	-	1	-	-	-	-	4	8	20	19
2001	233	-	-	-	-	-	-	1	1	2	3	5	15	17
2000	239	-	-	-	-	-	-	-	-	3	3	6	13	15
1999	243	-	-	-	-	-	-	-	-	-	3	12	11	22
Non-Hispanic[1]														
Both sexes														
2002	10,787	1	-	-	-	1	-	3	6	32	92	245	424	660
2001	10,618	-	-	-	-	-	-	-	6	33	105	230	395	680
2000	10,561	-	-	-	1	-	-	-	6	34	99	221	434	579
1999	10,532	-	-	-	-	-	-	1	17	41	103	220	394	675
Male														
2002	5,623	-	-	-	-	-	-	-	3	20	48	142	252	372
2001	5,418	-	-	-	-	-	-	-	3	19	65	145	233	386
2000	5,300	-	-	-	-	-	-	-	4	17	62	125	240	305
1999	5,352	-	-	-	-	-	-	-	12	27	59	123	227	382
Female														
2002	5,164	1	-	-	-	1	-	3	3	12	44	103	172	288
2001	5,200	-	-	-	-	-	-	-	3	14	40	85	162	294
2000	5,261	-	-	-	1	-	-	-	2	17	37	96	194	274
1999	5,180	-	-	-	-	-	-	1	5	14	44	97	167	293

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
All origins										
Female										
2002	395	600	830	977	976	664	267	47	2	-
2001	415	591	865	1,012	948	677	248	48	-	-
2000	432	623	859	1,010	954	647	264	51	7	-
1999	410	618	885	997	945	590	256	59	2	-
Hispanic										
Both sexes										
2002	49	94	75	94	66	44	13	2	1	-
2001	66	77	92	80	57	31	15	1	-	-
2000	59	78	83	82	42	33	13	1	-	-
1999	60	70	69	60	51	25	13	2	-	-
Male										
2002	32	51	41	51	26	26	5	-	1	-
2001	28	49	53	48	32	12	8	-	-	-
2000	28	47	40	38	23	10	5	1	-	-
1999	31	37	30	27	20	3	5	2	-	-
Female										
2002	17	43	34	43	40	18	8	2	-	-
2001	38	28	39	32	25	19	7	1	-	-
2000	31	31	43	44	19	23	8	-	-	-
1999	29	33	39	33	31	22	8	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	906	1,249	1,749	1,963	1,835	1,124	412	81	4	-
2001	879	1,254	1,752	2,013	1,723	1,090	385	70	3	-
2000	856	1,293	1,717	2,012	1,755	1,074	387	83	10	-
1999	906	1,275	1,755	1,959	1,711	1,012	375	83	5	-
Male										
2002	530	695	954	1,034	900	480	154	37	2	-
2001	504	692	928	1,036	804	433	144	23	3	-
2000	456	701	904	1,047	821	452	131	32	3	-
1999	526	691	910	997	799	445	127	24	3	-
Female										
2002	376	554	795	929	935	644	258	44	2	-
2001	375	562	824	977	919	657	241	47	-	-
2000	400	592	813	965	934	622	256	51	7	-
1999	380	584	845	962	912	567	248	59	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
Non-Hispanic White														
Both sexes														
2002	8,843	-	-	-	-	1	-	2	5	19	59	164	292	486
2001	8,641	-	-	-	-	-	-	-	2	24	57	160	287	495
2000	8,582	-	-	-	1	-	-	-	3	20	64	148	310	412
1999	8,547	-	-	-	-	-	-	-	10	29	63	162	270	480
Male														
2002	4,708	-	-	-	-	-	-	-	2	12	33	99	188	280
2001	4,472	-	-	-	-	-	-	-	2	12	42	104	176	283
2000	4,400	-	-	-	-	-	-	-	2	13	45	88	173	224
1999	4,430	-	-	-	-	-	-	-	6	20	36	94	170	278
Female														
2002	4,135	-	-	-	-	1	-	2	3	7	26	65	104	206
2001	4,169	-	-	-	-	-	-	-	-	12	15	56	111	212
2000	4,182	-	-	-	1	-	-	-	1	7	19	60	137	188
1999	4,117	-	-	-	-	-	-	-	4	9	27	68	100	202
Non-Hispanic Black														
Both sexes														
2002	1,750	-	-	-	-	-	-	1	1	12	28	75	117	157
2001	1,795	-	-	-	-	-	-	-	4	9	44	65	94	167
2000	1,824	-	-	-	-	-	-	-	3	13	31	68	118	156
1999	1,786	-	-	-	-	-	-	1	6	10	39	55	116	177
Male														
2002	830	-	-	-	-	-	-	-	1	7	14	42	57	82
2001	853	-	-	-	-	-	-	-	1	7	21	39	48	93
2000	829	-	-	-	-	-	-	-	2	3	14	33	62	75
1999	824	-	-	-	-	-	-	-	6	6	23	27	53	94
Female														
2002	920	-	-	-	-	-	-	1	-	5	14	33	60	75
2001	942	-	-	-	-	-	-	-	3	2	23	26	46	74
2000	995	-	-	-	-	-	-	-	1	10	17	35	56	81
1999	962	-	-	-	-	-	-	1	-	4	16	28	63	83
Origin not stated[2]														
Both sexes														
2002	33	-	-	-	-	-	-	-	-	-	-	1	-	2
2001	30	-	-	-	-	-	-	-	-	1	-	-	3	-
2000	24	-	-	-	-	-	-	-	-	-	1	2	-	1
1999	16	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	16	-	-	-	-	-	-	-	-	-	-	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
Non-Hispanic White										
Both sexes										
2002	693	977	1,442	1,677	1,606	999	348	71	2	-
2001	649	985	1,442	1,715	1,479	946	339	59	2	-
2000	660	1,028	1,414	1,684	1,536	912	319	65	6	-
1999	665	1,022	1,443	1,636	1,497	870	325	73	2	-
Male										
2002	419	559	812	904	799	434	130	36	1	-
2001	383	555	779	904	699	384	127	20	2	-
2000	362	571	754	894	734	399	110	28	3	-
1999	410	573	756	855	708	393	107	23	1	-
Female										
2002	274	418	630	773	807	565	218	35	1	-
2001	266	430	663	811	780	562	212	39	-	-
2000	298	457	660	790	802	513	209	37	3	-
1999	255	449	687	781	789	477	218	50	1	-
Non-Hispanic Black										
Both sexes										
2002	186	247	277	259	206	116	57	9	2	-
2001	210	237	282	269	224	135	44	10	1	-
2000	180	247	279	297	203	150	62	14	3	-
1999	214	218	276	284	194	136	47	10	3	-
Male										
2002	100	125	132	114	91	42	21	1	1	-
2001	110	119	132	120	100	43	16	3	1	-
2000	88	119	142	140	82	47	19	3	-	-
1999	102	100	137	123	85	47	18	1	2	-
Female										
2002	86	122	145	145	115	74	36	8	1	-
2001	100	118	150	149	124	92	28	7	-	-
2000	92	128	137	157	121	103	43	11	3	-
1999	112	118	139	161	109	89	29	9	1	-
Origin not stated[2]										
Both sexes										
2002	3	5	4	10	3	3	1	1	-	-
2001	5	1	5	8	5	2	-	-	-	-
2000	4	3	5	3	2	2	1	-	-	-
1999	1	2	1	4	3	3	1	-	-	-
Male										
2002	1	2	3	5	2	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)														
Origin not stated[2]														
Male														
2001	15	-	-	-	-	-	-	-	-	-	-	-	2	-
2000	14	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	7	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	17	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	15	-	-	-	-	-	-	-	-	1	-	-	1	-
2000	10	-	-	-	-	-	-	-	-	-	1	-	-	1
1999	9	-	-	-	-	-	-	-	-	-	-	-	-	1
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)														
All origins														
Both sexes														
2002	73	3	-	-	-	-	1	-	-	-	-	2	5	2
2001	90	1	1	-	-	-	-	-	1	1	3	1	3	2
2000	100	-	2	-	1	-	1	-	-	2	2	5	3	6
1999	83	1	1	-	-	-	-	-	1	2	2	3	6	6
Male														
2002	44	3	-	-	-	-	1	-	-	-	-	2	2	1
2001	51	1	-	-	-	-	-	-	1	1	2	1	2	1
2000	60	-	2	-	1	-	1	-	-	1	2	4	2	5
1999	39	1	-	-	-	-	-	-	1	-	1	1	3	5
Female														
2002	29	-	-	-	-	-	-	-	-	-	-	-	3	1
2001	39	-	1	-	-	-	-	-	-	-	1	-	1	1
2000	40	-	-	-	-	-	-	-	-	1	-	1	1	1
1999	44	-	1	-	-	-	-	-	-	2	1	2	3	1
Hispanic														
Both sexes														
2002	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	6	-	-	-	-	-	-	-	-	-	-	1	-	1
1999	5	-	1	-	-	-	-	-	1	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Multiple myeloma and immunoproliferative neoplasms (C88,C90)										
Origin not stated[2]										
Male										
2001	3	-	3	5	1	1	-	-	-	-
2000	3	3	2	2	1	-	1	-	-	-
1999	-	1	-	2	1	2	1	-	-	-
Female										
2002	2	3	1	5	1	2	1	1	-	-
2001	2	1	2	3	4	1	-	-	-	-
2000	1	-	3	1	1	2	-	-	-	-
1999	1	1	1	2	2	1	-	-	-	-
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
All origins										
Both sexes										
2002	1	7	11	12	13	9	5	2	-	-
2001	6	11	12	11	17	11	8	1	-	-
2000	6	7	15	13	16	13	7	1	-	-
1999	6	10	7	14	11	9	4	-	-	-
Male										
2002	1	5	10	4	8	5	1	1	-	-
2001	4	8	10	5	9	4	2	-	-	-
2000	4	7	5	10	7	6	3	-	-	-
1999	2	5	3	9	5	3	-	-	-	-
Female										
2002	-	2	1	8	5	4	4	1	-	-
2001	2	3	2	6	8	7	6	1	-	-
2000	2	-	10	3	9	7	4	1	-	-
1999	4	5	4	5	6	6	4	-	-	-
Hispanic										
Both sexes										
2002	-	1	-	-	-	-	-	-	-	-
2001	-	2	-	-	-	-	-	-	-	-
2000	-	-	2	-	1	1	-	-	-	-
1999	-	-	1	2	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)														
Hispanic														
Male														
2000	3	-	-	-	-	-	-	-	-	-	-	1	-	1
1999	2	-	-	-	-	-	-	-	1	-	-	-	-	-
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	1	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	71	3	-	-	-	-	-	-	-	-	-	2	5	2
2001	88	1	1	-	-	-	-	-	1	1	3	1	3	2
2000	94	-	2	-	1	-	1	-	-	2	2	4	3	5
1999	78	1	-	-	-	-	-	-	-	2	2	3	6	6
Male														
2002	43	3	-	-	-	-	-	-	-	-	-	2	2	1
2001	50	1	-	-	-	-	-	-	1	1	2	1	2	1
2000	57	-	2	-	1	-	1	-	-	1	2	3	2	4
1999	37	1	-	-	-	-	-	-	-	-	1	1	3	5
Female														
2002	28	-	-	-	-	-	-	-	-	-	-	-	3	1
2001	38	-	1	-	-	-	-	-	-	-	1	-	1	1
2000	37	-	-	-	-	-	-	-	-	1	-	1	1	1
1999	41	-	-	-	-	-	-	-	-	2	1	2	3	1
Non-Hispanic White														
Both sexes														
2002	59	1	-	-	-	-	-	-	-	-	-	2	4	2
2001	75	1	1	-	-	-	-	-	-	-	3	1	3	-
2000	83	-	2	-	1	-	-	-	-	1	2	2	3	4
1999	71	1	-	-	-	-	-	-	-	2	1	2	5	5
Male														
2002	37	1	-	-	-	-	-	-	-	-	-	2	2	1
2001	43	1	-	-	-	-	-	-	-	-	2	1	2	-
2000	48	-	2	-	1	-	-	-	-	-	2	1	2	3
1999	34	1	-	-	-	-	-	-	-	-	1	1	2	4
Female														
2002	22	-	-	-	-	-	-	-	-	-	-	-	2	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
Hispanic										
Male										
2000	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Female										
2002	-	1	-	-	-	-	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-
2000	-	-	2	-	1	-	-	-	-	-
1999	-	-	1	1	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1	6	11	12	13	9	5	2	-	-
2001	6	9	12	11	17	11	8	1	-	-
2000	6	7	13	13	15	12	7	1	-	-
1999	6	10	6	12	11	9	4	-	-	-
Male										
2002	1	5	10	4	8	5	1	1	-	-
2001	4	7	10	5	9	4	2	-	-	-
2000	4	7	5	10	7	5	3	-	-	-
1999	2	5	3	8	5	3	-	-	-	-
Female										
2002	-	1	1	8	5	4	4	1	-	-
2001	2	2	2	6	8	7	6	1	-	-
2000	2	-	8	3	8	7	4	1	-	-
1999	4	5	3	4	6	6	4	-	-	-
Non-Hispanic White										
Both sexes										
2002	1	5	11	9	11	9	2	2	-	-
2001	5	8	11	9	14	10	8	1	-	-
2000	6	7	12	11	14	11	6	1	-	-
1999	6	10	6	11	10	8	4	-	-	-
Male										
2002	1	4	10	4	6	5	-	1	-	-
2001	4	6	10	4	8	3	2	-	-	-
2000	4	7	4	9	7	4	2	-	-	-
1999	2	5	3	7	5	3	-	-	-	-
Female										
2002	-	1	1	5	5	4	2	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)														
Non-Hispanic White														
Female														
2001	32	-	1	-	-	-	-	-	-	-	1	-	1	-
2000	35	-	-	-	-	-	-	-	-	1	-	1	1	1
1999	37	-	-	-	-	-	-	-	-	2	-	1	3	1
Non-Hispanic Black														
Both sexes														
2002	6	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	12	-	-	-	-	-	-	-	1	1	-	-	-	2
2000	9	-	-	-	-	-	1	-	-	-	-	2	-	1
1999	6	-	-	-	-	-	-	-	-	-	1	1	1	1
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	7	-	-	-	-	-	-	-	1	1	-	-	-	1
2000	7	-	-	-	-	-	1	-	-	-	-	2	-	1
1999	3	-	-	-	-	-	-	-	-	-	-	-	1	1
Female														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	1	1	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)										
Non-Hispanic White										
Female										
2001	1	2	1	5	6	7	6	1	-	-
2000	2	-	8	2	7	7	4	1	-	-
1999	4	5	3	4	5	5	4	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	1	-	2	1	-	2	-	-	-
2001	1	1	1	1	3	1	-	-	-	-
2000	-	-	-	2	1	1	1	-	-	-
1999	-	-	-	1	-	1	-	-	-	-
Male										
2002	-	1	-	-	1	-	-	-	-	-
2001	-	1	-	1	1	1	-	-	-	-
2000	-	-	-	1	-	1	1	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Female										
2002	-	-	-	2	-	-	2	-	-	-
2001	1	-	1	-	2	-	-	-	-	-
2000	-	-	-	1	1	-	-	-	-	-
1999	-	-	-	-	-	1	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)														
All origins														
Both sexes														
2002	62,969	24	124	133	155	233	294	289	445	716	1,327	2,213	3,280	4,320
2001	62,708	17	117	116	163	242	274	316	412	718	1,337	2,187	3,319	4,278
2000	63,060	24	129	123	137	243	268	292	409	771	1,319	2,114	3,193	4,088
1999	66,251	14	116	150	141	246	273	307	455	833	1,481	2,247	3,230	4,249
Male														
2002	32,122	15	65	77	75	140	200	194	265	407	703	1,225	1,906	2,399
2001	31,663	7	73	64	89	152	194	190	252	385	723	1,231	1,933	2,398
2000	31,728	13	75	64	76	142	177	189	243	424	717	1,186	1,773	2,286
1999	33,449	8	64	80	80	158	171	204	286	464	836	1,274	1,824	2,416
Female														
2002	30,847	9	59	56	80	93	94	95	180	309	624	988	1,374	1,921
2001	31,045	10	44	52	74	90	80	126	160	333	614	956	1,386	1,880
2000	31,332	11	54	59	61	101	91	103	166	347	602	928	1,420	1,802
1999	32,802	6	52	70	61	88	102	103	169	369	645	973	1,406	1,833
Hispanic														
Both sexes														
2002	3,005	11	17	23	32	38	69	41	66	71	105	143	203	232
2001	2,844	2	15	13	29	41	44	36	46	82	106	143	191	200
2000	2,753	3	24	19	28	44	38	46	54	58	83	118	172	226
1999	2,747	2	11	23	22	39	45	48	51	68	107	144	170	216
Male														
2002	1,566	7	9	14	16	25	44	31	38	39	55	76	113	137
2001	1,455	-	9	6	16	25	39	21	35	42	53	78	103	115
2000	1,394	2	16	11	15	25	24	29	38	33	45	60	88	113
1999	1,399	2	3	10	13	27	35	33	30	36	54	75	87	115
Female														
2002	1,439	4	8	9	16	13	25	10	28	32	50	67	90	95
2001	1,389	2	6	7	13	16	5	15	11	40	53	65	88	85
2000	1,359	1	8	8	13	19	14	17	16	25	38	58	84	113
1999	1,348	-	8	13	9	12	10	15	21	32	53	69	83	101
Non-Hispanic[1]														
Both sexes														
2002	59,797	13	106	110	123	194	225	248	377	645	1,222	2,058	3,071	4,078
2001	59,685	15	102	103	134	201	230	280	365	632	1,226	2,037	3,118	4,065
2000	60,117	21	103	103	109	197	230	244	352	711	1,228	1,988	3,009	3,851
1999	63,328	12	105	127	118	207	227	257	398	761	1,369	2,095	3,054	4,021

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)										
All origins										
Both sexes										
2002	5,403	6,321	8,436	9,718	8,967	6,452	3,129	853	136	1
2001	5,163	6,406	8,617	9,902	8,798	6,343	3,019	845	119	-
2000	5,205	6,819	8,897	10,087	8,773	6,214	2,937	874	140	4
1999	5,614	7,265	9,569	10,451	9,205	6,485	2,955	851	112	2
Male										
2002	3,122	3,548	4,593	4,969	4,177	2,690	1,100	226	25	1
2001	2,928	3,501	4,676	4,962	4,070	2,569	1,001	239	26	-
2000	2,937	3,791	4,781	5,098	4,063	2,460	989	218	22	4
1999	3,181	3,984	5,173	5,240	4,199	2,586	986	213	21	1
Female										
2002	2,281	2,773	3,843	4,749	4,790	3,762	2,029	627	111	-
2001	2,235	2,905	3,941	4,940	4,728	3,774	2,018	606	93	-
2000	2,268	3,028	4,116	4,989	4,710	3,754	1,948	656	118	-
1999	2,433	3,281	4,396	5,211	5,006	3,899	1,969	638	91	1
Hispanic										
Both sexes										
2002	270	318	394	385	291	191	83	20	2	-
2001	281	295	395	347	268	189	83	35	3	-
2000	229	317	373	357	274	167	88	32	3	-
1999	255	325	346	336	251	183	80	20	5	-
Male										
2002	140	164	198	199	139	81	33	8	-	-
2001	147	141	198	191	124	79	20	12	1	-
2000	122	163	186	187	122	64	36	14	1	-
1999	141	161	190	159	119	73	32	4	-	-
Female										
2002	130	154	196	186	152	110	50	12	2	-
2001	134	154	197	156	144	110	63	23	2	-
2000	107	154	187	170	152	103	52	18	2	-
1999	114	164	156	177	132	110	48	16	5	-
Non-Hispanic[1]										
Both sexes										
2002	5,114	5,981	8,024	9,308	8,657	6,241	3,037	830	134	1
2001	4,870	6,098	8,180	9,534	8,508	6,140	2,923	808	116	-
2000	4,962	6,484	8,496	9,693	8,479	6,031	2,844	841	137	4
1999	5,340	6,926	9,195	10,091	8,929	6,286	2,871	831	106	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)														
Non-Hispanic[1]														
Male														
2002	30,463	8	56	63	59	114	156	163	226	368	648	1,140	1,789	2,258
2001	30,107	7	64	58	73	127	155	169	216	339	667	1,149	1,822	2,275
2000	30,237	11	57	53	61	116	153	158	204	391	668	1,121	1,682	2,164
1999	31,956	6	61	70	67	131	135	169	254	427	779	1,194	1,734	2,295
Female														
2002	29,334	5	50	47	64	80	69	85	151	277	574	918	1,282	1,820
2001	29,578	8	38	45	61	74	75	111	149	293	559	888	1,296	1,790
2000	29,880	10	46	50	48	81	77	86	148	320	560	867	1,327	1,687
1999	31,372	6	44	57	51	76	92	88	144	334	590	901	1,320	1,726
Non-Hispanic White														
Both sexes														
2002	51,731	11	73	79	92	153	166	185	280	494	939	1,590	2,442	3,367
2001	51,581	13	83	79	97	156	182	219	273	482	948	1,558	2,466	3,314
2000	52,223	16	75	87	81	142	186	203	276	537	915	1,532	2,376	3,201
1999	54,977	10	87	95	83	158	166	207	306	587	1,039	1,624	2,453	3,273
Male														
2002	26,457	8	40	45	43	88	113	123	172	284	485	892	1,440	1,871
2001	26,007	6	50	44	52	96	124	132	164	266	526	890	1,436	1,845
2000	26,258	9	46	44	45	84	120	134	169	304	493	864	1,349	1,804
1999	27,780	5	52	55	49	94	96	143	195	339	604	915	1,390	1,885
Female														
2002	25,274	3	33	34	49	65	53	62	108	210	454	698	1,002	1,496
2001	25,574	7	33	35	45	60	58	87	109	216	422	668	1,030	1,469
2000	25,965	7	29	43	36	58	66	69	107	233	422	668	1,027	1,397
1999	27,197	5	35	40	34	64	70	64	111	248	435	709	1,063	1,388
Non-Hispanic Black														
Both sexes														
2002	6,756	2	26	26	26	33	46	51	74	131	231	398	526	613
2001	6,725	1	16	19	27	37	39	52	72	122	217	409	554	623
2000	6,617	4	22	16	23	47	36	34	53	152	275	392	535	552
1999	7,009	1	14	22	27	39	44	41	65	144	277	416	512	629
Male														
2002	3,340	-	12	15	13	22	32	31	36	70	130	214	295	335
2001	3,392	-	13	12	14	23	28	31	38	58	108	215	328	364
2000	3,337	1	9	9	15	27	29	20	26	74	153	227	283	308
1999	3,514	-	8	9	13	29	27	20	40	70	149	244	295	361

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)										
Non-Hispanic[1]										
Male										
2002	2,967	3,370	4,385	4,757	4,025	2,604	1,063	218	25	1
2001	2,771	3,351	4,449	4,765	3,936	2,484	978	227	25	-
2000	2,808	3,616	4,579	4,892	3,931	2,390	953	204	21	4
1999	3,030	3,816	4,963	5,065	4,067	2,509	953	209	21	1
Female										
2002	2,147	2,611	3,639	4,551	4,632	3,637	1,974	612	109	-
2001	2,099	2,747	3,731	4,769	4,572	3,656	1,945	581	91	-
2000	2,154	2,868	3,917	4,801	4,548	3,641	1,891	637	116	-
1999	2,310	3,110	4,232	5,026	4,862	3,777	1,918	622	85	1
Non-Hispanic White										
Both sexes										
2002	4,268	5,072	6,999	8,306	7,853	5,705	2,785	750	121	1
2001	4,028	5,171	7,121	8,536	7,697	5,641	2,685	732	100	-
2000	4,129	5,583	7,436	8,644	7,751	5,548	2,638	752	112	3
1999	4,449	6,006	8,070	9,032	8,126	5,756	2,606	750	92	2
Male										
2002	2,491	2,920	3,870	4,268	3,698	2,402	981	198	24	1
2001	2,294	2,857	3,932	4,292	3,590	2,292	887	211	21	-
2000	2,307	3,159	4,049	4,356	3,633	2,203	883	181	19	3
1999	2,538	3,321	4,393	4,580	3,742	2,313	866	184	20	1
Female										
2002	1,777	2,152	3,129	4,038	4,155	3,303	1,804	552	97	-
2001	1,734	2,314	3,189	4,244	4,107	3,349	1,798	521	79	-
2000	1,822	2,424	3,387	4,288	4,118	3,345	1,755	571	93	-
1999	1,911	2,685	3,677	4,452	4,384	3,443	1,740	566	72	1
Non-Hispanic Black										
Both sexes										
2002	716	769	843	850	657	444	215	67	12	-
2001	710	782	877	807	661	415	205	66	14	-
2000	703	767	869	865	607	396	166	82	20	1
1999	778	766	938	876	663	451	223	69	14	-
Male										
2002	404	387	426	405	270	161	65	16	1	-
2001	395	429	437	383	267	158	73	14	4	-
2000	425	387	440	440	247	144	52	19	1	1
1999	440	418	477	401	266	153	71	22	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)														
Non-Hispanic Black														
Female														
2002	3,416	2	14	11	13	11	14	20	38	61	101	184	231	278
2001	3,333	1	3	7	13	14	11	21	34	64	109	194	226	259
2000	3,280	3	13	7	8	20	7	14	27	78	122	165	252	244
1999	3,495	1	6	13	14	10	17	21	25	74	128	172	217	268
Origin not stated[2]														
Both sexes														
2002	167	-	1	-	-	1	-	-	2	-	-	12	6	10
2001	179	-	-	-	-	-	-	-	1	4	5	7	10	13
2000	190	-	2	1	-	2	-	2	3	2	8	8	12	11
1999	176	-	-	-	1	-	1	2	6	4	5	8	6	12
Male														
2002	93	-	-	-	-	1	-	-	1	-	-	9	4	4
2001	101	-	-	-	-	-	-	-	1	4	3	4	8	8
2000	97	-	2	-	-	1	-	2	1	-	4	5	3	9
1999	94	-	-	-	-	-	1	2	2	1	3	5	3	6
Female														
2002	74	-	1	-	-	-	-	-	1	-	-	3	2	6
2001	78	-	-	-	-	-	-	-	-	-	2	3	2	5
2000	93	-	-	1	-	1	-	-	2	2	4	3	9	2
1999	82	-	-	-	1	-	-	-	4	3	2	3	3	6
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
All origins														
Both sexes														
2002	13,299	69	60	44	45	41	46	62	101	137	229	327	387	561
2001	13,676	51	58	52	53	47	55	60	91	156	210	304	415	553
2000	13,546	60	53	62	37	40	55	68	97	155	213	293	397	513
1999	13,227	58	63	64	37	52	46	55	116	140	202	276	386	461
Male														
2002	6,536	30	30	22	27	21	29	33	51	77	119	169	196	330
2001	6,735	26	30	26	26	20	33	33	44	79	114	173	218	279
2000	6,559	32	33	32	17	27	35	30	52	86	102	160	225	270
1999	6,520	30	33	30	19	26	28	32	64	79	118	142	220	255

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31, C37-C41,C44-C49,C51-C52, C57-C60,C62-C63,C66,C68-C69, C73-C80,C97)										
Non-Hispanic Black										
Female										
2002	312	382	417	445	387	283	150	51	11	-
2001	315	353	440	424	394	257	132	52	10	-
2000	278	380	429	425	360	252	114	63	19	-
1999	338	348	461	475	397	298	152	47	13	-
Origin not stated[2]										
Both sexes										
2002	19	22	18	25	19	20	9	3	-	-
2001	12	13	42	21	22	14	13	2	-	-
2000	14	18	28	37	20	16	5	1	-	-
1999	19	14	28	24	25	16	4	-	1	-
Male										
2002	15	14	10	13	13	5	4	-	-	-
2001	10	9	29	6	10	6	3	-	-	-
2000	7	12	16	19	10	6	-	-	-	-
1999	10	7	20	16	13	4	1	-	-	-
Female										
2002	4	8	8	12	6	15	5	3	-	-
2001	2	4	13	15	12	8	10	2	-	-
2000	7	6	12	18	10	10	5	1	-	-
1999	9	7	8	8	12	12	3	-	1	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All origins										
Both sexes										
2002	632	928	1,420	2,137	2,460	2,052	1,162	366	33	-
2001	700	1,000	1,574	2,192	2,462	2,102	1,141	351	49	-
2000	711	1,007	1,539	2,142	2,503	2,103	1,143	314	40	1
1999	672	1,029	1,655	2,151	2,321	1,998	1,059	337	49	-
Male										
2002	350	543	772	1,145	1,187	900	398	99	8	-
2001	394	563	876	1,199	1,204	882	404	102	10	-
2000	387	587	828	1,120	1,199	881	362	86	7	1
1999	401	590	932	1,119	1,099	839	367	82	15	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
All origins														
Female														
2002	6,763	39	30	22	18	20	17	29	50	60	110	158	191	231
2001	6,941	25	28	26	27	27	22	27	47	77	96	131	197	274
2000	6,987	28	20	30	20	13	20	38	45	69	111	133	172	243
1999	6,707	28	30	34	18	26	18	23	52	61	84	134	166	206
Hispanic														
Both sexes														
2002	563	17	7	16	10	4	4	5	11	14	20	30	24	20
2001	543	5	13	14	7	4	4	8	12	14	25	22	28	25
2000	521	6	7	5	8	3	7	8	17	14	23	20	23	23
1999	497	8	12	11	6	7	5	8	17	6	24	15	24	31
Male														
2002	282	9	4	7	8	2	3	3	5	7	8	16	10	14
2001	276	2	7	9	3	2	4	4	2	6	18	14	16	10
2000	237	4	2	4	4	2	6	2	13	7	7	9	9	10
1999	233	5	4	5	3	5	3	7	7	4	17	9	17	14
Female														
2002	281	8	3	9	2	2	1	2	6	7	12	14	14	6
2001	267	3	6	5	4	2	-	4	10	8	7	8	12	15
2000	284	2	5	1	4	1	1	6	4	7	16	11	14	13
1999	264	3	8	6	3	2	2	1	10	2	7	6	7	17
Non-Hispanic[1]														
Both sexes														
2002	12,708	52	53	28	35	37	41	57	90	123	205	297	362	539
2001	13,098	46	45	38	46	43	51	52	79	142	185	282	385	527
2000	12,990	53	46	57	28	37	48	60	80	141	189	271	374	487
1999	12,699	50	51	52	31	44	41	47	99	134	177	261	359	428
Male														
2002	6,237	21	26	15	19	19	25	30	46	70	109	153	186	316
2001	6,442	24	23	17	23	18	29	29	42	73	96	159	201	269
2000	6,302	28	31	28	13	25	29	28	39	79	94	149	216	257
1999	6,276	25	29	25	16	21	25	25	57	75	101	133	201	239
Female														
2002	6,471	31	27	13	16	18	16	27	44	53	96	144	176	223
2001	6,656	22	22	21	23	25	22	23	37	69	89	123	184	258
2000	6,688	25	15	29	15	12	19	32	41	62	95	122	158	230
1999	6,423	25	22	27	15	23	16	22	42	59	76	128	158	189

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
All origins										
Female										
2002	282	385	648	992	1,273	1,152	764	267	25	-
2001	306	437	698	993	1,258	1,220	737	249	39	-
2000	324	420	711	1,022	1,304	1,222	781	228	33	-
1999	271	439	723	1,032	1,222	1,159	692	255	34	-
Hispanic										
Both sexes										
2002	40	41	64	65	83	51	31	5	1	-
2001	42	47	53	67	61	44	34	11	3	-
2000	30	40	58	67	56	63	29	12	2	-
1999	29	39	70	57	54	38	26	9	1	-
Male										
2002	21	23	31	31	46	24	9	-	1	-
2001	25	21	29	29	38	18	15	3	1	-
2000	15	16	30	36	18	26	12	5	-	-
1999	16	13	34	24	23	14	6	2	1	-
Female										
2002	19	18	33	34	37	27	22	5	-	-
2001	17	26	24	38	23	26	19	8	2	-
2000	15	24	28	31	38	37	17	7	2	-
1999	13	26	36	33	31	24	20	7	-	-
Non-Hispanic[1]										
Both sexes										
2002	592	887	1,354	2,065	2,371	1,999	1,128	361	32	-
2001	655	951	1,517	2,120	2,390	2,055	1,103	340	46	-
2000	680	966	1,479	2,072	2,439	2,033	1,109	302	38	1
1999	642	988	1,584	2,088	2,261	1,957	1,030	327	48	-
Male										
2002	329	520	740	1,107	1,138	874	388	99	7	-
2001	367	540	845	1,166	1,162	863	388	99	9	-
2000	372	570	797	1,081	1,177	851	349	81	7	1
1999	384	576	898	1,093	1,075	825	360	79	14	-
Female										
2002	263	367	614	958	1,233	1,125	740	262	25	-
2001	288	411	672	954	1,228	1,192	715	241	37	-
2000	308	396	682	991	1,262	1,182	760	221	31	-
1999	258	412	686	995	1,186	1,132	670	248	34	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
Non-Hispanic White														
Both sexes														
2002	11,332	42	36	20	21	30	29	37	68	98	153	245	285	449
2001	11,701	32	36	24	32	29	37	45	61	110	142	235	326	451
2000	11,636	41	37	41	22	26	37	40	65	110	145	206	310	423
1999	11,335	34	36	40	26	30	28	31	68	101	138	199	295	357
Male														
2002	5,607	18	17	9	13	16	16	21	41	58	91	131	152	261
2001	5,801	18	18	10	15	10	21	25	33	58	77	135	177	234
2000	5,653	21	23	20	11	16	24	22	33	61	71	118	183	229
1999	5,653	17	18	19	13	12	18	14	39	59	84	107	169	200
Female														
2002	5,725	24	19	11	8	14	13	16	27	40	62	114	133	188
2001	5,900	14	18	14	17	19	16	20	28	52	65	100	149	217
2000	5,983	20	14	21	11	10	13	18	32	49	74	88	127	194
1999	5,682	17	18	21	13	18	10	17	29	42	54	92	126	157
Non-Hispanic Black														
Both sexes														
2002	1,134	9	14	7	11	6	9	13	18	20	47	49	63	77
2001	1,143	11	5	13	13	12	9	6	12	26	35	35	54	63
2000	1,122	9	9	13	5	8	8	18	15	24	38	54	58	55
1999	1,161	14	13	12	5	12	13	9	21	30	35	50	54	64
Male														
2002	496	2	8	6	3	3	6	5	3	8	17	19	27	50
2001	522	5	3	6	7	7	6	3	8	13	17	16	22	28
2000	514	4	8	6	2	7	2	5	6	13	19	23	31	25
1999	516	7	10	6	3	8	7	8	12	15	15	19	25	36
Female														
2002	638	7	6	1	8	3	3	8	15	12	30	30	36	27
2001	621	6	2	7	6	5	3	3	4	13	18	19	32	35
2000	608	5	1	7	3	1	6	13	9	11	19	31	27	30
1999	645	7	3	6	2	4	6	1	9	15	20	31	29	28
Origin not stated[2]														
Both sexes														
2002	28	-	-	-	-	-	1	-	-	-	4	-	1	2
2001	35	-	-	-	-	-	-	-	-	-	-	-	2	1
2000	35	1	-	-	1	-	-	-	-	-	1	2	-	3
1999	31	-	-	1	-	1	-	-	-	-	1	-	3	2
Male														
2002	17	-	-	-	-	-	1	-	-	-	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
Non-Hispanic White										
Both sexes										
2002	513	772	1,213	1,890	2,173	1,869	1,043	319	27	-
2001	555	837	1,348	1,945	2,205	1,897	1,015	303	36	-
2000	581	847	1,297	1,894	2,277	1,892	1,033	279	33	-
1999	542	846	1,430	1,899	2,102	1,829	967	294	43	-
Male										
2002	286	457	678	1,023	1,046	824	357	85	7	-
2001	313	483	767	1,078	1,074	796	361	91	7	-
2000	310	500	702	994	1,114	789	328	77	7	-
1999	336	503	823	1,002	1,011	778	345	73	13	-
Female										
2002	227	315	535	867	1,127	1,045	686	234	20	-
2001	242	354	581	867	1,131	1,101	654	212	29	-
2000	271	347	595	900	1,163	1,103	705	202	26	-
1999	206	343	607	897	1,091	1,051	622	221	30	-
Non-Hispanic Black										
Both sexes										
2002	67	99	112	141	159	108	67	34	4	-
2001	84	99	137	141	155	116	75	33	9	-
2000	78	96	149	149	133	116	62	20	4	1
1999	84	124	132	162	133	108	52	30	4	-
Male										
2002	33	50	48	64	70	41	22	11	-	-
2001	45	51	68	68	73	45	22	7	2	-
2000	49	60	70	72	46	44	18	3	-	1
1999	39	65	63	75	49	38	10	6	-	-
Female										
2002	34	49	64	77	89	67	45	23	4	-
2001	39	48	69	73	82	71	53	26	7	-
2000	29	36	79	77	87	72	44	17	4	-
1999	45	59	69	87	84	70	42	24	4	-
Origin not stated[2]										
Both sexes										
2002	-	-	2	7	6	2	3	-	-	-
2001	3	2	4	5	11	3	4	-	-	-
2000	1	1	2	3	8	7	5	-	-	-
1999	1	2	1	6	6	3	3	1	-	-
Male										
2002	-	-	1	7	3	2	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)														
Origin not stated[2]														
Male														
2001	17	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	20	-	-	-	-	-	-	-	-	-	1	2	-	3
1999	11	-	-	-	-	-	-	-	-	-	-	-	2	2
Female														
2002	11	-	-	-	-	-	-	-	-	-	2	-	1	2
2001	18	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	15	1	-	-	1	-	-	-	-	-	-	-	-	-
1999	20	-	-	1	-	1	-	-	-	-	1	-	1	-
Anemias (D50-D64)														
All origins														
Both sexes														
2002	4,614	18	21	29	24	42	69	83	97	99	97	131	119	121
2001	4,627	16	22	25	30	48	61	65	95	83	132	112	120	119
2000	4,494	10	35	24	28	27	58	62	91	85	105	110	117	127
1999	4,503	12	30	24	29	43	78	64	96	90	104	134	98	105
Male														
2002	1,781	12	9	13	13	23	44	53	48	53	43	57	52	57
2001	1,774	11	12	12	15	26	33	40	55	42	49	57	58	53
2000	1,669	6	16	17	12	9	30	29	41	34	44	52	56	66
1999	1,746	8	17	8	15	24	43	33	48	47	54	68	48	43
Female														
2002	2,833	6	12	16	11	19	25	30	49	46	54	74	67	64
2001	2,853	5	10	13	15	22	28	25	40	41	83	55	62	66
2000	2,825	4	19	7	16	18	28	33	50	51	61	58	61	61
1999	2,757	4	13	16	14	19	35	31	48	43	50	66	50	62
Hispanic														
Both sexes														
2002	208	2	3	3	1	3	5	5	6	6	5	8	4	3
2001	185	-	3	2	3	6	5	4	8	5	7	6	5	11
2000	144	1	1	2	5	1	4	9	4	4	4	5	5	6
1999	164	-	6	2	1	2	4	4	4	5	5	6	3	5
Male														
2002	106	1	1	2	1	1	4	2	3	5	4	5	2	3
2001	82	-	2	2	3	2	4	3	5	3	3	2	2	4
2000	55	1	-	1	2	-	1	4	1	-	1	1	3	1
1999	79	-	4	1	-	1	2	2	3	3	3	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)										
Origin not stated[2]										
Male										
2001	2	2	2	4	4	1	1	-	-	-
2000	-	1	1	3	4	4	1	-	-	-
1999	1	1	-	2	1	-	1	1	-	-
Female										
2002	-	-	1	-	3	-	2	-	-	-
2001	1	-	2	1	7	2	3	-	-	-
2000	1	-	1	-	4	3	4	-	-	-
1999	-	1	1	4	5	3	2	-	-	-
Anemias (D50-D64)										
All origins										
Both sexes										
2002	143	184	324	474	687	787	665	322	78	-
2001	130	186	306	525	701	808	659	294	90	-
2000	133	210	305	482	689	790	638	290	78	-
1999	126	175	311	497	667	775	647	316	82	-
Male										
2002	74	77	167	204	250	256	182	81	13	-
2001	58	91	146	220	276	276	177	52	15	-
2000	64	95	141	198	267	237	183	58	14	-
1999	62	92	134	217	267	261	172	73	12	-
Female										
2002	69	107	157	270	437	531	483	241	65	-
2001	72	95	160	305	425	532	482	242	75	-
2000	69	115	164	284	422	553	455	232	64	-
1999	64	83	177	280	400	514	475	243	70	-
Hispanic										
Both sexes										
2002	11	14	23	18	26	21	26	14	1	-
2001	6	14	11	24	19	23	15	6	2	-
2000	7	9	15	13	23	11	12	3	-	-
1999	5	7	20	16	19	21	17	11	1	-
Male										
2002	5	6	11	9	9	11	14	7	-	-
2001	3	9	3	12	9	7	2	1	1	-
2000	3	4	8	4	11	5	4	-	-	-
1999	3	4	13	6	11	7	7	5	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Anemias (D50-D64)														
Hispanic														
Female														
2002	102	1	2	1	-	2	1	3	3	1	1	3	2	-
2001	103	-	1	-	-	4	1	1	3	2	4	4	3	7
2000	89	-	1	1	3	1	3	5	3	4	3	4	2	5
1999	85	-	2	1	1	1	2	2	1	2	2	2	3	5
Non-Hispanic[1]														
Both sexes														
2002	4,391	14	17	26	23	39	64	78	89	91	91	122	115	118
2001	4,424	15	19	23	27	41	56	61	85	77	125	105	114	107
2000	4,336	8	34	21	23	26	53	53	87	81	100	105	110	121
1999	4,321	12	24	22	28	41	74	60	92	85	98	126	95	100
Male														
2002	1,671	9	8	11	12	22	40	51	45	47	39	52	50	54
2001	1,687	10	10	10	12	24	29	37	49	38	46	54	56	49
2000	1,608	4	16	16	10	9	28	25	40	34	43	51	53	65
1999	1,659	8	13	7	15	23	41	31	45	44	51	63	48	43
Female														
2002	2,720	5	9	15	11	17	24	27	44	44	52	70	65	64
2001	2,737	5	9	13	15	17	27	24	36	39	79	51	58	58
2000	2,728	4	18	5	13	17	25	28	47	47	57	54	57	56
1999	2,662	4	11	15	13	18	33	29	47	41	47	63	47	57
Non-Hispanic White														
Both sexes														
2002	3,253	5	10	12	7	10	13	11	22	16	18	42	48	68
2001	3,350	5	8	9	13	11	11	11	16	11	30	25	56	57
2000	3,299	3	14	7	7	9	7	6	10	13	23	33	49	78
1999	3,222	6	4	5	12	16	12	3	7	17	20	48	38	41
Male														
2002	1,164	2	4	4	4	6	7	9	11	10	8	20	25	33
2001	1,220	4	3	4	7	6	6	6	6	6	9	10	27	27
2000	1,178	1	8	6	3	2	5	5	5	7	13	15	24	44
1999	1,162	4	2	-	6	8	3	-	2	7	11	26	20	16
Female														
2002	2,089	3	6	8	3	4	6	2	11	6	10	22	23	35
2001	2,130	1	5	5	6	5	5	5	10	5	21	15	29	30
2000	2,121	2	6	1	4	7	2	1	5	6	10	18	25	34
1999	2,060	2	2	5	6	8	9	3	5	10	9	22	18	25
Non-Hispanic Black														
Both sexes														
2002	1,048	5	7	12	14	27	51	64	63	72	70	78	63	47

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)										
Hispanic										
Female										
2002	6	8	12	9	17	10	12	7	1	-
2001	3	5	8	12	10	16	13	5	1	-
2000	4	5	7	9	12	6	8	3	-	-
1999	2	3	7	10	8	14	10	6	1	-
Non-Hispanic[1]										
Both sexes										
2002	132	169	301	455	659	765	639	307	77	-
2001	124	172	294	500	677	783	644	288	87	-
2000	125	201	288	468	664	779	625	287	77	-
1999	120	165	290	478	647	751	630	302	81	-
Male										
2002	69	71	156	195	240	245	168	74	13	-
2001	55	82	143	208	266	269	175	51	14	-
2000	61	91	131	193	255	232	179	58	14	-
1999	58	87	120	209	256	254	165	66	12	-
Female										
2002	63	98	145	260	419	520	471	233	64	-
2001	69	90	151	292	411	514	469	237	73	-
2000	64	110	157	275	409	547	446	229	63	-
1999	62	78	170	269	391	497	465	236	69	-
Non-Hispanic White										
Both sexes										
2002	79	126	235	368	576	686	559	270	72	-
2001	93	132	236	422	591	703	584	260	66	-
2000	77	151	233	401	593	707	553	258	67	-
1999	71	126	234	410	585	668	566	266	67	-
Male										
2002	42	55	117	164	207	220	143	61	12	-
2001	41	65	110	181	238	252	157	45	10	-
2000	39	77	108	165	227	207	152	52	13	-
1999	29	70	93	183	233	223	153	61	12	-
Female										
2002	37	71	118	204	369	466	416	209	60	-
2001	52	67	126	241	353	451	427	215	56	-
2000	38	74	125	236	366	500	401	206	54	-
1999	42	56	141	227	352	445	413	205	55	-
Non-Hispanic Black										
Both sexes										
2002	47	41	59	80	73	70	68	33	4	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Anemias (D50-D64)														
Non-Hispanic Black														
Both sexes														
2001	993	6	11	13	13	27	45	50	67	62	92	78	57	46
2000	961	4	19	14	16	16	45	47	77	65	74	69	59	40
1999	1,010	3	20	17	16	23	61	55	84	66	76	73	52	55
Male														
2002	472	4	4	7	7	16	33	39	33	35	31	31	24	20
2001	443	4	7	5	5	18	23	31	42	30	36	44	29	21
2000	392	3	8	10	7	7	22	20	35	26	29	35	29	18
1999	453	2	11	7	9	13	37	29	43	35	38	35	26	25
Female														
2002	576	1	3	5	7	11	18	25	30	37	39	47	39	27
2001	550	2	4	8	8	9	22	19	25	32	56	34	28	25
2000	569	1	11	4	9	9	23	27	42	39	45	34	30	22
1999	557	1	9	10	7	10	24	26	41	31	38	38	26	30
Origin not stated[2]														
Both sexes														
2002	15	2	1	-	-	-	-	-	2	2	1	1	-	-
2001	18	1	-	-	-	1	-	-	2	1	-	1	1	1
2000	14	1	-	1	-	-	1	-	-	-	1	-	2	-
1999	18	-	-	-	-	-	-	-	-	-	1	2	-	-
Male														
2002	4	2	-	-	-	-	-	-	-	1	-	-	-	-
2001	5	1	-	-	-	-	-	-	1	1	-	1	-	-
2000	6	1	-	-	-	-	1	-	-	-	-	-	-	-
1999	8	-	-	-	-	-	-	-	-	-	-	1	-	-
Female														
2002	11	-	1	-	-	-	-	-	2	1	1	1	-	-
2001	13	-	-	-	-	1	-	-	1	-	-	-	1	1
2000	8	-	-	1	-	-	-	-	-	-	1	-	2	-
1999	10	-	-	-	-	-	-	-	-	-	1	1	-	-
Diabetes mellitus (E10-E14)														
All origins														
Both sexes														
2002	73,249	1	3	5	29	55	116	229	413	776	1,388	2,106	3,390	4,510
2001	71,372	2	6	7	32	48	103	215	380	735	1,223	2,063	3,280	4,188
2000	69,301	4	6	8	18	48	114	206	417	684	1,242	1,917	3,037	3,992
1999	68,399	2	8	9	23	51	104	189	393	730	1,212	1,942	2,793	3,865
Male														
2002	34,301	-	1	4	18	30	68	128	257	452	878	1,248	1,977	2,616
2001	32,841	1	2	2	21	27	58	114	227	436	745	1,234	1,920	2,362

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Anemias (D50-D64)										
Non-Hispanic Black										
Both sexes										
2001	29	36	54	71	78	66	50	24	18	-
2000	43	43	46	58	63	64	61	28	10	-
1999	45	33	48	63	56	67	57	28	12	-
Male										
2002	26	16	35	28	30	21	22	9	1	-
2001	14	15	31	26	25	15	13	6	3	-
2000	19	13	18	23	22	20	22	5	1	-
1999	28	14	23	22	19	25	9	3	-	-
Female										
2002	21	25	24	52	43	49	46	24	3	-
2001	15	21	23	45	53	51	37	18	15	-
2000	24	30	28	35	41	44	39	23	9	-
1999	17	19	25	41	37	42	48	25	12	-
Origin not stated[2]										
Both sexes										
2002	-	1	-	1	2	1	-	1	-	-
2001	-	-	1	1	5	2	-	-	1	-
2000	1	-	2	1	2	-	1	-	1	-
1999	1	3	1	3	1	3	-	3	-	-
Male										
2002	-	-	-	-	1	-	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	-	2	1	1	-	-	-	-	-
1999	1	1	1	2	-	-	-	2	-	-
Female										
2002	-	1	-	1	1	1	-	1	-	-
2001	-	-	1	1	4	2	-	-	1	-
2000	1	-	-	-	1	-	1	-	1	-
1999	-	2	-	1	1	3	-	1	-	-
Diabetes mellitus (E10-E14)										
All origins										
Both sexes										
2002	5,512	7,249	9,460	11,703	11,579	8,797	4,375	1,357	195	1
2001	5,382	7,195	9,536	11,546	11,259	8,366	4,259	1,350	196	1
2000	5,194	7,054	9,620	11,607	10,577	8,002	4,104	1,250	200	-
1999	5,232	7,215	9,693	11,269	10,488	7,893	3,943	1,165	177	3
Male										
2002	3,067	3,742	4,675	5,402	4,893	3,248	1,254	314	28	1
2001	2,879	3,734	4,703	5,244	4,674	2,966	1,188	269	34	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diabetes mellitus (E10-E14)														
All origins														
Male														
2000	31,602	4	2	5	8	31	63	118	258	396	755	1,124	1,697	2,173
1999	31,150	-	4	4	12	25	55	94	225	437	702	1,189	1,602	2,120
Female														
2002	38,948	1	2	1	11	25	48	101	156	324	510	858	1,413	1,894
2001	38,531	1	4	5	11	21	45	101	153	299	478	829	1,360	1,826
2000	37,699	-	4	3	10	17	51	88	159	288	487	793	1,340	1,819
1999	37,249	2	4	5	11	26	49	95	168	293	510	753	1,191	1,745
Hispanic														
Both sexes														
2002	5,912	-	-	-	1	5	10	32	24	73	132	191	322	432
2001	5,663	1	1	-	2	8	11	20	29	69	98	229	278	404
2000	5,328	1	-	1	3	5	8	21	43	49	105	174	265	436
1999	5,182	-	-	-	4	2	12	14	29	56	101	169	244	368
Male														
2002	2,779	-	-	-	1	3	7	19	15	41	83	107	198	236
2001	2,590	-	1	-	2	5	6	9	19	40	69	139	172	217
2000	2,507	1	-	-	2	4	3	16	32	28	67	98	142	241
1999	2,335	-	-	-	2	-	8	8	13	31	53	105	147	202
Female														
2002	3,133	-	-	-	-	2	3	13	9	32	49	84	124	196
2001	3,073	1	-	-	-	3	5	11	10	29	29	90	106	187
2000	2,821	-	-	1	1	1	5	5	11	21	38	76	123	195
1999	2,847	-	-	-	2	2	4	6	16	25	48	64	97	166
Non-Hispanic[1]														
Both sexes														
2002	67,112	1	3	5	28	50	105	196	388	697	1,249	1,902	3,059	4,065
2001	65,498	1	5	7	30	40	91	193	349	665	1,122	1,825	2,985	3,769
2000	63,769	3	6	7	15	43	106	185	372	631	1,131	1,741	2,760	3,545
1999	63,017	2	8	9	19	48	92	173	361	672	1,109	1,769	2,536	3,484
Male														
2002	31,399	-	1	4	17	27	60	108	241	407	789	1,132	1,774	2,374
2001	30,144	1	1	2	19	22	52	105	206	395	674	1,088	1,740	2,136
2000	28,994	3	2	5	6	27	60	102	226	364	684	1,025	1,549	1,924
1999	28,719	-	4	4	10	25	47	85	210	406	647	1,082	1,443	1,911
Female														
2002	35,713	1	2	1	11	23	45	88	147	290	460	770	1,285	1,691
2001	35,354	-	4	5	11	18	39	88	143	270	448	737	1,245	1,633
2000	34,775	-	4	2	9	16	46	83	146	267	447	716	1,211	1,621
1999	34,298	2	4	5	9	23	45	88	151	266	462	687	1,093	1,573

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)										
All origins										
Male										
2000	2,770	3,556	4,670	5,288	4,450	2,789	1,165	260	20	-
1999	2,736	3,630	4,649	5,190	4,349	2,816	1,071	210	28	2
Female										
2002	2,445	3,507	4,785	6,301	6,686	5,549	3,121	1,043	167	-
2001	2,503	3,461	4,833	6,302	6,585	5,400	3,071	1,081	162	-
2000	2,424	3,498	4,950	6,319	6,127	5,213	2,939	990	180	-
1999	2,496	3,585	5,044	6,079	6,139	5,077	2,872	955	149	1
Hispanic										
Both sexes										
2002	597	711	880	964	748	457	249	72	12	-
2001	578	764	840	899	649	477	212	88	6	-
2000	514	684	908	817	593	412	213	66	10	-
1999	559	721	830	782	587	459	179	60	6	-
Male										
2002	315	327	425	442	306	162	71	16	5	-
2001	293	374	384	385	253	146	56	19	1	-
2000	279	338	424	362	247	146	60	17	-	-
1999	278	331	378	336	226	160	46	11	-	-
Female										
2002	282	384	455	522	442	295	178	56	7	-
2001	285	390	456	514	396	331	156	69	5	-
2000	235	346	484	455	346	266	153	49	10	-
1999	281	390	452	446	361	299	133	49	6	-
Non-Hispanic[1]										
Both sexes										
2002	4,893	6,513	8,557	10,698	10,801	8,320	4,121	1,277	183	1
2001	4,793	6,404	8,662	10,616	10,582	7,869	4,039	1,261	189	1
2000	4,662	6,349	8,682	10,755	9,962	7,557	3,886	1,181	190	-
1999	4,654	6,477	8,841	10,455	9,870	7,414	3,752	1,101	169	2
Male										
2002	2,738	3,400	4,239	4,941	4,570	3,075	1,183	295	23	1
2001	2,579	3,345	4,298	4,844	4,411	2,812	1,131	250	32	1
2000	2,484	3,206	4,230	4,906	4,192	2,634	1,103	242	20	-
1999	2,447	3,288	4,260	4,842	4,113	2,648	1,021	197	28	1
Female										
2002	2,155	3,113	4,318	5,757	6,231	5,245	2,938	982	160	-
2001	2,214	3,059	4,364	5,772	6,171	5,057	2,908	1,011	157	-
2000	2,178	3,143	4,452	5,849	5,770	4,923	2,783	939	170	-
1999	2,207	3,189	4,581	5,613	5,757	4,766	2,731	904	141	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diabetes mellitus (E10-E14)														
Non-Hispanic White														
Both sexes														
2002	52,463	1	3	4	15	32	63	120	262	474	871	1,289	2,096	2,896
2001	51,482	-	5	3	16	20	46	105	230	442	791	1,250	2,082	2,664
2000	50,163	2	6	4	9	28	64	107	250	441	819	1,175	1,903	2,526
1999	49,347	2	6	8	11	36	58	122	241	462	769	1,197	1,743	2,382
Male														
2002	25,273	-	1	4	12	16	35	69	162	284	565	792	1,262	1,767
2001	24,302	-	1	1	12	14	28	54	130	262	505	761	1,253	1,585
2000	23,465	2	2	3	4	18	39	60	160	253	513	715	1,127	1,417
1999	23,177	-	3	4	5	21	31	63	140	289	458	760	1,041	1,379
Female														
2002	27,190	1	2	-	3	16	28	51	100	190	306	497	834	1,129
2001	27,180	-	4	2	4	6	18	51	100	180	286	489	829	1,079
2000	26,698	-	4	1	5	10	25	47	90	188	306	460	776	1,109
1999	26,170	2	3	4	6	15	27	59	101	173	311	437	702	1,003
Non-Hispanic Black														
Both sexes														
2002	12,583	-	-	1	13	16	41	73	121	208	346	543	850	1,003
2001	12,167	1	-	4	11	17	39	83	115	202	304	507	799	974
2000	11,918	1	-	3	6	14	41	75	112	176	282	513	757	889
1999	11,830	-	2	1	8	12	31	48	112	192	310	513	706	948
Male														
2002	5,158	-	-	-	5	9	25	37	75	114	202	299	445	512
2001	4,983	1	-	1	5	7	20	49	73	119	156	285	416	483
2000	4,720	1	-	2	2	8	20	40	58	99	153	277	371	434
1999	4,718	-	1	-	5	4	15	21	64	107	169	290	351	457
Female														
2002	7,425	-	-	1	8	7	16	36	46	94	144	244	405	491
2001	7,184	-	-	3	6	10	19	34	42	83	148	222	383	491
2000	7,198	-	-	1	4	6	21	35	54	77	129	236	386	455
1999	7,112	-	1	1	3	8	16	27	48	85	141	223	355	491
Origin not stated[2]														
Both sexes														
2002	225	-	-	-	-	-	1	1	1	6	7	13	9	13
2001	211	-	-	-	-	-	1	2	2	1	3	9	17	15
2000	204	-	-	-	-	-	-	-	2	4	6	2	12	11
1999	200	-	-	-	-	1	-	2	3	2	2	4	13	13
Male														
2002	123	-	-	-	-	-	1	1	1	4	6	9	5	6
2001	107	-	-	-	-	-	-	-	2	1	2	7	8	9
2000	101	-	-	-	-	-	-	-	-	4	4	1	6	8

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)										
Non-Hispanic White										
Both sexes										
2002	3,507	4,717	6,620	8,632	9,060	7,066	3,536	1,064	135	-
2001	3,454	4,736	6,728	8,585	8,922	6,742	3,467	1,057	136	1
2000	3,397	4,714	6,761	8,725	8,315	6,472	3,309	998	138	-
1999	3,292	4,820	6,844	8,523	8,298	6,280	3,190	934	129	-
Male										
2002	2,048	2,595	3,446	4,179	3,972	2,720	1,071	255	18	-
2001	1,939	2,607	3,535	4,063	3,811	2,515	990	210	25	1
2000	1,881	2,510	3,457	4,135	3,661	2,320	962	210	16	-
1999	1,821	2,568	3,498	4,116	3,606	2,310	876	166	22	-
Female										
2002	1,459	2,122	3,174	4,453	5,088	4,346	2,465	809	117	-
2001	1,515	2,129	3,193	4,522	5,111	4,227	2,477	847	111	-
2000	1,516	2,204	3,304	4,590	4,654	4,152	2,347	788	122	-
1999	1,471	2,252	3,346	4,407	4,692	3,970	2,314	768	107	-
Non-Hispanic Black										
Both sexes										
2002	1,201	1,513	1,639	1,728	1,485	1,045	516	195	45	1
2001	1,159	1,440	1,670	1,735	1,397	975	507	179	49	-
2000	1,106	1,420	1,663	1,769	1,424	953	505	160	49	-
1999	1,188	1,438	1,682	1,635	1,360	970	494	141	37	2
Male										
2002	592	658	662	634	501	259	90	33	5	1
2001	548	629	649	655	495	236	120	29	7	-
2000	518	583	665	652	448	258	108	21	2	-
1999	529	630	632	588	428	278	121	22	5	1
Female										
2002	609	855	977	1,094	984	786	426	162	40	-
2001	611	811	1,021	1,080	902	739	387	150	42	-
2000	588	837	998	1,117	976	695	397	139	47	-
1999	659	808	1,050	1,047	932	692	373	119	32	1
Origin not stated[2]										
Both sexes										
2002	22	25	23	41	30	20	5	8	-	-
2001	11	27	34	31	28	20	8	1	1	-
2000	18	21	30	35	22	33	5	3	-	-
1999	19	17	22	32	31	20	12	4	2	1
Male										
2002	14	15	11	19	17	11	-	3	-	-
2001	7	15	21	15	10	8	1	-	1	-
2000	7	12	16	20	11	9	2	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diabetes mellitus (E10-E14)														
Origin not stated[2]														
Male														
1999	96	-	-	-	-	-	-	1	2	-	2	2	12	7
Female														
2002	102	-	-	-	-	-	-	-	-	2	1	4	4	7
2001	104	-	-	-	-	-	1	2	-	-	1	2	9	6
2000	103	-	-	-	-	-	-	-	2	-	2	1	6	3
1999	104	-	-	-	-	1	-	1	1	2	-	2	1	6
Nutritional deficiencies (E40-E64)														
All origins														
Both sexes														
2002	3,779	4	1	-	1	1	6	5	11	27	33	39	63	77
2001	3,704	4	4	-	2	1	9	5	9	16	28	48	62	64
2000	4,242	14	2	1	4	4	6	9	9	30	26	46	53	70
1999	4,289	9	2	1	-	2	6	4	8	20	44	45	61	76
Male														
2002	1,321	3	1	-	-	-	4	3	4	12	19	20	28	43
2001	1,233	4	1	-	2	-	7	2	2	8	13	27	29	35
2000	1,412	10	2	-	2	1	1	5	4	9	7	25	21	45
1999	1,486	4	2	1	-	2	3	1	3	8	20	22	30	42
Female														
2002	2,458	1	-	-	1	1	2	2	7	15	14	19	35	34
2001	2,471	-	3	-	-	1	2	3	7	8	15	21	33	29
2000	2,830	4	-	1	2	3	5	4	5	21	19	21	32	25
1999	2,803	5	-	-	-	-	3	3	5	12	24	23	31	34
Hispanic														
Both sexes														
2002	141	2	-	-	-	-	3	1	-	1	2	2	3	1
2001	136	-	1	-	1	-	-	-	-	4	3	1	2	2
2000	147	-	-	-	-	-	1	2	1	4	-	2	3	3
1999	138	-	-	-	-	-	-	-	-	1	3	1	5	5
Male														
2002	61	1	-	-	-	-	1	1	-	1	2	2	1	1
2001	53	-	-	-	1	-	-	-	-	1	1	-	1	2
2000	57	-	-	-	-	-	-	2	-	1	-	1	2	3
1999	54	-	-	-	-	-	-	-	-	1	1	1	4	1
Female														
2002	80	1	-	-	-	-	2	-	-	-	-	-	2	-
2001	83	-	1	-	-	-	-	-	-	3	2	1	1	-
2000	90	-	-	-	-	-	1	-	1	3	-	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diabetes mellitus (E10-E14)										
Origin not stated[2]										
Male										
1999	11	11	11	12	10	8	4	2	-	1
Female										
2002	8	10	12	22	13	9	5	5	-	-
2001	4	12	13	16	18	12	7	1	-	-
2000	11	9	14	15	11	24	3	2	-	-
1999	8	6	11	20	21	12	8	2	2	-
Nutritional deficiencies (E40-E64)										
All origins										
Both sexes										
2002	91	150	232	452	680	834	667	312	93	-
2001	89	131	243	439	661	798	684	324	83	-
2000	103	192	268	505	780	947	781	302	90	-
1999	94	149	280	472	776	944	822	380	94	-
Male										
2002	50	70	116	193	241	283	169	49	13	-
2001	46	63	111	172	254	228	171	55	3	-
2000	52	87	134	207	264	289	182	52	13	-
1999	56	72	149	205	289	301	196	71	9	-
Female										
2002	41	80	116	259	439	551	498	263	80	-
2001	43	68	132	267	407	570	513	269	80	-
2000	51	105	134	298	516	658	599	250	77	-
1999	38	77	131	267	487	643	626	309	85	-
Hispanic										
Both sexes										
2002	3	7	14	25	24	22	19	8	4	-
2001	3	5	7	22	23	25	25	12	-	-
2000	4	14	10	18	25	20	29	8	3	-
1999	2	4	13	16	15	34	20	15	4	-
Male										
2002	1	5	8	9	10	8	7	3	-	-
2001	2	1	1	11	9	6	10	7	-	-
2000	-	4	9	8	9	7	9	1	1	-
1999	1	1	7	10	3	10	7	6	1	-
Female										
2002	2	2	6	16	14	14	12	5	4	-
2001	1	4	6	11	14	19	15	5	-	-
2000	4	10	1	10	16	13	20	7	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nutritional deficiencies (E40-E64)														
Hispanic														
Female														
1999	84	-	-	-	-	-	-	-	-	-	2	-	1	4
Non-Hispanic[1]														
Both sexes														
2002	3,621	2	1	-	1	1	3	4	11	25	31	37	60	76
2001	3,563	4	3	-	1	1	8	5	9	12	25	47	60	62
2000	4,080	14	2	1	4	4	5	7	8	26	26	44	50	67
1999	4,142	9	2	1	-	2	6	4	8	19	41	44	56	70
Male														
2002	1,255	2	1	-	-	-	3	2	4	11	17	18	27	42
2001	1,178	4	1	-	1	-	6	2	2	7	12	27	28	33
2000	1,348	10	2	-	2	1	1	3	4	8	7	24	19	42
1999	1,428	4	2	1	-	2	3	1	3	7	19	21	26	40
Female														
2002	2,366	-	-	-	1	1	-	2	7	14	14	19	33	34
2001	2,385	-	2	-	-	1	2	3	7	5	13	20	32	29
2000	2,732	4	-	1	2	3	4	4	4	18	19	20	31	25
1999	2,714	5	-	-	-	-	3	3	5	12	22	23	30	30
Non-Hispanic White														
Both sexes														
2002	3,064	1	1	-	-	1	2	4	8	13	24	30	46	53
2001	3,015	3	3	-	-	1	6	5	5	7	15	33	37	43
2000	3,469	8	1	-	2	4	3	3	3	17	19	35	33	49
1999	3,486	5	1	-	-	2	3	3	6	14	31	27	43	40
Male														
2002	1,039	1	1	-	-	-	2	2	3	6	13	12	19	26
2001	948	3	1	-	-	-	6	2	1	4	6	14	13	20
2000	1,112	6	1	-	2	1	1	1	1	6	4	19	11	27
1999	1,142	2	1	-	-	2	1	-	3	4	14	15	19	19
Female														
2002	2,025	-	-	-	-	1	-	2	5	7	11	18	27	27
2001	2,067	-	2	-	-	1	-	3	4	3	9	19	24	23
2000	2,357	2	-	-	-	3	2	2	2	11	15	16	22	22
1999	2,344	3	-	-	-	-	2	3	3	10	17	12	24	21
Non-Hispanic Black														
Both sexes														
2002	495	-	-	-	1	-	1	-	3	9	7	7	13	23
2001	472	1	-	-	1	-	2	-	4	5	10	12	21	17
2000	537	5	1	1	2	-	2	4	4	9	7	8	16	18
1999	585	4	1	1	-	-	3	1	2	5	10	15	12	26

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional deficiencies (E40-E64)										
Hispanic										
Female										
1999	1	3	6	6	12	24	13	9	3	-
Non-Hispanic[1]										
Both sexes										
2002	88	143	215	422	655	809	645	303	89	-
2001	86	126	236	416	638	772	657	312	83	-
2000	99	178	257	482	751	926	749	293	87	-
1999	92	145	266	455	760	907	800	365	90	-
Male										
2002	49	65	108	181	231	274	162	45	13	-
2001	44	62	110	160	245	222	161	48	3	-
2000	52	83	125	195	253	281	173	51	12	-
1999	55	71	142	194	285	290	189	65	8	-
Female										
2002	39	78	107	241	424	535	483	258	76	-
2001	42	64	126	256	393	550	496	264	80	-
2000	47	95	132	287	498	645	576	242	75	-
1999	37	74	124	261	475	617	611	300	82	-
Non-Hispanic White										
Both sexes										
2002	64	107	173	357	565	703	572	267	73	-
2001	70	93	198	342	536	690	578	274	76	-
2000	74	145	200	404	658	815	681	247	68	-
1999	70	111	196	375	643	797	731	312	76	-
Male										
2002	31	50	87	157	191	242	147	39	10	-
2001	32	42	89	133	207	197	135	41	2	-
2000	35	66	93	160	222	248	158	40	10	-
1999	42	51	99	154	238	254	169	49	6	-
Female										
2002	33	57	86	200	374	461	425	228	63	-
2001	38	51	109	209	329	493	443	233	74	-
2000	39	79	107	244	436	567	523	207	58	-
1999	28	60	97	221	405	543	562	263	70	-
Non-Hispanic Black										
Both sexes										
2002	23	33	37	61	78	88	66	32	13	-
2001	12	28	33	62	87	72	68	30	7	-
2000	23	29	49	67	80	96	60	40	16	-
1999	21	29	61	69	106	96	63	46	14	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nutritional deficiencies (E40-E64)														
Non-Hispanic Black														
Male														
2002	185	-	-	-	-	-	1	-	1	3	4	6	8	16
2001	202	1	-	-	1	-	-	-	1	3	6	11	14	12
2000	214	3	1	-	-	-	-	2	3	2	3	5	8	15
1999	250	2	1	1	-	-	2	1	-	3	5	5	7	18
Female														
2002	310	-	-	-	1	-	-	-	2	6	3	1	5	7
2001	270	-	-	-	-	-	2	-	3	2	4	1	7	5
2000	323	2	-	1	2	-	2	2	1	7	4	3	8	3
1999	335	2	-	-	-	-	1	-	2	2	5	10	5	8
Origin not stated[2]														
Both sexes														
2002	17	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	5	-	-	-	-	-	1	-	-	-	-	-	-	-
2000	15	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	9	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	5	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2000	7	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	12	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	8	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Malnutrition (E40-E46)														
All origins														
Both sexes														
2002	3,510	2	1	-	1	1	5	4	9	25	30	36	55	73
2001	3,454	2	-	-	2	1	8	4	9	16	23	42	57	62
2000	3,926	11	1	1	2	3	5	9	7	29	24	44	46	61
1999	3,985	7	2	1	-	2	5	4	7	20	41	42	57	66
Male														
2002	1,233	1	1	-	-	-	4	2	3	11	17	18	24	39
2001	1,148	2	-	-	2	-	6	2	2	8	11	22	25	33
2000	1,302	8	1	-	1	1	1	5	3	9	6	23	18	39
1999	1,371	4	2	1	-	2	2	1	3	8	19	21	27	38

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nutritional deficiencies (E40-E64)										
Non-Hispanic Black										
Male										
2002	17	14	17	21	35	26	12	2	2	-
2001	9	18	20	25	30	24	21	5	1	-
2000	15	15	29	31	27	31	12	11	1	-
1999	12	17	38	34	40	30	19	13	2	-
Female										
2002	6	19	20	40	43	62	54	30	11	-
2001	3	10	13	37	57	48	47	25	6	-
2000	8	14	20	36	53	65	48	29	15	-
1999	9	12	23	35	66	66	44	33	12	-
Origin not stated[2]										
Both sexes										
2002	-	-	3	5	1	3	3	1	-	-
2001	-	-	-	1	-	1	2	-	-	-
2000	-	-	1	5	4	1	3	1	-	-
1999	-	-	1	1	1	3	2	-	-	-
Male										
2002	-	-	-	3	-	1	-	1	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	4	2	1	-	-	-	-
1999	-	-	-	1	1	1	-	-	-	-
Female										
2002	-	-	3	2	1	2	3	-	-	-
2001	-	-	-	-	-	1	2	-	-	-
2000	-	-	1	1	2	-	3	1	-	-
1999	-	-	1	-	-	2	2	-	-	-
Malnutrition (E40-E46)										
All origins										
Both sexes										
2002	83	136	218	419	648	778	616	283	87	-
2001	84	123	224	406	619	756	643	302	71	-
2000	97	178	245	478	721	888	712	280	84	-
1999	88	141	266	439	720	883	755	353	86	-
Male										
2002	45	64	110	176	230	267	163	45	13	-
2001	42	57	100	156	244	220	164	50	2	-
2000	47	79	128	194	240	275	165	46	13	-
1999	54	68	139	186	266	282	175	65	8	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malnutrition (E40-E46)														
All origins														
Female														
2002	2,277	1	-	-	1	1	1	2	6	14	13	18	31	34
2001	2,306	-	-	-	-	1	2	2	7	8	12	20	32	29
2000	2,624	3	-	1	1	2	4	4	4	20	18	21	28	22
1999	2,614	3	-	-	-	-	3	3	4	12	22	21	30	28
Hispanic														
Both sexes														
2002	136	1	-	-	-	-	2	1	-	1	2	1	3	1
2001	132	-	-	-	1	-	-	-	-	4	2	1	2	2
2000	137	-	-	-	-	-	1	2	-	4	-	2	2	3
1999	132	-	-	-	-	-	-	-	-	1	3	1	5	5
Male														
2002	58	-	-	-	-	-	1	1	-	1	2	1	1	1
2001	52	-	-	-	1	-	-	-	-	1	1	-	1	2
2000	52	-	-	-	-	-	-	2	-	1	-	1	1	3
1999	50	-	-	-	-	-	-	-	-	1	1	1	4	1
Female														
2002	78	1	-	-	-	-	1	-	-	-	-	-	2	-
2001	80	-	-	-	-	-	-	-	-	3	1	1	1	-
2000	85	-	-	-	-	-	1	-	-	3	-	1	1	-
1999	82	-	-	-	-	-	-	-	-	-	2	-	1	4
Non-Hispanic[1]														
Both sexes														
2002	3,358	1	1	-	1	1	3	3	9	23	28	35	52	72
2001	3,317	2	-	-	1	1	7	4	9	12	21	41	55	60
2000	3,774	11	1	1	2	3	4	7	7	25	24	42	44	58
1999	3,844	7	2	1	-	2	5	4	7	19	38	41	52	60
Male														
2002	1,171	1	1	-	-	-	3	1	3	10	15	17	23	38
2001	1,094	2	-	-	1	-	5	2	2	7	10	22	24	31
2000	1,243	8	1	-	1	1	1	3	3	8	6	22	17	36
1999	1,317	4	2	1	-	2	2	1	3	7	18	20	23	36
Female														
2002	2,187	-	-	-	1	1	-	2	6	13	13	18	29	34
2001	2,223	-	-	-	-	1	2	2	7	5	11	19	31	29
2000	2,531	3	-	1	1	2	3	4	4	17	18	20	27	22
1999	2,527	3	-	-	-	-	3	3	4	12	20	21	29	24
Non-Hispanic White														
Both sexes														
2002	2,836	1	1	-	-	1	2	3	6	13	21	28	39	50

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)										
All origins										
Female										
2002	38	72	108	243	418	511	453	238	74	-
2001	42	66	124	250	375	536	479	252	69	-
2000	50	99	117	284	481	613	547	234	71	-
1999	34	73	127	253	454	601	580	288	78	-
Hispanic										
Both sexes										
2002	3	7	14	25	24	21	19	8	3	-
2001	3	5	7	22	22	25	25	11	-	-
2000	4	12	10	16	23	20	27	8	3	-
1999	2	4	13	15	15	32	17	15	4	-
Male										
2002	1	5	8	9	10	7	7	3	-	-
2001	2	1	1	11	9	6	10	6	-	-
2000	-	3	9	7	9	7	7	1	1	-
1999	1	1	7	9	3	9	5	6	1	-
Female										
2002	2	2	6	16	14	14	12	5	3	-
2001	1	4	6	11	13	19	15	5	-	-
2000	4	9	1	9	14	13	20	7	2	-
1999	1	3	6	6	12	23	12	9	3	-
Non-Hispanic[1]										
Both sexes										
2002	80	129	201	390	623	754	594	274	84	-
2001	81	118	217	383	597	730	616	291	71	-
2000	93	166	234	457	694	867	682	271	81	-
1999	86	137	252	423	704	848	736	338	82	-
Male										
2002	44	59	102	165	220	259	156	41	13	-
2001	40	56	99	144	235	214	154	44	2	-
2000	47	76	119	183	229	267	158	45	12	-
1999	53	67	132	176	262	272	170	59	7	-
Female										
2002	36	70	99	225	403	495	438	233	71	-
2001	41	62	118	239	362	516	462	247	69	-
2000	46	90	115	274	465	600	524	226	69	-
1999	33	70	120	247	442	576	566	279	75	-
Non-Hispanic White										
Both sexes										
2002	58	98	164	327	537	653	525	240	69	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malnutrition (E40-E46)														
Non-Hispanic White														
Both sexes														
2001	2,806	1	-	-	-	1	5	4	5	7	12	29	35	43
2000	3,204	7	-	-	1	3	3	3	3	16	17	33	29	42
1999	3,232	4	1	-	-	2	3	3	5	14	29	26	41	35
Male														
2002	970	1	1	-	-	-	2	1	2	6	11	11	16	23
2001	881	1	-	-	-	-	5	2	1	4	4	11	12	20
2000	1,021	5	-	-	1	1	1	1	1	6	3	17	10	23
1999	1,047	2	1	-	-	2	1	-	3	4	13	14	17	18
Female														
2002	1,866	-	-	-	-	1	-	2	4	7	10	17	23	27
2001	1,925	-	-	-	-	1	-	2	4	3	8	18	23	23
2000	2,183	2	-	-	-	2	2	2	2	10	14	16	19	19
1999	2,185	2	-	-	-	-	2	3	2	10	16	12	24	17
Non-Hispanic Black														
Both sexes														
2002	465	-	-	-	1	-	1	-	3	7	7	7	12	22
2001	438	1	-	-	1	-	2	-	4	5	9	10	18	15
2000	505	3	1	1	1	-	1	4	3	9	7	8	15	16
1999	543	3	1	1	-	-	2	1	2	5	9	13	11	21
Male														
2002	172	-	-	-	-	-	1	-	1	2	4	6	7	15
2001	186	1	-	-	1	-	-	-	1	3	6	9	11	10
2000	202	2	1	-	-	-	-	2	2	2	3	5	7	13
1999	234	2	1	1	-	-	1	1	-	3	5	5	6	15
Female														
2002	293	-	-	-	1	-	-	-	2	5	3	1	5	7
2001	252	-	-	-	-	-	2	-	3	2	3	1	7	5
2000	303	1	-	1	1	-	1	2	1	7	4	3	8	3
1999	309	1	-	-	-	-	1	-	2	2	4	8	5	6
Origin not stated[2]														
Both sexes														
2002	16	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	5	-	-	-	-	-	1	-	-	-	-	-	-	-
2000	15	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	9	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2000	7	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)										
Non-Hispanic White										
Both sexes										
2001	66	88	182	315	501	653	540	255	64	-
2000	70	134	178	386	607	760	621	227	64	-
1999	64	103	184	350	595	745	672	288	68	-
Male										
2002	28	46	83	142	183	228	141	35	10	-
2001	29	39	80	118	198	189	129	38	1	-
2000	32	60	87	152	198	234	145	34	10	-
1999	40	47	91	142	217	237	150	43	5	-
Female										
2002	30	52	81	185	354	425	384	205	59	-
2001	37	49	102	197	303	464	411	217	63	-
2000	38	74	91	234	409	526	476	193	54	-
1999	24	56	93	208	378	508	522	245	63	-
Non-Hispanic Black										
Both sexes										
2002	22	28	32	59	75	84	63	30	12	-
2001	11	26	30	58	81	67	65	28	7	-
2000	21	28	49	62	74	94	56	38	14	-
1999	21	29	59	62	99	89	58	43	14	-
Male										
2002	16	12	15	20	32	25	12	2	2	-
2001	8	16	18	24	29	24	20	4	1	-
2000	13	14	29	29	27	31	10	11	1	-
1999	12	17	36	28	38	29	19	13	2	-
Female										
2002	6	16	17	39	43	59	51	28	10	-
2001	3	10	12	34	52	43	45	24	6	-
2000	8	14	20	33	47	63	46	27	13	-
1999	9	12	23	34	61	60	39	30	12	-
Origin not stated[2]										
Both sexes										
2002	-	-	3	4	1	3	3	1	-	-
2001	-	-	-	1	-	1	2	-	-	-
2000	-	-	1	5	4	1	3	1	-	-
1999	-	-	1	1	1	3	2	-	-	-
Male										
2002	-	-	-	2	-	1	-	1	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	4	2	1	-	-	-	-
1999	-	-	-	1	1	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Malnutrition (E40-E46)														
Origin not stated[2]														
Female														
2002	12	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	8	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Other nutritional deficiencies (E50-E64)														
All origins														
Both sexes														
2002	269	2	-	-	-	-	1	1	2	2	3	3	8	4
2001	250	2	4	-	-	-	1	1	-	-	5	6	5	2
2000	316	3	1	-	2	1	1	-	2	1	2	2	7	9
1999	304	2	-	-	-	-	1	-	1	-	3	3	4	10
Male														
2002	88	2	-	-	-	-	-	1	1	1	2	2	4	4
2001	85	2	1	-	-	-	1	-	-	-	2	5	4	2
2000	110	2	1	-	1	-	-	-	1	-	1	2	3	6
1999	115	-	-	-	-	-	1	-	-	-	1	1	3	4
Female														
2002	181	-	-	-	-	-	1	-	1	1	1	1	4	-
2001	165	-	3	-	-	-	-	1	-	-	3	1	1	-
2000	206	1	-	-	1	1	1	-	1	1	1	-	4	3
1999	189	2	-	-	-	-	-	-	1	-	2	2	1	6
Hispanic														
Both sexes														
2002	5	1	-	-	-	-	1	-	-	-	-	1	-	-
2001	4	-	1	-	-	-	-	-	-	-	1	-	-	-
2000	10	-	-	-	-	-	-	-	1	-	-	-	1	-
1999	6	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	3	1	-	-	-	-	-	-	-	-	-	1	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	5	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	3	-	1	-	-	-	-	-	-	-	1	-	-	-
2000	5	-	-	-	-	-	-	-	1	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Malnutrition (E40-E46)										
Origin not stated[2]										
Female										
2002	-	-	3	2	1	2	3	-	-	-
2001	-	-	-	-	-	1	2	-	-	-
2000	-	-	1	1	2	-	3	1	-	-
1999	-	-	1	-	-	2	2	-	-	-
Other nutritional deficiencies (E50-E64)										
All origins										
Both sexes										
2002	8	14	14	33	32	56	51	29	6	-
2001	5	8	19	33	42	42	41	22	12	-
2000	6	14	23	27	59	59	69	22	6	-
1999	6	8	14	33	56	61	67	27	8	-
Male										
2002	5	6	6	17	11	16	6	4	-	-
2001	4	6	11	16	10	8	7	5	1	-
2000	5	8	6	13	24	14	17	6	-	-
1999	2	4	10	19	23	19	21	6	1	-
Female										
2002	3	8	8	16	21	40	45	25	6	-
2001	1	2	8	17	32	34	34	17	11	-
2000	1	6	17	14	35	45	52	16	6	-
1999	4	4	4	14	33	42	46	21	7	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	1	-	-	1	-
2001	-	-	-	-	1	-	-	1	-	-
2000	-	2	-	2	2	-	2	-	-	-
1999	-	-	-	1	-	2	3	-	-	-
Male										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	-	-	-	1	-	-
2000	-	1	-	1	-	-	2	-	-	-
1999	-	-	-	1	-	1	2	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	1	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	1	-	1	2	-	-	-	-	-
1999	-	-	-	-	-	1	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other nutritional deficiencies (E50-E64)														
Non-Hispanic[1]														
Both sexes														
2002	263	1	-	-	-	-	-	1	2	2	3	2	8	4
2001	246	2	3	-	-	-	1	1	-	-	4	6	5	2
2000	306	3	1	-	2	1	1	-	1	1	2	2	6	9
1999	298	2	-	-	-	-	1	-	1	-	3	3	4	10
Male														
2002	84	1	-	-	-	-	-	1	1	1	2	1	4	4
2001	84	2	1	-	-	-	1	-	-	-	2	5	4	2
2000	105	2	1	-	1	-	-	-	1	-	1	2	2	6
1999	111	-	-	-	-	-	1	-	-	-	1	1	3	4
Female														
2002	179	-	-	-	-	-	-	-	1	1	1	1	4	-
2001	162	-	2	-	-	-	-	1	-	-	2	1	1	-
2000	201	1	-	-	1	1	1	-	-	1	1	-	4	3
1999	187	2	-	-	-	-	-	-	1	-	2	2	1	6
Non-Hispanic White														
Both sexes														
2002	228	-	-	-	-	-	-	1	2	-	3	2	7	3
2001	209	2	3	-	-	-	1	1	-	-	3	4	2	-
2000	265	1	1	-	1	1	-	-	-	1	2	2	4	7
1999	254	1	-	-	-	-	-	-	1	-	2	1	2	5
Male														
2002	69	-	-	-	-	-	-	1	1	-	2	1	3	3
2001	67	2	1	-	-	-	1	-	-	-	2	3	1	-
2000	91	1	1	-	1	-	-	-	-	-	1	2	1	4
1999	95	-	-	-	-	-	-	-	-	-	1	1	2	1
Female														
2002	159	-	-	-	-	-	-	-	1	-	1	1	4	-
2001	142	-	2	-	-	-	-	1	-	-	1	1	1	-
2000	174	-	-	-	-	1	-	-	-	1	1	-	3	3
1999	159	1	-	-	-	-	-	-	1	-	1	-	-	4
Non-Hispanic Black														
Both sexes														
2002	30	-	-	-	-	-	-	-	-	2	-	-	1	1
2001	34	-	-	-	-	-	-	-	-	-	1	2	3	2
2000	32	2	-	-	1	-	1	-	1	-	-	-	1	2
1999	42	1	-	-	-	-	1	-	-	-	1	2	1	5
Male														
2002	13	-	-	-	-	-	-	-	-	1	-	-	1	1
2001	16	-	-	-	-	-	-	-	-	-	-	2	3	2
2000	12	1	-	-	-	-	-	-	1	-	-	-	1	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other nutritional deficiencies (E50-E64)										
Non-Hispanic[1]										
Both sexes										
2002	8	14	14	32	32	55	51	29	5	-
2001	5	8	19	33	41	42	41	21	12	-
2000	6	12	23	25	57	59	67	22	6	-
1999	6	8	14	32	56	59	64	27	8	-
Male										
2002	5	6	6	16	11	15	6	4	-	-
2001	4	6	11	16	10	8	7	4	1	-
2000	5	7	6	12	24	14	15	6	-	-
1999	2	4	10	18	23	18	19	6	1	-
Female										
2002	3	8	8	16	21	40	45	25	5	-
2001	1	2	8	17	31	34	34	17	11	-
2000	1	5	17	13	33	45	52	16	6	-
1999	4	4	4	14	33	41	45	21	7	-
Non-Hispanic White										
Both sexes										
2002	6	9	9	30	28	50	47	27	4	-
2001	4	5	16	27	35	37	38	19	12	-
2000	4	11	22	18	51	55	60	20	4	-
1999	6	8	12	25	48	52	59	24	8	-
Male										
2002	3	4	4	15	8	14	6	4	-	-
2001	3	3	9	15	9	8	6	3	1	-
2000	3	6	6	8	24	14	13	6	-	-
1999	2	4	8	12	21	17	19	6	1	-
Female										
2002	3	5	5	15	20	36	41	23	4	-
2001	1	2	7	12	26	29	32	16	11	-
2000	1	5	16	10	27	41	47	14	4	-
1999	4	4	4	13	27	35	40	18	7	-
Non-Hispanic Black										
Both sexes										
2002	1	5	5	2	3	4	3	2	1	-
2001	1	2	3	4	6	5	3	2	-	-
2000	2	1	-	5	6	2	4	2	2	-
1999	-	-	2	7	7	7	5	3	-	-
Male										
2002	1	2	2	1	3	1	-	-	-	-
2001	1	2	2	1	1	-	1	1	-	-
2000	2	1	-	2	-	-	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other nutritional deficiencies (E50-E64)														
Non-Hispanic Black														
Male														
1999	16	-	-	-	-	-	1	-	-	-	-	-	1	3
Female														
2002	17	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	18	-	-	-	-	-	-	-	-	-	1	-	-	-
2000	20	1	-	-	1	-	1	-	-	-	-	-	-	-
1999	26	1	-	-	-	-	-	-	-	-	1	2	-	2
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Meningitis (G00,G03)														
All origins														
Both sexes														
2002	700	74	19	10	21	12	14	16	20	23	38	63	53	48
2001	727	78	26	15	9	19	21	13	28	31	38	55	50	58
2000	758	91	36	12	18	7	8	16	27	35	37	45	55	43
1999	850	116	45	14	13	19	22	21	27	33	43	62	46	41
Male														
2002	389	41	13	5	11	8	7	9	10	17	26	41	30	29
2001	394	39	16	9	6	12	11	11	14	23	19	30	30	30
2000	405	48	20	6	10	2	2	7	13	22	24	25	31	26
1999	442	61	26	9	9	10	16	8	16	18	27	39	29	24
Female														
2002	311	33	6	5	10	4	7	7	10	6	12	22	23	19
2001	333	39	10	6	3	7	10	2	14	8	19	25	20	28
2000	353	43	16	6	8	5	6	9	14	13	13	20	24	17

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other nutritional deficiencies (E50-E64)										
Non-Hispanic Black										
Male										
1999	-	-	2	6	2	1	-	-	-	-
Female										
2002	-	3	3	1	-	3	3	2	1	-
2001	-	-	1	3	5	5	2	1	-	-
2000	-	-	-	3	6	2	2	2	2	-
1999	-	-	-	1	5	6	5	3	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Meningitis (G00,G03)										
All origins										
Both sexes										
2002	40	46	63	47	37	36	16	3	-	1
2001	55	46	48	46	50	25	12	4	-	-
2000	47	48	60	68	55	32	16	2	-	-
1999	49	49	66	65	57	40	20	2	-	-
Male										
2002	19	26	36	21	20	11	8	-	-	1
2001	31	24	27	26	21	11	3	1	-	-
2000	28	26	34	41	21	13	5	1	-	-
1999	21	28	32	26	23	13	7	-	-	-
Female										
2002	21	20	27	26	17	25	8	3	-	-
2001	24	22	21	20	29	14	9	3	-	-
2000	19	22	26	27	34	19	11	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningitis (G00,G03)														
All origins														
Female														
1999	408	55	19	5	4	9	6	13	11	15	16	23	17	17
Hispanic														
Both sexes														
2002	70	8	5	-	-	3	3	4	1	5	4	5	3	6
2001	58	8	4	2	1	-	3	-	4	4	3	5	1	4
2000	52	9	4	2	1	-	3	-	5	5	3	1	5	2
1999	76	17	7	1	-	3	2	2	4	2	2	8	4	3
Male														
2002	42	6	4	-	-	3	2	3	-	2	3	3	2	4
2001	33	5	3	1	1	-	1	-	1	4	2	3	-	2
2000	32	6	2	1	-	-	2	-	3	3	2	1	4	1
1999	41	13	5	1	-	-	1	1	2	1	2	4	2	3
Female														
2002	28	2	1	-	-	-	1	1	1	3	1	2	1	2
2001	25	3	1	1	-	-	2	-	3	-	1	2	1	2
2000	20	3	2	1	1	-	1	-	2	2	1	-	1	1
1999	35	4	2	-	-	3	1	1	2	1	-	4	2	-
Non-Hispanic[1]														
Both sexes														
2002	625	66	14	10	21	9	11	12	19	17	34	57	50	41
2001	666	70	22	13	8	19	18	13	24	27	35	49	49	54
2000	705	82	32	10	17	7	5	16	21	30	34	44	50	41
1999	767	97	38	12	13	16	18	19	23	31	40	54	42	37
Male														
2002	344	35	9	5	11	5	5	6	10	14	23	38	28	24
2001	361	34	13	8	5	12	10	11	13	19	17	27	30	28
2000	372	42	18	5	10	2	-	7	9	19	22	24	27	25
1999	396	48	21	7	9	10	13	7	14	17	24	35	27	20
Female														
2002	281	31	5	5	10	4	6	6	9	3	11	19	22	17
2001	305	36	9	5	3	7	8	2	11	8	18	22	19	26
2000	333	40	14	5	7	5	5	9	12	11	12	20	23	16
1999	371	49	17	5	4	6	5	12	9	14	16	19	15	17
Non-Hispanic White														
Both sexes														
2002	427	44	10	9	11	7	5	5	10	13	17	33	29	26
2001	469	39	15	8	7	14	10	8	17	11	22	32	34	36
2000	486	47	19	9	9	4	3	8	12	19	17	34	31	27
1999	524	51	26	8	10	7	8	14	14	20	25	31	29	28

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningitis (G00,G03)										
All origins										
Female										
1999	28	21	34	39	34	27	13	2	-	-
Hispanic										
Both sexes										
2002	2	4	7	5	4	1	-	-	-	-
2001	3	2	4	3	5	1	1	-	-	-
2000	3	2	3	3	1	-	-	-	-	-
1999	4	6	3	4	-	4	-	-	-	-
Male										
2002	-	1	5	1	2	1	-	-	-	-
2001	1	2	3	1	3	-	-	-	-	-
2000	2	1	-	3	1	-	-	-	-	-
1999	-	2	3	-	-	1	-	-	-	-
Female										
2002	2	3	2	4	2	-	-	-	-	-
2001	2	-	1	2	2	1	1	-	-	-
2000	1	1	3	-	-	-	-	-	-	-
1999	4	4	-	4	-	3	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	38	42	56	41	32	35	16	3	-	1
2001	52	44	44	43	45	22	11	4	-	-
2000	44	46	57	65	54	32	16	2	-	-
1999	45	43	63	61	57	36	20	2	-	-
Male										
2002	19	25	31	19	18	10	8	-	-	1
2001	30	22	24	25	18	11	3	1	-	-
2000	26	25	34	38	20	13	5	1	-	-
1999	21	26	29	26	23	12	7	-	-	-
Female										
2002	19	17	25	22	14	25	8	3	-	-
2001	22	22	20	18	27	11	8	3	-	-
2000	18	21	23	27	34	19	11	1	-	-
1999	24	17	34	35	34	24	13	2	-	-
Non-Hispanic White										
Both sexes										
2002	23	34	43	31	27	34	12	3	-	1
2001	41	33	40	37	37	17	9	2	-	-
2000	33	30	44	53	47	24	14	2	-	-
1999	35	32	48	44	48	32	12	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningitis (G00,G03)														
Non-Hispanic White														
Male														
2002	227	23	6	5	6	4	1	2	4	11	9	21	16	15
2001	243	17	10	4	4	7	5	7	10	6	12	17	21	19
2000	264	27	12	5	4	1	-	3	4	12	10	20	18	19
1999	265	33	15	4	7	5	5	3	7	11	13	18	17	16
Female														
2002	200	21	4	4	5	3	4	3	6	2	8	12	13	11
2001	226	22	5	4	3	7	5	1	7	5	10	15	13	17
2000	222	20	7	4	5	3	3	5	8	7	7	14	13	8
1999	259	18	11	4	3	2	3	11	7	9	12	13	12	12
Non-Hispanic Black														
Both sexes														
2002	175	20	4	1	9	1	5	7	8	4	15	21	19	12
2001	172	29	5	4	-	4	6	5	6	13	13	17	13	16
2000	202	34	13	1	8	3	2	8	9	10	17	9	17	13
1999	217	40	11	4	3	9	8	5	6	10	11	22	13	8
Male														
2002	105	11	3	-	5	1	4	4	5	3	12	15	11	7
2001	104	16	2	3	-	4	4	4	2	12	5	10	8	8
2000	100	15	6	-	6	1	-	4	5	6	12	3	8	6
1999	119	13	6	3	2	5	6	4	5	6	8	17	10	4
Female														
2002	70	9	1	1	4	-	1	3	3	1	3	6	8	5
2001	68	13	3	1	-	-	2	1	4	1	8	7	5	8
2000	102	19	7	1	2	2	2	4	4	4	5	6	9	7
1999	98	27	5	1	1	4	2	1	1	4	3	5	3	4
Origin not stated[2]														
Both sexes														
2002	5	-	-	-	-	-	-	-	-	1	-	1	-	1
2001	3	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-	-
1999	7	2	-	1	-	-	2	-	-	-	1	-	-	1
Male														
2002	3	-	-	-	-	-	-	-	-	1	-	-	-	1
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	1	-	-	-	-	-
1999	5	-	-	1	-	-	2	-	-	-	1	-	-	1
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	1	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningitis (G00,G03)										
Non-Hispanic White										
Male										
2002	12	21	24	15	17	9	5	-	-	1
2001	23	15	22	20	15	7	2	-	-	-
2000	20	18	29	30	18	9	4	1	-	-
1999	17	19	22	17	21	11	4	-	-	-
Female										
2002	11	13	19	16	10	25	7	3	-	-
2001	18	18	18	17	22	10	7	2	-	-
2000	13	12	15	23	29	15	10	1	-	-
1999	18	13	26	27	27	21	8	2	-	-
Non-Hispanic Black										
Both sexes										
2002	11	8	13	9	4	1	3	-	-	-
2001	9	9	4	5	6	4	2	2	-	-
2000	9	15	12	8	5	7	2	-	-	-
1999	10	9	12	16	8	4	8	-	-	-
Male										
2002	5	4	7	4	1	1	2	-	-	-
2001	6	6	2	4	2	4	1	1	-	-
2000	5	6	5	5	2	4	1	-	-	-
1999	4	6	6	8	2	1	3	-	-	-
Female										
2002	6	4	6	5	3	-	1	-	-	-
2001	3	3	2	1	4	-	1	1	-	-
2000	4	9	7	3	3	3	1	-	-	-
1999	6	3	6	8	6	3	5	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	1	1	-	-	-	-	-
2001	-	-	-	-	-	2	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	1	-	-	-	-	-
2001	-	-	-	-	-	2	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Meningitis (G00,G03)														
Origin not stated[2]														
Female														
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	2	-	-	-	-	-	-	-	-	-	-	-	-
Parkinson's disease (G20-G21)														
All origins														
Both sexes														
2002	16,959	-	-	-	-	-	-	-	-	1	5	19	39	95
2001	16,544	-	-	-	-	-	-	1	1	3	12	13	32	100
2000	15,682	-	1	-	-	-	-	2	2	4	3	16	33	82
1999	14,593	1	-	-	-	-	2	1	1	1	4	8	35	73
Male														
2002	9,593	-	-	-	-	-	-	-	-	-	3	14	26	57
2001	9,405	-	-	-	-	-	-	1	-	1	10	8	25	69
2000	8,804	-	-	-	-	-	-	1	2	3	2	10	19	55
1999	8,262	-	-	-	-	-	2	1	1	1	2	5	25	44
Female														
2002	7,366	-	-	-	-	-	-	-	-	1	2	5	13	38
2001	7,139	-	-	-	-	-	-	-	1	2	2	5	7	31
2000	6,878	-	1	-	-	-	-	1	-	1	1	6	14	27
1999	6,331	1	-	-	-	-	-	-	-	-	2	3	10	29
Hispanic														
Both sexes														
2002	491	-	-	-	-	-	-	-	-	-	1	4	2	8
2001	483	-	-	-	-	-	-	1	-	-	-	1	2	5
2000	425	-	-	-	-	-	-	-	-	-	-	3	2	4
1999	389	-	-	-	-	-	-	-	-	-	1	-	2	2
Male														
2002	281	-	-	-	-	-	-	-	-	-	1	3	-	5
2001	274	-	-	-	-	-	-	1	-	-	-	1	1	1
2000	240	-	-	-	-	-	-	-	-	-	-	1	1	3
1999	222	-	-	-	-	-	-	-	-	-	1	-	2	1
Female														
2002	210	-	-	-	-	-	-	-	-	-	-	1	2	3
2001	209	-	-	-	-	-	-	-	-	-	-	-	1	4
2000	185	-	-	-	-	-	-	-	-	-	-	2	1	1
1999	167	-	-	-	-	-	-	-	-	-	-	-	-	1
Non-Hispanic[1]														
Both sexes														
2002	16,435	-	-	-	-	-	-	-	-	1	4	15	37	87

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Meningitis (G00,G03)										
Origin not stated[2]										
Female										
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Parkinson's disease (G20-G21)										
All origins										
Both sexes										
2002	223	626	1,600	3,368	4,771	3,959	1,781	440	32	-
2001	201	554	1,593	3,446	4,679	3,786	1,655	425	43	-
2000	188	626	1,482	3,322	4,330	3,567	1,627	369	27	1
1999	169	539	1,481	3,066	4,044	3,332	1,492	313	31	-
Male										
2002	136	433	1,054	2,065	2,746	2,109	793	147	10	-
2001	145	362	1,032	2,104	2,733	2,012	759	134	10	-
2000	115	398	940	2,052	2,478	1,879	726	115	8	1
1999	117	338	965	1,940	2,367	1,699	639	110	6	-
Female										
2002	87	193	546	1,303	2,025	1,850	988	293	22	-
2001	56	192	561	1,342	1,946	1,774	896	291	33	-
2000	73	228	542	1,270	1,852	1,688	901	254	19	-
1999	52	201	516	1,126	1,677	1,633	853	203	25	-
Hispanic										
Both sexes										
2002	13	39	63	104	119	96	33	7	2	-
2001	8	30	68	108	122	94	34	8	2	-
2000	11	34	50	89	102	81	42	5	2	-
1999	6	24	60	97	87	63	36	9	2	-
Male										
2002	11	26	45	63	59	53	14	1	-	-
2001	6	19	37	63	78	50	14	3	-	-
2000	6	19	27	53	59	49	18	2	2	-
1999	4	19	33	60	53	31	14	4	-	-
Female										
2002	2	13	18	41	60	43	19	6	2	-
2001	2	11	31	45	44	44	20	5	2	-
2000	5	15	23	36	43	32	24	3	-	-
1999	2	5	27	37	34	32	22	5	2	-
Non-Hispanic[1]										
Both sexes										
2002	209	586	1,534	3,257	4,642	3,856	1,745	432	30	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Parkinson's disease (G20-G21)														
Non-Hispanic[1]														
Both sexes														
2001	16,019	-	-	-	-	-	-	-	1	3	12	12	29	95
2000	15,223	-	1	-	-	-	-	2	2	4	3	12	31	78
1999	14,177	1	-	-	-	-	2	1	1	1	3	8	33	71
Male														
2002	9,290	-	-	-	-	-	-	-	-	-	2	11	26	52
2001	9,104	-	-	-	-	-	-	-	-	1	10	7	23	68
2000	8,544	-	-	-	-	-	-	1	2	3	2	8	18	52
1999	8,029	-	-	-	-	-	2	1	1	1	1	5	23	43
Female														
2002	7,145	-	-	-	-	-	-	-	-	1	2	4	11	35
2001	6,915	-	-	-	-	-	-	-	1	2	2	5	6	27
2000	6,679	-	1	-	-	-	-	1	-	1	1	4	13	26
1999	6,148	1	-	-	-	-	-	-	-	-	2	3	10	28
Non-Hispanic White														
Both sexes														
2002	15,626	-	-	-	-	-	-	-	-	-	2	12	36	74
2001	15,239	-	-	-	-	-	-	-	1	2	9	9	27	85
2000	14,497	-	-	-	-	-	-	1	2	3	1	9	27	71
1999	13,475	1	-	-	-	-	2	1	-	1	3	8	29	63
Male														
2002	8,838	-	-	-	-	-	-	-	-	-	1	8	25	45
2001	8,685	-	-	-	-	-	-	-	-	-	8	6	21	61
2000	8,116	-	-	-	-	-	-	1	2	2	1	7	15	46
1999	7,648	-	-	-	-	-	2	1	-	1	1	5	22	40
Female														
2002	6,788	-	-	-	-	-	-	-	-	-	1	4	11	29
2001	6,554	-	-	-	-	-	-	-	1	2	1	3	6	24
2000	6,381	-	-	-	-	-	-	-	-	1	-	2	12	25
1999	5,827	1	-	-	-	-	-	-	-	-	2	3	7	23
Non-Hispanic Black														
Both sexes														
2002	547	-	-	-	-	-	-	-	-	1	-	2	1	8
2001	547	-	-	-	-	-	-	-	-	1	3	3	2	5
2000	528	-	-	-	-	-	-	1	-	1	2	2	4	5
1999	502	-	-	-	-	-	-	-	-	-	-	-	4	7
Male														
2002	306	-	-	-	-	-	-	-	-	-	-	2	1	3
2001	291	-	-	-	-	-	-	-	-	1	2	1	2	4
2000	304	-	-	-	-	-	-	-	-	1	1	1	3	4
1999	269	-	-	-	-	-	-	-	-	-	-	-	1	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Parkinson's disease (G20-G21)										
Non-Hispanic[1]										
Both sexes										
2001	193	523	1,523	3,329	4,544	3,682	1,617	415	41	-
2000	177	591	1,429	3,223	4,219	3,480	1,581	364	25	1
1999	163	511	1,419	2,967	3,950	3,263	1,452	303	28	-
Male										
2002	124	406	1,007	1,996	2,681	2,054	776	145	10	-
2001	139	342	993	2,034	2,647	1,957	743	130	10	-
2000	109	378	912	1,992	2,415	1,826	706	113	6	1
1999	113	318	930	1,878	2,311	1,667	623	106	6	-
Female										
2002	85	180	527	1,261	1,961	1,802	969	287	20	-
2001	54	181	530	1,295	1,897	1,725	874	285	31	-
2000	68	213	517	1,231	1,804	1,654	875	251	19	-
1999	50	193	489	1,089	1,639	1,596	829	197	22	-
Non-Hispanic White										
Both sexes										
2002	195	530	1,429	3,084	4,461	3,695	1,672	408	28	-
2001	172	483	1,428	3,172	4,349	3,528	1,547	392	35	-
2000	165	550	1,345	3,070	4,036	3,332	1,515	345	24	1
1999	144	471	1,336	2,811	3,783	3,124	1,387	285	26	-
Male										
2002	118	369	943	1,896	2,577	1,969	741	136	10	-
2001	129	319	933	1,950	2,536	1,881	709	123	9	-
2000	103	352	862	1,887	2,304	1,752	669	107	5	1
1999	102	294	888	1,780	2,225	1,593	590	98	6	-
Female										
2002	77	161	486	1,188	1,884	1,726	931	272	18	-
2001	43	164	495	1,222	1,813	1,647	838	269	26	-
2000	62	198	483	1,183	1,732	1,580	846	238	19	-
1999	42	177	448	1,031	1,558	1,531	797	187	20	-
Non-Hispanic Black										
Both sexes										
2002	10	41	70	120	123	103	49	18	1	-
2001	15	25	66	109	129	113	54	18	4	-
2000	9	28	65	116	130	102	47	15	1	-
1999	15	32	63	109	112	93	51	14	2	-
Male										
2002	6	29	47	66	72	56	19	5	-	-
2001	9	15	44	55	72	53	29	4	-	-
2000	5	19	41	78	76	44	24	6	1	-
1999	10	22	36	67	56	47	22	5	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Parkinson's disease (G20-G21)														
Non-Hispanic Black														
Female														
2002	241	-	-	-	-	-	-	-	-	1	-	-	-	5
2001	256	-	-	-	-	-	-	-	-	-	1	2	-	1
2000	224	-	-	-	-	-	-	1	-	-	1	1	1	1
1999	233	-	-	-	-	-	-	-	-	-	-	-	3	4
Origin not stated[2]														
Both sexes														
2002	33	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	42	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	34	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	27	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	22	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	27	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	20	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	11	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	11	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	15	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	14	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	16	-	-	-	-	-	-	-	-	-	-	-	-	-
Alzheimer's disease (G30)														
All origins														
Both sexes														
2002	58,866	-	-	-	1	-	-	-	2	3	9	12	40	135
2001	53,852	-	-	-	-	-	-	-	2	1	2	17	56	165
2000	49,558	-	-	-	-	-	-	1	1	2	4	17	47	142
1999	44,536	-	-	-	-	-	1	-	-	1	6	20	40	125
Male														
2002	16,989	-	-	-	-	-	-	-	2	3	5	5	22	55
2001	15,762	-	-	-	-	-	-	-	2	1	2	11	27	61
2000	14,438	-	-	-	-	-	-	1	-	1	3	10	24	60
1999	13,391	-	-	-	-	-	1	-	-	-	3	10	18	68
Female														
2002	41,877	-	-	-	1	-	-	-	-	-	4	7	18	80
2001	38,090	-	-	-	-	-	-	-	-	-	-	6	29	104
2000	35,120	-	-	-	-	-	-	-	1	1	1	7	23	82
1999	31,145	-	-	-	-	-	-	-	-	1	3	10	22	57

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Parkinson's disease (G20-G21)										
Non-Hispanic Black										
Female										
2002	4	12	23	54	51	47	30	13	1	-
2001	6	10	22	54	57	60	25	14	4	-
2000	4	9	24	38	54	58	23	9	-	-
1999	5	10	27	42	56	46	29	9	2	-
Origin not stated[2]										
Both sexes										
2002	1	1	3	7	10	7	3	1	-	-
2001	-	1	2	9	13	10	4	2	-	-
2000	-	1	3	10	9	6	4	-	-	-
1999	-	4	2	2	7	6	4	1	1	-
Male										
2002	1	1	2	6	6	2	3	1	-	-
2001	-	1	2	7	8	5	2	1	-	-
2000	-	1	1	7	4	4	2	-	-	-
1999	-	1	2	2	3	1	2	-	-	-
Female										
2002	-	-	1	1	4	5	-	-	-	-
2001	-	-	-	2	5	5	2	1	-	-
2000	-	-	2	3	5	2	2	-	-	-
1999	-	3	-	-	4	5	2	1	1	-
Alzheimer's disease (G30)										
All origins										
Both sexes										
2002	375	939	2,663	7,078	13,057	16,267	12,267	5,140	878	-
2001	364	858	2,564	6,568	11,974	14,843	11,168	4,474	796	-
2000	349	868	2,563	6,319	10,934	13,337	10,257	4,014	701	2
1999	323	830	2,374	5,769	10,067	11,862	9,041	3,473	604	-
Male										
2002	173	408	1,139	2,793	4,434	4,569	2,570	732	79	-
2001	167	397	1,075	2,616	4,179	4,225	2,312	608	79	-
2000	150	420	1,134	2,433	3,795	3,687	2,101	564	54	1
1999	154	389	1,048	2,332	3,572	3,382	1,861	505	48	-
Female										
2002	202	531	1,524	4,285	8,623	11,698	9,697	4,408	799	-
2001	197	461	1,489	3,952	7,795	10,618	8,856	3,866	717	-
2000	199	448	1,429	3,886	7,139	9,650	8,156	3,450	647	1
1999	169	441	1,326	3,437	6,495	8,480	7,180	2,968	556	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alzheimer's disease (G30)														
Hispanic														
Both sexes														
2002	1,474	-	-	-	-	-	-	-	-	-	1	-	-	3
2001	1,239	-	-	-	-	-	-	-	-	-	1	2	1	6
2000	1,127	-	-	-	-	-	-	-	-	-	-	2	4	5
1999	981	-	-	-	-	-	1	-	-	-	-	-	3	2
Male														
2002	464	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	401	-	-	-	-	-	-	-	-	-	1	2	1	-
2000	383	-	-	-	-	-	-	-	-	-	-	1	3	2
1999	314	-	-	-	-	-	1	-	-	-	-	-	-	-
Female														
2002	1,010	-	-	-	-	-	-	-	-	-	-	-	-	3
2001	838	-	-	-	-	-	-	-	-	-	-	-	-	6
2000	744	-	-	-	-	-	-	-	-	-	-	1	1	3
1999	667	-	-	-	-	-	-	-	-	-	-	-	3	2
Non-Hispanic[1]														
Both sexes														
2002	57,246	-	-	-	1	-	-	-	2	3	8	12	40	132
2001	52,516	-	-	-	-	-	-	-	2	1	1	15	55	159
2000	48,331	-	-	-	-	-	-	1	1	2	4	15	43	137
1999	43,471	-	-	-	-	-	-	-	-	1	6	20	37	123
Male														
2002	16,491	-	-	-	-	-	-	-	2	3	4	5	22	55
2001	15,334	-	-	-	-	-	-	-	2	1	1	9	26	61
2000	14,034	-	-	-	-	-	-	1	-	1	3	9	21	58
1999	13,052	-	-	-	-	-	-	-	-	-	3	10	18	68
Female														
2002	40,755	-	-	-	1	-	-	-	-	-	4	7	18	77
2001	37,182	-	-	-	-	-	-	-	-	-	-	6	29	98
2000	34,297	-	-	-	-	-	-	-	1	1	1	6	22	79
1999	30,419	-	-	-	-	-	-	-	-	1	3	10	19	55
Non-Hispanic White														
Both sexes														
2002	53,486	-	-	-	1	-	-	-	1	1	5	12	37	124
2001	49,030	-	-	-	-	-	-	-	2	1	1	11	53	152
2000	45,246	-	-	-	-	-	-	1	1	2	4	13	37	127
1999	40,835	-	-	-	-	-	-	-	-	1	4	18	33	111
Male														
2002	15,399	-	-	-	-	-	-	-	1	1	3	5	19	51
2001	14,295	-	-	-	-	-	-	-	2	1	1	7	24	56
2000	13,123	-	-	-	-	-	-	1	-	1	3	7	18	53

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alzheimer's disease (G30)										
Hispanic										
Both sexes										
2002	17	39	88	211	324	359	297	118	17	-
2001	13	32	80	194	271	313	197	106	23	-
2000	14	43	85	180	221	288	201	76	8	-
1999	12	25	75	166	196	244	174	74	9	-
Male										
2002	8	18	37	85	111	102	72	25	5	-
2001	6	16	28	89	97	88	50	20	3	-
2000	7	16	47	75	91	83	40	16	2	-
1999	5	17	34	67	61	72	38	18	1	-
Female										
2002	9	21	51	126	213	257	225	93	12	-
2001	7	16	52	105	174	225	147	86	20	-
2000	7	27	38	105	130	205	161	60	6	-
1999	7	8	41	99	135	172	136	56	8	-
Non-Hispanic[1]										
Both sexes										
2002	356	895	2,568	6,849	12,703	15,872	11,942	5,002	861	-
2001	350	826	2,479	6,362	11,689	14,497	10,949	4,360	771	-
2000	332	823	2,473	6,130	10,699	13,013	10,039	3,929	689	1
1999	311	804	2,295	5,594	9,853	11,596	8,845	3,392	594	-
Male										
2002	165	388	1,099	2,699	4,316	4,462	2,493	704	74	-
2001	161	381	1,046	2,520	4,080	4,125	2,258	588	75	-
2000	142	404	1,085	2,356	3,702	3,596	2,057	548	51	-
1999	149	372	1,012	2,262	3,506	3,302	1,816	487	47	-
Female										
2002	191	507	1,469	4,150	8,387	11,410	9,449	4,298	787	-
2001	189	445	1,433	3,842	7,609	10,372	8,691	3,772	696	-
2000	190	419	1,388	3,774	6,997	9,417	7,982	3,381	638	1
1999	162	432	1,283	3,332	6,347	8,294	7,029	2,905	547	-
Non-Hispanic White										
Both sexes										
2002	324	804	2,345	6,332	11,895	14,914	11,241	4,684	766	-
2001	328	736	2,284	5,829	10,913	13,619	10,349	4,070	682	-
2000	301	751	2,253	5,676	10,028	12,242	9,507	3,681	622	-
1999	288	731	2,090	5,195	9,258	10,953	8,422	3,189	542	-
Male										
2002	152	345	1,008	2,510	4,034	4,217	2,343	651	59	-
2001	149	338	967	2,329	3,808	3,868	2,125	557	63	-
2000	130	373	994	2,195	3,491	3,376	1,930	509	42	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alzheimer's disease (G30)														
Non-Hispanic White														
Male														
1999	12,253	-	-	-	-	-	-	-	-	-	2	10	16	61
Female														
2002	38,087	-	-	-	1	-	-	-	-	-	2	7	18	73
2001	34,735	-	-	-	-	-	-	-	-	-	-	4	29	96
2000	32,123	-	-	-	-	-	-	-	1	1	1	6	19	74
1999	28,582	-	-	-	-	-	-	-	-	1	2	8	17	50
Non-Hispanic Black														
Both sexes														
2002	3,303	-	-	-	-	-	-	-	1	2	2	-	3	3
2001	3,104	-	-	-	-	-	-	-	-	-	-	3	2	6
2000	2,719	-	-	-	-	-	-	-	-	-	-	1	4	7
1999	2,325	-	-	-	-	-	-	-	-	-	1	2	2	10
Male														
2002	940	-	-	-	-	-	-	-	1	2	1	-	3	2
2001	906	-	-	-	-	-	-	-	-	-	-	1	2	5
2000	774	-	-	-	-	-	-	-	-	-	-	1	2	5
1999	689	-	-	-	-	-	-	-	-	-	-	-	2	6
Female														
2002	2,363	-	-	-	-	-	-	-	-	-	1	-	-	1
2001	2,198	-	-	-	-	-	-	-	-	-	-	2	-	1
2000	1,945	-	-	-	-	-	-	-	-	-	-	-	2	2
1999	1,636	-	-	-	-	-	-	-	-	-	1	2	-	4
Origin not stated[2]														
Both sexes														
2002	146	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	97	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	100	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	84	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	34	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	27	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	21	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	25	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	112	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	70	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	79	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	59	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alzheimer's disease (G30)										
Non-Hispanic White										
Male										
1999	139	338	925	2,113	3,319	3,120	1,719	451	40	-
Female										
2002	172	459	1,337	3,822	7,861	10,697	8,898	4,033	707	-
2001	179	398	1,317	3,500	7,105	9,751	8,224	3,513	619	-
2000	171	378	1,259	3,481	6,537	8,866	7,577	3,172	580	-
1999	149	393	1,165	3,082	5,939	7,833	6,703	2,738	502	-
Non-Hispanic Black										
Both sexes										
2002	28	77	200	468	705	834	611	288	81	-
2001	20	80	173	499	677	782	522	260	80	-
2000	23	67	194	399	605	680	463	218	57	1
1999	19	62	187	358	525	567	362	182	48	-
Male										
2002	12	36	86	175	243	199	118	50	12	-
2001	12	38	72	182	230	219	109	25	11	-
2000	11	30	83	136	184	181	99	36	6	-
1999	9	31	81	130	160	157	75	31	7	-
Female										
2002	16	41	114	293	462	635	493	238	69	-
2001	8	42	101	317	447	563	413	235	69	-
2000	12	37	111	263	421	499	364	182	51	1
1999	10	31	106	228	365	410	287	151	41	-
Origin not stated[2]										
Both sexes										
2002	2	5	7	18	30	36	28	20	-	-
2001	1	-	5	12	14	33	22	8	2	-
2000	3	2	5	9	14	36	17	9	4	1
1999	-	1	4	9	18	22	22	7	1	-
Male										
2002	-	2	3	9	7	5	5	3	-	-
2001	-	-	1	7	2	12	4	-	1	-
2000	1	-	2	2	2	8	4	-	1	1
1999	-	-	2	3	5	8	7	-	-	-
Female										
2002	2	3	4	9	23	31	23	17	-	-
2001	1	-	4	5	12	21	18	8	1	-
2000	2	2	3	7	12	28	13	9	3	-
1999	-	1	2	6	13	14	15	7	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Major cardiovascular diseases (I00-I78)														
All origins														
Both sexes														
2002	918,628	640	223	127	231	493	790	1,348	2,638	5,541	11,421	18,716	27,084	35,004
2001	922,334	600	285	139	226	443	837	1,351	2,687	5,568	11,118	17,768	26,682	33,335
2000	936,923	636	234	136	226	493	816	1,295	2,497	5,491	11,133	17,785	25,798	33,454
1999	950,314	644	234	144	210	548	767	1,270	2,603	5,918	11,092	17,354	25,158	33,344
Male														
2002	429,682	343	122	57	137	328	517	869	1,699	3,602	7,754	12,998	18,930	23,721
2001	428,065	315	149	65	119	278	529	840	1,715	3,672	7,508	12,418	18,442	22,489
2000	435,912	341	128	78	128	295	488	800	1,578	3,538	7,572	12,544	17,991	22,817
1999	441,827	347	127	84	115	345	454	769	1,673	3,898	7,573	12,150	17,595	22,653
Female														
2002	488,946	297	101	70	94	165	273	479	939	1,939	3,667	5,718	8,154	11,283
2001	494,269	285	136	74	107	165	308	511	972	1,896	3,610	5,350	8,240	10,846
2000	501,011	295	106	58	98	198	328	495	919	1,953	3,561	5,241	7,807	10,637
1999	508,487	297	107	60	95	203	313	501	930	2,020	3,519	5,204	7,563	10,691
Hispanic														
Both sexes														
2002	36,543	109	36	17	34	57	119	176	269	435	749	1,169	1,521	1,956
2001	35,565	98	44	25	31	65	97	157	285	420	661	1,037	1,524	1,745
2000	34,059	106	37	21	26	60	109	162	257	410	684	995	1,356	1,711
1999	33,665	115	41	22	25	67	93	149	259	426	631	994	1,392	1,696
Male														
2002	18,824	52	16	7	14	36	86	134	188	289	540	825	1,051	1,346
2001	18,138	53	21	14	15	40	68	109	199	296	462	748	1,058	1,170
2000	17,442	54	21	10	20	41	69	109	174	268	471	711	960	1,174
1999	17,304	49	21	14	15	41	62	107	185	296	459	696	1,004	1,124
Female														
2002	17,719	57	20	10	20	21	33	42	81	146	209	344	470	610
2001	17,427	45	23	11	16	25	29	48	86	124	199	289	466	575
2000	16,617	52	16	11	6	19	40	53	83	142	213	284	396	537
1999	16,361	66	20	8	10	26	31	42	74	130	172	298	388	572
Non-Hispanic[1]														
Both sexes														
2002	879,170	528	187	110	197	436	669	1,166	2,356	5,081	10,626	17,480	25,433	32,910
2001	883,949	499	239	113	195	375	733	1,187	2,391	5,128	10,408	16,651	25,044	31,434
2000	899,801	526	196	114	199	433	704	1,126	2,231	5,055	10,378	16,695	24,326	31,592
1999	913,476	528	192	122	184	479	673	1,116	2,337	5,465	10,404	16,278	23,657	31,488
Male														
2002	409,343	289	106	50	123	292	429	733	1,501	3,295	7,182	12,126	17,775	22,267
2001	408,437	260	128	51	104	235	457	725	1,506	3,362	7,004	11,613	17,303	21,204
2000	416,837	284	107	68	108	254	418	687	1,401	3,250	7,044	11,763	16,939	21,532

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Major cardiovascular diseases (I00-I78)										
All origins										
Both sexes										
2002	43,426	57,731	86,340	126,160	161,255	163,421	117,826	47,620	10,542	51
2001	42,889	59,066	89,343	130,631	161,364	164,000	116,097	47,232	10,601	72
2000	44,028	61,769	94,418	135,296	160,906	165,313	117,092	47,427	10,618	62
1999	44,489	64,775	98,653	138,680	163,872	166,070	116,257	47,656	10,527	49
Male										
2002	28,316	35,603	49,716	65,751	73,019	61,344	33,439	9,860	1,521	36
2001	27,831	36,221	51,535	67,972	72,817	59,914	32,127	9,565	1,491	53
2000	28,606	37,721	54,789	70,196	72,601	60,379	32,104	9,636	1,539	43
1999	28,872	39,978	56,926	72,052	73,515	60,030	31,573	9,558	1,502	38
Female										
2002	15,110	22,128	36,624	60,409	88,236	102,077	84,387	37,760	9,021	15
2001	15,058	22,845	37,808	62,659	88,547	104,086	83,970	37,667	9,110	19
2000	15,422	24,048	39,629	65,100	88,305	104,934	84,988	37,791	9,079	19
1999	15,617	24,797	41,727	66,628	90,357	106,040	84,684	38,098	9,025	11
Hispanic										
Both sexes										
2002	2,504	3,097	4,245	5,078	5,284	4,558	3,306	1,493	330	1
2001	2,251	3,094	4,231	5,154	4,985	4,684	3,247	1,405	318	7
2000	2,330	3,003	4,111	4,742	4,638	4,566	3,165	1,299	266	5
1999	2,277	3,147	4,003	4,708	4,583	4,442	3,041	1,290	258	6
Male										
2002	1,612	1,849	2,455	2,635	2,404	1,745	1,075	389	75	1
2001	1,412	1,898	2,371	2,652	2,314	1,798	1,019	343	72	6
2000	1,483	1,766	2,357	2,505	2,117	1,692	1,014	358	64	4
1999	1,436	1,890	2,322	2,523	1,996	1,645	992	358	63	6
Female										
2002	892	1,248	1,790	2,443	2,880	2,813	2,231	1,104	255	-
2001	839	1,196	1,860	2,502	2,671	2,886	2,228	1,062	246	1
2000	847	1,237	1,754	2,237	2,521	2,874	2,151	941	202	1
1999	841	1,257	1,681	2,185	2,587	2,797	2,049	932	195	-
Non-Hispanic[1]										
Both sexes										
2002	40,740	54,399	81,776	120,688	155,544	158,407	114,222	46,000	10,177	38
2001	40,441	55,778	84,840	125,069	155,957	158,886	112,579	45,695	10,260	47
2000	41,472	58,562	89,955	130,119	155,841	160,276	113,625	46,004	10,329	43
1999	41,985	61,383	94,260	133,532	158,822	161,174	112,886	46,241	10,236	34
Male										
2002	26,584	33,601	47,065	62,889	70,414	59,421	32,288	9,448	1,440	25
2001	26,265	34,192	48,986	65,093	70,302	57,972	31,035	9,192	1,412	36
2000	26,963	35,803	52,202	67,446	70,308	58,511	31,008	9,245	1,468	28

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Major cardiovascular diseases (I00-I78)														
Non-Hispanic[1]														
Male														
1999	422,824	298	106	70	99	303	391	659	1,483	3,581	7,068	11,392	16,515	21,413
Female														
2002	469,827	239	81	60	74	144	240	433	855	1,786	3,444	5,354	7,658	10,643
2001	475,512	239	111	62	91	140	276	462	885	1,766	3,404	5,038	7,741	10,230
2000	482,964	242	89	46	91	179	286	439	830	1,805	3,334	4,932	7,387	10,060
1999	490,652	230	86	52	85	176	282	457	854	1,884	3,336	4,886	7,142	10,075
Non-Hispanic White														
Both sexes														
2002	758,738	314	125	75	120	289	406	681	1,440	3,374	7,464	12,325	18,473	25,150
2001	764,021	289	140	75	129	241	430	720	1,520	3,432	7,092	11,734	18,233	23,979
2000	780,445	330	126	74	121	272	418	681	1,403	3,392	7,164	11,764	18,016	24,261
1999	793,945	324	121	86	117	318	412	683	1,501	3,684	7,123	11,537	17,597	24,071
Male														
2002	352,520	177	77	35	76	192	270	443	944	2,256	5,303	8,993	13,442	17,508
2001	352,013	160	75	36	72	148	277	454	987	2,322	5,057	8,582	13,141	16,667
2000	360,757	180	75	47	71	161	257	428	933	2,296	5,126	8,783	13,002	16,982
1999	366,310	189	70	50	60	196	234	408	992	2,505	5,101	8,494	12,706	16,870
Female														
2002	406,218	137	48	40	44	97	136	238	496	1,118	2,161	3,332	5,031	7,642
2001	412,008	129	65	39	57	93	153	266	533	1,110	2,035	3,152	5,092	7,312
2000	419,688	150	51	27	50	111	161	253	470	1,096	2,038	2,981	5,014	7,279
1999	427,635	135	51	36	57	122	178	275	509	1,179	2,022	3,043	4,891	7,201
Non-Hispanic Black														
Both sexes														
2002	103,054	183	50	26	67	122	234	434	818	1,517	2,846	4,650	6,273	6,902
2001	103,213	184	86	33	52	123	272	429	780	1,500	3,006	4,432	6,107	6,601
2000	103,239	166	60	33	62	143	253	403	757	1,498	2,898	4,481	5,628	6,524
1999	103,629	172	60	35	58	147	241	390	741	1,615	2,995	4,273	5,448	6,612
Male														
2002	47,656	96	24	10	40	79	138	258	481	897	1,654	2,778	3,857	4,200
2001	47,646	93	45	12	27	80	160	251	455	898	1,727	2,683	3,703	3,956
2000	47,408	88	27	17	30	79	139	230	422	846	1,698	2,675	3,468	4,019
1999	47,982	91	32	19	34	98	142	228	431	955	1,761	2,585	3,399	4,001
Female														
2002	55,398	87	26	16	27	43	96	176	337	620	1,192	1,872	2,416	2,702
2001	55,567	91	41	21	25	43	112	178	325	602	1,279	1,749	2,404	2,645
2000	55,831	78	33	16	32	64	114	173	335	652	1,200	1,806	2,160	2,505
1999	55,647	81	28	16	24	49	99	162	310	660	1,234	1,688	2,049	2,611

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Major cardiovascular diseases (I00-I78)										
Non-Hispanic[1]										
Male										
1999	27,262	37,928	54,345	69,278	71,310	58,202	30,487	9,173	1,436	25
Female										
2002	14,156	20,798	34,711	57,799	85,130	98,986	81,934	36,552	8,737	13
2001	14,176	21,586	35,854	59,976	85,655	100,914	81,544	36,503	8,848	11
2000	14,509	22,759	37,753	62,673	85,533	101,765	82,617	36,759	8,861	15
1999	14,723	23,455	39,915	64,254	87,512	102,972	82,399	37,068	8,800	9
Non-Hispanic White										
Both sexes										
2002	31,715	43,606	68,682	104,667	139,407	144,795	104,762	41,989	8,853	26
2001	31,501	44,581	71,517	108,821	140,101	145,157	103,653	41,714	8,934	28
2000	32,405	47,591	76,117	113,698	140,391	146,434	104,746	41,938	9,074	29
1999	33,023	50,223	79,888	116,966	143,429	147,187	104,261	42,308	9,068	18
Male										
2002	21,228	27,612	40,404	55,507	63,824	54,839	29,602	8,570	1,200	18
2001	21,059	28,119	42,094	57,548	63,784	53,354	28,550	8,310	1,195	22
2000	21,699	29,895	45,131	59,825	64,009	53,818	28,511	8,257	1,251	20
1999	21,981	31,816	47,037	61,605	65,165	53,309	27,980	8,288	1,241	13
Female										
2002	10,487	15,994	28,278	49,160	75,583	89,956	75,160	33,419	7,653	8
2001	10,442	16,462	29,423	51,273	76,317	91,803	75,103	33,404	7,739	6
2000	10,706	17,696	30,986	53,873	76,382	92,616	76,235	33,681	7,823	9
1999	11,042	18,407	32,851	55,361	78,264	93,878	76,281	34,020	7,827	5
Non-Hispanic Black										
Both sexes										
2002	7,932	9,321	11,140	13,488	13,444	11,215	7,835	3,396	1,150	11
2001	7,882	9,697	11,387	13,796	13,229	11,446	7,562	3,442	1,150	17
2000	7,978	9,534	12,027	13,905	13,049	11,736	7,495	3,464	1,131	14
1999	7,875	9,789	12,453	14,150	12,979	11,800	7,298	3,457	1,026	15
Male										
2002	4,662	5,119	5,651	6,031	5,292	3,521	2,020	655	187	6
2001	4,533	5,226	5,829	6,257	5,279	3,627	1,914	694	184	13
2000	4,549	5,053	6,021	6,267	5,167	3,735	1,929	753	188	8
1999	4,608	5,329	6,238	6,332	4,997	3,853	1,935	736	167	11
Female										
2002	3,270	4,202	5,489	7,457	8,152	7,694	5,815	2,741	963	5
2001	3,349	4,471	5,558	7,539	7,950	7,819	5,648	2,748	966	4
2000	3,429	4,481	6,006	7,638	7,882	8,001	5,566	2,711	943	6
1999	3,267	4,460	6,215	7,818	7,982	7,947	5,363	2,721	859	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Major cardiovascular diseases (I00-I78)														
Origin not stated[2]														
Both sexes														
2002	2,915	3	-	-	-	-	2	6	13	25	46	67	130	138
2001	2,820	3	2	1	-	3	7	7	11	20	49	80	114	156
2000	3,063	4	1	1	1	-	3	7	9	26	71	95	116	151
1999	3,173	1	1	-	1	2	1	5	7	27	57	82	109	160
Male														
2002	1,515	2	-	-	-	-	2	2	10	18	32	47	104	108
2001	1,490	2	-	-	-	3	4	6	10	14	42	57	81	115
2000	1,633	3	-	-	-	-	1	4	3	20	57	70	92	111
1999	1,699	-	-	-	1	1	1	3	5	21	46	62	76	116
Female														
2002	1,400	1	-	-	-	-	-	4	3	7	14	20	26	30
2001	1,330	1	2	1	-	-	3	1	1	6	7	23	33	41
2000	1,430	1	1	1	1	-	2	3	6	6	14	25	24	40
1999	1,474	1	1	-	-	1	-	2	2	6	11	20	33	44
Diseases of heart (I00-I09,I11,I13, I20-I51)														
All origins														
Both sexes														
2002	696,947	500	165	92	163	405	617	1,076	2,089	4,448	9,240	15,179	22,391	28,895
2001	700,142	479	225	98	174	347	652	1,052	2,108	4,419	8,907	14,390	22,009	27,564
2000	710,760	493	181	106	165	403	628	1,013	1,945	4,333	8,848	14,313	21,167	27,676
1999	725,192	524	183	116	161	463	606	992	2,074	4,703	8,897	14,096	20,898	27,605
Male														
2002	340,933	267	87	40	101	275	411	716	1,415	3,048	6,590	10,983	16,256	20,199
2001	339,095	243	121	43	95	223	424	672	1,396	3,038	6,308	10,491	15,789	19,200
2000	344,807	260	107	64	99	248	390	640	1,282	2,945	6,341	10,596	15,331	19,456
1999	351,617	270	104	66	91	297	367	618	1,370	3,252	6,388	10,346	15,143	19,370
Female														
2002	356,014	233	78	52	62	130	206	360	674	1,400	2,650	4,196	6,135	8,696
2001	361,047	236	104	55	79	124	228	380	712	1,381	2,599	3,899	6,220	8,364
2000	365,953	233	74	42	66	155	238	373	663	1,388	2,507	3,717	5,836	8,220
1999	373,575	254	79	50	70	166	239	374	704	1,451	2,509	3,750	5,755	8,235
Hispanic														
Both sexes														
2002	27,887	85	24	12	18	39	89	126	195	313	539	851	1,169	1,528
2001	27,090	79	33	17	21	45	67	115	199	302	448	736	1,139	1,362
2000	25,819	86	29	17	21	49	76	112	168	284	476	709	1,045	1,355
1999	25,866	96	33	18	18	52	64	93	193	320	441	716	1,090	1,331

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Major cardiovascular diseases (I00-I78)										
Origin not stated[2]										
Both sexes										
2002	182	235	319	394	427	456	298	127	35	12
2001	197	194	272	408	422	430	271	132	23	18
2000	226	204	352	435	427	471	302	124	23	14
1999	227	245	390	440	467	454	330	125	33	9
Male										
2002	120	153	196	227	201	178	76	23	6	10
2001	154	131	178	227	201	144	73	30	7	11
2000	160	152	230	245	176	176	82	33	7	11
1999	174	160	259	251	209	183	94	27	3	7
Female										
2002	62	82	123	167	226	278	222	104	29	2
2001	43	63	94	181	221	286	198	102	16	7
2000	66	52	122	190	251	295	220	91	16	3
1999	53	85	131	189	258	271	236	98	30	2
Diseases of heart (I00-I09,I11,I13, I20-I51)										
All origins										
Both sexes										
2002	35,339	45,948	66,599	94,844	118,737	119,565	86,848	35,664	8,096	47
2001	34,922	47,131	69,168	98,099	118,893	119,848	85,878	35,479	8,234	66
2000	35,723	49,178	73,227	101,764	118,296	121,003	86,424	35,605	8,210	59
1999	36,562	52,206	77,047	104,841	121,311	122,126	85,752	35,797	8,185	47
Male										
2002	23,701	29,045	39,613	51,211	56,006	46,515	25,549	7,663	1,208	34
2001	23,319	29,643	40,994	52,960	55,616	45,298	24,606	7,394	1,175	47
2000	23,829	30,890	43,699	54,421	55,272	45,756	24,475	7,430	1,235	41
1999	24,292	33,118	45,718	56,178	56,292	45,667	23,999	7,442	1,192	37
Female										
2002	11,638	16,903	26,986	43,633	62,731	73,050	61,299	28,001	6,888	13
2001	11,603	17,488	28,174	45,139	63,277	74,550	61,272	28,085	7,059	19
2000	11,894	18,288	29,528	47,343	63,024	75,247	61,949	28,175	6,975	18
1999	12,270	19,088	31,329	48,663	65,019	76,459	61,753	28,355	6,993	10
Hispanic										
Both sexes										
2002	1,964	2,472	3,247	3,859	3,920	3,441	2,548	1,197	250	1
2001	1,790	2,466	3,264	3,875	3,757	3,506	2,531	1,080	252	6
2000	1,781	2,328	3,125	3,583	3,484	3,438	2,438	995	215	5
1999	1,824	2,464	3,093	3,618	3,428	3,363	2,349	1,050	206	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of heart (I00-I09,I11,I13, I20-I51)														
Hispanic														
Male														
2002	14,798	41	11	6	8	26	68	98	144	215	410	621	846	1,087
2001	14,195	43	16	9	10	27	49	84	142	228	337	565	832	956
2000	13,566	44	21	9	17	36	51	79	120	198	352	539	768	952
1999	13,554	40	20	11	12	31	42	77	142	235	337	531	817	924
Female														
2002	13,089	44	13	6	10	13	21	28	51	98	129	230	323	441
2001	12,895	36	17	8	11	18	18	31	57	74	111	171	307	406
2000	12,253	42	8	8	4	13	25	33	48	86	124	170	277	403
1999	12,312	56	13	7	6	21	22	16	51	85	104	185	273	407
Non-Hispanic[1]														
Both sexes														
2002	666,690	413	141	80	145	366	526	945	1,882	4,116	8,664	14,272	21,104	27,246
2001	670,787	397	191	81	153	299	578	933	1,901	4,100	8,415	13,588	20,771	26,068
2000	682,470	403	151	88	144	354	550	895	1,769	4,027	8,314	13,530	20,021	26,192
1999	696,719	427	149	98	142	409	541	894	1,876	4,358	8,409	13,314	19,715	26,140
Male														
2002	324,848	225	76	34	93	249	341	616	1,262	2,818	6,154	10,321	15,315	19,015
2001	323,673	198	105	34	85	193	371	585	1,247	2,799	5,934	9,876	14,883	18,146
2000	329,888	213	86	55	82	212	338	557	1,159	2,730	5,943	10,002	14,485	18,410
1999	336,615	230	84	55	78	265	324	538	1,224	2,997	6,012	9,762	14,257	18,345
Female														
2002	341,842	188	65	46	52	117	185	329	620	1,298	2,510	3,951	5,789	8,231
2001	347,114	199	86	47	68	106	207	348	654	1,301	2,481	3,712	5,888	7,922
2000	352,582	190	65	33	62	142	212	338	610	1,297	2,371	3,528	5,536	7,782
1999	360,104	197	65	43	64	144	217	356	652	1,361	2,397	3,552	5,458	7,795
Non-Hispanic White														
Both sexes														
2002	577,761	249	91	53	93	240	307	554	1,164	2,818	6,296	10,407	15,799	21,225
2001	582,349	234	108	52	100	186	336	563	1,212	2,803	5,931	9,888	15,656	20,256
2000	594,465	258	98	54	86	216	315	523	1,130	2,789	5,954	9,905	15,248	20,515
1999	607,704	259	92	65	91	269	316	542	1,213	2,998	5,938	9,762	15,061	20,346
Male														
2002	281,442	139	54	24	61	163	207	369	808	1,984	4,673	7,866	11,908	15,237
2001	280,713	127	59	23	57	117	219	367	817	1,984	4,411	7,516	11,662	14,516
2000	287,210	139	59	34	53	126	200	336	779	1,979	4,467	7,699	11,402	14,800
1999	293,206	145	55	36	48	167	190	331	821	2,134	4,456	7,501	11,246	14,675
Female														
2002	296,319	110	37	29	32	77	100	185	356	834	1,623	2,541	3,891	5,988

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13, I20-I51)										
Hispanic										
Male										
2002	1,306	1,511	1,930	2,054	1,844	1,372	820	327	52	1
2001	1,148	1,544	1,880	2,031	1,777	1,385	808	265	54	5
2000	1,176	1,409	1,836	1,912	1,613	1,312	793	275	50	4
1999	1,167	1,522	1,833	1,957	1,526	1,250	741	286	47	6
Female										
2002	658	961	1,317	1,805	2,076	2,069	1,728	870	198	-
2001	642	922	1,384	1,844	1,980	2,121	1,723	815	198	1
2000	605	919	1,289	1,671	1,871	2,126	1,645	720	165	1
1999	657	942	1,260	1,661	1,902	2,113	1,608	764	159	-
Non-Hispanic[1]										
Both sexes										
2002	33,219	43,274	63,077	90,673	114,483	115,769	84,079	34,366	7,816	34
2001	32,970	44,501	65,680	93,905	114,799	116,016	83,143	34,294	7,962	42
2000	33,751	46,667	69,809	97,840	114,488	117,193	83,757	34,509	7,977	41
1999	34,544	49,540	73,621	100,854	117,505	118,399	83,155	34,644	7,953	32
Male										
2002	22,290	27,403	37,509	48,972	54,000	44,997	24,670	7,314	1,151	23
2001	22,042	27,987	38,962	50,746	53,683	43,799	23,744	7,107	1,116	31
2000	22,514	29,341	41,669	52,315	53,519	44,305	23,618	7,128	1,180	27
1999	22,976	31,461	43,657	54,006	54,599	44,256	23,191	7,132	1,142	24
Female										
2002	10,929	15,871	25,568	41,701	60,483	70,772	59,409	27,052	6,665	11
2001	10,928	16,514	26,718	43,159	61,116	72,217	59,399	27,187	6,846	11
2000	11,237	17,326	28,140	45,525	60,969	72,888	60,139	27,381	6,797	14
1999	11,568	18,079	29,964	46,848	62,906	74,143	59,964	27,512	6,811	8
Non-Hispanic White										
Both sexes										
2002	26,269	35,099	53,429	78,991	102,984	106,117	77,340	31,405	6,808	23
2001	26,177	36,014	55,841	82,060	103,476	106,333	76,775	31,359	6,963	26
2000	26,777	38,325	59,423	85,862	103,504	107,375	77,430	31,597	7,052	29
1999	27,604	40,913	62,837	88,727	106,460	108,389	76,983	31,757	7,064	18
Male										
2002	18,076	22,786	32,479	43,461	49,155	41,657	22,700	6,647	972	16
2001	17,988	23,298	33,761	45,093	48,903	40,491	21,898	6,442	944	20
2000	18,379	24,751	36,273	46,643	48,933	40,903	21,817	6,412	1,006	20
1999	18,791	26,627	38,072	48,308	50,114	40,681	21,346	6,460	989	13
Female										
2002	8,193	12,313	20,950	35,530	53,829	64,460	54,640	24,758	5,836	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of heart (I00-I09,I11,I13, I20-I51)														
Non-Hispanic White														
Female														
2001	301,636	107	49	29	43	69	117	196	395	819	1,520	2,372	3,994	5,740
2000	307,255	119	39	20	33	90	115	187	351	810	1,487	2,206	3,846	5,715
1999	314,498	114	37	29	43	102	126	211	392	864	1,482	2,261	3,815	5,671
Non-Hispanic Black														
Both sexes														
2002	76,694	142	40	19	46	105	194	352	641	1,167	2,145	3,506	4,812	5,389
2001	76,794	140	73	27	43	104	218	339	622	1,162	2,262	3,345	4,611	5,178
2000	76,706	118	47	29	46	124	207	343	586	1,118	2,141	3,320	4,286	5,076
1999	77,713	141	49	32	45	128	208	318	595	1,246	2,260	3,239	4,209	5,202
Male														
2002	36,596	76	18	6	27	68	117	220	392	730	1,301	2,184	3,030	3,338
2001	36,541	65	40	10	24	69	136	199	379	715	1,356	2,087	2,868	3,170
2000	36,282	59	24	17	24	74	118	200	343	667	1,305	2,069	2,718	3,193
1999	37,077	71	27	18	27	90	122	191	355	773	1,392	2,024	2,692	3,245
Female														
2002	40,098	66	22	13	19	37	77	132	249	437	844	1,322	1,782	2,051
2001	40,253	75	33	17	19	35	82	140	243	447	906	1,258	1,743	2,008
2000	40,424	59	23	12	22	50	89	143	243	451	836	1,251	1,568	1,883
1999	40,636	70	22	14	18	38	86	127	240	473	868	1,215	1,517	1,957
Origin not stated[2]														
Both sexes														
2002	2,370	2	-	-	-	-	2	5	12	19	37	56	118	121
2001	2,265	3	1	-	-	3	7	4	8	17	44	66	99	134
2000	2,471	4	1	1	-	-	2	6	8	22	58	74	101	129
1999	2,607	1	1	-	1	2	1	5	5	25	47	66	93	134
Male														
2002	1,287	1	-	-	-	-	2	2	9	15	26	41	95	97
2001	1,227	2	-	-	-	3	4	3	7	11	37	50	74	98
2000	1,353	3	-	-	-	-	1	4	3	17	46	55	78	94
1999	1,448	-	-	-	1	1	1	3	4	20	39	53	69	101
Female														
2002	1,083	1	-	-	-	-	-	3	3	4	11	15	23	24
2001	1,038	1	1	-	-	-	3	1	1	6	7	16	25	36
2000	1,118	1	1	1	-	-	1	2	5	5	12	19	23	35
1999	1,159	1	1	-	-	1	-	2	1	5	8	13	24	33

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of heart (I00-I09,I11,I13, I20-I51)										
Non-Hispanic White										
Female										
2001	8,189	12,716	22,080	36,967	54,573	65,842	54,877	24,917	6,019	6
2000	8,398	13,574	23,150	39,219	54,571	66,472	55,613	25,185	6,046	9
1999	8,813	14,286	24,765	40,419	56,346	67,708	55,637	25,297	6,075	5
Non-Hispanic Black										
Both sexes										
2002	6,165	7,109	8,302	9,863	9,657	8,003	5,623	2,529	875	10
2001	6,031	7,418	8,522	10,132	9,549	8,157	5,431	2,542	874	14
2000	6,181	7,302	9,111	10,227	9,373	8,381	5,367	2,479	832	12
1999	6,144	7,622	9,397	10,398	9,410	8,483	5,242	2,546	785	14
Male										
2002	3,690	3,965	4,282	4,503	3,925	2,580	1,496	507	135	6
2001	3,543	4,044	4,447	4,705	3,916	2,630	1,451	528	149	10
2000	3,593	3,929	4,631	4,692	3,784	2,740	1,398	544	153	7
1999	3,662	4,226	4,790	4,725	3,691	2,831	1,433	553	128	11
Female										
2002	2,475	3,144	4,020	5,360	5,732	5,423	4,127	2,022	740	4
2001	2,488	3,374	4,075	5,427	5,633	5,527	3,980	2,014	725	4
2000	2,588	3,373	4,480	5,535	5,589	5,641	3,969	1,935	679	5
1999	2,482	3,396	4,607	5,673	5,719	5,652	3,809	1,993	657	3
Origin not stated[2]										
Both sexes										
2002	156	202	275	312	334	355	221	101	30	12
2001	162	164	224	319	337	326	204	105	20	18
2000	191	183	293	341	324	372	229	101	18	13
1999	194	202	333	369	378	364	248	103	26	9
Male										
2002	105	131	174	185	162	146	59	22	5	10
2001	129	112	152	183	156	114	54	22	5	11
2000	139	140	194	194	140	139	64	27	5	10
1999	149	135	228	215	167	161	67	24	3	7
Female										
2002	51	71	101	127	172	209	162	79	25	2
2001	33	52	72	136	181	212	150	83	15	7
2000	52	43	99	147	184	233	165	74	13	3
1999	45	67	105	154	211	203	181	79	23	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)														
All origins														
Both sexes														
2002	3,579	4	1	4	2	8	10	10	26	38	57	97	140	190
2001	3,489	3	1	1	4	3	8	11	23	45	85	107	125	168
2000	3,582	6	2	2	2	9	8	10	23	49	92	119	133	183
1999	3,676	6	1	2	3	5	8	12	30	47	77	101	144	189
Male														
2002	1,078	3	1	1	1	3	5	6	12	19	26	40	55	73
2001	1,031	3	1	-	2	2	5	4	11	18	36	46	41	73
2000	1,072	5	2	1	2	5	6	2	10	25	36	49	55	72
1999	1,042	2	1	1	1	4	6	5	12	19	30	40	50	64
Female														
2002	2,501	1	-	3	1	5	5	4	14	19	31	57	85	117
2001	2,458	-	-	1	2	1	3	7	12	27	49	61	84	95
2000	2,510	1	-	1	-	4	2	8	13	24	56	70	78	111
1999	2,634	4	-	1	2	1	2	7	18	28	47	61	94	125
Hispanic														
Both sexes														
2002	198	-	-	-	-	-	2	2	9	6	5	13	11	15
2001	156	-	1	-	1	-	2	-	2	5	7	9	7	9
2000	154	1	-	-	-	1	-	1	5	7	6	11	8	14
1999	166	2	-	-	-	-	3	1	3	9	7	10	13	12
Male														
2002	55	-	-	-	-	-	1	-	5	2	2	4	5	5
2001	51	-	1	-	1	-	1	-	2	1	4	3	2	4
2000	49	1	-	-	-	-	-	-	1	2	2	2	7	5
1999	38	-	-	-	-	-	3	-	3	2	1	4	2	2
Female														
2002	143	-	-	-	-	-	1	2	4	4	3	9	6	10
2001	105	-	-	-	-	-	1	-	-	4	3	6	5	5
2000	105	-	-	-	-	1	-	1	4	5	4	9	1	9
1999	128	2	-	-	-	-	-	1	-	7	6	6	11	10
Non-Hispanic[1]														
Both sexes														
2002	3,375	4	1	4	2	8	8	8	17	32	52	84	128	175
2001	3,327	3	-	1	3	2	6	11	21	40	78	97	117	159
2000	3,424	5	2	2	2	8	8	9	18	42	86	108	125	169
1999	3,498	4	1	2	3	5	5	11	27	37	69	91	130	176
Male														
2002	1,021	3	1	1	1	3	4	6	7	17	24	36	49	68
2001	978	3	-	-	1	1	4	4	9	17	32	42	39	69

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
All origins										
Both sexes										
2002	243	272	431	620	607	462	263	82	12	-
2001	193	307	437	599	589	468	235	67	10	-
2000	237	327	492	601	575	416	231	54	11	-
1999	222	338	516	614	613	442	235	63	8	-
Male										
2002	83	84	161	177	164	100	55	8	1	-
2001	62	106	125	182	148	106	50	10	-	-
2000	94	98	139	177	144	96	40	13	1	-
1999	69	107	183	164	161	89	26	8	-	-
Female										
2002	160	188	270	443	443	362	208	74	11	-
2001	131	201	312	417	441	362	185	57	10	-
2000	143	229	353	424	431	320	191	41	10	-
1999	153	231	333	450	452	353	209	55	8	-
Hispanic										
Both sexes										
2002	20	21	33	22	20	11	6	2	-	-
2001	18	23	25	17	13	10	6	-	1	-
2000	16	19	28	13	14	7	3	-	-	-
1999	14	25	19	22	13	8	2	3	-	-
Male										
2002	8	6	9	2	2	2	2	-	-	-
2001	3	11	4	8	3	2	1	-	-	-
2000	7	5	8	5	1	2	1	-	-	-
1999	6	6	4	3	2	-	-	-	-	-
Female										
2002	12	15	24	20	18	9	4	2	-	-
2001	15	12	21	9	10	8	5	-	1	-
2000	9	14	20	8	13	5	2	-	-	-
1999	8	19	15	19	11	8	2	3	-	-
Non-Hispanic[1]										
Both sexes										
2002	222	250	397	597	586	451	257	80	12	-
2001	174	284	412	581	576	457	229	67	9	-
2000	219	308	464	588	561	409	226	54	11	-
1999	207	311	495	590	600	433	233	60	8	-
Male										
2002	74	78	152	175	162	98	53	8	1	-
2001	59	95	121	174	145	104	49	10	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)														
Non-Hispanic[1]														
Male														
2000	1,022	4	2	1	2	5	6	2	9	23	34	47	48	67
1999	999	2	1	1	1	4	3	5	9	16	29	36	47	61
Female														
2002	2,354	1	-	3	1	5	4	2	10	15	28	48	79	107
2001	2,349	-	-	1	2	1	2	7	12	23	46	55	78	90
2000	2,402	1	-	1	-	3	2	7	9	19	52	61	77	102
1999	2,499	2	-	1	2	1	2	6	18	21	40	55	83	115
Non-Hispanic White														
Both sexes														
2002	3,007	4	1	1	2	3	5	4	10	26	31	57	99	134
2001	2,924	1	-	1	1	2	5	5	8	23	50	60	79	118
2000	3,073	4	2	2	2	5	4	6	13	27	56	79	99	134
1999	3,130	2	1	1	2	2	3	7	14	24	44	69	94	143
Male														
2002	897	3	1	1	1	-	3	3	4	13	13	26	37	51
2001	857	1	-	-	-	1	4	2	5	11	24	28	30	57
2000	917	3	2	1	2	3	3	1	8	17	25	37	41	57
1999	892	1	1	-	-	1	1	4	5	11	20	28	37	50
Female														
2002	2,110	1	-	-	1	3	2	1	6	13	18	31	62	83
2001	2,067	-	-	1	1	1	1	3	3	12	26	32	49	61
2000	2,156	1	-	1	-	2	1	5	5	10	31	42	58	77
1999	2,238	1	-	1	2	1	2	3	9	13	24	41	57	93
Non-Hispanic Black														
Both sexes														
2002	249	-	-	2	-	3	2	3	6	4	16	21	23	30
2001	302	2	-	-	1	-	1	3	10	15	26	27	33	31
2000	263	1	-	-	-	3	4	3	5	11	24	23	20	26
1999	269	1	-	1	1	2	2	4	9	11	19	20	29	29
Male														
2002	91	-	-	-	-	2	1	2	2	2	10	7	10	12
2001	86	2	-	-	-	-	-	-	3	6	7	11	9	10
2000	77	1	-	-	-	2	3	1	1	5	7	8	5	7
1999	86	1	-	1	1	2	2	1	2	4	8	8	10	9
Female														
2002	158	-	-	2	-	1	1	1	4	2	6	14	13	18
2001	216	-	-	-	1	-	1	3	7	9	19	16	24	21
2000	186	-	-	-	-	1	1	2	4	6	17	15	15	19

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
Non-Hispanic[1]										
Male										
2000	86	93	131	172	143	94	39	13	1	-
1999	62	101	178	161	159	89	26	8	-	-
Female										
2002	148	172	245	422	424	353	204	72	11	-
2001	115	189	291	407	431	353	180	57	9	-
2000	133	215	333	416	418	315	187	41	10	-
1999	145	210	317	429	441	344	207	52	8	-
Non-Hispanic White										
Both sexes										
2002	179	209	366	547	562	428	251	77	11	-
2001	147	238	363	540	554	436	222	62	9	-
2000	186	274	413	562	540	393	211	50	11	-
1999	171	276	454	555	570	412	225	54	7	-
Male										
2002	62	64	141	161	158	95	52	8	-	-
2001	51	77	110	164	138	99	47	8	-	-
2000	79	83	114	164	139	91	35	11	1	-
1999	52	94	166	148	153	87	26	7	-	-
Female										
2002	117	145	225	386	404	333	199	69	11	-
2001	96	161	253	376	416	337	175	54	9	-
2000	107	191	299	398	401	302	176	39	10	-
1999	119	182	288	407	417	325	199	47	7	-
Non-Hispanic Black										
Both sexes										
2002	32	29	19	28	14	12	2	2	1	-
2001	22	31	38	28	12	13	4	5	-	-
2000	22	31	27	21	15	12	11	4	-	-
1999	22	24	30	23	16	16	5	4	1	-
Male										
2002	9	12	8	8	3	2	-	-	1	-
2001	7	11	6	5	3	4	-	2	-	-
2000	5	8	8	7	1	3	3	2	-	-
1999	7	6	9	10	4	1	-	-	-	-
Female										
2002	23	17	11	20	11	10	2	2	-	-
2001	15	20	32	23	9	9	4	3	-	-
2000	17	23	19	14	14	9	8	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)														
Non-Hispanic Black														
Female														
1999	183	-	-	-	-	-	-	3	7	7	11	12	19	20
Origin not stated[2]														
Both sexes														
2002	6	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	6	-	-	-	-	1	-	-	-	-	-	1	1	-
2000	4	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	12	-	-	-	-	-	-	-	-	1	1	-	1	1
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	2	-	-	-	-	1	-	-	-	-	-	1	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	1	-	-	1	1
Female														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	7	-	-	-	-	-	-	-	-	-	1	-	-	-
Hypertensive heart disease (I11)														
All origins														
Both sexes														
2002	26,551	-	-	-	2	4	26	81	216	402	811	1,266	1,616	1,661
2001	24,689	-	-	1	-	6	18	65	181	352	739	1,050	1,414	1,380
2000	23,761	-	1	-	2	5	27	69	144	323	658	952	1,152	1,249
1999	22,702	1	-	-	-	6	27	50	160	347	617	889	1,041	1,260
Male														
2002	11,660	-	-	-	1	4	22	64	151	289	560	844	1,094	1,107
2001	10,451	-	-	-	-	4	13	40	129	234	506	715	931	879
2000	9,876	-	-	-	1	3	21	43	96	210	433	625	811	789
1999	9,316	-	-	-	-	6	19	41	108	232	428	586	673	800
Female														
2002	14,891	-	-	-	1	-	4	17	65	113	251	422	522	554
2001	14,238	-	-	1	-	2	5	25	52	118	233	335	483	501
2000	13,885	-	1	-	1	2	6	26	48	113	225	327	341	460
1999	13,386	1	-	-	-	-	8	9	52	115	189	303	368	460
Hispanic														
Both sexes														
2002	1,424	-	-	-	-	-	4	8	23	38	59	83	97	122

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
Non-Hispanic Black										
Female										
1999	15	18	21	13	12	15	5	4	1	-
Origin not stated[2]										
Both sexes										
2002	1	1	1	1	1	-	-	-	-	-
2001	1	-	-	1	-	1	-	-	-	-
2000	2	-	-	-	-	-	2	-	-	-
1999	1	2	2	2	-	1	-	-	-	-
Male										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	1	-	1	-	-	-	-	-	-	-
Female										
2002	-	1	1	1	1	-	-	-	-	-
2001	1	-	-	1	-	1	-	-	-	-
2000	1	-	-	-	-	-	2	-	-	-
1999	-	2	1	2	-	1	-	-	-	-
Hypertensive heart disease (I11)										
All origins										
Both sexes										
2002	1,659	1,807	2,088	2,816	3,505	3,716	3,115	1,442	314	4
2001	1,389	1,557	2,062	2,719	3,400	3,658	2,958	1,398	331	11
2000	1,423	1,570	2,035	2,694	3,268	3,684	2,886	1,330	286	3
1999	1,277	1,553	1,964	2,672	3,158	3,498	2,710	1,188	283	1
Male										
2002	1,052	1,046	1,042	1,259	1,250	1,004	629	194	44	4
2001	870	906	1,055	1,188	1,226	957	565	197	27	9
2000	895	909	1,062	1,135	1,134	960	545	181	21	2
1999	760	858	958	1,185	1,067	912	499	160	23	1
Female										
2002	607	761	1,046	1,557	2,255	2,712	2,486	1,248	270	-
2001	519	651	1,007	1,531	2,174	2,701	2,393	1,201	304	2
2000	528	661	973	1,559	2,134	2,724	2,341	1,149	265	1
1999	517	695	1,006	1,487	2,091	2,586	2,211	1,028	260	-
Hispanic										
Both sexes										
2002	119	130	134	162	146	153	90	43	13	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart disease (I11)														
Hispanic														
Both sexes														
2001	1,211	-	-	-	-	1	4	9	13	42	51	59	106	90
2000	991	-	-	-	-	-	1	9	11	26	38	55	63	56
1999	933	-	-	-	-	1	2	5	13	31	30	43	56	77
Male														
2002	781	-	-	-	-	-	4	7	18	28	50	65	70	84
2001	648	-	-	-	-	1	3	6	9	37	38	43	75	57
2000	495	-	-	-	-	-	1	9	7	18	25	42	42	37
1999	455	-	-	-	-	1	1	5	11	24	21	34	35	51
Female														
2002	643	-	-	-	-	-	-	1	5	10	9	18	27	38
2001	563	-	-	-	-	-	1	3	4	5	13	16	31	33
2000	496	-	-	-	-	-	-	-	4	8	13	13	21	19
1999	478	-	-	-	-	-	1	-	2	7	9	9	21	26
Non-Hispanic[1]														
Both sexes														
2002	24,976	-	-	-	2	4	22	72	193	362	747	1,174	1,505	1,529
2001	23,343	-	-	1	-	5	13	56	166	307	684	982	1,288	1,272
2000	22,646	-	1	-	2	5	26	60	133	295	610	888	1,080	1,185
1999	21,639	1	-	-	-	5	25	45	147	316	582	837	978	1,170
Male														
2002	10,790	-	-	-	1	4	18	57	133	259	506	770	1,014	1,017
2001	9,725	-	-	-	-	3	9	34	118	196	464	667	842	811
2000	9,298	-	-	-	1	3	20	34	89	190	401	576	762	745
1999	8,790	-	-	-	-	5	18	36	97	208	402	545	635	742
Female														
2002	14,186	-	-	-	1	-	4	15	60	103	241	404	491	512
2001	13,618	-	-	1	-	2	4	22	48	111	220	315	446	461
2000	13,348	-	1	-	1	2	6	26	44	105	209	312	318	440
1999	12,849	1	-	-	-	-	7	9	50	108	180	292	343	428
Non-Hispanic White														
Both sexes														
2002	18,060	-	-	-	1	1	9	31	79	186	401	666	853	912
2001	16,991	-	-	-	-	3	2	27	75	156	337	512	701	749
2000	16,373	-	-	-	2	1	13	22	53	147	329	462	580	661
1999	15,654	1	-	-	-	1	8	19	72	139	289	423	531	648
Male														
2002	7,427	-	-	-	1	1	8	25	58	137	299	470	618	644
2001	6,728	-	-	-	-	2	1	18	62	114	253	369	497	511
2000	6,390	-	-	-	1	1	10	13	40	107	234	336	435	444
1999	6,013	-	-	-	-	1	6	17	58	101	222	303	373	436

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)										
Hispanic										
Both sexes										
2001	94	106	133	133	127	115	78	41	9	-
2000	85	80	118	116	117	108	71	30	7	-
1999	69	95	79	103	112	99	80	30	8	-
Male										
2002	75	72	77	81	68	46	25	9	2	-
2001	63	63	69	57	51	42	24	7	3	-
2000	49	44	62	59	44	30	17	7	2	-
1999	45	52	43	44	38	24	18	6	2	-
Female										
2002	44	58	57	81	78	107	65	34	11	-
2001	31	43	64	76	76	73	54	34	6	-
2000	36	36	56	57	73	78	54	23	5	-
1999	24	43	36	59	74	75	62	24	6	-
Non-Hispanic[1]										
Both sexes										
2002	1,531	1,650	1,941	2,642	3,343	3,543	3,020	1,393	301	2
2001	1,282	1,447	1,917	2,571	3,266	3,530	2,872	1,354	321	9
2000	1,333	1,475	1,899	2,561	3,145	3,557	2,811	1,299	279	2
1999	1,193	1,446	1,869	2,559	3,032	3,388	2,619	1,151	275	1
Male										
2002	971	957	959	1,170	1,174	949	603	184	42	2
2001	797	841	974	1,125	1,174	910	539	190	24	7
2000	841	852	987	1,065	1,087	923	528	174	19	1
1999	703	798	904	1,135	1,022	884	480	154	21	1
Female										
2002	560	693	982	1,472	2,169	2,594	2,417	1,209	259	-
2001	485	606	943	1,446	2,092	2,620	2,333	1,164	297	2
2000	492	623	912	1,496	2,058	2,634	2,283	1,125	260	1
1999	490	648	965	1,424	2,010	2,504	2,139	997	254	-
Non-Hispanic White										
Both sexes										
2002	920	1,045	1,335	1,941	2,633	2,970	2,610	1,218	248	1
2001	771	898	1,300	1,913	2,627	2,952	2,528	1,183	252	5
2000	780	940	1,253	1,887	2,499	3,000	2,413	1,101	230	-
1999	723	904	1,243	1,868	2,426	2,874	2,280	982	223	-
Male										
2002	595	620	674	866	927	795	508	151	29	1
2001	505	551	686	848	933	740	468	155	11	4
2000	512	561	681	790	850	779	439	140	17	-
1999	446	504	618	835	812	730	411	123	17	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart disease (I11)														
Non-Hispanic White														
Female														
2002	10,633	-	-	-	-	-	1	6	21	49	102	196	235	268
2001	10,263	-	-	-	-	1	1	9	13	42	84	143	204	238
2000	9,983	-	-	-	1	-	3	9	13	40	95	126	145	217
1999	9,641	1	-	-	-	-	2	2	14	38	67	120	158	212
Non-Hispanic Black														
Both sexes														
2002	6,385	-	-	-	1	3	12	39	110	166	325	489	626	585
2001	5,880	-	-	1	-	2	11	28	86	147	327	451	556	493
2000	5,828	-	1	-	-	4	13	37	77	144	270	415	471	504
1999	5,556	-	-	-	-	3	17	26	72	169	278	403	435	491
Male														
2002	3,089	-	-	-	-	3	9	30	71	114	193	285	377	353
2001	2,760	-	-	-	-	1	8	15	52	79	196	282	327	281
2000	2,685	-	-	-	-	2	10	20	46	80	160	233	305	288
1999	2,571	-	-	-	-	3	12	19	36	101	166	235	252	287
Female														
2002	3,296	-	-	-	1	-	3	9	39	52	132	204	249	232
2001	3,120	-	-	1	-	1	3	13	34	68	131	169	229	212
2000	3,143	-	1	-	-	2	3	17	31	64	110	182	166	216
1999	2,985	-	-	-	-	-	5	7	36	68	112	168	183	204
Origin not stated[2]														
Both sexes														
2002	151	-	-	-	-	-	-	1	-	2	5	9	14	10
2001	135	-	-	-	-	-	1	-	2	3	4	9	20	18
2000	124	-	-	-	-	-	-	-	-	2	10	9	9	8
1999	130	-	-	-	-	-	-	-	-	-	5	9	7	13
Male														
2002	89	-	-	-	-	-	-	-	-	2	4	9	10	6
2001	78	-	-	-	-	-	1	-	2	1	4	5	14	11
2000	83	-	-	-	-	-	-	-	-	2	7	7	7	7
1999	71	-	-	-	-	-	-	-	-	-	5	7	3	7
Female														
2002	62	-	-	-	-	-	-	1	-	-	1	-	4	4
2001	57	-	-	-	-	-	-	-	-	2	-	4	6	7
2000	41	-	-	-	-	-	-	-	-	-	3	2	2	1
1999	59	-	-	-	-	-	-	-	-	-	-	2	4	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)										
Non-Hispanic White										
Female										
2002	325	425	661	1,075	1,706	2,175	2,102	1,067	219	-
2001	266	347	614	1,065	1,694	2,212	2,060	1,028	241	1
2000	268	379	572	1,097	1,649	2,221	1,974	961	213	-
1999	277	400	625	1,033	1,614	2,144	1,869	859	206	-
Non-Hispanic Black										
Both sexes										
2002	573	561	563	636	631	506	354	158	47	-
2001	479	510	568	599	565	520	313	155	65	4
2000	521	491	602	617	577	505	359	172	46	2
1999	441	508	576	632	550	455	299	154	46	1
Male										
2002	354	308	266	270	214	133	75	24	10	-
2001	270	273	265	246	212	148	62	28	12	3
2000	306	267	280	243	209	132	77	24	2	1
1999	241	274	262	270	188	131	60	30	3	1
Female										
2002	219	253	297	366	417	373	279	134	37	-
2001	209	237	303	353	353	372	251	127	53	1
2000	215	224	322	374	368	373	282	148	44	1
1999	200	234	314	362	362	324	239	124	43	-
Origin not stated[2]										
Both sexes										
2002	9	27	13	12	16	20	5	6	-	2
2001	13	4	12	15	7	13	8	3	1	2
2000	5	15	18	17	6	19	4	1	-	1
1999	15	12	16	10	14	11	11	7	-	-
Male										
2002	6	17	6	8	8	9	1	1	-	2
2001	10	2	12	6	1	5	2	-	-	2
2000	5	13	13	11	3	7	-	-	-	1
1999	12	8	11	6	7	4	1	-	-	-
Female										
2002	3	10	7	4	8	11	4	5	-	-
2001	3	2	-	9	6	8	6	3	1	-
2000	-	2	5	6	3	12	4	1	-	-
1999	3	4	5	4	7	7	10	7	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart and renal disease (I13)														
All origins														
Both sexes														
2002	2,895	-	-	-	-	1	4	11	17	28	49	75	90	134
2001	2,826	1	-	-	-	2	5	10	18	40	59	64	82	109
2000	2,785	-	-	-	1	1	5	11	17	22	61	68	79	103
1999	3,327	-	-	2	-	2	4	5	16	43	71	80	76	116
Male														
2002	1,205	-	-	-	-	-	3	9	10	19	25	49	56	82
2001	1,169	1	-	-	-	-	4	4	13	20	43	46	53	61
2000	1,163	-	-	-	-	1	3	7	11	12	37	43	44	57
1999	1,390	-	-	1	-	1	2	3	13	24	43	50	50	66
Female														
2002	1,690	-	-	-	-	1	1	2	7	9	24	26	34	52
2001	1,657	-	-	-	-	2	1	6	5	20	16	18	29	48
2000	1,622	-	-	-	1	-	2	4	6	10	24	25	35	46
1999	1,937	-	-	1	-	1	2	2	3	19	28	30	26	50
Hispanic														
Both sexes														
2002	172	-	-	-	-	1	1	2	1	1	3	7	4	10
2001	160	-	-	-	-	-	1	1	1	4	5	2	8	6
2000	138	-	-	-	-	-	1	3	-	1	7	6	8	6
1999	144	-	-	-	-	-	-	-	1	5	4	8	3	9
Male														
2002	80	-	-	-	-	-	1	1	-	1	1	2	3	7
2001	76	-	-	-	-	-	1	-	-	2	3	1	5	4
2000	67	-	-	-	-	-	1	2	-	1	3	3	4	2
1999	65	-	-	-	-	-	-	-	-	2	4	5	1	5
Female														
2002	92	-	-	-	-	1	-	1	1	-	2	5	1	3
2001	84	-	-	-	-	-	-	1	1	2	2	1	3	2
2000	71	-	-	-	-	-	-	1	-	-	4	3	4	4
1999	79	-	-	-	-	-	-	-	1	3	-	3	2	4
Non-Hispanic[1]														
Both sexes														
2002	2,714	-	-	-	-	-	3	9	16	27	46	68	85	124
2001	2,650	1	-	-	-	2	4	9	17	36	51	61	74	102
2000	2,637	-	-	-	1	1	4	8	17	21	54	62	71	96
1999	3,171	-	-	2	-	2	4	5	15	37	65	69	73	107
Male														
2002	1,120	-	-	-	-	-	2	8	10	18	24	47	52	75
2001	1,086	1	-	-	-	-	3	4	13	18	38	44	48	56

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)										
All origins										
Both sexes										
2002	139	156	246	363	503	511	386	150	32	-
2001	138	169	248	374	481	499	348	155	24	-
2000	105	149	268	376	487	494	361	153	24	-
1999	146	201	269	487	600	600	411	162	36	-
Male										
2002	76	67	131	158	199	171	108	38	4	-
2001	65	94	112	158	181	174	102	35	3	-
2000	51	74	132	152	198	188	109	37	7	-
1999	80	99	124	221	239	205	123	41	5	-
Female										
2002	63	89	115	205	304	340	278	112	28	-
2001	73	75	136	216	300	325	246	120	21	-
2000	54	75	136	224	289	306	252	116	17	-
1999	66	102	145	266	361	395	288	121	31	-
Hispanic										
Both sexes										
2002	16	16	18	25	26	16	19	6	-	-
2001	15	15	19	18	23	15	16	11	-	-
2000	9	7	20	16	16	20	12	6	-	-
1999	10	17	15	20	19	17	12	3	1	-
Male										
2002	11	6	12	9	11	8	4	3	-	-
2001	10	10	7	4	13	6	5	5	-	-
2000	3	3	16	11	7	7	3	1	-	-
1999	7	6	8	8	6	10	1	1	1	-
Female										
2002	5	10	6	16	15	8	15	3	-	-
2001	5	5	12	14	10	9	11	6	-	-
2000	6	4	4	5	9	13	9	5	-	-
1999	3	11	7	12	13	7	11	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	123	139	226	336	476	493	367	144	32	-
2001	123	153	227	355	455	482	331	143	24	-
2000	96	141	248	357	469	473	349	146	23	-
1999	135	183	253	466	581	581	399	159	35	-
Male										
2002	65	61	117	149	187	162	104	35	4	-
2001	55	84	104	154	167	167	97	30	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart and renal disease (I13)														
Non-Hispanic[1]														
Male														
2000	1,089	-	-	-	-	1	2	5	11	11	34	40	40	54
1999	1,318	-	-	1	-	1	2	3	13	21	38	42	49	61
Female														
2002	1,594	-	-	-	-	-	1	1	6	9	22	21	33	49
2001	1,564	-	-	-	-	2	1	5	4	18	13	17	26	46
2000	1,548	-	-	-	1	-	2	3	6	10	20	22	31	42
1999	1,853	-	-	1	-	1	2	2	2	16	27	27	24	46
Non-Hispanic White														
Both sexes														
2002	1,752	-	-	-	-	-	-	4	4	4	13	12	17	35
2001	1,759	-	-	-	-	-	2	1	4	3	13	16	27	39
2000	1,769	-	-	-	-	-	-	3	2	8	22	15	25	40
1999	2,091	-	-	1	-	-	-	1	5	8	11	15	28	38
Male														
2002	679	-	-	-	-	-	-	3	2	4	9	11	8	23
2001	687	-	-	-	-	-	2	1	2	1	11	11	16	20
2000	710	-	-	-	-	-	-	2	2	4	14	9	15	23
1999	843	-	-	1	-	-	-	1	5	4	7	8	18	21
Female														
2002	1,073	-	-	-	-	-	-	1	2	-	4	1	9	12
2001	1,072	-	-	-	-	-	-	-	2	2	2	5	11	19
2000	1,059	-	-	-	-	-	-	1	-	4	8	6	10	17
1999	1,248	-	-	-	-	-	-	-	-	4	4	7	10	17
Non-Hispanic Black														
Both sexes														
2002	897	-	-	-	-	-	2	5	12	22	32	56	68	88
2001	821	1	-	-	-	2	2	8	13	29	36	45	47	61
2000	797	-	-	-	1	1	4	4	15	13	32	45	46	54
1999	989	-	-	1	-	2	4	3	10	28	54	52	40	66
Male														
2002	402	-	-	-	-	-	1	5	8	13	14	36	44	51
2001	370	1	-	-	-	-	1	3	11	15	26	33	32	35
2000	348	-	-	-	-	1	2	3	9	7	20	30	25	30
1999	426	-	-	-	-	1	2	1	8	16	31	32	28	37
Female														
2002	495	-	-	-	-	-	1	-	4	9	18	20	24	37
2001	451	-	-	-	-	2	1	5	2	14	10	12	15	26
2000	449	-	-	-	1	-	2	1	6	6	12	15	21	24

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)										
Non-Hispanic[1]										
Male										
2000	48	70	116	140	189	180	106	36	6	-
1999	73	92	116	213	233	194	122	40	4	-
Female										
2002	58	78	109	187	289	331	263	109	28	-
2001	68	69	123	201	288	315	234	113	21	-
2000	48	71	132	217	280	293	243	110	17	-
1999	62	91	137	253	348	387	277	119	31	-
Non-Hispanic White										
Both sexes										
2002	47	57	127	228	376	383	300	121	24	-
2001	56	71	133	241	360	379	278	120	16	-
2000	35	65	155	249	344	378	294	122	12	-
1999	53	83	132	324	444	455	339	126	28	-
Male										
2002	24	26	70	107	149	126	86	29	2	-
2001	22	40	61	110	140	139	85	24	2	-
2000	18	35	74	111	145	142	87	27	2	-
1999	27	47	61	153	193	156	107	30	4	-
Female										
2002	23	31	57	121	227	257	214	92	22	-
2001	34	31	72	131	220	240	193	96	14	-
2000	17	30	81	138	199	236	207	95	10	-
1999	26	36	71	171	251	299	232	96	24	-
Non-Hispanic Black										
Both sexes										
2002	71	74	91	100	87	99	60	22	8	-
2001	64	78	86	104	83	83	51	21	7	-
2000	58	74	87	94	109	82	49	19	10	-
1999	77	91	114	121	122	113	52	32	7	-
Male										
2002	37	29	41	37	29	32	17	6	2	-
2001	31	42	39	39	23	22	11	5	1	-
2000	27	35	39	26	39	29	16	6	4	-
1999	44	41	51	49	34	29	13	9	-	-
Female										
2002	34	45	50	63	58	67	43	16	6	-
2001	33	36	47	65	60	61	40	16	6	-
2000	31	39	48	68	70	53	33	13	6	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hypertensive heart and renal disease (I13)														
Non-Hispanic Black														
Female														
1999	563	-	-	1	-	1	2	2	2	12	23	20	12	29
Origin not stated[2]														
Both sexes														
2002	9	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	16	-	-	-	-	-	-	-	-	-	3	1	-	1
2000	10	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	12	-	-	-	-	-	-	-	-	1	2	3	-	-
Male														
2002	5	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	7	-	-	-	-	-	-	-	-	-	2	1	-	1
2000	7	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	7	-	-	-	-	-	-	-	-	1	1	3	-	-
Female														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	9	-	-	-	-	-	-	-	-	-	1	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	-	1	-	-	-
Ischemic heart diseases (I20-I25)														
All origins														
Both sexes														
2002	494,382	23	7	4	16	30	103	251	745	2,211	5,472	9,892	15,683	20,997
2001	502,189	26	9	4	16	31	132	279	837	2,212	5,322	9,461	15,612	20,270
2000	515,204	30	10	8	8	33	90	236	699	2,233	5,286	9,638	15,252	20,724
1999	529,659	27	5	13	9	52	85	245	769	2,387	5,364	9,524	15,251	20,575
Male														
2002	252,760	15	4	1	12	18	74	191	550	1,670	4,170	7,582	11,969	15,350
2001	254,005	17	3	2	11	24	95	211	619	1,667	4,045	7,323	11,818	14,794
2000	260,574	17	7	5	4	21	60	181	523	1,665	4,122	7,601	11,548	15,154
1999	267,268	13	3	10	7	36	57	169	570	1,801	4,169	7,447	11,614	15,021
Female														
2002	241,622	8	3	3	4	12	29	60	195	541	1,302	2,310	3,714	5,647
2001	248,184	9	6	2	5	7	37	68	218	545	1,277	2,138	3,794	5,476
2000	254,630	13	3	3	4	12	30	55	176	568	1,164	2,037	3,704	5,570
1999	262,391	14	2	3	2	16	28	76	199	586	1,195	2,077	3,637	5,554
Hispanic														
Both sexes														
2002	20,941	4	1	1	2	4	14	29	59	144	298	557	832	1,137

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)										
Non-Hispanic Black										
Female										
1999	33	50	63	72	88	84	39	23	7	-
Origin not stated[2]										
Both sexes										
2002	-	1	2	2	1	2	-	-	-	-
2001	-	1	2	1	3	2	1	1	-	-
2000	-	1	-	3	2	1	-	1	1	-
1999	1	1	1	1	-	2	-	-	-	-
Male										
2002	-	-	2	-	1	1	-	-	-	-
2001	-	-	1	-	1	1	-	-	-	-
2000	-	1	-	1	2	1	-	-	1	-
1999	-	1	-	-	-	1	-	-	-	-
Female										
2002	-	1	-	2	-	1	-	-	-	-
2001	-	1	1	1	2	1	1	1	-	-
2000	-	-	-	2	-	-	-	1	-	-
1999	1	-	1	1	-	1	-	-	-	-
Ischemic heart diseases (I20-I25)										
All origins										
Both sexes										
2002	26,265	34,336	49,531	69,769	85,622	84,082	59,567	24,211	5,528	37
2001	26,304	35,502	51,961	73,082	86,772	84,880	59,558	24,206	5,665	48
2000	27,013	37,257	55,628	76,569	87,123	86,890	60,269	24,477	5,683	48
1999	27,802	40,080	58,760	79,379	89,917	88,158	60,502	24,983	5,733	39
Male										
2002	18,306	22,573	30,563	39,154	42,085	34,073	18,199	5,353	821	27
2001	18,212	23,181	31,891	40,894	42,059	33,466	17,702	5,124	814	33
2000	18,641	24,225	34,329	42,452	42,295	34,024	17,603	5,216	847	34
1999	19,102	26,184	36,003	44,041	43,264	34,168	17,472	5,245	841	31
Female										
2002	7,959	11,763	18,968	30,615	43,537	50,009	41,368	18,858	4,707	10
2001	8,092	12,321	20,070	32,188	44,713	51,414	41,856	19,082	4,851	15
2000	8,372	13,032	21,299	34,117	44,828	52,866	42,666	19,261	4,836	14
1999	8,700	13,896	22,757	35,338	46,653	53,990	43,030	19,738	4,892	8
Hispanic										
Both sexes										
2002	1,546	1,928	2,582	3,022	3,038	2,662	1,970	926	184	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Ischemic heart diseases (I20-I25)														
Hispanic														
Both sexes														
2001	20,664	4	1	-	4	5	13	32	77	128	240	495	819	1,033
2000	19,744	6	1	-	1	2	10	20	49	121	271	486	754	1,055
1999	19,936	2	-	2	-	6	5	24	63	128	238	502	826	1,018
Male														
2002	11,295	2	-	1	1	1	11	26	50	107	239	433	625	846
2001	11,019	3	1	-	1	4	9	27	63	105	190	398	618	764
2000	10,562	4	1	-	1	2	7	13	43	98	224	397	576	765
1999	10,604	-	-	2	-	5	3	23	49	112	198	391	651	740
Female														
2002	9,646	2	1	-	1	3	3	3	9	37	59	124	207	291
2001	9,645	1	-	-	3	1	4	5	14	23	50	97	201	269
2000	9,182	2	-	-	-	-	3	7	6	23	47	89	178	290
1999	9,332	2	-	-	-	1	2	1	14	16	40	111	175	278
Non-Hispanic[1]														
Both sexes														
2002	471,655	19	6	3	14	26	88	220	680	2,059	5,155	9,300	14,769	19,768
2001	479,842	21	8	4	12	26	118	247	757	2,077	5,058	8,932	14,731	19,141
2000	493,541	23	9	8	7	31	80	214	647	2,100	4,982	9,105	14,418	19,570
1999	507,700	25	5	11	9	45	80	220	706	2,245	5,099	8,980	14,351	19,458
Male														
2002	240,462	13	4	-	11	17	62	163	495	1,557	3,918	7,125	11,275	14,425
2001	242,054	13	2	2	10	20	85	184	553	1,558	3,834	6,896	11,151	13,952
2000	248,932	12	6	5	3	19	53	167	478	1,558	3,868	7,167	10,907	14,315
1999	255,524	13	3	8	7	31	54	145	521	1,677	3,946	7,021	10,907	14,202
Female														
2002	231,193	6	2	3	3	9	26	57	185	502	1,237	2,175	3,494	5,343
2001	237,788	8	6	2	2	6	33	63	204	519	1,224	2,036	3,580	5,189
2000	244,609	11	3	3	4	12	27	47	169	542	1,114	1,938	3,511	5,255
1999	252,176	12	2	3	2	14	26	75	185	568	1,153	1,959	3,444	5,256
Non-Hispanic White														
Both sexes														
2002	413,230	11	4	2	10	20	56	154	496	1,557	4,008	7,219	11,580	15,910
2001	420,959	14	5	3	7	17	69	167	551	1,582	3,874	6,918	11,600	15,364
2000	434,505	16	9	6	6	26	54	138	468	1,619	3,818	7,064	11,467	15,772
1999	447,960	16	4	7	5	31	45	155	510	1,724	3,883	7,017	11,408	15,660
Male														
2002	211,391	8	2	-	8	13	41	121	372	1,201	3,145	5,736	9,120	11,901
2001	212,928	10	1	2	6	12	53	130	409	1,208	3,042	5,546	9,044	11,484
2000	219,732	9	6	3	3	16	36	110	363	1,234	3,085	5,778	8,945	11,787
1999	225,909	8	3	5	5	21	36	100	382	1,310	3,106	5,680	8,905	11,699

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)										
Hispanic										
Both sexes										
2001	1,371	1,933	2,620	3,113	2,982	2,783	1,981	830	196	4
2000	1,402	1,873	2,491	2,854	2,772	2,706	1,914	783	169	4
1999	1,445	1,971	2,521	2,937	2,718	2,685	1,852	828	160	5
Male										
2002	1,061	1,221	1,554	1,646	1,453	1,083	642	251	41	1
2001	903	1,247	1,557	1,683	1,433	1,120	639	206	45	3
2000	961	1,167	1,494	1,573	1,315	1,052	616	211	39	3
1999	937	1,252	1,518	1,629	1,240	1,000	583	234	32	5
Female										
2002	485	707	1,028	1,376	1,585	1,579	1,328	675	143	-
2001	468	686	1,063	1,430	1,549	1,663	1,342	624	151	1
2000	441	706	997	1,281	1,457	1,654	1,298	572	130	1
1999	508	719	1,003	1,308	1,478	1,685	1,269	594	128	-
Non-Hispanic[1]										
Both sexes										
2002	24,591	32,256	46,731	66,495	82,339	81,150	57,423	23,213	5,321	29
2001	24,805	33,437	49,167	69,719	83,520	81,853	57,426	23,298	5,455	30
2000	25,452	35,235	52,902	73,441	84,082	83,895	58,189	23,617	5,501	33
1999	26,208	37,948	55,971	76,140	86,895	85,197	58,459	24,071	5,551	26
Male										
2002	17,156	21,251	28,865	37,356	40,510	32,871	17,507	5,086	776	19
2001	17,208	21,840	30,219	39,063	40,486	32,260	17,031	4,900	765	22
2000	17,565	22,946	32,677	40,716	40,862	32,863	16,934	4,984	804	23
1999	18,047	24,820	34,297	42,234	41,893	33,046	16,835	4,991	806	20
Female										
2002	7,435	11,005	17,866	29,139	41,829	48,279	39,916	18,127	4,545	10
2001	7,597	11,597	18,948	30,656	43,034	49,593	40,395	18,398	4,690	8
2000	7,887	12,289	20,225	32,725	43,220	51,032	41,255	18,633	4,697	10
1999	8,161	13,128	21,674	33,906	45,002	52,151	41,624	19,080	4,745	6
Non-Hispanic White										
Both sexes										
2002	19,889	26,553	40,000	58,243	74,305	74,501	52,858	21,199	4,635	20
2001	20,057	27,459	42,222	61,307	75,508	75,163	53,018	21,266	4,770	18
2000	20,595	29,352	45,515	64,841	76,270	77,010	53,898	21,670	4,867	24
1999	21,406	31,841	48,288	67,511	79,083	78,186	54,124	22,094	4,947	15
Male										
2002	14,207	17,941	25,264	33,367	37,003	30,493	16,159	4,618	658	13
2001	14,292	18,431	26,450	34,963	37,073	29,942	15,720	4,445	651	14
2000	14,611	19,650	28,724	36,526	37,516	30,438	15,690	4,499	686	17
1999	15,055	21,341	30,212	38,098	38,665	30,521	15,504	4,538	704	11

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Ischemic heart diseases (I20-I25)														
Non-Hispanic White														
Female														
2002	201,839	3	2	2	2	7	15	33	124	356	863	1,483	2,460	4,009
2001	208,031	4	4	1	1	5	16	37	142	374	832	1,372	2,556	3,880
2000	214,773	7	3	3	3	10	18	28	105	385	733	1,286	2,522	3,985
1999	222,051	8	1	2	-	10	9	55	128	414	777	1,337	2,503	3,961
Non-Hispanic Black														
Both sexes														
2002	49,522	7	2	1	4	4	27	55	155	430	1,011	1,835	2,830	3,399
2001	50,295	7	3	1	4	9	43	72	174	434	1,040	1,772	2,761	3,302
2000	50,659	4	-	2	-	4	20	66	163	426	1,025	1,824	2,594	3,340
1999	51,368	8	1	4	4	12	32	53	160	466	1,086	1,748	2,614	3,359
Male														
2002	23,976	4	2	-	3	2	16	34	97	293	654	1,189	1,864	2,186
2001	24,255	3	1	-	3	8	27	48	119	297	677	1,156	1,826	2,098
2000	24,289	2	-	2	-	2	12	48	102	277	669	1,208	1,678	2,192
1999	24,736	4	-	3	2	8	16	38	110	317	725	1,160	1,744	2,173
Female														
2002	25,546	3	-	1	1	2	11	21	58	137	357	646	966	1,213
2001	26,040	4	2	1	1	1	16	24	55	137	363	616	935	1,204
2000	26,370	2	-	-	-	2	8	18	61	149	356	616	916	1,148
1999	26,632	4	1	1	2	4	16	15	50	149	361	588	870	1,186
Origin not stated[2]														
Both sexes														
2002	1,786	-	-	-	-	-	1	2	6	8	19	35	82	92
2001	1,683	1	-	-	-	-	1	-	3	7	24	34	62	96
2000	1,919	1	-	-	-	-	-	2	3	12	33	47	80	99
1999	2,023	-	-	-	-	1	-	1	-	14	27	42	74	99
Male														
2002	1,003	-	-	-	-	-	1	2	5	6	13	24	69	79
2001	932	1	-	-	-	-	1	-	3	4	21	29	49	78
2000	1,080	1	-	-	-	-	-	1	2	9	30	37	65	74
1999	1,140	-	-	-	-	-	-	1	-	12	25	35	56	79
Female														
2002	783	-	-	-	-	-	-	-	1	2	6	11	13	13
2001	751	-	-	-	-	-	-	-	-	3	3	5	13	18
2000	839	-	-	-	-	-	-	1	1	3	3	10	15	25
1999	883	-	-	-	-	1	-	-	-	2	2	7	18	20

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)										
Non-Hispanic White										
Female										
2002	5,682	8,612	14,736	24,876	37,302	44,008	36,699	16,581	3,977	7
2001	5,765	9,028	15,772	26,344	38,435	45,221	37,298	16,821	4,119	4
2000	5,984	9,702	16,791	28,315	38,754	46,572	38,208	17,171	4,181	7
1999	6,351	10,500	18,076	29,413	40,418	47,665	38,620	17,556	4,243	4
Non-Hispanic Black										
Both sexes										
2002	4,105	4,864	5,734	6,871	6,670	5,456	3,758	1,699	596	9
2001	4,149	5,151	5,956	7,097	6,685	5,555	3,717	1,756	597	10
2000	4,247	5,070	6,415	7,217	6,592	5,809	3,610	1,655	567	9
1999	4,183	5,331	6,598	7,306	6,584	5,862	3,662	1,743	541	11
Male										
2002	2,536	2,788	3,023	3,213	2,800	1,817	1,006	356	87	6
2001	2,498	2,888	3,201	3,358	2,757	1,796	1,032	358	97	7
2000	2,521	2,759	3,362	3,404	2,714	1,912	949	366	104	6
1999	2,564	2,996	3,441	3,380	2,611	1,971	1,011	368	85	9
Female										
2002	1,569	2,076	2,711	3,658	3,870	3,639	2,752	1,343	509	3
2001	1,651	2,263	2,755	3,739	3,928	3,759	2,685	1,398	500	3
2000	1,726	2,311	3,053	3,813	3,878	3,897	2,661	1,289	463	3
1999	1,619	2,335	3,157	3,926	3,973	3,891	2,651	1,375	456	2
Origin not stated[2]										
Both sexes										
2002	128	152	218	252	245	270	174	72	23	7
2001	128	132	174	250	270	244	151	78	14	14
2000	159	149	235	274	269	289	166	77	13	11
1999	149	161	268	302	304	276	191	84	22	8
Male										
2002	89	101	144	152	122	119	50	16	4	7
2001	101	94	115	148	140	86	32	18	4	8
2000	115	112	158	163	118	109	53	21	4	8
1999	118	112	188	178	131	122	54	20	3	6
Female										
2002	39	51	74	100	123	151	124	56	19	-
2001	27	38	59	102	130	158	119	60	10	6
2000	44	37	77	111	151	180	113	56	9	3
1999	31	49	80	124	173	154	137	64	19	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute myocardial infarction (I21-I22)														
All origins														
Both sexes														
2002	179,514	11	5	2	5	15	47	91	309	882	2,279	4,055	6,564	8,758
2001	184,757	15	2	1	6	15	49	105	370	924	2,252	4,097	6,720	8,704
2000	192,898	20	1	5	1	14	38	108	267	904	2,228	4,230	6,647	9,030
1999	199,454	13	2	5	6	25	35	101	333	935	2,314	4,225	6,598	9,123
Male														
2002	93,830	7	2	-	2	10	31	67	225	656	1,691	3,066	4,929	6,287
2001	95,875	10	2	-	4	14	34	69	264	691	1,693	3,146	5,012	6,271
2000	100,306	10	1	3	1	8	22	75	194	657	1,730	3,319	4,946	6,490
1999	104,169	6	1	3	4	16	24	68	238	707	1,763	3,261	5,039	6,654
Female														
2002	85,684	4	3	2	3	5	16	24	84	226	588	989	1,635	2,471
2001	88,882	5	-	1	2	1	15	36	106	233	559	951	1,708	2,433
2000	92,592	10	-	2	-	6	16	33	73	247	498	911	1,701	2,540
1999	95,285	7	1	2	2	9	11	33	95	228	551	964	1,559	2,469
Hispanic														
Both sexes														
2002	7,631	3	1	-	1	3	4	9	25	59	118	198	320	456
2001	7,403	4	1	-	1	3	3	13	33	45	94	201	302	381
2000	7,106	5	-	-	-	-	5	12	19	47	107	196	290	400
1999	7,158	2	-	-	-	2	2	10	22	46	96	216	312	352
Male														
2002	4,125	2	-	-	-	1	4	7	22	39	87	151	231	332
2001	3,972	3	1	-	-	3	3	10	27	34	76	166	217	267
2000	3,825	3	-	-	-	-	4	6	18	35	90	156	207	286
1999	3,836	-	-	-	-	2	1	10	17	38	78	159	232	248
Female														
2002	3,506	1	1	-	1	2	-	2	3	20	31	47	89	124
2001	3,431	1	-	-	1	-	-	3	6	11	18	35	85	114
2000	3,281	2	-	-	-	-	1	6	1	12	17	40	83	114
1999	3,322	2	-	-	-	-	1	-	5	8	18	57	80	104
Non-Hispanic[1]														
Both sexes														
2002	171,441	8	4	2	4	12	43	82	283	822	2,153	3,848	6,228	8,286
2001	176,895	11	1	1	5	12	46	92	336	877	2,146	3,885	6,409	8,298
2000	185,283	14	1	5	1	14	33	96	248	853	2,110	4,027	6,336	8,598
1999	191,729	11	2	5	6	23	33	91	311	883	2,207	3,996	6,268	8,730
Male														
2002	89,480	5	2	-	2	9	27	60	202	617	1,598	2,910	4,687	5,943
2001	91,655	7	1	-	4	11	31	59	236	655	1,606	2,971	4,787	5,984
2000	96,218	6	1	3	1	8	18	69	176	619	1,631	3,160	4,726	6,184

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)										
All origins										
Both sexes										
2002	11,175	14,230	19,773	26,154	30,436	28,423	18,352	6,623	1,320	5
2001	11,126	14,690	20,690	27,766	31,488	29,242	18,525	6,606	1,354	10
2000	11,983	15,898	22,561	30,059	32,151	29,922	18,701	6,755	1,367	8
1999	12,283	17,333	24,108	31,053	33,658	30,711	18,472	6,765	1,349	7
Male										
2002	7,679	9,214	12,050	14,316	14,603	11,478	5,741	1,559	213	4
2001	7,587	9,446	12,517	15,151	14,974	11,411	5,778	1,576	219	6
2000	8,148	10,177	13,646	16,295	15,336	11,765	5,701	1,539	238	5
1999	8,375	11,154	14,554	16,991	15,952	12,066	5,566	1,523	202	2
Female										
2002	3,496	5,016	7,723	11,838	15,833	16,945	12,611	5,064	1,107	1
2001	3,539	5,244	8,173	12,615	16,514	17,831	12,747	5,030	1,135	4
2000	3,835	5,721	8,915	13,764	16,815	18,157	13,000	5,216	1,129	3
1999	3,908	6,179	9,554	14,062	17,706	18,645	12,906	5,242	1,147	5
Hispanic										
Both sexes										
2002	633	737	989	1,117	1,067	950	620	270	51	-
2001	533	715	986	1,161	1,029	962	626	245	63	2
2000	541	712	949	1,084	965	896	615	220	43	-
1999	567	750	949	1,131	951	936	548	225	41	-
Male										
2002	423	460	583	600	521	397	182	71	12	-
2001	336	458	578	599	494	386	227	72	14	1
2000	357	437	547	590	457	358	199	64	11	-
1999	359	470	558	655	417	325	194	64	9	-
Female										
2002	210	277	406	517	546	553	438	199	39	-
2001	197	257	408	562	535	576	399	173	49	1
2000	184	275	402	494	508	538	416	156	32	-
1999	208	280	391	476	534	611	354	161	32	-
Non-Hispanic[1]										
Both sexes										
2002	10,515	13,456	18,716	24,985	29,300	27,407	17,685	6,334	1,263	5
2001	10,546	13,946	19,658	26,537	30,386	28,202	17,858	6,347	1,289	7
2000	11,402	15,146	21,549	28,908	31,112	28,944	18,040	6,518	1,321	7
1999	11,679	16,535	23,079	29,839	32,627	29,701	17,871	6,526	1,302	4
Male										
2002	7,240	8,733	11,428	13,685	14,051	11,046	5,548	1,483	200	4
2001	7,218	8,968	11,907	14,514	14,442	11,004	5,541	1,500	204	5
2000	7,766	9,712	13,060	15,666	14,850	11,377	5,485	1,471	225	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute myocardial infarction (I21-I22)														
Non-Hispanic[1]														
Male														
1999	100,014	6	1	3	4	14	23	58	221	664	1,674	3,092	4,795	6,374
Female														
2002	81,961	3	2	2	2	3	16	22	81	205	555	938	1,541	2,343
2001	85,240	4	-	1	1	1	15	33	100	222	540	914	1,622	2,314
2000	89,065	8	-	2	-	6	15	27	72	234	479	867	1,610	2,414
1999	91,715	5	1	2	2	9	10	33	90	219	533	904	1,473	2,356
Non-Hispanic White														
Both sexes														
2002	150,051	6	4	1	3	10	26	54	213	639	1,706	3,048	5,046	6,883
2001	155,312	8	1	-	1	9	27	59	254	684	1,683	3,117	5,185	6,865
2000	163,203	10	1	3	1	11	23	59	184	694	1,642	3,249	5,161	7,157
1999	169,545	7	2	3	3	17	18	66	227	710	1,736	3,229	5,175	7,279
Male														
2002	79,037	4	2	-	2	7	18	43	157	493	1,300	2,395	3,911	5,066
2001	81,175	6	1	-	1	8	20	37	182	520	1,314	2,485	4,006	5,095
2000	85,412	5	1	1	1	6	14	44	139	519	1,320	2,643	3,992	5,261
1999	89,142	3	1	1	3	11	15	41	161	546	1,361	2,591	4,080	5,420
Female														
2002	71,014	2	2	1	1	3	8	11	56	146	406	653	1,135	1,817
2001	74,137	2	-	-	-	1	7	22	72	164	369	632	1,179	1,770
2000	77,791	5	-	2	-	5	9	15	45	175	322	606	1,169	1,896
1999	80,403	4	1	2	-	6	3	25	66	164	375	638	1,095	1,859
Non-Hispanic Black														
Both sexes														
2002	18,162	2	-	1	1	-	15	25	63	157	399	702	1,054	1,248
2001	18,399	3	-	1	3	3	18	28	69	168	404	681	1,059	1,254
2000	18,934	2	-	2	-	2	9	32	61	145	415	691	1,029	1,263
1999	19,120	4	-	2	3	6	13	22	66	159	421	685	968	1,262
Male														
2002	8,607	1	-	-	-	-	7	15	39	102	257	435	674	764
2001	8,628	1	-	-	2	3	10	18	44	110	244	415	653	761
2000	8,959	1	-	2	-	1	4	21	34	89	266	445	623	789
1999	9,106	3	-	2	1	3	7	16	45	104	270	434	621	812
Female														
2002	9,555	1	-	1	1	-	8	10	24	55	142	267	380	484
2001	9,771	2	-	1	1	-	8	10	25	58	160	266	406	493
2000	9,975	1	-	-	-	1	5	11	27	56	149	246	406	474
1999	10,014	1	-	-	2	3	6	6	21	55	151	251	347	450

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)										
Non-Hispanic[1]										
Male										
1999	7,989	10,658	13,940	16,288	15,497	11,711	5,355	1,454	192	1
Female										
2002	3,275	4,723	7,288	11,300	15,249	16,361	12,137	4,851	1,063	1
2001	3,328	4,978	7,751	12,023	15,944	17,198	12,317	4,847	1,085	2
2000	3,636	5,434	8,489	13,242	16,262	17,567	12,555	5,047	1,096	3
1999	3,690	5,877	9,139	13,551	17,130	17,990	12,516	5,072	1,110	3
Non-Hispanic White										
Both sexes										
2002	8,741	11,314	16,153	21,961	26,356	24,999	16,126	5,698	1,060	4
2001	8,812	11,690	17,059	23,472	27,436	25,780	16,332	5,728	1,106	4
2000	9,521	12,841	18,727	25,620	28,220	26,433	16,629	5,882	1,129	6
1999	9,837	14,202	20,137	26,593	29,696	27,116	16,434	5,930	1,126	2
Male										
2002	6,200	7,547	10,126	12,260	12,776	10,161	5,074	1,330	162	3
2001	6,208	7,742	10,560	13,064	13,218	10,137	5,058	1,337	172	4
2000	6,658	8,465	11,597	14,129	13,620	10,464	5,037	1,296	197	3
1999	6,899	9,342	12,449	14,792	14,312	10,743	4,907	1,307	157	-
Female										
2002	2,541	3,767	6,027	9,701	13,580	14,838	11,052	4,368	898	1
2001	2,604	3,948	6,499	10,408	14,218	15,643	11,274	4,391	934	-
2000	2,863	4,376	7,130	11,491	14,600	15,969	11,592	4,586	932	3
1999	2,938	4,860	7,688	11,801	15,384	16,373	11,527	4,623	969	2
Non-Hispanic Black										
Both sexes										
2002	1,552	1,810	2,183	2,508	2,431	1,982	1,279	570	179	1
2001	1,498	1,935	2,238	2,585	2,454	2,007	1,292	539	157	3
2000	1,647	1,969	2,446	2,734	2,455	2,126	1,191	536	178	1
1999	1,600	2,011	2,531	2,751	2,505	2,164	1,248	538	159	2
Male										
2002	887	987	1,099	1,150	1,004	684	353	123	25	1
2001	845	1,028	1,135	1,173	982	671	372	132	28	1
2000	942	1,032	1,247	1,239	1,008	717	343	128	27	1
1999	931	1,113	1,262	1,236	975	758	355	127	30	1
Female										
2002	665	823	1,084	1,358	1,427	1,298	926	447	154	-
2001	653	907	1,103	1,412	1,472	1,336	920	407	129	2
2000	705	937	1,199	1,495	1,447	1,409	848	408	151	-
1999	669	898	1,269	1,515	1,530	1,406	893	411	129	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute myocardial infarction (I21-I22)														
Origin not stated[2]														
Both sexes														
2002	442	-	-	-	-	-	-	-	1	1	8	9	16	16
2001	459	-	-	-	-	-	-	-	1	2	12	11	9	25
2000	509	1	-	-	-	-	-	-	-	4	11	7	21	32
1999	567	-	-	-	-	-	-	-	-	6	11	13	18	41
Male														
2002	225	-	-	-	-	-	-	-	1	-	6	5	11	12
2001	248	-	-	-	-	-	-	-	1	2	11	9	8	20
2000	263	1	-	-	-	-	-	-	-	3	9	3	13	20
1999	319	-	-	-	-	-	-	-	-	5	11	10	12	32
Female														
2002	217	-	-	-	-	-	-	-	-	1	2	4	5	4
2001	211	-	-	-	-	-	-	-	-	-	1	2	1	5
2000	246	-	-	-	-	-	-	-	-	1	2	4	8	12
1999	248	-	-	-	-	-	-	-	-	1	-	3	6	9
Other acute ischemic heart diseases (I24)														
All origins														
Both sexes														
2002	3,407	2	-	1	-	2	2	4	14	17	60	119	226	210
2001	3,351	3	-	-	2	-	7	2	12	32	62	120	187	200
2000	3,363	1	1	-	-	4	4	4	12	30	62	93	189	219
1999	3,232	3	1	1	-	-	3	2	4	25	56	92	160	209
Male														
2002	1,767	1	-	1	-	-	1	2	10	9	41	88	162	153
2001	1,720	2	-	-	1	-	6	2	8	18	47	89	141	134
2000	1,749	1	1	-	-	3	3	3	10	24	47	76	141	163
1999	1,665	1	1	1	-	-	3	1	3	19	44	69	109	159
Female														
2002	1,640	1	-	-	-	2	1	2	4	8	19	31	64	57
2001	1,631	1	-	-	1	-	1	-	4	14	15	31	46	66
2000	1,614	-	-	-	-	1	1	1	2	6	15	17	48	56
1999	1,567	2	-	-	-	-	-	1	1	6	12	23	51	50
Hispanic														
Both sexes														
2002	64	-	-	1	-	1	-	1	-	-	1	5	5	4
2001	63	-	-	-	-	-	-	-	-	-	1	1	3	6
2000	69	-	1	-	-	1	-	1	-	-	1	4	2	6
1999	53	-	-	1	-	-	-	-	-	-	1	1	6	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)										
Origin not stated[2]										
Both sexes										
2002	27	37	68	52	69	66	47	19	6	-
2001	47	29	46	68	73	78	41	14	2	1
2000	40	40	63	67	74	82	46	17	3	1
1999	37	48	80	83	80	74	53	14	6	3
Male										
2002	16	21	39	31	31	35	11	5	1	-
2001	33	20	32	38	38	21	10	4	1	-
2000	25	28	39	39	29	30	17	4	2	1
1999	27	26	56	48	38	30	17	5	1	1
Female										
2002	11	16	29	21	38	31	36	14	5	-
2001	14	9	14	30	35	57	31	10	1	1
2000	15	12	24	28	45	52	29	13	1	-
1999	10	22	24	35	42	44	36	9	5	2
Other acute ischemic heart diseases (I24)										
All origins										
Both sexes										
2002	250	289	342	420	478	452	339	153	27	-
2001	263	281	364	397	474	455	317	142	31	-
2000	265	294	350	440	466	439	318	136	35	1
1999	239	294	380	406	449	474	296	116	22	-
Male										
2002	167	193	197	216	228	172	87	35	4	-
2001	186	180	213	206	218	165	72	25	7	-
2000	178	194	204	232	214	153	68	28	5	1
1999	156	191	219	229	194	170	75	19	2	-
Female										
2002	83	96	145	204	250	280	252	118	23	-
2001	77	101	151	191	256	290	245	117	24	-
2000	87	100	146	208	252	286	250	108	30	-
1999	83	103	161	177	255	304	221	97	20	-
Hispanic										
Both sexes										
2002	8	4	10	9	6	4	2	2	1	-
2001	6	9	10	10	4	7	3	3	-	-
2000	4	4	11	9	8	10	4	2	1	-
1999	9	5	3	10	2	5	5	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute ischemic heart diseases (I24)														
Hispanic														
Male														
2002	40	-	-	1	-	-	-	1	-	-	1	4	4	3
2001	42	-	-	-	-	-	-	-	-	-	1	-	2	5
2000	39	-	1	-	-	1	-	1	-	-	1	2	1	3
1999	26	-	-	1	-	-	-	-	-	-	1	1	3	2
Female														
2002	24	-	-	-	-	1	-	-	-	-	-	1	1	1
2001	21	-	-	-	-	-	-	-	-	-	-	1	1	1
2000	30	-	-	-	-	-	-	-	-	-	-	2	1	3
1999	27	-	-	-	-	-	-	-	-	-	-	-	3	2
Non-Hispanic[1]														
Both sexes														
2002	3,339	2	-	-	-	1	1	3	14	17	58	114	221	206
2001	3,284	3	-	-	2	-	7	2	12	32	61	119	184	194
2000	3,282	1	-	-	-	3	4	3	12	30	60	89	184	213
1999	3,167	3	1	-	-	-	3	2	4	25	55	91	154	204
Male														
2002	1,725	1	-	-	-	-	-	1	10	9	40	84	158	150
2001	1,675	2	-	-	1	-	6	2	8	18	46	89	139	129
2000	1,705	1	-	-	-	2	3	2	10	24	45	74	138	160
1999	1,635	1	1	-	-	-	3	1	3	19	43	68	106	156
Female														
2002	1,614	1	-	-	-	1	1	2	4	8	18	30	63	56
2001	1,609	1	-	-	1	-	1	-	4	14	15	30	45	65
2000	1,577	-	-	-	-	1	1	1	2	6	15	15	46	53
1999	1,532	2	-	-	-	-	-	1	1	6	12	23	48	48
Non-Hispanic White														
Both sexes														
2002	2,843	1	-	-	-	1	-	2	9	15	39	85	166	170
2001	2,756	3	-	-	2	-	4	2	9	23	39	88	134	148
2000	2,767	1	-	-	-	3	2	3	8	23	43	65	142	154
1999	2,689	1	1	-	-	-	3	2	3	13	39	55	114	158
Male														
2002	1,454	1	-	-	-	-	-	1	6	8	33	60	119	124
2001	1,386	2	-	-	1	-	3	2	7	14	31	69	101	102
2000	1,412	1	-	-	-	2	1	2	6	19	33	59	106	121
1999	1,361	1	1	-	-	-	3	1	3	10	30	41	82	120
Female														
2002	1,389	-	-	-	-	1	-	1	3	7	6	25	47	46

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)										
Hispanic										
Male										
2002	6	1	9	4	3	2	1	-	-	-
2001	4	6	8	5	2	6	-	3	-	-
2000	2	4	7	5	4	6	-	1	-	-
1999	4	3	1	7	1	2	-	-	-	-
Female										
2002	2	3	1	5	3	2	1	2	1	-
2001	2	3	2	5	2	1	3	-	-	-
2000	2	-	4	4	4	4	4	1	1	-
1999	5	2	2	3	1	3	5	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	241	285	332	411	472	447	337	151	26	-
2001	256	271	354	387	468	448	314	139	31	-
2000	260	289	337	430	457	428	314	133	34	1
1999	229	288	376	395	446	467	287	115	22	-
Male										
2002	161	192	188	212	225	169	86	35	4	-
2001	181	173	205	201	215	159	72	22	7	-
2000	176	189	197	226	210	147	68	27	5	1
1999	151	188	218	222	192	167	75	19	2	-
Female										
2002	80	93	144	199	247	278	251	116	22	-
2001	75	98	149	186	253	289	242	117	24	-
2000	84	100	140	204	247	281	246	106	29	-
1999	78	100	158	173	254	300	212	96	20	-
Non-Hispanic White										
Both sexes										
2002	186	233	277	361	421	404	310	141	22	-
2001	194	209	302	336	421	399	295	122	26	-
2000	203	232	282	363	411	390	284	127	31	-
1999	186	241	312	343	393	431	270	103	21	-
Male										
2002	127	158	155	190	200	159	77	32	4	-
2001	138	128	179	174	200	142	66	20	7	-
2000	140	150	164	193	189	136	60	26	4	-
1999	120	157	182	199	167	156	69	17	2	-
Female										
2002	59	75	122	171	221	245	233	109	18	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute ischemic heart diseases (I24)														
Non-Hispanic White														
Female														
2001	1,370	1	-	-	1	-	1	-	2	9	8	19	33	46
2000	1,355	-	-	-	-	1	1	1	2	4	10	6	36	33
1999	1,328	-	-	-	-	-	-	1	-	3	9	14	32	38
Non-Hispanic Black														
Both sexes														
2002	443	1	-	-	-	-	1	1	5	2	17	27	51	31
2001	453	-	-	-	-	-	3	-	2	8	21	27	42	40
2000	449	-	-	-	-	-	1	-	4	7	15	23	38	51
1999	429	2	-	-	-	-	-	-	1	12	14	34	39	42
Male														
2002	242	-	-	-	-	-	-	-	4	1	5	23	35	21
2001	249	-	-	-	-	-	3	-	-	4	15	17	33	22
2000	255	-	-	-	-	-	1	-	4	5	11	15	29	35
1999	246	-	-	-	-	-	-	-	-	9	11	26	23	33
Female														
2002	201	1	-	-	-	-	1	1	1	1	12	4	16	10
2001	204	-	-	-	-	-	-	-	2	4	6	10	9	18
2000	194	-	-	-	-	-	-	-	-	2	4	8	9	16
1999	183	2	-	-	-	-	-	-	1	3	3	8	16	9
Origin not stated[2]														
Both sexes														
2002	4	-	-	-	-	-	1	-	-	-	1	-	-	-
2001	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	12	-	-	-	-	-	-	-	-	-	1	-	3	-
1999	12	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	5	-	-	-	-	-	-	-	-	-	1	-	2	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	2	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	7	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	8	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)										
Non-Hispanic White										
Female										
2001	56	81	123	162	221	257	229	102	19	-
2000	63	82	118	170	222	254	224	101	27	-
1999	66	84	130	144	226	275	201	86	19	-
Non-Hispanic Black										
Both sexes										
2002	47	47	53	44	45	38	21	8	4	-
2001	58	49	46	39	40	45	12	16	5	-
2000	52	46	45	56	41	34	26	6	3	1
1999	41	39	54	49	48	28	15	10	1	-
Male										
2002	29	32	32	19	21	9	8	3	-	-
2001	40	36	23	24	11	15	4	2	-	-
2000	32	31	27	28	18	9	7	1	1	1
1999	30	26	30	21	20	10	5	2	-	-
Female										
2002	18	15	21	25	24	29	13	5	4	-
2001	18	13	23	15	29	30	8	14	5	-
2000	20	15	18	28	23	25	19	5	2	-
1999	11	13	24	28	28	18	10	8	1	-
Origin not stated[2]										
Both sexes										
2002	1	-	-	-	-	1	-	-	-	-
2001	1	1	-	-	2	-	-	-	-	-
2000	1	1	2	1	1	1	-	1	-	-
1999	1	1	1	1	1	2	4	-	-	-
Male										
2002	-	-	-	-	-	1	-	-	-	-
2001	1	1	-	-	1	-	-	-	-	-
2000	-	1	-	1	-	-	-	-	-	-
1999	1	-	-	-	1	1	-	-	-	-
Female										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	1	-	2	-	1	1	-	1	-	-
1999	-	1	1	1	-	1	4	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other forms of chronic ischemic heart disease (I20,I25)														
All origins														
Both sexes														
2002	311,461	10	2	1	11	13	54	156	422	1,312	3,133	5,718	8,893	12,029
2001	314,081	8	7	3	8	16	76	172	455	1,256	3,008	5,244	8,705	11,366
2000	318,943	9	8	3	7	15	48	124	420	1,299	2,996	5,315	8,416	11,475
1999	326,973	11	2	7	3	27	47	142	432	1,427	2,994	5,207	8,493	11,243
Male														
2002	157,163	7	2	-	10	8	42	122	315	1,005	2,438	4,428	6,878	8,910
2001	156,410	5	1	2	6	10	55	140	347	958	2,305	4,088	6,665	8,389
2000	158,519	6	5	2	3	10	35	103	319	984	2,345	4,206	6,461	8,501
1999	161,434	6	1	6	3	20	30	100	329	1,075	2,362	4,117	6,466	8,208
Female														
2002	154,298	3	-	1	1	5	12	34	107	307	695	1,290	2,015	3,119
2001	157,671	3	6	1	2	6	21	32	108	298	703	1,156	2,040	2,977
2000	160,424	3	3	1	4	5	13	21	101	315	651	1,109	1,955	2,974
1999	165,539	5	1	1	-	7	17	42	103	352	632	1,090	2,027	3,035
Hispanic														
Both sexes														
2002	13,246	1	-	-	1	-	10	19	34	85	179	354	507	677
2001	13,198	-	-	-	3	2	10	19	44	83	145	293	514	646
2000	12,569	1	-	-	1	1	5	7	30	74	163	286	462	649
1999	12,725	-	-	1	-	4	3	14	41	82	141	285	508	662
Male														
2002	7,130	-	-	-	1	-	7	18	28	68	151	278	390	511
2001	7,005	-	-	-	1	1	6	17	36	71	113	232	399	492
2000	6,698	1	-	-	1	1	3	6	25	63	133	239	368	476
1999	6,742	-	-	1	-	3	2	13	32	74	119	231	416	490
Female														
2002	6,116	1	-	-	-	-	3	1	6	17	28	76	117	166
2001	6,193	-	-	-	2	1	4	2	8	12	32	61	115	154
2000	5,871	-	-	-	-	-	2	1	5	11	30	47	94	173
1999	5,983	-	-	-	-	1	1	1	9	8	22	54	92	172
Non-Hispanic[1]														
Both sexes														
2002	296,875	9	2	1	10	13	44	135	383	1,220	2,944	5,338	8,320	11,276
2001	299,663	7	7	3	5	14	65	153	409	1,168	2,851	4,928	8,138	10,649
2000	304,976	8	8	3	6	14	43	115	387	1,217	2,812	4,989	7,898	10,759
1999	312,804	11	2	6	3	22	44	127	391	1,337	2,837	4,893	7,929	10,524
Male														
2002	149,257	7	2	-	9	8	35	102	283	931	2,280	4,131	6,430	8,332
2001	148,724	4	1	2	5	9	48	123	309	885	2,182	3,836	6,225	7,839

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)										
All origins										
Both sexes										
2002	14,840	19,817	29,416	43,195	54,708	55,207	40,876	17,435	4,181	32
2001	14,915	20,531	30,907	44,919	54,810	55,183	40,716	17,458	4,280	38
2000	14,765	21,065	32,717	46,070	54,506	56,529	41,250	17,586	4,281	39
1999	15,280	22,453	34,272	47,920	55,810	56,973	41,734	18,102	4,362	32
Male										
2002	10,460	13,166	18,316	24,622	27,254	22,423	12,371	3,759	604	23
2001	10,439	13,555	19,161	25,537	26,867	21,890	11,852	3,523	588	27
2000	10,315	13,854	20,479	25,925	26,745	22,106	11,834	3,649	604	28
1999	10,571	14,839	21,230	26,821	27,118	21,932	11,831	3,703	637	29
Female										
2002	4,380	6,651	11,100	18,573	27,454	32,784	28,505	13,676	3,577	9
2001	4,476	6,976	11,746	19,382	27,943	33,293	28,864	13,935	3,692	11
2000	4,450	7,211	12,238	20,145	27,761	34,423	29,416	13,937	3,677	11
1999	4,709	7,614	13,042	21,099	28,692	35,041	29,903	14,399	3,725	3
Hispanic										
Both sexes										
2002	905	1,187	1,583	1,896	1,965	1,708	1,348	654	132	1
2001	832	1,209	1,624	1,942	1,949	1,814	1,352	582	133	2
2000	857	1,157	1,531	1,761	1,799	1,800	1,295	561	125	4
1999	869	1,216	1,569	1,796	1,765	1,744	1,299	602	119	5
Male										
2002	632	760	962	1,042	929	684	459	180	29	1
2001	563	783	971	1,079	937	728	412	131	31	2
2000	602	726	940	978	854	688	417	146	28	3
1999	574	779	959	967	822	673	389	170	23	5
Female										
2002	273	427	621	854	1,036	1,024	889	474	103	-
2001	269	426	653	863	1,012	1,086	940	451	102	-
2000	255	431	591	783	945	1,112	878	415	97	1
1999	295	437	610	829	943	1,071	910	432	96	-
Non-Hispanic[1]										
Both sexes										
2002	13,835	18,515	27,683	41,099	52,567	53,296	39,401	16,728	4,032	24
2001	14,003	19,220	29,155	42,795	52,666	53,203	39,254	16,812	4,135	23
2000	13,790	19,800	31,016	44,103	52,513	54,523	39,835	16,966	4,146	25
1999	14,300	21,125	32,516	45,906	53,822	55,029	40,301	17,430	4,227	22
Male										
2002	9,755	12,326	17,249	23,459	26,234	21,656	11,873	3,568	572	15
2001	9,809	12,699	18,107	24,348	25,829	21,097	11,418	3,378	554	17

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other forms of chronic ischemic heart disease (I20,I25)														
Non-Hispanic[1]														
Male														
2000	151,009	5	5	2	2	9	32	96	292	915	2,192	3,933	6,043	7,971
1999	153,875	6	1	5	3	17	28	86	297	994	2,229	3,861	6,006	7,672
Female														
2002	147,618	2	-	1	1	5	9	33	100	289	664	1,207	1,890	2,944
2001	150,939	3	6	1	-	5	17	30	100	283	669	1,092	1,913	2,810
2000	153,967	3	3	1	4	5	11	19	95	302	620	1,056	1,855	2,788
1999	158,929	5	1	1	-	5	16	41	94	343	608	1,032	1,923	2,852
Non-Hispanic White														
Both sexes														
2002	260,336	4	-	1	7	9	30	98	274	903	2,263	4,086	6,368	8,857
2001	262,891	3	4	3	4	8	38	106	288	875	2,152	3,713	6,281	8,351
2000	268,535	5	8	3	5	12	29	76	276	902	2,133	3,750	6,164	8,461
1999	275,726	8	1	4	2	14	24	87	280	1,001	2,108	3,733	6,119	8,223
Male														
2002	130,900	3	-	-	6	6	23	77	209	700	1,812	3,281	5,090	6,711
2001	130,367	2	-	2	4	4	30	91	220	674	1,697	2,992	4,937	6,287
2000	132,908	3	5	2	2	8	21	64	218	696	1,732	3,076	4,847	6,405
1999	135,406	4	1	4	2	10	18	58	218	754	1,715	3,048	4,743	6,159
Female														
2002	129,436	1	-	1	1	3	7	21	65	203	451	805	1,278	2,146
2001	132,524	1	4	1	-	4	8	15	68	201	455	721	1,344	2,064
2000	135,627	2	3	1	3	4	8	12	58	206	401	674	1,317	2,056
1999	140,320	4	-	-	-	4	6	29	62	247	393	685	1,376	2,064
Non-Hispanic Black														
Both sexes														
2002	30,917	4	2	-	3	4	11	29	87	271	595	1,106	1,725	2,120
2001	31,443	4	3	-	1	6	22	44	103	258	615	1,064	1,660	2,008
2000	31,276	2	-	-	-	2	10	34	98	274	595	1,110	1,527	2,026
1999	31,819	2	1	2	1	6	19	31	93	295	651	1,029	1,607	2,055
Male														
2002	15,127	3	2	-	3	2	9	19	54	190	392	731	1,155	1,401
2001	15,378	2	1	-	1	5	14	30	75	183	418	724	1,140	1,315
2000	15,075	1	-	-	-	1	7	27	64	183	392	748	1,026	1,368
1999	15,384	1	-	1	1	5	9	22	65	204	444	700	1,100	1,328
Female														
2002	15,790	1	-	-	-	2	2	10	33	81	203	375	570	719
2001	16,065	2	2	-	-	1	8	14	28	75	197	340	520	693
2000	16,201	1	-	-	-	1	3	7	34	91	203	362	501	658

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)										
Non-Hispanic[1]										
Male										
2000	9,623	13,045	19,420	24,824	25,802	21,339	11,381	3,486	574	18
1999	9,907	13,974	20,139	25,724	26,204	21,168	11,405	3,518	612	19
Female										
2002	4,080	6,189	10,434	17,640	26,333	31,640	27,528	13,160	3,460	9
2001	4,194	6,521	11,048	18,447	26,837	32,106	27,836	13,434	3,581	6
2000	4,167	6,755	11,596	19,279	26,711	33,184	28,454	13,480	3,572	7
1999	4,393	7,151	12,377	20,182	27,618	33,861	28,896	13,912	3,615	3
Non-Hispanic White										
Both sexes										
2002	10,962	15,006	23,570	35,921	47,528	49,098	36,422	15,360	3,553	16
2001	11,051	15,560	24,861	37,499	47,651	48,984	36,391	15,416	3,638	14
2000	10,871	16,279	26,506	38,858	47,639	50,187	36,985	15,661	3,707	18
1999	11,383	17,398	27,839	40,575	48,994	50,639	37,420	16,061	3,800	13
Male										
2002	7,880	10,236	14,983	20,917	24,027	20,173	11,008	3,256	492	10
2001	7,946	10,561	15,711	21,725	23,655	19,663	10,596	3,088	472	10
2000	7,813	11,035	16,963	22,204	23,707	19,838	10,593	3,177	485	14
1999	8,036	11,842	17,581	23,107	24,186	19,622	10,528	3,214	545	11
Female										
2002	3,082	4,770	8,587	15,004	23,501	28,925	25,414	12,104	3,061	6
2001	3,105	4,999	9,150	15,774	23,996	29,321	25,795	12,328	3,166	4
2000	3,058	5,244	9,543	16,654	23,932	30,349	26,392	12,484	3,222	4
1999	3,347	5,556	10,258	17,468	24,808	31,017	26,892	12,847	3,255	2
Non-Hispanic Black										
Both sexes										
2002	2,506	3,007	3,498	4,319	4,194	3,436	2,458	1,121	413	8
2001	2,593	3,167	3,672	4,473	4,191	3,503	2,413	1,201	435	7
2000	2,548	3,055	3,924	4,427	4,096	3,649	2,393	1,113	386	7
1999	2,542	3,281	4,013	4,506	4,031	3,670	2,399	1,195	381	9
Male										
2002	1,620	1,769	1,892	2,044	1,775	1,124	645	230	62	5
2001	1,613	1,824	2,043	2,161	1,764	1,110	656	224	69	6
2000	1,547	1,696	2,088	2,137	1,688	1,186	599	237	76	4
1999	1,603	1,857	2,149	2,123	1,616	1,203	651	239	55	8
Female										
2002	886	1,238	1,606	2,275	2,419	2,312	1,813	891	351	3
2001	980	1,343	1,629	2,312	2,427	2,393	1,757	977	366	1
2000	1,001	1,359	1,836	2,290	2,408	2,463	1,794	876	310	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other forms of chronic ischemic heart disease (I20,I25)														
Non-Hispanic Black														
Female														
1999	16,435	1	1	1	-	1	10	9	28	91	207	329	507	727
Origin not stated[2]														
Both sexes														
2002	1,340	-	-	-	-	-	-	2	5	7	10	26	66	76
2001	1,220	1	-	-	-	-	1	-	2	5	12	23	53	71
2000	1,398	-	-	-	-	-	-	2	3	8	21	40	56	67
1999	1,444	-	-	-	-	1	-	1	-	8	16	29	56	57
Male														
2002	776	-	-	-	-	-	-	2	4	6	7	19	58	67
2001	681	1	-	-	-	-	1	-	2	2	10	20	41	58
2000	812	-	-	-	-	-	-	1	2	6	20	34	50	54
1999	817	-	-	-	-	-	-	1	-	7	14	25	44	46
Female														
2002	564	-	-	-	-	-	-	-	1	1	3	7	8	9
2001	539	-	-	-	-	-	-	-	-	3	2	3	12	13
2000	586	-	-	-	-	-	-	1	1	2	1	6	6	13
1999	627	-	-	-	-	1	-	-	-	1	2	4	12	11
Atherosclerotic cardiovascular disease, so described (I25.0)														
All origins														
Both sexes														
2002	68,129	2	-	-	1	4	13	39	142	410	1,105	2,081	3,285	4,189
2001	68,103	-	-	-	-	3	17	39	140	402	1,055	1,838	3,209	3,924
2000	69,554	1	-	-	1	-	15	33	111	391	985	1,894	3,086	3,841
1999	71,926	-	1	-	1	-	13	32	128	453	1,035	1,792	2,945	3,747
Male														
2002	36,047	1	-	-	-	3	10	31	113	325	891	1,634	2,605	3,250
2001	35,440	-	-	-	-	2	14	30	103	315	815	1,488	2,549	2,978
2000	35,896	-	-	-	1	-	12	26	89	312	787	1,533	2,444	2,978
1999	36,633	-	-	-	1	-	10	21	98	355	837	1,438	2,314	2,842
Female														
2002	32,082	1	-	-	1	1	3	8	29	85	214	447	680	939
2001	32,663	-	-	-	-	1	3	9	37	87	240	350	660	946
2000	33,658	1	-	-	-	-	3	7	22	79	198	361	642	863
1999	35,293	-	1	-	-	-	3	11	30	98	198	354	631	905

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)										
Non-Hispanic Black										
Female										
1999	939	1,424	1,864	2,383	2,415	2,467	1,748	956	326	1
Origin not stated[2]										
Both sexes										
2002	100	115	150	200	176	203	127	53	17	7
2001	80	102	128	182	195	166	110	64	12	13
2000	118	108	170	206	194	206	120	59	10	10
1999	111	112	187	218	223	200	134	70	16	5
Male										
2002	73	80	105	121	91	83	39	11	3	7
2001	67	73	83	110	101	65	22	14	3	8
2000	90	83	119	123	89	79	36	17	2	7
1999	90	86	132	130	92	91	37	15	2	5
Female										
2002	27	35	45	79	85	120	88	42	14	-
2001	13	29	45	72	94	101	88	50	9	5
2000	28	25	51	83	105	127	84	42	8	3
1999	21	26	55	88	131	109	97	55	14	-
Atherosclerotic cardiovascular disease, so described (I25.0)										
All origins										
Both sexes										
2002	4,604	5,163	6,654	8,649	9,969	10,007	7,555	3,361	873	23
2001	4,514	5,272	6,733	9,041	10,155	9,984	7,556	3,323	873	25
2000	4,401	5,309	7,258	9,271	10,274	10,397	7,953	3,429	885	19
1999	4,636	5,691	7,704	9,666	10,655	10,745	8,204	3,591	863	24
Male										
2002	3,299	3,498	4,119	4,746	4,670	3,814	2,165	722	134	17
2001	3,267	3,483	4,185	4,897	4,774	3,630	2,167	616	108	19
2000	3,147	3,557	4,512	5,086	4,639	3,740	2,205	691	122	15
1999	3,281	3,842	4,704	5,200	4,763	3,895	2,196	692	122	22
Female										
2002	1,305	1,665	2,535	3,903	5,299	6,193	5,390	2,639	739	6
2001	1,247	1,789	2,548	4,144	5,381	6,354	5,389	2,707	765	6
2000	1,254	1,752	2,746	4,185	5,635	6,657	5,748	2,738	763	4
1999	1,355	1,849	3,000	4,466	5,892	6,850	6,008	2,899	741	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerotic cardiovascular disease, so described (I25.0)														
Hispanic														
Both sexes														
2002	3,166	1	-	-	-	-	4	9	17	30	78	141	186	258
2001	3,150	-	-	-	-	-	1	8	17	26	55	120	223	247
2000	2,953	-	-	-	-	-	2	2	5	28	55	111	188	230
1999	3,063	-	-	-	-	-	1	4	10	23	50	110	198	247
Male														
2002	1,946	-	-	-	-	-	3	8	16	22	67	112	158	220
2001	1,928	-	-	-	-	-	-	8	13	22	41	96	192	196
2000	1,794	-	-	-	-	-	2	1	5	25	46	97	160	182
1999	1,847	-	-	-	-	-	1	3	9	22	41	89	171	184
Female														
2002	1,220	1	-	-	-	-	1	1	1	8	11	29	28	38
2001	1,222	-	-	-	-	-	1	-	4	4	14	24	31	51
2000	1,159	-	-	-	-	-	-	1	-	3	9	14	28	48
1999	1,216	-	-	-	-	-	-	1	1	1	9	21	27	63
Non-Hispanic[1]														
Both sexes														
2002	64,528	1	-	-	1	4	9	28	124	378	1,025	1,928	3,059	3,888
2001	64,534	-	-	-	-	3	16	31	123	373	993	1,706	2,958	3,637
2000	66,141	1	-	-	1	-	13	31	106	360	923	1,764	2,867	3,573
1999	68,373	-	1	-	1	-	12	28	118	427	977	1,671	2,716	3,472
Male														
2002	33,809	1	-	-	-	3	7	21	97	302	822	1,514	2,413	2,989
2001	33,247	-	-	-	-	2	14	22	90	292	767	1,381	2,336	2,751
2000	33,786	-	-	-	1	-	10	25	84	285	734	1,419	2,255	2,765
1999	34,451	-	-	-	1	-	9	18	89	330	788	1,339	2,118	2,635
Female														
2002	30,719	-	-	-	1	1	2	7	27	76	203	414	646	899
2001	31,287	-	-	-	-	1	2	9	33	81	226	325	622	886
2000	32,355	1	-	-	-	-	3	6	22	75	189	345	612	808
1999	33,922	-	1	-	-	-	3	10	29	97	189	332	598	837
Non-Hispanic White														
Both sexes														
2002	53,161	1	-	-	1	4	7	21	84	265	755	1,458	2,245	2,892
2001	53,031	-	-	-	-	2	9	20	88	275	724	1,244	2,177	2,632
2000	54,544	-	-	-	1	-	11	17	78	261	683	1,274	2,133	2,606
1999	56,375	-	-	-	1	-	5	16	80	310	707	1,191	1,964	2,508
Male														
2002	27,493	1	-	-	-	3	5	16	68	216	628	1,178	1,830	2,264
2001	26,960	-	-	-	-	1	7	15	67	221	573	1,042	1,767	2,059

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)										
Hispanic										
Both sexes										
2002	328	327	395	395	349	264	240	112	31	1
2001	301	346	402	396	339	314	218	116	19	2
2000	292	324	383	331	318	331	233	94	24	2
1999	293	364	383	384	324	293	243	107	26	3
Male										
2002	241	221	254	219	156	112	91	36	9	1
2001	220	246	256	234	171	133	66	26	6	2
2000	224	201	237	215	153	133	75	28	9	1
1999	200	268	259	220	152	105	83	32	5	3
Female										
2002	87	106	141	176	193	152	149	76	22	-
2001	81	100	146	162	168	181	152	90	13	-
2000	68	123	146	116	165	198	158	66	15	1
1999	93	96	124	164	172	188	160	75	21	-
Non-Hispanic[1]										
Both sexes										
2002	4,221	4,794	6,198	8,189	9,582	9,707	7,293	3,242	840	17
2001	4,179	4,879	6,280	8,576	9,770	9,635	7,312	3,200	852	11
2000	4,054	4,945	6,811	8,862	9,907	10,029	7,693	3,331	859	11
1999	4,282	5,281	7,243	9,198	10,267	10,403	7,946	3,477	835	18
Male										
2002	3,015	3,249	3,827	4,486	4,491	3,684	2,068	684	125	11
2001	3,019	3,202	3,891	4,622	4,578	3,483	2,096	589	102	10
2000	2,881	3,324	4,225	4,819	4,462	3,589	2,122	663	113	10
1999	3,028	3,538	4,384	4,924	4,580	3,767	2,111	659	117	16
Female										
2002	1,206	1,545	2,371	3,703	5,091	6,023	5,225	2,558	715	6
2001	1,160	1,677	2,389	3,954	5,192	6,152	5,216	2,611	750	1
2000	1,173	1,621	2,586	4,043	5,445	6,440	5,571	2,668	746	1
1999	1,254	1,743	2,859	4,274	5,687	6,636	5,835	2,818	718	2
Non-Hispanic White										
Both sexes										
2002	3,151	3,551	4,893	6,693	8,279	8,663	6,572	2,908	708	10
2001	3,017	3,566	4,938	7,008	8,513	8,594	6,637	2,856	724	7
2000	2,931	3,747	5,332	7,331	8,540	8,891	6,948	3,017	735	8
1999	3,118	3,954	5,711	7,580	8,937	9,244	7,180	3,136	723	10
Male										
2002	2,295	2,441	3,085	3,707	3,876	3,304	1,859	609	102	6
2001	2,232	2,397	3,099	3,824	4,025	3,132	1,891	522	80	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerotic cardiovascular disease, so described (I25.0)														
Non-Hispanic White														
Male														
2000	27,461	-	-	-	1	-	8	13	64	211	564	1,060	1,726	2,053
1999	27,943	-	-	-	1	-	5	11	62	244	594	984	1,574	1,960
Female														
2002	25,668	-	-	-	1	1	2	5	16	49	127	280	415	628
2001	26,071	-	-	-	-	1	2	5	21	54	151	202	410	573
2000	27,083	-	-	-	-	-	3	4	14	50	119	214	407	553
1999	28,432	-	-	-	-	-	-	5	18	66	113	207	390	548
Non-Hispanic Black														
Both sexes														
2002	10,024	-	-	-	-	-	2	5	34	99	234	426	730	880
2001	10,156	-	-	-	-	1	6	11	27	89	235	408	707	889
2000	10,283	1	-	-	-	-	1	12	24	86	206	451	669	870
1999	10,643	-	1	-	-	-	7	8	33	110	248	433	672	876
Male														
2002	5,483	-	-	-	-	-	2	3	24	73	160	298	508	633
2001	5,476	-	-	-	-	1	6	7	17	64	166	291	508	604
2000	5,528	-	-	-	-	-	1	10	16	64	143	324	478	634
1999	5,692	-	-	-	-	-	4	5	22	80	174	313	479	604
Female														
2002	4,541	-	-	-	-	-	-	2	10	26	74	128	222	247
2001	4,680	-	-	-	-	-	-	4	10	25	69	117	199	285
2000	4,755	1	-	-	-	-	-	2	8	22	63	127	191	236
1999	4,951	-	1	-	-	-	3	3	11	30	74	120	193	272
Origin not stated[2]														
Both sexes														
2002	435	-	-	-	-	-	-	2	1	2	2	12	40	43
2001	419	-	-	-	-	-	-	-	-	3	7	12	28	40
2000	460	-	-	-	-	-	-	-	-	3	7	19	31	38
1999	490	-	-	-	-	-	-	-	-	3	8	11	31	28
Male														
2002	292	-	-	-	-	-	-	2	-	1	2	8	34	41
2001	265	-	-	-	-	-	-	-	-	1	7	11	21	31
2000	316	-	-	-	-	-	-	-	-	2	7	17	29	31
1999	335	-	-	-	-	-	-	-	-	3	8	10	25	23
Female														
2002	143	-	-	-	-	-	-	-	1	1	-	4	6	2
2001	154	-	-	-	-	-	-	-	-	2	-	1	7	9
2000	144	-	-	-	-	-	-	-	-	1	-	2	2	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)										
Non-Hispanic White										
Male										
2000	2,157	2,590	3,359	3,996	3,868	3,200	1,916	583	84	8
1999	2,248	2,715	3,512	4,093	4,004	3,358	1,884	583	103	8
Female										
2002	856	1,110	1,808	2,986	4,403	5,359	4,713	2,299	606	4
2001	785	1,169	1,839	3,184	4,488	5,462	4,746	2,334	644	1
2000	774	1,157	1,973	3,335	4,672	5,691	5,032	2,434	651	-
1999	870	1,239	2,199	3,487	4,933	5,886	5,296	2,553	620	2
Non-Hispanic Black										
Both sexes										
2002	964	1,107	1,151	1,328	1,138	892	620	290	117	7
2001	1,038	1,169	1,192	1,382	1,090	908	590	298	113	3
2000	1,007	1,068	1,312	1,329	1,193	1,002	659	277	113	3
1999	1,028	1,194	1,359	1,435	1,158	1,004	666	302	101	8
Male										
2002	645	712	645	684	539	302	169	63	18	5
2001	695	710	702	702	472	289	172	50	17	3
2000	648	638	760	704	500	339	174	67	26	2
1999	674	746	772	717	493	344	185	63	9	8
Female										
2002	319	395	506	644	599	590	451	227	99	2
2001	343	459	490	680	618	619	418	248	96	-
2000	359	430	552	625	693	663	485	210	87	1
1999	354	448	587	718	665	660	481	239	92	-
Origin not stated[2]										
Both sexes										
2002	55	42	61	65	38	36	22	7	2	5
2001	34	47	51	69	46	35	26	7	2	12
2000	55	40	64	78	49	37	27	4	2	6
1999	61	46	78	84	64	49	15	7	2	3
Male										
2002	43	28	38	41	23	18	6	2	-	5
2001	28	35	38	41	25	14	5	1	-	7
2000	42	32	50	52	24	18	8	-	-	4
1999	53	36	61	56	31	23	2	1	-	3
Female										
2002	12	14	23	24	15	18	16	5	2	-
2001	6	12	13	28	21	21	21	6	2	5
2000	13	8	14	26	25	19	19	4	2	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerotic cardiovascular disease, so described (I25.0)														
Origin not stated[2]														
Female														
1999	155	-	-	-	-	-	-	-	-	-	-	1	6	5
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)														
All origins														
Both sexes														
2002	243,332	8	2	1	10	9	41	117	280	902	2,028	3,637	5,608	7,840
2001	245,978	8	7	3	8	13	59	133	315	854	1,953	3,406	5,496	7,442
2000	249,389	8	8	3	6	15	33	91	309	908	2,011	3,421	5,330	7,634
1999	255,047	11	1	7	2	27	34	110	304	974	1,959	3,415	5,548	7,496
Male														
2002	121,116	6	2	-	10	5	32	91	202	680	1,547	2,794	4,273	5,660
2001	120,970	5	1	2	6	8	41	110	244	643	1,490	2,600	4,116	5,411
2000	122,623	6	5	2	2	10	23	77	230	672	1,558	2,673	4,017	5,523
1999	124,801	6	1	6	2	20	20	79	231	720	1,525	2,679	4,152	5,366
Female														
2002	122,216	2	-	1	-	4	9	26	78	222	481	843	1,335	2,180
2001	125,008	3	6	1	2	5	18	23	71	211	463	806	1,380	2,031
2000	126,766	2	3	1	4	5	10	14	79	236	453	748	1,313	2,111
1999	130,246	5	-	1	-	7	14	31	73	254	434	736	1,396	2,130
Hispanic														
Both sexes														
2002	10,080	-	-	-	1	-	6	10	17	55	101	213	321	419
2001	10,048	-	-	-	3	2	9	11	27	57	90	173	291	399
2000	9,616	1	-	-	1	1	3	5	25	46	108	175	274	419
1999	9,662	-	-	1	-	4	2	10	31	59	91	175	310	415
Male														
2002	5,184	-	-	-	1	-	4	10	12	46	84	166	232	291
2001	5,077	-	-	-	1	1	6	9	23	49	72	136	207	296
2000	4,904	1	-	-	1	1	1	5	20	38	87	142	208	294
1999	4,895	-	-	1	-	3	1	10	23	52	78	142	245	306
Female														
2002	4,896	-	-	-	-	-	2	-	5	9	17	47	89	128
2001	4,971	-	-	-	2	1	3	2	4	8	18	37	84	103
2000	4,712	-	-	-	-	-	2	-	5	8	21	33	66	125
1999	4,767	-	-	-	-	1	1	-	8	7	13	33	65	109

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)										
Origin not stated[2]										
Female										
1999	8	10	17	28	33	26	13	6	2	-
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)										
All origins										
Both sexes										
2002	10,236	14,654	22,762	34,546	44,739	45,200	33,321	14,074	3,308	9
2001	10,401	15,259	24,174	35,878	44,655	45,199	33,160	14,135	3,407	13
2000	10,364	15,756	25,459	36,799	44,232	46,132	33,297	14,157	3,396	20
1999	10,644	16,762	26,568	38,254	45,155	46,228	33,530	14,511	3,499	8
Male										
2002	7,161	9,668	14,197	19,876	22,584	18,609	10,206	3,037	470	6
2001	7,172	10,072	14,976	20,640	22,093	18,260	9,685	2,907	480	8
2000	7,168	10,297	15,967	20,839	22,106	18,366	9,629	2,958	482	13
1999	7,290	10,997	16,526	21,621	22,355	18,037	9,635	3,011	515	7
Female										
2002	3,075	4,986	8,565	14,670	22,155	26,591	23,115	11,037	2,838	3
2001	3,229	5,187	9,198	15,238	22,562	26,939	23,475	11,228	2,927	5
2000	3,196	5,459	9,492	15,960	22,126	27,766	23,668	11,199	2,914	7
1999	3,354	5,765	10,042	16,633	22,800	28,191	23,895	11,500	2,984	1
Hispanic										
Both sexes										
2002	577	860	1,188	1,501	1,616	1,444	1,108	542	101	-
2001	531	863	1,222	1,546	1,610	1,500	1,134	466	114	-
2000	565	833	1,148	1,430	1,481	1,469	1,062	467	101	2
1999	576	852	1,186	1,412	1,441	1,451	1,056	495	93	2
Male										
2002	391	539	708	823	773	572	368	144	20	-
2001	343	537	715	845	766	595	346	105	25	-
2000	378	525	703	763	701	555	342	118	19	2
1999	374	511	700	747	670	568	306	138	18	2
Female										
2002	186	321	480	678	843	872	740	398	81	-
2001	188	326	507	701	844	905	788	361	89	-
2000	187	308	445	667	780	914	720	349	82	-
1999	202	341	486	665	771	883	750	357	75	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)														
Non-Hispanic[1]														
Both sexes														
2002	232,347	8	2	1	9	9	35	107	259	842	1,919	3,410	5,261	7,388
2001	235,129	7	7	3	5	11	49	122	286	795	1,858	3,222	5,180	7,012
2000	238,835	7	8	3	5	14	30	84	281	857	1,889	3,225	5,031	7,186
1999	244,431	11	1	6	2	22	32	99	273	910	1,860	3,222	5,213	7,052
Male														
2002	115,448	6	2	-	9	5	28	81	186	629	1,458	2,617	4,017	5,343
2001	115,477	4	1	2	5	7	34	101	219	593	1,415	2,455	3,889	5,088
2000	117,223	5	5	2	1	9	22	71	208	630	1,458	2,514	3,788	5,206
1999	119,424	6	1	5	2	17	19	68	208	664	1,441	2,522	3,888	5,037
Female														
2002	116,899	2	-	1	-	4	7	26	73	213	461	793	1,244	2,045
2001	119,652	3	6	1	-	4	15	21	67	202	443	767	1,291	1,924
2000	121,612	2	3	1	4	5	8	13	73	227	431	711	1,243	1,980
1999	125,007	5	-	1	-	5	13	31	65	246	419	700	1,325	2,015
Non-Hispanic White														
Both sexes														
2002	207,175	3	-	1	6	5	23	77	190	638	1,508	2,628	4,123	5,965
2001	209,860	3	4	3	4	6	29	86	200	600	1,428	2,469	4,104	5,719
2000	213,991	5	8	3	4	12	18	59	198	641	1,450	2,476	4,031	5,855
1999	219,351	8	1	4	1	14	19	71	200	691	1,401	2,542	4,155	5,715
Male														
2002	103,407	2	-	-	6	3	18	61	141	484	1,184	2,103	3,260	4,447
2001	103,407	2	-	2	4	3	23	76	153	453	1,124	1,950	3,170	4,228
2000	105,447	3	5	2	1	8	13	51	154	485	1,168	2,016	3,121	4,352
1999	107,463	4	1	4	1	10	13	47	156	510	1,121	2,064	3,169	4,199
Female														
2002	103,768	1	-	1	-	2	5	16	49	154	324	525	863	1,518
2001	106,453	1	4	1	-	3	6	10	47	147	304	519	934	1,491
2000	108,544	2	3	1	3	4	5	8	44	156	282	460	910	1,503
1999	111,888	4	-	-	-	4	6	24	44	181	280	478	986	1,516
Non-Hispanic Black														
Both sexes														
2002	20,893	4	2	-	3	4	9	24	53	172	361	680	995	1,240
2001	21,287	4	3	-	1	5	16	33	76	169	380	656	953	1,119
2000	20,993	1	-	-	-	2	9	22	74	188	389	659	858	1,156
1999	21,176	2	-	2	1	6	12	23	60	185	403	596	935	1,179
Male														
2002	9,644	3	2	-	3	2	7	16	30	117	232	433	647	768
2001	9,902	2	1	-	1	4	8	23	58	119	252	433	632	711

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)										
Non-Hispanic[1]										
Both sexes										
2002	9,614	13,721	21,485	32,910	42,985	43,589	32,108	13,486	3,192	7
2001	9,824	14,341	22,875	34,219	42,896	43,568	31,942	13,612	3,283	12
2000	9,736	14,855	24,205	35,241	42,606	44,494	32,142	13,635	3,287	14
1999	10,018	15,844	25,273	36,708	43,555	44,626	32,355	13,953	3,392	4
Male										
2002	6,740	9,077	13,422	18,973	21,743	17,972	9,805	2,884	447	4
2001	6,790	9,497	14,216	19,726	21,251	17,614	9,322	2,789	452	7
2000	6,742	9,721	15,195	20,005	21,340	17,750	9,259	2,823	461	8
1999	6,879	10,436	15,755	20,800	21,624	17,401	9,294	2,859	495	3
Female										
2002	2,874	4,644	8,063	13,937	21,242	25,617	22,303	10,602	2,745	3
2001	3,034	4,844	8,659	14,493	21,645	25,954	22,620	10,823	2,831	5
2000	2,994	5,134	9,010	15,236	21,266	26,744	22,883	10,812	2,826	6
1999	3,139	5,408	9,518	15,908	21,931	27,225	23,061	11,094	2,897	1
Non-Hispanic White										
Both sexes										
2002	7,811	11,455	18,677	29,228	39,249	40,435	29,850	12,452	2,845	6
2001	8,034	11,994	19,923	30,491	39,138	40,390	29,754	12,560	2,914	7
2000	7,940	12,532	21,174	31,527	39,099	41,296	30,037	12,644	2,972	10
1999	8,265	13,444	22,128	32,995	40,057	41,395	30,240	12,925	3,077	3
Male										
2002	5,585	7,795	11,898	17,210	20,151	16,869	9,149	2,647	390	4
2001	5,714	8,164	12,612	17,901	19,630	16,531	8,705	2,566	392	4
2000	5,656	8,445	13,604	18,208	19,839	16,638	8,677	2,594	401	6
1999	5,788	9,127	14,069	19,014	20,182	16,264	8,644	2,631	442	3
Female										
2002	2,226	3,660	6,779	12,018	19,098	23,566	20,701	9,805	2,455	2
2001	2,320	3,830	7,311	12,590	19,508	23,859	21,049	9,994	2,522	3
2000	2,284	4,087	7,570	13,319	19,260	24,658	21,360	10,050	2,571	4
1999	2,477	4,317	8,059	13,981	19,875	25,131	21,596	10,294	2,635	-
Non-Hispanic Black										
Both sexes										
2002	1,542	1,900	2,347	2,991	3,056	2,544	1,838	831	296	1
2001	1,555	1,998	2,480	3,091	3,101	2,595	1,823	903	322	4
2000	1,541	1,987	2,612	3,098	2,903	2,647	1,734	836	273	4
1999	1,514	2,087	2,654	3,071	2,873	2,666	1,733	893	280	1
Male										
2002	975	1,057	1,247	1,360	1,236	822	476	167	44	-
2001	918	1,114	1,341	1,459	1,292	821	484	174	52	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)														
Non-Hispanic Black														
Male														
2000	9,547	1	-	-	-	1	6	17	48	119	249	424	548	734
1999	9,692	1	-	1	1	5	5	17	43	124	270	387	621	724
Female														
2002	11,249	1	-	-	-	2	2	8	23	55	129	247	348	472
2001	11,385	2	2	-	-	1	8	10	18	50	128	223	321	408
2000	11,446	-	-	-	-	1	3	5	26	69	140	235	310	422
1999	11,484	1	-	1	-	1	7	6	17	61	133	209	314	455
Origin not stated[2]														
Both sexes														
2002	905	-	-	-	-	-	-	-	4	5	8	14	26	33
2001	801	1	-	-	-	-	1	-	2	2	5	11	25	31
2000	938	-	-	-	-	-	-	2	3	5	14	21	25	29
1999	954	-	-	-	-	1	-	1	-	5	8	18	25	29
Male														
2002	484	-	-	-	-	-	-	-	4	5	5	11	24	26
2001	416	1	-	-	-	-	1	-	2	1	3	9	20	27
2000	496	-	-	-	-	-	-	1	2	4	13	17	21	23
1999	482	-	-	-	-	-	-	1	-	4	6	15	19	23
Female														
2002	421	-	-	-	-	-	-	-	-	-	3	3	2	7
2001	385	-	-	-	-	-	-	-	-	1	2	2	5	4
2000	442	-	-	-	-	-	-	1	1	1	1	4	4	6
1999	472	-	-	-	-	1	-	-	-	1	2	3	6	6
Other heart diseases (I26-I51)														
All origins														
Both sexes														
2002	169,540	473	157	84	143	362	474	723	1,085	1,769	2,851	3,849	4,862	5,913
2001	166,949	449	215	92	154	305	489	687	1,049	1,770	2,702	3,708	4,776	5,637
2000	165,428	457	168	96	152	355	498	687	1,062	1,706	2,751	3,536	4,551	5,417
1999	165,828	490	177	99	149	398	482	680	1,099	1,879	2,768	3,502	4,386	5,465
Male														
2002	74,230	249	82	38	87	250	307	446	692	1,051	1,809	2,468	3,082	3,587
2001	72,439	222	117	41	82	193	307	413	624	1,099	1,678	2,361	2,946	3,393
2000	72,122	238	98	58	92	218	300	407	642	1,033	1,713	2,278	2,873	3,384
1999	72,601	255	100	54	83	250	283	400	667	1,176	1,718	2,223	2,756	3,419
Female														
2002	95,310	224	75	46	56	112	167	277	393	718	1,042	1,381	1,780	2,326

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20, I25.1-I25.9)										
Non-Hispanic Black										
Male										
2000	899	1,058	1,328	1,433	1,188	847	425	170	50	2
1999	929	1,111	1,377	1,406	1,123	859	466	176	46	-
Female										
2002	567	843	1,100	1,631	1,820	1,722	1,362	664	252	1
2001	637	884	1,139	1,632	1,809	1,774	1,339	729	270	1
2000	642	929	1,284	1,665	1,715	1,800	1,309	666	223	2
1999	585	976	1,277	1,665	1,750	1,807	1,267	717	234	1
Origin not stated[2]										
Both sexes										
2002	45	73	89	135	138	167	105	46	15	2
2001	46	55	77	113	149	131	84	57	10	1
2000	63	68	106	128	145	169	93	55	8	4
1999	50	66	109	134	159	151	119	63	14	2
Male										
2002	30	52	67	80	68	65	33	9	3	2
2001	39	38	45	69	76	51	17	13	3	1
2000	48	51	69	71	65	61	28	17	2	3
1999	37	50	71	74	61	68	35	14	2	2
Female										
2002	15	21	22	55	70	102	72	37	12	-
2001	7	17	32	44	73	80	67	44	7	-
2000	15	17	37	57	80	108	65	38	6	1
1999	13	16	38	60	98	83	84	49	12	-
Other heart diseases (I26-I51)										
All origins										
Both sexes										
2002	7,033	9,377	14,303	21,276	28,500	30,794	23,517	9,779	2,210	6
2001	6,898	9,596	14,460	21,325	27,651	30,343	22,779	9,653	2,204	7
2000	6,945	9,875	14,804	21,524	26,843	29,519	22,677	9,591	2,206	8
1999	7,115	10,034	15,538	21,689	27,023	29,428	21,894	9,401	2,125	7
Male										
2002	4,184	5,275	7,716	10,463	12,308	11,167	6,558	2,070	338	3
2001	4,110	5,356	7,811	10,538	12,002	10,595	6,187	2,028	331	5
2000	4,148	5,584	8,037	10,505	11,501	10,488	6,178	1,983	359	5
1999	4,281	5,870	8,450	10,567	11,561	10,293	5,879	1,988	323	5
Female										
2002	2,849	4,102	6,587	10,813	16,192	19,627	16,959	7,709	1,872	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other heart diseases (I26-I51)														
All origins														
Female														
2001	94,510	227	98	51	72	112	182	274	425	671	1,024	1,347	1,830	2,244
2000	93,306	219	70	38	60	137	198	280	420	673	1,038	1,258	1,678	2,033
1999	93,227	235	77	45	66	148	199	280	432	703	1,050	1,279	1,630	2,046
Hispanic														
Both sexes														
2002	5,152	81	23	11	16	34	68	85	103	124	174	191	225	244
2001	4,899	75	31	17	16	39	47	73	106	123	145	171	199	224
2000	4,792	79	28	17	20	46	64	79	103	129	154	151	212	224
1999	4,687	92	33	16	18	45	54	63	113	147	162	153	192	215
Male														
2002	2,587	39	11	5	7	25	51	64	71	77	118	117	143	145
2001	2,401	40	14	9	8	22	35	51	68	83	102	120	132	127
2000	2,393	39	20	9	16	34	42	55	69	79	98	95	139	143
1999	2,392	40	20	9	12	25	35	49	79	95	113	97	128	126
Female														
2002	2,565	42	12	6	9	9	17	21	32	47	56	74	82	99
2001	2,498	35	17	8	8	17	12	22	38	40	43	51	67	97
2000	2,399	40	8	8	4	12	22	24	34	50	56	56	73	81
1999	2,295	52	13	7	6	20	19	14	34	52	49	56	64	89
Non-Hispanic[1]														
Both sexes														
2002	163,970	390	134	73	127	328	405	636	976	1,636	2,664	3,646	4,617	5,650
2001	161,625	372	183	75	138	264	437	610	940	1,640	2,544	3,516	4,561	5,394
2000	160,222	375	139	78	132	309	432	604	954	1,569	2,582	3,367	4,327	5,172
1999	160,711	397	143	83	130	352	427	613	981	1,723	2,594	3,337	4,183	5,229
Male														
2002	71,455	209	71	33	80	225	255	382	617	967	1,682	2,343	2,925	3,430
2001	69,830	181	103	32	74	169	270	359	554	1,010	1,566	2,227	2,803	3,258
2000	69,547	197	78	49	76	184	257	349	572	948	1,606	2,172	2,728	3,229
1999	69,984	215	80	45	70	224	247	349	584	1,075	1,597	2,118	2,619	3,279
Female														
2002	92,515	181	63	40	47	103	150	254	359	669	982	1,303	1,692	2,220
2001	91,795	191	80	43	64	95	167	251	386	630	978	1,289	1,758	2,136
2000	90,675	178	61	29	56	125	175	255	382	621	976	1,195	1,599	1,943
1999	90,727	182	63	38	60	128	180	264	397	648	997	1,219	1,564	1,950
Non-Hispanic White														
Both sexes														
2002	141,712	234	86	50	80	216	237	361	575	1,045	1,843	2,453	3,250	4,234
2001	139,716	219	103	48	92	164	258	363	574	1,039	1,657	2,382	3,249	3,986

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)										
All origins										
Female										
2001	2,788	4,240	6,649	10,787	15,649	19,748	16,592	7,625	1,873	2
2000	2,797	4,291	6,767	11,019	15,342	19,031	16,499	7,608	1,847	3
1999	2,834	4,164	7,088	11,122	15,462	19,135	16,015	7,413	1,802	2
Hispanic										
Both sexes										
2002	263	377	480	628	690	599	463	220	53	-
2001	292	389	467	594	612	583	450	198	46	2
2000	269	349	468	584	565	597	438	176	39	1
1999	286	356	459	536	566	554	403	186	37	1
Male										
2002	151	206	278	316	310	233	147	64	9	-
2001	169	213	243	279	277	215	139	47	6	2
2000	156	190	256	264	246	221	156	56	9	1
1999	172	206	260	273	240	216	139	45	12	1
Female										
2002	112	171	202	312	380	366	316	156	44	-
2001	123	176	224	315	335	368	311	151	40	-
2000	113	159	212	320	319	376	282	120	30	-
1999	114	150	199	263	326	338	264	141	25	-
Non-Hispanic[1]										
Both sexes										
2002	6,752	8,979	13,782	20,603	27,739	30,132	23,012	9,536	2,150	3
2001	6,586	9,180	13,957	20,679	26,982	29,694	22,285	9,432	2,153	3
2000	6,651	9,508	14,296	20,893	26,231	28,859	22,182	9,393	2,163	6
1999	6,801	9,652	15,033	21,099	26,397	28,800	21,445	9,203	2,084	5
Male										
2002	4,024	5,056	7,416	10,122	11,967	10,917	6,403	2,001	328	2
2001	3,923	5,127	7,544	10,230	11,711	10,358	6,028	1,977	324	2
2000	3,974	5,380	7,758	10,222	11,238	10,245	6,011	1,921	350	3
1999	4,091	5,650	8,162	10,263	11,292	10,043	5,728	1,939	311	3
Female										
2002	2,728	3,923	6,366	10,481	15,772	19,215	16,609	7,535	1,822	1
2001	2,663	4,053	6,413	10,449	15,271	19,336	16,257	7,455	1,829	1
2000	2,677	4,128	6,538	10,671	14,993	18,614	16,171	7,472	1,813	3
1999	2,710	4,002	6,871	10,836	15,105	18,757	15,717	7,264	1,773	2
Non-Hispanic White										
Both sexes										
2002	5,234	7,235	11,601	18,032	25,108	27,835	21,321	8,790	1,890	2
2001	5,146	7,348	11,823	18,059	24,427	27,403	20,729	8,728	1,916	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other heart diseases (I26-I51)														
Non-Hispanic White														
Both sexes														
2000	138,745	238	87	46	76	184	244	354	594	988	1,729	2,285	3,077	3,908
1999	138,869	240	87	56	84	235	260	360	612	1,103	1,711	2,238	3,000	3,857
Male														
2002	61,048	128	51	23	51	149	155	217	372	629	1,207	1,623	2,125	2,618
2001	59,513	116	58	21	51	102	159	216	339	650	1,081	1,562	2,075	2,444
2000	59,461	127	51	30	47	106	151	210	366	617	1,109	1,539	1,966	2,489
1999	59,549	136	51	30	43	144	147	209	371	708	1,101	1,482	1,913	2,469
Female														
2002	80,664	106	35	27	29	67	82	144	203	416	636	830	1,125	1,616
2001	80,203	103	45	27	41	62	99	147	235	389	576	820	1,174	1,542
2000	79,284	111	36	16	29	78	93	144	228	371	620	746	1,111	1,419
1999	79,320	104	36	26	41	91	113	151	241	395	610	756	1,087	1,388
Non-Hispanic Black														
Both sexes														
2002	19,641	135	38	16	41	95	151	250	358	545	761	1,105	1,265	1,287
2001	19,496	130	70	25	38	91	161	228	339	537	833	1,050	1,214	1,291
2000	19,159	113	46	27	45	112	166	233	326	524	790	1,013	1,155	1,152
1999	19,531	132	48	26	40	109	153	232	344	572	823	1,016	1,091	1,257
Male														
2002	9,038	72	16	6	24	61	90	149	214	308	430	667	735	736
2001	9,070	59	39	10	21	60	100	133	194	318	450	605	674	746
2000	8,883	56	24	15	24	67	91	128	185	298	449	590	705	676
1999	9,258	66	27	14	24	76	90	132	199	335	462	589	658	739
Female														
2002	10,603	63	22	10	17	34	61	101	144	237	331	438	530	551
2001	10,426	71	31	15	17	31	61	95	145	219	383	445	540	545
2000	10,276	57	22	12	21	45	75	105	141	226	341	423	450	476
1999	10,273	66	21	12	16	33	63	100	145	237	361	427	433	518
Origin not stated[2]														
Both sexes														
2002	418	2	-	-	-	-	1	2	6	9	13	12	20	19
2001	425	2	1	-	-	2	5	4	3	7	13	21	16	19
2000	414	3	1	1	-	-	2	4	5	8	15	18	12	21
1999	430	1	1	-	1	1	1	4	5	9	12	12	11	21
Male														
2002	188	1	-	-	-	-	1	-	4	7	9	8	14	12
2001	208	1	-	-	-	2	2	3	2	6	10	14	11	8
2000	182	2	-	-	-	-	1	3	1	6	9	11	6	12
1999	225	-	-	-	1	1	1	2	4	6	8	8	9	14

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)										
Non-Hispanic White										
Both sexes										
2000	5,181	7,694	12,087	18,323	23,851	26,594	20,614	8,654	1,932	5
1999	5,251	7,809	12,720	18,469	23,937	26,462	20,015	8,501	1,859	3
Male										
2002	3,188	4,135	6,330	8,960	10,918	10,148	5,895	1,841	283	2
2001	3,118	4,199	6,454	9,008	10,619	9,571	5,578	1,810	280	2
2000	3,159	4,422	6,680	9,052	10,283	9,453	5,566	1,735	300	3
1999	3,211	4,641	7,015	9,074	10,291	9,187	5,298	1,762	264	2
Female										
2002	2,046	3,100	5,271	9,072	14,190	17,687	15,426	6,949	1,607	-
2001	2,028	3,149	5,369	9,051	13,808	17,832	15,151	6,918	1,636	1
2000	2,022	3,272	5,407	9,271	13,568	17,141	15,048	6,919	1,632	2
1999	2,040	3,168	5,705	9,395	13,646	17,275	14,717	6,739	1,595	1
Non-Hispanic Black										
Both sexes										
2002	1,384	1,581	1,895	2,228	2,255	1,930	1,449	648	223	1
2001	1,317	1,648	1,874	2,304	2,204	1,986	1,346	605	205	-
2000	1,333	1,636	1,980	2,278	2,080	1,973	1,338	629	209	1
1999	1,421	1,668	2,079	2,316	2,138	2,037	1,224	613	190	2
Male										
2002	754	828	944	975	879	596	398	121	35	-
2001	737	830	936	1,057	921	660	346	135	39	-
2000	734	860	942	1,012	821	664	353	146	43	-
1999	806	909	1,027	1,016	854	699	349	146	40	1
Female										
2002	630	753	951	1,253	1,376	1,334	1,051	527	188	1
2001	580	818	938	1,247	1,283	1,326	1,000	470	166	-
2000	599	776	1,038	1,266	1,259	1,309	985	483	166	1
1999	615	759	1,052	1,300	1,284	1,338	875	467	150	1
Origin not stated[2]										
Both sexes										
2002	18	21	41	45	71	63	42	23	7	3
2001	20	27	36	52	57	66	44	23	5	2
2000	25	18	40	47	47	63	57	22	4	1
1999	28	26	46	54	60	74	46	12	4	1
Male										
2002	9	13	22	25	31	17	8	5	1	1
2001	18	16	24	29	14	22	20	4	1	1
2000	18	14	23	19	17	22	11	6	-	1
1999	18	14	28	31	29	34	12	4	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other heart diseases (I26-I51)														
Origin not stated[2]														
Female														
2002	230	1	-	-	-	-	-	2	2	2	4	4	6	7
2001	217	1	1	-	-	-	3	1	1	1	3	7	5	11
2000	232	1	1	1	-	-	1	1	4	2	6	7	6	9
1999	205	1	1	-	-	-	-	2	1	3	4	4	2	7
Acute and subacute endocarditis (I33)														
All origins														
Both sexes														
2002	1,154	2	2	1	1	2	5	16	28	26	64	84	90	73
2001	1,178	1	-	1	2	3	5	9	20	46	71	93	93	72
2000	1,172	3	1	2	1	5	8	13	27	41	75	89	97	88
1999	1,116	2	1	-	-	2	5	12	18	54	60	68	75	93
Male														
2002	643	1	1	-	1	2	4	9	22	19	41	55	62	42
2001	668	1	-	-	1	2	3	5	12	25	37	68	61	39
2000	665	2	-	1	-	1	6	7	17	24	49	59	61	52
1999	639	2	1	-	-	-	3	9	15	34	41	49	54	62
Female														
2002	511	1	1	1	-	-	1	7	6	7	23	29	28	31
2001	510	-	-	1	1	1	2	4	8	21	34	25	32	33
2000	507	1	1	1	1	4	2	6	10	17	26	30	36	36
1999	477	-	-	-	-	2	2	3	3	20	19	19	21	31
Hispanic														
Both sexes														
2002	69	1	1	-	-	-	-	2	5	2	3	9	4	6
2001	56	-	-	1	-	-	1	1	5	4	4	5	5	3
2000	70	-	1	1	-	-	-	2	4	5	6	4	12	4
1999	63	-	1	-	-	1	1	1	5	4	6	3	4	6
Male														
2002	45	-	-	-	-	-	-	2	4	2	3	7	4	4
2001	34	-	-	-	-	-	1	1	4	4	2	4	4	-
2000	39	-	-	-	-	-	-	1	3	3	4	2	8	3
1999	42	-	1	-	-	-	-	1	4	2	4	2	4	4
Female														
2002	24	1	1	-	-	-	-	-	1	-	-	2	-	2
2001	22	-	-	1	-	-	-	-	1	-	2	1	1	3
2000	31	-	1	1	-	-	-	1	1	2	2	2	4	1
1999	21	-	-	-	-	1	1	-	1	2	2	1	-	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)										
Origin not stated[2]										
Female										
2002	9	8	19	20	40	46	34	18	6	2
2001	2	11	12	23	43	44	24	19	4	1
2000	7	4	17	28	30	41	46	16	4	-
1999	10	12	18	23	31	40	34	8	4	-
Acute and subacute endocarditis (I33)										
All origins										
Both sexes										
2002	95	107	131	161	145	76	36	8	1	-
2001	88	102	145	162	154	79	30	2	-	-
2000	77	105	119	166	136	83	30	5	1	-
1999	78	103	137	155	134	92	21	5	1	-
Male										
2002	49	66	67	78	75	31	15	2	1	-
2001	56	49	83	94	83	37	11	1	-	-
2000	44	59	63	95	74	40	9	2	-	-
1999	41	51	67	83	71	46	10	-	-	-
Female										
2002	46	41	64	83	70	45	21	6	-	-
2001	32	53	62	68	71	42	19	1	-	-
2000	33	46	56	71	62	43	21	3	1	-
1999	37	52	70	72	63	46	11	5	1	-
Hispanic										
Both sexes										
2002	6	7	7	7	6	1	1	1	-	-
2001	6	3	6	7	3	1	1	-	-	-
2000	4	8	10	6	1	1	1	-	-	-
1999	4	6	7	9	3	2	-	-	-	-
Male										
2002	3	4	5	3	3	1	-	-	-	-
2001	4	1	4	4	-	-	1	-	-	-
2000	1	5	6	1	1	-	1	-	-	-
1999	1	3	6	6	2	2	-	-	-	-
Female										
2002	3	3	2	4	3	-	1	1	-	-
2001	2	2	2	3	3	1	-	-	-	-
2000	3	3	4	5	-	1	-	-	-	-
1999	3	3	1	3	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and subacute endocarditis (I33)														
Non-Hispanic[1]														
Both sexes														
2002	1,082	1	1	1	1	2	5	14	23	24	61	74	85	66
2001	1,117	1	-	-	2	3	4	8	15	42	66	87	87	68
2000	1,096	3	-	1	1	5	8	11	23	34	68	85	85	84
1999	1,049	2	-	-	-	1	4	10	13	50	54	64	71	86
Male														
2002	596	1	1	-	1	2	4	7	18	17	38	48	57	37
2001	631	1	-	-	1	2	2	4	8	21	34	63	57	38
2000	625	2	-	1	-	1	6	6	14	20	45	57	53	49
1999	594	2	-	-	-	-	3	7	11	32	37	46	50	57
Female														
2002	486	-	-	1	-	-	1	7	5	7	23	26	28	29
2001	486	-	-	-	1	1	2	4	7	21	32	24	30	30
2000	471	1	-	-	1	4	2	5	9	14	23	28	32	35
1999	455	-	-	-	-	1	1	3	2	18	17	18	21	29
Non-Hispanic White														
Both sexes														
2002	862	-	1	-	-	2	4	9	18	19	43	39	56	40
2001	859	1	-	-	2	1	4	7	8	26	39	59	55	49
2000	852	1	-	-	1	1	5	8	13	20	46	53	53	55
1999	814	1	-	-	-	1	3	8	9	28	31	38	44	51
Male														
2002	481	-	1	-	-	2	3	5	15	13	28	27	41	21
2001	482	1	-	-	1	1	2	3	4	12	22	41	38	27
2000	496	1	-	-	-	-	3	5	10	10	32	37	36	35
1999	466	1	-	-	-	-	2	5	7	19	20	30	29	37
Female														
2002	381	-	-	-	-	-	1	4	3	6	15	12	15	19
2001	377	-	-	-	1	-	2	4	4	14	17	18	17	22
2000	356	-	-	-	1	1	2	3	3	10	14	16	17	20
1999	348	-	-	-	-	1	1	3	2	9	11	8	15	14
Non-Hispanic Black														
Both sexes														
2002	197	-	-	1	-	-	1	5	5	5	17	29	28	25
2001	233	-	-	-	-	1	-	1	7	15	26	27	30	19
2000	213	2	-	-	-	3	3	3	8	11	20	30	27	26
1999	208	-	-	-	-	-	1	2	4	20	21	25	23	30
Male														
2002	104	-	-	-	-	-	1	2	3	4	10	18	15	16
2001	134	-	-	-	-	-	-	1	4	8	12	21	18	11
2000	111	1	-	-	-	1	3	1	2	9	12	20	15	12

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)										
Non-Hispanic[1]										
Both sexes										
2002	89	100	124	154	139	75	35	7	1	-
2001	82	99	139	154	151	78	29	2	-	-
2000	73	97	108	159	134	82	29	5	1	-
1999	74	97	129	146	131	90	21	5	1	-
Male										
2002	46	62	62	75	72	30	15	2	1	-
2001	52	48	79	90	83	37	10	1	-	-
2000	43	54	57	94	73	40	8	2	-	-
1999	40	48	61	77	69	44	10	-	-	-
Female										
2002	43	38	62	79	67	45	20	5	-	-
2001	30	51	60	64	68	41	19	1	-	-
2000	30	43	51	65	61	42	21	3	1	-
1999	34	49	68	69	62	46	11	5	1	-
Non-Hispanic White										
Both sexes										
2002	73	76	103	138	132	69	32	7	1	-
2001	58	71	111	128	137	75	26	2	-	-
2000	59	81	92	141	120	73	25	5	-	-
1999	56	76	106	133	119	85	20	5	-	-
Male										
2002	40	49	53	71	68	27	14	2	1	-
2001	38	34	59	79	75	35	9	1	-	-
2000	37	42	51	86	65	37	7	2	-	-
1999	31	42	51	75	64	43	10	-	-	-
Female										
2002	33	27	50	67	64	42	18	5	-	-
2001	20	37	52	49	62	40	17	1	-	-
2000	22	39	41	55	55	36	18	3	-	-
1999	25	34	55	58	55	42	10	5	-	-
Non-Hispanic Black										
Both sexes										
2002	16	22	16	14	5	5	3	-	-	-
2001	22	21	22	25	12	2	3	-	-	-
2000	13	15	15	16	10	6	4	-	1	-
1999	17	21	21	10	7	5	1	-	-	-
Male										
2002	6	11	8	3	3	3	1	-	-	-
2001	13	10	16	11	7	1	1	-	-	-
2000	6	11	5	6	5	1	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and subacute endocarditis (I33)														
Non-Hispanic Black														
Male														
1999	115	-	-	-	-	-	1	2	4	12	15	15	19	18
Female														
2002	93	-	-	1	-	-	-	3	2	1	7	11	13	9
2001	99	-	-	-	-	1	-	-	3	7	14	6	12	8
2000	102	1	-	-	-	2	-	2	6	2	8	10	12	14
1999	93	-	-	-	-	-	-	-	-	8	6	10	4	12
Origin not stated[2]														
Both sexes														
2002	3	-	-	-	-	-	-	-	-	-	-	1	1	1
2001	5	-	-	-	-	-	-	-	-	-	1	1	1	1
2000	6	-	-	-	-	-	-	-	-	2	1	-	-	-
1999	4	-	-	-	-	-	-	1	-	-	-	1	-	1
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	1	1
2001	3	-	-	-	-	-	-	-	-	-	1	1	-	1
2000	1	-	-	-	-	-	-	-	-	1	-	-	-	-
1999	3	-	-	-	-	-	-	1	-	-	-	1	-	1
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	1	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	5	-	-	-	-	-	-	-	-	1	1	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Diseases of pericardium and acute myocarditis (I30-I31,I40)														
All origins														
Both sexes														
2002	848	26	22	12	10	17	19	16	30	35	40	56	51	66
2001	839	32	26	12	9	14	26	20	30	43	50	57	54	54
2000	767	20	24	9	11	17	21	29	24	45	41	45	40	47
1999	890	27	25	11	10	12	20	19	31	54	60	59	46	63
Male														
2002	434	9	11	4	3	12	12	9	23	17	32	40	29	34
2001	401	18	15	2	5	7	15	16	17	24	17	30	29	28
2000	404	10	13	4	5	10	11	17	12	30	28	28	18	31
1999	442	13	14	5	2	8	12	14	20	32	32	31	25	40
Female														
2002	414	17	11	8	7	5	7	7	7	18	8	16	22	32
2001	438	14	11	10	4	7	11	4	13	19	33	27	25	26

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)										
Non-Hispanic Black										
Male										
1999	8	6	9	1	4	1	-	-	-	-
Female										
2002	10	11	8	11	2	2	2	-	-	-
2001	9	11	6	14	5	1	2	-	-	-
2000	7	4	10	10	5	5	3	-	1	-
1999	9	15	12	9	3	4	1	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	1	1	1	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	1	1	1	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All origins										
Both sexes										
2002	52	55	88	84	73	65	25	6	-	-
2001	52	47	73	87	67	49	28	7	1	1
2000	42	52	79	82	63	43	24	8	1	-
1999	62	69	77	88	72	56	25	4	-	-
Male										
2002	35	29	42	34	26	25	6	2	-	-
2001	33	28	31	39	22	14	8	2	-	1
2000	21	24	43	42	31	17	7	1	1	-
1999	30	34	35	38	27	19	11	-	-	-
Female										
2002	17	26	46	50	47	40	19	4	-	-
2001	19	19	42	48	45	35	20	5	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)														
All origins														
Female														
2000	363	10	11	5	6	7	10	12	12	15	13	17	22	16
1999	448	14	11	6	8	4	8	5	11	22	28	28	21	23
Hispanic														
Both sexes														
2002	64	4	2	2	1	1	4	-	3	4	2	6	4	7
2001	58	2	4	1	1	1	3	5	6	1	4	3	3	3
2000	61	5	4	1	-	3	2	6	3	5	5	2	2	5
1999	64	2	3	2	2	3	7	2	6	7	2	8	1	1
Male														
2002	35	3	-	-	-	1	2	-	3	2	1	4	3	5
2001	25	1	2	-	1	-	2	5	1	1	2	1	-	-
2000	29	2	4	-	-	-	1	2	2	3	3	1	1	2
1999	30	1	1	2	2	3	2	2	4	5	2	1	1	1
Female														
2002	29	1	2	2	1	-	2	-	-	2	1	2	1	2
2001	33	1	2	1	-	1	1	-	5	-	2	2	3	3
2000	32	3	-	1	-	3	1	4	1	2	2	1	1	3
1999	34	1	2	-	-	-	5	-	2	2	-	7	-	-
Non-Hispanic[1]														
Both sexes														
2002	783	22	20	10	9	16	15	16	27	31	38	50	47	58
2001	779	30	22	11	8	13	23	14	24	41	46	54	51	51
2000	700	15	19	8	11	14	18	23	21	40	36	43	38	42
1999	824	25	22	9	8	9	13	17	24	47	57	51	45	62
Male														
2002	398	6	11	4	3	11	10	9	20	15	31	36	26	28
2001	375	17	13	2	4	7	13	10	16	23	15	29	29	28
2000	374	8	9	4	5	10	10	15	10	27	25	27	17	29
1999	410	12	13	3	-	5	10	12	15	27	29	30	24	39
Female														
2002	385	16	9	6	6	5	5	7	7	16	7	14	21	30
2001	404	13	9	9	4	6	10	4	8	18	31	25	22	23
2000	326	7	10	4	6	4	8	8	11	13	11	16	21	13
1999	414	13	9	6	8	4	3	5	9	20	28	21	21	23
Non-Hispanic White														
Both sexes														
2002	601	9	11	6	3	9	12	14	16	18	31	37	32	41
2001	600	13	10	8	6	9	14	9	17	26	29	40	35	39

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All origins										
Female										
2000	21	28	36	40	32	26	17	7	-	-
1999	32	35	42	50	45	37	14	4	-	-
Hispanic										
Both sexes										
2002	7	3	4	3	2	5	-	-	-	-
2001	3	3	3	4	3	3	2	-	-	-
2000	4	3	2	3	4	1	-	1	-	-
1999	1	3	4	4	4	1	-	1	-	-
Male										
2002	4	2	1	-	1	3	-	-	-	-
2001	3	1	2	2	-	1	-	-	-	-
2000	1	-	-	2	4	-	-	1	-	-
1999	-	-	1	1	1	-	-	-	-	-
Female										
2002	3	1	3	3	1	2	-	-	-	-
2001	-	2	1	2	3	2	2	-	-	-
2000	3	3	2	1	-	1	-	-	-	-
1999	1	3	3	3	3	1	-	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	45	52	84	81	71	60	25	6	-	-
2001	49	44	70	83	64	46	26	7	1	1
2000	36	49	77	77	59	42	24	7	1	-
1999	61	66	73	84	68	55	25	3	-	-
Male										
2002	31	27	41	34	25	22	6	2	-	-
2001	30	27	29	37	22	13	8	2	-	1
2000	19	24	43	40	27	17	7	-	1	-
1999	30	34	34	37	26	19	11	-	-	-
Female										
2002	14	25	43	47	46	38	19	4	-	-
2001	19	17	41	46	42	33	18	5	1	-
2000	17	25	34	37	32	25	17	7	-	-
1999	31	32	39	47	42	36	14	3	-	-
Non-Hispanic White										
Both sexes										
2002	35	45	71	64	62	57	23	5	-	-
2001	40	33	65	76	58	40	25	7	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)														
Non-Hispanic White														
Both sexes														
2000	528	6	11	7	2	12	8	15	10	23	26	24	31	34
1999	639	10	10	5	4	6	8	11	15	34	36	39	34	52
Male														
2002	308	3	7	2	1	6	8	8	13	7	26	25	18	20
2001	286	10	6	2	3	3	7	6	12	14	13	20	21	22
2000	280	4	7	4	1	8	4	9	5	16	18	16	14	26
1999	320	6	4	2	-	4	6	8	9	19	21	23	17	33
Female														
2002	293	6	4	4	2	3	4	6	3	11	5	12	14	21
2001	314	3	4	6	3	6	7	3	5	12	16	20	14	17
2000	248	2	4	3	1	4	4	6	5	7	8	8	17	8
1999	319	4	6	3	4	2	2	3	6	15	15	16	17	19
Non-Hispanic Black														
Both sexes														
2002	154	12	7	2	6	5	3	1	9	10	7	12	14	14
2001	157	14	12	3	1	4	9	5	5	10	14	14	16	11
2000	140	7	7	1	7	1	6	6	10	14	10	14	7	7
1999	158	10	11	4	4	3	3	6	8	12	18	9	11	9
Male														
2002	75	3	3	1	2	3	2	-	6	5	5	10	7	8
2001	78	5	7	-	1	4	6	4	3	5	2	9	8	5
2000	72	4	2	-	3	1	2	5	5	9	7	7	3	2
1999	76	4	9	1	-	1	2	4	6	7	6	5	7	6
Female														
2002	79	9	4	1	4	2	1	1	3	5	2	2	7	6
2001	79	9	5	3	-	-	3	1	2	5	12	5	8	6
2000	68	3	5	1	4	-	4	1	5	5	3	7	4	5
1999	82	6	2	3	4	2	1	2	2	5	12	4	4	3
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	2	-	-	-	-	-	-	1	-	1	-	-	-	-
2000	6	-	1	-	-	-	1	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	1	-	1	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	1	-	-	-	-	-	-	1	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	1	-	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
Non-Hispanic White										
Both sexes										
2000	28	38	65	65	57	37	21	7	1	-
1999	52	54	61	73	61	49	22	3	-	-
Male										
2002	25	23	37	30	21	22	5	1	-	-
2001	23	19	29	33	20	12	8	2	-	1
2000	15	15	37	32	27	15	6	-	1	-
1999	27	27	29	34	24	17	10	-	-	-
Female										
2002	10	22	34	34	41	35	18	4	-	-
2001	17	14	36	43	38	28	17	5	-	-
2000	13	23	28	33	30	22	15	7	-	-
1999	25	27	32	39	37	32	12	3	-	-
Non-Hispanic Black										
Both sexes										
2002	8	7	12	14	6	3	2	-	-	-
2001	8	9	4	6	5	5	1	-	1	-
2000	8	10	8	8	2	5	2	-	-	-
1999	8	10	10	10	6	4	2	-	-	-
Male										
2002	5	4	3	4	3	-	1	-	-	-
2001	6	8	-	3	2	-	-	-	-	-
2000	4	8	3	5	-	2	-	-	-	-
1999	3	6	5	2	2	-	-	-	-	-
Female										
2002	3	3	9	10	3	3	1	-	-	-
2001	2	1	4	3	3	5	1	-	1	-
2000	4	2	5	3	2	3	2	-	-	-
1999	5	4	5	8	4	4	2	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	2	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)														
Origin not stated[2]														
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	5	-	1	-	-	-	1	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Heart failure (I50)														
All origins														
Both sexes														
2002	56,494	23	11	3	7	10	17	33	68	102	180	355	638	906
2001	56,934	24	17	8	7	19	15	41	51	115	192	353	601	926
2000	55,704	31	9	10	11	21	13	27	36	77	185	283	528	828
1999	54,913	11	10	5	7	14	17	27	38	88	184	298	484	853
Male														
2002	21,698	14	6	1	4	6	9	22	38	56	117	210	407	526
2001	21,632	16	5	5	3	11	5	25	31	76	106	227	361	556
2000	21,175	22	7	4	5	11	9	15	21	40	119	191	305	479
1999	20,755	5	7	3	3	12	11	13	19	52	112	186	300	506
Female														
2002	34,796	9	5	2	3	4	8	11	30	46	63	145	231	380
2001	35,302	8	12	3	4	8	10	16	20	39	86	126	240	370
2000	34,529	9	2	6	6	10	4	12	15	37	66	92	223	349
1999	34,158	6	3	2	4	2	6	14	19	36	72	112	184	347
Hispanic														
Both sexes														
2002	1,412	4	-	-	2	-	-	2	1	4	9	14	29	35
2001	1,364	8	3	-	-	1	-	3	6	11	10	18	25	31
2000	1,270	7	1	2	-	-	-	1	2	4	6	7	15	38
1999	1,242	3	4	-	1	1	2	2	2	7	11	10	19	32
Male														
2002	593	2	-	-	2	-	-	2	-	-	6	4	16	22
2001	565	6	1	-	-	1	-	2	4	9	6	15	16	17
2000	535	5	1	1	-	-	-	1	1	1	5	3	6	24
1999	522	1	3	-	-	1	2	2	1	5	7	6	10	19
Female														
2002	819	2	-	-	-	-	-	-	1	4	3	10	13	13
2001	799	2	2	-	-	-	-	1	2	2	4	3	9	14
2000	735	2	-	1	-	-	-	-	1	3	1	4	9	14
1999	720	2	1	-	1	-	-	-	1	2	4	4	9	13

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
Origin not stated[2]										
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	2	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Heart failure (I50)										
All origins										
Both sexes										
2002	1,436	2,192	3,976	6,603	10,307	12,620	10,776	5,031	1,200	-
2001	1,448	2,270	4,004	6,879	10,255	12,793	10,735	4,971	1,209	1
2000	1,383	2,284	4,119	6,936	9,884	12,315	10,543	4,969	1,208	4
1999	1,254	2,240	4,284	6,926	9,845	12,169	10,197	4,850	1,111	1
Male										
2002	819	1,200	2,054	3,192	4,381	4,423	3,000	1,035	178	-
2001	791	1,235	2,109	3,300	4,372	4,320	2,886	1,020	172	-
2000	785	1,243	2,171	3,286	4,212	4,256	2,819	982	191	2
1999	722	1,232	2,294	3,264	4,054	4,155	2,636	991	177	1
Female										
2002	617	992	1,922	3,411	5,926	8,197	7,776	3,996	1,022	-
2001	657	1,035	1,895	3,579	5,883	8,473	7,849	3,951	1,037	1
2000	598	1,041	1,948	3,650	5,672	8,059	7,724	3,987	1,017	2
1999	532	1,008	1,990	3,662	5,791	8,014	7,561	3,859	934	-
Hispanic										
Both sexes										
2002	56	93	142	200	251	232	194	113	31	-
2001	72	90	123	178	226	231	214	93	21	-
2000	62	73	130	171	194	243	219	74	21	-
1999	55	71	131	167	186	232	201	92	13	-
Male										
2002	32	51	70	100	109	85	56	31	5	-
2001	35	55	54	69	100	86	61	26	2	-
2000	38	44	65	71	80	86	76	21	6	-
1999	28	40	72	87	72	78	66	20	2	-
Female										
2002	24	42	72	100	142	147	138	82	26	-
2001	37	35	69	109	126	145	153	67	19	-
2000	24	29	65	100	114	157	143	53	15	-
1999	27	31	59	80	114	154	135	72	11	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Heart failure (I50)														
Non-Hispanic[1]														
Both sexes														
2002	54,932	19	11	3	5	10	17	31	66	97	171	339	604	869
2001	55,437	16	14	8	7	18	15	37	45	104	182	331	575	887
2000	54,307	24	8	8	11	21	13	26	34	73	179	276	513	789
1999	53,528	8	6	5	6	13	15	25	36	80	173	286	464	818
Male														
2002	21,050	12	6	1	2	6	9	20	38	55	111	204	388	502
2001	21,012	10	4	5	3	10	5	22	27	67	100	210	345	536
2000	20,599	17	6	3	5	11	9	14	20	39	114	188	299	455
1999	20,170	4	4	3	3	11	9	11	18	47	105	179	290	485
Female														
2002	33,882	7	5	2	3	4	8	11	28	42	60	135	216	367
2001	34,425	6	10	3	4	8	10	15	18	37	82	121	230	351
2000	33,708	7	2	5	6	10	4	12	14	34	65	88	214	334
1999	33,358	4	2	2	3	2	6	14	18	33	68	107	174	333
Non-Hispanic White														
Both sexes														
2002	49,162	11	8	3	2	5	11	14	35	56	104	196	397	644
2001	49,788	11	12	5	6	12	5	22	27	65	102	215	393	656
2000	48,782	13	4	5	9	12	5	14	21	40	114	173	363	588
1999	47,977	5	4	3	5	8	7	11	19	50	106	173	307	594
Male														
2002	18,705	8	5	1	1	2	5	9	19	33	72	125	262	381
2001	18,730	6	4	3	2	8	1	10	15	40	63	149	243	410
2000	18,386	10	4	2	4	5	5	8	12	24	79	124	218	359
1999	17,903	2	2	2	3	7	4	4	10	29	71	110	196	369
Female														
2002	30,457	3	3	2	1	3	6	5	16	23	32	71	135	263
2001	31,058	5	8	2	4	4	4	12	12	25	39	66	150	246
2000	30,396	3	-	3	5	7	-	6	9	16	35	49	145	229
1999	30,074	3	2	1	2	1	3	7	9	21	35	63	111	225
Non-Hispanic Black														
Both sexes														
2002	5,143	8	3	-	3	3	6	16	29	40	64	139	192	204
2001	5,094	5	2	3	-	6	10	15	17	35	77	111	173	221
2000	4,936	11	4	2	2	8	5	12	12	32	62	98	143	189
1999	4,961	3	2	2	1	4	8	13	17	28	65	109	148	212
Male														
2002	2,055	4	1	-	1	2	4	10	17	22	37	77	115	108
2001	2,036	4	-	2	-	2	4	12	11	26	35	57	96	120
2000	1,942	7	2	1	1	5	3	6	7	14	33	60	78	90

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)										
Non-Hispanic[1]										
Both sexes										
2002	1,376	2,095	3,819	6,388	10,028	12,353	10,561	4,907	1,163	-
2001	1,371	2,170	3,874	6,682	10,006	12,538	10,506	4,866	1,184	1
2000	1,316	2,206	3,979	6,750	9,667	12,050	10,290	4,886	1,185	3
1999	1,189	2,164	4,144	6,740	9,637	11,902	9,969	4,750	1,097	1
Male										
2002	786	1,147	1,977	3,088	4,256	4,328	2,939	1,003	172	-
2001	751	1,176	2,049	3,221	4,268	4,226	2,817	991	169	-
2000	742	1,195	2,101	3,210	4,125	4,164	2,736	960	185	1
1999	688	1,189	2,217	3,168	3,972	4,058	2,564	969	175	1
Female										
2002	590	948	1,842	3,300	5,772	8,025	7,622	3,904	991	-
2001	620	994	1,825	3,461	5,738	8,312	7,689	3,875	1,015	1
2000	574	1,011	1,878	3,540	5,542	7,886	7,554	3,926	1,000	2
1999	501	975	1,927	3,572	5,665	7,844	7,405	3,781	922	-
Non-Hispanic White										
Both sexes										
2002	1,075	1,695	3,219	5,618	9,107	11,477	9,844	4,600	1,041	-
2001	1,074	1,751	3,301	5,918	9,115	11,652	9,836	4,541	1,068	1
2000	1,023	1,795	3,381	5,998	8,829	11,171	9,599	4,547	1,075	3
1999	943	1,751	3,551	5,924	8,778	10,984	9,358	4,391	1,005	-
Male										
2002	619	933	1,687	2,752	3,911	4,044	2,739	945	152	-
2001	609	963	1,771	2,886	3,899	3,945	2,635	919	149	-
2000	583	1,003	1,825	2,884	3,787	3,870	2,536	883	160	1
1999	553	969	1,951	2,828	3,644	3,725	2,388	885	151	-
Female										
2002	456	762	1,532	2,866	5,196	7,433	7,105	3,655	889	-
2001	465	788	1,530	3,032	5,216	7,707	7,201	3,622	919	1
2000	440	792	1,556	3,114	5,042	7,301	7,063	3,664	915	2
1999	390	782	1,600	3,096	5,134	7,259	6,970	3,506	854	-
Non-Hispanic Black										
Both sexes										
2002	283	361	530	680	815	767	620	272	108	-
2001	273	380	525	690	785	784	588	295	99	-
2000	264	371	538	670	737	781	603	290	102	-
1999	234	380	542	724	744	817	514	315	78	1
Male										
2002	158	190	260	294	308	227	156	47	17	-
2001	128	194	253	299	319	245	147	63	19	-
2000	141	174	246	285	287	254	166	60	22	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Heart failure (I50)														
Non-Hispanic Black														
Male														
1999	2,005	2	2	1	-	3	5	7	8	16	34	67	89	111
Female														
2002	3,088	4	2	-	2	1	2	6	12	18	27	62	77	96
2001	3,058	1	2	1	-	4	6	3	6	9	42	54	77	101
2000	2,994	4	2	1	1	3	2	6	5	18	29	38	65	99
1999	2,956	1	-	1	1	1	3	6	9	12	31	42	59	101
Origin not stated[2]														
Both sexes														
2002	150	-	-	-	-	-	-	-	1	1	-	2	5	2
2001	133	-	-	-	-	-	-	1	-	-	-	4	1	8
2000	127	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	143	-	-	-	-	-	-	-	-	1	-	2	1	3
Male														
2002	55	-	-	-	-	-	-	-	-	1	-	2	3	2
2001	55	-	-	-	-	-	-	1	-	-	-	2	-	3
2000	41	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	63	-	-	-	-	-	-	-	-	-	-	1	-	2
Female														
2002	95	-	-	-	-	-	-	-	1	-	-	-	2	-
2001	78	-	-	-	-	-	-	-	-	-	-	2	1	5
2000	86	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	80	-	-	-	-	-	-	-	-	1	-	1	1	1
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
All origins														
Both sexes														
2002	111,044	422	122	68	125	333	433	658	959	1,606	2,567	3,354	4,083	4,868
2001	107,998	392	172	71	136	269	443	617	948	1,566	2,389	3,205	4,028	4,585
2000	107,785	403	134	75	129	312	456	618	975	1,543	2,450	3,119	3,886	4,454
1999	108,909	450	141	83	132	370	440	622	1,012	1,683	2,464	3,077	3,781	4,456
Male														
2002	51,455	225	64	33	79	230	282	406	609	959	1,619	2,163	2,584	2,985
2001	49,738	187	97	34	73	173	284	367	564	974	1,518	2,036	2,495	2,770
2000	49,878	204	78	49	82	196	274	368	592	939	1,517	2,000	2,489	2,822
1999	50,765	235	78	46	78	230	257	364	613	1,058	1,533	1,957	2,377	2,811
Female														
2002	59,589	197	58	35	46	103	151	252	350	647	948	1,191	1,499	1,883
2001	58,260	205	75	37	63	96	159	250	384	592	871	1,169	1,533	1,815

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)										
Non-Hispanic Black										
Male										
1999	127	198	244	304	273	284	141	69	19	1
Female										
2002	125	171	270	386	507	540	464	225	91	-
2001	145	186	272	391	466	539	441	232	80	-
2000	123	197	292	385	450	527	437	230	80	-
1999	107	182	298	420	471	533	373	246	59	-
Origin not stated[2]										
Both sexes										
2002	4	4	15	15	28	35	21	11	6	-
2001	5	10	7	19	23	24	15	12	4	-
2000	5	5	10	15	23	22	34	9	2	1
1999	10	5	9	19	22	35	27	8	1	-
Male										
2002	1	2	7	4	16	10	5	1	1	-
2001	5	4	6	10	4	8	8	3	1	-
2000	5	4	5	5	7	6	7	1	-	1
1999	6	3	5	9	10	19	6	2	-	-
Female										
2002	3	2	8	11	12	25	16	10	5	-
2001	-	6	1	9	19	16	7	9	3	-
2000	-	1	5	10	16	16	27	8	2	-
1999	4	2	4	10	12	16	21	6	1	-
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All origins										
Both sexes										
2002	5,450	7,023	10,108	14,428	17,975	18,033	12,680	4,734	1,009	6
2001	5,310	7,177	10,238	14,197	17,175	17,422	11,986	4,673	994	5
2000	5,443	7,434	10,487	14,340	16,760	17,078	12,080	4,609	996	4
1999	5,721	7,622	11,040	14,520	16,972	17,111	11,651	4,542	1,013	6
Male										
2002	3,281	3,980	5,553	7,159	7,826	6,688	3,537	1,031	159	3
2001	3,230	4,044	5,588	7,105	7,525	6,224	3,282	1,005	159	4
2000	3,298	4,258	5,760	7,082	7,184	6,175	3,343	998	167	3
1999	3,488	4,553	6,054	7,182	7,409	6,073	3,222	997	146	4
Female										
2002	2,169	3,043	4,555	7,269	10,149	11,345	9,143	3,703	850	3
2001	2,080	3,133	4,650	7,092	9,650	11,198	8,704	3,668	835	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
All origins														
Female														
2000	57,907	199	56	26	47	116	182	250	383	604	933	1,119	1,397	1,632
1999	58,144	215	63	37	54	140	183	258	399	625	931	1,120	1,404	1,645
Hispanic														
Both sexes														
2002	3,607	72	20	9	13	33	64	81	94	114	160	162	188	196
2001	3,421	65	24	15	15	37	43	64	89	107	127	145	166	187
2000	3,391	67	22	13	20	43	62	70	94	115	137	138	183	177
1999	3,318	87	25	14	15	40	44	58	100	129	143	132	168	176
Male														
2002	1,914	34	11	5	5	24	49	60	64	73	108	102	120	114
2001	1,777	33	11	9	7	21	32	43	59	69	92	100	112	110
2000	1,790	32	15	8	16	34	41	51	63	72	86	89	124	114
1999	1,798	38	15	7	10	21	31	44	70	83	100	88	113	102
Female														
2002	1,693	38	9	4	8	9	15	21	30	41	52	60	68	82
2001	1,644	32	13	6	8	16	11	21	30	38	35	45	54	77
2000	1,601	35	7	5	4	9	21	19	31	43	51	49	59	63
1999	1,520	49	10	7	5	19	13	14	30	46	43	44	55	74
Non-Hispanic[1]														
Both sexes														
2002	107,173	348	102	59	112	300	368	575	860	1,484	2,394	3,183	3,881	4,657
2001	104,292	325	147	56	121	230	395	551	856	1,453	2,250	3,044	3,848	4,388
2000	104,119	333	112	61	109	269	393	544	876	1,422	2,299	2,963	3,691	4,257
1999	105,310	362	115	69	116	329	395	561	908	1,546	2,310	2,936	3,603	4,263
Male														
2002	49,411	190	53	28	74	206	232	346	541	880	1,502	2,055	2,454	2,863
2001	47,812	153	86	25	66	150	250	323	503	899	1,417	1,925	2,372	2,656
2000	47,949	170	63	41	66	162	232	314	528	862	1,422	1,900	2,359	2,696
1999	48,810	197	63	39	67	208	225	319	540	969	1,426	1,863	2,255	2,698
Female														
2002	57,762	158	49	31	38	94	136	229	319	604	892	1,128	1,427	1,794
2001	56,480	172	61	31	55	80	145	228	353	554	833	1,119	1,476	1,732
2000	56,170	163	49	20	43	107	161	230	348	560	877	1,063	1,332	1,561
1999	56,500	165	52	30	49	121	170	242	368	577	884	1,073	1,348	1,565
Non-Hispanic White														
Both sexes														
2002	91,087	214	66	41	75	200	210	324	506	952	1,665	2,181	2,765	3,509
2001	88,469	194	81	35	78	142	235	325	522	922	1,487	2,068	2,766	3,242

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All origins										
Female										
2000	2,145	3,176	4,727	7,258	9,576	10,903	8,737	3,611	829	1
1999	2,233	3,069	4,986	7,338	9,563	11,038	8,429	3,545	867	2
Hispanic										
Both sexes										
2002	194	274	327	418	431	361	268	106	22	-
2001	211	293	335	405	380	348	233	105	25	2
2000	199	265	326	404	366	352	218	101	18	1
1999	226	276	317	356	373	319	202	93	24	1
Male										
2002	112	149	202	213	197	144	91	33	4	-
2001	127	156	183	204	177	128	77	21	4	2
2000	116	141	185	190	161	135	79	34	3	1
1999	143	163	181	179	165	136	73	25	10	1
Female										
2002	82	125	125	205	234	217	177	73	18	-
2001	84	137	152	201	203	220	156	84	21	-
2000	83	124	141	214	205	217	139	67	15	-
1999	83	113	136	177	208	183	129	68	14	-
Non-Hispanic[1]										
Both sexes										
2002	5,242	6,732	9,755	13,980	17,501	17,644	12,391	4,616	986	3
2001	5,084	6,867	9,874	13,760	16,761	17,032	11,724	4,557	968	1
2000	5,226	7,156	10,132	13,907	16,371	16,685	11,839	4,495	976	3
1999	5,477	7,325	10,687	14,129	16,561	16,753	11,430	4,445	986	4
Male										
2002	3,161	3,820	5,336	6,925	7,614	6,537	3,443	994	155	2
2001	3,090	3,876	5,387	6,882	7,338	6,082	3,193	983	155	1
2000	3,170	4,107	5,557	6,878	7,013	6,024	3,260	959	164	2
1999	3,333	4,379	5,850	6,981	7,225	5,922	3,143	970	136	2
Female										
2002	2,081	2,912	4,419	7,055	9,887	11,107	8,948	3,622	831	1
2001	1,994	2,991	4,487	6,878	9,423	10,950	8,531	3,574	813	-
2000	2,056	3,049	4,575	7,029	9,358	10,661	8,579	3,536	812	1
1999	2,144	2,946	4,837	7,148	9,336	10,831	8,287	3,475	850	2
Non-Hispanic White										
Both sexes										
2002	4,051	5,419	8,208	12,212	15,807	16,232	11,422	4,178	848	2
2001	3,974	5,493	8,346	11,937	15,117	15,636	10,842	4,178	848	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
Non-Hispanic White														
Both sexes														
2000	88,583	218	72	34	64	159	226	317	550	905	1,543	2,035	2,630	3,231
1999	89,439	224	73	48	75	220	242	330	569	991	1,538	1,988	2,615	3,160
Male														
2002	41,554	117	38	20	49	139	139	195	325	576	1,081	1,446	1,804	2,196
2001	40,015	99	48	16	45	90	149	197	308	584	983	1,352	1,773	1,985
2000	40,299	112	40	24	42	93	139	188	339	567	980	1,362	1,698	2,069
1999	40,860	127	45	26	40	133	135	192	345	641	989	1,319	1,671	2,030
Female														
2002	49,533	97	28	21	26	61	71	129	181	376	584	735	961	1,313
2001	48,454	95	33	19	33	52	86	128	214	338	504	716	993	1,257
2000	48,284	106	32	10	22	66	87	129	211	338	563	673	932	1,162
1999	48,579	97	28	22	35	87	107	138	224	350	549	669	944	1,130
Non-Hispanic Black														
Both sexes														
2002	14,147	115	28	13	32	87	141	228	315	490	673	925	1,031	1,044
2001	14,012	111	56	19	37	80	142	207	310	477	716	898	995	1,040
2000	13,870	93	35	24	36	100	152	212	296	467	698	871	978	930
1999	14,204	119	35	20	35	102	141	211	315	512	719	873	909	1,006
Male														
2002	6,804	65	12	5	21	56	83	137	188	277	378	562	598	604
2001	6,822	50	32	8	20	54	90	116	176	279	401	518	552	610
2000	6,758	44	20	14	20	60	83	116	171	266	397	503	609	572
1999	7,062	60	16	12	24	72	82	119	181	300	407	502	543	604
Female														
2002	7,343	50	16	8	11	31	58	91	127	213	295	363	433	440
2001	7,190	61	24	11	17	26	52	91	134	198	315	380	443	430
2000	7,112	49	15	10	16	40	69	96	125	201	301	368	369	358
1999	7,142	59	19	8	11	30	59	92	134	212	312	371	366	402
Origin not stated[2]														
Both sexes														
2002	264	2	-	-	-	-	1	2	5	8	13	9	14	15
2001	285	2	1	-	-	2	5	2	3	6	12	16	14	10
2000	275	3	-	1	-	-	1	4	5	6	14	18	12	20
1999	281	1	1	-	1	1	1	3	4	8	11	9	10	17
Male														
2002	130	1	-	-	-	-	1	-	4	6	9	6	10	8
2001	149	1	-	-	-	2	2	1	2	6	9	11	11	4
2000	139	2	-	-	-	-	1	3	1	5	9	11	6	12
1999	157	-	-	-	1	1	1	1	3	6	7	6	9	11

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
Non-Hispanic White										
Both sexes										
2000	4,071	5,780	8,549	12,119	14,845	15,313	10,969	4,095	856	2
1999	4,200	5,928	9,002	12,339	14,979	15,344	10,615	4,102	854	3
Male										
2002	2,504	3,130	4,553	6,107	6,918	6,055	3,137	893	130	2
2001	2,448	3,183	4,595	6,010	6,625	5,579	2,926	888	131	1
2000	2,524	3,362	4,767	6,050	6,404	5,531	3,017	850	139	2
1999	2,600	3,603	4,984	6,137	6,559	5,402	2,890	877	113	2
Female										
2002	1,547	2,289	3,655	6,105	8,889	10,177	8,285	3,285	718	-
2001	1,526	2,310	3,751	5,927	8,492	10,057	7,916	3,290	717	-
2000	1,547	2,418	3,782	6,069	8,441	9,782	7,952	3,245	717	-
1999	1,600	2,325	4,018	6,202	8,420	9,942	7,725	3,225	741	1
Non-Hispanic Black										
Both sexes										
2002	1,077	1,191	1,337	1,520	1,429	1,155	824	376	115	1
2001	1,014	1,238	1,323	1,583	1,402	1,195	754	310	105	-
2000	1,048	1,240	1,419	1,584	1,331	1,181	729	339	106	1
1999	1,162	1,257	1,506	1,572	1,381	1,211	707	298	112	1
Male										
2002	585	623	673	674	565	366	240	74	18	-
2001	590	618	667	744	593	414	198	72	20	-
2000	583	667	688	716	529	407	186	86	21	-
1999	668	699	769	709	575	414	208	77	21	-
Female										
2002	492	568	664	846	864	789	584	302	97	1
2001	424	620	656	839	809	781	556	238	85	-
2000	465	573	731	868	802	774	543	253	85	1
1999	494	558	737	863	806	797	499	221	91	1
Origin not stated[2]										
Both sexes										
2002	14	17	26	30	43	28	21	12	1	3
2001	15	17	29	32	34	42	29	11	1	2
2000	18	13	29	29	23	41	23	13	2	-
1999	18	21	36	35	38	39	19	4	3	1
Male										
2002	8	11	15	21	15	7	3	4	-	1
2001	13	12	18	19	10	14	12	1	-	1
2000	12	10	18	14	10	16	4	5	-	-
1999	12	11	23	22	19	15	6	2	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)														
Origin not stated[2]														
Female														
2002	134	1	-	-	-	-	-	2	1	2	4	3	4	7
2001	136	1	1	-	-	-	3	1	1	-	3	5	3	6
2000	136	1	-	1	-	-	-	1	4	1	5	7	6	8
1999	124	1	1	-	-	-	-	2	1	2	4	3	1	6
Essential (primary) hypertension and hypertensive renal disease (I10,I12)														
All origins														
Both sexes														
2002	20,261	5	1	1	-	4	17	36	49	126	228	382	541	681
2001	19,250	-	1	-	1	5	18	32	72	101	230	385	546	630
2000	18,073	1	1	2	2	6	11	21	49	105	243	352	526	597
1999	16,968	3	3	-	1	3	10	24	54	105	208	329	470	538
Male														
2002	7,647	2	1	1	-	-	11	21	26	75	144	238	322	398
2001	7,254	-	-	-	-	2	14	20	39	59	147	239	330	354
2000	6,895	1	-	1	2	-	4	12	27	61	155	224	331	352
1999	6,488	2	1	-	1	2	6	13	35	69	126	196	286	316
Female														
2002	12,614	3	-	-	-	4	6	15	23	51	84	144	219	283
2001	11,996	-	1	-	1	3	4	12	33	42	83	146	216	276
2000	11,178	-	1	1	-	6	7	9	22	44	88	128	195	245
1999	10,480	1	2	-	-	1	4	11	19	36	82	133	184	222
Hispanic														
Both sexes														
2002	942	2	-	-	-	2	3	4	5	12	18	17	22	50
2001	889	-	-	-	-	1	2	1	9	5	14	28	32	31
2000	821	1	1	-	-	1	3	2	5	11	16	18	37	34
1999	753	1	1	-	-	1	2	4	4	3	16	29	35	29
Male														
2002	385	1	-	-	-	-	1	3	2	8	10	13	9	30
2001	365	-	-	-	-	1	1	1	7	4	12	21	18	13
2000	386	1	-	-	-	-	2	1	4	8	12	16	28	20
1999	340	-	-	-	-	1	1	2	3	3	11	16	19	11
Female														
2002	557	1	-	-	-	2	2	1	3	4	8	4	13	20
2001	524	-	-	-	-	-	1	-	2	1	2	7	14	18
2000	435	-	1	-	-	1	1	1	1	3	4	2	9	14
1999	413	1	1	-	-	-	1	2	1	-	5	13	16	18

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
Origin not stated[2]										
Female										
2002	6	6	11	9	28	21	18	8	1	2
2001	2	5	11	13	24	28	17	10	1	1
2000	6	3	11	15	13	25	19	8	2	-
1999	6	10	13	13	19	24	13	2	3	-
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
All origins										
Both sexes										
2002	845	1,126	1,796	2,592	3,546	3,801	2,915	1,276	293	-
2001	832	1,167	1,665	2,638	3,355	3,606	2,573	1,136	257	-
2000	843	1,094	1,690	2,507	3,116	3,273	2,375	1,017	242	-
1999	780	1,146	1,653	2,400	2,924	2,949	2,208	942	218	-
Male										
2002	452	563	856	1,103	1,302	1,182	707	215	28	-
2001	430	603	776	1,100	1,182	1,129	610	192	28	-
2000	466	504	804	1,049	1,155	991	569	155	32	-
1999	416	589	783	996	1,071	878	521	161	20	-
Female										
2002	393	563	940	1,489	2,244	2,619	2,208	1,061	265	-
2001	402	564	889	1,538	2,173	2,477	1,963	944	229	-
2000	377	590	886	1,458	1,961	2,282	1,806	862	210	-
1999	364	557	870	1,404	1,853	2,071	1,687	781	198	-
Hispanic										
Both sexes										
2002	73	66	117	130	153	116	96	43	13	-
2001	52	72	115	135	136	129	82	40	5	-
2000	75	65	97	118	114	122	55	38	8	-
1999	53	71	87	110	122	80	80	19	6	-
Male										
2002	36	26	54	61	52	33	32	9	5	-
2001	21	30	54	58	56	37	24	6	1	-
2000	45	26	46	57	50	38	21	7	4	-
1999	29	42	42	56	43	26	28	7	-	-
Female										
2002	37	40	63	69	101	83	64	34	8	-
2001	31	42	61	77	80	92	58	34	4	-
2000	30	39	51	61	64	84	34	31	4	-
1999	24	29	45	54	79	54	52	12	6	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Essential (primary) hypertension and hypertensive renal disease (I10,I12)														
Non-Hispanic[1]														
Both sexes														
2002	19,254	3	1	1	-	2	14	31	44	114	208	364	517	628
2001	18,306	-	1	-	1	4	16	31	63	96	214	354	514	594
2000	17,182	-	-	2	2	5	8	19	44	93	225	328	487	557
1999	16,158	2	2	-	1	2	8	20	49	101	191	299	435	506
Male														
2002	7,232	1	1	1	-	-	10	18	24	67	133	225	311	365
2001	6,856	-	-	-	-	1	13	19	32	55	133	216	312	337
2000	6,475	-	-	1	2	-	2	11	23	52	142	203	301	328
1999	6,123	2	1	-	1	1	5	11	31	65	114	180	267	303
Female														
2002	12,022	2	-	-	-	2	4	13	20	47	75	139	206	263
2001	11,450	-	1	-	1	3	3	12	31	41	81	138	202	257
2000	10,707	-	-	1	-	5	6	8	21	41	83	125	186	229
1999	10,035	-	1	-	-	1	3	9	18	36	77	119	168	203
Non-Hispanic White														
Both sexes														
2002	14,617	2	1	1	-	1	8	11	15	33	93	168	281	339
2001	13,837	-	1	-	1	3	8	12	23	40	81	155	249	323
2000	12,917	-	-	1	2	1	2	10	15	30	102	149	232	307
1999	12,048	1	-	-	-	2	3	8	20	34	78	143	200	275
Male														
2002	5,325	1	1	1	-	-	7	7	9	17	65	110	180	210
2001	4,970	-	-	-	-	1	6	7	13	23	54	98	155	192
2000	4,721	-	-	1	2	-	1	7	10	22	67	100	155	190
1999	4,395	1	-	-	-	1	1	5	17	26	51	93	127	181
Female														
2002	9,292	1	-	-	-	1	1	4	6	16	28	58	101	129
2001	8,867	-	1	-	1	2	2	5	10	17	27	57	94	131
2000	8,196	-	-	-	-	1	1	3	5	8	35	49	77	117
1999	7,653	-	-	-	-	1	2	3	3	8	27	50	73	94
Non-Hispanic Black														
Both sexes														
2002	4,165	1	-	-	-	1	6	17	27	79	110	188	225	272
2001	4,012	-	-	-	-	1	8	18	37	53	124	188	251	253
2000	3,862	-	-	1	-	4	6	7	28	61	115	168	237	240
1999	3,708	1	1	-	1	-	5	11	26	65	110	144	226	221
Male														
2002	1,701	-	-	-	-	-	3	10	14	48	64	109	125	145
2001	1,698	-	-	-	-	-	7	12	17	30	72	113	150	136

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
Non-Hispanic[1]										
Both sexes										
2002	772	1,053	1,672	2,457	3,383	3,673	2,811	1,228	278	-
2001	776	1,091	1,545	2,492	3,214	3,467	2,485	1,096	252	-
2000	763	1,028	1,588	2,377	2,996	3,140	2,309	978	233	-
1999	726	1,071	1,562	2,287	2,786	2,857	2,121	920	212	-
Male										
2002	416	530	798	1,039	1,246	1,144	674	206	23	-
2001	406	570	718	1,037	1,123	1,088	583	186	27	-
2000	418	477	755	986	1,102	949	547	148	28	-
1999	386	545	740	940	1,018	850	490	153	20	-
Female										
2002	356	523	874	1,418	2,137	2,529	2,137	1,022	255	-
2001	370	521	827	1,455	2,091	2,379	1,902	910	225	-
2000	345	551	833	1,391	1,894	2,191	1,762	830	205	-
1999	340	526	822	1,347	1,768	2,007	1,631	767	192	-
Non-Hispanic White										
Both sexes										
2002	466	675	1,163	1,835	2,735	3,077	2,404	1,075	234	-
2001	453	679	1,068	1,908	2,591	2,922	2,156	958	206	-
2000	429	644	1,100	1,798	2,454	2,644	1,992	818	187	-
1999	411	683	1,105	1,710	2,230	2,380	1,818	775	172	-
Male										
2002	267	350	570	787	997	977	576	176	17	-
2001	247	364	509	804	897	917	508	155	20	-
2000	244	315	537	763	908	801	462	116	20	-
1999	229	361	536	714	812	694	413	117	16	-
Female										
2002	199	325	593	1,048	1,738	2,100	1,828	899	217	-
2001	206	315	559	1,104	1,694	2,005	1,648	803	186	-
2000	185	329	563	1,035	1,546	1,843	1,530	702	167	-
1999	182	322	569	996	1,418	1,686	1,405	658	156	-
Non-Hispanic Black										
Both sexes										
2002	278	339	456	559	574	512	351	130	40	-
2001	300	372	417	512	544	478	287	126	43	-
2000	315	362	439	518	462	435	281	142	41	-
1999	285	342	422	528	478	416	266	125	35	-
Male										
2002	129	159	200	228	214	138	83	26	6	-
2001	148	190	184	202	197	146	60	27	7	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Essential (primary) hypertension and hypertensive renal disease (I10,I12)														
Non-Hispanic Black														
Male														
2000	1,576	-	-	-	-	-	1	3	13	29	69	96	133	135
1999	1,550	1	1	-	1	-	4	5	12	37	60	83	135	114
Female														
2002	2,464	1	-	-	-	1	3	7	13	31	46	79	100	127
2001	2,314	-	-	-	-	1	1	6	20	23	52	75	101	117
2000	2,286	-	-	1	-	4	5	4	15	32	46	72	104	105
1999	2,158	-	-	-	-	-	1	6	14	28	50	61	91	107
Origin not stated[2]														
Both sexes														
2002	65	-	-	-	-	-	-	1	-	-	2	1	2	3
2001	55	-	-	-	-	-	-	-	-	-	2	3	-	5
2000	70	-	-	-	-	-	-	-	-	1	2	6	2	6
1999	57	-	-	-	-	-	-	-	1	1	1	1	-	3
Male														
2002	30	-	-	-	-	-	-	-	-	-	1	-	2	3
2001	33	-	-	-	-	-	-	-	-	-	2	2	-	4
2000	34	-	-	-	-	-	-	-	-	1	1	5	2	4
1999	25	-	-	-	-	-	-	-	1	1	1	-	-	2
Female														
2002	35	-	-	-	-	-	-	1	-	-	1	1	-	-
2001	22	-	-	-	-	-	-	-	-	-	-	1	-	1
2000	36	-	-	-	-	-	-	-	-	-	1	1	-	2
1999	32	-	-	-	-	-	-	-	-	-	-	1	-	1
Cerebrovascular diseases (I60-I69)														
All origins														
Both sexes														
2002	162,672	117	53	33	58	53	118	183	384	797	1,628	2,615	3,440	4,293
2001	163,538	108	54	38	42	68	128	199	402	840	1,651	2,507	3,403	4,093
2000	167,661	124	47	25	51	67	132	209	393	880	1,719	2,625	3,386	4,145
1999	167,366	103	46	26	39	67	115	202	378	916	1,658	2,455	3,108	4,152
Male														
2002	62,622	67	30	16	31	34	69	100	184	389	789	1,402	1,876	2,373
2001	63,177	63	25	21	21	36	67	105	206	435	832	1,354	1,851	2,251
2000	64,769	74	19	12	21	38	68	111	197	430	867	1,401	1,834	2,294
1999	64,485	65	22	17	17	36	59	102	197	449	859	1,282	1,714	2,256
Female														
2002	100,050	50	23	17	27	19	49	83	200	408	839	1,213	1,564	1,920

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension and hypertensive renal disease (I10,I12)										
Non-Hispanic Black										
Male										
2000	166	153	191	195	163	123	74	27	5	-
1999	142	165	183	204	177	131	61	30	4	-
Female										
2002	149	180	256	331	360	374	268	104	34	-
2001	152	182	233	310	347	332	227	99	36	-
2000	149	209	248	323	299	312	207	115	36	-
1999	143	177	239	324	301	285	205	95	31	-
Origin not stated[2]										
Both sexes										
2002	-	7	7	5	10	12	8	5	2	-
2001	4	4	5	11	5	10	6	-	-	-
2000	5	1	5	12	6	11	11	1	1	-
1999	1	4	4	3	16	12	7	3	-	-
Male										
2002	-	7	4	3	4	5	1	-	-	-
2001	3	3	4	5	3	4	3	-	-	-
2000	3	1	3	6	3	4	1	-	-	-
1999	1	2	1	-	10	2	3	1	-	-
Female										
2002	-	-	3	2	6	7	7	5	2	-
2001	1	1	1	6	2	6	3	-	-	-
2000	2	-	2	6	3	7	10	1	1	-
1999	-	2	3	3	6	10	4	2	-	-
Cerebrovascular diseases (I60-I69)										
All origins										
Both sexes										
2002	5,604	8,026	13,966	22,911	31,978	33,197	23,110	8,521	1,584	3
2001	5,515	8,170	14,428	23,772	32,050	33,465	22,650	8,398	1,553	4
2000	5,811	8,671	14,978	24,678	32,342	34,030	23,205	8,530	1,611	2
1999	5,500	8,694	15,398	25,007	32,420	33,841	23,149	8,571	1,519	2
Male										
2002	3,083	4,305	6,902	10,283	12,372	10,884	5,713	1,510	209	1
2001	3,026	4,345	7,304	10,671	12,630	10,731	5,482	1,506	211	4
2000	3,202	4,513	7,529	11,192	12,754	10,864	5,588	1,564	196	1
1999	3,049	4,522	7,698	11,349	12,739	10,818	5,569	1,467	198	1
Female										
2002	2,521	3,721	7,064	12,628	19,606	22,313	17,397	7,011	1,375	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cerebrovascular diseases (I60-I69)														
All origins														
Female														
2001	100,361	45	29	17	21	32	61	94	196	405	819	1,153	1,552	1,842
2000	102,892	50	28	13	30	29	64	98	196	450	852	1,224	1,552	1,851
1999	102,881	38	24	9	22	31	56	100	181	467	799	1,173	1,394	1,896
Hispanic														
Both sexes														
2002	6,451	20	12	5	15	10	19	40	53	94	162	269	291	321
2001	6,416	17	10	7	9	13	27	34	65	102	173	239	300	301
2000	6,187	16	6	4	4	5	26	37	71	95	172	245	232	271
1999	5,907	16	7	4	5	10	22	46	56	89	152	229	235	286
Male														
2002	3,003	9	5	1	6	6	11	29	32	57	97	169	171	196
2001	2,982	9	4	5	5	8	18	20	40	58	93	141	174	171
2000	2,865	8	-	1	2	2	15	22	43	53	92	144	136	172
1999	2,809	8	1	3	3	7	14	25	37	49	97	136	149	157
Female														
2002	3,448	11	7	4	9	4	8	11	21	37	65	100	120	125
2001	3,434	8	6	2	4	5	9	14	25	44	80	98	126	130
2000	3,322	8	6	3	2	3	11	15	28	42	80	101	96	99
1999	3,098	8	6	1	2	3	8	21	19	40	55	93	86	129
Non-Hispanic[1]														
Both sexes														
2002	155,852	96	41	28	43	43	99	143	331	699	1,462	2,337	3,140	3,961
2001	156,741	91	43	30	33	55	101	163	335	735	1,476	2,258	3,092	3,779
2000	161,057	108	41	21	46	62	105	172	321	783	1,540	2,366	3,143	3,861
1999	161,057	87	39	22	34	57	93	156	321	826	1,497	2,213	2,858	3,848
Male														
2002	59,470	57	25	15	25	28	58	71	152	330	689	1,227	1,699	2,172
2001	60,030	54	21	16	16	28	49	83	164	374	737	1,209	1,673	2,071
2000	61,715	66	19	11	19	36	53	89	154	376	769	1,248	1,688	2,111
1999	61,499	57	21	14	14	29	45	77	160	400	756	1,139	1,558	2,088
Female														
2002	96,382	39	16	13	18	15	41	72	179	369	773	1,110	1,441	1,789
2001	96,711	37	22	14	17	27	52	80	171	361	739	1,049	1,419	1,708
2000	99,342	42	22	10	27	26	52	83	167	407	771	1,118	1,455	1,750
1999	99,558	30	18	8	20	28	48	79	161	426	741	1,074	1,300	1,760
Non-Hispanic White														
Both sexes														
2002	133,118	51	29	20	24	28	73	85	200	412	876	1,407	1,913	2,733
2001	133,879	46	28	21	21	40	58	101	219	452	881	1,363	1,837	2,606

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)										
All origins										
Female										
2001	2,489	3,825	7,124	13,101	19,420	22,734	17,168	6,892	1,342	-
2000	2,609	4,158	7,449	13,486	19,588	23,166	17,617	6,966	1,415	1
1999	2,451	4,172	7,700	13,658	19,681	23,023	17,580	7,104	1,321	1
Hispanic										
Both sexes										
2002	387	462	753	901	1,017	838	535	202	45	-
2001	353	471	735	959	933	868	532	221	47	-
2000	410	498	721	859	883	852	539	209	32	-
1999	331	524	687	823	849	828	505	170	33	-
Male										
2002	225	256	396	421	404	282	177	42	11	-
2001	207	269	368	456	408	300	161	55	12	-
2000	220	267	378	421	378	282	159	63	7	-
1999	197	273	368	423	335	284	183	50	10	-
Female										
2002	162	206	357	480	613	556	358	160	34	-
2001	146	202	367	503	525	568	371	166	35	-
2000	190	231	343	438	505	570	380	146	25	-
1999	134	251	319	400	514	544	322	120	23	-
Non-Hispanic[1]										
Both sexes										
2002	5,199	7,546	13,185	21,958	30,892	32,291	22,516	8,302	1,537	3
2001	5,137	7,683	13,663	22,752	31,058	32,525	22,071	8,154	1,503	4
2000	5,377	8,160	14,213	23,754	31,375	33,112	22,615	8,305	1,576	1
1999	5,143	8,145	14,669	24,125	31,518	32,951	22,585	8,386	1,482	2
Male										
2002	2,846	4,039	6,493	9,834	11,939	10,583	5,522	1,467	198	1
2001	2,801	4,066	6,923	10,187	12,190	10,412	5,309	1,446	197	4
2000	2,968	4,239	7,125	10,737	12,347	10,558	5,418	1,496	188	-
1999	2,834	4,234	7,308	10,892	12,383	10,518	5,366	1,417	188	1
Female										
2002	2,353	3,507	6,692	12,124	18,953	21,708	16,994	6,835	1,339	2
2001	2,336	3,617	6,740	12,565	18,868	22,113	16,762	6,708	1,306	-
2000	2,409	3,921	7,088	13,017	19,028	22,554	17,197	6,809	1,388	1
1999	2,309	3,911	7,361	13,233	19,135	22,433	17,219	6,969	1,294	1
Non-Hispanic White										
Both sexes										
2002	3,699	5,654	10,701	18,821	27,506	29,412	20,578	7,553	1,341	2
2001	3,602	5,715	11,123	19,553	27,751	29,529	20,201	7,436	1,295	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cerebrovascular diseases (I60-I69)														
Non-Hispanic White														
Both sexes														
2000	138,210	59	25	17	29	46	72	123	190	464	914	1,380	2,053	2,657
1999	138,716	55	27	19	21	37	68	100	201	514	892	1,302	1,841	2,649
Male														
2002	49,908	32	18	10	13	17	45	47	89	195	422	765	1,032	1,490
2001	50,382	27	14	12	12	20	35	49	112	223	461	739	1,006	1,421
2000	52,000	38	14	11	12	31	36	67	94	229	465	752	1,104	1,431
1999	51,968	36	15	13	8	21	30	48	103	252	460	664	998	1,452
Female														
2002	83,210	19	11	10	11	11	28	38	111	217	454	642	881	1,243
2001	83,497	19	14	9	9	20	23	52	107	229	420	624	831	1,185
2000	86,210	21	11	6	17	15	36	56	96	235	449	628	949	1,226
1999	86,748	19	12	6	13	16	38	52	98	262	432	638	843	1,197
Non-Hispanic Black														
Both sexes														
2002	18,691	36	10	7	16	13	25	50	116	237	504	807	1,067	1,048
2001	18,844	42	12	6	8	13	38	57	96	233	525	790	1,096	992
2000	19,053	46	13	2	13	13	29	41	117	281	548	868	951	1,032
1999	18,731	27	10	3	12	18	22	49	100	267	541	780	882	1,017
Male														
2002	7,754	19	6	4	11	10	12	20	52	102	229	398	588	592
2001	7,828	26	5	2	3	8	12	33	42	118	237	412	586	554
2000	7,945	27	3	-	5	4	15	18	52	127	267	440	515	587
1999	7,823	17	4	1	6	7	12	24	48	123	264	409	491	541
Female														
2002	10,937	17	4	3	5	3	13	30	64	135	275	409	479	456
2001	11,016	16	7	4	5	5	26	24	54	115	288	378	510	438
2000	11,108	19	10	2	8	9	14	23	65	154	281	428	436	445
1999	10,908	10	6	2	6	11	10	25	52	144	277	371	391	476
Origin not stated[2]														
Both sexes														
2002	369	1	-	-	-	-	-	-	-	4	4	9	9	11
2001	381	-	1	1	-	-	-	2	2	3	2	10	11	13
2000	417	-	-	-	1	-	1	-	1	2	7	14	11	13
1999	402	-	-	-	-	-	-	-	1	1	9	13	15	18
Male														
2002	149	1	-	-	-	-	-	-	-	2	3	6	6	5
2001	165	-	-	-	-	-	-	2	2	3	2	4	4	9
2000	189	-	-	-	-	-	-	-	-	1	6	9	10	11
1999	177	-	-	-	-	-	-	-	-	-	6	7	7	11

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)										
Non-Hispanic White										
Both sexes										
2000	3,898	6,277	11,712	20,501	28,055	30,109	20,776	7,491	1,362	-
1999	3,662	6,342	11,990	20,866	28,312	29,995	20,832	7,676	1,315	-
Male										
2002	2,021	3,032	5,336	8,484	10,697	9,680	5,016	1,313	153	1
2001	1,979	3,066	5,687	8,818	10,944	9,439	4,851	1,294	172	1
2000	2,174	3,274	5,902	9,308	11,071	9,594	4,928	1,301	164	-
1999	2,027	3,320	6,017	9,436	11,182	9,548	4,884	1,286	168	-
Female										
2002	1,678	2,622	5,365	10,337	16,809	19,732	15,562	6,240	1,188	1
2001	1,623	2,649	5,436	10,735	16,807	20,090	15,350	6,142	1,123	-
2000	1,724	3,003	5,810	11,193	16,984	20,515	15,848	6,190	1,198	-
1999	1,635	3,022	5,973	11,430	17,130	20,447	15,948	6,390	1,147	-
Non-Hispanic Black										
Both sexes										
2002	1,255	1,584	2,016	2,576	2,701	2,298	1,546	610	168	1
2001	1,297	1,640	2,044	2,627	2,630	2,391	1,530	604	170	3
2000	1,233	1,573	2,087	2,644	2,700	2,472	1,507	685	197	1
1999	1,252	1,529	2,263	2,715	2,599	2,435	1,452	614	143	1
Male										
2002	696	848	978	1,084	947	671	349	100	38	-
2001	698	854	990	1,119	949	726	319	114	18	3
2000	653	812	1,005	1,138	1,017	727	367	145	21	-
1999	694	782	1,075	1,169	931	735	363	110	17	-
Female										
2002	559	736	1,038	1,492	1,754	1,627	1,197	510	130	1
2001	599	786	1,054	1,508	1,681	1,665	1,211	490	152	-
2000	580	761	1,082	1,506	1,683	1,745	1,140	540	176	1
1999	558	747	1,188	1,546	1,668	1,700	1,089	504	126	1
Origin not stated[2]										
Both sexes										
2002	18	18	28	52	69	68	59	17	2	-
2001	25	16	30	61	59	72	47	23	3	-
2000	24	13	44	65	84	66	51	16	3	1
1999	26	25	42	59	53	62	59	15	4	-
Male										
2002	12	10	13	28	29	19	14	1	-	-
2001	18	10	13	28	32	19	12	5	2	-
2000	14	7	26	34	29	24	11	5	1	1
1999	18	15	22	34	21	16	20	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cerebrovascular diseases (I60-I69)														
Origin not stated[2]														
Female														
2002	220	-	-	-	-	-	-	-	-	2	1	3	3	6
2001	216	-	1	1	-	-	-	-	-	-	-	6	7	4
2000	228	-	-	-	1	-	1	-	1	1	1	5	1	2
1999	225	-	-	-	-	-	-	-	1	1	3	6	8	7
Atherosclerosis (I70)														
All origins														
Both sexes														
2002	13,821	4	-	-	-	-	3	2	7	16	34	61	121	207
2001	14,086	1	1	1	-	-	1	1	2	14	26	65	121	198
2000	14,393	2	-	-	-	-	3	3	5	16	32	71	105	170
1999	14,979	2	-	-	-	1	1	1	7	16	29	56	129	172
Male														
2002	5,171	1	-	-	-	-	2	2	3	9	21	44	75	127
2001	5,143	1	1	1	-	-	1	-	2	8	18	43	75	118
2000	5,357	-	-	-	-	-	2	1	5	10	25	48	71	108
1999	5,443	1	-	-	-	-	-	1	4	11	21	37	84	114
Female														
2002	8,650	3	-	-	-	-	1	-	4	7	13	17	46	80
2001	8,943	-	-	-	-	-	-	1	-	6	8	22	46	80
2000	9,036	2	-	-	-	-	1	2	-	6	7	23	34	62
1999	9,536	1	-	-	-	1	1	-	3	5	8	19	45	58
Hispanic														
Both sexes														
2002	416	1	-	-	-	-	2	-	-	1	1	3	6	9
2001	416	-	-	-	-	-	-	-	-	-	-	3	11	12
2000	414	-	-	-	-	-	-	-	-	2	-	4	5	6
1999	410	-	-	-	-	-	-	-	-	1	2	5	9	11
Male														
2002	175	1	-	-	-	-	2	-	-	-	1	3	3	7
2001	172	-	-	-	-	-	-	-	-	-	-	1	8	8
2000	173	-	-	-	-	-	-	-	-	2	-	3	4	2
1999	185	-	-	-	-	-	-	-	-	-	1	2	5	7
Female														
2002	241	-	-	-	-	-	-	-	-	1	-	-	3	2
2001	244	-	-	-	-	-	-	-	-	-	-	2	3	4
2000	241	-	-	-	-	-	-	-	-	-	-	1	1	4
1999	225	-	-	-	-	-	-	-	-	1	1	3	4	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)										
Origin not stated[2]										
Female										
2002	6	8	15	24	40	49	45	16	2	-
2001	7	6	17	33	27	53	35	18	1	-
2000	10	6	18	31	55	42	40	11	2	-
1999	8	10	20	25	32	46	39	15	4	-
Atherosclerosis (I70)										
All origins										
Both sexes										
2002	280	494	806	1,521	2,541	3,090	2,802	1,416	415	1
2001	275	471	864	1,624	2,532	3,214	2,786	1,471	417	1
2000	281	495	928	1,685	2,592	3,188	2,870	1,518	429	-
1999	314	485	991	1,770	2,615	3,362	2,987	1,581	460	-
Male										
2002	178	282	446	748	1,055	1,088	755	281	53	1
2001	168	273	470	765	1,026	1,123	710	285	54	1
2000	186	290	510	838	1,097	1,090	738	283	55	-
1999	193	277	518	845	1,082	1,100	773	316	66	-
Female										
2002	102	212	360	773	1,486	2,002	2,047	1,135	362	-
2001	107	198	394	859	1,506	2,091	2,076	1,186	363	-
2000	95	205	418	847	1,495	2,098	2,132	1,235	374	-
1999	121	208	473	925	1,533	2,262	2,214	1,265	394	-
Hispanic										
Both sexes										
2002	11	24	26	58	78	78	74	30	14	-
2001	7	14	32	60	68	99	59	39	12	-
2000	10	24	39	53	71	83	77	32	8	-
1999	14	15	39	60	70	79	61	35	9	-
Male										
2002	5	12	15	29	36	29	23	6	3	-
2001	3	8	16	31	30	39	13	11	4	-
2000	6	13	26	34	31	27	17	7	1	-
1999	9	9	22	30	30	37	18	11	4	-
Female										
2002	6	12	11	29	42	49	51	24	11	-
2001	4	6	16	29	38	60	46	28	8	-
2000	4	11	13	19	40	56	60	25	7	-
1999	5	6	17	30	40	42	43	24	5	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerosis (I70)														
Non-Hispanic[1]														
Both sexes														
2002	13,369	3	-	-	-	-	1	2	7	15	33	58	115	198
2001	13,626	1	1	1	-	-	1	1	2	14	26	62	110	185
2000	13,945	2	-	-	-	-	3	3	5	14	32	67	100	164
1999	14,519	2	-	-	-	1	1	1	7	15	27	49	120	160
Male														
2002	4,984	-	-	-	-	-	-	2	3	9	20	41	72	120
2001	4,952	1	1	1	-	-	1	-	2	8	18	42	67	109
2000	5,166	-	-	-	-	-	2	1	5	8	25	45	67	106
1999	5,236	1	-	-	-	-	-	1	4	11	20	33	79	106
Female														
2002	8,385	3	-	-	-	-	1	-	4	6	13	17	43	78
2001	8,674	-	-	-	-	-	-	1	-	6	8	20	43	76
2000	8,779	2	-	-	-	-	1	2	-	6	7	22	33	58
1999	9,283	1	-	-	-	1	1	-	3	4	7	16	41	54
Non-Hispanic White														
Both sexes														
2002	12,191	2	-	-	-	-	1	2	4	13	25	42	79	161
2001	12,517	1	1	1	-	-	-	-	2	11	22	51	86	153
2000	12,740	2	-	-	-	-	3	1	4	10	24	46	68	134
1999	13,286	1	-	-	-	1	1	1	5	13	17	41	94	129
Male														
2002	4,508	-	-	-	-	-	-	2	2	7	16	32	47	99
2001	4,511	1	1	1	-	-	-	-	2	5	16	34	51	92
2000	4,688	-	-	-	-	-	2	1	4	7	19	33	46	91
1999	4,727	1	-	-	-	-	-	1	3	10	13	28	65	83
Female														
2002	7,683	2	-	-	-	-	1	-	2	6	9	10	32	62
2001	8,006	-	-	-	-	-	-	-	-	6	6	17	35	61
2000	8,052	2	-	-	-	-	1	-	-	3	5	13	22	43
1999	8,559	-	-	-	-	1	1	-	2	3	4	13	29	46
Non-Hispanic Black														
Both sexes														
2002	1,009	1	-	-	-	-	-	-	3	2	7	15	31	29
2001	957	-	-	-	-	-	1	1	-	2	4	11	20	28
2000	1,049	-	-	-	-	-	-	2	1	4	8	19	28	28
1999	1,061	1	-	-	-	-	-	-	2	2	7	6	23	26
Male														
2002	409	-	-	-	-	-	-	-	1	2	4	8	22	17
2001	365	-	-	-	-	-	1	-	-	2	2	8	13	14
2000	412	-	-	-	-	-	-	-	1	1	6	11	17	14

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)										
Non-Hispanic[1]										
Both sexes										
2002	268	470	778	1,455	2,456	3,001	2,722	1,386	400	1
2001	267	455	830	1,556	2,455	3,104	2,720	1,429	405	1
2000	270	469	888	1,627	2,518	3,095	2,787	1,481	420	-
1999	297	465	951	1,706	2,536	3,273	2,916	1,544	448	-
Male										
2002	172	270	430	716	1,017	1,056	731	275	49	1
2001	165	264	454	729	992	1,080	695	272	50	1
2000	179	275	484	801	1,064	1,058	717	276	53	-
1999	181	266	495	814	1,047	1,059	753	304	62	-
Female										
2002	96	200	348	739	1,439	1,945	1,991	1,111	351	-
2001	102	191	376	827	1,463	2,024	2,025	1,157	355	-
2000	91	194	404	826	1,454	2,037	2,070	1,205	367	-
1999	116	199	456	892	1,489	2,214	2,163	1,240	386	-
Non-Hispanic White										
Both sexes										
2002	215	411	671	1,282	2,236	2,828	2,560	1,301	357	1
2001	208	391	728	1,408	2,267	2,926	2,561	1,333	366	1
2000	215	405	776	1,479	2,303	2,892	2,610	1,394	374	-
1999	245	394	847	1,534	2,346	3,050	2,725	1,440	402	-
Male										
2002	138	239	374	631	930	999	688	260	43	1
2001	133	228	403	656	914	1,025	652	256	40	1
2000	147	237	436	740	976	983	666	254	46	-
1999	152	227	440	724	966	981	702	279	52	-
Female										
2002	77	172	297	651	1,306	1,829	1,872	1,041	314	-
2001	75	163	325	752	1,353	1,901	1,909	1,077	326	-
2000	68	168	340	739	1,327	1,909	1,944	1,140	328	-
1999	93	167	407	810	1,380	2,069	2,023	1,161	350	-
Non-Hispanic Black										
Both sexes										
2002	48	49	85	154	195	150	135	68	37	-
2001	56	57	88	131	154	149	137	86	32	-
2000	51	51	96	130	190	172	153	76	40	-
1999	50	62	88	152	160	190	162	91	39	-
Male										
2002	32	25	45	76	77	48	35	13	4	-
2001	29	30	44	65	59	44	35	12	7	-
2000	29	33	41	52	79	63	41	19	5	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Atherosclerosis (I70)														
Non-Hispanic Black														
Male														
1999	430	-	-	-	-	-	-	-	1	1	5	4	11	19
Female														
2002	600	1	-	-	-	-	-	-	2	-	3	7	9	12
2001	592	-	-	-	-	-	-	1	-	-	2	3	7	14
2000	637	-	-	-	-	-	-	2	-	3	2	8	11	14
1999	631	1	-	-	-	-	-	-	1	1	2	2	12	7
Origin not stated[2]														
Both sexes														
2002	36	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	44	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	34	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	50	-	-	-	-	-	-	-	-	-	-	2	-	1
Male														
2002	12	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	19	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	18	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	22	-	-	-	-	-	-	-	-	-	-	2	-	1
Female														
2002	24	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	25	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	16	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	28	-	-	-	-	-	-	-	-	-	-	-	-	-
Other diseases of circulatory system (I71-I78)														
All origins														
Both sexes														
2002	24,927	14	4	1	10	31	35	51	109	154	291	479	591	928
2001	25,318	12	4	2	9	23	38	67	103	194	304	421	603	850
2000	26,036	16	5	3	8	17	42	49	105	157	291	424	614	866
1999	25,809	12	2	2	9	14	35	51	90	178	300	418	553	877
Male														
2002	13,309	6	4	-	5	19	24	30	71	81	210	331	401	624
2001	13,396	8	2	-	3	17	23	43	72	132	203	291	397	566
2000	14,084	6	2	1	6	9	24	36	67	92	184	275	424	607
1999	13,794	9	-	1	6	10	22	35	67	117	179	289	368	597
Female														
2002	11,618	8	-	1	5	12	11	21	38	73	81	148	190	304
2001	11,922	4	2	2	6	6	15	24	31	62	101	130	206	284

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)										
Non-Hispanic Black										
Male										
1999	27	34	46	78	64	68	39	23	10	-
Female										
2002	16	24	40	78	118	102	100	55	33	-
2001	27	27	44	66	95	105	102	74	25	-
2000	22	18	55	78	111	109	112	57	35	-
1999	23	28	42	74	96	122	123	68	29	-
Origin not stated[2]										
Both sexes										
2002	1	-	2	8	7	11	6	-	1	-
2001	1	2	2	8	9	11	7	3	-	-
2000	1	2	1	5	3	10	6	5	1	-
1999	3	5	1	4	9	10	10	2	3	-
Male										
2002	1	-	1	3	2	3	1	-	1	-
2001	-	1	-	5	4	4	2	2	-	-
2000	1	2	-	3	2	5	4	-	1	-
1999	3	2	1	1	5	4	2	1	-	-
Female										
2002	-	-	1	5	5	8	5	-	-	-
2001	1	1	2	3	5	7	5	1	-	-
2000	-	-	1	2	1	5	2	5	-	-
1999	-	3	-	3	4	6	8	1	3	-
Other diseases of circulatory system (I71-I78)										
All origins										
Both sexes										
2002	1,358	2,137	3,173	4,292	4,453	3,768	2,151	743	154	-
2001	1,345	2,127	3,218	4,498	4,534	3,867	2,210	748	140	1
2000	1,370	2,331	3,595	4,662	4,560	3,819	2,218	757	126	1
1999	1,333	2,244	3,564	4,662	4,602	3,792	2,161	765	145	-
Male										
2002	902	1,408	1,899	2,406	2,284	1,675	715	191	23	-
2001	888	1,357	1,991	2,476	2,363	1,633	719	188	23	1
2000	923	1,524	2,247	2,696	2,323	1,678	734	204	21	1
1999	922	1,472	2,209	2,684	2,331	1,567	711	172	26	-
Female										
2002	456	729	1,274	1,886	2,169	2,093	1,436	552	131	-
2001	457	770	1,227	2,022	2,171	2,234	1,491	560	117	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of circulatory system (I71-I78)														
All origins														
Female														
2000	11,952	10	3	2	2	8	18	13	38	65	107	149	190	259
1999	12,015	3	2	1	3	4	13	16	23	61	121	129	185	280
Hispanic														
Both sexes														
2002	847	1	-	-	1	6	6	6	16	15	29	29	33	48
2001	754	2	1	1	1	6	1	7	12	11	26	31	42	39
2000	818	3	1	-	1	5	4	11	13	18	20	19	37	45
1999	729	2	-	-	2	4	5	6	6	13	20	15	23	39
Male														
2002	463	-	-	-	-	4	4	4	10	9	22	19	22	26
2001	424	1	1	-	-	4	-	4	10	6	20	20	26	22
2000	452	1	-	-	1	3	1	7	7	7	15	9	24	28
1999	416	1	-	-	-	2	5	3	3	9	13	11	14	25
Female														
2002	384	1	-	-	1	2	2	2	6	6	7	10	11	22
2001	330	1	-	1	1	2	1	3	2	5	6	11	16	17
2000	366	2	1	-	-	2	3	4	6	11	5	10	13	17
1999	313	1	-	-	2	2	-	3	3	4	7	4	9	14
Non-Hispanic[1]														
Both sexes														
2002	24,005	13	4	1	9	25	29	45	92	137	259	449	557	877
2001	24,489	10	3	1	8	17	37	59	90	183	277	389	557	808
2000	25,147	13	4	3	7	12	38	37	92	138	267	404	575	818
1999	25,023	10	2	2	7	10	30	45	84	165	280	403	529	834
Male														
2002	12,809	6	4	-	5	15	20	26	60	71	186	312	378	595
2001	12,926	7	1	-	3	13	23	38	61	126	182	270	368	541
2000	13,593	5	2	1	5	6	23	29	60	84	165	265	398	577
1999	13,351	8	-	1	6	8	17	32	64	108	166	278	354	571
Female														
2002	11,196	7	-	1	4	10	9	19	32	66	73	137	179	282
2001	11,563	3	2	1	5	4	14	21	29	57	95	119	189	267
2000	11,554	8	2	2	2	6	15	8	32	54	102	139	177	241
1999	11,672	2	2	1	1	2	13	13	20	57	114	125	175	263
Non-Hispanic White														
Both sexes														
2002	21,051	10	4	1	3	20	17	29	57	98	174	301	401	692
2001	21,439	8	2	1	7	12	28	44	64	126	177	277	405	641

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of circulatory system (I71-I78)										
All origins										
Female										
2000	447	807	1,348	1,966	2,237	2,141	1,484	553	105	-
1999	411	772	1,355	1,978	2,271	2,225	1,450	593	119	-
Hispanic										
Both sexes										
2002	69	73	102	130	116	85	53	21	8	-
2001	49	71	85	125	91	82	43	25	2	1
2000	54	88	129	129	86	71	56	25	3	-
1999	55	73	97	97	114	92	46	16	4	-
Male										
2002	40	44	60	70	68	29	23	5	4	-
2001	33	47	53	76	43	37	13	6	1	1
2000	36	51	71	81	45	33	24	6	2	-
1999	34	44	57	57	62	48	22	4	2	-
Female										
2002	29	29	42	60	48	56	30	16	4	-
2001	16	24	32	49	48	45	30	19	1	-
2000	18	37	58	48	41	38	32	19	1	-
1999	21	29	40	40	52	44	24	12	2	-
Non-Hispanic[1]										
Both sexes										
2002	1,282	2,056	3,064	4,145	4,330	3,673	2,094	718	146	-
2001	1,291	2,048	3,122	4,364	4,431	3,774	2,160	722	138	-
2000	1,311	2,238	3,457	4,521	4,464	3,736	2,157	731	123	1
1999	1,275	2,162	3,457	4,560	4,477	3,694	2,109	747	141	-
Male										
2002	860	1,359	1,835	2,328	2,212	1,641	691	186	19	-
2001	851	1,305	1,929	2,394	2,314	1,593	704	181	22	-
2000	884	1,471	2,169	2,607	2,276	1,641	708	197	19	1
1999	885	1,422	2,145	2,626	2,263	1,519	687	167	24	-
Female										
2002	422	697	1,229	1,817	2,118	2,032	1,403	532	127	-
2001	440	743	1,193	1,970	2,117	2,181	1,456	541	116	-
2000	427	767	1,288	1,914	2,188	2,095	1,449	534	104	-
1999	390	740	1,312	1,934	2,214	2,175	1,422	580	117	-
Non-Hispanic White										
Both sexes										
2002	1,066	1,767	2,718	3,738	3,946	3,361	1,880	655	113	-
2001	1,061	1,782	2,757	3,892	4,016	3,447	1,960	628	104	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of circulatory system (I71-I78)														
Non-Hispanic White														
Both sexes														
2000	22,113	11	3	2	4	9	26	24	64	99	170	284	415	648
1999	22,191	8	2	2	5	9	24	32	62	125	198	289	401	672
Male														
2002	11,337	5	4	-	2	12	11	18	36	53	127	220	275	472
2001	11,437	5	1	-	3	10	17	31	43	87	115	195	267	446
2000	12,138	3	2	1	4	4	18	17	46	59	108	199	295	470
1999	12,014	6	-	1	4	7	13	23	48	83	121	208	270	479
Female														
2002	9,714	5	-	1	1	8	6	11	21	45	47	81	126	220
2001	10,002	3	1	1	4	2	11	13	21	39	62	82	138	195
2000	9,975	8	1	1	-	5	8	7	18	40	62	85	120	178
1999	10,177	2	2	1	1	2	11	9	14	42	77	81	131	193
Non-Hispanic Black														
Both sexes														
2002	2,495	3	-	-	5	3	9	15	31	32	80	134	138	164
2001	2,606	2	1	-	1	5	7	14	25	50	91	98	129	150
2000	2,569	2	-	1	3	2	11	10	25	34	86	106	126	148
1999	2,416	2	-	-	-	1	6	12	18	35	77	104	108	146
Male														
2002	1,196	1	-	-	2	1	6	8	22	15	56	79	92	108
2001	1,214	2	-	-	-	3	4	7	17	33	60	63	86	82
2000	1,193	2	-	-	1	1	5	9	13	22	51	59	85	90
1999	1,102	2	-	-	-	1	4	8	15	21	40	65	70	82
Female														
2002	1,299	2	-	-	3	2	3	7	9	17	24	55	46	56
2001	1,392	-	1	-	1	2	3	7	8	17	31	35	43	68
2000	1,376	-	-	1	2	1	6	1	12	12	35	47	41	58
1999	1,314	-	-	-	-	-	2	4	3	14	37	39	38	64
Origin not stated[2]														
Both sexes														
2002	75	-	-	-	-	-	-	-	1	2	3	1	1	3
2001	75	-	-	-	-	-	-	1	1	-	1	1	4	3
2000	71	-	-	-	-	-	-	1	-	1	4	1	2	3
1999	57	-	-	-	-	-	-	-	-	-	-	-	1	4
Male														
2002	37	-	-	-	-	-	-	-	1	1	2	-	1	3
2001	46	-	-	-	-	-	-	1	1	-	1	1	3	3
2000	39	-	-	-	-	-	-	-	-	1	4	1	2	2
1999	27	-	-	-	-	-	-	-	-	-	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of circulatory system (I71-I78)										
Non-Hispanic White										
Both sexes										
2000	1,086	1,940	3,106	4,058	4,075	3,414	1,938	638	99	-
1999	1,101	1,891	3,109	4,129	4,081	3,373	1,903	660	115	-
Male										
2002	726	1,205	1,645	2,144	2,045	1,526	622	174	15	-
2001	712	1,163	1,734	2,177	2,126	1,482	641	163	19	-
2000	755	1,318	1,983	2,371	2,121	1,537	638	174	15	-
1999	782	1,281	1,972	2,423	2,091	1,405	635	146	16	-
Female										
2002	340	562	1,073	1,594	1,901	1,835	1,258	481	98	-
2001	349	619	1,023	1,715	1,890	1,965	1,319	465	85	-
2000	331	622	1,123	1,687	1,954	1,877	1,300	464	84	-
1999	319	610	1,137	1,706	1,990	1,968	1,268	514	99	-
Non-Hispanic Black										
Both sexes										
2002	186	240	281	336	317	252	180	59	30	-
2001	198	210	316	394	352	271	177	84	31	-
2000	198	246	294	386	324	276	187	82	21	1
1999	144	234	283	357	332	276	176	81	24	-
Male										
2002	115	122	146	140	129	84	57	9	4	-
2001	115	108	164	166	158	81	49	13	3	-
2000	108	126	153	190	124	82	49	18	4	1
1999	83	122	144	156	134	88	39	20	8	-
Female										
2002	71	118	135	196	188	168	123	50	26	-
2001	83	102	152	228	194	190	128	71	28	-
2000	90	120	141	196	200	194	138	64	17	-
1999	61	112	139	201	198	188	137	61	16	-
Origin not stated[2]										
Both sexes										
2002	7	8	7	17	7	10	4	4	-	-
2001	5	8	11	9	12	11	7	1	-	-
2000	5	5	9	12	10	12	5	1	-	-
1999	3	9	10	5	11	6	6	2	-	-
Male										
2002	2	5	4	8	4	5	1	-	-	-
2001	4	5	9	6	6	3	2	1	-	-
2000	3	2	7	8	2	4	2	1	-	-
1999	3	6	7	1	6	-	2	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of circulatory system (I71-I78)														
Origin not stated[2]														
Female														
2002	38	-	-	-	-	-	-	-	-	1	1	1	-	-
2001	29	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	32	-	-	-	-	-	-	1	-	-	-	-	-	1
1999	30	-	-	-	-	-	-	-	-	-	-	-	1	3
Aortic aneurysm and dissection (I71)														
All origins														
Both sexes														
2002	14,818	2	-	1	4	20	19	30	80	104	201	327	383	598
2001	15,234	-	2	-	4	15	28	46	71	134	198	287	374	532
2000	15,810	-	-	-	5	10	26	38	69	103	200	268	394	585
1999	15,807	1	1	-	3	8	23	33	60	134	195	283	355	570
Male														
2002	9,013	-	-	-	3	17	15	25	57	67	162	249	291	458
2001	9,114	-	2	-	3	12	19	34	56	105	159	219	287	398
2000	9,661	-	-	-	3	7	16	30	48	72	147	191	301	438
1999	9,615	1	-	-	2	6	14	25	53	98	134	216	270	430
Female														
2002	5,805	2	-	1	1	3	4	5	23	37	39	78	92	140
2001	6,120	-	-	-	1	3	9	12	15	29	39	68	87	134
2000	6,149	-	-	-	2	3	10	8	21	31	53	77	93	147
1999	6,192	-	1	-	1	2	9	8	7	36	61	67	85	140
Hispanic														
Both sexes														
2002	431	-	-	-	-	4	4	3	13	9	15	17	19	31
2001	403	-	1	-	-	3	1	4	10	8	16	18	23	20
2000	465	-	-	-	1	5	3	8	7	10	11	13	28	26
1999	384	-	-	-	1	2	2	3	3	10	13	10	17	24
Male														
2002	284	-	-	-	-	3	2	3	9	8	12	15	14	21
2001	279	-	1	-	-	2	-	4	10	6	14	14	17	12
2000	294	-	-	-	1	3	1	6	5	6	8	8	20	19
1999	256	-	-	-	-	1	2	2	2	7	10	9	13	17
Female														
2002	147	-	-	-	-	1	2	-	4	1	3	2	5	10
2001	124	-	-	-	-	1	1	-	-	2	2	4	6	8
2000	171	-	-	-	-	2	2	2	2	4	3	5	8	7
1999	128	-	-	-	1	1	-	1	1	3	3	1	4	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of circulatory system (I71-I78)										
Origin not stated[2]										
Female										
2002	5	3	3	9	3	5	3	4	-	-
2001	1	3	2	3	6	8	5	-	-	-
2000	2	3	2	4	8	8	3	-	-	-
1999	-	3	3	4	5	6	4	1	-	-
Aortic aneurysm and dissection (I71)										
All origins										
Both sexes										
2002	862	1,404	2,078	2,779	2,739	2,013	914	239	21	-
2001	862	1,430	2,087	2,899	2,852	2,122	1,022	242	26	1
2000	908	1,552	2,460	3,043	2,792	2,134	985	213	25	-
1999	867	1,578	2,423	3,095	2,841	2,118	935	257	27	-
Male										
2002	634	1,020	1,349	1,688	1,497	1,034	368	74	5	-
2001	633	981	1,401	1,722	1,591	1,000	406	78	7	1
2000	673	1,097	1,633	1,885	1,558	1,063	425	68	6	-
1999	658	1,114	1,598	1,905	1,596	1,008	398	83	6	-
Female										
2002	228	384	729	1,091	1,242	979	546	165	16	-
2001	229	449	686	1,177	1,261	1,122	616	164	19	-
2000	235	455	827	1,158	1,234	1,071	560	145	19	-
1999	209	464	825	1,190	1,245	1,110	537	174	21	-
Hispanic										
Both sexes										
2002	44	42	47	72	55	32	17	6	1	-
2001	27	45	48	75	52	35	13	3	-	1
2000	32	46	75	83	54	36	22	5	-	-
1999	31	42	53	57	58	41	12	5	-	-
Male										
2002	30	28	35	40	38	15	9	1	1	-
2001	21	35	36	52	26	23	4	1	-	1
2000	23	33	43	57	29	19	12	1	-	-
1999	24	30	37	31	34	29	7	1	-	-
Female										
2002	14	14	12	32	17	17	8	5	-	-
2001	6	10	12	23	26	12	9	2	-	-
2000	9	13	32	26	25	17	10	4	-	-
1999	7	12	16	26	24	12	5	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Aortic aneurysm and dissection (I71)														
Non-Hispanic[1]														
Both sexes														
2002	14,341	2	-	1	4	16	15	27	66	93	184	309	363	564
2001	14,782	-	1	-	4	12	27	41	61	126	181	268	349	509
2000	15,296	-	-	-	4	5	23	29	62	92	185	254	364	558
1999	15,393	1	1	-	2	6	21	30	57	124	182	273	338	543
Male														
2002	8,701	-	-	-	3	14	13	22	47	58	149	234	276	434
2001	8,798	-	1	-	3	10	19	29	46	99	144	204	268	383
2000	9,338	-	-	-	2	4	15	24	43	65	135	182	279	418
1999	9,342	1	-	-	2	5	12	23	51	91	124	207	257	412
Female														
2002	5,640	2	-	1	1	2	2	5	19	35	35	75	87	130
2001	5,984	-	-	-	1	2	8	12	15	27	37	64	81	126
2000	5,958	-	-	-	2	1	8	5	19	27	50	72	85	140
1999	6,051	-	1	-	-	1	9	7	6	33	58	66	81	131
Non-Hispanic White														
Both sexes														
2002	12,831	2	-	1	1	14	9	18	39	71	121	203	254	451
2001	13,217	-	1	-	4	8	20	32	45	88	115	189	246	410
2000	13,722	-	-	-	1	4	14	18	43	64	118	180	274	451
1999	13,934	1	1	-	1	6	17	22	42	95	122	184	256	450
Male														
2002	7,834	-	-	-	1	12	7	15	28	43	99	158	197	349
2001	7,924	-	1	-	3	7	14	24	34	71	92	148	194	320
2000	8,488	-	-	-	1	3	11	13	32	45	89	139	216	346
1999	8,551	1	-	-	1	5	9	17	37	70	85	145	195	353
Female														
2002	4,997	2	-	1	-	2	2	3	11	28	22	45	57	102
2001	5,293	-	-	-	1	1	6	8	11	17	23	41	52	90
2000	5,234	-	-	-	-	1	3	5	11	19	29	41	58	105
1999	5,383	-	1	-	-	1	8	5	5	25	37	39	61	97
Non-Hispanic Black														
Both sexes														
2002	1,200	-	-	-	3	1	4	9	24	19	60	95	97	99
2001	1,238	-	-	-	-	4	6	8	15	33	58	67	87	86
2000	1,262	-	-	-	3	1	8	8	17	25	61	67	68	95
1999	1,172	-	-	-	-	-	4	8	13	24	56	81	71	82
Male														
2002	664	-	-	-	2	1	4	7	17	13	48	66	72	73
2001	659	-	-	-	-	3	4	5	11	24	45	45	63	53
2000	661	-	-	-	1	1	4	8	11	18	42	38	51	61

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)										
Non-Hispanic[1]										
Both sexes										
2002	815	1,356	2,028	2,695	2,678	1,977	895	233	20	-
2001	832	1,379	2,029	2,818	2,792	2,084	1,005	238	26	-
2000	873	1,503	2,378	2,949	2,733	2,091	960	208	25	-
1999	835	1,528	2,365	3,036	2,775	2,076	921	252	27	-
Male										
2002	602	987	1,312	1,643	1,456	1,016	358	73	4	-
2001	609	942	1,357	1,665	1,560	976	400	76	7	-
2000	648	1,063	1,585	1,821	1,528	1,042	411	67	6	-
1999	633	1,079	1,556	1,873	1,559	979	390	82	6	-
Female										
2002	213	369	716	1,052	1,222	961	537	160	16	-
2001	223	437	672	1,153	1,232	1,108	605	162	19	-
2000	225	440	793	1,128	1,205	1,049	549	141	19	-
1999	202	449	809	1,163	1,216	1,097	531	170	21	-
Non-Hispanic White										
Both sexes										
2002	705	1,216	1,841	2,499	2,489	1,844	825	213	15	-
2001	707	1,228	1,851	2,561	2,594	1,951	926	217	24	-
2000	750	1,335	2,191	2,705	2,545	1,948	875	183	23	-
1999	745	1,373	2,175	2,814	2,593	1,929	858	228	22	-
Male										
2002	526	904	1,201	1,542	1,375	968	335	71	3	-
2001	521	856	1,253	1,534	1,468	927	379	71	7	-
2000	568	974	1,479	1,685	1,451	994	375	61	6	-
1999	575	992	1,459	1,774	1,469	917	369	73	5	-
Female										
2002	179	312	640	957	1,114	876	490	142	12	-
2001	186	372	598	1,027	1,126	1,024	547	146	17	-
2000	182	361	712	1,020	1,094	954	500	122	17	-
1999	170	381	716	1,040	1,124	1,012	489	155	17	-
Non-Hispanic Black										
Both sexes										
2002	93	107	137	143	142	94	48	20	5	-
2001	103	105	135	195	157	97	64	16	2	-
2000	101	129	150	188	143	113	64	20	1	-
1999	70	126	145	164	139	118	44	22	5	-
Male										
2002	65	58	74	66	53	27	15	2	1	-
2001	71	57	77	90	68	28	13	2	-	-
2000	62	68	83	99	54	35	21	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Aortic aneurysm and dissection (I71)														
Non-Hispanic Black														
Male														
1999	615	-	-	-	-	-	3	6	13	17	35	57	52	52
Female														
2002	536	-	-	-	1	-	-	2	7	6	12	29	25	26
2001	579	-	-	-	-	1	2	3	4	9	13	22	24	33
2000	601	-	-	-	2	-	4	-	6	7	19	29	17	34
1999	557	-	-	-	-	-	1	2	-	7	21	24	19	30
Origin not stated[2]														
Both sexes														
2002	46	-	-	-	-	-	-	-	1	2	2	1	1	3
2001	49	-	-	-	-	-	-	1	-	-	1	1	2	3
2000	49	-	-	-	-	-	-	1	-	1	4	1	2	1
1999	30	-	-	-	-	-	-	-	-	-	-	-	-	3
Male														
2002	28	-	-	-	-	-	-	-	1	1	1	-	1	3
2001	37	-	-	-	-	-	-	1	-	-	1	1	2	3
2000	29	-	-	-	-	-	-	-	-	1	4	1	2	1
1999	17	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	18	-	-	-	-	-	-	-	-	1	1	1	-	-
2001	12	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	20	-	-	-	-	-	-	1	-	-	-	-	-	-
1999	13	-	-	-	-	-	-	-	-	-	-	-	-	2
Other diseases of arteries, arterioles and capillaries (I72-I78)														
All origins														
Both sexes														
2002	10,109	12	4	-	6	11	16	21	29	50	90	152	208	330
2001	10,084	12	2	2	5	8	10	21	32	60	106	134	229	318
2000	10,226	16	5	3	3	7	16	11	36	54	91	156	220	281
1999	10,002	11	1	2	6	6	12	18	30	44	105	135	198	307
Male														
2002	4,296	6	4	-	2	2	9	5	14	14	48	82	110	166
2001	4,282	8	-	-	-	5	4	9	16	27	44	72	110	168
2000	4,423	6	2	1	3	2	8	6	19	20	37	84	123	169
1999	4,179	8	-	1	4	4	8	10	14	19	45	73	98	167
Female														
2002	5,813	6	-	-	4	9	7	16	15	36	42	70	98	164
2001	5,802	4	2	2	5	3	6	12	16	33	62	62	119	150

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)										
Non-Hispanic Black										
Male										
1999	44	73	74	62	62	44	12	8	1	-
Female										
2002	28	49	63	77	89	67	33	18	4	-
2001	32	48	58	105	89	69	51	14	2	-
2000	39	61	67	89	89	78	43	16	1	-
1999	26	53	71	102	77	74	32	14	4	-
Origin not stated[2]										
Both sexes										
2002	3	6	3	12	6	4	2	-	-	-
2001	3	6	10	6	8	3	4	1	-	-
2000	3	3	7	11	5	7	3	-	-	-
1999	1	8	5	2	8	1	2	-	-	-
Male										
2002	2	5	2	5	3	3	1	-	-	-
2001	3	4	8	5	5	1	2	1	-	-
2000	2	1	5	7	1	2	2	-	-	-
1999	1	5	5	1	3	-	1	-	-	-
Female										
2002	1	1	1	7	3	1	1	-	-	-
2001	-	2	2	1	3	2	2	-	-	-
2000	1	2	2	4	4	5	1	-	-	-
1999	-	3	-	1	5	1	1	-	-	-
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All origins										
Both sexes										
2002	496	733	1,095	1,513	1,714	1,755	1,237	504	133	-
2001	483	697	1,131	1,599	1,682	1,745	1,188	506	114	-
2000	462	779	1,135	1,619	1,768	1,685	1,233	544	101	1
1999	466	666	1,141	1,567	1,761	1,674	1,226	508	118	-
Male										
2002	268	388	550	718	787	641	347	117	18	-
2001	255	376	590	754	772	633	313	110	16	-
2000	250	427	614	811	765	615	309	136	15	1
1999	264	358	611	779	735	559	313	89	20	-
Female										
2002	228	345	545	795	927	1,114	890	387	115	-
2001	228	321	541	845	910	1,112	875	396	98	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of arteries, arterioles and capillaries (I72-I78)														
All origins														
Female														
2000	5,803	10	3	2	-	5	8	5	17	34	54	72	97	112
1999	5,823	3	1	1	2	2	4	8	16	25	60	62	100	140
Hispanic														
Both sexes														
2002	416	1	-	-	1	2	2	3	3	6	14	12	14	17
2001	351	2	-	1	1	3	-	3	2	3	10	13	19	19
2000	353	3	1	-	-	-	1	3	6	8	9	6	9	19
1999	345	2	-	-	1	2	3	3	3	3	7	5	6	15
Male														
2002	179	-	-	-	-	1	2	1	1	1	10	4	8	5
2001	145	1	-	-	-	2	-	-	-	-	6	6	9	10
2000	158	1	-	-	-	-	-	1	2	1	7	1	4	9
1999	160	1	-	-	-	1	3	1	1	2	3	2	1	8
Female														
2002	237	1	-	-	1	1	-	2	2	5	4	8	6	12
2001	206	1	-	1	1	1	-	3	2	3	4	7	10	9
2000	195	2	1	-	-	-	1	2	4	7	2	5	5	10
1999	185	1	-	-	1	1	-	2	2	1	4	3	5	7
Non-Hispanic[1]														
Both sexes														
2002	9,664	11	4	-	5	9	14	18	26	44	75	140	194	313
2001	9,707	10	2	1	4	5	10	18	29	57	96	121	208	299
2000	9,851	13	4	3	3	7	15	8	30	46	82	150	211	260
1999	9,630	9	1	2	5	4	9	15	27	41	98	130	191	291
Male														
2002	4,108	6	4	-	2	1	7	4	13	13	37	78	102	161
2001	4,128	7	-	-	-	3	4	9	15	27	38	66	100	158
2000	4,255	5	2	1	3	2	8	5	17	19	30	83	119	159
1999	4,009	7	-	1	4	3	5	9	13	17	42	71	97	159
Female														
2002	5,556	5	-	-	3	8	7	14	13	31	38	62	92	152
2001	5,579	3	2	1	4	2	6	9	14	30	58	55	108	141
2000	5,596	8	2	2	-	5	7	3	13	27	52	67	92	101
1999	5,621	2	1	1	1	1	4	6	14	24	56	59	94	132
Non-Hispanic White														
Both sexes														
2002	8,220	8	4	-	2	6	8	11	18	27	53	98	147	241
2001	8,222	8	1	1	3	4	8	12	19	38	62	88	159	231

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All origins										
Female										
2000	212	352	521	808	1,003	1,070	924	408	86	-
1999	202	308	530	788	1,026	1,115	913	419	98	-
Hispanic										
Both sexes										
2002	25	31	55	58	61	53	36	15	7	-
2001	22	26	37	50	39	47	30	22	2	-
2000	22	42	54	46	32	35	34	20	3	-
1999	24	31	44	40	56	51	34	11	4	-
Male										
2002	10	16	25	30	30	14	14	4	3	-
2001	12	12	17	24	17	14	9	5	1	-
2000	13	18	28	24	16	14	12	5	2	-
1999	10	14	20	26	28	19	15	3	2	-
Female										
2002	15	15	30	28	31	39	22	11	4	-
2001	10	14	20	26	22	33	21	17	1	-
2000	9	24	26	22	16	21	22	15	1	-
1999	14	17	24	14	28	32	19	8	2	-
Non-Hispanic[1]										
Both sexes										
2002	467	700	1,036	1,450	1,652	1,696	1,199	485	126	-
2001	459	669	1,093	1,546	1,639	1,690	1,155	484	112	-
2000	438	735	1,079	1,572	1,731	1,645	1,197	523	98	1
1999	440	634	1,092	1,524	1,702	1,618	1,188	495	114	-
Male										
2002	258	372	523	685	756	625	333	113	15	-
2001	242	363	572	729	754	617	304	105	15	-
2000	236	408	584	786	748	599	297	130	13	1
1999	252	343	589	753	704	540	297	85	18	-
Female										
2002	209	328	513	765	896	1,071	866	372	111	-
2001	217	306	521	817	885	1,073	851	379	97	-
2000	202	327	495	786	983	1,046	900	393	85	-
1999	188	291	503	771	998	1,078	891	410	96	-
Non-Hispanic White										
Both sexes										
2002	361	551	877	1,239	1,457	1,517	1,055	442	98	-
2001	354	554	906	1,331	1,422	1,496	1,034	411	80	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of arteries, arterioles and capillaries (I72-I78)														
Non-Hispanic White														
Both sexes														
2000	8,391	11	3	2	3	5	12	6	21	35	52	104	141	197
1999	8,257	7	1	2	4	3	7	10	20	30	76	105	145	222
Male														
2002	3,503	5	4	-	1	-	4	3	8	10	28	62	78	123
2001	3,513	5	-	-	-	3	3	7	9	16	23	47	73	126
2000	3,650	3	2	1	3	1	7	4	14	14	19	60	79	124
1999	3,463	5	-	1	3	2	4	6	11	13	36	63	75	126
Female														
2002	4,717	3	-	-	1	6	4	8	10	17	25	36	69	118
2001	4,709	3	1	1	3	1	5	5	10	22	39	41	86	105
2000	4,741	8	1	1	-	4	5	2	7	21	33	44	62	73
1999	4,794	2	1	1	1	1	3	4	9	17	40	42	70	96
Non-Hispanic Black														
Both sexes														
2002	1,295	3	-	-	2	2	5	6	7	13	20	39	41	65
2001	1,368	2	1	-	1	1	1	6	10	17	33	31	42	64
2000	1,307	2	-	1	-	1	3	2	8	9	25	39	58	53
1999	1,244	2	-	-	-	1	2	4	5	11	21	23	37	64
Male														
2002	532	1	-	-	-	-	2	1	5	2	8	13	20	35
2001	555	2	-	-	-	-	-	2	6	9	15	18	23	29
2000	532	2	-	-	-	-	1	1	2	4	9	21	34	29
1999	487	2	-	-	-	1	1	2	2	4	5	8	18	30
Female														
2002	763	2	-	-	2	2	3	5	2	11	12	26	21	30
2001	813	-	1	-	1	1	1	4	4	8	18	13	19	35
2000	775	-	-	1	-	1	2	1	6	5	16	18	24	24
1999	757	-	-	-	-	-	1	2	3	7	16	15	19	34
Origin not stated[2]														
Both sexes														
2002	29	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	26	-	-	-	-	-	-	-	1	-	-	-	2	-
2000	22	-	-	-	-	-	-	-	-	-	-	-	-	2
1999	27	-	-	-	-	-	-	-	-	-	-	-	1	1
Male														
2002	9	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	9	-	-	-	-	-	-	-	1	-	-	-	1	-
2000	10	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	10	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)										
Non-Hispanic White										
Both sexes										
2000	336	605	915	1,353	1,530	1,466	1,063	455	76	-
1999	356	518	934	1,315	1,488	1,444	1,045	432	93	-
Male										
2002	200	301	444	602	670	558	287	103	12	-
2001	191	307	481	643	658	555	262	92	12	-
2000	187	344	504	686	670	543	263	113	9	-
1999	207	289	513	649	622	488	266	73	11	-
Female										
2002	161	250	433	637	787	959	768	339	86	-
2001	163	247	425	688	764	941	772	319	68	-
2000	149	261	411	667	860	923	800	342	67	-
1999	149	229	421	666	866	956	779	359	82	-
Non-Hispanic Black										
Both sexes										
2002	93	133	144	193	175	158	132	39	25	-
2001	95	105	181	199	195	174	113	68	29	-
2000	97	117	144	198	181	163	123	62	20	1
1999	74	108	138	193	193	158	132	59	19	-
Male										
2002	50	64	72	74	76	57	42	7	3	-
2001	44	51	87	76	90	53	36	11	3	-
2000	46	58	70	91	70	47	28	14	4	1
1999	39	49	70	94	72	44	27	12	7	-
Female										
2002	43	69	72	119	99	101	90	32	22	-
2001	51	54	94	123	105	121	77	57	26	-
2000	51	59	74	107	111	116	95	48	16	-
1999	35	59	68	99	121	114	105	47	12	-
Origin not stated[2]										
Both sexes										
2002	4	2	4	5	1	6	2	4	-	-
2001	2	2	1	3	4	8	3	-	-	-
2000	2	2	2	1	5	5	2	1	-	-
1999	2	1	5	3	3	5	4	2	-	-
Male										
2002	-	-	2	3	1	2	-	-	-	-
2001	1	1	1	1	1	2	-	-	-	-
2000	1	1	2	1	1	2	-	1	-	-
1999	2	1	2	-	3	-	1	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of arteries, arterioles and capillaries (I72-I78)														
Origin not stated[2]														
Female														
2002	20	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	17	-	-	-	-	-	-	-	-	-	-	-	1	-
2000	12	-	-	-	-	-	-	-	-	-	-	-	-	1
1999	17	-	-	-	-	-	-	-	-	-	-	-	1	1
Other disorders of circulatory system (I80-I99)														
All origins														
Both sexes														
2002	4,711	27	4	1	1	12	23	38	101	128	177	251	280	302
2001	4,665	22	5	-	-	10	28	45	79	100	185	222	232	272
2000	4,603	27	4	5	7	10	27	58	75	125	194	249	266	231
1999	4,025	23	1	1	1	11	19	30	63	98	173	202	176	208
Male														
2002	1,944	15	1	-	-	9	12	12	57	67	87	137	147	168
2001	1,981	11	5	-	-	5	12	18	35	51	98	127	135	149
2000	1,981	16	2	3	3	6	10	18	39	70	120	145	150	126
1999	1,714	13	1	1	-	6	7	14	30	55	104	102	96	106
Female														
2002	2,767	12	3	1	1	3	11	26	44	61	90	114	133	134
2001	2,684	11	-	-	-	5	16	27	44	49	87	95	97	123
2000	2,622	11	2	2	4	4	17	40	36	55	74	104	116	105
1999	2,311	10	-	-	1	5	12	16	33	43	69	100	80	102
Hispanic														
Both sexes														
2002	219	7	1	-	-	2	3	6	9	10	11	17	18	14
2001	219	5	-	-	-	1	2	3	4	5	10	18	14	8
2000	229	6	-	2	2	-	3	1	10	14	18	14	20	8
1999	186	2	-	1	-	2	2	-	6	4	8	20	14	9
Male														
2002	95	5	-	-	-	1	2	3	5	5	4	7	10	7
2001	104	4	-	-	-	1	1	2	2	4	3	12	10	4
2000	116	3	-	1	1	-	1	1	6	10	10	8	11	4
1999	83	-	-	1	-	2	-	-	4	3	5	9	6	4
Female														
2002	124	2	1	-	-	1	1	3	4	5	7	10	8	7
2001	115	1	-	-	-	-	1	1	2	1	7	6	4	4
2000	113	3	-	1	1	-	2	-	4	4	8	6	9	4
1999	103	2	-	-	-	-	2	-	2	1	3	11	8	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)										
Origin not stated[2]										
Female										
2002	4	2	2	2	-	4	2	4	-	-
2001	1	1	-	2	3	6	3	-	-	-
2000	1	1	-	-	4	3	2	-	-	-
1999	-	-	3	3	-	5	3	1	-	-
Other disorders of circulatory system (I80-I99)										
All origins										
Both sexes										
2002	293	322	375	552	701	576	390	134	23	-
2001	291	366	475	572	611	568	414	140	27	1
2000	285	326	469	573	588	571	343	149	21	-
1999	237	312	380	498	619	509	322	116	25	1
Male										
2002	141	160	163	216	267	173	86	25	1	-
2001	155	174	231	254	234	164	97	19	6	1
2000	138	170	220	245	227	175	72	26	-	-
1999	118	163	187	225	238	155	66	19	8	-
Female										
2002	152	162	212	336	434	403	304	109	22	-
2001	136	192	244	318	377	404	317	121	21	-
2000	147	156	249	328	361	396	271	123	21	-
1999	119	149	193	273	381	354	256	97	17	1
Hispanic										
Both sexes										
2002	13	13	24	18	20	18	11	4	-	-
2001	8	21	27	21	20	29	21	2	-	-
2000	15	17	18	28	16	18	11	7	1	-
1999	15	15	27	8	18	25	8	1	1	-
Male										
2002	9	8	8	9	7	3	2	-	-	-
2001	5	11	9	10	10	8	8	-	-	-
2000	9	7	9	12	10	10	1	2	-	-
1999	7	5	12	6	7	9	2	-	1	-
Female										
2002	4	5	16	9	13	15	9	4	-	-
2001	3	10	18	11	10	21	13	2	-	-
2000	6	10	9	16	6	8	10	5	1	-
1999	8	10	15	2	11	16	6	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of circulatory system (I80-I99)														
Non-Hispanic[1]														
Both sexes														
2002	4,479	20	3	1	1	10	20	32	92	117	164	234	260	288
2001	4,425	17	5	-	-	9	26	42	75	94	175	204	214	264
2000	4,359	20	4	3	5	10	24	57	63	110	173	234	245	222
1999	3,825	21	1	-	1	9	17	30	56	92	164	181	162	198
Male														
2002	1,842	10	1	-	-	8	10	9	52	61	82	130	135	161
2001	1,869	7	5	-	-	4	11	16	33	46	95	115	124	145
2000	1,856	12	2	2	2	6	9	17	32	60	107	136	138	121
1999	1,623	13	1	-	-	4	7	14	26	50	98	92	90	101
Female														
2002	2,637	10	2	1	1	2	10	23	40	56	82	104	125	127
2001	2,556	10	-	-	-	5	15	26	42	48	80	89	90	119
2000	2,503	8	2	1	3	4	15	40	31	50	66	98	107	101
1999	2,202	8	-	-	1	5	10	16	30	42	66	89	72	97
Non-Hispanic White														
Both sexes														
2002	3,630	13	2	-	1	6	10	17	52	72	108	168	195	224
2001	3,617	12	3	-	-	4	14	27	46	56	119	145	161	205
2000	3,554	13	3	2	3	4	15	32	38	72	123	174	184	166
1999	3,149	17	-	-	1	3	9	21	36	60	108	126	117	167
Male														
2002	1,450	8	-	-	-	5	4	3	30	39	53	90	104	123
2001	1,496	4	3	-	-	2	6	9	21	28	64	90	92	114
2000	1,496	9	1	2	1	3	5	8	18	37	81	102	105	89
1999	1,310	9	-	-	-	2	5	10	16	32	63	64	63	87
Female														
2002	2,180	5	2	-	1	1	6	14	22	33	55	78	91	101
2001	2,121	8	-	-	-	2	8	18	25	28	55	55	69	91
2000	2,058	4	2	-	2	1	10	24	20	35	42	72	79	77
1999	1,839	8	-	-	1	1	4	11	20	28	45	62	54	80
Non-Hispanic Black														
Both sexes														
2002	789	7	1	1	-	4	9	15	39	42	52	60	60	62
2001	729	5	2	-	-	3	11	15	28	35	52	52	46	55
2000	751	5	1	1	2	6	8	23	25	36	50	58	58	53
1999	613	3	1	-	-	6	8	9	19	29	52	54	41	27
Male														
2002	359	2	1	-	-	3	6	6	21	20	28	36	27	36
2001	335	3	2	-	-	1	4	7	11	16	30	20	28	30

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)										
Non-Hispanic[1]										
Both sexes										
2002	278	308	351	534	678	557	379	129	23	-
2001	280	342	446	549	587	538	393	137	27	1
2000	269	307	451	545	572	553	332	140	20	-
1999	222	296	352	488	600	482	313	115	24	1
Male										
2002	131	151	155	207	259	170	84	25	1	-
2001	148	162	222	243	223	155	89	19	6	1
2000	129	162	211	233	217	165	71	24	-	-
1999	111	157	174	219	231	145	64	19	7	-
Female										
2002	147	157	196	327	419	387	295	104	22	-
2001	132	180	224	306	364	383	304	118	21	-
2000	140	145	240	312	355	388	261	116	20	-
1999	111	139	178	269	369	337	249	96	17	1
Non-Hispanic White										
Both sexes										
2002	206	247	291	455	586	497	344	115	21	-
2001	209	270	372	459	528	490	349	125	23	-
2000	207	225	377	462	513	498	300	123	20	-
1999	175	227	297	419	530	424	282	106	23	1
Male										
2002	101	119	129	178	218	151	73	21	1	-
2001	106	128	185	205	200	141	79	15	4	-
2000	99	124	183	203	195	148	63	20	-	-
1999	93	118	150	185	203	128	57	18	7	-
Female										
2002	105	128	162	277	368	346	271	94	20	-
2001	103	142	187	254	328	349	270	110	19	-
2000	108	101	194	259	318	350	237	103	20	-
1999	82	109	147	234	327	296	225	88	16	1
Non-Hispanic Black										
Both sexes										
2002	66	59	54	74	85	53	32	12	2	-
2001	67	64	65	79	54	41	39	12	3	1
2000	56	77	68	80	50	51	28	15	-	-
1999	42	62	47	64	61	51	28	8	1	-
Male										
2002	28	30	23	26	37	15	10	4	-	-
2001	42	31	31	34	19	11	8	4	2	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of circulatory system (I80-I99)														
Non-Hispanic Black														
Male														
2000	333	2	1	-	1	3	4	8	14	22	26	34	32	29
1999	281	3	1	-	-	2	2	4	10	16	33	28	24	11
Female														
2002	430	5	-	1	-	1	3	9	18	22	24	24	33	26
2001	394	2	-	-	-	2	7	8	17	19	22	32	18	25
2000	418	3	-	1	1	3	4	15	11	14	24	24	26	24
1999	332	-	-	-	-	4	6	5	9	13	19	26	17	16
Origin not stated[2]														
Both sexes														
2002	13	-	-	-	-	-	-	-	-	1	2	-	2	-
2001	21	-	-	-	-	-	-	-	-	1	-	-	4	-
2000	15	1	-	-	-	-	-	-	2	1	3	1	1	1
1999	14	-	-	-	-	-	-	-	1	2	1	1	-	1
Male														
2002	7	-	-	-	-	-	-	-	-	1	1	-	2	-
2001	8	-	-	-	-	-	-	-	-	1	-	-	1	-
2000	9	1	-	-	-	-	-	-	1	-	3	1	1	1
1999	8	-	-	-	-	-	-	-	-	2	1	1	-	1
Female														
2002	6	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	13	-	-	-	-	-	-	-	-	-	-	-	3	-
2000	6	-	-	-	-	-	-	-	1	1	-	-	-	-
1999	6	-	-	-	-	-	-	-	1	-	-	-	-	-
Influenza and pneumonia (J10-J18)														
All origins														
Both sexes														
2002	65,681	263	110	38	53	75	92	135	210	367	604	887	1,031	1,271
2001	62,034	299	112	46	46	66	115	124	215	393	590	810	991	1,155
2000	65,313	289	103	47	40	65	124	142	222	405	663	783	991	1,198
1999	63,730	320	130	46	47	73	106	135	204	462	601	798	899	1,098
Male														
2002	28,918	162	63	15	23	41	56	68	122	197	349	519	619	742
2001	27,342	167	57	23	24	36	70	64	127	219	337	497	602	686
2000	28,658	163	54	27	21	35	67	82	122	211	419	464	602	710
1999	27,718	171	66	27	26	34	56	76	121	251	341	500	559	611
Female														
2002	36,763	101	47	23	30	34	36	67	88	170	255	368	412	529

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)										
Non-Hispanic Black										
Male										
2000	27	35	26	27	16	14	8	4	-	-
1999	17	36	20	31	24	13	5	1	-	-
Female										
2002	38	29	31	48	48	38	22	8	2	-
2001	25	33	34	45	35	30	31	8	1	-
2000	29	42	42	53	34	37	20	11	-	-
1999	25	26	27	33	37	38	23	7	1	-
Origin not stated[2]										
Both sexes										
2002	2	1	-	-	3	1	-	1	-	-
2001	3	3	2	2	4	1	-	1	-	-
2000	1	2	-	-	-	-	-	2	-	-
1999	-	1	1	2	1	2	1	-	-	-
Male										
2002	1	1	-	-	1	-	-	-	-	-
2001	2	1	-	1	1	1	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	1	1	-	-	1	-	-	-	-
Female										
2002	1	-	-	-	2	1	-	1	-	-
2001	1	2	2	1	3	-	-	1	-	-
2000	1	1	-	-	-	-	-	2	-	-
1999	-	-	-	2	1	1	1	-	-	-
Influenza and pneumonia (J10-J18)										
All origins										
Both sexes										
2002	1,716	2,420	4,427	7,905	12,079	13,977	11,490	5,172	1,356	3
2001	1,549	2,323	4,327	7,510	11,167	13,382	10,692	4,836	1,281	5
2000	1,681	2,533	4,656	7,912	11,909	13,752	11,318	5,116	1,361	3
1999	1,527	2,382	4,479	7,776	11,416	13,499	11,151	5,159	1,420	2
Male										
2002	996	1,384	2,466	4,114	5,781	5,716	3,875	1,352	256	2
2001	901	1,338	2,394	3,995	5,299	5,373	3,587	1,282	260	4
2000	970	1,447	2,566	4,084	5,667	5,575	3,724	1,373	273	2
1999	851	1,394	2,499	4,020	5,438	5,501	3,595	1,295	285	1
Female										
2002	720	1,036	1,961	3,791	6,298	8,261	7,615	3,820	1,100	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza and pneumonia (J10-J18)														
All origins														
Female														
2001	34,692	132	55	23	22	30	45	60	88	174	253	313	389	469
2000	36,655	126	49	20	19	30	57	60	100	194	244	319	389	488
1999	36,012	149	64	19	21	39	50	59	83	211	260	298	340	487
Hispanic														
Both sexes														
2002	2,824	48	17	6	8	9	10	18	25	37	40	61	63	95
2001	2,722	57	32	5	5	7	11	10	13	32	41	56	65	72
2000	2,625	60	17	13	8	4	17	19	29	33	49	54	66	82
1999	2,246	64	17	5	5	10	15	10	14	46	28	48	56	58
Male														
2002	1,398	33	11	2	4	6	5	12	14	23	29	41	45	60
2001	1,309	31	16	3	3	6	9	7	7	20	28	47	38	50
2000	1,303	30	5	7	3	1	9	10	17	21	35	41	41	50
1999	1,092	40	10	4	2	5	6	6	9	34	20	32	39	32
Female														
2002	1,426	15	6	4	4	3	5	6	11	14	11	20	18	35
2001	1,413	26	16	2	2	1	2	3	6	12	13	9	27	22
2000	1,322	30	12	6	5	3	8	9	12	12	14	13	25	32
1999	1,154	24	7	1	3	5	9	4	5	12	8	16	17	26
Non-Hispanic[1]														
Both sexes														
2002	62,657	213	92	32	44	65	82	116	184	328	561	820	963	1,172
2001	59,103	240	79	41	41	59	104	112	200	360	546	752	917	1,072
2000	62,440	228	84	34	32	60	107	121	193	366	608	723	916	1,107
1999	61,218	255	110	41	42	63	90	125	190	413	566	741	836	1,029
Male														
2002	27,417	128	52	13	19	34	51	55	107	174	317	473	570	679
2001	25,926	135	41	20	21	30	61	57	118	198	306	448	559	630
2000	27,218	132	47	20	18	33	58	70	105	187	379	419	553	652
1999	26,489	131	55	23	24	29	50	70	112	214	317	461	513	573
Female														
2002	35,240	85	40	19	25	31	31	61	77	154	244	347	393	493
2001	33,177	105	38	21	20	29	43	55	82	162	240	304	358	442
2000	35,222	96	37	14	14	27	49	51	88	179	229	304	363	455
1999	34,729	124	55	18	18	34	40	55	78	199	249	280	323	456
Non-Hispanic White														
Both sexes														
2002	55,419	114	46	23	32	42	61	83	124	219	398	611	718	887
2001	51,952	117	48	28	29	39	69	72	135	243	362	516	677	831

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza and pneumonia (J10-J18)										
All origins										
Female										
2001	648	985	1,933	3,515	5,868	8,009	7,105	3,554	1,021	1
2000	711	1,086	2,090	3,828	6,242	8,177	7,594	3,743	1,088	1
1999	676	988	1,980	3,756	5,978	7,998	7,556	3,864	1,135	1
Hispanic										
Both sexes										
2002	112	150	249	385	470	443	358	177	43	-
2001	96	140	262	346	415	465	382	173	36	1
2000	109	138	267	313	377	440	339	161	30	-
1999	97	143	185	252	317	383	296	150	47	-
Male										
2002	70	88	141	189	233	186	137	55	14	-
2001	49	71	133	180	208	177	145	69	11	1
2000	69	75	143	166	201	164	144	62	9	-
1999	48	82	105	129	155	155	118	49	12	-
Female										
2002	42	62	108	196	237	257	221	122	29	-
2001	47	69	129	166	207	288	237	104	25	-
2000	40	63	124	147	176	276	195	99	21	-
1999	49	61	80	123	162	228	178	101	35	-
Non-Hispanic[1]										
Both sexes										
2002	1,601	2,260	4,164	7,492	11,567	13,504	11,101	4,983	1,310	3
2001	1,444	2,172	4,037	7,140	10,723	12,879	10,292	4,652	1,239	2
2000	1,567	2,378	4,372	7,565	11,486	13,278	10,947	4,938	1,329	1
1999	1,417	2,223	4,276	7,494	11,059	13,060	10,823	4,995	1,369	1
Male										
2002	924	1,287	2,313	3,912	5,525	5,523	3,726	1,293	240	2
2001	845	1,259	2,241	3,802	5,079	5,181	3,436	1,210	248	1
2000	897	1,361	2,411	3,899	5,440	5,395	3,571	1,306	264	1
1999	794	1,303	2,384	3,875	5,263	5,318	3,467	1,241	272	-
Female										
2002	677	973	1,851	3,580	6,042	7,981	7,375	3,690	1,070	1
2001	599	913	1,796	3,338	5,644	7,698	6,856	3,442	991	1
2000	670	1,017	1,961	3,666	6,046	7,883	7,376	3,632	1,065	-
1999	623	920	1,892	3,619	5,796	7,742	7,356	3,754	1,097	1
Non-Hispanic White										
Both sexes										
2002	1,265	1,866	3,553	6,568	10,433	12,381	10,228	4,585	1,179	3
2001	1,132	1,773	3,441	6,188	9,649	11,821	9,463	4,242	1,076	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza and pneumonia (J10-J18)														
Non-Hispanic White														
Both sexes														
2000	55,135	114	51	23	26	43	73	79	134	268	410	501	664	844
1999	54,280	121	58	31	31	44	51	75	124	283	354	503	606	797
Male														
2002	23,934	73	24	8	12	24	38	38	73	113	227	344	424	508
2001	22,389	71	26	14	14	18	46	33	79	131	202	311	396	484
2000	23,609	68	24	15	14	25	43	41	70	137	257	280	387	480
1999	23,097	64	22	17	19	21	30	44	75	148	186	309	361	436
Female														
2002	31,485	41	22	15	20	18	23	45	51	106	171	267	294	379
2001	29,563	46	22	14	15	21	23	39	56	112	160	205	281	347
2000	31,526	46	27	8	12	18	30	38	64	131	153	221	277	364
1999	31,183	57	36	14	12	23	21	31	49	135	168	194	245	361
Non-Hispanic Black														
Both sexes														
2002	5,803	86	40	7	11	18	20	27	53	101	149	189	210	254
2001	5,686	103	24	9	8	16	29	31	56	109	165	219	211	213
2000	5,920	101	29	7	4	14	30	37	55	83	171	197	226	237
1999	5,794	114	40	8	7	14	35	47	60	116	189	220	205	208
Male														
2002	2,734	47	22	4	6	8	12	13	30	54	82	116	123	158
2001	2,769	52	13	4	5	11	12	17	35	62	93	128	142	129
2000	2,880	55	20	4	3	6	15	26	32	42	109	124	147	155
1999	2,786	58	25	5	3	6	17	25	33	58	115	140	137	125
Female														
2002	3,069	39	18	3	5	10	8	14	23	47	67	73	87	96
2001	2,917	51	11	5	3	5	17	14	21	47	72	91	69	84
2000	3,040	46	9	3	1	8	15	11	23	41	62	73	79	82
1999	3,008	56	15	3	4	8	18	22	27	58	74	80	68	83
Origin not stated[2]														
Both sexes														
2002	200	2	1	-	1	1	-	1	1	2	3	6	5	4
2001	209	2	1	-	-	-	-	2	2	1	3	2	9	11
2000	248	1	2	-	-	1	-	2	-	6	6	6	9	9
1999	266	1	3	-	-	-	1	-	-	3	7	9	7	11
Male														
2002	103	1	-	-	-	1	-	1	1	-	3	5	4	3
2001	107	1	-	-	-	-	-	-	2	1	3	2	5	6
2000	137	1	2	-	-	1	-	2	-	3	5	4	8	8
1999	137	-	1	-	-	-	-	-	-	3	4	7	7	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza and pneumonia (J10-J18)										
Non-Hispanic White										
Both sexes										
2000	1,217	1,884	3,662	6,696	10,372	12,218	10,117	4,554	1,184	1
1999	1,097	1,843	3,638	6,686	10,099	11,998	9,993	4,610	1,237	1
Male										
2002	720	1,059	1,991	3,427	4,975	5,057	3,416	1,164	217	2
2001	648	1,013	1,906	3,268	4,562	4,746	3,119	1,089	212	1
2000	696	1,065	2,019	3,446	4,900	4,952	3,263	1,194	232	1
1999	613	1,075	2,012	3,454	4,801	4,862	3,160	1,139	249	-
Female										
2002	545	807	1,562	3,141	5,458	7,324	6,812	3,421	962	1
2001	484	760	1,535	2,920	5,087	7,075	6,344	3,153	864	-
2000	521	819	1,643	3,250	5,472	7,266	6,854	3,360	952	-
1999	484	768	1,626	3,232	5,298	7,136	6,833	3,471	988	1
Non-Hispanic Black										
Both sexes										
2002	300	327	510	708	876	849	651	310	107	-
2001	258	315	481	756	815	784	629	320	135	-
2000	294	408	576	689	861	815	653	311	122	-
1999	278	327	553	659	751	869	671	311	112	-
Male										
2002	180	188	266	373	416	318	208	92	18	-
2001	164	195	277	429	379	289	224	82	27	-
2000	167	244	318	360	403	321	218	86	25	-
1999	157	198	320	336	352	352	224	81	19	-
Female										
2002	120	139	244	335	460	531	443	218	89	-
2001	94	120	204	327	436	495	405	238	108	-
2000	127	164	258	329	458	494	435	225	97	-
1999	121	129	233	323	399	517	447	230	93	-
Origin not stated[2]										
Both sexes										
2002	3	10	14	28	42	30	31	12	3	-
2001	9	11	28	24	29	38	18	11	6	2
2000	5	17	17	34	46	34	32	17	2	2
1999	13	16	18	30	40	56	32	14	4	1
Male										
2002	2	9	12	13	23	7	12	4	2	-
2001	7	8	20	13	12	15	6	3	1	2
2000	4	11	12	19	26	16	9	5	-	1
1999	9	9	10	16	20	28	10	5	1	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza and pneumonia (J10-J18)														
Origin not stated[2]														
Female														
2002	97	1	1	-	1	-	-	-	-	2	-	1	1	1
2001	102	1	1	-	-	-	-	2	-	-	-	-	4	5
2000	111	-	-	-	-	-	-	-	-	3	1	2	1	1
1999	129	1	2	-	-	-	1	-	-	-	3	2	-	5
Influenza (J10-J11)														
All origins														
Both sexes														
2002	727	7	5	5	7	5	2	3	1	3	5	8	9	13
2001	257	7	6	9	3	4	3	4	6	1	5	8	7	8
2000	1,765	9	10	3	8	8	5	8	5	8	14	24	24	39
1999	1,665	13	12	6	5	7	4	2	7	8	8	13	13	29
Male														
2002	289	5	3	1	1	2	2	2	-	2	2	7	5	7
2001	101	6	2	3	1	3	3	3	3	1	1	5	4	4
2000	664	5	4	1	4	3	3	4	1	5	9	11	14	20
1999	602	6	9	4	1	1	1	-	5	5	4	8	7	13
Female														
2002	438	2	2	4	6	3	-	1	1	1	3	1	4	6
2001	156	1	4	6	2	1	-	1	3	-	4	3	3	4
2000	1,101	4	6	2	4	5	2	4	4	3	5	13	10	19
1999	1,063	7	3	2	4	6	3	2	2	3	4	5	6	16
Hispanic														
Both sexes														
2002	19	2	-	3	1	-	-	1	-	1	-	1	-	-
2001	13	-	3	1	-	1	-	-	-	-	1	1	-	-
2000	47	2	1	1	2	1	-	3	1	2	-	2	-	1
1999	45	4	3	-	-	1	1	-	-	1	1	-	2	2
Male														
2002	10	2	-	1	1	-	-	1	-	1	-	1	-	-
2001	7	-	2	1	-	1	-	-	-	-	-	1	-	-
2000	21	-	1	-	1	1	-	1	-	1	-	1	-	1
1999	21	-	3	-	-	-	-	-	-	1	1	-	2	2
Female														
2002	9	-	-	2	-	-	-	-	-	-	-	-	-	-
2001	6	-	1	-	-	-	-	-	-	-	1	-	-	-
2000	26	2	-	1	1	-	-	2	1	1	-	1	-	-
1999	24	4	-	-	-	1	1	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza and pneumonia (J10-J18)										
Origin not stated[2]										
Female										
2002	1	1	2	15	19	23	19	8	1	-
2001	2	3	8	11	17	23	12	8	5	-
2000	1	6	5	15	20	18	23	12	2	1
1999	4	7	8	14	20	28	22	9	3	-
Influenza (J10-J11)										
All origins										
Both sexes										
2002	17	19	37	59	118	174	144	59	27	-
2001	13	10	11	18	38	37	39	14	6	-
2000	54	76	123	189	303	372	305	130	48	-
1999	42	60	104	154	297	368	303	162	48	-
Male										
2002	13	13	19	28	53	66	45	10	3	-
2001	5	3	5	6	14	15	8	4	2	-
2000	24	43	63	81	115	127	86	35	6	-
1999	24	27	47	71	131	127	70	35	6	-
Female										
2002	4	6	18	31	65	108	99	49	24	-
2001	8	7	6	12	24	22	31	10	4	-
2000	30	33	60	108	188	245	219	95	42	-
1999	18	33	57	83	166	241	233	127	42	-
Hispanic										
Both sexes										
2002	-	1	1	2	1	-	5	-	-	-
2001	-	-	1	3	2	-	-	-	-	-
2000	1	3	1	3	7	10	3	3	-	-
1999	3	4	4	5	4	3	5	2	-	-
Male										
2002	-	1	-	-	1	-	1	-	-	-
2001	-	-	-	-	2	-	-	-	-	-
2000	1	2	-	1	4	3	1	2	-	-
1999	2	4	3	1	1	-	-	1	-	-
Female										
2002	-	-	1	2	-	-	4	-	-	-
2001	-	-	1	3	-	-	-	-	-	-
2000	-	1	1	2	3	7	2	1	-	-
1999	1	-	1	4	3	3	5	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza (J10-J11)														
Non-Hispanic[1]														
Both sexes														
2002	704	5	5	2	6	5	2	2	1	2	5	7	9	13
2001	244	7	3	8	3	3	3	4	6	1	4	7	7	8
2000	1,716	7	9	2	6	7	5	5	4	6	14	22	24	38
1999	1,612	9	9	6	5	6	3	2	7	7	7	13	11	27
Male														
2002	277	3	3	-	-	2	2	1	-	1	2	6	5	7
2001	94	6	-	2	1	2	3	3	3	1	1	4	4	4
2000	642	5	3	1	3	2	3	3	1	4	9	10	14	19
1999	580	6	6	4	1	1	1	-	5	4	3	8	5	11
Female														
2002	427	2	2	2	6	3	-	1	1	1	3	1	4	6
2001	150	1	3	6	2	1	-	1	3	-	3	3	3	4
2000	1,074	2	6	1	3	5	2	2	3	2	5	12	10	19
1999	1,032	3	3	2	4	5	2	2	2	3	4	5	6	16
Non-Hispanic White														
Both sexes														
2002	663	3	2	1	5	3	2	2	1	2	5	7	8	11
2001	216	7	2	4	2	2	2	2	5	1	2	7	7	7
2000	1,613	5	8	1	5	7	4	4	3	5	11	16	22	36
1999	1,522	3	5	6	4	5	3	1	7	3	5	12	8	24
Male														
2002	258	1	1	-	-	-	2	1	-	1	2	6	5	5
2001	81	6	-	1	1	1	2	1	2	1	1	4	4	3
2000	595	5	3	1	2	2	2	2	-	3	6	6	12	17
1999	547	1	3	4	1	1	1	-	5	2	2	7	4	11
Female														
2002	405	2	1	1	5	3	-	1	1	1	3	1	3	6
2001	135	1	2	3	1	1	-	1	3	-	1	3	3	4
2000	1,018	-	5	-	3	5	2	2	3	2	5	10	10	19
1999	975	2	2	2	3	4	2	1	2	1	3	5	4	13
Non-Hispanic Black														
Both sexes														
2002	30	1	3	1	1	2	-	-	-	-	-	-	-	1
2001	20	-	-	2	-	1	-	1	1	-	1	-	-	1
2000	81	2	1	-	1	-	1	1	-	-	3	5	2	2
1999	65	4	1	-	-	1	-	1	-	3	2	1	2	3
Male														
2002	15	1	2	-	-	2	-	-	-	-	-	-	-	1
2001	10	-	-	-	-	1	-	1	1	-	-	-	-	1
2000	35	-	-	-	1	-	1	1	-	-	3	3	2	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)										
Non-Hispanic[1]										
Both sexes										
2002	17	18	36	56	117	174	137	58	27	-
2001	13	10	10	15	36	37	39	14	6	-
2000	53	73	122	186	295	361	302	127	48	-
1999	39	55	99	149	290	364	296	160	48	-
Male										
2002	13	12	19	27	52	66	43	10	3	-
2001	5	3	5	6	12	15	8	4	2	-
2000	23	41	63	80	110	124	85	33	6	-
1999	22	23	43	70	130	127	70	34	6	-
Female										
2002	4	6	17	29	65	108	94	48	24	-
2001	8	7	5	9	24	22	31	10	4	-
2000	30	32	59	106	185	237	217	94	42	-
1999	17	32	56	79	160	237	226	126	42	-
Non-Hispanic White										
Both sexes										
2002	14	12	33	54	114	172	129	57	26	-
2001	10	8	10	15	32	34	38	13	6	-
2000	49	67	109	177	281	342	292	122	47	-
1999	36	51	88	141	276	352	291	155	46	-
Male										
2002	11	8	19	26	50	65	42	10	3	-
2001	3	1	5	6	12	14	7	4	2	-
2000	22	38	59	74	107	116	82	30	6	-
1999	21	20	39	69	123	124	70	33	6	-
Female										
2002	3	4	14	28	64	107	87	47	23	-
2001	7	7	5	9	20	20	31	9	4	-
2000	27	29	50	103	174	226	210	92	41	-
1999	15	31	49	72	153	228	221	122	40	-
Non-Hispanic Black										
Both sexes										
2002	3	4	2	1	3	1	5	1	1	-
2001	2	2	-	-	4	3	1	1	-	-
2000	4	5	11	5	8	17	8	4	1	-
1999	2	2	10	5	10	8	4	4	2	-
Male										
2002	2	3	-	-	2	1	1	-	-	-
2001	2	2	-	-	-	1	1	-	-	-
2000	1	2	3	4	1	7	1	3	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Influenza (J10-J11)														
Non-Hispanic Black														
Male														
1999	24	3	1	-	-	-	-	-	-	1	1	1	1	-
Female														
2002	15	-	1	1	1	-	-	-	-	-	-	-	-	-
2001	10	-	-	2	-	-	-	-	-	-	1	-	-	-
2000	46	2	1	-	-	-	-	-	-	-	-	2	-	-
1999	41	1	-	-	-	1	-	1	-	2	1	-	1	3
Origin not stated[2]														
Both sexes														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	8	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	7	-	-	-	-	-	-	-	-	-	-	-	-	-
Pneumonia (J12-J18)														
All origins														
Both sexes														
2002	64,954	256	105	33	46	70	90	132	209	364	599	879	1,022	1,258
2001	61,777	292	106	37	43	62	112	120	209	392	585	802	984	1,147
2000	63,548	280	93	44	32	57	119	134	217	397	649	759	967	1,159
1999	62,065	307	118	40	42	66	102	133	197	454	593	785	886	1,069
Male														
2002	28,629	157	60	14	22	39	54	66	122	195	347	512	614	735
2001	27,241	161	55	20	23	33	67	61	124	218	336	492	598	682
2000	27,994	158	50	26	17	32	64	78	121	206	410	453	588	690
1999	27,116	165	57	23	25	33	55	76	116	246	337	492	552	598
Female														
2002	36,325	99	45	19	24	31	36	66	87	169	252	367	408	523
2001	34,536	131	51	17	20	29	45	59	85	174	249	310	386	465
2000	35,554	122	43	18	15	25	55	56	96	191	239	306	379	469

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Influenza (J10-J11)										
Non-Hispanic Black										
Male										
1999	1	2	4	1	5	2	-	1	-	-
Female										
2002	1	1	2	1	1	-	4	1	1	-
2001	-	-	-	-	4	2	-	1	-	-
2000	3	3	8	1	7	10	7	1	1	-
1999	1	-	6	4	5	6	4	3	2	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	1	-	-	2	1	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	1	-	-	-	-
1999	-	1	1	-	3	1	2	-	-	-
Male										
2002	-	-	-	1	-	-	1	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	1	1	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	1	-	-	-	-
1999	-	1	-	-	3	1	2	-	-	-
Pneumonia (J12-J18)										
All origins										
Both sexes										
2002	1,699	2,401	4,390	7,846	11,961	13,803	11,346	5,113	1,329	3
2001	1,536	2,313	4,316	7,492	11,129	13,345	10,653	4,822	1,275	5
2000	1,627	2,457	4,533	7,723	11,606	13,380	11,013	4,986	1,313	3
1999	1,485	2,322	4,375	7,622	11,119	13,131	10,848	4,997	1,372	2
Male										
2002	983	1,371	2,447	4,086	5,728	5,650	3,830	1,342	253	2
2001	896	1,335	2,389	3,989	5,285	5,358	3,579	1,278	258	4
2000	946	1,404	2,503	4,003	5,552	5,448	3,638	1,338	267	2
1999	827	1,367	2,452	3,949	5,307	5,374	3,525	1,260	279	1
Female										
2002	716	1,030	1,943	3,760	6,233	8,153	7,516	3,771	1,076	1
2001	640	978	1,927	3,503	5,844	7,987	7,074	3,544	1,017	1
2000	681	1,053	2,030	3,720	6,054	7,932	7,375	3,648	1,046	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonia (J12-J18)														
All origins														
Female														
1999	34,949	142	61	17	17	33	47	57	81	208	256	293	334	471
Hispanic														
Both sexes														
2002	2,805	46	17	3	7	9	10	17	25	36	40	60	63	95
2001	2,709	57	29	4	5	6	11	10	13	32	40	55	65	72
2000	2,578	58	16	12	6	3	17	16	28	31	49	52	66	81
1999	2,201	60	14	5	5	9	14	10	14	45	27	48	54	56
Male														
2002	1,388	31	11	1	3	6	5	11	14	22	29	40	45	60
2001	1,302	31	14	2	3	5	9	7	7	20	28	46	38	50
2000	1,282	30	4	7	2	-	9	9	17	20	35	40	41	49
1999	1,071	40	7	4	2	5	6	6	9	33	19	32	37	30
Female														
2002	1,417	15	6	2	4	3	5	6	11	14	11	20	18	35
2001	1,407	26	15	2	2	1	2	3	6	12	12	9	27	22
2000	1,296	28	12	5	4	3	8	7	11	11	14	12	25	32
1999	1,130	20	7	1	3	4	8	4	5	12	8	16	17	26
Non-Hispanic[1]														
Both sexes														
2002	61,953	208	87	30	38	60	80	114	183	326	556	813	954	1,159
2001	58,859	233	76	33	38	56	101	108	194	359	542	745	910	1,064
2000	60,724	221	75	32	26	53	102	116	189	360	594	701	892	1,069
1999	59,606	246	101	35	37	57	87	123	183	406	559	728	825	1,002
Male														
2002	27,140	125	49	13	19	32	49	54	107	173	315	467	565	672
2001	25,832	129	41	18	20	28	58	54	115	197	305	444	555	626
2000	26,576	127	44	19	15	31	55	67	104	183	370	409	539	633
1999	25,909	125	49	19	23	28	49	70	107	210	314	453	508	562
Female														
2002	34,813	83	38	17	19	28	31	60	76	153	241	346	389	487
2001	33,027	104	35	15	18	28	43	54	79	162	237	301	355	438
2000	34,148	94	31	13	11	22	47	49	85	177	224	292	353	436
1999	33,697	121	52	16	14	29	38	53	76	196	245	275	317	440
Non-Hispanic White														
Both sexes														
2002	54,756	111	44	22	27	39	59	81	123	217	393	604	710	876
2001	51,736	110	46	24	27	37	67	70	130	242	360	509	670	824
2000	53,522	109	43	22	21	36	69	75	131	263	399	485	642	808
1999	52,758	118	53	25	27	39	48	74	117	280	349	491	598	773

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)										
All origins										
Female										
1999	658	955	1,923	3,673	5,812	7,757	7,323	3,737	1,093	1
Hispanic										
Both sexes										
2002	112	149	248	383	469	443	353	177	43	-
2001	96	140	261	343	413	465	382	173	36	1
2000	108	135	266	310	370	430	336	158	30	-
1999	94	139	181	247	313	380	291	148	47	-
Male										
2002	70	87	141	189	232	186	136	55	14	-
2001	49	71	133	180	206	177	145	69	11	1
2000	68	73	143	165	197	161	143	60	9	-
1999	46	78	102	128	154	155	118	48	12	-
Female										
2002	42	62	107	194	237	257	217	122	29	-
2001	47	69	128	163	207	288	237	104	25	-
2000	40	62	123	145	173	269	193	98	21	-
1999	48	61	79	119	159	225	173	100	35	-
Non-Hispanic[1]										
Both sexes										
2002	1,584	2,242	4,128	7,436	11,450	13,330	10,964	4,925	1,283	3
2001	1,431	2,162	4,027	7,125	10,687	12,842	10,253	4,638	1,233	2
2000	1,514	2,305	4,250	7,379	11,191	12,917	10,645	4,811	1,281	1
1999	1,378	2,168	4,177	7,345	10,769	12,696	10,527	4,835	1,321	1
Male										
2002	911	1,275	2,294	3,885	5,473	5,457	3,683	1,283	237	2
2001	840	1,256	2,236	3,796	5,067	5,166	3,428	1,206	246	1
2000	874	1,320	2,348	3,819	5,330	5,271	3,486	1,273	258	1
1999	772	1,280	2,341	3,805	5,133	5,191	3,397	1,207	266	-
Female										
2002	673	967	1,834	3,551	5,977	7,873	7,281	3,642	1,046	1
2001	591	906	1,791	3,329	5,620	7,676	6,825	3,432	987	1
2000	640	985	1,902	3,560	5,861	7,646	7,159	3,538	1,023	-
1999	606	888	1,836	3,540	5,636	7,505	7,130	3,628	1,055	1
Non-Hispanic White										
Both sexes										
2002	1,251	1,854	3,520	6,514	10,319	12,209	10,099	4,528	1,153	3
2001	1,122	1,765	3,431	6,173	9,617	11,787	9,425	4,229	1,070	1
2000	1,168	1,817	3,553	6,519	10,091	11,876	9,825	4,432	1,137	1
1999	1,061	1,792	3,550	6,545	9,823	11,646	9,702	4,455	1,191	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonia (J12-J18)														
Non-Hispanic White														
Male														
2002	23,676	72	23	8	12	24	36	37	73	112	225	338	419	503
2001	22,308	65	26	13	13	17	44	32	77	130	201	307	392	481
2000	23,014	63	21	14	12	23	41	39	70	134	251	274	375	463
1999	22,550	63	19	13	18	20	29	44	70	146	184	302	357	425
Female														
2002	31,080	39	21	14	15	15	23	44	50	105	168	266	291	373
2001	29,428	45	20	11	14	20	23	38	53	112	159	202	278	343
2000	30,508	46	22	8	9	13	28	36	61	129	148	211	267	345
1999	30,208	55	34	12	9	19	19	30	47	134	165	189	241	348
Non-Hispanic Black														
Both sexes														
2002	5,773	85	37	6	10	16	20	27	53	101	149	189	210	253
2001	5,666	103	24	7	8	15	29	30	55	109	164	219	211	212
2000	5,839	99	28	7	3	14	29	36	55	83	168	192	224	235
1999	5,729	110	39	8	7	13	35	46	60	113	187	219	203	205
Male														
2002	2,719	46	20	4	6	6	12	13	30	54	82	116	123	157
2001	2,759	52	13	4	5	10	12	16	34	62	93	128	142	128
2000	2,845	55	20	4	2	6	14	25	32	42	106	121	145	153
1999	2,762	55	24	5	3	6	17	25	33	57	114	139	136	125
Female														
2002	3,054	39	17	2	4	10	8	14	23	47	67	73	87	96
2001	2,907	51	11	3	3	5	17	14	21	47	71	91	69	84
2000	2,994	44	8	3	1	8	15	11	23	41	62	71	79	82
1999	2,967	55	15	3	4	7	18	21	27	56	73	80	67	80
Origin not stated[2]														
Both sexes														
2002	196	2	1	-	1	1	-	1	1	2	3	6	5	4
2001	209	2	1	-	-	-	-	2	2	1	3	2	9	11
2000	246	1	2	-	-	1	-	2	-	6	6	6	9	9
1999	258	1	3	-	-	-	1	-	-	3	7	9	7	11
Male														
2002	101	1	-	-	-	1	-	1	1	-	3	5	4	3
2001	107	1	-	-	-	-	-	-	2	1	3	2	5	6
2000	136	1	2	-	-	1	-	2	-	3	5	4	8	8
1999	136	-	1	-	-	-	-	-	-	3	4	7	7	6
Female														
2002	95	1	1	-	1	-	-	-	-	2	-	1	1	1
2001	102	1	1	-	-	-	-	2	-	-	-	-	4	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)										
Non-Hispanic White										
Male										
2002	709	1,051	1,972	3,401	4,925	4,992	3,374	1,154	214	2
2001	645	1,012	1,901	3,262	4,550	4,732	3,112	1,085	210	1
2000	674	1,027	1,960	3,372	4,793	4,836	3,181	1,164	226	1
1999	592	1,055	1,973	3,385	4,678	4,738	3,090	1,106	243	-
Female										
2002	542	803	1,548	3,113	5,394	7,217	6,725	3,374	939	1
2001	477	753	1,530	2,911	5,067	7,055	6,313	3,144	860	-
2000	494	790	1,593	3,147	5,298	7,040	6,644	3,268	911	-
1999	469	737	1,577	3,160	5,145	6,908	6,612	3,349	948	1
Non-Hispanic Black										
Both sexes										
2002	297	323	508	707	873	848	646	309	106	-
2001	256	313	481	756	811	781	628	319	135	-
2000	290	403	565	684	853	798	645	307	121	-
1999	276	325	543	654	741	861	667	307	110	-
Male										
2002	178	185	266	373	414	317	207	92	18	-
2001	162	193	277	429	379	288	223	82	27	-
2000	166	242	315	356	402	314	217	83	25	-
1999	156	196	316	335	347	350	224	80	19	-
Female										
2002	119	138	242	334	459	531	439	217	88	-
2001	94	120	204	327	432	493	405	237	108	-
2000	124	161	250	328	451	484	428	224	96	-
1999	120	129	227	319	394	511	443	227	91	-
Origin not stated[2]										
Both sexes										
2002	3	10	14	27	42	30	29	11	3	-
2001	9	11	28	24	29	38	18	11	6	2
2000	5	17	17	34	45	33	32	17	2	2
1999	13	15	17	30	37	55	30	14	4	1
Male										
2002	2	9	12	12	23	7	11	4	2	-
2001	7	8	20	13	12	15	6	3	1	2
2000	4	11	12	19	25	16	9	5	-	1
1999	9	9	9	16	20	28	10	5	1	1
Female										
2002	1	1	2	15	19	23	18	7	1	-
2001	2	3	8	11	17	23	12	8	5	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonia (J12-J18)														
Origin not stated[2]														
Female														
2000	110	-	-	-	-	-	-	-	-	3	1	2	1	1
1999	122	1	2	-	-	-	1	-	-	-	3	2	-	5
Other acute lower respiratory infections (J20-J22)														
All origins														
Both sexes														
2002	386	52	23	7	1	1	2	3	2	2	6	6	12	11
2001	347	50	18	3	-	-	1	1	1	7	5	4	10	4
2000	425	43	11	3	4	4	1	5	3	3	9	8	4	6
1999	540	63	25	2	3	3	2	2	2	6	5	7	8	8
Male														
2002	154	39	9	7	-	-	-	1	-	1	2	1	9	3
2001	117	26	8	2	-	-	1	-	-	4	3	1	6	1
2000	160	25	4	2	1	4	-	5	1	1	6	2	2	2
1999	196	36	15	-	2	1	-	2	1	3	2	5	3	4
Female														
2002	232	13	14	-	1	1	2	2	2	1	4	5	3	8
2001	230	24	10	1	-	-	-	1	1	3	2	3	4	3
2000	265	18	7	1	3	-	1	-	2	2	3	6	2	4
1999	344	27	10	2	1	2	2	-	1	3	3	2	5	4
Hispanic														
Both sexes														
2002	21	10	2	1	1	-	-	1	-	-	-	-	-	1
2001	22	8	1	-	-	-	-	-	1	-	-	-	2	-
2000	24	9	2	-	-	-	-	1	-	-	-	1	-	-
1999	21	10	3	-	1	-	-	-	-	-	-	-	1	-
Male														
2002	9	5	1	1	-	-	-	-	-	-	-	-	-	1
2001	9	6	-	-	-	-	-	-	-	-	-	-	1	-
2000	15	6	1	-	-	-	-	1	-	-	-	1	-	-
1999	10	5	2	-	1	-	-	-	-	-	-	-	-	-
Female														
2002	12	5	1	-	1	-	-	1	-	-	-	-	-	-
2001	13	2	1	-	-	-	-	-	1	-	-	-	1	-
2000	9	3	1	-	-	-	-	-	-	-	-	-	-	-
1999	11	5	1	-	-	-	-	-	-	-	-	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)										
Origin not stated[2]										
Female										
2000	1	6	5	15	20	17	23	12	2	1
1999	4	6	8	14	17	27	20	9	3	-
Other acute lower respiratory infections (J20-J22)										
All origins										
Both sexes										
2002	7	10	23	21	45	70	42	27	13	-
2001	12	10	18	24	34	60	55	25	5	-
2000	4	19	24	41	41	80	77	23	12	-
1999	13	14	21	44	63	86	96	46	21	-
Male										
2002	3	4	9	11	16	21	11	5	2	-
2001	5	3	7	10	10	15	10	4	1	-
2000	1	12	10	21	14	26	15	6	-	-
1999	6	6	11	22	29	19	18	6	5	-
Female										
2002	4	6	14	10	29	49	31	22	11	-
2001	7	7	11	14	24	45	45	21	4	-
2000	3	7	14	20	27	54	62	17	12	-
1999	7	8	10	22	34	67	78	40	16	-
Hispanic										
Both sexes										
2002	1	-	-	-	-	2	1	1	-	-
2001	-	1	1	1	3	1	1	1	1	-
2000	-	-	1	1	1	2	5	1	-	-
1999	-	-	1	-	-	4	1	-	-	-
Male										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	1	-	1	-	-	-	-	-
2000	-	-	1	1	1	2	-	1	-	-
1999	-	-	1	-	-	1	-	-	-	-
Female										
2002	1	-	-	-	-	1	1	1	-	-
2001	-	1	-	1	2	1	1	1	1	-
2000	-	-	-	-	-	-	5	-	-	-
1999	-	-	-	-	-	3	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute lower respiratory infections (J20-J22)														
Non-Hispanic[1]														
Both sexes														
2002	365	42	21	6	-	1	2	2	2	2	6	6	12	10
2001	325	42	17	3	-	-	1	1	-	7	5	4	8	4
2000	398	33	9	3	4	4	1	4	3	3	9	7	4	6
1999	518	53	22	2	2	3	2	2	2	5	5	7	7	8
Male														
2002	145	34	8	6	-	-	-	1	-	1	2	1	9	2
2001	108	20	8	2	-	-	1	-	-	4	3	1	5	1
2000	144	19	3	2	1	4	-	4	1	1	6	1	2	2
1999	185	31	13	-	1	1	-	2	1	2	2	5	3	4
Female														
2002	220	8	13	-	-	1	2	1	2	1	4	5	3	8
2001	217	22	9	1	-	-	-	1	-	3	2	3	3	3
2000	254	14	6	1	3	-	1	-	2	2	3	6	2	4
1999	333	22	9	2	1	2	2	-	1	3	3	2	4	4
Non-Hispanic White														
Both sexes														
2002	314	25	18	6	-	1	1	1	2	2	5	4	10	8
2001	278	24	12	3	-	-	1	-	-	7	5	4	4	3
2000	345	15	6	2	1	4	1	3	1	2	6	4	4	6
1999	440	26	12	1	2	2	2	1	2	5	3	6	4	6
Male														
2002	122	22	6	6	-	-	-	-	-	1	2	1	8	2
2001	86	11	5	2	-	-	1	-	-	4	3	1	2	1
2000	118	6	3	2	-	4	-	3	-	1	5	1	2	2
1999	145	14	6	-	1	1	-	1	1	2	1	5	3	3
Female														
2002	192	3	12	-	-	1	1	1	2	1	3	3	2	6
2001	192	13	7	1	-	-	-	-	-	3	2	3	2	2
2000	227	9	3	-	1	-	1	-	1	1	1	3	2	4
1999	295	12	6	1	1	1	2	-	1	3	2	1	1	3
Non-Hispanic Black														
Both sexes														
2002	44	15	2	-	-	-	1	1	-	-	1	2	2	1
2001	33	14	2	-	-	-	-	1	-	-	-	-	4	1
2000	46	14	3	-	3	-	-	1	1	1	3	3	-	-
1999	64	24	6	-	-	1	-	1	-	-	2	1	3	1
Male														
2002	22	12	1	-	-	-	-	1	-	-	-	-	1	-
2001	15	8	1	-	-	-	-	-	-	-	-	-	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)										
Non-Hispanic[1]										
Both sexes										
2002	6	10	23	21	45	68	41	26	13	-
2001	12	9	17	23	31	59	54	24	4	-
2000	4	19	23	40	39	77	72	22	12	-
1999	13	14	20	44	63	82	95	46	21	-
Male										
2002	3	4	9	11	16	20	11	5	2	-
2001	5	3	6	10	9	15	10	4	1	-
2000	1	12	9	20	12	24	15	5	-	-
1999	6	6	10	22	29	18	18	6	5	-
Female										
2002	3	6	14	10	29	48	30	21	11	-
2001	7	6	11	13	22	44	44	20	3	-
2000	3	7	14	20	27	53	57	17	12	-
1999	7	8	10	22	34	64	77	40	16	-
Non-Hispanic White										
Both sexes										
2002	4	9	19	19	40	62	40	26	12	-
2001	11	8	12	20	30	57	51	22	4	-
2000	4	18	22	37	37	72	67	21	12	-
1999	11	11	15	41	58	77	92	44	19	-
Male										
2002	1	3	9	11	14	18	11	5	2	-
2001	4	3	4	8	9	14	9	4	1	-
2000	1	11	8	20	10	22	12	5	-	-
1999	6	4	7	21	27	15	17	6	4	-
Female										
2002	3	6	10	8	26	44	29	21	10	-
2001	7	5	8	12	21	43	42	18	3	-
2000	3	7	14	17	27	50	55	16	12	-
1999	5	7	8	20	31	62	75	38	15	-
Non-Hispanic Black										
Both sexes										
2002	2	1	4	2	5	4	-	-	1	-
2001	-	-	4	-	-	2	3	2	-	-
2000	-	1	1	3	2	5	4	1	-	-
1999	1	3	5	2	4	4	2	2	2	-
Male										
2002	2	1	-	-	2	2	-	-	-	-
2001	-	-	1	-	-	1	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other acute lower respiratory infections (J20-J22)														
Non-Hispanic Black														
Male														
2000	22	10	-	-	1	-	-	1	1	-	1	-	-	-
1999	30	15	3	-	-	-	-	1	-	-	1	-	-	1
Female														
2002	22	3	1	-	-	-	1	-	-	-	1	2	1	1
2001	18	6	1	-	-	-	-	1	-	-	-	-	1	1
2000	24	4	3	-	2	-	-	-	-	1	2	3	-	-
1999	34	9	3	-	-	1	-	-	-	-	1	1	3	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Acute bronchitis and bronchiolitis (J20-J21)														
All origins														
Both sexes														
2002	279	49	23	7	1	-	2	3	2	2	4	6	11	8
2001	263	49	18	3	-	-	1	1	1	6	5	4	8	3
2000	290	42	11	2	4	3	1	3	3	2	8	8	1	4
1999	388	62	24	2	2	3	2	1	2	6	4	7	7	6
Male														
2002	122	37	9	7	-	-	-	1	-	1	2	1	8	3
2001	89	25	8	2	-	-	1	-	-	3	3	1	5	1
2000	119	24	4	1	1	3	-	3	1	1	5	2	1	2
1999	154	36	14	-	2	1	-	1	1	3	1	5	2	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute lower respiratory infections (J20-J22)										
Non-Hispanic Black										
Male										
2000	-	1	1	-	2	2	2	-	-	-
1999	-	2	3	-	1	2	-	-	1	-
Female										
2002	-	-	4	2	3	2	-	-	1	-
2001	-	-	3	-	-	1	2	2	-	-
2000	-	-	-	3	-	3	2	1	-	-
1999	1	1	2	2	3	2	2	2	1	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Acute bronchitis and bronchiolitis (J20-J21)										
All origins										
Both sexes										
2002	5	10	13	15	24	39	31	15	9	-
2001	11	9	14	18	21	38	36	13	4	-
2000	3	15	16	30	23	45	44	15	7	-
1999	12	13	16	28	41	54	55	30	11	-
Male										
2002	2	4	6	8	10	12	7	3	1	-
2001	5	3	5	6	4	9	5	2	1	-
2000	1	9	5	18	9	14	10	5	-	-
1999	5	6	9	15	19	12	12	5	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute bronchitis and bronchiolitis (J20-J21)														
All origins														
Female														
2002	157	12	14	-	1	-	2	2	2	1	2	5	3	5
2001	174	24	10	1	-	-	-	1	1	3	2	3	3	2
2000	171	18	7	1	3	-	1	-	2	1	3	6	-	2
1999	234	26	10	2	-	2	2	-	1	3	3	2	5	3
Hispanic														
Both sexes														
2002	19	9	2	1	1	-	-	1	-	-	-	-	-	1
2001	17	8	1	-	-	-	-	-	1	-	-	-	2	-
2000	22	9	2	-	-	-	-	1	-	-	-	1	-	-
1999	20	10	3	-	1	-	-	-	-	-	-	-	1	-
Male														
2002	9	5	1	1	-	-	-	-	-	-	-	-	-	1
2001	7	6	-	-	-	-	-	-	-	-	-	-	1	-
2000	15	6	1	-	-	-	-	1	-	-	-	1	-	-
1999	10	5	2	-	1	-	-	-	-	-	-	-	-	-
Female														
2002	10	4	1	-	1	-	-	1	-	-	-	-	-	-
2001	10	2	1	-	-	-	-	-	1	-	-	-	1	-
2000	7	3	1	-	-	-	-	-	-	-	-	-	-	-
1999	10	5	1	-	-	-	-	-	-	-	-	-	1	-
Non-Hispanic[1]														
Both sexes														
2002	260	40	21	6	-	-	2	2	2	2	4	6	11	7
2001	246	41	17	3	-	-	1	1	-	6	5	4	6	3
2000	267	32	9	2	4	3	1	2	3	2	8	7	1	4
1999	367	52	21	2	1	3	2	1	2	5	4	7	6	6
Male														
2002	113	32	8	6	-	-	-	1	-	1	2	1	8	2
2001	82	19	8	2	-	-	1	-	-	3	3	1	4	1
2000	104	18	3	1	1	3	-	2	1	1	5	1	1	2
1999	143	31	12	-	1	1	-	1	1	2	1	5	2	3
Female														
2002	147	8	13	-	-	-	2	1	2	1	2	5	3	5
2001	164	22	9	1	-	-	-	1	-	3	2	3	2	2
2000	163	14	6	1	3	-	1	-	2	1	3	6	-	2
1999	224	21	9	2	-	2	2	-	1	3	3	2	4	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)										
All origins										
Female										
2002	3	6	7	7	14	27	24	12	8	-
2001	6	6	9	12	17	29	31	11	3	-
2000	2	6	11	12	14	31	34	10	7	-
1999	7	7	7	13	22	42	43	25	9	-
Hispanic										
Both sexes										
2002	1	-	-	-	-	2	1	-	-	-
2001	-	1	-	1	1	-	-	1	1	-
2000	-	-	1	1	1	2	3	1	-	-
1999	-	-	1	-	-	3	1	-	-	-
Male										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	1	1	1	2	-	1	-	-
1999	-	-	1	-	-	1	-	-	-	-
Female										
2002	1	-	-	-	-	1	1	-	-	-
2001	-	1	-	1	1	-	-	1	1	-
2000	-	-	-	-	-	-	3	-	-	-
1999	-	-	-	-	-	2	1	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	4	10	13	15	24	37	30	15	9	-
2001	11	8	14	17	20	38	36	12	3	-
2000	3	15	15	29	22	43	41	14	7	-
1999	12	13	15	28	41	51	54	30	11	-
Male										
2002	2	4	6	8	10	11	7	3	1	-
2001	5	3	5	6	4	9	5	2	1	-
2000	1	9	4	17	8	12	10	4	-	-
1999	5	6	8	15	19	11	12	5	2	-
Female										
2002	2	6	7	7	14	26	23	12	8	-
2001	6	5	9	11	16	29	31	10	2	-
2000	2	6	11	12	14	31	31	10	7	-
1999	7	7	7	13	22	40	42	25	9	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute bronchitis and bronchiolitis (J20-J21)														
Non-Hispanic White														
Both sexes														
2002	219	23	18	6	-	-	1	1	2	2	4	4	9	5
2001	202	23	12	3	-	-	1	-	-	6	5	4	2	2
2000	221	15	6	1	1	3	1	1	1	1	5	4	1	4
1999	299	25	12	1	1	2	2	1	2	5	2	6	3	4
Male														
2002	92	20	6	6	-	-	-	-	-	1	2	1	7	2
2001	61	10	5	2	-	-	1	-	-	3	3	1	1	1
2000	83	6	3	1	-	3	-	1	-	1	4	1	1	2
1999	108	14	6	-	1	1	-	1	1	2	-	5	2	2
Female														
2002	127	3	12	-	-	-	1	1	2	1	2	3	2	3
2001	141	13	7	1	-	-	-	-	-	3	2	3	1	1
2000	138	9	3	-	1	-	1	-	1	-	1	3	-	2
1999	191	11	6	1	-	1	2	-	1	3	2	1	1	2
Non-Hispanic Black														
Both sexes														
2002	35	15	2	-	-	-	1	1	-	-	-	2	2	1
2001	31	14	2	-	-	-	-	1	-	-	-	-	4	1
2000	40	13	3	-	3	-	-	1	1	1	3	3	-	-
1999	55	24	5	-	-	1	-	-	-	-	2	1	3	1
Male														
2002	20	12	1	-	-	-	-	1	-	-	-	-	1	-
2001	14	8	1	-	-	-	-	-	-	-	-	-	3	-
2000	18	9	-	-	1	-	-	1	1	-	1	-	-	-
1999	26	15	2	-	-	-	-	-	-	-	1	-	-	1
Female														
2002	15	3	1	-	-	-	1	-	-	-	-	2	1	1
2001	17	6	1	-	-	-	-	1	-	-	-	-	1	1
2000	22	4	3	-	2	-	-	-	-	1	2	3	-	-
1999	29	9	3	-	-	1	-	-	-	-	1	1	3	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)										
Non-Hispanic White										
Both sexes										
2002	3	9	12	14	22	32	29	15	8	-
2001	10	7	10	14	20	37	33	10	3	-
2000	3	14	14	26	21	40	39	13	7	-
1999	10	10	10	25	38	48	52	29	11	-
Male										
2002	1	3	6	8	9	9	7	3	1	-
2001	4	3	3	4	4	9	4	2	1	-
2000	1	8	3	17	7	12	8	4	-	-
1999	5	4	5	14	18	9	11	5	2	-
Female										
2002	2	6	6	6	13	23	22	12	7	-
2001	6	4	7	10	16	28	29	8	2	-
2000	2	6	11	9	14	28	31	9	7	-
1999	5	6	5	11	20	39	41	24	9	-
Non-Hispanic Black										
Both sexes										
2002	1	1	1	1	2	4	-	-	1	-
2001	-	-	3	-	-	1	3	2	-	-
2000	-	1	1	3	1	3	2	1	-	-
1999	1	3	5	2	2	3	1	1	-	-
Male										
2002	1	1	-	-	1	2	-	-	-	-
2001	-	-	1	-	-	-	1	-	-	-
2000	-	1	1	-	1	-	2	-	-	-
1999	-	2	3	-	-	2	-	-	-	-
Female										
2002	-	-	1	1	1	2	-	-	1	-
2001	-	-	2	-	-	1	2	2	-	-
2000	-	-	-	3	-	3	-	1	-	-
1999	1	1	2	2	2	1	1	1	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute bronchitis and bronchiolitis (J20-J21)														
Origin not stated[2]														
Male														
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	1	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unspecified acute lower respiratory infection (J22)														
All origins														
Both sexes														
2002	107	3	-	-	-	1	-	-	-	-	2	-	1	3
2001	84	1	-	-	-	-	-	-	-	1	-	-	2	1
2000	135	1	-	1	-	1	-	2	-	1	1	-	3	2
1999	152	1	1	-	1	-	-	1	-	-	1	-	1	2
Male														
2002	32	2	-	-	-	-	-	-	-	-	-	-	1	-
2001	28	1	-	-	-	-	-	-	-	1	-	-	1	-
2000	41	1	-	1	-	1	-	2	-	-	1	-	1	-
1999	42	-	1	-	-	-	-	1	-	-	1	-	1	1
Female														
2002	75	1	-	-	-	1	-	-	-	-	2	-	-	3
2001	56	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	94	-	-	-	-	-	-	-	-	1	-	-	2	2
1999	110	1	-	-	1	-	-	-	-	-	-	-	-	1
Hispanic														
Both sexes														
2002	2	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute bronchitis and bronchiolitis (J20-J21)										
Origin not stated[2]										
Male										
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Unspecified acute lower respiratory infection (J22)										
All origins										
Both sexes										
2002	2	-	10	6	21	31	11	12	4	-
2001	1	1	4	6	13	22	19	12	1	-
2000	1	4	8	11	18	35	33	8	5	-
1999	1	1	5	16	22	32	41	16	10	-
Male										
2002	1	-	3	3	6	9	4	2	1	-
2001	-	-	2	4	6	6	5	2	-	-
2000	-	3	5	3	5	12	5	1	-	-
1999	1	-	2	7	10	7	6	1	3	-
Female										
2002	1	-	7	3	15	22	7	10	3	-
2001	1	1	2	2	7	16	14	10	1	-
2000	1	1	3	8	13	23	28	7	5	-
1999	-	1	3	9	12	25	35	15	7	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	1	-	2	1	1	-	-	-
2000	-	-	-	-	-	-	2	-	-	-
1999	-	-	-	-	-	1	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	1	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Unspecified acute lower respiratory infection (J22)														
Hispanic														
Female														
2002	2	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	105	2	-	-	-	1	-	-	-	-	2	-	1	3
2001	79	1	-	-	-	-	-	-	-	1	-	-	2	1
2000	131	1	-	1	-	1	-	2	-	1	1	-	3	2
1999	151	1	1	-	1	-	-	1	-	-	1	-	1	2
Male														
2002	32	2	-	-	-	-	-	-	-	-	-	-	1	-
2001	26	1	-	-	-	-	-	-	-	1	-	-	1	-
2000	40	1	-	1	-	1	-	2	-	-	1	-	1	-
1999	42	-	1	-	-	-	-	1	-	-	1	-	1	1
Female														
2002	73	-	-	-	-	1	-	-	-	-	2	-	-	3
2001	53	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	91	-	-	-	-	-	-	-	-	1	-	-	2	2
1999	109	1	-	-	1	-	-	-	-	-	-	-	-	1
Non-Hispanic White														
Both sexes														
2002	95	2	-	-	-	1	-	-	-	-	1	-	1	3
2001	76	1	-	-	-	-	-	-	-	1	-	-	2	1
2000	124	-	-	1	-	1	-	2	-	1	1	-	3	2
1999	141	1	-	-	1	-	-	-	-	-	1	-	1	2
Male														
2002	30	2	-	-	-	-	-	-	-	-	-	-	1	-
2001	25	1	-	-	-	-	-	-	-	1	-	-	1	-
2000	35	-	-	1	-	1	-	2	-	-	1	-	1	-
1999	37	-	-	-	-	-	-	-	-	-	1	-	1	1
Female														
2002	65	-	-	-	-	1	-	-	-	-	1	-	-	3
2001	51	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	89	-	-	-	-	-	-	-	-	1	-	-	2	2
1999	104	1	-	-	1	-	-	-	-	-	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)										
Hispanic										
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	1	1	1	-	-	-
2000	-	-	-	-	-	-	2	-	-	-
1999	-	-	-	-	-	1	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	2	-	10	6	21	31	11	11	4	-
2001	1	1	3	6	11	21	18	12	1	-
2000	1	4	8	11	17	34	31	8	5	-
1999	1	1	5	16	22	31	41	16	10	-
Male										
2002	1	-	3	3	6	9	4	2	1	-
2001	-	-	1	4	5	6	5	2	-	-
2000	-	3	5	3	4	12	5	1	-	-
1999	1	-	2	7	10	7	6	1	3	-
Female										
2002	1	-	7	3	15	22	7	9	3	-
2001	1	1	2	2	6	15	13	10	1	-
2000	1	1	3	8	13	22	26	7	5	-
1999	-	1	3	9	12	24	35	15	7	-
Non-Hispanic White										
Both sexes										
2002	1	-	7	5	18	30	11	11	4	-
2001	1	1	2	6	10	20	18	12	1	-
2000	1	4	8	11	16	32	28	8	5	-
1999	1	1	5	16	20	29	40	15	8	-
Male										
2002	-	-	3	3	5	9	4	2	1	-
2001	-	-	1	4	5	5	5	2	-	-
2000	-	3	5	3	3	10	4	1	-	-
1999	1	-	2	7	9	6	6	1	2	-
Female										
2002	1	-	4	2	13	21	7	9	3	-
2001	1	1	1	2	5	15	13	10	1	-
2000	1	1	3	8	13	22	24	7	5	-
1999	-	1	3	9	11	23	34	14	6	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Unspecified acute lower respiratory infection (J22)														
Non-Hispanic Black														
Both sexes														
2002	9	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	6	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	9	-	1	-	-	-	-	1	-	-	-	-	-	-
Male														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	4	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	1	-	-	-	-	1	-	-	-	-	-	-
Female														
2002	7	-	-	-	-	-	-	-	-	-	1	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Chronic lower respiratory diseases (J40-J47)														
All origins														
Both sexes														
2002	124,816	39	65	41	95	93	99	116	185	340	668	1,226	2,249	4,220
2001	123,013	41	43	42	62	74	97	102	188	355	614	1,127	2,197	4,176
2000	122,009	36	51	48	91	86	104	125	171	310	620	1,174	2,077	3,985
1999	124,181	36	54	49	90	107	102	145	185	302	588	1,079	2,031	4,179

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Unspecified acute lower respiratory infection (J22)										
Non-Hispanic Black										
Both sexes										
2002	1	-	3	1	3	-	-	-	-	-
2001	-	-	1	-	-	1	-	-	-	-
2000	-	-	-	-	1	2	2	-	-	-
1999	-	-	-	-	2	1	1	1	2	-
Male										
2002	1	-	-	-	1	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	-
2000	-	-	-	-	1	2	-	-	-	-
1999	-	-	-	-	1	-	-	-	1	-
Female										
2002	-	-	3	1	2	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	2	-	-	-
1999	-	-	-	-	1	1	1	1	1	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	1	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Chronic lower respiratory diseases (J40-J47)										
All origins										
Both sexes										
2002	7,060	11,405	18,383	24,495	24,746	17,971	8,508	2,430	375	7
2001	6,990	11,838	18,913	24,442	23,320	17,542	8,125	2,345	379	1
2000	6,754	11,833	19,324	24,309	23,413	17,085	7,823	2,238	350	2
1999	7,118	12,441	20,203	24,978	23,657	16,831	7,510	2,154	338	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic lower respiratory diseases (J40-J47)														
All origins														
Male														
2002	60,713	23	44	24	50	68	52	70	95	159	338	617	1,178	2,141
2001	59,697	22	29	24	37	43	56	55	91	171	284	591	1,111	2,159
2000	60,004	23	28	26	54	58	62	72	93	145	292	617	1,044	2,072
1999	62,415	19	30	29	64	60	63	69	83	142	268	547	1,004	2,169
Female														
2002	64,103	16	21	17	45	25	47	46	90	181	330	609	1,071	2,079
2001	63,316	19	14	18	25	31	41	47	97	184	330	536	1,086	2,017
2000	62,005	13	23	22	37	28	42	53	78	165	328	557	1,033	1,913
1999	61,766	17	24	20	26	47	39	76	102	160	320	532	1,027	2,010
Hispanic														
Both sexes														
2002	3,058	4	9	6	7	6	9	9	22	25	45	50	77	102
2001	2,832	7	8	2	7	7	3	10	12	25	27	47	77	100
2000	2,689	7	9	5	10	6	10	7	13	21	34	45	60	87
1999	2,859	6	8	7	11	8	7	17	14	34	45	52	63	95
Male														
2002	1,625	2	4	4	4	5	7	8	11	12	19	28	42	64
2001	1,482	4	4	1	5	4	2	6	7	12	9	22	32	53
2000	1,451	4	5	2	7	5	8	4	8	11	13	21	32	40
1999	1,547	4	3	3	8	4	5	6	7	14	22	21	31	50
Female														
2002	1,433	2	5	2	3	1	2	1	11	13	26	22	35	38
2001	1,350	3	4	1	2	3	1	4	5	13	18	25	45	47
2000	1,238	3	4	3	3	1	2	3	5	10	21	24	28	47
1999	1,312	2	5	4	3	4	2	11	7	20	23	31	32	45
Non-Hispanic[1]														
Both sexes														
2002	121,424	34	56	35	86	86	90	107	160	314	616	1,170	2,165	4,104
2001	119,839	34	35	40	54	67	94	92	175	328	581	1,073	2,107	4,055
2000	118,963	29	42	42	81	80	93	115	157	288	584	1,120	2,002	3,882
1999	120,968	30	46	42	79	99	95	128	168	267	540	1,023	1,955	4,075
Male														
2002	58,896	20	40	20	44	62	45	62	81	146	314	585	1,134	2,072
2001	58,028	18	25	23	32	39	54	49	84	158	270	565	1,070	2,092
2000	58,349	19	23	24	47	53	54	65	85	133	278	589	1,004	2,022
1999	60,669	15	27	26	56	56	58	63	75	127	245	523	963	2,113
Female														
2002	62,528	14	16	15	42	24	45	45	79	168	302	585	1,031	2,032

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)										
All origins										
Male										
2002	3,642	5,838	9,501	12,457	11,975	8,140	3,448	762	86	5
2001	3,543	6,046	9,831	12,433	11,258	7,806	3,279	733	94	1
2000	3,492	6,149	10,061	12,305	11,533	7,843	3,165	776	92	2
1999	3,789	6,642	10,835	13,021	11,684	7,853	3,154	779	108	3
Female										
2002	3,418	5,567	8,882	12,038	12,771	9,831	5,060	1,668	289	2
2001	3,447	5,792	9,082	12,009	12,062	9,736	4,846	1,612	285	-
2000	3,262	5,684	9,263	12,004	11,880	9,242	4,658	1,462	258	-
1999	3,329	5,799	9,368	11,957	11,973	8,978	4,356	1,375	230	1
Hispanic										
Both sexes										
2002	172	252	383	510	545	439	280	89	17	-
2001	158	265	369	513	427	399	271	80	18	-
2000	157	240	332	468	453	400	217	94	14	-
1999	144	262	387	483	452	415	249	91	9	-
Male										
2002	100	152	210	312	281	201	122	34	3	-
2001	87	161	217	286	215	203	112	34	6	-
2000	106	136	207	260	249	190	95	45	3	-
1999	85	136	237	278	245	216	119	48	5	-
Female										
2002	72	100	173	198	264	238	158	55	14	-
2001	71	104	152	227	212	196	159	46	12	-
2000	51	104	125	208	204	210	122	49	11	-
1999	59	126	150	205	207	199	130	43	4	-
Non-Hispanic[1]										
Both sexes										
2002	6,871	11,118	17,951	23,913	24,145	17,491	8,212	2,338	356	6
2001	6,805	11,545	18,483	23,869	22,844	17,108	7,830	2,260	360	-
2000	6,565	11,550	18,936	23,777	22,910	16,652	7,586	2,136	335	1
1999	6,943	12,145	19,757	24,425	23,153	16,368	7,239	2,059	329	3
Male										
2002	3,532	5,665	9,259	12,098	11,663	7,918	3,321	728	83	4
2001	3,438	5,869	9,580	12,120	11,014	7,588	3,156	696	88	-
2000	3,366	5,985	9,820	12,009	11,257	7,637	3,061	729	88	1
1999	3,684	6,482	10,558	12,709	11,415	7,616	3,024	729	103	2
Female										
2002	3,339	5,453	8,692	11,815	12,482	9,573	4,891	1,610	273	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic lower respiratory diseases (J40-J47)														
Non-Hispanic[1]														
Female														
2001	61,811	16	10	17	22	28	40	43	91	170	311	508	1,037	1,963
2000	60,614	10	19	18	34	27	39	50	72	155	306	531	998	1,860
1999	60,299	15	19	16	23	43	37	65	93	140	295	500	992	1,962
Non-Hispanic White														
Both sexes														
2002	112,128	20	23	11	31	40	54	59	98	201	423	917	1,769	3,600
2001	110,753	16	13	16	23	33	42	43	91	187	426	831	1,739	3,548
2000	109,901	16	19	17	30	34	53	59	85	174	396	830	1,633	3,369
1999	111,620	10	23	23	32	60	50	57	89	167	366	747	1,625	3,555
Male														
2002	53,662	13	17	8	15	25	23	29	52	84	220	459	924	1,793
2001	52,952	10	9	9	13	15	24	23	39	97	196	456	875	1,823
2000	53,231	12	8	10	14	21	31	31	40	74	193	445	814	1,744
1999	55,314	6	11	13	23	32	27	26	40	77	162	386	788	1,830
Female														
2002	58,466	7	6	3	16	15	31	30	46	117	203	458	845	1,807
2001	57,801	6	4	7	10	18	18	20	52	90	230	375	864	1,725
2000	56,670	4	11	7	16	13	22	28	45	100	203	385	819	1,625
1999	56,306	4	12	10	9	28	23	31	49	90	204	361	837	1,725
Non-Hispanic Black														
Both sexes														
2002	7,730	14	29	21	53	43	35	43	59	109	183	230	359	454
2001	7,504	16	22	23	30	30	47	47	81	131	142	222	327	446
2000	7,528	11	21	24	46	44	36	49	65	102	170	258	335	443
1999	7,840	18	18	19	47	36	42	67	73	93	165	256	300	472
Male														
2002	4,282	7	20	11	28	36	22	28	28	60	90	115	189	250
2001	4,143	6	16	14	18	22	26	26	43	58	66	101	170	235
2000	4,184	6	14	14	29	31	20	29	41	51	78	125	164	238
1999	4,452	9	13	13	33	23	29	34	33	44	77	126	158	254
Female														
2002	3,448	7	9	10	25	7	13	15	31	49	93	115	170	204
2001	3,361	10	6	9	12	8	21	21	38	73	76	121	157	211
2000	3,344	5	7	10	17	13	16	20	24	51	92	133	171	205
1999	3,388	9	5	6	14	13	13	33	40	49	88	130	142	218
Origin not stated[2]														
Both sexes														
2002	334	1	-	-	2	1	-	-	3	1	7	6	7	14

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)										
Non-Hispanic[1]										
Female										
2001	3,367	5,676	8,903	11,749	11,830	9,520	4,674	1,564	272	-
2000	3,199	5,565	9,116	11,768	11,653	9,015	4,525	1,407	247	-
1999	3,259	5,663	9,199	11,716	11,738	8,752	4,215	1,330	226	1
Non-Hispanic White										
Both sexes										
2002	6,157	10,170	16,638	22,378	22,739	16,549	7,760	2,169	318	4
2001	6,130	10,528	17,165	22,403	21,569	16,195	7,328	2,108	319	-
2000	5,884	10,523	17,599	22,319	21,705	15,765	7,125	1,970	295	1
1999	6,231	11,099	18,301	22,770	21,892	15,486	6,810	1,928	297	2
Male										
2002	3,116	5,124	8,491	11,168	10,866	7,394	3,109	659	70	3
2001	3,062	5,295	8,794	11,220	10,276	7,101	2,916	629	70	-
2000	2,984	5,367	9,031	11,134	10,569	7,141	2,836	657	74	1
1999	3,261	5,865	9,694	11,705	10,672	7,142	2,795	666	92	1
Female										
2002	3,041	5,046	8,147	11,210	11,873	9,155	4,651	1,510	248	1
2001	3,068	5,233	8,371	11,183	11,293	9,094	4,412	1,479	249	-
2000	2,900	5,156	8,568	11,185	11,136	8,624	4,289	1,313	221	-
1999	2,970	5,234	8,607	11,065	11,220	8,344	4,015	1,262	205	1
Non-Hispanic Black										
Both sexes										
2002	613	802	1,090	1,240	1,126	710	347	138	31	1
2001	573	867	1,095	1,178	1,012	697	372	117	29	-
2000	620	872	1,121	1,176	952	663	358	131	31	-
1999	613	899	1,191	1,351	1,013	694	337	110	25	1
Male										
2002	357	461	632	735	615	378	155	54	10	1
2001	319	496	651	703	582	362	166	51	12	-
2000	349	526	669	681	526	356	171	58	8	-
1999	370	533	705	810	592	362	171	52	10	1
Female										
2002	256	341	458	505	511	332	192	84	21	-
2001	254	371	444	475	430	335	206	66	17	-
2000	271	346	452	495	426	307	187	73	23	-
1999	243	366	486	541	421	332	166	58	15	-
Origin not stated[2]										
Both sexes										
2002	17	35	49	72	56	41	16	3	2	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic lower respiratory diseases (J40-J47)														
Origin not stated[2]														
Both sexes														
2001	342	-	-	-	1	-	-	-	1	2	6	7	13	21
2000	357	-	-	1	-	-	1	3	1	1	2	9	15	16
1999	354	-	-	-	-	-	-	-	3	1	3	4	13	9
Male														
2002	192	1	-	-	2	1	-	-	3	1	5	4	2	5
2001	187	-	-	-	-	-	-	-	-	1	5	4	9	14
2000	204	-	-	-	-	-	-	3	-	1	1	7	8	10
1999	199	-	-	-	-	-	-	-	1	1	1	3	10	6
Female														
2002	142	-	-	-	-	-	-	-	-	-	2	2	5	9
2001	155	-	-	-	1	-	-	-	1	1	1	3	4	7
2000	153	-	-	1	-	-	1	-	1	-	1	2	7	6
1999	155	-	-	-	-	-	-	-	2	-	2	1	3	3
Bronchitis, chronic and unspecified (J40-J42)														
All origins														
Both sexes														
2002	955	24	13	3	2	3	3	2	4	12	6	16	20	22
2001	959	21	8	-	-	3	2	4	2	9	12	21	21	25
2000	1,167	18	14	2	3	3	2	3	5	7	13	19	19	20
1999	1,172	20	19	3	1	3	4	3	8	10	8	14	24	35
Male														
2002	420	15	8	2	-	2	1	-	2	9	3	13	14	10
2001	396	11	5	-	-	2	1	2	-	3	5	12	9	16
2000	503	9	7	2	1	2	2	3	4	4	10	8	8	9
1999	498	9	11	2	1	2	2	-	3	3	2	9	10	24
Female														
2002	535	9	5	1	2	1	2	2	2	3	3	3	6	12
2001	563	10	3	-	-	1	1	2	2	6	7	9	12	9
2000	664	9	7	-	2	1	-	-	1	3	3	11	11	11
1999	674	11	8	1	-	1	2	3	5	7	6	5	14	11
Hispanic														
Both sexes														
2002	51	3	1	-	-	-	2	-	1	2	1	1	2	-
2001	37	4	1	-	-	1	-	1	-	1	1	1	-	2
2000	52	4	3	-	-	-	-	1	1	1	2	1	-	1
1999	52	3	3	-	1	-	-	2	-	4	3	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic lower respiratory diseases (J40-J47)										
Origin not stated[2]										
Both sexes										
2001	27	28	61	60	49	35	24	5	1	1
2000	32	43	56	64	50	33	20	8	1	1
1999	31	34	59	70	52	48	22	4	-	1
Male										
2002	10	21	32	47	31	21	5	-	-	1
2001	18	16	34	27	29	15	11	3	-	1
2000	20	28	34	36	27	16	9	2	1	1
1999	20	24	40	34	24	21	11	2	-	1
Female										
2002	7	14	17	25	25	20	11	3	2	-
2001	9	12	27	33	20	20	13	2	1	-
2000	12	15	22	28	23	17	11	6	-	-
1999	11	10	19	36	28	27	11	2	-	-
Bronchitis, chronic and unspecified (J40-J42)										
All origins										
Both sexes										
2002	32	51	84	131	155	162	114	69	27	-
2001	36	56	77	129	132	188	123	67	23	-
2000	47	67	91	156	177	236	161	80	24	-
1999	62	63	110	139	200	193	148	82	23	-
Male										
2002	18	32	42	69	58	62	35	20	5	-
2001	20	35	41	67	48	70	33	14	2	-
2000	23	41	38	84	90	92	46	19	1	-
1999	27	36	58	73	84	71	45	20	6	-
Female										
2002	14	19	42	62	97	100	79	49	22	-
2001	16	21	36	62	84	118	90	53	21	-
2000	24	26	53	72	87	144	115	61	23	-
1999	35	27	52	66	116	122	103	62	17	-
Hispanic										
Both sexes										
2002	2	2	-	7	7	9	8	3	-	-
2001	-	1	2	8	1	4	5	1	3	-
2000	-	2	7	4	9	6	4	5	1	-
1999	2	2	2	4	9	5	10	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Bronchitis, chronic and unspecified (J40-J42)														
Hispanic														
Male														
2002	20	1	1	-	-	-	1	-	1	1	1	1	2	-
2001	15	2	-	-	-	-	-	1	-	-	1	-	-	1
2000	28	2	2	-	-	-	-	1	1	1	1	-	-	1
1999	17	1	2	-	1	-	-	-	-	1	1	-	-	-
Female														
2002	31	2	-	-	-	-	1	-	-	1	-	-	-	-
2001	22	2	1	-	-	1	-	-	-	1	-	1	-	1
2000	24	2	1	-	-	-	-	-	-	-	1	1	-	-
1999	35	2	1	-	-	-	-	2	-	3	2	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	900	20	12	3	2	3	1	2	3	10	5	15	18	22
2001	919	17	7	-	-	2	2	3	2	8	11	19	21	22
2000	1,112	14	11	2	3	3	2	2	4	6	11	18	19	19
1999	1,117	17	16	3	-	3	4	1	8	6	5	14	24	34
Male														
2002	398	13	7	2	-	2	-	-	1	8	2	12	12	10
2001	380	9	5	-	-	2	1	1	-	3	4	12	9	14
2000	474	7	5	2	1	2	2	2	3	3	9	8	8	8
1999	478	8	9	2	-	2	2	-	3	2	1	9	10	23
Female														
2002	502	7	5	1	2	1	1	2	2	2	3	3	6	12
2001	539	8	2	-	-	-	1	2	2	5	7	7	12	8
2000	638	7	6	-	2	1	-	-	1	3	2	10	11	11
1999	639	9	7	1	-	1	2	1	5	4	4	5	14	11
Non-Hispanic White														
Both sexes														
2002	821	12	6	2	1	1	1	1	3	9	4	11	15	18
2001	839	7	5	-	-	1	2	2	2	8	8	15	20	21
2000	1,018	5	4	2	2	2	2	2	2	6	5	16	15	16
1999	1,002	8	12	3	-	3	3	-	6	6	3	9	15	31
Male														
2002	354	9	2	2	-	1	-	-	1	7	1	8	9	8
2001	338	5	5	-	-	1	1	-	-	3	3	9	8	14
2000	429	3	1	2	-	1	2	2	1	3	3	8	7	7
1999	421	5	6	2	-	2	1	-	2	2	1	5	6	21
Female														
2002	467	3	4	-	1	-	1	1	2	2	3	3	6	10

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)										
Hispanic										
Male										
2002	1	1	-	1	3	3	1	1	-	-
2001	-	-	1	6	-	1	2	-	-	-
2000	-	1	6	1	5	3	1	2	-	-
1999	-	2	1	1	2	3	1	-	1	-
Female										
2002	1	1	-	6	4	6	7	2	-	-
2001	-	1	1	2	1	3	3	1	3	-
2000	-	1	1	3	4	3	3	3	1	-
1999	2	-	1	3	7	2	9	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	30	48	84	123	148	153	106	66	26	-
2001	36	55	75	121	130	184	118	66	20	-
2000	47	65	84	152	168	230	157	72	23	-
1999	60	60	108	135	191	187	138	81	22	-
Male										
2002	17	31	42	67	55	59	34	19	5	-
2001	20	35	40	61	48	69	31	14	2	-
2000	23	40	32	83	85	89	45	16	1	-
1999	27	33	57	72	82	67	44	20	5	-
Female										
2002	13	17	42	56	93	94	72	47	21	-
2001	16	20	35	60	82	115	87	52	18	-
2000	24	25	52	69	83	141	112	56	22	-
1999	33	27	51	63	109	120	94	61	17	-
Non-Hispanic White										
Both sexes										
2002	26	43	81	117	135	147	100	63	25	-
2001	33	48	65	110	122	178	111	62	19	-
2000	44	56	75	140	157	223	153	69	22	-
1999	55	55	95	119	174	175	133	78	19	-
Male										
2002	14	27	40	63	50	58	31	19	4	-
2001	18	31	35	54	43	66	29	11	2	-
2000	20	36	28	77	82	85	45	15	1	-
1999	24	31	50	64	71	63	42	19	4	-
Female										
2002	12	16	41	54	85	89	69	44	21	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Bronchitis, chronic and unspecified (J40-J42)														
Non-Hispanic White														
Female														
2001	501	2	-	-	-	-	1	2	2	5	5	6	12	7
2000	589	2	3	-	2	1	-	-	1	3	2	8	8	9
1999	581	3	6	1	-	1	2	-	4	4	2	4	9	10
Non-Hispanic Black														
Both sexes														
2002	66	8	6	1	1	2	-	1	-	1	1	4	3	3
2001	62	10	2	-	-	1	-	1	-	-	3	3	1	1
2000	80	7	6	-	1	1	-	-	1	-	6	2	4	1
1999	88	7	1	-	-	-	1	1	2	-	2	4	8	3
Male														
2002	35	4	5	-	-	1	-	-	-	1	1	4	3	1
2001	32	4	-	-	-	1	-	1	-	-	1	3	1	-
2000	35	3	3	-	1	1	-	-	1	-	6	-	1	-
1999	43	3	1	-	-	-	1	-	1	-	-	3	4	2
Female														
2002	31	4	1	1	1	1	-	1	-	-	-	-	-	2
2001	30	6	2	-	-	-	-	-	-	-	2	-	-	1
2000	45	4	3	-	-	-	-	-	-	-	-	2	3	1
1999	45	4	-	-	-	-	-	1	1	-	2	1	4	1
Origin not stated[2]														
Both sexes														
2002	4	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	1	-	1
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	2	1	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Bronchitis, chronic and unspecified (J40-J42)										
Non-Hispanic White										
Female										
2001	15	17	30	56	79	112	82	51	17	-
2000	24	20	47	63	75	138	108	54	21	-
1999	31	24	45	55	103	112	91	59	15	-
Non-Hispanic Black										
Both sexes										
2002	2	4	2	4	9	6	5	2	1	-
2001	3	7	8	8	3	5	3	3	-	-
2000	3	7	9	10	10	5	3	3	1	-
1999	4	4	11	12	14	8	3	1	2	-
Male										
2002	1	3	1	3	2	1	3	-	1	-
2001	2	4	4	4	2	2	-	3	-	-
2000	3	2	4	4	3	2	-	1	-	-
1999	2	1	6	6	10	1	1	-	1	-
Female										
2002	1	1	1	1	7	5	2	2	-	-
2001	1	3	4	4	1	3	3	-	-	-
2000	-	5	5	6	7	3	3	2	1	-
1999	2	3	5	6	4	7	2	1	1	-
Origin not stated[2]										
Both sexes										
2002	-	1	-	1	-	-	-	-	1	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	3	-	-
1999	-	1	-	-	-	1	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	1	-	-
1999	-	1	-	-	-	1	-	-	-	-
Female										
2002	-	1	-	-	-	-	-	-	1	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	2	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Emphysema (J43)														
All origins														
Both sexes														
2002	15,489	1	3	-	-	1	1	2	8	21	73	156	298	643
2001	16,242	3	-	-	1	1	1	1	6	24	51	141	307	705
2000	16,700	3	2	-	-	1	4	6	5	23	59	162	312	680
1999	17,787	1	-	1	-	-	1	1	8	19	78	136	332	723
Male														
2002	8,084	-	3	-	-	-	-	1	7	14	52	95	172	334
2001	8,415	1	-	-	1	1	-	-	4	17	35	89	177	395
2000	8,671	3	-	-	-	1	4	1	4	15	31	101	179	366
1999	9,480	-	-	-	-	-	1	-	6	9	40	85	183	394
Female														
2002	7,405	1	-	-	-	1	1	1	1	7	21	61	126	309
2001	7,827	2	-	-	-	-	1	1	2	7	16	52	130	310
2000	8,029	-	2	-	-	-	-	5	1	8	28	61	133	314
1999	8,307	1	-	1	-	-	-	1	2	10	38	51	149	329
Hispanic														
Both sexes														
2002	295	-	-	-	-	-	-	-	2	-	1	1	4	9
2001	286	1	-	-	-	-	-	-	-	-	-	5	3	10
2000	266	-	-	-	-	-	1	-	-	1	2	3	8	8
1999	292	-	-	1	-	-	-	-	2	2	4	5	7	6
Male														
2002	168	-	-	-	-	-	-	-	1	-	1	1	3	5
2001	175	-	-	-	-	-	-	-	-	-	-	2	3	7
2000	167	-	-	-	-	-	1	-	-	1	2	2	6	2
1999	183	-	-	-	-	-	-	-	1	-	2	4	4	3
Female														
2002	127	-	-	-	-	-	-	-	1	-	-	-	1	4
2001	111	1	-	-	-	-	-	-	-	-	-	3	-	3
2000	99	-	-	-	-	-	-	-	-	-	-	1	2	6
1999	109	-	-	1	-	-	-	-	1	2	2	1	3	3
Non-Hispanic[1]														
Both sexes														
2002	15,161	1	3	-	-	1	1	2	6	21	72	154	294	633
2001	15,914	2	-	-	1	1	1	1	6	24	50	135	304	691
2000	16,382	3	2	-	-	1	3	6	5	22	57	155	302	670
1999	17,441	1	-	-	-	-	1	1	6	16	72	130	320	716
Male														
2002	7,895	-	3	-	-	-	-	1	6	14	51	93	169	328
2001	8,213	1	-	-	1	1	-	-	4	17	34	86	174	384
2000	8,483	3	-	-	-	1	3	1	4	14	29	96	172	363

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)										
All origins										
Both sexes										
2002	1,060	1,665	2,723	3,202	2,907	1,826	726	157	15	1
2001	1,124	1,892	2,912	3,481	2,856	1,827	727	161	21	-
2000	1,127	1,976	3,144	3,522	2,989	1,832	685	150	18	-
1999	1,296	2,246	3,410	3,768	3,187	1,785	622	146	27	-
Male										
2002	591	875	1,460	1,723	1,505	866	329	52	4	1
2001	571	988	1,566	1,809	1,446	929	326	54	6	-
2000	577	1,077	1,661	1,839	1,534	898	317	58	5	-
1999	691	1,230	1,906	2,027	1,650	886	295	68	9	-
Female										
2002	469	790	1,263	1,479	1,402	960	397	105	11	-
2001	553	904	1,346	1,672	1,410	898	401	107	15	-
2000	550	899	1,483	1,683	1,455	934	368	92	13	-
1999	605	1,016	1,504	1,741	1,537	899	327	78	18	-
Hispanic										
Both sexes										
2002	20	25	54	41	72	34	25	6	1	-
2001	16	33	42	61	52	32	22	6	3	-
2000	18	26	39	60	53	29	13	5	-	-
1999	15	28	46	56	57	37	18	7	1	-
Male										
2002	12	20	28	27	38	18	11	3	-	-
2001	9	19	27	41	35	21	7	2	2	-
2000	15	17	26	35	31	18	8	3	-	-
1999	13	18	33	34	35	21	9	6	-	-
Female										
2002	8	5	26	14	34	16	14	3	1	-
2001	7	14	15	20	17	11	15	4	1	-
2000	3	9	13	25	22	11	5	2	-	-
1999	2	10	13	22	22	16	9	1	1	-
Non-Hispanic[1]										
Both sexes										
2002	1,039	1,637	2,661	3,155	2,831	1,784	701	151	14	-
2001	1,103	1,855	2,863	3,412	2,798	1,794	701	154	18	-
2000	1,107	1,945	3,091	3,450	2,932	1,799	669	145	18	-
1999	1,276	2,213	3,358	3,701	3,124	1,740	601	139	26	-
Male										
2002	579	853	1,428	1,690	1,466	843	318	49	4	-
2001	559	966	1,534	1,764	1,408	907	318	51	4	-
2000	562	1,058	1,627	1,800	1,503	879	308	55	5	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Emphysema (J43)														
Non-Hispanic[1]														
Male														
1999	9,260	-	-	-	-	-	1	-	5	8	37	80	175	390
Female														
2002	7,266	1	-	-	-	1	1	1	-	7	21	61	125	305
2001	7,701	1	-	-	-	-	1	1	2	7	16	49	130	307
2000	7,899	-	2	-	-	-	-	5	1	8	28	59	130	307
1999	8,181	1	-	-	-	-	-	1	1	8	35	50	145	326
Non-Hispanic White														
Both sexes														
2002	14,157	1	1	-	-	1	-	2	4	17	49	127	258	576
2001	14,930	1	-	-	-	1	-	1	3	22	42	121	253	612
2000	15,301	3	-	-	-	-	1	6	4	17	43	115	252	599
1999	16,338	-	-	-	-	-	-	1	4	13	51	98	274	648
Male														
2002	7,241	-	1	-	-	-	-	1	4	11	36	76	143	293
2001	7,607	-	-	-	-	1	-	-	2	16	28	79	143	335
2000	7,805	3	-	-	-	-	1	1	3	10	21	67	139	322
1999	8,540	-	-	-	-	-	-	-	3	5	28	59	140	344
Female														
2002	6,916	1	-	-	-	1	-	1	-	6	13	51	115	283
2001	7,323	1	-	-	-	-	-	1	1	6	14	42	110	277
2000	7,496	-	-	-	-	-	-	5	1	7	22	48	113	277
1999	7,798	-	-	-	-	-	-	1	1	8	23	39	134	304
Non-Hispanic Black														
Both sexes														
2002	842	-	1	-	-	-	1	-	2	4	22	26	30	49
2001	800	1	-	-	1	-	1	-	3	1	8	13	46	69
2000	889	-	1	-	-	1	2	-	1	4	13	37	47	63
1999	923	1	-	-	-	-	-	-	-	2	20	32	43	60
Male														
2002	538	-	1	-	-	-	-	-	2	3	15	16	22	30
2001	480	1	-	-	1	-	-	-	2	1	6	6	27	43
2000	551	-	-	-	-	1	2	-	1	3	8	26	30	36
1999	594	-	-	-	-	-	-	-	-	2	9	21	33	38
Female														
2002	304	-	-	-	-	-	1	-	-	1	7	10	8	19
2001	320	-	-	-	-	-	1	-	1	-	2	7	19	26
2000	338	-	1	-	-	-	-	-	-	1	5	11	17	27
1999	329	1	-	-	-	-	-	-	-	-	11	11	10	22

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)										
Non-Hispanic[1]										
Male										
1999	673	1,208	1,868	1,986	1,612	862	284	62	9	-
Female										
2002	460	784	1,233	1,465	1,365	941	383	102	10	-
2001	544	889	1,329	1,648	1,390	887	383	103	14	-
2000	545	887	1,464	1,650	1,429	920	361	90	13	-
1999	603	1,005	1,490	1,715	1,512	878	317	77	17	-
Non-Hispanic White										
Both sexes										
2002	953	1,509	2,483	2,987	2,676	1,691	664	147	11	-
2001	1,021	1,736	2,678	3,238	2,675	1,711	655	143	17	-
2000	1,017	1,788	2,916	3,273	2,795	1,699	628	129	16	-
1999	1,177	2,073	3,165	3,482	2,987	1,647	569	128	21	-
Male										
2002	517	759	1,322	1,574	1,372	781	302	47	2	-
2001	506	900	1,410	1,656	1,330	852	297	48	4	-
2000	508	947	1,523	1,698	1,423	808	284	44	3	-
1999	609	1,115	1,728	1,855	1,516	812	262	57	7	-
Female										
2002	436	750	1,161	1,413	1,304	910	362	100	9	-
2001	515	836	1,268	1,582	1,345	859	358	95	13	-
2000	509	841	1,393	1,575	1,372	891	344	85	13	-
1999	568	958	1,437	1,627	1,471	835	307	71	14	-
Non-Hispanic Black										
Both sexes										
2002	68	111	156	135	131	67	32	4	3	-
2001	69	106	148	123	96	65	40	9	1	-
2000	83	135	141	135	108	73	31	14	-	-
1999	86	126	153	175	108	76	27	11	3	-
Male										
2002	47	81	92	90	79	42	14	2	2	-
2001	44	61	94	73	61	40	18	2	-	-
2000	51	97	86	72	62	47	20	9	-	-
1999	56	87	108	103	75	38	17	5	2	-
Female										
2002	21	30	64	45	52	25	18	2	1	-
2001	25	45	54	50	35	25	22	7	1	-
2000	32	38	55	63	46	26	11	5	-	-
1999	30	39	45	72	33	38	10	6	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Emphysema (J43)														
Origin not stated[2]														
Both sexes														
2002	33	-	-	-	-	-	-	-	-	-	-	1	-	1
2001	42	-	-	-	-	-	-	-	-	-	1	1	-	4
2000	52	-	-	-	-	-	-	-	-	-	-	4	2	2
1999	54	-	-	-	-	-	-	-	-	1	2	1	5	1
Male														
2002	21	-	-	-	-	-	-	-	-	-	-	1	-	1
2001	27	-	-	-	-	-	-	-	-	-	1	1	-	4
2000	21	-	-	-	-	-	-	-	-	-	-	3	1	1
1999	37	-	-	-	-	-	-	-	-	1	1	1	4	1
Female														
2002	12	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	15	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	31	-	-	-	-	-	-	-	-	-	-	1	1	1
1999	17	-	-	-	-	-	-	-	-	-	1	-	1	-
Asthma (J45-J46)														
All origins														
Both sexes														
2002	4,261	4	43	35	88	84	85	102	133	207	265	303	305	288
2001	4,269	10	31	41	58	60	80	88	145	207	248	279	324	265
2000	4,487	8	32	43	86	78	89	105	144	183	275	318	296	270
1999	4,657	5	27	40	86	95	87	121	141	206	241	312	295	343
Male														
2002	1,580	3	30	20	49	63	47	63	66	92	106	113	120	108
2001	1,479	6	22	24	33	35	45	49	67	93	102	103	116	84
2000	1,632	6	20	22	52	52	50	64	78	81	115	128	110	106
1999	1,620	3	16	25	60	53	55	64	64	94	90	96	101	119
Female														
2002	2,681	1	13	15	39	21	38	39	67	115	159	190	185	180
2001	2,790	4	9	17	25	25	35	39	78	114	146	176	208	181
2000	2,855	2	12	21	34	26	39	41	66	102	160	190	186	164
1999	3,037	2	11	15	26	42	32	57	77	112	151	216	194	224
Hispanic														
Both sexes														
2002	287	1	6	6	7	5	6	8	15	18	24	25	19	14
2001	274	2	5	2	6	5	3	8	9	16	15	20	29	26
2000	292	1	5	3	10	6	8	4	12	13	16	19	22	15
1999	320	-	3	4	10	8	6	12	9	24	22	25	23	23
Male														
2002	114	1	3	4	4	4	5	7	7	8	6	14	8	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Emphysema (J43)										
Origin not stated[2]										
Both sexes										
2002	1	3	8	6	4	8	-	-	-	1
2001	5	4	7	8	6	1	4	1	-	-
2000	2	5	14	12	4	4	3	-	-	-
1999	5	5	6	11	6	8	3	-	-	-
Male										
2002	-	2	4	6	1	5	-	-	-	1
2001	3	3	5	4	3	1	1	1	-	-
2000	-	2	8	4	-	1	1	-	-	-
1999	5	4	5	7	3	3	2	-	-	-
Female										
2002	1	1	4	-	3	3	-	-	-	-
2001	2	1	2	4	3	-	3	-	-	-
2000	2	3	6	8	4	3	2	-	-	-
1999	-	1	1	4	3	5	1	-	-	-
Asthma (J45-J46)										
All origins										
Both sexes										
2002	248	257	326	401	411	345	225	86	19	1
2001	288	288	346	384	418	371	239	77	22	-
2000	270	328	406	420	429	388	227	81	11	-
1999	240	353	420	454	471	381	224	97	18	-
Male										
2002	94	89	109	131	125	97	41	10	4	-
2001	98	104	116	116	109	86	57	11	3	-
2000	94	121	132	120	123	104	39	15	-	-
1999	88	115	152	144	129	81	52	17	2	-
Female										
2002	154	168	217	270	286	248	184	76	15	1
2001	190	184	230	268	309	285	182	66	19	-
2000	176	207	274	300	306	284	188	66	11	-
1999	152	238	268	310	342	300	172	80	16	-
Hispanic										
Both sexes										
2002	24	25	20	22	20	11	7	3	1	-
2001	19	19	24	26	18	8	10	4	-	-
2000	33	34	26	25	15	16	6	3	-	-
1999	16	32	28	27	23	19	3	3	-	-
Male										
2002	10	4	6	10	5	2	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Asthma (J45-J46)														
Hispanic														
Male														
2001	103	2	3	1	4	4	2	5	5	8	5	6	8	12
2000	129	1	2	1	7	5	6	2	7	7	7	8	12	5
1999	119	-	-	2	7	4	4	6	5	9	13	9	11	9
Female														
2002	173	-	3	2	3	1	1	1	8	10	18	11	11	9
2001	171	-	2	1	2	1	1	3	4	8	10	14	21	14
2000	163	-	3	2	3	1	2	2	5	6	9	11	10	10
1999	201	-	3	2	3	4	2	6	4	15	9	16	12	14
Non-Hispanic[1]														
Both sexes														
2002	3,950	3	37	29	79	78	79	94	116	188	237	276	285	272
2001	3,976	8	26	39	51	55	77	80	135	190	231	258	292	239
2000	4,164	7	27	39	76	72	80	98	131	170	258	298	270	252
1999	4,324	5	24	36	76	87	81	109	129	182	218	286	271	319
Male														
2002	1,448	2	27	16	43	58	42	56	57	83	96	98	111	103
2001	1,365	4	19	23	29	31	43	44	62	84	96	96	106	72
2000	1,486	5	18	21	45	47	44	59	71	74	107	119	96	98
1999	1,495	3	16	23	53	49	51	58	58	85	77	86	90	110
Female														
2002	2,502	1	10	13	36	20	37	38	59	105	141	178	174	169
2001	2,611	4	7	16	22	24	34	36	73	106	135	162	186	167
2000	2,678	2	9	18	31	25	36	39	60	96	151	179	174	154
1999	2,829	2	8	13	23	38	30	51	71	97	141	200	181	209
Non-Hispanic White														
Both sexes														
2002	2,720	1	14	9	25	35	44	49	72	103	119	161	155	159
2001	2,717	4	7	15	22	26	33	35	62	84	135	144	182	146
2000	2,836	4	13	14	26	30	45	44	70	83	144	158	159	145
1999	3,011	1	8	17	30	51	41	44	63	99	123	167	160	205
Male														
2002	895	-	12	6	14	23	20	24	35	35	45	54	63	58
2001	834	3	4	9	12	11	18	20	22	40	51	52	61	45
2000	920	2	7	7	13	19	25	26	34	30	60	70	60	50
1999	930	1	5	10	21	28	24	22	26	44	36	45	52	72
Female														
2002	1,825	1	2	3	11	12	24	25	37	68	74	107	92	101
2001	1,883	1	3	6	10	15	15	15	40	44	84	92	121	101
2000	1,916	2	6	7	13	11	20	18	36	53	84	88	99	95

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)										
Hispanic										
Male										
2001	5	9	9	8	5	-	2	-	-	-
2000	14	14	11	8	4	6	1	1	-	-
1999	3	10	9	7	5	4	1	1	-	-
Female										
2002	14	21	14	12	15	9	6	3	1	-
2001	14	10	15	18	13	8	8	4	-	-
2000	19	20	15	17	11	10	5	2	-	-
1999	13	22	19	20	18	15	2	2	-	-
Non-Hispanic[1]										
Both sexes										
2002	223	230	306	377	390	332	217	83	18	1
2001	268	267	320	356	400	362	227	73	22	-
2000	234	291	377	392	413	370	220	78	11	-
1999	224	320	391	427	447	359	221	94	18	-
Male										
2002	84	83	103	119	120	93	40	10	4	-
2001	93	95	105	107	104	85	53	11	3	-
2000	78	106	120	110	119	98	37	14	-	-
1999	85	104	142	137	123	76	51	16	2	-
Female										
2002	139	147	203	258	270	239	177	73	14	1
2001	175	172	215	249	296	277	174	62	19	-
2000	156	185	257	282	294	272	183	64	11	-
1999	139	216	249	290	324	283	170	78	16	-
Non-Hispanic White										
Both sexes										
2002	152	155	222	318	341	304	197	72	13	-
2001	160	195	237	285	343	316	201	67	18	-
2000	151	194	290	320	363	320	189	66	8	-
1999	146	223	288	355	392	305	191	85	17	-
Male										
2002	55	52	73	95	101	84	37	7	2	-
2001	50	67	74	83	84	75	43	9	1	-
2000	47	64	95	92	101	81	29	8	-	-
1999	52	67	97	108	110	56	39	13	2	-
Female										
2002	97	103	149	223	240	220	160	65	11	-
2001	110	128	163	202	259	241	158	58	17	-
2000	104	130	195	228	262	239	160	58	8	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Asthma (J45-J46)														
Non-Hispanic White														
Female														
1999	2,081	-	3	7	9	23	17	22	37	55	87	122	108	133
Non-Hispanic Black														
Both sexes														
2002	1,083	2	21	17	52	40	34	41	43	82	114	103	119	105
2001	1,092	3	19	23	28	27	40	44	71	99	87	106	97	84
2000	1,145	3	14	24	45	40	33	47	55	78	100	128	97	88
1999	1,134	4	14	19	46	33	38	62	64	77	89	110	101	96
Male														
2002	488	2	14	9	28	34	22	28	22	46	50	39	44	40
2001	450	-	15	14	16	19	22	24	38	41	38	42	41	23
2000	472	3	11	14	28	27	17	28	34	38	41	44	27	37
1999	474	2	10	13	32	20	26	34	32	36	36	38	34	30
Female														
2002	595	-	7	8	24	6	12	13	21	36	64	64	75	65
2001	642	3	4	9	12	8	18	20	33	58	49	64	56	61
2000	673	-	3	10	17	13	16	19	21	40	59	84	70	51
1999	660	2	4	6	14	13	12	28	32	41	53	72	67	66
Origin not stated[2]														
Both sexes														
2002	24	-	-	-	2	1	-	-	2	1	4	2	1	2
2001	19	-	-	-	1	-	-	-	1	1	2	1	3	-
2000	31	-	-	1	-	-	1	3	1	-	1	1	4	3
1999	13	-	-	-	-	-	-	-	3	-	1	1	1	1
Male														
2002	18	-	-	-	2	1	-	-	2	1	4	1	1	-
2001	11	-	-	-	-	-	-	-	-	1	1	1	2	-
2000	17	-	-	-	-	-	-	3	-	-	1	1	2	3
1999	6	-	-	-	-	-	-	-	1	-	-	1	-	-
Female														
2002	6	-	-	-	-	-	-	-	-	-	-	1	-	2
2001	8	-	-	-	1	-	-	-	1	-	1	-	1	-
2000	14	-	-	1	-	-	1	-	1	-	-	-	2	-
1999	7	-	-	-	-	-	-	-	2	-	1	-	1	1
Other chronic lower respiratory diseases (J44,J47)														
All origins														
Both sexes														
2002	104,111	10	6	3	5	5	10	10	40	100	324	751	1,626	3,267

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Asthma (J45-J46)										
Non-Hispanic White										
Female										
1999	94	156	191	247	282	249	152	72	15	-
Non-Hispanic Black										
Both sexes										
2002	64	57	75	41	34	20	10	7	2	-
2001	86	57	69	58	41	34	13	3	3	-
2000	77	80	72	60	34	32	26	9	3	-
1999	66	76	81	55	39	37	22	5	-	-
Male										
2002	27	25	26	13	10	6	1	2	-	-
2001	29	18	28	17	12	8	3	1	1	-
2000	28	33	20	10	11	9	7	5	-	-
1999	27	26	35	20	7	8	7	1	-	-
Female										
2002	37	32	49	28	24	14	9	5	2	-
2001	57	39	41	41	29	26	10	2	2	-
2000	49	47	52	50	23	23	19	4	3	-
1999	39	50	46	35	32	29	15	4	-	-
Origin not stated[2]										
Both sexes										
2002	1	2	-	2	1	2	1	-	-	-
2001	1	2	2	2	-	1	2	-	-	-
2000	3	3	3	3	1	2	1	-	-	-
1999	-	1	1	-	1	3	-	-	-	-
Male										
2002	-	2	-	2	-	2	-	-	-	-
2001	-	-	2	1	-	1	2	-	-	-
2000	2	1	1	2	-	-	1	-	-	-
1999	-	1	1	-	1	1	-	-	-	-
Female										
2002	1	-	-	-	1	-	1	-	-	-
2001	1	2	-	1	-	-	-	-	-	-
2000	1	2	2	1	1	2	-	-	-	-
1999	-	-	-	-	-	2	-	-	-	-
Other chronic lower respiratory diseases (J44,J47)										
All origins										
Both sexes										
2002	5,720	9,432	15,250	20,761	21,273	15,638	7,443	2,118	314	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic lower respiratory diseases (J44,J47)														
All origins														
Both sexes														
2001	101,543	7	4	1	3	10	14	9	35	115	303	686	1,545	3,181
2000	99,655	7	3	3	2	4	9	11	17	97	273	675	1,450	3,015
1999	100,565	10	8	5	3	9	10	20	28	67	261	617	1,380	3,078
Male														
2002	50,629	5	3	2	1	3	4	6	20	44	177	396	872	1,689
2001	49,407	4	2	-	3	5	10	4	20	58	142	387	809	1,664
2000	49,198	5	1	2	1	3	6	4	7	45	136	380	747	1,591
1999	50,817	7	3	2	3	5	5	5	10	36	136	357	710	1,632
Female														
2002	53,482	5	3	1	4	2	6	4	20	56	147	355	754	1,578
2001	52,136	3	2	1	-	5	4	5	15	57	161	299	736	1,517
2000	50,457	2	2	1	1	1	3	7	10	52	137	295	703	1,424
1999	49,748	3	5	3	-	4	5	15	18	31	125	260	670	1,446
Hispanic														
Both sexes														
2002	2,425	-	2	-	-	1	1	1	4	5	19	23	52	79
2001	2,235	-	2	-	1	1	-	1	3	8	11	21	45	62
2000	2,079	2	1	2	-	-	1	2	-	6	14	22	30	63
1999	2,195	3	2	2	-	-	1	3	3	4	16	22	33	66
Male														
2002	1,323	-	-	-	-	1	1	1	2	3	11	12	29	54
2001	1,189	-	1	-	1	-	-	-	2	4	3	14	21	33
2000	1,127	1	1	1	-	-	1	1	-	2	3	11	14	32
1999	1,228	3	1	1	-	-	1	-	1	4	6	8	16	38
Female														
2002	1,102	-	2	-	-	-	-	-	2	2	8	11	23	25
2001	1,046	-	1	-	-	1	-	1	1	4	8	7	24	29
2000	952	1	-	1	-	-	-	1	-	4	11	11	16	31
1999	967	-	1	1	-	-	-	3	2	-	10	14	17	28
Non-Hispanic[1]														
Both sexes														
2002	101,413	10	4	3	5	4	9	9	35	95	302	725	1,568	3,177
2001	99,030	7	2	1	2	9	14	8	32	106	289	661	1,490	3,103
2000	97,305	5	2	1	2	4	8	9	17	90	258	649	1,411	2,941
1999	98,086	7	6	3	3	9	9	17	25	63	245	593	1,340	3,006
Male														
2002	49,155	5	3	2	1	2	3	5	17	41	165	382	842	1,631
2001	48,070	4	1	-	2	5	10	4	18	54	136	371	781	1,622
2000	47,906	4	-	1	1	3	5	3	7	42	133	366	728	1,553

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)										
All origins										
Both sexes										
2001	5,542	9,602	15,578	20,448	19,914	15,156	7,036	2,040	313	1
2000	5,310	9,462	15,683	20,211	19,818	14,629	6,750	1,927	297	2
1999	5,520	9,779	16,263	20,617	19,799	14,472	6,516	1,829	270	4
Male										
2002	2,939	4,842	7,890	10,534	10,287	7,115	3,043	680	73	4
2001	2,854	4,919	8,108	10,441	9,655	6,721	2,863	654	83	1
2000	2,798	4,910	8,230	10,262	9,786	6,749	2,763	684	86	2
1999	2,983	5,261	8,719	10,777	9,821	6,815	2,762	674	91	3
Female										
2002	2,781	4,590	7,360	10,227	10,986	8,523	4,400	1,438	241	1
2001	2,688	4,683	7,470	10,007	10,259	8,435	4,173	1,386	230	-
2000	2,512	4,552	7,453	9,949	10,032	7,880	3,987	1,243	211	-
1999	2,537	4,518	7,544	9,840	9,978	7,657	3,754	1,155	179	1
Hispanic										
Both sexes										
2002	126	200	309	440	446	385	240	77	15	-
2001	123	212	301	418	356	355	234	69	12	-
2000	106	178	260	379	376	349	194	81	13	-
1999	111	200	311	396	363	354	218	80	7	-
Male										
2002	77	127	176	274	235	178	109	30	3	-
2001	73	133	180	231	175	181	101	32	4	-
2000	77	104	164	216	209	163	85	39	3	-
1999	69	106	194	236	203	188	108	41	4	-
Female										
2002	49	73	133	166	211	207	131	47	12	-
2001	50	79	121	187	181	174	133	37	8	-
2000	29	74	96	163	167	186	109	42	10	-
1999	42	94	117	160	160	166	110	39	3	-
Non-Hispanic[1]										
Both sexes										
2002	5,579	9,203	14,900	20,258	20,776	15,222	7,188	2,038	298	5
2001	5,398	9,368	15,225	19,980	19,516	14,768	6,784	1,967	300	-
2000	5,177	9,249	15,384	19,783	19,397	14,253	6,540	1,841	283	1
1999	5,383	9,552	15,900	20,162	19,391	14,082	6,279	1,745	263	3
Male										
2002	2,852	4,698	7,686	10,222	10,022	6,923	2,929	650	70	4
2001	2,766	4,773	7,901	10,188	9,454	6,527	2,754	620	79	-
2000	2,703	4,781	8,041	10,016	9,550	6,571	2,671	644	82	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic lower respiratory diseases (J44,J47)														
Non-Hispanic[1]														
Male														
1999	49,436	4	2	1	3	5	4	5	9	32	130	348	688	1,590
Female														
2002	52,258	5	1	1	4	2	6	4	18	54	137	343	726	1,546
2001	50,960	3	1	1	-	4	4	4	14	52	153	290	709	1,481
2000	49,399	1	2	-	1	1	3	6	10	48	125	283	683	1,388
1999	48,650	3	4	2	-	4	5	12	16	31	115	245	652	1,416
Non-Hispanic White														
Both sexes														
2002	94,430	6	2	-	5	3	9	7	19	72	251	618	1,341	2,847
2001	92,267	4	1	1	1	5	7	5	24	73	241	551	1,284	2,769
2000	90,746	4	2	1	2	2	5	7	9	68	204	541	1,207	2,609
1999	91,269	1	3	3	2	6	6	12	16	49	189	473	1,176	2,671
Male														
2002	45,172	4	2	-	1	1	3	4	12	31	138	321	709	1,434
2001	44,173	2	-	-	1	2	5	3	15	38	114	316	663	1,429
2000	44,077	4	-	1	1	1	3	2	2	31	109	300	608	1,365
1999	45,423	-	-	1	2	2	2	4	9	26	97	277	590	1,393
Female														
2002	49,258	2	-	-	4	2	6	3	7	41	113	297	632	1,413
2001	48,094	2	1	1	-	3	2	2	9	35	127	235	621	1,340
2000	46,669	-	2	-	1	1	2	5	7	37	95	241	599	1,244
1999	45,846	1	3	2	-	4	4	8	7	23	92	196	586	1,278
Non-Hispanic Black														
Both sexes														
2002	5,739	4	1	3	-	1	-	1	14	22	46	97	207	297
2001	5,550	2	1	-	1	2	6	2	7	31	44	100	183	292
2000	5,414	1	-	-	-	2	1	2	8	20	51	91	187	291
1999	5,695	6	3	-	1	3	3	4	7	14	54	110	148	313
Male														
2002	3,221	1	-	2	-	1	-	-	4	10	24	56	120	179
2001	3,181	1	1	-	1	2	4	1	3	16	21	50	101	169
2000	3,126	-	-	-	-	2	1	1	5	10	23	55	106	165
1999	3,341	4	2	-	1	3	2	-	-	6	32	64	87	184
Female														
2002	2,518	3	1	1	-	-	-	1	10	12	22	41	87	118
2001	2,369	1	-	-	-	-	2	1	4	15	23	50	82	123
2000	2,288	1	-	-	-	-	-	1	3	10	28	36	81	126
1999	2,354	2	1	-	-	-	1	4	7	8	22	46	61	129

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)										
Non-Hispanic[1]										
Male										
1999	2,899	5,137	8,491	10,514	9,598	6,611	2,645	631	87	2
Female										
2002	2,727	4,505	7,214	10,036	10,754	8,299	4,259	1,388	228	1
2001	2,632	4,595	7,324	9,792	10,062	8,241	4,030	1,347	221	-
2000	2,474	4,468	7,343	9,767	9,847	7,682	3,869	1,197	201	-
1999	2,484	4,415	7,409	9,648	9,793	7,471	3,634	1,114	176	1
Non-Hispanic White										
Both sexes										
2002	5,026	8,463	13,852	18,956	19,587	14,407	6,799	1,887	269	4
2001	4,916	8,549	14,185	18,770	18,429	13,990	6,361	1,836	265	-
2000	4,672	8,485	14,318	18,586	18,390	13,523	6,155	1,706	249	1
1999	4,853	8,748	14,753	18,814	18,339	13,359	5,917	1,637	240	2
Male										
2002	2,530	4,286	7,056	9,436	9,343	6,471	2,739	586	62	3
2001	2,488	4,297	7,275	9,427	8,819	6,108	2,547	561	63	-
2000	2,409	4,320	7,385	9,267	8,963	6,167	2,478	590	70	1
1999	2,576	4,652	7,819	9,678	8,975	6,211	2,452	577	79	1
Female										
2002	2,496	4,177	6,796	9,520	10,244	7,936	4,060	1,301	207	1
2001	2,428	4,252	6,910	9,343	9,610	7,882	3,814	1,275	202	-
2000	2,263	4,165	6,933	9,319	9,427	7,356	3,677	1,116	179	-
1999	2,277	4,096	6,934	9,136	9,364	7,148	3,465	1,060	161	1
Non-Hispanic Black										
Both sexes										
2002	479	630	857	1,060	952	617	300	125	25	1
2001	415	697	870	989	872	593	316	102	25	-
2000	457	650	899	971	800	553	298	105	27	-
1999	457	693	946	1,109	852	573	285	93	20	1
Male										
2002	282	352	513	629	524	329	137	50	7	1
2001	244	413	525	609	507	312	145	45	11	-
2000	267	394	559	595	450	298	144	43	8	-
1999	285	419	556	681	500	315	146	46	7	1
Female										
2002	197	278	344	431	428	288	163	75	18	-
2001	171	284	345	380	365	281	171	57	14	-
2000	190	256	340	376	350	255	154	62	19	-
1999	172	274	390	428	352	258	139	47	13	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic lower respiratory diseases (J44,J47)														
Origin not stated[2]														
Both sexes														
2002	273	-	-	-	-	-	-	-	1	-	3	3	6	11
2001	278	-	-	-	-	-	-	-	-	1	3	4	10	16
2000	271	-	-	-	-	-	-	-	-	1	1	4	9	11
1999	284	-	-	-	-	-	-	-	-	-	-	2	7	6
Male														
2002	151	-	-	-	-	-	-	-	1	-	1	2	1	4
2001	148	-	-	-	-	-	-	-	-	-	3	2	7	9
2000	165	-	-	-	-	-	-	-	-	1	-	3	5	6
1999	153	-	-	-	-	-	-	-	-	-	-	1	6	4
Female														
2002	122	-	-	-	-	-	-	-	-	-	2	1	5	7
2001	130	-	-	-	-	-	-	-	-	1	-	2	3	7
2000	106	-	-	-	-	-	-	-	-	-	1	1	4	5
1999	131	-	-	-	-	-	-	-	-	-	-	1	1	2
Pneumoconioses and chemical effects (J60-J66,J68)														
All origins														
Both sexes														
2002	1,114	2	-	-	-	-	-	-	-	-	5	5	12	26
2001	1,136	-	1	-	-	-	-	-	1	3	2	3	8	27
2000	1,161	-	-	-	-	-	-	1	2	2	5	5	7	21
1999	1,113	-	3	-	-	1	-	-	-	3	4	3	12	19
Male														
2002	1,057	-	-	-	-	-	-	-	-	-	4	4	9	24
2001	1,095	-	1	-	-	-	-	-	1	3	2	2	7	26
2000	1,113	-	-	-	-	-	-	1	2	1	5	5	7	19
1999	1,079	-	3	-	-	1	-	-	-	2	3	3	12	19
Female														
2002	57	2	-	-	-	-	-	-	-	-	1	1	3	2
2001	41	-	-	-	-	-	-	-	-	-	-	1	1	1
2000	48	-	-	-	-	-	-	-	-	1	-	-	-	2
1999	34	-	-	-	-	-	-	-	-	1	1	-	-	-
Hispanic														
Both sexes														
2002	33	-	-	-	-	-	-	-	-	-	1	1	1	2
2001	26	-	1	-	-	-	-	-	-	1	1	1	1	1
2000	27	-	-	-	-	-	-	-	1	-	1	-	1	1
1999	13	-	-	-	-	-	-	-	-	-	1	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic lower respiratory diseases (J44,J47)										
Origin not stated[2]										
Both sexes										
2002	15	29	41	63	51	31	15	3	1	-
2001	21	22	52	50	42	33	18	4	1	1
2000	27	35	39	49	45	27	16	5	1	1
1999	26	27	52	59	45	36	19	4	-	1
Male										
2002	10	17	28	38	30	14	5	-	-	-
2001	15	13	27	22	26	13	8	2	-	1
2000	18	25	25	30	27	15	7	1	1	1
1999	15	18	34	27	20	16	9	2	-	1
Female										
2002	5	12	13	25	21	17	10	3	1	-
2001	6	9	25	28	16	20	10	2	1	-
2000	9	10	14	19	18	12	9	4	-	-
1999	11	9	18	32	25	20	10	2	-	-
Pneumoconioses and chemical effects (J60-J66,J68)										
All origins										
Both sexes										
2002	49	80	143	256	224	217	76	17	2	-
2001	37	91	175	252	264	188	66	16	2	-
2000	36	85	196	249	235	217	80	20	-	-
1999	38	104	170	251	252	180	54	18	1	-
Male										
2002	45	77	137	245	212	207	75	17	1	-
2001	36	90	171	239	255	181	63	16	2	-
2000	35	79	186	241	226	208	79	19	-	-
1999	38	99	170	243	243	172	53	17	1	-
Female										
2002	4	3	6	11	12	10	1	-	1	-
2001	1	1	4	13	9	7	3	-	-	-
2000	1	6	10	8	9	9	1	1	-	-
1999	-	5	-	8	9	8	1	1	-	-
Hispanic										
Both sexes										
2002	2	6	6	1	4	4	5	-	-	-
2001	1	1	6	4	2	2	2	1	1	-
2000	-	4	1	3	10	4	-	1	-	-
1999	1	2	3	2	-	2	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumoconioses and chemical effects (J60-J66,J68)														
Hispanic														
Male														
2002	31	-	-	-	-	-	-	-	-	-	1	1	-	2
2001	25	-	1	-	-	-	-	-	-	1	1	1	1	1
2000	26	-	-	-	-	-	-	-	1	-	1	-	1	1
1999	11	-	-	-	-	-	-	-	-	-	1	-	1	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	1	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	1,078	2	-	-	-	-	-	-	-	-	4	4	11	23
2001	1,108	-	-	-	-	-	-	-	1	2	1	2	7	26
2000	1,133	-	-	-	-	-	-	1	1	2	4	5	6	20
1999	1,100	-	3	-	-	1	-	-	-	3	3	3	11	19
Male														
2002	1,023	-	-	-	-	-	-	-	-	-	3	3	9	21
2001	1,068	-	-	-	-	-	-	-	1	2	1	1	6	25
2000	1,086	-	-	-	-	-	-	1	1	1	4	5	6	18
1999	1,068	-	3	-	-	1	-	-	-	2	2	3	11	19
Female														
2002	55	2	-	-	-	-	-	-	-	-	1	1	2	2
2001	40	-	-	-	-	-	-	-	-	-	-	1	1	1
2000	47	-	-	-	-	-	-	-	-	1	-	-	-	2
1999	32	-	-	-	-	-	-	-	-	1	1	-	-	-
Non-Hispanic White														
Both sexes														
2002	1,014	1	-	-	-	-	-	-	-	-	3	3	10	21
2001	1,046	-	-	-	-	-	-	-	1	2	1	-	5	23
2000	1,072	-	-	-	-	-	-	1	1	1	2	5	6	20
1999	1,024	-	-	-	-	-	-	-	-	1	2	3	7	19
Male														
2002	966	-	-	-	-	-	-	-	-	-	2	2	8	21
2001	1,008	-	-	-	-	-	-	-	1	2	1	-	5	22
2000	1,027	-	-	-	-	-	-	1	1	-	2	5	6	18
1999	995	-	-	-	-	-	-	-	-	1	1	3	7	19
Female														
2002	48	1	-	-	-	-	-	-	-	-	1	1	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)										
Hispanic										
Male										
2002	2	6	5	1	4	4	5	-	-	-
2001	1	1	6	4	2	2	1	1	1	-
2000	-	4	1	3	10	3	-	1	-	-
1999	1	2	3	2	-	-	1	-	-	-
Female										
2002	-	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	2	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	47	74	137	255	218	213	71	17	2	-
2001	36	90	169	248	261	186	63	15	1	-
2000	36	81	195	245	225	213	80	19	-	-
1999	37	102	167	249	252	178	53	18	1	-
Male										
2002	43	71	132	244	206	203	70	17	1	-
2001	35	89	165	235	252	179	61	15	1	-
2000	35	75	185	237	216	205	79	18	-	-
1999	37	97	167	241	243	172	52	17	1	-
Female										
2002	4	3	5	11	12	10	1	-	1	-
2001	1	1	4	13	9	7	2	-	-	-
2000	1	6	10	8	9	8	1	1	-	-
1999	-	5	-	8	9	6	1	1	-	-
Non-Hispanic White										
Both sexes										
2002	41	71	130	238	207	208	66	14	1	-
2001	30	86	159	240	247	175	61	15	1	-
2000	34	76	185	230	212	206	76	17	-	-
1999	27	93	157	237	245	166	52	15	-	-
Male										
2002	38	68	125	228	196	199	65	14	-	-
2001	29	85	155	227	238	168	59	15	1	-
2000	33	70	175	223	204	198	75	16	-	-
1999	27	89	157	229	236	161	51	14	-	-
Female										
2002	3	3	5	10	11	9	1	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumoconioses and chemical effects (J60-J66,J68)														
Non-Hispanic White														
Female														
2001	38	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	45	-	-	-	-	-	-	-	-	1	-	-	-	2
1999	29	-	-	-	-	-	-	-	-	-	1	-	-	-
Non-Hispanic Black														
Both sexes														
2002	56	1	-	-	-	-	-	-	-	-	1	1	1	2
2001	56	-	-	-	-	-	-	-	-	-	-	2	2	3
2000	54	-	-	-	-	-	-	-	-	1	2	-	-	-
1999	67	-	3	-	-	1	-	-	-	2	1	-	4	-
Male														
2002	50	-	-	-	-	-	-	-	-	-	1	1	1	-
2001	54	-	-	-	-	-	-	-	-	-	-	1	1	3
2000	52	-	-	-	-	-	-	-	-	1	2	-	-	-
1999	64	-	3	-	-	1	-	-	-	1	1	-	4	-
Female														
2002	6	1	-	-	-	-	-	-	-	-	-	-	-	2
2001	2	-	-	-	-	-	-	-	-	-	-	1	1	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	1	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	3	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	3	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumoconioses and chemical effects (J60-J66,J68)										
Non-Hispanic White										
Female										
2001	1	1	4	13	9	7	2	-	-	-
2000	1	6	10	7	8	8	1	1	-	-
1999	-	4	-	8	9	5	1	1	-	-
Non-Hispanic Black										
Both sexes										
2002	4	3	7	12	10	5	5	3	1	-
2001	5	4	8	8	12	10	2	-	-	-
2000	2	5	10	13	10	7	3	1	-	-
1999	8	8	8	9	7	11	1	3	1	-
Male										
2002	4	3	7	11	9	4	5	3	1	-
2001	5	4	8	8	12	10	2	-	-	-
2000	2	5	10	12	9	7	3	1	-	-
1999	8	7	8	9	7	10	1	3	1	-
Female										
2002	-	-	-	1	1	1	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	1	1	-	-	-	-	-
1999	-	1	-	-	-	1	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	2	-	-	-	-	-
2001	-	-	-	-	1	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	2	-	-	-	-	-
2001	-	-	-	-	1	-	1	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonitis due to solids and liquids (J69)														
All origins														
Both sexes														
2002	17,593	2	7	7	9	16	28	31	43	72	120	143	208	271
2001	17,301	12	14	5	10	10	24	31	48	61	118	142	236	259
2000	16,636	10	11	8	5	12	25	22	40	65	116	157	209	257
1999	15,268	16	12	4	12	9	23	33	45	59	107	129	155	259
Male														
2002	8,876	-	3	3	6	10	17	17	25	41	78	90	136	162
2001	8,576	6	6	3	4	5	17	15	25	42	75	81	151	160
2000	8,390	4	5	5	4	8	15	11	24	45	70	99	126	153
1999	7,755	9	3	-	6	7	15	21	26	35	62	83	82	151
Female														
2002	8,717	2	4	4	3	6	11	14	18	31	42	53	72	109
2001	8,725	6	8	2	6	5	7	16	23	19	43	61	85	99
2000	8,246	6	6	3	1	4	10	11	16	20	46	58	83	104
1999	7,513	7	9	4	6	2	8	12	19	24	45	46	73	108
Hispanic														
Both sexes														
2002	547	1	-	2	1	3	4	1	2	4	6	8	7	12
2001	555	-	3	1	4	2	1	-	4	4	6	7	11	16
2000	496	3	2	1	-	1	2	3	4	6	9	11	9	15
1999	413	2	2	2	3	1	1	3	1	1	2	4	4	8
Male														
2002	282	-	-	1	-	2	3	-	2	1	4	4	4	9
2001	283	-	1	1	1	-	1	-	3	2	3	4	6	14
2000	253	-	-	-	-	1	1	1	1	4	4	8	7	10
1999	195	1	1	-	1	-	-	2	1	1	-	3	1	4
Female														
2002	265	1	-	1	1	1	1	1	-	3	2	4	3	3
2001	272	-	2	-	3	2	-	-	1	2	3	3	5	2
2000	243	3	2	1	-	-	1	2	3	2	5	3	2	5
1999	218	1	1	2	2	1	1	1	-	-	2	1	3	4
Non-Hispanic[1]														
Both sexes														
2002	17,006	1	7	5	8	13	23	30	41	68	114	135	201	257
2001	16,700	12	11	4	6	8	23	31	44	56	111	133	225	243
2000	16,094	7	9	7	5	11	23	19	36	59	107	145	198	239
1999	14,812	14	10	2	9	8	22	30	44	57	105	124	150	250
Male														
2002	8,565	-	3	2	6	8	13	17	23	40	74	86	132	151
2001	8,270	6	5	2	3	5	16	15	22	39	71	76	145	146
2000	8,107	4	5	5	4	7	14	10	23	41	66	90	117	142

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)										
All origins										
Both sexes										
2002	400	627	1,166	2,323	3,577	4,100	2,995	1,219	229	-
2001	400	589	1,241	2,259	3,500	3,958	2,960	1,160	264	-
2000	341	642	1,257	2,258	3,247	3,825	2,779	1,111	239	-
1999	327	582	1,172	2,068	2,962	3,443	2,586	1,053	212	-
Male										
2002	231	380	703	1,332	1,911	1,994	1,267	415	55	-
2001	224	362	744	1,253	1,892	1,893	1,180	379	59	-
2000	201	405	790	1,291	1,747	1,839	1,116	367	65	-
1999	214	367	716	1,252	1,623	1,659	1,006	372	46	-
Female										
2002	169	247	463	991	1,666	2,106	1,728	804	174	-
2001	176	227	497	1,006	1,608	2,065	1,780	781	205	-
2000	140	237	467	967	1,500	1,986	1,663	744	174	-
1999	113	215	456	816	1,339	1,784	1,580	681	166	-
Hispanic										
Both sexes										
2002	23	24	38	73	101	110	86	35	6	-
2001	17	27	42	67	85	112	97	41	8	-
2000	18	21	43	68	73	87	78	38	4	-
1999	18	21	36	57	61	85	67	28	6	-
Male										
2002	15	14	18	42	63	46	41	11	2	-
2001	8	17	25	37	44	56	41	16	3	-
2000	10	10	25	44	43	41	33	10	-	-
1999	10	12	19	34	31	39	22	13	-	-
Female										
2002	8	10	20	31	38	64	45	24	4	-
2001	9	10	17	30	41	56	56	25	5	-
2000	8	11	18	24	30	46	45	28	4	-
1999	8	9	17	23	30	46	45	15	6	-
Non-Hispanic[1]										
Both sexes										
2002	377	602	1,121	2,249	3,468	3,980	2,904	1,180	222	-
2001	381	562	1,194	2,188	3,405	3,837	2,854	1,116	256	-
2000	323	621	1,212	2,183	3,163	3,726	2,696	1,070	235	-
1999	309	560	1,131	2,004	2,891	3,349	2,513	1,024	206	-
Male										
2002	216	366	679	1,289	1,841	1,939	1,224	403	53	-
2001	215	345	716	1,213	1,844	1,832	1,137	361	56	-
2000	191	395	763	1,240	1,697	1,793	1,081	354	65	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonitis due to solids and liquids (J69)														
Non-Hispanic[1]														
Male														
1999	7,534	8	2	-	5	7	15	19	25	33	62	79	80	146
Female														
2002	8,441	1	4	3	2	5	10	13	18	28	40	49	69	106
2001	8,430	6	6	2	3	3	7	16	22	17	40	57	80	97
2000	7,987	3	4	2	1	4	9	9	13	18	41	55	81	97
1999	7,278	6	8	2	4	1	7	11	19	24	43	45	70	104
Non-Hispanic White														
Both sexes														
2002	15,203	1	5	4	7	10	17	26	30	51	87	111	151	215
2001	14,843	9	9	3	3	6	16	26	32	41	90	107	173	205
2000	14,343	6	7	7	4	9	15	14	26	37	91	104	155	196
1999	13,139	11	6	2	5	7	13	20	33	42	76	93	116	195
Male														
2002	7,654	-	3	2	5	6	9	16	17	30	56	70	98	123
2001	7,411	6	3	2	1	4	13	14	18	28	59	67	112	125
2000	7,214	4	4	5	3	6	10	7	14	28	57	64	90	120
1999	6,642	6	1	-	4	6	8	11	19	25	45	62	60	113
Female														
2002	7,549	1	2	2	2	4	8	10	13	21	31	41	53	92
2001	7,432	3	6	1	2	2	3	12	14	13	31	40	61	80
2000	7,129	2	3	2	1	3	5	7	12	9	34	40	65	76
1999	6,497	5	5	2	1	1	5	9	14	17	31	31	56	82
Non-Hispanic Black														
Both sexes														
2002	1,528	-	2	1	1	2	6	4	10	11	27	20	44	35
2001	1,584	3	2	1	3	1	6	5	11	12	21	23	48	34
2000	1,507	1	2	-	1	2	7	5	9	21	16	38	41	39
1999	1,495	2	4	-	3	1	9	9	9	13	27	28	31	51
Male														
2002	755	-	-	-	1	1	4	1	5	5	18	12	30	24
2001	731	-	2	-	2	-	2	1	3	8	12	7	31	20
2000	757	-	1	-	1	1	3	3	8	12	9	25	27	20
1999	768	1	1	-	1	1	7	7	5	7	15	15	18	30
Female														
2002	773	-	2	1	-	1	2	3	5	6	9	8	14	11
2001	853	3	-	1	1	1	4	4	8	4	9	16	17	14
2000	750	1	1	-	-	1	4	2	1	9	7	13	14	19
1999	727	1	3	-	2	-	2	2	4	6	12	13	13	21

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)										
Non-Hispanic[1]										
Male										
1999	204	354	692	1,214	1,589	1,615	981	358	46	-
Female										
2002	161	236	442	960	1,627	2,041	1,680	777	169	-
2001	166	217	478	975	1,561	2,005	1,717	755	200	-
2000	132	226	449	943	1,466	1,933	1,615	716	170	-
1999	105	206	439	790	1,302	1,734	1,532	666	160	-
Non-Hispanic White										
Both sexes										
2002	304	480	957	1,969	3,162	3,664	2,686	1,067	199	-
2001	313	463	1,006	1,933	3,079	3,503	2,601	1,011	214	-
2000	253	509	1,006	1,934	2,859	3,437	2,482	985	207	-
1999	251	458	942	1,758	2,603	3,086	2,317	919	186	-
Male										
2002	180	295	579	1,131	1,691	1,800	1,124	367	52	-
2001	173	290	612	1,079	1,686	1,688	1,048	333	50	-
2000	150	319	626	1,091	1,557	1,676	995	328	60	-
1999	163	286	574	1,063	1,429	1,492	908	323	44	-
Female										
2002	124	185	378	838	1,471	1,864	1,562	700	147	-
2001	140	173	394	854	1,393	1,815	1,553	678	164	-
2000	103	190	380	843	1,302	1,761	1,487	657	147	-
1999	88	172	368	695	1,174	1,594	1,409	596	142	-
Non-Hispanic Black										
Both sexes										
2002	63	110	140	243	256	259	184	93	17	-
2001	66	86	165	221	276	271	203	92	34	-
2000	63	102	187	216	252	232	178	74	21	-
1999	54	90	174	225	244	233	176	95	17	-
Male										
2002	32	67	85	135	118	105	81	30	1	-
2001	40	49	91	120	132	115	69	22	5	-
2000	35	67	123	129	114	88	66	21	4	-
1999	38	59	109	135	126	103	58	30	2	-
Female										
2002	31	43	55	108	138	154	103	63	16	-
2001	26	37	74	101	144	156	134	70	29	-
2000	28	35	64	87	138	144	112	53	17	-
1999	16	31	65	90	118	130	118	65	15	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pneumonitis due to solids and liquids (J69)														
Origin not stated[2]														
Both sexes														
2002	40	-	-	-	-	-	1	-	-	-	-	-	-	2
2001	46	-	-	-	-	-	-	-	-	1	1	2	-	-
2000	46	-	-	-	-	-	-	-	-	-	-	1	2	3
1999	43	-	-	-	-	-	-	-	-	1	-	1	1	1
Male														
2002	29	-	-	-	-	-	1	-	-	-	-	-	-	2
2001	23	-	-	-	-	-	-	-	-	1	1	1	-	-
2000	30	-	-	-	-	-	-	-	-	-	-	1	2	1
1999	26	-	-	-	-	-	-	-	-	1	-	1	1	1
Female														
2002	11	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	23	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	16	-	-	-	-	-	-	-	-	-	-	-	-	2
1999	17	-	-	-	-	-	-	-	-	-	-	-	-	-
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)														
All origins														
Both sexes														
2002	25,039	278	85	41	43	51	75	70	115	191	337	522	766	1,055
2001	26,178	294	93	30	48	55	86	97	119	218	352	542	814	1,030
2000	25,535	273	87	36	38	60	71	99	154	231	384	527	805	1,044
1999	25,031	252	91	37	39	59	56	97	160	228	348	525	694	904
Male														
2002	12,057	164	56	21	18	29	39	31	57	101	180	263	393	573
2001	12,495	186	47	18	25	30	46	45	59	105	199	295	442	530
2000	12,391	154	45	21	17	30	43	48	79	132	203	281	431	572
1999	12,110	145	55	20	19	37	32	45	82	125	192	269	368	467
Female														
2002	12,982	114	29	20	25	22	36	39	58	90	157	259	373	482
2001	13,683	108	46	12	23	25	40	52	60	113	153	247	372	500
2000	13,144	119	42	15	21	30	28	51	75	99	181	246	374	472
1999	12,921	107	36	17	20	22	24	52	78	103	156	256	326	437
Hispanic														
Both sexes														
2002	1,273	43	18	8	6	6	5	11	16	13	26	45	54	71
2001	1,275	41	17	2	5	2	16	4	13	14	25	42	64	83
2000	1,225	43	17	3	7	6	9	14	20	27	29	39	42	83
1999	1,157	38	10	5	3	9	6	9	14	12	33	39	55	53

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonitis due to solids and liquids (J69)										
Origin not stated[2]										
Both sexes										
2002	-	1	7	1	8	10	5	4	1	-
2001	2	-	5	4	10	9	9	3	-	-
2000	-	-	2	7	11	12	5	3	-	-
1999	-	1	5	7	10	9	6	1	-	-
Male										
2002	-	-	6	1	7	9	2	1	-	-
2001	1	-	3	3	4	5	2	2	-	-
2000	-	-	2	7	7	5	2	3	-	-
1999	-	1	5	4	3	5	3	1	-	-
Female										
2002	-	1	1	-	1	1	3	3	1	-
2001	1	-	2	1	6	4	7	1	-	-
2000	-	-	-	-	4	7	3	-	-	-
1999	-	-	-	3	7	4	3	-	-	-
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
All origins										
Both sexes										
2002	1,385	1,971	3,266	4,212	4,446	3,445	1,908	639	138	-
2001	1,402	2,174	3,336	4,590	4,612	3,627	1,895	643	118	3
2000	1,367	2,184	3,264	4,487	4,413	3,394	1,850	629	135	3
1999	1,356	2,143	3,266	4,369	4,277	3,498	1,823	667	142	-
Male										
2002	766	1,084	1,845	2,166	2,095	1,395	625	136	20	-
2001	792	1,197	1,828	2,330	2,123	1,415	618	143	21	1
2000	752	1,179	1,782	2,353	2,062	1,409	612	156	28	2
1999	746	1,199	1,806	2,255	2,073	1,417	551	178	29	-
Female										
2002	619	887	1,421	2,046	2,351	2,050	1,283	503	118	-
2001	610	977	1,508	2,260	2,489	2,212	1,277	500	97	2
2000	615	1,005	1,482	2,134	2,351	1,985	1,238	473	107	1
1999	610	944	1,460	2,114	2,204	2,081	1,272	489	113	-
Hispanic										
Both sexes										
2002	93	123	185	185	163	118	57	22	5	-
2001	82	125	162	186	177	128	61	22	4	-
2000	84	123	150	174	151	108	67	21	8	-
1999	104	131	138	174	133	110	57	22	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)														
Hispanic														
Male														
2002	635	21	14	4	4	4	5	5	13	9	14	23	25	43
2001	643	21	8	-	3	1	8	4	8	10	16	24	34	43
2000	608	23	8	1	3	2	6	9	10	18	16	22	24	44
1999	583	20	6	-	3	4	3	5	7	6	23	22	23	31
Female														
2002	638	22	4	4	2	2	-	6	3	4	12	22	29	28
2001	632	20	9	2	2	1	8	-	5	4	9	18	30	40
2000	617	20	9	2	4	4	3	5	10	9	13	17	18	39
1999	574	18	4	5	-	5	3	4	7	6	10	17	32	22
Non-Hispanic[1]														
Both sexes														
2002	23,706	235	67	33	37	45	70	59	99	177	310	472	705	981
2001	24,835	251	76	28	43	53	70	93	105	204	324	497	748	946
2000	24,236	228	69	33	31	53	62	85	133	203	353	486	760	956
1999	23,802	211	81	32	36	50	50	88	145	216	313	484	636	843
Male														
2002	11,397	143	42	17	14	25	34	26	44	92	166	238	364	527
2001	11,818	165	39	18	22	29	38	41	51	95	181	270	407	486
2000	11,752	130	37	20	14	28	37	39	69	113	185	259	404	528
1999	11,483	123	49	20	16	33	29	40	75	119	167	246	343	431
Female														
2002	12,309	92	25	16	23	20	36	33	55	85	144	234	341	454
2001	13,017	86	37	10	21	24	32	52	54	109	143	227	341	460
2000	12,484	98	32	13	17	25	25	46	64	90	168	227	356	428
1999	12,319	88	32	12	20	17	21	48	70	97	146	238	293	412
Non-Hispanic White														
Both sexes														
2002	20,879	109	39	21	22	30	40	39	66	118	221	318	541	790
2001	21,835	116	44	20	32	36	46	57	65	123	227	344	539	765
2000	21,397	122	46	21	22	33	38	58	90	125	260	336	592	782
1999	20,959	111	52	20	27	38	37	56	99	143	208	335	491	662
Male														
2002	10,043	67	26	13	6	15	24	17	28	61	117	162	284	434
2001	10,433	76	20	14	17	20	26	22	30	63	139	191	307	407
2000	10,349	77	22	13	11	15	23	26	48	69	136	178	315	435
1999	10,154	66	28	13	13	27	24	34	52	82	118	171	272	330
Female														
2002	10,836	42	13	8	16	15	16	22	38	57	104	156	257	356

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
Hispanic										
Male										
2002	49	72	109	78	78	45	15	3	2	-
2001	49	66	85	95	80	57	22	6	3	-
2000	52	61	78	92	73	38	20	5	3	-
1999	58	69	70	91	67	48	21	5	1	-
Female										
2002	44	51	76	107	85	73	42	19	3	-
2001	33	59	77	91	97	71	39	16	1	-
2000	32	62	72	82	78	70	47	16	5	-
1999	46	62	68	83	66	62	36	17	1	-
Non-Hispanic[1]										
Both sexes										
2002	1,287	1,840	3,076	4,022	4,274	3,322	1,848	615	132	-
2001	1,319	2,044	3,159	4,390	4,425	3,495	1,830	619	113	3
2000	1,282	2,054	3,104	4,304	4,248	3,278	1,779	606	127	2
1999	1,249	2,008	3,119	4,183	4,132	3,380	1,762	644	140	-
Male										
2002	716	1,009	1,734	2,086	2,012	1,347	610	133	18	-
2001	742	1,128	1,732	2,228	2,037	1,357	596	137	18	1
2000	699	1,115	1,700	2,254	1,985	1,368	591	151	25	1
1999	686	1,128	1,729	2,157	1,996	1,366	529	173	28	-
Female										
2002	571	831	1,342	1,936	2,262	1,975	1,238	482	114	-
2001	577	916	1,427	2,162	2,388	2,138	1,234	482	95	2
2000	583	939	1,404	2,050	2,263	1,910	1,188	455	102	1
1999	563	880	1,390	2,026	2,136	2,014	1,233	471	112	-
Non-Hispanic White										
Both sexes										
2002	1,060	1,598	2,731	3,688	3,931	3,099	1,726	572	120	-
2001	1,095	1,768	2,829	4,020	4,113	3,258	1,688	549	100	1
2000	1,065	1,765	2,792	3,943	3,923	3,078	1,641	553	112	-
1999	1,056	1,743	2,768	3,817	3,825	3,133	1,641	574	123	-
Male										
2002	589	902	1,546	1,920	1,862	1,269	563	122	16	-
2001	611	993	1,581	2,060	1,903	1,270	549	118	16	-
2000	586	975	1,533	2,068	1,845	1,287	536	133	18	-
1999	589	999	1,562	1,988	1,860	1,267	486	151	22	-
Female										
2002	471	696	1,185	1,768	2,069	1,830	1,163	450	104	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)														
Non-Hispanic White														
Female														
2001	11,402	40	24	6	15	16	20	35	35	60	88	153	232	358
2000	11,048	45	24	8	11	18	15	32	42	56	124	158	277	347
1999	10,805	45	24	7	14	11	13	22	47	61	90	164	219	332
Non-Hispanic Black														
Both sexes														
2002	2,257	112	28	12	13	15	26	18	29	51	82	138	140	161
2001	2,462	116	26	6	11	15	23	29	38	72	91	141	185	154
2000	2,336	102	20	10	8	17	22	22	36	69	83	140	144	155
1999	2,339	90	26	10	9	10	11	29	44	68	92	133	125	154
Male														
2002	1,076	67	16	4	6	10	9	7	16	26	47	66	70	80
2001	1,116	76	14	3	5	8	11	14	20	27	41	71	86	67
2000	1,126	50	14	7	3	11	12	12	18	40	44	78	73	81
1999	1,081	52	19	6	3	5	4	5	22	34	40	70	62	81
Female														
2002	1,181	45	12	8	7	5	17	11	13	25	35	72	70	81
2001	1,346	40	12	3	6	7	12	15	18	45	50	70	99	87
2000	1,210	52	6	3	5	6	10	10	18	29	39	62	71	74
1999	1,258	38	7	4	6	5	7	24	22	34	52	63	63	73
Origin not stated[2]														
Both sexes														
2002	60	-	-	-	-	-	-	-	-	1	1	5	7	3
2001	68	2	-	-	-	-	-	-	1	-	3	3	2	1
2000	74	2	1	-	-	1	-	-	1	1	2	2	3	5
1999	72	3	-	-	-	-	-	-	1	-	2	2	3	8
Male														
2002	25	-	-	-	-	-	-	-	-	-	-	2	4	3
2001	34	-	-	-	-	-	-	-	-	-	2	1	1	1
2000	31	1	-	-	-	-	-	-	-	1	2	-	3	-
1999	44	2	-	-	-	-	-	-	-	-	2	1	2	5
Female														
2002	35	-	-	-	-	-	-	-	-	1	1	3	3	-
2001	34	2	-	-	-	-	-	-	1	-	1	2	1	-
2000	43	1	1	-	-	1	-	-	1	-	-	2	-	5
1999	28	1	-	-	-	-	-	-	1	-	-	1	1	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)										
Non-Hispanic White										
Female										
2001	484	775	1,248	1,960	2,210	1,988	1,139	431	84	1
2000	479	790	1,259	1,875	2,078	1,791	1,105	420	94	-
1999	467	744	1,206	1,829	1,965	1,866	1,155	423	101	-
Non-Hispanic Black										
Both sexes										
2002	192	195	251	236	258	157	98	34	11	-
2001	189	220	262	284	243	171	114	57	13	2
2000	170	249	246	277	250	148	111	42	13	2
1999	163	223	280	295	228	192	89	54	14	-
Male										
2002	107	89	133	116	113	50	33	9	2	-
2001	106	107	115	126	104	65	34	13	2	1
2000	85	121	121	143	97	54	41	15	5	1
1999	83	108	129	139	96	74	30	14	5	-
Female										
2002	85	106	118	120	145	107	65	25	9	-
2001	83	113	147	158	139	106	80	44	11	1
2000	85	128	125	134	153	94	70	27	8	1
1999	80	115	151	156	132	118	59	40	9	-
Origin not stated[2]										
Both sexes										
2002	5	8	5	5	9	5	3	2	1	-
2001	1	5	15	14	10	4	4	2	1	-
2000	1	7	10	9	14	8	4	2	-	1
1999	3	4	9	12	12	8	4	1	-	-
Male										
2002	1	3	2	2	5	3	-	-	-	-
2001	1	3	11	7	6	1	-	-	-	-
2000	1	3	4	7	4	3	1	-	-	1
1999	2	2	7	7	10	3	1	-	-	-
Female										
2002	4	5	3	3	4	2	3	2	1	-
2001	-	2	4	7	4	3	4	2	1	-
2000	-	4	6	2	10	5	3	2	-	-
1999	1	2	2	5	2	5	3	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Peptic ulcer (K25-K28)														
All origins														
Both sexes														
2002	4,079	1	1	-	1	-	-	5	17	40	73	118	154	209
2001	4,491	-	1	3	2	3	3	13	19	36	82	137	147	183
2000	4,558	-	-	-	-	4	5	3	11	42	71	130	177	200
1999	4,604	2	2	-	-	1	1	7	14	38	85	113	142	186
Male														
2002	1,951	-	1	-	1	-	-	2	12	28	44	71	106	129
2001	2,074	-	1	2	-	2	1	10	11	25	62	91	95	123
2000	2,201	-	-	-	-	3	4	1	5	31	48	86	131	134
1999	2,189	-	-	-	-	1	1	5	7	27	49	67	102	111
Female														
2002	2,128	1	-	-	-	-	-	3	5	12	29	47	48	80
2001	2,417	-	-	1	2	1	2	3	8	11	20	46	52	60
2000	2,357	-	-	-	-	1	1	2	6	11	23	44	46	66
1999	2,415	2	2	-	-	-	-	2	7	11	36	46	40	75
Hispanic														
Both sexes														
2002	187	-	-	-	-	-	-	1	-	2	3	10	10	15
2001	166	-	-	-	-	-	1	-	-	-	1	6	10	12
2000	186	-	-	-	-	-	-	-	-	5	4	7	10	11
1999	172	-	1	-	-	-	-	4	4	1	3	7	8	14
Male														
2002	115	-	-	-	-	-	-	1	-	1	2	9	8	11
2001	99	-	-	-	-	-	1	-	-	-	1	4	7	10
2000	97	-	-	-	-	-	-	-	-	3	3	7	7	8
1999	101	-	-	-	-	-	-	3	2	1	2	3	7	9
Female														
2002	72	-	-	-	-	-	-	-	-	1	1	1	2	4
2001	67	-	-	-	-	-	-	-	-	-	-	2	3	2
2000	89	-	-	-	-	-	-	-	-	2	1	-	3	3
1999	71	-	1	-	-	-	-	1	2	-	1	4	1	5
Non-Hispanic[1]														
Both sexes														
2002	3,879	1	1	-	1	-	-	4	17	38	70	107	144	191
2001	4,308	-	1	3	2	3	2	13	19	36	81	131	136	170
2000	4,357	-	-	-	-	4	5	3	11	37	65	123	167	188
1999	4,415	2	1	-	-	1	1	3	10	37	81	106	133	170
Male														
2002	1,829	-	1	-	1	-	-	1	12	27	42	61	98	117
2001	1,966	-	1	2	-	2	-	10	11	25	61	87	88	113
2000	2,099	-	-	-	-	3	4	1	5	28	44	79	124	126

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)										
All origins										
Both sexes										
2002	202	254	411	592	699	714	407	148	33	-
2001	235	294	424	615	823	788	458	184	40	1
2000	243	299	449	642	773	801	498	177	33	-
1999	236	314	522	665	835	771	470	164	34	2
Male										
2002	118	142	236	287	335	279	116	39	5	-
2001	150	179	218	315	334	277	125	47	5	1
2000	150	172	270	341	326	291	150	49	9	-
1999	154	196	308	331	381	279	138	29	2	1
Female										
2002	84	112	175	305	364	435	291	109	28	-
2001	85	115	206	300	489	511	333	137	35	-
2000	93	127	179	301	447	510	348	128	24	-
1999	82	118	214	334	454	492	332	135	32	1
Hispanic										
Both sexes										
2002	17	21	18	27	26	21	10	6	-	-
2001	10	10	23	24	30	22	13	2	1	1
2000	17	13	23	19	31	22	17	5	2	-
1999	11	16	21	20	24	22	12	3	1	-
Male										
2002	12	16	12	14	13	8	5	3	-	-
2001	9	5	13	17	20	6	4	1	-	1
2000	15	8	13	11	6	12	2	2	-	-
1999	9	10	12	14	13	9	5	2	-	-
Female										
2002	5	5	6	13	13	13	5	3	-	-
2001	1	5	10	7	10	16	9	1	1	-
2000	2	5	10	8	25	10	15	3	2	-
1999	2	6	9	6	11	13	7	1	1	-
Non-Hispanic[1]										
Both sexes										
2002	183	233	392	564	671	691	396	142	33	-
2001	223	280	401	589	791	763	445	180	39	-
2000	226	285	424	622	739	778	478	171	31	-
1999	223	297	499	642	809	746	458	161	33	2
Male										
2002	105	126	223	272	322	270	110	36	5	-
2001	139	170	205	297	312	271	121	46	5	-
2000	135	163	256	329	319	279	148	47	9	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Peptic ulcer (K25-K28)														
Non-Hispanic[1]														
Male														
1999	2,079	-	-	-	-	1	1	2	5	26	46	64	94	101
Female														
2002	2,050	1	-	-	-	-	-	3	5	11	28	46	46	74
2001	2,342	-	-	1	2	1	2	3	8	11	20	44	48	57
2000	2,258	-	-	-	-	1	1	2	6	9	21	44	43	62
1999	2,336	2	1	-	-	-	-	1	5	11	35	42	39	69
Non-Hispanic White														
Both sexes														
2002	3,394	1	1	-	-	-	-	1	10	28	44	84	108	146
2001	3,766	-	1	3	1	2	-	5	11	19	58	96	101	126
2000	3,825	-	-	-	-	4	5	3	6	26	44	88	122	151
1999	3,859	1	1	-	-	1	1	2	5	20	53	73	98	136
Male														
2002	1,569	-	1	-	-	-	-	1	8	19	28	47	69	87
2001	1,649	-	1	2	-	1	-	4	7	12	43	61	61	82
2000	1,799	-	-	-	-	3	4	1	3	19	30	61	89	102
1999	1,783	-	-	-	-	1	1	2	1	15	26	42	69	78
Female														
2002	1,825	1	-	-	-	-	-	-	2	9	16	37	39	59
2001	2,117	-	-	1	1	1	-	1	4	7	15	35	40	44
2000	2,026	-	-	-	-	1	1	2	3	7	14	27	33	49
1999	2,076	1	1	-	-	-	-	-	4	5	27	31	29	58
Non-Hispanic Black														
Both sexes														
2002	376	-	-	-	1	-	-	3	7	9	26	23	33	35
2001	403	-	-	-	1	1	1	8	7	16	19	27	32	34
2000	420	-	-	-	-	-	-	-	4	10	19	32	38	34
1999	443	1	-	-	-	-	-	-	4	16	27	28	32	32
Male														
2002	209	-	-	-	1	-	-	-	4	7	14	14	27	24
2001	238	-	-	-	-	1	-	6	4	13	14	21	25	25
2000	228	-	-	-	-	-	-	-	2	8	13	17	28	21
1999	240	-	-	-	-	-	-	-	3	11	19	20	23	22
Female														
2002	167	-	-	-	-	-	-	3	3	2	12	9	6	11
2001	165	-	-	-	1	-	1	2	3	3	5	6	7	9
2000	192	-	-	-	-	-	-	-	2	2	6	15	10	13
1999	203	1	-	-	-	-	-	-	1	5	8	8	9	10

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)										
Non-Hispanic[1]										
Male										
1999	143	186	295	315	367	270	133	27	2	1
Female										
2002	78	107	169	292	349	421	286	106	28	-
2001	84	110	196	292	479	492	324	134	34	-
2000	91	122	168	293	420	499	330	124	22	-
1999	80	111	204	327	442	476	325	134	31	1
Non-Hispanic White										
Both sexes										
2002	142	193	344	497	626	639	370	132	28	-
2001	182	232	351	522	726	706	418	168	38	-
2000	192	236	373	550	683	709	448	158	27	-
1999	186	242	435	578	735	690	427	147	27	1
Male										
2002	82	104	192	239	300	251	103	33	5	-
2001	112	140	171	259	284	251	111	42	5	-
2000	115	137	220	289	291	251	136	40	8	-
1999	117	150	259	287	335	247	128	22	2	1
Female										
2002	60	89	152	258	326	388	267	99	23	-
2001	70	92	180	263	442	455	307	126	33	-
2000	77	99	153	261	392	458	312	118	19	-
1999	69	92	176	291	400	443	299	125	25	-
Non-Hispanic Black										
Both sexes										
2002	33	30	39	50	25	37	13	8	4	-
2001	34	32	41	45	38	38	20	8	1	-
2000	29	37	42	52	39	49	23	8	4	-
1999	33	45	47	42	56	40	24	11	4	1
Male										
2002	21	17	24	25	13	14	2	2	-	-
2001	22	20	27	23	16	15	5	1	-	-
2000	19	18	30	26	19	18	5	3	1	-
1999	24	29	26	18	21	18	3	3	-	-
Female										
2002	12	13	15	25	12	23	11	6	4	-
2001	12	12	14	22	22	23	15	7	1	-
2000	10	19	12	26	20	31	18	5	3	-
1999	9	16	21	24	35	22	21	8	4	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Peptic ulcer (K25-K28)														
Origin not stated[2]														
Both sexes														
2002	13	-	-	-	-	-	-	-	-	-	-	1	-	3
2001	17	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	15	-	-	-	-	-	-	-	-	-	2	-	-	1
1999	17	-	-	-	-	-	-	-	-	-	1	-	1	2
Male														
2002	7	-	-	-	-	-	-	-	-	-	-	1	-	1
2001	9	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	5	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	9	-	-	-	-	-	-	-	-	-	1	-	1	1
Female														
2002	6	-	-	-	-	-	-	-	-	-	-	-	-	2
2001	8	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	10	-	-	-	-	-	-	-	-	-	1	-	-	1
1999	8	-	-	-	-	-	-	-	-	-	-	-	-	1
Diseases of appendix (K35-K38)														
All origins														
Both sexes														
2002	480	3	-	3	4	7	5	4	6	8	19	24	30	28
2001	428	1	8	3	3	5	10	7	8	9	17	25	15	22
2000	435	3	5	4	11	5	6	3	8	11	16	18	25	28
1999	390	3	2	12	14	2	3	3	3	13	11	25	24	22
Male														
2002	275	1	-	1	2	5	2	4	5	7	10	19	20	18
2001	252	-	7	3	2	3	5	5	6	7	10	17	11	13
2000	237	2	2	1	3	2	5	3	7	5	11	12	17	20
1999	236	1	2	6	9	2	3	3	2	10	7	18	19	17
Female														
2002	205	2	-	2	2	2	3	-	1	1	9	5	10	10
2001	176	1	1	-	1	2	5	2	2	2	7	8	4	9
2000	198	1	3	3	8	3	1	-	1	6	5	6	8	8
1999	154	2	-	6	5	-	-	-	1	3	4	7	5	5
Hispanic														
Both sexes														
2002	28	1	-	1	1	1	2	-	1	1	4	-	1	3
2001	30	1	4	-	1	1	1	-	1	1	-	4	1	-
2000	25	1	1	-	4	3	2	-	2	1	-	1	-	4
1999	29	1	-	5	3	-	2	-	-	4	-	2	2	1
Male														
2002	16	-	-	1	1	1	-	-	1	1	1	-	1	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Peptic ulcer (K25-K28)										
Origin not stated[2]										
Both sexes										
2002	2	-	1	1	2	2	1	-	-	-
2001	2	4	-	2	2	3	-	2	-	-
2000	-	1	2	1	3	1	3	1	-	-
1999	2	1	2	3	2	3	-	-	-	-
Male										
2002	1	-	1	1	-	1	1	-	-	-
2001	2	4	-	1	2	-	-	-	-	-
2000	-	1	1	1	1	-	-	-	-	-
1999	2	-	1	2	1	-	-	-	-	-
Female										
2002	1	-	-	-	2	1	-	-	-	-
2001	-	-	-	1	-	3	-	2	-	-
2000	-	-	1	-	2	1	3	1	-	-
1999	-	1	1	1	1	3	-	-	-	-
Diseases of appendix (K35-K38)										
All origins										
Both sexes										
2002	36	32	57	61	59	56	25	10	3	-
2001	23	34	55	37	53	51	30	11	1	-
2000	32	26	45	53	68	39	24	4	1	-
1999	26	30	39	43	53	40	18	3	1	-
Male										
2002	27	24	32	35	32	22	6	1	2	-
2001	17	23	34	22	30	20	12	4	1	-
2000	23	15	25	30	30	15	9	-	-	-
1999	17	18	20	28	19	23	9	2	1	-
Female										
2002	9	8	25	26	27	34	19	9	1	-
2001	6	11	21	15	23	31	18	7	-	-
2000	9	11	20	23	38	24	15	4	1	-
1999	9	12	19	15	34	17	9	1	-	-
Hispanic										
Both sexes										
2002	3	1	2	3	1	1	1	-	-	-
2001	2	2	3	3	4	1	-	-	-	-
2000	1	-	2	1	2	-	-	-	-	-
1999	2	2	1	2	1	1	-	-	-	-
Male										
2002	3	1	1	2	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of appendix (K35-K38)														
Hispanic														
Male														
2001	16	-	3	-	1	-	-	-	-	-	-	2	1	-
2000	16	-	1	-	3	1	2	-	2	-	-	1	-	4
1999	20	1	-	2	2	-	2	-	-	4	-	1	2	1
Female														
2002	12	1	-	-	-	-	2	-	-	-	3	-	-	1
2001	14	1	1	-	-	1	1	-	1	1	-	2	-	-
2000	9	1	-	-	1	2	-	-	-	1	-	-	-	-
1999	9	-	-	3	1	-	-	-	-	-	-	1	-	-
Non-Hispanic[1]														
Both sexes														
2002	451	2	-	2	3	6	3	4	5	7	15	24	29	25
2001	398	-	4	3	2	4	9	7	7	8	17	21	14	22
2000	410	2	4	4	7	2	4	3	6	10	16	17	25	24
1999	358	2	2	7	10	2	1	3	3	9	11	22	22	21
Male														
2002	258	1	-	-	1	4	2	4	4	6	9	19	19	16
2001	236	-	4	3	1	3	5	5	6	7	10	15	10	13
2000	221	2	1	1	-	1	3	3	5	5	11	11	17	16
1999	213	-	2	4	6	2	1	3	2	6	7	16	17	16
Female														
2002	193	1	-	2	2	2	1	-	1	1	6	5	10	9
2001	162	-	-	-	1	1	4	2	1	1	7	6	4	9
2000	189	-	3	3	7	1	1	-	1	5	5	6	8	8
1999	145	2	-	3	4	-	-	-	1	3	4	6	5	5
Non-Hispanic White														
Both sexes														
2002	377	1	-	1	1	3	1	3	4	4	9	17	22	22
2001	313	-	1	2	1	4	4	3	4	6	10	11	13	13
2000	323	2	3	3	6	2	1	1	1	6	11	9	19	17
1999	284	2	-	3	7	1	-	3	1	6	9	13	20	14
Male														
2002	210	-	-	-	-	1	1	3	3	3	8	13	15	15
2001	182	-	1	2	-	3	2	2	3	6	6	8	9	10
2000	167	2	1	1	-	1	1	1	1	3	6	6	12	10
1999	163	-	-	2	4	1	-	3	-	4	5	9	15	11
Female														
2002	167	1	-	1	1	2	-	-	1	1	1	4	7	7
2001	131	-	-	-	1	1	2	1	1	-	4	3	4	3
2000	156	-	2	2	6	1	-	-	-	3	5	3	7	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)										
Hispanic										
Male										
2001	1	2	3	1	2	-	-	-	-	-
2000	-	-	-	1	1	-	-	-	-	-
1999	1	2	-	1	-	1	-	-	-	-
Female										
2002	-	-	1	1	1	1	1	-	-	-
2001	1	-	-	2	2	1	-	-	-	-
2000	1	-	2	-	1	-	-	-	-	-
1999	1	-	1	1	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	32	31	55	58	58	55	24	10	3	-
2001	21	32	52	34	49	50	30	11	1	-
2000	31	26	43	52	66	39	24	4	1	-
1999	24	28	37	41	52	39	18	3	1	-
Male										
2002	23	23	31	33	32	22	6	1	2	-
2001	16	21	31	21	28	20	12	4	1	-
2000	23	15	25	29	29	15	9	-	-	-
1999	16	16	19	27	19	22	9	2	1	-
Female										
2002	9	8	24	25	26	33	18	9	1	-
2001	5	11	21	13	21	30	18	7	-	-
2000	8	11	18	23	37	24	15	4	1	-
1999	8	12	18	14	33	17	9	1	-	-
Non-Hispanic White										
Both sexes										
2002	28	25	44	53	55	51	21	10	2	-
2001	14	26	49	30	40	43	28	10	1	-
2000	28	20	32	41	56	38	22	4	1	-
1999	13	23	31	36	45	36	17	3	1	-
Male										
2002	20	18	25	30	30	19	4	1	1	-
2001	11	17	30	19	24	14	11	3	1	-
2000	20	13	18	25	24	14	8	-	-	-
1999	11	13	15	23	15	20	9	2	1	-
Female										
2002	8	7	19	23	25	32	17	9	1	-
2001	3	9	19	11	16	29	17	7	-	-
2000	8	7	14	16	32	24	14	4	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Diseases of appendix (K35-K38)														
Non-Hispanic White														
Female														
1999	121	2	-	1	3	-	-	-	1	2	4	4	5	3
Non-Hispanic Black														
Both sexes														
2002	61	1	-	1	2	3	2	1	1	-	5	7	6	3
2001	72	-	2	1	1	-	5	3	3	2	6	8	1	9
2000	77	-	-	1	1	-	2	2	5	4	5	8	4	7
1999	67	-	2	4	3	-	1	-	2	3	2	8	2	7
Male														
2002	40	1	-	-	1	3	1	1	1	-	1	6	4	1
2001	47	-	2	1	1	-	3	3	3	1	4	6	1	3
2000	48	-	-	-	-	-	1	2	4	2	5	5	3	6
1999	47	-	2	2	2	-	1	-	2	2	2	7	2	5
Female														
2002	21	-	-	1	1	-	1	-	-	-	4	1	2	2
2001	25	-	-	-	-	-	2	-	-	1	2	2	-	6
2000	29	-	-	1	1	-	1	-	1	2	-	3	1	1
1999	20	-	-	2	1	-	-	-	-	1	-	1	-	2
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	1	-	-	-	-	-	-	1	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	1	-	-	-	-	-	-	1	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Hernia (K40-K46)														
All origins														
Both sexes														
2002	1,595	29	3	4	3	2	3	4	7	14	21	33	44	55
2001	1,522	43	2	1	2	2	1	6	9	13	21	24	58	57

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of appendix (K35-K38)										
Non-Hispanic White										
Female										
1999	2	10	16	13	30	16	8	1	-	-
Non-Hispanic Black										
Both sexes										
2002	3	5	8	3	2	4	3	-	1	-
2001	7	6	1	3	8	3	2	1	-	-
2000	3	6	10	9	7	1	2	-	-	-
1999	10	3	6	5	6	2	1	-	-	-
Male										
2002	2	4	4	2	2	3	2	-	1	-
2001	5	4	-	2	3	3	1	1	-	-
2000	3	2	7	3	3	1	1	-	-	-
1999	4	2	4	4	4	2	-	-	-	-
Female										
2002	1	1	4	1	-	1	1	-	-	-
2001	2	2	1	1	5	-	1	-	-	-
2000	-	4	3	6	4	-	1	-	-	-
1999	6	1	2	1	2	-	1	-	-	-
Origin not stated[2]										
Both sexes										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Male										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hernia (K40-K46)										
All origins										
Both sexes										
2002	71	90	135	208	281	320	172	85	11	-
2001	74	81	132	185	258	269	190	88	6	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hernia (K40-K46)														
All origins														
Both sexes														
2000	1,522	46	4	1	2	2	3	2	3	13	18	26	58	51
1999	1,464	44	3	1	1	1	2	6	10	15	24	16	39	36
Male														
2002	599	18	3	3	2	1	3	2	5	8	9	16	19	27
2001	587	28	1	-	-	1	-	4	6	5	11	13	31	23
2000	586	28	3	1	2	1	2	-	1	8	5	13	32	20
1999	597	26	2	1	-	1	2	5	3	10	13	12	20	21
Female														
2002	996	11	-	1	1	1	-	2	2	6	12	17	25	28
2001	935	15	1	1	2	1	1	2	3	8	10	11	27	34
2000	936	18	1	-	-	1	1	2	2	5	13	13	26	31
1999	867	18	1	-	1	-	-	1	7	5	11	4	19	15
Hispanic														
Both sexes														
2002	81	3	-	1	1	-	2	-	1	1	4	2	4	5
2001	92	9	-	-	-	1	1	1	-	3	4	3	5	3
2000	85	9	-	-	1	1	-	-	1	1	-	2	5	4
1999	66	3	1	-	-	-	-	-	-	2	1	1	3	-
Male														
2002	35	2	-	1	1	-	2	-	1	-	2	2	2	2
2001	32	7	-	-	-	1	-	-	-	1	2	1	2	-
2000	33	6	-	-	1	-	-	-	1	1	-	2	3	3
1999	27	2	1	-	-	-	-	-	-	2	-	1	2	-
Female														
2002	46	1	-	-	-	-	-	-	-	1	2	-	2	3
2001	60	2	-	-	-	-	1	1	-	2	2	2	3	3
2000	52	3	-	-	-	1	-	-	-	-	-	-	2	1
1999	39	1	-	-	-	-	-	-	-	-	1	-	1	-
Non-Hispanic[1]														
Both sexes														
2002	1,504	26	3	3	2	2	1	4	6	12	17	31	40	50
2001	1,425	33	2	1	2	1	-	5	9	10	17	21	53	54
2000	1,431	37	4	1	1	1	3	2	2	12	18	23	53	47
1999	1,390	41	2	1	1	1	2	6	10	13	23	14	36	36
Male														
2002	558	16	3	2	1	1	1	2	4	7	7	14	17	25
2001	552	20	1	-	-	-	-	4	6	4	9	12	29	23
2000	550	22	3	1	1	1	2	-	-	7	5	11	29	17
1999	566	24	1	1	-	1	2	5	3	8	13	10	18	21

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)										
All origins										
Both sexes										
2000	62	84	126	224	242	294	188	63	10	-
1999	71	77	138	186	265	269	196	56	8	-
Male										
2002	29	40	63	84	105	97	45	20	-	-
2001	41	40	54	71	102	88	48	19	1	-
2000	29	39	53	91	104	96	42	12	4	-
1999	39	34	59	75	117	96	51	8	2	-
Female										
2002	42	50	72	124	176	223	127	65	11	-
2001	33	41	78	114	156	181	142	69	5	-
2000	33	45	73	133	138	198	146	51	6	-
1999	32	43	79	111	148	173	145	48	6	-
Hispanic										
Both sexes										
2002	9	6	8	6	13	7	6	2	-	-
2001	6	3	9	12	12	13	6	-	1	-
2000	5	7	7	15	8	10	8	1	-	-
1999	5	7	9	6	11	8	8	1	-	-
Male										
2002	5	2	2	2	5	1	3	-	-	-
2001	3	2	2	1	5	2	3	-	-	-
2000	-	-	3	6	4	2	1	-	-	-
1999	4	1	5	2	4	1	2	-	-	-
Female										
2002	4	4	6	4	8	6	3	2	-	-
2001	3	1	7	11	7	11	3	-	1	-
2000	5	7	4	9	4	8	7	1	-	-
1999	1	6	4	4	7	7	6	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	61	82	127	199	266	312	166	83	11	-
2001	67	78	123	172	245	256	183	88	5	-
2000	56	77	117	209	233	284	179	62	10	-
1999	66	70	128	177	254	260	187	54	8	-
Male										
2002	24	37	61	80	98	96	42	20	-	-
2001	37	38	52	69	97	86	45	19	1	-
2000	28	39	49	85	99	94	41	12	4	-
1999	35	33	53	72	113	94	49	8	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hernia (K40-K46)														
Non-Hispanic[1]														
Female														
2002	946	10	-	1	1	1	-	2	2	5	10	17	23	25
2001	873	13	1	1	2	1	-	1	3	6	8	9	24	31
2000	881	15	1	-	-	-	1	2	2	5	13	12	24	30
1999	824	17	1	-	1	-	-	1	7	5	10	4	18	15
Non-Hispanic White														
Both sexes														
2002	1,343	21	2	3	2	2	1	2	3	9	11	22	30	42
2001	1,248	22	2	-	1	-	-	5	7	7	11	17	42	42
2000	1,266	27	2	1	1	1	1	2	1	8	13	17	42	41
1999	1,227	31	2	1	1	1	1	4	6	12	15	9	27	32
Male														
2002	493	12	2	2	1	1	1	-	2	5	6	11	15	22
2001	469	11	1	-	-	-	-	4	4	4	6	8	21	16
2000	486	17	2	1	1	1	-	-	-	5	4	7	22	15
1999	505	18	1	1	-	1	1	3	2	7	11	7	15	21
Female														
2002	850	9	-	1	1	1	-	2	1	4	5	11	15	20
2001	779	11	1	-	1	-	-	1	3	3	5	9	21	26
2000	780	10	-	-	-	-	1	2	1	3	9	10	20	26
1999	722	13	1	-	1	-	-	1	4	5	4	2	12	11
Non-Hispanic Black														
Both sexes														
2002	137	4	1	-	-	-	-	2	1	3	5	9	8	8
2001	155	10	-	1	-	1	-	-	2	3	6	4	11	12
2000	142	7	2	-	-	-	2	-	1	3	5	5	8	5
1999	139	9	-	-	-	-	1	-	4	1	8	5	7	4
Male														
2002	56	3	1	-	-	-	-	2	-	2	1	3	2	3
2001	74	8	-	-	-	-	-	-	2	-	3	4	8	7
2000	52	3	1	-	-	-	2	-	-	1	1	4	5	1
1999	53	6	-	-	-	-	1	-	1	1	2	3	3	-
Female														
2002	81	1	-	-	-	-	-	-	1	1	4	6	6	5
2001	81	2	-	1	-	1	-	-	-	3	3	-	3	5
2000	90	4	1	-	-	-	-	-	1	2	4	1	3	4
1999	86	3	-	-	-	-	-	-	3	-	6	2	4	4
Origin not stated[2]														
Both sexes														
2002	10	-	-	-	-	-	-	-	-	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)										
Non-Hispanic[1]										
Female										
2002	37	45	66	119	168	216	124	63	11	-
2001	30	40	71	103	148	170	138	69	4	-
2000	28	38	68	124	134	190	138	50	6	-
1999	31	37	75	105	141	166	138	46	6	-
Non-Hispanic White										
Both sexes										
2002	46	63	113	188	243	288	161	81	10	-
2001	55	66	107	151	223	238	172	77	3	-
2000	49	61	108	184	215	260	167	56	9	-
1999	57	53	114	156	234	239	174	52	6	-
Male										
2002	19	28	53	76	91	88	39	19	-	-
2001	30	34	47	62	87	73	42	18	1	-
2000	25	31	47	79	91	86	38	10	4	-
1999	32	28	51	66	105	84	42	7	2	-
Female										
2002	27	35	60	112	152	200	122	62	10	-
2001	25	32	60	89	136	165	130	59	2	-
2000	24	30	61	105	124	174	129	46	5	-
1999	25	25	63	90	129	155	132	45	4	-
Non-Hispanic Black										
Both sexes										
2002	13	15	12	10	20	19	4	2	1	-
2001	11	10	15	17	16	16	9	9	2	-
2000	7	14	7	21	15	24	12	3	1	-
1999	8	12	12	19	18	15	12	2	2	-
Male										
2002	4	8	7	4	7	6	2	1	-	-
2001	6	2	5	6	8	11	3	1	-	-
2000	3	7	2	4	7	8	3	-	-	-
1999	2	5	2	5	8	7	6	1	-	-
Female										
2002	9	7	5	6	13	13	2	1	1	-
2001	5	8	10	11	8	5	6	8	2	-
2000	4	7	5	17	8	16	9	3	1	-
1999	6	7	10	14	10	8	6	1	2	-
Origin not stated[2]										
Both sexes										
2002	1	2	-	3	2	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hernia (K40-K46)														
Origin not stated[2]														
Both sexes														
2001	5	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	6	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	8	-	-	-	-	-	-	-	-	-	-	1	-	-
Male														
2002	6	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	3	1	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	1	-	-
Female														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	1	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Chronic liver disease and cirrhosis (K70,K73-K74)														
All origins														
Both sexes														
2002	27,257	6	3	1	5	7	24	77	297	929	2,225	3,404	3,812	3,258
2001	27,035	3	-	1	1	6	20	77	310	1,108	2,228	3,557	3,702	2,999
2000	26,552	3	3	-	3	7	29	88	327	1,083	2,288	3,374	3,280	2,947
1999	26,259	5	2	1	1	9	17	70	337	1,103	2,199	3,228	3,140	2,803
Male														
2002	17,401	6	1	1	2	3	11	46	190	605	1,474	2,442	2,843	2,323
2001	17,393	1	-	1	1	2	10	53	206	713	1,493	2,581	2,714	2,161
2000	17,214	3	1	-	2	3	16	63	217	716	1,598	2,515	2,460	2,066
1999	17,115	4	-	1	-	3	8	41	216	716	1,598	2,446	2,383	2,051
Female														
2002	9,856	-	2	-	3	4	13	31	107	324	751	962	969	935
2001	9,642	2	-	-	-	4	10	24	104	395	735	976	988	838
2000	9,338	-	2	-	1	4	13	25	110	367	690	859	820	881
1999	9,144	1	2	-	1	6	9	29	121	387	601	782	757	752
Hispanic														
Both sexes														
2002	3,409	1	1	1	1	2	7	18	63	162	345	478	505	422
2001	3,301	-	-	-	-	1	4	16	65	187	343	480	471	378
2000	3,187	-	1	-	-	-	7	16	76	170	331	440	414	380
1999	2,972	-	-	-	-	4	4	7	73	190	328	423	364	325
Male														
2002	2,437	1	-	1	-	-	5	13	49	127	283	397	408	333

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hernia (K40-K46)										
Origin not stated[2]										
Both sexes										
2001	1	-	-	1	1	-	1	-	-	-
2000	1	-	2	-	1	-	1	-	-	-
1999	-	-	1	3	-	1	1	1	-	-
Male										
2002	-	1	-	2	2	-	-	-	-	-
2001	1	-	-	1	-	-	-	-	-	-
2000	1	-	1	-	1	-	-	-	-	-
1999	-	-	1	1	-	1	-	-	-	-
Female										
2002	1	1	-	1	-	1	-	-	-	-
2001	-	-	-	-	1	-	1	-	-	-
2000	-	-	1	-	-	-	1	-	-	-
1999	-	-	-	2	-	-	1	1	-	-
Chronic liver disease and cirrhosis (K70,K73-K74)										
All origins										
Both sexes										
2002	2,839	2,670	2,711	2,412	1,588	728	230	24	3	4
2001	2,751	2,752	2,734	2,325	1,476	714	221	43	1	6
2000	2,827	2,751	2,731	2,367	1,462	742	203	30	4	3
1999	2,834	2,833	2,809	2,396	1,505	702	226	33	4	2
Male										
2002	1,905	1,652	1,545	1,211	742	310	77	8	-	4
2001	1,867	1,749	1,586	1,151	682	307	89	19	1	6
2000	1,939	1,729	1,588	1,240	634	325	83	12	1	3
1999	1,892	1,837	1,617	1,208	697	293	86	14	2	2
Female										
2002	934	1,018	1,166	1,201	846	418	153	16	3	-
2001	884	1,003	1,148	1,174	794	407	132	24	-	-
2000	888	1,022	1,143	1,127	828	417	120	18	3	-
1999	942	996	1,192	1,188	808	409	140	19	2	-
Hispanic										
Both sexes										
2002	336	332	331	198	123	58	21	3	-	1
2001	343	322	296	219	116	46	12	1	-	1
2000	328	333	295	215	108	51	19	2	1	-
1999	313	310	273	187	98	51	20	1	1	-
Male										
2002	228	226	181	95	62	18	8	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic liver disease and cirrhosis (K70,K73-K74)														
Hispanic														
Male														
2001	2,410	-	-	-	-	-	2	15	52	150	275	403	379	295
2000	2,312	-	-	-	-	-	5	14	68	134	265	365	342	295
1999	2,143	-	-	-	-	3	2	6	56	154	267	354	293	257
Female														
2002	972	-	1	-	1	2	2	5	14	35	62	81	97	89
2001	891	-	-	-	-	1	2	1	13	37	68	77	92	83
2000	875	-	1	-	-	-	2	2	8	36	66	75	72	85
1999	829	-	-	-	-	1	2	1	17	36	61	69	71	68
Non-Hispanic[1]														
Both sexes														
2002	23,733	5	2	-	4	5	17	58	233	760	1,872	2,907	3,291	2,819
2001	23,629	3	-	-	1	5	16	61	244	919	1,877	3,064	3,216	2,607
2000	23,251	3	2	-	3	7	22	72	247	909	1,947	2,916	2,846	2,557
1999	23,185	5	2	1	1	5	13	63	263	909	1,858	2,791	2,756	2,467
Male														
2002	14,876	5	1	-	2	3	6	32	140	474	1,184	2,029	2,423	1,977
2001	14,914	1	-	-	1	2	8	38	154	563	1,214	2,170	2,323	1,854
2000	14,814	3	1	-	2	3	11	49	146	578	1,325	2,137	2,100	1,762
1999	14,896	4	-	1	-	-	6	35	159	560	1,321	2,080	2,075	1,784
Female														
2002	8,857	-	1	-	2	2	11	26	93	286	688	878	868	842
2001	8,715	2	-	-	-	3	8	23	90	356	663	894	893	753
2000	8,437	-	1	-	1	4	11	23	101	331	622	779	746	795
1999	8,289	1	2	-	1	5	7	28	104	349	537	711	681	683
Non-Hispanic White														
Both sexes														
2002	20,353	4	2	-	3	5	12	37	167	603	1,545	2,353	2,670	2,364
2001	20,065	3	-	-	1	1	4	35	192	733	1,479	2,478	2,557	2,127
2000	19,739	-	2	-	2	2	12	42	175	702	1,538	2,365	2,272	2,121
1999	19,602	4	-	-	1	3	4	45	198	712	1,436	2,170	2,260	2,013
Male														
2002	12,800	4	1	-	2	3	5	21	109	376	995	1,682	1,988	1,674
2001	12,609	1	-	-	1	-	4	23	122	443	960	1,780	1,865	1,511
2000	12,640	-	1	-	1	1	5	33	107	453	1,065	1,785	1,692	1,480
1999	12,613	4	-	-	-	-	3	29	122	445	1,033	1,647	1,718	1,467
Female														
2002	7,553	-	1	-	1	2	7	16	58	227	550	671	682	690
2001	7,456	2	-	-	-	1	-	12	70	290	519	698	692	616

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)										
Hispanic										
Male										
2001	260	220	166	120	50	16	6	-	-	1
2000	236	233	162	121	42	17	12	1	-	-
1999	224	198	162	93	43	20	10	-	1	-
Female										
2002	108	106	150	103	61	40	13	2	-	-
2001	83	102	130	99	66	30	6	1	-	-
2000	92	100	133	94	66	34	7	1	1	-
1999	89	112	111	94	55	31	10	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	2,488	2,329	2,371	2,207	1,461	669	209	21	3	2
2001	2,395	2,417	2,429	2,099	1,356	667	209	40	1	3
2000	2,484	2,405	2,428	2,148	1,350	691	181	28	3	2
1999	2,511	2,514	2,530	2,201	1,405	649	204	32	3	2
Male										
2002	1,664	1,420	1,358	1,110	678	292	69	7	-	2
2001	1,596	1,518	1,416	1,030	631	290	83	18	1	3
2000	1,690	1,486	1,422	1,117	591	308	69	11	1	2
1999	1,660	1,631	1,450	1,111	654	272	76	14	1	2
Female										
2002	824	909	1,013	1,097	783	377	140	14	3	-
2001	799	899	1,013	1,069	725	377	126	22	-	-
2000	794	919	1,006	1,031	759	383	112	17	2	-
1999	851	883	1,080	1,090	751	377	128	18	2	-
Non-Hispanic White										
Both sexes										
2002	2,119	2,047	2,158	2,061	1,356	628	194	20	3	2
2001	2,062	2,112	2,195	1,945	1,277	628	196	37	1	2
2000	2,114	2,093	2,192	2,008	1,256	646	166	27	3	1
1999	2,104	2,191	2,291	2,027	1,320	602	187	29	3	2
Male										
2002	1,423	1,246	1,243	1,050	631	275	63	7	-	2
2001	1,364	1,318	1,288	956	601	276	76	17	1	2
2000	1,455	1,314	1,288	1,040	557	288	62	11	1	1
1999	1,418	1,428	1,318	1,021	620	255	69	13	1	2
Female										
2002	696	801	915	1,011	725	353	131	13	3	-
2001	698	794	907	989	676	352	120	20	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic liver disease and cirrhosis (K70,K73-K74)														
Non-Hispanic White														
Female														
2000	7,099	-	1	-	1	1	7	9	68	249	473	580	580	641
1999	6,989	-	-	-	1	3	1	16	76	267	403	523	542	546
Non-Hispanic Black														
Both sexes														
2002	2,545	1	-	-	1	-	5	12	37	84	229	440	516	360
2001	2,733	-	-	-	-	2	8	18	27	117	303	463	530	390
2000	2,705	2	-	-	1	4	8	20	36	142	304	452	477	341
1999	2,793	-	2	1	-	2	8	16	38	129	324	507	398	363
Male														
2002	1,592	1	-	-	-	-	1	7	19	49	131	274	363	243
2001	1,819	-	-	-	-	1	1	10	15	77	196	319	383	278
2000	1,709	2	-	-	1	1	5	11	15	90	189	289	344	230
1999	1,812	-	-	1	-	-	3	5	22	75	218	357	292	254
Female														
2002	953	-	-	-	1	-	4	5	18	35	98	166	153	117
2001	914	-	-	-	-	1	7	8	12	40	107	144	147	112
2000	996	-	-	-	-	3	3	9	21	52	115	163	133	111
1999	981	-	2	-	-	2	5	11	16	54	106	150	106	109
Origin not stated[2]														
Both sexes														
2002	115	-	-	-	-	-	-	1	1	7	8	19	16	17
2001	105	-	-	1	-	-	-	-	1	2	8	13	15	14
2000	114	-	-	-	-	-	-	-	4	4	10	18	20	10
1999	102	-	-	-	-	-	-	-	1	4	13	14	20	11
Male														
2002	88	-	-	-	-	-	-	1	1	4	7	16	12	13
2001	69	-	-	1	-	-	-	-	-	-	4	8	12	12
2000	88	-	-	-	-	-	-	-	3	4	8	13	18	9
1999	76	-	-	-	-	-	-	-	1	2	10	12	15	10
Female														
2002	27	-	-	-	-	-	-	-	-	3	1	3	4	4
2001	36	-	-	-	-	-	-	-	1	2	4	5	3	2
2000	26	-	-	-	-	-	-	-	1	-	2	5	2	1
1999	26	-	-	-	-	-	-	-	-	2	3	2	5	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic liver disease and cirrhosis (K70,K73-K74)										
Non-Hispanic White										
Female										
2000	659	779	904	968	699	358	104	16	2	-
1999	686	763	973	1,006	700	347	118	16	2	-
Non-Hispanic Black										
Both sexes										
2002	289	213	145	112	61	28	12	-	-	-
2001	259	245	167	116	53	22	10	2	-	1
2000	281	246	183	101	62	30	13	1	-	1
1999	328	250	182	128	65	37	13	2	-	-
Male										
2002	193	137	84	47	28	11	4	-	-	-
2001	184	168	100	57	18	7	4	-	-	1
2000	178	144	111	58	23	12	5	-	-	1
1999	199	166	107	67	29	13	4	-	-	-
Female										
2002	96	76	61	65	33	17	8	-	-	-
2001	75	77	67	59	35	15	6	2	-	-
2000	103	102	72	43	39	18	8	1	-	-
1999	129	84	75	61	36	24	9	2	-	-
Origin not stated[2]										
Both sexes										
2002	15	9	9	7	4	1	-	-	-	1
2001	13	13	9	7	4	1	-	2	-	2
2000	15	13	8	4	4	-	3	-	-	1
1999	10	9	6	8	2	2	2	-	-	-
Male										
2002	13	6	6	6	2	-	-	-	-	1
2001	11	11	4	1	1	1	-	1	-	2
2000	13	10	4	2	1	-	2	-	-	1
1999	8	8	5	4	-	1	-	-	-	-
Female										
2002	2	3	3	1	2	1	-	-	-	-
2001	2	2	5	6	3	-	-	1	-	-
2000	2	3	4	2	3	-	1	-	-	-
1999	2	1	1	4	2	1	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcoholic liver disease (K70)														
All origins														
Both sexes														
2002	12,121	-	1	-	-	-	8	43	199	607	1,364	1,998	2,139	1,712
2001	12,207	-	-	-	-	1	7	45	219	745	1,400	2,121	2,082	1,582
2000	12,109	-	-	-	-	-	8	56	223	741	1,485	2,000	1,888	1,575
1999	11,958	-	-	-	-	-	8	37	222	763	1,423	1,965	1,837	1,511
Male														
2002	8,814	-	1	-	-	-	6	32	130	392	913	1,450	1,624	1,277
2001	8,932	-	-	-	-	-	5	31	148	492	954	1,563	1,536	1,235
2000	8,985	-	-	-	-	-	5	45	157	486	1,047	1,510	1,458	1,172
1999	8,922	-	-	-	-	-	6	24	141	491	1,029	1,501	1,426	1,179
Female														
2002	3,307	-	-	-	-	-	2	11	69	215	451	548	515	435
2001	3,275	-	-	-	-	1	2	14	71	253	446	558	546	347
2000	3,124	-	-	-	-	-	3	11	66	255	438	490	430	403
1999	3,036	-	-	-	-	-	2	13	81	272	394	464	411	332
Hispanic														
Both sexes														
2002	1,679	-	-	-	-	-	4	12	41	112	209	282	305	229
2001	1,729	-	-	-	-	-	-	14	51	123	218	308	274	220
2000	1,680	-	-	-	-	-	4	13	53	122	231	273	253	218
1999	1,547	-	-	-	-	-	2	4	46	128	214	270	229	196
Male														
2002	1,418	-	-	-	-	-	4	12	35	94	177	236	257	200
2001	1,457	-	-	-	-	-	-	13	42	107	175	265	229	189
2000	1,442	-	-	-	-	-	4	12	50	94	188	237	218	191
1999	1,294	-	-	-	-	-	2	4	37	101	175	232	194	174
Female														
2002	261	-	-	-	-	-	-	-	6	18	32	46	48	29
2001	272	-	-	-	-	-	-	1	9	16	43	43	45	31
2000	238	-	-	-	-	-	-	1	3	28	43	36	35	27
1999	253	-	-	-	-	-	-	-	9	27	39	38	35	22
Non-Hispanic[1]														
Both sexes														
2002	10,377	-	1	-	-	-	4	31	157	490	1,149	1,703	1,824	1,469
2001	10,423	-	-	-	-	1	7	31	167	621	1,180	1,806	1,797	1,350
2000	10,370	-	-	-	-	-	4	43	168	616	1,249	1,713	1,624	1,351
1999	10,353	-	-	-	-	-	6	33	175	631	1,200	1,686	1,597	1,309
Male														
2002	7,342	-	1	-	-	-	2	20	94	296	731	1,203	1,359	1,066
2001	7,431	-	-	-	-	-	5	18	106	385	777	1,294	1,298	1,036
2000	7,492	-	-	-	-	-	1	33	106	389	854	1,262	1,230	976

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)										
All origins										
Both sexes										
2002	1,371	1,023	826	495	235	79	16	3	-	2
2001	1,331	1,062	786	515	202	86	12	6	-	5
2000	1,394	1,085	806	533	218	76	17	2	-	2
1999	1,348	1,104	831	533	256	94	21	3	-	2
Male										
2002	1,060	771	596	334	161	54	10	1	-	2
2001	1,007	789	590	370	135	60	8	4	-	5
2000	1,044	821	612	419	142	52	12	1	-	2
1999	1,015	867	630	364	174	59	13	1	-	2
Female										
2002	311	252	230	161	74	25	6	2	-	-
2001	324	273	196	145	67	26	4	2	-	-
2000	350	264	194	114	76	24	5	1	-	-
1999	333	237	201	169	82	35	8	2	-	-
Hispanic										
Both sexes										
2002	163	139	107	44	19	8	3	1	-	1
2001	176	140	100	68	26	8	2	-	-	1
2000	182	153	94	62	16	4	1	1	-	-
1999	160	124	89	51	18	14	2	-	-	-
Male										
2002	139	122	89	30	14	4	3	1	-	1
2001	149	124	86	50	18	7	2	-	-	1
2000	153	139	82	56	13	3	1	1	-	-
1999	133	103	77	38	14	10	-	-	-	-
Female										
2002	24	17	18	14	5	4	-	-	-	-
2001	27	16	14	18	8	1	-	-	-	-
2000	29	14	12	6	3	1	-	-	-	-
1999	27	21	12	13	4	4	2	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,202	879	718	448	215	71	13	2	-	1
2001	1,149	914	682	446	176	78	10	6	-	2
2000	1,203	927	712	469	202	72	15	1	-	1
1999	1,182	975	738	479	238	80	19	3	-	2
Male										
2002	915	644	506	301	146	50	7	-	-	1
2001	852	658	500	320	117	53	6	4	-	2
2000	883	678	530	361	129	49	10	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcoholic liver disease (K70)														
Non-Hispanic[1]														
Male														
1999	7,580	-	-	-	-	-	4	20	103	388	847	1,261	1,224	1,000
Female														
2002	3,035	-	-	-	-	-	2	11	63	194	418	500	465	403
2001	2,992	-	-	-	-	1	2	13	61	236	403	512	499	314
2000	2,878	-	-	-	-	-	3	10	62	227	395	451	394	375
1999	2,773	-	-	-	-	-	2	13	72	243	353	425	373	309
Non-Hispanic White														
Both sexes														
2002	8,713	-	1	-	-	-	1	19	114	386	949	1,408	1,499	1,230
2001	8,635	-	-	-	-	-	2	18	128	491	940	1,504	1,431	1,105
2000	8,556	-	-	-	-	-	2	29	121	474	980	1,388	1,315	1,109
1999	8,470	-	-	-	-	-	3	27	128	492	922	1,335	1,314	1,065
Male														
2002	6,242	-	1	-	-	-	1	13	74	231	607	1,012	1,131	915
2001	6,209	-	-	-	-	-	2	13	83	302	620	1,098	1,054	849
2000	6,296	-	-	-	-	-	-	24	81	303	676	1,056	1,012	821
1999	6,288	-	-	-	-	-	2	17	78	307	660	1,021	1,020	819
Female														
2002	2,471	-	-	-	-	-	-	6	40	155	342	396	368	315
2001	2,426	-	-	-	-	-	-	5	45	189	320	406	377	256
2000	2,260	-	-	-	-	-	2	5	40	171	304	332	303	288
1999	2,182	-	-	-	-	-	1	10	50	185	262	314	294	246
Non-Hispanic Black														
Both sexes														
2002	1,173	-	-	-	-	-	3	6	20	46	123	209	252	180
2001	1,331	-	-	-	-	-	1	7	19	74	170	218	290	198
2000	1,370	-	-	-	-	-	-	7	27	92	193	258	247	184
1999	1,445	-	-	-	-	-	2	4	24	84	197	285	222	192
Male														
2002	783	-	-	-	-	-	1	4	10	27	76	136	178	112
2001	926	-	-	-	-	-	-	2	10	50	110	147	202	149
2000	910	-	-	-	-	-	-	5	12	59	126	164	175	121
1999	1,001	-	-	-	-	-	2	2	12	49	129	197	161	145
Female														
2002	390	-	-	-	-	-	2	2	10	19	47	73	74	68
2001	405	-	-	-	-	-	1	5	9	24	60	71	88	49
2000	460	-	-	-	-	-	-	2	15	33	67	94	72	63
1999	444	-	-	-	-	-	-	2	12	35	68	88	61	47

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)										
Non-Hispanic[1]										
Male										
1999	877	759	549	323	160	49	13	1	-	2
Female										
2002	287	235	212	147	69	21	6	2	-	-
2001	297	256	182	126	59	25	4	2	-	-
2000	320	249	182	108	73	23	5	1	-	-
1999	305	216	189	156	78	31	6	2	-	-
Non-Hispanic White										
Both sexes										
2002	1,025	762	641	406	195	63	11	2	-	1
2001	987	773	608	404	162	66	8	6	-	2
2000	1,017	805	634	423	178	66	14	1	-	-
1999	980	829	651	421	215	67	16	3	-	2
Male										
2002	786	560	452	275	134	44	5	-	-	1
2001	734	553	444	293	108	46	4	4	-	2
2000	758	594	471	329	117	45	9	-	-	-
1999	747	645	486	284	147	40	12	1	-	2
Female										
2002	239	202	189	131	61	19	6	2	-	-
2001	253	220	164	111	54	20	4	2	-	-
2000	259	211	163	94	61	21	5	1	-	-
1999	233	184	165	137	68	27	4	2	-	-
Non-Hispanic Black										
Both sexes										
2002	135	88	55	33	15	7	1	-	-	-
2001	126	118	58	32	11	8	1	-	-	-
2000	137	106	63	35	15	4	1	-	-	1
1999	166	117	74	45	19	11	3	-	-	-
Male										
2002	100	64	39	21	9	5	1	-	-	-
2001	90	90	45	19	7	4	1	-	-	-
2000	87	73	50	24	9	3	1	-	-	1
1999	109	93	53	30	11	7	1	-	-	-
Female										
2002	35	24	16	12	6	2	-	-	-	-
2001	36	28	13	13	4	4	-	-	-	-
2000	50	33	13	11	6	1	-	-	-	-
1999	57	24	21	15	8	4	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcoholic liver disease (K70)														
Origin not stated[2]														
Both sexes														
2002	65	-	-	-	-	-	-	-	1	5	6	13	10	14
2001	55	-	-	-	-	-	-	-	1	1	2	7	11	12
2000	59	-	-	-	-	-	-	-	2	3	5	14	11	6
1999	58	-	-	-	-	-	-	-	1	4	9	9	11	6
Male														
2002	54	-	-	-	-	-	-	-	1	2	5	11	8	11
2001	44	-	-	-	-	-	-	-	-	-	2	4	9	10
2000	51	-	-	-	-	-	-	-	1	3	5	11	10	5
1999	48	-	-	-	-	-	-	-	1	2	7	8	8	5
Female														
2002	11	-	-	-	-	-	-	-	-	3	1	2	2	3
2001	11	-	-	-	-	-	-	-	1	1	-	3	2	2
2000	8	-	-	-	-	-	-	-	1	-	-	3	1	1
1999	10	-	-	-	-	-	-	-	-	2	2	1	3	1
Other chronic liver disease and cirrhosis (K73-K74)														
All origins														
Both sexes														
2002	15,136	6	2	1	5	7	16	34	98	322	861	1,406	1,673	1,546
2001	14,828	3	-	1	1	5	13	32	91	363	828	1,436	1,620	1,417
2000	14,443	3	3	-	3	7	21	32	104	342	803	1,374	1,392	1,372
1999	14,301	5	2	1	1	9	9	33	115	340	776	1,263	1,303	1,292
Male														
2002	8,587	6	-	1	2	3	5	14	60	213	561	992	1,219	1,046
2001	8,461	1	-	1	1	2	5	22	58	221	539	1,018	1,178	926
2000	8,229	3	1	-	2	3	11	18	60	230	551	1,005	1,002	894
1999	8,193	4	-	1	-	3	2	17	75	225	569	945	957	872
Female														
2002	6,549	-	2	-	3	4	11	20	38	109	300	414	454	500
2001	6,367	2	-	-	-	3	8	10	33	142	289	418	442	491
2000	6,214	-	2	-	1	4	10	14	44	112	252	369	390	478
1999	6,108	1	2	-	1	6	7	16	40	115	207	318	346	420
Hispanic														
Both sexes														
2002	1,730	1	1	1	1	2	3	6	22	50	136	196	200	193
2001	1,572	-	-	-	-	1	4	2	14	64	125	172	197	158
2000	1,507	-	1	-	-	-	3	3	23	48	100	167	161	162
1999	1,425	-	-	-	-	4	2	3	27	62	114	153	135	129

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)										
Origin not stated[2]										
Both sexes										
2002	6	5	1	3	1	-	-	-	-	-
2001	6	8	4	1	-	-	-	-	-	2
2000	9	5	-	2	-	-	1	-	-	1
1999	6	5	4	3	-	-	-	-	-	-
Male										
2002	6	5	1	3	1	-	-	-	-	-
2001	6	7	4	-	-	-	-	-	-	2
2000	8	4	-	2	-	-	1	-	-	1
1999	5	5	4	3	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	-	1	-	-	-	-	-	-
2000	1	1	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-
Other chronic liver disease and cirrhosis (K73-K74)										
All origins										
Both sexes										
2002	1,468	1,647	1,885	1,917	1,353	649	214	21	3	2
2001	1,420	1,690	1,948	1,810	1,274	628	209	37	1	1
2000	1,433	1,666	1,925	1,834	1,244	666	186	28	4	1
1999	1,486	1,729	1,978	1,863	1,249	608	205	30	4	-
Male										
2002	845	881	949	877	581	256	67	7	-	2
2001	860	960	996	781	547	247	81	15	1	1
2000	895	908	976	821	492	273	71	11	1	1
1999	877	970	987	844	523	234	73	13	2	-
Female										
2002	623	766	936	1,040	772	393	147	14	3	-
2001	560	730	952	1,029	727	381	128	22	-	-
2000	538	758	949	1,013	752	393	115	17	3	-
1999	609	759	991	1,019	726	374	132	17	2	-
Hispanic										
Both sexes										
2002	173	193	224	154	104	50	18	2	-	-
2001	167	182	196	151	90	38	10	1	-	-
2000	146	180	201	153	92	47	18	1	1	-
1999	153	186	184	136	80	37	18	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic liver disease and cirrhosis (K73-K74)														
Hispanic														
Male														
2002	1,019	1	-	1	-	-	1	1	14	33	106	161	151	133
2001	953	-	-	-	-	-	2	2	10	43	100	138	150	106
2000	870	-	-	-	-	-	1	2	18	40	77	128	124	104
1999	849	-	-	-	-	3	-	2	19	53	92	122	99	83
Female														
2002	711	-	1	-	1	2	2	5	8	17	30	35	49	60
2001	619	-	-	-	-	1	2	-	4	21	25	34	47	52
2000	637	-	1	-	-	-	2	1	5	8	23	39	37	58
1999	576	-	-	-	-	1	2	1	8	9	22	31	36	46
Non-Hispanic[1]														
Both sexes														
2002	13,356	5	1	-	4	5	13	27	76	270	723	1,204	1,467	1,350
2001	13,206	3	-	-	1	4	9	30	77	298	697	1,258	1,419	1,257
2000	12,881	3	2	-	3	7	18	29	79	293	698	1,203	1,222	1,206
1999	12,832	5	2	1	1	5	7	30	88	278	658	1,105	1,159	1,158
Male														
2002	7,534	5	-	-	2	3	4	12	46	178	453	826	1,064	911
2001	7,483	1	-	-	1	2	3	20	48	178	437	876	1,025	818
2000	7,322	3	1	-	2	3	10	16	40	189	471	875	870	786
1999	7,316	4	-	1	-	-	2	15	56	172	474	819	851	784
Female														
2002	5,822	-	1	-	2	2	9	15	30	92	270	378	403	439
2001	5,723	2	-	-	-	2	6	10	29	120	260	382	394	439
2000	5,559	-	1	-	1	4	8	13	39	104	227	328	352	420
1999	5,516	1	2	-	1	5	5	15	32	106	184	286	308	374
Non-Hispanic White														
Both sexes														
2002	11,640	4	1	-	3	5	11	18	53	217	596	945	1,171	1,134
2001	11,430	3	-	-	1	1	2	17	64	242	539	974	1,126	1,022
2000	11,183	-	2	-	2	2	10	13	54	228	558	977	957	1,012
1999	11,132	4	-	-	1	3	1	18	70	220	514	835	946	948
Male														
2002	6,558	4	-	-	2	3	4	8	35	145	388	670	857	759
2001	6,400	1	-	-	1	-	2	10	39	141	340	682	811	662
2000	6,344	-	1	-	1	1	5	9	26	150	389	729	680	659
1999	6,325	4	-	-	-	-	1	12	44	138	373	626	698	648
Female														
2002	5,082	-	1	-	1	2	7	10	18	72	208	275	314	375

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)										
Hispanic										
Male										
2002	89	104	92	65	48	14	5	-	-	-
2001	111	96	80	70	32	9	4	-	-	-
2000	83	94	80	65	29	14	11	-	-	-
1999	91	95	85	55	29	10	10	-	1	-
Female										
2002	84	89	132	89	56	36	13	2	-	-
2001	56	86	116	81	58	29	6	1	-	-
2000	63	86	121	88	63	33	7	1	1	-
1999	62	91	99	81	51	27	8	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,286	1,450	1,653	1,759	1,246	598	196	19	3	1
2001	1,246	1,503	1,747	1,653	1,180	589	199	34	1	1
2000	1,281	1,478	1,716	1,679	1,148	619	166	27	3	1
1999	1,329	1,539	1,792	1,722	1,167	569	185	29	3	-
Male										
2002	749	776	852	809	532	242	62	7	-	1
2001	744	860	916	710	514	237	77	14	1	1
2000	807	808	892	756	462	259	59	11	1	1
1999	783	872	901	788	494	223	63	13	1	-
Female										
2002	537	674	801	950	714	356	134	12	3	-
2001	502	643	831	943	666	352	122	20	-	-
2000	474	670	824	923	686	360	107	16	2	-
1999	546	667	891	934	673	346	122	16	2	-
Non-Hispanic White										
Both sexes										
2002	1,094	1,285	1,517	1,655	1,161	565	183	18	3	1
2001	1,075	1,339	1,587	1,541	1,115	562	188	31	1	-
2000	1,097	1,288	1,558	1,585	1,078	580	152	26	3	1
1999	1,124	1,362	1,640	1,606	1,105	535	171	26	3	-
Male										
2002	637	686	791	775	497	231	58	7	-	1
2001	630	765	844	663	493	230	72	13	1	-
2000	697	720	817	711	440	243	53	11	1	1
1999	671	783	832	737	473	215	57	12	1	-
Female										
2002	457	599	726	880	664	334	125	11	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other chronic liver disease and cirrhosis (K73-K74)														
Non-Hispanic White														
Female														
2001	5,030	2	-	-	-	1	-	7	25	101	199	292	315	360
2000	4,839	-	1	-	1	1	5	4	28	78	169	248	277	353
1999	4,807	-	-	-	1	3	-	6	26	82	141	209	248	300
Non-Hispanic Black														
Both sexes														
2002	1,372	1	-	-	1	-	2	6	17	38	106	231	264	180
2001	1,402	-	-	-	-	2	7	11	8	43	133	245	240	192
2000	1,335	2	-	-	1	4	8	13	9	50	111	194	230	157
1999	1,348	-	2	1	-	2	6	12	14	45	127	222	176	171
Male														
2002	809	1	-	-	-	-	-	3	9	22	55	138	185	131
2001	893	-	-	-	-	1	1	8	5	27	86	172	181	129
2000	799	2	-	-	1	1	5	6	3	31	63	125	169	109
1999	811	-	-	1	-	-	1	3	10	26	89	160	131	109
Female														
2002	563	-	-	-	1	-	2	3	8	16	51	93	79	49
2001	509	-	-	-	-	1	6	3	3	16	47	73	59	63
2000	536	-	-	-	-	3	3	7	6	19	48	69	61	48
1999	537	-	2	-	-	2	5	9	4	19	38	62	45	62
Origin not stated[2]														
Both sexes														
2002	50	-	-	-	-	-	-	1	-	2	2	6	6	3
2001	50	-	-	1	-	-	-	-	-	1	6	6	4	2
2000	55	-	-	-	-	-	-	-	2	1	5	4	9	4
1999	44	-	-	-	-	-	-	-	-	-	4	5	9	5
Male														
2002	34	-	-	-	-	-	-	1	-	2	2	5	4	2
2001	25	-	-	1	-	-	-	-	-	-	2	4	3	2
2000	37	-	-	-	-	-	-	-	2	1	3	2	8	4
1999	28	-	-	-	-	-	-	-	-	-	3	4	7	5
Female														
2002	16	-	-	-	-	-	-	-	-	-	-	1	2	1
2001	25	-	-	-	-	-	-	-	-	1	4	2	1	-
2000	18	-	-	-	-	-	-	-	-	-	2	2	1	-
1999	16	-	-	-	-	-	-	-	-	-	1	1	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic liver disease and cirrhosis (K73-K74)										
Non-Hispanic White										
Female										
2001	445	574	743	878	622	332	116	18	-	-
2000	400	568	741	874	638	337	99	15	2	-
1999	453	579	808	869	632	320	114	14	2	-
Non-Hispanic Black										
Both sexes										
2002	154	125	90	79	46	21	11	-	-	-
2001	133	127	109	84	42	14	9	2	-	1
2000	144	140	120	66	47	26	12	1	-	-
1999	162	133	108	83	46	26	10	2	-	-
Male										
2002	93	73	45	26	19	6	3	-	-	-
2001	94	78	55	38	11	3	3	-	-	1
2000	91	71	61	34	14	9	4	-	-	-
1999	90	73	54	37	18	6	3	-	-	-
Female										
2002	61	52	45	53	27	15	8	-	-	-
2001	39	49	54	46	31	11	6	2	-	-
2000	53	69	59	32	33	17	8	1	-	-
1999	72	60	54	46	28	20	7	2	-	-
Origin not stated[2]										
Both sexes										
2002	9	4	8	4	3	1	-	-	-	1
2001	7	5	5	6	4	1	-	2	-	-
2000	6	8	8	2	4	-	2	-	-	-
1999	4	4	2	5	2	2	2	-	-	-
Male										
2002	7	1	5	3	1	-	-	-	-	1
2001	5	4	-	1	1	1	-	1	-	-
2000	5	6	4	-	1	-	1	-	-	-
1999	3	3	1	1	-	1	-	-	-	-
Female										
2002	2	3	3	1	2	1	-	-	-	-
2001	2	1	5	5	3	-	-	1	-	-
2000	1	2	4	2	3	-	1	-	-	-
1999	1	1	1	4	2	1	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cholelithiasis and other disorders of gallbladder (K80-K82)														
All origins														
Both sexes														
2002	2,979	-	-	-	-	2	7	9	12	15	39	52	77	93
2001	2,965	1	-	-	1	2	5	11	8	21	23	69	68	94
2000	2,810	-	-	1	1	2	7	6	13	22	31	43	66	80
1999	2,830	2	-	-	1	4	6	5	11	25	34	55	69	95
Male														
2002	1,341	-	-	-	-	-	3	4	7	7	22	35	48	59
2001	1,341	1	-	-	-	1	2	5	4	9	16	39	40	61
2000	1,244	-	-	-	1	1	3	2	5	12	15	23	30	48
1999	1,248	1	-	-	1	2	3	-	4	14	16	28	30	54
Female														
2002	1,638	-	-	-	-	2	4	5	5	8	17	17	29	34
2001	1,624	-	-	-	1	1	3	6	4	12	7	30	28	33
2000	1,566	-	-	1	-	1	4	4	8	10	16	20	36	32
1999	1,582	1	-	-	-	2	3	5	7	11	18	27	39	41
Hispanic														
Both sexes														
2002	196	-	-	-	-	-	2	2	4	3	8	5	6	9
2001	197	-	-	-	-	1	1	1	2	6	2	8	5	12
2000	183	-	-	-	-	-	2	1	2	2	3	7	11	8
1999	185	-	-	-	-	-	-	1	2	6	5	8	6	9
Male														
2002	93	-	-	-	-	-	-	-	2	2	5	4	6	8
2001	92	-	-	-	-	-	-	-	1	2	1	2	3	9
2000	99	-	-	-	-	-	-	1	-	2	3	4	5	6
1999	84	-	-	-	-	-	-	-	1	3	3	2	4	3
Female														
2002	103	-	-	-	-	-	2	2	2	1	3	1	-	1
2001	105	-	-	-	-	1	1	1	1	4	1	6	2	3
2000	84	-	-	-	-	-	2	-	2	-	-	3	6	2
1999	101	-	-	-	-	-	-	1	1	3	2	6	2	6
Non-Hispanic[1]														
Both sexes														
2002	2,776	-	-	-	-	2	5	7	8	12	31	47	71	84
2001	2,764	1	-	-	1	1	4	10	6	15	21	61	63	82
2000	2,619	-	-	1	1	2	5	5	11	20	28	36	55	72
1999	2,639	2	-	-	1	4	6	4	9	19	29	47	63	86
Male														
2002	1,244	-	-	-	-	-	3	4	5	5	17	31	42	51
2001	1,248	1	-	-	-	1	2	5	3	7	15	37	37	52

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All origins										
Both sexes										
2002	123	194	310	420	572	525	351	150	28	-
2001	148	193	333	432	513	542	364	106	31	-
2000	109	184	266	419	507	528	358	143	24	-
1999	108	171	304	417	531	521	324	120	27	-
Male										
2002	63	103	155	229	271	195	112	26	2	-
2001	81	96	173	231	233	209	111	21	8	-
2000	62	100	142	224	237	204	105	27	3	-
1999	66	89	163	208	239	203	101	22	4	-
Female										
2002	60	91	155	191	301	330	239	124	26	-
2001	67	97	160	201	280	333	253	85	23	-
2000	47	84	124	195	270	324	253	116	21	-
1999	42	82	141	209	292	318	223	98	23	-
Hispanic										
Both sexes										
2002	16	17	25	29	26	24	11	8	1	-
2001	11	17	34	22	25	22	23	2	3	-
2000	11	22	19	26	25	25	12	7	-	-
1999	11	14	20	30	27	26	10	8	2	-
Male										
2002	6	7	14	15	8	9	5	2	-	-
2001	7	7	18	13	12	6	7	1	3	-
2000	9	12	13	18	14	6	4	2	-	-
1999	9	7	8	18	11	9	5	1	-	-
Female										
2002	10	10	11	14	18	15	6	6	1	-
2001	4	10	16	9	13	16	16	1	-	-
2000	2	10	6	8	11	19	8	5	-	-
1999	2	7	12	12	16	17	5	7	2	-
Non-Hispanic[1]										
Both sexes										
2002	106	176	284	391	544	500	339	142	27	-
2001	137	174	299	410	486	520	341	104	28	-
2000	98	161	244	392	481	502	345	136	24	-
1999	97	157	281	387	502	494	314	112	25	-
Male										
2002	56	96	140	214	262	185	107	24	2	-
2001	74	88	155	218	221	203	104	20	5	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cholelithiasis and other disorders of gallbladder (K80-K82)														
Non-Hispanic[1]														
Male														
2000	1,141	-	-	-	1	1	3	1	5	10	12	19	25	42
1999	1,163	1	-	-	1	2	3	-	3	11	13	26	26	51
Female														
2002	1,532	-	-	-	-	2	2	3	3	7	14	16	29	33
2001	1,516	-	-	-	1	-	2	5	3	8	6	24	26	30
2000	1,478	-	-	1	-	1	2	4	6	10	16	17	30	30
1999	1,476	1	-	-	-	2	3	4	6	8	16	21	37	35
Non-Hispanic White														
Both sexes														
2002	2,420	-	-	-	-	2	3	4	6	7	26	33	49	68
2001	2,368	1	-	-	1	1	2	8	4	8	17	41	45	60
2000	2,272	-	-	-	1	1	4	2	5	15	18	20	38	56
1999	2,299	-	-	-	1	3	3	3	7	11	15	36	49	64
Male														
2002	1,087	-	-	-	-	-	3	2	4	3	14	20	29	42
2001	1,065	1	-	-	-	1	1	3	2	5	12	24	28	39
2000	1,012	-	-	-	1	1	2	1	3	8	9	13	18	35
1999	1,017	-	-	-	1	1	3	-	1	7	7	19	21	35
Female														
2002	1,333	-	-	-	-	2	-	2	2	4	12	13	20	26
2001	1,303	-	-	-	1	-	1	5	2	3	5	17	17	21
2000	1,260	-	-	-	-	-	2	1	2	7	9	7	20	21
1999	1,282	-	-	-	-	2	-	3	6	4	8	17	28	29
Non-Hispanic Black														
Both sexes														
2002	266	-	-	-	-	-	1	3	2	5	3	10	19	12
2001	309	-	-	-	-	-	2	2	2	7	4	14	17	20
2000	274	-	-	1	-	1	1	3	5	4	10	13	14	13
1999	257	2	-	-	-	1	3	1	2	7	13	8	13	14
Male														
2002	106	-	-	-	-	-	-	2	1	2	2	8	10	6
2001	130	-	-	-	-	-	1	2	1	2	3	9	9	12
2000	92	-	-	-	-	-	1	-	2	1	3	4	6	5
1999	102	1	-	-	-	1	-	-	2	3	6	5	4	10
Female														
2002	160	-	-	-	-	-	1	1	1	3	1	2	9	6
2001	179	-	-	-	-	-	1	-	1	5	1	5	8	8
2000	182	-	-	1	-	1	-	3	3	3	7	9	8	8

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)										
Non-Hispanic[1]										
Male										
2000	53	87	128	205	223	198	100	25	3	-
1999	57	82	154	190	228	194	96	21	4	-
Female										
2002	50	80	144	177	282	315	232	118	25	-
2001	63	86	144	192	265	317	237	84	23	-
2000	45	74	116	187	258	304	245	111	21	-
1999	40	75	127	197	274	300	218	91	21	-
Non-Hispanic White										
Both sexes										
2002	84	150	240	334	484	458	315	134	23	-
2001	108	131	250	357	430	471	312	95	26	-
2000	76	131	194	347	428	463	322	128	23	-
1999	70	120	241	348	457	454	289	105	23	-
Male										
2002	45	85	122	190	234	169	100	23	2	-
2001	57	65	135	196	199	181	94	18	4	-
2000	42	76	105	185	206	187	94	24	2	-
1999	44	65	136	173	216	175	88	21	4	-
Female										
2002	39	65	118	144	250	289	215	111	21	-
2001	51	66	115	161	231	290	218	77	22	-
2000	34	55	89	162	222	276	228	104	21	-
1999	26	55	105	175	241	279	201	84	19	-
Non-Hispanic Black										
Both sexes										
2002	16	20	34	44	47	26	14	6	4	-
2001	22	32	46	35	42	33	21	8	2	-
2000	16	25	36	36	39	33	18	6	-	-
1999	21	29	30	26	39	25	15	6	2	-
Male										
2002	10	7	10	16	21	8	3	-	-	-
2001	12	16	18	10	11	15	6	2	1	-
2000	9	8	15	14	12	9	3	-	-	-
1999	9	12	13	12	11	9	4	-	-	-
Female										
2002	6	13	24	28	26	18	11	6	4	-
2001	10	16	28	25	31	18	15	6	1	-
2000	7	17	21	22	27	24	15	6	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Cholelithiasis and other disorders of gallbladder (K80-K82)														
Non-Hispanic Black														
Female														
1999	155	1	-	-	-	-	3	1	-	4	7	3	9	4
Origin not stated[2]														
Both sexes														
2002	7	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	8	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	6	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	4	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	4	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	-	-	-	-	-
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
All origins														
Both sexes														
2002	40,974	173	14	9	12	21	63	107	162	274	475	790	1,103	1,505
2001	39,480	133	16	9	9	21	71	82	143	299	469	749	1,073	1,404
2000	37,251	162	14	11	14	32	56	80	164	278	462	687	964	1,269
1999	35,525	166	15	13	18	24	48	97	155	268	439	608	872	1,176
Male														
2002	19,695	109	7	5	7	10	31	44	85	148	259	454	632	797
2001	18,852	76	7	3	3	14	31	45	67	158	256	445	608	751
2000	17,811	103	7	7	10	18	25	44	85	152	277	433	549	697
1999	17,016	90	9	8	10	11	22	43	82	152	250	365	481	627
Female														
2002	21,279	64	7	4	5	11	32	63	77	126	216	336	471	708
2001	20,628	57	9	6	6	7	40	37	76	141	213	304	465	653
2000	19,440	59	7	4	4	14	31	36	79	126	185	254	415	572
1999	18,509	76	6	5	8	13	26	54	73	116	189	243	391	549

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cholelithiasis and other disorders of gallbladder (K80-K82)										
Non-Hispanic Black										
Female										
1999	12	17	17	14	28	16	11	6	2	-
Origin not stated[2]										
Both sexes										
2002	1	1	1	-	2	1	1	-	-	-
2001	-	2	-	-	2	-	-	-	-	-
2000	-	1	3	1	1	1	1	-	-	-
1999	-	-	3	-	2	1	-	-	-	-
Male										
2002	1	-	1	-	1	1	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-
2000	-	1	1	1	-	-	1	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	-	1	-	-	1	-	1	-	-	-
2001	-	1	-	-	2	-	-	-	-	-
2000	-	-	2	-	1	1	-	-	-	-
1999	-	-	2	-	2	1	-	-	-	-
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All origins										
Both sexes										
2002	1,950	2,820	4,344	6,338	7,558	6,959	4,450	1,549	298	-
2001	1,880	3,026	4,330	6,105	6,992	6,612	4,225	1,528	303	1
2000	1,831	2,811	4,179	5,893	6,565	6,150	3,915	1,443	269	2
1999	1,688	2,694	4,147	5,707	6,220	5,825	3,703	1,388	254	-
Male										
2002	1,027	1,491	2,311	3,248	3,672	3,088	1,693	495	82	-
2001	963	1,610	2,216	3,074	3,411	2,996	1,619	435	64	-
2000	954	1,444	2,120	3,039	3,174	2,672	1,473	453	73	2
1999	903	1,366	2,138	2,986	3,045	2,584	1,353	424	67	-
Female										
2002	923	1,329	2,033	3,090	3,886	3,871	2,757	1,054	216	-
2001	917	1,416	2,114	3,031	3,581	3,616	2,606	1,093	239	1
2000	877	1,367	2,059	2,854	3,391	3,478	2,442	990	196	-
1999	785	1,328	2,009	2,721	3,175	3,241	2,350	964	187	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
Hispanic														
Both sexes														
2002	1,898	35	5	2	3	-	9	12	21	23	42	57	100	119
2001	1,857	18	2	2	2	4	12	10	15	24	50	51	81	117
2000	1,646	25	4	2	1	7	7	6	20	19	34	45	64	87
1999	1,584	20	3	4	5	1	5	10	11	23	39	50	57	96
Male														
2002	937	21	1	1	2	-	5	3	9	12	26	29	59	64
2001	892	13	-	1	1	4	5	4	6	14	32	27	37	63
2000	805	16	3	1	1	3	4	4	7	9	27	32	37	45
1999	814	15	2	3	4	-	2	4	6	12	19	30	34	58
Female														
2002	961	14	4	1	1	-	4	9	12	11	16	28	41	55
2001	965	5	2	1	1	-	7	6	9	10	18	24	44	54
2000	841	9	1	1	-	4	3	2	13	10	7	13	27	42
1999	770	5	1	1	1	1	3	6	5	11	20	20	23	38
Non-Hispanic[1]														
Both sexes														
2002	38,933	138	9	7	9	21	54	94	141	250	431	728	997	1,377
2001	37,511	113	14	7	7	17	59	72	126	272	419	692	985	1,282
2000	35,509	134	10	9	13	24	49	74	143	259	427	642	894	1,180
1999	33,830	145	12	9	13	23	43	87	144	245	398	557	811	1,073
Male														
2002	18,682	88	6	4	5	10	26	41	76	135	232	422	570	730
2001	17,908	61	7	2	2	10	26	41	59	142	224	415	567	687
2000	16,958	84	4	6	9	14	21	40	78	143	249	401	507	651
1999	16,143	74	7	5	6	11	20	39	76	140	230	334	443	565
Female														
2002	20,251	50	3	3	4	11	28	53	65	115	199	306	427	647
2001	19,603	52	7	5	5	7	33	31	67	130	195	277	418	595
2000	18,551	50	6	3	4	10	28	34	65	116	178	241	387	529
1999	17,687	71	5	4	7	12	23	48	68	105	168	223	368	508
Non-Hispanic White														
Both sexes														
2002	30,669	77	5	5	5	10	23	44	75	127	226	403	519	822
2001	29,449	58	11	3	3	11	34	35	65	142	217	352	552	763
2000	27,924	61	4	6	11	12	36	43	67	126	233	334	495	704
1999	26,418	81	10	7	6	15	21	42	74	123	196	272	445	645
Male														
2002	14,883	49	3	3	3	5	12	20	44	70	121	234	299	438
2001	14,331	31	4	2	1	7	14	18	27	89	122	209	334	425

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
Hispanic										
Both sexes										
2002	148	216	211	281	251	179	120	45	19	-
2001	120	218	252	296	228	180	121	44	10	-
2000	145	195	209	235	194	182	127	31	6	1
1999	134	172	197	239	187	167	111	48	5	-
Male										
2002	77	115	99	144	129	72	41	21	7	-
2001	69	100	117	150	101	82	48	15	3	-
2000	70	90	104	116	90	79	52	10	4	1
1999	79	84	94	133	79	88	46	21	1	-
Female										
2002	71	101	112	137	122	107	79	24	12	-
2001	51	118	135	146	127	98	73	29	7	-
2000	75	105	105	119	104	103	75	21	2	-
1999	55	88	103	106	108	79	65	27	4	-
Non-Hispanic[1]										
Both sexes										
2002	1,794	2,597	4,113	6,035	7,287	6,760	4,316	1,496	279	-
2001	1,754	2,803	4,063	5,796	6,742	6,418	4,093	1,484	293	-
2000	1,674	2,611	3,956	5,643	6,357	5,958	3,784	1,405	263	-
1999	1,548	2,514	3,937	5,459	6,015	5,636	3,577	1,335	249	-
Male										
2002	947	1,372	2,197	3,089	3,531	3,008	1,648	470	75	-
2001	890	1,507	2,091	2,920	3,300	2,908	1,568	420	61	-
2000	878	1,351	2,006	2,918	3,077	2,591	1,421	440	69	-
1999	819	1,276	2,036	2,849	2,959	2,484	1,302	402	66	-
Female										
2002	847	1,225	1,916	2,946	3,756	3,752	2,668	1,026	204	-
2001	864	1,296	1,972	2,876	3,442	3,510	2,525	1,064	232	-
2000	796	1,260	1,950	2,725	3,280	3,367	2,363	965	194	-
1999	729	1,238	1,901	2,610	3,056	3,152	2,275	933	183	-
Non-Hispanic White										
Both sexes										
2002	1,156	1,794	3,076	4,879	6,231	5,901	3,787	1,280	224	-
2001	1,098	1,907	3,077	4,686	5,712	5,617	3,572	1,293	241	-
2000	1,072	1,776	2,965	4,616	5,410	5,202	3,318	1,223	210	-
1999	991	1,738	2,980	4,440	5,064	4,807	3,126	1,140	195	-
Male										
2002	607	983	1,711	2,583	3,104	2,679	1,454	406	55	-
2001	587	1,072	1,657	2,454	2,856	2,598	1,399	372	53	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
Non-Hispanic White														
Male														
2000	13,567	41	2	5	7	7	18	24	38	71	135	208	283	400
1999	12,793	46	5	4	4	9	8	20	40	71	117	162	244	365
Female														
2002	15,786	28	2	2	2	5	11	24	31	57	105	169	220	384
2001	15,118	27	7	1	2	4	20	17	38	53	95	143	218	338
2000	14,357	20	2	1	4	5	18	19	29	55	98	126	212	304
1999	13,625	35	5	3	2	6	13	22	34	52	79	110	201	280
Non-Hispanic Black														
Both sexes														
2002	7,410	55	4	2	3	10	26	48	60	114	192	296	449	497
2001	7,214	52	3	4	3	5	22	35	58	114	192	307	395	459
2000	6,853	69	5	3	1	12	12	28	71	129	176	300	364	433
1999	6,659	58	1	2	6	8	22	40	63	118	192	269	346	378
Male														
2002	3,389	35	3	1	1	4	11	20	29	59	102	172	256	267
2001	3,158	28	3	-	-	2	10	22	32	45	94	188	211	240
2000	3,045	40	2	1	1	7	3	13	37	71	104	191	203	226
1999	2,984	25	1	1	2	2	12	17	30	67	106	163	190	171
Female														
2002	4,021	20	1	1	2	6	15	28	31	55	90	124	193	230
2001	4,056	24	-	4	3	3	12	13	26	69	98	119	184	219
2000	3,808	29	3	2	-	5	9	15	34	58	72	109	161	207
1999	3,675	33	-	1	4	6	10	23	33	51	86	106	156	207
Origin not stated[2]														
Both sexes														
2002	143	-	-	-	-	-	-	1	-	1	2	5	6	9
2001	112	2	-	-	-	-	-	-	2	3	-	6	7	5
2000	96	3	-	-	-	1	-	-	1	-	1	-	6	2
1999	111	1	-	-	-	-	-	-	-	-	2	1	4	7
Male														
2002	76	-	-	-	-	-	-	-	-	1	1	3	3	3
2001	52	2	-	-	-	-	-	-	2	2	-	3	4	1
2000	48	3	-	-	-	1	-	-	-	-	1	-	5	1
1999	59	1	-	-	-	-	-	-	-	-	1	1	4	4
Female														
2002	67	-	-	-	-	-	-	1	-	-	1	2	3	6
2001	60	-	-	-	-	-	-	-	-	1	-	3	3	4
2000	48	-	-	-	-	-	-	-	1	-	-	-	1	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
Non-Hispanic White										
Male										
2000	590	983	1,591	2,449	2,692	2,318	1,268	375	62	-
1999	538	922	1,610	2,359	2,548	2,144	1,161	362	54	-
Female										
2002	549	811	1,365	2,296	3,127	3,222	2,333	874	169	-
2001	511	835	1,420	2,232	2,856	3,019	2,173	921	188	-
2000	482	793	1,374	2,167	2,718	2,884	2,050	848	148	-
1999	453	816	1,370	2,081	2,516	2,663	1,965	778	141	-
Non-Hispanic Black										
Both sexes										
2002	575	737	933	1,024	917	759	462	195	52	-
2001	589	817	881	982	909	707	454	176	50	-
2000	547	748	889	923	853	667	411	168	44	-
1999	501	698	856	911	839	727	397	173	54	-
Male										
2002	306	353	441	444	365	282	165	54	19	-
2001	269	397	385	402	382	260	142	39	7	-
2000	258	330	376	418	341	233	126	59	5	-
1999	255	310	381	435	368	292	114	30	12	-
Female										
2002	269	384	492	580	552	477	297	141	33	-
2001	320	420	496	580	527	447	312	137	43	-
2000	289	418	513	505	512	434	285	109	39	-
1999	246	388	475	476	471	435	283	143	42	-
Origin not stated[2]										
Both sexes										
2002	8	7	20	22	20	20	14	8	-	-
2001	6	5	15	13	22	14	11	-	-	1
2000	12	5	14	15	14	10	4	7	-	1
1999	6	8	13	9	18	22	15	5	-	-
Male										
2002	3	4	15	15	12	8	4	4	-	-
2001	4	3	8	4	10	6	3	-	-	-
2000	6	3	10	5	7	2	-	3	-	1
1999	5	6	8	4	7	12	5	1	-	-
Female										
2002	5	3	5	7	8	12	10	4	-	-
2001	2	2	7	9	12	8	8	-	-	1
2000	6	2	4	10	7	8	4	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)														
Origin not stated[2]														
Female														
1999	52	-	-	-	-	-	-	-	-	-	1	-	-	3
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)														
All origins														
Both sexes														
2002	166	6	5	1	1	2	1	-	1	2	2	5	4	6
2001	156	8	4	1	-	-	-	2	1	4	4	1	5	9
2000	177	8	3	1	1	-	1	-	4	1	2	5	11	2
1999	182	3	2	3	2	-	2	3	2	2	3	6	5	5
Male														
2002	70	3	2	1	1	1	-	-	1	-	-	4	1	5
2001	73	4	3	-	-	-	-	2	-	4	2	1	4	6
2000	91	3	2	1	1	-	1	-	3	-	1	2	9	2
1999	90	3	2	1	1	-	2	2	2	-	2	3	3	2
Female														
2002	96	3	3	-	-	1	1	-	-	2	2	1	3	1
2001	83	4	1	1	-	-	-	-	1	-	2	-	1	3
2000	86	5	1	-	-	-	-	-	1	1	1	3	2	-
1999	92	-	-	2	1	-	-	1	-	2	1	3	2	3
Hispanic														
Both sexes														
2002	10	-	2	-	-	-	-	-	-	1	1	-	-	1
2001	13	1	-	-	-	-	-	-	1	1	-	-	2	1
2000	9	2	1	-	-	-	-	-	-	-	-	-	2	-
1999	12	1	-	1	1	-	-	-	-	-	-	1	-	-
Male														
2002	4	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	7	-	-	-	-	-	-	-	-	1	-	-	2	1
2000	4	-	1	-	-	-	-	-	-	-	-	-	1	-
1999	4	1	-	-	1	-	-	-	-	-	-	-	-	-
Female														
2002	6	-	2	-	-	-	-	-	-	1	1	-	-	-
2001	6	1	-	-	-	-	-	-	1	-	-	-	-	-
2000	5	2	-	-	-	-	-	-	-	-	-	-	1	-
1999	8	-	-	1	-	-	-	-	-	-	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
Origin not stated[2]										
Female										
1999	1	2	5	5	11	10	10	4	-	-
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)										
All origins										
Both sexes										
2002	10	13	14	19	32	23	13	5	1	-
2001	4	13	14	28	20	24	12	1	1	-
2000	8	13	16	16	36	27	13	8	1	-
1999	11	10	27	37	25	22	7	5	-	-
Male										
2002	6	9	5	5	12	10	2	1	1	-
2001	3	6	4	10	9	10	5	-	-	-
2000	6	7	5	11	15	13	5	4	-	-
1999	4	5	12	15	17	9	1	4	-	-
Female										
2002	4	4	9	14	20	13	11	4	-	-
2001	1	7	10	18	11	14	7	1	1	-
2000	2	6	11	5	21	14	8	4	1	-
1999	7	5	15	22	8	13	6	1	-	-
Hispanic										
Both sexes										
2002	1	2	-	-	2	-	-	-	-	-
2001	1	1	2	-	1	-	2	-	-	-
2000	-	1	1	-	1	1	-	-	-	-
1999	2	3	1	1	-	1	-	-	-	-
Male										
2002	1	1	-	-	1	-	-	-	-	-
2001	1	1	1	-	-	-	-	-	-	-
2000	-	-	-	-	1	1	-	-	-	-
1999	1	-	-	-	-	1	-	-	-	-
Female										
2002	-	1	-	-	1	-	-	-	-	-
2001	-	-	1	-	1	-	2	-	-	-
2000	-	1	1	-	-	-	-	-	-	-
1999	1	3	1	1	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)														
Non-Hispanic[1]														
Both sexes														
2002	155	6	3	1	1	2	1	-	1	1	1	5	4	5
2001	142	7	4	1	-	-	-	2	-	3	4	1	3	8
2000	168	6	2	1	1	-	1	-	4	1	2	5	9	2
1999	170	2	2	2	1	-	2	3	2	2	3	5	5	5
Male														
2002	66	3	2	1	1	1	-	-	1	-	-	4	1	4
2001	65	4	3	-	-	-	-	2	-	3	2	1	2	5
2000	87	3	1	1	1	-	1	-	3	-	1	2	8	2
1999	86	2	2	1	-	-	2	2	2	-	2	3	3	2
Female														
2002	89	3	1	-	-	1	1	-	-	1	1	1	3	1
2001	77	3	1	1	-	-	-	-	-	-	2	-	1	3
2000	81	3	1	-	-	-	-	-	1	1	1	3	1	-
1999	84	-	-	1	1	-	-	1	-	2	1	2	2	3
Non-Hispanic White														
Both sexes														
2002	119	5	2	1	1	1	-	-	-	-	1	3	2	4
2001	116	6	3	-	-	-	-	-	-	3	4	1	2	6
2000	132	4	-	1	-	-	1	-	3	-	1	2	2	2
1999	132	2	2	2	-	-	1	1	1	2	1	3	3	3
Male														
2002	52	2	1	1	1	1	-	-	-	-	-	3	1	3
2001	52	3	2	-	-	-	-	-	-	3	2	1	2	3
2000	68	3	-	1	-	-	1	-	2	-	1	2	2	2
1999	63	2	2	1	-	-	1	-	1	-	1	2	2	1
Female														
2002	67	3	1	-	-	-	-	-	-	-	1	-	1	1
2001	64	3	1	-	-	-	-	-	-	-	2	-	-	3
2000	64	1	-	-	-	-	-	-	1	-	-	-	-	-
1999	69	-	-	1	-	-	-	1	-	2	-	1	1	2
Non-Hispanic Black														
Both sexes														
2002	29	1	1	-	-	1	1	-	1	1	-	2	2	1
2001	21	-	1	1	-	-	-	2	-	-	-	-	1	1
2000	30	2	2	-	-	-	-	-	1	1	1	3	7	-
1999	28	-	-	-	1	-	1	2	-	-	2	2	2	1
Male														
2002	11	1	1	-	-	-	-	-	1	-	-	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)										
Non-Hispanic[1]										
Both sexes										
2002	9	11	14	19	30	23	12	5	1	-
2001	3	12	12	28	18	24	10	1	1	-
2000	8	12	15	16	35	26	13	8	1	-
1999	9	7	26	36	25	21	7	5	-	-
Male										
2002	5	8	5	5	11	10	2	1	1	-
2001	2	5	3	10	8	10	5	-	-	-
2000	6	7	5	11	14	12	5	4	-	-
1999	3	5	12	15	17	8	1	4	-	-
Female										
2002	4	3	9	14	19	13	10	4	-	-
2001	1	7	9	18	10	14	5	1	1	-
2000	2	5	10	5	21	14	8	4	1	-
1999	6	2	14	21	8	13	6	1	-	-
Non-Hispanic White										
Both sexes										
2002	8	7	12	17	23	17	9	5	1	-
2001	3	8	10	25	16	20	7	1	1	-
2000	8	7	13	12	33	22	12	8	1	-
1999	4	6	22	31	20	16	7	5	-	-
Male										
2002	5	5	5	4	9	8	1	1	1	-
2001	2	2	2	10	8	8	4	-	-	-
2000	6	4	5	9	13	9	4	4	-	-
1999	1	4	10	13	13	4	1	4	-	-
Female										
2002	3	2	7	13	14	9	8	4	-	-
2001	1	6	8	15	8	12	3	1	1	-
2000	2	3	8	3	20	13	8	4	1	-
1999	3	2	12	18	7	12	6	1	-	-
Non-Hispanic Black										
Both sexes										
2002	-	3	2	2	4	5	2	-	-	-
2001	-	4	2	3	2	2	2	-	-	-
2000	-	4	2	3	2	2	-	-	-	-
1999	4	-	3	3	3	4	-	-	-	-
Male										
2002	-	2	-	1	1	2	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)														
Non-Hispanic Black														
Male														
2001	9	-	1	-	-	-	-	2	-	-	-	-	-	1
2000	13	-	1	-	-	-	-	-	1	-	-	-	6	-
1999	16	-	-	-	-	-	1	2	-	-	1	1	1	-
Female														
2002	18	-	-	-	-	1	1	-	-	1	-	1	2	-
2001	12	-	-	1	-	-	-	-	-	-	-	-	1	-
2000	17	2	1	-	-	-	-	-	-	1	1	3	1	-
1999	12	-	-	-	1	-	-	-	-	-	1	1	1	1
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
All origins														
Both sexes														
2002	553	2	-	1	2	2	2	10	13	13	13	28	25	33
2001	511	-	2	-	1	3	5	4	8	16	11	15	26	17
2000	556	1	-	1	3	2	7	8	9	14	20	18	21	31
1999	589	-	1	1	-	3	4	10	5	15	13	16	28	26
Male														
2002	278	1	-	-	1	1	2	4	9	7	7	10	15	20

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01, N04)										
Non-Hispanic Black										
Male										
2001	-	3	1	-	-	1	-	-	-	-
2000	-	2	-	1	1	1	-	-	-	-
1999	2	-	2	1	2	3	-	-	-	-
Female										
2002	-	1	2	1	3	3	2	-	-	-
2001	-	1	1	3	2	1	2	-	-	-
2000	-	2	2	2	1	1	-	-	-	-
1999	2	-	1	2	1	1	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	1	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	1	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)										
All origins										
Both sexes										
2002	19	29	56	86	92	74	38	14	1	-
2001	23	47	64	69	78	70	38	13	1	-
2000	28	43	60	78	94	78	23	15	2	-
1999	34	40	61	104	96	83	37	10	2	-
Male										
2002	7	14	35	50	42	32	19	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
All origins														
Male														
2001	264	-	1	-	-	3	2	-	1	7	6	9	15	10
2000	284	-	-	-	1	-	2	5	6	8	12	13	14	16
1999	304	-	-	1	-	3	3	7	3	9	9	10	15	13
Female														
2002	275	1	-	1	1	1	-	6	4	6	6	18	10	13
2001	247	-	1	-	1	-	3	4	7	9	5	6	11	7
2000	272	1	-	1	2	2	5	3	3	6	8	5	7	15
1999	285	-	1	-	-	-	1	3	2	6	4	6	13	13
Hispanic														
Both sexes														
2002	32	-	-	1	1	-	-	1	2	1	1	2	1	3
2001	34	-	1	-	-	1	-	-	1	4	2	3	1	2
2000	28	-	-	-	-	-	-	-	2	-	1	2	2	2
1999	33	-	1	-	-	-	1	2	1	-	2	2	-	3
Male														
2002	15	-	-	-	-	-	-	1	2	-	-	-	-	3
2001	17	-	-	-	-	1	-	-	-	3	2	-	-	1
2000	17	-	-	-	-	-	-	-	1	-	-	2	1	1
1999	18	-	-	-	-	-	1	2	1	-	2	1	-	2
Female														
2002	17	-	-	1	1	-	-	-	-	1	1	2	1	-
2001	17	-	1	-	-	-	-	-	1	1	-	3	1	1
2000	11	-	-	-	-	-	-	-	1	-	1	-	1	1
1999	15	-	1	-	-	-	-	-	-	-	-	1	-	1
Non-Hispanic[1]														
Both sexes														
2002	520	2	-	-	1	2	2	8	11	12	12	26	24	30
2001	477	-	1	-	1	2	5	4	7	12	9	12	25	15
2000	528	1	-	1	3	2	7	8	7	14	19	16	19	29
1999	555	-	-	1	-	3	3	8	4	15	11	14	28	23
Male														
2002	263	1	-	-	1	1	2	3	7	7	7	10	15	17
2001	247	-	1	-	-	2	2	-	1	4	4	9	15	9
2000	267	-	-	-	1	-	2	5	5	8	12	11	13	15
1999	285	-	-	1	-	3	2	5	2	9	7	9	15	11

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)										
All origins										
Male										
2001	15	23	40	39	45	32	10	6	-	-
2000	15	24	30	45	38	40	9	6	-	-
1999	19	20	25	58	49	38	17	5	-	-
Female										
2002	12	15	21	36	50	42	19	12	1	-
2001	8	24	24	30	33	38	28	7	1	-
2000	13	19	30	33	56	38	14	9	2	-
1999	15	20	36	46	47	45	20	5	2	-
Hispanic										
Both sexes										
2002	2	2	1	7	2	3	2	-	-	-
2001	2	2	4	4	2	4	1	-	-	-
2000	2	2	3	3	4	5	-	-	-	-
1999	3	5	3	2	3	2	2	1	-	-
Male										
2002	1	-	-	5	1	1	1	-	-	-
2001	2	1	1	2	2	2	-	-	-	-
2000	2	-	2	2	1	5	-	-	-	-
1999	2	4	-	1	-	1	-	1	-	-
Female										
2002	1	2	1	2	1	2	1	-	-	-
2001	-	1	3	2	-	2	1	-	-	-
2000	-	2	1	1	3	-	-	-	-	-
1999	1	1	3	1	3	1	2	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	17	27	55	79	90	71	36	14	1	-
2001	21	45	60	65	76	66	37	13	1	-
2000	26	41	57	75	90	73	23	15	2	-
1999	30	35	58	102	93	81	35	9	2	-
Male										
2002	6	14	35	45	41	31	18	2	-	-
2001	13	22	39	37	43	30	10	6	-	-
2000	13	24	28	43	37	35	9	6	-	-
1999	16	16	25	57	49	37	17	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
Non-Hispanic[1]														
Female														
2002	257	1	-	-	-	1	-	5	4	5	5	16	9	13
2001	230	-	-	-	1	-	3	4	6	8	5	3	10	6
2000	261	1	-	1	2	2	5	3	2	6	7	5	6	14
1999	270	-	-	-	-	-	1	3	2	6	4	5	13	12
Non-Hispanic White														
Both sexes														
2002	419	-	-	-	-	2	1	5	7	10	10	17	14	21
2001	384	-	1	-	-	1	5	3	5	4	5	5	12	11
2000	411	-	-	-	3	1	6	4	5	7	12	13	9	22
1999	436	-	-	1	-	2	1	6	2	8	5	12	18	15
Male														
2002	218	-	-	-	-	1	1	2	5	6	5	8	8	13
2001	200	-	1	-	-	1	2	-	1	2	1	3	7	6
2000	209	-	-	-	1	-	1	3	3	5	7	9	6	12
1999	227	-	-	1	-	2	-	4	1	4	4	8	10	7
Female														
2002	201	-	-	-	-	1	-	3	2	4	5	9	6	8
2001	184	-	-	-	-	-	3	3	4	2	4	2	5	5
2000	202	-	-	-	2	1	5	1	2	2	5	4	3	10
1999	209	-	-	-	-	-	1	2	1	4	1	4	8	8
Non-Hispanic Black														
Both sexes														
2002	80	2	-	-	-	-	1	3	3	1	2	9	9	5
2001	68	-	-	-	1	-	-	1	1	4	4	6	9	2
2000	90	1	-	1	-	1	1	3	1	7	3	3	9	5
1999	92	-	-	-	-	1	2	-	2	6	5	1	9	6
Male														
2002	37	1	-	-	-	-	1	1	2	-	2	2	6	3
2001	36	-	-	-	-	-	-	-	-	2	3	5	7	2
2000	44	-	-	-	-	-	1	1	1	3	2	2	6	2
1999	46	-	-	-	-	1	2	-	1	4	2	1	5	3
Female														
2002	43	1	-	-	-	-	-	2	1	1	-	7	3	2
2001	32	-	-	-	1	-	-	1	1	2	1	1	2	-
2000	46	1	-	1	-	1	-	2	-	4	1	1	3	3
1999	46	-	-	-	-	-	-	-	1	2	3	-	4	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)										
Non-Hispanic[1]										
Female										
2002	11	13	20	34	49	40	18	12	1	-
2001	8	23	21	28	33	36	27	7	1	-
2000	13	17	29	32	53	38	14	9	2	-
1999	14	19	33	45	44	44	18	5	2	-
Non-Hispanic White										
Both sexes										
2002	13	26	50	61	80	57	33	12	-	-
2001	13	33	54	57	66	62	35	12	-	-
2000	17	30	50	64	79	59	17	12	1	-
1999	21	24	48	89	80	68	27	7	2	-
Male										
2002	6	14	31	35	37	27	17	2	-	-
2001	8	16	36	32	39	29	10	6	-	-
2000	8	18	25	37	35	29	6	4	-	-
1999	12	13	20	51	42	31	14	3	-	-
Female										
2002	7	12	19	26	43	30	16	10	-	-
2001	5	17	18	25	27	33	25	6	-	-
2000	9	12	25	27	44	30	11	8	1	-
1999	9	11	28	38	38	37	13	4	2	-
Non-Hispanic Black										
Both sexes										
2002	3	1	4	13	7	12	2	2	1	-
2001	6	10	4	6	7	3	2	1	1	-
2000	7	9	5	9	10	9	4	2	-	-
1999	5	10	7	11	8	12	5	2	-	-
Male										
2002	-	-	4	8	3	3	1	-	-	-
2001	4	5	2	3	2	1	-	-	-	-
2000	5	6	2	4	2	4	2	1	-	-
1999	3	3	3	6	4	5	2	1	-	-
Female										
2002	3	1	-	5	4	9	1	2	1	-
2001	2	5	2	3	5	2	2	1	1	-
2000	2	3	3	5	8	5	2	1	-	-
1999	2	7	4	5	4	7	3	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)														
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	1	-	-	-	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Renal failure (N17-N19)														
All origins														
Both sexes														
2002	40,222	165	9	6	9	17	60	95	148	258	460	757	1,072	1,464
2001	38,784	125	9	8	8	18	64	76	133	278	454	729	1,041	1,376
2000	36,471	152	11	9	10	30	48	72	149	262	439	658	932	1,234
1999	34,719	163	12	9	16	21	42	83	147	250	423	586	836	1,144
Male														
2002	19,341	105	5	3	5	8	29	39	75	141	252	440	615	772
2001	18,509	72	3	3	3	11	29	43	66	146	248	434	588	735
2000	17,416	100	5	6	8	18	22	39	76	144	263	415	526	678
1999	16,611	87	7	6	9	8	17	34	76	143	239	352	462	612
Female														
2002	20,881	60	4	3	4	9	31	56	73	117	208	317	457	692
2001	20,275	53	6	5	5	7	35	33	67	132	206	295	453	641
2000	19,055	52	6	3	2	12	26	33	73	118	176	243	406	556
1999	18,108	76	5	3	7	13	25	49	71	107	184	234	374	532
Hispanic														
Both sexes														
2002	1,856	35	3	1	2	-	9	11	19	21	40	55	99	115
2001	1,806	17	1	2	2	3	12	10	13	18	48	47	78	114
2000	1,605	22	3	2	1	7	7	6	18	19	32	43	60	85

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic glomerulonephritis, nephritis and nephropathy, not specified as acute or chronic, and renal sclerosis, unspecified (N02-N03,N05-N07, N26)										
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Renal failure (N17-N19)										
All origins										
Both sexes										
2002	1,919	2,774	4,272	6,228	7,429	6,856	4,398	1,530	296	-
2001	1,850	2,963	4,248	6,007	6,891	6,517	4,174	1,514	300	1
2000	1,789	2,754	4,094	5,794	6,429	6,040	3,878	1,420	265	2
1999	1,640	2,641	4,056	5,560	6,092	5,718	3,656	1,372	252	-
Male										
2002	1,014	1,468	2,271	3,190	3,618	3,046	1,672	492	81	-
2001	945	1,581	2,171	3,025	3,356	2,953	1,604	429	64	-
2000	929	1,412	2,080	2,981	3,121	2,617	1,458	443	73	2
1999	880	1,340	2,100	2,912	2,976	2,535	1,334	415	67	-
Female										
2002	905	1,306	2,001	3,038	3,811	3,810	2,726	1,038	215	-
2001	905	1,382	2,077	2,982	3,535	3,564	2,570	1,085	236	1
2000	860	1,342	2,014	2,813	3,308	3,423	2,420	977	192	-
1999	760	1,301	1,956	2,648	3,116	3,183	2,322	957	185	-
Hispanic										
Both sexes										
2002	145	212	210	274	247	176	118	45	19	-
2001	116	214	246	292	225	176	118	44	10	-
2000	142	192	205	231	189	176	127	31	6	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Renal failure (N17-N19)														
Hispanic														
Both sexes														
1999	1,535	19	2	3	4	1	4	8	10	23	37	47	57	93
Male														
2002	918	21	1	1	2	-	5	2	7	12	26	29	59	60
2001	867	13	-	1	1	3	5	4	6	9	30	27	35	61
2000	782	16	2	1	1	3	4	4	6	9	26	30	35	44
1999	790	14	2	3	3	-	1	2	5	12	17	29	34	56
Female														
2002	938	14	2	-	-	-	4	9	12	9	14	26	40	55
2001	939	4	1	1	1	-	7	6	7	9	18	20	43	53
2000	823	6	1	1	-	4	3	2	12	10	6	13	25	41
1999	745	5	-	-	1	1	3	6	5	11	20	18	23	37
Non-Hispanic[1]														
Both sexes														
2002	38,225	130	6	5	7	17	51	84	129	236	418	697	967	1,340
2001	36,867	106	8	6	6	15	52	66	118	257	406	676	956	1,257
2000	34,771	127	8	7	9	22	41	66	130	243	406	615	866	1,147
1999	33,074	143	10	6	12	20	38	75	137	227	384	538	775	1,044
Male														
2002	18,347	84	4	2	3	8	24	37	68	128	225	408	553	709
2001	17,591	57	3	2	2	8	24	39	58	135	218	404	549	673
2000	16,586	81	3	5	7	14	18	35	70	135	236	385	486	633
1999	15,763	72	5	3	6	8	16	32	71	131	221	322	424	552
Female														
2002	19,878	46	2	3	4	9	27	47	61	108	193	289	414	631
2001	19,276	49	5	4	4	7	28	27	60	122	188	272	407	584
2000	18,185	46	5	2	2	8	23	31	60	108	170	230	380	514
1999	17,311	71	5	3	6	12	22	43	66	96	163	216	351	492
Non-Hispanic White														
Both sexes														
2002	30,104	72	3	4	4	7	22	37	68	116	215	383	502	795
2001	28,932	52	6	3	3	10	28	32	59	135	208	344	537	745
2000	27,350	57	4	5	8	11	29	39	58	119	220	316	484	678
1999	25,831	79	8	4	6	13	19	34	70	112	190	257	422	627
Male														
2002	14,610	47	2	2	2	3	11	17	39	64	116	223	290	422
2001	14,076	28	1	2	1	6	12	18	26	84	119	204	324	416
2000	13,275	38	2	4	6	7	16	21	33	66	127	196	275	385
1999	12,496	44	3	2	4	7	7	16	37	67	112	152	231	357

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)										
Hispanic										
Both sexes										
1999	129	163	193	234	183	164	109	47	5	-
Male										
2002	75	114	99	139	127	71	40	21	7	-
2001	66	98	115	148	99	80	48	15	3	-
2000	67	90	102	114	88	73	52	10	4	1
1999	76	80	94	131	78	86	46	20	1	-
Female										
2002	70	98	111	135	120	105	78	24	12	-
2001	50	116	131	144	126	96	70	29	7	-
2000	75	102	103	117	101	103	75	21	2	-
1999	53	83	99	103	105	78	63	27	4	-
Non-Hispanic[1]										
Both sexes										
2002	1,766	2,555	4,042	5,932	7,162	6,660	4,267	1,477	277	-
2001	1,728	2,744	3,987	5,702	6,645	6,327	4,045	1,470	290	-
2000	1,636	2,557	3,875	5,548	6,226	5,854	3,747	1,382	259	-
1999	1,506	2,470	3,850	5,317	5,891	5,532	3,532	1,320	247	-
Male										
2002	936	1,350	2,157	3,036	3,479	2,967	1,628	467	74	-
2001	875	1,480	2,048	2,873	3,248	2,867	1,553	414	61	-
2000	856	1,319	1,968	2,862	3,026	2,542	1,406	430	69	-
1999	800	1,254	1,998	2,777	2,891	2,437	1,283	394	66	-
Female										
2002	830	1,205	1,885	2,896	3,683	3,693	2,639	1,010	203	-
2001	853	1,264	1,939	2,829	3,397	3,460	2,492	1,056	229	-
2000	780	1,238	1,907	2,686	3,200	3,312	2,341	952	190	-
1999	706	1,216	1,852	2,540	3,000	3,095	2,249	926	181	-
Non-Hispanic White										
Both sexes										
2002	1,133	1,757	3,012	4,797	6,125	5,822	3,744	1,263	223	-
2001	1,080	1,864	3,009	4,604	5,629	5,535	3,529	1,280	240	-
2000	1,044	1,738	2,894	4,537	5,293	5,117	3,288	1,203	208	-
1999	965	1,707	2,907	4,317	4,962	4,721	3,091	1,127	193	-
Male										
2002	596	964	1,675	2,542	3,058	2,644	1,436	403	54	-
2001	577	1,054	1,618	2,412	2,809	2,561	1,385	366	53	-
2000	574	960	1,556	2,401	2,644	2,278	1,257	367	62	-
1999	525	905	1,579	2,295	2,492	2,107	1,145	355	54	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Renal failure (N17-N19)														
Non-Hispanic White														
Female														
2002	15,494	25	1	2	2	4	11	20	29	52	99	160	212	373
2001	14,856	24	5	1	2	4	16	14	33	51	89	140	213	329
2000	14,075	19	2	1	2	4	13	18	25	53	93	120	209	293
1999	13,335	35	5	2	2	6	12	18	33	45	78	105	191	270
Non-Hispanic Black														
Both sexes														
2002	7,297	52	3	1	3	9	24	45	56	112	190	285	437	491
2001	7,119	52	2	3	2	5	21	32	57	110	188	301	385	455
2000	6,724	66	3	2	1	11	11	25	68	120	172	291	348	428
1999	6,527	58	1	2	5	7	19	38	61	112	185	266	334	370
Male														
2002	3,339	33	2	-	1	4	10	19	26	59	100	169	249	263
2001	3,111	28	2	-	-	2	10	20	32	43	91	183	204	237
2000	2,986	40	1	1	1	7	2	12	35	68	102	187	191	224
1999	2,920	25	1	1	2	1	9	15	29	63	103	161	184	168
Female														
2002	3,958	19	1	1	2	5	14	26	30	53	90	116	188	228
2001	4,008	24	-	3	2	3	11	12	25	67	97	118	181	218
2000	3,738	26	2	1	-	4	9	13	33	52	70	104	157	204
1999	3,607	33	-	1	3	6	10	23	32	49	82	105	150	202
Origin not stated[2]														
Both sexes														
2002	141	-	-	-	-	-	-	-	-	1	2	5	6	9
2001	111	2	-	-	-	-	-	-	2	3	-	6	7	5
2000	95	3	-	-	-	1	-	-	1	-	1	-	6	2
1999	110	1	-	-	-	-	-	-	-	-	2	1	4	7
Male														
2002	76	-	-	-	-	-	-	-	-	1	1	3	3	3
2001	51	2	-	-	-	-	-	-	2	2	-	3	4	1
2000	48	3	-	-	-	1	-	-	-	-	1	-	5	1
1999	58	1	-	-	-	-	-	-	-	-	1	1	4	4
Female														
2002	65	-	-	-	-	-	-	-	-	-	1	2	3	6
2001	60	-	-	-	-	-	-	-	-	1	-	3	3	4
2000	47	-	-	-	-	-	-	-	1	-	-	-	1	1
1999	52	-	-	-	-	-	-	-	-	-	1	-	-	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)										
Non-Hispanic White										
Female										
2002	537	793	1,337	2,255	3,067	3,178	2,308	860	169	-
2001	503	810	1,391	2,192	2,820	2,974	2,144	914	187	-
2000	470	778	1,338	2,136	2,649	2,839	2,031	836	146	-
1999	440	802	1,328	2,022	2,470	2,614	1,946	772	139	-
Non-Hispanic Black										
Both sexes										
2002	572	733	927	1,009	905	741	458	193	51	-
2001	583	803	875	972	899	701	450	175	48	-
2000	540	735	881	910	841	655	407	166	43	-
1999	490	687	846	896	824	711	390	171	54	-
Male										
2002	306	351	437	435	361	277	164	54	19	-
2001	265	389	382	399	379	257	142	39	7	-
2000	253	322	374	413	338	228	124	58	5	-
1999	250	306	376	428	361	284	112	29	12	-
Female										
2002	266	382	490	574	544	464	294	139	32	-
2001	318	414	493	573	520	444	308	136	41	-
2000	287	413	507	497	503	427	283	108	38	-
1999	240	381	470	468	463	427	278	142	42	-
Origin not stated[2]										
Both sexes										
2002	8	7	20	22	20	20	13	8	-	-
2001	6	5	15	13	21	14	11	-	-	1
2000	11	5	14	15	14	10	4	7	-	1
1999	5	8	13	9	18	22	15	5	-	-
Male										
2002	3	4	15	15	12	8	4	4	-	-
2001	4	3	8	4	9	6	3	-	-	-
2000	6	3	10	5	7	2	-	3	-	1
1999	4	6	8	4	7	12	5	1	-	-
Female										
2002	5	3	5	7	8	12	9	4	-	-
2001	2	2	7	9	12	8	8	-	-	1
2000	5	2	4	10	7	8	4	4	-	-
1999	1	2	5	5	11	10	10	4	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of kidney (N25,N27)														
All origins														
Both sexes														
2002	33	-	-	1	-	-	-	2	-	1	-	-	2	2
2001	29	-	1	-	-	-	2	-	1	1	-	4	1	2
2000	47	1	-	-	-	-	-	-	2	1	1	6	-	2
1999	35	-	-	-	-	-	-	1	1	1	-	-	3	1
Male														
2002	6	-	-	1	-	-	-	1	-	-	-	-	1	-
2001	6	-	-	-	-	-	-	-	-	1	-	1	1	-
2000	20	-	-	-	-	-	-	-	-	-	1	3	-	1
1999	11	-	-	-	-	-	-	-	1	-	-	-	1	-
Female														
2002	27	-	-	-	-	-	-	1	-	1	-	-	1	2
2001	23	-	1	-	-	-	2	-	1	-	-	3	-	2
2000	27	1	-	-	-	-	-	-	2	1	-	3	-	1
1999	24	-	-	-	-	-	-	1	-	1	-	-	2	1
Hispanic														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	-	-	-	-	-	-	-	-	1	-	1	-	-
2000	4	1	-	-	-	-	-	-	-	-	1	-	-	-
1999	4	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	2	1	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	33	-	-	1	-	-	-	2	-	1	-	-	2	2
2001	25	-	1	-	-	-	2	-	1	-	-	3	1	2
2000	42	-	-	-	-	-	-	-	2	1	-	6	-	2
1999	31	-	-	-	-	-	-	1	1	1	-	-	3	1
Male														
2002	6	-	-	1	-	-	-	1	-	-	-	-	1	-
2001	5	-	-	-	-	-	-	-	-	-	-	1	1	-
2000	18	-	-	-	-	-	-	-	-	-	-	3	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of kidney (N25,N27)										
All origins										
Both sexes										
2002	2	4	2	5	5	6	1	-	-	-
2001	3	3	4	1	3	1	1	-	1	-
2000	6	1	9	5	6	5	1	-	1	-
1999	3	3	3	6	7	2	3	1	-	-
Male										
2002	-	-	-	3	-	-	-	-	-	-
2001	-	-	1	-	1	1	-	-	-	-
2000	4	1	5	2	-	2	1	-	-	-
1999	-	1	1	1	3	2	1	-	-	-
Female										
2002	2	4	2	2	5	6	1	-	-	-
2001	3	3	3	1	2	-	1	-	1	-
2000	2	-	4	3	6	3	-	-	1	-
1999	3	2	2	5	4	-	2	1	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-
2000	1	-	-	1	-	-	-	-	-	-
1999	-	1	-	2	1	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	1	1	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	1	-	-	-	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	-	1	-	1	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	2	4	2	5	5	6	1	-	-	-
2001	2	2	4	1	3	1	1	-	1	-
2000	4	1	9	4	6	5	1	-	1	-
1999	3	2	3	4	6	2	3	1	-	-
Male										
2002	-	-	-	3	-	-	-	-	-	-
2001	-	-	1	-	1	1	-	-	-	-
2000	3	1	5	2	-	2	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of kidney (N25,N27)														
Non-Hispanic[1]														
Male														
1999	9	-	-	-	-	-	-	-	1	-	-	-	1	-
Female														
2002	27	-	-	-	-	-	-	1	-	1	-	-	1	2
2001	20	-	1	-	-	-	2	-	1	-	-	2	-	2
2000	24	-	-	-	-	-	-	-	2	1	-	3	-	1
1999	22	-	-	-	-	-	-	1	-	1	-	-	2	1
Non-Hispanic White														
Both sexes														
2002	27	-	-	-	-	-	-	2	-	1	-	-	1	2
2001	17	-	1	-	-	-	1	-	1	-	-	2	1	1
2000	31	-	-	-	-	-	-	-	1	-	-	3	-	2
1999	19	-	-	-	-	-	-	1	1	1	-	-	2	-
Male														
2002	3	-	-	-	-	-	-	1	-	-	-	-	-	-
2001	3	-	-	-	-	-	-	-	-	-	-	1	1	-
2000	15	-	-	-	-	-	-	-	-	-	-	1	-	1
1999	7	-	-	-	-	-	-	-	1	-	-	-	1	-
Female														
2002	24	-	-	-	-	-	-	1	-	1	-	-	1	2
2001	14	-	1	-	-	-	1	-	1	-	-	1	-	1
2000	16	-	-	-	-	-	-	-	1	-	-	2	-	1
1999	12	-	-	-	-	-	-	1	-	1	-	-	1	-
Non-Hispanic Black														
Both sexes														
2002	4	-	-	1	-	-	-	-	-	-	-	-	1	-
2001	6	-	-	-	-	-	1	-	-	-	-	-	-	1
2000	9	-	-	-	-	-	-	-	1	1	-	3	-	-
1999	12	-	-	-	-	-	-	-	-	-	-	-	1	1
Male														
2002	2	-	-	1	-	-	-	-	-	-	-	-	1	-
2001	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	2	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	4	-	-	-	-	-	1	-	-	-	-	-	-	1
2000	7	-	-	-	-	-	-	-	1	1	-	1	-	-
1999	10	-	-	-	-	-	-	-	-	-	-	-	1	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of kidney (N25,N27)										
Non-Hispanic[1]										
Male										
1999	-	1	1	-	2	2	1	-	-	-
Female										
2002	2	4	2	2	5	6	1	-	-	-
2001	2	2	3	1	2	-	1	-	1	-
2000	1	-	4	2	6	3	-	-	1	-
1999	3	1	2	4	4	-	2	1	-	-
Non-Hispanic White										
Both sexes										
2002	2	4	2	4	3	5	1	-	-	-
2001	2	2	4	-	1	-	1	-	-	-
2000	3	1	8	3	5	4	1	-	-	-
1999	1	1	3	3	2	2	1	1	-	-
Male										
2002	-	-	-	2	-	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	2	1	5	2	-	2	1	-	-	-
1999	-	-	1	-	1	2	1	-	-	-
Female										
2002	2	4	2	2	3	5	1	-	-	-
2001	2	2	3	-	1	-	1	-	-	-
2000	1	-	3	1	5	2	-	-	-	-
1999	1	1	2	3	1	-	-	1	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	-	-	1	1	1	-	-	1	-
2000	-	-	1	1	-	1	-	-	1	-
1999	2	1	-	1	4	-	2	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	1	1	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	1	-	-	-	-	-
Female										
2002	-	-	-	-	1	1	-	-	-	-
2001	-	-	-	1	-	-	-	-	1	-
2000	-	-	1	1	-	1	-	-	1	-
1999	2	-	-	1	3	-	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other disorders of kidney (N25,N27)														
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Infections of kidney (N10-N12,N13.6, N15.1)														
All origins														
Both sexes														
2002	788	2	-	-	3	2	3	5	8	11	17	19	21	26
2001	748	2	4	-	-	1	4	5	6	16	17	21	25	23
2000	813	5	3	-	-	1	1	4	5	9	12	21	21	25
1999	846	3	1	1	-	3	1	5	4	13	15	22	33	30
Male														
2002	251	2	-	-	1	-	1	3	3	3	9	5	10	10
2001	213	1	2	-	-	-	2	-	3	4	6	6	5	6
2000	214	3	-	-	-	1	-	2	1	2	6	6	9	5
1999	263	1	-	-	-	1	-	-	-	2	5	8	11	14
Female														
2002	537	-	-	-	2	2	2	2	5	8	8	14	11	16
2001	535	1	2	-	-	1	2	5	3	12	11	15	20	17
2000	599	2	3	-	-	-	1	2	4	7	6	15	12	20
1999	583	2	1	1	-	2	1	5	4	11	10	14	22	16
Hispanic														
Both sexes														
2002	47	1	-	-	2	-	-	2	1	2	-	1	2	-
2001	33	-	-	-	-	-	-	1	-	1	2	2	3	1
2000	39	1	-	-	-	-	-	-	-	1	1	3	1	1
1999	55	-	-	-	-	-	-	1	-	4	3	5	1	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of kidney (N25,N27)										
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Infections of kidney (N10-N12,N13.6, N15.1)										
All origins										
Both sexes										
2002	33	48	62	118	120	146	105	36	3	-
2001	23	35	52	126	123	129	100	33	3	-
2000	34	38	66	114	134	145	114	44	17	-
1999	29	52	81	109	133	154	102	49	6	-
Male										
2002	11	19	28	44	28	42	25	7	-	-
2001	6	19	21	36	40	32	16	8	-	-
2000	10	13	13	34	40	32	30	5	2	-
1999	13	19	25	37	46	45	24	12	-	-
Female										
2002	22	29	34	74	92	104	80	29	3	-
2001	17	16	31	90	83	97	84	25	3	-
2000	24	25	53	80	94	113	84	39	15	-
1999	16	33	56	72	87	109	78	37	6	-
Hispanic										
Both sexes										
2002	2	2	10	9	5	2	5	1	-	-
2001	-	4	4	4	5	3	2	1	-	-
2000	3	4	3	4	6	4	6	1	-	-
1999	3	3	5	6	9	4	4	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Infections of kidney (N10-N12,N13.6, N15.1)														
Hispanic														
Male														
2002	19	1	-	-	-	-	-	2	-	1	-	-	2	-
2001	10	-	-	-	-	-	-	-	-	-	-	2	1	-
2000	11	1	-	-	-	-	-	-	-	-	-	2	1	-
1999	21	-	-	-	-	-	-	-	-	1	1	2	-	3
Female														
2002	28	-	-	-	2	-	-	-	1	1	-	1	-	-
2001	23	-	-	-	-	-	-	1	-	1	2	-	2	1
2000	28	-	-	-	-	-	-	-	-	1	1	1	-	1
1999	34	-	-	-	-	-	-	1	-	3	2	3	1	3
Non-Hispanic[1]														
Both sexes														
2002	740	1	-	-	1	2	3	3	7	9	17	18	19	26
2001	714	2	4	-	-	1	4	4	6	15	15	19	22	22
2000	771	4	3	-	-	1	1	4	5	8	11	18	20	24
1999	786	3	1	1	-	3	1	4	4	8	12	17	32	24
Male														
2002	232	1	-	-	1	-	1	1	3	2	9	5	8	10
2001	203	1	2	-	-	-	2	-	3	4	6	4	4	6
2000	201	2	-	-	-	1	-	2	1	2	6	4	8	5
1999	240	1	-	-	-	1	-	-	-	-	4	6	11	11
Female														
2002	508	-	-	-	-	2	2	2	4	7	8	13	11	16
2001	511	1	2	-	-	1	2	4	3	11	9	15	18	16
2000	570	2	3	-	-	-	1	2	4	6	5	14	12	19
1999	546	2	1	1	-	2	1	4	4	8	8	11	21	13
Non-Hispanic White														
Both sexes														
2002	645	1	-	-	-	-	1	2	5	5	15	9	14	22
2001	614	1	2	-	-	-	2	3	4	10	13	13	16	17
2000	647	4	3	-	-	-	1	4	3	4	6	14	12	21
1999	689	1	1	1	-	1	1	3	4	3	9	14	24	20
Male														
2002	200	1	-	-	-	-	1	1	3	2	7	3	7	9
2001	166	-	1	-	-	-	1	-	2	3	6	3	3	6
2000	169	2	-	-	-	-	-	2	1	-	4	4	5	5
1999	209	1	-	-	-	1	-	-	-	-	2	4	8	9
Female														
2002	445	-	-	-	-	-	-	1	2	3	8	6	7	13

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of kidney (N10-N12,N13.6, N15.1)										
Hispanic										
Male										
2002	1	-	4	5	-	1	1	1	-	-
2001	-	1	2	1	-	2	1	-	-	-
2000	1	2	1	-	2	1	-	-	-	-
1999	2	2	2	2	4	-	1	1	-	-
Female										
2002	1	2	6	4	5	1	4	-	-	-
2001	-	3	2	3	5	1	1	1	-	-
2000	2	2	2	4	4	3	6	1	-	-
1999	1	1	3	4	5	4	3	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	31	46	52	109	115	143	100	35	3	-
2001	23	31	48	122	117	126	98	32	3	-
2000	31	33	63	109	127	141	108	43	17	-
1999	25	49	76	101	124	150	98	47	6	-
Male										
2002	10	19	24	39	28	41	24	6	-	-
2001	6	18	19	35	40	30	15	8	-	-
2000	9	10	12	34	37	31	30	5	2	-
1999	11	17	23	34	42	45	23	11	-	-
Female										
2002	21	27	28	70	87	102	76	29	3	-
2001	17	13	29	87	77	96	83	24	3	-
2000	22	23	51	75	90	110	78	38	15	-
1999	14	32	53	67	82	105	75	36	6	-
Non-Hispanic White										
Both sexes										
2002	22	39	50	96	107	131	93	30	3	-
2001	19	22	43	111	99	114	90	32	3	-
2000	23	26	48	91	107	130	98	37	15	-
1999	18	40	62	91	115	140	93	43	5	-
Male										
2002	7	13	23	33	25	36	24	5	-	-
2001	3	11	16	29	35	26	13	8	-	-
2000	7	6	7	30	32	31	27	4	2	-
1999	8	14	16	32	38	43	22	11	-	-
Female										
2002	15	26	27	63	82	95	69	25	3	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Infections of kidney (N10-N12,N13.6, N15.1)														
Non-Hispanic White														
Female														
2001	448	1	1	-	-	-	1	3	2	7	7	10	13	11
2000	478	2	3	-	-	-	1	2	2	4	2	10	7	16
1999	480	-	1	1	-	-	1	3	4	3	7	10	16	11
Non-Hispanic Black														
Both sexes														
2002	73	-	-	-	-	1	2	-	2	4	2	6	2	4
2001	80	1	-	-	-	-	1	1	2	5	2	4	6	3
2000	101	-	-	-	-	1	-	-	2	3	4	3	8	2
1999	85	1	-	-	-	2	-	1	-	3	3	3	7	4
Male														
2002	22	-	-	-	-	-	-	-	-	-	2	1	-	1
2001	28	1	-	-	-	-	1	-	1	1	-	1	1	-
2000	27	-	-	-	-	1	-	-	-	1	1	-	3	-
1999	29	-	-	-	-	-	-	-	-	-	2	2	3	2
Female														
2002	51	-	-	-	-	1	2	-	2	4	-	5	2	3
2001	52	-	-	-	-	-	-	1	1	4	2	3	5	3
2000	74	-	-	-	-	-	-	-	2	2	3	3	5	2
1999	56	1	-	-	-	2	-	1	-	3	1	1	4	2
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	1	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	1	-	-	-	-
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Infections of kidney (N10-N12,N13.6, N15.1)										
Non-Hispanic White										
Female										
2001	16	11	27	82	64	88	77	24	3	-
2000	16	20	41	61	75	99	71	33	13	-
1999	10	26	46	59	77	97	71	32	5	-
Non-Hispanic Black										
Both sexes										
2002	6	7	1	11	4	11	5	5	-	-
2001	4	8	5	8	14	10	6	-	-	-
2000	6	7	12	16	12	10	8	6	1	-
1999	7	7	13	8	8	9	5	3	1	-
Male										
2002	2	6	-	4	1	4	-	1	-	-
2001	3	6	3	3	3	3	1	-	-	-
2000	2	4	4	4	3	-	3	1	-	-
1999	3	3	7	1	3	2	1	-	-	-
Female										
2002	4	1	1	7	3	7	5	4	-	-
2001	1	2	2	5	11	7	5	-	-	-
2000	4	3	8	12	9	10	5	5	1	-
1999	4	4	6	7	5	7	4	3	1	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	1	-	1	1	-	-	-	-	-
1999	1	-	-	2	-	-	-	1	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	1	-	-	-	-	-
1999	-	-	-	1	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	1	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	-	-	1	-	-	-	-	-	-
1999	1	-	-	1	-	-	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hyperplasia of prostate (N40)														
All origins														
Both sexes														
2002	437	-	-	-	-	-	-	-	-	1	-	-	-	1
2001	439	-	-	-	-	-	-	-	-	-	-	-	2	3
2000	433	-	-	-	-	-	-	-	-	-	1	1	-	4
1999	430	-	-	-	-	-	-	-	-	-	-	-	1	4
Male														
2002	437	-	-	-	-	-	-	-	-	1	-	-	-	1
2001	439	-	-	-	-	-	-	-	-	-	-	-	2	3
2000	433	-	-	-	-	-	-	-	-	-	1	1	-	4
1999	430	-	-	-	-	-	-	-	-	-	-	-	1	4
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic														
Both sexes														
2002	14	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	11	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	19	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	11	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	14	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	11	-	-	-	-	-	-	-	-	-	-	-	-	1
2000	19	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	11	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic[1]														
Both sexes														
2002	417	-	-	-	-	-	-	-	-	1	-	-	-	1
2001	423	-	-	-	-	-	-	-	-	-	-	-	2	2
2000	414	-	-	-	-	-	-	-	-	-	1	1	-	4
1999	417	-	-	-	-	-	-	-	-	-	-	-	-	3
Male														
2002	417	-	-	-	-	-	-	-	-	1	-	-	-	1
2001	423	-	-	-	-	-	-	-	-	-	-	-	2	2
2000	414	-	-	-	-	-	-	-	-	-	1	1	-	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hyperplasia of prostate (N40)										
All origins										
Both sexes										
2002	9	9	32	48	116	93	85	31	11	1
2001	6	14	31	60	81	110	94	33	5	-
2000	7	10	28	64	95	106	83	29	4	1
1999	9	15	32	59	101	109	67	24	9	-
Male										
2002	9	9	32	48	116	93	85	31	11	1
2001	6	14	31	60	81	110	94	33	5	-
2000	7	10	28	64	95	106	83	29	4	1
1999	9	15	32	59	101	109	67	24	9	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Hispanic										
Both sexes										
2002	1	1	2	3	6	-	1	-	-	-
2001	-	2	2	1	3	1	-	1	-	-
2000	1	2	3	3	-	6	2	2	-	-
1999	-	2	1	-	4	1	1	-	1	-
Male										
2002	1	1	2	3	6	-	1	-	-	-
2001	-	2	2	1	3	1	-	1	-	-
2000	1	2	3	3	-	6	2	2	-	-
1999	-	2	1	-	4	1	1	-	1	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic[1]										
Both sexes										
2002	7	7	29	45	110	92	83	30	11	1
2001	6	11	28	57	77	109	94	32	5	-
2000	6	8	25	61	95	100	81	27	4	1
1999	9	13	30	59	97	108	66	24	8	-
Male										
2002	7	7	29	45	110	92	83	30	11	1
2001	6	11	28	57	77	109	94	32	5	-
2000	6	8	25	61	95	100	81	27	4	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hyperplasia of prostate (N40)														
Non-Hispanic[1]														
Male														
1999	417	-	-	-	-	-	-	-	-	-	-	-	-	3
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White														
Both sexes														
2002	370	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	388	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	370	-	-	-	-	-	-	-	-	-	-	1	-	2
1999	377	-	-	-	-	-	-	-	-	-	-	-	-	2
Male														
2002	370	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	388	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	370	-	-	-	-	-	-	-	-	-	-	1	-	2
1999	377	-	-	-	-	-	-	-	-	-	-	-	-	2
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black														
Both sexes														
2002	38	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	31	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	37	-	-	-	-	-	-	-	-	-	1	-	-	2
1999	38	-	-	-	-	-	-	-	-	-	-	-	-	1
Male														
2002	38	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	31	-	-	-	-	-	-	-	-	-	-	-	1	1
2000	37	-	-	-	-	-	-	-	-	-	1	-	-	2
1999	38	-	-	-	-	-	-	-	-	-	-	-	-	1
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hyperplasia of prostate (N40)										
Non-Hispanic[1]										
Male										
1999	9	13	30	59	97	108	66	24	8	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic White										
Both sexes										
2002	4	6	23	36	101	82	80	27	10	-
2001	3	10	25	52	70	101	90	30	5	-
2000	4	8	22	54	88	90	72	25	4	-
1999	6	10	26	56	87	98	62	24	6	-
Male										
2002	4	6	23	36	101	82	80	27	10	-
2001	3	10	25	52	70	101	90	30	5	-
2000	4	8	22	54	88	90	72	25	4	-
1999	6	10	26	56	87	98	62	24	6	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Non-Hispanic Black										
Both sexes										
2002	3	1	6	6	7	9	2	1	1	1
2001	3	1	2	5	7	6	4	1	-	-
2000	2	-	2	7	7	7	7	1	-	1
1999	3	3	3	3	10	9	4	-	2	-
Male										
2002	3	1	6	6	7	9	2	1	1	1
2001	3	1	2	5	7	6	4	1	-	-
2000	2	-	2	7	7	7	7	1	-	1
1999	3	3	3	3	10	9	4	-	2	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Hyperplasia of prostate (N40)														
Origin not stated[2]														
Both sexes														
2002	6	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	1	-
Male														
2002	6	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	1	-
Female														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Inflammatory diseases of female pelvic organs (N70-N76)														
All origins														
Both sexes														
2002	114	1	-	-	-	1	1	-	4	12	5	4	7	4
2001	127	1	-	-	-	-	2	2	1	2	8	9	3	4
2000	120	2	-	-	1	-	2	2	4	9	6	6	3	4
1999	108	2	1	-	-	1	2	3	3	4	8	4	2	6
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	114	1	-	-	-	1	1	-	4	12	5	4	7	4
2001	127	1	-	-	-	-	2	2	1	2	8	9	3	4
2000	120	2	-	-	1	-	2	2	4	9	6	6	3	4
1999	108	2	1	-	-	1	2	3	3	4	8	4	2	6
Hispanic														
Both sexes														
2002	3	-	-	-	-	-	-	-	1	-	2	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	1	1	-
2000	7	-	-	-	-	-	-	-	1	1	-	-	-	1
1999	7	-	-	-	-	-	-	-	-	-	1	1	1	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hyperplasia of prostate (N40)										
Origin not stated[2]										
Both sexes										
2002	1	1	1	-	-	1	1	1	-	-
2001	-	1	1	2	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Male										
2002	1	1	1	-	-	1	1	1	-	-
2001	-	1	1	2	1	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	1	-	-	-	-	-	-	-
Female										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Inflammatory diseases of female pelvic organs (N70-N76)										
All origins										
Both sexes										
2002	2	4	4	17	12	22	11	3	-	-
2001	8	8	10	13	18	23	11	4	-	-
2000	4	11	10	17	12	11	12	3	1	-
1999	7	4	11	15	17	6	6	5	1	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	2	4	4	17	12	22	11	3	-	-
2001	8	8	10	13	18	23	11	4	-	-
2000	4	11	10	17	12	11	12	3	1	-
1999	7	4	11	15	17	6	6	5	1	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	1	-	-	1	-	-	-	-
2000	1	2	-	-	-	-	1	-	-	-
1999	-	-	-	1	-	2	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Inflammatory diseases of female pelvic organs (N70-N76)														
Hispanic														
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	3	-	-	-	-	-	-	-	1	-	2	-	-	-
2001	5	-	-	-	-	-	-	-	-	-	-	1	1	-
2000	7	-	-	-	-	-	-	-	1	1	-	-	-	1
1999	7	-	-	-	-	-	-	-	-	-	1	1	1	1
Non-Hispanic[1]														
Both sexes														
2002	110	1	-	-	-	1	1	-	3	12	3	4	7	4
2001	122	1	-	-	-	-	2	2	1	2	8	8	2	4
2000	111	2	-	-	1	-	2	2	3	8	6	4	3	3
1999	101	2	1	-	-	1	2	3	3	4	7	3	1	5
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	110	1	-	-	-	1	1	-	3	12	3	4	7	4
2001	122	1	-	-	-	-	2	2	1	2	8	8	2	4
2000	111	2	-	-	1	-	2	2	3	8	6	4	3	3
1999	101	2	1	-	-	1	2	3	3	4	7	3	1	5
Non-Hispanic White														
Both sexes														
2002	88	1	-	-	-	-	-	-	2	4	3	2	5	3
2001	98	1	-	-	-	-	2	2	1	-	6	5	1	4
2000	87	1	-	-	-	-	2	2	1	4	2	3	3	2
1999	73	1	1	-	-	1	1	1	2	1	2	1	1	5
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	88	1	-	-	-	-	-	-	2	4	3	2	5	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)										
Hispanic										
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	1	-	-	1	-	-	-	-
2000	1	2	-	-	-	-	1	-	-	-
1999	-	-	-	1	-	2	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	2	4	4	17	11	22	11	3	-	-
2001	8	7	9	13	18	22	11	4	-	-
2000	3	9	10	17	12	11	11	3	1	-
1999	7	4	11	14	17	4	6	5	1	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	2	4	4	17	11	22	11	3	-	-
2001	8	7	9	13	18	22	11	4	-	-
2000	3	9	10	17	12	11	11	3	1	-
1999	7	4	11	14	17	4	6	5	1	-
Non-Hispanic White										
Both sexes										
2002	2	4	3	16	11	19	10	3	-	-
2001	4	5	7	11	16	21	8	4	-	-
2000	3	7	8	15	11	9	11	2	1	-
1999	6	2	8	9	16	4	6	5	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	2	4	3	16	11	19	10	3	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Inflammatory diseases of female pelvic organs (N70-N76)														
Non-Hispanic White														
Female														
2001	98	1	-	-	-	-	2	2	1	-	6	5	1	4
2000	87	1	-	-	-	-	2	2	1	4	2	3	3	2
1999	73	1	1	-	-	1	1	1	2	1	2	1	1	5
Non-Hispanic Black														
Both sexes														
2002	20	-	-	-	-	1	1	-	1	8	-	1	2	1
2001	23	-	-	-	-	-	-	-	-	2	2	3	1	-
2000	21	1	-	-	1	-	-	-	1	3	4	1	-	1
1999	26	1	-	-	-	-	1	2	-	2	5	2	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	20	-	-	-	-	1	1	-	1	8	-	1	2	1
2001	23	-	-	-	-	-	-	-	-	2	2	3	1	-
2000	21	1	-	-	1	-	-	-	1	3	4	1	-	1
1999	26	1	-	-	-	-	1	2	-	2	5	2	-	-
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	2	-	-	-	-	-	-	-	-	-	-	2	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Inflammatory diseases of female pelvic organs (N70-N76)										
Non-Hispanic White										
Female										
2001	4	5	7	11	16	21	8	4	-	-
2000	3	7	8	15	11	9	11	2	1	-
1999	6	2	8	9	16	4	6	5	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	1	1	-	2	1	-	-	-
2001	4	2	2	2	2	-	3	-	-	-
2000	-	2	2	2	1	2	-	-	-	-
1999	1	2	3	5	1	-	-	-	1	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	1	1	-	2	1	-	-	-
2001	4	2	2	2	2	-	3	-	-	-
2000	-	2	2	2	1	2	-	-	-	-
1999	1	2	3	5	1	-	-	-	1	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	1	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	1	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy, childbirth and the puerperium (O00-O99)														
All origins														
Both sexes														
2002	379	. . .	. . .	. . .	1	28	61	82	93	79	26	8	-	-
2001	416	. . .	. . .	. . .	3	37	73	93	97	73	33	5	1	-
2000	404	. . .	. . .	. . .	-	18	77	87	95	86	35	5	1	-
1999	406	. . .	. . .	. . .	-	33	64	89	92	88	33	5	2	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	379	. . .	. . .	. . .	1	28	61	82	93	79	26	8	-	-
2001	416	. . .	. . .	. . .	3	37	73	93	97	73	33	5	1	-
2000	404	. . .	. . .	. . .	-	18	77	87	95	86	35	5	1	-
1999	406	. . .	. . .	. . .	-	33	64	89	92	88	33	5	2	-
Hispanic														
Both sexes														
2002	70	. . .	. . .	. . .	-	6	13	10	18	16	5	2	-	-
2001	83	. . .	. . .	. . .	1	9	17	21	17	10	7	-	1	-
2000	84	. . .	. . .	. . .	-	2	22	20	16	19	4	1	-	-
1999	70	. . .	. . .	. . .	-	6	13	18	8	21	4	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	70	. . .	. . .	. . .	-	6	13	10	18	16	5	2	-	-
2001	83	. . .	. . .	. . .	1	9	17	21	17	10	7	-	1	-
2000	84	. . .	. . .	. . .	-	2	22	20	16	19	4	1	-	-
1999	70	. . .	. . .	. . .	-	6	13	18	8	21	4	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	305	. . .	. . .	. . .	1	22	47	71	74	62	21	6	-	-
2001	332	. . .	. . .	. . .	2	28	56	72	79	63	26	5	-	-
2000	319	. . .	. . .	. . .	-	16	55	67	79	66	31	4	1	-
1999	333	. . .	. . .	. . .	-	27	49	70	84	67	29	5	2	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)										
All origins										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy, childbirth and the puerperium (O00-O99)														
Non-Hispanic[1]														
Male														
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	305	. . .	. . .	. . .	1	22	47	71	74	62	21	6	-	-
2001	332	. . .	. . .	. . .	2	28	56	72	79	63	26	5	-	-
2000	319	. . .	. . .	. . .	-	16	55	67	79	66	31	4	1	-
1999	333	. . .	. . .	. . .	-	27	49	70	84	67	29	5	2	-
Non-Hispanic White														
Both sexes														
2002	133	. . .	. . .	. . .	-	6	15	31	38	25	14	3	-	-
2001	156	. . .	. . .	. . .	1	13	25	31	46	25	12	2	-	-
2000	161	. . .	. . .	. . .	-	9	24	33	39	37	17	1	1	-
1999	152	. . .	. . .	. . .	-	8	18	29	45	35	13	4	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	133	. . .	. . .	. . .	-	6	15	31	38	25	14	3	-	-
2001	156	. . .	. . .	. . .	1	13	25	31	46	25	12	2	-	-
2000	161	. . .	. . .	. . .	-	9	24	33	39	37	17	1	1	-
1999	152	. . .	. . .	. . .	-	8	18	29	45	35	13	4	-	-
Non-Hispanic Black														
Both sexes														
2002	153	. . .	. . .	. . .	1	16	29	37	27	34	7	2	-	-
2001	154	. . .	. . .	. . .	1	14	31	35	30	28	12	3	-	-
2000	139	. . .	. . .	. . .	-	4	30	31	34	25	12	3	-	-
1999	158	. . .	. . .	. . .	-	18	28	37	33	27	13	-	2	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	153	. . .	. . .	. . .	1	16	29	37	27	34	7	2	-	-
2001	154	. . .	. . .	. . .	1	14	31	35	30	28	12	3	-	-
2000	139	. . .	. . .	. . .	-	4	30	31	34	25	12	3	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)										
Non-Hispanic[1]										
Male										
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy, childbirth and the puerperium (O00-O99)														
Non-Hispanic Black														
Female														
1999	158	. . .	. . .	. . .	-	18	28	37	33	27	13	-	2	-
Origin not stated[2]														
Both sexes														
2002	4	. . .	. . .	. . .	-	-	1	1	1	1	-	-	-	-
2001	1	. . .	. . .	. . .	-	-	-	-	1	-	-	-	-	-
2000	1	. . .	. . .	. . .	-	-	-	-	-	1	-	-	-	-
1999	3	. . .	. . .	. . .	-	-	2	1	-	-	-	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	4	. . .	. . .	. . .	-	-	1	1	1	1	-	-	-	-
2001	1	. . .	. . .	. . .	-	-	-	-	1	-	-	-	-	-
2000	1	. . .	. . .	. . .	-	-	-	-	-	1	-	-	-	-
1999	3	. . .	. . .	. . .	-	-	2	1	-	-	-	-	-	-
Pregnancy with abortive outcome (O00-O07)														
All origins														
Both sexes														
2002	22	. . .	. . .	. . .	1	-	4	4	5	6	1	1	-	-
2001	38	. . .	. . .	. . .	2	6	7	6	8	8	1	-	-	-
2000	37	. . .	. . .	. . .	-	-	6	12	5	10	4	-	-	-
1999	31	. . .	. . .	. . .	-	3	5	5	10	6	2	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	22	. . .	. . .	. . .	1	-	4	4	5	6	1	1	-	-
2001	38	. . .	. . .	. . .	2	6	7	6	8	8	1	-	-	-
2000	37	. . .	. . .	. . .	-	-	6	12	5	10	4	-	-	-
1999	31	. . .	. . .	. . .	-	3	5	5	10	6	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pregnancy, childbirth and the puerperium (O00-O99)										
Non-Hispanic Black										
Female										
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Pregnancy with abortive outcome (O00-O07)										
All origins										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy with abortive outcome (O00-O07)														
Hispanic														
Both sexes														
2002	1	. . .	. . .	. . .	-	-	-	1	-	-	-	-	-	-
2001	1	. . .	. . .	. . .	1	-	-	-	-	-	-	-	-	-
2000	6	. . .	. . .	. . .	-	-	2	3	-	1	-	-	-	-
1999	4	. . .	. . .	. . .	-	2	-	-	-	2	-	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	1	. . .	. . .	. . .	-	-	-	1	-	-	-	-	-	-
2001	1	. . .	. . .	. . .	1	-	-	-	-	-	-	-	-	-
2000	6	. . .	. . .	. . .	-	-	2	3	-	1	-	-	-	-
1999	4	. . .	. . .	. . .	-	2	-	-	-	2	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	20	. . .	. . .	. . .	1	-	4	3	5	5	1	1	-	-
2001	37	. . .	. . .	. . .	1	6	7	6	8	8	1	-	-	-
2000	30	. . .	. . .	. . .	-	-	4	9	5	8	4	-	-	-
1999	26	. . .	. . .	. . .	-	1	4	5	10	4	2	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	20	. . .	. . .	. . .	1	-	4	3	5	5	1	1	-	-
2001	37	. . .	. . .	. . .	1	6	7	6	8	8	1	-	-	-
2000	30	. . .	. . .	. . .	-	-	4	9	5	8	4	-	-	-
1999	26	. . .	. . .	. . .	-	1	4	5	10	4	2	-	-	-
Non-Hispanic White														
Both sexes														
2002	6	. . .	. . .	. . .	-	-	1	1	3	-	1	-	-	-
2001	19	. . .	. . .	. . .	-	3	4	4	4	3	1	-	-	-
2000	12	. . .	. . .	. . .	-	-	-	4	1	4	3	-	-	-
1999	11	. . .	. . .	. . .	-	-	2	1	5	1	2	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)										
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy with abortive outcome (O00-O07)														
Non-Hispanic White														
Male														
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	6	. . .	. . .	. . .	-	-	1	1	3	-	1	-	-	-
2001	19	. . .	. . .	. . .	-	3	4	4	4	3	1	-	-	-
2000	12	. . .	. . .	. . .	-	-	-	4	1	4	3	-	-	-
1999	11	. . .	. . .	. . .	-	-	2	1	5	1	2	-	-	-
Non-Hispanic Black														
Both sexes														
2002	11	. . .	. . .	. . .	1	-	2	2	2	4	-	-	-	-
2001	16	. . .	. . .	. . .	1	3	3	2	3	4	-	-	-	-
2000	18	. . .	. . .	. . .	-	-	4	5	4	4	1	-	-	-
1999	14	. . .	. . .	. . .	-	1	2	4	4	3	-	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	11	. . .	. . .	. . .	1	-	2	2	2	4	-	-	-	-
2001	16	. . .	. . .	. . .	1	3	3	2	3	4	-	-	-	-
2000	18	. . .	. . .	. . .	-	-	4	5	4	4	1	-	-	-
1999	14	. . .	. . .	. . .	-	1	2	4	4	3	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	1	. . .	. . .	. . .	-	-	-	-	-	1	-	-	-	-
2001	-	. . .	. . .	. . .	-	-	-	-	-	-	-	-	-	-
2000	1	. . .	. . .	. . .	-	-	-	-	-	1	-	-	-	-
1999	1	. . .	. . .	. . .	-	-	1	-	-	-	-	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	1	. . .	. . .	. . .	-	-	-	-	-	1	-	-	-	-
2001	-	. . .	. . .	. . .	-	-	-	-	-	-	-	-	-	-
2000	1	. . .	. . .	. . .	-	-	-	-	-	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)										
Non-Hispanic White										
Male										
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Pregnancy with abortive outcome (O00-O07)														
Origin not stated[2]														
Female														
1999	1	. . .	. . .	. . .	-	-	1	-	-	-	-	-	-	-
Other complications of pregnancy, childbirth and the puerperium (O10-O99)														
All origins														
Both sexes														
2002	357	. . .	. . .	. . .	-	28	57	78	88	73	25	7	-	-
2001	378	. . .	. . .	. . .	1	31	66	87	89	65	32	5	1	-
2000	367	. . .	. . .	. . .	-	18	71	75	90	76	31	5	1	-
1999	375	. . .	. . .	. . .	-	30	59	84	82	82	31	5	2	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	357	. . .	. . .	. . .	-	28	57	78	88	73	25	7	-	-
2001	378	. . .	. . .	. . .	1	31	66	87	89	65	32	5	1	-
2000	367	. . .	. . .	. . .	-	18	71	75	90	76	31	5	1	-
1999	375	. . .	. . .	. . .	-	30	59	84	82	82	31	5	2	-
Hispanic														
Both sexes														
2002	69	. . .	. . .	. . .	-	6	13	9	18	16	5	2	-	-
2001	82	. . .	. . .	. . .	-	9	17	21	17	10	7	-	1	-
2000	78	. . .	. . .	. . .	-	2	20	17	16	18	4	1	-	-
1999	66	. . .	. . .	. . .	-	4	13	18	8	19	4	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	69	. . .	. . .	. . .	-	6	13	9	18	16	5	2	-	-
2001	82	. . .	. . .	. . .	-	9	17	21	17	10	7	-	1	-
2000	78	. . .	. . .	. . .	-	2	20	17	16	18	4	1	-	-
1999	66	. . .	. . .	. . .	-	4	13	18	8	19	4	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pregnancy with abortive outcome (O00-O07)										
Origin not stated[2]										
Female										
1999	-	-	-	-	-	-	-	-	-	-
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
All origins										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)														
Non-Hispanic[1]														
Both sexes														
2002	285	. . .	. . .	. . .	-	22	43	68	69	57	20	5	-	-
2001	295	. . .	. . .	. . .	1	22	49	66	71	55	25	5	-	-
2000	289	. . .	. . .	. . .	-	16	51	58	74	58	27	4	1	-
1999	307	. . .	. . .	. . .	-	26	45	65	74	63	27	5	2	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	285	. . .	. . .	. . .	-	22	43	68	69	57	20	5	-	-
2001	295	. . .	. . .	. . .	1	22	49	66	71	55	25	5	-	-
2000	289	. . .	. . .	. . .	-	16	51	58	74	58	27	4	1	-
1999	307	. . .	. . .	. . .	-	26	45	65	74	63	27	5	2	-
Non-Hispanic White														
Both sexes														
2002	127	. . .	. . .	. . .	-	6	14	30	35	25	13	3	-	-
2001	137	. . .	. . .	. . .	1	10	21	27	42	22	11	2	-	-
2000	149	. . .	. . .	. . .	-	9	24	29	38	33	14	1	1	-
1999	141	. . .	. . .	. . .	-	8	16	28	40	34	11	4	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	127	. . .	. . .	. . .	-	6	14	30	35	25	13	3	-	-
2001	137	. . .	. . .	. . .	1	10	21	27	42	22	11	2	-	-
2000	149	. . .	. . .	. . .	-	9	24	29	38	33	14	1	1	-
1999	141	. . .	. . .	. . .	-	8	16	28	40	34	11	4	-	-
Non-Hispanic Black														
Both sexes														
2002	142	. . .	. . .	. . .	-	16	27	35	25	30	7	2	-	-
2001	138	. . .	. . .	. . .	-	11	28	33	27	24	12	3	-	-
2000	121	. . .	. . .	. . .	-	4	26	26	30	21	11	3	-	-
1999	144	. . .	. . .	. . .	-	17	26	33	29	24	13	-	2	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
Non-Hispanic[1]										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)														
Non-Hispanic Black														
Male														
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	142	. . .	. . .	. . .	-	16	27	35	25	30	7	2	-	-
2001	138	. . .	. . .	. . .	-	11	28	33	27	24	12	3	-	-
2000	121	. . .	. . .	. . .	-	4	26	26	30	21	11	3	-	-
1999	144	. . .	. . .	. . .	-	17	26	33	29	24	13	-	2	-
Origin not stated[2]														
Both sexes														
2002	3	. . .	. . .	. . .	-	-	1	1	1	-	-	-	-	-
2001	1	. . .	. . .	. . .	-	-	-	-	1	-	-	-	-	-
2000	-	. . .	. . .	. . .	-	-	-	-	-	-	-	-	-	-
1999	2	. . .	. . .	. . .	-	-	1	1	-	-	-	-	-	-
Male														
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female														
2002	3	. . .	. . .	. . .	-	-	1	1	1	-	-	-	-	-
2001	1	. . .	. . .	. . .	-	-	-	-	1	-	-	-	-	-
2000	-	. . .	. . .	. . .	-	-	-	-	-	-	-	-	-	-
1999	2	. . .	. . .	. . .	-	-	1	1	-	-	-	-	-	-
Certain conditions originating in the perinatal period (P00-P96)														
All origins														
Both sexes														
2002	14,254	14,106	65	15	16	14	7	6	4	7	3	2	-	1
2001	13,887	13,734	72	22	14	8	9	5	1	6	3	3	2	2
2000	14,069	13,918	79	25	16	10	5	5	2	2	2	1	1	1
1999	14,259	14,084	92	25	18	11	5	2	5	2	1	2	-	1
Male														
2002	8,047	7,960	40	10	9	9	3	3	1	4	1	2	-	1
2001	7,868	7,775	44	16	8	3	4	2	-	4	2	3	2	1
2000	7,968	7,872	52	13	10	9	2	4	2	-	2	1	1	-
1999	8,044	7,941	56	14	9	6	4	-	2	2	-	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other complications of pregnancy, childbirth and the puerperium (O10-O99)										
Non-Hispanic Black										
Male										
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2001	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
2000	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
1999	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .	. . .
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Certain conditions originating in the perinatal period (P00-P96)										
All origins										
Both sexes										
2002	-	1	2	-	2	-	-	1	-	2
2001	-	2	1	1	-	1	-	-	-	1
2000	1	-	-	-	-	-	-	-	-	1
1999	2	1	1	1	-	-	1	-	-	5
Male										
2002	-	-	1	-	1	-	-	-	-	2
2001	-	1	1	1	-	-	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	1	-	-	1	-	-	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain conditions originating in the perinatal period (P00-P96)														
All origins														
Female														
2002	6,207	6,146	25	5	7	5	4	3	3	3	2	-	-	-
2001	6,019	5,959	28	6	6	5	5	3	1	2	1	-	-	1
2000	6,101	6,046	27	12	6	1	3	1	-	2	-	-	-	1
1999	6,215	6,143	36	11	9	5	1	2	3	-	1	-	-	1
Hispanic														
Both sexes														
2002	2,402	2,375	14	4	2	2	1	1	-	1	1	-	-	-
2001	2,227	2,203	14	3	-	-	2	4	-	-	-	1	-	-
2000	2,145	2,120	12	7	3	1	1	-	-	1	-	-	-	-
1999	2,153	2,127	18	2	-	3	1	1	-	-	-	-	-	-
Male														
2002	1,352	1,335	9	3	2	2	-	1	-	-	-	-	-	-
2001	1,268	1,258	6	2	-	-	-	1	-	-	-	1	-	-
2000	1,194	1,180	7	4	2	1	-	-	-	-	-	-	-	-
1999	1,192	1,181	5	2	-	2	1	-	-	-	-	-	-	-
Female														
2002	1,050	1,040	5	1	-	-	1	-	-	1	1	-	-	-
2001	959	945	8	1	-	-	2	3	-	-	-	-	-	-
2000	951	940	5	3	1	-	1	-	-	1	-	-	-	-
1999	961	946	13	-	-	1	-	1	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	11,673	11,553	50	11	14	12	6	5	4	6	2	2	-	1
2001	11,500	11,371	58	19	14	8	7	1	1	6	3	2	2	2
2000	11,672	11,546	67	18	13	9	4	5	2	1	2	1	1	1
1999	11,815	11,668	73	23	18	8	4	1	5	2	1	2	-	1
Male														
2002	6,596	6,527	30	7	7	7	3	2	1	4	1	2	-	1
2001	6,511	6,428	38	14	8	3	4	1	-	4	2	2	2	1
2000	6,637	6,555	45	9	8	8	2	4	2	-	2	1	1	-
1999	6,680	6,589	50	12	9	4	3	-	2	2	-	2	-	-
Female														
2002	5,077	5,026	20	4	7	5	3	3	3	2	1	-	-	-
2001	4,989	4,943	20	5	6	5	3	-	1	2	1	-	-	1
2000	5,035	4,991	22	9	5	1	2	1	-	1	-	-	-	1
1999	5,135	5,079	23	11	9	4	1	1	3	-	1	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)										
All origins										
Female										
2002	-	1	1	-	1	-	-	1	-	-
2001	-	1	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	1
1999	1	1	-	-	-	-	-	-	-	1
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	1
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	1
Female										
2002	-	-	-	-	-	-	-	1	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	-	1	2	-	2	-	-	-	-	2
2001	-	2	1	1	-	1	-	-	-	1
2000	1	-	-	-	-	-	-	-	-	1
1999	2	1	1	1	-	-	1	-	-	3
Male										
2002	-	-	1	-	1	-	-	-	-	2
2001	-	1	1	1	-	-	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	1	-	-	1	-	-	3
Female										
2002	-	1	1	-	1	-	-	-	-	-
2001	-	1	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	1
1999	1	1	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain conditions originating in the perinatal period (P00-P96)														
Non-Hispanic White														
Both sexes														
2002	6,384	6,313	22	5	10	11	3	4	2	4	2	2	-	1
2001	6,323	6,239	34	11	7	5	6	1	1	6	2	2	2	1
2000	6,318	6,242	34	11	9	7	4	2	1	1	2	1	1	1
1999	6,384	6,286	37	17	14	8	3	1	5	2	1	2	-	1
Male														
2002	3,663	3,628	10	4	3	6	1	2	1	2	1	2	-	1
2001	3,574	3,518	22	9	5	2	4	1	-	4	1	2	2	-
2000	3,605	3,553	23	6	8	7	2	1	1	-	2	1	1	-
1999	3,627	3,570	24	8	7	4	3	-	2	2	-	2	-	-
Female														
2002	2,721	2,685	12	1	7	5	2	2	1	2	1	-	-	-
2001	2,749	2,721	12	2	2	3	2	-	1	2	1	-	-	1
2000	2,713	2,689	11	5	1	-	2	1	-	1	-	-	-	1
1999	2,757	2,716	13	9	7	4	-	1	3	-	1	-	-	1
Non-Hispanic Black														
Both sexes														
2002	4,777	4,735	26	4	4	1	2	1	1	1	-	-	-	-
2001	4,713	4,675	22	6	5	3	1	-	-	-	1	-	-	-
2000	4,878	4,836	27	6	4	1	-	3	1	-	-	-	-	-
1999	5,000	4,963	29	2	3	-	1	-	-	-	-	-	-	-
Male														
2002	2,640	2,611	18	2	4	1	1	-	-	1	-	-	-	-
2001	2,674	2,653	14	3	2	1	-	-	-	-	1	-	-	-
2000	2,760	2,734	18	3	-	1	-	3	1	-	-	-	-	-
1999	2,818	2,794	19	2	1	-	-	-	-	-	-	-	-	-
Female														
2002	2,137	2,124	8	2	-	-	1	1	1	-	-	-	-	-
2001	2,039	2,022	8	3	3	2	1	-	-	-	-	-	-	-
2000	2,118	2,102	9	3	4	-	-	-	-	-	-	-	-	-
1999	2,182	2,169	10	-	2	-	1	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	179	178	1	-	-	-	-	-	-	-	-	-	-	-
2001	160	160	-	-	-	-	-	-	-	-	-	-	-	-
2000	252	252	-	-	-	-	-	-	-	-	-	-	-	-
1999	291	289	1	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	99	98	1	-	-	-	-	-	-	-	-	-	-	-
2001	89	89	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)										
Non-Hispanic White										
Both sexes										
2002	-	1	1	-	2	-	-	-	-	1
2001	-	2	1	1	-	1	-	-	-	1
2000	1	-	-	-	-	-	-	-	-	1
1999	2	1	1	1	-	-	1	-	-	1
Male										
2002	-	-	-	-	1	-	-	-	-	1
2001	-	1	1	1	-	-	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	1	-	-	1	-	-	1
Female										
2002	-	1	1	-	1	-	-	-	-	-
2001	-	1	-	-	-	1	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	1
1999	1	1	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	1	-	-	-	-	-	-	1
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	2
Male										
2002	-	-	1	-	-	-	-	-	-	1
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	2
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	1
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Certain conditions originating in the perinatal period (P00-P96)														
Origin not stated[2]														
Male														
2000	137	137	-	-	-	-	-	-	-	-	-	-	-	-
1999	172	171	1	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	80	80	-	-	-	-	-	-	-	-	-	-	-	-
2001	71	71	-	-	-	-	-	-	-	-	-	-	-	-
2000	115	115	-	-	-	-	-	-	-	-	-	-	-	-
1999	119	118	-	-	-	-	-	-	-	-	-	-	-	-
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)														
All origins														
Both sexes														
2002	10,687	5,623	530	199	218	248	244	213	262	264	308	346	363	351
2001	10,442	5,513	557	182	194	255	250	225	233	259	327	303	333	298
2000	10,578	5,743	495	198	201	225	216	208	269	277	292	287	325	296
1999	10,393	5,473	549	207	221	222	212	212	253	279	273	342	305	304
Male														
2002	5,515	2,942	282	99	116	146	136	132	152	132	164	195	186	170
2001	5,448	2,954	282	101	109	161	145	121	128	137	180	156	167	146
2000	5,517	3,028	259	105	102	137	131	120	159	154	162	157	190	130
1999	5,346	2,875	310	102	113	139	123	110	132	152	144	185	162	157
Female														
2002	5,172	2,681	248	100	102	102	108	81	110	132	144	151	177	181
2001	4,994	2,559	275	81	85	94	105	104	105	122	147	147	166	152
2000	5,061	2,715	236	93	99	88	85	88	110	123	130	130	135	166
1999	5,047	2,598	239	105	108	83	89	102	121	127	129	157	143	147
Hispanic														
Both sexes														
2002	1,715	1,257	111	40	30	37	30	23	30	23	17	21	21	13
2001	1,632	1,195	103	28	30	32	41	26	19	19	23	24	12	10
2000	1,612	1,182	111	33	19	24	30	19	27	24	19	20	16	15
1999	1,498	1,072	101	31	26	27	29	28	26	15	19	26	16	13
Male														
2002	903	653	60	18	17	22	17	14	22	11	8	12	11	7
2001	892	639	54	17	19	17	22	17	9	11	16	19	6	4
2000	857	605	60	16	12	19	20	11	15	16	11	11	11	8
1999	756	534	63	16	13	17	13	15	10	8	7	13	11	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Certain conditions originating in the perinatal period (P00-P96)										
Origin not stated[2]										
Male										
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	1
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
All origins										
Both sexes										
2002	283	224	216	232	257	184	86	33	2	1
2001	283	214	234	244	221	182	87	38	10	-
2000	273	248	216	242	246	178	97	39	6	1
1999	253	251	218	249	242	187	100	34	5	2
Male										
2002	151	107	106	105	96	65	22	9	1	1
2001	130	109	119	110	93	70	17	9	4	-
2000	134	120	102	99	98	74	41	13	2	-
1999	126	106	103	102	102	60	35	6	1	1
Female										
2002	132	117	110	127	161	119	64	24	1	-
2001	153	105	115	134	128	112	70	29	6	-
2000	139	128	114	143	148	104	56	26	4	1
1999	127	145	115	147	140	127	65	28	4	1
Hispanic										
Both sexes										
2002	12	12	6	12	11	5	2	1	1	-
2001	14	10	12	17	7	6	4	-	-	-
2000	16	14	7	13	11	4	5	3	-	-
1999	14	16	14	7	9	8	1	-	-	-
Male										
2002	8	4	2	6	7	3	-	-	1	-
2001	10	5	9	8	5	5	-	-	-	-
2000	9	5	6	9	6	3	1	3	-	-
1999	2	7	10	4	3	2	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)														
Hispanic														
Female														
2002	812	604	51	22	13	15	13	9	8	12	9	9	10	6
2001	740	556	49	11	11	15	19	9	10	8	7	5	6	6
2000	755	577	51	17	7	5	10	8	12	8	8	9	5	7
1999	742	538	38	15	13	10	16	13	16	7	12	13	5	6
Non-Hispanic[1]														
Both sexes														
2002	8,904	4,316	415	159	188	210	212	190	232	241	289	324	338	337
2001	8,748	4,271	452	153	164	223	209	198	214	238	304	277	319	288
2000	8,900	4,514	384	164	181	200	186	187	240	252	273	266	308	279
1999	8,825	4,349	447	175	195	195	183	183	227	263	253	313	288	291
Male														
2002	4,577	2,262	221	81	99	124	117	118	130	121	155	182	173	163
2001	4,527	2,297	227	83	90	144	123	103	119	125	164	135	159	142
2000	4,623	2,400	199	88	89	118	111	107	143	137	151	145	178	121
1999	4,550	2,311	246	85	100	122	110	95	122	144	136	169	151	150
Female														
2002	4,327	2,054	194	78	89	86	95	72	102	120	134	142	165	174
2001	4,221	1,974	225	70	74	79	86	95	95	113	140	142	160	146
2000	4,277	2,114	185	76	92	82	75	80	97	115	122	121	130	158
1999	4,275	2,038	201	90	95	73	73	88	105	119	117	144	137	141
Non-Hispanic White														
Both sexes														
2002	6,830	3,068	281	117	136	153	162	146	185	191	232	270	292	290
2001	6,709	3,053	287	111	115	157	145	152	178	187	252	223	280	257
2000	6,792	3,199	240	113	127	134	145	144	200	201	218	224	277	238
1999	6,782	3,127	284	113	131	144	133	151	183	216	210	265	249	262
Male														
2002	3,510	1,643	151	58	67	90	88	90	105	96	127	150	145	139
2001	3,487	1,664	144	64	71	100	83	87	99	102	134	110	141	126
2000	3,519	1,720	124	55	64	74	89	82	121	112	121	120	156	105
1999	3,498	1,693	154	53	66	93	82	79	99	120	110	142	129	136
Female														
2002	3,320	1,425	130	59	69	63	74	56	80	95	105	120	147	151
2001	3,222	1,389	143	47	44	57	62	65	79	85	118	113	139	131
2000	3,273	1,479	116	58	63	60	56	62	79	89	97	104	121	133
1999	3,284	1,434	130	60	65	51	51	72	84	96	100	123	120	126

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
Hispanic										
Female										
2002	4	8	4	6	4	2	2	1	-	-
2001	4	5	3	9	2	1	4	-	-	-
2000	7	9	1	4	5	1	4	-	-	-
1999	12	9	4	3	6	6	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	271	211	210	220	245	179	84	31	1	1
2001	269	201	221	227	214	176	83	37	10	-
2000	253	234	209	227	235	174	91	36	6	1
1999	238	232	203	241	232	177	99	34	5	2
Male										
2002	143	102	104	99	89	62	22	9	-	1
2001	120	102	109	102	88	65	17	9	4	-
2000	122	115	96	89	92	71	39	10	2	-
1999	124	98	93	97	99	56	34	6	1	1
Female										
2002	128	109	106	121	156	117	62	22	1	-
2001	149	99	112	125	126	111	66	28	6	-
2000	131	119	113	138	143	103	52	26	4	1
1999	114	134	110	144	133	121	65	28	4	1
Non-Hispanic White										
Both sexes										
2002	237	189	185	200	220	165	81	28	1	1
2001	246	181	197	209	196	166	77	34	6	-
2000	222	206	184	210	221	164	88	31	5	1
1999	211	207	176	213	214	161	91	34	5	2
Male										
2002	124	94	89	89	76	58	22	8	-	1
2001	109	96	100	98	76	59	14	9	1	-
2000	112	94	83	84	88	67	38	9	1	-
1999	112	88	81	78	93	50	32	6	1	1
Female										
2002	113	95	96	111	144	107	59	20	1	-
2001	137	85	97	111	120	107	63	25	5	-
2000	110	112	101	126	133	97	50	22	4	1
1999	99	119	95	135	121	111	59	28	4	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)														
Non-Hispanic Black														
Both sexes														
2002	1,716	1,015	113	38	43	52	46	35	40	40	52	46	37	41
2001	1,674	999	127	32	41	61	52	35	32	43	46	44	32	27
2000	1,739	1,062	123	44	45	56	35	36	38	46	45	39	28	32
1999	1,680	994	137	49	56	42	40	28	32	42	34	41	32	25
Male														
2002	887	506	58	22	25	33	26	22	21	18	27	26	21	22
2001	861	525	65	15	18	41	31	11	19	18	28	18	16	13
2000	911	545	67	29	21	39	19	22	21	23	26	22	19	11
1999	880	518	79	24	29	26	24	14	18	19	22	25	17	12
Female														
2002	829	509	55	16	18	19	20	13	19	22	25	20	16	19
2001	813	474	62	17	23	20	21	24	13	25	18	26	16	14
2000	828	517	56	15	24	17	16	14	17	23	19	17	9	21
1999	800	476	58	25	27	16	16	14	14	23	12	16	15	13
Origin not stated[2]														
Both sexes														
2002	68	50	4	-	-	1	2	-	-	-	2	1	4	1
2001	62	47	2	1	-	-	-	1	-	2	-	2	2	-
2000	66	47	-	1	1	1	-	2	2	1	-	1	1	2
1999	70	52	1	1	-	-	-	1	-	1	1	3	1	-
Male														
2002	35	27	1	-	-	-	2	-	-	-	1	1	2	-
2001	29	18	1	1	-	-	-	1	-	1	-	2	2	-
2000	37	23	-	1	1	-	-	2	1	1	-	1	1	1
1999	40	30	1	1	-	-	-	-	-	-	1	3	-	-
Female														
2002	33	23	3	-	-	1	-	-	-	-	1	-	2	1
2001	33	29	1	-	-	-	-	-	-	1	-	-	-	-
2000	29	24	-	-	-	1	-	-	1	-	-	-	-	1
1999	30	22	-	-	-	-	-	1	-	1	-	-	1	-
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
All origins														
Both sexes														
2002	29,975	3,456	248	61	78	256	484	562	757	1,013	1,389	1,368	1,304	1,042
2001	32,326	3,491	272	79	102	361	588	663	867	1,379	1,747	1,882	1,622	1,192
2000	31,876	3,639	248	85	92	317	551	663	891	1,284	1,703	1,727	1,415	1,152
1999	26,672	3,538	236	65	72	228	381	446	649	940	1,122	1,168	983	908

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
Non-Hispanic Black										
Both sexes										
2002	28	20	20	14	18	13	2	3	-	-
2001	21	14	21	17	11	7	5	3	4	-
2000	26	25	18	14	11	7	3	5	1	-
1999	21	21	25	22	16	15	8	-	-	-
Male										
2002	18	8	13	7	9	4	-	1	-	-
2001	10	4	8	4	8	4	2	-	3	-
2000	6	18	8	5	4	3	1	1	1	-
1999	10	6	10	14	5	6	2	-	-	-
Female										
2002	10	12	7	7	9	9	2	2	-	-
2001	11	10	13	13	3	3	3	3	1	-
2000	20	7	10	9	7	4	2	4	-	-
1999	11	15	15	8	11	9	6	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	1	-	-	1	-	-	1	-	-
2001	-	3	1	-	-	-	-	1	-	-
2000	4	-	-	2	-	-	1	-	-	-
1999	1	3	1	1	1	2	-	-	-	-
Male										
2002	-	1	-	-	-	-	-	-	-	-
2001	-	2	1	-	-	-	-	-	-	-
2000	3	-	-	1	-	-	1	-	-	-
1999	-	1	-	1	-	2	-	-	-	-
Female										
2002	-	-	-	-	1	-	-	1	-	-
2001	-	1	-	-	-	-	-	1	-	-
2000	1	-	-	1	-	-	-	-	-	-
1999	1	2	1	-	1	-	-	-	-	-
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All origins										
Both sexes										
2002	960	995	1,403	2,103	2,833	3,350	3,406	1,944	854	109
2001	1,100	1,207	1,567	2,209	2,794	3,239	3,220	1,882	748	115
2000	1,101	1,353	1,687	2,235	2,794	3,286	3,014	1,798	726	115
1999	952	1,140	1,558	1,945	2,468	2,818	2,799	1,592	589	75

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
All origins														
Male														
2002	14,269	2,032	145	37	45	172	329	391	514	670	870	895	828	652
2001	15,779	2,041	162	44	55	249	405	461	579	888	1,119	1,220	1,066	758
2000	15,863	2,156	137	48	51	196	385	456	604	857	1,137	1,157	966	743
1999	12,877	2,072	136	35	42	153	269	287	434	599	740	796	650	578
Female														
2002	15,706	1,424	103	24	33	84	155	171	243	343	519	473	476	390
2001	16,547	1,450	110	35	47	112	183	202	288	491	628	662	556	434
2000	16,013	1,483	111	37	41	121	166	207	287	427	566	570	449	409
1999	13,795	1,466	100	30	30	75	112	159	215	341	382	372	333	330
Hispanic														
Both sexes														
2002	1,548	472	40	4	7	34	57	68	81	81	96	65	79	44
2001	2,190	496	47	16	20	73	111	132	154	175	152	173	110	71
2000	2,071	506	51	15	16	50	90	118	142	177	165	144	107	72
1999	1,281	415	30	6	10	30	53	58	80	80	68	64	29	29
Male														
2002	926	283	26	3	4	20	42	51	65	60	60	40	45	26
2001	1,410	280	28	8	7	56	83	105	116	130	109	135	77	50
2000	1,369	310	31	7	9	29	73	93	108	137	125	110	82	41
1999	768	237	14	5	5	21	43	42	57	59	51	52	16	19
Female														
2002	622	189	14	1	3	14	15	17	16	21	36	25	34	18
2001	780	216	19	8	13	17	28	27	38	45	43	38	33	21
2000	702	196	20	8	7	21	17	25	34	40	40	34	25	31
1999	513	178	16	1	5	9	10	16	23	21	17	12	13	10
Non-Hispanic[1]														
Both sexes														
2002	28,211	2,968	207	56	70	220	423	492	670	925	1,281	1,292	1,212	993
2001	29,931	2,978	225	62	82	287	470	520	707	1,201	1,581	1,694	1,499	1,118
2000	29,514	3,108	196	69	75	265	455	540	739	1,091	1,513	1,567	1,282	1,058
1999	25,181	3,097	204	59	62	198	323	385	566	848	1,037	1,087	936	868
Male														
2002	13,206	1,739	119	34	40	150	284	338	445	605	803	849	770	621
2001	14,246	1,750	134	36	48	193	317	350	457	757	997	1,076	982	705
2000	14,288	1,834	106	41	41	166	308	360	488	708	992	1,037	864	683
1999	11,962	1,817	121	30	37	132	223	244	375	532	677	729	618	550

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
All origins										
Male										
2002	621	563	761	1,010	1,218	1,103	857	379	100	77
2001	699	685	812	1,014	1,170	1,017	760	392	105	78
2000	686	832	876	1,100	1,144	1,078	733	331	100	90
1999	603	665	848	949	981	894	699	306	85	56
Female										
2002	339	432	642	1,093	1,615	2,247	2,549	1,565	754	32
2001	401	522	755	1,195	1,624	2,222	2,460	1,490	643	37
2000	415	521	811	1,135	1,650	2,208	2,281	1,467	626	25
1999	349	475	710	996	1,487	1,924	2,100	1,286	504	19
Hispanic										
Both sexes										
2002	51	36	47	44	68	59	59	30	17	9
2001	56	55	57	51	56	72	50	32	19	12
2000	42	49	58	54	50	61	51	25	17	11
1999	34	36	37	43	50	47	43	21	13	5
Male										
2002	33	23	27	24	33	25	19	8	1	8
2001	42	36	35	27	29	22	8	13	3	11
2000	31	35	33	27	25	28	19	4	2	10
1999	23	22	16	28	18	11	15	9	1	4
Female										
2002	18	13	20	20	35	34	40	22	16	1
2001	14	19	22	24	27	50	42	19	16	1
2000	11	14	25	27	25	33	32	21	15	1
1999	11	14	21	15	32	36	28	12	12	1
Non-Hispanic[1]										
Both sexes										
2002	897	954	1,353	2,050	2,754	3,282	3,332	1,904	833	43
2001	1,036	1,149	1,504	2,149	2,737	3,158	3,168	1,843	727	36
2000	1,051	1,291	1,617	2,171	2,737	3,216	2,960	1,770	708	35
1999	912	1,090	1,511	1,896	2,414	2,765	2,752	1,567	574	30
Male										
2002	577	535	732	981	1,175	1,074	836	369	97	33
2001	650	648	774	984	1,141	993	750	376	102	26
2000	648	787	834	1,064	1,114	1,047	713	326	97	30
1999	575	638	825	916	960	881	683	296	84	19

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
Non-Hispanic[1]														
Female														
2002	15,005	1,229	88	22	30	70	139	154	225	320	478	443	442	372
2001	15,685	1,228	91	26	34	94	153	170	250	444	584	618	517	413
2000	15,226	1,274	90	28	34	99	147	180	251	383	521	530	418	375
1999	13,219	1,280	83	29	25	66	100	141	191	316	360	358	318	318
Non-Hispanic White														
Both sexes														
2002	23,099	1,777	124	30	41	157	311	354	495	689	975	955	867	770
2001	24,596	1,752	139	37	52	217	346	380	527	914	1,261	1,342	1,168	884
2000	24,064	1,805	134	41	47	191	327	374	544	849	1,178	1,211	962	822
1999	20,200	1,812	129	43	38	133	215	253	407	613	728	773	686	658
Male														
2002	10,432	1,056	66	19	25	108	218	252	329	454	617	629	548	478
2001	11,404	1,046	79	18	34	143	233	271	343	573	814	857	777	548
2000	11,351	1,067	79	28	24	128	228	255	370	568	798	815	650	520
1999	9,249	1,083	82	21	24	89	155	168	263	400	484	509	454	413
Female														
2002	12,667	721	58	11	16	49	93	102	166	235	358	326	319	292
2001	13,192	706	60	19	18	74	113	109	184	341	447	485	391	336
2000	12,713	738	55	13	23	63	99	119	174	281	380	396	312	302
1999	10,951	729	47	22	14	44	60	85	144	213	244	264	232	245
Non-Hispanic Black														
Both sexes														
2002	4,556	1,070	66	24	26	57	95	119	156	216	274	302	324	201
2001	4,606	1,081	69	20	22	52	106	121	145	239	279	315	290	202
2000	4,892	1,186	54	21	25	65	114	140	168	208	306	318	291	207
1999	4,508	1,149	67	15	18	54	98	114	146	225	279	294	226	192
Male														
2002	2,482	615	44	14	14	39	53	71	103	138	167	198	209	129
2001	2,414	614	44	15	11	37	72	67	83	144	156	192	172	138
2000	2,614	694	26	11	17	32	73	89	102	113	170	194	196	146
1999	2,460	658	35	9	11	38	62	67	105	127	172	205	147	128
Female														
2002	2,074	455	22	10	12	18	42	48	53	78	107	104	115	72
2001	2,192	467	25	5	11	15	34	54	62	95	123	123	118	64
2000	2,278	492	28	10	8	33	41	51	66	95	136	124	95	61
1999	2,048	491	32	6	7	16	36	47	41	98	107	89	79	64

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
Non-Hispanic[1]										
Female										
2002	320	419	621	1,069	1,579	2,208	2,496	1,535	736	10
2001	386	501	730	1,165	1,596	2,165	2,418	1,467	625	10
2000	403	504	783	1,107	1,623	2,169	2,247	1,444	611	5
1999	337	452	686	980	1,454	1,884	2,069	1,271	490	11
Non-Hispanic White										
Both sexes										
2002	723	769	1,165	1,795	2,480	3,012	3,082	1,762	735	31
2001	829	921	1,270	1,897	2,453	2,907	2,926	1,693	652	29
2000	830	1,016	1,378	1,903	2,455	2,946	2,748	1,652	629	22
1999	716	878	1,288	1,634	2,141	2,533	2,552	1,442	506	22
Male										
2002	467	433	632	844	1,053	997	772	337	74	24
2001	510	517	667	885	1,035	912	686	342	92	22
2000	516	618	710	939	1,006	958	667	303	85	19
1999	457	514	708	793	845	797	625	271	78	16
Female										
2002	256	336	533	951	1,427	2,015	2,310	1,425	661	7
2001	319	404	603	1,012	1,418	1,995	2,240	1,351	560	7
2000	314	398	668	964	1,449	1,988	2,081	1,349	544	3
1999	259	364	580	841	1,296	1,736	1,927	1,171	428	6
Non-Hispanic Black										
Both sexes										
2002	155	165	170	226	245	235	213	120	87	10
2001	171	197	204	221	245	212	214	131	65	5
2000	204	250	222	247	258	238	184	95	78	13
1999	179	194	201	238	250	206	180	113	64	6
Male										
2002	100	89	91	122	112	66	52	28	21	7
2001	124	114	93	87	87	69	54	28	10	3
2000	120	154	113	115	99	75	39	14	11	11
1999	110	113	105	113	101	71	53	22	6	2
Female										
2002	55	76	79	104	133	169	161	92	66	3
2001	47	83	111	134	158	143	160	103	55	2
2000	84	96	109	132	159	163	145	81	67	2
1999	69	81	96	125	149	135	127	91	58	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)														
Origin not stated[2]														
Both sexes														
2002	216	16	1	1	1	2	4	2	6	7	12	11	13	5
2001	205	17	-	1	-	1	7	11	6	3	14	15	13	3
2000	291	25	1	1	1	2	6	5	10	16	25	16	26	22
1999	210	26	2	-	-	-	5	3	3	12	17	17	18	11
Male														
2002	137	10	-	-	1	2	3	2	4	5	7	6	13	5
2001	123	11	-	-	-	-	5	6	6	1	13	9	7	3
2000	206	12	-	-	1	1	4	3	8	12	20	10	20	19
1999	147	18	1	-	-	-	3	1	2	8	12	15	16	9
Female														
2002	79	6	1	1	-	-	1	-	2	2	5	5	-	-
2001	82	6	-	1	-	1	2	5	-	2	1	6	6	-
2000	85	13	1	1	-	1	2	2	2	4	5	6	6	3
1999	63	8	1	-	-	-	2	2	1	4	5	2	2	2
All other diseases (Residual)														
All origins														
Both sexes														
2002	194,591	1,144	661	386	482	717	1,136	1,256	1,880	3,099	5,407	7,159	7,992	8,431
2001	186,406	1,131	646	389	471	772	1,027	1,225	1,847	3,145	4,939	6,667	7,536	7,549
2000	178,378	1,105	609	376	503	732	1,032	1,257	1,840	3,161	4,952	6,227	7,099	7,152
1999	171,683	1,148	647	436	485	785	986	1,235	1,827	3,203	4,786	5,877	6,506	6,676
Male														
2002	79,007	690	363	220	266	423	649	714	1,123	1,800	3,219	4,305	4,810	4,675
2001	75,162	649	358	187	267	463	601	699	1,030	1,869	2,905	3,970	4,427	4,112
2000	72,227	629	338	198	276	432	625	737	1,074	1,873	2,928	3,811	4,100	3,967
1999	69,784	680	354	229	254	483	562	717	1,073	1,912	2,923	3,510	3,762	3,762
Female														
2002	115,584	454	298	166	216	294	487	542	757	1,299	2,188	2,854	3,182	3,756
2001	111,244	482	288	202	204	309	426	526	817	1,276	2,034	2,697	3,109	3,437
2000	106,151	476	271	178	227	300	407	520	766	1,288	2,024	2,416	2,999	3,185
1999	101,899	468	293	207	231	302	424	518	754	1,291	1,863	2,367	2,744	2,914
Hispanic														
Both sexes														
2002	8,377	209	120	74	77	100	150	173	236	307	450	530	476	493
2001	7,847	187	112	63	59	102	144	162	192	289	382	438	454	444
2000	7,325	196	109	57	56	82	132	168	213	311	395	440	422	434
1999	7,291	200	106	50	57	95	123	166	243	306	384	437	400	395

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)										
Origin not stated[2]										
Both sexes										
2002	12	5	3	9	11	9	15	10	4	57
2001	8	3	6	9	1	9	2	7	2	67
2000	8	13	12	10	7	9	3	3	1	69
1999	6	14	10	6	4	6	4	4	2	40
Male										
2002	11	5	2	5	10	4	2	2	2	36
2001	7	1	3	3	-	2	2	3	-	41
2000	7	10	9	9	5	3	1	1	1	50
1999	5	5	7	5	3	2	1	1	-	33
Female										
2002	1	-	1	4	1	5	13	8	2	21
2001	1	2	3	6	1	7	-	4	2	26
2000	1	3	3	1	2	6	2	2	-	19
1999	1	9	3	1	1	4	3	3	2	7
All other diseases (Residual)										
All origins										
Both sexes										
2002	8,947	10,576	15,113	22,098	29,520	31,671	24,416	10,231	2,256	13
2001	8,499	10,676	15,127	21,941	27,770	30,312	22,714	9,872	2,134	17
2000	8,004	10,108	14,899	21,425	26,594	28,517	21,457	9,192	2,126	11
1999	7,714	10,272	14,893	20,657	24,941	27,390	20,562	8,680	1,967	10
Male										
2002	4,853	5,453	7,214	9,509	11,158	9,835	5,691	1,711	315	11
2001	4,540	5,480	7,252	9,406	10,411	9,265	5,258	1,740	257	16
2000	4,269	5,174	7,147	9,237	9,865	8,712	4,966	1,593	265	11
1999	4,108	5,154	7,178	8,976	9,519	8,208	4,722	1,443	250	5
Female										
2002	4,094	5,123	7,899	12,589	18,362	21,836	18,725	8,520	1,941	2
2001	3,959	5,196	7,875	12,535	17,359	21,047	17,456	8,132	1,877	1
2000	3,735	4,934	7,752	12,188	16,729	19,805	16,491	7,599	1,861	-
1999	3,606	5,118	7,715	11,681	15,422	19,182	15,840	7,237	1,717	5
Hispanic										
Both sexes										
2002	465	597	716	796	836	749	543	240	39	1
2001	474	529	689	741	798	781	516	236	52	3
2000	379	505	608	711	727	685	427	215	49	4
1999	447	513	620	627	638	728	494	213	47	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other diseases (Residual)														
Hispanic														
Male														
2002	4,221	116	66	47	42	54	94	109	152	201	316	355	292	283
2001	3,849	106	66	31	34	65	82	97	125	197	254	295	289	259
2000	3,663	100	54	29	34	54	86	100	146	211	272	295	257	270
1999	3,655	124	50	28	26	59	69	103	172	218	261	296	245	230
Female														
2002	4,156	93	54	27	35	46	56	64	84	106	134	175	184	210
2001	3,998	81	46	32	25	37	62	65	67	92	128	143	165	185
2000	3,662	96	55	28	22	28	46	68	67	100	123	145	165	164
1999	3,636	76	56	22	31	36	54	63	71	88	123	141	155	165
Non-Hispanic[1]														
Both sexes														
2002	185,608	927	538	311	404	615	985	1,076	1,638	2,774	4,937	6,591	7,469	7,898
2001	177,985	934	532	325	411	667	878	1,060	1,649	2,842	4,528	6,192	7,038	7,064
2000	170,462	898	497	317	445	646	895	1,080	1,617	2,836	4,530	5,738	6,638	6,685
1999	163,832	935	537	386	428	688	859	1,068	1,577	2,878	4,369	5,397	6,070	6,247
Male														
2002	74,500	572	294	172	224	368	555	601	967	1,589	2,886	3,923	4,487	4,366
2001	71,014	539	291	156	232	396	516	599	900	1,664	2,629	3,645	4,106	3,822
2000	68,263	522	282	169	242	374	537	630	926	1,650	2,637	3,481	3,813	3,680
1999	65,835	546	302	201	228	423	490	613	896	1,679	2,637	3,186	3,490	3,509
Female														
2002	111,108	355	244	139	180	247	430	475	671	1,185	2,051	2,668	2,982	3,532
2001	106,971	395	241	169	179	271	362	461	749	1,178	1,899	2,547	2,932	3,242
2000	102,199	376	215	148	203	272	358	450	691	1,186	1,893	2,257	2,825	3,005
1999	97,997	389	235	185	200	265	369	455	681	1,199	1,732	2,211	2,580	2,738
Non-Hispanic White														
Both sexes														
2002	159,945	578	352	221	284	451	651	720	1,111	1,938	3,607	4,895	5,672	6,294
2001	153,156	556	362	246	295	474	615	699	1,109	1,983	3,218	4,511	5,303	5,553
2000	146,518	544	316	219	318	476	637	707	1,100	1,955	3,247	4,107	5,079	5,241
1999	141,002	554	375	265	304	513	587	749	1,096	1,949	3,087	3,858	4,579	4,912
Male														
2002	63,109	376	191	124	155	284	393	420	693	1,179	2,207	2,985	3,479	3,499
2001	60,022	311	209	121	165	294	383	411	629	1,208	1,922	2,727	3,146	3,007
2000	57,565	310	183	123	184	286	397	446	659	1,183	1,945	2,600	2,969	2,918
1999	55,517	321	214	136	155	332	349	456	654	1,194	1,950	2,344	2,685	2,786
Female														
2002	96,836	202	161	97	129	167	258	300	418	759	1,400	1,910	2,193	2,795
2001	93,134	245	153	125	130	180	232	288	480	775	1,296	1,784	2,157	2,546

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases (Residual)										
Hispanic										
Male										
2002	244	322	337	365	342	258	156	60	9	1
2001	254	271	316	301	311	280	130	67	16	3
2000	194	284	284	300	257	230	128	59	15	4
1999	251	250	314	282	248	227	132	60	9	1
Female										
2002	221	275	379	431	494	491	387	180	30	-
2001	220	258	373	440	487	501	386	169	36	-
2000	185	221	324	411	470	455	299	156	34	-
1999	196	263	306	345	390	501	362	153	38	1
Non-Hispanic[1]										
Both sexes										
2002	8,449	9,948	14,359	21,240	28,595	30,850	23,814	9,967	2,215	8
2001	7,999	10,110	14,396	21,150	26,909	29,462	22,146	9,608	2,074	11
2000	7,598	9,566	14,233	20,649	25,809	27,767	20,978	8,964	2,071	5
1999	7,232	9,721	14,235	19,977	24,255	26,591	20,022	8,440	1,915	5
Male										
2002	4,587	5,108	6,856	9,121	10,784	9,559	5,524	1,644	306	7
2001	4,265	5,189	6,915	9,077	10,078	8,966	5,111	1,667	241	10
2000	4,054	4,868	6,832	8,905	9,588	8,459	4,827	1,533	249	5
1999	3,832	4,882	6,841	8,672	9,252	7,956	4,579	1,377	241	3
Female										
2002	3,862	4,840	7,503	12,119	17,811	21,291	18,290	8,323	1,909	1
2001	3,734	4,921	7,481	12,073	16,831	20,496	17,035	7,941	1,833	1
2000	3,544	4,698	7,401	11,744	16,221	19,308	16,151	7,431	1,822	-
1999	3,400	4,839	7,394	11,305	15,003	18,635	15,443	7,063	1,674	2
Non-Hispanic White										
Both sexes										
2002	6,822	8,269	12,261	18,720	25,769	28,310	22,014	9,077	1,924	5
2001	6,429	8,415	12,392	18,647	24,343	26,949	20,435	8,830	1,786	6
2000	6,091	7,954	12,234	18,262	23,300	25,412	19,296	8,212	1,809	2
1999	5,837	8,027	12,240	17,624	22,039	24,396	18,595	7,727	1,688	1
Male										
2002	3,731	4,242	5,867	8,035	9,713	8,765	5,056	1,461	249	5
2001	3,469	4,333	5,979	7,992	9,148	8,203	4,674	1,494	192	5
2000	3,231	4,051	5,882	7,874	8,645	7,710	4,375	1,382	210	2
1999	3,120	4,054	5,923	7,621	8,392	7,226	4,177	1,221	206	1
Female										
2002	3,091	4,027	6,394	10,685	16,056	19,545	16,958	7,616	1,675	-
2001	2,960	4,082	6,413	10,655	15,195	18,746	15,761	7,336	1,594	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
All other diseases (Residual)														
Non-Hispanic White														
Female														
2000	88,953	234	133	96	134	190	240	261	441	772	1,302	1,507	2,110	2,323
1999	85,485	233	161	129	149	181	238	293	442	755	1,137	1,514	1,894	2,126
Non-Hispanic Black														
Both sexes														
2002	22,297	298	146	72	100	133	283	317	458	735	1,200	1,528	1,619	1,416
2001	21,752	329	131	65	100	163	223	301	457	770	1,174	1,529	1,577	1,341
2000	21,081	296	144	79	110	146	227	343	467	789	1,159	1,495	1,402	1,278
1999	20,137	332	129	100	101	144	241	282	422	839	1,158	1,399	1,329	1,195
Male														
2002	9,722	167	80	37	58	66	139	155	233	354	594	834	899	771
2001	9,561	192	62	26	57	86	110	154	227	403	627	838	875	722
2000	9,351	180	77	35	52	76	121	165	235	412	621	804	758	689
1999	8,967	192	67	56	62	75	127	135	209	434	606	765	722	646
Female														
2002	12,575	131	66	35	42	67	144	162	225	381	606	694	720	645
2001	12,191	137	69	39	43	77	113	147	230	367	547	691	702	619
2000	11,730	116	67	44	58	70	106	178	232	377	538	691	644	589
1999	11,170	140	62	44	39	69	114	147	213	405	552	634	607	549
Origin not stated[2]														
Both sexes														
2002	606	8	3	1	1	2	1	7	6	18	20	38	47	40
2001	574	10	2	1	1	3	5	3	6	14	29	37	44	41
2000	591	11	3	2	2	4	5	9	10	14	27	49	39	33
1999	560	13	4	-	-	2	4	1	7	19	33	43	36	34
Male														
2002	286	2	3	1	-	1	-	4	4	10	17	27	31	26
2001	299	4	1	-	1	2	3	3	5	8	22	30	32	31
2000	301	7	2	-	-	4	2	7	2	12	19	35	30	17
1999	294	10	2	-	-	1	3	1	5	15	25	28	27	23
Female														
2002	320	6	-	-	1	1	1	3	2	8	3	11	16	14
2001	275	6	1	1	-	1	2	-	1	6	7	7	12	10
2000	290	4	1	2	2	-	3	2	8	2	8	14	9	16
1999	266	3	2	-	-	1	1	-	2	4	8	15	9	11

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases (Residual)										
Non-Hispanic White										
Female										
2000	2,860	3,903	6,352	10,388	14,655	17,702	14,921	6,830	1,599	-
1999	2,717	3,973	6,317	10,003	13,647	17,170	14,418	6,506	1,482	-
Non-Hispanic Black										
Both sexes										
2002	1,411	1,450	1,797	2,128	2,431	2,161	1,578	778	255	3
2001	1,350	1,498	1,738	2,170	2,245	2,160	1,480	684	262	5
2000	1,329	1,412	1,741	2,075	2,173	2,054	1,466	656	237	3
1999	1,242	1,498	1,753	2,053	1,910	1,903	1,273	633	197	4
Male										
2002	730	735	856	898	897	646	389	138	44	2
2001	689	767	803	948	819	622	346	141	42	5
2000	722	717	831	897	814	618	360	130	34	3
1999	624	725	793	909	721	608	335	130	24	2
Female										
2002	681	715	941	1,230	1,534	1,515	1,189	640	211	1
2001	661	731	935	1,222	1,426	1,538	1,134	543	220	-
2000	607	695	910	1,178	1,359	1,436	1,106	526	203	-
1999	618	773	960	1,144	1,189	1,295	938	503	173	2
Origin not stated[2]										
Both sexes										
2002	33	31	38	62	89	72	59	24	2	4
2001	26	37	42	50	63	69	52	28	8	3
2000	27	37	58	65	58	65	52	13	6	2
1999	35	38	38	53	48	71	46	27	5	3
Male										
2002	22	23	21	23	32	18	11	7	-	3
2001	21	20	21	28	22	19	17	6	-	3
2000	21	22	31	32	20	23	11	1	1	2
1999	25	22	23	22	19	25	11	6	-	1
Female										
2002	11	8	17	39	57	54	48	17	2	1
2001	5	17	21	22	41	50	35	22	8	-
2000	6	15	27	33	38	42	41	12	5	-
1999	10	16	15	31	29	46	35	21	5	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)														
All origins														
Both sexes														
2002	106,742	946	1,641	1,176	1,542	7,137	8,275	6,122	6,447	7,642	9,068	8,241	6,434	4,557
2001	101,537	976	1,714	1,283	1,553	6,646	7,765	5,935	5,904	7,589	8,356	7,469	5,875	4,159
2000	97,900	881	1,826	1,391	1,588	6,755	7,358	5,898	5,871	7,471	7,942	6,870	5,408	4,021
1999	97,860	845	1,898	1,459	1,632	6,688	6,968	5,881	6,009	7,366	7,865	6,696	4,943	3,944
Male														
2002	69,257	553	1,024	720	993	4,981	6,457	4,768	4,867	5,535	6,477	5,940	4,552	3,255
2001	66,060	541	1,045	786	1,017	4,722	6,110	4,605	4,487	5,502	6,110	5,519	4,224	2,872
2000	63,817	501	1,098	824	1,037	4,757	5,703	4,606	4,453	5,537	5,907	5,103	3,960	2,804
1999	63,535	494	1,127	897	1,077	4,624	5,416	4,555	4,542	5,411	5,837	4,996	3,606	2,753
Female														
2002	37,485	393	617	456	549	2,156	1,818	1,354	1,580	2,107	2,591	2,301	1,882	1,302
2001	35,477	435	669	497	536	1,924	1,655	1,330	1,417	2,087	2,246	1,950	1,651	1,287
2000	34,083	380	728	567	551	1,998	1,655	1,292	1,418	1,934	2,035	1,767	1,448	1,217
1999	34,325	351	771	562	555	2,064	1,552	1,326	1,467	1,955	2,028	1,700	1,337	1,191
Hispanic														
Both sexes														
2002	10,106	127	341	199	190	984	1,429	1,082	945	898	943	685	483	311
2001	9,523	154	310	186	193	879	1,303	1,073	865	942	818	627	451	313
2000	8,830	118	306	205	178	884	1,211	944	832	856	762	542	411	272
1999	8,650	134	341	170	192	800	1,146	936	854	844	763	546	380	250
Male														
2002	7,698	83	215	120	124	763	1,195	917	792	712	756	522	366	235
2001	7,157	75	195	114	110	669	1,104	908	711	763	655	481	341	213
2000	6,696	67	174	105	105	689	1,028	804	695	682	635	435	315	193
1999	6,578	80	197	107	120	608	967	787	707	692	615	443	291	183
Female														
2002	2,408	44	126	79	66	221	234	165	153	186	187	163	117	76
2001	2,366	79	115	72	83	210	199	165	154	179	163	146	110	100
2000	2,134	51	132	100	73	195	183	140	137	174	127	107	96	79
1999	2,072	54	144	63	72	192	179	149	147	152	148	103	89	67
Non-Hispanic[1]														
Both sexes														
2002	96,175	816	1,296	972	1,346	6,136	6,820	5,019	5,470	6,705	8,087	7,509	5,913	4,214
2001	91,522	817	1,399	1,091	1,354	5,748	6,439	4,846	5,011	6,602	7,487	6,793	5,391	3,816
2000	88,558	759	1,512	1,180	1,405	5,848	6,120	4,927	5,008	6,580	7,126	6,274	4,945	3,723
1999	88,669	710	1,550	1,286	1,435	5,865	5,793	4,913	5,121	6,481	7,048	6,099	4,523	3,652
Male														
2002	61,201	468	806	596	865	4,208	5,240	3,835	4,047	4,790	5,692	5,380	4,157	2,988
2001	58,528	462	845	668	902	4,038	4,986	3,684	3,754	4,701	5,416	4,999	3,853	2,636

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All origins										
Both sexes										
2002	3,788	3,672	4,414	5,938	6,966	6,507	4,255	1,595	294	85
2001	3,499	3,506	4,329	5,972	6,716	6,266	4,074	1,519	312	120
2000	3,484	3,375	4,323	5,596	6,162	5,933	3,939	1,419	304	85
1999	3,341	3,692	4,516	5,772	6,510	6,138	3,954	1,380	257	106
Male										
2002	2,526	2,369	2,662	3,268	3,419	2,795	1,488	480	54	74
2001	2,356	2,200	2,551	3,310	3,342	2,730	1,486	381	60	104
2000	2,329	2,117	2,522	3,039	3,024	2,563	1,391	397	62	83
1999	2,262	2,336	2,656	3,096	3,276	2,647	1,397	376	60	94
Female										
2002	1,262	1,303	1,752	2,670	3,547	3,712	2,767	1,115	240	11
2001	1,143	1,306	1,778	2,662	3,374	3,536	2,588	1,138	252	16
2000	1,155	1,258	1,801	2,557	3,138	3,370	2,548	1,022	242	2
1999	1,079	1,356	1,860	2,676	3,234	3,491	2,557	1,004	197	12
Hispanic										
Both sexes										
2002	255	228	238	262	202	130	92	47	7	28
2001	251	205	249	222	187	140	84	36	6	29
2000	220	187	231	193	172	146	94	30	8	28
1999	216	245	195	183	151	140	83	28	9	44
Male										
2002	172	166	141	152	106	66	43	21	4	27
2001	178	130	145	136	99	59	35	8	2	26
2000	154	126	145	117	74	69	42	11	3	28
1999	151	162	126	100	82	67	37	15	1	40
Female										
2002	83	62	97	110	96	64	49	26	3	1
2001	73	75	104	86	88	81	49	28	4	3
2000	66	61	86	76	98	77	52	19	5	-
1999	65	83	69	83	69	73	46	13	8	4
Non-Hispanic[1]										
Both sexes										
2002	3,514	3,428	4,156	5,655	6,739	6,362	4,154	1,544	287	33
2001	3,226	3,286	4,061	5,738	6,507	6,114	3,980	1,475	305	36
2000	3,243	3,175	4,072	5,386	5,975	5,771	3,836	1,386	295	12
1999	3,101	3,430	4,308	5,567	6,329	5,977	3,859	1,348	248	26
Male										
2002	2,337	2,189	2,504	3,105	3,293	2,720	1,444	457	50	30
2001	2,159	2,062	2,393	3,166	3,231	2,667	1,447	372	58	29

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)														
Non-Hispanic[1]														
Male														
2000	56,736	431	921	717	931	4,053	4,654	3,781	3,731	4,826	5,228	4,626	3,605	2,590
1999	56,551	414	926	787	952	4,003	4,425	3,745	3,805	4,687	5,184	4,510	3,281	2,537
Female														
2002	34,974	348	490	376	481	1,928	1,580	1,184	1,423	1,915	2,395	2,129	1,756	1,226
2001	32,994	355	554	423	452	1,710	1,453	1,162	1,257	1,901	2,071	1,794	1,538	1,180
2000	31,822	328	591	463	474	1,795	1,466	1,146	1,277	1,754	1,898	1,648	1,340	1,133
1999	32,118	296	624	499	483	1,862	1,368	1,168	1,316	1,794	1,864	1,589	1,242	1,115
Non-Hispanic White														
Both sexes														
2002	80,605	462	865	616	965	5,099	5,565	3,921	4,423	5,358	6,677	6,080	4,779	3,459
2001	76,262	500	1,006	722	966	4,692	5,175	3,819	3,979	5,314	6,026	5,361	4,343	3,132
2000	73,536	455	1,033	796	1,047	4,852	4,883	3,788	3,942	5,302	5,706	4,978	3,904	3,044
1999	73,326	435	1,042	843	1,065	4,840	4,570	3,854	4,001	5,186	5,658	4,789	3,603	2,968
Male														
2002	50,652	279	527	378	619	3,466	4,280	2,992	3,281	3,822	4,728	4,412	3,333	2,427
2001	48,180	281	624	424	642	3,266	4,031	2,908	2,998	3,794	4,394	3,941	3,093	2,142
2000	46,470	266	638	487	689	3,326	3,739	2,899	2,962	3,914	4,199	3,680	2,841	2,089
1999	46,113	258	617	505	700	3,250	3,499	2,953	2,979	3,770	4,179	3,531	2,611	2,040
Female														
2002	29,953	183	338	238	346	1,633	1,285	929	1,142	1,536	1,949	1,668	1,446	1,032
2001	28,082	219	382	298	324	1,426	1,144	911	981	1,520	1,632	1,420	1,250	990
2000	27,066	189	395	309	358	1,526	1,144	889	980	1,388	1,507	1,298	1,063	955
1999	27,213	177	425	338	365	1,590	1,071	901	1,022	1,416	1,479	1,258	992	928
Non-Hispanic Black														
Both sexes														
2002	12,285	324	350	298	309	740	940	853	823	1,084	1,143	1,211	934	618
2001	12,239	272	331	313	317	760	953	784	816	1,026	1,256	1,212	891	568
2000	12,087	268	396	330	289	721	941	877	863	1,042	1,211	1,096	865	561
1999	12,533	251	428	392	309	764	930	841	897	1,063	1,182	1,138	765	562
Male														
2002	8,428	172	231	180	211	541	719	665	614	786	776	836	687	470
2001	8,381	158	188	212	220	564	721	592	600	718	890	918	656	416
2000	8,380	145	232	204	202	530	703	681	635	759	890	808	646	429
1999	8,629	140	260	254	213	568	713	647	666	754	874	864	577	418
Female														
2002	3,857	152	119	118	98	199	221	188	209	298	367	375	247	148
2001	3,858	114	143	101	97	196	232	192	216	308	366	294	235	152
2000	3,707	123	164	126	87	191	238	196	228	283	321	288	219	132

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Non-Hispanic[1]										
Male										
2000	2,157	1,979	2,365	2,913	2,942	2,487	1,344	385	59	11
1999	2,090	2,161	2,520	2,984	3,174	2,571	1,354	359	59	23
Female										
2002	1,177	1,239	1,652	2,550	3,446	3,642	2,710	1,087	237	3
2001	1,067	1,224	1,668	2,572	3,276	3,447	2,533	1,103	247	7
2000	1,086	1,196	1,707	2,473	3,033	3,284	2,492	1,001	236	1
1999	1,011	1,269	1,788	2,583	3,155	3,406	2,505	989	189	3
Non-Hispanic White										
Both sexes										
2002	2,934	2,840	3,641	5,082	6,234	5,977	3,894	1,448	260	26
2001	2,644	2,784	3,525	5,160	5,997	5,711	3,729	1,376	276	25
2000	2,709	2,659	3,522	4,834	5,522	5,410	3,594	1,284	264	8
1999	2,487	2,875	3,693	4,946	5,811	5,537	3,625	1,260	225	13
Male										
2002	1,947	1,795	2,196	2,778	3,027	2,544	1,332	425	41	23
2001	1,763	1,748	2,089	2,838	2,962	2,482	1,348	339	51	22
2000	1,782	1,636	2,056	2,589	2,703	2,326	1,246	345	51	7
1999	1,690	1,805	2,157	2,632	2,919	2,368	1,251	334	53	12
Female										
2002	987	1,045	1,445	2,304	3,207	3,433	2,562	1,023	219	3
2001	881	1,036	1,436	2,322	3,035	3,229	2,381	1,037	225	3
2000	927	1,023	1,466	2,245	2,819	3,084	2,348	939	213	1
1999	797	1,070	1,536	2,314	2,892	3,169	2,374	926	172	1
Non-Hispanic Black										
Both sexes										
2002	457	435	383	433	366	289	195	76	20	4
2001	477	415	421	447	383	295	196	73	25	8
2000	422	414	443	419	349	279	189	80	28	4
1999	485	454	489	499	426	349	197	81	19	12
Male										
2002	321	295	232	254	194	129	81	25	5	4
2001	334	268	237	256	205	122	73	21	6	6
2000	307	282	255	250	182	120	77	32	7	4
1999	324	297	299	282	201	162	79	22	5	10
Female										
2002	136	140	151	179	172	160	114	51	15	-
2001	143	147	184	191	178	173	123	52	19	2
2000	115	132	188	169	167	159	112	48	21	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)														
Non-Hispanic Black														
Female														
1999	3,904	111	168	138	96	196	217	194	231	309	308	274	188	144
Origin not stated[2]														
Both sexes														
2002	461	3	4	5	6	17	26	21	32	39	38	47	38	32
2001	492	5	5	6	6	19	23	16	28	45	51	49	33	30
2000	512	4	8	6	5	23	27	27	31	35	54	54	52	26
1999	541	1	7	3	5	23	29	32	34	41	54	51	40	42
Male														
2002	358	2	3	4	4	10	22	16	28	33	29	38	29	32
2001	375	4	5	4	5	15	20	13	22	38	39	39	30	23
2000	385	3	3	2	1	15	21	21	27	29	44	42	40	21
1999	406	-	4	3	5	13	24	23	30	32	38	43	34	33
Female														
2002	103	1	1	1	2	7	4	5	4	6	9	9	9	-
2001	117	1	-	2	1	4	3	3	6	7	12	10	3	7
2000	127	1	5	4	4	8	6	6	4	6	10	12	12	5
1999	135	1	3	-	-	10	5	9	4	9	16	8	6	9
Transport accidents (V01-V99,Y85)														
All origins														
Both sexes														
2002	48,366	127	640	705	1,033	5,782	6,034	3,891	3,593	3,750	3,938	3,593	2,970	2,229
2001	47,288	151	678	769	1,054	5,393	5,746	3,953	3,501	3,881	3,946	3,379	2,872	2,152
2000	46,749	174	691	842	1,080	5,475	5,553	3,957	3,443	3,889	3,772	3,248	2,717	2,115
1999	46,423	189	712	897	1,078	5,452	5,237	3,922	3,512	3,869	3,684	3,121	2,524	2,131
Male														
2002	33,533	72	381	405	634	3,875	4,596	2,990	2,686	2,703	2,796	2,638	2,089	1,615
2001	32,794	68	392	447	668	3,694	4,417	3,015	2,646	2,744	2,869	2,452	2,032	1,467
2000	32,166	94	391	464	674	3,693	4,197	3,016	2,554	2,802	2,758	2,338	1,922	1,456
1999	31,642	101	400	532	674	3,593	3,965	2,952	2,559	2,741	2,635	2,234	1,782	1,454
Female														
2002	14,833	55	259	300	399	1,907	1,438	901	907	1,047	1,142	955	881	614
2001	14,494	83	286	322	386	1,699	1,329	938	855	1,137	1,077	927	840	685
2000	14,583	80	300	378	406	1,782	1,356	941	889	1,087	1,014	910	795	659
1999	14,781	88	312	365	404	1,859	1,272	970	953	1,128	1,049	887	742	677
Hispanic														
Both sexes														
2002	5,921	26	182	138	139	785	1,090	760	580	480	450	317	237	155

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Non-Hispanic Black										
Female										
1999	161	157	190	217	225	187	118	59	14	2
Origin not stated[2]										
Both sexes										
2002	19	16	20	21	25	15	9	4	-	24
2001	22	15	19	12	22	12	10	8	1	55
2000	21	13	20	17	15	16	9	3	1	45
1999	24	17	13	22	30	21	12	4	-	36
Male										
2002	17	14	17	11	20	9	1	2	-	17
2001	19	8	13	8	12	4	4	1	-	49
2000	18	12	12	9	8	7	5	1	-	44
1999	21	13	10	12	20	9	6	2	-	31
Female										
2002	2	2	3	10	5	6	8	2	-	7
2001	3	7	6	4	10	8	6	7	1	6
2000	3	1	8	8	7	9	4	2	1	1
1999	3	4	3	10	10	12	6	2	-	5
Transport accidents (V01-V99,Y85)										
All origins										
Both sexes										
2002	1,936	1,597	1,754	1,855	1,578	936	330	47	9	39
2001	1,709	1,567	1,715	1,943	1,564	903	304	63	2	43
2000	1,768	1,599	1,716	1,860	1,482	985	298	49	3	33
1999	1,681	1,727	1,852	1,902	1,566	964	311	50	5	37
Male										
2002	1,262	1,039	1,062	1,060	857	527	180	26	4	36
2001	1,164	974	1,041	1,106	877	492	163	28	1	37
2000	1,161	996	1,025	1,050	808	539	167	27	1	33
1999	1,107	1,091	1,071	1,089	870	551	179	25	3	34
Female										
2002	674	558	692	795	721	409	150	21	5	3
2001	545	593	674	837	687	411	141	35	1	6
2000	607	603	691	810	674	446	131	22	2	-
1999	574	636	781	813	696	413	132	25	2	3
Hispanic										
Both sexes										
2002	145	109	108	103	54	28	12	5	1	17

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Transport accidents (V01-V99,Y85)														
Hispanic														
Both sexes														
2001	5,774	43	159	120	145	718	1,022	768	525	514	428	313	244	180
2000	5,217	42	154	144	135	695	921	658	468	442	389	281	227	153
1999	4,880	40	157	108	142	615	842	640	485	392	366	246	199	148
Male														
2002	4,481	16	113	80	93	589	890	624	470	372	350	231	163	114
2001	4,334	16	92	71	84	527	856	641	426	402	329	216	174	107
2000	3,921	24	85	71	84	528	757	552	369	337	302	204	169	107
1999	3,630	22	85	65	86	452	684	522	382	307	283	184	146	94
Female														
2002	1,440	10	69	58	46	196	200	136	110	108	100	86	74	41
2001	1,440	27	67	49	61	191	166	127	99	112	99	97	70	73
2000	1,296	18	69	73	51	167	164	106	99	105	87	77	58	46
1999	1,250	18	72	43	56	163	158	118	103	85	83	62	53	54
Non-Hispanic[1]														
Both sexes														
2002	42,268	101	457	565	891	4,986	4,926	3,121	2,999	3,254	3,476	3,254	2,721	2,063
2001	41,339	108	517	648	906	4,659	4,712	3,177	2,964	3,351	3,497	3,054	2,614	1,964
2000	41,341	132	533	695	942	4,760	4,615	3,283	2,961	3,436	3,365	2,957	2,473	1,952
1999	41,330	149	553	786	933	4,820	4,372	3,260	3,012	3,461	3,305	2,859	2,316	1,967
Male														
2002	28,906	56	267	323	539	3,280	3,690	2,359	2,203	2,318	2,437	2,388	1,917	1,490
2001	28,327	52	298	375	582	3,155	3,551	2,367	2,211	2,330	2,525	2,226	1,845	1,356
2000	28,096	70	304	392	590	3,151	3,429	2,452	2,172	2,456	2,440	2,126	1,739	1,341
1999	27,850	79	313	464	585	3,133	3,263	2,415	2,163	2,422	2,342	2,037	1,629	1,346
Female														
2002	13,362	45	190	242	352	1,706	1,236	762	796	936	1,039	866	804	573
2001	13,012	56	219	273	324	1,504	1,161	810	753	1,021	972	828	769	608
2000	13,245	62	229	303	352	1,609	1,186	831	789	980	925	831	734	611
1999	13,480	70	240	322	348	1,687	1,109	845	849	1,039	963	822	687	621
Non-Hispanic White														
Both sexes														
2002	34,689	59	300	392	684	4,195	3,951	2,345	2,358	2,547	2,842	2,659	2,250	1,738
2001	33,753	74	353	440	682	3,851	3,712	2,472	2,299	2,664	2,830	2,431	2,180	1,650
2000	33,808	79	357	507	739	4,020	3,649	2,461	2,298	2,760	2,711	2,394	2,008	1,636
1999	33,803	98	349	557	734	4,044	3,449	2,528	2,361	2,737	2,688	2,307	1,904	1,643
Male														
2002	23,628	35	168	227	416	2,741	2,966	1,769	1,732	1,814	2,002	1,984	1,586	1,250
2001	23,047	36	208	250	436	2,584	2,817	1,842	1,719	1,844	2,043	1,772	1,554	1,136
2000	22,855	44	205	291	457	2,637	2,724	1,826	1,692	1,970	1,965	1,738	1,422	1,113
1999	22,682	58	192	325	457	2,595	2,586	1,876	1,685	1,917	1,910	1,644	1,348	1,127

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Transport accidents (V01-V99,Y85)										
Hispanic										
Both sexes										
2001	151	114	115	99	68	27	9	2	-	10
2000	115	98	122	76	45	26	10	2	-	14
1999	105	132	96	76	44	26	9	1	-	11
Male										
2002	93	82	67	63	28	15	8	3	1	16
2001	107	73	68	67	47	16	6	1	-	8
2000	82	63	77	53	21	13	7	2	-	14
1999	67	85	59	46	31	13	5	1	-	11
Female										
2002	52	27	41	40	26	13	4	2	-	1
2001	44	41	47	32	21	11	3	1	-	2
2000	33	35	45	23	24	13	3	-	-	-
1999	38	47	37	30	13	13	4	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,781	1,482	1,638	1,750	1,514	906	318	42	8	15
2001	1,549	1,452	1,592	1,841	1,489	874	294	60	2	15
2000	1,643	1,495	1,587	1,782	1,432	957	288	47	3	3
1999	1,568	1,589	1,749	1,820	1,509	936	301	49	5	11
Male										
2002	1,161	952	988	995	821	510	172	23	3	14
2001	1,049	901	968	1,038	827	475	157	26	1	12
2000	1,070	928	943	997	782	525	160	25	1	3
1999	1,033	1,002	1,007	1,040	829	537	174	24	3	10
Female										
2002	620	530	650	755	693	396	146	19	5	1
2001	500	551	624	803	662	399	137	34	1	3
2000	573	567	644	785	650	432	128	22	2	-
1999	535	587	742	780	680	399	127	25	2	1
Non-Hispanic White										
Both sexes										
2002	1,508	1,246	1,449	1,580	1,394	840	298	39	5	10
2001	1,287	1,235	1,393	1,668	1,376	814	277	55	2	8
2000	1,403	1,272	1,377	1,607	1,330	887	265	42	3	3
1999	1,303	1,361	1,536	1,636	1,384	860	272	44	4	4
Male										
2002	990	792	869	884	746	467	156	22	3	9
2001	873	765	852	931	767	436	149	25	1	7
2000	901	778	819	890	719	490	147	23	1	3
1999	859	863	882	929	757	489	158	19	2	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Transport accidents (V01-V99,Y85)														
Non-Hispanic White														
Female														
2002	11,061	24	132	165	268	1,454	985	576	626	733	840	675	664	488
2001	10,706	38	145	190	246	1,267	895	630	580	820	787	659	626	514
2000	10,953	35	152	216	282	1,383	925	635	606	790	746	656	586	523
1999	11,121	40	157	232	277	1,449	863	652	676	820	778	663	556	516
Non-Hispanic Black														
Both sexes														
2002	5,665	36	121	145	157	554	721	598	491	547	486	469	341	249
2001	5,803	18	127	177	177	578	744	541	517	521	550	494	335	249
2000	5,757	40	139	151	158	528	734	625	527	523	536	453	356	242
1999	5,810	44	168	199	161	571	691	569	506	583	491	443	315	244
Male														
2002	4,051	18	79	79	98	381	536	465	372	397	336	334	248	193
2001	4,143	7	69	112	122	414	552	408	389	369	410	381	234	182
2000	4,127	21	77	83	108	369	538	481	394	393	403	318	253	186
1999	4,097	18	96	124	105	400	509	436	381	409	360	327	230	171
Female														
2002	1,614	18	42	66	59	173	185	133	119	150	150	135	93	56
2001	1,660	11	58	65	55	164	192	133	128	152	140	113	101	67
2000	1,630	19	62	68	50	159	196	144	133	130	133	135	103	56
1999	1,713	26	72	75	56	171	182	133	125	174	131	116	85	73
Origin not stated[2]														
Both sexes														
2002	177	-	1	2	3	11	18	10	14	16	12	22	12	11
2001	175	-	2	1	3	16	12	8	12	16	21	12	14	8
2000	191	-	4	3	3	20	17	16	14	11	18	10	17	10
1999	213	-	2	3	3	17	23	22	15	16	13	16	9	16
Male														
2002	146	-	1	2	2	6	16	7	13	13	9	19	9	11
2001	133	-	2	1	2	12	10	7	9	12	15	10	13	4
2000	149	-	2	1	-	14	11	12	13	9	16	8	14	8
1999	162	-	2	3	3	8	18	15	14	12	10	13	7	14
Female														
2002	31	-	-	-	1	5	2	3	1	3	3	3	3	-
2001	42	-	-	-	1	4	2	1	3	4	6	2	1	4
2000	42	-	2	2	3	6	6	4	1	2	2	2	3	2
1999	51	-	-	-	-	9	5	7	1	4	3	3	2	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Transport accidents (V01-V99,Y85)										
Non-Hispanic White										
Female										
2002	518	454	580	696	648	373	142	17	2	1
2001	414	470	541	737	609	378	128	30	1	1
2000	502	494	558	717	611	397	118	19	2	-
1999	444	498	654	707	627	371	114	25	2	-
Non-Hispanic Black										
Both sexes										
2002	195	158	126	117	89	44	13	2	3	3
2001	206	171	136	126	72	38	14	5	-	7
2000	175	164	158	107	72	50	16	3	-	-
1999	186	161	153	136	96	59	21	5	1	7
Male										
2002	134	111	85	83	56	32	10	1	-	3
2001	150	111	74	84	39	23	7	1	-	5
2000	128	119	99	72	46	27	10	2	-	-
1999	132	103	93	84	55	41	11	5	1	6
Female										
2002	61	47	41	34	33	12	3	1	3	-
2001	56	60	62	42	33	15	7	4	-	2
2000	47	45	59	35	26	23	6	1	-	-
1999	54	58	60	52	41	18	10	-	-	1
Origin not stated[2]										
Both sexes										
2002	10	6	8	2	10	2	-	-	-	7
2001	9	1	8	3	7	2	1	1	-	18
2000	10	6	7	2	5	2	-	-	-	16
1999	8	6	7	6	13	2	1	-	-	15
Male										
2002	8	5	7	2	8	2	-	-	-	6
2001	8	-	5	1	3	1	-	1	-	17
2000	9	5	5	-	5	1	-	-	-	16
1999	7	4	5	3	10	1	-	-	-	13
Female										
2002	2	1	1	-	2	-	-	-	-	1
2001	1	1	3	2	4	1	1	-	-	1
2000	1	1	2	2	-	1	-	-	-	-
1999	1	2	2	3	3	1	1	-	-	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)														
All origins														
Both sexes														
2002	45,380	123	610	664	950	5,634	5,825	3,731	3,361	3,462	3,615	3,240	2,697	2,004
2001	43,788	144	626	718	968	5,208	5,517	3,710	3,227	3,545	3,538	3,011	2,553	1,939
2000	43,354	168	651	780	992	5,251	5,309	3,715	3,169	3,545	3,382	2,936	2,425	1,917
1999	42,401	184	650	802	969	5,198	4,930	3,604	3,174	3,460	3,278	2,747	2,225	1,885
Male														
2002	31,064	68	361	385	578	3,765	4,413	2,855	2,494	2,459	2,525	2,328	1,861	1,423
2001	30,064	66	356	418	611	3,546	4,232	2,826	2,410	2,473	2,543	2,164	1,774	1,296
2000	29,451	90	368	429	614	3,527	3,989	2,804	2,331	2,507	2,433	2,091	1,689	1,293
1999	28,552	97	366	469	593	3,412	3,715	2,690	2,303	2,424	2,311	1,936	1,537	1,250
Female														
2002	14,316	55	249	279	372	1,869	1,412	876	867	1,003	1,090	912	836	581
2001	13,724	78	270	300	357	1,662	1,285	884	817	1,072	995	847	779	643
2000	13,903	78	283	351	378	1,724	1,320	911	838	1,038	949	845	736	624
1999	13,849	87	284	333	376	1,786	1,215	914	871	1,036	967	811	688	635
Hispanic														
Both sexes														
2002	5,659	26	176	129	130	763	1,047	737	557	458	427	294	215	144
2001	5,272	40	142	112	134	701	976	716	481	458	376	266	204	152
2000	4,900	40	144	135	125	671	886	627	439	406	350	262	210	141
1999	4,525	40	148	100	131	587	797	593	442	354	330	224	181	141
Male														
2002	4,262	16	108	74	89	571	850	604	448	353	330	213	145	105
2001	3,987	15	82	67	77	513	817	600	392	357	296	192	148	93
2000	3,658	24	78	65	77	510	726	523	346	304	270	186	155	98
1999	3,337	22	82	59	77	431	646	480	348	275	255	165	129	87
Female														
2002	1,397	10	68	55	41	192	197	133	109	105	97	81	70	39
2001	1,285	25	60	45	57	188	159	116	89	101	80	74	56	59
2000	1,242	16	66	70	48	161	160	104	93	102	80	76	55	43
1999	1,188	18	66	41	54	156	151	113	94	79	75	59	52	54
Non-Hispanic[1]														
Both sexes														
2002	39,567	97	433	533	817	4,861	4,760	2,985	2,792	2,990	3,179	2,927	2,473	1,851
2001	38,362	104	483	605	831	4,491	4,530	2,987	2,735	3,073	3,144	2,735	2,337	1,781
2000	38,295	128	505	643	864	4,561	4,408	3,075	2,717	3,130	3,017	2,665	2,201	1,767
1999	37,684	144	500	699	835	4,594	4,113	2,992	2,720	3,090	2,936	2,509	2,037	1,728

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)										
All origins										
Both sexes										
2002	1,760	1,469	1,644	1,763	1,508	909	321	47	8	35
2001	1,526	1,409	1,581	1,823	1,489	868	294	59	2	33
2000	1,589	1,436	1,602	1,764	1,409	953	285	47	3	26
1999	1,485	1,565	1,711	1,772	1,471	920	291	46	4	30
Male										
2002	1,117	923	965	992	801	512	176	26	4	33
2001	1,013	846	940	1,027	832	478	157	28	1	27
2000	1,013	870	932	980	759	517	161	27	1	26
1999	950	969	962	1,012	815	524	164	24	2	27
Female										
2002	643	546	679	771	707	397	145	21	4	2
2001	513	563	641	796	657	390	137	31	1	6
2000	576	566	670	784	650	436	124	20	2	-
1999	535	596	749	760	656	396	127	22	2	3
Hispanic										
Both sexes										
2002	139	106	105	96	50	28	11	5	1	15
2001	127	96	105	83	63	22	8	2	-	8
2000	105	85	113	71	42	22	10	2	-	14
1999	97	119	85	72	41	23	9	1	-	10
Male										
2002	87	79	65	59	25	15	7	3	1	15
2001	91	61	62	56	43	13	5	1	-	6
2000	72	54	69	50	18	10	7	2	-	14
1999	59	75	49	42	29	11	5	1	-	10
Female										
2002	52	27	40	37	25	13	4	2	-	-
2001	36	35	43	27	20	9	3	1	-	2
2000	33	31	44	21	24	12	3	-	-	-
1999	38	44	36	30	12	12	4	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,613	1,358	1,532	1,665	1,449	879	310	42	7	14
2001	1,392	1,312	1,468	1,737	1,419	844	285	56	2	11
2000	1,476	1,346	1,483	1,691	1,363	929	275	45	3	3
1999	1,382	1,440	1,619	1,694	1,418	895	281	45	4	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)														
Non-Hispanic[1]														
Male														
2002	26,677	52	252	309	487	3,188	3,547	2,245	2,035	2,095	2,189	2,098	1,710	1,309
2001	25,962	51	273	350	532	3,021	3,405	2,220	2,010	2,106	2,234	1,964	1,615	1,200
2000	25,670	66	288	363	537	3,004	3,253	2,270	1,973	2,196	2,150	1,898	1,523	1,188
1999	25,072	75	282	407	513	2,973	3,054	2,198	1,943	2,137	2,047	1,760	1,403	1,149
Female														
2002	12,890	45	181	224	330	1,673	1,213	740	757	895	990	829	763	542
2001	12,400	53	210	255	299	1,470	1,125	767	725	967	910	771	722	581
2000	12,625	62	217	280	327	1,557	1,155	805	744	934	867	767	678	579
1999	12,612	69	218	292	322	1,621	1,059	794	777	953	889	749	634	579
Non-Hispanic White														
Both sexes														
2002	32,405	56	280	370	630	4,090	3,807	2,231	2,181	2,326	2,597	2,386	2,035	1,559
2001	31,283	71	323	412	624	3,711	3,555	2,310	2,114	2,444	2,536	2,179	1,942	1,500
2000	31,219	75	340	464	672	3,861	3,469	2,281	2,098	2,508	2,415	2,155	1,770	1,475
1999	30,743	94	315	491	653	3,860	3,239	2,301	2,124	2,426	2,371	2,011	1,667	1,434
Male														
2002	21,743	32	155	216	378	2,664	2,841	1,675	1,587	1,625	1,798	1,739	1,409	1,100
2001	21,073	35	185	232	399	2,470	2,694	1,716	1,554	1,666	1,802	1,560	1,355	1,011
2000	20,789	40	197	269	411	2,523	2,571	1,669	1,527	1,754	1,719	1,550	1,231	982
1999	20,362	55	175	284	399	2,467	2,417	1,693	1,508	1,678	1,660	1,410	1,155	954
Female														
2002	10,662	24	125	154	252	1,426	966	556	594	701	799	647	626	459
2001	10,210	36	138	180	225	1,241	861	594	560	778	734	619	587	489
2000	10,430	35	143	195	261	1,338	898	612	571	754	696	605	539	493
1999	10,381	39	140	207	254	1,393	822	608	616	748	711	601	512	480
Non-Hispanic Black														
Both sexes														
2002	5,338	35	118	135	143	538	705	581	468	509	450	419	315	222
2001	5,423	18	123	163	161	557	728	519	488	484	501	441	306	223
2000	5,424	40	132	143	151	503	713	606	489	481	496	412	338	223
1999	5,388	43	152	182	149	544	665	540	466	534	451	403	282	222
Male														
2002	3,774	17	77	76	87	368	523	449	353	367	306	292	224	168
2001	3,842	7	67	105	109	398	537	391	366	339	367	338	212	157
2000	3,857	21	72	76	103	348	520	463	365	358	369	286	238	169

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)										
Non-Hispanic[1]										
Male										
2002	1,024	840	894	931	769	495	169	23	3	13
2001	916	785	873	970	786	464	152	26	1	8
2000	934	812	859	930	737	506	154	25	1	3
1999	885	890	908	967	777	512	159	23	2	8
Female										
2002	589	518	638	734	680	384	141	19	4	1
2001	476	527	595	767	633	380	133	30	1	3
2000	542	534	624	761	626	423	121	20	2	-
1999	497	550	711	727	641	383	122	22	2	1
Non-Hispanic White										
Both sexes										
2002	1,359	1,142	1,354	1,508	1,336	816	290	39	4	9
2001	1,152	1,120	1,278	1,578	1,313	789	270	52	2	8
2000	1,253	1,145	1,286	1,527	1,264	862	252	41	3	3
1999	1,145	1,233	1,422	1,527	1,305	824	254	40	4	3
Male										
2002	868	698	786	833	699	454	153	22	3	8
2001	753	671	765	870	730	427	145	25	1	7
2000	777	679	745	830	675	472	141	23	1	3
1999	735	764	797	863	712	468	145	18	2	3
Female										
2002	491	444	568	675	637	362	137	17	1	1
2001	399	449	513	708	583	362	125	27	1	1
2000	476	466	541	697	589	390	111	18	2	-
1999	410	469	625	664	593	356	109	22	2	-
Non-Hispanic Black										
Both sexes										
2002	182	145	118	107	84	43	13	2	3	3
2001	187	152	131	115	69	37	13	4	-	3
2000	165	149	149	100	69	47	16	2	-	-
1999	165	148	143	126	88	54	20	5	-	6
Male										
2002	122	98	77	73	52	31	10	1	-	3
2001	138	94	69	78	38	23	7	1	-	1
2000	120	106	93	67	45	26	10	2	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)														
Non-Hispanic Black														
Male														
1999	3,763	17	84	109	94	379	490	412	349	371	325	296	203	153
Female														
2002	1,564	18	41	59	56	170	182	132	115	142	144	127	91	54
2001	1,581	11	56	58	52	159	191	128	122	145	134	103	94	66
2000	1,567	19	60	67	48	155	193	143	124	123	127	126	100	54
1999	1,625	26	68	73	55	165	175	128	117	163	126	107	79	69
Origin not stated[2]														
Both sexes														
2002	154	-	1	2	3	10	18	9	12	14	9	19	9	9
2001	154	-	1	1	3	16	11	7	11	14	18	10	12	6
2000	159	-	2	2	3	19	15	13	13	9	15	9	14	9
1999	192	-	2	3	3	17	20	19	12	16	12	14	7	16
Male														
2002	125	-	1	2	2	6	16	6	11	11	6	17	6	9
2001	115	-	1	1	2	12	10	6	8	10	13	8	11	3
2000	123	-	2	1	-	13	10	11	12	7	13	7	11	7
1999	143	-	2	3	3	8	15	12	12	12	9	11	5	14
Female														
2002	29	-	-	-	1	4	2	3	1	3	3	2	3	-
2001	39	-	-	-	1	4	1	1	3	4	5	2	1	3
2000	36	-	-	1	3	6	5	2	1	2	2	2	3	2
1999	49	-	-	-	-	9	5	7	-	4	3	3	2	2
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
All origins														
Both sexes														
2002	1,086	2	17	21	49	63	96	59	79	101	122	121	90	61
2001	1,405	3	28	25	55	95	103	85	92	141	144	142	105	67
2000	1,492	2	31	39	59	128	115	94	118	149	168	125	109	64
1999	2,118	4	48	62	66	172	190	152	165	205	203	179	125	101

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04, V09.0,V09.2,V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79,V80.3-V80.5, V81.0-V81.1,V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8,V89.0, V89.2)										
Non-Hispanic Black										
Male										
1999	114	94	84	81	51	37	10	5	-	5
Female										
2002	60	47	41	34	32	12	3	1	3	-
2001	49	58	62	37	31	14	6	3	-	2
2000	45	43	56	33	24	21	6	-	-	-
1999	51	54	59	45	37	17	10	-	-	1
Origin not stated[2]										
Both sexes										
2002	8	5	7	2	9	2	-	-	-	6
2001	7	1	8	3	7	2	1	1	-	14
2000	8	5	6	2	4	2	-	-	-	9
1999	6	6	7	6	12	2	1	-	-	11
Male										
2002	6	4	6	2	7	2	-	-	-	5
2001	6	-	5	1	3	1	-	1	-	13
2000	7	4	4	-	4	1	-	-	-	9
1999	6	4	5	3	9	1	-	-	-	9
Female										
2002	2	1	1	-	2	-	-	-	-	1
2001	1	1	3	2	4	1	1	-	-	1
2000	1	1	2	2	-	1	-	-	-	-
1999	-	2	2	3	3	1	1	-	-	2
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)										
All origins										
Both sexes										
2002	48	33	37	35	31	11	5	-	1	4
2001	66	64	53	59	39	20	7	2	-	10
2000	58	52	48	51	40	22	12	1	-	7
1999	81	79	66	92	67	34	16	3	1	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
All origins														
Male														
2002	882	2	10	11	36	49	85	47	61	85	102	107	78	53
2001	1,067	-	20	13	36	72	86	71	81	117	118	105	84	56
2000	1,125	2	17	21	40	99	95	81	90	121	137	92	74	48
1999	1,559	4	26	43	48	119	160	126	118	157	162	135	99	77
Female														
2002	204	-	7	10	13	14	11	12	18	16	20	14	12	8
2001	338	3	8	12	19	23	17	14	11	24	26	37	21	11
2000	367	-	14	18	19	29	20	13	28	28	31	33	35	16
1999	559	-	22	19	18	53	30	26	47	48	41	44	26	24
Hispanic														
Both sexes														
2002	144	-	2	5	5	14	28	11	10	13	15	12	10	4
2001	200	-	8	2	5	13	26	24	14	20	22	19	10	7
2000	201	-	10	7	8	15	19	15	21	19	25	15	10	10
1999	264	-	8	6	8	20	35	29	31	28	29	16	15	2
Male														
2002	119	-	1	4	2	12	26	9	9	12	13	10	7	3
2001	173	-	6	-	4	12	25	23	12	19	20	13	9	6
2000	166	-	7	5	6	13	18	15	17	17	19	14	8	7
1999	220	-	3	5	7	16	29	28	25	24	23	13	14	2
Female														
2002	25	-	1	1	3	2	2	2	1	1	2	2	3	1
2001	27	-	2	2	1	1	1	1	2	1	2	6	1	1
2000	35	-	3	2	2	2	1	-	4	2	6	1	2	3
1999	44	-	5	1	1	4	6	1	6	4	6	3	1	-
Non-Hispanic[1]														
Both sexes														
2002	929	2	15	16	44	48	68	48	69	88	104	107	77	55
2001	1,187	3	19	23	50	82	76	60	77	120	121	121	93	58
2000	1,262	2	19	31	51	112	94	76	96	128	140	110	97	53
1999	1,836	4	40	56	58	152	152	121	131	177	173	162	109	99
Male														
2002	752	2	9	7	34	37	59	38	52	73	86	96	68	48
2001	878	-	13	13	32	60	61	47	68	97	97	90	73	49

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)										
All origins										
Male										
2002	39	26	30	25	25	6	2	-	-	3
2001	48	50	34	38	18	6	4	-	-	10
2000	50	38	37	33	24	14	5	-	-	7
1999	58	52	45	51	37	20	13	1	1	7
Female										
2002	9	7	7	10	6	5	3	-	1	1
2001	18	14	19	21	21	14	3	2	-	-
2000	8	14	11	18	16	8	7	1	-	-
1999	23	27	21	41	30	14	3	2	-	-
Hispanic										
Both sexes										
2002	3	1	1	5	3	-	-	-	-	2
2001	7	3	2	11	2	2	1	-	-	2
2000	4	9	5	4	1	4	-	-	-	-
1999	6	13	8	4	3	2	-	-	-	1
Male										
2002	3	1	1	2	3	-	-	-	-	1
2001	6	2	2	9	1	1	1	-	-	2
2000	4	6	4	2	1	3	-	-	-	-
1999	6	10	7	4	2	1	-	-	-	1
Female										
2002	-	-	-	3	-	-	-	-	-	1
2001	1	1	-	2	1	1	-	-	-	-
2000	-	3	1	2	-	1	-	-	-	-
1999	-	3	1	-	1	1	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	45	32	36	30	27	11	5	-	1	1
2001	57	61	51	48	37	18	6	2	-	4
2000	52	42	43	47	38	18	12	1	-	-
1999	73	66	58	88	63	32	16	3	1	2
Male										
2002	36	25	29	23	21	6	2	-	-	1
2001	40	48	32	29	17	5	3	-	-	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
Non-Hispanic[1]														
Male														
2000	936	2	10	16	34	85	76	65	72	102	115	78	64	40
1999	1,323	4	23	38	41	103	128	96	91	133	138	121	84	75
Female														
2002	177	-	6	9	10	11	9	10	17	15	18	11	9	7
2001	309	3	6	10	18	22	15	13	9	23	24	31	20	9
2000	326	-	9	15	17	27	18	11	24	26	25	32	33	13
1999	513	-	17	18	17	49	24	25	40	44	35	41	25	24
Non-Hispanic White														
Both sexes														
2002	719	1	11	12	32	39	56	38	56	67	73	77	64	39
2001	901	2	16	14	35	62	64	50	53	79	86	88	77	47
2000	997	2	9	26	41	94	74	59	73	97	104	80	81	39
1999	1,480	3	25	39	47	122	117	96	99	141	141	130	89	88
Male														
2002	582	1	7	7	25	31	50	29	42	57	59	68	56	34
2001	673	-	12	9	21	48	51	40	49	65	68	67	62	39
2000	741	2	3	13	28	70	60	51	55	79	86	57	53	28
1999	1,051	3	12	25	32	81	96	75	68	106	112	95	68	67
Female														
2002	137	-	4	5	7	8	6	9	14	10	14	9	8	5
2001	228	2	4	5	14	14	13	10	4	14	18	21	15	8
2000	256	-	6	13	13	24	14	8	18	18	18	23	28	11
1999	429	-	13	14	15	41	21	21	31	35	29	35	21	21
Non-Hispanic Black														
Both sexes														
2002	168	1	3	4	9	9	8	10	9	18	22	29	12	12
2001	216	-	3	9	14	15	6	8	16	27	32	26	12	9
2000	203	-	7	4	7	13	17	11	21	26	28	24	10	11
1999	248	1	12	14	6	18	15	18	24	29	23	23	17	11
Male														
2002	143	1	2	-	7	6	6	9	8	14	20	28	11	10
2001	159	-	1	4	11	10	5	6	12	21	27	20	8	8
2000	155	-	5	3	5	10	14	10	15	21	22	19	7	9
1999	192	1	9	13	6	14	13	16	18	22	19	18	14	8

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)										
Non-Hispanic[1]										
Male										
2000	44	31	33	31	22	11	5	-	-	-
1999	51	42	38	47	34	19	13	1	1	2
Female										
2002	9	7	7	7	6	5	3	-	1	-
2001	17	13	19	19	20	13	3	2	-	-
2000	8	11	10	16	16	7	7	1	-	-
1999	22	24	20	41	29	13	3	2	-	-
Non-Hispanic White										
Both sexes										
2002	35	24	31	23	25	9	5	-	1	1
2001	42	45	47	41	32	15	4	2	-	-
2000	45	33	37	40	35	16	12	-	-	-
1999	63	53	49	75	56	29	14	3	-	1
Male										
2002	29	19	24	16	20	5	2	-	-	1
2001	34	35	28	26	14	3	2	-	-	-
2000	40	23	28	28	21	11	5	-	-	-
1999	43	36	31	41	30	17	11	1	-	1
Female										
2002	6	5	7	7	5	4	3	-	1	-
2001	8	10	19	15	18	12	2	2	-	-
2000	5	10	9	12	14	5	7	-	-	-
1999	20	17	18	34	26	12	3	2	-	-
Non-Hispanic Black										
Both sexes										
2002	6	5	4	5	2	-	-	-	-	-
2001	13	12	3	5	1	-	1	-	-	4
2000	4	7	3	4	3	2	-	1	-	-
1999	6	8	6	8	3	3	1	-	1	1
Male										
2002	6	5	4	5	1	-	-	-	-	-
2001	6	10	3	3	-	-	-	-	-	4
2000	3	7	2	2	1	-	-	-	-	-
1999	4	4	5	2	1	2	1	-	1	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)														
Non-Hispanic Black														
Female														
2002	25	-	1	4	2	3	2	1	1	4	2	1	1	2
2001	57	-	2	5	3	5	1	2	4	6	5	6	4	1
2000	48	-	2	1	2	3	3	1	6	5	6	5	3	2
1999	56	-	3	1	-	4	2	2	6	7	4	5	3	3
Origin not stated[2]														
Both sexes														
2002	13	-	-	-	-	1	-	-	-	-	3	2	3	2
2001	18	-	1	-	-	-	1	1	1	1	1	2	2	2
2000	29	-	2	1	-	1	2	3	1	2	3	-	2	1
1999	18	-	-	-	-	-	3	2	3	-	1	1	1	-
Male														
2002	11	-	-	-	-	-	-	-	-	-	3	1	3	2
2001	16	-	1	-	-	-	-	1	1	1	1	2	2	1
2000	23	-	-	-	-	1	1	1	1	2	3	-	2	1
1999	16	-	-	-	-	-	3	2	2	-	1	1	1	-
Female														
2002	2	-	-	-	-	1	-	-	-	-	-	1	-	-
2001	2	-	-	-	-	-	1	-	-	-	-	-	-	1
2000	6	-	2	1	-	-	1	2	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	1	-	-	-	-	-
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
All origins														
Both sexes														
2002	1,900	2	13	20	34	85	113	101	153	187	201	232	183	164
2001	2,095	4	24	26	31	90	126	158	182	195	264	226	214	146
2000	1,903	4	9	23	29	96	129	148	156	195	222	187	183	134
1999	1,904	1	14	33	43	82	117	166	173	204	203	195	174	145
Male														
2002	1,587	2	10	9	20	61	98	88	131	159	169	203	150	139
2001	1,663	2	16	16	21	76	99	118	155	154	208	183	174	115
2000	1,590	2	6	14	20	67	113	131	133	174	188	155	159	115
1999	1,531	-	8	20	33	62	90	136	138	160	162	163	146	127

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3, V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9,V88.9,V89.1, V89.3,V89.9)										
Non-Hispanic Black										
Female										
2002	-	-	-	-	1	-	-	-	-	-
2001	7	2	-	2	1	-	1	-	-	-
2000	1	-	1	2	2	2	-	1	-	-
1999	2	4	1	6	2	1	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	1	-	-	-	-	1
2001	2	-	-	-	-	-	-	-	-	4
2000	2	1	-	-	1	-	-	-	-	7
1999	2	-	-	-	1	-	-	-	-	4
Male										
2002	-	-	-	-	1	-	-	-	-	1
2001	2	-	-	-	-	-	-	-	-	4
2000	2	1	-	-	1	-	-	-	-	7
1999	1	-	-	-	1	-	-	-	-	4
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
All origins										
Both sexes										
2002	128	95	73	57	39	16	4	-	-	-
2001	117	94	81	61	36	15	3	2	-	-
2000	121	111	66	45	33	10	1	1	-	-
1999	115	83	75	38	28	10	4	1	-	-
Male										
2002	106	90	67	43	31	9	2	-	-	-
2001	103	78	67	41	27	8	2	-	-	-
2000	98	88	56	37	25	8	1	-	-	-
1999	99	70	64	26	18	7	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
All origins														
Female														
2002	313	-	3	11	14	24	15	13	22	28	32	29	33	25
2001	432	2	8	10	10	14	27	40	27	41	56	43	40	31
2000	313	2	3	9	9	29	16	17	23	21	34	32	24	19
1999	373	1	6	13	10	20	27	30	35	44	41	32	28	18
Hispanic														
Both sexes														
2002	118	-	4	4	4	8	15	12	13	9	8	11	12	7
2001	302	3	9	6	6	4	20	28	30	36	30	28	30	21
2000	116	2	-	2	2	9	16	16	8	17	14	4	7	2
1999	91	-	1	2	3	8	10	18	12	10	7	6	3	5
Male														
2002	100	-	4	2	2	6	14	11	13	7	7	8	11	6
2001	174	1	4	4	3	2	14	18	22	26	13	11	17	8
2000	97	-	-	1	1	5	13	14	6	16	13	4	6	2
1999	73	-	-	1	2	5	9	14	9	8	5	6	3	5
Female														
2002	18	-	-	2	2	2	1	1	-	2	1	3	1	1
2001	128	2	5	2	3	2	6	10	8	10	17	17	13	13
2000	19	2	-	1	1	4	3	2	2	1	1	-	1	-
1999	18	-	1	1	1	3	1	4	3	2	2	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	1,772	2	9	16	30	77	98	88	138	176	193	220	171	157
2001	1,790	1	15	20	25	86	106	130	152	158	232	198	184	125
2000	1,784	2	9	21	27	87	113	132	148	178	208	182	175	132
1999	1,810	1	13	31	40	74	107	147	161	194	196	188	170	140
Male														
2002	1,477	2	6	7	18	55	84	76	116	150	162	194	139	133
2001	1,487	1	12	12	18	74	85	100	133	127	194	172	157	107
2000	1,490	2	6	13	19	62	100	117	127	158	175	150	152	113
1999	1,455	-	8	19	31	57	81	121	129	152	157	156	142	122
Female														
2002	295	-	3	9	12	22	14	12	22	26	31	26	32	24
2001	303	-	3	8	7	12	21	30	19	31	38	26	27	18
2000	294	-	3	8	8	25	13	15	21	20	33	32	23	19
1999	355	1	5	12	9	17	26	26	32	42	39	32	28	18

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
All origins										
Female										
2002	22	5	6	14	8	7	2	-	-	-
2001	14	16	14	20	9	7	1	2	-	-
2000	23	23	10	8	8	2	-	1	-	-
1999	16	13	11	12	10	3	2	1	-	-
Hispanic										
Both sexes										
2002	3	2	2	2	1	-	1	-	-	-
2001	17	15	8	5	3	3	-	-	-	-
2000	6	4	4	1	2	-	-	-	-	-
1999	2	-	3	-	-	1	-	-	-	-
Male										
2002	3	2	1	2	-	-	1	-	-	-
2001	10	10	4	2	3	2	-	-	-	-
2000	6	3	4	1	2	-	-	-	-	-
1999	2	-	3	-	-	1	-	-	-	-
Female										
2002	-	-	1	-	1	-	-	-	-	-
2001	7	5	4	3	-	1	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	123	92	70	55	38	16	3	-	-	-
2001	100	79	73	56	33	12	3	2	-	-
2000	115	107	61	44	31	10	1	1	-	-
1999	113	83	72	38	28	9	4	1	-	-
Male										
2002	101	87	65	41	31	9	1	-	-	-
2001	93	68	63	39	24	6	2	-	-	-
2000	92	85	51	36	23	8	1	-	-	-
1999	97	70	61	26	18	6	2	-	-	-
Female										
2002	22	5	5	14	7	7	2	-	-	-
2001	7	11	10	17	9	6	1	2	-	-
2000	23	22	10	8	8	2	-	1	-	-
1999	16	13	11	12	10	3	2	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
Non-Hispanic White														
Both sexes														
2002	1,565	2	9	10	22	66	88	76	121	154	172	196	151	140
2001	1,569	1	14	14	23	78	93	112	132	141	208	164	161	103
2000	1,592	2	8	17	26	65	106	121	127	155	192	159	157	122
1999	1,580	1	9	27	34	62	93	131	138	170	176	166	148	121
Male														
2002	1,303	2	6	4	13	46	75	65	103	132	145	177	121	116
2001	1,301	1	11	9	16	66	72	86	116	113	173	145	137	86
2000	1,325	2	5	9	18	44	93	106	110	137	160	131	138	103
1999	1,269	-	5	16	26	47	73	108	109	133	138	139	125	106
Female														
2002	262	-	3	6	9	20	13	11	18	22	27	19	30	24
2001	268	-	3	5	7	12	21	26	16	28	35	19	24	17
2000	267	-	3	8	8	21	13	15	17	18	32	28	19	19
1999	311	1	4	11	8	15	20	23	29	37	38	27	23	15
Non-Hispanic Black														
Both sexes														
2002	159	-	-	6	5	7	8	7	14	20	14	21	14	15
2001	164	-	1	5	2	6	10	14	13	10	17	27	17	17
2000	130	-	-	4	-	12	4	8	17	16	12	17	8	8
1999	174	-	4	3	6	9	11	11	16	20	17	17	16	11
Male														
2002	134	-	-	3	4	7	7	7	11	16	10	14	13	15
2001	142	-	1	3	2	6	10	11	11	9	16	23	14	17
2000	115	-	-	4	-	11	4	8	14	14	12	13	8	8
1999	142	-	3	2	5	7	6	8	14	16	16	13	13	10
Female														
2002	25	-	-	3	1	-	1	-	3	4	4	7	1	-
2001	22	-	-	2	-	-	-	3	2	1	1	4	3	-
2000	15	-	-	-	-	1	-	-	3	2	-	4	-	-
1999	32	-	1	1	1	2	5	3	2	4	1	4	3	1
Origin not stated[2]														
Both sexes														
2002	10	-	-	-	-	-	-	1	2	2	-	1	-	-
2001	3	-	-	-	-	-	-	-	-	1	2	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	1	1	-
1999	3	-	-	-	-	-	-	1	-	-	-	1	1	-
Male														
2002	10	-	-	-	-	-	-	1	2	2	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
Non-Hispanic White										
Both sexes										
2002	114	80	64	49	33	15	3	-	-	-
2001	93	70	68	49	31	10	3	1	-	-
2000	105	94	54	40	31	9	1	1	-	-
1999	95	75	65	34	23	7	4	1	-	-
Male										
2002	93	75	59	35	27	8	1	-	-	-
2001	86	59	59	35	23	6	2	-	-	-
2000	84	76	46	32	23	7	1	-	-	-
1999	81	63	54	25	15	4	2	-	-	-
Female										
2002	21	5	5	14	6	7	2	-	-	-
2001	7	11	9	14	8	4	1	1	-	-
2000	21	18	8	8	8	2	-	1	-	-
1999	14	12	11	9	8	3	2	1	-	-
Non-Hispanic Black										
Both sexes										
2002	7	8	4	5	3	1	-	-	-	-
2001	6	7	2	6	2	1	-	1	-	-
2000	6	8	6	3	-	1	-	-	-	-
1999	15	5	4	2	5	2	-	-	-	-
Male										
2002	6	8	4	5	3	1	-	-	-	-
2001	6	7	2	3	1	-	-	-	-	-
2000	5	6	4	3	-	1	-	-	-	-
1999	14	5	4	1	3	2	-	-	-	-
Female										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	3	1	1	-	1	-	-
2000	1	2	2	-	-	-	-	-	-	-
1999	1	-	-	1	2	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	2	1	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	1	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	2	1	1	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)														
Origin not stated[2]														
Male														
2001	2	-	-	-	-	-	-	-	-	1	1	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	1	1	-
1999	3	-	-	-	-	-	-	1	-	-	-	1	1	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	1	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nontransport accidents (W00-X59,Y86)														
All origins														
Both sexes														
2002	58,376	819	1,001	471	509	1,355	2,241	2,231	2,854	3,892	5,130	4,648	3,464	2,328
2001	54,249	825	1,036	514	499	1,253	2,019	1,982	2,403	3,708	4,410	4,090	3,003	2,007
2000	51,151	707	1,135	549	508	1,280	1,805	1,941	2,428	3,582	4,170	3,622	2,691	1,906
1999	51,437	656	1,186	562	554	1,236	1,731	1,959	2,497	3,497	4,181	3,575	2,419	1,813
Male														
2002	35,724	481	643	315	359	1,106	1,861	1,778	2,181	2,832	3,681	3,302	2,463	1,640
2001	33,266	473	653	339	349	1,028	1,693	1,590	1,841	2,758	3,241	3,067	2,192	1,405
2000	31,651	407	707	360	363	1,064	1,506	1,590	1,899	2,735	3,149	2,765	2,038	1,348
1999	31,893	393	727	365	403	1,031	1,451	1,603	1,983	2,670	3,202	2,762	1,824	1,299
Female														
2002	22,652	338	358	156	150	249	380	453	673	1,060	1,449	1,346	1,001	688
2001	20,983	352	383	175	150	225	326	392	562	950	1,169	1,023	811	602
2000	19,500	300	428	189	145	216	299	351	529	847	1,021	857	653	558
1999	19,544	263	459	197	151	205	280	356	514	827	979	813	595	514
Hispanic														
Both sexes														
2002	4,185	101	159	61	51	199	339	322	365	418	493	368	246	156
2001	3,749	111	151	66	48	161	281	305	340	428	390	314	207	133
2000	3,613	76	152	61	43	189	290	286	364	414	373	261	184	119
1999	3,770	94	184	62	50	185	304	296	369	452	397	300	181	102
Male														
2002	3,217	67	102	40	31	174	305	293	322	340	406	291	203	121
2001	2,823	59	103	43	26	142	248	267	285	361	326	265	167	106
2000	2,775	43	89	34	21	161	271	252	326	345	333	231	146	86
1999	2,948	58	112	42	34	156	283	265	325	385	332	259	145	89

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)										
Origin not stated[2]										
Male										
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	1	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Nontransport accidents (W00-X59,Y86)										
All origins										
Both sexes										
2002	1,852	2,075	2,660	4,083	5,388	5,571	3,925	1,548	285	46
2001	1,790	1,939	2,614	4,029	5,152	5,363	3,770	1,456	310	77
2000	1,716	1,776	2,607	3,736	4,680	4,948	3,641	1,370	301	52
1999	1,660	1,965	2,664	3,870	4,944	5,174	3,643	1,330	252	69
Male										
2002	1,264	1,330	1,600	2,208	2,562	2,268	1,308	454	50	38
2001	1,192	1,226	1,510	2,204	2,465	2,238	1,323	353	59	67
2000	1,168	1,121	1,497	1,989	2,216	2,024	1,224	370	61	50
1999	1,155	1,245	1,585	2,007	2,406	2,096	1,218	351	57	60
Female										
2002	588	745	1,060	1,875	2,826	3,303	2,617	1,094	235	8
2001	598	713	1,104	1,825	2,687	3,125	2,447	1,103	251	10
2000	548	655	1,110	1,747	2,464	2,924	2,417	1,000	240	2
1999	505	720	1,079	1,863	2,538	3,078	2,425	979	195	9
Hispanic										
Both sexes										
2002	110	119	130	159	148	102	80	42	6	11
2001	100	91	134	123	119	113	75	34	6	19
2000	105	89	109	117	127	120	84	28	8	14
1999	111	113	99	107	107	114	74	27	9	33
Male										
2002	79	84	74	89	78	51	35	18	3	11
2001	71	57	77	69	52	43	29	7	2	18
2000	72	63	68	64	53	56	35	9	3	14
1999	84	77	67	54	51	54	32	14	1	29

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nontransport accidents (W00-X59,Y86)														
Hispanic														
Female														
2002	968	34	57	21	20	25	34	29	43	78	87	77	43	35
2001	926	52	48	23	22	19	33	38	55	67	64	49	40	27
2000	838	33	63	27	22	28	19	34	38	69	40	30	38	33
1999	822	36	72	20	16	29	21	31	44	67	65	41	36	13
Non-Hispanic[1]														
Both sexes														
2002	53,907	715	839	407	455	1,150	1,894	1,898	2,471	3,451	4,611	4,255	3,192	2,151
2001	50,183	709	882	443	448	1,089	1,727	1,669	2,047	3,251	3,990	3,739	2,777	1,852
2000	47,217	627	979	485	463	1,088	1,505	1,644	2,047	3,144	3,761	3,317	2,472	1,771
1999	47,339	561	997	500	502	1,045	1,421	1,653	2,109	3,020	3,743	3,240	2,207	1,685
Male														
2002	32,295	412	539	273	326	928	1,550	1,476	1,844	2,472	3,255	2,992	2,240	1,498
2001	30,201	410	547	293	320	883	1,435	1,317	1,543	2,371	2,891	2,773	2,008	1,280
2000	28,640	361	617	325	341	902	1,225	1,329	1,559	2,370	2,788	2,500	1,866	1,249
1999	28,701	335	613	323	367	870	1,162	1,330	1,642	2,265	2,842	2,473	1,652	1,191
Female														
2002	21,612	303	300	134	129	222	344	422	627	979	1,356	1,263	952	653
2001	19,982	299	335	150	128	206	292	352	504	880	1,099	966	769	572
2000	18,577	266	362	160	122	186	280	315	488	774	973	817	606	522
1999	18,638	226	384	177	135	175	259	323	467	755	901	767	555	494
Non-Hispanic White														
Both sexes														
2002	45,916	403	565	224	281	904	1,614	1,576	2,065	2,811	3,835	3,421	2,529	1,721
2001	42,509	426	653	282	284	841	1,463	1,347	1,680	2,650	3,196	2,930	2,163	1,482
2000	39,728	376	676	289	308	832	1,234	1,327	1,644	2,542	2,995	2,584	1,896	1,408
1999	39,523	337	693	286	331	796	1,121	1,326	1,640	2,449	2,970	2,482	1,699	1,325
Male														
2002	27,024	244	359	151	203	725	1,314	1,223	1,549	2,008	2,726	2,428	1,747	1,177
2001	25,133	245	416	174	206	682	1,214	1,066	1,279	1,950	2,351	2,169	1,539	1,006
2000	23,615	222	433	196	232	689	1,015	1,073	1,270	1,944	2,234	1,942	1,419	976
1999	23,431	200	425	180	243	655	913	1,077	1,294	1,853	2,269	1,887	1,263	913
Female														
2002	18,892	159	206	73	78	179	300	353	516	803	1,109	993	782	544
2001	17,376	181	237	108	78	159	249	281	401	700	845	761	624	476
2000	16,113	154	243	93	76	143	219	254	374	598	761	642	477	432
1999	16,092	137	268	106	88	141	208	249	346	596	701	595	436	412
Non-Hispanic Black														
Both sexes														
2002	6,620	288	229	153	152	186	219	255	332	537	657	742	593	369

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)										
Hispanic										
Female										
2002	31	35	56	70	70	51	45	24	3	-
2001	29	34	57	54	67	70	46	27	4	1
2000	33	26	41	53	74	64	49	19	5	-
1999	27	36	32	53	56	60	42	13	8	4
Non-Hispanic[1]										
Both sexes										
2002	1,733	1,946	2,518	3,905	5,225	5,456	3,836	1,502	279	18
2001	1,677	1,834	2,469	3,897	5,018	5,240	3,686	1,415	303	21
2000	1,600	1,680	2,485	3,604	4,543	4,814	3,548	1,339	292	9
1999	1,533	1,841	2,559	3,747	4,820	5,041	3,558	1,299	243	15
Male										
2002	1,176	1,237	1,516	2,110	2,472	2,210	1,272	434	47	16
2001	1,110	1,161	1,425	2,128	2,404	2,192	1,290	346	57	17
2000	1,087	1,051	1,422	1,916	2,160	1,962	1,184	360	58	8
1999	1,057	1,159	1,513	1,944	2,345	2,034	1,180	335	56	13
Female										
2002	557	709	1,002	1,795	2,753	3,246	2,564	1,068	232	2
2001	567	673	1,044	1,769	2,614	3,048	2,396	1,069	246	4
2000	513	629	1,063	1,688	2,383	2,852	2,364	979	234	1
1999	476	682	1,046	1,803	2,475	3,007	2,378	964	187	2
Non-Hispanic White										
Both sexes										
2002	1,426	1,594	2,192	3,502	4,840	5,137	3,596	1,409	255	16
2001	1,357	1,549	2,132	3,492	4,621	4,897	3,452	1,321	274	17
2000	1,306	1,387	2,145	3,227	4,192	4,523	3,329	1,242	261	5
1999	1,184	1,514	2,157	3,310	4,427	4,677	3,353	1,216	221	9
Male										
2002	957	1,003	1,327	1,894	2,281	2,077	1,176	403	38	14
2001	890	983	1,237	1,907	2,195	2,046	1,199	314	50	15
2000	881	858	1,237	1,699	1,984	1,836	1,099	322	50	4
1999	831	942	1,275	1,703	2,162	1,879	1,093	315	51	8
Female										
2002	469	591	865	1,608	2,559	3,060	2,420	1,006	217	2
2001	467	566	895	1,585	2,426	2,851	2,253	1,007	224	2
2000	425	529	908	1,528	2,208	2,687	2,230	920	211	1
1999	353	572	882	1,607	2,265	2,798	2,260	901	170	1
Non-Hispanic Black										
Both sexes										
2002	262	277	257	316	277	245	182	74	17	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Nontransport accidents (W00-X59,Y86)														
Non-Hispanic Black														
Both sexes														
2001	6,436	254	204	136	140	182	209	243	299	505	706	718	556	319
2000	6,330	228	257	179	131	193	207	252	336	519	675	643	509	319
1999	6,723	207	260	193	148	193	239	272	391	480	691	695	450	318
Male														
2002	4,377	154	152	101	113	160	183	200	242	389	440	502	439	277
2001	4,238	151	119	100	98	150	169	184	211	349	480	537	422	234
2000	4,253	124	155	121	94	161	165	200	241	366	487	490	393	243
1999	4,532	122	164	130	108	168	204	211	285	345	514	537	347	247
Female														
2002	2,243	134	77	52	39	26	36	55	90	148	217	240	154	92
2001	2,198	103	85	36	42	32	40	59	88	156	226	181	134	85
2000	2,077	104	102	58	37	32	42	52	95	153	188	153	116	76
1999	2,191	85	96	63	40	25	35	61	106	135	177	158	103	71
Origin not stated[2]														
Both sexes														
2002	284	3	3	3	3	6	8	11	18	23	26	25	26	21
2001	317	5	3	5	3	3	11	8	16	29	30	37	19	22
2000	321	4	4	3	2	3	10	11	17	24	36	44	35	16
1999	328	1	5	-	2	6	6	10	19	25	41	35	31	26
Male														
2002	212	2	2	2	2	4	6	9	15	20	20	19	20	21
2001	242	4	3	3	3	3	10	6	13	26	24	29	17	19
2000	236	3	1	1	1	1	10	9	14	20	28	34	26	13
1999	244	-	2	-	2	5	6	8	16	20	28	30	27	19
Female														
2002	72	1	1	1	1	2	2	2	3	3	6	6	6	-
2001	75	1	-	2	-	-	1	2	3	3	6	8	2	3
2000	85	1	3	2	1	2	-	2	3	4	8	10	9	3
1999	84	1	3	-	-	1	-	2	3	5	13	5	4	7
Falls (W00-W19)														
All origins														
Both sexes														
2002	16,257	16	37	18	24	83	164	130	177	256	408	487	530	515
2001	15,019	23	32	33	33	88	168	160	180	272	375	520	504	463
2000	13,322	8	36	16	21	99	138	138	165	238	370	409	462	448
1999	13,162	12	55	25	28	112	130	156	187	276	369	447	377	398
Male														
2002	8,463	9	28	9	18	65	146	118	151	191	331	378	394	379
2001	8,089	15	25	22	24	79	153	142	153	233	311	426	395	344

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nontransport accidents (W00-X59,Y86)										
Non-Hispanic Black										
Both sexes										
2001	271	244	285	321	311	257	182	68	25	1
2000	247	250	285	312	277	229	173	77	28	4
1999	299	293	336	363	330	290	176	76	18	5
Male										
2002	187	184	147	171	138	97	71	24	5	1
2001	184	157	163	172	166	99	66	20	6	1
2000	179	163	156	178	136	93	67	30	7	4
1999	192	194	206	198	146	121	68	17	4	4
Female										
2002	75	93	110	145	139	148	111	50	12	-
2001	87	87	122	149	145	158	116	48	19	-
2000	68	87	129	134	141	136	106	47	21	-
1999	107	99	130	165	184	169	108	59	14	1
Origin not stated[2]										
Both sexes										
2002	9	10	12	19	15	13	9	4	-	17
2001	13	14	11	9	15	10	9	7	1	37
2000	11	7	13	15	10	14	9	3	1	29
1999	16	11	6	16	17	19	11	4	-	21
Male										
2002	9	9	10	9	12	7	1	2	-	11
2001	11	8	8	7	9	3	4	-	-	32
2000	9	7	7	9	3	6	5	1	-	28
1999	14	9	5	9	10	8	6	2	-	18
Female										
2002	-	1	2	10	3	6	8	2	-	6
2001	2	6	3	2	6	7	5	7	1	5
2000	2	-	6	6	7	8	4	2	1	1
1999	2	2	1	7	7	11	5	2	-	3
Falls (W00-W19)										
All origins										
Both sexes										
2002	574	759	1,208	2,011	2,869	3,032	2,045	784	129	1
2001	541	747	1,086	1,873	2,567	2,647	1,850	718	135	4
2000	501	595	1,065	1,641	2,200	2,352	1,682	618	120	-
1999	489	663	1,000	1,636	2,220	2,308	1,625	533	112	4
Male										
2002	392	474	728	1,079	1,365	1,262	691	231	24	-
2001	374	469	621	1,041	1,236	1,149	655	188	31	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Falls (W00-W19)														
All origins														
Male														
2000	7,122	6	23	14	16	88	115	122	150	199	297	332	358	326
1999	7,109	8	37	19	21	95	114	137	157	225	303	366	295	296
Female														
2002	7,794	7	9	9	6	18	18	12	26	65	77	109	136	136
2001	6,930	8	7	11	9	9	15	18	27	39	64	94	109	119
2000	6,200	2	13	2	5	11	23	16	15	39	73	77	104	122
1999	6,053	4	18	6	7	17	16	19	30	51	66	81	82	102
Hispanic														
Both sexes														
2002	832	4	9	4	2	12	39	32	37	40	54	44	46	45
2001	757	7	6	10	4	15	29	35	38	44	56	52	36	40
2000	693	3	9	1	2	18	28	36	37	31	35	39	38	35
1999	652	4	9	1	2	13	32	35	33	33	37	34	31	30
Male														
2002	563	3	7	1	2	12	37	30	35	31	50	35	35	36
2001	525	4	5	9	4	15	29	34	37	40	53	49	26	32
2000	485	2	8	1	2	18	26	34	36	27	35	35	34	27
1999	464	4	6	1	2	12	31	34	31	30	32	28	28	28
Female														
2002	269	1	2	3	-	-	2	2	2	9	4	9	11	9
2001	232	3	1	1	-	-	-	1	1	4	3	3	10	8
2000	208	1	1	-	-	-	2	2	1	4	-	4	4	8
1999	188	-	3	-	-	1	1	1	2	3	5	6	3	2
Non-Hispanic[1]														
Both sexes														
2002	15,365	12	28	14	22	71	125	98	140	215	353	439	477	467
2001	14,204	15	26	22	29	73	139	125	141	227	317	461	468	421
2000	12,585	5	27	15	19	81	110	101	128	205	332	366	418	409
1999	12,447	8	46	24	26	99	98	121	152	242	331	408	345	363
Male														
2002	7,856	6	21	8	16	53	109	88	116	159	280	339	352	340
2001	7,527	10	20	13	20	64	124	108	115	192	256	370	369	310
2000	6,603	4	15	13	14	70	89	87	114	170	259	293	319	295
1999	6,602	4	31	18	19	83	83	103	124	194	270	333	266	264
Female														
2002	7,509	6	7	6	6	18	16	10	24	56	73	100	125	127
2001	6,677	5	6	9	9	9	15	17	26	35	61	91	99	111
2000	5,982	1	12	2	5	11	21	14	14	35	73	73	99	114
1999	5,845	4	15	6	7	16	15	18	28	48	61	75	79	99

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)										
All origins										
Male										
2000	354	378	613	895	1,052	992	592	173	27	-
1999	354	433	594	840	1,110	964	568	140	29	4
Female										
2002	182	285	480	932	1,504	1,770	1,354	553	105	1
2001	167	278	465	832	1,331	1,498	1,195	530	104	1
2000	147	217	452	746	1,148	1,360	1,090	445	93	-
1999	135	230	406	796	1,110	1,344	1,057	393	83	-
Hispanic										
Both sexes										
2002	42	46	72	80	95	57	46	26	-	-
2001	38	35	68	67	64	53	38	17	4	1
2000	39	40	46	60	74	67	40	11	4	-
1999	33	47	37	55	61	65	41	12	5	2
Male										
2002	29	33	39	40	50	32	16	10	-	-
2001	30	21	36	33	29	20	14	3	1	1
2000	26	28	32	34	28	31	16	3	2	-
1999	27	28	24	29	31	32	17	6	1	2
Female										
2002	13	13	33	40	45	25	30	16	-	-
2001	8	14	32	34	35	33	24	14	3	-
2000	13	12	14	26	46	36	24	8	2	-
1999	6	19	13	26	30	33	24	6	4	-
Non-Hispanic[1]										
Both sexes										
2002	528	710	1,132	1,920	2,765	2,967	1,997	756	129	-
2001	500	704	1,014	1,805	2,494	2,588	1,808	697	130	-
2000	460	553	1,016	1,577	2,123	2,281	1,637	606	116	-
1999	450	611	961	1,571	2,150	2,235	1,579	519	107	1
Male										
2002	359	438	685	1,034	1,308	1,226	675	220	24	-
2001	341	443	581	1,007	1,202	1,127	640	185	30	-
2000	327	348	580	858	1,023	959	572	169	25	-
1999	323	400	569	805	1,073	928	549	134	28	1
Female										
2002	169	272	447	886	1,457	1,741	1,322	536	105	-
2001	159	261	433	798	1,292	1,461	1,168	512	100	-
2000	133	205	436	719	1,100	1,322	1,065	437	91	-
1999	127	211	392	766	1,077	1,307	1,030	385	79	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Falls (W00-W19)														
Non-Hispanic White														
Both sexes														
2002	14,157	8	15	11	17	60	108	87	116	176	290	364	398	391
2001	13,037	8	23	17	25	65	128	106	120	182	253	373	395	350
2000	11,537	2	15	11	16	73	99	88	104	163	275	311	335	336
1999	11,295	4	34	17	22	87	81	101	115	206	271	323	270	291
Male														
2002	7,107	3	11	6	13	44	93	78	97	128	231	284	290	284
2001	6,785	5	17	11	16	56	117	93	98	154	206	295	308	259
2000	5,949	1	6	9	13	64	80	77	92	136	218	253	252	239
1999	5,860	3	23	13	16	74	70	87	94	163	217	265	209	209
Female														
2002	7,050	5	4	5	4	16	15	9	19	48	59	80	108	107
2001	6,252	3	6	6	9	9	11	13	22	28	47	78	87	91
2000	5,588	1	9	2	3	9	19	11	12	27	57	58	83	97
1999	5,435	1	11	4	6	13	11	14	21	43	54	58	61	82
Non-Hispanic Black														
Both sexes														
2002	821	3	7	3	3	8	9	8	15	31	44	63	65	56
2001	837	6	3	4	2	4	9	9	17	33	57	62	66	55
2000	745	2	12	4	3	5	7	5	17	28	43	45	66	60
1999	865	2	11	6	4	8	13	13	29	26	47	73	63	57
Male														
2002	518	2	6	2	3	6	9	7	13	25	33	47	51	41
2001	537	4	3	2	2	4	5	6	13	27	45	54	55	40
2000	471	2	9	4	1	3	7	3	15	22	32	35	53	47
1999	554	1	8	4	3	6	10	11	23	23	42	59	48	43
Female														
2002	303	1	1	1	-	2	-	1	2	6	11	16	14	15
2001	300	2	-	2	-	-	4	3	4	6	12	8	11	15
2000	274	-	3	-	2	2	-	2	2	6	11	10	13	13
1999	311	1	3	2	1	2	3	2	6	3	5	14	15	14
Origin not stated[2]														
Both sexes														
2002	60	-	-	-	-	-	-	-	-	1	1	4	7	3
2001	58	1	-	1	-	-	-	-	1	1	2	7	-	2
2000	44	-	-	-	-	-	-	1	-	2	3	4	6	4
1999	63	-	-	-	-	-	-	-	2	1	1	5	1	5
Male														
2002	44	-	-	-	-	-	-	-	-	1	1	4	7	3
2001	37	1	-	-	-	-	-	-	1	1	2	7	-	2
2000	34	-	-	-	-	-	-	1	-	2	3	4	5	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)										
Non-Hispanic White										
Both sexes										
2002	462	621	1,023	1,784	2,635	2,846	1,897	726	122	-
2001	430	618	922	1,675	2,369	2,465	1,727	666	120	-
2000	389	473	920	1,475	2,012	2,191	1,570	572	107	-
1999	362	524	855	1,456	2,044	2,133	1,508	492	98	1
Male										
2002	312	381	621	959	1,240	1,170	631	209	22	-
2001	292	389	532	933	1,134	1,068	607	169	26	-
2000	280	297	532	796	968	918	540	155	23	-
1999	262	339	507	739	1,020	879	518	125	27	1
Female										
2002	150	240	402	825	1,395	1,676	1,266	517	100	-
2001	138	229	390	742	1,235	1,397	1,120	497	94	-
2000	109	176	388	679	1,044	1,273	1,030	417	84	-
1999	100	185	348	717	1,024	1,254	990	367	71	-
Non-Hispanic Black										
Both sexes										
2002	54	56	79	85	69	71	65	22	5	-
2001	53	72	71	94	79	69	49	16	7	-
2000	54	63	71	65	72	50	45	21	7	-
1999	70	80	70	73	80	59	53	21	7	-
Male										
2002	37	36	45	48	37	33	28	8	1	-
2001	36	46	39	52	45	29	19	8	3	-
2000	40	40	36	39	31	20	22	9	1	-
1999	47	57	43	41	36	23	19	6	1	-
Female										
2002	17	20	34	37	32	38	37	14	4	-
2001	17	26	32	42	34	40	30	8	4	-
2000	14	23	35	26	41	30	23	12	6	-
1999	23	23	27	32	44	36	34	15	6	-
Origin not stated[2]										
Both sexes										
2002	4	3	4	11	9	8	2	2	-	1
2001	3	8	4	1	9	6	4	4	1	3
2000	2	2	3	4	3	4	5	1	-	-
1999	6	5	2	10	9	8	5	2	-	1
Male										
2002	4	3	4	5	7	4	-	1	-	-
2001	3	5	4	1	5	2	1	-	-	2
2000	1	2	1	3	1	2	4	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Falls (W00-W19)														
Origin not stated[2]														
Male														
1999	43	-	-	-	-	-	-	-	2	1	1	5	1	4
Female														
2002	16	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	21	-	-	1	-	-	-	-	-	-	-	-	-	-
2000	10	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	20	-	-	-	-	-	-	-	-	-	-	-	-	1
Accidental discharge of firearms (W32-W34)														
All origins														
Both sexes														
2002	762	1	11	14	34	107	103	75	68	52	71	52	43	36
2001	802	-	15	18	39	110	96	71	51	70	76	61	42	44
2000	776	1	18	18	49	107	95	74	57	75	78	38	40	36
1999	824	-	12	19	57	126	125	76	67	67	69	51	41	25
Male														
2002	667	1	7	10	29	101	94	70	59	42	56	45	39	32
2001	690	-	12	14	35	102	91	63	40	57	64	51	36	34
2000	671	-	11	16	46	97	86	65	44	60	65	31	38	30
1999	707	-	7	13	47	115	114	62	60	56	56	46	32	22
Female														
2002	95	-	4	4	5	6	9	5	9	10	15	7	4	4
2001	112	-	3	4	4	8	5	8	11	13	12	10	6	10
2000	105	1	7	2	3	10	9	9	13	15	13	7	2	6
1999	117	-	5	6	10	11	11	14	7	11	13	5	9	3
Hispanic														
Both sexes														
2002	60	-	3	3	2	13	11	11	7	4	1	2	1	1
2001	86	-	1	1	3	19	17	12	9	7	5	3	3	-
2000	59	-	2	2	4	12	13	7	5	6	5	2	-	1
1999	66	-	-	-	3	15	16	7	10	4	4	4	-	1
Male														
2002	51	-	1	1	2	11	10	11	7	3	-	2	1	1
2001	76	-	1	1	2	17	16	10	6	7	5	2	3	-
2000	50	-	1	1	3	11	13	7	3	4	5	2	-	-
1999	61	-	-	-	3	14	15	6	10	3	4	3	-	1
Female														
2002	9	-	2	2	-	2	1	-	-	1	1	-	-	-
2001	10	-	-	-	1	2	1	2	3	-	-	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)										
Origin not stated[2]										
Male										
1999	4	5	1	6	6	4	2	-	-	1
Female										
2002	-	-	-	6	2	4	2	1	-	1
2001	-	3	-	-	4	4	3	4	1	1
2000	1	-	2	1	2	2	1	-	-	-
1999	2	-	1	4	3	4	3	2	-	-
Accidental discharge of firearms (W32-W34)										
All origins										
Both sexes										
2002	20	17	19	18	14	5	1	1	-	-
2001	25	19	14	24	17	6	2	-	-	2
2000	14	24	17	17	7	7	4	-	-	-
1999	23	15	18	15	8	8	2	-	-	-
Male										
2002	18	15	17	15	11	4	1	1	-	-
2001	21	16	10	22	14	5	2	-	-	1
2000	14	20	14	16	7	7	4	-	-	-
1999	19	13	18	13	7	5	2	-	-	-
Female										
2002	2	2	2	3	3	1	-	-	-	-
2001	4	3	4	2	3	1	-	-	-	1
2000	-	4	3	1	-	-	-	-	-	-
1999	4	2	-	2	1	3	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	1	-	-	-	-	-
2001	1	1	-	2	-	1	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	-	2	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	1	-	-	-	-	-
2001	1	1	-	2	-	1	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	-	2	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental discharge of firearms (W32-W34)														
Hispanic														
Female														
2000	9	-	1	1	1	1	-	-	2	2	-	-	-	1
1999	5	-	-	-	-	1	1	1	-	1	-	1	-	-
Non-Hispanic[1]														
Both sexes														
2002	700	1	8	11	32	93	92	64	61	48	70	50	42	35
2001	715	-	14	17	36	90	79	59	42	63	71	58	39	44
2000	717	1	16	16	45	95	82	67	52	69	73	36	40	35
1999	758	-	12	19	54	111	109	69	57	63	65	47	41	24
Male														
2002	614	1	6	9	27	89	84	59	52	39	56	43	38	31
2001	613	-	11	13	33	84	75	53	34	50	59	49	33	34
2000	621	-	10	15	43	86	73	58	41	56	60	29	38	30
1999	646	-	7	13	44	101	99	56	50	53	52	43	32	21
Female														
2002	86	-	2	2	5	4	8	5	9	9	14	7	4	4
2001	102	-	3	4	3	6	4	6	8	13	12	9	6	10
2000	96	1	6	1	2	9	9	9	11	13	13	7	2	5
1999	112	-	5	6	10	10	10	13	7	10	13	4	9	3
Non-Hispanic White														
Both sexes														
2002	531	1	6	7	24	56	62	40	42	35	62	42	36	31
2001	591	-	9	13	28	61	56	43	39	57	59	54	38	38
2000	555	1	9	9	37	65	52	46	37	54	64	28	37	33
1999	585	-	5	9	45	73	72	55	43	52	56	41	36	20
Male														
2002	459	1	4	5	20	53	57	35	35	29	48	36	33	28
2001	506	-	7	10	28	56	53	38	33	45	51	45	33	29
2000	483	-	5	9	35	60	46	38	28	45	54	24	35	28
1999	495	-	3	4	39	66	65	43	37	43	44	38	27	18
Female														
2002	72	-	2	2	4	3	5	5	7	6	14	6	3	3
2001	85	-	2	3	-	5	3	5	6	12	8	9	5	9
2000	72	1	4	-	2	5	6	8	9	9	10	4	2	5
1999	90	-	2	5	6	7	7	12	6	9	12	3	9	2
Non-Hispanic Black														
Both sexes														
2002	153	-	2	3	6	35	25	23	17	13	7	8	5	4
2001	110	-	5	2	6	25	23	14	3	4	11	4	1	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental discharge of firearms (W32-W34)										
Hispanic										
Female										
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	20	17	19	17	13	5	1	1	-	-
2001	24	18	14	22	17	5	2	-	-	1
2000	14	24	17	17	7	7	4	-	-	-
1999	23	13	18	15	8	8	2	-	-	-
Male										
2002	18	15	17	14	10	4	1	1	-	-
2001	20	15	10	20	14	4	2	-	-	-
2000	14	20	14	16	7	7	4	-	-	-
1999	19	11	18	13	7	5	2	-	-	-
Female										
2002	2	2	2	3	3	1	-	-	-	-
2001	4	3	4	2	3	1	-	-	-	1
2000	-	4	3	1	-	-	-	-	-	-
1999	4	2	-	2	1	3	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	18	16	19	14	13	5	1	1	-	-
2001	23	17	12	21	15	5	2	-	-	1
2000	13	22	15	16	6	7	4	-	-	-
1999	20	11	17	13	7	8	2	-	-	-
Male										
2002	17	14	17	11	10	4	1	1	-	-
2001	19	14	8	19	12	4	2	-	-	-
2000	13	19	12	15	6	7	4	-	-	-
1999	18	9	17	11	6	5	2	-	-	-
Female										
2002	1	2	2	3	3	1	-	-	-	-
2001	4	3	4	2	3	1	-	-	-	1
2000	-	3	3	1	-	-	-	-	-	-
1999	2	2	-	2	1	3	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	2	-	-	3	-	-	-	-	-	-
2001	1	1	2	1	2	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental discharge of firearms (W32-W34)														
Non-Hispanic Black														
Both sexes														
2000	137	-	5	5	7	28	24	18	14	13	6	7	3	2
1999	156	-	7	9	8	36	31	12	11	9	9	6	5	4
Male														
2002	140	-	2	3	5	34	23	23	15	10	7	7	4	3
2001	97	-	4	1	5	24	22	13	1	4	8	4	-	4
2000	117	-	3	5	7	25	21	17	12	9	5	4	3	2
1999	135	-	4	8	4	33	28	12	10	8	8	5	5	3
Female														
2002	13	-	-	-	1	1	2	-	2	3	-	1	1	1
2001	13	-	1	1	1	1	1	1	2	-	3	-	1	1
2000	20	-	2	-	-	3	3	1	2	4	1	3	-	-
1999	21	-	3	1	4	3	3	-	1	1	1	1	-	1
Origin not stated[2]														
Both sexes														
2002	2	-	-	-	-	1	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	2	-	-	-	-	1	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Accidental drowning and submersion (W65-W74)														
All origins														
Both sexes														
2002	3,447	63	454	159	162	320	309	227	206	263	263	205	169	118
2001	3,281	68	458	168	165	322	274	212	162	221	241	214	145	109
2000	3,482	75	493	201	174	371	275	213	206	250	230	182	172	120
1999	3,529	68	490	192	177	359	288	212	234	250	225	198	172	122
Male														
2002	2,761	40	312	126	118	288	290	201	179	223	216	164	137	93

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental discharge of firearms (W32-W34)										
Non-Hispanic Black										
Both sexes										
2000	1	1	2	-	1	-	-	-	-	-
1999	3	2	1	2	1	-	-	-	-	-
Male										
2002	1	-	-	3	-	-	-	-	-	-
2001	1	1	2	1	2	-	-	-	-	-
2000	1	-	2	-	1	-	-	-	-	-
1999	1	2	1	2	1	-	-	-	-	-
Female										
2002	1	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	1	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Accidental drowning and submersion (W65-W74)										
All origins										
Both sexes										
2002	94	81	88	88	79	48	25	5	1	20
2001	97	78	81	85	72	56	15	4	-	34
2000	97	80	99	71	85	37	19	8	-	24
1999	97	84	101	88	70	45	16	7	-	34
Male										
2002	68	64	68	56	50	31	15	4	-	18

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental drowning and submersion (W65-W74)														
All origins														
Male														
2001	2,560	30	291	120	117	295	249	186	142	181	195	182	117	83
2000	2,735	43	316	137	130	339	244	184	180	212	194	156	143	102
1999	2,794	39	310	139	136	330	262	193	204	215	187	166	145	95
Female														
2002	686	23	142	33	44	32	19	26	27	40	47	41	32	25
2001	721	38	167	48	48	27	25	26	20	40	46	32	28	26
2000	747	32	177	64	44	32	31	29	26	38	36	26	29	18
1999	735	29	180	53	41	29	26	19	30	35	38	32	27	27
Hispanic														
Both sexes														
2002	464	18	65	16	13	69	74	45	35	29	26	18	12	8
2001	421	16	78	27	13	42	59	39	35	36	21	21	4	2
2000	433	9	75	29	14	73	74	34	28	25	16	11	9	7
1999	480	16	78	25	21	69	76	39	40	27	18	12	13	4
Male														
2002	409	13	48	15	8	64	73	43	35	24	24	16	12	6
2001	348	9	53	21	7	42	57	33	32	30	18	18	3	1
2000	350	6	44	16	6	65	70	32	28	21	15	11	7	6
1999	405	12	43	18	17	67	73	37	38	24	14	10	13	3
Female														
2002	55	5	17	1	5	5	1	2	-	5	2	2	-	2
2001	73	7	25	6	6	-	2	6	3	6	3	3	1	1
2000	83	3	31	13	8	8	4	2	-	4	1	-	2	1
1999	75	4	35	7	4	2	3	2	2	3	4	2	-	1
Non-Hispanic[1]														
Both sexes														
2002	2,952	45	387	141	148	250	235	178	169	232	235	187	155	108
2001	2,832	51	380	139	152	279	214	173	127	184	219	190	139	106
2000	3,018	65	417	171	160	297	199	177	178	224	213	169	163	112
1999	3,029	52	409	167	156	289	211	173	190	222	207	184	159	118
Male														
2002	2,326	27	262	110	109	223	217	155	143	197	190	148	123	85
2001	2,188	21	238	97	110	252	191	153	110	150	176	163	112	81
2000	2,358	37	272	120	124	273	172	150	152	190	178	143	136	95
1999	2,370	27	265	121	119	262	188	156	162	190	173	154	132	92
Female														
2002	626	18	125	31	39	27	18	23	26	35	45	39	32	23
2001	644	30	142	42	42	27	23	20	17	34	43	27	27	25

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental drowning and submersion (W65-W74)										
All origins										
Male										
2001	72	61	63	55	47	37	8	-	-	29
2000	77	63	70	38	50	22	7	5	-	23
1999	75	64	65	63	48	19	6	3	-	30
Female										
2002	26	17	20	32	29	17	10	1	1	2
2001	25	17	18	30	25	19	7	4	-	5
2000	20	17	29	33	35	15	12	3	-	1
1999	22	20	36	25	22	26	10	4	-	4
Hispanic										
Both sexes										
2002	7	4	8	8	2	1	1	-	-	5
2001	5	3	2	3	1	2	-	-	-	12
2000	4	3	9	1	3	-	1	-	-	8
1999	4	3	4	5	1	-	-	-	-	25
Male										
2002	4	4	6	6	1	1	1	-	-	5
2001	5	1	2	3	-	2	-	-	-	11
2000	2	3	7	1	2	-	-	-	-	8
1999	4	3	4	4	-	-	-	-	-	21
Female										
2002	3	-	2	2	1	-	-	-	-	-
2001	-	2	-	-	1	-	-	-	-	1
2000	2	-	2	-	1	-	1	-	-	-
1999	-	-	-	1	1	-	-	-	-	4
Non-Hispanic[1]										
Both sexes										
2002	87	77	79	80	75	47	24	5	1	7
2001	92	74	79	82	71	54	15	4	-	8
2000	93	77	89	70	82	36	18	7	-	1
1999	92	81	97	83	69	45	16	6	-	3
Male										
2002	64	60	61	50	47	30	14	4	-	7
2001	67	59	61	52	47	35	8	-	-	5
2000	75	60	62	37	48	21	7	5	-	1
1999	70	61	61	59	48	19	6	2	-	3
Female										
2002	23	17	18	30	28	17	10	1	1	-
2001	25	15	18	30	24	19	7	4	-	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental drowning and submersion (W65-W74)														
Non-Hispanic[1]														
Female														
2000	660	28	145	51	36	24	27	27	26	34	35	26	27	17
1999	659	25	144	46	37	27	23	17	28	32	34	30	27	26
Non-Hispanic White														
Both sexes														
2002	2,143	32	303	71	63	153	163	118	127	170	190	141	119	94
2001	2,082	36	321	70	65	165	157	113	89	138	168	146	119	82
2000	2,235	38	342	96	80	181	135	126	132	180	164	134	121	97
1999	2,230	30	337	81	61	178	139	113	137	178	171	161	130	96
Male														
2002	1,669	18	203	52	44	136	148	101	108	141	153	109	96	74
2001	1,577	16	205	40	47	149	135	97	74	109	136	126	94	61
2000	1,737	22	226	68	61	165	113	107	115	151	136	112	100	82
1999	1,708	14	217	53	46	159	120	99	117	148	142	131	110	74
Female														
2002	474	14	100	19	19	17	15	17	19	29	37	32	23	20
2001	505	20	116	30	18	16	22	16	15	29	32	20	25	21
2000	498	16	116	28	19	16	22	19	17	29	28	22	21	15
1999	522	16	120	28	15	19	19	14	20	30	29	30	20	22
Non-Hispanic Black														
Both sexes														
2002	604	10	62	57	77	70	56	42	32	43	31	33	24	11
2001	570	11	47	55	78	89	44	33	29	33	37	34	17	21
2000	580	25	54	66	68	84	44	34	24	35	42	23	29	9
1999	624	21	59	76	82	84	53	47	39	31	27	17	19	19
Male														
2002	502	6	43	46	61	68	54	39	26	38	28	28	19	10
2001	471	3	28	46	56	82	44	33	28	29	29	27	15	17
2000	470	14	33	47	54	81	42	29	23	31	36	21	24	8
1999	521	12	40	60	64	77	51	44	34	29	24	17	17	16
Female														
2002	102	4	19	11	16	2	2	3	6	5	3	5	5	1
2001	99	8	19	9	22	7	-	-	1	4	8	7	2	4
2000	110	11	21	19	14	3	2	5	1	4	6	2	5	1
1999	103	9	19	16	18	7	2	3	5	2	3	-	2	3
Origin not stated[2]														
Both sexes														
2002	31	-	2	2	1	1	-	4	2	2	2	-	2	2
2001	28	1	-	2	-	1	1	-	-	1	1	3	2	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental drowning and submersion (W65-W74)										
Non-Hispanic[1]										
Female										
2000	18	17	27	33	34	15	11	2	-	-
1999	22	20	36	24	21	26	10	4	-	-
Non-Hispanic White										
Both sexes										
2002	71	64	62	67	64	40	21	3	-	7
2001	76	62	71	73	61	48	14	4	-	4
2000	78	67	77	57	77	33	15	5	-	-
1999	78	69	78	69	63	39	16	4	-	2
Male										
2002	51	50	49	44	41	28	13	3	-	7
2001	53	50	55	47	41	31	8	-	-	3
2000	68	52	52	30	48	19	7	3	-	-
1999	59	50	49	49	44	17	6	2	-	2
Female										
2002	20	14	13	23	23	12	8	-	-	-
2001	23	12	16	26	20	17	6	4	-	1
2000	10	15	25	27	29	14	8	2	-	-
1999	19	19	29	20	19	22	10	2	-	-
Non-Hispanic Black										
Both sexes										
2002	10	9	9	10	7	6	2	2	1	-
2001	11	7	6	5	8	3	1	-	-	1
2000	8	7	8	9	3	3	2	2	-	1
1999	9	9	12	11	4	3	-	2	-	-
Male										
2002	8	8	7	5	4	2	1	1	-	-
2001	9	7	5	4	5	3	-	-	-	1
2000	5	6	7	4	-	2	-	2	-	1
1999	6	8	9	8	3	2	-	-	-	-
Female										
2002	2	1	2	5	3	4	1	1	1	-
2001	2	-	1	1	3	-	1	-	-	-
2000	3	1	1	5	3	1	2	-	-	-
1999	3	1	3	3	1	1	-	2	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	1	-	2	-	-	-	-	8
2001	-	1	-	-	-	-	-	-	-	14

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental drowning and submersion (W65-W74)														
Origin not stated[2]														
Both sexes														
2000	31	1	1	1	-	1	2	2	-	1	1	2	-	1
1999	20	-	3	-	-	1	1	-	4	1	-	2	-	-
Male														
2002	26	-	2	1	1	1	-	3	1	2	2	-	2	2
2001	24	-	-	2	-	1	1	-	-	1	1	1	2	1
2000	27	-	-	1	-	1	2	2	-	1	1	2	-	1
1999	19	-	2	-	-	1	1	-	4	1	-	2	-	-
Female														
2002	5	-	-	1	-	-	-	1	1	-	-	-	-	-
2001	4	1	-	-	-	-	-	-	-	-	-	2	-	-
2000	4	1	1	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	1	-	-	-	-	-	-	-	-	-	-	-
Accidental exposure to smoke, fire and flames (X00-X09)														
All origins														
Both sexes														
2002	3,159	36	221	152	101	86	107	112	154	147	214	213	235	183
2001	3,309	50	225	164	88	76	133	114	133	190	251	213	206	215
2000	3,377	37	290	182	84	79	113	120	121	188	214	214	225	187
1999	3,348	41	302	170	92	88	113	142	140	200	221	198	170	157
Male														
2002	1,935	21	126	89	62	61	71	75	95	101	150	146	146	111
2001	1,998	30	142	104	48	45	92	73	88	132	176	151	137	142
2000	2,025	20	174	109	36	44	77	78	79	133	135	145	158	126
1999	2,025	21	173	95	52	58	75	93	104	138	162	144	110	98
Female														
2002	1,224	15	95	63	39	25	36	37	59	46	64	67	89	72
2001	1,311	20	83	60	40	31	41	41	45	58	75	62	69	73
2000	1,352	17	116	73	48	35	36	42	42	55	79	69	67	61
1999	1,323	20	129	75	40	30	38	49	36	62	59	54	60	59
Hispanic														
Both sexes														
2002	223	5	36	17	10	11	14	7	10	14	13	12	10	15
2001	176	9	18	17	6	3	16	11	10	13	15	12	5	7
2000	178	4	21	15	4	14	7	13	6	8	17	2	9	7
1999	192	5	41	16	7	12	12	6	11	9	12	10	6	3
Male														
2002	144	3	18	10	7	9	10	4	5	14	7	9	9	10

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental drowning and submersion (W65-W74)										
Origin not stated[2]										
Both sexes										
2000	-	-	1	-	-	1	-	1	-	15
1999	1	-	-	-	-	-	-	1	-	6
Male										
2002	-	-	1	-	2	-	-	-	-	6
2001	-	1	-	-	-	-	-	-	-	13
2000	-	-	1	-	-	1	-	-	-	14
1999	1	-	-	-	-	-	-	1	-	6
Female										
2002	-	-	-	-	-	-	-	-	-	2
2001	-	-	-	-	-	-	-	-	-	1
2000	-	-	-	-	-	-	-	1	-	1
1999	-	-	-	-	-	-	-	-	-	-
Accidental exposure to smoke, fire and flames (X00-X09)										
All origins										
Both sexes										
2002	157	174	211	211	221	129	70	18	4	3
2001	163	190	221	225	198	159	72	18	2	3
2000	182	188	213	257	215	158	78	24	3	5
1999	162	193	223	258	229	150	74	18	3	4
Male										
2002	108	117	125	108	123	55	31	9	2	3
2001	94	112	117	106	98	69	30	8	1	3
2000	115	107	125	136	108	69	34	11	1	5
1999	96	111	133	128	119	67	34	9	1	4
Female										
2002	49	57	86	103	98	74	39	9	2	-
2001	69	78	104	119	100	90	42	10	1	-
2000	67	81	88	121	107	89	44	13	2	-
1999	66	82	90	130	110	83	40	9	2	-
Hispanic										
Both sexes										
2002	2	12	8	7	9	9	-	1	-	1
2001	6	5	4	4	5	9	-	1	-	-
2000	10	5	13	10	7	5	1	-	-	-
1999	11	8	3	8	7	3	2	-	-	-
Male										
2002	2	9	5	4	4	3	-	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental exposure to smoke, fire and flames (X00-X09)														
Hispanic														
Male														
2001	109	5	13	9	2	2	8	9	7	9	10	10	1	6
2000	105	2	11	9	1	5	6	9	5	6	11	1	6	5
1999	127	3	23	10	2	7	9	4	8	6	10	9	3	1
Female														
2002	79	2	18	7	3	2	4	3	5	-	6	3	1	5
2001	67	4	5	8	4	1	8	2	3	4	5	2	4	1
2000	73	2	10	6	3	9	1	4	1	2	6	1	3	2
1999	65	2	18	6	5	5	3	2	3	3	2	1	3	2
Non-Hispanic[1]														
Both sexes														
2002	2,918	31	184	135	90	75	93	104	143	133	201	200	224	164
2001	3,106	41	205	145	81	73	117	101	122	177	233	200	198	206
2000	3,178	31	268	167	79	65	104	107	115	179	196	211	215	178
1999	3,133	36	260	154	85	73	100	136	128	187	208	186	161	152
Male														
2002	1,780	18	108	79	54	52	61	70	89	87	143	137	137	97
2001	1,873	25	127	94	45	43	84	64	81	123	163	140	134	134
2000	1,907	16	163	100	35	39	69	69	74	127	123	143	152	120
1999	1,879	18	150	85	50	48	65	89	95	128	151	133	105	96
Female														
2002	1,138	13	76	56	36	23	32	34	54	46	58	63	87	67
2001	1,233	16	78	51	36	30	33	37	41	54	70	60	64	72
2000	1,271	15	105	67	44	26	35	38	41	52	73	68	63	58
1999	1,254	18	110	69	35	25	35	47	33	59	57	53	56	56
Non-Hispanic White														
Both sexes														
2002	2,055	16	90	61	51	50	63	64	107	96	162	154	163	122
2001	2,204	26	124	85	51	53	86	60	93	127	170	143	141	146
2000	2,194	17	139	90	49	46	74	71	80	126	141	153	167	122
1999	2,172	21	137	80	56	55	70	101	98	129	157	134	119	118
Male														
2002	1,271	12	52	38	35	35	43	47	68	61	112	108	96	73
2001	1,322	14	82	54	28	31	65	44	61	86	121	98	92	97
2000	1,330	7	88	53	23	27	52	45	54	91	88	105	117	85
1999	1,325	11	80	42	34	37	46	70	75	88	116	96	77	72
Female														
2002	784	4	38	23	16	15	20	17	39	35	50	46	67	49
2001	882	12	42	31	23	22	21	16	32	41	49	45	49	49

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)										
Hispanic										
Male										
2001	3	3	3	3	2	4	-	-	-	-
2000	4	4	6	6	4	4	-	-	-	-
1999	10	6	2	5	4	3	2	-	-	-
Female										
2002	-	3	3	3	5	6	-	-	-	-
2001	3	2	1	1	3	5	-	1	-	-
2000	6	1	7	4	3	1	1	-	-	-
1999	1	2	1	3	3	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	153	161	201	202	212	120	70	17	4	1
2001	155	183	215	219	193	150	71	17	2	2
2000	171	183	199	244	207	152	77	24	3	3
1999	150	185	219	249	221	147	72	18	3	3
Male										
2002	104	107	120	104	119	52	31	8	2	1
2001	90	108	114	103	96	65	29	8	1	2
2000	110	103	119	128	104	64	34	11	1	3
1999	85	105	130	122	115	64	32	9	1	3
Female										
2002	49	54	81	98	93	68	39	9	2	-
2001	65	75	101	116	97	85	42	9	1	-
2000	61	80	80	116	103	88	43	13	2	-
1999	65	80	89	127	106	83	40	9	2	-
Non-Hispanic White										
Both sexes										
2002	105	119	149	155	170	95	49	10	3	1
2001	115	133	158	164	147	117	53	9	1	2
2000	120	134	144	179	149	123	50	17	-	3
1999	100	128	153	178	165	105	51	13	2	2
Male										
2002	73	78	88	81	101	44	20	4	1	1
2001	63	77	80	73	76	53	21	4	-	2
2000	75	72	87	94	78	56	22	8	-	3
1999	64	71	89	83	91	49	24	7	1	2
Female										
2002	32	41	61	74	69	51	29	6	2	-
2001	52	56	78	91	71	64	32	5	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental exposure to smoke, fire and flames (X00-X09)														
Non-Hispanic White														
Female														
2000	864	10	51	37	26	19	22	26	26	35	53	48	50	37
1999	847	10	57	38	22	18	24	31	23	41	41	38	42	46
Non-Hispanic Black														
Both sexes														
2002	765	13	88	64	31	21	27	31	30	29	31	40	58	38
2001	829	14	80	58	27	14	28	36	27	45	58	54	51	53
2000	915	14	120	74	27	16	26	30	34	44	53	51	46	51
1999	887	15	110	74	28	16	24	29	23	54	46	46	40	34
Male														
2002	454	6	53	37	16	14	17	17	18	21	24	25	39	22
2001	501	10	44	38	15	8	16	16	18	34	39	40	39	32
2000	539	9	70	46	11	10	15	19	20	30	33	34	33	33
1999	507	7	63	43	15	9	15	15	14	38	31	32	27	24
Female														
2002	311	7	35	27	15	7	10	14	12	8	7	15	19	16
2001	328	4	36	20	12	6	12	20	9	11	19	14	12	21
2000	376	5	50	28	16	6	11	11	14	14	20	17	13	18
1999	380	8	47	31	13	7	9	14	9	16	15	14	13	10
Origin not stated[2]														
Both sexes														
2002	18	-	1	-	1	-	-	1	1	-	-	1	1	4
2001	27	-	2	2	1	-	-	2	1	-	3	1	3	2
2000	21	2	1	-	1	-	2	-	-	1	1	1	1	2
1999	23	-	1	-	-	3	1	-	1	4	1	2	3	2
Male														
2002	11	-	-	-	1	-	-	1	1	-	-	-	-	4
2001	16	-	2	1	1	-	-	-	-	-	3	1	2	2
2000	13	2	-	-	-	-	2	-	-	-	1	1	-	1
1999	19	-	-	-	-	3	1	-	1	4	1	2	2	1
Female														
2002	7	-	1	-	-	-	-	-	-	-	-	1	1	-
2001	11	-	-	1	-	-	-	2	1	-	-	-	1	-
2000	8	-	1	-	1	-	-	-	-	1	-	-	1	1
1999	4	-	1	-	-	-	-	-	-	-	-	-	1	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental exposure to smoke, fire and flames (X00-X09)										
Non-Hispanic White										
Female										
2000	45	62	57	85	71	67	28	9	-	-
1999	36	57	64	95	74	56	27	6	1	-
Non-Hispanic Black										
Both sexes										
2002	46	36	47	42	40	25	20	7	1	-
2001	36	46	53	49	43	31	17	8	1	-
2000	49	45	53	61	56	29	26	7	3	-
1999	46	52	61	67	53	41	21	5	1	1
Male										
2002	30	25	28	20	18	8	11	4	1	-
2001	24	28	33	26	18	11	7	4	1	-
2000	34	29	31	33	25	8	12	3	1	-
1999	20	30	39	36	23	15	8	2	-	1
Female										
2002	16	11	19	22	22	17	9	3	-	-
2001	12	18	20	23	25	20	10	4	-	-
2000	15	16	22	28	31	21	14	4	2	-
1999	26	22	22	31	30	26	13	3	1	-
Origin not stated[2]										
Both sexes										
2002	2	1	2	2	-	-	-	-	-	1
2001	2	2	2	2	-	-	1	-	-	1
2000	1	-	1	3	1	1	-	-	-	2
1999	1	-	1	1	1	-	-	-	-	1
Male										
2002	2	1	-	-	-	-	-	-	-	1
2001	1	1	-	-	-	-	1	-	-	1
2000	1	-	-	2	-	1	-	-	-	2
1999	1	-	1	1	-	-	-	-	-	1
Female										
2002	-	-	2	2	-	-	-	-	-	-
2001	1	1	2	2	-	-	-	-	-	-
2000	-	-	1	1	1	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental poisoning and exposure to noxious substances (X40-X49)														
All origins														
Both sexes														
2002	17,550	26	31	15	28	486	1,193	1,310	1,806	2,597	3,410	2,946	1,736	754
2001	14,078	15	31	18	32	406	956	1,078	1,429	2,307	2,729	2,266	1,281	530
2000	12,757	14	32	17	28	351	809	980	1,400	2,159	2,504	1,992	1,069	447
1999	12,186	12	34	12	28	260	704	952	1,403	2,082	2,467	1,926	918	449
Male														
2002	12,059	21	14	10	18	366	940	1,003	1,316	1,807	2,315	1,960	1,176	491
2001	9,885	8	21	7	16	311	769	832	1,030	1,626	1,896	1,610	893	341
2000	9,138	11	21	12	15	273	647	779	1,041	1,554	1,830	1,455	768	260
1999	8,887	11	17	6	19	195	559	757	1,071	1,523	1,826	1,456	650	298
Female														
2002	5,491	5	17	5	10	120	253	307	490	790	1,095	986	560	263
2001	4,193	7	10	11	16	95	187	246	399	681	833	656	388	189
2000	3,619	3	11	5	13	78	162	201	359	605	674	537	301	187
1999	3,299	1	17	6	9	65	145	195	332	559	641	470	268	151
Hispanic														
Both sexes														
2002	1,652	6	4	4	5	57	138	160	204	262	324	243	125	53
2001	1,324	1	5	2	8	48	96	153	181	251	224	165	98	49
2000	1,308	2	4	1	3	38	94	133	211	267	226	158	93	38
1999	1,502	4	3	2	4	41	105	155	228	311	274	191	98	37
Male														
2002	1,324	5	-	2	2	44	117	141	171	206	260	184	104	39
2001	1,061	1	3	1	1	35	76	131	143	205	180	135	83	40
2000	1,090	2	2	1	1	31	82	112	183	214	201	139	72	24
1999	1,271	4	3	1	3	28	95	133	199	262	231	165	75	33
Female														
2002	328	1	4	2	3	13	21	19	33	56	64	59	21	14
2001	263	-	2	1	7	13	20	22	38	46	44	30	15	9
2000	218	-	2	-	2	7	12	21	28	53	25	19	21	14
1999	231	-	-	1	1	13	10	22	29	49	43	26	23	4
Non-Hispanic[1]														
Both sexes														
2002	15,789	20	27	11	23	428	1,048	1,144	1,591	2,318	3,064	2,687	1,603	695
2001	12,631	14	26	16	24	357	853	921	1,236	2,034	2,485	2,079	1,171	471
2000	11,311	12	27	14	24	312	710	841	1,177	1,878	2,253	1,805	953	401
1999	10,542	8	31	10	23	219	595	790	1,165	1,753	2,154	1,715	801	402
Male														
2002	10,649	16	14	8	16	322	818	857	1,136	1,587	2,039	1,763	1,068	446
2001	8,720	7	18	6	15	275	687	697	877	1,402	1,701	1,458	799	293

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)										
All origins										
Both sexes										
2002	344	218	165	154	141	103	64	13	1	9
2001	268	161	130	153	126	91	47	12	2	10
2000	241	146	132	118	127	109	60	14	1	7
1999	219	158	142	138	127	100	31	12	-	12
Male										
2002	199	126	83	71	65	38	25	8	-	7
2001	159	88	82	83	54	28	18	2	1	10
2000	141	79	68	49	53	51	23	2	-	6
1999	150	98	74	63	53	38	10	5	-	8
Female										
2002	145	92	82	83	76	65	39	5	1	2
2001	109	73	48	70	72	63	29	10	1	-
2000	100	67	64	69	74	58	37	12	1	1
1999	69	60	68	75	74	62	21	7	-	4
Hispanic										
Both sexes										
2002	26	15	11	8	5	1	1	-	-	-
2001	15	10	9	2	5	-	-	1	-	1
2000	15	9	6	2	3	3	2	-	-	-
1999	21	12	6	4	1	1	1	-	-	3
Male										
2002	21	11	7	6	3	-	1	-	-	-
2001	10	7	6	2	1	-	-	-	-	1
2000	11	7	3	1	1	1	2	-	-	-
1999	18	11	4	2	-	1	-	-	-	3
Female										
2002	5	4	4	2	2	1	-	-	-	-
2001	5	3	3	-	4	-	-	1	-	-
2000	4	2	3	1	2	2	-	-	-	-
1999	3	1	2	2	1	-	1	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	316	199	152	144	136	102	63	13	1	4
2001	248	150	120	150	121	91	47	11	2	4
2000	223	135	124	116	124	106	58	14	1	3
1999	194	144	136	134	124	97	30	12	-	5
Male										
2002	176	111	74	63	62	38	24	8	-	3
2001	144	80	75	80	53	28	18	2	1	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental poisoning and exposure to noxious substances (X40-X49)														
Non-Hispanic[1]														
Male														
2000	7,944	9	18	11	13	242	560	663	849	1,328	1,610	1,297	678	230
1999	7,510	7	14	5	15	167	460	618	864	1,248	1,569	1,275	558	259
Female														
2002	5,140	4	13	3	7	106	230	287	455	731	1,025	924	535	249
2001	3,911	7	8	10	9	82	166	224	359	632	784	621	372	178
2000	3,367	3	9	3	11	70	150	178	328	550	643	508	275	171
1999	3,032	1	17	5	8	52	135	172	301	505	585	440	243	143
Non-Hispanic White														
Both sexes														
2002	13,165	7	15	8	20	395	962	1,004	1,371	1,932	2,564	2,156	1,246	527
2001	10,269	8	16	12	18	314	764	797	1,039	1,678	2,020	1,620	848	362
2000	9,124	8	15	7	17	281	635	713	970	1,540	1,791	1,390	681	304
1999	8,311	6	17	3	18	190	504	663	918	1,416	1,680	1,290	577	309
Male														
2002	8,849	7	6	6	15	299	749	750	987	1,331	1,714	1,418	807	317
2001	7,111	4	12	5	12	251	613	601	757	1,170	1,409	1,128	560	208
2000	6,413	6	9	6	10	220	516	563	709	1,112	1,287	990	474	158
1999	5,921	5	6	1	12	145	391	523	694	1,023	1,242	952	385	182
Female														
2002	4,316	-	9	2	5	96	213	254	384	601	850	738	439	210
2001	3,158	4	4	7	6	63	151	196	282	508	611	492	288	154
2000	2,711	2	6	1	7	61	119	150	261	428	504	400	207	146
1999	2,390	1	11	2	6	45	113	140	224	393	438	338	192	127
Non-Hispanic Black														
Both sexes														
2002	2,326	12	11	3	3	19	67	113	186	336	450	489	336	154
2001	2,122	6	8	4	2	26	67	103	170	316	433	431	305	101
2000	1,985	4	11	6	2	19	57	111	183	309	423	385	260	91
1999	2,008	1	12	5	2	18	75	112	216	295	434	404	209	83
Male														
2002	1,601	8	8	2	1	14	52	86	127	226	291	320	244	119
2001	1,452	3	5	1	1	14	53	81	105	205	272	314	226	78
2000	1,400	3	8	5	1	13	36	87	122	200	296	286	194	69
1999	1,450	1	7	3	2	15	61	85	146	199	304	308	163	70
Female														
2002	725	4	3	1	2	5	15	27	59	110	159	169	92	35
2001	670	3	3	3	1	12	14	22	65	111	161	117	79	23
2000	585	1	3	1	1	6	21	24	61	109	127	99	66	22

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)										
Non-Hispanic[1]										
Male										
2000	127	70	64	48	52	50	21	2	-	2
1999	128	85	70	61	53	36	10	5	-	3
Female										
2002	140	88	78	81	74	64	39	5	1	1
2001	104	70	45	70	68	63	29	9	1	-
2000	96	65	60	68	72	56	37	12	1	1
1999	66	59	66	73	71	61	20	7	-	2
Non-Hispanic White										
Both sexes										
2002	245	166	133	128	122	93	57	11	1	2
2001	184	125	101	128	101	80	41	9	-	4
2000	178	112	104	109	115	89	52	11	-	2
1999	135	116	108	124	114	82	28	10	-	3
Male										
2002	119	88	62	57	51	34	24	7	-	1
2001	96	66	63	63	46	25	17	1	-	4
2000	93	52	52	45	48	42	19	1	-	1
1999	81	67	57	56	49	34	10	4	-	2
Female										
2002	126	78	71	71	71	59	33	4	1	1
2001	88	59	38	65	55	55	24	8	-	-
2000	85	60	52	64	67	47	33	10	-	1
1999	54	49	51	68	65	48	18	6	-	1
Non-Hispanic Black										
Both sexes										
2002	67	26	15	14	12	6	5	1	-	1
2001	56	23	16	17	17	11	6	2	2	-
2000	41	23	18	7	8	16	6	3	1	1
1999	56	25	24	10	8	13	2	2	-	2
Male										
2002	55	20	11	6	9	1	-	-	-	1
2001	43	14	11	14	6	3	1	1	1	-
2000	32	18	11	3	4	8	2	1	-	1
1999	44	17	12	5	4	2	-	1	-	1
Female										
2002	12	6	4	8	3	5	5	1	-	-
2001	13	9	5	3	11	8	5	1	1	-
2000	9	5	7	4	4	8	4	2	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Accidental poisoning and exposure to noxious substances (X40-X49)														
Non-Hispanic Black														
Female														
1999	558	-	5	2	-	3	14	27	70	96	130	96	46	13
Origin not stated[2]														
Both sexes														
2002	109	-	-	-	-	1	7	6	11	17	22	16	8	6
2001	123	-	-	-	-	1	7	4	12	22	20	22	12	10
2000	138	-	1	2	1	1	5	6	12	14	25	29	23	8
1999	142	-	-	-	1	-	4	7	10	18	39	20	19	10
Male														
2002	86	-	-	-	-	-	5	5	9	14	16	13	4	6
2001	104	-	-	-	-	1	6	4	10	19	15	17	11	8
2000	104	-	1	-	1	-	5	4	9	12	19	19	18	6
1999	106	-	-	-	1	-	4	6	8	13	26	16	17	6
Female														
2002	23	-	-	-	-	1	2	1	2	3	6	3	4	-
2001	19	-	-	-	-	-	1	-	2	3	5	5	1	2
2000	34	-	-	2	-	1	-	2	3	2	6	10	5	2
1999	36	-	-	-	-	-	-	1	2	5	13	4	2	4
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
All origins														
Both sexes														
2002	17,201	677	247	113	160	273	365	377	443	577	764	745	751	722
2001	17,760	669	275	113	142	251	392	347	448	648	738	816	825	646
2000	17,437	572	266	115	152	273	375	416	479	672	774	787	723	668
1999	18,388	523	293	144	172	291	371	421	466	622	830	755	741	662
Male														
2002	9,839	389	156	71	114	225	320	311	381	468	613	609	571	534
2001	10,044	390	162	72	109	196	339	294	388	529	599	647	614	461
2000	9,960	327	162	72	120	223	337	362	405	577	628	646	573	504
1999	10,371	314	183	93	128	238	327	361	387	513	668	584	592	490
Female														
2002	7,362	288	91	42	46	48	45	66	62	109	151	136	180	188
2001	7,716	279	113	41	33	55	53	53	60	119	139	169	211	185
2000	7,477	245	104	43	32	50	38	54	74	95	146	141	150	164
1999	8,017	209	110	51	44	53	44	60	79	109	162	171	149	172

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Accidental poisoning and exposure to noxious substances (X40-X49)										
Non-Hispanic Black										
Female										
1999	12	8	12	5	4	11	2	1	-	1
Origin not stated[2]										
Both sexes										
2002	2	4	2	2	-	-	-	-	-	5
2001	5	1	1	1	-	-	-	-	-	5
2000	3	2	2	-	-	-	-	-	-	4
1999	4	2	-	-	2	2	-	-	-	4
Male										
2002	2	4	2	2	-	-	-	-	-	4
2001	5	1	1	1	-	-	-	-	-	5
2000	3	2	1	-	-	-	-	-	-	4
1999	4	2	-	-	-	1	-	-	-	2
Female										
2002	-	-	-	-	-	-	-	-	-	1
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	1	-	-	-	-	-	-	-
1999	-	-	-	-	2	1	-	-	-	2
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)										
All origins										
Both sexes										
2002	663	826	969	1,601	2,064	2,254	1,720	727	150	13
2001	696	744	1,082	1,669	2,172	2,404	1,784	704	171	24
2000	681	743	1,081	1,632	2,046	2,285	1,798	706	177	16
1999	670	852	1,180	1,735	2,290	2,563	1,895	760	137	15
Male										
2002	479	534	579	879	948	878	545	201	24	10
2001	472	480	617	897	1,016	950	610	155	26	21
2000	467	474	607	855	946	883	564	179	33	16
1999	461	526	701	900	1,069	1,003	598	194	27	14
Female										
2002	184	292	390	722	1,116	1,376	1,175	526	126	3
2001	224	264	465	772	1,156	1,454	1,174	549	145	3
2000	214	269	474	777	1,100	1,402	1,234	527	144	-
1999	209	326	479	835	1,221	1,560	1,297	566	110	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
Hispanic														
Both sexes														
2002	954	68	42	17	19	37	63	67	72	69	75	49	52	34
2001	985	78	43	9	14	34	64	55	67	77	69	61	61	35
2000	942	58	41	13	16	34	74	63	77	77	74	49	35	31
1999	878	65	53	18	13	35	63	54	47	68	52	49	33	27
Male														
2002	726	43	28	11	10	34	58	64	69	62	65	45	42	29
2001	704	40	28	2	10	31	62	50	60	70	60	51	51	27
2000	695	31	23	6	8	31	74	58	71	73	66	43	27	24
1999	620	35	37	12	7	28	60	51	39	60	41	44	26	23
Female														
2002	228	25	14	6	9	3	5	3	3	7	10	4	10	5
2001	281	38	15	7	4	3	2	5	7	7	9	10	10	8
2000	247	27	18	7	8	3	-	5	6	4	8	6	8	7
1999	258	30	16	6	6	7	3	3	8	8	11	5	7	4
Non-Hispanic[1]														
Both sexes														
2002	16,183	606	205	95	140	233	301	310	367	505	688	692	691	682
2001	16,695	588	231	104	126	217	325	290	379	566	665	751	762	604
2000	16,408	513	224	102	136	238	300	351	397	589	694	730	683	636
1999	17,430	457	239	126	158	254	308	364	417	553	778	700	700	626
Male														
2002	9,070	344	128	59	104	189	261	247	308	403	547	562	522	499
2001	9,280	347	133	70	97	165	274	242	326	454	536	593	561	428
2000	9,207	295	139	66	112	192	262	302	329	499	558	595	543	479
1999	9,694	279	146	81	120	209	267	308	347	452	627	535	559	459
Female														
2002	7,113	262	77	36	36	44	40	63	59	102	141	130	169	183
2001	7,415	241	98	34	29	52	51	48	53	112	129	158	201	176
2000	7,201	218	85	36	24	46	38	49	68	90	136	135	140	157
1999	7,736	178	93	45	38	45	41	56	70	101	151	165	141	167
Non-Hispanic White														
Both sexes														
2002	13,865	339	136	66	106	190	256	263	302	402	567	564	567	556
2001	14,326	348	160	85	97	183	272	228	300	468	526	594	622	504
2000	14,083	310	156	76	109	186	239	283	321	479	560	568	555	516
1999	14,930	276	163	96	129	213	255	293	329	468	635	533	567	491

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)										
Hispanic										
Both sexes										
2002	33	42	31	56	36	34	32	15	6	5
2001	35	37	51	45	44	48	37	15	2	4
2000	37	32	35	44	40	45	40	17	4	6
1999	42	41	49	35	37	45	30	15	4	3
Male										
2002	23	27	17	33	19	15	17	7	3	5
2001	22	24	30	26	20	16	15	4	1	4
2000	29	21	20	22	18	20	17	6	1	6
1999	25	27	33	14	16	18	13	8	-	3
Female										
2002	10	15	14	23	17	19	15	8	3	-
2001	13	13	21	19	24	32	22	11	1	-
2000	8	11	15	22	22	25	23	11	3	-
1999	17	14	16	21	21	27	17	7	4	-
Non-Hispanic[1]										
Both sexes										
2002	629	782	935	1,542	2,024	2,215	1,681	710	144	6
2001	658	705	1,027	1,619	2,122	2,352	1,743	686	169	6
2000	639	708	1,040	1,580	2,000	2,232	1,754	688	172	2
1999	624	807	1,128	1,695	2,248	2,509	1,859	744	133	3
Male										
2002	455	506	559	845	926	860	527	193	21	5
2001	448	456	584	866	992	933	593	151	25	6
2000	434	450	583	829	926	861	546	173	32	2
1999	432	497	665	884	1,049	982	581	185	27	3
Female										
2002	174	276	376	697	1,098	1,355	1,154	517	123	1
2001	210	249	443	753	1,130	1,419	1,150	535	144	-
2000	205	258	457	751	1,074	1,371	1,208	515	140	-
1999	192	310	463	811	1,199	1,527	1,278	559	106	-
Non-Hispanic White										
Both sexes										
2002	525	608	806	1,354	1,836	2,058	1,571	658	129	6
2001	529	594	868	1,431	1,928	2,182	1,615	633	153	6
2000	528	579	885	1,391	1,833	2,080	1,638	637	154	-
1999	489	666	946	1,470	2,034	2,310	1,748	697	121	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
Non-Hispanic White														
Male														
2002	7,669	203	83	44	76	158	224	212	254	318	468	473	425	401
2001	7,832	206	93	54	75	139	231	193	256	386	428	477	452	352
2000	7,703	186	99	51	90	153	208	243	272	409	451	458	441	384
1999	8,122	167	96	67	96	174	221	255	277	388	508	405	455	358
Female														
2002	6,196	136	53	22	30	32	32	51	48	84	99	91	142	155
2001	6,494	142	67	31	22	44	41	35	44	82	98	117	170	152
2000	6,380	124	57	25	19	33	31	40	49	70	109	110	114	132
1999	6,808	109	67	29	33	39	34	38	52	80	127	128	112	133
Non-Hispanic Black														
Both sexes														
2002	1,951	250	59	23	32	33	35	38	52	85	94	109	105	106
2001	1,968	217	61	13	25	24	38	48	53	74	110	133	116	84
2000	1,968	183	55	24	24	41	49	54	64	90	108	132	105	106
1999	2,183	168	61	23	24	31	43	59	73	65	128	149	114	121
Male														
2002	1,162	132	40	11	27	24	28	28	43	69	57	75	82	82
2001	1,180	131	35	12	19	18	29	35	46	50	87	98	87	63
2000	1,256	96	32	14	20	29	44	45	49	74	85	110	86	84
1999	1,365	101	42	12	20	28	39	44	58	48	105	116	87	91
Female														
2002	789	118	19	12	5	9	7	10	9	16	37	34	23	24
2001	788	86	26	1	6	6	9	13	7	24	23	35	29	21
2000	712	87	23	10	4	12	5	9	15	16	23	22	19	22
1999	818	67	19	11	4	3	4	15	15	17	23	33	27	30
Origin not stated[2]														
Both sexes														
2002	64	3	-	1	1	3	1	-	4	3	1	4	8	6
2001	80	3	1	-	2	-	3	2	2	5	4	4	2	7
2000	87	1	1	-	-	1	1	2	5	6	6	8	5	1
1999	80	1	1	-	1	2	-	3	2	1	-	6	8	9
Male														
2002	43	2	-	1	-	2	1	-	4	3	1	2	7	6
2001	60	3	1	-	2	-	3	2	2	5	3	3	2	6
2000	58	1	-	-	-	-	1	2	5	5	4	8	3	1
1999	57	-	-	-	1	1	-	2	1	1	-	5	7	8

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)										
Non-Hispanic White										
Male										
2002	385	392	490	742	838	797	487	179	15	5
2001	367	387	499	772	886	865	544	140	24	6
2000	352	366	502	719	836	794	507	155	27	-
1999	347	406	556	765	952	895	533	177	23	1
Female										
2002	140	216	316	612	998	1,261	1,084	479	114	1
2001	162	207	369	659	1,042	1,317	1,071	493	129	-
2000	176	213	383	672	997	1,286	1,131	482	127	-
1999	142	260	390	705	1,082	1,415	1,215	520	98	-
Non-Hispanic Black										
Both sexes										
2002	83	150	107	162	149	137	90	42	10	-
2001	114	95	137	155	162	143	109	42	15	-
2000	94	111	133	170	137	131	94	44	17	2
1999	115	125	168	200	184	174	100	46	10	2
Male										
2002	56	95	56	89	70	53	31	11	3	-
2001	71	61	73	75	90	53	39	7	1	-
2000	67	70	69	99	75	55	31	15	5	2
1999	74	80	102	106	79	79	41	8	3	2
Female										
2002	27	55	51	73	79	84	59	31	7	-
2001	43	34	64	80	72	90	70	35	14	-
2000	27	41	64	71	62	76	63	29	12	-
1999	41	45	66	94	105	95	59	38	7	-
Origin not stated[2]										
Both sexes										
2002	1	2	3	3	4	5	7	2	-	2
2001	3	2	4	5	6	4	4	3	-	14
2000	5	3	6	8	6	8	4	1	1	8
1999	4	4	3	5	5	9	6	1	-	9
Male										
2002	1	1	3	1	3	3	1	1	-	-
2001	2	-	3	5	4	1	2	-	-	11
2000	4	3	4	4	2	2	1	-	-	8
1999	4	2	3	2	4	3	4	1	-	8

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)														
Origin not stated[2]														
Female														
2002	21	1	-	-	1	1	-	-	-	-	-	2	1	-
2001	20	-	-	-	-	-	-	-	-	-	1	1	-	1
2000	29	-	1	-	-	1	-	-	-	1	2	-	2	-
1999	23	1	1	-	-	1	-	1	1	-	-	1	1	1
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)														
All origins														
Both sexes														
2002	31,655	. . .	. . .	4	260	1,513	2,497	2,423	2,623	3,141	3,710	3,473	2,835	2,186
2001[3]	30,622	. . .	. . .	7	272	1,611	2,360	2,389	2,681	3,176	3,459	3,260	2,682	1,985
2000	29,350	. . .	. . .	7	300	1,621	2,373	2,341	2,451	3,112	3,450	2,987	2,450	1,708
1999	29,199	. . .	. . .	2	242	1,615	2,286	2,398	2,708	3,286	3,180	2,817	2,264	1,678
Male														
2002	25,409	. . .	. . .	3	196	1,280	2,152	2,003	2,132	2,484	2,821	2,608	2,188	1,708
2001[3]	24,672	. . .	. . .	7	207	1,345	2,064	2,015	2,184	2,518	2,662	2,484	2,020	1,534
2000	23,618	. . .	. . .	6	238	1,351	2,073	1,956	1,982	2,457	2,657	2,307	1,842	1,306
1999	23,458	. . .	. . .	2	192	1,347	1,979	2,005	2,189	2,590	2,429	2,152	1,702	1,284
Female														
2002	6,246	. . .	. . .	1	64	233	345	420	491	657	889	865	647	478
2001[3]	5,950	. . .	. . .	-	65	266	296	374	497	658	797	776	662	451
2000	5,732	. . .	. . .	1	62	270	300	385	469	655	793	680	608	402
1999	5,741	. . .	. . .	-	50	268	307	393	519	696	751	665	562	394
Hispanic														
Both sexes														
2002	1,954	. . .	. . .	2	35	181	274	248	236	222	185	149	111	90
2001[3]	1,850	. . .	. . .	-	37	166	248	259	231	224	173	130	109	65
2000	1,787	. . .	. . .	2	27	178	269	227	193	194	191	141	103	62
1999	1,695	. . .	. . .	-	30	175	216	220	203	228	141	121	89	65
Male														
2002	1,651	. . .	. . .	1	20	151	237	222	204	188	152	115	91	79
2001[3]	1,576	. . .	. . .	-	26	129	214	223	196	200	150	106	86	56
2000	1,525	. . .	. . .	2	20	143	244	203	167	158	162	112	82	54
1999	1,430	. . .	. . .	-	21	146	192	187	177	188	119	104	65	55
Female														
2002	303	. . .	. . .	1	15	30	37	26	32	34	33	34	20	11
2001[3]	274	. . .	. . .	-	11	37	34	36	35	24	23	24	23	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99, X10-X39,X50-X59,Y86)										
Origin not stated[2]										
Female										
2002	-	1	-	2	1	2	6	1	-	2
2001	1	2	1	-	2	3	2	3	-	3
2000	1	-	2	4	4	6	3	1	1	-
1999	-	2	-	3	1	6	2	-	-	1
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All origins										
Both sexes										
2002	1,432	1,197	1,266	1,231	1,028	573	211	41	1	10
2001[3]	1,332	1,212	1,220	1,219	973	538	200	28	3	15
2000	1,237	1,073	1,219	1,227	954	590	205	34	4	7
1999	1,218	1,187	1,284	1,280	937	580	182	39	-	16
Male										
2002	1,129	975	1,078	1,054	884	492	179	33	-	10
2001[3]	1,029	1,001	1,039	1,033	852	468	168	25	3	14
2000	959	870	1,018	1,062	822	507	168	26	4	7
1999	971	972	1,079	1,073	795	501	150	30	-	16
Female										
2002	303	222	188	177	144	81	32	8	1	-
2001[3]	303	211	181	186	121	70	32	3	-	1
2000	278	203	201	165	132	83	37	8	-	-
1999	247	215	205	207	142	79	32	9	-	-
Hispanic										
Both sexes										
2002	52	45	44	30	25	17	4	2	-	2
2001[3]	49	42	44	24	25	14	6	2	-	2
2000	40	43	41	32	27	11	3	2	-	1
1999	47	46	27	35	28	12	7	3	-	2
Male										
2002	44	39	38	25	22	16	4	1	-	2
2001[3]	44	39	38	23	22	14	6	2	-	2
2000	35	39	35	27	26	10	3	2	-	1
1999	39	41	22	30	24	10	6	2	-	2
Female										
2002	8	6	6	5	3	1	-	1	-	-
2001[3]	5	3	6	1	3	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)														
Hispanic														
Female														
2000	262	. . .	. . .	-	7	35	25	24	26	36	29	29	21	8
1999	265	. . .	. . .	-	9	29	24	33	26	40	22	17	24	10
Non-Hispanic[1]														
Both sexes														
2002	29,543	. . .	. . .	2	224	1,322	2,214	2,165	2,373	2,902	3,511	3,309	2,706	2,086
2001[3]	28,647	. . .	. . .	7	235	1,440	2,105	2,119	2,441	2,943	3,270	3,118	2,564	1,909
2000	27,426	. . .	. . .	5	271	1,436	2,093	2,101	2,249	2,904	3,241	2,829	2,336	1,638
1999	27,379	. . .	. . .	1	211	1,434	2,066	2,167	2,498	3,048	3,020	2,683	2,159	1,608
Male														
2002	23,623	. . .	. . .	2	175	1,120	1,906	1,772	1,917	2,282	2,659	2,480	2,083	1,621
2001[3]	22,983	. . .	. . .	7	181	1,212	1,843	1,782	1,980	2,311	2,498	2,367	1,925	1,468
2000	21,981	. . .	. . .	4	217	1,203	1,820	1,744	1,809	2,288	2,479	2,180	1,751	1,244
1999	21,929	. . .	. . .	1	170	1,198	1,785	1,809	2,007	2,395	2,294	2,038	1,625	1,225
Female														
2002	5,920	. . .	. . .	-	49	202	308	393	456	620	852	829	623	465
2001[3]	5,664	. . .	. . .	-	54	228	262	337	461	632	772	751	639	441
2000	5,445	. . .	. . .	1	54	233	273	357	440	616	762	649	585	394
1999	5,450	. . .	. . .	-	41	236	281	358	491	653	726	645	534	383
Non-Hispanic White														
Both sexes														
2002	26,691	. . .	. . .	1	176	1,124	1,827	1,771	2,038	2,570	3,225	3,084	2,519	1,965
2001[3]	25,813	. . .	. . .	5	178	1,217	1,709	1,755	2,116	2,621	2,983	2,897	2,395	1,787
2000	24,616	. . .	. . .	5	220	1,175	1,693	1,709	1,949	2,599	2,941	2,611	2,163	1,536
1999	24,534	. . .	. . .	1	173	1,169	1,661	1,776	2,151	2,699	2,770	2,468	2,019	1,501
Male														
2002	21,323	. . .	. . .	1	137	954	1,573	1,427	1,638	2,013	2,449	2,296	1,956	1,524
2001[3]	20,710	. . .	. . .	5	139	1,030	1,492	1,479	1,711	2,046	2,283	2,215	1,801	1,375
2000	19,711	. . .	. . .	4	175	987	1,480	1,424	1,565	2,038	2,249	2,008	1,619	1,165
1999	19,620	. . .	. . .	1	140	978	1,438	1,478	1,727	2,098	2,112	1,876	1,512	1,141
Female														
2002	5,368	. . .	. . .	-	39	170	254	344	400	557	776	788	563	441
2001[3]	5,103	. . .	. . .	-	39	187	217	276	405	575	700	682	594	412
2000	4,905	. . .	. . .	1	45	188	213	285	384	561	692	603	544	371
1999	4,914	. . .	. . .	-	33	191	223	298	424	601	658	592	507	360
Non-Hispanic Black														
Both sexes														
2002	1,896	. . .	. . .	1	34	122	270	271	226	229	203	149	116	68
2001[3]	1,904	. . .	. . .	2	38	134	291	243	229	227	191	152	116	78

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
Hispanic										
Female										
2000	5	4	6	5	1	1	-	-	-	-
1999	8	5	5	5	4	2	1	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,369	1,141	1,219	1,198	1,001	554	205	37	1	4
2001[3]	1,275	1,162	1,172	1,193	945	521	194	26	3	5
2000	1,189	1,025	1,173	1,194	925	577	201	31	4	4
1999	1,165	1,133	1,255	1,240	906	565	175	36	-	9
Male										
2002	1,075	926	1,037	1,026	861	474	173	30	-	4
2001[3]	978	955	997	1,008	827	451	162	23	3	5
2000	917	826	979	1,034	794	496	164	24	4	4
1999	927	923	1,055	1,039	768	489	144	28	-	9
Female										
2002	294	215	182	172	140	80	32	7	1	-
2001[3]	297	207	175	185	118	70	32	3	-	-
2000	272	199	194	160	131	81	37	7	-	-
1999	238	210	200	201	138	76	31	8	-	-
Non-Hispanic White										
Both sexes										
2002	1,289	1,074	1,154	1,131	974	533	197	34	1	4
2001[3]	1,192	1,085	1,102	1,144	905	501	189	25	3	4
2000	1,123	969	1,106	1,141	887	561	194	29	2	3
1999	1,079	1,050	1,199	1,186	872	548	171	35	-	6
Male										
2002	1,022	870	995	974	838	458	167	27	-	4
2001[3]	909	899	937	966	797	438	158	23	3	4
2000	866	779	922	992	769	484	158	22	2	3
1999	860	862	1,007	997	743	475	142	27	-	6
Female										
2002	267	204	159	157	136	75	30	7	1	-
2001[3]	283	186	165	178	108	63	31	2	-	-
2000	257	190	184	149	118	77	36	7	-	-
1999	219	188	192	189	129	73	29	8	-	-
Non-Hispanic Black										
Both sexes										
2002	57	33	47	38	13	15	2	2	-	-
2001[3]	48	42	48	29	23	8	4	1	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)														
Non-Hispanic Black														
Both sexes														
2000	1,923	. . .	. . .	-	39	170	301	262	207	205	222	150	116	60
1999	1,924	. . .	. . .	-	29	176	287	258	239	244	186	144	82	67
Male														
2002	1,595	. . .	. . .	1	28	107	235	244	197	191	154	129	83	57
2001[3]	1,578	. . .	. . .	2	29	115	273	212	194	188	141	109	87	62
2000	1,601	. . .	. . .	-	34	148	259	222	175	172	174	123	87	49
1999	1,630	. . .	. . .	-	22	152	256	229	203	215	138	114	69	54
Female														
2002	301	. . .	. . .	-	6	15	35	27	29	38	49	20	33	11
2001[3]	326	. . .	. . .	-	9	19	18	31	35	39	50	43	29	16
2000	322	. . .	. . .	-	5	22	42	40	32	33	48	27	29	11
1999	294	. . .	. . .	-	7	24	31	29	36	29	48	30	13	13
Origin not stated[2]														
Both sexes														
2002	158	. . .	. . .	-	1	10	9	10	14	17	14	15	18	10
2001[3]	125	. . .	. . .	-	-	5	7	11	9	9	16	12	9	11
2000	137	. . .	. . .	-	2	7	11	13	9	14	18	17	11	8
1999	125	. . .	. . .	1	1	6	4	11	7	10	19	13	16	5
Male														
2002	135	. . .	. . .	-	1	9	9	9	11	14	10	13	14	8
2001[3]	113	. . .	. . .	-	-	4	7	10	8	7	14	11	9	10
2000	112	. . .	. . .	-	1	5	9	9	6	11	16	15	9	8
1999	99	. . .	. . .	1	1	3	2	9	5	7	16	10	12	4
Female														
2002	23	. . .	. . .	-	-	1	-	1	3	3	4	2	4	2
2001[3]	12	. . .	. . .	-	-	1	-	1	1	2	2	1	-	1
2000	25	. . .	. . .	-	1	2	2	4	3	3	2	2	2	-
1999	26	. . .	. . .	-	-	3	2	2	2	3	3	3	4	1
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
All origins														
Both sexes														
2002	17,108	. . .	. . .	-	86	742	1,346	1,217	1,182	1,450	1,707	1,652	1,484	1,295
2001	16,869	. . .	. . .	-	90	838	1,292	1,263	1,301	1,429	1,601	1,552	1,471	1,195
2000	16,586	. . .	. . .	-	110	897	1,370	1,223	1,230	1,521	1,670	1,491	1,367	1,001
1999	16,599	. . .	. . .	-	103	975	1,340	1,292	1,329	1,617	1,480	1,462	1,215	1,058
Male														
2002	15,045	. . .	. . .	-	68	668	1,229	1,068	1,010	1,234	1,438	1,394	1,283	1,115

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
Non-Hispanic Black										
Both sexes										
2000	41	39	46	30	23	9	2	-	1	-
1999	49	58	36	35	17	10	4	1	-	2
Male										
2002	40	31	35	33	13	13	2	2	-	-
2001[3]	40	31	42	25	19	5	4	-	-	-
2000	31	36	40	25	15	8	2	-	1	-
1999	41	47	31	30	15	9	2	1	-	2
Female										
2002	17	2	12	5	-	2	-	-	-	-
2001[3]	8	11	6	4	4	3	-	1	-	-
2000	10	3	6	5	8	1	-	-	-	-
1999	8	11	5	5	2	1	2	-	-	-
Origin not stated[2]										
Both sexes										
2002	11	11	3	3	2	2	2	2	-	4
2001[3]	8	8	4	2	3	3	-	-	-	8
2000	8	5	5	1	2	2	1	1	-	2
1999	6	8	2	5	3	3	-	-	-	5
Male										
2002	10	10	3	3	1	2	2	2	-	4
2001[3]	7	7	4	2	3	3	-	-	-	7
2000	7	5	4	1	2	1	1	-	-	2
1999	5	8	2	4	3	2	-	-	-	5
Female										
2002	1	1	-	-	1	-	-	-	-	-
2001[3]	1	1	-	-	-	-	-	-	-	1
2000	1	-	1	-	-	1	-	1	-	-
1999	1	-	-	1	-	1	-	-	-	-
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All origins										
Both sexes										
2002	939	846	930	918	762	392	133	25	-	2
2001	888	857	901	920	728	382	135	18	2	6
2000	832	791	909	917	694	410	129	18	1	5
1999	802	827	964	946	660	399	102	23	-	5
Male										
2002	815	753	867	854	722	374	127	24	-	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
All origins														
Male														
2001	14,758	. . .	. . .	-	69	743	1,192	1,122	1,124	1,200	1,351	1,309	1,242	1,020
2000	14,454	. . .	. . .	-	90	793	1,250	1,084	1,039	1,255	1,402	1,263	1,146	854
1999	14,479	. . .	. . .	-	80	867	1,216	1,137	1,133	1,349	1,237	1,221	1,030	882
Female														
2002	2,063	. . .	. . .	-	18	74	117	149	172	216	269	258	201	180
2001	2,111	. . .	. . .	-	21	95	100	141	177	229	250	243	229	175
2000	2,132	. . .	. . .	-	20	104	120	139	191	266	268	228	221	147
1999	2,120	. . .	. . .	-	23	108	124	155	196	268	243	241	185	176
Hispanic														
Both sexes														
2002	834	. . .	. . .	-	10	88	136	112	86	80	65	59	39	40
2001	798	. . .	. . .	-	5	67	126	129	95	75	60	54	48	32
2000	813	. . .	. . .	-	7	88	136	104	77	65	79	72	52	28
1999	794	. . .	. . .	-	13	94	125	104	78	91	57	64	32	29
Male														
2002	763	. . .	. . .	-	5	81	127	101	82	72	56	51	37	37
2001	731	. . .	. . .	-	4	58	114	116	87	70	56	48	42	32
2000	737	. . .	. . .	-	7	78	127	96	69	55	71	61	45	27
1999	705	. . .	. . .	-	7	81	115	92	72	76	51	55	26	26
Female														
2002	71	. . .	. . .	-	5	7	9	11	4	8	9	8	2	3
2001	67	. . .	. . .	-	1	9	12	13	8	5	4	6	6	-
2000	76	. . .	. . .	-	-	10	9	8	8	10	8	11	7	1
1999	89	. . .	. . .	-	6	13	10	12	6	15	6	9	6	3
Non-Hispanic[1]														
Both sexes														
2002	16,213	. . .	. . .	-	76	649	1,206	1,102	1,093	1,363	1,639	1,585	1,439	1,251
2001	16,020	. . .	. . .	-	85	770	1,163	1,130	1,205	1,351	1,536	1,495	1,421	1,157
2000	15,717	. . .	. . .	-	103	805	1,230	1,112	1,152	1,452	1,584	1,412	1,310	971
1999	15,755	. . .	. . .	-	90	879	1,213	1,184	1,247	1,520	1,417	1,395	1,178	1,028
Male														
2002	14,226	. . .	. . .	-	63	582	1,098	964	926	1,155	1,381	1,336	1,240	1,075
2001	13,978	. . .	. . .	-	65	684	1,075	1,002	1,037	1,127	1,290	1,259	1,198	982
2000	13,664	. . .	. . .	-	83	712	1,119	982	969	1,196	1,324	1,195	1,096	825
1999	13,732	. . .	. . .	-	73	784	1,100	1,043	1,058	1,268	1,180	1,164	1,000	855
Female														
2002	1,987	. . .	. . .	-	13	67	108	138	167	208	258	249	199	176
2001	2,042	. . .	. . .	-	20	86	88	128	168	224	246	236	223	175

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All origins										
Male										
2001	751	761	831	846	685	360	126	18	2	6
2000	714	698	830	856	649	388	119	18	1	5
1999	712	742	873	875	624	376	99	21	-	5
Female										
2002	124	93	63	64	40	18	6	1	-	-
2001	137	96	70	74	43	22	9	-	-	-
2000	118	93	79	61	45	22	10	-	-	-
1999	90	85	91	71	36	23	3	2	-	-
Hispanic										
Both sexes										
2002	27	26	26	14	14	9	2	1	-	-
2001	27	24	21	12	14	4	3	1	-	1
2000	19	24	25	16	11	7	1	1	-	1
1999	19	24	13	24	15	5	5	1	-	1
Male										
2002	25	24	25	14	14	9	2	1	-	-
2001	25	23	21	12	14	4	3	1	-	1
2000	19	23	23	15	11	7	1	1	-	1
1999	19	22	13	24	14	5	5	1	-	1
Female										
2002	2	2	1	-	-	-	-	-	-	-
2001	2	1	-	-	-	-	-	-	-	-
2000	-	1	2	1	-	-	-	-	-	-
1999	-	2	-	-	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	908	813	903	902	747	382	130	24	-	1
2001	854	829	877	906	711	375	132	17	2	4
2000	808	764	881	900	683	402	127	17	1	3
1999	778	800	949	918	645	392	97	22	-	3
Male										
2002	786	722	841	838	707	364	124	23	-	1
2001	719	734	807	832	668	353	123	17	2	4
2000	690	672	804	840	638	381	117	17	1	3
1999	688	717	858	847	610	370	94	20	-	3
Female										
2002	122	91	62	64	40	18	6	1	-	-
2001	135	95	70	74	43	22	9	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
Non-Hispanic[1]														
Female														
2000	2,053	. . .	. . .	-	20	93	111	130	183	256	260	217	214	146
1999	2,023	. . .	. . .	-	17	95	113	141	189	252	237	231	178	173
Non-Hispanic White														
Both sexes														
2002	14,865	. . .	. . .	-	63	540	981	898	955	1,219	1,530	1,493	1,353	1,195
2001	14,648	. . .	. . .	-	69	650	915	942	1,059	1,220	1,409	1,413	1,347	1,101
2000	14,333	. . .	. . .	-	96	651	1,004	904	1,012	1,316	1,464	1,315	1,237	912
1999	14,316	. . .	. . .	-	81	726	958	977	1,094	1,357	1,319	1,295	1,118	970
Male														
2002	13,014	. . .	. . .	-	53	485	890	767	806	1,028	1,291	1,255	1,168	1,023
2001	12,771	. . .	. . .	-	55	573	841	833	906	1,011	1,188	1,196	1,137	933
2000	12,448	. . .	. . .	-	79	574	915	793	855	1,075	1,221	1,114	1,037	772
1999	12,441	. . .	. . .	-	65	645	869	854	922	1,120	1,093	1,084	945	802
Female														
2002	1,851	. . .	. . .	-	10	55	91	131	149	191	239	238	185	172
2001	1,877	. . .	. . .	-	14	77	74	109	153	209	221	217	210	168
2000	1,885	. . .	. . .	-	17	77	89	111	157	241	243	201	200	140
1999	1,875	. . .	. . .	-	16	81	89	123	172	237	226	211	173	168
Non-Hispanic Black														
Both sexes														
2002	1,041	. . .	. . .	-	9	77	173	162	110	114	85	73	57	39
2001	1,069	. . .	. . .	-	12	88	205	147	111	107	91	62	60	41
2000	1,073	. . .	. . .	-	4	117	187	159	109	100	93	73	58	38
1999	1,112	. . .	. . .	-	5	118	205	153	115	123	74	76	42	46
Male														
2002	945	. . .	. . .	-	8	68	161	157	101	100	70	67	48	36
2001	949	. . .	. . .	-	9	80	197	131	102	96	73	48	50	37
2000	949	. . .	. . .	-	4	109	168	144	93	88	80	62	45	34
1999	1,007	. . .	. . .	-	4	107	187	142	105	112	66	63	39	42
Female														
2002	96	. . .	. . .	-	1	9	12	5	9	14	15	6	9	3
2001	120	. . .	. . .	-	3	8	8	16	9	11	18	14	10	4
2000	124	. . .	. . .	-	-	8	19	15	16	12	13	11	13	4
1999	105	. . .	. . .	-	1	11	18	11	10	11	8	13	3	4
Origin not stated[2]														
Both sexes														
2002	61	. . .	. . .	-	-	5	4	3	3	7	3	8	6	4
2001	51	. . .	. . .	-	-	1	3	4	1	3	5	3	2	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
Non-Hispanic[1]										
Female										
2000	118	92	77	60	45	21	10	-	-	-
1999	90	83	91	71	35	22	3	2	-	-
Non-Hispanic White										
Both sexes										
2002	868	784	869	862	733	371	128	22	-	1
2001	818	792	836	876	684	367	128	17	2	3
2000	767	732	840	877	668	395	124	16	1	2
1999	733	758	914	889	625	383	95	21	-	3
Male										
2002	751	695	812	800	693	353	122	21	-	1
2001	686	704	770	805	646	346	119	17	2	3
2000	655	641	764	820	626	374	114	16	1	2
1999	649	682	823	821	591	362	92	19	-	3
Female										
2002	117	89	57	62	40	18	6	1	-	-
2001	132	88	66	71	38	21	9	-	-	-
2000	112	91	76	57	42	21	10	-	-	-
1999	84	76	91	68	34	21	3	2	-	-
Non-Hispanic Black										
Both sexes										
2002	32	22	32	31	10	11	2	2	-	-
2001	26	27	35	25	22	6	4	-	-	-
2000	33	28	34	19	13	6	2	-	-	-
1999	35	36	29	28	15	9	2	1	-	-
Male										
2002	27	20	28	29	10	11	2	2	-	-
2001	24	22	31	22	18	5	4	-	-	-
2000	27	27	33	17	10	6	2	-	-	-
1999	31	30	29	25	14	8	2	1	-	-
Female										
2002	5	2	4	2	-	-	-	-	-	-
2001	2	5	4	3	4	1	-	-	-	-
2000	6	1	1	2	3	-	-	-	-	-
1999	4	6	-	3	1	1	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	4	7	1	2	1	1	1	-	-	1
2001	7	4	3	2	3	3	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)														
Origin not stated[2]														
Both sexes														
2000	56	. . .	. . .	-	-	4	4	7	1	4	7	7	5	2
1999	50	. . .	. . .	-	-	2	2	4	4	6	6	3	5	1
Male														
2002	56	. . .	. . .	-	-	5	4	3	2	7	1	7	6	3
2001	49	. . .	. . .	-	-	1	3	4	-	3	5	2	2	6
2000	53	. . .	. . .	-	-	3	4	6	1	4	7	7	5	2
1999	42	. . .	. . .	-	-	2	1	2	3	5	6	2	4	1
Female														
2002	5	. . .	. . .	-	-	-	-	-	1	-	2	1	-	1
2001	2	. . .	. . .	-	-	-	-	-	1	-	-	1	-	-
2000	3	. . .	. . .	-	-	1	-	1	-	-	-	-	-	-
1999	8	. . .	. . .	-	-	-	1	2	1	1	-	1	1	-
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)														
All origins														
Both sexes														
2002	14,547	. . .	. . .	4	174	771	1,151	1,206	1,441	1,691	2,003	1,821	1,351	891
2001[3]	13,753	. . .	. . .	7	182	773	1,068	1,126	1,380	1,747	1,858	1,708	1,211	790
2000	12,764	. . .	. . .	7	190	724	1,003	1,118	1,221	1,591	1,780	1,496	1,083	707
1999	12,600	. . .	. . .	2	139	640	946	1,106	1,379	1,669	1,700	1,355	1,049	620
Male														
2002	10,364	. . .	. . .	3	128	612	923	935	1,122	1,250	1,383	1,214	905	593
2001[3]	9,914	. . .	. . .	7	138	602	872	893	1,060	1,318	1,311	1,175	778	514
2000	9,164	. . .	. . .	6	148	558	823	872	943	1,202	1,255	1,044	696	452
1999	8,979	. . .	. . .	2	112	480	763	868	1,056	1,241	1,192	931	672	402
Female														
2002	4,183	. . .	. . .	1	46	159	228	271	319	441	620	607	446	298
2001[3]	3,839	. . .	. . .	-	44	171	196	233	320	429	547	533	433	276
2000	3,600	. . .	. . .	1	42	166	180	246	278	389	525	452	387	255
1999	3,621	. . .	. . .	-	27	160	183	238	323	428	508	424	377	218
Hispanic														
Both sexes														
2002	1,120	. . .	. . .	2	25	93	138	136	150	142	120	90	72	50
2001[3]	1,052	. . .	. . .	-	32	99	122	130	136	149	113	76	61	33
2000	974	. . .	. . .	2	20	90	133	123	116	129	112	69	51	34
1999	901	. . .	. . .	-	17	81	91	116	125	137	84	57	57	36

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
Origin not stated[2]										
Both sexes										
2000	5	3	3	1	-	1	1	-	-	1
1999	5	3	2	4	-	2	-	-	-	1
Male										
2002	4	7	1	2	1	1	1	-	-	1
2001	7	4	3	2	3	3	-	-	-	1
2000	5	3	3	1	-	-	1	-	-	1
1999	5	3	2	4	-	1	-	-	-	1
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	1	-	-	-	-
1999	-	-	-	-	-	1	-	-	-	-
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
All origins										
Both sexes										
2002	493	351	336	313	266	181	78	16	1	8
2001[3]	444	355	319	299	245	156	65	10	1	9
2000	405	282	310	310	260	180	76	16	3	2
1999	416	360	320	334	277	181	80	16	-	11
Male										
2002	314	222	211	200	162	118	52	9	-	8
2001[3]	278	240	208	187	167	108	42	7	1	8
2000	245	172	188	206	173	119	49	8	3	2
1999	259	230	206	198	171	125	51	9	-	11
Female										
2002	179	129	125	113	104	63	26	7	1	-
2001[3]	166	115	111	112	78	48	23	3	-	1
2000	160	110	122	104	87	61	27	8	-	-
1999	157	130	114	136	106	56	29	7	-	-
Hispanic										
Both sexes										
2002	25	19	18	16	11	8	2	1	-	2
2001[3]	22	18	23	12	11	10	3	1	-	1
2000	21	19	16	16	16	4	2	1	-	-
1999	28	22	14	11	13	7	2	2	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)														
Hispanic														
Male														
2002	888	. . .	. . .	1	15	70	110	121	122	116	96	64	54	42
2001[3]	845	. . .	. . .	-	22	71	100	107	109	130	94	58	44	24
2000	788	. . .	. . .	2	13	65	117	107	98	103	91	51	37	27
1999	725	. . .	. . .	-	14	65	77	95	105	112	68	49	39	29
Female														
2002	232	. . .	. . .	1	10	23	28	15	28	26	24	26	18	8
2001[3]	207	. . .	. . .	-	10	28	22	23	27	19	19	18	17	9
2000	186	. . .	. . .	-	7	25	16	16	18	26	21	18	14	7
1999	176	. . .	. . .	-	3	16	14	21	20	25	16	8	18	7
Non-Hispanic[1]														
Both sexes														
2002	13,330	. . .	. . .	2	148	673	1,008	1,063	1,280	1,539	1,872	1,724	1,267	835
2001[3]	12,627	. . .	. . .	7	150	670	942	989	1,236	1,592	1,734	1,623	1,143	752
2000	11,709	. . .	. . .	5	168	631	863	989	1,097	1,452	1,657	1,417	1,026	667
1999	11,624	. . .	. . .	1	121	555	853	983	1,251	1,528	1,603	1,288	981	580
Male														
2002	9,397	. . .	. . .	2	112	538	808	808	991	1,127	1,278	1,144	843	546
2001[3]	9,005	. . .	. . .	7	116	528	768	780	943	1,184	1,208	1,108	727	486
2000	8,317	. . .	. . .	4	134	491	701	762	840	1,092	1,155	985	655	419
1999	8,197	. . .	. . .	1	97	414	685	766	949	1,127	1,114	874	625	370
Female														
2002	3,933	. . .	. . .	-	36	135	200	255	289	412	594	580	424	289
2001[3]	3,622	. . .	. . .	-	34	142	174	209	293	408	526	515	416	266
2000	3,392	. . .	. . .	1	34	140	162	227	257	360	502	432	371	248
1999	3,427	. . .	. . .	-	24	141	168	217	302	401	489	414	356	210
Non-Hispanic White														
Both sexes														
2002	11,826	. . .	. . .	1	113	584	846	873	1,083	1,351	1,695	1,591	1,166	770
2001[3]	11,165	. . .	. . .	5	109	567	794	813	1,057	1,401	1,574	1,484	1,048	686
2000	10,283	. . .	. . .	5	124	524	689	805	937	1,283	1,477	1,296	926	624
1999	10,218	. . .	. . .	1	92	443	703	799	1,057	1,342	1,451	1,173	901	531
Male														
2002	8,309	. . .	. . .	1	84	469	683	660	832	985	1,158	1,041	788	501
2001[3]	7,939	. . .	. . .	5	84	457	651	646	805	1,035	1,095	1,019	664	442
2000	7,263	. . .	. . .	4	96	413	565	631	710	963	1,028	894	582	393
1999	7,179	. . .	. . .	1	75	333	569	624	805	978	1,019	792	567	339

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
Hispanic										
Male										
2002	19	15	13	11	8	7	2	-	-	2
2001[3]	19	16	17	11	8	10	3	1	-	1
2000	16	16	12	12	15	3	2	1	-	-
1999	20	19	9	6	10	5	1	1	-	1
Female										
2002	6	4	5	5	3	1	-	1	-	-
2001[3]	3	2	6	1	3	-	-	-	-	-
2000	5	3	4	4	1	1	-	-	-	-
1999	8	3	5	5	3	2	1	1	-	-
Non-Hispanic[1]										
Both sexes										
2002	461	328	316	296	254	172	75	13	1	3
2001[3]	421	333	295	287	234	146	62	9	1	1
2000	381	261	292	294	242	175	74	14	3	1
1999	387	333	306	322	261	173	78	14	-	6
Male										
2002	289	204	196	188	154	110	49	7	-	3
2001[3]	259	221	190	176	159	98	39	6	1	1
2000	227	154	175	194	156	115	47	7	3	1
1999	239	206	197	192	158	119	50	8	-	6
Female										
2002	172	124	120	108	100	62	26	6	1	-
2001[3]	162	112	105	111	75	48	23	3	-	-
2000	154	107	117	100	86	60	27	7	-	-
1999	148	127	109	130	103	54	28	6	-	-
Non-Hispanic White										
Both sexes										
2002	421	290	285	269	241	162	69	12	1	3
2001[3]	374	293	266	268	221	134	61	8	1	1
2000	356	237	266	264	219	166	70	13	1	1
1999	346	292	285	297	247	165	76	14	-	3
Male										
2002	271	175	183	174	145	105	45	6	-	3
2001[3]	223	195	167	161	151	92	39	6	1	1
2000	211	138	158	172	143	110	44	6	1	1
1999	211	180	184	176	152	113	50	8	-	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)														
Non-Hispanic White														
Female														
2002	3,517	. . .	. . .	-	29	115	163	213	251	366	537	550	378	269
2001[3]	3,226	. . .	. . .	-	25	110	143	167	252	366	479	465	384	244
2000	3,020	. . .	. . .	1	28	111	124	174	227	320	449	402	344	231
1999	3,039	. . .	. . .	-	17	110	134	175	252	364	432	381	334	192
Non-Hispanic Black														
Both sexes														
2002	855	. . .	. . .	1	25	45	97	109	116	115	118	76	59	29
2001[3]	835	. . .	. . .	2	26	46	86	96	118	120	100	90	56	37
2000	850	. . .	. . .	-	35	53	114	103	98	105	129	77	58	22
1999	812	. . .	. . .	-	24	58	82	105	124	121	112	68	40	21
Male														
2002	650	. . .	. . .	1	20	39	74	87	96	91	84	62	35	21
2001[3]	629	. . .	. . .	2	20	35	76	81	92	92	68	61	37	25
2000	652	. . .	. . .	-	30	39	91	78	82	84	94	61	42	15
1999	623	. . .	. . .	-	18	45	69	87	98	103	72	51	30	12
Female														
2002	205	. . .	. . .	-	5	6	23	22	20	24	34	14	24	8
2001[3]	206	. . .	. . .	-	6	11	10	15	26	28	32	29	19	12
2000	198	. . .	. . .	-	5	14	23	25	16	21	35	16	16	7
1999	189	. . .	. . .	-	6	13	13	18	26	18	40	17	10	9
Origin not stated[2]														
Both sexes														
2002	97	. . .	. . .	-	1	5	5	7	11	10	11	7	12	6
2001[3]	74	. . .	. . .	-	-	4	4	7	8	6	11	9	7	5
2000	81	. . .	. . .	-	2	3	7	6	8	10	11	10	6	6
1999	75	. . .	. . .	1	1	4	2	7	3	4	13	10	11	4
Male														
2002	79	. . .	. . .	-	1	4	5	6	9	7	9	6	8	5
2001[3]	64	. . .	. . .	-	-	3	4	6	8	4	9	9	7	4
2000	59	. . .	. . .	-	1	2	5	3	5	7	9	8	4	6
1999	57	. . .	. . .	1	1	1	1	7	2	2	10	8	8	3
Female														
2002	18	. . .	. . .	-	-	1	-	1	2	3	2	1	4	1
2001[3]	10	. . .	. . .	-	-	1	-	1	-	2	2	-	-	1
2000	22	. . .	. . .	-	1	1	2	3	3	3	2	2	2	-
1999	18	. . .	. . .	-	-	3	1	-	1	2	3	2	3	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
Non-Hispanic White										
Female										
2002	150	115	102	95	96	57	24	6	1	-
2001[3]	151	98	99	107	70	42	22	2	-	-
2000	145	99	108	92	76	56	26	7	-	-
1999	135	112	101	121	95	52	26	6	-	-
Non-Hispanic Black										
Both sexes										
2002	25	11	15	7	3	4	-	-	-	-
2001[3]	22	15	13	4	1	2	-	1	-	-
2000	8	11	12	11	10	3	-	-	1	-
1999	14	22	7	7	2	1	2	-	-	2
Male										
2002	13	11	7	4	3	2	-	-	-	-
2001[3]	16	9	11	3	1	-	-	-	-	-
2000	4	9	7	8	5	2	-	-	1	-
1999	10	17	2	5	1	1	-	-	-	2
Female										
2002	12	-	8	3	-	2	-	-	-	-
2001[3]	6	6	2	1	-	2	-	1	-	-
2000	4	2	5	3	5	1	-	-	-	-
1999	4	5	5	2	1	-	2	-	-	-
Origin not stated[2]										
Both sexes										
2002	7	4	2	1	1	1	1	2	-	3
2001[3]	1	4	1	-	-	-	-	-	-	7
2000	3	2	2	-	2	1	-	1	-	1
1999	1	5	-	1	3	1	-	-	-	4
Male										
2002	6	3	2	1	-	1	1	2	-	3
2001[3]	-	3	1	-	-	-	-	-	-	6
2000	2	2	1	-	2	1	-	-	-	1
1999	-	5	-	-	3	1	-	-	-	4
Female										
2002	1	1	-	-	1	-	-	-	-	-
2001[3]	1	1	-	-	-	-	-	-	-	1
2000	1	-	1	-	-	-	-	1	-	-
1999	1	-	-	1	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)														
All origins														
Both sexes														
2002	17,638	303	423	140	216	1,892	3,327	2,514	1,975	1,734	1,505	1,155	760	523
2001[3]	20,308	332	415	137	189	1,899	3,398	2,817	2,387	2,306	1,962	1,463	1,004	628
2000	16,765	349	356	140	231	1,914	3,025	2,312	1,852	1,732	1,487	1,059	696	455
1999	16,889	331	376	186	246	2,093	2,905	2,362	1,869	1,751	1,455	998	670	420
Male														
2002	13,640	164	234	64	133	1,604	2,870	2,121	1,548	1,288	1,098	841	553	383
2001[3]	15,555	197	238	75	118	1,632	2,909	2,308	1,849	1,698	1,424	1,085	747	461
2000	12,820	203	198	77	150	1,613	2,590	1,918	1,443	1,235	1,084	773	511	338
1999	12,785	187	194	95	155	1,748	2,443	1,922	1,424	1,257	1,050	746	483	323
Female														
2002	3,998	139	189	76	83	288	457	393	427	446	407	314	207	140
2001[3]	4,753	135	177	62	71	267	489	509	538	608	538	378	257	167
2000	3,945	146	158	63	81	301	435	394	409	497	403	286	185	117
1999	4,104	144	182	91	91	345	462	440	445	494	405	252	187	97
Hispanic														
Both sexes														
2002	3,129	52	81	28	36	472	728	554	349	273	191	129	87	56
2001[3]	3,331	55	69	20	36	436	786	573	387	329	238	156	92	46
2000	2,917	54	57	24	52	476	650	463	347	248	199	135	76	50
1999	2,864	53	52	29	60	482	637	482	319	247	185	133	64	47
Male														
2002	2,635	28	49	15	22	425	656	493	296	216	151	104	62	47
2001[3]	2,756	33	38	12	26	397	702	488	312	258	193	121	69	33
2000	2,431	26	30	19	37	434	580	405	292	199	160	98	60	40
1999	2,399	26	25	16	43	425	565	428	270	201	153	100	50	37
Female														
2002	494	24	32	13	14	47	72	61	53	57	40	25	25	9
2001[3]	575	22	31	8	10	39	84	85	75	71	45	35	23	13
2000	486	28	27	5	15	42	70	58	55	49	39	37	16	10
1999	465	27	27	13	17	57	72	54	49	46	32	33	14	10
Non-Hispanic[1]														
Both sexes														
2002	14,346	246	340	112	177	1,406	2,577	1,940	1,608	1,451	1,295	1,016	666	463
2001[3]	16,726	270	343	116	153	1,449	2,579	2,213	1,965	1,954	1,698	1,283	899	571
2000	13,717	291	299	114	179	1,424	2,352	1,845	1,492	1,469	1,280	913	613	400
1999	13,873	270	323	155	186	1,601	2,259	1,870	1,532	1,487	1,256	858	594	367
Male														
2002	10,881	131	184	49	108	1,168	2,194	1,614	1,238	1,065	937	728	485	333
2001[3]	12,608	161	199	62	92	1,222	2,179	1,796	1,513	1,421	1,211	945	667	422

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All origins										
Both sexes										
2002	318	229	192	169	127	64	24	6	1	41
2001[3]	390	310	222	183	129	71	29	4	1	32
2000	283	233	210	172	129	68	23	8	1	30
1999	302	259	214	185	119	65	24	10	-	49
Male										
2002	231	152	125	100	58	32	6	3	-	32
2001[3]	262	204	128	98	57	33	7	1	1	23
2000	199	147	132	85	69	25	10	5	-	15
1999	218	183	121	101	61	28	8	5	-	33
Female										
2002	87	77	67	69	69	32	18	3	1	9
2001[3]	128	106	94	85	72	38	22	3	-	9
2000	84	86	78	87	60	43	13	3	1	15
1999	84	76	93	84	58	37	16	5	-	16
Hispanic										
Both sexes										
2002	31	19	10	8	11	2	-	1	-	11
2001[3]	36	31	14	7	9	2	1	-	-	8
2000	27	15	21	12	4	3	-	1	-	3
1999	22	20	13	6	5	2	-	1	-	5
Male										
2002	25	16	9	4	5	1	-	1	-	10
2001[3]	26	24	9	4	4	1	1	-	-	5
2000	17	10	12	7	1	1	-	1	-	2
1999	19	17	9	5	3	1	-	1	-	5
Female										
2002	6	3	1	4	6	1	-	-	-	1
2001[3]	10	7	5	3	5	1	-	-	-	3
2000	10	5	9	5	3	2	-	-	-	1
1999	3	3	4	1	2	1	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	284	210	178	159	114	62	24	5	1	12
2001[3]	345	274	206	176	120	69	28	3	1	11
2000	252	213	189	160	125	65	23	6	1	12
1999	275	236	201	178	114	63	24	8	-	16
Male										
2002	203	136	112	95	53	31	6	2	-	9
2001[3]	229	176	118	94	53	32	6	-	1	9

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)														
Non-Hispanic[1]														
Male														
2000	10,290	174	168	57	113	1,170	1,989	1,509	1,141	1,022	917	669	446	295
1999	10,275	158	168	79	112	1,316	1,872	1,484	1,140	1,043	887	641	422	282
Female														
2002	3,465	115	156	63	69	238	383	326	370	386	358	288	181	130
2001[3]	4,118	109	144	54	61	227	400	417	452	533	487	338	232	149
2000	3,427	117	131	57	66	254	363	336	351	447	363	244	167	105
1999	3,598	112	155	76	74	285	387	386	392	444	369	217	172	85
Non-Hispanic White														
Both sexes														
2002	5,571	117	169	56	77	366	592	464	513	618	650	533	365	309
2001[3]	7,928	139	160	82	82	382	676	750	892	1,066	1,015	792	581	408
2000	5,429	141	144	72	92	352	522	490	533	656	617	479	355	257
1999	5,748	143	156	78	98	409	553	531	604	696	631	455	359	236
Male														
2002	3,661	70	89	21	40	263	422	323	337	411	463	339	247	205
2001[3]	5,453	84	95	40	43	263	466	542	635	788	710	570	420	292
2000	3,510	96	86	39	54	231	357	321	356	407	429	325	235	183
1999	3,747	94	83	41	52	283	361	360	393	455	426	328	251	171
Female														
2002	1,910	47	80	35	37	103	170	141	176	207	187	194	118	104
2001[3]	2,475	55	65	42	39	119	210	208	257	278	305	222	161	116
2000	1,919	45	58	33	38	121	165	169	177	249	188	154	120	74
1999	2,001	49	73	37	46	126	192	171	211	241	205	127	108	65
Non-Hispanic Black														
Both sexes														
2002	8,147	114	156	45	87	967	1,887	1,412	1,013	770	587	442	268	130
2001[3]	8,077	119	166	28	64	1,009	1,808	1,370	965	817	610	433	271	138
2000	7,748	139	143	40	82	1,012	1,759	1,291	902	754	605	390	225	121
1999	7,536	114	151	65	80	1,117	1,635	1,262	860	730	575	364	193	115
Male														
2002	6,780	54	89	24	60	845	1,695	1,246	846	607	435	359	213	112
2001[3]	6,658	68	100	19	47	911	1,640	1,192	806	584	452	333	211	109
2000	6,382	73	73	16	54	886	1,573	1,138	745	584	443	310	188	96
1999	6,131	58	74	34	55	969	1,464	1,070	701	548	432	285	146	99
Female														
2002	1,367	60	67	21	27	122	192	166	167	163	152	83	55	18
2001[3]	1,419	51	66	9	17	98	168	178	159	233	158	100	60	29
2000	1,366	66	70	24	28	126	186	153	157	170	162	80	37	25

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
Non-Hispanic[1]										
Male										
2000	178	134	120	78	68	24	10	3	-	5
1999	194	163	112	95	58	27	8	4	-	10
Female										
2002	81	74	66	64	61	31	18	3	1	3
2001[3]	116	98	88	82	67	37	22	3	-	2
2000	74	79	69	82	57	41	13	3	1	7
1999	81	73	89	83	56	36	16	4	-	6
Non-Hispanic White										
Both sexes										
2002	187	141	134	110	87	51	18	4	-	10
2001[3]	244	191	151	132	93	61	21	3	-	7
2000	158	132	132	126	85	54	18	5	1	8
1999	182	172	147	126	88	49	21	6	-	8
Male										
2002	127	89	77	62	37	27	3	2	-	7
2001[3]	162	116	81	69	41	27	4	-	-	5
2000	105	75	78	60	41	19	7	2	-	4
1999	122	111	77	64	45	18	6	3	-	3
Female										
2002	60	52	57	48	50	24	15	2	-	3
2001[3]	82	75	70	63	52	34	17	3	-	2
2000	53	57	54	66	44	35	11	3	1	4
1999	60	61	70	62	43	31	15	3	-	5
Non-Hispanic Black										
Both sexes										
2002	86	55	40	47	22	11	5	1	-	2
2001[3]	81	69	48	40	23	7	6	-	1	4
2000	80	72	51	30	34	8	5	1	-	4
1999	79	59	44	46	21	13	3	2	-	8
Male										
2002	68	40	32	32	15	4	2	-	-	2
2001[3]	56	50	35	25	9	4	2	-	1	4
2000	63	56	37	16	23	3	3	1	-	1
1999	60	49	30	29	10	8	2	1	-	7
Female										
2002	18	15	8	15	7	7	3	1	-	-
2001[3]	25	19	13	15	14	3	4	-	-	-
2000	17	16	14	14	11	5	2	-	-	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)														
Non-Hispanic Black														
Female														
1999	1,405	56	77	31	25	148	171	192	159	182	143	79	47	16
Origin not stated[2]														
Both sexes														
2002	163	5	2	-	3	14	22	20	18	10	19	10	7	4
2001[3]	251	7	3	1	-	14	33	31	35	23	26	24	13	11
2000	131	4	-	2	-	14	23	4	13	15	8	11	7	5
1999	152	8	1	2	-	10	9	10	18	17	14	7	12	6
Male														
2002	124	5	1	-	3	11	20	14	14	7	10	9	6	3
2001[3]	191	3	1	1	-	13	28	24	24	19	20	19	11	6
2000	99	3	-	1	-	9	21	4	10	14	7	6	5	3
1999	111	3	1	-	-	7	6	10	14	13	10	5	11	4
Female														
2002	39	-	1	-	-	3	2	6	4	3	9	1	1	1
2001[3]	60	4	2	-	-	1	5	7	11	4	6	5	2	5
2000	32	1	-	1	-	5	2	-	3	1	1	5	2	2
1999	41	5	-	2	-	3	3	-	4	4	4	2	1	2
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
All origins														
Both sexes														
2002	11,829	9	49	55	150	1,567	2,750	2,018	1,447	1,135	907	604	421	263
2001	11,348	11	55	59	121	1,525	2,675	1,940	1,368	1,142	836	552	382	213
2000	10,801	12	28	50	137	1,549	2,414	1,789	1,285	1,064	842	582	377	235
1999	10,828	8	50	61	163	1,708	2,330	1,795	1,253	1,054	814	551	342	206
Male														
2002	9,899	3	30	26	101	1,384	2,467	1,790	1,227	918	705	445	316	197
2001	9,532	7	27	35	90	1,377	2,419	1,708	1,156	892	636	434	292	159
2000	9,006	9	17	28	110	1,381	2,167	1,574	1,076	842	645	445	285	172
1999	8,944	5	26	37	118	1,498	2,071	1,568	1,041	807	607	426	250	167
Female														
2002	1,930	6	19	29	49	183	283	228	220	217	202	159	105	66
2001	1,816	4	28	24	31	148	256	232	212	250	200	118	90	54
2000	1,795	3	11	22	27	168	247	215	209	222	197	137	92	63
1999	1,884	3	24	24	45	210	259	227	212	247	207	125	92	39

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
Non-Hispanic Black										
Female										
1999	19	10	14	17	11	5	1	1	-	1
Origin not stated[2]										
Both sexes										
2002	3	-	4	2	2	-	-	-	-	18
2001[3]	9	5	2	-	-	-	-	1	-	13
2000	4	5	-	-	-	-	-	1	-	15
1999	5	3	-	1	-	-	-	1	-	28
Male										
2002	3	-	4	1	-	-	-	-	-	13
2001[3]	7	4	1	-	-	-	-	1	-	9
2000	4	3	-	-	-	-	-	1	-	8
1999	5	3	-	1	-	-	-	-	-	18
Female										
2002	-	-	-	1	2	-	-	-	-	5
2001[3]	2	1	1	-	-	-	-	-	-	4
2000	-	2	-	-	-	-	-	-	-	7
1999	-	-	-	-	-	-	-	1	-	10
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All origins										
Both sexes										
2002	146	92	69	72	43	14	4	1	-	13
2001	151	114	78	61	38	12	2	2	-	11
2000	132	99	85	54	27	23	4	1	-	12
1999	167	102	91	61	34	17	4	2	-	15
Male										
2002	104	61	40	44	20	8	-	1	-	12
2001	105	79	44	38	16	6	-	1	-	11
2000	94	58	50	23	14	6	2	1	-	7
1999	127	68	56	34	14	7	2	1	-	14
Female										
2002	42	31	29	28	23	6	4	-	-	1
2001	46	35	34	23	22	6	2	1	-	-
2000	38	41	35	31	13	17	2	-	-	5
1999	40	34	35	27	20	10	2	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
Hispanic														
Both sexes														
2002	2,168	2	12	14	26	402	574	434	245	171	123	65	51	21
2001	2,123	4	12	5	25	371	603	416	238	172	134	66	36	14
2000	1,958	1	5	8	33	399	505	350	231	136	113	80	46	22
1999	1,939	2	5	11	41	410	498	344	206	148	110	80	32	23
Male														
2002	1,942	-	8	11	18	371	538	397	222	145	100	53	38	18
2001	1,892	3	5	4	19	339	559	383	213	146	111	53	28	11
2000	1,738	-	3	6	28	370	467	319	205	111	96	62	37	19
1999	1,709	2	3	6	33	373	451	319	180	129	89	58	26	18
Female														
2002	226	2	4	3	8	31	36	37	23	26	23	12	13	3
2001	231	1	7	1	6	32	44	33	25	26	23	13	8	3
2000	220	1	2	2	5	29	38	31	26	25	17	18	9	3
1999	230	-	2	5	8	37	47	25	26	19	21	22	6	5
Non-Hispanic[1]														
Both sexes														
2002	9,575	7	37	41	123	1,152	2,157	1,569	1,191	958	777	535	367	241
2001	9,134	7	43	53	96	1,141	2,045	1,509	1,122	965	697	483	344	198
2000	8,767	11	23	40	104	1,140	1,892	1,436	1,044	917	723	499	329	212
1999	8,821	6	45	50	122	1,288	1,827	1,441	1,035	898	699	469	307	181
Male														
2002	7,884	3	22	15	82	1,002	1,911	1,382	996	768	599	388	276	179
2001	7,559	4	22	30	71	1,026	1,837	1,310	936	743	521	379	262	147
2000	7,205	9	14	21	82	1,002	1,684	1,252	863	721	543	381	247	152
1999	7,177	3	23	31	85	1,118	1,616	1,239	852	671	513	366	222	147
Female														
2002	1,691	4	15	26	41	150	246	187	195	190	178	147	91	62
2001	1,575	3	21	23	25	115	208	199	186	222	176	104	82	51
2000	1,562	2	9	19	22	138	208	184	181	196	180	118	82	60
1999	1,644	3	22	19	37	170	211	202	183	227	186	103	85	34
Non-Hispanic White														
Both sexes														
2002	3,052	1	22	21	45	253	389	308	328	363	372	280	203	170
2001	3,085	4	21	42	48	230	388	330	351	411	348	260	200	129
2000	2,861	4	12	29	43	222	322	301	323	383	346	259	186	143
1999	2,995	2	21	23	57	256	335	336	358	405	344	237	179	115
Male														
2002	2,139	1	12	7	27	191	297	230	235	256	275	175	146	117
2001	2,172	2	12	24	32	176	292	249	246	298	238	190	145	84

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Hispanic										
Both sexes										
2002	10	7	2	1	3	-	-	1	-	4
2001	6	11	3	-	3	1	-	-	-	3
2000	12	5	9	1	-	-	-	-	-	2
1999	12	7	2	3	2	-	-	1	-	2
Male										
2002	8	5	2	1	2	-	-	1	-	4
2001	4	9	1	-	-	1	-	-	-	3
2000	7	2	5	-	-	-	-	-	-	1
1999	9	4	2	3	1	-	-	1	-	2
Female										
2002	2	2	-	-	1	-	-	-	-	-
2001	2	2	2	-	3	-	-	-	-	-
2000	5	3	4	1	-	-	-	-	-	1
1999	3	3	-	-	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	135	85	66	70	40	14	4	-	-	6
2001	143	101	74	61	35	11	2	1	-	3
2000	118	92	76	53	27	23	4	1	-	3
1999	154	95	89	57	32	17	4	1	-	4
Male										
2002	95	56	37	42	18	8	-	-	-	5
2001	99	68	42	38	16	5	-	-	-	3
2000	85	56	45	23	14	6	2	1	-	2
1999	117	64	54	30	13	7	2	-	-	4
Female										
2002	40	29	29	28	22	6	4	-	-	1
2001	44	33	32	23	19	6	2	1	-	-
2000	33	36	31	30	13	17	2	-	-	1
1999	37	31	35	27	19	10	2	1	-	-
Non-Hispanic White										
Both sexes										
2002	83	55	52	54	33	13	3	-	-	4
2001	106	73	53	47	28	11	2	1	-	2
2000	81	63	55	46	17	19	4	1	-	2
1999	100	73	67	39	28	16	4	-	-	-
Male										
2002	57	33	25	31	13	8	-	-	-	3
2001	71	42	27	27	10	5	-	-	-	2

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
Non-Hispanic White														
Male														
2000	1,967	2	6	15	32	165	236	214	230	267	243	185	126	98
1999	2,119	2	11	15	35	197	246	253	262	277	237	174	134	87
Female														
2002	913	-	10	14	18	62	92	78	93	107	97	105	57	53
2001	913	2	9	18	16	54	96	81	105	113	110	70	55	45
2000	894	2	6	14	11	57	86	87	93	116	103	74	60	45
1999	876	-	10	8	22	59	89	83	96	128	107	63	45	28
Non-Hispanic Black														
Both sexes														
2002	6,181	6	14	13	67	851	1,696	1,222	826	558	373	239	146	61
2001	5,790	2	18	11	45	885	1,605	1,137	737	536	328	203	129	60
2000	5,622	7	8	11	57	876	1,520	1,090	700	504	350	221	126	56
1999	5,508	2	18	22	59	979	1,447	1,053	642	463	333	216	104	59
Male														
2002	5,482	2	9	6	48	768	1,557	1,123	731	481	303	202	115	55
2001	5,190	1	9	6	37	826	1,503	1,032	665	431	269	172	108	54
2000	5,011	7	5	6	46	800	1,406	1,002	614	435	280	181	109	43
1999	4,826	1	7	14	46	874	1,334	947	565	374	262	179	77	55
Female														
2002	699	4	5	7	19	83	139	99	95	77	70	37	31	6
2001	600	1	9	5	8	59	102	105	72	105	59	31	21	6
2000	611	-	3	5	11	76	114	88	86	69	70	40	17	13
1999	682	1	11	8	13	105	113	106	77	89	71	37	27	4
Origin not stated[2]														
Both sexes														
2002	86	-	-	-	1	13	19	15	11	6	7	4	3	1
2001	91	-	-	1	-	13	27	15	8	5	5	3	2	1
2000	76	-	-	2	-	10	17	3	10	11	6	3	2	1
1999	68	-	-	-	-	10	5	10	12	8	5	2	3	2
Male														
2002	73	-	-	-	1	11	18	11	9	5	6	4	2	-
2001	81	-	-	1	-	12	23	15	7	3	4	2	2	1
2000	63	-	-	1	-	9	16	3	8	10	6	2	1	1
1999	58	-	-	-	-	7	4	10	9	7	5	2	2	2
Female														
2002	13	-	-	-	-	2	1	4	2	1	1	-	1	1
2001	10	-	-	-	-	1	4	-	1	2	1	1	-	-
2000	13	-	-	1	-	1	1	-	2	1	-	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Non-Hispanic White										
Male										
2000	51	35	28	20	5	5	2	1	-	1
1999	70	46	38	16	11	6	2	-	-	-
Female										
2002	26	22	27	23	20	5	3	-	-	1
2001	35	31	26	20	18	6	2	1	-	-
2000	30	28	27	26	12	14	2	-	-	1
1999	30	27	29	23	17	10	2	-	-	-
Non-Hispanic Black										
Both sexes										
2002	46	26	13	14	6	1	1	-	-	2
2001	32	25	18	13	5	-	-	-	-	1
2000	29	25	20	7	10	4	-	-	-	1
1999	45	20	19	17	4	1	-	1	-	4
Male										
2002	33	21	11	10	5	-	-	-	-	2
2001	24	23	14	11	4	-	-	-	-	1
2000	26	20	17	3	9	1	-	-	-	1
1999	39	17	14	14	2	1	-	-	-	4
Female										
2002	13	5	2	4	1	1	1	-	-	-
2001	8	2	4	2	1	-	-	-	-	-
2000	3	5	3	4	1	3	-	-	-	-
1999	6	3	5	3	2	-	-	1	-	-
Origin not stated[2]										
Both sexes										
2002	1	-	1	1	-	-	-	-	-	3
2001	2	2	1	-	-	-	-	1	-	5
2000	2	2	-	-	-	-	-	-	-	7
1999	1	-	-	1	-	-	-	-	-	9
Male										
2002	1	-	1	1	-	-	-	-	-	3
2001	2	2	1	-	-	-	-	1	-	5
2000	2	-	-	-	-	-	-	-	-	4
1999	1	-	-	1	-	-	-	-	-	8
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	2	-	-	-	-	-	-	-	3

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)														
Origin not stated[2]														
Female														
1999	10	-	-	-	-	3	1	-	3	1	-	-	1	-
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)														
All origins														
Both sexes														
2002	5,809	294	374	85	66	325	577	496	528	599	598	551	339	260
2001[3]	8,960	321	360	78	68	374	723	877	1,019	1,164	1,126	911	622	415
2000	5,964	337	328	90	94	365	611	523	567	668	645	477	319	220
1999	6,061	323	326	125	83	385	575	567	616	697	641	447	328	214
Male														
2002	3,741	161	204	38	32	220	403	331	321	370	393	396	237	186
2001[3]	6,023	190	211	40	28	255	490	600	693	806	788	651	455	302
2000	3,814	194	181	49	40	232	423	344	367	393	439	328	226	166
1999	3,841	182	168	58	37	250	372	354	383	450	443	320	233	156
Female														
2002	2,068	133	170	47	34	105	174	165	207	229	205	155	102	74
2001[3]	2,937	131	149	38	40	119	233	277	326	358	338	260	167	113
2000	2,150	143	147	41	54	133	188	179	200	275	206	149	93	54
1999	2,220	141	158	67	46	135	203	213	233	247	198	127	95	58
Hispanic														
Both sexes														
2002	961	50	69	14	10	70	154	120	104	102	68	64	36	35
2001[3]	1,208	51	57	15	11	65	183	157	149	157	104	90	56	32
2000	959	53	52	16	19	77	145	113	116	112	86	55	30	28
1999	925	51	47	18	19	72	139	138	113	99	75	53	32	24
Male														
2002	693	28	41	4	4	54	118	96	74	71	51	51	24	29
2001[3]	864	30	33	8	7	58	143	105	99	112	82	68	41	22
2000	693	26	27	13	9	64	113	86	87	88	64	36	23	21
1999	690	24	22	10	10	52	114	109	90	72	64	42	24	19
Female														
2002	268	22	28	10	6	16	36	24	30	31	17	13	12	6
2001[3]	344	21	24	7	4	7	40	52	50	45	22	22	15	10
2000	266	27	25	3	10	13	32	27	29	24	22	19	7	7
1999	235	27	25	8	9	20	25	29	23	27	11	11	8	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Origin not stated[2]										
Female										
1999	-	-	-	-	-	-	-	-	-	1
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)										
All origins										
Both sexes										
2002	172	137	123	97	84	50	20	5	1	28
2001[3]	239	196	144	122	91	59	27	2	1	21
2000	151	134	125	118	102	45	19	7	1	18
1999	135	157	123	124	85	48	20	8	-	34
Male										
2002	127	91	85	56	38	24	6	2	-	20
2001[3]	157	125	84	60	41	27	7	-	1	12
2000	105	89	82	62	55	19	8	4	-	8
1999	91	115	65	67	47	21	6	4	-	19
Female										
2002	45	46	38	41	46	26	14	3	1	8
2001[3]	82	71	60	62	50	32	20	2	-	9
2000	46	45	43	56	47	26	11	3	1	10
1999	44	42	58	57	38	27	14	4	-	15
Hispanic										
Both sexes										
2002	21	12	8	7	8	2	-	-	-	7
2001[3]	30	20	11	7	6	1	1	-	-	5
2000	15	10	12	11	4	3	-	1	-	1
1999	10	13	11	3	3	2	-	-	-	3
Male										
2002	17	11	7	3	3	1	-	-	-	6
2001[3]	22	15	8	4	4	-	1	-	-	2
2000	10	8	7	7	1	1	-	1	-	1
1999	10	13	7	2	2	1	-	-	-	3
Female										
2002	4	1	1	4	5	1	-	-	-	1
2001[3]	8	5	3	3	2	1	-	-	-	3
2000	5	2	5	4	3	2	-	-	-	-
1999	-	-	4	1	1	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)														
Non-Hispanic[1]														
Both sexes														
2002	4,771	239	303	71	54	254	420	371	417	493	518	481	299	222
2001[3]	7,592	263	300	63	57	308	534	704	843	989	1,001	800	555	373
2000	4,950	280	276	74	75	284	460	409	448	552	557	414	284	188
1999	5,052	264	278	105	64	313	432	429	497	589	557	389	287	186
Male														
2002	2,997	128	162	34	26	166	283	232	242	297	338	340	209	154
2001[3]	5,049	157	177	32	21	196	342	486	577	678	690	566	405	275
2000	3,085	165	154	36	31	168	305	257	278	301	374	288	199	143
1999	3,098	155	145	48	27	198	256	245	288	372	374	275	200	135
Female														
2002	1,774	111	141	37	28	88	137	139	175	196	180	141	90	68
2001[3]	2,543	106	123	31	36	112	192	218	266	311	311	234	150	98
2000	1,865	115	122	38	44	116	155	152	170	251	183	126	85	45
1999	1,954	109	133	57	37	115	176	184	209	217	183	114	87	51
Non-Hispanic White														
Both sexes														
2002	2,519	116	147	35	32	113	203	156	185	255	278	253	162	139
2001[3]	4,843	135	139	40	34	152	288	420	541	655	667	532	381	279
2000	2,568	137	132	43	49	130	200	189	210	273	271	220	169	114
1999	2,753	141	135	55	41	153	218	195	246	291	287	218	180	121
Male														
2002	1,522	69	77	14	13	72	125	93	102	155	188	164	101	88
2001[3]	3,281	82	83	16	11	87	174	293	389	490	472	380	275	208
2000	1,543	94	80	24	22	66	121	107	126	140	186	140	109	85
1999	1,628	92	72	26	17	86	115	107	131	178	189	154	117	84
Female														
2002	997	47	70	21	19	41	78	63	83	100	90	89	61	51
2001[3]	1,562	53	56	24	23	65	114	127	152	165	195	152	106	71
2000	1,025	43	52	19	27	64	79	82	84	133	85	80	60	29
1999	1,125	49	63	29	24	67	103	88	115	113	98	64	63	37
Non-Hispanic Black														
Both sexes														
2002	1,966	108	142	32	20	116	191	190	187	212	214	203	122	69
2001[3]	2,287	117	148	17	19	124	203	233	228	281	282	230	142	78
2000	2,126	132	135	29	25	136	239	201	202	250	255	169	99	65
1999	2,028	112	133	43	21	138	188	209	218	267	242	148	89	56

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)										
Non-Hispanic[1]										
Both sexes										
2002	149	125	112	89	74	48	20	5	1	6
2001[3]	202	173	132	115	85	58	26	2	1	8
2000	134	121	113	107	98	42	19	5	1	9
1999	121	141	112	121	82	46	20	7	-	12
Male										
2002	108	80	75	53	35	23	6	2	-	4
2001[3]	130	108	76	56	37	27	6	-	1	6
2000	93	78	75	55	54	18	8	2	-	3
1999	77	99	58	65	45	20	6	4	-	6
Female										
2002	41	45	37	36	39	25	14	3	1	2
2001[3]	72	65	56	59	48	31	20	2	-	2
2000	41	43	38	52	44	24	11	3	1	6
1999	44	42	54	56	37	26	14	3	-	6
Non-Hispanic White										
Both sexes										
2002	104	86	82	56	54	38	15	4	-	6
2001[3]	138	118	98	85	65	50	19	2	-	5
2000	77	69	77	80	68	35	14	4	1	6
1999	82	99	80	87	60	33	17	6	-	8
Male										
2002	70	56	52	31	24	19	3	2	-	4
2001[3]	91	74	54	42	31	22	4	-	-	3
2000	54	40	50	40	36	14	5	1	-	3
1999	52	65	39	48	34	12	4	3	-	3
Female										
2002	34	30	30	25	30	19	12	2	-	2
2001[3]	47	44	44	43	34	28	15	2	-	2
2000	23	29	27	40	32	21	9	3	1	3
1999	30	34	41	39	26	21	13	3	-	5
Non-Hispanic Black										
Both sexes										
2002	40	29	27	33	16	10	4	1	-	-
2001[3]	49	44	30	27	18	7	6	-	1	3
2000	51	47	31	23	24	4	5	1	-	3
1999	34	39	25	29	17	12	3	1	-	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)														
Non-Hispanic Black														
Male														
2002	1,298	52	80	18	12	77	138	123	115	126	132	157	98	57
2001[3]	1,468	67	91	13	10	85	137	160	141	153	183	161	103	55
2000	1,371	66	68	10	8	86	167	136	131	149	163	129	79	53
1999	1,305	57	67	20	9	95	130	123	136	174	170	106	69	44
Female														
2002	668	56	62	14	8	39	53	67	72	86	82	46	24	12
2001[3]	819	50	57	4	9	39	66	73	87	128	99	69	39	23
2000	755	66	67	19	17	50	72	65	71	101	92	40	20	12
1999	723	55	66	23	12	43	58	86	82	93	72	42	20	12
Origin not stated[2]														
Both sexes														
2002	77	5	2	-	2	1	3	5	7	4	12	6	4	3
2001[3]	160	7	3	-	-	1	6	16	27	18	21	21	11	10
2000	55	4	-	-	-	4	6	1	3	4	2	8	5	4
1999	84	8	1	2	-	-	4	-	6	9	9	5	9	4
Male														
2002	51	5	1	-	2	-	2	3	5	2	4	5	4	3
2001[3]	110	3	1	-	-	1	5	9	17	16	16	17	9	5
2000	36	3	-	-	-	-	5	1	2	4	1	4	4	2
1999	53	3	1	-	-	-	2	-	5	6	5	3	9	2
Female														
2002	26	-	1	-	-	1	1	2	2	2	8	1	-	-
2001[3]	50	4	2	-	-	-	1	7	10	2	5	4	2	5
2000	19	1	-	-	-	4	1	-	1	-	1	4	1	2
1999	31	5	-	2	-	-	2	-	1	3	4	2	-	2
Legal intervention (Y35,Y89.0)														
All origins														
Both sexes														
2002	384	-	-	-	3	24	67	51	51	64	45	43	13	10
2001	396	-	-	-	-	26	62	47	60	65	50	33	18	15
2000	359	-	-	-	3	27	46	59	51	60	41	38	17	6
1999	398	-	-	-	1	20	47	66	70	72	49	19	31	9
Male														
2002	369	-	-	-	3	23	64	51	48	60	45	40	12	10
2001	380	-	-	-	-	25	59	46	58	63	45	33	18	15

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)										
Non-Hispanic Black										
Male										
2002	35	19	21	22	10	4	2	-	-	-
2001[3]	32	27	21	14	5	4	2	-	1	3
2000	37	36	20	13	14	2	3	1	-	-
1999	21	32	16	15	8	7	2	1	-	3
Female										
2002	5	10	6	11	6	6	2	1	-	-
2001[3]	17	17	9	13	13	3	4	-	-	-
2000	14	11	11	10	10	2	2	-	-	3
1999	13	7	9	14	9	5	1	-	-	1
Origin not stated[2]										
Both sexes										
2002	2	-	3	1	2	-	-	-	-	15
2001[3]	7	3	1	-	-	-	-	-	-	8
2000	2	3	-	-	-	-	-	1	-	8
1999	4	3	-	-	-	-	-	1	-	19
Male										
2002	2	-	3	-	-	-	-	-	-	10
2001[3]	5	2	-	-	-	-	-	-	-	4
2000	2	3	-	-	-	-	-	1	-	4
1999	4	3	-	-	-	-	-	-	-	10
Female										
2002	-	-	-	1	2	-	-	-	-	5
2001[3]	2	1	1	-	-	-	-	-	-	4
2000	-	-	-	-	-	-	-	-	-	4
1999	-	-	-	-	-	-	-	1	-	9
Legal intervention (Y35,Y89.0)										
All origins										
Both sexes										
2002	6	4	2	-	1	-	-	-	-	-
2001	9	4	2	3	1	1	-	-	-	-
2000	4	2	3	1	-	1	-	-	-	-
1999	11	2	-	1	-	-	-	-	-	-
Male										
2002	6	4	2	-	1	-	-	-	-	-
2001	7	4	2	3	1	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Legal intervention (Y35,Y89.0)														
All origins														
Male														
2000	348	-	-	-	3	27	44	58	50	58	38	37	17	6
1999	392	-	-	-	1	20	47	66	69	69	48	19	30	9
Female														
2002	15	-	-	-	-	1	3	-	3	4	-	3	1	-
2001	16	-	-	-	-	1	3	1	2	2	5	-	-	-
2000	11	-	-	-	-	-	2	1	1	2	3	1	-	-
1999	6	-	-	-	-	-	-	-	1	3	1	-	1	-
Hispanic														
Both sexes														
2002	68	-	-	-	1	5	17	13	9	9	5	2	5	1
2001	65	-	-	-	-	7	16	15	9	6	7	2	2	-
2000	48	-	-	-	1	5	13	6	9	5	4	2	2	-
1999	58	-	-	-	-	5	8	10	13	10	4	1	4	-
Male														
2002	67	-	-	-	1	5	17	13	8	9	5	2	5	1
2001	65	-	-	-	-	7	16	15	9	6	7	2	2	-
2000	47	-	-	-	1	5	12	6	9	5	4	2	2	-
1999	58	-	-	-	-	5	8	10	13	10	4	1	4	-
Female														
2002	1	-	-	-	-	-	-	-	1	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	1	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	315	-	-	-	2	19	50	38	42	54	40	41	8	9
2001	330	-	-	-	-	19	46	32	51	59	43	30	16	15
2000	308	-	-	-	2	22	32	53	42	53	37	36	15	6
1999	338	-	-	-	1	15	39	55	57	61	45	18	27	9
Male														
2002	301	-	-	-	2	18	47	38	40	50	40	38	7	9
2001	314	-	-	-	-	18	43	31	49	57	38	30	16	15
2000	298	-	-	-	2	22	31	52	41	51	34	35	15	6
1999	332	-	-	-	1	15	39	55	56	58	44	18	26	9
Female														
2002	14	-	-	-	-	1	3	-	2	4	-	3	1	-
2001	16	-	-	-	-	1	3	1	2	2	5	-	-	-
2000	10	-	-	-	-	-	1	1	1	2	3	1	-	-
1999	6	-	-	-	-	-	-	-	1	3	1	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Legal intervention (Y35,Y89.0)										
All origins										
Male										
2000	3	2	3	1	-	1	-	-	-	-
1999	11	2	-	1	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	1	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	-	-
Male										
2002	-	1	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	2	1	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	6	3	2	-	1	-	-	-	-	-
2001	8	4	2	3	1	1	-	-	-	-
2000	3	2	3	1	-	1	-	-	-	-
1999	9	1	-	1	-	-	-	-	-	-
Male										
2002	6	3	2	-	1	-	-	-	-	-
2001	6	4	2	3	1	1	-	-	-	-
2000	2	2	3	1	-	1	-	-	-	-
1999	9	1	-	1	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	2	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Legal intervention (Y35,Y89.0)														
Non-Hispanic White														
Both sexes														
2002	191	-	-	-	-	9	19	16	32	36	28	26	8	7
2001	230	-	-	-	-	8	29	20	36	40	34	21	15	10
2000	187	-	-	-	-	6	20	30	27	31	21	30	9	5
1999	197	-	-	-	-	5	15	28	34	36	29	14	21	7
Male														
2002	181	-	-	-	-	8	17	16	31	34	28	23	7	7
2001	221	-	-	-	-	7	28	19	36	38	31	21	15	10
2000	182	-	-	-	-	6	19	29	26	30	21	30	9	5
1999	192	-	-	-	-	5	15	28	34	33	28	14	20	7
Female														
2002	10	-	-	-	-	1	2	-	1	2	-	3	1	-
2001	9	-	-	-	-	1	1	1	-	2	3	-	-	-
2000	5	-	-	-	-	-	1	1	1	1	-	-	-	-
1999	5	-	-	-	-	-	-	-	-	3	1	-	1	-
Non-Hispanic Black														
Both sexes														
2002	109	-	-	-	2	10	27	19	9	17	11	11	-	1
2001	82	-	-	-	-	10	16	9	12	15	7	8	1	3
2000	110	-	-	-	2	15	12	19	14	20	15	6	5	-
1999	125	-	-	-	1	10	21	24	18	21	15	4	6	2
Male														
2002	105	-	-	-	2	10	26	19	8	15	11	11	-	1
2001	77	-	-	-	-	10	14	9	11	15	5	8	1	3
2000	107	-	-	-	2	15	12	19	14	20	13	5	5	-
1999	124	-	-	-	1	10	21	24	17	21	15	4	6	2
Female														
2002	4	-	-	-	-	-	1	-	1	2	-	-	-	-
2001	5	-	-	-	-	-	2	-	1	-	2	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	2	1	-	-
1999	1	-	-	-	-	-	-	-	1	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	3	-	-	-	-	-	1	-	-	2	-	-	-	-
1999	2	-	-	-	-	-	-	1	-	1	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	-	1	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	1	-	-
2000	3	-	-	-	-	-	1	-	-	2	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Legal intervention (Y35,Y89.0)										
Non-Hispanic White										
Both sexes										
2002	5	3	1	-	1	-	-	-	-	-
2001	7	4	2	3	1	-	-	-	-	-
2000	3	1	2	1	-	1	-	-	-	-
1999	6	1	-	1	-	-	-	-	-	-
Male										
2002	5	3	1	-	1	-	-	-	-	-
2001	6	4	2	3	1	-	-	-	-	-
2000	2	1	2	1	-	1	-	-	-	-
1999	6	1	-	1	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	1	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	-
2000	-	1	1	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-
Male										
2002	1	-	1	-	-	-	-	-	-	-
2001	-	-	-	-	-	1	-	-	-	-
2000	-	1	1	-	-	-	-	-	-	-
1999	3	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Legal intervention (Y35,Y89.0)														
Origin not stated[2]														
Male														
1999	2	-	-	-	-	-	-	1	-	1	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)														
All origins														
Both sexes														
2002	4,830	101	57	26	42	143	344	352	476	727	789	709	431	207
2001	4,198	83	44	17	34	132	280	311	417	674	722	619	371	152
2000	3,819	65	39	18	42	120	250	306	373	599	687	543	298	154
1999	3,917	55	43	12	37	159	271	313	430	644	651	528	265	136
Male														
2002	3,114	52	28	11	27	108	250	256	329	494	487	432	264	131
2001	2,833	49	19	10	28	102	214	231	284	438	493	433	229	105
2000	2,635	27	26	13	37	93	192	223	266	415	470	374	196	105
1999	2,692	33	21	6	32	132	213	230	293	446	441	357	175	94
Female														
2002	1,716	49	29	15	15	35	94	96	147	233	302	277	167	76
2001	1,365	34	25	7	6	30	66	80	133	236	229	186	142	47
2000	1,184	38	13	5	5	27	58	83	107	184	217	169	102	49
1999	1,225	22	22	6	5	27	58	83	137	198	210	171	90	42
Hispanic														
Both sexes														
2002	320	14	11	3	1	13	38	29	44	53	37	34	15	6
2001	283	7	6	1	3	13	21	31	32	49	43	31	23	5
2000	215	5	5	-	1	11	23	29	32	41	23	19	6	7
1999	249	6	9	1	3	20	32	32	37	36	25	18	10	8
Male														
2002	242	6	7	1	-	12	33	23	35	40	24	28	12	5
2001	216	4	3	1	3	11	15	25	27	35	38	19	17	5
2000	165	2	2	-	1	10	20	22	27	31	16	16	4	6
1999	200	3	6	1	2	17	29	26	28	29	20	15	8	7
Female														
2002	78	8	4	2	1	1	5	6	9	13	13	6	3	1
2001	67	3	3	-	-	2	6	6	5	14	5	12	6	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Legal intervention (Y35,Y89.0)										
Origin not stated[2]										
Male										
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All origins										
Both sexes										
2002	113	80	59	44	51	39	17	8	1	14
2001	73	58	41	53	55	22	11	9	2	18
2000	74	55	49	43	36	26	25	5	1	11
1999	86	72	42	51	45	37	18	2	1	19
Male										
2002	72	46	37	24	22	20	6	5	-	13
2001	41	40	24	28	29	12	6	2	-	16
2000	47	38	32	29	20	14	5	2	-	11
1999	54	47	25	31	24	20	4	1	-	13
Female										
2002	41	34	22	20	29	19	11	3	1	1
2001	32	18	17	25	26	10	5	7	2	2
2000	27	17	17	14	16	12	20	3	1	-
1999	32	25	17	20	21	17	14	1	1	6
Hispanic										
Both sexes										
2002	9	4	5	1	1	-	-	-	-	2
2001	4	4	2	1	1	2	2	-	-	2
2000	5	4	1	-	2	-	-	-	-	1
1999	5	2	1	2	1	-	-	-	-	1
Male										
2002	9	2	2	-	1	-	-	-	-	2
2001	2	3	2	-	1	2	1	-	-	2
2000	3	2	-	-	2	-	-	-	-	1
1999	4	2	-	2	-	-	-	-	-	1
Female										
2002	-	2	3	1	-	-	-	-	-	-
2001	2	1	-	1	-	-	1	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)														
Hispanic														
Female														
2000	50	3	3	-	-	1	3	7	5	10	7	3	2	1
1999	49	3	3	-	1	3	3	6	9	7	5	3	2	1
Non-Hispanic[1]														
Both sexes														
2002	4,470	87	44	23	40	127	305	320	429	671	749	671	415	198
2001	3,876	76	38	16	31	118	259	279	382	622	676	586	341	143
2000	3,571	58	34	18	41	109	227	275	339	553	661	522	290	143
1999	3,630	49	34	11	34	139	237	276	390	606	624	507	250	126
Male														
2002	2,836	46	21	10	26	94	216	230	291	452	460	400	251	123
2001	2,586	45	16	9	25	90	199	205	254	400	453	414	208	96
2000	2,445	25	24	13	36	83	172	199	237	382	452	356	190	97
1999	2,463	30	15	5	30	115	182	200	262	415	419	340	163	85
Female														
2002	1,634	41	23	13	14	33	89	90	138	219	289	271	164	75
2001	1,290	31	22	7	6	28	60	74	128	222	223	172	133	47
2000	1,126	33	10	5	5	26	55	76	102	171	209	166	100	46
1999	1,167	19	19	6	4	24	55	76	128	191	205	167	87	41
Non-Hispanic White														
Both sexes														
2002	3,627	43	33	13	31	107	263	262	345	546	614	544	329	170
2001	3,155	40	25	8	27	103	215	229	320	499	555	478	271	121
2000	2,823	35	22	9	28	93	178	222	268	448	517	415	232	105
1999	2,848	30	15	9	29	106	184	218	313	480	480	390	188	102
Male														
2002	2,265	23	16	4	20	77	187	187	232	359	380	321	183	102
2001	2,054	25	10	3	22	78	160	165	212	312	361	327	158	77
2000	1,900	16	15	6	25	70	131	158	183	309	352	277	148	66
1999	1,887	19	8	4	26	90	142	150	207	319	327	247	116	62
Female														
2002	1,362	20	17	9	11	30	76	75	113	187	234	223	146	68
2001	1,101	15	15	5	5	25	55	64	108	187	194	151	113	44
2000	923	19	7	3	3	23	47	64	85	139	165	138	84	39
1999	961	11	7	5	3	16	42	68	106	161	153	143	72	40
Non-Hispanic Black														
Both sexes														
2002	738	37	7	9	6	14	38	51	67	115	125	116	79	23
2001	629	36	13	7	4	12	31	37	53	110	112	98	65	20

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
Hispanic										
Female										
2000	2	2	1	-	-	-	-	-	-	-
1999	1	-	1	-	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	103	75	53	43	50	38	17	6	1	5
2001	67	53	38	51	54	20	9	9	2	6
2000	68	49	47	43	34	26	24	5	1	4
1999	81	68	39	49	44	36	18	1	1	10
Male										
2002	62	43	34	24	21	19	6	3	-	4
2001	38	36	21	27	28	10	5	2	-	5
2000	43	34	31	29	18	14	4	2	-	4
1999	50	44	23	29	24	20	4	-	-	8
Female										
2002	41	32	19	19	29	19	11	3	1	1
2001	29	17	17	24	26	10	4	7	2	1
2000	25	15	16	14	16	12	20	3	1	-
1999	31	24	16	20	20	16	14	1	1	2
Non-Hispanic White										
Both sexes										
2002	83	61	44	36	41	36	16	5	1	4
2001	55	41	34	46	47	18	9	8	2	4
2000	56	39	37	38	32	22	20	5	-	2
1999	70	60	32	45	41	32	16	1	1	6
Male										
2002	49	32	25	19	19	18	6	3	-	3
2001	29	28	17	25	26	9	5	2	-	3
2000	35	25	23	27	16	11	3	2	-	2
1999	39	39	18	26	21	18	4	-	-	5
Female										
2002	34	29	19	17	22	18	10	2	1	1
2001	26	13	17	21	21	9	4	6	2	1
2000	21	14	14	11	16	11	17	3	-	-
1999	31	21	14	19	20	14	12	1	1	1
Non-Hispanic Black										
Both sexes										
2002	16	9	7	7	9	1	1	1	-	-
2001	10	10	2	2	3	2	-	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)														
Non-Hispanic Black														
Both sexes														
2000	654	14	11	9	12	12	38	44	59	97	133	100	53	31
1999	686	18	15	2	5	25	45	51	68	113	126	107	59	20
Male														
2002	504	18	3	5	6	12	27	38	46	87	74	74	62	17
2001	467	20	6	5	3	10	27	30	35	79	86	79	48	18
2000	478	3	8	7	10	12	33	34	44	67	93	75	38	24
1999	514	10	7	1	4	20	35	44	47	86	84	86	46	19
Female														
2002	234	19	4	4	-	2	11	13	21	28	51	42	17	6
2001	162	16	7	2	1	2	4	7	18	31	26	19	17	2
2000	176	11	3	2	2	-	5	10	15	30	40	25	15	7
1999	172	8	8	1	1	5	10	7	21	27	42	21	13	1
Origin not stated[2]														
Both sexes														
2002	40	-	2	-	1	3	1	3	3	3	3	4	1	3
2001	39	-	-	-	-	1	-	1	3	3	3	2	7	4
2000	33	2	-	-	-	-	-	2	2	5	3	2	2	4
1999	38	-	-	-	-	-	2	5	3	2	2	3	5	2
Male														
2002	36	-	-	-	1	2	1	3	3	2	3	4	1	3
2001	31	-	-	-	-	1	-	1	3	3	2	-	4	4
2000	25	-	-	-	-	-	-	2	2	2	2	2	2	2
1999	29	-	-	-	-	-	2	4	3	2	2	2	4	2
Female														
2002	4	-	2	-	-	1	-	-	-	1	-	-	-	-
2001	8	-	-	-	-	-	-	-	-	-	1	2	3	-
2000	8	2	-	-	-	-	-	-	-	3	1	-	-	2
1999	9	-	-	-	-	-	-	1	-	-	-	1	1	-
Discharge of firearms, undetermined intent (Y22-Y24)														
All origins														
Both sexes														
2002	243	-	1	2	4	35	44	27	19	31	19	19	10	9
2001	231	-	-	2	4	24	40	32	16	33	17	16	13	9
2000	230	-	-	2	8	26	45	32	20	18	21	9	14	12
1999	324	-	3	-	12	68	59	32	19	28	29	18	10	6
Male														
2002	199	-	1	2	3	34	37	23	16	22	17	14	9	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
Non-Hispanic Black										
Both sexes										
2000	11	8	7	4	2	3	3	-	1	2
1999	8	7	5	4	1	3	-	-	-	4
Male										
2002	13	8	7	5	2	-	-	-	-	-
2001	8	8	2	-	1	1	-	-	-	1
2000	8	7	6	2	2	2	1	-	-	2
1999	8	5	4	3	1	1	-	-	-	3
Female										
2002	3	1	-	2	7	1	1	1	-	-
2001	2	2	-	2	2	1	-	1	-	-
2000	3	1	1	2	-	1	2	-	1	-
1999	-	2	1	1	-	2	-	-	-	1
Origin not stated[2]										
Both sexes										
2002	1	1	1	-	-	1	-	2	-	7
2001	2	1	1	1	-	-	-	-	-	10
2000	1	2	1	-	-	-	1	-	-	6
1999	-	2	2	-	-	1	-	1	-	8
Male										
2002	1	1	1	-	-	1	-	2	-	7
2001	1	1	1	1	-	-	-	-	-	9
2000	1	2	1	-	-	-	1	-	-	6
1999	-	1	2	-	-	-	-	1	-	4
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	1	-	-	-	4
Discharge of firearms, undetermined intent (Y22-Y24)										
All origins										
Both sexes										
2002	3	8	7	3	2	-	-	-	-	-
2001	3	6	4	6	5	1	-	-	-	-
2000	3	7	4	5	1	2	-	-	-	1
1999	8	9	6	6	7	4	-	-	-	-
Male										
2002	2	4	6	3	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Discharge of firearms, undetermined intent (Y22-Y24)														
All origins														
Male														
2001	190	-	-	2	4	23	36	27	15	20	13	12	9	8
2000	190	-	-	2	7	25	37	26	17	12	18	6	12	10
1999	277	-	3	-	12	61	50	28	16	24	21	15	6	5
Female														
2002	44	-	-	-	1	1	7	4	3	9	2	5	1	4
2001	41	-	-	-	-	1	4	5	1	13	4	4	4	1
2000	40	-	-	-	1	1	8	6	3	6	3	3	2	2
1999	47	-	-	-	-	7	9	4	3	4	8	3	4	1
Hispanic														
Both sexes														
2002	23	-	-	-	-	6	8	4	2	3	-	-	-	-
2001	21	-	-	-	1	4	4	3	1	4	-	2	1	1
2000	20	-	-	-	-	7	6	3	1	2	-	-	-	-
1999	31	-	1	-	1	9	9	2	1	2	1	2	-	1
Male														
2002	21	-	-	-	-	6	7	4	2	2	-	-	-	-
2001	16	-	-	-	1	4	3	2	1	2	-	1	1	1
2000	17	-	-	-	-	7	6	2	-	1	-	-	-	-
1999	26	-	1	-	1	8	8	1	1	2	-	1	-	1
Female														
2002	2	-	-	-	-	-	1	-	-	1	-	-	-	-
2001	5	-	-	-	-	-	1	1	-	2	-	1	-	-
2000	3	-	-	-	-	-	-	1	1	1	-	-	-	-
1999	5	-	-	-	-	1	1	1	-	-	1	1	-	-
Non-Hispanic[1]														
Both sexes														
2002	215	-	1	2	3	27	36	22	17	28	19	19	10	8
2001	208	-	-	2	3	20	36	29	14	29	17	14	12	8
2000	209	-	-	2	8	19	39	29	19	16	21	9	14	12
1999	293	-	2	-	11	59	50	30	18	26	28	16	10	5
Male														
2002	173	-	1	2	2	26	30	18	14	20	17	14	9	4
2001	172	-	-	2	3	19	33	25	13	18	13	11	8	7
2000	172	-	-	2	7	18	31	24	17	11	18	6	12	10
1999	251	-	2	-	11	53	42	27	15	22	21	14	6	4
Female														
2002	42	-	-	-	1	1	6	4	3	8	2	5	1	4
2001	36	-	-	-	-	1	3	4	1	11	4	3	4	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)										
All origins										
Male										
2001	3	6	2	6	3	1	-	-	-	-
2000	1	5	4	4	1	2	-	-	-	1
1999	6	8	6	5	7	4	-	-	-	-
Female										
2002	1	4	1	-	1	-	-	-	-	-
2001	-	-	2	-	2	-	-	-	-	-
2000	2	2	-	1	-	-	-	-	-	-
1999	2	1	-	1	-	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	2	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	2	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	3	8	7	3	2	-	-	-	-	-
2001	3	5	4	6	5	1	-	-	-	-
2000	3	6	4	5	1	2	-	-	-	-
1999	8	7	6	6	7	4	-	-	-	-
Male										
2002	2	4	6	3	1	-	-	-	-	-
2001	3	5	2	6	3	1	-	-	-	-
2000	1	4	4	4	1	2	-	-	-	-
1999	6	6	6	5	7	4	-	-	-	-
Female										
2002	1	4	1	-	1	-	-	-	-	-
2001	-	-	2	-	2	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Discharge of firearms, undetermined intent (Y22-Y24)														
Non-Hispanic[1]														
Female														
2000	37	-	-	-	1	1	8	5	2	5	3	3	2	2
1999	42	-	-	-	-	6	8	3	3	4	7	2	4	1
Non-Hispanic White														
Both sexes														
2002	169	-	-	-	3	18	29	16	11	22	16	18	9	7
2001	166	-	-	-	3	13	23	23	12	27	14	13	10	5
2000	148	-	-	-	5	14	22	20	10	12	18	8	12	8
1999	217	-	1	-	10	33	34	20	13	23	24	11	10	4
Male														
2002	132	-	-	-	2	17	23	12	9	16	14	13	8	4
2001	133	-	-	-	3	12	21	20	11	16	11	10	6	4
2000	120	-	-	-	5	13	17	16	10	7	17	5	10	6
1999	187	-	1	-	10	31	29	18	11	19	20	9	6	3
Female														
2002	37	-	-	-	1	1	6	4	2	6	2	5	1	3
2001	33	-	-	-	-	1	2	3	1	11	3	3	4	1
2000	28	-	-	-	-	1	5	4	-	5	1	3	2	2
1999	30	-	-	-	-	2	5	2	2	4	4	2	4	1
Non-Hispanic Black														
Both sexes														
2002	37	-	1	2	-	7	7	6	4	4	2	-	1	1
2001	31	-	-	2	-	7	8	4	1	-	2	1	2	3
2000	53	-	-	2	3	5	14	8	7	4	3	1	1	3
1999	65	-	1	-	1	20	15	10	4	3	3	4	-	1
Male														
2002	35	-	1	2	-	7	7	6	4	3	2	-	1	-
2001	28	-	-	2	-	7	7	3	1	-	1	1	2	3
2000	46	-	-	2	2	5	13	7	5	4	1	1	1	3
1999	55	-	1	-	1	17	12	9	3	3	1	4	-	1
Female														
2002	2	-	-	-	-	-	-	-	-	1	-	-	-	1
2001	3	-	-	-	-	-	1	1	-	-	1	-	-	-
2000	7	-	-	-	1	-	1	1	2	-	2	-	-	-
1999	10	-	-	-	-	3	3	1	1	-	2	-	-	-
Origin not stated[2]														
Both sexes														
2002	5	-	-	-	1	2	-	1	-	-	-	-	-	1
2001	2	-	-	-	-	-	-	-	1	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)										
Non-Hispanic[1]										
Female										
2000	2	2	-	1	-	-	-	-	-	-
1999	2	1	-	1	-	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	2	8	6	2	2	-	-	-	-	-
2001	3	5	4	6	4	1	-	-	-	-
2000	3	5	3	5	1	2	-	-	-	-
1999	7	7	4	6	6	4	-	-	-	-
Male										
2002	2	4	5	2	1	-	-	-	-	-
2001	3	5	2	6	2	1	-	-	-	-
2000	1	3	3	4	1	2	-	-	-	-
1999	5	6	4	5	6	4	-	-	-	-
Female										
2002	-	4	1	-	1	-	-	-	-	-
2001	-	-	2	-	2	-	-	-	-	-
2000	2	2	-	1	-	-	-	-	-	-
1999	2	1	-	1	-	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	1	1	-	-	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	1	1	-	-	-	-	-	-	-
1999	-	-	2	-	1	-	-	-	-	-
Male										
2002	-	-	1	1	-	-	-	-	-	-
2001	-	-	-	-	1	-	-	-	-	-
2000	-	1	1	-	-	-	-	-	-	-
1999	-	-	2	-	1	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Discharge of firearms, undetermined intent (Y22-Y24)														
Origin not stated[2]														
Both sexes														
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	5	-	-	-	1	2	-	1	-	-	-	-	-	1
2001	2	-	-	-	-	-	-	-	1	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)														
All origins														
Both sexes														
2002	4,587	101	56	24	38	108	300	325	457	696	770	690	421	198
2001	3,967	83	44	15	30	108	240	279	401	641	705	603	358	143
2000	3,589	65	39	16	34	94	205	274	353	581	666	534	284	142
1999	3,593	55	40	12	25	91	212	281	411	616	622	510	255	130
Male														
2002	2,915	52	27	9	24	74	213	233	313	472	470	418	255	126
2001	2,643	49	19	8	24	79	178	204	269	418	480	421	220	97
2000	2,445	27	26	11	30	68	155	197	249	403	452	368	184	95
1999	2,415	33	18	6	20	71	163	202	277	422	420	342	169	89
Female														
2002	1,672	49	29	15	14	34	87	92	144	224	300	272	166	72
2001	1,324	34	25	7	6	29	62	75	132	223	225	182	138	46
2000	1,144	38	13	5	4	26	50	77	104	178	214	166	100	47
1999	1,178	22	22	6	5	20	49	79	134	194	202	168	86	41
Hispanic														
Both sexes														
2002	297	14	11	3	1	7	30	25	42	50	37	34	15	6
2001	262	7	6	1	2	9	17	28	31	45	43	29	22	4
2000	195	5	5	-	1	4	17	26	31	39	23	19	6	7
1999	218	6	8	1	2	11	23	30	36	34	24	16	10	7

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Discharge of firearms, undetermined intent (Y22-Y24)										
Origin not stated[2]										
Both sexes										
2000	-	-	-	-	-	-	-	-	-	1
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	1	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	1
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
All origins										
Both sexes										
2002	110	72	52	41	49	39	17	8	1	14
2001	70	52	37	47	50	21	11	9	2	18
2000	71	48	45	38	35	24	25	5	1	10
1999	78	63	36	45	38	33	18	2	1	19
Male										
2002	70	42	31	21	21	20	6	5	-	13
2001	38	34	22	22	26	11	6	2	-	16
2000	46	33	28	25	19	12	5	2	-	10
1999	48	39	19	26	17	16	4	1	-	13
Female										
2002	40	30	21	20	28	19	11	3	1	1
2001	32	18	15	25	24	10	5	7	2	2
2000	25	15	17	13	16	12	20	3	1	-
1999	30	24	17	19	21	17	14	1	1	6
Hispanic										
Both sexes										
2002	9	4	5	1	1	-	-	-	-	2
2001	4	4	2	1	1	2	2	-	-	2
2000	5	3	1	-	2	-	-	-	-	1
1999	5	-	1	2	1	-	-	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)														
Hispanic														
Male														
2002	221	6	7	1	-	6	26	19	33	38	24	28	12	5
2001	200	4	3	1	2	7	12	23	26	33	38	18	16	4
2000	148	2	2	-	1	3	14	20	27	30	16	16	4	6
1999	174	3	5	1	1	9	21	25	27	27	20	14	8	6
Female														
2002	76	8	4	2	1	1	4	6	9	12	13	6	3	1
2001	62	3	3	-	-	2	5	5	5	12	5	11	6	-
2000	47	3	3	-	-	1	3	6	4	9	7	3	2	1
1999	44	3	3	-	1	2	2	5	9	7	4	2	2	1
Non-Hispanic[1]														
Both sexes														
2002	4,255	87	43	21	37	100	269	298	412	643	730	652	405	190
2001	3,668	76	38	14	28	98	223	250	368	593	659	572	329	135
2000	3,362	58	34	16	33	90	188	246	320	537	640	513	276	131
1999	3,337	49	32	11	23	80	187	246	372	580	596	491	240	121
Male														
2002	2,663	46	20	8	24	68	186	212	277	432	443	386	242	119
2001	2,414	45	16	7	22	71	166	180	241	382	440	403	200	89
2000	2,273	25	24	11	29	65	141	175	220	371	434	350	178	87
1999	2,212	30	13	5	19	62	140	173	247	393	398	326	157	81
Female														
2002	1,592	41	23	13	13	32	83	86	135	211	287	266	163	71
2001	1,254	31	22	7	6	27	57	70	127	211	219	169	129	46
2000	1,089	33	10	5	4	25	47	71	100	166	206	163	98	44
1999	1,125	19	19	6	4	18	47	73	125	187	198	165	83	40
Non-Hispanic White														
Both sexes														
2002	3,458	43	33	13	28	89	234	246	334	524	598	526	320	163
2001	2,989	40	25	8	24	90	192	206	308	472	541	465	261	116
2000	2,675	35	22	9	23	79	156	202	258	436	499	407	220	97
1999	2,631	30	14	9	19	73	150	198	300	457	456	379	178	98
Male														
2002	2,133	23	16	4	18	60	164	175	223	343	366	308	175	98
2001	1,921	25	10	3	19	66	139	145	201	296	350	317	152	73
2000	1,780	16	15	6	20	57	114	142	173	302	335	272	138	60
1999	1,700	19	7	4	16	59	113	132	196	300	307	238	110	59

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
Hispanic										
Male										
2002	9	2	2	-	1	-	-	-	-	2
2001	2	3	2	-	1	2	1	-	-	2
2000	3	1	-	-	2	-	-	-	-	1
1999	4	-	-	2	-	-	-	-	-	1
Female										
2002	-	2	3	1	-	-	-	-	-	-
2001	2	1	-	1	-	-	1	-	-	-
2000	2	2	1	-	-	-	-	-	-	-
1999	1	-	1	-	1	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	100	67	46	40	48	38	17	6	1	5
2001	64	48	34	45	49	19	9	9	2	6
2000	65	43	43	38	33	24	24	5	1	4
1999	73	61	33	43	37	32	18	1	1	10
Male										
2002	60	39	28	21	20	19	6	3	-	4
2001	35	31	19	21	25	9	5	2	-	5
2000	42	30	27	25	17	12	4	2	-	4
1999	44	38	17	24	17	16	4	-	-	8
Female										
2002	40	28	18	19	28	19	11	3	1	1
2001	29	17	15	24	24	10	4	7	2	1
2000	23	13	16	13	16	12	20	3	1	-
1999	29	23	16	19	20	16	14	1	1	2
Non-Hispanic White										
Both sexes										
2002	81	53	38	34	39	36	16	5	1	4
2001	52	36	30	40	43	17	9	8	2	4
2000	53	34	34	33	31	20	20	5	-	2
1999	63	53	28	39	35	28	16	1	1	6
Male										
2002	47	28	20	17	18	18	6	3	-	3
2001	26	23	15	19	24	8	5	2	-	3
2000	34	22	20	23	15	9	3	2	-	2
1999	34	33	14	21	15	14	4	-	-	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)														
Non-Hispanic White														
Female														
2002	1,325	20	17	9	10	29	70	71	111	181	232	218	145	65
2001	1,068	15	15	5	5	24	53	61	107	176	191	148	109	43
2000	895	19	7	3	3	22	42	60	85	134	164	135	82	37
1999	931	11	7	5	3	14	37	66	104	157	149	141	68	39
Non-Hispanic Black														
Both sexes														
2002	701	37	6	7	6	7	31	45	63	111	123	116	78	22
2001	598	36	13	5	4	5	23	33	52	110	110	97	63	17
2000	601	14	11	7	9	7	24	36	52	93	130	99	52	28
1999	621	18	14	2	4	5	30	41	64	110	123	103	59	19
Male														
2002	469	18	2	3	6	5	20	32	42	84	72	74	61	17
2001	439	20	6	3	3	3	20	27	34	79	85	78	46	15
2000	432	3	8	5	8	7	20	27	39	63	92	74	37	21
1999	459	10	6	1	3	3	23	35	44	83	83	82	46	18
Female														
2002	232	19	4	4	-	2	11	13	21	27	51	42	17	5
2001	159	16	7	2	1	2	3	6	18	31	25	19	17	2
2000	169	11	3	2	1	-	4	9	13	30	38	25	15	7
1999	162	8	8	1	1	2	7	6	20	27	40	21	13	1
Origin not stated[2]														
Both sexes														
2002	35	-	2	-	-	1	1	2	3	3	3	4	1	2
2001	37	-	-	-	-	1	-	1	2	3	3	2	7	4
2000	32	2	-	-	-	-	-	2	2	5	3	2	2	4
1999	38	-	-	-	-	-	2	5	3	2	2	3	5	2
Male														
2002	31	-	-	-	-	-	1	2	3	2	3	4	1	2
2001	29	-	-	-	-	1	-	1	2	3	2	-	4	4
2000	24	-	-	-	-	-	-	2	2	2	2	2	2	2
1999	29	-	-	-	-	-	2	4	3	2	2	2	4	2
Female														
2002	4	-	2	-	-	1	-	-	-	1	-	-	-	-
2001	8	-	-	-	-	-	-	-	-	-	1	2	3	-
2000	8	2	-	-	-	-	-	-	-	3	1	-	-	2
1999	9	-	-	-	-	-	-	1	-	-	-	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)										
Non-Hispanic White										
Female										
2002	34	25	18	17	21	18	10	2	1	1
2001	26	13	15	21	19	9	4	6	2	1
2000	19	12	14	10	16	11	17	3	-	-
1999	29	20	14	18	20	14	12	1	1	1
Non-Hispanic Black										
Both sexes										
2002	16	9	6	6	9	1	1	1	-	-
2001	10	10	2	2	2	2	-	1	-	1
2000	11	7	6	4	2	3	3	-	1	2
1999	8	7	3	4	-	3	-	-	-	4
Male										
2002	13	8	6	4	2	-	-	-	-	-
2001	8	8	2	-	-	1	-	-	-	1
2000	8	6	5	2	2	2	1	-	-	2
1999	8	5	2	3	-	1	-	-	-	3
Female										
2002	3	1	-	2	7	1	1	1	-	-
2001	2	2	-	2	2	1	-	1	-	-
2000	3	1	1	2	-	1	2	-	1	-
1999	-	2	1	1	-	2	-	-	-	1
Origin not stated[2]										
Both sexes										
2002	1	1	1	-	-	1	-	2	-	7
2001	2	-	1	1	-	-	-	-	-	10
2000	1	2	1	-	-	-	1	-	-	5
1999	-	2	2	-	-	1	-	1	-	8
Male										
2002	1	1	1	-	-	1	-	2	-	7
2001	1	-	1	1	-	-	-	-	-	9
2000	1	2	1	-	-	-	1	-	-	5
1999	-	1	2	-	-	-	-	1	-	4
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	1
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	1	-	-	-	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Operations of war and their sequelae (Y36,Y89.1)														
All origins														
Both sexes														
2002	20	-	-	-	-	-	-	-	1	-	1	1	-	2
2001	17	-	-	-	-	-	1	-	-	1	-	1	1	-
2000	16	-	-	-	-	-	-	-	-	-	-	-	2	-
1999	23	-	-	-	-	-	-	-	-	1	-	-	2	1
Male														
2002	20	-	-	-	-	-	-	-	1	-	1	1	-	2
2001	16	-	-	-	-	-	1	-	-	1	-	1	1	-
2000	16	-	-	-	-	-	-	-	-	-	-	-	2	-
1999	22	-	-	-	-	-	-	-	-	1	-	-	2	1
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic														
Both sexes														
2002	1	-	-	-	-	-	-	-	1	-	-	-	-	-
2001	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	1	-	-	-	-	-	-	-	1	-	-	-	-	-
2001	2	-	-	-	-	-	1	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	2	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]														
Both sexes														
2002	19	-	-	-	-	-	-	-	-	-	1	1	-	2
2001	15	-	-	-	-	-	-	-	-	1	-	1	1	-
2000	15	-	-	-	-	-	-	-	-	-	-	-	2	-
1999	21	-	-	-	-	-	-	-	-	1	-	-	2	1
Male														
2002	19	-	-	-	-	-	-	-	-	-	1	1	-	2
2001	14	-	-	-	-	-	-	-	-	1	-	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)										
All origins										
Both sexes										
2002	1	2	1	4	6	1	-	-	-	-
2001	-	-	4	7	1	1	-	-	-	-
2000	1	1	2	4	5	1	-	-	-	-
1999	-	2	6	4	4	2	1	-	-	-
Male										
2002	1	2	1	4	6	1	-	-	-	-
2001	-	-	4	6	1	1	-	-	-	-
2000	1	1	2	4	5	1	-	-	-	-
1999	-	2	6	4	3	2	1	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Hispanic										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	1	1	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	1	-	-	-	-	-	-	-	-
1999	-	-	-	-	1	1	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1	2	1	4	6	1	-	-	-	-
2001	-	-	4	6	1	1	-	-	-	-
2000	1	-	2	4	5	1	-	-	-	-
1999	-	2	6	4	3	1	1	-	-	-
Male										
2002	1	2	1	4	6	1	-	-	-	-
2001	-	-	4	5	1	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Operations of war and their sequelae (Y36,Y89.1)														
Non-Hispanic[1]														
Male														
2000	15	-	-	-	-	-	-	-	-	-	-	-	2	-
1999	20	-	-	-	-	-	-	-	-	1	-	-	2	1
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic White														
Both sexes														
2002	14	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	14	-	-	-	-	-	-	-	-	1	-	1	1	-
2000	14	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	20	-	-	-	-	-	-	-	-	1	-	-	2	1
Male														
2002	14	-	-	-	-	-	-	-	-	-	-	-	-	1
2001	13	-	-	-	-	-	-	-	-	1	-	1	1	-
2000	14	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	19	-	-	-	-	-	-	-	-	1	-	-	2	1
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic Black														
Both sexes														
2002	5	-	-	-	-	-	-	-	-	-	1	1	-	1
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	5	-	-	-	-	-	-	-	-	-	1	1	-	1
2001	1	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	1	-	-	-	-	-	-	-	-	-	-	-	1	-
1999	1	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)										
Non-Hispanic[1]										
Male										
2000	1	-	2	4	5	1	-	-	-	-
1999	-	2	6	4	2	1	1	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Non-Hispanic White										
Both sexes										
2002	1	2	-	4	5	1	-	-	-	-
2001	-	-	3	6	1	1	-	-	-	-
2000	1	-	2	4	5	1	-	-	-	-
1999	-	1	6	4	3	1	1	-	-	-
Male										
2002	1	2	-	4	5	1	-	-	-	-
2001	-	-	3	5	1	1	-	-	-	-
2000	1	-	2	4	5	1	-	-	-	-
1999	-	1	6	4	2	1	1	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	1	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Non-Hispanic Black										
Both sexes										
2002	-	-	1	-	1	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	-	-	-	-	-
Male										
2002	-	-	1	-	1	-	-	-	-	-
2001	-	-	1	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	1	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Operations of war and their sequelae (Y36,Y89.1)														
Non-Hispanic Black														
Female														
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]														
Both sexes														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Male														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Female														
2002	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Complications of medical and surgical care (Y40-Y84,Y88)														
All origins														
Both sexes														
2002	2,843	15	19	13	22	16	19	27	52	73	98	114	151	164
2001	3,021	26	24	5	9	18	25	24	55	71	101	142	141	178
2000	3,059	19	22	18	13	10	22	37	48	65	84	128	167	157
1999	2,823	54	30	7	11	17	19	29	49	66	86	91	123	138
Male														
2002	1,236	8	11	8	18	10	9	16	31	35	47	60	67	84
2001	1,357	15	14	2	6	6	13	11	31	39	44	75	68	91
2000	1,430	12	11	11	4	5	11	23	24	33	43	59	84	77
1999	1,308	31	12	4	5	8	13	9	34	37	41	37	61	77
Female														
2002	1,607	7	8	5	4	6	10	11	21	38	51	54	84	80
2001	1,664	11	10	3	3	12	12	13	24	32	57	67	73	87
2000	1,629	7	11	7	9	5	11	14	24	32	41	69	83	80
1999	1,515	23	18	3	6	9	6	20	15	29	45	54	62	61

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Operations of war and their sequelae (Y36,Y89.1)										
Non-Hispanic Black										
Female										
1999	-	-	-	-	-	-	-	-	-	-
Origin not stated[2]										
Both sexes										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Male										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	-	-	-	-	-	-	-	-	-
1999	-	-	-	-	-	-	-	-	-	-
Complications of medical and surgical care (Y40-Y84,Y88)										
All origins										
Both sexes										
2002	187	225	335	382	410	288	167	59	7	-
2001	206	247	320	439	425	315	186	56	7	1
2000	219	263	384	427	398	337	176	54	11	-
1999	177	229	357	401	410	312	153	52	12	-
Male										
2002	85	110	146	166	154	110	40	18	3	-
2001	105	116	149	192	191	119	55	13	2	-
2000	104	123	208	212	190	134	48	13	1	-
1999	88	102	177	193	182	128	49	17	3	-
Female										
2002	102	115	189	216	256	178	127	41	4	-
2001	101	131	171	247	234	196	131	43	5	1
2000	115	140	176	215	208	203	128	41	10	-
1999	89	127	180	208	228	184	104	35	9	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Complications of medical and surgical care (Y40-Y84,Y88)														
Hispanic														
Both sexes														
2002	137	2	7	1	4	5	1	4	4	7	6	4	10	8
2001	145	3	6	1	3	4	4	5	2	4	8	14	8	5
2000	136	3	4	3	3	2	5	2	2	5	6	8	5	4
1999	100	5	6	2	1	2	2	2	6	6	2	5	6	6
Male														
2002	57	2	4	-	4	3	-	1	-	3	2	3	5	2
2001	65	-	3	1	2	1	2	2	1	2	2	9	2	1
2000	70	2	2	2	2	1	5	1	1	2	3	3	1	4
1999	52	3	3	1	1	1	1	-	3	5	2	2	4	4
Female														
2002	80	-	3	1	-	2	1	3	4	4	4	1	5	6
2001	80	3	3	-	1	3	2	3	1	2	6	5	6	4
2000	66	1	2	1	1	1	-	1	1	3	3	5	4	-
1999	48	2	3	1	-	1	1	2	3	1	-	3	2	2
Non-Hispanic[1]														
Both sexes														
2002	2,695	13	12	12	18	11	18	23	48	66	92	109	140	156
2001	2,861	23	18	4	6	13	21	19	53	67	91	128	133	173
2000	2,914	16	18	15	10	8	17	35	46	60	77	120	161	153
1999	2,714	49	24	5	10	15	17	27	43	59	82	86	116	131
Male														
2002	1,175	6	7	8	14	7	9	15	31	32	45	56	61	82
2001	1,286	15	11	1	4	4	11	9	30	37	41	66	66	90
2000	1,354	10	9	9	2	4	6	22	23	31	39	56	83	73
1999	1,253	28	9	3	4	7	12	9	31	32	38	35	56	73
Female														
2002	1,520	7	5	4	4	4	9	8	17	34	47	53	79	74
2001	1,575	8	7	3	2	9	10	10	23	30	50	62	67	83
2000	1,560	6	9	6	8	4	11	13	23	29	38	64	78	80
1999	1,461	21	15	2	6	8	5	18	12	27	44	51	60	58
Non-Hispanic White														
Both sexes														
2002	2,188	6	7	6	10	8	13	16	33	45	62	75	94	114
2001	2,294	14	10	2	4	8	17	12	39	43	62	86	99	132
2000	2,378	12	8	9	8	6	11	20	36	47	56	89	129	116
1999	2,212	30	16	4	7	13	13	14	33	45	58	64	84	91
Male														
2002	966	3	5	4	8	6	7	11	23	25	35	41	44	63
2001	1,016	9	7	-	2	3	9	5	21	22	28	44	47	65

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)										
Hispanic										
Both sexes										
2002	11	14	14	16	13	2	3	-	1	-
2001	8	17	8	18	11	5	8	2	-	1
2000	7	22	13	15	11	7	6	3	-	-
1999	10	8	7	7	5	3	7	1	1	-
Male										
2002	4	5	6	9	4	-	-	-	-	-
2001	7	6	2	8	6	2	4	2	-	-
2000	4	7	9	9	6	4	1	1	-	-
1999	5	4	1	6	2	1	2	1	-	-
Female										
2002	7	9	8	7	9	2	3	-	1	-
2001	1	11	6	10	5	3	4	-	-	1
2000	3	15	4	6	5	3	5	2	-	-
1999	5	4	6	1	3	2	5	-	1	-
Non-Hispanic[1]										
Both sexes										
2002	176	209	321	365	393	285	163	59	6	-
2001	196	228	309	419	412	310	177	54	7	-
2000	210	241	369	411	386	329	170	51	11	-
1999	167	220	350	394	404	308	145	51	11	-
Male										
2002	81	104	140	157	149	110	40	18	3	-
2001	98	108	146	184	184	117	51	11	2	-
2000	99	116	197	202	183	130	47	12	1	-
1999	83	98	176	187	179	127	47	16	3	-
Female										
2002	95	105	181	208	244	175	123	41	3	-
2001	98	120	163	235	228	193	126	43	5	-
2000	111	125	172	209	203	199	123	39	10	-
1999	84	122	174	207	225	181	98	35	8	-
Non-Hispanic White										
Both sexes										
2002	138	162	278	315	349	258	142	51	6	-
2001	147	180	237	367	368	262	151	48	6	-
2000	156	193	305	346	345	289	149	38	10	-
1999	133	185	293	337	351	262	129	43	7	-
Male										
2002	66	78	122	138	138	97	35	14	3	-
2001	72	88	109	163	170	101	41	9	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Complications of medical and surgical care (Y40-Y84,Y88)														
Non-Hispanic White														
Male														
2000	1,135	9	6	5	2	3	4	12	19	25	33	42	69	60
1999	1,025	19	8	2	3	6	9	4	25	25	27	24	41	51
Female														
2002	1,222	3	2	2	2	2	6	5	10	20	27	34	50	51
2001	1,278	5	3	2	2	5	8	7	18	21	34	42	52	67
2000	1,243	3	2	4	6	3	7	8	17	22	23	47	60	56
1999	1,187	11	8	2	4	7	4	10	8	20	31	40	43	40
Non-Hispanic Black														
Both sexes														
2002	445	7	5	6	8	3	2	6	11	19	29	31	44	38
2001	515	9	7	1	2	3	4	7	14	22	26	39	31	38
2000	484	4	8	4	1	2	5	13	10	13	18	27	30	35
1999	442	18	8	1	3	2	4	10	8	13	22	20	31	34
Male														
2002	182	3	2	4	6	1	1	4	5	6	10	13	16	18
2001	247	6	4	-	2	-	2	4	9	14	12	20	18	22
2000	192	1	2	2	-	1	1	9	4	6	6	12	13	12
1999	195	8	1	1	1	1	3	4	6	6	10	11	15	16
Female														
2002	263	4	3	2	2	2	1	2	6	13	19	18	28	20
2001	268	3	3	1	-	3	2	3	5	8	14	19	13	16
2000	292	3	6	2	1	1	4	4	6	7	12	15	17	23
1999	247	10	7	-	2	1	1	6	2	7	12	9	16	18
Origin not stated[2]														
Both sexes														
2002	11	-	-	-	-	-	-	-	-	-	-	1	1	-
2001	15	-	-	-	-	1	-	-	-	-	2	-	-	-
2000	9	-	-	-	-	-	-	-	-	-	1	-	1	-
1999	9	-	-	-	-	-	-	-	-	1	2	-	1	1
Male														
2002	4	-	-	-	-	-	-	-	-	-	-	1	1	-
2001	6	-	-	-	-	1	-	-	-	-	1	-	-	-
2000	6	-	-	-	-	-	-	-	-	-	1	-	-	-
1999	3	-	-	-	-	-	-	-	-	-	1	-	1	-
Female														
2002	7	-	-	-	-	-	-	-	-	-	-	-	-	-
2001	9	-	-	-	-	-	-	-	-	-	1	-	-	-
2000	3	-	-	-	-	-	-	-	-	-	-	-	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)										
Non-Hispanic White										
Male										
2000	75	91	164	177	170	115	43	10	1	-
1999	66	83	147	164	157	107	41	15	1	-
Female										
2002	72	84	156	177	211	161	107	37	3	-
2001	75	92	128	204	198	161	110	39	5	-
2000	81	102	141	169	175	174	106	28	9	-
1999	67	102	146	173	194	155	88	28	6	-
Non-Hispanic Black										
Both sexes										
2002	32	38	39	44	36	21	19	7	-	-
2001	46	42	62	46	40	46	23	6	1	-
2000	51	42	59	58	37	34	20	12	1	-
1999	27	31	53	52	42	37	16	7	3	-
Male										
2002	12	20	17	14	11	10	5	4	-	-
2001	25	20	33	17	13	14	9	2	1	-
2000	22	24	28	21	12	12	3	1	-	-
1999	14	13	27	21	17	12	6	1	1	-
Female										
2002	20	18	22	30	25	11	14	3	-	-
2001	21	22	29	29	27	32	14	4	-	-
2000	29	18	31	37	25	22	17	11	1	-
1999	13	18	26	31	25	25	10	6	2	-
Origin not stated[2]										
Both sexes										
2002	-	2	-	1	4	1	1	-	-	-
2001	2	2	3	2	2	-	1	-	-	-
2000	2	-	2	1	1	1	-	-	-	-
1999	-	1	-	-	1	1	1	-	-	-
Male										
2002	-	1	-	-	1	-	-	-	-	-
2001	-	2	1	-	1	-	-	-	-	-
2000	1	-	2	1	1	-	-	-	-	-
1999	-	-	-	-	1	-	-	-	-	-
Female										
2002	-	1	-	1	3	1	1	-	-	-
2001	2	-	2	2	1	-	1	-	-	-
2000	1	-	-	-	-	1	-	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Complications of medical and surgical care (Y40-Y84,Y88)														
Origin not stated[2]														
Female														
1999	6	-	-	-	-	-	-	-	-	1	1	-	-	1
Alcohol-induced causes (F10,G31.2, G62.1,I42.6,K29.2,K70,R78.0,X45, X65,Y15)[4]														
All origins														
Both sexes														
2002	19,928	1	2	-	3	33	68	154	427	1,170	2,409	3,259	3,384	2,697
2001	19,817	2	-	-	2	41	64	161	463	1,305	2,381	3,372	3,255	2,459
2000	19,358	3	1	-	4	33	60	166	467	1,322	2,427	3,157	2,909	2,423
1999	19,171	-	-	-	2	25	73	140	475	1,382	2,373	3,103	2,808	2,340
Male														
2002	15,036	-	1	-	2	28	55	128	311	813	1,714	2,435	2,652	2,092
2001	14,923	2	-	-	-	31	56	119	330	920	1,705	2,549	2,484	1,940
2000	14,770	2	-	-	4	27	47	138	350	958	1,778	2,439	2,293	1,851
1999	14,665	-	-	-	1	20	58	115	323	967	1,791	2,404	2,232	1,853
Female														
2002	4,892	1	1	-	1	5	13	26	116	357	695	824	732	605
2001	4,894	-	-	-	2	10	8	42	133	385	676	823	771	519
2000	4,588	1	1	-	-	6	13	28	117	364	649	718	616	572
1999	4,506	-	-	-	1	5	15	25	152	415	582	699	576	487
Hispanic														
Both sexes														
2002	2,381	-	-	-	1	11	22	47	93	186	314	383	384	303
2001	2,363	-	-	-	-	15	16	51	90	184	304	411	351	286
2000	2,290	-	-	-	1	7	24	41	90	199	319	361	327	287
1999	2,166	-	-	-	1	6	20	32	95	204	296	357	309	253
Male														
2002	2,040	-	-	-	-	8	22	44	83	156	269	328	328	266
2001	2,009	-	-	-	-	10	15	40	77	163	249	353	296	247
2000	1,994	-	-	-	1	6	22	39	85	166	267	314	283	254
1999	1,849	-	-	-	1	4	20	30	80	171	252	308	265	221
Female														
2002	341	-	-	-	1	3	-	3	10	30	45	55	56	37
2001	354	-	-	-	-	5	1	11	13	21	55	58	55	39
2000	296	-	-	-	-	1	2	2	5	33	52	47	44	33
1999	317	-	-	-	-	2	-	2	15	33	44	49	44	32

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Complications of medical and surgical care (Y40-Y84,Y88)										
Origin not stated[2]										
Female										
1999	-	1	-	-	-	1	1	-	-	-
Alcohol-induced causes (F10,G31.2, G62.1,I42.6,K29.2,K70,R78.0,X45, X65,Y15)[4]										
All origins										
Both sexes										
2002	2,123	1,569	1,239	800	383	153	38	8	-	8
2001	2,071	1,621	1,190	808	387	165	39	14	1	16
2000	2,066	1,631	1,250	819	389	158	55	7	3	8
1999	2,042	1,657	1,235	866	413	174	43	11	1	8
Male										
2002	1,686	1,228	908	571	270	104	25	5	-	8
2001	1,614	1,240	916	587	266	114	26	8	1	15
2000	1,602	1,264	961	634	271	106	32	4	1	8
1999	1,579	1,306	943	627	298	114	24	3	-	7
Female										
2002	437	341	331	229	113	49	13	3	-	-
2001	457	381	274	221	121	51	13	6	-	1
2000	464	367	289	185	118	52	23	3	2	-
1999	463	351	292	239	115	60	19	8	1	1
Hispanic										
Both sexes										
2002	215	181	134	61	26	12	5	2	-	1
2001	233	175	120	79	34	8	2	-	-	4
2000	215	194	115	72	24	7	3	1	-	3
1999	214	162	106	64	26	17	3	-	-	1
Male										
2002	185	158	114	45	20	7	5	1	-	1
2001	200	157	104	61	24	7	2	-	-	4
2000	184	177	100	65	21	4	2	1	-	3
1999	183	137	93	48	21	13	1	-	-	1
Female										
2002	30	23	20	16	6	5	-	1	-	-
2001	33	18	16	18	10	1	-	-	-	-
2000	31	17	15	7	3	3	1	-	-	-
1999	31	25	13	16	5	4	2	-	-	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcohol-induced causes (F10,G31.2, G62.1,I42.6,K29.2,K70,R78.0,X45, X65,Y15)[4]														
Non-Hispanic[1]														
Both sexes														
2002	17,401	1	2	-	2	22	45	106	332	971	2,080	2,847	2,977	2,372
2001	17,315	2	-	-	2	26	48	110	369	1,116	2,063	2,942	2,872	2,150
2000	16,929	3	1	-	3	26	35	124	374	1,113	2,093	2,768	2,555	2,124
1999	16,863	-	-	-	1	19	51	107	376	1,166	2,056	2,725	2,474	2,074
Male														
2002	12,870	-	1	-	2	20	32	83	226	649	1,433	2,081	2,304	1,808
2001	12,801	2	-	-	-	21	41	79	250	753	1,443	2,182	2,161	1,675
2000	12,654	2	-	-	3	21	24	98	263	782	1,497	2,102	1,985	1,587
1999	12,691	-	-	-	-	16	36	84	239	786	1,522	2,077	1,946	1,621
Female														
2002	4,531	1	1	-	-	2	13	23	106	322	647	766	673	564
2001	4,514	-	-	-	2	5	7	31	119	363	620	760	711	475
2000	4,275	1	1	-	-	5	11	26	111	331	596	666	570	537
1999	4,172	-	-	-	1	3	15	23	137	380	534	648	528	453
Non-Hispanic White														
Both sexes														
2002	14,297	-	1	-	1	20	31	74	247	753	1,708	2,337	2,426	1,926
2001	13,991	1	-	-	2	17	29	71	279	880	1,615	2,388	2,239	1,709
2000	13,645	3	-	-	2	21	23	88	269	869	1,636	2,185	2,032	1,692
1999	13,434	-	-	-	1	11	37	84	280	884	1,574	2,116	1,965	1,633
Male														
2002	10,625	-	1	-	1	19	23	58	172	507	1,191	1,721	1,897	1,477
2001	10,351	1	-	-	-	15	25	55	192	587	1,128	1,789	1,709	1,321
2000	10,280	2	-	-	2	17	17	72	190	620	1,182	1,693	1,593	1,276
1999	10,185	-	-	-	-	9	27	68	181	603	1,186	1,638	1,562	1,271
Female														
2002	3,672	-	-	-	-	1	8	16	75	246	517	616	529	449
2001	3,640	-	-	-	2	2	4	16	87	293	487	599	530	388
2000	3,365	1	-	-	-	4	6	16	79	249	454	492	439	416
1999	3,249	-	-	-	1	2	10	16	99	281	388	478	403	362
Non-Hispanic Black														
Both sexes														
2002	2,336	1	1	-	-	-	8	16	48	123	255	386	442	364
2001	2,598	1	-	-	-	5	12	19	47	144	336	431	524	365
2000	2,598	-	1	-	1	2	5	23	66	154	339	485	438	347
1999	2,719	-	-	-	-	4	6	12	54	197	344	495	413	371
Male														
2002	1,720	-	-	-	-	-	6	14	33	80	167	275	330	271

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcohol-induced causes (F10,G31.2, G62.1,I42.6,K29.2,K70,R78.0,X45, X65,Y15)[4]										
Non-Hispanic[1]										
Both sexes										
2002	1,894	1,377	1,100	736	354	140	33	6	-	4
2001	1,825	1,433	1,063	725	353	157	37	14	1	7
2000	1,837	1,423	1,130	740	365	151	51	6	3	4
1999	1,810	1,486	1,120	797	387	157	40	11	1	5
Male										
2002	1,488	1,059	789	523	248	96	20	4	-	4
2001	1,403	1,074	806	523	242	107	24	8	1	6
2000	1,406	1,076	857	562	250	102	29	3	1	4
1999	1,379	1,160	842	574	277	101	23	3	-	5
Female										
2002	406	318	311	213	106	44	13	2	-	-
2001	422	359	257	202	111	50	13	6	-	1
2000	431	347	273	178	115	49	22	3	2	-
1999	431	326	278	223	110	56	17	8	1	-
Non-Hispanic White										
Both sexes										
2002	1,543	1,162	940	654	312	123	30	6	-	3
2001	1,509	1,175	931	639	322	133	32	14	1	5
2000	1,491	1,183	979	656	326	135	45	5	3	2
1999	1,457	1,226	950	687	343	134	35	11	1	5
Male										
2002	1,216	886	666	466	218	81	18	4	-	3
2001	1,157	867	700	460	221	91	20	8	1	4
2000	1,142	893	739	496	226	91	24	2	1	2
1999	1,127	946	709	494	250	84	22	3	-	5
Female										
2002	327	276	274	188	94	42	12	2	-	-
2001	352	308	231	179	101	42	12	6	-	1
2000	349	290	240	160	100	44	21	3	2	-
1999	330	280	241	193	93	50	13	8	1	-
Non-Hispanic Black										
Both sexes										
2002	275	170	127	69	34	14	2	-	-	1
2001	264	222	106	74	27	16	3	-	-	2
2000	278	211	127	71	28	13	6	1	-	2
1999	301	220	150	90	39	19	4	-	-	-
Male										
2002	217	140	100	48	25	12	1	-	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Alcohol-induced causes (F10,G31.2, G62.1,I42.6,K29.2,K70,R78.0,X45, X65,Y15)[4]														
Non-Hispanic Black														
Male														
2001	1,957	1	-	-	-	3	10	14	29	108	238	315	386	292
2000	1,901	-	-	-	1	2	4	19	41	104	232	343	332	257
1999	2,008	-	-	-	-	4	4	8	32	131	235	358	314	299
Female														
2002	616	1	1	-	-	-	2	2	15	43	88	111	112	93
2001	641	-	-	-	-	2	2	5	18	36	98	116	138	73
2000	697	-	1	-	-	-	1	4	25	50	107	142	106	90
1999	711	-	-	-	-	-	2	4	22	66	109	137	99	72
Origin not stated[2]														
Both sexes														
2002	146	-	-	-	-	-	1	1	2	13	15	29	23	22
2001	139	-	-	-	-	-	-	-	4	5	14	19	32	23
2000	139	-	-	-	-	-	1	1	3	10	15	28	27	12
1999	142	-	-	-	-	-	2	1	4	12	21	21	25	13
Male														
2002	126	-	-	-	-	-	1	1	2	8	12	26	20	18
2001	113	-	-	-	-	-	-	-	3	4	13	14	27	18
2000	122	-	-	-	-	-	1	1	2	10	14	23	25	10
1999	125	-	-	-	-	-	2	1	4	10	17	19	21	11
Female														
2002	20	-	-	-	-	-	-	-	-	5	3	3	3	4
2001	26	-	-	-	-	-	-	-	1	1	1	5	5	5
2000	17	-	-	-	-	-	-	-	1	-	1	5	2	2
1999	17	-	-	-	-	-	-	-	-	2	4	2	4	2
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]														
All origins														
Both sexes														
2002	26,018	29	30	7	35	625	1,590	1,850	2,599	3,837	5,020	4,450	2,731	1,234
2001	21,683	22	26	7	37	538	1,274	1,552	2,187	3,448	4,219	3,660	2,133	901
2000	19,698	18	15	9	35	452	1,115	1,450	2,076	3,270	3,935	3,170	1,814	793
1999	19,102	21	25	7	23	367	983	1,452	2,127	3,240	3,850	3,042	1,581	760

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcohol-induced causes (F10,G31.2, G62.1,I42.6,K29.2,K70,R78.0,X45, X65,Y15)[4]										
Non-Hispanic Black										
Male										
2001	208	181	86	55	18	9	2	-	-	2
2000	209	161	103	57	19	9	5	1	-	2
1999	220	183	117	64	25	13	1	-	-	-
Female										
2002	58	30	27	21	9	2	1	-	-	-
2001	56	41	20	19	9	7	1	-	-	-
2000	69	50	24	14	9	4	1	-	-	-
1999	81	37	33	26	14	6	3	-	-	-
Origin not stated[2]										
Both sexes										
2002	14	11	5	3	3	1	-	-	-	3
2001	13	13	7	4	-	-	-	-	-	5
2000	14	14	5	7	-	-	1	-	-	1
1999	18	9	9	5	-	-	-	-	-	2
Male										
2002	13	11	5	3	2	1	-	-	-	3
2001	11	9	6	3	-	-	-	-	-	5
2000	12	11	4	7	-	-	1	-	-	1
1999	17	9	8	5	-	-	-	-	-	1
Female										
2002	1	-	-	-	1	-	-	-	-	-
2001	2	4	1	1	-	-	-	-	-	-
2000	2	3	1	-	-	-	-	-	-	-
1999	1	-	1	-	-	-	-	-	-	1
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]										
All origins										
Both sexes										
2002	591	362	285	261	216	157	83	15	1	10
2001	455	294	255	253	182	134	69	17	4	16
2000	393	242	238	230	194	146	74	15	-	14
1999	399	301	252	257	199	131	49	17	1	18

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]														
All origins														
Male														
2002	16,724	20	9	6	18	447	1,175	1,346	1,793	2,534	3,194	2,764	1,752	724
2001	14,244	14	21	2	19	386	964	1,142	1,498	2,302	2,775	2,440	1,352	530
2000	13,125	12	13	6	19	318	848	1,057	1,458	2,242	2,647	2,170	1,187	433
1999	12,873	14	12	1	15	251	729	1,067	1,512	2,237	2,666	2,121	1,012	451
Female														
2002	9,294	9	21	1	17	178	415	504	806	1,303	1,826	1,686	979	510
2001	7,439	8	5	5	18	152	310	410	689	1,146	1,444	1,220	781	371
2000	6,573	6	2	3	16	134	267	393	618	1,028	1,288	1,000	627	360
1999	6,229	7	13	6	8	116	254	385	615	1,003	1,184	921	569	309
Hispanic														
Both sexes														
2002	2,137	5	4	-	4	58	159	188	268	358	419	330	175	74
2001	1,731	-	4	-	5	54	108	176	232	330	314	231	142	66
2000	1,700	2	2	-	4	40	108	170	279	345	306	215	117	61
1999	1,965	7	4	1	2	50	122	211	303	400	341	254	144	46
Male														
2002	1,647	4	-	-	1	44	128	160	211	272	329	247	138	54
2001	1,335	-	3	-	1	39	80	146	174	259	245	178	113	53
2000	1,348	2	2	-	-	30	91	130	237	264	257	178	87	41
1999	1,605	4	4	-	2	34	108	174	257	320	286	217	103	40
Female														
2002	490	1	4	-	3	14	31	28	57	86	90	83	37	20
2001	396	-	1	-	4	15	28	30	58	71	69	53	29	13
2000	352	-	-	-	4	10	17	40	42	81	49	37	30	20
1999	360	3	-	1	-	16	14	37	46	80	55	37	41	6
Non-Hispanic[1]														
Both sexes														
2002	23,734	24	26	7	31	565	1,424	1,654	2,319	3,456	4,572	4,097	2,541	1,151
2001	19,778	22	22	7	32	482	1,159	1,372	1,941	3,091	3,876	3,396	1,968	821
2000	17,813	16	12	9	30	410	1,000	1,271	1,781	2,908	3,600	2,910	1,666	721
1999	16,940	14	21	6	21	315	858	1,233	1,813	2,814	3,460	2,755	1,412	696
Male														
2002	14,968	16	9	6	17	403	1,042	1,179	1,572	2,246	2,843	2,499	1,607	662

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]										
All origins										
Male										
2002	315	180	128	118	93	61	33	7	-	7
2001	230	163	128	116	73	44	23	5	1	16
2000	198	126	105	106	78	61	24	4	-	13
1999	236	164	120	103	79	49	16	7	-	11
Female										
2002	276	182	157	143	123	96	50	8	1	3
2001	225	131	127	137	109	90	46	12	3	-
2000	195	116	133	124	116	85	50	11	-	1
1999	163	137	132	154	120	82	33	10	1	7
Hispanic										
Both sexes										
2002	37	19	17	12	8	1	1	-	-	-
2001	22	17	12	7	5	2	-	2	-	2
2000	18	11	9	5	3	2	3	-	-	-
1999	38	17	11	5	2	1	1	-	-	5
Male										
2002	27	10	8	9	4	-	1	-	-	-
2001	16	12	6	5	1	1	-	1	-	2
2000	11	8	3	2	1	1	3	-	-	-
1999	29	16	5	1	-	1	-	-	-	4
Female										
2002	10	9	9	3	4	1	-	-	-	-
2001	6	5	6	2	4	1	-	1	-	-
2000	7	3	6	3	2	1	-	-	-	-
1999	9	1	6	4	2	-	1	-	-	1
Non-Hispanic[1]										
Both sexes										
2002	551	337	265	246	208	156	82	14	1	7
2001	426	274	242	245	177	132	69	15	4	5
2000	369	230	226	225	191	144	71	15	-	8
1999	354	281	240	252	195	128	48	17	1	6
Male										
2002	285	165	117	107	89	61	32	6	-	5

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]														
Non-Hispanic[1]														
Male														
2001	12,769	14	18	2	18	346	878	992	1,312	2,020	2,510	2,236	1,222	466
2000	11,644	10	10	6	18	287	751	923	1,210	1,965	2,368	1,963	1,077	384
1999	11,124	10	8	1	13	216	618	886	1,246	1,898	2,347	1,879	889	399
Female														
2002	8,766	8	17	1	14	162	382	475	747	1,210	1,729	1,598	934	489
2001	7,009	8	4	5	14	136	281	380	629	1,071	1,366	1,160	746	355
2000	6,169	6	2	3	12	123	249	348	571	943	1,232	947	589	337
1999	5,816	4	13	5	8	99	240	347	567	916	1,113	876	523	297
Non-Hispanic White														
Both sexes														
2002	19,929	9	14	4	28	514	1,305	1,444	1,985	2,894	3,827	3,329	1,998	936
2001	16,349	10	13	5	27	436	1,048	1,191	1,644	2,564	3,164	2,666	1,521	671
2000	14,568	9	5	6	27	376	890	1,083	1,471	2,386	2,877	2,265	1,265	584
1999	13,624	6	10	3	19	282	736	1,020	1,473	2,288	2,710	2,106	1,073	556
Male														
2002	12,468	7	3	3	16	368	957	1,030	1,363	1,883	2,393	2,017	1,212	502
2001	10,456	4	12	2	15	324	788	859	1,133	1,688	2,060	1,719	894	353
2000	9,431	5	3	5	16	267	677	784	1,006	1,633	1,891	1,499	784	284
1999	8,822	5	4	-	12	192	534	727	1,023	1,547	1,851	1,404	636	289
Female														
2002	7,461	2	11	1	12	146	348	414	622	1,011	1,434	1,312	786	434
2001	5,893	6	1	3	12	112	260	332	511	876	1,104	947	627	318
2000	5,137	4	2	1	11	109	213	299	465	753	986	766	481	300
1999	4,802	1	6	3	7	90	202	293	450	741	859	702	437	267
Non-Hispanic Black														
Both sexes														
2002	3,402	13	10	2	3	31	96	170	281	490	691	713	516	201
2001	3,097	11	8	2	3	27	85	147	257	478	662	681	427	139
2000	2,975	6	5	3	1	27	86	160	279	479	664	605	385	129
1999	3,024	8	11	2	2	23	100	183	305	472	696	623	316	123
Male														
2002	2,264	8	6	2	1	22	68	120	180	322	418	455	378	152
2001	2,113	9	5	-	2	14	71	112	157	300	421	488	314	105

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]										
Non-Hispanic[1]										
Male										
2001	207	149	121	110	72	43	23	4	1	5
2000	181	117	101	104	77	60	21	4	-	7
1999	200	145	114	102	79	47	16	7	-	4
Female										
2002	266	172	148	139	119	95	50	8	1	2
2001	219	125	121	135	105	89	46	11	3	-
2000	188	113	125	121	114	84	50	11	-	1
1999	154	136	126	150	116	81	32	10	1	2
Non-Hispanic White										
Both sexes										
2002	447	293	243	225	197	142	77	12	1	5
2001	344	236	217	225	165	120	64	11	2	5
2000	315	197	210	214	179	129	65	11	-	4
1999	283	243	216	236	187	112	47	15	1	2
Male										
2002	205	132	106	97	78	56	32	5	-	3
2001	145	126	104	93	69	38	23	2	-	5
2000	144	95	89	100	72	54	18	2	-	3
1999	142	119	103	92	75	44	16	6	-	1
Female										
2002	242	161	137	128	119	86	45	7	1	2
2001	199	110	113	132	96	82	41	9	2	-
2000	171	102	121	114	107	75	47	9	-	1
1999	141	124	113	144	112	68	31	9	1	1
Non-Hispanic Black										
Both sexes										
2002	99	33	15	16	8	8	4	1	-	1
2001	74	30	21	17	8	10	5	3	2	-
2000	49	32	15	11	11	14	6	4	-	4
1999	66	34	20	12	7	14	1	2	-	4
Male										
2002	79	26	8	8	8	2	-	-	-	1
2001	58	20	15	14	1	5	-	1	1	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]														
Non-Hispanic Black														
Male														
2000	2,049	4	5	1	-	16	60	120	183	309	440	440	283	96
1999	2,131	5	4	1	1	19	74	140	201	320	465	458	237	101
Female														
2002	1,138	5	4	-	2	9	28	50	101	168	273	258	138	49
2001	984	2	3	2	1	13	14	35	100	178	241	193	113	34
2000	926	2	-	2	1	11	26	40	96	170	224	165	102	33
1999	893	3	7	1	1	4	26	43	104	152	231	165	79	22
Origin not stated[2]														
Both sexes														
2002	147	-	-	-	-	2	7	8	12	23	29	23	15	9
2001	174	-	-	-	-	2	7	4	14	27	29	33	23	14
2000	185	-	1	-	1	2	7	9	16	17	29	45	31	11
1999	197	-	-	-	-	2	3	8	11	26	49	33	25	18
Male														
2002	109	-	-	-	-	-	5	7	10	16	22	18	7	8
2001	140	-	-	-	-	1	6	4	12	23	20	26	17	11
2000	133	-	1	-	1	1	6	4	11	13	22	29	23	8
1999	144	-	-	-	-	1	3	7	9	19	33	25	20	12
Female														
2002	38	-	-	-	-	2	2	1	2	7	7	5	8	1
2001	34	-	-	-	-	1	1	-	2	4	9	7	6	3
2000	52	-	-	-	-	1	1	5	5	4	7	16	8	3
1999	53	-	-	-	-	1	-	1	2	7	16	8	5	6
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]														
All origins														
Both sexes														
2002	30,242	10	61	71	277	2,474	4,306	3,380	2,760	2,712	2,734	2,358	1,963	1,608
2001	29,573	11	70	79	254	2,523	4,164	3,350	2,787	2,723	2,565	2,203	1,917	1,471
2000	28,663	13	46	70	307	2,606	3,969	3,165	2,624	2,718	2,640	2,146	1,805	1,289
1999	28,874	8	65	80	336	2,896	3,899	3,248	2,727	2,812	2,422	2,096	1,626	1,299

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Drug-induced causes (F11.0-11.5, F11.7-F11.9,F12.0-F12.5,F12.7-F12.9, F13.0-F13.5,F13.7-F13.9,F14.0-F14.5, F14.7-F14.9,F15.0-F15.5,F15.7-F15.9, F16.0-F16.5,F16.7-F16.9,F17.0, F17.3-F17.5,F17.7-F17.9,F18.0-F18.5, F18.7-F18.9,F19.0-F19.5,F19.7-F19.9, X40-X44,X60-X64,X85,Y10-Y14)[4]										
Non-Hispanic Black										
Male										
2000	35	22	11	4	5	6	3	2	-	4
1999	53	24	10	7	4	3	-	1	-	3
Female										
2002	20	7	7	8	-	6	4	1	-	-
2001	16	10	6	3	7	5	5	2	1	-
2000	14	10	4	7	6	8	3	2	-	-
1999	13	10	10	5	3	11	1	1	-	1
Origin not stated[2]										
Both sexes										
2002	3	6	3	3	-	-	-	1	-	3
2001	7	3	1	1	-	-	-	-	-	9
2000	6	1	3	-	-	-	-	-	-	6
1999	7	3	1	-	2	2	-	-	-	7
Male										
2002	3	5	3	2	-	-	-	1	-	2
2001	7	2	1	1	-	-	-	-	-	9
2000	6	1	1	-	-	-	-	-	-	6
1999	7	3	1	-	-	1	-	-	-	3
Female										
2002	-	1	-	1	-	-	-	-	-	1
2001	-	1	-	-	-	-	-	-	-	-
2000	-	-	2	-	-	-	-	-	-	-
1999	-	-	-	-	2	1	-	-	-	4
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]										
All origins										
Both sexes										
2002	1,111	966	1,027	1,011	822	411	138	27	-	15
2001	1,073	999	999	1,014	789	402	139	20	2	19
2000	983	923	1,018	994	729	443	137	19	1	18
1999	1,007	955	1,079	1,029	709	428	108	25	-	20

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]														
All origins														
Male														
2002	26,098	4	38	38	204	2,209	3,887	2,994	2,353	2,257	2,246	1,927	1,652	1,354
2001	25,480	7	39	51	198	2,270	3,796	2,963	2,384	2,216	2,096	1,828	1,588	1,231
2000	24,582	9	28	46	256	2,323	3,583	2,796	2,207	2,207	2,156	1,770	1,488	1,071
1999	24,700	5	36	50	258	2,560	3,496	2,848	2,308	2,279	1,950	1,722	1,335	1,080
Female														
2002	4,144	6	23	33	73	265	419	386	407	455	488	431	311	254
2001	4,093	4	31	28	56	253	368	387	403	507	469	375	329	240
2000	4,081	4	18	24	51	283	386	369	417	511	484	376	317	218
1999	4,174	3	29	30	78	336	403	400	419	533	472	374	291	219
Hispanic														
Both sexes														
2002	3,143	2	15	17	39	514	746	572	348	267	192	127	93	62
2001	3,087	4	13	6	34	468	765	575	351	264	204	127	88	47
2000	2,891	1	7	10	45	511	673	469	319	213	201	156	99	51
1999	2,878	2	6	11	58	533	656	466	304	251	176	151	67	54
Male														
2002	2,834	-	9	12	26	474	699	524	320	231	159	107	78	56
2001	2,774	3	6	5	26	425	707	526	315	231	177	106	74	44
2000	2,582	-	4	7	39	471	625	429	282	175	176	127	83	46
1999	2,549	2	4	6	44	481	597	427	272	216	148	118	55	46
Female														
2002	309	2	6	5	13	40	47	48	28	36	33	20	15	6
2001	313	1	7	1	8	43	58	49	36	33	27	21	14	3
2000	309	1	3	3	6	40	48	40	37	38	25	29	16	5
1999	329	-	2	5	14	52	59	39	32	35	28	33	12	8
Non-Hispanic[1]														
Both sexes														
2002	26,944	8	46	54	236	1,939	3,537	2,789	2,398	2,431	2,532	2,219	1,861	1,540
2001	26,341	7	57	72	220	2,040	3,369	2,756	2,426	2,451	2,351	2,070	1,825	1,417
2000	25,637	12	39	58	262	2,081	3,274	2,686	2,294	2,489	2,426	1,980	1,699	1,235
1999	25,877	6	59	69	278	2,351	3,236	2,767	2,407	2,547	2,235	1,940	1,551	1,242
Male														
2002	23,127	4	29	26	176	1,716	3,166	2,455	2,022	2,013	2,080	1,809	1,566	1,294
2001	22,573	4	33	45	172	1,831	3,063	2,418	2,061	1,979	1,910	1,718	1,510	1,180
2000	21,881	9	24	38	217	1,840	2,937	2,358	1,916	2,017	1,967	1,634	1,399	1,022
1999	22,050	3	32	44	214	2,070	2,894	2,408	2,024	2,051	1,791	1,600	1,274	1,031
Female														
2002	3,817	4	17	28	60	223	371	334	376	418	452	410	295	246

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]										
All origins										
Male										
2002	942	836	932	916	755	386	128	26	-	14
2001	885	865	889	915	719	373	128	19	2	18
2000	825	783	901	900	671	404	125	19	1	13
1999	871	833	953	928	652	392	103	22	-	19
Female										
2002	169	130	95	95	67	25	10	1	-	1
2001	188	134	110	99	70	29	11	1	-	1
2000	158	140	117	94	58	39	12	-	-	5
1999	136	122	126	101	57	36	5	3	-	1
Hispanic										
Both sexes										
2002	37	34	28	15	18	9	2	2	-	4
2001	35	36	24	14	17	6	3	1	-	5
2000	32	30	34	17	11	7	1	1	-	3
1999	33	36	15	27	17	5	5	2	-	3
Male										
2002	33	30	27	15	17	9	2	2	-	4
2001	31	33	22	14	14	6	3	1	-	5
2000	27	26	28	15	11	7	1	1	-	2
1999	30	31	15	27	15	5	5	2	-	3
Female										
2002	4	4	1	-	1	-	-	-	-	-
2001	4	3	2	-	3	-	-	-	-	-
2000	5	4	6	2	-	-	-	-	-	1
1999	3	5	-	-	2	-	-	-	-	-
Non-Hispanic[1]										
Both sexes										
2002	1,069	925	997	992	803	401	135	25	-	7
2001	1,029	956	971	998	769	393	136	18	2	8
2000	944	888	981	976	718	435	135	18	1	6
1999	968	916	1,062	997	692	421	103	23	-	7
Male										
2002	904	799	903	897	737	376	125	24	-	6
2001	845	825	863	899	702	364	125	17	2	7
2000	791	754	870	884	660	397	123	18	1	5
1999	835	799	936	896	637	386	98	20	-	7
Female										
2002	165	126	94	95	66	25	10	1	-	1

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]														
Non-Hispanic[1]														
Female														
2001	3,768	3	24	27	48	209	306	338	365	472	441	352	315	237
2000	3,756	3	15	20	45	241	337	328	378	472	459	346	300	213
1999	3,827	3	27	25	64	281	342	359	383	496	444	340	277	211
Non-Hispanic White														
Both sexes														
2002	18,762	2	28	28	135	876	1,480	1,276	1,363	1,662	2,000	1,853	1,604	1,406
2001	18,676	4	30	55	148	962	1,411	1,355	1,492	1,746	1,855	1,757	1,604	1,278
2000	18,042	5	21	38	181	958	1,419	1,298	1,404	1,786	1,908	1,630	1,476	1,100
1999	18,260	2	27	32	193	1,092	1,412	1,412	1,537	1,863	1,761	1,594	1,357	1,112
Male														
2002	15,881	2	16	12	102	754	1,284	1,058	1,111	1,350	1,648	1,497	1,358	1,175
2001	15,760	2	19	34	118	824	1,235	1,156	1,227	1,399	1,511	1,458	1,330	1,055
2000	15,160	2	11	24	151	818	1,232	1,088	1,144	1,414	1,551	1,348	1,212	908
1999	15,384	2	15	19	149	943	1,222	1,192	1,261	1,482	1,411	1,315	1,125	913
Female														
2002	2,881	-	12	16	33	122	196	218	252	312	352	356	246	231
2001	2,916	2	11	21	30	138	176	199	265	347	344	299	274	223
2000	2,882	3	10	14	30	140	187	210	260	372	357	282	264	192
1999	2,876	-	12	13	44	149	190	220	276	381	350	279	232	199
Non-Hispanic Black														
Both sexes														
2002	7,494	6	17	18	84	979	1,925	1,428	965	699	473	326	209	106
2001	7,063	2	23	15	63	1,015	1,857	1,311	861	656	436	272	192	112
2000	6,958	7	13	18	73	1,041	1,757	1,287	834	633	460	306	190	99
1999	6,933	2	26	31	74	1,163	1,719	1,246	788	610	427	305	152	111
Male														
2002	6,681	2	12	11	63	886	1,771	1,324	858	603	388	282	168	95
2001	6,323	1	13	9	51	947	1,743	1,188	777	540	354	227	160	101
2000	6,193	7	8	13	61	954	1,620	1,182	728	548	372	251	160	82
1999	6,114	1	12	22	56	1,041	1,582	1,128	698	509	345	254	122	102
Female														
2002	813	4	5	7	21	93	154	104	107	96	85	44	41	11
2001	740	1	10	6	12	68	114	123	84	116	82	45	32	11
2000	765	-	5	5	12	87	137	105	106	85	88	55	30	17
1999	819	1	14	9	18	122	137	118	90	101	82	51	30	9
Origin not stated[2]														
Both sexes														
2002	155	-	-	-	2	21	23	19	14	14	10	12	9	6

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]										
Non-Hispanic[1]										
Female										
2001	184	131	108	99	67	29	11	1	-	1
2000	153	134	111	92	58	38	12	-	-	1
1999	133	117	126	101	55	35	5	3	-	-
Non-Hispanic White										
Both sexes										
2002	974	865	947	932	782	389	132	23	-	5
2001	955	890	907	953	732	384	132	18	2	6
2000	865	823	915	945	692	424	132	17	1	4
1999	864	850	1,002	948	666	411	101	21	-	3
Male										
2002	830	748	860	844	718	365	123	22	-	4
2001	783	768	809	860	671	356	121	17	2	5
2000	721	699	809	860	638	389	120	17	1	3
1999	746	744	882	854	614	377	96	19	-	3
Female										
2002	144	117	87	88	64	24	9	1	-	1
2001	172	122	98	93	61	28	11	1	-	1
2000	144	124	106	85	54	35	12	-	-	1
1999	118	106	120	94	52	34	5	2	-	-
Non-Hispanic Black										
Both sexes										
2002	80	48	47	49	16	12	3	2	-	2
2001	59	53	55	39	30	7	4	-	-	1
2000	63	56	58	26	24	10	2	-	-	1
1999	84	58	51	47	21	10	2	2	-	4
Male										
2002	61	41	41	43	15	11	2	2	-	2
2001	49	46	47	34	25	6	4	-	-	1
2000	54	49	54	20	20	7	2	-	-	1
1999	72	49	46	41	18	9	2	1	-	4
Female										
2002	19	7	6	6	1	1	1	-	-	-
2001	10	7	8	5	5	1	-	-	-	-
2000	9	7	4	6	4	3	-	-	-	-
1999	12	9	5	6	3	1	-	1	-	-
Origin not stated[2]										
Both sexes										
2002	5	7	2	4	1	1	1	-	-	4

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years	50-54 years	55-59 years
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]														
Origin not stated[2]														
Both sexes														
2001	145	-	-	1	-	15	30	19	10	8	10	6	4	7
2000	135	-	-	2	-	14	22	10	11	16	13	10	7	3
1999	119	-	-	-	-	12	7	15	16	14	11	5	8	3
Male														
2002	137	-	-	-	2	19	22	15	11	13	7	11	8	4
2001	133	-	-	1	-	14	26	19	8	6	9	4	4	7
2000	119	-	-	1	-	12	21	9	9	15	13	9	6	3
1999	101	-	-	-	-	9	5	13	12	12	11	4	6	3
Female														
2002	18	-	-	-	-	2	1	4	3	1	3	1	1	2
2001	12	-	-	-	-	1	4	-	2	2	1	2	-	-
2000	16	-	-	1	-	2	1	1	2	1	-	1	1	-
1999	18	-	-	-	-	3	2	2	4	2	-	1	2	-

See footnotes at end of table.

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2002 - Con.

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2002"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), Hispanic origin, race for non-Hispanic population, sex, and year	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Injury by firearms (*U01.4,W32-W34, X72-X74,X93-X95,Y22-Y24,Y35.0)[4]										
Origin not stated[2]										
Both sexes										
2001	9	7	4	2	3	3	-	1	-	6
2000	7	5	3	1	-	1	1	-	-	9
1999	6	3	2	5	-	2	-	-	-	10
Male										
2002	5	7	2	4	1	1	1	-	-	4
2001	9	7	4	2	3	3	-	1	-	6
2000	7	3	3	1	-	-	1	-	-	6
1999	6	3	2	5	-	1	-	-	-	9
Female										
2002	-	-	-	-	-	-	-	-	-	-
2001	-	-	-	-	-	-	-	-	-	-
2000	-	2	-	-	-	1	-	-	-	3
1999	-	-	-	-	-	1	-	-	-	1

- Quantity zero.
. . . Category not applicable.
1 Includes races other than white and black.
2 Includes deaths for which Hispanic origin was not reported on the death certificate.
3 Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
4 Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.