

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
All origins, both sexes												
2003.....	2,448,288	28,025	4,965	2,898	4,056	13,595	19,973	18,250	23,050	34,055	55,406	78,173
2002.....	2,443,387	28,034	4,858	3,018	4,132	13,812	19,234	17,959	23,396	35,347	55,793	76,065
2001.....	2,416,425	27,568	5,107	3,093	4,002	13,555	18,697	18,211	23,472	36,895	54,779	73,684
2000.....	2,403,351	28,035	4,979	3,253	4,160	13,563	17,744	17,681	22,770	36,140	53,658	70,832
1999.....	2,391,399	27,937	5,249	3,474	4,121	13,778	16,878	17,703	23,363	36,804	52,452	68,120
All origins, male												
2003.....	1,201,964	15,902	2,826	1,640	2,510	9,706	14,964	13,043	15,559	21,741	34,694	49,139
2002.....	1,199,264	15,717	2,806	1,702	2,496	9,844	14,572	12,954	15,782	22,626	34,967	47,701
2001.....	1,183,421	15,477	2,899	1,727	2,441	9,766	14,197	13,006	15,751	23,727	34,437	46,567
2000.....	1,177,578	15,718	2,824	1,850	2,551	9,697	13,374	12,619	15,271	23,252	34,045	45,121
1999.....	1,175,460	15,646	2,975	1,964	2,528	9,780	12,634	12,542	15,734	23,681	33,437	43,303
All origins, female												
2003.....	1,246,324	12,123	2,139	1,258	1,546	3,889	5,009	5,207	7,491	12,314	20,712	29,034
2002.....	1,244,123	12,317	2,052	1,316	1,636	3,968	4,662	5,005	7,614	12,721	20,826	28,364
2001.....	1,233,004	12,091	2,208	1,366	1,561	3,789	4,500	5,205	7,721	13,168	20,342	27,117
2000.....	1,225,773	12,317	2,155	1,403	1,609	3,866	4,370	5,062	7,499	12,888	19,613	25,711
1999.....	1,215,939	12,291	2,274	1,510	1,593	3,998	4,244	5,161	7,629	13,123	19,015	24,817
Hispanic, both sexes												
2003.....	122,026	5,281	993	538	635	2,183	3,211	3,088	3,101	3,683	4,893	5,917
2002.....	117,135	4,943	933	538	586	2,086	3,169	2,803	3,013	3,644	4,823	5,527
2001.....	113,413	4,680	917	480	558	1,998	3,060	2,863	2,934	3,859	4,582	5,397
2000.....	107,254	4,598	872	510	555	1,953	2,840	2,598	2,911	3,656	4,429	4,990
1999.....	103,740	4,412	882	464	550	1,871	2,634	2,527	2,870	3,624	4,174	4,767

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Hispanic, male</b>												
2003.....	68,119	2,940	569	314	392	1,659	2,585	2,421	2,321	2,587	3,300	4,010
2002.....	65,703	2,746	546	303	344	1,614	2,568	2,239	2,235	2,514	3,314	3,695
2001.....	63,317	2,601	518	281	319	1,530	2,489	2,223	2,143	2,739	3,156	3,655
2000.....	60,172	2,514	475	277	345	1,528	2,309	2,014	2,186	2,543	3,054	3,357
1999.....	57,991	2,411	481	266	326	1,432	2,115	1,956	2,128	2,601	2,875	3,213
<b>Hispanic, female</b>												
2003.....	53,907	2,341	424	224	243	524	626	667	780	1,096	1,593	1,907
2002.....	51,432	2,197	387	235	242	472	601	564	778	1,130	1,509	1,832
2001.....	50,096	2,079	399	199	239	468	571	640	791	1,120	1,426	1,742
2000.....	47,082	2,084	397	233	210	425	531	584	725	1,113	1,375	1,633
1999.....	45,749	2,001	401	198	224	439	519	571	742	1,023	1,299	1,554
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	2,319,476	22,475	3,956	2,349	3,412	11,357	16,687	15,087	19,868	30,220	50,283	71,939
2002.....	2,318,269	22,816	3,904	2,472	3,530	11,671	15,987	15,073	20,256	31,512	50,713	70,178
2001.....	2,295,244	22,629	4,171	2,599	3,434	11,508	15,551	15,252	20,404	32,857	49,906	67,927
2000.....	2,287,846	23,074	4,086	2,725	3,591	11,550	14,824	15,000	19,741	32,283	48,899	65,454
1999.....	2,279,325	23,126	4,343	3,001	3,559	11,861	14,182	15,091	20,380	32,997	47,952	62,981
<b>Non-Hispanic 1/, male</b>												
2003.....	1,129,927	12,799	2,247	1,318	2,114	8,003	12,325	10,565	13,175	19,048	31,236	44,906
2002.....	1,129,090	12,815	2,250	1,394	2,140	8,192	11,937	10,657	13,457	19,976	31,475	43,743
2001.....	1,115,683	12,739	2,371	1,438	2,116	8,197	11,639	10,715	13,510	20,859	31,059	42,656
2000.....	1,112,704	13,007	2,338	1,569	2,201	8,131	11,005	10,542	13,006	20,556	30,748	41,490
1999.....	1,112,718	12,998	2,479	1,692	2,193	8,323	10,474	10,525	13,522	20,947	30,325	39,813

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	1,189,549	9,676	1,709	1,031	1,298	3,354	4,362	4,522	6,693	11,172	19,047	27,033
2002.....	1,189,179	10,001	1,654	1,078	1,390	3,479	4,050	4,416	6,799	11,536	19,238	26,435
2001.....	1,179,561	9,890	1,800	1,161	1,318	3,311	3,912	4,537	6,894	11,998	18,847	25,271
2000.....	1,175,142	10,067	1,748	1,156	1,390	3,419	3,819	4,458	6,735	11,727	18,151	23,964
1999.....	1,166,607	10,128	1,864	1,309	1,366	3,538	3,708	4,566	6,858	12,050	17,627	23,168
<b>Non-Hispanic White, both sexes</b>												
2003.....	1,979,465	13,223	2,539	1,579	2,305	8,230	11,410	9,937	13,620	21,279	36,816	53,153
2002.....	1,981,973	13,463	2,502	1,643	2,388	8,552	10,906	9,900	13,858	22,230	37,299	52,044
2001.....	1,962,810	13,311	2,772	1,777	2,374	8,264	10,466	10,130	14,061	23,334	36,212	50,151
2000.....	1,959,919	13,513	2,641	1,850	2,547	8,327	9,934	9,822	13,369	22,912	35,319	48,005
1999.....	1,953,197	13,553	2,819	1,990	2,498	8,490	9,378	10,048	13,938	23,345	34,285	45,974
<b>Non-Hispanic White, male</b>												
2003.....	956,194	7,591	1,423	893	1,426	5,661	8,361	6,942	9,155	13,718	23,405	33,819
2002.....	957,645	7,706	1,431	923	1,442	5,884	8,073	6,988	9,351	14,299	23,750	33,102
2001.....	945,967	7,508	1,592	957	1,471	5,756	7,786	7,168	9,445	15,102	23,111	32,045
2000.....	944,781	7,660	1,541	1,077	1,575	5,726	7,340	6,920	9,014	14,922	22,700	31,133
1999.....	944,913	7,718	1,605	1,115	1,528	5,799	6,878	7,056	9,393	15,144	22,147	29,448
<b>Non-Hispanic White, female</b>												
2003.....	1,023,271	5,632	1,116	686	879	2,569	3,049	2,995	4,465	7,561	13,411	19,334
2002.....	1,024,328	5,757	1,071	720	946	2,668	2,833	2,912	4,507	7,931	13,549	18,942
2001.....	1,016,843	5,803	1,180	820	903	2,508	2,680	2,962	4,616	8,232	13,101	18,106
2000.....	1,015,138	5,853	1,100	773	972	2,601	2,594	2,902	4,355	7,990	12,619	16,872
1999.....	1,008,284	5,835	1,214	875	970	2,691	2,500	2,992	4,545	8,201	12,138	16,526

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	287,968	8,159	1,169	628	902	2,530	4,491	4,447	5,348	7,746	11,815	16,498
2002.....	286,573	8,284	1,163	686	948	2,542	4,362	4,487	5,522	8,102	11,853	16,002
2001.....	284,343	8,277	1,156	685	872	2,662	4,362	4,388	5,471	8,287	12,149	15,657
2000.....	282,676	8,525	1,217	745	875	2,672	4,254	4,489	5,614	8,247	12,032	15,545
1999.....	281,979	8,589	1,283	869	902	2,813	4,167	4,390	5,602	8,563	12,207	15,131
<b>Non-Hispanic Black, male</b>												
2003.....	146,136	4,601	681	352	559	1,938	3,405	3,161	3,431	4,571	6,834	9,713
2002.....	144,802	4,520	681	392	585	1,914	3,334	3,179	3,544	4,906	6,755	9,372
2001.....	143,971	4,651	655	400	541	2,041	3,336	3,056	3,492	4,959	7,012	9,354
2000.....	143,297	4,757	672	425	534	2,012	3,212	3,127	3,522	4,947	7,114	9,225
1999.....	143,883	4,765	734	508	573	2,133	3,153	3,038	3,595	5,104	7,280	9,266
<b>Non-Hispanic Black, female</b>												
2003.....	141,832	3,558	488	276	343	592	1,086	1,286	1,917	3,175	4,981	6,785
2002.....	141,771	3,764	482	294	363	628	1,028	1,308	1,978	3,196	5,098	6,630
2001.....	140,372	3,626	501	285	331	621	1,026	1,332	1,979	3,328	5,137	6,303
2000.....	139,379	3,768	545	320	341	660	1,042	1,362	2,092	3,300	4,918	6,320
1999.....	138,096	3,824	549	361	329	680	1,014	1,352	2,007	3,459	4,927	5,865
<b>Origin not stated 2/, both sexes</b>												
2003.....	6,786	269	16	11	9	55	75	75	81	152	230	317
2002.....	7,983	275	21	8	16	55	78	83	127	191	257	360
2001.....	7,768	259	19	14	10	49	86	96	134	179	291	360
2000.....	8,251	363	21	18	14	60	80	83	118	201	330	388
1999.....	8,334	399	24	9	12	46	62	85	113	183	326	372

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Origin not stated 2/, male												
2003.....	3,918	163	10	8	4	44	54	57	63	106	158	223
2002.....	4,471	156	10	5	12	38	67	58	90	136	178	263
2001.....	4,421	137	10	8	6	39	69	68	98	129	222	256
2000.....	4,702	197	11	4	5	38	60	63	79	153	243	274
1999.....	4,751	237	15	6	9	25	45	61	84	133	237	277
Origin not stated 2/, female												
2003.....	2,868	106	6	3	5	11	21	18	18	46	72	94
2002.....	3,512	119	11	3	4	17	11	25	37	55	79	97
2001.....	3,347	122	9	6	4	10	17	28	36	50	69	104
2000.....	3,549	166	10	14	9	22	20	20	39	48	87	114
1999.....	3,583	162	9	3	3	21	17	24	29	50	89	95



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Salmonella infections (A01-A02)

Hispanic, male

2003.....	3	-	-	-	-	-	-	-	-	-	1	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	3	-	-	1	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	37	3	-	-	-	-	-	1	-	-	2	1
2002.....	18	3	-	-	-	-	-	-	-	-	2	1
2001.....	36	2	-	-	-	-	-	-	2	1	-	1
2000.....	26	1	2	-	-	1	-	-	1	-	-	1
1999.....	34	3	-	-	-	-	2	-	-	-	2	-

Non-Hispanic 1/, male

2003.....	20	1	-	-	-	-	-	-	-	-	1	1
2002.....	9	1	-	-	-	-	-	-	-	-	-	-
2001.....	16	-	-	-	-	-	-	-	1	1	-	1
2000.....	14	1	-	-	-	1	-	-	1	-	-	1
1999.....	18	2	-	-	-	-	2	-	-	-	1	-

















Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain other intestinal infections (A04,A07-A09)

All origins, both sexes

2003.....	3,090	10	4	4	3	1	3	-	8	4	10	21
2002.....	2,465	13	8	1	2	4	2	3	1	5	13	25
2001.....	1,537	10	18	1	3	-	-	1	3	4	8	9
2000.....	1,328	13	10	-	-	-	1	2	1	7	12	11
1999.....	1,108	12	8	2	1	1	4	3	3	2	5	11

All origins, male

2003.....	1,109	6	4	1	1	1	1	-	4	3	2	11
2002.....	947	3	5	1	-	3	-	1	-	2	6	12
2001.....	572	7	11	1	1	-	-	-	2	4	4	4
2000.....	524	10	6	-	-	-	-	2	1	4	6	7
1999.....	420	8	4	2	1	1	2	2	3	1	3	5

All origins, female

2003.....	1,981	4	-	3	2	-	2	-	4	1	8	10
2002.....	1,518	10	3	-	2	1	2	2	1	3	7	13
2001.....	965	3	7	-	2	-	-	1	1	-	4	5
2000.....	804	3	4	-	-	-	1	-	-	3	6	4
1999.....	688	4	4	-	-	-	2	1	-	1	2	6

Hispanic, both sexes

2003.....	99	2	2	2	1	-	-	-	1	-	-	4
2002.....	78	5	5	-	-	-	1	-	-	-	1	-
2001.....	53	1	1	-	-	-	-	-	-	1	3	-
2000.....	46	3	2	-	-	-	-	-	-	1	1	-
1999.....	34	3	1	-	-	1	1	1	-	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain other intestinal infections (A04,A07-A09)

Hispanic, male

2003.....	47	-	2	1	-	-	-	-	-	-	-	2
2002.....	31	1	3	-	-	-	-	-	-	-	-	-
2001.....	19	1	1	-	-	-	-	-	-	1	2	-
2000.....	23	2	1	-	-	-	-	-	-	1	-	-
1999.....	17	2	-	-	-	1	1	1	-	-	-	1

Hispanic, female

2003.....	52	2	-	1	1	-	-	-	1	-	-	2
2002.....	47	4	2	-	-	-	1	-	-	-	1	-
2001.....	34	-	-	-	-	-	-	-	-	-	1	-
2000.....	23	1	1	-	-	-	-	-	-	-	1	-
1999.....	17	1	1	-	-	-	-	-	-	-	-	1

Non-Hispanic 1/, both sexes

2003.....	2,986	8	2	2	2	1	3	-	7	4	10	17
2002.....	2,381	8	3	1	2	4	1	3	1	5	12	25
2001.....	1,474	9	16	1	3	-	-	1	3	3	5	9
2000.....	1,277	10	8	-	-	-	1	2	1	6	11	11
1999.....	1,071	9	7	2	1	-	3	2	3	2	5	9

Non-Hispanic 1/, male

2003.....	1,061	6	2	-	1	1	1	-	4	3	2	9
2002.....	915	2	2	1	-	3	-	1	-	2	6	12
2001.....	550	6	10	1	1	-	-	-	2	3	2	4
2000.....	497	8	5	-	-	-	-	2	1	3	6	7
1999.....	403	6	4	2	1	-	1	1	3	1	3	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic 1/, female

2003.....	1,925	2	-	2	1	-	2	-	3	1	8	8
2002.....	1,466	6	1	-	2	1	1	2	1	3	6	13
2001.....	924	3	6	-	2	-	-	1	1	-	3	5
2000.....	780	2	3	-	-	-	1	-	-	3	5	4
1999.....	668	3	3	-	-	-	2	1	-	1	2	5

Non-Hispanic White, both sexes

2003.....	2,776	3	1	2	1	1	2	-	5	4	7	15
2002.....	2,205	2	1	1	2	4	1	2	-	4	10	21
2001.....	1,357	5	13	1	2	-	-	-	2	2	4	6
2000.....	1,175	5	5	-	-	-	1	1	1	2	7	8
1999.....	990	5	6	-	1	-	2	2	3	2	2	6

Non-Hispanic White, male

2003.....	989	2	1	-	1	1	1	-	3	3	2	8
2002.....	840	-	1	1	-	3	-	1	-	1	5	11
2001.....	497	3	8	1	-	-	-	-	2	2	1	2
2000.....	453	4	4	-	-	-	-	1	1	1	5	5
1999.....	373	4	3	-	1	-	1	1	3	1	2	2

Non-Hispanic White, female

2003.....	1,787	1	-	2	-	-	1	-	2	1	5	7
2002.....	1,365	2	-	-	2	1	1	1	-	3	5	10
2001.....	860	2	5	-	2	-	-	-	-	-	3	4
2000.....	722	1	1	-	-	-	1	-	-	1	2	3
1999.....	617	1	3	-	-	-	1	1	-	1	-	4





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Tuberculosis (A16-A19)</b>												
All origins, both sexes												
2003.....	711	-	1	-	2	3	7	8	10	11	24	54
2002.....	784	1	-	-	-	-	9	7	11	18	34	35
2001.....	764	2	1	-	-	6	6	11	14	18	22	37
2000.....	776	1	6	-	1	2	9	7	13	17	25	38
1999.....	930	2	2	1	2	1	10	3	16	25	32	51
All origins, male												
2003.....	427	-	1	-	2	1	2	6	7	8	17	39
2002.....	462	-	-	-	-	-	5	4	7	11	27	20
2001.....	453	-	-	-	-	5	3	7	7	12	15	27
2000.....	468	-	1	-	-	2	5	3	11	11	17	35
1999.....	584	-	2	-	2	1	3	1	10	20	23	42
All origins, female												
2003.....	284	-	-	-	-	2	5	2	3	3	7	15
2002.....	322	1	-	-	-	-	4	3	4	7	7	15
2001.....	311	2	1	-	-	1	3	4	7	6	7	10
2000.....	308	1	5	-	1	-	4	4	2	6	8	3
1999.....	346	2	-	1	-	-	7	2	6	5	9	9
Hispanic, both sexes												
2003.....	89	-	1	-	-	2	3	2	4	4	4	6
2002.....	95	-	-	-	-	-	2	2	2	5	6	3
2001.....	97	-	-	-	-	4	2	6	3	4	5	3
2000.....	83	-	2	-	1	-	3	4	3	1	2	2
1999.....	125	1	-	-	1	1	3	1	2	7	6	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Tuberculosis (A16-A19)</b>												
<b>Hispanic, male</b>												
2003.....	60	-	1	-	-	1	2	2	2	4	3	5
2002.....	52	-	-	-	-	-	1	2	1	4	5	1
2001.....	67	-	-	-	-	4	2	5	2	2	3	3
2000.....	46	-	-	-	-	-	1	2	2	1	1	2
1999.....	78	-	-	-	1	1	2	1	2	5	4	8
<b>Hispanic, female</b>												
2003.....	29	-	-	-	-	1	1	-	2	-	1	1
2002.....	43	-	-	-	-	-	1	-	1	1	1	2
2001.....	30	-	-	-	-	-	-	1	1	2	2	-
2000.....	37	-	2	-	1	-	2	2	1	-	1	-
1999.....	47	1	-	-	-	-	1	-	-	2	2	2
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	617	-	-	-	2	1	4	6	6	6	20	48
2002.....	682	1	-	-	-	-	6	5	9	11	28	32
2001.....	660	2	1	-	-	2	4	5	11	13	17	34
2000.....	683	1	4	-	-	2	6	3	10	16	23	34
1999.....	788	1	2	1	1	-	6	2	14	18	24	39
<b>Non-Hispanic 1/, male</b>												
2003.....	364	-	-	-	2	-	-	4	5	4	14	34
2002.....	404	-	-	-	-	-	4	2	6	5	22	19
2001.....	379	-	-	-	-	1	1	2	5	9	12	24
2000.....	414	-	1	-	-	2	4	1	9	10	16	31
1999.....	492	-	2	-	1	-	-	-	8	15	17	32

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Tuberculosis (A16-A19)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	253	-	-	-	-	1	4	2	1	2	6	14
2002.....	278	1	-	-	-	-	2	3	3	6	6	13
2001.....	281	2	1	-	-	1	3	3	6	4	5	10
2000.....	269	1	3	-	-	-	2	2	1	6	7	3
1999.....	296	1	-	1	-	-	6	2	6	3	7	7
<b>Non-Hispanic White, both sexes</b>												
2003.....	347	-	-	-	2	-	1	2	-	-	3	16
2002.....	366	-	-	-	-	-	1	1	1	2	8	10
2001.....	368	1	1	-	-	-	2	1	1	2	10	17
2000.....	353	-	2	-	-	1	2	1	3	7	3	10
1999.....	430	-	-	-	-	-	-	1	1	5	6	14
<b>Non-Hispanic White, male</b>												
2003.....	195	-	-	-	2	-	-	1	-	-	2	13
2002.....	218	-	-	-	-	-	1	-	-	-	7	6
2001.....	200	-	-	-	-	-	-	1	1	2	8	11
2000.....	205	-	-	-	-	1	2	-	3	7	2	9
1999.....	263	-	-	-	-	-	-	-	1	4	5	12
<b>Non-Hispanic White, female</b>												
2003.....	152	-	-	-	-	-	1	1	-	-	1	3
2002.....	148	-	-	-	-	-	-	1	1	2	1	4
2001.....	168	1	1	-	-	-	2	-	-	-	2	6
2000.....	148	-	2	-	-	-	-	1	-	-	1	1
1999.....	167	-	-	-	-	-	-	1	-	1	1	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Tuberculosis (A16-A19)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	168	-	-	-	-	1	3	1	5	4	11	24
2002.....	198	-	-	-	-	-	3	4	4	8	15	17
2001.....	200	-	-	-	-	-	1	3	6	9	7	14
2000.....	238	-	2	-	-	-	2	2	5	8	17	20
1999.....	239	1	2	1	-	-	4	1	8	10	14	20
<b>Non-Hispanic Black, male</b>												
2003.....	111	-	-	-	-	-	-	1	4	3	8	17
2002.....	123	-	-	-	-	-	2	2	3	4	12	11
2001.....	124	-	-	-	-	-	-	1	2	6	4	11
2000.....	156	-	1	-	-	-	1	1	4	2	12	18
1999.....	154	-	2	-	-	-	-	-	4	8	10	17
<b>Non-Hispanic Black, female</b>												
2003.....	57	-	-	-	-	1	3	-	1	1	3	7
2002.....	75	-	-	-	-	-	1	2	1	4	3	6
2001.....	76	-	-	-	-	-	1	2	4	3	3	3
2000.....	82	-	1	-	-	-	1	1	1	6	5	2
1999.....	85	1	-	1	-	-	4	1	4	2	4	3
<b>Origin not stated 2/, both sexes</b>												
2003.....	5	-	-	-	-	-	-	-	-	1	-	-
2002.....	7	-	-	-	-	-	1	-	-	2	-	-
2001.....	7	-	-	-	-	-	-	-	-	1	-	-
2000.....	10	-	-	-	-	-	-	-	-	-	-	2
1999.....	17	-	-	-	-	-	1	-	-	-	2	2



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Respiratory tuberculosis (A16)</b>												
<b>All origins, both sexes</b>												
2003.....	562	-	1	-	2	2	7	2	4	5	21	44
2002.....	626	-	-	-	-	-	6	6	8	13	27	28
2001.....	600	2	1	-	-	3	4	8	9	14	14	27
2000.....	615	-	3	-	-	2	5	5	6	15	20	30
1999.....	764	1	1	-	1	-	8	2	12	19	26	38
<b>All origins, male</b>												
2003.....	344	-	1	-	2	1	2	2	4	3	14	32
2002.....	372	-	-	-	-	-	4	4	5	9	22	16
2001.....	371	-	-	-	-	3	2	5	5	9	9	20
2000.....	393	-	1	-	-	2	2	3	5	11	15	29
1999.....	486	-	1	-	1	-	2	-	8	15	18	32
<b>All origins, female</b>												
2003.....	218	-	-	-	-	1	5	-	-	2	7	12
2002.....	254	-	-	-	-	-	2	2	3	4	5	12
2001.....	229	2	1	-	-	-	2	3	4	5	5	7
2000.....	222	-	2	-	-	-	3	2	1	4	5	1
1999.....	278	1	-	-	-	-	6	2	4	4	8	6
<b>Hispanic, both sexes</b>												
2003.....	77	-	1	-	-	2	3	2	2	3	4	4
2002.....	78	-	-	-	-	-	1	2	2	5	5	3
2001.....	77	-	-	-	-	2	1	5	3	2	3	2
2000.....	60	-	1	-	-	-	2	2	1	1	1	-
1999.....	93	-	-	-	-	-	2	-	1	5	5	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Respiratory tuberculosis (A16)</b>												
<b>Hispanic, male</b>												
2003.....	55	-	1	-	-	1	2	2	2	3	3	4
2002.....	44	-	-	-	-	-	1	2	1	4	4	1
2001.....	56	-	-	-	-	2	1	4	2	1	2	2
2000.....	40	-	-	-	-	-	1	2	1	1	1	-
1999.....	55	-	-	-	-	-	1	-	1	3	3	5
<b>Hispanic, female</b>												
2003.....	22	-	-	-	-	1	1	-	-	-	1	-
2002.....	34	-	-	-	-	-	-	-	1	1	1	2
2001.....	21	-	-	-	-	-	-	1	1	1	1	-
2000.....	20	-	1	-	-	-	1	-	-	-	-	-
1999.....	38	-	-	-	-	-	1	-	-	2	2	1
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	481	-	-	-	2	-	4	-	2	1	17	40
2002.....	543	-	-	-	-	-	5	4	6	6	22	25
2001.....	517	2	1	-	-	1	3	3	6	11	11	25
2000.....	546	-	2	-	-	2	3	3	5	14	19	28
1999.....	657	1	1	-	1	-	5	2	11	14	19	31
<b>Non-Hispanic 1/, male</b>												
2003.....	287	-	-	-	2	-	-	-	2	-	11	28
2002.....	323	-	-	-	-	-	3	2	4	3	18	15
2001.....	309	-	-	-	-	1	1	1	3	7	7	18
2000.....	345	-	1	-	-	2	1	1	4	10	14	27
1999.....	419	-	1	-	1	-	-	-	7	12	13	26

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Respiratory tuberculosis (A16)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	194	-	-	-	-	-	4	-	-	1	6	12
2002.....	220	-	-	-	-	-	2	2	2	3	4	10
2001.....	208	2	1	-	-	-	2	2	3	4	4	7
2000.....	201	-	1	-	-	-	2	2	1	4	5	1
1999.....	238	1	-	-	-	-	5	2	4	2	6	5
<b>Non-Hispanic White, both sexes</b>												
2003.....	270	-	-	-	2	-	1	-	-	-	2	14
2002.....	296	-	-	-	-	-	1	-	1	2	7	9
2001.....	288	1	1	-	-	-	1	-	-	1	7	12
2000.....	288	-	-	-	-	1	1	1	1	7	2	8
1999.....	371	-	-	-	-	-	-	1	1	3	6	13
<b>Non-Hispanic White, male</b>												
2003.....	154	-	-	-	2	-	-	-	-	-	1	12
2002.....	177	-	-	-	-	-	1	-	-	-	7	5
2001.....	162	-	-	-	-	-	-	-	-	1	5	7
2000.....	174	-	-	-	-	1	1	-	1	7	2	8
1999.....	234	-	-	-	-	-	-	-	1	3	5	11
<b>Non-Hispanic White, female</b>												
2003.....	116	-	-	-	-	-	1	-	-	-	1	2
2002.....	119	-	-	-	-	-	-	-	1	2	-	4
2001.....	126	1	1	-	-	-	1	-	-	-	2	5
2000.....	114	-	-	-	-	-	-	1	-	-	-	-
1999.....	137	-	-	-	-	-	-	1	-	-	1	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Respiratory tuberculosis (A16)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	134	-	-	-	-	-	3	-	2	-	10	20
2002.....	160	-	-	-	-	-	2	4	4	3	11	13
2001.....	157	-	-	-	-	-	1	3	3	8	4	12
2000.....	187	-	2	-	-	-	1	2	4	6	14	16
1999.....	194	1	1	-	-	-	4	1	7	9	10	15
<b>Non-Hispanic Black, male</b>												
2003.....	90	-	-	-	-	-	-	-	2	-	7	13
2002.....	101	-	-	-	-	-	1	2	3	2	8	8
2001.....	103	-	-	-	-	-	-	1	1	5	2	10
2000.....	129	-	1	-	-	-	-	1	3	2	10	15
1999.....	122	-	1	-	-	-	-	-	3	7	6	13
<b>Non-Hispanic Black, female</b>												
2003.....	44	-	-	-	-	-	3	-	-	-	3	7
2002.....	59	-	-	-	-	-	1	2	1	1	3	5
2001.....	54	-	-	-	-	-	1	2	2	3	2	2
2000.....	58	-	1	-	-	-	1	1	1	4	4	1
1999.....	72	1	-	-	-	-	4	1	4	2	4	2
<b>Origin not stated 2/, both sexes</b>												
2003.....	4	-	-	-	-	-	-	-	-	1	-	-
2002.....	5	-	-	-	-	-	-	-	-	2	-	-
2001.....	6	-	-	-	-	-	-	-	-	1	-	-
2000.....	9	-	-	-	-	-	-	-	-	-	-	2
1999.....	14	-	-	-	-	-	1	-	-	-	2	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
All origins, both sexes												
2003.....	149	-	-	-	-	1	-	6	6	6	3	10
2002.....	158	1	-	-	-	-	3	1	3	5	7	7
2001.....	164	-	-	-	-	3	2	3	5	4	8	10
2000.....	161	1	3	-	1	-	4	2	7	2	5	8
1999.....	166	1	1	1	1	1	2	1	4	6	6	13
All origins, male												
2003.....	83	-	-	-	-	-	-	4	3	5	3	7
2002.....	90	-	-	-	-	-	1	-	2	2	5	4
2001.....	82	-	-	-	-	2	1	2	2	3	6	7
2000.....	75	-	-	-	-	-	3	-	6	-	2	6
1999.....	98	-	1	-	1	1	1	1	2	5	5	10
All origins, female												
2003.....	66	-	-	-	-	1	-	2	3	1	-	3
2002.....	68	1	-	-	-	-	2	1	1	3	2	3
2001.....	82	-	-	-	-	1	1	1	3	1	2	3
2000.....	86	1	3	-	1	-	1	2	1	2	3	2
1999.....	68	1	-	1	-	-	1	-	2	1	1	3
Hispanic, both sexes												
2003.....	12	-	-	-	-	-	-	-	2	1	-	2
2002.....	17	-	-	-	-	-	1	-	-	-	1	-
2001.....	20	-	-	-	-	2	1	1	-	2	2	1
2000.....	23	-	1	-	1	-	1	2	2	-	1	2
1999.....	32	1	-	-	1	1	1	1	1	2	1	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Hispanic, male												
2003.....	5	-	-	-	-	-	-	-	-	1	-	1
2002.....	8	-	-	-	-	-	-	-	-	-	1	-
2001.....	11	-	-	-	-	2	1	1	-	1	1	1
2000.....	6	-	-	-	-	-	-	-	1	-	-	2
1999.....	23	-	-	-	1	1	1	1	1	2	1	3
Hispanic, female												
2003.....	7	-	-	-	-	-	-	-	2	-	-	1
2002.....	9	-	-	-	-	-	1	-	-	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	1	1	-
2000.....	17	-	1	-	1	-	1	2	1	-	1	-
1999.....	9	1	-	-	-	-	-	-	-	-	-	1
Non-Hispanic 1/, both sexes												
2003.....	136	-	-	-	-	1	-	6	4	5	3	8
2002.....	139	1	-	-	-	-	1	1	3	5	6	7
2001.....	143	-	-	-	-	1	1	2	5	2	6	9
2000.....	137	1	2	-	-	-	3	-	5	2	4	6
1999.....	131	-	1	1	-	-	1	-	3	4	5	8
Non-Hispanic 1/, male												
2003.....	77	-	-	-	-	-	-	4	3	4	3	6
2002.....	81	-	-	-	-	-	1	-	2	2	4	4
2001.....	70	-	-	-	-	-	-	1	2	2	5	6
2000.....	69	-	-	-	-	-	3	-	5	-	2	4
1999.....	73	-	1	-	-	-	-	-	1	3	4	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other tuberculosis (A17-A19)												
Non-Hispanic 1/, female												
2003.....	59	-	-	-	-	1	-	2	1	1	-	2
2002.....	58	1	-	-	-	-	-	1	1	3	2	3
2001.....	73	-	-	-	-	1	1	1	3	-	1	3
2000.....	68	1	2	-	-	-	-	-	-	2	2	2
1999.....	58	-	-	1	-	-	1	-	2	1	1	2
Non-Hispanic White, both sexes												
2003.....	77	-	-	-	-	-	-	2	-	-	1	2
2002.....	70	-	-	-	-	-	-	1	-	-	1	1
2001.....	80	-	-	-	-	-	1	1	1	1	3	5
2000.....	65	-	2	-	-	-	1	-	2	-	1	2
1999.....	59	-	-	-	-	-	-	-	-	2	-	1
Non-Hispanic White, male												
2003.....	41	-	-	-	-	-	-	1	-	-	1	1
2002.....	41	-	-	-	-	-	-	-	-	-	-	1
2001.....	38	-	-	-	-	-	-	1	1	1	3	4
2000.....	31	-	-	-	-	-	1	-	2	-	-	1
1999.....	29	-	-	-	-	-	-	-	-	1	-	1
Non-Hispanic White, female												
2003.....	36	-	-	-	-	-	-	1	-	-	-	1
2002.....	29	-	-	-	-	-	-	1	-	-	1	-
2001.....	42	-	-	-	-	-	1	-	-	-	-	1
2000.....	34	-	2	-	-	-	-	-	-	-	1	1
1999.....	30	-	-	-	-	-	-	-	-	1	-	-

























Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningococcal infection (A39)</b>												
All origins, both sexes												
2003.....	161	12	18	2	9	18	21	8	4	10	9	8
2002.....	161	17	12	6	14	21	12	12	4	4	5	10
2001.....	199	16	19	12	14	43	12	6	4	8	5	9
2000.....	211	22	19	12	6	34	22	9	6	7	6	13
1999.....	227	32	30	13	12	28	11	11	12	12	15	11
All origins, male												
2003.....	75	9	10	1	2	8	10	3	4	4	5	5
2002.....	83	8	9	5	8	11	6	8	2	2	3	5
2001.....	94	6	9	6	9	20	7	4	2	3	3	6
2000.....	106	14	12	5	2	24	11	3	3	6	3	5
1999.....	101	14	19	7	6	13	6	8	4	3	6	1
All origins, female												
2003.....	86	3	8	1	7	10	11	5	-	6	4	3
2002.....	78	9	3	1	6	10	6	4	2	2	2	5
2001.....	105	10	10	6	5	23	5	2	2	5	2	3
2000.....	105	8	7	7	4	10	11	6	3	1	3	8
1999.....	126	18	11	6	6	15	5	3	8	9	9	10
Hispanic, both sexes												
2003.....	23	5	2	-	1	1	1	3	2	2	-	-
2002.....	19	7	-	1	-	3	2	2	1	-	1	1
2001.....	21	4	5	2	-	2	-	1	1	1	1	1
2000.....	34	9	3	2	-	3	6	3	2	-	-	2
1999.....	27	7	1	1	2	-	2	2	4	3	2	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningococcal infection (A39)</b>												
<b>Hispanic, male</b>												
2003.....	13	4	1	-	-	-	1	1	2	1	-	-
2002.....	9	3	-	1	-	1	1	1	1	-	-	-
2001.....	11	-	3	2	-	2	-	1	-	-	1	1
2000.....	16	4	2	-	-	2	4	2	-	-	-	1
1999.....	11	5	1	1	1	-	1	1	1	-	-	-
<b>Hispanic, female</b>												
2003.....	10	1	1	-	1	1	-	2	-	1	-	-
2002.....	10	4	-	-	-	2	1	1	-	-	1	1
2001.....	10	4	2	-	-	-	-	-	1	1	-	-
2000.....	18	5	1	2	-	1	2	1	2	-	-	1
1999.....	16	2	-	-	1	-	1	1	3	3	2	2
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	138	7	16	2	8	17	20	5	2	8	9	8
2002.....	142	10	12	5	14	18	10	10	3	4	4	9
2001.....	178	12	14	10	14	41	12	5	3	7	4	8
2000.....	177	13	16	10	6	31	16	6	4	7	6	11
1999.....	198	25	28	12	10	28	9	9	8	9	13	8
<b>Non-Hispanic 1/, male</b>												
2003.....	62	5	9	1	2	8	9	2	2	3	5	5
2002.....	74	5	9	4	8	10	5	7	1	2	3	5
2001.....	83	6	6	4	9	18	7	3	2	3	2	5
2000.....	90	10	10	5	2	22	7	1	3	6	3	4
1999.....	89	9	17	6	5	13	5	7	3	3	6	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningococcal infection (A39)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	76	2	7	1	6	9	11	3	-	5	4	3
2002.....	68	5	3	1	6	8	5	3	2	2	1	4
2001.....	95	6	8	6	5	23	5	2	1	4	2	3
2000.....	87	3	6	5	4	9	9	5	1	1	3	7
1999.....	109	16	11	6	5	15	4	2	5	6	7	7
<b>Non-Hispanic White, both sexes</b>												
2003.....	99	7	10	1	5	13	15	3	-	7	5	6
2002.....	106	7	8	4	5	14	8	7	3	1	3	8
2001.....	138	10	10	6	12	29	9	4	1	5	4	7
2000.....	129	10	11	3	5	27	10	5	1	3	4	5
1999.....	155	23	24	9	6	20	7	4	4	5	9	7
<b>Non-Hispanic White, male</b>												
2003.....	44	5	6	-	2	6	7	2	-	3	3	4
2002.....	58	5	6	4	4	7	5	6	1	1	2	5
2001.....	62	5	5	2	8	13	5	2	1	1	2	4
2000.....	60	7	6	2	2	18	4	1	-	2	3	1
1999.....	67	9	15	4	3	8	4	3	1	2	4	1
<b>Non-Hispanic White, female</b>												
2003.....	55	2	4	1	3	7	8	1	-	4	2	2
2002.....	48	2	2	-	1	7	3	1	2	-	1	3
2001.....	76	5	5	4	4	16	4	2	-	4	2	3
2000.....	69	3	5	1	3	9	6	4	1	1	1	4
1999.....	88	14	9	5	3	12	3	1	3	3	5	6





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Septicemia (A40-A41)</b>												
All origins, both sexes												
2003.....	34,069	278	85	39	38	59	95	123	186	315	595	916
2002.....	33,865	296	79	42	53	49	69	119	195	318	538	904
2001.....	32,238	312	108	29	40	57	76	96	185	309	520	837
2000.....	31,224	274	99	38	25	40	60	104	176	324	553	781
1999.....	30,680	283	87	47	30	32	89	106	169	314	498	731
All origins, male												
2003.....	14,987	151	54	15	19	35	47	56	105	167	288	498
2002.....	14,947	158	41	26	26	22	37	63	103	159	282	495
2001.....	14,307	174	60	17	22	31	44	50	89	170	281	439
2000.....	13,537	139	53	24	14	23	37	62	96	162	289	439
1999.....	13,395	175	48	22	14	17	50	53	94	170	290	406
All origins, female												
2003.....	19,082	127	31	24	19	24	48	67	81	148	307	418
2002.....	18,918	138	38	16	27	27	32	56	92	159	256	409
2001.....	17,931	138	48	12	18	26	32	46	96	139	239	398
2000.....	17,687	135	46	14	11	17	23	42	80	162	264	342
1999.....	17,285	108	39	25	16	15	39	53	75	144	208	325
Hispanic, both sexes												
2003.....	1,505	50	10	5	2	7	10	12	23	31	55	74
2002.....	1,521	65	16	7	7	7	8	11	27	27	40	56
2001.....	1,386	50	18	8	10	5	8	9	14	28	44	70
2000.....	1,226	62	15	3	4	3	9	17	14	28	51	51
1999.....	1,215	43	14	6	2	5	4	7	19	30	44	35

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Septicemia (A40-A41)</b>												
<b>Hispanic, male</b>												
2003.....	721	29	6	3	1	4	4	3	17	18	30	48
2002.....	744	34	9	6	2	5	3	8	17	17	30	40
2001.....	684	29	5	7	7	4	5	5	10	20	23	37
2000.....	625	33	9	3	2	2	4	9	12	16	30	36
1999.....	605	27	8	3	1	3	2	3	11	20	31	22
<b>Hispanic, female</b>												
2003.....	784	21	4	2	1	3	6	9	6	13	25	26
2002.....	777	31	7	1	5	2	5	3	10	10	10	16
2001.....	702	21	13	1	3	1	3	4	4	8	21	33
2000.....	601	29	6	-	2	1	5	8	2	12	21	15
1999.....	610	16	6	3	1	2	2	4	8	10	13	13
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	32,476	227	75	34	36	52	85	110	162	284	539	842
2002.....	32,225	227	63	35	46	42	60	107	168	291	496	843
2001.....	30,754	261	90	21	29	52	68	87	170	280	475	766
2000.....	29,864	208	83	35	21	37	51	86	162	294	499	728
1999.....	29,351	239	73	41	28	27	85	98	150	283	453	695
<b>Non-Hispanic 1/, male</b>												
2003.....	14,226	122	48	12	18	31	43	52	87	149	257	450
2002.....	14,136	120	32	20	24	17	33	55	86	142	250	451
2001.....	13,573	145	55	10	15	27	39	45	78	150	258	402
2000.....	12,842	104	44	21	12	21	33	53	84	144	257	402
1999.....	12,737	147	40	19	13	14	48	49	83	149	259	384

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Septicemia (A40-A41)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	18,250	105	27	22	18	21	42	58	75	135	282	392
2002.....	18,089	107	31	15	22	25	27	52	82	149	246	392
2001.....	17,181	116	35	11	14	25	29	42	92	130	217	364
2000.....	17,022	104	39	14	9	16	18	33	78	150	242	326
1999.....	16,614	92	33	22	15	13	37	49	67	134	194	311
<b>Non-Hispanic White, both sexes</b>												
2003.....	25,664	96	43	21	26	30	56	62	93	171	342	544
2002.....	25,545	129	42	24	31	31	41	63	102	184	326	572
2001.....	24,389	122	63	17	18	32	37	52	93	162	296	478
2000.....	23,517	108	51	25	17	21	27	44	90	175	323	445
1999.....	23,023	121	47	29	18	21	50	58	97	165	274	421
<b>Non-Hispanic White, male</b>												
2003.....	11,181	46	29	8	14	21	27	28	46	94	161	298
2002.....	11,148	67	25	15	19	11	26	37	49	87	162	312
2001.....	10,668	71	41	7	11	18	19	28	41	95	171	250
2000.....	10,056	55	26	16	9	13	20	25	49	93	169	254
1999.....	9,863	79	25	15	9	11	33	33	54	89	154	234
<b>Non-Hispanic White, female</b>												
2003.....	14,483	50	14	13	12	9	29	34	47	77	181	246
2002.....	14,397	62	17	9	12	20	15	26	53	97	164	260
2001.....	13,721	51	22	10	7	14	18	24	52	67	125	228
2000.....	13,461	53	25	9	8	8	7	19	41	82	154	191
1999.....	13,160	42	22	14	9	10	17	25	43	76	120	187

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Septicemia (A40-A41)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	6,141	119	24	10	5	18	23	42	60	101	182	274
2002.....	6,074	85	17	7	13	9	15	41	61	96	158	238
2001.....	5,829	124	21	2	8	19	22	28	70	109	159	261
2000.....	5,863	96	25	8	3	12	21	39	68	109	160	258
1999.....	5,772	106	16	7	7	5	27	37	48	109	170	251
<b>Non-Hispanic Black, male</b>												
2003.....	2,748	70	14	4	2	8	11	20	36	48	86	143
2002.....	2,668	46	5	3	4	5	5	15	32	50	80	120
2001.....	2,611	67	12	1	2	9	16	13	33	51	77	133
2000.....	2,559	48	13	4	2	6	12	27	32	48	77	136
1999.....	2,594	60	10	4	3	3	11	14	27	55	99	136
<b>Non-Hispanic Black, female</b>												
2003.....	3,393	49	10	6	3	10	12	22	24	53	96	131
2002.....	3,406	39	12	4	9	4	10	26	29	46	78	118
2001.....	3,218	57	9	1	6	10	6	15	37	58	82	128
2000.....	3,304	48	12	4	1	6	9	12	36	61	83	122
1999.....	3,178	46	6	3	4	2	16	23	21	54	71	115
<b>Origin not stated 2/, both sexes</b>												
2003.....	88	1	-	-	-	-	-	1	1	-	1	-
2002.....	119	4	-	-	-	-	1	1	-	-	2	5
2001.....	98	1	-	-	1	-	-	-	1	1	1	1
2000.....	134	4	1	-	-	-	-	1	-	2	3	2
1999.....	114	1	-	-	-	-	-	1	-	1	1	1





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Syphilis (A50-A53)

Hispanic, male

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	5	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	1

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	2	-	-	-	-	-	-	-	-	-	1

Non-Hispanic 1/, both sexes

2003.....	32	-	-	-	-	-	-	-	-	1	2	2
2002.....	39	1	1	-	-	-	-	-	-	1	1	1
2001.....	28	-	-	-	1	-	-	-	1	3	-	2
2000.....	39	1	-	-	-	-	-	-	1	-	-	2
1999.....	27	-	-	-	-	-	-	-	1	-	1	1

Non-Hispanic 1/, male

2003.....	21	-	-	-	-	-	-	-	-	1	2	2
2002.....	20	-	1	-	-	-	-	-	-	1	1	-
2001.....	14	-	-	-	1	-	-	-	-	2	-	2
2000.....	24	1	-	-	-	-	-	-	1	-	-	1
1999.....	11	-	-	-	-	-	-	-	1	-	-	1





































Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Viral hepatitis (B15-B19)</b>												
All origins, both sexes												
2003.....	5,431	1	-	-	-	5	9	21	43	174	478	1,053
2002.....	5,793	2	3	-	2	12	11	37	57	216	524	1,192
2001.....	5,585	1	-	3	-	9	11	19	55	202	595	1,101
2000.....	5,357	6	1	2	-	4	14	25	54	219	560	1,044
1999.....	4,853	2	3	1	2	3	11	19	69	204	518	871
All origins, male												
2003.....	3,546	-	-	-	-	2	5	12	30	126	335	763
2002.....	3,770	1	3	-	1	4	9	20	43	154	365	887
2001.....	3,539	1	-	2	-	6	5	7	35	133	420	833
2000.....	3,381	2	1	1	-	2	9	13	33	151	389	787
1999.....	3,043	2	2	-	2	3	7	9	44	139	381	651
All origins, female												
2003.....	1,885	1	-	-	-	3	4	9	13	48	143	290
2002.....	2,023	1	-	-	1	8	2	17	14	62	159	305
2001.....	2,046	-	-	1	-	3	6	12	20	69	175	268
2000.....	1,976	4	-	1	-	2	5	12	21	68	171	257
1999.....	1,810	-	1	1	-	-	4	10	25	65	137	220
Hispanic, both sexes												
2003.....	710	-	-	-	-	2	-	6	4	29	78	124
2002.....	758	-	-	-	-	2	3	4	13	30	71	142
2001.....	791	-	-	2	-	3	3	2	8	35	91	155
2000.....	708	3	-	-	-	1	2	2	12	39	87	119
1999.....	656	-	1	-	-	-	3	3	13	38	66	99

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Viral hepatitis (B15-B19)</b>												
<b>Hispanic, male</b>												
2003.....	465	-	-	-	-	2	-	4	4	20	60	97
2002.....	491	-	-	-	-	2	3	4	10	22	53	113
2001.....	487	-	-	2	-	1	1	2	1	23	66	119
2000.....	456	1	-	-	-	1	1	2	9	30	70	91
1999.....	406	-	1	-	-	-	1	-	11	30	54	76
<b>Hispanic, female</b>												
2003.....	245	-	-	-	-	-	-	2	-	9	18	27
2002.....	267	-	-	-	-	-	-	-	3	8	18	29
2001.....	304	-	-	-	-	2	2	-	7	12	25	36
2000.....	252	2	-	-	-	-	1	-	3	9	17	28
1999.....	250	-	-	-	-	-	2	3	2	8	12	23
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	4,704	1	-	-	-	3	9	15	39	145	400	925
2002.....	5,004	2	3	-	2	10	8	33	44	184	450	1,044
2001.....	4,776	1	-	1	-	6	8	17	47	165	504	940
2000.....	4,629	3	1	2	-	3	12	23	42	180	472	920
1999.....	4,176	2	2	1	2	3	8	16	55	166	447	765
<b>Non-Hispanic 1/, male</b>												
2003.....	3,068	-	-	-	-	-	5	8	26	106	275	663
2002.....	3,256	1	3	-	1	2	6	16	33	130	309	768
2001.....	3,038	1	-	-	-	5	4	5	34	108	354	709
2000.....	2,909	1	1	1	-	1	8	11	24	121	319	692
1999.....	2,620	2	1	-	2	3	6	9	32	109	323	570

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Viral hepatitis (B15-B19)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	1,636	1	-	-	-	3	4	7	13	39	125	262
2002.....	1,748	1	-	-	1	8	2	17	11	54	141	276
2001.....	1,738	-	-	1	-	1	4	12	13	57	150	231
2000.....	1,720	2	-	1	-	2	4	12	18	59	153	228
1999.....	1,556	-	1	1	-	-	2	7	23	57	124	195
<b>Non-Hispanic White, both sexes</b>												
2003.....	3,585	-	-	-	-	3	4	11	28	117	308	753
2002.....	3,857	1	-	-	2	7	8	17	32	147	349	820
2001.....	3,663	1	-	1	-	3	4	10	33	131	391	742
2000.....	3,513	1	-	2	-	1	4	15	22	142	359	707
1999.....	3,176	1	2	1	1	3	3	8	43	131	333	580
<b>Non-Hispanic White, male</b>												
2003.....	2,381	-	-	-	-	-	1	6	16	84	216	549
2002.....	2,551	1	-	-	1	-	6	8	25	110	241	613
2001.....	2,410	1	-	-	-	3	1	3	25	88	280	570
2000.....	2,234	-	-	1	-	1	3	7	12	94	248	545
1999.....	2,026	1	1	-	1	3	3	4	25	88	245	423
<b>Non-Hispanic White, female</b>												
2003.....	1,204	-	-	-	-	3	3	5	12	33	92	204
2002.....	1,306	-	-	-	1	7	2	9	7	37	108	207
2001.....	1,253	-	-	1	-	-	3	7	8	43	111	172
2000.....	1,279	1	-	1	-	-	1	8	10	48	111	162
1999.....	1,150	-	1	1	-	-	-	4	18	43	88	157

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Viral hepatitis (B15-B19)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	842	-	-	-	-	-	4	2	9	20	77	136
2002.....	849	1	1	-	-	2	-	15	7	31	80	180
2001.....	819	-	-	-	-	2	4	7	8	27	94	162
2000.....	777	2	-	-	-	2	8	7	14	29	88	170
1999.....	718	1	-	-	1	-	4	6	11	25	94	160
<b>Non-Hispanic Black, male</b>												
2003.....	536	-	-	-	-	-	3	1	8	15	48	92
2002.....	532	-	1	-	-	1	-	7	4	16	51	121
2001.....	479	-	-	-	-	1	3	2	5	15	64	112
2000.....	490	1	-	-	-	-	5	3	8	21	56	119
1999.....	433	1	-	-	1	-	2	3	6	15	64	125
<b>Non-Hispanic Black, female</b>												
2003.....	306	-	-	-	-	-	1	1	1	5	29	44
2002.....	317	1	-	-	-	1	-	8	3	15	29	59
2001.....	340	-	-	-	-	1	1	5	3	12	30	50
2000.....	287	1	-	-	-	2	3	4	6	8	32	51
1999.....	285	-	-	-	-	-	2	3	5	10	30	35
<b>Origin not stated 2/, both sexes</b>												
2003.....	17	-	-	-	-	-	-	-	-	-	-	4
2002.....	31	-	-	-	-	-	-	-	-	2	3	6
2001.....	18	-	-	-	-	-	-	-	-	2	-	6
2000.....	20	-	-	-	-	-	-	-	-	-	1	5
1999.....	21	-	-	-	-	-	-	-	1	-	5	7



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
All origins, both sexes												
2003.....	13,658	5	2	15	21	32	146	444	1,144	2,401	2,939	2,636
2002.....	14,095	7	5	4	21	38	140	500	1,339	2,618	3,089	2,680
2001.....	14,175	10	13	17	21	42	183	584	1,517	2,741	3,126	2,511
2000.....	14,478	10	14	17	29	35	144	647	1,790	2,896	3,023	2,551
1999.....	14,802	14	24	38	28	31	167	763	1,966	2,981	3,251	2,440
All origins, male												
2003.....	10,119	2	2	7	12	15	80	272	766	1,710	2,174	2,013
2002.....	10,468	4	1	4	10	17	76	288	899	1,943	2,255	2,039
2001.....	10,590	7	8	11	12	21	84	354	1,067	2,042	2,318	1,942
2000.....	10,868	8	6	9	13	19	80	371	1,231	2,172	2,269	2,008
1999.....	11,234	9	16	20	16	15	80	504	1,410	2,224	2,482	1,936
All origins, female												
2003.....	3,539	3	-	8	9	17	66	172	378	691	765	623
2002.....	3,627	3	4	-	11	21	64	212	440	675	834	641
2001.....	3,585	3	5	6	9	21	99	230	450	699	808	569
2000.....	3,610	2	8	8	16	16	64	276	559	724	754	543
1999.....	3,568	5	8	18	12	16	87	259	556	757	769	504
Hispanic, both sexes												
2003.....	1,890	-	1	1	1	7	24	75	169	311	416	368
2002.....	1,855	4	-	-	3	6	23	80	182	353	426	320
2001.....	1,853	1	-	1	2	4	28	80	220	355	404	316
2000.....	1,908	-	-	5	4	7	16	98	267	382	380	322
1999.....	1,958	3	5	5	4	4	29	115	292	422	433	287

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
Hispanic, male												
2003.....	1,465	-	1	-	-	4	16	55	129	239	311	306
2002.....	1,440	3	-	-	3	1	18	59	142	281	326	251
2001.....	1,437	-	-	1	1	2	19	56	166	273	316	241
2000.....	1,493	-	-	3	2	4	14	64	210	309	294	252
1999.....	1,525	1	4	2	1	3	17	87	227	338	328	224
Hispanic, female												
2003.....	425	-	-	1	1	3	8	20	40	72	105	62
2002.....	415	1	-	-	-	5	5	21	40	72	100	69
2001.....	416	1	-	-	1	2	9	24	54	82	88	75
2000.....	415	-	-	2	2	3	2	34	57	73	86	70
1999.....	433	2	1	3	3	1	12	28	65	84	105	63
Non-Hispanic 1/, both sexes												
2003.....	11,629	5	1	14	20	25	122	362	971	2,065	2,496	2,245
2002.....	12,094	3	5	4	18	32	117	417	1,147	2,237	2,634	2,324
2001.....	12,141	9	13	16	19	38	155	499	1,280	2,363	2,681	2,163
2000.....	12,389	10	14	12	25	28	128	545	1,508	2,476	2,601	2,200
1999.....	12,638	11	19	33	23	27	138	640	1,661	2,535	2,770	2,108
Non-Hispanic 1/, male												
2003.....	8,550	2	1	7	12	11	64	212	634	1,454	1,845	1,687
2002.....	8,911	1	1	4	7	16	58	226	751	1,640	1,909	1,756
2001.....	9,011	7	8	10	11	19	65	295	892	1,754	1,971	1,671
2000.....	9,224	8	6	6	11	15	66	304	1,009	1,834	1,944	1,728
1999.....	9,538	8	12	18	15	12	63	412	1,172	1,867	2,117	1,673

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic 1/, female												
2003.....	3,079	3	-	7	8	14	58	150	337	611	651	558
2002.....	3,183	2	4	-	11	16	59	191	396	597	725	568
2001.....	3,130	2	5	6	8	19	90	204	388	609	710	492
2000.....	3,165	2	8	6	14	13	62	241	499	642	657	472
1999.....	3,100	3	7	15	8	15	75	228	489	668	653	435
Non-Hispanic White, both sexes												
2003.....	4,126	1	-	1	2	5	23	63	312	732	971	828
2002.....	4,224	-	-	-	3	6	17	105	324	842	948	864
2001.....	4,286	2	3	6	2	2	28	110	409	914	997	771
2000.....	4,562	1	1	3	10	6	24	139	480	1,007	993	826
1999.....	4,744	2	2	7	8	6	28	188	606	1,044	1,038	766
Non-Hispanic White, male												
2003.....	3,487	1	-	-	2	2	15	50	244	603	826	702
2002.....	3,582	-	-	-	1	4	8	77	258	710	807	739
2001.....	3,669	2	-	4	1	-	15	79	331	781	858	674
2000.....	3,837	1	-	1	4	4	16	95	380	844	834	720
1999.....	4,043	2	1	4	6	4	18	139	504	880	888	666
Non-Hispanic White, female												
2003.....	639	-	-	1	-	3	8	13	68	129	145	126
2002.....	642	-	-	-	2	2	9	28	66	132	141	125
2001.....	617	-	3	2	1	2	13	31	78	133	139	97
2000.....	725	-	1	2	6	2	8	44	100	163	159	106
1999.....	701	-	1	3	2	2	10	49	102	164	150	100

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Human immunodeficiency virus (HIV) disease (B20-B24)												
Non-Hispanic Black, both sexes												
2003.....	7,351	3	1	13	18	20	97	295	643	1,291	1,495	1,388
2002.....	7,714	3	5	4	15	25	99	306	804	1,368	1,643	1,441
2001.....	7,710	7	10	9	17	34	127	381	853	1,417	1,655	1,372
2000.....	7,708	9	12	9	14	22	103	405	1,009	1,447	1,581	1,360
1999.....	7,751	9	17	26	15	21	109	446	1,033	1,458	1,696	1,322
Non-Hispanic Black, male												
2003.....	4,947	1	1	7	10	9	48	160	377	817	997	962
2002.....	5,203	1	1	4	6	11	49	145	479	911	1,067	999
2001.....	5,227	5	8	5	10	18	50	210	543	947	1,093	981
2000.....	5,291	7	6	5	7	11	49	208	614	973	1,087	995
1999.....	5,377	6	11	14	9	8	44	268	651	957	1,200	992
Non-Hispanic Black, female												
2003.....	2,404	2	-	6	8	11	49	135	266	474	498	426
2002.....	2,511	2	4	-	9	14	50	161	325	457	576	442
2001.....	2,483	2	2	4	7	16	77	171	310	470	562	391
2000.....	2,417	2	6	4	7	11	54	197	395	474	494	365
1999.....	2,374	3	6	12	6	13	65	178	382	501	496	330
Origin not stated 2/, both sexes												
2003.....	139	-	-	-	-	-	-	7	4	25	27	23
2002.....	146	-	-	-	-	-	-	3	10	28	29	36
2001.....	181	-	-	-	-	-	-	5	17	23	41	32
2000.....	181	-	-	-	-	-	-	4	15	38	42	29
1999.....	206	-	-	-	1	-	-	8	13	24	48	45

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Origin not stated 2/, male

2003.....	104	-	-	-	-	-	-	5	3	17	18	20
2002.....	117	-	-	-	-	-	-	3	6	22	20	32
2001.....	142	-	-	-	-	-	-	3	9	15	31	30
2000.....	151	-	-	-	-	-	-	3	12	29	31	28
1999.....	171	-	-	-	-	-	-	5	11	19	37	39

Origin not stated 2/, female

2003.....	35	-	-	-	-	-	-	2	1	8	9	3
2002.....	29	-	-	-	-	-	-	-	4	6	9	4
2001.....	39	-	-	-	-	-	-	2	8	8	10	2
2000.....	30	-	-	-	-	-	-	1	3	9	11	1
1999.....	35	-	-	-	1	-	-	3	2	5	11	6



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malaria (B50-B54)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	12	-	-	-	-	-	1	-	1	-	1	1
2001.....	9	-	-	-	1	-	-	-	-	2	-	2
2000.....	3	-	-	-	-	-	1	-	-	-	-	-
1999.....	7	1	-	-	-	-	-	2	-	-	1	-

Non-Hispanic 1/, male

2003.....	4	-	-	-	-	-	-	-	-	-	1	-
2002.....	8	-	-	-	-	-	-	-	-	-	1	1
2001.....	5	-	-	-	1	-	-	-	-	-	-	1
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	1	-	-	-	-	-	1	-	-	-	-







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins, both sexes

2003.....	7,427	174	85	29	22	49	39	54	86	158	289	526
2002.....	6,707	224	61	41	43	42	55	67	102	163	290	403
2001.....	5,715	191	64	34	30	40	46	58	89	162	207	307
2000.....	5,532	167	71	30	37	44	52	56	85	162	188	277
1999.....	7,279	203	81	40	39	43	47	92	150	234	311	392

All origins, male

2003.....	3,964	86	43	14	11	30	22	30	53	96	188	351
2002.....	3,622	125	39	16	24	20	32	38	58	97	194	278
2001.....	2,931	101	37	14	15	23	28	31	52	93	134	211
2000.....	2,861	94	41	16	19	25	29	29	46	105	115	164
1999.....	3,655	119	51	20	24	23	26	46	85	149	184	231

All origins, female

2003.....	3,463	88	42	15	11	19	17	24	33	62	101	175
2002.....	3,085	99	22	25	19	22	23	29	44	66	96	125
2001.....	2,784	90	27	20	15	17	18	27	37	69	73	96
2000.....	2,671	73	30	14	18	19	23	27	39	57	73	113
1999.....	3,624	84	30	20	15	20	21	46	65	85	127	161

Hispanic, both sexes

2003.....	675	23	17	7	7	9	3	7	14	27	50	71
2002.....	593	41	11	8	9	7	12	18	33	22	45	49
2001.....	423	41	7	6	5	6	6	14	19	24	22	40
2000.....	412	23	18	6	7	5	14	12	14	22	28	30
1999.....	497	36	13	8	6	5	8	17	22	37	30	33

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Hispanic, male

2003.....	416	9	8	4	3	7	1	6	8	21	36	54
2002.....	363	27	7	5	6	3	4	9	22	13	35	36
2001.....	250	21	4	1	4	4	5	9	13	16	15	26
2000.....	237	14	12	4	4	2	10	7	13	21	16	16
1999.....	260	22	10	4	3	3	3	10	12	29	20	17

Hispanic, female

2003.....	259	14	9	3	4	2	2	1	6	6	14	17
2002.....	230	14	4	3	3	4	8	9	11	9	10	13
2001.....	173	20	3	5	1	2	1	5	6	8	7	14
2000.....	175	9	6	2	3	3	4	5	1	1	12	14
1999.....	237	14	3	4	3	2	5	7	10	8	10	16

Non-Hispanic 1/, both sexes

2003.....	6,730	151	66	22	15	40	35	47	72	131	237	451
2002.....	6,097	181	49	33	34	35	42	48	69	141	243	354
2001.....	5,271	149	57	28	25	34	39	44	70	137	182	264
2000.....	5,100	144	53	24	30	37	38	44	71	139	158	247
1999.....	6,753	166	68	32	31	38	39	75	127	197	278	356

Non-Hispanic 1/, male

2003.....	3,536	77	33	10	8	23	20	24	45	75	152	296
2002.....	3,250	97	31	11	18	17	27	29	36	84	158	242
2001.....	2,666	79	33	13	11	19	23	22	39	76	117	182
2000.....	2,613	80	29	12	15	23	19	22	33	84	98	148
1999.....	3,379	96	41	16	20	20	23	36	72	120	162	213

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 1/, female

2003.....	3,194	74	33	12	7	17	15	23	27	56	85	155
2002.....	2,847	84	18	22	16	18	15	19	33	57	85	112
2001.....	2,605	70	24	15	14	15	16	22	31	61	65	82
2000.....	2,487	64	24	12	15	14	19	22	38	55	60	99
1999.....	3,374	70	27	16	11	18	16	39	55	77	116	143

Non-Hispanic White, both sexes

2003.....	5,509	87	36	15	10	27	24	30	46	83	174	348
2002.....	4,939	113	34	24	25	24	23	28	38	82	161	267
2001.....	4,257	80	37	22	16	21	24	25	38	87	124	194
2000.....	4,131	79	33	18	22	26	25	22	36	81	108	183
1999.....	5,288	101	43	24	18	27	24	39	65	121	166	227

Non-Hispanic White, male

2003.....	2,871	42	17	8	4	17	14	16	27	50	113	235
2002.....	2,608	57	21	9	15	13	14	15	18	52	109	185
2001.....	2,148	40	21	10	7	10	14	15	24	49	80	137
2000.....	2,092	45	16	8	10	14	15	12	16	47	68	111
1999.....	2,644	64	27	14	10	13	15	17	38	82	108	145

Non-Hispanic White, female

2003.....	2,638	45	19	7	6	10	10	14	19	33	61	113
2002.....	2,331	56	13	15	10	11	9	13	20	30	52	82
2001.....	2,109	40	16	12	9	11	10	10	14	38	44	57
2000.....	2,039	34	17	10	12	12	10	10	20	34	40	72
1999.....	2,644	37	16	10	8	14	9	22	27	39	58	82

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic Black, both sexes

2003.....	942	53	23	5	4	11	7	13	20	40	59	92
2002.....	927	58	13	5	5	9	16	18	28	52	69	79
2001.....	815	61	16	6	7	9	14	17	28	41	50	62
2000.....	815	60	14	5	5	8	13	16	34	48	47	61
1999.....	1,247	57	23	7	10	7	13	31	58	73	96	116

Non-Hispanic Black, male

2003.....	515	30	12	2	3	4	5	5	15	21	36	56
2002.....	516	33	9	1	2	3	10	13	15	27	41	52
2001.....	419	36	10	3	3	7	8	7	12	22	34	41
2000.....	428	31	10	3	2	6	4	6	16	30	28	35
1999.....	617	29	14	2	8	5	7	14	34	36	45	62

Non-Hispanic Black, female

2003.....	427	23	11	3	1	7	2	8	5	19	23	36
2002.....	411	25	4	4	3	6	6	5	13	25	28	27
2001.....	396	25	6	3	4	2	6	10	16	19	16	21
2000.....	387	29	4	2	3	2	9	10	18	18	19	26
1999.....	630	28	9	5	2	2	6	17	24	37	51	54

Origin not stated 2/, both sexes

2003.....	22	-	2	-	-	-	1	-	-	-	2	4
2002.....	17	2	1	-	-	-	1	1	-	-	2	-
2001.....	21	1	-	-	-	-	1	-	-	1	3	3
2000.....	20	-	-	-	-	2	-	-	-	1	2	-
1999.....	29	1	-	-	2	-	-	-	1	-	3	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Origin not stated 2/, male

2003.....	12	-	2	-	-	-	1	-	-	-	-	1
2002.....	9	1	1	-	-	-	1	-	-	-	1	-
2001.....	15	1	-	-	-	-	-	-	-	1	2	3
2000.....	11	-	-	-	-	-	-	-	-	-	1	-
1999.....	16	1	-	-	1	-	-	-	1	-	2	1

Origin not stated 2/, female

2003.....	10	-	-	-	-	-	-	-	-	-	2	3
2002.....	8	1	-	-	-	-	-	1	-	-	1	-
2001.....	6	-	-	-	-	-	1	-	-	-	1	-
2000.....	9	-	-	-	-	2	-	-	-	1	1	-
1999.....	13	-	-	-	1	-	-	-	-	-	1	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms (C00-C97)</b>												
<b>All origins, both sexes</b>												
2003.....	556,902	75	392	516	560	690	961	1,307	2,434	4,919	10,590	19,552
2002.....	557,271	74	402	537	535	723	1,007	1,258	2,614	5,150	10,935	19,295
2001.....	553,768	66	420	493	515	732	972	1,389	2,605	5,466	11,103	19,199
2000.....	553,091	92	420	489	525	745	968	1,359	2,557	5,565	10,955	18,803
1999.....	549,838	68	418	509	503	745	979	1,429	2,576	5,764	10,968	18,412
<b>All origins, male</b>												
2003.....	287,990	34	227	293	298	438	541	664	1,140	2,070	4,755	9,727
2002.....	288,768	42	216	297	303	424	592	678	1,190	2,166	4,890	9,418
2001.....	287,075	31	229	253	280	447	577	694	1,166	2,331	4,966	9,377
2000.....	286,082	50	233	269	300	446	576	692	1,162	2,402	4,942	9,243
1999.....	285,832	36	227	279	276	448	575	755	1,204	2,467	5,004	9,151
<b>All origins, female</b>												
2003.....	268,912	41	165	223	262	252	420	643	1,294	2,849	5,835	9,825
2002.....	268,503	32	186	240	232	299	415	580	1,424	2,984	6,045	9,877
2001.....	266,693	35	191	240	235	285	395	695	1,439	3,135	6,137	9,822
2000.....	267,009	42	187	220	225	299	392	667	1,395	3,163	6,013	9,560
1999.....	264,006	32	191	230	227	297	404	674	1,372	3,297	5,964	9,261
<b>Hispanic, both sexes</b>												
2003.....	24,070	13	98	111	106	124	164	208	314	513	842	1,316
2002.....	23,141	18	72	103	107	121	183	177	325	500	846	1,190
2001.....	22,371	14	84	87	90	143	146	204	293	545	851	1,186
2000.....	21,160	17	75	91	107	122	153	196	280	515	780	1,141
1999.....	20,233	16	72	87	89	94	165	189	277	471	733	1,093

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms (C00-C97)</b>												
<b>Hispanic, male</b>												
2003.....	12,671	4	60	70	53	91	95	116	158	214	352	647
2002.....	12,235	10	35	51	57	81	110	103	149	198	359	566
2001.....	11,825	4	44	49	41	95	104	100	133	236	379	579
2000.....	11,138	10	43	55	67	72	90	105	136	204	330	540
1999.....	10,668	11	34	45	47	55	103	98	124	198	306	539
<b>Hispanic, female</b>												
2003.....	11,399	9	38	41	53	33	69	92	156	299	490	669
2002.....	10,906	8	37	52	50	40	73	74	176	302	487	624
2001.....	10,546	10	40	38	49	48	42	104	160	309	472	607
2000.....	10,022	7	32	36	40	50	63	91	144	311	450	601
1999.....	9,565	5	38	42	42	39	62	91	153	273	427	554
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	531,638	62	293	404	453	565	794	1,096	2,111	4,397	9,723	18,189
2002.....	532,712	55	329	433	428	598	820	1,077	2,273	4,636	10,057	18,043
2001.....	530,058	52	333	404	424	588	825	1,179	2,306	4,901	10,220	17,961
2000.....	530,531	75	341	395	417	619	813	1,159	2,272	5,036	10,134	17,602
1999.....	528,218	52	343	422	413	650	812	1,236	2,289	5,281	10,201	17,265
<b>Non-Hispanic 1/, male</b>												
2003.....	274,651	30	166	222	245	346	444	546	974	1,853	4,390	9,057
2002.....	275,770	31	181	246	246	342	479	573	1,037	1,962	4,513	8,817
2001.....	274,492	27	183	203	239	352	472	593	1,030	2,084	4,568	8,775
2000.....	274,185	40	186	214	232	372	486	584	1,023	2,192	4,595	8,675
1999.....	274,419	25	190	234	229	393	471	655	1,077	2,266	4,680	8,581

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms (C00-C97)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	256,987	32	127	182	208	219	350	550	1,137	2,544	5,333	9,132
2002.....	256,942	24	148	187	182	256	341	504	1,236	2,674	5,544	9,226
2001.....	255,566	25	150	201	185	236	353	586	1,276	2,817	5,652	9,186
2000.....	256,346	35	155	181	185	247	327	575	1,249	2,844	5,539	8,927
1999.....	253,799	27	153	188	184	257	341	581	1,212	3,015	5,521	8,684
<b>Non-Hispanic White, both sexes</b>												
2003.....	457,127	51	221	315	322	442	597	808	1,546	3,308	7,415	13,917
2002.....	458,754	37	236	331	311	454	643	795	1,688	3,509	7,725	13,970
2001.....	456,709	42	268	302	305	448	628	865	1,729	3,728	7,823	13,929
2000.....	458,200	60	256	307	320	452	607	884	1,715	3,802	7,656	13,598
1999.....	456,315	37	281	311	297	476	594	953	1,729	4,025	7,823	13,404
<b>Non-Hispanic White, male</b>												
2003.....	236,150	25	123	177	171	269	338	426	733	1,467	3,395	6,970
2002.....	237,323	24	134	182	173	264	375	427	791	1,511	3,530	6,912
2001.....	236,004	21	144	146	164	270	362	449	811	1,621	3,539	6,840
2000.....	235,932	34	146	161	177	270	366	452	807	1,705	3,496	6,745
1999.....	236,384	20	160	180	165	283	346	510	831	1,789	3,636	6,622
<b>Non-Hispanic White, female</b>												
2003.....	220,977	26	98	138	151	173	259	382	813	1,841	4,020	6,947
2002.....	221,431	13	102	149	138	190	268	368	897	1,998	4,195	7,058
2001.....	220,705	21	124	156	141	178	266	416	918	2,107	4,284	7,089
2000.....	222,268	26	110	146	143	182	241	432	908	2,097	4,160	6,853
1999.....	219,931	17	121	131	132	193	248	443	898	2,236	4,187	6,782

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms (C00-C97)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	62,054	7	55	73	100	93	152	226	430	873	1,914	3,604
2002.....	61,996	12	71	81	87	110	132	211	452	907	1,963	3,420
2001.....	61,580	5	46	78	94	110	147	241	450	949	2,001	3,370
2000.....	61,390	11	66	71	76	135	162	213	428	1,006	2,066	3,398
1999.....	61,409	11	47	84	95	136	178	227	433	1,049	2,000	3,278
<b>Non-Hispanic Black, male</b>												
2003.....	32,131	4	30	40	56	56	85	89	174	288	824	1,767
2002.....	32,287	4	33	52	51	64	73	102	185	345	813	1,612
2001.....	32,335	2	29	43	57	65	83	106	161	365	836	1,628
2000.....	32,493	4	31	42	44	82	96	93	166	377	915	1,648
1999.....	32,542	3	24	40	54	84	97	113	181	382	873	1,687
<b>Non-Hispanic Black, female</b>												
2003.....	29,923	3	25	33	44	37	67	137	256	585	1,090	1,837
2002.....	29,709	8	38	29	36	46	59	109	267	562	1,150	1,808
2001.....	29,245	3	17	35	37	45	64	135	289	584	1,165	1,742
2000.....	28,897	7	35	29	32	53	66	120	262	629	1,151	1,750
1999.....	28,867	8	23	44	41	52	81	114	252	667	1,127	1,591
<b>Origin not stated 2/, both sexes</b>												
2003.....	1,194	-	1	1	1	1	3	3	9	9	25	47
2002.....	1,418	1	1	1	-	4	4	4	16	14	32	62
2001.....	1,339	-	3	2	1	1	1	6	6	20	32	52
2000.....	1,400	-	4	3	1	4	2	4	5	14	41	60
1999.....	1,387	-	3	-	1	1	2	4	10	12	34	54

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms (C00-C97)</b>												
Origin not stated 2/, male												
2003.....	668	-	1	1	-	1	2	2	8	3	13	23
2002.....	763	1	-	-	-	1	3	2	4	6	18	35
2001.....	758	-	2	1	-	-	1	1	3	11	19	23
2000.....	759	-	4	-	1	2	-	3	3	6	17	28
1999.....	745	-	3	-	-	-	1	2	3	3	18	31
Origin not stated 2/, female												
2003.....	526	-	-	-	1	-	1	1	1	6	12	24
2002.....	655	-	1	1	-	3	1	2	12	8	14	27
2001.....	581	-	1	1	1	1	-	5	3	9	13	29
2000.....	641	-	-	3	-	2	2	1	2	8	24	32
1999.....	642	-	-	-	1	1	1	2	7	9	16	23

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All origins, both sexes												
2003.....	7,778	-	4	4	4	11	19	19	40	64	210	445
2002.....	7,737	-	3	1	7	12	11	22	35	77	181	431
2001.....	7,702	-	-	2	2	10	19	19	37	93	190	427
2000.....	7,492	-	-	-	4	9	17	18	32	88	205	408
1999.....	7,486	-	2	1	5	9	18	14	29	96	195	449
All origins, male												
2003.....	5,298	-	2	1	3	8	13	11	28	40	155	346
2002.....	5,184	-	2	1	2	7	4	14	25	47	132	336
2001.....	5,156	-	-	1	-	3	9	12	23	61	141	325
2000.....	4,952	-	-	-	4	6	7	12	16	60	155	309
1999.....	4,944	-	1	-	3	6	14	10	17	58	146	346
All origins, female												
2003.....	2,480	-	2	3	1	3	6	8	12	24	55	99
2002.....	2,553	-	1	-	5	5	7	8	10	30	49	95
2001.....	2,546	-	-	1	2	7	10	7	14	32	49	102
2000.....	2,540	-	-	-	-	3	10	6	16	28	50	99
1999.....	2,542	-	1	1	2	3	4	4	12	38	49	103
Hispanic, both sexes												
2003.....	321	-	1	-	-	-	4	2	3	3	11	25
2002.....	284	-	-	-	2	2	1	1	3	4	13	16
2001.....	311	-	-	-	-	1	3	-	1	6	8	25
2000.....	287	-	-	-	1	-	1	2	4	5	8	15
1999.....	282	-	-	-	1	-	-	-	1	9	9	19

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)</b>												
<b>Hispanic, male</b>												
2003.....	233	-	1	-	-	-	2	1	3	1	7	18
2002.....	215	-	-	-	-	1	-	-	3	4	10	14
2001.....	219	-	-	-	-	1	1	-	1	3	8	17
2000.....	218	-	-	-	1	-	-	1	3	3	8	13
1999.....	204	-	-	-	1	-	-	-	-	4	5	16
<b>Hispanic, female</b>												
2003.....	88	-	-	-	-	-	2	1	-	2	4	7
2002.....	69	-	-	-	2	1	1	1	-	-	3	2
2001.....	92	-	-	-	-	-	2	-	-	3	-	8
2000.....	69	-	-	-	-	-	1	1	1	2	-	2
1999.....	78	-	-	-	-	-	-	-	1	5	4	3
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	7,437	-	3	4	4	11	15	17	36	61	199	420
2002.....	7,431	-	3	1	5	10	10	21	32	73	165	413
2001.....	7,370	-	-	2	2	9	16	19	36	87	181	402
2000.....	7,181	-	-	-	3	9	15	16	28	83	197	390
1999.....	7,181	-	2	1	4	9	18	14	28	86	186	429
<b>Non-Hispanic 1/, male</b>												
2003.....	5,050	-	1	1	3	8	11	10	24	39	148	328
2002.....	4,950	-	2	1	2	6	4	14	22	43	119	320
2001.....	4,920	-	-	1	-	2	8	12	22	58	132	308
2000.....	4,719	-	-	-	3	6	7	11	13	57	147	294
1999.....	4,725	-	1	-	2	6	14	10	17	53	141	329

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic 1/, female												
2003.....	2,387	-	2	3	1	3	4	7	12	22	51	92
2002.....	2,481	-	1	-	3	4	6	7	10	30	46	93
2001.....	2,450	-	-	1	2	7	8	7	14	29	49	94
2000.....	2,462	-	-	-	-	3	8	5	15	26	50	96
1999.....	2,456	-	1	1	2	3	4	4	11	33	45	100
Non-Hispanic White, both sexes												
2003.....	6,079	-	2	4	1	5	10	12	24	44	138	297
2002.....	6,141	-	2	1	2	6	6	13	22	48	116	286
2001.....	6,052	-	-	1	1	7	11	14	22	49	116	272
2000.....	5,843	-	-	-	1	6	10	10	19	55	131	275
1999.....	5,826	-	2	1	2	5	9	10	19	51	125	269
Non-Hispanic White, male												
2003.....	4,043	-	-	1	1	4	7	7	14	31	114	225
2002.....	4,035	-	2	1	1	4	3	9	16	31	83	224
2001.....	3,961	-	-	1	-	1	5	10	12	34	86	211
2000.....	3,726	-	-	-	1	3	4	7	9	35	93	209
1999.....	3,746	-	1	-	1	3	6	6	11	30	95	203
Non-Hispanic White, female												
2003.....	2,036	-	2	3	-	1	3	5	10	13	24	72
2002.....	2,106	-	-	-	1	2	3	4	6	17	33	62
2001.....	2,091	-	-	-	1	6	6	4	10	15	30	61
2000.....	2,117	-	-	-	-	3	6	3	10	20	38	66
1999.....	2,080	-	1	1	1	2	3	4	8	21	30	66

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Non-Hispanic Black, both sexes												
2003.....	1,100	-	1	-	3	5	4	4	6	14	49	104
2002.....	1,034	-	-	-	2	4	2	3	8	22	41	106
2001.....	1,056	-	-	1	-	1	3	2	4	24	47	105
2000.....	1,089	-	-	-	2	3	4	5	2	15	45	94
1999.....	1,131	-	-	-	2	4	4	2	6	25	42	133
Non-Hispanic Black, male												
2003.....	831	-	1	-	2	3	3	3	6	6	26	86
2002.....	757	-	-	-	1	2	1	2	5	10	31	80
2001.....	773	-	-	-	-	1	1	1	4	14	31	79
2000.....	821	-	-	-	2	3	2	3	1	11	35	67
1999.....	819	-	-	-	1	3	3	2	4	17	30	107
Non-Hispanic Black, female												
2003.....	269	-	-	-	1	2	1	1	-	8	23	18
2002.....	277	-	-	-	1	2	1	1	3	12	10	26
2001.....	283	-	-	1	-	-	2	1	-	10	16	26
2000.....	268	-	-	-	-	-	2	2	1	4	10	27
1999.....	312	-	-	-	1	1	1	-	2	8	12	26
Origin not stated 2/, both sexes												
2003.....	20	-	-	-	-	-	-	-	1	-	-	-
2002.....	22	-	-	-	-	-	-	-	-	-	3	2
2001.....	21	-	-	-	-	-	-	-	-	-	1	-
2000.....	24	-	-	-	-	-	1	-	-	-	-	3
1999.....	23	-	-	-	-	-	-	-	-	1	-	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of esophagus (C15)</b>												
<b>All origins, both sexes</b>												
2003.....	12,860	-	-	-	-	2	3	11	18	76	212	526
2002.....	12,701	-	-	-	1	-	6	4	15	84	227	503
2001.....	12,530	-	-	-	-	2	4	9	20	68	241	492
2000.....	12,232	1	-	-	-	1	2	9	30	78	219	478
1999.....	11,918	-	-	-	-	-	-	10	27	75	218	469
<b>All origins, male</b>												
2003.....	9,921	-	-	-	-	2	2	8	15	66	174	445
2002.....	9,789	-	-	-	-	-	3	4	10	71	186	426
2001.....	9,607	-	-	-	-	1	4	8	18	52	203	417
2000.....	9,279	1	-	-	-	-	2	7	24	69	196	415
1999.....	9,094	-	-	-	-	-	-	7	22	65	190	410
<b>All origins, female</b>												
2003.....	2,939	-	-	-	-	-	1	3	3	10	38	81
2002.....	2,912	-	-	-	1	-	3	-	5	13	41	77
2001.....	2,923	-	-	-	-	1	-	1	2	16	38	75
2000.....	2,953	-	-	-	-	1	-	2	6	9	23	63
1999.....	2,824	-	-	-	-	-	-	3	5	10	28	59
<b>Hispanic, both sexes</b>												
2003.....	421	-	-	-	-	-	1	3	-	-	17	20
2002.....	411	-	-	-	-	-	1	-	1	5	18	22
2001.....	401	-	-	-	-	-	-	2	-	3	9	21
2000.....	384	-	-	-	-	-	-	-	2	3	7	15
1999.....	367	-	-	-	-	-	-	-	2	6	10	22

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of esophagus (C15)</b>												
<b>Hispanic, male</b>												
2003.....	322	-	-	-	-	-	-	2	-	-	16	19
2002.....	329	-	-	-	-	-	1	-	1	5	14	22
2001.....	316	-	-	-	-	-	-	2	-	2	7	19
2000.....	292	-	-	-	-	-	-	-	2	3	5	13
1999.....	294	-	-	-	-	-	-	-	2	5	10	20
<b>Hispanic, female</b>												
2003.....	99	-	-	-	-	-	1	1	-	-	1	1
2002.....	82	-	-	-	-	-	-	-	-	-	4	-
2001.....	85	-	-	-	-	-	-	-	-	1	2	2
2000.....	92	-	-	-	-	-	-	-	-	-	2	2
1999.....	73	-	-	-	-	-	-	-	-	1	-	2
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	12,415	-	-	-	-	2	2	8	18	76	195	506
2002.....	12,254	-	-	-	1	-	5	3	14	79	208	480
2001.....	12,085	-	-	-	-	2	4	7	20	65	231	468
2000.....	11,819	1	-	-	-	1	2	9	28	75	212	463
1999.....	11,520	-	-	-	-	-	-	10	25	69	207	447
<b>Non-Hispanic 1/, male</b>												
2003.....	9,580	-	-	-	-	2	2	6	15	66	158	426
2002.....	9,431	-	-	-	-	-	2	3	9	66	171	403
2001.....	9,254	-	-	-	-	1	4	6	18	50	195	395
2000.....	8,964	1	-	-	-	-	2	7	22	66	191	402
1999.....	8,775	-	-	-	-	-	-	7	20	60	179	390

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of esophagus (C15)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	2,835	-	-	-	-	-	-	2	3	10	37	80
2002.....	2,823	-	-	-	1	-	3	-	5	13	37	77
2001.....	2,831	-	-	-	-	1	-	1	2	15	36	73
2000.....	2,855	-	-	-	-	1	-	2	6	9	21	61
1999.....	2,745	-	-	-	-	-	-	3	5	9	28	57
<b>Non-Hispanic White, both sexes</b>												
2003.....	10,631	-	-	-	-	2	2	6	16	65	157	387
2002.....	10,484	-	-	-	1	-	5	2	13	68	159	382
2001.....	10,193	-	-	-	-	2	4	5	17	56	182	365
2000.....	10,005	-	-	-	-	1	2	9	27	71	168	362
1999.....	9,703	-	-	-	-	-	-	9	21	61	152	340
<b>Non-Hispanic White, male</b>												
2003.....	8,292	-	-	-	-	2	2	5	14	58	130	331
2002.....	8,169	-	-	-	-	-	2	2	9	60	136	332
2001.....	7,863	-	-	-	-	1	4	4	15	43	157	316
2000.....	7,680	-	-	-	-	-	2	7	22	64	157	328
1999.....	7,440	-	-	-	-	-	-	6	16	55	133	303
<b>Non-Hispanic White, female</b>												
2003.....	2,339	-	-	-	-	-	-	1	2	7	27	56
2002.....	2,315	-	-	-	1	-	3	-	4	8	23	50
2001.....	2,330	-	-	-	-	1	-	1	2	13	25	49
2000.....	2,325	-	-	-	-	1	-	2	5	7	11	34
1999.....	2,263	-	-	-	-	-	-	3	5	6	19	37





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of stomach (C16)</b>												
<b>All origins, both sexes</b>												
2003.....	12,110	-	-	-	1	5	14	34	93	178	302	493
2002.....	12,198	-	1	-	-	7	22	48	90	160	276	443
2001.....	12,319	-	-	-	-	5	15	33	89	178	286	452
2000.....	12,645	-	-	-	1	3	11	36	88	167	317	468
1999.....	12,711	-	-	-	1	5	8	47	89	189	316	450
<b>All origins, male</b>												
2003.....	7,055	-	-	-	1	3	4	16	47	91	171	305
2002.....	7,133	-	1	-	-	-	8	31	49	82	157	285
2001.....	7,191	-	-	-	-	4	5	20	51	98	155	288
2000.....	7,328	-	-	-	-	-	6	22	47	100	184	290
1999.....	7,459	-	-	-	1	2	4	22	54	101	182	303
<b>All origins, female</b>												
2003.....	5,055	-	-	-	-	2	10	18	46	87	131	188
2002.....	5,065	-	-	-	-	7	14	17	41	78	119	158
2001.....	5,128	-	-	-	-	1	10	13	38	80	131	164
2000.....	5,317	-	-	-	1	3	5	14	41	67	133	178
1999.....	5,252	-	-	-	-	3	4	25	35	88	134	147
<b>Hispanic, both sexes</b>												
2003.....	1,253	-	-	-	1	2	1	11	23	44	48	88
2002.....	1,185	-	-	-	-	1	5	11	27	45	52	71
2001.....	1,189	-	-	-	-	1	3	16	25	40	58	78
2000.....	1,111	-	-	-	1	-	4	10	14	48	57	73
1999.....	1,055	-	-	-	-	1	3	14	24	34	51	50

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of stomach (C16)</b>												
<b>Hispanic, male</b>												
2003.....	708	-	-	-	1	1	1	5	12	23	23	51
2002.....	671	-	-	-	-	-	2	8	18	15	29	42
2001.....	680	-	-	-	-	1	1	7	12	23	31	43
2000.....	655	-	-	-	-	-	1	7	9	26	26	35
1999.....	607	-	-	-	-	-	1	8	17	14	27	35
<b>Hispanic, female</b>												
2003.....	545	-	-	-	-	1	-	6	11	21	25	37
2002.....	514	-	-	-	-	1	3	3	9	30	23	29
2001.....	509	-	-	-	-	-	2	9	13	17	27	35
2000.....	456	-	-	-	1	-	3	3	5	22	31	38
1999.....	448	-	-	-	-	1	2	6	7	20	24	15
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	10,827	-	-	-	-	3	13	22	69	134	254	404
2002.....	10,974	-	1	-	-	6	17	36	63	114	223	372
2001.....	11,099	-	-	-	-	4	12	17	64	138	227	374
2000.....	11,491	-	-	-	-	3	7	25	74	119	259	394
1999.....	11,615	-	-	-	1	4	5	33	65	155	265	395
<b>Non-Hispanic 1/, male</b>												
2003.....	6,326	-	-	-	-	2	3	11	34	68	148	253
2002.....	6,438	-	1	-	-	-	6	23	31	66	128	243
2001.....	6,492	-	-	-	-	3	4	13	39	75	124	245
2000.....	6,648	-	-	-	-	-	5	15	38	74	157	254
1999.....	6,827	-	-	-	1	2	3	14	37	87	155	263

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of stomach (C16)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	4,501	-	-	-	-	1	10	11	35	66	106	151
2002.....	4,536	-	-	-	-	6	11	13	32	48	95	129
2001.....	4,607	-	-	-	-	1	8	4	25	63	103	129
2000.....	4,843	-	-	-	-	3	2	10	36	45	102	140
1999.....	4,788	-	-	-	-	2	2	19	28	68	110	132
<b>Non-Hispanic White, both sexes</b>												
2003.....	8,035	-	-	-	-	3	9	12	40	79	155	242
2002.....	8,124	-	1	-	-	4	11	21	34	75	130	246
2001.....	8,308	-	-	-	-	1	5	9	34	72	139	244
2000.....	8,600	-	-	-	-	2	3	10	43	65	148	242
1999.....	8,712	-	-	-	-	1	3	15	35	95	142	267
<b>Non-Hispanic White, male</b>												
2003.....	4,718	-	-	-	-	2	3	7	21	48	92	155
2002.....	4,864	-	1	-	-	-	3	13	18	44	79	167
2001.....	4,884	-	-	-	-	1	1	7	22	44	78	164
2000.....	5,033	-	-	-	-	-	2	6	23	40	89	158
1999.....	5,171	-	-	-	-	1	2	7	23	57	85	184
<b>Non-Hispanic White, female</b>												
2003.....	3,317	-	-	-	-	1	6	5	19	31	63	87
2002.....	3,260	-	-	-	-	4	8	8	16	31	51	79
2001.....	3,424	-	-	-	-	-	4	2	12	28	61	80
2000.....	3,567	-	-	-	-	2	1	4	20	25	59	84
1999.....	3,541	-	-	-	-	-	1	8	12	38	57	83





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms of colon, rectum and anus (C18-C21)</b>												
<b>All origins, both sexes</b>												
2003.....	55,958	-	-	1	3	8	32	96	195	445	870	1,724
2002.....	56,741	1	-	-	1	9	41	88	202	439	986	1,755
2001.....	56,887	-	-	-	-	11	35	81	218	505	928	1,744
2000.....	57,477	-	-	-	3	13	27	79	193	480	889	1,656
1999.....	57,155	1	-	-	1	11	38	87	196	439	894	1,623
<b>All origins, male</b>												
2003.....	28,007	-	-	-	3	6	19	59	99	235	479	965
2002.....	28,501	1	-	-	1	6	28	48	113	233	564	958
2001.....	28,225	-	-	-	-	6	18	38	119	290	491	967
2000.....	28,462	-	-	-	2	8	16	42	111	271	488	893
1999.....	28,240	-	-	-	1	5	22	57	103	226	452	876
<b>All origins, female</b>												
2003.....	27,951	-	-	1	-	2	13	37	96	210	391	759
2002.....	28,240	-	-	-	-	3	13	40	89	206	422	797
2001.....	28,662	-	-	-	-	5	17	43	99	215	437	777
2000.....	29,015	-	-	-	1	5	11	37	82	209	401	763
1999.....	28,915	1	-	-	-	6	16	30	93	213	442	747
<b>Hispanic, both sexes</b>												
2003.....	2,471	-	-	-	-	1	9	16	18	54	68	132
2002.....	2,364	-	-	-	1	1	1	12	15	44	77	104
2001.....	2,250	-	-	-	-	-	6	5	15	44	83	88
2000.....	2,117	-	-	-	-	3	5	8	25	47	71	111
1999.....	2,031	-	-	-	-	1	6	7	24	33	53	113

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms of colon, rectum and anus (C18-C21)</b>												
<b>Hispanic, male</b>												
2003.....	1,363	-	-	-	-	1	7	12	8	28	44	80
2002.....	1,287	-	-	-	1	-	-	7	6	23	39	66
2001.....	1,185	-	-	-	-	-	3	4	4	28	48	44
2000.....	1,124	-	-	-	-	1	2	6	11	20	32	53
1999.....	1,062	-	-	-	-	1	4	3	11	15	24	59
<b>Hispanic, female</b>												
2003.....	1,108	-	-	-	-	-	2	4	10	26	24	52
2002.....	1,077	-	-	-	-	1	1	5	9	21	38	38
2001.....	1,065	-	-	-	-	-	3	1	11	16	35	44
2000.....	993	-	-	-	-	2	3	2	14	27	39	58
1999.....	969	-	-	-	-	-	2	4	13	18	29	54
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	53,372	-	-	1	3	7	23	80	177	391	802	1,585
2002.....	54,244	1	-	-	-	8	40	75	184	394	905	1,647
2001.....	54,513	-	-	-	-	11	29	76	203	458	843	1,650
2000.....	55,244	-	-	-	3	10	22	71	167	432	815	1,538
1999.....	54,976	1	-	-	1	10	31	80	172	404	837	1,506
<b>Non-Hispanic 1/, male</b>												
2003.....	26,579	-	-	-	3	5	12	47	91	207	435	882
2002.....	27,140	1	-	-	-	6	28	41	105	209	521	888
2001.....	26,978	-	-	-	-	6	15	34	115	260	441	920
2000.....	27,287	-	-	-	2	7	14	36	99	250	455	837
1999.....	27,103	-	-	-	1	4	18	54	92	211	425	816

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Non-Hispanic 1/, female												
2003.....	26,793	-	-	1	-	2	11	33	86	184	367	703
2002.....	27,104	-	-	-	-	2	12	34	79	185	384	759
2001.....	27,535	-	-	-	-	5	14	42	88	198	402	730
2000.....	27,957	-	-	-	1	3	8	35	68	182	360	701
1999.....	27,873	1	-	-	-	6	13	26	80	193	412	690
Non-Hispanic White, both sexes												
2003.....	45,205	-	-	1	3	6	13	56	128	275	570	1,177
2002.....	46,147	1	-	-	-	6	30	57	149	292	655	1,211
2001.....	46,450	-	-	-	-	7	21	55	148	339	627	1,235
2000.....	47,317	-	-	-	3	9	14	49	118	303	595	1,137
1999.....	47,275	-	-	-	1	6	20	56	126	303	630	1,152
Non-Hispanic White, male												
2003.....	22,624	-	-	-	3	4	8	33	66	149	318	668
2002.....	23,119	1	-	-	-	4	20	32	82	151	383	670
2001.....	23,128	-	-	-	-	5	8	24	87	196	327	701
2000.....	23,369	-	-	-	2	6	7	26	75	184	331	637
1999.....	23,446	-	-	-	1	2	10	38	67	166	334	634
Non-Hispanic White, female												
2003.....	22,581	-	-	1	-	2	5	23	62	126	252	509
2002.....	23,028	-	-	-	-	2	10	25	67	141	272	541
2001.....	23,322	-	-	-	-	2	13	31	61	143	300	534
2000.....	23,948	-	-	-	1	3	7	23	43	119	264	500
1999.....	23,829	-	-	-	-	4	10	18	59	137	296	518

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Non-Hispanic Black, both sexes												
2003.....	6,881	-	-	-	-	1	7	17	33	96	191	345
2002.....	6,817	-	-	-	-	2	8	11	27	75	208	367
2001.....	6,847	-	-	-	-	2	6	13	43	97	183	353
2000.....	6,820	-	-	-	-	-	6	12	42	109	181	357
1999.....	6,709	1	-	-	-	4	10	20	35	90	178	301
Non-Hispanic Black, male												
2003.....	3,307	-	-	-	-	1	3	9	16	44	97	185
2002.....	3,322	-	-	-	-	2	6	5	18	43	113	184
2001.....	3,221	-	-	-	-	1	6	6	24	53	99	186
2000.....	3,307	-	-	-	-	-	5	4	22	56	104	180
1999.....	3,160	-	-	-	-	2	7	13	17	42	81	158
Non-Hispanic Black, female												
2003.....	3,574	-	-	-	-	-	4	8	17	52	94	160
2002.....	3,495	-	-	-	-	-	2	6	9	32	95	183
2001.....	3,626	-	-	-	-	1	-	7	19	44	84	167
2000.....	3,513	-	-	-	-	-	1	8	20	53	77	177
1999.....	3,549	1	-	-	-	2	3	7	18	48	97	143
Origin not stated 2/, both sexes												
2003.....	115	-	-	-	-	-	-	-	-	-	-	7
2002.....	133	-	-	-	-	-	-	1	3	1	4	4
2001.....	124	-	-	-	-	-	-	-	-	3	2	6
2000.....	116	-	-	-	-	-	-	-	1	1	3	7
1999.....	148	-	-	-	-	-	1	-	-	2	4	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Origin not stated 2/, male

2003.....	65	-	-	-	-	-	-	-	-	-	-	3
2002.....	74	-	-	-	-	-	-	-	2	1	4	4
2001.....	62	-	-	-	-	-	-	-	-	2	2	3
2000.....	51	-	-	-	-	-	-	-	1	1	1	3
1999.....	75	-	-	-	-	-	-	-	-	-	3	1

Origin not stated 2/, female

2003.....	50	-	-	-	-	-	-	-	-	-	-	4
2002.....	59	-	-	-	-	-	-	1	1	-	-	-
2001.....	62	-	-	-	-	-	-	-	-	1	-	3
2000.....	65	-	-	-	-	-	-	-	-	-	2	4
1999.....	73	-	-	-	-	-	1	-	-	2	1	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All origins, both sexes												
2003.....	14,706	6	23	6	8	10	26	25	64	113	282	815
2002.....	14,047	3	28	7	11	10	17	33	72	123	291	731
2001.....	13,351	5	18	7	8	14	15	29	46	137	264	795
2000.....	12,916	2	23	9	11	15	15	28	58	113	316	742
1999.....	12,382	8	29	5	6	19	24	33	45	127	317	666
All origins, male												
2003.....	9,482	2	18	5	4	6	14	12	41	83	205	643
2002.....	9,011	2	19	4	8	9	9	20	42	85	223	596
2001.....	8,483	3	7	4	3	6	9	16	27	100	202	618
2000.....	8,176	1	14	6	2	11	9	18	41	74	223	595
1999.....	7,689	5	20	5	4	12	12	22	26	88	230	533
All origins, female												
2003.....	5,224	4	5	1	4	4	12	13	23	30	77	172
2002.....	5,036	1	9	3	3	1	8	13	30	38	68	135
2001.....	4,868	2	11	3	5	8	6	13	19	37	62	177
2000.....	4,740	1	9	3	9	4	6	10	17	39	93	147
1999.....	4,693	3	9	-	2	7	12	11	19	39	87	133
Hispanic, both sexes												
2003.....	1,511	-	10	1	-	1	-	3	8	16	32	111
2002.....	1,363	-	3	-	-	-	1	1	7	11	37	95
2001.....	1,256	1	3	1	2	1	1	5	9	19	27	107
2000.....	1,176	1	3	3	4	2	2	2	5	14	45	92
1999.....	1,084	1	4	1	1	2	2	3	4	13	30	76

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms of liver and intrahepatic bile ducts (C22)</b>												
<b>Hispanic, male</b>												
2003.....	986	-	8	1	-	1	-	1	5	14	23	90
2002.....	891	-	2	-	-	-	1	-	4	7	26	74
2001.....	812	1	1	-	-	1	-	2	5	12	24	96
2000.....	762	1	2	3	1	2	1	2	3	10	35	72
1999.....	688	1	3	1	1	1	2	3	2	6	21	58
<b>Hispanic, female</b>												
2003.....	525	-	2	-	-	-	-	2	3	2	9	21
2002.....	472	-	1	-	-	-	-	1	3	4	11	21
2001.....	444	-	2	1	2	-	1	3	4	7	3	11
2000.....	414	-	1	-	3	-	1	-	2	4	10	20
1999.....	396	-	1	-	-	1	-	-	2	7	9	18
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	13,157	6	13	5	8	9	26	22	55	97	249	700
2002.....	12,638	3	25	7	11	10	16	31	65	112	253	632
2001.....	12,053	4	15	6	6	13	14	24	37	116	236	686
2000.....	11,702	1	20	6	7	13	13	25	53	98	270	646
1999.....	11,265	7	23	4	5	17	22	30	41	114	284	588
<b>Non-Hispanic 1/, male</b>												
2003.....	8,469	2	10	4	4	5	14	11	35	69	181	549
2002.....	8,085	2	17	4	8	9	8	19	38	78	196	518
2001.....	7,640	2	6	4	3	5	9	14	22	86	177	520
2000.....	7,384	-	12	3	1	9	8	15	38	63	188	519
1999.....	6,976	4	15	4	3	11	10	19	24	82	208	474

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Non-Hispanic 1/, female												
2003.....	4,688	4	3	1	4	4	12	11	20	28	68	151
2002.....	4,553	1	8	3	3	1	8	12	27	34	57	114
2001.....	4,413	2	9	2	3	8	5	10	15	30	59	166
2000.....	4,318	1	8	3	6	4	5	10	15	35	82	127
1999.....	4,289	3	8	-	2	6	12	11	17	32	76	114
Non-Hispanic White, both sexes												
2003.....	10,149	4	9	3	4	6	16	9	25	58	158	447
2002.....	9,878	2	19	7	6	6	12	17	32	63	164	427
2001.....	9,433	4	11	1	4	9	11	14	20	62	127	445
2000.....	9,121	-	14	6	2	8	8	10	27	51	150	420
1999.....	8,890	5	20	2	2	12	13	14	15	60	178	371
Non-Hispanic White, male												
2003.....	6,504	2	7	3	3	3	9	2	15	37	109	354
2002.....	6,215	2	12	4	3	5	4	8	15	41	127	353
2001.....	5,956	2	4	1	3	4	7	7	10	43	94	334
2000.....	5,679	-	9	3	1	5	5	6	16	27	107	331
1999.....	5,455	2	13	2	1	9	5	8	8	39	134	293
Non-Hispanic White, female												
2003.....	3,645	2	2	-	1	3	7	7	10	21	49	93
2002.....	3,663	-	7	3	3	1	8	9	17	22	37	74
2001.....	3,477	2	7	-	1	5	4	7	10	19	33	111
2000.....	3,442	-	5	3	1	3	3	4	11	24	43	89
1999.....	3,435	3	7	-	1	3	8	6	7	21	44	78





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of pancreas (C25)</b>												
<b>All origins, both sexes</b>												
2003.....	30,777	-	-	-	1	-	10	20	40	153	395	891
2002.....	30,264	1	-	-	-	2	5	7	32	156	440	889
2001.....	29,803	-	1	1	1	1	1	11	44	167	407	833
2000.....	29,332	2	1	-	1	1	4	17	38	140	408	834
1999.....	29,082	1	-	-	-	-	3	9	46	178	416	887
<b>All origins, male</b>												
2003.....	15,060	-	-	-	1	-	6	11	24	97	232	556
2002.....	14,877	-	-	-	-	2	3	3	23	89	275	555
2001.....	14,467	-	1	-	-	1	1	5	25	99	222	503
2000.....	14,238	1	1	-	-	1	2	11	22	87	237	531
1999.....	14,176	-	-	-	-	-	1	7	26	107	262	575
<b>All origins, female</b>												
2003.....	15,717	-	-	-	-	-	4	9	16	56	163	335
2002.....	15,387	1	-	-	-	-	2	4	9	67	165	334
2001.....	15,336	-	-	1	1	-	-	6	19	68	185	330
2000.....	15,094	1	-	-	1	-	2	6	16	53	171	303
1999.....	14,906	1	-	-	-	-	2	2	20	71	154	312
<b>Hispanic, both sexes</b>												
2003.....	1,485	-	-	-	-	-	-	4	6	16	33	64
2002.....	1,363	-	-	-	-	-	1	-	4	17	34	59
2001.....	1,336	-	-	-	-	1	-	2	4	16	30	47
2000.....	1,243	1	-	-	-	-	2	2	4	13	24	53
1999.....	1,135	-	-	-	-	-	-	1	5	18	26	63

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of pancreas (C25)</b>												
<b>Hispanic, male</b>												
2003.....	744	-	-	-	-	-	-	2	3	9	21	39
2002.....	667	-	-	-	-	-	1	-	2	14	21	39
2001.....	643	-	-	-	-	1	-	1	2	9	17	28
2000.....	620	1	-	-	-	-	1	1	1	8	13	37
1999.....	563	-	-	-	-	-	-	-	2	12	18	43
<b>Hispanic, female</b>												
2003.....	741	-	-	-	-	-	-	2	3	7	12	25
2002.....	696	-	-	-	-	-	-	-	2	3	13	20
2001.....	693	-	-	-	-	-	-	1	2	7	13	19
2000.....	623	-	-	-	-	-	1	1	3	5	11	16
1999.....	572	-	-	-	-	-	-	1	3	6	8	20
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	29,233	-	-	-	1	-	10	16	34	137	361	826
2002.....	28,839	1	-	-	-	2	4	7	28	139	406	827
2001.....	28,407	-	1	1	1	-	1	9	39	151	375	786
2000.....	28,026	1	1	-	1	1	2	15	34	127	383	778
1999.....	27,869	1	-	-	-	-	3	8	41	160	387	820
<b>Non-Hispanic 1/, male</b>												
2003.....	14,285	-	-	-	1	-	6	9	21	88	211	517
2002.....	14,181	-	-	-	-	2	2	3	21	75	254	516
2001.....	13,792	-	1	-	-	-	1	4	23	90	204	475
2000.....	13,590	-	1	-	-	1	1	10	21	79	223	492
1999.....	13,572	-	-	-	-	-	1	7	24	95	243	528

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of pancreas (C25)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	14,948	-	-	-	-	-	4	7	13	49	150	309
2002.....	14,658	1	-	-	-	-	2	4	7	64	152	311
2001.....	14,615	-	-	1	1	-	-	5	16	61	171	311
2000.....	14,436	1	-	-	1	-	1	5	13	48	160	286
1999.....	14,297	1	-	-	-	-	2	1	17	65	144	292
<b>Non-Hispanic White, both sexes</b>												
2003.....	24,919	-	-	-	-	-	7	15	24	105	267	636
2002.....	24,647	-	-	-	-	2	4	4	21	103	321	656
2001.....	24,424	-	1	1	1	-	1	5	31	119	296	606
2000.....	24,034	1	1	-	1	1	1	13	26	97	294	625
1999.....	23,903	-	-	-	-	-	2	6	28	119	302	663
<b>Non-Hispanic White, male</b>												
2003.....	12,277	-	-	-	-	-	3	8	13	69	153	397
2002.....	12,242	-	-	-	-	2	2	1	16	52	205	413
2001.....	11,994	-	1	-	-	-	1	3	19	69	164	376
2000.....	11,764	-	1	-	-	1	1	8	17	61	175	399
1999.....	11,684	-	-	-	-	-	1	6	17	73	193	430
<b>Non-Hispanic White, female</b>												
2003.....	12,642	-	-	-	-	-	4	7	11	36	114	239
2002.....	12,405	-	-	-	-	-	2	3	5	51	116	243
2001.....	12,430	-	-	1	1	-	-	2	12	50	132	230
2000.....	12,270	1	-	-	1	-	-	5	9	36	119	226
1999.....	12,219	-	-	-	-	-	1	-	11	46	109	233







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of larynx (C32)</b>												
<b>Hispanic, male</b>												
2003.....	148	-	-	-	-	-	-	-	-	-	-	8
2002.....	139	-	-	-	-	-	-	-	-	3	4	6
2001.....	141	-	-	-	-	-	-	-	-	-	3	13
2000.....	124	-	-	-	-	-	-	-	-	1	2	5
1999.....	149	-	-	-	-	-	-	-	-	-	4	15
<b>Hispanic, female</b>												
2003.....	14	-	-	-	-	-	-	-	-	-	-	1
2002.....	26	-	-	-	-	-	-	-	1	-	1	1
2001.....	22	-	-	-	-	-	-	-	-	-	-	1
2000.....	19	-	-	-	1	-	-	-	1	-	1	-
1999.....	22	-	-	-	-	-	-	-	-	-	1	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	3,610	-	-	-	-	-	-	-	1	10	68	147
2002.....	3,548	-	-	-	-	-	-	3	7	16	52	149
2001.....	3,614	-	-	-	-	-	-	1	2	18	64	135
2000.....	3,699	-	-	-	1	1	-	-	3	8	51	150
1999.....	3,627	-	-	-	-	-	-	-	6	20	53	135
<b>Non-Hispanic 1/, male</b>												
2003.....	2,853	-	-	-	-	-	-	-	1	7	52	120
2002.....	2,775	-	-	-	-	-	-	3	3	13	38	124
2001.....	2,858	-	-	-	-	-	-	1	1	12	50	110
2000.....	2,901	-	-	-	-	-	-	-	2	5	40	116
1999.....	2,832	-	-	-	-	-	-	-	5	18	42	114

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of larynx (C32)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	757	-	-	-	-	-	-	-	-	3	16	27
2002.....	773	-	-	-	-	-	-	-	4	3	14	25
2001.....	756	-	-	-	-	-	-	-	1	6	14	25
2000.....	798	-	-	-	1	1	-	-	1	3	11	34
1999.....	795	-	-	-	-	-	-	-	1	2	11	21
<b>Non-Hispanic White, both sexes</b>												
2003.....	2,879	-	-	-	-	-	-	-	1	9	52	100
2002.....	2,838	-	-	-	-	-	-	2	4	7	33	97
2001.....	2,876	-	-	-	-	-	-	1	2	11	47	84
2000.....	2,940	-	-	-	1	1	-	-	3	7	37	102
1999.....	2,924	-	-	-	-	-	-	-	4	14	26	85
<b>Non-Hispanic White, male</b>												
2003.....	2,251	-	-	-	-	-	-	-	1	6	42	85
2002.....	2,205	-	-	-	-	-	-	2	1	5	25	82
2001.....	2,267	-	-	-	-	-	-	1	1	10	37	66
2000.....	2,291	-	-	-	-	-	-	-	2	4	27	80
1999.....	2,263	-	-	-	-	-	-	-	3	14	20	71
<b>Non-Hispanic White, female</b>												
2003.....	628	-	-	-	-	-	-	-	-	3	10	15
2002.....	633	-	-	-	-	-	-	-	3	2	8	15
2001.....	609	-	-	-	-	-	-	-	1	1	10	18
2000.....	649	-	-	-	1	1	-	-	1	3	10	22
1999.....	661	-	-	-	-	-	-	-	1	-	6	14





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All origins, both sexes												
2003.....	158,086	-	2	-	2	7	18	29	125	520	1,958	4,617
2002.....	157,713	-	1	2	2	4	17	35	130	583	2,115	4,341
2001.....	156,058	-	5	1	1	6	14	36	121	640	2,156	4,255
2000.....	155,521	2	2	-	1	3	19	37	160	664	2,092	4,220
1999.....	152,156	-	-	3	2	6	14	43	136	668	2,056	3,916
All origins, male												
2003.....	89,964	-	1	-	1	4	10	9	69	258	1,086	2,659
2002.....	90,171	-	1	1	2	2	14	19	71	290	1,096	2,471
2001.....	90,426	-	4	1	-	2	10	17	60	315	1,189	2,487
2000.....	90,469	2	1	-	-	3	11	28	74	370	1,172	2,540
1999.....	89,453	-	-	2	2	5	8	24	73	323	1,151	2,346
All origins, female												
2003.....	68,122	-	1	-	1	3	8	20	56	262	872	1,958
2002.....	67,542	-	-	1	-	2	3	16	59	293	1,019	1,870
2001.....	65,632	-	1	-	1	4	4	19	61	325	967	1,768
2000.....	65,052	-	1	-	1	-	8	9	86	294	920	1,680
1999.....	62,703	-	-	1	-	1	6	19	63	345	905	1,570
Hispanic, both sexes												
2003.....	4,136	-	-	-	-	1	3	7	13	26	66	143
2002.....	4,038	-	-	2	-	-	4	13	14	23	67	124
2001.....	3,816	-	2	-	1	1	6	6	10	27	79	127
2000.....	3,643	-	-	-	-	1	2	4	13	28	62	124
1999.....	3,545	-	-	-	-	-	3	3	14	32	66	111

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Hispanic, male												
2003.....	2,640	-	-	-	-	1	2	1	6	14	38	77
2002.....	2,617	-	-	1	-	-	2	5	9	8	37	71
2001.....	2,495	-	2	-	-	-	6	3	7	22	46	83
2000.....	2,411	-	-	-	-	1	2	2	8	20	36	81
1999.....	2,373	-	-	-	-	-	2	3	7	17	41	74
Hispanic, female												
2003.....	1,496	-	-	-	-	-	1	6	7	12	28	66
2002.....	1,421	-	-	1	-	-	2	8	5	15	30	53
2001.....	1,321	-	-	-	1	1	-	3	3	5	33	44
2000.....	1,232	-	-	-	-	-	-	2	5	8	26	43
1999.....	1,172	-	-	-	-	-	1	-	7	15	25	37
Non-Hispanic 1/, both sexes												
2003.....	153,616	-	2	-	2	6	15	22	110	493	1,885	4,461
2002.....	153,292	-	1	-	2	4	13	22	113	559	2,042	4,204
2001.....	151,882	-	3	1	-	5	8	30	111	611	2,067	4,119
2000.....	151,483	2	2	-	1	2	17	33	147	635	2,018	4,081
1999.....	148,206	-	-	3	2	6	11	40	121	635	1,983	3,795
Non-Hispanic 1/, male												
2003.....	87,111	-	1	-	1	3	8	8	61	244	1,043	2,576
2002.....	87,335	-	1	-	2	2	12	14	62	281	1,056	2,392
2001.....	87,706	-	2	1	-	2	4	14	53	293	1,138	2,400
2000.....	87,816	2	1	-	-	2	9	26	66	350	1,129	2,453
1999.....	86,822	-	-	2	2	5	6	21	66	306	1,105	2,266

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasms of trachea, bronchus and lung (C33-C34)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	66,505	-	1	-	1	3	7	14	49	249	842	1,885
2002.....	65,957	-	-	-	-	2	1	8	51	278	986	1,812
2001.....	64,176	-	1	-	-	3	4	16	58	318	929	1,719
2000.....	63,667	-	1	-	1	-	8	7	81	285	889	1,628
1999.....	61,384	-	-	1	-	1	5	19	55	329	878	1,529
<b>Non-Hispanic White, both sexes</b>												
2003.....	134,437	-	2	-	1	6	12	17	90	382	1,470	3,427
2002.....	134,435	-	1	-	1	3	10	12	85	441	1,585	3,294
2001.....	133,266	-	2	1	-	4	5	23	91	494	1,606	3,201
2000.....	133,061	1	2	-	1	-	11	23	116	470	1,526	3,134
1999.....	129,967	-	-	2	1	3	6	30	101	479	1,546	2,910
<b>Non-Hispanic White, male</b>												
2003.....	75,468	-	1	-	-	3	6	7	48	189	797	1,965
2002.....	75,754	-	1	-	1	2	9	9	49	213	803	1,835
2001.....	76,066	-	1	1	-	2	2	12	43	232	863	1,827
2000.....	76,218	1	1	-	-	-	7	18	55	257	819	1,869
1999.....	75,338	-	-	1	1	3	3	14	57	231	847	1,708
<b>Non-Hispanic White, female</b>												
2003.....	58,969	-	1	-	1	3	6	10	42	193	673	1,462
2002.....	58,681	-	-	-	-	1	1	3	36	228	782	1,459
2001.....	57,200	-	1	-	-	2	3	11	48	262	743	1,374
2000.....	56,843	-	1	-	1	-	4	5	61	213	707	1,265
1999.....	54,629	-	-	1	-	-	3	16	44	248	699	1,202

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Non-Hispanic Black, both sexes												
2003.....	16,252	-	-	-	1	-	3	2	16	81	360	922
2002.....	16,170	-	-	-	1	1	3	4	19	96	407	822
2001.....	15,883	-	1	-	-	1	2	3	13	102	413	800
2000.....	15,924	1	-	-	-	2	3	8	23	142	436	833
1999.....	15,834	-	-	1	1	3	5	5	15	134	397	795
Non-Hispanic Black, male												
2003.....	9,914	-	-	-	1	-	2	1	11	39	212	549
2002.....	9,983	-	-	-	1	-	3	2	12	57	230	504
2001.....	10,030	-	1	-	-	-	1	1	6	53	245	511
2000.....	10,056	1	-	-	-	2	-	6	8	80	279	518
1999.....	10,039	-	-	1	1	2	3	4	7	63	238	512
Non-Hispanic Black, female												
2003.....	6,338	-	-	-	-	-	1	1	5	42	148	373
2002.....	6,187	-	-	-	-	1	-	2	7	39	177	318
2001.....	5,853	-	-	-	-	1	1	2	7	49	168	289
2000.....	5,868	-	-	-	-	-	3	2	15	62	157	315
1999.....	5,795	-	-	-	-	1	2	1	8	71	159	283
Origin not stated 2/, both sexes												
2003.....	334	-	-	-	-	-	-	-	2	1	7	13
2002.....	383	-	-	-	-	-	-	-	3	1	6	13
2001.....	360	-	-	-	-	-	-	-	-	2	10	9
2000.....	395	-	-	-	-	-	-	-	-	1	12	15
1999.....	405	-	-	-	-	-	-	-	1	1	7	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Origin not stated 2/, male

2003.....	213	-	-	-	-	-	-	-	2	-	5	6
2002.....	219	-	-	-	-	-	-	-	-	1	3	8
2001.....	225	-	-	-	-	-	-	-	-	-	5	4
2000.....	242	-	-	-	-	-	-	-	-	-	7	6
1999.....	258	-	-	-	-	-	-	-	-	-	5	6

Origin not stated 2/, female

2003.....	121	-	-	-	-	-	-	-	-	1	2	7
2002.....	164	-	-	-	-	-	-	-	3	-	3	5
2001.....	135	-	-	-	-	-	-	-	-	2	5	5
2000.....	153	-	-	-	-	-	-	-	-	1	5	9
1999.....	147	-	-	-	-	-	-	-	1	1	2	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant melanoma of skin (C43)</b>												
<b>All origins, both sexes</b>												
2003.....	7,818	-	1	1	1	9	29	63	128	189	372	541
2002.....	7,514	-	3	-	1	7	34	70	124	219	385	495
2001.....	7,542	-	-	-	-	11	30	72	114	213	409	536
2000.....	7,420	-	-	-	1	7	34	58	128	234	392	534
1999.....	7,215	1	-	1	-	9	30	62	132	239	386	518
<b>All origins, male</b>												
2003.....	4,963	-	-	-	-	6	18	40	79	114	222	328
2002.....	4,725	-	2	-	1	3	17	38	81	120	243	343
2001.....	4,763	-	-	-	-	9	15	40	63	130	250	331
2000.....	4,592	-	-	-	1	4	17	33	77	118	237	321
1999.....	4,529	1	-	-	-	7	16	38	72	151	246	337
<b>All origins, female</b>												
2003.....	2,855	-	1	1	1	3	11	23	49	75	150	213
2002.....	2,789	-	1	-	-	4	17	32	43	99	142	152
2001.....	2,779	-	-	-	-	2	15	32	51	83	159	205
2000.....	2,828	-	-	-	-	3	17	25	51	116	155	213
1999.....	2,686	-	-	1	-	2	14	24	60	88	140	181
<b>Hispanic, both sexes</b>												
2003.....	159	-	-	1	-	1	1	3	3	4	11	13
2002.....	147	-	-	-	-	-	2	2	5	7	14	13
2001.....	145	-	-	-	-	1	1	3	3	8	16	11
2000.....	144	-	-	-	-	-	2	2	3	14	12	15
1999.....	133	1	-	1	-	1	1	1	4	6	8	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant melanoma of skin (C43)</b>												
<b>Hispanic, male</b>												
2003.....	91	-	-	-	-	1	-	2	2	1	7	4
2002.....	77	-	-	-	-	-	1	1	5	4	9	8
2001.....	78	-	-	-	-	-	-	1	1	7	10	8
2000.....	92	-	-	-	-	-	1	1	3	9	8	11
1999.....	79	1	-	-	-	1	-	1	3	4	5	7
<b>Hispanic, female</b>												
2003.....	68	-	-	1	-	-	1	1	1	3	4	9
2002.....	70	-	-	-	-	-	1	1	-	3	5	5
2001.....	67	-	-	-	-	1	1	2	2	1	6	3
2000.....	52	-	-	-	-	-	1	1	-	5	4	4
1999.....	54	-	-	1	-	-	1	-	1	2	3	2
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	7,640	-	1	-	1	8	28	60	123	185	359	528
2002.....	7,349	-	3	-	1	7	31	68	118	211	370	480
2001.....	7,387	-	-	-	-	10	29	69	110	204	393	525
2000.....	7,263	-	-	-	1	7	32	56	125	219	380	519
1999.....	7,062	-	-	-	-	8	29	61	128	233	378	507
<b>Non-Hispanic 1/, male</b>												
2003.....	4,859	-	-	-	-	5	18	38	75	113	213	324
2002.....	4,640	-	2	-	1	3	16	37	76	116	234	334
2001.....	4,678	-	-	-	-	9	15	39	61	123	240	323
2000.....	4,490	-	-	-	1	4	16	32	74	108	229	310
1999.....	4,438	-	-	-	-	6	16	37	69	147	241	328

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant melanoma of skin (C43)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	2,781	-	1	-	1	3	10	22	48	72	146	204
2002.....	2,709	-	1	-	-	4	15	31	42	95	136	146
2001.....	2,709	-	-	-	-	1	14	30	49	81	153	202
2000.....	2,773	-	-	-	-	3	16	24	51	111	151	209
1999.....	2,624	-	-	-	-	2	13	24	59	86	137	179
<b>Non-Hispanic White, both sexes</b>												
2003.....	7,471	-	1	-	1	8	26	58	120	182	351	515
2002.....	7,190	-	3	-	1	6	31	66	118	202	366	471
2001.....	7,250	-	-	-	-	9	29	67	110	202	388	518
2000.....	7,090	-	-	-	1	6	32	53	122	214	365	510
1999.....	6,936	-	-	-	-	7	27	60	123	230	374	498
<b>Non-Hispanic White, male</b>												
2003.....	4,784	-	-	-	-	5	18	38	73	112	211	318
2002.....	4,555	-	2	-	1	2	16	35	76	114	230	329
2001.....	4,618	-	-	-	-	8	15	37	61	122	237	320
2000.....	4,414	-	-	-	1	4	16	31	73	106	221	308
1999.....	4,370	-	-	-	-	5	14	36	66	145	240	322
<b>Non-Hispanic White, female</b>												
2003.....	2,687	-	1	-	1	3	8	20	47	70	140	197
2002.....	2,635	-	1	-	-	4	15	31	42	88	136	142
2001.....	2,632	-	-	-	-	1	14	30	49	80	151	198
2000.....	2,676	-	-	-	-	2	16	22	49	108	144	202
1999.....	2,566	-	-	-	-	2	13	24	57	85	134	176





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of breast (C50)</b>												
<b>All origins, both sexes</b>												
2003.....	42,000	-	-	-	-	1	15	55	352	912	1,804	2,734
2002.....	41,883	-	-	-	-	1	9	63	362	904	1,818	2,747
2001.....	41,809	-	-	-	-	1	8	88	375	960	1,863	2,790
2000.....	42,300	-	-	-	-	1	10	83	384	982	1,859	2,730
1999.....	41,528	-	1	-	1	3	10	86	353	986	1,755	2,619
<b>All origins, male</b>												
2003.....	380	-	-	-	-	-	-	1	1	2	7	14
2002.....	369	-	-	-	-	-	-	2	1	4	8	16
2001.....	415	-	-	-	-	-	-	-	2	5	9	13
2000.....	428	-	-	-	-	-	-	2	1	12	7	18
1999.....	384	-	-	-	-	-	-	-	3	6	9	11
<b>All origins, female</b>												
2003.....	41,620	-	-	-	-	1	15	54	351	910	1,797	2,720
2002.....	41,514	-	-	-	-	1	9	61	361	900	1,810	2,731
2001.....	41,394	-	-	-	-	1	8	88	373	955	1,854	2,777
2000.....	41,872	-	-	-	-	1	10	81	383	970	1,852	2,712
1999.....	41,144	-	1	-	1	3	10	86	350	980	1,746	2,608
<b>Hispanic, both sexes</b>												
2003.....	1,893	-	-	-	-	-	2	11	44	86	170	182
2002.....	1,736	-	-	-	-	-	2	9	45	76	142	185
2001.....	1,760	-	-	-	-	-	1	15	38	104	140	184
2000.....	1,689	-	-	-	-	-	1	16	38	90	128	184
1999.....	1,551	-	-	-	-	1	1	13	34	74	123	160

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of breast (C50)</b>												
<b>Hispanic, male</b>												
2003.....	9	-	-	-	-	-	-	-	-	-	-	1
2002.....	12	-	-	-	-	-	-	1	-	-	-	1
2001.....	14	-	-	-	-	-	-	-	-	1	1	-
2000.....	19	-	-	-	-	-	-	-	-	1	1	-
1999.....	13	-	-	-	-	-	-	-	-	-	-	1
<b>Hispanic, female</b>												
2003.....	1,884	-	-	-	-	-	2	11	44	86	170	181
2002.....	1,724	-	-	-	-	-	2	8	45	76	142	184
2001.....	1,746	-	-	-	-	-	1	15	38	103	139	184
2000.....	1,670	-	-	-	-	-	1	16	38	89	127	184
1999.....	1,538	-	-	-	-	1	1	13	34	74	123	159
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	40,003	-	-	-	-	1	13	44	307	824	1,629	2,549
2002.....	40,037	-	-	-	-	1	7	54	314	825	1,670	2,554
2001.....	39,957	-	-	-	-	1	7	73	336	855	1,721	2,597
2000.....	40,506	-	-	-	-	1	9	67	346	891	1,724	2,541
1999.....	39,881	-	1	-	1	2	9	72	318	911	1,626	2,453
<b>Non-Hispanic 1/, male</b>												
2003.....	371	-	-	-	-	-	-	1	1	2	7	13
2002.....	355	-	-	-	-	-	-	1	1	4	8	14
2001.....	399	-	-	-	-	-	-	-	2	4	8	13
2000.....	409	-	-	-	-	-	-	2	1	11	6	18
1999.....	370	-	-	-	-	-	-	-	3	6	9	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of breast (C50)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	39,632	-	-	-	-	1	13	43	306	822	1,622	2,536
2002.....	39,682	-	-	-	-	1	7	53	313	821	1,662	2,540
2001.....	39,558	-	-	-	-	1	7	73	334	851	1,713	2,584
2000.....	40,097	-	-	-	-	1	9	65	345	880	1,718	2,523
1999.....	39,511	-	1	-	1	2	9	72	315	905	1,617	2,443
<b>Non-Hispanic White, both sexes</b>												
2003.....	33,451	-	-	-	-	1	8	17	199	567	1,161	1,875
2002.....	33,611	-	-	-	-	-	4	31	199	558	1,175	1,876
2001.....	33,629	-	-	-	-	-	3	43	212	596	1,232	1,947
2000.....	34,386	-	-	-	-	1	3	45	230	628	1,249	1,887
1999.....	33,790	-	-	-	1	2	4	46	211	627	1,175	1,820
<b>Non-Hispanic White, male</b>												
2003.....	317	-	-	-	-	-	-	-	1	1	3	8
2002.....	288	-	-	-	-	-	-	1	-	2	7	10
2001.....	329	-	-	-	-	-	-	-	1	3	3	10
2000.....	341	-	-	-	-	-	-	-	1	7	2	13
1999.....	299	-	-	-	-	-	-	-	2	-	8	8
<b>Non-Hispanic White, female</b>												
2003.....	33,134	-	-	-	-	1	8	17	198	566	1,158	1,867
2002.....	33,323	-	-	-	-	-	4	30	199	556	1,168	1,866
2001.....	33,300	-	-	-	-	-	3	43	211	593	1,229	1,937
2000.....	34,045	-	-	-	-	1	3	45	229	621	1,247	1,874
1999.....	33,491	-	-	-	1	2	4	46	209	627	1,167	1,812

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of breast (C50)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	5,677	-	-	-	-	-	5	23	94	226	406	588
2002.....	5,579	-	-	-	-	1	2	22	101	232	438	578
2001.....	5,528	-	-	-	-	1	4	30	110	219	422	552
2000.....	5,384	-	-	-	-	-	5	20	100	227	414	556
1999.....	5,386	-	1	-	-	-	5	24	100	259	402	545
<b>Non-Hispanic Black, male</b>												
2003.....	51	-	-	-	-	-	-	1	-	1	4	4
2002.....	62	-	-	-	-	-	-	-	1	2	1	4
2001.....	64	-	-	-	-	-	-	-	1	-	5	3
2000.....	63	-	-	-	-	-	-	2	-	4	4	4
1999.....	67	-	-	-	-	-	-	-	1	6	1	1
<b>Non-Hispanic Black, female</b>												
2003.....	5,626	-	-	-	-	-	5	22	94	225	402	584
2002.....	5,517	-	-	-	-	1	2	22	100	230	437	574
2001.....	5,464	-	-	-	-	1	4	30	109	219	417	549
2000.....	5,321	-	-	-	-	-	5	18	100	223	410	552
1999.....	5,319	-	1	-	-	-	5	24	99	253	401	544
<b>Origin not stated 2/, both sexes</b>												
2003.....	104	-	-	-	-	-	-	-	1	2	5	3
2002.....	110	-	-	-	-	-	-	-	3	3	6	8
2001.....	92	-	-	-	-	-	-	-	1	1	2	9
2000.....	105	-	-	-	-	-	-	-	-	1	7	5
1999.....	96	-	-	-	-	-	-	1	1	1	6	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	1
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	104	-	-	-	-	-	-	-	1	2	5	3
2002.....	108	-	-	-	-	-	-	-	3	3	6	7
2001.....	90	-	-	-	-	-	-	-	1	1	2	9
2000.....	105	-	-	-	-	-	-	-	-	1	7	5
1999.....	95	-	-	-	-	-	-	1	1	1	6	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of cervix uteri (C53)												
All origins, both sexes												
2003.....	3,919	-	-	-	-	1	15	59	142	250	383	440
2002.....	3,952	-	-	-	-	1	16	47	161	275	391	439
2001.....	4,092	-	-	-	1	-	12	51	158	291	420	475
2000.....	4,200	-	-	-	1	3	14	77	153	294	400	448
1999.....	4,205	-	-	-	-	2	10	62	155	292	407	462
All origins, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female												
2003.....	3,919	-	-	-	-	1	15	59	142	250	383	440
2002.....	3,952	-	-	-	-	1	16	47	161	275	391	439
2001.....	4,092	-	-	-	1	-	12	51	158	291	420	475
2000.....	4,200	-	-	-	1	3	14	77	153	294	400	448
1999.....	4,205	-	-	-	-	2	10	62	155	292	407	462
Hispanic, both sexes												
2003.....	391	-	-	-	-	1	2	13	18	38	47	54
2002.....	388	-	-	-	-	-	2	8	22	42	44	42
2001.....	412	-	-	-	-	-	3	10	26	42	60	60
2000.....	434	-	-	-	-	-	4	13	20	54	54	47
1999.....	378	-	-	-	-	-	2	7	24	34	53	55



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of cervix uteri (C53)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	3,519	-	-	-	-	-	13	46	124	212	334	383
2002.....	3,551	-	-	-	-	1	14	39	138	231	346	396
2001.....	3,669	-	-	-	1	-	9	40	132	247	359	412
2000.....	3,755	-	-	-	1	3	10	64	133	239	344	399
1999.....	3,817	-	-	-	-	2	8	55	131	258	353	407
<b>Non-Hispanic White, both sexes</b>												
2003.....	2,564	-	-	-	-	-	10	27	83	144	248	279
2002.....	2,580	-	-	-	-	-	10	26	103	168	249	281
2001.....	2,725	-	-	-	-	-	8	29	96	182	266	311
2000.....	2,725	-	-	-	-	3	6	50	91	170	233	288
1999.....	2,854	-	-	-	-	-	7	47	100	188	263	316
<b>Non-Hispanic White, male</b>												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Non-Hispanic White, female</b>												
2003.....	2,564	-	-	-	-	-	10	27	83	144	248	279
2002.....	2,580	-	-	-	-	-	10	26	103	168	249	281
2001.....	2,725	-	-	-	-	-	8	29	96	182	266	311
2000.....	2,725	-	-	-	-	3	6	50	91	170	233	288
1999.....	2,854	-	-	-	-	-	7	47	100	188	263	316





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
All origins, both sexes												
2003.....	6,899	-	-	-	-	1	3	8	19	37	117	171
2002.....	6,853	-	-	-	-	-	-	2	21	46	100	189
2001.....	6,784	-	-	-	-	-	-	10	28	47	92	192
2000.....	6,586	-	-	-	-	-	3	2	25	41	97	190
1999.....	6,468	-	-	-	-	-	2	11	12	38	104	182
All origins, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female												
2003.....	6,899	-	-	-	-	1	3	8	19	37	117	171
2002.....	6,853	-	-	-	-	-	-	2	21	46	100	189
2001.....	6,784	-	-	-	-	-	-	10	28	47	92	192
2000.....	6,586	-	-	-	-	-	3	2	25	41	97	190
1999.....	6,468	-	-	-	-	-	2	11	12	38	104	182
Hispanic, both sexes												
2003.....	321	-	-	-	-	-	1	1	2	5	19	16
2002.....	337	-	-	-	-	-	-	-	5	7	13	18
2001.....	319	-	-	-	-	-	-	-	7	3	10	16
2000.....	277	-	-	-	-	-	-	-	2	4	15	11
1999.....	272	-	-	-	-	-	1	2	1	4	11	10



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Non-Hispanic 1/, female												
2003.....	6,561	-	-	-	-	1	2	7	17	32	98	154
2002.....	6,487	-	-	-	-	-	-	2	15	39	87	171
2001.....	6,449	-	-	-	-	-	-	10	21	44	82	176
2000.....	6,294	-	-	-	-	-	3	2	23	37	82	178
1999.....	6,188	-	-	-	-	-	1	9	11	34	93	172
Non-Hispanic White, both sexes												
2003.....	5,321	-	-	-	-	1	1	4	9	24	62	119
2002.....	5,213	-	-	-	-	-	-	1	10	27	70	130
2001.....	5,283	-	-	-	-	-	-	3	16	30	64	134
2000.....	5,113	-	-	-	-	-	1	2	16	29	59	147
1999.....	5,114	-	-	-	-	-	1	6	9	28	68	140
Non-Hispanic White, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic White, female												
2003.....	5,321	-	-	-	-	1	1	4	9	24	62	119
2002.....	5,213	-	-	-	-	-	-	1	10	27	70	130
2001.....	5,283	-	-	-	-	-	-	3	16	30	64	134
2000.....	5,113	-	-	-	-	-	1	2	16	29	59	147
1999.....	5,114	-	-	-	-	-	1	6	9	28	68	140





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of ovary (C56)</b>												
<b>All origins, both sexes</b>												
2003.....	14,657	-	-	-	-	10	18	31	53	134	318	632
2002.....	14,682	1	1	1	1	12	25	33	64	150	319	688
2001.....	14,414	-	-	-	2	5	16	37	65	152	320	610
2000.....	14,060	-	-	2	-	7	16	36	65	145	327	617
1999.....	13,627	-	1	-	2	9	18	26	51	162	334	616
<b>All origins, male</b>												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
<b>All origins, female</b>												
2003.....	14,657	-	-	-	-	10	18	31	53	134	318	632
2002.....	14,682	1	1	1	1	12	25	33	64	150	319	688
2001.....	14,414	-	-	-	2	5	16	37	65	152	320	610
2000.....	14,060	-	-	2	-	7	16	36	65	145	327	617
1999.....	13,627	-	1	-	2	9	18	26	51	162	334	616
<b>Hispanic, both sexes</b>												
2003.....	645	-	-	-	-	1	2	3	2	17	31	51
2002.....	622	-	-	1	1	2	1	1	9	10	37	45
2001.....	603	-	-	-	1	-	-	9	4	13	30	39
2000.....	578	-	-	1	-	-	4	3	6	11	19	42
1999.....	478	-	1	-	-	1	1	5	1	11	18	29



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of ovary (C56)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	13,985	-	-	-	-	9	16	28	51	117	286	577
2002.....	14,031	1	1	-	-	10	24	32	55	140	281	640
2001.....	13,783	-	-	-	1	5	16	28	61	139	289	569
2000.....	13,454	-	-	1	-	7	12	33	59	133	308	571
1999.....	13,103	-	-	-	2	8	17	21	50	150	316	585
<b>Non-Hispanic White, both sexes</b>												
2003.....	12,477	-	-	-	-	7	11	20	44	87	242	483
2002.....	12,560	1	1	-	-	7	20	28	39	112	231	535
2001.....	12,338	-	-	-	-	3	11	23	45	107	240	496
2000.....	12,138	-	-	-	-	5	8	29	46	109	258	498
1999.....	11,795	-	-	-	2	7	13	15	36	124	270	506
<b>Non-Hispanic White, male</b>												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Non-Hispanic White, female</b>												
2003.....	12,477	-	-	-	-	7	11	20	44	87	242	483
2002.....	12,560	1	1	-	-	7	20	28	39	112	231	535
2001.....	12,338	-	-	-	-	3	11	23	45	107	240	496
2000.....	12,138	-	-	-	-	5	8	29	46	109	258	498
1999.....	11,795	-	-	-	2	7	13	15	36	124	270	506

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of ovary (C56)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	1,180	-	-	-	-	2	4	7	5	22	34	72
2002.....	1,155	-	-	-	-	3	4	3	10	24	35	72
2001.....	1,140	-	-	-	1	1	3	5	10	25	40	47
2000.....	1,065	-	-	1	-	-	2	4	8	16	43	50
1999.....	1,065	-	-	-	-	1	3	6	7	16	31	54
<b>Non-Hispanic Black, male</b>												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Non-Hispanic Black, female</b>												
2003.....	1,180	-	-	-	-	2	4	7	5	22	34	72
2002.....	1,155	-	-	-	-	3	4	3	10	24	35	72
2001.....	1,140	-	-	-	1	1	3	5	10	25	40	47
2000.....	1,065	-	-	1	-	-	2	4	8	16	43	50
1999.....	1,065	-	-	-	-	1	3	6	7	16	31	54
<b>Origin not stated 2/, both sexes</b>												
2003.....	27	-	-	-	-	-	-	-	-	-	1	4
2002.....	29	-	-	-	-	-	-	-	-	-	1	3
2001.....	28	-	-	-	-	-	-	-	-	-	1	2
2000.....	28	-	-	-	-	-	-	-	-	1	-	4
1999.....	46	-	-	-	-	-	-	-	-	1	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasm of ovary (C56)												
Origin not stated 2/, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
Origin not stated 2/, female												
2003.....	27	-	-	-	-	-	-	-	-	-	1	4
2002.....	29	-	-	-	-	-	-	-	-	-	1	3
2001.....	28	-	-	-	-	-	-	-	-	-	1	2
2000.....	28	-	-	-	-	-	-	-	-	1	-	4
1999.....	46	-	-	-	-	-	-	-	-	1	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of prostate (C61)</b>												
All origins, both sexes												
2003.....	29,554	-	-	1	-	1	-	1	-	3	22	99
2002.....	30,446	-	-	-	-	-	1	-	1	7	25	80
2001.....	30,719	-	-	-	-	-	1	1	-	3	27	99
2000.....	31,078	-	-	-	-	-	-	1	-	6	19	108
1999.....	31,729	1	-	-	-	-	1	1	1	5	28	100
All origins, male												
2003.....	29,554	-	-	1	-	1	-	1	-	3	22	99
2002.....	30,446	-	-	-	-	-	1	-	1	7	25	80
2001.....	30,719	-	-	-	-	-	1	1	-	3	27	99
2000.....	31,078	-	-	-	-	-	-	1	-	6	19	108
1999.....	31,729	1	-	-	-	-	1	1	1	5	28	100
All origins, female												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
Hispanic, both sexes												
2003.....	1,219	-	-	-	-	-	-	-	-	-	1	7
2002.....	1,209	-	-	-	-	-	-	-	1	-	1	7
2001.....	1,226	-	-	-	-	-	-	-	-	-	-	5
2000.....	1,033	-	-	-	-	-	-	-	-	-	1	7
1999.....	1,050	-	-	-	-	-	1	1	-	-	2	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of prostate (C61)</b>												
<b>Hispanic, male</b>												
2003.....	1,219	-	-	-	-	-	-	-	-	-	1	7
2002.....	1,209	-	-	-	-	-	-	-	1	-	1	7
2001.....	1,226	-	-	-	-	-	-	-	-	-	-	5
2000.....	1,033	-	-	-	-	-	-	-	-	-	1	7
1999.....	1,050	-	-	-	-	-	1	1	-	-	2	2
<b>Hispanic, female</b>												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	28,279	-	-	1	-	1	-	1	-	3	21	92
2002.....	29,151	-	-	-	-	-	1	-	-	7	24	72
2001.....	29,407	-	-	-	-	-	1	1	-	3	26	94
2000.....	29,964	-	-	-	-	-	-	1	-	6	18	101
1999.....	30,614	1	-	-	-	-	-	-	1	5	26	97
<b>Non-Hispanic 1/, male</b>												
2003.....	28,279	-	-	1	-	1	-	1	-	3	21	92
2002.....	29,151	-	-	-	-	-	1	-	-	7	24	72
2001.....	29,407	-	-	-	-	-	1	1	-	3	26	94
2000.....	29,964	-	-	-	-	-	-	1	-	6	18	101
1999.....	30,614	1	-	-	-	-	-	-	1	5	26	97







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All origins, both sexes												
2003.....	12,286	1	14	28	11	11	13	19	37	81	240	438
2002.....	12,165	4	8	23	7	7	13	14	34	91	215	514
2001.....	12,078	1	22	30	5	8	12	19	33	74	284	498
2000.....	11,736	2	15	21	8	8	12	14	28	83	251	450
1999.....	11,116	2	19	23	10	8	9	25	29	105	237	427
All origins, male												
2003.....	7,667	1	4	15	4	7	9	9	27	56	175	324
2002.....	7,587	1	-	14	5	2	5	8	23	63	154	355
2001.....	7,524	1	10	7	2	7	9	11	16	58	182	355
2000.....	7,249	1	8	12	3	4	6	7	19	53	175	309
1999.....	6,867	1	11	12	4	4	7	15	22	70	166	310
All origins, female												
2003.....	4,619	-	10	13	7	4	4	10	10	25	65	114
2002.....	4,578	3	8	9	2	5	8	6	11	28	61	159
2001.....	4,554	-	12	23	3	1	3	8	17	16	102	143
2000.....	4,487	1	7	9	5	4	6	7	9	30	76	141
1999.....	4,249	1	8	11	6	4	2	10	7	35	71	117
Hispanic, both sexes												
2003.....	698	-	2	4	-	3	3	-	3	11	19	37
2002.....	659	2	3	1	1	-	1	2	4	9	28	43
2001.....	640	-	6	5	1	-	-	4	4	7	31	49
2000.....	567	-	3	2	1	-	3	3	4	7	25	42
1999.....	547	-	1	4	-	1	-	1	3	9	22	32

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Hispanic, male												
2003.....	426	-	-	2	-	3	2	-	3	8	15	28
2002.....	426	-	-	1	1	-	-	1	3	5	21	27
2001.....	426	-	1	3	1	-	-	3	2	7	27	37
2000.....	372	-	1	2	1	-	2	1	3	6	21	30
1999.....	353	-	1	2	-	-	-	-	2	7	16	28
Hispanic, female												
2003.....	272	-	2	2	-	-	1	-	-	3	4	9
2002.....	233	2	3	-	-	-	1	1	1	4	7	16
2001.....	214	-	5	2	-	-	-	1	2	-	4	12
2000.....	195	-	2	-	-	-	1	2	1	1	4	12
1999.....	194	-	-	2	-	1	-	1	1	2	6	4
Non-Hispanic 1/, both sexes												
2003.....	11,566	1	12	24	11	8	9	19	34	70	221	401
2002.....	11,477	2	5	22	6	7	12	12	30	82	186	471
2001.....	11,415	1	16	25	4	8	12	15	29	66	253	448
2000.....	11,124	2	12	19	7	8	9	11	24	76	226	408
1999.....	10,542	2	18	19	10	7	9	23	26	94	215	393
Non-Hispanic 1/, male												
2003.....	7,224	1	4	13	4	4	6	9	24	48	160	296
2002.....	7,141	1	-	13	4	2	5	7	20	58	132	328
2001.....	7,084	1	9	4	1	7	9	8	14	50	155	317
2000.....	6,847	1	7	10	2	4	4	6	16	47	154	279
1999.....	6,501	1	10	10	4	4	7	15	20	62	150	281

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Non-Hispanic 1/, female												
2003.....	4,342	-	8	11	7	4	3	10	10	22	61	105
2002.....	4,336	1	5	9	2	5	7	5	10	24	54	143
2001.....	4,331	-	7	21	3	1	3	7	15	16	98	131
2000.....	4,277	1	5	9	5	4	5	5	8	29	72	129
1999.....	4,041	1	8	9	6	3	2	8	6	32	65	112
Non-Hispanic White, both sexes												
2003.....	10,225	1	12	20	6	5	5	12	20	47	172	322
2002.....	10,179	2	5	18	5	5	10	8	16	57	154	385
2001.....	10,073	-	11	16	1	2	6	6	17	43	192	375
2000.....	9,884	-	7	10	6	6	5	5	19	52	187	339
1999.....	9,319	2	11	15	6	4	5	13	15	65	173	313
Non-Hispanic White, male												
2003.....	6,408	1	4	10	2	2	4	4	13	35	124	246
2002.....	6,363	1	-	11	3	1	4	4	11	40	110	277
2001.....	6,263	-	6	1	-	1	4	4	7	33	124	275
2000.....	6,082	-	4	4	2	3	3	1	12	36	129	236
1999.....	5,754	1	7	8	1	4	3	7	11	43	122	226
Non-Hispanic White, female												
2003.....	3,817	-	8	10	4	3	1	8	7	12	48	76
2002.....	3,816	1	5	7	2	4	6	4	5	17	44	108
2001.....	3,810	-	5	15	1	1	2	2	10	10	68	100
2000.....	3,802	-	3	6	4	3	2	4	7	16	58	103
1999.....	3,565	1	4	7	5	-	2	6	4	22	51	87

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Non-Hispanic Black, both sexes												
2003.....	1,105	-	-	4	5	3	4	7	12	19	44	65
2002.....	1,062	-	-	4	1	2	1	3	14	21	24	72
2001.....	1,105	-	3	9	3	5	5	9	10	21	51	63
2000.....	1,033	1	5	7	1	2	4	6	4	21	30	58
1999.....	1,028	-	4	3	3	3	4	10	10	27	34	67
Non-Hispanic Black, male												
2003.....	667	-	-	3	2	2	2	5	9	9	34	44
2002.....	626	-	-	2	1	1	1	2	9	15	16	41
2001.....	673	-	1	3	1	5	5	4	6	15	25	39
2000.....	625	-	3	4	-	1	1	5	3	9	18	35
1999.....	618	-	2	1	2	-	4	8	8	19	23	44
Non-Hispanic Black, female												
2003.....	438	-	-	1	3	1	2	2	3	10	10	21
2002.....	436	-	-	2	-	1	-	1	5	6	8	31
2001.....	432	-	2	6	2	-	-	5	4	6	26	24
2000.....	408	1	2	3	1	1	3	1	1	12	12	23
1999.....	410	-	2	2	1	3	-	2	2	8	11	23
Origin not stated 2/, both sexes												
2003.....	22	-	-	-	-	-	1	-	-	-	-	-
2002.....	29	-	-	-	-	-	-	-	-	-	1	-
2001.....	23	-	-	-	-	-	-	-	-	1	-	1
2000.....	45	-	-	-	-	-	-	-	-	-	-	-
1999.....	27	-	-	-	-	-	-	1	-	2	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Origin not stated 2/, male

2003.....	17	-	-	-	-	-	1	-	-	-	-	-
2002.....	20	-	-	-	-	-	-	-	-	-	1	-
2001.....	14	-	-	-	-	-	-	-	-	1	-	1
2000.....	30	-	-	-	-	-	-	-	-	-	-	-
1999.....	13	-	-	-	-	-	-	-	-	1	-	1

Origin not stated 2/, female

2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	9	-	-	-	-	-	-	-	-	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	-	-	-
2000.....	15	-	-	-	-	-	-	-	-	-	-	-
1999.....	14	-	-	-	-	-	-	1	-	1	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of bladder (C67)</b>												
<b>All origins, both sexes</b>												
2003.....	12,483	-	-	-	-	-	1	1	3	24	74	181
2002.....	12,628	-	-	2	1	-	1	3	11	31	92	182
2001.....	12,225	-	-	1	-	1	2	6	9	35	93	161
2000.....	12,002	-	-	-	1	2	1	1	9	40	71	160
1999.....	11,911	-	-	-	1	-	1	6	10	32	68	140
<b>All origins, male</b>												
2003.....	8,576	-	-	-	-	-	1	1	1	15	49	149
2002.....	8,504	-	-	1	-	-	-	2	7	21	57	129
2001.....	8,343	-	-	-	-	1	2	1	7	14	56	118
2000.....	8,163	-	-	-	-	2	1	-	2	21	41	110
1999.....	8,105	-	-	-	1	-	-	3	6	20	43	107
<b>All origins, female</b>												
2003.....	3,907	-	-	-	-	-	-	-	2	9	25	32
2002.....	4,124	-	-	1	1	-	1	1	4	10	35	53
2001.....	3,882	-	-	1	-	-	-	5	2	21	37	43
2000.....	3,839	-	-	-	1	-	-	1	7	19	30	50
1999.....	3,806	-	-	-	-	-	1	3	4	12	25	33
<b>Hispanic, both sexes</b>												
2003.....	394	-	-	-	-	-	-	-	1	-	4	6
2002.....	387	-	-	1	-	-	-	-	-	-	5	11
2001.....	336	-	-	-	-	1	1	-	-	2	2	12
2000.....	307	-	-	-	-	-	-	-	1	3	5	8
1999.....	301	-	-	-	-	-	-	-	2	1	3	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of bladder (C67)</b>												
<b>Hispanic, male</b>												
2003.....	265	-	-	-	-	-	-	-	-	-	2	5
2002.....	252	-	-	-	-	-	-	-	-	-	2	7
2001.....	223	-	-	-	-	1	1	-	-	-	1	7
2000.....	216	-	-	-	-	-	-	-	-	3	3	5
1999.....	222	-	-	-	-	-	-	-	2	1	3	4
<b>Hispanic, female</b>												
2003.....	129	-	-	-	-	-	-	-	1	-	2	1
2002.....	135	-	-	1	-	-	-	-	-	-	3	4
2001.....	113	-	-	-	-	-	-	-	-	2	1	5
2000.....	91	-	-	-	-	-	-	-	1	-	2	3
1999.....	79	-	-	-	-	-	-	-	-	-	-	1
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	12,067	-	-	-	-	-	1	1	2	24	70	174
2002.....	12,211	-	-	1	1	-	1	3	11	31	87	170
2001.....	11,858	-	-	1	-	-	1	6	9	33	91	147
2000.....	11,665	-	-	-	1	2	1	1	8	36	66	152
1999.....	11,582	-	-	-	1	-	1	6	8	31	65	135
<b>Non-Hispanic 1/, male</b>												
2003.....	8,294	-	-	-	-	-	1	1	1	15	47	143
2002.....	8,235	-	-	1	-	-	-	2	7	21	55	122
2001.....	8,100	-	-	-	-	-	1	1	7	14	55	110
2000.....	7,925	-	-	-	-	2	1	-	2	18	38	105
1999.....	7,865	-	-	-	1	-	-	3	4	19	40	103

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malignant neoplasm of bladder (C67)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	3,773	-	-	-	-	-	-	-	1	9	23	31
2002.....	3,976	-	-	-	1	-	1	1	4	10	32	48
2001.....	3,758	-	-	1	-	-	-	5	2	19	36	37
2000.....	3,740	-	-	-	1	-	-	1	6	18	28	47
1999.....	3,717	-	-	-	-	-	1	3	4	12	25	32
<b>Non-Hispanic White, both sexes</b>												
2003.....	11,003	-	-	-	-	-	1	1	2	21	51	138
2002.....	11,109	-	-	1	1	-	-	2	8	29	71	151
2001.....	10,823	-	-	1	-	-	1	6	8	28	72	121
2000.....	10,712	-	-	-	1	1	1	1	7	33	58	129
1999.....	10,548	-	-	-	1	-	-	3	5	23	54	109
<b>Non-Hispanic White, male</b>												
2003.....	7,692	-	-	-	-	-	1	1	1	13	35	117
2002.....	7,649	-	-	1	-	-	-	1	5	19	47	109
2001.....	7,541	-	-	-	-	-	1	1	7	13	43	95
2000.....	7,390	-	-	-	-	1	1	-	2	18	34	91
1999.....	7,290	-	-	-	1	-	-	1	2	13	33	83
<b>Non-Hispanic White, female</b>												
2003.....	3,311	-	-	-	-	-	-	-	1	8	16	21
2002.....	3,460	-	-	-	1	-	-	1	3	10	24	42
2001.....	3,282	-	-	1	-	-	-	5	1	15	29	26
2000.....	3,322	-	-	-	1	-	-	1	5	15	24	38
1999.....	3,258	-	-	-	-	-	-	2	3	10	21	26





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All origins, both sexes												
2003.....	12,901	18	104	175	161	119	117	165	240	411	668	857
2002.....	12,830	12	114	202	135	108	102	129	259	375	634	900
2001.....	12,609	16	100	170	130	104	120	134	282	399	629	830
2000.....	12,655	20	104	166	150	105	120	159	247	454	642	815
1999.....	12,765	10	113	184	124	115	103	177	267	443	622	842
All origins, male												
2003.....	7,122	9	56	89	86	68	68	85	147	246	393	544
2002.....	7,125	8	63	112	79	65	71	67	156	222	396	560
2001.....	7,015	8	52	81	64	63	66	88	174	245	393	500
2000.....	7,011	10	58	89	87	63	69	94	170	293	411	481
1999.....	7,001	8	56	96	58	69	62	104	172	268	370	507
All origins, female												
2003.....	5,779	9	48	86	75	51	49	80	93	165	275	313
2002.....	5,705	4	51	90	56	43	31	62	103	153	238	340
2001.....	5,594	8	48	89	66	41	54	46	108	154	236	330
2000.....	5,644	10	46	77	63	42	51	65	77	161	231	334
1999.....	5,764	2	57	88	66	46	41	73	95	175	252	335
Hispanic, both sexes												
2003.....	669	5	20	39	23	16	7	20	24	24	39	44
2002.....	649	2	21	29	23	15	9	14	19	21	34	50
2001.....	617	4	20	22	15	18	10	16	22	39	38	42
2000.....	631	4	16	36	27	17	11	20	25	30	46	41
1999.....	568	4	27	30	18	2	16	20	11	31	34	37

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Hispanic, male												
2003.....	368	2	10	21	15	9	6	10	16	15	23	30
2002.....	359	2	12	14	13	11	5	6	10	9	21	30
2001.....	335	1	10	12	4	13	6	11	10	21	20	29
2000.....	346	1	10	24	17	9	7	11	16	18	22	23
1999.....	293	4	11	13	9	2	8	12	5	20	25	18
Hispanic, female												
2003.....	301	3	10	18	8	7	1	10	8	9	16	14
2002.....	290	-	9	15	10	4	4	8	9	12	13	20
2001.....	282	3	10	10	11	5	4	5	12	18	18	13
2000.....	285	3	6	12	10	8	4	9	9	12	24	18
1999.....	275	-	16	17	9	-	8	8	6	11	9	19
Non-Hispanic 1/, both sexes												
2003.....	12,213	13	84	136	137	103	110	145	216	384	628	813
2002.....	12,152	10	93	172	112	91	93	115	240	352	599	848
2001.....	11,967	12	79	147	114	86	110	117	260	359	591	787
2000.....	11,995	16	87	130	123	88	109	139	222	423	596	773
1999.....	12,173	6	86	154	106	112	87	157	256	412	588	803
Non-Hispanic 1/, male												
2003.....	6,746	7	46	68	71	59	62	75	131	229	369	514
2002.....	6,752	6	51	98	66	54	66	61	146	212	374	530
2001.....	6,667	7	41	68	60	50	60	77	164	223	373	471
2000.....	6,646	9	47	65	70	54	62	83	154	274	389	457
1999.....	6,696	4	45	83	49	67	54	92	167	248	345	488

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic 1/, female												
2003.....	5,467	6	38	68	66	44	48	70	85	155	259	299
2002.....	5,400	4	42	74	46	37	27	54	94	140	225	318
2001.....	5,300	5	38	79	54	36	50	40	96	136	218	316
2000.....	5,349	7	40	65	53	34	47	56	68	149	207	316
1999.....	5,477	2	41	71	57	45	33	65	89	164	243	315
Non-Hispanic White, both sexes												
2003.....	11,179	10	65	101	108	83	91	125	189	339	572	725
2002.....	11,139	5	59	131	83	73	77	95	210	318	547	774
2001.....	11,004	8	63	121	84	70	85	103	232	334	530	711
2000.....	11,026	12	65	99	95	68	88	120	195	376	520	704
1999.....	11,238	3	70	110	85	83	69	145	227	377	525	736
Non-Hispanic White, male												
2003.....	6,178	6	38	49	54	46	51	65	116	200	333	466
2002.....	6,195	3	34	70	46	44	57	50	125	193	347	484
2001.....	6,154	5	35	53	43	41	47	70	147	210	337	432
2000.....	6,128	8	35	48	53	43	50	71	129	247	344	426
1999.....	6,193	2	39	63	41	48	43	85	148	229	304	444
Non-Hispanic White, female												
2003.....	5,001	4	27	52	54	37	40	60	73	139	239	259
2002.....	4,944	2	25	61	37	29	20	45	85	125	200	290
2001.....	4,850	3	28	68	41	29	38	33	85	124	193	279
2000.....	4,898	4	30	51	42	25	38	49	66	129	176	278
1999.....	5,045	1	31	47	44	35	26	60	79	148	221	292

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Non-Hispanic Black, both sexes												
2003.....	799	2	17	27	21	17	17	18	15	35	47	68
2002.....	766	3	27	34	20	14	12	11	22	30	38	49
2001.....	736	1	10	19	25	12	17	8	20	18	45	64
2000.....	762	2	18	26	25	16	19	15	21	32	56	51
1999.....	754	3	11	36	18	23	14	10	26	26	50	59
Non-Hispanic Black, male												
2003.....	440	1	6	16	11	11	10	9	9	23	32	37
2002.....	400	2	12	23	12	9	5	5	17	16	19	25
2001.....	378	-	3	12	13	6	10	4	10	8	25	34
2000.....	407	1	10	15	15	9	12	9	19	17	35	23
1999.....	399	2	4	17	7	15	9	6	17	15	32	39
Non-Hispanic Black, female												
2003.....	359	1	11	11	10	6	7	9	6	12	15	31
2002.....	366	1	15	11	8	5	7	6	5	14	19	24
2001.....	358	1	7	7	12	6	7	4	10	10	20	30
2000.....	355	1	8	11	10	7	7	6	2	15	21	28
1999.....	355	1	7	19	11	8	5	4	9	11	18	20
Origin not stated 2/, both sexes												
2003.....	19	-	-	-	1	-	-	-	-	3	1	-
2002.....	29	-	-	1	-	2	-	-	-	2	1	2
2001.....	25	-	1	1	1	-	-	1	-	1	-	1
2000.....	29	-	1	-	-	-	-	-	-	1	-	1
1999.....	24	-	-	-	-	1	-	-	-	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Origin not stated 2/, male

2003.....	8	-	-	-	-	-	-	-	-	2	1	-
2002.....	14	-	-	-	-	-	-	-	-	1	1	-
2001.....	13	-	1	1	-	-	-	-	-	1	-	-
2000.....	19	-	1	-	-	-	-	-	-	1	-	1
1999.....	12	-	-	-	-	-	-	-	-	-	-	1

Origin not stated 2/, female

2003.....	11	-	-	-	1	-	-	-	-	1	-	-
2002.....	15	-	-	1	-	2	-	-	-	1	-	2
2001.....	12	-	-	-	1	-	-	1	-	-	-	1
2000.....	10	-	-	-	-	-	-	-	-	-	-	-
1999.....	12	-	-	-	-	1	-	-	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All origins, both sexes												
2003.....	55,679	31	138	173	221	252	354	373	492	655	1,073	1,557
2002.....	56,225	28	119	166	213	310	393	368	548	695	1,056	1,599
2001.....	56,341	27	157	165	202	311	394	436	552	768	1,089	1,674
2000.....	56,518	39	146	168	204	323	395	412	506	776	1,077	1,675
1999.....	56,318	30	137	142	209	303	417	423	537	837	1,076	1,648
All origins, male												
2003.....	30,076	10	80	102	123	166	209	220	315	380	663	947
2002.....	30,302	15	63	87	130	188	229	225	320	409	629	953
2001.....	30,060	12	82	95	122	192	234	246	328	464	669	1,002
2000.....	29,888	21	76	98	125	202	253	226	313	438	637	1,015
1999.....	29,721	12	75	84	122	180	257	241	316	497	647	986
All origins, female												
2003.....	25,603	21	58	71	98	86	145	153	177	275	410	610
2002.....	25,923	13	56	79	83	122	164	143	228	286	427	646
2001.....	26,281	15	75	70	80	119	160	190	224	304	420	672
2000.....	26,630	18	70	70	79	121	142	186	193	338	440	660
1999.....	26,597	18	62	58	87	123	160	182	221	340	429	662
Hispanic, both sexes												
2003.....	2,836	6	44	42	55	48	83	60	87	83	119	159
2002.....	2,829	3	28	46	47	62	83	62	77	105	120	135
2001.....	2,747	7	38	46	41	77	67	75	79	90	121	118
2000.....	2,643	8	29	30	44	55	74	65	59	85	115	138
1999.....	2,538	8	28	28	47	45	83	63	72	88	102	141

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Hispanic, male												
2003.....	1,528	1	26	31	25	39	50	40	56	55	75	91
2002.....	1,518	1	12	21	26	44	53	43	49	62	70	76
2001.....	1,577	2	21	28	20	52	47	45	54	59	83	72
2000.....	1,460	5	14	15	32	34	49	44	39	43	72	95
1999.....	1,319	3	16	19	23	23	50	34	41	57	51	84
Hispanic, female												
2003.....	1,308	5	18	11	30	9	33	20	31	28	44	68
2002.....	1,311	2	16	25	21	18	30	19	28	43	50	59
2001.....	1,170	5	17	18	21	25	20	30	25	31	38	46
2000.....	1,183	3	15	15	12	21	25	21	20	42	43	43
1999.....	1,219	5	12	9	24	22	33	29	31	31	51	57
Non-Hispanic 1/, both sexes												
2003.....	52,736	25	94	130	166	203	271	312	405	572	953	1,396
2002.....	53,249	24	91	120	166	247	307	306	469	587	931	1,459
2001.....	53,458	20	117	118	161	233	326	357	471	675	965	1,549
2000.....	53,749	31	116	136	159	266	320	347	446	688	957	1,532
1999.....	53,669	22	108	114	162	258	334	360	463	749	970	1,503
Non-Hispanic 1/, male												
2003.....	28,485	9	54	70	98	126	159	179	259	325	587	854
2002.....	28,698	13	51	66	104	144	173	182	270	345	555	873
2001.....	28,410	10	60	67	102	140	186	200	273	404	583	925
2000.....	28,358	16	61	83	92	167	204	182	273	393	563	917
1999.....	28,347	9	58	65	99	157	207	207	274	440	592	900

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic 1/, female												
2003.....	24,251	16	40	60	68	77	112	133	146	247	366	542
2002.....	24,551	11	40	54	62	103	134	124	199	242	376	586
2001.....	25,048	10	57	51	59	93	140	157	198	271	382	624
2000.....	25,391	15	55	53	67	99	116	165	173	295	394	615
1999.....	25,322	13	50	49	63	101	127	153	189	309	378	603
Non-Hispanic White, both sexes												
2003.....	46,553	23	72	103	115	154	201	231	293	427	739	1,098
2002.....	47,058	15	72	94	119	183	247	225	345	441	742	1,138
2001.....	47,190	17	97	81	117	178	244	240	355	519	736	1,241
2000.....	47,695	30	92	105	127	192	228	251	334	531	752	1,204
1999.....	47,608	16	91	86	113	188	249	271	347	590	762	1,220
Non-Hispanic White, male												
2003.....	25,390	7	42	60	67	97	117	134	190	246	462	683
2002.....	25,523	9	42	50	75	112	142	137	196	256	445	695
2001.....	25,162	8	47	45	66	110	142	137	215	300	448	758
2000.....	25,272	16	50	62	72	120	148	136	202	311	464	733
1999.....	25,219	9	48	51	69	114	163	153	204	351	465	733
Non-Hispanic White, female												
2003.....	21,163	16	30	43	48	57	84	97	103	181	277	415
2002.....	21,535	6	30	44	44	71	105	88	149	185	297	443
2001.....	22,028	9	50	36	51	68	102	103	140	219	288	483
2000.....	22,423	14	42	43	55	72	80	115	132	220	288	471
1999.....	22,389	7	43	35	44	74	86	118	143	239	297	487

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Non-Hispanic Black, both sexes												
2003.....	5,059	1	14	20	37	34	54	65	90	117	182	249
2002.....	5,080	5	14	17	34	43	44	62	94	104	162	267
2001.....	5,162	3	13	27	35	46	61	84	97	133	189	255
2000.....	5,026	1	19	21	22	58	72	74	84	128	174	288
1999.....	5,038	4	14	21	40	53	80	71	86	134	173	236
Non-Hispanic Black, male												
2003.....	2,517	1	8	9	22	19	33	32	55	63	107	145
2002.....	2,593	2	6	12	20	24	20	35	58	66	96	155
2001.....	2,642	2	9	14	29	27	30	47	51	90	109	134
2000.....	2,535	-	8	14	12	36	43	33	57	64	85	159
1999.....	2,566	-	8	11	27	32	41	43	54	74	106	138
Non-Hispanic Black, female												
2003.....	2,542	-	6	11	15	15	21	33	35	54	75	104
2002.....	2,487	3	8	5	14	19	24	27	36	38	66	112
2001.....	2,520	1	4	13	6	19	31	37	46	43	80	121
2000.....	2,491	1	11	7	10	22	29	41	27	64	89	129
1999.....	2,472	4	6	10	13	21	39	28	32	60	67	98
Origin not stated 2/, both sexes												
2003.....	107	-	-	1	-	1	-	1	-	-	1	2
2002.....	147	1	-	-	-	1	3	-	2	3	5	5
2001.....	136	-	2	1	-	1	1	4	2	3	3	7
2000.....	126	-	1	2	1	2	1	-	1	3	5	5
1999.....	111	-	1	-	-	-	-	-	2	-	4	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Origin not stated 2/, male

2003.....	63	-	-	1	-	1	-	1	-	-	1	2
2002.....	86	1	-	-	-	-	3	-	1	2	4	4
2001.....	73	-	1	-	-	-	1	1	1	1	3	5
2000.....	70	-	1	-	1	1	-	-	1	2	2	3
1999.....	55	-	1	-	-	-	-	-	1	-	4	2

Origin not stated 2/, female

2003.....	44	-	-	-	-	-	-	-	-	-	-	-
2002.....	61	-	-	-	-	1	-	-	1	1	1	1
2001.....	63	-	1	1	-	1	-	3	1	2	-	2
2000.....	56	-	-	2	-	1	1	-	-	1	3	2
1999.....	56	-	-	-	-	-	-	-	1	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
All origins, both sexes												
2003.....	1,347	-	-	-	5	16	48	71	96	76	78	71
2002.....	1,352	-	-	3	3	27	59	64	88	81	78	83
2001.....	1,323	-	-	1	5	25	53	76	91	81	80	91
2000.....	1,287	-	-	3	5	21	52	60	85	78	73	73
1999.....	1,403	-	1	-	9	31	63	77	71	90	75	82
All origins, male												
2003.....	725	-	-	-	4	7	23	35	53	43	52	44
2002.....	751	-	-	2	3	15	26	37	44	46	46	53
2001.....	725	-	-	1	2	15	28	39	51	50	51	54
2000.....	686	-	-	1	3	6	29	31	48	42	40	46
1999.....	785	-	1	-	3	20	35	42	44	56	47	48
All origins, female												
2003.....	622	-	-	-	1	9	25	36	43	33	26	27
2002.....	601	-	-	1	-	12	33	27	44	35	32	30
2001.....	598	-	-	-	3	10	25	37	40	31	29	37
2000.....	601	-	-	2	2	15	23	29	37	36	33	27
1999.....	618	-	-	-	6	11	28	35	27	34	28	34
Hispanic, both sexes												
2003.....	91	-	-	-	1	2	6	6	10	3	3	7
2002.....	99	-	-	3	1	2	7	5	10	6	6	8
2001.....	99	-	-	1	-	4	3	7	6	6	7	3
2000.....	95	-	-	-	-	3	6	9	5	3	5	6
1999.....	113	-	-	-	3	2	6	2	5	5	7	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
Hispanic, male												
2003.....	52	-	-	-	1	2	3	4	4	3	2	3
2002.....	53	-	-	2	1	2	2	4	3	4	4	4
2001.....	66	-	-	1	-	3	1	3	5	3	5	1
2000.....	63	-	-	-	-	1	6	3	4	2	4	6
1999.....	65	-	-	-	2	2	4	1	2	4	4	3
Hispanic, female												
2003.....	39	-	-	-	-	-	3	2	6	-	1	4
2002.....	46	-	-	1	-	-	5	1	7	2	2	4
2001.....	33	-	-	-	-	1	2	4	1	3	2	2
2000.....	32	-	-	-	-	2	-	6	1	1	1	-
1999.....	48	-	-	-	1	-	2	1	3	1	3	3
Non-Hispanic 1/, both sexes												
2003.....	1,253	-	-	-	4	14	42	65	86	73	74	64
2002.....	1,248	-	-	-	2	25	51	59	77	74	72	75
2001.....	1,221	-	-	-	5	21	50	69	85	75	72	87
2000.....	1,189	-	-	3	5	18	46	51	80	75	68	67
1999.....	1,286	-	1	-	6	29	57	75	66	85	68	75
Non-Hispanic 1/, male												
2003.....	672	-	-	-	3	5	20	31	49	40	49	41
2002.....	693	-	-	-	2	13	23	33	40	41	42	49
2001.....	657	-	-	-	2	12	27	36	46	47	45	52
2000.....	621	-	-	1	3	5	23	28	44	40	36	40
1999.....	719	-	1	-	1	18	31	41	42	52	43	45

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hodgkin's disease (C81)												
Non-Hispanic 1/, female												
2003.....	581	-	-	-	1	9	22	34	37	33	25	23
2002.....	555	-	-	-	-	12	28	26	37	33	30	26
2001.....	564	-	-	-	3	9	23	33	39	28	27	35
2000.....	568	-	-	2	2	13	23	23	36	35	32	27
1999.....	567	-	-	-	5	11	26	34	24	33	25	30
Non-Hispanic White, both sexes												
2003.....	1,099	-	-	-	4	13	30	50	68	62	63	57
2002.....	1,093	-	-	-	1	19	44	47	65	58	60	64
2001.....	1,069	-	-	-	4	17	38	50	63	63	61	73
2000.....	1,059	-	-	2	3	14	34	46	71	63	56	60
1999.....	1,149	-	1	-	5	24	43	58	58	75	59	66
Non-Hispanic White, male												
2003.....	585	-	-	-	3	5	14	26	40	36	42	37
2002.....	599	-	-	-	1	11	21	24	33	32	33	40
2001.....	577	-	-	-	2	10	24	25	37	38	38	44
2000.....	551	-	-	1	1	3	20	25	38	34	30	36
1999.....	643	-	1	-	1	15	24	31	38	45	39	39
Non-Hispanic White, female												
2003.....	514	-	-	-	1	8	16	24	28	26	21	20
2002.....	494	-	-	-	-	8	23	23	32	26	27	24
2001.....	492	-	-	-	2	7	14	25	26	25	23	29
2000.....	508	-	-	1	2	11	14	21	33	29	26	24
1999.....	506	-	-	-	4	9	19	27	20	30	20	27





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
All origins, both sexes												
2003.....	21,475	-	5	17	28	53	87	96	156	233	397	649
2002.....	21,910	2	15	15	42	44	98	111	170	253	434	638
2001.....	22,305	1	1	11	27	61	87	129	169	301	455	711
2000.....	22,729	1	7	23	24	60	120	124	169	290	461	730
1999.....	22,802	2	16	12	25	51	93	109	180	324	465	696
All origins, male												
2003.....	11,399	-	5	12	14	34	54	59	106	153	253	414
2002.....	11,538	1	6	10	30	31	62	71	107	161	278	404
2001.....	11,710	1	1	8	21	43	51	72	107	195	305	450
2000.....	11,812	-	5	16	12	46	80	64	108	192	290	466
1999.....	11,794	-	9	9	21	30	61	69	109	204	306	459
All origins, female												
2003.....	10,076	-	-	5	14	19	33	37	50	80	144	235
2002.....	10,372	1	9	5	12	13	36	40	63	92	156	234
2001.....	10,595	-	-	3	6	18	36	57	62	106	150	261
2000.....	10,917	1	2	7	12	14	40	60	61	98	171	264
1999.....	11,008	2	7	3	4	21	32	40	71	120	159	237
Hispanic, both sexes												
2003.....	1,057	-	3	4	6	7	20	16	30	35	47	60
2002.....	1,068	-	2	3	4	5	20	19	22	32	46	52
2001.....	1,008	-	-	2	4	11	11	20	17	33	44	55
2000.....	1,026	-	2	2	-	4	17	19	19	26	57	51
1999.....	958	-	1	1	2	5	14	10	27	25	43	64

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
Hispanic, male												
2003.....	569	-	3	2	3	6	13	14	19	25	31	34
2002.....	549	-	-	1	2	5	14	14	11	19	26	34
2001.....	579	-	-	2	4	7	8	11	13	22	34	35
2000.....	560	-	2	1	-	4	13	11	12	16	35	34
1999.....	493	-	-	1	1	4	6	5	13	19	24	42
Hispanic, female												
2003.....	488	-	-	2	3	1	7	2	11	10	16	26
2002.....	519	-	2	2	2	-	6	5	11	13	20	18
2001.....	429	-	-	-	-	4	3	9	4	11	10	20
2000.....	466	-	-	1	-	-	4	8	7	10	22	17
1999.....	465	-	1	-	1	1	8	5	14	6	19	22
Non-Hispanic 1/, both sexes												
2003.....	20,379	-	2	13	22	45	67	80	126	198	350	588
2002.....	20,780	2	13	12	38	38	77	92	148	219	384	583
2001.....	21,246	1	1	9	23	49	76	108	150	268	410	653
2000.....	21,657	1	5	21	24	56	103	105	149	264	404	677
1999.....	21,792	2	15	11	23	46	79	99	153	299	419	630
Non-Hispanic 1/, male												
2003.....	10,810	-	2	10	11	27	41	45	87	128	222	379
2002.....	10,953	1	6	9	28	26	47	57	96	141	248	368
2001.....	11,105	1	1	6	17	36	43	60	93	173	270	413
2000.....	11,225	-	3	15	12	42	67	53	95	176	255	432
1999.....	11,271	-	9	8	20	26	55	64	96	185	279	416

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic 1/, female												
2003.....	9,569	-	-	3	11	18	26	35	39	70	128	209
2002.....	9,827	1	7	3	10	12	30	35	52	78	136	215
2001.....	10,141	-	-	3	6	13	33	48	57	95	140	240
2000.....	10,432	1	2	6	12	14	36	52	54	88	149	245
1999.....	10,521	2	6	3	3	20	24	35	57	114	140	214
Non-Hispanic White, both sexes												
2003.....	18,526	-	2	10	12	32	53	50	82	148	269	464
2002.....	18,901	1	12	12	26	23	57	65	97	164	299	479
2001.....	19,300	1	1	8	18	33	54	69	113	201	322	528
2000.....	19,843	-	2	16	22	42	68	70	111	202	329	553
1999.....	19,956	-	12	7	16	31	60	69	108	228	331	517
Non-Hispanic White, male												
2003.....	9,871	-	2	8	4	21	32	30	58	100	171	298
2002.....	9,961	-	6	9	20	18	37	44	62	104	193	314
2001.....	10,052	1	1	5	12	25	29	40	75	127	212	342
2000.....	10,264	-	2	11	12	34	45	39	70	136	217	355
1999.....	10,259	-	8	6	15	18	43	43	69	144	218	344
Non-Hispanic White, female												
2003.....	8,655	-	-	2	8	11	21	20	24	48	98	166
2002.....	8,940	1	6	3	6	5	20	21	35	60	106	165
2001.....	9,248	-	-	3	6	8	25	29	38	74	110	186
2000.....	9,579	-	-	5	10	8	23	31	41	66	112	198
1999.....	9,697	-	4	1	1	13	17	26	39	84	113	173





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Leukemia (C91-C95)</b>												
<b>All origins, both sexes</b>												
2003.....	21,535	30	133	156	186	183	219	205	228	321	475	595
2002.....	21,498	21	104	148	168	237	235	190	281	325	442	606
2001.....	21,451	25	155	153	170	225	254	230	283	346	433	630
2000.....	21,339	38	137	142	173	242	222	228	245	368	433	622
1999.....	21,014	27	119	130	175	221	261	234	266	374	425	626
<b>All origins, male</b>												
2003.....	12,073	9	75	90	105	125	132	125	149	167	277	347
2002.....	12,016	11	57	75	97	142	140	117	163	178	251	335
2001.....	11,850	10	81	86	99	134	155	135	165	196	236	346
2000.....	11,777	21	69	81	109	150	143	131	152	185	238	356
1999.....	11,519	11	65	75	98	130	161	128	148	204	231	346
<b>All origins, female</b>												
2003.....	9,462	21	58	66	81	58	87	80	79	154	198	248
2002.....	9,482	10	47	73	71	95	95	73	118	147	191	271
2001.....	9,601	15	74	67	71	91	99	95	118	150	197	284
2000.....	9,562	17	68	61	64	92	79	97	93	183	195	266
1999.....	9,495	16	54	55	77	91	100	106	118	170	194	280
<b>Hispanic, both sexes</b>												
2003.....	1,156	6	41	38	48	39	57	38	46	40	60	77
2002.....	1,088	2	26	40	42	54	55	38	42	63	58	51
2001.....	1,114	7	38	43	37	62	53	47	54	46	57	49
2000.....	1,038	8	27	28	44	48	51	37	34	52	45	58
1999.....	994	8	26	27	42	38	63	49	37	52	46	50

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Leukemia (C91-C95)</b>												
<b>Hispanic, male</b>												
2003.....	649	1	23	29	21	31	34	22	32	23	35	46
2002.....	601	1	12	18	23	37	36	25	32	35	34	22
2001.....	640	2	21	25	16	42	38	31	35	31	34	30
2000.....	595	5	12	14	32	29	30	30	22	24	28	38
1999.....	534	3	16	18	20	17	40	26	23	28	20	30
<b>Hispanic, female</b>												
2003.....	507	5	18	9	27	8	23	16	14	17	25	31
2002.....	487	1	14	22	19	17	19	13	10	28	24	29
2001.....	474	5	17	18	21	20	15	16	19	15	23	19
2000.....	443	3	15	14	12	19	21	7	12	28	17	20
1999.....	460	5	10	9	22	21	23	23	14	24	26	20
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	20,340	24	92	117	138	144	162	166	182	281	415	517
2002.....	20,363	18	78	108	126	183	179	152	238	262	383	554
2001.....	20,285	18	115	109	133	163	200	180	229	298	375	578
2000.....	20,248	30	109	112	128	192	170	191	211	313	384	563
1999.....	19,981	19	92	103	133	183	198	185	227	322	378	575
<b>Non-Hispanic 1/, male</b>												
2003.....	11,394	8	52	60	84	94	98	102	117	144	242	300
2002.....	11,386	9	45	57	74	105	103	92	131	143	217	312
2001.....	11,180	8	59	61	83	92	116	104	130	164	201	314
2000.....	11,155	16	56	67	76	120	113	101	130	159	208	317
1999.....	10,968	8	48	57	78	113	121	102	124	176	210	315

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Leukemia (C91-C95)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	8,946	16	40	57	54	50	64	64	65	137	173	217
2002.....	8,977	9	33	51	52	78	76	60	107	119	166	242
2001.....	9,105	10	56	48	50	71	84	76	99	134	174	264
2000.....	9,093	14	53	45	52	72	57	90	81	154	176	246
1999.....	9,013	11	44	46	55	70	77	83	103	146	168	260
<b>Non-Hispanic White, both sexes</b>												
2003.....	18,190	22	70	93	97	109	118	130	135	205	329	432
2002.....	18,162	13	60	82	92	140	146	111	178	200	324	429
2001.....	18,105	15	95	73	95	128	152	121	177	231	293	479
2000.....	18,128	30	88	87	100	136	126	135	149	245	301	441
1999.....	17,885	15	78	79	92	133	146	144	171	256	308	473
<b>Non-Hispanic White, male</b>												
2003.....	10,278	6	40	52	60	71	71	77	87	101	195	259
2002.....	10,218	8	36	41	54	83	84	69	99	108	186	240
2001.....	10,018	6	46	40	52	75	89	72	101	123	154	267
2000.....	10,009	16	46	50	58	83	83	72	92	128	170	253
1999.....	9,853	8	39	45	53	81	96	79	91	142	171	255
<b>Non-Hispanic White, female</b>												
2003.....	7,912	16	30	41	37	38	47	53	48	104	134	173
2002.....	7,944	5	24	41	38	57	62	42	79	92	138	189
2001.....	8,087	9	49	33	43	53	63	49	76	108	139	212
2000.....	8,119	14	42	37	42	53	43	63	57	117	131	188
1999.....	8,032	7	39	34	39	52	50	65	80	114	137	218

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Leukemia (C91-C95)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	1,745	1	14	17	30	23	32	26	38	58	72	67
2002.....	1,752	4	13	17	23	29	24	28	40	42	47	93
2001.....	1,741	3	13	26	30	26	34	37	38	53	65	75
2000.....	1,729	-	17	16	20	46	34	39	49	49	67	107
1999.....	1,694	2	12	18	34	35	47	29	38	52	52	79
<b>Non-Hispanic Black, male</b>												
2003.....	908	1	8	7	16	15	20	17	23	33	40	33
2002.....	914	1	6	12	12	18	12	16	21	25	23	57
2001.....	915	2	9	13	25	14	17	21	23	32	35	32
2000.....	923	-	8	11	11	31	23	18	32	20	32	54
1999.....	887	-	7	9	23	23	22	16	22	25	29	47
<b>Non-Hispanic Black, female</b>												
2003.....	837	-	6	10	14	8	12	9	15	25	32	34
2002.....	838	3	7	5	11	11	12	12	19	17	24	36
2001.....	826	1	4	13	5	12	17	16	15	21	30	43
2000.....	806	-	9	5	9	15	11	21	17	29	35	53
1999.....	807	2	5	9	11	12	25	13	16	27	23	32
<b>Origin not stated 2/, both sexes</b>												
2003.....	39	-	-	1	-	-	-	1	-	-	-	1
2002.....	47	1	-	-	-	-	1	-	1	-	1	1
2001.....	52	-	2	1	-	-	1	3	-	2	1	3
2000.....	53	-	1	2	1	2	1	-	-	3	4	1
1999.....	39	-	1	-	-	-	-	-	2	-	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Leukemia (C91-C95)												
Origin not stated 2/, male												
2003.....	30	-	-	1	-	-	-	1	-	-	-	1
2002.....	29	1	-	-	-	-	1	-	-	-	-	1
2001.....	30	-	1	-	-	-	1	-	-	1	1	2
2000.....	27	-	1	-	1	1	-	-	-	2	2	1
1999.....	17	-	1	-	-	-	-	-	1	-	1	1
Origin not stated 2/, female												
2003.....	9	-	-	-	-	-	-	-	-	-	-	-
2002.....	18	-	-	-	-	-	-	-	1	-	1	-
2001.....	22	-	1	1	-	-	-	3	-	1	-	1
2000.....	26	-	-	2	-	1	1	-	-	1	2	-
1999.....	22	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All origins, both sexes												
2003.....	11,267	-	-	-	1	-	-	1	11	25	122	238
2002.....	11,392	2	-	-	-	2	-	3	9	36	102	270
2001.....	11,172	-	-	-	-	-	-	1	8	39	118	241
2000.....	11,063	-	-	-	1	-	-	-	7	38	108	245
1999.....	11,016	-	-	-	-	-	-	3	19	47	109	241
All origins, male												
2003.....	5,848	-	-	-	-	-	-	1	6	17	80	139
2002.....	5,953	-	-	-	-	-	-	-	6	24	54	159
2001.....	5,724	-	-	-	-	-	-	-	4	22	75	151
2000.....	5,553	-	-	-	-	-	-	-	5	18	67	143
1999.....	5,584	-	-	-	-	-	-	2	14	33	62	132
All origins, female												
2003.....	5,419	-	-	-	1	-	-	-	5	8	42	99
2002.....	5,439	2	-	-	-	2	-	3	3	12	48	111
2001.....	5,448	-	-	-	-	-	-	1	4	17	43	90
2000.....	5,510	-	-	-	1	-	-	-	2	20	41	102
1999.....	5,432	-	-	-	-	-	-	1	5	14	47	109
Hispanic, both sexes												
2003.....	531	-	-	-	-	-	-	-	-	5	9	15
2002.....	572	1	-	-	-	1	-	-	3	4	10	24
2001.....	524	-	-	-	-	-	-	1	2	5	13	11
2000.....	478	-	-	-	-	-	-	-	1	4	8	22
1999.....	468	-	-	-	-	-	-	2	2	6	6	21

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Hispanic, male												
2003.....	257	-	-	-	-	-	-	-	-	4	7	8
2002.....	314	-	-	-	-	-	-	-	3	4	6	16
2001.....	291	-	-	-	-	-	-	-	1	3	10	6
2000.....	239	-	-	-	-	-	-	-	1	1	5	16
1999.....	225	-	-	-	-	-	-	2	2	6	3	9
Hispanic, female												
2003.....	274	-	-	-	-	-	-	-	-	1	2	7
2002.....	258	1	-	-	-	1	-	-	-	-	4	8
2001.....	233	-	-	-	-	-	-	1	1	2	3	5
2000.....	239	-	-	-	-	-	-	-	-	3	3	6
1999.....	243	-	-	-	-	-	-	-	-	-	3	12
Non-Hispanic 1/, both sexes												
2003.....	10,710	-	-	-	1	-	-	1	11	20	113	223
2002.....	10,787	1	-	-	-	1	-	3	6	32	92	245
2001.....	10,618	-	-	-	-	-	-	-	6	33	105	230
2000.....	10,561	-	-	-	1	-	-	-	6	34	99	221
1999.....	10,532	-	-	-	-	-	-	1	17	41	103	220
Non-Hispanic 1/, male												
2003.....	5,579	-	-	-	-	-	-	1	6	13	73	131
2002.....	5,623	-	-	-	-	-	-	-	3	20	48	142
2001.....	5,418	-	-	-	-	-	-	-	3	19	65	145
2000.....	5,300	-	-	-	-	-	-	-	4	17	62	125
1999.....	5,352	-	-	-	-	-	-	-	12	27	59	123

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Non-Hispanic 1/, female												
2003.....	5,131	-	-	-	1	-	-	-	5	7	40	92
2002.....	5,164	1	-	-	-	1	-	3	3	12	44	103
2001.....	5,200	-	-	-	-	-	-	-	3	14	40	85
2000.....	5,261	-	-	-	1	-	-	-	2	17	37	96
1999.....	5,180	-	-	-	-	-	-	1	5	14	44	97
Non-Hispanic White, both sexes												
2003.....	8,689	-	-	-	1	-	-	1	8	12	77	143
2002.....	8,843	-	-	-	-	1	-	2	5	19	59	164
2001.....	8,641	-	-	-	-	-	-	-	2	24	57	160
2000.....	8,582	-	-	-	1	-	-	-	3	20	64	148
1999.....	8,547	-	-	-	-	-	-	-	10	29	63	162
Non-Hispanic White, male												
2003.....	4,629	-	-	-	-	-	-	1	5	9	53	88
2002.....	4,708	-	-	-	-	-	-	-	2	12	33	99
2001.....	4,472	-	-	-	-	-	-	-	2	12	42	104
2000.....	4,400	-	-	-	-	-	-	-	2	13	45	88
1999.....	4,430	-	-	-	-	-	-	-	6	20	36	94
Non-Hispanic White, female												
2003.....	4,060	-	-	-	1	-	-	-	3	3	24	55
2002.....	4,135	-	-	-	-	1	-	2	3	7	26	65
2001.....	4,169	-	-	-	-	-	-	-	-	12	15	56
2000.....	4,182	-	-	-	1	-	-	-	1	7	19	60
1999.....	4,117	-	-	-	-	-	-	-	4	9	27	68





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All origins, both sexes

2003.....	55	1	-	-	1	-	-	-	1	-	1	4
2002.....	73	3	-	-	-	-	1	-	-	-	-	2
2001.....	90	1	1	-	-	-	-	-	1	1	3	1
2000.....	100	-	2	-	1	-	1	-	-	2	2	5
1999.....	83	1	1	-	-	-	-	-	1	2	2	3

All origins, male

2003.....	31	1	-	-	-	-	-	-	1	-	1	3
2002.....	44	3	-	-	-	-	1	-	-	-	-	2
2001.....	51	1	-	-	-	-	-	-	1	1	2	1
2000.....	60	-	2	-	1	-	1	-	-	1	2	4
1999.....	39	1	-	-	-	-	-	-	1	-	1	1

All origins, female

2003.....	24	-	-	-	1	-	-	-	-	-	-	1
2002.....	29	-	-	-	-	-	-	-	-	-	-	-
2001.....	39	-	1	-	-	-	-	-	-	-	1	-
2000.....	40	-	-	-	-	-	-	-	-	1	-	1
1999.....	44	-	1	-	-	-	-	-	-	2	1	2

Hispanic, both sexes

2003.....	1	-	-	-	-	-	-	-	1	-	-	-
2002.....	2	-	-	-	-	-	1	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	-	-	-	-	1
1999.....	5	-	1	-	-	-	-	-	1	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Non-Hispanic 1/, female												
2003.....	24	-	-	-	1	-	-	-	-	-	-	1
2002.....	28	-	-	-	-	-	-	-	-	-	-	-
2001.....	38	-	1	-	-	-	-	-	-	-	1	-
2000.....	37	-	-	-	-	-	-	-	-	1	-	1
1999.....	41	-	-	-	-	-	-	-	-	2	1	2
Non-Hispanic White, both sexes												
2003.....	49	1	-	-	1	-	-	-	-	-	1	2
2002.....	59	1	-	-	-	-	-	-	-	-	-	2
2001.....	75	1	1	-	-	-	-	-	-	-	3	1
2000.....	83	-	2	-	1	-	-	-	-	1	2	2
1999.....	71	1	-	-	-	-	-	-	-	2	1	2
Non-Hispanic White, male												
2003.....	27	1	-	-	-	-	-	-	-	-	1	1
2002.....	37	1	-	-	-	-	-	-	-	-	-	2
2001.....	43	1	-	-	-	-	-	-	-	-	2	1
2000.....	48	-	2	-	1	-	-	-	-	-	2	1
1999.....	34	1	-	-	-	-	-	-	-	-	1	1
Non-Hispanic White, female												
2003.....	22	-	-	-	1	-	-	-	-	-	-	1
2002.....	22	-	-	-	-	-	-	-	-	-	-	-
2001.....	32	-	1	-	-	-	-	-	-	-	1	-
2000.....	35	-	-	-	-	-	-	-	-	1	-	1
1999.....	37	-	-	-	-	-	-	-	-	2	-	1





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
All origins, both sexes												
2003.....	62,639	19	106	127	147	242	274	298	392	664	1,221	2,234
2002.....	62,969	24	124	133	155	233	294	289	445	716	1,327	2,213
2001.....	62,708	17	117	116	163	242	274	316	412	718	1,337	2,187
2000.....	63,060	24	129	123	137	243	268	292	409	771	1,319	2,114
1999.....	66,251	14	116	150	141	246	273	307	455	833	1,481	2,247
All origins, male												
2003.....	31,845	12	66	80	72	161	168	181	246	377	669	1,274
2002.....	32,122	15	65	77	75	140	200	194	265	407	703	1,225
2001.....	31,663	7	73	64	89	152	194	190	252	385	723	1,231
2000.....	31,728	13	75	64	76	142	177	189	243	424	717	1,186
1999.....	33,449	8	64	80	80	158	171	204	286	464	836	1,274
All origins, female												
2003.....	30,794	7	40	47	75	81	106	117	146	287	552	960
2002.....	30,847	9	59	56	80	93	94	95	180	309	624	988
2001.....	31,045	10	44	52	74	90	80	126	160	333	614	956
2000.....	31,332	11	54	59	61	101	91	103	166	347	602	928
1999.....	32,802	6	52	70	61	88	102	103	169	369	645	973
Hispanic, both sexes												
2003.....	3,085	2	21	24	27	49	45	51	59	86	107	155
2002.....	3,005	11	17	23	32	38	69	41	66	71	105	143
2001.....	2,844	2	15	13	29	41	44	36	46	82	106	143
2000.....	2,753	3	24	19	28	44	38	46	54	58	83	118
1999.....	2,747	2	11	23	22	39	45	48	51	68	107	144

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Hispanic, male												
2003.....	1,621	1	15	15	12	35	25	40	44	46	57	99
2002.....	1,566	7	9	14	16	25	44	31	38	39	55	76
2001.....	1,455	-	9	6	16	25	39	21	35	42	53	78
2000.....	1,394	2	16	11	15	25	24	29	38	33	45	60
1999.....	1,399	2	3	10	13	27	35	33	30	36	54	75
Hispanic, female												
2003.....	1,464	1	6	9	15	14	20	11	15	40	50	56
2002.....	1,439	4	8	9	16	13	25	10	28	32	50	67
2001.....	1,389	2	6	7	13	16	5	15	11	40	53	65
2000.....	1,359	1	8	8	13	19	14	17	16	25	38	58
1999.....	1,348	-	8	13	9	12	10	15	21	32	53	69
Non-Hispanic 1/, both sexes												
2003.....	59,402	17	84	103	120	193	227	246	332	575	1,111	2,073
2002.....	59,797	13	106	110	123	194	225	248	377	645	1,222	2,058
2001.....	59,685	15	102	103	134	201	230	280	365	632	1,226	2,037
2000.....	60,117	21	103	103	109	197	230	244	352	711	1,228	1,988
1999.....	63,328	12	105	127	118	207	227	257	398	761	1,369	2,095
Non-Hispanic 1/, male												
2003.....	30,140	11	50	65	60	126	142	140	201	330	610	1,170
2002.....	30,463	8	56	63	59	114	156	163	226	368	648	1,140
2001.....	30,107	7	64	58	73	127	155	169	216	339	667	1,149
2000.....	30,237	11	57	53	61	116	153	158	204	391	668	1,121
1999.....	31,956	6	61	70	67	131	135	169	254	427	779	1,194

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Non-Hispanic 1/, female												
2003.....	29,262	6	34	38	60	67	85	106	131	245	501	903
2002.....	29,334	5	50	47	64	80	69	85	151	277	574	918
2001.....	29,578	8	38	45	61	74	75	111	149	293	559	888
2000.....	29,880	10	46	50	48	81	77	86	148	320	560	867
1999.....	31,372	6	44	57	51	76	92	88	144	334	590	901
Non-Hispanic White, both sexes												
2003.....	51,534	13	58	82	83	154	174	185	239	451	836	1,586
2002.....	51,731	11	73	79	92	153	166	185	280	494	939	1,590
2001.....	51,581	13	83	79	97	156	182	219	273	482	948	1,558
2000.....	52,223	16	75	87	81	142	186	203	276	537	915	1,532
1999.....	54,977	10	87	95	83	158	166	207	306	587	1,039	1,624
Non-Hispanic White, male												
2003.....	26,189	9	31	53	41	100	109	114	147	271	458	888
2002.....	26,457	8	40	45	43	88	113	123	172	284	485	892
2001.....	26,007	6	50	44	52	96	124	132	164	266	526	890
2000.....	26,258	9	46	44	45	84	120	134	169	304	493	864
1999.....	27,780	5	52	55	49	94	96	143	195	339	604	915
Non-Hispanic White, female												
2003.....	25,345	4	27	29	42	54	65	71	92	180	378	698
2002.....	25,274	3	33	34	49	65	53	62	108	210	454	698
2001.....	25,574	7	33	35	45	60	58	87	109	216	422	668
2000.....	25,965	7	29	43	36	58	66	69	107	233	422	668
1999.....	27,197	5	35	40	34	64	70	64	111	248	435	709

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Non-Hispanic Black, both sexes												
2003.....	6,537	2	22	20	31	29	41	49	73	104	222	415
2002.....	6,756	2	26	26	26	33	46	51	74	131	231	398
2001.....	6,725	1	16	19	27	37	39	52	72	122	217	409
2000.....	6,617	4	22	16	23	47	36	34	53	152	275	392
1999.....	7,009	1	14	22	27	39	44	41	65	144	277	416
Non-Hispanic Black, male												
2003.....	3,274	2	15	11	16	19	26	21	39	47	124	239
2002.....	3,340	-	12	15	13	22	32	31	36	70	130	214
2001.....	3,392	-	13	12	14	23	28	31	38	58	108	215
2000.....	3,337	1	9	9	15	27	29	20	26	74	153	227
1999.....	3,514	-	8	9	13	29	27	20	40	70	149	244
Non-Hispanic Black, female												
2003.....	3,263	-	7	9	15	10	15	28	34	57	98	176
2002.....	3,416	2	14	11	13	11	14	20	38	61	101	184
2001.....	3,333	1	3	7	13	14	11	21	34	64	109	194
2000.....	3,280	3	13	7	8	20	7	14	27	78	122	165
1999.....	3,495	1	6	13	14	10	17	21	25	74	128	172
Origin not stated 2/, both sexes												
2003.....	152	-	1	-	-	-	2	1	1	3	3	6
2002.....	167	-	1	-	-	1	-	-	2	-	-	12
2001.....	179	-	-	-	-	-	-	-	1	4	5	7
2000.....	190	-	2	1	-	2	-	2	3	2	8	8
1999.....	176	-	-	-	1	-	1	2	6	4	5	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Origin not stated 2/, male

2003.....	84	-	1	-	-	-	1	1	1	1	2	5
2002.....	93	-	-	-	-	1	-	-	1	-	-	9
2001.....	101	-	-	-	-	-	-	-	1	4	3	4
2000.....	97	-	2	-	-	1	-	2	1	-	4	5
1999.....	94	-	-	-	-	-	1	2	2	1	3	5

Origin not stated 2/, female

2003.....	68	-	-	-	-	-	1	-	-	2	1	1
2002.....	74	-	1	-	-	-	-	-	1	-	-	3
2001.....	78	-	-	-	-	-	-	-	-	-	2	3
2000.....	93	-	-	1	-	1	-	-	2	2	4	3
1999.....	82	-	-	-	1	-	-	-	4	3	2	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins, both sexes

2003.....	13,563	62	51	38	41	41	55	69	109	142	197	295
2002.....	13,299	69	60	44	45	41	46	62	101	137	229	327
2001.....	13,676	51	58	52	53	47	55	60	91	156	210	304
2000.....	13,546	60	53	62	37	40	55	68	97	155	213	293
1999.....	13,227	58	63	64	37	52	46	55	116	140	202	276

All origins, male

2003.....	6,687	32	31	21	22	24	30	40	58	79	116	148
2002.....	6,536	30	30	22	27	21	29	33	51	77	119	169
2001.....	6,735	26	30	26	26	20	33	33	44	79	114	173
2000.....	6,559	32	33	32	17	27	35	30	52	86	102	160
1999.....	6,520	30	33	30	19	26	28	32	64	79	118	142

All origins, female

2003.....	6,876	30	20	17	19	17	25	29	51	63	81	147
2002.....	6,763	39	30	22	18	20	17	29	50	60	110	158
2001.....	6,941	25	28	26	27	27	22	27	47	77	96	131
2000.....	6,987	28	20	30	20	13	20	38	45	69	111	133
1999.....	6,707	28	30	34	18	26	18	23	52	61	84	134

Hispanic, both sexes

2003.....	603	15	7	8	5	6	15	13	19	10	15	23
2002.....	563	17	7	16	10	4	4	5	11	14	20	30
2001.....	543	5	13	14	7	4	4	8	12	14	25	22
2000.....	521	6	7	5	8	3	7	8	17	14	23	20
1999.....	497	8	12	11	6	7	5	8	17	6	24	15

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, male

2003.....	301	6	3	4	3	5	8	10	11	5	9	9
2002.....	282	9	4	7	8	2	3	3	5	7	8	16
2001.....	276	2	7	9	3	2	4	4	2	6	18	14
2000.....	237	4	2	4	4	2	6	2	13	7	7	9
1999.....	233	5	4	5	3	5	3	7	7	4	17	9

Hispanic, female

2003.....	302	9	4	4	2	1	7	3	8	5	6	14
2002.....	281	8	3	9	2	2	1	2	6	7	12	14
2001.....	267	3	6	5	4	2	-	4	10	8	7	8
2000.....	284	2	5	1	4	1	1	6	4	7	16	11
1999.....	264	3	8	6	3	2	2	1	10	2	7	6

Non-Hispanic 1/, both sexes

2003.....	12,938	46	44	30	36	35	40	56	90	132	181	272
2002.....	12,708	52	53	28	35	37	41	57	90	123	205	297
2001.....	13,098	46	45	38	46	43	51	52	79	142	185	282
2000.....	12,990	53	46	57	28	37	48	60	80	141	189	271
1999.....	12,699	50	51	52	31	44	41	47	99	134	177	261

Non-Hispanic 1/, male

2003.....	6,372	26	28	17	19	19	22	30	47	74	106	139
2002.....	6,237	21	26	15	19	19	25	30	46	70	109	153
2001.....	6,442	24	23	17	23	18	29	29	42	73	96	159
2000.....	6,302	28	31	28	13	25	29	28	39	79	94	149
1999.....	6,276	25	29	25	16	21	25	25	57	75	101	133

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Non-Hispanic 1/, female												
2003.....	6,566	20	16	13	17	16	18	26	43	58	75	133
2002.....	6,471	31	27	13	16	18	16	27	44	53	96	144
2001.....	6,656	22	22	21	23	25	22	23	37	69	89	123
2000.....	6,688	25	15	29	15	12	19	32	41	62	95	122
1999.....	6,423	25	22	27	15	23	16	22	42	59	76	128
Non-Hispanic White, both sexes												
2003.....	11,572	33	32	19	24	26	32	42	71	91	145	209
2002.....	11,332	42	36	20	21	30	29	37	68	98	153	245
2001.....	11,701	32	36	24	32	29	37	45	61	110	142	235
2000.....	11,636	41	37	41	22	26	37	40	65	110	145	206
1999.....	11,335	34	36	40	26	30	28	31	68	101	138	199
Non-Hispanic White, male												
2003.....	5,732	20	19	10	14	15	19	22	35	57	86	112
2002.....	5,607	18	17	9	13	16	16	21	41	58	91	131
2001.....	5,801	18	18	10	15	10	21	25	33	58	77	135
2000.....	5,653	21	23	20	11	16	24	22	33	61	71	118
1999.....	5,653	17	18	19	13	12	18	14	39	59	84	107
Non-Hispanic White, female												
2003.....	5,840	13	13	9	10	11	13	20	36	34	59	97
2002.....	5,725	24	19	11	8	14	13	16	27	40	62	114
2001.....	5,900	14	18	14	17	19	16	20	28	52	65	100
2000.....	5,983	20	14	21	11	10	13	18	32	49	74	88
1999.....	5,682	17	18	21	13	18	10	17	29	42	54	92

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic Black, both sexes

2003.....	1,111	11	9	9	8	7	8	13	15	37	30	58
2002.....	1,134	9	14	7	11	6	9	13	18	20	47	49
2001.....	1,143	11	5	13	13	12	9	6	12	26	35	35
2000.....	1,122	9	9	13	5	8	8	18	15	24	38	54
1999.....	1,161	14	13	12	5	12	13	9	21	30	35	50

Non-Hispanic Black, male

2003.....	516	5	8	5	3	3	3	8	9	15	16	25
2002.....	496	2	8	6	3	3	6	5	3	8	17	19
2001.....	522	5	3	6	7	7	6	3	8	13	17	16
2000.....	514	4	8	6	2	7	2	5	6	13	19	23
1999.....	516	7	10	6	3	8	7	8	12	15	15	19

Non-Hispanic Black, female

2003.....	595	6	1	4	5	4	5	5	6	22	14	33
2002.....	638	7	6	1	8	3	3	8	15	12	30	30
2001.....	621	6	2	7	6	5	3	3	4	13	18	19
2000.....	608	5	1	7	3	1	6	13	9	11	19	31
1999.....	645	7	3	6	2	4	6	1	9	15	20	31

Origin not stated 2/, both sexes

2003.....	22	1	-	-	-	-	-	-	-	-	1	-
2002.....	28	-	-	-	-	-	1	-	-	-	4	-
2001.....	35	-	-	-	-	-	-	-	-	-	-	-
2000.....	35	1	-	-	1	-	-	-	-	-	1	2
1999.....	31	-	-	1	-	1	-	-	-	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Origin not stated 2/, male

2003.....	14	-	-	-	-	-	-	-	-	-	1	-
2002.....	17	-	-	-	-	-	1	-	-	-	2	-
2001.....	17	-	-	-	-	-	-	-	-	-	-	-
2000.....	20	-	-	-	-	-	-	-	-	-	1	2
1999.....	11	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	8	1	-	-	-	-	-	-	-	-	-	-
2002.....	11	-	-	-	-	-	-	-	-	-	2	-
2001.....	18	-	-	-	-	-	-	-	-	-	-	-
2000.....	15	1	-	-	1	-	-	-	-	-	-	-
1999.....	20	-	-	1	-	1	-	-	-	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Anemias (D50-D64)</b>												
All origins, both sexes												
2003.....	4,594	22	24	16	31	37	68	69	88	81	102	137
2002.....	4,614	18	21	29	24	42	69	83	97	99	97	131
2001.....	4,627	16	22	25	30	48	61	65	95	83	132	112
2000.....	4,494	10	35	24	28	27	58	62	91	85	105	110
1999.....	4,503	12	30	24	29	43	78	64	96	90	104	134
All origins, male												
2003.....	1,832	11	14	8	15	19	33	37	46	37	54	77
2002.....	1,781	12	9	13	13	23	44	53	48	53	43	57
2001.....	1,774	11	12	12	15	26	33	40	55	42	49	57
2000.....	1,669	6	16	17	12	9	30	29	41	34	44	52
1999.....	1,746	8	17	8	15	24	43	33	48	47	54	68
All origins, female												
2003.....	2,762	11	10	8	16	18	35	32	42	44	48	60
2002.....	2,833	6	12	16	11	19	25	30	49	46	54	74
2001.....	2,853	5	10	13	15	22	28	25	40	41	83	55
2000.....	2,825	4	19	7	16	18	28	33	50	51	61	58
1999.....	2,757	4	13	16	14	19	35	31	48	43	50	66
Hispanic, both sexes												
2003.....	191	2	5	2	2	2	3	2	5	3	6	7
2002.....	208	2	3	3	1	3	5	5	6	6	5	8
2001.....	185	-	3	2	3	6	5	4	8	5	7	6
2000.....	144	1	1	2	5	1	4	9	4	4	4	5
1999.....	164	-	6	2	1	2	4	4	4	5	5	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Anemias (D50-D64)</b>												
<b>Hispanic, male</b>												
2003.....	84	2	2	1	1	1	1	1	4	1	2	5
2002.....	106	1	1	2	1	1	4	2	3	5	4	5
2001.....	82	-	2	2	3	2	4	3	5	3	3	2
2000.....	55	1	-	1	2	-	1	4	1	-	1	1
1999.....	79	-	4	1	-	1	2	2	3	3	3	4
<b>Hispanic, female</b>												
2003.....	107	-	3	1	1	1	2	1	1	2	4	2
2002.....	102	1	2	1	-	2	1	3	3	1	1	3
2001.....	103	-	1	-	-	4	1	1	3	2	4	4
2000.....	89	-	1	1	3	1	3	5	3	4	3	4
1999.....	85	-	2	1	1	1	2	2	1	2	2	2
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	4,395	20	19	14	29	35	64	67	83	77	96	130
2002.....	4,391	14	17	26	23	39	64	78	89	91	91	122
2001.....	4,424	15	19	23	27	41	56	61	85	77	125	105
2000.....	4,336	8	34	21	23	26	53	53	87	81	100	105
1999.....	4,321	12	24	22	28	41	74	60	92	85	98	126
<b>Non-Hispanic 1/, male</b>												
2003.....	1,745	9	12	7	14	18	31	36	42	35	52	72
2002.....	1,671	9	8	11	12	22	40	51	45	47	39	52
2001.....	1,687	10	10	10	12	24	29	37	49	38	46	54
2000.....	1,608	4	16	16	10	9	28	25	40	34	43	51
1999.....	1,659	8	13	7	15	23	41	31	45	44	51	63

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Anemias (D50-D64)</b>												
Non-Hispanic 1/, female												
2003.....	2,650	11	7	7	15	17	33	31	41	42	44	58
2002.....	2,720	5	9	15	11	17	24	27	44	44	52	70
2001.....	2,737	5	9	13	15	17	27	24	36	39	79	51
2000.....	2,728	4	18	5	13	17	25	28	47	47	57	54
1999.....	2,662	4	11	15	13	18	33	29	47	41	47	63
Non-Hispanic White, both sexes												
2003.....	3,287	7	5	2	15	13	5	14	18	14	22	40
2002.....	3,253	5	10	12	7	10	13	11	22	16	18	42
2001.....	3,350	5	8	9	13	11	11	11	16	11	30	25
2000.....	3,299	3	14	7	7	9	7	6	10	13	23	33
1999.....	3,222	6	4	5	12	16	12	3	7	17	20	48
Non-Hispanic White, male												
2003.....	1,236	2	3	2	7	5	4	7	10	7	12	24
2002.....	1,164	2	4	4	4	6	7	9	11	10	8	20
2001.....	1,220	4	3	4	7	6	6	6	6	6	9	10
2000.....	1,178	1	8	6	3	2	5	5	5	7	13	15
1999.....	1,162	4	2	-	6	8	3	-	2	7	11	26
Non-Hispanic White, female												
2003.....	2,051	5	2	-	8	8	1	7	8	7	10	16
2002.....	2,089	3	6	8	3	4	6	2	11	6	10	22
2001.....	2,130	1	5	5	6	5	5	5	10	5	21	15
2000.....	2,121	2	6	1	4	7	2	1	5	6	10	18
1999.....	2,060	2	2	5	6	8	9	3	5	10	9	22





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diabetes mellitus (E10-E14)</b>												
All origins, both sexes												
2003.....	74,219	3	8	10	21	46	114	227	430	715	1,334	2,245
2002.....	73,249	1	3	5	29	55	116	229	413	776	1,388	2,106
2001.....	71,372	2	6	7	32	48	103	215	380	735	1,223	2,063
2000.....	69,301	4	6	8	18	48	114	206	417	684	1,242	1,917
1999.....	68,399	2	8	9	23	51	104	189	393	730	1,212	1,942
All origins, male												
2003.....	35,438	-	6	6	10	28	68	119	260	420	844	1,354
2002.....	34,301	-	1	4	18	30	68	128	257	452	878	1,248
2001.....	32,841	1	2	2	21	27	58	114	227	436	745	1,234
2000.....	31,602	4	2	5	8	31	63	118	258	396	755	1,124
1999.....	31,150	-	4	4	12	25	55	94	225	437	702	1,189
All origins, female												
2003.....	38,781	3	2	4	11	18	46	108	170	295	490	891
2002.....	38,948	1	2	1	11	25	48	101	156	324	510	858
2001.....	38,531	1	4	5	11	21	45	101	153	299	478	829
2000.....	37,699	-	4	3	10	17	51	88	159	288	487	793
1999.....	37,249	2	4	5	11	26	49	95	168	293	510	753
Hispanic, both sexes												
2003.....	6,179	-	1	1	2	6	13	28	36	59	138	219
2002.....	5,912	-	-	-	1	5	10	32	24	73	132	191
2001.....	5,663	1	1	-	2	8	11	20	29	69	98	229
2000.....	5,328	1	-	1	3	5	8	21	43	49	105	174
1999.....	5,182	-	-	-	4	2	12	14	29	56	101	169

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diabetes mellitus (E10-E14)</b>												
<b>Hispanic, male</b>												
2003.....	2,976	-	1	-	1	4	7	14	19	37	84	127
2002.....	2,779	-	-	-	1	3	7	19	15	41	83	107
2001.....	2,590	-	1	-	2	5	6	9	19	40	69	139
2000.....	2,507	1	-	-	2	4	3	16	32	28	67	98
1999.....	2,335	-	-	-	2	-	8	8	13	31	53	105
<b>Hispanic, female</b>												
2003.....	3,203	-	-	1	1	2	6	14	17	22	54	92
2002.....	3,133	-	-	-	-	2	3	13	9	32	49	84
2001.....	3,073	1	-	-	-	3	5	11	10	29	29	90
2000.....	2,821	-	-	1	1	1	5	5	11	21	38	76
1999.....	2,847	-	-	-	2	2	4	6	16	25	48	64
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	67,862	3	7	9	18	39	99	199	393	655	1,192	2,020
2002.....	67,112	1	3	5	28	50	105	196	388	697	1,249	1,902
2001.....	65,498	1	5	7	30	40	91	193	349	665	1,122	1,825
2000.....	63,769	3	6	7	15	43	106	185	372	631	1,131	1,741
1999.....	63,017	2	8	9	19	48	92	173	361	672	1,109	1,769
<b>Non-Hispanic 1/, male</b>												
2003.....	32,368	-	5	6	9	23	59	105	240	382	757	1,222
2002.....	31,399	-	1	4	17	27	60	108	241	407	789	1,132
2001.....	30,144	1	1	2	19	22	52	105	206	395	674	1,088
2000.....	28,994	3	2	5	6	27	60	102	226	364	684	1,025
1999.....	28,719	-	4	4	10	25	47	85	210	406	647	1,082

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diabetes mellitus (E10-E14)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	35,494	3	2	3	9	16	40	94	153	273	435	798
2002.....	35,713	1	2	1	11	23	45	88	147	290	460	770
2001.....	35,354	-	4	5	11	18	39	88	143	270	448	737
2000.....	34,775	-	4	2	9	16	46	83	146	267	447	716
1999.....	34,298	2	4	5	9	23	45	88	151	266	462	687
<b>Non-Hispanic White, both sexes</b>												
2003.....	52,894	2	5	8	11	22	68	107	259	428	812	1,368
2002.....	52,463	1	3	4	15	32	63	120	262	474	871	1,289
2001.....	51,482	-	5	3	16	20	46	105	230	442	791	1,250
2000.....	50,163	2	6	4	9	28	64	107	250	441	819	1,175
1999.....	49,347	2	6	8	11	36	58	122	241	462	769	1,197
<b>Non-Hispanic White, male</b>												
2003.....	25,940	-	4	5	7	13	41	49	163	258	536	868
2002.....	25,273	-	1	4	12	16	35	69	162	284	565	792
2001.....	24,302	-	1	1	12	14	28	54	130	262	505	761
2000.....	23,465	2	2	3	4	18	39	60	160	253	513	715
1999.....	23,177	-	3	4	5	21	31	63	140	289	458	760
<b>Non-Hispanic White, female</b>												
2003.....	26,954	2	1	3	4	9	27	58	96	170	276	500
2002.....	27,190	1	2	-	3	16	28	51	100	190	306	497
2001.....	27,180	-	4	2	4	6	18	51	100	180	286	489
2000.....	26,698	-	4	1	5	10	25	47	90	188	306	460
1999.....	26,170	2	3	4	6	15	27	59	101	173	311	437

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diabetes mellitus (E10-E14)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	12,787	1	2	1	6	16	29	85	123	208	339	578
2002.....	12,583	-	-	1	13	16	41	73	121	208	346	543
2001.....	12,167	1	-	4	11	17	39	83	115	202	304	507
2000.....	11,918	1	-	3	6	14	41	75	112	176	282	513
1999.....	11,830	-	2	1	8	12	31	48	112	192	310	513
<b>Non-Hispanic Black, male</b>												
2003.....	5,380	-	1	1	2	10	16	53	69	113	196	310
2002.....	5,158	-	-	-	5	9	25	37	75	114	202	299
2001.....	4,983	1	-	1	5	7	20	49	73	119	156	285
2000.....	4,720	1	-	2	2	8	20	40	58	99	153	277
1999.....	4,718	-	1	-	5	4	15	21	64	107	169	290
<b>Non-Hispanic Black, female</b>												
2003.....	7,407	1	1	-	4	6	13	32	54	95	143	268
2002.....	7,425	-	-	1	8	7	16	36	46	94	144	244
2001.....	7,184	-	-	3	6	10	19	34	42	83	148	222
2000.....	7,198	-	-	1	4	6	21	35	54	77	129	236
1999.....	7,112	-	1	1	3	8	16	27	48	85	141	223
<b>Origin not stated 2/, both sexes</b>												
2003.....	178	-	-	-	1	1	2	-	1	1	4	6
2002.....	225	-	-	-	-	-	1	1	1	6	7	13
2001.....	211	-	-	-	-	-	1	2	2	1	3	9
2000.....	204	-	-	-	-	-	-	-	2	4	6	2
1999.....	200	-	-	-	-	1	-	2	3	2	2	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

Origin not stated 2/, male

2003.....	94	-	-	-	-	1	2	-	1	1	3	5
2002.....	123	-	-	-	-	-	1	1	1	4	6	9
2001.....	107	-	-	-	-	-	-	-	2	1	2	7
2000.....	101	-	-	-	-	-	-	-	-	4	4	1
1999.....	96	-	-	-	-	-	-	1	2	-	2	2

Origin not stated 2/, female

2003.....	84	-	-	-	1	-	-	-	-	-	1	1
2002.....	102	-	-	-	-	-	-	-	-	2	1	4
2001.....	104	-	-	-	-	-	1	2	-	-	1	2
2000.....	103	-	-	-	-	-	-	-	2	-	2	1
1999.....	104	-	-	-	-	1	-	1	1	2	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Nutritional deficiencies (E40-E64)</b>												
All origins, both sexes												
2003.....	3,338	10	2	2	-	3	4	5	5	14	21	39
2002.....	3,779	4	1	-	1	1	6	5	11	27	33	39
2001.....	3,704	4	4	-	2	1	9	5	9	16	28	48
2000.....	4,242	14	2	1	4	4	6	9	9	30	26	46
1999.....	4,289	9	2	1	-	2	6	4	8	20	44	45
All origins, male												
2003.....	1,176	7	2	2	-	1	3	4	3	7	12	19
2002.....	1,321	3	1	-	-	-	4	3	4	12	19	20
2001.....	1,233	4	1	-	2	-	7	2	2	8	13	27
2000.....	1,412	10	2	-	2	1	1	5	4	9	7	25
1999.....	1,486	4	2	1	-	2	3	1	3	8	20	22
All origins, female												
2003.....	2,162	3	-	-	-	2	1	1	2	7	9	20
2002.....	2,458	1	-	-	1	1	2	2	7	15	14	19
2001.....	2,471	-	3	-	-	1	2	3	7	8	15	21
2000.....	2,830	4	-	1	2	3	5	4	5	21	19	21
1999.....	2,803	5	-	-	-	-	3	3	5	12	24	23
Hispanic, both sexes												
2003.....	126	2	2	-	-	-	-	-	1	1	-	2
2002.....	141	2	-	-	-	-	3	1	-	1	2	2
2001.....	136	-	1	-	1	-	-	-	-	4	3	1
2000.....	147	-	-	-	-	-	1	2	1	4	-	2
1999.....	138	-	-	-	-	-	-	-	-	1	3	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Nutritional deficiencies (E40-E64)</b>												
<b>Hispanic, male</b>												
2003.....	54	2	2	-	-	-	-	-	1	-	-	1
2002.....	61	1	-	-	-	-	1	1	-	1	2	2
2001.....	53	-	-	-	1	-	-	-	-	1	1	-
2000.....	57	-	-	-	-	-	-	2	-	1	-	1
1999.....	54	-	-	-	-	-	-	-	-	1	1	1
<b>Hispanic, female</b>												
2003.....	72	-	-	-	-	-	-	-	-	1	-	1
2002.....	80	1	-	-	-	-	2	-	-	-	-	-
2001.....	83	-	1	-	-	-	-	-	-	3	2	1
2000.....	90	-	-	-	-	-	1	-	1	3	-	1
1999.....	84	-	-	-	-	-	-	-	-	-	2	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	3,204	8	-	2	-	3	4	5	4	13	21	37
2002.....	3,621	2	1	-	1	1	3	4	11	25	31	37
2001.....	3,563	4	3	-	1	1	8	5	9	12	25	47
2000.....	4,080	14	2	1	4	4	5	7	8	26	26	44
1999.....	4,142	9	2	1	-	2	6	4	8	19	41	44
<b>Non-Hispanic 1/, male</b>												
2003.....	1,118	5	-	2	-	1	3	4	2	7	12	18
2002.....	1,255	2	1	-	-	-	3	2	4	11	17	18
2001.....	1,178	4	1	-	1	-	6	2	2	7	12	27
2000.....	1,348	10	2	-	2	1	1	3	4	8	7	24
1999.....	1,428	4	2	1	-	2	3	1	3	7	19	21

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

#### Nutritional deficiencies (E40-E64)

##### Non-Hispanic 1/, female

2003.....	2,086	3	-	-	-	2	1	1	2	6	9	19
2002.....	2,366	-	-	-	1	1	-	2	7	14	14	19
2001.....	2,385	-	2	-	-	1	2	3	7	5	13	20
2000.....	2,732	4	-	1	2	3	4	4	4	18	19	20
1999.....	2,714	5	-	-	-	-	3	3	5	12	22	23

##### Non-Hispanic White, both sexes

2003.....	2,699	4	-	2	-	1	4	4	3	10	18	25
2002.....	3,064	1	1	-	-	1	2	4	8	13	24	30
2001.....	3,015	3	3	-	-	1	6	5	5	7	15	33
2000.....	3,469	8	1	-	2	4	3	3	3	17	19	35
1999.....	3,486	5	1	-	-	2	3	3	6	14	31	27

##### Non-Hispanic White, male

2003.....	915	3	-	2	-	-	3	4	1	6	10	11
2002.....	1,039	1	1	-	-	-	2	2	3	6	13	12
2001.....	948	3	1	-	-	-	6	2	1	4	6	14
2000.....	1,112	6	1	-	2	1	1	1	1	6	4	19
1999.....	1,142	2	1	-	-	2	1	-	3	4	14	15

##### Non-Hispanic White, female

2003.....	1,784	1	-	-	-	1	1	-	2	4	8	14
2002.....	2,025	-	-	-	-	1	-	2	5	7	11	18
2001.....	2,067	-	2	-	-	1	-	3	4	3	9	19
2000.....	2,357	2	-	-	-	3	2	2	2	11	15	16
1999.....	2,344	3	-	-	-	-	2	3	3	10	17	12





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malnutrition (E40-E46)</b>												
<b>All origins, both sexes</b>												
2003.....	3,153	5	1	1	-	2	4	4	4	12	21	36
2002.....	3,510	2	1	-	1	1	5	4	9	25	30	36
2001.....	3,454	2	-	-	2	1	8	4	9	16	23	42
2000.....	3,926	11	1	1	2	3	5	9	7	29	24	44
1999.....	3,985	7	2	1	-	2	5	4	7	20	41	42
<b>All origins, male</b>												
2003.....	1,098	4	1	1	-	-	3	3	3	7	12	19
2002.....	1,233	1	1	-	-	-	4	2	3	11	17	18
2001.....	1,148	2	-	-	2	-	6	2	2	8	11	22
2000.....	1,302	8	1	-	1	1	1	5	3	9	6	23
1999.....	1,371	4	2	1	-	2	2	1	3	8	19	21
<b>All origins, female</b>												
2003.....	2,055	1	-	-	-	2	1	1	1	5	9	17
2002.....	2,277	1	-	-	1	1	1	2	6	14	13	18
2001.....	2,306	-	-	-	-	1	2	2	7	8	12	20
2000.....	2,624	3	-	1	1	2	4	4	4	20	18	21
1999.....	2,614	3	-	-	-	-	3	3	4	12	22	21
<b>Hispanic, both sexes</b>												
2003.....	112	1	1	-	-	-	-	-	1	-	-	2
2002.....	136	1	-	-	-	-	2	1	-	1	2	1
2001.....	132	-	-	-	1	-	-	-	-	4	2	1
2000.....	137	-	-	-	-	-	1	2	-	4	-	2
1999.....	132	-	-	-	-	-	-	-	-	1	3	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malnutrition (E40-E46)</b>												
<b>Hispanic, male</b>												
2003.....	47	1	1	-	-	-	-	-	1	-	-	1
2002.....	58	-	-	-	-	-	1	1	-	1	2	1
2001.....	52	-	-	-	1	-	-	-	-	1	1	-
2000.....	52	-	-	-	-	-	-	2	-	1	-	1
1999.....	50	-	-	-	-	-	-	-	-	1	1	1
<b>Hispanic, female</b>												
2003.....	65	-	-	-	-	-	-	-	-	-	-	1
2002.....	78	1	-	-	-	-	1	-	-	-	-	-
2001.....	80	-	-	-	-	-	-	-	-	3	1	1
2000.....	85	-	-	-	-	-	1	-	-	3	-	1
1999.....	82	-	-	-	-	-	-	-	-	-	2	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	3,034	4	-	1	-	2	4	4	3	12	21	34
2002.....	3,358	1	1	-	1	1	3	3	9	23	28	35
2001.....	3,317	2	-	-	1	1	7	4	9	12	21	41
2000.....	3,774	11	1	1	2	3	4	7	7	25	24	42
1999.....	3,844	7	2	1	-	2	5	4	7	19	38	41
<b>Non-Hispanic 1/, male</b>												
2003.....	1,047	3	-	1	-	-	3	3	2	7	12	18
2002.....	1,171	1	1	-	-	-	3	1	3	10	15	17
2001.....	1,094	2	-	-	1	-	5	2	2	7	10	22
2000.....	1,243	8	1	-	1	1	1	3	3	8	6	22
1999.....	1,317	4	2	1	-	2	2	1	3	7	18	20

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Malnutrition (E40-E46)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	1,987	1	-	-	-	2	1	1	1	5	9	16
2002.....	2,187	-	-	-	1	1	-	2	6	13	13	18
2001.....	2,223	-	-	-	-	1	2	2	7	5	11	19
2000.....	2,531	3	-	1	1	2	3	4	4	17	18	20
1999.....	2,527	3	-	-	-	-	3	3	4	12	20	21
<b>Non-Hispanic White, both sexes</b>												
2003.....	2,545	1	-	1	-	1	4	3	2	10	18	23
2002.....	2,836	1	1	-	-	1	2	3	6	13	21	28
2001.....	2,806	1	-	-	-	1	5	4	5	7	12	29
2000.....	3,204	7	-	-	1	3	3	3	3	16	17	33
1999.....	3,232	4	1	-	-	2	3	3	5	14	29	26
<b>Non-Hispanic White, male</b>												
2003.....	852	1	-	1	-	-	3	3	1	6	10	11
2002.....	970	1	1	-	-	-	2	1	2	6	11	11
2001.....	881	1	-	-	-	-	5	2	1	4	4	11
2000.....	1,021	5	-	-	1	1	1	1	1	6	3	17
1999.....	1,047	2	1	-	-	2	1	-	3	4	13	14
<b>Non-Hispanic White, female</b>												
2003.....	1,693	-	-	-	-	1	1	-	1	4	8	12
2002.....	1,866	-	-	-	-	1	-	2	4	7	10	17
2001.....	1,925	-	-	-	-	1	-	2	4	3	8	18
2000.....	2,183	2	-	-	-	2	2	2	2	10	14	16
1999.....	2,185	2	-	-	-	-	2	3	2	10	16	12







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other nutritional deficiencies (E50-E64)

Hispanic, male

2003.....	7	1	1	-	-	-	-	-	-	-	-	-
2002.....	3	1	-	-	-	-	-	-	-	-	-	1
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	5	-	-	-	-	-	-	-	-	-	-	-
1999.....	4	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	7	-	-	-	-	-	-	-	-	1	-	-
2002.....	2	-	-	-	-	-	1	-	-	-	-	-
2001.....	3	-	1	-	-	-	-	-	-	-	1	-
2000.....	5	-	-	-	-	-	-	-	1	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	170	4	-	1	-	1	-	1	1	1	-	3
2002.....	263	1	-	-	-	-	-	1	2	2	3	2
2001.....	246	2	3	-	-	-	1	1	-	-	4	6
2000.....	306	3	1	-	2	1	1	-	1	1	2	2
1999.....	298	2	-	-	-	-	1	-	1	-	3	3

Non-Hispanic 1/, male

2003.....	71	2	-	1	-	1	-	1	-	-	-	-
2002.....	84	1	-	-	-	-	-	1	1	1	2	1
2001.....	84	2	1	-	-	-	1	-	-	-	2	5
2000.....	105	2	1	-	1	-	-	-	1	-	1	2
1999.....	111	-	-	-	-	-	1	-	-	-	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other nutritional deficiencies (E50-E64)												
Non-Hispanic 1/, female												
2003.....	99	2	-	-	-	-	-	-	1	1	-	3
2002.....	179	-	-	-	-	-	-	-	1	1	1	1
2001.....	162	-	2	-	-	-	-	1	-	-	2	1
2000.....	201	1	-	-	1	1	1	-	-	1	1	-
1999.....	187	2	-	-	-	-	-	-	1	-	2	2
Non-Hispanic White, both sexes												
2003.....	154	3	-	1	-	-	-	1	1	-	-	2
2002.....	228	-	-	-	-	-	-	1	2	-	3	2
2001.....	209	2	3	-	-	-	1	1	-	-	3	4
2000.....	265	1	1	-	1	1	-	-	-	1	2	2
1999.....	254	1	-	-	-	-	-	-	1	-	2	1
Non-Hispanic White, male												
2003.....	63	2	-	1	-	-	-	1	-	-	-	-
2002.....	69	-	-	-	-	-	-	1	1	-	2	1
2001.....	67	2	1	-	-	-	1	-	-	-	2	3
2000.....	91	1	1	-	1	-	-	-	-	-	1	2
1999.....	95	-	-	-	-	-	-	-	-	-	1	1
Non-Hispanic White, female												
2003.....	91	1	-	-	-	-	-	-	1	-	-	2
2002.....	159	-	-	-	-	-	-	-	1	-	1	1
2001.....	142	-	2	-	-	-	-	1	-	-	1	1
2000.....	174	-	-	-	-	1	-	-	-	1	1	-
1999.....	159	1	-	-	-	-	-	-	1	-	1	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningitis (G00,G03)</b>												
All origins, both sexes												
2003.....	730	77	24	10	7	14	23	18	26	31	36	59
2002.....	700	74	19	10	21	12	14	16	20	23	38	63
2001.....	727	78	26	15	9	19	21	13	28	31	38	55
2000.....	758	91	36	12	18	7	8	16	27	35	37	45
1999.....	850	116	45	14	13	19	22	21	27	33	43	62
All origins, male												
2003.....	387	38	12	7	3	9	11	10	18	21	19	35
2002.....	389	41	13	5	11	8	7	9	10	17	26	41
2001.....	394	39	16	9	6	12	11	11	14	23	19	30
2000.....	405	48	20	6	10	2	2	7	13	22	24	25
1999.....	442	61	26	9	9	10	16	8	16	18	27	39
All origins, female												
2003.....	343	39	12	3	4	5	12	8	8	10	17	24
2002.....	311	33	6	5	10	4	7	7	10	6	12	22
2001.....	333	39	10	6	3	7	10	2	14	8	19	25
2000.....	353	43	16	6	8	5	6	9	14	13	13	20
1999.....	408	55	19	5	4	9	6	13	11	15	16	23
Hispanic, both sexes												
2003.....	82	18	4	1	2	2	5	3	8	3	4	6
2002.....	70	8	5	-	-	3	3	4	1	5	4	5
2001.....	58	8	4	2	1	-	3	-	4	4	3	5
2000.....	52	9	4	2	1	-	3	-	5	5	3	1
1999.....	76	17	7	1	-	3	2	2	4	2	2	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningitis (G00,G03)</b>												
<b>Hispanic, male</b>												
2003.....	46	12	2	1	2	1	4	-	6	1	3	3
2002.....	42	6	4	-	-	3	2	3	-	2	3	3
2001.....	33	5	3	1	1	-	1	-	1	4	2	3
2000.....	32	6	2	1	-	-	2	-	3	3	2	1
1999.....	41	13	5	1	-	-	1	1	2	1	2	4
<b>Hispanic, female</b>												
2003.....	36	6	2	-	-	1	1	3	2	2	1	3
2002.....	28	2	1	-	-	-	1	1	1	3	1	2
2001.....	25	3	1	1	-	-	2	-	3	-	1	2
2000.....	20	3	2	1	1	-	1	-	2	2	1	-
1999.....	35	4	2	-	-	3	1	1	2	1	-	4
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	646	59	20	9	5	12	18	15	18	28	32	53
2002.....	625	66	14	10	21	9	11	12	19	17	34	57
2001.....	666	70	22	13	8	19	18	13	24	27	35	49
2000.....	705	82	32	10	17	7	5	16	21	30	34	44
1999.....	767	97	38	12	13	16	18	19	23	31	40	54
<b>Non-Hispanic 1/, male</b>												
2003.....	339	26	10	6	1	8	7	10	12	20	16	32
2002.....	344	35	9	5	11	5	5	6	10	14	23	38
2001.....	361	34	13	8	5	12	10	11	13	19	17	27
2000.....	372	42	18	5	10	2	-	7	9	19	22	24
1999.....	396	48	21	7	9	10	13	7	14	17	24	35

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningitis (G00,G03)</b>												
Non-Hispanic 1/, female												
2003.....	307	33	10	3	4	4	11	5	6	8	16	21
2002.....	281	31	5	5	10	4	6	6	9	3	11	19
2001.....	305	36	9	5	3	7	8	2	11	8	18	22
2000.....	333	40	14	5	7	5	5	9	12	11	12	20
1999.....	371	49	17	5	4	6	5	12	9	14	16	19
Non-Hispanic White, both sexes												
2003.....	444	35	14	7	3	8	10	7	8	14	22	36
2002.....	427	44	10	9	11	7	5	5	10	13	17	33
2001.....	469	39	15	8	7	14	10	8	17	11	22	32
2000.....	486	47	19	9	9	4	3	8	12	19	17	34
1999.....	524	51	26	8	10	7	8	14	14	20	25	31
Non-Hispanic White, male												
2003.....	247	21	7	5	1	5	6	6	5	11	11	24
2002.....	227	23	6	5	6	4	1	2	4	11	9	21
2001.....	243	17	10	4	4	7	5	7	10	6	12	17
2000.....	264	27	12	5	4	1	-	3	4	12	10	20
1999.....	265	33	15	4	7	5	5	3	7	11	13	18
Non-Hispanic White, female												
2003.....	197	14	7	2	2	3	4	1	3	3	11	12
2002.....	200	21	4	4	5	3	4	3	6	2	8	12
2001.....	226	22	5	4	3	7	5	1	7	5	10	15
2000.....	222	20	7	4	5	3	3	5	8	7	7	14
1999.....	259	18	11	4	3	2	3	11	7	9	12	13

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Meningitis (G00,G03)</b>												
Non-Hispanic Black, both sexes												
2003.....	183	22	5	2	1	4	8	8	10	13	9	15
2002.....	175	20	4	1	9	1	5	7	8	4	15	21
2001.....	172	29	5	4	-	4	6	5	6	13	13	17
2000.....	202	34	13	1	8	3	2	8	9	10	17	9
1999.....	217	40	11	4	3	9	8	5	6	10	11	22
Non-Hispanic Black, male												
2003.....	82	5	2	1	-	3	1	4	7	8	4	7
2002.....	105	11	3	-	5	1	4	4	5	3	12	15
2001.....	104	16	2	3	-	4	4	4	2	12	5	10
2000.....	100	15	6	-	6	1	-	4	5	6	12	3
1999.....	119	13	6	3	2	5	6	4	5	6	8	17
Non-Hispanic Black, female												
2003.....	101	17	3	1	1	1	7	4	3	5	5	8
2002.....	70	9	1	1	4	-	1	3	3	1	3	6
2001.....	68	13	3	1	-	-	2	1	4	1	8	7
2000.....	102	19	7	1	2	2	2	4	4	4	5	6
1999.....	98	27	5	1	1	4	2	1	1	4	3	5
Origin not stated 2/, both sexes												
2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	5	-	-	-	-	-	-	-	-	1	-	1
2001.....	3	-	-	-	-	-	-	-	-	-	-	1
2000.....	1	-	-	-	-	-	-	-	1	-	-	-
1999.....	7	2	-	1	-	-	2	-	-	-	1	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Parkinson's disease (G20-G21)

Hispanic, male

2003.....	313	-	-	-	-	-	-	-	-	1	-	-
2002.....	281	-	-	-	-	-	-	-	-	-	1	3
2001.....	274	-	-	-	-	-	-	1	-	-	-	1
2000.....	240	-	-	-	-	-	-	-	-	-	-	1
1999.....	222	-	-	-	-	-	-	-	-	-	1	-

Hispanic, female

2003.....	208	-	-	-	-	-	-	-	-	-	-	-
2002.....	210	-	-	-	-	-	-	-	-	-	-	1
2001.....	209	-	-	-	-	-	-	-	-	-	-	-
2000.....	185	-	-	-	-	-	-	-	-	-	-	2
1999.....	167	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	17,443	-	-	1	-	-	-	-	2	2	8	18
2002.....	16,435	-	-	-	-	-	-	-	-	1	4	15
2001.....	16,019	-	-	-	-	-	-	-	1	3	12	12
2000.....	15,223	-	1	-	-	-	-	2	2	4	3	12
1999.....	14,177	1	-	-	-	-	2	1	1	1	3	8

Non-Hispanic 1/, male

2003.....	9,846	-	-	-	-	-	-	-	-	1	4	10
2002.....	9,290	-	-	-	-	-	-	-	-	-	2	11
2001.....	9,104	-	-	-	-	-	-	-	-	1	10	7
2000.....	8,544	-	-	-	-	-	-	1	2	3	2	8
1999.....	8,029	-	-	-	-	-	2	1	1	1	1	5







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Alzheimer's disease (G30)</b>												
All origins, both sexes												
2003.....	63,457	-	-	-	-	-	-	-	-	2	9	21
2002.....	58,866	-	-	-	1	-	-	-	2	3	9	12
2001.....	53,852	-	-	-	-	-	-	-	2	1	2	17
2000.....	49,558	-	-	-	-	-	-	1	1	2	4	17
1999.....	44,536	-	-	-	-	-	1	-	-	1	6	20
All origins, male												
2003.....	18,335	-	-	-	-	-	-	-	-	1	3	9
2002.....	16,989	-	-	-	-	-	-	-	2	3	5	5
2001.....	15,762	-	-	-	-	-	-	-	2	1	2	11
2000.....	14,438	-	-	-	-	-	-	1	-	1	3	10
1999.....	13,391	-	-	-	-	-	1	-	-	-	3	10
All origins, female												
2003.....	45,122	-	-	-	-	-	-	-	-	1	6	12
2002.....	41,877	-	-	-	1	-	-	-	-	-	4	7
2001.....	38,090	-	-	-	-	-	-	-	-	-	-	6
2000.....	35,120	-	-	-	-	-	-	-	1	1	1	7
1999.....	31,145	-	-	-	-	-	-	-	-	1	3	10
Hispanic, both sexes												
2003.....	1,835	-	-	-	-	-	-	-	-	-	1	2
2002.....	1,474	-	-	-	-	-	-	-	-	-	1	-
2001.....	1,239	-	-	-	-	-	-	-	-	-	1	2
2000.....	1,127	-	-	-	-	-	-	-	-	-	-	2
1999.....	981	-	-	-	-	-	1	-	-	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Alzheimer's disease (G30)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	43,840	-	-	-	-	-	-	-	-	1	5	12
2002.....	40,755	-	-	-	1	-	-	-	-	-	4	7
2001.....	37,182	-	-	-	-	-	-	-	-	-	-	6
2000.....	34,297	-	-	-	-	-	-	-	1	1	1	6
1999.....	30,419	-	-	-	-	-	-	-	-	1	3	10
<b>Non-Hispanic White, both sexes</b>												
2003.....	57,268	-	-	-	-	-	-	-	-	2	8	17
2002.....	53,486	-	-	-	1	-	-	-	1	1	5	12
2001.....	49,030	-	-	-	-	-	-	-	2	1	1	11
2000.....	45,246	-	-	-	-	-	-	1	1	2	4	13
1999.....	40,835	-	-	-	-	-	-	-	-	1	4	18
<b>Non-Hispanic White, male</b>												
2003.....	16,427	-	-	-	-	-	-	-	-	1	3	6
2002.....	15,399	-	-	-	-	-	-	-	1	1	3	5
2001.....	14,295	-	-	-	-	-	-	-	2	1	1	7
2000.....	13,123	-	-	-	-	-	-	1	-	1	3	7
1999.....	12,253	-	-	-	-	-	-	-	-	-	2	10
<b>Non-Hispanic White, female</b>												
2003.....	40,841	-	-	-	-	-	-	-	-	1	5	11
2002.....	38,087	-	-	-	1	-	-	-	-	-	2	7
2001.....	34,735	-	-	-	-	-	-	-	-	-	-	4
2000.....	32,123	-	-	-	-	-	-	-	1	1	1	6
1999.....	28,582	-	-	-	-	-	-	-	-	1	2	8





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Major cardiovascular diseases (I00-I78)</b>												
<b>All origins, both sexes</b>												
2003.....	902,443	555	234	134	213	519	928	1,352	2,737	5,375	11,555	19,091
2002.....	918,628	640	223	127	231	493	790	1,348	2,638	5,541	11,421	18,716
2001.....	922,334	600	285	139	226	443	837	1,351	2,687	5,568	11,118	17,768
2000.....	936,923	636	234	136	226	493	816	1,295	2,497	5,491	11,133	17,785
1999.....	950,314	644	234	144	210	548	767	1,270	2,603	5,918	11,092	17,354
<b>All origins, male</b>												
2003.....	423,810	310	116	75	129	323	570	830	1,779	3,585	7,717	13,215
2002.....	429,682	343	122	57	137	328	517	869	1,699	3,602	7,754	12,998
2001.....	428,065	315	149	65	119	278	529	840	1,715	3,672	7,508	12,418
2000.....	435,912	341	128	78	128	295	488	800	1,578	3,538	7,572	12,544
1999.....	441,827	347	127	84	115	345	454	769	1,673	3,898	7,573	12,150
<b>All origins, female</b>												
2003.....	478,633	245	118	59	84	196	358	522	958	1,790	3,838	5,876
2002.....	488,946	297	101	70	94	165	273	479	939	1,939	3,667	5,718
2001.....	494,269	285	136	74	107	165	308	511	972	1,896	3,610	5,350
2000.....	501,011	295	106	58	98	198	328	495	919	1,953	3,561	5,241
1999.....	508,487	297	107	60	95	203	313	501	930	2,020	3,519	5,204
<b>Hispanic, both sexes</b>												
2003.....	37,266	95	43	14	32	62	114	183	287	428	783	1,185
2002.....	36,543	109	36	17	34	57	119	176	269	435	749	1,169
2001.....	35,565	98	44	25	31	65	97	157	285	420	661	1,037
2000.....	34,059	106	37	21	26	60	109	162	257	410	684	995
1999.....	33,665	115	41	22	25	67	93	149	259	426	631	994

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Major cardiovascular diseases (I00-I78)</b>												
<b>Hispanic, male</b>												
2003.....	19,066	54	26	8	19	41	82	119	196	308	556	839
2002.....	18,824	52	16	7	14	36	86	134	188	289	540	825
2001.....	18,138	53	21	14	15	40	68	109	199	296	462	748
2000.....	17,442	54	21	10	20	41	69	109	174	268	471	711
1999.....	17,304	49	21	14	15	41	62	107	185	296	459	696
<b>Hispanic, female</b>												
2003.....	18,200	41	17	6	13	21	32	64	91	120	227	346
2002.....	17,719	57	20	10	20	21	33	42	81	146	209	344
2001.....	17,427	45	23	11	16	25	29	48	86	124	199	289
2000.....	16,617	52	16	11	6	19	40	53	83	142	213	284
1999.....	16,361	66	20	8	10	26	31	42	74	130	172	298
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	862,744	456	190	119	181	453	812	1,163	2,439	4,927	10,720	17,827
2002.....	879,170	528	187	110	197	436	669	1,166	2,356	5,081	10,626	17,480
2001.....	883,949	499	239	113	195	375	733	1,187	2,391	5,128	10,408	16,651
2000.....	899,801	526	196	114	199	433	704	1,126	2,231	5,055	10,378	16,695
1999.....	913,476	528	192	122	184	479	673	1,116	2,337	5,465	10,404	16,278
<b>Non-Hispanic 1/, male</b>												
2003.....	403,411	255	90	66	110	280	486	709	1,574	3,262	7,125	12,315
2002.....	409,343	289	106	50	123	292	429	733	1,501	3,295	7,182	12,126
2001.....	408,437	260	128	51	104	235	457	725	1,506	3,362	7,004	11,613
2000.....	416,837	284	107	68	108	254	418	687	1,401	3,250	7,044	11,763
1999.....	422,824	298	106	70	99	303	391	659	1,483	3,581	7,068	11,392

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Major cardiovascular diseases (I00-I78)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	459,333	201	100	53	71	173	326	454	865	1,665	3,595	5,512
2002.....	469,827	239	81	60	74	144	240	433	855	1,786	3,444	5,354
2001.....	475,512	239	111	62	91	140	276	462	885	1,766	3,404	5,038
2000.....	482,964	242	89	46	91	179	286	439	830	1,805	3,334	4,932
1999.....	490,652	230	86	52	85	176	282	457	854	1,884	3,336	4,886
<b>Non-Hispanic White, both sexes</b>												
2003.....	741,886	250	107	73	118	293	487	681	1,526	3,245	7,355	12,472
2002.....	758,738	314	125	75	120	289	406	681	1,440	3,374	7,464	12,325
2001.....	764,021	289	140	75	129	241	430	720	1,520	3,432	7,092	11,734
2000.....	780,445	330	126	74	121	272	418	681	1,403	3,392	7,164	11,764
1999.....	793,945	324	121	86	117	318	412	683	1,501	3,684	7,123	11,537
<b>Non-Hispanic White, male</b>												
2003.....	345,832	133	43	39	76	180	296	427	1,019	2,245	5,137	9,022
2002.....	352,520	177	77	35	76	192	270	443	944	2,256	5,303	8,993
2001.....	352,013	160	75	36	72	148	277	454	987	2,322	5,057	8,582
2000.....	360,757	180	75	47	71	161	257	428	933	2,296	5,126	8,783
1999.....	366,310	189	70	50	60	196	234	408	992	2,505	5,101	8,494
<b>Non-Hispanic White, female</b>												
2003.....	396,054	117	64	34	42	113	191	254	507	1,000	2,218	3,450
2002.....	406,218	137	48	40	44	97	136	238	496	1,118	2,161	3,332
2001.....	412,008	129	65	39	57	93	153	266	533	1,110	2,035	3,152
2000.....	419,688	150	51	27	50	111	161	253	470	1,096	2,038	2,981
1999.....	427,635	135	51	36	57	122	178	275	509	1,179	2,022	3,043

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Major cardiovascular diseases (I00-I78)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	103,027	168	70	39	56	138	290	429	802	1,487	3,002	4,796
2002.....	103,054	183	50	26	67	122	234	434	818	1,517	2,846	4,650
2001.....	103,213	184	86	33	52	123	272	429	780	1,500	3,006	4,432
2000.....	103,239	166	60	33	62	143	253	403	757	1,498	2,898	4,481
1999.....	103,629	172	60	35	58	147	241	390	741	1,615	2,995	4,273
<b>Non-Hispanic Black, male</b>												
2003.....	48,237	102	41	23	30	90	172	251	470	876	1,752	2,906
2002.....	47,656	96	24	10	40	79	138	258	481	897	1,654	2,778
2001.....	47,646	93	45	12	27	80	160	251	455	898	1,727	2,683
2000.....	47,408	88	27	17	30	79	139	230	422	846	1,698	2,675
1999.....	47,982	91	32	19	34	98	142	228	431	955	1,761	2,585
<b>Non-Hispanic Black, female</b>												
2003.....	54,790	66	29	16	26	48	118	178	332	611	1,250	1,890
2002.....	55,398	87	26	16	27	43	96	176	337	620	1,192	1,872
2001.....	55,567	91	41	21	25	43	112	178	325	602	1,279	1,749
2000.....	55,831	78	33	16	32	64	114	173	335	652	1,200	1,806
1999.....	55,647	81	28	16	24	49	99	162	310	660	1,234	1,688
<b>Origin not stated 2/, both sexes</b>												
2003.....	2,433	4	1	1	-	4	2	6	11	20	52	79
2002.....	2,915	3	-	-	-	-	2	6	13	25	46	67
2001.....	2,820	3	2	1	-	3	7	7	11	20	49	80
2000.....	3,063	4	1	1	1	-	3	7	9	26	71	95
1999.....	3,173	1	1	-	1	2	1	5	7	27	57	82

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

Origin not stated 2/, male

2003.....	1,333	1	-	1	-	2	2	2	9	15	36	61
2002.....	1,515	2	-	-	-	-	2	2	10	18	32	47
2001.....	1,490	2	-	-	-	3	4	6	10	14	42	57
2000.....	1,633	3	-	-	-	-	1	4	3	20	57	70
1999.....	1,699	-	-	-	1	1	1	3	5	21	46	62

Origin not stated 2/, female

2003.....	1,100	3	1	-	-	2	-	4	2	5	16	18
2002.....	1,400	1	-	-	-	-	-	4	3	7	14	20
2001.....	1,330	1	2	1	-	-	3	1	1	6	7	23
2000.....	1,430	1	1	1	1	-	2	3	6	6	14	25
1999.....	1,474	1	1	-	-	1	-	2	2	6	11	20

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>All origins, both sexes</b>												
2003.....	685,089	439	186	104	160	393	740	1,075	2,175	4,310	9,290	15,520
2002.....	696,947	500	165	92	163	405	617	1,076	2,089	4,448	9,240	15,179
2001.....	700,142	479	225	98	174	347	652	1,052	2,108	4,419	8,907	14,390
2000.....	710,760	493	181	106	165	403	628	1,013	1,945	4,333	8,848	14,313
1999.....	725,192	524	183	116	161	463	606	992	2,074	4,703	8,897	14,096
<b>All origins, male</b>												
2003.....	336,095	247	89	56	99	249	469	677	1,452	2,977	6,486	11,164
2002.....	340,933	267	87	40	101	275	411	716	1,415	3,048	6,590	10,983
2001.....	339,095	243	121	43	95	223	424	672	1,396	3,038	6,308	10,491
2000.....	344,807	260	107	64	99	248	390	640	1,282	2,945	6,341	10,596
1999.....	351,617	270	104	66	91	297	367	618	1,370	3,252	6,388	10,346
<b>All origins, female</b>												
2003.....	348,994	192	97	48	61	144	271	398	723	1,333	2,804	4,356
2002.....	356,014	233	78	52	62	130	206	360	674	1,400	2,650	4,196
2001.....	361,047	236	104	55	79	124	228	380	712	1,381	2,599	3,899
2000.....	365,953	233	74	42	66	155	238	373	663	1,388	2,507	3,717
1999.....	373,575	254	79	50	70	166	239	374	704	1,451	2,509	3,750
<b>Hispanic, both sexes</b>												
2003.....	28,298	75	35	9	22	46	89	134	213	327	552	865
2002.....	27,887	85	24	12	18	39	89	126	195	313	539	851
2001.....	27,090	79	33	17	21	45	67	115	199	302	448	736
2000.....	25,819	86	29	17	21	49	76	112	168	284	476	709
1999.....	25,866	96	33	18	18	52	64	93	193	320	441	716

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>Hispanic, male</b>												
2003.....	14,867	42	19	5	14	30	71	93	151	244	410	637
2002.....	14,798	41	11	6	8	26	68	98	144	215	410	621
2001.....	14,195	43	16	9	10	27	49	84	142	228	337	565
2000.....	13,566	44	21	9	17	36	51	79	120	198	352	539
1999.....	13,554	40	20	11	12	31	42	77	142	235	337	531
<b>Hispanic, female</b>												
2003.....	13,431	33	16	4	8	16	18	41	62	83	142	228
2002.....	13,089	44	13	6	10	13	21	28	51	98	129	230
2001.....	12,895	36	17	8	11	18	18	31	57	74	111	171
2000.....	12,253	42	8	8	4	13	25	33	48	86	124	170
1999.....	12,312	56	13	7	6	21	22	16	51	85	104	185
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	654,828	362	150	94	138	343	650	937	1,954	3,969	8,693	14,586
2002.....	666,690	413	141	80	145	366	526	945	1,882	4,116	8,664	14,272
2001.....	670,787	397	191	81	153	299	578	933	1,901	4,100	8,415	13,588
2000.....	682,470	403	151	88	144	354	550	895	1,769	4,027	8,314	13,530
1999.....	696,719	427	149	98	142	409	541	894	1,876	4,358	8,409	13,314
<b>Non-Hispanic 1/, male</b>												
2003.....	320,095	205	70	50	85	217	397	583	1,295	2,723	6,042	10,473
2002.....	324,848	225	76	34	93	249	341	616	1,262	2,818	6,154	10,321
2001.....	323,673	198	105	34	85	193	371	585	1,247	2,799	5,934	9,876
2000.....	329,888	213	86	55	82	212	338	557	1,159	2,730	5,943	10,002
1999.....	336,615	230	84	55	78	265	324	538	1,224	2,997	6,012	9,762

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	334,733	157	80	44	53	126	253	354	659	1,246	2,651	4,113
2002.....	341,842	188	65	46	52	117	185	329	620	1,298	2,510	3,951
2001.....	347,114	199	86	47	68	106	207	348	654	1,301	2,481	3,712
2000.....	352,582	190	65	33	62	142	212	338	610	1,297	2,371	3,528
1999.....	360,104	197	65	43	64	144	217	356	652	1,361	2,397	3,552
<b>Non-Hispanic White, both sexes</b>												
2003.....	565,808	200	83	56	90	218	387	549	1,239	2,722	6,172	10,546
2002.....	577,761	249	91	53	93	240	307	554	1,164	2,818	6,296	10,407
2001.....	582,349	234	108	52	100	186	336	563	1,212	2,803	5,931	9,888
2000.....	594,465	258	98	54	86	216	315	523	1,130	2,789	5,954	9,905
1999.....	607,704	259	92	65	91	269	316	542	1,213	2,998	5,938	9,762
<b>Non-Hispanic White, male</b>												
2003.....	276,241	110	34	29	59	137	231	353	859	1,943	4,486	7,883
2002.....	281,442	139	54	24	61	163	207	369	808	1,984	4,673	7,866
2001.....	280,713	127	59	23	57	117	219	367	817	1,984	4,411	7,516
2000.....	287,210	139	59	34	53	126	200	336	779	1,979	4,467	7,699
1999.....	293,206	145	55	36	48	167	190	331	821	2,134	4,456	7,501
<b>Non-Hispanic White, female</b>												
2003.....	289,567	90	49	27	31	81	156	196	380	779	1,686	2,663
2002.....	296,319	110	37	29	32	77	100	185	356	834	1,623	2,541
2001.....	301,636	107	49	29	43	69	117	196	395	819	1,520	2,372
2000.....	307,255	119	39	20	33	90	115	187	351	810	1,487	2,206
1999.....	314,498	114	37	29	43	102	126	211	392	864	1,482	2,261

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	76,452	129	55	33	43	108	238	345	629	1,118	2,274	3,642
2002.....	76,694	142	40	19	46	105	194	352	641	1,167	2,145	3,506
2001.....	76,794	140	73	27	43	104	218	339	622	1,162	2,262	3,345
2000.....	76,706	118	47	29	46	124	207	343	586	1,118	2,141	3,320
1999.....	77,713	141	49	32	45	128	208	318	595	1,246	2,260	3,239
<b>Non-Hispanic Black, male</b>												
2003.....	36,962	79	31	18	24	72	153	202	370	682	1,384	2,280
2002.....	36,596	76	18	6	27	68	117	220	392	730	1,301	2,184
2001.....	36,541	65	40	10	24	69	136	199	379	715	1,356	2,087
2000.....	36,282	59	24	17	24	74	118	200	343	667	1,305	2,069
1999.....	37,077	71	27	18	27	90	122	191	355	773	1,392	2,024
<b>Non-Hispanic Black, female</b>												
2003.....	39,490	50	24	15	19	36	85	143	259	436	890	1,362
2002.....	40,098	66	22	13	19	37	77	132	249	437	844	1,322
2001.....	40,253	75	33	17	19	35	82	140	243	447	906	1,258
2000.....	40,424	59	23	12	22	50	89	143	243	451	836	1,251
1999.....	40,636	70	22	14	18	38	86	127	240	473	868	1,215
<b>Origin not stated 2/, both sexes</b>												
2003.....	1,963	2	1	1	-	4	1	4	8	14	45	69
2002.....	2,370	2	-	-	-	-	2	5	12	19	37	56
2001.....	2,265	3	1	-	-	3	7	4	8	17	44	66
2000.....	2,471	4	1	1	-	-	2	6	8	22	58	74
1999.....	2,607	1	1	-	1	2	1	5	5	25	47	66

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Origin not stated 2/, male												
2003.....	1,133	-	-	1	-	2	1	1	6	10	34	54
2002.....	1,287	1	-	-	-	-	2	2	9	15	26	41
2001.....	1,227	2	-	-	-	3	4	3	7	11	37	50
2000.....	1,353	3	-	-	-	-	1	4	3	17	46	55
1999.....	1,448	-	-	-	1	1	1	3	4	20	39	53
Origin not stated 2/, female												
2003.....	830	2	1	-	-	2	-	3	2	4	11	15
2002.....	1,083	1	-	-	-	-	-	3	3	4	11	15
2001.....	1,038	1	1	-	-	-	3	1	1	6	7	16
2000.....	1,118	1	1	1	-	-	1	2	5	5	12	19
1999.....	1,159	1	1	-	-	1	-	2	1	5	8	13

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All origins, both sexes												
2003.....	3,624	1	1	2	4	4	7	11	14	41	58	107
2002.....	3,579	4	1	4	2	8	10	10	26	38	57	97
2001.....	3,489	3	1	1	4	3	8	11	23	45	85	107
2000.....	3,582	6	2	2	2	9	8	10	23	49	92	119
1999.....	3,676	6	1	2	3	5	8	12	30	47	77	101
All origins, male												
2003.....	1,153	1	-	2	2	3	4	8	6	19	25	39
2002.....	1,078	3	1	1	1	3	5	6	12	19	26	40
2001.....	1,031	3	1	-	2	2	5	4	11	18	36	46
2000.....	1,072	5	2	1	2	5	6	2	10	25	36	49
1999.....	1,042	2	1	1	1	4	6	5	12	19	30	40
All origins, female												
2003.....	2,471	-	1	-	2	1	3	3	8	22	33	68
2002.....	2,501	1	-	3	1	5	5	4	14	19	31	57
2001.....	2,458	-	-	1	2	1	3	7	12	27	49	61
2000.....	2,510	1	-	1	-	4	2	8	13	24	56	70
1999.....	2,634	4	-	1	2	1	2	7	18	28	47	61
Hispanic, both sexes												
2003.....	159	1	-	-	1	-	2	4	3	-	7	10
2002.....	198	-	-	-	-	-	2	2	9	6	5	13
2001.....	156	-	1	-	1	-	2	-	2	5	7	9
2000.....	154	1	-	-	-	1	-	1	5	7	6	11
1999.....	166	2	-	-	-	-	3	1	3	9	7	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hispanic, male												
2003.....	57	1	-	-	1	-	2	3	1	-	4	5
2002.....	55	-	-	-	-	-	1	-	5	2	2	4
2001.....	51	-	1	-	1	-	1	-	2	1	4	3
2000.....	49	1	-	-	-	-	-	-	1	2	2	2
1999.....	38	-	-	-	-	-	3	-	3	2	1	4
Hispanic, female												
2003.....	102	-	-	-	-	-	-	1	2	-	3	5
2002.....	143	-	-	-	-	-	1	2	4	4	3	9
2001.....	105	-	-	-	-	-	1	-	-	4	3	6
2000.....	105	-	-	-	-	1	-	1	4	5	4	9
1999.....	128	2	-	-	-	-	-	1	-	7	6	6
Non-Hispanic 1/, both sexes												
2003.....	3,460	-	1	2	3	4	5	7	11	41	51	97
2002.....	3,375	4	1	4	2	8	8	8	17	32	52	84
2001.....	3,327	3	-	1	3	2	6	11	21	40	78	97
2000.....	3,424	5	2	2	2	8	8	9	18	42	86	108
1999.....	3,498	4	1	2	3	5	5	11	27	37	69	91
Non-Hispanic 1/, male												
2003.....	1,095	-	-	2	1	3	2	5	5	19	21	34
2002.....	1,021	3	1	1	1	3	4	6	7	17	24	36
2001.....	978	3	-	-	1	1	4	4	9	17	32	42
2000.....	1,022	4	2	1	2	5	6	2	9	23	34	47
1999.....	999	2	1	1	1	4	3	5	9	16	29	36

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic 1/, female												
2003.....	2,365	-	1	-	2	1	3	2	6	22	30	63
2002.....	2,354	1	-	3	1	5	4	2	10	15	28	48
2001.....	2,349	-	-	1	2	1	2	7	12	23	46	55
2000.....	2,402	1	-	1	-	3	2	7	9	19	52	61
1999.....	2,499	2	-	1	2	1	2	6	18	21	40	55
Non-Hispanic White, both sexes												
2003.....	3,066	-	1	2	1	-	3	5	7	25	30	61
2002.....	3,007	4	1	1	2	3	5	4	10	26	31	57
2001.....	2,924	1	-	1	1	2	5	5	8	23	50	60
2000.....	3,073	4	2	2	2	5	4	6	13	27	56	79
1999.....	3,130	2	1	1	2	2	3	7	14	24	44	69
Non-Hispanic White, male												
2003.....	967	-	-	2	-	-	2	3	3	12	11	24
2002.....	897	3	1	1	1	-	3	3	4	13	13	26
2001.....	857	1	-	-	-	1	4	2	5	11	24	28
2000.....	917	3	2	1	2	3	3	1	8	17	25	37
1999.....	892	1	1	-	-	1	1	4	5	11	20	28
Non-Hispanic White, female												
2003.....	2,099	-	1	-	1	-	1	2	4	13	19	37
2002.....	2,110	1	-	-	1	3	2	1	6	13	18	31
2001.....	2,067	-	-	1	1	1	1	3	3	12	26	32
2000.....	2,156	1	-	1	-	2	1	5	5	10	31	42
1999.....	2,238	1	-	1	2	1	2	3	9	13	24	41

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Non-Hispanic Black, both sexes												
2003.....	305	-	-	-	2	2	2	2	3	13	16	32
2002.....	249	-	-	2	-	3	2	3	6	4	16	21
2001.....	302	2	-	-	1	-	1	3	10	15	26	27
2000.....	263	1	-	-	-	3	4	3	5	11	24	23
1999.....	269	1	-	1	1	2	2	4	9	11	19	20
Non-Hispanic Black, male												
2003.....	109	-	-	-	1	2	-	2	1	6	9	9
2002.....	91	-	-	-	-	2	1	2	2	2	10	7
2001.....	86	2	-	-	-	-	-	-	3	6	7	11
2000.....	77	1	-	-	-	2	3	1	1	5	7	8
1999.....	86	1	-	1	1	2	2	1	2	4	8	8
Non-Hispanic Black, female												
2003.....	196	-	-	-	1	-	2	-	2	7	7	23
2002.....	158	-	-	2	-	1	1	1	4	2	6	14
2001.....	216	-	-	-	1	-	1	3	7	9	19	16
2000.....	186	-	-	-	-	1	1	2	4	6	17	15
1999.....	183	-	-	-	-	-	-	3	7	7	11	12
Origin not stated 2/, both sexes												
2003.....	5	-	-	-	-	-	-	-	-	-	-	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	-
2001.....	6	-	-	-	-	1	-	-	-	-	-	1
2000.....	4	-	-	-	-	-	-	-	-	-	-	-
1999.....	12	-	-	-	-	-	-	-	-	1	1	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart disease (I11)</b>												
<b>All origins, both sexes</b>												
2003.....	28,345	-	1	-	-	9	44	90	215	409	897	1,446
2002.....	26,551	-	-	-	2	4	26	81	216	402	811	1,266
2001.....	24,689	-	-	1	-	6	18	65	181	352	739	1,050
2000.....	23,761	-	1	-	2	5	27	69	144	323	658	952
1999.....	22,702	1	-	-	-	6	27	50	160	347	617	889
<b>All origins, male</b>												
2003.....	12,536	-	1	-	-	6	34	64	144	285	592	962
2002.....	11,660	-	-	-	1	4	22	64	151	289	560	844
2001.....	10,451	-	-	-	-	4	13	40	129	234	506	715
2000.....	9,876	-	-	-	1	3	21	43	96	210	433	625
1999.....	9,316	-	-	-	-	6	19	41	108	232	428	586
<b>All origins, female</b>												
2003.....	15,809	-	-	-	-	3	10	26	71	124	305	484
2002.....	14,891	-	-	-	1	-	4	17	65	113	251	422
2001.....	14,238	-	-	1	-	2	5	25	52	118	233	335
2000.....	13,885	-	1	-	1	2	6	26	48	113	225	327
1999.....	13,386	1	-	-	-	-	8	9	52	115	189	303
<b>Hispanic, both sexes</b>												
2003.....	1,586	-	-	-	-	3	9	10	17	38	71	109
2002.....	1,424	-	-	-	-	-	4	8	23	38	59	83
2001.....	1,211	-	-	-	-	1	4	9	13	42	51	59
2000.....	991	-	-	-	-	-	1	9	11	26	38	55
1999.....	933	-	-	-	-	1	2	5	13	31	30	43

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart disease (I11)</b>												
<b>Hispanic, male</b>												
2003.....	876	-	-	-	-	2	8	8	14	33	58	79
2002.....	781	-	-	-	-	-	4	7	18	28	50	65
2001.....	648	-	-	-	-	1	3	6	9	37	38	43
2000.....	495	-	-	-	-	-	1	9	7	18	25	42
1999.....	455	-	-	-	-	1	1	5	11	24	21	34
<b>Hispanic, female</b>												
2003.....	710	-	-	-	-	1	1	2	3	5	13	30
2002.....	643	-	-	-	-	-	-	1	5	10	9	18
2001.....	563	-	-	-	-	-	1	3	4	5	13	16
2000.....	496	-	-	-	-	-	-	-	4	8	13	13
1999.....	478	-	-	-	-	-	1	-	2	7	9	9
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	26,632	-	1	-	-	6	35	79	198	367	822	1,323
2002.....	24,976	-	-	-	2	4	22	72	193	362	747	1,174
2001.....	23,343	-	-	1	-	5	13	56	166	307	684	982
2000.....	22,646	-	1	-	2	5	26	60	133	295	610	888
1999.....	21,639	1	-	-	-	5	25	45	147	316	582	837
<b>Non-Hispanic 1/, male</b>												
2003.....	11,588	-	1	-	-	4	26	56	130	249	531	872
2002.....	10,790	-	-	-	1	4	18	57	133	259	506	770
2001.....	9,725	-	-	-	-	3	9	34	118	196	464	667
2000.....	9,298	-	-	-	1	3	20	34	89	190	401	576
1999.....	8,790	-	-	-	-	5	18	36	97	208	402	545

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart disease (I11)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	15,044	-	-	-	-	2	9	23	68	118	291	451
2002.....	14,186	-	-	-	1	-	4	15	60	103	241	404
2001.....	13,618	-	-	1	-	2	4	22	48	111	220	315
2000.....	13,348	-	1	-	1	2	6	26	44	105	209	312
1999.....	12,849	1	-	-	-	-	7	9	50	108	180	292
<b>Non-Hispanic White, both sexes</b>												
2003.....	19,286	-	-	-	-	1	14	39	94	191	432	673
2002.....	18,060	-	-	-	1	1	9	31	79	186	401	666
2001.....	16,991	-	-	-	-	3	2	27	75	156	337	512
2000.....	16,373	-	-	-	2	1	13	22	53	147	329	462
1999.....	15,654	1	-	-	-	1	8	19	72	139	289	423
<b>Non-Hispanic White, male</b>												
2003.....	8,056	-	-	-	-	-	11	33	68	144	319	491
2002.....	7,427	-	-	-	1	1	8	25	58	137	299	470
2001.....	6,728	-	-	-	-	2	1	18	62	114	253	369
2000.....	6,390	-	-	-	1	1	10	13	40	107	234	336
1999.....	6,013	-	-	-	-	1	6	17	58	101	222	303
<b>Non-Hispanic White, female</b>												
2003.....	11,230	-	-	-	-	1	3	6	26	47	113	182
2002.....	10,633	-	-	-	-	-	1	6	21	49	102	196
2001.....	10,263	-	-	-	-	1	1	9	13	42	84	143
2000.....	9,983	-	-	-	1	-	3	9	13	40	95	126
1999.....	9,641	1	-	-	-	-	2	2	14	38	67	120





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart and renal disease (I13)</b>												
All origins, both sexes												
2003.....	3,108	-	1	-	-	-	5	8	20	35	58	87
2002.....	2,895	-	-	-	-	1	4	11	17	28	49	75
2001.....	2,826	1	-	-	-	2	5	10	18	40	59	64
2000.....	2,785	-	-	-	1	1	5	11	17	22	61	68
1999.....	3,327	-	-	2	-	2	4	5	16	43	71	80
All origins, male												
2003.....	1,326	-	1	-	-	-	4	4	9	26	36	63
2002.....	1,205	-	-	-	-	-	3	9	10	19	25	49
2001.....	1,169	1	-	-	-	-	4	4	13	20	43	46
2000.....	1,163	-	-	-	-	1	3	7	11	12	37	43
1999.....	1,390	-	-	1	-	1	2	3	13	24	43	50
All origins, female												
2003.....	1,782	-	-	-	-	-	1	4	11	9	22	24
2002.....	1,690	-	-	-	-	1	1	2	7	9	24	26
2001.....	1,657	-	-	-	-	2	1	6	5	20	16	18
2000.....	1,622	-	-	-	1	-	2	4	6	10	24	25
1999.....	1,937	-	-	1	-	1	2	2	3	19	28	30
Hispanic, both sexes												
2003.....	158	-	1	-	-	-	1	-	-	1	5	7
2002.....	172	-	-	-	-	1	1	2	1	1	3	7
2001.....	160	-	-	-	-	-	1	1	1	4	5	2
2000.....	138	-	-	-	-	-	1	3	-	1	7	6
1999.....	144	-	-	-	-	-	-	-	1	5	4	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart and renal disease (I13)</b>												
<b>Hispanic, male</b>												
2003.....	75	-	1	-	-	-	1	-	-	1	-	5
2002.....	80	-	-	-	-	-	1	1	-	1	1	2
2001.....	76	-	-	-	-	-	1	-	-	2	3	1
2000.....	67	-	-	-	-	-	1	2	-	1	3	3
1999.....	65	-	-	-	-	-	-	-	-	2	4	5
<b>Hispanic, female</b>												
2003.....	83	-	-	-	-	-	-	-	-	-	5	2
2002.....	92	-	-	-	-	1	-	1	1	-	2	5
2001.....	84	-	-	-	-	-	-	1	1	2	2	1
2000.....	71	-	-	-	-	-	-	1	-	-	4	3
1999.....	79	-	-	-	-	-	-	-	1	3	-	3
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	2,944	-	-	-	-	-	4	8	20	33	53	80
2002.....	2,714	-	-	-	-	-	3	9	16	27	46	68
2001.....	2,650	1	-	-	-	2	4	9	17	36	51	61
2000.....	2,637	-	-	-	1	1	4	8	17	21	54	62
1999.....	3,171	-	-	2	-	2	4	5	15	37	65	69
<b>Non-Hispanic 1/, male</b>												
2003.....	1,248	-	-	-	-	-	3	4	9	24	36	58
2002.....	1,120	-	-	-	-	-	2	8	10	18	24	47
2001.....	1,086	1	-	-	-	-	3	4	13	18	38	44
2000.....	1,089	-	-	-	-	1	2	5	11	11	34	40
1999.....	1,318	-	-	1	-	1	2	3	13	21	38	42

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart and renal disease (I13)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	1,696	-	-	-	-	-	1	4	11	9	17	22
2002.....	1,594	-	-	-	-	-	1	1	6	9	22	21
2001.....	1,564	-	-	-	-	2	1	5	4	18	13	17
2000.....	1,548	-	-	-	1	-	2	3	6	10	20	22
1999.....	1,853	-	-	1	-	1	2	2	2	16	27	27
<b>Non-Hispanic White, both sexes</b>												
2003.....	1,942	-	-	-	-	-	1	4	3	6	8	18
2002.....	1,752	-	-	-	-	-	-	4	4	4	13	12
2001.....	1,759	-	-	-	-	-	2	1	4	3	13	16
2000.....	1,769	-	-	-	-	-	-	3	2	8	22	15
1999.....	2,091	-	-	1	-	-	-	1	5	8	11	15
<b>Non-Hispanic White, male</b>												
2003.....	789	-	-	-	-	-	-	2	-	6	6	14
2002.....	679	-	-	-	-	-	-	3	2	4	9	11
2001.....	687	-	-	-	-	-	2	1	2	1	11	11
2000.....	710	-	-	-	-	-	-	2	2	4	14	9
1999.....	843	-	-	1	-	-	-	1	5	4	7	8
<b>Non-Hispanic White, female</b>												
2003.....	1,153	-	-	-	-	-	1	2	3	-	2	4
2002.....	1,073	-	-	-	-	-	-	1	2	-	4	1
2001.....	1,072	-	-	-	-	-	-	-	2	2	2	5
2000.....	1,059	-	-	-	-	-	-	1	-	4	8	6
1999.....	1,248	-	-	-	-	-	-	-	-	4	4	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hypertensive heart and renal disease (I13)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	910	-	-	-	-	-	3	4	16	27	44	60
2002.....	897	-	-	-	-	-	2	5	12	22	32	56
2001.....	821	1	-	-	-	2	2	8	13	29	36	45
2000.....	797	-	-	-	1	1	4	4	15	13	32	45
1999.....	989	-	-	1	-	2	4	3	10	28	54	52
<b>Non-Hispanic Black, male</b>												
2003.....	411	-	-	-	-	-	3	2	8	18	29	42
2002.....	402	-	-	-	-	-	1	5	8	13	14	36
2001.....	370	1	-	-	-	-	1	3	11	15	26	33
2000.....	348	-	-	-	-	1	2	3	9	7	20	30
1999.....	426	-	-	-	-	1	2	1	8	16	31	32
<b>Non-Hispanic Black, female</b>												
2003.....	499	-	-	-	-	-	-	2	8	9	15	18
2002.....	495	-	-	-	-	-	1	-	4	9	18	20
2001.....	451	-	-	-	-	2	1	5	2	14	10	12
2000.....	449	-	-	-	1	-	2	1	6	6	12	15
1999.....	563	-	-	1	-	1	2	2	2	12	23	20
<b>Origin not stated 2/, both sexes</b>												
2003.....	6	-	-	-	-	-	-	-	-	1	-	-
2002.....	9	-	-	-	-	-	-	-	-	-	-	-
2001.....	16	-	-	-	-	-	-	-	-	-	3	1
2000.....	10	-	-	-	-	-	-	-	-	-	-	-
1999.....	12	-	-	-	-	-	-	-	-	1	2	3



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
All origins, both sexes												
2003.....	480,028	22	12	12	12	32	128	274	808	2,152	5,395	9,989
2002.....	494,382	23	7	4	16	30	103	251	745	2,211	5,472	9,892
2001.....	502,189	26	9	4	16	31	132	279	837	2,212	5,322	9,461
2000.....	515,204	30	10	8	8	33	90	236	699	2,233	5,286	9,638
1999.....	529,659	27	5	13	9	52	85	245	769	2,387	5,364	9,524
All origins, male												
2003.....	246,342	13	6	4	6	21	86	197	616	1,630	4,049	7,661
2002.....	252,760	15	4	1	12	18	74	191	550	1,670	4,170	7,582
2001.....	254,005	17	3	2	11	24	95	211	619	1,667	4,045	7,323
2000.....	260,574	17	7	5	4	21	60	181	523	1,665	4,122	7,601
1999.....	267,268	13	3	10	7	36	57	169	570	1,801	4,169	7,447
All origins, female												
2003.....	233,686	9	6	8	6	11	42	77	192	522	1,346	2,328
2002.....	241,622	8	3	3	4	12	29	60	195	541	1,302	2,310
2001.....	248,184	9	6	2	5	7	37	68	218	545	1,277	2,138
2000.....	254,630	13	3	3	4	12	30	55	176	568	1,164	2,037
1999.....	262,391	14	2	3	2	16	28	76	199	586	1,195	2,077
Hispanic, both sexes												
2003.....	20,783	2	1	3	1	5	17	32	72	151	295	517
2002.....	20,941	4	1	1	2	4	14	29	59	144	298	557
2001.....	20,664	4	1	-	4	5	13	32	77	128	240	495
2000.....	19,744	6	1	-	1	2	10	20	49	121	271	486
1999.....	19,936	2	-	2	-	6	5	24	63	128	238	502

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Hispanic, male												
2003.....	11,054	1	1	2	1	5	15	26	57	115	226	395
2002.....	11,295	2	-	1	1	1	11	26	50	107	239	433
2001.....	11,019	3	1	-	1	4	9	27	63	105	190	398
2000.....	10,562	4	1	-	1	2	7	13	43	98	224	397
1999.....	10,604	-	-	2	-	5	3	23	49	112	198	391
Hispanic, female												
2003.....	9,729	1	-	1	-	-	2	6	15	36	69	122
2002.....	9,646	2	1	-	1	3	3	3	9	37	59	124
2001.....	9,645	1	-	-	3	1	4	5	14	23	50	97
2000.....	9,182	2	-	-	-	-	3	7	6	23	47	89
1999.....	9,332	2	-	-	-	1	2	1	14	16	40	111
Non-Hispanic 1/, both sexes												
2003.....	457,794	20	11	9	11	27	110	241	733	1,996	5,074	9,431
2002.....	471,655	19	6	3	14	26	88	220	680	2,059	5,155	9,300
2001.....	479,842	21	8	4	12	26	118	247	757	2,077	5,058	8,932
2000.....	493,541	23	9	8	7	31	80	214	647	2,100	4,982	9,105
1999.....	507,700	25	5	11	9	45	80	220	706	2,245	5,099	8,980
Non-Hispanic 1/, male												
2003.....	234,433	12	5	2	5	16	70	170	557	1,512	3,804	7,236
2002.....	240,462	13	4	-	11	17	62	163	495	1,557	3,918	7,125
2001.....	242,054	13	2	2	10	20	85	184	553	1,558	3,834	6,896
2000.....	248,932	12	6	5	3	19	53	167	478	1,558	3,868	7,167
1999.....	255,524	13	3	8	7	31	54	145	521	1,677	3,946	7,021

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Non-Hispanic 1/, female												
2003.....	223,361	8	6	7	6	11	40	71	176	484	1,270	2,195
2002.....	231,193	6	2	3	3	9	26	57	185	502	1,237	2,175
2001.....	237,788	8	6	2	2	6	33	63	204	519	1,224	2,036
2000.....	244,609	11	3	3	4	12	27	47	169	542	1,114	1,938
1999.....	252,176	12	2	3	2	14	26	75	185	568	1,153	1,959
Non-Hispanic White, both sexes												
2003.....	400,101	14	5	4	7	20	71	159	536	1,530	3,904	7,281
2002.....	413,230	11	4	2	10	20	56	154	496	1,557	4,008	7,219
2001.....	420,959	14	5	3	7	17	69	167	551	1,582	3,874	6,918
2000.....	434,505	16	9	6	6	26	54	138	468	1,619	3,818	7,064
1999.....	447,960	16	4	7	5	31	45	155	510	1,724	3,883	7,017
Non-Hispanic White, male												
2003.....	205,310	8	2	-	4	12	45	116	411	1,190	3,004	5,746
2002.....	211,391	8	2	-	8	13	41	121	372	1,201	3,145	5,736
2001.....	212,928	10	1	2	6	12	53	130	409	1,208	3,042	5,546
2000.....	219,732	9	6	3	3	16	36	110	363	1,234	3,085	5,778
1999.....	225,909	8	3	5	5	21	36	100	382	1,310	3,106	5,680
Non-Hispanic White, female												
2003.....	194,791	6	3	4	3	8	26	43	125	340	900	1,535
2002.....	201,839	3	2	2	2	7	15	33	124	356	863	1,483
2001.....	208,031	4	4	1	1	5	16	37	142	374	832	1,372
2000.....	214,773	7	3	3	3	10	18	28	105	385	733	1,286
1999.....	222,051	8	1	2	-	10	9	55	128	414	777	1,337

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Non-Hispanic Black, both sexes												
2003.....	48,617	4	5	4	4	7	34	64	165	396	1,011	1,865
2002.....	49,522	7	2	1	4	4	27	55	155	430	1,011	1,835
2001.....	50,295	7	3	1	4	9	43	72	174	434	1,040	1,772
2000.....	50,659	4	-	2	-	4	20	66	163	426	1,025	1,824
1999.....	51,368	8	1	4	4	12	32	53	160	466	1,086	1,748
Non-Hispanic Black, male												
2003.....	23,919	3	3	2	1	4	22	40	119	261	677	1,254
2002.....	23,976	4	2	-	3	2	16	34	97	293	654	1,189
2001.....	24,255	3	1	-	3	8	27	48	119	297	677	1,156
2000.....	24,289	2	-	2	-	2	12	48	102	277	669	1,208
1999.....	24,736	4	-	3	2	8	16	38	110	317	725	1,160
Non-Hispanic Black, female												
2003.....	24,698	1	2	2	3	3	12	24	46	135	334	611
2002.....	25,546	3	-	1	1	2	11	21	58	137	357	646
2001.....	26,040	4	2	1	1	1	16	24	55	137	363	616
2000.....	26,370	2	-	-	-	2	8	18	61	149	356	616
1999.....	26,632	4	1	1	2	4	16	15	50	149	361	588
Origin not stated 2/, both sexes												
2003.....	1,451	-	-	-	-	-	1	1	3	5	26	41
2002.....	1,786	-	-	-	-	-	1	2	6	8	19	35
2001.....	1,683	1	-	-	-	-	1	-	3	7	24	34
2000.....	1,919	1	-	-	-	-	-	2	3	12	33	47
1999.....	2,023	-	-	-	-	1	-	1	-	14	27	42

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Origin not stated 2/, male												
2003.....	855	-	-	-	-	-	1	1	2	3	19	30
2002.....	1,003	-	-	-	-	-	1	2	5	6	13	24
2001.....	932	1	-	-	-	-	1	-	3	4	21	29
2000.....	1,080	1	-	-	-	-	-	1	2	9	30	37
1999.....	1,140	-	-	-	-	-	-	1	-	12	25	35
Origin not stated 2/, female												
2003.....	596	-	-	-	-	-	-	-	1	2	7	11
2002.....	783	-	-	-	-	-	-	-	1	2	6	11
2001.....	751	-	-	-	-	-	-	-	-	3	3	5
2000.....	839	-	-	-	-	-	-	1	1	3	3	10
1999.....	883	-	-	-	-	1	-	-	-	2	2	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
All origins, both sexes												
2003.....	170,564	9	6	6	6	17	56	105	308	881	2,213	4,092
2002.....	179,514	11	5	2	5	15	47	91	309	882	2,279	4,055
2001.....	184,757	15	2	1	6	15	49	105	370	924	2,252	4,097
2000.....	192,898	20	1	5	1	14	38	108	267	904	2,228	4,230
1999.....	199,454	13	2	5	6	25	35	101	333	935	2,314	4,225
All origins, male												
2003.....	89,515	4	2	2	4	11	35	75	225	664	1,637	3,130
2002.....	93,830	7	2	-	2	10	31	67	225	656	1,691	3,066
2001.....	95,875	10	2	-	4	14	34	69	264	691	1,693	3,146
2000.....	100,306	10	1	3	1	8	22	75	194	657	1,730	3,319
1999.....	104,169	6	1	3	4	16	24	68	238	707	1,763	3,261
All origins, female												
2003.....	81,049	5	4	4	2	6	21	30	83	217	576	962
2002.....	85,684	4	3	2	3	5	16	24	84	226	588	989
2001.....	88,882	5	-	1	2	1	15	36	106	233	559	951
2000.....	92,592	10	-	2	-	6	16	33	73	247	498	911
1999.....	95,285	7	1	2	2	9	11	33	95	228	551	964
Hispanic, both sexes												
2003.....	7,370	-	-	2	-	3	8	8	29	59	116	180
2002.....	7,631	3	1	-	1	3	4	9	25	59	118	198
2001.....	7,403	4	1	-	1	3	3	13	33	45	94	201
2000.....	7,106	5	-	-	-	5	5	12	19	47	107	196
1999.....	7,158	2	-	-	-	2	2	10	22	46	96	216

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Hispanic, male												
2003.....	3,903	-	-	1	-	3	7	7	20	42	85	134
2002.....	4,125	2	-	-	-	1	4	7	22	39	87	151
2001.....	3,972	3	1	-	-	3	3	10	27	34	76	166
2000.....	3,825	3	-	-	-	-	4	6	18	35	90	156
1999.....	3,836	-	-	-	-	2	1	10	17	38	78	159
Hispanic, female												
2003.....	3,467	-	-	1	-	-	1	1	9	17	31	46
2002.....	3,506	1	1	-	1	2	-	2	3	20	31	47
2001.....	3,431	1	-	-	1	-	-	3	6	11	18	35
2000.....	3,281	2	-	-	-	-	1	6	1	12	17	40
1999.....	3,322	2	-	-	-	-	1	-	5	8	18	57
Non-Hispanic 1/, both sexes												
2003.....	162,813	9	6	4	6	14	48	97	278	821	2,088	3,901
2002.....	171,441	8	4	2	4	12	43	82	283	822	2,153	3,848
2001.....	176,895	11	1	1	5	12	46	92	336	877	2,146	3,885
2000.....	185,283	14	1	5	1	14	33	96	248	853	2,110	4,027
1999.....	191,729	11	2	5	6	23	33	91	311	883	2,207	3,996
Non-Hispanic 1/, male												
2003.....	85,386	4	2	1	4	8	28	68	204	622	1,546	2,989
2002.....	89,480	5	2	-	2	9	27	60	202	617	1,598	2,910
2001.....	91,655	7	1	-	4	11	31	59	236	655	1,606	2,971
2000.....	96,218	6	1	3	1	8	18	69	176	619	1,631	3,160
1999.....	100,014	6	1	3	4	14	23	58	221	664	1,674	3,092

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Non-Hispanic 1/, female												
2003.....	77,427	5	4	3	2	6	20	29	74	199	542	912
2002.....	81,961	3	2	2	2	3	16	22	81	205	555	938
2001.....	85,240	4	-	1	1	1	15	33	100	222	540	914
2000.....	89,065	8	-	2	-	6	15	27	72	234	479	867
1999.....	91,715	5	1	2	2	9	10	33	90	219	533	904
Non-Hispanic White, both sexes												
2003.....	142,402	7	3	3	4	11	32	67	206	648	1,629	3,080
2002.....	150,051	6	4	1	3	10	26	54	213	639	1,706	3,048
2001.....	155,312	8	1	-	1	9	27	59	254	684	1,683	3,117
2000.....	163,203	10	1	3	1	11	23	59	184	694	1,642	3,249
1999.....	169,545	7	2	3	3	17	18	66	227	710	1,736	3,229
Non-Hispanic White, male												
2003.....	75,157	3	1	-	3	7	19	51	150	508	1,240	2,435
2002.....	79,037	4	2	-	2	7	18	43	157	493	1,300	2,395
2001.....	81,175	6	1	-	1	8	20	37	182	520	1,314	2,485
2000.....	85,412	5	1	1	1	6	14	44	139	519	1,320	2,643
1999.....	89,142	3	1	1	3	11	15	41	161	546	1,361	2,591
Non-Hispanic White, female												
2003.....	67,245	4	2	3	1	4	13	16	56	140	389	645
2002.....	71,014	2	2	1	1	3	8	11	56	146	406	653
2001.....	74,137	2	-	-	-	1	7	22	72	164	369	632
2000.....	77,791	5	-	2	-	5	9	15	45	175	322	606
1999.....	80,403	4	1	2	-	6	3	25	66	164	375	638

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Non-Hispanic Black, both sexes												
2003.....	17,268	1	2	1	2	3	15	24	63	159	414	722
2002.....	18,162	2	-	1	1	-	15	25	63	157	399	702
2001.....	18,399	3	-	1	3	3	18	28	69	168	404	681
2000.....	18,934	2	-	2	-	2	9	32	61	145	415	691
1999.....	19,120	4	-	2	3	6	13	22	66	159	421	685
Non-Hispanic Black, male												
2003.....	8,381	-	1	1	1	1	9	13	46	101	270	471
2002.....	8,607	1	-	-	-	-	7	15	39	102	257	435
2001.....	8,628	1	-	-	2	3	10	18	44	110	244	415
2000.....	8,959	1	-	2	-	1	4	21	34	89	266	445
1999.....	9,106	3	-	2	1	3	7	16	45	104	270	434
Non-Hispanic Black, female												
2003.....	8,887	1	1	-	1	2	6	11	17	58	144	251
2002.....	9,555	1	-	1	1	-	8	10	24	55	142	267
2001.....	9,771	2	-	1	1	-	8	10	25	58	160	266
2000.....	9,975	1	-	-	-	1	5	11	27	56	149	246
1999.....	10,014	1	-	-	2	3	6	6	21	55	151	251
Origin not stated 2/, both sexes												
2003.....	381	-	-	-	-	-	-	-	1	1	9	11
2002.....	442	-	-	-	-	-	-	-	1	1	8	9
2001.....	459	-	-	-	-	-	-	-	1	2	12	11
2000.....	509	1	-	-	-	-	-	-	-	4	11	7
1999.....	567	-	-	-	-	-	-	-	-	6	11	13

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Origin not stated 2/, male												
2003.....	226	-	-	-	-	-	-	-	1	-	6	7
2002.....	225	-	-	-	-	-	-	-	1	-	6	5
2001.....	248	-	-	-	-	-	-	-	1	2	11	9
2000.....	263	1	-	-	-	-	-	-	-	3	9	3
1999.....	319	-	-	-	-	-	-	-	-	5	11	10
Origin not stated 2/, female												
2003.....	155	-	-	-	-	-	-	-	-	1	3	4
2002.....	217	-	-	-	-	-	-	-	-	1	2	4
2001.....	211	-	-	-	-	-	-	-	-	-	1	2
2000.....	246	-	-	-	-	-	-	-	-	1	2	4
1999.....	248	-	-	-	-	-	-	-	-	1	-	3



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Hispanic, male												
2003.....	21	-	-	-	-	-	1	1	-	-	1	1
2002.....	40	-	-	1	-	-	-	1	-	-	1	4
2001.....	42	-	-	-	-	-	-	-	-	-	1	-
2000.....	39	-	1	-	-	1	-	1	-	-	1	2
1999.....	26	-	-	1	-	-	-	-	-	-	1	1
Hispanic, female												
2003.....	26	-	-	-	-	-	-	-	-	-	-	2
2002.....	24	-	-	-	-	1	-	-	-	-	-	1
2001.....	21	-	-	-	-	-	-	-	-	-	-	1
2000.....	30	-	-	-	-	-	-	-	-	-	-	2
1999.....	27	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes												
2003.....	3,104	4	1	1	-	-	4	2	8	27	70	103
2002.....	3,339	2	-	-	-	1	1	3	14	17	58	114
2001.....	3,284	3	-	-	2	-	7	2	12	32	61	119
2000.....	3,282	1	-	-	-	3	4	3	12	30	60	89
1999.....	3,167	3	1	-	-	-	3	2	4	25	55	91
Non-Hispanic 1/, male												
2003.....	1,595	2	1	-	-	-	4	1	5	20	52	74
2002.....	1,725	1	-	-	-	-	-	1	10	9	40	84
2001.....	1,675	2	-	-	1	-	6	2	8	18	46	89
2000.....	1,705	1	-	-	-	2	3	2	10	24	45	74
1999.....	1,635	1	1	-	-	-	3	1	3	19	43	68

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Non-Hispanic 1/, female												
2003.....	1,509	2	-	1	-	-	-	1	3	7	18	29
2002.....	1,614	1	-	-	-	1	1	2	4	8	18	30
2001.....	1,609	1	-	-	1	-	1	-	4	14	15	30
2000.....	1,577	-	-	-	-	1	1	1	2	6	15	15
1999.....	1,532	2	-	-	-	-	-	1	1	6	12	23
Non-Hispanic White, both sexes												
2003.....	2,613	2	-	1	-	-	3	1	4	20	48	70
2002.....	2,843	1	-	-	-	1	-	2	9	15	39	85
2001.....	2,756	3	-	-	2	-	4	2	9	23	39	88
2000.....	2,767	1	-	-	-	3	2	3	8	23	43	65
1999.....	2,689	1	1	-	-	-	3	2	3	13	39	55
Non-Hispanic White, male												
2003.....	1,314	1	-	-	-	-	3	1	3	16	39	50
2002.....	1,454	1	-	-	-	-	-	1	6	8	33	60
2001.....	1,386	2	-	-	1	-	3	2	7	14	31	69
2000.....	1,412	1	-	-	-	2	1	2	6	19	33	59
1999.....	1,361	1	1	-	-	-	3	1	3	10	30	41
Non-Hispanic White, female												
2003.....	1,299	1	-	1	-	-	-	-	1	4	9	20
2002.....	1,389	-	-	-	-	1	-	1	3	7	6	25
2001.....	1,370	1	-	-	1	-	1	-	2	9	8	19
2000.....	1,355	-	-	-	-	1	1	1	2	4	10	6
1999.....	1,328	-	-	-	-	-	-	1	-	3	9	14





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
All origins, both sexes												
2003.....	306,306	9	5	5	6	15	67	166	492	1,244	3,111	5,791
2002.....	311,461	10	2	1	11	13	54	156	422	1,312	3,133	5,718
2001.....	314,081	8	7	3	8	16	76	172	455	1,256	3,008	5,244
2000.....	318,943	9	8	3	7	15	48	124	420	1,299	2,996	5,315
1999.....	326,973	11	2	7	3	27	47	142	432	1,427	2,994	5,207
All origins, male												
2003.....	155,206	7	3	2	2	10	46	120	386	946	2,359	4,456
2002.....	157,163	7	2	-	10	8	42	122	315	1,005	2,438	4,428
2001.....	156,410	5	1	2	6	10	55	140	347	958	2,305	4,088
2000.....	158,519	6	5	2	3	10	35	103	319	984	2,345	4,206
1999.....	161,434	6	1	6	3	20	30	100	329	1,075	2,362	4,117
All origins, female												
2003.....	151,100	2	2	3	4	5	21	46	106	298	752	1,335
2002.....	154,298	3	-	1	1	5	12	34	107	307	695	1,290
2001.....	157,671	3	6	1	2	6	21	32	108	298	703	1,156
2000.....	160,424	3	3	1	4	5	13	21	101	315	651	1,109
1999.....	165,539	5	1	1	-	7	17	42	103	352	632	1,090
Hispanic, both sexes												
2003.....	13,366	2	1	1	1	2	8	23	43	92	178	334
2002.....	13,246	1	-	-	1	-	10	19	34	85	179	354
2001.....	13,198	-	-	-	3	2	10	19	44	83	145	293
2000.....	12,569	1	-	-	1	1	5	7	30	74	163	286
1999.....	12,725	-	-	1	-	4	3	14	41	82	141	285

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Hispanic, male												
2003.....	7,130	1	1	1	1	2	7	18	37	73	140	260
2002.....	7,130	-	-	-	1	-	7	18	28	68	151	278
2001.....	7,005	-	-	-	1	1	6	17	36	71	113	232
2000.....	6,698	1	-	-	1	1	3	6	25	63	133	239
1999.....	6,742	-	-	1	-	3	2	13	32	74	119	231
Hispanic, female												
2003.....	6,236	1	-	-	-	-	1	5	6	19	38	74
2002.....	6,116	1	-	-	-	-	3	1	6	17	28	76
2001.....	6,193	-	-	-	2	1	4	2	8	12	32	61
2000.....	5,871	-	-	-	-	-	2	1	5	11	30	47
1999.....	5,983	-	-	-	-	1	1	1	9	8	22	54
Non-Hispanic 1/, both sexes												
2003.....	291,877	7	4	4	5	13	58	142	447	1,148	2,916	5,427
2002.....	296,875	9	2	1	10	13	44	135	383	1,220	2,944	5,338
2001.....	299,663	7	7	3	5	14	65	153	409	1,168	2,851	4,928
2000.....	304,976	8	8	3	6	14	43	115	387	1,217	2,812	4,989
1999.....	312,804	11	2	6	3	22	44	127	391	1,337	2,837	4,893
Non-Hispanic 1/, male												
2003.....	147,452	6	2	1	1	8	38	101	348	870	2,206	4,173
2002.....	149,257	7	2	-	9	8	35	102	283	931	2,280	4,131
2001.....	148,724	4	1	2	5	9	48	123	309	885	2,182	3,836
2000.....	151,009	5	5	2	2	9	32	96	292	915	2,192	3,933
1999.....	153,875	6	1	5	3	17	28	86	297	994	2,229	3,861

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic 1/, female												
2003.....	144,425	1	2	3	4	5	20	41	99	278	710	1,254
2002.....	147,618	2	-	1	1	5	9	33	100	289	664	1,207
2001.....	150,939	3	6	1	-	5	17	30	100	283	669	1,092
2000.....	153,967	3	3	1	4	5	11	19	95	302	620	1,056
1999.....	158,929	5	1	1	-	5	16	41	94	343	608	1,032
Non-Hispanic White, both sexes												
2003.....	255,086	5	2	-	3	9	36	91	326	862	2,227	4,131
2002.....	260,336	4	-	1	7	9	30	98	274	903	2,263	4,086
2001.....	262,891	3	4	3	4	8	38	106	288	875	2,152	3,713
2000.....	268,535	5	8	3	5	12	29	76	276	902	2,133	3,750
1999.....	275,726	8	1	4	2	14	24	87	280	1,001	2,108	3,733
Non-Hispanic White, male												
2003.....	128,839	4	1	-	1	5	23	64	258	666	1,725	3,261
2002.....	130,900	3	-	-	6	6	23	77	209	700	1,812	3,281
2001.....	130,367	2	-	2	4	4	30	91	220	674	1,697	2,992
2000.....	132,908	3	5	2	2	8	21	64	218	696	1,732	3,076
1999.....	135,406	4	1	4	2	10	18	58	218	754	1,715	3,048
Non-Hispanic White, female												
2003.....	126,247	1	1	-	2	4	13	27	68	196	502	870
2002.....	129,436	1	-	1	1	3	7	21	65	203	451	805
2001.....	132,524	1	4	1	-	4	8	15	68	201	455	721
2000.....	135,627	2	3	1	3	4	8	12	58	206	401	674
1999.....	140,320	4	-	-	-	4	6	29	62	247	393	685

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Non-Hispanic Black, both sexes												
2003.....	30,917	2	2	3	2	4	18	39	100	230	577	1,113
2002.....	30,917	4	2	-	3	4	11	29	87	271	595	1,106
2001.....	31,443	4	3	-	1	6	22	44	103	258	615	1,064
2000.....	31,276	2	-	-	-	2	10	34	98	274	595	1,110
1999.....	31,819	2	1	2	1	6	19	31	93	295	651	1,029
Non-Hispanic Black, male												
2003.....	15,284	2	1	1	-	3	12	27	72	156	395	762
2002.....	15,127	3	2	-	3	2	9	19	54	190	392	731
2001.....	15,378	2	1	-	1	5	14	30	75	183	418	724
2000.....	15,075	1	-	-	-	1	7	27	64	183	392	748
1999.....	15,384	1	-	1	1	5	9	22	65	204	444	700
Non-Hispanic Black, female												
2003.....	15,633	-	1	2	2	1	6	12	28	74	182	351
2002.....	15,790	1	-	-	-	2	2	10	33	81	203	375
2001.....	16,065	2	2	-	-	1	8	14	28	75	197	340
2000.....	16,201	1	-	-	-	1	3	7	34	91	203	362
1999.....	16,435	1	1	1	-	1	10	9	28	91	207	329
Origin not stated 2/, both sexes												
2003.....	1,063	-	-	-	-	-	1	1	2	4	17	30
2002.....	1,340	-	-	-	-	-	-	2	5	7	10	26
2001.....	1,220	1	-	-	-	-	1	-	2	5	12	23
2000.....	1,398	-	-	-	-	-	-	2	3	8	21	40
1999.....	1,444	-	-	-	-	1	-	1	-	8	16	29

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Origin not stated 2/, male												
2003.....	624	-	-	-	-	-	1	1	1	3	13	23
2002.....	776	-	-	-	-	-	-	2	4	6	7	19
2001.....	681	1	-	-	-	-	1	-	2	2	10	20
2000.....	812	-	-	-	-	-	-	1	2	6	20	34
1999.....	817	-	-	-	-	-	-	1	-	7	14	25
Origin not stated 2/, female												
2003.....	439	-	-	-	-	-	-	-	1	1	4	7
2002.....	564	-	-	-	-	-	-	-	1	1	3	7
2001.....	539	-	-	-	-	-	-	-	-	3	2	3
2000.....	586	-	-	-	-	-	-	1	1	2	1	6
1999.....	627	-	-	-	-	1	-	-	-	1	2	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
All origins, both sexes												
2003.....	67,232	2	-	-	-	2	14	50	161	414	1,074	2,151
2002.....	68,129	2	-	-	1	4	13	39	142	410	1,105	2,081
2001.....	68,103	-	-	-	-	3	17	39	140	402	1,055	1,838
2000.....	69,554	1	-	-	1	-	15	33	111	391	985	1,894
1999.....	71,926	-	1	-	1	-	13	32	128	453	1,035	1,792
All origins, male												
2003.....	36,042	2	-	-	-	1	11	40	141	315	842	1,715
2002.....	36,047	1	-	-	-	3	10	31	113	325	891	1,634
2001.....	35,440	-	-	-	-	2	14	30	103	315	815	1,488
2000.....	35,896	-	-	-	1	-	12	26	89	312	787	1,533
1999.....	36,633	-	-	-	1	-	10	21	98	355	837	1,438
All origins, female												
2003.....	31,190	-	-	-	-	1	3	10	20	99	232	436
2002.....	32,082	1	-	-	1	1	3	8	29	85	214	447
2001.....	32,663	-	-	-	-	1	3	9	37	87	240	350
2000.....	33,658	1	-	-	-	-	3	7	22	79	198	361
1999.....	35,293	-	1	-	-	-	3	11	30	98	198	354
Hispanic, both sexes												
2003.....	3,295	-	-	-	-	-	3	5	16	30	63	155
2002.....	3,166	1	-	-	-	-	4	9	17	30	78	141
2001.....	3,150	-	-	-	-	-	1	8	17	26	55	120
2000.....	2,953	-	-	-	-	-	2	2	5	28	55	111
1999.....	3,063	-	-	-	-	-	1	4	10	23	50	110

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hispanic, male												
2003.....	2,094	-	-	-	-	-	3	5	15	25	55	132
2002.....	1,946	-	-	-	-	-	3	8	16	22	67	112
2001.....	1,928	-	-	-	-	-	-	8	13	22	41	96
2000.....	1,794	-	-	-	-	-	2	1	5	25	46	97
1999.....	1,847	-	-	-	-	-	1	3	9	22	41	89
Hispanic, female												
2003.....	1,201	-	-	-	-	-	-	-	1	5	8	23
2002.....	1,220	1	-	-	-	-	1	1	1	8	11	29
2001.....	1,222	-	-	-	-	-	1	-	4	4	14	24
2000.....	1,159	-	-	-	-	-	-	1	-	3	9	14
1999.....	1,216	-	-	-	-	-	-	1	1	1	9	21
Non-Hispanic 1/, both sexes												
2003.....	63,550	2	-	-	-	2	11	44	145	381	1,004	1,976
2002.....	64,528	1	-	-	1	4	9	28	124	378	1,025	1,928
2001.....	64,534	-	-	-	-	3	16	31	123	373	993	1,706
2000.....	66,141	1	-	-	1	-	13	31	106	360	923	1,764
1999.....	68,373	-	1	-	1	-	12	28	118	427	977	1,671
Non-Hispanic 1/, male												
2003.....	33,677	2	-	-	-	1	8	34	126	287	780	1,566
2002.....	33,809	1	-	-	-	3	7	21	97	302	822	1,514
2001.....	33,247	-	-	-	-	2	14	22	90	292	767	1,381
2000.....	33,786	-	-	-	1	-	10	25	84	285	734	1,419
1999.....	34,451	-	-	-	1	-	9	18	89	330	788	1,339

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic 1/, female												
2003.....	29,873	-	-	-	-	1	3	10	19	94	224	410
2002.....	30,719	-	-	-	1	1	2	7	27	76	203	414
2001.....	31,287	-	-	-	-	1	2	9	33	81	226	325
2000.....	32,355	1	-	-	-	-	3	6	22	75	189	345
1999.....	33,922	-	1	-	-	-	3	10	29	97	189	332
Non-Hispanic White, both sexes												
2003.....	52,268	1	-	-	-	2	7	29	107	293	754	1,442
2002.....	53,161	1	-	-	1	4	7	21	84	265	755	1,458
2001.....	53,031	-	-	-	-	2	9	20	88	275	724	1,244
2000.....	54,544	-	-	-	1	-	11	17	78	261	683	1,274
1999.....	56,375	-	-	-	1	-	5	16	80	310	707	1,191
Non-Hispanic White, male												
2003.....	27,329	1	-	-	-	1	5	21	93	220	594	1,180
2002.....	27,493	1	-	-	-	3	5	16	68	216	628	1,178
2001.....	26,960	-	-	-	-	1	7	15	67	221	573	1,042
2000.....	27,461	-	-	-	1	-	8	13	64	211	564	1,060
1999.....	27,943	-	-	-	1	-	5	11	62	244	594	984
Non-Hispanic White, female												
2003.....	24,939	-	-	-	-	1	2	8	14	73	160	262
2002.....	25,668	-	-	-	1	1	2	5	16	49	127	280
2001.....	26,071	-	-	-	-	1	2	5	21	54	151	202
2000.....	27,083	-	-	-	-	-	3	4	14	50	119	214
1999.....	28,432	-	-	-	-	-	-	5	18	66	113	207

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Non-Hispanic Black, both sexes												
2003.....	9,764	1	-	-	-	-	3	12	34	72	209	452
2002.....	10,024	-	-	-	-	-	2	5	34	99	234	426
2001.....	10,156	-	-	-	-	1	6	11	27	89	235	408
2000.....	10,283	1	-	-	-	-	1	12	24	86	206	451
1999.....	10,643	-	1	-	-	-	7	8	33	110	248	433
Non-Hispanic Black, male												
2003.....	5,431	1	-	-	-	-	2	10	29	51	152	318
2002.....	5,483	-	-	-	-	-	2	3	24	73	160	298
2001.....	5,476	-	-	-	-	1	6	7	17	64	166	291
2000.....	5,528	-	-	-	-	-	1	10	16	64	143	324
1999.....	5,692	-	-	-	-	-	4	5	22	80	174	313
Non-Hispanic Black, female												
2003.....	4,333	-	-	-	-	-	1	2	5	21	57	134
2002.....	4,541	-	-	-	-	-	-	2	10	26	74	128
2001.....	4,680	-	-	-	-	-	-	4	10	25	69	117
2000.....	4,755	1	-	-	-	-	-	2	8	22	63	127
1999.....	4,951	-	1	-	-	-	3	3	11	30	74	120
Origin not stated 2/, both sexes												
2003.....	387	-	-	-	-	-	-	1	-	3	7	20
2002.....	435	-	-	-	-	-	-	2	1	2	2	12
2001.....	419	-	-	-	-	-	-	-	-	3	7	12
2000.....	460	-	-	-	-	-	-	-	-	3	7	19
1999.....	490	-	-	-	-	-	-	-	-	3	8	11



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All origins, both sexes												
2003.....	239,074	7	5	5	6	13	53	116	331	830	2,037	3,640
2002.....	243,332	8	2	1	10	9	41	117	280	902	2,028	3,637
2001.....	245,978	8	7	3	8	13	59	133	315	854	1,953	3,406
2000.....	249,389	8	8	3	6	15	33	91	309	908	2,011	3,421
1999.....	255,047	11	1	7	2	27	34	110	304	974	1,959	3,415
All origins, male												
2003.....	119,164	5	3	2	2	9	35	80	245	631	1,517	2,741
2002.....	121,116	6	2	-	10	5	32	91	202	680	1,547	2,794
2001.....	120,970	5	1	2	6	8	41	110	244	643	1,490	2,600
2000.....	122,623	6	5	2	2	10	23	77	230	672	1,558	2,673
1999.....	124,801	6	1	6	2	20	20	79	231	720	1,525	2,679
All origins, female												
2003.....	119,910	2	2	3	4	4	18	36	86	199	520	899
2002.....	122,216	2	-	1	-	4	9	26	78	222	481	843
2001.....	125,008	3	6	1	2	5	18	23	71	211	463	806
2000.....	126,766	2	3	1	4	5	10	14	79	236	453	748
1999.....	130,246	5	-	1	-	7	14	31	73	254	434	736
Hispanic, both sexes												
2003.....	10,071	2	1	1	1	2	5	18	27	62	115	179
2002.....	10,080	-	-	-	1	-	6	10	17	55	101	213
2001.....	10,048	-	-	-	3	2	9	11	27	57	90	173
2000.....	9,616	1	-	-	1	1	3	5	25	46	108	175
1999.....	9,662	-	-	1	-	4	2	10	31	59	91	175

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hispanic, male												
2003.....	5,036	1	1	1	1	2	4	13	22	48	85	128
2002.....	5,184	-	-	-	1	-	4	10	12	46	84	166
2001.....	5,077	-	-	-	1	1	6	9	23	49	72	136
2000.....	4,904	1	-	-	1	1	1	5	20	38	87	142
1999.....	4,895	-	-	1	-	3	1	10	23	52	78	142
Hispanic, female												
2003.....	5,035	1	-	-	-	-	1	5	5	14	30	51
2002.....	4,896	-	-	-	-	-	2	-	5	9	17	47
2001.....	4,971	-	-	-	2	1	3	2	4	8	18	37
2000.....	4,712	-	-	-	-	-	2	-	5	8	21	33
1999.....	4,767	-	-	-	-	1	1	-	8	7	13	33
Non-Hispanic 1/, both sexes												
2003.....	228,327	5	4	4	5	11	47	98	302	767	1,912	3,451
2002.....	232,347	8	2	1	9	9	35	107	259	842	1,919	3,410
2001.....	235,129	7	7	3	5	11	49	122	286	795	1,858	3,222
2000.....	238,835	7	8	3	5	14	30	84	281	857	1,889	3,225
1999.....	244,431	11	1	6	2	22	32	99	273	910	1,860	3,222
Non-Hispanic 1/, male												
2003.....	113,775	4	2	1	1	7	30	67	222	583	1,426	2,607
2002.....	115,448	6	2	-	9	5	28	81	186	629	1,458	2,617
2001.....	115,477	4	1	2	5	7	34	101	219	593	1,415	2,455
2000.....	117,223	5	5	2	1	9	22	71	208	630	1,458	2,514
1999.....	119,424	6	1	5	2	17	19	68	208	664	1,441	2,522

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Non-Hispanic 1/, female												
2003.....	114,552	1	2	3	4	4	17	31	80	184	486	844
2002.....	116,899	2	-	1	-	4	7	26	73	213	461	793
2001.....	119,652	3	6	1	-	4	15	21	67	202	443	767
2000.....	121,612	2	3	1	4	5	8	13	73	227	431	711
1999.....	125,007	5	-	1	-	5	13	31	65	246	419	700
Non-Hispanic White, both sexes												
2003.....	202,818	4	2	-	3	7	29	62	219	569	1,473	2,689
2002.....	207,175	3	-	1	6	5	23	77	190	638	1,508	2,628
2001.....	209,860	3	4	3	4	6	29	86	200	600	1,428	2,469
2000.....	213,991	5	8	3	4	12	18	59	198	641	1,450	2,476
1999.....	219,351	8	1	4	1	14	19	71	200	691	1,401	2,542
Non-Hispanic White, male												
2003.....	101,510	3	1	-	1	4	18	43	165	446	1,131	2,081
2002.....	103,407	2	-	-	6	3	18	61	141	484	1,184	2,103
2001.....	103,407	2	-	2	4	3	23	76	153	453	1,124	1,950
2000.....	105,447	3	5	2	1	8	13	51	154	485	1,168	2,016
1999.....	107,463	4	1	4	1	10	13	47	156	510	1,121	2,064
Non-Hispanic White, female												
2003.....	101,308	1	1	-	2	3	11	19	54	123	342	608
2002.....	103,768	1	-	1	-	2	5	16	49	154	324	525
2001.....	106,453	1	4	1	-	3	6	10	47	147	304	519
2000.....	108,544	2	3	1	3	4	5	8	44	156	282	460
1999.....	111,888	4	-	-	-	4	6	24	44	181	280	478

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Non-Hispanic Black, both sexes												
2003.....	21,153	1	2	3	2	4	15	27	66	158	368	661
2002.....	20,893	4	2	-	3	4	9	24	53	172	361	680
2001.....	21,287	4	3	-	1	5	16	33	76	169	380	656
2000.....	20,993	1	-	-	-	2	9	22	74	188	389	659
1999.....	21,176	2	-	2	1	6	12	23	60	185	403	596
Non-Hispanic Black, male												
2003.....	9,853	1	1	1	-	3	10	17	43	105	243	444
2002.....	9,644	3	2	-	3	2	7	16	30	117	232	433
2001.....	9,902	2	1	-	1	4	8	23	58	119	252	433
2000.....	9,547	1	-	-	-	1	6	17	48	119	249	424
1999.....	9,692	1	-	1	1	5	5	17	43	124	270	387
Non-Hispanic Black, female												
2003.....	11,300	-	1	2	2	1	5	10	23	53	125	217
2002.....	11,249	1	-	-	-	2	2	8	23	55	129	247
2001.....	11,385	2	2	-	-	1	8	10	18	50	128	223
2000.....	11,446	-	-	-	-	1	3	5	26	69	140	235
1999.....	11,484	1	-	1	-	1	7	6	17	61	133	209
Origin not stated 2/, both sexes												
2003.....	676	-	-	-	-	-	1	-	2	1	10	10
2002.....	905	-	-	-	-	-	-	-	4	5	8	14
2001.....	801	1	-	-	-	-	1	-	2	2	5	11
2000.....	938	-	-	-	-	-	-	2	3	5	14	21
1999.....	954	-	-	-	-	1	-	1	-	5	8	18

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Origin not stated 2/, male												
2003.....	353	-	-	-	-	-	1	-	1	-	6	6
2002.....	484	-	-	-	-	-	-	-	4	5	5	11
2001.....	416	1	-	-	-	-	1	-	2	1	3	9
2000.....	496	-	-	-	-	-	-	1	2	4	13	17
1999.....	482	-	-	-	-	-	-	1	-	4	6	15
Origin not stated 2/, female												
2003.....	323	-	-	-	-	-	-	-	1	1	4	4
2002.....	421	-	-	-	-	-	-	-	-	-	3	3
2001.....	385	-	-	-	-	-	-	-	-	1	2	2
2000.....	442	-	-	-	-	-	-	1	1	1	1	4
1999.....	472	-	-	-	-	1	-	-	-	1	2	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Other heart diseases (I26-I51)</b>												
<b>All origins, both sexes</b>												
2003.....	169,984	416	171	90	144	348	556	692	1,118	1,673	2,882	3,891
2002.....	169,540	473	157	84	143	362	474	723	1,085	1,769	2,851	3,849
2001.....	166,949	449	215	92	154	305	489	687	1,049	1,770	2,702	3,708
2000.....	165,428	457	168	96	152	355	498	687	1,062	1,706	2,751	3,536
1999.....	165,828	490	177	99	149	398	482	680	1,099	1,879	2,768	3,502
<b>All origins, male</b>												
2003.....	74,738	233	81	50	91	219	341	404	677	1,017	1,784	2,439
2002.....	74,230	249	82	38	87	250	307	446	692	1,051	1,809	2,468
2001.....	72,439	222	117	41	82	193	307	413	624	1,099	1,678	2,361
2000.....	72,122	238	98	58	92	218	300	407	642	1,033	1,713	2,278
1999.....	72,601	255	100	54	83	250	283	400	667	1,176	1,718	2,223
<b>All origins, female</b>												
2003.....	95,246	183	90	40	53	129	215	288	441	656	1,098	1,452
2002.....	95,310	224	75	46	56	112	167	277	393	718	1,042	1,381
2001.....	94,510	227	98	51	72	112	182	274	425	671	1,024	1,347
2000.....	93,306	219	70	38	60	137	198	280	420	673	1,038	1,258
1999.....	93,227	235	77	45	66	148	199	280	432	703	1,050	1,279
<b>Hispanic, both sexes</b>												
2003.....	5,612	72	33	6	20	38	60	88	121	137	174	222
2002.....	5,152	81	23	11	16	34	68	85	103	124	174	191
2001.....	4,899	75	31	17	16	39	47	73	106	123	145	171
2000.....	4,792	79	28	17	20	46	64	79	103	129	154	151
1999.....	4,687	92	33	16	18	45	54	63	113	147	162	153

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Other heart diseases (I26-I51)</b>												
<b>Hispanic, male</b>												
2003.....	2,805	40	17	3	12	23	45	56	79	95	122	153
2002.....	2,587	39	11	5	7	25	51	64	71	77	118	117
2001.....	2,401	40	14	9	8	22	35	51	68	83	102	120
2000.....	2,393	39	20	9	16	34	42	55	69	79	98	95
1999.....	2,392	40	20	9	12	25	35	49	79	95	113	97
<b>Hispanic, female</b>												
2003.....	2,807	32	16	3	8	15	15	32	42	42	52	69
2002.....	2,565	42	12	6	9	9	17	21	32	47	56	74
2001.....	2,498	35	17	8	8	17	12	22	38	40	43	51
2000.....	2,399	40	8	8	4	12	22	24	34	50	56	56
1999.....	2,295	52	13	7	6	20	19	14	34	52	49	56
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	163,998	342	137	83	124	306	496	602	992	1,532	2,693	3,655
2002.....	163,970	390	134	73	127	328	405	636	976	1,636	2,664	3,646
2001.....	161,625	372	183	75	138	264	437	610	940	1,640	2,544	3,516
2000.....	160,222	375	139	78	132	309	432	604	954	1,569	2,582	3,367
1999.....	160,711	397	143	83	130	352	427	613	981	1,723	2,594	3,337
<b>Non-Hispanic 1/, male</b>												
2003.....	71,731	193	64	46	79	194	296	348	594	919	1,650	2,273
2002.....	71,455	209	71	33	80	225	255	382	617	967	1,682	2,343
2001.....	69,830	181	103	32	74	169	270	359	554	1,010	1,566	2,227
2000.....	69,547	197	78	49	76	184	257	349	572	948	1,606	2,172
1999.....	69,984	215	80	45	70	224	247	349	584	1,075	1,597	2,118

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Non-Hispanic 1/, female												
2003.....	92,267	149	73	37	45	112	200	254	398	613	1,043	1,382
2002.....	92,515	181	63	40	47	103	150	254	359	669	982	1,303
2001.....	91,795	191	80	43	64	95	167	251	386	630	978	1,289
2000.....	90,675	178	61	29	56	125	175	255	382	621	976	1,195
1999.....	90,727	182	63	38	60	128	180	264	397	648	997	1,219
Non-Hispanic White, both sexes												
2003.....	141,413	186	77	50	82	197	298	342	599	970	1,798	2,513
2002.....	141,712	234	86	50	80	216	237	361	575	1,045	1,843	2,453
2001.....	139,716	219	103	48	92	164	258	363	574	1,039	1,657	2,382
2000.....	138,745	238	87	46	76	184	244	354	594	988	1,729	2,285
1999.....	138,869	240	87	56	84	235	260	360	612	1,103	1,711	2,238
Non-Hispanic White, male												
2003.....	61,119	102	32	27	55	125	173	199	377	591	1,146	1,608
2002.....	61,048	128	51	23	51	149	155	217	372	629	1,207	1,623
2001.....	59,513	116	58	21	51	102	159	216	339	650	1,081	1,562
2000.....	59,461	127	51	30	47	106	151	210	366	617	1,109	1,539
1999.....	59,549	136	51	30	43	144	147	209	371	708	1,101	1,482
Non-Hispanic White, female												
2003.....	80,294	84	45	23	27	72	125	143	222	379	652	905
2002.....	80,664	106	35	27	29	67	82	144	203	416	636	830
2001.....	80,203	103	45	27	41	62	99	147	235	389	576	820
2000.....	79,284	111	36	16	29	78	93	144	228	371	620	746
1999.....	79,320	104	36	26	41	91	113	151	241	395	610	756

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Other heart diseases (I26-I51)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	19,861	125	49	29	37	94	179	236	350	514	832	1,067
2002.....	19,641	135	38	16	41	95	151	250	358	545	761	1,105
2001.....	19,496	130	70	25	38	91	161	228	339	537	833	1,050
2000.....	19,159	113	46	27	45	112	166	233	326	524	790	1,013
1999.....	19,531	132	48	26	40	109	153	232	344	572	823	1,016
<b>Non-Hispanic Black, male</b>												
2003.....	9,275	76	27	16	22	62	113	136	186	300	467	619
2002.....	9,038	72	16	6	24	61	90	149	214	308	430	667
2001.....	9,070	59	39	10	21	60	100	133	194	318	450	605
2000.....	8,883	56	24	15	24	67	91	128	185	298	449	590
1999.....	9,258	66	27	14	24	76	90	132	199	335	462	589
<b>Non-Hispanic Black, female</b>												
2003.....	10,586	49	22	13	15	32	66	100	164	214	365	448
2002.....	10,603	63	22	10	17	34	61	101	144	237	331	438
2001.....	10,426	71	31	15	17	31	61	95	145	219	383	445
2000.....	10,276	57	22	12	21	45	75	105	141	226	341	423
1999.....	10,273	66	21	12	16	33	63	100	145	237	361	427
<b>Origin not stated 2/, both sexes</b>												
2003.....	374	2	1	1	-	4	-	2	5	4	15	14
2002.....	418	2	-	-	-	-	1	2	6	9	13	12
2001.....	425	2	1	-	-	2	5	4	3	7	13	21
2000.....	414	3	1	1	-	-	2	4	5	8	15	18
1999.....	430	1	1	-	1	1	1	4	5	9	12	12

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Origin not stated 2/, male												
2003.....	202	-	-	1	-	2	-	-	4	3	12	13
2002.....	188	1	-	-	-	-	1	-	4	7	9	8
2001.....	208	1	-	-	-	2	2	3	2	6	10	14
2000.....	182	2	-	-	-	-	1	3	1	6	9	11
1999.....	225	-	-	-	1	1	1	2	4	6	8	8
Origin not stated 2/, female												
2003.....	172	2	1	-	-	2	-	2	1	1	3	1
2002.....	230	1	-	-	-	-	-	2	2	2	4	4
2001.....	217	1	1	-	-	-	3	1	1	1	3	7
2000.....	232	1	1	1	-	-	1	1	4	2	6	7
1999.....	205	1	1	-	-	-	-	2	1	3	4	4

Worktable Orig291F. Deaths from I13 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Acute and subacute endocarditis (I33)</b>												
<b>All origins, both sexes</b>												
2003.....	1,245	1	-	-	1	3	13	18	15	43	69	89
2002.....	1,154	2	2	1	1	2	5	16	28	26	64	84
2001.....	1,178	1	-	1	2	3	5	9	20	46	71	93
2000.....	1,172	3	1	2	1	5	8	13	27	41	75	89
1999.....	1,116	2	1	-	-	2	5	12	18	54	60	68
<b>All origins, male</b>												
2003.....	682	1	-	-	1	-	6	11	8	22	41	60
2002.....	643	1	1	-	1	2	4	9	22	19	41	55
2001.....	668	1	-	-	1	2	3	5	12	25	37	68
2000.....	665	2	-	1	-	1	6	7	17	24	49	59
1999.....	639	2	1	-	-	-	3	9	15	34	41	49
<b>All origins, female</b>												
2003.....	563	-	-	-	-	3	7	7	7	21	28	29
2002.....	511	1	1	1	-	-	1	7	6	7	23	29
2001.....	510	-	-	1	1	1	2	4	8	21	34	25
2000.....	507	1	1	1	1	4	2	6	10	17	26	30
1999.....	477	-	-	-	-	2	2	3	3	20	19	19
<b>Hispanic, both sexes</b>												
2003.....	80	1	-	-	1	1	2	2	2	4	6	9
2002.....	69	1	1	-	-	-	-	2	5	2	3	9
2001.....	56	-	-	1	-	-	1	1	5	4	4	5
2000.....	70	-	1	1	-	-	-	2	4	5	6	4
1999.....	63	-	1	-	-	1	1	1	5	4	6	3

Worktable Orig291F. Deaths from I13 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Acute and subacute endocarditis (I33)</b>												
<b>Hispanic, male</b>												
2003.....	42	1	-	-	1	-	1	1	2	3	4	5
2002.....	45	-	-	-	-	-	-	2	4	2	3	7
2001.....	34	-	-	-	-	-	1	1	4	4	2	4
2000.....	39	-	-	-	-	-	-	1	3	3	4	2
1999.....	42	-	1	-	-	-	-	1	4	2	4	2
<b>Hispanic, female</b>												
2003.....	38	-	-	-	-	1	1	1	-	1	2	4
2002.....	24	1	1	-	-	-	-	-	1	-	-	2
2001.....	22	-	-	1	-	-	-	-	1	-	2	1
2000.....	31	-	1	1	-	-	-	1	1	2	2	2
1999.....	21	-	-	-	-	1	1	-	1	2	2	1
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	1,163	-	-	-	-	2	11	16	13	39	63	79
2002.....	1,082	1	1	1	1	2	5	14	23	24	61	74
2001.....	1,117	1	-	-	2	3	4	8	15	42	66	87
2000.....	1,096	3	-	1	1	5	8	11	23	34	68	85
1999.....	1,049	2	-	-	-	1	4	10	13	50	54	64
<b>Non-Hispanic 1/, male</b>												
2003.....	638	-	-	-	-	-	5	10	6	19	37	54
2002.....	596	1	1	-	1	2	4	7	18	17	38	48
2001.....	631	1	-	-	1	2	2	4	8	21	34	63
2000.....	625	2	-	1	-	1	6	6	14	20	45	57
1999.....	594	2	-	-	-	-	3	7	11	32	37	46

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Acute and subacute endocarditis (I33)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	525	-	-	-	-	2	6	6	7	20	26	25
2002.....	486	-	-	1	-	-	1	7	5	7	23	26
2001.....	486	-	-	-	1	1	2	4	7	21	32	24
2000.....	471	1	-	-	1	4	2	5	9	14	23	28
1999.....	455	-	-	-	-	1	1	3	2	18	17	18
<b>Non-Hispanic White, both sexes</b>												
2003.....	912	-	-	-	-	1	10	13	9	23	45	48
2002.....	862	-	1	-	-	2	4	9	18	19	43	39
2001.....	859	1	-	-	2	1	4	7	8	26	39	59
2000.....	852	1	-	-	1	1	5	8	13	20	46	53
1999.....	814	1	-	-	-	1	3	8	9	28	31	38
<b>Non-Hispanic White, male</b>												
2003.....	500	-	-	-	-	-	5	8	3	11	28	31
2002.....	481	-	1	-	-	2	3	5	15	13	28	27
2001.....	482	1	-	-	1	1	2	3	4	12	22	41
2000.....	496	1	-	-	-	-	3	5	10	10	32	37
1999.....	466	1	-	-	-	-	2	5	7	19	20	30
<b>Non-Hispanic White, female</b>												
2003.....	412	-	-	-	-	1	5	5	6	12	17	17
2002.....	381	-	-	-	-	-	1	4	3	6	15	12
2001.....	377	-	-	-	1	-	2	4	4	14	17	18
2000.....	356	-	-	-	1	1	2	3	3	10	14	16
1999.....	348	-	-	-	-	1	1	3	2	9	11	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Acute and subacute endocarditis (I33)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	222	-	-	-	-	1	1	2	3	15	17	28
2002.....	197	-	-	1	-	-	1	5	5	5	17	29
2001.....	233	-	-	-	-	1	-	1	7	15	26	27
2000.....	213	2	-	-	-	3	3	3	8	11	20	30
1999.....	208	-	-	-	-	-	1	2	4	20	21	25
<b>Non-Hispanic Black, male</b>												
2003.....	119	-	-	-	-	-	-	2	2	7	8	20
2002.....	104	-	-	-	-	-	1	2	3	4	10	18
2001.....	134	-	-	-	-	-	-	1	4	8	12	21
2000.....	111	1	-	-	-	1	3	1	2	9	12	20
1999.....	115	-	-	-	-	-	1	2	4	12	15	15
<b>Non-Hispanic Black, female</b>												
2003.....	103	-	-	-	-	1	1	-	1	8	9	8
2002.....	93	-	-	1	-	-	-	3	2	1	7	11
2001.....	99	-	-	-	-	1	-	-	3	7	14	6
2000.....	102	1	-	-	-	2	-	2	6	2	8	10
1999.....	93	-	-	-	-	-	-	-	-	8	6	10
<b>Origin not stated 2/, both sexes</b>												
2003.....	2	-	-	-	-	-	-	-	-	-	-	1
2002.....	3	-	-	-	-	-	-	-	-	-	-	1
2001.....	5	-	-	-	-	-	-	-	-	-	1	1
2000.....	6	-	-	-	-	-	-	-	-	2	1	-
1999.....	4	-	-	-	-	-	-	1	-	-	-	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All origins, both sexes												
2003.....	868	33	22	14	14	21	34	23	25	32	54	63
2002.....	848	26	22	12	10	17	19	16	30	35	40	56
2001.....	839	32	26	12	9	14	26	20	30	43	50	57
2000.....	767	20	24	9	11	17	21	29	24	45	41	45
1999.....	890	27	25	11	10	12	20	19	31	54	60	59
All origins, male												
2003.....	449	15	8	4	8	14	20	13	16	21	31	41
2002.....	434	9	11	4	3	12	12	9	23	17	32	40
2001.....	401	18	15	2	5	7	15	16	17	24	17	30
2000.....	404	10	13	4	5	10	11	17	12	30	28	28
1999.....	442	13	14	5	2	8	12	14	20	32	32	31
All origins, female												
2003.....	419	18	14	10	6	7	14	10	9	11	23	22
2002.....	414	17	11	8	7	5	7	7	7	18	8	16
2001.....	438	14	11	10	4	7	11	4	13	19	33	27
2000.....	363	10	11	5	6	7	10	12	12	15	13	17
1999.....	448	14	11	6	8	4	8	5	11	22	28	28
Hispanic, both sexes												
2003.....	57	5	5	1	3	2	5	3	3	3	2	6
2002.....	64	4	2	2	1	1	4	-	3	4	2	6
2001.....	58	2	4	1	1	1	3	5	6	1	4	3
2000.....	61	5	4	1	-	3	2	6	3	5	5	2
1999.....	64	2	3	2	2	3	7	2	6	7	2	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hispanic, male												
2003.....	33	3	1	1	1	2	2	1	3	3	-	5
2002.....	35	3	-	-	-	1	2	-	3	2	1	4
2001.....	25	1	2	-	1	-	2	5	1	1	2	1
2000.....	29	2	4	-	-	-	1	2	2	3	3	1
1999.....	30	1	1	2	2	3	2	2	4	5	2	1
Hispanic, female												
2003.....	24	2	4	-	2	-	3	2	-	-	2	1
2002.....	29	1	2	2	1	-	2	-	-	2	1	2
2001.....	33	1	2	1	-	1	1	-	5	-	2	2
2000.....	32	3	-	1	-	3	1	4	1	2	2	1
1999.....	34	1	2	-	-	-	5	-	2	2	-	7
Non-Hispanic 1/, both sexes												
2003.....	806	28	17	13	11	19	29	19	22	29	52	57
2002.....	783	22	20	10	9	16	15	16	27	31	38	50
2001.....	779	30	22	11	8	13	23	14	24	41	46	54
2000.....	700	15	19	8	11	14	18	23	21	40	36	43
1999.....	824	25	22	9	8	9	13	17	24	47	57	51
Non-Hispanic 1/, male												
2003.....	414	12	7	3	7	12	18	12	13	18	31	36
2002.....	398	6	11	4	3	11	10	9	20	15	31	36
2001.....	375	17	13	2	4	7	13	10	16	23	15	29
2000.....	374	8	9	4	5	10	10	15	10	27	25	27
1999.....	410	12	13	3	-	5	10	12	15	27	29	30

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic 1/, female												
2003.....	392	16	10	10	4	7	11	7	9	11	21	21
2002.....	385	16	9	6	6	5	5	7	7	16	7	14
2001.....	404	13	9	9	4	6	10	4	8	18	31	25
2000.....	326	7	10	4	6	4	8	8	11	13	11	16
1999.....	414	13	9	6	8	4	3	5	9	20	28	21
Non-Hispanic White, both sexes												
2003.....	616	10	8	8	5	15	16	7	11	14	36	40
2002.....	601	9	11	6	3	9	12	14	16	18	31	37
2001.....	600	13	10	8	6	9	14	9	17	26	29	40
2000.....	528	6	11	7	2	12	8	15	10	23	26	24
1999.....	639	10	10	5	4	6	8	11	15	34	36	39
Non-Hispanic White, male												
2003.....	324	6	3	1	4	11	10	4	5	10	23	26
2002.....	308	3	7	2	1	6	8	8	13	7	26	25
2001.....	286	10	6	2	3	3	7	6	12	14	13	20
2000.....	280	4	7	4	1	8	4	9	5	16	18	16
1999.....	320	6	4	2	-	4	6	8	9	19	21	23
Non-Hispanic White, female												
2003.....	292	4	5	7	1	4	6	3	6	4	13	14
2002.....	293	6	4	4	2	3	4	6	3	11	5	12
2001.....	314	3	4	6	3	6	7	3	5	12	16	20
2000.....	248	2	4	3	1	4	4	6	5	7	8	8
1999.....	319	4	6	3	4	2	2	3	6	15	15	16

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Non-Hispanic Black, both sexes												
2003.....	168	14	9	4	5	3	12	12	10	14	16	16
2002.....	154	12	7	2	6	5	3	1	9	10	7	12
2001.....	157	14	12	3	1	4	9	5	5	10	14	14
2000.....	140	7	7	1	7	1	6	6	10	14	10	14
1999.....	158	10	11	4	4	3	3	6	8	12	18	9
Non-Hispanic Black, male												
2003.....	81	4	4	2	3	1	8	8	7	8	8	9
2002.....	75	3	3	1	2	3	2	-	6	5	5	10
2001.....	78	5	7	-	1	4	6	4	3	5	2	9
2000.....	72	4	2	-	3	1	2	5	5	9	7	7
1999.....	76	4	9	1	-	1	2	4	6	7	6	5
Non-Hispanic Black, female												
2003.....	87	10	5	2	2	2	4	4	3	6	8	7
2002.....	79	9	4	1	4	2	1	1	3	5	2	2
2001.....	79	9	5	3	-	-	3	1	2	5	12	5
2000.....	68	3	5	1	4	-	4	1	5	5	3	7
1999.....	82	6	2	3	4	2	1	2	2	5	12	4
Origin not stated 2/, both sexes												
2003.....	5	-	-	-	-	-	-	1	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	1	-	1	-	-
2000.....	6	-	1	-	-	-	1	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	1	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Heart failure (I50)</b>												
<b>All origins, both sexes</b>												
2003.....	57,448	38	5	5	7	18	20	28	69	91	224	358
2002.....	56,494	23	11	3	7	10	17	33	68	102	180	355
2001.....	56,934	24	17	8	7	19	15	41	51	115	192	353
2000.....	55,704	31	9	10	11	21	13	27	36	77	185	283
1999.....	54,913	11	10	5	7	14	17	27	38	88	184	298
<b>All origins, male</b>												
2003.....	22,427	28	3	4	3	13	11	16	44	50	143	223
2002.....	21,698	14	6	1	4	6	9	22	38	56	117	210
2001.....	21,632	16	5	5	3	11	5	25	31	76	106	227
2000.....	21,175	22	7	4	5	11	9	15	21	40	119	191
1999.....	20,755	5	7	3	3	12	11	13	19	52	112	186
<b>All origins, female</b>												
2003.....	35,021	10	2	1	4	5	9	12	25	41	81	135
2002.....	34,796	9	5	2	3	4	8	11	30	46	63	145
2001.....	35,302	8	12	3	4	8	10	16	20	39	86	126
2000.....	34,529	9	2	6	6	10	4	12	15	37	66	92
1999.....	34,158	6	3	2	4	2	6	14	19	36	72	112
<b>Hispanic, both sexes</b>												
2003.....	1,606	7	-	-	1	3	1	4	6	2	10	22
2002.....	1,412	4	-	-	2	-	-	2	1	4	9	14
2001.....	1,364	8	3	-	-	1	-	3	6	11	10	18
2000.....	1,270	7	1	2	-	-	-	1	2	4	6	7
1999.....	1,242	3	4	-	1	1	2	2	2	7	11	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Hispanic, male												
2003.....	714	5	-	-	-	3	1	3	4	-	10	15
2002.....	593	2	-	-	2	-	-	2	-	-	6	4
2001.....	565	6	1	-	-	1	-	2	4	9	6	15
2000.....	535	5	1	1	-	-	-	1	1	1	5	3
1999.....	522	1	3	-	-	1	2	2	1	5	7	6
Hispanic, female												
2003.....	892	2	-	-	1	-	-	1	2	2	-	7
2002.....	819	2	-	-	-	-	-	-	1	4	3	10
2001.....	799	2	2	-	-	-	-	1	2	2	4	3
2000.....	735	2	-	1	-	-	-	-	1	3	1	4
1999.....	720	2	1	-	1	-	-	-	1	2	4	4
Non-Hispanic 1/, both sexes												
2003.....	55,732	31	5	5	6	15	19	24	63	88	214	336
2002.....	54,932	19	11	3	5	10	17	31	66	97	171	339
2001.....	55,437	16	14	8	7	18	15	37	45	104	182	331
2000.....	54,307	24	8	8	11	21	13	26	34	73	179	276
1999.....	53,528	8	6	5	6	13	15	25	36	80	173	286
Non-Hispanic 1/, male												
2003.....	21,658	23	3	4	3	10	10	13	40	49	133	208
2002.....	21,050	12	6	1	2	6	9	20	38	55	111	204
2001.....	21,012	10	4	5	3	10	5	22	27	67	100	210
2000.....	20,599	17	6	3	5	11	9	14	20	39	114	188
1999.....	20,170	4	4	3	3	11	9	11	18	47	105	179

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Heart failure (I50)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	34,074	8	2	1	3	5	9	11	23	39	81	128
2002.....	33,882	7	5	2	3	4	8	11	28	42	60	135
2001.....	34,425	6	10	3	4	8	10	15	18	37	82	121
2000.....	33,708	7	2	5	6	10	4	12	14	34	65	88
1999.....	33,358	4	2	2	3	2	6	14	18	33	68	107
<b>Non-Hispanic White, both sexes</b>												
2003.....	49,788	13	4	3	3	6	11	15	33	53	112	207
2002.....	49,162	11	8	3	2	5	11	14	35	56	104	196
2001.....	49,788	11	12	5	6	12	5	22	27	65	102	215
2000.....	48,782	13	4	5	9	12	5	14	21	40	114	173
1999.....	47,977	5	4	3	5	8	7	11	19	50	106	173
<b>Non-Hispanic White, male</b>												
2003.....	19,239	8	2	2	2	2	7	10	21	29	76	136
2002.....	18,705	8	5	1	1	2	5	9	19	33	72	125
2001.....	18,730	6	4	3	2	8	1	10	15	40	63	149
2000.....	18,386	10	4	2	4	5	5	8	12	24	79	124
1999.....	17,903	2	2	2	3	7	4	4	10	29	71	110
<b>Non-Hispanic White, female</b>												
2003.....	30,549	5	2	1	1	4	4	5	12	24	36	71
2002.....	30,457	3	3	2	1	3	6	5	16	23	32	71
2001.....	31,058	5	8	2	4	4	4	12	12	25	39	66
2000.....	30,396	3	-	3	5	7	-	6	9	16	35	49
1999.....	30,074	3	2	1	2	1	3	7	9	21	35	63

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Heart failure (I50)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	5,294	15	-	2	3	9	8	8	29	34	97	124
2002.....	5,143	8	3	-	3	3	6	16	29	40	64	139
2001.....	5,094	5	2	3	-	6	10	15	17	35	77	111
2000.....	4,936	11	4	2	2	8	5	12	12	32	62	98
1999.....	4,961	3	2	2	1	4	8	13	17	28	65	109
<b>Non-Hispanic Black, male</b>												
2003.....	2,132	13	-	2	1	8	3	2	18	20	53	68
2002.....	2,055	4	1	-	1	2	4	10	17	22	37	77
2001.....	2,036	4	-	2	-	2	4	12	11	26	35	57
2000.....	1,942	7	2	1	1	5	3	6	7	14	33	60
1999.....	2,005	2	2	1	-	3	5	7	8	16	34	67
<b>Non-Hispanic Black, female</b>												
2003.....	3,162	2	-	-	2	1	5	6	11	14	44	56
2002.....	3,088	4	2	-	2	1	2	6	12	18	27	62
2001.....	3,058	1	2	1	-	4	6	3	6	9	42	54
2000.....	2,994	4	2	1	1	3	2	6	5	18	29	38
1999.....	2,956	1	-	1	1	1	3	6	9	12	31	42
<b>Origin not stated 2/, both sexes</b>												
2003.....	110	-	-	-	-	-	-	-	-	1	-	-
2002.....	150	-	-	-	-	-	-	-	1	1	-	2
2001.....	133	-	-	-	-	-	-	1	-	-	-	4
2000.....	127	-	-	-	-	-	-	-	-	-	-	-
1999.....	143	-	-	-	-	-	-	-	-	1	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Origin not stated 2/, male												
2003.....	55	-	-	-	-	-	-	-	-	1	-	-
2002.....	55	-	-	-	-	-	-	-	-	1	-	2
2001.....	55	-	-	-	-	-	-	1	-	-	-	2
2000.....	41	-	-	-	-	-	-	-	-	-	-	-
1999.....	63	-	-	-	-	-	-	-	-	-	-	1
Origin not stated 2/, female												
2003.....	55	-	-	-	-	-	-	-	-	-	-	-
2002.....	95	-	-	-	-	-	-	-	1	-	-	-
2001.....	78	-	-	-	-	-	-	-	-	-	-	2
2000.....	86	-	-	-	-	-	-	-	-	-	-	-
1999.....	80	-	-	-	-	-	-	-	-	1	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All origins, both sexes												
2003.....	110,423	344	144	71	122	306	489	623	1,009	1,507	2,535	3,381
2002.....	111,044	422	122	68	125	333	433	658	959	1,606	2,567	3,354
2001.....	107,998	392	172	71	136	269	443	617	948	1,566	2,389	3,205
2000.....	107,785	403	134	75	129	312	456	618	975	1,543	2,450	3,119
1999.....	108,909	450	141	83	132	370	440	622	1,012	1,683	2,464	3,077
All origins, male												
2003.....	51,180	189	70	42	79	192	304	364	609	924	1,569	2,115
2002.....	51,455	225	64	33	79	230	282	406	609	959	1,619	2,163
2001.....	49,738	187	97	34	73	173	284	367	564	974	1,518	2,036
2000.....	49,878	204	78	49	82	196	274	368	592	939	1,517	2,000
1999.....	50,765	235	78	46	78	230	257	364	613	1,058	1,533	1,957
All origins, female												
2003.....	59,243	155	74	29	43	114	185	259	400	583	966	1,266
2002.....	59,589	197	58	35	46	103	151	252	350	647	948	1,191
2001.....	58,260	205	75	37	63	96	159	250	384	592	871	1,169
2000.....	57,907	199	56	26	47	116	182	250	383	604	933	1,119
1999.....	58,144	215	63	37	54	140	183	258	399	625	931	1,120
Hispanic, both sexes												
2003.....	3,869	59	28	5	15	32	52	79	110	128	156	185
2002.....	3,607	72	20	9	13	33	64	81	94	114	160	162
2001.....	3,421	65	24	15	15	37	43	64	89	107	127	145
2000.....	3,391	67	22	13	20	43	62	70	94	115	137	138
1999.....	3,318	87	25	14	15	40	44	58	100	129	143	132

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Hispanic, male												
2003.....	2,016	31	16	2	10	18	41	51	70	89	108	128
2002.....	1,914	34	11	5	5	24	49	60	64	73	108	102
2001.....	1,777	33	11	9	7	21	32	43	59	69	92	100
2000.....	1,790	32	15	8	16	34	41	51	63	72	86	89
1999.....	1,798	38	15	7	10	21	31	44	70	83	100	88
Hispanic, female												
2003.....	1,853	28	12	3	5	14	11	28	40	39	48	57
2002.....	1,693	38	9	4	8	9	15	21	30	41	52	60
2001.....	1,644	32	13	6	8	16	11	21	30	38	35	45
2000.....	1,601	35	7	5	4	9	21	19	31	43	51	49
1999.....	1,520	49	10	7	5	19	13	14	30	46	43	44
Non-Hispanic 1/, both sexes												
2003.....	106,297	283	115	65	107	270	437	543	894	1,376	2,364	3,183
2002.....	107,173	348	102	59	112	300	368	575	860	1,484	2,394	3,183
2001.....	104,292	325	147	56	121	230	395	551	856	1,453	2,250	3,044
2000.....	104,119	333	112	61	109	269	393	544	876	1,422	2,299	2,963
1999.....	105,310	362	115	69	116	329	395	561	908	1,546	2,310	2,936
Non-Hispanic 1/, male												
2003.....	49,021	158	54	39	69	172	263	313	535	833	1,449	1,975
2002.....	49,411	190	53	28	74	206	232	346	541	880	1,502	2,055
2001.....	47,812	153	86	25	66	150	250	323	503	899	1,417	1,925
2000.....	47,949	170	63	41	66	162	232	314	528	862	1,422	1,900
1999.....	48,810	197	63	39	67	208	225	319	540	969	1,426	1,863

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic 1/, female												
2003.....	57,276	125	61	26	38	98	174	230	359	543	915	1,208
2002.....	57,762	158	49	31	38	94	136	229	319	604	892	1,128
2001.....	56,480	172	61	31	55	80	145	228	353	554	833	1,119
2000.....	56,170	163	49	20	43	107	161	230	348	560	877	1,063
1999.....	56,500	165	52	30	49	121	170	242	368	577	884	1,073
Non-Hispanic White, both sexes												
2003.....	90,097	163	65	39	74	175	261	307	546	880	1,605	2,218
2002.....	91,087	214	66	41	75	200	210	324	506	952	1,665	2,181
2001.....	88,469	194	81	35	78	142	235	325	522	922	1,487	2,068
2000.....	88,583	218	72	34	64	159	226	317	550	905	1,543	2,035
1999.....	89,439	224	73	48	75	220	242	330	569	991	1,538	1,988
Non-Hispanic White, male												
2003.....	41,056	88	27	24	49	112	151	177	348	541	1,019	1,415
2002.....	41,554	117	38	20	49	139	139	195	325	576	1,081	1,446
2001.....	40,015	99	48	16	45	90	149	197	308	584	983	1,352
2000.....	40,299	112	40	24	42	93	139	188	339	567	980	1,362
1999.....	40,860	127	45	26	40	133	135	192	345	641	989	1,319
Non-Hispanic White, female												
2003.....	49,041	75	38	15	25	63	110	130	198	339	586	803
2002.....	49,533	97	28	21	26	61	71	129	181	376	584	735
2001.....	48,454	95	33	19	33	52	86	128	214	338	504	716
2000.....	48,284	106	32	10	22	66	87	129	211	338	563	673
1999.....	48,579	97	28	22	35	87	107	138	224	350	549	669

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Non-Hispanic Black, both sexes												
2003.....	14,177	96	40	23	29	81	158	214	308	451	702	899
2002.....	14,147	115	28	13	32	87	141	228	315	490	673	925
2001.....	14,012	111	56	19	37	80	142	207	310	477	716	898
2000.....	13,870	93	35	24	36	100	152	212	296	467	698	871
1999.....	14,204	119	35	20	35	102	141	211	315	512	719	873
Non-Hispanic Black, male												
2003.....	6,943	59	23	12	18	53	102	124	159	265	398	522
2002.....	6,804	65	12	5	21	56	83	137	188	277	378	562
2001.....	6,822	50	32	8	20	54	90	116	176	279	401	518
2000.....	6,758	44	20	14	20	60	83	116	171	266	397	503
1999.....	7,062	60	16	12	24	72	82	119	181	300	407	502
Non-Hispanic Black, female												
2003.....	7,234	37	17	11	11	28	56	90	149	186	304	377
2002.....	7,343	50	16	8	11	31	58	91	127	213	295	363
2001.....	7,190	61	24	11	17	26	52	91	134	198	315	380
2000.....	7,112	49	15	10	16	40	69	96	125	201	301	368
1999.....	7,142	59	19	8	11	30	59	92	134	212	312	371
Origin not stated 2/, both sexes												
2003.....	257	2	1	1	-	4	-	1	5	3	15	13
2002.....	264	2	-	-	-	-	1	2	5	8	13	9
2001.....	285	2	1	-	-	2	5	2	3	6	12	16
2000.....	275	3	-	1	-	-	1	4	5	6	14	18
1999.....	281	1	1	-	1	1	1	3	4	8	11	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Origin not stated 2/, male												
2003.....	143	-	-	1	-	2	-	-	4	2	12	12
2002.....	130	1	-	-	-	-	1	-	4	6	9	6
2001.....	149	1	-	-	-	2	2	1	2	6	9	11
2000.....	139	2	-	-	-	-	1	3	1	5	9	11
1999.....	157	-	-	-	1	1	1	1	3	6	7	6
Origin not stated 2/, female												
2003.....	114	2	1	-	-	2	-	1	1	1	3	1
2002.....	134	1	-	-	-	-	-	2	1	2	4	3
2001.....	136	1	1	-	-	-	3	1	1	-	3	5
2000.....	136	1	-	1	-	-	-	1	4	1	5	7
1999.....	124	1	1	-	-	-	-	2	1	2	4	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Essential (primary) hypertension and hypertensive renal disease (I10,I12)</b>												
<b>All origins, both sexes</b>												
2003.....	21,940	1	2	-	2	7	15	27	71	126	246	400
2002.....	20,261	5	1	1	-	4	17	36	49	126	228	382
2001.....	19,250	-	1	-	1	5	18	32	72	101	230	385
2000.....	18,073	1	1	2	2	6	11	21	49	105	243	352
1999.....	16,968	3	3	-	1	3	10	24	54	105	208	329
<b>All origins, male</b>												
2003.....	8,213	1	2	-	1	5	7	9	49	72	143	265
2002.....	7,647	2	1	1	-	-	11	21	26	75	144	238
2001.....	7,254	-	-	-	-	2	14	20	39	59	147	239
2000.....	6,895	1	-	1	2	-	4	12	27	61	155	224
1999.....	6,488	2	1	-	1	2	6	13	35	69	126	196
<b>All origins, female</b>												
2003.....	13,727	-	-	-	1	2	8	18	22	54	103	135
2002.....	12,614	3	-	-	-	4	6	15	23	51	84	144
2001.....	11,996	-	1	-	1	3	4	12	33	42	83	146
2000.....	11,178	-	1	1	-	6	7	9	22	44	88	128
1999.....	10,480	1	2	-	-	1	4	11	19	36	82	133
<b>Hispanic, both sexes</b>												
2003.....	1,053	-	-	-	-	2	2	4	7	6	17	37
2002.....	942	2	-	-	-	2	3	4	5	12	18	17
2001.....	889	-	1	-	-	1	2	1	9	5	14	28
2000.....	821	1	1	-	-	1	3	2	5	11	16	18
1999.....	753	1	1	-	-	1	2	4	4	3	16	29

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Essential (primary) hypertension and hypertensive renal disease (I10,I12)</b>												
<b>Hispanic, male</b>												
2003.....	437	-	-	-	-	1	1	2	3	4	13	26
2002.....	385	1	-	-	-	-	1	3	2	8	10	13
2001.....	365	-	-	-	-	1	1	1	7	4	12	21
2000.....	386	1	-	-	-	-	2	1	4	8	12	16
1999.....	340	-	-	-	-	1	1	2	3	3	11	16
<b>Hispanic, female</b>												
2003.....	616	-	-	-	-	1	1	2	4	2	4	11
2002.....	557	1	-	-	-	2	2	1	3	4	8	4
2001.....	524	-	-	-	-	-	1	-	2	1	2	7
2000.....	435	-	1	-	-	1	1	1	1	3	4	2
1999.....	413	1	1	-	-	-	1	2	1	-	5	13
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	20,831	1	2	-	2	5	13	23	62	120	229	362
2002.....	19,254	3	1	1	-	2	14	31	44	114	208	364
2001.....	18,306	-	1	-	1	4	16	31	63	96	214	354
2000.....	17,182	-	-	2	2	5	8	19	44	93	225	328
1999.....	16,158	2	2	-	1	2	8	20	49	101	191	299
<b>Non-Hispanic 1/, male</b>												
2003.....	7,751	1	2	-	1	4	6	7	44	68	130	238
2002.....	7,232	1	1	1	-	-	10	18	24	67	133	225
2001.....	6,856	-	-	-	-	1	13	19	32	55	133	216
2000.....	6,475	-	-	1	2	-	2	11	23	52	142	203
1999.....	6,123	2	1	-	1	1	5	11	31	65	114	180

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic 1/, female

2003.....	13,080	-	-	-	1	1	7	16	18	52	99	124
2002.....	12,022	2	-	-	-	2	4	13	20	47	75	139
2001.....	11,450	-	1	-	1	3	3	12	31	41	81	138
2000.....	10,707	-	-	1	-	5	6	8	21	41	83	125
1999.....	10,035	-	1	-	-	1	3	9	18	36	77	119

Non-Hispanic White, both sexes

2003.....	15,878	-	1	-	1	3	5	8	24	52	113	173
2002.....	14,617	2	1	1	-	1	8	11	15	33	93	168
2001.....	13,837	-	1	-	1	3	8	12	23	40	81	155
2000.....	12,917	-	-	1	2	1	2	10	15	30	102	149
1999.....	12,048	1	-	-	-	2	3	8	20	34	78	143

Non-Hispanic White, male

2003.....	5,673	-	1	-	-	2	3	2	14	35	67	120
2002.....	5,325	1	1	1	-	-	7	7	9	17	65	110
2001.....	4,970	-	-	-	-	1	6	7	13	23	54	98
2000.....	4,721	-	-	1	2	-	1	7	10	22	67	100
1999.....	4,395	1	-	-	-	1	1	5	17	26	51	93

Non-Hispanic White, female

2003.....	10,205	-	-	-	1	1	2	6	10	17	46	53
2002.....	9,292	1	-	-	-	1	1	4	6	16	28	58
2001.....	8,867	-	1	-	1	2	2	5	10	17	27	57
2000.....	8,196	-	-	-	-	1	1	3	5	8	35	49
1999.....	7,653	-	-	-	-	1	2	3	3	8	27	50

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic Black, both sexes

2003.....	4,445	1	1	-	-	2	8	15	36	63	111	183
2002.....	4,165	1	-	-	-	1	6	17	27	79	110	188
2001.....	4,012	-	-	-	-	1	8	18	37	53	124	188
2000.....	3,862	-	-	1	-	4	6	7	28	61	115	168
1999.....	3,708	1	1	-	1	-	5	11	26	65	110	144

Non-Hispanic Black, male

2003.....	1,875	1	1	-	-	2	3	5	28	31	61	117
2002.....	1,701	-	-	-	-	-	3	10	14	48	64	109
2001.....	1,698	-	-	-	-	-	7	12	17	30	72	113
2000.....	1,576	-	-	-	-	-	1	3	13	29	69	96
1999.....	1,550	1	1	-	1	-	4	5	12	37	60	83

Non-Hispanic Black, female

2003.....	2,570	-	-	-	-	-	5	10	8	32	50	66
2002.....	2,464	1	-	-	-	1	3	7	13	31	46	79
2001.....	2,314	-	-	-	-	1	1	6	20	23	52	75
2000.....	2,286	-	-	1	-	4	5	4	15	32	46	72
1999.....	2,158	-	-	-	-	-	1	6	14	28	50	61

Origin not stated 2/, both sexes

2003.....	56	-	-	-	-	-	-	-	2	-	-	1
2002.....	65	-	-	-	-	-	-	1	-	-	2	1
2001.....	55	-	-	-	-	-	-	-	-	-	2	3
2000.....	70	-	-	-	-	-	-	-	-	1	2	6
1999.....	57	-	-	-	-	-	-	-	1	1	1	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Cerebrovascular diseases (I60-I69)</b>												
All origins, both sexes												
2003.....	157,689	101	46	29	40	88	133	188	395	761	1,699	2,577
2002.....	162,672	117	53	33	58	53	118	183	384	797	1,628	2,615
2001.....	163,538	108	54	38	42	68	128	199	402	840	1,651	2,507
2000.....	167,661	124	47	25	51	67	132	209	393	880	1,719	2,625
1999.....	167,366	103	46	26	39	67	115	202	378	916	1,658	2,455
All origins, male												
2003.....	61,426	57	25	19	22	51	65	101	213	411	872	1,389
2002.....	62,622	67	30	16	31	34	69	100	184	389	789	1,402
2001.....	63,177	63	25	21	21	36	67	105	206	435	832	1,354
2000.....	64,769	74	19	12	21	38	68	111	197	430	867	1,401
1999.....	64,485	65	22	17	17	36	59	102	197	449	859	1,282
All origins, female												
2003.....	96,263	44	21	10	18	37	68	87	182	350	827	1,188
2002.....	100,050	50	23	17	27	19	49	83	200	408	839	1,213
2001.....	100,361	45	29	17	21	32	61	94	196	405	819	1,153
2000.....	102,892	50	28	13	30	29	64	98	196	450	852	1,224
1999.....	102,881	38	24	9	22	31	56	100	181	467	799	1,173
Hispanic, both sexes												
2003.....	6,658	17	8	5	9	9	19	35	54	80	187	251
2002.....	6,451	20	12	5	15	10	19	40	53	94	162	269
2001.....	6,416	17	10	7	9	13	27	34	65	102	173	239
2000.....	6,187	16	6	4	4	5	26	37	71	95	172	245
1999.....	5,907	16	7	4	5	10	22	46	56	89	152	229

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Hispanic, male</b>												
2003.....	3,070	11	7	3	5	7	8	18	33	48	114	156
2002.....	3,003	9	5	1	6	6	11	29	32	57	97	169
2001.....	2,982	9	4	5	5	8	18	20	40	58	93	141
2000.....	2,865	8	-	1	2	2	15	22	43	53	92	144
1999.....	2,809	8	1	3	3	7	14	25	37	49	97	136
<b>Hispanic, female</b>												
2003.....	3,588	6	1	2	4	2	11	17	21	32	73	95
2002.....	3,448	11	7	4	9	4	8	11	21	37	65	100
2001.....	3,434	8	6	2	4	5	9	14	25	44	80	98
2000.....	3,322	8	6	3	2	3	11	15	28	42	80	101
1999.....	3,098	8	6	1	2	3	8	21	19	40	55	93
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	150,719	82	38	24	31	79	113	151	340	676	1,506	2,318
2002.....	155,852	96	41	28	43	43	99	143	331	699	1,462	2,337
2001.....	156,741	91	43	30	33	55	101	163	335	735	1,476	2,258
2000.....	161,057	108	41	21	46	62	105	172	321	783	1,540	2,366
1999.....	161,057	87	39	22	34	57	93	156	321	826	1,497	2,213
<b>Non-Hispanic 1/, male</b>												
2003.....	58,226	45	18	16	17	44	56	82	179	359	756	1,228
2002.....	59,470	57	25	15	25	28	58	71	152	330	689	1,227
2001.....	60,030	54	21	16	16	28	49	83	164	374	737	1,209
2000.....	61,715	66	19	11	19	36	53	89	154	376	769	1,248
1999.....	61,499	57	21	14	14	29	45	77	160	400	756	1,139

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	92,493	37	20	8	14	35	57	69	161	317	750	1,090
2002.....	96,382	39	16	13	18	15	41	72	179	369	773	1,110
2001.....	96,711	37	22	14	17	27	52	80	171	361	739	1,049
2000.....	99,342	42	22	10	27	26	52	83	167	407	771	1,118
1999.....	99,558	30	18	8	20	28	48	79	161	426	741	1,074
<b>Non-Hispanic White, both sexes</b>												
2003.....	127,969	46	23	16	21	54	70	90	208	375	864	1,364
2002.....	133,118	51	29	20	24	28	73	85	200	412	876	1,407
2001.....	133,879	46	28	21	21	40	58	101	219	452	881	1,363
2000.....	138,210	59	25	17	29	46	72	123	190	464	914	1,380
1999.....	138,716	55	27	19	21	37	68	100	201	514	892	1,302
<b>Non-Hispanic White, male</b>												
2003.....	48,549	22	8	10	13	29	43	48	111	195	442	747
2002.....	49,908	32	18	10	13	17	45	47	89	195	422	765
2001.....	50,382	27	14	12	12	20	35	49	112	223	461	739
2000.....	52,000	38	14	11	12	31	36	67	94	229	465	752
1999.....	51,968	36	15	13	8	21	30	48	103	252	460	664
<b>Non-Hispanic White, female</b>												
2003.....	79,420	24	15	6	8	25	27	42	97	180	422	617
2002.....	83,210	19	11	10	11	11	28	38	111	217	454	642
2001.....	83,497	19	14	9	9	20	23	52	107	229	420	624
2000.....	86,210	21	11	6	17	15	36	56	96	235	449	628
1999.....	86,748	19	12	6	13	16	38	52	98	262	432	638

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Cerebrovascular diseases (I60-I69)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	18,649	31	14	6	10	21	37	53	114	248	538	820
2002.....	18,691	36	10	7	16	13	25	50	116	237	504	807
2001.....	18,844	42	12	6	8	13	38	57	96	233	525	790
2000.....	19,053	46	13	2	13	13	29	41	117	281	548	868
1999.....	18,731	27	10	3	12	18	22	49	100	267	541	780
<b>Non-Hispanic Black, male</b>												
2003.....	7,798	19	9	5	4	13	12	32	54	129	256	415
2002.....	7,754	19	6	4	11	10	12	20	52	102	229	398
2001.....	7,828	26	5	2	3	8	12	33	42	118	237	412
2000.....	7,945	27	3	-	5	4	15	18	52	127	267	440
1999.....	7,823	17	4	1	6	7	12	24	48	123	264	409
<b>Non-Hispanic Black, female</b>												
2003.....	10,851	12	5	1	6	8	25	21	60	119	282	405
2002.....	10,937	17	4	3	5	3	13	30	64	135	275	409
2001.....	11,016	16	7	4	5	5	26	24	54	115	288	378
2000.....	11,108	19	10	2	8	9	14	23	65	154	281	428
1999.....	10,908	10	6	2	6	11	10	25	52	144	277	371
<b>Origin not stated 2/, both sexes</b>												
2003.....	312	2	-	-	-	-	1	2	1	5	6	8
2002.....	369	1	-	-	-	-	-	-	-	4	4	9
2001.....	381	-	1	1	-	-	-	2	2	3	2	10
2000.....	417	-	-	-	1	-	1	-	1	2	7	14
1999.....	402	-	-	-	-	-	-	-	1	1	9	13

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

Origin not stated 2/, male

2003.....	130	1	-	-	-	-	1	1	1	4	2	5
2002.....	149	1	-	-	-	-	-	-	-	2	3	6
2001.....	165	-	-	-	-	-	-	2	2	3	2	4
2000.....	189	-	-	-	-	-	-	-	-	1	6	9
1999.....	177	-	-	-	-	-	-	-	-	-	6	7

Origin not stated 2/, female

2003.....	182	1	-	-	-	-	-	1	-	1	4	3
2002.....	220	-	-	-	-	-	-	-	-	2	1	3
2001.....	216	-	1	1	-	-	-	-	-	-	-	6
2000.....	228	-	-	-	1	-	1	-	1	1	1	5
1999.....	225	-	-	-	-	-	-	-	1	1	3	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Atherosclerosis (I70)</b>												
All origins, both sexes												
2003.....	13,053	3	-	-	-	1	2	-	7	11	36	70
2002.....	13,821	4	-	-	-	-	3	2	7	16	34	61
2001.....	14,086	1	1	1	-	-	1	1	2	14	26	65
2000.....	14,393	2	-	-	-	-	3	3	5	16	32	71
1999.....	14,979	2	-	-	-	1	1	1	7	16	29	56
All origins, male												
2003.....	4,859	2	-	-	-	1	1	-	5	8	27	54
2002.....	5,171	1	-	-	-	-	2	2	3	9	21	44
2001.....	5,143	1	1	1	-	-	1	-	2	8	18	43
2000.....	5,357	-	-	-	-	-	2	1	5	10	25	48
1999.....	5,443	1	-	-	-	-	-	1	4	11	21	37
All origins, female												
2003.....	8,194	1	-	-	-	-	1	-	2	3	9	16
2002.....	8,650	3	-	-	-	-	1	-	4	7	13	17
2001.....	8,943	-	-	-	-	-	-	1	-	6	8	22
2000.....	9,036	2	-	-	-	-	1	2	-	6	7	23
1999.....	9,536	1	-	-	-	1	1	-	3	5	8	19
Hispanic, both sexes												
2003.....	379	1	-	-	-	-	-	-	1	-	3	1
2002.....	416	1	-	-	-	-	2	-	-	1	1	3
2001.....	416	-	-	-	-	-	-	-	-	-	-	3
2000.....	414	-	-	-	-	-	-	-	-	2	-	4
1999.....	410	-	-	-	-	-	-	-	-	1	2	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Atherosclerosis (I70)</b>												
Hispanic, male												
2003.....	182	-	-	-	-	-	-	-	-	-	3	1
2002.....	175	1	-	-	-	-	2	-	-	-	1	3
2001.....	172	-	-	-	-	-	-	-	-	-	-	1
2000.....	173	-	-	-	-	-	-	-	-	2	-	3
1999.....	185	-	-	-	-	-	-	-	-	-	1	2
Hispanic, female												
2003.....	197	1	-	-	-	-	-	-	1	-	-	-
2002.....	241	-	-	-	-	-	-	-	-	1	-	-
2001.....	244	-	-	-	-	-	-	-	-	-	-	2
2000.....	241	-	-	-	-	-	-	-	-	-	-	1
1999.....	225	-	-	-	-	-	-	-	-	1	1	3
Non-Hispanic 1/, both sexes												
2003.....	12,627	2	-	-	-	1	2	-	6	11	33	69
2002.....	13,369	3	-	-	-	-	1	2	7	15	33	58
2001.....	13,626	1	1	1	-	-	1	1	2	14	26	62
2000.....	13,945	2	-	-	-	-	3	3	5	14	32	67
1999.....	14,519	2	-	-	-	1	1	1	7	15	27	49
Non-Hispanic 1/, male												
2003.....	4,660	2	-	-	-	1	1	-	5	8	24	53
2002.....	4,984	-	-	-	-	-	-	2	3	9	20	41
2001.....	4,952	1	1	1	-	-	1	-	2	8	18	42
2000.....	5,166	-	-	-	-	-	2	1	5	8	25	45
1999.....	5,236	1	-	-	-	-	-	1	4	11	20	33

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

#### Atherosclerosis (I70)

##### Non-Hispanic 1/, female

2003.....	7,967	-	-	-	-	-	1	-	1	3	9	16
2002.....	8,385	3	-	-	-	-	1	-	4	6	13	17
2001.....	8,674	-	-	-	-	-	-	1	-	6	8	20
2000.....	8,779	2	-	-	-	-	1	2	-	6	7	22
1999.....	9,283	1	-	-	-	1	1	-	3	4	7	16

##### Non-Hispanic White, both sexes

2003.....	11,475	-	-	-	-	-	2	-	4	7	29	57
2002.....	12,191	2	-	-	-	-	1	2	4	13	25	42
2001.....	12,517	1	1	1	-	-	-	-	2	11	22	51
2000.....	12,740	2	-	-	-	-	3	1	4	10	24	46
1999.....	13,286	1	-	-	-	1	1	1	5	13	17	41

##### Non-Hispanic White, male

2003.....	4,181	-	-	-	-	-	1	-	4	5	20	42
2002.....	4,508	-	-	-	-	-	-	2	2	7	16	32
2001.....	4,511	1	1	1	-	-	-	-	2	5	16	34
2000.....	4,688	-	-	-	-	-	2	1	4	7	19	33
1999.....	4,727	1	-	-	-	-	-	1	3	10	13	28

##### Non-Hispanic White, female

2003.....	7,294	-	-	-	-	-	1	-	-	2	9	15
2002.....	7,683	2	-	-	-	-	1	-	2	6	9	10
2001.....	8,006	-	-	-	-	-	-	-	-	6	6	17
2000.....	8,052	2	-	-	-	-	1	-	-	3	5	13
1999.....	8,559	-	-	-	-	1	1	-	2	3	4	13





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of circulatory system (I71-I78)												
All origins, both sexes												
2003.....	24,672	11	-	1	11	30	38	62	89	167	284	524
2002.....	24,927	14	4	1	10	31	35	51	109	154	291	479
2001.....	25,318	12	4	2	9	23	38	67	103	194	304	421
2000.....	26,036	16	5	3	8	17	42	49	105	157	291	424
1999.....	25,809	12	2	2	9	14	35	51	90	178	300	418
All origins, male												
2003.....	13,217	3	-	-	7	17	28	43	60	117	189	343
2002.....	13,309	6	4	-	5	19	24	30	71	81	210	331
2001.....	13,396	8	2	-	3	17	23	43	72	132	203	291
2000.....	14,084	6	2	1	6	9	24	36	67	92	184	275
1999.....	13,794	9	-	1	6	10	22	35	67	117	179	289
All origins, female												
2003.....	11,455	8	-	1	4	13	10	19	29	50	95	181
2002.....	11,618	8	-	1	5	12	11	21	38	73	81	148
2001.....	11,922	4	2	2	6	6	15	24	31	62	101	130
2000.....	11,952	10	3	2	2	8	18	13	38	65	107	149
1999.....	12,015	3	2	1	3	4	13	16	23	61	121	129
Hispanic, both sexes												
2003.....	878	2	-	-	1	5	4	10	12	15	24	31
2002.....	847	1	-	-	1	6	6	6	16	15	29	29
2001.....	754	2	1	1	1	6	1	7	12	11	26	31
2000.....	818	3	1	-	1	5	4	11	13	18	20	19
1999.....	729	2	-	-	2	4	5	6	6	13	20	15

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of circulatory system (I71-I78)												
Hispanic, male												
2003.....	510	1	-	-	-	3	2	6	9	12	16	19
2002.....	463	-	-	-	-	4	4	4	10	9	22	19
2001.....	424	1	1	-	-	4	-	4	10	6	20	20
2000.....	452	1	-	-	1	3	1	7	7	7	15	9
1999.....	416	1	-	-	-	2	5	3	3	9	13	11
Hispanic, female												
2003.....	368	1	-	-	1	2	2	4	3	3	8	12
2002.....	384	1	-	-	1	2	2	2	6	6	7	10
2001.....	330	1	-	1	1	2	1	3	2	5	6	11
2000.....	366	2	1	-	-	2	3	4	6	11	5	10
1999.....	313	1	-	-	2	2	-	3	3	4	7	4
Non-Hispanic 1/, both sexes												
2003.....	23,739	9	-	1	10	25	34	52	77	151	259	492
2002.....	24,005	13	4	1	9	25	29	45	92	137	259	449
2001.....	24,489	10	3	1	8	17	37	59	90	183	277	389
2000.....	25,147	13	4	3	7	12	38	37	92	138	267	404
1999.....	25,023	10	2	2	7	10	30	45	84	165	280	403
Non-Hispanic 1/, male												
2003.....	12,679	2	-	-	7	14	26	37	51	104	173	323
2002.....	12,809	6	4	-	5	15	20	26	60	71	186	312
2001.....	12,926	7	1	-	3	13	23	38	61	126	182	270
2000.....	13,593	5	2	1	5	6	23	29	60	84	165	265
1999.....	13,351	8	-	1	6	8	17	32	64	108	166	278

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of circulatory system (I71-I78)												
Non-Hispanic 1/, female												
2003.....	11,060	7	-	1	3	11	8	15	26	47	86	169
2002.....	11,196	7	-	1	4	10	9	19	32	66	73	137
2001.....	11,563	3	2	1	5	4	14	21	29	57	95	119
2000.....	11,554	8	2	2	2	6	15	8	32	54	102	139
1999.....	11,672	2	2	1	1	2	13	13	20	57	114	125
Non-Hispanic White, both sexes												
2003.....	20,756	4	-	1	6	18	23	34	51	89	177	332
2002.....	21,051	10	4	1	3	20	17	29	57	98	174	301
2001.....	21,439	8	2	1	7	12	28	44	64	126	177	277
2000.....	22,113	11	3	2	4	9	26	24	64	99	170	284
1999.....	22,191	8	2	2	5	9	24	32	62	125	198	289
Non-Hispanic White, male												
2003.....	11,188	1	-	-	4	12	18	24	31	67	122	230
2002.....	11,337	5	4	-	2	12	11	18	36	53	127	220
2001.....	11,437	5	1	-	3	10	17	31	43	87	115	195
2000.....	12,138	3	2	1	4	4	18	17	46	59	108	199
1999.....	12,014	6	-	1	4	7	13	23	48	83	121	208
Non-Hispanic White, female												
2003.....	9,568	3	-	1	2	6	5	10	20	22	55	102
2002.....	9,714	5	-	1	1	8	6	11	21	45	47	81
2001.....	10,002	3	1	1	4	2	11	13	21	39	62	82
2000.....	9,975	8	1	1	-	5	8	7	18	40	62	85
1999.....	10,177	2	2	1	1	2	11	9	14	42	77	81





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Aortic aneurysm and dissection (I71)</b>												
<b>All origins, both sexes</b>												
2003.....	14,810	1	-	-	3	16	25	43	69	124	192	368
2002.....	14,818	2	-	1	4	20	19	30	80	104	201	327
2001.....	15,234	-	2	-	4	15	28	46	71	134	198	287
2000.....	15,810	-	-	-	5	10	26	38	69	103	200	268
1999.....	15,807	1	1	-	3	8	23	33	60	134	195	283
<b>All origins, male</b>												
2003.....	8,976	-	-	-	2	10	21	32	51	96	150	269
2002.....	9,013	-	-	-	3	17	15	25	57	67	162	249
2001.....	9,114	-	2	-	3	12	19	34	56	105	159	219
2000.....	9,661	-	-	-	3	7	16	30	48	72	147	191
1999.....	9,615	1	-	-	2	6	14	25	53	98	134	216
<b>All origins, female</b>												
2003.....	5,834	1	-	-	1	6	4	11	18	28	42	99
2002.....	5,805	2	-	1	1	3	4	5	23	37	39	78
2001.....	6,120	-	-	-	1	3	9	12	15	29	39	68
2000.....	6,149	-	-	-	2	3	10	8	21	31	53	77
1999.....	6,192	-	1	-	1	2	9	8	7	36	61	67
<b>Hispanic, both sexes</b>												
2003.....	515	-	-	-	-	4	2	5	11	9	16	18
2002.....	431	-	-	-	-	4	4	3	13	9	15	17
2001.....	403	-	1	-	-	3	1	4	10	8	16	18
2000.....	465	-	-	-	1	5	3	8	7	10	11	13
1999.....	384	-	-	-	1	2	2	3	3	10	13	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Aortic aneurysm and dissection (I71)</b>												
<b>Hispanic, male</b>												
2003.....	333	-	-	-	-	3	1	3	9	6	13	12
2002.....	284	-	-	-	-	3	2	3	9	8	12	15
2001.....	279	-	1	-	-	2	-	4	10	6	14	14
2000.....	294	-	-	-	1	3	1	6	5	6	8	8
1999.....	256	-	-	-	-	1	2	2	2	7	10	9
<b>Hispanic, female</b>												
2003.....	182	-	-	-	-	1	1	2	2	3	3	6
2002.....	147	-	-	-	-	1	2	-	4	1	3	2
2001.....	124	-	-	-	-	1	1	-	-	2	2	4
2000.....	171	-	-	-	-	2	2	2	2	4	3	5
1999.....	128	-	-	-	1	1	-	1	1	3	3	1
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	14,259	1	-	-	3	12	23	38	58	114	175	349
2002.....	14,341	2	-	1	4	16	15	27	66	93	184	309
2001.....	14,782	-	1	-	4	12	27	41	61	126	181	268
2000.....	15,296	-	-	-	4	5	23	29	62	92	185	254
1999.....	15,393	1	1	-	2	6	21	30	57	124	182	273
<b>Non-Hispanic 1/, male</b>												
2003.....	8,622	-	-	-	2	7	20	29	42	89	137	256
2002.....	8,701	-	-	-	3	14	13	22	47	58	149	234
2001.....	8,798	-	1	-	3	10	19	29	46	99	144	204
2000.....	9,338	-	-	-	2	4	15	24	43	65	135	182
1999.....	9,342	1	-	-	2	5	12	23	51	91	124	207

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Aortic aneurysm and dissection (I71)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	5,637	1	-	-	1	5	3	9	16	25	38	93
2002.....	5,640	2	-	1	1	2	2	5	19	35	35	75
2001.....	5,984	-	-	-	1	2	8	12	15	27	37	64
2000.....	5,958	-	-	-	2	1	8	5	19	27	50	72
1999.....	6,051	-	1	-	-	1	9	7	6	33	58	66
<b>Non-Hispanic White, both sexes</b>												
2003.....	12,712	-	-	-	1	10	17	23	38	68	120	233
2002.....	12,831	2	-	1	1	14	9	18	39	71	121	203
2001.....	13,217	-	1	-	4	8	20	32	45	88	115	189
2000.....	13,722	-	-	-	1	4	14	18	43	64	118	180
1999.....	13,934	1	1	-	1	6	17	22	42	95	122	184
<b>Non-Hispanic White, male</b>												
2003.....	7,731	-	-	-	-	7	14	17	25	58	97	176
2002.....	7,834	-	-	-	1	12	7	15	28	43	99	158
2001.....	7,924	-	1	-	3	7	14	24	34	71	92	148
2000.....	8,488	-	-	-	1	3	11	13	32	45	89	139
1999.....	8,551	1	-	-	1	5	9	17	37	70	85	145
<b>Non-Hispanic White, female</b>												
2003.....	4,981	-	-	-	1	3	3	6	13	10	23	57
2002.....	4,997	2	-	1	-	2	2	3	11	28	22	45
2001.....	5,293	-	-	-	1	1	6	8	11	17	23	41
2000.....	5,234	-	-	-	1	3	3	5	11	19	29	41
1999.....	5,383	-	1	-	-	1	8	5	5	25	37	39





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All origins, both sexes												
2003.....	9,862	10	-	1	8	14	13	19	20	43	92	156
2002.....	10,109	12	4	-	6	11	16	21	29	50	90	152
2001.....	10,084	12	2	2	5	8	10	21	32	60	106	134
2000.....	10,226	16	5	3	3	7	16	11	36	54	91	156
1999.....	10,002	11	1	2	6	6	12	18	30	44	105	135
All origins, male												
2003.....	4,241	3	-	-	5	7	7	11	9	21	39	74
2002.....	4,296	6	4	-	2	2	9	5	14	14	48	82
2001.....	4,282	8	-	-	-	5	4	9	16	27	44	72
2000.....	4,423	6	2	1	3	2	8	6	19	20	37	84
1999.....	4,179	8	-	1	4	4	8	10	14	19	45	73
All origins, female												
2003.....	5,621	7	-	1	3	7	6	8	11	22	53	82
2002.....	5,813	6	-	-	4	9	7	16	15	36	42	70
2001.....	5,802	4	2	2	5	3	6	12	16	33	62	62
2000.....	5,803	10	3	2	-	5	8	5	17	34	54	72
1999.....	5,823	3	1	1	2	2	4	8	16	25	60	62
Hispanic, both sexes												
2003.....	363	2	-	-	1	1	2	5	1	6	8	13
2002.....	416	1	-	-	1	2	2	3	3	6	14	12
2001.....	351	2	-	1	1	3	-	3	2	3	10	13
2000.....	353	3	1	-	-	-	1	3	6	8	9	6
1999.....	345	2	-	-	1	2	3	3	3	3	7	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hispanic, male												
2003.....	177	1	-	-	-	-	1	3	-	6	3	7
2002.....	179	-	-	-	-	1	2	1	1	1	10	4
2001.....	145	1	-	-	-	2	-	-	-	-	6	6
2000.....	158	1	-	-	-	-	-	1	2	1	7	1
1999.....	160	1	-	-	-	1	3	1	1	2	3	2
Hispanic, female												
2003.....	186	1	-	-	1	1	1	2	1	-	5	6
2002.....	237	1	-	-	1	1	-	2	2	5	4	8
2001.....	206	1	-	1	1	1	-	3	2	3	4	7
2000.....	195	2	1	-	-	-	1	2	4	7	2	5
1999.....	185	1	-	-	1	1	-	2	2	1	4	3
Non-Hispanic 1/, both sexes												
2003.....	9,480	8	-	1	7	13	11	14	19	37	84	143
2002.....	9,664	11	4	-	5	9	14	18	26	44	75	140
2001.....	9,707	10	2	1	4	5	10	18	29	57	96	121
2000.....	9,851	13	4	3	3	7	15	8	30	46	82	150
1999.....	9,630	9	1	2	5	4	9	15	27	41	98	130
Non-Hispanic 1/, male												
2003.....	4,057	2	-	-	5	7	6	8	9	15	36	67
2002.....	4,108	6	4	-	2	1	7	4	13	13	37	78
2001.....	4,128	7	-	-	-	3	4	9	15	27	38	66
2000.....	4,255	5	2	1	3	2	8	5	17	19	30	83
1999.....	4,009	7	-	1	4	3	5	9	13	17	42	71

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Non-Hispanic 1/, female												
2003.....	5,423	6	-	1	2	6	5	6	10	22	48	76
2002.....	5,556	5	-	-	3	8	7	14	13	31	38	62
2001.....	5,579	3	2	1	4	2	6	9	14	30	58	55
2000.....	5,596	8	2	2	-	5	7	3	13	27	52	67
1999.....	5,621	2	1	1	1	1	4	6	14	24	56	59
Non-Hispanic White, both sexes												
2003.....	8,044	4	-	1	5	8	6	11	13	21	57	99
2002.....	8,220	8	4	-	2	6	8	11	18	27	53	98
2001.....	8,222	8	1	1	3	4	8	12	19	38	62	88
2000.....	8,391	11	3	2	3	5	12	6	21	35	52	104
1999.....	8,257	7	1	2	4	3	7	10	20	30	76	105
Non-Hispanic White, male												
2003.....	3,457	1	-	-	4	5	4	7	6	9	25	54
2002.....	3,503	5	4	-	1	-	4	3	8	10	28	62
2001.....	3,513	5	-	-	-	3	3	7	9	16	23	47
2000.....	3,650	3	2	1	3	1	7	4	14	14	19	60
1999.....	3,463	5	-	1	3	2	4	6	11	13	36	63
Non-Hispanic White, female												
2003.....	4,587	3	-	1	1	3	2	4	7	12	32	45
2002.....	4,717	3	-	-	1	6	4	8	10	17	25	36
2001.....	4,709	3	1	1	3	1	5	5	10	22	39	41
2000.....	4,741	8	1	1	-	4	5	2	7	21	33	44
1999.....	4,794	2	1	1	1	1	3	4	9	17	40	42





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
All origins, both sexes												
2003.....	4,737	36	4	1	3	14	36	53	86	126	216	273
2002.....	4,711	27	4	1	1	12	23	38	101	128	177	251
2001.....	4,665	22	5	-	-	10	28	45	79	100	185	222
2000.....	4,603	27	4	5	7	10	27	58	75	125	194	249
1999.....	4,025	23	1	1	1	11	19	30	63	98	173	202
All origins, male												
2003.....	1,966	21	3	1	1	5	15	23	37	57	110	147
2002.....	1,944	15	1	-	-	9	12	12	57	67	87	137
2001.....	1,981	11	5	-	-	5	12	18	35	51	98	127
2000.....	1,981	16	2	3	3	6	10	18	39	70	120	145
1999.....	1,714	13	1	1	-	6	7	14	30	55	104	102
All origins, female												
2003.....	2,771	15	1	-	2	9	21	30	49	69	106	126
2002.....	2,767	12	3	1	1	3	11	26	44	61	90	114
2001.....	2,684	11	-	-	-	5	16	27	44	49	87	95
2000.....	2,622	11	2	2	4	4	17	40	36	55	74	104
1999.....	2,311	10	-	-	1	5	12	16	33	43	69	100
Hispanic, both sexes												
2003.....	239	11	2	1	-	3	4	5	10	8	18	18
2002.....	219	7	1	-	-	2	3	6	9	10	11	17
2001.....	219	5	-	-	-	1	2	3	4	5	10	18
2000.....	229	6	-	2	2	-	3	1	10	14	18	14
1999.....	186	2	-	1	-	2	2	-	6	4	8	20

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

Hispanic, male

2003.....	120	5	1	1	-	1	1	1	4	6	13	9
2002.....	95	5	-	-	-	1	2	3	5	5	4	7
2001.....	104	4	-	-	-	1	1	2	2	4	3	12
2000.....	116	3	-	1	1	-	1	1	6	10	10	8
1999.....	83	-	-	1	-	2	-	-	4	3	5	9

Hispanic, female

2003.....	119	6	1	-	-	2	3	4	6	2	5	9
2002.....	124	2	1	-	-	1	1	3	4	5	7	10
2001.....	115	1	-	-	-	-	1	1	2	1	7	6
2000.....	113	3	-	1	1	-	2	-	4	4	8	6
1999.....	103	2	-	-	-	-	2	-	2	1	3	11

Non-Hispanic 1/, both sexes

2003.....	4,485	25	2	-	3	11	32	47	76	118	197	254
2002.....	4,479	20	3	1	1	10	20	32	92	117	164	234
2001.....	4,425	17	5	-	-	9	26	42	75	94	175	204
2000.....	4,359	20	4	3	5	10	24	57	63	110	173	234
1999.....	3,825	21	1	-	1	9	17	30	56	92	164	181

Non-Hispanic 1/, male

2003.....	1,840	16	2	-	1	4	14	21	33	51	97	137
2002.....	1,842	10	1	-	-	8	10	9	52	61	82	130
2001.....	1,869	7	5	-	-	4	11	16	33	46	95	115
2000.....	1,856	12	2	2	2	6	9	17	32	60	107	136
1999.....	1,623	13	1	-	-	4	7	14	26	50	98	92

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Non-Hispanic 1/, female												
2003.....	2,645	9	-	-	2	7	18	26	43	67	100	117
2002.....	2,637	10	2	1	1	2	10	23	40	56	82	104
2001.....	2,556	10	-	-	-	5	15	26	42	48	80	89
2000.....	2,503	8	2	1	3	4	15	40	31	50	66	98
1999.....	2,202	8	-	-	1	5	10	16	30	42	66	89
Non-Hispanic White, both sexes												
2003.....	3,607	18	2	-	2	6	17	31	41	70	128	183
2002.....	3,630	13	2	-	1	6	10	17	52	72	108	168
2001.....	3,617	12	3	-	-	4	14	27	46	56	119	145
2000.....	3,554	13	3	2	3	4	15	32	38	72	123	174
1999.....	3,149	17	-	-	1	3	9	21	36	60	108	126
Non-Hispanic White, male												
2003.....	1,449	11	2	-	-	2	7	10	21	39	71	93
2002.....	1,450	8	-	-	-	5	4	3	30	39	53	90
2001.....	1,496	4	3	-	-	2	6	9	21	28	64	90
2000.....	1,496	9	1	2	1	3	5	8	18	37	81	102
1999.....	1,310	9	-	-	-	2	5	10	16	32	63	64
Non-Hispanic White, female												
2003.....	2,158	7	-	-	2	4	10	21	20	31	57	90
2002.....	2,180	5	2	-	1	1	6	14	22	33	55	78
2001.....	2,121	8	-	-	-	2	8	18	25	28	55	55
2000.....	2,058	4	2	-	2	1	10	24	20	35	42	72
1999.....	1,839	8	-	-	1	1	4	11	20	28	45	62

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Non-Hispanic Black, both sexes												
2003.....	812	6	-	-	-	5	14	16	32	45	66	69
2002.....	789	7	1	1	-	4	9	15	39	42	52	60
2001.....	729	5	2	-	-	3	11	15	28	35	52	52
2000.....	751	5	1	1	2	6	8	23	25	36	50	58
1999.....	613	3	1	-	-	6	8	9	19	29	52	54
Non-Hispanic Black, male												
2003.....	363	5	-	-	-	2	7	11	11	11	25	42
2002.....	359	2	1	-	-	3	6	6	21	20	28	36
2001.....	335	3	2	-	-	1	4	7	11	16	30	20
2000.....	333	2	1	-	1	3	4	8	14	22	26	34
1999.....	281	3	1	-	-	2	2	4	10	16	33	28
Non-Hispanic Black, female												
2003.....	449	1	-	-	-	3	7	5	21	34	41	27
2002.....	430	5	-	1	-	1	3	9	18	22	24	24
2001.....	394	2	-	-	-	2	7	8	17	19	22	32
2000.....	418	3	-	1	1	3	4	15	11	14	24	24
1999.....	332	-	-	-	-	4	6	5	9	13	19	26
Origin not stated 2/, both sexes												
2003.....	13	-	-	-	-	-	-	1	-	-	1	1
2002.....	13	-	-	-	-	-	-	-	-	1	2	-
2001.....	21	-	-	-	-	-	-	-	-	1	-	-
2000.....	15	1	-	-	-	-	-	-	2	1	3	1
1999.....	14	-	-	-	-	-	-	-	1	2	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

Origin not stated 2/, male

2003.....	6	-	-	-	-	-	-	1	-	-	-	1
2002.....	7	-	-	-	-	-	-	-	-	1	1	-
2001.....	8	-	-	-	-	-	-	-	-	1	-	-
2000.....	9	1	-	-	-	-	-	-	1	-	3	1
1999.....	8	-	-	-	-	-	-	-	-	2	1	1

Origin not stated 2/, female

2003.....	7	-	-	-	-	-	-	-	-	-	1	-
2002.....	6	-	-	-	-	-	-	-	-	-	1	-
2001.....	13	-	-	-	-	-	-	-	-	-	-	-
2000.....	6	-	-	-	-	-	-	-	1	1	-	-
1999.....	6	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Influenza and pneumonia (J10-J18)</b>												
All origins, both sexes												
2003.....	65,163	322	163	75	72	80	144	149	224	376	616	918
2002.....	65,681	263	110	38	53	75	92	135	210	367	604	887
2001.....	62,034	299	112	46	46	66	115	124	215	393	590	810
2000.....	65,313	289	103	47	40	65	124	142	222	405	663	783
1999.....	63,730	320	130	46	47	73	106	135	204	462	601	798
All origins, male												
2003.....	28,778	192	91	37	33	50	74	90	139	208	352	528
2002.....	28,918	162	63	15	23	41	56	68	122	197	349	519
2001.....	27,342	167	57	23	24	36	70	64	127	219	337	497
2000.....	28,658	163	54	27	21	35	67	82	122	211	419	464
1999.....	27,718	171	66	27	26	34	56	76	121	251	341	500
All origins, female												
2003.....	36,385	130	72	38	39	30	70	59	85	168	264	390
2002.....	36,763	101	47	23	30	34	36	67	88	170	255	368
2001.....	34,692	132	55	23	22	30	45	60	88	174	253	313
2000.....	36,655	126	49	20	19	30	57	60	100	194	244	319
1999.....	36,012	149	64	19	21	39	50	59	83	211	260	298
Hispanic, both sexes												
2003.....	2,948	66	37	13	9	11	25	24	30	27	47	63
2002.....	2,824	48	17	6	8	9	10	18	25	37	40	61
2001.....	2,722	57	32	5	5	7	11	10	13	32	41	56
2000.....	2,625	60	17	13	8	4	17	19	29	33	49	54
1999.....	2,246	64	17	5	5	10	15	10	14	46	28	48

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Influenza and pneumonia (J10-J18)</b>												
<b>Hispanic, male</b>												
2003.....	1,429	40	24	5	7	6	14	16	19	12	29	38
2002.....	1,398	33	11	2	4	6	5	12	14	23	29	41
2001.....	1,309	31	16	3	3	6	9	7	7	20	28	47
2000.....	1,303	30	5	7	3	1	9	10	17	21	35	41
1999.....	1,092	40	10	4	2	5	6	6	9	34	20	32
<b>Hispanic, female</b>												
2003.....	1,519	26	13	8	2	5	11	8	11	15	18	25
2002.....	1,426	15	6	4	4	3	5	6	11	14	11	20
2001.....	1,413	26	16	2	2	1	2	3	6	12	13	9
2000.....	1,322	30	12	6	5	3	8	9	12	12	14	13
1999.....	1,154	24	7	1	3	5	9	4	5	12	8	16
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	62,042	254	126	61	63	69	119	125	194	349	568	848
2002.....	62,657	213	92	32	44	65	82	116	184	328	561	820
2001.....	59,103	240	79	41	41	59	104	112	200	360	546	752
2000.....	62,440	228	84	34	32	60	107	121	193	366	608	723
1999.....	61,218	255	110	41	42	63	90	125	190	413	566	741
<b>Non-Hispanic 1/, male</b>												
2003.....	27,255	150	67	32	26	44	60	74	120	196	322	483
2002.....	27,417	128	52	13	19	34	51	55	107	174	317	473
2001.....	25,926	135	41	20	21	30	61	57	118	198	306	448
2000.....	27,218	132	47	20	18	33	58	70	105	187	379	419
1999.....	26,489	131	55	23	24	29	50	70	112	214	317	461

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Influenza and pneumonia (J10-J18)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	34,787	104	59	29	37	25	59	51	74	153	246	365
2002.....	35,240	85	40	19	25	31	31	61	77	154	244	347
2001.....	33,177	105	38	21	20	29	43	55	82	162	240	304
2000.....	35,222	96	37	14	14	27	49	51	88	179	229	304
1999.....	34,729	124	55	18	18	34	40	55	78	199	249	280
<b>Non-Hispanic White, both sexes</b>												
2003.....	54,617	143	80	37	49	46	91	78	128	245	392	601
2002.....	55,419	114	46	23	32	42	61	83	124	219	398	611
2001.....	51,952	117	48	28	29	39	69	72	135	243	362	516
2000.....	55,135	114	51	23	26	43	73	79	134	268	410	501
1999.....	54,280	121	58	31	31	44	51	75	124	283	354	503
<b>Non-Hispanic White, male</b>												
2003.....	23,541	87	44	18	21	29	46	52	79	141	215	343
2002.....	23,934	73	24	8	12	24	38	38	73	113	227	344
2001.....	22,389	71	26	14	14	18	46	33	79	131	202	311
2000.....	23,609	68	24	15	14	25	43	41	70	137	257	280
1999.....	23,097	64	22	17	19	21	30	44	75	148	186	309
<b>Non-Hispanic White, female</b>												
2003.....	31,076	56	36	19	28	17	45	26	49	104	177	258
2002.....	31,485	41	22	15	20	18	23	45	51	106	171	267
2001.....	29,563	46	22	14	15	21	23	39	56	112	160	205
2000.....	31,526	46	27	8	12	18	30	38	64	131	153	221
1999.....	31,183	57	36	14	12	23	21	31	49	135	168	194

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Influenza and pneumonia (J10-J18)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	5,798	91	27	19	9	17	25	43	58	93	158	221
2002.....	5,803	86	40	7	11	18	20	27	53	101	149	189
2001.....	5,686	103	24	9	8	16	29	31	56	109	165	219
2000.....	5,920	101	29	7	4	14	30	37	55	83	171	197
1999.....	5,794	114	40	8	7	14	35	47	60	116	189	220
<b>Non-Hispanic Black, male</b>												
2003.....	2,861	52	13	12	3	10	14	21	37	47	95	124
2002.....	2,734	47	22	4	6	8	12	13	30	54	82	116
2001.....	2,769	52	13	4	5	11	12	17	35	62	93	128
2000.....	2,880	55	20	4	3	6	15	26	32	42	109	124
1999.....	2,786	58	25	5	3	6	17	25	33	58	115	140
<b>Non-Hispanic Black, female</b>												
2003.....	2,937	39	14	7	6	7	11	22	21	46	63	97
2002.....	3,069	39	18	3	5	10	8	14	23	47	67	73
2001.....	2,917	51	11	5	3	5	17	14	21	47	72	91
2000.....	3,040	46	9	3	1	8	15	11	23	41	62	73
1999.....	3,008	56	15	3	4	8	18	22	27	58	74	80
<b>Origin not stated 2/, both sexes</b>												
2003.....	173	2	-	1	-	-	-	-	-	-	1	7
2002.....	200	2	1	-	1	1	-	1	1	2	3	6
2001.....	209	2	1	-	-	-	-	2	2	1	3	2
2000.....	248	1	2	-	-	1	-	2	-	6	6	6
1999.....	266	1	3	-	-	-	1	-	-	3	7	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

Origin not stated 2/, male

2003.....	94	2	-	-	-	-	-	-	-	-	1	7
2002.....	103	1	-	-	-	1	-	1	1	-	3	5
2001.....	107	1	-	-	-	-	-	-	2	1	3	2
2000.....	137	1	2	-	-	1	-	2	-	3	5	4
1999.....	137	-	1	-	-	-	-	-	-	3	4	7

Origin not stated 2/, female

2003.....	79	-	-	1	-	-	-	-	-	-	-	-
2002.....	97	1	1	-	1	-	-	-	-	2	-	1
2001.....	102	1	1	-	-	-	-	2	-	-	-	-
2000.....	111	-	-	-	-	-	-	-	-	3	1	2
1999.....	129	1	2	-	-	-	1	-	-	-	3	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Influenza (J10-J11)</b>												
<b>All origins, both sexes</b>												
2003.....	1,792	31	59	28	18	18	12	5	8	10	14	29
2002.....	727	7	5	5	7	5	2	3	1	3	5	8
2001.....	257	7	6	9	3	4	3	4	6	1	5	8
2000.....	1,765	9	10	3	8	8	5	8	5	8	14	24
1999.....	1,665	13	12	6	5	7	4	2	7	8	8	13
<b>All origins, male</b>												
2003.....	699	18	33	13	6	9	3	4	5	5	5	15
2002.....	289	5	3	1	1	2	2	2	-	2	2	7
2001.....	101	6	2	3	1	3	3	3	3	1	1	5
2000.....	664	5	4	1	4	3	3	4	1	5	9	11
1999.....	602	6	9	4	1	1	1	-	5	5	4	8
<b>All origins, female</b>												
2003.....	1,093	13	26	15	12	9	9	1	3	5	9	14
2002.....	438	2	2	4	6	3	-	1	1	1	3	1
2001.....	156	1	4	6	2	1	-	1	3	-	4	3
2000.....	1,101	4	6	2	4	5	2	4	4	3	5	13
1999.....	1,063	7	3	2	4	6	3	2	2	3	4	5
<b>Hispanic, both sexes</b>												
2003.....	75	10	14	6	3	4	-	-	2	-	3	1
2002.....	19	2	-	3	1	-	-	1	-	1	-	1
2001.....	13	-	3	1	-	1	-	-	-	-	1	1
2000.....	47	2	1	1	2	1	-	3	1	2	-	2
1999.....	45	4	3	-	-	1	1	-	-	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Influenza (J10-J11)</b>												
<b>Hispanic, male</b>												
2003.....	40	7	10	2	1	1	-	-	1	-	2	-
2002.....	10	2	-	1	1	-	-	1	-	1	-	1
2001.....	7	-	2	1	-	1	-	-	-	-	-	1
2000.....	21	-	1	-	1	1	-	1	-	1	-	1
1999.....	21	-	3	-	-	-	-	-	-	1	1	-
<b>Hispanic, female</b>												
2003.....	35	3	4	4	2	3	-	-	1	-	1	1
2002.....	9	-	-	2	-	-	-	-	-	-	-	-
2001.....	6	-	1	-	-	-	-	-	-	-	1	-
2000.....	26	2	-	1	1	-	-	2	1	1	-	1
1999.....	24	4	-	-	-	1	1	-	-	-	-	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	1,714	21	45	22	15	14	12	5	6	10	11	28
2002.....	704	5	5	2	6	5	2	2	1	2	5	7
2001.....	244	7	3	8	3	3	3	4	6	1	4	7
2000.....	1,716	7	9	2	6	7	5	5	4	6	14	22
1999.....	1,612	9	9	6	5	6	3	2	7	7	7	13
<b>Non-Hispanic 1/, male</b>												
2003.....	657	11	23	11	5	8	3	4	4	5	3	15
2002.....	277	3	3	-	-	2	2	1	-	1	2	6
2001.....	94	6	-	2	1	2	3	3	3	1	1	4
2000.....	642	5	3	1	3	2	3	3	1	4	9	10
1999.....	580	6	6	4	1	1	1	-	5	4	3	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Influenza (J10-J11)												
Non-Hispanic 1/, female												
2003.....	1,057	10	22	11	10	6	9	1	2	5	8	13
2002.....	427	2	2	2	6	3	-	1	1	1	3	1
2001.....	150	1	3	6	2	1	-	1	3	-	3	3
2000.....	1,074	2	6	1	3	5	2	2	3	2	5	12
1999.....	1,032	3	3	2	4	5	2	2	2	3	4	5
Non-Hispanic White, both sexes												
2003.....	1,578	13	29	14	8	7	9	3	5	8	9	21
2002.....	663	3	2	1	5	3	2	2	1	2	5	7
2001.....	216	7	2	4	2	2	2	2	5	1	2	7
2000.....	1,613	5	8	1	5	7	4	4	3	5	11	16
1999.....	1,522	3	5	6	4	5	3	1	7	3	5	12
Non-Hispanic White, male												
2003.....	599	7	15	7	2	4	2	3	4	3	1	11
2002.....	258	1	1	-	-	-	2	1	-	1	2	6
2001.....	81	6	-	1	1	1	2	1	2	1	1	4
2000.....	595	5	3	1	2	2	2	2	-	3	6	6
1999.....	547	1	3	4	1	1	1	-	5	2	2	7
Non-Hispanic White, female												
2003.....	979	6	14	7	6	3	7	-	1	5	8	10
2002.....	405	2	1	1	5	3	-	1	1	1	3	1
2001.....	135	1	2	3	1	1	-	1	3	-	1	3
2000.....	1,018	-	5	-	3	5	2	2	3	2	5	10
1999.....	975	2	2	2	3	4	2	1	2	1	3	5





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Pneumonia (J12-J18)</b>												
<b>All origins, both sexes</b>												
2003.....	63,371	291	104	47	54	62	132	144	216	366	602	889
2002.....	64,954	256	105	33	46	70	90	132	209	364	599	879
2001.....	61,777	292	106	37	43	62	112	120	209	392	585	802
2000.....	63,548	280	93	44	32	57	119	134	217	397	649	759
1999.....	62,065	307	118	40	42	66	102	133	197	454	593	785
<b>All origins, male</b>												
2003.....	28,079	174	58	24	27	41	71	86	134	203	347	513
2002.....	28,629	157	60	14	22	39	54	66	122	195	347	512
2001.....	27,241	161	55	20	23	33	67	61	124	218	336	492
2000.....	27,994	158	50	26	17	32	64	78	121	206	410	453
1999.....	27,116	165	57	23	25	33	55	76	116	246	337	492
<b>All origins, female</b>												
2003.....	35,292	117	46	23	27	21	61	58	82	163	255	376
2002.....	36,325	99	45	19	24	31	36	66	87	169	252	367
2001.....	34,536	131	51	17	20	29	45	59	85	174	249	310
2000.....	35,554	122	43	18	15	25	55	56	96	191	239	306
1999.....	34,949	142	61	17	17	33	47	57	81	208	256	293
<b>Hispanic, both sexes</b>												
2003.....	2,873	56	23	7	6	7	25	24	28	27	44	62
2002.....	2,805	46	17	3	7	9	10	17	25	36	40	60
2001.....	2,709	57	29	4	5	6	11	10	13	32	40	55
2000.....	2,578	58	16	12	6	3	17	16	28	31	49	52
1999.....	2,201	60	14	5	5	9	14	10	14	45	27	48

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Pneumonia (J12-J18)</b>												
<b>Hispanic, male</b>												
2003.....	1,389	33	14	3	6	5	14	16	18	12	27	38
2002.....	1,388	31	11	1	3	6	5	11	14	22	29	40
2001.....	1,302	31	14	2	3	5	9	7	7	20	28	46
2000.....	1,282	30	4	7	2	-	9	9	17	20	35	40
1999.....	1,071	40	7	4	2	5	6	6	9	33	19	32
<b>Hispanic, female</b>												
2003.....	1,484	23	9	4	-	2	11	8	10	15	17	24
2002.....	1,417	15	6	2	4	3	5	6	11	14	11	20
2001.....	1,407	26	15	2	2	1	2	3	6	12	12	9
2000.....	1,296	28	12	5	4	3	8	7	11	11	14	12
1999.....	1,130	20	7	1	3	4	8	4	5	12	8	16
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	60,328	233	81	39	48	55	107	120	188	339	557	820
2002.....	61,953	208	87	30	38	60	80	114	183	326	556	813
2001.....	58,859	233	76	33	38	56	101	108	194	359	542	745
2000.....	60,724	221	75	32	26	53	102	116	189	360	594	701
1999.....	59,606	246	101	35	37	57	87	123	183	406	559	728
<b>Non-Hispanic 1/, male</b>												
2003.....	26,598	139	44	21	21	36	57	70	116	191	319	468
2002.....	27,140	125	49	13	19	32	49	54	107	173	315	467
2001.....	25,832	129	41	18	20	28	58	54	115	197	305	444
2000.....	26,576	127	44	19	15	31	55	67	104	183	370	409
1999.....	25,909	125	49	19	23	28	49	70	107	210	314	453

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Pneumonia (J12-J18)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	33,730	94	37	18	27	19	50	50	72	148	238	352
2002.....	34,813	83	38	17	19	28	31	60	76	153	241	346
2001.....	33,027	104	35	15	18	28	43	54	79	162	237	301
2000.....	34,148	94	31	13	11	22	47	49	85	177	224	292
1999.....	33,697	121	52	16	14	29	38	53	76	196	245	275
<b>Non-Hispanic White, both sexes</b>												
2003.....	53,039	130	51	23	41	39	82	75	123	237	383	580
2002.....	54,756	111	44	22	27	39	59	81	123	217	393	604
2001.....	51,736	110	46	24	27	37	67	70	130	242	360	509
2000.....	53,522	109	43	22	21	36	69	75	131	263	399	485
1999.....	52,758	118	53	25	27	39	48	74	117	280	349	491
<b>Non-Hispanic White, male</b>												
2003.....	22,942	80	29	11	19	25	44	49	75	138	214	332
2002.....	23,676	72	23	8	12	24	36	37	73	112	225	338
2001.....	22,308	65	26	13	13	17	44	32	77	130	201	307
2000.....	23,014	63	21	14	12	23	41	39	70	134	251	274
1999.....	22,550	63	19	13	18	20	29	44	70	146	184	302
<b>Non-Hispanic White, female</b>												
2003.....	30,097	50	22	12	22	14	38	26	48	99	169	248
2002.....	31,080	39	21	14	15	15	23	44	50	105	168	266
2001.....	29,428	45	20	11	14	20	23	38	53	112	159	202
2000.....	30,508	46	22	8	9	13	28	36	61	129	148	211
1999.....	30,208	55	34	12	9	19	19	30	47	134	165	189

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Pneumonia (J12-J18)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	5,703	84	21	13	5	13	22	41	57	93	157	214
2002.....	5,773	85	37	6	10	16	20	27	53	101	149	189
2001.....	5,666	103	24	7	8	15	29	30	55	109	164	219
2000.....	5,839	99	28	7	3	14	29	36	55	83	168	192
1999.....	5,729	110	39	8	7	13	35	46	60	113	187	219
<b>Non-Hispanic Black, male</b>												
2003.....	2,825	49	10	9	2	8	13	20	37	47	94	120
2002.....	2,719	46	20	4	6	6	12	13	30	54	82	116
2001.....	2,759	52	13	4	5	10	12	16	34	62	93	128
2000.....	2,845	55	20	4	2	6	14	25	32	42	106	121
1999.....	2,762	55	24	5	3	6	17	25	33	57	114	139
<b>Non-Hispanic Black, female</b>												
2003.....	2,878	35	11	4	3	5	9	21	20	46	63	94
2002.....	3,054	39	17	2	4	10	8	14	23	47	67	73
2001.....	2,907	51	11	3	3	5	17	14	21	47	71	91
2000.....	2,994	44	8	3	1	8	15	11	23	41	62	71
1999.....	2,967	55	15	3	4	7	18	21	27	56	73	80
<b>Origin not stated 2/, both sexes</b>												
2003.....	170	2	-	1	-	-	-	-	-	-	1	7
2002.....	196	2	1	-	1	1	-	1	1	2	3	6
2001.....	209	2	1	-	-	-	-	2	2	1	3	2
2000.....	246	1	2	-	-	1	-	2	-	6	6	6
1999.....	258	1	3	-	-	-	1	-	-	3	7	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonia (J12-J18)												
Origin not stated 2/, male												
2003.....	92	2	-	-	-	-	-	-	-	-	1	7
2002.....	101	1	-	-	-	1	-	1	1	-	3	5
2001.....	107	1	-	-	-	-	-	-	2	1	3	2
2000.....	136	1	2	-	-	1	-	2	-	3	5	4
1999.....	136	-	1	-	-	-	-	-	-	3	4	7
Origin not stated 2/, female												
2003.....	78	-	-	1	-	-	-	-	-	-	-	-
2002.....	95	1	1	-	1	-	-	-	-	2	-	1
2001.....	102	1	1	-	-	-	-	2	-	-	-	-
2000.....	110	-	-	-	-	-	-	-	-	3	1	2
1999.....	122	1	2	-	-	-	1	-	-	-	3	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

All origins, both sexes

2003.....	421	54	28	3	3	1	5	-	2	4	7	7
2002.....	386	52	23	7	1	1	2	3	2	2	6	6
2001.....	347	50	18	3	-	-	1	1	1	7	5	4
2000.....	425	43	11	3	4	4	1	5	3	3	9	8
1999.....	540	63	25	2	3	3	2	2	2	6	5	7

All origins, male

2003.....	170	35	17	2	-	-	2	-	1	3	3	2
2002.....	154	39	9	7	-	-	-	1	-	1	2	1
2001.....	117	26	8	2	-	-	1	-	-	4	3	1
2000.....	160	25	4	2	1	4	-	5	1	1	6	2
1999.....	196	36	15	-	2	1	-	2	1	3	2	5

All origins, female

2003.....	251	19	11	1	3	1	3	-	1	1	4	5
2002.....	232	13	14	-	1	1	2	2	2	1	4	5
2001.....	230	24	10	1	-	-	-	1	1	3	2	3
2000.....	265	18	7	1	3	-	1	-	2	2	3	6
1999.....	344	27	10	2	1	2	2	-	1	3	3	2

Hispanic, both sexes

2003.....	31	17	6	-	-	-	-	-	-	-	1	-
2002.....	21	10	2	1	1	-	-	1	-	-	-	-
2001.....	22	8	1	-	-	-	-	-	1	-	-	-
2000.....	24	9	2	-	-	-	-	1	-	-	-	1
1999.....	21	10	3	-	1	-	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

Hispanic, male

2003.....	20	12	4	-	-	-	-	-	-	-	-	-
2002.....	9	5	1	1	-	-	-	-	-	-	-	-
2001.....	9	6	-	-	-	-	-	-	-	-	-	-
2000.....	15	6	1	-	-	-	-	1	-	-	-	1
1999.....	10	5	2	-	1	-	-	-	-	-	-	-

Hispanic, female

2003.....	11	5	2	-	-	-	-	-	-	-	1	-
2002.....	12	5	1	-	1	-	-	1	-	-	-	-
2001.....	13	2	1	-	-	-	-	-	1	-	-	-
2000.....	9	3	1	-	-	-	-	-	-	-	-	-
1999.....	11	5	1	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	388	37	22	3	3	1	5	-	2	4	6	7
2002.....	365	42	21	6	-	1	2	2	2	2	6	6
2001.....	325	42	17	3	-	-	1	1	-	7	5	4
2000.....	398	33	9	3	4	4	1	4	3	3	9	7
1999.....	518	53	22	2	2	3	2	2	2	5	5	7

Non-Hispanic 1/, male

2003.....	148	23	13	2	-	-	2	-	1	3	3	2
2002.....	145	34	8	6	-	-	-	1	-	1	2	1
2001.....	108	20	8	2	-	-	1	-	-	4	3	1
2000.....	144	19	3	2	1	4	-	4	1	1	6	1
1999.....	185	31	13	-	1	1	-	2	1	2	2	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

Non-Hispanic 1/, female

2003.....	240	14	9	1	3	1	3	-	1	1	3	5
2002.....	220	8	13	-	-	1	2	1	2	1	4	5
2001.....	217	22	9	1	-	-	-	1	-	3	2	3
2000.....	254	14	6	1	3	-	1	-	2	2	3	6
1999.....	333	22	9	2	1	2	2	-	1	3	3	2

Non-Hispanic White, both sexes

2003.....	342	25	11	3	2	1	2	-	1	4	6	6
2002.....	314	25	18	6	-	1	1	1	2	2	5	4
2001.....	278	24	12	3	-	-	1	-	-	7	5	4
2000.....	345	15	6	2	1	4	1	3	1	2	6	4
1999.....	440	26	12	1	2	2	2	1	2	5	3	6

Non-Hispanic White, male

2003.....	125	15	6	2	-	-	-	-	1	3	3	2
2002.....	122	22	6	6	-	-	-	-	-	1	2	1
2001.....	86	11	5	2	-	-	1	-	-	4	3	1
2000.....	118	6	3	2	-	4	-	3	-	1	5	1
1999.....	145	14	6	-	1	1	-	1	1	2	1	5

Non-Hispanic White, female

2003.....	217	10	5	1	2	1	2	-	-	1	3	4
2002.....	192	3	12	-	-	1	1	1	2	1	3	3
2001.....	192	13	7	1	-	-	-	-	-	3	2	3
2000.....	227	9	3	-	1	-	1	-	1	1	1	3
1999.....	295	12	6	1	1	1	2	-	1	3	2	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other acute lower respiratory infections (J20-J22)

Non-Hispanic Black, both sexes

2003.....	40	12	7	-	1	-	3	-	1	-	-	1
2002.....	44	15	2	-	-	-	1	1	-	-	1	2
2001.....	33	14	2	-	-	-	-	1	-	-	-	-
2000.....	46	14	3	-	3	-	-	1	1	1	3	3
1999.....	64	24	6	-	-	1	-	1	-	-	2	1

Non-Hispanic Black, male

2003.....	21	8	6	-	-	-	2	-	-	-	-	-
2002.....	22	12	1	-	-	-	-	1	-	-	-	-
2001.....	15	8	1	-	-	-	-	-	-	-	-	-
2000.....	22	10	-	-	1	-	-	1	1	-	1	-
1999.....	30	15	3	-	-	-	-	1	-	-	1	-

Non-Hispanic Black, female

2003.....	19	4	1	-	1	-	1	-	1	-	-	1
2002.....	22	3	1	-	-	-	1	-	-	-	1	2
2001.....	18	6	1	-	-	-	-	1	-	-	-	-
2000.....	24	4	3	-	2	-	-	-	-	1	2	3
1999.....	34	9	3	-	-	1	-	-	-	-	1	1

Origin not stated 2/, both sexes

2003.....	2	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	3	1	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
All origins, both sexes												
2003.....	304	53	27	3	3	1	5	-	2	4	5	7
2002.....	279	49	23	7	1	-	2	3	2	2	4	6
2001.....	263	49	18	3	-	-	1	1	1	6	5	4
2000.....	290	42	11	2	4	3	1	3	3	2	8	8
1999.....	388	62	24	2	2	3	2	1	2	6	4	7
All origins, male												
2003.....	126	34	16	2	-	-	2	-	1	3	2	2
2002.....	122	37	9	7	-	-	-	1	-	1	2	1
2001.....	89	25	8	2	-	-	1	-	-	3	3	1
2000.....	119	24	4	1	1	3	-	3	1	1	5	2
1999.....	154	36	14	-	2	1	-	1	1	3	1	5
All origins, female												
2003.....	178	19	11	1	3	1	3	-	1	1	3	5
2002.....	157	12	14	-	1	-	2	2	2	1	2	5
2001.....	174	24	10	1	-	-	-	1	1	3	2	3
2000.....	171	18	7	1	3	-	1	-	2	1	3	6
1999.....	234	26	10	2	-	2	2	-	1	3	3	2
Hispanic, both sexes												
2003.....	31	17	6	-	-	-	-	-	-	-	1	-
2002.....	19	9	2	1	1	-	-	1	-	-	-	-
2001.....	17	8	1	-	-	-	-	-	1	-	-	-
2000.....	22	9	2	-	-	-	-	1	-	-	-	1
1999.....	20	10	3	-	1	-	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Acute bronchitis and bronchiolitis (J20-J21)</b>												
<b>Hispanic, male</b>												
2003.....	20	12	4	-	-	-	-	-	-	-	-	-
2002.....	9	5	1	1	-	-	-	-	-	-	-	-
2001.....	7	6	-	-	-	-	-	-	-	-	-	-
2000.....	15	6	1	-	-	-	-	1	-	-	-	1
1999.....	10	5	2	-	1	-	-	-	-	-	-	-
<b>Hispanic, female</b>												
2003.....	11	5	2	-	-	-	-	-	-	-	1	-
2002.....	10	4	1	-	1	-	-	1	-	-	-	-
2001.....	10	2	1	-	-	-	-	-	1	-	-	-
2000.....	7	3	1	-	-	-	-	-	-	-	-	-
1999.....	10	5	1	-	-	-	-	-	-	-	-	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	272	36	21	3	3	1	5	-	2	4	4	7
2002.....	260	40	21	6	-	-	2	2	2	2	4	6
2001.....	246	41	17	3	-	-	1	1	-	6	5	4
2000.....	267	32	9	2	4	3	1	2	3	2	8	7
1999.....	367	52	21	2	1	3	2	1	2	5	4	7
<b>Non-Hispanic 1/, male</b>												
2003.....	105	22	12	2	-	-	2	-	1	3	2	2
2002.....	113	32	8	6	-	-	-	1	-	1	2	1
2001.....	82	19	8	2	-	-	1	-	-	3	3	1
2000.....	104	18	3	1	1	3	-	2	1	1	5	1
1999.....	143	31	12	-	1	1	-	1	1	2	1	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute bronchitis and bronchiolitis (J20-J21)												
Non-Hispanic 1/, female												
2003.....	167	14	9	1	3	1	3	-	1	1	2	5
2002.....	147	8	13	-	-	-	2	1	2	1	2	5
2001.....	164	22	9	1	-	-	-	1	-	3	2	3
2000.....	163	14	6	1	3	-	1	-	2	1	3	6
1999.....	224	21	9	2	-	2	2	-	1	3	3	2
Non-Hispanic White, both sexes												
2003.....	231	24	10	3	2	1	2	-	1	4	4	6
2002.....	219	23	18	6	-	-	1	1	2	2	4	4
2001.....	202	23	12	3	-	-	1	-	-	6	5	4
2000.....	221	15	6	1	1	3	1	1	1	1	5	4
1999.....	299	25	12	1	1	2	2	1	2	5	2	6
Non-Hispanic White, male												
2003.....	83	14	5	2	-	-	-	-	1	3	2	2
2002.....	92	20	6	6	-	-	-	-	-	1	2	1
2001.....	61	10	5	2	-	-	1	-	-	3	3	1
2000.....	83	6	3	1	-	3	-	1	-	1	4	1
1999.....	108	14	6	-	1	1	-	1	1	2	-	5
Non-Hispanic White, female												
2003.....	148	10	5	1	2	1	2	-	-	1	2	4
2002.....	127	3	12	-	-	-	1	1	2	1	2	3
2001.....	141	13	7	1	-	-	-	-	-	3	2	3
2000.....	138	9	3	-	1	-	1	-	1	-	1	3
1999.....	191	11	6	1	-	1	2	-	1	3	2	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute bronchitis and bronchiolitis (J20-J21)

Non-Hispanic Black, both sexes

2003.....	35	12	7	-	1	-	3	-	1	-	-	1
2002.....	35	15	2	-	-	-	1	1	-	-	-	2
2001.....	31	14	2	-	-	-	-	1	-	-	-	-
2000.....	40	13	3	-	3	-	-	1	1	1	3	3
1999.....	55	24	5	-	-	1	-	-	-	-	2	1

Non-Hispanic Black, male

2003.....	20	8	6	-	-	-	2	-	-	-	-	-
2002.....	20	12	1	-	-	-	-	1	-	-	-	-
2001.....	14	8	1	-	-	-	-	-	-	-	-	-
2000.....	18	9	-	-	1	-	-	1	1	-	1	-
1999.....	26	15	2	-	-	-	-	-	-	-	1	-

Non-Hispanic Black, female

2003.....	15	4	1	-	1	-	1	-	1	-	-	1
2002.....	15	3	1	-	-	-	1	-	-	-	-	2
2001.....	17	6	1	-	-	-	-	1	-	-	-	-
2000.....	22	4	3	-	2	-	-	-	-	1	2	3
1999.....	29	9	3	-	-	1	-	-	-	-	1	1

Origin not stated 2/, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	1	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified acute lower respiratory infection (J22)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	1	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	116	1	1	-	-	-	-	-	-	-	2	-
2002.....	105	2	-	-	-	1	-	-	-	-	2	-
2001.....	79	1	-	-	-	-	-	-	-	1	-	-
2000.....	131	1	-	1	-	1	-	2	-	1	1	-
1999.....	151	1	1	-	1	-	-	1	-	-	1	-

Non-Hispanic 1/, male

2003.....	43	1	1	-	-	-	-	-	-	-	1	-
2002.....	32	2	-	-	-	-	-	-	-	-	-	-
2001.....	26	1	-	-	-	-	-	-	-	1	-	-
2000.....	40	1	-	1	-	1	-	2	-	-	1	-
1999.....	42	-	1	-	-	-	-	1	-	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Unspecified acute lower respiratory infection (J22)

Non-Hispanic 1/, female

2003.....	73	-	-	-	-	-	-	-	-	-	1	-
2002.....	73	-	-	-	-	1	-	-	-	-	2	-
2001.....	53	-	-	-	-	-	-	-	-	-	-	-
2000.....	91	-	-	-	-	-	-	-	-	1	-	-
1999.....	109	1	-	-	1	-	-	-	-	-	-	-

Non-Hispanic White, both sexes

2003.....	111	1	1	-	-	-	-	-	-	-	2	-
2002.....	95	2	-	-	-	1	-	-	-	-	1	-
2001.....	76	1	-	-	-	-	-	-	-	1	-	-
2000.....	124	-	-	1	-	1	-	2	-	1	1	-
1999.....	141	1	-	-	1	-	-	-	-	-	1	-

Non-Hispanic White, male

2003.....	42	1	1	-	-	-	-	-	-	-	1	-
2002.....	30	2	-	-	-	-	-	-	-	-	-	-
2001.....	25	1	-	-	-	-	-	-	-	1	-	-
2000.....	35	-	-	1	-	1	-	2	-	-	1	-
1999.....	37	-	-	-	-	-	-	-	-	-	1	-

Non-Hispanic White, female

2003.....	69	-	-	-	-	-	-	-	-	-	1	-
2002.....	65	-	-	-	-	1	-	-	-	-	1	-
2001.....	51	-	-	-	-	-	-	-	-	-	-	-
2000.....	89	-	-	-	-	-	-	-	-	1	-	-
1999.....	104	1	-	-	1	-	-	-	-	-	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic lower respiratory diseases (J40-J47)</b>												
<b>All origins, both sexes</b>												
2003.....	126,382	31	55	37	81	84	107	118	164	301	649	1,276
2002.....	124,816	39	65	41	95	93	99	116	185	340	668	1,226
2001.....	123,013	41	43	42	62	74	97	102	188	355	614	1,127
2000.....	122,009	36	51	48	91	86	104	125	171	310	620	1,174
1999.....	124,181	36	54	49	90	107	102	145	185	302	588	1,079
<b>All origins, male</b>												
2003.....	60,714	23	38	23	52	50	64	67	89	142	287	681
2002.....	60,713	23	44	24	50	68	52	70	95	159	338	617
2001.....	59,697	22	29	24	37	43	56	55	91	171	284	591
2000.....	60,004	23	28	26	54	58	62	72	93	145	292	617
1999.....	62,415	19	30	29	64	60	63	69	83	142	268	547
<b>All origins, female</b>												
2003.....	65,668	8	17	14	29	34	43	51	75	159	362	595
2002.....	64,103	16	21	17	45	25	47	46	90	181	330	609
2001.....	63,316	19	14	18	25	31	41	47	97	184	330	536
2000.....	62,005	13	23	22	37	28	42	53	78	165	328	557
1999.....	61,766	17	24	20	26	47	39	76	102	160	320	532
<b>Hispanic, both sexes</b>												
2003.....	3,174	7	11	4	11	12	8	14	15	25	33	61
2002.....	3,058	4	9	6	7	6	9	9	22	25	45	50
2001.....	2,832	7	8	2	7	7	3	10	12	25	27	47
2000.....	2,689	7	9	5	10	6	10	7	13	21	34	45
1999.....	2,859	6	8	7	11	8	7	17	14	34	45	52

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

#### Chronic lower respiratory diseases (J40-J47)

##### Hispanic, male

2003.....	1,697	5	8	2	8	7	5	9	6	12	18	31
2002.....	1,625	2	4	4	4	5	7	8	11	12	19	28
2001.....	1,482	4	4	1	5	4	2	6	7	12	9	22
2000.....	1,451	4	5	2	7	5	8	4	8	11	13	21
1999.....	1,547	4	3	3	8	4	5	6	7	14	22	21

##### Hispanic, female

2003.....	1,477	2	3	2	3	5	3	5	9	13	15	30
2002.....	1,433	2	5	2	3	1	2	1	11	13	26	22
2001.....	1,350	3	4	1	2	3	1	4	5	13	18	25
2000.....	1,238	3	4	3	3	1	2	3	5	10	21	24
1999.....	1,312	2	5	4	3	4	2	11	7	20	23	31

##### Non-Hispanic 1/, both sexes

2003.....	122,896	24	43	32	70	71	99	104	149	276	614	1,211
2002.....	121,424	34	56	35	86	86	90	107	160	314	616	1,170
2001.....	119,839	34	35	40	54	67	94	92	175	328	581	1,073
2000.....	118,963	29	42	42	81	80	93	115	157	288	584	1,120
1999.....	120,968	30	46	42	79	99	95	128	168	267	540	1,023

##### Non-Hispanic 1/, male

2003.....	58,848	18	29	21	44	42	59	58	83	130	267	647
2002.....	58,896	20	40	20	44	62	45	62	81	146	314	585
2001.....	58,028	18	25	23	32	39	54	49	84	158	270	565
2000.....	58,349	19	23	24	47	53	54	65	85	133	278	589
1999.....	60,669	15	27	26	56	56	58	63	75	127	245	523

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic lower respiratory diseases (J40-J47)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	64,048	6	14	11	26	29	40	46	66	146	347	564
2002.....	62,528	14	16	15	42	24	45	45	79	168	302	585
2001.....	61,811	16	10	17	22	28	40	43	91	170	311	508
2000.....	60,614	10	19	18	34	27	39	50	72	155	306	531
1999.....	60,299	15	19	16	23	43	37	65	93	140	295	500
<b>Non-Hispanic White, both sexes</b>												
2003.....	113,541	17	15	10	29	27	55	56	84	186	431	952
2002.....	112,128	20	23	11	31	40	54	59	98	201	423	917
2001.....	110,753	16	13	16	23	33	42	43	91	187	426	831
2000.....	109,901	16	19	17	30	34	53	59	85	174	396	830
1999.....	111,620	10	23	23	32	60	50	57	89	167	366	747
<b>Non-Hispanic White, male</b>												
2003.....	53,593	16	7	5	16	14	34	25	39	88	187	518
2002.....	53,662	13	17	8	15	25	23	29	52	84	220	459
2001.....	52,952	10	9	9	13	15	24	23	39	97	196	456
2000.....	53,231	12	8	10	14	21	31	31	40	74	193	445
1999.....	55,314	6	11	13	23	32	27	26	40	77	162	386
<b>Non-Hispanic White, female</b>												
2003.....	59,948	1	8	5	13	13	21	31	45	98	244	434
2002.....	58,466	7	6	3	16	15	31	30	46	117	203	458
2001.....	57,801	6	4	7	10	18	18	20	52	90	230	375
2000.....	56,670	4	11	7	16	13	22	28	45	100	203	385
1999.....	56,306	4	12	10	9	28	23	31	49	90	204	361

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic lower respiratory diseases (J40-J47)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	7,632	7	24	22	40	40	41	45	59	81	164	239
2002.....	7,730	14	29	21	53	43	35	43	59	109	183	230
2001.....	7,504	16	22	23	30	30	47	47	81	131	142	222
2000.....	7,528	11	21	24	46	44	36	49	65	102	170	258
1999.....	7,840	18	18	19	47	36	42	67	73	93	165	256
<b>Non-Hispanic Black, male</b>												
2003.....	4,202	2	19	16	27	25	23	31	41	38	68	119
2002.....	4,282	7	20	11	28	36	22	28	28	60	90	115
2001.....	4,143	6	16	14	18	22	26	26	43	58	66	101
2000.....	4,184	6	14	14	29	31	20	29	41	51	78	125
1999.....	4,452	9	13	13	33	23	29	34	33	44	77	126
<b>Non-Hispanic Black, female</b>												
2003.....	3,430	5	5	6	13	15	18	14	18	43	96	120
2002.....	3,448	7	9	10	25	7	13	15	31	49	93	115
2001.....	3,361	10	6	9	12	8	21	21	38	73	76	121
2000.....	3,344	5	7	10	17	13	16	20	24	51	92	133
1999.....	3,388	9	5	6	14	13	13	33	40	49	88	130
<b>Origin not stated 2/, both sexes</b>												
2003.....	312	-	1	1	-	1	-	-	-	-	2	4
2002.....	334	1	-	-	2	1	-	-	3	1	7	6
2001.....	342	-	-	-	1	-	-	-	1	2	6	7
2000.....	357	-	-	1	-	-	1	3	1	1	2	9
1999.....	354	-	-	-	-	-	-	-	3	1	3	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

Origin not stated 2/, male

2003.....	169	-	1	-	-	1	-	-	-	-	2	3
2002.....	192	1	-	-	2	1	-	-	3	1	5	4
2001.....	187	-	-	-	-	-	-	-	-	1	5	4
2000.....	204	-	-	-	-	-	-	3	-	1	1	7
1999.....	199	-	-	-	-	-	-	-	1	1	1	3

Origin not stated 2/, female

2003.....	143	-	-	1	-	-	-	-	-	-	-	1
2002.....	142	-	-	-	-	-	-	-	-	-	2	2
2001.....	155	-	-	-	1	-	-	-	1	1	1	3
2000.....	153	-	-	1	-	-	1	-	1	-	1	2
1999.....	155	-	-	-	-	-	-	-	2	-	2	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified (J40-J42)												
All origins, both sexes												
2003.....	850	20	13	1	5	2	5	5	5	8	9	17
2002.....	955	24	13	3	2	3	3	2	4	12	6	16
2001.....	959	21	8	-	-	3	2	4	2	9	12	21
2000.....	1,167	18	14	2	3	3	2	3	5	7	13	19
1999.....	1,172	20	19	3	1	3	4	3	8	10	8	14
All origins, male												
2003.....	351	15	6	-	3	2	4	1	2	5	4	13
2002.....	420	15	8	2	-	2	1	-	2	9	3	13
2001.....	396	11	5	-	-	2	1	2	-	3	5	12
2000.....	503	9	7	2	1	2	2	3	4	4	10	8
1999.....	498	9	11	2	1	2	2	-	3	3	2	9
All origins, female												
2003.....	499	5	7	1	2	-	1	4	3	3	5	4
2002.....	535	9	5	1	2	1	2	2	2	3	3	3
2001.....	563	10	3	-	-	1	1	2	2	6	7	9
2000.....	664	9	7	-	2	1	-	-	1	3	3	11
1999.....	674	11	8	1	-	1	2	3	5	7	6	5
Hispanic, both sexes												
2003.....	40	5	5	-	-	-	1	-	1	-	-	1
2002.....	51	3	1	-	-	-	2	-	1	2	1	1
2001.....	37	4	1	-	-	1	-	1	-	1	1	1
2000.....	52	4	3	-	-	-	-	1	1	1	2	1
1999.....	52	3	3	-	1	-	-	2	-	4	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified (J40-J42)												
Hispanic, male												
2003.....	20	3	3	-	-	-	1	-	-	-	-	-
2002.....	20	1	1	-	-	-	1	-	1	1	1	1
2001.....	15	2	-	-	-	-	-	1	-	-	1	-
2000.....	28	2	2	-	-	-	-	1	1	1	1	-
1999.....	17	1	2	-	1	-	-	-	-	1	1	-
Hispanic, female												
2003.....	20	2	2	-	-	-	-	-	1	-	-	1
2002.....	31	2	-	-	-	-	1	-	-	1	-	-
2001.....	22	2	1	-	-	1	-	-	-	1	-	1
2000.....	24	2	1	-	-	-	-	-	-	-	1	1
1999.....	35	2	1	-	-	-	-	2	-	3	2	-
Non-Hispanic 1/, both sexes												
2003.....	808	15	8	1	5	2	4	5	4	8	9	16
2002.....	900	20	12	3	2	3	1	2	3	10	5	15
2001.....	919	17	7	-	-	2	2	3	2	8	11	19
2000.....	1,112	14	11	2	3	3	2	2	4	6	11	18
1999.....	1,117	17	16	3	-	3	4	1	8	6	5	14
Non-Hispanic 1/, male												
2003.....	330	12	3	-	3	2	3	1	2	5	4	13
2002.....	398	13	7	2	-	2	-	-	1	8	2	12
2001.....	380	9	5	-	-	2	1	1	-	3	4	12
2000.....	474	7	5	2	1	2	2	2	3	3	9	8
1999.....	478	8	9	2	-	2	2	-	3	2	1	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Bronchitis, chronic and unspecified (J40-J42)												
Non-Hispanic 1/, female												
2003.....	478	3	5	1	2	-	1	4	2	3	5	3
2002.....	502	7	5	1	2	1	1	2	2	2	3	3
2001.....	539	8	2	-	-	-	1	2	2	5	7	7
2000.....	638	7	6	-	2	1	-	-	1	3	2	10
1999.....	639	9	7	1	-	1	2	1	5	4	4	5
Non-Hispanic White, both sexes												
2003.....	712	11	4	1	2	1	4	1	4	6	7	10
2002.....	821	12	6	2	1	1	1	1	3	9	4	11
2001.....	839	7	5	-	-	1	2	2	2	8	8	15
2000.....	1,018	5	4	2	2	2	2	2	2	6	5	16
1999.....	1,002	8	12	3	-	3	3	-	6	6	3	9
Non-Hispanic White, male												
2003.....	290	10	2	-	1	1	3	-	2	4	3	9
2002.....	354	9	2	2	-	1	-	-	1	7	1	8
2001.....	338	5	5	-	-	1	1	-	-	3	3	9
2000.....	429	3	1	2	-	1	2	2	1	3	3	8
1999.....	421	5	6	2	-	2	1	-	2	2	1	5
Non-Hispanic White, female												
2003.....	422	1	2	1	1	-	1	1	2	2	4	1
2002.....	467	3	4	-	1	-	1	1	2	2	3	3
2001.....	501	2	-	-	-	-	1	2	2	5	5	6
2000.....	589	2	3	-	2	1	-	-	1	3	2	8
1999.....	581	3	6	1	-	1	2	-	4	4	2	4





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Emphysema (J43)</b>												
All origins, both sexes												
2003.....	14,861	-	1	-	-	1	3	2	3	21	66	157
2002.....	15,489	1	3	-	-	1	1	2	8	21	73	156
2001.....	16,242	3	-	-	1	1	1	1	6	24	51	141
2000.....	16,700	3	2	-	-	1	4	6	5	23	59	162
1999.....	17,787	1	-	1	-	-	1	1	8	19	78	136
All origins, male												
2003.....	7,549	-	1	-	-	-	2	2	1	14	35	105
2002.....	8,084	-	3	-	-	-	-	1	7	14	52	95
2001.....	8,415	1	-	-	1	1	-	-	4	17	35	89
2000.....	8,671	3	-	-	-	1	4	1	4	15	31	101
1999.....	9,480	-	-	-	-	-	1	-	6	9	40	85
All origins, female												
2003.....	7,312	-	-	-	-	1	1	-	2	7	31	52
2002.....	7,405	1	-	-	-	1	1	1	1	7	21	61
2001.....	7,827	2	-	-	-	-	1	1	2	7	16	52
2000.....	8,029	-	2	-	-	-	-	5	1	8	28	61
1999.....	8,307	1	-	1	-	-	-	1	2	10	38	51
Hispanic, both sexes												
2003.....	312	-	-	-	-	-	-	-	-	1	5	4
2002.....	295	-	-	-	-	-	-	-	2	-	1	1
2001.....	286	1	-	-	-	-	-	-	-	-	-	5
2000.....	266	-	-	-	-	-	1	-	-	1	2	3
1999.....	292	-	-	1	-	-	-	-	2	2	4	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Emphysema (J43)</b>												
<b>Hispanic, male</b>												
2003.....	184	-	-	-	-	-	-	-	-	1	4	3
2002.....	168	-	-	-	-	-	-	-	1	-	1	1
2001.....	175	-	-	-	-	-	-	-	-	-	-	2
2000.....	167	-	-	-	-	-	1	-	-	1	2	2
1999.....	183	-	-	-	-	-	-	-	1	-	2	4
<b>Hispanic, female</b>												
2003.....	128	-	-	-	-	-	-	-	-	-	1	1
2002.....	127	-	-	-	-	-	-	-	1	-	-	-
2001.....	111	1	-	-	-	-	-	-	-	-	-	3
2000.....	99	-	-	-	-	-	-	-	-	-	-	1
1999.....	109	-	-	1	-	-	-	-	1	2	2	1
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	14,514	-	1	-	-	1	3	2	3	20	61	152
2002.....	15,161	1	3	-	-	1	1	2	6	21	72	154
2001.....	15,914	2	-	-	1	1	1	1	6	24	50	135
2000.....	16,382	3	2	-	-	1	3	6	5	22	57	155
1999.....	17,441	1	-	-	-	-	1	1	6	16	72	130
<b>Non-Hispanic 1/, male</b>												
2003.....	7,342	-	1	-	-	-	2	2	1	13	31	101
2002.....	7,895	-	3	-	-	-	-	1	6	14	51	93
2001.....	8,213	1	-	-	1	1	-	-	4	17	34	86
2000.....	8,483	3	-	-	-	1	3	1	4	14	29	96
1999.....	9,260	-	-	-	-	-	1	-	5	8	37	80

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Emphysema (J43)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	7,172	-	-	-	-	1	1	-	2	7	30	51
2002.....	7,266	1	-	-	-	1	1	1	-	7	21	61
2001.....	7,701	1	-	-	-	-	1	1	2	7	16	49
2000.....	7,899	-	2	-	-	-	-	5	1	8	28	59
1999.....	8,181	1	-	-	-	-	-	1	1	8	35	50
<b>Non-Hispanic White, both sexes</b>												
2003.....	13,518	-	1	-	-	1	3	1	2	16	48	124
2002.....	14,157	1	1	-	-	1	-	2	4	17	49	127
2001.....	14,930	1	-	-	-	1	-	1	3	22	42	121
2000.....	15,301	3	-	-	-	-	1	6	4	17	43	115
1999.....	16,338	-	-	-	-	-	-	1	4	13	51	98
<b>Non-Hispanic White, male</b>												
2003.....	6,714	-	1	-	-	-	2	1	-	10	24	82
2002.....	7,241	-	1	-	-	-	-	1	4	11	36	76
2001.....	7,607	-	-	-	-	1	-	-	2	16	28	79
2000.....	7,805	3	-	-	-	-	1	1	3	10	21	67
1999.....	8,540	-	-	-	-	-	-	-	3	5	28	59
<b>Non-Hispanic White, female</b>												
2003.....	6,804	-	-	-	-	1	1	-	2	6	24	42
2002.....	6,916	1	-	-	-	1	-	1	-	6	13	51
2001.....	7,323	1	-	-	-	-	-	1	1	6	14	42
2000.....	7,496	-	-	-	-	-	-	5	1	7	22	48
1999.....	7,798	-	-	-	-	-	-	1	1	8	23	39

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Emphysema (J43)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	805	-	-	-	-	-	-	1	1	3	12	27
2002.....	842	-	1	-	-	-	1	-	2	4	22	26
2001.....	800	1	-	-	1	-	1	-	3	1	8	13
2000.....	889	-	1	-	-	1	2	-	1	4	13	37
1999.....	923	1	-	-	-	-	-	-	-	2	20	32
<b>Non-Hispanic Black, male</b>												
2003.....	495	-	-	-	-	-	-	1	1	2	6	18
2002.....	538	-	1	-	-	-	-	-	2	3	15	16
2001.....	480	1	-	-	1	-	-	-	2	1	6	6
2000.....	551	-	-	-	-	1	2	-	1	3	8	26
1999.....	594	-	-	-	-	-	-	-	-	2	9	21
<b>Non-Hispanic Black, female</b>												
2003.....	310	-	-	-	-	-	-	-	-	1	6	9
2002.....	304	-	-	-	-	-	1	-	-	1	7	10
2001.....	320	-	-	-	-	-	1	-	1	-	2	7
2000.....	338	-	1	-	-	-	-	-	-	1	5	11
1999.....	329	1	-	-	-	-	-	-	-	-	11	11
<b>Origin not stated 2/, both sexes</b>												
2003.....	35	-	-	-	-	-	-	-	-	-	-	1
2002.....	33	-	-	-	-	-	-	-	-	-	-	1
2001.....	42	-	-	-	-	-	-	-	-	-	1	1
2000.....	52	-	-	-	-	-	-	-	-	-	-	4
1999.....	54	-	-	-	-	-	-	-	-	1	2	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Asthma (J45-J46)												
All origins, both sexes												
2003.....	4,099	7	37	35	75	70	88	98	129	152	259	317
2002.....	4,261	4	43	35	88	84	85	102	133	207	265	303
2001.....	4,269	10	31	41	58	60	80	88	145	207	248	279
2000.....	4,487	8	32	43	86	78	89	105	144	183	275	318
1999.....	4,657	5	27	40	86	95	87	121	141	206	241	312
All origins, male												
2003.....	1,493	4	29	23	48	40	51	58	69	60	100	127
2002.....	1,580	3	30	20	49	63	47	63	66	92	106	113
2001.....	1,479	6	22	24	33	35	45	49	67	93	102	103
2000.....	1,632	6	20	22	52	52	50	64	78	81	115	128
1999.....	1,620	3	16	25	60	53	55	64	64	94	90	96
All origins, female												
2003.....	2,606	3	8	12	27	30	37	40	60	92	159	190
2002.....	2,681	1	13	15	39	21	38	39	67	115	159	190
2001.....	2,790	4	9	17	25	25	35	39	78	114	146	176
2000.....	2,855	2	12	21	34	26	39	41	66	102	160	190
1999.....	3,037	2	11	15	26	42	32	57	77	112	151	216
Hispanic, both sexes												
2003.....	299	-	6	4	11	7	7	13	10	13	14	35
2002.....	287	1	6	6	7	5	6	8	15	18	24	25
2001.....	274	2	5	2	6	5	3	8	9	16	15	20
2000.....	292	1	5	3	10	6	8	4	12	13	16	19
1999.....	320	-	3	4	10	8	6	12	9	24	22	25

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Asthma (J45-J46)</b>												
<b>Hispanic, male</b>												
2003.....	131	-	5	2	8	4	4	8	4	4	9	16
2002.....	114	1	3	4	4	4	5	7	7	8	6	14
2001.....	103	2	3	1	4	4	2	5	5	8	5	6
2000.....	129	1	2	1	7	5	6	2	7	7	7	8
1999.....	119	-	-	2	7	4	4	6	5	9	13	9
<b>Hispanic, female</b>												
2003.....	168	-	1	2	3	3	3	5	6	9	5	19
2002.....	173	-	3	2	3	1	1	1	8	10	18	11
2001.....	171	-	2	1	2	1	1	3	4	8	10	14
2000.....	163	-	3	2	3	1	2	2	5	6	9	11
1999.....	201	-	3	2	3	4	2	6	4	15	9	16
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	3,786	7	30	30	64	62	81	85	119	139	243	282
2002.....	3,950	3	37	29	79	78	79	94	116	188	237	276
2001.....	3,976	8	26	39	51	55	77	80	135	190	231	258
2000.....	4,164	7	27	39	76	72	80	98	131	170	258	298
1999.....	4,324	5	24	36	76	87	81	109	129	182	218	286
<b>Non-Hispanic 1/, male</b>												
2003.....	1,356	4	23	21	40	35	47	50	65	56	89	111
2002.....	1,448	2	27	16	43	58	42	56	57	83	96	98
2001.....	1,365	4	19	23	29	31	43	44	62	84	96	96
2000.....	1,486	5	18	21	45	47	44	59	71	74	107	119
1999.....	1,495	3	16	23	53	49	51	58	58	85	77	86

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Asthma (J45-J46)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	2,430	3	7	9	24	27	34	35	54	83	154	171
2002.....	2,502	1	10	13	36	20	37	38	59	105	141	178
2001.....	2,611	4	7	16	22	24	34	36	73	106	135	162
2000.....	2,678	2	9	18	31	25	36	39	60	96	151	179
1999.....	2,829	2	8	13	23	38	30	51	71	97	141	200
<b>Non-Hispanic White, both sexes</b>												
2003.....	2,589	4	8	8	26	24	40	46	62	77	138	166
2002.....	2,720	1	14	9	25	35	44	49	72	103	119	161
2001.....	2,717	4	7	15	22	26	33	35	62	84	135	144
2000.....	2,836	4	13	14	26	30	45	44	70	83	144	158
1999.....	3,011	1	8	17	30	51	41	44	63	99	123	167
<b>Non-Hispanic White, male</b>												
2003.....	831	4	4	5	14	12	24	21	27	28	50	64
2002.....	895	-	12	6	14	23	20	24	35	35	45	54
2001.....	834	3	4	9	12	11	18	20	22	40	51	52
2000.....	920	2	7	7	13	19	25	26	34	30	60	70
1999.....	930	1	5	10	21	28	24	22	26	44	36	45
<b>Non-Hispanic White, female</b>												
2003.....	1,758	-	4	3	12	12	16	25	35	49	88	102
2002.....	1,825	1	2	3	11	12	24	25	37	68	74	107
2001.....	1,883	1	3	6	10	15	15	15	40	44	84	92
2000.....	1,916	2	6	7	13	11	20	18	36	53	84	88
1999.....	2,081	-	3	7	9	23	17	22	37	55	87	122

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Asthma (J45-J46)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	1,019	3	20	22	37	35	39	37	53	57	98	107
2002.....	1,083	2	21	17	52	40	34	41	43	82	114	103
2001.....	1,092	3	19	23	28	27	40	44	71	99	87	106
2000.....	1,145	3	14	24	45	40	33	47	55	78	100	128
1999.....	1,134	4	14	19	46	33	38	62	64	77	89	110
<b>Non-Hispanic Black, male</b>												
2003.....	439	-	18	16	25	21	21	27	36	26	35	42
2002.....	488	2	14	9	28	34	22	28	22	46	50	39
2001.....	450	-	15	14	16	19	22	24	38	41	38	42
2000.....	472	3	11	14	28	27	17	28	34	38	41	44
1999.....	474	2	10	13	32	20	26	34	32	36	36	38
<b>Non-Hispanic Black, female</b>												
2003.....	580	3	2	6	12	14	18	10	17	31	63	65
2002.....	595	-	7	8	24	6	12	13	21	36	64	64
2001.....	642	3	4	9	12	8	18	20	33	58	49	64
2000.....	673	-	3	10	17	13	16	19	21	40	59	84
1999.....	660	2	4	6	14	13	12	28	32	41	53	72
<b>Origin not stated 2/, both sexes</b>												
2003.....	14	-	1	1	-	1	-	-	-	-	2	-
2002.....	24	-	-	-	2	1	-	-	2	1	4	2
2001.....	19	-	-	-	1	-	-	-	1	1	2	1
2000.....	31	-	-	1	-	-	1	3	1	-	1	1
1999.....	13	-	-	-	-	-	-	-	3	-	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Asthma (J45-J46)

Origin not stated 2/, male

2003.....	6	-	1	-	-	1	-	-	-	-	2	-
2002.....	18	-	-	-	2	1	-	-	2	1	4	1
2001.....	11	-	-	-	-	-	-	-	-	1	1	1
2000.....	17	-	-	-	-	-	-	3	-	-	1	1
1999.....	6	-	-	-	-	-	-	-	1	-	-	1

Origin not stated 2/, female

2003.....	8	-	-	1	-	-	-	-	-	-	-	-
2002.....	6	-	-	-	-	-	-	-	-	-	-	1
2001.....	8	-	-	-	1	-	-	-	1	-	1	-
2000.....	14	-	-	1	-	-	1	-	1	-	-	-
1999.....	7	-	-	-	-	-	-	-	2	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic lower respiratory diseases (J44,J47)												
All origins, both sexes												
2003.....	106,572	4	4	1	1	11	11	13	27	120	315	785
2002.....	104,111	10	6	3	5	5	10	10	40	100	324	751
2001.....	101,543	7	4	1	3	10	14	9	35	115	303	686
2000.....	99,655	7	3	3	2	4	9	11	17	97	273	675
1999.....	100,565	10	8	5	3	9	10	20	28	67	261	617
All origins, male												
2003.....	51,321	4	2	-	1	8	7	6	17	63	148	436
2002.....	50,629	5	3	2	1	3	4	6	20	44	177	396
2001.....	49,407	4	2	-	3	5	10	4	20	58	142	387
2000.....	49,198	5	1	2	1	3	6	4	7	45	136	380
1999.....	50,817	7	3	2	3	5	5	5	10	36	136	357
All origins, female												
2003.....	55,251	-	2	1	-	3	4	7	10	57	167	349
2002.....	53,482	5	3	1	4	2	6	4	20	56	147	355
2001.....	52,136	3	2	1	-	5	4	5	15	57	161	299
2000.....	50,457	2	2	1	1	1	3	7	10	52	137	295
1999.....	49,748	3	5	3	-	4	5	15	18	31	125	260
Hispanic, both sexes												
2003.....	2,523	2	-	-	-	5	-	1	4	11	14	21
2002.....	2,425	-	2	-	-	1	1	1	4	5	19	23
2001.....	2,235	-	2	-	1	1	-	1	3	8	11	21
2000.....	2,079	2	1	2	-	-	1	2	-	6	14	22
1999.....	2,195	3	2	2	-	-	1	3	3	4	16	22

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic lower respiratory diseases (J44,J47)												
Hispanic, male												
2003.....	1,362	2	-	-	-	3	-	1	2	7	5	12
2002.....	1,323	-	-	-	-	1	1	1	2	3	11	12
2001.....	1,189	-	1	-	1	-	-	-	2	4	3	14
2000.....	1,127	1	1	1	-	-	1	1	-	2	3	11
1999.....	1,228	3	1	1	-	-	1	-	1	4	6	8
Hispanic, female												
2003.....	1,161	-	-	-	-	2	-	-	2	4	9	9
2002.....	1,102	-	2	-	-	-	-	-	2	2	8	11
2001.....	1,046	-	1	-	-	1	-	1	1	4	8	7
2000.....	952	1	-	1	-	-	-	1	-	4	11	11
1999.....	967	-	1	1	-	-	-	3	2	-	10	14
Non-Hispanic 1/, both sexes												
2003.....	103,788	2	4	1	1	6	11	12	23	109	301	761
2002.....	101,413	10	4	3	5	4	9	9	35	95	302	725
2001.....	99,030	7	2	1	2	9	14	8	32	106	289	661
2000.....	97,305	5	2	1	2	4	8	9	17	90	258	649
1999.....	98,086	7	6	3	3	9	9	17	25	63	245	593
Non-Hispanic 1/, male												
2003.....	49,820	2	2	-	1	5	7	5	15	56	143	422
2002.....	49,155	5	3	2	1	2	3	5	17	41	165	382
2001.....	48,070	4	1	-	2	5	10	4	18	54	136	371
2000.....	47,906	4	-	1	1	3	5	3	7	42	133	366
1999.....	49,436	4	2	1	3	5	4	5	9	32	130	348

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic lower respiratory diseases (J44,J47)												
Non-Hispanic 1/, female												
2003.....	53,968	-	2	1	-	1	4	7	8	53	158	339
2002.....	52,258	5	1	1	4	2	6	4	18	54	137	343
2001.....	50,960	3	1	1	-	4	4	4	14	52	153	290
2000.....	49,399	1	2	-	1	1	3	6	10	48	125	283
1999.....	48,650	3	4	2	-	4	5	12	16	31	115	245
Non-Hispanic White, both sexes												
2003.....	96,722	2	2	1	1	1	8	8	16	87	238	652
2002.....	94,430	6	2	-	5	3	9	7	19	72	251	618
2001.....	92,267	4	1	1	1	5	7	5	24	73	241	551
2000.....	90,746	4	2	1	2	2	5	7	9	68	204	541
1999.....	91,269	1	3	3	2	6	6	12	16	49	189	473
Non-Hispanic White, male												
2003.....	45,758	2	-	-	1	1	5	3	10	46	110	363
2002.....	45,172	4	2	-	1	1	3	4	12	31	138	321
2001.....	44,173	2	-	-	1	2	5	3	15	38	114	316
2000.....	44,077	4	-	1	1	1	3	2	2	31	109	300
1999.....	45,423	-	-	1	2	2	2	4	9	26	97	277
Non-Hispanic White, female												
2003.....	50,964	-	2	1	-	-	3	5	6	41	128	289
2002.....	49,258	2	-	-	4	2	6	3	7	41	113	297
2001.....	48,094	2	1	1	-	3	2	2	9	35	127	235
2000.....	46,669	-	2	-	1	2	2	5	7	37	95	241
1999.....	45,846	1	3	2	-	4	4	8	7	23	92	196







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumoconioses and chemical effects (J60-J66,J68)

Hispanic, male

2003.....	19	-	-	-	-	-	1	-	-	-	-	-
2002.....	31	-	-	-	-	-	-	-	-	-	1	1
2001.....	25	-	1	-	-	-	-	-	-	1	1	1
2000.....	26	-	-	-	-	-	-	-	1	-	1	-
1999.....	11	-	-	-	-	-	-	-	-	-	1	-

Hispanic, female

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	2	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,093	-	-	-	-	-	-	-	1	2	5	4
2002.....	1,078	2	-	-	-	-	-	-	-	-	4	4
2001.....	1,108	-	-	-	-	-	-	-	1	2	1	2
2000.....	1,133	-	-	-	-	-	-	1	1	2	4	5
1999.....	1,100	-	3	-	-	1	-	-	-	3	3	3

Non-Hispanic 1/, male

2003.....	1,045	-	-	-	-	-	-	-	1	2	4	4
2002.....	1,023	-	-	-	-	-	-	-	-	-	3	3
2001.....	1,068	-	-	-	-	-	-	-	1	2	1	1
2000.....	1,086	-	-	-	-	-	-	1	1	1	4	5
1999.....	1,068	-	3	-	-	1	-	-	-	2	2	3







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Pneumonitis due to solids and liquids (J69)</b>												
<b>All origins, both sexes</b>												
2003.....	17,335	13	10	4	12	13	33	26	43	66	100	188
2002.....	17,593	2	7	7	9	16	28	31	43	72	120	143
2001.....	17,301	12	14	5	10	10	24	31	48	61	118	142
2000.....	16,636	10	11	8	5	12	25	22	40	65	116	157
1999.....	15,268	16	12	4	12	9	23	33	45	59	107	129
<b>All origins, male</b>												
2003.....	8,702	8	5	4	5	6	25	16	26	39	50	110
2002.....	8,876	-	3	3	6	10	17	17	25	41	78	90
2001.....	8,576	6	6	3	4	5	17	15	25	42	75	81
2000.....	8,390	4	5	5	4	8	15	11	24	45	70	99
1999.....	7,755	9	3	-	6	7	15	21	26	35	62	83
<b>All origins, female</b>												
2003.....	8,633	5	5	-	7	7	8	10	17	27	50	78
2002.....	8,717	2	4	4	3	6	11	14	18	31	42	53
2001.....	8,725	6	8	2	6	5	7	16	23	19	43	61
2000.....	8,246	6	6	3	1	4	10	11	16	20	46	58
1999.....	7,513	7	9	4	6	2	8	12	19	24	45	46
<b>Hispanic, both sexes</b>												
2003.....	524	2	3	1	1	1	5	1	4	6	3	11
2002.....	547	1	-	2	1	3	4	1	2	4	6	8
2001.....	555	-	3	1	4	2	1	-	4	4	6	7
2000.....	496	3	2	1	-	1	2	3	4	6	9	11
1999.....	413	2	2	2	3	1	1	3	1	1	2	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Hispanic, male												
2003.....	263	2	2	1	-	1	4	1	2	4	2	7
2002.....	282	-	-	1	-	2	3	-	2	1	4	4
2001.....	283	-	1	1	1	-	1	-	3	2	3	4
2000.....	253	-	-	-	-	1	1	1	1	4	4	8
1999.....	195	1	1	-	1	-	-	2	1	1	-	3
Hispanic, female												
2003.....	261	-	1	-	1	-	1	-	2	2	1	4
2002.....	265	1	-	1	1	1	1	1	-	3	2	4
2001.....	272	-	2	-	3	2	-	-	1	2	3	3
2000.....	243	3	2	1	-	-	1	2	3	2	5	3
1999.....	218	1	1	2	2	1	1	1	-	-	2	1
Non-Hispanic 1/, both sexes												
2003.....	16,769	10	7	3	11	12	28	25	39	60	97	176
2002.....	17,006	1	7	5	8	13	23	30	41	68	114	135
2001.....	16,700	12	11	4	6	8	23	31	44	56	111	133
2000.....	16,094	7	9	7	5	11	23	19	36	59	107	145
1999.....	14,812	14	10	2	9	8	22	30	44	57	105	124
Non-Hispanic 1/, male												
2003.....	8,415	5	3	3	5	5	21	15	24	35	48	102
2002.....	8,565	-	3	2	6	8	13	17	23	40	74	86
2001.....	8,270	6	5	2	3	5	16	15	22	39	71	76
2000.....	8,107	4	5	5	4	7	14	10	23	41	66	90
1999.....	7,534	8	2	-	5	7	15	19	25	33	62	79

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pneumonitis due to solids and liquids (J69)												
Non-Hispanic 1/, female												
2003.....	8,354	5	4	-	6	7	7	10	15	25	49	74
2002.....	8,441	1	4	3	2	5	10	13	18	28	40	49
2001.....	8,430	6	6	2	3	3	7	16	22	17	40	57
2000.....	7,987	3	4	2	1	4	9	9	13	18	41	55
1999.....	7,278	6	8	2	4	1	7	11	19	24	43	45
Non-Hispanic White, both sexes												
2003.....	14,845	4	6	2	9	7	16	18	33	44	71	140
2002.....	15,203	1	5	4	7	10	17	26	30	51	87	111
2001.....	14,843	9	9	3	3	6	16	26	32	41	90	107
2000.....	14,343	6	7	7	4	9	15	14	26	37	91	104
1999.....	13,139	11	6	2	5	7	13	20	33	42	76	93
Non-Hispanic White, male												
2003.....	7,454	1	3	2	4	3	13	10	20	27	38	81
2002.....	7,654	-	3	2	5	6	9	16	17	30	56	70
2001.....	7,411	6	3	2	1	4	13	14	18	28	59	67
2000.....	7,214	4	4	5	3	6	10	7	14	28	57	64
1999.....	6,642	6	1	-	4	6	8	11	19	25	45	62
Non-Hispanic White, female												
2003.....	7,391	3	3	-	5	4	3	8	13	17	33	59
2002.....	7,549	1	2	2	2	4	8	10	13	21	31	41
2001.....	7,432	3	6	1	2	2	3	12	14	13	31	40
2000.....	7,129	2	3	2	1	3	5	7	12	9	34	40
1999.....	6,497	5	5	2	1	1	5	9	14	17	31	31

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Pneumonitis due to solids and liquids (J69)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	1,598	4	1	1	2	4	10	6	6	15	26	34
2002.....	1,528	-	2	1	1	2	6	4	10	11	27	20
2001.....	1,584	3	2	1	3	1	6	5	11	12	21	23
2000.....	1,507	1	2	-	1	2	7	5	9	21	16	38
1999.....	1,495	2	4	-	3	1	9	9	9	13	27	28
<b>Non-Hispanic Black, male</b>												
2003.....	776	3	-	1	1	2	6	5	4	7	10	21
2002.....	755	-	-	-	1	1	4	1	5	5	18	12
2001.....	731	-	2	-	2	-	2	1	3	8	12	7
2000.....	757	-	1	-	1	1	3	3	8	12	9	25
1999.....	768	1	1	-	1	1	7	7	5	7	15	15
<b>Non-Hispanic Black, female</b>												
2003.....	822	1	1	-	1	2	4	1	2	8	16	13
2002.....	773	-	2	1	-	1	2	3	5	6	9	8
2001.....	853	3	-	1	1	1	4	4	8	4	9	16
2000.....	750	1	1	-	-	1	4	2	1	9	7	13
1999.....	727	1	3	-	2	-	2	2	4	6	12	13
<b>Origin not stated 2/, both sexes</b>												
2003.....	42	1	-	-	-	-	-	-	-	-	-	1
2002.....	40	-	-	-	-	-	1	-	-	-	-	-
2001.....	46	-	-	-	-	-	-	-	-	1	1	2
2000.....	46	-	-	-	-	-	-	-	-	-	-	1
1999.....	43	-	-	-	-	-	-	-	-	1	-	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
All origins, both sexes												
2003.....	25,520	279	86	25	35	59	77	80	130	188	307	573
2002.....	25,039	278	85	41	43	51	75	70	115	191	337	522
2001.....	26,178	294	93	30	48	55	86	97	119	218	352	542
2000.....	25,535	273	87	36	38	60	71	99	154	231	384	527
1999.....	25,031	252	91	37	39	59	56	97	160	228	348	525
All origins, male												
2003.....	12,391	163	49	16	22	31	45	49	66	97	154	301
2002.....	12,057	164	56	21	18	29	39	31	57	101	180	263
2001.....	12,495	186	47	18	25	30	46	45	59	105	199	295
2000.....	12,391	154	45	21	17	30	43	48	79	132	203	281
1999.....	12,110	145	55	20	19	37	32	45	82	125	192	269
All origins, female												
2003.....	13,129	116	37	9	13	28	32	31	64	91	153	272
2002.....	12,982	114	29	20	25	22	36	39	58	90	157	259
2001.....	13,683	108	46	12	23	25	40	52	60	113	153	247
2000.....	13,144	119	42	15	21	30	28	51	75	99	181	246
1999.....	12,921	107	36	17	20	22	24	52	78	103	156	256
Hispanic, both sexes												
2003.....	1,374	57	14	8	8	6	10	10	17	20	32	44
2002.....	1,273	43	18	8	6	6	5	11	16	13	26	45
2001.....	1,275	41	17	2	5	2	16	4	13	14	25	42
2000.....	1,225	43	17	3	7	6	9	14	20	27	29	39
1999.....	1,157	38	10	5	3	9	6	9	14	12	33	39

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, male

2003.....	686	29	8	5	4	2	7	5	7	14	19	25
2002.....	635	21	14	4	4	4	5	5	13	9	14	23
2001.....	643	21	8	-	3	1	8	4	8	10	16	24
2000.....	608	23	8	1	3	2	6	9	10	18	16	22
1999.....	583	20	6	-	3	4	3	5	7	6	23	22

Hispanic, female

2003.....	688	28	6	3	4	4	3	5	10	6	13	19
2002.....	638	22	4	4	2	2	-	6	3	4	12	22
2001.....	632	20	9	2	2	1	8	-	5	4	9	18
2000.....	617	20	9	2	4	4	3	5	10	9	13	17
1999.....	574	18	4	5	-	5	3	4	7	6	10	17

Non-Hispanic 1/, both sexes

2003.....	24,094	221	72	17	26	53	67	70	110	168	275	529
2002.....	23,706	235	67	33	37	45	70	59	99	177	310	472
2001.....	24,835	251	76	28	43	53	70	93	105	204	324	497
2000.....	24,236	228	69	33	31	53	62	85	133	203	353	486
1999.....	23,802	211	81	32	36	50	50	88	145	216	313	484

Non-Hispanic 1/, male

2003.....	11,675	133	41	11	18	29	38	44	57	83	135	276
2002.....	11,397	143	42	17	14	25	34	26	44	92	166	238
2001.....	11,818	165	39	18	22	29	38	41	51	95	181	270
2000.....	11,752	130	37	20	14	28	37	39	69	113	185	259
1999.....	11,483	123	49	20	16	33	29	40	75	119	167	246

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
Non-Hispanic 1/, female												
2003.....	12,419	88	31	6	8	24	29	26	53	85	140	253
2002.....	12,309	92	25	16	23	20	36	33	55	85	144	234
2001.....	13,017	86	37	10	21	24	32	52	54	109	143	227
2000.....	12,484	98	32	13	17	25	25	46	64	90	168	227
1999.....	12,319	88	32	12	20	17	21	48	70	97	146	238
Non-Hispanic White, both sexes												
2003.....	21,311	113	39	13	17	34	39	46	74	109	175	359
2002.....	20,879	109	39	21	22	30	40	39	66	118	221	318
2001.....	21,835	116	44	20	32	36	46	57	65	123	227	344
2000.....	21,397	122	46	21	22	33	38	58	90	125	260	336
1999.....	20,959	111	52	20	27	38	37	56	99	143	208	335
Non-Hispanic White, male												
2003.....	10,314	64	20	10	10	15	22	29	41	56	93	200
2002.....	10,043	67	26	13	6	15	24	17	28	61	117	162
2001.....	10,433	76	20	14	17	20	26	22	30	63	139	191
2000.....	10,349	77	22	13	11	15	23	26	48	69	136	178
1999.....	10,154	66	28	13	13	27	24	34	52	82	118	171
Non-Hispanic White, female												
2003.....	10,997	49	19	3	7	19	17	17	33	53	82	159
2002.....	10,836	42	13	8	16	15	16	22	38	57	104	156
2001.....	11,402	40	24	6	15	16	20	35	35	60	88	153
2000.....	11,048	45	24	8	11	18	15	32	42	56	124	158
1999.....	10,805	45	24	7	14	11	13	22	47	61	90	164

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic Black, both sexes

2003.....	2,207	97	27	2	5	15	26	21	33	51	88	148
2002.....	2,257	112	28	12	13	15	26	18	29	51	82	138
2001.....	2,462	116	26	6	11	15	23	29	38	72	91	141
2000.....	2,336	102	20	10	8	17	22	22	36	69	83	140
1999.....	2,339	90	26	10	9	10	11	29	44	68	92	133

Non-Hispanic Black, male

2003.....	1,083	60	18	1	5	12	15	13	14	25	34	67
2002.....	1,076	67	16	4	6	10	9	7	16	26	47	66
2001.....	1,116	76	14	3	5	8	11	14	20	27	41	71
2000.....	1,126	50	14	7	3	11	12	12	18	40	44	78
1999.....	1,081	52	19	6	3	5	4	5	22	34	40	70

Non-Hispanic Black, female

2003.....	1,124	37	9	1	-	3	11	8	19	26	54	81
2002.....	1,181	45	12	8	7	5	17	11	13	25	35	72
2001.....	1,346	40	12	3	6	7	12	15	18	45	50	70
2000.....	1,210	52	6	3	5	6	10	10	18	29	39	62
1999.....	1,258	38	7	4	6	5	7	24	22	34	52	63

Origin not stated 2/, both sexes

2003.....	52	1	-	-	1	-	-	-	3	-	-	-
2002.....	60	-	-	-	-	-	-	-	-	1	1	5
2001.....	68	2	-	-	-	-	-	-	1	-	3	3
2000.....	74	2	1	-	-	1	-	-	1	1	2	2
1999.....	72	3	-	-	-	-	-	-	1	-	2	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Origin not stated 2/, male

2003.....	30	1	-	-	-	-	-	-	2	-	-	-
2002.....	25	-	-	-	-	-	-	-	-	-	-	2
2001.....	34	-	-	-	-	-	-	-	-	-	2	1
2000.....	31	1	-	-	-	-	-	-	-	1	2	-
1999.....	44	2	-	-	-	-	-	-	-	-	2	1

Origin not stated 2/, female

2003.....	22	-	-	-	1	-	-	-	1	-	-	-
2002.....	35	-	-	-	-	-	-	-	-	1	1	3
2001.....	34	2	-	-	-	-	-	-	1	-	1	2
2000.....	43	1	1	-	-	1	-	-	1	-	-	2
1999.....	28	1	-	-	-	-	-	-	1	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Peptic ulcer (K25-K28)												
All origins, both sexes												
2003.....	3,913	2	1	-	3	3	4	8	17	33	68	122
2002.....	4,079	1	1	-	1	-	-	5	17	40	73	118
2001.....	4,491	-	1	3	2	3	3	13	19	36	82	137
2000.....	4,558	-	-	-	-	4	5	3	11	42	71	130
1999.....	4,604	2	2	-	-	1	1	7	14	38	85	113
All origins, male												
2003.....	1,880	-	1	-	1	3	3	4	9	20	40	82
2002.....	1,951	-	1	-	1	-	-	2	12	28	44	71
2001.....	2,074	-	1	2	-	2	1	10	11	25	62	91
2000.....	2,201	-	-	-	-	3	4	1	5	31	48	86
1999.....	2,189	-	-	-	-	1	1	5	7	27	49	67
All origins, female												
2003.....	2,033	2	-	-	2	-	1	4	8	13	28	40
2002.....	2,128	1	-	-	-	-	-	3	5	12	29	47
2001.....	2,417	-	-	1	2	1	2	3	8	11	20	46
2000.....	2,357	-	-	-	-	1	1	2	6	11	23	44
1999.....	2,415	2	2	-	-	-	-	2	7	11	36	46
Hispanic, both sexes												
2003.....	200	1	-	-	-	1	-	3	2	1	5	13
2002.....	187	-	-	-	-	-	-	1	-	2	3	10
2001.....	166	-	-	-	-	-	1	-	-	-	1	6
2000.....	186	-	-	-	-	-	-	-	-	5	4	7
1999.....	172	-	1	-	-	-	-	4	4	1	3	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Peptic ulcer (K25-K28)</b>												
<b>Hispanic, male</b>												
2003.....	114	-	-	-	-	1	-	1	2	1	5	10
2002.....	115	-	-	-	-	-	-	1	-	1	2	9
2001.....	99	-	-	-	-	-	1	-	-	-	1	4
2000.....	97	-	-	-	-	-	-	-	-	3	3	7
1999.....	101	-	-	-	-	-	-	3	2	1	2	3
<b>Hispanic, female</b>												
2003.....	86	1	-	-	-	-	-	2	-	-	-	3
2002.....	72	-	-	-	-	-	-	-	-	1	1	1
2001.....	67	-	-	-	-	-	-	-	-	-	-	2
2000.....	89	-	-	-	-	-	-	-	-	2	1	-
1999.....	71	-	1	-	-	-	-	1	2	-	1	4
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	3,692	1	1	-	3	2	4	5	15	31	63	108
2002.....	3,879	1	1	-	1	-	-	4	17	38	70	107
2001.....	4,308	-	1	3	2	3	2	13	19	36	81	131
2000.....	4,357	-	-	-	-	4	5	3	11	37	65	123
1999.....	4,415	2	1	-	-	1	1	3	10	37	81	106
<b>Non-Hispanic 1/, male</b>												
2003.....	1,753	-	1	-	1	2	3	3	7	18	35	71
2002.....	1,829	-	1	-	1	-	-	1	12	27	42	61
2001.....	1,966	-	1	2	-	2	-	10	11	25	61	87
2000.....	2,099	-	-	-	-	3	4	1	5	28	44	79
1999.....	2,079	-	-	-	-	1	1	2	5	26	46	64

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Peptic ulcer (K25-K28)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	1,939	1	-	-	2	-	1	2	8	13	28	37
2002.....	2,050	1	-	-	-	-	-	3	5	11	28	46
2001.....	2,342	-	-	1	2	1	2	3	8	11	20	44
2000.....	2,258	-	-	-	-	1	1	2	6	9	21	44
1999.....	2,336	2	1	-	-	-	-	1	5	11	35	42
<b>Non-Hispanic White, both sexes</b>												
2003.....	3,197	-	1	-	2	1	2	3	12	20	45	77
2002.....	3,394	1	1	-	-	-	-	1	10	28	44	84
2001.....	3,766	-	1	3	1	2	-	5	11	19	58	96
2000.....	3,825	-	-	-	-	4	5	3	6	26	44	88
1999.....	3,859	1	1	-	-	1	1	2	5	20	53	73
<b>Non-Hispanic White, male</b>												
2003.....	1,471	-	1	-	1	1	1	3	4	11	24	51
2002.....	1,569	-	1	-	-	-	-	1	8	19	28	47
2001.....	1,649	-	1	2	-	1	-	4	7	12	43	61
2000.....	1,799	-	-	-	-	3	4	1	3	19	30	61
1999.....	1,783	-	-	-	-	1	1	2	1	15	26	42
<b>Non-Hispanic White, female</b>												
2003.....	1,726	-	-	-	1	-	1	-	8	9	21	26
2002.....	1,825	1	-	-	-	-	-	-	2	9	16	37
2001.....	2,117	-	-	1	1	1	-	1	4	7	15	35
2000.....	2,026	-	-	-	-	1	1	2	3	7	14	27
1999.....	2,076	1	1	-	-	-	-	-	4	5	27	31





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
All origins, both sexes												
2003.....	439	4	5	6	5	6	5	5	11	10	15	25
2002.....	480	3	-	3	4	7	5	4	6	8	19	24
2001.....	428	1	8	3	3	5	10	7	8	9	17	25
2000.....	435	3	5	4	11	5	6	3	8	11	16	18
1999.....	390	3	2	12	14	2	3	3	3	13	11	25
All origins, male												
2003.....	258	2	2	2	3	4	5	2	7	5	11	14
2002.....	275	1	-	1	2	5	2	4	5	7	10	19
2001.....	252	-	7	3	2	3	5	5	6	7	10	17
2000.....	237	2	2	1	3	2	5	3	7	5	11	12
1999.....	236	1	2	6	9	2	3	3	2	10	7	18
All origins, female												
2003.....	181	2	3	4	2	2	-	3	4	5	4	11
2002.....	205	2	-	2	2	2	3	-	1	1	9	5
2001.....	176	1	1	-	1	2	5	2	2	2	7	8
2000.....	198	1	3	3	8	3	1	-	1	6	5	6
1999.....	154	2	-	6	5	-	-	-	1	3	4	7
Hispanic, both sexes												
2003.....	31	1	-	4	1	-	1	-	2	-	1	4
2002.....	28	1	-	1	1	1	2	-	1	1	4	-
2001.....	30	1	4	-	1	1	1	-	1	1	-	4
2000.....	25	1	1	-	4	3	2	-	2	1	-	1
1999.....	29	1	-	5	3	-	2	-	-	4	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Hispanic, male												
2003.....	20	1	-	1	-	-	1	-	2	-	1	2
2002.....	16	-	-	1	1	1	-	-	1	1	1	-
2001.....	16	-	3	-	1	-	-	-	-	-	-	2
2000.....	16	-	1	-	3	1	2	-	2	-	-	1
1999.....	20	1	-	2	2	-	2	-	-	4	-	1
Hispanic, female												
2003.....	11	-	-	3	1	-	-	-	-	-	-	2
2002.....	12	1	-	-	-	-	2	-	-	-	3	-
2001.....	14	1	1	-	-	1	1	-	1	1	-	2
2000.....	9	1	-	-	1	2	-	-	-	1	-	-
1999.....	9	-	-	3	1	-	-	-	-	-	-	1
Non-Hispanic 1/, both sexes												
2003.....	407	3	5	2	4	6	4	5	9	10	14	20
2002.....	451	2	-	2	3	6	3	4	5	7	15	24
2001.....	398	-	4	3	2	4	9	7	7	8	17	21
2000.....	410	2	4	4	7	2	4	3	6	10	16	17
1999.....	358	2	2	7	10	2	1	3	3	9	11	22
Non-Hispanic 1/, male												
2003.....	237	1	2	1	3	4	4	2	5	5	10	11
2002.....	258	1	-	-	1	4	2	4	4	6	9	19
2001.....	236	-	4	3	1	3	5	5	6	7	10	15
2000.....	221	2	1	1	-	1	3	3	5	5	11	11
1999.....	213	-	2	4	6	2	1	3	2	6	7	16

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of appendix (K35-K38)												
Non-Hispanic 1/, female												
2003.....	170	2	3	1	1	2	-	3	4	5	4	9
2002.....	193	1	-	2	2	2	1	-	1	1	6	5
2001.....	162	-	-	-	1	1	4	2	1	1	7	6
2000.....	189	-	3	3	7	1	1	-	1	5	5	6
1999.....	145	2	-	3	4	-	-	-	1	3	4	6
Non-Hispanic White, both sexes												
2003.....	323	2	3	-	1	4	2	3	5	6	7	14
2002.....	377	1	-	1	1	3	1	3	4	4	9	17
2001.....	313	-	1	2	1	4	4	3	4	6	10	11
2000.....	323	2	3	3	6	2	1	1	1	6	11	9
1999.....	284	2	-	3	7	1	-	3	1	6	9	13
Non-Hispanic White, male												
2003.....	192	1	2	-	-	3	2	1	2	5	6	8
2002.....	210	-	-	-	-	1	1	3	3	3	8	13
2001.....	182	-	1	2	-	3	2	2	3	6	6	8
2000.....	167	2	1	1	-	1	1	1	1	3	6	6
1999.....	163	-	-	2	4	1	-	3	-	4	5	9
Non-Hispanic White, female												
2003.....	131	1	1	-	1	1	-	2	3	1	1	6
2002.....	167	1	-	1	1	2	-	-	1	1	1	4
2001.....	131	-	-	-	1	1	2	1	1	-	4	3
2000.....	156	-	2	2	6	1	-	-	-	3	5	3
1999.....	121	2	-	1	3	-	-	-	1	2	4	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of appendix (K35-K38)

Non-Hispanic Black, both sexes

2003.....	70	-	2	2	3	2	2	2	3	4	7	5
2002.....	61	1	-	1	2	3	2	1	1	-	5	7
2001.....	72	-	2	1	1	-	5	3	3	2	6	8
2000.....	77	-	-	1	1	-	2	2	5	4	5	8
1999.....	67	-	2	4	3	-	1	-	2	3	2	8

Non-Hispanic Black, male

2003.....	37	-	-	1	3	1	2	1	2	-	4	3
2002.....	40	1	-	-	1	3	1	1	1	-	1	6
2001.....	47	-	2	1	1	-	3	3	3	1	4	6
2000.....	48	-	-	-	-	-	1	2	4	2	5	5
1999.....	47	-	2	2	2	-	1	-	2	2	2	7

Non-Hispanic Black, female

2003.....	33	-	2	1	-	1	-	1	1	4	3	2
2002.....	21	-	-	1	1	-	1	-	-	-	4	1
2001.....	25	-	-	-	-	-	2	-	-	1	2	2
2000.....	29	-	-	1	1	-	1	-	1	2	-	3
1999.....	20	-	-	2	1	-	-	-	-	1	-	1

Origin not stated 2/, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	1
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	3	-	-	-	1	-	-	-	-	-	-	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hernia (K40-K46)</b>												
All origins, both sexes												
2003.....	1,613	37	6	4	2	-	3	1	4	7	21	45
2002.....	1,595	29	3	4	3	2	3	4	7	14	21	33
2001.....	1,522	43	2	1	2	2	1	6	9	13	21	24
2000.....	1,522	46	4	1	2	2	3	2	3	13	18	26
1999.....	1,464	44	3	1	1	1	2	6	10	15	24	16
All origins, male												
2003.....	668	20	4	2	2	-	1	1	3	6	10	24
2002.....	599	18	3	3	2	1	3	2	5	8	9	16
2001.....	587	28	1	-	-	1	-	4	6	5	11	13
2000.....	586	28	3	1	2	1	2	-	1	8	5	13
1999.....	597	26	2	1	-	1	2	5	3	10	13	12
All origins, female												
2003.....	945	17	2	2	-	-	2	-	1	1	11	21
2002.....	996	11	-	1	1	1	-	2	2	6	12	17
2001.....	935	15	1	1	2	1	1	2	3	8	10	11
2000.....	936	18	1	-	-	1	1	2	2	5	13	13
1999.....	867	18	1	-	1	-	-	1	7	5	11	4
Hispanic, both sexes												
2003.....	88	3	2	-	-	-	-	-	-	-	2	6
2002.....	81	3	-	1	1	-	2	-	1	1	4	2
2001.....	92	9	-	-	-	1	1	1	-	3	4	3
2000.....	85	9	-	-	1	1	-	-	1	1	-	2
1999.....	66	3	1	-	-	-	-	-	-	2	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hernia (K40-K46)</b>												
Hispanic, male												
2003.....	38	2	2	-	-	-	-	-	-	-	1	3
2002.....	35	2	-	1	1	-	2	-	1	-	2	2
2001.....	32	7	-	-	-	1	-	-	-	1	2	1
2000.....	33	6	-	-	1	-	-	-	1	1	-	2
1999.....	27	2	1	-	-	-	-	-	-	2	-	1
Hispanic, female												
2003.....	50	1	-	-	-	-	-	-	-	-	1	3
2002.....	46	1	-	-	-	-	-	-	-	1	2	-
2001.....	60	2	-	-	-	-	1	1	-	2	2	2
2000.....	52	3	-	-	-	1	-	-	-	-	-	-
1999.....	39	1	-	-	-	-	-	-	-	-	1	-
Non-Hispanic 1/, both sexes												
2003.....	1,518	34	4	4	2	-	3	1	4	7	19	38
2002.....	1,504	26	3	3	2	2	1	4	6	12	17	31
2001.....	1,425	33	2	1	2	1	-	5	9	10	17	21
2000.....	1,431	37	4	1	1	1	3	2	2	12	18	23
1999.....	1,390	41	2	1	1	1	2	6	10	13	23	14
Non-Hispanic 1/, male												
2003.....	626	18	2	2	2	-	1	1	3	6	9	20
2002.....	558	16	3	2	1	1	1	2	4	7	7	14
2001.....	552	20	1	-	-	-	-	4	6	4	9	12
2000.....	550	22	3	1	1	1	2	-	-	7	5	11
1999.....	566	24	1	1	-	1	2	5	3	8	13	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Hernia (K40-K46)</b>												
Non-Hispanic 1/, female												
2003.....	892	16	2	2	-	-	2	-	1	1	10	18
2002.....	946	10	-	1	1	1	-	2	2	5	10	17
2001.....	873	13	1	1	2	1	-	1	3	6	8	9
2000.....	881	15	1	-	-	-	1	2	2	5	13	12
1999.....	824	17	1	-	1	-	-	1	7	5	10	4
Non-Hispanic White, both sexes												
2003.....	1,327	26	1	2	1	-	2	1	2	4	14	30
2002.....	1,343	21	2	3	2	2	1	2	3	9	11	22
2001.....	1,248	22	2	-	1	-	-	5	7	7	11	17
2000.....	1,266	27	2	1	1	1	1	2	1	8	13	17
1999.....	1,227	31	2	1	1	1	1	4	6	12	15	9
Non-Hispanic White, male												
2003.....	544	14	1	1	1	-	1	1	1	3	7	15
2002.....	493	12	2	2	1	1	1	-	2	5	6	11
2001.....	469	11	1	-	-	-	-	4	4	4	6	8
2000.....	486	17	2	1	1	1	-	-	-	5	4	7
1999.....	505	18	1	1	-	1	1	3	2	7	11	7
Non-Hispanic White, female												
2003.....	783	12	-	1	-	-	1	-	1	1	7	15
2002.....	850	9	-	1	1	1	-	2	1	4	5	11
2001.....	779	11	1	-	1	-	-	1	3	3	5	9
2000.....	780	10	-	-	-	-	1	2	1	3	9	10
1999.....	722	13	1	-	1	-	-	1	4	5	4	2





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>												
<b>All origins, both sexes</b>												
2003.....	27,503	2	-	-	-	3	13	78	280	882	2,138	3,457
2002.....	27,257	6	3	1	5	7	24	77	297	929	2,225	3,404
2001.....	27,035	3	-	1	1	6	20	77	310	1,108	2,228	3,557
2000.....	26,552	3	3	-	3	7	29	88	327	1,083	2,288	3,374
1999.....	26,259	5	2	1	1	9	17	70	337	1,103	2,199	3,228
<b>All origins, male</b>												
2003.....	17,912	2	-	-	-	1	7	54	198	574	1,459	2,510
2002.....	17,401	6	1	1	2	3	11	46	190	605	1,474	2,442
2001.....	17,393	1	-	1	1	2	10	53	206	713	1,493	2,581
2000.....	17,214	3	1	-	2	3	16	63	217	716	1,598	2,515
1999.....	17,115	4	-	1	-	3	8	41	216	716	1,598	2,446
<b>All origins, female</b>												
2003.....	9,591	-	-	-	-	2	6	24	82	308	679	947
2002.....	9,856	-	2	-	3	4	13	31	107	324	751	962
2001.....	9,642	2	-	-	-	4	10	24	104	395	735	976
2000.....	9,338	-	2	-	1	4	13	25	110	367	690	859
1999.....	9,144	1	2	-	1	6	9	29	121	387	601	782
<b>Hispanic, both sexes</b>												
2003.....	3,382	1	-	-	-	1	1	13	87	156	313	464
2002.....	3,409	1	1	1	1	2	7	18	63	162	345	478
2001.....	3,301	-	-	-	-	1	4	16	65	187	343	480
2000.....	3,187	-	1	-	-	-	7	16	76	170	331	440
1999.....	2,972	-	-	-	-	4	4	7	73	190	328	423

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>												
<b>Hispanic, male</b>												
2003.....	2,387	1	-	-	-	1	1	12	67	133	247	389
2002.....	2,437	1	-	1	-	-	5	13	49	127	283	397
2001.....	2,410	-	-	-	-	-	2	15	52	150	275	403
2000.....	2,312	-	-	-	-	-	5	14	68	134	265	365
1999.....	2,143	-	-	-	-	3	2	6	56	154	267	354
<b>Hispanic, female</b>												
2003.....	995	-	-	-	-	-	-	1	20	23	66	75
2002.....	972	-	1	-	1	2	2	5	14	35	62	81
2001.....	891	-	-	-	-	1	2	1	13	37	68	77
2000.....	875	-	1	-	-	-	2	2	8	36	66	75
1999.....	829	-	-	-	-	1	2	1	17	36	61	69
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	24,030	1	-	-	-	2	12	65	192	725	1,816	2,979
2002.....	23,733	5	2	-	4	5	17	58	233	760	1,872	2,907
2001.....	23,629	3	-	-	1	5	16	61	244	919	1,877	3,064
2000.....	23,251	3	2	-	3	7	22	72	247	909	1,947	2,916
1999.....	23,185	5	2	1	1	5	13	63	263	909	1,858	2,791
<b>Non-Hispanic 1/, male</b>												
2003.....	15,452	1	-	-	-	-	6	42	130	441	1,205	2,108
2002.....	14,876	5	1	-	2	3	6	32	140	474	1,184	2,029
2001.....	14,914	1	-	-	1	2	8	38	154	563	1,214	2,170
2000.....	14,814	3	1	-	2	3	11	49	146	578	1,325	2,137
1999.....	14,896	4	-	1	-	-	6	35	159	560	1,321	2,080

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	8,578	-	-	-	-	2	6	23	62	284	611	871
2002.....	8,857	-	1	-	2	2	11	26	93	286	688	878
2001.....	8,715	2	-	-	-	3	8	23	90	356	663	894
2000.....	8,437	-	1	-	1	4	11	23	101	331	622	779
1999.....	8,289	1	2	-	1	5	7	28	104	349	537	711
<b>Non-Hispanic White, both sexes</b>												
2003.....	20,602	-	-	-	-	1	6	40	146	573	1,495	2,444
2002.....	20,353	4	2	-	3	5	12	37	167	603	1,545	2,353
2001.....	20,065	3	-	-	1	1	4	35	192	733	1,479	2,478
2000.....	19,739	-	2	-	2	2	12	42	175	702	1,538	2,365
1999.....	19,602	4	-	-	1	3	4	45	198	712	1,436	2,170
<b>Non-Hispanic White, male</b>												
2003.....	13,254	-	-	-	-	-	3	26	100	354	1,012	1,756
2002.....	12,800	4	1	-	2	3	5	21	109	376	995	1,682
2001.....	12,609	1	-	-	1	-	4	23	122	443	960	1,780
2000.....	12,640	-	1	-	1	1	5	33	107	453	1,065	1,785
1999.....	12,613	4	-	-	-	-	3	29	122	445	1,033	1,647
<b>Non-Hispanic White, female</b>												
2003.....	7,348	-	-	-	-	1	3	14	46	219	483	688
2002.....	7,553	-	1	-	1	2	7	16	58	227	550	671
2001.....	7,456	2	-	-	-	1	-	12	70	290	519	698
2000.....	7,099	-	1	-	1	1	7	9	68	249	473	580
1999.....	6,989	-	-	-	1	3	1	16	76	267	403	523

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	2,582	1	-	-	-	-	3	18	23	101	222	390
2002.....	2,545	1	-	-	1	-	5	12	37	84	229	440
2001.....	2,733	-	-	-	-	2	8	18	27	117	303	463
2000.....	2,705	2	-	-	1	4	8	20	36	142	304	452
1999.....	2,793	-	2	1	-	2	8	16	38	129	324	507
<b>Non-Hispanic Black, male</b>												
2003.....	1,696	1	-	-	-	-	2	11	14	58	130	261
2002.....	1,592	1	-	-	-	-	1	7	19	49	131	274
2001.....	1,819	-	-	-	-	1	1	10	15	77	196	319
2000.....	1,709	2	-	-	1	1	5	11	15	90	189	289
1999.....	1,812	-	-	1	-	-	3	5	22	75	218	357
<b>Non-Hispanic Black, female</b>												
2003.....	886	-	-	-	-	-	1	7	9	43	92	129
2002.....	953	-	-	-	1	-	4	5	18	35	98	166
2001.....	914	-	-	-	-	1	7	8	12	40	107	144
2000.....	996	-	-	-	-	3	3	9	21	52	115	163
1999.....	981	-	2	-	-	2	5	11	16	54	106	150
<b>Origin not stated 2/, both sexes</b>												
2003.....	91	-	-	-	-	-	-	-	1	1	9	14
2002.....	115	-	-	-	-	-	-	1	1	7	8	19
2001.....	105	-	-	1	-	-	-	-	1	2	8	13
2000.....	114	-	-	-	-	-	-	-	4	4	10	18
1999.....	102	-	-	-	-	-	-	-	1	4	13	14

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Origin not stated 2/, male

2003.....	73	-	-	-	-	-	-	-	1	-	7	13
2002.....	88	-	-	-	-	-	-	1	1	4	7	16
2001.....	69	-	-	1	-	-	-	-	-	-	4	8
2000.....	88	-	-	-	-	-	-	-	3	4	8	13
1999.....	76	-	-	-	-	-	-	-	1	2	10	12

Origin not stated 2/, female

2003.....	18	-	-	-	-	-	-	-	-	1	2	1
2002.....	27	-	-	-	-	-	-	-	-	3	1	3
2001.....	36	-	-	-	-	-	-	-	1	2	4	5
2000.....	26	-	-	-	-	-	-	-	1	-	2	5
1999.....	26	-	-	-	-	-	-	-	-	2	3	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Alcoholic liver disease (K70)</b>												
<b>All origins, both sexes</b>												
2003.....	12,360	-	-	-	-	1	5	49	194	612	1,359	2,050
2002.....	12,121	-	1	-	-	-	8	43	199	607	1,364	1,998
2001.....	12,207	-	-	-	-	1	7	45	219	745	1,400	2,121
2000.....	12,109	-	-	-	-	-	8	56	223	741	1,485	2,000
1999.....	11,958	-	-	-	-	-	8	37	222	763	1,423	1,965
<b>All origins, male</b>												
2003.....	9,104	-	-	-	-	-	3	39	143	402	915	1,524
2002.....	8,814	-	1	-	-	-	6	32	130	392	913	1,450
2001.....	8,932	-	-	-	-	-	5	31	148	492	954	1,563
2000.....	8,985	-	-	-	-	-	5	45	157	486	1,047	1,510
1999.....	8,922	-	-	-	-	-	6	24	141	491	1,029	1,501
<b>All origins, female</b>												
2003.....	3,256	-	-	-	-	1	2	10	51	210	444	526
2002.....	3,307	-	-	-	-	-	2	11	69	215	451	548
2001.....	3,275	-	-	-	-	1	2	14	71	253	446	558
2000.....	3,124	-	-	-	-	-	3	11	66	255	438	490
1999.....	3,036	-	-	-	-	-	2	13	81	272	394	464
<b>Hispanic, both sexes</b>												
2003.....	1,712	-	-	-	-	-	1	10	57	120	212	287
2002.....	1,679	-	-	-	-	-	4	12	41	112	209	282
2001.....	1,729	-	-	-	-	-	-	14	51	123	218	308
2000.....	1,680	-	-	-	-	-	4	13	53	122	231	273
1999.....	1,547	-	-	-	-	-	2	4	46	128	214	270

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Alcoholic liver disease (K70)</b>												
<b>Hispanic, male</b>												
2003.....	1,425	-	-	-	-	-	1	9	44	106	167	253
2002.....	1,418	-	-	-	-	-	4	12	35	94	177	236
2001.....	1,457	-	-	-	-	-	-	13	42	107	175	265
2000.....	1,442	-	-	-	-	-	4	12	50	94	188	237
1999.....	1,294	-	-	-	-	-	2	4	37	101	175	232
<b>Hispanic, female</b>												
2003.....	287	-	-	-	-	-	-	1	13	14	45	34
2002.....	261	-	-	-	-	-	-	-	6	18	32	46
2001.....	272	-	-	-	-	-	-	1	9	16	43	43
2000.....	238	-	-	-	-	-	-	1	3	28	43	36
1999.....	253	-	-	-	-	-	-	-	9	27	39	38
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	10,599	-	-	-	-	1	4	39	137	491	1,142	1,754
2002.....	10,377	-	1	-	-	-	4	31	157	490	1,149	1,703
2001.....	10,423	-	-	-	-	1	7	31	167	621	1,180	1,806
2000.....	10,370	-	-	-	-	-	4	43	168	616	1,249	1,713
1999.....	10,353	-	-	-	-	-	6	33	175	631	1,200	1,686
<b>Non-Hispanic 1/, male</b>												
2003.....	7,638	-	-	-	-	-	2	30	99	296	745	1,263
2002.....	7,342	-	1	-	-	-	2	20	94	296	731	1,203
2001.....	7,431	-	-	-	-	-	5	18	106	385	777	1,294
2000.....	7,492	-	-	-	-	-	1	33	106	389	854	1,262
1999.....	7,580	-	-	-	-	-	4	20	103	388	847	1,261

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Alcoholic liver disease (K70)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	2,961	-	-	-	-	1	2	9	38	195	397	491
2002.....	3,035	-	-	-	-	-	2	11	63	194	418	500
2001.....	2,992	-	-	-	-	1	2	13	61	236	403	512
2000.....	2,878	-	-	-	-	-	3	10	62	227	395	451
1999.....	2,773	-	-	-	-	-	2	13	72	243	353	425
<b>Non-Hispanic White, both sexes</b>												
2003.....	8,902	-	-	-	-	-	3	27	104	379	946	1,456
2002.....	8,713	-	1	-	-	-	1	19	114	386	949	1,408
2001.....	8,635	-	-	-	-	-	2	18	128	491	940	1,504
2000.....	8,556	-	-	-	-	-	2	29	121	474	980	1,388
1999.....	8,470	-	-	-	-	-	3	27	128	492	922	1,335
<b>Non-Hispanic White, male</b>												
2003.....	6,467	-	-	-	-	-	1	22	74	234	624	1,050
2002.....	6,242	-	1	-	-	-	1	13	74	231	607	1,012
2001.....	6,209	-	-	-	-	-	2	13	83	302	620	1,098
2000.....	6,296	-	-	-	-	-	-	24	81	303	676	1,056
1999.....	6,288	-	-	-	-	-	2	17	78	307	660	1,021
<b>Non-Hispanic White, female</b>												
2003.....	2,435	-	-	-	-	-	2	5	30	145	322	406
2002.....	2,471	-	-	-	-	-	-	6	40	155	342	396
2001.....	2,426	-	-	-	-	-	-	5	45	189	320	406
2000.....	2,260	-	-	-	-	-	2	5	40	171	304	332
1999.....	2,182	-	-	-	-	-	1	10	50	185	262	314

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Alcoholic liver disease (K70)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	1,206	-	-	-	-	-	-	6	13	67	120	196
2002.....	1,173	-	-	-	-	-	3	6	20	46	123	209
2001.....	1,331	-	-	-	-	-	1	7	19	74	170	218
2000.....	1,370	-	-	-	-	-	-	7	27	92	193	258
1999.....	1,445	-	-	-	-	-	2	4	24	84	197	285
<b>Non-Hispanic Black, male</b>												
2003.....	850	-	-	-	-	-	-	4	10	37	73	145
2002.....	783	-	-	-	-	-	1	4	10	27	76	136
2001.....	926	-	-	-	-	-	-	2	10	50	110	147
2000.....	910	-	-	-	-	-	-	5	12	59	126	164
1999.....	1,001	-	-	-	-	-	2	2	12	49	129	197
<b>Non-Hispanic Black, female</b>												
2003.....	356	-	-	-	-	-	-	2	3	30	47	51
2002.....	390	-	-	-	-	-	2	2	10	19	47	73
2001.....	405	-	-	-	-	-	1	5	9	24	60	71
2000.....	460	-	-	-	-	-	-	2	15	33	67	94
1999.....	444	-	-	-	-	-	-	2	12	35	68	88
<b>Origin not stated 2/, both sexes</b>												
2003.....	49	-	-	-	-	-	-	-	-	1	5	9
2002.....	65	-	-	-	-	-	-	-	1	5	6	13
2001.....	55	-	-	-	-	-	-	-	1	1	2	7
2000.....	59	-	-	-	-	-	-	-	2	3	5	14
1999.....	58	-	-	-	-	-	-	-	1	4	9	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcoholic liver disease (K70)

Origin not stated 2/, male

2003.....	41	-	-	-	-	-	-	-	-	-	3	8
2002.....	54	-	-	-	-	-	-	-	1	2	5	11
2001.....	44	-	-	-	-	-	-	-	-	-	2	4
2000.....	51	-	-	-	-	-	-	-	1	3	5	11
1999.....	48	-	-	-	-	-	-	-	1	2	7	8

Origin not stated 2/, female

2003.....	8	-	-	-	-	-	-	-	-	1	2	1
2002.....	11	-	-	-	-	-	-	-	-	3	1	2
2001.....	11	-	-	-	-	-	-	-	1	1	-	3
2000.....	8	-	-	-	-	-	-	-	1	-	-	3
1999.....	10	-	-	-	-	-	-	-	-	2	2	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Other chronic liver disease and cirrhosis (K73-K74)</b>												
<b>All origins, both sexes</b>												
2003.....	15,143	2	-	-	-	2	8	29	86	270	779	1,407
2002.....	15,136	6	2	1	5	7	16	34	98	322	861	1,406
2001.....	14,828	3	-	1	1	5	13	32	91	363	828	1,436
2000.....	14,443	3	3	-	3	7	21	32	104	342	803	1,374
1999.....	14,301	5	2	1	1	9	9	33	115	340	776	1,263
<b>All origins, male</b>												
2003.....	8,808	2	-	-	-	1	4	15	55	172	544	986
2002.....	8,587	6	-	1	2	3	5	14	60	213	561	992
2001.....	8,461	1	-	1	1	2	5	22	58	221	539	1,018
2000.....	8,229	3	1	-	2	3	11	18	60	230	551	1,005
1999.....	8,193	4	-	1	-	3	2	17	75	225	569	945
<b>All origins, female</b>												
2003.....	6,335	-	-	-	-	1	4	14	31	98	235	421
2002.....	6,549	-	2	-	3	4	11	20	38	109	300	414
2001.....	6,367	2	-	-	-	3	8	10	33	142	289	418
2000.....	6,214	-	2	-	1	4	10	14	44	112	252	369
1999.....	6,108	1	2	-	1	6	7	16	40	115	207	318
<b>Hispanic, both sexes</b>												
2003.....	1,670	1	-	-	-	1	-	3	30	36	101	177
2002.....	1,730	1	1	1	1	2	3	6	22	50	136	196
2001.....	1,572	-	-	-	-	1	4	2	14	64	125	172
2000.....	1,507	-	1	-	-	-	3	3	23	48	100	167
1999.....	1,425	-	-	-	-	4	2	3	27	62	114	153

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic liver disease and cirrhosis (K73-K74)												
Hispanic, male												
2003.....	962	1	-	-	-	1	-	3	23	27	80	136
2002.....	1,019	1	-	1	-	-	1	1	14	33	106	161
2001.....	953	-	-	-	-	-	2	2	10	43	100	138
2000.....	870	-	-	-	-	-	1	2	18	40	77	128
1999.....	849	-	-	-	-	3	-	2	19	53	92	122
Hispanic, female												
2003.....	708	-	-	-	-	-	-	-	7	9	21	41
2002.....	711	-	1	-	1	2	2	5	8	17	30	35
2001.....	619	-	-	-	-	1	2	-	4	21	25	34
2000.....	637	-	1	-	-	-	2	1	5	8	23	39
1999.....	576	-	-	-	-	1	2	1	8	9	22	31
Non-Hispanic 1/, both sexes												
2003.....	13,431	1	-	-	-	1	8	26	55	234	674	1,225
2002.....	13,356	5	1	-	4	5	13	27	76	270	723	1,204
2001.....	13,206	3	-	-	1	4	9	30	77	298	697	1,258
2000.....	12,881	3	2	-	3	7	18	29	79	293	698	1,203
1999.....	12,832	5	2	1	1	5	7	30	88	278	658	1,105
Non-Hispanic 1/, male												
2003.....	7,814	1	-	-	-	-	4	12	31	145	460	845
2002.....	7,534	5	-	-	2	3	4	12	46	178	453	826
2001.....	7,483	1	-	-	1	2	3	20	48	178	437	876
2000.....	7,322	3	1	-	2	3	10	16	40	189	471	875
1999.....	7,316	4	-	1	-	-	2	15	56	172	474	819

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic liver disease and cirrhosis (K73-K74)												
Non-Hispanic 1/, female												
2003.....	5,617	-	-	-	-	1	4	14	24	89	214	380
2002.....	5,822	-	1	-	2	2	9	15	30	92	270	378
2001.....	5,723	2	-	-	-	2	6	10	29	120	260	382
2000.....	5,559	-	1	-	1	4	8	13	39	104	227	328
1999.....	5,516	1	2	-	1	5	5	15	32	106	184	286
Non-Hispanic White, both sexes												
2003.....	11,700	-	-	-	-	1	3	13	42	194	549	988
2002.....	11,640	4	1	-	3	5	11	18	53	217	596	945
2001.....	11,430	3	-	-	1	1	2	17	64	242	539	974
2000.....	11,183	-	2	-	2	2	10	13	54	228	558	977
1999.....	11,132	4	-	-	1	3	1	18	70	220	514	835
Non-Hispanic White, male												
2003.....	6,787	-	-	-	-	-	2	4	26	120	388	706
2002.....	6,558	4	-	-	2	3	4	8	35	145	388	670
2001.....	6,400	1	-	-	1	-	2	10	39	141	340	682
2000.....	6,344	-	1	-	1	1	5	9	26	150	389	729
1999.....	6,325	4	-	-	-	-	1	12	44	138	373	626
Non-Hispanic White, female												
2003.....	4,913	-	-	-	-	1	1	9	16	74	161	282
2002.....	5,082	-	1	-	1	2	7	10	18	72	208	275
2001.....	5,030	2	-	-	-	1	-	7	25	101	199	292
2000.....	4,839	-	1	-	1	1	5	4	28	78	169	248
1999.....	4,807	-	-	-	1	3	-	6	26	82	141	209





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Cholelithiasis and other disorders of gallbladder (K80-K82)</b>												
All origins, both sexes												
2003.....	2,948	-	-	-	1	1	3	9	16	14	26	46
2002.....	2,979	-	-	-	-	2	7	9	12	15	39	52
2001.....	2,965	1	-	-	1	2	5	11	8	21	23	69
2000.....	2,810	-	-	1	1	2	7	6	13	22	31	43
1999.....	2,830	2	-	-	1	4	6	5	11	25	34	55
All origins, male												
2003.....	1,318	-	-	-	-	1	3	3	8	7	14	20
2002.....	1,341	-	-	-	-	-	3	4	7	7	22	35
2001.....	1,341	1	-	-	-	1	2	5	4	9	16	39
2000.....	1,244	-	-	-	1	1	3	2	5	12	15	23
1999.....	1,248	1	-	-	1	2	3	-	4	14	16	28
All origins, female												
2003.....	1,630	-	-	-	1	-	-	6	8	7	12	26
2002.....	1,638	-	-	-	-	2	4	5	5	8	17	17
2001.....	1,624	-	-	-	1	1	3	6	4	12	7	30
2000.....	1,566	-	-	1	-	1	4	4	8	10	16	20
1999.....	1,582	1	-	-	-	2	3	5	7	11	18	27
Hispanic, both sexes												
2003.....	178	-	-	-	-	1	1	2	-	4	2	6
2002.....	196	-	-	-	-	-	2	2	4	3	8	5
2001.....	197	-	-	-	-	1	1	1	2	6	2	8
2000.....	183	-	-	-	-	-	2	1	2	2	3	7
1999.....	185	-	-	-	-	-	-	1	2	6	5	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

#### Cholelithiasis and other disorders of gallbladder (K80-K82)

##### Hispanic, male

2003.....	94	-	-	-	-	1	1	-	-	2	1	4
2002.....	93	-	-	-	-	-	-	-	2	2	5	4
2001.....	92	-	-	-	-	-	-	-	1	2	1	2
2000.....	99	-	-	-	-	-	-	1	-	2	3	4
1999.....	84	-	-	-	-	-	-	-	1	3	3	2

##### Hispanic, female

2003.....	84	-	-	-	-	-	-	2	-	2	1	2
2002.....	103	-	-	-	-	-	2	2	2	1	3	1
2001.....	105	-	-	-	-	1	1	1	1	4	1	6
2000.....	84	-	-	-	-	-	2	-	2	-	-	3
1999.....	101	-	-	-	-	-	-	1	1	3	2	6

##### Non-Hispanic 1/, both sexes

2003.....	2,765	-	-	-	1	-	2	7	16	10	24	40
2002.....	2,776	-	-	-	-	2	5	7	8	12	31	47
2001.....	2,764	1	-	-	1	1	4	10	6	15	21	61
2000.....	2,619	-	-	1	1	2	5	5	11	20	28	36
1999.....	2,639	2	-	-	1	4	6	4	9	19	29	47

##### Non-Hispanic 1/, male

2003.....	1,223	-	-	-	-	-	2	3	8	5	13	16
2002.....	1,244	-	-	-	-	-	3	4	5	5	17	31
2001.....	1,248	1	-	-	-	1	2	5	3	7	15	37
2000.....	1,141	-	-	-	1	1	3	1	5	10	12	19
1999.....	1,163	1	-	-	1	2	3	-	3	11	13	26

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Cholelithiasis and other disorders of gallbladder (K80-K82)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	1,542	-	-	-	1	-	-	4	8	5	11	24
2002.....	1,532	-	-	-	-	2	2	3	3	7	14	16
2001.....	1,516	-	-	-	1	-	2	5	3	8	6	24
2000.....	1,478	-	-	1	-	1	2	4	6	10	16	17
1999.....	1,476	1	-	-	-	2	3	4	6	8	16	21
<b>Non-Hispanic White, both sexes</b>												
2003.....	2,391	-	-	-	1	-	1	6	10	7	15	27
2002.....	2,420	-	-	-	-	2	3	4	6	7	26	33
2001.....	2,368	1	-	-	1	1	2	8	4	8	17	41
2000.....	2,272	-	-	-	1	1	4	2	5	15	18	20
1999.....	2,299	-	-	-	1	3	3	3	7	11	15	36
<b>Non-Hispanic White, male</b>												
2003.....	1,056	-	-	-	-	-	1	2	4	4	6	12
2002.....	1,087	-	-	-	-	-	3	2	4	3	14	20
2001.....	1,065	1	-	-	-	1	1	3	2	5	12	24
2000.....	1,012	-	-	-	1	1	2	1	3	8	9	13
1999.....	1,017	-	-	-	1	1	3	-	1	7	7	19
<b>Non-Hispanic White, female</b>												
2003.....	1,335	-	-	-	1	-	-	4	6	3	9	15
2002.....	1,333	-	-	-	-	2	-	2	2	4	12	13
2001.....	1,303	-	-	-	1	-	1	5	2	3	5	17
2000.....	1,260	-	-	-	-	-	2	1	2	7	9	7
1999.....	1,282	-	-	-	-	2	-	3	6	4	8	17





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)</b>												
All origins, both sexes												
2003.....	42,453	181	14	7	15	29	66	103	179	270	526	835
2002.....	40,974	173	14	9	12	21	63	107	162	274	475	790
2001.....	39,480	133	16	9	9	21	71	82	143	299	469	749
2000.....	37,251	162	14	11	14	32	56	80	164	278	462	687
1999.....	35,525	166	15	13	18	24	48	97	155	268	439	608
All origins, male												
2003.....	20,481	106	7	3	6	19	40	53	94	148	311	482
2002.....	19,695	109	7	5	7	10	31	44	85	148	259	454
2001.....	18,852	76	7	3	3	14	31	45	67	158	256	445
2000.....	17,811	103	7	7	10	18	25	44	85	152	277	433
1999.....	17,016	90	9	8	10	11	22	43	82	152	250	365
All origins, female												
2003.....	21,972	75	7	4	9	10	26	50	85	122	215	353
2002.....	21,279	64	7	4	5	11	32	63	77	126	216	336
2001.....	20,628	57	9	6	6	7	40	37	76	141	213	304
2000.....	19,440	59	7	4	4	14	31	36	79	126	185	254
1999.....	18,509	76	6	5	8	13	26	54	73	116	189	243
Hispanic, both sexes												
2003.....	2,170	35	1	1	3	8	11	15	15	25	51	70
2002.....	1,898	35	5	2	3	-	9	12	21	23	42	57
2001.....	1,857	18	2	2	2	4	12	10	15	24	50	51
2000.....	1,646	25	4	2	1	7	7	6	20	19	34	45
1999.....	1,584	20	3	4	5	1	5	10	11	23	39	50

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)</b>												
<b>Hispanic, male</b>												
2003.....	1,069	19	-	-	1	6	7	8	8	14	35	39
2002.....	937	21	1	1	2	-	5	3	9	12	26	29
2001.....	892	13	-	1	1	4	5	4	6	14	32	27
2000.....	805	16	3	1	1	3	4	4	7	9	27	32
1999.....	814	15	2	3	4	-	2	4	6	12	19	30
<b>Hispanic, female</b>												
2003.....	1,101	16	1	1	2	2	4	7	7	11	16	31
2002.....	961	14	4	1	1	-	4	9	12	11	16	28
2001.....	965	5	2	1	1	-	7	6	9	10	18	24
2000.....	841	9	1	1	-	4	3	2	13	10	7	13
1999.....	770	5	1	1	1	1	3	6	5	11	20	20
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	40,172	143	13	6	12	21	55	88	164	244	473	764
2002.....	38,933	138	9	7	9	21	54	94	141	250	431	728
2001.....	37,511	113	14	7	7	17	59	72	126	272	419	692
2000.....	35,509	134	10	9	13	24	49	74	143	259	427	642
1999.....	33,830	145	12	9	13	23	43	87	144	245	398	557
<b>Non-Hispanic 1/, male</b>												
2003.....	19,358	87	7	3	5	13	33	45	86	134	275	442
2002.....	18,682	88	6	4	5	10	26	41	76	135	232	422
2001.....	17,908	61	7	2	2	10	26	41	59	142	224	415
2000.....	16,958	84	4	6	9	14	21	40	78	143	249	401
1999.....	16,143	74	7	5	6	11	20	39	76	140	230	334

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	20,814	56	6	3	7	8	22	43	78	110	198	322
2002.....	20,251	50	3	3	4	11	28	53	65	115	199	306
2001.....	19,603	52	7	5	5	7	33	31	67	130	195	277
2000.....	18,551	50	6	3	4	10	28	34	65	116	178	241
1999.....	17,687	71	5	4	7	12	23	48	68	105	168	223
<b>Non-Hispanic White, both sexes</b>												
2003.....	31,497	80	9	5	6	10	28	48	80	113	227	402
2002.....	30,669	77	5	5	5	10	23	44	75	127	226	403
2001.....	29,449	58	11	3	3	11	34	35	65	142	217	352
2000.....	27,924	61	4	6	11	12	36	43	67	126	233	334
1999.....	26,418	81	10	7	6	15	21	42	74	123	196	272
<b>Non-Hispanic White, male</b>												
2003.....	15,323	52	5	2	4	8	16	25	39	60	145	241
2002.....	14,883	49	3	3	3	5	12	20	44	70	121	234
2001.....	14,331	31	4	2	1	7	14	18	27	89	122	209
2000.....	13,567	41	2	5	7	7	18	24	38	71	135	208
1999.....	12,793	46	5	4	4	9	8	20	40	71	117	162
<b>Non-Hispanic White, female</b>												
2003.....	16,174	28	4	3	2	2	12	23	41	53	82	161
2002.....	15,786	28	2	2	2	5	11	24	31	57	105	169
2001.....	15,118	27	7	1	2	4	20	17	38	53	95	143
2000.....	14,357	20	2	1	4	5	18	19	29	55	98	126
1999.....	13,625	35	5	3	2	6	13	22	34	52	79	110





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All origins, both sexes

2003.....	148	8	2	1	-	1	3	1	1	1	3	3
2002.....	166	6	5	1	1	2	1	-	1	2	2	5
2001.....	156	8	4	1	-	-	-	2	1	4	4	1
2000.....	177	8	3	1	1	-	1	-	4	1	2	5
1999.....	182	3	2	3	2	-	2	3	2	2	3	6

All origins, male

2003.....	65	7	1	-	-	1	3	1	-	-	2	2
2002.....	70	3	2	1	1	1	-	-	1	-	-	4
2001.....	73	4	3	-	-	-	-	2	-	4	2	1
2000.....	91	3	2	1	1	-	1	-	3	-	1	2
1999.....	90	3	2	1	1	-	2	2	2	-	2	3

All origins, female

2003.....	83	1	1	1	-	-	-	-	1	1	1	1
2002.....	96	3	3	-	-	1	1	-	-	2	2	1
2001.....	83	4	1	1	-	-	-	-	1	-	2	-
2000.....	86	5	1	-	-	-	-	-	1	1	1	3
1999.....	92	-	-	2	1	-	-	1	-	2	1	3

Hispanic, both sexes

2003.....	13	1	-	1	-	1	2	-	-	-	-	-
2002.....	10	-	2	-	-	-	-	-	-	1	1	-
2001.....	13	1	-	-	-	-	-	-	1	1	-	-
2000.....	9	2	1	-	-	-	-	-	-	-	-	-
1999.....	12	1	-	1	1	-	-	-	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Hispanic, male

2003.....	7	1	-	-	-	1	2	-	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	-
2001.....	7	-	-	-	-	-	-	-	-	1	-	-
2000.....	4	-	1	-	-	-	-	-	-	-	-	-
1999.....	4	1	-	-	1	-	-	-	-	-	-	-

Hispanic, female

2003.....	6	-	-	1	-	-	-	-	-	-	-	-
2002.....	6	-	2	-	-	-	-	-	-	1	1	-
2001.....	6	1	-	-	-	-	-	-	1	-	-	-
2000.....	5	2	-	-	-	-	-	-	-	-	-	-
1999.....	8	-	-	1	-	-	-	-	-	-	-	1

Non-Hispanic 1/, both sexes

2003.....	135	7	2	-	-	-	1	1	1	1	3	3
2002.....	155	6	3	1	1	2	1	-	1	1	1	5
2001.....	142	7	4	1	-	-	-	2	-	3	4	1
2000.....	168	6	2	1	1	-	1	-	4	1	2	5
1999.....	170	2	2	2	1	-	2	3	2	2	3	5

Non-Hispanic 1/, male

2003.....	58	6	1	-	-	-	1	1	-	-	2	2
2002.....	66	3	2	1	1	1	-	-	1	-	-	4
2001.....	65	4	3	-	-	-	-	2	-	3	2	1
2000.....	87	3	1	1	1	-	1	-	3	-	1	2
1999.....	86	2	2	1	-	-	2	2	2	-	2	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Non-Hispanic 1/, female

2003.....	77	1	1	-	-	-	-	-	1	1	1	1
2002.....	89	3	1	-	-	1	1	-	-	1	1	1
2001.....	77	3	1	1	-	-	-	-	-	-	2	-
2000.....	81	3	1	-	-	-	-	-	1	1	1	3
1999.....	84	-	-	1	1	-	-	1	-	2	1	2

Non-Hispanic White, both sexes

2003.....	107	5	1	-	-	-	-	-	1	1	2	3
2002.....	119	5	2	1	1	1	-	-	-	-	1	3
2001.....	116	6	3	-	-	-	-	-	-	3	4	1
2000.....	132	4	-	1	-	-	1	-	3	-	1	2
1999.....	132	2	2	2	-	-	1	1	1	2	1	3

Non-Hispanic White, male

2003.....	40	4	-	-	-	-	-	-	-	-	2	2
2002.....	52	2	1	1	1	1	-	-	-	-	-	3
2001.....	52	3	2	-	-	-	-	-	-	3	2	1
2000.....	68	3	-	1	-	-	1	-	2	-	1	2
1999.....	63	2	2	1	-	-	1	-	1	-	1	2

Non-Hispanic White, female

2003.....	67	1	1	-	-	-	-	-	1	1	-	1
2002.....	67	3	1	-	-	-	-	-	-	-	1	-
2001.....	64	3	1	-	-	-	-	-	-	-	2	-
2000.....	64	1	-	-	-	-	-	-	1	-	-	-
1999.....	69	-	-	1	-	-	-	1	-	2	-	1





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
All origins, both sexes												
2003.....	537	3	-	-	-	3	5	7	7	8	12	15
2002.....	553	2	-	1	2	2	2	10	13	13	13	28
2001.....	511	-	2	-	1	3	5	4	8	16	11	15
2000.....	556	1	-	1	3	2	7	8	9	14	20	18
1999.....	589	-	1	1	-	3	4	10	5	15	13	16
All origins, male												
2003.....	269	1	-	-	-	1	4	3	2	6	8	12
2002.....	278	1	-	-	1	1	2	4	9	7	7	10
2001.....	264	-	1	-	-	3	2	-	1	7	6	9
2000.....	284	-	-	-	1	-	2	5	6	8	12	13
1999.....	304	-	-	1	-	3	3	7	3	9	9	10
All origins, female												
2003.....	268	2	-	-	-	2	1	4	5	2	4	3
2002.....	275	1	-	1	1	1	-	6	4	6	6	18
2001.....	247	-	1	-	1	-	3	4	7	9	5	6
2000.....	272	1	-	1	2	2	5	3	3	6	8	5
1999.....	285	-	1	-	-	-	1	3	2	6	4	6
Hispanic, both sexes												
2003.....	35	1	-	-	-	1	-	1	1	-	2	2
2002.....	32	-	-	1	1	-	-	1	2	1	1	2
2001.....	34	-	1	-	-	1	-	-	1	4	2	3
2000.....	28	-	-	-	-	-	-	-	2	-	1	2
1999.....	33	-	1	-	-	-	1	2	1	-	2	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Hispanic, male

2003.....	17	-	-	-	-	-	-	1	-	-	1	2
2002.....	15	-	-	-	-	-	-	1	2	-	-	-
2001.....	17	-	-	-	-	1	-	-	-	3	2	-
2000.....	17	-	-	-	-	-	-	-	1	-	-	2
1999.....	18	-	-	-	-	-	1	2	1	-	2	1

Hispanic, female

2003.....	18	1	-	-	-	1	-	-	1	-	1	-
2002.....	17	-	-	1	1	-	-	-	-	1	1	2
2001.....	17	-	1	-	-	-	-	-	1	1	-	3
2000.....	11	-	-	-	-	-	-	-	1	-	1	-
1999.....	15	-	1	-	-	-	-	-	-	-	-	1

Non-Hispanic 1/, both sexes

2003.....	501	2	-	-	-	2	5	6	6	8	9	13
2002.....	520	2	-	-	1	2	2	8	11	12	12	26
2001.....	477	-	1	-	1	2	5	4	7	12	9	12
2000.....	528	1	-	1	3	2	7	8	7	14	19	16
1999.....	555	-	-	1	-	3	3	8	4	15	11	14

Non-Hispanic 1/, male

2003.....	251	1	-	-	-	1	4	2	2	6	6	10
2002.....	263	1	-	-	1	1	2	3	7	7	7	10
2001.....	247	-	1	-	-	2	2	-	1	4	4	9
2000.....	267	-	-	-	1	-	2	5	5	8	12	11
1999.....	285	-	-	1	-	3	2	5	2	9	7	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic 1/, female

2003.....	250	1	-	-	-	1	1	4	4	2	3	3
2002.....	257	1	-	-	-	1	-	5	4	5	5	16
2001.....	230	-	-	-	1	-	3	4	6	8	5	3
2000.....	261	1	-	1	2	2	5	3	2	6	7	5
1999.....	270	-	-	-	-	-	1	3	2	6	4	5

Non-Hispanic White, both sexes

2003.....	394	2	-	-	-	-	2	4	2	8	5	10
2002.....	419	-	-	-	-	2	1	5	7	10	10	17
2001.....	384	-	1	-	-	1	5	3	5	4	5	5
2000.....	411	-	-	-	3	1	6	4	5	7	12	13
1999.....	436	-	-	1	-	2	1	6	2	8	5	12

Non-Hispanic White, male

2003.....	196	1	-	-	-	-	1	1	1	6	4	7
2002.....	218	-	-	-	-	1	1	2	5	6	5	8
2001.....	200	-	1	-	-	1	2	-	1	2	1	3
2000.....	209	-	-	-	1	-	1	3	3	5	7	9
1999.....	227	-	-	1	-	2	-	4	1	4	4	8

Non-Hispanic White, female

2003.....	198	1	-	-	-	-	1	3	1	2	1	3
2002.....	201	-	-	-	-	1	-	3	2	4	5	9
2001.....	184	-	-	-	-	-	3	3	4	2	4	2
2000.....	202	-	-	-	2	1	5	1	2	2	5	4
1999.....	209	-	-	-	-	-	1	2	1	4	1	4





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Renal failure (N17-N19)</b>												
<b>All origins, both sexes</b>												
2003.....	41,737	170	11	6	15	25	57	95	171	261	510	813
2002.....	40,222	165	9	6	9	17	60	95	148	258	460	757
2001.....	38,784	125	9	8	8	18	64	76	133	278	454	729
2000.....	36,471	152	11	9	10	30	48	72	149	262	439	658
1999.....	34,719	163	12	9	16	21	42	83	147	250	423	586
<b>All origins, male</b>												
2003.....	20,135	98	5	3	6	17	32	49	92	142	301	465
2002.....	19,341	105	5	3	5	8	29	39	75	141	252	440
2001.....	18,509	72	3	3	3	11	29	43	66	146	248	434
2000.....	17,416	100	5	6	8	18	22	39	76	144	263	415
1999.....	16,611	87	7	6	9	8	17	34	76	143	239	352
<b>All origins, female</b>												
2003.....	21,602	72	6	3	9	8	25	46	79	119	209	348
2002.....	20,881	60	4	3	4	9	31	56	73	117	208	317
2001.....	20,275	53	6	5	5	7	35	33	67	132	206	295
2000.....	19,055	52	6	3	2	12	26	33	73	118	176	243
1999.....	18,108	76	5	3	7	13	25	49	71	107	184	234
<b>Hispanic, both sexes</b>												
2003.....	2,121	33	1	-	3	6	9	14	14	25	49	67
2002.....	1,856	35	3	1	2	-	9	11	19	21	40	55
2001.....	1,806	17	1	2	2	3	12	10	13	18	48	47
2000.....	1,605	22	3	2	1	7	7	6	18	19	32	43
1999.....	1,535	19	2	3	4	1	4	8	10	23	37	47

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Renal failure (N17-N19)</b>												
<b>Hispanic, male</b>												
2003.....	1,044	18	-	-	1	5	5	7	8	14	34	36
2002.....	918	21	1	1	2	-	5	2	7	12	26	29
2001.....	867	13	-	1	1	3	5	4	6	9	30	27
2000.....	782	16	2	1	1	3	4	4	6	9	26	30
1999.....	790	14	2	3	3	-	1	2	5	12	17	29
<b>Hispanic, female</b>												
2003.....	1,077	15	1	-	2	1	4	7	6	11	15	31
2002.....	938	14	2	-	-	-	4	9	12	9	14	26
2001.....	939	4	1	1	1	-	7	6	7	9	18	20
2000.....	823	6	1	1	-	4	3	2	12	10	6	13
1999.....	745	5	-	-	1	1	3	6	5	11	20	18
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	39,506	134	10	6	12	19	48	81	157	235	460	745
2002.....	38,225	130	6	5	7	17	51	84	129	236	418	697
2001.....	36,867	106	8	6	6	15	52	66	118	257	406	676
2000.....	34,771	127	8	7	9	22	41	66	130	243	406	615
1999.....	33,074	143	10	6	12	20	38	75	137	227	384	538
<b>Non-Hispanic 1/, male</b>												
2003.....	19,038	80	5	3	5	12	27	42	84	128	267	428
2002.....	18,347	84	4	2	3	8	24	37	68	128	225	408
2001.....	17,591	57	3	2	2	8	24	39	58	135	218	404
2000.....	16,586	81	3	5	7	14	18	35	70	135	236	385
1999.....	15,763	72	5	3	6	8	16	32	71	131	221	322

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Renal failure (N17-N19)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	20,468	54	5	3	7	7	21	39	73	107	193	317
2002.....	19,878	46	2	3	4	9	27	47	61	108	193	289
2001.....	19,276	49	5	4	4	7	28	27	60	122	188	272
2000.....	18,185	46	5	2	2	8	23	31	60	108	170	230
1999.....	17,311	71	5	3	6	12	22	43	66	96	163	216
<b>Non-Hispanic White, both sexes</b>												
2003.....	30,973	73	7	5	6	10	26	44	77	104	219	387
2002.....	30,104	72	3	4	4	7	22	37	68	116	215	383
2001.....	28,932	52	6	3	3	10	28	32	59	135	208	344
2000.....	27,350	57	4	5	8	11	29	39	58	119	220	316
1999.....	25,831	79	8	4	6	13	19	34	70	112	190	257
<b>Non-Hispanic White, male</b>												
2003.....	15,080	47	4	2	4	8	15	24	38	54	139	231
2002.....	14,610	47	2	2	2	3	11	17	39	64	116	223
2001.....	14,076	28	1	2	1	6	12	18	26	84	119	204
2000.....	13,275	38	2	4	6	7	16	21	33	66	127	196
1999.....	12,496	44	3	2	4	7	7	16	37	67	112	152
<b>Non-Hispanic White, female</b>												
2003.....	15,893	26	3	3	2	2	11	20	39	50	80	156
2002.....	15,494	25	1	2	2	4	11	20	29	52	99	160
2001.....	14,856	24	5	1	2	4	16	14	33	51	89	140
2000.....	14,075	19	2	1	2	4	13	18	25	53	93	120
1999.....	13,335	35	5	2	2	6	12	18	33	45	78	105







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of kidney (N25,N27)

Hispanic, male

2003.....	1	-	-	-	-	-	-	-	-	-	-	1
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	1	-	-
2000.....	2	-	-	-	-	-	-	-	-	-	1	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	1
2000.....	2	1	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	30	-	1	-	-	-	1	-	-	-	1	3
2002.....	33	-	-	1	-	-	-	2	-	1	-	-
2001.....	25	-	1	-	-	-	2	-	1	-	-	3
2000.....	42	-	-	-	-	-	-	-	2	1	-	6
1999.....	31	-	-	-	-	-	-	1	1	1	-	-

Non-Hispanic 1/, male

2003.....	11	-	1	-	-	-	1	-	-	-	-	2
2002.....	6	-	-	1	-	-	-	1	-	-	-	-
2001.....	5	-	-	-	-	-	-	-	-	-	-	1
2000.....	18	-	-	-	-	-	-	-	-	-	-	3
1999.....	9	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of kidney (N25,N27)

Non-Hispanic 1/, female

2003.....	19	-	-	-	-	-	-	-	-	-	1	1
2002.....	27	-	-	-	-	-	-	1	-	1	-	-
2001.....	20	-	1	-	-	-	2	-	1	-	-	2
2000.....	24	-	-	-	-	-	-	-	2	1	-	3
1999.....	22	-	-	-	-	-	-	1	-	1	-	-

Non-Hispanic White, both sexes

2003.....	23	-	1	-	-	-	-	-	-	-	1	2
2002.....	27	-	-	-	-	-	-	2	-	1	-	-
2001.....	17	-	1	-	-	-	1	-	1	-	-	2
2000.....	31	-	-	-	-	-	-	-	1	-	-	3
1999.....	19	-	-	-	-	-	-	1	1	1	-	-

Non-Hispanic White, male

2003.....	7	-	1	-	-	-	-	-	-	-	-	1
2002.....	3	-	-	-	-	-	-	1	-	-	-	-
2001.....	3	-	-	-	-	-	-	-	-	-	-	1
2000.....	15	-	-	-	-	-	-	-	-	-	-	1
1999.....	7	-	-	-	-	-	-	-	1	-	-	-

Non-Hispanic White, female

2003.....	16	-	-	-	-	-	-	-	-	-	1	1
2002.....	24	-	-	-	-	-	-	1	-	1	-	-
2001.....	14	-	1	-	-	-	1	-	1	-	-	1
2000.....	16	-	-	-	-	-	-	-	1	-	-	2
1999.....	12	-	-	-	-	-	-	1	-	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Infections of kidney (N10-N12,N13.6,N15.1)												
All origins, both sexes												
2003.....	823	4	2	-	1	2	8	7	9	10	26	32
2002.....	788	2	-	-	3	2	3	5	8	11	17	19
2001.....	748	2	4	-	-	1	4	5	6	16	17	21
2000.....	813	5	3	-	-	1	1	4	5	9	12	21
1999.....	846	3	1	1	-	3	1	5	4	13	15	22
All origins, male												
2003.....	246	3	-	-	1	-	2	-	1	3	7	14
2002.....	251	2	-	-	1	-	1	3	3	3	9	5
2001.....	213	1	2	-	-	-	2	-	3	4	6	6
2000.....	214	3	-	-	-	1	-	2	1	2	6	6
1999.....	263	1	-	-	-	1	-	-	-	2	5	8
All origins, female												
2003.....	577	1	2	-	-	2	6	7	8	7	19	18
2002.....	537	-	-	-	2	2	2	2	5	8	8	14
2001.....	535	1	2	-	-	1	2	5	3	12	11	15
2000.....	599	2	3	-	-	-	1	2	4	7	6	15
1999.....	583	2	1	1	-	2	1	5	4	11	10	14
Hispanic, both sexes												
2003.....	46	-	-	-	-	-	1	-	-	-	2	1
2002.....	47	1	-	-	2	-	-	2	1	2	-	1
2001.....	33	-	-	-	-	-	-	1	-	1	2	2
2000.....	39	1	-	-	-	-	-	-	-	1	1	3
1999.....	55	-	-	-	-	-	-	1	-	4	3	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Infections of kidney (N10-N12,N13.6,N15.1)</b>												
<b>Hispanic, male</b>												
2003.....	11	-	-	-	-	-	-	-	-	-	-	1
2002.....	19	1	-	-	-	-	-	2	-	1	-	-
2001.....	10	-	-	-	-	-	-	-	-	-	-	2
2000.....	11	1	-	-	-	-	-	-	-	-	-	2
1999.....	21	-	-	-	-	-	-	-	-	1	1	2
<b>Hispanic, female</b>												
2003.....	35	-	-	-	-	-	1	-	-	-	2	-
2002.....	28	-	-	-	2	-	-	-	1	1	-	1
2001.....	23	-	-	-	-	-	-	1	-	1	2	-
2000.....	28	-	-	-	-	-	-	-	-	1	1	1
1999.....	34	-	-	-	-	-	-	1	-	3	2	3
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	774	4	2	-	1	2	7	7	9	10	23	30
2002.....	740	1	-	-	1	2	3	3	7	9	17	18
2001.....	714	2	4	-	-	1	4	4	6	15	15	19
2000.....	771	4	3	-	-	1	1	4	5	8	11	18
1999.....	786	3	1	1	-	3	1	4	4	8	12	17
<b>Non-Hispanic 1/, male</b>												
2003.....	233	3	-	-	1	-	2	-	1	3	6	12
2002.....	232	1	-	-	1	-	1	1	3	2	9	5
2001.....	203	1	2	-	-	-	2	-	3	4	6	4
2000.....	201	2	-	-	-	1	-	2	1	2	6	4
1999.....	240	1	-	-	-	1	-	-	-	-	4	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Infections of kidney (N10-N12,N13.6,N15.1)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	541	1	2	-	-	2	5	7	8	7	17	18
2002.....	508	-	-	-	-	2	2	2	4	7	8	13
2001.....	511	1	2	-	-	1	2	4	3	11	9	15
2000.....	570	2	3	-	-	-	1	2	4	6	5	14
1999.....	546	2	1	1	-	2	1	4	4	8	8	11
<b>Non-Hispanic White, both sexes</b>												
2003.....	656	1	1	-	1	2	5	3	8	6	19	21
2002.....	645	1	-	-	-	-	1	2	5	5	15	9
2001.....	614	1	2	-	-	-	2	3	4	10	13	13
2000.....	647	4	3	-	-	-	1	4	3	4	6	14
1999.....	689	1	1	1	-	1	1	3	4	3	9	14
<b>Non-Hispanic White, male</b>												
2003.....	202	-	-	-	1	-	2	-	1	2	6	10
2002.....	200	1	-	-	-	-	1	1	3	2	7	3
2001.....	166	-	1	-	-	-	1	-	2	3	6	3
2000.....	169	2	-	-	-	-	-	2	1	-	4	4
1999.....	209	1	-	-	-	1	-	-	-	-	2	4
<b>Non-Hispanic White, female</b>												
2003.....	454	1	1	-	-	2	3	3	7	4	13	11
2002.....	445	-	-	-	-	-	-	1	2	3	8	6
2001.....	448	1	1	-	-	-	1	3	2	7	7	10
2000.....	478	2	3	-	-	-	1	2	2	4	2	10
1999.....	480	-	1	1	-	-	1	3	4	3	7	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic Black, both sexes

2003.....	95	3	1	-	-	-	2	2	-	3	4	7
2002.....	73	-	-	-	-	1	2	-	2	4	2	6
2001.....	80	1	-	-	-	-	1	1	2	5	2	4
2000.....	101	-	-	-	-	1	-	-	2	3	4	3
1999.....	85	1	-	-	-	2	-	1	-	3	3	3

Non-Hispanic Black, male

2003.....	29	3	-	-	-	-	-	-	-	1	-	2
2002.....	22	-	-	-	-	-	-	-	-	-	2	1
2001.....	28	1	-	-	-	-	1	-	1	1	-	1
2000.....	27	-	-	-	-	1	-	-	-	1	1	-
1999.....	29	-	-	-	-	-	-	-	-	-	2	2

Non-Hispanic Black, female

2003.....	66	-	1	-	-	-	2	2	-	2	4	5
2002.....	51	-	-	-	-	1	2	-	2	4	-	5
2001.....	52	-	-	-	-	-	-	1	1	4	2	3
2000.....	74	-	-	-	-	-	-	-	2	2	3	3
1999.....	56	1	-	-	-	2	-	1	-	3	1	1

Origin not stated 2/, both sexes

2003.....	3	-	-	-	-	-	-	-	-	-	1	1
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	5	-	-	-	-	-	-	-	-	1	-	-

















Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

#### Inflammatory diseases of female pelvic organs (N70-N76)

##### Non-Hispanic 1/, female

2003.....	115	1	-	-	1	1	4	1	1	6	7	5
2002.....	110	1	-	-	-	1	1	-	3	12	3	4
2001.....	122	1	-	-	-	-	2	2	1	2	8	8
2000.....	111	2	-	-	1	-	2	2	3	8	6	4
1999.....	101	2	1	-	-	1	2	3	3	4	7	3

##### Non-Hispanic White, both sexes

2003.....	93	-	-	-	1	1	2	1	-	5	4	4
2002.....	88	1	-	-	-	-	-	-	2	4	3	2
2001.....	98	1	-	-	-	-	2	2	1	-	6	5
2000.....	87	1	-	-	-	-	2	2	1	4	2	3
1999.....	73	1	1	-	-	1	1	1	2	1	2	1

##### Non-Hispanic White, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

##### Non-Hispanic White, female

2003.....	93	-	-	-	1	1	2	1	-	5	4	4
2002.....	88	1	-	-	-	-	-	-	2	4	3	2
2001.....	98	1	-	-	-	-	2	2	1	-	6	5
2000.....	87	1	-	-	-	-	2	2	1	4	2	3
1999.....	73	1	1	-	-	1	1	1	2	1	2	1





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

All origins, both sexes

2003.....	545	...	...	...	1	27	89	110	119	106	60	19
2002.....	379	...	...	...	1	28	61	82	93	79	26	8
2001.....	416	...	...	...	3	37	73	93	97	73	33	5
2000.....	404	...	...	...	-	18	77	87	95	86	35	5
1999.....	406	...	...	...	-	33	64	89	92	88	33	5

All origins, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins, female

2003.....	545	...	...	...	1	27	89	110	119	106	60	19
2002.....	379	...	...	...	1	28	61	82	93	79	26	8
2001.....	416	...	...	...	3	37	73	93	97	73	33	5
2000.....	404	...	...	...	-	18	77	87	95	86	35	5
1999.....	406	...	...	...	-	33	64	89	92	88	33	5

Hispanic, both sexes

2003.....	104	...	...	...	-	8	16	31	15	17	11	4
2002.....	70	...	...	...	-	6	13	10	18	16	5	2
2001.....	83	...	...	...	1	9	17	21	17	10	7	-
2000.....	84	...	...	...	-	2	22	20	16	19	4	1
1999.....	70	...	...	...	-	6	13	18	8	21	4	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pregnancy, childbirth and the puerperium (000-099)												
Non-Hispanic 1/, female												
2003.....	437	...	...	...	1	19	70	79	103	89	49	15
2002.....	305	...	...	...	1	22	47	71	74	62	21	6
2001.....	332	...	...	...	2	28	56	72	79	63	26	5
2000.....	319	...	...	...	-	16	55	67	79	66	31	4
1999.....	333	...	...	...	-	27	49	70	84	67	29	5
Non-Hispanic White, both sexes												
2003.....	213	...	...	...	-	7	34	43	44	39	27	10
2002.....	133	...	...	...	-	6	15	31	38	25	14	3
2001.....	156	...	...	...	1	13	25	31	46	25	12	2
2000.....	161	...	...	...	-	9	24	33	39	37	17	1
1999.....	152	...	...	...	-	8	18	29	45	35	13	4
Non-Hispanic White, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic White, female												
2003.....	213	...	...	...	-	7	34	43	44	39	27	10
2002.....	133	...	...	...	-	6	15	31	38	25	14	3
2001.....	156	...	...	...	1	13	25	31	46	25	12	2
2000.....	161	...	...	...	-	9	24	33	39	37	17	1
1999.....	152	...	...	...	-	8	18	29	45	35	13	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

Non-Hispanic Black, both sexes

2003.....	190	...	...	...	1	12	30	32	50	40	18	5
2002.....	153	...	...	...	1	16	29	37	27	34	7	2
2001.....	154	...	...	...	1	14	31	35	30	28	12	3
2000.....	139	...	...	...	-	4	30	31	34	25	12	3
1999.....	158	...	...	...	-	18	28	37	33	27	13	-

Non-Hispanic Black, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic Black, female

2003.....	190	...	...	...	1	12	30	32	50	40	18	5
2002.....	153	...	...	...	1	16	29	37	27	34	7	2
2001.....	154	...	...	...	1	14	31	35	30	28	12	3
2000.....	139	...	...	...	-	4	30	31	34	25	12	3
1999.....	158	...	...	...	-	18	28	37	33	27	13	-

Origin not stated 2/, both sexes

2003.....	4	...	...	...	-	-	3	-	1	-	-	-
2002.....	4	...	...	...	-	-	1	1	1	1	-	-
2001.....	1	...	...	...	-	-	-	-	1	-	-	-
2000.....	1	...	...	...	-	-	-	-	-	1	-	-
1999.....	3	...	...	...	-	-	2	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

Origin not stated 2/, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Origin not stated 2/, female

2003.....	4	...	...	...	-	-	3	-	1	-	-	-
2002.....	4	...	...	...	-	-	1	1	1	1	-	-
2001.....	1	...	...	...	-	-	-	-	1	-	-	-
2000.....	1	...	...	...	-	-	-	-	-	1	-	-
1999.....	3	...	...	...	-	-	2	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

All origins, both sexes

2003.....	40	...	...	...	-	6	5	8	11	6	3	1
2002.....	22	...	...	...	1	-	4	4	5	6	1	1
2001.....	38	...	...	...	2	6	7	6	8	8	1	-
2000.....	37	...	...	...	-	-	6	12	5	10	4	-
1999.....	31	...	...	...	-	3	5	5	10	6	2	-

All origins, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All origins, female

2003.....	40	...	...	...	-	6	5	8	11	6	3	1
2002.....	22	...	...	...	1	-	4	4	5	6	1	1
2001.....	38	...	...	...	2	6	7	6	8	8	1	-
2000.....	37	...	...	...	-	-	6	12	5	10	4	-
1999.....	31	...	...	...	-	3	5	5	10	6	2	-

Hispanic, both sexes

2003.....	8	...	...	...	-	-	1	4	2	1	-	-
2002.....	1	...	...	...	-	-	-	1	-	-	-	-
2001.....	1	...	...	...	1	-	-	-	-	-	-	-
2000.....	6	...	...	...	-	-	2	3	-	1	-	-
1999.....	4	...	...	...	-	2	-	-	-	2	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Non-Hispanic 1/, female

2003.....	32	...	...	...	-	6	4	4	9	5	3	1
2002.....	20	...	...	...	1	-	4	3	5	5	1	1
2001.....	37	...	...	...	1	6	7	6	8	8	1	-
2000.....	30	...	...	...	-	-	4	9	5	8	4	-
1999.....	26	...	...	...	-	1	4	5	10	4	2	-

Non-Hispanic White, both sexes

2003.....	11	...	...	...	-	1	1	4	2	2	1	-
2002.....	6	...	...	...	-	-	1	1	3	-	1	-
2001.....	19	...	...	...	-	3	4	4	4	3	1	-
2000.....	12	...	...	...	-	-	-	4	1	4	3	-
1999.....	11	...	...	...	-	-	2	1	5	1	2	-

Non-Hispanic White, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic White, female

2003.....	11	...	...	...	-	1	1	4	2	2	1	-
2002.....	6	...	...	...	-	-	1	1	3	-	1	-
2001.....	19	...	...	...	-	3	4	4	4	3	1	-
2000.....	12	...	...	...	-	-	-	4	1	4	3	-
1999.....	11	...	...	...	-	-	2	1	5	1	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pregnancy with abortive outcome (000-007)												
Non-Hispanic Black, both sexes												
2003.....	18	...	...	...	-	5	3	-	5	2	2	1
2002.....	11	...	...	...	1	-	2	2	2	4	-	-
2001.....	16	...	...	...	1	3	3	2	3	4	-	-
2000.....	18	...	...	...	-	-	4	5	4	4	1	-
1999.....	14	...	...	...	-	1	2	4	4	3	-	-
Non-Hispanic Black, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic Black, female												
2003.....	18	...	...	...	-	5	3	-	5	2	2	1
2002.....	11	...	...	...	1	-	2	2	2	4	-	-
2001.....	16	...	...	...	1	3	3	2	3	4	-	-
2000.....	18	...	...	...	-	-	4	5	4	4	1	-
1999.....	14	...	...	...	-	1	2	4	4	3	-	-
Origin not stated 2/, both sexes												
2003.....	-	...	...	...	-	-	-	-	-	-	-	-
2002.....	1	...	...	...	-	-	-	-	-	1	-	-
2001.....	-	...	...	...	-	-	-	-	-	-	-	-
2000.....	1	...	...	...	-	-	-	-	-	1	-	-
1999.....	1	...	...	...	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

Origin not stated 2/, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Origin not stated 2/, female

2003.....	-	...	...	...	-	-	-	-	-	-	-	-
2002.....	1	...	...	...	-	-	-	-	-	1	-	-
2001.....	-	...	...	...	-	-	-	-	-	-	-	-
2000.....	1	...	...	...	-	-	-	-	-	1	-	-
1999.....	1	...	...	...	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
All origins, both sexes												
2003.....	505	...	...	...	1	21	84	102	108	100	57	18
2002.....	357	...	...	...	-	28	57	78	88	73	25	7
2001.....	378	...	...	...	1	31	66	87	89	65	32	5
2000.....	367	...	...	...	-	18	71	75	90	76	31	5
1999.....	375	...	...	...	-	30	59	84	82	82	31	5
All origins, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
All origins, female												
2003.....	505	...	...	...	1	21	84	102	108	100	57	18
2002.....	357	...	...	...	-	28	57	78	88	73	25	7
2001.....	378	...	...	...	1	31	66	87	89	65	32	5
2000.....	367	...	...	...	-	18	71	75	90	76	31	5
1999.....	375	...	...	...	-	30	59	84	82	82	31	5
Hispanic, both sexes												
2003.....	96	...	...	...	-	8	15	27	13	16	11	4
2002.....	69	...	...	...	-	6	13	9	18	16	5	2
2001.....	82	...	...	...	-	9	17	21	17	10	7	-
2000.....	78	...	...	...	-	2	20	17	16	18	4	1
1999.....	66	...	...	...	-	4	13	18	8	19	4	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Non-Hispanic 1/, female												
2003.....	405	...	...	...	1	13	66	75	94	84	46	14
2002.....	285	...	...	...	-	22	43	68	69	57	20	5
2001.....	295	...	...	...	1	22	49	66	71	55	25	5
2000.....	289	...	...	...	-	16	51	58	74	58	27	4
1999.....	307	...	...	...	-	26	45	65	74	63	27	5
Non-Hispanic White, both sexes												
2003.....	202	...	...	...	-	6	33	39	42	37	26	10
2002.....	127	...	...	...	-	6	14	30	35	25	13	3
2001.....	137	...	...	...	1	10	21	27	42	22	11	2
2000.....	149	...	...	...	-	9	24	29	38	33	14	1
1999.....	141	...	...	...	-	8	16	28	40	34	11	4
Non-Hispanic White, male												
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...
Non-Hispanic White, female												
2003.....	202	...	...	...	-	6	33	39	42	37	26	10
2002.....	127	...	...	...	-	6	14	30	35	25	13	3
2001.....	137	...	...	...	1	10	21	27	42	22	11	2
2000.....	149	...	...	...	-	9	24	29	38	33	14	1
1999.....	141	...	...	...	-	8	16	28	40	34	11	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Non-Hispanic Black, both sexes

2003.....	172	...	...	...	1	7	27	32	45	38	16	4
2002.....	142	...	...	...	-	16	27	35	25	30	7	2
2001.....	138	...	...	...	-	11	28	33	27	24	12	3
2000.....	121	...	...	...	-	4	26	26	30	21	11	3
1999.....	144	...	...	...	-	17	26	33	29	24	13	-

Non-Hispanic Black, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic Black, female

2003.....	172	...	...	...	1	7	27	32	45	38	16	4
2002.....	142	...	...	...	-	16	27	35	25	30	7	2
2001.....	138	...	...	...	-	11	28	33	27	24	12	3
2000.....	121	...	...	...	-	4	26	26	30	21	11	3
1999.....	144	...	...	...	-	17	26	33	29	24	13	-

Origin not stated 2/, both sexes

2003.....	4	...	...	...	-	-	3	-	1	-	-	-
2002.....	3	...	...	...	-	-	1	1	1	-	-	-
2001.....	1	...	...	...	-	-	-	-	1	-	-	-
2000.....	-	...	...	...	-	-	-	-	-	-	-	-
1999.....	2	...	...	...	-	-	1	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (O10-O99)

Origin not stated 2/, male

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Origin not stated 2/, female

2003.....	4	...	...	...	-	-	3	-	1	-	-	-
2002.....	3	...	...	...	-	-	1	1	1	-	-	-
2001.....	1	...	...	...	-	-	-	-	1	-	-	-
2000.....	-	...	...	...	-	-	-	-	-	-	-	-
1999.....	2	...	...	...	-	-	1	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

All origins, both sexes

2003.....	14,378	14,254	79	10	9	12	4	2	1	1	2	1
2002.....	14,254	14,106	65	15	16	14	7	6	4	7	3	2
2001.....	13,887	13,734	72	22	14	8	9	5	1	6	3	3
2000.....	14,069	13,918	79	25	16	10	5	5	2	2	2	1
1999.....	14,259	14,084	92	25	18	11	5	2	5	2	1	2

All origins, male

2003.....	8,197	8,129	43	5	6	6	2	1	1	1	-	-
2002.....	8,047	7,960	40	10	9	9	3	3	1	4	1	2
2001.....	7,868	7,775	44	16	8	3	4	2	-	4	2	3
2000.....	7,968	7,872	52	13	10	9	2	4	2	-	2	1
1999.....	8,044	7,941	56	14	9	6	4	-	2	2	-	2

All origins, female

2003.....	6,181	6,125	36	5	3	6	2	1	-	-	2	1
2002.....	6,207	6,146	25	5	7	5	4	3	3	3	2	-
2001.....	6,019	5,959	28	6	6	5	5	3	1	2	1	-
2000.....	6,101	6,046	27	12	6	1	3	1	-	2	-	-
1999.....	6,215	6,143	36	11	9	5	1	2	3	-	1	-

Hispanic, both sexes

2003.....	2,628	2,599	15	4	4	2	1	1	-	1	-	-
2002.....	2,402	2,375	14	4	2	2	1	1	-	1	1	-
2001.....	2,227	2,203	14	3	-	-	2	4	-	-	-	1
2000.....	2,145	2,120	12	7	3	1	1	-	-	1	-	-
1999.....	2,153	2,127	18	2	-	3	1	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

Hispanic, male

2003.....	1,459	1,442	8	2	2	1	1	1	-	1	-	-
2002.....	1,352	1,335	9	3	2	2	-	1	-	-	-	-
2001.....	1,268	1,258	6	2	-	-	-	1	-	-	-	1
2000.....	1,194	1,180	7	4	2	1	-	-	-	-	-	-
1999.....	1,192	1,181	5	2	-	2	1	-	-	-	-	-

Hispanic, female

2003.....	1,169	1,157	7	2	2	1	-	-	-	-	-	-
2002.....	1,050	1,040	5	1	-	-	1	-	-	1	1	-
2001.....	959	945	8	1	-	-	2	3	-	-	-	-
2000.....	951	940	5	3	1	-	1	-	-	1	-	-
1999.....	961	946	13	-	-	1	-	1	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	11,559	11,464	64	6	5	10	3	1	1	-	2	1
2002.....	11,673	11,553	50	11	14	12	6	5	4	6	2	2
2001.....	11,500	11,371	58	19	14	8	7	1	1	6	3	2
2000.....	11,672	11,546	67	18	13	9	4	5	2	1	2	1
1999.....	11,815	11,668	73	23	18	8	4	1	5	2	1	2

Non-Hispanic 1/, male

2003.....	6,622	6,571	35	3	4	5	1	-	1	-	-	-
2002.....	6,596	6,527	30	7	7	7	3	2	1	4	1	2
2001.....	6,511	6,428	38	14	8	3	4	1	-	4	2	2
2000.....	6,637	6,555	45	9	8	8	2	4	2	-	2	1
1999.....	6,680	6,589	50	12	9	4	3	-	2	2	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

Non-Hispanic 1/, female

2003.....	4,937	4,893	29	3	1	5	2	1	-	-	2	1
2002.....	5,077	5,026	20	4	7	5	3	3	3	2	1	-
2001.....	4,989	4,943	20	5	6	5	3	-	1	2	1	-
2000.....	5,035	4,991	22	9	5	1	2	1	-	1	-	-
1999.....	5,135	5,079	23	11	9	4	1	1	3	-	1	-

Non-Hispanic White, both sexes

2003.....	6,359	6,303	31	5	3	10	2	1	1	-	2	1
2002.....	6,384	6,313	22	5	10	11	3	4	2	4	2	2
2001.....	6,323	6,239	34	11	7	5	6	1	1	6	2	2
2000.....	6,318	6,242	34	11	9	7	4	2	1	1	2	1
1999.....	6,384	6,286	37	17	14	8	3	1	5	2	1	2

Non-Hispanic White, male

2003.....	3,673	3,643	17	3	3	5	1	-	1	-	-	-
2002.....	3,663	3,628	10	4	3	6	1	2	1	2	1	2
2001.....	3,574	3,518	22	9	5	2	4	1	-	4	1	2
2000.....	3,605	3,553	23	6	8	7	2	1	1	-	2	1
1999.....	3,627	3,570	24	8	7	4	3	-	2	2	-	2

Non-Hispanic White, female

2003.....	2,686	2,660	14	2	-	5	1	1	-	-	2	1
2002.....	2,721	2,685	12	1	7	5	2	2	1	2	1	-
2001.....	2,749	2,721	12	2	2	3	2	-	1	2	1	-
2000.....	2,713	2,689	11	5	1	-	2	1	-	1	-	-
1999.....	2,757	2,716	13	9	7	4	-	1	3	-	1	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All origins, both sexes												
2003.....	10,518	5,621	541	180	206	223	228	198	228	242	322	324
2002.....	10,687	5,623	530	199	218	248	244	213	262	264	308	346
2001.....	10,442	5,513	557	182	194	255	250	225	233	259	327	303
2000.....	10,578	5,743	495	198	201	225	216	208	269	277	292	287
1999.....	10,393	5,473	549	207	221	222	212	212	253	279	273	342
All origins, male												
2003.....	5,460	3,020	275	94	104	144	128	119	114	129	175	177
2002.....	5,515	2,942	282	99	116	146	136	132	152	132	164	195
2001.....	5,448	2,954	282	101	109	161	145	121	128	137	180	156
2000.....	5,517	3,028	259	105	102	137	131	120	159	154	162	157
1999.....	5,346	2,875	310	102	113	139	123	110	132	152	144	185
All origins, female												
2003.....	5,058	2,601	266	86	102	79	100	79	114	113	147	147
2002.....	5,172	2,681	248	100	102	102	108	81	110	132	144	151
2001.....	4,994	2,559	275	81	85	94	105	104	105	122	147	147
2000.....	5,061	2,715	236	93	99	88	85	88	110	123	130	130
1999.....	5,047	2,598	239	105	108	83	89	102	121	127	129	157
Hispanic, both sexes												
2003.....	1,787	1,324	122	31	34	27	30	35	23	16	28	18
2002.....	1,715	1,257	111	40	30	37	30	23	30	23	17	21
2001.....	1,632	1,195	103	28	30	32	41	26	19	19	23	24
2000.....	1,612	1,182	111	33	19	24	30	19	27	24	19	20
1999.....	1,498	1,072	101	31	26	27	29	28	26	15	19	26

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Hispanic, male												
2003.....	937	693	62	15	13	17	13	22	15	10	15	11
2002.....	903	653	60	18	17	22	17	14	22	11	8	12
2001.....	892	639	54	17	19	17	22	17	9	11	16	19
2000.....	857	605	60	16	12	19	20	11	15	16	11	11
1999.....	756	534	63	16	13	17	13	15	10	8	7	13
Hispanic, female												
2003.....	850	631	60	16	21	10	17	13	8	6	13	7
2002.....	812	604	51	22	13	15	13	9	8	12	9	9
2001.....	740	556	49	11	11	15	19	9	10	8	7	5
2000.....	755	577	51	17	7	5	10	8	12	8	8	9
1999.....	742	538	38	15	13	10	16	13	16	7	12	13
Non-Hispanic 1/, both sexes												
2003.....	8,687	4,269	418	147	170	195	198	162	205	226	290	305
2002.....	8,904	4,316	415	159	188	210	212	190	232	241	289	324
2001.....	8,748	4,271	452	153	164	223	209	198	214	238	304	277
2000.....	8,900	4,514	384	164	181	200	186	187	240	252	273	266
1999.....	8,825	4,349	447	175	195	195	183	183	227	263	253	313
Non-Hispanic 1/, male												
2003.....	4,500	2,312	213	77	90	126	115	97	99	119	157	166
2002.....	4,577	2,262	221	81	99	124	117	118	130	121	155	182
2001.....	4,527	2,297	227	83	90	144	123	103	119	125	164	135
2000.....	4,623	2,400	199	88	89	118	111	107	143	137	151	145
1999.....	4,550	2,311	246	85	100	122	110	95	122	144	136	169

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic 1/, female												
2003.....	4,187	1,957	205	70	80	69	83	65	106	107	133	139
2002.....	4,327	2,054	194	78	89	86	95	72	102	120	134	142
2001.....	4,221	1,974	225	70	74	79	86	95	95	113	140	142
2000.....	4,277	2,114	185	76	92	82	75	80	97	115	122	121
1999.....	4,275	2,038	201	90	95	73	73	88	105	119	117	144
Non-Hispanic White, both sexes												
2003.....	6,594	2,977	272	98	124	143	142	123	162	183	247	245
2002.....	6,830	3,068	281	117	136	153	162	146	185	191	232	270
2001.....	6,709	3,053	287	111	115	157	145	152	178	187	252	223
2000.....	6,792	3,199	240	113	127	134	145	144	200	201	218	224
1999.....	6,782	3,127	284	113	131	144	133	151	183	216	210	265
Non-Hispanic White, male												
2003.....	3,353	1,596	136	50	67	92	83	73	79	98	135	127
2002.....	3,510	1,643	151	58	67	90	88	90	105	96	127	150
2001.....	3,487	1,664	144	64	71	100	83	87	99	102	134	110
2000.....	3,519	1,720	124	55	64	74	89	82	121	112	121	120
1999.....	3,498	1,693	154	53	66	93	82	79	99	120	110	142
Non-Hispanic White, female												
2003.....	3,241	1,381	136	48	57	51	59	50	83	85	112	118
2002.....	3,320	1,425	130	59	69	63	74	56	80	95	105	120
2001.....	3,222	1,389	143	47	44	57	62	65	79	85	118	113
2000.....	3,273	1,479	116	58	63	60	56	62	79	89	97	104
1999.....	3,284	1,434	130	60	65	51	51	72	84	96	100	123

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Non-Hispanic Black, both sexes												
2003.....	1,671	1,015	118	36	37	40	52	33	42	34	35	51
2002.....	1,716	1,015	113	38	43	52	46	35	40	40	52	46
2001.....	1,674	999	127	32	41	61	52	35	32	43	46	44
2000.....	1,739	1,062	123	44	45	56	35	36	38	46	45	39
1999.....	1,680	994	137	49	56	42	40	28	32	42	34	41
Non-Hispanic Black, male												
2003.....	905	558	62	20	18	27	28	19	19	15	17	32
2002.....	887	506	58	22	25	33	26	22	21	18	27	26
2001.....	861	525	65	15	18	41	31	11	19	18	28	18
2000.....	911	545	67	29	21	39	19	22	21	23	26	22
1999.....	880	518	79	24	29	26	24	14	18	19	22	25
Non-Hispanic Black, female												
2003.....	766	457	56	16	19	13	24	14	23	19	18	19
2002.....	829	509	55	16	18	19	20	13	19	22	25	20
2001.....	813	474	62	17	23	20	21	24	13	25	18	26
2000.....	828	517	56	15	24	17	16	14	17	23	19	17
1999.....	800	476	58	25	27	16	16	14	14	23	12	16
Origin not stated 2/, both sexes												
2003.....	44	28	1	2	2	1	-	1	-	-	4	1
2002.....	68	50	4	-	-	1	2	-	-	-	2	1
2001.....	62	47	2	1	-	-	-	1	-	2	-	2
2000.....	66	47	-	1	1	1	-	2	2	1	-	1
1999.....	70	52	1	1	-	-	-	1	-	1	1	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Origin not stated 2/, male

2003.....	23	15	-	2	1	1	-	-	-	-	3	-
2002.....	35	27	1	-	-	-	2	-	-	-	1	1
2001.....	29	18	1	1	-	-	-	1	-	1	-	2
2000.....	37	23	-	1	1	-	-	2	1	1	-	1
1999.....	40	30	1	1	-	-	-	-	-	-	1	3

Origin not stated 2/, female

2003.....	21	13	1	-	1	-	-	1	-	-	1	1
2002.....	33	23	3	-	-	1	-	-	-	-	1	-
2001.....	33	29	1	-	-	-	-	-	-	1	-	-
2000.....	29	24	-	-	-	1	-	-	1	-	-	-
1999.....	30	22	-	-	-	-	-	1	-	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins, both sexes

2003.....	31,444	3,318	256	75	103	300	537	617	710	1,037	1,345	1,578
2002.....	29,975	3,456	248	61	78	256	484	562	757	1,013	1,389	1,368
2001.....	32,326	3,491	272	79	102	361	588	663	867	1,379	1,747	1,882
2000.....	31,876	3,639	248	85	92	317	551	663	891	1,284	1,703	1,727
1999.....	26,672	3,538	236	65	72	228	381	446	649	940	1,122	1,168

All origins, male

2003.....	14,583	1,974	143	30	62	197	366	434	473	685	841	1,011
2002.....	14,269	2,032	145	37	45	172	329	391	514	670	870	895
2001.....	15,779	2,041	162	44	55	249	405	461	579	888	1,119	1,220
2000.....	15,863	2,156	137	48	51	196	385	456	604	857	1,137	1,157
1999.....	12,877	2,072	136	35	42	153	269	287	434	599	740	796

All origins, female

2003.....	16,861	1,344	113	45	41	103	171	183	237	352	504	567
2002.....	15,706	1,424	103	24	33	84	155	171	243	343	519	473
2001.....	16,547	1,450	110	35	47	112	183	202	288	491	628	662
2000.....	16,013	1,483	111	37	41	121	166	207	287	427	566	570
1999.....	13,795	1,466	100	30	30	75	112	159	215	341	382	372

Hispanic, both sexes

2003.....	1,623	492	30	11	9	44	54	90	80	94	87	75
2002.....	1,548	472	40	4	7	34	57	68	81	81	96	65
2001.....	2,190	496	47	16	20	73	111	132	154	175	152	173
2000.....	2,071	506	51	15	16	50	90	118	142	177	165	144
1999.....	1,281	415	30	6	10	30	53	58	80	80	68	64

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Hispanic, male												
2003.....	998	309	16	6	6	27	38	73	62	72	63	63
2002.....	926	283	26	3	4	20	42	51	65	60	60	40
2001.....	1,410	280	28	8	7	56	83	105	116	130	109	135
2000.....	1,369	310	31	7	9	29	73	93	108	137	125	110
1999.....	768	237	14	5	5	21	43	42	57	59	51	52
Hispanic, female												
2003.....	625	183	14	5	3	17	16	17	18	22	24	12
2002.....	622	189	14	1	3	14	15	17	16	21	36	25
2001.....	780	216	19	8	13	17	28	27	38	45	43	38
2000.....	702	196	20	8	7	21	17	25	34	40	40	34
1999.....	513	178	16	1	5	9	10	16	23	21	17	12
Non-Hispanic 1/, both sexes												
2003.....	29,617	2,805	226	64	93	253	478	524	627	933	1,248	1,492
2002.....	28,211	2,968	207	56	70	220	423	492	670	925	1,281	1,292
2001.....	29,931	2,978	225	62	82	287	470	520	707	1,201	1,581	1,694
2000.....	29,514	3,108	196	69	75	265	455	540	739	1,091	1,513	1,567
1999.....	25,181	3,097	204	59	62	198	323	385	566	848	1,037	1,087
Non-Hispanic 1/, male												
2003.....	13,448	1,650	127	24	56	169	326	359	408	605	770	939
2002.....	13,206	1,739	119	34	40	150	284	338	445	605	803	849
2001.....	14,246	1,750	134	36	48	193	317	350	457	757	997	1,076
2000.....	14,288	1,834	106	41	41	166	308	360	488	708	992	1,037
1999.....	11,962	1,817	121	30	37	132	223	244	375	532	677	729

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Non-Hispanic 1/, female												
2003.....	16,169	1,155	99	40	37	84	152	165	219	328	478	553
2002.....	15,005	1,229	88	22	30	70	139	154	225	320	478	443
2001.....	15,685	1,228	91	26	34	94	153	170	250	444	584	618
2000.....	15,226	1,274	90	28	34	99	147	180	251	383	521	530
1999.....	13,219	1,280	83	29	25	66	100	141	191	316	360	358
Non-Hispanic White, both sexes												
2003.....	24,452	1,665	136	47	52	200	362	374	469	708	953	1,109
2002.....	23,099	1,777	124	30	41	157	311	354	495	689	975	955
2001.....	24,596	1,752	139	37	52	217	346	380	527	914	1,261	1,342
2000.....	24,064	1,805	134	41	47	191	327	374	544	849	1,178	1,211
1999.....	20,200	1,812	129	43	38	133	215	253	407	613	728	773
Non-Hispanic White, male												
2003.....	10,752	1,025	75	21	30	136	253	256	314	460	590	706
2002.....	10,432	1,056	66	19	25	108	218	252	329	454	617	629
2001.....	11,404	1,046	79	18	34	143	233	271	343	573	814	857
2000.....	11,351	1,067	79	28	24	128	228	255	370	568	798	815
1999.....	9,249	1,083	82	21	24	89	155	168	263	400	484	509
Non-Hispanic White, female												
2003.....	13,700	640	61	26	22	64	109	118	155	248	363	403
2002.....	12,667	721	58	11	16	49	93	102	166	235	358	326
2001.....	13,192	706	60	19	18	74	113	109	184	341	447	485
2000.....	12,713	738	55	13	23	63	99	119	174	281	380	396
1999.....	10,951	729	47	22	14	44	60	85	144	213	244	264

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic Black, both sexes

2003.....	4,628	1,022	80	13	36	46	104	130	137	198	278	349
2002.....	4,556	1,070	66	24	26	57	95	119	156	216	274	302
2001.....	4,606	1,081	69	20	22	52	106	121	145	239	279	315
2000.....	4,892	1,186	54	21	25	65	114	140	168	208	306	318
1999.....	4,508	1,149	67	15	18	54	98	114	146	225	279	294

Non-Hispanic Black, male

2003.....	2,404	554	44	2	23	28	64	92	81	124	166	211
2002.....	2,482	615	44	14	14	39	53	71	103	138	167	198
2001.....	2,414	614	44	15	11	37	72	67	83	144	156	192
2000.....	2,614	694	26	11	17	32	73	89	102	113	170	194
1999.....	2,460	658	35	9	11	38	62	67	105	127	172	205

Non-Hispanic Black, female

2003.....	2,224	468	36	11	13	18	40	38	56	74	112	138
2002.....	2,074	455	22	10	12	18	42	48	53	78	107	104
2001.....	2,192	467	25	5	11	15	34	54	62	95	123	123
2000.....	2,278	492	28	10	8	33	41	51	66	95	136	124
1999.....	2,048	491	32	6	7	16	36	47	41	98	107	89

Origin not stated 2/, both sexes

2003.....	204	21	-	-	1	3	5	3	3	10	10	11
2002.....	216	16	1	1	1	2	4	2	6	7	12	11
2001.....	205	17	-	1	-	1	7	11	6	3	14	15
2000.....	291	25	1	1	1	2	6	5	10	16	25	16
1999.....	210	26	2	-	-	-	5	3	3	12	17	17

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Origin not stated 2/, male

2003.....	137	15	-	-	-	1	2	2	3	8	8	9
2002.....	137	10	-	-	1	2	3	2	4	5	7	6
2001.....	123	11	-	-	-	-	5	6	6	1	13	9
2000.....	206	12	-	-	1	1	4	3	8	12	20	10
1999.....	147	18	1	-	-	-	3	1	2	8	12	15

Origin not stated 2/, female

2003.....	67	6	-	-	1	2	3	1	-	2	2	2
2002.....	79	6	1	1	-	-	1	-	2	2	5	5
2001.....	82	6	-	1	-	1	2	5	-	2	1	6
2000.....	85	13	1	1	-	1	2	2	2	4	5	6
1999.....	63	8	1	-	-	-	2	2	1	4	5	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
All origins, both sexes												
2003.....	201,676	1,190	635	408	519	830	1,193	1,413	1,917	3,099	5,308	7,248
2002.....	194,591	1,144	661	386	482	717	1,136	1,256	1,880	3,099	5,407	7,159
2001.....	186,406	1,131	646	389	471	772	1,027	1,225	1,847	3,145	4,939	6,667
2000.....	178,378	1,105	609	376	503	732	1,032	1,257	1,840	3,161	4,952	6,227
1999.....	171,683	1,148	647	436	485	785	986	1,235	1,827	3,203	4,786	5,877
All origins, male												
2003.....	81,907	711	342	234	307	489	700	802	1,083	1,765	3,167	4,320
2002.....	79,007	690	363	220	266	423	649	714	1,123	1,800	3,219	4,305
2001.....	75,162	649	358	187	267	463	601	699	1,030	1,869	2,905	3,970
2000.....	72,227	629	338	198	276	432	625	737	1,074	1,873	2,928	3,811
1999.....	69,784	680	354	229	254	483	562	717	1,073	1,912	2,923	3,510
All origins, female												
2003.....	119,769	479	293	174	212	341	493	611	834	1,334	2,141	2,928
2002.....	115,584	454	298	166	216	294	487	542	757	1,299	2,188	2,854
2001.....	111,244	482	288	202	204	309	426	526	817	1,276	2,034	2,697
2000.....	106,151	476	271	178	227	300	407	520	766	1,288	2,024	2,416
1999.....	101,899	468	293	207	231	302	424	518	754	1,291	1,863	2,367
Hispanic, both sexes												
2003.....	9,218	235	140	69	90	136	151	196	256	353	476	548
2002.....	8,377	209	120	74	77	100	150	173	236	307	450	530
2001.....	7,847	187	112	63	59	102	144	162	192	289	382	438
2000.....	7,325	196	109	57	56	82	132	168	213	311	395	440
1999.....	7,291	200	106	50	57	95	123	166	243	306	384	437

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Hispanic, male

2003.....	4,571	144	75	37	50	75	90	127	168	212	307	357
2002.....	4,221	116	66	47	42	54	94	109	152	201	316	355
2001.....	3,849	106	66	31	34	65	82	97	125	197	254	295
2000.....	3,663	100	54	29	34	54	86	100	146	211	272	295
1999.....	3,655	124	50	28	26	59	69	103	172	218	261	296

Hispanic, female

2003.....	4,647	91	65	32	40	61	61	69	88	141	169	191
2002.....	4,156	93	54	27	35	46	56	64	84	106	134	175
2001.....	3,998	81	46	32	25	37	62	65	67	92	128	143
2000.....	3,662	96	55	28	22	28	46	68	67	100	123	145
1999.....	3,636	76	56	22	31	36	54	63	71	88	123	141

Non-Hispanic 1/, both sexes

2003.....	191,961	946	493	337	428	688	1,039	1,210	1,655	2,731	4,813	6,669
2002.....	185,608	927	538	311	404	615	985	1,076	1,638	2,774	4,937	6,591
2001.....	177,985	934	532	325	411	667	878	1,060	1,649	2,842	4,528	6,192
2000.....	170,462	898	497	317	445	646	895	1,080	1,617	2,836	4,530	5,738
1999.....	163,832	935	537	386	428	688	859	1,068	1,577	2,878	4,369	5,397

Non-Hispanic 1/, male

2003.....	77,081	561	266	196	256	409	607	668	912	1,543	2,850	3,946
2002.....	74,500	572	294	172	224	368	555	601	967	1,589	2,886	3,923
2001.....	71,014	539	291	156	232	396	516	599	900	1,664	2,629	3,645
2000.....	68,263	522	282	169	242	374	537	630	926	1,650	2,637	3,481
1999.....	65,835	546	302	201	228	423	490	613	896	1,679	2,637	3,186

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 1/, female

2003.....	114,880	385	227	141	172	279	432	542	743	1,188	1,963	2,723
2002.....	111,108	355	244	139	180	247	430	475	671	1,185	2,051	2,668
2001.....	106,971	395	241	169	179	271	362	461	749	1,178	1,899	2,547
2000.....	102,199	376	215	148	203	272	358	450	691	1,186	1,893	2,257
1999.....	97,997	389	235	185	200	265	369	455	681	1,199	1,732	2,211

Non-Hispanic White, both sexes

2003.....	165,261	565	324	241	290	497	748	809	1,101	1,873	3,531	4,969
2002.....	159,945	578	352	221	284	451	651	720	1,111	1,938	3,607	4,895
2001.....	153,156	556	362	246	295	474	615	699	1,109	1,983	3,218	4,511
2000.....	146,518	544	316	219	318	476	637	707	1,100	1,955	3,247	4,107
1999.....	141,002	554	375	265	304	513	587	749	1,096	1,949	3,087	3,858

Non-Hispanic White, male

2003.....	65,306	357	168	150	169	314	474	480	639	1,104	2,176	3,041
2002.....	63,109	376	191	124	155	284	393	420	693	1,179	2,207	2,985
2001.....	60,022	311	209	121	165	294	383	411	629	1,208	1,922	2,727
2000.....	57,565	310	183	123	184	286	397	446	659	1,183	1,945	2,600
1999.....	55,517	321	214	136	155	332	349	456	654	1,194	1,950	2,344

Non-Hispanic White, female

2003.....	99,955	208	156	91	121	183	274	329	462	769	1,355	1,928
2002.....	96,836	202	161	97	129	167	258	300	418	759	1,400	1,910
2001.....	93,134	245	153	125	130	180	232	288	480	775	1,296	1,784
2000.....	88,953	234	133	96	134	190	240	261	441	772	1,302	1,507
1999.....	85,485	233	161	129	149	181	238	293	442	755	1,137	1,514

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic Black, both sexes

2003.....	22,997	333	133	69	113	155	246	349	464	763	1,117	1,521
2002.....	22,297	298	146	72	100	133	283	317	458	735	1,200	1,528
2001.....	21,752	329	131	65	100	163	223	301	457	770	1,174	1,529
2000.....	21,081	296	144	79	110	146	227	343	467	789	1,159	1,495
1999.....	20,137	332	129	100	101	144	241	282	422	839	1,158	1,399

Non-Hispanic Black, male

2003.....	10,019	183	78	29	70	75	107	163	223	382	570	800
2002.....	9,722	167	80	37	58	66	139	155	233	354	594	834
2001.....	9,561	192	62	26	57	86	110	154	227	403	627	838
2000.....	9,351	180	77	35	52	76	121	165	235	412	621	804
1999.....	8,967	192	67	56	62	75	127	135	209	434	606	765

Non-Hispanic Black, female

2003.....	12,978	150	55	40	43	80	139	186	241	381	547	721
2002.....	12,575	131	66	35	42	67	144	162	225	381	606	694
2001.....	12,191	137	69	39	43	77	113	147	230	367	547	691
2000.....	11,730	116	67	44	58	70	106	178	232	377	538	691
1999.....	11,170	140	62	44	39	69	114	147	213	405	552	634

Origin not stated 2/, both sexes

2003.....	497	9	2	2	1	6	3	7	6	15	19	31
2002.....	606	8	3	1	1	2	1	7	6	18	20	38
2001.....	574	10	2	1	1	3	5	3	6	14	29	37
2000.....	591	11	3	2	2	4	5	9	10	14	27	49
1999.....	560	13	4	-	-	2	4	1	7	19	33	43

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Origin not stated 2/, male

2003.....	255	6	1	1	1	5	3	7	3	10	10	17
2002.....	286	2	3	1	-	1	-	4	4	10	17	27
2001.....	299	4	1	-	1	2	3	3	5	8	22	30
2000.....	301	7	2	-	-	4	2	7	2	12	19	35
1999.....	294	10	2	-	-	1	3	1	5	15	25	28

Origin not stated 2/, female

2003.....	242	3	1	1	-	1	-	-	3	5	9	14
2002.....	320	6	-	-	1	1	1	3	2	8	3	11
2001.....	275	6	1	1	-	1	2	-	1	6	7	7
2000.....	290	4	1	2	2	-	3	2	8	2	8	14
1999.....	266	3	2	-	-	1	1	-	2	4	8	15

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All origins, both sexes												
2003.....	109,277	945	1,717	1,096	1,522	6,755	8,517	6,284	6,257	7,495	9,271	8,865
2002.....	106,742	946	1,641	1,176	1,542	7,137	8,275	6,122	6,447	7,642	9,068	8,241
2001.....	101,537	976	1,714	1,283	1,553	6,646	7,765	5,935	5,904	7,589	8,356	7,469
2000.....	97,900	881	1,826	1,391	1,588	6,755	7,358	5,898	5,871	7,471	7,942	6,870
1999.....	97,860	845	1,898	1,459	1,632	6,688	6,968	5,881	6,009	7,366	7,865	6,696
All origins, male												
2003.....	70,532	539	1,048	650	1,012	4,706	6,585	4,826	4,708	5,390	6,613	6,325
2002.....	69,257	553	1,024	720	993	4,981	6,457	4,768	4,867	5,535	6,477	5,940
2001.....	66,060	541	1,045	786	1,017	4,722	6,110	4,605	4,487	5,502	6,110	5,519
2000.....	63,817	501	1,098	824	1,037	4,757	5,703	4,606	4,453	5,537	5,907	5,103
1999.....	63,535	494	1,127	897	1,077	4,624	5,416	4,555	4,542	5,411	5,837	4,996
All origins, female												
2003.....	38,745	406	669	446	510	2,049	1,932	1,458	1,549	2,105	2,658	2,540
2002.....	37,485	393	617	456	549	2,156	1,818	1,354	1,580	2,107	2,591	2,301
2001.....	35,477	435	669	497	536	1,924	1,655	1,330	1,417	2,087	2,246	1,950
2000.....	34,083	380	728	567	551	1,998	1,655	1,292	1,418	1,934	2,035	1,767
1999.....	34,325	351	771	562	555	2,064	1,552	1,326	1,467	1,955	2,028	1,700
Hispanic, both sexes												
2003.....	10,418	131	325	218	213	1,005	1,361	1,208	959	948	929	735
2002.....	10,106	127	341	199	190	984	1,429	1,082	945	898	943	685
2001.....	9,523	154	310	186	193	879	1,303	1,073	865	942	818	627
2000.....	8,830	118	306	205	178	884	1,211	944	832	856	762	542
1999.....	8,650	134	341	170	192	800	1,146	936	854	844	763	546

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidents (unintentional injuries) (V01-X59,Y85-Y86)</b>												
<b>Hispanic, male</b>												
2003.....	7,899	77	200	138	151	756	1,149	1,022	805	763	725	585
2002.....	7,698	83	215	120	124	763	1,195	917	792	712	756	522
2001.....	7,157	75	195	114	110	669	1,104	908	711	763	655	481
2000.....	6,696	67	174	105	105	689	1,028	804	695	682	635	435
1999.....	6,578	80	197	107	120	608	967	787	707	692	615	443
<b>Hispanic, female</b>												
2003.....	2,519	54	125	80	62	249	212	186	154	185	204	150
2002.....	2,408	44	126	79	66	221	234	165	153	186	187	163
2001.....	2,366	79	115	72	83	210	199	165	154	179	163	146
2000.....	2,134	51	132	100	73	195	183	140	137	174	127	107
1999.....	2,072	54	144	63	72	192	179	149	147	152	148	103
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	98,406	812	1,386	875	1,308	5,732	7,134	5,051	5,275	6,512	8,301	8,082
2002.....	96,175	816	1,296	972	1,346	6,136	6,820	5,019	5,470	6,705	8,087	7,509
2001.....	91,522	817	1,399	1,091	1,354	5,748	6,439	4,846	5,011	6,602	7,487	6,793
2000.....	88,558	759	1,512	1,180	1,405	5,848	6,120	4,927	5,008	6,580	7,126	6,274
1999.....	88,669	710	1,550	1,286	1,435	5,865	5,793	4,913	5,121	6,481	7,048	6,099
<b>Non-Hispanic 1/, male</b>												
2003.....	62,290	460	844	509	860	3,936	5,419	3,785	3,885	4,600	5,854	5,707
2002.....	61,201	468	806	596	865	4,208	5,240	3,835	4,047	4,790	5,692	5,380
2001.....	58,528	462	845	668	902	4,038	4,986	3,684	3,754	4,701	5,416	4,999
2000.....	56,736	431	921	717	931	4,053	4,654	3,781	3,731	4,826	5,228	4,626
1999.....	56,551	414	926	787	952	4,003	4,425	3,745	3,805	4,687	5,184	4,510

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidents (unintentional injuries) (V01-X59,Y85-Y86)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	36,116	352	542	366	448	1,796	1,715	1,266	1,390	1,912	2,447	2,375
2002.....	34,974	348	490	376	481	1,928	1,580	1,184	1,423	1,915	2,395	2,129
2001.....	32,994	355	554	423	452	1,710	1,453	1,162	1,257	1,901	2,071	1,794
2000.....	31,822	328	591	463	474	1,795	1,466	1,146	1,277	1,754	1,898	1,648
1999.....	32,118	296	624	499	483	1,862	1,368	1,168	1,316	1,794	1,864	1,589
<b>Non-Hispanic White, both sexes</b>												
2003.....	82,815	496	985	592	943	4,774	5,864	4,020	4,218	5,259	6,847	6,593
2002.....	80,605	462	865	616	965	5,099	5,565	3,921	4,423	5,358	6,677	6,080
2001.....	76,262	500	1,006	722	966	4,692	5,175	3,819	3,979	5,314	6,026	5,361
2000.....	73,536	455	1,033	796	1,047	4,852	4,883	3,788	3,942	5,302	5,706	4,978
1999.....	73,326	435	1,042	843	1,065	4,840	4,570	3,854	4,001	5,186	5,658	4,789
<b>Non-Hispanic White, male</b>												
2003.....	51,888	283	610	338	616	3,251	4,447	3,026	3,112	3,703	4,871	4,643
2002.....	50,652	279	527	378	619	3,466	4,280	2,992	3,281	3,822	4,728	4,412
2001.....	48,180	281	624	424	642	3,266	4,031	2,908	2,998	3,794	4,394	3,941
2000.....	46,470	266	638	487	689	3,326	3,739	2,899	2,962	3,914	4,199	3,680
1999.....	46,113	258	617	505	700	3,250	3,499	2,953	2,979	3,770	4,179	3,531
<b>Non-Hispanic White, female</b>												
2003.....	30,927	213	375	254	327	1,523	1,417	994	1,106	1,556	1,976	1,950
2002.....	29,953	183	338	238	346	1,633	1,285	929	1,142	1,536	1,949	1,668
2001.....	28,082	219	382	298	324	1,426	1,144	911	981	1,520	1,632	1,420
2000.....	27,066	189	395	309	358	1,526	1,144	889	980	1,388	1,507	1,298
1999.....	27,213	177	425	338	365	1,590	1,071	901	1,022	1,416	1,479	1,258

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidents (unintentional injuries) (V01-X59,Y85-Y86)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	12,154	283	333	233	284	656	934	777	825	997	1,198	1,247
2002.....	12,285	324	350	298	309	740	940	853	823	1,084	1,143	1,211
2001.....	12,239	272	331	313	317	760	953	784	816	1,026	1,256	1,212
2000.....	12,087	268	396	330	289	721	941	877	863	1,042	1,211	1,096
1999.....	12,533	251	428	392	309	764	930	841	897	1,063	1,182	1,138
<b>Non-Hispanic Black, male</b>												
2003.....	8,237	158	197	141	189	482	723	583	614	725	814	896
2002.....	8,428	172	231	180	211	541	719	665	614	786	776	836
2001.....	8,381	158	188	212	220	564	721	592	600	718	890	918
2000.....	8,380	145	232	204	202	530	703	681	635	759	890	808
1999.....	8,629	140	260	254	213	568	713	647	666	754	874	864
<b>Non-Hispanic Black, female</b>												
2003.....	3,917	125	136	92	95	174	211	194	211	272	384	351
2002.....	3,857	152	119	118	98	199	221	188	209	298	367	375
2001.....	3,858	114	143	101	97	196	232	192	216	308	366	294
2000.....	3,707	123	164	126	87	191	238	196	228	283	321	288
1999.....	3,904	111	168	138	96	196	217	194	231	309	308	274
<b>Origin not stated 2/, both sexes</b>												
2003.....	453	2	6	3	1	18	22	25	23	35	41	48
2002.....	461	3	4	5	6	17	26	21	32	39	38	47
2001.....	492	5	5	6	6	19	23	16	28	45	51	49
2000.....	512	4	8	6	5	23	27	27	31	35	54	54
1999.....	541	1	7	3	5	23	29	32	34	41	54	51

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Origin not stated 2/, male

2003.....	343	2	4	3	1	14	17	19	18	27	34	33
2002.....	358	2	3	4	4	10	22	16	28	33	29	38
2001.....	375	4	5	4	5	15	20	13	22	38	39	39
2000.....	385	3	3	2	1	15	21	21	27	29	44	42
1999.....	406	-	4	3	5	13	24	23	30	32	38	43

Origin not stated 2/, female

2003.....	110	-	2	-	-	4	5	6	5	8	7	15
2002.....	103	1	1	1	2	7	4	5	4	6	9	9
2001.....	117	1	-	2	1	4	3	3	6	7	12	10
2000.....	127	1	5	4	4	8	6	6	4	6	10	12
1999.....	135	1	3	-	-	10	5	9	4	9	16	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
All origins, both sexes												
2003.....	48,071	148	649	694	1,071	5,471	5,961	3,855	3,425	3,579	4,015	3,656
2002.....	48,366	127	640	705	1,033	5,782	6,034	3,891	3,593	3,750	3,938	3,593
2001.....	47,288	151	678	769	1,054	5,393	5,746	3,953	3,501	3,881	3,946	3,379
2000.....	46,749	174	691	842	1,080	5,475	5,553	3,957	3,443	3,889	3,772	3,248
1999.....	46,423	189	712	897	1,078	5,452	5,237	3,922	3,512	3,869	3,684	3,121
All origins, male												
2003.....	33,291	81	366	401	683	3,653	4,518	2,915	2,596	2,585	2,912	2,667
2002.....	33,533	72	381	405	634	3,875	4,596	2,990	2,686	2,703	2,796	2,638
2001.....	32,794	68	392	447	668	3,694	4,417	3,015	2,646	2,744	2,869	2,452
2000.....	32,166	94	391	464	674	3,693	4,197	3,016	2,554	2,802	2,758	2,338
1999.....	31,642	101	400	532	674	3,593	3,965	2,952	2,559	2,741	2,635	2,234
All origins, female												
2003.....	14,780	67	283	293	388	1,818	1,443	940	829	994	1,103	989
2002.....	14,833	55	259	300	399	1,907	1,438	901	907	1,047	1,142	955
2001.....	14,494	83	286	322	386	1,699	1,329	938	855	1,137	1,077	927
2000.....	14,583	80	300	378	406	1,782	1,356	941	889	1,087	1,014	910
1999.....	14,781	88	312	365	404	1,859	1,272	970	953	1,128	1,049	887
Hispanic, both sexes												
2003.....	6,045	39	139	147	161	817	1,017	821	591	505	468	330
2002.....	5,921	26	182	138	139	785	1,090	760	580	480	450	317
2001.....	5,774	43	159	120	145	718	1,022	768	525	514	428	313
2000.....	5,217	42	154	144	135	695	921	658	468	442	389	281
1999.....	4,880	40	157	108	142	615	842	640	485	392	366	246

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Hispanic, male												
2003.....	4,574	25	77	95	114	592	845	685	490	390	350	251
2002.....	4,481	16	113	80	93	589	890	624	470	372	350	231
2001.....	4,334	16	92	71	84	527	856	641	426	402	329	216
2000.....	3,921	24	85	71	84	528	757	552	369	337	302	204
1999.....	3,630	22	85	65	86	452	684	522	382	307	283	184
Hispanic, female												
2003.....	1,471	14	62	52	47	225	172	136	101	115	118	79
2002.....	1,440	10	69	58	46	196	200	136	110	108	100	86
2001.....	1,440	27	67	49	61	191	166	127	99	112	99	97
2000.....	1,296	18	69	73	51	167	164	106	99	105	87	77
1999.....	1,250	18	72	43	56	163	158	118	103	85	83	62
Non-Hispanic 1/, both sexes												
2003.....	41,856	109	505	545	909	4,642	4,928	3,022	2,824	3,061	3,536	3,313
2002.....	42,268	101	457	565	891	4,986	4,926	3,121	2,999	3,254	3,476	3,254
2001.....	41,339	108	517	648	906	4,659	4,712	3,177	2,964	3,351	3,497	3,054
2000.....	41,341	132	533	695	942	4,760	4,615	3,283	2,961	3,436	3,365	2,957
1999.....	41,330	149	553	786	933	4,820	4,372	3,260	3,012	3,461	3,305	2,859
Non-Hispanic 1/, male												
2003.....	28,591	56	286	304	568	3,052	3,661	2,221	2,099	2,184	2,553	2,408
2002.....	28,906	56	267	323	539	3,280	3,690	2,359	2,203	2,318	2,437	2,388
2001.....	28,327	52	298	375	582	3,155	3,551	2,367	2,211	2,330	2,525	2,226
2000.....	28,096	70	304	392	590	3,151	3,429	2,452	2,172	2,456	2,440	2,126
1999.....	27,850	79	313	464	585	3,133	3,263	2,415	2,163	2,422	2,342	2,037

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Non-Hispanic 1/, female												
2003.....	13,265	53	219	241	341	1,590	1,267	801	725	877	983	905
2002.....	13,362	45	190	242	352	1,706	1,236	762	796	936	1,039	866
2001.....	13,012	56	219	273	324	1,504	1,161	810	753	1,021	972	828
2000.....	13,245	62	229	303	352	1,609	1,186	831	789	980	925	831
1999.....	13,480	70	240	322	348	1,687	1,109	845	849	1,039	963	822
Non-Hispanic White, both sexes												
2003.....	34,220	73	352	392	703	3,880	3,946	2,314	2,166	2,414	2,862	2,708
2002.....	34,689	59	300	392	684	4,195	3,951	2,345	2,358	2,547	2,842	2,659
2001.....	33,753	74	353	440	682	3,851	3,712	2,472	2,299	2,664	2,830	2,431
2000.....	33,808	79	357	507	739	4,020	3,649	2,461	2,298	2,760	2,711	2,394
1999.....	33,803	98	349	557	734	4,044	3,449	2,528	2,361	2,737	2,688	2,307
Non-Hispanic White, male												
2003.....	23,332	39	208	211	437	2,526	2,913	1,703	1,616	1,715	2,081	1,968
2002.....	23,628	35	168	227	416	2,741	2,966	1,769	1,732	1,814	2,002	1,984
2001.....	23,047	36	208	250	436	2,584	2,817	1,842	1,719	1,844	2,043	1,772
2000.....	22,855	44	205	291	457	2,637	2,724	1,826	1,692	1,970	1,965	1,738
1999.....	22,682	58	192	325	457	2,595	2,586	1,876	1,685	1,917	1,910	1,644
Non-Hispanic White, female												
2003.....	10,888	34	144	181	266	1,354	1,033	611	550	699	781	740
2002.....	11,061	24	132	165	268	1,454	985	576	626	733	840	675
2001.....	10,706	38	145	190	246	1,267	895	630	580	820	787	659
2000.....	10,953	35	152	216	282	1,383	925	635	606	790	746	656
1999.....	11,121	40	157	232	277	1,449	863	652	676	820	778	663

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Transport accidents (V01-V99,Y85)												
Non-Hispanic Black, both sexes												
2003.....	5,684	28	125	117	159	512	709	532	507	502	538	472
2002.....	5,665	36	121	145	157	554	721	598	491	547	486	469
2001.....	5,803	18	127	177	177	578	744	541	517	521	550	494
2000.....	5,757	40	139	151	158	528	734	625	527	523	536	453
1999.....	5,810	44	168	199	161	571	691	569	506	583	491	443
Non-Hispanic Black, male												
2003.....	4,053	12	63	73	100	365	552	400	386	376	379	355
2002.....	4,051	18	79	79	98	381	536	465	372	397	336	334
2001.....	4,143	7	69	112	122	414	552	408	389	369	410	381
2000.....	4,127	21	77	83	108	369	538	481	394	393	403	318
1999.....	4,097	18	96	124	105	400	509	436	381	409	360	327
Non-Hispanic Black, female												
2003.....	1,631	16	62	44	59	147	157	132	121	126	159	117
2002.....	1,614	18	42	66	59	173	185	133	119	150	150	135
2001.....	1,660	11	58	65	55	164	192	133	128	152	140	113
2000.....	1,630	19	62	68	50	159	196	144	133	130	133	135
1999.....	1,713	26	72	75	56	171	182	133	125	174	131	116
Origin not stated 2/, both sexes												
2003.....	170	-	5	2	1	12	16	12	10	13	11	13
2002.....	177	-	1	2	3	11	18	10	14	16	12	22
2001.....	175	-	2	1	3	16	12	8	12	16	21	12
2000.....	191	-	4	3	3	20	17	16	14	11	18	10
1999.....	213	-	2	3	3	17	23	22	15	16	13	16

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

Origin not stated 2/, male

2003.....	126	-	3	2	1	9	12	9	7	11	9	8
2002.....	146	-	1	2	2	6	16	7	13	13	9	19
2001.....	133	-	2	1	2	12	10	7	9	12	15	10
2000.....	149	-	2	1	-	14	11	12	13	9	16	8
1999.....	162	-	2	3	3	8	18	15	14	12	10	13

Origin not stated 2/, female

2003.....	44	-	2	-	-	3	4	3	3	2	2	5
2002.....	31	-	-	-	1	5	2	3	1	3	3	3
2001.....	42	-	-	-	1	4	2	1	3	4	6	2
2000.....	42	-	2	2	3	6	6	4	1	2	2	2
1999.....	51	-	-	-	-	9	5	7	1	4	3	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All origins, both sexes												
2003.....	44,757	145	621	646	996	5,269	5,703	3,644	3,189	3,289	3,672	3,286
2002.....	45,380	123	610	664	950	5,634	5,825	3,731	3,361	3,462	3,615	3,240
2001.....	43,788	144	626	718	968	5,208	5,517	3,710	3,227	3,545	3,538	3,011
2000.....	43,354	168	651	780	992	5,251	5,309	3,715	3,169	3,545	3,382	2,936
1999.....	42,401	184	650	802	969	5,198	4,930	3,604	3,174	3,460	3,278	2,747
All origins, male												
2003.....	30,655	80	353	373	636	3,500	4,306	2,753	2,403	2,350	2,634	2,366
2002.....	31,064	68	361	385	578	3,765	4,413	2,855	2,494	2,459	2,525	2,328
2001.....	30,064	66	356	418	611	3,546	4,232	2,826	2,410	2,473	2,543	2,164
2000.....	29,451	90	368	429	614	3,527	3,989	2,804	2,331	2,507	2,433	2,091
1999.....	28,552	97	366	469	593	3,412	3,715	2,690	2,303	2,424	2,311	1,936
All origins, female												
2003.....	14,102	65	268	273	360	1,769	1,397	891	786	939	1,038	920
2002.....	14,316	55	249	279	372	1,869	1,412	876	867	1,003	1,090	912
2001.....	13,724	78	270	300	357	1,662	1,285	884	817	1,072	995	847
2000.....	13,903	78	283	351	378	1,724	1,320	911	838	1,038	949	845
1999.....	13,849	87	284	333	376	1,786	1,215	914	871	1,036	967	811
Hispanic, both sexes												
2003.....	5,753	39	134	141	154	799	983	772	560	475	436	307
2002.....	5,659	26	176	129	130	763	1,047	737	557	458	427	294
2001.....	5,272	40	142	112	134	701	976	716	481	458	376	266
2000.....	4,900	40	144	135	125	671	886	627	439	406	350	262
1999.....	4,525	40	148	100	131	587	797	593	442	354	330	224

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Hispanic, male												
2003.....	4,340	25	73	91	107	578	819	646	463	364	321	232
2002.....	4,262	16	108	74	89	571	850	604	448	353	330	213
2001.....	3,987	15	82	67	77	513	817	600	392	357	296	192
2000.....	3,658	24	78	65	77	510	726	523	346	304	270	186
1999.....	3,337	22	82	59	77	431	646	480	348	275	255	165
Hispanic, female												
2003.....	1,413	14	61	50	47	221	164	126	97	111	115	75
2002.....	1,397	10	68	55	41	192	197	133	109	105	97	81
2001.....	1,285	25	60	45	57	188	159	116	89	101	80	74
2000.....	1,242	16	66	70	48	161	160	104	93	102	80	76
1999.....	1,188	18	66	41	54	156	151	113	94	79	75	59
Non-Hispanic 1/, both sexes												
2003.....	38,849	106	482	503	841	4,460	4,705	2,861	2,619	2,803	3,228	2,966
2002.....	39,567	97	433	533	817	4,861	4,760	2,985	2,792	2,990	3,179	2,927
2001.....	38,362	104	483	605	831	4,491	4,530	2,987	2,735	3,073	3,144	2,735
2000.....	38,295	128	505	643	864	4,561	4,408	3,075	2,717	3,130	3,017	2,665
1999.....	37,684	144	500	699	835	4,594	4,113	2,992	2,720	3,090	2,936	2,509
Non-Hispanic 1/, male												
2003.....	26,202	55	277	280	528	2,915	3,476	2,099	1,933	1,977	2,306	2,126
2002.....	26,677	52	252	309	487	3,188	3,547	2,245	2,035	2,095	2,189	2,098
2001.....	25,962	51	273	350	532	3,021	3,405	2,220	2,010	2,106	2,234	1,964
2000.....	25,670	66	288	363	537	3,004	3,253	2,270	1,973	2,196	2,150	1,898
1999.....	25,072	75	282	407	513	2,973	3,054	2,198	1,943	2,137	2,047	1,760

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic 1/, female												
2003.....	12,647	51	205	223	313	1,545	1,229	762	686	826	922	840
2002.....	12,890	45	181	224	330	1,673	1,213	740	757	895	990	829
2001.....	12,400	53	210	255	299	1,470	1,125	767	725	967	910	771
2000.....	12,625	62	217	280	327	1,557	1,155	805	744	934	867	767
1999.....	12,612	69	218	292	322	1,621	1,059	794	777	953	889	749
Non-Hispanic White, both sexes												
2003.....	31,662	70	333	361	644	3,731	3,764	2,180	2,002	2,195	2,597	2,423
2002.....	32,405	56	280	370	630	4,090	3,807	2,231	2,181	2,326	2,597	2,386
2001.....	31,283	71	323	412	624	3,711	3,555	2,310	2,114	2,444	2,536	2,179
2000.....	31,219	75	340	464	672	3,861	3,469	2,281	2,098	2,508	2,415	2,155
1999.....	30,743	94	315	491	653	3,860	3,239	2,301	2,124	2,426	2,371	2,011
Non-Hispanic White, male												
2003.....	21,309	38	200	196	406	2,414	2,763	1,601	1,484	1,541	1,867	1,737
2002.....	21,743	32	155	216	378	2,664	2,841	1,675	1,587	1,625	1,798	1,739
2001.....	21,073	35	185	232	399	2,470	2,694	1,716	1,554	1,666	1,802	1,560
2000.....	20,789	40	197	269	411	2,523	2,571	1,669	1,527	1,754	1,719	1,550
1999.....	20,362	55	175	284	399	2,467	2,417	1,693	1,508	1,678	1,660	1,410
Non-Hispanic White, female												
2003.....	10,353	32	133	165	238	1,317	1,001	579	518	654	730	686
2002.....	10,662	24	125	154	252	1,426	966	556	594	701	799	647
2001.....	10,210	36	138	180	225	1,241	861	594	560	778	734	619
2000.....	10,430	35	143	195	261	1,338	898	612	571	754	696	605
1999.....	10,381	39	140	207	254	1,393	822	608	616	748	711	601

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic Black, both sexes												
2003.....	5,349	28	121	110	151	488	680	511	480	471	503	424
2002.....	5,338	35	118	135	143	538	705	581	468	509	450	419
2001.....	5,423	18	123	163	161	557	728	519	488	484	501	441
2000.....	5,424	40	132	143	151	503	713	606	489	481	496	412
1999.....	5,388	43	152	182	149	544	665	540	466	534	451	403
Non-Hispanic Black, male												
2003.....	3,771	12	62	67	92	346	525	384	363	349	353	315
2002.....	3,774	17	77	76	87	368	523	449	353	367	306	292
2001.....	3,842	7	67	105	109	398	537	391	366	339	367	338
2000.....	3,857	21	72	76	103	348	520	463	365	358	369	286
1999.....	3,763	17	84	109	94	379	490	412	349	371	325	296
Non-Hispanic Black, female												
2003.....	1,578	16	59	43	59	142	155	127	117	122	150	109
2002.....	1,564	18	41	59	56	170	182	132	115	142	144	127
2001.....	1,581	11	56	58	52	159	191	128	122	145	134	103
2000.....	1,567	19	60	67	48	155	193	143	124	123	127	126
1999.....	1,625	26	68	73	55	165	175	128	117	163	126	107
Origin not stated 2/, both sexes												
2003.....	155	-	5	2	1	10	15	11	10	11	8	13
2002.....	154	-	1	2	3	10	18	9	12	14	9	19
2001.....	154	-	1	1	3	16	11	7	11	14	18	10
2000.....	159	-	2	2	3	19	15	13	13	9	15	9
1999.....	192	-	2	3	3	17	20	19	12	16	12	14

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Origin not stated 2/, male

2003.....	113	-	3	2	1	7	11	8	7	9	7	8
2002.....	125	-	1	2	2	6	16	6	11	11	6	17
2001.....	115	-	1	1	2	12	10	6	8	10	13	8
2000.....	123	-	2	1	-	13	10	11	12	7	13	7
1999.....	143	-	2	3	3	8	15	12	12	12	9	11

Origin not stated 2/, female

2003.....	42	-	2	-	-	3	4	3	3	2	1	5
2002.....	29	-	-	-	1	4	2	3	1	3	3	2
2001.....	39	-	-	-	1	4	1	1	3	4	5	2
2000.....	36	-	-	1	3	6	5	2	1	2	2	2
1999.....	49	-	-	-	-	9	5	7	-	4	3	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
All origins, both sexes												
2003.....	1,357	2	17	23	37	129	135	100	104	107	132	137
2002.....	1,086	2	17	21	49	63	96	59	79	101	122	121
2001.....	1,405	3	28	25	55	95	103	85	92	141	144	142
2000.....	1,492	2	31	39	59	128	115	94	118	149	168	125
1999.....	2,118	4	48	62	66	172	190	152	165	205	203	179
All origins, male												
2003.....	1,044	-	9	12	26	95	110	80	86	85	101	114
2002.....	882	2	10	11	36	49	85	47	61	85	102	107
2001.....	1,067	-	20	13	36	72	86	71	81	117	118	105
2000.....	1,125	2	17	21	40	99	95	81	90	121	137	92
1999.....	1,559	4	26	43	48	119	160	126	118	157	162	135
All origins, female												
2003.....	313	2	8	11	11	34	25	20	18	22	31	23
2002.....	204	-	7	10	13	14	11	12	18	16	20	14
2001.....	338	3	8	12	19	23	17	14	11	24	26	37
2000.....	367	-	14	18	19	29	20	13	28	28	31	33
1999.....	559	-	22	19	18	53	30	26	47	48	41	44
Hispanic, both sexes												
2003.....	167	-	3	6	4	11	21	29	19	16	18	12
2002.....	144	-	2	5	5	14	28	11	10	13	15	12
2001.....	200	-	8	2	5	13	26	24	14	20	22	19
2000.....	201	-	10	7	8	15	19	15	21	19	25	15
1999.....	264	-	8	6	8	20	35	29	31	28	29	16

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Hispanic, male												
2003.....	139	-	2	4	4	7	17	25	18	15	17	11
2002.....	119	-	1	4	2	12	26	9	9	12	13	10
2001.....	173	-	6	-	4	12	25	23	12	19	20	13
2000.....	166	-	7	5	6	13	18	15	17	17	19	14
1999.....	220	-	3	5	7	16	29	28	25	24	23	13
Hispanic, female												
2003.....	28	-	1	2	-	4	4	4	1	1	1	1
2002.....	25	-	1	1	3	2	2	2	1	1	2	2
2001.....	27	-	2	2	1	1	1	1	2	1	2	6
2000.....	35	-	3	2	2	2	1	-	4	2	6	1
1999.....	44	-	5	1	1	4	6	1	6	4	6	3
Non-Hispanic 1/, both sexes												
2003.....	1,185	2	14	17	33	118	114	71	85	90	113	125
2002.....	929	2	15	16	44	48	68	48	69	88	104	107
2001.....	1,187	3	19	23	50	82	76	60	77	120	121	121
2000.....	1,262	2	19	31	51	112	94	76	96	128	140	110
1999.....	1,836	4	40	56	58	152	152	121	131	177	173	162
Non-Hispanic 1/, male												
2003.....	901	-	7	8	22	88	93	55	68	69	84	103
2002.....	752	2	9	7	34	37	59	38	52	73	86	96
2001.....	878	-	13	13	32	60	61	47	68	97	97	90
2000.....	936	2	10	16	34	85	76	65	72	102	115	78
1999.....	1,323	4	23	38	41	103	128	96	91	133	138	121

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Non-Hispanic 1/, female												
2003.....	284	2	7	9	11	30	21	16	17	21	29	22
2002.....	177	-	6	9	10	11	9	10	17	15	18	11
2001.....	309	3	6	10	18	22	15	13	9	23	24	31
2000.....	326	-	9	15	17	27	18	11	24	26	25	32
1999.....	513	-	17	18	17	49	24	25	40	44	35	41
Non-Hispanic White, both sexes												
2003.....	952	2	11	10	29	99	96	56	61	71	88	91
2002.....	719	1	11	12	32	39	56	38	56	67	73	77
2001.....	901	2	16	14	35	62	64	50	53	79	86	88
2000.....	997	2	9	26	41	94	74	59	73	97	104	80
1999.....	1,480	3	25	39	47	122	117	96	99	141	141	130
Non-Hispanic White, male												
2003.....	715	-	6	3	18	74	79	42	49	53	69	71
2002.....	582	1	7	7	25	31	50	29	42	57	59	68
2001.....	673	-	12	9	21	48	51	40	49	65	68	67
2000.....	741	2	3	13	28	70	60	51	55	79	86	57
1999.....	1,051	3	12	25	32	81	96	75	68	106	112	95
Non-Hispanic White, female												
2003.....	237	2	5	7	11	25	17	14	12	18	19	20
2002.....	137	-	4	5	7	8	6	9	14	10	14	9
2001.....	228	2	4	5	14	14	13	10	4	14	18	21
2000.....	256	-	6	13	13	24	14	8	18	18	18	23
1999.....	429	-	13	14	15	41	21	21	31	35	29	35

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Non-Hispanic Black, both sexes												
2003.....	188	-	3	5	3	14	13	13	17	16	23	29
2002.....	168	1	3	4	9	9	8	10	9	18	22	29
2001.....	216	-	3	9	14	15	6	8	16	27	32	26
2000.....	203	-	7	4	7	13	17	11	21	26	28	24
1999.....	248	1	12	14	6	18	15	18	24	29	23	23
Non-Hispanic Black, male												
2003.....	154	-	1	4	3	10	11	11	14	14	14	27
2002.....	143	1	2	-	7	6	6	9	8	14	20	28
2001.....	159	-	1	4	11	10	5	6	12	21	27	20
2000.....	155	-	5	3	5	10	14	10	15	21	22	19
1999.....	192	1	9	13	6	14	13	16	18	22	19	18
Non-Hispanic Black, female												
2003.....	34	-	2	1	-	4	2	2	3	2	9	2
2002.....	25	-	1	4	2	3	2	1	1	4	2	1
2001.....	57	-	2	5	3	5	1	2	4	6	5	6
2000.....	48	-	2	1	2	3	3	1	6	5	6	5
1999.....	56	-	3	1	-	4	2	2	6	7	4	5
Origin not stated 2/, both sexes												
2003.....	5	-	-	-	-	-	-	-	-	1	1	-
2002.....	13	-	-	-	-	1	-	-	-	-	3	2
2001.....	18	-	1	-	-	-	1	1	1	1	1	2
2000.....	29	-	2	1	-	1	2	3	1	2	3	-
1999.....	18	-	-	-	-	-	3	2	3	-	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Origin not stated 2/, male

2003.....	4	-	-	-	-	-	-	-	-	1	-	-
2002.....	11	-	-	-	-	-	-	-	-	-	3	1
2001.....	16	-	1	-	-	-	1	1	1	1	1	2
2000.....	23	-	-	-	-	1	1	1	1	2	3	-
1999.....	16	-	-	-	-	-	3	2	2	-	1	1

Origin not stated 2/, female

2003.....	1	-	-	-	-	-	-	-	-	-	1	-
2002.....	2	-	-	-	-	1	-	-	-	-	-	1
2001.....	2	-	-	-	-	-	1	-	-	-	-	-
2000.....	6	-	2	1	-	-	1	2	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All origins, both sexes												
2003.....	1,957	1	11	25	38	73	123	111	132	183	211	233
2002.....	1,900	2	13	20	34	85	113	101	153	187	201	232
2001.....	2,095	4	24	26	31	90	126	158	182	195	264	226
2000.....	1,903	4	9	23	29	96	129	148	156	195	222	187
1999.....	1,904	1	14	33	43	82	117	166	173	204	203	195
All origins, male												
2003.....	1,592	1	4	16	21	58	102	82	107	150	177	187
2002.....	1,587	2	10	9	20	61	98	88	131	159	169	203
2001.....	1,663	2	16	16	21	76	99	118	155	154	208	183
2000.....	1,590	2	6	14	20	67	113	131	133	174	188	155
1999.....	1,531	-	8	20	33	62	90	136	138	160	162	163
All origins, female												
2003.....	365	-	7	9	17	15	21	29	25	33	34	46
2002.....	313	-	3	11	14	24	15	13	22	28	32	29
2001.....	432	2	8	10	10	14	27	40	27	41	56	43
2000.....	313	2	3	9	9	29	16	17	23	21	34	32
1999.....	373	1	6	13	10	20	27	30	35	44	41	32
Hispanic, both sexes												
2003.....	125	-	2	-	3	7	13	20	12	14	14	11
2002.....	118	-	4	4	4	8	15	12	13	9	8	11
2001.....	302	3	9	6	6	4	20	28	30	36	30	28
2000.....	116	2	-	2	2	9	16	16	8	17	14	4
1999.....	91	-	1	2	3	8	10	18	12	10	7	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Hispanic, male												
2003.....	95	-	2	-	3	7	9	14	9	11	12	8
2002.....	100	-	4	2	2	6	14	11	13	7	7	8
2001.....	174	1	4	4	3	2	14	18	22	26	13	11
2000.....	97	-	-	1	1	5	13	14	6	16	13	4
1999.....	73	-	-	1	2	5	9	14	9	8	5	6
Hispanic, female												
2003.....	30	-	-	-	-	-	4	6	3	3	2	3
2002.....	18	-	-	2	2	2	1	1	-	2	1	3
2001.....	128	2	5	2	3	2	6	10	8	10	17	17
2000.....	19	2	-	1	1	4	3	2	2	1	1	-
1999.....	18	-	1	1	1	3	1	4	3	2	2	-
Non-Hispanic 1/, both sexes												
2003.....	1,822	1	9	25	35	64	109	90	120	168	195	222
2002.....	1,772	2	9	16	30	77	98	88	138	176	193	220
2001.....	1,790	1	15	20	25	86	106	130	152	158	232	198
2000.....	1,784	2	9	21	27	87	113	132	148	178	208	182
1999.....	1,810	1	13	31	40	74	107	147	161	194	196	188
Non-Hispanic 1/, male												
2003.....	1,488	1	2	16	18	49	92	67	98	138	163	179
2002.....	1,477	2	6	7	18	55	84	76	116	150	162	194
2001.....	1,487	1	12	12	18	74	85	100	133	127	194	172
2000.....	1,490	2	6	13	19	62	100	117	127	158	175	150
1999.....	1,455	-	8	19	31	57	81	121	129	152	157	156

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic 1/, female												
2003.....	334	-	7	9	17	15	17	23	22	30	32	43
2002.....	295	-	3	9	12	22	14	12	22	26	31	26
2001.....	303	-	3	8	7	12	21	30	19	31	38	26
2000.....	294	-	3	8	8	25	13	15	21	20	33	32
1999.....	355	1	5	12	9	17	26	26	32	42	39	32
Non-Hispanic White, both sexes												
2003.....	1,606	1	8	21	30	50	86	78	103	148	177	194
2002.....	1,565	2	9	10	22	66	88	76	121	154	172	196
2001.....	1,569	1	14	14	23	78	93	112	132	141	208	164
2000.....	1,592	2	8	17	26	65	106	121	127	155	192	159
1999.....	1,580	1	9	27	34	62	93	131	138	170	176	166
Non-Hispanic White, male												
2003.....	1,308	1	2	12	13	38	71	60	83	121	145	160
2002.....	1,303	2	6	4	13	46	75	65	103	132	145	177
2001.....	1,301	1	11	9	16	66	72	86	116	113	173	145
2000.....	1,325	2	5	9	18	44	93	106	110	137	160	131
1999.....	1,269	-	5	16	26	47	73	108	109	133	138	139
Non-Hispanic White, female												
2003.....	298	-	6	9	17	12	15	18	20	27	32	34
2002.....	262	-	3	6	9	20	13	11	18	22	27	19
2001.....	268	-	3	5	7	12	21	26	16	28	35	19
2000.....	267	-	3	8	8	21	13	15	17	18	32	28
1999.....	311	1	4	11	8	15	20	23	29	37	38	27

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Non-Hispanic Black, both sexes												
2003.....	147	-	1	2	5	10	16	8	10	15	12	19
2002.....	159	-	-	6	5	7	8	7	14	20	14	21
2001.....	164	-	1	5	2	6	10	14	13	10	17	27
2000.....	130	-	-	4	-	12	4	8	17	16	12	17
1999.....	174	-	4	3	6	9	11	11	16	20	17	17
Non-Hispanic Black, male												
2003.....	128	-	-	2	5	9	16	5	9	13	12	13
2002.....	134	-	-	3	4	7	7	7	11	16	10	14
2001.....	142	-	1	3	2	6	10	11	11	9	16	23
2000.....	115	-	-	4	-	11	4	8	14	14	12	13
1999.....	142	-	3	2	5	7	6	8	14	16	16	13
Non-Hispanic Black, female												
2003.....	19	-	1	-	-	1	-	3	1	2	-	6
2002.....	25	-	-	3	1	-	1	-	3	4	4	7
2001.....	22	-	-	2	-	-	-	3	2	1	1	4
2000.....	15	-	-	-	-	1	-	-	3	2	-	4
1999.....	32	-	1	1	1	2	5	3	2	4	1	4
Origin not stated 2/, both sexes												
2003.....	10	-	-	-	-	2	1	1	-	1	2	-
2002.....	10	-	-	-	-	-	-	1	2	2	-	1
2001.....	3	-	-	-	-	-	-	-	-	1	2	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	1
1999.....	3	-	-	-	-	-	-	1	-	-	-	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
All origins, both sexes												
2003.....	61,206	797	1,068	402	451	1,284	2,556	2,429	2,832	3,916	5,256	5,209
2002.....	58,376	819	1,001	471	509	1,355	2,241	2,231	2,854	3,892	5,130	4,648
2001.....	54,249	825	1,036	514	499	1,253	2,019	1,982	2,403	3,708	4,410	4,090
2000.....	51,151	707	1,135	549	508	1,280	1,805	1,941	2,428	3,582	4,170	3,622
1999.....	51,437	656	1,186	562	554	1,236	1,731	1,959	2,497	3,497	4,181	3,575
All origins, male												
2003.....	37,241	458	682	249	329	1,053	2,067	1,911	2,112	2,805	3,701	3,658
2002.....	35,724	481	643	315	359	1,106	1,861	1,778	2,181	2,832	3,681	3,302
2001.....	33,266	473	653	339	349	1,028	1,693	1,590	1,841	2,758	3,241	3,067
2000.....	31,651	407	707	360	363	1,064	1,506	1,590	1,899	2,735	3,149	2,765
1999.....	31,893	393	727	365	403	1,031	1,451	1,603	1,983	2,670	3,202	2,762
All origins, female												
2003.....	23,965	339	386	153	122	231	489	518	720	1,111	1,555	1,551
2002.....	22,652	338	358	156	150	249	380	453	673	1,060	1,449	1,346
2001.....	20,983	352	383	175	150	225	326	392	562	950	1,169	1,023
2000.....	19,500	300	428	189	145	216	299	351	529	847	1,021	857
1999.....	19,544	263	459	197	151	205	280	356	514	827	979	813
Hispanic, both sexes												
2003.....	4,373	92	186	71	52	188	344	387	368	443	461	405
2002.....	4,185	101	159	61	51	199	339	322	365	418	493	368
2001.....	3,749	111	151	66	48	161	281	305	340	428	390	314
2000.....	3,613	76	152	61	43	189	290	286	364	414	373	261
1999.....	3,770	94	184	62	50	185	304	296	369	452	397	300

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
Hispanic, male												
2003.....	3,325	52	123	43	37	164	304	337	315	373	375	334
2002.....	3,217	67	102	40	31	174	305	293	322	340	406	291
2001.....	2,823	59	103	43	26	142	248	267	285	361	326	265
2000.....	2,775	43	89	34	21	161	271	252	326	345	333	231
1999.....	2,948	58	112	42	34	156	283	265	325	385	332	259
Hispanic, female												
2003.....	1,048	40	63	28	15	24	40	50	53	70	86	71
2002.....	968	34	57	21	20	25	34	29	43	78	87	77
2001.....	926	52	48	23	22	19	33	38	55	67	64	49
2000.....	838	33	63	27	22	28	19	34	38	69	40	30
1999.....	822	36	72	20	16	29	21	31	44	67	65	41
Non-Hispanic 1/, both sexes												
2003.....	56,550	703	881	330	399	1,090	2,206	2,029	2,451	3,451	4,765	4,769
2002.....	53,907	715	839	407	455	1,150	1,894	1,898	2,471	3,451	4,611	4,255
2001.....	50,183	709	882	443	448	1,089	1,727	1,669	2,047	3,251	3,990	3,739
2000.....	47,217	627	979	485	463	1,088	1,505	1,644	2,047	3,144	3,761	3,317
1999.....	47,339	561	997	500	502	1,045	1,421	1,653	2,109	3,020	3,743	3,240
Non-Hispanic 1/, male												
2003.....	33,699	404	558	205	292	884	1,758	1,564	1,786	2,416	3,301	3,299
2002.....	32,295	412	539	273	326	928	1,550	1,476	1,844	2,472	3,255	2,992
2001.....	30,201	410	547	293	320	883	1,435	1,317	1,543	2,371	2,891	2,773
2000.....	28,640	361	617	325	341	902	1,225	1,329	1,559	2,370	2,788	2,500
1999.....	28,701	335	613	323	367	870	1,162	1,330	1,642	2,265	2,842	2,473

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic 1/, female												
2003.....	22,851	299	323	125	107	206	448	465	665	1,035	1,464	1,470
2002.....	21,612	303	300	134	129	222	344	422	627	979	1,356	1,263
2001.....	19,982	299	335	150	128	206	292	352	504	880	1,099	966
2000.....	18,577	266	362	160	122	186	280	315	488	774	973	817
1999.....	18,638	226	384	177	135	175	259	323	467	755	901	767
Non-Hispanic White, both sexes												
2003.....	48,595	423	633	200	240	894	1,918	1,706	2,052	2,845	3,985	3,885
2002.....	45,916	403	565	224	281	904	1,614	1,576	2,065	2,811	3,835	3,421
2001.....	42,509	426	653	282	284	841	1,463	1,347	1,680	2,650	3,196	2,930
2000.....	39,728	376	676	289	308	832	1,234	1,327	1,644	2,542	2,995	2,584
1999.....	39,523	337	693	286	331	796	1,121	1,326	1,640	2,449	2,970	2,482
Non-Hispanic White, male												
2003.....	28,556	244	402	127	179	725	1,534	1,323	1,496	1,988	2,790	2,675
2002.....	27,024	244	359	151	203	725	1,314	1,223	1,549	2,008	2,726	2,428
2001.....	25,133	245	416	174	206	682	1,214	1,066	1,279	1,950	2,351	2,169
2000.....	23,615	222	433	196	232	689	1,015	1,073	1,270	1,944	2,234	1,942
1999.....	23,431	200	425	180	243	655	913	1,077	1,294	1,853	2,269	1,887
Non-Hispanic White, female												
2003.....	20,039	179	231	73	61	169	384	383	556	857	1,195	1,210
2002.....	18,892	159	206	73	78	179	300	353	516	803	1,109	993
2001.....	17,376	181	237	108	78	159	249	281	401	700	845	761
2000.....	16,113	154	243	93	76	143	219	254	374	598	761	642
1999.....	16,092	137	268	106	88	141	208	249	346	596	701	595

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
Non-Hispanic Black, both sexes												
2003.....	6,470	255	208	116	125	144	225	245	318	495	660	775
2002.....	6,620	288	229	153	152	186	219	255	332	537	657	742
2001.....	6,436	254	204	136	140	182	209	243	299	505	706	718
2000.....	6,330	228	257	179	131	193	207	252	336	519	675	643
1999.....	6,723	207	260	193	148	193	239	272	391	480	691	695
Non-Hispanic Black, male												
2003.....	4,184	146	134	68	89	117	171	183	228	349	435	541
2002.....	4,377	154	152	101	113	160	183	200	242	389	440	502
2001.....	4,238	151	119	100	98	150	169	184	211	349	480	537
2000.....	4,253	124	155	121	94	161	165	200	241	366	487	490
1999.....	4,532	122	164	130	108	168	204	211	285	345	514	537
Non-Hispanic Black, female												
2003.....	2,286	109	74	48	36	27	54	62	90	146	225	234
2002.....	2,243	134	77	52	39	26	36	55	90	148	217	240
2001.....	2,198	103	85	36	42	32	40	59	88	156	226	181
2000.....	2,077	104	102	58	37	32	42	52	95	153	188	153
1999.....	2,191	85	96	63	40	25	35	61	106	135	177	158
Origin not stated 2/, both sexes												
2003.....	283	2	1	1	-	6	6	13	13	22	30	35
2002.....	284	3	3	3	3	6	8	11	18	23	26	25
2001.....	317	5	3	5	3	3	11	8	16	29	30	37
2000.....	321	4	4	3	2	3	10	11	17	24	36	44
1999.....	328	1	5	-	2	6	6	10	19	25	41	35

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nontransport accidents (W00-X59,Y86)												
Origin not stated 2/, male												
2003.....	217	2	1	1	-	5	5	10	11	16	25	25
2002.....	212	2	2	2	2	4	6	9	15	20	20	19
2001.....	242	4	3	3	3	3	10	6	13	26	24	29
2000.....	236	3	1	1	1	1	10	9	14	20	28	34
1999.....	244	-	2	-	2	5	6	8	16	20	28	30
Origin not stated 2/, female												
2003.....	66	-	-	-	-	1	1	3	2	6	5	10
2002.....	72	1	1	1	1	2	2	2	3	3	6	6
2001.....	75	1	-	2	-	-	1	2	3	3	6	8
2000.....	85	1	3	2	1	2	-	2	3	4	8	10
1999.....	84	1	3	-	-	1	-	2	3	5	13	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
All origins, both sexes												
2003.....	17,229	14	54	14	31	90	140	139	146	255	381	508
2002.....	16,257	16	37	18	24	83	164	130	177	256	408	487
2001.....	15,019	23	32	33	33	88	168	160	180	272	375	520
2000.....	13,322	8	36	16	21	99	138	138	165	238	370	409
1999.....	13,162	12	55	25	28	112	130	156	187	276	369	447
All origins, male												
2003.....	8,910	9	39	9	21	71	115	123	114	218	307	391
2002.....	8,463	9	28	9	18	65	146	118	151	191	331	378
2001.....	8,089	15	25	22	24	79	153	142	153	233	311	426
2000.....	7,122	6	23	14	16	88	115	122	150	199	297	332
1999.....	7,109	8	37	19	21	95	114	137	157	225	303	366
All origins, female												
2003.....	8,319	5	15	5	10	19	25	16	32	37	74	117
2002.....	7,794	7	9	9	6	18	18	12	26	65	77	109
2001.....	6,930	8	7	11	9	9	15	18	27	39	64	94
2000.....	6,200	2	13	2	5	11	23	16	15	39	73	77
1999.....	6,053	4	18	6	7	17	16	19	30	51	66	81
Hispanic, both sexes												
2003.....	840	1	11	4	6	10	28	36	24	33	50	50
2002.....	832	4	9	4	2	12	39	32	37	40	54	44
2001.....	757	7	6	10	4	15	29	35	38	44	56	52
2000.....	693	3	9	1	2	18	28	36	37	31	35	39
1999.....	652	4	9	1	2	13	32	35	33	33	37	34

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Falls (W00-W19)</b>												
<b>Hispanic, male</b>												
2003.....	557	1	8	2	3	9	24	34	22	31	49	43
2002.....	563	3	7	1	2	12	37	30	35	31	50	35
2001.....	525	4	5	9	4	15	29	34	37	40	53	49
2000.....	485	2	8	1	2	18	26	34	36	27	35	35
1999.....	464	4	6	1	2	12	31	34	31	30	32	28
<b>Hispanic, female</b>												
2003.....	283	-	3	2	3	1	4	2	2	2	1	7
2002.....	269	1	2	3	-	-	2	2	2	9	4	9
2001.....	232	3	1	1	-	-	-	1	1	4	3	3
2000.....	208	1	1	-	-	-	2	2	1	4	-	4
1999.....	188	-	3	-	-	1	1	1	2	3	5	6
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	16,337	13	42	10	25	80	112	102	121	221	327	453
2002.....	15,365	12	28	14	22	71	125	98	140	215	353	439
2001.....	14,204	15	26	22	29	73	139	125	141	227	317	461
2000.....	12,585	5	27	15	19	81	110	101	128	205	332	366
1999.....	12,447	8	46	24	26	99	98	121	152	242	331	408
<b>Non-Hispanic 1/, male</b>												
2003.....	8,320	8	30	7	18	62	91	88	91	186	254	345
2002.....	7,856	6	21	8	16	53	109	88	116	159	280	339
2001.....	7,527	10	20	13	20	64	124	108	115	192	256	370
2000.....	6,603	4	15	13	14	70	89	87	114	170	259	293
1999.....	6,602	4	31	18	19	83	83	103	124	194	270	333

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
Non-Hispanic 1/, female												
2003.....	8,017	5	12	3	7	18	21	14	30	35	73	108
2002.....	7,509	6	7	6	6	18	16	10	24	56	73	100
2001.....	6,677	5	6	9	9	9	15	17	26	35	61	91
2000.....	5,982	1	12	2	5	11	21	14	14	35	73	73
1999.....	5,845	4	15	6	7	16	15	18	28	48	61	75
Non-Hispanic White, both sexes												
2003.....	14,995	9	28	9	20	69	98	87	103	177	269	362
2002.....	14,157	8	15	11	17	60	108	87	116	176	290	364
2001.....	13,037	8	23	17	25	65	128	106	120	182	253	373
2000.....	11,537	2	15	11	16	73	99	88	104	163	275	311
1999.....	11,295	4	34	17	22	87	81	101	115	206	271	323
Non-Hispanic White, male												
2003.....	7,531	5	22	6	14	54	82	75	79	151	211	279
2002.....	7,107	3	11	6	13	44	93	78	97	128	231	284
2001.....	6,785	5	17	11	16	56	117	93	98	154	206	295
2000.....	5,949	1	6	9	13	64	80	77	92	136	218	253
1999.....	5,860	3	23	13	16	74	70	87	94	163	217	265
Non-Hispanic White, female												
2003.....	7,464	4	6	3	6	15	16	12	24	26	58	83
2002.....	7,050	5	4	5	4	16	15	9	19	48	59	80
2001.....	6,252	3	6	6	9	9	11	13	22	28	47	78
2000.....	5,588	1	9	2	3	9	19	11	12	27	57	58
1999.....	5,435	1	11	4	6	13	11	14	21	43	54	58

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Falls (W00-W19)												
Non-Hispanic Black, both sexes												
2003.....	898	2	11	1	2	7	10	12	13	31	43	75
2002.....	821	3	7	3	3	8	9	8	15	31	44	63
2001.....	837	6	3	4	2	4	9	9	17	33	57	62
2000.....	745	2	12	4	3	5	7	5	17	28	43	45
1999.....	865	2	11	6	4	8	13	13	29	26	47	73
Non-Hispanic Black, male												
2003.....	519	1	7	1	2	5	5	10	8	23	34	56
2002.....	518	2	6	2	3	6	9	7	13	25	33	47
2001.....	537	4	3	2	2	4	5	6	13	27	45	54
2000.....	471	2	9	4	1	3	7	3	15	22	32	35
1999.....	554	1	8	4	3	6	10	11	23	23	42	59
Non-Hispanic Black, female												
2003.....	379	1	4	-	-	2	5	2	5	8	9	19
2002.....	303	1	1	1	-	2	-	1	2	6	11	16
2001.....	300	2	-	2	-	-	4	3	4	6	12	8
2000.....	274	-	3	-	2	2	-	2	2	6	11	10
1999.....	311	1	3	2	1	2	3	2	6	3	5	14
Origin not stated 2/, both sexes												
2003.....	52	-	1	-	-	-	-	1	1	1	4	5
2002.....	60	-	-	-	-	-	-	-	-	1	1	4
2001.....	58	1	-	1	-	-	-	-	1	1	2	7
2000.....	44	-	-	-	-	-	-	1	-	2	3	4
1999.....	63	-	-	-	-	-	-	-	2	1	1	5



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
All origins, both sexes												
2003.....	730	-	7	13	36	95	105	74	46	57	61	48
2002.....	762	1	11	14	34	107	103	75	68	52	71	52
2001.....	802	-	15	18	39	110	96	71	51	70	76	61
2000.....	776	1	18	18	49	107	95	74	57	75	78	38
1999.....	824	-	12	19	57	126	125	76	67	67	69	51
All origins, male												
2003.....	656	-	7	11	35	88	94	69	41	50	55	43
2002.....	667	1	7	10	29	101	94	70	59	42	56	45
2001.....	690	-	12	14	35	102	91	63	40	57	64	51
2000.....	671	-	11	16	46	97	86	65	44	60	65	31
1999.....	707	-	7	13	47	115	114	62	60	56	56	46
All origins, female												
2003.....	74	-	-	2	1	7	11	5	5	7	6	5
2002.....	95	-	4	4	5	6	9	5	9	10	15	7
2001.....	112	-	3	4	4	8	5	8	11	13	12	10
2000.....	105	1	7	2	3	10	9	9	13	15	13	7
1999.....	117	-	5	6	10	11	11	14	7	11	13	5
Hispanic, both sexes												
2003.....	71	-	-	4	4	17	16	9	5	3	5	1
2002.....	60	-	3	3	2	13	11	11	7	4	1	2
2001.....	86	-	1	1	3	19	17	12	9	7	5	3
2000.....	59	-	2	2	4	12	13	7	5	6	5	2
1999.....	66	-	-	-	3	15	16	7	10	4	4	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidental discharge of firearms (W32-W34)</b>												
<b>Hispanic, male</b>												
2003.....	64	-	-	2	4	16	14	9	5	2	5	1
2002.....	51	-	1	1	2	11	10	11	7	3	-	2
2001.....	76	-	1	1	2	17	16	10	6	7	5	2
2000.....	50	-	1	1	3	11	13	7	3	4	5	2
1999.....	61	-	-	-	3	14	15	6	10	3	4	3
<b>Hispanic, female</b>												
2003.....	7	-	-	2	-	1	2	-	-	1	-	-
2002.....	9	-	2	2	-	2	1	-	-	1	1	-
2001.....	10	-	-	-	1	2	1	2	3	-	-	1
2000.....	9	-	1	1	1	1	-	-	2	2	-	-
1999.....	5	-	-	-	-	1	1	1	-	1	-	1
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	658	-	7	9	32	78	89	65	41	53	56	47
2002.....	700	1	8	11	32	93	92	64	61	48	70	50
2001.....	715	-	14	17	36	90	79	59	42	63	71	58
2000.....	717	1	16	16	45	95	82	67	52	69	73	36
1999.....	758	-	12	19	54	111	109	69	57	63	65	47
<b>Non-Hispanic 1/, male</b>												
2003.....	591	-	7	9	31	72	80	60	36	47	50	42
2002.....	614	1	6	9	27	89	84	59	52	39	56	43
2001.....	613	-	11	13	33	84	75	53	34	50	59	49
2000.....	621	-	10	15	43	86	73	58	41	56	60	29
1999.....	646	-	7	13	44	101	99	56	50	53	52	43

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental discharge of firearms (W32-W34)												
Non-Hispanic 1/, female												
2003.....	67	-	-	-	1	6	9	5	5	6	6	5
2002.....	86	-	2	2	5	4	8	5	9	9	14	7
2001.....	102	-	3	4	3	6	4	6	8	13	12	9
2000.....	96	1	6	1	2	9	9	9	11	13	13	7
1999.....	112	-	5	6	10	10	10	13	7	10	13	4
Non-Hispanic White, both sexes												
2003.....	515	-	4	6	24	55	60	38	31	45	50	38
2002.....	531	1	6	7	24	56	62	40	42	35	62	42
2001.....	591	-	9	13	28	61	56	43	39	57	59	54
2000.....	555	1	9	9	37	65	52	46	37	54	64	28
1999.....	585	-	5	9	45	73	72	55	43	52	56	41
Non-Hispanic White, male												
2003.....	461	-	4	6	23	50	53	34	27	41	46	33
2002.....	459	1	4	5	20	53	57	35	35	29	48	36
2001.....	506	-	7	10	28	56	53	38	33	45	51	45
2000.....	483	-	5	9	35	60	46	38	28	45	54	24
1999.....	495	-	3	4	39	66	65	43	37	43	44	38
Non-Hispanic White, female												
2003.....	54	-	-	-	1	5	7	4	4	4	4	5
2002.....	72	-	2	2	4	3	5	5	7	6	14	6
2001.....	85	-	2	3	-	5	3	5	6	12	8	9
2000.....	72	1	4	-	2	5	6	8	9	9	10	4
1999.....	90	-	2	5	6	7	7	12	6	9	12	3





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
All origins, both sexes												
2003.....	3,306	58	456	126	142	280	287	191	165	217	245	211
2002.....	3,447	63	454	159	162	320	309	227	206	263	263	205
2001.....	3,281	68	458	168	165	322	274	212	162	221	241	214
2000.....	3,482	75	493	201	174	371	275	213	206	250	230	182
1999.....	3,529	68	490	192	177	359	288	212	234	250	225	198
All origins, male												
2003.....	2,632	34	301	90	109	262	263	177	146	186	206	173
2002.....	2,761	40	312	126	118	288	290	201	179	223	216	164
2001.....	2,560	30	291	120	117	295	249	186	142	181	195	182
2000.....	2,735	43	316	137	130	339	244	184	180	212	194	156
1999.....	2,794	39	310	139	136	330	262	193	204	215	187	166
All origins, female												
2003.....	674	24	155	36	33	18	24	14	19	31	39	38
2002.....	686	23	142	33	44	32	19	26	27	40	47	41
2001.....	721	38	167	48	48	27	25	26	20	40	46	32
2000.....	747	32	177	64	44	32	31	29	26	38	36	26
1999.....	735	29	180	53	41	29	26	19	30	35	38	32
Hispanic, both sexes												
2003.....	458	12	82	23	22	46	66	54	31	28	23	19
2002.....	464	18	65	16	13	69	74	45	35	29	26	18
2001.....	421	16	78	27	13	42	59	39	35	36	21	21
2000.....	433	9	75	29	14	73	74	34	28	25	16	11
1999.....	480	16	78	25	21	69	76	39	40	27	18	12

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Hispanic, male												
2003.....	399	7	67	17	16	45	64	51	28	26	21	19
2002.....	409	13	48	15	8	64	73	43	35	24	24	16
2001.....	348	9	53	21	7	42	57	33	32	30	18	18
2000.....	350	6	44	16	6	65	70	32	28	21	15	11
1999.....	405	12	43	18	17	67	73	37	38	24	14	10
Hispanic, female												
2003.....	59	5	15	6	6	1	2	3	3	2	2	-
2002.....	55	5	17	1	5	5	1	2	-	5	2	2
2001.....	73	7	25	6	6	-	2	6	3	6	3	3
2000.....	83	3	31	13	8	8	4	2	-	4	1	-
1999.....	75	4	35	7	4	2	3	2	2	3	4	2
Non-Hispanic 1/, both sexes												
2003.....	2,816	46	374	102	120	231	221	137	134	188	219	190
2002.....	2,952	45	387	141	148	250	235	178	169	232	235	187
2001.....	2,832	51	380	139	152	279	214	173	127	184	219	190
2000.....	3,018	65	417	171	160	297	199	177	178	224	213	169
1999.....	3,029	52	409	167	156	289	211	173	190	222	207	184
Non-Hispanic 1/, male												
2003.....	2,202	27	234	72	93	214	199	126	118	159	183	152
2002.....	2,326	27	262	110	109	223	217	155	143	197	190	148
2001.....	2,188	21	238	97	110	252	191	153	110	150	176	163
2000.....	2,358	37	272	120	124	273	172	150	152	190	178	143
1999.....	2,370	27	265	121	119	262	188	156	162	190	173	154

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Non-Hispanic 1/, female												
2003.....	614	19	140	30	27	17	22	11	16	29	36	38
2002.....	626	18	125	31	39	27	18	23	26	35	45	39
2001.....	644	30	142	42	42	27	23	20	17	34	43	27
2000.....	660	28	145	51	36	24	27	27	26	34	35	26
1999.....	659	25	144	46	37	27	23	17	28	32	34	30
Non-Hispanic White, both sexes												
2003.....	2,152	32	306	64	54	154	152	93	94	148	172	154
2002.....	2,143	32	303	71	63	153	163	118	127	170	190	141
2001.....	2,082	36	321	70	65	165	157	113	89	138	168	146
2000.....	2,235	38	342	96	80	181	135	126	132	180	164	134
1999.....	2,230	30	337	81	61	178	139	113	137	178	171	161
Non-Hispanic White, male												
2003.....	1,656	17	187	45	41	142	137	86	80	121	146	119
2002.....	1,669	18	203	52	44	136	148	101	108	141	153	109
2001.....	1,577	16	205	40	47	149	135	97	74	109	136	126
2000.....	1,737	22	226	68	61	165	113	107	115	151	136	112
1999.....	1,708	14	217	53	46	159	120	99	117	148	142	131
Non-Hispanic White, female												
2003.....	496	15	119	19	13	12	15	7	14	27	26	35
2002.....	474	14	100	19	19	17	15	17	19	29	37	32
2001.....	505	20	116	30	18	16	22	16	15	29	32	20
2000.....	498	16	116	28	19	16	22	19	17	29	28	22
1999.....	522	16	120	28	15	19	19	14	20	30	29	30

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental drowning and submersion (W65-W74)												
Non-Hispanic Black, both sexes												
2003.....	466	13	52	32	55	57	43	28	31	24	24	23
2002.....	604	10	62	57	77	70	56	42	32	43	31	33
2001.....	570	11	47	55	78	89	44	33	29	33	37	34
2000.....	580	25	54	66	68	84	44	34	24	35	42	23
1999.....	624	21	59	76	82	84	53	47	39	31	27	17
Non-Hispanic Black, male												
2003.....	394	9	37	23	43	54	40	25	29	23	22	22
2002.....	502	6	43	46	61	68	54	39	26	38	28	28
2001.....	471	3	28	46	56	82	44	33	28	29	29	27
2000.....	470	14	33	47	54	81	42	29	23	31	36	21
1999.....	521	12	40	60	64	77	51	44	34	29	24	17
Non-Hispanic Black, female												
2003.....	72	4	15	9	12	3	3	3	2	1	2	1
2002.....	102	4	19	11	16	2	2	3	6	5	3	5
2001.....	99	8	19	9	22	7	-	-	1	4	8	7
2000.....	110	11	21	19	14	3	2	5	1	4	6	2
1999.....	103	9	19	16	18	7	2	3	5	2	3	-
Origin not stated 2/, both sexes												
2003.....	32	-	-	1	-	3	-	-	-	1	3	2
2002.....	31	-	2	2	1	1	-	4	2	2	2	-
2001.....	28	1	-	2	-	1	1	-	-	1	1	3
2000.....	31	1	1	1	-	1	2	2	-	1	1	2
1999.....	20	-	3	-	-	1	1	-	4	1	-	2



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
All origins, both sexes												
2003.....	3,369	30	224	137	78	75	124	118	134	167	228	270
2002.....	3,159	36	221	152	101	86	107	112	154	147	214	213
2001.....	3,309	50	225	164	88	76	133	114	133	190	251	213
2000.....	3,377	37	290	182	84	79	113	120	121	188	214	214
1999.....	3,348	41	302	170	92	88	113	142	140	200	221	198
All origins, male												
2003.....	1,948	8	131	72	40	43	78	70	84	115	142	181
2002.....	1,935	21	126	89	62	61	71	75	95	101	150	146
2001.....	1,998	30	142	104	48	45	92	73	88	132	176	151
2000.....	2,025	20	174	109	36	44	77	78	79	133	135	145
1999.....	2,025	21	173	95	52	58	75	93	104	138	162	144
All origins, female												
2003.....	1,421	22	93	65	38	32	46	48	50	52	86	89
2002.....	1,224	15	95	63	39	25	36	37	59	46	64	67
2001.....	1,311	20	83	60	40	31	41	41	45	58	75	62
2000.....	1,352	17	116	73	48	35	36	42	42	55	79	69
1999.....	1,323	20	129	75	40	30	38	49	36	62	59	54
Hispanic, both sexes												
2003.....	203	4	27	18	8	8	7	12	13	13	11	13
2002.....	223	5	36	17	10	11	14	7	10	14	13	12
2001.....	176	9	18	17	6	3	16	11	10	13	15	12
2000.....	178	4	21	15	4	14	7	13	6	8	17	2
1999.....	192	5	41	16	7	12	12	6	11	9	12	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Hispanic, male												
2003.....	128	2	11	9	3	5	4	8	10	9	8	11
2002.....	144	3	18	10	7	9	10	4	5	14	7	9
2001.....	109	5	13	9	2	2	8	9	7	9	10	10
2000.....	105	2	11	9	1	5	6	9	5	6	11	1
1999.....	127	3	23	10	2	7	9	4	8	6	10	9
Hispanic, female												
2003.....	75	2	16	9	5	3	3	4	3	4	3	2
2002.....	79	2	18	7	3	2	4	3	5	-	6	3
2001.....	67	4	5	8	4	1	8	2	3	4	5	2
2000.....	73	2	10	6	3	9	1	4	1	2	6	1
1999.....	65	2	18	6	5	5	3	2	3	3	2	1
Non-Hispanic 1/, both sexes												
2003.....	3,147	26	197	119	70	67	116	106	121	150	217	257
2002.....	2,918	31	184	135	90	75	93	104	143	133	201	200
2001.....	3,106	41	205	145	81	73	117	101	122	177	233	200
2000.....	3,178	31	268	167	79	65	104	107	115	179	196	211
1999.....	3,133	36	260	154	85	73	100	136	128	187	208	186
Non-Hispanic 1/, male												
2003.....	1,809	6	120	63	37	38	73	62	74	106	134	170
2002.....	1,780	18	108	79	54	52	61	70	89	87	143	137
2001.....	1,873	25	127	94	45	43	84	64	81	123	163	140
2000.....	1,907	16	163	100	35	39	69	69	74	127	123	143
1999.....	1,879	18	150	85	50	48	65	89	95	128	151	133

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic 1/, female												
2003.....	1,338	20	77	56	33	29	43	44	47	44	83	87
2002.....	1,138	13	76	56	36	23	32	34	54	46	58	63
2001.....	1,233	16	78	51	36	30	33	37	41	54	70	60
2000.....	1,271	15	105	67	44	26	35	38	41	52	73	68
1999.....	1,254	18	110	69	35	25	35	47	33	59	57	53
Non-Hispanic White, both sexes												
2003.....	2,313	18	118	56	33	46	95	76	94	122	170	194
2002.....	2,055	16	90	61	51	50	63	64	107	96	162	154
2001.....	2,204	26	124	85	51	53	86	60	93	127	170	143
2000.....	2,194	17	139	90	49	46	74	71	80	126	141	153
1999.....	2,172	21	137	80	56	55	70	101	98	129	157	134
Non-Hispanic White, male												
2003.....	1,344	5	74	30	16	29	62	44	58	87	110	127
2002.....	1,271	12	52	38	35	35	43	47	68	61	112	108
2001.....	1,322	14	82	54	28	31	65	44	61	86	121	98
2000.....	1,330	7	88	53	23	27	52	45	54	91	88	105
1999.....	1,325	11	80	42	34	37	46	70	75	88	116	96
Non-Hispanic White, female												
2003.....	969	13	44	26	17	17	33	32	36	35	60	67
2002.....	784	4	38	23	16	15	20	17	39	35	50	46
2001.....	882	12	42	31	23	22	21	16	32	41	49	45
2000.....	864	10	51	37	26	19	22	26	26	35	53	48
1999.....	847	10	57	38	22	18	24	31	23	41	41	38

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental exposure to smoke, fire and flames (X00-X09)												
Non-Hispanic Black, both sexes												
2003.....	767	7	75	60	34	19	19	25	22	26	40	61
2002.....	765	13	88	64	31	21	27	31	30	29	31	40
2001.....	829	14	80	58	27	14	28	36	27	45	58	54
2000.....	915	14	120	74	27	16	26	30	34	44	53	51
1999.....	887	15	110	74	28	16	24	29	23	54	46	46
Non-Hispanic Black, male												
2003.....	430	1	42	31	19	8	10	16	12	18	21	41
2002.....	454	6	53	37	16	14	17	17	18	21	24	25
2001.....	501	10	44	38	15	8	16	16	18	34	39	40
2000.....	539	9	70	46	11	10	15	19	20	30	33	34
1999.....	507	7	63	43	15	9	15	15	14	38	31	32
Non-Hispanic Black, female												
2003.....	337	6	33	29	15	11	9	9	10	8	19	20
2002.....	311	7	35	27	15	7	10	14	12	8	7	15
2001.....	328	4	36	20	12	6	12	20	9	11	19	14
2000.....	376	5	50	28	16	6	11	11	14	14	20	17
1999.....	380	8	47	31	13	7	9	14	9	16	15	14
Origin not stated 2/, both sexes												
2003.....	19	-	-	-	-	-	1	-	-	4	-	-
2002.....	18	-	1	-	1	-	-	1	1	-	-	1
2001.....	27	-	2	2	1	-	-	2	1	-	3	1
2000.....	21	2	1	-	1	-	2	-	-	1	1	1
1999.....	23	-	1	-	-	3	1	-	1	4	1	2



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
All origins, both sexes												
2003.....	19,457	20	49	18	43	520	1,479	1,517	1,918	2,674	3,556	3,351
2002.....	17,550	26	31	15	28	486	1,193	1,310	1,806	2,597	3,410	2,946
2001.....	14,078	15	31	18	32	406	956	1,078	1,429	2,307	2,729	2,266
2000.....	12,757	14	32	17	28	351	809	980	1,400	2,159	2,504	1,992
1999.....	12,186	12	34	12	28	260	704	952	1,403	2,082	2,467	1,926
All origins, male												
2003.....	13,176	9	27	4	27	398	1,164	1,145	1,369	1,804	2,362	2,230
2002.....	12,059	21	14	10	18	366	940	1,003	1,316	1,807	2,315	1,960
2001.....	9,885	8	21	7	16	311	769	832	1,030	1,626	1,896	1,610
2000.....	9,138	11	21	12	15	273	647	779	1,041	1,554	1,830	1,455
1999.....	8,887	11	17	6	19	195	559	757	1,071	1,523	1,826	1,456
All origins, female												
2003.....	6,281	11	22	14	16	122	315	372	549	870	1,194	1,121
2002.....	5,491	5	17	5	10	120	253	307	490	790	1,095	986
2001.....	4,193	7	10	11	16	95	187	246	399	681	833	656
2000.....	3,619	3	11	5	13	78	162	201	359	605	674	537
1999.....	3,299	1	17	6	9	65	145	195	332	559	641	470
Hispanic, both sexes												
2003.....	1,802	4	13	1	4	68	151	196	225	307	303	259
2002.....	1,652	6	4	4	5	57	138	160	204	262	324	243
2001.....	1,324	1	5	2	8	48	96	153	181	251	224	165
2000.....	1,308	2	4	1	3	38	94	133	211	267	226	158
1999.....	1,502	4	3	2	4	41	105	155	228	311	274	191

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidental poisoning and exposure to noxious substances (X40-X49)</b>												
<b>Hispanic, male</b>												
2003.....	1,428	2	5	-	3	55	124	161	186	251	232	206
2002.....	1,324	5	-	2	2	44	117	141	171	206	260	184
2001.....	1,061	1	3	1	1	35	76	131	143	205	180	135
2000.....	1,090	2	2	1	1	31	82	112	183	214	201	139
1999.....	1,271	4	3	1	3	28	95	133	199	262	231	165
<b>Hispanic, female</b>												
2003.....	374	2	8	1	1	13	27	35	39	56	71	53
2002.....	328	1	4	2	3	13	21	19	33	56	64	59
2001.....	263	-	2	1	7	13	20	22	38	46	44	30
2000.....	218	-	2	-	2	7	12	21	28	53	25	19
1999.....	231	-	-	1	1	13	10	22	29	49	43	26
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	17,537	16	36	17	39	450	1,326	1,311	1,683	2,353	3,237	3,069
2002.....	15,789	20	27	11	23	428	1,048	1,144	1,591	2,318	3,064	2,687
2001.....	12,631	14	26	16	24	357	853	921	1,236	2,034	2,485	2,079
2000.....	11,311	12	27	14	24	312	710	841	1,177	1,878	2,253	1,805
1999.....	10,542	8	31	10	23	219	595	790	1,165	1,753	2,154	1,715
<b>Non-Hispanic 1/, male</b>												
2003.....	11,655	7	22	4	24	342	1,039	976	1,175	1,541	2,118	2,009
2002.....	10,649	16	14	8	16	322	818	857	1,136	1,587	2,039	1,763
2001.....	8,720	7	18	6	15	275	687	697	877	1,402	1,701	1,458
2000.....	7,944	9	18	11	13	242	560	663	849	1,328	1,610	1,297
1999.....	7,510	7	14	5	15	167	460	618	864	1,248	1,569	1,275

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidental poisoning and exposure to noxious substances (X40-X49)</b>												
<b>Non-Hispanic 1/, female</b>												
2003.....	5,882	9	14	13	15	108	287	335	508	812	1,119	1,060
2002.....	5,140	4	13	3	7	106	230	287	455	731	1,025	924
2001.....	3,911	7	8	10	9	82	166	224	359	632	784	621
2000.....	3,367	3	9	3	11	70	150	178	328	550	643	508
1999.....	3,032	1	17	5	8	52	135	172	301	505	585	440
<b>Non-Hispanic White, both sexes</b>												
2003.....	14,829	8	19	11	20	420	1,239	1,168	1,443	1,964	2,744	2,538
2002.....	13,165	7	15	8	20	395	962	1,004	1,371	1,932	2,564	2,156
2001.....	10,269	8	16	12	18	314	764	797	1,039	1,678	2,020	1,620
2000.....	9,124	8	15	7	17	281	635	713	970	1,540	1,791	1,390
1999.....	8,311	6	17	3	18	190	504	663	918	1,416	1,680	1,290
<b>Non-Hispanic White, male</b>												
2003.....	9,855	6	11	3	14	323	969	878	1,014	1,288	1,810	1,654
2002.....	8,849	7	6	6	15	299	749	750	987	1,331	1,714	1,418
2001.....	7,111	4	12	5	12	251	613	601	757	1,170	1,409	1,128
2000.....	6,413	6	9	6	10	220	516	563	709	1,112	1,287	990
1999.....	5,921	5	6	1	12	145	391	523	694	1,023	1,242	952
<b>Non-Hispanic White, female</b>												
2003.....	4,974	2	8	8	6	97	270	290	429	676	934	884
2002.....	4,316	-	9	2	5	96	213	254	384	601	850	738
2001.....	3,158	4	4	7	6	63	151	196	282	508	611	492
2000.....	2,711	2	6	1	7	61	119	150	261	428	504	400
1999.....	2,390	1	11	2	6	45	113	140	224	393	438	338

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Accidental poisoning and exposure to noxious substances (X40-X49)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	2,354	7	15	4	13	17	69	103	200	330	440	490
2002.....	2,326	12	11	3	3	19	67	113	186	336	450	489
2001.....	2,122	6	8	4	2	26	67	103	170	316	433	431
2000.....	1,985	4	11	6	2	19	57	111	183	309	423	385
1999.....	2,008	1	12	5	2	18	75	112	216	295	434	404
<b>Non-Hispanic Black, male</b>												
2003.....	1,579	1	9	-	7	10	54	70	134	219	278	323
2002.....	1,601	8	8	2	1	14	52	86	127	226	291	320
2001.....	1,452	3	5	1	1	14	53	81	105	205	272	314
2000.....	1,400	3	8	5	1	13	36	87	122	200	296	286
1999.....	1,450	1	7	3	2	15	61	85	146	199	304	308
<b>Non-Hispanic Black, female</b>												
2003.....	775	6	6	4	6	7	15	33	66	111	162	167
2002.....	725	4	3	1	2	5	15	27	59	110	159	169
2001.....	670	3	3	3	1	12	14	22	65	111	161	117
2000.....	585	1	3	1	1	6	21	24	61	109	127	99
1999.....	558	-	5	2	-	3	14	27	70	96	130	96
<b>Origin not stated 2/, both sexes</b>												
2003.....	118	-	-	-	-	2	2	10	10	14	16	23
2002.....	109	-	-	-	-	1	7	6	11	17	22	16
2001.....	123	-	-	-	-	1	7	4	12	22	20	22
2000.....	138	-	1	2	1	1	5	6	12	14	25	29
1999.....	142	-	-	-	1	-	4	7	10	18	39	20

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Accidental poisoning and exposure to noxious substances (X40-X49)												
Origin not stated 2/, male												
2003.....	93	-	-	-	-	1	1	8	8	12	12	15
2002.....	86	-	-	-	-	-	5	5	9	14	16	13
2001.....	104	-	-	-	-	1	6	4	10	19	15	17
2000.....	104	-	1	-	1	-	5	4	9	12	19	19
1999.....	106	-	-	-	1	-	4	6	8	13	26	16
Origin not stated 2/, female												
2003.....	25	-	-	-	-	1	1	2	2	2	4	8
2002.....	23	-	-	-	-	1	2	1	2	3	6	3
2001.....	19	-	-	-	-	-	1	-	2	3	5	5
2000.....	34	-	-	2	-	1	-	2	3	2	6	10
1999.....	36	-	-	-	-	-	-	1	2	5	13	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
All origins, both sexes												
2003.....	17,115	675	278	94	121	224	421	390	423	546	785	821
2002.....	17,201	677	247	113	160	273	365	377	443	577	764	745
2001.....	17,760	669	275	113	142	251	392	347	448	648	738	816
2000.....	17,437	572	266	115	152	273	375	416	479	672	774	787
1999.....	18,388	523	293	144	172	291	371	421	466	622	830	755
All origins, male												
2003.....	9,919	398	177	63	97	191	353	327	358	432	629	640
2002.....	9,839	389	156	71	114	225	320	311	381	468	613	609
2001.....	10,044	390	162	72	109	196	339	294	388	529	599	647
2000.....	9,960	327	162	72	120	223	337	362	405	577	628	646
1999.....	10,371	314	183	93	128	238	327	361	387	513	668	584
All origins, female												
2003.....	7,196	277	101	31	24	33	68	63	65	114	156	181
2002.....	7,362	288	91	42	46	48	45	66	62	109	151	136
2001.....	7,716	279	113	41	33	55	53	53	60	119	139	169
2000.....	7,477	245	104	43	32	50	38	54	74	95	146	141
1999.....	8,017	209	110	51	44	53	44	60	79	109	162	171
Hispanic, both sexes												
2003.....	999	71	53	21	8	39	76	80	70	59	69	63
2002.....	954	68	42	17	19	37	63	67	72	69	75	49
2001.....	985	78	43	9	14	34	64	55	67	77	69	61
2000.....	942	58	41	13	16	34	74	63	77	77	74	49
1999.....	878	65	53	18	13	35	63	54	47	68	52	49

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Hispanic, male												
2003.....	749	40	32	13	8	34	74	74	64	54	60	54
2002.....	726	43	28	11	10	34	58	64	69	62	65	45
2001.....	704	40	28	2	10	31	62	50	60	70	60	51
2000.....	695	31	23	6	8	31	74	58	71	73	66	43
1999.....	620	35	37	12	7	28	60	51	39	60	41	44
Hispanic, female												
2003.....	250	31	21	8	-	5	2	6	6	5	9	9
2002.....	228	25	14	6	9	3	5	3	3	7	10	4
2001.....	281	38	15	7	4	3	2	5	7	7	9	10
2000.....	247	27	18	7	8	3	-	5	6	4	8	6
1999.....	258	30	16	6	6	7	3	3	8	8	11	5
Non-Hispanic 1/, both sexes												
2003.....	16,055	602	225	73	113	184	342	308	351	486	709	753
2002.....	16,183	606	205	95	140	233	301	310	367	505	688	692
2001.....	16,695	588	231	104	126	217	325	290	379	566	665	751
2000.....	16,408	513	224	102	136	238	300	351	397	589	694	730
1999.....	17,430	457	239	126	158	254	308	364	417	553	778	700
Non-Hispanic 1/, male												
2003.....	9,122	356	145	50	89	156	276	252	292	377	562	581
2002.....	9,070	344	128	59	104	189	261	247	308	403	547	562
2001.....	9,280	347	133	70	97	165	274	242	326	454	536	593
2000.....	9,207	295	139	66	112	192	262	302	329	499	558	595
1999.....	9,694	279	146	81	120	209	267	308	347	452	627	535

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Non-Hispanic 1/, female												
2003.....	6,933	246	80	23	24	28	66	56	59	109	147	172
2002.....	7,113	262	77	36	36	44	40	63	59	102	141	130
2001.....	7,415	241	98	34	29	52	51	48	53	112	129	158
2000.....	7,201	218	85	36	24	46	38	49	68	90	136	135
1999.....	7,736	178	93	45	38	45	41	56	70	101	151	165
Non-Hispanic White, both sexes												
2003.....	13,791	356	158	54	89	150	274	244	287	389	580	599
2002.....	13,865	339	136	66	106	190	256	263	302	402	567	564
2001.....	14,326	348	160	85	97	183	272	228	300	468	526	594
2000.....	14,083	310	156	76	109	186	239	283	321	479	560	568
1999.....	14,930	276	163	96	129	213	255	293	329	468	635	533
Non-Hispanic White, male												
2003.....	7,709	211	104	37	71	127	231	206	238	300	467	463
2002.....	7,669	203	83	44	76	158	224	212	254	318	468	473
2001.....	7,832	206	93	54	75	139	231	193	256	386	428	477
2000.....	7,703	186	99	51	90	153	208	243	272	409	451	458
1999.....	8,122	167	96	67	96	174	221	255	277	388	508	405
Non-Hispanic White, female												
2003.....	6,082	145	54	17	18	23	43	38	49	89	113	136
2002.....	6,196	136	53	22	30	32	32	51	48	84	99	91
2001.....	6,494	142	67	31	22	44	41	35	44	82	98	117
2000.....	6,380	124	57	25	19	33	31	40	49	70	109	110
1999.....	6,808	109	67	29	33	39	34	38	52	80	127	128

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Non-Hispanic Black, both sexes												
2003.....	1,858	226	52	16	15	22	57	51	44	78	107	122
2002.....	1,951	250	59	23	32	33	35	38	52	85	94	109
2001.....	1,968	217	61	13	25	24	38	48	53	74	110	133
2000.....	1,968	183	55	24	24	41	49	54	64	90	108	132
1999.....	2,183	168	61	23	24	31	43	59	73	65	128	149
Non-Hispanic Black, male												
2003.....	1,147	134	36	10	12	19	37	37	37	62	76	95
2002.....	1,162	132	40	11	27	24	28	28	43	69	57	75
2001.....	1,180	131	35	12	19	18	29	35	46	50	87	98
2000.....	1,256	96	32	14	20	29	44	45	49	74	85	110
1999.....	1,365	101	42	12	20	28	39	44	58	48	105	116
Non-Hispanic Black, female												
2003.....	711	92	16	6	3	3	20	14	7	16	31	27
2002.....	789	118	19	12	5	9	7	10	9	16	37	34
2001.....	788	86	26	1	6	6	9	13	7	24	23	35
2000.....	712	87	23	10	4	12	5	9	15	16	23	22
1999.....	818	67	19	11	4	3	4	15	15	17	23	33
Origin not stated 2/, both sexes												
2003.....	61	2	-	-	-	1	3	2	2	1	7	5
2002.....	64	3	-	1	1	3	1	-	4	3	1	4
2001.....	80	3	1	-	2	-	3	2	2	5	4	4
2000.....	87	1	1	-	-	1	1	2	5	6	6	8
1999.....	80	1	1	-	1	2	-	3	2	1	-	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Origin not stated 2/, male

2003.....	48	2	-	-	-	1	3	1	2	1	7	5
2002.....	43	2	-	1	-	2	1	-	4	3	1	2
2001.....	60	3	1	-	2	-	3	2	2	5	3	3
2000.....	58	1	-	-	-	-	1	2	5	5	4	8
1999.....	57	-	-	-	1	1	-	2	1	1	-	5

Origin not stated 2/, female

2003.....	13	-	-	-	-	-	-	1	-	-	-	-
2002.....	21	1	-	-	1	1	-	-	-	-	-	2
2001.....	20	-	-	-	-	-	-	-	-	-	1	1
2000.....	29	-	1	-	-	1	-	-	-	1	2	-
1999.....	23	1	1	-	-	1	-	1	1	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All origins, both sexes												
2003.....	31,484	...	...	6	244	1,487	2,501	2,276	2,789	2,958	3,644	3,485
2002.....	31,655	...	...	4	260	1,513	2,497	2,423	2,623	3,141	3,710	3,473
2001 3/.....	30,622	...	...	7	272	1,611	2,360	2,389	2,681	3,176	3,459	3,260
2000.....	29,350	...	...	7	300	1,621	2,373	2,341	2,451	3,112	3,450	2,987
1999.....	29,199	...	...	2	242	1,615	2,286	2,398	2,708	3,286	3,180	2,817
All origins, male												
2003.....	25,203	...	...	6	188	1,222	2,159	1,901	2,255	2,347	2,791	2,616
2002.....	25,409	...	...	3	196	1,280	2,152	2,003	2,132	2,484	2,821	2,608
2001 3/.....	24,672	...	...	7	207	1,345	2,064	2,015	2,184	2,518	2,662	2,484
2000.....	23,618	...	...	6	238	1,351	2,073	1,956	1,982	2,457	2,657	2,307
1999.....	23,458	...	...	2	192	1,347	1,979	2,005	2,189	2,590	2,429	2,152
All origins, female												
2003.....	6,281	...	...	-	56	265	342	375	534	611	853	869
2002.....	6,246	...	...	1	64	233	345	420	491	657	889	865
2001 3/.....	5,950	...	...	-	65	266	296	374	497	658	797	776
2000.....	5,732	...	...	1	62	270	300	385	469	655	793	680
1999.....	5,741	...	...	-	50	268	307	393	519	696	751	665
Hispanic, both sexes												
2003.....	2,007	...	...	-	43	192	299	251	248	203	199	161
2002.....	1,954	...	...	2	35	181	274	248	236	222	185	149
2001 3/.....	1,850	...	...	-	37	166	248	259	231	224	173	130
2000.....	1,787	...	...	2	27	178	269	227	193	194	191	141
1999.....	1,695	...	...	-	30	175	216	220	203	228	141	121

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Hispanic, male												
2003.....	1,711	...	...	-	31	155	266	220	214	173	168	130
2002.....	1,651	...	...	1	20	151	237	222	204	188	152	115
2001 3/.....	1,576	...	...	-	26	129	214	223	196	200	150	106
2000.....	1,525	...	...	2	20	143	244	203	167	158	162	112
1999.....	1,430	...	...	-	21	146	192	187	177	188	119	104
Hispanic, female												
2003.....	296	...	...	-	12	37	33	31	34	30	31	31
2002.....	303	...	...	1	15	30	37	26	32	34	33	34
2001 3/.....	274	...	...	-	11	37	34	36	35	24	23	24
2000.....	262	...	...	-	7	35	25	24	26	36	29	29
1999.....	265	...	...	-	9	29	24	33	26	40	22	17
Non-Hispanic 1/, both sexes												
2003.....	29,367	...	...	6	200	1,292	2,191	2,019	2,535	2,741	3,430	3,309
2002.....	29,543	...	...	2	224	1,322	2,214	2,165	2,373	2,902	3,511	3,309
2001 3/.....	28,647	...	...	7	235	1,440	2,105	2,119	2,441	2,943	3,270	3,118
2000.....	27,426	...	...	5	271	1,436	2,093	2,101	2,249	2,904	3,241	2,829
1999.....	27,379	...	...	1	211	1,434	2,066	2,167	2,498	3,048	3,020	2,683
Non-Hispanic 1/, male												
2003.....	23,406	...	...	6	156	1,065	1,885	1,677	2,035	2,162	2,613	2,475
2002.....	23,623	...	...	2	175	1,120	1,906	1,772	1,917	2,282	2,659	2,480
2001 3/.....	22,983	...	...	7	181	1,212	1,843	1,782	1,980	2,311	2,498	2,367
2000.....	21,981	...	...	4	217	1,203	1,820	1,744	1,809	2,288	2,479	2,180
1999.....	21,929	...	...	1	170	1,198	1,785	1,809	2,007	2,395	2,294	2,038

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic 1/, female												
2003.....	5,961	...	...	-	44	227	306	342	500	579	817	834
2002.....	5,920	...	...	-	49	202	308	393	456	620	852	829
2001 3/.....	5,664	...	...	-	54	228	262	337	461	632	772	751
2000.....	5,445	...	...	1	54	233	273	357	440	616	762	649
1999.....	5,450	...	...	-	41	236	281	358	491	653	726	645
Non-Hispanic White, both sexes												
2003.....	26,434	...	...	5	146	1,088	1,744	1,647	2,164	2,416	3,147	3,082
2002.....	26,691	...	...	1	176	1,124	1,827	1,771	2,038	2,570	3,225	3,084
2001 3/.....	25,813	...	...	5	178	1,217	1,709	1,755	2,116	2,621	2,983	2,897
2000.....	24,616	...	...	5	220	1,175	1,693	1,709	1,949	2,599	2,941	2,611
1999.....	24,534	...	...	1	173	1,169	1,661	1,776	2,151	2,699	2,770	2,468
Non-Hispanic White, male												
2003.....	21,085	...	...	5	117	896	1,516	1,363	1,736	1,898	2,398	2,309
2002.....	21,323	...	...	1	137	954	1,573	1,427	1,638	2,013	2,449	2,296
2001 3/.....	20,710	...	...	5	139	1,030	1,492	1,479	1,711	2,046	2,283	2,215
2000.....	19,711	...	...	4	175	987	1,480	1,424	1,565	2,038	2,249	2,008
1999.....	19,620	...	...	1	140	978	1,438	1,478	1,727	2,098	2,112	1,876
Non-Hispanic White, female												
2003.....	5,349	...	...	-	29	192	228	284	428	518	749	773
2002.....	5,368	...	...	-	39	170	254	344	400	557	776	788
2001 3/.....	5,103	...	...	-	39	187	217	276	405	575	700	682
2000.....	4,905	...	...	1	45	188	213	285	384	561	692	603
1999.....	4,914	...	...	-	33	191	223	298	424	601	658	592

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Non-Hispanic Black, both sexes												
2003.....	1,918	...	...	1	42	119	319	249	249	209	200	140
2002.....	1,896	...	...	1	34	122	270	271	226	229	203	149
2001 3/.....	1,904	...	...	2	38	134	291	243	229	227	191	152
2000.....	1,923	...	...	-	39	170	301	262	207	205	222	150
1999.....	1,924	...	...	-	29	176	287	258	239	244	186	144
Non-Hispanic Black, male												
2003.....	1,566	...	...	1	33	105	272	220	204	171	153	108
2002.....	1,595	...	...	1	28	107	235	244	197	191	154	129
2001 3/.....	1,578	...	...	2	29	115	273	212	194	188	141	109
2000.....	1,601	...	...	-	34	148	259	222	175	172	174	123
1999.....	1,630	...	...	-	22	152	256	229	203	215	138	114
Non-Hispanic Black, female												
2003.....	352	...	...	-	9	14	47	29	45	38	47	32
2002.....	301	...	...	-	6	15	35	27	29	38	49	20
2001 3/.....	326	...	...	-	9	19	18	31	35	39	50	43
2000.....	322	...	...	-	5	22	42	40	32	33	48	27
1999.....	294	...	...	-	7	24	31	29	36	29	48	30
Origin not stated 2/, both sexes												
2003.....	110	...	...	-	1	3	11	6	6	14	15	15
2002.....	158	...	...	-	1	10	9	10	14	17	14	15
2001 3/.....	125	...	...	-	-	5	7	11	9	9	16	12
2000.....	137	...	...	-	2	7	11	13	9	14	18	17
1999.....	125	...	...	1	1	6	4	11	7	10	19	13

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Origin not stated 2/, male

2003.....	86	...	...	-	1	2	8	4	6	12	10	11
2002.....	135	...	...	-	1	9	9	9	11	14	10	13
2001 3/.....	113	...	...	-	-	4	7	10	8	7	14	11
2000.....	112	...	...	-	1	5	9	9	6	11	16	15
1999.....	99	...	...	1	1	3	2	9	5	7	16	10

Origin not stated 2/, female

2003.....	24	...	...	-	-	1	3	2	-	2	5	4
2002.....	23	...	...	-	-	1	-	1	3	3	4	2
2001 3/.....	12	...	...	-	-	1	-	1	1	2	2	1
2000.....	25	...	...	-	1	2	2	4	3	3	2	2
1999.....	26	...	...	-	-	3	2	2	2	3	3	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All origins, both sexes												
2003.....	16,907	...	...	1	73	736	1,339	1,114	1,267	1,280	1,647	1,705
2002.....	17,108	...	...	-	86	742	1,346	1,217	1,182	1,450	1,707	1,652
2001.....	16,869	...	...	-	90	838	1,292	1,263	1,301	1,429	1,601	1,552
2000.....	16,586	...	...	-	110	897	1,370	1,223	1,230	1,521	1,670	1,491
1999.....	16,599	...	...	-	103	975	1,340	1,292	1,329	1,617	1,480	1,462
All origins, male												
2003.....	14,827	...	...	1	62	659	1,220	990	1,109	1,092	1,378	1,419
2002.....	15,045	...	...	-	68	668	1,229	1,068	1,010	1,234	1,438	1,394
2001.....	14,758	...	...	-	69	743	1,192	1,122	1,124	1,200	1,351	1,309
2000.....	14,454	...	...	-	90	793	1,250	1,084	1,039	1,255	1,402	1,263
1999.....	14,479	...	...	-	80	867	1,216	1,137	1,133	1,349	1,237	1,221
All origins, female												
2003.....	2,080	...	...	-	11	77	119	124	158	188	269	286
2002.....	2,063	...	...	-	18	74	117	149	172	216	269	258
2001.....	2,111	...	...	-	21	95	100	141	177	229	250	243
2000.....	2,132	...	...	-	20	104	120	139	191	266	268	228
1999.....	2,120	...	...	-	23	108	124	155	196	268	243	241
Hispanic, both sexes												
2003.....	835	...	...	-	9	83	145	121	87	76	69	67
2002.....	834	...	...	-	10	88	136	112	86	80	65	59
2001.....	798	...	...	-	5	67	126	129	95	75	60	54
2000.....	813	...	...	-	7	88	136	104	77	65	79	72
1999.....	794	...	...	-	13	94	125	104	78	91	57	64

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Hispanic, male												
2003.....	774	...	...	-	7	73	137	113	81	68	65	58
2002.....	763	...	...	-	5	81	127	101	82	72	56	51
2001.....	731	...	...	-	4	58	114	116	87	70	56	48
2000.....	737	...	...	-	7	78	127	96	69	55	71	61
1999.....	705	...	...	-	7	81	115	92	72	76	51	55
Hispanic, female												
2003.....	61	...	...	-	2	10	8	8	6	8	4	9
2002.....	71	...	...	-	5	7	9	11	4	8	9	8
2001.....	67	...	...	-	1	9	12	13	8	5	4	6
2000.....	76	...	...	-	-	10	9	8	8	10	8	11
1999.....	89	...	...	-	6	13	10	12	6	15	6	9
Non-Hispanic 1/, both sexes												
2003.....	16,035	...	...	1	64	652	1,191	992	1,179	1,201	1,571	1,633
2002.....	16,213	...	...	-	76	649	1,206	1,102	1,093	1,363	1,639	1,585
2001.....	16,020	...	...	-	85	770	1,163	1,130	1,205	1,351	1,536	1,495
2000.....	15,717	...	...	-	103	805	1,230	1,112	1,152	1,452	1,584	1,412
1999.....	15,755	...	...	-	90	879	1,213	1,184	1,247	1,520	1,417	1,395
Non-Hispanic 1/, male												
2003.....	14,025	...	...	1	55	585	1,082	876	1,027	1,021	1,309	1,358
2002.....	14,226	...	...	-	63	582	1,098	964	926	1,155	1,381	1,336
2001.....	13,978	...	...	-	65	684	1,075	1,002	1,037	1,127	1,290	1,259
2000.....	13,664	...	...	-	83	712	1,119	982	969	1,196	1,324	1,195
1999.....	13,732	...	...	-	73	784	1,100	1,043	1,058	1,268	1,180	1,164

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic 1/, female												
2003.....	2,010	...	...	-	9	67	109	116	152	180	262	275
2002.....	1,987	...	...	-	13	67	108	138	167	208	258	249
2001.....	2,042	...	...	-	20	86	88	128	168	224	246	236
2000.....	2,053	...	...	-	20	93	111	130	183	256	260	217
1999.....	2,023	...	...	-	17	95	113	141	189	252	237	231
Non-Hispanic White, both sexes												
2003.....	14,737	...	...	1	55	556	947	806	1,007	1,088	1,476	1,556
2002.....	14,865	...	...	-	63	540	981	898	955	1,219	1,530	1,493
2001.....	14,648	...	...	-	69	650	915	942	1,059	1,220	1,409	1,413
2000.....	14,333	...	...	-	96	651	1,004	904	1,012	1,316	1,464	1,315
1999.....	14,316	...	...	-	81	726	958	977	1,094	1,357	1,319	1,295
Non-Hispanic White, male												
2003.....	12,865	...	...	1	48	497	864	702	871	915	1,230	1,291
2002.....	13,014	...	...	-	53	485	890	767	806	1,028	1,291	1,255
2001.....	12,771	...	...	-	55	573	841	833	906	1,011	1,188	1,196
2000.....	12,448	...	...	-	79	574	915	793	855	1,075	1,221	1,114
1999.....	12,441	...	...	-	65	645	869	854	922	1,120	1,093	1,084
Non-Hispanic White, female												
2003.....	1,872	...	...	-	7	59	83	104	136	173	246	265
2002.....	1,851	...	...	-	10	55	91	131	149	191	239	238
2001.....	1,877	...	...	-	14	77	74	109	153	209	221	217
2000.....	1,885	...	...	-	17	77	89	111	157	241	243	201
1999.....	1,875	...	...	-	16	81	89	123	172	237	226	211

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Non-Hispanic Black, both sexes												
2003.....	993	...	...	-	7	65	199	147	132	80	70	50
2002.....	1,041	...	...	-	9	77	173	162	110	114	85	73
2001.....	1,069	...	...	-	12	88	205	147	111	107	91	62
2000.....	1,073	...	...	-	4	117	187	159	109	100	93	73
1999.....	1,112	...	...	-	5	118	205	153	115	123	74	76
Non-Hispanic Black, male												
2003.....	892	...	...	-	5	62	181	141	119	74	57	46
2002.....	945	...	...	-	8	68	161	157	101	100	70	67
2001.....	949	...	...	-	9	80	197	131	102	96	73	48
2000.....	949	...	...	-	4	109	168	144	93	88	80	62
1999.....	1,007	...	...	-	4	107	187	142	105	112	66	63
Non-Hispanic Black, female												
2003.....	101	...	...	-	2	3	18	6	13	6	13	4
2002.....	96	...	...	-	1	9	12	5	9	14	15	6
2001.....	120	...	...	-	3	8	8	16	9	11	18	14
2000.....	124	...	...	-	-	8	19	15	16	12	13	11
1999.....	105	...	...	-	1	11	18	11	10	11	8	13
Origin not stated 2/, both sexes												
2003.....	37	...	...	-	-	1	3	1	1	3	7	5
2002.....	61	...	...	-	-	5	4	3	3	7	3	8
2001.....	51	...	...	-	-	1	3	4	1	3	5	3
2000.....	56	...	...	-	-	4	4	7	1	4	7	7
1999.....	50	...	...	-	-	2	2	4	4	6	6	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Origin not stated 2/, male

2003.....	28	...	...	-	-	1	1	1	1	3	4	3
2002.....	56	...	...	-	-	5	4	3	2	7	1	7
2001.....	49	...	...	-	-	1	3	4	-	3	5	2
2000.....	53	...	...	-	-	3	4	6	1	4	7	7
1999.....	42	...	...	-	-	2	1	2	3	5	6	2

Origin not stated 2/, female

2003.....	9	...	...	-	-	-	2	-	-	-	3	2
2002.....	5	...	...	-	-	-	-	-	1	-	2	1
2001.....	2	...	...	-	-	-	-	-	1	-	-	1
2000.....	3	...	...	-	-	1	-	1	-	-	-	-
1999.....	8	...	...	-	-	-	1	2	1	1	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
All origins, both sexes												
2003.....	14,577	...	...	5	171	751	1,162	1,162	1,522	1,678	1,997	1,780
2002.....	14,547	...	...	4	174	771	1,151	1,206	1,441	1,691	2,003	1,821
2001 3/.....	13,753	...	...	7	182	773	1,068	1,126	1,380	1,747	1,858	1,708
2000.....	12,764	...	...	7	190	724	1,003	1,118	1,221	1,591	1,780	1,496
1999.....	12,600	...	...	2	139	640	946	1,106	1,379	1,669	1,700	1,355
All origins, male												
2003.....	10,376	...	...	5	126	563	939	911	1,146	1,255	1,413	1,197
2002.....	10,364	...	...	3	128	612	923	935	1,122	1,250	1,383	1,214
2001 3/.....	9,914	...	...	7	138	602	872	893	1,060	1,318	1,311	1,175
2000.....	9,164	...	...	6	148	558	823	872	943	1,202	1,255	1,044
1999.....	8,979	...	...	2	112	480	763	868	1,056	1,241	1,192	931
All origins, female												
2003.....	4,201	...	...	-	45	188	223	251	376	423	584	583
2002.....	4,183	...	...	1	46	159	228	271	319	441	620	607
2001 3/.....	3,839	...	...	-	44	171	196	233	320	429	547	533
2000.....	3,600	...	...	1	42	166	180	246	278	389	525	452
1999.....	3,621	...	...	-	27	160	183	238	323	428	508	424
Hispanic, both sexes												
2003.....	1,172	...	...	-	34	109	154	130	161	127	130	94
2002.....	1,120	...	...	2	25	93	138	136	150	142	120	90
2001 3/.....	1,052	...	...	-	32	99	122	130	136	149	113	76
2000.....	974	...	...	2	20	90	133	123	116	129	112	69
1999.....	901	...	...	-	17	81	91	116	125	137	84	57

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Hispanic, male												
2003.....	937	...	...	-	24	82	129	107	133	105	103	72
2002.....	888	...	...	1	15	70	110	121	122	116	96	64
2001 3/.....	845	...	...	-	22	71	100	107	109	130	94	58
2000.....	788	...	...	2	13	65	117	107	98	103	91	51
1999.....	725	...	...	-	14	65	77	95	105	112	68	49
Hispanic, female												
2003.....	235	...	...	-	10	27	25	23	28	22	27	22
2002.....	232	...	...	1	10	23	28	15	28	26	24	26
2001 3/.....	207	...	...	-	10	28	22	23	27	19	19	18
2000.....	186	...	...	-	7	25	16	16	18	26	21	18
1999.....	176	...	...	-	3	16	14	21	20	25	16	8
Non-Hispanic 1/, both sexes												
2003.....	13,332	...	...	5	136	640	1,000	1,027	1,356	1,540	1,859	1,676
2002.....	13,330	...	...	2	148	673	1,008	1,063	1,280	1,539	1,872	1,724
2001 3/.....	12,627	...	...	7	150	670	942	989	1,236	1,592	1,734	1,623
2000.....	11,709	...	...	5	168	631	863	989	1,097	1,452	1,657	1,417
1999.....	11,624	...	...	1	121	555	853	983	1,251	1,528	1,603	1,288
Non-Hispanic 1/, male												
2003.....	9,381	...	...	5	101	480	803	801	1,008	1,141	1,304	1,117
2002.....	9,397	...	...	2	112	538	808	808	991	1,127	1,278	1,144
2001 3/.....	9,005	...	...	7	116	528	768	780	943	1,184	1,208	1,108
2000.....	8,317	...	...	4	134	491	701	762	840	1,092	1,155	985
1999.....	8,197	...	...	1	97	414	685	766	949	1,127	1,114	874

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Non-Hispanic 1/, female												
2003.....	3,951	...	...	-	35	160	197	226	348	399	555	559
2002.....	3,933	...	...	-	36	135	200	255	289	412	594	580
2001 3/.....	3,622	...	...	-	34	142	174	209	293	408	526	515
2000.....	3,392	...	...	1	34	140	162	227	257	360	502	432
1999.....	3,427	...	...	-	24	141	168	217	302	401	489	414
Non-Hispanic White, both sexes												
2003.....	11,697	...	...	4	91	532	797	841	1,157	1,328	1,671	1,526
2002.....	11,826	...	...	1	113	584	846	873	1,083	1,351	1,695	1,591
2001 3/.....	11,165	...	...	5	109	567	794	813	1,057	1,401	1,574	1,484
2000.....	10,283	...	...	5	124	524	689	805	937	1,283	1,477	1,296
1999.....	10,218	...	...	1	92	443	703	799	1,057	1,342	1,451	1,173
Non-Hispanic White, male												
2003.....	8,220	...	...	4	69	399	652	661	865	983	1,168	1,018
2002.....	8,309	...	...	1	84	469	683	660	832	985	1,158	1,041
2001 3/.....	7,939	...	...	5	84	457	651	646	805	1,035	1,095	1,019
2000.....	7,263	...	...	4	96	413	565	631	710	963	1,028	894
1999.....	7,179	...	...	1	75	333	569	624	805	978	1,019	792
Non-Hispanic White, female												
2003.....	3,477	...	...	-	22	133	145	180	292	345	503	508
2002.....	3,517	...	...	-	29	115	163	213	251	366	537	550
2001 3/.....	3,226	...	...	-	25	110	143	167	252	366	479	465
2000.....	3,020	...	...	1	28	111	124	174	227	320	449	402
1999.....	3,039	...	...	-	17	110	134	175	252	364	432	381

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Non-Hispanic Black, both sexes												
2003.....	925	...	...	1	35	54	120	102	117	129	130	90
2002.....	855	...	...	1	25	45	97	109	116	115	118	76
2001 3/.....	835	...	...	2	26	46	86	96	118	120	100	90
2000.....	850	...	...	-	35	53	114	103	98	105	129	77
1999.....	812	...	...	-	24	58	82	105	124	121	112	68
Non-Hispanic Black, male												
2003.....	674	...	...	1	28	43	91	79	85	97	96	62
2002.....	650	...	...	1	20	39	74	87	96	91	84	62
2001 3/.....	629	...	...	2	20	35	76	81	92	92	68	61
2000.....	652	...	...	-	30	39	91	78	82	84	94	61
1999.....	623	...	...	-	18	45	69	87	98	103	72	51
Non-Hispanic Black, female												
2003.....	251	...	...	-	7	11	29	23	32	32	34	28
2002.....	205	...	...	-	5	6	23	22	20	24	34	14
2001 3/.....	206	...	...	-	6	11	10	15	26	28	32	29
2000.....	198	...	...	-	5	14	23	25	16	21	35	16
1999.....	189	...	...	-	6	13	13	18	26	18	40	17
Origin not stated 2/, both sexes												
2003.....	73	...	...	-	1	2	8	5	5	11	8	10
2002.....	97	...	...	-	1	5	5	7	11	10	11	7
2001 3/.....	74	...	...	-	-	4	4	7	8	6	11	9
2000.....	81	...	...	-	2	3	7	6	8	10	11	10
1999.....	75	...	...	1	1	4	2	7	3	4	13	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Origin not stated 2/, male

2003.....	58	...	...	-	1	1	7	3	5	9	6	8
2002.....	79	...	...	-	1	4	5	6	9	7	9	6
2001 3/.....	64	...	...	-	-	3	4	6	8	4	9	9
2000.....	59	...	...	-	1	2	5	3	5	7	9	8
1999.....	57	...	...	1	1	1	1	7	2	2	10	8

Origin not stated 2/, female

2003.....	15	...	...	-	-	1	1	2	-	2	2	2
2002.....	18	...	...	-	-	1	-	1	2	3	2	1
2001 3/.....	10	...	...	-	-	1	-	1	-	2	2	-
2000.....	22	...	...	-	1	1	2	3	3	3	2	2
1999.....	18	...	...	-	-	3	1	-	1	2	3	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All origins, both sexes												
2003.....	17,732	341	376	122	202	1,938	3,430	2,566	1,950	1,636	1,474	1,219
2002.....	17,638	303	423	140	216	1,892	3,327	2,514	1,975	1,734	1,505	1,155
2001 3/.....	20,308	332	415	137	189	1,899	3,398	2,817	2,387	2,306	1,962	1,463
2000.....	16,765	349	356	140	231	1,914	3,025	2,312	1,852	1,732	1,487	1,059
1999.....	16,889	331	376	186	246	2,093	2,905	2,362	1,869	1,751	1,455	998
All origins, male												
2003.....	13,882	205	202	70	145	1,677	2,947	2,191	1,550	1,224	1,084	893
2002.....	13,640	164	234	64	133	1,604	2,870	2,121	1,548	1,288	1,098	841
2001 3/.....	15,555	197	238	75	118	1,632	2,909	2,308	1,849	1,698	1,424	1,085
2000.....	12,820	203	198	77	150	1,613	2,590	1,918	1,443	1,235	1,084	773
1999.....	12,785	187	194	95	155	1,748	2,443	1,922	1,424	1,257	1,050	746
All origins, female												
2003.....	3,850	136	174	52	57	261	483	375	400	412	390	326
2002.....	3,998	139	189	76	83	288	457	393	427	446	407	314
2001 3/.....	4,753	135	177	62	71	267	489	509	538	608	538	378
2000.....	3,945	146	158	63	81	301	435	394	409	497	403	286
1999.....	4,104	144	182	91	91	345	462	440	445	494	405	252
Hispanic, both sexes												
2003.....	3,355	60	82	16	37	471	815	592	379	296	206	158
2002.....	3,129	52	81	28	36	472	728	554	349	273	191	129
2001 3/.....	3,331	55	69	20	36	436	786	573	387	329	238	156
2000.....	2,917	54	57	24	52	476	650	463	347	248	199	135
1999.....	2,864	53	52	29	60	482	637	482	319	247	185	133

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Hispanic, male												
2003.....	2,809	35	36	7	29	423	716	527	325	245	156	123
2002.....	2,635	28	49	15	22	425	656	493	296	216	151	104
2001 3/.....	2,756	33	38	12	26	397	702	488	312	258	193	121
2000.....	2,431	26	30	19	37	434	580	405	292	199	160	98
1999.....	2,399	26	25	16	43	425	565	428	270	201	153	100
Hispanic, female												
2003.....	546	25	46	9	8	48	99	65	54	51	50	35
2002.....	494	24	32	13	14	47	72	61	53	57	40	25
2001 3/.....	575	22	31	8	10	39	84	85	75	71	45	35
2000.....	486	28	27	5	15	42	70	58	55	49	39	37
1999.....	465	27	27	13	17	57	72	54	49	46	32	33
Non-Hispanic 1/, both sexes												
2003.....	14,245	278	293	106	165	1,451	2,594	1,960	1,560	1,326	1,262	1,052
2002.....	14,346	246	340	112	177	1,406	2,577	1,940	1,608	1,451	1,295	1,016
2001 3/.....	16,726	270	343	116	153	1,449	2,579	2,213	1,965	1,954	1,698	1,283
2000.....	13,717	291	299	114	179	1,424	2,352	1,845	1,492	1,469	1,280	913
1999.....	13,873	270	323	155	186	1,601	2,259	1,870	1,532	1,487	1,256	858
Non-Hispanic 1/, male												
2003.....	10,969	168	166	63	116	1,239	2,216	1,651	1,217	971	923	762
2002.....	10,881	131	184	49	108	1,168	2,194	1,614	1,238	1,065	937	728
2001 3/.....	12,608	161	199	62	92	1,222	2,179	1,796	1,513	1,421	1,211	945
2000.....	10,290	174	168	57	113	1,170	1,989	1,509	1,141	1,022	917	669
1999.....	10,275	158	168	79	112	1,316	1,872	1,484	1,140	1,043	887	641

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic 1/, female												
2003.....	3,276	110	127	43	49	212	378	309	343	355	339	290
2002.....	3,465	115	156	63	69	238	383	326	370	386	358	288
2001 3/.....	4,118	109	144	54	61	227	400	417	452	533	487	338
2000.....	3,427	117	131	57	66	254	363	336	351	447	363	244
1999.....	3,598	112	155	76	74	285	387	386	392	444	369	217
Non-Hispanic White, both sexes												
2003.....	5,367	156	123	45	64	331	576	459	494	555	592	543
2002.....	5,571	117	169	56	77	366	592	464	513	618	650	533
2001 3/.....	7,928	139	160	82	82	382	676	750	892	1,066	1,015	792
2000.....	5,429	141	144	72	92	352	522	490	533	656	617	479
1999.....	5,748	143	156	78	98	409	553	531	604	696	631	455
Non-Hispanic White, male												
2003.....	3,550	94	61	27	45	242	423	312	336	355	392	362
2002.....	3,661	70	89	21	40	263	422	323	337	411	463	339
2001 3/.....	5,453	84	95	40	43	263	466	542	635	788	710	570
2000.....	3,510	96	86	39	54	231	357	321	356	407	429	325
1999.....	3,747	94	83	41	52	283	361	360	393	455	426	328
Non-Hispanic White, female												
2003.....	1,817	62	62	18	19	89	153	147	158	200	200	181
2002.....	1,910	47	80	35	37	103	170	141	176	207	187	194
2001 3/.....	2,475	55	65	42	39	119	210	208	257	278	305	222
2000.....	1,919	45	58	33	38	121	165	169	177	249	188	154
1999.....	2,001	49	73	37	46	126	192	171	211	241	205	127

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Non-Hispanic Black, both sexes												
2003.....	8,276	106	155	54	93	1,052	1,903	1,443	1,000	715	620	460
2002.....	8,147	114	156	45	87	967	1,887	1,412	1,013	770	587	442
2001 3/.....	8,077	119	166	28	64	1,009	1,808	1,370	965	817	610	433
2000.....	7,748	139	143	40	82	1,012	1,759	1,291	902	754	605	390
1999.....	7,536	114	151	65	80	1,117	1,635	1,262	860	730	575	364
Non-Hispanic Black, male												
2003.....	6,980	60	96	33	65	946	1,696	1,292	836	580	498	363
2002.....	6,780	54	89	24	60	845	1,695	1,246	846	607	435	359
2001 3/.....	6,658	68	100	19	47	911	1,640	1,192	806	584	452	333
2000.....	6,382	73	73	16	54	886	1,573	1,138	745	584	443	310
1999.....	6,131	58	74	34	55	969	1,464	1,070	701	548	432	285
Non-Hispanic Black, female												
2003.....	1,296	46	59	21	28	106	207	151	164	135	122	97
2002.....	1,367	60	67	21	27	122	192	166	167	163	152	83
2001 3/.....	1,419	51	66	9	17	98	168	178	159	233	158	100
2000.....	1,366	66	70	24	28	126	186	153	157	170	162	80
1999.....	1,405	56	77	31	25	148	171	192	159	182	143	79
Origin not stated 2/, both sexes												
2003.....	132	3	1	-	-	16	21	14	11	14	6	9
2002.....	163	5	2	-	3	14	22	20	18	10	19	10
2001 3/.....	251	7	3	1	-	14	33	31	35	23	26	24
2000.....	131	4	-	2	-	14	23	4	13	15	8	11
1999.....	152	8	1	2	-	10	9	10	18	17	14	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Origin not stated 2/, male

2003.....	104	2	-	-	-	15	15	13	8	8	5	8
2002.....	124	5	1	-	3	11	20	14	14	7	10	9
2001 3/.....	191	3	1	1	-	13	28	24	24	19	20	19
2000.....	99	3	-	1	-	9	21	4	10	14	7	6
1999.....	111	3	1	-	-	7	6	10	14	13	10	5

Origin not stated 2/, female

2003.....	28	1	1	-	-	1	6	1	3	6	1	1
2002.....	39	-	1	-	-	3	2	6	4	3	9	1
2001 3/.....	60	4	2	-	-	1	5	7	11	4	6	5
2000.....	32	1	-	1	-	5	2	-	3	1	1	5
1999.....	41	5	-	2	-	3	3	-	4	4	4	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All origins, both sexes												
2003.....	11,920	8	40	48	139	1,587	2,823	2,070	1,470	1,066	875	683
2002.....	11,829	9	49	55	150	1,567	2,750	2,018	1,447	1,135	907	604
2001.....	11,348	11	55	59	121	1,525	2,675	1,940	1,368	1,142	836	552
2000.....	10,801	12	28	50	137	1,549	2,414	1,789	1,285	1,064	842	582
1999.....	10,828	8	50	61	163	1,708	2,330	1,795	1,253	1,054	814	551
All origins, male												
2003.....	10,126	7	20	27	107	1,436	2,541	1,867	1,258	860	675	534
2002.....	9,899	3	30	26	101	1,384	2,467	1,790	1,227	918	705	445
2001.....	9,532	7	27	35	90	1,377	2,419	1,708	1,156	892	636	434
2000.....	9,006	9	17	28	110	1,381	2,167	1,574	1,076	842	645	445
1999.....	8,944	5	26	37	118	1,498	2,071	1,568	1,041	807	607	426
All origins, female												
2003.....	1,794	1	20	21	32	151	282	203	212	206	200	149
2002.....	1,930	6	19	29	49	183	283	228	220	217	202	159
2001.....	1,816	4	28	24	31	148	256	232	212	250	200	118
2000.....	1,795	3	11	22	27	168	247	215	209	222	197	137
1999.....	1,884	3	24	24	45	210	259	227	212	247	207	125
Hispanic, both sexes												
2003.....	2,316	1	9	6	25	386	658	453	277	193	120	94
2002.....	2,168	2	12	14	26	402	574	434	245	171	123	65
2001.....	2,123	4	12	5	25	371	603	416	238	172	134	66
2000.....	1,958	1	5	8	33	399	505	350	231	136	113	80
1999.....	1,939	2	5	11	41	410	498	344	206	148	110	80

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Hispanic, male												
2003.....	2,069	1	5	2	20	355	602	421	252	164	95	80
2002.....	1,942	-	8	11	18	371	538	397	222	145	100	53
2001.....	1,892	3	5	4	19	339	559	383	213	146	111	53
2000.....	1,738	-	3	6	28	370	467	319	205	111	96	62
1999.....	1,709	2	3	6	33	373	451	319	180	129	89	58
Hispanic, female												
2003.....	247	-	4	4	5	31	56	32	25	29	25	14
2002.....	226	2	4	3	8	31	36	37	23	26	23	12
2001.....	231	1	7	1	6	32	44	33	25	26	23	13
2000.....	220	1	2	2	5	29	38	31	26	25	17	18
1999.....	230	-	2	5	8	37	47	25	26	19	21	22
Non-Hispanic 1/, both sexes												
2003.....	9,536	7	31	42	114	1,190	2,149	1,607	1,185	866	750	587
2002.....	9,575	7	37	41	123	1,152	2,157	1,569	1,191	958	777	535
2001.....	9,134	7	43	53	96	1,141	2,045	1,509	1,122	965	697	483
2000.....	8,767	11	23	40	104	1,140	1,892	1,436	1,044	917	723	499
1999.....	8,821	6	45	50	122	1,288	1,827	1,441	1,035	898	699	469
Non-Hispanic 1/, male												
2003.....	7,997	6	15	25	87	1,071	1,925	1,436	1,000	692	575	452
2002.....	7,884	3	22	15	82	1,002	1,911	1,382	996	768	599	388
2001.....	7,559	4	22	30	71	1,026	1,837	1,310	936	743	521	379
2000.....	7,205	9	14	21	82	1,002	1,684	1,252	863	721	543	381
1999.....	7,177	3	23	31	85	1,118	1,616	1,239	852	671	513	366

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic 1/, female												
2003.....	1,539	1	16	17	27	119	224	171	185	174	175	135
2002.....	1,691	4	15	26	41	150	246	187	195	190	178	147
2001.....	1,575	3	21	23	25	115	208	199	186	222	176	104
2000.....	1,562	2	9	19	22	138	208	184	181	196	180	118
1999.....	1,644	3	22	19	37	170	211	202	183	227	186	103
Non-Hispanic White, both sexes												
2003.....	2,883	1	11	16	35	221	389	297	316	324	325	294
2002.....	3,052	1	22	21	45	253	389	308	328	363	372	280
2001.....	3,085	4	21	42	48	230	388	330	351	411	348	260
2000.....	2,861	4	12	29	43	222	322	301	323	383	346	259
1999.....	2,995	2	21	23	57	256	335	336	358	405	344	237
Non-Hispanic White, male												
2003.....	2,068	-	3	8	29	177	312	228	228	220	225	201
2002.....	2,139	1	12	7	27	191	297	230	235	256	275	175
2001.....	2,172	2	12	24	32	176	292	249	246	298	238	190
2000.....	1,967	2	6	15	32	165	236	214	230	267	243	185
1999.....	2,119	2	11	15	35	197	246	253	262	277	237	174
Non-Hispanic White, female												
2003.....	815	1	8	8	6	44	77	69	88	104	100	93
2002.....	913	-	10	14	18	62	92	78	93	107	97	105
2001.....	913	2	9	18	16	54	96	81	105	113	110	70
2000.....	894	2	6	14	11	57	86	87	93	116	103	74
1999.....	876	-	10	8	22	59	89	83	96	128	107	63

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Non-Hispanic Black, both sexes												
2003.....	6,319	6	19	23	72	921	1,685	1,271	829	512	397	268
2002.....	6,181	6	14	13	67	851	1,696	1,222	826	558	373	239
2001.....	5,790	2	18	11	45	885	1,605	1,137	737	536	328	203
2000.....	5,622	7	8	11	57	876	1,520	1,090	700	504	350	221
1999.....	5,508	2	18	22	59	979	1,447	1,053	642	463	333	216
Non-Hispanic Black, male												
2003.....	5,666	6	11	15	52	858	1,546	1,176	738	451	329	233
2002.....	5,482	2	9	6	48	768	1,557	1,123	731	481	303	202
2001.....	5,190	1	9	6	37	826	1,503	1,032	665	431	269	172
2000.....	5,011	7	5	6	46	800	1,406	1,002	614	435	280	181
1999.....	4,826	1	7	14	46	874	1,334	947	565	374	262	179
Non-Hispanic Black, female												
2003.....	653	-	8	8	20	63	139	95	91	61	68	35
2002.....	699	4	5	7	19	83	139	99	95	77	70	37
2001.....	600	1	9	5	8	59	102	105	72	105	59	31
2000.....	611	-	3	5	11	76	114	88	86	69	70	40
1999.....	682	1	11	8	13	105	113	106	77	89	71	37
Origin not stated 2/, both sexes												
2003.....	68	-	-	-	-	11	16	10	8	7	5	2
2002.....	86	-	-	-	1	13	19	15	11	6	7	4
2001.....	91	-	-	1	-	13	27	15	8	5	5	3
2000.....	76	-	-	2	-	10	17	3	10	11	6	3
1999.....	68	-	-	-	-	10	5	10	12	8	5	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Origin not stated 2/, male

2003.....	60	-	-	-	-	10	14	10	6	4	5	2
2002.....	73	-	-	-	1	11	18	11	9	5	6	4
2001.....	81	-	-	1	-	12	23	15	7	3	4	2
2000.....	63	-	-	1	-	9	16	3	8	10	6	2
1999.....	58	-	-	-	-	7	4	10	9	7	5	2

Origin not stated 2/, female

2003.....	8	-	-	-	-	1	2	-	2	3	-	-
2002.....	13	-	-	-	-	2	1	4	2	1	1	-
2001.....	10	-	-	-	-	1	4	-	1	2	1	1
2000.....	13	-	-	1	-	1	1	-	2	1	-	1
1999.....	10	-	-	-	-	3	1	-	3	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)												
All origins, both sexes												
2003.....	5,812	333	336	74	63	351	607	496	480	570	599	536
2002.....	5,809	294	374	85	66	325	577	496	528	599	598	551
2001 3/.....	8,960	321	360	78	68	374	723	877	1,019	1,164	1,126	911
2000.....	5,964	337	328	90	94	365	611	523	567	668	645	477
1999.....	6,061	323	326	125	83	385	575	567	616	697	641	447
All origins, male												
2003.....	3,756	198	182	43	38	241	406	324	292	364	409	359
2002.....	3,741	161	204	38	32	220	403	331	321	370	393	396
2001 3/.....	6,023	190	211	40	28	255	490	600	693	806	788	651
2000.....	3,814	194	181	49	40	232	423	344	367	393	439	328
1999.....	3,841	182	168	58	37	250	372	354	383	450	443	320
All origins, female												
2003.....	2,056	135	154	31	25	110	201	172	188	206	190	177
2002.....	2,068	133	170	47	34	105	174	165	207	229	205	155
2001 3/.....	2,937	131	149	38	40	119	233	277	326	358	338	260
2000.....	2,150	143	147	41	54	133	188	179	200	275	206	149
1999.....	2,220	141	158	67	46	135	203	213	233	247	198	127
Hispanic, both sexes												
2003.....	1,039	59	73	10	12	85	157	139	102	103	86	64
2002.....	961	50	69	14	10	70	154	120	104	102	68	64
2001 3/.....	1,208	51	57	15	11	65	183	157	149	157	104	90
2000.....	959	53	52	16	19	77	145	113	116	112	86	55
1999.....	925	51	47	18	19	72	139	138	113	99	75	53

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)												
Hispanic, male												
2003.....	740	34	31	5	9	68	114	106	73	81	61	43
2002.....	693	28	41	4	4	54	118	96	74	71	51	51
2001 3/.....	864	30	33	8	7	58	143	105	99	112	82	68
2000.....	693	26	27	13	9	64	113	86	87	88	64	36
1999.....	690	24	22	10	10	52	114	109	90	72	64	42
Hispanic, female												
2003.....	299	25	42	5	3	17	43	33	29	22	25	21
2002.....	268	22	28	10	6	16	36	24	30	31	17	13
2001 3/.....	344	21	24	7	4	7	40	52	50	45	22	22
2000.....	266	27	25	3	10	13	32	27	29	24	22	19
1999.....	235	27	25	8	9	20	25	29	23	27	11	11
Non-Hispanic 1/, both sexes												
2003.....	4,709	271	262	64	51	261	445	353	375	460	512	465
2002.....	4,771	239	303	71	54	254	420	371	417	493	518	481
2001 3/.....	7,592	263	300	63	57	308	534	704	843	989	1,001	800
2000.....	4,950	280	276	74	75	284	460	409	448	552	557	414
1999.....	5,052	264	278	105	64	313	432	429	497	589	557	389
Non-Hispanic 1/, male												
2003.....	2,972	162	151	38	29	168	291	215	217	279	348	310
2002.....	2,997	128	162	34	26	166	283	232	242	297	338	340
2001 3/.....	5,049	157	177	32	21	196	342	486	577	678	690	566
2000.....	3,085	165	154	36	31	168	305	257	278	301	374	288
1999.....	3,098	155	145	48	27	198	256	245	288	372	374	275

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)												
Non-Hispanic 1/, female												
2003.....	1,737	109	111	26	22	93	154	138	158	181	164	155
2002.....	1,774	111	141	37	28	88	137	139	175	196	180	141
2001 3/.....	2,543	106	123	31	36	112	192	218	266	311	311	234
2000.....	1,865	115	122	38	44	116	155	152	170	251	183	126
1999.....	1,954	109	133	57	37	115	176	184	209	217	183	114
Non-Hispanic White, both sexes												
2003.....	2,484	155	112	29	29	110	187	162	178	231	267	249
2002.....	2,519	116	147	35	32	113	203	156	185	255	278	253
2001 3/.....	4,843	135	139	40	34	152	288	420	541	655	667	532
2000.....	2,568	137	132	43	49	130	200	189	210	273	271	220
1999.....	2,753	141	135	55	41	153	218	195	246	291	287	218
Non-Hispanic White, male												
2003.....	1,482	94	58	19	16	65	111	84	108	135	167	161
2002.....	1,522	69	77	14	13	72	125	93	102	155	188	164
2001 3/.....	3,281	82	83	16	11	87	174	293	389	490	472	380
2000.....	1,543	94	80	24	22	66	121	107	126	140	186	140
1999.....	1,628	92	72	26	17	86	115	107	131	178	189	154
Non-Hispanic White, female												
2003.....	1,002	61	54	10	13	45	76	78	70	96	100	88
2002.....	997	47	70	21	19	41	78	63	83	100	90	89
2001 3/.....	1,562	53	56	24	23	65	114	127	152	165	195	152
2000.....	1,025	43	52	19	27	64	79	82	84	133	85	80
1999.....	1,125	49	63	29	24	67	103	88	115	113	98	64

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02.X85-X92,X96-Y09,Y87.1)												
Non-Hispanic Black, both sexes												
2003.....	1,957	100	136	31	21	131	218	172	171	203	223	192
2002.....	1,966	108	142	32	20	116	191	190	187	212	214	203
2001 3/.....	2,287	117	148	17	19	124	203	233	228	281	282	230
2000.....	2,126	132	135	29	25	136	239	201	202	250	255	169
1999.....	2,028	112	133	43	21	138	188	209	218	267	242	148
Non-Hispanic Black, male												
2003.....	1,314	54	85	18	13	88	150	116	98	129	169	130
2002.....	1,298	52	80	18	12	77	138	123	115	126	132	157
2001 3/.....	1,468	67	91	13	10	85	137	160	141	153	183	161
2000.....	1,371	66	68	10	8	86	167	136	131	149	163	129
1999.....	1,305	57	67	20	9	95	130	123	136	174	170	106
Non-Hispanic Black, female												
2003.....	643	46	51	13	8	43	68	56	73	74	54	62
2002.....	668	56	62	14	8	39	53	67	72	86	82	46
2001 3/.....	819	50	57	4	9	39	66	73	87	128	99	69
2000.....	755	66	67	19	17	50	72	65	71	101	92	40
1999.....	723	55	66	23	12	43	58	86	82	93	72	42
Origin not stated 2/, both sexes												
2003.....	64	3	1	-	-	5	5	4	3	7	1	7
2002.....	77	5	2	-	2	1	3	5	7	4	12	6
2001 3/.....	160	7	3	-	-	1	6	16	27	18	21	21
2000.....	55	4	-	-	-	4	6	1	3	4	2	8
1999.....	84	8	1	2	-	-	4	-	6	9	9	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Origin not stated 2/, male

2003.....	44	2	-	-	-	5	1	3	2	4	-	6
2002.....	51	5	1	-	2	-	2	3	5	2	4	5
2001 3/.....	110	3	1	-	-	1	5	9	17	16	16	17
2000.....	36	3	-	-	-	-	5	1	2	4	1	4
1999.....	53	3	1	-	-	-	2	-	5	6	5	3

Origin not stated 2/, female

2003.....	20	1	1	-	-	-	4	1	1	3	1	1
2002.....	26	-	1	-	-	1	1	2	2	2	8	1
2001 3/.....	50	4	2	-	-	-	1	7	10	2	5	4
2000.....	19	1	-	-	-	4	1	-	1	-	1	4
1999.....	31	5	-	2	-	-	2	-	1	3	4	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Legal intervention (Y35,Y89.0)</b>												
<b>All origins, both sexes</b>												
2003.....	423	-	-	-	2	20	73	63	72	66	51	29
2002.....	384	-	-	-	3	24	67	51	51	64	45	43
2001.....	396	-	-	-	-	26	62	47	60	65	50	33
2000.....	359	-	-	-	3	27	46	59	51	60	41	38
1999.....	398	-	-	-	1	20	47	66	70	72	49	19
<b>All origins, male</b>												
2003.....	405	-	-	-	1	19	70	61	70	65	49	27
2002.....	369	-	-	-	3	23	64	51	48	60	45	40
2001.....	380	-	-	-	-	25	59	46	58	63	45	33
2000.....	348	-	-	-	3	27	44	58	50	58	38	37
1999.....	392	-	-	-	1	20	47	66	69	69	48	19
<b>All origins, female</b>												
2003.....	18	-	-	-	1	1	3	2	2	1	2	2
2002.....	15	-	-	-	-	1	3	-	3	4	-	3
2001.....	16	-	-	-	-	1	3	1	2	2	5	-
2000.....	11	-	-	-	-	-	2	1	1	2	3	1
1999.....	6	-	-	-	-	-	-	-	1	3	1	-
<b>Hispanic, both sexes</b>												
2003.....	74	-	-	-	1	6	22	14	18	6	4	1
2002.....	68	-	-	-	1	5	17	13	9	9	5	2
2001.....	65	-	-	-	-	7	16	15	9	6	7	2
2000.....	48	-	-	-	1	5	13	6	9	5	4	2
1999.....	58	-	-	-	-	5	8	10	13	10	4	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Legal intervention (Y35,Y89.0)												
Hispanic, male												
2003.....	73	-	-	-	-	6	22	14	18	6	4	1
2002.....	67	-	-	-	1	5	17	13	8	9	5	2
2001.....	65	-	-	-	-	7	16	15	9	6	7	2
2000.....	47	-	-	-	1	5	12	6	9	5	4	2
1999.....	58	-	-	-	-	5	8	10	13	10	4	1
Hispanic, female												
2003.....	1	-	-	-	1	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-
Non-Hispanic 1/, both sexes												
2003.....	347	-	-	-	1	14	50	49	54	60	47	28
2002.....	315	-	-	-	2	19	50	38	42	54	40	41
2001.....	330	-	-	-	-	19	46	32	51	59	43	30
2000.....	308	-	-	-	2	22	32	53	42	53	37	36
1999.....	338	-	-	-	1	15	39	55	57	61	45	18
Non-Hispanic 1/, male												
2003.....	330	-	-	-	1	13	47	47	52	59	45	26
2002.....	301	-	-	-	2	18	47	38	40	50	40	38
2001.....	314	-	-	-	-	18	43	31	49	57	38	30
2000.....	298	-	-	-	2	22	31	52	41	51	34	35
1999.....	332	-	-	-	1	15	39	55	56	58	44	18

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Legal intervention (Y35,Y89.0)												
Non-Hispanic 1/, female												
2003.....	17	-	-	-	-	1	3	2	2	1	2	2
2002.....	14	-	-	-	-	1	3	-	2	4	-	3
2001.....	16	-	-	-	-	1	3	1	2	2	5	-
2000.....	10	-	-	-	-	-	1	1	1	2	3	1
1999.....	6	-	-	-	-	-	-	-	1	3	1	-
Non-Hispanic White, both sexes												
2003.....	211	-	-	-	-	7	20	17	28	46	34	21
2002.....	191	-	-	-	-	9	19	16	32	36	28	26
2001.....	230	-	-	-	-	8	29	20	36	40	34	21
2000.....	187	-	-	-	-	6	20	30	27	31	21	30
1999.....	197	-	-	-	-	5	15	28	34	36	29	14
Non-Hispanic White, male												
2003.....	199	-	-	-	-	6	19	17	26	45	32	19
2002.....	181	-	-	-	-	8	17	16	31	34	28	23
2001.....	221	-	-	-	-	7	28	19	36	38	31	21
2000.....	182	-	-	-	-	6	19	29	26	30	21	30
1999.....	192	-	-	-	-	5	15	28	34	33	28	14
Non-Hispanic White, female												
2003.....	12	-	-	-	-	1	1	-	2	1	2	2
2002.....	10	-	-	-	-	1	2	-	1	2	-	3
2001.....	9	-	-	-	-	1	1	1	-	2	3	-
2000.....	5	-	-	-	-	-	1	1	1	1	-	-
1999.....	5	-	-	-	-	-	-	-	-	3	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

Non-Hispanic Black, both sexes

2003.....	117	-	-	-	1	6	27	28	24	12	10	6
2002.....	109	-	-	-	2	10	27	19	9	17	11	11
2001.....	82	-	-	-	-	10	16	9	12	15	7	8
2000.....	110	-	-	-	2	15	12	19	14	20	15	6
1999.....	125	-	-	-	1	10	21	24	18	21	15	4

Non-Hispanic Black, male

2003.....	114	-	-	-	1	6	25	27	24	12	10	6
2002.....	105	-	-	-	2	10	26	19	8	15	11	11
2001.....	77	-	-	-	-	10	14	9	11	15	5	8
2000.....	107	-	-	-	2	15	12	19	14	20	13	5
1999.....	124	-	-	-	1	10	21	24	17	21	15	4

Non-Hispanic Black, female

2003.....	3	-	-	-	-	-	2	1	-	-	-	-
2002.....	4	-	-	-	-	-	1	-	1	2	-	-
2001.....	5	-	-	-	-	-	2	-	1	-	2	-
2000.....	3	-	-	-	-	-	-	-	-	-	2	1
1999.....	1	-	-	-	-	-	-	-	1	-	-	-

Origin not stated 2/, both sexes

2003.....	2	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	1	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	1
2000.....	3	-	-	-	-	-	1	-	-	2	-	-
1999.....	2	-	-	-	-	-	-	1	-	1	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All origins, both sexes												
2003.....	5,072	75	31	16	39	169	394	351	484	686	895	777
2002.....	4,830	101	57	26	42	143	344	352	476	727	789	709
2001.....	4,198	83	44	17	34	132	280	311	417	674	722	619
2000.....	3,819	65	39	18	42	120	250	306	373	599	687	543
1999.....	3,917	55	43	12	37	159	271	313	430	644	651	528
All origins, male												
2003.....	3,295	50	17	9	32	124	288	249	331	450	563	493
2002.....	3,114	52	28	11	27	108	250	256	329	494	487	432
2001.....	2,833	49	19	10	28	102	214	231	284	438	493	433
2000.....	2,635	27	26	13	37	93	192	223	266	415	470	374
1999.....	2,692	33	21	6	32	132	213	230	293	446	441	357
All origins, female												
2003.....	1,777	25	14	7	7	45	106	102	153	236	332	284
2002.....	1,716	49	29	15	15	35	94	96	147	233	302	277
2001.....	1,365	34	25	7	6	30	66	80	133	236	229	186
2000.....	1,184	38	13	5	5	27	58	83	107	184	217	169
1999.....	1,225	22	22	6	5	27	58	83	137	198	210	171
Hispanic, both sexes												
2003.....	339	6	6	4	9	15	36	35	50	42	38	40
2002.....	320	14	11	3	1	13	38	29	44	53	37	34
2001.....	283	7	6	1	3	13	21	31	32	49	43	31
2000.....	215	5	5	-	1	11	23	29	32	41	23	19
1999.....	249	6	9	1	3	20	32	32	37	36	25	18

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Hispanic, male												
2003.....	258	1	4	2	7	10	26	29	40	31	34	30
2002.....	242	6	7	1	-	12	33	23	35	40	24	28
2001.....	216	4	3	1	3	11	15	25	27	35	38	19
2000.....	165	2	2	-	1	10	20	22	27	31	16	16
1999.....	200	3	6	1	2	17	29	26	28	29	20	15
Hispanic, female												
2003.....	81	5	2	2	2	5	10	6	10	11	4	10
2002.....	78	8	4	2	1	1	5	6	9	13	13	6
2001.....	67	3	3	-	-	2	6	6	5	14	5	12
2000.....	50	3	3	-	-	1	3	7	5	10	7	3
1999.....	49	3	3	-	1	3	3	6	9	7	5	3
Non-Hispanic 1/, both sexes												
2003.....	4,701	68	25	12	30	154	358	315	433	640	851	733
2002.....	4,470	87	44	23	40	127	305	320	429	671	749	671
2001.....	3,876	76	38	16	31	118	259	279	382	622	676	586
2000.....	3,571	58	34	18	41	109	227	275	339	553	661	522
1999.....	3,630	49	34	11	34	139	237	276	390	606	624	507
Non-Hispanic 1/, male												
2003.....	3,016	48	13	7	25	114	262	219	291	416	526	462
2002.....	2,836	46	21	10	26	94	216	230	291	452	460	400
2001.....	2,586	45	16	9	25	90	199	205	254	400	453	414
2000.....	2,445	25	24	13	36	83	172	199	237	382	452	356
1999.....	2,463	30	15	5	30	115	182	200	262	415	419	340

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Non-Hispanic 1/, female												
2003.....	1,685	20	12	5	5	40	96	96	142	224	325	271
2002.....	1,634	41	23	13	14	33	89	90	138	219	289	271
2001.....	1,290	31	22	7	6	28	60	74	128	222	223	172
2000.....	1,126	33	10	5	5	26	55	76	102	171	209	166
1999.....	1,167	19	19	6	4	24	55	76	128	191	205	167
Non-Hispanic White, both sexes												
2003.....	3,898	45	15	3	21	135	304	266	369	533	714	610
2002.....	3,627	43	33	13	31	107	263	262	345	546	614	544
2001.....	3,155	40	25	8	27	103	215	229	320	499	555	478
2000.....	2,823	35	22	9	28	93	178	222	268	448	517	415
1999.....	2,848	30	15	9	29	106	184	218	313	480	480	390
Non-Hispanic White, male												
2003.....	2,441	28	7	1	19	98	218	181	241	347	433	374
2002.....	2,265	23	16	4	20	77	187	187	232	359	380	321
2001.....	2,054	25	10	3	22	78	160	165	212	312	361	327
2000.....	1,900	16	15	6	25	70	131	158	183	309	352	277
1999.....	1,887	19	8	4	26	90	142	150	207	319	327	247
Non-Hispanic White, female												
2003.....	1,457	17	8	2	2	37	86	85	128	186	281	236
2002.....	1,362	20	17	9	11	30	76	75	113	187	234	223
2001.....	1,101	15	15	5	5	25	55	64	108	187	194	151
2000.....	923	19	7	3	3	23	47	64	85	139	165	138
1999.....	961	11	7	5	3	16	42	68	106	161	153	143

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic Black, both sexes

2003.....	675	21	9	7	7	16	34	36	57	91	122	115
2002.....	738	37	7	9	6	14	38	51	67	115	125	116
2001.....	629	36	13	7	4	12	31	37	53	110	112	98
2000.....	654	14	11	9	12	12	38	44	59	97	133	100
1999.....	686	18	15	2	5	25	45	51	68	113	126	107

Non-Hispanic Black, male

2003.....	487	19	5	4	4	13	29	30	44	58	83	83
2002.....	504	18	3	5	6	12	27	38	46	87	74	74
2001.....	467	20	6	5	3	10	27	30	35	79	86	79
2000.....	478	3	8	7	10	12	33	34	44	67	93	75
1999.....	514	10	7	1	4	20	35	44	47	86	84	86

Non-Hispanic Black, female

2003.....	188	2	4	3	3	3	5	6	13	33	39	32
2002.....	234	19	4	4	-	2	11	13	21	28	51	42
2001.....	162	16	7	2	1	2	4	7	18	31	26	19
2000.....	176	11	3	2	2	-	5	10	15	30	40	25
1999.....	172	8	8	1	1	5	10	7	21	27	42	21

Origin not stated 2/, both sexes

2003.....	32	1	-	-	-	-	-	1	1	4	6	4
2002.....	40	-	2	-	1	3	1	3	3	3	3	4
2001.....	39	-	-	-	-	1	-	1	3	3	3	2
2000.....	33	2	-	-	-	-	-	2	2	5	3	2
1999.....	38	-	-	-	-	-	2	5	3	2	2	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Origin not stated 2/, male

2003.....	21	1	-	-	-	-	-	1	-	3	3	1
2002.....	36	-	-	-	1	2	1	3	3	2	3	4
2001.....	31	-	-	-	-	1	-	1	3	3	2	-
2000.....	25	-	-	-	-	-	-	2	2	2	2	2
1999.....	29	-	-	-	-	-	2	4	3	2	2	2

Origin not stated 2/, female

2003.....	11	-	-	-	-	-	-	-	1	1	3	3
2002.....	4	-	2	-	-	1	-	-	-	1	-	-
2001.....	8	-	-	-	-	-	-	-	-	-	1	2
2000.....	8	2	-	-	-	-	-	-	-	3	1	-
1999.....	9	-	-	-	-	-	-	1	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Discharge of firearms, undetermined intent (Y22-Y24)												
All origins, both sexes												
2003.....	232	-	1	1	11	31	38	21	17	21	22	19
2002.....	243	-	1	2	4	35	44	27	19	31	19	19
2001.....	231	-	-	2	4	24	40	32	16	33	17	16
2000.....	230	-	-	2	8	26	45	32	20	18	21	9
1999.....	324	-	3	-	12	68	59	32	19	28	29	18
All origins, male												
2003.....	186	-	1	1	11	29	31	18	15	12	14	11
2002.....	199	-	1	2	3	34	37	23	16	22	17	14
2001.....	190	-	-	2	4	23	36	27	15	20	13	12
2000.....	190	-	-	2	7	25	37	26	17	12	18	6
1999.....	277	-	3	-	12	61	50	28	16	24	21	15
All origins, female												
2003.....	46	-	-	-	-	2	7	3	2	9	8	8
2002.....	44	-	-	-	1	1	7	4	3	9	2	5
2001.....	41	-	-	-	-	1	4	5	1	13	4	4
2000.....	40	-	-	-	1	1	8	6	3	6	3	3
1999.....	47	-	-	-	-	7	9	4	3	4	8	3
Hispanic, both sexes												
2003.....	31	-	-	-	3	6	7	4	2	2	3	3
2002.....	23	-	-	-	-	6	8	4	2	3	-	-
2001.....	21	-	-	-	1	4	4	3	1	4	-	2
2000.....	20	-	-	-	-	7	6	3	1	2	-	-
1999.....	31	-	1	-	1	9	9	2	1	2	1	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Discharge of firearms, undetermined intent (Y22-Y24)												
Hispanic, male												
2003.....	26	-	-	-	3	4	5	4	2	2	3	2
2002.....	21	-	-	-	-	6	7	4	2	2	-	-
2001.....	16	-	-	-	1	4	3	2	1	2	-	1
2000.....	17	-	-	-	-	7	6	2	-	1	-	-
1999.....	26	-	1	-	1	8	8	1	1	2	-	1
Hispanic, female												
2003.....	5	-	-	-	-	2	2	-	-	-	-	1
2002.....	2	-	-	-	-	-	1	-	-	1	-	-
2001.....	5	-	-	-	-	-	1	1	-	2	-	1
2000.....	3	-	-	-	-	-	-	1	1	1	-	-
1999.....	5	-	-	-	-	1	1	1	-	-	1	1
Non-Hispanic 1/, both sexes												
2003.....	201	-	1	1	8	25	31	17	15	19	19	16
2002.....	215	-	1	2	3	27	36	22	17	28	19	19
2001.....	208	-	-	2	3	20	36	29	14	29	17	14
2000.....	209	-	-	2	8	19	39	29	19	16	21	9
1999.....	293	-	2	-	11	59	50	30	18	26	28	16
Non-Hispanic 1/, male												
2003.....	160	-	1	1	8	25	26	14	13	10	11	9
2002.....	173	-	1	2	2	26	30	18	14	20	17	14
2001.....	172	-	-	2	3	19	33	25	13	18	13	11
2000.....	172	-	-	2	7	18	31	24	17	11	18	6
1999.....	251	-	2	-	11	53	42	27	15	22	21	14

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Discharge of firearms, undetermined intent (Y22-Y24)												
Non-Hispanic 1/, female												
2003.....	41	-	-	-	-	-	5	3	2	9	8	7
2002.....	42	-	-	-	1	1	6	4	3	8	2	5
2001.....	36	-	-	-	-	1	3	4	1	11	4	3
2000.....	37	-	-	-	1	1	8	5	2	5	3	3
1999.....	42	-	-	-	-	6	8	3	3	4	7	2
Non-Hispanic White, both sexes												
2003.....	152	-	-	-	4	17	17	10	11	16	17	15
2002.....	169	-	-	-	3	18	29	16	11	22	16	18
2001.....	166	-	-	-	3	13	23	23	12	27	14	13
2000.....	148	-	-	-	5	14	22	20	10	12	18	8
1999.....	217	-	1	-	10	33	34	20	13	23	24	11
Non-Hispanic White, male												
2003.....	118	-	-	-	4	17	14	8	9	8	9	9
2002.....	132	-	-	-	2	17	23	12	9	16	14	13
2001.....	133	-	-	-	3	12	21	20	11	16	11	10
2000.....	120	-	-	-	5	13	17	16	10	7	17	5
1999.....	187	-	1	-	10	31	29	18	11	19	20	9
Non-Hispanic White, female												
2003.....	34	-	-	-	-	-	3	2	2	8	8	6
2002.....	37	-	-	-	1	1	6	4	2	6	2	5
2001.....	33	-	-	-	-	1	2	3	1	11	3	3
2000.....	28	-	-	-	-	1	5	4	-	5	1	3
1999.....	30	-	-	-	-	2	5	2	2	4	4	2





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
All origins, both sexes												
2003.....	4,840	75	30	15	28	138	356	330	467	665	873	758
2002.....	4,587	101	56	24	38	108	300	325	457	696	770	690
2001.....	3,967	83	44	15	30	108	240	279	401	641	705	603
2000.....	3,589	65	39	16	34	94	205	274	353	581	666	534
1999.....	3,593	55	40	12	25	91	212	281	411	616	622	510
All origins, male												
2003.....	3,109	50	16	8	21	95	257	231	316	438	549	482
2002.....	2,915	52	27	9	24	74	213	233	313	472	470	418
2001.....	2,643	49	19	8	24	79	178	204	269	418	480	421
2000.....	2,445	27	26	11	30	68	155	197	249	403	452	368
1999.....	2,415	33	18	6	20	71	163	202	277	422	420	342
All origins, female												
2003.....	1,731	25	14	7	7	43	99	99	151	227	324	276
2002.....	1,672	49	29	15	14	34	87	92	144	224	300	272
2001.....	1,324	34	25	7	6	29	62	75	132	223	225	182
2000.....	1,144	38	13	5	4	26	50	77	104	178	214	166
1999.....	1,178	22	22	6	5	20	49	79	134	194	202	168
Hispanic, both sexes												
2003.....	308	6	6	4	6	9	29	31	48	40	35	37
2002.....	297	14	11	3	1	7	30	25	42	50	37	34
2001.....	262	7	6	1	2	9	17	28	31	45	43	29
2000.....	195	5	5	-	1	4	17	26	31	39	23	19
1999.....	218	6	8	1	2	11	23	30	36	34	24	16

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Hispanic, male												
2003.....	232	1	4	2	4	6	21	25	38	29	31	28
2002.....	221	6	7	1	-	6	26	19	33	38	24	28
2001.....	200	4	3	1	2	7	12	23	26	33	38	18
2000.....	148	2	2	-	1	3	14	20	27	30	16	16
1999.....	174	3	5	1	1	9	21	25	27	27	20	14
Hispanic, female												
2003.....	76	5	2	2	2	3	8	6	10	11	4	9
2002.....	76	8	4	2	1	1	4	6	9	12	13	6
2001.....	62	3	3	-	-	2	5	5	5	12	5	11
2000.....	47	3	3	-	-	1	3	6	4	9	7	3
1999.....	44	3	3	-	1	2	2	5	9	7	4	2
Non-Hispanic 1/, both sexes												
2003.....	4,500	68	24	11	22	129	327	298	418	621	832	717
2002.....	4,255	87	43	21	37	100	269	298	412	643	730	652
2001.....	3,668	76	38	14	28	98	223	250	368	593	659	572
2000.....	3,362	58	34	16	33	90	188	246	320	537	640	513
1999.....	3,337	49	32	11	23	80	187	246	372	580	596	491
Non-Hispanic 1/, male												
2003.....	2,856	48	12	6	17	89	236	205	278	406	515	453
2002.....	2,663	46	20	8	24	68	186	212	277	432	443	386
2001.....	2,414	45	16	7	22	71	166	180	241	382	440	403
2000.....	2,273	25	24	11	29	65	141	175	220	371	434	350
1999.....	2,212	30	13	5	19	62	140	173	247	393	398	326

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Non-Hispanic 1/, female												
2003.....	1,644	20	12	5	5	40	91	93	140	215	317	264
2002.....	1,592	41	23	13	13	32	83	86	135	211	287	266
2001.....	1,254	31	22	7	6	27	57	70	127	211	219	169
2000.....	1,089	33	10	5	4	25	47	71	100	166	206	163
1999.....	1,125	19	19	6	4	18	47	73	125	187	198	165
Non-Hispanic White, both sexes												
2003.....	3,746	45	15	3	17	118	287	256	358	517	697	595
2002.....	3,458	43	33	13	28	89	234	246	334	524	598	526
2001.....	2,989	40	25	8	24	90	192	206	308	472	541	465
2000.....	2,675	35	22	9	23	79	156	202	258	436	499	407
1999.....	2,631	30	14	9	19	73	150	198	300	457	456	379
Non-Hispanic White, male												
2003.....	2,323	28	7	1	15	81	204	173	232	339	424	365
2002.....	2,133	23	16	4	18	60	164	175	223	343	366	308
2001.....	1,921	25	10	3	19	66	139	145	201	296	350	317
2000.....	1,780	16	15	6	20	57	114	142	173	302	335	272
1999.....	1,700	19	7	4	16	59	113	132	196	300	307	238
Non-Hispanic White, female												
2003.....	1,423	17	8	2	2	37	83	83	126	178	273	230
2002.....	1,325	20	17	9	10	29	70	71	111	181	232	218
2001.....	1,068	15	15	5	5	24	53	61	107	176	191	148
2000.....	895	19	7	3	3	22	42	60	85	134	164	135
1999.....	931	11	7	5	3	14	37	66	104	157	149	141

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Non-Hispanic Black, both sexes												
2003.....	640	21	8	6	4	9	24	32	54	89	121	114
2002.....	701	37	6	7	6	7	31	45	63	111	123	116
2001.....	598	36	13	5	4	5	23	33	52	110	110	97
2000.....	601	14	11	7	9	7	24	36	52	93	130	99
1999.....	621	18	14	2	4	5	30	41	64	110	123	103
Non-Hispanic Black, male												
2003.....	455	19	4	3	1	6	20	26	41	57	82	83
2002.....	469	18	2	3	6	5	20	32	42	84	72	74
2001.....	439	20	6	3	3	3	20	27	34	79	85	78
2000.....	432	3	8	5	8	7	20	27	39	63	92	74
1999.....	459	10	6	1	3	3	23	35	44	83	83	82
Non-Hispanic Black, female												
2003.....	185	2	4	3	3	3	4	6	13	32	39	31
2002.....	232	19	4	4	-	2	11	13	21	27	51	42
2001.....	159	16	7	2	1	2	3	6	18	31	25	19
2000.....	169	11	3	2	1	-	4	9	13	30	38	25
1999.....	162	8	8	1	1	2	7	6	20	27	40	21
Origin not stated 2/, both sexes												
2003.....	32	1	-	-	-	-	-	1	1	4	6	4
2002.....	35	-	2	-	-	1	1	2	3	3	3	4
2001.....	37	-	-	-	-	1	-	1	2	3	3	2
2000.....	32	2	-	-	-	-	-	2	2	5	3	2
1999.....	38	-	-	-	-	-	2	5	3	2	2	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Origin not stated 2/, male

2003.....	21	1	-	-	-	-	-	1	-	3	3	1
2002.....	31	-	-	-	-	-	1	2	3	2	3	4
2001.....	29	-	-	-	-	1	-	1	2	3	2	-
2000.....	24	-	-	-	-	-	-	2	2	2	2	2
1999.....	29	-	-	-	-	-	2	4	3	2	2	2

Origin not stated 2/, female

2003.....	11	-	-	-	-	-	-	-	1	1	3	3
2002.....	4	-	2	-	-	1	-	-	-	1	-	-
2001.....	8	-	-	-	-	-	-	-	-	-	1	2
2000.....	8	2	-	-	-	-	-	-	-	3	1	-
1999.....	9	-	-	-	-	-	-	1	-	-	-	1



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Operations of war and their sequelae (Y36,Y89.1)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	1	-	-	-
2001.....	2	-	-	-	-	-	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	2	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	14	-	-	-	-	-	1	-	-	-	-	-
2002.....	19	-	-	-	-	-	-	-	-	-	1	1
2001.....	15	-	-	-	-	-	-	-	-	1	-	1
2000.....	15	-	-	-	-	-	-	-	-	-	-	-
1999.....	21	-	-	-	-	-	-	-	-	1	-	-

Non-Hispanic 1/, male

2003.....	14	-	-	-	-	-	1	-	-	-	-	-
2002.....	19	-	-	-	-	-	-	-	-	-	1	1
2001.....	14	-	-	-	-	-	-	-	-	1	-	1
2000.....	15	-	-	-	-	-	-	-	-	-	-	-
1999.....	20	-	-	-	-	-	-	-	-	1	-	-







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
All origins, both sexes												
2003.....	2,855	15	24	6	6	21	23	26	43	75	104	145
2002.....	2,843	15	19	13	22	16	19	27	52	73	98	114
2001.....	3,021	26	24	5	9	18	25	24	55	71	101	142
2000.....	3,059	19	22	18	13	10	22	37	48	65	84	128
1999.....	2,823	54	30	7	11	17	19	29	49	66	86	91
All origins, male												
2003.....	1,333	8	16	2	2	18	8	14	22	36	60	80
2002.....	1,236	8	11	8	18	10	9	16	31	35	47	60
2001.....	1,357	15	14	2	6	6	13	11	31	39	44	75
2000.....	1,430	12	11	11	4	5	11	23	24	33	43	59
1999.....	1,308	31	12	4	5	8	13	9	34	37	41	37
All origins, female												
2003.....	1,522	7	8	4	4	3	15	12	21	39	44	65
2002.....	1,607	7	8	5	4	6	10	11	21	38	51	54
2001.....	1,664	11	10	3	3	12	12	13	24	32	57	67
2000.....	1,629	7	11	7	9	5	11	14	24	32	41	69
1999.....	1,515	23	18	3	6	9	6	20	15	29	45	54
Hispanic, both sexes												
2003.....	155	3	4	-	1	3	2	6	3	11	11	8
2002.....	137	2	7	1	4	5	1	4	4	7	6	4
2001.....	145	3	6	1	3	4	4	5	2	4	8	14
2000.....	136	3	4	3	3	2	5	2	2	5	6	8
1999.....	100	5	6	2	1	2	2	2	6	6	2	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
Hispanic, male												
2003.....	76	1	4	-	-	2	1	2	-	6	9	7
2002.....	57	2	4	-	4	3	-	1	-	3	2	3
2001.....	65	-	3	1	2	1	2	2	1	2	2	9
2000.....	70	2	2	2	2	1	5	1	1	2	3	3
1999.....	52	3	3	1	1	1	1	-	3	5	2	2
Hispanic, female												
2003.....	79	2	-	-	1	1	1	4	3	5	2	1
2002.....	80	-	3	1	-	2	1	3	4	4	4	1
2001.....	80	3	3	-	1	3	2	3	1	2	6	5
2000.....	66	1	2	1	1	1	-	1	1	3	3	5
1999.....	48	2	3	1	-	1	1	2	3	1	-	3
Non-Hispanic 1/, both sexes												
2003.....	2,693	12	19	6	5	17	21	20	40	64	92	135
2002.....	2,695	13	12	12	18	11	18	23	48	66	92	109
2001.....	2,861	23	18	4	6	13	21	19	53	67	91	128
2000.....	2,914	16	18	15	10	8	17	35	46	60	77	120
1999.....	2,714	49	24	5	10	15	17	27	43	59	82	86
Non-Hispanic 1/, male												
2003.....	1,252	7	11	2	2	15	7	12	22	30	50	72
2002.....	1,175	6	7	8	14	7	9	15	31	32	45	56
2001.....	1,286	15	11	1	4	4	11	9	30	37	41	66
2000.....	1,354	10	9	9	2	4	6	22	23	31	39	56
1999.....	1,253	28	9	3	4	7	12	9	31	32	38	35

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic 1/, female												
2003.....	1,441	5	8	4	3	2	14	8	18	34	42	63
2002.....	1,520	7	5	4	4	4	9	8	17	34	47	53
2001.....	1,575	8	7	3	2	9	10	10	23	30	50	62
2000.....	1,560	6	9	6	8	4	11	13	23	29	38	64
1999.....	1,461	21	15	2	6	8	5	18	12	27	44	51
Non-Hispanic White, both sexes												
2003.....	2,202	6	9	5	2	14	17	10	29	46	73	94
2002.....	2,188	6	7	6	10	8	13	16	33	45	62	75
2001.....	2,294	14	10	2	4	8	17	12	39	43	62	86
2000.....	2,378	12	8	9	8	6	11	20	36	47	56	89
1999.....	2,212	30	16	4	7	13	13	14	33	45	58	64
Non-Hispanic White, male												
2003.....	1,044	4	5	2	1	12	6	6	17	23	43	48
2002.....	966	3	5	4	8	6	7	11	23	25	35	41
2001.....	1,016	9	7	-	2	3	9	5	21	22	28	44
2000.....	1,135	9	6	5	2	3	4	12	19	25	33	42
1999.....	1,025	19	8	2	3	6	9	4	25	25	27	24
Non-Hispanic White, female												
2003.....	1,158	2	4	3	1	2	11	4	12	23	30	46
2002.....	1,222	3	2	2	2	2	6	5	10	20	27	34
2001.....	1,278	5	3	2	2	5	8	7	18	21	34	42
2000.....	1,243	3	2	4	6	3	7	8	17	22	23	47
1999.....	1,187	11	8	2	4	7	4	10	8	20	31	40

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Complications of medical and surgical care (Y40-Y84,Y88)												
Non-Hispanic Black, both sexes												
2003.....	425	5	9	-	3	2	4	7	9	17	16	33
2002.....	445	7	5	6	8	3	2	6	11	19	29	31
2001.....	515	9	7	1	2	3	4	7	14	22	26	39
2000.....	484	4	8	4	1	2	5	13	10	13	18	27
1999.....	442	18	8	1	3	2	4	10	8	13	22	20
Non-Hispanic Black, male												
2003.....	174	3	5	-	1	2	1	3	5	6	5	19
2002.....	182	3	2	4	6	1	1	4	5	6	10	13
2001.....	247	6	4	-	2	-	2	4	9	14	12	20
2000.....	192	1	2	2	-	1	1	9	4	6	6	12
1999.....	195	8	1	1	1	1	3	4	6	6	10	11
Non-Hispanic Black, female												
2003.....	251	2	4	-	2	-	3	4	4	11	11	14
2002.....	263	4	3	2	2	2	1	2	6	13	19	18
2001.....	268	3	3	1	-	3	2	3	5	8	14	19
2000.....	292	3	6	2	1	1	4	4	6	7	12	15
1999.....	247	10	7	-	2	1	1	6	2	7	12	9
Origin not stated 2/, both sexes												
2003.....	7	-	1	-	-	1	-	-	-	-	1	2
2002.....	11	-	-	-	-	-	-	-	-	-	-	1
2001.....	15	-	-	-	-	1	-	-	-	-	2	-
2000.....	9	-	-	-	-	-	-	-	-	-	1	-
1999.....	9	-	-	-	-	-	-	-	-	1	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

Origin not stated 2/, male

2003.....	5	-	1	-	-	1	-	-	-	-	1	1
2002.....	4	-	-	-	-	-	-	-	-	-	-	1
2001.....	6	-	-	-	-	1	-	-	-	-	1	-
2000.....	6	-	-	-	-	-	-	-	-	-	1	-
1999.....	3	-	-	-	-	-	-	-	-	-	1	-

Origin not stated 2/, female

2003.....	2	-	-	-	-	-	-	-	-	-	-	1
2002.....	7	-	-	-	-	-	-	-	-	-	-	-
2001.....	9	-	-	-	-	-	-	-	-	-	1	-
2000.....	3	-	-	-	-	-	-	-	-	-	-	-
1999.....	6	-	-	-	-	-	-	-	-	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/												
All origins, both sexes												
2003.....	20,687	-	-	-	5	49	75	167	429	1,144	2,428	3,472
2002 5/.....	20,218	1	2	-	3	33	72	160	441	1,191	2,445	3,321
2001 5/.....	20,114	2	-	-	2	41	65	168	477	1,329	2,417	3,422
2000 5/.....	19,643	3	1	-	4	33	62	175	475	1,355	2,482	3,203
1999 5/.....	19,469	1	-	-	2	25	75	147	494	1,416	2,413	3,149
All origins, male												
2003.....	15,630	-	-	-	3	37	64	139	333	799	1,735	2,627
2002 5/.....	15,272	-	1	-	2	28	58	133	325	829	1,742	2,485
2001 5/.....	15,149	2	-	-	-	31	57	124	342	940	1,725	2,593
2000 5/.....	14,993	2	-	-	4	27	49	146	358	986	1,822	2,477
1999 5/.....	14,894	1	-	-	1	20	60	121	338	994	1,822	2,436
All origins, female												
2003.....	5,057	-	-	-	2	12	11	28	96	345	693	845
2002 5/.....	4,946	1	1	-	1	5	14	27	116	362	703	836
2001 5/.....	4,965	-	-	-	2	10	8	44	135	389	692	829
2000 5/.....	4,650	1	1	-	-	6	13	29	117	369	660	726
1999 5/.....	4,575	-	-	-	1	5	15	26	156	422	591	713
Hispanic, both sexes												
2003.....	2,422	-	-	-	1	17	22	38	95	195	298	396
2002 5/.....	2,408	-	-	-	1	11	22	50	97	187	319	387
2001 5/.....	2,381	-	-	-	-	15	16	53	93	187	306	413
2000 5/.....	2,323	-	-	-	1	7	25	43	93	206	325	365
1999 5/.....	2,184	-	-	-	1	6	21	33	98	207	298	358

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/												
Hispanic, male												
2003.....	2,048	-	-	-	-	13	21	35	78	169	241	344
2002 5/.....	2,065	-	-	-	-	8	22	47	87	157	274	331
2001 5/.....	2,026	-	-	-	-	10	15	42	80	166	251	355
2000 5/.....	2,024	-	-	-	1	6	23	41	88	172	273	318
1999 5/.....	1,864	-	-	-	1	4	21	31	83	173	253	309
Hispanic, female												
2003.....	374	-	-	-	1	4	1	3	17	26	57	52
2002 5/.....	343	-	-	-	1	3	-	3	10	30	45	56
2001 5/.....	355	-	-	-	-	5	1	11	13	21	55	58
2000 5/.....	299	-	-	-	-	1	2	2	5	34	52	47
1999 5/.....	320	-	-	-	-	2	-	2	15	34	45	49
Non-Hispanic 1/, both sexes												
2003.....	18,160	-	-	-	4	31	53	127	333	946	2,120	3,062
2002 5/.....	17,661	1	2	-	2	22	49	109	341	991	2,111	2,905
2001 5/.....	17,593	2	-	-	2	26	49	115	380	1,137	2,097	2,989
2000 5/.....	17,177	3	1	-	3	26	36	130	379	1,139	2,141	2,809
1999 5/.....	17,143	1	-	-	1	19	52	113	392	1,197	2,094	2,770
Non-Hispanic 1/, male												
2003.....	13,490	-	-	-	3	23	43	102	254	628	1,486	2,271
2002 5/.....	13,078	-	1	-	2	20	35	85	235	664	1,456	2,128
2001 5/.....	13,009	2	-	-	-	21	42	82	259	770	1,461	2,223
2000 5/.....	12,843	2	-	-	3	21	25	103	268	804	1,534	2,135
1999 5/.....	12,905	1	-	-	-	16	37	89	251	811	1,552	2,108

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/												
Non-Hispanic 1/, female												
2003.....	4,670	-	-	-	1	8	10	25	79	318	634	791
2002 5/.....	4,583	1	1	-	-	2	14	24	106	327	655	777
2001 5/.....	4,584	-	-	-	2	5	7	33	121	367	636	766
2000 5/.....	4,334	1	1	-	-	5	11	27	111	335	607	674
1999 5/.....	4,238	-	-	-	1	3	15	24	141	386	542	662
Non-Hispanic White, both sexes												
2003.....	14,977	-	-	-	3	20	37	93	254	734	1,731	2,517
2002 5/.....	14,494	-	1	-	1	20	33	76	251	770	1,735	2,378
2001 5/.....	14,186	1	-	-	2	17	30	75	285	895	1,638	2,422
2000 5/.....	13,815	3	-	-	2	21	23	93	272	888	1,667	2,211
1999 5/.....	13,633	1	-	-	1	11	38	88	292	903	1,604	2,141
Non-Hispanic White, male												
2003.....	11,133	-	-	-	3	18	30	76	191	492	1,226	1,864
2002 5/.....	10,783	-	1	-	1	19	24	60	176	519	1,211	1,756
2001 5/.....	10,497	1	-	-	-	15	26	57	197	599	1,141	1,819
2000 5/.....	10,408	2	-	-	2	17	17	76	193	635	1,205	1,714
1999 5/.....	10,337	1	-	-	-	9	28	72	189	619	1,211	1,654
Non-Hispanic White, female												
2003.....	3,844	-	-	-	-	2	7	17	63	242	505	653
2002 5/.....	3,711	-	-	-	-	1	9	16	75	251	524	622
2001 5/.....	3,689	-	-	-	2	2	4	18	88	296	497	603
2000 5/.....	3,407	1	-	-	-	4	6	17	79	253	462	497
1999 5/.....	3,296	-	-	-	1	2	10	16	103	284	393	487

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Non-Hispanic Black, both sexes

2003.....	2,367	-	-	-	1	2	5	24	36	135	253	390
2002 5/.....	2,396	1	1	-	-	-	10	17	53	125	259	402
2001 5/.....	2,677	1	-	-	-	5	12	20	50	150	346	444
2000 5/.....	2,672	-	1	-	1	2	6	24	68	160	356	500
1999 5/.....	2,794	-	-	-	-	4	6	14	58	208	351	513

Non-Hispanic Black, male

2003.....	1,787	-	-	-	-	2	3	19	29	87	171	298
2002 5/.....	1,768	-	-	-	-	-	8	14	38	82	170	287
2001 5/.....	2,016	1	-	-	-	3	10	15	32	113	242	326
2000 5/.....	1,959	-	-	-	1	2	5	20	43	110	246	355
1999 5/.....	2,066	-	-	-	-	4	4	9	36	140	240	371

Non-Hispanic Black, female

2003.....	580	-	-	-	1	-	2	5	7	48	82	92
2002 5/.....	628	1	1	-	-	-	2	3	15	43	89	115
2001 5/.....	661	-	-	-	-	2	2	5	18	37	104	118
2000 5/.....	713	-	1	-	-	-	1	4	25	50	110	145
1999 5/.....	728	-	-	-	-	-	2	5	22	68	111	142

Origin not stated 2/, both sexes

2003.....	105	-	-	-	-	1	-	2	1	3	10	14
2002 5/.....	149	-	-	-	-	-	1	1	3	13	15	29
2001 5/.....	140	-	-	-	-	-	-	-	4	5	14	20
2000 5/.....	143	-	-	-	-	-	1	2	3	10	16	29
1999 5/.....	142	-	-	-	-	-	2	1	4	12	21	21

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Origin not stated 2/, male

2003.....	92	-	-	-	-	1	-	2	1	2	8	12
2002 5/.....	129	-	-	-	-	-	1	1	3	8	12	26
2001 5/.....	114	-	-	-	-	-	-	-	3	4	13	15
2000 5/.....	126	-	-	-	-	-	1	2	2	10	15	24
1999 5/.....	125	-	-	-	-	-	2	1	4	10	17	19

Origin not stated 2/, female

2003.....	13	-	-	-	-	-	-	-	-	1	2	2
2002 5/.....	20	-	-	-	-	-	-	-	-	5	3	3
2001 5/.....	26	-	-	-	-	-	-	-	1	1	1	5
2000 5/.....	17	-	-	-	-	-	-	-	1	-	1	5
1999 5/.....	17	-	-	-	-	-	-	-	-	2	4	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
All origins, both sexes												
2003.....	28,723	25	35	9	32	669	1,950	2,086	2,759	3,918	5,240	4,946
2002 5/.....	26,040	29	30	7	35	625	1,590	1,850	2,599	3,837	5,021	4,451
2001 5/.....	21,705	22	26	8	37	539	1,274	1,552	2,187	3,448	4,221	3,660
2000 5/.....	19,720	18	15	9	35	452	1,115	1,450	2,077	3,270	3,936	3,171
1999 5/.....	19,128	21	25	7	23	367	984	1,452	2,127	3,242	3,853	3,042
All origins, male												
2003.....	18,426	17	21	2	18	477	1,478	1,508	1,853	2,549	3,324	3,131
2002 5/.....	16,734	20	9	6	18	447	1,175	1,346	1,793	2,534	3,194	2,765
2001 5/.....	14,253	14	21	3	19	386	964	1,142	1,498	2,302	2,777	2,440
2000 5/.....	13,137	12	13	6	19	318	848	1,057	1,458	2,242	2,648	2,170
1999 5/.....	12,885	14	12	1	15	251	730	1,067	1,512	2,238	2,669	2,121
All origins, female												
2003.....	10,297	8	14	7	14	192	472	578	906	1,369	1,916	1,815
2002 5/.....	9,306	9	21	1	17	178	415	504	806	1,303	1,827	1,686
2001 5/.....	7,452	8	5	5	18	153	310	410	689	1,146	1,444	1,220
2000 5/.....	6,583	6	2	3	16	134	267	393	619	1,028	1,288	1,001
1999 5/.....	6,243	7	13	6	8	116	254	385	615	1,004	1,184	921
Hispanic, both sexes												
2003.....	2,358	4	5	1	6	68	175	227	306	380	409	356
2002 5/.....	2,137	5	4	-	4	58	159	188	268	358	419	330
2001 5/.....	1,731	-	4	-	5	54	108	176	232	330	314	231
2000 5/.....	1,700	2	2	-	4	40	108	170	279	345	306	215
1999 5/.....	1,965	7	4	1	2	50	122	211	303	400	341	254

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/												
Hispanic, male												
2003.....	1,800	3	3	-	1	51	143	184	237	297	309	267
2002 5/.....	1,647	4	-	-	1	44	128	160	211	272	329	247
2001 5/.....	1,335	-	3	-	1	39	80	146	174	259	245	178
2000 5/.....	1,348	2	2	-	-	30	91	130	237	264	257	178
1999 5/.....	1,605	4	4	-	2	34	108	174	257	320	286	217
Hispanic, female												
2003.....	558	1	2	1	5	17	32	43	69	83	100	89
2002 5/.....	490	1	4	-	3	14	31	28	57	86	90	83
2001 5/.....	396	-	1	-	4	15	28	30	58	71	69	53
2000 5/.....	352	-	-	-	4	10	17	40	42	81	49	37
1999 5/.....	360	3	-	1	-	16	14	37	46	80	55	37
Non-Hispanic 1/, both sexes												
2003.....	26,199	21	30	8	26	599	1,772	1,846	2,441	3,515	4,806	4,559
2002 5/.....	23,756	24	26	7	31	565	1,424	1,654	2,319	3,456	4,573	4,098
2001 5/.....	19,799	22	22	8	32	483	1,159	1,372	1,941	3,091	3,878	3,396
2000 5/.....	17,835	16	12	9	30	410	1,000	1,271	1,782	2,908	3,601	2,911
1999 5/.....	16,966	14	21	6	21	315	859	1,233	1,813	2,816	3,463	2,755
Non-Hispanic 1/, male												
2003.....	16,497	14	18	2	17	425	1,333	1,315	1,606	2,232	2,999	2,845
2002 5/.....	14,978	16	9	6	17	403	1,042	1,179	1,572	2,246	2,843	2,500
2001 5/.....	12,778	14	18	3	18	346	878	992	1,312	2,020	2,512	2,236
2000 5/.....	11,656	10	10	6	18	287	751	923	1,210	1,965	2,369	1,963
1999 5/.....	11,136	10	8	1	13	216	619	886	1,246	1,899	2,350	1,879

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Non-Hispanic 1/, female

2003.....	9,702	7	12	6	9	174	439	531	835	1,283	1,807	1,714
2002 5/.....	8,778	8	17	1	14	162	382	475	747	1,210	1,730	1,598
2001 5/.....	7,021	8	4	5	14	137	281	380	629	1,071	1,366	1,160
2000 5/.....	6,179	6	2	3	12	123	249	348	572	943	1,232	948
1999 5/.....	5,830	4	13	5	8	99	240	347	567	917	1,113	876

Non-Hispanic White, both sexes

2003.....	22,245	12	17	6	19	562	1,646	1,638	2,101	2,944	4,050	3,766
2002 5/.....	19,949	9	14	4	28	514	1,305	1,444	1,985	2,894	3,828	3,330
2001 5/.....	16,367	10	13	6	27	436	1,048	1,191	1,644	2,564	3,166	2,666
2000 5/.....	14,585	9	5	6	27	376	890	1,083	1,471	2,386	2,878	2,266
1999 5/.....	13,644	6	10	3	19	282	737	1,020	1,473	2,288	2,713	2,106

Non-Hispanic White, male

2003.....	13,959	11	11	2	14	406	1,245	1,186	1,392	1,878	2,536	2,330
2002 5/.....	12,478	7	3	3	16	368	957	1,030	1,363	1,883	2,393	2,018
2001 5/.....	10,465	4	12	3	15	324	788	859	1,133	1,688	2,062	1,719
2000 5/.....	9,439	5	3	5	16	267	677	784	1,006	1,633	1,892	1,499
1999 5/.....	8,831	5	4	-	12	192	535	727	1,023	1,547	1,854	1,404

Non-Hispanic White, female

2003.....	8,286	1	6	4	5	156	401	452	709	1,066	1,514	1,436
2002 5/.....	7,471	2	11	1	12	146	348	414	622	1,011	1,435	1,312
2001 5/.....	5,902	6	1	3	12	112	260	332	511	876	1,104	947
2000 5/.....	5,146	4	2	1	11	109	213	299	465	753	986	767
1999 5/.....	4,813	1	6	3	7	90	202	293	450	741	859	702

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Non-Hispanic Black, both sexes

2003.....	3,466	9	13	2	6	23	89	148	283	501	680	737
2002 5/.....	3,404	13	10	2	3	31	96	170	281	490	691	713
2001 5/.....	3,099	11	8	2	3	27	85	147	257	478	662	681
2000 5/.....	2,977	6	5	3	1	27	86	160	280	479	664	605
1999 5/.....	3,030	8	11	2	2	23	100	183	305	474	696	623

Non-Hispanic Black, male

2003.....	2,256	3	7	-	3	10	67	94	176	318	418	478
2002 5/.....	2,264	8	6	2	1	22	68	120	180	322	418	455
2001 5/.....	2,113	9	5	-	2	14	71	112	157	300	421	488
2000 5/.....	2,050	4	5	1	-	16	60	120	183	309	440	440
1999 5/.....	2,134	5	4	1	1	19	74	140	201	321	465	458

Non-Hispanic Black, female

2003.....	1,210	6	6	2	3	13	22	54	107	183	262	259
2002 5/.....	1,140	5	4	-	2	9	28	50	101	168	273	258
2001 5/.....	986	2	3	2	1	13	14	35	100	178	241	193
2000 5/.....	927	2	-	2	1	11	26	40	97	170	224	165
1999 5/.....	896	3	7	1	1	4	26	43	104	153	231	165

Origin not stated 2/, both sexes

2003.....	166	-	-	-	-	2	3	13	12	23	25	31
2002 5/.....	147	-	-	-	-	2	7	8	12	23	29	23
2001 5/.....	175	-	-	-	-	2	7	4	14	27	29	33
2000 5/.....	185	-	1	-	1	2	7	9	16	17	29	45
1999 5/.....	197	-	-	-	-	2	3	8	11	26	49	33

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Origin not stated 2/, male

2003.....	129	-	-	-	-	1	2	9	10	20	16	19
2002 5/.....	109	-	-	-	-	-	5	7	10	16	22	18
2001 5/.....	140	-	-	-	-	1	6	4	12	23	20	26
2000 5/.....	133	-	1	-	1	1	6	4	11	13	22	29
1999 5/.....	144	-	-	-	-	1	3	7	9	19	33	25

Origin not stated 2/, female

2003.....	37	-	-	-	-	1	1	4	2	3	9	12
2002 5/.....	38	-	-	-	-	2	2	1	2	7	7	5
2001 5/.....	35	-	-	-	-	1	1	-	2	4	9	7
2000 5/.....	52	-	-	-	-	1	1	5	5	4	7	16
1999 5/.....	53	-	-	-	-	1	-	1	2	7	16	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/												
All origins, both sexes												
2003.....	30,136	8	48	63	261	2,469	4,377	3,336	2,852	2,476	2,642	2,475
2002.....	30,242	10	61	71	277	2,474	4,306	3,380	2,760	2,712	2,734	2,358
2001.....	29,573	11	70	79	254	2,523	4,164	3,350	2,787	2,723	2,565	2,203
2000.....	28,663	13	46	70	307	2,606	3,969	3,165	2,624	2,718	2,640	2,146
1999.....	28,874	8	65	80	336	2,896	3,899	3,248	2,727	2,812	2,422	2,096
All origins, male												
2003.....	26,124	7	28	40	216	2,231	3,955	2,999	2,473	2,065	2,157	2,025
2002.....	26,098	4	38	38	204	2,209	3,887	2,994	2,353	2,257	2,246	1,927
2001.....	25,480	7	39	51	198	2,270	3,796	2,963	2,384	2,216	2,096	1,828
2000.....	24,582	9	28	46	256	2,323	3,583	2,796	2,207	2,207	2,156	1,770
1999.....	24,700	5	36	50	258	2,560	3,496	2,848	2,308	2,279	1,950	1,722
All origins, female												
2003.....	4,012	1	20	23	45	238	422	337	379	411	485	450
2002.....	4,144	6	23	33	73	265	419	386	407	455	488	431
2001.....	4,093	4	31	28	56	253	368	387	403	507	469	375
2000.....	4,081	4	18	24	51	283	386	369	417	511	484	376
1999.....	4,174	3	29	30	78	336	403	400	419	533	472	374
Hispanic, both sexes												
2003.....	3,319	1	9	10	42	498	848	599	386	278	200	166
2002.....	3,143	2	15	17	39	514	746	572	348	267	192	127
2001.....	3,087	4	13	6	34	468	765	575	351	264	204	127
2000.....	2,891	1	7	10	45	511	673	469	319	213	201	156
1999.....	2,878	2	6	11	58	533	656	466	304	251	176	151

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/												
Hispanic, male												
2003.....	2,998	1	5	4	34	454	780	559	355	240	171	142
2002.....	2,834	-	9	12	26	474	699	524	320	231	159	107
2001.....	2,774	3	6	5	26	425	707	526	315	231	177	106
2000.....	2,582	-	4	7	39	471	625	429	282	175	176	127
1999.....	2,549	2	4	6	44	481	597	427	272	216	148	118
Hispanic, female												
2003.....	321	-	4	6	8	44	68	40	31	38	29	24
2002.....	309	2	6	5	13	40	47	48	28	36	33	20
2001.....	313	1	7	1	8	43	58	49	36	33	27	21
2000.....	309	1	3	3	6	40	48	40	37	38	25	29
1999.....	329	-	2	5	14	52	59	39	32	35	28	33
Non-Hispanic 1/, both sexes												
2003.....	26,710	7	39	53	219	1,959	3,509	2,726	2,457	2,187	2,430	2,302
2002.....	26,944	8	46	54	236	1,939	3,537	2,789	2,398	2,431	2,532	2,219
2001.....	26,341	7	57	72	220	2,040	3,369	2,756	2,426	2,451	2,351	2,070
2000.....	25,637	12	39	58	262	2,081	3,274	2,686	2,294	2,489	2,426	1,980
1999.....	25,877	6	59	69	278	2,351	3,236	2,767	2,407	2,547	2,235	1,940
Non-Hispanic 1/, male												
2003.....	23,036	6	23	36	182	1,766	3,159	2,429	2,111	1,817	1,977	1,878
2002.....	23,127	4	29	26	176	1,716	3,166	2,455	2,022	2,013	2,080	1,809
2001.....	22,573	4	33	45	172	1,831	3,063	2,418	2,061	1,979	1,910	1,718
2000.....	21,881	9	24	38	217	1,840	2,937	2,358	1,916	2,017	1,967	1,634
1999.....	22,050	3	32	44	214	2,070	2,894	2,408	2,024	2,051	1,791	1,600

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/												
Non-Hispanic 1/, female												
2003.....	3,674	1	16	17	37	193	350	297	346	370	453	424
2002.....	3,817	4	17	28	60	223	371	334	376	418	452	410
2001.....	3,768	3	24	27	48	209	306	338	365	472	441	352
2000.....	3,756	3	15	20	45	241	337	328	378	472	459	346
1999.....	3,827	3	27	25	64	281	342	359	383	496	444	340
Non-Hispanic White, both sexes												
2003.....	18,457	1	15	23	118	856	1,433	1,167	1,385	1,512	1,892	1,918
2002.....	18,762	2	28	28	135	876	1,480	1,276	1,363	1,662	2,000	1,853
2001.....	18,676	4	30	55	148	962	1,411	1,355	1,492	1,746	1,855	1,757
2000.....	18,042	5	21	38	181	958	1,419	1,298	1,404	1,786	1,908	1,630
1999.....	18,260	2	27	32	193	1,092	1,412	1,412	1,537	1,863	1,761	1,594
Non-Hispanic White, male												
2003.....	15,670	-	7	15	104	747	1,262	988	1,153	1,222	1,532	1,547
2002.....	15,881	2	16	12	102	754	1,284	1,058	1,111	1,350	1,648	1,497
2001.....	15,760	2	19	34	118	824	1,235	1,156	1,227	1,399	1,511	1,458
2000.....	15,160	2	11	24	151	818	1,232	1,088	1,144	1,414	1,551	1,348
1999.....	15,384	2	15	19	149	943	1,222	1,192	1,261	1,482	1,411	1,315
Non-Hispanic White, female												
2003.....	2,787	1	8	8	14	109	171	179	232	290	360	371
2002.....	2,881	-	12	16	33	122	196	218	252	312	352	356
2001.....	2,916	2	11	21	30	138	176	199	265	347	344	299
2000.....	2,882	3	10	14	30	140	187	210	260	372	357	282
1999.....	2,876	-	12	13	44	149	190	220	276	381	350	279

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Non-Hispanic Black, both sexes

2003.....	7,566	6	23	27	89	1,021	1,947	1,473	987	607	481	326
2002.....	7,494	6	17	18	84	979	1,925	1,428	965	699	473	326
2001.....	7,063	2	23	15	63	1,015	1,857	1,311	861	656	436	272
2000.....	6,958	7	13	18	73	1,041	1,757	1,287	834	633	460	306
1999.....	6,933	2	26	31	74	1,163	1,719	1,246	788	610	427	305

Non-Hispanic Black, male

2003.....	6,794	6	15	19	67	954	1,785	1,370	883	537	398	286
2002.....	6,681	2	12	11	63	886	1,771	1,324	858	603	388	282
2001.....	6,323	1	13	9	51	947	1,743	1,188	777	540	354	227
2000.....	6,193	7	8	13	61	954	1,620	1,182	728	548	372	251
1999.....	6,114	1	12	22	56	1,041	1,582	1,128	698	509	345	254

Non-Hispanic Black, female

2003.....	772	-	8	8	22	67	162	103	104	70	83	40
2002.....	813	4	5	7	21	93	154	104	107	96	85	44
2001.....	740	1	10	6	12	68	114	123	84	116	82	45
2000.....	765	-	5	5	12	87	137	105	106	85	88	55
1999.....	819	1	14	9	18	122	137	118	90	101	82	51

Origin not stated 2/, both sexes

2003.....	107	-	-	-	-	12	20	11	9	11	12	7
2002.....	155	-	-	-	2	21	23	19	14	14	10	12
2001.....	145	-	-	1	-	15	30	19	10	8	10	6
2000.....	135	-	-	2	-	14	22	10	11	16	13	10
1999.....	119	-	-	-	-	12	7	15	16	14	11	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Origin not stated 2/, male

2003.....	90	-	-	-	-	11	16	11	7	8	9	5
2002.....	137	-	-	-	2	19	22	15	11	13	7	11
2001.....	133	-	-	1	-	14	26	19	8	6	9	4
2000.....	119	-	-	1	-	12	21	9	9	15	13	9
1999.....	101	-	-	-	-	9	5	13	12	12	11	4

Origin not stated 2/, female

2003.....	17	-	-	-	-	1	4	-	2	3	3	2
2002.....	18	-	-	-	-	2	1	4	3	1	3	1
2001.....	12	-	-	-	-	1	4	-	2	2	1	2
2000.....	16	-	-	1	-	2	1	1	2	1	-	1
1999.....	18	-	-	-	-	3	2	2	4	2	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>All origins, both sexes</b>												
2003.....	98,608	119,339	143,180	174,817	238,680	325,185	377,839	346,847	231,403	90,271	19,331	342
2002.....	96,320	115,441	137,901	175,591	247,399	330,140	377,514	345,385	228,587	88,282	18,822	357
2001.....	94,381	109,523	134,616	178,232	252,728	333,677	368,252	339,775	221,314	85,954	18,488	422
2000.....	89,509	106,751	134,095	181,739	259,470	337,700	362,745	335,813	219,283	84,641	18,434	356
1999.....	84,854	104,884	134,095	186,695	265,905	338,349	360,241	330,903	213,991	83,334	17,913	356
<b>All origins, male</b>												
2003.....	61,543	72,188	84,273	100,315	131,106	167,354	174,978	135,177	70,162	19,872	3,001	271
2002.....	60,021	69,830	81,533	100,590	136,431	169,625	173,879	134,637	68,817	19,522	2,930	282
2001.....	58,281	65,801	79,157	102,110	139,471	171,311	169,431	130,405	65,527	18,732	2,869	331
2000.....	55,277	64,425	78,896	103,935	143,473	173,327	166,892	128,877	64,439	18,552	2,874	289
1999.....	52,356	63,441	79,283	107,529	147,391	174,853	166,117	126,531	62,511	18,111	2,836	277
<b>All origins, female</b>												
2003.....	37,065	47,151	58,907	74,502	107,574	157,831	202,861	211,670	161,241	70,399	16,330	71
2002.....	36,299	45,611	56,368	75,001	110,968	160,515	203,635	210,748	159,770	68,760	15,892	75
2001.....	36,100	43,722	55,459	76,122	113,257	162,366	198,821	209,370	155,787	67,222	15,619	91
2000.....	34,232	42,326	55,199	77,804	115,997	164,373	195,853	206,936	154,844	66,089	15,560	67
1999.....	32,498	41,443	54,812	79,166	118,514	163,496	194,124	204,372	151,480	65,223	15,077	79
<b>Hispanic, both sexes</b>												
2003.....	6,599	7,177	7,736	9,230	11,366	13,406	12,813	9,960	6,835	2,703	631	47
2002.....	6,218	6,634	7,581	9,060	11,416	12,797	12,041	9,591	6,460	2,668	546	58
2001.....	5,965	6,114	7,223	8,910	11,262	12,560	11,077	9,601	6,177	2,582	547	67
2000.....	5,366	5,876	6,857	8,546	10,874	11,578	10,373	9,170	5,853	2,334	461	54
1999.....	5,109	5,495	6,824	8,843	10,358	11,081	9,831	8,953	5,662	2,285	458	66

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## All causes

## Hispanic, male

2003.....	4,326	4,463	4,659	5,235	6,230	6,788	6,116	3,926	2,276	804	159	39
2002.....	4,024	4,247	4,515	5,214	6,262	6,654	5,637	3,846	2,226	768	137	55
2001.....	3,822	3,760	4,266	5,129	6,102	6,535	5,224	3,821	2,051	754	141	58
2000.....	3,451	3,656	4,104	4,838	6,004	6,069	4,904	3,617	2,035	725	116	51
1999.....	3,284	3,409	4,081	5,026	5,757	5,860	4,457	3,483	1,960	703	107	60

## Hispanic, female

2003.....	2,273	2,714	3,077	3,995	5,136	6,618	6,697	6,034	4,559	1,899	472	8
2002.....	2,194	2,387	3,066	3,846	5,154	6,143	6,404	5,745	4,234	1,900	409	3
2001.....	2,143	2,354	2,957	3,781	5,160	6,025	5,853	5,780	4,126	1,828	406	9
2000.....	1,915	2,220	2,753	3,708	4,870	5,509	5,469	5,553	3,818	1,609	345	3
1999.....	1,825	2,086	2,743	3,817	4,601	5,221	5,374	5,470	3,702	1,582	351	6

## Non-Hispanic 1/, both sexes

2003.....	91,594	111,711	134,954	165,087	226,691	311,065	364,244	336,218	224,096	87,370	18,666	147
2002.....	89,665	108,370	129,815	165,934	235,248	316,397	364,481	334,920	221,569	85,363	18,226	169
2001.....	87,989	102,952	126,860	168,796	240,704	320,213	356,299	329,357	214,629	83,144	17,890	173
2000.....	83,687	100,433	126,685	172,601	247,768	325,151	351,481	325,765	212,888	82,088	17,932	140
1999.....	79,321	98,934	126,725	177,240	254,712	326,279	349,451	321,050	207,750	80,833	17,402	155

## Non-Hispanic 1/, male

2003.....	56,922	67,414	79,282	94,743	124,497	160,166	168,474	130,988	67,726	19,027	2,834	118
2002.....	55,690	65,283	76,683	94,990	129,716	162,434	167,757	130,450	66,440	18,694	2,783	134
2001.....	54,159	61,716	74,498	96,644	132,895	164,283	163,796	126,292	63,325	17,924	2,715	137
2000.....	51,494	60,462	74,428	98,692	136,956	166,709	161,575	124,927	62,247	17,768	2,746	107
1999.....	48,754	59,720	74,820	102,119	141,091	168,453	161,229	122,676	60,373	17,354	2,725	113

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes												
Non-Hispanic 1/, female												
2003.....	34,672	44,297	55,672	70,344	102,194	150,899	195,770	205,230	156,370	68,343	15,832	29
2002.....	33,975	43,087	53,132	70,944	105,532	153,963	196,724	204,470	155,129	66,669	15,443	35
2001.....	33,830	41,236	52,362	72,152	107,809	155,930	192,503	203,065	151,304	65,220	15,175	36
2000.....	32,193	39,971	52,257	73,909	110,812	158,442	189,906	200,838	150,641	64,320	15,186	33
1999.....	30,567	39,214	51,905	75,121	113,621	157,826	188,222	198,374	147,377	63,479	14,677	42
Non-Hispanic White, both sexes												
2003.....	68,679	88,382	109,216	136,254	193,316	273,142	327,421	307,581	205,431	79,709	16,122	121
2002.....	67,768	85,916	104,927	137,191	201,770	278,144	328,287	306,287	203,219	77,698	15,859	122
2001.....	66,935	81,505	102,604	139,503	206,886	282,018	321,125	301,010	197,069	75,705	15,485	113
2000.....	63,959	79,771	102,577	143,765	213,104	287,117	317,541	297,730	195,687	74,671	15,669	89
1999.....	60,834	78,170	102,623	148,130	219,360	288,018	316,326	292,962	191,334	73,705	15,326	91
Non-Hispanic White, male												
2003.....	43,287	53,763	64,620	78,944	107,168	141,742	152,032	120,507	62,124	17,126	2,391	96
2002.....	42,622	52,169	62,366	79,363	112,481	143,909	151,854	119,953	60,743	16,825	2,313	98
2001.....	41,686	49,212	60,781	80,805	115,353	145,725	148,288	115,799	57,884	16,125	2,276	92
2000.....	39,655	48,255	60,776	83,126	118,912	148,070	146,772	114,484	56,864	15,857	2,332	70
1999.....	37,703	47,570	61,122	86,299	122,856	149,671	146,857	111,859	55,125	15,589	2,365	66
Non-Hispanic White, female												
2003.....	25,392	34,619	44,596	57,310	86,148	131,400	175,389	187,074	143,307	62,583	13,731	25
2002.....	25,146	33,747	42,561	57,828	89,289	134,235	176,433	186,334	142,476	60,873	13,546	24
2001.....	25,249	32,293	41,823	58,698	91,533	136,293	172,837	185,211	139,185	59,580	13,209	21
2000.....	24,304	31,516	41,801	60,639	94,192	139,047	170,769	183,246	138,823	58,814	13,337	19
1999.....	23,131	30,600	41,501	61,831	96,504	138,347	169,469	181,103	136,209	58,116	12,961	25

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## All causes

## Non-Hispanic Black, both sexes

2003.....	20,086	20,251	22,099	24,501	28,082	31,540	30,230	23,345	15,434	6,458	2,189	20
2002.....	19,209	19,483	21,395	24,508	28,240	32,076	30,107	23,674	15,316	6,521	2,054	39
2001.....	18,441	18,646	20,877	25,103	28,700	32,243	29,349	23,592	14,846	6,423	2,107	50
2000.....	17,366	17,965	20,902	24,831	29,814	32,219	28,655	23,696	14,579	6,362	2,023	49
1999.....	16,246	18,176	20,887	25,217	30,350	32,618	28,016	23,831	13,993	6,229	1,842	58

## Non-Hispanic Black, male

2003.....	11,995	11,905	12,563	13,395	14,562	15,124	13,173	8,119	4,264	1,433	340	17
2002.....	11,467	11,414	12,292	13,292	14,567	15,246	12,891	8,195	4,407	1,441	379	29
2001.....	10,953	10,845	11,755	13,617	14,829	15,429	12,654	8,313	4,249	1,415	375	40
2000.....	10,422	10,649	11,753	13,355	15,390	15,491	12,224	8,341	4,218	1,523	349	35
1999.....	9,760	10,627	11,866	13,720	15,539	15,706	11,846	8,736	4,134	1,451	307	42

## Non-Hispanic Black, female

2003.....	8,091	8,346	9,536	11,106	13,520	16,416	17,057	15,226	11,170	5,025	1,849	3
2002.....	7,742	8,069	9,103	11,216	13,673	16,830	17,216	15,479	10,909	5,080	1,675	10
2001.....	7,488	7,801	9,122	11,486	13,871	16,814	16,695	15,279	10,597	5,008	1,732	10
2000.....	6,944	7,316	9,149	11,476	14,424	16,728	16,431	15,355	10,361	4,839	1,674	14
1999.....	6,486	7,549	9,021	11,497	14,811	16,912	16,170	15,095	9,859	4,778	1,535	16

## Origin not stated 2/, both sexes

2003.....	415	451	490	500	623	714	782	669	472	198	34	148
2002.....	437	437	505	597	735	946	992	874	558	251	50	130
2001.....	427	457	533	526	762	904	876	817	508	228	51	182
2000.....	456	442	553	592	828	971	891	878	542	219	41	162
1999.....	424	455	546	612	835	989	959	900	579	216	53	135

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## All causes

## Origin not stated 2/, male

2003.....	295	311	332	337	379	400	388	263	160	41	8	114
2002.....	307	300	335	386	453	537	485	341	151	60	10	93
2001.....	300	325	393	337	474	493	411	292	151	54	13	136
2000.....	332	307	364	405	513	549	413	333	157	59	12	131
1999.....	318	312	382	384	543	540	431	372	178	54	4	104

## Origin not stated 2/, female

2003.....	120	140	158	163	244	314	394	406	312	157	26	34
2002.....	130	137	170	211	282	409	507	533	407	191	40	37
2001.....	127	132	140	189	288	411	465	525	357	174	38	46
2000.....	124	135	189	187	315	422	478	545	385	160	29	31
1999.....	106	143	164	228	292	449	528	528	401	162	49	31

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Salmonella infections (A01-A02)

All origins, both sexes

2003.....	1	4	2	2	6	8	2	5	3	1	-	-
2002.....	2	-	-	3	1	3	5	1	-	-	-	-
2001.....	2	2	1	4	3	5	9	6	3	-	-	-
2000.....	3	2	-	1	6	3	2	1	2	-	-	-
1999.....	2	1	2	2	5	8	4	4	2	1	-	-

All origins, male

2003.....	1	3	1	2	2	4	-	4	1	1	-	-
2002.....	1	-	-	1	1	1	4	1	-	-	-	-
2001.....	2	2	-	3	3	3	4	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	-	-	-
1999.....	2	-	2	2	3	2	3	1	1	-	-	-

All origins, female

2003.....	-	1	1	-	4	4	2	1	2	-	-	-
2002.....	1	-	-	2	-	2	1	-	-	-	-	-
2001.....	-	-	1	1	-	2	5	5	3	-	-	-
2000.....	2	-	-	-	2	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	6	1	3	1	1	-	-

Hispanic, both sexes

2003.....	1	-	-	-	2	1	-	-	-	-	-	-
2002.....	1	-	-	2	-	-	-	-	-	-	-	-
2001.....	1	1	-	1	-	-	2	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	1	-	1	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Salmonella infections (A01-A02)

Hispanic, male

2003.....	1	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	1	1	-	1	-	-	2	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	1	-	-	1	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	1	1	-	-	-	-	-	-
2002.....	1	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	-	4	2	2	4	7	2	5	3	1	-	-
2002.....	1	-	-	1	1	3	5	1	-	-	-	-
2001.....	1	1	1	3	3	5	7	6	3	-	-	-
2000.....	3	2	-	1	6	3	2	1	2	-	-	-
1999.....	2	1	1	1	5	7	3	4	2	1	-	-

Non-Hispanic 1/, male

2003.....	-	3	1	2	1	4	-	4	1	1	-	-
2002.....	1	-	-	-	1	1	4	1	-	-	-	-
2001.....	1	1	-	2	3	3	2	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	4	-	-
1999.....	2	-	1	1	3	2	2	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Salmonella infections (A01-A02)

Non-Hispanic 1/, female

2003.....	-	1	1	-	3	3	2	1	2	-	-	-
2002.....	-	-	-	1	-	2	1	-	-	-	-	-
2001.....	-	-	1	1	-	2	5	5	3	-	-	-
2000.....	2	-	-	-	2	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	5	1	3	1	1	-	-

Non-Hispanic White, both sexes

2003.....	-	3	2	2	2	4	2	4	3	1	-	-
2002.....	1	-	-	1	-	2	5	1	-	-	-	-
2001.....	1	1	1	1	2	5	5	6	2	-	-	-
2000.....	3	2	-	1	5	3	2	1	2	-	-	-
1999.....	2	1	-	-	4	6	2	4	2	1	-	-

Non-Hispanic White, male

2003.....	-	2	1	2	1	2	-	3	1	1	-	-
2002.....	1	-	-	-	-	1	4	1	-	-	-	-
2001.....	1	1	-	1	2	3	2	1	-	-	-	-
2000.....	1	2	-	1	4	1	-	-	1	-	-	-
1999.....	2	-	-	-	2	2	1	1	1	-	-	-

Non-Hispanic White, female

2003.....	-	1	1	-	1	2	2	1	2	-	-	-
2002.....	-	-	-	1	-	1	1	-	-	-	-	-
2001.....	-	-	1	-	-	2	3	5	2	-	-	-
2000.....	2	-	-	-	1	2	2	1	1	-	-	-
1999.....	-	1	-	-	2	4	1	3	1	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Shigellosis and amebiasis (A03,A06)

##### All origins, both sexes

2003.....	-	1	1	-	1	1	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	2	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	1	-	2	5	-	-	-	1	-
1999.....	-	-	1	-	1	1	-	1	1	2	-	-

##### All origins, male

2003.....	-	-	1	-	1	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	1	-	-	-

##### All origins, female

2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	1	-	1	3	-	-	-	1	-
1999.....	-	-	-	-	-	1	-	1	-	2	-	-

##### Hispanic, both sexes

2003.....	-	1	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Shigellosis and amebiasis (A03,A06)

##### Hispanic, male

2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

##### Hispanic, female

2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

##### Non-Hispanic 1/, both sexes

2003.....	-	-	1	-	-	1	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	2	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	1	-	2	5	-	-	-	1	-
1999.....	-	-	1	-	1	-	-	1	1	2	-	-

##### Non-Hispanic 1/, male

2003.....	-	-	1	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Shigellosis and amebiasis (A03,A06)

##### Non-Hispanic 1/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	1	-	1	3	-	-	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	2	-	-

##### Non-Hispanic White, both sexes

2003.....	-	-	1	-	-	1	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	2	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	-	-	2	5	-	-	-	1	-
1999.....	-	-	1	-	1	-	-	1	1	2	-	-

##### Non-Hispanic White, male

2003.....	-	-	1	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	2	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	2	-	-	-	-	-
1999.....	-	-	1	-	1	-	-	-	1	-	-	-

##### Non-Hispanic White, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	3	-	-	-	1	-
1999.....	-	-	-	-	-	-	-	1	-	2	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

All origins, both sexes

2003.....	43	70	106	161	283	524	668	647	383	120	17	-
2002.....	31	63	79	133	229	374	515	509	318	114	23	-
2001.....	33	28	53	73	162	237	329	293	187	75	10	-
2000.....	18	33	37	85	137	202	260	267	168	53	11	-
1999.....	12	15	34	67	109	168	222	211	146	58	14	-

All origins, male

2003.....	21	35	47	68	117	213	265	198	86	23	2	-
2002.....	17	29	42	67	118	156	202	160	91	27	5	-
2001.....	11	14	21	32	76	107	120	93	50	14	-	-
2000.....	9	20	19	44	61	88	107	81	48	9	2	-
1999.....	6	10	16	28	46	80	89	54	44	13	2	-

All origins, female

2003.....	22	35	59	93	166	311	403	449	297	97	15	-
2002.....	14	34	37	66	111	218	313	349	227	87	18	-
2001.....	22	14	32	41	86	130	209	200	137	61	10	-
2000.....	9	13	18	41	76	114	153	186	120	44	9	-
1999.....	6	5	18	39	63	88	133	157	102	45	12	-

Hispanic, both sexes

2003.....	2	1	9	7	9	18	13	18	8	2	-	-
2002.....	3	-	6	4	6	15	16	7	6	2	1	-
2001.....	2	2	2	3	5	8	8	8	7	2	-	-
2000.....	-	5	3	5	4	6	5	6	3	1	1	-
1999.....	2	2	3	4	3	3	1	1	4	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Hispanic, male

2003.....	2	1	6	5	2	9	4	7	5	1	-	-
2002.....	2	-	4	2	3	6	6	3	-	-	1	-
2001.....	-	1	1	-	1	4	1	3	3	-	-	-
2000.....	-	3	2	3	3	3	1	2	1	-	1	-
1999.....	1	-	2	2	1	2	-	1	1	1	-	-

Hispanic, female

2003.....	-	-	3	2	7	9	9	11	3	1	-	-
2002.....	1	-	2	2	3	9	10	4	6	2	-	-
2001.....	2	1	1	3	4	4	7	5	4	2	-	-
2000.....	-	2	1	2	1	3	4	4	2	1	-	-
1999.....	1	2	1	2	2	1	1	-	3	1	-	-

Non-Hispanic 1/, both sexes

2003.....	40	68	96	154	274	505	655	629	374	118	17	-
2002.....	27	63	72	129	222	358	499	500	312	112	22	-
2001.....	31	26	51	69	156	228	318	284	179	72	10	-
2000.....	18	28	34	80	131	196	253	260	165	52	10	-
1999.....	10	13	31	63	106	165	221	209	142	54	14	-

Non-Hispanic 1/, male

2003.....	19	34	41	63	115	204	261	191	80	22	2	-
2002.....	14	29	38	65	115	150	196	157	91	27	4	-
2001.....	11	13	20	31	74	103	119	90	46	14	-	-
2000.....	9	17	17	41	57	85	104	78	47	9	1	-
1999.....	5	10	14	26	45	78	89	53	43	12	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic 1/, female

2003.....	21	34	55	91	159	301	394	438	294	96	15	-
2002.....	13	34	34	64	107	208	303	343	221	85	18	-
2001.....	20	13	31	38	82	125	199	194	133	58	10	-
2000.....	9	11	17	39	74	111	149	182	118	43	9	-
1999.....	5	3	17	37	61	87	132	156	99	42	12	-

Non-Hispanic White, both sexes

2003.....	29	60	80	140	257	459	629	601	354	110	16	-
2002.....	24	47	57	120	199	336	478	473	297	107	19	-
2001.....	29	22	40	56	146	214	295	275	170	65	10	-
2000.....	14	25	29	71	120	184	246	240	159	48	9	-
1999.....	9	11	25	58	97	152	208	202	135	51	13	-

Non-Hispanic White, male

2003.....	15	27	34	59	107	191	251	182	79	20	2	-
2002.....	13	22	30	59	100	139	189	149	85	27	4	-
2001.....	10	11	18	26	68	95	108	85	45	12	-	-
2000.....	7	14	15	37	52	80	100	73	45	8	1	-
1999.....	4	9	11	25	44	73	85	52	40	11	1	-

Non-Hispanic White, female

2003.....	14	33	46	81	150	268	378	419	275	90	14	-
2002.....	11	25	27	61	99	197	289	324	212	80	15	-
2001.....	19	11	22	30	78	119	187	190	125	53	10	-
2000.....	7	11	14	34	68	104	146	167	114	40	8	-
1999.....	5	2	14	33	53	79	123	150	95	40	12	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Non-Hispanic Black, both sexes

2003.....	8	7	14	13	12	40	24	26	18	7	1	-
2002.....	2	15	14	9	22	18	17	25	12	4	3	-
2001.....	2	4	9	12	10	10	18	7	7	5	-	-
2000.....	4	3	3	8	7	10	6	17	5	4	-	-
1999.....	1	2	4	5	9	13	12	5	7	3	1	-

Non-Hispanic Black, male

2003.....	3	6	6	4	5	13	10	8	1	2	-	-
2002.....	1	7	8	6	14	8	5	7	5	-	-	-
2001.....	1	2	2	4	6	4	8	3	1	2	-	-
2000.....	2	3	1	3	3	4	4	4	2	1	-	-
1999.....	1	1	1	1	1	5	4	-	3	1	1	-

Non-Hispanic Black, female

2003.....	5	1	8	9	7	27	14	18	17	5	1	-
2002.....	1	8	6	3	8	10	12	18	7	4	3	-
2001.....	1	2	7	8	4	6	10	4	6	3	-	-
2000.....	2	-	2	5	4	6	2	13	3	3	-	-
1999.....	-	1	3	4	8	8	8	5	4	2	-	-

Origin not stated 2/, both sexes

2003.....	1	1	1	-	-	1	-	-	1	-	-	-
2002.....	1	-	1	-	1	1	-	2	-	-	-	-
2001.....	-	-	-	1	1	1	3	1	1	1	-	-
2000.....	-	-	-	-	2	-	2	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain other intestinal infections (A04,A07-A09)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	1	-	-	-	1	-	-	-
2000.....	-	-	-	-	1	-	2	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	1	1	1	-	-	1	-	-	-	-	-	-
2002.....	-	-	1	-	1	1	-	2	-	-	-	-
2001.....	-	-	-	-	-	1	3	1	-	1	-	-
2000.....	-	-	-	-	1	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Tuberculosis (A16-A19)

##### All origins, both sexes

2003.....	51	63	39	48	66	106	111	70	33	4	-	-
2002.....	46	44	63	82	79	114	120	68	42	9	2	-
2001.....	47	63	53	60	92	102	104	85	30	10	-	1
2000.....	53	31	50	70	85	132	120	74	31	10	-	1
1999.....	55	67	63	70	103	134	119	115	41	16	1	1

##### All origins, male

2003.....	40	46	25	28	44	61	55	32	11	2	-	-
2002.....	35	36	39	54	48	56	65	36	16	2	1	-
2001.....	36	47	38	40	58	53	47	34	18	5	-	1
2000.....	41	23	36	46	41	69	70	38	13	5	-	1
1999.....	45	48	48	49	67	71	66	58	16	11	-	1

##### All origins, female

2003.....	11	17	14	20	22	45	56	38	22	2	-	-
2002.....	11	8	24	28	31	58	55	32	26	7	1	-
2001.....	11	16	15	20	34	49	57	51	12	5	-	-
2000.....	12	8	14	24	44	63	50	36	18	5	-	-
1999.....	10	19	15	21	36	63	53	57	25	5	1	-

##### Hispanic, both sexes

2003.....	7	10	3	4	11	11	9	6	2	-	-	-
2002.....	7	4	13	11	10	6	13	5	5	1	-	-
2001.....	6	9	9	8	10	10	10	4	3	1	-	-
2000.....	7	4	8	5	8	14	9	8	1	1	-	-
1999.....	9	9	11	11	13	17	10	9	2	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Tuberculosis (A16-A19)

##### Hispanic, male

2003.....	6	7	2	1	10	6	5	3	-	-	-	-
2002.....	5	2	4	6	5	4	7	3	2	-	-	-
2001.....	6	8	5	6	8	4	6	3	-	-	-	-
2000.....	3	1	6	4	4	8	6	3	1	1	-	-
1999.....	8	3	8	8	8	9	4	4	2	-	-	-

##### Hispanic, female

2003.....	1	3	1	3	1	5	4	3	2	-	-	-
2002.....	2	2	9	5	5	2	6	2	3	1	-	-
2001.....	-	1	4	2	2	6	4	1	3	1	-	-
2000.....	4	3	2	1	4	6	3	5	-	-	-	-
1999.....	1	6	3	3	5	8	6	5	-	2	-	-

##### Non-Hispanic 1/, both sexes

2003.....	44	51	36	43	55	95	102	64	30	4	-	-
2002.....	38	40	48	70	69	108	107	63	37	8	2	-
2001.....	39	52	44	52	82	92	94	80	27	9	-	-
2000.....	46	27	41	64	76	115	111	66	29	9	-	-
1999.....	43	57	52	59	89	116	106	105	39	14	-	-

##### Non-Hispanic 1/, male

2003.....	34	37	23	26	34	55	50	29	11	2	-	-
2002.....	29	34	33	47	43	52	58	33	14	2	1	-
2001.....	28	37	33	34	50	49	41	30	18	5	-	-
2000.....	38	22	29	42	36	58	64	35	12	4	-	-
1999.....	34	44	40	41	58	61	61	53	14	11	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

Non-Hispanic 1/, female

2003.....	10	14	13	17	21	40	52	35	19	2	-	-
2002.....	9	6	15	23	26	56	49	30	23	6	1	-
2001.....	11	15	11	18	32	43	53	50	9	4	-	-
2000.....	8	5	12	22	40	57	47	31	17	5	-	-
1999.....	9	13	12	18	31	55	45	52	25	3	-	-

Non-Hispanic White, both sexes

2003.....	16	29	18	18	42	58	72	45	22	3	-	-
2002.....	16	16	25	29	37	72	73	43	25	6	1	-
2001.....	15	24	24	30	42	63	57	54	18	6	-	-
2000.....	24	5	17	32	40	64	67	45	24	6	-	-
1999.....	16	27	26	37	47	68	74	77	22	9	-	-

Non-Hispanic White, male

2003.....	13	22	12	10	24	30	36	21	7	2	-	-
2002.....	14	13	18	24	28	33	39	24	9	1	1	-
2001.....	9	19	18	18	21	34	25	18	11	4	-	-
2000.....	19	5	12	22	22	31	32	24	11	3	-	-
1999.....	15	23	18	25	32	33	44	38	7	6	-	-

Non-Hispanic White, female

2003.....	3	7	6	8	18	28	36	24	15	1	-	-
2002.....	2	3	7	5	9	39	34	19	16	5	-	-
2001.....	6	5	6	12	21	29	32	36	7	2	-	-
2000.....	5	-	5	10	18	33	35	21	13	3	-	-
1999.....	1	4	8	12	15	35	30	39	15	3	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

Non-Hispanic Black, both sexes

2003.....	23	19	10	14	6	16	17	10	4	-	-	-
2002.....	17	18	13	25	18	17	20	10	9	-	-	-
2001.....	22	23	14	19	25	16	19	11	8	3	-	-
2000.....	21	17	16	25	27	27	31	13	2	3	-	-
1999.....	21	22	17	15	27	31	19	14	8	4	-	-

Non-Hispanic Black, male

2003.....	18	14	5	10	5	12	8	4	2	-	-	-
2002.....	12	15	8	16	9	11	11	3	4	-	-	-
2001.....	18	16	10	13	17	7	7	5	6	1	-	-
2000.....	18	13	13	17	11	15	23	6	-	1	-	-
1999.....	16	15	16	11	17	17	8	6	3	4	-	-

Non-Hispanic Black, female

2003.....	5	5	5	4	1	4	9	6	2	-	-	-
2002.....	5	3	5	9	9	6	9	7	5	-	-	-
2001.....	4	7	4	6	8	9	12	6	2	2	-	-
2000.....	3	4	3	8	16	12	8	7	2	2	-	-
1999.....	5	7	1	4	10	14	11	8	5	-	-	-

Origin not stated 2/, both sexes

2003.....	-	2	-	1	-	-	-	-	1	-	-	-
2002.....	1	-	2	1	-	-	-	-	-	-	-	-
2001.....	2	2	-	-	-	-	-	1	-	-	-	1
2000.....	-	-	1	1	1	3	-	-	1	-	-	1
1999.....	3	1	-	-	1	1	3	1	-	-	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

Origin not stated 2/, male

2003.....	-	2	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	2	1	-	-	-	-	-	-	-	-
2001.....	2	2	-	-	-	-	-	1	-	-	-	1
2000.....	-	-	1	-	1	3	-	-	-	-	-	1
1999.....	3	1	-	-	1	1	1	1	-	-	-	1

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	2	-	-	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Respiratory tuberculosis (A16)

##### All origins, both sexes

2003.....	42	52	29	36	52	89	88	55	28	3	-	-
2002.....	36	37	47	61	59	87	105	60	37	7	2	-
2001.....	34	51	45	47	68	82	84	71	28	7	-	1
2000.....	42	24	36	52	67	107	102	64	26	8	-	1
1999.....	38	53	56	58	81	112	104	100	36	16	1	1

##### All origins, male

2003.....	32	38	19	23	38	52	45	26	9	1	-	-
2002.....	27	30	30	44	36	40	56	32	15	1	1	-
2001.....	29	41	37	31	45	46	38	29	17	4	-	1
2000.....	35	21	29	38	35	58	58	33	13	4	-	1
1999.....	32	39	44	41	56	60	60	52	13	11	-	1

##### All origins, female

2003.....	10	14	10	13	14	37	43	29	19	2	-	-
2002.....	9	7	17	17	23	47	49	28	22	6	1	-
2001.....	5	10	8	16	23	36	46	42	11	3	-	-
2000.....	7	3	7	14	32	49	44	31	13	4	-	-
1999.....	6	14	12	17	25	52	44	48	23	5	1	-

##### Hispanic, both sexes

2003.....	6	10	1	3	11	11	8	5	1	-	-	-
2002.....	5	4	8	9	7	3	13	5	5	1	-	-
2001.....	5	8	7	7	10	7	10	3	2	-	-	-
2000.....	5	3	6	5	7	10	8	6	1	1	-	-
1999.....	4	6	10	7	11	16	8	8	2	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Hispanic, male

2003.....	5	7	-	1	10	6	5	3	-	-	-	-
2002.....	3	2	3	6	3	2	7	3	2	-	-	-
2001.....	5	7	5	5	8	4	6	2	-	-	-	-
2000.....	3	1	5	4	4	7	5	3	1	1	-	-
1999.....	4	2	7	5	8	8	3	3	2	-	-	-

Hispanic, female

2003.....	1	3	1	2	1	5	3	2	1	-	-	-
2002.....	2	2	5	3	4	1	6	2	3	1	-	-
2001.....	-	1	2	2	2	3	4	1	2	-	-	-
2000.....	2	2	1	1	3	3	3	3	-	-	-	-
1999.....	-	4	3	2	3	8	5	5	-	2	-	-

Non-Hispanic 1/, both sexes

2003.....	36	40	28	33	41	78	80	50	26	3	-	-
2002.....	31	33	37	51	52	84	92	55	32	6	2	-
2001.....	28	41	38	40	58	75	74	67	26	7	-	-
2000.....	37	21	29	46	59	94	94	58	25	7	-	-
1999.....	32	46	46	51	69	95	94	91	34	14	-	-

Non-Hispanic 1/, male

2003.....	27	29	19	22	28	46	40	23	9	1	-	-
2002.....	24	28	25	37	33	38	49	29	13	1	1	-
2001.....	23	32	32	26	37	42	32	26	17	4	-	-
2000.....	32	20	23	34	30	48	53	30	12	3	-	-
1999.....	26	36	37	36	47	51	56	48	11	11	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Non-Hispanic 1/, female

2003.....	9	11	9	11	13	32	40	27	17	2	-	-
2002.....	7	5	12	14	19	46	43	26	19	5	1	-
2001.....	5	9	6	14	21	33	42	41	9	3	-	-
2000.....	5	1	6	12	29	46	41	28	13	4	-	-
1999.....	6	10	9	15	22	44	38	43	23	3	-	-

Non-Hispanic White, both sexes

2003.....	11	21	16	13	30	48	58	34	18	2	-	-
2002.....	12	13	19	24	29	56	58	37	22	5	1	-
2001.....	11	20	21	22	30	52	45	42	17	5	-	-
2000.....	21	4	13	21	31	54	58	39	21	5	-	-
1999.....	14	20	25	31	38	54	65	70	21	9	-	-

Non-Hispanic White, male

2003.....	8	16	12	9	19	24	29	16	5	1	-	-
2002.....	11	10	14	21	23	23	30	21	9	1	1	-
2001.....	9	17	18	13	16	30	19	14	10	3	-	-
2000.....	18	4	10	16	18	28	26	21	11	2	-	-
1999.....	13	18	18	22	26	28	39	37	7	6	-	-

Non-Hispanic White, female

2003.....	3	5	4	4	11	24	29	18	13	1	-	-
2002.....	1	3	5	3	6	33	28	16	13	4	-	-
2001.....	2	3	3	9	14	22	26	28	7	2	-	-
2000.....	3	-	3	5	13	26	32	18	10	3	-	-
1999.....	1	2	7	9	12	26	26	33	14	3	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Non-Hispanic Black, both sexes

2003.....	21	16	7	11	6	12	14	8	4	-	-	-
2002.....	16	15	11	19	13	14	20	8	7	-	-	-
2001.....	16	18	12	15	17	13	15	10	8	2	-	-
2000.....	15	12	10	21	22	22	24	12	2	2	-	-
1999.....	14	20	15	13	19	26	17	11	7	4	-	-

Non-Hispanic Black, male

2003.....	16	12	4	8	5	10	7	4	2	-	-	-
2002.....	12	13	7	12	7	10	11	2	3	-	-	-
2001.....	14	14	9	10	13	6	6	5	6	1	-	-
2000.....	13	12	10	16	10	11	19	5	-	1	-	-
1999.....	11	13	15	9	12	14	8	4	2	4	-	-

Non-Hispanic Black, female

2003.....	5	4	3	3	1	2	7	4	2	-	-	-
2002.....	4	2	4	7	6	4	9	6	4	-	-	-
2001.....	2	4	3	5	4	7	9	5	2	1	-	-
2000.....	2	-	-	5	12	11	5	7	2	1	-	-
1999.....	3	7	-	4	7	12	9	7	5	-	-	-

Origin not stated 2/, both sexes

2003.....	-	2	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	2	1	-	-	-	-	-	-	-	-
2001.....	1	2	-	-	-	-	-	1	-	-	-	1
2000.....	-	-	1	1	1	3	-	-	-	-	-	1
1999.....	2	1	-	-	1	1	2	1	-	-	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Origin not stated 2/, male

2003.....	-	2	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	2	1	-	-	-	-	-	-	-	-
2001.....	1	2	-	-	-	-	-	1	-	-	-	1
2000.....	-	-	1	-	1	3	-	-	-	-	-	1
1999.....	2	1	-	-	1	1	1	1	-	-	-	1

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

All origins, both sexes

2003.....	9	11	10	12	14	17	23	15	5	1	-	-
2002.....	10	7	16	21	20	27	15	8	5	2	-	-
2001.....	13	12	8	13	24	20	20	14	2	3	-	-
2000.....	11	7	14	18	18	25	18	10	5	2	-	-
1999.....	17	14	7	12	22	22	15	15	5	-	-	-

All origins, male

2003.....	8	8	6	5	6	9	10	6	2	1	-	-
2002.....	8	6	9	10	12	16	9	4	1	1	-	-
2001.....	7	6	1	9	13	7	9	5	1	1	-	-
2000.....	6	2	7	8	6	11	12	5	-	1	-	-
1999.....	13	9	4	8	11	11	6	6	3	-	-	-

All origins, female

2003.....	1	3	4	7	8	8	13	9	3	-	-	-
2002.....	2	1	7	11	8	11	6	4	4	1	-	-
2001.....	6	6	7	4	11	13	11	9	1	2	-	-
2000.....	5	5	7	10	12	14	6	5	5	1	-	-
1999.....	4	5	3	4	11	11	9	9	2	-	-	-

Hispanic, both sexes

2003.....	1	-	2	1	-	-	1	1	1	-	-	-
2002.....	2	-	5	2	3	3	-	-	-	-	-	-
2001.....	1	1	2	1	-	3	-	1	1	1	-	-
2000.....	2	1	2	-	1	4	1	2	-	-	-	-
1999.....	5	3	1	4	2	1	2	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Hispanic, male

2003.....	1	-	2	-	-	-	-	-	-	-	-	-
2002.....	2	-	1	-	2	2	-	-	-	-	-	-
2001.....	1	1	-	1	-	-	-	1	-	-	-	-
2000.....	-	-	1	-	-	1	1	-	-	-	-	-
1999.....	4	1	1	3	-	1	1	1	-	-	-	-

Hispanic, female

2003.....	-	-	-	1	-	-	1	1	1	-	-	-
2002.....	-	-	4	2	1	1	-	-	-	-	-	-
2001.....	-	-	2	-	-	3	-	-	1	1	-	-
2000.....	2	1	1	-	1	3	-	2	-	-	-	-
1999.....	1	2	-	1	2	-	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	8	11	8	10	14	17	22	14	4	1	-	-
2002.....	7	7	11	19	17	24	15	8	5	2	-	-
2001.....	11	11	6	12	24	17	20	13	1	2	-	-
2000.....	9	6	12	18	17	21	17	8	4	2	-	-
1999.....	11	11	6	8	20	21	12	14	5	-	-	-

Non-Hispanic 1/, male

2003.....	7	8	4	4	6	9	10	6	2	1	-	-
2002.....	5	6	8	10	10	14	9	4	1	1	-	-
2001.....	5	5	1	8	13	7	9	4	1	1	-	-
2000.....	6	2	6	8	6	10	11	5	-	1	-	-
1999.....	8	8	3	5	11	10	5	5	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Non-Hispanic 1/, female

2003.....	1	3	4	6	8	8	12	8	2	-	-	-
2002.....	2	1	3	9	7	10	6	4	4	1	-	-
2001.....	6	6	5	4	11	10	11	9	-	1	-	-
2000.....	3	4	6	10	11	11	6	3	4	1	-	-
1999.....	3	3	3	3	9	11	7	9	2	-	-	-

Non-Hispanic White, both sexes

2003.....	5	8	2	5	12	10	14	11	4	1	-	-
2002.....	4	3	6	5	8	16	15	6	3	1	-	-
2001.....	4	4	3	8	12	11	12	12	1	1	-	-
2000.....	3	1	4	11	9	10	9	6	3	1	-	-
1999.....	2	7	1	6	9	14	9	7	1	-	-	-

Non-Hispanic White, male

2003.....	5	6	-	1	5	6	7	5	2	1	-	-
2002.....	3	3	4	3	5	10	9	3	-	-	-	-
2001.....	-	2	-	5	5	4	6	4	1	1	-	-
2000.....	1	1	2	6	4	3	6	3	-	1	-	-
1999.....	2	5	-	3	6	5	5	1	-	-	-	-

Non-Hispanic White, female

2003.....	-	2	2	4	7	4	7	6	2	-	-	-
2002.....	1	-	2	2	3	6	6	3	3	1	-	-
2001.....	4	2	3	3	7	7	6	8	-	-	-	-
2000.....	2	-	2	5	5	7	3	3	3	-	-	-
1999.....	-	2	1	3	3	9	4	6	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Non-Hispanic Black, both sexes

2003.....	2	3	3	3	-	4	3	2	-	-	-	-
2002.....	1	3	2	6	5	3	-	2	2	-	-	-
2001.....	6	5	2	4	8	3	4	1	-	1	-	-
2000.....	6	5	6	4	5	5	7	1	-	1	-	-
1999.....	7	2	2	2	8	5	2	3	1	-	-	-

Non-Hispanic Black, male

2003.....	2	2	1	2	-	2	1	-	-	-	-	-
2002.....	-	2	1	4	2	1	-	1	1	-	-	-
2001.....	4	2	1	3	4	1	1	-	-	-	-	-
2000.....	5	1	3	1	1	4	4	1	-	-	-	-
1999.....	5	2	1	2	5	3	-	2	1	-	-	-

Non-Hispanic Black, female

2003.....	-	1	2	1	-	2	2	2	-	-	-	-
2002.....	1	1	1	2	3	2	-	1	1	-	-	-
2001.....	2	3	1	1	4	2	3	1	-	1	-	-
2000.....	1	4	3	3	4	1	3	-	-	1	-	-
1999.....	2	-	1	-	3	2	2	1	-	-	-	-

Origin not stated 2/, both sexes

2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Origin not stated 2/, male

2003.....	-	-	-	1	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-















Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Scarlet fever and erysipelas (A38,A46)

Non-Hispanic 1/, female

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	1	1	-	-	1	-	1	-

Non-Hispanic White, both sexes

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	1	-	1	-

Non-Hispanic White, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic White, female

2003.....	-	-	-	-	-	-	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	1	-	1	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Meningococcal infection (A39)

##### All origins, both sexes

2003.....	10	9	2	3	6	4	5	3	-	-	-	-
2002.....	9	9	4	3	4	3	7	2	-	3	-	-
2001.....	4	11	4	7	8	7	1	6	1	2	-	-
2000.....	16	6	3	3	5	7	8	2	4	1	-	-
1999.....	6	3	1	9	-	7	6	3	4	1	-	-

##### All origins, male

2003.....	5	4	1	1	1	-	1	1	-	-	-	-
2002.....	5	2	2	1	1	1	2	1	-	1	-	-
2001.....	2	5	2	2	3	2	1	1	-	1	-	-
2000.....	6	1	1	-	2	4	3	-	-	1	-	-
1999.....	3	-	-	5	-	2	2	-	1	1	-	-

##### All origins, female

2003.....	5	5	1	2	5	4	4	2	-	-	-	-
2002.....	4	7	2	2	3	2	5	1	-	2	-	-
2001.....	2	6	2	5	5	5	-	5	1	1	-	-
2000.....	10	5	2	3	3	3	5	2	4	-	-	-
1999.....	3	3	1	4	-	5	4	3	3	-	-	-

##### Hispanic, both sexes

2003.....	1	-	-	2	2	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	2	-	-	-	1	-	-	-	-	-	-
2000.....	1	-	1	-	-	1	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningococcal infection (A39)

Hispanic, male

2003.....	1	-	-	1	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	1	2	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	2	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	1	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	9	9	2	1	4	4	5	2	-	-	-	-
2002.....	9	9	3	3	4	3	7	2	-	3	-	-
2001.....	4	9	4	7	8	6	1	6	1	2	-	-
2000.....	15	6	2	3	5	6	7	2	4	1	-	-
1999.....	6	3	1	9	-	6	6	3	4	1	-	-

Non-Hispanic 1/, male

2003.....	4	4	1	-	1	-	1	-	-	-	-	-
2002.....	5	2	1	1	1	1	2	1	-	1	-	-
2001.....	2	5	2	2	3	1	1	1	-	1	-	-
2000.....	6	1	1	-	2	3	3	-	-	1	-	-
1999.....	3	-	-	5	-	2	2	-	1	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningococcal infection (A39)

Non-Hispanic 1/, female

2003.....	5	5	1	1	3	4	4	2	-	-	-	-
2002.....	4	7	2	2	3	2	5	1	-	2	-	-
2001.....	2	4	2	5	5	5	-	5	1	1	-	-
2000.....	9	5	1	3	3	3	4	2	4	-	-	-
1999.....	3	3	1	4	-	4	4	3	3	-	-	-

Non-Hispanic White, both sexes

2003.....	4	6	1	1	4	4	5	2	-	-	-	-
2002.....	7	7	3	2	4	3	7	2	-	3	-	-
2001.....	4	8	1	6	8	5	1	5	1	2	-	-
2000.....	11	5	2	2	5	6	7	2	4	1	-	-
1999.....	5	3	1	9	-	6	5	3	4	1	-	-

Non-Hispanic White, male

2003.....	2	2	-	-	1	-	1	-	-	-	-	-
2002.....	3	2	1	-	1	1	2	1	-	1	-	-
2001.....	2	4	1	2	3	-	1	-	-	1	-	-
2000.....	3	1	1	-	2	3	3	-	-	1	-	-
1999.....	2	-	-	5	-	2	2	-	1	1	-	-

Non-Hispanic White, female

2003.....	2	4	1	1	3	4	4	2	-	-	-	-
2002.....	4	5	2	2	3	2	5	1	-	2	-	-
2001.....	2	4	-	4	5	5	-	5	1	1	-	-
2000.....	8	4	1	2	3	3	4	2	4	-	-	-
1999.....	3	3	1	4	-	4	3	3	3	-	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Septicemia (A40-A41)

##### All origins, both sexes

2003.....	1,241	1,631	2,020	2,459	3,511	5,096	5,836	5,178	3,064	1,112	189	3
2002.....	1,170	1,499	1,861	2,589	3,747	5,195	5,815	5,026	2,988	1,111	199	3
2001.....	1,140	1,392	1,719	2,409	3,589	5,000	5,351	4,856	3,001	1,020	192	-
2000.....	1,064	1,260	1,639	2,303	3,401	4,745	5,193	4,900	2,949	1,079	216	1
1999.....	954	1,178	1,536	2,352	3,398	4,686	5,024	4,854	2,984	1,133	195	-

##### All origins, male

2003.....	657	864	1,023	1,299	1,724	2,377	2,517	1,920	894	247	28	2
2002.....	629	767	981	1,292	1,862	2,476	2,552	1,857	834	247	35	3
2001.....	604	728	878	1,227	1,817	2,402	2,360	1,788	869	225	32	-
2000.....	600	657	856	1,182	1,685	2,249	2,233	1,690	812	199	35	1
1999.....	513	613	799	1,218	1,701	2,252	2,174	1,712	817	236	21	-

##### All origins, female

2003.....	584	767	997	1,160	1,787	2,719	3,319	3,258	2,170	865	161	1
2002.....	541	732	880	1,297	1,885	2,719	3,263	3,169	2,154	864	164	-
2001.....	536	664	841	1,182	1,772	2,598	2,991	3,068	2,132	795	160	-
2000.....	464	603	783	1,121	1,716	2,496	2,960	3,210	2,137	880	181	-
1999.....	441	565	737	1,134	1,697	2,434	2,850	3,142	2,167	897	174	-

##### Hispanic, both sexes

2003.....	74	90	100	130	140	199	198	143	108	38	4	2
2002.....	80	71	94	132	209	206	176	144	96	41	1	-
2001.....	63	57	94	129	159	191	155	136	93	35	10	-
2000.....	71	61	84	97	138	157	143	119	70	24	5	-
1999.....	57	63	68	119	146	162	134	131	83	37	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Septicemia (A40-A41)

##### Hispanic, male

2003.....	46	53	42	70	67	93	82	60	32	12	-	1
2002.....	54	35	49	61	100	96	80	52	32	13	1	-
2001.....	43	32	54	69	80	101	69	49	25	9	1	-
2000.....	42	37	43	60	66	79	64	41	24	11	2	-
1999.....	34	37	41	67	67	90	54	46	28	10	-	-

##### Hispanic, female

2003.....	28	37	58	60	73	106	116	83	76	26	4	1
2002.....	26	36	45	71	109	110	96	92	64	28	-	-
2001.....	20	25	40	60	79	90	86	87	68	26	9	-
2000.....	29	24	41	37	72	78	79	78	46	13	3	-
1999.....	23	26	27	52	79	72	80	85	55	27	6	-

##### Non-Hispanic 1/, both sexes

2003.....	1,164	1,534	1,911	2,322	3,358	4,890	5,622	5,024	2,946	1,073	185	1
2002.....	1,088	1,420	1,762	2,442	3,527	4,971	5,622	4,865	2,884	1,066	197	3
2001.....	1,071	1,327	1,614	2,275	3,415	4,798	5,179	4,712	2,901	983	180	-
2000.....	985	1,194	1,545	2,194	3,248	4,568	5,033	4,765	2,868	1,049	210	1
1999.....	890	1,108	1,462	2,224	3,242	4,505	4,873	4,708	2,887	1,092	188	-

##### Non-Hispanic 1/, male

2003.....	609	808	976	1,225	1,651	2,279	2,429	1,855	861	235	28	1
2002.....	574	727	930	1,221	1,754	2,367	2,463	1,799	801	233	34	3
2001.....	555	694	816	1,154	1,728	2,294	2,281	1,738	843	216	30	-
2000.....	555	615	809	1,113	1,611	2,160	2,156	1,641	787	187	32	1
1999.....	474	573	757	1,147	1,630	2,149	2,112	1,659	785	225	21	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Septicemia (A40-A41)

##### Non-Hispanic 1/, female

2003.....	555	726	935	1,097	1,707	2,611	3,193	3,169	2,085	838	157	-
2002.....	514	693	832	1,221	1,773	2,604	3,159	3,066	2,083	833	163	-
2001.....	516	633	798	1,121	1,687	2,504	2,898	2,974	2,058	767	150	-
2000.....	430	579	736	1,081	1,637	2,408	2,877	3,124	2,081	862	178	-
1999.....	416	535	705	1,077	1,612	2,356	2,761	3,049	2,102	867	167	-

##### Non-Hispanic White, both sexes

2003.....	751	1,088	1,383	1,688	2,627	4,019	4,784	4,302	2,516	890	132	-
2002.....	702	996	1,245	1,805	2,778	4,043	4,763	4,168	2,466	883	149	2
2001.....	703	919	1,162	1,704	2,681	3,928	4,424	4,040	2,499	825	134	-
2000.....	647	836	1,083	1,610	2,563	3,690	4,211	4,075	2,468	848	159	1
1999.....	594	749	1,043	1,646	2,494	3,628	4,089	3,974	2,452	917	136	-

##### Non-Hispanic White, male

2003.....	400	562	718	895	1,312	1,886	2,086	1,599	735	195	21	-
2002.....	368	524	680	903	1,407	1,921	2,085	1,557	676	192	23	2
2001.....	365	470	601	868	1,355	1,892	1,959	1,492	712	184	18	-
2000.....	359	437	578	826	1,305	1,748	1,831	1,401	668	150	23	1
1999.....	317	381	523	847	1,264	1,762	1,776	1,396	659	184	18	-

##### Non-Hispanic White, female

2003.....	351	526	665	793	1,315	2,133	2,698	2,703	1,781	695	111	-
2002.....	334	472	565	902	1,371	2,122	2,678	2,611	1,790	691	126	-
2001.....	338	449	561	836	1,326	2,036	2,465	2,548	1,787	641	116	-
2000.....	288	399	505	784	1,258	1,942	2,380	2,674	1,800	698	136	-
1999.....	277	368	520	799	1,230	1,866	2,313	2,578	1,793	733	118	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Septicemia (A40-A41)

##### Non-Hispanic Black, both sexes

2003.....	376	396	485	575	657	778	758	654	384	170	49	1
2002.....	355	400	468	581	670	845	788	632	381	166	47	1
2001.....	334	381	423	526	660	797	704	626	364	148	43	-
2000.....	314	324	432	537	636	817	759	631	378	185	51	-
1999.....	267	329	387	520	692	792	721	673	394	162	52	-

##### Non-Hispanic Black, male

2003.....	195	226	236	301	300	348	315	229	112	36	7	1
2002.....	192	191	227	286	312	395	334	212	108	35	10	1
2001.....	171	211	200	259	327	365	291	225	110	27	11	-
2000.....	183	165	216	262	283	384	293	216	109	34	9	-
1999.....	143	173	219	274	335	343	305	234	106	37	3	-

##### Non-Hispanic Black, female

2003.....	181	170	249	274	357	430	443	425	272	134	42	-
2002.....	163	209	241	295	358	450	454	420	273	131	37	-
2001.....	163	170	223	267	333	432	413	401	254	121	32	-
2000.....	131	159	216	275	353	433	466	415	269	151	42	-
1999.....	124	156	168	246	357	449	416	439	288	125	49	-

##### Origin not stated 2/, both sexes

2003.....	3	7	9	7	13	7	16	11	10	1	-	-
2002.....	2	8	5	15	11	18	17	17	8	4	1	-
2001.....	6	8	11	5	15	11	17	8	7	2	2	-
2000.....	8	5	10	12	15	20	17	16	11	6	1	-
1999.....	7	7	6	9	10	19	17	15	14	4	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Septicemia (A40-A41)

Origin not stated 2/, male

2003.....	2	3	5	4	6	5	6	5	1	-	-	-
2002.....	1	5	2	10	8	13	9	6	1	1	-	-
2001.....	6	2	8	4	9	7	10	1	1	-	1	-
2000.....	3	5	4	9	8	10	13	8	1	1	1	-
1999.....	5	3	1	4	4	13	8	7	4	1	-	-

Origin not stated 2/, female

2003.....	1	4	4	3	7	2	10	6	9	1	-	-
2002.....	1	3	3	5	3	5	8	11	7	3	1	-
2001.....	-	6	3	1	6	4	7	7	6	2	1	-
2000.....	5	-	6	3	7	10	4	8	10	5	-	-
1999.....	2	4	5	5	6	6	9	8	10	3	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Syphilis (A50-A53)

##### All origins, both sexes

2003.....	3	2	1	3	3	6	2	5	3	1	-	-
2002.....	4	2	3	4	2	5	7	6	3	-	-	-
2001.....	1	3	1	3	2	4	9	2	2	1	-	-
2000.....	4	2	1	1	6	7	6	6	3	1	-	-
1999.....	2	1	3	2	4	6	4	2	1	1	-	-

##### All origins, male

2003.....	3	1	1	2	3	2	2	2	1	-	-	-
2002.....	4	1	1	4	1	3	1	2	1	-	-	-
2001.....	1	2	1	1	2	2	3	1	-	-	-	-
2000.....	4	2	1	1	1	3	4	3	2	-	-	-
1999.....	1	1	2	2	3	-	-	1	-	-	-	-

##### All origins, female

2003.....	-	1	-	1	-	4	-	3	2	1	-	-
2002.....	-	1	2	-	1	2	6	4	2	-	-	-
2001.....	-	1	-	2	-	2	6	1	2	1	-	-
2000.....	-	-	-	-	5	4	2	3	1	1	-	-
1999.....	1	-	1	-	1	6	4	1	1	1	-	-

##### Hispanic, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	2	-	-	-	1	2	1	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Syphilis (A50-A53)

Hispanic, male

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	2	-	-	-	1	-	1	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	2	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	2	2	1	2	3	6	2	5	3	1	-	-
2002.....	4	2	3	4	2	4	7	5	3	-	-	-
2001.....	1	-	1	3	2	3	7	1	2	1	-	-
2000.....	4	2	1	1	6	5	6	6	3	1	-	-
1999.....	1	1	3	2	4	6	3	2	1	1	-	-

Non-Hispanic 1/, male

2003.....	2	1	1	2	3	2	2	2	1	-	-	-
2002.....	4	1	1	4	1	2	1	2	1	-	-	-
2001.....	1	-	1	1	2	1	3	-	-	-	-	-
2000.....	4	2	1	1	1	3	4	3	2	-	-	-
1999.....	-	1	2	2	3	-	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Syphilis (A50-A53)

Non-Hispanic 1/, female

2003.....	-	1	-	-	-	4	-	3	2	1	-	-
2002.....	-	1	2	-	1	2	6	3	2	-	-	-
2001.....	-	-	-	2	-	2	4	1	2	1	-	-
2000.....	-	-	-	-	5	2	2	3	1	1	-	-
1999.....	1	-	1	-	1	6	3	1	1	1	-	-

Non-Hispanic White, both sexes

2003.....	-	2	-	2	-	4	1	2	3	1	-	-
2002.....	2	-	1	2	1	-	4	3	2	-	-	-
2001.....	1	-	1	1	1	3	4	1	-	1	-	-
2000.....	1	-	-	-	3	2	4	4	3	1	-	-
1999.....	1	-	3	2	1	2	1	-	-	1	-	-

Non-Hispanic White, male

2003.....	-	1	-	2	-	2	1	1	1	-	-	-
2002.....	2	-	-	2	1	-	-	2	-	-	-	-
2001.....	1	-	1	-	1	1	1	-	-	-	-	-
2000.....	1	-	-	-	-	-	2	1	2	-	-	-
1999.....	-	-	2	2	-	-	-	-	-	-	-	-

Non-Hispanic White, female

2003.....	-	1	-	-	-	2	-	1	2	1	-	-
2002.....	-	-	1	-	-	-	4	1	2	-	-	-
2001.....	-	-	-	1	-	2	3	1	-	1	-	-
2000.....	-	-	-	-	3	2	2	3	1	1	-	-
1999.....	1	-	1	-	1	2	1	-	-	1	-	-

















Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Arthropod-borne viral encephalitis (A83-A84,A85.2)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1	-	3	-	2	4	1	-	-	-	-	-
2002.....	2	-	1	-	1	1	1	1	-	-	-	-
2001.....	1	-	-	-	5	1	1	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	-	-	1	-	-	-	-

Non-Hispanic 1/, male

2003.....	1	-	2	-	2	2	-	-	-	-	-	-
2002.....	1	-	1	-	1	1	-	-	-	-	-	-
2001.....	-	-	-	-	4	1	1	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	1	-	-	1	-	-	-	-

















Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Viral hepatitis (B15-B19)

##### All origins, both sexes

2003.....	1,206	709	376	330	368	323	220	87	24	4	-	-
2002.....	1,139	644	428	369	393	352	235	136	34	7	-	-
2001.....	959	586	428	407	429	375	238	120	41	5	-	1
2000.....	850	492	412	410	435	401	259	123	35	10	-	1
1999.....	635	438	348	427	454	389	262	145	45	5	2	-

##### All origins, male

2003.....	903	506	223	169	183	147	102	30	8	2	-	-
2002.....	868	448	254	193	180	174	96	56	11	3	-	-
2001.....	737	406	237	207	189	156	107	42	13	2	-	1
2000.....	638	338	239	202	194	198	113	55	11	4	-	1
1999.....	487	298	193	217	209	185	132	61	20	1	-	-

##### All origins, female

2003.....	303	203	153	161	185	176	118	57	16	2	-	-
2002.....	271	196	174	176	213	178	139	80	23	4	-	-
2001.....	222	180	191	200	240	219	131	78	28	3	-	-
2000.....	212	154	173	208	241	203	146	68	24	6	-	-
1999.....	148	140	155	210	245	204	130	84	25	4	2	-

##### Hispanic, both sexes

2003.....	142	93	58	59	47	39	20	7	2	-	-	-
2002.....	149	92	72	58	44	43	20	12	1	2	-	-
2001.....	111	117	75	52	61	43	18	10	5	-	-	-
2000.....	99	67	80	62	58	42	24	9	1	1	-	-
1999.....	104	69	58	68	61	42	23	6	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

Hispanic, male

2003.....	110	67	36	28	14	14	6	2	1	-	-	-
2002.....	105	65	43	22	18	20	5	6	-	-	-	-
2001.....	87	85	38	17	18	15	8	3	1	-	-	-
2000.....	74	43	51	28	24	17	8	5	-	1	-	-
1999.....	77	43	32	32	27	14	4	4	-	-	-	-

Hispanic, female

2003.....	32	26	22	31	33	25	14	5	1	-	-	-
2002.....	44	27	29	36	26	23	15	6	1	2	-	-
2001.....	24	32	37	35	43	28	10	7	4	-	-	-
2000.....	25	24	29	34	34	25	16	4	1	-	-	-
1999.....	27	26	26	36	34	28	19	2	2	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,059	614	317	269	318	284	200	80	22	4	-	-
2002.....	986	543	355	309	346	309	214	124	33	5	-	-
2001.....	845	465	352	354	367	332	220	110	36	5	-	1
2000.....	745	423	328	348	376	359	235	113	34	9	-	1
1999.....	529	366	290	358	392	346	239	139	43	5	2	-

Non-Hispanic 1/, male

2003.....	788	438	187	139	167	133	96	28	7	2	-	-
2002.....	761	376	210	169	162	154	91	50	11	3	-	-
2001.....	648	317	198	190	171	141	99	39	12	2	-	1
2000.....	558	293	186	174	169	181	105	49	11	3	-	1
1999.....	408	253	161	184	181	170	128	57	20	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

Non-Hispanic 1/, female

2003.....	271	176	130	130	151	151	104	52	15	2	-	-
2002.....	225	167	145	140	184	155	123	74	22	2	-	-
2001.....	197	148	154	164	196	191	121	71	24	3	-	-
2000.....	187	130	142	174	207	178	130	64	23	6	-	-
1999.....	121	113	129	174	211	176	111	82	23	4	2	-

Non-Hispanic White, both sexes

2003.....	813	423	219	185	242	225	165	67	19	3	-	-
2002.....	770	388	247	219	270	256	184	109	29	2	-	-
2001.....	622	330	256	263	281	279	188	94	29	5	-	-
2000.....	536	286	234	258	301	308	198	101	29	8	-	1
1999.....	378	254	211	262	305	285	212	119	38	4	2	-

Non-Hispanic White, male

2003.....	613	310	135	101	131	103	83	25	6	2	-	-
2002.....	599	271	144	131	132	132	82	45	10	-	-	-
2001.....	492	233	151	153	143	128	90	38	9	2	-	-
2000.....	413	205	134	128	137	158	90	44	11	2	-	1
1999.....	294	180	128	144	151	146	122	49	17	1	-	-

Non-Hispanic White, female

2003.....	200	113	84	84	111	122	82	42	13	1	-	-
2002.....	171	117	103	88	138	124	102	64	19	2	-	-
2001.....	130	97	105	110	138	151	98	56	20	3	-	-
2000.....	123	81	100	130	164	150	108	57	18	6	-	-
1999.....	84	74	83	118	154	139	90	70	21	3	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Viral hepatitis (B15-B19)

Non-Hispanic Black, both sexes

2003.....	216	147	74	48	41	36	21	8	2	1	-	-
2002.....	188	113	71	56	42	34	15	7	4	2	-	-
2001.....	187	100	67	46	45	35	18	12	4	-	-	1
2000.....	159	90	48	47	46	32	25	8	2	-	-	-
1999.....	117	73	45	59	50	36	17	16	2	1	-	-

Non-Hispanic Black, male

2003.....	158	106	39	22	19	16	5	3	1	-	-	-
2002.....	141	82	40	31	17	14	3	-	1	2	-	-
2001.....	128	63	36	18	16	9	4	-	2	-	-	1
2000.....	115	61	25	26	22	13	12	3	-	-	-	-
1999.....	90	44	21	21	17	13	2	7	1	-	-	-

Non-Hispanic Black, female

2003.....	58	41	35	26	22	20	16	5	1	1	-	-
2002.....	47	31	31	25	25	20	12	7	3	-	-	-
2001.....	59	37	31	28	29	26	14	12	2	-	-	-
2000.....	44	29	23	21	24	19	13	5	2	-	-	-
1999.....	27	29	24	38	33	23	15	9	1	1	-	-

Origin not stated 2/, both sexes

2003.....	5	2	1	2	3	-	-	-	-	-	-	-
2002.....	4	9	1	2	3	-	1	-	-	-	-	-
2001.....	3	4	1	1	1	-	-	-	-	-	-	-
2000.....	6	2	4	-	1	-	-	1	-	-	-	-
1999.....	2	3	-	1	1	1	-	-	-	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins, both sexes

2003.....	1,806	1,017	500	277	161	71	24	11	3	-	-	3
2002.....	1,794	885	462	275	126	71	25	8	3	1	-	4
2001.....	1,609	821	496	258	126	68	20	5	3	2	-	2
2000.....	1,591	797	442	250	147	69	16	4	3	2	-	1
1999.....	1,467	777	372	261	137	52	18	7	5	1	1	1

All origins, male

2003.....	1,425	816	399	209	127	55	20	10	2	-	-	3
2002.....	1,446	710	373	223	100	53	18	4	-	1	-	4
2001.....	1,299	673	378	187	105	55	18	4	1	2	-	2
2000.....	1,288	655	358	199	114	54	11	1	1	-	-	1
1999.....	1,198	646	302	211	106	39	11	6	1	1	-	1

All origins, female

2003.....	381	201	101	68	34	16	4	1	1	-	-	-
2002.....	348	175	89	52	26	18	7	4	3	-	-	-
2001.....	310	148	118	71	21	13	2	1	2	-	-	-
2000.....	303	142	84	51	33	15	5	3	2	2	-	-
1999.....	269	131	70	50	31	13	7	1	4	-	1	-

Hispanic, both sexes

2003.....	238	150	54	42	22	8	2	-	1	-	-	-
2002.....	217	123	65	28	12	6	3	-	1	-	-	3
2001.....	210	111	58	38	13	9	1	1	-	1	-	-
2000.....	209	96	67	28	13	11	3	-	-	-	-	-
1999.....	168	89	48	35	14	2	2	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Hispanic, male

2003.....	193	118	35	31	19	5	2	-	1	-	-	-
2002.....	170	95	45	25	11	4	3	-	-	-	-	3
2001.....	179	88	41	32	11	8	1	1	-	1	-	-
2000.....	174	79	44	22	12	8	2	-	-	-	-	-
1999.....	130	76	42	28	13	2	2	-	-	-	-	-

Hispanic, female

2003.....	45	32	19	11	3	3	-	-	-	-	-	-
2002.....	47	28	20	3	1	2	-	-	1	-	-	-
2001.....	31	23	17	6	2	1	-	-	-	-	-	-
2000.....	35	17	23	6	1	3	1	-	-	-	-	-
1999.....	38	13	6	7	1	-	-	-	1	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,540	857	441	229	137	62	22	11	2	-	-	2
2002.....	1,550	755	393	245	114	65	22	8	2	1	-	1
2001.....	1,370	702	421	215	110	58	19	4	3	1	-	2
2000.....	1,362	688	370	213	130	57	13	4	3	2	-	-
1999.....	1,269	675	312	218	121	49	16	7	4	1	1	-

Non-Hispanic 1/, male

2003.....	1,213	689	361	172	106	49	18	10	1	-	-	2
2002.....	1,253	609	324	197	89	49	15	4	-	1	-	1
2001.....	1,095	579	323	150	91	46	17	3	1	1	-	2
2000.....	1,097	563	309	169	98	46	9	1	1	-	-	-
1999.....	1,042	557	249	177	91	36	9	6	1	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic 1/, female

2003.....	327	168	80	57	31	13	4	1	1	-	-	-
2002.....	297	146	69	48	25	16	7	4	2	-	-	-
2001.....	275	123	98	65	19	12	2	1	2	-	-	-
2000.....	265	125	61	44	32	11	4	3	2	2	-	-
1999.....	227	118	63	41	30	13	7	1	3	-	1	-

Non-Hispanic White, both sexes

2003.....	532	323	162	68	57	33	5	5	2	-	-	1
2002.....	502	302	158	78	34	24	9	5	2	1	-	-
2001.....	477	295	128	64	46	22	5	2	2	1	-	-
2000.....	499	291	129	80	42	23	5	2	1	-	-	-
1999.....	520	252	117	83	38	22	10	3	3	-	1	-

Non-Hispanic White, male

2003.....	470	280	148	55	48	30	5	4	1	-	-	1
2002.....	444	269	140	68	31	17	5	3	-	1	-	-
2001.....	432	262	108	50	45	19	5	1	1	1	-	-
2000.....	441	253	116	72	31	21	4	-	-	-	-	-
1999.....	470	224	107	75	29	19	5	2	-	-	-	-

Non-Hispanic White, female

2003.....	62	43	14	13	9	3	-	1	1	-	-	-
2002.....	58	33	18	10	3	7	4	2	2	-	-	-
2001.....	45	33	20	14	1	3	-	1	1	-	-	-
2000.....	58	38	13	8	11	2	1	2	1	-	-	-
1999.....	50	28	10	8	9	3	5	1	3	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Non-Hispanic Black, both sexes

2003.....	994	527	275	159	80	28	17	6	-	-	-	1
2002.....	1,028	445	230	163	77	41	13	3	-	-	-	1
2001.....	877	404	286	146	62	36	13	1	1	-	-	2
2000.....	851	388	237	130	84	33	8	2	2	2	-	-
1999.....	735	421	192	132	81	27	6	3	1	1	-	-

Non-Hispanic Black, male

2003.....	731	404	210	117	58	18	13	6	-	-	-	1
2002.....	792	332	180	125	57	32	10	1	-	-	-	1
2001.....	648	315	209	99	45	27	11	1	-	-	-	2
2000.....	647	303	189	95	64	24	5	1	1	-	-	-
1999.....	560	331	140	99	61	17	4	3	1	1	-	-

Non-Hispanic Black, female

2003.....	263	123	65	42	22	10	4	-	-	-	-	-
2002.....	236	113	50	38	20	9	3	2	-	-	-	-
2001.....	229	89	77	47	17	9	2	-	1	-	-	-
2000.....	204	85	48	35	20	9	3	1	1	2	-	-
1999.....	175	90	52	33	20	10	2	-	-	-	-	-

Origin not stated 2/, both sexes

2003.....	28	10	5	6	2	1	-	-	-	-	-	1
2002.....	27	7	4	2	-	-	-	-	-	-	-	-
2001.....	29	8	17	5	3	1	-	-	-	-	-	-
2000.....	20	13	5	9	4	1	-	-	-	-	-	1
1999.....	30	13	12	8	2	1	-	-	-	-	-	1





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malaria (B50-B54)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	-	2	-	1	-	-	-	-	-	-	-	-
2002.....	4	2	-	1	1	-	-	-	-	-	-	-
2001.....	-	-	2	1	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	1	1	-	-	-	-	-	-	-

Non-Hispanic 1/, male

2003.....	-	2	-	1	-	-	-	-	-	-	-	-
2002.....	3	1	-	1	1	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	1	-	-	-	-	-	-
2000.....	1	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins, both sexes

2003.....	659	590	546	577	767	913	802	594	335	113	20	-
2002.....	513	445	444	520	703	859	735	580	289	112	16	-
2001.....	356	338	433	537	606	698	632	483	288	98	18	-
2000.....	308	323	384	451	623	739	636	513	265	93	28	-
1999.....	444	427	471	652	791	927	812	633	330	138	22	-

All origins, male

2003.....	441	376	313	316	395	430	386	232	123	24	4	-
2002.....	348	273	259	290	373	429	356	230	110	30	3	-
2001.....	215	209	240	275	290	350	306	185	101	21	-	-
2000.....	194	179	204	253	344	367	298	216	97	22	4	-
1999.....	247	220	262	351	409	460	367	253	99	28	1	-

All origins, female

2003.....	218	214	233	261	372	483	416	362	212	89	16	-
2002.....	165	172	185	230	330	430	379	350	179	82	13	-
2001.....	141	129	193	262	316	348	326	298	187	77	18	-
2000.....	114	144	180	198	279	372	338	297	168	71	24	-
1999.....	197	207	209	301	382	467	445	380	231	110	21	-

Hispanic, both sexes

2003.....	98	57	59	57	46	60	38	19	5	1	-	-
2002.....	78	52	33	41	34	33	35	22	7	2	1	-
2001.....	29	34	22	31	36	37	20	11	6	6	1	-
2000.....	31	30	24	27	35	31	26	19	6	4	-	-
1999.....	35	27	35	47	33	43	25	23	9	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Hispanic, male

2003.....	76	44	32	37	16	27	16	9	1	1	-	-
2002.....	59	34	18	24	16	15	15	12	3	-	-	-
2001.....	21	25	13	19	19	17	10	3	2	3	-	-
2000.....	17	21	14	10	18	12	15	10	-	1	-	-
1999.....	23	9	20	24	12	21	8	7	2	1	-	-

Hispanic, female

2003.....	22	13	27	20	30	33	22	10	4	-	-	-
2002.....	19	18	15	17	18	18	20	10	4	2	1	-
2001.....	8	9	9	12	17	20	10	8	4	3	1	-
2000.....	14	9	10	17	17	19	11	9	6	3	-	-
1999.....	12	18	15	23	21	22	17	16	7	4	-	-

Non-Hispanic 1/, both sexes

2003.....	559	531	487	518	717	851	763	575	330	112	20	-
2002.....	433	392	409	479	669	823	699	557	282	110	15	-
2001.....	325	302	409	505	568	660	612	471	281	92	17	-
2000.....	274	290	359	424	588	704	608	494	258	88	28	-
1999.....	407	397	434	603	755	880	786	610	319	133	22	-

Non-Hispanic 1/, male

2003.....	363	332	281	277	377	401	370	223	122	23	4	-
2002.....	288	239	240	266	357	412	341	217	107	30	3	-
2001.....	192	182	226	256	269	332	296	182	99	18	-	-
2000.....	174	157	190	243	326	352	281	206	96	21	4	-
1999.....	222	208	241	326	396	438	359	246	96	27	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic 1/, female

2003.....	196	199	206	241	340	450	393	352	208	89	16	-
2002.....	145	153	169	213	312	411	358	340	175	80	12	-
2001.....	133	120	183	249	299	328	316	289	182	74	17	-
2000.....	100	133	169	181	262	352	327	288	162	67	24	-
1999.....	185	189	193	277	359	442	427	364	223	106	21	-

Non-Hispanic White, both sexes

2003.....	425	412	382	434	622	720	679	524	305	107	19	-
2002.....	315	285	321	403	570	716	628	514	253	102	13	-
2001.....	241	227	324	417	482	577	546	426	254	82	13	-
2000.....	200	232	273	353	498	621	544	451	229	72	25	-
1999.....	282	300	328	488	638	751	687	542	278	121	18	-

Non-Hispanic White, male

2003.....	277	259	219	232	322	346	329	203	114	23	4	-
2002.....	213	169	189	230	307	360	305	202	96	27	2	-
2001.....	144	139	181	211	232	294	268	166	90	16	-	-
2000.....	128	128	147	201	282	304	253	187	81	16	3	-
1999.....	158	166	179	266	342	382	299	214	80	25	-	-

Non-Hispanic White, female

2003.....	148	153	163	202	300	374	350	321	191	84	15	-
2002.....	102	116	132	173	263	356	323	312	157	75	11	-
2001.....	97	88	143	206	250	283	278	260	164	66	13	-
2000.....	72	104	126	152	216	317	291	264	148	56	22	-
1999.....	124	134	149	222	296	369	388	328	198	96	18	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Non-Hispanic Black, both sexes

2003.....	110	93	72	54	62	95	62	38	24	4	1	-
2002.....	100	89	62	50	72	87	43	35	27	8	2	-
2001.....	68	62	67	64	71	56	49	35	20	8	4	-
2000.....	63	52	64	55	74	64	56	33	25	15	3	-
1999.....	103	83	86	96	96	105	81	54	36	12	4	-

Non-Hispanic Black, male

2003.....	68	54	43	34	39	36	30	14	8	-	-	-
2002.....	65	62	37	26	36	39	19	12	10	3	1	-
2001.....	39	34	36	37	29	22	20	10	7	2	-	-
2000.....	40	27	35	33	37	35	23	10	11	5	1	-
1999.....	52	34	48	47	47	46	48	24	12	2	1	-

Non-Hispanic Black, female

2003.....	42	39	29	20	23	59	32	24	16	4	1	-
2002.....	35	27	25	24	36	48	24	23	17	5	1	-
2001.....	29	28	31	27	42	34	29	25	13	6	4	-
2000.....	23	25	29	22	37	29	33	23	14	10	2	-
1999.....	51	49	38	49	49	59	33	30	24	10	3	-

Origin not stated 2/, both sexes

2003.....	2	2	-	2	4	2	1	-	-	-	-	-
2002.....	2	1	2	-	-	3	1	1	-	-	-	-
2001.....	2	2	2	1	2	1	-	1	1	-	-	-
2000.....	3	3	1	-	-	4	2	-	1	1	-	-
1999.....	2	3	2	2	3	4	1	-	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Origin not stated 2/, male

2003.....	2	-	-	2	2	2	-	-	-	-	-	-
2002.....	1	-	1	-	-	2	-	1	-	-	-	-
2001.....	2	2	1	-	2	1	-	-	-	-	-	-
2000.....	3	1	-	-	-	3	2	-	1	-	-	-
1999.....	2	3	1	1	1	1	-	-	1	-	-	-

Origin not stated 2/, female

2003.....	-	2	-	-	2	-	1	-	-	-	-	-
2002.....	1	1	1	-	-	1	1	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	1	1	-	-	-
2000.....	-	2	1	-	-	1	-	-	-	1	-	-
1999.....	-	-	1	1	2	3	1	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Malignant neoplasms (C00-C97)

##### All origins, both sexes

2003.....	30,291	42,315	53,377	63,141	78,107	89,086	78,531	50,789	22,660	5,827	770	12
2002.....	30,342	41,487	51,904	63,598	81,159	89,565	77,497	50,298	22,323	5,772	789	7
2001.....	30,363	39,827	50,396	64,172	82,846	89,967	75,478	49,680	21,660	5,684	727	8
2000.....	29,231	38,930	50,075	65,843	84,288	90,655	74,444	49,155	21,564	5,653	764	11
1999.....	28,269	38,637	50,430	66,814	85,524	89,874	72,896	48,041	20,702	5,492	779	9

##### All origins, male

2003.....	15,815	22,671	29,256	35,048	42,745	47,516	39,866	23,531	9,124	2,006	217	8
2002.....	15,774	22,407	28,708	35,374	44,711	47,734	39,218	23,345	9,106	1,952	218	5
2001.....	15,694	21,390	27,854	35,900	45,799	47,951	38,175	23,140	8,628	1,951	235	7
2000.....	14,972	20,912	27,515	36,730	46,461	48,284	37,623	22,714	8,506	1,850	190	10
1999.....	14,483	20,891	27,942	37,315	47,229	48,184	36,876	22,147	8,287	1,826	224	6

##### All origins, female

2003.....	14,476	19,644	24,121	28,093	35,362	41,570	38,665	27,258	13,536	3,821	553	4
2002.....	14,568	19,080	23,196	28,224	36,448	41,831	38,279	26,953	13,217	3,820	571	2
2001.....	14,669	18,437	22,542	28,272	37,047	42,016	37,303	26,540	13,032	3,733	492	1
2000.....	14,259	18,018	22,560	29,113	37,827	42,371	36,821	26,441	13,058	3,803	574	1
1999.....	13,786	17,746	22,488	29,499	38,295	41,690	36,020	25,894	12,415	3,666	555	3

##### Hispanic, both sexes

2003.....	1,765	2,152	2,330	2,789	3,140	3,248	2,453	1,450	728	181	24	1
2002.....	1,669	1,971	2,276	2,712	3,145	3,120	2,245	1,446	707	188	20	-
2001.....	1,629	1,814	2,312	2,608	3,116	2,988	2,080	1,322	615	222	22	-
2000.....	1,434	1,735	2,045	2,532	3,025	2,783	2,044	1,292	586	183	24	-
1999.....	1,423	1,635	2,051	2,598	2,796	2,586	1,863	1,231	588	150	26	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Malignant neoplasms (C00-C97)

##### Hispanic, male

2003.....	903	1,134	1,308	1,568	1,779	1,751	1,284	656	342	75	11	-
2002.....	864	1,116	1,261	1,532	1,724	1,737	1,165	700	319	92	6	-
2001.....	793	948	1,254	1,465	1,754	1,718	1,096	642	271	109	11	-
2000.....	692	911	1,123	1,398	1,688	1,545	1,112	632	292	88	5	-
1999.....	698	901	1,159	1,469	1,576	1,436	967	564	271	59	8	-

##### Hispanic, female

2003.....	862	1,018	1,022	1,221	1,361	1,497	1,169	794	386	106	13	1
2002.....	805	855	1,015	1,180	1,421	1,383	1,080	746	388	96	14	-
2001.....	836	866	1,058	1,143	1,362	1,270	984	680	344	113	11	-
2000.....	742	824	922	1,134	1,337	1,238	932	660	294	95	19	-
1999.....	725	734	892	1,129	1,220	1,150	896	667	317	91	18	-

##### Non-Hispanic 1/, both sexes

2003.....	28,432	40,041	50,932	60,213	74,801	85,672	75,934	49,236	21,894	5,640	745	11
2002.....	28,585	39,406	49,480	60,718	77,827	86,232	75,071	48,738	21,564	5,569	768	5
2001.....	28,656	37,907	47,935	61,421	79,522	86,761	73,262	48,251	20,990	5,451	702	7
2000.....	27,703	37,088	47,889	63,135	81,065	87,661	72,245	47,750	20,926	5,456	739	11
1999.....	26,771	36,902	48,244	64,056	82,516	87,052	70,852	46,695	20,070	5,337	750	9

##### Non-Hispanic 1/, male

2003.....	14,852	21,472	27,881	33,391	40,869	45,673	38,506	22,829	8,763	1,928	206	8
2002.....	14,865	21,232	27,355	33,742	42,879	45,877	37,962	22,595	8,763	1,857	212	4
2001.....	14,856	20,368	26,503	34,350	43,916	46,112	37,009	22,450	8,335	1,839	222	6
2000.....	14,227	19,938	26,315	35,223	44,655	46,620	36,427	22,037	8,194	1,755	185	10
1999.....	13,742	19,936	26,708	35,752	45,519	46,616	35,827	21,532	7,999	1,765	216	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

Non-Hispanic 1/, female

2003.....	13,580	18,569	23,051	26,822	33,932	39,999	37,428	26,407	13,131	3,712	539	3
2002.....	13,720	18,174	22,125	26,976	34,948	40,355	37,109	26,143	12,801	3,712	556	1
2001.....	13,800	17,539	21,432	27,071	35,606	40,649	36,253	25,801	12,655	3,612	480	1
2000.....	13,476	17,150	21,574	27,912	36,410	41,041	35,818	25,713	12,732	3,701	554	1
1999.....	13,029	16,966	21,536	28,304	36,997	40,436	35,025	25,163	12,071	3,572	534	3

Non-Hispanic White, both sexes

2003.....	22,251	32,981	42,752	51,177	65,021	75,855	68,157	44,609	19,783	4,977	612	10
2002.....	22,639	32,418	41,391	51,562	67,902	76,462	67,495	44,205	19,422	4,915	642	2
2001.....	22,907	31,194	40,195	52,117	69,412	77,026	65,734	43,739	18,926	4,811	577	4
2000.....	22,263	30,664	40,201	54,052	70,790	77,899	65,180	43,210	18,871	4,800	606	7
1999.....	21,540	30,407	40,381	54,833	72,286	77,288	64,007	42,207	18,092	4,716	621	7

Non-Hispanic White, male

2003.....	11,631	17,674	23,352	28,381	35,638	40,432	34,536	20,704	7,872	1,660	168	8
2002.....	11,787	17,454	22,756	28,750	37,500	40,610	34,029	20,477	7,837	1,629	169	2
2001.....	11,839	16,696	22,188	29,183	38,371	40,816	33,069	20,248	7,441	1,599	183	4
2000.....	11,347	16,345	22,023	30,179	38,906	41,247	32,705	19,854	7,309	1,507	144	7
1999.....	10,966	16,371	22,287	30,626	39,892	41,239	32,296	19,315	7,148	1,519	179	4

Non-Hispanic White, female

2003.....	10,620	15,307	19,400	22,796	29,383	35,423	33,621	23,905	11,911	3,317	444	2
2002.....	10,852	14,964	18,635	22,812	30,402	35,852	33,466	23,728	11,585	3,286	473	-
2001.....	11,068	14,498	18,007	22,934	31,041	36,210	32,665	23,491	11,485	3,212	394	-
2000.....	10,916	14,319	18,178	23,873	31,884	36,652	32,475	23,356	11,562	3,293	462	-
1999.....	10,574	14,036	18,094	24,207	32,394	36,049	31,711	22,892	10,944	3,197	442	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

Non-Hispanic Black, both sexes

2003.....	5,198	5,990	6,913	7,577	8,089	8,071	6,411	3,814	1,789	560	115	-
2002.....	4,956	5,899	6,794	7,759	8,325	8,221	6,297	3,818	1,819	550	109	3
2001.....	4,805	5,694	6,502	7,877	8,515	8,163	6,291	3,776	1,780	567	116	3
2000.....	4,670	5,447	6,572	7,733	8,752	8,264	5,980	3,860	1,769	591	117	3
1999.....	4,471	5,587	6,735	7,888	8,737	8,347	5,845	3,905	1,703	531	121	1

Non-Hispanic Black, male

2003.....	2,766	3,287	3,848	4,208	4,385	4,289	3,251	1,710	731	216	27	-
2002.....	2,591	3,223	3,883	4,276	4,559	4,413	3,256	1,755	770	187	38	2
2001.....	2,568	3,133	3,619	4,430	4,704	4,432	3,268	1,830	734	203	37	2
2000.....	2,504	3,079	3,684	4,332	4,903	4,532	3,157	1,811	734	219	38	2
1999.....	2,411	3,117	3,816	4,423	4,805	4,603	3,000	1,887	710	198	33	1

Non-Hispanic Black, female

2003.....	2,432	2,703	3,065	3,369	3,704	3,782	3,160	2,104	1,058	344	88	-
2002.....	2,365	2,676	2,911	3,483	3,766	3,808	3,041	2,063	1,049	363	71	1
2001.....	2,237	2,561	2,883	3,447	3,811	3,731	3,023	1,946	1,046	364	79	1
2000.....	2,166	2,368	2,888	3,401	3,849	3,732	2,823	2,049	1,035	372	79	1
1999.....	2,060	2,470	2,919	3,465	3,932	3,744	2,845	2,018	993	333	88	-

Origin not stated 2/, both sexes

2003.....	94	122	115	139	166	166	144	103	38	6	1	-
2002.....	88	110	148	168	187	213	181	114	52	15	1	2
2001.....	78	106	149	143	208	218	136	107	55	11	3	1
2000.....	94	107	141	176	198	211	155	113	52	14	1	-
1999.....	75	100	135	160	212	236	181	115	44	5	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms (C00-C97)

Origin not stated 2/, male

2003.....	60	65	67	89	97	92	76	46	19	3	-	-
2002.....	45	59	92	100	108	120	91	50	24	3	-	1
2001.....	45	74	97	85	129	121	70	48	22	3	2	1
2000.....	53	63	77	109	118	119	84	45	20	7	-	-
1999.....	43	54	75	94	134	132	82	51	17	2	-	-

Origin not stated 2/, female

2003.....	34	57	48	50	69	74	68	57	19	3	1	-
2002.....	43	51	56	68	79	93	90	64	28	12	1	1
2001.....	33	32	52	58	79	97	66	59	33	8	1	-
2000.....	41	44	64	67	80	92	71	68	32	7	1	-
1999.....	32	46	60	66	78	104	99	64	27	3	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins, both sexes

2003.....	686	867	959	950	924	955	737	543	242	78	16	1
2002.....	670	875	936	846	1,052	973	748	505	262	77	13	-
2001.....	649	861	879	893	1,019	1,004	783	502	240	67	5	1
2000.....	660	770	872	942	1,017	933	693	488	249	74	13	-
1999.....	612	753	863	889	1,007	993	751	501	220	64	15	-

All origins, male

2003.....	575	694	729	678	626	589	410	255	108	20	6	1
2002.....	534	681	711	619	698	590	407	245	101	24	4	-
2001.....	521	687	685	631	696	598	427	232	83	18	2	1
2000.....	516	618	652	667	659	569	370	217	96	19	-	-
1999.....	474	568	663	629	679	605	402	229	79	14	1	-

All origins, female

2003.....	111	173	230	272	298	366	327	288	134	58	10	-
2002.....	136	194	225	227	354	383	341	260	161	53	9	-
2001.....	128	174	194	262	323	406	356	270	157	49	3	-
2000.....	144	152	220	275	358	364	323	271	153	55	13	-
1999.....	138	185	200	260	328	388	349	272	141	50	14	-

Hispanic, both sexes

2003.....	25	38	37	46	34	38	25	16	9	1	3	-
2002.....	31	38	28	30	32	34	20	14	11	3	1	-
2001.....	31	29	40	41	37	39	19	21	6	4	-	-
2000.....	27	40	35	40	33	27	21	16	10	2	-	-
1999.....	22	29	37	37	35	36	27	13	6	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Hispanic, male

2003.....	23	33	31	35	21	28	15	8	5	-	1	-
2002.....	25	32	21	23	28	25	11	11	5	2	-	-
2001.....	24	23	32	31	27	26	13	8	2	2	-	-
2000.....	21	34	28	32	26	15	13	9	9	2	-	-
1999.....	17	27	28	33	29	27	8	7	2	-	-	-

Hispanic, female

2003.....	2	5	6	11	13	10	10	8	4	1	2	-
2002.....	6	6	7	7	4	9	9	3	6	1	1	-
2001.....	7	6	8	10	10	13	6	13	4	2	-	-
2000.....	6	6	7	8	7	12	8	7	1	-	-	-
1999.....	5	2	9	4	6	9	19	6	4	-	1	-

Non-Hispanic 1/, both sexes

2003.....	659	825	918	904	887	916	711	525	231	77	13	1
2002.....	636	837	902	815	1,015	938	728	490	251	74	12	-
2001.....	616	828	838	849	980	960	764	479	234	63	5	-
2000.....	632	727	833	896	981	905	670	472	239	72	13	-
1999.....	589	720	822	849	969	954	724	487	214	64	12	-

Non-Hispanic 1/, male

2003.....	550	657	694	643	603	560	395	247	102	20	5	1
2002.....	506	649	686	595	666	564	396	233	96	22	4	-
2001.....	496	661	652	597	667	568	414	223	81	16	2	-
2000.....	494	582	622	631	632	553	355	208	87	17	-	-
1999.....	456	538	634	593	648	576	394	221	77	14	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic 1/, female

2003.....	109	168	224	261	284	356	316	278	129	57	8	-
2002.....	130	188	216	220	349	374	332	257	155	52	8	-
2001.....	120	167	186	252	313	392	350	256	153	47	3	-
2000.....	138	145	211	265	349	352	315	264	152	55	13	-
1999.....	133	182	188	256	321	378	330	266	137	50	11	-

Non-Hispanic White, both sexes

2003.....	477	664	721	736	747	793	626	487	205	72	13	1
2002.....	458	646	712	659	906	843	661	444	236	64	10	-
2001.....	464	632	656	679	845	859	701	440	216	62	5	-
2000.....	448	541	636	733	833	806	606	436	221	65	11	-
1999.....	408	530	650	699	828	837	660	443	207	60	11	-

Non-Hispanic White, male

2003.....	409	525	530	519	495	477	342	226	91	19	5	1
2002.....	363	506	536	476	591	510	357	210	92	17	3	-
2001.....	375	500	504	478	567	502	378	204	75	16	2	-
2000.....	337	434	470	507	529	480	317	196	79	16	-	-
1999.....	320	390	502	487	556	496	354	197	74	13	1	-

Non-Hispanic White, female

2003.....	68	139	191	217	252	316	284	261	114	53	8	-
2002.....	95	140	176	183	315	333	304	234	144	47	7	-
2001.....	89	132	152	201	278	357	323	236	141	46	3	-
2000.....	111	107	166	226	304	326	289	240	142	49	11	-
1999.....	88	140	148	212	272	341	306	246	133	47	10	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Non-Hispanic Black, both sexes

2003.....	138	139	170	147	114	87	63	29	20	3	-	-
2002.....	148	150	168	124	89	66	48	32	12	7	2	-
2001.....	124	172	146	140	116	80	49	25	16	1	-	-
2000.....	155	161	166	133	121	80	56	25	14	6	2	-
1999.....	154	162	154	128	120	96	54	37	5	2	1	-

Non-Hispanic Black, male

2003.....	111	116	144	111	90	58	41	15	9	-	-	-
2002.....	123	122	135	96	61	36	28	17	3	3	1	-
2001.....	100	141	122	98	86	52	26	12	5	-	-	-
2000.....	136	130	132	104	87	60	34	7	7	-	-	-
1999.....	117	125	118	91	78	69	33	19	1	1	-	-

Non-Hispanic Black, female

2003.....	27	23	26	36	24	29	22	14	11	3	-	-
2002.....	25	28	33	28	28	30	20	15	9	4	1	-
2001.....	24	31	24	42	30	28	23	13	11	1	-	-
2000.....	19	31	34	29	34	20	22	18	7	6	2	-
1999.....	37	37	36	37	42	27	21	18	4	1	1	-

Origin not stated 2/, both sexes

2003.....	2	4	4	-	3	1	1	2	2	-	-	-
2002.....	3	-	6	1	5	1	-	1	-	-	-	-
2001.....	2	4	1	3	2	5	-	2	-	-	-	1
2000.....	1	3	4	6	3	1	2	-	-	-	-	-
1999.....	1	4	4	3	3	3	-	1	-	-	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Origin not stated 2/, male

2003.....	2	4	4	-	2	1	-	-	1	-	-	-
2002.....	3	-	4	1	4	1	-	1	-	-	-	-
2001.....	1	3	1	3	2	4	-	1	-	-	-	1
2000.....	1	2	2	4	1	1	2	-	-	-	-	-
1999.....	1	3	1	3	2	2	-	1	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	1	-	1	2	1	-	-	-
2002.....	-	-	2	-	1	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	1	-	1	-	-	-	-
2000.....	-	1	2	2	2	-	-	-	-	-	-	-
1999.....	-	1	3	-	1	1	-	-	-	-	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

All origins, both sexes

2003.....	881	1,389	1,551	1,714	1,905	1,895	1,497	806	306	63	5	-
2002.....	887	1,286	1,477	1,736	1,955	1,909	1,480	766	295	65	4	1
2001.....	916	1,114	1,440	1,765	2,014	1,941	1,366	757	317	56	7	1
2000.....	870	1,180	1,430	1,711	1,960	1,817	1,346	764	274	55	7	-
1999.....	793	1,139	1,392	1,739	1,955	1,784	1,202	755	279	72	8	1

All origins, male

2003.....	757	1,175	1,308	1,385	1,466	1,415	1,049	476	156	22	-	-
2002.....	759	1,099	1,222	1,394	1,558	1,437	1,017	452	127	23	1	-
2001.....	787	940	1,197	1,436	1,553	1,423	942	452	153	18	2	1
2000.....	746	988	1,181	1,349	1,482	1,337	896	437	122	23	4	-
1999.....	684	969	1,112	1,396	1,529	1,328	808	421	127	22	3	1

All origins, female

2003.....	124	214	243	329	439	480	448	330	150	41	5	-
2002.....	128	187	255	342	397	472	463	314	168	42	3	1
2001.....	129	174	243	329	461	518	424	305	164	38	5	-
2000.....	124	192	249	362	478	480	450	327	152	32	3	-
1999.....	109	170	280	343	426	456	394	334	152	50	5	-

Hispanic, both sexes

2003.....	30	57	48	59	63	50	42	21	7	3	-	-
2002.....	31	36	44	58	62	54	37	23	14	5	-	-
2001.....	42	34	62	37	58	49	52	12	18	2	-	-
2000.....	23	37	49	56	64	62	36	20	6	3	1	-
1999.....	22	36	49	56	39	54	33	22	14	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

Hispanic, male

2003.....	26	44	35	47	46	41	33	9	3	1	-	-
2002.....	27	35	40	47	49	41	25	13	8	1	-	-
2001.....	42	25	49	33	47	33	35	7	15	-	-	-
2000.....	21	26	38	44	48	48	24	14	3	2	1	-
1999.....	19	34	44	42	32	42	21	16	7	-	-	-

Hispanic, female

2003.....	4	13	13	12	17	9	9	12	4	2	-	-
2002.....	4	1	4	11	13	13	12	10	6	4	-	-
2001.....	-	9	13	4	11	16	17	5	3	2	-	-
2000.....	2	11	11	12	16	14	12	6	3	1	-	-
1999.....	3	2	5	14	7	12	12	6	7	2	-	-

Non-Hispanic 1/, both sexes

2003.....	848	1,324	1,502	1,653	1,838	1,842	1,455	783	298	60	5	-
2002.....	854	1,243	1,428	1,675	1,885	1,853	1,443	740	278	60	4	1
2001.....	871	1,077	1,370	1,721	1,943	1,888	1,313	745	298	54	7	1
2000.....	845	1,143	1,378	1,649	1,889	1,751	1,307	742	266	52	6	-
1999.....	768	1,100	1,339	1,677	1,910	1,725	1,168	732	264	70	8	1

Non-Hispanic 1/, male

2003.....	728	1,124	1,272	1,336	1,417	1,373	1,016	466	152	21	-	-
2002.....	730	1,058	1,177	1,344	1,504	1,394	992	437	118	22	1	-
2001.....	743	912	1,140	1,396	1,498	1,386	906	445	138	18	2	1
2000.....	724	962	1,140	1,300	1,428	1,285	871	421	118	21	3	-
1999.....	663	932	1,064	1,349	1,492	1,281	787	405	120	22	3	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

Non-Hispanic 1/, female

2003.....	120	200	230	317	421	469	439	317	146	39	5	-
2002.....	124	185	251	331	381	459	451	303	160	38	3	1
2001.....	128	165	230	325	445	502	407	300	160	36	5	-
2000.....	121	181	238	349	461	466	436	321	148	31	3	-
1999.....	105	168	275	328	418	444	381	327	144	48	5	-

Non-Hispanic White, both sexes

2003.....	655	1,083	1,231	1,367	1,625	1,658	1,313	729	278	52	5	-
2002.....	694	1,002	1,171	1,384	1,643	1,645	1,316	679	261	56	3	-
2001.....	674	823	1,102	1,413	1,647	1,690	1,198	688	268	51	7	1
2000.....	651	897	1,100	1,380	1,637	1,535	1,179	687	245	48	6	-
1999.....	596	860	1,063	1,390	1,648	1,508	1,068	673	244	65	5	-

Non-Hispanic White, male

2003.....	577	941	1,063	1,112	1,271	1,251	939	437	143	16	-	-
2002.....	606	866	983	1,137	1,328	1,261	910	403	113	21	-	-
2001.....	581	713	932	1,162	1,296	1,255	831	413	120	17	2	1
2000.....	567	766	936	1,110	1,256	1,136	805	396	108	17	3	-
1999.....	524	739	859	1,148	1,303	1,127	730	369	108	18	2	-

Non-Hispanic White, female

2003.....	78	142	168	255	354	407	374	292	135	36	5	-
2002.....	88	136	188	247	315	384	406	276	148	35	3	-
2001.....	93	110	170	251	351	435	367	275	148	34	5	-
2000.....	84	131	164	270	381	399	374	291	137	31	3	-
1999.....	72	121	204	242	345	381	338	304	136	47	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

Non-Hispanic Black, both sexes

2003.....	177	219	244	249	186	157	116	49	16	7	-	-
2002.....	143	221	233	258	217	182	100	52	12	3	1	1
2001.....	172	225	241	272	262	176	102	51	23	3	-	-
2000.....	182	228	253	243	219	201	106	48	19	4	-	-
1999.....	162	222	250	257	229	198	82	49	17	5	2	1

Non-Hispanic Black, male

2003.....	136	164	187	189	124	103	59	25	7	4	-	-
2002.....	109	179	172	181	159	115	63	30	3	1	1	-
2001.....	141	176	187	202	177	118	67	27	13	1	-	-
2000.....	149	185	184	171	147	137	54	21	8	4	-	-
1999.....	130	177	188	176	164	141	42	28	10	4	-	1

Non-Hispanic Black, female

2003.....	41	55	57	60	62	54	57	24	9	3	-	-
2002.....	34	42	61	77	58	67	37	22	9	2	-	1
2001.....	31	49	54	70	85	58	35	24	10	2	-	-
2000.....	33	43	69	72	72	64	52	27	11	-	-	-
1999.....	32	45	62	81	65	57	40	21	7	1	2	-

Origin not stated 2/, both sexes

2003.....	3	8	1	2	4	3	-	2	1	-	-	-
2002.....	2	7	5	3	8	2	-	3	3	-	-	-
2001.....	3	3	8	7	13	4	1	-	1	-	-	-
2000.....	2	-	3	6	7	4	3	2	2	-	-	-
1999.....	3	3	4	6	6	5	1	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of esophagus (C15)

Origin not stated 2/, male

2003.....	3	7	1	2	3	1	-	1	1	-	-	-
2002.....	2	6	5	3	5	2	-	2	1	-	-	-
2001.....	2	3	8	7	8	4	1	-	-	-	-	-
2000.....	1	-	3	5	6	4	1	2	1	-	-	-
1999.....	2	3	4	5	5	5	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	1	-	-	1	2	-	1	-	-	-	-
2002.....	-	1	-	-	3	-	-	1	2	-	-	-
2001.....	1	-	-	-	5	-	-	-	1	-	-	-
2000.....	1	-	-	1	1	-	2	-	1	-	-	-
1999.....	1	-	-	1	1	-	1	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

All origins, both sexes

2003.....	717	830	1,044	1,211	1,582	1,805	1,744	1,263	619	161	14	-
2002.....	663	841	990	1,179	1,637	1,903	1,753	1,344	664	153	24	-
2001.....	675	751	1,020	1,308	1,713	2,004	1,789	1,216	620	149	16	-
2000.....	629	753	1,051	1,317	1,737	2,009	1,824	1,379	641	188	26	-
1999.....	651	808	1,052	1,359	1,793	2,064	1,796	1,277	607	170	29	-

All origins, male

2003.....	462	574	708	810	1,016	1,061	919	574	238	51	4	-
2002.....	440	556	659	804	1,063	1,118	946	628	258	38	10	-
2001.....	443	514	724	858	1,070	1,194	954	550	222	36	5	-
2000.....	413	506	722	860	1,106	1,201	943	617	247	56	8	-
1999.....	443	558	694	893	1,112	1,250	989	568	223	52	8	-

All origins, female

2003.....	255	256	336	401	566	744	825	689	381	110	10	-
2002.....	223	285	331	375	574	785	807	716	406	115	14	-
2001.....	232	237	296	450	643	810	835	666	398	113	11	-
2000.....	216	247	329	457	631	808	881	762	394	132	18	-
1999.....	208	250	358	466	681	814	807	709	384	118	21	-

Hispanic, both sexes

2003.....	105	113	126	135	156	170	109	75	38	8	-	-
2002.....	83	97	121	111	148	162	121	73	42	15	-	-
2001.....	104	93	116	135	165	142	111	60	30	11	1	-
2000.....	78	68	99	139	167	134	120	65	26	8	-	-
1999.....	71	83	109	134	137	137	102	63	32	6	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

Hispanic, male

2003.....	57	67	73	77	103	102	64	30	17	1	-	-
2002.....	43	59	69	71	101	87	63	39	20	5	-	-
2001.....	61	51	79	86	100	76	59	32	10	7	1	-
2000.....	43	43	67	93	112	73	71	34	12	3	-	-
1999.....	41	55	71	86	76	87	54	22	9	2	2	-

Hispanic, female

2003.....	48	46	53	58	53	68	45	45	21	7	-	-
2002.....	40	38	52	40	47	75	58	34	22	10	-	-
2001.....	43	42	37	49	65	66	52	28	20	4	-	-
2000.....	35	25	32	46	55	61	49	31	14	5	-	-
1999.....	30	28	38	48	61	50	48	41	23	4	2	-

Non-Hispanic 1/, both sexes

2003.....	610	715	915	1,072	1,424	1,630	1,630	1,186	579	153	14	-
2002.....	578	742	867	1,064	1,484	1,736	1,627	1,264	619	137	24	-
2001.....	569	656	902	1,171	1,541	1,853	1,676	1,155	588	137	15	-
2000.....	549	681	946	1,173	1,566	1,869	1,699	1,309	613	180	25	-
1999.....	575	724	938	1,221	1,646	1,924	1,691	1,212	572	164	25	-

Non-Hispanic 1/, male

2003.....	403	506	632	730	911	957	851	543	220	50	4	-
2002.....	396	495	589	729	959	1,028	880	585	236	33	10	-
2001.....	380	461	644	771	966	1,114	893	517	211	28	4	-
2000.....	368	461	652	762	991	1,123	871	582	234	53	8	-
1999.....	399	502	620	805	1,029	1,162	934	545	213	50	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

Non-Hispanic 1/, female

2003.....	207	209	283	342	513	673	779	643	359	103	10	-
2002.....	182	247	278	335	525	708	747	679	383	104	14	-
2001.....	189	195	258	400	575	739	783	638	377	109	11	-
2000.....	181	220	294	411	575	746	828	727	379	127	17	-
1999.....	176	222	318	416	617	762	757	667	359	114	19	-

Non-Hispanic White, both sexes

2003.....	395	510	662	788	1,073	1,248	1,292	952	461	106	8	-
2002.....	370	525	612	753	1,108	1,331	1,282	1,011	496	100	14	-
2001.....	371	474	636	827	1,183	1,448	1,331	948	471	105	10	-
2000.....	383	491	689	850	1,181	1,436	1,345	1,061	497	141	13	-
1999.....	379	502	654	930	1,251	1,499	1,360	976	463	125	15	-

Non-Hispanic White, male

2003.....	266	369	456	533	691	740	684	429	183	35	4	-
2002.....	264	366	425	534	731	815	708	481	187	24	4	-
2001.....	252	335	459	540	748	894	712	431	172	22	2	-
2000.....	266	336	489	567	763	877	703	476	192	42	4	-
1999.....	270	362	436	626	800	916	754	430	174	40	4	-

Non-Hispanic White, female

2003.....	129	141	206	255	382	508	608	523	278	71	4	-
2002.....	106	159	187	219	377	516	574	530	309	76	10	-
2001.....	119	139	177	287	435	554	619	517	299	83	8	-
2000.....	117	155	200	283	418	559	642	585	305	99	9	-
1999.....	109	140	218	304	451	583	606	546	289	85	11	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

Non-Hispanic Black, both sexes

2003.....	157	151	196	210	248	285	243	155	93	35	6	-
2002.....	156	155	184	229	289	306	239	196	96	30	9	-
2001.....	147	128	196	251	254	292	256	161	94	26	5	-
2000.....	130	147	188	239	284	316	263	196	86	30	11	-
1999.....	148	170	203	212	295	326	254	193	85	32	9	-

Non-Hispanic Black, male

2003.....	107	105	134	146	164	164	115	71	21	9	-	-
2002.....	105	102	121	145	178	159	117	73	40	7	5	-
2001.....	104	86	134	172	160	162	131	59	25	6	2	-
2000.....	83	98	124	145	169	173	122	80	27	8	3	-
1999.....	98	109	132	134	175	188	137	89	28	7	2	-

Non-Hispanic Black, female

2003.....	50	46	62	64	84	121	128	84	72	26	6	-
2002.....	51	53	63	84	111	147	122	123	56	23	4	-
2001.....	43	42	62	79	94	130	125	102	69	20	3	-
2000.....	47	49	64	94	115	143	141	116	59	22	8	-
1999.....	50	61	71	78	120	138	117	104	57	25	7	-

Origin not stated 2/, both sexes

2003.....	2	2	3	4	2	5	5	2	2	-	-	-
2002.....	2	2	2	4	5	5	5	7	3	1	-	-
2001.....	2	2	2	2	7	9	2	1	2	1	-	-
2000.....	2	4	6	5	4	6	5	5	2	-	1	-
1999.....	5	1	5	4	10	3	3	2	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of stomach (C16)

Origin not stated 2/, male

2003.....	2	1	3	3	2	2	4	1	1	-	-	-
2002.....	1	2	1	4	3	3	3	4	2	-	-	-
2001.....	2	2	1	1	4	4	2	1	1	1	-	-
2000.....	2	2	3	5	3	5	1	1	1	-	-	-
1999.....	3	1	3	2	7	1	1	1	1	-	-	-

Origin not stated 2/, female

2003.....	-	1	-	1	-	3	1	1	1	-	-	-
2002.....	1	-	1	-	2	2	2	3	1	1	-	-
2001.....	-	-	1	1	3	5	-	-	1	-	-	-
2000.....	-	2	3	-	1	1	4	4	1	-	1	-
1999.....	2	-	2	2	3	2	2	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins, both sexes

2003.....	2,718	3,631	4,673	5,775	7,159	8,720	8,611	6,590	3,547	1,024	135	1
2002.....	2,818	3,694	4,727	5,901	7,491	8,826	8,672	6,453	3,433	1,075	128	1
2001.....	2,804	3,610	4,374	5,997	7,794	8,988	8,539	6,618	3,482	1,028	131	-
2000.....	2,581	3,397	4,543	5,913	8,188	9,410	8,643	6,735	3,518	1,061	148	-
1999.....	2,434	3,309	4,658	6,095	8,154	9,473	8,456	6,672	3,503	973	138	-

All origins, male

2003.....	1,544	2,117	2,714	3,393	3,953	4,433	4,038	2,565	1,112	245	27	1
2002.....	1,606	2,185	2,850	3,455	4,125	4,536	3,968	2,508	1,024	267	24	1
2001.....	1,545	2,116	2,536	3,479	4,294	4,568	3,883	2,520	1,068	251	36	-
2000.....	1,466	1,989	2,619	3,407	4,517	4,852	3,908	2,571	1,012	260	30	-
1999.....	1,374	1,932	2,689	3,431	4,452	4,857	3,897	2,525	1,081	236	24	-

All origins, female

2003.....	1,174	1,514	1,959	2,382	3,206	4,287	4,573	4,025	2,435	779	108	-
2002.....	1,212	1,509	1,877	2,446	3,366	4,290	4,704	3,945	2,409	808	104	-
2001.....	1,259	1,494	1,838	2,518	3,500	4,420	4,656	4,098	2,414	777	95	-
2000.....	1,115	1,408	1,924	2,506	3,671	4,558	4,735	4,164	2,506	801	118	-
1999.....	1,060	1,377	1,969	2,664	3,702	4,616	4,559	4,147	2,422	737	114	-

Hispanic, both sexes

2003.....	182	250	243	286	321	330	282	161	93	24	1	-
2002.....	178	197	257	275	330	338	235	161	108	28	2	-
2001.....	157	176	239	255	337	317	218	169	96	41	4	-
2000.....	152	184	211	242	288	279	218	162	73	32	6	-
1999.....	132	132	212	271	310	274	211	153	72	25	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Hispanic, male

2003.....	111	144	151	167	184	168	146	64	37	11	-	-
2002.....	104	135	156	168	183	185	110	65	29	10	-	-
2001.....	80	99	146	150	183	167	94	76	39	19	1	-
2000.....	87	102	125	147	154	148	119	78	28	11	-	-
1999.....	74	72	113	152	179	147	110	58	31	8	1	-

Hispanic, female

2003.....	71	106	92	119	137	162	136	97	56	13	1	-
2002.....	74	62	101	107	147	153	125	96	79	18	2	-
2001.....	77	77	93	105	154	150	124	93	57	22	3	-
2000.....	65	82	86	95	134	131	99	84	45	21	6	-
1999.....	58	60	99	119	131	127	101	95	41	17	1	-

Non-Hispanic 1/, both sexes

2003.....	2,526	3,375	4,418	5,475	6,829	8,377	8,310	6,414	3,447	998	133	1
2002.....	2,634	3,488	4,460	5,615	7,143	8,466	8,416	6,284	3,312	1,045	126	1
2001.....	2,639	3,421	4,124	5,733	7,439	8,655	8,308	6,434	3,379	985	126	-
2000.....	2,425	3,201	4,322	5,656	7,888	9,114	8,413	6,562	3,436	1,027	142	-
1999.....	2,298	3,168	4,436	5,811	7,829	9,180	8,210	6,494	3,424	948	136	-

Non-Hispanic 1/, male

2003.....	1,425	1,968	2,557	3,218	3,763	4,256	3,882	2,493	1,073	234	27	1
2002.....	1,500	2,045	2,686	3,279	3,929	4,338	3,850	2,441	991	257	24	1
2001.....	1,462	2,008	2,382	3,322	4,097	4,396	3,785	2,441	1,028	232	34	-
2000.....	1,378	1,885	2,488	3,250	4,359	4,694	3,782	2,492	980	249	30	-
1999.....	1,297	1,854	2,570	3,271	4,267	4,699	3,768	2,458	1,047	228	23	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic 1/, female

2003.....	1,101	1,407	1,861	2,257	3,066	4,121	4,428	3,921	2,374	764	106	-
2002.....	1,134	1,443	1,774	2,336	3,214	4,128	4,566	3,843	2,321	788	102	-
2001.....	1,177	1,413	1,742	2,411	3,342	4,259	4,523	3,993	2,351	753	92	-
2000.....	1,047	1,316	1,834	2,406	3,529	4,420	4,631	4,070	2,456	778	112	-
1999.....	1,001	1,314	1,866	2,540	3,562	4,481	4,442	4,036	2,377	720	113	-

Non-Hispanic White, both sexes

2003.....	1,875	2,650	3,591	4,562	5,735	7,283	7,417	5,743	3,115	891	113	1
2002.....	2,060	2,788	3,613	4,658	6,124	7,384	7,437	5,694	2,954	925	108	1
2001.....	2,049	2,721	3,371	4,747	6,377	7,509	7,394	5,839	3,044	866	101	-
2000.....	1,897	2,546	3,539	4,739	6,804	8,034	7,485	5,923	3,100	899	123	-
1999.....	1,804	2,540	3,644	4,874	6,769	8,085	7,342	5,856	3,121	839	107	-

Non-Hispanic White, male

2003.....	1,082	1,588	2,092	2,719	3,223	3,734	3,489	2,256	964	204	23	1
2002.....	1,183	1,661	2,198	2,763	3,426	3,805	3,406	2,205	883	227	18	1
2001.....	1,173	1,630	1,986	2,793	3,576	3,871	3,392	2,211	922	198	28	-
2000.....	1,095	1,507	2,054	2,752	3,801	4,178	3,361	2,249	874	207	23	-
1999.....	1,045	1,500	2,138	2,791	3,768	4,150	3,406	2,215	957	203	21	-

Non-Hispanic White, female

2003.....	793	1,062	1,499	1,843	2,512	3,549	3,928	3,487	2,151	687	90	-
2002.....	877	1,127	1,415	1,895	2,698	3,579	4,031	3,489	2,071	698	90	-
2001.....	876	1,091	1,385	1,954	2,801	3,638	4,002	3,628	2,122	668	73	-
2000.....	802	1,039	1,485	1,987	3,003	3,856	4,124	3,674	2,226	692	100	-
1999.....	759	1,040	1,506	2,083	3,001	3,935	3,936	3,641	2,164	636	86	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Non-Hispanic Black, both sexes

2003.....	545	615	698	765	937	908	772	555	289	90	17	-
2002.....	490	591	706	809	847	927	829	505	308	94	13	-
2001.....	492	607	641	839	887	985	773	503	296	104	23	-
2000.....	462	534	681	782	938	936	796	556	293	117	18	-
1999.....	410	547	701	808	916	958	775	573	266	92	24	-

Non-Hispanic Black, male

2003.....	295	323	393	414	473	427	325	182	90	28	2	-
2002.....	268	317	400	433	407	452	370	195	86	19	4	-
2001.....	242	325	327	449	432	442	324	187	86	27	5	-
2000.....	250	309	367	417	476	440	352	198	81	39	7	-
1999.....	216	315	377	420	430	475	311	205	76	15	-	-

Non-Hispanic Black, female

2003.....	250	292	305	351	464	481	447	373	199	62	15	-
2002.....	222	274	306	376	440	475	459	310	222	75	9	-
2001.....	250	282	314	390	455	543	449	316	210	77	18	-
2000.....	212	225	314	365	462	496	444	358	212	78	11	-
1999.....	194	232	324	388	486	483	464	368	190	77	24	-

Origin not stated 2/, both sexes

2003.....	10	6	12	14	9	13	19	15	7	2	1	-
2002.....	6	9	10	11	18	22	21	8	13	2	-	-
2001.....	8	13	11	9	18	16	13	15	7	2	1	-
2000.....	4	12	10	15	12	17	12	11	9	2	-	-
1999.....	4	9	10	13	15	19	35	25	7	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Origin not stated 2/, male

2003.....	8	5	6	8	6	9	10	8	2	-	-	-
2002.....	2	5	8	8	13	13	8	2	4	-	-	-
2001.....	3	9	8	7	14	5	4	3	1	-	1	-
2000.....	1	2	6	10	4	10	7	1	4	-	-	-
1999.....	3	6	6	8	6	11	19	9	3	-	-	-

Origin not stated 2/, female

2003.....	2	1	6	6	3	4	9	7	5	2	1	-
2002.....	4	4	2	3	5	9	13	6	9	2	-	-
2001.....	5	4	3	2	4	11	9	12	6	2	-	-
2000.....	3	10	4	5	8	7	5	10	5	2	-	-
1999.....	1	3	4	5	9	8	16	16	4	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins, both sexes

2003.....	1,408	1,348	1,417	1,612	2,018	2,208	1,741	1,056	396	113	11	-
2002.....	1,245	1,208	1,350	1,618	1,963	2,049	1,679	1,094	419	80	16	-
2001.....	1,073	1,063	1,288	1,556	1,884	2,003	1,646	1,009	386	97	8	-
2000.....	944	1,040	1,216	1,526	1,913	2,011	1,547	943	359	81	4	-
1999.....	826	928	1,163	1,558	1,933	1,914	1,511	863	325	75	7	-

All origins, male

2003.....	1,136	999	996	1,054	1,276	1,354	947	495	154	36	2	-
2002.....	1,015	899	959	1,100	1,267	1,212	900	463	155	22	2	-
2001.....	872	780	911	1,036	1,188	1,200	847	481	140	31	2	-
2000.....	729	761	846	1,039	1,207	1,196	856	407	114	26	1	-
1999.....	628	671	794	1,041	1,243	1,103	774	352	109	17	-	-

All origins, female

2003.....	272	349	421	558	742	854	794	561	242	77	9	-
2002.....	230	309	391	518	696	837	779	631	264	58	14	-
2001.....	201	283	377	520	696	803	799	528	246	66	6	-
2000.....	215	279	370	487	706	815	691	536	245	55	3	-
1999.....	198	257	369	517	690	811	737	511	216	58	7	-

Hispanic, both sexes

2003.....	176	175	162	205	192	193	126	63	25	9	3	-
2002.....	122	146	164	198	205	165	100	68	38	2	-	-
2001.....	119	127	134	157	170	179	110	59	21	4	-	-
2000.....	84	111	123	145	190	150	112	61	18	9	-	-
1999.....	94	89	125	169	155	135	104	49	20	5	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Hispanic, male

2003.....	149	123	118	119	126	114	58	24	9	2	1	-
2002.....	105	114	111	130	126	97	49	31	13	1	-	-
2001.....	91	96	91	100	106	100	52	24	9	1	-	-
2000.....	69	82	83	95	113	88	64	23	10	3	-	-
1999.....	83	69	90	94	101	72	49	18	12	1	-	-

Hispanic, female

2003.....	27	52	44	86	66	79	68	39	16	7	2	-
2002.....	17	32	53	68	79	68	51	37	25	1	-	-
2001.....	28	31	43	57	64	79	58	35	12	3	-	-
2000.....	15	29	40	50	77	62	48	38	8	6	-	-
1999.....	11	20	35	75	54	63	55	31	8	4	2	-

Non-Hispanic 1/, both sexes

2003.....	1,227	1,169	1,249	1,398	1,824	2,013	1,613	992	370	104	8	-
2002.....	1,119	1,058	1,176	1,416	1,755	1,878	1,574	1,022	381	78	16	-
2001.....	951	929	1,150	1,395	1,708	1,817	1,535	947	364	93	7	-
2000.....	856	929	1,089	1,375	1,716	1,857	1,430	881	341	72	4	-
1999.....	732	833	1,036	1,385	1,773	1,777	1,403	812	304	70	5	-

Non-Hispanic 1/, male

2003.....	982	874	873	930	1,148	1,240	888	471	144	34	1	-
2002.....	908	782	839	967	1,139	1,111	846	431	142	21	2	-
2001.....	779	677	818	932	1,077	1,095	794	457	131	30	2	-
2000.....	656	679	761	938	1,090	1,105	788	383	104	23	1	-
1999.....	545	598	702	944	1,137	1,029	723	332	96	16	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic 1/, female

2003.....	245	295	376	468	676	773	725	521	226	70	7	-
2002.....	211	276	337	449	616	767	728	591	239	57	14	-
2001.....	172	252	332	463	631	722	741	490	233	63	5	-
2000.....	200	250	328	437	626	752	642	498	237	49	3	-
1999.....	187	235	334	441	636	748	680	480	208	54	5	-

Non-Hispanic White, both sexes

2003.....	824	837	904	1,065	1,441	1,687	1,384	860	323	81	4	-
2002.....	740	729	884	1,100	1,404	1,603	1,367	904	316	65	11	-
2001.....	644	640	865	1,084	1,393	1,552	1,305	835	323	78	6	-
2000.....	567	647	800	1,061	1,387	1,565	1,259	769	307	61	2	-
1999.....	509	571	769	1,089	1,457	1,524	1,231	715	270	58	5	-

Non-Hispanic White, male

2003.....	651	631	639	721	933	1,051	774	412	122	26	-	-
2002.....	598	537	630	759	913	953	728	387	118	17	1	-
2001.....	525	460	628	752	899	938	694	403	120	26	2	-
2000.....	428	462	558	733	891	957	698	329	95	18	-	-
1999.....	367	411	523	753	949	887	647	301	90	13	-	-

Non-Hispanic White, female

2003.....	173	206	265	344	508	636	610	448	201	55	4	-
2002.....	142	192	254	341	491	650	639	517	198	48	10	-
2001.....	119	180	237	332	494	614	611	432	203	52	4	-
2000.....	139	185	242	328	496	608	561	440	212	43	2	-
1999.....	142	160	246	336	508	637	584	414	180	45	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Non-Hispanic Black, both sexes

2003.....	303	229	215	175	210	175	129	90	30	18	3	-
2002.....	256	227	158	180	204	171	121	75	43	6	2	-
2001.....	212	215	158	182	175	148	151	71	31	10	1	-
2000.....	198	171	173	183	200	161	101	70	21	7	2	-
1999.....	142	156	156	189	206	148	117	69	29	10	-	-

Non-Hispanic Black, male

2003.....	250	169	147	106	128	94	61	38	14	7	1	-
2002.....	208	167	112	126	135	92	66	23	12	3	-	-
2001.....	171	162	104	108	96	81	61	31	6	-	-	-
2000.....	151	134	122	126	123	77	54	33	4	3	1	-
1999.....	114	109	101	122	123	84	49	18	6	2	-	-

Non-Hispanic Black, female

2003.....	53	60	68	69	82	81	68	52	16	11	2	-
2002.....	48	60	46	54	69	79	55	52	31	3	2	-
2001.....	41	53	54	74	79	67	90	40	25	10	1	-
2000.....	47	37	51	57	77	84	47	37	17	4	1	-
1999.....	28	47	55	67	83	64	68	51	23	8	-	-

Origin not stated 2/, both sexes

2003.....	5	4	6	9	2	2	2	1	1	-	-	-
2002.....	4	4	10	4	3	6	5	4	-	-	-	-
2001.....	3	7	4	4	6	7	1	3	1	-	1	-
2000.....	4	-	4	6	7	4	5	1	-	-	-	-
1999.....	-	6	2	4	5	2	4	2	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Origin not stated 2/, male

2003.....	5	2	5	5	2	-	1	-	1	-	-	-
2002.....	2	3	9	3	2	4	5	1	-	-	-	-
2001.....	2	7	2	4	5	5	1	-	-	-	-	-
2000.....	4	-	2	6	4	3	4	1	-	-	-	-
1999.....	-	4	2	3	5	2	2	2	1	-	-	-

Origin not stated 2/, female

2003.....	-	2	1	4	-	2	1	1	-	-	-	-
2002.....	2	1	1	1	1	2	-	3	-	-	-	-
2001.....	1	-	2	-	1	2	-	3	1	-	1	-
2000.....	-	-	2	-	3	1	1	-	-	-	-	-
1999.....	-	2	-	1	-	-	2	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

All origins, both sexes

2003.....	1,649	2,327	2,993	3,550	4,554	5,038	4,670	2,897	1,264	293	32	-
2002.....	1,544	2,321	2,784	3,564	4,623	5,210	4,345	2,824	1,217	263	36	1
2001.....	1,577	2,111	2,757	3,525	4,633	4,987	4,335	2,861	1,214	293	42	1
2000.....	1,528	2,030	2,678	3,546	4,557	5,223	4,236	2,571	1,196	286	34	1
1999.....	1,466	2,039	2,668	3,632	4,661	4,931	4,002	2,673	1,120	309	40	1

All origins, male

2003.....	992	1,391	1,715	1,924	2,327	2,358	2,044	992	327	55	8	-
2002.....	971	1,394	1,618	1,924	2,346	2,424	1,846	985	351	60	7	1
2001.....	964	1,269	1,605	1,921	2,294	2,360	1,831	984	310	64	7	1
2000.....	955	1,206	1,543	1,965	2,224	2,447	1,742	891	310	59	2	1
1999.....	907	1,204	1,517	1,976	2,371	2,281	1,659	910	311	59	3	-

All origins, female

2003.....	657	936	1,278	1,626	2,227	2,680	2,626	1,905	937	238	24	-
2002.....	573	927	1,166	1,640	2,277	2,786	2,499	1,839	866	203	29	-
2001.....	613	842	1,152	1,604	2,339	2,627	2,504	1,877	904	229	35	-
2000.....	573	824	1,135	1,581	2,333	2,776	2,494	1,680	886	227	32	-
1999.....	559	835	1,151	1,656	2,290	2,650	2,343	1,763	809	250	37	1

Hispanic, both sexes

2003.....	93	140	160	199	230	221	165	94	49	10	1	-
2002.....	89	127	146	176	234	202	131	97	35	10	1	-
2001.....	80	122	141	184	235	196	144	85	39	10	-	-
2000.....	75	92	142	176	221	185	128	74	44	7	-	-
1999.....	74	95	123	155	183	172	100	73	33	14	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

Hispanic, male

2003.....	56	88	88	102	115	105	70	32	12	2	-	-
2002.....	57	71	76	97	117	85	51	27	7	2	-	-
2001.....	45	68	73	102	110	94	60	22	7	4	-	-
2000.....	43	58	75	106	108	78	50	21	15	4	-	-
1999.....	41	58	59	85	97	68	40	28	7	5	-	-

Hispanic, female

2003.....	37	52	72	97	115	116	95	62	37	8	1	-
2002.....	32	56	70	79	117	117	80	70	28	8	1	-
2001.....	35	54	68	82	125	102	84	63	32	6	-	-
2000.....	32	34	67	70	113	107	78	53	29	3	-	-
1999.....	33	37	64	70	86	104	60	45	26	9	-	-

Non-Hispanic 1/, both sexes

2003.....	1,553	2,182	2,822	3,348	4,317	4,808	4,494	2,796	1,215	282	31	-
2002.....	1,451	2,187	2,634	3,380	4,383	4,996	4,204	2,721	1,180	253	35	1
2001.....	1,493	1,987	2,611	3,331	4,388	4,779	4,184	2,771	1,174	282	42	1
2000.....	1,450	1,934	2,531	3,365	4,327	5,029	4,095	2,490	1,149	278	34	1
1999.....	1,387	1,942	2,538	3,467	4,461	4,747	3,891	2,595	1,086	294	40	1

Non-Hispanic 1/, male

2003.....	935	1,300	1,621	1,821	2,207	2,247	1,969	957	315	52	8	-
2002.....	912	1,317	1,539	1,825	2,226	2,334	1,792	954	343	58	7	1
2001.....	918	1,200	1,528	1,813	2,176	2,262	1,769	959	302	59	7	1
2000.....	909	1,145	1,464	1,857	2,113	2,367	1,687	869	293	55	2	1
1999.....	863	1,145	1,453	1,887	2,263	2,206	1,616	880	304	54	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

Non-Hispanic 1/, female

2003.....	618	882	1,201	1,527	2,110	2,561	2,525	1,839	900	230	23	-
2002.....	539	870	1,095	1,555	2,157	2,662	2,412	1,767	837	195	28	-
2001.....	575	787	1,083	1,518	2,212	2,517	2,415	1,812	872	223	35	-
2000.....	541	789	1,067	1,508	2,214	2,662	2,408	1,621	856	223	32	-
1999.....	524	797	1,085	1,580	2,198	2,541	2,275	1,715	782	240	37	1

Non-Hispanic White, both sexes

2003.....	1,194	1,784	2,313	2,814	3,689	4,226	3,980	2,488	1,104	249	24	-
2002.....	1,133	1,795	2,176	2,798	3,735	4,369	3,765	2,450	1,067	219	29	-
2001.....	1,210	1,632	2,215	2,802	3,762	4,199	3,732	2,493	1,034	250	33	1
2000.....	1,172	1,584	2,106	2,824	3,721	4,416	3,650	2,218	1,013	242	27	1
1999.....	1,108	1,564	2,111	2,926	3,858	4,160	3,465	2,322	971	267	30	1

Non-Hispanic White, male

2003.....	728	1,080	1,354	1,576	1,914	2,007	1,770	863	289	45	8	-
2002.....	728	1,091	1,301	1,549	1,931	2,088	1,627	869	312	50	5	-
2001.....	755	1,014	1,310	1,544	1,897	2,026	1,598	883	273	55	5	1
2000.....	742	946	1,237	1,603	1,841	2,120	1,518	790	256	45	2	1
1999.....	684	920	1,224	1,613	1,992	1,958	1,454	792	273	51	3	-

Non-Hispanic White, female

2003.....	466	704	959	1,238	1,775	2,219	2,210	1,625	815	204	16	-
2002.....	405	704	875	1,249	1,804	2,281	2,138	1,581	755	169	24	-
2001.....	455	618	905	1,258	1,865	2,173	2,134	1,610	761	195	28	-
2000.....	430	638	869	1,221	1,880	2,296	2,132	1,428	757	197	25	-
1999.....	424	644	887	1,313	1,866	2,202	2,011	1,530	698	216	27	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

Non-Hispanic Black, both sexes

2003.....	310	323	420	447	511	484	419	249	97	29	6	-
2002.....	257	332	394	471	545	509	349	225	95	32	6	1
2001.....	243	291	347	442	519	481	384	231	122	29	9	-
2000.....	245	302	362	452	518	524	366	227	117	32	5	-
1999.....	247	327	342	459	513	486	360	234	99	25	10	-

Non-Hispanic Black, male

2003.....	183	185	218	203	245	192	151	73	22	5	-	-
2002.....	157	193	206	225	250	196	136	62	22	8	2	1
2001.....	140	153	186	231	229	200	135	59	22	4	2	-
2000.....	150	174	193	213	234	213	137	60	26	8	-	-
1999.....	158	200	186	230	227	210	134	72	22	3	-	-

Non-Hispanic Black, female

2003.....	127	138	202	244	266	292	268	176	75	24	6	-
2002.....	100	139	188	246	295	313	213	163	73	24	4	-
2001.....	103	138	161	211	290	281	249	172	100	25	7	-
2000.....	95	128	169	239	284	311	229	167	91	24	5	-
1999.....	89	127	156	229	286	276	226	162	77	22	10	-

Origin not stated 2/, both sexes

2003.....	3	5	11	3	7	9	11	7	-	1	-	-
2002.....	4	7	4	8	6	12	10	6	2	-	-	-
2001.....	4	2	5	10	10	12	7	5	1	1	-	-
2000.....	3	4	5	5	9	9	13	7	3	1	-	-
1999.....	5	2	7	10	17	12	11	5	1	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of pancreas (C25)

Origin not stated 2/, male

2003.....	1	3	6	1	5	6	5	3	-	1	-	-
2002.....	2	6	3	2	3	5	3	4	1	-	-	-
2001.....	1	1	4	6	8	4	2	3	1	1	-	-
2000.....	3	3	4	2	3	2	5	1	2	-	-	-
1999.....	3	1	5	4	11	7	3	2	-	-	-	-

Origin not stated 2/, female

2003.....	2	2	5	2	2	3	6	4	-	-	-	-
2002.....	2	1	1	6	3	7	7	2	1	-	-	-
2001.....	3	1	1	4	2	8	5	2	-	-	-	-
2000.....	-	1	1	3	6	7	8	6	1	1	-	-
1999.....	2	1	2	6	6	5	8	3	1	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

All origins, both sexes

2003.....	263	404	552	568	568	490	401	225	70	13	-	1
2002.....	269	412	492	508	576	588	381	172	76	5	-	1
2001.....	270	414	499	531	639	553	357	208	74	13	1	-
2000.....	270	384	489	574	645	595	401	191	73	11	1	-
1999.....	254	410	494	584	655	529	363	200	72	16	3	-

All origins, male

2003.....	225	324	453	464	435	391	317	164	47	10	-	-
2002.....	219	339	399	395	437	455	300	126	56	1	-	1
2001.....	223	339	398	417	502	442	268	171	58	8	1	-
2000.....	218	319	407	441	504	451	314	155	51	8	-	-
1999.....	216	339	396	458	490	397	284	154	46	10	3	-

All origins, female

2003.....	38	80	99	104	133	99	84	61	23	3	-	1
2002.....	50	73	93	113	139	133	81	46	20	4	-	-
2001.....	47	75	101	114	137	111	89	37	16	5	-	-
2000.....	52	65	82	133	141	144	87	36	22	3	1	-
1999.....	38	71	98	126	165	132	79	46	26	6	-	-

Hispanic, both sexes

2003.....	12	20	24	25	19	20	18	11	3	1	-	-
2002.....	12	21	22	21	27	24	9	7	6	-	-	-
2001.....	6	18	20	35	25	20	9	9	4	-	-	-
2000.....	12	13	16	27	18	15	20	8	3	-	-	-
1999.....	10	17	12	30	25	24	16	11	4	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

Hispanic, male

2003.....	11	20	23	23	18	17	14	10	3	1	-	-
2002.....	10	21	19	19	23	19	8	2	5	-	-	-
2001.....	5	16	18	29	24	16	8	5	4	-	-	-
2000.....	11	13	15	23	14	12	18	7	3	-	-	-
1999.....	9	16	11	26	22	20	11	10	4	1	-	-

Hispanic, female

2003.....	1	-	1	2	1	3	4	1	-	-	-	-
2002.....	2	-	3	2	4	5	1	5	1	-	-	-
2001.....	1	2	2	6	1	4	1	4	-	-	-	-
2000.....	1	-	1	4	4	3	2	1	-	-	-	-
1999.....	1	1	1	4	3	4	5	1	-	1	-	-

Non-Hispanic 1/, both sexes

2003.....	249	383	525	540	546	468	379	214	67	12	-	1
2002.....	256	388	469	486	547	563	371	165	70	5	-	1
2001.....	263	390	475	494	612	530	347	199	70	13	1	-
2000.....	256	370	470	543	623	579	380	182	70	11	1	-
1999.....	242	393	478	552	628	503	344	189	67	14	3	-

Non-Hispanic 1/, male

2003.....	212	303	427	438	415	372	299	154	44	9	-	-
2002.....	208	315	380	376	412	435	291	124	51	1	-	1
2001.....	217	318	376	386	476	423	259	166	54	8	1	-
2000.....	206	305	389	415	486	438	295	148	48	8	-	-
1999.....	206	323	383	430	467	375	271	144	42	9	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

Non-Hispanic 1/, female

2003.....	37	80	98	102	131	96	80	60	23	3	-	1
2002.....	48	73	89	110	135	128	80	41	19	4	-	-
2001.....	46	72	99	108	136	107	88	33	16	5	-	-
2000.....	50	65	81	128	137	141	85	34	22	3	1	-
1999.....	36	70	95	122	161	128	73	45	25	5	-	-

Non-Hispanic White, both sexes

2003.....	169	291	406	433	445	398	324	188	54	8	-	1
2002.....	179	293	360	394	456	469	324	151	65	4	-	-
2001.....	184	287	370	396	497	442	305	174	64	11	1	-
2000.....	179	266	346	436	500	485	337	162	67	10	1	-
1999.....	171	291	369	444	534	443	308	165	54	13	3	-

Non-Hispanic White, male

2003.....	142	226	333	341	335	312	252	134	36	6	-	-
2002.....	148	234	284	303	342	361	258	113	47	-	-	-
2001.....	150	233	296	309	388	347	223	147	51	7	1	-
2000.....	140	215	285	331	390	370	262	133	45	7	-	-
1999.....	148	236	298	345	394	330	238	123	32	8	3	-

Non-Hispanic White, female

2003.....	27	65	73	92	110	86	72	54	18	2	-	1
2002.....	31	59	76	91	114	108	66	38	18	4	-	-
2001.....	34	54	74	87	109	95	82	27	13	4	-	-
2000.....	39	51	61	105	110	115	75	29	22	3	1	-
1999.....	23	55	71	99	140	113	70	42	22	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

Non-Hispanic Black, both sexes

2003.....	76	91	115	99	94	65	49	24	11	3	-	-
2002.....	75	91	107	86	88	89	42	12	3	1	-	1
2001.....	76	98	97	87	101	77	34	20	6	1	-	-
2000.....	74	101	117	95	114	88	41	19	2	1	-	-
1999.....	69	98	105	102	86	57	32	15	10	1	-	-

Non-Hispanic Black, male

2003.....	66	76	91	90	76	56	42	18	6	2	-	-
2002.....	58	77	95	68	67	72	29	10	2	1	-	1
2001.....	64	80	76	68	76	66	28	15	3	-	-	-
2000.....	64	87	99	75	89	62	32	14	2	1	-	-
1999.....	56	85	83	80	67	42	30	13	7	1	-	-

Non-Hispanic Black, female

2003.....	10	15	24	9	18	9	7	6	5	1	-	-
2002.....	17	14	12	18	21	17	13	2	1	-	-	-
2001.....	12	18	21	19	25	11	6	5	3	1	-	-
2000.....	10	14	18	20	25	26	9	5	-	-	-	-
1999.....	13	13	22	22	19	15	2	2	3	-	-	-

Origin not stated 2/, both sexes

2003.....	2	1	3	3	3	2	4	-	-	-	-	-
2002.....	1	3	1	1	2	1	1	-	-	-	-	-
2001.....	1	6	4	2	2	3	1	-	-	-	-	-
2000.....	2	1	3	4	4	1	1	1	-	-	-	-
1999.....	2	-	4	2	2	2	3	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of larynx (C32)

Origin not stated 2/, male

2003.....	2	1	3	3	2	2	4	-	-	-	-	-
2002.....	1	3	-	-	2	1	1	-	-	-	-	-
2001.....	1	5	4	2	2	3	1	-	-	-	-	-
2000.....	1	1	3	3	4	1	1	-	-	-	-	-
1999.....	1	-	2	2	1	2	2	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	1	1	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	1	-	-	-	1	-	-	-	-
1999.....	1	-	2	-	1	-	1	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins, both sexes

2003.....	7,757	12,920	18,036	22,414	26,972	28,031	20,588	10,076	3,319	617	76	2
2002.....	7,809	12,930	17,754	22,358	27,898	27,969	20,113	9,781	3,214	576	81	-
2001.....	7,786	12,595	17,182	22,850	28,375	27,723	18,977	9,623	3,064	587	57	4
2000.....	7,695	12,453	17,268	23,438	28,820	27,305	18,531	9,302	2,902	542	60	5
1999.....	7,518	12,165	17,179	23,239	28,609	26,556	17,705	8,945	2,766	549	79	2

All origins, male

2003.....	4,654	7,758	10,589	13,218	15,717	15,934	11,130	5,099	1,485	255	26	2
2002.....	4,723	7,769	10,646	13,209	16,334	15,833	10,988	5,022	1,439	216	25	-
2001.....	4,703	7,523	10,350	13,798	16,901	15,968	10,438	5,026	1,370	237	23	4
2000.....	4,583	7,546	10,391	14,189	17,193	15,726	10,240	4,840	1,335	204	16	5
1999.....	4,505	7,465	10,548	14,186	17,101	15,461	9,936	4,739	1,314	234	28	2

All origins, female

2003.....	3,103	5,162	7,447	9,196	11,255	12,097	9,458	4,977	1,834	362	50	-
2002.....	3,086	5,161	7,108	9,149	11,564	12,136	9,125	4,759	1,775	360	56	-
2001.....	3,083	5,072	6,832	9,052	11,474	11,755	8,539	4,597	1,694	350	34	-
2000.....	3,112	4,907	6,877	9,249	11,627	11,579	8,291	4,462	1,567	338	44	-
1999.....	3,013	4,700	6,631	9,053	11,508	11,095	7,769	4,206	1,452	315	51	-

Hispanic, both sexes

2003.....	247	331	464	605	688	703	474	246	91	24	4	-
2002.....	238	364	444	574	729	681	420	224	97	16	4	-
2001.....	237	325	433	564	666	626	385	202	90	27	2	-
2000.....	218	303	369	545	646	595	411	204	91	21	6	-
1999.....	197	303	421	556	614	553	348	215	92	15	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Hispanic, male

2003.....	163	233	316	423	454	448	275	123	49	14	3	-
2002.....	158	256	311	393	453	463	266	121	54	6	3	-
2001.....	156	219	298	369	451	413	244	119	43	13	1	-
2000.....	148	210	261	351	442	412	254	117	55	10	1	-
1999.....	134	219	304	394	404	377	229	107	51	9	1	-

Hispanic, female

2003.....	84	98	148	182	234	255	199	123	42	10	1	-
2002.....	80	108	133	181	276	218	154	103	43	10	1	-
2001.....	81	106	135	195	215	213	141	83	47	14	1	-
2000.....	70	93	108	194	204	183	157	87	36	11	5	-
1999.....	63	84	117	162	210	176	119	108	41	6	1	-

Non-Hispanic 1/, both sexes

2003.....	7,490	12,556	17,545	21,753	26,224	27,272	20,080	9,813	3,221	592	72	2
2002.....	7,548	12,541	17,261	21,725	27,111	27,217	19,645	9,534	3,114	559	77	-
2001.....	7,523	12,239	16,697	22,236	27,651	27,041	18,558	9,396	2,968	560	54	4
2000.....	7,444	12,110	16,855	22,831	28,109	26,655	18,087	9,076	2,801	518	54	5
1999.....	7,292	11,829	16,720	22,620	27,916	25,928	17,315	8,710	2,667	534	77	2

Non-Hispanic 1/, male

2003.....	4,476	7,505	10,257	12,752	15,221	15,451	10,835	4,969	1,434	241	23	2
2002.....	4,550	7,498	10,308	12,776	15,852	15,326	10,700	4,889	1,383	209	22	-
2001.....	4,529	7,278	10,013	13,402	16,416	15,518	10,177	4,892	1,325	224	21	4
2000.....	4,413	7,307	10,102	13,802	16,709	15,279	9,966	4,712	1,277	191	15	5
1999.....	4,350	7,224	10,214	13,750	16,647	15,037	9,687	4,620	1,260	225	27	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic 1/, female

2003.....	3,014	5,051	7,288	9,001	11,003	11,821	9,245	4,844	1,787	351	49	-
2002.....	2,998	5,043	6,953	8,949	11,259	11,891	8,945	4,645	1,731	350	55	-
2001.....	2,994	4,961	6,684	8,834	11,235	11,523	8,381	4,504	1,643	336	33	-
2000.....	3,031	4,803	6,753	9,029	11,400	11,376	8,121	4,364	1,524	327	39	-
1999.....	2,942	4,605	6,506	8,870	11,269	10,891	7,628	4,090	1,407	309	50	-

Non-Hispanic White, both sexes

2003.....	5,964	10,520	15,093	18,919	23,285	24,520	18,232	8,978	2,930	525	62	2
2002.....	6,041	10,506	14,712	18,892	24,212	24,544	17,962	8,732	2,840	502	60	-
2001.....	6,100	10,333	14,380	19,332	24,640	24,359	16,864	8,593	2,707	487	42	2
2000.....	6,042	10,244	14,533	19,953	24,999	24,044	16,622	8,287	2,552	455	43	3
1999.....	5,922	10,004	14,325	19,694	24,883	23,335	15,818	7,934	2,427	483	63	1

Non-Hispanic White, male

2003.....	3,515	6,177	8,684	10,988	13,452	13,795	9,778	4,532	1,296	211	22	2
2002.....	3,580	6,189	8,629	11,013	14,089	13,707	9,701	4,458	1,253	193	20	-
2001.....	3,594	6,032	8,520	11,536	14,520	13,853	9,169	4,432	1,205	202	18	2
2000.....	3,486	6,053	8,629	11,945	14,759	13,629	9,089	4,275	1,149	164	10	3
1999.....	3,447	6,023	8,631	11,872	14,724	13,441	8,778	4,188	1,143	206	19	1

Non-Hispanic White, female

2003.....	2,449	4,343	6,409	7,931	9,833	10,725	8,454	4,446	1,634	314	40	-
2002.....	2,461	4,317	6,083	7,879	10,123	10,837	8,261	4,274	1,587	309	40	-
2001.....	2,506	4,301	5,860	7,796	10,120	10,506	7,695	4,161	1,502	285	24	-
2000.....	2,556	4,191	5,904	8,008	10,240	10,415	7,533	4,012	1,403	291	33	-
1999.....	2,475	3,981	5,694	7,822	10,159	9,894	7,040	3,746	1,284	277	44	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Non-Hispanic Black, both sexes

2003.....	1,328	1,804	2,151	2,428	2,464	2,259	1,480	656	232	56	9	-
2002.....	1,343	1,784	2,210	2,483	2,447	2,254	1,387	630	220	44	15	-
2001.....	1,260	1,674	2,011	2,509	2,567	2,264	1,359	635	195	61	11	2
2000.....	1,280	1,638	2,050	2,512	2,666	2,206	1,208	647	206	53	9	1
1999.....	1,234	1,628	2,115	2,532	2,579	2,237	1,246	660	196	38	13	-

Non-Hispanic Black, male

2003.....	850	1,189	1,381	1,516	1,493	1,343	849	341	113	23	1	-
2002.....	874	1,145	1,461	1,575	1,508	1,360	803	334	102	10	2	-
2001.....	841	1,099	1,301	1,635	1,631	1,410	816	373	86	15	3	2
2000.....	854	1,109	1,301	1,646	1,663	1,379	726	351	104	23	5	1
1999.....	826	1,088	1,412	1,637	1,630	1,371	756	376	92	13	7	-

Non-Hispanic Black, female

2003.....	478	615	770	912	971	916	631	315	119	33	8	-
2002.....	469	639	749	908	939	894	584	296	118	34	13	-
2001.....	419	575	710	874	936	854	543	262	109	46	8	-
2000.....	426	529	749	866	1,003	827	482	296	102	30	4	-
1999.....	408	540	703	895	949	866	490	284	104	25	6	-

Origin not stated 2/, both sexes

2003.....	20	33	27	56	60	56	34	17	7	1	-	-
2002.....	23	25	49	59	58	71	48	23	3	1	-	-
2001.....	26	31	52	50	58	56	34	25	6	-	1	-
2000.....	33	40	44	62	65	55	33	22	10	3	-	-
1999.....	29	33	38	63	79	75	42	20	7	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Origin not stated 2/, male

2003.....	15	20	16	43	42	35	20	7	2	-	-	-
2002.....	15	15	27	40	29	44	22	12	2	1	-	-
2001.....	18	26	39	27	34	37	17	15	2	-	1	-
2000.....	22	29	28	36	42	35	20	11	3	3	-	-
1999.....	21	22	30	42	50	47	20	12	3	-	-	-

Origin not stated 2/, female

2003.....	5	13	11	13	18	21	14	10	5	1	-	-
2002.....	8	10	22	19	29	27	26	11	1	-	-	-
2001.....	8	5	13	23	24	19	17	10	4	-	-	-
2000.....	11	11	16	26	23	20	13	11	7	-	-	-
1999.....	8	11	8	21	29	28	22	8	4	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

All origins, both sexes

2003.....	638	778	769	710	883	1,033	842	500	243	78	10	-
2002.....	640	659	692	748	868	954	794	512	233	58	17	1
2001.....	672	653	680	746	943	919	759	489	213	72	11	-
2000.....	665	627	676	753	884	887	737	495	252	52	4	-
1999.....	602	615	648	791	861	945	694	435	181	53	12	-

All origins, male

2003.....	425	528	514	471	579	692	511	291	110	32	3	-
2002.....	403	445	470	500	575	609	469	264	115	21	6	-
2001.....	425	430	463	524	632	600	463	256	99	31	2	-
2000.....	448	420	446	495	572	559	454	277	100	13	-	-
1999.....	380	425	426	526	570	596	410	223	90	11	4	-

All origins, female

2003.....	213	250	255	239	304	341	331	209	133	46	7	-
2002.....	237	214	222	248	293	345	325	248	118	37	11	1
2001.....	247	223	217	222	311	319	296	233	114	41	9	-
2000.....	217	207	230	258	312	328	283	218	152	39	4	-
1999.....	222	190	222	265	291	349	284	212	91	42	8	-

Hispanic, both sexes

2003.....	20	16	13	16	21	11	14	5	4	2	-	-
2002.....	14	10	14	9	10	16	13	7	7	3	1	-
2001.....	12	14	11	13	15	15	13	4	2	2	1	-
2000.....	13	17	10	9	17	11	8	7	2	2	-	-
1999.....	13	17	13	15	12	15	5	4	5	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

Hispanic, male

2003.....	15	9	7	10	11	5	10	2	4	1	-	-
2002.....	5	8	8	4	7	7	6	3	1	-	-	-
2001.....	3	9	6	10	9	9	4	-	-	1	-	-
2000.....	8	13	6	3	11	6	7	4	1	-	-	-
1999.....	9	11	8	10	7	5	4	1	2	-	-	-

Hispanic, female

2003.....	5	7	6	6	10	6	4	3	-	1	-	-
2002.....	9	2	6	5	3	9	7	4	6	3	1	-
2001.....	9	5	5	3	6	6	9	4	2	1	1	-
2000.....	5	4	4	6	6	5	1	3	1	2	-	-
1999.....	4	6	5	5	5	10	1	3	3	2	-	-

Non-Hispanic 1/, both sexes

2003.....	616	758	755	694	861	1,021	825	492	239	76	10	-
2002.....	626	648	675	739	857	935	779	504	226	55	16	-
2001.....	659	638	669	733	927	901	745	484	211	70	10	-
2000.....	648	610	666	742	866	875	727	487	249	50	4	-
1999.....	587	597	634	772	848	926	686	431	174	51	12	-

Non-Hispanic 1/, male

2003.....	408	518	507	461	567	687	498	287	106	31	3	-
2002.....	398	437	459	496	567	601	461	261	114	21	6	-
2001.....	421	420	457	514	622	588	459	256	99	30	2	-
2000.....	436	407	440	490	561	552	446	273	98	13	-	-
1999.....	370	414	417	514	562	589	405	222	86	11	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

Non-Hispanic 1/, female

2003.....	208	240	248	233	294	334	327	205	133	45	7	-
2002.....	228	211	216	243	290	334	318	243	112	34	10	-
2001.....	238	218	212	219	305	313	286	228	112	40	8	-
2000.....	212	203	226	252	305	323	281	214	151	37	4	-
1999.....	217	183	217	258	286	337	281	209	88	40	8	-

Non-Hispanic White, both sexes

2003.....	601	740	739	678	848	993	812	482	232	75	9	-
2002.....	614	631	660	722	832	919	768	493	220	52	15	-
2001.....	650	632	654	711	905	885	729	476	207	69	9	-
2000.....	637	594	653	726	844	859	707	475	240	48	4	-
1999.....	583	591	622	760	833	905	672	420	169	50	12	-

Non-Hispanic White, male

2003.....	400	509	498	451	565	674	491	284	103	31	3	-
2002.....	390	425	451	486	553	592	458	256	112	21	6	-
2001.....	417	417	450	503	613	580	454	253	99	30	2	-
2000.....	429	398	435	480	552	545	436	272	94	13	-	-
1999.....	367	410	409	508	556	577	399	218	84	10	4	-

Non-Hispanic White, female

2003.....	201	231	241	227	283	319	321	198	129	44	6	-
2002.....	224	206	209	236	279	327	310	237	108	31	9	-
2001.....	233	215	204	208	292	305	275	223	108	39	7	-
2000.....	208	196	218	246	292	314	271	203	146	35	4	-
1999.....	216	181	213	252	277	328	273	202	85	40	8	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

Non-Hispanic Black, both sexes

2003.....	8	12	11	16	10	20	8	7	5	-	1	-
2002.....	9	13	8	11	14	10	8	10	5	-	-	-
2001.....	6	5	8	18	18	9	15	5	4	1	1	-
2000.....	7	11	8	11	19	14	18	12	7	2	-	-
1999.....	3	5	9	9	14	14	11	8	5	1	-	-

Non-Hispanic Black, male

2003.....	4	6	6	10	2	7	3	-	2	-	-	-
2002.....	5	10	3	5	7	6	2	4	2	-	-	-
2001.....	1	3	1	9	7	2	4	1	-	-	-	-
2000.....	5	6	4	7	7	6	9	1	2	-	-	-
1999.....	2	3	7	3	6	8	4	3	2	1	-	-

Non-Hispanic Black, female

2003.....	4	6	5	6	8	13	5	7	3	-	1	-
2002.....	4	3	5	6	7	4	6	6	3	-	-	-
2001.....	5	2	7	9	11	7	11	4	4	1	1	-
2000.....	2	5	4	4	12	8	9	11	5	2	-	-
1999.....	1	2	2	6	8	6	7	5	3	-	-	-

Origin not stated 2/, both sexes

2003.....	2	4	1	-	1	1	3	3	-	-	-	-
2002.....	-	1	3	-	1	3	2	1	-	-	-	1
2001.....	1	1	-	-	1	3	1	1	-	-	-	-
2000.....	4	-	-	2	1	1	2	1	1	-	-	-
1999.....	2	1	1	4	1	4	3	-	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant melanoma of skin (C43)

Origin not stated 2/, male

2003.....	2	1	-	-	1	-	3	2	-	-	-	-
2002.....	-	-	3	-	1	1	2	-	-	-	-	-
2001.....	1	1	-	-	1	3	-	-	-	-	-	-
2000.....	4	-	-	2	-	1	1	-	1	-	-	-
1999.....	1	-	1	2	1	2	1	-	2	-	-	-

Origin not stated 2/, female

2003.....	-	3	1	-	-	1	-	1	-	-	-	-
2002.....	-	1	-	-	-	2	-	1	-	-	-	1
2001.....	-	-	-	-	-	-	1	1	-	-	-	-
2000.....	-	-	-	-	1	-	1	1	-	-	-	-
1999.....	1	1	-	2	-	2	2	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

All origins, both sexes

2003.....	3,631	4,206	4,061	3,887	4,451	4,916	4,728	3,530	1,965	630	120	2
2002.....	3,707	3,924	3,914	3,954	4,551	4,978	4,770	3,528	1,869	658	126	-
2001.....	3,803	3,843	3,789	3,972	4,743	5,130	4,533	3,330	1,848	644	89	-
2000.....	3,655	3,790	3,783	4,135	4,889	5,256	4,494	3,596	1,880	658	115	-
1999.....	3,555	3,682	3,717	4,147	4,962	5,140	4,511	3,428	1,837	611	124	-

All origins, male

2003.....	34	30	41	42	47	46	58	35	17	4	-	1
2002.....	28	29	38	35	49	57	55	31	13	3	-	-
2001.....	33	32	40	54	66	69	46	31	9	6	-	-
2000.....	28	44	41	47	68	65	44	36	13	1	1	-
1999.....	30	36	34	47	56	56	42	33	13	7	1	-

All origins, female

2003.....	3,597	4,176	4,020	3,845	4,404	4,870	4,670	3,495	1,948	626	120	1
2002.....	3,679	3,895	3,876	3,919	4,502	4,921	4,715	3,497	1,856	655	126	-
2001.....	3,770	3,811	3,749	3,918	4,677	5,061	4,487	3,299	1,839	638	89	-
2000.....	3,627	3,746	3,742	4,088	4,821	5,191	4,450	3,560	1,867	657	114	-
1999.....	3,525	3,646	3,683	4,100	4,906	5,084	4,469	3,395	1,824	604	123	-

Hispanic, both sexes

2003.....	220	241	187	188	143	169	100	95	41	12	2	-
2002.....	216	181	167	173	150	138	124	79	35	11	3	-
2001.....	224	216	191	158	160	137	82	55	36	17	2	-
2000.....	197	204	155	183	154	135	96	64	34	7	3	-
1999.....	202	164	157	146	144	115	94	61	45	14	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

Hispanic, male

2003.....	1	2	-	-	2	-	-	1	2	-	-	-
2002.....	4	-	-	1	2	-	2	-	1	-	-	-
2001.....	1	-	1	4	1	3	2	-	-	-	-	-
2000.....	2	3	1	2	4	2	2	-	1	-	-	-
1999.....	2	3	-	2	3	1	1	-	-	-	-	-

Hispanic, female

2003.....	219	239	187	188	141	169	100	94	39	12	2	-
2002.....	212	181	167	172	148	138	122	79	34	11	3	-
2001.....	223	216	190	154	159	134	80	55	36	17	2	-
2000.....	195	201	154	181	150	133	94	64	33	7	3	-
1999.....	200	161	157	144	141	114	93	61	45	14	3	-

Non-Hispanic 1/, both sexes

2003.....	3,401	3,949	3,864	3,693	4,294	4,733	4,612	3,428	1,924	618	118	2
2002.....	3,479	3,729	3,741	3,770	4,391	4,830	4,634	3,442	1,829	645	122	-
2001.....	3,573	3,621	3,587	3,805	4,574	4,976	4,442	3,267	1,808	627	87	-
2000.....	3,448	3,580	3,612	3,941	4,729	5,105	4,389	3,521	1,842	648	112	-
1999.....	3,347	3,507	3,549	3,991	4,812	5,011	4,408	3,355	1,790	597	121	-

Non-Hispanic 1/, male

2003.....	33	28	41	42	45	46	58	34	15	4	-	1
2002.....	24	28	38	34	47	57	53	31	12	3	-	-
2001.....	32	32	38	49	65	66	44	31	9	6	-	-
2000.....	26	41	40	45	64	63	42	36	12	1	1	-
1999.....	28	33	34	45	52	55	41	33	13	7	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

Non-Hispanic 1/, female

2003.....	3,368	3,921	3,823	3,651	4,249	4,687	4,554	3,394	1,909	614	118	1
2002.....	3,455	3,701	3,703	3,736	4,344	4,773	4,581	3,411	1,817	642	122	-
2001.....	3,541	3,589	3,549	3,756	4,509	4,910	4,398	3,236	1,799	621	87	-
2000.....	3,422	3,539	3,572	3,896	4,665	5,042	4,347	3,485	1,830	647	111	-
1999.....	3,319	3,474	3,515	3,946	4,760	4,956	4,367	3,322	1,777	590	120	-

Non-Hispanic White, both sexes

2003.....	2,600	3,161	3,193	3,097	3,727	4,198	4,158	3,099	1,739	556	94	1
2002.....	2,683	3,021	3,124	3,188	3,813	4,320	4,136	3,135	1,665	578	105	-
2001.....	2,790	2,922	2,973	3,201	3,984	4,458	4,014	2,959	1,660	568	67	-
2000.....	2,733	2,895	3,016	3,394	4,187	4,586	4,005	3,194	1,658	576	99	-
1999.....	2,650	2,838	2,941	3,396	4,227	4,474	4,014	3,084	1,634	541	105	-

Non-Hispanic White, male

2003.....	29	22	35	36	38	42	53	30	14	4	-	1
2002.....	17	23	28	23	42	47	47	29	9	3	-	-
2001.....	23	27	32	41	54	56	40	26	9	4	-	-
2000.....	22	31	32	41	56	58	36	29	11	1	1	-
1999.....	20	26	25	38	41	48	36	30	11	5	1	-

Non-Hispanic White, female

2003.....	2,571	3,139	3,158	3,061	3,689	4,156	4,105	3,069	1,725	552	94	-
2002.....	2,666	2,998	3,096	3,165	3,771	4,273	4,089	3,106	1,656	575	105	-
2001.....	2,767	2,895	2,941	3,160	3,930	4,402	3,974	2,933	1,651	564	67	-
2000.....	2,711	2,864	2,984	3,353	4,131	4,528	3,969	3,165	1,647	575	98	-
1999.....	2,630	2,812	2,916	3,358	4,186	4,426	3,978	3,054	1,623	536	104	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

Non-Hispanic Black, both sexes

2003.....	670	678	560	513	488	466	411	298	169	59	23	-
2002.....	669	596	526	508	497	455	445	285	146	63	15	-
2001.....	657	598	527	532	521	473	389	274	141	59	19	-
2000.....	611	589	514	473	488	472	353	305	173	71	13	-
1999.....	605	577	518	522	515	490	362	251	140	54	16	-

Non-Hispanic Black, male

2003.....	4	6	6	6	7	3	4	4	1	-	-	-
2002.....	6	5	9	11	5	8	5	2	3	-	-	-
2001.....	9	4	6	7	10	9	4	4	-	2	-	-
2000.....	4	8	8	3	8	5	5	7	1	-	-	-
1999.....	8	7	9	6	10	6	5	3	2	2	-	-

Non-Hispanic Black, female

2003.....	666	672	554	507	481	463	407	294	168	59	23	-
2002.....	663	591	517	497	492	447	440	283	143	63	15	-
2001.....	648	594	521	525	511	464	385	270	141	57	19	-
2000.....	607	581	506	470	480	467	348	298	172	71	13	-
1999.....	597	570	509	516	505	484	357	248	138	52	16	-

Origin not stated 2/, both sexes

2003.....	10	16	10	6	14	14	16	7	-	-	-	-
2002.....	12	14	6	11	10	10	12	7	5	2	1	-
2001.....	6	6	11	9	9	17	9	8	4	-	-	-
2000.....	10	6	16	11	6	16	9	11	4	3	-	-
1999.....	6	11	11	10	6	14	9	12	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of breast (C50)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	10	16	10	6	14	14	16	7	-	-	-	-
2002.....	12	13	6	11	10	10	12	7	5	2	1	-
2001.....	6	6	10	8	9	17	9	8	4	-	-	-
2000.....	10	6	16	11	6	16	9	11	4	3	-	-
1999.....	6	11	11	10	5	14	9	12	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

All origins, both sexes

2003.....	450	445	348	308	256	274	278	166	78	22	4	-
2002.....	455	401	324	315	315	305	223	155	99	24	6	-
2001.....	478	371	352	279	306	298	285	193	90	30	2	-
2000.....	467	416	332	342	344	320	279	180	92	35	3	-
1999.....	452	393	376	350	346	345	223	199	98	26	6	1

All origins, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

All origins, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, both sexes

2003.....	55	42	23	21	22	22	20	8	4	1	-	-
2002.....	45	40	31	24	31	24	16	8	6	2	1	-
2001.....	42	41	34	16	23	21	18	6	8	1	1	-
2000.....	40	40	38	30	30	24	19	12	8	1	-	-
1999.....	44	34	30	27	24	20	11	8	4	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Non-Hispanic Black, both sexes

2003.....	82	80	72	67	58	66	45	29	19	2	-	-
2002.....	93	62	69	77	75	54	47	29	19	9	1	-
2001.....	87	65	66	72	57	55	56	39	18	4	-	-
2000.....	91	82	70	85	73	60	48	30	21	10	1	-
1999.....	72	80	87	77	96	68	33	35	21	9	2	-

Non-Hispanic Black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic Black, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Origin not stated 2/, both sexes

2003.....	1	-	1	1	1	-	-	-	-	-	-	-
2002.....	1	2	2	-	2	-	1	-	-	-	-	-
2001.....	2	1	1	-	-	-	-	-	-	-	-	-
2000.....	1	-	2	-	1	2	-	-	-	-	-	-
1999.....	2	1	2	1	-	2	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of cervix uteri (C53)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	1	-	1	1	1	-	-	-	-	-	-	-
2002.....	1	2	2	-	2	-	1	-	-	-	-	-
2001.....	2	1	1	-	-	-	-	-	-	-	-	-
2000.....	1	-	2	-	1	2	-	-	-	-	-	-
1999.....	2	1	2	1	-	2	-	-	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins, both sexes

2003.....	363	557	744	878	973	1,025	926	668	316	76	17	-
2002.....	376	531	668	809	1,026	1,051	906	694	314	109	11	-
2001.....	374	457	648	785	999	1,075	988	662	312	104	10	1
2000.....	317	478	644	800	981	1,039	919	617	333	84	15	1
1999.....	286	427	592	835	972	1,061	947	593	290	100	16	-

All origins, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

All origins, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, both sexes

2003.....	28	30	31	43	42	35	28	21	14	4	1	-
2002.....	21	38	26	54	52	40	26	26	7	4	-	-
2001.....	32	32	39	39	46	34	31	20	6	4	-	-
2000.....	24	31	31	35	40	35	19	20	7	2	1	-
1999.....	23	28	33	45	41	30	21	16	4	2	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Non-Hispanic Black, both sexes

2003.....	54	81	164	189	168	146	123	65	28	4	5	-
2002.....	61	98	123	187	186	160	108	81	31	17	2	-
2001.....	59	63	127	173	179	150	115	55	36	11	3	1
2000.....	44	70	131	177	194	156	124	56	36	10	4	1
1999.....	39	76	113	187	159	142	103	65	35	9	2	-

Non-Hispanic Black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic Black, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Origin not stated 2/, both sexes

2003.....	1	1	2	3	2	6	1	-	-	-	-	-
2002.....	2	5	6	2	2	4	3	3	1	-	-	-
2001.....	1	2	3	-	2	2	-	5	1	-	-	-
2000.....	2	1	2	-	4	-	3	2	-	-	-	-
1999.....	-	1	1	-	2	2	1	-	-	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	1	1	2	3	2	6	1	-	-	-	-	-
2002.....	2	5	6	2	2	4	3	3	1	-	-	-
2001.....	1	2	3	-	2	2	-	5	1	-	-	-
2000.....	2	1	2	-	4	-	3	2	-	-	-	-
1999.....	-	1	1	-	2	2	1	-	-	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of ovary (C56)

All origins, both sexes

2003.....	997	1,315	1,576	1,687	1,882	2,193	2,001	1,207	490	102	11	-
2002.....	1,029	1,249	1,499	1,662	1,939	2,249	2,010	1,174	447	115	14	-
2001.....	1,016	1,171	1,441	1,682	1,984	2,246	1,924	1,181	451	96	15	-
2000.....	966	1,123	1,321	1,669	1,932	2,295	1,803	1,176	453	98	9	-
1999.....	933	1,081	1,345	1,626	1,986	2,143	1,732	1,052	412	91	7	-

All origins, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

All origins, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Hispanic, both sexes

2003.....	53	70	64	74	81	83	58	35	16	4	-	-
2002.....	71	47	64	71	74	90	55	31	12	-	-	-
2001.....	52	57	74	89	75	58	53	36	12	1	-	-
2000.....	52	55	56	75	79	80	46	33	15	1	-	-
1999.....	51	41	59	51	75	59	41	24	9	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of ovary (C56)

Non-Hispanic Black, both sexes

2003.....	109	106	130	143	154	154	113	82	31	9	3	-
2002.....	86	113	124	139	148	153	124	78	27	9	3	-
2001.....	87	112	139	150	151	149	103	79	24	9	5	-
2000.....	88	86	119	149	146	147	97	65	31	12	1	-
1999.....	89	94	117	135	166	159	97	54	30	6	-	-

Non-Hispanic Black, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic Black, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Origin not stated 2/, both sexes

2003.....	2	6	2	4	4	-	3	-	1	-	-	-
2002.....	2	3	1	8	3	4	3	1	-	-	-	-
2001.....	-	2	4	4	3	6	1	3	1	1	-	-
2000.....	1	1	3	4	3	3	5	1	2	-	-	-
1999.....	1	5	5	5	7	11	8	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of ovary (C56)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	2	6	2	4	4	-	3	-	1	-	-	-
2002.....	2	3	1	8	3	4	3	1	-	-	-	-
2001.....	-	2	4	4	3	6	1	3	1	1	-	-
2000.....	1	1	3	4	3	3	5	1	2	-	-	-
1999.....	1	5	5	5	7	11	8	1	-	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of prostate (C61)

Hispanic, male

2003.....	18	38	73	127	185	230	247	170	92	28	3	-
2002.....	22	36	70	131	161	235	237	171	89	47	1	-
2001.....	22	40	68	142	190	278	209	155	78	37	2	-
2000.....	19	21	57	95	182	208	193	150	75	25	-	-
1999.....	20	40	52	135	160	209	177	142	82	23	4	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Non-Hispanic 1/, male

2003.....	299	630	1,329	2,158	3,547	5,428	6,359	5,180	2,527	624	79	-
2002.....	279	658	1,248	2,177	3,806	5,667	6,584	5,290	2,612	655	71	-
2001.....	291	603	1,187	2,231	3,966	5,820	6,629	5,335	2,486	651	83	-
2000.....	243	601	1,245	2,368	4,144	6,085	6,549	5,370	2,558	600	75	-
1999.....	254	595	1,253	2,573	4,431	6,308	6,717	5,263	2,383	619	87	1







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All origins, both sexes

2003.....	763	1,121	1,377	1,409	1,681	1,849	1,626	1,033	420	107	7	-
2002.....	808	1,075	1,265	1,533	1,706	1,829	1,526	967	429	90	7	-
2001.....	813	1,006	1,322	1,395	1,826	1,810	1,496	947	390	82	5	-
2000.....	789	1,026	1,169	1,432	1,714	1,819	1,463	945	397	83	7	-
1999.....	720	959	1,152	1,374	1,702	1,696	1,314	867	341	91	6	-

All origins, male

2003.....	535	798	989	916	1,059	1,114	922	486	175	40	2	-
2002.....	592	771	887	1,020	1,067	1,123	801	457	201	34	4	-
2001.....	586	722	912	928	1,165	1,052	814	500	164	23	-	-
2000.....	571	727	803	942	1,068	1,054	811	477	165	31	3	-
1999.....	506	671	803	894	1,064	1,016	701	421	140	29	-	-

All origins, female

2003.....	228	323	388	493	622	735	704	547	245	67	5	-
2002.....	216	304	378	513	639	706	725	510	228	56	3	-
2001.....	227	284	410	467	661	758	682	447	226	59	5	-
2000.....	218	299	366	490	646	765	652	468	232	52	4	-
1999.....	214	288	349	480	638	680	613	446	201	62	6	-

Hispanic, both sexes

2003.....	45	76	94	88	110	96	55	29	21	2	-	-
2002.....	60	75	85	78	74	88	58	30	14	3	-	-
2001.....	63	48	79	77	86	86	43	37	9	4	1	-
2000.....	46	60	75	65	67	70	55	28	8	3	-	-
1999.....	52	61	68	77	67	65	43	28	9	3	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Hispanic, male

2003.....	26	48	61	58	63	57	27	16	8	1	-	-
2002.....	45	54	52	50	47	64	31	16	7	1	-	-
2001.....	43	33	53	50	57	52	27	23	6	1	-	-
2000.....	30	46	46	40	45	47	31	16	3	1	-	-
1999.....	35	38	53	49	47	38	19	13	4	1	-	-

Hispanic, female

2003.....	19	28	33	30	47	39	28	13	13	1	-	-
2002.....	15	21	33	28	27	24	27	14	7	2	-	-
2001.....	20	15	26	27	29	34	16	14	3	3	1	-
2000.....	16	14	29	25	22	23	24	12	5	2	-	-
1999.....	17	23	15	28	20	27	24	15	5	2	1	-

Non-Hispanic 1/, both sexes

2003.....	716	1,043	1,282	1,319	1,565	1,751	1,569	1,003	396	105	7	-
2002.....	745	996	1,177	1,451	1,628	1,735	1,465	937	414	87	7	-
2001.....	748	958	1,240	1,314	1,737	1,722	1,450	910	379	76	4	-
2000.....	739	961	1,092	1,357	1,637	1,744	1,406	911	388	80	7	-
1999.....	667	897	1,082	1,295	1,632	1,624	1,268	838	330	88	5	-

Non-Hispanic 1/, male

2003.....	508	748	927	856	992	1,055	893	470	165	39	2	-
2002.....	545	713	833	967	1,017	1,057	768	441	193	33	4	-
2001.....	542	689	857	875	1,106	999	785	477	157	22	-	-
2000.....	539	678	755	895	1,016	1,003	779	457	162	30	3	-
1999.....	470	633	748	843	1,015	975	682	408	135	28	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic 1/, female

2003.....	208	295	355	463	573	696	676	533	231	66	5	-
2002.....	200	283	344	484	611	678	697	496	221	54	3	-
2001.....	206	269	383	439	631	723	665	433	222	54	4	-
2000.....	200	283	337	462	621	741	627	454	226	50	4	-
1999.....	197	264	334	452	617	649	586	430	195	60	5	-

Non-Hispanic White, both sexes

2003.....	614	890	1,114	1,164	1,406	1,589	1,442	927	356	96	5	-
2002.....	626	872	1,026	1,287	1,467	1,561	1,342	860	388	78	7	-
2001.....	623	833	1,071	1,178	1,563	1,554	1,326	836	345	71	4	-
2000.....	638	840	954	1,198	1,467	1,582	1,289	848	353	73	6	-
1999.....	564	782	958	1,149	1,464	1,470	1,166	773	293	74	4	-

Non-Hispanic White, male

2003.....	436	654	809	761	889	958	829	440	151	35	1	-
2002.....	464	624	731	865	920	967	701	408	187	30	4	-
2001.....	451	608	732	788	1,002	895	721	444	146	21	-	-
2000.....	465	590	668	782	915	914	714	425	149	28	2	-
1999.....	392	550	670	748	914	887	641	381	119	19	-	-

Non-Hispanic White, female

2003.....	178	236	305	403	517	631	613	487	205	61	4	-
2002.....	162	248	295	422	547	594	641	452	201	48	3	-
2001.....	172	225	339	390	561	659	605	392	199	50	4	-
2000.....	173	250	286	416	552	668	575	423	204	45	4	-
1999.....	172	232	288	401	550	583	525	392	174	55	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Non-Hispanic Black, both sexes

2003.....	88	129	132	126	120	129	103	70	36	7	2	-
2002.....	91	104	122	135	125	149	101	65	21	7	-	-
2001.....	100	104	136	108	141	137	104	63	28	5	-	-
2000.....	84	99	114	134	137	132	101	56	29	7	1	-
1999.....	87	99	103	122	141	129	84	55	31	11	1	-

Non-Hispanic Black, male

2003.....	63	81	88	77	77	76	53	26	13	2	1	-
2002.....	62	79	85	80	71	75	50	28	6	2	-	-
2001.....	72	67	101	67	86	83	53	30	9	1	-	-
2000.....	59	68	74	96	83	68	56	29	10	2	1	-
1999.....	65	72	65	77	82	72	32	21	13	8	-	-

Non-Hispanic Black, female

2003.....	25	48	44	49	43	53	50	44	23	5	1	-
2002.....	29	25	37	55	54	74	51	37	15	5	-	-
2001.....	28	37	35	41	55	54	51	33	19	4	-	-
2000.....	25	31	40	38	54	64	45	27	19	5	-	-
1999.....	22	27	38	45	59	57	52	34	18	3	1	-

Origin not stated 2/, both sexes

2003.....	2	2	1	2	6	2	2	1	3	-	-	-
2002.....	3	4	3	4	4	6	3	-	1	-	-	-
2001.....	2	-	3	4	3	2	3	-	2	2	-	-
2000.....	4	5	2	10	10	5	2	6	1	-	-	-
1999.....	1	1	2	2	3	7	3	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Origin not stated 2/, male

2003.....	1	2	1	2	4	2	2	-	2	-	-	-
2002.....	2	4	2	3	3	2	2	-	1	-	-	-
2001.....	1	-	2	3	2	1	2	-	1	-	-	-
2000.....	2	3	2	7	7	4	1	4	-	-	-	-
1999.....	1	-	2	2	2	3	-	-	1	-	-	-

Origin not stated 2/, female

2003.....	1	-	-	-	2	-	-	1	1	-	-	-
2002.....	1	-	1	1	1	4	1	-	-	-	-	-
2001.....	1	-	1	1	1	1	1	-	1	2	-	-
2000.....	2	2	-	3	3	1	1	2	1	-	-	-
1999.....	-	1	-	-	1	4	3	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

All origins, both sexes

2003.....	341	554	837	1,148	1,588	2,239	2,440	1,850	888	278	36	-
2002.....	321	572	738	1,151	1,756	2,366	2,341	1,864	908	255	32	1
2001.....	314	484	781	1,147	1,667	2,179	2,309	1,865	864	265	42	-
2000.....	280	486	737	1,119	1,736	2,155	2,252	1,790	875	256	31	-
1999.....	323	492	752	1,126	1,797	2,178	2,139	1,714	852	249	30	1

All origins, male

2003.....	252	419	630	865	1,133	1,603	1,651	1,157	508	131	11	-
2002.....	223	425	559	845	1,244	1,659	1,540	1,163	498	118	12	1
2001.....	233	359	597	859	1,229	1,554	1,562	1,129	482	123	17	-
2000.....	212	369	547	838	1,256	1,542	1,512	1,110	478	110	12	-
1999.....	243	374	559	849	1,283	1,575	1,433	1,039	438	116	15	1

All origins, female

2003.....	89	135	207	283	455	636	789	693	380	147	25	-
2002.....	98	147	179	306	512	707	801	701	410	137	20	-
2001.....	81	125	184	288	438	625	747	736	382	142	25	-
2000.....	68	117	190	281	480	613	740	680	397	146	19	-
1999.....	80	118	193	277	514	603	706	675	414	133	15	-

Hispanic, both sexes

2003.....	19	24	30	43	53	65	68	42	30	9	-	-
2002.....	13	27	30	40	51	53	57	55	39	4	1	-
2001.....	9	16	25	40	50	51	53	41	20	11	2	-
2000.....	9	25	25	21	50	46	43	39	21	8	3	-
1999.....	17	21	30	35	53	41	38	37	14	2	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

Hispanic, male

2003.....	10	17	23	37	34	40	47	23	24	3	-	-
2002.....	9	15	24	31	36	34	34	33	24	3	-	-
2001.....	4	15	22	27	32	31	36	21	15	8	2	-
2000.....	7	21	16	16	34	39	30	21	15	4	2	-
1999.....	12	18	25	30	39	28	24	26	9	1	-	-

Hispanic, female

2003.....	9	7	7	6	19	25	21	19	6	6	-	-
2002.....	4	12	6	9	15	19	23	22	15	1	1	-
2001.....	5	1	3	13	18	20	17	20	5	3	-	-
2000.....	2	4	9	5	16	7	13	18	6	4	1	-
1999.....	5	3	5	5	14	13	14	11	5	1	2	-

Non-Hispanic 1/, both sexes

2003.....	321	529	807	1,105	1,532	2,170	2,364	1,804	858	269	36	-
2002.....	306	543	705	1,108	1,701	2,312	2,279	1,803	867	251	31	-
2001.....	305	467	753	1,103	1,615	2,120	2,252	1,819	843	253	40	-
2000.....	271	458	711	1,097	1,683	2,103	2,201	1,747	852	247	28	-
1999.....	304	470	720	1,088	1,744	2,129	2,099	1,671	834	247	28	1

Non-Hispanic 1/, male

2003.....	241	401	607	828	1,097	1,559	1,598	1,132	484	128	11	-
2002.....	213	409	532	812	1,207	1,625	1,503	1,127	472	115	12	-
2001.....	229	344	573	829	1,196	1,518	1,523	1,104	466	115	15	-
2000.....	205	346	530	821	1,219	1,499	1,475	1,087	462	105	10	-
1999.....	229	355	533	816	1,244	1,542	1,408	1,010	427	115	15	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

Non-Hispanic 1/, female

2003.....	80	128	200	277	435	611	766	672	374	141	25	-
2002.....	93	134	173	296	494	687	776	676	395	136	19	-
2001.....	76	123	180	274	419	602	729	715	377	138	25	-
2000.....	66	112	181	276	464	604	726	660	390	142	18	-
1999.....	75	115	187	272	500	587	691	661	407	132	13	-

Non-Hispanic White, both sexes

2003.....	269	467	731	982	1,417	1,990	2,178	1,672	806	245	32	-
2002.....	265	474	638	989	1,542	2,126	2,106	1,663	790	225	28	-
2001.....	265	409	667	995	1,484	1,956	2,070	1,697	772	235	36	-
2000.....	245	412	644	998	1,535	1,926	2,031	1,624	809	233	24	-
1999.....	266	407	635	969	1,587	1,964	1,957	1,550	770	219	28	1

Non-Hispanic White, male

2003.....	211	360	561	743	1,034	1,453	1,499	1,074	461	119	9	-
2002.....	193	375	491	742	1,122	1,511	1,410	1,061	441	109	12	-
2001.....	201	308	522	768	1,116	1,428	1,428	1,044	440	113	13	-
2000.....	188	312	491	762	1,135	1,405	1,379	1,015	445	101	10	-
1999.....	198	321	479	746	1,161	1,445	1,326	957	402	106	15	1

Non-Hispanic White, female

2003.....	58	107	170	239	383	537	679	598	345	126	23	-
2002.....	72	99	147	247	420	615	696	602	349	116	16	-
2001.....	64	101	145	227	368	528	642	653	332	122	23	-
2000.....	57	100	153	236	400	521	652	609	364	132	14	-
1999.....	68	86	156	223	426	519	631	593	368	113	13	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

Non-Hispanic Black, both sexes

2003.....	47	50	59	109	95	155	160	114	42	22	3	-
2002.....	33	59	57	103	139	154	141	120	68	21	3	-
2001.....	35	48	77	93	104	140	151	97	61	17	3	-
2000.....	24	40	59	86	132	159	155	102	39	12	3	-
1999.....	37	59	71	105	139	140	122	109	54	25	-	-

Non-Hispanic Black, male

2003.....	27	32	34	74	49	89	83	50	17	7	1	-
2002.....	16	29	34	57	71	90	74	55	26	3	-	-
2001.....	26	27	45	50	61	72	72	44	20	2	2	-
2000.....	16	28	35	48	71	86	85	58	14	3	-	-
1999.....	30	32	44	59	69	81	68	48	18	7	-	-

Non-Hispanic Black, female

2003.....	20	18	25	35	46	66	77	64	25	15	2	-
2002.....	17	30	23	46	68	64	67	65	42	18	3	-
2001.....	9	21	32	43	43	68	79	53	41	15	1	-
2000.....	8	12	24	38	61	73	70	44	25	9	3	-
1999.....	7	27	27	46	70	59	54	61	36	18	-	-

Origin not stated 2/, both sexes

2003.....	1	1	-	-	3	4	8	4	-	-	-	-
2002.....	2	2	3	3	4	1	5	6	2	-	-	1
2001.....	-	1	3	4	2	8	4	5	1	1	-	-
2000.....	-	3	1	1	3	6	8	4	2	1	-	-
1999.....	2	1	2	3	-	8	2	6	4	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasm of bladder (C67)

Origin not stated 2/, male

2003.....	1	1	-	-	2	4	6	2	-	-	-	-
2002.....	1	1	3	2	1	-	3	3	2	-	-	1
2001.....	-	-	2	3	1	5	3	4	1	-	-	-
2000.....	-	2	1	1	3	4	7	2	1	1	-	-
1999.....	2	1	1	3	-	5	1	3	2	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	1	-	2	2	-	-	-	-
2002.....	1	1	-	1	3	1	2	3	-	-	-	-
2001.....	-	1	1	1	1	3	1	1	-	1	-	-
2000.....	-	1	-	-	-	2	1	2	1	-	-	-
1999.....	-	-	1	-	-	3	1	3	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins, both sexes

2003.....	1,105	1,337	1,361	1,349	1,487	1,449	1,061	513	173	28	3	-
2002.....	1,158	1,276	1,284	1,407	1,547	1,454	981	536	174	39	4	-
2001.....	1,121	1,174	1,290	1,354	1,469	1,506	1,060	528	157	31	5	-
2000.....	1,109	1,110	1,227	1,409	1,623	1,437	1,041	521	166	28	2	-
1999.....	1,038	1,206	1,247	1,498	1,621	1,543	949	496	144	22	1	-

All origins, male

2003.....	689	741	789	741	815	745	522	214	65	8	2	-
2002.....	665	773	721	819	859	750	460	213	54	10	2	-
2001.....	673	734	736	763	826	777	511	202	48	9	2	-
2000.....	667	660	719	789	889	705	482	212	53	9	1	-
1999.....	613	670	747	858	870	785	441	192	47	7	1	-

All origins, female

2003.....	416	596	572	608	672	704	539	299	108	20	1	-
2002.....	493	503	563	588	688	704	521	323	120	29	2	-
2001.....	448	440	554	591	643	729	549	326	109	22	3	-
2000.....	442	450	508	620	734	732	559	309	113	19	1	-
1999.....	425	536	500	640	751	758	508	304	97	15	-	-

Hispanic, both sexes

2003.....	53	57	66	56	60	64	34	15	3	-	-	-
2002.....	73	53	62	60	58	54	28	17	6	-	1	-
2001.....	45	47	62	60	57	47	35	14	4	-	-	-
2000.....	51	36	70	58	47	53	19	19	4	1	-	-
1999.....	51	47	52	47	57	38	29	12	3	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Hispanic, male

2003.....	31	28	34	31	33	25	18	9	2	-	-	-
2002.....	50	27	35	36	29	27	12	8	1	-	1	-
2001.....	21	31	28	32	40	24	16	6	-	-	-	-
2000.....	18	20	42	32	27	27	11	10	1	-	-	-
1999.....	24	24	36	25	23	19	12	1	1	1	-	-

Hispanic, female

2003.....	22	29	32	25	27	39	16	6	1	-	-	-
2002.....	23	26	27	24	29	27	16	9	5	-	-	-
2001.....	24	16	34	28	17	23	19	8	4	-	-	-
2000.....	33	16	28	26	20	26	8	9	3	1	-	-
1999.....	27	23	16	22	34	19	17	11	2	1	-	-

Non-Hispanic 1/, both sexes

2003.....	1,050	1,280	1,293	1,290	1,425	1,384	1,024	498	169	28	3	-
2002.....	1,081	1,219	1,219	1,345	1,486	1,396	953	518	168	39	3	-
2001.....	1,075	1,124	1,222	1,292	1,411	1,456	1,023	513	153	31	5	-
2000.....	1,055	1,070	1,152	1,348	1,575	1,381	1,017	500	162	27	2	-
1999.....	986	1,157	1,191	1,449	1,561	1,501	917	482	141	20	1	-

Non-Hispanic 1/, male

2003.....	657	713	755	708	781	720	503	205	63	8	2	-
2002.....	612	744	683	782	830	720	448	205	53	10	1	-
2001.....	651	700	706	731	785	751	494	196	48	9	2	-
2000.....	646	639	674	754	861	677	468	201	52	9	1	-
1999.....	588	644	709	832	846	764	427	191	46	6	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic 1/, female

2003.....	393	567	538	582	644	664	521	293	106	20	1	-
2002.....	469	475	536	563	656	676	505	313	115	29	2	-
2001.....	424	424	516	561	626	705	529	317	105	22	3	-
2000.....	409	431	478	594	714	704	549	299	110	18	1	-
1999.....	398	513	482	617	715	737	490	291	95	14	-	-

Non-Hispanic White, both sexes

2003.....	953	1,168	1,205	1,190	1,330	1,305	971	463	161	22	3	-
2002.....	982	1,112	1,131	1,251	1,395	1,310	899	490	158	36	3	-
2001.....	984	1,023	1,132	1,188	1,321	1,385	971	485	144	26	4	-
2000.....	974	980	1,067	1,261	1,495	1,306	954	466	154	26	1	-
1999.....	899	1,074	1,097	1,351	1,488	1,423	870	459	128	18	1	-

Non-Hispanic White, male

2003.....	593	654	708	653	732	677	476	192	60	7	2	-
2002.....	567	673	633	733	782	677	420	197	50	9	1	-
2001.....	598	642	651	685	742	710	471	184	43	7	1	-
2000.....	589	591	628	713	818	641	445	190	50	9	-	-
1999.....	536	599	653	781	810	725	411	182	44	5	1	-

Non-Hispanic White, female

2003.....	360	514	497	537	598	628	495	271	101	15	1	-
2002.....	415	439	498	518	613	633	479	293	108	27	2	-
2001.....	386	381	481	503	579	675	500	301	101	19	3	-
2000.....	385	389	439	548	677	665	509	276	104	17	1	-
1999.....	363	475	444	570	678	698	459	277	84	13	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Non-Hispanic Black, both sexes

2003.....	77	87	64	77	72	61	41	23	7	6	-	-
2002.....	76	82	62	72	70	69	43	20	9	3	-	-
2001.....	69	76	69	83	70	52	40	24	8	5	1	-
2000.....	68	72	66	67	66	58	51	25	6	1	1	-
1999.....	69	70	72	77	55	66	38	21	9	1	-	-

Non-Hispanic Black, male

2003.....	51	47	34	46	36	35	19	4	2	1	-	-
2002.....	31	54	38	35	35	32	23	4	2	1	-	-
2001.....	41	44	43	36	34	25	15	8	4	2	1	-
2000.....	48	39	37	29	37	25	18	6	2	-	1	-
1999.....	42	35	41	42	26	30	10	8	2	-	-	-

Non-Hispanic Black, female

2003.....	26	40	30	31	36	26	22	19	5	5	-	-
2002.....	45	28	24	37	35	37	20	16	7	2	-	-
2001.....	28	32	26	47	36	27	25	16	4	3	-	-
2000.....	20	33	29	38	29	33	33	19	4	1	-	-
1999.....	27	35	31	35	29	36	28	13	7	1	-	-

Origin not stated 2/, both sexes

2003.....	2	-	2	3	2	1	3	-	1	-	-	-
2002.....	4	4	3	2	3	4	-	1	-	-	-	-
2001.....	1	3	6	2	1	3	2	1	-	-	-	-
2000.....	3	4	5	3	1	3	5	2	-	-	-	-
1999.....	1	2	4	2	3	4	3	2	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Origin not stated 2/, male

2003.....	1	-	-	2	1	-	1	-	-	-	-	-
2002.....	3	2	3	1	-	3	-	-	-	-	-	-
2001.....	1	3	2	-	1	2	1	-	-	-	-	-
2000.....	3	1	3	3	1	1	3	1	-	-	-	-
1999.....	1	2	2	1	1	2	2	-	-	-	-	-

Origin not stated 2/, female

2003.....	1	-	2	1	1	1	2	-	1	-	-	-
2002.....	1	2	-	1	3	1	-	1	-	-	-	-
2001.....	-	-	4	2	-	1	1	1	-	-	-	-
2000.....	-	3	2	-	-	2	2	1	-	-	-	-
1999.....	-	-	2	1	2	2	1	2	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins, both sexes

2003.....	2,228	3,171	4,300	5,384	7,501	9,497	8,945	6,087	2,541	644	62	-
2002.....	2,361	3,217	4,283	5,672	7,836	9,319	8,970	6,003	2,433	574	62	-
2001.....	2,390	3,224	4,231	5,602	8,057	9,580	8,677	5,848	2,344	536	77	-
2000.....	2,350	3,156	4,128	5,932	8,109	9,743	8,705	5,716	2,329	559	70	-
1999.....	2,302	3,346	4,208	5,996	8,336	9,596	8,493	5,468	2,233	526	55	-

All origins, male

2003.....	1,347	1,929	2,628	3,255	4,295	5,068	4,456	2,722	943	202	16	-
2002.....	1,388	1,947	2,523	3,391	4,512	5,047	4,506	2,624	906	186	24	-
2001.....	1,440	1,900	2,512	3,316	4,543	5,065	4,262	2,537	846	168	25	-
2000.....	1,384	1,850	2,355	3,445	4,593	5,168	4,231	2,475	783	185	15	-
1999.....	1,382	1,957	2,469	3,436	4,631	5,102	3,998	2,337	816	155	21	-

All origins, female

2003.....	881	1,242	1,672	2,129	3,206	4,429	4,489	3,365	1,598	442	46	-
2002.....	973	1,270	1,760	2,281	3,324	4,272	4,464	3,379	1,527	388	38	-
2001.....	950	1,324	1,719	2,286	3,514	4,515	4,415	3,311	1,498	368	52	-
2000.....	966	1,306	1,773	2,487	3,516	4,575	4,474	3,241	1,546	374	55	-
1999.....	920	1,389	1,739	2,560	3,705	4,494	4,495	3,131	1,417	371	34	-

Hispanic, both sexes

2003.....	164	208	217	263	313	362	273	156	81	11	2	-
2002.....	147	206	231	311	323	337	267	164	58	15	2	-
2001.....	161	179	263	271	326	346	227	148	53	11	3	-
2000.....	142	172	255	274	359	317	206	143	53	19	1	-
1999.....	158	182	214	287	319	273	212	117	60	11	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Hispanic, male

2003.....	91	116	125	143	168	177	122	61	33	2	1	-
2002.....	87	116	129	167	164	169	121	79	22	6	1	-
2001.....	92	108	143	159	179	205	113	65	23	4	3	-
2000.....	77	106	141	156	182	155	103	64	25	9	-	-
1999.....	91	102	124	145	167	137	89	42	18	3	-	-

Hispanic, female

2003.....	73	92	92	120	145	185	151	95	48	9	1	-
2002.....	60	90	102	144	159	168	146	85	36	9	1	-
2001.....	69	71	120	112	147	141	114	83	30	7	-	-
2000.....	65	66	114	118	177	162	103	79	28	10	1	-
1999.....	67	80	90	142	152	136	123	75	42	8	-	-

Non-Hispanic 1/, both sexes

2003.....	2,054	2,955	4,074	5,110	7,174	9,117	8,660	5,916	2,457	632	60	-
2002.....	2,204	3,005	4,043	5,344	7,495	8,959	8,677	5,829	2,372	554	60	-
2001.....	2,223	3,040	3,954	5,317	7,710	9,211	8,434	5,691	2,287	525	74	-
2000.....	2,203	2,973	3,860	5,643	7,736	9,406	8,487	5,564	2,270	540	69	-
1999.....	2,139	3,158	3,985	5,698	8,006	9,299	8,262	5,343	2,167	514	55	-

Non-Hispanic 1/, male

2003.....	1,249	1,807	2,497	3,106	4,123	4,880	4,328	2,653	908	199	15	-
2002.....	1,295	1,826	2,391	3,213	4,333	4,863	4,373	2,542	884	179	23	-
2001.....	1,344	1,790	2,361	3,151	4,354	4,843	4,142	2,467	822	164	22	-
2000.....	1,302	1,736	2,205	3,279	4,403	5,005	4,122	2,408	756	176	15	-
1999.....	1,290	1,851	2,344	3,284	4,456	4,953	3,900	2,291	797	152	21	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic 1/, female

2003.....	805	1,148	1,577	2,004	3,051	4,237	4,332	3,263	1,549	433	45	-
2002.....	909	1,179	1,652	2,131	3,162	4,096	4,304	3,287	1,488	375	37	-
2001.....	879	1,250	1,593	2,166	3,356	4,368	4,292	3,224	1,465	361	52	-
2000.....	901	1,237	1,655	2,364	3,333	4,401	4,365	3,156	1,514	364	54	-
1999.....	849	1,307	1,641	2,414	3,550	4,346	4,362	3,052	1,370	362	34	-

Non-Hispanic White, both sexes

2003.....	1,629	2,501	3,495	4,433	6,346	8,269	7,984	5,530	2,284	575	51	-
2002.....	1,759	2,485	3,445	4,651	6,676	8,160	8,042	5,450	2,202	515	52	-
2001.....	1,816	2,523	3,355	4,581	6,911	8,395	7,804	5,296	2,131	484	69	-
2000.....	1,813	2,514	3,298	4,968	6,943	8,611	7,917	5,146	2,088	493	58	-
1999.....	1,760	2,654	3,388	5,004	7,204	8,487	7,685	4,971	2,004	471	47	-

Non-Hispanic White, male

2003.....	999	1,539	2,180	2,745	3,726	4,510	4,042	2,502	846	182	14	-
2002.....	1,060	1,541	2,056	2,833	3,913	4,480	4,077	2,398	818	169	19	-
2001.....	1,115	1,498	2,021	2,748	3,945	4,456	3,851	2,312	768	151	21	-
2000.....	1,070	1,478	1,901	2,939	3,960	4,623	3,868	2,246	698	161	14	-
1999.....	1,079	1,569	2,022	2,919	4,048	4,538	3,664	2,131	733	139	17	-

Non-Hispanic White, female

2003.....	630	962	1,315	1,688	2,620	3,759	3,942	3,028	1,438	393	37	-
2002.....	699	944	1,389	1,818	2,763	3,680	3,965	3,052	1,384	346	33	-
2001.....	701	1,025	1,334	1,833	2,966	3,939	3,953	2,984	1,363	333	48	-
2000.....	743	1,036	1,397	2,029	2,983	3,988	4,049	2,900	1,390	332	44	-
1999.....	681	1,085	1,366	2,085	3,156	3,949	4,021	2,840	1,271	332	30	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Non-Hispanic Black, both sexes

2003.....	371	368	485	561	676	680	543	313	148	43	8	-
2002.....	357	439	496	579	689	647	519	322	145	34	7	-
2001.....	329	427	488	618	670	661	522	329	134	37	4	-
2000.....	327	390	478	561	671	639	462	353	158	38	8	-
1999.....	318	431	486	582	665	665	477	317	138	39	8	-

Non-Hispanic Black, male

2003.....	221	226	265	296	317	296	225	117	49	11	-	-
2002.....	187	240	277	313	366	299	231	120	52	10	4	-
2001.....	192	239	273	343	336	300	238	123	44	11	1	-
2000.....	187	222	259	281	372	311	199	130	49	13	1	-
1999.....	180	242	258	309	336	333	188	122	50	10	4	-

Non-Hispanic Black, female

2003.....	150	142	220	265	359	384	318	196	99	32	8	-
2002.....	170	199	219	266	323	348	288	202	93	24	3	-
2001.....	137	188	215	275	334	361	284	206	90	26	3	-
2000.....	140	168	219	280	299	328	263	223	109	25	7	-
1999.....	138	189	228	273	329	332	289	195	88	29	4	-

Origin not stated 2/, both sexes

2003.....	10	8	9	11	14	18	12	15	3	1	-	-
2002.....	10	6	9	17	18	23	26	10	3	5	-	-
2001.....	6	5	14	14	21	23	16	9	4	-	-	-
2000.....	5	11	13	15	14	20	12	9	6	-	-	-
1999.....	5	6	9	11	11	24	19	8	6	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Origin not stated 2/, male

2003.....	7	6	6	6	4	11	6	8	2	1	-	-
2002.....	6	5	3	11	15	15	12	3	-	1	-	-
2001.....	4	2	8	6	10	17	7	5	1	-	-	-
2000.....	5	8	9	10	8	8	6	3	2	-	-	-
1999.....	1	4	1	7	8	12	9	4	1	-	-	-

Origin not stated 2/, female

2003.....	3	2	3	5	10	7	6	7	1	-	-	-
2002.....	4	1	6	6	3	8	14	7	3	4	-	-
2001.....	2	3	6	8	11	6	9	4	3	-	-	-
2000.....	-	3	4	5	6	12	6	6	4	-	-	-
1999.....	4	2	8	4	3	12	10	4	5	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hodgkin's disease (C81)

##### All origins, both sexes

2003.....	83	98	100	83	133	153	124	75	34	3	-	-
2002.....	94	74	94	94	138	126	121	83	35	7	-	-
2001.....	91	61	82	103	108	151	127	66	24	7	-	-
2000.....	88	64	94	92	140	134	121	70	24	10	-	-
1999.....	83	87	91	120	163	149	113	70	22	5	1	-

##### All origins, male

2003.....	47	57	62	52	76	73	59	31	7	-	-	-
2002.....	52	53	53	62	84	60	65	37	11	2	-	-
2001.....	56	40	45	65	64	75	55	26	7	1	-	-
2000.....	53	30	57	58	76	71	50	30	10	5	-	-
1999.....	50	54	61	72	88	69	50	28	16	1	-	-

##### All origins, female

2003.....	36	41	38	31	57	80	65	44	27	3	-	-
2002.....	42	21	41	32	54	66	56	46	24	5	-	-
2001.....	35	21	37	38	44	76	72	40	17	6	-	-
2000.....	35	34	37	34	64	63	71	40	14	5	-	-
1999.....	33	33	30	48	75	80	63	42	6	4	1	-

##### Hispanic, both sexes

2003.....	8	4	4	8	11	9	3	4	1	1	-	-
2002.....	7	7	3	6	7	7	11	2	1	-	-	-
2001.....	11	8	6	11	10	6	5	5	-	-	-	-
2000.....	7	1	12	4	7	12	11	2	1	1	-	-
1999.....	11	7	10	14	12	9	9	5	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

Hispanic, male

2003.....	5	2	3	4	7	6	2	1	-	-	-	-
2002.....	5	4	3	2	4	2	6	1	-	-	-	-
2001.....	10	8	3	8	7	3	2	3	-	-	-	-
2000.....	5	-	8	4	6	9	3	-	1	1	-	-
1999.....	8	4	5	7	6	5	6	2	-	-	-	-

Hispanic, female

2003.....	3	2	1	4	4	3	1	3	1	1	-	-
2002.....	2	3	-	4	3	5	5	1	1	-	-	-
2001.....	1	-	3	3	3	3	3	2	-	-	-	-
2000.....	2	1	4	-	1	3	8	2	-	-	-	-
1999.....	3	3	5	7	6	4	3	3	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	74	94	96	75	122	144	120	71	33	2	-	-
2002.....	86	67	91	88	131	118	110	81	34	7	-	-
2001.....	80	53	76	92	97	145	122	61	24	7	-	-
2000.....	81	62	81	88	133	122	109	68	23	9	-	-
1999.....	72	79	80	106	151	140	103	65	22	5	1	-

Non-Hispanic 1/, male

2003.....	42	55	59	48	69	67	57	30	7	-	-	-
2002.....	46	49	50	60	80	57	59	36	11	2	-	-
2001.....	46	32	42	57	57	72	53	23	7	1	-	-
2000.....	48	29	49	54	70	62	46	30	9	4	-	-
1999.....	42	49	56	65	82	64	44	26	16	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hodgkin's disease (C81)												
Non-Hispanic 1/, female												
2003.....	32	39	37	27	53	77	63	41	26	2	-	-
2002.....	40	18	41	28	51	61	51	45	23	5	-	-
2001.....	34	21	34	35	40	73	69	38	17	6	-	-
2000.....	33	33	32	34	63	60	63	38	14	5	-	-
1999.....	30	30	24	41	69	76	59	39	6	4	1	-
Non-Hispanic White, both sexes												
2003.....	66	75	87	63	114	134	112	67	32	2	-	-
2002.....	71	54	78	79	121	111	104	79	32	6	-	-
2001.....	73	44	66	88	90	132	117	59	24	7	-	-
2000.....	68	54	69	82	118	122	102	66	21	8	-	-
1999.....	63	70	74	92	142	135	100	57	21	5	1	-
Non-Hispanic White, male												
2003.....	36	43	53	39	62	61	54	27	7	-	-	-
2002.....	35	39	41	55	75	54	57	36	10	2	-	-
2001.....	41	27	36	55	54	67	50	21	7	1	-	-
2000.....	40	25	40	51	61	62	43	29	8	4	-	-
1999.....	34	44	53	57	78	62	43	23	15	1	-	-
Non-Hispanic White, female												
2003.....	30	32	34	24	52	73	58	40	25	2	-	-
2002.....	36	15	37	24	46	57	47	43	22	4	-	-
2001.....	32	17	30	33	36	65	67	38	17	6	-	-
2000.....	28	29	29	31	57	60	59	37	13	4	-	-
1999.....	29	26	21	35	64	73	57	34	6	4	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

Non-Hispanic Black, both sexes

2003.....	8	15	6	11	6	6	2	-	-	-	-	-
2002.....	10	10	10	6	9	5	4	1	1	1	-	-
2001.....	5	9	9	4	7	9	3	2	-	-	-	-
2000.....	10	8	9	4	11	-	5	1	2	-	-	-
1999.....	9	9	5	11	7	5	1	6	1	-	-	-

Non-Hispanic Black, male

2003.....	6	10	5	8	5	3	1	-	-	-	-	-
2002.....	7	8	7	3	5	3	1	-	1	-	-	-
2001.....	4	5	5	2	3	3	2	2	-	-	-	-
2000.....	5	4	7	2	5	-	3	-	1	-	-	-
1999.....	8	5	3	8	4	2	1	1	1	-	-	-

Non-Hispanic Black, female

2003.....	2	5	1	3	1	3	1	-	-	-	-	-
2002.....	3	2	3	3	4	2	3	1	-	1	-	-
2001.....	1	4	4	2	4	6	1	-	-	-	-	-
2000.....	5	4	2	2	6	-	2	1	1	-	-	-
1999.....	1	4	2	3	3	3	-	5	-	-	-	-

Origin not stated 2/, both sexes

2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	1	1	-	-	-	1	-	-	-	-	-
1999.....	-	1	1	-	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hodgkin's disease (C81)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	-	-	-	1	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	1	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	1	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)												
All origins, both sexes												
2003.....	899	1,272	1,735	2,104	2,815	3,674	3,543	2,520	949	213	30	-
2002.....	943	1,319	1,730	2,225	3,039	3,722	3,561	2,446	881	200	22	-
2001.....	1,021	1,367	1,731	2,193	3,165	3,776	3,593	2,392	904	186	24	-
2000.....	1,015	1,434	1,736	2,386	3,212	3,984	3,551	2,330	856	193	23	-
1999.....	1,045	1,414	1,740	2,441	3,407	4,009	3,545	2,198	834	178	18	-
All origins, male												
2003.....	558	773	1,075	1,260	1,569	1,882	1,681	1,065	345	78	9	-
2002.....	574	823	1,015	1,334	1,652	1,898	1,696	1,013	315	51	6	-
2001.....	663	817	1,008	1,286	1,739	1,873	1,685	1,020	305	55	5	-
2000.....	646	865	984	1,342	1,763	2,018	1,616	959	274	62	4	-
1999.....	652	851	996	1,355	1,821	2,052	1,544	892	292	53	9	-
All origins, female												
2003.....	341	499	660	844	1,246	1,792	1,862	1,455	604	135	21	-
2002.....	369	496	715	891	1,387	1,824	1,865	1,433	566	149	16	-
2001.....	358	550	723	907	1,426	1,903	1,908	1,372	599	131	19	-
2000.....	369	569	752	1,044	1,449	1,966	1,935	1,371	582	131	19	-
1999.....	393	563	744	1,086	1,586	1,957	2,001	1,306	542	125	9	-
Hispanic, both sexes												
2003.....	56	94	82	105	128	142	117	67	32	4	2	-
2002.....	56	76	110	122	146	155	107	65	19	6	1	-
2001.....	67	83	118	91	117	147	93	68	21	5	1	-
2000.....	62	78	112	111	166	128	82	63	18	9	-	-
1999.....	67	62	91	117	144	117	88	51	25	4	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

Hispanic, male

2003.....	31	55	49	58	71	67	51	19	16	1	1	-
2002.....	31	46	63	69	67	66	45	26	8	2	-	-
2001.....	39	50	65	55	67	82	49	28	5	2	1	-
2000.....	38	48	64	60	83	60	38	28	9	4	-	-
1999.....	42	38	53	56	69	59	33	20	8	-	-	-

Hispanic, female

2003.....	25	39	33	47	57	75	66	48	16	3	1	-
2002.....	25	30	47	53	79	89	62	39	11	4	1	-
2001.....	28	33	53	36	50	65	44	40	16	3	-	-
2000.....	24	30	48	51	83	68	44	35	9	5	-	-
1999.....	25	24	38	61	75	58	55	31	17	4	-	-

Non-Hispanic 1/, both sexes

2003.....	839	1,176	1,647	1,996	2,683	3,526	3,421	2,448	915	209	28	-
2002.....	883	1,240	1,617	2,095	2,888	3,561	3,438	2,377	861	193	21	-
2001.....	952	1,282	1,610	2,093	3,041	3,619	3,494	2,322	881	181	23	-
2000.....	949	1,353	1,617	2,271	3,040	3,848	3,464	2,263	836	184	23	-
1999.....	976	1,350	1,646	2,318	3,257	3,880	3,447	2,144	807	173	18	-

Non-Hispanic 1/, male

2003.....	525	716	1,023	1,200	1,497	1,812	1,628	1,044	328	77	8	-
2002.....	542	774	951	1,258	1,580	1,828	1,644	987	307	49	6	-
2001.....	623	767	941	1,227	1,670	1,783	1,633	991	300	53	4	-
2000.....	604	814	914	1,279	1,675	1,955	1,578	929	265	58	4	-
1999.....	609	812	942	1,296	1,748	1,986	1,504	870	284	53	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Non-Hodgkin's lymphoma (C82-C85)												
Non-Hispanic 1/, female												
2003.....	314	460	624	796	1,186	1,714	1,793	1,404	587	132	20	-
2002.....	341	466	666	837	1,308	1,733	1,794	1,390	554	144	15	-
2001.....	329	515	669	866	1,371	1,836	1,861	1,331	581	128	19	-
2000.....	345	539	703	992	1,365	1,893	1,886	1,334	571	126	19	-
1999.....	367	538	704	1,022	1,509	1,894	1,943	1,274	523	120	9	-
Non-Hispanic White, both sexes												
2003.....	694	1,044	1,456	1,800	2,442	3,289	3,244	2,351	870	187	27	-
2002.....	719	1,074	1,443	1,886	2,637	3,339	3,258	2,278	827	185	20	-
2001.....	809	1,105	1,430	1,870	2,813	3,400	3,302	2,197	832	173	21	-
2000.....	814	1,209	1,446	2,065	2,813	3,626	3,312	2,149	797	176	21	-
1999.....	855	1,188	1,479	2,129	3,016	3,632	3,303	2,037	762	160	16	-
Non-Hispanic White, male												
2003.....	440	642	914	1,103	1,387	1,710	1,557	1,009	309	68	8	-
2002.....	449	678	848	1,143	1,457	1,725	1,562	950	291	45	6	-
2001.....	531	661	836	1,108	1,547	1,678	1,548	936	284	50	4	-
2000.....	515	720	818	1,181	1,544	1,854	1,511	890	251	55	4	-
1999.....	535	713	841	1,190	1,632	1,847	1,455	818	266	47	7	-
Non-Hispanic White, female												
2003.....	254	402	542	697	1,055	1,579	1,687	1,342	561	119	19	-
2002.....	270	396	595	743	1,180	1,614	1,696	1,328	536	140	14	-
2001.....	278	444	594	762	1,266	1,722	1,754	1,261	548	123	17	-
2000.....	299	489	628	884	1,269	1,772	1,801	1,259	546	121	17	-
1999.....	320	475	638	939	1,384	1,785	1,848	1,219	496	113	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's Lymphoma (C82-C85)

Non-Hispanic Black, both sexes

2003.....	127	95	149	138	169	155	120	62	29	13	1	-
2002.....	126	136	141	165	191	153	131	74	24	5	-	-
2001.....	118	137	132	174	174	146	138	83	36	7	2	-
2000.....	102	119	137	150	168	145	99	79	30	6	-	-
1999.....	97	138	121	153	177	179	107	78	34	11	2	-

Non-Hispanic Black, male

2003.....	74	55	83	70	73	62	46	18	11	4	-	-
2002.....	71	76	83	91	102	68	54	27	9	4	-	-
2001.....	79	77	74	96	94	65	59	35	11	2	-	-
2000.....	67	82	75	74	95	65	38	22	12	3	-	-
1999.....	62	81	72	90	84	99	32	32	11	5	2	-

Non-Hispanic Black, female

2003.....	53	40	66	68	96	93	74	44	18	9	1	-
2002.....	55	60	58	74	89	85	77	47	15	1	-	-
2001.....	39	60	58	78	80	81	79	48	25	5	2	-
2000.....	35	37	62	76	73	80	61	57	18	3	-	-
1999.....	35	57	49	63	93	80	75	46	23	6	-	-

Origin not stated 2/, both sexes

2003.....	4	2	6	3	4	6	5	5	2	-	-	-
2002.....	4	3	3	8	5	6	16	4	1	1	-	-
2001.....	2	2	3	9	7	10	6	2	2	-	-	-
2000.....	4	3	7	4	6	8	5	4	2	-	-	-
1999.....	2	2	3	6	6	12	10	3	2	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Non-Hodgkin's lymphoma (C82-C85)

Origin not stated 2/, male

2003.....	2	2	3	2	1	3	2	2	1	-	-	-
2002.....	1	3	1	7	5	4	7	-	-	-	-	-
2001.....	1	-	2	4	2	8	3	1	-	-	-	-
2000.....	4	3	6	3	5	3	-	2	-	-	-	-
1999.....	1	1	1	3	4	7	7	2	-	-	-	-

Origin not stated 2/, female

2003.....	2	-	3	1	3	3	3	3	1	-	-	-
2002.....	3	-	2	1	-	2	9	4	1	1	-	-
2001.....	1	2	1	5	5	2	3	1	2	-	-	-
2000.....	-	-	1	1	1	5	5	2	2	-	-	-
1999.....	1	1	2	3	2	5	3	1	2	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>Leukemia (C91-C95)</b>												
<b>All origins, both sexes</b>												
2003.....	770	1,102	1,503	1,911	2,795	3,532	3,405	2,317	1,091	350	28	-
2002.....	853	1,111	1,500	1,998	2,820	3,392	3,371	2,294	1,086	281	35	-
2001.....	845	1,073	1,462	1,963	2,923	3,541	3,155	2,256	1,008	271	50	-
2000.....	789	1,041	1,373	2,073	2,937	3,515	3,218	2,194	1,041	271	37	-
1999.....	744	1,112	1,404	2,078	2,934	3,401	3,059	2,151	984	258	31	-
<b>All origins, male</b>												
2003.....	454	683	910	1,215	1,715	2,054	1,826	1,102	416	91	6	-
2002.....	486	667	891	1,242	1,768	1,995	1,809	1,062	420	95	15	-
2001.....	467	632	920	1,216	1,746	2,023	1,676	1,041	380	89	17	-
2000.....	435	629	823	1,287	1,803	1,982	1,713	1,018	359	85	8	-
1999.....	431	636	853	1,275	1,779	1,946	1,579	964	375	75	9	-
<b>All origins, female</b>												
2003.....	316	419	593	696	1,080	1,478	1,579	1,215	675	259	22	-
2002.....	367	444	609	756	1,052	1,397	1,562	1,232	666	186	20	-
2001.....	378	441	542	747	1,177	1,518	1,479	1,215	628	182	33	-
2000.....	354	412	550	786	1,134	1,533	1,505	1,176	682	186	29	-
1999.....	313	476	551	803	1,155	1,455	1,480	1,187	609	183	22	-
<b>Hispanic, both sexes</b>												
2003.....	62	57	75	81	89	115	96	49	38	4	-	-
2002.....	42	74	69	88	95	81	83	53	25	7	-	-
2001.....	51	47	73	90	107	113	72	44	17	5	2	-
2000.....	52	61	72	81	101	95	70	44	21	8	1	-
1999.....	50	62	53	86	93	85	64	36	22	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>Leukemia (C91-C95)</b>												
<b>Hispanic, male</b>												
2003.....	32	29	46	45	54	59	47	25	15	-	-	-
2002.....	29	36	31	45	52	50	44	26	9	4	-	-
2001.....	26	26	47	46	52	72	30	22	10	2	2	-
2000.....	26	41	41	45	53	48	39	25	10	3	-	-
1999.....	22	31	35	45	62	45	30	17	5	1	-	-
<b>Hispanic, female</b>												
2003.....	30	28	29	36	35	56	49	24	23	4	-	-
2002.....	13	38	38	43	43	31	39	27	16	3	-	-
2001.....	25	21	26	44	55	41	42	22	7	3	-	-
2000.....	26	20	31	36	48	47	31	19	11	5	1	-
1999.....	28	31	18	41	31	40	34	19	17	4	-	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	704	1,041	1,426	1,826	2,700	3,412	3,306	2,262	1,052	345	28	-
2002.....	806	1,036	1,428	1,906	2,716	3,305	3,281	2,238	1,060	271	35	-
2001.....	793	1,023	1,383	1,869	2,808	3,423	3,078	2,207	989	266	48	-
2000.....	736	974	1,300	1,984	2,833	3,411	3,144	2,147	1,017	263	36	-
1999.....	691	1,048	1,347	1,989	2,837	3,308	2,990	2,113	959	253	31	-
<b>Non-Hispanic 1/, male</b>												
2003.....	418	651	862	1,167	1,658	1,992	1,776	1,073	400	90	6	-
2002.....	453	630	859	1,195	1,709	1,940	1,762	1,034	411	90	15	-
2001.....	440	604	870	1,168	1,689	1,947	1,643	1,016	369	87	15	-
2000.....	408	584	782	1,238	1,749	1,931	1,670	992	348	82	8	-
1999.....	409	603	818	1,227	1,713	1,898	1,548	947	370	74	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
Non-Hispanic 1/, female												
2003.....	286	390	564	659	1,042	1,420	1,530	1,189	652	255	22	-
2002.....	353	406	569	711	1,007	1,365	1,519	1,204	649	181	20	-
2001.....	353	419	513	701	1,119	1,476	1,435	1,191	620	179	33	-
2000.....	328	390	518	746	1,084	1,480	1,474	1,155	669	181	28	-
1999.....	282	445	529	762	1,124	1,410	1,442	1,166	589	179	22	-
Non-Hispanic White, both sexes												
2003.....	572	904	1,247	1,620	2,434	3,149	3,072	2,125	980	324	23	-
2002.....	673	869	1,230	1,704	2,465	3,024	3,063	2,085	993	251	30	-
2001.....	644	879	1,205	1,630	2,555	3,139	2,892	2,084	928	244	46	-
2000.....	618	835	1,117	1,786	2,586	3,168	2,953	2,008	945	243	31	-
1999.....	567	911	1,164	1,751	2,597	3,073	2,775	1,999	892	233	28	-
Non-Hispanic White, male												
2003.....	342	563	772	1,045	1,515	1,865	1,675	1,018	374	84	6	-
2002.....	386	543	747	1,072	1,559	1,793	1,653	973	387	85	12	-
2001.....	365	527	762	1,024	1,555	1,803	1,546	968	348	80	15	-
2000.....	340	506	677	1,129	1,597	1,804	1,573	924	327	74	7	-
1999.....	338	530	716	1,094	1,579	1,767	1,453	894	345	68	9	-
Non-Hispanic White, female												
2003.....	230	341	475	575	919	1,284	1,397	1,107	606	240	17	-
2002.....	287	326	483	632	906	1,231	1,410	1,112	606	166	18	-
2001.....	279	352	443	606	1,000	1,336	1,346	1,116	580	164	31	-
2000.....	278	329	440	657	989	1,364	1,380	1,084	618	169	24	-
1999.....	229	381	448	657	1,018	1,306	1,322	1,105	547	165	19	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Leukemia (C91-C95)												
Non-Hispanic Black, both sexes												
2003.....	109	111	151	169	220	216	188	115	67	16	5	-
2002.....	104	136	159	160	212	228	177	131	61	19	5	-
2001.....	112	112	136	202	206	236	154	108	54	20	1	-
2000.....	97	106	152	160	213	195	154	122	63	18	5	-
1999.....	95	106	146	200	205	196	175	96	56	18	3	-
Non-Hispanic Black, male												
2003.....	66	75	75	99	119	108	79	46	23	5	-	-
2002.....	52	74	87	93	127	114	84	51	21	5	3	-
2001.....	61	63	84	125	107	111	76	42	17	6	-	-
2000.....	53	60	89	86	130	105	76	60	16	7	1	-
1999.....	56	61	81	111	111	108	70	42	20	4	-	-
Non-Hispanic Black, female												
2003.....	43	36	76	70	101	108	109	69	44	11	5	-
2002.....	52	62	72	67	85	114	93	80	40	14	2	-
2001.....	51	49	52	77	99	125	78	66	37	14	1	-
2000.....	44	46	63	74	83	90	78	62	47	11	4	-
1999.....	39	45	65	89	94	88	105	54	36	14	3	-
Origin not stated 2/, both sexes												
2003.....	4	4	2	4	6	5	3	6	1	1	-	-
2002.....	5	1	3	4	9	6	7	3	1	3	-	-
2001.....	1	3	6	4	8	5	5	5	2	-	-	-
2000.....	1	6	1	8	3	9	4	3	3	-	-	-
1999.....	3	2	4	3	4	8	5	2	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Leukemia (C91-C95)

Origin not stated 2/, male

2003.....	4	3	2	3	3	3	3	4	1	1	-	-
2002.....	4	1	1	2	7	5	3	2	-	1	-	-
2001.....	1	2	3	2	5	4	3	3	1	-	-	-
2000.....	1	4	-	4	1	3	4	1	1	-	-	-
1999.....	-	2	-	3	4	3	1	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	1	-	1	3	2	-	2	-	-	-	-
2002.....	1	-	2	2	2	1	4	1	1	2	-	-
2001.....	-	1	3	2	3	1	2	2	1	-	-	-
2000.....	-	2	1	4	2	6	-	2	2	-	-	-
1999.....	3	-	4	-	-	5	4	2	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All origins, both sexes

2003.....	473	698	961	1,280	1,749	2,127	1,870	1,165	464	78	4	-
2002.....	466	711	958	1,348	1,828	2,067	1,904	1,171	426	84	5	-
2001.....	430	721	950	1,332	1,849	2,101	1,785	1,123	400	71	3	-
2000.....	455	611	919	1,374	1,805	2,097	1,799	1,109	401	84	10	-
1999.....	424	727	967	1,347	1,825	2,023	1,765	1,040	389	85	5	-

All origins, male

2003.....	287	415	580	725	929	1,052	889	520	174	33	1	-
2002.....	274	403	563	748	998	1,090	928	507	159	37	3	-
2001.....	252	410	535	741	984	1,089	837	446	152	23	3	-
2000.....	248	321	487	751	946	1,087	845	462	137	33	3	-
1999.....	246	411	557	729	940	1,026	820	450	133	26	3	-

All origins, female

2003.....	186	283	381	555	820	1,075	981	645	290	45	3	-
2002.....	192	308	395	600	830	977	976	664	267	47	2	-
2001.....	178	311	415	591	865	1,012	948	677	248	48	-	-
2000.....	207	290	432	623	859	1,010	954	647	264	51	7	-
1999.....	178	316	410	618	885	997	945	590	256	59	2	-

Hispanic, both sexes

2003.....	38	53	56	69	85	96	57	36	10	2	-	-
2002.....	42	49	49	94	75	94	66	44	13	2	1	-
2001.....	32	41	66	77	92	80	57	31	15	1	-	-
2000.....	21	31	59	78	83	82	42	33	13	1	-	-
1999.....	30	51	60	70	69	60	51	25	13	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Hispanic, male

2003.....	23	30	27	36	36	45	22	16	2	1	-	-
2002.....	22	30	32	51	41	51	26	26	5	-	1	-
2001.....	17	24	28	49	53	48	32	12	8	-	-	-
2000.....	8	16	28	47	40	38	23	10	5	1	-	-
1999.....	19	29	31	37	30	27	20	3	5	2	-	-

Hispanic, female

2003.....	15	23	29	33	49	51	35	20	8	1	-	-
2002.....	20	19	17	43	34	43	40	18	8	2	-	-
2001.....	15	17	38	28	39	32	25	19	7	1	-	-
2000.....	13	15	31	31	43	44	19	23	8	-	-	-
1999.....	11	22	29	33	39	33	31	22	8	-	-	-

Non-Hispanic 1/, both sexes

2003.....	434	643	904	1,207	1,660	2,024	1,810	1,125	454	76	4	-
2002.....	424	660	906	1,249	1,749	1,963	1,835	1,124	412	81	4	-
2001.....	395	680	879	1,254	1,752	2,013	1,723	1,090	385	70	3	-
2000.....	434	579	856	1,293	1,717	2,012	1,755	1,074	387	83	10	-
1999.....	394	675	906	1,275	1,755	1,959	1,711	1,012	375	83	5	-

Non-Hispanic 1/, male

2003.....	263	384	552	688	893	1,002	866	502	172	32	1	-
2002.....	252	372	530	695	954	1,034	900	480	154	37	2	-
2001.....	233	386	504	692	928	1,036	804	433	144	23	3	-
2000.....	240	305	456	701	904	1,047	821	452	131	32	3	-
1999.....	227	382	526	691	910	997	799	445	127	24	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic 1/, female

2003.....	171	259	352	519	767	1,022	944	623	282	44	3	-
2002.....	172	288	376	554	795	929	935	644	258	44	2	-
2001.....	162	294	375	562	824	977	919	657	241	47	-	-
2000.....	194	274	400	592	813	965	934	622	256	51	7	-
1999.....	167	293	380	584	845	962	912	567	248	59	2	-

Non-Hispanic White, both sexes

2003.....	294	477	704	945	1,347	1,686	1,554	978	399	62	1	-
2002.....	292	486	693	977	1,442	1,677	1,606	999	348	71	2	-
2001.....	287	495	649	985	1,442	1,715	1,479	946	339	59	2	-
2000.....	310	412	660	1,028	1,414	1,684	1,536	912	319	65	6	-
1999.....	270	480	665	1,022	1,443	1,636	1,497	870	325	73	2	-

Non-Hispanic White, male

2003.....	180	290	440	555	756	867	756	444	155	30	-	-
2002.....	188	280	419	559	812	904	799	434	130	36	1	-
2001.....	176	283	383	555	779	904	699	384	127	20	2	-
2000.....	173	224	362	571	754	894	734	399	110	28	3	-
1999.....	170	278	410	573	756	855	708	393	107	23	1	-

Non-Hispanic White, female

2003.....	114	187	264	390	591	819	798	534	244	32	1	-
2002.....	104	206	274	418	630	773	807	565	218	35	1	-
2001.....	111	212	266	430	663	811	780	562	212	39	-	-
2000.....	137	188	298	457	660	790	802	513	209	37	3	-
1999.....	100	202	255	449	687	781	789	477	218	50	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Non-Hispanic Black, both sexes

2003.....	127	147	179	242	281	303	233	135	52	14	2	-
2002.....	117	157	186	247	277	259	206	116	57	9	2	-
2001.....	94	167	210	237	282	269	224	135	44	10	1	-
2000.....	118	156	180	247	279	297	203	150	62	14	3	-
1999.....	116	177	214	218	276	284	194	136	47	10	3	-

Non-Hispanic Black, male

2003.....	75	86	102	119	120	123	99	53	15	2	-	-
2002.....	57	82	100	125	132	114	91	42	21	1	1	-
2001.....	48	93	110	119	132	120	100	43	16	3	1	-
2000.....	62	75	88	119	142	140	82	47	19	3	-	-
1999.....	53	94	102	100	137	123	85	47	18	1	2	-

Non-Hispanic Black, female

2003.....	52	61	77	123	161	180	134	82	37	12	2	-
2002.....	60	75	86	122	145	145	115	74	36	8	1	-
2001.....	46	74	100	118	150	149	124	92	28	7	-	-
2000.....	56	81	92	128	137	157	121	103	43	11	3	-
1999.....	63	83	112	118	139	161	109	89	29	9	1	-

Origin not stated 2/, both sexes

2003.....	1	2	1	4	4	7	3	4	-	-	-	-
2002.....	-	2	3	5	4	10	3	3	1	1	-	-
2001.....	3	-	5	1	5	8	5	2	-	-	-	-
2000.....	-	1	4	3	5	3	2	2	1	-	-	-
1999.....	-	1	1	2	1	4	3	3	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Origin not stated 2/, male

2003.....	1	1	1	1	-	5	1	2	-	-	-	-
2002.....	-	1	1	2	3	5	2	1	-	-	-	-
2001.....	2	-	3	-	3	5	1	1	-	-	-	-
2000.....	-	-	3	3	2	2	1	-	1	-	-	-
1999.....	-	-	-	1	-	2	1	2	1	-	-	-

Origin not stated 2/, female

2003.....	-	1	-	3	4	2	2	2	-	-	-	-
2002.....	-	1	2	3	1	5	1	2	1	1	-	-
2001.....	1	-	2	1	2	3	4	1	-	-	-	-
2000.....	-	1	1	-	3	1	1	2	-	-	-	-
1999.....	-	1	1	1	1	2	2	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All origins, both sexes

2003.....	3	1	1	6	9	11	3	10	3	-	-	-
2002.....	5	2	1	7	11	12	13	9	5	2	-	-
2001.....	3	2	6	11	12	11	17	11	8	1	-	-
2000.....	3	6	6	7	15	13	16	13	7	1	-	-
1999.....	6	6	6	10	7	14	11	9	4	-	-	-

All origins, male

2003.....	1	1	1	3	6	7	1	4	1	-	-	-
2002.....	2	1	1	5	10	4	8	5	1	1	-	-
2001.....	2	1	4	8	10	5	9	4	2	-	-	-
2000.....	2	5	4	7	5	10	7	6	3	-	-	-
1999.....	3	5	2	5	3	9	5	3	-	-	-	-

All origins, female

2003.....	2	-	-	3	3	4	2	6	2	-	-	-
2002.....	3	1	-	2	1	8	5	4	4	1	-	-
2001.....	1	1	2	3	2	6	8	7	6	1	-	-
2000.....	1	1	2	-	10	3	9	7	4	1	-	-
1999.....	3	1	4	5	4	5	6	6	4	-	-	-

Hispanic, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	2	-	-	-	-	-	-	-	-
2000.....	-	1	-	-	2	-	1	1	-	-	-	-
1999.....	-	-	-	-	1	2	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	1	-	-	-	-	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	2	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	3	1	1	6	9	11	3	10	3	-	-	-
2002.....	5	2	1	6	11	12	13	9	5	2	-	-
2001.....	3	2	6	9	12	11	17	11	8	1	-	-
2000.....	3	5	6	7	13	13	15	12	7	1	-	-
1999.....	6	6	6	10	6	12	11	9	4	-	-	-

Non-Hispanic 1/, male

2003.....	1	1	1	3	6	7	1	4	1	-	-	-
2002.....	2	1	1	5	10	4	8	5	1	1	-	-
2001.....	2	1	4	7	10	5	9	4	2	-	-	-
2000.....	2	4	4	7	5	10	7	5	3	-	-	-
1999.....	3	5	2	5	3	8	5	3	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

Non-Hispanic 1/, female

2003.....	2	-	-	3	3	4	2	6	2	-	-	-
2002.....	3	1	-	1	1	8	5	4	4	1	-	-
2001.....	1	1	2	2	2	6	8	7	6	1	-	-
2000.....	1	1	2	-	8	3	8	7	4	1	-	-
1999.....	3	1	4	5	3	4	6	6	4	-	-	-

Non-Hispanic White, both sexes

2003.....	3	1	1	5	9	11	2	9	3	-	-	-
2002.....	4	2	1	5	11	9	11	9	2	2	-	-
2001.....	3	-	5	8	11	9	14	10	8	1	-	-
2000.....	3	4	6	7	12	11	14	11	6	1	-	-
1999.....	5	5	6	10	6	11	10	8	4	-	-	-

Non-Hispanic White, male

2003.....	1	1	1	3	6	7	-	4	1	-	-	-
2002.....	2	1	1	4	10	4	6	5	-	1	-	-
2001.....	2	-	4	6	10	4	8	3	2	-	-	-
2000.....	2	3	4	7	4	9	7	4	2	-	-	-
1999.....	2	4	2	5	3	7	5	3	-	-	-	-

Non-Hispanic White, female

2003.....	2	-	-	2	3	4	2	5	2	-	-	-
2002.....	2	1	-	1	1	5	5	4	2	1	-	-
2001.....	1	-	1	2	1	5	6	7	6	1	-	-
2000.....	1	1	2	-	8	2	7	7	4	1	-	-
1999.....	3	1	4	5	3	4	5	5	4	-	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins, both sexes

2003.....	3,377	4,446	5,374	6,299	7,978	9,798	9,082	6,421	3,159	847	129	5
2002.....	3,280	4,320	5,403	6,321	8,436	9,718	8,967	6,452	3,129	853	136	1
2001.....	3,319	4,278	5,163	6,406	8,617	9,902	8,798	6,343	3,019	845	119	-
2000.....	3,193	4,088	5,205	6,819	8,897	10,087	8,773	6,214	2,937	874	140	4
1999.....	3,230	4,249	5,614	7,265	9,569	10,451	9,205	6,485	2,955	851	112	2

All origins, male

2003.....	1,869	2,525	3,048	3,544	4,256	5,042	4,279	2,648	1,055	242	28	3
2002.....	1,906	2,399	3,122	3,548	4,593	4,969	4,177	2,690	1,100	226	25	1
2001.....	1,933	2,398	2,928	3,501	4,676	4,962	4,070	2,569	1,001	239	26	-
2000.....	1,773	2,286	2,937	3,791	4,781	5,098	4,063	2,460	989	218	22	4
1999.....	1,824	2,416	3,181	3,984	5,173	5,240	4,199	2,586	986	213	21	1

All origins, female

2003.....	1,508	1,921	2,326	2,755	3,722	4,756	4,803	3,773	2,104	605	101	2
2002.....	1,374	1,921	2,281	2,773	3,843	4,749	4,790	3,762	2,029	627	111	-
2001.....	1,386	1,880	2,235	2,905	3,941	4,940	4,728	3,774	2,018	606	93	-
2000.....	1,420	1,802	2,268	3,028	4,116	4,989	4,710	3,754	1,948	656	118	-
1999.....	1,406	1,833	2,433	3,281	4,396	5,211	5,006	3,899	1,969	638	91	1

Hispanic, both sexes

2003.....	220	226	268	310	407	386	315	187	107	28	4	1
2002.....	203	232	270	318	394	385	291	191	83	20	2	-
2001.....	191	200	281	295	395	347	268	189	83	35	3	-
2000.....	172	226	229	317	373	357	274	167	88	32	3	-
1999.....	170	216	255	325	346	336	251	183	80	20	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Hispanic, male

2003.....	115	124	150	169	216	194	138	74	42	8	2	-
2002.....	113	137	140	164	198	199	139	81	33	8	-	-
2001.....	103	115	147	141	198	191	124	79	20	12	1	-
2000.....	88	113	122	163	186	187	122	64	36	14	1	-
1999.....	87	115	141	161	190	159	119	73	32	4	-	-

Hispanic, female

2003.....	105	102	118	141	191	192	177	113	65	20	2	1
2002.....	90	95	130	154	196	186	152	110	50	12	2	-
2001.....	88	85	134	154	197	156	144	110	63	23	2	-
2000.....	84	113	107	154	187	170	152	103	52	18	2	-
1999.....	83	101	114	164	156	177	132	110	48	16	5	-

Non-Hispanic 1/, both sexes

2003.....	3,143	4,200	5,089	5,974	7,555	9,396	8,754	6,215	3,047	819	125	4
2002.....	3,071	4,078	5,114	5,981	8,024	9,308	8,657	6,241	3,037	830	134	1
2001.....	3,118	4,065	4,870	6,098	8,180	9,534	8,508	6,140	2,923	808	116	-
2000.....	3,009	3,851	4,962	6,484	8,496	9,693	8,479	6,031	2,844	841	137	4
1999.....	3,054	4,021	5,340	6,926	9,195	10,091	8,929	6,286	2,871	831	106	2

Non-Hispanic 1/, male

2003.....	1,746	2,390	2,885	3,364	4,032	4,842	4,134	2,568	1,011	234	26	3
2002.....	1,789	2,258	2,967	3,370	4,385	4,757	4,025	2,604	1,063	218	25	1
2001.....	1,822	2,275	2,771	3,351	4,449	4,765	3,936	2,484	978	227	25	-
2000.....	1,682	2,164	2,808	3,616	4,579	4,892	3,931	2,390	953	204	21	4
1999.....	1,734	2,295	3,030	3,816	4,963	5,065	4,067	2,509	953	209	21	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic 1/, female

2003.....	1,397	1,810	2,204	2,610	3,523	4,554	4,620	3,647	2,036	585	99	1
2002.....	1,282	1,820	2,147	2,611	3,639	4,551	4,632	3,637	1,974	612	109	-
2001.....	1,296	1,790	2,099	2,747	3,731	4,769	4,572	3,656	1,945	581	91	-
2000.....	1,327	1,687	2,154	2,868	3,917	4,801	4,548	3,641	1,891	637	116	-
1999.....	1,320	1,726	2,310	3,110	4,232	5,026	4,862	3,777	1,918	622	85	1

Non-Hispanic White, both sexes

2003.....	2,480	3,448	4,295	5,101	6,611	8,403	7,965	5,734	2,786	735	111	4
2002.....	2,442	3,367	4,268	5,072	6,999	8,306	7,853	5,705	2,785	750	121	1
2001.....	2,466	3,314	4,028	5,171	7,121	8,536	7,697	5,641	2,685	732	100	-
2000.....	2,376	3,201	4,129	5,583	7,436	8,644	7,751	5,548	2,638	752	112	3
1999.....	2,453	3,273	4,449	6,006	8,070	9,032	8,126	5,756	2,606	750	92	2

Non-Hispanic White, male

2003.....	1,393	1,965	2,454	2,889	3,551	4,367	3,802	2,388	927	207	22	3
2002.....	1,440	1,871	2,491	2,920	3,870	4,268	3,698	2,402	981	198	24	1
2001.....	1,436	1,845	2,294	2,857	3,932	4,292	3,590	2,292	887	211	21	-
2000.....	1,349	1,804	2,307	3,159	4,049	4,356	3,633	2,203	883	181	19	3
1999.....	1,390	1,885	2,538	3,321	4,393	4,580	3,742	2,313	866	184	20	1

Non-Hispanic White, female

2003.....	1,087	1,483	1,841	2,212	3,060	4,036	4,163	3,346	1,859	528	89	1
2002.....	1,002	1,496	1,777	2,152	3,129	4,038	4,155	3,303	1,804	552	97	-
2001.....	1,030	1,469	1,734	2,314	3,189	4,244	4,107	3,349	1,798	521	79	-
2000.....	1,027	1,397	1,822	2,424	3,387	4,288	4,118	3,345	1,755	571	93	-
1999.....	1,063	1,388	1,911	2,685	3,677	4,452	4,384	3,443	1,740	566	72	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Non-Hispanic Black, both sexes

2003.....	563	642	674	728	770	811	647	402	217	66	9	-
2002.....	526	613	716	769	843	850	657	444	215	67	12	-
2001.....	554	623	710	782	877	807	661	415	205	66	14	-
2000.....	535	552	703	767	869	865	607	396	166	82	20	1
1999.....	512	629	778	766	938	876	663	451	223	69	14	-

Non-Hispanic Black, male

2003.....	303	376	367	396	390	383	275	142	66	16	1	-
2002.....	295	335	404	387	426	405	270	161	65	16	1	-
2001.....	328	364	395	429	437	383	267	158	73	14	4	-
2000.....	283	308	425	387	440	440	247	144	52	19	1	1
1999.....	295	361	440	418	477	401	266	153	71	22	1	-

Non-Hispanic Black, female

2003.....	260	266	307	332	380	428	372	260	151	50	8	-
2002.....	231	278	312	382	417	445	387	283	150	51	11	-
2001.....	226	259	315	353	440	424	394	257	132	52	10	-
2000.....	252	244	278	380	429	425	360	252	114	63	19	-
1999.....	217	268	338	348	461	475	397	298	152	47	13	-

Origin not stated 2/, both sexes

2003.....	14	20	17	15	16	16	13	19	5	-	-	-
2002.....	6	10	19	22	18	25	19	20	9	3	-	-
2001.....	10	13	12	13	42	21	22	14	13	2	-	-
2000.....	12	11	14	18	28	37	20	16	5	1	-	-
1999.....	6	12	19	14	28	24	25	16	4	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Origin not stated 2/, male

2003.....	8	11	13	11	8	6	7	6	2	-	-	-
2002.....	4	4	15	14	10	13	13	5	4	-	-	-
2001.....	8	8	10	9	29	6	10	6	3	-	-	-
2000.....	3	9	7	12	16	19	10	6	-	-	-	-
1999.....	3	6	10	7	20	16	13	4	1	-	-	-

Origin not stated 2/, female

2003.....	6	9	4	4	8	10	6	13	3	-	-	-
2002.....	2	6	4	8	8	12	6	15	5	3	-	-
2001.....	2	5	2	4	13	15	12	8	10	2	-	-
2000.....	9	2	7	6	12	18	10	10	5	1	-	-
1999.....	3	6	9	7	8	8	12	12	3	-	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins, both sexes

2003.....	387	506	732	955	1,399	2,123	2,566	2,203	1,215	337	40	-
2002.....	387	561	632	928	1,420	2,137	2,460	2,052	1,162	366	33	-
2001.....	415	553	700	1,000	1,574	2,192	2,462	2,102	1,141	351	49	-
2000.....	397	513	711	1,007	1,539	2,142	2,503	2,103	1,143	314	40	1
1999.....	386	461	672	1,029	1,655	2,151	2,321	1,998	1,059	337	49	-

All origins, male

2003.....	218	279	406	524	810	1,114	1,265	955	411	94	10	-
2002.....	196	330	350	543	772	1,145	1,187	900	398	99	8	-
2001.....	218	279	394	563	876	1,199	1,204	882	404	102	10	-
2000.....	225	270	387	587	828	1,120	1,199	881	362	86	7	1
1999.....	220	255	401	590	932	1,119	1,099	839	367	82	15	-

All origins, female

2003.....	169	227	326	431	589	1,009	1,301	1,248	804	243	30	-
2002.....	191	231	282	385	648	992	1,273	1,152	764	267	25	-
2001.....	197	274	306	437	698	993	1,258	1,220	737	249	39	-
2000.....	172	243	324	420	711	1,022	1,304	1,222	781	228	33	-
1999.....	166	206	271	439	723	1,032	1,222	1,159	692	255	34	-

Hispanic, both sexes

2003.....	19	27	37	49	57	88	97	55	30	8	-	-
2002.....	24	20	40	41	64	65	83	51	31	5	1	-
2001.....	28	25	42	47	53	67	61	44	34	11	3	-
2000.....	23	23	30	40	58	67	56	63	29	12	2	-
1999.....	24	31	29	39	70	57	54	38	26	9	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Hispanic, male

2003.....	13	17	22	25	29	43	45	20	10	4	-	-
2002.....	10	14	21	23	31	31	46	24	9	-	1	-
2001.....	16	10	25	21	29	29	38	18	15	3	1	-
2000.....	9	10	15	16	30	36	18	26	12	5	-	-
1999.....	17	14	16	13	34	24	23	14	6	2	1	-

Hispanic, female

2003.....	6	10	15	24	28	45	52	35	20	4	-	-
2002.....	14	6	19	18	33	34	37	27	22	5	-	-
2001.....	12	15	17	26	24	38	23	26	19	8	2	-
2000.....	14	13	15	24	28	31	38	37	17	7	2	-
1999.....	7	17	13	26	36	33	31	24	20	7	-	-

Non-Hispanic 1/, both sexes

2003.....	368	479	691	904	1,337	2,032	2,467	2,144	1,185	329	40	-
2002.....	362	539	592	887	1,354	2,065	2,371	1,999	1,128	361	32	-
2001.....	385	527	655	951	1,517	2,120	2,390	2,055	1,103	340	46	-
2000.....	374	487	680	966	1,479	2,072	2,439	2,033	1,109	302	38	1
1999.....	359	428	642	988	1,584	2,088	2,261	1,957	1,030	327	48	-

Non-Hispanic 1/, male

2003.....	205	262	383	498	777	1,068	1,218	933	401	90	10	-
2002.....	186	316	329	520	740	1,107	1,138	874	388	99	7	-
2001.....	201	269	367	540	845	1,166	1,162	863	388	99	9	-
2000.....	216	257	372	570	797	1,081	1,177	851	349	81	7	1
1999.....	201	239	384	576	898	1,093	1,075	825	360	79	14	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic 1/, female

2003.....	163	217	308	406	560	964	1,249	1,211	784	239	30	-
2002.....	176	223	263	367	614	958	1,233	1,125	740	262	25	-
2001.....	184	258	288	411	672	954	1,228	1,192	715	241	37	-
2000.....	158	230	308	396	682	991	1,262	1,182	760	221	31	-
1999.....	158	189	258	412	686	995	1,186	1,132	670	248	34	-

Non-Hispanic White, both sexes

2003.....	299	397	598	770	1,200	1,870	2,277	2,006	1,098	303	30	-
2002.....	285	449	513	772	1,213	1,890	2,173	1,869	1,043	319	27	-
2001.....	326	451	555	837	1,348	1,945	2,205	1,897	1,015	303	36	-
2000.....	310	423	581	847	1,297	1,894	2,277	1,892	1,033	279	33	-
1999.....	295	357	542	846	1,430	1,899	2,102	1,829	967	294	43	-

Non-Hispanic White, male

2003.....	169	213	331	438	704	994	1,131	883	370	82	8	-
2002.....	152	261	286	457	678	1,023	1,046	824	357	85	7	-
2001.....	177	234	313	483	767	1,078	1,074	796	361	91	7	-
2000.....	183	229	310	500	702	994	1,114	789	328	77	7	-
1999.....	169	200	336	503	823	1,002	1,011	778	345	73	13	-

Non-Hispanic White, female

2003.....	130	184	267	332	496	876	1,146	1,123	728	221	22	-
2002.....	133	188	227	315	535	867	1,127	1,045	686	234	20	-
2001.....	149	217	242	354	581	867	1,131	1,101	654	212	29	-
2000.....	127	194	271	347	595	900	1,163	1,103	705	202	26	-
1999.....	126	157	206	343	607	897	1,091	1,051	622	221	30	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Non-Hispanic Black, both sexes

2003.....	59	66	77	107	115	129	155	98	70	22	8	-
2002.....	63	77	67	99	112	141	159	108	67	34	4	-
2001.....	54	63	84	99	137	141	155	116	75	33	9	-
2000.....	58	55	78	96	149	149	133	116	62	20	4	1
1999.....	54	64	84	124	132	162	133	108	52	30	4	-

Non-Hispanic Black, male

2003.....	29	39	43	50	64	57	70	34	23	6	1	-
2002.....	27	50	33	50	48	64	70	41	22	11	-	-
2001.....	22	28	45	51	68	68	73	45	22	7	2	-
2000.....	31	25	49	60	70	72	46	44	18	3	-	1
1999.....	25	36	39	65	63	75	49	38	10	6	-	-

Non-Hispanic Black, female

2003.....	30	27	34	57	51	72	85	64	47	16	7	-
2002.....	36	27	34	49	64	77	89	67	45	23	4	-
2001.....	32	35	39	48	69	73	82	71	53	26	7	-
2000.....	27	30	29	36	79	77	87	72	44	17	4	-
1999.....	29	28	45	59	69	87	84	70	42	24	4	-

Origin not stated 2/, both sexes

2003.....	-	-	4	2	5	3	2	4	-	-	-	-
2002.....	1	2	-	-	2	7	6	2	3	-	-	-
2001.....	2	1	3	2	4	5	11	3	4	-	-	-
2000.....	-	3	1	1	2	3	8	7	5	-	-	-
1999.....	3	2	1	2	1	6	6	3	3	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Origin not stated 2/, male

2003.....	-	-	1	1	4	3	2	2	-	-	-	-
2002.....	-	-	-	-	1	7	3	2	1	-	-	-
2001.....	1	-	2	2	2	4	4	1	1	-	-	-
2000.....	-	3	-	1	1	3	4	4	1	-	-	-
1999.....	2	2	1	1	-	2	1	-	1	1	-	-

Origin not stated 2/, female

2003.....	-	-	3	1	1	-	-	2	-	-	-	-
2002.....	1	2	-	-	1	-	3	-	2	-	-	-
2001.....	1	1	1	-	2	1	7	2	3	-	-	-
2000.....	-	-	1	-	1	-	4	3	4	-	-	-
1999.....	1	-	-	1	1	4	5	3	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Anemias (D50-D64)

##### All origins, both sexes

2003.....	104	121	155	184	300	447	689	784	721	342	72	-
2002.....	119	121	143	184	324	474	687	787	665	322	78	-
2001.....	120	119	130	186	306	525	701	808	659	294	90	-
2000.....	117	127	133	210	305	482	689	790	638	290	78	-
1999.....	98	105	126	175	311	497	667	775	647	316	82	-

##### All origins, male

2003.....	48	67	89	104	147	203	274	270	196	75	8	-
2002.....	52	57	74	77	167	204	250	256	182	81	13	-
2001.....	58	53	58	91	146	220	276	276	177	52	15	-
2000.....	56	66	64	95	141	198	267	237	183	58	14	-
1999.....	48	43	62	92	134	217	267	261	172	73	12	-

##### All origins, female

2003.....	56	54	66	80	153	244	415	514	525	267	64	-
2002.....	67	64	69	107	157	270	437	531	483	241	65	-
2001.....	62	66	72	95	160	305	425	532	482	242	75	-
2000.....	61	61	69	115	164	284	422	553	455	232	64	-
1999.....	50	62	64	83	177	280	400	514	475	243	70	-

##### Hispanic, both sexes

2003.....	5	12	6	12	13	18	30	23	22	9	2	-
2002.....	4	3	11	14	23	18	26	21	26	14	1	-
2001.....	5	11	6	14	11	24	19	23	15	6	2	-
2000.....	5	6	7	9	15	13	23	11	12	3	-	-
1999.....	3	5	5	7	20	16	19	21	17	11	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Anemias (D50-D64)

## Hispanic, male

2003.....	1	8	3	6	6	8	11	12	4	3	1	-
2002.....	2	3	5	6	11	9	9	11	14	7	-	-
2001.....	2	4	3	9	3	12	9	7	2	1	1	-
2000.....	3	1	3	4	8	4	11	5	4	-	-	-
1999.....	-	-	3	4	13	6	11	7	7	5	-	-

## Hispanic, female

2003.....	4	4	3	6	7	10	19	11	18	6	1	-
2002.....	2	-	6	8	12	9	17	10	12	7	1	-
2001.....	3	7	3	5	8	12	10	16	13	5	1	-
2000.....	2	5	4	5	7	9	12	6	8	3	-	-
1999.....	3	5	2	3	7	10	8	14	10	6	1	-

## Non-Hispanic 1/, both sexes

2003.....	98	109	149	172	285	428	659	760	698	333	70	-
2002.....	115	118	132	169	301	455	659	765	639	307	77	-
2001.....	114	107	124	172	294	500	677	783	644	288	87	-
2000.....	110	121	125	201	288	468	664	779	625	287	77	-
1999.....	95	100	120	165	290	478	647	751	630	302	81	-

## Non-Hispanic 1/, male

2003.....	46	59	86	98	141	195	263	258	192	72	7	-
2002.....	50	54	69	71	156	195	240	245	168	74	13	-
2001.....	56	49	55	82	143	208	266	269	175	51	14	-
2000.....	53	65	61	91	131	193	255	232	179	58	14	-
1999.....	48	43	58	87	120	209	256	254	165	66	12	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Anemias (D50-D64)

##### Non-Hispanic 1/, female

2003.....	52	50	63	74	144	233	396	502	506	261	63	-
2002.....	65	64	63	98	145	260	419	520	471	233	64	-
2001.....	58	58	69	90	151	292	411	514	469	237	73	-
2000.....	57	56	64	110	157	275	409	547	446	229	63	-
1999.....	47	57	62	78	170	269	391	497	465	236	69	-

##### Non-Hispanic White, both sexes

2003.....	40	44	91	124	224	361	578	694	616	297	63	-
2002.....	48	68	79	126	235	368	576	686	559	270	72	-
2001.....	56	57	93	132	236	422	591	703	584	260	66	-
2000.....	49	78	77	151	233	401	593	707	553	258	67	-
1999.....	38	41	71	126	234	410	585	668	566	266	67	-

##### Non-Hispanic White, male

2003.....	21	24	50	73	105	167	238	237	170	61	7	-
2002.....	25	33	42	55	117	164	207	220	143	61	12	-
2001.....	27	27	41	65	110	181	238	252	157	45	10	-
2000.....	24	44	39	77	108	165	227	207	152	52	13	-
1999.....	20	16	29	70	93	183	233	223	153	61	12	-

##### Non-Hispanic White, female

2003.....	19	20	41	51	119	194	340	457	446	236	56	-
2002.....	23	35	37	71	118	204	369	466	416	209	60	-
2001.....	29	30	52	67	126	241	353	451	427	215	56	-
2000.....	25	34	38	74	125	236	366	500	401	206	54	-
1999.....	18	25	42	56	141	227	352	445	413	205	55	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Anemias (D50-D64)

##### Non-Hispanic Black, both sexes

2003.....	57	64	52	40	54	63	76	59	75	30	5	-
2002.....	63	47	47	41	59	80	73	70	68	33	4	-
2001.....	57	46	29	36	54	71	78	66	50	24	18	-
2000.....	59	40	43	43	46	58	63	64	61	28	10	-
1999.....	52	55	45	33	48	63	56	67	57	28	12	-

##### Non-Hispanic Black, male

2003.....	25	35	32	22	33	28	25	19	19	9	-	-
2002.....	24	20	26	16	35	28	30	21	22	9	1	-
2001.....	29	21	14	15	31	26	25	15	13	6	3	-
2000.....	29	18	19	13	18	23	22	20	22	5	1	-
1999.....	26	25	28	14	23	22	19	25	9	3	-	-

##### Non-Hispanic Black, female

2003.....	32	29	20	18	21	35	51	40	56	21	5	-
2002.....	39	27	21	25	24	52	43	49	46	24	3	-
2001.....	28	25	15	21	23	45	53	51	37	18	15	-
2000.....	30	22	24	30	28	35	41	44	39	23	9	-
1999.....	26	30	17	19	25	41	37	42	48	25	12	-

##### Origin not stated 2/, both sexes

2003.....	1	-	-	-	2	1	-	1	1	-	-	-
2002.....	-	-	-	1	-	1	2	1	-	1	-	-
2001.....	1	1	-	-	1	1	5	2	-	-	1	-
2000.....	2	-	1	-	2	1	2	-	1	-	1	-
1999.....	-	-	1	3	1	3	1	3	-	3	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Anemias (D50-D64)

##### Origin not stated 2/, male

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	2	1	1	-	-	-	-	-
1999.....	-	-	1	1	1	2	-	-	-	2	-	-

##### Origin not stated 2/, female

2003.....	-	-	-	-	2	1	-	1	1	-	-	-
2002.....	-	-	-	1	-	1	1	1	-	1	-	-
2001.....	1	1	-	-	1	1	4	2	-	-	1	-
2000.....	2	-	1	-	-	-	1	-	1	-	1	-
1999.....	-	-	-	2	-	1	1	3	-	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Diabetes mellitus (E10-E14)

## All origins, both sexes

2003.....	3,413	4,806	5,925	7,310	9,346	11,574	11,725	8,758	4,572	1,424	210	3
2002.....	3,390	4,510	5,512	7,249	9,460	11,703	11,579	8,797	4,375	1,357	195	1
2001.....	3,280	4,188	5,382	7,195	9,536	11,546	11,259	8,366	4,259	1,350	196	1
2000.....	3,037	3,992	5,194	7,054	9,620	11,607	10,577	8,002	4,104	1,250	200	-
1999.....	2,793	3,865	5,232	7,215	9,693	11,269	10,488	7,893	3,943	1,165	177	3

## All origins, male

2003.....	1,987	2,715	3,329	3,929	4,758	5,547	5,132	3,260	1,318	306	39	3
2002.....	1,977	2,616	3,067	3,742	4,675	5,402	4,893	3,248	1,254	314	28	1
2001.....	1,920	2,362	2,879	3,734	4,703	5,244	4,674	2,966	1,188	269	34	1
2000.....	1,697	2,173	2,770	3,556	4,670	5,288	4,450	2,789	1,165	260	20	-
1999.....	1,602	2,120	2,736	3,630	4,649	5,190	4,349	2,816	1,071	210	28	2

## All origins, female

2003.....	1,426	2,091	2,596	3,381	4,588	6,027	6,593	5,498	3,254	1,118	171	-
2002.....	1,413	1,894	2,445	3,507	4,785	6,301	6,686	5,549	3,121	1,043	167	-
2001.....	1,360	1,826	2,503	3,461	4,833	6,302	6,585	5,400	3,071	1,081	162	-
2000.....	1,340	1,819	2,424	3,498	4,950	6,319	6,127	5,213	2,939	990	180	-
1999.....	1,191	1,745	2,496	3,585	5,044	6,079	6,139	5,077	2,872	955	149	1

## Hispanic, both sexes

2003.....	374	492	629	719	892	945	798	471	267	83	5	1
2002.....	322	432	597	711	880	964	748	457	249	72	12	-
2001.....	278	404	578	764	840	899	649	477	212	88	6	-
2000.....	265	436	514	684	908	817	593	412	213	66	10	-
1999.....	244	368	559	721	830	782	587	459	179	60	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Diabetes mellitus (E10-E14)

## Hispanic, male

2003.....	218	264	354	359	434	419	356	167	84	24	2	1
2002.....	198	236	315	327	425	442	306	162	71	16	5	-
2001.....	172	217	293	374	384	385	253	146	56	19	1	-
2000.....	142	241	279	338	424	362	247	146	60	17	-	-
1999.....	147	202	278	331	378	336	226	160	46	11	-	-

## Hispanic, female

2003.....	156	228	275	360	458	526	442	304	183	59	3	-
2002.....	124	196	282	384	455	522	442	295	178	56	7	-
2001.....	106	187	285	390	456	514	396	331	156	69	5	-
2000.....	123	195	235	346	484	455	346	266	153	49	10	-
1999.....	97	166	281	390	452	446	361	299	133	49	6	-

## Non-Hispanic 1/, both sexes

2003.....	3,026	4,301	5,282	6,575	8,439	10,600	10,898	8,270	4,294	1,338	203	2
2002.....	3,059	4,065	4,893	6,513	8,557	10,698	10,801	8,320	4,121	1,277	183	1
2001.....	2,985	3,769	4,793	6,404	8,662	10,616	10,582	7,869	4,039	1,261	189	1
2000.....	2,760	3,545	4,662	6,349	8,682	10,755	9,962	7,557	3,886	1,181	190	-
1999.....	2,536	3,484	4,654	6,477	8,841	10,455	9,870	7,414	3,752	1,101	169	2

## Non-Hispanic 1/, male

2003.....	1,764	2,442	2,966	3,560	4,316	5,112	4,763	3,086	1,230	282	37	2
2002.....	1,774	2,374	2,738	3,400	4,239	4,941	4,570	3,075	1,183	295	23	1
2001.....	1,740	2,136	2,579	3,345	4,298	4,844	4,411	2,812	1,131	250	32	1
2000.....	1,549	1,924	2,484	3,206	4,230	4,906	4,192	2,634	1,103	242	20	-
1999.....	1,443	1,911	2,447	3,288	4,260	4,842	4,113	2,648	1,021	197	28	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Diabetes mellitus (E10-E14)

## Non-Hispanic 1/, female

2003.....	1,262	1,859	2,316	3,015	4,123	5,488	6,135	5,184	3,064	1,056	166	-
2002.....	1,285	1,691	2,155	3,113	4,318	5,757	6,231	5,245	2,938	982	160	-
2001.....	1,245	1,633	2,214	3,059	4,364	5,772	6,171	5,057	2,908	1,011	157	-
2000.....	1,211	1,621	2,178	3,143	4,452	5,849	5,770	4,923	2,783	939	170	-
1999.....	1,093	1,573	2,207	3,189	4,581	5,613	5,757	4,766	2,731	904	141	1

## Non-Hispanic White, both sexes

2003.....	2,044	3,064	3,809	4,843	6,473	8,535	9,066	7,035	3,654	1,124	155	2
2002.....	2,096	2,896	3,507	4,717	6,620	8,632	9,060	7,066	3,536	1,064	135	-
2001.....	2,082	2,664	3,454	4,736	6,728	8,585	8,922	6,742	3,467	1,057	136	1
2000.....	1,903	2,526	3,397	4,714	6,761	8,725	8,315	6,472	3,309	998	138	-
1999.....	1,743	2,382	3,292	4,820	6,844	8,523	8,298	6,280	3,190	934	129	-

## Non-Hispanic White, male

2003.....	1,254	1,822	2,219	2,741	3,465	4,303	4,114	2,720	1,071	252	33	2
2002.....	1,262	1,767	2,048	2,595	3,446	4,179	3,972	2,720	1,071	255	18	-
2001.....	1,253	1,585	1,939	2,607	3,535	4,063	3,811	2,515	990	210	25	1
2000.....	1,127	1,417	1,881	2,510	3,457	4,135	3,661	2,320	962	210	16	-
1999.....	1,041	1,379	1,821	2,568	3,498	4,116	3,606	2,310	876	166	22	-

## Non-Hispanic White, female

2003.....	790	1,242	1,590	2,102	3,008	4,232	4,952	4,315	2,583	872	122	-
2002.....	834	1,129	1,459	2,122	3,174	4,453	5,088	4,346	2,465	809	117	-
2001.....	829	1,079	1,515	2,129	3,193	4,522	5,111	4,227	2,477	847	111	-
2000.....	776	1,109	1,516	2,204	3,304	4,590	4,654	4,152	2,347	788	122	-
1999.....	702	1,003	1,471	2,252	3,346	4,407	4,692	3,970	2,314	768	107	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Diabetes mellitus (E10-E14)

## Non-Hispanic Black, both sexes

2003.....	853	1,092	1,255	1,484	1,661	1,743	1,522	1,024	537	183	45	-
2002.....	850	1,003	1,201	1,513	1,639	1,728	1,485	1,045	516	195	45	1
2001.....	799	974	1,159	1,440	1,670	1,735	1,397	975	507	179	49	-
2000.....	757	889	1,106	1,420	1,663	1,769	1,424	953	505	160	49	-
1999.....	706	948	1,188	1,438	1,682	1,635	1,360	970	494	141	37	2

## Non-Hispanic Black, male

2003.....	431	545	624	691	701	667	521	285	119	22	3	-
2002.....	445	512	592	658	662	634	501	259	90	33	5	1
2001.....	416	483	548	629	649	655	495	236	120	29	7	-
2000.....	371	434	518	583	665	652	448	258	108	21	2	-
1999.....	351	457	529	630	632	588	428	278	121	22	5	1

## Non-Hispanic Black, female

2003.....	422	547	631	793	960	1,076	1,001	739	418	161	42	-
2002.....	405	491	609	855	977	1,094	984	786	426	162	40	-
2001.....	383	491	611	811	1,021	1,080	902	739	387	150	42	-
2000.....	386	455	588	837	998	1,117	976	695	397	139	47	-
1999.....	355	491	659	808	1,050	1,047	932	692	373	119	32	1

## Origin not stated 2/, both sexes

2003.....	13	13	14	16	15	29	29	17	11	3	2	-
2002.....	9	13	22	25	23	41	30	20	5	8	-	-
2001.....	17	15	11	27	34	31	28	20	8	1	1	-
2000.....	12	11	18	21	30	35	22	33	5	3	-	-
1999.....	13	13	19	17	22	32	31	20	12	4	2	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diabetes mellitus (E10-E14)

Origin not stated 2/, male

2003.....	5	9	9	10	8	16	13	7	4	-	-	-
2002.....	5	6	14	15	11	19	17	11	-	3	-	-
2001.....	8	9	7	15	21	15	10	8	1	-	1	-
2000.....	6	8	7	12	16	20	11	9	2	1	-	-
1999.....	12	7	11	11	11	12	10	8	4	2	-	1

Origin not stated 2/, female

2003.....	8	4	5	6	7	13	16	10	7	3	2	-
2002.....	4	7	8	10	12	22	13	9	5	5	-	-
2001.....	9	6	4	12	13	16	18	12	7	1	-	-
2000.....	6	3	11	9	14	15	11	24	3	2	-	-
1999.....	1	6	8	6	11	20	21	12	8	2	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Nutritional deficiencies (E40-E64)

##### All origins, both sexes

2003.....	61	77	91	125	215	428	564	727	570	301	74	-
2002.....	63	77	91	150	232	452	680	834	667	312	93	-
2001.....	62	64	89	131	243	439	661	798	684	324	83	-
2000.....	53	70	103	192	268	505	780	947	781	302	90	-
1999.....	61	76	94	149	280	472	776	944	822	380	94	-

##### All origins, male

2003.....	29	38	41	63	103	175	225	243	139	49	11	-
2002.....	28	43	50	70	116	193	241	283	169	49	13	-
2001.....	29	35	46	63	111	172	254	228	171	55	3	-
2000.....	21	45	52	87	134	207	264	289	182	52	13	-
1999.....	30	42	56	72	149	205	289	301	196	71	9	-

##### All origins, female

2003.....	32	39	50	62	112	253	339	484	431	252	63	-
2002.....	35	34	41	80	116	259	439	551	498	263	80	-
2001.....	33	29	43	68	132	267	407	570	513	269	80	-
2000.....	32	25	51	105	134	298	516	658	599	250	77	-
1999.....	31	34	38	77	131	267	487	643	626	309	85	-

##### Hispanic, both sexes

2003.....	1	4	3	3	16	17	19	24	15	13	3	-
2002.....	3	1	3	7	14	25	24	22	19	8	4	-
2001.....	2	2	3	5	7	22	23	25	25	12	-	-
2000.....	3	3	4	14	10	18	25	20	29	8	3	-
1999.....	5	5	2	4	13	16	15	34	20	15	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Nutritional deficiencies (E40-E64)

##### Hispanic, male

2003.....	1	4	2	2	8	6	7	10	4	3	1	-
2002.....	1	1	1	5	8	9	10	8	7	3	-	-
2001.....	1	2	2	1	1	11	9	6	10	7	-	-
2000.....	2	3	-	4	9	8	9	7	9	1	1	-
1999.....	4	1	1	1	7	10	3	10	7	6	1	-

##### Hispanic, female

2003.....	-	-	1	1	8	11	12	14	11	10	2	-
2002.....	2	-	2	2	6	16	14	14	12	5	4	-
2001.....	1	-	1	4	6	11	14	19	15	5	-	-
2000.....	1	-	4	10	1	10	16	13	20	7	2	-
1999.....	1	4	1	3	6	6	12	24	13	9	3	-

##### Non-Hispanic 1/, both sexes

2003.....	59	73	88	121	199	410	542	701	555	288	71	-
2002.....	60	76	88	143	215	422	655	809	645	303	89	-
2001.....	60	62	86	126	236	416	638	772	657	312	83	-
2000.....	50	67	99	178	257	482	751	926	749	293	87	-
1999.....	56	70	92	145	266	455	760	907	800	365	90	-

##### Non-Hispanic 1/, male

2003.....	27	34	39	60	95	169	216	233	135	46	10	-
2002.....	27	42	49	65	108	181	231	274	162	45	13	-
2001.....	28	33	44	62	110	160	245	222	161	48	3	-
2000.....	19	42	52	83	125	195	253	281	173	51	12	-
1999.....	26	40	55	71	142	194	285	290	189	65	8	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

Non-Hispanic 1/, female

2003.....	32	39	49	61	104	241	326	468	420	242	61	-
2002.....	33	34	39	78	107	241	424	535	483	258	76	-
2001.....	32	29	42	64	126	256	393	550	496	264	80	-
2000.....	31	25	47	95	132	287	498	645	576	242	75	-
1999.....	30	30	37	74	124	261	475	617	611	300	82	-

Non-Hispanic White, both sexes

2003.....	37	56	66	97	160	351	460	621	483	245	52	-
2002.....	46	53	64	107	173	357	565	703	572	267	73	-
2001.....	37	43	70	93	198	342	536	690	578	274	76	-
2000.....	33	49	74	145	200	404	658	815	681	247	68	-
1999.....	43	40	70	111	196	375	643	797	731	312	76	-

Non-Hispanic White, male

2003.....	18	22	28	50	71	138	179	208	116	39	6	-
2002.....	19	26	31	50	87	157	191	242	147	39	10	-
2001.....	13	20	32	42	89	133	207	197	135	41	2	-
2000.....	11	27	35	66	93	160	222	248	158	40	10	-
1999.....	19	19	42	51	99	154	238	254	169	49	6	-

Non-Hispanic White, female

2003.....	19	34	38	47	89	213	281	413	367	206	46	-
2002.....	27	27	33	57	86	200	374	461	425	228	63	-
2001.....	24	23	38	51	109	209	329	493	443	233	74	-
2000.....	22	22	39	79	107	244	436	567	523	207	58	-
1999.....	24	21	28	60	97	221	405	543	562	263	70	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Nutritional deficiencies (E40-E64)

##### Non-Hispanic Black, both sexes

2003.....	20	17	20	19	35	53	74	72	66	35	15	-
2002.....	13	23	23	33	37	61	78	88	66	32	13	-
2001.....	21	17	12	28	33	62	87	72	68	30	7	-
2000.....	16	18	23	29	49	67	80	96	60	40	16	-
1999.....	12	26	21	29	61	69	106	96	63	46	14	-

##### Non-Hispanic Black, male

2003.....	9	12	10	9	22	30	32	22	16	5	3	-
2002.....	8	16	17	14	17	21	35	26	12	2	2	-
2001.....	14	12	9	18	20	25	30	24	21	5	1	-
2000.....	8	15	15	15	29	31	27	31	12	11	1	-
1999.....	7	18	12	17	38	34	40	30	19	13	2	-

##### Non-Hispanic Black, female

2003.....	11	5	10	10	13	23	42	50	50	30	12	-
2002.....	5	7	6	19	20	40	43	62	54	30	11	-
2001.....	7	5	3	10	13	37	57	48	47	25	6	-
2000.....	8	3	8	14	20	36	53	65	48	29	15	-
1999.....	5	8	9	12	23	35	66	66	44	33	12	-

##### Origin not stated 2/, both sexes

2003.....	1	-	-	1	-	1	3	2	-	-	-	-
2002.....	-	-	-	-	3	5	1	3	3	1	-	-
2001.....	-	-	-	-	-	1	-	1	2	-	-	-
2000.....	-	-	-	-	1	5	4	1	3	1	-	-
1999.....	-	1	-	-	1	1	1	3	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nutritional deficiencies (E40-E64)

Origin not stated 2/, male

2003.....	1	-	-	1	-	-	2	-	-	-	-	-
2002.....	-	-	-	-	-	3	-	1	-	1	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	4	2	1	-	-	-	-
1999.....	-	1	-	-	-	1	1	1	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	1	1	2	-	-	-	-
2002.....	-	-	-	-	3	2	1	2	3	-	-	-
2001.....	-	-	-	-	-	-	-	1	2	-	-	-
2000.....	-	-	-	-	1	1	2	-	3	1	-	-
1999.....	-	-	-	-	1	-	-	2	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Malnutrition (E40-E46)

##### All origins, both sexes

2003.....	58	74	86	118	202	413	539	684	537	284	68	-
2002.....	55	73	83	136	218	419	648	778	616	283	87	-
2001.....	57	62	84	123	224	406	619	756	643	302	71	-
2000.....	46	61	97	178	245	478	721	888	712	280	84	-
1999.....	57	66	88	141	266	439	720	883	755	353	86	-

##### All origins, male

2003.....	28	37	38	59	96	166	213	222	130	46	10	-
2002.....	24	39	45	64	110	176	230	267	163	45	13	-
2001.....	25	33	42	57	100	156	244	220	164	50	2	-
2000.....	18	39	47	79	128	194	240	275	165	46	13	-
1999.....	27	38	54	68	139	186	266	282	175	65	8	-

##### All origins, female

2003.....	30	37	48	59	106	247	326	462	407	238	58	-
2002.....	31	34	38	72	108	243	418	511	453	238	74	-
2001.....	32	29	42	66	124	250	375	536	479	252	69	-
2000.....	28	22	50	99	117	284	481	613	547	234	71	-
1999.....	30	28	34	73	127	253	454	601	580	288	78	-

##### Hispanic, both sexes

2003.....	1	4	3	3	16	14	16	21	14	12	3	-
2002.....	3	1	3	7	14	25	24	21	19	8	3	-
2001.....	2	2	3	5	7	22	22	25	25	11	-	-
2000.....	2	3	4	12	10	16	23	20	27	8	3	-
1999.....	5	5	2	4	13	15	15	32	17	15	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Malnutrition (E40-E46)

##### Hispanic, male

2003.....	1	4	2	2	8	5	5	8	4	3	1	-
2002.....	1	1	1	5	8	9	10	7	7	3	-	-
2001.....	1	2	2	1	1	11	9	6	10	6	-	-
2000.....	1	3	-	3	9	7	9	7	7	1	1	-
1999.....	4	1	1	1	7	9	3	9	5	6	1	-

##### Hispanic, female

2003.....	-	-	1	1	8	9	11	13	10	9	2	-
2002.....	2	-	2	2	6	16	14	14	12	5	3	-
2001.....	1	-	1	4	6	11	13	19	15	5	-	-
2000.....	1	-	4	9	1	9	14	13	20	7	2	-
1999.....	1	4	1	3	6	6	12	23	12	9	3	-

##### Non-Hispanic 1/, both sexes

2003.....	56	70	83	114	186	398	520	662	523	272	65	-
2002.....	52	72	80	129	201	390	623	754	594	274	84	-
2001.....	55	60	81	118	217	383	597	730	616	291	71	-
2000.....	44	58	93	166	234	457	694	867	682	271	81	-
1999.....	52	60	86	137	252	423	704	848	736	338	82	-

##### Non-Hispanic 1/, male

2003.....	26	33	36	56	88	161	206	214	126	43	9	-
2002.....	23	38	44	59	102	165	220	259	156	41	13	-
2001.....	24	31	40	56	99	144	235	214	154	44	2	-
2000.....	17	36	47	76	119	183	229	267	158	45	12	-
1999.....	23	36	53	67	132	176	262	272	170	59	7	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Malnutrition (E40-E46)

##### Non-Hispanic 1/, female

2003.....	30	37	47	58	98	237	314	448	397	229	56	-
2002.....	29	34	36	70	99	225	403	495	438	233	71	-
2001.....	31	29	41	62	118	239	362	516	462	247	69	-
2000.....	27	22	46	90	115	274	465	600	524	226	69	-
1999.....	29	24	33	70	120	247	442	576	566	279	75	-

##### Non-Hispanic White, both sexes

2003.....	34	53	62	90	149	340	439	582	455	231	47	-
2002.....	39	50	58	98	164	327	537	653	525	240	69	-
2001.....	35	43	66	88	182	315	501	653	540	255	64	-
2000.....	29	42	70	134	178	386	607	760	621	227	64	-
1999.....	41	35	64	103	184	350	595	745	672	288	68	-

##### Non-Hispanic White, male

2003.....	17	21	26	46	65	131	170	189	108	37	6	-
2002.....	16	23	28	46	83	142	183	228	141	35	10	-
2001.....	12	20	29	39	80	118	198	189	129	38	1	-
2000.....	10	23	32	60	87	152	198	234	145	34	10	-
1999.....	17	18	40	47	91	142	217	237	150	43	5	-

##### Non-Hispanic White, female

2003.....	17	32	36	44	84	209	269	393	347	194	41	-
2002.....	23	27	30	52	81	185	354	425	384	205	59	-
2001.....	23	23	37	49	102	197	303	464	411	217	63	-
2000.....	19	19	38	74	91	234	409	526	476	193	54	-
1999.....	24	17	24	56	93	208	378	508	522	245	63	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Malnutrition (E40-E46)

##### Non-Hispanic Black, both sexes

2003.....	20	17	19	19	33	52	73	72	62	33	15	-
2002.....	12	22	22	28	32	59	75	84	63	30	12	-
2001.....	18	15	11	26	30	58	81	67	65	28	7	-
2000.....	15	16	21	28	49	62	74	94	56	38	14	-
1999.....	11	21	21	29	59	62	99	89	58	43	14	-

##### Non-Hispanic Black, male

2003.....	9	12	9	9	21	29	31	22	15	4	3	-
2002.....	7	15	16	12	15	20	32	25	12	2	2	-
2001.....	11	10	8	16	18	24	29	24	20	4	1	-
2000.....	7	13	13	14	29	29	27	31	10	11	1	-
1999.....	6	15	12	17	36	28	38	29	19	13	2	-

##### Non-Hispanic Black, female

2003.....	11	5	10	10	12	23	42	50	47	29	12	-
2002.....	5	7	6	16	17	39	43	59	51	28	10	-
2001.....	7	5	3	10	12	34	52	43	45	24	6	-
2000.....	8	3	8	14	20	33	47	63	46	27	13	-
1999.....	5	6	9	12	23	34	61	60	39	30	12	-

##### Origin not stated 2/, both sexes

2003.....	1	-	-	1	-	1	3	1	-	-	-	-
2002.....	-	-	-	-	3	4	1	3	3	1	-	-
2001.....	-	-	-	-	-	1	-	1	2	-	-	-
2000.....	-	-	-	-	1	5	4	1	3	1	-	-
1999.....	-	1	-	-	1	1	1	3	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Malnutrition (E40-E46)

Origin not stated 2/, male

2003.....	1	-	-	1	-	-	2	-	-	-	-	-
2002.....	-	-	-	-	-	2	-	1	-	1	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	4	2	1	-	-	-	-
1999.....	-	1	-	-	-	1	1	1	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	1	1	1	-	-	-	-
2002.....	-	-	-	-	3	2	1	2	3	-	-	-
2001.....	-	-	-	-	-	-	-	1	2	-	-	-
2000.....	-	-	-	-	1	1	2	-	3	1	-	-
1999.....	-	-	-	-	1	-	-	2	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

All origins, both sexes

2003.....	3	3	5	7	13	15	25	43	33	17	6	-
2002.....	8	4	8	14	14	33	32	56	51	29	6	-
2001.....	5	2	5	8	19	33	42	42	41	22	12	-
2000.....	7	9	6	14	23	27	59	59	69	22	6	-
1999.....	4	10	6	8	14	33	56	61	67	27	8	-

All origins, male

2003.....	1	1	3	4	7	9	12	21	9	3	1	-
2002.....	4	4	5	6	6	17	11	16	6	4	-	-
2001.....	4	2	4	6	11	16	10	8	7	5	1	-
2000.....	3	6	5	8	6	13	24	14	17	6	-	-
1999.....	3	4	2	4	10	19	23	19	21	6	1	-

All origins, female

2003.....	2	2	2	3	6	6	13	22	24	14	5	-
2002.....	4	-	3	8	8	16	21	40	45	25	6	-
2001.....	1	-	1	2	8	17	32	34	34	17	11	-
2000.....	4	3	1	6	17	14	35	45	52	16	6	-
1999.....	1	6	4	4	4	14	33	42	46	21	7	-

Hispanic, both sexes

2003.....	-	-	-	-	-	3	3	3	1	1	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	1	-
2001.....	-	-	-	-	-	-	1	-	-	1	-	-
2000.....	1	-	-	2	-	2	2	-	2	-	-	-
1999.....	-	-	-	-	-	1	-	2	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

Hispanic, male

2003.....	-	-	-	-	-	1	2	2	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	1	-	-
2000.....	1	-	-	1	-	1	-	-	2	-	-	-
1999.....	-	-	-	-	-	1	-	1	2	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	2	1	1	1	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	1	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	1	-	1	2	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	1	1	-	-	-

Non-Hispanic 1/, both sexes

2003.....	3	3	5	7	13	12	22	39	32	16	6	-
2002.....	8	4	8	14	14	32	32	55	51	29	5	-
2001.....	5	2	5	8	19	33	41	42	41	21	12	-
2000.....	6	9	6	12	23	25	57	59	67	22	6	-
1999.....	4	10	6	8	14	32	56	59	64	27	8	-

Non-Hispanic 1/, male

2003.....	1	1	3	4	7	8	10	19	9	3	1	-
2002.....	4	4	5	6	6	16	11	15	6	4	-	-
2001.....	4	2	4	6	11	16	10	8	7	4	1	-
2000.....	2	6	5	7	6	12	24	14	15	6	-	-
1999.....	3	4	2	4	10	18	23	18	19	6	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other nutritional deficiencies (E50-E64)

Non-Hispanic 1/, female

2003.....	2	2	2	3	6	4	12	20	23	13	5	-
2002.....	4	-	3	8	8	16	21	40	45	25	5	-
2001.....	1	-	1	2	8	17	31	34	34	17	11	-
2000.....	4	3	1	5	17	13	33	45	52	16	6	-
1999.....	1	6	4	4	4	14	33	41	45	21	7	-

Non-Hispanic White, both sexes

2003.....	3	3	4	7	11	11	21	39	28	14	5	-
2002.....	7	3	6	9	9	30	28	50	47	27	4	-
2001.....	2	-	4	5	16	27	35	37	38	19	12	-
2000.....	4	7	4	11	22	18	51	55	60	20	4	-
1999.....	2	5	6	8	12	25	48	52	59	24	8	-

Non-Hispanic White, male

2003.....	1	1	2	4	6	7	9	19	8	2	-	-
2002.....	3	3	3	4	4	15	8	14	6	4	-	-
2001.....	1	-	3	3	9	15	9	8	6	3	1	-
2000.....	1	4	3	6	6	8	24	14	13	6	-	-
1999.....	2	1	2	4	8	12	21	17	19	6	1	-

Non-Hispanic White, female

2003.....	2	2	2	3	5	4	12	20	20	12	5	-
2002.....	4	-	3	5	5	15	20	36	41	23	4	-
2001.....	1	-	1	2	7	12	26	29	32	16	11	-
2000.....	3	3	1	5	16	10	27	41	47	14	4	-
1999.....	-	4	4	4	4	13	27	35	40	18	7	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Meningitis (G00,G03)

##### All origins, both sexes

2003.....	58	55	53	57	39	55	35	33	18	1	-	1
2002.....	53	48	40	46	63	47	37	36	16	3	-	1
2001.....	50	58	55	46	48	46	50	25	12	4	-	-
2000.....	55	43	47	48	60	68	55	32	16	2	-	-
1999.....	46	41	49	49	66	65	57	40	20	2	-	-

##### All origins, male

2003.....	38	30	31	27	18	26	14	13	5	1	-	1
2002.....	30	29	19	26	36	21	20	11	8	-	-	1
2001.....	30	30	31	24	27	26	21	11	3	1	-	-
2000.....	31	26	28	26	34	41	21	13	5	1	-	-
1999.....	29	24	21	28	32	26	23	13	7	-	-	-

##### All origins, female

2003.....	20	25	22	30	21	29	21	20	13	-	-	-
2002.....	23	19	21	20	27	26	17	25	8	3	-	-
2001.....	20	28	24	22	21	20	29	14	9	3	-	-
2000.....	24	17	19	22	26	27	34	19	11	1	-	-
1999.....	17	17	28	21	34	39	34	27	13	2	-	-

##### Hispanic, both sexes

2003.....	5	4	7	4	1	2	-	1	2	-	-	-
2002.....	3	6	2	4	7	5	4	1	-	-	-	-
2001.....	1	4	3	2	4	3	5	1	1	-	-	-
2000.....	5	2	3	2	3	3	1	-	-	-	-	-
1999.....	4	3	4	6	3	4	-	4	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Meningitis (G00,G03)

## Hispanic, male

2003.....	4	2	2	2	-	-	-	1	-	-	-	-
2002.....	2	4	-	1	5	1	2	1	-	-	-	-
2001.....	-	2	1	2	3	1	3	-	-	-	-	-
2000.....	4	1	2	1	-	3	1	-	-	-	-	-
1999.....	2	3	-	2	3	-	-	1	-	-	-	-

## Hispanic, female

2003.....	1	2	5	2	1	2	-	-	2	-	-	-
2002.....	1	2	2	3	2	4	2	-	-	-	-	-
2001.....	1	2	2	-	1	2	2	1	1	-	-	-
2000.....	1	1	1	1	3	-	-	-	-	-	-	-
1999.....	2	-	4	4	-	4	-	3	-	-	-	-

## Non-Hispanic 1/, both sexes

2003.....	52	51	46	53	38	53	35	32	16	1	-	-
2002.....	50	41	38	42	56	41	32	35	16	3	-	1
2001.....	49	54	52	44	44	43	45	22	11	4	-	-
2000.....	50	41	44	46	57	65	54	32	16	2	-	-
1999.....	42	37	45	43	63	61	57	36	20	2	-	-

## Non-Hispanic 1/, male

2003.....	33	28	29	25	18	26	14	12	5	1	-	-
2002.....	28	24	19	25	31	19	18	10	8	-	-	1
2001.....	30	28	30	22	24	25	18	11	3	1	-	-
2000.....	27	25	26	25	34	38	20	13	5	1	-	-
1999.....	27	20	21	26	29	26	23	12	7	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Meningitis (G00,G03)

Non-Hispanic 1/, female

2003.....	19	23	17	28	20	27	21	20	11	-	-	-
2002.....	22	17	19	17	25	22	14	25	8	3	-	-
2001.....	19	26	22	22	20	18	27	11	8	3	-	-
2000.....	23	16	18	21	23	27	34	19	11	1	-	-
1999.....	15	17	24	17	34	35	34	24	13	2	-	-

Non-Hispanic White, both sexes

2003.....	32	32	29	42	29	47	30	27	11	1	-	-
2002.....	29	26	23	34	43	31	27	34	12	3	-	1
2001.....	34	36	41	33	40	37	37	17	9	2	-	-
2000.....	31	27	33	30	44	53	47	24	14	2	-	-
1999.....	29	28	35	32	48	44	48	32	12	2	-	-

Non-Hispanic White, male

2003.....	19	20	19	21	14	25	12	10	4	1	-	-
2002.....	16	15	12	21	24	15	17	9	5	-	-	1
2001.....	21	19	23	15	22	20	15	7	2	-	-	-
2000.....	18	19	20	18	29	30	18	9	4	1	-	-
1999.....	17	16	17	19	22	17	21	11	4	-	-	-

Non-Hispanic White, female

2003.....	13	12	10	21	15	22	18	17	7	-	-	-
2002.....	13	11	11	13	19	16	10	25	7	3	-	-
2001.....	13	17	18	18	18	17	22	10	7	2	-	-
2000.....	13	8	13	12	15	23	29	15	10	1	-	-
1999.....	12	12	18	13	26	27	27	21	8	2	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

All origins, both sexes

2003.....	46	112	240	624	1,706	3,671	5,052	4,157	1,900	425	31	-
2002.....	39	95	223	626	1,600	3,368	4,771	3,959	1,781	440	32	-
2001.....	32	100	201	554	1,593	3,446	4,679	3,786	1,655	425	43	-
2000.....	33	82	188	626	1,482	3,322	4,330	3,567	1,627	369	27	1
1999.....	35	73	169	539	1,481	3,066	4,044	3,332	1,492	313	31	-

All origins, male

2003.....	23	65	154	402	1,115	2,300	2,891	2,210	846	143	14	-
2002.....	26	57	136	433	1,054	2,065	2,746	2,109	793	147	10	-
2001.....	25	69	145	362	1,032	2,104	2,733	2,012	759	134	10	-
2000.....	19	55	115	398	940	2,052	2,478	1,879	726	115	8	1
1999.....	25	44	117	338	965	1,940	2,367	1,699	639	110	6	-

All origins, female

2003.....	23	47	86	222	591	1,371	2,161	1,947	1,054	282	17	-
2002.....	13	38	87	193	546	1,303	2,025	1,850	988	293	22	-
2001.....	7	31	56	192	561	1,342	1,946	1,774	896	291	33	-
2000.....	14	27	73	228	542	1,270	1,852	1,688	901	254	19	-
1999.....	10	29	52	201	516	1,126	1,677	1,633	853	203	25	-

Hispanic, both sexes

2003.....	1	7	8	25	71	112	146	92	44	14	-	-
2002.....	2	8	13	39	63	104	119	96	33	7	2	-
2001.....	2	5	8	30	68	108	122	94	34	8	2	-
2000.....	2	4	11	34	50	89	102	81	42	5	2	-
1999.....	2	2	6	24	60	97	87	63	36	9	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

Hispanic, male

2003.....	-	4	5	20	47	64	88	58	20	6	-	-
2002.....	-	5	11	26	45	63	59	53	14	1	-	-
2001.....	1	1	6	19	37	63	78	50	14	3	-	-
2000.....	1	3	6	19	27	53	59	49	18	2	2	-
1999.....	2	1	4	19	33	60	53	31	14	4	-	-

Hispanic, female

2003.....	1	3	3	5	24	48	58	34	24	8	-	-
2002.....	2	3	2	13	18	41	60	43	19	6	2	-
2001.....	1	4	2	11	31	45	44	44	20	5	2	-
2000.....	1	1	5	15	23	36	43	32	24	3	-	-
1999.....	-	1	2	5	27	37	34	32	22	5	2	-

Non-Hispanic 1/, both sexes

2003.....	45	105	231	598	1,634	3,550	4,894	4,062	1,851	411	31	-
2002.....	37	87	209	586	1,534	3,257	4,642	3,856	1,745	432	30	-
2001.....	29	95	193	523	1,523	3,329	4,544	3,682	1,617	415	41	-
2000.....	31	78	177	591	1,429	3,223	4,219	3,480	1,581	364	25	1
1999.....	33	71	163	511	1,419	2,967	3,950	3,263	1,452	303	28	-

Non-Hispanic 1/, male

2003.....	23	61	148	381	1,067	2,230	2,795	2,151	824	137	14	-
2002.....	26	52	124	406	1,007	1,996	2,681	2,054	776	145	10	-
2001.....	23	68	139	342	993	2,034	2,647	1,957	743	130	10	-
2000.....	18	52	109	378	912	1,992	2,415	1,826	706	113	6	1
1999.....	23	43	113	318	930	1,878	2,311	1,667	623	106	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

Non-Hispanic 1/, female

2003.....	22	44	83	217	567	1,320	2,099	1,911	1,027	274	17	-
2002.....	11	35	85	180	527	1,261	1,961	1,802	969	287	20	-
2001.....	6	27	54	181	530	1,295	1,897	1,725	874	285	31	-
2000.....	13	26	68	213	517	1,231	1,804	1,654	875	251	19	-
1999.....	10	28	50	193	489	1,089	1,639	1,596	829	197	22	-

Non-Hispanic White, both sexes

2003.....	36	91	201	540	1,521	3,352	4,668	3,869	1,775	391	26	-
2002.....	36	74	195	530	1,429	3,084	4,461	3,695	1,672	408	28	-
2001.....	27	85	172	483	1,428	3,172	4,349	3,528	1,547	392	35	-
2000.....	27	71	165	550	1,345	3,070	4,036	3,332	1,515	345	24	1
1999.....	29	63	144	471	1,336	2,811	3,783	3,124	1,387	285	26	-

Non-Hispanic White, male

2003.....	19	53	127	347	996	2,110	2,674	2,038	785	127	12	-
2002.....	25	45	118	369	943	1,896	2,577	1,969	741	136	10	-
2001.....	21	61	129	319	933	1,950	2,536	1,881	709	123	9	-
2000.....	15	46	103	352	862	1,887	2,304	1,752	669	107	5	1
1999.....	22	40	102	294	888	1,780	2,225	1,593	590	98	6	-

Non-Hispanic White, female

2003.....	17	38	74	193	525	1,242	1,994	1,831	990	264	14	-
2002.....	11	29	77	161	486	1,188	1,884	1,726	931	272	18	-
2001.....	6	24	43	164	495	1,222	1,813	1,647	838	269	26	-
2000.....	12	25	62	198	483	1,183	1,732	1,580	846	238	19	-
1999.....	7	23	42	177	448	1,031	1,558	1,531	797	187	20	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

Non-Hispanic Black, both sexes

2003.....	7	11	22	39	81	134	148	112	53	11	4	-
2002.....	1	8	10	41	70	120	123	103	49	18	1	-
2001.....	2	5	15	25	66	109	129	113	54	18	4	-
2000.....	4	5	9	28	65	116	130	102	47	15	1	-
1999.....	4	7	15	32	63	109	112	93	51	14	2	-

Non-Hispanic Black, male

2003.....	3	7	14	22	51	81	77	62	25	5	1	-
2002.....	1	3	6	29	47	66	72	56	19	5	-	-
2001.....	2	4	9	15	44	55	72	53	29	4	-	-
2000.....	3	4	5	19	41	78	76	44	24	6	1	-
1999.....	1	3	10	22	36	67	56	47	22	5	-	-

Non-Hispanic Black, female

2003.....	4	4	8	17	30	53	71	50	28	6	3	-
2002.....	-	5	4	12	23	54	51	47	30	13	1	-
2001.....	-	1	6	10	22	54	57	60	25	14	4	-
2000.....	1	1	4	9	24	38	54	58	23	9	-	-
1999.....	3	4	5	10	27	42	56	46	29	9	2	-

Origin not stated 2/, both sexes

2003.....	-	-	1	1	1	9	12	3	5	-	-	-
2002.....	-	-	1	1	3	7	10	7	3	1	-	-
2001.....	1	-	-	1	2	9	13	10	4	2	-	-
2000.....	-	-	-	1	3	10	9	6	4	-	-	-
1999.....	-	-	-	4	2	2	7	6	4	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Parkinson's disease (G20-G21)

Origin not stated 2/, male

2003.....	-	-	1	1	1	6	8	1	2	-	-	-
2002.....	-	-	1	1	2	6	6	2	3	1	-	-
2001.....	1	-	-	1	2	7	8	5	2	1	-	-
2000.....	-	-	-	1	1	7	4	4	2	-	-	-
1999.....	-	-	-	1	2	2	3	1	2	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	3	4	2	3	-	-	-
2002.....	-	-	-	-	1	1	4	5	-	-	-	-
2001.....	-	-	-	-	-	2	5	5	2	1	-	-
2000.....	-	-	-	-	2	3	5	2	2	-	-	-
1999.....	-	-	-	3	-	-	4	5	2	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Alzheimer's disease (G30)

##### All origins, both sexes

2003.....	57	152	402	979	2,857	7,348	13,809	17,368	13,774	5,704	975	-
2002.....	40	135	375	939	2,663	7,078	13,057	16,267	12,267	5,140	878	-
2001.....	56	165	364	858	2,564	6,568	11,974	14,843	11,168	4,474	796	-
2000.....	47	142	349	868	2,563	6,319	10,934	13,337	10,257	4,014	701	2
1999.....	40	125	323	830	2,374	5,769	10,067	11,862	9,041	3,473	604	-

##### All origins, male

2003.....	25	76	167	456	1,228	2,855	4,751	4,916	2,933	824	91	-
2002.....	22	55	173	408	1,139	2,793	4,434	4,569	2,570	732	79	-
2001.....	27	61	167	397	1,075	2,616	4,179	4,225	2,312	608	79	-
2000.....	24	60	150	420	1,134	2,433	3,795	3,687	2,101	564	54	1
1999.....	18	68	154	389	1,048	2,332	3,572	3,382	1,861	505	48	-

##### All origins, female

2003.....	32	76	235	523	1,629	4,493	9,058	12,452	10,841	4,880	884	-
2002.....	18	80	202	531	1,524	4,285	8,623	11,698	9,697	4,408	799	-
2001.....	29	104	197	461	1,489	3,952	7,795	10,618	8,856	3,866	717	-
2000.....	23	82	199	448	1,429	3,886	7,139	9,650	8,156	3,450	647	1
1999.....	22	57	169	441	1,326	3,437	6,495	8,480	7,180	2,968	556	-

##### Hispanic, both sexes

2003.....	10	5	22	44	135	268	396	402	364	155	31	-
2002.....	-	3	17	39	88	211	324	359	297	118	17	-
2001.....	1	6	13	32	80	194	271	313	197	106	23	-
2000.....	4	5	14	43	85	180	221	288	201	76	8	-
1999.....	3	2	12	25	75	166	196	244	174	74	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Alzheimer's disease (G30)

##### Hispanic, male

2003.....	4	3	8	18	57	114	150	123	102	48	4	-
2002.....	-	-	8	18	37	85	111	102	72	25	5	-
2001.....	1	-	6	16	28	89	97	88	50	20	3	-
2000.....	3	2	7	16	47	75	91	83	40	16	2	-
1999.....	-	-	5	17	34	67	61	72	38	18	1	-

##### Hispanic, female

2003.....	6	2	14	26	78	154	246	279	262	107	27	-
2002.....	-	3	9	21	51	126	213	257	225	93	12	-
2001.....	-	6	7	16	52	105	174	225	147	86	20	-
2000.....	1	3	7	27	38	105	130	205	161	60	6	-
1999.....	3	2	7	8	41	99	135	172	136	56	8	-

##### Non-Hispanic 1/, both sexes

2003.....	47	146	379	932	2,715	7,069	13,390	16,934	13,389	5,533	943	-
2002.....	40	132	356	895	2,568	6,849	12,703	15,872	11,942	5,002	861	-
2001.....	55	159	350	826	2,479	6,362	11,689	14,497	10,949	4,360	771	-
2000.....	43	137	332	823	2,473	6,130	10,699	13,013	10,039	3,929	689	1
1999.....	37	123	311	804	2,295	5,594	9,853	11,596	8,845	3,392	594	-

##### Non-Hispanic 1/, male

2003.....	21	72	159	436	1,167	2,740	4,592	4,781	2,827	773	87	-
2002.....	22	55	165	388	1,099	2,699	4,316	4,462	2,493	704	74	-
2001.....	26	61	161	381	1,046	2,520	4,080	4,125	2,258	588	75	-
2000.....	21	58	142	404	1,085	2,356	3,702	3,596	2,057	548	51	-
1999.....	18	68	149	372	1,012	2,262	3,506	3,302	1,816	487	47	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Alzheimer's disease (G30)

## Non-Hispanic 1/, female

2003.....	26	74	220	496	1,548	4,329	8,798	12,153	10,562	4,760	856	-
2002.....	18	77	191	507	1,469	4,150	8,387	11,410	9,449	4,298	787	-
2001.....	29	98	189	445	1,433	3,842	7,609	10,372	8,691	3,772	696	-
2000.....	22	79	190	419	1,388	3,774	6,997	9,417	7,982	3,381	638	1
1999.....	19	55	162	432	1,283	3,332	6,347	8,294	7,029	2,905	547	-

## Non-Hispanic White, both sexes

2003.....	45	136	353	815	2,454	6,469	12,468	15,885	12,589	5,184	843	-
2002.....	37	124	324	804	2,345	6,332	11,895	14,914	11,241	4,684	766	-
2001.....	53	152	328	736	2,284	5,829	10,913	13,619	10,349	4,070	682	-
2000.....	37	127	301	751	2,253	5,676	10,028	12,242	9,507	3,681	622	-
1999.....	33	111	288	731	2,090	5,195	9,258	10,953	8,422	3,189	542	-

## Non-Hispanic White, male

2003.....	20	68	144	385	1,058	2,523	4,273	4,510	2,651	708	77	-
2002.....	19	51	152	345	1,008	2,510	4,034	4,217	2,343	651	59	-
2001.....	24	56	149	338	967	2,329	3,808	3,868	2,125	557	63	-
2000.....	18	53	130	373	994	2,195	3,491	3,376	1,930	509	42	-
1999.....	16	61	139	338	925	2,113	3,319	3,120	1,719	451	40	-

## Non-Hispanic White, female

2003.....	25	68	209	430	1,396	3,946	8,195	11,375	9,938	4,476	766	-
2002.....	18	73	172	459	1,337	3,822	7,861	10,697	8,898	4,033	707	-
2001.....	29	96	179	398	1,317	3,500	7,105	9,751	8,224	3,513	619	-
2000.....	19	74	171	378	1,259	3,481	6,537	8,866	7,577	3,172	580	-
1999.....	17	50	149	393	1,165	3,082	5,939	7,833	6,703	2,738	502	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Alzheimer's disease (G30)

##### Non-Hispanic Black, both sexes

2003.....	2	8	21	100	226	535	812	900	694	304	89	-
2002.....	3	3	28	77	200	468	705	834	611	288	81	-
2001.....	2	6	20	80	173	499	677	782	522	260	80	-
2000.....	4	7	23	67	194	399	605	680	463	218	57	1
1999.....	2	10	19	62	187	358	525	567	362	182	48	-

##### Non-Hispanic Black, male

2003.....	1	3	14	45	97	203	278	233	147	49	8	-
2002.....	3	2	12	36	86	175	243	199	118	50	12	-
2001.....	2	5	12	38	72	182	230	219	109	25	11	-
2000.....	2	5	11	30	83	136	184	181	99	36	6	-
1999.....	2	6	9	31	81	130	160	157	75	31	7	-

##### Non-Hispanic Black, female

2003.....	1	5	7	55	129	332	534	667	547	255	81	-
2002.....	-	1	16	41	114	293	462	635	493	238	69	-
2001.....	-	1	8	42	101	317	447	563	413	235	69	-
2000.....	2	2	12	37	111	263	421	499	364	182	51	1
1999.....	-	4	10	31	106	228	365	410	287	151	41	-

##### Origin not stated 2/, both sexes

2003.....	-	1	1	3	7	11	23	32	21	16	1	-
2002.....	-	-	2	5	7	18	30	36	28	20	-	-
2001.....	-	-	1	-	5	12	14	33	22	8	2	-
2000.....	-	-	3	2	5	9	14	36	17	9	4	1
1999.....	-	-	-	1	4	9	18	22	22	7	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Alzheimer's disease (G30)

##### Origin not stated 2/, male

2003.....	-	1	-	2	4	1	9	12	4	3	-	-
2002.....	-	-	-	2	3	9	7	5	5	3	-	-
2001.....	-	-	-	-	1	7	2	12	4	-	1	-
2000.....	-	-	1	-	2	2	2	8	4	-	1	1
1999.....	-	-	-	-	2	3	5	8	7	-	-	-

##### Origin not stated 2/, female

2003.....	-	-	1	1	3	10	14	20	17	13	1	-
2002.....	-	-	2	3	4	9	23	31	23	17	-	-
2001.....	-	-	1	-	4	5	12	21	18	8	1	-
2000.....	-	-	2	2	3	7	12	28	13	9	3	-
1999.....	-	-	-	1	2	6	13	14	15	7	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Major cardiovascular diseases (I00-I78)

##### All origins, both sexes

2003.....	27,188	35,465	44,032	55,941	81,279	121,274	158,084	160,879	117,184	47,760	10,626	38
2002.....	27,084	35,004	43,426	57,731	86,340	126,160	161,255	163,421	117,826	47,620	10,542	51
2001.....	26,682	33,335	42,889	59,066	89,343	130,631	161,364	164,000	116,097	47,232	10,601	72
2000.....	25,798	33,454	44,028	61,769	94,418	135,296	160,906	165,313	117,092	47,427	10,618	62
1999.....	25,158	33,344	44,489	64,775	98,653	138,680	163,872	166,070	116,257	47,656	10,527	49

##### All origins, male

2003.....	19,045	24,105	28,699	34,660	46,852	63,584	72,284	60,672	33,792	9,930	1,509	29
2002.....	18,930	23,721	28,316	35,603	49,716	65,751	73,019	61,344	33,439	9,860	1,521	36
2001.....	18,442	22,489	27,831	36,221	51,535	67,972	72,817	59,914	32,127	9,565	1,491	53
2000.....	17,991	22,817	28,606	37,721	54,789	70,196	72,601	60,379	32,104	9,636	1,539	43
1999.....	17,595	22,653	28,872	39,978	56,926	72,052	73,515	60,030	31,573	9,558	1,502	38

##### All origins, female

2003.....	8,143	11,360	15,333	21,281	34,427	57,690	85,800	100,207	83,392	37,830	9,117	9
2002.....	8,154	11,283	15,110	22,128	36,624	60,409	88,236	102,077	84,387	37,760	9,021	15
2001.....	8,240	10,846	15,058	22,845	37,808	62,659	88,547	104,086	83,970	37,667	9,110	19
2000.....	7,807	10,637	15,422	24,048	39,629	65,100	88,305	104,934	84,988	37,791	9,079	19
1999.....	7,563	10,691	15,617	24,797	41,727	66,628	90,357	106,040	84,684	38,098	9,025	11

##### Hispanic, both sexes

2003.....	1,562	2,006	2,425	3,123	4,146	5,299	5,484	4,722	3,494	1,393	380	6
2002.....	1,521	1,956	2,504	3,097	4,245	5,078	5,284	4,558	3,306	1,493	330	1
2001.....	1,524	1,745	2,251	3,094	4,231	5,154	4,985	4,684	3,247	1,405	318	7
2000.....	1,356	1,711	2,330	3,003	4,111	4,742	4,638	4,566	3,165	1,299	266	5
1999.....	1,392	1,696	2,277	3,147	4,003	4,708	4,583	4,442	3,041	1,290	258	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Major cardiovascular diseases (I00-I78)

##### Hispanic, male

2003.....	1,125	1,363	1,565	1,873	2,351	2,703	2,533	1,778	1,057	379	86	5
2002.....	1,051	1,346	1,612	1,849	2,455	2,635	2,404	1,745	1,075	389	75	1
2001.....	1,058	1,170	1,412	1,898	2,371	2,652	2,314	1,798	1,019	343	72	6
2000.....	960	1,174	1,483	1,766	2,357	2,505	2,117	1,692	1,014	358	64	4
1999.....	1,004	1,124	1,436	1,890	2,322	2,523	1,996	1,645	992	358	63	6

##### Hispanic, female

2003.....	437	643	860	1,250	1,795	2,596	2,951	2,944	2,437	1,014	294	1
2002.....	470	610	892	1,248	1,790	2,443	2,880	2,813	2,231	1,104	255	-
2001.....	466	575	839	1,196	1,860	2,502	2,671	2,886	2,228	1,062	246	1
2000.....	396	537	847	1,237	1,754	2,237	2,521	2,874	2,151	941	202	1
1999.....	388	572	841	1,257	1,681	2,185	2,587	2,797	2,049	932	195	-

##### Non-Hispanic 1/, both sexes

2003.....	25,523	33,303	41,420	52,638	76,862	115,670	152,234	155,846	113,451	46,262	10,227	21
2002.....	25,433	32,910	40,740	54,399	81,776	120,688	155,544	158,407	114,222	46,000	10,177	38
2001.....	25,044	31,434	40,441	55,778	84,840	125,069	155,957	158,886	112,579	45,695	10,260	47
2000.....	24,326	31,592	41,472	58,562	89,955	130,119	155,841	160,276	113,625	46,004	10,329	43
1999.....	23,657	31,488	41,985	61,383	94,260	133,532	158,822	161,174	112,886	46,241	10,236	34

##### Non-Hispanic 1/, male

2003.....	17,846	22,621	26,993	32,665	44,336	60,707	69,569	58,775	32,665	9,528	1,419	15
2002.....	17,775	22,267	26,584	33,601	47,065	62,889	70,414	59,421	32,288	9,448	1,440	25
2001.....	17,303	21,204	26,265	34,192	48,986	65,093	70,302	57,972	31,035	9,192	1,412	36
2000.....	16,939	21,532	26,963	35,803	52,202	67,446	70,308	58,511	31,008	9,245	1,468	28
1999.....	16,515	21,413	27,262	37,928	54,345	69,278	71,310	58,202	30,487	9,173	1,436	25

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic 1/, female

2003.....	7,677	10,682	14,427	19,973	32,526	54,963	82,665	97,071	80,786	36,734	8,808	6
2002.....	7,658	10,643	14,156	20,798	34,711	57,799	85,130	98,986	81,934	36,552	8,737	13
2001.....	7,741	10,230	14,176	21,586	35,854	59,976	85,655	100,914	81,544	36,503	8,848	11
2000.....	7,387	10,060	14,509	22,759	37,753	62,673	85,533	101,765	82,617	36,759	8,861	15
1999.....	7,142	10,075	14,723	23,455	39,915	64,254	87,512	102,972	82,399	37,068	8,800	9

Non-Hispanic White, both sexes

2003.....	18,236	25,325	32,123	41,865	63,871	100,168	136,086	142,319	104,121	42,305	8,843	17
2002.....	18,473	25,150	31,715	43,606	68,682	104,667	139,407	144,795	104,762	41,989	8,853	26
2001.....	18,233	23,979	31,501	44,581	71,517	108,821	140,101	145,157	103,653	41,714	8,934	28
2000.....	18,016	24,261	32,405	47,591	76,117	113,698	140,391	146,434	104,746	41,938	9,074	29
1999.....	17,597	24,071	33,023	50,223	79,888	116,966	143,429	147,187	104,261	42,308	9,068	18

Non-Hispanic White, male

2003.....	13,279	17,721	21,540	26,670	37,584	53,459	62,795	54,176	30,141	8,642	1,197	11
2002.....	13,442	17,508	21,228	27,612	40,404	55,507	63,824	54,839	29,602	8,570	1,200	18
2001.....	13,141	16,667	21,059	28,119	42,094	57,548	63,784	53,354	28,550	8,310	1,195	22
2000.....	13,002	16,982	21,699	29,895	45,131	59,825	64,009	53,818	28,511	8,257	1,251	20
1999.....	12,706	16,870	21,981	31,816	47,037	61,605	65,165	53,309	27,980	8,288	1,241	13

Non-Hispanic White, female

2003.....	4,957	7,604	10,583	15,195	26,287	46,709	73,291	88,143	73,980	33,663	7,646	6
2002.....	5,031	7,642	10,487	15,994	28,278	49,160	75,583	89,956	75,160	33,419	7,653	8
2001.....	5,092	7,312	10,442	16,462	29,423	51,273	76,317	91,803	75,103	33,404	7,739	6
2000.....	5,014	7,279	10,706	17,696	30,986	53,873	76,382	92,616	76,235	33,681	7,823	9
1999.....	4,891	7,201	11,042	18,407	32,851	55,361	78,264	93,878	76,281	34,020	7,827	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

Non-Hispanic Black, both sexes

2003.....	6,553	7,059	8,115	9,269	11,113	13,078	13,285	11,026	7,680	3,369	1,201	2
2002.....	6,273	6,902	7,932	9,321	11,140	13,488	13,444	11,215	7,835	3,396	1,150	11
2001.....	6,107	6,601	7,882	9,697	11,387	13,796	13,229	11,446	7,562	3,442	1,150	17
2000.....	5,628	6,524	7,978	9,534	12,027	13,905	13,049	11,736	7,495	3,464	1,131	14
1999.....	5,448	6,612	7,875	9,789	12,453	14,150	12,979	11,800	7,298	3,457	1,026	15

Non-Hispanic Black, male

2003.....	4,053	4,293	4,713	5,104	5,705	6,037	5,398	3,513	1,869	668	169	2
2002.....	3,857	4,200	4,662	5,119	5,651	6,031	5,292	3,521	2,020	655	187	6
2001.....	3,703	3,956	4,533	5,226	5,829	6,257	5,279	3,627	1,914	694	184	13
2000.....	3,468	4,019	4,549	5,053	6,021	6,267	5,167	3,735	1,929	753	188	8
1999.....	3,399	4,001	4,608	5,329	6,238	6,332	4,997	3,853	1,935	736	167	11

Non-Hispanic Black, female

2003.....	2,500	2,766	3,402	4,165	5,408	7,041	7,887	7,513	5,811	2,701	1,032	-
2002.....	2,416	2,702	3,270	4,202	5,489	7,457	8,152	7,694	5,815	2,741	963	5
2001.....	2,404	2,645	3,349	4,471	5,558	7,539	7,950	7,819	5,648	2,748	966	4
2000.....	2,160	2,505	3,429	4,481	6,006	7,638	7,882	8,001	5,566	2,711	943	6
1999.....	2,049	2,611	3,267	4,460	6,215	7,818	7,982	7,947	5,363	2,721	859	4

Origin not stated 2/, both sexes

2003.....	103	156	187	180	271	305	366	311	239	105	19	11
2002.....	130	138	182	235	319	394	427	456	298	127	35	12
2001.....	114	156	197	194	272	408	422	430	271	132	23	18
2000.....	116	151	226	204	352	435	427	471	302	124	23	14
1999.....	109	160	227	245	390	440	467	454	330	125	33	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Major cardiovascular diseases (I00-I78)

Origin not stated 2/, male

2003.....	74	121	141	122	165	174	182	119	70	23	4	9
2002.....	104	108	120	153	196	227	201	178	76	23	6	10
2001.....	81	115	154	131	178	227	201	144	73	30	7	11
2000.....	92	111	160	152	230	245	176	176	82	33	7	11
1999.....	76	116	174	160	259	251	209	183	94	27	3	7

Origin not stated 2/, female

2003.....	29	35	46	58	106	131	184	192	169	82	15	2
2002.....	26	30	62	82	123	167	226	278	222	104	29	2
2001.....	33	41	43	63	94	181	221	286	198	102	16	7
2000.....	24	40	66	52	122	190	251	295	220	91	16	3
1999.....	33	44	53	85	131	189	258	271	236	98	30	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins, both sexes

2003.....	22,212	29,217	35,843	44,470	62,793	90,796	116,535	118,106	86,773	35,782	8,135	35
2002.....	22,391	28,895	35,339	45,948	66,599	94,844	118,737	119,565	86,848	35,664	8,096	47
2001.....	22,009	27,564	34,922	47,131	69,168	98,099	118,893	119,848	85,878	35,479	8,234	66
2000.....	21,167	27,676	35,723	49,178	73,227	101,764	118,296	121,003	86,424	35,605	8,210	59
1999.....	20,898	27,605	36,562	52,206	77,047	104,841	121,311	122,126	85,752	35,797	8,185	47

All origins, male

2003.....	16,140	20,538	23,984	28,270	37,298	49,241	55,405	46,400	25,955	7,662	1,210	27
2002.....	16,256	20,199	23,701	29,045	39,613	51,211	56,006	46,515	25,549	7,663	1,208	34
2001.....	15,789	19,200	23,319	29,643	40,994	52,960	55,616	45,298	24,606	7,394	1,175	47
2000.....	15,331	19,456	23,829	30,890	43,699	54,421	55,272	45,756	24,475	7,430	1,235	41
1999.....	15,143	19,370	24,292	33,118	45,718	56,178	56,292	45,667	23,999	7,442	1,192	37

All origins, female

2003.....	6,072	8,679	11,859	16,200	25,495	41,555	61,130	71,706	60,818	28,120	6,925	8
2002.....	6,135	8,696	11,638	16,903	26,986	43,633	62,731	73,050	61,299	28,001	6,888	13
2001.....	6,220	8,364	11,603	17,488	28,174	45,139	63,277	74,550	61,272	28,085	7,059	19
2000.....	5,836	8,220	11,894	18,288	29,528	47,343	63,024	75,247	61,949	28,175	6,975	18
1999.....	5,755	8,235	12,270	19,088	31,329	48,663	65,019	76,459	61,753	28,355	6,993	10

Hispanic, both sexes

2003.....	1,170	1,541	1,913	2,410	3,225	3,966	4,111	3,522	2,686	1,067	314	6
2002.....	1,169	1,528	1,964	2,472	3,247	3,859	3,920	3,441	2,548	1,197	250	1
2001.....	1,139	1,362	1,790	2,466	3,264	3,875	3,757	3,506	2,531	1,080	252	6
2000.....	1,045	1,355	1,781	2,328	3,125	3,583	3,484	3,438	2,438	995	215	5
1999.....	1,090	1,331	1,824	2,464	3,093	3,618	3,428	3,363	2,349	1,050	206	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Hispanic, male

2003.....	883	1,091	1,243	1,501	1,886	2,074	1,951	1,331	832	281	73	5
2002.....	846	1,087	1,306	1,511	1,930	2,054	1,844	1,372	820	327	52	1
2001.....	832	956	1,148	1,544	1,880	2,031	1,777	1,385	808	265	54	5
2000.....	768	952	1,176	1,409	1,836	1,912	1,613	1,312	793	275	50	4
1999.....	817	924	1,167	1,522	1,833	1,957	1,526	1,250	741	286	47	6

Hispanic, female

2003.....	287	450	670	909	1,339	1,892	2,160	2,191	1,854	786	241	1
2002.....	323	441	658	961	1,317	1,805	2,076	2,069	1,728	870	198	-
2001.....	307	406	642	922	1,384	1,844	1,980	2,121	1,723	815	198	1
2000.....	277	403	605	919	1,289	1,671	1,871	2,126	1,645	720	165	1
1999.....	273	407	657	942	1,260	1,661	1,902	2,113	1,608	764	159	-

Non-Hispanic 1/, both sexes

2003.....	20,959	27,536	33,769	41,905	59,356	86,578	112,135	114,347	83,906	34,635	7,806	20
2002.....	21,104	27,246	33,219	43,274	63,077	90,673	114,483	115,769	84,079	34,366	7,816	34
2001.....	20,771	26,068	32,970	44,501	65,680	93,905	114,799	116,016	83,143	34,294	7,962	42
2000.....	20,021	26,192	33,751	46,667	69,809	97,840	114,488	117,193	83,757	34,509	7,977	41
1999.....	19,715	26,140	34,544	49,540	73,621	100,854	117,505	118,399	83,155	34,644	7,953	32

Non-Hispanic 1/, male

2003.....	15,192	19,333	22,624	26,666	35,277	47,021	53,300	44,969	25,064	7,362	1,133	14
2002.....	15,315	19,015	22,290	27,403	37,509	48,972	54,000	44,997	24,670	7,314	1,151	23
2001.....	14,883	18,146	22,042	27,987	38,962	50,746	53,683	43,799	23,744	7,107	1,116	31
2000.....	14,485	18,410	22,514	29,341	41,669	52,315	53,519	44,305	23,618	7,128	1,180	27
1999.....	14,257	18,345	22,976	31,461	43,657	54,006	54,599	44,256	23,191	7,132	1,142	24

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic 1/, female

2003.....	5,767	8,203	11,145	15,239	24,079	39,557	58,835	69,378	58,842	27,273	6,673	6
2002.....	5,789	8,231	10,929	15,871	25,568	41,701	60,483	70,772	59,409	27,052	6,665	11
2001.....	5,888	7,922	10,928	16,514	26,718	43,159	61,116	72,217	59,399	27,187	6,846	11
2000.....	5,536	7,782	11,237	17,326	28,140	45,525	60,969	72,888	60,139	27,381	6,797	14
1999.....	5,458	7,795	11,568	18,079	29,964	46,848	62,906	74,143	59,964	27,512	6,811	8

Non-Hispanic White, both sexes

2003.....	15,436	21,427	26,664	33,729	49,750	75,414	100,621	104,708	77,242	31,761	6,778	16
2002.....	15,799	21,225	26,269	35,099	53,429	78,991	102,984	106,117	77,340	31,405	6,808	23
2001.....	15,656	20,256	26,177	36,014	55,841	82,060	103,476	106,333	76,775	31,359	6,963	26
2000.....	15,248	20,515	26,777	38,325	59,423	85,862	103,504	107,375	77,430	31,597	7,052	29
1999.....	15,061	20,346	27,604	40,913	62,837	88,727	106,460	108,389	76,983	31,757	7,064	18

Non-Hispanic White, male

2003.....	11,621	15,485	18,386	21,997	30,151	41,666	48,319	41,591	23,226	6,715	950	10
2002.....	11,908	15,237	18,076	22,786	32,479	43,461	49,155	41,657	22,700	6,647	972	16
2001.....	11,662	14,516	17,988	23,298	33,761	45,093	48,903	40,491	21,898	6,442	944	20
2000.....	11,402	14,800	18,379	24,751	36,273	46,643	48,933	40,903	21,817	6,412	1,006	20
1999.....	11,246	14,675	18,791	26,627	38,072	48,308	50,114	40,681	21,346	6,460	989	13

Non-Hispanic White, female

2003.....	3,815	5,942	8,278	11,732	19,599	33,748	52,302	63,117	54,016	25,046	5,828	6
2002.....	3,891	5,988	8,193	12,313	20,950	35,530	53,829	64,460	54,640	24,758	5,836	7
2001.....	3,994	5,740	8,189	12,716	22,080	36,967	54,573	65,842	54,877	24,917	6,019	6
2000.....	3,846	5,715	8,398	13,574	23,150	39,219	54,571	66,472	55,613	25,185	6,046	9
1999.....	3,815	5,671	8,813	14,286	24,765	40,419	56,346	67,708	55,637	25,297	6,075	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Non-Hispanic Black, both sexes

2003.....	4,983	5,440	6,224	7,093	8,316	9,512	9,534	7,888	5,490	2,466	890	2
2002.....	4,812	5,389	6,165	7,109	8,302	9,863	9,657	8,003	5,623	2,529	875	10
2001.....	4,611	5,178	6,031	7,418	8,522	10,132	9,549	8,157	5,431	2,542	874	14
2000.....	4,286	5,076	6,181	7,302	9,111	10,227	9,373	8,381	5,367	2,479	832	12
1999.....	4,209	5,202	6,144	7,622	9,397	10,398	9,410	8,483	5,242	2,546	785	14

Non-Hispanic Black, male

2003.....	3,163	3,382	3,658	3,999	4,378	4,503	3,985	2,587	1,374	497	139	2
2002.....	3,030	3,338	3,690	3,965	4,282	4,503	3,925	2,580	1,496	507	135	6
2001.....	2,868	3,170	3,543	4,044	4,447	4,705	3,916	2,630	1,451	528	149	10
2000.....	2,718	3,193	3,593	3,929	4,631	4,692	3,784	2,740	1,398	544	153	7
1999.....	2,692	3,245	3,662	4,226	4,790	4,725	3,691	2,831	1,433	553	128	11

Non-Hispanic Black, female

2003.....	1,820	2,058	2,566	3,094	3,938	5,009	5,549	5,301	4,116	1,969	751	-
2002.....	1,782	2,051	2,475	3,144	4,020	5,360	5,732	5,423	4,127	2,022	740	4
2001.....	1,743	2,008	2,488	3,374	4,075	5,427	5,633	5,527	3,980	2,014	725	4
2000.....	1,568	1,883	2,588	3,373	4,480	5,535	5,589	5,641	3,969	1,935	679	5
1999.....	1,517	1,957	2,482	3,396	4,607	5,673	5,719	5,652	3,809	1,993	657	3

Origin not stated 2/, both sexes

2003.....	83	140	161	155	212	252	289	237	181	80	15	9
2002.....	118	121	156	202	275	312	334	355	221	101	30	12
2001.....	99	134	162	164	224	319	337	326	204	105	20	18
2000.....	101	129	191	183	293	341	324	372	229	101	18	13
1999.....	93	134	194	202	333	369	378	364	248	103	26	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Origin not stated 2/, male

2003.....	65	114	117	103	135	146	154	100	59	19	4	8
2002.....	95	97	105	131	174	185	162	146	59	22	5	10
2001.....	74	98	129	112	152	183	156	114	54	22	5	11
2000.....	78	94	139	140	194	194	140	139	64	27	5	10
1999.....	69	101	149	135	228	215	167	161	67	24	3	7

Origin not stated 2/, female

2003.....	18	26	44	52	77	106	135	137	122	61	11	1
2002.....	23	24	51	71	101	127	172	209	162	79	25	2
2001.....	25	36	33	52	72	136	181	212	150	83	15	7
2000.....	23	35	52	43	99	147	184	233	165	74	13	3
1999.....	24	33	45	67	105	154	211	203	181	79	23	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All origins, both sexes

2003.....	117	151	216	295	398	603	630	557	291	93	23	-
2002.....	140	190	243	272	431	620	607	462	263	82	12	-
2001.....	125	168	193	307	437	599	589	468	235	67	10	-
2000.....	133	183	237	327	492	601	575	416	231	54	11	-
1999.....	144	189	222	338	516	614	613	442	235	63	8	-

All origins, male

2003.....	39	62	84	105	136	199	185	160	62	10	2	-
2002.....	55	73	83	84	161	177	164	100	55	8	1	-
2001.....	41	73	62	106	125	182	148	106	50	10	-	-
2000.....	55	72	94	98	139	177	144	96	40	13	1	-
1999.....	50	64	69	107	183	164	161	89	26	8	-	-

All origins, female

2003.....	78	89	132	190	262	404	445	397	229	83	21	-
2002.....	85	117	160	188	270	443	443	362	208	74	11	-
2001.....	84	95	131	201	312	417	441	362	185	57	10	-
2000.....	78	111	143	229	353	424	431	320	191	41	10	-
1999.....	94	125	153	231	333	450	452	353	209	55	8	-

Hispanic, both sexes

2003.....	11	10	11	23	17	23	20	13	3	-	-	-
2002.....	11	15	20	21	33	22	20	11	6	2	-	-
2001.....	7	9	18	23	25	17	13	10	6	-	1	-
2000.....	8	14	16	19	28	13	14	7	3	-	-	-
1999.....	13	12	14	25	19	22	13	8	2	3	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Hispanic, male

2003.....	3	5	3	9	4	8	3	5	-	-	-	-
2002.....	5	5	8	6	9	2	2	2	2	-	-	-
2001.....	2	4	3	11	4	8	3	2	1	-	-	-
2000.....	7	5	7	5	8	5	1	2	1	-	-	-
1999.....	2	2	6	6	4	3	2	-	-	-	-	-

Hispanic, female

2003.....	8	5	8	14	13	15	17	8	3	-	-	-
2002.....	6	10	12	15	24	20	18	9	4	2	-	-
2001.....	5	5	15	12	21	9	10	8	5	-	1	-
2000.....	1	9	9	14	20	8	13	5	2	-	-	-
1999.....	11	10	8	19	15	19	11	8	2	3	-	-

Non-Hispanic 1/, both sexes

2003.....	106	140	204	272	381	580	610	543	287	93	22	-
2002.....	128	175	222	250	397	597	586	451	257	80	12	-
2001.....	117	159	174	284	412	581	576	457	229	67	9	-
2000.....	125	169	219	308	464	588	561	409	226	54	11	-
1999.....	130	176	207	311	495	590	600	433	233	60	8	-

Non-Hispanic 1/, male

2003.....	36	56	81	96	132	191	182	155	62	10	2	-
2002.....	49	68	74	78	152	175	162	98	53	8	1	-
2001.....	39	69	59	95	121	174	145	104	49	10	-	-
2000.....	48	67	86	93	131	172	143	94	39	13	1	-
1999.....	47	61	62	101	178	161	159	89	26	8	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic 1/, female

2003.....	70	84	123	176	249	389	428	388	225	83	20	-
2002.....	79	107	148	172	245	422	424	353	204	72	11	-
2001.....	78	90	115	189	291	407	431	353	180	57	9	-
2000.....	77	102	133	215	333	416	418	315	187	41	10	-
1999.....	83	115	145	210	317	429	441	344	207	52	8	-

Non-Hispanic White, both sexes

2003.....	67	98	156	236	335	545	584	520	280	89	21	-
2002.....	99	134	179	209	366	547	562	428	251	77	11	-
2001.....	79	118	147	238	363	540	554	436	222	62	9	-
2000.....	99	134	186	274	413	562	540	393	211	50	11	-
1999.....	94	143	171	276	454	555	570	412	225	54	7	-

Non-Hispanic White, male

2003.....	24	37	61	82	121	185	179	149	60	10	2	-
2002.....	37	51	62	64	141	161	158	95	52	8	-	-
2001.....	30	57	51	77	110	164	138	99	47	8	-	-
2000.....	41	57	79	83	114	164	139	91	35	11	1	-
1999.....	37	50	52	94	166	148	153	87	26	7	-	-

Non-Hispanic White, female

2003.....	43	61	95	154	214	360	405	371	220	79	19	-
2002.....	62	83	117	145	225	386	404	333	199	69	11	-
2001.....	49	61	96	161	253	376	416	337	175	54	9	-
2000.....	58	77	107	191	299	398	401	302	176	39	10	-
1999.....	57	93	119	182	288	407	417	325	199	47	7	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Non-Hispanic Black, both sexes

2003.....	31	34	38	30	36	23	17	16	4	3	1	-
2002.....	23	30	32	29	19	28	14	12	2	2	1	-
2001.....	33	31	22	31	38	28	12	13	4	5	-	-
2000.....	20	26	22	31	27	21	15	12	11	4	-	-
1999.....	29	29	22	24	30	23	16	16	5	4	1	-

Non-Hispanic Black, male

2003.....	11	17	16	13	8	6	2	4	2	-	-	-
2002.....	10	12	9	12	8	8	3	2	-	-	1	-
2001.....	9	10	7	11	6	5	3	4	-	2	-	-
2000.....	5	7	5	8	8	7	1	3	3	2	-	-
1999.....	10	9	7	6	9	10	4	1	-	-	-	-

Non-Hispanic Black, female

2003.....	20	17	22	17	28	17	15	12	2	3	1	-
2002.....	13	18	23	17	11	20	11	10	2	2	-	-
2001.....	24	21	15	20	32	23	9	9	4	3	-	-
2000.....	15	19	17	23	19	14	14	9	8	2	-	-
1999.....	19	20	15	18	21	13	12	15	5	4	1	-

Origin not stated 2/, both sexes

2003.....	-	1	1	-	-	-	-	1	1	-	1	-
2002.....	1	-	1	1	1	1	1	-	-	-	-	-
2001.....	1	-	1	-	-	1	-	1	-	-	-	-
2000.....	-	-	2	-	-	-	-	-	2	-	-	-
1999.....	1	1	1	2	2	2	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Origin not stated 2/, male

2003.....	-	1	-	-	-	-	-	-	-	-	-	-
2002.....	1	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	1	1	1	-	1	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	1	-	-	-	-	1	1	-	1	-
2002.....	-	-	-	1	1	1	1	-	-	-	-	-
2001.....	1	-	1	-	-	1	-	1	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	2	-	-	-
1999.....	-	-	-	2	1	2	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hypertensive heart disease (I11)

##### All origins, both sexes

2003.....	1,784	1,852	1,799	1,892	2,153	2,945	3,620	3,976	3,285	1,590	338	-
2002.....	1,616	1,661	1,659	1,807	2,088	2,816	3,505	3,716	3,115	1,442	314	4
2001.....	1,414	1,380	1,389	1,557	2,062	2,719	3,400	3,658	2,958	1,398	331	11
2000.....	1,152	1,249	1,423	1,570	2,035	2,694	3,268	3,684	2,886	1,330	286	3
1999.....	1,041	1,260	1,277	1,553	1,964	2,672	3,158	3,498	2,710	1,188	283	1

##### All origins, male

2003.....	1,195	1,222	1,086	1,141	1,123	1,326	1,301	1,143	669	206	36	-
2002.....	1,094	1,107	1,052	1,046	1,042	1,259	1,250	1,004	629	194	44	4
2001.....	931	879	870	906	1,055	1,188	1,226	957	565	197	27	9
2000.....	811	789	895	909	1,062	1,135	1,134	960	545	181	21	2
1999.....	673	800	760	858	958	1,185	1,067	912	499	160	23	1

##### All origins, female

2003.....	589	630	713	751	1,030	1,619	2,319	2,833	2,616	1,384	302	-
2002.....	522	554	607	761	1,046	1,557	2,255	2,712	2,486	1,248	270	-
2001.....	483	501	519	651	1,007	1,531	2,174	2,701	2,393	1,201	304	2
2000.....	341	460	528	661	973	1,559	2,134	2,724	2,341	1,149	265	1
1999.....	368	460	517	695	1,006	1,487	2,091	2,586	2,211	1,028	260	-

##### Hispanic, both sexes

2003.....	109	119	128	135	145	169	178	156	119	58	13	-
2002.....	97	122	119	130	134	162	146	153	90	43	13	-
2001.....	106	90	94	106	133	133	127	115	78	41	9	-
2000.....	63	56	85	80	118	116	117	108	71	30	7	-
1999.....	56	77	69	95	79	103	112	99	80	30	8	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hypertensive heart disease (I11)

##### Hispanic, male

2003.....	79	84	84	79	88	85	81	52	32	9	1	-
2002.....	70	84	75	72	77	81	68	46	25	9	2	-
2001.....	75	57	63	63	69	57	51	42	24	7	3	-
2000.....	42	37	49	44	62	59	44	30	17	7	2	-
1999.....	35	51	45	52	43	44	38	24	18	6	2	-

##### Hispanic, female

2003.....	30	35	44	56	57	84	97	104	87	49	12	-
2002.....	27	38	44	58	57	81	78	107	65	34	11	-
2001.....	31	33	31	43	64	76	76	73	54	34	6	-
2000.....	21	19	36	36	56	57	73	78	54	23	5	-
1999.....	21	26	24	43	36	59	74	75	62	24	6	-

##### Non-Hispanic 1/, both sexes

2003.....	1,667	1,720	1,658	1,746	1,995	2,770	3,429	3,808	3,156	1,527	325	-
2002.....	1,505	1,529	1,531	1,650	1,941	2,642	3,343	3,543	3,020	1,393	301	2
2001.....	1,288	1,272	1,282	1,447	1,917	2,571	3,266	3,530	2,872	1,354	321	9
2000.....	1,080	1,185	1,333	1,475	1,899	2,561	3,145	3,557	2,811	1,299	279	2
1999.....	978	1,170	1,193	1,446	1,869	2,559	3,032	3,388	2,619	1,151	275	1

##### Non-Hispanic 1/, male

2003.....	1,109	1,127	994	1,055	1,027	1,237	1,215	1,087	636	197	35	-
2002.....	1,014	1,017	971	957	959	1,170	1,174	949	603	184	42	2
2001.....	842	811	797	841	974	1,125	1,174	910	539	190	24	7
2000.....	762	745	841	852	987	1,065	1,087	923	528	174	19	1
1999.....	635	742	703	798	904	1,135	1,022	884	480	154	21	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

Non-Hispanic 1/, female

2003.....	558	593	664	691	968	1,533	2,214	2,721	2,520	1,330	290	-
2002.....	491	512	560	693	982	1,472	2,169	2,594	2,417	1,209	259	-
2001.....	446	461	485	606	943	1,446	2,092	2,620	2,333	1,164	297	2
2000.....	318	440	492	623	912	1,496	2,058	2,634	2,283	1,125	260	1
1999.....	343	428	490	648	965	1,424	2,010	2,504	2,139	997	254	-

Non-Hispanic White, both sexes

2003.....	933	1,030	1,057	1,097	1,368	2,025	2,798	3,203	2,735	1,339	257	-
2002.....	853	912	920	1,045	1,335	1,941	2,633	2,970	2,610	1,218	248	1
2001.....	701	749	771	898	1,300	1,913	2,627	2,952	2,528	1,183	252	5
2000.....	580	661	780	940	1,253	1,887	2,499	3,000	2,413	1,101	230	-
1999.....	531	648	723	904	1,243	1,868	2,426	2,874	2,280	982	223	-

Non-Hispanic White, male

2003.....	681	718	665	682	707	915	993	906	527	167	29	-
2002.....	618	644	595	620	674	866	927	795	508	151	29	1
2001.....	497	511	505	551	686	848	933	740	468	155	11	4
2000.....	435	444	512	561	681	790	850	779	439	140	17	-
1999.....	373	436	446	504	618	835	812	730	411	123	17	-

Non-Hispanic White, female

2003.....	252	312	392	415	661	1,110	1,805	2,297	2,208	1,172	228	-
2002.....	235	268	325	425	661	1,075	1,706	2,175	2,102	1,067	219	-
2001.....	204	238	266	347	614	1,065	1,694	2,212	2,060	1,028	241	1
2000.....	145	217	268	379	572	1,097	1,649	2,221	1,974	961	213	-
1999.....	158	212	277	400	625	1,033	1,614	2,144	1,869	859	206	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

Non-Hispanic Black, both sexes

2003.....	703	651	559	594	576	665	568	528	365	169	64	-
2002.....	626	585	573	561	563	636	631	506	354	158	47	-
2001.....	556	493	479	510	568	599	565	520	313	155	65	4
2000.....	471	504	521	491	602	617	577	505	359	172	46	2
1999.....	435	491	441	508	576	632	550	455	299	154	46	1

Non-Hispanic Black, male

2003.....	406	384	303	346	300	283	199	149	93	26	6	-
2002.....	377	353	354	308	266	270	214	133	75	24	10	-
2001.....	327	281	270	273	265	246	212	148	62	28	12	3
2000.....	305	288	306	267	280	243	209	132	77	24	2	1
1999.....	252	287	241	274	262	270	188	131	60	30	3	1

Non-Hispanic Black, female

2003.....	297	267	256	248	276	382	369	379	272	143	58	-
2002.....	249	232	219	253	297	366	417	373	279	134	37	-
2001.....	229	212	209	237	303	353	353	372	251	127	53	1
2000.....	166	216	215	224	322	374	368	373	282	148	44	1
1999.....	183	204	200	234	314	362	362	324	239	124	43	-

Origin not stated 2/, both sexes

2003.....	8	13	13	11	13	6	13	12	10	5	-	-
2002.....	14	10	9	27	13	12	16	20	5	6	-	2
2001.....	20	18	13	4	12	15	7	13	8	3	1	2
2000.....	9	8	5	15	18	17	6	19	4	1	-	1
1999.....	7	13	15	12	16	10	14	11	11	7	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart disease (I11)

Origin not stated 2/, male

2003.....	7	11	8	7	8	4	5	4	1	-	-	-
2002.....	10	6	6	17	6	8	8	9	1	1	-	2
2001.....	14	11	10	2	12	6	1	5	2	-	-	2
2000.....	7	7	5	13	13	11	3	7	-	-	-	1
1999.....	3	7	12	8	11	6	7	4	1	-	-	-

Origin not stated 2/, female

2003.....	1	2	5	4	5	2	8	8	9	5	-	-
2002.....	4	4	3	10	7	4	8	11	4	5	-	-
2001.....	6	7	3	2	-	9	6	8	6	3	1	-
2000.....	2	1	-	2	5	6	3	12	4	1	-	-
1999.....	4	6	3	4	5	4	7	7	10	7	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hypertensive heart and renal disease (I13)

##### All origins, both sexes

2003.....	114	136	131	154	247	386	517	560	456	167	26	-
2002.....	90	134	139	156	246	363	503	511	386	150	32	-
2001.....	82	109	138	169	248	374	481	499	348	155	24	-
2000.....	79	103	105	149	268	376	487	494	361	153	24	-
1999.....	76	116	146	201	269	487	600	600	411	162	36	-

##### All origins, male

2003.....	67	86	73	75	114	174	208	207	128	45	6	-
2002.....	56	82	76	67	131	158	199	171	108	38	4	-
2001.....	53	61	65	94	112	158	181	174	102	35	3	-
2000.....	44	57	51	74	132	152	198	188	109	37	7	-
1999.....	50	66	80	99	124	221	239	205	123	41	5	-

##### All origins, female

2003.....	47	50	58	79	133	212	309	353	328	122	20	-
2002.....	34	52	63	89	115	205	304	340	278	112	28	-
2001.....	29	48	73	75	136	216	300	325	246	120	21	-
2000.....	35	46	54	75	136	224	289	306	252	116	17	-
1999.....	26	50	66	102	145	266	361	395	288	121	31	-

##### Hispanic, both sexes

2003.....	4	8	11	10	23	22	16	25	18	6	-	-
2002.....	4	10	16	16	18	25	26	16	19	6	-	-
2001.....	8	6	15	15	19	18	23	15	16	11	-	-
2000.....	8	6	9	7	20	16	16	20	12	6	-	-
1999.....	3	9	10	17	15	20	19	17	12	3	1	-

Worktable Orig291F. Deaths from I13 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hypertensive heart and renal disease (I13)

##### Hispanic, male

2003.....	2	6	5	5	7	10	9	14	6	3	-	-
2002.....	3	7	11	6	12	9	11	8	4	3	-	-
2001.....	5	4	10	10	7	4	13	6	5	5	-	-
2000.....	4	2	3	3	16	11	7	7	3	1	-	-
1999.....	1	5	7	6	8	8	6	10	1	1	1	-

##### Hispanic, female

2003.....	2	2	6	5	16	12	7	11	12	3	-	-
2002.....	1	3	5	10	6	16	15	8	15	3	-	-
2001.....	3	2	5	5	12	14	10	9	11	6	-	-
2000.....	4	4	6	4	4	5	9	13	9	5	-	-
1999.....	2	4	3	11	7	12	13	7	11	2	-	-

##### Non-Hispanic 1/, both sexes

2003.....	110	128	119	144	224	363	500	535	436	161	26	-
2002.....	85	124	123	139	226	336	476	493	367	144	32	-
2001.....	74	102	123	153	227	355	455	482	331	143	24	-
2000.....	71	96	96	141	248	357	469	473	349	146	23	-
1999.....	73	107	135	183	253	466	581	581	399	159	35	-

##### Non-Hispanic 1/, male

2003.....	65	80	67	70	107	164	198	193	122	42	6	-
2002.....	52	75	65	61	117	149	187	162	104	35	4	-
2001.....	48	56	55	84	104	154	167	167	97	30	3	-
2000.....	40	54	48	70	116	140	189	180	106	36	6	-
1999.....	49	61	73	92	116	213	233	194	122	40	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

Non-Hispanic 1/, female

2003.....	45	48	52	74	117	199	302	342	314	119	20	-
2002.....	33	49	58	78	109	187	289	331	263	109	28	-
2001.....	26	46	68	69	123	201	288	315	234	113	21	-
2000.....	31	42	48	71	132	217	280	293	243	110	17	-
1999.....	24	46	62	91	137	253	348	387	277	119	31	-

Non-Hispanic White, both sexes

2003.....	24	39	53	65	131	256	390	418	371	138	17	-
2002.....	17	35	47	57	127	228	376	383	300	121	24	-
2001.....	27	39	56	71	133	241	360	379	278	120	16	-
2000.....	25	40	35	65	155	249	344	378	294	122	12	-
1999.....	28	38	53	83	132	324	444	455	339	126	28	-

Non-Hispanic White, male

2003.....	16	24	33	38	68	125	163	147	109	36	2	-
2002.....	8	23	24	26	70	107	149	126	86	29	2	-
2001.....	16	20	22	40	61	110	140	139	85	24	2	-
2000.....	15	23	18	35	74	111	145	142	87	27	2	-
1999.....	18	21	27	47	61	153	193	156	107	30	4	-

Non-Hispanic White, female

2003.....	8	15	20	27	63	131	227	271	262	102	15	-
2002.....	9	12	23	31	57	121	227	257	214	92	22	-
2001.....	11	19	34	31	72	131	220	240	193	96	14	-
2000.....	10	17	17	30	81	138	199	236	207	95	10	-
1999.....	10	17	26	36	71	171	251	299	232	96	24	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

Non-Hispanic Black, both sexes

2003.....	82	84	64	71	83	97	94	99	54	21	7	-
2002.....	68	88	71	74	91	100	87	99	60	22	8	-
2001.....	47	61	64	78	86	104	83	83	51	21	7	-
2000.....	46	54	58	74	87	94	109	82	49	19	10	-
1999.....	40	66	77	91	114	121	122	113	52	32	7	-

Non-Hispanic Black, male

2003.....	46	54	32	27	36	34	32	35	7	4	2	-
2002.....	44	51	37	29	41	37	29	32	17	6	2	-
2001.....	32	35	31	42	39	39	23	22	11	5	1	-
2000.....	25	30	27	35	39	26	39	29	16	6	4	-
1999.....	28	37	44	41	51	49	34	29	13	9	-	-

Non-Hispanic Black, female

2003.....	36	30	32	44	47	63	62	64	47	17	5	-
2002.....	24	37	34	45	50	63	58	67	43	16	6	-
2001.....	15	26	33	36	47	65	60	61	40	16	6	-
2000.....	21	24	31	39	48	68	70	53	33	13	6	-
1999.....	12	29	33	50	63	72	88	84	39	23	7	-

Origin not stated 2/, both sexes

2003.....	-	-	1	-	-	1	1	-	2	-	-	-
2002.....	1	-	-	1	2	2	1	2	-	-	-	-
2001.....	-	1	-	1	2	1	3	2	1	1	-	-
2000.....	-	1	-	1	-	3	2	1	-	1	1	-
1999.....	-	-	1	1	1	1	-	2	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Hypertensive heart and renal disease (I13)

Origin not stated 2/, male

2003.....	-	-	1	-	-	-	1	-	-	-	-	-
2002.....	1	-	-	-	2	-	1	1	-	-	-	-
2001.....	-	1	-	-	1	-	1	1	-	-	-	-
2000.....	-	1	-	1	-	1	2	1	-	-	1	-
1999.....	-	-	-	1	-	-	-	1	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	1	-	-	2	-	-	-
2002.....	-	-	-	1	-	2	-	1	-	-	-	-
2001.....	-	-	-	1	1	1	2	1	1	1	-	-
2000.....	-	-	-	-	-	2	-	-	-	-	1	-
1999.....	-	-	1	-	1	1	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
All origins, both sexes												
2003.....	15,372	20,907	26,470	32,648	46,232	66,062	83,308	81,998	58,794	23,939	5,433	29
2002.....	15,683	20,997	26,265	34,336	49,531	69,769	85,622	84,082	59,567	24,211	5,528	37
2001.....	15,612	20,270	26,304	35,502	51,961	73,082	86,772	84,880	59,558	24,206	5,665	48
2000.....	15,252	20,724	27,013	37,257	55,628	76,569	87,123	86,890	60,269	24,477	5,683	48
1999.....	15,251	20,575	27,802	40,080	58,760	79,379	89,917	88,158	60,502	24,983	5,733	39
All origins, male												
2003.....	11,820	15,342	18,425	21,547	28,452	37,191	41,382	33,563	18,269	5,232	808	22
2002.....	11,969	15,350	18,306	22,573	30,563	39,154	42,085	34,073	18,199	5,353	821	27
2001.....	11,818	14,794	18,212	23,181	31,891	40,894	42,059	33,466	17,702	5,124	814	33
2000.....	11,548	15,154	18,641	24,225	34,329	42,452	42,295	34,024	17,603	5,216	847	34
1999.....	11,614	15,021	19,102	26,184	36,003	44,041	43,264	34,168	17,472	5,245	841	31
All origins, female												
2003.....	3,552	5,565	8,045	11,101	17,780	28,871	41,926	48,435	40,525	18,707	4,625	7
2002.....	3,714	5,647	7,959	11,763	18,968	30,615	43,537	50,009	41,368	18,858	4,707	10
2001.....	3,794	5,476	8,092	12,321	20,070	32,188	44,713	51,414	41,856	19,082	4,851	15
2000.....	3,704	5,570	8,372	13,032	21,299	34,117	44,828	52,866	42,666	19,261	4,836	14
1999.....	3,637	5,554	8,700	13,896	22,757	35,338	46,653	53,990	43,030	19,738	4,892	8
Hispanic, both sexes												
2003.....	835	1,112	1,434	1,823	2,514	3,058	3,174	2,688	1,998	810	235	6
2002.....	832	1,137	1,546	1,928	2,582	3,022	3,038	2,662	1,970	926	184	1
2001.....	819	1,033	1,371	1,933	2,620	3,113	2,982	2,783	1,981	830	196	4
2000.....	754	1,055	1,402	1,873	2,491	2,854	2,772	2,706	1,914	783	169	4
1999.....	826	1,018	1,445	1,971	2,521	2,937	2,718	2,685	1,852	828	160	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

Hispanic, male

2003.....	671	825	955	1,176	1,504	1,638	1,530	1,028	609	212	57	5
2002.....	625	846	1,061	1,221	1,554	1,646	1,453	1,083	642	251	41	1
2001.....	618	764	903	1,247	1,557	1,683	1,433	1,120	639	206	45	3
2000.....	576	765	961	1,167	1,494	1,573	1,315	1,052	616	211	39	3
1999.....	651	740	937	1,252	1,518	1,629	1,240	1,000	583	234	32	5

Hispanic, female

2003.....	164	287	479	647	1,010	1,420	1,644	1,660	1,389	598	178	1
2002.....	207	291	485	707	1,028	1,376	1,585	1,579	1,328	675	143	-
2001.....	201	269	468	686	1,063	1,430	1,549	1,663	1,342	624	151	1
2000.....	178	290	441	706	997	1,281	1,457	1,654	1,298	572	130	1
1999.....	175	278	508	719	1,003	1,308	1,478	1,685	1,269	594	128	-

Non-Hispanic 1/, both sexes

2003.....	14,482	19,689	24,916	30,702	43,554	62,794	79,917	79,135	56,671	23,070	5,185	16
2002.....	14,769	19,768	24,591	32,256	46,731	66,495	82,339	81,150	57,423	23,213	5,321	29
2001.....	14,731	19,141	24,805	33,437	49,167	69,719	83,520	81,853	57,426	23,298	5,455	30
2000.....	14,418	19,570	25,452	35,235	52,902	73,441	84,082	83,895	58,189	23,617	5,501	33
1999.....	14,351	19,458	26,208	37,948	55,971	76,140	86,895	85,197	58,459	24,071	5,551	26

Non-Hispanic 1/, male

2003.....	11,105	14,429	17,384	20,284	26,840	35,428	39,737	32,456	17,619	5,004	747	11
2002.....	11,275	14,425	17,156	21,251	28,865	37,356	40,510	32,871	17,507	5,086	776	19
2001.....	11,151	13,952	17,208	21,840	30,219	39,063	40,486	32,260	17,031	4,900	765	22
2000.....	10,907	14,315	17,565	22,946	32,677	40,716	40,862	32,863	16,934	4,984	804	23
1999.....	10,907	14,202	18,047	24,820	34,297	42,234	41,893	33,046	16,835	4,991	806	20

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

Non-Hispanic 1/, female

2003.....	3,377	5,260	7,532	10,418	16,714	27,366	40,180	46,679	39,052	18,066	4,438	5
2002.....	3,494	5,343	7,435	11,005	17,866	29,139	41,829	48,279	39,916	18,127	4,545	10
2001.....	3,580	5,189	7,597	11,597	18,948	30,656	43,034	49,593	40,395	18,398	4,690	8
2000.....	3,511	5,255	7,887	12,289	20,225	32,725	43,220	51,032	41,255	18,633	4,697	10
1999.....	3,444	5,256	8,161	13,128	21,674	33,906	45,002	52,151	41,624	19,080	4,745	6

Non-Hispanic White, both sexes

2003.....	11,227	15,872	20,137	25,159	36,951	54,982	71,835	72,577	52,173	21,155	4,489	13
2002.....	11,580	15,910	19,889	26,553	40,000	58,243	74,305	74,501	52,858	21,199	4,635	20
2001.....	11,600	15,364	20,057	27,459	42,222	61,307	75,508	75,163	53,018	21,266	4,770	18
2000.....	11,467	15,772	20,595	29,352	45,515	64,841	76,270	77,010	53,898	21,670	4,867	24
1999.....	11,408	15,660	21,406	31,841	48,288	67,511	79,083	78,186	54,124	22,094	4,947	15

Non-Hispanic White, male

2003.....	8,864	11,948	14,415	17,041	23,175	31,555	36,097	30,108	16,370	4,564	627	8
2002.....	9,120	11,901	14,207	17,941	25,264	33,367	37,003	30,493	16,159	4,618	658	13
2001.....	9,044	11,484	14,292	18,431	26,450	34,963	37,073	29,942	15,720	4,445	651	14
2000.....	8,945	11,787	14,611	19,650	28,724	36,526	37,516	30,438	15,690	4,499	686	17
1999.....	8,905	11,699	15,055	21,341	30,212	38,098	38,665	30,521	15,504	4,538	704	11

Non-Hispanic White, female

2003.....	2,363	3,924	5,722	8,118	13,776	23,427	35,738	42,469	35,803	16,591	3,862	5
2002.....	2,460	4,009	5,682	8,612	14,736	24,876	37,302	44,008	36,699	16,581	3,977	7
2001.....	2,556	3,880	5,765	9,028	15,772	26,344	38,435	45,221	37,298	16,821	4,119	4
2000.....	2,522	3,985	5,984	9,702	16,791	28,315	38,754	46,572	38,208	17,171	4,181	7
1999.....	2,503	3,961	6,351	10,500	18,076	29,413	40,418	47,665	38,620	17,556	4,243	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Non-Hispanic Black, both sexes												
2003.....	2,861	3,329	4,110	4,728	5,632	6,598	6,609	5,286	3,680	1,622	602	1
2002.....	2,830	3,399	4,105	4,864	5,734	6,871	6,670	5,456	3,758	1,699	596	9
2001.....	2,761	3,302	4,149	5,151	5,956	7,097	6,685	5,555	3,717	1,756	597	10
2000.....	2,594	3,340	4,247	5,070	6,415	7,217	6,592	5,809	3,610	1,655	567	9
1999.....	2,614	3,359	4,183	5,331	6,598	7,306	6,584	5,862	3,662	1,743	541	11
Non-Hispanic Black, male												
2003.....	1,924	2,121	2,503	2,708	3,076	3,234	2,860	1,764	924	329	89	1
2002.....	1,864	2,186	2,536	2,788	3,023	3,213	2,800	1,817	1,006	356	87	6
2001.....	1,826	2,098	2,498	2,888	3,201	3,358	2,757	1,796	1,032	358	97	7
2000.....	1,678	2,192	2,521	2,759	3,362	3,404	2,714	1,912	949	366	104	6
1999.....	1,744	2,173	2,564	2,996	3,441	3,380	2,611	1,971	1,011	368	85	9
Non-Hispanic Black, female												
2003.....	937	1,208	1,607	2,020	2,556	3,364	3,749	3,522	2,756	1,293	513	-
2002.....	966	1,213	1,569	2,076	2,711	3,658	3,870	3,639	2,752	1,343	509	3
2001.....	935	1,204	1,651	2,263	2,755	3,739	3,928	3,759	2,685	1,398	500	3
2000.....	916	1,148	1,726	2,311	3,053	3,813	3,878	3,897	2,661	1,289	463	3
1999.....	870	1,186	1,619	2,335	3,157	3,926	3,973	3,891	2,651	1,375	456	2
Origin not stated 2/, both sexes												
2003.....	55	106	120	123	164	210	217	175	125	59	13	7
2002.....	82	92	128	152	218	252	245	270	174	72	23	7
2001.....	62	96	128	132	174	250	270	244	151	78	14	14
2000.....	80	99	159	149	235	274	269	289	166	77	13	11
1999.....	74	99	149	161	268	302	304	276	191	84	22	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Ischemic heart diseases (I20-I25)

Origin not stated 2/, male

2003.....	44	88	86	87	108	125	115	79	41	16	4	6
2002.....	69	79	89	101	144	152	122	119	50	16	4	7
2001.....	49	78	101	94	115	148	140	86	32	18	4	8
2000.....	65	74	115	112	158	163	118	109	53	21	4	8
1999.....	56	79	118	112	188	178	131	122	54	20	3	6

Origin not stated 2/, female

2003.....	11	18	34	36	56	85	102	96	84	43	9	1
2002.....	13	13	39	51	74	100	123	151	124	56	19	-
2001.....	13	18	27	38	59	102	130	158	119	60	10	6
2000.....	15	25	44	37	77	111	151	180	113	56	9	3
1999.....	18	20	31	49	80	124	173	154	137	64	19	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

All origins, both sexes

2003.....	6,328	8,769	10,998	13,108	17,815	24,265	28,843	27,055	17,795	6,527	1,357	5
2002.....	6,564	8,758	11,175	14,230	19,773	26,154	30,436	28,423	18,352	6,623	1,320	5
2001.....	6,720	8,704	11,126	14,690	20,690	27,766	31,488	29,242	18,525	6,606	1,354	10
2000.....	6,647	9,030	11,983	15,898	22,561	30,059	32,151	29,922	18,701	6,755	1,367	8
1999.....	6,598	9,123	12,283	17,333	24,108	31,053	33,658	30,711	18,472	6,765	1,349	7

All origins, male

2003.....	4,854	6,386	7,586	8,531	10,759	13,324	14,026	10,866	5,704	1,469	217	4
2002.....	4,929	6,287	7,679	9,214	12,050	14,316	14,603	11,478	5,741	1,559	213	4
2001.....	5,012	6,271	7,587	9,446	12,517	15,151	14,974	11,411	5,778	1,576	219	6
2000.....	4,946	6,490	8,148	10,177	13,646	16,295	15,336	11,765	5,701	1,539	238	5
1999.....	5,039	6,654	8,375	11,154	14,554	16,991	15,952	12,066	5,566	1,523	202	2

All origins, female

2003.....	1,474	2,383	3,412	4,577	7,056	10,941	14,817	16,189	12,091	5,058	1,140	1
2002.....	1,635	2,471	3,496	5,016	7,723	11,838	15,833	16,945	12,611	5,064	1,107	1
2001.....	1,708	2,433	3,539	5,244	8,173	12,615	16,514	17,831	12,747	5,030	1,135	4
2000.....	1,701	2,540	3,835	5,721	8,915	13,764	16,815	18,157	13,000	5,216	1,129	3
1999.....	1,559	2,469	3,908	6,179	9,554	14,062	17,706	18,645	12,906	5,242	1,147	5

Hispanic, both sexes

2003.....	328	434	495	715	968	1,081	1,107	915	621	237	64	-
2002.....	320	456	633	737	989	1,117	1,067	950	620	270	51	-
2001.....	302	381	533	715	986	1,161	1,029	962	626	245	63	2
2000.....	290	400	541	712	949	1,084	965	896	615	220	43	-
1999.....	312	352	567	750	949	1,131	951	936	548	225	41	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Hispanic, male												
2003.....	260	310	325	447	569	549	518	352	200	58	16	-
2002.....	231	332	423	460	583	600	521	397	182	71	12	-
2001.....	217	267	336	458	578	599	494	386	227	72	14	1
2000.....	207	286	357	437	547	590	457	358	199	64	11	-
1999.....	232	248	359	470	558	655	417	325	194	64	9	-
Hispanic, female												
2003.....	68	124	170	268	399	532	589	563	421	179	48	-
2002.....	89	124	210	277	406	517	546	553	438	199	39	-
2001.....	85	114	197	257	408	562	535	576	399	173	49	1
2000.....	83	114	184	275	402	494	508	538	416	156	32	-
1999.....	80	104	208	280	391	476	534	611	354	161	32	-
Non-Hispanic 1/, both sexes												
2003.....	5,985	8,311	10,477	12,357	16,802	23,127	27,679	26,091	17,141	6,276	1,292	3
2002.....	6,228	8,286	10,515	13,456	18,716	24,985	29,300	27,407	17,685	6,334	1,263	5
2001.....	6,409	8,298	10,546	13,946	19,658	26,537	30,386	28,202	17,858	6,347	1,289	7
2000.....	6,336	8,598	11,402	15,146	21,549	28,908	31,112	28,944	18,040	6,518	1,321	7
1999.....	6,268	8,730	11,679	16,535	23,079	29,839	32,627	29,701	17,871	6,526	1,302	4
Non-Hispanic 1/, male												
2003.....	4,582	6,053	7,244	8,063	10,161	12,744	13,472	10,490	5,490	1,408	200	3
2002.....	4,687	5,943	7,240	8,733	11,428	13,685	14,051	11,046	5,548	1,483	200	4
2001.....	4,787	5,984	7,218	8,968	11,907	14,514	14,442	11,004	5,541	1,500	204	5
2000.....	4,726	6,184	7,766	9,712	13,060	15,666	14,850	11,377	5,485	1,471	225	4
1999.....	4,795	6,374	7,989	10,658	13,940	16,288	15,497	11,711	5,355	1,454	192	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Non-Hispanic 1/, female												
2003.....	1,403	2,258	3,233	4,294	6,641	10,383	14,207	15,601	11,651	4,868	1,092	-
2002.....	1,541	2,343	3,275	4,723	7,288	11,300	15,249	16,361	12,137	4,851	1,063	1
2001.....	1,622	2,314	3,328	4,978	7,751	12,023	15,944	17,198	12,317	4,847	1,085	2
2000.....	1,610	2,414	3,636	5,434	8,489	13,242	16,262	17,567	12,555	5,047	1,096	3
1999.....	1,473	2,356	3,690	5,877	9,139	13,551	17,130	17,990	12,516	5,072	1,110	3
Non-Hispanic White, both sexes												
2003.....	4,760	6,878	8,746	10,353	14,438	20,382	24,817	23,864	15,719	5,665	1,087	3
2002.....	5,046	6,883	8,741	11,314	16,153	21,961	26,356	24,999	16,126	5,698	1,060	4
2001.....	5,185	6,865	8,812	11,690	17,059	23,472	27,436	25,780	16,332	5,728	1,106	4
2000.....	5,161	7,157	9,521	12,841	18,727	25,620	28,220	26,433	16,629	5,882	1,129	6
1999.....	5,175	7,279	9,837	14,202	20,137	26,593	29,696	27,116	16,434	5,930	1,126	2
Non-Hispanic White, male												
2003.....	3,751	5,157	6,195	6,944	8,884	11,406	12,201	9,702	5,065	1,272	160	3
2002.....	3,911	5,066	6,200	7,547	10,126	12,260	12,776	10,161	5,074	1,330	162	3
2001.....	4,006	5,095	6,208	7,742	10,560	13,064	13,218	10,137	5,058	1,337	172	4
2000.....	3,992	5,261	6,658	8,465	11,597	14,129	13,620	10,464	5,037	1,296	197	3
1999.....	4,080	5,420	6,899	9,342	12,449	14,792	14,312	10,743	4,907	1,307	157	-
Non-Hispanic White, female												
2003.....	1,009	1,721	2,551	3,409	5,554	8,976	12,616	14,162	10,654	4,393	927	-
2002.....	1,135	1,817	2,541	3,767	6,027	9,701	13,580	14,838	11,052	4,368	898	1
2001.....	1,179	1,770	2,604	3,948	6,499	10,408	14,218	15,643	11,274	4,391	934	-
2000.....	1,169	1,896	2,863	4,376	7,130	11,491	14,600	15,969	11,592	4,586	932	3
1999.....	1,095	1,859	2,938	4,860	7,688	11,801	15,384	16,373	11,527	4,623	969	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

Non-Hispanic Black, both sexes

2003.....	1,068	1,242	1,457	1,702	2,012	2,335	2,336	1,806	1,188	536	180	-
2002.....	1,054	1,248	1,552	1,810	2,183	2,508	2,431	1,982	1,279	570	179	1
2001.....	1,059	1,254	1,498	1,935	2,238	2,585	2,454	2,007	1,292	539	157	3
2000.....	1,029	1,263	1,647	1,969	2,446	2,734	2,455	2,126	1,191	536	178	1
1999.....	968	1,262	1,600	2,011	2,531	2,751	2,505	2,164	1,248	538	159	2

Non-Hispanic Black, male

2003.....	701	761	863	925	1,061	1,119	984	592	324	103	34	-
2002.....	674	764	887	987	1,099	1,150	1,004	684	353	123	25	1
2001.....	653	761	845	1,028	1,135	1,173	982	671	372	132	28	1
2000.....	623	789	942	1,032	1,247	1,239	1,008	717	343	128	27	1
1999.....	621	812	931	1,113	1,262	1,236	975	758	355	127	30	1

Non-Hispanic Black, female

2003.....	367	481	594	777	951	1,216	1,352	1,214	864	433	146	-
2002.....	380	484	665	823	1,084	1,358	1,427	1,298	926	447	154	-
2001.....	406	493	653	907	1,103	1,412	1,472	1,336	920	407	129	2
2000.....	406	474	705	937	1,199	1,495	1,447	1,409	848	408	151	-
1999.....	347	450	669	898	1,269	1,515	1,530	1,406	893	411	129	1

Origin not stated 2/, both sexes

2003.....	15	24	26	36	45	57	57	49	33	14	1	2
2002.....	16	16	27	37	68	52	69	66	47	19	6	-
2001.....	9	25	47	29	46	68	73	78	41	14	2	1
2000.....	21	32	40	40	63	67	74	82	46	17	3	1
1999.....	18	41	37	48	80	83	80	74	53	14	6	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute myocardial infarction (I21-I22)

Origin not stated 2/, male

2003.....	12	23	17	21	29	31	36	24	14	3	1	1
2002.....	11	12	16	21	39	31	31	35	11	5	1	-
2001.....	8	20	33	20	32	38	38	21	10	4	1	-
2000.....	13	20	25	28	39	39	29	30	17	4	2	1
1999.....	12	32	27	26	56	48	38	30	17	5	1	1

Origin not stated 2/, female

2003.....	3	1	9	15	16	26	21	25	19	11	-	1
2002.....	5	4	11	16	29	21	38	31	36	14	5	-
2001.....	1	5	14	9	14	30	35	57	31	10	1	1
2000.....	8	12	15	12	24	28	45	52	29	13	1	-
1999.....	6	9	10	22	24	35	42	44	36	9	5	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

All origins, both sexes

2003.....	171	198	205	267	312	389	448	447	337	129	28	1
2002.....	226	210	250	289	342	420	478	452	339	153	27	-
2001.....	187	200	263	281	364	397	474	455	317	142	31	-
2000.....	189	219	265	294	350	440	466	439	318	136	35	1
1999.....	160	209	239	294	380	406	449	474	296	116	22	-

All origins, male

2003.....	122	137	141	173	188	220	194	159	93	21	9	1
2002.....	162	153	167	193	197	216	228	172	87	35	4	-
2001.....	141	134	186	180	213	206	218	165	72	25	7	-
2000.....	141	163	178	194	204	232	214	153	68	28	5	1
1999.....	109	159	156	191	219	229	194	170	75	19	2	-

All origins, female

2003.....	49	61	64	94	124	169	254	288	244	108	19	-
2002.....	64	57	83	96	145	204	250	280	252	118	23	-
2001.....	46	66	77	101	151	191	256	290	245	117	24	-
2000.....	48	56	87	100	146	208	252	286	250	108	30	-
1999.....	51	50	83	103	161	177	255	304	221	97	20	-

Hispanic, both sexes

2003.....	2	2	4	3	7	8	4	5	3	2	-	1
2002.....	5	4	8	4	10	9	6	4	2	2	1	-
2001.....	3	6	6	9	10	10	4	7	3	3	-	-
2000.....	2	6	4	4	11	9	8	10	4	2	1	-
1999.....	6	4	9	5	3	10	2	5	5	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Hispanic, male

2003.....	1	-	3	2	3	3	1	1	2	-	-	1
2002.....	4	3	6	1	9	4	3	2	1	-	-	-
2001.....	2	5	4	6	8	5	2	6	-	3	-	-
2000.....	1	3	2	4	7	5	4	6	-	1	-	-
1999.....	3	2	4	3	1	7	1	2	-	-	-	-

Hispanic, female

2003.....	1	2	1	1	4	5	3	4	1	2	-	-
2002.....	1	1	2	3	1	5	3	2	1	2	1	-
2001.....	1	1	2	3	2	5	2	1	3	-	-	-
2000.....	1	3	2	-	4	4	4	4	4	1	1	-
1999.....	3	2	5	2	2	3	1	3	5	1	-	-

Non-Hispanic 1/, both sexes

2003.....	169	194	201	262	304	381	443	441	334	127	28	-
2002.....	221	206	241	285	332	411	472	447	337	151	26	-
2001.....	184	194	256	271	354	387	468	448	314	139	31	-
2000.....	184	213	260	289	337	430	457	428	314	133	34	1
1999.....	154	204	229	288	376	395	446	467	287	115	22	-

Non-Hispanic 1/, male

2003.....	121	135	138	170	184	217	192	158	91	21	9	-
2002.....	158	150	161	192	188	212	225	169	86	35	4	-
2001.....	139	129	181	173	205	201	215	159	72	22	7	-
2000.....	138	160	176	189	197	226	210	147	68	27	5	1
1999.....	106	156	151	188	218	222	192	167	75	19	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Non-Hispanic 1/, female

2003.....	48	59	63	92	120	164	251	283	243	106	19	-
2002.....	63	56	80	93	144	199	247	278	251	116	22	-
2001.....	45	65	75	98	149	186	253	289	242	117	24	-
2000.....	46	53	84	100	140	204	247	281	246	106	29	-
1999.....	48	48	78	100	158	173	254	300	212	96	20	-

Non-Hispanic White, both sexes

2003.....	130	147	157	210	253	324	392	400	304	122	25	-
2002.....	166	170	186	233	277	361	421	404	310	141	22	-
2001.....	134	148	194	209	302	336	421	399	295	122	26	-
2000.....	142	154	203	232	282	363	411	390	284	127	31	-
1999.....	114	158	186	241	312	343	393	431	270	103	21	-

Non-Hispanic White, male

2003.....	96	101	108	139	157	182	164	143	84	21	6	-
2002.....	119	124	127	158	155	190	200	159	77	32	4	-
2001.....	101	102	138	128	179	174	200	142	66	20	7	-
2000.....	106	121	140	150	164	193	189	136	60	26	4	-
1999.....	82	120	120	157	182	199	167	156	69	17	2	-

Non-Hispanic White, female

2003.....	34	46	49	71	96	142	228	257	220	101	19	-
2002.....	47	46	59	75	122	171	221	245	233	109	18	-
2001.....	33	46	56	81	123	162	221	257	229	102	19	-
2000.....	36	33	63	82	118	170	222	254	224	101	27	-
1999.....	32	38	66	84	130	144	226	275	201	86	19	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Non-Hispanic Black, both sexes

2003.....	32	42	39	46	45	51	46	36	26	4	2	-
2002.....	51	31	47	47	53	44	45	38	21	8	4	-
2001.....	42	40	58	49	46	39	40	45	12	16	5	-
2000.....	38	51	52	46	45	56	41	34	26	6	3	1
1999.....	39	42	41	39	54	49	48	28	15	10	1	-

Non-Hispanic Black, male

2003.....	22	32	26	29	24	31	26	14	7	-	2	-
2002.....	35	21	29	32	32	19	21	9	8	3	-	-
2001.....	33	22	40	36	23	24	11	15	4	2	-	-
2000.....	29	35	32	31	27	28	18	9	7	1	1	1
1999.....	23	33	30	26	30	21	20	10	5	2	-	-

Non-Hispanic Black, female

2003.....	10	10	13	17	21	20	20	22	19	4	-	-
2002.....	16	10	18	15	21	25	24	29	13	5	4	-
2001.....	9	18	18	13	23	15	29	30	8	14	5	-
2000.....	9	16	20	15	18	28	23	25	19	5	2	-
1999.....	16	9	11	13	24	28	28	18	10	8	1	-

Origin not stated 2/, both sexes

2003.....	-	2	-	2	1	-	1	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	1	-	-	-	-
2001.....	-	-	1	1	-	-	2	-	-	-	-	-
2000.....	3	-	1	1	2	1	1	1	-	1	-	-
1999.....	-	1	1	1	1	1	1	2	4	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute ischemic heart diseases (I24)

Origin not stated 2/, male

2003.....	-	2	-	1	1	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	1	1	-	-	1	-	-	-	-	-
2000.....	2	-	-	1	-	1	-	-	-	-	-	-
1999.....	-	1	1	-	-	-	1	1	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	1	-	-	-	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	1	-	1	-	2	-	1	1	-	1	-	-
1999.....	-	-	-	1	1	1	-	1	4	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

All origins, both sexes

2003.....	8,873	11,940	15,267	19,273	28,105	41,408	54,017	54,496	40,662	17,283	4,048	23
2002.....	8,893	12,029	14,840	19,817	29,416	43,195	54,708	55,207	40,876	17,435	4,181	32
2001.....	8,705	11,366	14,915	20,531	30,907	44,919	54,810	55,183	40,716	17,458	4,280	38
2000.....	8,416	11,475	14,765	21,065	32,717	46,070	54,506	56,529	41,250	17,586	4,281	39
1999.....	8,493	11,243	15,280	22,453	34,272	47,920	55,810	56,973	41,734	18,102	4,362	32

All origins, male

2003.....	6,844	8,819	10,698	12,843	17,505	23,647	27,162	22,538	12,472	3,742	582	17
2002.....	6,878	8,910	10,460	13,166	18,316	24,622	27,254	22,423	12,371	3,759	604	23
2001.....	6,665	8,389	10,439	13,555	19,161	25,537	26,867	21,890	11,852	3,523	588	27
2000.....	6,461	8,501	10,315	13,854	20,479	25,925	26,745	22,106	11,834	3,649	604	28
1999.....	6,466	8,208	10,571	14,839	21,230	26,821	27,118	21,932	11,831	3,703	637	29

All origins, female

2003.....	2,029	3,121	4,569	6,430	10,600	17,761	26,855	31,958	28,190	13,541	3,466	6
2002.....	2,015	3,119	4,380	6,651	11,100	18,573	27,454	32,784	28,505	13,676	3,577	9
2001.....	2,040	2,977	4,476	6,976	11,746	19,382	27,943	33,293	28,864	13,935	3,692	11
2000.....	1,955	2,974	4,450	7,211	12,238	20,145	27,761	34,423	29,416	13,937	3,677	11
1999.....	2,027	3,035	4,709	7,614	13,042	21,099	28,692	35,041	29,903	14,399	3,725	3

Hispanic, both sexes

2003.....	505	676	935	1,105	1,539	1,969	2,063	1,768	1,374	571	171	5
2002.....	507	677	905	1,187	1,583	1,896	1,965	1,708	1,348	654	132	1
2001.....	514	646	832	1,209	1,624	1,942	1,949	1,814	1,352	582	133	2
2000.....	462	649	857	1,157	1,531	1,761	1,799	1,800	1,295	561	125	4
1999.....	508	662	869	1,216	1,569	1,796	1,765	1,744	1,299	602	119	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Hispanic, male

2003.....	410	515	627	727	932	1,086	1,011	675	407	154	41	4
2002.....	390	511	632	760	962	1,042	929	684	459	180	29	1
2001.....	399	492	563	783	971	1,079	937	728	412	131	31	2
2000.....	368	476	602	726	940	978	854	688	417	146	28	3
1999.....	416	490	574	779	959	967	822	673	389	170	23	5

Hispanic, female

2003.....	95	161	308	378	607	883	1,052	1,093	967	417	130	1
2002.....	117	166	273	427	621	854	1,036	1,024	889	474	103	-
2001.....	115	154	269	426	653	863	1,012	1,086	940	451	102	-
2000.....	94	173	255	431	591	783	945	1,112	878	415	97	1
1999.....	92	172	295	437	610	829	943	1,071	910	432	96	-

Non-Hispanic 1/, both sexes

2003.....	8,328	11,184	14,238	18,083	26,448	39,286	51,795	52,603	39,196	16,667	3,865	13
2002.....	8,320	11,276	13,835	18,515	27,683	41,099	52,567	53,296	39,401	16,728	4,032	24
2001.....	8,138	10,649	14,003	19,220	29,155	42,795	52,666	53,203	39,254	16,812	4,135	23
2000.....	7,898	10,759	13,790	19,800	31,016	44,103	52,513	54,523	39,835	16,966	4,146	25
1999.....	7,929	10,524	14,300	21,125	32,516	45,906	53,822	55,029	40,301	17,430	4,227	22

Non-Hispanic 1/, male

2003.....	6,402	8,241	10,002	12,051	16,495	22,467	26,073	21,808	12,038	3,575	538	8
2002.....	6,430	8,332	9,755	12,326	17,249	23,459	26,234	21,656	11,873	3,568	572	15
2001.....	6,225	7,839	9,809	12,699	18,107	24,348	25,829	21,097	11,418	3,378	554	17
2000.....	6,043	7,971	9,623	13,045	19,420	24,824	25,802	21,339	11,381	3,486	574	18
1999.....	6,006	7,672	9,907	13,974	20,139	25,724	26,204	21,168	11,405	3,518	612	19

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic 1/, female

2003.....	1,926	2,943	4,236	6,032	9,953	16,819	25,722	30,795	27,158	13,092	3,327	5
2002.....	1,890	2,944	4,080	6,189	10,434	17,640	26,333	31,640	27,528	13,160	3,460	9
2001.....	1,913	2,810	4,194	6,521	11,048	18,447	26,837	32,106	27,836	13,434	3,581	6
2000.....	1,855	2,788	4,167	6,755	11,596	19,279	26,711	33,184	28,454	13,480	3,572	7
1999.....	1,923	2,852	4,393	7,151	12,377	20,182	27,618	33,861	28,896	13,912	3,615	3

Non-Hispanic White, both sexes

2003.....	6,337	8,847	11,234	14,596	22,260	34,276	46,626	48,313	36,150	15,368	3,377	10
2002.....	6,368	8,857	10,962	15,006	23,570	35,921	47,528	49,098	36,422	15,360	3,553	16
2001.....	6,281	8,351	11,051	15,560	24,861	37,499	47,651	48,984	36,391	15,416	3,638	14
2000.....	6,164	8,461	10,871	16,279	26,506	38,858	47,639	50,187	36,985	15,661	3,707	18
1999.....	6,119	8,223	11,383	17,398	27,839	40,575	48,994	50,639	37,420	16,061	3,800	13

Non-Hispanic White, male

2003.....	5,017	6,690	8,112	9,958	14,134	19,967	23,732	20,263	11,221	3,271	461	5
2002.....	5,090	6,711	7,880	10,236	14,983	20,917	24,027	20,173	11,008	3,256	492	10
2001.....	4,937	6,287	7,946	10,561	15,711	21,725	23,655	19,663	10,596	3,088	472	10
2000.....	4,847	6,405	7,813	11,035	16,963	22,204	23,707	19,838	10,593	3,177	485	14
1999.....	4,743	6,159	8,036	11,842	17,581	23,107	24,186	19,622	10,528	3,214	545	11

Non-Hispanic White, female

2003.....	1,320	2,157	3,122	4,638	8,126	14,309	22,894	28,050	24,929	12,097	2,916	5
2002.....	1,278	2,146	3,082	4,770	8,587	15,004	23,501	28,925	25,414	12,104	3,061	6
2001.....	1,344	2,064	3,105	4,999	9,150	15,774	23,996	29,321	25,795	12,328	3,166	4
2000.....	1,317	2,056	3,058	5,244	9,543	16,654	23,932	30,349	26,392	12,484	3,222	4
1999.....	1,376	2,064	3,347	5,556	10,258	17,468	24,808	31,017	26,892	12,847	3,255	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Non-Hispanic Black, both sexes

2003.....	1,761	2,045	2,614	2,980	3,575	4,212	4,227	3,444	2,466	1,082	420	1
2002.....	1,725	2,120	2,506	3,007	3,498	4,319	4,194	3,436	2,458	1,121	413	8
2001.....	1,660	2,008	2,593	3,167	3,672	4,473	4,191	3,503	2,413	1,201	435	7
2000.....	1,527	2,026	2,548	3,055	3,924	4,427	4,096	3,649	2,393	1,113	386	7
1999.....	1,607	2,055	2,542	3,281	4,013	4,506	4,031	3,670	2,399	1,195	381	9

Non-Hispanic Black, male

2003.....	1,201	1,328	1,614	1,754	1,991	2,084	1,850	1,158	593	226	53	1
2002.....	1,155	1,401	1,620	1,769	1,892	2,044	1,775	1,124	645	230	62	5
2001.....	1,140	1,315	1,613	1,824	2,043	2,161	1,764	1,110	656	224	69	6
2000.....	1,026	1,368	1,547	1,696	2,088	2,137	1,688	1,186	599	237	76	4
1999.....	1,100	1,328	1,603	1,857	2,149	2,123	1,616	1,203	651	239	55	8

Non-Hispanic Black, female

2003.....	560	717	1,000	1,226	1,584	2,128	2,377	2,286	1,873	856	367	-
2002.....	570	719	886	1,238	1,606	2,275	2,419	2,312	1,813	891	351	3
2001.....	520	693	980	1,343	1,629	2,312	2,427	2,393	1,757	977	366	1
2000.....	501	658	1,001	1,359	1,836	2,290	2,408	2,463	1,794	876	310	3
1999.....	507	727	939	1,424	1,864	2,383	2,415	2,467	1,748	956	326	1

Origin not stated 2/, both sexes

2003.....	40	80	94	85	118	153	159	125	92	45	12	5
2002.....	66	76	100	115	150	200	176	203	127	53	17	7
2001.....	53	71	80	102	128	182	195	166	110	64	12	13
2000.....	56	67	118	108	170	206	194	206	120	59	10	10
1999.....	56	57	111	112	187	218	223	200	134	70	16	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other forms of chronic ischemic heart disease (I20,I25)

Origin not stated 2/, male

2003.....	32	63	69	65	78	94	78	55	27	13	3	5
2002.....	58	67	73	80	105	121	91	83	39	11	3	7
2001.....	41	58	67	73	83	110	101	65	22	14	3	8
2000.....	50	54	90	83	119	123	89	79	36	17	2	7
1999.....	44	46	90	86	132	130	92	91	37	15	2	5

Origin not stated 2/, female

2003.....	8	17	25	20	40	59	81	70	65	32	9	-
2002.....	8	9	27	35	45	79	85	120	88	42	14	-
2001.....	12	13	13	29	45	72	94	101	88	50	9	5
2000.....	6	13	28	25	51	83	105	127	84	42	8	3
1999.....	12	11	21	26	55	88	131	109	97	55	14	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All origins, both sexes

2003.....	3,533	4,208	4,902	5,194	6,300	8,253	10,086	9,600	7,316	3,135	823	14
2002.....	3,285	4,189	4,604	5,163	6,654	8,649	9,969	10,007	7,555	3,361	873	23
2001.....	3,209	3,924	4,514	5,272	6,733	9,041	10,155	9,984	7,556	3,323	873	25
2000.....	3,086	3,841	4,401	5,309	7,258	9,271	10,274	10,397	7,953	3,429	885	19
1999.....	2,945	3,747	4,636	5,691	7,704	9,666	10,655	10,745	8,204	3,591	863	24

All origins, male

2003.....	2,837	3,203	3,491	3,490	3,932	4,543	4,790	3,759	2,106	700	112	12
2002.....	2,605	3,250	3,299	3,498	4,119	4,746	4,670	3,814	2,165	722	134	17
2001.....	2,549	2,978	3,267	3,483	4,185	4,897	4,774	3,630	2,167	616	108	19
2000.....	2,444	2,978	3,147	3,557	4,512	5,086	4,639	3,740	2,205	691	122	15
1999.....	2,314	2,842	3,281	3,842	4,704	5,200	4,763	3,895	2,196	692	122	22

All origins, female

2003.....	696	1,005	1,411	1,704	2,368	3,710	5,296	5,841	5,210	2,435	711	2
2002.....	680	939	1,305	1,665	2,535	3,903	5,299	6,193	5,390	2,639	739	6
2001.....	660	946	1,247	1,789	2,548	4,144	5,381	6,354	5,389	2,707	765	6
2000.....	642	863	1,254	1,752	2,746	4,185	5,635	6,657	5,748	2,738	763	4
1999.....	631	905	1,355	1,849	3,000	4,466	5,892	6,850	6,008	2,899	741	2

Hispanic, both sexes

2003.....	219	269	345	318	382	417	381	328	237	85	39	3
2002.....	186	258	328	327	395	395	349	264	240	112	31	1
2001.....	223	247	301	346	402	396	339	314	218	116	19	2
2000.....	188	230	292	324	383	331	318	331	233	94	24	2
1999.....	198	247	293	364	383	384	324	293	243	107	26	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Hispanic, male

2003.....	195	223	259	228	252	245	197	143	80	25	10	2
2002.....	158	220	241	221	254	219	156	112	91	36	9	1
2001.....	192	196	220	246	256	234	171	133	66	26	6	2
2000.....	160	182	224	201	237	215	153	133	75	28	9	1
1999.....	171	184	200	268	259	220	152	105	83	32	5	3

Hispanic, female

2003.....	24	46	86	90	130	172	184	185	157	60	29	1
2002.....	28	38	87	106	141	176	193	152	149	76	22	-
2001.....	31	51	81	100	146	162	168	181	152	90	13	-
2000.....	28	48	68	123	146	116	165	198	158	66	15	1
1999.....	27	63	93	96	124	164	172	188	160	75	21	-

Non-Hispanic 1/, both sexes

2003.....	3,293	3,893	4,510	4,846	5,869	7,775	9,661	9,247	7,062	3,041	782	6
2002.....	3,059	3,888	4,221	4,794	6,198	8,189	9,582	9,707	7,293	3,242	840	17
2001.....	2,958	3,637	4,179	4,879	6,280	8,576	9,770	9,635	7,312	3,200	852	11
2000.....	2,867	3,573	4,054	4,945	6,811	8,862	9,907	10,029	7,693	3,331	859	11
1999.....	2,716	3,472	4,282	5,281	7,243	9,198	10,267	10,403	7,946	3,477	835	18

Non-Hispanic 1/, male

2003.....	2,627	2,945	3,191	3,238	3,645	4,258	4,570	3,603	2,019	670	102	5
2002.....	2,413	2,989	3,015	3,249	3,827	4,486	4,491	3,684	2,068	684	125	11
2001.....	2,336	2,751	3,019	3,202	3,891	4,622	4,578	3,483	2,096	589	102	10
2000.....	2,255	2,765	2,881	3,324	4,225	4,819	4,462	3,589	2,122	663	113	10
1999.....	2,118	2,635	3,028	3,538	4,384	4,924	4,580	3,767	2,111	659	117	16

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic 1/, female

2003.....	666	948	1,319	1,608	2,224	3,517	5,091	5,644	5,043	2,371	680	1
2002.....	646	899	1,206	1,545	2,371	3,703	5,091	6,023	5,225	2,558	715	6
2001.....	622	886	1,160	1,677	2,389	3,954	5,192	6,152	5,216	2,611	750	1
2000.....	612	808	1,173	1,621	2,586	4,043	5,445	6,440	5,571	2,668	746	1
1999.....	598	837	1,254	1,743	2,859	4,274	5,687	6,636	5,835	2,818	718	2

Non-Hispanic White, both sexes

2003.....	2,428	2,964	3,309	3,644	4,593	6,367	8,339	8,219	6,354	2,745	667	4
2002.....	2,245	2,892	3,151	3,551	4,893	6,693	8,279	8,663	6,572	2,908	708	10
2001.....	2,177	2,632	3,017	3,566	4,938	7,008	8,513	8,594	6,637	2,856	724	7
2000.....	2,133	2,606	2,931	3,747	5,332	7,331	8,540	8,891	6,948	3,017	735	8
1999.....	1,964	2,508	3,118	3,954	5,711	7,580	8,937	9,244	7,180	3,136	723	10

Non-Hispanic White, male

2003.....	1,981	2,291	2,389	2,479	2,859	3,534	3,956	3,225	1,821	593	83	3
2002.....	1,830	2,264	2,295	2,441	3,085	3,707	3,876	3,304	1,859	609	102	6
2001.....	1,767	2,059	2,232	2,397	3,099	3,824	4,025	3,132	1,891	522	80	6
2000.....	1,726	2,053	2,157	2,590	3,359	3,996	3,868	3,200	1,916	583	84	8
1999.....	1,574	1,960	2,248	2,715	3,512	4,093	4,004	3,358	1,884	583	103	8

Non-Hispanic White, female

2003.....	447	673	920	1,165	1,734	2,833	4,383	4,994	4,533	2,152	584	1
2002.....	415	628	856	1,110	1,808	2,986	4,403	5,359	4,713	2,299	606	4
2001.....	410	573	785	1,169	1,839	3,184	4,488	5,462	4,746	2,334	644	1
2000.....	407	553	774	1,157	1,973	3,335	4,672	5,691	5,032	2,434	651	-
1999.....	390	548	870	1,239	2,199	3,487	4,933	5,886	5,296	2,553	620	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Non-Hispanic Black, both sexes

2003.....	761	821	1,059	1,051	1,106	1,239	1,105	872	607	262	97	1
2002.....	730	880	964	1,107	1,151	1,328	1,138	892	620	290	117	7
2001.....	707	889	1,038	1,169	1,192	1,382	1,090	908	590	298	113	3
2000.....	669	870	1,007	1,068	1,312	1,329	1,193	1,002	659	277	113	3
1999.....	672	876	1,028	1,194	1,359	1,435	1,158	1,004	666	302	101	8

Non-Hispanic Black, male

2003.....	563	565	702	647	680	648	510	316	156	67	13	1
2002.....	508	633	645	712	645	684	539	302	169	63	18	5
2001.....	508	604	695	710	702	702	472	289	172	50	17	3
2000.....	478	634	648	638	760	704	500	339	174	67	26	2
1999.....	479	604	674	746	772	717	493	344	185	63	9	8

Non-Hispanic Black, female

2003.....	198	256	357	404	426	591	595	556	451	195	84	-
2002.....	222	247	319	395	506	644	599	590	451	227	99	2
2001.....	199	285	343	459	490	680	618	619	418	248	96	-
2000.....	191	236	359	430	552	625	693	663	485	210	87	1
1999.....	193	272	354	448	587	718	665	660	481	239	92	-

Origin not stated 2/, both sexes

2003.....	21	46	47	30	49	61	44	25	17	9	2	5
2002.....	40	43	55	42	61	65	38	36	22	7	2	5
2001.....	28	40	34	47	51	69	46	35	26	7	2	12
2000.....	31	38	55	40	64	78	49	37	27	4	2	6
1999.....	31	28	61	46	78	84	64	49	15	7	2	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Origin not stated 2/, male

2003.....	15	35	41	24	35	40	23	13	7	5	-	5
2002.....	34	41	43	28	38	41	23	18	6	2	-	5
2001.....	21	31	28	35	38	41	25	14	5	1	-	7
2000.....	29	31	42	32	50	52	24	18	8	-	-	4
1999.....	25	23	53	36	61	56	31	23	2	1	-	3

Origin not stated 2/, female

2003.....	6	11	6	6	14	21	21	12	10	4	2	-
2002.....	6	2	12	14	23	24	15	18	16	5	2	-
2001.....	7	9	6	12	13	28	21	21	21	6	2	5
2000.....	2	7	13	8	14	26	25	19	19	4	2	2
1999.....	6	5	8	10	17	28	33	26	13	6	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins, both sexes

2003.....	5,340	7,732	10,365	14,079	21,805	33,155	43,931	44,896	33,346	14,148	3,225	9
2002.....	5,608	7,840	10,236	14,654	22,762	34,546	44,739	45,200	33,321	14,074	3,308	9
2001.....	5,496	7,442	10,401	15,259	24,174	35,878	44,655	45,199	33,160	14,135	3,407	13
2000.....	5,330	7,634	10,364	15,756	25,459	36,799	44,232	46,132	33,297	14,157	3,396	20
1999.....	5,548	7,496	10,644	16,762	26,568	38,254	45,155	46,228	33,530	14,511	3,499	8

All origins, male

2003.....	4,007	5,616	7,207	9,353	13,573	19,104	22,372	18,779	10,366	3,042	470	5
2002.....	4,273	5,660	7,161	9,668	14,197	19,876	22,584	18,609	10,206	3,037	470	6
2001.....	4,116	5,411	7,172	10,072	14,976	20,640	22,093	18,260	9,685	2,907	480	8
2000.....	4,017	5,523	7,168	10,297	15,967	20,839	22,106	18,366	9,629	2,958	482	13
1999.....	4,152	5,366	7,290	10,997	16,526	21,621	22,355	18,037	9,635	3,011	515	7

All origins, female

2003.....	1,333	2,116	3,158	4,726	8,232	14,051	21,559	26,117	22,980	11,106	2,755	4
2002.....	1,335	2,180	3,075	4,986	8,565	14,670	22,155	26,591	23,115	11,037	2,838	3
2001.....	1,380	2,031	3,229	5,187	9,198	15,238	22,562	26,939	23,475	11,228	2,927	5
2000.....	1,313	2,111	3,196	5,459	9,492	15,960	22,126	27,766	23,668	11,199	2,914	7
1999.....	1,396	2,130	3,354	5,765	10,042	16,633	22,800	28,191	23,895	11,500	2,984	1

Hispanic, both sexes

2003.....	286	407	590	787	1,157	1,552	1,682	1,440	1,137	486	132	2
2002.....	321	419	577	860	1,188	1,501	1,616	1,444	1,108	542	101	-
2001.....	291	399	531	863	1,222	1,546	1,610	1,500	1,134	466	114	-
2000.....	274	419	565	833	1,148	1,430	1,481	1,469	1,062	467	101	2
1999.....	310	415	576	852	1,186	1,412	1,441	1,451	1,056	495	93	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Hispanic, male

2003.....	215	292	368	499	680	841	814	532	327	129	31	2
2002.....	232	291	391	539	708	823	773	572	368	144	20	-
2001.....	207	296	343	537	715	845	766	595	346	105	25	-
2000.....	208	294	378	525	703	763	701	555	342	118	19	2
1999.....	245	306	374	511	700	747	670	568	306	138	18	2

Hispanic, female

2003.....	71	115	222	288	477	711	868	908	810	357	101	-
2002.....	89	128	186	321	480	678	843	872	740	398	81	-
2001.....	84	103	188	326	507	701	844	905	788	361	89	-
2000.....	66	125	187	308	445	667	780	914	720	349	82	-
1999.....	65	109	202	341	486	665	771	883	750	357	75	-

Non-Hispanic 1/, both sexes

2003.....	5,035	7,291	9,728	13,237	20,579	31,511	42,134	43,356	32,134	13,626	3,083	7
2002.....	5,261	7,388	9,614	13,721	21,485	32,910	42,985	43,589	32,108	13,486	3,192	7
2001.....	5,180	7,012	9,824	14,341	22,875	34,219	42,896	43,568	31,942	13,612	3,283	12
2000.....	5,031	7,186	9,736	14,855	24,205	35,241	42,606	44,494	32,142	13,635	3,287	14
1999.....	5,213	7,052	10,018	15,844	25,273	36,708	43,555	44,626	32,355	13,953	3,392	4

Non-Hispanic 1/, male

2003.....	3,775	5,296	6,811	8,813	12,850	18,209	21,503	18,205	10,019	2,905	436	3
2002.....	4,017	5,343	6,740	9,077	13,422	18,973	21,743	17,972	9,805	2,884	447	4
2001.....	3,889	5,088	6,790	9,497	14,216	19,726	21,251	17,614	9,322	2,789	452	7
2000.....	3,788	5,206	6,742	9,721	15,195	20,005	21,340	17,750	9,259	2,823	461	8
1999.....	3,888	5,037	6,879	10,436	15,755	20,800	21,624	17,401	9,294	2,859	495	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic 1/, female

2003.....	1,260	1,995	2,917	4,424	7,729	13,302	20,631	25,151	22,115	10,721	2,647	4
2002.....	1,244	2,045	2,874	4,644	8,063	13,937	21,242	25,617	22,303	10,602	2,745	3
2001.....	1,291	1,924	3,034	4,844	8,659	14,493	21,645	25,954	22,620	10,823	2,831	5
2000.....	1,243	1,980	2,994	5,134	9,010	15,236	21,266	26,744	22,883	10,812	2,826	6
1999.....	1,325	2,015	3,139	5,408	9,518	15,908	21,931	27,225	23,061	11,094	2,897	1

Non-Hispanic White, both sexes

2003.....	3,909	5,883	7,925	10,952	17,667	27,909	38,287	40,094	29,796	12,623	2,710	6
2002.....	4,123	5,965	7,811	11,455	18,677	29,228	39,249	40,435	29,850	12,452	2,845	6
2001.....	4,104	5,719	8,034	11,994	19,923	30,491	39,138	40,390	29,754	12,560	2,914	7
2000.....	4,031	5,855	7,940	12,532	21,174	31,527	39,099	41,296	30,037	12,644	2,972	10
1999.....	4,155	5,715	8,265	13,444	22,128	32,995	40,057	41,395	30,240	12,925	3,077	3

Non-Hispanic White, male

2003.....	3,036	4,399	5,723	7,479	11,275	16,433	19,776	17,038	9,400	2,678	378	2
2002.....	3,260	4,447	5,585	7,795	11,898	17,210	20,151	16,869	9,149	2,647	390	4
2001.....	3,170	4,228	5,714	8,164	12,612	17,901	19,630	16,531	8,705	2,566	392	4
2000.....	3,121	4,352	5,656	8,445	13,604	18,208	19,839	16,638	8,677	2,594	401	6
1999.....	3,169	4,199	5,788	9,127	14,069	19,014	20,182	16,264	8,644	2,631	442	3

Non-Hispanic White, female

2003.....	873	1,484	2,202	3,473	6,392	11,476	18,511	23,056	20,396	9,945	2,332	4
2002.....	863	1,518	2,226	3,660	6,779	12,018	19,098	23,566	20,701	9,805	2,455	2
2001.....	934	1,491	2,320	3,830	7,311	12,590	19,508	23,859	21,049	9,994	2,522	3
2000.....	910	1,503	2,284	4,087	7,570	13,319	19,260	24,658	21,360	10,050	2,571	4
1999.....	986	1,516	2,477	4,317	8,059	13,981	19,875	25,131	21,596	10,294	2,635	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Non-Hispanic Black, both sexes

2003.....	1,000	1,224	1,555	1,929	2,469	2,973	3,122	2,572	1,859	820	323	-
2002.....	995	1,240	1,542	1,900	2,347	2,991	3,056	2,544	1,838	831	296	1
2001.....	953	1,119	1,555	1,998	2,480	3,091	3,101	2,595	1,823	903	322	4
2000.....	858	1,156	1,541	1,987	2,612	3,098	2,903	2,647	1,734	836	273	4
1999.....	935	1,179	1,514	2,087	2,654	3,071	2,873	2,666	1,733	893	280	1

Non-Hispanic Black, male

2003.....	638	763	912	1,107	1,311	1,436	1,340	842	437	159	40	-
2002.....	647	768	975	1,057	1,247	1,360	1,236	822	476	167	44	-
2001.....	632	711	918	1,114	1,341	1,459	1,292	821	484	174	52	3
2000.....	548	734	899	1,058	1,328	1,433	1,188	847	425	170	50	2
1999.....	621	724	929	1,111	1,377	1,406	1,123	859	466	176	46	-

Non-Hispanic Black, female

2003.....	362	461	643	822	1,158	1,537	1,782	1,730	1,422	661	283	-
2002.....	348	472	567	843	1,100	1,631	1,820	1,722	1,362	664	252	1
2001.....	321	408	637	884	1,139	1,632	1,809	1,774	1,339	729	270	1
2000.....	310	422	642	929	1,284	1,665	1,715	1,800	1,309	666	223	2
1999.....	314	455	585	976	1,277	1,665	1,750	1,807	1,267	717	234	1

Origin not stated 2/, both sexes

2003.....	19	34	47	55	69	92	115	100	75	36	10	-
2002.....	26	33	45	73	89	135	138	167	105	46	15	2
2001.....	25	31	46	55	77	113	149	131	84	57	10	1
2000.....	25	29	63	68	106	128	145	169	93	55	8	4
1999.....	25	29	50	66	109	134	159	151	119	63	14	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Origin not stated 2/, male

2003.....	17	28	28	41	43	54	55	42	20	8	3	-
2002.....	24	26	30	52	67	80	68	65	33	9	3	2
2001.....	20	27	39	38	45	69	76	51	17	13	3	1
2000.....	21	23	48	51	69	71	65	61	28	17	2	3
1999.....	19	23	37	50	71	74	61	68	35	14	2	2

Origin not stated 2/, female

2003.....	2	6	19	14	26	38	60	58	55	28	7	-
2002.....	2	7	15	21	22	55	70	102	72	37	12	-
2001.....	5	4	7	17	32	44	73	80	67	44	7	-
2000.....	4	6	15	17	37	57	80	108	65	38	6	1
1999.....	6	6	13	16	38	60	98	83	84	49	12	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

All origins, both sexes

2003.....	4,825	6,171	7,227	9,481	13,763	20,800	28,460	31,015	23,947	9,993	2,315	6
2002.....	4,862	5,913	7,033	9,377	14,303	21,276	28,500	30,794	23,517	9,779	2,210	6
2001.....	4,776	5,637	6,898	9,596	14,460	21,325	27,651	30,343	22,779	9,653	2,204	7
2000.....	4,551	5,417	6,945	9,875	14,804	21,524	26,843	29,519	22,677	9,591	2,206	8
1999.....	4,386	5,465	7,115	10,034	15,538	21,689	27,023	29,428	21,894	9,401	2,125	7

All origins, male

2003.....	3,019	3,826	4,316	5,402	7,473	10,351	12,329	11,327	6,827	2,169	358	5
2002.....	3,082	3,587	4,184	5,275	7,716	10,463	12,308	11,167	6,558	2,070	338	3
2001.....	2,946	3,393	4,110	5,356	7,811	10,538	12,002	10,595	6,187	2,028	331	5
2000.....	2,873	3,384	4,148	5,584	8,037	10,505	11,501	10,488	6,178	1,983	359	5
1999.....	2,756	3,419	4,281	5,870	8,450	10,567	11,561	10,293	5,879	1,988	323	5

All origins, female

2003.....	1,806	2,345	2,911	4,079	6,290	10,449	16,131	19,688	17,120	7,824	1,957	1
2002.....	1,780	2,326	2,849	4,102	6,587	10,813	16,192	19,627	16,959	7,709	1,872	3
2001.....	1,830	2,244	2,788	4,240	6,649	10,787	15,649	19,748	16,592	7,625	1,873	2
2000.....	1,678	2,033	2,797	4,291	6,767	11,019	15,342	19,031	16,499	7,608	1,847	3
1999.....	1,630	2,046	2,834	4,164	7,088	11,122	15,462	19,135	16,015	7,413	1,802	2

Hispanic, both sexes

2003.....	211	292	329	419	526	694	723	640	548	193	66	-
2002.....	225	244	263	377	480	628	690	599	463	220	53	-
2001.....	199	224	292	389	467	594	612	583	450	198	46	2
2000.....	212	224	269	349	468	584	565	597	438	176	39	1
1999.....	192	215	286	356	459	536	566	554	403	186	37	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

Hispanic, male

2003.....	128	171	196	232	283	333	328	232	185	57	15	-
2002.....	143	145	151	206	278	316	310	233	147	64	9	-
2001.....	132	127	169	213	243	279	277	215	139	47	6	2
2000.....	139	143	156	190	256	264	246	221	156	56	9	1
1999.....	128	126	172	206	260	273	240	216	139	45	12	1

Hispanic, female

2003.....	83	121	133	187	243	361	395	408	363	136	51	-
2002.....	82	99	112	171	202	312	380	366	316	156	44	-
2001.....	67	97	123	176	224	315	335	368	311	151	40	-
2000.....	73	81	113	159	212	320	319	376	282	120	30	-
1999.....	64	89	114	150	199	263	326	338	264	141	25	-

Non-Hispanic 1/, both sexes

2003.....	4,594	5,859	6,872	9,041	13,202	20,071	27,679	30,326	23,356	9,784	2,248	4
2002.....	4,617	5,650	6,752	8,979	13,782	20,603	27,739	30,132	23,012	9,536	2,150	3
2001.....	4,561	5,394	6,586	9,180	13,957	20,679	26,982	29,694	22,285	9,432	2,153	3
2000.....	4,327	5,172	6,651	9,508	14,296	20,893	26,231	28,859	22,182	9,393	2,163	6
1999.....	4,183	5,229	6,801	9,652	15,033	21,099	26,397	28,800	21,445	9,203	2,084	5

Non-Hispanic 1/, male

2003.....	2,877	3,641	4,098	5,161	7,171	10,001	11,968	11,078	6,625	2,109	343	3
2002.....	2,925	3,430	4,024	5,056	7,416	10,122	11,967	10,917	6,403	2,001	328	2
2001.....	2,803	3,258	3,923	5,127	7,544	10,230	11,711	10,358	6,028	1,977	324	2
2000.....	2,728	3,229	3,974	5,380	7,758	10,222	11,238	10,245	6,011	1,921	350	3
1999.....	2,619	3,279	4,091	5,650	8,162	10,263	11,292	10,043	5,728	1,939	311	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Non-Hispanic 1/, female												
2003.....	1,717	2,218	2,774	3,880	6,031	10,070	15,711	19,248	16,731	7,675	1,905	1
2002.....	1,692	2,220	2,728	3,923	6,366	10,481	15,772	19,215	16,609	7,535	1,822	1
2001.....	1,758	2,136	2,663	4,053	6,413	10,449	15,271	19,336	16,257	7,455	1,829	1
2000.....	1,599	1,943	2,677	4,128	6,538	10,671	14,993	18,614	16,171	7,472	1,813	3
1999.....	1,564	1,950	2,710	4,002	6,871	10,836	15,105	18,757	15,717	7,264	1,773	2
Non-Hispanic White, both sexes												
2003.....	3,185	4,388	5,261	7,172	10,965	17,606	25,014	27,990	21,683	9,040	1,994	3
2002.....	3,250	4,234	5,234	7,235	11,601	18,032	25,108	27,835	21,321	8,790	1,890	2
2001.....	3,249	3,986	5,146	7,348	11,823	18,059	24,427	27,403	20,729	8,728	1,916	3
2000.....	3,077	3,908	5,181	7,694	12,087	18,323	23,851	26,594	20,614	8,654	1,932	5
1999.....	3,000	3,857	5,251	7,809	12,720	18,469	23,937	26,462	20,015	8,501	1,859	3
Non-Hispanic White, male												
2003.....	2,036	2,758	3,212	4,154	6,080	8,886	10,887	10,281	6,160	1,938	290	2
2002.....	2,125	2,618	3,188	4,135	6,330	8,960	10,918	10,148	5,895	1,841	283	2
2001.....	2,075	2,444	3,118	4,199	6,454	9,008	10,619	9,571	5,578	1,810	280	2
2000.....	1,966	2,489	3,159	4,422	6,680	9,052	10,283	9,453	5,566	1,735	300	3
1999.....	1,913	2,469	3,211	4,641	7,015	9,074	10,291	9,187	5,298	1,762	264	2
Non-Hispanic White, female												
2003.....	1,149	1,630	2,049	3,018	4,885	8,720	14,127	17,709	15,523	7,102	1,704	1
2002.....	1,125	1,616	2,046	3,100	5,271	9,072	14,190	17,687	15,426	6,949	1,607	-
2001.....	1,174	1,542	2,028	3,149	5,369	9,051	13,808	17,832	15,151	6,918	1,636	1
2000.....	1,111	1,419	2,022	3,272	5,407	9,271	13,568	17,141	15,048	6,919	1,632	2
1999.....	1,087	1,388	2,040	3,168	5,705	9,395	13,646	17,275	14,717	6,739	1,595	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>Other heart diseases (I26-I51)</b>												
<b>Non-Hispanic Black, both sexes</b>												
2003.....	1,306	1,342	1,453	1,670	1,989	2,129	2,246	1,959	1,387	651	216	1
2002.....	1,265	1,287	1,384	1,581	1,895	2,228	2,255	1,930	1,449	648	223	1
2001.....	1,214	1,291	1,317	1,648	1,874	2,304	2,204	1,986	1,346	605	205	-
2000.....	1,155	1,152	1,333	1,636	1,980	2,278	2,080	1,973	1,338	629	209	1
1999.....	1,091	1,257	1,421	1,668	2,079	2,316	2,138	2,037	1,224	613	190	2
<b>Non-Hispanic Black, male</b>												
2003.....	776	806	804	905	958	946	892	635	348	138	42	1
2002.....	735	736	754	828	944	975	879	596	398	121	35	-
2001.....	674	746	737	830	936	1,057	921	660	346	135	39	-
2000.....	705	676	734	860	942	1,012	821	664	353	146	43	-
1999.....	658	739	806	909	1,027	1,016	854	699	349	146	40	1
<b>Non-Hispanic Black, female</b>												
2003.....	530	536	649	765	1,031	1,183	1,354	1,324	1,039	513	174	-
2002.....	530	551	630	753	951	1,253	1,376	1,334	1,051	527	188	1
2001.....	540	545	580	818	938	1,247	1,283	1,326	1,000	470	166	-
2000.....	450	476	599	776	1,038	1,266	1,259	1,309	985	483	166	1
1999.....	433	518	615	759	1,052	1,300	1,284	1,338	875	467	150	1
<b>Origin not stated 2/, both sexes</b>												
2003.....	20	20	26	21	35	35	58	49	43	16	1	2
2002.....	20	19	18	21	41	45	71	63	42	23	7	3
2001.....	16	19	20	27	36	52	57	66	44	23	5	2
2000.....	12	21	25	18	40	47	47	63	57	22	4	1
1999.....	11	21	28	26	46	54	60	74	46	12	4	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other heart diseases (I26-I51)

Origin not stated 2/, male

2003.....	14	14	22	9	19	17	33	17	17	3	-	2
2002.....	14	12	9	13	22	25	31	17	8	5	1	1
2001.....	11	8	18	16	24	29	14	22	20	4	1	1
2000.....	6	12	18	14	23	19	17	22	11	6	-	1
1999.....	9	14	18	14	28	31	29	34	12	4	-	1

Origin not stated 2/, female

2003.....	6	6	4	12	16	18	25	32	26	13	1	-
2002.....	6	7	9	8	19	20	40	46	34	18	6	2
2001.....	5	11	2	11	12	23	43	44	24	19	4	1
2000.....	6	9	7	4	17	28	30	41	46	16	4	-
1999.....	2	7	10	12	18	23	31	40	34	8	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

All origins, both sexes

2003.....	94	95	114	102	130	172	146	104	32	3	1	-
2002.....	90	73	95	107	131	161	145	76	36	8	1	-
2001.....	93	72	88	102	145	162	154	79	30	2	-	-
2000.....	97	88	77	105	119	166	136	83	30	5	1	-
1999.....	75	93	78	103	137	155	134	92	21	5	1	-

All origins, male

2003.....	65	60	63	57	62	89	77	43	15	1	-	-
2002.....	62	42	49	66	67	78	75	31	15	2	1	-
2001.....	61	39	56	49	83	94	83	37	11	1	-	-
2000.....	61	52	44	59	63	95	74	40	9	2	-	-
1999.....	54	62	41	51	67	83	71	46	10	-	-	-

All origins, female

2003.....	29	35	51	45	68	83	69	61	17	2	1	-
2002.....	28	31	46	41	64	83	70	45	21	6	-	-
2001.....	32	33	32	53	62	68	71	42	19	1	-	-
2000.....	36	36	33	46	56	71	62	43	21	3	1	-
1999.....	21	31	37	52	70	72	63	46	11	5	1	-

Hispanic, both sexes

2003.....	8	8	4	13	3	7	6	2	1	-	-	-
2002.....	4	6	6	7	7	7	6	1	1	1	-	-
2001.....	5	3	6	3	6	7	3	1	1	-	-	-
2000.....	12	4	4	8	10	6	1	1	1	-	-	-
1999.....	4	6	4	6	7	9	3	2	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

Hispanic, male

2003.....	4	5	1	9	-	2	1	1	1	-	-	-
2002.....	4	4	3	4	5	3	3	1	-	-	-	-
2001.....	4	-	4	1	4	4	-	-	1	-	-	-
2000.....	8	3	1	5	6	1	1	-	1	-	-	-
1999.....	4	4	1	3	6	6	2	2	-	-	-	-

Hispanic, female

2003.....	4	3	3	4	3	5	5	1	-	-	-	-
2002.....	-	2	3	3	2	4	3	-	1	1	-	-
2001.....	1	3	2	2	2	3	3	1	-	-	-	-
2000.....	4	1	3	3	4	5	-	1	-	-	-	-
1999.....	-	2	3	3	1	3	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	85	87	110	89	127	165	140	102	31	3	1	-
2002.....	85	66	89	100	124	154	139	75	35	7	1	-
2001.....	87	68	82	99	139	154	151	78	29	2	-	-
2000.....	85	84	73	97	108	159	134	82	29	5	1	-
1999.....	71	86	74	97	129	146	131	90	21	5	1	-

Non-Hispanic 1/, male

2003.....	60	55	62	48	62	87	76	42	14	1	-	-
2002.....	57	37	46	62	62	75	72	30	15	2	1	-
2001.....	57	38	52	48	79	90	83	37	10	1	-	-
2000.....	53	49	43	54	57	94	73	40	8	2	-	-
1999.....	50	57	40	48	61	77	69	44	10	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

Non-Hispanic 1/, female

2003.....	25	32	48	41	65	78	64	60	17	2	1	-
2002.....	28	29	43	38	62	79	67	45	20	5	-	-
2001.....	30	30	30	51	60	64	68	41	19	1	-	-
2000.....	32	35	30	43	51	65	61	42	21	3	1	-
1999.....	21	29	34	49	68	69	62	46	11	5	1	-

Non-Hispanic White, both sexes

2003.....	52	58	79	71	101	143	128	96	31	3	1	-
2002.....	56	40	73	76	103	138	132	69	32	7	1	-
2001.....	55	49	58	71	111	128	137	75	26	2	-	-
2000.....	53	55	59	81	92	141	120	73	25	5	-	-
1999.....	44	51	56	76	106	133	119	85	20	5	-	-

Non-Hispanic White, male

2003.....	36	35	50	39	50	77	71	41	14	1	-	-
2002.....	41	21	40	49	53	71	68	27	14	2	1	-
2001.....	38	27	38	34	59	79	75	35	9	1	-	-
2000.....	36	35	37	42	51	86	65	37	7	2	-	-
1999.....	29	37	31	42	51	75	64	43	10	-	-	-

Non-Hispanic White, female

2003.....	16	23	29	32	51	66	57	55	17	2	1	-
2002.....	15	19	33	27	50	67	64	42	18	5	-	-
2001.....	17	22	20	37	52	49	62	40	17	1	-	-
2000.....	17	20	22	39	41	55	55	36	18	3	-	-
1999.....	15	14	25	34	55	58	55	42	10	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

Non-Hispanic Black, both sexes

2003.....	32	27	26	16	21	19	10	4	-	-	-	-
2002.....	28	25	16	22	16	14	5	5	3	-	-	-
2001.....	30	19	22	21	22	25	12	2	3	-	-	-
2000.....	27	26	13	15	15	16	10	6	4	-	1	-
1999.....	23	30	17	21	21	10	7	5	1	-	-	-

Non-Hispanic Black, male

2003.....	23	18	9	8	9	8	5	-	-	-	-	-
2002.....	15	16	6	11	8	3	3	3	1	-	-	-
2001.....	18	11	13	10	16	11	7	1	1	-	-	-
2000.....	15	12	6	11	5	6	5	1	1	-	-	-
1999.....	19	18	8	6	9	1	4	1	-	-	-	-

Non-Hispanic Black, female

2003.....	9	9	17	8	12	11	5	4	-	-	-	-
2002.....	13	9	10	11	8	11	2	2	2	-	-	-
2001.....	12	8	9	11	6	14	5	1	2	-	-	-
2000.....	12	14	7	4	10	10	5	5	3	-	1	-
1999.....	4	12	9	15	12	9	3	4	1	-	-	-

Origin not stated 2/, both sexes

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	1	-	-	-	-	-
1999.....	-	1	-	-	1	-	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and subacute endocarditis (I33)

Origin not stated 2/, male

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	-	-	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All origins, both sexes

2003.....	48	59	45	80	71	70	83	45	27	5	-	-
2002.....	51	66	52	55	88	84	73	65	25	6	-	-
2001.....	54	54	52	47	73	87	67	49	28	7	1	1
2000.....	40	47	42	52	79	82	63	43	24	8	1	-
1999.....	46	63	62	69	77	88	72	56	25	4	-	-

All origins, male

2003.....	30	32	25	44	40	34	31	13	8	1	-	-
2002.....	29	34	35	29	42	34	26	25	6	2	-	-
2001.....	29	28	33	28	31	39	22	14	8	2	-	1
2000.....	18	31	21	24	43	42	31	17	7	1	1	-
1999.....	25	40	30	34	35	38	27	19	11	-	-	-

All origins, female

2003.....	18	27	20	36	31	36	52	32	19	4	-	-
2002.....	22	32	17	26	46	50	47	40	19	4	-	-
2001.....	25	26	19	19	42	48	45	35	20	5	1	-
2000.....	22	16	21	28	36	40	32	26	17	7	-	-
1999.....	21	23	32	35	42	50	45	37	14	4	-	-

Hispanic, both sexes

2003.....	-	4	3	6	1	2	3	-	-	-	-	-
2002.....	4	7	7	3	4	3	2	5	-	-	-	-
2001.....	3	3	3	3	3	4	3	3	2	-	-	-
2000.....	2	5	4	3	2	3	4	1	-	1	-	-
1999.....	1	1	1	3	4	4	4	1	-	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Hispanic, male

2003.....	-	1	3	5	1	1	-	-	-	-	-	-
2002.....	3	5	4	2	1	-	1	3	-	-	-	-
2001.....	-	-	3	1	2	2	-	1	-	-	-	-
2000.....	1	2	1	-	-	2	4	-	-	1	-	-
1999.....	1	1	-	-	1	1	1	-	-	-	-	-

Hispanic, female

2003.....	-	3	-	1	-	1	3	-	-	-	-	-
2002.....	1	2	3	1	3	3	1	2	-	-	-	-
2001.....	3	3	-	2	1	2	3	2	2	-	-	-
2000.....	1	3	3	3	2	1	-	1	-	-	-	-
1999.....	-	-	1	3	3	3	3	1	-	1	-	-

Non-Hispanic 1/, both sexes

2003.....	45	55	42	74	70	68	80	44	27	5	-	-
2002.....	47	58	45	52	84	81	71	60	25	6	-	-
2001.....	51	51	49	44	70	83	64	46	26	7	1	1
2000.....	38	42	36	49	77	77	59	42	24	7	1	-
1999.....	45	62	61	66	73	84	68	55	25	3	-	-

Non-Hispanic 1/, male

2003.....	28	31	22	39	39	33	31	13	8	1	-	-
2002.....	26	28	31	27	41	34	25	22	6	2	-	-
2001.....	29	28	30	27	29	37	22	13	8	2	-	1
2000.....	17	29	19	24	43	40	27	17	7	-	1	-
1999.....	24	39	30	34	34	37	26	19	11	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

Non-Hispanic 1/, female

2003.....	17	24	20	35	31	35	49	31	19	4	-	-
2002.....	21	30	14	25	43	47	46	38	19	4	-	-
2001.....	22	23	19	17	41	46	42	33	18	5	1	-
2000.....	21	13	17	25	34	37	32	25	17	7	-	-
1999.....	21	23	31	32	39	47	42	36	14	3	-	-

Non-Hispanic White, both sexes

2003.....	35	47	36	65	61	61	70	40	26	5	-	-
2002.....	32	41	35	45	71	64	62	57	23	5	-	-
2001.....	35	39	40	33	65	76	58	40	25	7	-	1
2000.....	31	34	28	38	65	65	57	37	21	7	1	-
1999.....	34	52	52	54	61	73	61	49	22	3	-	-

Non-Hispanic White, male

2003.....	23	25	20	36	35	30	30	13	8	1	-	-
2002.....	18	20	25	23	37	30	21	22	5	1	-	-
2001.....	21	22	23	19	29	33	20	12	8	2	-	1
2000.....	14	26	15	15	37	32	27	15	6	-	1	-
1999.....	17	33	27	27	29	34	24	17	10	-	-	-

Non-Hispanic White, female

2003.....	12	22	16	29	26	31	40	27	18	4	-	-
2002.....	14	21	10	22	34	34	41	35	18	4	-	-
2001.....	14	17	17	14	36	43	38	28	17	5	-	-
2000.....	17	8	13	23	28	33	30	22	15	7	-	-
1999.....	17	19	25	27	32	39	37	32	12	3	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>Heart failure (I50)</b>												
<b>All origins, both sexes</b>												
2003.....	623	1,007	1,462	2,302	3,836	6,697	10,439	12,844	11,082	5,071	1,222	-
2002.....	638	906	1,436	2,192	3,976	6,603	10,307	12,620	10,776	5,031	1,200	-
2001.....	601	926	1,448	2,270	4,004	6,879	10,255	12,793	10,735	4,971	1,209	1
2000.....	528	828	1,383	2,284	4,119	6,936	9,884	12,315	10,543	4,969	1,208	4
1999.....	484	853	1,254	2,240	4,284	6,926	9,845	12,169	10,197	4,850	1,111	1
<b>All origins, male</b>												
2003.....	364	605	849	1,281	2,032	3,262	4,396	4,668	3,115	1,111	206	-
2002.....	407	526	819	1,200	2,054	3,192	4,381	4,423	3,000	1,035	178	-
2001.....	361	556	791	1,235	2,109	3,300	4,372	4,320	2,886	1,020	172	-
2000.....	305	479	785	1,243	2,171	3,286	4,212	4,256	2,819	982	191	2
1999.....	300	506	722	1,232	2,294	3,264	4,054	4,155	2,636	991	177	1
<b>All origins, female</b>												
2003.....	259	402	613	1,021	1,804	3,435	6,043	8,176	7,967	3,960	1,016	-
2002.....	231	380	617	992	1,922	3,411	5,926	8,197	7,776	3,996	1,022	-
2001.....	240	370	657	1,035	1,895	3,579	5,883	8,473	7,849	3,951	1,037	1
2000.....	223	349	598	1,041	1,948	3,650	5,672	8,059	7,724	3,987	1,017	2
1999.....	184	347	532	1,008	1,990	3,662	5,791	8,014	7,561	3,859	934	-
<b>Hispanic, both sexes</b>												
2003.....	26	50	68	115	160	221	273	259	244	99	35	-
2002.....	29	35	56	93	142	200	251	232	194	113	31	-
2001.....	25	31	72	90	123	178	226	231	214	93	21	-
2000.....	15	38	62	73	130	171	194	243	219	74	21	-
1999.....	19	32	55	71	131	167	186	232	201	92	13	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Heart failure (I50)

## Hispanic, male

2003.....	10	31	46	56	85	113	129	85	86	25	7	-
2002.....	16	22	32	51	70	100	109	85	56	31	5	-
2001.....	16	17	35	55	54	69	100	86	61	26	2	-
2000.....	6	24	38	44	65	71	80	86	76	21	6	-
1999.....	10	19	28	40	72	87	72	78	66	20	2	-

## Hispanic, female

2003.....	16	19	22	59	75	108	144	174	158	74	28	-
2002.....	13	13	24	42	72	100	142	147	138	82	26	-
2001.....	9	14	37	35	69	109	126	145	153	67	19	-
2000.....	9	14	24	29	65	100	114	157	143	53	15	-
1999.....	9	13	27	31	59	80	114	154	135	72	11	-

## Non-Hispanic 1/, both sexes

2003.....	597	956	1,389	2,180	3,663	6,470	10,144	12,566	10,813	4,961	1,187	-
2002.....	604	869	1,376	2,095	3,819	6,388	10,028	12,353	10,561	4,907	1,163	-
2001.....	575	887	1,371	2,170	3,874	6,682	10,006	12,538	10,506	4,866	1,184	1
2000.....	513	789	1,316	2,206	3,979	6,750	9,667	12,050	10,290	4,886	1,185	3
1999.....	464	818	1,189	2,164	4,144	6,740	9,637	11,902	9,969	4,750	1,097	1

## Non-Hispanic 1/, male

2003.....	354	573	798	1,222	1,939	3,146	4,254	4,575	3,018	1,084	199	-
2002.....	388	502	786	1,147	1,977	3,088	4,256	4,328	2,939	1,003	172	-
2001.....	345	536	751	1,176	2,049	3,221	4,268	4,226	2,817	991	169	-
2000.....	299	455	742	1,195	2,101	3,210	4,125	4,164	2,736	960	185	1
1999.....	290	485	688	1,189	2,217	3,168	3,972	4,058	2,564	969	175	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Non-Hispanic 1/, female												
2003.....	243	383	591	958	1,724	3,324	5,890	7,991	7,795	3,877	988	-
2002.....	216	367	590	948	1,842	3,300	5,772	8,025	7,622	3,904	991	-
2001.....	230	351	620	994	1,825	3,461	5,738	8,312	7,689	3,875	1,015	1
2000.....	214	334	574	1,011	1,878	3,540	5,542	7,886	7,554	3,926	1,000	2
1999.....	174	333	501	975	1,927	3,572	5,665	7,844	7,405	3,781	922	-
Non-Hispanic White, both sexes												
2003.....	381	702	1,041	1,753	3,079	5,704	9,228	11,662	10,104	4,615	1,059	-
2002.....	397	644	1,075	1,695	3,219	5,618	9,107	11,477	9,844	4,600	1,041	-
2001.....	393	656	1,074	1,751	3,301	5,918	9,115	11,652	9,836	4,541	1,068	1
2000.....	363	588	1,023	1,795	3,381	5,998	8,829	11,171	9,599	4,547	1,075	3
1999.....	307	594	943	1,751	3,551	5,924	8,778	10,984	9,358	4,391	1,005	-
Non-Hispanic White, male												
2003.....	230	427	606	1,000	1,672	2,829	3,896	4,267	2,844	1,006	167	-
2002.....	262	381	619	933	1,687	2,752	3,911	4,044	2,739	945	152	-
2001.....	243	410	609	963	1,771	2,886	3,899	3,945	2,635	919	149	-
2000.....	218	359	583	1,003	1,825	2,884	3,787	3,870	2,536	883	160	1
1999.....	196	369	553	969	1,951	2,828	3,644	3,725	2,388	885	151	-
Non-Hispanic White, female												
2003.....	151	275	435	753	1,407	2,875	5,332	7,395	7,260	3,609	892	-
2002.....	135	263	456	762	1,532	2,866	5,196	7,433	7,105	3,655	889	-
2001.....	150	246	465	788	1,530	3,032	5,216	7,707	7,201	3,622	919	1
2000.....	145	229	440	792	1,556	3,114	5,042	7,301	7,063	3,664	915	2
1999.....	111	225	390	782	1,600	3,096	5,134	7,259	6,970	3,506	854	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Heart failure (I50)

## Non-Hispanic Black, both sexes

2003.....	204	239	324	392	529	671	795	786	612	306	107	-
2002.....	192	204	283	361	530	680	815	767	620	272	108	-
2001.....	173	221	273	380	525	690	785	784	588	295	99	-
2000.....	143	189	264	371	538	670	737	781	603	290	102	-
1999.....	148	212	234	380	542	724	744	817	514	315	78	1

## Non-Hispanic Black, male

2003.....	115	138	182	207	238	272	303	259	141	63	26	-
2002.....	115	108	158	190	260	294	308	227	156	47	17	-
2001.....	96	120	128	194	253	299	319	245	147	63	19	-
2000.....	78	90	141	174	246	285	287	254	166	60	22	-
1999.....	89	111	127	198	244	304	273	284	141	69	19	1

## Non-Hispanic Black, female

2003.....	89	101	142	185	291	399	492	527	471	243	81	-
2002.....	77	96	125	171	270	386	507	540	464	225	91	-
2001.....	77	101	145	186	272	391	466	539	441	232	80	-
2000.....	65	99	123	197	292	385	450	527	437	230	80	-
1999.....	59	101	107	182	298	420	471	533	373	246	59	-

## Origin not stated 2/, both sexes

2003.....	-	1	5	7	13	6	22	19	25	11	-	-
2002.....	5	2	4	4	15	15	28	35	21	11	6	-
2001.....	1	8	5	10	7	19	23	24	15	12	4	-
2000.....	-	1	5	5	10	15	23	22	34	9	2	1
1999.....	1	3	10	5	9	19	22	35	27	8	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Heart failure (I50)

## Origin not stated 2/, male

2003.....	-	1	5	3	8	3	13	8	11	2	-	-
2002.....	3	2	1	2	7	4	16	10	5	1	1	-
2001.....	-	3	5	4	6	10	4	8	8	3	1	-
2000.....	-	-	5	4	5	5	7	6	7	1	-	1
1999.....	-	2	6	3	5	9	10	19	6	2	-	-

## Origin not stated 2/, female

2003.....	-	-	-	4	5	3	9	11	14	9	-	-
2002.....	2	-	3	2	8	11	12	25	16	10	5	-
2001.....	1	5	-	6	1	9	19	16	7	9	3	-
2000.....	-	1	-	1	5	10	16	16	27	8	2	-
1999.....	1	1	4	2	4	10	12	16	21	6	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins, both sexes

2003.....	4,060	5,010	5,606	6,997	9,726	13,861	17,792	18,022	12,806	4,914	1,092	6
2002.....	4,083	4,868	5,450	7,023	10,108	14,428	17,975	18,033	12,680	4,734	1,009	6
2001.....	4,028	4,585	5,310	7,177	10,238	14,197	17,175	17,422	11,986	4,673	994	5
2000.....	3,886	4,454	5,443	7,434	10,487	14,340	16,760	17,078	12,080	4,609	996	4
1999.....	3,781	4,456	5,721	7,622	11,040	14,520	16,972	17,111	11,651	4,542	1,013	6

All origins, male

2003.....	2,560	3,129	3,379	4,020	5,339	6,966	7,825	6,603	3,689	1,056	152	5
2002.....	2,584	2,985	3,281	3,980	5,553	7,159	7,826	6,688	3,537	1,031	159	3
2001.....	2,495	2,770	3,230	4,044	5,588	7,105	7,525	6,224	3,282	1,005	159	4
2000.....	2,489	2,822	3,298	4,258	5,760	7,082	7,184	6,175	3,343	998	167	3
1999.....	2,377	2,811	3,488	4,553	6,054	7,182	7,409	6,073	3,222	997	146	4

All origins, female

2003.....	1,500	1,881	2,227	2,977	4,387	6,895	9,967	11,419	9,117	3,858	940	1
2002.....	1,499	1,883	2,169	3,043	4,555	7,269	10,149	11,345	9,143	3,703	850	3
2001.....	1,533	1,815	2,080	3,133	4,650	7,092	9,650	11,198	8,704	3,668	835	1
2000.....	1,397	1,632	2,145	3,176	4,727	7,258	9,576	10,903	8,737	3,611	829	1
1999.....	1,404	1,645	2,233	3,069	4,986	7,338	9,563	11,038	8,429	3,545	867	2

Hispanic, both sexes

2003.....	177	230	254	285	362	464	441	379	303	94	31	-
2002.....	188	196	194	274	327	418	431	361	268	106	22	-
2001.....	166	187	211	293	335	405	380	348	233	105	25	2
2000.....	183	177	199	265	326	404	366	352	218	101	18	1
1999.....	168	176	226	276	317	356	373	319	202	93	24	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Hispanic, male

2003.....	114	134	146	162	197	217	198	146	98	32	8	-
2002.....	120	114	112	149	202	213	197	144	91	33	4	-
2001.....	112	110	127	156	183	204	177	128	77	21	4	2
2000.....	124	114	116	141	185	190	161	135	79	34	3	1
1999.....	113	102	143	163	181	179	165	136	73	25	10	1

Hispanic, female

2003.....	63	96	108	123	165	247	243	233	205	62	23	-
2002.....	68	82	82	125	125	205	234	217	177	73	18	-
2001.....	54	77	84	137	152	201	203	220	156	84	21	-
2000.....	59	63	83	124	141	214	205	217	139	67	15	-
1999.....	55	74	83	113	136	177	208	183	129	68	14	-

Non-Hispanic 1/, both sexes

2003.....	3,867	4,761	5,331	6,698	9,342	13,368	17,315	17,614	12,485	4,815	1,060	4
2002.....	3,881	4,657	5,242	6,732	9,755	13,980	17,501	17,644	12,391	4,616	986	3
2001.....	3,848	4,388	5,084	6,867	9,874	13,760	16,761	17,032	11,724	4,557	968	1
2000.....	3,691	4,257	5,226	7,156	10,132	13,907	16,371	16,685	11,839	4,495	976	3
1999.....	3,603	4,263	5,477	7,325	10,687	14,129	16,561	16,753	11,430	4,445	986	4

Non-Hispanic 1/, male

2003.....	2,435	2,982	3,216	3,852	5,131	6,735	7,607	6,448	3,585	1,023	144	3
2002.....	2,454	2,863	3,161	3,820	5,336	6,925	7,614	6,537	3,443	994	155	2
2001.....	2,372	2,656	3,090	3,876	5,387	6,882	7,338	6,082	3,193	983	155	1
2000.....	2,359	2,696	3,170	4,107	5,557	6,878	7,013	6,024	3,260	959	164	2
1999.....	2,255	2,698	3,333	4,379	5,850	6,981	7,225	5,922	3,143	970	136	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic 1/, female

2003.....	1,432	1,779	2,115	2,846	4,211	6,633	9,708	11,166	8,900	3,792	916	1
2002.....	1,427	1,794	2,081	2,912	4,419	7,055	9,887	11,107	8,948	3,622	831	1
2001.....	1,476	1,732	1,994	2,991	4,487	6,878	9,423	10,950	8,531	3,574	813	-
2000.....	1,332	1,561	2,056	3,049	4,575	7,029	9,358	10,661	8,579	3,536	812	1
1999.....	1,348	1,565	2,144	2,946	4,837	7,148	9,336	10,831	8,287	3,475	850	2

Non-Hispanic White, both sexes

2003.....	2,717	3,581	4,105	5,283	7,724	11,698	15,588	16,192	11,522	4,417	934	3
2002.....	2,765	3,509	4,051	5,419	8,208	12,212	15,807	16,232	11,422	4,178	848	2
2001.....	2,766	3,242	3,974	5,493	8,346	11,937	15,117	15,636	10,842	4,178	848	1
2000.....	2,630	3,231	4,071	5,780	8,549	12,119	14,845	15,313	10,969	4,095	856	2
1999.....	2,615	3,160	4,200	5,928	9,002	12,339	14,979	15,344	10,615	4,102	854	3

Non-Hispanic White, male

2003.....	1,747	2,271	2,536	3,079	4,323	5,950	6,890	5,960	3,294	930	123	2
2002.....	1,804	2,196	2,504	3,130	4,553	6,107	6,918	6,055	3,137	893	130	2
2001.....	1,773	1,985	2,448	3,183	4,595	6,010	6,625	5,579	2,926	888	131	1
2000.....	1,698	2,069	2,524	3,362	4,767	6,050	6,404	5,531	3,017	850	139	2
1999.....	1,671	2,030	2,600	3,603	4,984	6,137	6,559	5,402	2,890	877	113	2

Non-Hispanic White, female

2003.....	970	1,310	1,569	2,204	3,401	5,748	8,698	10,232	8,228	3,487	811	1
2002.....	961	1,313	1,547	2,289	3,655	6,105	8,889	10,177	8,285	3,285	718	-
2001.....	993	1,257	1,526	2,310	3,751	5,927	8,492	10,057	7,916	3,290	717	-
2000.....	932	1,162	1,547	2,418	3,782	6,069	8,441	9,782	7,952	3,245	717	-
1999.....	944	1,130	1,600	2,325	4,018	6,202	8,420	9,942	7,725	3,225	741	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Non-Hispanic Black, both sexes

2003.....	1,063	1,069	1,098	1,255	1,430	1,434	1,433	1,165	774	345	109	1
2002.....	1,031	1,044	1,077	1,191	1,337	1,520	1,429	1,155	824	376	115	1
2001.....	995	1,040	1,014	1,238	1,323	1,583	1,402	1,195	754	310	105	-
2000.....	978	930	1,048	1,240	1,419	1,584	1,331	1,181	729	339	106	1
1999.....	909	1,006	1,162	1,257	1,506	1,572	1,381	1,211	707	298	112	1

Non-Hispanic Black, male

2003.....	634	645	611	689	707	664	583	376	207	75	16	1
2002.....	598	604	585	623	673	674	565	366	240	74	18	-
2001.....	552	610	590	618	667	744	593	414	198	72	20	-
2000.....	609	572	583	667	688	716	529	407	186	86	21	-
1999.....	543	604	668	699	769	709	575	414	208	77	21	-

Non-Hispanic Black, female

2003.....	429	424	487	566	723	770	850	789	567	270	93	-
2002.....	433	440	492	568	664	846	864	789	584	302	97	1
2001.....	443	430	424	620	656	839	809	781	556	238	85	-
2000.....	369	358	465	573	731	868	802	774	543	253	85	1
1999.....	366	402	494	558	737	863	806	797	499	221	91	1

Origin not stated 2/, both sexes

2003.....	16	19	21	14	22	29	36	29	18	5	1	2
2002.....	14	15	14	17	26	30	43	28	21	12	1	3
2001.....	14	10	15	17	29	32	34	42	29	11	1	2
2000.....	12	20	18	13	29	29	23	41	23	13	2	-
1999.....	10	17	18	21	36	35	38	39	19	4	3	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Origin not stated 2/, male

2003.....	11	13	17	6	11	14	20	9	6	1	-	2
2002.....	10	8	8	11	15	21	15	7	3	4	-	1
2001.....	11	4	13	12	18	19	10	14	12	1	-	1
2000.....	6	12	12	10	18	14	10	16	4	5	-	-
1999.....	9	11	12	11	23	22	19	15	6	2	-	1

Origin not stated 2/, female

2003.....	5	6	4	8	11	15	16	20	12	4	1	-
2002.....	4	7	6	6	11	9	28	21	18	8	1	2
2001.....	3	6	2	5	11	13	24	28	17	10	1	1
2000.....	6	8	6	3	11	15	13	25	19	8	2	-
1999.....	1	6	6	10	13	13	19	24	13	2	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All origins, both sexes

2003.....	631	785	970	1,272	1,832	2,864	3,785	3,949	3,233	1,348	374	-
2002.....	541	681	845	1,126	1,796	2,592	3,546	3,801	2,915	1,276	293	-
2001.....	546	630	832	1,167	1,665	2,638	3,355	3,606	2,573	1,136	257	-
2000.....	526	597	843	1,094	1,690	2,507	3,116	3,273	2,375	1,017	242	-
1999.....	470	538	780	1,146	1,653	2,400	2,924	2,949	2,208	942	218	-

All origins, male

2003.....	395	457	532	676	859	1,163	1,359	1,163	773	240	42	-
2002.....	322	398	452	563	856	1,103	1,302	1,182	707	215	28	-
2001.....	330	354	430	603	776	1,100	1,182	1,129	610	192	28	-
2000.....	331	352	466	504	804	1,049	1,155	991	569	155	32	-
1999.....	286	316	416	589	783	996	1,071	878	521	161	20	-

All origins, female

2003.....	236	328	438	596	973	1,701	2,426	2,786	2,460	1,108	332	-
2002.....	219	283	393	563	940	1,489	2,244	2,619	2,208	1,061	265	-
2001.....	216	276	402	564	889	1,538	2,173	2,477	1,963	944	229	-
2000.....	195	245	377	590	886	1,458	1,961	2,282	1,806	862	210	-
1999.....	184	222	364	557	870	1,404	1,853	2,071	1,687	781	198	-

Hispanic, both sexes

2003.....	38	52	69	86	118	136	149	153	107	52	18	-
2002.....	22	50	73	66	117	130	153	116	96	43	13	-
2001.....	32	31	52	72	115	135	136	129	82	40	5	-
2000.....	37	34	75	65	97	118	114	122	55	38	8	-
1999.....	35	29	53	71	87	110	122	80	80	19	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Hispanic, male

2003.....	27	26	47	34	50	55	61	44	25	14	4	-
2002.....	9	30	36	26	54	61	52	33	32	9	5	-
2001.....	18	13	21	30	54	58	56	37	24	6	1	-
2000.....	28	20	45	26	46	57	50	38	21	7	4	-
1999.....	19	11	29	42	42	56	43	26	28	7	-	-

Hispanic, female

2003.....	11	26	22	52	68	81	88	109	82	38	14	-
2002.....	13	20	37	40	63	69	101	83	64	34	8	-
2001.....	14	18	31	42	61	77	80	92	58	34	4	-
2000.....	9	14	30	39	51	61	64	84	34	31	4	-
1999.....	16	18	24	29	45	54	79	54	52	12	6	-

Non-Hispanic 1/, both sexes

2003.....	591	732	896	1,181	1,702	2,722	3,628	3,791	3,122	1,292	355	-
2002.....	517	628	772	1,053	1,672	2,457	3,383	3,673	2,811	1,228	278	-
2001.....	514	594	776	1,091	1,545	2,492	3,214	3,467	2,485	1,096	252	-
2000.....	487	557	763	1,028	1,588	2,377	2,996	3,140	2,309	978	233	-
1999.....	435	506	726	1,071	1,562	2,287	2,786	2,857	2,121	920	212	-

Non-Hispanic 1/, male

2003.....	367	431	480	638	803	1,105	1,296	1,119	748	225	38	-
2002.....	311	365	416	530	798	1,039	1,246	1,144	674	206	23	-
2001.....	312	337	406	570	718	1,037	1,123	1,088	583	186	27	-
2000.....	301	328	418	477	755	986	1,102	949	547	148	28	-
1999.....	267	303	386	545	740	940	1,018	850	490	153	20	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic 1/, female

2003.....	224	301	416	543	899	1,617	2,332	2,672	2,374	1,067	317	-
2002.....	206	263	356	523	874	1,418	2,137	2,529	2,137	1,022	255	-
2001.....	202	257	370	521	827	1,455	2,091	2,379	1,902	910	225	-
2000.....	186	229	345	551	833	1,391	1,894	2,191	1,762	830	205	-
1999.....	168	203	340	526	822	1,347	1,768	2,007	1,631	767	192	-

Non-Hispanic White, both sexes

2003.....	320	387	531	734	1,128	2,053	2,973	3,266	2,694	1,119	293	-
2002.....	281	339	466	675	1,163	1,835	2,735	3,077	2,404	1,075	234	-
2001.....	249	323	453	679	1,068	1,908	2,591	2,922	2,156	958	206	-
2000.....	232	307	429	644	1,100	1,798	2,454	2,644	1,992	818	187	-
1999.....	200	275	411	683	1,105	1,710	2,230	2,380	1,818	775	172	-

Non-Hispanic White, male

2003.....	199	245	282	421	555	857	1,059	954	639	189	29	-
2002.....	180	210	267	350	570	787	997	977	576	176	17	-
2001.....	155	192	247	364	509	804	897	917	508	155	20	-
2000.....	155	190	244	315	537	763	908	801	462	116	20	-
1999.....	127	181	229	361	536	714	812	694	413	117	16	-

Non-Hispanic White, female

2003.....	121	142	249	313	573	1,196	1,914	2,312	2,055	930	264	-
2002.....	101	129	199	325	593	1,048	1,738	2,100	1,828	899	217	-
2001.....	94	131	206	315	559	1,104	1,694	2,005	1,648	803	186	-
2000.....	77	117	185	329	563	1,035	1,546	1,843	1,530	702	167	-
1999.....	73	94	182	322	569	996	1,418	1,686	1,405	658	156	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Non-Hispanic Black, both sexes

2003.....	258	323	341	405	513	594	568	446	366	154	57	-
2002.....	225	272	278	339	456	559	574	512	351	130	40	-
2001.....	251	253	300	372	417	512	544	478	287	126	43	-
2000.....	237	240	315	362	439	518	462	435	281	142	41	-
1999.....	226	221	285	342	422	528	478	416	266	125	35	-

Non-Hispanic Black, male

2003.....	161	173	186	200	224	221	198	136	87	32	8	-
2002.....	125	145	129	159	200	228	214	138	83	26	6	-
2001.....	150	136	148	190	184	202	197	146	60	27	7	-
2000.....	133	135	166	153	191	195	163	123	74	27	5	-
1999.....	135	114	142	165	183	204	177	131	61	30	4	-

Non-Hispanic Black, female

2003.....	97	150	155	205	289	373	370	310	279	122	49	-
2002.....	100	127	149	180	256	331	360	374	268	104	34	-
2001.....	101	117	152	182	233	310	347	332	227	99	36	-
2000.....	104	105	149	209	248	323	299	312	207	115	36	-
1999.....	91	107	143	177	239	324	301	285	205	95	31	-

Origin not stated 2/, both sexes

2003.....	2	1	5	5	12	6	8	5	4	4	1	-
2002.....	2	3	-	7	7	5	10	12	8	5	2	-
2001.....	-	5	4	4	5	11	5	10	6	-	-	-
2000.....	2	6	5	1	5	12	6	11	11	1	1	-
1999.....	-	3	1	4	4	3	16	12	7	3	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

Origin not stated 2/, male

2003.....	1	-	5	4	6	3	2	-	-	1	-	-
2002.....	2	3	-	7	4	3	4	5	1	-	-	-
2001.....	-	4	3	3	4	5	3	4	3	-	-	-
2000.....	2	4	3	1	3	6	3	4	1	-	-	-
1999.....	-	2	1	2	1	-	10	2	3	1	-	-

Origin not stated 2/, female

2003.....	1	1	-	1	6	3	6	5	4	3	1	-
2002.....	-	-	-	-	3	2	6	7	7	5	2	-
2001.....	-	1	1	1	1	6	2	6	3	-	-	-
2000.....	-	2	2	-	2	6	3	7	10	1	1	-
1999.....	-	1	-	2	3	3	6	10	4	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cerebrovascular diseases (I60-I69)

##### All origins, both sexes

2003.....	3,550	4,392	5,554	7,765	12,943	21,917	30,930	32,139	22,367	8,493	1,580	2
2002.....	3,440	4,293	5,604	8,026	13,966	22,911	31,978	33,197	23,110	8,521	1,584	3
2001.....	3,403	4,093	5,515	8,170	14,428	23,772	32,050	33,465	22,650	8,398	1,553	4
2000.....	3,386	4,145	5,811	8,671	14,978	24,678	32,342	34,030	23,205	8,530	1,611	2
1999.....	3,108	4,152	5,500	8,694	15,398	25,007	32,420	33,841	23,149	8,571	1,519	2

##### All origins, male

2003.....	1,961	2,394	3,084	4,138	6,530	9,993	12,241	10,471	5,634	1,575	179	1
2002.....	1,876	2,373	3,083	4,305	6,902	10,283	12,372	10,884	5,713	1,510	209	1
2001.....	1,851	2,251	3,026	4,345	7,304	10,671	12,630	10,731	5,482	1,506	211	4
2000.....	1,834	2,294	3,202	4,513	7,529	11,192	12,754	10,864	5,588	1,564	196	1
1999.....	1,714	2,256	3,049	4,522	7,698	11,349	12,739	10,818	5,569	1,467	198	1

##### All origins, female

2003.....	1,589	1,998	2,470	3,627	6,413	11,924	18,689	21,668	16,733	6,918	1,401	1
2002.....	1,564	1,920	2,521	3,721	7,064	12,628	19,606	22,313	17,397	7,011	1,375	2
2001.....	1,552	1,842	2,489	3,825	7,124	13,101	19,420	22,734	17,168	6,892	1,342	-
2000.....	1,552	1,851	2,609	4,158	7,449	13,486	19,588	23,166	17,617	6,966	1,415	1
1999.....	1,394	1,896	2,451	4,172	7,700	13,658	19,681	23,023	17,580	7,104	1,321	1

##### Hispanic, both sexes

2003.....	318	353	383	529	673	1,001	1,028	863	581	222	33	-
2002.....	291	321	387	462	753	901	1,017	838	535	202	45	-
2001.....	300	301	353	471	735	959	933	868	532	221	47	-
2000.....	232	271	410	498	721	859	883	852	539	209	32	-
1999.....	235	286	331	524	687	823	849	828	505	170	33	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cerebrovascular diseases (I60-I69)

##### Hispanic, male

2003.....	189	203	228	280	337	454	425	319	155	65	5	-
2002.....	171	196	225	256	396	421	404	282	177	42	11	-
2001.....	174	171	207	269	368	456	408	300	161	55	12	-
2000.....	136	172	220	267	378	421	378	282	159	63	7	-
1999.....	149	157	197	273	368	423	335	284	183	50	10	-

##### Hispanic, female

2003.....	129	150	155	249	336	547	603	544	426	157	28	-
2002.....	120	125	162	206	357	480	613	556	358	160	34	-
2001.....	126	130	146	202	367	503	525	568	371	166	35	-
2000.....	96	99	190	231	343	438	505	570	380	146	25	-
1999.....	86	129	134	251	319	400	514	544	322	120	23	-

##### Non-Hispanic 1/, both sexes

2003.....	3,218	4,025	5,154	7,220	12,238	20,888	29,854	31,220	21,742	8,255	1,546	1
2002.....	3,140	3,961	5,199	7,546	13,185	21,958	30,892	32,291	22,516	8,302	1,537	3
2001.....	3,092	3,779	5,137	7,683	13,663	22,752	31,058	32,525	22,071	8,154	1,503	4
2000.....	3,143	3,861	5,377	8,160	14,213	23,754	31,375	33,112	22,615	8,305	1,576	1
1999.....	2,858	3,848	5,143	8,145	14,669	24,125	31,518	32,951	22,585	8,386	1,482	2

##### Non-Hispanic 1/, male

2003.....	1,765	2,185	2,841	3,846	6,177	9,525	11,797	10,139	5,469	1,507	174	1
2002.....	1,699	2,172	2,846	4,039	6,493	9,834	11,939	10,583	5,522	1,467	198	1
2001.....	1,673	2,071	2,801	4,066	6,923	10,187	12,190	10,412	5,309	1,446	197	4
2000.....	1,688	2,111	2,968	4,239	7,125	10,737	12,347	10,558	5,418	1,496	188	-
1999.....	1,558	2,088	2,834	4,234	7,308	10,892	12,383	10,518	5,366	1,417	188	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cerebrovascular diseases (I60-I69)

##### Non-Hispanic 1/, female

2003.....	1,453	1,840	2,313	3,374	6,061	11,363	18,057	21,081	16,273	6,748	1,372	-
2002.....	1,441	1,789	2,353	3,507	6,692	12,124	18,953	21,708	16,994	6,835	1,339	2
2001.....	1,419	1,708	2,336	3,617	6,740	12,565	18,868	22,113	16,762	6,708	1,306	-
2000.....	1,455	1,750	2,409	3,921	7,088	13,017	19,028	22,554	17,197	6,809	1,388	1
1999.....	1,300	1,760	2,309	3,911	7,361	13,233	19,135	22,433	17,219	6,969	1,294	1

##### Non-Hispanic White, both sexes

2003.....	1,930	2,693	3,600	5,407	9,852	17,759	26,518	28,371	19,865	7,521	1,321	1
2002.....	1,913	2,733	3,699	5,654	10,701	18,821	27,506	29,412	20,578	7,553	1,341	2
2001.....	1,837	2,606	3,602	5,715	11,123	19,553	27,751	29,529	20,201	7,436	1,295	1
2000.....	2,053	2,657	3,898	6,277	11,712	20,501	28,055	30,109	20,776	7,491	1,362	-
1999.....	1,841	2,649	3,662	6,342	11,990	20,866	28,312	29,995	20,832	7,676	1,315	-

##### Non-Hispanic White, male

2003.....	1,066	1,433	1,994	2,936	5,011	8,161	10,538	9,261	4,988	1,339	153	1
2002.....	1,032	1,490	2,021	3,032	5,336	8,484	10,697	9,680	5,016	1,313	153	1
2001.....	1,006	1,421	1,979	3,066	5,687	8,818	10,944	9,439	4,851	1,294	172	1
2000.....	1,104	1,431	2,174	3,274	5,902	9,308	11,071	9,594	4,928	1,301	164	-
1999.....	998	1,452	2,027	3,320	6,017	9,436	11,182	9,548	4,884	1,286	168	-

##### Non-Hispanic White, female

2003.....	864	1,260	1,606	2,471	4,841	9,598	15,980	19,110	14,877	6,182	1,168	-
2002.....	881	1,243	1,678	2,622	5,365	10,337	16,809	19,732	15,562	6,240	1,188	1
2001.....	831	1,185	1,623	2,649	5,436	10,735	16,807	20,090	15,350	6,142	1,123	-
2000.....	949	1,226	1,724	3,003	5,810	11,193	16,984	20,515	15,848	6,190	1,198	-
1999.....	843	1,197	1,635	3,022	5,973	11,430	17,130	20,447	15,948	6,390	1,147	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cerebrovascular diseases (I60-I69)

##### Non-Hispanic Black, both sexes

2003.....	1,129	1,132	1,317	1,483	1,928	2,530	2,646	2,273	1,525	598	196	-
2002.....	1,067	1,048	1,255	1,584	2,016	2,576	2,701	2,298	1,546	610	168	1
2001.....	1,096	992	1,297	1,640	2,044	2,627	2,630	2,391	1,530	604	170	3
2000.....	951	1,032	1,233	1,573	2,087	2,644	2,700	2,472	1,507	685	197	1
1999.....	882	1,017	1,252	1,529	2,263	2,715	2,599	2,435	1,452	614	143	1

##### Non-Hispanic Black, male

2003.....	616	639	724	739	934	1,094	977	656	343	113	15	-
2002.....	588	592	696	848	978	1,084	947	671	349	100	38	-
2001.....	586	554	698	854	990	1,119	949	726	319	114	18	3
2000.....	515	587	653	812	1,005	1,138	1,017	727	367	145	21	-
1999.....	491	541	694	782	1,075	1,169	931	735	363	110	17	-

##### Non-Hispanic Black, female

2003.....	513	493	593	744	994	1,436	1,669	1,617	1,182	485	181	-
2002.....	479	456	559	736	1,038	1,492	1,754	1,627	1,197	510	130	1
2001.....	510	438	599	786	1,054	1,508	1,681	1,665	1,211	490	152	-
2000.....	436	445	580	761	1,082	1,506	1,683	1,745	1,140	540	176	1
1999.....	391	476	558	747	1,188	1,546	1,668	1,700	1,089	504	126	1

##### Origin not stated 2/, both sexes

2003.....	14	14	17	16	32	28	48	56	44	16	1	1
2002.....	9	11	18	18	28	52	69	68	59	17	2	-
2001.....	11	13	25	16	30	61	59	72	47	23	3	-
2000.....	11	13	24	13	44	65	84	66	51	16	3	1
1999.....	15	18	26	25	42	59	53	62	59	15	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cerebrovascular diseases (I60-I69)

Origin not stated 2/, male

2003.....	7	6	15	12	16	14	19	13	10	3	-	-
2002.....	6	5	12	10	13	28	29	19	14	1	-	-
2001.....	4	9	18	10	13	28	32	19	12	5	2	-
2000.....	10	11	14	7	26	34	29	24	11	5	1	1
1999.....	7	11	18	15	22	34	21	16	20	-	-	-

Origin not stated 2/, female

2003.....	7	8	2	4	16	14	29	43	34	13	1	1
2002.....	3	6	6	8	15	24	40	49	45	16	2	-
2001.....	7	4	7	6	17	33	27	53	35	18	1	-
2000.....	1	2	10	6	18	31	55	42	40	11	2	-
1999.....	8	7	8	10	20	25	32	46	39	15	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Atherosclerosis (I70)

##### All origins, both sexes

2003.....	120	171	295	436	775	1,482	2,355	2,946	2,582	1,375	385	1
2002.....	121	207	280	494	806	1,521	2,541	3,090	2,802	1,416	415	1
2001.....	121	198	275	471	864	1,624	2,532	3,214	2,786	1,471	417	1
2000.....	105	170	281	495	928	1,685	2,592	3,188	2,870	1,518	429	-
1999.....	129	172	314	485	991	1,770	2,615	3,362	2,987	1,581	460	-

##### All origins, male

2003.....	77	113	185	245	404	742	992	1,009	668	272	53	1
2002.....	75	127	178	282	446	748	1,055	1,088	755	281	53	1
2001.....	75	118	168	273	470	765	1,026	1,123	710	285	54	1
2000.....	71	108	186	290	510	838	1,097	1,090	738	283	55	-
1999.....	84	114	193	277	518	845	1,082	1,100	773	316	66	-

##### All origins, female

2003.....	43	58	110	191	371	740	1,363	1,937	1,914	1,103	332	-
2002.....	46	80	102	212	360	773	1,486	2,002	2,047	1,135	362	-
2001.....	46	80	107	198	394	859	1,506	2,091	2,076	1,186	363	-
2000.....	34	62	95	205	418	847	1,495	2,098	2,132	1,235	374	-
1999.....	45	58	121	208	473	925	1,533	2,262	2,214	1,265	394	-

##### Hispanic, both sexes

2003.....	6	14	10	21	35	47	74	64	64	29	9	-
2002.....	6	9	11	24	26	58	78	78	74	30	14	-
2001.....	11	12	7	14	32	60	68	99	59	39	12	-
2000.....	5	6	10	24	39	53	71	83	77	32	8	-
1999.....	9	11	14	15	39	60	70	79	61	35	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Atherosclerosis (I70)

##### Hispanic, male

2003.....	3	9	8	11	22	31	32	23	23	13	3	-
2002.....	3	7	5	12	15	29	36	29	23	6	3	-
2001.....	8	8	3	8	16	31	30	39	13	11	4	-
2000.....	4	2	6	13	26	34	31	27	17	7	1	-
1999.....	5	7	9	9	22	30	30	37	18	11	4	-

##### Hispanic, female

2003.....	3	5	2	10	13	16	42	41	41	16	6	-
2002.....	3	2	6	12	11	29	42	49	51	24	11	-
2001.....	3	4	4	6	16	29	38	60	46	28	8	-
2000.....	1	4	4	11	13	19	40	56	60	25	7	-
1999.....	4	4	5	6	17	30	40	42	43	24	5	-

##### Non-Hispanic 1/, both sexes

2003.....	113	156	284	414	738	1,427	2,269	2,874	2,510	1,344	374	-
2002.....	115	198	268	470	778	1,455	2,456	3,001	2,722	1,386	400	1
2001.....	110	185	267	455	830	1,556	2,455	3,104	2,720	1,429	405	1
2000.....	100	164	270	469	888	1,627	2,518	3,095	2,787	1,481	420	-
1999.....	120	160	297	465	951	1,706	2,536	3,273	2,916	1,544	448	-

##### Non-Hispanic 1/, male

2003.....	73	103	176	234	381	707	956	982	645	259	50	-
2002.....	72	120	172	270	430	716	1,017	1,056	731	275	49	1
2001.....	67	109	165	264	454	729	992	1,080	695	272	50	1
2000.....	67	106	179	275	484	801	1,064	1,058	717	276	53	-
1999.....	79	106	181	266	495	814	1,047	1,059	753	304	62	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Atherosclerosis (I70)

##### Non-Hispanic 1/, female

2003.....	40	53	108	180	357	720	1,313	1,892	1,865	1,085	324	-
2002.....	43	78	96	200	348	739	1,439	1,945	1,991	1,111	351	-
2001.....	43	76	102	191	376	827	1,463	2,024	2,025	1,157	355	-
2000.....	33	58	91	194	404	826	1,454	2,037	2,070	1,205	367	-
1999.....	41	54	116	199	456	892	1,489	2,214	2,163	1,240	386	-

##### Non-Hispanic White, both sexes

2003.....	84	124	235	344	634	1,286	2,063	2,684	2,334	1,256	332	-
2002.....	79	161	215	411	671	1,282	2,236	2,828	2,560	1,301	357	1
2001.....	86	153	208	391	728	1,408	2,267	2,926	2,561	1,333	366	1
2000.....	68	134	215	405	776	1,479	2,303	2,892	2,610	1,394	374	-
1999.....	94	129	245	394	847	1,534	2,346	3,050	2,725	1,440	402	-

##### Non-Hispanic White, male

2003.....	56	83	147	196	324	630	865	917	602	244	45	-
2002.....	47	99	138	239	374	631	930	999	688	260	43	1
2001.....	51	92	133	228	403	656	914	1,025	652	256	40	1
2000.....	46	91	147	237	436	740	976	983	666	254	46	-
1999.....	65	83	152	227	440	724	966	981	702	279	52	-

##### Non-Hispanic White, female

2003.....	28	41	88	148	310	656	1,198	1,767	1,732	1,012	287	-
2002.....	32	62	77	172	297	651	1,306	1,829	1,872	1,041	314	-
2001.....	35	61	75	163	325	752	1,353	1,901	1,909	1,077	326	-
2000.....	22	43	68	168	340	739	1,327	1,909	1,944	1,140	328	-
1999.....	29	46	93	167	407	810	1,380	2,069	2,023	1,161	350	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Atherosclerosis (I70)

##### Non-Hispanic Black, both sexes

2003.....	27	30	42	61	91	120	176	153	148	78	34	-
2002.....	31	29	48	49	85	154	195	150	135	68	37	-
2001.....	20	28	56	57	88	131	154	149	137	86	32	-
2000.....	28	28	51	51	96	130	190	172	153	76	40	-
1999.....	23	26	50	62	88	152	160	190	162	91	39	-

##### Non-Hispanic Black, male

2003.....	16	19	25	32	51	62	78	50	33	12	4	-
2002.....	22	17	32	25	45	76	77	48	35	13	4	-
2001.....	13	14	29	30	44	65	59	44	35	12	7	-
2000.....	17	14	29	33	41	52	79	63	41	19	5	-
1999.....	11	19	27	34	46	78	64	68	39	23	10	-

##### Non-Hispanic Black, female

2003.....	11	11	17	29	40	58	98	103	115	66	30	-
2002.....	9	12	16	24	40	78	118	102	100	55	33	-
2001.....	7	14	27	27	44	66	95	105	102	74	25	-
2000.....	11	14	22	18	55	78	111	109	112	57	35	-
1999.....	12	7	23	28	42	74	96	122	123	68	29	-

##### Origin not stated 2/, both sexes

2003.....	1	1	1	1	2	8	12	8	8	2	2	1
2002.....	-	-	1	-	2	8	7	11	6	-	1	-
2001.....	-	1	1	2	2	8	9	11	7	3	-	-
2000.....	-	-	1	2	1	5	3	10	6	5	1	-
1999.....	-	1	3	5	1	4	9	10	10	2	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Atherosclerosis (I70)

##### Origin not stated 2/, male

2003.....	1	1	1	-	1	4	4	4	-	-	-	1
2002.....	-	-	1	-	1	3	2	3	1	-	1	-
2001.....	-	1	-	1	-	5	4	4	2	2	-	-
2000.....	-	-	1	2	-	3	2	5	4	-	1	-
1999.....	-	1	3	2	1	1	5	4	2	1	-	-

##### Origin not stated 2/, female

2003.....	-	-	-	1	1	4	8	4	8	2	2	-
2002.....	-	-	-	-	1	5	5	8	5	-	-	-
2001.....	-	-	1	1	2	3	5	7	5	1	-	-
2000.....	-	-	-	-	1	2	1	5	2	5	-	-
1999.....	-	-	-	3	-	3	4	6	8	1	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

All origins, both sexes

2003.....	675	900	1,370	1,998	2,936	4,215	4,479	3,739	2,229	762	152	-
2002.....	591	928	1,358	2,137	3,173	4,292	4,453	3,768	2,151	743	154	-
2001.....	603	850	1,345	2,127	3,218	4,498	4,534	3,867	2,210	748	140	1
2000.....	614	866	1,370	2,331	3,595	4,662	4,560	3,819	2,218	757	126	1
1999.....	553	877	1,333	2,244	3,564	4,662	4,602	3,792	2,161	765	145	-

All origins, male

2003.....	472	603	914	1,331	1,761	2,445	2,287	1,629	762	181	25	-
2002.....	401	624	902	1,408	1,899	2,406	2,284	1,675	715	191	23	-
2001.....	397	566	888	1,357	1,991	2,476	2,363	1,633	719	188	23	1
2000.....	424	607	923	1,524	2,247	2,696	2,323	1,678	734	204	21	1
1999.....	368	597	922	1,472	2,209	2,684	2,331	1,567	711	172	26	-

All origins, female

2003.....	203	297	456	667	1,175	1,770	2,192	2,110	1,467	581	127	-
2002.....	190	304	456	729	1,274	1,886	2,169	2,093	1,436	552	131	-
2001.....	206	284	457	770	1,227	2,022	2,171	2,234	1,491	560	117	-
2000.....	190	259	447	807	1,348	1,966	2,237	2,141	1,484	553	105	-
1999.....	185	280	411	772	1,355	1,978	2,271	2,225	1,450	593	119	-

Hispanic, both sexes

2003.....	30	46	50	77	95	149	122	120	56	23	6	-
2002.....	33	48	69	73	102	130	116	85	53	21	8	-
2001.....	42	39	49	71	85	125	91	82	43	25	2	1
2000.....	37	45	54	88	129	129	86	71	56	25	3	-
1999.....	23	39	55	73	97	97	114	92	46	16	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

Hispanic, male

2003.....	23	34	39	47	56	89	64	61	22	6	1	-
2002.....	22	26	40	44	60	70	68	29	23	5	4	-
2001.....	26	22	33	47	53	76	43	37	13	6	1	1
2000.....	24	28	36	51	71	81	45	33	24	6	2	-
1999.....	14	25	34	44	57	57	62	48	22	4	2	-

Hispanic, female

2003.....	7	12	11	30	39	60	58	59	34	17	5	-
2002.....	11	22	29	29	42	60	48	56	30	16	4	-
2001.....	16	17	16	24	32	49	48	45	30	19	1	-
2000.....	13	17	18	37	58	48	41	38	32	19	1	-
1999.....	9	14	21	29	40	40	52	44	24	12	2	-

Non-Hispanic 1/, both sexes

2003.....	642	854	1,317	1,918	2,828	4,055	4,348	3,614	2,171	736	146	-
2002.....	557	877	1,282	2,056	3,064	4,145	4,330	3,673	2,094	718	146	-
2001.....	557	808	1,291	2,048	3,122	4,364	4,431	3,774	2,160	722	138	-
2000.....	575	818	1,311	2,238	3,457	4,521	4,464	3,736	2,157	731	123	1
1999.....	529	834	1,275	2,162	3,457	4,560	4,477	3,694	2,109	747	141	-

Non-Hispanic 1/, male

2003.....	449	569	872	1,281	1,698	2,349	2,220	1,566	739	175	24	-
2002.....	378	595	860	1,359	1,835	2,328	2,212	1,641	691	186	19	-
2001.....	368	541	851	1,305	1,929	2,394	2,314	1,593	704	181	22	-
2000.....	398	577	884	1,471	2,169	2,607	2,276	1,641	708	197	19	1
1999.....	354	571	885	1,422	2,145	2,626	2,263	1,519	687	167	24	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic 1/, female

2003.....	193	285	445	637	1,130	1,706	2,128	2,048	1,432	561	122	-
2002.....	179	282	422	697	1,229	1,817	2,118	2,032	1,403	532	127	-
2001.....	189	267	440	743	1,193	1,970	2,117	2,181	1,456	541	116	-
2000.....	177	241	427	767	1,288	1,914	2,188	2,095	1,449	534	104	-
1999.....	175	263	390	740	1,312	1,934	2,214	2,175	1,422	580	117	-

Non-Hispanic White, both sexes

2003.....	466	694	1,093	1,651	2,507	3,656	3,911	3,290	1,986	648	119	-
2002.....	401	692	1,066	1,767	2,718	3,738	3,946	3,361	1,880	655	113	-
2001.....	405	641	1,061	1,782	2,757	3,892	4,016	3,447	1,960	628	104	-
2000.....	415	648	1,086	1,940	3,106	4,058	4,075	3,414	1,938	638	99	-
1999.....	401	672	1,101	1,891	3,109	4,129	4,081	3,373	1,903	660	115	-

Non-Hispanic White, male

2003.....	337	475	731	1,120	1,543	2,145	2,014	1,453	686	155	20	-
2002.....	275	472	726	1,205	1,645	2,144	2,045	1,526	622	174	15	-
2001.....	267	446	712	1,163	1,734	2,177	2,126	1,482	641	163	19	-
2000.....	295	470	755	1,318	1,983	2,371	2,121	1,537	638	174	15	-
1999.....	270	479	782	1,281	1,972	2,423	2,091	1,405	635	146	16	-

Non-Hispanic White, female

2003.....	129	219	362	531	964	1,511	1,897	1,837	1,300	493	99	-
2002.....	126	220	340	562	1,073	1,594	1,901	1,835	1,258	481	98	-
2001.....	138	195	349	619	1,023	1,715	1,890	1,965	1,319	465	85	-
2000.....	120	178	331	622	1,123	1,687	1,954	1,877	1,300	464	84	-
1999.....	131	193	319	610	1,137	1,706	1,990	1,968	1,268	514	99	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

Non-Hispanic Black, both sexes

2003.....	156	134	191	227	265	322	361	266	151	73	24	-
2002.....	138	164	186	240	281	336	317	252	180	59	30	-
2001.....	129	150	198	210	316	394	352	271	177	84	31	-
2000.....	126	148	198	246	294	386	324	276	187	82	21	1
1999.....	108	146	144	234	283	357	332	276	176	81	24	-

Non-Hispanic Black, male

2003.....	97	80	120	134	118	157	160	84	32	14	3	-
2002.....	92	108	115	122	146	140	129	84	57	9	4	-
2001.....	86	82	115	108	164	166	158	81	49	13	3	-
2000.....	85	90	108	126	153	190	124	82	49	18	4	1
1999.....	70	82	83	122	144	156	134	88	39	20	8	-

Non-Hispanic Black, female

2003.....	59	54	71	93	147	165	201	182	119	59	21	-
2002.....	46	56	71	118	135	196	188	168	123	50	26	-
2001.....	43	68	83	102	152	228	194	190	128	71	28	-
2000.....	41	58	90	120	141	196	200	194	138	64	17	-
1999.....	38	64	61	112	139	201	198	188	137	61	16	-

Origin not stated 2/, both sexes

2003.....	3	-	3	3	13	11	9	5	2	3	-	-
2002.....	1	3	7	8	7	17	7	10	4	4	-	-
2001.....	4	3	5	8	11	9	12	11	7	1	-	-
2000.....	2	3	5	5	9	12	10	12	5	1	-	-
1999.....	1	4	3	9	10	5	11	6	6	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of circulatory system (I71-I78)

Origin not stated 2/, male

2003.....	-	-	3	3	7	7	3	2	1	-	-	-
2002.....	1	3	2	5	4	8	4	5	1	-	-	-
2001.....	3	3	4	5	9	6	6	3	2	1	-	-
2000.....	2	2	3	2	7	8	2	4	2	1	-	-
1999.....	-	1	3	6	7	1	6	-	2	1	-	-

Origin not stated 2/, female

2003.....	3	-	-	-	6	4	6	3	1	3	-	-
2002.....	-	-	5	3	3	9	3	5	3	4	-	-
2001.....	1	-	1	3	2	3	6	8	5	-	-	-
2000.....	-	1	2	3	2	4	8	8	3	-	-	-
1999.....	1	3	-	3	3	4	5	6	4	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Aortic aneurysm and dissection (I71)

##### All origins, both sexes

2003.....	452	590	887	1,317	1,954	2,726	2,745	2,059	965	245	29	-
2002.....	383	598	862	1,404	2,078	2,779	2,739	2,013	914	239	21	-
2001.....	374	532	862	1,430	2,087	2,899	2,852	2,122	1,022	242	26	1
2000.....	394	585	908	1,552	2,460	3,043	2,792	2,134	985	213	25	-
1999.....	355	570	867	1,578	2,423	3,095	2,841	2,118	935	257	27	-

##### All origins, male

2003.....	347	448	659	912	1,262	1,667	1,525	1,033	411	74	7	-
2002.....	291	458	634	1,020	1,349	1,688	1,497	1,034	368	74	5	-
2001.....	287	398	633	981	1,401	1,722	1,591	1,000	406	78	7	1
2000.....	301	438	673	1,097	1,633	1,885	1,558	1,063	425	68	6	-
1999.....	270	430	658	1,114	1,598	1,905	1,596	1,008	398	83	6	-

##### All origins, female

2003.....	105	142	228	405	692	1,059	1,220	1,026	554	171	22	-
2002.....	92	140	228	384	729	1,091	1,242	979	546	165	16	-
2001.....	87	134	229	449	686	1,177	1,261	1,122	616	164	19	-
2000.....	93	147	235	455	827	1,158	1,234	1,071	560	145	19	-
1999.....	85	140	209	464	825	1,190	1,245	1,110	537	174	21	-

##### Hispanic, both sexes

2003.....	18	30	34	39	63	99	72	59	25	8	3	-
2002.....	19	31	44	42	47	72	55	32	17	6	1	-
2001.....	23	20	27	45	48	75	52	35	13	3	-	1
2000.....	28	26	32	46	75	83	54	36	22	5	-	-
1999.....	17	24	31	42	53	57	58	41	12	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Aortic aneurysm and dissection (I71)

##### Hispanic, male

2003.....	15	24	27	26	42	61	39	37	11	4	-	-
2002.....	14	21	30	28	35	40	38	15	9	1	1	-
2001.....	17	12	21	35	36	52	26	23	4	1	-	1
2000.....	20	19	23	33	43	57	29	19	12	1	-	-
1999.....	13	17	24	30	37	31	34	29	7	1	-	-

##### Hispanic, female

2003.....	3	6	7	13	21	38	33	22	14	4	3	-
2002.....	5	10	14	14	12	32	17	17	8	5	-	-
2001.....	6	8	6	10	12	23	26	12	9	2	-	-
2000.....	8	7	9	13	32	26	25	17	10	4	-	-
1999.....	4	7	7	12	16	26	24	12	5	4	-	-

##### Non-Hispanic 1/, both sexes

2003.....	432	560	851	1,276	1,883	2,619	2,667	1,997	938	237	26	-
2002.....	363	564	815	1,356	2,028	2,695	2,678	1,977	895	233	20	-
2001.....	349	509	832	1,379	2,029	2,818	2,792	2,084	1,005	238	26	-
2000.....	364	558	873	1,503	2,378	2,949	2,733	2,091	960	208	25	-
1999.....	338	543	835	1,528	2,365	3,036	2,775	2,076	921	252	27	-

##### Non-Hispanic 1/, male

2003.....	332	424	630	884	1,214	1,601	1,484	995	399	70	7	-
2002.....	276	434	602	987	1,312	1,643	1,456	1,016	358	73	4	-
2001.....	268	383	609	942	1,357	1,665	1,560	976	400	76	7	-
2000.....	279	418	648	1,063	1,585	1,821	1,528	1,042	411	67	6	-
1999.....	257	412	633	1,079	1,556	1,873	1,559	979	390	82	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Aortic aneurysm and dissection (I71)

##### Non-Hispanic 1/, female

2003.....	100	136	221	392	669	1,018	1,183	1,002	539	167	19	-
2002.....	87	130	213	369	716	1,052	1,222	961	537	160	16	-
2001.....	81	126	223	437	672	1,153	1,232	1,108	605	162	19	-
2000.....	85	140	225	440	793	1,128	1,205	1,049	549	141	19	-
1999.....	81	131	202	449	809	1,163	1,216	1,097	531	170	21	-

##### Non-Hispanic White, both sexes

2003.....	315	452	726	1,137	1,707	2,417	2,466	1,861	888	211	22	-
2002.....	254	451	705	1,216	1,841	2,499	2,489	1,844	825	213	15	-
2001.....	246	410	707	1,228	1,851	2,561	2,594	1,951	926	217	24	-
2000.....	274	451	750	1,335	2,191	2,705	2,545	1,948	875	183	23	-
1999.....	256	450	745	1,373	2,175	2,814	2,593	1,929	858	228	22	-

##### Non-Hispanic White, male

2003.....	252	355	542	801	1,125	1,497	1,380	940	376	63	6	-
2002.....	197	349	526	904	1,201	1,542	1,375	968	335	71	3	-
2001.....	194	320	521	856	1,253	1,534	1,468	927	379	71	7	-
2000.....	216	346	568	974	1,479	1,685	1,451	994	375	61	6	-
1999.....	195	353	575	992	1,459	1,774	1,469	917	369	73	5	-

##### Non-Hispanic White, female

2003.....	63	97	184	336	582	920	1,086	921	512	148	16	-
2002.....	57	102	179	312	640	957	1,114	876	490	142	12	-
2001.....	52	90	186	372	598	1,027	1,126	1,024	547	146	17	-
2000.....	58	105	182	361	712	1,020	1,094	954	500	122	17	-
1999.....	61	97	170	381	716	1,040	1,124	1,012	489	155	17	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Aortic aneurysm and dissection (I71)

##### Non-Hispanic Black, both sexes

2003.....	101	90	106	111	131	151	141	99	36	22	2	-
2002.....	97	99	93	107	137	143	142	94	48	20	5	-
2001.....	87	86	103	105	135	195	157	97	64	16	2	-
2000.....	68	95	101	129	150	188	143	113	64	20	1	-
1999.....	71	82	70	126	145	164	139	118	44	22	5	-

##### Non-Hispanic Black, male

2003.....	69	57	75	65	59	71	67	33	11	5	-	-
2002.....	72	73	65	58	74	66	53	27	15	2	1	-
2001.....	63	53	71	57	77	90	68	28	13	2	-	-
2000.....	51	61	62	68	83	99	54	35	21	4	-	-
1999.....	52	52	44	73	74	62	62	44	12	8	1	-

##### Non-Hispanic Black, female

2003.....	32	33	31	46	72	80	74	66	25	17	2	-
2002.....	25	26	28	49	63	77	89	67	33	18	4	-
2001.....	24	33	32	48	58	105	89	69	51	14	2	-
2000.....	17	34	39	61	67	89	89	78	43	16	1	-
1999.....	19	30	26	53	71	102	77	74	32	14	4	-

##### Origin not stated 2/, both sexes

2003.....	2	-	2	2	8	8	6	3	2	-	-	-
2002.....	1	3	3	6	3	12	6	4	2	-	-	-
2001.....	2	3	3	6	10	6	8	3	4	1	-	-
2000.....	2	1	3	3	7	11	5	7	3	-	-	-
1999.....	-	3	1	8	5	2	8	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Aortic aneurysm and dissection (I71)

Origin not stated 2/, male

2003.....	-	-	2	2	6	5	2	1	1	-	-	-
2002.....	1	3	2	5	2	5	3	3	1	-	-	-
2001.....	2	3	3	4	8	5	5	1	2	1	-	-
2000.....	2	1	2	1	5	7	1	2	2	-	-	-
1999.....	-	1	1	5	5	1	3	-	1	-	-	-

Origin not stated 2/, female

2003.....	2	-	-	-	2	3	4	2	1	-	-	-
2002.....	-	-	1	1	1	7	3	1	1	-	-	-
2001.....	-	-	-	2	2	1	3	2	2	-	-	-
2000.....	-	-	1	2	2	4	4	5	1	-	-	-
1999.....	-	2	-	3	-	1	5	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins, both sexes

2003.....	223	310	483	681	982	1,489	1,734	1,680	1,264	517	123	-
2002.....	208	330	496	733	1,095	1,513	1,714	1,755	1,237	504	133	-
2001.....	229	318	483	697	1,131	1,599	1,682	1,745	1,188	506	114	-
2000.....	220	281	462	779	1,135	1,619	1,768	1,685	1,233	544	101	1
1999.....	198	307	466	666	1,141	1,567	1,761	1,674	1,226	508	118	-

All origins, male

2003.....	125	155	255	419	499	778	762	596	351	107	18	-
2002.....	110	166	268	388	550	718	787	641	347	117	18	-
2001.....	110	168	255	376	590	754	772	633	313	110	16	-
2000.....	123	169	250	427	614	811	765	615	309	136	15	1
1999.....	98	167	264	358	611	779	735	559	313	89	20	-

All origins, female

2003.....	98	155	228	262	483	711	972	1,084	913	410	105	-
2002.....	98	164	228	345	545	795	927	1,114	890	387	115	-
2001.....	119	150	228	321	541	845	910	1,112	875	396	98	-
2000.....	97	112	212	352	521	808	1,003	1,070	924	408	86	-
1999.....	100	140	202	308	530	788	1,026	1,115	913	419	98	-

Hispanic, both sexes

2003.....	12	16	16	38	32	50	50	61	31	15	3	-
2002.....	14	17	25	31	55	58	61	53	36	15	7	-
2001.....	19	19	22	26	37	50	39	47	30	22	2	-
2000.....	9	19	22	42	54	46	32	35	34	20	3	-
1999.....	6	15	24	31	44	40	56	51	34	11	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Hispanic, male

2003.....	8	10	12	21	14	28	25	24	11	2	1	-
2002.....	8	5	10	16	25	30	30	14	14	4	3	-
2001.....	9	10	12	12	17	24	17	14	9	5	1	-
2000.....	4	9	13	18	28	24	16	14	12	5	2	-
1999.....	1	8	10	14	20	26	28	19	15	3	2	-

Hispanic, female

2003.....	4	6	4	17	18	22	25	37	20	13	2	-
2002.....	6	12	15	15	30	28	31	39	22	11	4	-
2001.....	10	9	10	14	20	26	22	33	21	17	1	-
2000.....	5	10	9	24	26	22	16	21	22	15	1	-
1999.....	5	7	14	17	24	14	28	32	19	8	2	-

Non-Hispanic 1/, both sexes

2003.....	210	294	466	642	945	1,436	1,681	1,617	1,233	499	120	-
2002.....	194	313	467	700	1,036	1,450	1,652	1,696	1,199	485	126	-
2001.....	208	299	459	669	1,093	1,546	1,639	1,690	1,155	484	112	-
2000.....	211	260	438	735	1,079	1,572	1,731	1,645	1,197	523	98	1
1999.....	191	291	440	634	1,092	1,524	1,702	1,618	1,188	495	114	-

Non-Hispanic 1/, male

2003.....	117	145	242	397	484	748	736	571	340	105	17	-
2002.....	102	161	258	372	523	685	756	625	333	113	15	-
2001.....	100	158	242	363	572	729	754	617	304	105	15	-
2000.....	119	159	236	408	584	786	748	599	297	130	13	1
1999.....	97	159	252	343	589	753	704	540	297	85	18	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic 1/, female

2003.....	93	149	224	245	461	688	945	1,046	893	394	103	-
2002.....	92	152	209	328	513	765	896	1,071	866	372	111	-
2001.....	108	141	217	306	521	817	885	1,073	851	379	97	-
2000.....	92	101	202	327	495	786	983	1,046	900	393	85	-
1999.....	94	132	188	291	503	771	998	1,078	891	410	96	-

Non-Hispanic White, both sexes

2003.....	151	242	367	514	800	1,239	1,445	1,429	1,098	437	97	-
2002.....	147	241	361	551	877	1,239	1,457	1,517	1,055	442	98	-
2001.....	159	231	354	554	906	1,331	1,422	1,496	1,034	411	80	-
2000.....	141	197	336	605	915	1,353	1,530	1,466	1,063	455	76	-
1999.....	145	222	356	518	934	1,315	1,488	1,444	1,045	432	93	-

Non-Hispanic White, male

2003.....	85	120	189	319	418	648	634	513	310	92	14	-
2002.....	78	123	200	301	444	602	670	558	287	103	12	-
2001.....	73	126	191	307	481	643	658	555	262	92	12	-
2000.....	79	124	187	344	504	686	670	543	263	113	9	-
1999.....	75	126	207	289	513	649	622	488	266	73	11	-

Non-Hispanic White, female

2003.....	66	122	178	195	382	591	811	916	788	345	83	-
2002.....	69	118	161	250	433	637	787	959	768	339	86	-
2001.....	86	105	163	247	425	688	764	941	772	319	68	-
2000.....	62	73	149	261	411	667	860	923	800	342	67	-
1999.....	70	96	149	229	421	666	866	956	779	359	82	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Non-Hispanic Black, both sexes

2003.....	55	44	85	116	134	171	220	167	115	51	22	-
2002.....	41	65	93	133	144	193	175	158	132	39	25	-
2001.....	42	64	95	105	181	199	195	174	113	68	29	-
2000.....	58	53	97	117	144	198	181	163	123	62	20	1
1999.....	37	64	74	108	138	193	193	158	132	59	19	-

Non-Hispanic Black, male

2003.....	28	23	45	69	59	86	93	51	21	9	3	-
2002.....	20	35	50	64	72	74	76	57	42	7	3	-
2001.....	23	29	44	51	87	76	90	53	36	11	3	-
2000.....	34	29	46	58	70	91	70	47	28	14	4	1
1999.....	18	30	39	49	70	94	72	44	27	12	7	-

Non-Hispanic Black, female

2003.....	27	21	40	47	75	85	127	116	94	42	19	-
2002.....	21	30	43	69	72	119	99	101	90	32	22	-
2001.....	19	35	51	54	94	123	105	121	77	57	26	-
2000.....	24	24	51	59	74	107	111	116	95	48	16	-
1999.....	19	34	35	59	68	99	121	114	105	47	12	-

Origin not stated 2/, both sexes

2003.....	1	-	1	1	5	3	3	2	-	3	-	-
2002.....	-	-	4	2	4	5	1	6	2	4	-	-
2001.....	2	-	2	2	1	3	4	8	3	-	-	-
2000.....	-	2	2	2	2	1	5	5	2	1	-	-
1999.....	1	1	2	1	5	3	3	5	4	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Origin not stated 2/, male

2003.....	-	-	1	1	1	2	1	1	-	-	-	-
2002.....	-	-	-	-	2	3	1	2	-	-	-	-
2001.....	1	-	1	1	1	1	1	2	-	-	-	-
2000.....	-	1	1	1	2	1	1	2	-	1	-	-
1999.....	-	-	2	1	2	-	3	-	1	1	-	-

Origin not stated 2/, female

2003.....	1	-	-	-	4	1	2	1	-	3	-	-
2002.....	-	-	4	2	2	2	-	4	2	4	-	-
2001.....	1	-	1	1	-	2	3	6	3	-	-	-
2000.....	-	1	1	1	-	-	4	3	2	-	-	-
1999.....	1	1	-	-	3	3	-	5	3	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

All origins, both sexes

2003.....	282	272	297	319	425	518	643	577	382	148	26	-
2002.....	280	302	293	322	375	552	701	576	390	134	23	-
2001.....	232	272	291	366	475	572	611	568	414	140	27	1
2000.....	266	231	285	326	469	573	588	571	343	149	21	-
1999.....	176	208	237	312	380	498	619	509	322	116	25	1

All origins, male

2003.....	165	166	154	160	188	214	223	163	82	27	4	-
2002.....	147	168	141	160	163	216	267	173	86	25	1	-
2001.....	135	149	155	174	231	254	234	164	97	19	6	1
2000.....	150	126	138	170	220	245	227	175	72	26	-	-
1999.....	96	106	118	163	187	225	238	155	66	19	8	-

All origins, female

2003.....	117	106	143	159	237	304	420	414	300	121	22	-
2002.....	133	134	152	162	212	336	434	403	304	109	22	-
2001.....	97	123	136	192	244	318	377	404	317	121	21	-
2000.....	116	105	147	156	249	328	361	396	271	123	21	-
1999.....	80	102	119	149	193	273	381	354	256	97	17	1

Hispanic, both sexes

2003.....	23	17	12	19	13	24	17	13	12	6	3	-
2002.....	18	14	13	13	24	18	20	18	11	4	-	-
2001.....	14	8	8	21	27	21	20	29	21	2	-	-
2000.....	20	8	15	17	18	28	16	18	11	7	1	-
1999.....	14	9	15	15	27	8	18	25	8	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Hispanic, male

2003.....	18	10	7	10	8	10	6	2	4	2	1	-
2002.....	10	7	9	8	8	9	7	3	2	-	-	-
2001.....	10	4	5	11	9	10	10	8	8	-	-	-
2000.....	11	4	9	7	9	12	10	10	1	2	-	-
1999.....	6	4	7	5	12	6	7	9	2	-	1	-

Hispanic, female

2003.....	5	7	5	9	5	14	11	11	8	4	2	-
2002.....	8	7	4	5	16	9	13	15	9	4	-	-
2001.....	4	4	3	10	18	11	10	21	13	2	-	-
2000.....	9	4	6	10	9	16	6	8	10	5	1	-
1999.....	8	5	8	10	15	2	11	16	6	1	-	-

Non-Hispanic 1/, both sexes

2003.....	259	254	283	299	410	492	625	563	370	142	23	-
2002.....	260	288	278	308	351	534	678	557	379	129	23	-
2001.....	214	264	280	342	446	549	587	538	393	137	27	1
2000.....	245	222	269	307	451	545	572	553	332	140	20	-
1999.....	162	198	222	296	352	488	600	482	313	115	24	1

Non-Hispanic 1/, male

2003.....	147	156	147	150	178	203	216	161	78	25	3	-
2002.....	135	161	131	151	155	207	259	170	84	25	1	-
2001.....	124	145	148	162	222	243	223	155	89	19	6	1
2000.....	138	121	129	162	211	233	217	165	71	24	-	-
1999.....	90	101	111	157	174	219	231	145	64	19	7	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic 1/, female

2003.....	112	98	136	149	232	289	409	402	292	117	20	-
2002.....	125	127	147	157	196	327	419	387	295	104	22	-
2001.....	90	119	132	180	224	306	364	383	304	118	21	-
2000.....	107	101	140	145	240	312	355	388	261	116	20	-
1999.....	72	97	111	139	178	269	369	337	249	96	17	1

Non-Hispanic White, both sexes

2003.....	198	195	218	236	339	404	548	502	327	124	18	-
2002.....	195	224	206	247	291	455	586	497	344	115	21	-
2001.....	161	205	209	270	372	459	528	490	349	125	23	-
2000.....	184	166	207	225	377	462	513	498	300	123	20	-
1999.....	117	167	175	227	297	419	530	424	282	106	23	1

Non-Hispanic White, male

2003.....	109	130	106	116	150	167	182	141	69	21	2	-
2002.....	104	123	101	119	129	178	218	151	73	21	1	-
2001.....	92	114	106	128	185	205	200	141	79	15	4	-
2000.....	105	89	99	124	183	203	195	148	63	20	-	-
1999.....	63	87	93	118	150	185	203	128	57	18	7	-

Non-Hispanic White, female

2003.....	89	65	112	120	189	237	366	361	258	103	16	-
2002.....	91	101	105	128	162	277	368	346	271	94	20	-
2001.....	69	91	103	142	187	254	328	349	270	110	19	-
2000.....	79	77	108	101	194	259	318	350	237	103	20	-
1999.....	54	80	82	109	147	234	327	296	225	88	16	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Non-Hispanic Black, both sexes

2003.....	55	57	61	58	66	80	69	51	41	16	5	-
2002.....	60	62	66	59	54	74	85	53	32	12	2	-
2001.....	46	55	67	64	65	79	54	41	39	12	3	1
2000.....	58	53	56	77	68	80	50	51	28	15	-	-
1999.....	41	27	42	62	47	64	61	51	28	8	1	-

Non-Hispanic Black, male

2003.....	34	26	38	31	27	34	28	18	8	4	1	-
2002.....	27	36	28	30	23	26	37	15	10	4	-	-
2001.....	28	30	42	31	31	34	19	11	8	4	2	1
2000.....	32	29	27	35	26	27	16	14	8	4	-	-
1999.....	24	11	17	36	20	31	24	13	5	1	-	-

Non-Hispanic Black, female

2003.....	21	31	23	27	39	46	41	33	33	12	4	-
2002.....	33	26	38	29	31	48	48	38	22	8	2	-
2001.....	18	25	25	33	34	45	35	30	31	8	1	-
2000.....	26	24	29	42	42	53	34	37	20	11	-	-
1999.....	17	16	25	26	27	33	37	38	23	7	1	-

Origin not stated 2/, both sexes

2003.....	-	1	2	1	2	2	1	1	-	-	-	-
2002.....	2	-	2	1	-	-	3	1	-	1	-	-
2001.....	4	-	3	3	2	2	4	1	-	1	-	-
2000.....	1	1	1	2	-	-	-	-	-	2	-	-
1999.....	-	1	-	1	1	2	1	2	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of circulatory system (I80-I99)

Origin not stated 2/, male

2003.....	-	-	-	-	2	1	1	-	-	-	-	-
2002.....	2	-	1	1	-	-	1	-	-	-	-	-
2001.....	1	-	2	1	-	1	1	1	-	-	-	-
2000.....	1	1	-	1	-	-	-	-	-	-	-	-
1999.....	-	1	-	1	1	-	-	1	-	-	-	-

Origin not stated 2/, female

2003.....	-	1	2	1	-	1	-	1	-	-	-	-
2002.....	-	-	1	-	-	-	2	1	-	1	-	-
2001.....	3	-	1	2	2	1	3	-	-	1	-	-
2000.....	-	-	1	1	-	-	-	-	-	2	-	-
1999.....	-	-	-	-	-	2	1	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Influenza and pneumonia (J10-J18)

##### All origins, both sexes

2003.....	1,222	1,292	1,838	2,588	4,243	7,698	11,744	13,653	11,200	5,172	1,372	2
2002.....	1,031	1,271	1,716	2,420	4,427	7,905	12,079	13,977	11,490	5,172	1,356	3
2001.....	991	1,155	1,549	2,323	4,327	7,510	11,167	13,382	10,692	4,836	1,281	5
2000.....	991	1,198	1,681	2,533	4,656	7,912	11,909	13,752	11,318	5,116	1,361	3
1999.....	899	1,098	1,527	2,382	4,479	7,776	11,416	13,499	11,151	5,159	1,420	2

##### All origins, male

2003.....	751	739	1,032	1,506	2,376	4,032	5,597	5,592	3,715	1,360	283	1
2002.....	619	742	996	1,384	2,466	4,114	5,781	5,716	3,875	1,352	256	2
2001.....	602	686	901	1,338	2,394	3,995	5,299	5,373	3,587	1,282	260	4
2000.....	602	710	970	1,447	2,566	4,084	5,667	5,575	3,724	1,373	273	2
1999.....	559	611	851	1,394	2,499	4,020	5,438	5,501	3,595	1,295	285	1

##### All origins, female

2003.....	471	553	806	1,082	1,867	3,666	6,147	8,061	7,485	3,812	1,089	1
2002.....	412	529	720	1,036	1,961	3,791	6,298	8,261	7,615	3,820	1,100	1
2001.....	389	469	648	985	1,933	3,515	5,868	8,009	7,105	3,554	1,021	1
2000.....	389	488	711	1,086	2,090	3,828	6,242	8,177	7,594	3,743	1,088	1
1999.....	340	487	676	988	1,980	3,756	5,978	7,998	7,556	3,864	1,135	1

##### Hispanic, both sexes

2003.....	88	96	128	168	260	371	489	415	344	188	49	-
2002.....	63	95	112	150	249	385	470	443	358	177	43	-
2001.....	65	72	96	140	262	346	415	465	382	173	36	1
2000.....	66	82	109	138	267	313	377	440	339	161	30	-
1999.....	56	58	97	143	185	252	317	383	296	150	47	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Influenza and pneumonia (J10-J18)

##### Hispanic, male

2003.....	62	53	62	95	140	184	248	171	128	57	19	-
2002.....	45	60	70	88	141	189	233	186	137	55	14	-
2001.....	38	50	49	71	133	180	208	177	145	69	11	1
2000.....	41	50	69	75	143	166	201	164	144	62	9	-
1999.....	39	32	48	82	105	129	155	155	118	49	12	-

##### Hispanic, female

2003.....	26	43	66	73	120	187	241	244	216	131	30	-
2002.....	18	35	42	62	108	196	237	257	221	122	29	-
2001.....	27	22	47	69	129	166	207	288	237	104	25	-
2000.....	25	32	40	63	124	147	176	276	195	99	21	-
1999.....	17	26	49	61	80	123	162	228	178	101	35	-

##### Non-Hispanic 1/, both sexes

2003.....	1,127	1,189	1,694	2,403	3,970	7,310	11,234	13,217	10,828	4,971	1,321	2
2002.....	963	1,172	1,601	2,260	4,164	7,492	11,567	13,504	11,101	4,983	1,310	3
2001.....	917	1,072	1,444	2,172	4,037	7,140	10,723	12,879	10,292	4,652	1,239	2
2000.....	916	1,107	1,567	2,378	4,372	7,565	11,486	13,278	10,947	4,938	1,329	1
1999.....	836	1,029	1,417	2,223	4,276	7,494	11,059	13,060	10,823	4,995	1,369	1

##### Non-Hispanic 1/, male

2003.....	684	680	961	1,399	2,227	3,838	5,340	5,413	3,575	1,300	263	1
2002.....	570	679	924	1,287	2,313	3,912	5,525	5,523	3,726	1,293	240	2
2001.....	559	630	845	1,259	2,241	3,802	5,079	5,181	3,436	1,210	248	1
2000.....	553	652	897	1,361	2,411	3,899	5,440	5,395	3,571	1,306	264	1
1999.....	513	573	794	1,303	2,384	3,875	5,263	5,318	3,467	1,241	272	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Influenza and pneumonia (J10-J18)

##### Non-Hispanic 1/, female

2003.....	443	509	733	1,004	1,743	3,472	5,894	7,804	7,253	3,671	1,058	1
2002.....	393	493	677	973	1,851	3,580	6,042	7,981	7,375	3,690	1,070	1
2001.....	358	442	599	913	1,796	3,338	5,644	7,698	6,856	3,442	991	1
2000.....	363	455	670	1,017	1,961	3,666	6,046	7,883	7,376	3,632	1,065	-
1999.....	323	456	623	920	1,892	3,619	5,796	7,742	7,356	3,754	1,097	1

##### Non-Hispanic White, both sexes

2003.....	823	894	1,316	1,929	3,333	6,406	10,139	12,184	9,962	4,577	1,163	1
2002.....	718	887	1,265	1,866	3,553	6,568	10,433	12,381	10,228	4,585	1,179	3
2001.....	677	831	1,132	1,773	3,441	6,188	9,649	11,821	9,463	4,242	1,076	1
2000.....	664	844	1,217	1,884	3,662	6,696	10,372	12,218	10,117	4,554	1,184	1
1999.....	606	797	1,097	1,843	3,638	6,686	10,099	11,998	9,993	4,610	1,237	1

##### Non-Hispanic White, male

2003.....	501	503	716	1,109	1,857	3,348	4,793	4,993	3,257	1,168	221	-
2002.....	424	508	720	1,059	1,991	3,427	4,975	5,057	3,416	1,164	217	2
2001.....	396	484	648	1,013	1,906	3,268	4,562	4,746	3,119	1,089	212	1
2000.....	387	480	696	1,065	2,019	3,446	4,900	4,952	3,263	1,194	232	1
1999.....	361	436	613	1,075	2,012	3,454	4,801	4,862	3,160	1,139	249	-

##### Non-Hispanic White, female

2003.....	322	391	600	820	1,476	3,058	5,346	7,191	6,705	3,409	942	1
2002.....	294	379	545	807	1,562	3,141	5,458	7,324	6,812	3,421	962	1
2001.....	281	347	484	760	1,535	2,920	5,087	7,075	6,344	3,153	864	-
2000.....	277	364	521	819	1,643	3,250	5,472	7,266	6,854	3,360	952	-
1999.....	245	361	484	768	1,626	3,232	5,298	7,136	6,833	3,471	988	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Influenza and pneumonia (J10-J18)

##### Non-Hispanic Black, both sexes

2003.....	266	247	318	391	524	712	802	750	635	278	113	1
2002.....	210	254	300	327	510	708	876	849	651	310	107	-
2001.....	211	213	258	315	481	756	815	784	629	320	135	-
2000.....	226	237	294	408	576	689	861	815	653	311	122	-
1999.....	205	208	278	327	553	659	751	869	671	311	112	-

##### Non-Hispanic Black, male

2003.....	161	152	204	245	302	384	381	283	214	76	30	1
2002.....	123	158	180	188	266	373	416	318	208	92	18	-
2001.....	142	129	164	195	277	429	379	289	224	82	27	-
2000.....	147	155	167	244	318	360	403	321	218	86	25	-
1999.....	137	125	157	198	320	336	352	352	224	81	19	-

##### Non-Hispanic Black, female

2003.....	105	95	114	146	222	328	421	467	421	202	83	-
2002.....	87	96	120	139	244	335	460	531	443	218	89	-
2001.....	69	84	94	120	204	327	436	495	405	238	108	-
2000.....	79	82	127	164	258	329	458	494	435	225	97	-
1999.....	68	83	121	129	233	323	399	517	447	230	93	-

##### Origin not stated 2/, both sexes

2003.....	7	7	16	17	13	17	21	21	28	13	2	-
2002.....	5	4	3	10	14	28	42	30	31	12	3	-
2001.....	9	11	9	11	28	24	29	38	18	11	6	2
2000.....	9	9	5	17	17	34	46	34	32	17	2	2
1999.....	7	11	13	16	18	30	40	56	32	14	4	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza and pneumonia (J10-J18)

Origin not stated 2/, male

2003.....	5	6	9	12	9	10	9	8	12	3	1	-
2002.....	4	3	2	9	12	13	23	7	12	4	2	-
2001.....	5	6	7	8	20	13	12	15	6	3	1	2
2000.....	8	8	4	11	12	19	26	16	9	5	-	1
1999.....	7	6	9	9	10	16	20	28	10	5	1	1

Origin not stated 2/, female

2003.....	2	1	7	5	4	7	12	13	16	10	1	-
2002.....	1	1	1	1	2	15	19	23	19	8	1	-
2001.....	4	5	2	3	8	11	17	23	12	8	5	-
2000.....	1	1	1	6	5	15	20	18	23	12	2	1
1999.....	-	5	4	7	8	14	20	28	22	9	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Influenza (J10-J11)

##### All origins, both sexes

2003.....	32	47	38	72	111	200	257	341	303	128	31	-
2002.....	9	13	17	19	37	59	118	174	144	59	27	-
2001.....	7	8	13	10	11	18	38	37	39	14	6	-
2000.....	24	39	54	76	123	189	303	372	305	130	48	-
1999.....	13	29	42	60	104	154	297	368	303	162	48	-

##### All origins, male

2003.....	18	26	16	38	51	89	100	132	84	26	3	-
2002.....	5	7	13	13	19	28	53	66	45	10	3	-
2001.....	4	4	5	3	5	6	14	15	8	4	2	-
2000.....	14	20	24	43	63	81	115	127	86	35	6	-
1999.....	7	13	24	27	47	71	131	127	70	35	6	-

##### All origins, female

2003.....	14	21	22	34	60	111	157	209	219	102	28	-
2002.....	4	6	4	6	18	31	65	108	99	49	24	-
2001.....	3	4	8	7	6	12	24	22	31	10	4	-
2000.....	10	19	30	33	60	108	188	245	219	95	42	-
1999.....	6	16	18	33	57	83	166	241	233	127	42	-

##### Hispanic, both sexes

2003.....	1	3	3	3	-	4	8	1	5	3	1	-
2002.....	-	-	-	1	1	2	1	-	5	-	-	-
2001.....	-	-	-	-	1	3	2	-	-	-	-	-
2000.....	-	1	1	3	1	3	7	10	3	3	-	-
1999.....	2	2	3	4	4	5	4	3	5	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Influenza (J10-J11)

## Hispanic, male

2003.....	-	1	1	3	-	1	6	1	1	2	-	-
2002.....	-	-	-	1	-	-	1	-	1	-	-	-
2001.....	-	-	-	-	-	-	2	-	-	-	-	-
2000.....	-	1	1	2	-	1	4	3	1	2	-	-
1999.....	2	2	2	4	3	1	1	-	-	1	-	-

## Hispanic, female

2003.....	1	2	2	-	-	3	2	-	4	1	1	-
2002.....	-	-	-	-	1	2	-	-	4	-	-	-
2001.....	-	-	-	-	1	3	-	-	-	-	-	-
2000.....	-	-	-	1	1	2	3	7	2	1	-	-
1999.....	-	-	1	-	1	4	3	3	5	1	-	-

## Non-Hispanic 1/, both sexes

2003.....	31	44	34	69	111	196	247	340	298	125	30	-
2002.....	9	13	17	18	36	56	117	174	137	58	27	-
2001.....	7	8	13	10	10	15	36	37	39	14	6	-
2000.....	24	38	53	73	122	186	295	361	302	127	48	-
1999.....	11	27	39	55	99	149	290	364	296	160	48	-

## Non-Hispanic 1/, male

2003.....	18	25	14	35	51	88	93	131	83	24	3	-
2002.....	5	7	13	12	19	27	52	66	43	10	3	-
2001.....	4	4	5	3	5	6	12	15	8	4	2	-
2000.....	14	19	23	41	63	80	110	124	85	33	6	-
1999.....	5	11	22	23	43	70	130	127	70	34	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

Non-Hispanic 1/, female

2003.....	13	19	20	34	60	108	154	209	215	101	27	-
2002.....	4	6	4	6	17	29	65	108	94	48	24	-
2001.....	3	4	8	7	5	9	24	22	31	10	4	-
2000.....	10	19	30	32	59	106	185	237	217	94	42	-
1999.....	6	16	17	32	56	79	160	237	226	126	42	-

Non-Hispanic White, both sexes

2003.....	26	38	31	61	106	187	239	329	287	120	28	-
2002.....	8	11	14	12	33	54	114	172	129	57	26	-
2001.....	7	7	10	8	10	15	32	34	38	13	6	-
2000.....	22	36	49	67	109	177	281	342	292	122	47	-
1999.....	8	24	36	51	88	141	276	352	291	155	46	-

Non-Hispanic White, male

2003.....	16	21	12	32	49	84	91	128	82	22	3	-
2002.....	5	5	11	8	19	26	50	65	42	10	3	-
2001.....	4	3	3	1	5	6	12	14	7	4	2	-
2000.....	12	17	22	38	59	74	107	116	82	30	6	-
1999.....	4	11	21	20	39	69	123	124	70	33	6	-

Non-Hispanic White, female

2003.....	10	17	19	29	57	103	148	201	205	98	25	-
2002.....	3	6	3	4	14	28	64	107	87	47	23	-
2001.....	3	4	7	7	5	9	20	20	31	9	4	-
2000.....	10	19	27	29	50	103	174	226	210	92	41	-
1999.....	4	13	15	31	49	72	153	228	221	122	40	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

Non-Hispanic Black, both sexes

2003.....	5	5	2	6	4	5	7	8	8	3	1	-
2002.....	-	1	3	4	2	1	3	1	5	1	1	-
2001.....	-	1	2	2	-	-	4	3	1	1	-	-
2000.....	2	2	4	5	11	5	8	17	8	4	1	-
1999.....	2	3	2	2	10	5	10	8	4	4	2	-

Non-Hispanic Black, male

2003.....	2	4	1	3	1	2	1	2	-	1	-	-
2002.....	-	1	2	3	-	-	2	1	1	-	-	-
2001.....	-	1	2	2	-	-	-	1	1	-	-	-
2000.....	2	2	1	2	3	4	1	7	1	3	-	-
1999.....	1	-	1	2	4	1	5	2	-	1	-	-

Non-Hispanic Black, female

2003.....	3	1	1	3	3	3	6	6	8	2	1	-
2002.....	-	-	1	1	2	1	1	-	4	1	1	-
2001.....	-	-	-	-	-	-	4	2	-	1	-	-
2000.....	-	-	3	3	8	1	7	10	7	1	1	-
1999.....	1	3	1	-	6	4	5	6	4	3	2	-

Origin not stated 2/, both sexes

2003.....	-	-	1	-	-	-	2	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	2	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	-	1	1	-	3	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Influenza (J10-J11)

Origin not stated 2/, male

2003.....	-	-	1	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	1	-	-	1	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	1	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	1	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	1	-	-	3	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>Pneumonia (J12-J18)</b>												
<b>All origins, both sexes</b>												
2003.....	1,190	1,245	1,800	2,516	4,132	7,498	11,487	13,312	10,897	5,044	1,341	2
2002.....	1,022	1,258	1,699	2,401	4,390	7,846	11,961	13,803	11,346	5,113	1,329	3
2001.....	984	1,147	1,536	2,313	4,316	7,492	11,129	13,345	10,653	4,822	1,275	5
2000.....	967	1,159	1,627	2,457	4,533	7,723	11,606	13,380	11,013	4,986	1,313	3
1999.....	886	1,069	1,485	2,322	4,375	7,622	11,119	13,131	10,848	4,997	1,372	2
<b>All origins, male</b>												
2003.....	733	713	1,016	1,468	2,325	3,943	5,497	5,460	3,631	1,334	280	1
2002.....	614	735	983	1,371	2,447	4,086	5,728	5,650	3,830	1,342	253	2
2001.....	598	682	896	1,335	2,389	3,989	5,285	5,358	3,579	1,278	258	4
2000.....	588	690	946	1,404	2,503	4,003	5,552	5,448	3,638	1,338	267	2
1999.....	552	598	827	1,367	2,452	3,949	5,307	5,374	3,525	1,260	279	1
<b>All origins, female</b>												
2003.....	457	532	784	1,048	1,807	3,555	5,990	7,852	7,266	3,710	1,061	1
2002.....	408	523	716	1,030	1,943	3,760	6,233	8,153	7,516	3,771	1,076	1
2001.....	386	465	640	978	1,927	3,503	5,844	7,987	7,074	3,544	1,017	1
2000.....	379	469	681	1,053	2,030	3,720	6,054	7,932	7,375	3,648	1,046	1
1999.....	334	471	658	955	1,923	3,673	5,812	7,757	7,323	3,737	1,093	1
<b>Hispanic, both sexes</b>												
2003.....	87	93	125	165	260	367	481	414	339	185	48	-
2002.....	63	95	112	149	248	383	469	443	353	177	43	-
2001.....	65	72	96	140	261	343	413	465	382	173	36	1
2000.....	66	81	108	135	266	310	370	430	336	158	30	-
1999.....	54	56	94	139	181	247	313	380	291	148	47	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>Pneumonia (J12-J18)</b>												
<b>Hispanic, male</b>												
2003.....	62	52	61	92	140	183	242	170	127	55	19	-
2002.....	45	60	70	87	141	189	232	186	136	55	14	-
2001.....	38	50	49	71	133	180	206	177	145	69	11	1
2000.....	41	49	68	73	143	165	197	161	143	60	9	-
1999.....	37	30	46	78	102	128	154	155	118	48	12	-
<b>Hispanic, female</b>												
2003.....	25	41	64	73	120	184	239	244	212	130	29	-
2002.....	18	35	42	62	107	194	237	257	217	122	29	-
2001.....	27	22	47	69	128	163	207	288	237	104	25	-
2000.....	25	32	40	62	123	145	173	269	193	98	21	-
1999.....	17	26	48	61	79	119	159	225	173	100	35	-
<b>Non-Hispanic 1/, both sexes</b>												
2003.....	1,096	1,145	1,660	2,334	3,859	7,114	10,987	12,877	10,530	4,846	1,291	2
2002.....	954	1,159	1,584	2,242	4,128	7,436	11,450	13,330	10,964	4,925	1,283	3
2001.....	910	1,064	1,431	2,162	4,027	7,125	10,687	12,842	10,253	4,638	1,233	2
2000.....	892	1,069	1,514	2,305	4,250	7,379	11,191	12,917	10,645	4,811	1,281	1
1999.....	825	1,002	1,378	2,168	4,177	7,345	10,769	12,696	10,527	4,835	1,321	1
<b>Non-Hispanic 1/, male</b>												
2003.....	666	655	947	1,364	2,176	3,750	5,247	5,282	3,492	1,276	260	1
2002.....	565	672	911	1,275	2,294	3,885	5,473	5,457	3,683	1,283	237	2
2001.....	555	626	840	1,256	2,236	3,796	5,067	5,166	3,428	1,206	246	1
2000.....	539	633	874	1,320	2,348	3,819	5,330	5,271	3,486	1,273	258	1
1999.....	508	562	772	1,280	2,341	3,805	5,133	5,191	3,397	1,207	266	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pneumonia (J12-J18)												
Non-Hispanic 1/, female												
2003.....	430	490	713	970	1,683	3,364	5,740	7,595	7,038	3,570	1,031	1
2002.....	389	487	673	967	1,834	3,551	5,977	7,873	7,281	3,642	1,046	1
2001.....	355	438	591	906	1,791	3,329	5,620	7,676	6,825	3,432	987	1
2000.....	353	436	640	985	1,902	3,560	5,861	7,646	7,159	3,538	1,023	-
1999.....	317	440	606	888	1,836	3,540	5,636	7,505	7,130	3,628	1,055	1
Non-Hispanic White, both sexes												
2003.....	797	856	1,285	1,868	3,227	6,219	9,900	11,855	9,675	4,457	1,135	1
2002.....	710	876	1,251	1,854	3,520	6,514	10,319	12,209	10,099	4,528	1,153	3
2001.....	670	824	1,122	1,765	3,431	6,173	9,617	11,787	9,425	4,229	1,070	1
2000.....	642	808	1,168	1,817	3,553	6,519	10,091	11,876	9,825	4,432	1,137	1
1999.....	598	773	1,061	1,792	3,550	6,545	9,823	11,646	9,702	4,455	1,191	1
Non-Hispanic White, male												
2003.....	485	482	704	1,077	1,808	3,264	4,702	4,865	3,175	1,146	218	-
2002.....	419	503	709	1,051	1,972	3,401	4,925	4,992	3,374	1,154	214	2
2001.....	392	481	645	1,012	1,901	3,262	4,550	4,732	3,112	1,085	210	1
2000.....	375	463	674	1,027	1,960	3,372	4,793	4,836	3,181	1,164	226	1
1999.....	357	425	592	1,055	1,973	3,385	4,678	4,738	3,090	1,106	243	-
Non-Hispanic White, female												
2003.....	312	374	581	791	1,419	2,955	5,198	6,990	6,500	3,311	917	1
2002.....	291	373	542	803	1,548	3,113	5,394	7,217	6,725	3,374	939	1
2001.....	278	343	477	753	1,530	2,911	5,067	7,055	6,313	3,144	860	-
2000.....	267	345	494	790	1,593	3,147	5,298	7,040	6,644	3,268	911	-
1999.....	241	348	469	737	1,577	3,160	5,145	6,908	6,612	3,349	948	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

Non-Hispanic Black, both sexes

2003.....	261	242	316	385	520	707	795	742	627	275	112	1
2002.....	210	253	297	323	508	707	873	848	646	309	106	-
2001.....	211	212	256	313	481	756	811	781	628	319	135	-
2000.....	224	235	290	403	565	684	853	798	645	307	121	-
1999.....	203	205	276	325	543	654	741	861	667	307	110	-

Non-Hispanic Black, male

2003.....	159	148	203	242	301	382	380	281	214	75	30	1
2002.....	123	157	178	185	266	373	414	317	207	92	18	-
2001.....	142	128	162	193	277	429	379	288	223	82	27	-
2000.....	145	153	166	242	315	356	402	314	217	83	25	-
1999.....	136	125	156	196	316	335	347	350	224	80	19	-

Non-Hispanic Black, female

2003.....	102	94	113	143	219	325	415	461	413	200	82	-
2002.....	87	96	119	138	242	334	459	531	439	217	88	-
2001.....	69	84	94	120	204	327	432	493	405	237	108	-
2000.....	79	82	124	161	250	328	451	484	428	224	96	-
1999.....	67	80	120	129	227	319	394	511	443	227	91	-

Origin not stated 2/, both sexes

2003.....	7	7	15	17	13	17	19	21	28	13	2	-
2002.....	5	4	3	10	14	27	42	30	29	11	3	-
2001.....	9	11	9	11	28	24	29	38	18	11	6	2
2000.....	9	9	5	17	17	34	45	33	32	17	2	2
1999.....	7	11	13	15	17	30	37	55	30	14	4	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonia (J12-J18)

Origin not stated 2/, male

2003.....	5	6	8	12	9	10	8	8	12	3	1	-
2002.....	4	3	2	9	12	12	23	7	11	4	2	-
2001.....	5	6	7	8	20	13	12	15	6	3	1	2
2000.....	8	8	4	11	12	19	25	16	9	5	-	1
1999.....	7	6	9	9	9	16	20	28	10	5	1	1

Origin not stated 2/, female

2003.....	2	1	7	5	4	7	11	13	16	10	1	-
2002.....	1	1	1	1	2	15	19	23	18	7	1	-
2001.....	4	5	2	3	8	11	17	23	12	8	5	-
2000.....	1	1	1	6	5	15	20	17	23	12	2	1
1999.....	-	5	4	6	8	14	17	27	20	9	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute lower respiratory infections (J20-J22)

All origins, both sexes

2003.....	12	11	10	13	17	35	45	57	60	32	15	-
2002.....	12	11	7	10	23	21	45	70	42	27	13	-
2001.....	10	4	12	10	18	24	34	60	55	25	5	-
2000.....	4	6	4	19	24	41	41	80	77	23	12	-
1999.....	8	8	13	14	21	44	63	86	96	46	21	-

All origins, male

2003.....	6	6	6	11	6	17	18	14	15	4	2	-
2002.....	9	3	3	4	9	11	16	21	11	5	2	-
2001.....	6	1	5	3	7	10	10	15	10	4	1	-
2000.....	2	2	1	12	10	21	14	26	15	6	-	-
1999.....	3	4	6	6	11	22	29	19	18	6	5	-

All origins, female

2003.....	6	5	4	2	11	18	27	43	45	28	13	-
2002.....	3	8	4	6	14	10	29	49	31	22	11	-
2001.....	4	3	7	7	11	14	24	45	45	21	4	-
2000.....	2	4	3	7	14	20	27	54	62	17	12	-
1999.....	5	4	7	8	10	22	34	67	78	40	16	-

Hispanic, both sexes

2003.....	-	1	-	1	-	1	1	2	1	-	-	-
2002.....	-	1	1	-	-	-	-	2	1	1	-	-
2001.....	2	-	-	1	1	1	3	1	1	1	1	-
2000.....	-	-	-	-	1	1	1	2	5	1	-	-
1999.....	1	-	-	-	1	-	-	4	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute lower respiratory infections (J20-J22)

Hispanic, male

2003.....	-	1	-	-	-	1	-	1	1	-	-	-
2002.....	-	1	-	-	-	-	-	1	-	-	-	-
2001.....	1	-	-	-	1	-	1	-	-	-	-	-
2000.....	-	-	-	-	1	1	1	2	-	1	-	-
1999.....	-	-	-	-	1	-	-	1	-	-	-	-

Hispanic, female

2003.....	-	-	-	1	-	-	1	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	1	1	1	-	-
2001.....	1	-	-	1	-	1	2	1	1	1	1	-
2000.....	-	-	-	-	-	-	-	-	5	-	-	-
1999.....	1	-	-	-	-	-	-	3	1	-	-	-

Non-Hispanic 1/, both sexes

2003.....	12	9	10	12	17	33	44	55	59	32	15	-
2002.....	12	10	6	10	23	21	45	68	41	26	13	-
2001.....	8	4	12	9	17	23	31	59	54	24	4	-
2000.....	4	6	4	19	23	40	39	77	72	22	12	-
1999.....	7	8	13	14	20	44	63	82	95	46	21	-

Non-Hispanic 1/, male

2003.....	6	4	6	11	6	15	18	13	14	4	2	-
2002.....	9	2	3	4	9	11	16	20	11	5	2	-
2001.....	5	1	5	3	6	10	9	15	10	4	1	-
2000.....	2	2	1	12	9	20	12	24	15	5	-	-
1999.....	3	4	6	6	10	22	29	18	18	6	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other acute lower respiratory infections (J20-J22)

Non-Hispanic 1/, female

2003.....	6	5	4	1	11	18	26	42	45	28	13	-
2002.....	3	8	3	6	14	10	29	48	30	21	11	-
2001.....	3	3	7	6	11	13	22	44	44	20	3	-
2000.....	2	4	3	7	14	20	27	53	57	17	12	-
1999.....	4	4	7	8	10	22	34	64	77	40	16	-

Non-Hispanic White, both sexes

2003.....	11	7	9	10	15	32	41	54	58	30	14	-
2002.....	10	8	4	9	19	19	40	62	40	26	12	-
2001.....	4	3	11	8	12	20	30	57	51	22	4	-
2000.....	4	6	4	18	22	37	37	72	67	21	12	-
1999.....	4	6	11	11	15	41	58	77	92	44	19	-

Non-Hispanic White, male

2003.....	6	3	5	9	6	14	17	13	14	4	2	-
2002.....	8	2	1	3	9	11	14	18	11	5	2	-
2001.....	2	1	4	3	4	8	9	14	9	4	1	-
2000.....	2	2	1	11	8	20	10	22	12	5	-	-
1999.....	3	3	6	4	7	21	27	15	17	6	4	-

Non-Hispanic White, female

2003.....	5	4	4	1	9	18	24	41	44	26	12	-
2002.....	2	6	3	6	10	8	26	44	29	21	10	-
2001.....	2	2	7	5	8	12	21	43	42	18	3	-
2000.....	2	4	3	7	14	17	27	50	55	16	12	-
1999.....	1	3	5	7	8	20	31	62	75	38	15	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

All origins, both sexes

2003.....	9	9	8	11	11	19	26	38	36	19	8	-
2002.....	11	8	5	10	13	15	24	39	31	15	9	-
2001.....	8	3	11	9	14	18	21	38	36	13	4	-
2000.....	1	4	3	15	16	30	23	45	44	15	7	-
1999.....	7	6	12	13	16	28	41	54	55	30	11	-

All origins, male

2003.....	5	5	6	9	3	7	10	6	9	3	1	-
2002.....	8	3	2	4	6	8	10	12	7	3	1	-
2001.....	5	1	5	3	5	6	4	9	5	2	1	-
2000.....	1	2	1	9	5	18	9	14	10	5	-	-
1999.....	2	3	5	6	9	15	19	12	12	5	2	-

All origins, female

2003.....	4	4	2	2	8	12	16	32	27	16	7	-
2002.....	3	5	3	6	7	7	14	27	24	12	8	-
2001.....	3	2	6	6	9	12	17	29	31	11	3	-
2000.....	-	2	2	6	11	12	14	31	34	10	7	-
1999.....	5	3	7	7	7	13	22	42	43	25	9	-

Hispanic, both sexes

2003.....	-	1	-	1	-	1	1	2	1	-	-	-
2002.....	-	1	1	-	-	-	-	2	1	-	-	-
2001.....	2	-	-	1	-	1	1	-	-	1	1	-
2000.....	-	-	-	-	1	1	1	2	3	1	-	-
1999.....	1	-	-	-	1	-	-	3	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Hispanic, male

2003.....	-	1	-	-	-	1	-	1	1	-	-	-
2002.....	-	1	-	-	-	-	-	1	-	-	-	-
2001.....	1	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	1	1	1	2	-	1	-	-
1999.....	-	-	-	-	1	-	-	1	-	-	-	-

Hispanic, female

2003.....	-	-	-	1	-	-	1	1	-	-	-	-
2002.....	-	-	1	-	-	-	-	1	1	-	-	-
2001.....	1	-	-	1	-	1	1	-	-	1	1	-
2000.....	-	-	-	-	-	-	-	-	3	-	-	-
1999.....	1	-	-	-	-	-	-	2	1	-	-	-

Non-Hispanic 1/, both sexes

2003.....	9	7	8	10	11	18	25	36	35	19	8	-
2002.....	11	7	4	10	13	15	24	37	30	15	9	-
2001.....	6	3	11	8	14	17	20	38	36	12	3	-
2000.....	1	4	3	15	15	29	22	43	41	14	7	-
1999.....	6	6	12	13	15	28	41	51	54	30	11	-

Non-Hispanic 1/, male

2003.....	5	3	6	9	3	6	10	5	8	3	1	-
2002.....	8	2	2	4	6	8	10	11	7	3	1	-
2001.....	4	1	5	3	5	6	4	9	5	2	1	-
2000.....	1	2	1	9	4	17	8	12	10	4	-	-
1999.....	2	3	5	6	8	15	19	11	12	5	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute bronchitis and bronchiolitis (J20-J21)

Non-Hispanic 1/, female

2003.....	4	4	2	1	8	12	15	31	27	16	7	-
2002.....	3	5	2	6	7	7	14	26	23	12	8	-
2001.....	2	2	6	5	9	11	16	29	31	10	2	-
2000.....	-	2	2	6	11	12	14	31	31	10	7	-
1999.....	4	3	7	7	7	13	22	40	42	25	9	-

Non-Hispanic White, both sexes

2003.....	9	5	7	8	9	18	23	35	34	18	8	-
2002.....	9	5	3	9	12	14	22	32	29	15	8	-
2001.....	2	2	10	7	10	14	20	37	33	10	3	-
2000.....	1	4	3	14	14	26	21	40	39	13	7	-
1999.....	3	4	10	10	10	25	38	48	52	29	11	-

Non-Hispanic White, male

2003.....	5	2	5	7	3	6	9	5	8	3	1	-
2002.....	7	2	1	3	6	8	9	9	7	3	1	-
2001.....	1	1	4	3	3	4	4	9	4	2	1	-
2000.....	1	2	1	8	3	17	7	12	8	4	-	-
1999.....	2	2	5	4	5	14	18	9	11	5	2	-

Non-Hispanic White, female

2003.....	4	3	2	1	6	12	14	30	26	15	7	-
2002.....	2	3	2	6	6	6	13	23	22	12	7	-
2001.....	1	1	6	4	7	10	16	28	29	8	2	-
2000.....	-	2	2	6	11	9	14	28	31	9	7	-
1999.....	1	2	5	6	5	11	20	39	41	24	9	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Unspecified acute lower respiratory infection (J22)

All origins, both sexes

2003.....	3	2	2	2	6	16	19	19	24	13	7	-
2002.....	1	3	2	-	10	6	21	31	11	12	4	-
2001.....	2	1	1	1	4	6	13	22	19	12	1	-
2000.....	3	2	1	4	8	11	18	35	33	8	5	-
1999.....	1	2	1	1	5	16	22	32	41	16	10	-

All origins, male

2003.....	1	1	-	2	3	10	8	8	6	1	1	-
2002.....	1	-	1	-	3	3	6	9	4	2	1	-
2001.....	1	-	-	-	2	4	6	6	5	2	-	-
2000.....	1	-	-	3	5	3	5	12	5	1	-	-
1999.....	1	1	1	-	2	7	10	7	6	1	3	-

All origins, female

2003.....	2	1	2	-	3	6	11	11	18	12	6	-
2002.....	-	3	1	-	7	3	15	22	7	10	3	-
2001.....	1	1	1	1	2	2	7	16	14	10	1	-
2000.....	2	2	1	1	3	8	13	23	28	7	5	-
1999.....	-	1	-	1	3	9	12	25	35	15	7	-

Hispanic, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	1	-	2	1	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Unspecified acute lower respiratory infection (J22)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	1	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	1	1	1	-	-	-
2000.....	-	-	-	-	-	-	-	-	2	-	-	-
1999.....	-	-	-	-	-	-	-	1	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	3	2	2	2	6	15	19	19	24	13	7	-
2002.....	1	3	2	-	10	6	21	31	11	11	4	-
2001.....	2	1	1	1	3	6	11	21	18	12	1	-
2000.....	3	2	1	4	8	11	17	34	31	8	5	-
1999.....	1	2	1	1	5	16	22	31	41	16	10	-

Non-Hispanic 1/, male

2003.....	1	1	-	2	3	9	8	8	6	1	1	-
2002.....	1	-	1	-	3	3	6	9	4	2	1	-
2001.....	1	-	-	-	1	4	5	6	5	2	-	-
2000.....	1	-	-	3	5	3	4	12	5	1	-	-
1999.....	1	1	1	-	2	7	10	7	6	1	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Unspecified acute lower respiratory infection (J22)

Non-Hispanic 1/, female

2003.....	2	1	2	-	3	6	11	11	18	12	6	-
2002.....	-	3	1	-	7	3	15	22	7	9	3	-
2001.....	1	1	1	1	2	2	6	15	13	10	1	-
2000.....	2	2	1	1	3	8	13	22	26	7	5	-
1999.....	-	1	-	1	3	9	12	24	35	15	7	-

Non-Hispanic White, both sexes

2003.....	2	2	2	2	6	14	18	19	24	12	6	-
2002.....	1	3	1	-	7	5	18	30	11	11	4	-
2001.....	2	1	1	1	2	6	10	20	18	12	1	-
2000.....	3	2	1	4	8	11	16	32	28	8	5	-
1999.....	1	2	1	1	5	16	20	29	40	15	8	-

Non-Hispanic White, male

2003.....	1	1	-	2	3	8	8	8	6	1	1	-
2002.....	1	-	-	-	3	3	5	9	4	2	1	-
2001.....	1	-	-	-	1	4	5	5	5	2	-	-
2000.....	1	-	-	3	5	3	3	10	4	1	-	-
1999.....	1	1	1	-	2	7	9	6	6	1	2	-

Non-Hispanic White, female

2003.....	1	1	2	-	3	6	10	11	18	11	5	-
2002.....	-	3	1	-	4	2	13	21	7	9	3	-
2001.....	1	1	1	1	1	2	5	15	13	10	1	-
2000.....	2	2	1	1	3	8	13	22	24	7	5	-
1999.....	-	1	-	1	3	9	11	23	34	14	6	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Chronic lower respiratory diseases (J40-J47)

##### All origins, both sexes

2003.....	2,261	4,344	7,733	11,805	18,114	24,593	24,693	18,193	8,874	2,516	351	2
2002.....	2,249	4,220	7,060	11,405	18,383	24,495	24,746	17,971	8,508	2,430	375	7
2001.....	2,197	4,176	6,990	11,838	18,913	24,442	23,320	17,542	8,125	2,345	379	1
2000.....	2,077	3,985	6,754	11,833	19,324	24,309	23,413	17,085	7,823	2,238	350	2
1999.....	2,031	4,179	7,118	12,441	20,203	24,978	23,657	16,831	7,510	2,154	338	4

##### All origins, male

2003.....	1,145	2,252	3,991	6,001	9,332	12,266	11,697	8,055	3,547	812	98	2
2002.....	1,178	2,141	3,642	5,838	9,501	12,457	11,975	8,140	3,448	762	86	5
2001.....	1,111	2,159	3,543	6,046	9,831	12,433	11,258	7,806	3,279	733	94	1
2000.....	1,044	2,072	3,492	6,149	10,061	12,305	11,533	7,843	3,165	776	92	2
1999.....	1,004	2,169	3,789	6,642	10,835	13,021	11,684	7,853	3,154	779	108	3

##### All origins, female

2003.....	1,116	2,092	3,742	5,804	8,782	12,327	12,996	10,138	5,327	1,704	253	-
2002.....	1,071	2,079	3,418	5,567	8,882	12,038	12,771	9,831	5,060	1,668	289	2
2001.....	1,086	2,017	3,447	5,792	9,082	12,009	12,062	9,736	4,846	1,612	285	-
2000.....	1,033	1,913	3,262	5,684	9,263	12,004	11,880	9,242	4,658	1,462	258	-
1999.....	1,027	2,010	3,329	5,799	9,368	11,957	11,973	8,978	4,356	1,375	230	1

##### Hispanic, both sexes

2003.....	73	114	152	252	399	515	580	491	277	103	17	-
2002.....	77	102	172	252	383	510	545	439	280	89	17	-
2001.....	77	100	158	265	369	513	427	399	271	80	18	-
2000.....	60	87	157	240	332	468	453	400	217	94	14	-
1999.....	63	95	144	262	387	483	452	415	249	91	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

Hispanic, male

2003.....	30	69	95	126	243	281	322	235	128	46	11	-
2002.....	42	64	100	152	210	312	281	201	122	34	3	-
2001.....	32	53	87	161	217	286	215	203	112	34	6	-
2000.....	32	40	106	136	207	260	249	190	95	45	3	-
1999.....	31	50	85	136	237	278	245	216	119	48	5	-

Hispanic, female

2003.....	43	45	57	126	156	234	258	256	149	57	6	-
2002.....	35	38	72	100	173	198	264	238	158	55	14	-
2001.....	45	47	71	104	152	227	212	196	159	46	12	-
2000.....	28	47	51	104	125	208	204	210	122	49	11	-
1999.....	32	45	59	126	150	205	207	199	130	43	4	-

Non-Hispanic 1/, both sexes

2003.....	2,171	4,214	7,547	11,516	17,676	24,036	24,057	17,664	8,581	2,406	334	1
2002.....	2,165	4,104	6,871	11,118	17,951	23,913	24,145	17,491	8,212	2,338	356	6
2001.....	2,107	4,055	6,805	11,545	18,483	23,869	22,844	17,108	7,830	2,260	360	-
2000.....	2,002	3,882	6,565	11,550	18,936	23,777	22,910	16,652	7,586	2,136	335	1
1999.....	1,955	4,075	6,943	12,145	19,757	24,425	23,153	16,368	7,239	2,059	329	3

Non-Hispanic 1/, male

2003.....	1,106	2,175	3,875	5,853	9,067	11,961	11,343	7,806	3,413	763	87	1
2002.....	1,134	2,072	3,532	5,665	9,259	12,098	11,663	7,918	3,321	728	83	4
2001.....	1,070	2,092	3,438	5,869	9,580	12,120	11,014	7,588	3,156	696	88	-
2000.....	1,004	2,022	3,366	5,985	9,820	12,009	11,257	7,637	3,061	729	88	1
1999.....	963	2,113	3,684	6,482	10,558	12,709	11,415	7,616	3,024	729	103	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

Non-Hispanic 1/, female

2003.....	1,065	2,039	3,672	5,663	8,609	12,075	12,714	9,858	5,168	1,643	247	-
2002.....	1,031	2,032	3,339	5,453	8,692	11,815	12,482	9,573	4,891	1,610	273	2
2001.....	1,037	1,963	3,367	5,676	8,903	11,749	11,830	9,520	4,674	1,564	272	-
2000.....	998	1,860	3,199	5,565	9,116	11,768	11,653	9,015	4,525	1,407	247	-
1999.....	992	1,962	3,259	5,663	9,199	11,716	11,738	8,752	4,215	1,330	226	1

Non-Hispanic White, both sexes

2003.....	1,773	3,650	6,815	10,564	16,338	22,460	22,717	16,739	8,091	2,236	295	1
2002.....	1,769	3,600	6,157	10,170	16,638	22,378	22,739	16,549	7,760	2,169	318	4
2001.....	1,739	3,548	6,130	10,528	17,165	22,403	21,569	16,195	7,328	2,108	319	-
2000.....	1,633	3,369	5,884	10,523	17,599	22,319	21,705	15,765	7,125	1,970	295	1
1999.....	1,625	3,555	6,231	11,099	18,301	22,770	21,892	15,486	6,810	1,928	297	2

Non-Hispanic White, male

2003.....	908	1,867	3,460	5,330	8,274	11,014	10,539	7,311	3,170	695	75	1
2002.....	924	1,793	3,116	5,124	8,491	11,168	10,866	7,394	3,109	659	70	3
2001.....	875	1,823	3,062	5,295	8,794	11,220	10,276	7,101	2,916	629	70	-
2000.....	814	1,744	2,984	5,367	9,031	11,134	10,569	7,141	2,836	657	74	1
1999.....	788	1,830	3,261	5,865	9,694	11,705	10,672	7,142	2,795	666	92	1

Non-Hispanic White, female

2003.....	865	1,783	3,355	5,234	8,064	11,446	12,178	9,428	4,921	1,541	220	-
2002.....	845	1,807	3,041	5,046	8,147	11,210	11,873	9,155	4,651	1,510	248	1
2001.....	864	1,725	3,068	5,233	8,371	11,183	11,293	9,094	4,412	1,479	249	-
2000.....	819	1,625	2,900	5,156	8,568	11,185	11,136	8,624	4,289	1,313	221	-
1999.....	837	1,725	2,970	5,234	8,607	11,065	11,220	8,344	4,015	1,262	205	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Chronic lower respiratory diseases (J40-J47)

##### Non-Hispanic Black, both sexes

2003.....	358	491	618	820	1,100	1,233	1,057	670	360	128	35	-
2002.....	359	454	613	802	1,090	1,240	1,126	710	347	138	31	1
2001.....	327	446	573	867	1,095	1,178	1,012	697	372	117	29	-
2000.....	335	443	620	872	1,121	1,176	952	663	358	131	31	-
1999.....	300	472	613	899	1,191	1,351	1,013	694	337	110	25	1

##### Non-Hispanic Black, male

2003.....	174	269	346	452	641	710	630	346	168	45	12	-
2002.....	189	250	357	461	632	735	615	378	155	54	10	1
2001.....	170	235	319	496	651	703	582	362	166	51	12	-
2000.....	164	238	349	526	669	681	526	356	171	58	8	-
1999.....	158	254	370	533	705	810	592	362	171	52	10	1

##### Non-Hispanic Black, female

2003.....	184	222	272	368	459	523	427	324	192	83	23	-
2002.....	170	204	256	341	458	505	511	332	192	84	21	-
2001.....	157	211	254	371	444	475	430	335	206	66	17	-
2000.....	171	205	271	346	452	495	426	307	187	73	23	-
1999.....	142	218	243	366	486	541	421	332	166	58	15	-

##### Origin not stated 2/, both sexes

2003.....	17	16	34	37	39	42	56	38	16	7	-	1
2002.....	7	14	17	35	49	72	56	41	16	3	2	1
2001.....	13	21	27	28	61	60	49	35	24	5	1	1
2000.....	15	16	32	43	56	64	50	33	20	8	1	1
1999.....	13	9	31	34	59	70	52	48	22	4	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic lower respiratory diseases (J40-J47)

Origin not stated 2/, male

2003.....	9	8	21	22	22	24	32	14	6	3	-	1
2002.....	2	5	10	21	32	47	31	21	5	-	-	1
2001.....	9	14	18	16	34	27	29	15	11	3	-	1
2000.....	8	10	20	28	34	36	27	16	9	2	1	1
1999.....	10	6	20	24	40	34	24	21	11	2	-	1

Origin not stated 2/, female

2003.....	8	8	13	15	17	18	24	24	10	4	-	-
2002.....	5	9	7	14	17	25	25	20	11	3	2	-
2001.....	4	7	9	12	27	33	20	20	13	2	1	-
2000.....	7	6	12	15	22	28	23	17	11	6	-	-
1999.....	3	3	11	10	19	36	28	27	11	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

All origins, both sexes

2003.....	18	21	26	40	64	107	139	151	129	51	14	-
2002.....	20	22	32	51	84	131	155	162	114	69	27	-
2001.....	21	25	36	56	77	129	132	188	123	67	23	-
2000.....	19	20	47	67	91	156	177	236	161	80	24	-
1999.....	24	35	62	63	110	139	200	193	148	82	23	-

All origins, male

2003.....	10	12	14	17	29	59	54	51	35	13	2	-
2002.....	14	10	18	32	42	69	58	62	35	20	5	-
2001.....	9	16	20	35	41	67	48	70	33	14	2	-
2000.....	8	9	23	41	38	84	90	92	46	19	1	-
1999.....	10	24	27	36	58	73	84	71	45	20	6	-

All origins, female

2003.....	8	9	12	23	35	48	85	100	94	38	12	-
2002.....	6	12	14	19	42	62	97	100	79	49	22	-
2001.....	12	9	16	21	36	62	84	118	90	53	21	-
2000.....	11	11	24	26	53	72	87	144	115	61	23	-
1999.....	14	11	35	27	52	66	116	122	103	62	17	-

Hispanic, both sexes

2003.....	-	-	1	-	3	3	6	7	3	4	-	-
2002.....	2	-	2	2	-	7	7	9	8	3	-	-
2001.....	-	2	-	1	2	8	1	4	5	1	3	-
2000.....	-	1	-	2	7	4	9	6	4	5	1	-
1999.....	-	-	2	2	2	4	9	5	10	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Hispanic, male

2003.....	-	-	-	-	2	2	3	2	3	1	-	-
2002.....	2	-	1	1	-	1	3	3	1	1	-	-
2001.....	-	1	-	-	1	6	-	1	2	-	-	-
2000.....	-	1	-	1	6	1	5	3	1	2	-	-
1999.....	-	-	-	2	1	1	2	3	1	-	1	-

Hispanic, female

2003.....	-	-	1	-	1	1	3	5	-	3	-	-
2002.....	-	-	1	1	-	6	4	6	7	2	-	-
2001.....	-	1	-	1	1	2	1	3	3	1	3	-
2000.....	-	-	-	1	1	3	4	3	3	3	1	-
1999.....	-	-	2	-	1	3	7	2	9	1	-	-

Non-Hispanic 1/, both sexes

2003.....	18	21	25	40	61	104	131	144	126	47	14	-
2002.....	18	22	30	48	84	123	148	153	106	66	26	-
2001.....	21	22	36	55	75	121	130	184	118	66	20	-
2000.....	19	19	47	65	84	152	168	230	157	72	23	-
1999.....	24	34	60	60	108	135	191	187	138	81	22	-

Non-Hispanic 1/, male

2003.....	10	12	14	17	27	57	50	49	32	12	2	-
2002.....	12	10	17	31	42	67	55	59	34	19	5	-
2001.....	9	14	20	35	40	61	48	69	31	14	2	-
2000.....	8	8	23	40	32	83	85	89	45	16	1	-
1999.....	10	23	27	33	57	72	82	67	44	20	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Non-Hispanic 1/, female

2003.....	8	9	11	23	34	47	81	95	94	35	12	-
2002.....	6	12	13	17	42	56	93	94	72	47	21	-
2001.....	12	8	16	20	35	60	82	115	87	52	18	-
2000.....	11	11	24	25	52	69	83	141	112	56	22	-
1999.....	14	11	33	27	51	63	109	120	94	61	17	-

Non-Hispanic White, both sexes

2003.....	13	14	23	38	54	97	117	132	118	42	13	-
2002.....	15	18	26	43	81	117	135	147	100	63	25	-
2001.....	20	21	33	48	65	110	122	178	111	62	19	-
2000.....	15	16	44	56	75	140	157	223	153	69	22	-
1999.....	15	31	55	55	95	119	174	175	133	78	19	-

Non-Hispanic White, male

2003.....	8	10	13	17	22	52	44	44	31	12	2	-
2002.....	9	8	14	27	40	63	50	58	31	19	4	-
2001.....	8	14	18	31	35	54	43	66	29	11	2	-
2000.....	7	7	20	36	28	77	82	85	45	15	1	-
1999.....	6	21	24	31	50	64	71	63	42	19	4	-

Non-Hispanic White, female

2003.....	5	4	10	21	32	45	73	88	87	30	11	-
2002.....	6	10	12	16	41	54	85	89	69	44	21	-
2001.....	12	7	15	17	30	56	79	112	82	51	17	-
2000.....	8	9	24	20	47	63	75	138	108	54	21	-
1999.....	9	10	31	24	45	55	103	112	91	59	15	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Bronchitis, chronic and unspecified (J40-J42)

Non-Hispanic Black, both sexes

2003.....	5	6	2	2	4	3	9	9	5	4	1	-
2002.....	3	3	2	4	2	4	9	6	5	2	1	-
2001.....	1	1	3	7	8	8	3	5	3	3	-	-
2000.....	4	1	3	7	9	10	10	5	3	3	1	-
1999.....	8	3	4	4	11	12	14	8	3	1	2	-

Non-Hispanic Black, male

2003.....	2	2	1	-	2	3	4	3	-	-	-	-
2002.....	3	1	1	3	1	3	2	1	3	-	1	-
2001.....	1	-	2	4	4	4	2	2	-	3	-	-
2000.....	1	-	3	2	4	4	3	2	-	1	-	-
1999.....	4	2	2	1	6	6	10	1	1	-	1	-

Non-Hispanic Black, female

2003.....	3	4	1	2	2	-	5	6	5	4	1	-
2002.....	-	2	1	1	1	1	7	5	2	2	-	-
2001.....	-	1	1	3	4	4	1	3	3	-	-	-
2000.....	3	1	-	5	5	6	7	3	3	2	1	-
1999.....	4	1	2	3	5	6	4	7	2	1	1	-

Origin not stated 2/, both sexes

2003.....	-	-	-	-	-	-	2	-	-	-	-	-
2002.....	-	-	-	1	-	1	-	-	-	-	1	-
2001.....	-	1	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	3	-	-
1999.....	-	1	-	1	-	-	-	1	-	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Emphysema (J43)

##### All origins, both sexes

2003.....	293	630	1,125	1,618	2,445	3,124	2,796	1,740	672	143	19	2
2002.....	298	643	1,060	1,665	2,723	3,202	2,907	1,826	726	157	15	1
2001.....	307	705	1,124	1,892	2,912	3,481	2,856	1,827	727	161	21	-
2000.....	312	680	1,127	1,976	3,144	3,522	2,989	1,832	685	150	18	-
1999.....	332	723	1,296	2,246	3,410	3,768	3,187	1,785	622	146	27	-

##### All origins, male

2003.....	174	339	593	827	1,314	1,608	1,387	792	300	48	5	2
2002.....	172	334	591	875	1,460	1,723	1,505	866	329	52	4	1
2001.....	177	395	571	988	1,566	1,809	1,446	929	326	54	6	-
2000.....	179	366	577	1,077	1,661	1,839	1,534	898	317	58	5	-
1999.....	183	394	691	1,230	1,906	2,027	1,650	886	295	68	9	-

##### All origins, female

2003.....	119	291	532	791	1,131	1,516	1,409	948	372	95	14	-
2002.....	126	309	469	790	1,263	1,479	1,402	960	397	105	11	-
2001.....	130	310	553	904	1,346	1,672	1,410	898	401	107	15	-
2000.....	133	314	550	899	1,483	1,683	1,455	934	368	92	13	-
1999.....	149	329	605	1,016	1,504	1,741	1,537	899	327	78	18	-

##### Hispanic, both sexes

2003.....	4	12	18	24	49	54	74	41	19	4	3	-
2002.....	4	9	20	25	54	41	72	34	25	6	1	-
2001.....	3	10	16	33	42	61	52	32	22	6	3	-
2000.....	8	8	18	26	39	60	53	29	13	5	-	-
1999.....	7	6	15	28	46	56	57	37	18	7	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Emphysema (J43)

## Hispanic, male

2003.....	3	6	15	10	31	36	40	21	12	1	1	-
2002.....	3	5	12	20	28	27	38	18	11	3	-	-
2001.....	3	7	9	19	27	41	35	21	7	2	2	-
2000.....	6	2	15	17	26	35	31	18	8	3	-	-
1999.....	4	3	13	18	33	34	35	21	9	6	-	-

## Hispanic, female

2003.....	1	6	3	14	18	18	34	20	7	3	2	-
2002.....	1	4	8	5	26	14	34	16	14	3	1	-
2001.....	-	3	7	14	15	20	17	11	15	4	1	-
2000.....	2	6	3	9	13	25	22	11	5	2	-	-
1999.....	3	3	2	10	13	22	22	16	9	1	1	-

## Non-Hispanic 1/, both sexes

2003.....	286	617	1,099	1,590	2,392	3,063	2,718	1,698	652	139	16	1
2002.....	294	633	1,039	1,637	2,661	3,155	2,831	1,784	701	151	14	-
2001.....	304	691	1,103	1,855	2,863	3,412	2,798	1,794	701	154	18	-
2000.....	302	670	1,107	1,945	3,091	3,450	2,932	1,799	669	145	18	-
1999.....	320	716	1,276	2,213	3,358	3,701	3,124	1,740	601	139	26	-

## Non-Hispanic 1/, male

2003.....	169	332	576	814	1,280	1,567	1,343	771	287	47	4	1
2002.....	169	328	579	853	1,428	1,690	1,466	843	318	49	4	-
2001.....	174	384	559	966	1,534	1,764	1,408	907	318	51	4	-
2000.....	172	363	562	1,058	1,627	1,800	1,503	879	308	55	5	-
1999.....	175	390	673	1,208	1,868	1,986	1,612	862	284	62	9	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

Non-Hispanic 1/, female

2003.....	117	285	523	776	1,112	1,496	1,375	927	365	92	12	-
2002.....	125	305	460	784	1,233	1,465	1,365	941	383	102	10	-
2001.....	130	307	544	889	1,329	1,648	1,390	887	383	103	14	-
2000.....	130	307	545	887	1,464	1,650	1,429	920	361	90	13	-
1999.....	145	326	603	1,005	1,490	1,715	1,512	878	317	77	17	-

Non-Hispanic White, both sexes

2003.....	252	556	1,001	1,481	2,236	2,875	2,563	1,615	603	125	14	1
2002.....	258	576	953	1,509	2,483	2,987	2,676	1,691	664	147	11	-
2001.....	253	612	1,021	1,736	2,678	3,238	2,675	1,711	655	143	17	-
2000.....	252	599	1,017	1,788	2,916	3,273	2,795	1,699	628	129	16	-
1999.....	274	648	1,177	2,073	3,165	3,482	2,987	1,647	569	128	21	-

Non-Hispanic White, male

2003.....	144	293	515	753	1,182	1,440	1,250	720	252	41	3	1
2002.....	143	293	517	759	1,322	1,574	1,372	781	302	47	2	-
2001.....	143	335	506	900	1,410	1,656	1,330	852	297	48	4	-
2000.....	139	322	508	947	1,523	1,698	1,423	808	284	44	3	-
1999.....	140	344	609	1,115	1,728	1,855	1,516	812	262	57	7	-

Non-Hispanic White, female

2003.....	108	263	486	728	1,054	1,435	1,313	895	351	84	11	-
2002.....	115	283	436	750	1,161	1,413	1,304	910	362	100	9	-
2001.....	110	277	515	836	1,268	1,582	1,345	859	358	95	13	-
2000.....	113	277	509	841	1,393	1,575	1,372	891	344	85	13	-
1999.....	134	304	568	958	1,437	1,627	1,471	835	307	71	14	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

Non-Hispanic Black, both sexes

2003.....	28	57	85	97	123	146	120	57	36	11	1	-
2002.....	30	49	68	111	156	135	131	67	32	4	3	-
2001.....	46	69	69	106	148	123	96	65	40	9	1	-
2000.....	47	63	83	135	141	135	108	73	31	14	-	-
1999.....	43	60	86	126	153	175	108	76	27	11	3	-

Non-Hispanic Black, male

2003.....	19	37	53	54	75	94	68	36	26	4	1	-
2002.....	22	30	47	81	92	90	79	42	14	2	2	-
2001.....	27	43	44	61	94	73	61	40	18	2	-	-
2000.....	30	36	51	97	86	72	62	47	20	9	-	-
1999.....	33	38	56	87	108	103	75	38	17	5	2	-

Non-Hispanic Black, female

2003.....	9	20	32	43	48	52	52	21	10	7	-	-
2002.....	8	19	21	30	64	45	52	25	18	2	1	-
2001.....	19	26	25	45	54	50	35	25	22	7	1	-
2000.....	17	27	32	38	55	63	46	26	11	5	-	-
1999.....	10	22	30	39	45	72	33	38	10	6	1	-

Origin not stated 2/, both sexes

2003.....	3	1	8	4	4	7	4	1	1	-	-	1
2002.....	-	1	1	3	8	6	4	8	-	-	-	1
2001.....	-	4	5	4	7	8	6	1	4	1	-	-
2000.....	2	2	2	5	14	12	4	4	3	-	-	-
1999.....	5	1	5	5	6	11	6	8	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Emphysema (J43)

Origin not stated 2/, male

2003.....	2	1	2	3	3	5	4	-	1	-	-	1
2002.....	-	1	-	2	4	6	1	5	-	-	-	1
2001.....	-	4	3	3	5	4	3	1	1	1	-	-
2000.....	1	1	-	2	8	4	-	1	1	-	-	-
1999.....	4	1	5	4	5	7	3	3	2	-	-	-

Origin not stated 2/, female

2003.....	1	-	6	1	1	2	-	1	-	-	-	-
2002.....	-	-	1	1	4	-	3	3	-	-	-	-
2001.....	-	-	2	1	2	4	3	-	3	-	-	-
2000.....	1	1	2	3	6	8	4	3	2	-	-	-
1999.....	1	-	-	1	1	4	3	5	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Asthma (J45-J46)

## All origins, both sexes

2003.....	315	299	263	262	270	364	388	335	236	75	25	-
2002.....	305	288	248	257	326	401	411	345	225	86	19	1
2001.....	324	265	288	288	346	384	418	371	239	77	22	-
2000.....	296	270	270	328	406	420	429	388	227	81	11	-
1999.....	295	343	240	353	420	454	471	381	224	97	18	-

## All origins, male

2003.....	109	117	97	80	93	116	118	78	57	14	5	-
2002.....	120	108	94	89	109	131	125	97	41	10	4	-
2001.....	116	84	98	104	116	116	109	86	57	11	3	-
2000.....	110	106	94	121	132	120	123	104	39	15	-	-
1999.....	101	119	88	115	152	144	129	81	52	17	2	-

## All origins, female

2003.....	206	182	166	182	177	248	270	257	179	61	20	-
2002.....	185	180	154	168	217	270	286	248	184	76	15	1
2001.....	208	181	190	184	230	268	309	285	182	66	19	-
2000.....	186	164	176	207	274	300	306	284	188	66	11	-
1999.....	194	224	152	238	268	310	342	300	172	80	16	-

## Hispanic, both sexes

2003.....	26	29	16	28	13	17	22	15	12	1	-	-
2002.....	19	14	24	25	20	22	20	11	7	3	1	-
2001.....	29	26	19	19	24	26	18	8	10	4	-	-
2000.....	22	15	33	34	26	25	15	16	6	3	-	-
1999.....	23	23	16	32	28	27	23	19	3	3	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Asthma (J45-J46)

## Hispanic, male

2003.....	11	16	6	7	6	4	11	4	2	-	-	-
2002.....	8	5	10	4	6	10	5	2	1	-	-	-
2001.....	8	12	5	9	9	8	5	-	2	-	-	-
2000.....	12	5	14	14	11	8	4	6	1	1	-	-
1999.....	11	9	3	10	9	7	5	4	1	1	-	-

## Hispanic, female

2003.....	15	13	10	21	7	13	11	11	10	1	-	-
2002.....	11	9	14	21	14	12	15	9	6	3	1	-
2001.....	21	14	14	10	15	18	13	8	8	4	-	-
2000.....	10	10	19	20	15	17	11	10	5	2	-	-
1999.....	12	14	13	22	19	20	18	15	2	2	-	-

## Non-Hispanic 1/, both sexes

2003.....	286	269	247	234	257	347	366	317	223	73	25	-
2002.....	285	272	223	230	306	377	390	332	217	83	18	1
2001.....	292	239	268	267	320	356	400	362	227	73	22	-
2000.....	270	252	234	291	377	392	413	370	220	78	11	-
1999.....	271	319	224	320	391	427	447	359	221	94	18	-

## Non-Hispanic 1/, male

2003.....	97	101	91	73	87	112	107	74	55	13	5	-
2002.....	111	103	84	83	103	119	120	93	40	10	4	-
2001.....	106	72	93	95	105	107	104	85	53	11	3	-
2000.....	96	98	78	106	120	110	119	98	37	14	-	-
1999.....	90	110	85	104	142	137	123	76	51	16	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Asthma (J45-J46)

## Non-Hispanic 1/, female

2003.....	189	168	156	161	170	235	259	243	168	60	20	-
2002.....	174	169	139	147	203	258	270	239	177	73	14	1
2001.....	186	167	175	172	215	249	296	277	174	62	19	-
2000.....	174	154	156	185	257	282	294	272	183	64	11	-
1999.....	181	209	139	216	249	290	324	283	170	78	16	-

## Non-Hispanic White, both sexes

2003.....	152	167	167	166	182	275	306	285	199	69	22	-
2002.....	155	159	152	155	222	318	341	304	197	72	13	-
2001.....	182	146	160	195	237	285	343	316	201	67	18	-
2000.....	159	145	151	194	290	320	363	320	189	66	8	-
1999.....	160	205	146	223	288	355	392	305	191	85	17	-

## Non-Hispanic White, male

2003.....	49	60	54	53	61	79	90	65	50	12	5	-
2002.....	63	58	55	52	73	95	101	84	37	7	2	-
2001.....	61	45	50	67	74	83	84	75	43	9	1	-
2000.....	60	50	47	64	95	92	101	81	29	8	-	-
1999.....	52	72	52	67	97	108	110	56	39	13	2	-

## Non-Hispanic White, female

2003.....	103	107	113	113	121	196	216	220	149	57	17	-
2002.....	92	101	97	103	149	223	240	220	160	65	11	-
2001.....	121	101	110	128	163	202	259	241	158	58	17	-
2000.....	99	95	104	130	195	228	262	239	160	58	8	-
1999.....	108	133	94	156	191	247	282	249	152	72	15	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Asthma (J45-J46)

## Non-Hispanic Black, both sexes

2003.....	122	81	59	63	57	48	41	18	16	4	2	-
2002.....	119	105	64	57	75	41	34	20	10	7	2	-
2001.....	97	84	86	57	69	58	41	34	13	3	3	-
2000.....	97	88	77	80	72	60	34	32	26	9	3	-
1999.....	101	96	66	76	81	55	39	37	22	5	-	-

## Non-Hispanic Black, male

2003.....	44	27	25	20	18	20	11	3	3	1	-	-
2002.....	44	40	27	25	26	13	10	6	1	2	-	-
2001.....	41	23	29	18	28	17	12	8	3	1	1	-
2000.....	27	37	28	33	20	10	11	9	7	5	-	-
1999.....	34	30	27	26	35	20	7	8	7	1	-	-

## Non-Hispanic Black, female

2003.....	78	54	34	43	39	28	30	15	13	3	2	-
2002.....	75	65	37	32	49	28	24	14	9	5	2	-
2001.....	56	61	57	39	41	41	29	26	10	2	2	-
2000.....	70	51	49	47	52	50	23	23	19	4	3	-
1999.....	67	66	39	50	46	35	32	29	15	4	-	-

## Origin not stated 2/, both sexes

2003.....	3	1	-	-	-	-	-	3	1	1	-	-
2002.....	1	2	1	2	-	2	1	2	1	-	-	-
2001.....	3	-	1	2	2	2	-	1	2	-	-	-
2000.....	4	3	3	3	3	3	1	2	1	-	-	-
1999.....	1	1	-	1	1	-	1	3	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Asthma (J45-J46)

## Origin not stated 2/, male

2003.....	1	-	-	-	-	-	-	-	-	1	-	-
2002.....	1	-	-	2	-	2	-	2	-	-	-	-
2001.....	2	-	-	-	2	1	-	1	2	-	-	-
2000.....	2	3	2	1	1	2	-	-	1	-	-	-
1999.....	-	-	-	1	1	-	1	1	-	-	-	-

## Origin not stated 2/, female

2003.....	2	1	-	-	-	-	-	3	1	-	-	-
2002.....	-	2	1	-	-	-	1	-	1	-	-	-
2001.....	1	-	1	2	-	1	-	-	-	-	-	-
2000.....	2	-	1	2	2	1	1	2	-	-	-	-
1999.....	1	1	-	-	-	-	-	2	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

All origins, both sexes

2003.....	1,635	3,394	6,319	9,885	15,335	20,998	21,370	15,967	7,837	2,247	293	-
2002.....	1,626	3,267	5,720	9,432	15,250	20,761	21,273	15,638	7,443	2,118	314	5
2001.....	1,545	3,181	5,542	9,602	15,578	20,448	19,914	15,156	7,036	2,040	313	1
2000.....	1,450	3,015	5,310	9,462	15,683	20,211	19,818	14,629	6,750	1,927	297	2
1999.....	1,380	3,078	5,520	9,779	16,263	20,617	19,799	14,472	6,516	1,829	270	4

All origins, male

2003.....	852	1,784	3,287	5,077	7,896	10,483	10,138	7,134	3,155	737	86	-
2002.....	872	1,689	2,939	4,842	7,890	10,534	10,287	7,115	3,043	680	73	4
2001.....	809	1,664	2,854	4,919	8,108	10,441	9,655	6,721	2,863	654	83	1
2000.....	747	1,591	2,798	4,910	8,230	10,262	9,786	6,749	2,763	684	86	2
1999.....	710	1,632	2,983	5,261	8,719	10,777	9,821	6,815	2,762	674	91	3

All origins, female

2003.....	783	1,610	3,032	4,808	7,439	10,515	11,232	8,833	4,682	1,510	207	-
2002.....	754	1,578	2,781	4,590	7,360	10,227	10,986	8,523	4,400	1,438	241	1
2001.....	736	1,517	2,688	4,683	7,470	10,007	10,259	8,435	4,173	1,386	230	-
2000.....	703	1,424	2,512	4,552	7,453	9,949	10,032	7,880	3,987	1,243	211	-
1999.....	670	1,446	2,537	4,518	7,544	9,840	9,978	7,657	3,754	1,155	179	1

Hispanic, both sexes

2003.....	43	73	117	200	334	441	478	428	243	94	14	-
2002.....	52	79	126	200	309	440	446	385	240	77	15	-
2001.....	45	62	123	212	301	418	356	355	234	69	12	-
2000.....	30	63	106	178	260	379	376	349	194	81	13	-
1999.....	33	66	111	200	311	396	363	354	218	80	7	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Hispanic, male

2003.....	16	47	74	109	204	239	268	208	111	44	10	-
2002.....	29	54	77	127	176	274	235	178	109	30	3	-
2001.....	21	33	73	133	180	231	175	181	101	32	4	-
2000.....	14	32	77	104	164	216	209	163	85	39	3	-
1999.....	16	38	69	106	194	236	203	188	108	41	4	-

Hispanic, female

2003.....	27	26	43	91	130	202	210	220	132	50	4	-
2002.....	23	25	49	73	133	166	211	207	131	47	12	-
2001.....	24	29	50	79	121	187	181	174	133	37	8	-
2000.....	16	31	29	74	96	163	167	186	109	42	10	-
1999.....	17	28	42	94	117	160	160	166	110	39	3	-

Non-Hispanic 1/, both sexes

2003.....	1,581	3,307	6,176	9,652	14,966	20,522	20,842	15,505	7,580	2,147	279	-
2002.....	1,568	3,177	5,579	9,203	14,900	20,258	20,776	15,222	7,188	2,038	298	5
2001.....	1,490	3,103	5,398	9,368	15,225	19,980	19,516	14,768	6,784	1,967	300	-
2000.....	1,411	2,941	5,177	9,249	15,384	19,783	19,397	14,253	6,540	1,841	283	1
1999.....	1,340	3,006	5,383	9,552	15,900	20,162	19,391	14,082	6,279	1,745	263	3

Non-Hispanic 1/, male

2003.....	830	1,730	3,194	4,949	7,673	10,225	9,843	6,912	3,039	691	76	-
2002.....	842	1,631	2,852	4,698	7,686	10,222	10,022	6,923	2,929	650	70	4
2001.....	781	1,622	2,766	4,773	7,901	10,188	9,454	6,527	2,754	620	79	-
2000.....	728	1,553	2,703	4,781	8,041	10,016	9,550	6,571	2,671	644	82	1
1999.....	688	1,590	2,899	5,137	8,491	10,514	9,598	6,611	2,645	631	87	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic 1/, female

2003.....	751	1,577	2,982	4,703	7,293	10,297	10,999	8,593	4,541	1,456	203	-
2002.....	726	1,546	2,727	4,505	7,214	10,036	10,754	8,299	4,259	1,388	228	1
2001.....	709	1,481	2,632	4,595	7,324	9,792	10,062	8,241	4,030	1,347	221	-
2000.....	683	1,388	2,474	4,468	7,343	9,767	9,847	7,682	3,869	1,197	201	-
1999.....	652	1,416	2,484	4,415	7,409	9,648	9,793	7,471	3,634	1,114	176	1

Non-Hispanic White, both sexes

2003.....	1,356	2,913	5,624	8,879	13,866	19,213	19,731	14,707	7,171	2,000	246	-
2002.....	1,341	2,847	5,026	8,463	13,852	18,956	19,587	14,407	6,799	1,887	269	4
2001.....	1,284	2,769	4,916	8,549	14,185	18,770	18,429	13,990	6,361	1,836	265	-
2000.....	1,207	2,609	4,672	8,485	14,318	18,586	18,390	13,523	6,155	1,706	249	1
1999.....	1,176	2,671	4,853	8,748	14,753	18,814	18,339	13,359	5,917	1,637	240	2

Non-Hispanic White, male

2003.....	707	1,504	2,878	4,507	7,009	9,443	9,155	6,482	2,837	630	65	-
2002.....	709	1,434	2,530	4,286	7,056	9,436	9,343	6,471	2,739	586	62	3
2001.....	663	1,429	2,488	4,297	7,275	9,427	8,819	6,108	2,547	561	63	-
2000.....	608	1,365	2,409	4,320	7,385	9,267	8,963	6,167	2,478	590	70	1
1999.....	590	1,393	2,576	4,652	7,819	9,678	8,975	6,211	2,452	577	79	1

Non-Hispanic White, female

2003.....	649	1,409	2,746	4,372	6,857	9,770	10,576	8,225	4,334	1,370	181	-
2002.....	632	1,413	2,496	4,177	6,796	9,520	10,244	7,936	4,060	1,301	207	1
2001.....	621	1,340	2,428	4,252	6,910	9,343	9,610	7,882	3,814	1,275	202	-
2000.....	599	1,244	2,263	4,165	6,933	9,319	9,427	7,356	3,677	1,116	179	-
1999.....	586	1,278	2,277	4,096	6,934	9,136	9,364	7,148	3,465	1,060	161	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Non-Hispanic Black, both sexes

2003.....	203	347	472	658	916	1,036	887	586	303	109	31	-
2002.....	207	297	479	630	857	1,060	952	617	300	125	25	1
2001.....	183	292	415	697	870	989	872	593	316	102	25	-
2000.....	187	291	457	650	899	971	800	553	298	105	27	-
1999.....	148	313	457	693	946	1,109	852	573	285	93	20	1

Non-Hispanic Black, male

2003.....	109	203	267	378	546	593	547	304	139	40	11	-
2002.....	120	179	282	352	513	629	524	329	137	50	7	1
2001.....	101	169	244	413	525	609	507	312	145	45	11	-
2000.....	106	165	267	394	559	595	450	298	144	43	8	-
1999.....	87	184	285	419	556	681	500	315	146	46	7	1

Non-Hispanic Black, female

2003.....	94	144	205	280	370	443	340	282	164	69	20	-
2002.....	87	118	197	278	344	431	428	288	163	75	18	-
2001.....	82	123	171	284	345	380	365	281	171	57	14	-
2000.....	81	126	190	256	340	376	350	255	154	62	19	-
1999.....	61	129	172	274	390	428	352	258	139	47	13	-

Origin not stated 2/, both sexes

2003.....	11	14	26	33	35	35	50	34	14	6	-	-
2002.....	6	11	15	29	41	63	51	31	15	3	1	-
2001.....	10	16	21	22	52	50	42	33	18	4	1	1
2000.....	9	11	27	35	39	49	45	27	16	5	1	1
1999.....	7	6	26	27	52	59	45	36	19	4	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic lower respiratory diseases (J44,J47)

Origin not stated 2/, male

2003.....	6	7	19	19	19	19	27	14	5	2	-	-
2002.....	1	4	10	17	28	38	30	14	5	-	-	-
2001.....	7	9	15	13	27	22	26	13	8	2	-	1
2000.....	5	6	18	25	25	30	27	15	7	1	1	1
1999.....	6	4	15	18	34	27	20	16	9	2	-	1

Origin not stated 2/, female

2003.....	5	7	7	14	16	16	23	20	9	4	-	-
2002.....	5	7	5	12	13	25	21	17	10	3	1	-
2001.....	3	7	6	9	25	28	16	20	10	2	1	-
2000.....	4	5	9	10	14	19	18	12	9	4	-	-
1999.....	1	2	11	9	18	32	25	20	10	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

All origins, both sexes

2003.....	10	27	43	91	142	220	267	203	84	12	2	-
2002.....	12	26	49	80	143	256	224	217	76	17	2	-
2001.....	8	27	37	91	175	252	264	188	66	16	2	-
2000.....	7	21	36	85	196	249	235	217	80	20	-	-
1999.....	12	19	38	104	170	251	252	180	54	18	1	-

All origins, male

2003.....	9	26	41	90	132	213	256	192	80	12	2	-
2002.....	9	24	45	77	137	245	212	207	75	17	1	-
2001.....	7	26	36	90	171	239	255	181	63	16	2	-
2000.....	7	19	35	79	186	241	226	208	79	19	-	-
1999.....	12	19	38	99	170	243	243	172	53	17	1	-

All origins, female

2003.....	1	1	2	1	10	7	11	11	4	-	-	-
2002.....	3	2	4	3	6	11	12	10	1	-	1	-
2001.....	1	1	1	1	4	13	9	7	3	-	-	-
2000.....	-	2	1	6	10	8	9	9	1	1	-	-
1999.....	-	-	-	5	-	8	9	8	1	1	-	-

Hispanic, both sexes

2003.....	-	2	1	2	1	6	6	-	1	-	-	-
2002.....	1	2	2	6	6	1	4	4	5	-	-	-
2001.....	1	1	1	1	6	4	2	2	2	1	1	-
2000.....	1	1	-	4	1	3	10	4	-	1	-	-
1999.....	1	-	1	2	3	2	-	2	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

Hispanic, male

2003.....	-	2	1	2	1	6	6	-	-	-	-	-
2002.....	-	2	2	6	5	1	4	4	5	-	-	-
2001.....	1	1	1	1	6	4	2	2	1	1	1	-
2000.....	1	1	-	4	1	3	10	3	-	1	-	-
1999.....	1	-	1	2	3	2	-	-	1	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	1	-	-	-
2002.....	1	-	-	-	1	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	1	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	-	-	-	-	-	-	-	2	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	10	25	42	89	141	214	260	203	83	12	2	-
2002.....	11	23	47	74	137	255	218	213	71	17	2	-
2001.....	7	26	36	90	169	248	261	186	63	15	1	-
2000.....	6	20	36	81	195	245	225	213	80	19	-	-
1999.....	11	19	37	102	167	249	252	178	53	18	1	-

Non-Hispanic 1/, male

2003.....	9	24	40	88	131	207	249	192	80	12	2	-
2002.....	9	21	43	71	132	244	206	203	70	17	1	-
2001.....	6	25	35	89	165	235	252	179	61	15	1	-
2000.....	6	18	35	75	185	237	216	205	79	18	-	-
1999.....	11	19	37	97	167	241	243	172	52	17	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumoconioses and chemical effects (J60-J66,J68)

Non-Hispanic 1/, female

2003.....	1	1	2	1	10	7	11	11	3	-	-	-
2002.....	2	2	4	3	5	11	12	10	1	-	1	-
2001.....	1	1	1	1	4	13	9	7	2	-	-	-
2000.....	-	2	1	6	10	8	9	8	1	1	-	-
1999.....	-	-	-	5	-	8	9	6	1	1	-	-

Non-Hispanic White, both sexes

2003.....	9	23	41	83	135	202	245	198	79	12	2	-
2002.....	10	21	41	71	130	238	207	208	66	14	1	-
2001.....	5	23	30	86	159	240	247	175	61	15	1	-
2000.....	6	20	34	76	185	230	212	206	76	17	-	-
1999.....	7	19	27	93	157	237	245	166	52	15	-	-

Non-Hispanic White, male

2003.....	8	22	39	82	125	195	235	188	76	12	2	-
2002.....	8	21	38	68	125	228	196	199	65	14	-	-
2001.....	5	22	29	85	155	227	238	168	59	15	1	-
2000.....	6	18	33	70	175	223	204	198	75	16	-	-
1999.....	7	19	27	89	157	229	236	161	51	14	-	-

Non-Hispanic White, female

2003.....	1	1	2	1	10	7	10	10	3	-	-	-
2002.....	2	-	3	3	5	10	11	9	1	-	1	-
2001.....	-	1	1	1	4	13	9	7	2	-	-	-
2000.....	-	2	1	6	10	7	8	8	1	1	-	-
1999.....	-	-	-	4	-	8	9	5	1	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

All origins, both sexes

2003.....	204	317	456	610	1,144	2,245	3,544	3,908	2,945	1,204	250	-
2002.....	208	271	400	627	1,166	2,323	3,577	4,100	2,995	1,219	229	-
2001.....	236	259	400	589	1,241	2,259	3,500	3,958	2,960	1,160	264	-
2000.....	209	257	341	642	1,257	2,258	3,247	3,825	2,779	1,111	239	-
1999.....	155	259	327	582	1,172	2,068	2,962	3,443	2,586	1,053	212	-

All origins, male

2003.....	129	196	275	388	681	1,285	1,896	1,861	1,223	407	67	-
2002.....	136	162	231	380	703	1,332	1,911	1,994	1,267	415	55	-
2001.....	151	160	224	362	744	1,253	1,892	1,893	1,180	379	59	-
2000.....	126	153	201	405	790	1,291	1,747	1,839	1,116	367	65	-
1999.....	82	151	214	367	716	1,252	1,623	1,659	1,006	372	46	-

All origins, female

2003.....	75	121	181	222	463	960	1,648	2,047	1,722	797	183	-
2002.....	72	109	169	247	463	991	1,666	2,106	1,728	804	174	-
2001.....	85	99	176	227	497	1,006	1,608	2,065	1,780	781	205	-
2000.....	83	104	140	237	467	967	1,500	1,986	1,663	744	174	-
1999.....	73	108	113	215	456	816	1,339	1,784	1,580	681	166	-

Hispanic, both sexes

2003.....	7	20	15	29	51	63	83	89	73	46	10	-
2002.....	7	12	23	24	38	73	101	110	86	35	6	-
2001.....	11	16	17	27	42	67	85	112	97	41	8	-
2000.....	9	15	18	21	43	68	73	87	78	38	4	-
1999.....	4	8	18	21	36	57	61	85	67	28	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Hispanic, male

2003.....	6	10	10	19	28	34	50	35	27	16	2	-
2002.....	4	9	15	14	18	42	63	46	41	11	2	-
2001.....	6	14	8	17	25	37	44	56	41	16	3	-
2000.....	7	10	10	10	25	44	43	41	33	10	-	-
1999.....	1	4	10	12	19	34	31	39	22	13	-	-

Hispanic, female

2003.....	1	10	5	10	23	29	33	54	46	30	8	-
2002.....	3	3	8	10	20	31	38	64	45	24	4	-
2001.....	5	2	9	10	17	30	41	56	56	25	5	-
2000.....	2	5	8	11	18	24	30	46	45	28	4	-
1999.....	3	4	8	9	17	23	30	46	45	15	6	-

Non-Hispanic 1/, both sexes

2003.....	197	296	440	579	1,089	2,176	3,454	3,813	2,861	1,156	240	-
2002.....	201	257	377	602	1,121	2,249	3,468	3,980	2,904	1,180	222	-
2001.....	225	243	381	562	1,194	2,188	3,405	3,837	2,854	1,116	256	-
2000.....	198	239	323	621	1,212	2,183	3,163	3,726	2,696	1,070	235	-
1999.....	150	250	309	560	1,131	2,004	2,891	3,349	2,513	1,024	206	-

Non-Hispanic 1/, male

2003.....	123	185	264	368	650	1,247	1,842	1,824	1,190	391	65	-
2002.....	132	151	216	366	679	1,289	1,841	1,939	1,224	403	53	-
2001.....	145	146	215	345	716	1,213	1,844	1,832	1,137	361	56	-
2000.....	117	142	191	395	763	1,240	1,697	1,793	1,081	354	65	-
1999.....	80	146	204	354	692	1,214	1,589	1,615	981	358	46	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Non-Hispanic 1/, female

2003.....	74	111	176	211	439	929	1,612	1,989	1,671	765	175	-
2002.....	69	106	161	236	442	960	1,627	2,041	1,680	777	169	-
2001.....	80	97	166	217	478	975	1,561	2,005	1,717	755	200	-
2000.....	81	97	132	226	449	943	1,466	1,933	1,615	716	170	-
1999.....	70	104	105	206	439	790	1,302	1,734	1,532	666	160	-

Non-Hispanic White, both sexes

2003.....	167	247	332	467	930	1,900	3,078	3,496	2,616	1,055	207	-
2002.....	151	215	304	480	957	1,969	3,162	3,664	2,686	1,067	199	-
2001.....	173	205	313	463	1,006	1,933	3,079	3,503	2,601	1,011	214	-
2000.....	155	196	253	509	1,006	1,934	2,859	3,437	2,482	985	207	-
1999.....	116	195	251	458	942	1,758	2,603	3,086	2,317	919	186	-

Non-Hispanic White, male

2003.....	106	154	200	304	553	1,092	1,649	1,683	1,097	354	60	-
2002.....	98	123	180	295	579	1,131	1,691	1,800	1,124	367	52	-
2001.....	112	125	173	290	612	1,079	1,686	1,688	1,048	333	50	-
2000.....	90	120	150	319	626	1,091	1,557	1,676	995	328	60	-
1999.....	60	113	163	286	574	1,063	1,429	1,492	908	323	44	-

Non-Hispanic White, female

2003.....	61	93	132	163	377	808	1,429	1,813	1,519	701	147	-
2002.....	53	92	124	185	378	838	1,471	1,864	1,562	700	147	-
2001.....	61	80	140	173	394	854	1,393	1,815	1,553	678	164	-
2000.....	65	76	103	190	380	843	1,302	1,761	1,487	657	147	-
1999.....	56	82	88	172	368	695	1,174	1,594	1,409	596	142	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Non-Hispanic Black, both sexes

2003.....	27	42	99	103	128	234	299	252	204	77	24	-
2002.....	44	35	63	110	140	243	256	259	184	93	17	-
2001.....	48	34	66	86	165	221	276	271	203	92	34	-
2000.....	41	39	63	102	187	216	252	232	178	74	21	-
1999.....	31	51	54	90	174	225	244	233	176	95	17	-

Non-Hispanic Black, male

2003.....	15	27	59	58	79	130	146	102	70	26	4	-
2002.....	30	24	32	67	85	135	118	105	81	30	1	-
2001.....	31	20	40	49	91	120	132	115	69	22	5	-
2000.....	27	20	35	67	123	129	114	88	66	21	4	-
1999.....	18	30	38	59	109	135	126	103	58	30	2	-

Non-Hispanic Black, female

2003.....	12	15	40	45	49	104	153	150	134	51	20	-
2002.....	14	11	31	43	55	108	138	154	103	63	16	-
2001.....	17	14	26	37	74	101	144	156	134	70	29	-
2000.....	14	19	28	35	64	87	138	144	112	53	17	-
1999.....	13	21	16	31	65	90	118	130	118	65	15	-

Origin not stated 2/, both sexes

2003.....	-	1	1	2	4	6	7	6	11	2	-	-
2002.....	-	2	-	1	7	1	8	10	5	4	1	-
2001.....	-	-	2	-	5	4	10	9	9	3	-	-
2000.....	2	3	-	-	2	7	11	12	5	3	-	-
1999.....	1	1	-	1	5	7	10	9	6	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Pneumonitis due to solids and liquids (J69)

Origin not stated 2/, male

2003.....	-	1	1	1	3	4	4	2	6	-	-	-
2002.....	-	2	-	-	6	1	7	9	2	1	-	-
2001.....	-	-	1	-	3	3	4	5	2	2	-	-
2000.....	2	1	-	-	2	7	7	5	2	3	-	-
1999.....	1	1	-	1	5	4	3	5	3	1	-	-

Origin not stated 2/, female

2003.....	-	-	-	1	1	2	3	4	5	2	-	-
2002.....	-	-	-	1	1	-	1	1	3	3	1	-
2001.....	-	-	1	-	2	1	6	4	7	1	-	-
2000.....	-	2	-	-	-	-	4	7	3	-	-	-
1999.....	-	-	-	-	-	3	7	4	3	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins, both sexes

2003.....	762	1,037	1,405	2,050	3,106	4,272	4,538	3,721	1,926	723	140	1
2002.....	766	1,055	1,385	1,971	3,266	4,212	4,446	3,445	1,908	639	138	-
2001.....	814	1,030	1,402	2,174	3,336	4,590	4,612	3,627	1,895	643	118	3
2000.....	805	1,044	1,367	2,184	3,264	4,487	4,413	3,394	1,850	629	135	3
1999.....	694	904	1,356	2,143	3,266	4,369	4,277	3,498	1,823	667	142	-

All origins, male

2003.....	403	573	794	1,174	1,675	2,226	2,169	1,531	633	187	33	-
2002.....	393	573	766	1,084	1,845	2,166	2,095	1,395	625	136	20	-
2001.....	442	530	792	1,197	1,828	2,330	2,123	1,415	618	143	21	1
2000.....	431	572	752	1,179	1,782	2,353	2,062	1,409	612	156	28	2
1999.....	368	467	746	1,199	1,806	2,255	2,073	1,417	551	178	29	-

All origins, female

2003.....	359	464	611	876	1,431	2,046	2,369	2,190	1,293	536	107	1
2002.....	373	482	619	887	1,421	2,046	2,351	2,050	1,283	503	118	-
2001.....	372	500	610	977	1,508	2,260	2,489	2,212	1,277	500	97	2
2000.....	374	472	615	1,005	1,482	2,134	2,351	1,985	1,238	473	107	1
1999.....	326	437	610	944	1,460	2,114	2,204	2,081	1,272	489	113	-

Hispanic, both sexes

2003.....	50	91	94	126	185	194	179	124	63	38	4	-
2002.....	54	71	93	123	185	185	163	118	57	22	5	-
2001.....	64	83	82	125	162	186	177	128	61	22	4	-
2000.....	42	83	84	123	150	174	151	108	67	21	8	-
1999.....	55	53	104	131	138	174	133	110	57	22	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Hispanic, male

2003.....	34	52	60	67	100	93	72	52	19	11	1	-
2002.....	25	43	49	72	109	78	78	45	15	3	2	-
2001.....	34	43	49	66	85	95	80	57	22	6	3	-
2000.....	24	44	52	61	78	92	73	38	20	5	3	-
1999.....	23	31	58	69	70	91	67	48	21	5	1	-

Hispanic, female

2003.....	16	39	34	59	85	101	107	72	44	27	3	-
2002.....	29	28	44	51	76	107	85	73	42	19	3	-
2001.....	30	40	33	59	77	91	97	71	39	16	1	-
2000.....	18	39	32	62	72	82	78	70	47	16	5	-
1999.....	32	22	46	62	68	83	66	62	36	17	1	-

Non-Hispanic 1/, both sexes

2003.....	708	944	1,307	1,919	2,914	4,070	4,355	3,588	1,859	685	136	1
2002.....	705	981	1,287	1,840	3,076	4,022	4,274	3,322	1,848	615	132	-
2001.....	748	946	1,319	2,044	3,159	4,390	4,425	3,495	1,830	619	113	3
2000.....	760	956	1,282	2,054	3,104	4,304	4,248	3,278	1,779	606	127	2
1999.....	636	843	1,249	2,008	3,119	4,183	4,132	3,380	1,762	644	140	-

Non-Hispanic 1/, male

2003.....	366	520	730	1,103	1,571	2,128	2,094	1,477	613	176	32	-
2002.....	364	527	716	1,009	1,734	2,086	2,012	1,347	610	133	18	-
2001.....	407	486	742	1,128	1,732	2,228	2,037	1,357	596	137	18	1
2000.....	404	528	699	1,115	1,700	2,254	1,985	1,368	591	151	25	1
1999.....	343	431	686	1,128	1,729	2,157	1,996	1,366	529	173	28	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic 1/, female

2003.....	342	424	577	816	1,343	1,942	2,261	2,111	1,246	509	104	1
2002.....	341	454	571	831	1,342	1,936	2,262	1,975	1,238	482	114	-
2001.....	341	460	577	916	1,427	2,162	2,388	2,138	1,234	482	95	2
2000.....	356	428	583	939	1,404	2,050	2,263	1,910	1,188	455	102	1
1999.....	293	412	563	880	1,390	2,026	2,136	2,014	1,233	471	112	-

Non-Hispanic White, both sexes

2003.....	537	754	1,104	1,671	2,614	3,681	4,078	3,373	1,732	629	119	1
2002.....	541	790	1,060	1,598	2,731	3,688	3,931	3,099	1,726	572	120	-
2001.....	539	765	1,095	1,768	2,829	4,020	4,113	3,258	1,688	549	100	1
2000.....	592	782	1,065	1,765	2,792	3,943	3,923	3,078	1,641	553	112	-
1999.....	491	662	1,056	1,743	2,768	3,817	3,825	3,133	1,641	574	123	-

Non-Hispanic White, male

2003.....	286	417	621	968	1,423	1,929	1,968	1,388	563	159	32	-
2002.....	284	434	589	902	1,546	1,920	1,862	1,269	563	122	16	-
2001.....	307	407	611	993	1,581	2,060	1,903	1,270	549	118	16	-
2000.....	315	435	586	975	1,533	2,068	1,845	1,287	536	133	18	-
1999.....	272	330	589	999	1,562	1,988	1,860	1,267	486	151	22	-

Non-Hispanic White, female

2003.....	251	337	483	703	1,191	1,752	2,110	1,985	1,169	470	87	1
2002.....	257	356	471	696	1,185	1,768	2,069	1,830	1,163	450	104	-
2001.....	232	358	484	775	1,248	1,960	2,210	1,988	1,139	431	84	1
2000.....	277	347	479	790	1,259	1,875	2,078	1,791	1,105	420	94	-
1999.....	219	332	467	744	1,206	1,829	1,965	1,866	1,155	423	101	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Non-Hispanic Black, both sexes

2003.....	156	162	167	191	219	297	196	153	86	50	17	-
2002.....	140	161	192	195	251	236	258	157	98	34	11	-
2001.....	185	154	189	220	262	284	243	171	114	57	13	2
2000.....	144	155	170	249	246	277	250	148	111	42	13	2
1999.....	125	154	163	223	280	295	228	192	89	54	14	-

Non-Hispanic Black, male

2003.....	71	89	94	102	115	151	88	66	29	14	-	-
2002.....	70	80	107	89	133	116	113	50	33	9	2	-
2001.....	86	67	106	107	115	126	104	65	34	13	2	1
2000.....	73	81	85	121	121	143	97	54	41	15	5	1
1999.....	62	81	83	108	129	139	96	74	30	14	5	-

Non-Hispanic Black, female

2003.....	85	73	73	89	104	146	108	87	57	36	17	-
2002.....	70	81	85	106	118	120	145	107	65	25	9	-
2001.....	99	87	83	113	147	158	139	106	80	44	11	1
2000.....	71	74	85	128	125	134	153	94	70	27	8	1
1999.....	63	73	80	115	151	156	132	118	59	40	9	-

Origin not stated 2/, both sexes

2003.....	4	2	4	5	7	8	4	9	4	-	-	-
2002.....	7	3	5	8	5	5	9	5	3	2	1	-
2001.....	2	1	1	5	15	14	10	4	4	2	1	-
2000.....	3	5	1	7	10	9	14	8	4	2	-	1
1999.....	3	8	3	4	9	12	12	8	4	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Origin not stated 2/, male

2003.....	3	1	4	4	4	5	3	2	1	-	-	-
2002.....	4	3	1	3	2	2	5	3	-	-	-	-
2001.....	1	1	1	3	11	7	6	1	-	-	-	-
2000.....	3	-	1	3	4	7	4	3	1	-	-	1
1999.....	2	5	2	2	7	7	10	3	1	-	-	-

Origin not stated 2/, female

2003.....	1	1	-	1	3	3	1	7	3	-	-	-
2002.....	3	-	4	5	3	3	4	2	3	2	1	-
2001.....	1	-	-	2	4	7	4	3	4	2	1	-
2000.....	-	5	-	4	6	2	10	5	3	2	-	-
1999.....	1	3	1	2	2	5	2	5	3	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

All origins, both sexes

2003.....	159	183	200	228	359	545	734	685	391	138	30	-
2002.....	154	209	202	254	411	592	699	714	407	148	33	-
2001.....	147	183	235	294	424	615	823	788	458	184	40	1
2000.....	177	200	243	299	449	642	773	801	498	177	33	-
1999.....	142	186	236	314	522	665	835	771	470	164	34	2

All origins, male

2003.....	110	105	132	141	205	273	333	262	126	25	5	-
2002.....	106	129	118	142	236	287	335	279	116	39	5	-
2001.....	95	123	150	179	218	315	334	277	125	47	5	1
2000.....	131	134	150	172	270	341	326	291	150	49	9	-
1999.....	102	111	154	196	308	331	381	279	138	29	2	1

All origins, female

2003.....	49	78	68	87	154	272	401	423	265	113	25	-
2002.....	48	80	84	112	175	305	364	435	291	109	28	-
2001.....	52	60	85	115	206	300	489	511	333	137	35	-
2000.....	46	66	93	127	179	301	447	510	348	128	24	-
1999.....	40	75	82	118	214	334	454	492	332	135	32	1

Hispanic, both sexes

2003.....	17	15	11	17	25	28	28	18	11	4	-	-
2002.....	10	15	17	21	18	27	26	21	10	6	-	-
2001.....	10	12	10	10	23	24	30	22	13	2	1	1
2000.....	10	11	17	13	23	19	31	22	17	5	2	-
1999.....	8	14	11	16	21	20	24	22	12	3	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

Hispanic, male

2003.....	11	8	8	13	17	15	15	5	1	1	-	-
2002.....	8	11	12	16	12	14	13	8	5	3	-	-
2001.....	7	10	9	5	13	17	20	6	4	1	-	1
2000.....	7	8	15	8	13	11	6	12	2	2	-	-
1999.....	7	9	9	10	12	14	13	9	5	2	-	-

Hispanic, female

2003.....	6	7	3	4	8	13	13	13	10	3	-	-
2002.....	2	4	5	5	6	13	13	13	5	3	-	-
2001.....	3	2	1	5	10	7	10	16	9	1	1	-
2000.....	3	3	2	5	10	8	25	10	15	3	2	-
1999.....	1	5	2	6	9	6	11	13	7	1	1	-

Non-Hispanic 1/, both sexes

2003.....	138	167	187	211	333	517	698	665	380	133	30	-
2002.....	144	191	183	233	392	564	671	691	396	142	33	-
2001.....	136	170	223	280	401	589	791	763	445	180	39	-
2000.....	167	188	226	285	424	622	739	778	478	171	31	-
1999.....	133	170	223	297	499	642	809	746	458	161	33	2

Non-Hispanic 1/, male

2003.....	95	96	123	128	187	258	316	255	125	24	5	-
2002.....	98	117	105	126	223	272	322	270	110	36	5	-
2001.....	88	113	139	170	205	297	312	271	121	46	5	-
2000.....	124	126	135	163	256	329	319	279	148	47	9	-
1999.....	94	101	143	186	295	315	367	270	133	27	2	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

Non-Hispanic 1/, female

2003.....	43	71	64	83	146	259	382	410	255	109	25	-
2002.....	46	74	78	107	169	292	349	421	286	106	28	-
2001.....	48	57	84	110	196	292	479	492	324	134	34	-
2000.....	43	62	91	122	168	293	420	499	330	124	22	-
1999.....	39	69	80	111	204	327	442	476	325	134	31	1

Non-Hispanic White, both sexes

2003.....	97	134	153	180	292	448	619	611	354	122	24	-
2002.....	108	146	142	193	344	497	626	639	370	132	28	-
2001.....	101	126	182	232	351	522	726	706	418	168	38	-
2000.....	122	151	192	236	373	550	683	709	448	158	27	-
1999.....	98	136	186	242	435	578	735	690	427	147	27	1

Non-Hispanic White, male

2003.....	63	75	100	107	162	223	273	234	114	20	3	-
2002.....	69	87	82	104	192	239	300	251	103	33	5	-
2001.....	61	82	112	140	171	259	284	251	111	42	5	-
2000.....	89	102	115	137	220	289	291	251	136	40	8	-
1999.....	69	78	117	150	259	287	335	247	128	22	2	1

Non-Hispanic White, female

2003.....	34	59	53	73	130	225	346	377	240	102	21	-
2002.....	39	59	60	89	152	258	326	388	267	99	23	-
2001.....	40	44	70	92	180	263	442	455	307	126	33	-
2000.....	33	49	77	99	153	261	392	458	312	118	19	-
1999.....	29	58	69	92	176	291	400	443	299	125	25	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

Non-Hispanic Black, both sexes

2003.....	37	31	28	20	33	50	57	34	16	4	4	-
2002.....	33	35	33	30	39	50	25	37	13	8	4	-
2001.....	32	34	34	32	41	45	38	38	20	8	1	-
2000.....	38	34	29	37	42	52	39	49	23	8	4	-
1999.....	32	32	33	45	47	42	56	40	24	11	4	1

Non-Hispanic Black, male

2003.....	29	21	20	15	21	25	32	13	5	1	-	-
2002.....	27	24	21	17	24	25	13	14	2	2	-	-
2001.....	25	25	22	20	27	23	16	15	5	1	-	-
2000.....	28	21	19	18	30	26	19	18	5	3	1	-
1999.....	23	22	24	29	26	18	21	18	3	3	-	-

Non-Hispanic Black, female

2003.....	8	10	8	5	12	25	25	21	11	3	4	-
2002.....	6	11	12	13	15	25	12	23	11	6	4	-
2001.....	7	9	12	12	14	22	22	23	15	7	1	-
2000.....	10	13	10	19	12	26	20	31	18	5	3	-
1999.....	9	10	9	16	21	24	35	22	21	8	4	1

Origin not stated 2/, both sexes

2003.....	4	1	2	-	1	-	8	2	-	1	-	-
2002.....	-	3	2	-	1	1	2	2	1	-	-	-
2001.....	1	1	2	4	-	2	2	3	-	2	-	-
2000.....	-	1	-	1	2	1	3	1	3	1	-	-
1999.....	1	2	2	1	2	3	2	3	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Peptic ulcer (K25-K28)

Origin not stated 2/, male

2003.....	4	1	1	-	1	-	2	2	-	-	-	-
2002.....	-	1	1	-	1	1	-	1	1	-	-	-
2001.....	-	-	2	4	-	1	2	-	-	-	-	-
2000.....	-	-	-	1	1	1	1	-	-	-	-	-
1999.....	1	1	2	-	1	2	1	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	1	-	-	-	6	-	-	1	-	-
2002.....	-	2	1	-	-	-	2	1	-	-	-	-
2001.....	1	1	-	-	-	1	-	3	-	2	-	-
2000.....	-	1	-	-	1	-	2	1	3	1	-	-
1999.....	-	1	-	1	1	1	1	3	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

All origins, both sexes

2003.....	25	18	25	29	35	56	76	42	28	7	1	-
2002.....	30	28	36	32	57	61	59	56	25	10	3	-
2001.....	15	22	23	34	55	37	53	51	30	11	1	-
2000.....	25	28	32	26	45	53	68	39	24	4	1	-
1999.....	24	22	26	30	39	43	53	40	18	3	1	-

All origins, male

2003.....	16	13	17	20	26	28	47	21	10	3	-	-
2002.....	20	18	27	24	32	35	32	22	6	1	2	-
2001.....	11	13	17	23	34	22	30	20	12	4	1	-
2000.....	17	20	23	15	25	30	30	15	9	-	-	-
1999.....	19	17	17	18	20	28	19	23	9	2	1	-

All origins, female

2003.....	9	5	8	9	9	28	29	21	18	4	1	-
2002.....	10	10	9	8	25	26	27	34	19	9	1	-
2001.....	4	9	6	11	21	15	23	31	18	7	-	-
2000.....	8	8	9	11	20	23	38	24	15	4	1	-
1999.....	5	5	9	12	19	15	34	17	9	1	-	-

Hispanic, both sexes

2003.....	4	1	-	2	2	1	3	2	1	1	-	-
2002.....	1	3	3	1	2	3	1	1	1	-	-	-
2001.....	1	-	2	2	3	3	4	1	-	-	-	-
2000.....	-	4	1	-	2	1	2	-	-	-	-	-
1999.....	2	1	2	2	1	2	1	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

Hispanic, male

2003.....	2	1	-	1	2	1	2	1	1	1	-	-
2002.....	1	2	3	1	1	2	-	-	-	-	-	-
2001.....	1	-	1	2	3	1	2	-	-	-	-	-
2000.....	-	4	-	-	-	1	1	-	-	-	-	-
1999.....	2	1	1	2	-	1	-	1	-	-	-	-

Hispanic, female

2003.....	2	-	-	1	-	-	1	1	-	-	-	-
2002.....	-	1	-	-	1	1	1	1	1	-	-	-
2001.....	-	-	1	-	-	2	2	1	-	-	-	-
2000.....	-	-	1	-	2	-	1	-	-	-	-	-
1999.....	-	-	1	-	1	1	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	21	17	25	27	33	55	73	40	27	6	1	-
2002.....	29	25	32	31	55	58	58	55	24	10	3	-
2001.....	14	22	21	32	52	34	49	50	30	11	1	-
2000.....	25	24	31	26	43	52	66	39	24	4	1	-
1999.....	22	21	24	28	37	41	52	39	18	3	1	-

Non-Hispanic 1/, male

2003.....	14	12	17	19	24	27	45	20	9	2	-	-
2002.....	19	16	23	23	31	33	32	22	6	1	2	-
2001.....	10	13	16	21	31	21	28	20	12	4	1	-
2000.....	17	16	23	15	25	29	29	15	9	-	-	-
1999.....	17	16	16	16	19	27	19	22	9	2	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

Non-Hispanic 1/, female

2003.....	7	5	8	8	9	28	28	20	18	4	1	-
2002.....	10	9	9	8	24	25	26	33	18	9	1	-
2001.....	4	9	5	11	21	13	21	30	18	7	-	-
2000.....	8	8	8	11	18	23	37	24	15	4	1	-
1999.....	5	5	8	12	18	14	33	17	9	1	-	-

Non-Hispanic White, both sexes

2003.....	15	14	17	21	28	45	68	37	24	6	1	-
2002.....	22	22	28	25	44	53	55	51	21	10	2	-
2001.....	13	13	14	26	49	30	40	43	28	10	1	-
2000.....	19	17	28	20	32	41	56	38	22	4	1	-
1999.....	20	14	13	23	31	36	45	36	17	3	1	-

Non-Hispanic White, male

2003.....	11	10	12	15	20	22	42	19	9	2	-	-
2002.....	15	15	20	18	25	30	30	19	4	1	1	-
2001.....	9	10	11	17	30	19	24	14	11	3	1	-
2000.....	12	10	20	13	18	25	24	14	8	-	-	-
1999.....	15	11	11	13	15	23	15	20	9	2	1	-

Non-Hispanic White, female

2003.....	4	4	5	6	8	23	26	18	15	4	1	-
2002.....	7	7	8	7	19	23	25	32	17	9	1	-
2001.....	4	3	3	9	19	11	16	29	17	7	-	-
2000.....	7	7	8	7	14	16	32	24	14	4	1	-
1999.....	5	3	2	10	16	13	30	16	8	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of appendix (K35-K38)

Non-Hispanic Black, both sexes

2003.....	3	3	7	4	5	9	2	3	2	-	-	-
2002.....	6	3	3	5	8	3	2	4	3	-	1	-
2001.....	1	9	7	6	1	3	8	3	2	1	-	-
2000.....	4	7	3	6	10	9	7	1	2	-	-	-
1999.....	2	7	10	3	6	5	6	2	1	-	-	-

Non-Hispanic Black, male

2003.....	1	2	4	3	4	4	1	1	-	-	-	-
2002.....	4	1	2	4	4	2	2	3	2	-	1	-
2001.....	1	3	5	4	-	2	3	3	1	1	-	-
2000.....	3	6	3	2	7	3	3	1	1	-	-	-
1999.....	2	5	4	2	4	4	4	2	-	-	-	-

Non-Hispanic Black, female

2003.....	2	1	3	1	1	5	1	2	2	-	-	-
2002.....	2	2	1	1	4	1	-	1	1	-	-	-
2001.....	-	6	2	2	1	1	5	-	1	-	-	-
2000.....	1	1	-	4	3	6	4	-	1	-	-	-
1999.....	-	2	6	1	2	1	2	-	1	-	-	-

Origin not stated 2/, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hernia (K40-K46)

##### All origins, both sexes

2003.....	60	76	85	101	136	203	259	296	202	56	9	-
2002.....	44	55	71	90	135	208	281	320	172	85	11	-
2001.....	58	57	74	81	132	185	258	269	190	88	6	-
2000.....	58	51	62	84	126	224	242	294	188	63	10	-
1999.....	39	36	71	77	138	186	265	269	196	56	8	-

##### All origins, male

2003.....	35	30	42	48	63	93	106	95	62	17	4	-
2002.....	19	27	29	40	63	84	105	97	45	20	-	-
2001.....	31	23	41	40	54	71	102	88	48	19	1	-
2000.....	32	20	29	39	53	91	104	96	42	12	4	-
1999.....	20	21	39	34	59	75	117	96	51	8	2	-

##### All origins, female

2003.....	25	46	43	53	73	110	153	201	140	39	5	-
2002.....	25	28	42	50	72	124	176	223	127	65	11	-
2001.....	27	34	33	41	78	114	156	181	142	69	5	-
2000.....	26	31	33	45	73	133	138	198	146	51	6	-
1999.....	19	15	32	43	79	111	148	173	145	48	6	-

##### Hispanic, both sexes

2003.....	8	11	4	10	5	11	12	9	4	-	1	-
2002.....	4	5	9	6	8	6	13	7	6	2	-	-
2001.....	5	3	6	3	9	12	12	13	6	-	1	-
2000.....	5	4	5	7	7	15	8	10	8	1	-	-
1999.....	3	-	5	7	9	6	11	8	8	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hernia (K40-K46)

##### Hispanic, male

2003.....	5	4	1	4	3	2	5	4	2	-	-	-
2002.....	2	2	5	2	2	2	5	1	3	-	-	-
2001.....	2	-	3	2	2	1	5	2	3	-	-	-
2000.....	3	3	-	-	3	6	4	2	1	-	-	-
1999.....	2	-	4	1	5	2	4	1	2	-	-	-

##### Hispanic, female

2003.....	3	7	3	6	2	9	7	5	2	-	1	-
2002.....	2	3	4	4	6	4	8	6	3	2	-	-
2001.....	3	3	3	1	7	11	7	11	3	-	1	-
2000.....	2	1	5	7	4	9	4	8	7	1	-	-
1999.....	1	-	1	6	4	4	7	7	6	1	-	-

##### Non-Hispanic 1/, both sexes

2003.....	52	65	81	90	131	189	246	287	197	56	8	-
2002.....	40	50	61	82	127	199	266	312	166	83	11	-
2001.....	53	54	67	78	123	172	245	256	183	88	5	-
2000.....	53	47	56	77	117	209	233	284	179	62	10	-
1999.....	36	36	66	70	128	177	254	260	187	54	8	-

##### Non-Hispanic 1/, male

2003.....	30	26	41	43	60	89	101	91	60	17	4	-
2002.....	17	25	24	37	61	80	98	96	42	20	-	-
2001.....	29	23	37	38	52	69	97	86	45	19	1	-
2000.....	29	17	28	39	49	85	99	94	41	12	4	-
1999.....	18	21	35	33	53	72	113	94	49	8	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hernia (K40-K46)

##### Non-Hispanic 1/, female

2003.....	22	39	40	47	71	100	145	196	137	39	4	-
2002.....	23	25	37	45	66	119	168	216	124	63	11	-
2001.....	24	31	30	40	71	103	148	170	138	69	4	-
2000.....	24	30	28	38	68	124	134	190	138	50	6	-
1999.....	18	15	31	37	75	105	141	166	138	46	6	-

##### Non-Hispanic White, both sexes

2003.....	40	49	67	77	115	172	217	267	179	55	6	-
2002.....	30	42	46	63	113	188	243	288	161	81	10	-
2001.....	42	42	55	66	107	151	223	238	172	77	3	-
2000.....	42	41	49	61	108	184	215	260	167	56	9	-
1999.....	27	32	57	53	114	156	234	239	174	52	6	-

##### Non-Hispanic White, male

2003.....	25	23	34	37	51	83	89	83	55	17	2	-
2002.....	15	22	19	28	53	76	91	88	39	19	-	-
2001.....	21	16	30	34	47	62	87	73	42	18	1	-
2000.....	22	15	25	31	47	79	91	86	38	10	4	-
1999.....	15	21	32	28	51	66	105	84	42	7	2	-

##### Non-Hispanic White, female

2003.....	15	26	33	40	64	89	128	184	124	38	4	-
2002.....	15	20	27	35	60	112	152	200	122	62	10	-
2001.....	21	26	25	32	60	89	136	165	130	59	2	-
2000.....	20	26	24	30	61	105	124	174	129	46	5	-
1999.....	12	11	25	25	63	90	129	155	132	45	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hernia (K40-K46)

##### Non-Hispanic Black, both sexes

2003.....	11	16	12	12	12	16	22	14	15	1	1	-
2002.....	8	8	13	15	12	10	20	19	4	2	1	-
2001.....	11	12	11	10	15	17	16	16	9	9	2	-
2000.....	8	5	7	14	7	21	15	24	12	3	1	-
1999.....	7	4	8	12	12	19	18	15	12	2	2	-

##### Non-Hispanic Black, male

2003.....	5	3	7	6	7	5	9	6	4	-	1	-
2002.....	2	3	4	8	7	4	7	6	2	1	-	-
2001.....	8	7	6	2	5	6	8	11	3	1	-	-
2000.....	5	1	3	7	2	4	7	8	3	-	-	-
1999.....	3	-	2	5	2	5	8	7	6	1	-	-

##### Non-Hispanic Black, female

2003.....	6	13	5	6	5	11	13	8	11	1	-	-
2002.....	6	5	9	7	5	6	13	13	2	1	1	-
2001.....	3	5	5	8	10	11	8	5	6	8	2	-
2000.....	3	4	4	7	5	17	8	16	9	3	1	-
1999.....	4	4	6	7	10	14	10	8	6	1	2	-

##### Origin not stated 2/, both sexes

2003.....	-	-	-	1	-	3	1	-	1	-	-	-
2002.....	-	-	1	2	-	3	2	1	-	-	-	-
2001.....	-	-	1	-	-	1	1	-	1	-	-	-
2000.....	-	-	1	-	2	-	1	-	1	-	-	-
1999.....	-	-	-	-	1	3	-	1	1	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Hernia (K40-K46)

##### Origin not stated 2/, male

2003.....	-	-	-	1	-	2	-	-	-	-	-	-
2002.....	-	-	-	1	-	2	2	-	-	-	-	-
2001.....	-	-	1	-	-	1	-	-	-	-	-	-
2000.....	-	-	1	-	1	-	1	-	-	-	-	-
1999.....	-	-	-	-	1	1	-	1	-	-	-	-

##### Origin not stated 2/, female

2003.....	-	-	-	-	-	1	1	-	1	-	-	-
2002.....	-	-	1	1	-	1	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	1	-	-	-
2000.....	-	-	-	-	1	-	-	-	1	-	-	-
1999.....	-	-	-	-	-	2	-	-	1	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Chronic liver disease and cirrhosis (K70,K73-K74)

##### All origins, both sexes

2003.....	4,009	3,476	2,952	2,729	2,677	2,314	1,544	712	204	29	1	3
2002.....	3,812	3,258	2,839	2,670	2,711	2,412	1,588	728	230	24	3	4
2001.....	3,702	2,999	2,751	2,752	2,734	2,325	1,476	714	221	43	1	6
2000.....	3,280	2,947	2,827	2,751	2,731	2,367	1,462	742	203	30	4	3
1999.....	3,140	2,803	2,834	2,833	2,809	2,396	1,505	702	226	33	4	2

##### All origins, male

2003.....	3,004	2,485	2,052	1,736	1,529	1,181	720	304	80	13	-	3
2002.....	2,843	2,323	1,905	1,652	1,545	1,211	742	310	77	8	-	4
2001.....	2,714	2,161	1,867	1,749	1,586	1,151	682	307	89	19	1	6
2000.....	2,460	2,066	1,939	1,729	1,588	1,240	634	325	83	12	1	3
1999.....	2,383	2,051	1,892	1,837	1,617	1,208	697	293	86	14	2	2

##### All origins, female

2003.....	1,005	991	900	993	1,148	1,133	824	408	124	16	1	-
2002.....	969	935	934	1,018	1,166	1,201	846	418	153	16	3	-
2001.....	988	838	884	1,003	1,148	1,174	794	407	132	24	-	-
2000.....	820	881	888	1,022	1,143	1,127	828	417	120	18	3	-
1999.....	757	752	942	996	1,192	1,188	808	409	140	19	2	-

##### Hispanic, both sexes

2003.....	463	433	369	322	302	265	132	43	13	4	-	-
2002.....	505	422	336	332	331	198	123	58	21	3	-	1
2001.....	471	378	343	322	296	219	116	46	12	1	-	1
2000.....	414	380	328	333	295	215	108	51	19	2	1	-
1999.....	364	325	313	310	273	187	98	51	20	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Hispanic, male

2003.....	379	331	285	184	148	131	55	14	8	1	-	-
2002.....	408	333	228	226	181	95	62	18	8	1	-	1
2001.....	379	295	260	220	166	120	50	16	6	-	-	1
2000.....	342	295	236	233	162	121	42	17	12	1	-	-
1999.....	293	257	224	198	162	93	43	20	10	-	1	-

Hispanic, female

2003.....	84	102	84	138	154	134	77	29	5	3	-	-
2002.....	97	89	108	106	150	103	61	40	13	2	-	-
2001.....	92	83	83	102	130	99	66	30	6	1	-	-
2000.....	72	85	92	100	133	94	66	34	7	1	1	-
1999.....	71	68	89	112	111	94	55	31	10	1	-	-

Non-Hispanic 1/, both sexes

2003.....	3,529	3,030	2,566	2,399	2,372	2,045	1,409	669	191	25	1	2
2002.....	3,291	2,819	2,488	2,329	2,371	2,207	1,461	669	209	21	3	2
2001.....	3,216	2,607	2,395	2,417	2,429	2,099	1,356	667	209	40	1	3
2000.....	2,846	2,557	2,484	2,405	2,428	2,148	1,350	691	181	28	3	2
1999.....	2,756	2,467	2,511	2,514	2,530	2,201	1,405	649	204	32	3	2

Non-Hispanic 1/, male

2003.....	2,611	2,142	1,752	1,547	1,378	1,049	664	290	72	12	-	2
2002.....	2,423	1,977	1,664	1,420	1,358	1,110	678	292	69	7	-	2
2001.....	2,323	1,854	1,596	1,518	1,416	1,030	631	290	83	18	1	3
2000.....	2,100	1,762	1,690	1,486	1,422	1,117	591	308	69	11	1	2
1999.....	2,075	1,784	1,660	1,631	1,450	1,111	654	272	76	14	1	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic 1/, female

2003.....	918	888	814	852	994	996	745	379	119	13	1	-
2002.....	868	842	824	909	1,013	1,097	783	377	140	14	3	-
2001.....	893	753	799	899	1,013	1,069	725	377	126	22	-	-
2000.....	746	795	794	919	1,006	1,031	759	383	112	17	2	-
1999.....	681	683	851	883	1,080	1,090	751	377	128	18	2	-

Non-Hispanic White, both sexes

2003.....	2,866	2,524	2,182	2,099	2,157	1,918	1,321	627	178	22	1	2
2002.....	2,670	2,364	2,119	2,047	2,158	2,061	1,356	628	194	20	3	2
2001.....	2,557	2,127	2,062	2,112	2,195	1,945	1,277	628	196	37	1	2
2000.....	2,272	2,121	2,114	2,093	2,192	2,008	1,256	646	166	27	3	1
1999.....	2,260	2,013	2,104	2,191	2,291	2,027	1,320	602	187	29	3	2

Non-Hispanic White, male

2003.....	2,138	1,788	1,497	1,364	1,263	986	617	272	66	10	-	2
2002.....	1,988	1,674	1,423	1,246	1,243	1,050	631	275	63	7	-	2
2001.....	1,865	1,511	1,364	1,318	1,288	956	601	276	76	17	1	2
2000.....	1,692	1,480	1,455	1,314	1,288	1,040	557	288	62	11	1	1
1999.....	1,718	1,467	1,418	1,428	1,318	1,021	620	255	69	13	1	2

Non-Hispanic White, female

2003.....	728	736	685	735	894	932	704	355	112	12	1	-
2002.....	682	690	696	801	915	1,011	725	353	131	13	3	-
2001.....	692	616	698	794	907	989	676	352	120	20	-	-
2000.....	580	641	659	779	904	968	699	358	104	16	2	-
1999.....	542	546	686	763	973	1,006	700	347	118	16	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Non-Hispanic Black, both sexes

2003.....	546	409	291	239	162	79	56	31	9	2	-	-
2002.....	516	360	289	213	145	112	61	28	12	-	-	-
2001.....	530	390	259	245	167	116	53	22	10	2	-	1
2000.....	477	341	281	246	183	101	62	30	13	1	-	1
1999.....	398	363	328	250	182	128	65	37	13	2	-	-

Non-Hispanic Black, male

2003.....	403	294	198	147	86	40	30	16	4	1	-	-
2002.....	363	243	193	137	84	47	28	11	4	-	-	-
2001.....	383	278	184	168	100	57	18	7	4	-	-	1
2000.....	344	230	178	144	111	58	23	12	5	-	-	1
1999.....	292	254	199	166	107	67	29	13	4	-	-	-

Non-Hispanic Black, female

2003.....	143	115	93	92	76	39	26	15	5	1	-	-
2002.....	153	117	96	76	61	65	33	17	8	-	-	-
2001.....	147	112	75	77	67	59	35	15	6	2	-	-
2000.....	133	111	103	102	72	43	39	18	8	1	-	-
1999.....	106	109	129	84	75	61	36	24	9	2	-	-

Origin not stated 2/, both sexes

2003.....	17	13	17	8	3	4	3	-	-	-	-	1
2002.....	16	17	15	9	9	7	4	1	-	-	-	1
2001.....	15	14	13	13	9	7	4	1	-	2	-	2
2000.....	20	10	15	13	8	4	4	-	3	-	-	1
1999.....	20	11	10	9	6	8	2	2	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Origin not stated 2/, male

2003.....	14	12	15	5	3	1	1	-	-	-	-	1
2002.....	12	13	13	6	6	6	2	-	-	-	-	1
2001.....	12	12	11	11	4	1	1	1	-	1	-	2
2000.....	18	9	13	10	4	2	1	-	2	-	-	1
1999.....	15	10	8	8	5	4	-	1	-	-	-	-

Origin not stated 2/, female

2003.....	3	1	2	3	-	3	2	-	-	-	-	-
2002.....	4	4	2	3	3	1	2	1	-	-	-	-
2001.....	3	2	2	2	5	6	3	-	-	1	-	-
2000.....	2	1	2	3	4	2	3	-	1	-	-	-
1999.....	5	1	2	1	1	4	2	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Alcoholic liver disease (K70)												
All origins, both sexes												
2003.....	2,238	1,837	1,410	1,075	713	491	238	67	19	-	-	2
2002.....	2,139	1,712	1,371	1,023	826	495	235	79	16	3	-	2
2001.....	2,082	1,582	1,331	1,062	786	515	202	86	12	6	-	5
2000.....	1,888	1,575	1,394	1,085	806	533	218	76	17	2	-	2
1999.....	1,837	1,511	1,348	1,104	831	533	256	94	21	3	-	2
All origins, male												
2003.....	1,704	1,386	1,073	800	543	349	169	40	12	-	-	2
2002.....	1,624	1,277	1,060	771	596	334	161	54	10	1	-	2
2001.....	1,536	1,235	1,007	789	590	370	135	60	8	4	-	5
2000.....	1,458	1,172	1,044	821	612	419	142	52	12	1	-	2
1999.....	1,426	1,179	1,015	867	630	364	174	59	13	1	-	2
All origins, female												
2003.....	534	451	337	275	170	142	69	27	7	-	-	-
2002.....	515	435	311	252	230	161	74	25	6	2	-	-
2001.....	546	347	324	273	196	145	67	26	4	2	-	-
2000.....	430	403	350	264	194	114	76	24	5	1	-	-
1999.....	411	332	333	237	201	169	82	35	8	2	-	-
Hispanic, both sexes												
2003.....	264	255	192	137	79	64	26	7	1	-	-	-
2002.....	305	229	163	139	107	44	19	8	3	1	-	1
2001.....	274	220	176	140	100	68	26	8	2	-	-	1
2000.....	253	218	182	153	94	62	16	4	1	1	-	-
1999.....	229	196	160	124	89	51	18	14	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

Hispanic, male

2003.....	234	215	165	98	59	50	19	4	1	-	-	-
2002.....	257	200	139	122	89	30	14	4	3	1	-	1
2001.....	229	189	149	124	86	50	18	7	2	-	-	1
2000.....	218	191	153	139	82	56	13	3	1	1	-	-
1999.....	194	174	133	103	77	38	14	10	-	-	-	-

Hispanic, female

2003.....	30	40	27	39	20	14	7	3	-	-	-	-
2002.....	48	29	24	17	18	14	5	4	-	-	-	-
2001.....	45	31	27	16	14	18	8	1	-	-	-	-
2000.....	35	27	29	14	12	6	3	1	-	-	-	-
1999.....	35	22	27	21	12	13	4	4	2	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,964	1,575	1,208	934	632	426	212	60	18	-	-	2
2002.....	1,824	1,469	1,202	879	718	448	215	71	13	2	-	1
2001.....	1,797	1,350	1,149	914	682	446	176	78	10	6	-	2
2000.....	1,624	1,351	1,203	927	712	469	202	72	15	1	-	1
1999.....	1,597	1,309	1,182	975	738	479	238	80	19	3	-	2

Non-Hispanic 1/, male

2003.....	1,462	1,165	898	699	482	298	150	36	11	-	-	2
2002.....	1,359	1,066	915	644	506	301	146	50	7	-	-	1
2001.....	1,298	1,036	852	658	500	320	117	53	6	4	-	2
2000.....	1,230	976	883	678	530	361	129	49	10	-	-	1
1999.....	1,224	1,000	877	759	549	323	160	49	13	1	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

Non-Hispanic 1/, female

2003.....	502	410	310	235	150	128	62	24	7	-	-	-
2002.....	465	403	287	235	212	147	69	21	6	2	-	-
2001.....	499	314	297	256	182	126	59	25	4	2	-	-
2000.....	394	375	320	249	182	108	73	23	5	1	-	-
1999.....	373	309	305	216	189	156	78	31	6	2	-	-

Non-Hispanic White, both sexes

2003.....	1,612	1,332	1,010	804	564	398	194	55	16	-	-	2
2002.....	1,499	1,230	1,025	762	641	406	195	63	11	2	-	1
2001.....	1,431	1,105	987	773	608	404	162	66	8	6	-	2
2000.....	1,315	1,109	1,017	805	634	423	178	66	14	1	-	-
1999.....	1,314	1,065	980	829	651	421	215	67	16	3	-	2

Non-Hispanic White, male

2003.....	1,212	992	761	606	433	279	134	33	10	-	-	2
2002.....	1,131	915	786	560	452	275	134	44	5	-	-	1
2001.....	1,054	849	734	553	444	293	108	46	4	4	-	2
2000.....	1,012	821	758	594	471	329	117	45	9	-	-	-
1999.....	1,020	819	747	645	486	284	147	40	12	1	-	2

Non-Hispanic White, female

2003.....	400	340	249	198	131	119	60	22	6	-	-	-
2002.....	368	315	239	202	189	131	61	19	6	2	-	-
2001.....	377	256	253	220	164	111	54	20	4	2	-	-
2000.....	303	288	259	211	163	94	61	21	5	1	-	-
1999.....	294	246	233	184	165	137	68	27	4	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcoholic liver disease (K70)

Non-Hispanic Black, both sexes

2003.....	269	189	151	105	52	18	15	4	1	-	-	-
2002.....	252	180	135	88	55	33	15	7	1	-	-	-
2001.....	290	198	126	118	58	32	11	8	1	-	-	-
2000.....	247	184	137	106	63	35	15	4	1	-	-	1
1999.....	222	192	166	117	74	45	19	11	3	-	-	-

Non-Hispanic Black, male

2003.....	199	136	106	74	38	11	13	3	1	-	-	-
2002.....	178	112	100	64	39	21	9	5	1	-	-	-
2001.....	202	149	90	90	45	19	7	4	1	-	-	-
2000.....	175	121	87	73	50	24	9	3	1	-	-	1
1999.....	161	145	109	93	53	30	11	7	1	-	-	-

Non-Hispanic Black, female

2003.....	70	53	45	31	14	7	2	1	-	-	-	-
2002.....	74	68	35	24	16	12	6	2	-	-	-	-
2001.....	88	49	36	28	13	13	4	4	-	-	-	-
2000.....	72	63	50	33	13	11	6	1	-	-	-	-
1999.....	61	47	57	24	21	15	8	4	2	-	-	-

Origin not stated 2/, both sexes

2003.....	10	7	10	4	2	1	-	-	-	-	-	-
2002.....	10	14	6	5	1	3	1	-	-	-	-	-
2001.....	11	12	6	8	4	1	-	-	-	-	-	2
2000.....	11	6	9	5	-	2	-	-	1	-	-	1
1999.....	11	6	6	5	4	3	-	-	-	-	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

All origins, both sexes

2003.....	1,771	1,639	1,542	1,654	1,964	1,823	1,306	645	185	29	1	1
2002.....	1,673	1,546	1,468	1,647	1,885	1,917	1,353	649	214	21	3	2
2001.....	1,620	1,417	1,420	1,690	1,948	1,810	1,274	628	209	37	1	1
2000.....	1,392	1,372	1,433	1,666	1,925	1,834	1,244	666	186	28	4	1
1999.....	1,303	1,292	1,486	1,729	1,978	1,863	1,249	608	205	30	4	-

All origins, male

2003.....	1,300	1,099	979	936	986	832	551	264	68	13	-	1
2002.....	1,219	1,046	845	881	949	877	581	256	67	7	-	2
2001.....	1,178	926	860	960	996	781	547	247	81	15	1	1
2000.....	1,002	894	895	908	976	821	492	273	71	11	1	1
1999.....	957	872	877	970	987	844	523	234	73	13	2	-

All origins, female

2003.....	471	540	563	718	978	991	755	381	117	16	1	-
2002.....	454	500	623	766	936	1,040	772	393	147	14	3	-
2001.....	442	491	560	730	952	1,029	727	381	128	22	-	-
2000.....	390	478	538	758	949	1,013	752	393	115	17	3	-
1999.....	346	420	609	759	991	1,019	726	374	132	17	2	-

Hispanic, both sexes

2003.....	199	178	177	185	223	201	106	36	12	4	-	-
2002.....	200	193	173	193	224	154	104	50	18	2	-	-
2001.....	197	158	167	182	196	151	90	38	10	1	-	-
2000.....	161	162	146	180	201	153	92	47	18	1	1	-
1999.....	135	129	153	186	184	136	80	37	18	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Hispanic, male

2003.....	145	116	120	86	89	81	36	10	7	1	-	-
2002.....	151	133	89	104	92	65	48	14	5	-	-	-
2001.....	150	106	111	96	80	70	32	9	4	-	-	-
2000.....	124	104	83	94	80	65	29	14	11	-	-	-
1999.....	99	83	91	95	85	55	29	10	10	-	1	-

Hispanic, female

2003.....	54	62	57	99	134	120	70	26	5	3	-	-
2002.....	49	60	84	89	132	89	56	36	13	2	-	-
2001.....	47	52	56	86	116	81	58	29	6	1	-	-
2000.....	37	58	63	86	121	88	63	33	7	1	1	-
1999.....	36	46	62	91	99	81	51	27	8	1	-	-

Non-Hispanic 1/, both sexes

2003.....	1,565	1,455	1,358	1,465	1,740	1,619	1,197	609	173	25	1	-
2002.....	1,467	1,350	1,286	1,450	1,653	1,759	1,246	598	196	19	3	1
2001.....	1,419	1,257	1,246	1,503	1,747	1,653	1,180	589	199	34	1	1
2000.....	1,222	1,206	1,281	1,478	1,716	1,679	1,148	619	166	27	3	1
1999.....	1,159	1,158	1,329	1,539	1,792	1,722	1,167	569	185	29	3	-

Non-Hispanic 1/, male

2003.....	1,149	977	854	848	896	751	514	254	61	12	-	-
2002.....	1,064	911	749	776	852	809	532	242	62	7	-	1
2001.....	1,025	818	744	860	916	710	514	237	77	14	1	1
2000.....	870	786	807	808	892	756	462	259	59	11	1	1
1999.....	851	784	783	872	901	788	494	223	63	13	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic 1/, female

2003.....	416	478	504	617	844	868	683	355	112	13	1	-
2002.....	403	439	537	674	801	950	714	356	134	12	3	-
2001.....	394	439	502	643	831	943	666	352	122	20	-	-
2000.....	352	420	474	670	824	923	686	360	107	16	2	-
1999.....	308	374	546	667	891	934	673	346	122	16	2	-

Non-Hispanic White, both sexes

2003.....	1,254	1,192	1,172	1,295	1,593	1,520	1,127	572	162	22	1	-
2002.....	1,171	1,134	1,094	1,285	1,517	1,655	1,161	565	183	18	3	1
2001.....	1,126	1,022	1,075	1,339	1,587	1,541	1,115	562	188	31	1	-
2000.....	957	1,012	1,097	1,288	1,558	1,585	1,078	580	152	26	3	1
1999.....	946	948	1,124	1,362	1,640	1,606	1,105	535	171	26	3	-

Non-Hispanic White, male

2003.....	926	796	736	758	830	707	483	239	56	10	-	-
2002.....	857	759	637	686	791	775	497	231	58	7	-	1
2001.....	811	662	630	765	844	663	493	230	72	13	1	-
2000.....	680	659	697	720	817	711	440	243	53	11	1	1
1999.....	698	648	671	783	832	737	473	215	57	12	1	-

Non-Hispanic White, female

2003.....	328	396	436	537	763	813	644	333	106	12	1	-
2002.....	314	375	457	599	726	880	664	334	125	11	3	-
2001.....	315	360	445	574	743	878	622	332	116	18	-	-
2000.....	277	353	400	568	741	874	638	337	99	15	2	-
1999.....	248	300	453	579	808	869	632	320	114	14	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Non-Hispanic Black, both sexes

2003.....	277	220	140	134	110	61	41	27	8	2	-	-
2002.....	264	180	154	125	90	79	46	21	11	-	-	-
2001.....	240	192	133	127	109	84	42	14	9	2	-	1
2000.....	230	157	144	140	120	66	47	26	12	1	-	-
1999.....	176	171	162	133	108	83	46	26	10	2	-	-

Non-Hispanic Black, male

2003.....	204	158	92	73	48	29	17	13	3	1	-	-
2002.....	185	131	93	73	45	26	19	6	3	-	-	-
2001.....	181	129	94	78	55	38	11	3	3	-	-	1
2000.....	169	109	91	71	61	34	14	9	4	-	-	-
1999.....	131	109	90	73	54	37	18	6	3	-	-	-

Non-Hispanic Black, female

2003.....	73	62	48	61	62	32	24	14	5	1	-	-
2002.....	79	49	61	52	45	53	27	15	8	-	-	-
2001.....	59	63	39	49	54	46	31	11	6	2	-	-
2000.....	61	48	53	69	59	32	33	17	8	1	-	-
1999.....	45	62	72	60	54	46	28	20	7	2	-	-

Origin not stated 2/, both sexes

2003.....	7	6	7	4	1	3	3	-	-	-	-	1
2002.....	6	3	9	4	8	4	3	1	-	-	-	1
2001.....	4	2	7	5	5	6	4	1	-	2	-	-
2000.....	9	4	6	8	8	2	4	-	2	-	-	-
1999.....	9	5	4	4	2	5	2	2	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other chronic liver disease and cirrhosis (K73-K74)

Origin not stated 2/, male

2003.....	6	6	5	2	1	-	1	-	-	-	-	1
2002.....	4	2	7	1	5	3	1	-	-	-	-	1
2001.....	3	2	5	4	-	1	1	1	-	1	-	-
2000.....	8	4	5	6	4	-	1	-	1	-	-	-
1999.....	7	5	3	3	1	1	-	1	-	-	-	-

Origin not stated 2/, female

2003.....	1	-	2	2	-	3	2	-	-	-	-	-
2002.....	2	1	2	3	3	1	2	1	-	-	-	-
2001.....	1	-	2	1	5	5	3	-	-	1	-	-
2000.....	1	-	1	2	4	2	3	-	1	-	-	-
1999.....	2	-	1	1	1	4	2	1	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cholelithiasis and other disorders of gallbladder (K80-K82)

##### All origins, both sexes

2003.....	80	77	105	193	296	440	614	498	371	133	25	-
2002.....	77	93	123	194	310	420	572	525	351	150	28	-
2001.....	68	94	148	193	333	432	513	542	364	106	31	-
2000.....	66	80	109	184	266	419	507	528	358	143	24	-
1999.....	69	95	108	171	304	417	531	521	324	120	27	-

##### All origins, male

2003.....	44	45	61	101	164	233	286	199	99	27	3	-
2002.....	48	59	63	103	155	229	271	195	112	26	2	-
2001.....	40	61	81	96	173	231	233	209	111	21	8	-
2000.....	30	48	62	100	142	224	237	204	105	27	3	-
1999.....	30	54	66	89	163	208	239	203	101	22	4	-

##### All origins, female

2003.....	36	32	44	92	132	207	328	299	272	106	22	-
2002.....	29	34	60	91	155	191	301	330	239	124	26	-
2001.....	28	33	67	97	160	201	280	333	253	85	23	-
2000.....	36	32	47	84	124	195	270	324	253	116	21	-
1999.....	39	41	42	82	141	209	292	318	223	98	23	-

##### Hispanic, both sexes

2003.....	7	7	7	17	27	30	31	18	13	3	2	-
2002.....	6	9	16	17	25	29	26	24	11	8	1	-
2001.....	5	12	11	17	34	22	25	22	23	2	3	-
2000.....	11	8	11	22	19	26	25	25	12	7	-	-
1999.....	6	9	11	14	20	30	27	26	10	8	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cholelithiasis and other disorders of gallbladder (K80-K82)

##### Hispanic, male

2003.....	7	4	3	11	13	15	21	6	4	-	1	-
2002.....	6	8	6	7	14	15	8	9	5	2	-	-
2001.....	3	9	7	7	18	13	12	6	7	1	3	-
2000.....	5	6	9	12	13	18	14	6	4	2	-	-
1999.....	4	3	9	7	8	18	11	9	5	1	-	-

##### Hispanic, female

2003.....	-	3	4	6	14	15	10	12	9	3	1	-
2002.....	-	1	10	10	11	14	18	15	6	6	1	-
2001.....	2	3	4	10	16	9	13	16	16	1	-	-
2000.....	6	2	2	10	6	8	11	19	8	5	-	-
1999.....	2	6	2	7	12	12	16	17	5	7	2	-

##### Non-Hispanic 1/, both sexes

2003.....	72	70	98	176	269	409	581	479	358	130	23	-
2002.....	71	84	106	176	284	391	544	500	339	142	27	-
2001.....	63	82	137	174	299	410	486	520	341	104	28	-
2000.....	55	72	98	161	244	392	481	502	345	136	24	-
1999.....	63	86	97	157	281	387	502	494	314	112	25	-

##### Non-Hispanic 1/, male

2003.....	36	41	58	90	151	218	265	193	95	27	2	-
2002.....	42	51	56	96	140	214	262	185	107	24	2	-
2001.....	37	52	74	88	155	218	221	203	104	20	5	-
2000.....	25	42	53	87	128	205	223	198	100	25	3	-
1999.....	26	51	57	82	154	190	228	194	96	21	4	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cholelithiasis and other disorders of gallbladder (K80-K82)

##### Non-Hispanic 1/, female

2003.....	36	29	40	86	118	191	316	286	263	103	21	-
2002.....	29	33	50	80	144	177	282	315	232	118	25	-
2001.....	26	30	63	86	144	192	265	317	237	84	23	-
2000.....	30	30	45	74	116	187	258	304	245	111	21	-
1999.....	37	35	40	75	127	197	274	300	218	91	21	-

##### Non-Hispanic White, both sexes

2003.....	45	52	76	137	240	359	519	436	322	118	20	-
2002.....	49	68	84	150	240	334	484	458	315	134	23	-
2001.....	45	60	108	131	250	357	430	471	312	95	26	-
2000.....	38	56	76	131	194	347	428	463	322	128	23	-
1999.....	49	64	70	120	241	348	457	454	289	105	23	-

##### Non-Hispanic White, male

2003.....	24	31	44	71	137	193	237	178	86	25	1	-
2002.....	29	42	45	85	122	190	234	169	100	23	2	-
2001.....	28	39	57	65	135	196	199	181	94	18	4	-
2000.....	18	35	42	76	105	185	206	187	94	24	2	-
1999.....	21	35	44	65	136	173	216	175	88	21	4	-

##### Non-Hispanic White, female

2003.....	21	21	32	66	103	166	282	258	236	93	19	-
2002.....	20	26	39	65	118	144	250	289	215	111	21	-
2001.....	17	21	51	66	115	161	231	290	218	77	22	-
2000.....	20	21	34	55	89	162	222	276	228	104	21	-
1999.....	28	29	26	55	105	175	241	279	201	84	19	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Cholelithiasis and other disorders of gallbladder (K80-K82)

##### Non-Hispanic Black, both sexes

2003.....	21	15	18	29	21	35	43	35	21	12	3	-
2002.....	19	12	16	20	34	44	47	26	14	6	4	-
2001.....	17	20	22	32	46	35	42	33	21	8	2	-
2000.....	14	13	16	25	36	36	39	33	18	6	-	-
1999.....	13	14	21	29	30	26	39	25	15	6	2	-

##### Non-Hispanic Black, male

2003.....	10	8	10	15	8	18	19	11	6	2	1	-
2002.....	10	6	10	7	10	16	21	8	3	-	-	-
2001.....	9	12	12	16	18	10	11	15	6	2	1	-
2000.....	6	5	9	8	15	14	12	9	3	-	-	-
1999.....	4	10	9	12	13	12	11	9	4	-	-	-

##### Non-Hispanic Black, female

2003.....	11	7	8	14	13	17	24	24	15	10	2	-
2002.....	9	6	6	13	24	28	26	18	11	6	4	-
2001.....	8	8	10	16	28	25	31	18	15	6	1	-
2000.....	8	8	7	17	21	22	27	24	15	6	-	-
1999.....	9	4	12	17	17	14	28	16	11	6	2	-

##### Origin not stated 2/, both sexes

2003.....	1	-	-	-	-	1	2	1	-	-	-	-
2002.....	-	-	1	1	1	-	2	1	1	-	-	-
2001.....	-	-	-	2	-	-	2	-	-	-	-	-
2000.....	-	-	-	1	3	1	1	1	1	-	-	-
1999.....	-	-	-	-	3	-	2	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Origin not stated 2/, male

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	1	-	1	-	1	1	-	-	-	-
2001.....	-	-	-	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	1	1	-	-	1	-	-	-
1999.....	-	-	-	-	1	-	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	1	2	1	-	-	-	-
2002.....	-	-	-	1	-	-	1	-	1	-	-	-
2001.....	-	-	-	1	-	-	2	-	-	-	-	-
2000.....	-	-	-	-	2	-	1	1	-	-	-	-
1999.....	-	-	-	-	2	-	2	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins, both sexes

2003.....	1,166	1,638	2,168	2,996	4,349	6,298	7,795	7,277	4,584	1,641	314	2
2002.....	1,103	1,505	1,950	2,820	4,344	6,338	7,558	6,959	4,450	1,549	298	-
2001.....	1,073	1,404	1,880	3,026	4,330	6,105	6,992	6,612	4,225	1,528	303	1
2000.....	964	1,269	1,831	2,811	4,179	5,893	6,565	6,150	3,915	1,443	269	2
1999.....	872	1,176	1,688	2,694	4,147	5,707	6,220	5,825	3,703	1,388	254	-

All origins, male

2003.....	711	936	1,129	1,565	2,220	3,188	3,861	3,232	1,795	492	81	2
2002.....	632	797	1,027	1,491	2,311	3,248	3,672	3,088	1,693	495	82	-
2001.....	608	751	963	1,610	2,216	3,074	3,411	2,996	1,619	435	64	-
2000.....	549	697	954	1,444	2,120	3,039	3,174	2,672	1,473	453	73	2
1999.....	481	627	903	1,366	2,138	2,986	3,045	2,584	1,353	424	67	-

All origins, female

2003.....	455	702	1,039	1,431	2,129	3,110	3,934	4,045	2,789	1,149	233	-
2002.....	471	708	923	1,329	2,033	3,090	3,886	3,871	2,757	1,054	216	-
2001.....	465	653	917	1,416	2,114	3,031	3,581	3,616	2,606	1,093	239	1
2000.....	415	572	877	1,367	2,059	2,854	3,391	3,478	2,442	990	196	-
1999.....	391	549	785	1,328	2,009	2,721	3,175	3,241	2,350	964	187	-

Hispanic, both sexes

2003.....	104	146	172	225	259	320	258	248	136	56	10	1
2002.....	100	119	148	216	211	281	251	179	120	45	19	-
2001.....	81	117	120	218	252	296	228	180	121	44	10	-
2000.....	64	87	145	195	209	235	194	182	127	31	6	1
1999.....	57	96	134	172	197	239	187	167	111	48	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Hispanic, male

2003.....	68	75	85	111	119	151	129	116	49	23	5	1
2002.....	59	64	77	115	99	144	129	72	41	21	7	-
2001.....	37	63	69	100	117	150	101	82	48	15	3	-
2000.....	37	45	70	90	104	116	90	79	52	10	4	1
1999.....	34	58	79	84	94	133	79	88	46	21	1	-

Hispanic, female

2003.....	36	71	87	114	140	169	129	132	87	33	5	-
2002.....	41	55	71	101	112	137	122	107	79	24	12	-
2001.....	44	54	51	118	135	146	127	98	73	29	7	-
2000.....	27	42	75	105	105	119	104	103	75	21	2	-
1999.....	23	38	55	88	103	106	108	79	65	27	4	-

Non-Hispanic 1/, both sexes

2003.....	1,057	1,487	1,985	2,764	4,084	5,961	7,520	7,011	4,434	1,582	303	1
2002.....	997	1,377	1,794	2,597	4,113	6,035	7,287	6,760	4,316	1,496	279	-
2001.....	985	1,282	1,754	2,803	4,063	5,796	6,742	6,418	4,093	1,484	293	-
2000.....	894	1,180	1,674	2,611	3,956	5,643	6,357	5,958	3,784	1,405	263	-
1999.....	811	1,073	1,548	2,514	3,937	5,459	6,015	5,636	3,577	1,335	249	-

Non-Hispanic 1/, male

2003.....	640	857	1,038	1,448	2,098	3,029	3,724	3,109	1,740	468	76	1
2002.....	570	730	947	1,372	2,197	3,089	3,531	3,008	1,648	470	75	-
2001.....	567	687	890	1,507	2,091	2,920	3,300	2,908	1,568	420	61	-
2000.....	507	651	878	1,351	2,006	2,918	3,077	2,591	1,421	440	69	-
1999.....	443	565	819	1,276	2,036	2,849	2,959	2,484	1,302	402	66	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic 1/, female

2003.....	417	630	947	1,316	1,986	2,932	3,796	3,902	2,694	1,114	227	-
2002.....	427	647	847	1,225	1,916	2,946	3,756	3,752	2,668	1,026	204	-
2001.....	418	595	864	1,296	1,972	2,876	3,442	3,510	2,525	1,064	232	-
2000.....	387	529	796	1,260	1,950	2,725	3,280	3,367	2,363	965	194	-
1999.....	368	508	729	1,238	1,901	2,610	3,056	3,152	2,275	933	183	-

Non-Hispanic White, both sexes

2003.....	575	907	1,290	1,886	3,005	4,813	6,366	6,147	3,881	1,372	247	-
2002.....	519	822	1,156	1,794	3,076	4,879	6,231	5,901	3,787	1,280	224	-
2001.....	552	763	1,098	1,907	3,077	4,686	5,712	5,617	3,572	1,293	241	-
2000.....	495	704	1,072	1,776	2,965	4,616	5,410	5,202	3,318	1,223	210	-
1999.....	445	645	991	1,738	2,980	4,440	5,064	4,807	3,126	1,140	195	-

Non-Hispanic White, male

2003.....	336	528	688	1,027	1,587	2,540	3,210	2,812	1,537	399	62	-
2002.....	299	438	607	983	1,711	2,583	3,104	2,679	1,454	406	55	-
2001.....	334	425	587	1,072	1,657	2,454	2,856	2,598	1,399	372	53	-
2000.....	283	400	590	983	1,591	2,449	2,692	2,318	1,268	375	62	-
1999.....	244	365	538	922	1,610	2,359	2,548	2,144	1,161	362	54	-

Non-Hispanic White, female

2003.....	239	379	602	859	1,418	2,273	3,156	3,335	2,344	973	185	-
2002.....	220	384	549	811	1,365	2,296	3,127	3,222	2,333	874	169	-
2001.....	218	338	511	835	1,420	2,232	2,856	3,019	2,173	921	188	-
2000.....	212	304	482	793	1,374	2,167	2,718	2,884	2,050	848	148	-
1999.....	201	280	453	816	1,370	2,081	2,516	2,663	1,965	778	141	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Non-Hispanic Black, both sexes

2003.....	448	538	634	795	963	998	1,017	769	494	190	48	1
2002.....	449	497	575	737	933	1,024	917	759	462	195	52	-
2001.....	395	459	589	817	881	982	909	707	454	176	50	-
2000.....	364	433	547	748	889	923	853	667	411	168	44	-
1999.....	346	378	501	698	856	911	839	727	397	173	54	-

Non-Hispanic Black, male

2003.....	286	306	321	381	472	417	445	252	171	58	13	1
2002.....	256	267	306	353	441	444	365	282	165	54	19	-
2001.....	211	240	269	397	385	402	382	260	142	39	7	-
2000.....	203	226	258	330	376	418	341	233	126	59	5	-
1999.....	190	171	255	310	381	435	368	292	114	30	12	-

Non-Hispanic Black, female

2003.....	162	232	313	414	491	581	572	517	323	132	35	-
2002.....	193	230	269	384	492	580	552	477	297	141	33	-
2001.....	184	219	320	420	496	580	527	447	312	137	43	-
2000.....	161	207	289	418	513	505	512	434	285	109	39	-
1999.....	156	207	246	388	475	476	471	435	283	143	42	-

Origin not stated 2/, both sexes

2003.....	5	5	11	7	6	17	17	18	14	3	1	-
2002.....	6	9	8	7	20	22	20	20	14	8	-	-
2001.....	7	5	6	5	15	13	22	14	11	-	-	1
2000.....	6	2	12	5	14	15	14	10	4	7	-	1
1999.....	4	7	6	8	13	9	18	22	15	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Origin not stated 2/, male

2003.....	3	4	6	6	3	8	8	7	6	1	-	-
2002.....	3	3	3	4	15	15	12	8	4	4	-	-
2001.....	4	1	4	3	8	4	10	6	3	-	-	-
2000.....	5	1	6	3	10	5	7	2	-	3	-	1
1999.....	4	4	5	6	8	4	7	12	5	1	-	-

Origin not stated 2/, female

2003.....	2	1	5	1	3	9	9	11	8	2	1	-
2002.....	3	6	5	3	5	7	8	12	10	4	-	-
2001.....	3	4	2	2	7	9	12	8	8	-	-	1
2000.....	1	1	6	2	4	10	7	8	4	4	-	-
1999.....	-	3	1	2	5	5	11	10	10	4	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All origins, both sexes

2003.....	4	5	9	11	19	18	32	17	5	2	2	-
2002.....	4	6	10	13	14	19	32	23	13	5	1	-
2001.....	5	9	4	13	14	28	20	24	12	1	1	-
2000.....	11	2	8	13	16	16	36	27	13	8	1	-
1999.....	5	5	11	10	27	37	25	22	7	5	-	-

All origins, male

2003.....	1	3	4	6	9	7	13	2	3	-	-	-
2002.....	1	5	6	9	5	5	12	10	2	1	1	-
2001.....	4	6	3	6	4	10	9	10	5	-	-	-
2000.....	9	2	6	7	5	11	15	13	5	4	-	-
1999.....	3	2	4	5	12	15	17	9	1	4	-	-

All origins, female

2003.....	3	2	5	5	10	11	19	15	2	2	2	-
2002.....	3	1	4	4	9	14	20	13	11	4	-	-
2001.....	1	3	1	7	10	18	11	14	7	1	1	-
2000.....	2	-	2	6	11	5	21	14	8	4	1	-
1999.....	2	3	7	5	15	22	8	13	6	1	-	-

Hispanic, both sexes

2003.....	-	-	1	1	2	1	1	1	-	1	-	-
2002.....	-	1	1	2	-	-	2	-	-	-	-	-
2001.....	2	1	1	1	2	-	1	-	2	-	-	-
2000.....	2	-	-	1	1	-	1	1	-	-	-	-
1999.....	-	-	2	3	1	1	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Hispanic, male

2003.....	-	-	-	-	-	1	1	1	-	-	-	-
2002.....	-	1	1	1	-	-	1	-	-	-	-	-
2001.....	2	1	1	1	1	-	-	-	-	-	-	-
2000.....	1	-	-	-	-	-	1	1	-	-	-	-
1999.....	-	-	1	-	-	-	-	1	-	-	-	-

Hispanic, female

2003.....	-	-	1	1	2	-	-	-	-	1	-	-
2002.....	-	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	-	1	-	1	-	2	-	-	-
2000.....	1	-	-	1	1	-	-	-	-	-	-	-
1999.....	-	-	1	3	1	1	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	4	5	8	10	17	17	31	16	5	1	2	-
2002.....	4	5	9	11	14	19	30	23	12	5	1	-
2001.....	3	8	3	12	12	28	18	24	10	1	1	-
2000.....	9	2	8	12	15	16	35	26	13	8	1	-
1999.....	5	5	9	7	26	36	25	21	7	5	-	-

Non-Hispanic 1/, male

2003.....	1	3	4	6	9	6	12	1	3	-	-	-
2002.....	1	4	5	8	5	5	11	10	2	1	1	-
2001.....	2	5	2	5	3	10	8	10	5	-	-	-
2000.....	8	2	6	7	5	11	14	12	5	4	-	-
1999.....	3	2	3	5	12	15	17	8	1	4	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Non-Hispanic 1/, female

2003.....	3	2	4	4	8	11	19	15	2	1	2	-
2002.....	3	1	4	3	9	14	19	13	10	4	-	-
2001.....	1	3	1	7	9	18	10	14	5	1	1	-
2000.....	1	-	2	5	10	5	21	14	8	4	1	-
1999.....	2	3	6	2	14	21	8	13	6	1	-	-

Non-Hispanic White, both sexes

2003.....	4	3	5	8	15	15	25	13	5	1	-	-
2002.....	2	4	8	7	12	17	23	17	9	5	1	-
2001.....	2	6	3	8	10	25	16	20	7	1	1	-
2000.....	2	2	8	7	13	12	33	22	12	8	1	-
1999.....	3	3	4	6	22	31	20	16	7	5	-	-

Non-Hispanic White, male

2003.....	1	2	1	4	8	4	8	1	3	-	-	-
2002.....	1	3	5	5	5	4	9	8	1	1	1	-
2001.....	2	3	2	2	2	10	8	8	4	-	-	-
2000.....	2	2	6	4	5	9	13	9	4	4	-	-
1999.....	2	1	1	4	10	13	13	4	1	4	-	-

Non-Hispanic White, female

2003.....	3	1	4	4	7	11	17	12	2	1	-	-
2002.....	1	1	3	2	7	13	14	9	8	4	-	-
2001.....	-	3	1	6	8	15	8	12	3	1	1	-
2000.....	-	-	2	3	8	3	20	13	8	4	1	-
1999.....	1	2	3	2	12	18	7	12	6	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All origins, both sexes

2003.....	30	22	28	42	60	77	88	72	37	16	5	-
2002.....	25	33	19	29	56	86	92	74	38	14	1	-
2001.....	26	17	23	47	64	69	78	70	38	13	1	-
2000.....	21	31	28	43	60	78	94	78	23	15	2	-
1999.....	28	26	34	40	61	104	96	83	37	10	2	-

All origins, male

2003.....	23	10	12	24	24	39	41	32	20	6	1	-
2002.....	15	20	7	14	35	50	42	32	19	2	-	-
2001.....	15	10	15	23	40	39	45	32	10	6	-	-
2000.....	14	16	15	24	30	45	38	40	9	6	-	-
1999.....	15	13	19	20	25	58	49	38	17	5	-	-

All origins, female

2003.....	7	12	16	18	36	38	47	40	17	10	4	-
2002.....	10	13	12	15	21	36	50	42	19	12	1	-
2001.....	11	7	8	24	24	30	33	38	28	7	1	-
2000.....	7	15	13	19	30	33	56	38	14	9	2	-
1999.....	13	13	15	20	36	46	47	45	20	5	2	-

Hispanic, both sexes

2003.....	3	5	2	3	2	4	5	1	1	1	-	-
2002.....	1	3	2	2	1	7	2	3	2	-	-	-
2001.....	1	2	2	2	4	4	2	4	1	-	-	-
2000.....	2	2	2	2	3	3	4	5	-	-	-	-
1999.....	-	3	3	5	3	2	3	2	2	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Hispanic, male

2003.....	3	3	-	2	1	1	3	-	-	-	-	-
2002.....	-	3	1	-	-	5	1	1	1	-	-	-
2001.....	-	1	2	1	1	2	2	2	-	-	-	-
2000.....	1	1	2	-	2	2	1	5	-	-	-	-
1999.....	-	2	2	4	-	1	-	1	-	1	-	-

Hispanic, female

2003.....	-	2	2	1	1	3	2	1	1	1	-	-
2002.....	1	-	1	2	1	2	1	2	1	-	-	-
2001.....	1	1	-	1	3	2	-	2	1	-	-	-
2000.....	1	1	-	2	1	1	3	-	-	-	-	-
1999.....	-	1	1	1	3	1	3	1	2	-	-	-

Non-Hispanic 1/, both sexes

2003.....	27	17	26	39	58	73	83	71	36	15	5	-
2002.....	24	30	17	27	55	79	90	71	36	14	1	-
2001.....	25	15	21	45	60	65	76	66	37	13	1	-
2000.....	19	29	26	41	57	75	90	73	23	15	2	-
1999.....	28	23	30	35	58	102	93	81	35	9	2	-

Non-Hispanic 1/, male

2003.....	20	7	12	22	23	38	38	32	20	6	1	-
2002.....	15	17	6	14	35	45	41	31	18	2	-	-
2001.....	15	9	13	22	39	37	43	30	10	6	-	-
2000.....	13	15	13	24	28	43	37	35	9	6	-	-
1999.....	15	11	16	16	25	57	49	37	17	4	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

Non-Hispanic 1/, female

2003.....	7	10	14	17	35	35	45	39	16	9	4	-
2002.....	9	13	11	13	20	34	49	40	18	12	1	-
2001.....	10	6	8	23	21	28	33	36	27	7	1	-
2000.....	6	14	13	17	29	32	53	38	14	9	2	-
1999.....	13	12	14	19	33	45	44	44	18	5	2	-

Non-Hispanic White, both sexes

2003.....	16	16	19	33	41	58	70	63	27	14	4	-
2002.....	14	21	13	26	50	61	80	57	33	12	-	-
2001.....	12	11	13	33	54	57	66	62	35	12	-	-
2000.....	9	22	17	30	50	64	79	59	17	12	1	-
1999.....	18	15	21	24	48	89	80	68	27	7	2	-

Non-Hispanic White, male

2003.....	13	7	9	19	16	29	32	28	15	6	1	-
2002.....	8	13	6	14	31	35	37	27	17	2	-	-
2001.....	7	6	8	16	36	32	39	29	10	6	-	-
2000.....	6	12	8	18	25	37	35	29	6	4	-	-
1999.....	10	7	12	13	20	51	42	31	14	3	-	-

Non-Hispanic White, female

2003.....	3	9	10	14	25	29	38	35	12	8	3	-
2002.....	6	8	7	12	19	26	43	30	16	10	-	-
2001.....	5	5	5	17	18	25	27	33	25	6	-	-
2000.....	3	10	9	12	25	27	44	30	11	8	1	-
1999.....	8	8	9	11	28	38	38	37	13	4	2	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Renal failure (N17-N19)

## All origins, both sexes

2003.....	1,131	1,611	2,130	2,939	4,267	6,198	7,672	7,184	4,541	1,621	307	2
2002.....	1,072	1,464	1,919	2,774	4,272	6,228	7,429	6,856	4,398	1,530	296	-
2001.....	1,041	1,376	1,850	2,963	4,248	6,007	6,891	6,517	4,174	1,514	300	1
2000.....	932	1,234	1,789	2,754	4,094	5,794	6,429	6,040	3,878	1,420	265	2
1999.....	836	1,144	1,640	2,641	4,056	5,560	6,092	5,718	3,656	1,372	252	-

## All origins, male

2003.....	686	923	1,113	1,532	2,187	3,140	3,807	3,198	1,771	486	80	2
2002.....	615	772	1,014	1,468	2,271	3,190	3,618	3,046	1,672	492	81	-
2001.....	588	735	945	1,581	2,171	3,025	3,356	2,953	1,604	429	64	-
2000.....	526	678	929	1,412	2,080	2,981	3,121	2,617	1,458	443	73	2
1999.....	462	612	880	1,340	2,100	2,912	2,976	2,535	1,334	415	67	-

## All origins, female

2003.....	445	688	1,017	1,407	2,080	3,058	3,865	3,986	2,770	1,135	227	-
2002.....	457	692	905	1,306	2,001	3,038	3,811	3,810	2,726	1,038	215	-
2001.....	453	641	905	1,382	2,077	2,982	3,535	3,564	2,570	1,085	236	1
2000.....	406	556	860	1,342	2,014	2,813	3,308	3,423	2,420	977	192	-
1999.....	374	532	760	1,301	1,956	2,648	3,116	3,183	2,322	957	185	-

## Hispanic, both sexes

2003.....	101	141	169	221	255	315	252	246	135	54	10	1
2002.....	99	115	145	212	210	274	247	176	118	45	19	-
2001.....	78	114	116	214	246	292	225	176	118	44	10	-
2000.....	60	85	142	192	205	231	189	176	127	31	6	1
1999.....	57	93	129	163	193	234	183	164	109	47	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Renal failure (N17-N19)

## Hispanic, male

2003.....	65	72	85	109	118	149	125	115	49	23	5	1
2002.....	59	60	75	114	99	139	127	71	40	21	7	-
2001.....	35	61	66	98	115	148	99	80	48	15	3	-
2000.....	35	44	67	90	102	114	88	73	52	10	4	1
1999.....	34	56	76	80	94	131	78	86	46	20	1	-

## Hispanic, female

2003.....	36	69	84	112	137	166	127	131	86	31	5	-
2002.....	40	55	70	98	111	135	120	105	78	24	12	-
2001.....	43	53	50	116	131	144	126	96	70	29	7	-
2000.....	25	41	75	102	103	117	101	103	75	21	2	-
1999.....	23	37	53	83	99	103	105	78	63	27	4	-

## Non-Hispanic 1/, both sexes

2003.....	1,025	1,465	1,950	2,711	4,006	5,866	7,403	6,920	4,392	1,564	296	1
2002.....	967	1,340	1,766	2,555	4,042	5,932	7,162	6,660	4,267	1,477	277	-
2001.....	956	1,257	1,728	2,744	3,987	5,702	6,645	6,327	4,045	1,470	290	-
2000.....	866	1,147	1,636	2,557	3,875	5,548	6,226	5,854	3,747	1,382	259	-
1999.....	775	1,044	1,506	2,470	3,850	5,317	5,891	5,532	3,532	1,320	247	-

## Non-Hispanic 1/, male

2003.....	618	847	1,022	1,417	2,066	2,983	3,674	3,076	1,716	462	75	1
2002.....	553	709	936	1,350	2,157	3,036	3,479	2,967	1,628	467	74	-
2001.....	549	673	875	1,480	2,048	2,873	3,248	2,867	1,553	414	61	-
2000.....	486	633	856	1,319	1,968	2,862	3,026	2,542	1,406	430	69	-
1999.....	424	552	800	1,254	1,998	2,777	2,891	2,437	1,283	394	66	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Renal failure (N17-N19)

## Non-Hispanic 1/, female

2003.....	407	618	928	1,294	1,940	2,883	3,729	3,844	2,676	1,102	221	-
2002.....	414	631	830	1,205	1,885	2,896	3,683	3,693	2,639	1,010	203	-
2001.....	407	584	853	1,264	1,939	2,829	3,397	3,460	2,492	1,056	229	-
2000.....	380	514	780	1,238	1,907	2,686	3,200	3,312	2,341	952	190	-
1999.....	351	492	706	1,216	1,852	2,540	3,000	3,095	2,249	926	181	-

## Non-Hispanic White, both sexes

2003.....	554	888	1,265	1,844	2,947	4,735	6,269	6,067	3,848	1,355	243	-
2002.....	502	795	1,133	1,757	3,012	4,797	6,125	5,822	3,744	1,263	223	-
2001.....	537	745	1,080	1,864	3,009	4,604	5,629	5,535	3,529	1,280	240	-
2000.....	484	678	1,044	1,738	2,894	4,537	5,293	5,117	3,288	1,203	208	-
1999.....	422	627	965	1,707	2,907	4,317	4,962	4,721	3,091	1,127	193	-

## Non-Hispanic White, male

2003.....	321	519	678	1,003	1,563	2,505	3,170	2,783	1,518	393	61	-
2002.....	290	422	596	964	1,675	2,542	3,058	2,644	1,436	403	54	-
2001.....	324	416	577	1,054	1,618	2,412	2,809	2,561	1,385	366	53	-
2000.....	275	385	574	960	1,556	2,401	2,644	2,278	1,257	367	62	-
1999.....	231	357	525	905	1,579	2,295	2,492	2,107	1,145	355	54	-

## Non-Hispanic White, female

2003.....	233	369	587	841	1,384	2,230	3,099	3,284	2,330	962	182	-
2002.....	212	373	537	793	1,337	2,255	3,067	3,178	2,308	860	169	-
2001.....	213	329	503	810	1,391	2,192	2,820	2,974	2,144	914	187	-
2000.....	209	293	470	778	1,338	2,136	2,649	2,839	2,031	836	146	-
1999.....	191	270	440	802	1,328	2,022	2,470	2,614	1,946	772	139	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Renal failure (N17-N19)

## Non-Hispanic Black, both sexes

2003.....	439	536	627	786	950	985	1,004	765	486	190	46	1
2002.....	437	491	572	733	927	1,009	905	741	458	193	51	-
2001.....	385	455	583	803	875	972	899	701	450	175	48	-
2000.....	348	428	540	735	881	910	841	655	407	166	43	-
1999.....	334	370	490	687	846	896	824	711	390	171	54	-

## Non-Hispanic Black, male

2003.....	280	306	317	375	466	409	438	251	167	58	13	1
2002.....	249	263	306	351	437	435	361	277	164	54	19	-
2001.....	204	237	265	389	382	399	379	257	142	39	7	-
2000.....	191	224	253	322	374	413	338	228	124	58	5	-
1999.....	184	168	250	306	376	428	361	284	112	29	12	-

## Non-Hispanic Black, female

2003.....	159	230	310	411	484	576	566	514	319	132	33	-
2002.....	188	228	266	382	490	574	544	464	294	139	32	-
2001.....	181	218	318	414	493	573	520	444	308	136	41	-
2000.....	157	204	287	413	507	497	503	427	283	108	38	-
1999.....	150	202	240	381	470	468	463	427	278	142	42	-

## Origin not stated 2/, both sexes

2003.....	5	5	11	7	6	17	17	18	14	3	1	-
2002.....	6	9	8	7	20	22	20	20	13	8	-	-
2001.....	7	5	6	5	15	13	21	14	11	-	-	1
2000.....	6	2	11	5	14	15	14	10	4	7	-	1
1999.....	4	7	5	8	13	9	18	22	15	5	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Renal failure (N17-N19)

Origin not stated 2/, male

2003.....	3	4	6	6	3	8	8	7	6	1	-	-
2002.....	3	3	3	4	15	15	12	8	4	4	-	-
2001.....	4	1	4	3	8	4	9	6	3	-	-	-
2000.....	5	1	6	3	10	5	7	2	-	3	-	1
1999.....	4	4	4	6	8	4	7	12	5	1	-	-

Origin not stated 2/, female

2003.....	2	1	5	1	3	9	9	11	8	2	1	-
2002.....	3	6	5	3	5	7	8	12	9	4	-	-
2001.....	3	4	2	2	7	9	12	8	8	-	-	1
2000.....	1	1	5	2	4	10	7	8	4	4	-	-
1999.....	-	3	1	2	5	5	11	10	10	4	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of kidney (N25,N27)

All origins, both sexes

2003.....	1	-	1	4	3	5	3	4	1	2	-	-
2002.....	2	2	2	4	2	5	5	6	1	-	-	-
2001.....	1	2	3	3	4	1	3	1	1	-	1	-
2000.....	-	2	6	1	9	5	6	5	1	-	1	-
1999.....	3	1	3	3	3	6	7	2	3	1	-	-

All origins, male

2003.....	1	-	-	3	-	2	-	-	1	-	-	-
2002.....	1	-	-	-	-	3	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	1	1	-	-	-	-
2000.....	-	1	4	1	5	2	-	2	1	-	-	-
1999.....	1	-	-	1	1	1	3	2	1	-	-	-

All origins, female

2003.....	-	-	1	1	3	3	3	4	-	2	-	-
2002.....	1	2	2	4	2	2	5	6	1	-	-	-
2001.....	-	2	3	3	3	1	2	-	1	-	1	-
2000.....	-	1	2	-	4	3	6	3	-	-	1	-
1999.....	2	1	3	2	2	5	4	-	2	1	-	-

Hispanic, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	-	2	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of kidney (N25,N27)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	1	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	1	1	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	1	1	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	-	1	-	1	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1	-	1	4	3	5	3	4	1	2	-	-
2002.....	2	2	2	4	2	5	5	6	1	-	-	-
2001.....	1	2	2	2	4	1	3	1	1	-	1	-
2000.....	-	2	4	1	9	4	6	5	1	-	1	-
1999.....	3	1	3	2	3	4	6	2	3	1	-	-

Non-Hispanic 1/, male

2003.....	1	-	-	3	-	2	-	-	1	-	-	-
2002.....	1	-	-	-	-	3	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	1	1	-	-	-	-
2000.....	-	1	3	1	5	2	-	2	1	-	-	-
1999.....	1	-	-	1	1	-	2	2	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other disorders of kidney (N25,N27)

Non-Hispanic 1/, female

2003.....	-	-	1	1	3	3	3	4	-	2	-	-
2002.....	1	2	2	4	2	2	5	6	1	-	-	-
2001.....	-	2	2	2	3	1	2	-	1	-	1	-
2000.....	-	1	1	-	4	2	6	3	-	-	1	-
1999.....	2	1	3	1	2	4	4	-	2	1	-	-

Non-Hispanic White, both sexes

2003.....	1	-	1	1	2	5	2	4	1	2	-	-
2002.....	1	2	2	4	2	4	3	5	1	-	-	-
2001.....	1	1	2	2	4	-	1	-	1	-	-	-
2000.....	-	2	3	1	8	3	5	4	1	-	-	-
1999.....	2	-	1	1	3	3	2	2	1	1	-	-

Non-Hispanic White, male

2003.....	1	-	-	1	-	2	-	-	1	-	-	-
2002.....	-	-	-	-	-	2	-	-	-	-	-	-
2001.....	1	-	-	-	1	-	-	-	-	-	-	-
2000.....	-	1	2	1	5	2	-	2	1	-	-	-
1999.....	1	-	-	-	1	-	1	2	1	-	-	-

Non-Hispanic White, female

2003.....	-	-	1	-	2	3	2	4	-	2	-	-
2002.....	1	2	2	4	2	2	3	5	1	-	-	-
2001.....	-	1	2	2	3	-	1	-	1	-	-	-
2000.....	-	1	1	-	3	1	5	2	-	-	-	-
1999.....	1	-	1	1	2	3	1	-	-	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Infections of kidney (N10-N12,N13.6,N15.1)

##### All origins, both sexes

2003.....	30	34	28	56	49	95	152	141	91	40	6	-
2002.....	21	26	33	48	62	118	120	146	105	36	3	-
2001.....	25	23	23	35	52	126	123	129	100	33	3	-
2000.....	21	25	34	38	66	114	134	145	114	44	17	-
1999.....	33	30	29	52	81	109	133	154	102	49	6	-

##### All origins, male

2003.....	11	11	10	15	16	25	49	44	27	7	-	-
2002.....	10	10	11	19	28	44	28	42	25	7	-	-
2001.....	5	6	6	19	21	36	40	32	16	8	-	-
2000.....	9	5	10	13	13	34	40	32	30	5	2	-
1999.....	11	14	13	19	25	37	46	45	24	12	-	-

##### All origins, female

2003.....	19	23	18	41	33	70	103	97	64	33	6	-
2002.....	11	16	22	29	34	74	92	104	80	29	3	-
2001.....	20	17	17	16	31	90	83	97	84	25	3	-
2000.....	12	20	24	25	53	80	94	113	84	39	15	-
1999.....	22	16	16	33	56	72	87	109	78	37	6	-

##### Hispanic, both sexes

2003.....	1	7	1	6	5	7	6	4	2	3	-	-
2002.....	2	-	2	2	10	9	5	2	5	1	-	-
2001.....	3	1	-	4	4	4	5	3	2	1	-	-
2000.....	1	1	3	4	3	4	6	4	6	1	-	-
1999.....	1	6	3	3	5	6	9	4	4	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Infections of kidney (N10-N12,N13.6,N15.1)

##### Hispanic, male

2003.....	-	1	-	1	-	3	1	1	2	1	-	-
2002.....	2	-	1	-	4	5	-	1	1	1	-	-
2001.....	1	-	-	1	2	1	-	2	1	-	-	-
2000.....	1	-	1	2	1	-	2	1	-	-	-	-
1999.....	-	3	2	2	2	2	4	-	1	1	-	-

##### Hispanic, female

2003.....	1	6	1	5	5	4	5	3	-	2	-	-
2002.....	-	-	1	2	6	4	5	1	4	-	-	-
2001.....	2	1	-	3	2	3	5	1	1	1	-	-
2000.....	-	1	2	2	2	4	4	3	6	1	-	-
1999.....	1	3	1	1	3	4	5	4	3	-	-	-

##### Non-Hispanic 1/, both sexes

2003.....	29	27	27	50	44	87	146	137	89	37	6	-
2002.....	19	26	31	46	52	109	115	143	100	35	3	-
2001.....	22	22	23	31	48	122	117	126	98	32	3	-
2000.....	20	24	31	33	63	109	127	141	108	43	17	-
1999.....	32	24	25	49	76	101	124	150	98	47	6	-

##### Non-Hispanic 1/, male

2003.....	11	10	10	14	16	22	48	43	25	6	-	-
2002.....	8	10	10	19	24	39	28	41	24	6	-	-
2001.....	4	6	6	18	19	35	40	30	15	8	-	-
2000.....	8	5	9	10	12	34	37	31	30	5	2	-
1999.....	11	11	11	17	23	34	42	45	23	11	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic 1/, female

2003.....	18	17	17	36	28	65	98	94	64	31	6	-
2002.....	11	16	21	27	28	70	87	102	76	29	3	-
2001.....	18	16	17	13	29	87	77	96	83	24	3	-
2000.....	12	19	22	23	51	75	90	110	78	38	15	-
1999.....	21	13	14	32	53	67	82	105	75	36	6	-

Non-Hispanic White, both sexes

2003.....	23	19	19	42	40	75	126	128	80	33	4	-
2002.....	14	22	22	39	50	96	107	131	93	30	3	-
2001.....	16	17	19	22	43	111	99	114	90	32	3	-
2000.....	12	21	23	26	48	91	107	130	98	37	15	-
1999.....	24	20	18	40	62	91	115	140	93	43	5	-

Non-Hispanic White, male

2003.....	9	8	6	12	15	19	39	43	23	6	-	-
2002.....	7	9	7	13	23	33	25	36	24	5	-	-
2001.....	3	6	3	11	16	29	35	26	13	8	-	-
2000.....	5	5	7	6	7	30	32	31	27	4	2	-
1999.....	8	9	8	14	16	32	38	43	22	11	-	-

Non-Hispanic White, female

2003.....	14	11	13	30	25	56	87	85	57	27	4	-
2002.....	7	13	15	26	27	63	82	95	69	25	3	-
2001.....	13	11	16	11	27	82	64	88	77	24	3	-
2000.....	7	16	16	20	41	61	75	99	71	33	13	-
1999.....	16	11	10	26	46	59	77	97	71	32	5	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Non-Hispanic Black, both sexes

2003.....	5	7	7	6	4	9	15	8	8	2	2	-
2002.....	2	4	6	7	1	11	4	11	5	5	-	-
2001.....	6	3	4	8	5	8	14	10	6	-	-	-
2000.....	8	2	6	7	12	16	12	10	8	6	1	-
1999.....	7	4	7	7	13	8	8	9	5	3	1	-

Non-Hispanic Black, male

2003.....	2	2	3	2	1	3	8	-	2	-	-	-
2002.....	-	1	2	6	-	4	1	4	-	1	-	-
2001.....	1	-	3	6	3	3	3	3	1	-	-	-
2000.....	3	-	2	4	4	4	3	-	3	1	-	-
1999.....	3	2	3	3	7	1	3	2	1	-	-	-

Non-Hispanic Black, female

2003.....	3	5	4	4	3	6	7	8	6	2	2	-
2002.....	2	3	4	1	1	7	3	7	5	4	-	-
2001.....	5	3	1	2	2	5	11	7	5	-	-	-
2000.....	5	2	4	3	8	12	9	10	5	5	1	-
1999.....	4	2	4	4	6	7	5	7	4	3	1	-

Origin not stated 2/, both sexes

2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	1	-	1	1	-	-	-	-	-
1999.....	-	-	1	-	-	2	-	-	-	1	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

Origin not stated 2/, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	1	-	-	-	-	-
1999.....	-	-	-	-	-	1	-	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	1	-	-	-	-
2001.....	-	-	-	-	-	-	1	-	-	-	-	-
2000.....	-	-	-	-	-	1	-	-	-	-	-	-
1999.....	-	-	1	-	-	1	-	-	-	1	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Hyperplasia of prostate (N40)

## Hispanic, male

2003.....	-	-	-	1	5	6	5	5	1	1	-	-
2002.....	-	-	1	1	2	3	6	-	1	-	-	-
2001.....	-	1	-	2	2	1	3	1	-	1	-	-
2000.....	-	-	1	2	3	3	-	6	2	2	-	-
1999.....	-	1	-	2	1	-	4	1	1	-	1	-

## Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

## Non-Hispanic 1/, both sexes

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

## Non-Hispanic 1/, male

2003.....	-	1	10	13	29	52	101	107	95	37	2	-
2002.....	-	1	7	7	29	45	110	92	83	30	11	1
2001.....	2	2	6	11	28	57	77	109	94	32	5	-
2000.....	-	4	6	8	25	61	95	100	81	27	4	1
1999.....	-	3	9	13	30	59	97	108	66	24	8	-







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

#### Inflammatory diseases of female pelvic organs (N70-N76)

##### All origins, both sexes

2003.....	7	3	7	6	10	17	12	18	9	4	-	-
2002.....	7	4	2	4	4	17	12	22	11	3	-	-
2001.....	3	4	8	8	10	13	18	23	11	4	-	-
2000.....	3	4	4	11	10	17	12	11	12	3	1	-
1999.....	2	6	7	4	11	15	17	6	6	5	1	-

##### All origins, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

##### All origins, female

2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

##### Hispanic, both sexes

2003.....	-	-	-	1	2	1	-	-	-	1	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	-	-	1	1	-	-	1	-	-	-	-
2000.....	-	1	1	2	-	-	-	-	1	-	-	-
1999.....	1	1	-	-	-	1	-	2	-	-	-	-









































Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

Hispanic, male

2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	1

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	1	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	1	2	-	2	-	-	-	-	2
2001.....	2	2	-	2	1	1	-	1	-	-	-	1
2000.....	1	1	1	-	-	-	-	-	-	-	-	1
1999.....	-	1	2	1	1	1	-	-	1	-	-	3

Non-Hispanic 1/, male

2003.....	1	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	1	-	1	-	-	-	-	2
2001.....	2	1	-	1	1	1	-	-	-	-	-	1
2000.....	1	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	1	-	1	1	-	-	1	-	-	3







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins, both sexes

2003.....	343	376	290	223	190	224	240	170	100	40	9	-
2002.....	363	351	283	224	216	232	257	184	86	33	2	1
2001.....	333	298	283	214	234	244	221	182	87	38	10	-
2000.....	325	296	273	248	216	242	246	178	97	39	6	1
1999.....	305	304	253	251	218	249	242	187	100	34	5	2

All origins, male

2003.....	172	189	147	108	90	86	92	57	27	12	1	-
2002.....	186	170	151	107	106	105	96	65	22	9	1	1
2001.....	167	146	130	109	119	110	93	70	17	9	4	-
2000.....	190	130	134	120	102	99	98	74	41	13	2	-
1999.....	162	157	126	106	103	102	102	60	35	6	1	1

All origins, female

2003.....	171	187	143	115	100	138	148	113	73	28	8	-
2002.....	177	181	132	117	110	127	161	119	64	24	1	-
2001.....	166	152	153	105	115	134	128	112	70	29	6	-
2000.....	135	166	139	128	114	143	148	104	56	26	4	1
1999.....	143	147	127	145	115	147	140	127	65	28	4	1

Hispanic, both sexes

2003.....	20	17	18	9	8	9	4	7	4	2	1	-
2002.....	21	13	12	12	6	12	11	5	2	1	1	-
2001.....	12	10	14	10	12	17	7	6	4	-	-	-
2000.....	16	15	16	14	7	13	11	4	5	3	-	-
1999.....	16	13	14	16	14	7	9	8	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Hispanic, male

2003.....	11	11	8	4	6	4	2	5	-	-	-	-
2002.....	11	7	8	4	2	6	7	3	-	-	1	-
2001.....	6	4	10	5	9	8	5	5	-	-	-	-
2000.....	11	8	9	5	6	9	6	3	1	3	-	-
1999.....	11	7	2	7	10	4	3	2	1	-	-	-

Hispanic, female

2003.....	9	6	10	5	2	5	2	2	4	2	1	-
2002.....	10	6	4	8	4	6	4	2	2	1	-	-
2001.....	6	6	4	5	3	9	2	1	4	-	-	-
2000.....	5	7	7	9	1	4	5	1	4	-	-	-
1999.....	5	6	12	9	4	3	6	6	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	323	357	272	214	181	215	236	163	95	38	8	-
2002.....	338	337	271	211	210	220	245	179	84	31	1	1
2001.....	319	288	269	201	221	227	214	176	83	37	10	-
2000.....	308	279	253	234	209	227	235	174	91	36	6	1
1999.....	288	291	238	232	203	241	232	177	99	34	5	2

Non-Hispanic 1/, male

2003.....	161	178	139	104	83	82	90	52	27	12	1	-
2002.....	173	163	143	102	104	99	89	62	22	9	-	1
2001.....	159	142	120	102	109	102	88	65	17	9	4	-
2000.....	178	121	122	115	96	89	92	71	39	10	2	-
1999.....	151	150	124	98	93	97	99	56	34	6	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic 1/, female

2003.....	162	179	133	110	98	133	146	111	68	26	7	-
2002.....	165	174	128	109	106	121	156	117	62	22	1	-
2001.....	160	146	149	99	112	125	126	111	66	28	6	-
2000.....	130	158	131	119	113	138	143	103	52	26	4	1
1999.....	137	141	114	134	110	144	133	121	65	28	4	1

Non-Hispanic White, both sexes

2003.....	285	317	243	182	160	195	216	153	85	34	8	-
2002.....	292	290	237	189	185	200	220	165	81	28	1	1
2001.....	280	257	246	181	197	209	196	166	77	34	6	-
2000.....	277	238	222	206	184	210	221	164	88	31	5	1
1999.....	249	262	211	207	176	213	214	161	91	34	5	2

Non-Hispanic White, male

2003.....	144	158	121	88	70	74	77	50	24	10	1	-
2002.....	145	139	124	94	89	89	76	58	22	8	-	1
2001.....	141	126	109	96	100	98	76	59	14	9	1	-
2000.....	156	105	112	94	83	84	88	67	38	9	1	-
1999.....	129	136	112	88	81	78	93	50	32	6	1	1

Non-Hispanic White, female

2003.....	141	159	122	94	90	121	139	103	61	24	7	-
2002.....	147	151	113	95	96	111	144	107	59	20	1	-
2001.....	139	131	137	85	97	111	120	107	63	25	5	-
2000.....	121	133	110	112	101	126	133	97	50	22	4	1
1999.....	120	126	99	119	95	135	121	111	59	28	4	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Non-Hispanic Black, both sexes

2003.....	28	34	26	24	16	15	15	8	8	4	-	-
2002.....	37	41	28	20	20	14	18	13	2	3	-	-
2001.....	32	27	21	14	21	17	11	7	5	3	4	-
2000.....	28	32	26	25	18	14	11	7	3	5	1	-
1999.....	32	25	21	21	25	22	16	15	8	-	-	-

Non-Hispanic Black, male

2003.....	10	18	15	14	11	5	11	2	2	2	-	-
2002.....	21	22	18	8	13	7	9	4	-	1	-	-
2001.....	16	13	10	4	8	4	8	4	2	-	3	-
2000.....	19	11	6	18	8	5	4	3	1	1	1	-
1999.....	17	12	10	6	10	14	5	6	2	-	-	-

Non-Hispanic Black, female

2003.....	18	16	11	10	5	10	4	6	6	2	-	-
2002.....	16	19	10	12	7	7	9	9	2	2	-	-
2001.....	16	14	11	10	13	13	3	3	3	3	1	-
2000.....	9	21	20	7	10	9	7	4	2	4	-	-
1999.....	15	13	11	15	15	8	11	9	6	-	-	-

Origin not stated 2/, both sexes

2003.....	-	2	-	-	1	-	-	-	1	-	-	-
2002.....	4	1	-	1	-	-	1	-	-	1	-	-
2001.....	2	-	-	3	1	-	-	-	-	1	-	-
2000.....	1	2	4	-	-	2	-	-	1	-	-	-
1999.....	1	-	1	3	1	1	1	2	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Origin not stated 2/, male

2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	2	-	-	1	-	-	-	-	-	-	-	-
2001.....	2	-	-	2	1	-	-	-	-	-	-	-
2000.....	1	1	3	-	-	1	-	-	1	-	-	-
1999.....	-	-	-	1	-	1	-	2	-	-	-	-

Origin not stated 2/, female

2003.....	-	2	-	-	-	-	-	-	1	-	-	-
2002.....	2	1	-	-	-	-	1	-	-	1	-	-
2001.....	-	-	-	1	-	-	-	-	-	1	-	-
2000.....	-	1	1	-	-	1	-	-	-	-	-	-
1999.....	1	-	1	2	1	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins, both sexes

2003.....	1,296	1,077	990	1,033	1,489	1,989	3,017	3,723	3,702	2,225	915	112
2002.....	1,304	1,042	960	995	1,403	2,103	2,833	3,350	3,406	1,944	854	109
2001.....	1,622	1,192	1,100	1,207	1,567	2,209	2,794	3,239	3,220	1,882	748	115
2000.....	1,415	1,152	1,101	1,353	1,687	2,235	2,794	3,286	3,014	1,798	726	115
1999.....	983	908	952	1,140	1,558	1,945	2,468	2,818	2,799	1,592	589	75

All origins, male

2003.....	829	682	612	618	813	967	1,193	1,148	919	385	118	83
2002.....	828	652	621	563	761	1,010	1,218	1,103	857	379	100	77
2001.....	1,066	758	699	685	812	1,014	1,170	1,017	760	392	105	78
2000.....	966	743	686	832	876	1,100	1,144	1,078	733	331	100	90
1999.....	650	578	603	665	848	949	981	894	699	306	85	56

All origins, female

2003.....	467	395	378	415	676	1,022	1,824	2,575	2,783	1,840	797	29
2002.....	476	390	339	432	642	1,093	1,615	2,247	2,549	1,565	754	32
2001.....	556	434	401	522	755	1,195	1,624	2,222	2,460	1,490	643	37
2000.....	449	409	415	521	811	1,135	1,650	2,208	2,281	1,467	626	25
1999.....	333	330	349	475	710	996	1,487	1,924	2,100	1,286	504	19

Hispanic, both sexes

2003.....	66	42	31	36	39	58	60	83	79	39	13	11
2002.....	79	44	51	36	47	44	68	59	59	30	17	9
2001.....	110	71	56	55	57	51	56	72	50	32	19	12
2000.....	107	72	42	49	58	54	50	61	51	25	17	11
1999.....	29	29	34	36	37	43	50	47	43	21	13	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Hispanic, male

2003.....	45	33	22	20	24	25	24	26	26	10	-	8
2002.....	45	26	33	23	27	24	33	25	19	8	1	8
2001.....	77	50	42	36	35	27	29	22	8	13	3	11
2000.....	82	41	31	35	33	27	25	28	19	4	2	10
1999.....	16	19	23	22	16	28	18	11	15	9	1	4

Hispanic, female

2003.....	21	9	9	16	15	33	36	57	53	29	13	3
2002.....	34	18	18	13	20	20	35	34	40	22	16	1
2001.....	33	21	14	19	22	24	27	50	42	19	16	1
2000.....	25	31	11	14	25	27	25	33	32	21	15	1
1999.....	13	10	11	14	21	15	32	36	28	12	12	1

Non-Hispanic 1/, both sexes

2003.....	1,216	1,023	952	994	1,445	1,925	2,951	3,632	3,616	2,175	901	44
2002.....	1,212	993	897	954	1,353	2,050	2,754	3,282	3,332	1,904	833	43
2001.....	1,499	1,118	1,036	1,149	1,504	2,149	2,737	3,158	3,168	1,843	727	36
2000.....	1,282	1,058	1,051	1,291	1,617	2,171	2,737	3,216	2,960	1,770	708	35
1999.....	936	868	912	1,090	1,511	1,896	2,414	2,765	2,752	1,567	574	30

Non-Hispanic 1/, male

2003.....	772	640	585	595	784	938	1,164	1,116	893	373	117	38
2002.....	770	621	577	535	732	981	1,175	1,074	836	369	97	33
2001.....	982	705	650	648	774	984	1,141	993	750	376	102	26
2000.....	864	683	648	787	834	1,064	1,114	1,047	713	326	97	30
1999.....	618	550	575	638	825	916	960	881	683	296	84	19

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic 1/, female

2003.....	444	383	367	399	661	987	1,787	2,516	2,723	1,802	784	6
2002.....	442	372	320	419	621	1,069	1,579	2,208	2,496	1,535	736	10
2001.....	517	413	386	501	730	1,165	1,596	2,165	2,418	1,467	625	10
2000.....	418	375	403	504	783	1,107	1,623	2,169	2,247	1,444	611	5
1999.....	318	318	337	452	686	980	1,454	1,884	2,069	1,271	490	11

Non-Hispanic White, both sexes

2003.....	913	806	756	807	1,206	1,694	2,673	3,353	3,319	2,016	799	35
2002.....	867	770	723	769	1,165	1,795	2,480	3,012	3,082	1,762	735	31
2001.....	1,168	884	829	921	1,270	1,897	2,453	2,907	2,926	1,693	652	29
2000.....	962	822	830	1,016	1,378	1,903	2,455	2,946	2,748	1,652	629	22
1999.....	686	658	716	878	1,288	1,634	2,141	2,533	2,552	1,442	506	22

Non-Hispanic White, male

2003.....	591	497	461	481	662	833	1,048	1,025	814	343	100	31
2002.....	548	478	467	433	632	844	1,053	997	772	337	74	24
2001.....	777	548	510	517	667	885	1,035	912	686	342	92	22
2000.....	650	520	516	618	710	939	1,006	958	667	303	85	19
1999.....	454	413	457	514	708	793	845	797	625	271	78	16

Non-Hispanic White, female

2003.....	322	309	295	326	544	861	1,625	2,328	2,505	1,673	699	4
2002.....	319	292	256	336	533	951	1,427	2,015	2,310	1,425	661	7
2001.....	391	336	319	404	603	1,012	1,418	1,995	2,240	1,351	560	7
2000.....	312	302	314	398	668	964	1,449	1,988	2,081	1,349	544	3
1999.....	232	245	259	364	580	841	1,296	1,736	1,927	1,171	428	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Non-Hispanic Black, both sexes

2003.....	283	202	183	171	229	203	236	242	256	134	89	7
2002.....	324	201	155	165	170	226	245	235	213	120	87	10
2001.....	290	202	171	197	204	221	245	212	214	131	65	5
2000.....	291	207	204	250	222	247	258	238	184	95	78	13
1999.....	226	192	179	194	201	238	250	206	180	113	64	6

Non-Hispanic Black, male

2003.....	172	130	117	105	118	90	93	80	69	22	14	5
2002.....	209	129	100	89	91	122	112	66	52	28	21	7
2001.....	172	138	124	114	93	87	87	69	54	28	10	3
2000.....	196	146	120	154	113	115	99	75	39	14	11	11
1999.....	147	128	110	113	105	113	101	71	53	22	6	2

Non-Hispanic Black, female

2003.....	111	72	66	66	111	113	143	162	187	112	75	2
2002.....	115	72	55	76	79	104	133	169	161	92	66	3
2001.....	118	64	47	83	111	134	158	143	160	103	55	2
2000.....	95	61	84	96	109	132	159	163	145	81	67	2
1999.....	79	64	69	81	96	125	149	135	127	91	58	4

Origin not stated 2/, both sexes

2003.....	14	12	7	3	5	6	6	8	7	11	1	57
2002.....	13	5	12	5	3	9	11	9	15	10	4	57
2001.....	13	3	8	3	6	9	1	9	2	7	2	67
2000.....	26	22	8	13	12	10	7	9	3	3	1	69
1999.....	18	11	6	14	10	6	4	6	4	4	2	40

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Origin not stated 2/, male

2003.....	12	9	5	3	5	4	5	6	-	2	1	37
2002.....	13	5	11	5	2	5	10	4	2	2	2	36
2001.....	7	3	7	1	3	3	-	2	2	3	-	41
2000.....	20	19	7	10	9	9	5	3	1	1	1	50
1999.....	16	9	5	5	7	5	3	2	1	1	-	33

Origin not stated 2/, female

2003.....	2	3	2	-	-	2	1	2	7	9	-	20
2002.....	-	-	1	-	1	4	1	5	13	8	2	21
2001.....	6	-	1	2	3	6	1	7	-	4	2	26
2000.....	6	3	1	3	3	1	2	6	2	2	-	19
1999.....	2	2	1	9	3	1	1	4	3	3	2	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins, both sexes

2003.....	8,635	8,940	9,794	11,160	15,315	22,635	30,320	32,746	24,949	10,915	2,490	17
2002.....	7,992	8,431	8,947	10,576	15,113	22,098	29,520	31,671	24,416	10,231	2,256	13
2001.....	7,536	7,549	8,499	10,676	15,127	21,941	27,770	30,312	22,714	9,872	2,134	17
2000.....	7,099	7,152	8,004	10,108	14,899	21,425	26,594	28,517	21,457	9,192	2,126	11
1999.....	6,506	6,676	7,714	10,272	14,893	20,657	24,941	27,390	20,562	8,680	1,967	10

All origins, male

2003.....	5,136	5,027	5,247	5,689	7,304	9,807	11,586	10,145	5,900	1,838	295	13
2002.....	4,810	4,675	4,853	5,453	7,214	9,509	11,158	9,835	5,691	1,711	315	11
2001.....	4,427	4,112	4,540	5,480	7,252	9,406	10,411	9,265	5,258	1,740	257	16
2000.....	4,100	3,967	4,269	5,174	7,147	9,237	9,865	8,712	4,966	1,593	265	11
1999.....	3,762	3,762	4,108	5,154	7,178	8,976	9,519	8,208	4,722	1,443	250	5

All origins, female

2003.....	3,499	3,913	4,547	5,471	8,011	12,828	18,734	22,601	19,049	9,077	2,195	4
2002.....	3,182	3,756	4,094	5,123	7,899	12,589	18,362	21,836	18,725	8,520	1,941	2
2001.....	3,109	3,437	3,959	5,196	7,875	12,535	17,359	21,047	17,456	8,132	1,877	1
2000.....	2,999	3,185	3,735	4,934	7,752	12,188	16,729	19,805	16,491	7,599	1,861	-
1999.....	2,744	2,914	3,606	5,118	7,715	11,681	15,422	19,182	15,840	7,237	1,717	5

Hispanic, both sexes

2003.....	608	563	554	594	712	872	950	771	607	269	65	3
2002.....	476	493	465	597	716	796	836	749	543	240	39	1
2001.....	454	444	474	529	689	741	798	781	516	236	52	3
2000.....	422	434	379	505	608	711	727	685	427	215	49	4
1999.....	400	395	447	513	620	627	638	728	494	213	47	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Hispanic, male

2003.....	377	347	293	300	331	377	410	244	170	66	12	2
2002.....	292	283	244	322	337	365	342	258	156	60	9	1
2001.....	289	259	254	271	316	301	311	280	130	67	16	3
2000.....	257	270	194	284	284	300	257	230	128	59	15	4
1999.....	245	230	251	250	314	282	248	227	132	60	9	1

Hispanic, female

2003.....	231	216	261	294	381	495	540	527	437	203	53	1
2002.....	184	210	221	275	379	431	494	491	387	180	30	-
2001.....	165	185	220	258	373	440	487	501	386	169	36	-
2000.....	165	164	185	221	324	411	470	455	299	156	34	-
1999.....	155	165	196	263	306	345	390	501	362	153	38	1

Non-Hispanic 1/, both sexes

2003.....	7,993	8,351	9,207	10,534	14,575	21,714	29,330	31,917	24,288	10,618	2,418	7
2002.....	7,469	7,898	8,449	9,948	14,359	21,240	28,595	30,850	23,814	9,967	2,215	8
2001.....	7,038	7,064	7,999	10,110	14,396	21,150	26,909	29,462	22,146	9,608	2,074	11
2000.....	6,638	6,685	7,598	9,566	14,233	20,649	25,809	27,767	20,978	8,964	2,071	5
1999.....	6,070	6,247	7,232	9,721	14,235	19,977	24,255	26,591	20,022	8,440	1,915	5

Non-Hispanic 1/, male

2003.....	4,734	4,665	4,930	5,368	6,958	9,407	11,160	9,883	5,707	1,769	281	5
2002.....	4,487	4,366	4,587	5,108	6,856	9,121	10,784	9,559	5,524	1,644	306	7
2001.....	4,106	3,822	4,265	5,189	6,915	9,077	10,078	8,966	5,111	1,667	241	10
2000.....	3,813	3,680	4,054	4,868	6,832	8,905	9,588	8,459	4,827	1,533	249	5
1999.....	3,490	3,509	3,832	4,882	6,841	8,672	9,252	7,956	4,579	1,377	241	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic 1/, female

2003.....	3,259	3,686	4,277	5,166	7,617	12,307	18,170	22,034	18,581	8,849	2,137	2
2002.....	2,982	3,532	3,862	4,840	7,503	12,119	17,811	21,291	18,290	8,323	1,909	1
2001.....	2,932	3,242	3,734	4,921	7,481	12,073	16,831	20,496	17,035	7,941	1,833	1
2000.....	2,825	3,005	3,544	4,698	7,401	11,744	16,221	19,308	16,151	7,431	1,822	-
1999.....	2,580	2,738	3,400	4,839	7,394	11,305	15,003	18,635	15,443	7,063	1,674	2

Non-Hispanic White, both sexes

2003.....	5,996	6,633	7,438	8,765	12,459	19,102	26,408	29,385	22,339	9,694	2,089	5
2002.....	5,672	6,294	6,822	8,269	12,261	18,720	25,769	28,310	22,014	9,077	1,924	5
2001.....	5,303	5,553	6,429	8,415	12,392	18,647	24,343	26,949	20,435	8,830	1,786	6
2000.....	5,079	5,241	6,091	7,954	12,234	18,262	23,300	25,412	19,296	8,212	1,809	2
1999.....	4,579	4,912	5,837	8,027	12,240	17,624	22,039	24,396	18,595	7,727	1,688	1

Non-Hispanic White, male

2003.....	3,608	3,740	4,002	4,465	5,951	8,270	10,004	9,111	5,261	1,589	230	3
2002.....	3,479	3,499	3,731	4,242	5,867	8,035	9,713	8,765	5,056	1,461	249	5
2001.....	3,146	3,007	3,469	4,333	5,979	7,992	9,148	8,203	4,674	1,494	192	5
2000.....	2,969	2,918	3,231	4,051	5,882	7,874	8,645	7,710	4,375	1,382	210	2
1999.....	2,685	2,786	3,120	4,054	5,923	7,621	8,392	7,226	4,177	1,221	206	1

Non-Hispanic White, female

2003.....	2,388	2,893	3,436	4,300	6,508	10,832	16,404	20,274	17,078	8,105	1,859	2
2002.....	2,193	2,795	3,091	4,027	6,394	10,685	16,056	19,545	16,958	7,616	1,675	-
2001.....	2,157	2,546	2,960	4,082	6,413	10,655	15,195	18,746	15,761	7,336	1,594	1
2000.....	2,110	2,323	2,860	3,903	6,352	10,388	14,655	17,702	14,921	6,830	1,599	-
1999.....	1,894	2,126	2,717	3,973	6,317	10,003	13,647	17,170	14,418	6,506	1,482	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Non-Hispanic Black, both sexes

2003.....	1,787	1,509	1,539	1,528	1,807	2,227	2,455	2,130	1,663	795	292	2
2002.....	1,619	1,416	1,411	1,450	1,797	2,128	2,431	2,161	1,578	778	255	3
2001.....	1,577	1,341	1,350	1,498	1,738	2,170	2,245	2,160	1,480	684	262	5
2000.....	1,402	1,278	1,329	1,412	1,741	2,075	2,173	2,054	1,466	656	237	3
1999.....	1,329	1,195	1,242	1,498	1,753	2,053	1,910	1,903	1,273	633	197	4

Non-Hispanic Black, male

2003.....	1,002	801	809	770	873	958	957	627	355	145	40	2
2002.....	899	771	730	735	856	898	897	646	389	138	44	2
2001.....	875	722	689	767	803	948	819	622	346	141	42	5
2000.....	758	689	722	717	831	897	814	618	360	130	34	3
1999.....	722	646	624	725	793	909	721	608	335	130	24	2

Non-Hispanic Black, female

2003.....	785	708	730	758	934	1,269	1,498	1,503	1,308	650	252	-
2002.....	720	645	681	715	941	1,230	1,534	1,515	1,189	640	211	1
2001.....	702	619	661	731	935	1,222	1,426	1,538	1,134	543	220	-
2000.....	644	589	607	695	910	1,178	1,359	1,436	1,106	526	203	-
1999.....	607	549	618	773	960	1,144	1,189	1,295	938	503	173	2

Origin not stated 2/, both sexes

2003.....	34	26	33	32	28	49	40	58	54	28	7	7
2002.....	47	40	33	31	38	62	89	72	59	24	2	4
2001.....	44	41	26	37	42	50	63	69	52	28	8	3
2000.....	39	33	27	37	58	65	58	65	52	13	6	2
1999.....	36	34	35	38	38	53	48	71	46	27	5	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Origin not stated 2/, male

2003.....	25	15	24	21	15	23	16	18	23	3	2	6
2002.....	31	26	22	23	21	23	32	18	11	7	-	3
2001.....	32	31	21	20	21	28	22	19	17	6	-	3
2000.....	30	17	21	22	31	32	20	23	11	1	1	2
1999.....	27	23	25	22	23	22	19	25	11	6	-	1

Origin not stated 2/, female

2003.....	9	11	9	11	13	26	24	40	31	25	5	1
2002.....	16	14	11	8	17	39	57	54	48	17	2	1
2001.....	12	10	5	17	21	22	41	50	35	22	8	-
2000.....	9	16	6	15	27	33	38	42	41	12	5	-
1999.....	9	11	10	16	15	31	29	46	35	21	5	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins, both sexes

2003.....	6,972	5,186	3,984	3,711	4,370	5,875	7,233	6,839	4,352	1,620	335	76
2002.....	6,434	4,557	3,788	3,672	4,414	5,938	6,966	6,507	4,255	1,595	294	85
2001.....	5,875	4,159	3,499	3,506	4,329	5,972	6,716	6,266	4,074	1,519	312	120
2000.....	5,408	4,021	3,484	3,375	4,323	5,596	6,162	5,933	3,939	1,419	304	85
1999.....	4,943	3,944	3,341	3,692	4,516	5,772	6,510	6,138	3,954	1,380	257	106

All origins, male

2003.....	4,871	3,613	2,711	2,365	2,556	3,251	3,600	2,991	1,595	440	67	70
2002.....	4,552	3,255	2,526	2,369	2,662	3,268	3,419	2,795	1,488	480	54	74
2001.....	4,224	2,872	2,356	2,200	2,551	3,310	3,342	2,730	1,486	381	60	104
2000.....	3,960	2,804	2,329	2,117	2,522	3,039	3,024	2,563	1,391	397	62	83
1999.....	3,606	2,753	2,262	2,336	2,656	3,096	3,276	2,647	1,397	376	60	94

All origins, female

2003.....	2,101	1,573	1,273	1,346	1,814	2,624	3,633	3,848	2,757	1,180	268	6
2002.....	1,882	1,302	1,262	1,303	1,752	2,670	3,547	3,712	2,767	1,115	240	11
2001.....	1,651	1,287	1,143	1,306	1,778	2,662	3,374	3,536	2,588	1,138	252	16
2000.....	1,448	1,217	1,155	1,258	1,801	2,557	3,138	3,370	2,548	1,022	242	2
1999.....	1,337	1,191	1,079	1,356	1,860	2,676	3,234	3,491	2,557	1,004	197	12

Hispanic, both sexes

2003.....	519	338	301	239	227	234	220	157	92	37	7	15
2002.....	483	311	255	228	238	262	202	130	92	47	7	28
2001.....	451	313	251	205	249	222	187	140	84	36	6	29
2000.....	411	272	220	187	231	193	172	146	94	30	8	28
1999.....	380	250	216	245	195	183	151	140	83	28	9	44

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Hispanic, male

2003.....	381	246	203	153	136	146	121	77	37	11	2	15
2002.....	366	235	172	166	141	152	106	66	43	21	4	27
2001.....	341	213	178	130	145	136	99	59	35	8	2	26
2000.....	315	193	154	126	145	117	74	69	42	11	3	28
1999.....	291	183	151	162	126	100	82	67	37	15	1	40

Hispanic, female

2003.....	138	92	98	86	91	88	99	80	55	26	5	-
2002.....	117	76	83	62	97	110	96	64	49	26	3	1
2001.....	110	100	73	75	104	86	88	81	49	28	4	3
2000.....	96	79	66	61	86	76	98	77	52	19	5	-
1999.....	89	67	65	83	69	83	69	73	46	13	8	4

Non-Hispanic 1/, both sexes

2003.....	6,406	4,821	3,664	3,454	4,126	5,624	6,996	6,663	4,252	1,581	328	23
2002.....	5,913	4,214	3,514	3,428	4,156	5,655	6,739	6,362	4,154	1,544	287	33
2001.....	5,391	3,816	3,226	3,286	4,061	5,738	6,507	6,114	3,980	1,475	305	36
2000.....	4,945	3,723	3,243	3,175	4,072	5,386	5,975	5,771	3,836	1,386	295	12
1999.....	4,523	3,652	3,101	3,430	4,308	5,567	6,329	5,977	3,859	1,348	248	26

Non-Hispanic 1/, male

2003.....	4,451	3,345	2,494	2,197	2,407	3,092	3,472	2,905	1,554	429	65	20
2002.....	4,157	2,988	2,337	2,189	2,504	3,105	3,293	2,720	1,444	457	50	30
2001.....	3,853	2,636	2,159	2,062	2,393	3,166	3,231	2,667	1,447	372	58	29
2000.....	3,605	2,590	2,157	1,979	2,365	2,913	2,942	2,487	1,344	385	59	11
1999.....	3,281	2,537	2,090	2,161	2,520	2,984	3,174	2,571	1,354	359	59	23

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic 1/, female

2003.....	1,955	1,476	1,170	1,257	1,719	2,532	3,524	3,758	2,698	1,152	263	3
2002.....	1,756	1,226	1,177	1,239	1,652	2,550	3,446	3,642	2,710	1,087	237	3
2001.....	1,538	1,180	1,067	1,224	1,668	2,572	3,276	3,447	2,533	1,103	247	7
2000.....	1,340	1,133	1,086	1,196	1,707	2,473	3,033	3,284	2,492	1,001	236	1
1999.....	1,242	1,115	1,011	1,269	1,788	2,583	3,155	3,406	2,505	989	189	3

Non-Hispanic White, both sexes

2003.....	5,128	3,978	3,065	2,874	3,592	5,065	6,422	6,282	3,991	1,505	300	22
2002.....	4,779	3,459	2,934	2,840	3,641	5,082	6,234	5,977	3,894	1,448	260	26
2001.....	4,343	3,132	2,644	2,784	3,525	5,160	5,997	5,711	3,729	1,376	276	25
2000.....	3,904	3,044	2,709	2,659	3,522	4,834	5,522	5,410	3,594	1,284	264	8
1999.....	3,603	2,968	2,487	2,875	3,693	4,946	5,811	5,537	3,625	1,260	225	13

Non-Hispanic White, male

2003.....	3,576	2,752	2,089	1,833	2,119	2,784	3,173	2,735	1,445	405	58	19
2002.....	3,333	2,427	1,947	1,795	2,196	2,778	3,027	2,544	1,332	425	41	23
2001.....	3,093	2,142	1,763	1,748	2,089	2,838	2,962	2,482	1,348	339	51	22
2000.....	2,841	2,089	1,782	1,636	2,056	2,589	2,703	2,326	1,246	345	51	7
1999.....	2,611	2,040	1,690	1,805	2,157	2,632	2,919	2,368	1,251	334	53	12

Non-Hispanic White, female

2003.....	1,552	1,226	976	1,041	1,473	2,281	3,249	3,547	2,546	1,100	242	3
2002.....	1,446	1,032	987	1,045	1,445	2,304	3,207	3,433	2,562	1,023	219	3
2001.....	1,250	990	881	1,036	1,436	2,322	3,035	3,229	2,381	1,037	225	3
2000.....	1,063	955	927	1,023	1,466	2,245	2,819	3,084	2,348	939	213	1
1999.....	992	928	797	1,070	1,536	2,314	2,892	3,169	2,374	926	172	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Non-Hispanic Black, both sexes

2003.....	1,042	683	469	451	399	393	405	276	189	59	20	1
2002.....	934	618	457	435	383	433	366	289	195	76	20	4
2001.....	891	568	477	415	421	447	383	295	196	73	25	8
2000.....	865	561	422	414	443	419	349	279	189	80	28	4
1999.....	765	562	485	454	489	499	426	349	197	81	19	12

Non-Hispanic Black, male

2003.....	729	493	331	298	220	227	202	121	71	17	5	1
2002.....	687	470	321	295	232	254	194	129	81	25	5	4
2001.....	656	416	334	268	237	256	205	122	73	21	6	6
2000.....	646	429	307	282	255	250	182	120	77	32	7	4
1999.....	577	418	324	297	299	282	201	162	79	22	5	10

Non-Hispanic Black, female

2003.....	313	190	138	153	179	166	203	155	118	42	15	-
2002.....	247	148	136	140	151	179	172	160	114	51	15	-
2001.....	235	152	143	147	184	191	178	173	123	52	19	2
2000.....	219	132	115	132	188	169	167	159	112	48	21	-
1999.....	188	144	161	157	190	217	225	187	118	59	14	2

Origin not stated 2/, both sexes

2003.....	47	27	19	18	17	17	17	19	8	2	-	38
2002.....	38	32	19	16	20	21	25	15	9	4	-	24
2001.....	33	30	22	15	19	12	22	12	10	8	1	55
2000.....	52	26	21	13	20	17	15	16	9	3	1	45
1999.....	40	42	24	17	13	22	30	21	12	4	-	36

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Origin not stated 2/, male

2003.....	39	22	14	15	13	13	7	9	4	-	-	35
2002.....	29	32	17	14	17	11	20	9	1	2	-	17
2001.....	30	23	19	8	13	8	12	4	4	1	-	49
2000.....	40	21	18	12	12	9	8	7	5	1	-	44
1999.....	34	33	21	13	10	12	20	9	6	2	-	31

Origin not stated 2/, female

2003.....	8	5	5	3	4	4	10	10	4	2	-	3
2002.....	9	-	2	2	3	10	5	6	8	2	-	7
2001.....	3	7	3	7	6	4	10	8	6	7	1	6
2000.....	12	5	3	1	8	8	7	9	4	2	1	1
1999.....	6	9	3	4	3	10	10	12	6	2	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

All origins, both sexes

2003.....	3,098	2,507	1,934	1,578	1,632	1,782	1,595	994	350	55	2	20
2002.....	2,970	2,229	1,936	1,597	1,754	1,855	1,578	936	330	47	9	39
2001.....	2,872	2,152	1,709	1,567	1,715	1,943	1,564	903	304	63	2	43
2000.....	2,717	2,115	1,768	1,599	1,716	1,860	1,482	985	298	49	3	33
1999.....	2,524	2,131	1,681	1,727	1,852	1,902	1,566	964	311	50	5	37

All origins, male

2003.....	2,198	1,778	1,309	1,019	956	1,019	866	534	187	30	-	18
2002.....	2,089	1,615	1,262	1,039	1,062	1,060	857	527	180	26	4	36
2001.....	2,032	1,467	1,164	974	1,041	1,106	877	492	163	28	1	37
2000.....	1,922	1,456	1,161	996	1,025	1,050	808	539	167	27	1	33
1999.....	1,782	1,454	1,107	1,091	1,071	1,089	870	551	179	25	3	34

All origins, female

2003.....	900	729	625	559	676	763	729	460	163	25	2	2
2002.....	881	614	674	558	692	795	721	409	150	21	5	3
2001.....	840	685	545	593	674	837	687	411	141	35	1	6
2000.....	795	659	607	603	691	810	674	446	131	22	2	-
1999.....	742	677	574	636	781	813	696	413	132	25	2	3

Hispanic, both sexes

2003.....	247	177	152	121	106	83	68	27	16	5	-	8
2002.....	237	155	145	109	108	103	54	28	12	5	1	17
2001.....	244	180	151	114	115	99	68	27	9	2	-	10
2000.....	227	153	115	98	122	76	45	26	10	2	-	14
1999.....	199	148	105	132	96	76	44	26	9	1	-	11

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

Hispanic, male

2003.....	176	119	93	78	64	50	42	15	11	4	-	8
2002.....	163	114	93	82	67	63	28	15	8	3	1	16
2001.....	174	107	107	73	68	67	47	16	6	1	-	8
2000.....	169	107	82	63	77	53	21	13	7	2	-	14
1999.....	146	94	67	85	59	46	31	13	5	1	-	11

Hispanic, female

2003.....	71	58	59	43	42	33	26	12	5	1	-	-
2002.....	74	41	52	27	41	40	26	13	4	2	-	1
2001.....	70	73	44	41	47	32	21	11	3	1	-	2
2000.....	58	46	33	35	45	23	24	13	3	-	-	-
1999.....	53	54	38	47	37	30	13	13	4	-	-	-

Non-Hispanic 1/, both sexes

2003.....	2,837	2,318	1,772	1,449	1,517	1,696	1,521	962	334	50	2	4
2002.....	2,721	2,063	1,781	1,482	1,638	1,750	1,514	906	318	42	8	15
2001.....	2,614	1,964	1,549	1,452	1,592	1,841	1,489	874	294	60	2	15
2000.....	2,473	1,952	1,643	1,495	1,587	1,782	1,432	957	288	47	3	3
1999.....	2,316	1,967	1,568	1,589	1,749	1,820	1,509	936	301	49	5	11

Non-Hispanic 1/, male

2003.....	2,010	1,650	1,208	934	886	967	822	516	176	26	-	4
2002.....	1,917	1,490	1,161	952	988	995	821	510	172	23	3	14
2001.....	1,845	1,356	1,049	901	968	1,038	827	475	157	26	1	12
2000.....	1,739	1,341	1,070	928	943	997	782	525	160	25	1	3
1999.....	1,629	1,346	1,033	1,002	1,007	1,040	829	537	174	24	3	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

Non-Hispanic 1/, female

2003.....	827	668	564	515	631	729	699	446	158	24	2	-
2002.....	804	573	620	530	650	755	693	396	146	19	5	1
2001.....	769	608	500	551	624	803	662	399	137	34	1	3
2000.....	734	611	573	567	644	785	650	432	128	22	2	-
1999.....	687	621	535	587	742	780	680	399	127	25	2	1

Non-Hispanic White, both sexes

2003.....	2,311	1,927	1,487	1,212	1,331	1,501	1,391	893	306	46	1	4
2002.....	2,250	1,738	1,508	1,246	1,449	1,580	1,394	840	298	39	5	10
2001.....	2,180	1,650	1,287	1,235	1,393	1,668	1,376	814	277	55	2	8
2000.....	2,008	1,636	1,403	1,272	1,377	1,607	1,330	887	265	42	3	3
1999.....	1,904	1,643	1,303	1,361	1,536	1,636	1,384	860	272	44	4	4

Non-Hispanic White, male

2003.....	1,652	1,381	1,030	790	794	856	750	478	157	23	-	4
2002.....	1,586	1,250	990	792	869	884	746	467	156	22	3	9
2001.....	1,554	1,136	873	765	852	931	767	436	149	25	1	7
2000.....	1,422	1,113	901	778	819	890	719	490	147	23	1	3
1999.....	1,348	1,127	859	863	882	929	757	489	158	19	2	4

Non-Hispanic White, female

2003.....	659	546	457	422	537	645	641	415	149	23	1	-
2002.....	664	488	518	454	580	696	648	373	142	17	2	1
2001.....	626	514	414	470	541	737	609	378	128	30	1	1
2000.....	586	523	502	494	558	717	611	397	118	19	2	-
1999.....	556	516	444	498	654	707	627	371	114	25	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

Non-Hispanic Black, both sexes

2003.....	390	295	206	174	129	133	84	48	22	2	-	-
2002.....	341	249	195	158	126	117	89	44	13	2	3	3
2001.....	335	249	206	171	136	126	72	38	14	5	-	7
2000.....	356	242	175	164	158	107	72	50	16	3	-	-
1999.....	315	244	186	161	153	136	96	59	21	5	1	7

Non-Hispanic Black, male

2003.....	282	213	141	114	69	82	50	27	13	1	-	-
2002.....	248	193	134	111	85	83	56	32	10	1	-	3
2001.....	234	182	150	111	74	84	39	23	7	1	-	5
2000.....	253	186	128	119	99	72	46	27	10	2	-	-
1999.....	230	171	132	103	93	84	55	41	11	5	1	6

Non-Hispanic Black, female

2003.....	108	82	65	60	60	51	34	21	9	1	-	-
2002.....	93	56	61	47	41	34	33	12	3	1	3	-
2001.....	101	67	56	60	62	42	33	15	7	4	-	2
2000.....	103	56	47	45	59	35	26	23	6	1	-	-
1999.....	85	73	54	58	60	52	41	18	10	-	-	1

Origin not stated 2/, both sexes

2003.....	14	12	10	8	9	3	6	5	-	-	-	8
2002.....	12	11	10	6	8	2	10	2	-	-	-	7
2001.....	14	8	9	1	8	3	7	2	1	1	-	18
2000.....	17	10	10	6	7	2	5	2	-	-	-	16
1999.....	9	16	8	6	7	6	13	2	1	-	-	15

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Transport accidents (V01-V99,Y85)

Origin not stated 2/, male

2003.....	12	9	8	7	6	2	2	3	-	-	-	6
2002.....	9	11	8	5	7	2	8	2	-	-	-	6
2001.....	13	4	8	-	5	1	3	1	-	1	-	17
2000.....	14	8	9	5	5	-	5	1	-	-	-	16
1999.....	7	14	7	4	5	3	10	1	-	-	-	13

Origin not stated 2/, female

2003.....	2	3	2	1	3	1	4	2	-	-	-	2
2002.....	3	-	2	1	1	-	2	-	-	-	-	1
2001.....	1	4	1	1	3	2	4	1	1	-	-	1
2000.....	3	2	1	1	2	2	-	1	-	-	-	-
1999.....	2	2	1	2	2	3	3	1	1	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
All origins, both sexes												
2003.....	2,788	2,232	1,726	1,444	1,523	1,680	1,530	961	343	53	1	16
2002.....	2,697	2,004	1,760	1,469	1,644	1,763	1,508	909	321	47	8	35
2001.....	2,553	1,939	1,526	1,409	1,581	1,823	1,489	868	294	59	2	33
2000.....	2,425	1,917	1,589	1,436	1,602	1,764	1,409	953	285	47	3	26
1999.....	2,225	1,885	1,485	1,565	1,711	1,772	1,471	920	291	46	4	30
All origins, male												
2003.....	1,937	1,553	1,130	908	870	944	823	511	182	29	-	14
2002.....	1,861	1,423	1,117	923	965	992	801	512	176	26	4	33
2001.....	1,774	1,296	1,013	846	940	1,027	832	478	157	28	1	27
2000.....	1,689	1,293	1,013	870	932	980	759	517	161	27	1	26
1999.....	1,537	1,250	950	969	962	1,012	815	524	164	24	2	27
All origins, female												
2003.....	851	679	596	536	653	736	707	450	161	24	1	2
2002.....	836	581	643	546	679	771	707	397	145	21	4	2
2001.....	779	643	513	563	641	796	657	390	137	31	1	6
2000.....	736	624	576	566	670	784	650	436	124	20	2	-
1999.....	688	635	535	596	749	760	656	396	127	22	2	3
Hispanic, both sexes												
2003.....	234	163	141	112	101	81	67	26	16	5	-	7
2002.....	215	144	139	106	105	96	50	28	11	5	1	15
2001.....	204	152	127	96	105	83	63	22	8	2	-	8
2000.....	210	141	105	85	113	71	42	22	10	2	-	14
1999.....	181	141	97	119	85	72	41	23	9	1	-	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Hispanic, male												
2003.....	165	110	86	72	62	48	41	15	11	4	-	7
2002.....	145	105	87	79	65	59	25	15	7	3	1	15
2001.....	148	93	91	61	62	56	43	13	5	1	-	6
2000.....	155	98	72	54	69	50	18	10	7	2	-	14
1999.....	129	87	59	75	49	42	29	11	5	1	-	10
Hispanic, female												
2003.....	69	53	55	40	39	33	26	11	5	1	-	-
2002.....	70	39	52	27	40	37	25	13	4	2	-	-
2001.....	56	59	36	35	43	27	20	9	3	1	-	2
2000.....	55	43	33	31	44	21	24	12	3	-	-	-
1999.....	52	54	38	44	36	30	12	12	4	-	-	-
Non-Hispanic 1/, both sexes												
2003.....	2,541	2,059	1,577	1,325	1,413	1,596	1,457	930	327	48	1	1
2002.....	2,473	1,851	1,613	1,358	1,532	1,665	1,449	879	310	42	7	14
2001.....	2,337	1,781	1,392	1,312	1,468	1,737	1,419	844	285	56	2	11
2000.....	2,201	1,767	1,476	1,346	1,483	1,691	1,363	929	275	45	3	3
1999.....	2,037	1,728	1,382	1,440	1,619	1,694	1,418	895	281	45	4	9
Non-Hispanic 1/, male												
2003.....	1,761	1,436	1,037	830	802	894	780	493	171	25	-	1
2002.....	1,710	1,309	1,024	840	894	931	769	495	169	23	3	13
2001.....	1,615	1,200	916	785	873	970	786	464	152	26	1	8
2000.....	1,523	1,188	934	812	859	930	737	506	154	25	1	3
1999.....	1,403	1,149	885	890	908	967	777	512	159	23	2	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)												
Non-Hispanic 1/, female												
2003.....	780	623	540	495	611	702	677	437	156	23	1	-
2002.....	763	542	589	518	638	734	680	384	141	19	4	1
2001.....	722	581	476	527	595	767	633	380	133	30	1	3
2000.....	678	579	542	534	624	761	626	423	121	20	2	-
1999.....	634	579	497	550	711	727	641	383	122	22	2	1
Non-Hispanic White, both sexes												
2003.....	2,056	1,694	1,314	1,109	1,234	1,414	1,334	862	300	44	-	1
2002.....	2,035	1,559	1,359	1,142	1,354	1,508	1,336	816	290	39	4	9
2001.....	1,942	1,500	1,152	1,120	1,278	1,578	1,313	789	270	52	2	8
2000.....	1,770	1,475	1,253	1,145	1,286	1,527	1,264	862	252	41	3	3
1999.....	1,667	1,434	1,145	1,233	1,422	1,527	1,305	824	254	40	4	3
Non-Hispanic White, male												
2003.....	1,439	1,190	877	704	715	791	714	456	153	22	-	1
2002.....	1,409	1,100	868	698	786	833	699	454	153	22	3	8
2001.....	1,355	1,011	753	671	765	870	730	427	145	25	1	7
2000.....	1,231	982	777	679	745	830	675	472	141	23	1	3
1999.....	1,155	954	735	764	797	863	712	468	145	18	2	3
Non-Hispanic White, female												
2003.....	617	504	437	405	519	623	620	406	147	22	-	-
2002.....	626	459	491	444	568	675	637	362	137	17	1	1
2001.....	587	489	399	449	513	708	583	362	125	27	1	1
2000.....	539	493	476	466	541	697	589	390	111	18	2	-
1999.....	512	480	410	469	625	664	593	356	109	22	2	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Non-Hispanic Black, both sexes

2003.....	360	274	189	160	125	124	79	48	21	2	-	-
2002.....	315	222	182	145	118	107	84	43	13	2	3	3
2001.....	306	223	187	152	131	115	69	37	13	4	-	3
2000.....	338	223	165	149	149	100	69	47	16	2	-	-
1999.....	282	222	165	148	143	126	88	54	20	5	-	6

Non-Hispanic Black, male

2003.....	254	194	125	101	66	77	46	27	12	1	-	-
2002.....	224	168	122	98	77	73	52	31	10	1	-	3
2001.....	212	157	138	94	69	78	38	23	7	1	-	1
2000.....	238	169	120	106	93	67	45	26	10	2	-	-
1999.....	203	153	114	94	84	81	51	37	10	5	-	5

Non-Hispanic Black, female

2003.....	106	80	64	59	59	47	33	21	9	1	-	-
2002.....	91	54	60	47	41	34	32	12	3	1	3	-
2001.....	94	66	49	58	62	37	31	14	6	3	-	2
2000.....	100	54	45	43	56	33	24	21	6	-	-	-
1999.....	79	69	51	54	59	45	37	17	10	-	-	1

Origin not stated 2/, both sexes

2003.....	13	10	8	7	9	3	6	5	-	-	-	8
2002.....	9	9	8	5	7	2	9	2	-	-	-	6
2001.....	12	6	7	1	8	3	7	2	1	1	-	14
2000.....	14	9	8	5	6	2	4	2	-	-	-	9
1999.....	7	16	6	6	7	6	12	2	1	-	-	11

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Origin not stated 2/, male

2003.....	11	7	7	6	6	2	2	3	-	-	-	6
2002.....	6	9	6	4	6	2	7	2	-	-	-	5
2001.....	11	3	6	-	5	1	3	1	-	1	-	13
2000.....	11	7	7	4	4	-	4	1	-	-	-	9
1999.....	5	14	6	4	5	3	9	1	-	-	-	9

Origin not stated 2/, female

2003.....	2	3	1	1	3	1	4	2	-	-	-	2
2002.....	3	-	2	1	1	-	2	-	-	-	-	1
2001.....	1	3	1	1	3	2	4	1	1	-	-	1
2000.....	3	2	1	1	2	2	-	1	-	-	-	-
1999.....	2	2	-	2	2	3	3	1	1	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All origins, both sexes

2003.....	97	87	60	42	39	42	38	18	5	1	1	4
2002.....	90	61	48	33	37	35	31	11	5	-	1	4
2001.....	105	67	66	64	53	59	39	20	7	2	-	10
2000.....	109	64	58	52	48	51	40	22	12	1	-	7
1999.....	125	101	81	79	66	92	67	34	16	3	1	7

All origins, male

2003.....	78	68	45	35	28	25	27	11	4	1	-	4
2002.....	78	53	39	26	30	25	25	6	2	-	-	3
2001.....	84	56	48	50	34	38	18	6	4	-	-	10
2000.....	74	48	50	38	37	33	24	14	5	-	-	7
1999.....	99	77	58	52	45	51	37	20	13	1	1	7

All origins, female

2003.....	19	19	15	7	11	17	11	7	1	-	1	-
2002.....	12	8	9	7	7	10	6	5	3	-	1	1
2001.....	21	11	18	14	19	21	21	14	3	2	-	-
2000.....	35	16	8	14	11	18	16	8	7	1	-	-
1999.....	26	24	23	27	21	41	30	14	3	2	-	-

Hispanic, both sexes

2003.....	5	6	7	4	3	-	1	1	-	-	-	1
2002.....	10	4	3	1	1	5	3	-	-	-	-	2
2001.....	10	7	7	3	2	11	2	2	1	-	-	2
2000.....	10	10	4	9	5	4	1	4	-	-	-	-
1999.....	15	2	6	13	8	4	3	2	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Hispanic, male

2003.....	4	6	3	3	1	-	1	-	-	-	-	1
2002.....	7	3	3	1	1	2	3	-	-	-	-	1
2001.....	9	6	6	2	2	9	1	1	1	-	-	2
2000.....	8	7	4	6	4	2	1	3	-	-	-	-
1999.....	14	2	6	10	7	4	2	1	-	-	-	1

Hispanic, female

2003.....	1	-	4	1	2	-	-	1	-	-	-	-
2002.....	3	1	-	-	-	3	-	-	-	-	-	1
2001.....	1	1	1	1	-	2	1	1	-	-	-	-
2000.....	2	3	-	3	1	2	-	1	-	-	-	-
1999.....	1	-	-	3	1	-	1	1	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	91	80	53	37	36	42	37	17	5	1	1	3
2002.....	77	55	45	32	36	30	27	11	5	-	1	1
2001.....	93	58	57	61	51	48	37	18	6	2	-	4
2000.....	97	53	52	42	43	47	38	18	12	1	-	-
1999.....	109	99	73	66	58	88	63	32	16	3	1	2

Non-Hispanic 1/, male

2003.....	73	61	42	31	27	25	26	11	4	1	-	3
2002.....	68	48	36	25	29	23	21	6	2	-	-	1
2001.....	73	49	40	48	32	29	17	5	3	-	-	4
2000.....	64	40	44	31	33	31	22	11	5	-	-	-
1999.....	84	75	51	42	38	47	34	19	13	1	1	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic 1/, female

2003.....	18	19	11	6	9	17	11	6	1	-	1	-
2002.....	9	7	9	7	7	7	6	5	3	-	1	-
2001.....	20	9	17	13	19	19	20	13	3	2	-	-
2000.....	33	13	8	11	10	16	16	7	7	1	-	-
1999.....	25	24	22	24	20	41	29	13	3	2	-	-

Non-Hispanic White, both sexes

2003.....	71	67	45	30	31	37	32	16	4	1	1	3
2002.....	64	39	35	24	31	23	25	9	5	-	1	1
2001.....	77	47	42	45	47	41	32	15	4	2	-	-
2000.....	81	39	45	33	37	40	35	16	12	-	-	-
1999.....	89	88	63	53	49	75	56	29	14	3	-	1

Non-Hispanic White, male

2003.....	57	50	35	25	23	22	22	10	3	1	-	3
2002.....	56	34	29	19	24	16	20	5	2	-	-	1
2001.....	62	39	34	35	28	26	14	3	2	-	-	-
2000.....	53	28	40	23	28	28	21	11	5	-	-	-
1999.....	68	67	43	36	31	41	30	17	11	1	-	1

Non-Hispanic White, female

2003.....	14	17	10	5	8	15	10	6	1	-	1	-
2002.....	8	5	6	5	7	7	5	4	3	-	1	-
2001.....	15	8	8	10	19	15	18	12	2	2	-	-
2000.....	28	11	5	10	9	12	14	5	7	-	-	-
1999.....	21	21	20	17	18	34	26	12	3	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

Non-Hispanic Black, both sexes

2003.....	17	12	7	5	3	4	3	-	1	-	-	-
2002.....	12	12	6	5	4	5	2	-	-	-	-	-
2001.....	12	9	13	12	3	5	1	-	1	-	-	4
2000.....	10	11	4	7	3	4	3	2	-	1	-	-
1999.....	17	11	6	8	6	8	3	3	1	-	1	1

Non-Hispanic Black, male

2003.....	15	10	7	5	3	2	2	-	1	-	-	-
2002.....	11	10	6	5	4	5	1	-	-	-	-	-
2001.....	8	8	6	10	3	3	-	-	-	-	-	4
2000.....	7	9	3	7	2	2	1	-	-	-	-	-
1999.....	14	8	4	4	5	2	1	2	1	-	1	1

Non-Hispanic Black, female

2003.....	2	2	-	-	-	2	1	-	-	-	-	-
2002.....	1	2	-	-	-	-	1	-	-	-	-	-
2001.....	4	1	7	2	-	2	1	-	1	-	-	-
2000.....	3	2	1	-	1	2	2	2	-	1	-	-
1999.....	3	3	2	4	1	6	2	1	-	-	-	-

Origin not stated 2/, both sexes

2003.....	1	1	-	1	-	-	-	-	-	-	-	-
2002.....	3	2	-	-	-	-	1	-	-	-	-	1
2001.....	2	2	2	-	-	-	-	-	-	-	-	4
2000.....	2	1	2	1	-	-	1	-	-	-	-	7
1999.....	1	-	2	-	-	-	1	-	-	-	-	4



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All origins, both sexes

2003.....	213	188	148	92	70	60	27	15	2	1	-	-
2002.....	183	164	128	95	73	57	39	16	4	-	-	-
2001.....	214	146	117	94	81	61	36	15	3	2	-	-
2000.....	183	134	121	111	66	45	33	10	1	1	-	-
1999.....	174	145	115	83	75	38	28	10	4	1	-	-

All origins, male

2003.....	183	157	134	76	58	50	16	12	1	-	-	-
2002.....	150	139	106	90	67	43	31	9	2	-	-	-
2001.....	174	115	103	78	67	41	27	8	2	-	-	-
2000.....	159	115	98	88	56	37	25	8	1	-	-	-
1999.....	146	127	99	70	64	26	18	7	2	-	-	-

All origins, female

2003.....	30	31	14	16	12	10	11	3	1	1	-	-
2002.....	33	25	22	5	6	14	8	7	2	-	-	-
2001.....	40	31	14	16	14	20	9	7	1	2	-	-
2000.....	24	19	23	23	10	8	8	2	-	1	-	-
1999.....	28	18	16	13	11	12	10	3	2	1	-	-

Hispanic, both sexes

2003.....	8	8	4	5	2	2	-	-	-	-	-	-
2002.....	12	7	3	2	2	2	1	-	1	-	-	-
2001.....	30	21	17	15	8	5	3	3	-	-	-	-
2000.....	7	2	6	4	4	1	2	-	-	-	-	-
1999.....	3	5	2	-	3	-	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Hispanic, male

2003.....	7	3	4	3	1	2	-	-	-	-	-	-
2002.....	11	6	3	2	1	2	-	-	1	-	-	-
2001.....	17	8	10	10	4	2	3	2	-	-	-	-
2000.....	6	2	6	3	4	1	2	-	-	-	-	-
1999.....	3	5	2	-	3	-	-	1	-	-	-	-

Hispanic, female

2003.....	1	5	-	2	1	-	-	-	-	-	-	-
2002.....	1	1	-	-	1	-	1	-	-	-	-	-
2001.....	13	13	7	5	4	3	-	1	-	-	-	-
2000.....	1	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	205	179	142	87	68	58	27	15	2	1	-	-
2002.....	171	157	123	92	70	55	38	16	3	-	-	-
2001.....	184	125	100	79	73	56	33	12	3	2	-	-
2000.....	175	132	115	107	61	44	31	10	1	1	-	-
1999.....	170	140	113	83	72	38	28	9	4	1	-	-

Non-Hispanic 1/, male

2003.....	176	153	129	73	57	48	16	12	1	-	-	-
2002.....	139	133	101	87	65	41	31	9	1	-	-	-
2001.....	157	107	93	68	63	39	24	6	2	-	-	-
2000.....	152	113	92	85	51	36	23	8	1	-	-	-
1999.....	142	122	97	70	61	26	18	6	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

Non-Hispanic 1/, female

2003.....	29	26	13	14	11	10	11	3	1	1	-	-
2002.....	32	24	22	5	5	14	7	7	2	-	-	-
2001.....	27	18	7	11	10	17	9	6	1	2	-	-
2000.....	23	19	23	22	10	8	8	2	-	1	-	-
1999.....	28	18	16	13	11	12	10	3	2	1	-	-

Non-Hispanic White, both sexes

2003.....	184	166	128	73	66	50	25	15	2	1	-	-
2002.....	151	140	114	80	64	49	33	15	3	-	-	-
2001.....	161	103	93	70	68	49	31	10	3	1	-	-
2000.....	157	122	105	94	54	40	31	9	1	1	-	-
1999.....	148	121	95	75	65	34	23	7	4	1	-	-

Non-Hispanic White, male

2003.....	156	141	118	61	56	43	14	12	1	-	-	-
2002.....	121	116	93	75	59	35	27	8	1	-	-	-
2001.....	137	86	86	59	59	35	23	6	2	-	-	-
2000.....	138	103	84	76	46	32	23	7	1	-	-	-
1999.....	125	106	81	63	54	25	15	4	2	-	-	-

Non-Hispanic White, female

2003.....	28	25	10	12	10	7	11	3	1	1	-	-
2002.....	30	24	21	5	5	14	6	7	2	-	-	-
2001.....	24	17	7	11	9	14	8	4	1	1	-	-
2000.....	19	19	21	18	8	8	8	2	-	1	-	-
1999.....	23	15	14	12	11	9	8	3	2	1	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

All origins, both sexes

2003.....	3,874	2,679	2,050	2,133	2,738	4,093	5,638	5,845	4,002	1,565	333	56
2002.....	3,464	2,328	1,852	2,075	2,660	4,083	5,388	5,571	3,925	1,548	285	46
2001.....	3,003	2,007	1,790	1,939	2,614	4,029	5,152	5,363	3,770	1,456	310	77
2000.....	2,691	1,906	1,716	1,776	2,607	3,736	4,680	4,948	3,641	1,370	301	52
1999.....	2,419	1,813	1,660	1,965	2,664	3,870	4,944	5,174	3,643	1,330	252	69

All origins, male

2003.....	2,673	1,835	1,402	1,346	1,600	2,232	2,734	2,457	1,408	410	67	52
2002.....	2,463	1,640	1,264	1,330	1,600	2,208	2,562	2,268	1,308	454	50	38
2001.....	2,192	1,405	1,192	1,226	1,510	2,204	2,465	2,238	1,323	353	59	67
2000.....	2,038	1,348	1,168	1,121	1,497	1,989	2,216	2,024	1,224	370	61	50
1999.....	1,824	1,299	1,155	1,245	1,585	2,007	2,406	2,096	1,218	351	57	60

All origins, female

2003.....	1,201	844	648	787	1,138	1,861	2,904	3,388	2,594	1,155	266	4
2002.....	1,001	688	588	745	1,060	1,875	2,826	3,303	2,617	1,094	235	8
2001.....	811	602	598	713	1,104	1,825	2,687	3,125	2,447	1,103	251	10
2000.....	653	558	548	655	1,110	1,747	2,464	2,924	2,417	1,000	240	2
1999.....	595	514	505	720	1,079	1,863	2,538	3,078	2,425	979	195	9

Hispanic, both sexes

2003.....	272	161	149	118	121	151	152	130	76	32	7	7
2002.....	246	156	110	119	130	159	148	102	80	42	6	11
2001.....	207	133	100	91	134	123	119	113	75	34	6	19
2000.....	184	119	105	89	109	117	127	120	84	28	8	14
1999.....	181	102	111	113	99	107	107	114	74	27	9	33

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

Hispanic, male

2003.....	205	127	110	75	72	96	79	62	26	7	2	7
2002.....	203	121	79	84	74	89	78	51	35	18	3	11
2001.....	167	106	71	57	77	69	52	43	29	7	2	18
2000.....	146	86	72	63	68	64	53	56	35	9	3	14
1999.....	145	89	84	77	67	54	51	54	32	14	1	29

Hispanic, female

2003.....	67	34	39	43	49	55	73	68	50	25	5	-
2002.....	43	35	31	35	56	70	70	51	45	24	3	-
2001.....	40	27	29	34	57	54	67	70	46	27	4	1
2000.....	38	33	33	26	41	53	74	64	49	19	5	-
1999.....	36	13	27	36	32	53	56	60	42	13	8	4

Non-Hispanic 1/, both sexes

2003.....	3,569	2,503	1,892	2,005	2,609	3,928	5,475	5,701	3,918	1,531	326	19
2002.....	3,192	2,151	1,733	1,946	2,518	3,905	5,225	5,456	3,836	1,502	279	18
2001.....	2,777	1,852	1,677	1,834	2,469	3,897	5,018	5,240	3,686	1,415	303	21
2000.....	2,472	1,771	1,600	1,680	2,485	3,604	4,543	4,814	3,548	1,339	292	9
1999.....	2,207	1,685	1,533	1,841	2,559	3,747	4,820	5,041	3,558	1,299	243	15

Non-Hispanic 1/, male

2003.....	2,441	1,695	1,286	1,263	1,521	2,125	2,650	2,389	1,378	403	65	16
2002.....	2,240	1,498	1,176	1,237	1,516	2,110	2,472	2,210	1,272	434	47	16
2001.....	2,008	1,280	1,110	1,161	1,425	2,128	2,404	2,192	1,290	346	57	17
2000.....	1,866	1,249	1,087	1,051	1,422	1,916	2,160	1,962	1,184	360	58	8
1999.....	1,652	1,191	1,057	1,159	1,513	1,944	2,345	2,034	1,180	335	56	13

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

Non-Hispanic 1/, female

2003.....	1,128	808	606	742	1,088	1,803	2,825	3,312	2,540	1,128	261	3
2002.....	952	653	557	709	1,002	1,795	2,753	3,246	2,564	1,068	232	2
2001.....	769	572	567	673	1,044	1,769	2,614	3,048	2,396	1,069	246	4
2000.....	606	522	513	629	1,063	1,688	2,383	2,852	2,364	979	234	1
1999.....	555	494	476	682	1,046	1,803	2,475	3,007	2,378	964	187	2

Non-Hispanic White, both sexes

2003.....	2,817	2,051	1,578	1,662	2,261	3,564	5,031	5,389	3,685	1,459	299	18
2002.....	2,529	1,721	1,426	1,594	2,192	3,502	4,840	5,137	3,596	1,409	255	16
2001.....	2,163	1,482	1,357	1,549	2,132	3,492	4,621	4,897	3,452	1,321	274	17
2000.....	1,896	1,408	1,306	1,387	2,145	3,227	4,192	4,523	3,329	1,242	261	5
1999.....	1,699	1,325	1,184	1,514	2,157	3,310	4,427	4,677	3,353	1,216	221	9

Non-Hispanic White, male

2003.....	1,924	1,371	1,059	1,043	1,325	1,928	2,423	2,257	1,288	382	58	15
2002.....	1,747	1,177	957	1,003	1,327	1,894	2,281	2,077	1,176	403	38	14
2001.....	1,539	1,006	890	983	1,237	1,907	2,195	2,046	1,199	314	50	15
2000.....	1,419	976	881	858	1,237	1,699	1,984	1,836	1,099	322	50	4
1999.....	1,263	913	831	942	1,275	1,703	2,162	1,879	1,093	315	51	8

Non-Hispanic White, female

2003.....	893	680	519	619	936	1,636	2,608	3,132	2,397	1,077	241	3
2002.....	782	544	469	591	865	1,608	2,559	3,060	2,420	1,006	217	2
2001.....	624	476	467	566	895	1,585	2,426	2,851	2,253	1,007	224	2
2000.....	477	432	425	529	908	1,528	2,208	2,687	2,230	920	211	1
1999.....	436	412	353	572	882	1,607	2,265	2,798	2,260	901	170	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

Non-Hispanic Black, both sexes

2003.....	652	388	263	277	270	260	321	228	167	57	20	1
2002.....	593	369	262	277	257	316	277	245	182	74	17	1
2001.....	556	319	271	244	285	321	311	257	182	68	25	1
2000.....	509	319	247	250	285	312	277	229	173	77	28	4
1999.....	450	318	299	293	336	363	330	290	176	76	18	5

Non-Hispanic Black, male

2003.....	447	280	190	184	151	145	152	94	58	16	5	1
2002.....	439	277	187	184	147	171	138	97	71	24	5	1
2001.....	422	234	184	157	163	172	166	99	66	20	6	1
2000.....	393	243	179	163	156	178	136	93	67	30	7	4
1999.....	347	247	192	194	206	198	146	121	68	17	4	4

Non-Hispanic Black, female

2003.....	205	108	73	93	119	115	169	134	109	41	15	-
2002.....	154	92	75	93	110	145	139	148	111	50	12	-
2001.....	134	85	87	87	122	149	145	158	116	48	19	-
2000.....	116	76	68	87	129	134	141	136	106	47	21	-
1999.....	103	71	107	99	130	165	184	169	108	59	14	1

Origin not stated 2/, both sexes

2003.....	33	15	9	10	8	14	11	14	8	2	-	30
2002.....	26	21	9	10	12	19	15	13	9	4	-	17
2001.....	19	22	13	14	11	9	15	10	9	7	1	37
2000.....	35	16	11	7	13	15	10	14	9	3	1	29
1999.....	31	26	16	11	6	16	17	19	11	4	-	21

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Nontransport accidents (W00-X59,Y86)

Origin not stated 2/, male

2003.....	27	13	6	8	7	11	5	6	4	-	-	29
2002.....	20	21	9	9	10	9	12	7	1	2	-	11
2001.....	17	19	11	8	8	7	9	3	4	-	-	32
2000.....	26	13	9	7	7	9	3	6	5	1	-	28
1999.....	27	19	14	9	5	9	10	8	6	2	-	18

Origin not stated 2/, female

2003.....	6	2	3	2	1	3	6	8	4	2	-	1
2002.....	6	-	-	1	2	10	3	6	8	2	-	6
2001.....	2	3	2	6	3	2	6	7	5	7	1	5
2000.....	9	3	2	-	6	6	7	8	4	2	1	1
1999.....	4	7	2	2	1	7	7	11	5	2	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Falls (W00-W19)												
All origins, both sexes												
2003.....	535	579	641	787	1,261	2,136	3,113	3,220	2,221	806	157	1
2002.....	530	515	574	759	1,208	2,011	2,869	3,032	2,045	784	129	1
2001.....	504	463	541	747	1,086	1,873	2,567	2,647	1,850	718	135	4
2000.....	462	448	501	595	1,065	1,641	2,200	2,352	1,682	618	120	-
1999.....	377	398	489	663	1,000	1,636	2,220	2,308	1,625	533	112	4
All origins, male												
2003.....	409	408	419	485	724	1,145	1,516	1,359	796	206	25	1
2002.....	394	379	392	474	728	1,079	1,365	1,262	691	231	24	-
2001.....	395	344	374	469	621	1,041	1,236	1,149	655	188	31	3
2000.....	358	326	354	378	613	895	1,052	992	592	173	27	-
1999.....	295	296	354	433	594	840	1,110	964	568	140	29	4
All origins, female												
2003.....	126	171	222	302	537	991	1,597	1,861	1,425	600	132	-
2002.....	136	136	182	285	480	932	1,504	1,770	1,354	553	105	1
2001.....	109	119	167	278	465	832	1,331	1,498	1,195	530	104	1
2000.....	104	122	147	217	452	746	1,148	1,360	1,090	445	93	-
1999.....	82	102	135	230	406	796	1,110	1,344	1,057	393	83	-
Hispanic, both sexes												
2003.....	50	41	64	42	71	79	84	78	51	23	4	-
2002.....	46	45	42	46	72	80	95	57	46	26	-	-
2001.....	36	40	38	35	68	67	64	53	38	17	4	1
2000.....	38	35	39	40	46	60	74	67	40	11	4	-
1999.....	31	30	33	47	37	55	61	65	41	12	5	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Falls (W00-W19)

## Hispanic, male

2003.....	40	30	44	25	39	51	43	38	15	5	1	-
2002.....	35	36	29	33	39	40	50	32	16	10	-	-
2001.....	26	32	30	21	36	33	29	20	14	3	1	1
2000.....	34	27	26	28	32	34	28	31	16	3	2	-
1999.....	28	28	27	28	24	29	31	32	17	6	1	2

## Hispanic, female

2003.....	10	11	20	17	32	28	41	40	36	18	3	-
2002.....	11	9	13	13	33	40	45	25	30	16	-	-
2001.....	10	8	8	14	32	34	35	33	24	14	3	-
2000.....	4	8	13	12	14	26	46	36	24	8	2	-
1999.....	3	2	6	19	13	26	30	33	24	6	4	-

## Non-Hispanic 1/, both sexes

2003.....	484	535	574	743	1,187	2,051	3,022	3,136	2,165	781	153	-
2002.....	477	467	528	710	1,132	1,920	2,765	2,967	1,997	756	129	-
2001.....	468	421	500	704	1,014	1,805	2,494	2,588	1,808	697	130	-
2000.....	418	409	460	553	1,016	1,577	2,123	2,281	1,637	606	116	-
1999.....	345	363	450	611	961	1,571	2,150	2,235	1,579	519	107	1

## Non-Hispanic 1/, male

2003.....	368	376	374	458	682	1,089	1,471	1,318	779	201	24	-
2002.....	352	340	359	438	685	1,034	1,308	1,226	675	220	24	-
2001.....	369	310	341	443	581	1,007	1,202	1,127	640	185	30	-
2000.....	319	295	327	348	580	858	1,023	959	572	169	25	-
1999.....	266	264	323	400	569	805	1,073	928	549	134	28	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Falls (W00-W19)

## Non-Hispanic 1/, female

2003.....	116	159	200	285	505	962	1,551	1,818	1,386	580	129	-
2002.....	125	127	169	272	447	886	1,457	1,741	1,322	536	105	-
2001.....	99	111	159	261	433	798	1,292	1,461	1,168	512	100	-
2000.....	99	114	133	205	436	719	1,100	1,322	1,065	437	91	-
1999.....	79	99	127	211	392	766	1,077	1,307	1,030	385	79	-

## Non-Hispanic White, both sexes

2003.....	382	449	494	644	1,067	1,912	2,835	3,010	2,066	759	146	-
2002.....	398	391	462	621	1,023	1,784	2,635	2,846	1,897	726	122	-
2001.....	395	350	430	618	922	1,675	2,369	2,465	1,727	666	120	-
2000.....	335	336	389	473	920	1,475	2,012	2,191	1,570	572	107	-
1999.....	270	291	362	524	855	1,456	2,044	2,133	1,508	492	98	1

## Non-Hispanic White, male

2003.....	287	317	321	390	618	1,019	1,376	1,265	741	195	24	-
2002.....	290	284	312	381	621	959	1,240	1,170	631	209	22	-
2001.....	308	259	292	389	532	933	1,134	1,068	607	169	26	-
2000.....	252	239	280	297	532	796	968	918	540	155	23	-
1999.....	209	209	262	339	507	739	1,020	879	518	125	27	1

## Non-Hispanic White, female

2003.....	95	132	173	254	449	893	1,459	1,745	1,325	564	122	-
2002.....	108	107	150	240	402	825	1,395	1,676	1,266	517	100	-
2001.....	87	91	138	229	390	742	1,235	1,397	1,120	497	94	-
2000.....	83	97	109	176	388	679	1,044	1,273	1,030	417	84	-
1999.....	61	82	100	185	348	717	1,024	1,254	990	367	71	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Falls (W00-W19)

## Non-Hispanic Black, both sexes

2003.....	75	68	63	71	79	76	109	74	58	14	4	-
2002.....	65	56	54	56	79	85	69	71	65	22	5	-
2001.....	66	55	53	72	71	94	79	69	49	16	7	-
2000.....	66	60	54	63	71	65	72	50	45	21	7	-
1999.....	63	57	70	80	70	73	80	59	53	21	7	-

## Non-Hispanic Black, male

2003.....	56	46	40	51	36	38	47	32	18	3	-	-
2002.....	51	41	37	36	45	48	37	33	28	8	1	-
2001.....	55	40	36	46	39	52	45	29	19	8	3	-
2000.....	53	47	40	40	36	39	31	20	22	9	1	-
1999.....	48	43	47	57	43	41	36	23	19	6	1	-

## Non-Hispanic Black, female

2003.....	19	22	23	20	43	38	62	42	40	11	4	-
2002.....	14	15	17	20	34	37	32	38	37	14	4	-
2001.....	11	15	17	26	32	42	34	40	30	8	4	-
2000.....	13	13	14	23	35	26	41	30	23	12	6	-
1999.....	15	14	23	23	27	32	44	36	34	15	6	-

## Origin not stated 2/, both sexes

2003.....	1	3	3	2	3	6	7	6	5	2	-	1
2002.....	7	3	4	3	4	11	9	8	2	2	-	1
2001.....	-	2	3	8	4	1	9	6	4	4	1	3
2000.....	6	4	2	2	3	4	3	4	5	1	-	-
1999.....	1	5	6	5	2	10	9	8	5	2	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Falls (W00-W19)

Origin not stated 2/, male

2003.....	1	2	1	2	3	5	2	3	2	-	-	1
2002.....	7	3	4	3	4	5	7	4	-	1	-	-
2001.....	-	2	3	5	4	1	5	2	1	-	-	2
2000.....	5	4	1	2	1	3	1	2	4	1	-	-
1999.....	1	4	4	5	1	6	6	4	2	-	-	1

Origin not stated 2/, female

2003.....	-	1	2	-	-	1	5	3	3	2	-	-
2002.....	-	-	-	-	-	6	2	4	2	1	-	1
2001.....	-	-	-	3	-	-	4	4	3	4	1	1
2000.....	1	-	1	-	2	1	2	2	1	-	-	-
1999.....	-	1	2	-	1	4	3	4	3	2	-	-



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental discharge of firearms (W32-W34)

Hispanic, male

2003.....	1	1	2	1	-	1	-	-	-	-	-	-
2002.....	1	1	-	-	-	-	1	-	-	-	-	-
2001.....	3	-	1	1	-	2	-	1	-	-	-	1
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	1	-	2	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	1	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	1	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	42	35	24	22	16	15	21	3	3	-	-	-
2002.....	42	35	20	17	19	17	13	5	1	1	-	-
2001.....	39	44	24	18	14	22	17	5	2	-	-	1
2000.....	40	35	14	24	17	17	7	7	4	-	-	-
1999.....	41	24	23	13	18	15	8	8	2	-	-	-

Non-Hispanic 1/, male

2003.....	34	34	22	19	10	14	18	3	3	-	-	-
2002.....	38	31	18	15	17	14	10	4	1	1	-	-
2001.....	33	34	20	15	10	20	14	4	2	-	-	-
2000.....	38	30	14	20	14	16	7	7	4	-	-	-
1999.....	32	21	19	11	18	13	7	5	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental discharge of firearms (W32-W34)

Non-Hispanic 1/, female

2003.....	8	1	2	3	6	1	3	-	-	-	-	-
2002.....	4	4	2	2	2	3	3	1	-	-	-	-
2001.....	6	10	4	3	4	2	3	1	-	-	-	1
2000.....	2	5	-	4	3	1	-	-	-	-	-	-
1999.....	9	3	4	2	-	2	1	3	-	-	-	-

Non-Hispanic White, both sexes

2003.....	39	33	23	21	12	15	17	2	2	-	-	-
2002.....	36	31	18	16	19	14	13	5	1	1	-	-
2001.....	38	38	23	17	12	21	15	5	2	-	-	1
2000.....	37	33	13	22	15	16	6	7	4	-	-	-
1999.....	36	20	20	11	17	13	7	8	2	-	-	-

Non-Hispanic White, male

2003.....	32	33	21	18	8	14	14	2	2	-	-	-
2002.....	33	28	17	14	17	11	10	4	1	1	-	-
2001.....	33	29	19	14	8	19	12	4	2	-	-	-
2000.....	35	28	13	19	12	15	6	7	4	-	-	-
1999.....	27	18	18	9	17	11	6	5	2	-	-	-

Non-Hispanic White, female

2003.....	7	-	2	3	4	1	3	-	-	-	-	-
2002.....	3	3	1	2	2	3	3	1	-	-	-	-
2001.....	5	9	4	3	4	2	3	1	-	-	-	1
2000.....	2	5	-	3	3	1	-	-	-	-	-	-
1999.....	9	2	2	2	-	2	1	3	-	-	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

All origins, both sexes

2003.....	186	153	112	102	100	94	68	58	20	5	-	30
2002.....	169	118	94	81	88	88	79	48	25	5	1	20
2001.....	145	109	97	78	81	85	72	56	15	4	-	34
2000.....	172	120	97	80	99	71	85	37	19	8	-	24
1999.....	172	122	97	84	101	88	70	45	16	7	-	34

All origins, male

2003.....	147	125	86	73	72	57	47	31	15	2	-	30
2002.....	137	93	68	64	68	56	50	31	15	4	-	18
2001.....	117	83	72	61	63	55	47	37	8	-	-	29
2000.....	143	102	77	63	70	38	50	22	7	5	-	23
1999.....	145	95	75	64	65	63	48	19	6	3	-	30

All origins, female

2003.....	39	28	26	29	28	37	21	27	5	3	-	-
2002.....	32	25	26	17	20	32	29	17	10	1	1	2
2001.....	28	26	25	17	18	30	25	19	7	4	-	5
2000.....	29	18	20	17	29	33	35	15	12	3	-	1
1999.....	27	27	22	20	36	25	22	26	10	4	-	4

Hispanic, both sexes

2003.....	8	10	5	10	4	3	3	1	-	1	-	7
2002.....	12	8	7	4	8	8	2	1	1	-	-	5
2001.....	4	2	5	3	2	3	1	2	-	-	-	12
2000.....	9	7	4	3	9	1	3	-	1	-	-	8
1999.....	13	4	4	3	4	5	1	-	-	-	-	25

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

Hispanic, male

2003.....	6	9	5	6	1	2	1	1	-	-	-	7
2002.....	12	6	4	4	6	6	1	1	1	-	-	5
2001.....	3	1	5	1	2	3	-	2	-	-	-	11
2000.....	7	6	2	3	7	1	2	-	-	-	-	8
1999.....	13	3	4	3	4	4	-	-	-	-	-	21

Hispanic, female

2003.....	2	1	-	4	3	1	2	-	-	1	-	-
2002.....	-	2	3	-	2	2	1	-	-	-	-	-
2001.....	1	1	-	2	-	-	1	-	-	-	-	1
2000.....	2	1	2	-	2	-	1	-	1	-	-	-
1999.....	-	1	-	-	-	1	1	-	-	-	-	4

Non-Hispanic 1/, both sexes

2003.....	174	143	105	92	95	91	65	57	20	4	-	8
2002.....	155	108	87	77	79	80	75	47	24	5	1	7
2001.....	139	106	92	74	79	82	71	54	15	4	-	8
2000.....	163	112	93	77	89	70	82	36	18	7	-	1
1999.....	159	118	92	81	97	83	69	45	16	6	-	3

Non-Hispanic 1/, male

2003.....	137	116	79	67	70	55	46	30	15	2	-	8
2002.....	123	85	64	60	61	50	47	30	14	4	-	7
2001.....	112	81	67	59	61	52	47	35	8	-	-	5
2000.....	136	95	75	60	62	37	48	21	7	5	-	1
1999.....	132	92	70	61	61	59	48	19	6	2	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

Non-Hispanic 1/, female

2003.....	37	27	26	25	25	36	19	27	5	2	-	-
2002.....	32	23	23	17	18	30	28	17	10	1	1	-
2001.....	27	25	25	15	18	30	24	19	7	4	-	3
2000.....	27	17	18	17	27	33	34	15	11	2	-	-
1999.....	27	26	22	20	36	24	21	26	10	4	-	-

Non-Hispanic White, both sexes

2003.....	143	121	87	82	81	77	62	48	17	3	-	8
2002.....	119	94	71	64	62	67	64	40	21	3	-	7
2001.....	119	82	76	62	71	73	61	48	14	4	-	4
2000.....	121	97	78	67	77	57	77	33	15	5	-	-
1999.....	130	96	78	69	78	69	63	39	16	4	-	2

Non-Hispanic White, male

2003.....	111	96	64	60	61	49	44	28	12	2	-	8
2002.....	96	74	51	50	49	44	41	28	13	3	-	7
2001.....	94	61	53	50	55	47	41	31	8	-	-	3
2000.....	100	82	68	52	52	30	48	19	7	3	-	-
1999.....	110	74	59	50	49	49	44	17	6	2	-	2

Non-Hispanic White, female

2003.....	32	25	23	22	20	28	18	20	5	1	-	-
2002.....	23	20	20	14	13	23	23	12	8	-	-	-
2001.....	25	21	23	12	16	26	20	17	6	4	-	1
2000.....	21	15	10	15	25	27	29	14	8	2	-	-
1999.....	20	22	19	19	29	20	19	22	10	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental drowning and submersion (W65-W74)

Non-Hispanic Black, both sexes

2003.....	22	19	14	5	11	6	2	4	1	-	-	-
2002.....	24	11	10	9	9	10	7	6	2	2	1	-
2001.....	17	21	11	7	6	5	8	3	1	-	-	1
2000.....	29	9	8	7	8	9	3	3	2	2	-	1
1999.....	19	19	9	9	12	11	4	3	-	2	-	-

Non-Hispanic Black, male

2003.....	20	17	12	3	7	3	2	2	1	-	-	-
2002.....	19	10	8	8	7	5	4	2	1	1	-	-
2001.....	15	17	9	7	5	4	5	3	-	-	-	1
2000.....	24	8	5	6	7	4	-	2	-	2	-	1
1999.....	17	16	6	8	9	8	3	2	-	-	-	-

Non-Hispanic Black, female

2003.....	2	2	2	2	4	3	-	2	-	-	-	-
2002.....	5	1	2	1	2	5	3	4	1	1	1	-
2001.....	2	4	2	-	1	1	3	-	1	-	-	-
2000.....	5	1	3	1	1	5	3	1	2	-	-	-
1999.....	2	3	3	1	3	3	1	1	-	2	-	-

Origin not stated 2/, both sexes

2003.....	4	-	2	-	1	-	-	-	-	-	-	15
2002.....	2	2	-	-	1	-	2	-	-	-	-	8
2001.....	2	1	-	1	-	-	-	-	-	-	-	14
2000.....	-	1	-	-	1	-	-	1	-	1	-	15
1999.....	-	-	1	-	-	-	-	-	-	1	-	6



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

All origins, both sexes

2003.....	256	204	188	198	221	226	234	157	67	24	5	4
2002.....	235	183	157	174	211	211	221	129	70	18	4	3
2001.....	206	215	163	190	221	225	198	159	72	18	2	3
2000.....	225	187	182	188	213	257	215	158	78	24	3	5
1999.....	170	157	162	193	223	258	229	150	74	18	3	4

All origins, male

2003.....	148	138	116	113	131	124	104	72	27	8	-	3
2002.....	146	111	108	117	125	108	123	55	31	9	2	3
2001.....	137	142	94	112	117	106	98	69	30	8	1	3
2000.....	158	126	115	107	125	136	108	69	34	11	1	5
1999.....	110	98	96	111	133	128	119	67	34	9	1	4

All origins, female

2003.....	108	66	72	85	90	102	130	85	40	16	5	1
2002.....	89	72	49	57	86	103	98	74	39	9	2	-
2001.....	69	73	69	78	104	119	100	90	42	10	1	-
2000.....	67	61	67	81	88	121	107	89	44	13	2	-
1999.....	60	59	66	82	90	130	110	83	40	9	2	-

Hispanic, both sexes

2003.....	19	7	7	7	5	10	5	7	1	1	-	-
2002.....	10	15	2	12	8	7	9	9	-	1	-	1
2001.....	5	7	6	5	4	4	5	9	-	1	-	-
2000.....	9	7	10	5	13	10	7	5	1	-	-	-
1999.....	6	3	11	8	3	8	7	3	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Hispanic, male

2003.....	15	6	4	5	4	5	4	3	1	1	-	-
2002.....	9	10	2	9	5	4	4	3	-	1	-	1
2001.....	1	6	3	3	3	3	2	4	-	-	-	-
2000.....	6	5	4	4	6	6	4	4	-	-	-	-
1999.....	3	1	10	6	2	5	4	3	2	-	-	-

Hispanic, female

2003.....	4	1	3	2	1	5	1	4	-	-	-	-
2002.....	1	5	-	3	3	3	5	6	-	-	-	-
2001.....	4	1	3	2	1	1	3	5	-	1	-	-
2000.....	3	2	6	1	7	4	3	1	1	-	-	-
1999.....	3	2	1	2	1	3	3	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	234	195	180	189	215	214	229	149	66	23	5	2
2002.....	224	164	153	161	201	202	212	120	70	17	4	1
2001.....	198	206	155	183	215	219	193	150	71	17	2	2
2000.....	215	178	171	183	199	244	207	152	77	24	3	3
1999.....	161	152	150	185	219	249	221	147	72	18	3	3

Non-Hispanic 1/, male

2003.....	132	130	111	107	126	118	100	68	26	7	-	1
2002.....	137	97	104	107	120	104	119	52	31	8	2	1
2001.....	134	134	90	108	114	103	96	65	29	8	1	2
2000.....	152	120	110	103	119	128	104	64	34	11	1	3
1999.....	105	96	85	105	130	122	115	64	32	9	1	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic 1/, female

2003.....	102	65	69	82	89	96	129	81	40	16	5	1
2002.....	87	67	49	54	81	98	93	68	39	9	2	-
2001.....	64	72	65	75	101	116	97	85	42	9	1	-
2000.....	63	58	61	80	80	116	103	88	43	13	2	-
1999.....	56	56	65	80	89	127	106	83	40	9	2	-

Non-Hispanic White, both sexes

2003.....	168	147	139	143	162	164	176	112	58	18	2	2
2002.....	163	122	105	119	149	155	170	95	49	10	3	1
2001.....	141	146	115	133	158	164	147	117	53	9	1	2
2000.....	167	122	120	134	144	179	149	123	50	17	-	3
1999.....	119	118	100	128	153	178	165	105	51	13	2	2

Non-Hispanic White, male

2003.....	100	95	88	81	97	85	74	54	22	5	-	1
2002.....	96	73	73	78	88	81	101	44	20	4	1	1
2001.....	92	97	63	77	80	73	76	53	21	4	-	2
2000.....	117	85	75	72	87	94	78	56	22	8	-	3
1999.....	77	72	64	71	89	83	91	49	24	7	1	2

Non-Hispanic White, female

2003.....	68	52	51	62	65	79	102	58	36	13	2	1
2002.....	67	49	32	41	61	74	69	51	29	6	2	-
2001.....	49	49	52	56	78	91	71	64	32	5	1	-
2000.....	50	37	45	62	57	85	71	67	28	9	-	-
1999.....	42	46	36	57	64	95	74	56	27	6	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Non-Hispanic Black, both sexes

2003.....	61	45	34	42	48	50	49	35	8	4	3	-
2002.....	58	38	46	36	47	42	40	25	20	7	1	-
2001.....	51	53	36	46	53	49	43	31	17	8	1	-
2000.....	46	51	49	45	53	61	56	29	26	7	3	-
1999.....	40	34	46	52	61	67	53	41	21	5	1	1

Non-Hispanic Black, male

2003.....	31	33	20	26	27	33	23	13	4	1	-	-
2002.....	39	22	30	25	28	20	18	8	11	4	1	-
2001.....	39	32	24	28	33	26	18	11	7	4	1	-
2000.....	33	33	34	29	31	33	25	8	12	3	1	-
1999.....	27	24	20	30	39	36	23	15	8	2	-	1

Non-Hispanic Black, female

2003.....	30	12	14	16	21	17	26	22	4	3	3	-
2002.....	19	16	16	11	19	22	22	17	9	3	-	-
2001.....	12	21	12	18	20	23	25	20	10	4	-	-
2000.....	13	18	15	16	22	28	31	21	14	4	2	-
1999.....	13	10	26	22	22	31	30	26	13	3	1	-

Origin not stated 2/, both sexes

2003.....	3	2	1	2	1	2	-	1	-	-	-	2
2002.....	1	4	2	1	2	2	-	-	-	-	-	1
2001.....	3	2	2	2	2	2	-	-	1	-	-	1
2000.....	1	2	1	-	1	3	1	1	-	-	-	2
1999.....	3	2	1	-	1	1	1	-	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental exposure to smoke, fire and flames (X00-X09)

Origin not stated 2/, male

2003.....	1	2	1	1	1	1	-	1	-	-	-	2
2002.....	-	4	2	1	-	-	-	-	-	-	-	1
2001.....	2	2	1	1	-	-	-	-	1	-	-	1
2000.....	-	1	1	-	-	2	-	1	-	-	-	2
1999.....	2	1	1	-	1	1	-	-	-	-	-	1

Origin not stated 2/, female

2003.....	2	-	-	1	-	1	-	-	-	-	-	-
2002.....	1	-	-	-	2	2	-	-	-	-	-	-
2001.....	1	-	1	1	2	2	-	-	-	-	-	-
2000.....	1	1	-	-	1	1	1	-	-	-	-	-
1999.....	1	1	-	-	-	-	1	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins, both sexes

2003.....	2,083	957	413	229	174	146	138	105	47	10	4	6
2002.....	1,736	754	344	218	165	154	141	103	64	13	1	9
2001.....	1,281	530	268	161	130	153	126	91	47	12	2	10
2000.....	1,069	447	241	146	132	118	127	109	60	14	1	7
1999.....	918	449	219	158	142	138	127	100	31	12	-	12

All origins, male

2003.....	1,368	584	257	137	86	71	71	34	19	4	1	5
2002.....	1,176	491	199	126	83	71	65	38	25	8	-	7
2001.....	893	341	159	88	82	83	54	28	18	2	1	10
2000.....	768	260	141	79	68	49	53	51	23	2	-	6
1999.....	650	298	150	98	74	63	53	38	10	5	-	8

All origins, female

2003.....	715	373	156	92	88	75	67	71	28	6	3	1
2002.....	560	263	145	92	82	83	76	65	39	5	1	2
2001.....	388	189	109	73	48	70	72	63	29	10	1	-
2000.....	301	187	100	67	64	69	74	58	37	12	1	1
1999.....	268	151	69	60	68	75	74	62	21	7	-	4

Hispanic, both sexes

2003.....	142	63	28	12	10	7	3	6	-	-	-	-
2002.....	125	53	26	15	11	8	5	1	1	-	-	-
2001.....	98	49	15	10	9	2	5	-	-	1	-	1
2000.....	93	38	15	9	6	2	3	3	2	-	-	-
1999.....	98	37	21	12	6	4	1	1	1	-	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Hispanic, male

2003.....	107	50	23	10	7	2	1	3	-	-	-	-
2002.....	104	39	21	11	7	6	3	-	1	-	-	-
2001.....	83	40	10	7	6	2	1	-	-	-	-	1
2000.....	72	24	11	7	3	1	1	1	2	-	-	-
1999.....	75	33	18	11	4	2	-	1	-	-	-	3

Hispanic, female

2003.....	35	13	5	2	3	5	2	3	-	-	-	-
2002.....	21	14	5	4	4	2	2	1	-	-	-	-
2001.....	15	9	5	3	3	-	4	-	-	1	-	-
2000.....	21	14	4	2	3	1	2	2	-	-	-	-
1999.....	23	4	3	1	2	2	1	-	1	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,917	886	383	214	163	138	135	99	47	10	4	4
2002.....	1,603	695	316	199	152	144	136	102	63	13	1	4
2001.....	1,171	471	248	150	120	150	121	91	47	11	2	4
2000.....	953	401	223	135	124	116	124	106	58	14	1	3
1999.....	801	402	194	144	136	134	124	97	30	12	-	5

Non-Hispanic 1/, male

2003.....	1,241	526	233	124	78	68	70	31	19	4	1	3
2002.....	1,068	446	176	111	74	63	62	38	24	8	-	3
2001.....	799	293	144	80	75	80	53	28	18	2	1	4
2000.....	678	230	127	70	64	48	52	50	21	2	-	2
1999.....	558	259	128	85	70	61	53	36	10	5	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic 1/, female

2003.....	676	360	150	90	85	70	65	68	28	6	3	1
2002.....	535	249	140	88	78	81	74	64	39	5	1	1
2001.....	372	178	104	70	45	70	68	63	29	9	1	-
2000.....	275	171	96	65	60	68	72	56	37	12	1	1
1999.....	243	143	66	59	66	73	71	61	20	7	-	2

Non-Hispanic White, both sexes

2003.....	1,520	723	321	153	139	125	128	89	41	9	4	3
2002.....	1,246	527	245	166	133	128	122	93	57	11	1	2
2001.....	848	362	184	125	101	128	101	80	41	9	-	4
2000.....	681	304	178	112	104	109	115	89	52	11	-	2
1999.....	577	309	135	116	108	124	114	82	28	10	-	3

Non-Hispanic White, male

2003.....	971	409	187	78	66	61	66	26	15	3	1	2
2002.....	807	317	119	88	62	57	51	34	24	7	-	1
2001.....	560	208	96	66	63	63	46	25	17	1	-	4
2000.....	474	158	93	52	52	45	48	42	19	1	-	1
1999.....	385	182	81	67	57	56	49	34	10	4	-	2

Non-Hispanic White, female

2003.....	549	314	134	75	73	64	62	63	26	6	3	1
2002.....	439	210	126	78	71	71	71	59	33	4	1	1
2001.....	288	154	88	59	38	65	55	55	24	8	-	-
2000.....	207	146	85	60	52	64	67	47	33	10	-	1
1999.....	192	127	54	49	51	68	65	48	18	6	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Non-Hispanic Black, both sexes

2003.....	358	148	56	51	17	13	6	9	6	1	-	1
2002.....	336	154	67	26	15	14	12	6	5	1	-	1
2001.....	305	101	56	23	16	17	17	11	6	2	2	-
2000.....	260	91	41	23	18	7	8	16	6	3	1	1
1999.....	209	83	56	25	24	10	8	13	2	2	-	2

Non-Hispanic Black, male

2003.....	247	109	45	41	11	7	3	5	4	1	-	1
2002.....	244	119	55	20	11	6	9	1	-	-	-	1
2001.....	226	78	43	14	11	14	6	3	1	1	1	-
2000.....	194	69	32	18	11	3	4	8	2	1	-	1
1999.....	163	70	44	17	12	5	4	2	-	1	-	1

Non-Hispanic Black, female

2003.....	111	39	11	10	6	6	3	4	2	-	-	-
2002.....	92	35	12	6	4	8	3	5	5	1	-	-
2001.....	79	23	13	9	5	3	11	8	5	1	1	-
2000.....	66	22	9	5	7	4	4	8	4	2	1	-
1999.....	46	13	12	8	12	5	4	11	2	1	-	1

Origin not stated 2/, both sexes

2003.....	24	8	2	3	1	1	-	-	-	-	-	2
2002.....	8	6	2	4	2	2	-	-	-	-	-	5
2001.....	12	10	5	1	1	1	-	-	-	-	-	5
2000.....	23	8	3	2	2	-	-	-	-	-	-	4
1999.....	19	10	4	2	-	-	2	2	-	-	-	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Origin not stated 2/, male

2003.....	20	8	1	3	1	1	-	-	-	-	-	2
2002.....	4	6	2	4	2	2	-	-	-	-	-	4
2001.....	11	8	5	1	1	1	-	-	-	-	-	5
2000.....	18	6	3	2	1	-	-	-	-	-	-	4
1999.....	17	6	4	2	-	-	-	1	-	-	-	2

Origin not stated 2/, female

2003.....	4	-	1	-	-	-	-	-	-	-	-	-
2002.....	4	-	-	-	-	-	-	-	-	-	-	1
2001.....	1	2	-	-	-	-	-	-	-	-	-	-
2000.....	5	2	-	-	1	-	-	-	-	-	-	-
1999.....	2	4	-	-	-	-	2	1	-	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins, both sexes

2003.....	770	750	670	794	966	1,475	2,064	2,302	1,644	720	167	15
2002.....	751	722	663	826	969	1,601	2,064	2,254	1,720	727	150	13
2001.....	825	646	696	744	1,082	1,669	2,172	2,404	1,784	704	171	24
2000.....	723	668	681	743	1,081	1,632	2,046	2,285	1,798	706	177	16
1999.....	741	662	670	852	1,180	1,735	2,290	2,563	1,895	760	137	15

All origins, male

2003.....	566	545	500	518	577	820	978	958	548	190	41	13
2002.....	571	534	479	534	579	879	948	878	545	201	24	10
2001.....	614	461	472	480	617	897	1,016	950	610	155	26	21
2000.....	573	504	467	474	607	855	946	883	564	179	33	16
1999.....	592	490	461	526	701	900	1,069	1,003	598	194	27	14

All origins, female

2003.....	204	205	170	276	389	655	1,086	1,344	1,096	530	126	2
2002.....	180	188	184	292	390	722	1,116	1,376	1,175	526	126	3
2001.....	211	185	224	264	465	772	1,156	1,454	1,174	549	145	3
2000.....	150	164	214	269	474	777	1,100	1,402	1,234	527	144	-
1999.....	149	172	209	326	479	835	1,221	1,560	1,297	566	110	1

Hispanic, both sexes

2003.....	51	39	43	46	31	51	57	38	24	7	3	-
2002.....	52	34	33	42	31	56	36	34	32	15	6	5
2001.....	61	35	35	37	51	45	44	48	37	15	2	4
2000.....	35	31	37	32	35	44	40	45	40	17	4	6
1999.....	33	27	42	41	49	35	37	45	30	15	4	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Hispanic, male

2003.....	36	31	32	28	21	35	30	17	10	1	1	-
2002.....	42	29	23	27	17	33	19	15	17	7	3	5
2001.....	51	27	22	24	30	26	20	16	15	4	1	4
2000.....	27	24	29	21	20	22	18	20	17	6	1	6
1999.....	26	23	25	27	33	14	16	18	13	8	-	3

Hispanic, female

2003.....	15	8	11	18	10	16	27	21	14	6	2	-
2002.....	10	5	10	15	14	23	17	19	15	8	3	-
2001.....	10	8	13	13	21	19	24	32	22	11	1	-
2000.....	8	7	8	11	15	22	22	25	23	11	3	-
1999.....	7	4	17	14	16	21	21	27	17	7	4	-

Non-Hispanic 1/, both sexes

2003.....	718	709	626	745	933	1,419	2,003	2,257	1,617	713	164	5
2002.....	691	682	629	782	935	1,542	2,024	2,215	1,681	710	144	6
2001.....	762	604	658	705	1,027	1,619	2,122	2,352	1,743	686	169	6
2000.....	683	636	639	708	1,040	1,580	2,000	2,232	1,754	688	172	2
1999.....	700	626	624	807	1,128	1,695	2,248	2,509	1,859	744	133	3

Non-Hispanic 1/, male

2003.....	529	513	467	488	555	781	945	939	536	189	40	4
2002.....	522	499	455	506	559	845	926	860	527	193	21	5
2001.....	561	428	448	456	584	866	992	933	593	151	25	6
2000.....	543	479	434	450	583	829	926	861	546	173	32	2
1999.....	559	459	432	497	665	884	1,049	982	581	185	27	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic 1/, female

2003.....	189	196	159	257	378	638	1,058	1,318	1,081	524	124	1
2002.....	169	183	174	276	376	697	1,098	1,355	1,154	517	123	1
2001.....	201	176	210	249	443	753	1,130	1,419	1,150	535	144	-
2000.....	140	157	205	258	457	751	1,074	1,371	1,208	515	140	-
1999.....	141	167	192	310	463	811	1,199	1,527	1,278	559	106	-

Non-Hispanic White, both sexes

2003.....	565	578	514	619	800	1,271	1,813	2,128	1,501	670	147	5
2002.....	567	556	525	608	806	1,354	1,836	2,058	1,571	658	129	6
2001.....	622	504	529	594	868	1,431	1,928	2,182	1,615	633	153	6
2000.....	555	516	528	579	885	1,391	1,833	2,080	1,638	637	154	-
1999.....	567	491	489	666	946	1,470	2,034	2,310	1,748	697	121	1

Non-Hispanic White, male

2003.....	423	421	378	416	475	700	849	882	496	177	33	4
2002.....	425	401	385	392	490	742	838	797	487	179	15	5
2001.....	452	352	367	387	499	772	886	865	544	140	24	6
2000.....	441	384	352	366	502	719	836	794	507	155	27	-
1999.....	455	358	347	406	556	765	952	895	533	177	23	1

Non-Hispanic White, female

2003.....	142	157	136	203	325	571	964	1,246	1,005	493	114	1
2002.....	142	155	140	216	316	612	998	1,261	1,084	479	114	1
2001.....	170	152	162	207	369	659	1,042	1,317	1,071	493	129	-
2000.....	114	132	176	213	383	672	997	1,286	1,131	482	127	-
1999.....	112	133	142	260	390	705	1,082	1,415	1,215	520	98	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Non-Hispanic Black, both sexes

2003.....	133	106	95	107	111	115	152	105	93	38	13	-
2002.....	105	106	83	150	107	162	149	137	90	42	10	-
2001.....	116	84	114	95	137	155	162	143	109	42	15	-
2000.....	105	106	94	111	133	170	137	131	94	44	17	2
1999.....	114	121	115	125	168	200	184	174	100	46	10	2

Non-Hispanic Black, male

2003.....	91	74	72	62	68	64	74	41	30	11	5	-
2002.....	82	82	56	95	56	89	70	53	31	11	3	-
2001.....	87	63	71	61	73	75	90	53	39	7	1	-
2000.....	86	84	67	70	69	99	75	55	31	15	5	2
1999.....	87	91	74	80	102	106	79	79	41	8	3	2

Non-Hispanic Black, female

2003.....	42	32	23	45	43	51	78	64	63	27	8	-
2002.....	23	24	27	55	51	73	79	84	59	31	7	-
2001.....	29	21	43	34	64	80	72	90	70	35	14	-
2000.....	19	22	27	41	64	71	62	76	63	29	12	-
1999.....	27	30	41	45	66	94	105	95	59	38	7	-

Origin not stated 2/, both sexes

2003.....	1	2	1	3	2	5	4	7	3	-	-	10
2002.....	8	6	1	2	3	3	4	5	7	2	-	2
2001.....	2	7	3	2	4	5	6	4	4	3	-	14
2000.....	5	1	5	3	6	8	6	8	4	1	1	8
1999.....	8	9	4	4	3	5	5	9	6	1	-	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Origin not stated 2/, male

2003.....	1	1	1	2	1	4	3	2	2	-	-	9
2002.....	7	6	1	1	3	1	3	3	1	1	-	-
2001.....	2	6	2	-	3	5	4	1	2	-	-	11
2000.....	3	1	4	3	4	4	2	2	1	-	-	8
1999.....	7	8	4	2	3	2	4	3	4	1	-	8

Origin not stated 2/, female

2003.....	-	1	-	1	1	1	1	5	1	-	-	1
2002.....	1	-	-	1	-	2	1	2	6	1	-	2
2001.....	-	1	1	2	1	-	2	3	2	3	-	3
2000.....	2	-	1	-	2	4	4	6	3	1	1	-
1999.....	1	1	-	2	-	3	1	6	2	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

All origins, both sexes

2003.....	2,996	2,314	1,529	1,167	1,168	1,159	956	558	205	34	1	7
2002.....	2,835	2,186	1,432	1,197	1,266	1,231	1,028	573	211	41	1	10
2001 3/.....	2,682	1,985	1,332	1,212	1,220	1,219	973	538	200	28	3	15
2000.....	2,450	1,708	1,237	1,073	1,219	1,227	954	590	205	34	4	7
1999.....	2,264	1,678	1,218	1,187	1,284	1,280	937	580	182	39	-	16

All origins, male

2003.....	2,271	1,771	1,216	958	996	985	824	484	178	28	-	7
2002.....	2,188	1,708	1,129	975	1,078	1,054	884	492	179	33	-	10
2001 3/.....	2,020	1,534	1,029	1,001	1,039	1,033	852	468	168	25	3	14
2000.....	1,842	1,306	959	870	1,018	1,062	822	507	168	26	4	7
1999.....	1,702	1,284	971	972	1,079	1,073	795	501	150	30	-	16

All origins, female

2003.....	725	543	313	209	172	174	132	74	27	6	1	-
2002.....	647	478	303	222	188	177	144	81	32	8	1	-
2001 3/.....	662	451	303	211	181	186	121	70	32	3	-	1
2000.....	608	402	278	203	201	165	132	83	37	8	-	-
1999.....	562	394	247	215	205	207	142	79	32	9	-	-

Hispanic, both sexes

2003.....	114	88	57	44	36	29	22	15	5	1	-	-
2002.....	111	90	52	45	44	30	25	17	4	2	-	2
2001 3/.....	109	65	49	42	44	24	25	14	6	2	-	2
2000.....	103	62	40	43	41	32	27	11	3	2	-	1
1999.....	89	65	47	46	27	35	28	12	7	3	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Hispanic, male

2003.....	90	78	51	38	33	26	21	13	3	1	-	-
2002.....	91	79	44	39	38	25	22	16	4	1	-	2
2001 3/.....	86	56	44	39	38	23	22	14	6	2	-	2
2000.....	82	54	35	39	35	27	26	10	3	2	-	1
1999.....	65	55	39	41	22	30	24	10	6	2	-	2

Hispanic, female

2003.....	24	10	6	6	3	3	1	2	2	-	-	-
2002.....	20	11	8	6	6	5	3	1	-	1	-	-
2001 3/.....	23	9	5	3	6	1	3	-	-	-	-	-
2000.....	21	8	5	4	6	5	1	1	-	-	-	-
1999.....	24	10	8	5	5	5	4	2	1	1	-	-

Non-Hispanic 1/, both sexes

2003.....	2,875	2,215	1,467	1,118	1,129	1,129	934	541	200	33	1	2
2002.....	2,706	2,086	1,369	1,141	1,219	1,198	1,001	554	205	37	1	4
2001 3/.....	2,564	1,909	1,275	1,162	1,172	1,193	945	521	194	26	3	5
2000.....	2,336	1,638	1,189	1,025	1,173	1,194	925	577	201	31	4	4
1999.....	2,159	1,608	1,165	1,133	1,255	1,240	906	565	175	36	-	9

Non-Hispanic 1/, male

2003.....	2,176	1,683	1,161	917	961	958	803	469	175	27	-	2
2002.....	2,083	1,621	1,075	926	1,037	1,026	861	474	173	30	-	4
2001 3/.....	1,925	1,468	978	955	997	1,008	827	451	162	23	3	5
2000.....	1,751	1,244	917	826	979	1,034	794	496	164	24	4	4
1999.....	1,625	1,225	927	923	1,055	1,039	768	489	144	28	-	9

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Non-Hispanic 1/, female

2003.....	699	532	306	201	168	171	131	72	25	6	1	-
2002.....	623	465	294	215	182	172	140	80	32	7	1	-
2001 3/.....	639	441	297	207	175	185	118	70	32	3	-	-
2000.....	585	394	272	199	194	160	131	81	37	7	-	-
1999.....	534	383	238	210	200	201	138	76	31	8	-	-

Non-Hispanic White, both sexes

2003.....	2,692	2,085	1,377	1,043	1,067	1,079	901	518	199	31	1	2
2002.....	2,519	1,965	1,289	1,074	1,154	1,131	974	533	197	34	1	4
2001 3/.....	2,395	1,787	1,192	1,085	1,102	1,144	905	501	189	25	3	4
2000.....	2,163	1,536	1,123	969	1,106	1,141	887	561	194	29	2	3
1999.....	2,019	1,501	1,079	1,050	1,199	1,186	872	548	171	35	-	6

Non-Hispanic White, male

2003.....	2,048	1,588	1,090	860	909	919	777	453	174	27	-	2
2002.....	1,956	1,524	1,022	870	995	974	838	458	167	27	-	4
2001 3/.....	1,801	1,375	909	899	937	966	797	438	158	23	3	4
2000.....	1,619	1,165	866	779	922	992	769	484	158	22	2	3
1999.....	1,512	1,141	860	862	1,007	997	743	475	142	27	-	6

Non-Hispanic White, female

2003.....	644	497	287	183	158	160	124	65	25	4	1	-
2002.....	563	441	267	204	159	157	136	75	30	7	1	-
2001 3/.....	594	412	283	186	165	178	108	63	31	2	-	-
2000.....	544	371	257	190	184	149	118	77	36	7	-	-
1999.....	507	360	219	188	192	189	129	73	29	8	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Non-Hispanic Black, both sexes

2003.....	126	73	59	43	34	25	18	11	-	1	-	-
2002.....	116	68	57	33	47	38	13	15	2	2	-	-
2001 3/.....	116	78	48	42	48	29	23	8	4	1	-	-
2000.....	116	60	41	39	46	30	23	9	2	-	1	-
1999.....	82	67	49	58	36	35	17	10	4	1	-	2

Non-Hispanic Black, male

2003.....	91	54	48	32	27	23	16	8	-	-	-	-
2002.....	83	57	40	31	35	33	13	13	2	2	-	-
2001 3/.....	87	62	40	31	42	25	19	5	4	-	-	-
2000.....	87	49	31	36	40	25	15	8	2	-	1	-
1999.....	69	54	41	47	31	30	15	9	2	1	-	2

Non-Hispanic Black, female

2003.....	35	19	11	11	7	2	2	3	-	1	-	-
2002.....	33	11	17	2	12	5	-	2	-	-	-	-
2001 3/.....	29	16	8	11	6	4	4	3	-	1	-	-
2000.....	29	11	10	3	6	5	8	1	-	-	-	-
1999.....	13	13	8	11	5	5	2	1	2	-	-	-

Origin not stated 2/, both sexes

2003.....	7	11	5	5	3	1	-	2	-	-	-	5
2002.....	18	10	11	11	3	3	2	2	2	2	-	4
2001 3/.....	9	11	8	8	4	2	3	3	-	-	-	8
2000.....	11	8	8	5	5	1	2	2	1	1	-	2
1999.....	16	5	6	8	2	5	3	3	-	-	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Origin not stated 2/, male

2003.....	5	10	4	3	2	1	-	2	-	-	-	5
2002.....	14	8	10	10	3	3	1	2	2	2	-	4
2001 3/.....	9	10	7	7	4	2	3	3	-	-	-	7
2000.....	9	8	7	5	4	1	2	1	1	-	-	2
1999.....	12	4	5	8	2	4	3	2	-	-	-	5

Origin not stated 2/, female

2003.....	2	1	1	2	1	-	-	-	-	-	-	-
2002.....	4	2	1	1	-	-	1	-	-	-	-	-
2001 3/.....	-	1	1	1	-	-	-	-	-	-	-	1
2000.....	2	-	1	-	1	-	-	1	-	1	-	-
1999.....	4	1	1	-	-	1	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins, both sexes

2003.....	1,574	1,329	988	833	867	886	709	399	136	24	-	-
2002.....	1,484	1,295	939	846	930	918	762	392	133	25	-	2
2001.....	1,471	1,195	888	857	901	920	728	382	135	18	2	6
2000.....	1,367	1,001	832	791	909	917	694	410	129	18	1	5
1999.....	1,215	1,058	802	827	964	946	660	399	102	23	-	5

All origins, male

2003.....	1,349	1,122	879	747	791	815	665	375	132	22	-	-
2002.....	1,283	1,115	815	753	867	854	722	374	127	24	-	2
2001.....	1,242	1,020	751	761	831	846	685	360	126	18	2	6
2000.....	1,146	854	714	698	830	856	649	388	119	18	1	5
1999.....	1,030	882	712	742	873	875	624	376	99	21	-	5

All origins, female

2003.....	225	207	109	86	76	71	44	24	4	2	-	-
2002.....	201	180	124	93	63	64	40	18	6	1	-	-
2001.....	229	175	137	96	70	74	43	22	9	-	-	-
2000.....	221	147	118	93	79	61	45	22	10	-	-	-
1999.....	185	176	90	85	91	71	36	23	3	2	-	-

Hispanic, both sexes

2003.....	46	32	25	18	20	17	9	9	2	-	-	-
2002.....	39	40	27	26	26	14	14	9	2	1	-	-
2001.....	48	32	27	24	21	12	14	4	3	1	-	1
2000.....	52	28	19	24	25	16	11	7	1	1	-	1
1999.....	32	29	19	24	13	24	15	5	5	1	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Hispanic, male

2003.....	44	30	25	18	20	16	9	8	2	-	-	-
2002.....	37	37	25	24	25	14	14	9	2	1	-	-
2001.....	42	32	25	23	21	12	14	4	3	1	-	1
2000.....	45	27	19	23	23	15	11	7	1	1	-	1
1999.....	26	26	19	22	13	24	14	5	5	1	-	1

Hispanic, female

2003.....	2	2	-	-	-	1	-	1	-	-	-	-
2002.....	2	3	2	2	1	-	-	-	-	-	-	-
2001.....	6	-	2	1	-	-	-	-	-	-	-	-
2000.....	7	1	-	1	2	1	-	-	-	-	-	-
1999.....	6	3	-	2	-	-	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,525	1,293	961	812	845	869	700	388	134	24	-	-
2002.....	1,439	1,251	908	813	903	902	747	382	130	24	-	1
2001.....	1,421	1,157	854	829	877	906	711	375	132	17	2	4
2000.....	1,310	971	808	764	881	900	683	402	127	17	1	3
1999.....	1,178	1,028	778	800	949	918	645	392	97	22	-	3

Non-Hispanic 1/, male

2003.....	1,302	1,088	852	727	770	799	656	365	130	22	-	-
2002.....	1,240	1,075	786	722	841	838	707	364	124	23	-	1
2001.....	1,198	982	719	734	807	832	668	353	123	17	2	4
2000.....	1,096	825	690	672	804	840	638	381	117	17	1	3
1999.....	1,000	855	688	717	858	847	610	370	94	20	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic 1/, female

2003.....	223	205	109	85	75	70	44	23	4	2	-	-
2002.....	199	176	122	91	62	64	40	18	6	1	-	-
2001.....	223	175	135	95	70	74	43	22	9	-	-	-
2000.....	214	146	118	92	77	60	45	21	10	-	-	-
1999.....	178	173	90	83	91	71	35	22	3	2	-	-

Non-Hispanic White, both sexes

2003.....	1,440	1,240	918	776	813	841	679	381	134	23	-	-
2002.....	1,353	1,195	868	784	869	862	733	371	128	22	-	1
2001.....	1,347	1,101	818	792	836	876	684	367	128	17	2	3
2000.....	1,237	912	767	732	840	877	668	395	124	16	1	2
1999.....	1,118	970	733	758	914	889	625	383	95	21	-	3

Non-Hispanic White, male

2003.....	1,230	1,045	812	697	741	773	638	358	130	22	-	-
2002.....	1,168	1,023	751	695	812	800	693	353	122	21	-	1
2001.....	1,137	933	686	704	770	805	646	346	119	17	2	3
2000.....	1,037	772	655	641	764	820	626	374	114	16	1	2
1999.....	945	802	649	682	823	821	591	362	92	19	-	3

Non-Hispanic White, female

2003.....	210	195	106	79	72	68	41	23	4	1	-	-
2002.....	185	172	117	89	57	62	40	18	6	1	-	-
2001.....	210	168	132	88	66	71	38	21	9	-	-	-
2000.....	200	140	112	91	76	57	42	21	10	-	-	-
1999.....	173	168	84	76	91	68	34	21	3	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Non-Hispanic Black, both sexes

2003.....	67	39	36	32	24	22	16	6	-	1	-	-
2002.....	57	39	32	22	32	31	10	11	2	2	-	-
2001.....	60	41	26	27	35	25	22	6	4	-	-	-
2000.....	58	38	33	28	34	19	13	6	2	-	-	-
1999.....	42	46	35	36	29	28	15	9	2	1	-	-

Non-Hispanic Black, male

2003.....	57	30	33	26	21	20	14	6	-	-	-	-
2002.....	48	36	27	20	28	29	10	11	2	2	-	-
2001.....	50	37	24	22	31	22	18	5	4	-	-	-
2000.....	45	34	27	27	33	17	10	6	2	-	-	-
1999.....	39	42	31	30	29	25	14	8	2	1	-	-

Non-Hispanic Black, female

2003.....	10	9	3	6	3	2	2	-	-	1	-	-
2002.....	9	3	5	2	4	2	-	-	-	-	-	-
2001.....	10	4	2	5	4	3	4	1	-	-	-	-
2000.....	13	4	6	1	1	2	3	-	-	-	-	-
1999.....	3	4	4	6	-	3	1	1	-	-	-	-

Origin not stated 2/, both sexes

2003.....	3	4	2	3	2	-	-	2	-	-	-	-
2002.....	6	4	4	7	1	2	1	1	1	-	-	1
2001.....	2	6	7	4	3	2	3	3	-	-	-	1
2000.....	5	2	5	3	3	1	-	1	1	-	-	1
1999.....	5	1	5	3	2	4	-	2	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Origin not stated 2/, male

2003.....	3	4	2	2	1	-	-	2	-	-	-	-
2002.....	6	3	4	7	1	2	1	1	1	-	-	1
2001.....	2	6	7	4	3	2	3	3	-	-	-	1
2000.....	5	2	5	3	3	1	-	-	1	-	-	1
1999.....	4	1	5	3	2	4	-	1	-	-	-	1

Origin not stated 2/, female

2003.....	-	-	-	1	1	-	-	-	-	-	-	-
2002.....	-	1	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	1	-	-	-	-
1999.....	1	-	-	-	-	-	-	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

All origins, both sexes

2003.....	1,422	985	541	334	301	273	247	159	69	10	1	7
2002.....	1,351	891	493	351	336	313	266	181	78	16	1	8
2001 3/.....	1,211	790	444	355	319	299	245	156	65	10	1	9
2000.....	1,083	707	405	282	310	310	260	180	76	16	3	2
1999.....	1,049	620	416	360	320	334	277	181	80	16	-	11

All origins, male

2003.....	922	649	337	211	205	170	159	109	46	6	-	7
2002.....	905	593	314	222	211	200	162	118	52	9	-	8
2001 3/.....	778	514	278	240	208	187	167	108	42	7	1	8
2000.....	696	452	245	172	188	206	173	119	49	8	3	2
1999.....	672	402	259	230	206	198	171	125	51	9	-	11

All origins, female

2003.....	500	336	204	123	96	103	88	50	23	4	1	-
2002.....	446	298	179	129	125	113	104	63	26	7	1	-
2001 3/.....	433	276	166	115	111	112	78	48	23	3	-	1
2000.....	387	255	160	110	122	104	87	61	27	8	-	-
1999.....	377	218	157	130	114	136	106	56	29	7	-	-

Hispanic, both sexes

2003.....	68	56	32	26	16	12	13	6	3	1	-	-
2002.....	72	50	25	19	18	16	11	8	2	1	-	2
2001 3/.....	61	33	22	18	23	12	11	10	3	1	-	1
2000.....	51	34	21	19	16	16	16	4	2	1	-	-
1999.....	57	36	28	22	14	11	13	7	2	2	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Hispanic, male

2003.....	46	48	26	20	13	10	12	5	1	1	-	-
2002.....	54	42	19	15	13	11	8	7	2	-	-	2
2001 3/.....	44	24	19	16	17	11	8	10	3	1	-	1
2000.....	37	27	16	16	12	12	15	3	2	1	-	-
1999.....	39	29	20	19	9	6	10	5	1	1	-	1

Hispanic, female

2003.....	22	8	6	6	3	2	1	1	2	-	-	-
2002.....	18	8	6	4	5	5	3	1	-	1	-	-
2001 3/.....	17	9	3	2	6	1	3	-	-	-	-	-
2000.....	14	7	5	3	4	4	1	1	-	-	-	-
1999.....	18	7	8	3	5	5	3	2	1	1	-	-

Non-Hispanic 1/, both sexes

2003.....	1,350	922	506	306	284	260	234	153	66	9	1	2
2002.....	1,267	835	461	328	316	296	254	172	75	13	1	3
2001 3/.....	1,143	752	421	333	295	287	234	146	62	9	1	1
2000.....	1,026	667	381	261	292	294	242	175	74	14	3	1
1999.....	981	580	387	333	306	322	261	173	78	14	-	6

Non-Hispanic 1/, male

2003.....	874	595	309	190	191	159	147	104	45	5	-	2
2002.....	843	546	289	204	196	188	154	110	49	7	-	3
2001 3/.....	727	486	259	221	190	176	159	98	39	6	1	1
2000.....	655	419	227	154	175	194	156	115	47	7	3	1
1999.....	625	370	239	206	197	192	158	119	50	8	-	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic 1/, female

2003.....	476	327	197	116	93	101	87	49	21	4	1	-
2002.....	424	289	172	124	120	108	100	62	26	6	1	-
2001 3/.....	416	266	162	112	105	111	75	48	23	3	-	-
2000.....	371	248	154	107	117	100	86	60	27	7	-	-
1999.....	356	210	148	127	109	130	103	54	28	6	-	-

Non-Hispanic White, both sexes

2003.....	1,252	845	459	267	254	238	222	137	65	8	1	2
2002.....	1,166	770	421	290	285	269	241	162	69	12	1	3
2001 3/.....	1,048	686	374	293	266	268	221	134	61	8	1	1
2000.....	926	624	356	237	266	264	219	166	70	13	1	1
1999.....	901	531	346	292	285	297	247	165	76	14	-	3

Non-Hispanic White, male

2003.....	818	543	278	163	168	146	139	95	44	5	-	2
2002.....	788	501	271	175	183	174	145	105	45	6	-	3
2001 3/.....	664	442	223	195	167	161	151	92	39	6	1	1
2000.....	582	393	211	138	158	172	143	110	44	6	1	1
1999.....	567	339	211	180	184	176	152	113	50	8	-	3

Non-Hispanic White, female

2003.....	434	302	181	104	86	92	83	42	21	3	1	-
2002.....	378	269	150	115	102	95	96	57	24	6	1	-
2001 3/.....	384	244	151	98	99	107	70	42	22	2	-	-
2000.....	344	231	145	99	108	92	76	56	26	7	-	-
1999.....	334	192	135	112	101	121	95	52	26	6	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Non-Hispanic Black, both sexes

2003.....	59	34	23	11	10	3	2	5	-	-	-	-
2002.....	59	29	25	11	15	7	3	4	-	-	-	-
2001 3/.....	56	37	22	15	13	4	1	2	-	1	-	-
2000.....	58	22	8	11	12	11	10	3	-	-	1	-
1999.....	40	21	14	22	7	7	2	1	2	-	-	2

Non-Hispanic Black, male

2003.....	34	24	15	6	6	3	2	2	-	-	-	-
2002.....	35	21	13	11	7	4	3	2	-	-	-	-
2001 3/.....	37	25	16	9	11	3	1	-	-	-	-	-
2000.....	42	15	4	9	7	8	5	2	-	-	1	-
1999.....	30	12	10	17	2	5	1	1	-	-	-	2

Non-Hispanic Black, female

2003.....	25	10	8	5	4	-	-	3	-	-	-	-
2002.....	24	8	12	-	8	3	-	2	-	-	-	-
2001 3/.....	19	12	6	6	2	1	-	2	-	1	-	-
2000.....	16	7	4	2	5	3	5	1	-	-	-	-
1999.....	10	9	4	5	5	2	1	-	2	-	-	-

Origin not stated 2/, both sexes

2003.....	4	7	3	2	1	1	-	-	-	-	-	5
2002.....	12	6	7	4	2	1	1	1	1	2	-	3
2001 3/.....	7	5	1	4	1	-	-	-	-	-	-	7
2000.....	6	6	3	2	2	-	2	1	-	1	-	1
1999.....	11	4	1	5	-	1	3	1	-	-	-	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Origin not stated 2/, male

2003.....	2	6	2	1	1	1	-	-	-	-	-	5
2002.....	8	5	6	3	2	1	-	1	1	2	-	3
2001 3/.....	7	4	-	3	1	-	-	-	-	-	-	6
2000.....	4	6	2	2	1	-	2	1	-	-	-	1
1999.....	8	3	-	5	-	-	3	1	-	-	-	4

Origin not stated 2/, female

2003.....	2	1	1	1	-	-	-	-	-	-	-	-
2002.....	4	1	1	1	-	-	1	-	-	-	-	-
2001 3/.....	-	1	1	1	-	-	-	-	-	-	-	1
2000.....	2	-	1	-	1	-	-	-	-	1	-	-
1999.....	3	1	1	-	-	1	-	-	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

All origins, both sexes

2003.....	798	490	296	227	205	187	130	77	22	5	-	41
2002.....	760	523	318	229	192	169	127	64	24	6	1	41
2001 3/.....	1,004	628	390	310	222	183	129	71	29	4	1	32
2000.....	696	455	283	233	210	172	129	68	23	8	1	30
1999.....	670	420	302	259	214	185	119	65	24	10	-	49

All origins, male

2003.....	595	352	216	156	129	110	61	32	13	1	-	29
2002.....	553	383	231	152	125	100	58	32	6	3	-	32
2001 3/.....	747	461	262	204	128	98	57	33	7	1	1	23
2000.....	511	338	199	147	132	85	69	25	10	5	-	15
1999.....	483	323	218	183	121	101	61	28	8	5	-	33

All origins, female

2003.....	203	138	80	71	76	77	69	45	9	4	-	12
2002.....	207	140	87	77	67	69	69	32	18	3	1	9
2001 3/.....	257	167	128	106	94	85	72	38	22	3	-	9
2000.....	185	117	84	86	78	87	60	43	13	3	1	15
1999.....	187	97	84	76	93	84	58	37	16	5	-	16

Hispanic, both sexes

2003.....	88	37	38	24	22	15	8	3	2	-	-	6
2002.....	87	56	31	19	10	8	11	2	-	1	-	11
2001 3/.....	92	46	36	31	14	7	9	2	1	-	-	8
2000.....	76	50	27	15	21	12	4	3	-	1	-	3
1999.....	64	47	22	20	13	6	5	2	-	1	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Hispanic, male

2003.....	71	25	34	20	17	8	5	-	2	-	-	5
2002.....	62	47	25	16	9	4	5	1	-	1	-	10
2001 3/.....	69	33	26	24	9	4	4	1	1	-	-	5
2000.....	60	40	17	10	12	7	1	1	-	1	-	2
1999.....	50	37	19	17	9	5	3	1	-	1	-	5

Hispanic, female

2003.....	17	12	4	4	5	7	3	3	-	-	-	1
2002.....	25	9	6	3	1	4	6	1	-	-	-	1
2001 3/.....	23	13	10	7	5	3	5	1	-	-	-	3
2000.....	16	10	10	5	9	5	3	2	-	-	-	1
1999.....	14	10	3	3	4	1	2	1	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	705	447	258	201	180	170	121	73	19	5	-	19
2002.....	666	463	284	210	178	159	114	62	24	5	1	12
2001 3/.....	899	571	345	274	206	176	120	69	28	3	1	11
2000.....	613	400	252	213	189	160	125	65	23	6	1	12
1999.....	594	367	275	236	201	178	114	63	24	8	-	16

Non-Hispanic 1/, male

2003.....	520	321	182	134	109	100	56	31	10	1	-	13
2002.....	485	333	203	136	112	95	53	31	6	2	-	9
2001 3/.....	667	422	229	176	118	94	53	32	6	-	1	9
2000.....	446	295	178	134	120	78	68	24	10	3	-	5
1999.....	422	282	194	163	112	95	58	27	8	4	-	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Non-Hispanic 1/, female

2003.....	185	126	76	67	71	70	65	42	9	4	-	6
2002.....	181	130	81	74	66	64	61	31	18	3	1	3
2001 3/.....	232	149	116	98	88	82	67	37	22	3	-	2
2000.....	167	105	74	79	69	82	57	41	13	3	1	7
1999.....	172	85	81	73	89	83	56	36	16	4	-	6

Non-Hispanic White, both sexes

2003.....	404	289	166	134	120	128	92	62	15	3	-	16
2002.....	365	309	187	141	134	110	87	51	18	4	-	10
2001 3/.....	581	408	244	191	151	132	93	61	21	3	-	7
2000.....	355	257	158	132	132	126	85	54	18	5	1	8
1999.....	359	236	182	172	147	126	88	49	21	6	-	8

Non-Hispanic White, male

2003.....	283	189	114	83	70	74	41	27	8	1	-	11
2002.....	247	205	127	89	77	62	37	27	3	2	-	7
2001 3/.....	420	292	162	116	81	69	41	27	4	-	-	5
2000.....	235	183	105	75	78	60	41	19	7	2	-	4
1999.....	251	171	122	111	77	64	45	18	6	3	-	3

Non-Hispanic White, female

2003.....	121	100	52	51	50	54	51	35	7	2	-	5
2002.....	118	104	60	52	57	48	50	24	15	2	-	3
2001 3/.....	161	116	82	75	70	63	52	34	17	3	-	2
2000.....	120	74	53	57	54	66	44	35	11	3	1	4
1999.....	108	65	60	61	70	62	43	31	15	3	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Non-Hispanic Black, both sexes

2003.....	271	139	79	54	51	37	26	10	4	2	-	2
2002.....	268	130	86	55	40	47	22	11	5	1	-	2
2001 3/.....	271	138	81	69	48	40	23	7	6	-	1	4
2000.....	225	121	80	72	51	30	34	8	5	1	-	4
1999.....	193	115	79	59	44	46	21	13	3	2	-	8

Non-Hispanic Black, male

2003.....	213	120	58	44	34	25	14	4	2	-	-	1
2002.....	213	112	68	40	32	32	15	4	2	-	-	2
2001 3/.....	211	109	56	50	35	25	9	4	2	-	1	4
2000.....	188	96	63	56	37	16	23	3	3	1	-	1
1999.....	146	99	60	49	30	29	10	8	2	1	-	7

Non-Hispanic Black, female

2003.....	58	19	21	10	17	12	12	6	2	2	-	1
2002.....	55	18	18	15	8	15	7	7	3	1	-	-
2001 3/.....	60	29	25	19	13	15	14	3	4	-	-	-
2000.....	37	25	17	16	14	14	11	5	2	-	-	3
1999.....	47	16	19	10	14	17	11	5	1	1	-	1

Origin not stated 2/, both sexes

2003.....	5	6	-	2	3	2	1	1	1	-	-	16
2002.....	7	4	3	-	4	2	2	-	-	-	-	18
2001 3/.....	13	11	9	5	2	-	-	-	-	1	-	13
2000.....	7	5	4	5	-	-	-	-	-	1	-	15
1999.....	12	6	5	3	-	1	-	-	-	1	-	28

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Origin not stated 2/, male

2003.....	4	6	-	2	3	2	-	1	1	-	-	11
2002.....	6	3	3	-	4	1	-	-	-	-	-	13
2001 3/.....	11	6	7	4	1	-	-	-	-	1	-	9
2000.....	5	3	4	3	-	-	-	-	-	1	-	8
1999.....	11	4	5	3	-	1	-	-	-	-	-	18

Origin not stated 2/, female

2003.....	1	-	-	-	-	-	1	-	-	-	-	5
2002.....	1	1	-	-	-	1	2	-	-	-	-	5
2001 3/.....	2	5	2	1	1	-	-	-	-	-	-	4
2000.....	2	2	-	2	-	-	-	-	-	-	-	7
1999.....	1	2	-	-	-	-	-	-	-	1	-	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

All origins, both sexes

2003.....	427	250	144	83	78	55	33	17	5	1	-	18
2002.....	421	263	146	92	69	72	43	14	4	1	-	13
2001.....	382	213	151	114	78	61	38	12	2	2	-	11
2000.....	377	235	132	99	85	54	27	23	4	1	-	12
1999.....	342	206	167	102	91	61	34	17	4	2	-	15

All origins, male

2003.....	326	186	106	63	52	27	8	9	1	-	-	16
2002.....	316	197	104	61	40	44	20	8	-	1	-	12
2001.....	292	159	105	79	44	38	16	6	-	1	-	11
2000.....	285	172	94	58	50	23	14	6	2	1	-	7
1999.....	250	167	127	68	56	34	14	7	2	1	-	14

All origins, female

2003.....	101	64	38	20	26	28	25	8	4	1	-	2
2002.....	105	66	42	31	29	28	23	6	4	-	-	1
2001.....	90	54	46	35	34	23	22	6	2	1	-	-
2000.....	92	63	38	41	35	31	13	17	2	-	-	5
1999.....	92	39	40	34	35	27	20	10	2	1	-	1

Hispanic, both sexes

2003.....	44	10	16	9	8	3	-	-	-	-	-	4
2002.....	51	21	10	7	2	1	3	-	-	1	-	4
2001.....	36	14	6	11	3	-	3	1	-	-	-	3
2000.....	46	22	12	5	9	1	-	-	-	-	-	2
1999.....	32	23	12	7	2	3	2	-	-	1	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Hispanic, male

2003.....	32	9	13	8	5	2	-	-	-	-	-	3
2002.....	38	18	8	5	2	1	2	-	-	1	-	4
2001.....	28	11	4	9	1	-	-	1	-	-	-	3
2000.....	37	19	7	2	5	-	-	-	-	-	-	1
1999.....	26	18	9	4	2	3	1	-	-	1	-	2

Hispanic, female

2003.....	12	1	3	1	3	1	-	-	-	-	-	1
2002.....	13	3	2	2	-	-	1	-	-	-	-	-
2001.....	8	3	2	2	2	-	3	-	-	-	-	-
2000.....	9	3	5	3	4	1	-	-	-	-	-	1
1999.....	6	5	3	3	-	-	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	383	237	128	74	70	51	33	17	5	1	-	9
2002.....	367	241	135	85	66	70	40	14	4	-	-	6
2001.....	344	198	143	101	74	61	35	11	2	1	-	3
2000.....	329	212	118	92	76	53	27	23	4	1	-	3
1999.....	307	181	154	95	89	57	32	17	4	1	-	4

Non-Hispanic 1/, male

2003.....	294	174	93	55	47	24	8	9	1	-	-	8
2002.....	276	179	95	56	37	42	18	8	-	-	-	5
2001.....	262	147	99	68	42	38	16	5	-	-	-	3
2000.....	247	152	85	56	45	23	14	6	2	1	-	2
1999.....	222	147	117	64	54	30	13	7	2	-	-	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Non-Hispanic 1/, female

2003.....	89	63	35	19	23	27	25	8	4	1	-	1
2002.....	91	62	40	29	29	28	22	6	4	-	-	1
2001.....	82	51	44	33	32	23	19	6	2	1	-	-
2000.....	82	60	33	36	31	30	13	17	2	-	-	1
1999.....	85	34	37	31	35	27	19	10	2	1	-	-

Non-Hispanic White, both sexes

2003.....	222	157	85	49	50	38	27	17	3	-	-	6
2002.....	203	170	83	55	52	54	33	13	3	-	-	4
2001.....	200	129	106	73	53	47	28	11	2	1	-	2
2000.....	186	143	81	63	55	46	17	19	4	1	-	2
1999.....	179	115	100	73	67	39	28	16	4	-	-	-

Non-Hispanic White, male

2003.....	165	106	60	35	33	18	5	9	-	-	-	6
2002.....	146	117	57	33	25	31	13	8	-	-	-	3
2001.....	145	84	71	42	27	27	10	5	-	-	-	2
2000.....	126	98	51	35	28	20	5	5	2	1	-	1
1999.....	134	87	70	46	38	16	11	6	2	-	-	-

Non-Hispanic White, female

2003.....	57	51	25	14	17	20	22	8	3	-	-	-
2002.....	57	53	26	22	27	23	20	5	3	-	-	1
2001.....	55	45	35	31	26	20	18	6	2	1	-	-
2000.....	60	45	30	28	27	26	12	14	2	-	-	1
1999.....	45	28	30	27	29	23	17	10	2	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Non-Hispanic Black, both sexes

2003.....	147	70	36	23	19	10	6	-	2	1	-	2
2002.....	146	61	46	26	13	14	6	1	1	-	-	2
2001.....	129	60	32	25	18	13	5	-	-	-	-	1
2000.....	126	56	29	25	20	7	10	4	-	-	-	1
1999.....	104	59	45	20	19	17	4	1	-	1	-	4

Non-Hispanic Black, male

2003.....	118	63	26	19	14	6	3	-	1	-	-	1
2002.....	115	55	33	21	11	10	5	-	-	-	-	2
2001.....	108	54	24	23	14	11	4	-	-	-	-	1
2000.....	109	43	26	20	17	3	9	1	-	-	-	1
1999.....	77	55	39	17	14	14	2	1	-	-	-	4

Non-Hispanic Black, female

2003.....	29	7	10	4	5	4	3	-	1	1	-	1
2002.....	31	6	13	5	2	4	1	1	1	-	-	-
2001.....	21	6	8	2	4	2	1	-	-	-	-	-
2000.....	17	13	3	5	3	4	1	3	-	-	-	-
1999.....	27	4	6	3	5	3	2	-	-	1	-	-

Origin not stated 2/, both sexes

2003.....	-	3	-	-	-	1	-	-	-	-	-	5
2002.....	3	1	1	-	1	1	-	-	-	-	-	3
2001.....	2	1	2	2	1	-	-	-	-	1	-	5
2000.....	2	1	2	2	-	-	-	-	-	-	-	7
1999.....	3	2	1	-	-	1	-	-	-	-	-	9



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02.X85-X92,X96-Y09,Y87.1)

All origins, both sexes

2003.....	371	240	152	144	127	132	97	60	17	4	-	23
2002.....	339	260	172	137	123	97	84	50	20	5	1	28
2001 3/.....	622	415	239	196	144	122	91	59	27	2	1	21
2000.....	319	220	151	134	125	118	102	45	19	7	1	18
1999.....	328	214	135	157	123	124	85	48	20	8	-	34

All origins, male

2003.....	269	166	110	93	77	83	53	23	12	1	-	13
2002.....	237	186	127	91	85	56	38	24	6	2	-	20
2001 3/.....	455	302	157	125	84	60	41	27	7	-	1	12
2000.....	226	166	105	89	82	62	55	19	8	4	-	8
1999.....	233	156	91	115	65	67	47	21	6	4	-	19

All origins, female

2003.....	102	74	42	51	50	49	44	37	5	3	-	10
2002.....	102	74	45	46	38	41	46	26	14	3	1	8
2001 3/.....	167	113	82	71	60	62	50	32	20	2	-	9
2000.....	93	54	46	45	43	56	47	26	11	3	1	10
1999.....	95	58	44	42	58	57	38	27	14	4	-	15

Hispanic, both sexes

2003.....	44	27	22	15	14	12	8	3	2	-	-	2
2002.....	36	35	21	12	8	7	8	2	-	-	-	7
2001 3/.....	56	32	30	20	11	7	6	1	1	-	-	5
2000.....	30	28	15	10	12	11	4	3	-	1	-	1
1999.....	32	24	10	13	11	3	3	2	-	-	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02.X85-X92,X96-Y09,Y87.1)

Hispanic, male

2003.....	39	16	21	12	12	6	5	-	2	-	-	2
2002.....	24	29	17	11	7	3	3	1	-	-	-	6
2001 3/.....	41	22	22	15	8	4	4	-	1	-	-	2
2000.....	23	21	10	8	7	7	1	1	-	1	-	1
1999.....	24	19	10	13	7	2	2	1	-	-	-	3

Hispanic, female

2003.....	5	11	1	3	2	6	3	3	-	-	-	-
2002.....	12	6	4	1	1	4	5	1	-	-	-	1
2001 3/.....	15	10	8	5	3	3	2	1	-	-	-	3
2000.....	7	7	5	2	5	4	3	2	-	-	-	-
1999.....	8	5	-	-	4	1	1	1	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	322	210	130	127	110	119	88	56	14	4	-	10
2002.....	299	222	149	125	112	89	74	48	20	5	1	6
2001 3/.....	555	373	202	173	132	115	85	58	26	2	1	8
2000.....	284	188	134	121	113	107	98	42	19	5	1	9
1999.....	287	186	121	141	112	121	82	46	20	7	-	12

Non-Hispanic 1/, male

2003.....	226	147	89	79	62	76	48	22	9	1	-	5
2002.....	209	154	108	80	75	53	35	23	6	2	-	4
2001 3/.....	405	275	130	108	76	56	37	27	6	-	1	6
2000.....	199	143	93	78	75	55	54	18	8	2	-	3
1999.....	200	135	77	99	58	65	45	20	6	4	-	6

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02.X85-X92,X96-Y09,Y87.1)

Non-Hispanic 1/, female

2003.....	96	63	41	48	48	43	40	34	5	3	-	5
2002.....	90	68	41	45	37	36	39	25	14	3	1	2
2001 3/.....	150	98	72	65	56	59	48	31	20	2	-	2
2000.....	85	45	41	43	38	52	44	24	11	3	1	6
1999.....	87	51	44	42	54	56	37	26	14	3	-	6

Non-Hispanic White, both sexes

2003.....	182	132	81	85	70	90	65	45	12	3	-	10
2002.....	162	139	104	86	82	56	54	38	15	4	-	6
2001 3/.....	381	279	138	118	98	85	65	50	19	2	-	5
2000.....	169	114	77	69	77	80	68	35	14	4	1	6
1999.....	180	121	82	99	80	87	60	33	17	6	-	8

Non-Hispanic White, male

2003.....	118	83	54	48	37	56	36	18	8	1	-	5
2002.....	101	88	70	56	52	31	24	19	3	2	-	4
2001 3/.....	275	208	91	74	54	42	31	22	4	-	-	3
2000.....	109	85	54	40	50	40	36	14	5	1	-	3
1999.....	117	84	52	65	39	48	34	12	4	3	-	3

Non-Hispanic White, female

2003.....	64	49	27	37	33	34	29	27	4	2	-	5
2002.....	61	51	34	30	30	25	30	19	12	2	-	2
2001 3/.....	106	71	47	44	44	43	34	28	15	2	-	2
2000.....	60	29	23	29	27	40	32	21	9	3	1	3
1999.....	63	37	30	34	41	39	26	21	13	3	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Non-Hispanic Black, both sexes

2003.....	124	69	43	31	32	27	20	10	2	1	-	-
2002.....	122	69	40	29	27	33	16	10	4	1	-	-
2001 3/.....	142	78	49	44	30	27	18	7	6	-	1	3
2000.....	99	65	51	47	31	23	24	4	5	1	-	3
1999.....	89	56	34	39	25	29	17	12	3	1	-	4

Non-Hispanic Black, male

2003.....	95	57	32	25	20	19	11	4	1	-	-	-
2002.....	98	57	35	19	21	22	10	4	2	-	-	-
2001 3/.....	103	55	32	27	21	14	5	4	2	-	1	3
2000.....	79	53	37	36	20	13	14	2	3	1	-	-
1999.....	69	44	21	32	16	15	8	7	2	1	-	3

Non-Hispanic Black, female

2003.....	29	12	11	6	12	8	9	6	1	1	-	-
2002.....	24	12	5	10	6	11	6	6	2	1	-	-
2001 3/.....	39	23	17	17	9	13	13	3	4	-	-	-
2000.....	20	12	14	11	11	10	10	2	2	-	-	3
1999.....	20	12	13	7	9	14	9	5	1	-	-	1

Origin not stated 2/, both sexes

2003.....	5	3	-	2	3	1	1	1	1	-	-	11
2002.....	4	3	2	-	3	1	2	-	-	-	-	15
2001 3/.....	11	10	7	3	1	-	-	-	-	-	-	8
2000.....	5	4	2	3	-	-	-	-	-	1	-	8
1999.....	9	4	4	3	-	-	-	-	-	1	-	19

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Origin not stated 2/, male

2003.....	4	3	-	2	3	1	-	1	1	-	-	6
2002.....	4	3	2	-	3	-	-	-	-	-	-	10
2001 3/.....	9	5	5	2	-	-	-	-	-	-	-	4
2000.....	4	2	2	3	-	-	-	-	-	1	-	4
1999.....	9	2	4	3	-	-	-	-	-	-	-	10

Origin not stated 2/, female

2003.....	1	-	-	-	-	-	1	-	-	-	-	5
2002.....	-	-	-	-	-	1	2	-	-	-	-	5
2001 3/.....	2	5	2	1	1	-	-	-	-	-	-	4
2000.....	1	2	-	-	-	-	-	-	-	-	-	4
1999.....	-	2	-	-	-	-	-	-	-	1	-	9



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Legal intervention (Y35,Y89.0)

Hispanic, male

2003.....	1	-	-	1	-	-	-	-	-	-	-	-
2002.....	5	1	-	1	-	-	-	-	-	-	-	-
2001.....	2	-	1	-	-	-	-	-	-	-	-	-
2000.....	2	-	1	-	-	-	-	-	-	-	-	-
1999.....	4	-	2	1	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	21	11	7	1	2	2	-	-	-	-	-	-
2002.....	8	9	6	3	2	-	1	-	-	-	-	-
2001.....	16	15	8	4	2	3	1	1	-	-	-	-
2000.....	15	6	3	2	3	1	-	1	-	-	-	-
1999.....	27	9	9	1	-	1	-	-	-	-	-	-

Non-Hispanic 1/, male

2003.....	18	10	7	1	2	2	-	-	-	-	-	-
2002.....	7	9	6	3	2	-	1	-	-	-	-	-
2001.....	16	15	6	4	2	3	1	1	-	-	-	-
2000.....	15	6	2	2	3	1	-	1	-	-	-	-
1999.....	26	9	9	1	-	1	-	-	-	-	-	-







Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins, both sexes

2003.....	490	245	123	74	57	45	51	36	12	4	-	18
2002.....	431	207	113	80	59	44	51	39	17	8	1	14
2001.....	371	152	73	58	41	53	55	22	11	9	2	18
2000.....	298	154	74	55	49	43	36	26	25	5	1	11
1999.....	265	136	86	72	42	51	45	37	18	2	1	19

All origins, male

2003.....	318	153	69	37	28	26	23	15	4	2	-	14
2002.....	264	131	72	46	37	24	22	20	6	5	-	13
2001.....	229	105	41	40	24	28	29	12	6	2	-	16
2000.....	196	105	47	38	32	29	20	14	5	2	-	11
1999.....	175	94	54	47	25	31	24	20	4	1	-	13

All origins, female

2003.....	172	92	54	37	29	19	28	21	8	2	-	4
2002.....	167	76	41	34	22	20	29	19	11	3	1	1
2001.....	142	47	32	18	17	25	26	10	5	7	2	2
2000.....	102	49	27	17	17	14	16	12	20	3	1	-
1999.....	90	42	32	25	17	20	21	17	14	1	1	6

Hispanic, both sexes

2003.....	26	12	7	2	4	2	3	2	-	-	-	-
2002.....	15	6	9	4	5	1	1	-	-	-	-	2
2001.....	23	5	4	4	2	1	1	2	2	-	-	2
2000.....	6	7	5	4	1	-	2	-	-	-	-	1
1999.....	10	8	5	2	1	2	1	-	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Hispanic, male

2003.....	20	9	4	2	3	1	3	2	-	-	-	-
2002.....	12	5	9	2	2	-	1	-	-	-	-	2
2001.....	17	5	2	3	2	-	1	2	1	-	-	2
2000.....	4	6	3	2	-	-	2	-	-	-	-	1
1999.....	8	7	4	2	-	2	-	-	-	-	-	1

Hispanic, female

2003.....	6	3	3	-	1	1	-	-	-	-	-	-
2002.....	3	1	-	2	3	1	-	-	-	-	-	-
2001.....	6	-	2	1	-	1	-	-	1	-	-	-
2000.....	2	1	2	2	1	-	-	-	-	-	-	-
1999.....	2	1	1	-	1	-	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	463	231	115	72	52	43	48	34	12	4	-	8
2002.....	415	198	103	75	53	43	50	38	17	6	1	5
2001.....	341	143	67	53	38	51	54	20	9	9	2	6
2000.....	290	143	68	49	47	43	34	26	24	5	1	4
1999.....	250	126	81	68	39	49	44	36	18	1	1	10

Non-Hispanic 1/, male

2003.....	297	142	64	35	24	25	20	13	4	2	-	7
2002.....	251	123	62	43	34	24	21	19	6	3	-	4
2001.....	208	96	38	36	21	27	28	10	5	2	-	5
2000.....	190	97	43	34	31	29	18	14	4	2	-	4
1999.....	163	85	50	44	23	29	24	20	4	-	-	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic 1/, female

2003.....	166	89	51	37	28	18	28	21	8	2	-	1
2002.....	164	75	41	32	19	19	29	19	11	3	1	1
2001.....	133	47	29	17	17	24	26	10	4	7	2	1
2000.....	100	46	25	15	16	14	16	12	20	3	1	-
1999.....	87	41	31	24	16	20	20	16	14	1	1	2

Non-Hispanic White, both sexes

2003.....	376	189	88	57	39	38	42	32	12	4	-	6
2002.....	329	170	83	61	44	36	41	36	16	5	1	4
2001.....	271	121	55	41	34	46	47	18	9	8	2	4
2000.....	232	105	56	39	37	38	32	22	20	5	-	2
1999.....	188	102	70	60	32	45	41	32	16	1	1	6

Non-Hispanic White, male

2003.....	230	109	46	26	19	23	18	12	4	2	-	5
2002.....	183	102	49	32	25	19	19	18	6	3	-	3
2001.....	158	77	29	28	17	25	26	9	5	2	-	3
2000.....	148	66	35	25	23	27	16	11	3	2	-	2
1999.....	116	62	39	39	18	26	21	18	4	-	-	5

Non-Hispanic White, female

2003.....	146	80	42	31	20	15	24	20	8	2	-	1
2002.....	146	68	34	29	19	17	22	18	10	2	1	1
2001.....	113	44	26	13	17	21	21	9	4	6	2	1
2000.....	84	39	21	14	14	11	16	11	17	3	-	-
1999.....	72	40	31	21	14	19	20	14	12	1	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Non-Hispanic Black, both sexes

2003.....	75	31	22	9	12	3	4	2	-	-	-	2
2002.....	79	23	16	9	7	7	9	1	1	1	-	-
2001.....	65	20	10	10	2	2	3	2	-	1	-	1
2000.....	53	31	11	8	7	4	2	3	3	-	1	2
1999.....	59	20	8	7	5	4	1	3	-	-	-	4

Non-Hispanic Black, male

2003.....	58	25	15	8	4	1	1	1	-	-	-	2
2002.....	62	17	13	8	7	5	2	-	-	-	-	-
2001.....	48	18	8	8	2	-	1	1	-	-	-	1
2000.....	38	24	8	7	6	2	2	2	1	-	-	2
1999.....	46	19	8	5	4	3	1	1	-	-	-	3

Non-Hispanic Black, female

2003.....	17	6	7	1	8	2	3	1	-	-	-	-
2002.....	17	6	3	1	-	2	7	1	1	1	-	-
2001.....	17	2	2	2	-	2	2	1	-	1	-	-
2000.....	15	7	3	1	1	2	-	1	2	-	1	-
1999.....	13	1	-	2	1	1	-	2	-	-	-	1

Origin not stated 2/, both sexes

2003.....	1	2	1	-	1	-	-	-	-	-	-	10
2002.....	1	3	1	1	1	-	-	1	-	2	-	7
2001.....	7	4	2	1	1	1	-	-	-	-	-	10
2000.....	2	4	1	2	1	-	-	-	1	-	-	6
1999.....	5	2	-	2	2	-	-	1	-	1	-	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Origin not stated 2/, male

2003.....	1	2	1	-	1	-	-	-	-	-	-	7
2002.....	1	3	1	1	1	-	-	1	-	2	-	7
2001.....	4	4	1	1	1	1	-	-	-	-	-	9
2000.....	2	2	1	2	1	-	-	-	1	-	-	6
1999.....	4	2	-	1	2	-	-	-	-	1	-	4

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	3
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	-	-	-	-	-	-	-	-	1
2000.....	-	2	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	1	-	-	-	4



Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

Hispanic, male

2003.....	-	-	-	-	1	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	1	1	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	1	-	2	-	-	-	-	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	12	14	6	4	3	4	5	1	-	-	-	-
2002.....	10	8	3	8	7	3	2	-	-	-	-	-
2001.....	12	8	3	5	4	6	5	1	-	-	-	-
2000.....	14	12	3	6	4	5	1	2	-	-	-	-
1999.....	10	5	8	7	6	6	7	4	-	-	-	-

Non-Hispanic 1/, male

2003.....	11	12	4	3	3	4	4	1	-	-	-	-
2002.....	9	4	2	4	6	3	1	-	-	-	-	-
2001.....	8	7	3	5	2	6	3	1	-	-	-	-
2000.....	12	10	1	4	4	4	1	2	-	-	-	-
1999.....	6	4	6	6	6	5	7	4	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

Non-Hispanic 1/, female

2003.....	1	2	2	1	-	-	1	-	-	-	-	-
2002.....	1	4	1	4	1	-	1	-	-	-	-	-
2001.....	4	1	-	-	2	-	2	-	-	-	-	-
2000.....	2	2	2	2	-	1	-	-	-	-	-	-
1999.....	4	1	2	1	-	1	-	-	-	-	-	-

Non-Hispanic White, both sexes

2003.....	11	12	5	4	3	4	5	1	-	-	-	-
2002.....	9	7	2	8	6	2	2	-	-	-	-	-
2001.....	10	5	3	5	4	6	4	1	-	-	-	-
2000.....	12	8	3	5	3	5	1	2	-	-	-	-
1999.....	10	4	7	7	4	6	6	4	-	-	-	-

Non-Hispanic White, male

2003.....	10	11	4	3	3	4	4	1	-	-	-	-
2002.....	8	4	2	4	5	2	1	-	-	-	-	-
2001.....	6	4	3	5	2	6	2	1	-	-	-	-
2000.....	10	6	1	3	3	4	1	2	-	-	-	-
1999.....	6	3	5	6	4	5	6	4	-	-	-	-

Non-Hispanic White, female

2003.....	1	1	1	1	-	-	1	-	-	-	-	-
2002.....	1	3	-	4	1	-	1	-	-	-	-	-
2001.....	4	1	-	-	2	-	2	-	-	-	-	-
2000.....	2	2	2	2	-	1	-	-	-	-	-	-
1999.....	4	1	2	1	-	1	-	-	-	-	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins, both sexes

2003.....	478	231	117	70	53	41	46	35	12	4	-	18
2002.....	421	198	110	72	52	41	49	39	17	8	1	14
2001.....	358	143	70	52	37	47	50	21	11	9	2	18
2000.....	284	142	71	48	45	38	35	24	25	5	1	10
1999.....	255	130	78	63	36	45	38	33	18	2	1	19

All origins, male

2003.....	307	141	65	34	24	22	19	14	4	2	-	14
2002.....	255	126	70	42	31	21	21	20	6	5	-	13
2001.....	220	97	38	34	22	22	26	11	6	2	-	16
2000.....	184	95	46	33	28	25	19	12	5	2	-	10
1999.....	169	89	48	39	19	26	17	16	4	1	-	13

All origins, female

2003.....	171	90	52	36	29	19	27	21	8	2	-	4
2002.....	166	72	40	30	21	20	28	19	11	3	1	1
2001.....	138	46	32	18	15	25	24	10	5	7	2	2
2000.....	100	47	25	15	17	13	16	12	20	3	1	-
1999.....	86	41	30	24	17	19	21	17	14	1	1	6

Hispanic, both sexes

2003.....	26	12	7	2	3	2	3	2	-	-	-	-
2002.....	15	6	9	4	5	1	1	-	-	-	-	2
2001.....	22	4	4	4	2	1	1	2	2	-	-	2
2000.....	6	7	5	3	1	-	2	-	-	-	-	1
1999.....	10	7	5	-	1	2	1	-	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Hispanic, male

2003.....	20	9	4	2	2	1	3	2	-	-	-	-
2002.....	12	5	9	2	2	-	1	-	-	-	-	2
2001.....	16	4	2	3	2	-	1	2	1	-	-	2
2000.....	4	6	3	1	-	-	2	-	-	-	-	1
1999.....	8	6	4	-	-	2	-	-	-	-	-	1

Hispanic, female

2003.....	6	3	3	-	1	1	-	-	-	-	-	-
2002.....	3	1	-	2	3	1	-	-	-	-	-	-
2001.....	6	-	2	1	-	1	-	-	1	-	-	-
2000.....	2	1	2	2	1	-	-	-	-	-	-	-
1999.....	2	1	1	-	1	-	1	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	451	217	109	68	49	39	43	33	12	4	-	8
2002.....	405	190	100	67	46	40	48	38	17	6	1	5
2001.....	329	135	64	48	34	45	49	19	9	9	2	6
2000.....	276	131	65	43	43	38	33	24	24	5	1	4
1999.....	240	121	73	61	33	43	37	32	18	1	1	10

Non-Hispanic 1/, male

2003.....	286	130	60	32	21	21	16	12	4	2	-	7
2002.....	242	119	60	39	28	21	20	19	6	3	-	4
2001.....	200	89	35	31	19	21	25	9	5	2	-	5
2000.....	178	87	42	30	27	25	17	12	4	2	-	4
1999.....	157	81	44	38	17	24	17	16	4	-	-	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic 1/, female

2003.....	165	87	49	36	28	18	27	21	8	2	-	1
2002.....	163	71	40	28	18	19	28	19	11	3	1	1
2001.....	129	46	29	17	15	24	24	10	4	7	2	1
2000.....	98	44	23	13	16	13	16	12	20	3	1	-
1999.....	83	40	29	23	16	19	20	16	14	1	1	2

Non-Hispanic White, both sexes

2003.....	365	177	83	53	36	34	37	31	12	4	-	6
2002.....	320	163	81	53	38	34	39	36	16	5	1	4
2001.....	261	116	52	36	30	40	43	17	9	8	2	4
2000.....	220	97	53	34	34	33	31	20	20	5	-	2
1999.....	178	98	63	53	28	39	35	28	16	1	1	6

Non-Hispanic White, male

2003.....	220	98	42	23	16	19	14	11	4	2	-	5
2002.....	175	98	47	28	20	17	18	18	6	3	-	3
2001.....	152	73	26	23	15	19	24	8	5	2	-	3
2000.....	138	60	34	22	20	23	15	9	3	2	-	2
1999.....	110	59	34	33	14	21	15	14	4	-	-	5

Non-Hispanic White, female

2003.....	145	79	41	30	20	15	23	20	8	2	-	1
2002.....	145	65	34	25	18	17	21	18	10	2	1	1
2001.....	109	43	26	13	15	21	19	9	4	6	2	1
2000.....	82	37	19	12	14	10	16	11	17	3	-	-
1999.....	68	39	29	20	14	18	20	14	12	1	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Non-Hispanic Black, both sexes

2003.....	74	30	22	9	12	3	4	2	-	-	-	2
2002.....	78	22	16	9	6	6	9	1	1	1	-	-
2001.....	63	17	10	10	2	2	2	2	-	1	-	1
2000.....	52	28	11	7	6	4	2	3	3	-	1	2
1999.....	59	19	8	7	3	4	-	3	-	-	-	4

Non-Hispanic Black, male

2003.....	57	24	15	8	4	1	1	1	-	-	-	2
2002.....	61	17	13	8	6	4	2	-	-	-	-	-
2001.....	46	15	8	8	2	-	-	1	-	-	-	1
2000.....	37	21	8	6	5	2	2	2	1	-	-	2
1999.....	46	18	8	5	2	3	-	1	-	-	-	3

Non-Hispanic Black, female

2003.....	17	6	7	1	8	2	3	1	-	-	-	-
2002.....	17	5	3	1	-	2	7	1	1	1	-	-
2001.....	17	2	2	2	-	2	2	1	-	1	-	-
2000.....	15	7	3	1	1	2	-	1	2	-	1	-
1999.....	13	1	-	2	1	1	-	2	-	-	-	1

Origin not stated 2/, both sexes

2003.....	1	2	1	-	1	-	-	-	-	-	-	10
2002.....	1	2	1	1	1	-	-	1	-	2	-	7
2001.....	7	4	2	-	1	1	-	-	-	-	-	10
2000.....	2	4	1	2	1	-	-	-	1	-	-	5
1999.....	5	2	-	2	2	-	-	1	-	1	-	8

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Origin not stated 2/, male

2003.....	1	2	1	-	1	-	-	-	-	-	-	7
2002.....	1	2	1	1	1	-	-	1	-	2	-	7
2001.....	4	4	1	-	1	1	-	-	-	-	-	9
2000.....	2	2	1	2	1	-	-	-	1	-	-	5
1999.....	4	2	-	1	2	-	-	-	-	1	-	4

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	3
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	3	-	1	-	-	-	-	-	-	-	-	1
2000.....	-	2	-	-	-	-	-	-	-	-	-	-
1999.....	1	-	-	1	-	-	-	1	-	-	-	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

All origins, both sexes

2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	7	1	1	-	-	-	-
2000.....	2	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	4	2	1	-	-	-

All origins, male

2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	6	1	1	-	-	-	-
2000.....	2	-	1	1	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	3	2	1	-	-	-

All origins, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Hispanic, both sexes

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

Hispanic, male

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	1	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	1	-	-	-	-

Hispanic, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	-	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	6	1	1	-	-	-	-
2000.....	2	-	1	-	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	3	1	1	-	-	-

Non-Hispanic 1/, male

2003.....	1	5	-	-	2	2	3	-	-	-	-	-
2002.....	-	2	1	2	1	4	6	1	-	-	-	-
2001.....	1	-	-	-	4	5	1	1	-	-	-	-
2000.....	2	-	1	-	2	4	5	1	-	-	-	-
1999.....	2	1	-	2	6	4	2	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

Non-Hispanic 1/, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-

Non-Hispanic White, both sexes

2003.....	1	4	-	-	2	2	3	-	-	-	-	-
2002.....	-	1	1	2	-	4	5	1	-	-	-	-
2001.....	1	-	-	-	3	6	1	1	-	-	-	-
2000.....	1	-	1	-	2	4	5	1	-	-	-	-
1999.....	2	1	-	1	6	4	3	1	1	-	-	-

Non-Hispanic White, male

2003.....	1	4	-	-	2	2	3	-	-	-	-	-
2002.....	-	1	1	2	-	4	5	1	-	-	-	-
2001.....	1	-	-	-	3	5	1	1	-	-	-	-
2000.....	1	-	1	-	2	4	5	1	-	-	-	-
1999.....	2	1	-	1	6	4	2	1	1	-	-	-

Non-Hispanic White, female

2003.....	-	-	-	-	-	-	-	-	-	-	-	-
2002.....	-	-	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	1	-	-	-	-	-	-
2000.....	-	-	-	-	-	-	-	-	-	-	-	-
1999.....	-	-	-	-	-	-	1	-	-	-	-	-





Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

All origins, both sexes

2003.....	128	178	202	214	320	398	396	307	152	58	13	1
2002.....	151	164	187	225	335	382	410	288	167	59	7	-
2001.....	141	178	206	247	320	439	425	315	186	56	7	1
2000.....	167	157	219	263	384	427	398	337	176	54	11	-
1999.....	123	138	177	229	357	401	410	312	153	52	12	-

All origins, male

2003.....	66	101	94	103	161	174	181	129	46	8	3	1
2002.....	67	84	85	110	146	166	154	110	40	18	3	-
2001.....	68	91	105	116	149	192	191	119	55	13	2	-
2000.....	84	77	104	123	208	212	190	134	48	13	1	-
1999.....	61	77	88	102	177	193	182	128	49	17	3	-

All origins, female

2003.....	62	77	108	111	159	224	215	178	106	50	10	-
2002.....	84	80	102	115	189	216	256	178	127	41	4	-
2001.....	73	87	101	131	171	247	234	196	131	43	5	1
2000.....	83	80	115	140	176	215	208	203	128	41	10	-
1999.....	62	61	89	127	180	208	228	184	104	35	9	-

Hispanic, both sexes

2003.....	6	7	14	14	24	11	13	7	2	4	-	1
2002.....	10	8	11	14	14	16	13	2	3	-	1	-
2001.....	8	5	8	17	8	18	11	5	8	2	-	1
2000.....	5	4	7	22	13	15	11	7	6	3	-	-
1999.....	6	6	10	8	7	7	5	3	7	1	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Hispanic, male

2003.....	3	4	5	6	11	6	4	4	-	-	-	1
2002.....	5	2	4	5	6	9	4	-	-	-	-	-
2001.....	2	1	7	6	2	8	6	2	4	2	-	-
2000.....	1	4	4	7	9	9	6	4	1	1	-	-
1999.....	4	4	5	4	1	6	2	1	2	1	-	-

Hispanic, female

2003.....	3	3	9	8	13	5	9	3	2	4	-	-
2002.....	5	6	7	9	8	7	9	2	3	-	1	-
2001.....	6	4	1	11	6	10	5	3	4	-	-	1
2000.....	4	-	3	15	4	6	5	3	5	2	-	-
1999.....	2	2	5	4	6	1	3	2	5	-	1	-

Non-Hispanic 1/, both sexes

2003.....	122	171	188	200	296	386	383	299	150	54	13	-
2002.....	140	156	176	209	321	365	393	285	163	59	6	-
2001.....	133	173	196	228	309	419	412	310	177	54	7	-
2000.....	161	153	210	241	369	411	386	329	170	51	11	-
1999.....	116	131	167	220	350	394	404	308	145	51	11	-

Non-Hispanic 1/, male

2003.....	63	97	89	97	150	167	177	125	46	8	3	-
2002.....	61	82	81	104	140	157	149	110	40	18	3	-
2001.....	66	90	98	108	146	184	184	117	51	11	2	-
2000.....	83	73	99	116	197	202	183	130	47	12	1	-
1999.....	56	73	83	98	176	187	179	127	47	16	3	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic 1/, female

2003.....	59	74	99	103	146	219	206	174	104	46	10	-
2002.....	79	74	95	105	181	208	244	175	123	41	3	-
2001.....	67	83	98	120	163	235	228	193	126	43	5	-
2000.....	78	80	111	125	172	209	203	199	123	39	10	-
1999.....	60	58	84	122	174	207	225	181	98	35	8	-

Non-Hispanic White, both sexes

2003.....	84	126	150	164	246	324	347	261	134	50	11	-
2002.....	94	114	138	162	278	315	349	258	142	51	6	-
2001.....	99	132	147	180	237	367	368	262	151	48	6	-
2000.....	129	116	156	193	305	346	345	289	149	38	10	-
1999.....	84	91	133	185	293	337	351	262	129	43	7	-

Non-Hispanic White, male

2003.....	40	74	74	84	130	147	165	110	43	7	3	-
2002.....	44	63	66	78	122	138	138	97	35	14	3	-
2001.....	47	65	72	88	109	163	170	101	41	9	1	-
2000.....	69	60	75	91	164	177	170	115	43	10	1	-
1999.....	41	51	66	83	147	164	157	107	41	15	1	-

Non-Hispanic White, female

2003.....	44	52	76	80	116	177	182	151	91	43	8	-
2002.....	50	51	72	84	156	177	211	161	107	37	3	-
2001.....	52	67	75	92	128	204	198	161	110	39	5	-
2000.....	60	56	81	102	141	169	175	174	106	28	9	-
1999.....	43	40	67	102	146	173	194	155	88	28	6	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Non-Hispanic Black, both sexes

2003.....	33	39	35	34	42	56	30	31	14	4	2	-
2002.....	44	38	32	38	39	44	36	21	19	7	-	-
2001.....	31	38	46	42	62	46	40	46	23	6	1	-
2000.....	30	35	51	42	59	58	37	34	20	12	1	-
1999.....	31	34	27	31	53	52	42	37	16	7	3	-

Non-Hispanic Black, male

2003.....	21	20	15	13	14	17	10	10	3	1	-	-
2002.....	16	18	12	20	17	14	11	10	5	4	-	-
2001.....	18	22	25	20	33	17	13	14	9	2	1	-
2000.....	13	12	22	24	28	21	12	12	3	1	-	-
1999.....	15	16	14	13	27	21	17	12	6	1	1	-

Non-Hispanic Black, female

2003.....	12	19	20	21	28	39	20	21	11	3	2	-
2002.....	28	20	20	18	22	30	25	11	14	3	-	-
2001.....	13	16	21	22	29	29	27	32	14	4	-	-
2000.....	17	23	29	18	31	37	25	22	17	11	1	-
1999.....	16	18	13	18	26	31	25	25	10	6	2	-

Origin not stated 2/, both sexes

2003.....	-	-	-	-	-	1	-	1	-	-	-	-
2002.....	1	-	-	2	-	1	4	1	1	-	-	-
2001.....	-	-	2	2	3	2	2	-	1	-	-	-
2000.....	1	-	2	-	2	1	1	1	-	-	-	-
1999.....	1	1	-	1	-	-	1	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Complications of medical and surgical care (Y40-Y84,Y88)

Origin not stated 2/, male

2003.....	-	-	-	-	-	1	-	-	-	-	-	-
2002.....	1	-	-	1	-	-	1	-	-	-	-	-
2001.....	-	-	-	2	1	-	1	-	-	-	-	-
2000.....	-	-	1	-	2	1	1	-	-	-	-	-
1999.....	1	-	-	-	-	-	1	-	-	-	-	-

Origin not stated 2/, female

2003.....	-	-	-	-	-	-	-	1	-	-	-	-
2002.....	-	-	-	1	-	1	3	1	1	-	-	-
2001.....	-	-	2	-	2	2	1	-	1	-	-	-
2000.....	1	-	1	-	-	-	-	1	-	-	-	-
1999.....	-	1	-	1	-	-	-	1	1	-	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

All origins, both sexes

2003.....	3,591	2,891	2,277	1,641	1,130	770	413	149	38	7	-	11
2002 5/.....	3,427	2,734	2,154	1,582	1,253	806	384	154	39	8	-	8
2001 5/.....	3,305	2,495	2,104	1,645	1,201	816	388	166	40	14	1	16
2000 5/.....	2,940	2,455	2,087	1,644	1,266	831	392	162	55	7	3	8
1999 5/.....	2,853	2,374	2,065	1,676	1,250	876	415	175	43	11	1	8

All origins, male

2003.....	2,787	2,233	1,764	1,249	871	558	308	90	20	3	-	10
2002 5/.....	2,686	2,123	1,714	1,240	918	575	270	104	26	5	-	8
2001 5/.....	2,523	1,969	1,638	1,257	924	592	266	115	27	8	1	15
2000 5/.....	2,315	1,874	1,615	1,273	976	641	274	109	32	4	1	8
1999 5/.....	2,266	1,880	1,601	1,317	953	637	299	114	24	3	-	7

All origins, female

2003.....	804	658	513	392	259	212	105	59	18	4	-	1
2002 5/.....	741	611	440	342	335	231	114	50	13	3	-	-
2001 5/.....	782	526	466	388	277	224	122	51	13	6	-	1
2000 5/.....	625	581	472	371	290	190	118	53	23	3	2	-
1999 5/.....	587	494	464	359	297	239	116	61	19	8	1	1

Hispanic, both sexes

2003.....	376	337	239	178	103	78	36	9	1	-	-	3
2002 5/.....	387	306	215	184	135	61	26	12	5	2	-	1
2001 5/.....	353	288	234	175	121	79	34	8	2	-	-	4
2000 5/.....	333	288	216	195	116	72	24	7	3	1	-	3
1999 5/.....	310	255	215	165	106	64	26	17	3	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Hispanic, male

2003.....	329	294	206	136	83	62	29	5	1	-	-	2
2002 5/.....	330	269	185	161	115	45	20	7	5	1	-	1
2001 5/.....	297	249	201	157	105	61	24	7	2	-	-	4
2000 5/.....	288	255	185	177	101	65	21	4	2	1	-	3
1999 5/.....	266	222	184	140	93	48	21	13	1	-	-	1

Hispanic, female

2003.....	47	43	33	42	20	16	7	4	-	-	-	1
2002 5/.....	57	37	30	23	20	16	6	5	-	1	-	-
2001 5/.....	56	39	33	18	16	18	10	1	-	-	-	-
2000 5/.....	45	33	31	18	15	7	3	3	1	-	-	-
1999 5/.....	44	33	31	25	13	16	5	4	2	-	-	-

Non-Hispanic 1/, both sexes

2003.....	3,197	2,537	2,019	1,453	1,022	691	376	140	37	7	-	5
2002 5/.....	3,016	2,406	1,924	1,387	1,113	742	355	141	34	6	-	4
2001 5/.....	2,920	2,184	1,857	1,457	1,073	733	354	158	38	14	1	7
2000 5/.....	2,580	2,155	1,857	1,434	1,145	752	368	155	51	6	3	4
1999 5/.....	2,518	2,106	1,832	1,502	1,135	807	389	158	40	11	1	5

Non-Hispanic 1/, male

2003.....	2,443	1,924	1,539	1,105	784	495	278	85	19	3	-	5
2002 5/.....	2,335	1,836	1,515	1,068	798	527	248	96	21	4	-	4
2001 5/.....	2,199	1,702	1,426	1,091	813	528	242	108	25	8	1	6
2000 5/.....	2,002	1,609	1,418	1,084	871	569	253	105	29	3	1	4
1999 5/.....	1,979	1,647	1,400	1,168	852	584	278	101	23	3	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Non-Hispanic 1/, female

2003.....	754	613	480	348	238	196	98	55	18	4	-	-
2002 5/.....	681	570	409	319	315	215	107	45	13	2	-	-
2001 5/.....	721	482	431	366	260	205	112	50	13	6	-	1
2000 5/.....	578	546	439	350	274	183	115	50	22	3	2	-
1999 5/.....	539	459	432	334	283	223	111	57	17	8	1	-

Non-Hispanic White, both sexes

2003.....	2,588	2,105	1,641	1,238	886	624	335	127	34	6	-	4
2002 5/.....	2,456	1,954	1,564	1,170	949	659	313	124	31	6	-	3
2001 5/.....	2,269	1,733	1,534	1,191	940	644	323	134	33	14	1	5
2000 5/.....	2,048	1,714	1,505	1,192	990	664	328	139	45	5	3	2
1999 5/.....	2,001	1,655	1,474	1,238	962	694	344	134	35	11	1	5

Non-Hispanic White, male

2003.....	1,973	1,603	1,242	946	682	445	242	76	17	3	-	4
2002 5/.....	1,922	1,500	1,236	893	671	469	218	81	19	4	-	3
2001 5/.....	1,734	1,338	1,174	878	707	464	221	92	21	8	1	4
2000 5/.....	1,603	1,294	1,150	899	749	501	228	94	24	2	1	2
1999 5/.....	1,589	1,288	1,143	952	717	501	250	84	22	3	-	5

Non-Hispanic White, female

2003.....	615	502	399	292	204	179	93	51	17	3	-	-
2002 5/.....	534	454	328	277	278	190	95	43	12	2	-	-
2001 5/.....	535	395	360	313	233	180	102	42	12	6	-	1
2000 5/.....	445	420	355	293	241	163	100	45	21	3	2	-
1999 5/.....	412	367	331	286	245	193	94	50	13	8	1	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Non-Hispanic Black, both sexes

2003.....	477	348	309	178	107	53	34	12	1	1	-	1
2002 5/.....	451	370	283	172	131	70	34	14	2	-	-	1
2001 5/.....	542	375	270	230	107	77	27	16	3	-	-	2
2000 5/.....	446	355	284	212	131	75	29	13	6	1	-	2
1999 5/.....	420	381	306	224	153	92	40	20	4	-	-	-

Non-Hispanic Black, male

2003.....	379	259	248	131	83	38	29	9	1	-	-	1
2002 5/.....	336	276	223	142	104	49	25	12	1	-	-	1
2001 5/.....	399	302	213	187	86	56	18	9	2	-	-	2
2000 5/.....	338	261	213	162	107	59	20	9	5	1	-	2
1999 5/.....	319	308	225	185	119	66	26	13	1	-	-	-

Non-Hispanic Black, female

2003.....	98	89	61	47	24	15	5	3	-	1	-	-
2002 5/.....	115	94	60	30	27	21	9	2	1	-	-	-
2001 5/.....	143	73	57	43	21	21	9	7	1	-	-	-
2000 5/.....	108	94	71	50	24	16	9	4	1	-	-	-
1999 5/.....	101	73	81	39	34	26	14	7	3	-	-	-

Origin not stated 2/, both sexes

2003.....	18	17	19	10	5	1	1	-	-	-	-	3
2002 5/.....	24	22	15	11	5	3	3	1	-	-	-	3
2001 5/.....	32	23	13	13	7	4	-	-	-	-	-	5
2000 5/.....	27	12	14	15	5	7	-	-	1	-	-	1
1999 5/.....	25	13	18	9	9	5	-	-	-	-	-	2

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

Origin not stated 2/, male

2003.....	15	15	19	8	4	1	1	-	-	-	-	3
2002 5/.....	21	18	14	11	5	3	2	1	-	-	-	3
2001 5/.....	27	18	11	9	6	3	-	-	-	-	-	5
2000 5/.....	25	10	12	12	4	7	-	-	1	-	-	1
1999 5/.....	21	11	17	9	8	5	-	-	-	-	-	1

Origin not stated 2/, female

2003.....	3	2	-	2	1	-	-	-	-	-	-	-
2002 5/.....	3	4	1	-	-	-	1	-	-	-	-	-
2001 5/.....	5	5	2	4	1	1	-	-	-	-	-	-
2000 5/.....	2	2	2	3	1	-	-	-	-	-	-	-
1999 5/.....	4	2	1	-	1	-	-	-	-	-	-	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

All origins, both sexes

2003.....	3,228	1,546	693	441	314	288	249	187	79	13	4	12
2002 5/.....	2,731	1,235	593	365	288	264	220	159	84	15	2	10
2001 5/.....	2,136	901	460	296	257	254	185	136	69	17	4	16
2000 5/.....	1,815	794	393	244	242	235	195	150	75	15	-	14
1999 5/.....	1,582	760	401	301	255	264	203	134	49	17	1	18

All origins, male

2003.....	2,022	883	383	234	150	145	108	72	35	4	1	11
2002 5/.....	1,752	724	316	181	130	120	95	62	33	7	-	7
2001 5/.....	1,353	530	230	164	129	116	75	45	23	5	1	16
2000 5/.....	1,188	433	198	127	108	109	78	63	25	4	-	13
1999 5/.....	1,012	451	237	164	122	104	80	51	16	7	-	11

All origins, female

2003.....	1,206	663	310	207	164	143	141	115	44	9	3	1
2002 5/.....	979	511	277	184	158	144	125	97	51	8	2	3
2001 5/.....	783	371	230	132	128	138	110	91	46	12	3	-
2000 5/.....	627	361	195	117	134	126	117	87	50	11	-	1
1999 5/.....	570	309	164	137	133	160	123	83	33	10	1	7

Hispanic, both sexes

2003.....	214	92	43	26	12	12	12	8	2	-	-	-
2002 5/.....	175	74	37	19	17	12	8	1	1	-	-	-
2001 5/.....	142	66	22	17	12	7	5	2	-	2	-	2
2000 5/.....	117	61	18	11	9	5	3	2	3	-	-	-
1999 5/.....	144	46	38	17	11	5	2	1	1	-	-	5

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Hispanic, male

2003.....	155	72	32	18	10	6	7	4	1	-	-	-
2002 5/.....	138	54	27	10	8	9	4	-	1	-	-	-
2001 5/.....	113	53	16	12	6	5	1	1	-	1	-	2
2000 5/.....	87	41	11	8	3	2	1	1	3	-	-	-
1999 5/.....	103	40	29	16	5	1	-	1	-	-	-	4

Hispanic, female

2003.....	59	20	11	8	2	6	5	4	1	-	-	-
2002 5/.....	37	20	10	9	9	3	4	1	-	-	-	-
2001 5/.....	29	13	6	5	6	2	4	1	-	1	-	-
2000 5/.....	30	20	7	3	6	3	2	1	-	-	-	-
1999 5/.....	41	6	9	1	6	4	2	-	1	-	-	1

Non-Hispanic 1/, both sexes

2003.....	2,987	1,445	642	411	300	275	236	179	77	13	4	7
2002 5/.....	2,541	1,152	553	340	268	249	212	158	83	14	2	7
2001 5/.....	1,971	821	431	276	244	246	180	133	69	15	4	5
2000 5/.....	1,667	722	369	232	230	230	192	148	72	15	-	8
1999 5/.....	1,413	696	356	281	243	259	199	131	48	17	1	6

Non-Hispanic 1/, male

2003.....	1,844	802	344	212	138	138	100	68	34	4	1	6
2002 5/.....	1,607	662	286	166	119	109	91	62	32	6	-	5
2001 5/.....	1,223	466	207	150	122	110	74	44	23	4	1	5
2000 5/.....	1,078	384	181	118	104	107	77	62	22	4	-	7
1999 5/.....	889	399	201	145	116	103	80	49	16	7	-	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Non-Hispanic 1/, female

2003.....	1,143	643	298	199	162	137	136	111	43	9	3	1
2002 5/.....	934	490	267	174	149	140	121	96	51	8	2	2
2001 5/.....	748	355	224	126	122	136	106	89	46	11	3	-
2000 5/.....	589	338	188	114	126	123	115	86	50	11	-	1
1999 5/.....	524	297	155	136	127	156	119	82	32	10	1	2

Non-Hispanic White, both sexes

2003.....	2,392	1,192	538	342	270	263	225	168	74	12	4	4
2002 5/.....	1,998	937	448	296	246	228	200	144	78	12	2	5
2001 5/.....	1,524	671	348	237	219	226	168	121	64	11	2	5
2000 5/.....	1,266	585	315	198	212	219	180	133	65	11	-	4
1999 5/.....	1,073	556	284	243	218	243	190	115	47	15	1	2

Non-Hispanic White, male

2003.....	1,437	629	271	163	123	130	95	62	31	3	1	3
2002 5/.....	1,212	502	206	133	108	99	80	57	32	5	-	3
2001 5/.....	895	353	145	127	105	93	71	39	23	2	-	5
2000 5/.....	785	284	144	95	90	103	72	56	18	2	-	3
1999 5/.....	636	289	142	119	104	93	76	46	16	6	-	1

Non-Hispanic White, female

2003.....	955	563	267	179	147	133	130	106	43	9	3	1
2002 5/.....	786	435	242	163	138	129	120	87	46	7	2	2
2001 5/.....	629	318	203	110	114	133	97	82	41	9	2	-
2000 5/.....	481	301	171	103	122	116	108	77	47	9	-	1
1999 5/.....	437	267	142	124	114	150	114	69	31	9	1	1

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

Non-Hispanic Black, both sexes

2003.....	541	227	88	60	22	12	9	9	3	1	-	3
2002 5/.....	516	201	100	33	15	16	9	8	4	1	-	1
2001 5/.....	427	139	75	31	21	17	8	10	5	3	2	-
2000 5/.....	385	129	49	32	16	11	11	14	6	4	-	4
1999 5/.....	317	123	67	34	21	12	8	14	1	2	-	4

Non-Hispanic Black, male

2003.....	373	162	65	46	12	8	4	5	3	1	-	3
2002 5/.....	378	152	79	26	8	8	8	2	-	-	-	1
2001 5/.....	314	105	58	20	15	14	1	5	-	1	1	-
2000 5/.....	283	96	35	22	12	4	5	6	3	2	-	4
1999 5/.....	237	101	54	24	11	7	4	3	-	1	-	3

Non-Hispanic Black, female

2003.....	168	65	23	14	10	4	5	4	-	-	-	-
2002 5/.....	138	49	21	7	7	8	1	6	4	1	-	-
2001 5/.....	113	34	17	11	6	3	7	5	5	2	1	-
2000 5/.....	102	33	14	10	4	7	6	8	3	2	-	-
1999 5/.....	80	22	13	10	10	5	4	11	1	1	-	1

Origin not stated 2/, both sexes

2003.....	27	9	8	4	2	1	1	-	-	-	-	5
2002 5/.....	15	9	3	6	3	3	-	-	-	1	-	3
2001 5/.....	23	14	7	3	1	1	-	1	-	-	-	9
2000 5/.....	31	11	6	1	3	-	-	-	-	-	-	6
1999 5/.....	25	18	7	3	1	-	2	2	-	-	-	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

## Origin not stated 2/, male

2003.....	23	9	7	4	2	1	1	-	-	-	-	5
2002 5/.....	7	8	3	5	3	2	-	-	-	1	-	2
2001 5/.....	17	11	7	2	1	1	-	-	-	-	-	9
2000 5/.....	23	8	6	1	1	-	-	-	-	-	-	6
1999 5/.....	20	12	7	3	1	-	-	1	-	-	-	3

## Origin not stated 2/, female

2003.....	4	-	1	-	-	-	-	-	-	-	-	-
2002 5/.....	8	1	-	1	-	1	-	-	-	-	-	1
2001 5/.....	6	3	-	1	-	-	-	1	-	-	-	-
2000 5/.....	8	3	-	-	2	-	-	-	-	-	-	-
1999 5/.....	5	6	-	-	-	-	2	1	-	-	-	4

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

All origins, both sexes

2003.....	2,075	1,637	1,167	945	967	963	768	420	144	25	-	18
2002.....	1,963	1,608	1,111	966	1,027	1,011	822	411	138	27	-	15
2001.....	1,917	1,471	1,073	999	999	1,014	789	402	139	20	2	19
2000.....	1,805	1,289	983	923	1,018	994	729	443	137	19	1	18
1999.....	1,626	1,299	1,007	955	1,079	1,029	709	428	108	25	-	20

All origins, male

2003.....	1,736	1,362	1,016	835	859	863	695	388	136	22	-	16
2002.....	1,652	1,354	942	836	932	916	755	386	128	26	-	14
2001.....	1,588	1,231	885	865	889	915	719	373	128	19	2	18
2000.....	1,488	1,071	825	783	901	900	671	404	125	19	1	13
1999.....	1,335	1,080	871	833	953	928	652	392	103	22	-	19

All origins, female

2003.....	339	275	151	110	108	100	73	32	8	3	-	2
2002.....	311	254	169	130	95	95	67	25	10	1	-	1
2001.....	329	240	188	134	110	99	70	29	11	1	-	1
2000.....	317	218	158	140	117	94	58	39	12	-	-	5
1999.....	291	219	136	122	126	101	57	36	5	3	-	1

Hispanic, both sexes

2003.....	93	43	43	29	29	21	9	9	2	-	-	4
2002.....	93	62	37	34	28	15	18	9	2	2	-	4
2001.....	88	47	35	36	24	14	17	6	3	1	-	5
2000.....	99	51	32	30	34	17	11	7	1	1	-	3
1999.....	67	54	33	36	15	27	17	5	5	2	-	3

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Hispanic, male

2003.....	78	40	40	28	26	19	9	8	2	-	-	3
2002.....	78	56	33	30	27	15	17	9	2	2	-	4
2001.....	74	44	31	33	22	14	14	6	3	1	-	5
2000.....	83	46	27	26	28	15	11	7	1	1	-	2
1999.....	55	46	30	31	15	27	15	5	5	2	-	3

Hispanic, female

2003.....	15	3	3	1	3	2	-	1	-	-	-	1
2002.....	15	6	4	4	1	-	1	-	-	-	-	-
2001.....	14	3	4	3	2	-	3	-	-	-	-	-
2000.....	16	5	5	4	6	2	-	-	-	-	-	1
1999.....	12	8	3	5	-	-	2	-	-	-	-	-

Non-Hispanic 1/, both sexes

2003.....	1,979	1,587	1,122	913	936	941	759	409	142	25	-	9
2002.....	1,861	1,540	1,069	925	997	992	803	401	135	25	-	7
2001.....	1,825	1,417	1,029	956	971	998	769	393	136	18	2	8
2000.....	1,699	1,235	944	888	981	976	718	435	135	18	1	6
1999.....	1,551	1,242	968	916	1,062	997	692	421	103	23	-	7

Non-Hispanic 1/, male

2003.....	1,655	1,315	974	805	832	843	686	378	134	22	-	8
2002.....	1,566	1,294	904	799	903	897	737	376	125	24	-	6
2001.....	1,510	1,180	845	825	863	899	702	364	125	17	2	7
2000.....	1,399	1,022	791	754	870	884	660	397	123	18	1	5
1999.....	1,274	1,031	835	799	936	896	637	386	98	20	-	7

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Non-Hispanic 1/, female

2003.....	324	272	148	108	104	98	73	31	8	3	-	1
2002.....	295	246	165	126	94	95	66	25	10	1	-	1
2001.....	315	237	184	131	108	99	67	29	11	1	-	1
2000.....	300	213	153	134	111	92	58	38	12	-	-	1
1999.....	277	211	133	117	126	101	55	35	5	3	-	-

Non-Hispanic White, both sexes

2003.....	1,727	1,449	1,034	851	880	899	728	401	139	23	-	6
2002.....	1,604	1,406	974	865	947	932	782	389	132	23	-	5
2001.....	1,604	1,278	955	890	907	953	732	384	132	18	2	6
2000.....	1,476	1,100	865	823	915	945	692	424	132	17	1	4
1999.....	1,357	1,112	864	850	1,002	948	666	411	101	21	-	3

Non-Hispanic White, male

2003.....	1,450	1,201	900	754	787	810	661	370	132	22	-	6
2002.....	1,358	1,175	830	748	860	844	718	365	123	22	-	4
2001.....	1,330	1,055	783	768	809	860	671	356	121	17	2	5
2000.....	1,212	908	721	699	809	860	638	389	120	17	1	3
1999.....	1,125	913	746	744	882	854	614	377	96	19	-	3

Non-Hispanic White, female

2003.....	277	248	134	97	93	89	67	31	7	1	-	-
2002.....	246	231	144	117	87	88	64	24	9	1	-	1
2001.....	274	223	172	122	98	93	61	28	11	1	-	1
2000.....	264	192	144	124	106	85	54	35	12	-	-	1
1999.....	232	199	118	106	120	94	52	34	5	2	-	-

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Non-Hispanic Black, both sexes

2003.....	218	113	73	56	47	33	25	7	3	2	-	2
2002.....	209	106	80	48	47	49	16	12	3	2	-	2
2001.....	192	112	59	53	55	39	30	7	4	-	-	1
2000.....	190	99	63	56	58	26	24	10	2	-	-	1
1999.....	152	111	84	58	51	47	21	10	2	2	-	4

Non-Hispanic Black, male

2003.....	178	96	60	46	37	27	20	7	2	-	-	1
2002.....	168	95	61	41	41	43	15	11	2	2	-	2
2001.....	160	101	49	46	47	34	25	6	4	-	-	1
2000.....	160	82	54	49	54	20	20	7	2	-	-	1
1999.....	122	102	72	49	46	41	18	9	2	1	-	4

Non-Hispanic Black, female

2003.....	40	17	13	10	10	6	5	-	1	2	-	1
2002.....	41	11	19	7	6	6	1	1	1	-	-	-
2001.....	32	11	10	7	8	5	5	1	-	-	-	-
2000.....	30	17	9	7	4	6	4	3	-	-	-	-
1999.....	30	9	12	9	5	6	3	1	-	1	-	-

Origin not stated 2/, both sexes

2003.....	3	7	2	3	2	1	-	2	-	-	-	5
2002.....	9	6	5	7	2	4	1	1	1	-	-	4
2001.....	4	7	9	7	4	2	3	3	-	1	-	6
2000.....	7	3	7	5	3	1	-	1	1	-	-	9
1999.....	8	3	6	3	2	5	-	2	-	-	-	10

Worktable Orig291F. Deaths from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2003

[Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget guidelines. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

Origin not stated 2/, male

2003.....	3	7	2	2	1	1	-	2	-	-	-	5
2002.....	8	4	5	7	2	4	1	1	1	-	-	4
2001.....	4	7	9	7	4	2	3	3	-	1	-	6
2000.....	6	3	7	3	3	1	-	-	1	-	-	6
1999.....	6	3	6	3	2	5	-	1	-	-	-	9

Origin not stated 2/, female

2003.....	-	-	-	1	1	-	-	-	-	-	-	-
2002.....	1	2	-	-	-	-	-	-	-	-	-	-
2001.....	-	-	-	-	-	-	-	-	-	-	-	-
2000.....	1	-	-	2	-	-	-	1	-	-	-	3
1999.....	2	-	-	-	-	-	-	1	-	-	-	1

- Quantity zero.

... Category not applicable.

1/ Includes races other than white and black.

2/ Includes data for which Hispanic origin was not reported on the death certificate.

3/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

4/ Included in selected categories.

5/ Figures may differ from those previously shown as they are based on a modified list of causes; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003."

Source: CDC/NCHS, National Vital Statistics System, Mortality.