Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
All races 3/, both sexes												
2003	841.9	700.0	31.5	17.0	81.5	103.6	201.6	433.2	940.9	2,255.0	5,463.1	14,593.3
2002	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6
All races, male												
2003	840.3	777.4	35.1	19.8	116.5	141.4	255.0	552.2	1,165.5	2,771.7	6,641.8	15,794.0
2002	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2
All races, female												
2003	843.4	619.1	27.8	14.0	44.4	64.6	148.5	318.4	732.7	1,823.0	4,675.5	14,062.5
2002	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
White, both sexes												
2003	890.1	591.3	28.5	15.8	77.1	94.3	185.7	397.5	892.2	2,207.0	5,457.8	14,792.3
2002	895.7	586.7	28.1	16.1	77.1	93.5	186.6	395.3	902.5	2,267.6	5,544.2	15,015.9
2001	895.1	570.9	30.7	16.2	75.5	95.3	185.6	393.2	913.5	2,298.9	5,560.9	15,287.5
2000	900.2	610.6	29.2	17.0	74.5	90.3	179.9	388.3	940.2	2,341.5	5,637.7	15,707.0
1999	901.4	606.1	30.8	17.2	73.2	91.4	177.3	380.4	948.1	2,396.2	5,679.5	15,718.0
White, male												
2003	877.6	658.7	31.5	18.4	108.9	128.8	237.6	509.2	1,102.3	2,707.3	6,620.6	16,037.9
2002	884.0	650.9	31.5	18.4	109.7	128.3	239.3	505.4	1,118.6	2,795.4	6,738.8	16,473.2
2001	881.9	627.6	34.2	18.4	108.0	130.3	239.7	501.3	1,127.5	2,842.3	6,799.7	16,935.4
2000	887.8	667.6	32.6	19.8	105.8	124.1	233.6	496.9	1,163.3	2,905.7	6,933.1	17,716.4
1999	892.1	667.0	33.9	19.7	102.8	125.4	230.8	484.6	1,179.7	2,998.7	7,040.1	17,752.9
White, female												
2003	902.3	520.7	25.5	13.1	43.2	58.0	132.9	287.2	693.7	1,780.6	4,671.6	14,240.6
2002	907.0	519.4	24.5	13.7	42.4	56.9	133.2	286.8	698.7	1,819.7	4,742.5	14,382.8
2001	907.9	511.5	27.1	13.9	40.8	58.6	131.0	286.8	712.2	1,840.2	4,738.5	14,597.6
2000	912.3	550.5	25.5	14.1	41.1	55.1	125.7	281.4	730.9	1,868.3	4,785.3	14,890.7
1999	910.4	542.0	27.5	14.6	41.5	56.2	123.2	277.9	731.0	1,893.9	4,787.8	14,900.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes												
Black, both sexes												
2003	763.6	1,273.8	46.8	22.9	113.1	181.6	343.6	771.4	1,549.9	3,108.4	6,353.9	13,976.7
2002	768.4	1,263.6	47.1	24.5	113.7	186.8	348.6	765.8	1,565.2	3,159.3	6,518.9	14,260.8
2001	773.5	1,304.5	47.5	23.3	117.7	185.0	356.5	771.9	1,576.8	3,257.6	6,585.6	14,550.8
2000	781.1	1,426.1	49.9	24.2	119.4	187.8	356.9	786.9	1,607.8	3,335.6	6,701.3	14,714.3
1999	788.1	1,457.3	52.6	26.5	121.2	184.4	368.4	782.4	1,655.6	3,413.6	6,753.3	14,757.9
Black, male												
2003	813.7	1,410.0	53.7	26.8	171.3	256.2	426.7	991.4	2,011.2	3,981.3	8,067.0	14,903.4
2002	816.7	1,351.5	54.4	28.9	172.6	264.5	434.7	983.0	2,039.2	4,024.5	8,169.6	15,635.5
2001	823.9	1,426.4	53.2	27.7	180.7	259.2	445.0	996.1	2,025.3	4,166.6	8,355.9	16,439.9
2000	834.1	1,567.6	54.5	28.2	181.4	261.0	453.0	1,017.7	2,080.1	4,253.5	8,486.0	16,791.0
1999	847.4	1,592.8	59.0	32.0	184.6	258.6	469.2	1,030.7	2,145.6	4,352.3	8,559.1	17,304.5
Black, female												
2003	717.9	1,132.2	39.7	18.9	54.0	113.5	270.0	582.3	1,178.8	2,487.2	5,385.5	13,616.7
2002	724.4	1,172.0	39.5	19.9	54.4	116.4	272.3	579.4	1,184.2	2,545.0	5,584.4	13,734.2
2001	727.7	1,176.3	41.7	18.7	54.9	117.9	278.0	579.1	1,215.5	2,616.1	5,591.6	13,849.9
2000	733.0	1,279.8	45.3	20.0	58.3	121.8	271.9	588.3	1,227.2	2,689.6	5,696.5	13,941.3
1999	734.3	1,317.4	46.1	20.9	58.6	117.6	279.4	569.8	1,262.7	2,751.5	5,742.4	13,805.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
All races 3/, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
White, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Salmonella infections (A01-A02)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Shigellosis and amebiasis (A03,A06)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
All races 3/, both sexes												
2003	1.1	*	*	*	*	*	*	0.2	0.6	2.4	9.3	24.8
2002	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.0	21.0
2001	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3
All races, male												
2003	0.8	*	*	*	*	*	*	0.2	0.6	2.2	9.3	21.4
2002	0.7	*	*	*	*	*	*	0.1	0.6	2.2	7.0	20.4
2001	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5
All races, female												
2003	1.3	*	*	*	*	*	*	0.2	0.6	2.6	9.3	26.3
2002	1.0	*	*	*	*	*	*	0.1	0.5	1.8	6.9	21.3
2001	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
White, both sexes												
2003	1.2	*	*	*	*	*	*	0.1	0.6	2.6	9.7	26.1
2002	1.0	*	*	*	*	*	*	0.1	0.5	2.1	7.4	22.0
2001	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.7	13.5
2000	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.2
1999	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.9
White, male												
2003	0.9	*	*	*	*	*	*	0.2	0.6	2.4	9.8	22.7
2002	0.8	*	*	*	*	*	*	0.2	0.5	2.3	7.4	21.4
2001	0.5	*	*	*	*	*	*	*	0.3	1.3	4.6	12.7
2000	0.4	*	*	*	*	*	*	*	0.3	1.3	4.2	11.9
1999	0.3	*	*	*	*	*	*	*	0.2	1.0	3.7	9.9
White, female												
2003	1.5	*	*	*	*	*	*	0.1	0.7	2.8	9.7	27.6
2002	1.2	*	*	*	*	*	*	0.1	0.5	1.9	7.4	22.3
2001	0.8	*	*	*	*	*	*	0.2	0.3	1.3	4.7	13.9
2000	0.6	*	*	*	*	*	*	*	0.3	1.2	3.8	12.3
1999	0.5	*	*	*	*	*	*	*	*	1.0	3.1	11.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain other intestinal infections (A04,A07-A09)												
Black, both sexes												
2003	0.5	*	*	*	*	*	*	*	0.8	1.5	6.5	15.2
2002	0.4	*	*	*	*	*	*	*	1.1	1.8	3.7	13.1
2001	0.3	*	*	*	*	*	*	*	*	1.4	3.2	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	8.1
1999	0.2	*	*	*	*	*	*	*	*	*	2.8	*
Black, male												
2003	0.4	*	*	*	*	*	*	*	*	*	6.5	*
2002	0.4	*	*	*	*	*	*	*	*	2.9	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.6	*	*	*	*	*	*	*	*	*	6.5	16.6
2002	0.5	*	*	*	*	*	*	*	*	*	3.7	13.1
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
All races 3/, both sexes												
2003	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.7	2.3
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2
All races, male												
2003	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	2.3	3.1
2002	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1
All races, female												
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
White, both sexes												
2003	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.3	1.8
2002	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4	2.1
2001	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.2
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
1999	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.2
White, male												
2003	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.7	2.4
2002	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.8	3.2
2001	0.2	*	*	*	*	*	*	0.2	0.5	0.7	1.5	3.1
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.7	1.8	3.8
1999	0.3	*	*	*	*	*	*	0.3	0.5	1.0	2.1	5.4
White, female												
2003	0.2	*	*	*	*	*	*	*	*	0.3	1.1	1.6
2002	0.2	*	*	*	*	*	*	*	0.2	0.3	1.2	1.6
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	2.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Tuberculosis (A16-A19)												
Black, both sexes												
2003	0.5	*	*	*	*	*	*	1.0	1.1	1.2	3.4	*
2002	0.5	*	*	*	*	*	0.4	0.8	1.3	2.5	3.8	*
2001	0.5	*	*	*	*	*	*	0.8	1.5	2.6	3.7	6.8
2000	0.7	*	*	*	*	*	0.4	1.0	1.4	3.1	6.5	*
1999	0.7	*	*	*	*	*	0.4	1.0	1.7	2.6	5.5	8.6
Black, male												
2003	0.6	*	*	*	*	*	*	1.6	1.7	*	5.6	*
2002	0.7	*	*	*	*	*	*	1.1	2.1	3.6	6.3	*
2001	0.7	*	*	*	*	*	*	1.5	2.4	4.3	*	*
2000	0.9	*	*	*	*	*	*	1.9	2.4	4.1	12.1	*
1999	0.9	*	*	*	*	*	*	1.8	3.0	4.3	7.7	*
Black, female												
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	3.5	*
2000	0.4	*	*	*	*	*	*	*	*	2.5	3.4	*
1999	0.5	*	*	*	*	*	*	*	*	*	4.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
All races 3/, both sexes												
2003	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.4	1.8
2002	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7
All races, male												
2003	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.9	2.5
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4
All races, female												
2003	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.5
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
White, both sexes												
2003	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.4
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	1.8
2001	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8
2000	0.2	*	*	*	*	*	*	0.1	0.1	0.4	1.2	1.9
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.3	3.0
White, male												
2003	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.4	1.8
2002	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.4	2.9
2001	0.2	*	*	*	*	*	*	0.1	0.5	0.6	1.3	2.6
2000	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.5	3.4
1999	0.3	*	*	*	*	*	*	0.2	0.5	0.8	1.8	5.2
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2002	0.1	*	*	*	*	*	*	*	*	*	1.0	1.3
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.4
2000	0.1	*	*	*	*	*	*	*	*	0.3	1.0	1.2
1999	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Respiratory tuberculosis (A16)												
Black, both sexes												
2003	0.4	*	*	*	*	*	*	0.9	0.9	*	2.6	*
2002	0.4	*	*	*	*	*	*	0.6	1.1	1.9	3.5	*
2001	0.4	*	*	*	*	*	*	0.6	1.2	1.9	3.0	6.1
2000	0.5	*	*	*	*	*	0.3	0.8	0.9	2.6	5.2	*
1999	0.6	*	*	*	*	*	0.4	0.7	1.5	2.0	4.7	7.3
Black, male												
2003	0.5	*	*	*	*	*	*	1.3	*	*	*	*
2002	0.6	*	*	*	*	*	*	0.9	1.9	*	6.0	*
2001	0.6	*	*	*	*	*	*	1.2	2.1	3.3	*	*
2000	0.8	*	*	*	*	*	*	1.5	2.0	3.8	9.7	*
1999	0.7	*	*	*	*	*	*	1.3	2.7	3.2	6.8	*
Black, female												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	3.6	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
All races 3/, both sexes												
2003	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.4
2002	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	*
2000	0.1	*	*	*	*	*	*	*	0.1	0.2	0.3	*
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.3	0.5
All races, male												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.5	*
2001	0.1	*	*	*	*	*	*	*	*	0.3	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	0.1	*	*	*	*
All races, female												
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.3	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
White, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
White, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.5	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other tuberculosis (A17-A19)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Whooping cough (A37)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Scarlet fever and erysipelas (A38,A46)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
All races 3/, both sexes												
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*
All races, male												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.1	*	*	*	0.1	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
White, both sexes												
2003	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2001	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999	0.1	1.0	0.2	*	0.1	*	*	*	*	*	*	*
White, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningococcal infection (A39)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
All races 3/, both sexes												
2003	11.7	6.9	0.5	0.2	0.4	0.8	2.1	5.3	13.1	32.6	85.0	202.5
2002	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7
All races, male												
2003	10.5	7.4	0.7	0.2	0.4	0.8	2.1	5.8	14.1	36.2	95.0	213.8
2002	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5
All races, female												
2003	12.9	6.5	0.4	0.2	0.4	0.8	2.0	4.8	12.2	29.5	78.3	197.5
2002	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
White, both sexes												
2003	11.5	4.6	0.4	0.2	0.3	0.6	1.7	4.3	11.2	28.9	80.1	191.1
2002	11.5	6.1	0.5	0.2	0.3	0.6	1.6	4.2	10.6	31.1	80.7	191.9
2001	11.1	5.4	0.7	0.2	0.3	0.5	1.4	4.0	10.3	29.3	77.1	195.3
2000	10.8	5.7	0.6	0.2	0.2	0.5	1.6	3.9	9.9	27.4	73.9	203.0
1999	10.6	5.5	0.5	0.2	0.3	0.6	1.4	3.6	9.4	27.3	72.8	205.9
White, male												
2003	10.2	4.6	0.6	0.2	0.3	0.6	1.7	4.7	11.9	32.1	89.5	203.3
2002	10.3	6.2	0.5	0.3	0.3	0.7	1.6	4.7	11.7	34.1	91.5	202.9
2001	9.9	6.2	0.8	0.2	0.3	0.5	1.7	4.3	11.1	32.5	89.4	211.8
2000	9.4	5.8	0.6	0.2	0.2	0.6	1.7	4.4	10.9	30.8	84.6	209.8
1999	9.3	6.9	0.5	0.2	0.3	0.6	1.6	4.0	10.0	30.6	84.6	217.4
White, female												
2003	12.8	4.5	*	0.2	0.3	0.6	1.7	3.8	10.5	26.2	73.8	185.8
2002	12.8	6.0	0.4	0.2	0.3	0.6	1.5	3.8	9.6	28.6	73.4	187.1
2001	12.3	4.6	0.6	0.1	0.2	0.5	1.2	3.7	9.7	26.6	69.0	188.3
2000	12.1	5.6	0.5	*	0.1	0.4	1.5	3.3	9.0	24.6	66.8	200.2
1999	11.9	4.0	0.5	0.2	0.2	0.5	1.2	3.2	9.0	24.5	65.1	201.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Septicemia (A40-A41)												
Black, both sexes												
2003	16.3	18.6	0.9	*	0.6	1.9	4.9	13.7	32.2	73.0	157.5	372.0
2002	16.3	13.5	*	0.3	0.4	1.9	4.4	12.9	33.2	74.7	171.2	367.0
2001	15.8	19.3	0.9	*	0.7	1.8	4.6	13.4	32.1	71.8	160.2	364.8
2000	16.2	16.4	1.0	*	0.6	2.0	4.7	13.6	31.4	71.9	173.4	392.6
1999	16.1	17.7	*	*	0.6	1.5	4.9	12.9	30.3	74.4	168.3	412.3
Black, male												
2003	15.3	21.1	*	*	*	2.1	5.0	15.3	37.8	85.7	189.0	404.5
2002	15.0	14.8	*	*	*	1.9	4.9	14.7	35.9	86.1	212.3	393.5
2001	14.9	20.1	*	*	0.8	1.8	4.7	14.9	36.9	85.6	195.1	424.4
2000	14.9	16.0	*	*	*	2.3	4.6	16.3	35.5	81.2	207.1	427.8
1999	15.3	19.8	*	*	*	1.6	5.8	15.1	37.1	90.3	201.7	453.5
Black, female												
2003	17.2	16.1	*	*	0.7	1.6	4.9	12.3	27.8	63.9	139.7	359.3
2002	17.4	12.1	*	*	*	1.9	4.1	11.4	30.9	66.6	148.0	356.9
2001	16.7	18.6	*	*	*	1.8	4.6	12.1	28.3	62.0	140.6	342.7
2000	17.4	16.9	*	*	*	1.7	4.8	11.3	28.2	65.3	154.4	379.5
1999	16.9	15.4	*	*	*	1.5	4.2	11.1	24.8	63.2	149.6	396.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
All races 3/, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Syphilis (A50-A53)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute poliomyelitis (A80)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Arthropod-borne viral encephalitis (A83-A84,A85.2)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Measles (B05)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
All races 3/, both sexes												
2003	1.9	*	*	*	*	0.2	1.5	5.5	3.9	3.8	4.2	2.4
2002	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7
All races, male												
2003	2.5	*	*	*	*	0.2	2.1	8.3	5.4	4.2	4.8	2.8
2002	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9
All races, female												
2003	1.3	*	*	*	*	0.1	0.9	2.9	2.5	3.5	3.8	2.3
2002	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
White, both sexes												
2003	1.8	*	*	*	*	0.2	1.5	5.4	3.3	3.4	3.9	2.3
2002	2.0	*	*	*	0.1	0.2	1.6	5.6	3.5	3.7	4.4	3.7
2001	1.9	*	*	*	*	0.2	1.8	5.0	3.6	4.1	4.7	3.6
2000	1.8	*	*	*	*	0.2	1.7	4.6	3.2	4.2	5.1	3.9
1999	1.7	*	*	*	*	0.2	1.5	3.8	2.9	4.3	5.1	4.5
White, male												
2003	2.4	*	*	*	*	0.2	2.1	8.2	4.7	3.8	4.4	2.8
2002	2.6	*	*	*	*	0.3	2.3	8.7	4.8	4.1	5.2	4.9
2001	2.5	*	*	*	*	0.2	2.5	7.8	4.8	4.5	5.4	4.5
2000	2.4	*	*	*	*	0.2	2.4	7.2	4.3	4.3	6.2	5.7
1999	2.2	*	*	*	*	0.3	2.3	5.7	3.9	4.8	6.5	6.5
White, female												
2003	1.2	*	*	*	*	*	0.9	2.7	2.0	3.0	3.5	2.1
2002	1.3	*	*	*	*	*	0.9	2.7	2.4	3.3	3.9	3.2
2001	1.3	*	*	*	*	0.1	1.0	2.2	2.4	3.7	4.2	3.2
2000	1.3	*	*	*	*	0.1	1.0	2.1	2.2	4.1	4.5	3.2
1999	1.2	*	*	*	*	0.2	0.8	1.9	2.0	3.9	4.1	3.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Viral hepatitis (B15-B19)												
Black, both sexes												
2003	2.2	*	*	*	*	*	1.7	7.4	8.0	5.4	6.0	*
2002	2.3	*	*	*	*	0.4	1.9	8.0	7.1	6.1	5.2	*
2001	2.2	*	*	*	*	*	2.1	7.9	6.7	5.6	5.7	*
2000	2.2	*	*	*	*	0.4	2.0	7.9	5.9	5.7	6.2	*
1999	2.0	*	*	*	*	*	2.1	6.9	5.0	6.9	6.1	6.4
Black, male												
2003	3.0	*	*	*	*	*	2.4	11.5	11.8	6.1	6.2	*
2002	3.0	*	*	*	*	*	2.5	12.4	10.5	7.1	*	*
2001	2.7	*	*	*	*	*	2.9	11.8	8.9	4.9	*	*
2000	2.9	*	*	*	*	*	2.9	12.3	8.1	7.0	7.6	*
1999	2.6	*	*	*	*	*	3.0	11.6	6.2	5.9	*	*
Black, female												
2003	1.6	*	*	*	*	*	1.1	4.0	5.0	4.9	5.9	*
2002	1.6	*	*	*	*	*	1.4	4.3	4.3	5.4	5.4	*
2001	1.8	*	*	*	*	*	1.4	4.7	4.9	6.0	6.8	*
2000	1.5	*	*	*	*	*	1.3	4.2	4.1	4.7	5.5	*
1999	1.5	*	*	*	*	*	1.3	2.8	4.1	7.6	6.5	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
All races 3/, both sexes												
2003	4.7	*	*	0.1	0.4	4.0	12.0	10.9	5.4	2.4	0.7	*
2002	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*
All races, male												
2003	7.1	*	*	*	0.4	5.1	17.5	17.2	9.1	4.0	1.5	*
2002	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*
All races, female												
2003	2.4	*	*	*	0.4	2.8	6.5	4.8	2.1	1.0	0.3	*
2002	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
White, both sexes												
2003	2.5	*	*	*	0.2	2.0	6.8	5.8	2.9	1.2	0.4	*
2002	2.6	*	*	*	0.2	2.2	7.0	5.8	2.8	1.0	0.4	*
2001	2.7	*	*	*	0.2	2.6	7.3	5.5	2.8	1.0	0.3	*
2000	2.8	*	*	0.1	0.2	3.1	7.5	5.9	2.8	1.0	0.4	*
1999	3.0	*	*	0.1	0.2	3.7	8.0	5.7	2.5	1.1	0.3	*
White, male												
2003	4.2	*	*	*	0.2	3.0	11.0	10.0	5.0	2.1	0.9	*
2002	4.3	*	*	*	0.2	3.3	11.6	9.8	4.9	1.8	0.6	*
2001	4.5	*	*	*	0.2	3.9	12.1	9.5	4.8	1.9	0.7	*
2000	4.7	*	*	*	0.2	4.6	12.4	10.1	5.0	1.9	0.7	*
1999	5.0	*	*	*	0.3	5.8	13.2	9.9	4.7	2.0	0.7	*
White, female												
2003	0.9	*	*	*	0.1	0.9	2.5	1.7	0.9	0.4	*	*
2002	0.9	*	*	*	0.1	1.0	2.5	1.8	0.8	*	*	*
2001	0.9	*	*	*	0.2	1.2	2.4	1.5	0.8	0.3	*	*
2000	1.0	*	*	*	*	1.5	2.6	1.7	0.8	0.3	*	*
1999	1.0	*	*	*	0.2	1.6	2.7	1.6	0.6	0.3	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)												
Black, both sexes												
2003	19.6	*	*	0.5	1.9	17.3	49.2	50.6	29.6	14.5	4.7	*
2002	20.8	*	*	*	2.0	20.6	52.8	53.9	26.1	14.4	5.6	*
2001	21.1	*	*	0.4	2.7	23.2	53.7	51.2	27.7	12.8	5.3	*
2000	21.4	*	*	0.3	2.1	26.3	53.5	53.4	26.0	13.5	4.8	*
1999	21.8	*	*	0.6	2.3	27.3	56.1	51.8	26.2	13.5	3.6	*
Black, male												
2003	27.7	*	*	*	1.9	20.8	68.0	77.8	50.7	25.8	9.0	*
2002	29.5	*	*	*	1.9	24.5	74.0	85.0	44.4	26.3	12.0	*
2001	30.1	*	*	*	2.3	29.8	75.9	80.7	47.1	21.7	11.5	*
2000	31.0	*	*	*	2.0	32.3	77.5	86.0	46.0	24.6	9.4	*
1999	31.9	*	*	0.7	1.9	36.0	81.9	85.0	45.5	24.7	6.5	*
Black, female												
2003	12.3	*	*	*	1.9	14.2	32.5	27.3	12.5	6.4	*	*
2002	12.8	*	*	*	2.1	17.1	34.0	27.2	11.3	6.0	*	*
2001	12.9	*	*	*	3.1	17.3	34.1	25.9	12.1	6.5	*	*
2000	12.8	*	*	*	2.2	20.8	32.2	25.3	10.0	5.8	*	*
1999	12.6	*	*	*	2.7	19.5	33.4	23.4	10.7	5.6	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malaria (B50-B54)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
All races 3/, both sexes												
2003	2.6	4.3	0.5	0.1	0.2	0.4	1.0	2.9	4.1	7.3	13.3	22.5
2002	2.3	5.6	0.4	0.2	0.2	0.4	1.0	2.3	3.3	6.7	12.5	21.7
2001	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0
All races, male												
2003	2.8	4.2	0.5	0.1	0.2	0.4	1.3	4.0	5.1	8.5	15.8	26.5
2002	2.6	6.1	0.5	0.2	0.2	0.5	1.3	3.2	4.2	8.0	15.4	26.8
2001	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9
All races, female												
2003	2.3	4.5	0.5	0.1	0.2	0.3	0.7	1.9	3.1	6.3	11.7	20.8
2002	2.1	5.0	0.3	0.2	0.2	0.4	0.7	1.4	2.6	5.6	10.6	19.5
2001	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
White, both sexes												
2003	2.6	3.5	0.4	0.1	0.2	0.3	0.9	2.8	3.8	7.3	13.0	23.0
2002	2.4	5.0	0.4	0.2	0.2	0.4	0.8	2.1	3.0	6.6	12.4	22.0
2001	2.0	3.8	0.4	0.2	0.2	0.3	0.7	1.6	2.8	6.0	10.5	20.1
2000	2.0	3.4	0.4	0.2	0.2	0.3	0.7	1.4	2.7	5.7	11.0	21.0
1999	2.5	4.5	0.5	0.2	0.2	0.4	1.0	1.9	3.4	7.5	13.7	26.5
White, male												
2003	2.8	3.2	0.4	*	0.2	0.4	1.2	3.8	4.8	8.3	15.5	27.2
2002	2.6	5.3	0.5	0.2	0.2	0.4	1.1	3.0	3.7	7.9	15.2	27.2
2001	2.1	3.9	0.4	0.1	0.2	0.4	0.9	2.0	3.4	6.6	13.1	23.8
2000	2.1	3.9	0.4	0.2	0.3	0.3	0.8	1.7	3.1	7.0	13.3	26.9
1999	2.6	5.6	0.6	0.2	0.2	0.5	1.3	2.3	3.8	8.8	16.3	30.6
White, female												
2003	2.4	3.9	0.5	0.1	0.1	0.3	0.6	1.8	2.9	6.4	11.3	21.2
2002	2.2	4.6	*	0.2	0.2	0.4	0.6	1.3	2.4	5.5	10.5	19.8
2001	1.9	3.8	*	0.2	0.2	0.2	0.5	1.1	2.2	5.6	8.7	18.5
2000	1.9	2.9	0.4	0.2	0.2	0.2	0.5	1.1	2.3	4.6	9.5	18.7
1999	2.5	3.4	*	0.2	0.2	0.4	0.6	1.5	3.0	6.4	12.0	24.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)												
Black, both sexes												
2003	2.5	8.0	1.0	*	*	0.6	1.8	4.3	6.1	7.0	16.2	19.5
2002	2.5	8.7	*	*	0.4	0.9	2.2	3.9	5.8	7.2	13.8	21.7
2001	2.2	9.5	*	*	0.4	0.8	1.6	2.9	5.1	8.3	11.1	20.6
2000	2.3	9.9	*	*	0.4	0.9	1.6	3.0	4.9	7.8	13.3	24.1
1999	3.5	9.9	0.9	*	0.3	1.6	3.0	5.4	7.2	11.8	20.5	34.1
Black, male												
2003	2.9	8.9	*	*	*	0.8	2.1	5.7	8.0	10.7	18.6	22.9
2002	2.9	9.6	*	*	*	1.1	2.6	5.5	8.4	8.8	16.9	28.9
2001	2.4	10.8	*	*	*	*	2.2	3.9	6.3	10.0	12.4	*
2000	2.5	9.9	*	*	*	0.9	2.1	3.9	5.7	10.2	18.2	32.3
1999	3.6	10.1	*	*	*	1.8	3.1	6.1	7.8	14.0	28.9	45.7
Black, female												
2003	2.2	7.1	*	*	*	*	1.4	3.1	4.6	4.3	14.8	18.2
2002	2.1	7.9	*	*	*	*	1.8	2.5	3.6	6.1	12.0	18.9
2001	2.0	8.2	*	*	*	0.9	1.1	2.1	4.2	7.2	10.4	20.2
2000	2.1	9.9	*	*	*	1.0	1.2	2.2	4.2	6.1	10.6	21.0
1999	3.4	9.7	*	*	*	1.4	2.9	4.8	6.7	10.3	15.8	29.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
All races 3/, both sexes												
2003	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
2002	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8
All races, male												
2003	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
2002	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2
All races, female												
2003	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
2002	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
White, both sexes												
2003	203.7	2.1	2.6	2.7	4.1	9.1	33.6	116.0	336.6	769.5	1,303.4	1,701.4
2002	205.5	1.8	2.5	2.7	4.4	9.5	34.5	118.5	343.3	790.1	1,311.0	1,725.7
2001	206.4	1.7	3.0	2.4	4.3	9.8	35.3	121.2	349.1	797.8	1,311.1	1,765.2
2000	208.6	2.5	2.8	2.6	4.3	9.7	34.5	121.8	359.4	811.0	1,330.3	1,815.7
1999	208.7	1.8	2.9	2.4	4.3	9.8	35.3	121.9	365.2	820.7	1,323.7	1,798.8
White, male												
2003	213.1	1.8	2.9	2.9	4.7	8.9	29.9	119.9	375.6	922.7	1,683.6	2,412.1
2002	215.5	2.2	2.7	2.8	5.0	9.1	30.5	121.8	386.0	954.8	1,695.3	2,486.8
2001	216.4	1.5	3.1	2.4	5.1	9.3	31.3	123.6	392.1	969.4	1,704.6	2,597.6
2000	218.1	2.8	3.1	2.8	5.0	9.2	30.9	123.5	401.9	984.3	1,736.0	2,693.7
1999	219.5	2.0	3.2	2.7	5.0	9.5	31.9	123.5	412.7	1,002.3	1,741.8	2,696.3
White, female												
2003	194.6	2.3	2.3	2.5	3.4	9.4	37.3	112.1	299.8	638.9	1,046.3	1,386.5
2002	195.8	1.4	2.3	2.5	3.7	9.9	38.5	115.3	303.1	650.4	1,053.1	1,395.1
2001	196.7	2.0	2.8	2.5	3.5	10.4	39.3	118.9	308.6	652.9	1,049.8	1,416.7
2000	199.4	2.3	2.4	2.3	3.6	10.1	38.2	120.1	319.7	665.6	1,063.4	1,459.1
1999	198.2	1.5	2.7	2.2	3.7	10.1	38.7	120.2	320.7	669.4	1,049.6	1,438.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms (C00-C97)												
Black, both sexes												
2003	164.3	*	2.2	2.6	3.9	12.2	48.9	185.6	471.0	925.4	1,487.9	1,853.1
2002	165.9	*	2.9	2.5	4.0	12.3	50.0	181.7	485.6	962.6	1,520.7	1,885.7
2001	167.1	*	1.9	2.6	4.3	12.9	51.4	184.4	486.1	991.2	1,543.9	1,931.7
2000	169.3	*	2.8	2.2	5.1	11.9	54.0	192.6	496.0	1,005.3	1,566.3	1,998.9
1999	171.3	*	2.0	2.7	5.5	12.2	53.9	192.9	521.4	1,019.7	1,579.3	2,008.4
Black, male												
2003	178.3	*	2.4	2.8	4.5	10.3	41.7	207.0	583.8	1,221.5	2,144.2	2,825.5
2002	181.5	*	2.6	3.1	4.5	11.2	43.0	197.3	610.3	1,274.7	2,223.0	2,976.1
2001	184.5	*	2.4	2.9	5.0	10.5	44.6	204.8	604.2	1,335.3	2,290.0	3,209.9
2000	188.5	*	2.7	2.5	6.1	10.1	48.4	214.2	626.4	1,363.8	2,351.8	3,264.8
1999	191.0	*	2.1	2.8	6.4	11.4	47.3	221.3	659.0	1,369.8	2,359.4	3,340.7
Black, female												
2003	151.4	*	2.1	2.3	3.3	13.9	55.4	167.2	380.4	714.6	1,116.9	1,475.3
2002	151.7	*	3.2	2.0	3.6	13.3	56.2	168.2	385.4	741.1	1,123.1	1,468.0
2001	151.3	*	*	2.2	3.6	15.0	57.3	166.8	390.9	748.4	1,125.0	1,457.5
2000	151.8	*	2.9	1.9	4.1	13.5	58.9	173.9	391.0	753.1	1,124.0	1,527.7
1999	153.3	*	1.9	2.6	4.6	13.0	59.8	168.5	411.1	772.7	1,142.6	1,510.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
All races 3/, both sexes												
2003	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
2002	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3
All races, male												
2003	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
2002	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1
All races, female												
2003	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
2002	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
White, both sexes												
2003	2.7	*	*	*	*	0.1	0.5	2.4	6.2	9.8	12.9	19.0
2002	2.7	*	*	*	*	0.1	0.5	2.4	6.3	10.2	13.7	18.9
2001	2.7	*	*	*	0.1	0.1	0.5	2.4	6.3	10.0	14.4	19.0
2000	2.7	*	*	*	*	0.1	0.5	2.4	6.0	10.2	13.1	19.7
1999	2.7	*	*	*	*	0.1	0.5	2.3	6.1	9.9	14.1	19.8
White, male												
2003	3.7	*	*	*	*	0.2	0.8	4.0	9.7	14.6	18.6	27.2
2002	3.7	*	*	*	*	0.2	0.7	3.8	9.9	15.3	19.7	27.0
2001	3.6	*	*	*	*	0.1	0.7	3.9	10.1	15.1	20.4	26.4
2000	3.5	*	*	*	*	0.1	0.7	3.7	9.6	14.9	18.7	27.6
1999	3.5	*	*	*	*	*	0.7	3.6	9.6	15.1	20.3	27.3
White, female												
2003	1.8	*	*	*	*	*	0.2	0.9	2.8	5.7	9.1	15.4
2002	1.8	*	*	*	*	*	0.3	1.0	2.8	5.9	9.6	15.3
2001	1.9	*	*	*	*	*	0.3	1.0	2.7	5.8	10.3	15.9
2000	1.9	*	*	*	*	*	0.3	1.2	2.7	6.3	9.4	16.5
1999	1.9	*	*	*	*	*	0.3	1.0	2.8	5.6	10.1	16.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)												
Black, both sexes												
2003	2.9	*	*	*	*	*	1.1	5.1	11.2	15.4	15.3	15.5
2002	2.8	*	*	*	*	*	1.1	5.5	12.1	12.9	11.8	16.0
2001	2.9	*	*	*	*	*	1.2	5.1	12.6	15.5	13.9	12.9
2000	3.0	*	*	*	*	*	1.1	6.0	13.5	15.5	15.1	15.3
1999	3.2	*	*	*	*	*	1.2	7.1	13.5	15.2	16.9	14.3
Black, male												
2003	4.6	*	*	*	*	*	1.2	9.0	21.1	28.7	27.9	25.0
2002	4.3	*	*	*	*	*	1.6	9.5	21.9	23.0	18.4	26.8
2001	4.4	*	*	*	*	*	1.7	8.7	23.3	27.1	23.5	*
2000	4.8	*	*	*	*	*	1.7	10.5	24.3	28.2	28.8	*
1999	4.8	*	*	*	*	*	1.8	12.1	23.2	25.1	31.7	24.6
Black, female												
2003	1.4	*	*	*	*	*	1.0	1.7	3.2	6.0	8.1	11.7
2002	1.4	*	*	*	*	*	0.7	2.0	4.2	5.8	8.1	11.9
2001	1.5	*	*	*	*	*	0.8	2.1	3.9	7.4	8.4	10.5
2000	1.4	*	*	*	*	*	*	2.0	4.8	6.5	7.3	14.2
1999	1.7	*	*	*	*	*	0.7	2.9	5.7	8.2	8.6	10.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
All races 3/, both sexes												
2003	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
2002	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8
All races, male												
2003	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
2002	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0
All races, female												
2003	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
2002	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
White, both sexes												
2003	4.7	*	*	*	*	0.1	0.7	3.2	10.2	19.6	26.7	25.8
2002	4.6	*	*	*	*	*	0.7	3.4	9.9	19.8	26.7	25.1
2001	4.6	*	*	*	*	0.1	0.7	3.4	9.4	19.8	26.5	26.3
2000	4.5	*	*	*	*	0.1	0.7	3.3	10.0	19.5	25.3	26.6
1999	4.4	*	*	*	*	0.1	0.6	3.2	9.8	19.4	24.2	27.2
White, male												
2003	7.4	*	*	*	*	0.1	1.1	5.7	18.0	33.9	48.8	46.7
2002	7.3	*	*	*	*	*	1.2	6.0	17.5	35.2	48.8	44.6
2001	7.2	*	*	*	*	0.1	1.1	5.9	16.4	34.9	47.9	48.8
2000	7.0	*	*	*	*	0.2	1.2	5.9	17.5	33.6	45.6	49.3
1999	6.9	*	*	*	*	0.1	1.1	5.7	17.0	34.4	44.0	48.0
White, female												
2003	2.0	*	*	*	*	*	0.2	0.8	2.7	7.5	11.7	16.5
2002	2.0	*	*	*	*	*	0.2	0.8	2.8	6.8	11.9	16.7
2001	2.1	*	*	*	*	*	0.2	0.9	2.7	7.1	12.3	16.8
2000	2.1	*	*	*	*	*	0.1	0.8	2.9	7.8	11.9	17.3
1999	2.0	*	*	*	*	*	0.1	0.7	3.1	6.9	11.2	18.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of esophagus (C15)												
Black, both sexes												
2003	4.2	*	*	*	*	*	0.8	6.0	16.9	25.6	27.9	21.3
2002	4.2	*	*	*	*	*	1.0	5.1	17.3	28.5	29.6	20.8
2001	4.5	*	*	*	*	*	0.9	5.9	18.5	32.3	29.5	23.6
2000	4.5	*	*	*	*	*	0.8	6.6	19.8	28.2	33.7	22.2
1999	4.6	*	*	*	*	*	0.9	6.5	20.0	29.9	31.1	23.9
Black, male												
2003	6.2	*	*	*	*	*	1.3	10.2	28.7	44.3	46.0	38.6
2002	6.3	*	*	*	*	*	1.4	8.3	30.1	49.2	51.9	37.5
2001	7.0	*	*	*	*	*	1.5	10.2	32.5	55.3	54.7	46.5
2000	6.7	*	*	*	*	*	1.2	11.2	34.0	47.0	58.4	38.1
1999	7.0	*	*	*	*	*	1.7	11.3	34.8	50.6	56.9	51.6
Black, female												
2003	2.3	*	*	*	*	*	*	2.5	7.4	12.2	17.7	14.6
2002	2.3	*	*	*	*	*	*	2.4	7.0	13.8	16.9	14.4
2001	2.3	*	*	*	*	*	*	2.2	7.3	16.1	15.4	15.2
2000	2.5	*	*	*	*	*	*	2.7	8.3	14.9	19.8	16.3
1999	2.3	*	*	*	*	*	*	2.4	8.2	15.2	16.7	13.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
All races 3/, both sexes												
2003	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
2002	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1
All races, male												
2003	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
2002	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3
All races, female												
2003	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
2002	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
White, both sexes												
2003	3.9	*	*	*	*	0.3	0.9	2.4	5.9	13.5	24.6	38.7
2002	4.0	*	*	*	0.1	0.3	0.8	2.3	5.9	13.4	25.4	42.2
2001	4.1	*	*	*	*	0.3	0.8	2.4	6.1	14.5	26.9	41.1
2000	4.2	*	*	*	*	0.2	0.9	2.5	6.5	14.5	27.3	47.3
1999	4.3	*	*	*	*	0.3	0.9	2.5	6.6	15.2	28.1	44.8
White, male												
2003	4.6	*	*	*	*	0.3	1.0	3.1	8.3	19.2	34.3	53.5
2002	4.8	*	*	*	*	0.4	0.9	3.1	8.3	19.7	36.6	60.5
2001	4.9	*	*	*	*	0.3	1.0	3.2	8.8	20.2	38.7	57.7
2000	5.0	*	*	*	*	0.3	1.0	3.2	9.3	20.9	39.0	68.9
1999	5.1	*	*	*	*	0.3	1.0	3.5	9.3	21.7	41.5	63.4
White, female												
2003	3.2	*	*	*	*	0.3	0.8	1.7	3.6	8.7	17.9	32.1
2002	3.2	*	*	*	*	0.2	0.7	1.5	3.7	8.0	17.9	34.2
2001	3.3	*	*	*	*	0.2	0.7	1.7	3.5	9.7	19.0	34.1
2000	3.5	*	*	*	*	0.2	0.7	1.7	3.9	9.2	19.6	38.6
1999	3.4	*	*	*	*	0.2	0.8	1.5	4.0	9.8	19.3	37.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of stomach (C16)												
Black, both sexes												
2003	5.4	*	*	*	*	0.4	1.8	5.7	12.7	27.1	54.3	85.8
2002	5.6	*	*	*	*	0.6	1.6	5.4	12.9	30.9	56.8	100.4
2001	5.6	*	*	*	*	0.5	2.1	5.4	12.9	30.5	58.4	88.1
2000	5.9	*	*	*	*	0.7	2.0	5.7	13.8	32.0	63.9	101.7
1999	6.1	*	*	*	*	0.7	2.2	5.9	15.9	31.1	64.6	102.4
Black, male												
2003	6.5	*	*	*	*	*	2.0	8.3	19.6	44.0	79.5	106.3
2002	6.6	*	*	*	*	0.8	1.8	7.7	18.9	46.7	79.8	137.2
2001	6.7	*	*	*	*	*	2.3	7.9	19.7	48.2	87.1	104.4
2000	7.0	*	*	*	*	0.8	2.9	7.9	20.6	46.7	91.1	136.1
1999	7.3	*	*	*	*	*	2.7	8.5	23.1	46.0	100.5	147.6
Black, female												
2003	4.3	*	*	*	*	*	1.6	3.5	7.1	15.0	40.0	77.8
2002	4.7	*	*	*	*	*	1.4	3.4	8.0	19.8	43.9	86.3
2001	4.5	*	*	*	*	*	1.9	3.2	7.5	18.0	42.3	82.1
2000	5.0	*	*	*	*	*	1.3	3.7	8.4	21.6	48.6	88.9
1999	5.0	*	*	*	*	*	1.8	3.7	10.1	20.6	44.4	85.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
All races 3/, both sexes												
2003	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
2002	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7
All races, male												
2003	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
2002	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0
All races, female												
2003	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
2002	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
White, both sexes												
2003	20.2	*	*	*	0.1	0.7	2.7	10.0	28.2	68.7	133.3	238.5
2002	20.7	*	*	*	0.1	0.7	2.9	10.7	30.1	71.8	135.2	240.7
2001	21.0	*	*	*	0.1	0.7	3.0	10.8	30.1	73.4	136.9	255.3
2000	21.5	*	*	*	0.1	0.6	2.7	10.4	31.2	75.1	144.0	269.2
1999	21.6	*	*	*	0.1	0.7	2.8	10.4	32.0	75.6	144.5	270.9
White, male												
2003	20.5	*	*	*	0.1	0.7	3.0	11.5	34.3	86.2	162.8	272.9
2002	21.1	*	*	*	0.1	0.8	3.3	12.2	37.6	89.8	164.2	273.6
2001	21.2	*	*	*	*	0.7	3.2	12.3	36.8	91.8	167.0	297.3
2000	21.6	*	*	*	*	0.7	3.1	12.0	37.6	93.4	176.9	313.3
1999	21.8	*	*	*	*	0.7	2.9	11.9	38.8	93.7	179.3	324.5
White, female												
2003	19.8	*	*	*	*	0.6	2.4	8.4	22.5	53.8	113.4	223.2
2002	20.3	*	*	*	*	0.7	2.6	9.1	23.1	56.4	115.7	226.4
2001	20.7	*	*	*	*	0.7	2.7	9.3	23.7	57.9	116.8	237.7
2000	21.4	*	*	*	*	0.5	2.4	9.0	25.2	59.7	122.3	251.3
1999	21.4	*	*	*	*	0.6	2.6	8.9	25.7	60.5	121.7	249.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of colon, rectum and anus (C18-C21)												
Black, both sexes												
2003	18.2	*	*	*	*	0.9	5.0	18.7	48.1	100.4	172.0	281.5
2002	18.2	*	*	*	*	0.7	4.9	18.6	49.5	99.2	183.6	274.7
2001	18.6	*	*	*	*	1.0	4.9	19.0	49.8	104.0	187.6	286.5
2000	18.8	*	*	*	*	1.0	5.1	19.5	50.1	104.7	190.1	310.0
1999	18.7	*	*	*	*	1.0	4.8	17.7	52.7	106.2	193.1	306.1
Black, male												
2003	18.3	*	*	*	*	1.0	5.2	22.0	58.9	125.8	213.0	317.0
2002	18.6	*	*	*	*	1.0	5.8	21.2	61.2	121.6	237.5	328.1
2001	18.4	*	*	*	*	1.2	5.7	20.8	58.4	128.1	228.4	347.2
2000	19.1	*	*	*	*	1.0	6.0	22.0	62.5	131.5	241.0	377.1
1999	18.6	*	*	*	*	1.2	4.7	20.1	65.8	126.6	244.4	350.4
Black, female												
2003	18.1	*	*	*	*	0.9	4.8	16.0	39.4	82.3	148.8	267.8
2002	17.8	*	*	*	*	*	4.1	16.3	40.2	83.3	153.0	254.2
2001	18.8	*	*	*	*	0.9	4.3	17.5	42.8	87.0	164.7	264.0
2000	18.5	*	*	*	*	1.0	4.3	17.2	40.2	85.8	161.4	285.0
1999	18.9	*	*	*	*	0.9	4.9	15.6	42.2	91.8	164.4	289.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
All races 3/, both sexes												
2003	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
2002	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6
All races, male												
2003	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
2002	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1
All races, female												
2003	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
2002	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
White, both sexes												
2003	4.9	*	*	*	0.1	0.1	0.7	4.6	8.7	18.3	29.5	32.1
2002	4.8	*	0.2	*	*	0.2	0.8	4.1	8.5	18.3	28.4	33.8
2001	4.6	*	*	*	0.1	0.2	0.6	4.0	8.2	17.6	27.9	33.3
2000	4.5	*	*	*	0.1	0.1	0.7	3.7	8.1	17.3	27.8	32.0
1999	4.4	*	0.2	*	0.1	0.1	0.8	3.4	7.6	17.8	27.2	29.9
White, male												
2003	6.4	*	*	*	*	0.1	1.0	7.4	13.1	26.0	43.0	45.6
2002	6.1	*	*	*	*	0.2	1.1	6.8	12.6	26.5	40.0	45.1
2001	5.9	*	*	*	*	0.1	0.9	6.4	12.2	25.4	39.6	49.7
2000	5.7	*	*	*	*	0.2	1.0	5.7	11.8	25.0	41.1	43.1
1999	5.5	*	*	*	*	0.1	1.1	5.2	11.1	25.8	38.0	40.6
White, female												
2003	3.5	*	*	*	*	0.1	0.5	1.8	4.6	11.7	20.3	26.2
2002	3.5	*	*	*	*	0.2	0.4	1.5	4.5	11.4	20.6	29.0
2001	3.3	*	*	*	*	0.2	0.3	1.6	4.4	10.9	20.1	26.5
2000	3.3	*	*	*	*	*	0.4	1.7	4.6	10.9	19.0	27.4
1999	3.3	*	*	*	*	*	0.5	1.6	4.4	11.0	20.1	25.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of liver and intrahepatic bile ducts (C22)												
Black, both sexes												
2003	4.9	*	*	*	*	0.5	1.4	9.8	16.3	22.9	31.3	41.7
2002	4.6	*	*	*	*	0.5	1.4	8.7	14.8	23.0	30.9	38.0
2001	4.5	*	*	*	*	*	1.7	8.7	14.8	21.7	32.4	35.0
2000	4.4	*	*	*	*	0.4	1.8	8.6	14.1	23.6	28.7	31.6
1999	4.2	*	*	*	*	0.4	1.6	7.2	13.2	24.2	29.4	34.4
Black, male												
2003	6.8	*	*	*	*	*	2.2	17.1	25.9	33.5	44.3	63.6
2002	6.5	*	*	*	*	*	2.2	15.1	24.2	37.7	47.0	41.8
2001	5.9	*	*	*	*	*	2.6	14.8	23.6	30.0	43.0	42.0
2000	6.1	*	*	*	*	*	2.4	14.8	23.5	37.3	39.7	48.4
1999	5.5	*	*	*	*	*	2.4	12.8	20.0	36.5	41.2	30.5
Black, female												
2003	3.2	*	*	*	*	*	*	3.5	8.6	15.3	23.9	33.2
2002	2.9	*	*	*	*	*	0.7	3.2	7.3	12.6	21.8	36.6
2001	3.2	*	*	*	*	*	0.8	3.5	7.7	15.9	26.4	32.4
2000	2.8	*	*	*	*	*	1.2	3.3	6.5	14.0	22.5	25.3
1999	3.0	*	*	*	*	*	0.9	2.4	7.8	15.6	22.7	35.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
All races 3/, both sexes												
2003	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
2002	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7
All races, male												
2003	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
2002	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5
All races, female												
2003	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
2002	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
White, both sexes												
2003	11.2	*	*	*	*	0.2	1.2	5.9	18.4	43.7	74.8	94.4
2002	11.1	*	*	*	*	0.1	1.3	5.8	18.7	43.7	74.3	94.3
2001	11.1	*	*	*	*	0.1	1.3	5.9	19.0	43.8	73.3	99.0
2000	11.0	*	*	*	*	0.1	1.2	6.1	18.9	43.1	75.3	94.7
1999	11.0	*	*	*	*	0.1	1.3	6.2	19.1	44.2	71.7	98.6
White, male												
2003	11.2	*	*	*	*	0.2	1.4	7.2	22.5	50.8	85.4	96.0
2002	11.1	*	*	*	*	*	1.6	7.5	23.0	50.7	84.2	101.6
2001	11.0	*	*	*	*	0.2	1.4	7.4	23.5	50.0	83.8	106.5
2000	10.9	*	*	*	*	0.2	1.4	7.8	23.0	49.8	85.2	102.4
1999	10.9	*	*	*	*	0.2	1.6	7.9	22.9	51.6	80.8	107.4
White, female												
2003	11.2	*	*	*	*	0.1	1.0	4.5	14.6	37.6	67.7	93.8
2002	11.0	*	*	*	*	*	1.0	4.2	14.5	37.8	67.7	91.1
2001	11.2	*	*	*	*	*	1.1	4.5	14.7	38.5	66.3	95.9
2000	11.1	*	*	*	*	*	0.9	4.4	15.0	37.5	68.8	91.5
1999	11.0	*	*	*	*	*	0.9	4.6	15.5	37.9	65.8	95.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of pancreas (C25)												
Black, both sexes												
2003	9.5	*	*	*	*	*	1.8	10.0	27.3	56.3	93.1	112.9
2002	9.3	*	*	*	*	*	1.8	8.7	27.6	60.7	89.9	106.6
2001	9.1	*	*	*	*	*	1.6	9.0	25.5	58.4	92.7	121.3
2000	9.3	*	*	*	*	*	1.7	9.0	27.5	59.1	97.8	119.8
1999	9.4	*	*	*	*	*	2.0	9.6	28.2	59.5	94.0	118.6
Black, male												
2003	9.2	*	*	*	*	*	2.4	13.3	33.2	63.2	97.9	104.3
2002	9.0	*	*	*	*	*	2.1	11.4	34.1	68.4	96.1	100.8
2001	8.6	*	*	*	*	*	1.7	11.1	30.3	68.1	100.3	99.8
2000	9.0	*	*	*	*	*	2.0	11.8	34.2	65.8	106.9	109.6
1999	9.4	*	*	*	*	*	2.4	13.1	36.5	67.9	106.7	114.8
Black, female												
2003	9.7	*	*	*	*	*	1.3	7.2	22.5	51.5	90.4	116.3
2002	9.5	*	*	*	*	*	1.4	6.4	22.5	55.3	86.4	108.8
2001	9.6	*	*	*	*	*	1.4	7.1	21.6	51.6	88.4	129.2
2000	9.7	*	*	*	*	*	1.4	6.6	22.1	54.4	92.7	123.6
1999	9.4	*	*	*	*	*	1.6	6.6	21.6	53.6	86.9	120.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
All races 3/, both sexes												
2003	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
2002	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0
All races, male												
2003	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
2002	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8
All races, female												
2003	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
2002	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
White, both sexes												
2003	1.3	*	*	*	*	*	0.2	0.9	3.1	5.8	6.6	6.2
2002	1.3	*	*	*	*	*	0.1	0.9	3.1	5.7	7.3	5.6
2001	1.3	*	*	*	*	*	0.2	0.9	3.2	6.0	6.9	6.6
2000	1.3	*	*	*	*	*	0.1	0.9	3.1	6.1	7.7	6.6
1999	1.4	*	*	*	*	*	0.1	0.9	3.4	6.4	7.2	6.7
White, male												
2003	2.1	*	*	*	*	*	0.3	1.5	5.2	9.8	12.9	14.5
2002	2.0	*	*	*	*	*	0.2	1.5	5.1	9.5	14.1	13.3
2001	2.1	*	*	*	*	*	0.3	1.4	5.4	10.3	13.3	18.3
2000	2.1	*	*	*	*	*	0.2	1.5	5.3	10.4	15.0	17.6
1999	2.1	*	*	*	*	*	0.2	1.6	5.7	10.7	13.8	16.8
White, female												
2003	0.5	*	*	*	*	*	*	0.3	1.1	2.4	2.4	2.5
2002	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.6	2.3
2001	0.5	*	*	*	*	*	*	0.3	1.2	2.3	2.7	1.7
2000	0.6	*	*	*	*	*	*	0.4	1.1	2.6	2.9	2.1
1999	0.6	*	*	*	*	*	*	0.3	1.2	2.8	2.9	2.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of larynx (C32)												
Black, both sexes												
2003	1.8	*	*	*	*	*	*	2.5	7.5	11.5	11.7	11.1
2002	1.8	*	*	*	*	*	0.5	2.6	7.5	10.3	13.7	*
2001	1.8	*	*	*	*	*	0.4	2.8	7.9	11.4	11.7	8.3
2000	2.0	*	*	*	*	*	*	2.9	9.0	12.8	14.1	6.9
1999	1.8	*	*	*	*	*	0.6	2.9	8.6	11.6	10.0	8.3
Black, male												
2003	3.1	*	*	*	*	*	*	4.5	13.6	23.8	27.9	27.1
2002	3.0	*	*	*	*	*	0.7	4.5	14.6	19.3	29.3	*
2001	3.1	*	*	*	*	*	*	5.1	14.2	21.3	27.7	*
2000	3.3	*	*	*	*	*	*	5.1	17.2	24.3	28.5	*
1999	3.1	*	*	*	*	*	0.9	5.2	15.9	21.8	22.8	24.6
Black, female												
2003	0.6	*	*	*	*	*	*	0.8	2.5	2.8	*	*
2002	0.7	*	*	*	*	*	*	1.1	1.8	4.0	4.9	*
2001	0.7	*	*	*	*	*	*	*	2.8	4.5	*	*
2000	0.7	*	*	*	*	*	*	1.0	2.4	4.6	6.0	*
1999	0.7	*	*	*	*	*	*	0.9	2.7	4.4	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
All races 3/, both sexes												
2003	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
2002	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0
All races, male												
2003	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
2002	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2
All races, female												
2003	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
2002	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
White, both sexes												
2003	58.7	*	*	*	0.1	0.4	5.4	28.9	110.8	274.2	382.6	302.4
2002	59.1	*	*	*	*	0.4	5.8	29.1	114.5	280.0	382.9	300.9
2001	59.1	*	*	*	*	0.4	6.0	29.6	117.8	283.4	374.7	305.4
2000	59.5	*	*	*	*	0.5	5.7	30.2	122.5	287.0	374.7	304.5
1999	58.5	*	*	*	*	0.5	5.8	29.7	122.9	283.5	363.8	298.8
White, male												
2003	66.9	*	*	*	*	0.4	5.7	34.1	133.1	347.1	524.6	478.5
2002	67.7	*	*	*	*	0.4	5.8	34.2	139.3	356.4	527.8	486.6
2001	68.6	*	*	*	*	0.4	6.3	34.9	143.8	368.1	526.4	513.9
2000	69.4	*	*	*	*	0.5	6.1	35.7	150.8	374.9	529.9	522.4
1999	69.1	*	*	*	*	0.5	6.1	35.2	153.9	373.4	523.5	531.5
White, female												
2003	50.7	*	*	*	*	0.4	5.1	23.8	89.8	212.2	286.6	224.4
2002	50.7	*	*	*	*	0.4	5.8	24.2	91.0	215.1	285.8	220.2
2001	49.8	*	*	*	*	0.4	5.7	24.3	93.3	211.9	274.0	218.1
2000	49.9	*	*	*	*	0.5	5.2	24.8	96.1	213.2	272.7	215.9
1999	48.2	*	*	*	*	0.4	5.4	24.4	93.8	208.6	259.1	205.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of trachea, bronchus and lung (C33-C34)												
Black, both sexes												
2003	42.9	*	*	*	*	*	7.7	47.4	144.0	288.6	383.1	280.9
2002	43.2	*	*	*	*	0.4	8.7	46.9	152.4	295.3	381.3	272.3
2001	43.1	*	*	*	*	*	8.9	46.5	147.0	306.5	385.4	280.7
2000	43.9	*	*	*	*	0.6	10.1	50.4	152.5	315.8	375.0	288.1
1999	44.1	*	*	*	*	0.4	9.3	50.6	158.1	313.1	387.9	290.5
Black, male												
2003	54.9	*	*	*	*	*	9.3	63.6	210.0	427.1	622.2	502.5
2002	56.0	*	*	*	*	*	10.6	64.6	223.6	444.6	626.2	484.6
2001	57.2	*	*	*	*	*	11.0	66.2	215.2	476.4	659.8	549.2
2000	58.3	*	*	*	*	*	13.4	70.7	223.5	488.8	642.5	562.8
1999	58.9	*	*	*	*	*	11.3	72.6	237.6	484.4	661.0	574.2
Black, female												
2003	32.0	*	*	*	*	*	6.2	33.4	91.0	190.0	248.0	194.8
2002	31.6	*	*	*	*	*	7.0	31.7	95.3	189.3	242.6	191.0
2001	30.2	*	*	*	*	*	7.1	29.5	92.0	186.7	231.3	181.0
2000	30.8	*	*	*	*	*	7.3	32.9	95.3	194.1	224.3	185.9
1999	30.7	*	*	*	*	*	7.6	31.9	94.4	192.3	234.9	184.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
All races 3/, both sexes												
2003	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
2002	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4
All races, male												
2003	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
2002	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5
All races, female												
2003	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
2002	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
White, both sexes												
2003	3.2	*	*	*	0.1	0.6	1.5	3.4	6.3	9.8	16.0	19.1
2002	3.1	*	*	*	0.1	0.6	1.6	3.3	5.8	9.9	15.1	19.2
2001	3.2	*	*	*	0.1	0.6	1.7	3.6	6.1	10.3	14.6	19.3
2000	3.1	*	*	*	0.1	0.6	1.6	3.7	6.1	9.9	14.3	20.3
1999	3.1	*	*	*	0.1	0.6	1.7	3.6	6.1	10.0	14.5	17.6
White, male												
2003	4.2	*	*	*	0.1	0.7	1.8	4.4	8.8	14.2	25.5	32.9
2002	4.0	*	*	*	*	0.7	1.9	4.4	8.1	14.4	23.3	31.7
2001	4.1	*	*	*	0.1	0.6	2.0	4.6	8.4	15.5	23.3	32.7
2000	4.0	*	*	*	0.1	0.7	1.9	4.8	8.5	14.3	22.5	34.7
1999	4.0	*	*	*	0.1	0.6	2.1	4.6	8.5	14.7	22.6	29.7
White, female												
2003	2.3	*	*	*	*	0.5	1.2	2.4	4.0	6.1	9.5	12.9
2002	2.3	*	*	*	0.1	0.5	1.3	2.3	3.6	6.1	9.6	13.8
2001	2.3	*	*	*	*	0.5	1.3	2.7	3.8	5.9	8.8	13.7
2000	2.3	*	*	*	*	0.5	1.4	2.6	3.9	6.3	8.8	14.5
1999	2.3	*	*	*	*	0.5	1.2	2.6	3.9	6.1	9.2	12.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant melanoma of skin (C43)												
Black, both sexes												
2003	0.3	*	*	*	*	*	*	*	0.8	1.5	2.9	*
2002	0.3	*	*	*	*	*	*	*	0.8	1.5	*	*
2001	0.3	*	*	*	*	*	*	*	*	2.2	2.6	*
2000	0.4	*	*	*	*	*	*	*	0.8	1.8	3.5	6.6
1999	0.3	*	*	*	*	*	*	*	*	1.5	2.8	*
Black, male												
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	2.0	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
All races 3/, both sexes												
2003	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
2002	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4
All races, male												
2003	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
2002	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5
All races, female												
2003	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
2002	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
White, both sexes												
2003	15.0	*	*	*	*	0.9	5.5	14.4	28.4	45.1	75.1	132.5
2002	15.1	*	*	*	*	0.9	5.3	14.9	28.5	46.2	76.6	135.3
2001	15.2	*	*	*	*	1.0	5.6	15.7	29.1	47.0	77.1	134.8
2000	15.7	*	*	*	*	1.0	5.7	15.8	30.2	49.2	79.4	147.3
1999	15.5	*	*	*	*	0.9	5.4	15.7	29.9	49.0	79.0	146.0
White, male												
2003	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.0	3.9
2002	0.3	*	*	*	*	*	*	0.2	0.5	0.9	2.1	3.3
2001	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.3
2000	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	3.9
1999	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.0	4.4
White, female												
2003	29.3	*	*	*	*	1.8	11.1	28.4	54.9	82.6	124.6	189.4
2002	29.5	*	*	*	*	1.8	10.7	29.4	55.0	84.6	126.5	192.6
2001	29.8	*	*	*	*	2.0	11.3	31.0	56.0	85.6	126.9	189.9
2000	30.7	*	*	*	*	2.1	11.3	31.2	57.9	89.3	130.2	205.5
1999	30.2	*	*	*	*	1.9	10.8	30.9	57.5	88.9	129.5	202.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of breast (C50)												
Black, both sexes												
2003	15.1	*	*	*	*	2.2	11.1	26.5	45.4	59.3	90.8	161.6
2002	14.9	*	*	*	*	2.3	11.7	27.0	43.1	60.1	94.0	152.6
2001	15.0	*	*	*	*	2.6	11.1	27.3	44.6	63.5	92.3	152.0
2000	14.8	*	*	*	*	2.3	11.2	27.8	45.4	58.6	90.5	176.4
1999	15.0	*	*	*	*	2.3	11.6	28.7	46.3	63.3	94.5	147.3
Black, male												
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	28.5	*	*	*	*	4.1	20.8	48.9	81.1	100.1	141.0	222.4
2002	28.2	*	*	*	*	4.3	22.0	49.8	76.6	101.1	145.0	209.1
2001	28.3	*	*	*	*	5.0	20.8	50.3	79.8	106.6	141.8	205.9
2000	27.9	*	*	*	*	4.2	20.9	51.5	80.9	98.6	139.8	238.7
1999	28.2	*	*	*	*	4.3	21.7	53.0	82.2	106.2	145.4	199.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
All races 3/, both sexes												
2003	1.3	*	*	*	*	0.5	1.4	2.2	2.8	3.1	4.3	5.7
2002	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	2.7	*	*	*	*	1.0	2.8	4.3	5.5	5.6	7.2	8.3
2002	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
White, both sexes												
2003	1.2	*	*	*	*	0.4	1.3	2.0	2.6	2.6	3.6	4.9
2002	1.3	*	*	*	*	0.5	1.4	2.0	2.4	2.9	3.6	5.1
2001	1.3	*	*	*	*	0.5	1.5	2.2	2.6	2.7	4.0	6.2
2000	1.4	*	*	*	*	0.5	1.4	2.2	2.7	3.1	4.3	6.3
1999	1.4	*	*	*	*	0.5	1.4	2.4	2.8	3.1	4.0	6.8
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	2.5	*	*	*	*	0.9	2.7	4.0	5.0	4.8	6.0	7.1
2002	2.5	*	*	*	*	1.0	2.8	4.0	4.7	5.3	6.0	7.3
2001	2.7	*	*	*	*	1.1	3.0	4.5	5.0	5.0	6.6	8.8
2000	2.7	*	*	*	*	1.1	2.8	4.3	5.2	5.7	7.1	8.8
1999	2.8	*	*	*	*	1.1	2.9	4.7	5.4	5.7	6.7	9.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of cervix uteri (C53)												
Black, both sexes												
2003	2.1	*	*	*	*	0.9	2.2	3.6	5.5	7.4	11.3	14.6
2002	2.2	*	*	*	*	0.8	2.5	4.1	5.0	9.0	10.8	17.5
2001	2.1	*	*	*	*	0.8	2.4	3.8	5.2	7.8	11.9	19.0
2000	2.4	*	*	*	*	0.9	2.6	4.3	6.3	9.5	11.9	19.7
1999	2.3	*	*	*	*	0.7	2.5	3.4	7.1	10.7	11.3	21.7
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	4.0	*	*	*	*	1.7	4.2	6.8	10.0	12.6	17.7	20.3
2002	4.2	*	*	*	*	1.5	4.7	7.6	8.9	15.4	16.9	24.2
2001	4.0	*	*	*	*	1.5	4.5	7.0	9.3	13.3	18.5	26.1
2000	4.5	*	*	*	*	1.8	4.9	7.9	11.3	16.3	18.6	27.0
1999	4.4	*	*	*	*	1.2	4.7	6.4	12.8	18.3	17.6	29.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
All races 3/, both sexes												
2003	2.4	*	*	*	*	0.1	0.3	1.3	4.7	10.1	15.2	22.8
2002	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	4.7	*	*	*	*	0.1	0.7	2.6	9.0	18.5	25.3	33.0
2002	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
White, both sexes												
2003	2.4	*	*	*	*	*	0.3	1.3	4.3	9.1	14.4	22.7
2002	2.4	*	*	*	*	*	0.3	1.3	4.1	8.9	14.4	23.8
2001	2.4	*	*	*	*	0.1	0.3	1.4	4.0	8.8	15.7	24.5
2000	2.3	*	*	*	*	0.1	0.3	1.3	4.3	8.5	14.8	24.4
1999	2.4	*	*	*	*	*	0.3	1.3	4.0	8.9	15.8	23.4
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	4.7	*	*	*	*	*	0.6	2.5	8.3	16.9	24.1	32.8
2002	4.7	*	*	*	*	*	0.6	2.6	8.0	16.5	24.1	34.2
2001	4.8	*	*	*	*	0.2	0.6	2.7	7.8	16.2	26.2	34.7
2000	4.6	*	*	*	*	0.1	0.6	2.6	8.3	15.7	24.6	34.3
1999	4.6	*	*	*	*	*	0.6	2.5	7.7	16.3	26.1	32.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)												
Black, both sexes												
2003	2.9	*	*	*	*	*	0.5	1.8	9.0	21.0	27.8	30.0
2002	3.0	*	*	*	*	*	0.4	1.9	8.4	22.3	28.3	38.9
2001	2.8	*	*	*	*	*	0.4	2.1	7.7	21.2	28.4	32.2
2000	2.9	*	*	*	*	*	0.4	1.6	8.3	22.7	30.5	33.2
1999	2.7	*	*	*	*	*	0.5	1.4	7.9	21.3	27.3	35.4
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	5.5	*	*	*	*	*	1.0	3.3	16.2	36.0	43.5	41.7
2002	5.7	*	*	*	*	*	0.7	3.6	15.2	38.1	44.3	53.8
2001	5.4	*	*	*	*	*	0.7	3.9	13.9	36.2	44.3	44.2
2000	5.5	*	*	*	*	*	0.7	3.0	14.9	38.6	47.7	45.5
1999	5.2	*	*	*	*	*	0.9	2.6	14.3	36.3	42.5	48.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
All races 3/, both sexes												
2003	5.0	*	*	*	0.1	0.2	1.0	4.0	10.4	19.5	32.6	38.4
2002	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	9.9	*	*	*	0.1	0.4	2.0	7.8	20.0	35.7	54.4	55.4
2002	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
White, both sexes												
2003	5.6	*	*	*	0.1	0.2	1.0	4.1	10.8	20.0	33.6	39.1
2002	5.6	*	*	*	0.1	0.2	1.1	4.5	10.7	20.4	34.3	39.0
2001	5.6	*	*	*	*	0.3	1.1	4.4	10.5	20.6	34.2	40.4
2000	5.5	*	*	*	*	0.3	1.1	4.4	10.5	20.1	34.3	42.0
1999	5.4	*	*	*	0.1	0.2	1.1	4.4	10.6	20.1	32.4	38.7
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	11.0	*	*	*	0.1	0.5	2.1	8.1	21.0	37.1	56.3	56.4
2002	11.1	*	*	*	0.2	0.5	2.2	8.9	20.8	37.7	57.4	55.9
2001	11.0	*	*	*	*	0.5	2.1	8.6	20.5	38.0	56.8	57.3
2000	10.9	*	*	*	*	0.5	2.2	8.7	20.4	37.0	56.8	59.0
1999	10.6	*	*	*	0.1	0.4	2.3	8.7	20.5	36.9	53.6	54.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of ovary (C56)												
Black, both sexes												
2003	3.1	*	*	*	*	*	1.0	3.8	8.6	17.7	27.5	36.8
2002	3.1	*	*	*	*	*	1.0	3.4	9.1	17.1	28.8	34.7
2001	3.1	*	*	*	*	*	1.1	3.0	10.0	18.3	26.9	36.5
2000	2.9	*	*	*	*	*	1.0	3.3	8.4	18.1	26.6	34.4
1999	3.0	*	*	*	*	*	0.8	3.5	9.1	18.5	28.7	28.7
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	6.0	*	*	*	*	*	1.9	7.1	15.5	30.2	43.1	51.0
2002	5.9	*	*	*	*	*	1.9	6.4	16.4	29.3	45.2	48.1
2001	5.9	*	*	*	*	*	2.1	5.7	18.0	31.2	42.0	50.1
2000	5.6	*	*	*	*	*	2.0	6.2	15.2	30.8	41.6	47.2
1999	5.7	*	*	*	*	*	1.5	6.6	16.4	31.5	44.8	39.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
All races 3/, both sexes												
2003	10.2	*	*	*	*	*	0.1	1.0	7.4	32.9	95.5	184.9
2002	10.6	*	*	*	*	*	0.1	1.0	7.6	34.5	100.1	195.0
2001	10.8	*	*	*	*	*	0.1	1.1	7.5	35.7	103.2	200.9
2000	11.0	*	*	*	*	*	0.1	1.0	7.9	37.0	105.7	209.3
1999	11.4	*	*	*	*	*	0.1	1.0	8.2	39.7	109.9	207.5
All races, male												
2003	20.7	*	*	*	*	*	0.1	2.1	15.4	72.3	238.3	603.3
2002	21.5	*	*	*	*	*	0.1	1.9	15.8	75.9	251.0	644.5
2001	22.0	*	*	*	*	*	0.1	2.1	15.7	78.9	260.2	681.2
2000	22.5	*	*	*	*	*	0.1	2.0	16.6	82.0	267.9	723.1
1999	23.2	*	*	*	*	*	0.1	2.1	17.1	88.2	279.1	722.4
All races, female												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
White, both sexes												
2003	10.3	*	*	*	*	*	*	0.9	6.3	29.5	88.5	177.5
2002	10.6	*	*	*	*	*	0.1	0.8	6.4	30.6	92.6	186.7
2001	10.8	*	*	*	*	*	*	0.9	6.4	31.9	95.1	189.4
2000	11.0	*	*	*	*	*	*	0.8	6.8	32.5	97.0	198.6
1999	11.4	*	*	*	*	*	0.1	0.9	6.9	35.3	101.9	195.2
White, male												
2003	20.7	*	*	*	*	*	*	1.7	13.0	64.2	219.4	578.2
2002	21.5	*	*	*	*	*	0.1	1.5	13.2	66.7	230.7	616.3
2001	21.9	*	*	*	*	*	*	1.8	13.3	69.6	238.2	641.8
2000	22.3	*	*	*	*	*	*	1.7	13.9	71.2	244.5	687.5
1999	23.1	*	*	*	*	*	0.1	1.7	14.2	77.7	257.4	681.0
White, female												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)												
Black, both sexes												
2003	12.8	*	*	*	*	*	*	2.6	19.6	73.8	196.2	301.1
2002	13.6	*	*	*	*	*	*	2.6	19.4	81.2	210.5	328.4
2001	14.2	*	*	*	*	*	*	2.8	19.2	82.9	219.7	362.9
2000	14.6	*	*	*	*	*	*	2.4	20.3	90.1	229.8	356.6
1999	14.8	*	*	*	*	*	*	2.6	21.6	93.3	225.8	366.4
Black, male												
2003	26.9	*	*	*	*	*	*	5.6	43.9	177.6	543.2	1,076.0
2002	28.6	*	*	*	*	*	*	5.5	43.5	195.5	582.4	1,185.7
2001	29.7	*	*	*	*	*	*	6.0	43.1	200.4	610.9	1,341.2
2000	30.7	*	*	*	*	*	*	5.2	45.6	218.2	638.0	1,314.7
1999	31.2	*	*	*	*	*	*	5.7	48.5	225.6	629.0	1,346.3
Black, female												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
All races 3/, both sexes												
2003	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
2002	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4
All races, male												
2003	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
2002	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4
All races, female												
2003	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
2002	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
White, both sexes												
2003	4.6	*	*	0.1	*	0.1	0.7	3.0	9.1	17.5	27.7	33.8
2002	4.6	*	*	0.1	*	0.1	0.7	3.3	9.1	18.3	26.8	33.2
2001	4.6	*	*	0.1	*	0.1	0.7	3.4	9.4	18.2	26.7	32.8
2000	4.6	*	*	*	*	0.1	0.7	3.4	9.3	17.5	27.0	34.5
1999	4.3	*	*	0.1	*	0.1	0.7	3.1	9.1	17.1	24.9	31.5
White, male												
2003	5.9	*	*	*	*	0.1	1.0	4.4	13.6	24.3	40.4	50.0
2002	5.9	*	*	*	*	*	1.0	4.9	13.3	25.9	38.6	51.9
2001	5.8	*	*	*	*	*	1.0	5.0	13.6	26.0	37.7	54.4
2000	5.7	*	*	*	*	*	1.0	4.9	13.4	24.4	38.7	56.6
1999	5.4	*	*	*	*	0.1	1.0	4.5	13.3	23.9	36.3	49.8
White, female												
2003	3.4	*	*	*	*	*	0.4	1.7	4.9	11.6	19.1	26.6
2002	3.4	*	*	*	*	*	0.4	1.8	5.1	11.9	18.9	25.1
2001	3.4	*	*	*	*	*	0.4	1.8	5.4	11.6	19.4	23.8
2000	3.4	*	*	*	*	*	0.4	1.9	5.4	11.6	19.2	25.5
1999	3.2	*	*	*	*	*	0.4	1.8	5.3	11.3	17.5	24.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of kidney and renal pelvis (C64-C65)												
Black, both sexes												
2003	2.9	*	*	*	*	*	1.1	3.2	9.5	14.6	23.8	33.8
2002	2.8	*	*	*	*	*	0.8	3.5	8.6	15.4	26.1	28.2
2001	3.0	*	*	*	*	*	1.2	3.7	9.5	15.0	25.6	29.8
2000	2.9	*	*	*	*	*	0.9	3.4	8.8	16.5	25.6	29.7
1999	2.9	*	*	*	*	0.4	1.1	3.8	8.6	16.1	23.5	31.2
Black, male												
2003	3.7	*	*	*	*	*	1.6	4.9	13.8	22.2	37.0	43.8
2002	3.5	*	*	*	*	*	1.2	4.8	13.9	21.6	36.1	39.7
2001	3.8	*	*	*	*	*	1.5	5.4	14.9	22.3	40.3	46.5
2000	3.6	*	*	*	*	*	1.0	4.9	13.1	26.3	38.2	49.6
1999	3.6	*	*	*	*	*	1.6	5.8	13.2	23.5	32.0	49.2
Black, female												
2003	2.2	*	*	*	*	*	0.7	1.8	6.0	9.2	16.4	30.0
2002	2.2	*	*	*	*	*	*	2.4	4.2	11.0	20.5	23.8
2001	2.2	*	*	*	*	*	1.0	2.2	5.2	9.9	17.4	23.6
2000	2.1	*	*	*	*	*	0.8	2.2	5.2	9.6	18.6	22.3
1999	2.2	*	*	*	*	*	*	2.1	5.0	10.9	18.8	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
All races 3/, both sexes												
2003	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
2002	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5
All races, male												
2003	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
2002	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8
All races, female												
2003	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
2002	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
White, both sexes												
2003	4.8	*	*	*	*	*	0.2	1.3	5.3	15.7	37.5	66.6
2002	4.9	*	*	*	*	*	0.3	1.3	5.1	16.5	38.1	67.7
2001	4.8	*	*	*	*	*	0.3	1.2	5.2	16.1	36.7	70.8
2000	4.8	*	*	*	*	*	0.3	1.2	5.3	16.2	36.4	72.0
1999	4.8	*	*	*	*	*	0.2	1.3	5.4	16.3	36.3	69.8
White, male												
2003	6.8	*	*	*	*	*	0.3	2.0	8.3	25.3	65.7	131.0
2002	6.8	*	*	*	*	*	0.4	1.9	8.2	26.5	65.3	134.1
2001	6.8	*	*	*	*	*	0.3	1.9	8.3	26.6	65.0	141.1
2000	6.7	*	*	*	*	*	0.3	1.9	8.4	26.5	64.7	145.7
1999	6.7	*	*	*	*	*	0.3	1.9	8.5	26.9	64.8	140.7
White, female												
2003	2.9	*	*	*	*	*	0.1	0.5	2.4	7.6	18.4	38.1
2002	3.0	*	*	*	*	*	0.2	0.7	2.3	8.1	19.8	38.9
2001	2.9	*	*	*	*	*	0.3	0.6	2.2	7.2	17.8	41.3
2000	2.9	*	*	*	*	*	0.2	0.6	2.5	7.5	17.8	42.1
1999	2.9	*	*	*	*	*	0.2	0.6	2.4	7.6	17.7	41.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of bladder (C67)												
Black, both sexes												
2003	2.4	*	*	*	*	*	0.4	1.6	3.9	12.0	32.3	53.4
2002	2.5	*	*	*	*	*	*	1.1	4.5	14.3	31.0	63.3
2001	2.4	*	*	*	*	*	0.4	1.3	4.9	11.9	31.2	55.0
2000	2.3	*	*	*	*	*	*	1.0	4.1	13.3	34.7	49.7
1999	2.5	*	*	*	*	*	*	1.5	5.6	15.0	29.1	60.3
Black, male												
2003	2.8	*	*	*	*	*	*	2.2	5.4	17.5	48.8	80.3
2002	2.7	*	*	*	*	*	*	1.2	5.6	18.3	47.6	91.1
2001	2.5	*	*	*	*	*	*	1.7	6.4	16.1	43.0	78.3
2000	2.7	*	*	*	*	*	*	1.4	5.8	17.6	52.7	88.8
1999	2.9	*	*	*	*	*	*	2.6	7.4	19.1	46.1	86.7
Black, female												
2003	2.1	*	*	*	*	*	*	1.2	2.8	8.1	23.0	42.9
2002	2.3	*	*	*	*	*	*	0.9	3.7	11.6	21.6	52.6
2001	2.2	*	*	*	*	*	*	0.8	3.8	8.9	24.6	46.3
2000	2.0	*	*	*	*	*	*	*	2.7	10.3	24.5	35.2
1999	2.2	*	*	*	*	*	*	*	4.2	12.1	19.6	50.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
All races 3/, both sexes												
2003	4.4	*	0.7	0.8	0.6	1.0	2.4	4.8	9.7	15.5	19.5	15.2
2002	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0
All races, male												
2003	5.0	*	0.7	0.8	0.6	1.1	2.9	6.2	11.4	18.6	24.6	20.0
2002	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7
All races, female												
2003	3.9	*	0.6	0.8	0.5	0.9	2.0	3.5	8.1	12.8	16.1	13.1
2002	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
White, both sexes												
2003	5.0	*	0.7	0.9	0.6	1.1	2.7	5.2	10.5	16.6	20.7	15.7
2002	5.0	*	0.7	0.8	0.5	1.1	2.5	5.6	10.4	17.4	20.1	17.1
2001	5.0	*	0.7	0.8	0.6	1.2	2.6	5.4	10.5	16.5	21.6	17.0
2000	5.1	*	0.7	0.8	0.6	1.1	2.6	5.6	10.4	17.8	21.0	17.5
1999	5.2	*	0.8	0.8	0.6	1.3	2.6	5.6	11.1	18.2	21.4	16.6
White, male												
2003	5.6	*	0.8	0.9	0.7	1.3	3.2	6.7	12.3	19.8	25.8	20.7
2002	5.7	*	0.7	0.9	0.7	1.2	3.1	6.8	12.4	21.6	24.8	21.2
2001	5.7	*	0.8	0.7	0.7	1.5	3.2	6.7	12.9	20.5	27.1	20.5
2000	5.7	*	0.7	0.9	0.7	1.4	3.4	6.8	12.8	21.7	25.5	23.5
1999	5.8	*	0.8	0.8	0.6	1.5	3.1	6.7	13.3	22.3	26.8	21.8
White, female												
2003	4.4	*	0.6	0.9	0.5	1.0	2.3	3.8	8.7	13.9	17.2	13.4
2002	4.4	*	0.6	0.8	0.4	1.0	1.9	4.5	8.5	13.8	16.9	15.4
2001	4.4	*	0.7	0.8	0.5	0.9	1.9	4.3	8.2	13.0	18.0	15.6
2000	4.4	*	0.6	0.7	0.5	0.9	1.8	4.5	8.2	14.5	18.0	15.0
1999	4.6	*	0.8	0.8	0.5	1.0	2.1	4.5	9.1	14.8	17.9	14.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)												
Black, both sexes												
2003	2.1	*	*	0.7	0.5	0.7	1.4	3.1	5.5	8.9	10.5	10.5
2002	2.1	*	1.1	0.8	0.5	0.6	1.2	2.7	5.6	8.7	11.8	9.5
2001	2.0	*	*	0.6	0.5	0.5	1.1	3.0	5.8	9.2	9.9	11.7
2000	2.1	*	*	0.8	0.6	0.7	1.6	2.9	5.7	8.1	11.9	10.6
1999	2.1	*	*	0.8	0.6	0.7	1.3	3.2	6.0	8.2	11.8	9.9
Black, male												
2003	2.5	*	*	0.8	0.7	*	2.1	4.0	6.7	12.0	15.2	*
2002	2.3	*	*	1.0	*	0.8	1.3	2.7	8.1	10.4	16.4	*
2001	2.2	*	*	0.7	*	*	1.2	3.6	7.9	10.1	12.4	*
2000	2.4	*	*	0.9	0.7	1.1	2.0	3.7	7.1	9.8	13.0	*
1999	2.3	*	*	0.7	0.8	0.9	1.8	4.4	7.2	10.0	12.6	*
Black, female												
2003	1.8	*	*	0.6	*	*	0.9	2.2	4.6	6.7	7.8	11.7
2002	1.9	*	*	*	*	*	1.1	2.8	3.7	7.4	9.3	10.3
2001	1.8	*	*	*	*	*	1.0	2.4	4.2	8.5	8.6	9.7
2000	1.9	*	*	0.6	*	*	1.2	2.2	4.6	6.9	11.3	10.7
1999	1.9	*	*	0.9	*	*	1.0	2.1	5.0	6.9	11.4	9.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
All races 3/, both sexes												
2003	19.1	0.8	0.9	1.0	1.5	2.2	3.9	9.3	26.8	70.3	143.3	198.0
2002	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4
All races, male												
2003	21.0	*	1.0	1.1	1.8	2.6	4.7	11.4	33.9	90.4	184.8	268.7
2002	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0
All races, female												
2003	17.3	1.1	0.8	0.8	1.2	1.7	3.1	7.2	20.1	53.4	115.6	166.8
2002	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
White, both sexes												
2003	20.9	0.9	0.9	1.0	1.5	2.1	3.8	9.0	27.0	71.5	146.9	204.5
2002	21.3	*	0.8	1.0	1.8	2.2	3.9	9.6	28.0	75.4	147.6	204.0
2001	21.5	0.7	1.1	0.9	1.8	2.4	4.0	10.2	29.2	75.8	148.7	205.9
2000	21.9	1.2	1.0	1.0	1.8	2.2	4.0	10.4	30.1	78.0	153.3	208.7
1999	21.9	0.8	1.0	0.9	1.8	2.3	4.2	10.7	31.5	79.3	151.3	204.1
White, male												
2003	23.1	*	1.1	1.1	1.8	2.6	4.6	11.1	34.2	92.8	191.0	279.5
2002	23.4	*	0.9	1.0	2.1	2.6	4.5	11.6	34.8	97.3	193.7	279.4
2001	23.3	*	1.1	1.0	2.1	2.8	4.8	12.6	35.9	96.3	191.8	284.8
2000	23.6	*	1.0	1.1	2.2	2.6	4.8	12.6	36.1	98.7	198.2	290.2
1999	23.6	*	1.0	1.0	2.2	2.6	5.0	13.0	38.6	99.1	193.4	285.8
White, female												
2003	18.8	1.4	0.8	0.8	1.2	1.6	3.0	7.0	20.1	53.4	117.2	171.3
2002	19.2	*	0.8	0.9	1.4	1.8	3.2	7.5	21.5	56.8	116.7	171.2
2001	19.7	*	1.1	0.8	1.4	2.0	3.2	7.9	22.8	58.5	120.1	172.8
2000	20.3	*	1.0	0.8	1.3	1.8	3.2	8.3	24.4	60.6	123.8	175.5
1999	20.4	*	0.9	0.7	1.5	2.0	3.3	8.4	24.9	62.8	123.7	171.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)												
Black, both sexes												
2003	13.4	*	*	0.8	1.4	2.8	5.2	13.0	31.1	73.1	126.1	150.5
2002	13.6	*	*	0.8	1.5	2.9	4.8	13.6	35.9	76.0	122.3	152.3
2001	14.1	*	*	1.0	1.8	3.4	5.6	13.2	36.7	78.0	126.9	155.7
2000	13.9	*	0.8	0.6	2.2	2.9	5.3	14.7	35.8	75.0	121.0	175.5
1999	14.1	*	*	0.9	2.3	2.9	5.5	13.8	38.9	76.2	126.8	162.0
Black, male												
2003	13.9	*	*	0.9	1.6	3.4	6.4	16.6	40.0	87.0	148.9	184.5
2002	14.6	*	*	1.0	1.4	3.6	6.2	16.1	44.4	97.8	152.6	202.6
2001	15.1	*	*	1.3	2.0	3.9	7.3	16.0	46.0	99.2	159.5	206.5
2000	14.7	*	*	0.8	2.7	3.5	5.7	17.9	44.5	96.0	155.0	224.9
1999	15.1	*	*	1.1	2.6	3.8	6.8	17.3	47.4	95.4	161.1	220.3
Black, female												
2003	12.9	*	*	0.8	1.1	2.4	4.2	9.9	23.8	63.3	113.2	137.3
2002	12.8	*	*	0.6	1.5	2.2	3.5	11.4	29.1	60.5	105.1	133.1
2001	13.1	*	*	0.6	1.6	2.9	4.0	10.8	29.2	63.0	108.6	136.8
2000	13.1	*	*	*	1.8	2.4	5.0	12.0	28.8	60.2	101.8	157.1
1999	13.2	*	*	0.7	2.0	2.1	4.4	10.8	32.0	62.6	107.6	140.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
All races 3/, both sexes												
2003	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.2	2.4
2002	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4
All races, male												
2003	0.5	*	*	*	0.1	0.4	0.4	0.5	0.9	1.5	2.6	2.6
2002	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8
All races, female												
2003	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.9	2.3
2002	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
White, both sexes												
2003	0.5	*	*	*	0.2	0.4	0.4	0.4	0.7	1.2	2.2	2.5
2002	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.1	2.9
2001	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.3	2.4
2000	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.2	2.6
1999	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.6	2.3	2.4
White, male												
2003	0.5	*	*	*	0.1	0.5	0.5	0.5	0.9	1.5	2.7	2.7
2002	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.9	2.6	3.9
2001	0.6	*	*	*	0.2	0.4	0.5	0.6	0.7	1.7	2.7	2.7
2000	0.5	*	*	*	0.2	0.4	0.4	0.6	0.7	1.7	2.7	3.9
1999	0.6	*	*	*	0.3	0.4	0.5	0.5	1.1	2.0	2.7	3.8
White, female												
2003	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	2.0	2.4
2002	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.7	2.5
2001	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	2.0	2.2
2000	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.9	2.1
1999	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.3	2.1	1.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hodgkin's disease (C81)												
Black, both sexes												
2003	0.3	*	*	*	*	0.6	*	*	0.8	*	*	*
2002	0.3	*	*	*	*	0.4	0.4	0.4	0.8	*	*	*
2001	0.4	*	*	*	*	0.7	0.4	*	*	*	*	*
2000	0.3	*	*	*	*	*	0.4	*	*	*	*	*
1999	0.4	*	*	*	*	0.5	*	*	*	*	*	*
Black, male												
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.5	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	0.8	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
All races 3/, both sexes												
2003	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
2002	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7
All races, male												
2003	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
2002	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4
All races, female												
2003	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
2002	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
White, both sexes												
2003	8.3	*	*	0.1	0.3	0.6	1.4	3.8	11.2	28.2	59.1	83.3
2002	8.5	*	*	0.1	0.3	0.6	1.5	3.9	11.9	30.2	60.3	82.1
2001	8.8	*	*	0.1	0.3	0.7	1.6	4.5	12.6	30.7	61.6	83.4
2000	9.1	*	*	0.1	0.4	0.7	1.7	4.7	13.7	32.1	64.3	84.3
1999	9.2	*	*	0.1	0.4	0.7	1.7	4.9	13.8	33.5	64.9	81.2
White, male												
2003	8.9	*	*	*	0.4	0.7	1.8	4.8	14.4	35.9	73.0	109.7
2002	9.1	*	*	0.2	0.5	0.8	1.9	5.0	14.8	37.7	74.4	105.6
2001	9.3	*	*	0.1	0.4	0.9	2.1	5.8	15.3	38.0	74.7	111.5
2000	9.6	*	*	0.1	0.6	0.8	2.2	6.0	16.5	39.2	78.4	111.8
1999	9.6	*	*	0.1	0.5	0.8	2.2	6.3	16.6	40.2	78.0	108.0
White, female												
2003	7.7	*	*	*	0.3	0.4	1.0	2.8	8.3	21.7	49.7	71.6
2002	8.0	*	*	*	0.2	0.5	1.1	2.9	9.1	23.9	50.8	71.8
2001	8.2	*	*	*	0.3	0.5	1.1	3.1	10.1	24.5	52.9	71.6
2000	8.6	*	*	*	0.2	0.6	1.1	3.4	11.1	26.1	55.0	73.1
1999	8.8	*	*	*	0.3	0.5	1.2	3.5	11.1	27.9	56.3	70.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)												
Black, both sexes												
2003	3.6	*	*	*	0.4	1.1	1.9	4.8	8.9	18.2	28.3	31.2
2002	3.9	*	*	*	0.4	1.2	2.0	4.7	10.8	21.2	30.2	30.6
2001	4.0	*	*	*	0.5	1.2	2.2	5.0	10.9	21.0	30.5	39.3
2000	3.7	*	*	*	0.6	1.0	2.1	5.0	10.5	19.3	26.8	36.3
1999	4.0	*	*	*	0.5	1.1	2.4	4.7	11.1	20.1	31.8	40.5
Black, male												
2003	3.8	*	*	*	*	1.3	2.3	6.5	11.2	20.4	31.0	34.4
2002	4.3	*	*	*	*	1.5	2.9	5.5	13.7	27.8	35.3	42.9
2001	4.5	*	*	*	0.7	1.3	3.2	6.7	13.6	27.7	36.8	54.5
2000	4.2	*	*	*	*	1.2	2.6	6.9	14.5	24.9	31.2	43.8
1999	4.6	*	*	*	*	1.5	3.2	6.6	14.6	25.6	40.6	59.8
Black, female												
2003	3.4	*	*	*	*	0.8	1.5	3.3	7.1	16.6	26.8	30.0
2002	3.5	*	*	*	*	0.8	1.3	4.0	8.5	16.5	27.3	25.9
2001	3.6	*	*	*	*	1.1	1.3	3.6	8.7	16.3	26.9	33.7
2000	3.3	*	*	*	*	0.9	1.6	3.3	7.3	15.4	24.4	33.5
1999	3.5	*	*	*	*	*	1.7	3.2	8.2	16.3	26.9	33.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
All races 3/, both sexes												
2003	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
2002	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4
All races, male												
2003	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
2002	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2
All races, female												
2003	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
2002	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
White, both sexes												
2003	8.2	0.9	0.9	0.9	1.0	1.1	1.8	3.4	9.6	26.6	56.0	83.4
2002	8.2	*	0.7	0.8	1.2	1.2	1.8	3.6	9.9	27.5	54.9	83.0
2001	8.3	0.6	1.1	0.8	1.2	1.3	1.7	3.7	10.2	27.5	55.1	84.7
2000	8.3	1.2	1.0	0.8	1.2	1.1	1.8	3.7	10.0	28.3	56.6	86.0
1999	8.3	0.8	0.9	0.7	1.2	1.2	1.8	3.7	10.8	28.0	54.5	85.4
White, male												
2003	9.4	*	1.0	1.0	1.2	1.3	1.9	4.1	12.2	36.4	78.7	116.9
2002	9.3	*	0.8	0.8	1.5	1.4	2.0	4.1	12.3	37.5	77.4	119.1
2001	9.3	*	1.1	0.8	1.5	1.5	1.9	4.3	13.0	36.7	76.8	123.1
2000	9.4	*	0.9	0.9	1.4	1.3	1.9	4.2	12.6	38.5	78.5	123.6
1999	9.2	*	0.9	0.8	1.5	1.3	1.9	4.2	13.3	37.9	75.5	124.1
White, female												
2003	7.0	1.4	0.8	0.7	0.7	0.9	1.6	2.7	7.1	18.2	40.6	68.6
2002	7.1	*	0.6	0.8	1.0	0.9	1.6	3.1	7.6	18.9	39.8	67.3
2001	7.3	*	1.1	0.7	1.0	1.1	1.6	3.2	7.6	19.7	40.8	68.6
2000	7.4	*	1.0	0.7	0.9	0.9	1.6	3.2	7.7	19.8	42.1	70.8
1999	7.3	*	0.8	0.7	1.0	1.2	1.6	3.2	8.4	19.8	40.6	69.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Leukemia (C91-C95)												
Black, both sexes												
2003	4.6	*	*	0.7	0.9	1.2	2.3	3.7	9.6	23.1	41.6	59.2
2002	4.7	*	*	0.6	0.9	1.2	1.6	4.3	11.2	22.3	42.1	64.7
2001	4.7	*	*	0.9	1.1	1.4	2.1	4.2	9.8	24.9	41.6	57.1
2000	4.8	*	*	0.5	1.4	1.6	2.0	4.8	10.7	22.7	38.2	66.0
1999	4.7	*	*	0.8	1.4	1.2	1.9	4.3	10.6	24.8	41.2	55.8
Black, male												
2003	5.0	*	*	0.7	1.1	1.6	2.8	4.5	12.3	31.1	53.3	77.2
2002	5.1	*	*	0.7	1.0	1.4	1.8	5.2	13.7	31.5	56.8	86.8
2001	5.2	*	*	1.1	1.2	1.7	2.5	4.5	13.0	34.0	55.0	77.2
2000	5.3	*	*	0.6	1.9	1.9	2.0	5.5	13.8	31.6	54.8	98.0
1999	5.2	*	*	1.0	1.6	1.5	2.0	5.5	13.4	32.9	54.7	77.3
Black, female												
2003	4.2	*	*	0.7	0.6	0.8	1.9	3.0	7.4	17.4	34.9	52.3
2002	4.3	*	*	*	0.7	1.1	1.4	3.6	9.2	15.7	33.8	56.3
2001	4.3	*	*	0.6	1.0	1.1	1.7	3.9	7.3	18.4	34.0	49.7
2000	4.2	*	*	*	0.9	1.3	2.1	4.3	8.1	16.4	28.8	54.1
1999	4.3	*	*	0.6	1.3	1.0	1.7	3.3	8.3	19.0	33.6	47.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
All races 3/, both sexes												
2003	3.9	*	*	*	*	*	0.3	1.7	5.9	16.5	31.1	36.3
2002	4.0	*	*	*	*	*	0.3	1.8	6.3	17.4	31.2	36.7
2001	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6
All races, male												
2003	4.1	*	*	*	*	*	0.4	2.1	7.4	19.8	37.7	50.4
2002	4.2	*	*	*	*	*	0.3	2.2	7.6	21.0	39.7	50.8
2001	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3
All races, female												
2003	3.7	*	*	*	*	*	0.2	1.4	4.6	13.8	26.7	30.1
2002	3.7	*	*	*	*	*	0.3	1.5	5.1	14.3	25.5	30.6
2001	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
White, both sexes												
2003	3.9	*	*	*	*	*	0.3	1.4	5.4	15.4	29.5	35.0
2002	4.0	*	*	*	*	*	0.2	1.6	5.6	16.3	30.2	35.7
2001	3.9	*	*	*	*	*	0.3	1.5	5.8	16.3	29.5	35.0
2000	3.9	*	*	*	*	*	0.3	1.6	5.6	16.2	30.0	35.2
1999	3.9	*	*	*	*	*	0.3	1.6	6.1	16.1	29.4	34.8
White, male												
2003	4.2	*	*	*	*	*	0.4	1.8	6.8	18.9	36.5	49.7
2002	4.3	*	*	*	*	*	0.3	2.0	6.9	20.1	39.0	50.3
2001	4.2	*	*	*	*	*	0.4	1.9	6.8	19.7	37.4	47.1
2000	4.1	*	*	*	*	*	0.3	1.8	6.3	19.3	38.2	50.2
1999	4.1	*	*	*	*	*	0.3	1.9	7.6	19.0	36.9	49.7
White, female												
2003	3.6	*	*	*	*	*	0.2	1.1	4.1	12.4	24.8	28.5
2002	3.7	*	*	*	*	*	0.2	1.2	4.4	13.1	24.3	29.4
2001	3.7	*	*	*	*	*	0.2	1.1	4.8	13.4	24.3	29.9
2000	3.8	*	*	*	*	*	0.2	1.4	5.0	13.6	24.6	29.1
1999	3.8	*	*	*	*	*	0.2	1.2	4.8	13.7	24.5	28.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Multiple myeloma and immunoproliferative neoplasms (C88,C90)												
Black, both sexes												
2003	4.8	*	*	*	*	*	0.7	4.2	11.8	30.8	55.3	59.8
2002	4.7	*	*	*	*	*	0.7	4.1	13.1	31.5	48.7	55.5
2001	4.9	*	*	*	*	*	0.9	3.6	15.1	31.3	53.1	58.3
2000	5.0	*	*	*	*	*	0.8	4.5	13.9	32.1	55.1	71.6
1999	5.0	*	*	*	*	*	0.9	4.3	16.6	30.2	53.1	63.1
Black, male												
2003	4.7	*	*	*	*	*	0.9	5.2	15.3	33.6	63.5	73.0
2002	4.7	*	*	*	*	*	0.8	4.6	15.7	37.1	59.1	71.8
2001	4.9	*	*	*	*	*	1.0	4.3	18.4	36.6	65.6	71.5
2000	4.8	*	*	*	*	*	*	5.0	15.0	38.5	67.8	79.6
1999	4.9	*	*	*	*	*	1.1	4.4	18.6	35.1	64.6	80.9
Black, female												
2003	4.9	*	*	*	*	*	*	3.3	9.0	28.7	50.7	54.7
2002	4.7	*	*	*	*	*	*	3.7	11.1	27.6	42.9	49.3
2001	4.9	*	*	*	*	*	0.8	3.0	12.5	27.6	46.1	53.5
2000	5.2	*	*	*	*	*	0.9	4.0	12.9	27.6	47.9	68.7
1999	5.1	*	*	*	*	*	0.7	4.2	15.0	26.8	46.6	56.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
All races 3/, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.5
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
All races, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
White, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	*
2001	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	*
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
White, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
All races 3/, both sexes												
2003	21.5	*	0.7	0.7	1.3	1.7	4.2	13.8	35.2	77.9	146.7	224.0
2002	21.8	0.6	0.8	0.7	1.3	1.8	4.5	13.7	36.6	80.8	146.7	230.1
2001	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4
All races, male												
2003	22.3	*	0.8	0.7	1.6	2.1	4.7	15.7	41.5	93.4	180.8	275.0
2002	22.7	*	0.8	0.7	1.6	2.3	5.0	15.9	43.2	98.1	180.0	290.8
2001	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9
All races, female												
2003	20.8	*	0.5	0.6	0.9	1.3	3.8	11.9	29.3	64.8	123.9	201.4
2002	21.0	*	0.8	0.7	0.9	1.4	4.1	11.6	30.4	66.3	124.6	203.8
2001	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
White, both sexes												
2003	23.1	*	0.6	0.7	1.3	1.7	4.1	13.1	34.6	78.3	148.6	227.9
2002	23.3	0.7	0.7	0.7	1.3	1.8	4.4	13.1	35.8	80.7	147.8	233.0
2001	23.5	*	0.8	0.7	1.3	1.8	4.4	13.3	36.2	81.4	149.5	238.1
2000	23.9	*	0.8	0.7	1.3	1.8	4.3	13.3	37.5	85.3	153.3	243.7
1999	25.3	*	0.8	0.7	1.3	1.9	4.9	14.3	40.2	91.3	161.2	252.4
White, male												
2003	23.8	*	0.7	0.7	1.6	2.1	4.6	14.8	40.6	93.5	183.4	280.9
2002	24.2	*	0.8	0.7	1.6	2.3	4.7	15.3	42.0	98.4	181.6	296.5
2001	24.0	*	1.0	0.7	1.7	2.2	4.8	15.5	42.0	97.7	182.1	300.2
2000	24.4	*	1.0	0.7	1.6	2.3	4.7	15.1	43.2	103.1	188.2	307.0
1999	25.9	*	0.9	0.8	1.6	2.4	5.6	16.2	47.4	109.9	197.5	323.6
White, female												
2003	22.5	*	0.5	0.6	1.0	1.2	3.6	11.4	28.9	65.4	125.0	204.4
2002	22.5	*	0.7	0.7	1.0	1.4	4.1	11.1	29.9	65.6	125.1	205.5
2001	22.9	*	0.7	0.6	0.9	1.4	4.0	11.2	30.7	67.7	127.8	212.1
2000	23.5	*	0.6	0.6	1.0	1.3	3.9	11.6	32.1	70.3	130.3	218.0
1999	24.6	*	0.7	0.6	1.1	1.4	4.2	12.4	33.4	75.8	137.4	223.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)												
Black, both sexes												
2003	17.3	*	0.9	0.8	1.1	2.3	5.8	20.7	48.2	88.2	150.1	206.3
2002	18.1	*	1.1	0.8	1.3	2.3	6.3	20.1	51.0	96.2	157.6	222.1
2001	18.3	*	*	0.7	1.3	2.3	6.0	21.7	52.9	100.8	156.9	217.1
2000	18.3	*	1.0	0.6	1.4	1.6	7.5	22.1	51.9	100.1	162.0	210.8
1999	19.5	*	*	0.7	1.4	2.0	7.5	22.9	59.6	104.7	170.9	242.6
Black, male												
2003	18.3	*	*	0.8	1.4	2.4	6.5	25.0	61.1	111.3	187.6	241.9
2002	18.8	*	*	0.8	1.8	2.6	7.4	24.0	63.8	116.7	195.6	267.0
2001	19.4	*	*	0.8	1.7	2.7	6.3	26.6	67.5	127.9	193.6	283.7
2000	19.4	*	*	0.7	1.9	1.8	8.5	26.2	68.1	122.6	210.4	253.7
1999	20.6	*	*	0.6	2.0	2.3	8.3	28.9	76.1	133.2	206.3	292.9
Black, female												
2003	16.5	*	*	0.7	0.8	2.2	5.1	17.0	37.8	71.8	128.8	192.4
2002	17.4	*	*	0.7	0.8	2.0	5.3	16.7	40.7	81.7	136.1	205.0
2001	17.3	*	*	0.6	0.8	1.9	5.7	17.6	41.2	81.7	136.3	192.4
2000	17.3	*	*	*	1.0	1.5	6.6	18.6	38.8	84.3	134.7	194.9
1999	18.5	*	*	0.8	0.9	1.6	6.7	17.8	46.4	84.5	151.1	223.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
All races 3/, both sexes												
2003	4.7	1.5	0.3	0.2	0.2	0.4	0.8	1.7	4.4	12.8	36.4	80.5
2002	4.6	1.7	0.4	0.2	0.2	0.4	0.8	1.8	4.5	12.8	36.1	78.7
2001	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9
All races, male												
2003	4.7	1.6	0.4	0.2	0.3	0.5	0.9	1.8	5.1	16.0	46.2	101.7
2002	4.6	1.5	0.4	0.2	0.2	0.4	0.9	1.9	5.3	15.8	45.9	101.1
2001	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2
All races, female												
2003	4.7	1.5	0.3	0.2	0.2	0.4	0.6	1.5	3.8	10.2	29.9	71.1
2002	4.6	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.7	10.4	29.6	68.9
2001	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
White, both sexes												
2003	5.2	1.5	0.3	0.2	0.2	0.5	0.7	1.6	4.4	13.1	37.7	83.0
2002	5.1	1.9	0.4	0.2	0.2	0.4	0.8	1.7	4.5	13.2	37.0	80.7
2001	5.3	1.2	0.4	0.2	0.2	0.4	0.8	1.9	5.0	14.3	38.0	83.9
2000	5.3	1.6	0.4	0.2	0.2	0.4	0.8	1.8	5.1	13.9	38.6	87.3
1999	5.2	1.4	0.4	0.3	0.2	0.4	0.7	1.7	4.7	14.8	37.4	85.3
White, male												
2003	5.2	1.6	0.3	0.2	0.3	0.5	0.9	1.8	5.0	16.4	47.8	105.5
2002	5.1	1.7	0.3	0.2	0.2	0.4	0.9	1.9	5.3	16.3	47.0	104.2
2001	5.3	1.2	0.4	0.2	0.2	0.4	0.9	2.1	5.6	17.8	49.4	109.9
2000	5.2	1.6	0.4	0.2	0.3	0.4	0.8	2.0	5.6	17.0	49.1	112.6
1999	5.2	1.4	0.4	0.2	0.2	0.4	0.9	2.0	5.8	18.7	47.2	114.3
White, female												
2003	5.1	1.5	*	0.2	0.2	0.4	0.6	1.5	3.9	10.3	30.9	73.0
2002	5.1	2.1	0.4	0.2	0.2	0.3	0.7	1.6	3.8	10.5	30.3	70.5
2001	5.2	*	0.4	0.3	0.2	0.4	0.7	1.6	4.4	11.4	30.4	73.0
2000	5.4	1.6	*	0.2	0.2	0.4	0.8	1.5	4.6	11.3	31.8	77.1
1999	5.1	1.4	0.4	0.3	0.2	0.4	0.6	1.5	3.7	11.5	30.9	73.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)												
Black, both sexes												
2003	2.9	*	*	*	*	0.5	1.2	2.5	5.2	13.0	29.5	57.8
2002	3.0	*	*	*	*	0.6	1.2	2.5	5.5	12.6	31.3	63.6
2001	3.1	*	*	0.4	0.3	*	1.1	2.0	5.9	14.3	31.5	73.4
2000	3.1	*	*	*	*	0.6	1.1	2.7	5.5	15.0	31.0	64.1
1999	3.2	*	*	*	0.4	0.5	1.2	2.6	6.2	15.7	33.0	61.9
Black, male												
2003	2.9	*	*	*	*	*	1.1	2.5	6.7	16.2	36.7	66.7
2002	2.8	*	*	*	*	*	1.0	2.2	7.1	14.1	38.7	79.3
2001	3.0	*	*	*	*	*	1.1	1.8	6.5	17.2	41.5	88.5
2000	3.0	*	*	*	*	*	1.2	2.8	7.0	19.2	36.0	76.1
1999	3.0	*	*	*	*	0.8	1.1	2.5	7.1	19.0	38.4	63.3
Black, female												
2003	3.0	*	*	*	*	*	1.2	2.4	4.0	10.8	25.5	54.3
2002	3.3	*	*	*	*	0.8	1.4	2.7	4.2	11.5	27.1	57.5
2001	3.2	*	*	*	*	*	1.0	2.2	5.3	12.3	25.9	67.8
2000	3.2	*	*	*	*	0.8	1.0	2.6	4.4	12.0	28.1	59.7
1999	3.4	*	*	*	*	*	1.2	2.7	5.6	13.4	29.9	61.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
All races 3/, both sexes												
2003	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
2002	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8
All races, male												
2003	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
2002	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4
All races, female												
2003	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
2002	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
White, both sexes												
2003	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.6	2.4	8.6	40.6
2002	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.7	2.5	8.7	39.7
2001	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	41.7
2000	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.3	42.0
1999	1.5	*	*	0.1	0.1	*	0.1	0.3	0.6	2.4	9.4	43.0
White, male												
2003	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.6	9.1	37.9
2002	1.1	*	*	*	*	0.1	0.1	0.3	0.8	2.6	8.5	37.2
2001	1.1	*	*	*	*	0.1	0.1	0.2	0.7	2.6	9.8	40.4
2000	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	39.1
1999	1.1	*	*	*	*	*	0.1	0.3	0.5	2.5	9.9	43.4
White, female												
2003	1.8	*	*	*	*	*	0.1	0.2	0.6	2.2	8.2	41.8
2002	1.8	*	*	*	*	*	*	0.3	0.7	2.4	8.8	40.8
2001	1.9	*	*	*	*	*	0.2	0.3	0.8	2.4	9.1	42.3
2000	1.9	*	*	*	*	*	0.1	0.3	0.8	2.4	9.3	43.2
1999	1.9	*	*	*	0.1	*	0.1	0.3	0.7	2.3	9.0	42.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Anemias (D50-D64)												
Black, both sexes												
2003	2.7	*	*	0.4	1.3	2.1	2.3	3.0	4.2	5.6	14.5	49.3
2002	2.8	*	*	0.4	1.3	2.4	2.5	3.1	3.6	5.9	16.0	52.3
2001	2.7	*	*	0.4	1.2	2.2	2.7	3.1	3.0	5.5	16.0	48.5
2000	2.6	*	*	0.5	1.1	2.3	2.4	3.0	3.4	5.4	13.3	51.0
1999	2.8	*	0.8	0.5	1.4	2.6	2.5	3.1	4.2	5.0	13.4	52.9
Black, male												
2003	2.7	*	*	*	1.3	2.2	2.6	3.3	5.4	7.9	14.9	49.0
2002	2.7	*	*	*	1.6	2.8	2.5	2.6	3.9	7.3	16.9	56.8
2001	2.5	*	*	*	1.4	2.9	2.4	3.6	3.1	6.7	15.0	42.0
2000	2.3	*	*	*	1.0	2.1	2.0	3.3	3.4	4.5	13.9	55.4
1999	2.7	*	*	*	1.7	2.8	2.7	3.3	5.1	5.4	12.9	44.5
Black, female												
2003	2.8	*	*	*	1.2	2.0	2.1	2.8	3.2	3.9	14.2	49.4
2002	3.0	*	*	*	0.9	2.0	2.5	3.6	3.3	5.0	15.4	50.5
2001	2.8	*	*	*	1.1	1.6	2.9	2.6	2.8	4.6	16.5	50.9
2000	3.0	*	*	*	1.1	2.4	2.8	2.8	3.4	6.0	13.0	49.4
1999	3.0	*	*	*	1.2	2.3	2.3	3.0	3.6	4.7	13.6	56.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
All races 3/, both sexes												
2003	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
2002	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2
All races, male												
2003	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
2002	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7
All races, female												
2003	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	307.2
2002	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
White, both sexes												
2003	25.0	*	*	0.1	0.3	1.4	4.0	11.8	33.5	81.4	168.4	300.8
2002	24.9	*	*	0.1	0.3	1.4	4.3	11.7	32.7	81.5	170.4	303.7
2001	24.6	*	*	0.1	0.3	1.2	3.8	11.7	32.7	81.9	168.9	306.0
2000	24.1	*	*	*	0.3	1.3	3.8	11.1	33.1	81.2	166.0	303.5
1999	23.9	*	*	0.1	0.4	1.3	3.7	10.9	32.4	81.8	165.1	298.9
White, male												
2003	24.8	*	*	*	0.4	1.5	5.1	14.7	40.2	95.8	198.6	333.4
2002	24.2	*	*	*	0.4	1.6	5.3	14.3	39.6	93.3	194.7	344.3
2001	23.5	*	*	*	0.3	1.3	4.7	14.3	38.4	94.6	189.2	337.3
2000	22.9	*	*	*	0.4	1.6	4.7	13.3	38.0	91.8	190.5	337.4
1999	22.7	*	*	*	0.4	1.4	4.5	13.5	37.3	92.3	189.9	333.4
White, female												
2003	25.2	*	*	*	0.3	1.2	2.9	9.0	27.2	69.2	148.0	286.3
2002	25.6	*	*	*	0.3	1.1	3.2	9.2	26.2	71.5	154.0	286.1
2001	25.7	*	*	*	0.2	1.1	2.9	9.2	27.4	71.2	155.5	293.0
2000	25.3	*	*	*	0.3	1.0	3.0	9.0	28.5	72.4	149.8	289.7
1999	25.0	*	*	*	0.3	1.2	3.0	8.4	27.8	73.1	148.9	285.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diabetes mellitus (E10-E14)												
Black, both sexes												
2003	33.8	*	*	*	0.7	3.8	9.5	30.1	85.7	184.9	336.1	527.5
2002	33.6	*	*	*	1.0	3.6	9.6	30.1	84.2	188.1	336.9	537.8
2001	33.1	*	*	*	1.0	3.7	8.8	29.6	85.0	188.4	335.2	529.7
2000	32.8	*	*	*	0.9	3.5	8.0	30.2	82.5	188.4	350.6	524.9
1999	33.0	*	*	*	0.8	3.0	8.9	30.3	90.2	190.8	333.5	529.0
Black, male												
2003	29.8	*	*	*	0.8	4.7	11.5	33.7	96.0	197.2	337.3	453.5
2002	29.0	*	*	*	1.2	4.4	11.7	34.8	94.7	190.1	329.9	417.0
2001	28.5	*	*	*	1.0	4.8	10.2	34.5	92.2	187.7	341.9	450.5
2000	27.4	*	*	*	1.0	3.8	9.4	33.4	88.2	184.9	337.0	452.1
1999	27.7	*	*	*	*	3.3	10.3	34.7	93.8	186.5	314.8	506.2
Black, female												
2003	37.4	*	*	*	*	3.0	7.8	27.0	77.4	176.1	335.4	556.2
2002	37.8	*	*	*	0.7	2.9	7.8	26.0	75.8	186.7	340.9	584.1
2001	37.2	*	*	*	1.0	2.8	7.6	25.4	79.1	188.9	331.5	559.1
2000	37.8	*	*	*	0.9	3.1	6.8	27.5	77.8	190.9	358.3	552.0
1999	37.8	*	*	*	0.9	2.6	7.6	26.6	87.4	193.9	343.9	537.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
All races 3/, both sexes												
2003	1.1	*	*	*	*	*	0.1	0.2	0.6	1.9	7.7	35.5
2002	1.3	*	*	*	*	*	0.1	0.3	0.6	2.1	8.9	41.5
2001	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9
All races, male												
2003	0.8	*	*	*	*	*	*	0.2	0.6	2.0	7.8	30.6
2002	0.9	*	*	*	*	*	0.1	0.2	0.7	2.2	8.5	37.0
2001	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4
All races, female												
2003	1.5	*	*	*	*	*	*	0.3	0.6	1.7	7.7	37.6
2002	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	9.1	43.5
2001	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
White, both sexes												
2003	1.2	*	*	*	*	*	0.1	0.2	0.5	1.7	7.4	34.2
2002	1.4	*	*	*	*	*	0.1	0.2	0.5	1.9	8.6	40.2
2001	1.4	*	*	*	*	*	0.1	0.2	0.5	1.9	8.2	42.2
2000	1.6	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	48.8
1999	1.6	*	*	*	*	*	0.1	0.2	0.6	2.0	9.5	52.8
White, male												
2003	0.8	*	*	*	*	*	*	0.2	0.5	1.8	7.1	29.6
2002	1.0	*	*	*	*	*	0.1	0.2	0.5	2.1	8.1	36.3
2001	0.9	*	*	*	*	*	*	0.2	0.5	1.8	8.0	33.8
2000	1.0	*	*	*	*	*	*	0.2	0.6	2.3	9.1	42.8
1999	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	9.3	46.5
White, female												
2003	1.6	*	*	*	*	*	*	0.2	0.6	1.7	7.5	36.3
2002	1.8	*	*	*	*	*	*	0.3	0.5	1.8	8.9	41.9
2001	1.8	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.7
2000	2.1	*	*	*	*	*	0.2	0.3	0.6	2.3	10.5	51.3
1999	2.1	*	*	*	*	*	0.2	0.2	0.5	1.9	9.6	55.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nutritional deficiencies (E40-E64)												
Black, both sexes												
2003	1.2	*	*	*	*	*	*	0.7	1.4	3.2	13.2	55.1
2002	1.3	*	*	*	*	*	*	0.4	1.7	4.2	14.5	60.6
2001	1.3	*	*	*	*	*	*	0.7	1.2	3.7	15.9	54.7
2000	1.5	*	*	*	*	*	*	0.6	1.7	4.7	16.1	66.9
1999	1.6	*	*	*	*	*	*	0.7	2.0	5.5	19.4	70.8
Black, male												
2003	1.0	*	*	*	*	*	*	*	1.9	4.4	17.8	48.0
2002	1.0	*	*	*	*	*	*	*	2.8	4.4	16.4	47.2
2001	1.2	*	*	*	*	*	*	1.2	1.9	5.5	16.5	57.9
2000	1.2	*	*	*	*	*	*	*	2.8	6.4	17.9	63.4
1999	1.5	*	*	*	*	*	*	*	2.8	8.1	22.8	75.0
Black, female												
2003	1.4	*	*	*	*	*	*	*	*	2.3	10.7	57.9
2002	1.6	*	*	*	*	*	*	*	*	4.1	13.5	65.7
2001	1.4	*	*	*	*	*	*	*	*	2.4	15.5	53.5
2000	1.7	*	*	*	*	*	*	*	*	3.5	15.2	68.3
1999	1.8	*	*	*	*	*	*	*	*	3.6	17.6	69.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
All races 3/, both sexes												
2003	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	33.4
2002	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0
All races, male												
2003	0.8	*	*	*	*	*	*	0.2	0.6	1.9	7.4	28.2
2002	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4
All races, female												
2003	1.4	*	*	*	*	*	*	0.2	0.6	1.7	7.4	35.6
2002	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
White, both sexes												
2003	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	7.0	32.1
2002	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.0
2001	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	7.6	39.5
2000	1.5	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.2
1999	1.5	*	*	*	*	*	0.1	0.2	0.5	1.9	8.8	48.9
White, male												
2003	0.8	*	*	*	*	*	*	0.2	0.4	1.7	6.7	27.2
2002	0.9	*	*	*	*	*	0.1	0.2	0.5	1.9	7.5	34.3
2001	0.8	*	*	*	*	*	*	0.1	0.5	1.7	7.4	32.2
2000	1.0	*	*	*	*	*	*	0.2	0.6	2.2	8.4	39.7
1999	1.0	*	*	*	*	*	*	0.2	0.6	2.0	8.5	42.2
White, female												
2003	1.5	*	*	*	*	*	*	0.2	0.6	1.6	7.3	34.2
2002	1.6	*	*	*	*	*	*	0.3	0.5	1.7	8.4	38.2
2001	1.7	*	*	*	*	*	*	0.3	0.5	1.9	7.7	42.6
2000	1.9	*	*	*	*	*	0.1	0.2	0.6	2.0	9.9	47.4
1999	2.0	*	*	*	*	*	0.2	0.2	0.4	1.8	9.0	51.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malnutrition (E40-E46)												
Black, both sexes												
2003	1.2	*	*	*	*	*	*	0.7	1.4	3.0	13.0	53.4
2002	1.3	*	*	*	*	*	*	*	1.7	3.6	14.0	57.6
2001	1.2	*	*	*	*	*	*	0.6	1.1	3.4	14.8	51.6
2000	1.4	*	*	*	*	*	*	0.5	1.5	4.7	14.9	63.8
1999	1.5	*	*	*	*	*	*	0.6	1.8	5.4	17.9	66.0
Black, male												
2003	1.0	*	*	*	*	*	*	*	1.9	4.2	17.2	45.9
2002	1.0	*	*	*	*	*	*	*	2.6	3.9	15.2	46.1
2001	1.1	*	*	*	*	*	*	1.0	*	4.9	15.9	55.6
2000	1.2	*	*	*	*	*	*	*	2.4	6.3	17.3	61.1
1999	1.4	*	*	*	*	*	*	*	2.5	7.8	20.3	73.8
Black, female												
2003	1.3	*	*	*	*	*	*	*	*	2.2	10.7	56.3
2002	1.5	*	*	*	*	*	*	*	*	3.4	13.3	62.0
2001	1.3	*	*	*	*	*	*	*	*	2.2	14.2	50.1
2000	1.6	*	*	*	*	*	*	*	*	3.5	13.6	64.8
1999	1.6	*	*	*	*	*	*	*	*	3.6	16.5	63.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
All races 3/, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	0.1	0.3	2.1
2002	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.1
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
All races, male												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2002	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9
All races, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	2.0
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
White, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.1
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9
White, male												
2003	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.8	4.3
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2002	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000	0.2	*	*	*	*	*	*	*	*	0.3	0.6	3.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	3.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other nutritional deficiencies (E50-E64)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
All races 3/, both sexes												
2003	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5
All races, male												
2003	0.3	1.9	*	*	0.1	0.1	0.2	0.4	0.5	0.5	0.8	*
2002	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7
All races, female												
2003	0.2	2.0	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0
2002	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
White, both sexes												
2003	0.2	1.7	*	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.0
2002	0.2	1.7	*	0.1	*	0.1	0.1	0.2	0.3	0.6	0.6	1.2
2001	0.2	1.5	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.8
2000	0.2	1.8	0.2	0.1	*	0.1	0.1	0.2	0.3	0.5	0.9	1.0
1999	0.3	2.3	0.3	*	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.3
White, male												
2003	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.5	0.8	*
2002	0.2	1.8	*	*	*	*	0.1	0.3	0.3	0.7	0.8	*
2001	0.2	1.4	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
2000	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.7	1.2	*
1999	0.3	3.0	0.3	*	*	*	0.1	0.3	0.4	0.6	0.9	*
White, female												
2003	0.2	1.3	*	*	*	*	*	0.2	0.2	0.5	0.6	0.9
2002	0.2	1.5	*	*	*	*	*	0.2	0.2	0.4	0.5	1.2
2001	0.2	1.6	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8
2000	0.2	1.6	*	*	*	*	*	0.2	0.2	0.4	0.8	1.0
1999	0.3	1.5	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Meningitis (G00,G03)												
Black, both sexes												
2003	0.5	3.5	*	*	*	*	0.4	0.7	1.3	*	*	*
2002	0.5	3.0	*	*	*	*	0.3	0.9	0.9	1.2	*	*
2001	0.5	4.5	*	*	*	*	0.4	0.7	1.0	*	*	*
2000	0.6	5.7	*	*	*	*	0.5	0.6	0.9	1.6	*	*
1999	0.6	6.9	*	*	*	*	0.4	0.9	*	1.3	2.8	*
Black, male												
2003	0.5	*	*	*	*	*	*	*	*	*	*	*
2002	0.6	*	*	*	*	*	*	1.2	*	*	*	*
2001	0.6	*	*	*	*	*	*	*	*	*	*	*
2000	0.6	*	*	*	*	*	*	*	*	*	*	*
1999	0.7	*	*	*	*	*	*	1.4	*	*	*	*
Black, female												
2003	0.5	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	9.7	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
All races 3/, both sexes												
2003	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
2002	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4
All races, male												
2003	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
2002	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6
All races, female												
2003	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
2002	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
White, both sexes												
2003	7.2	*	*	*	*	*	*	0.2	1.3	13.6	72.2	146.0
2002	6.9	*	*	*	*	*	*	0.2	1.3	13.0	68.2	143.3
2001	6.8	*	*	*	*	*	*	0.1	1.2	12.6	68.8	141.8
2000	6.5	*	*	*	*	*	*	0.1	1.2	12.3	65.6	139.5
1999	6.1	*	*	*	*	*	*	0.1	1.1	11.7	61.5	131.2
White, male												
2003	8.2	*	*	*	*	*	*	0.2	1.6	19.3	106.6	233.3
2002	7.9	*	*	*	*	*	*	0.2	1.6	19.0	100.5	232.7
2001	7.8	*	*	*	*	*	*	0.2	1.9	17.9	103.0	237.6
2000	7.4	*	*	*	*	*	*	0.2	1.6	17.2	97.5	235.3
1999	7.0	*	*	*	*	*	*	0.2	1.5	16.8	94.4	216.7
White, female												
2003	6.2	*	*	*	*	*	*	0.2	1.0	8.7	48.8	107.4
2002	5.9	*	*	*	*	*	*	*	0.9	7.9	46.5	104.4
2001	5.8	*	*	*	*	*	*	*	0.7	8.1	46.2	101.7
2000	5.6	*	*	*	*	*	*	*	0.9	8.2	44.7	100.6
1999	5.2	*	*	*	*	*	*	*	0.6	7.5	40.0	96.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Parkinson's disease (G20-G21)												
Black, both sexes												
2003	1.6	*	*	*	*	*	*	*	1.2	7.0	29.0	52.8
2002	1.5	*	*	*	*	*	*	*	*	6.8	25.4	51.7
2001	1.5	*	*	*	*	*	*	*	0.8	5.5	25.2	59.0
2000	1.5	*	*	*	*	*	*	*	*	5.7	27.4	51.6
1999	1.4	*	*	*	*	*	*	*	0.9	5.9	24.4	52.0
Black, male												
2003	1.9	*	*	*	*	*	*	*	1.7	10.3	45.4	97.0
2002	1.7	*	*	*	*	*	*	*	*	11.1	39.6	89.0
2001	1.7	*	*	*	*	*	*	*	*	8.5	37.4	99.8
2000	1.8	*	*	*	*	*	*	*	*	8.8	47.8	86.5
1999	1.6	*	*	*	*	*	*	*	*	8.7	37.8	86.7
Black, female												
2003	1.4	*	*	*	*	*	*	*	*	4.7	19.8	35.6
2002	1.2	*	*	*	*	*	*	*	*	3.6	17.4	37.4
2001	1.3	*	*	*	*	*	*	*	*	3.3	18.3	43.8
2000	1.2	*	*	*	*	*	*	*	*	3.5	15.9	38.6
1999	1.2	*	*	*	*	*	*	*	*	3.9	16.9	39.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
All races 3/, both sexes												
2003	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
2002	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3
All races, male												
2003	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
2002	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7
All races, female												
2003	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
2002	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
White, both sexes												
2003	25.0	*	*	*	*	*	*	0.2	2.2	21.7	170.7	834.0
2002	23.5	*	*	*	*	*	*	0.1	2.1	20.7	164.8	781.8
2001	21.7	*	*	*	*	*	*	0.2	2.3	19.6	152.6	738.1
2000	20.2	*	*	*	*	*	*	0.2	2.2	19.5	144.8	695.3
1999	18.3	*	*	*	*	*	*	0.2	2.0	18.1	134.5	627.9
White, male												
2003	14.6	*	*	*	*	*	*	0.2	1.9	20.8	152.4	630.3
2002	13.7	*	*	*	*	*	*	0.1	1.9	19.3	147.3	594.8
2001	12.8	*	*	*	*	*	*	0.2	2.0	18.5	140.5	577.2
2000	11.9	*	*	*	*	*	*	0.2	1.9	19.5	132.5	542.1
1999	11.2	*	*	*	*	*	*	0.2	2.1	17.9	127.4	506.9
White, female												
2003	35.2	*	*	*	*	*	*	0.2	2.4	22.5	183.1	924.3
2002	33.0	*	*	*	*	*	*	0.1	2.2	21.8	176.5	863.0
2001	30.3	*	*	*	*	*	*	0.2	2.6	20.6	160.6	805.4
2000	28.2	*	*	*	*	*	*	0.2	2.4	19.5	152.9	757.6
1999	25.3	*	*	*	*	*	*	0.2	2.0	18.3	139.1	676.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alzheimer's disease (G30)												
Black, both sexes												
2003	9.7	*	*	*	*	*	*	*	1.0	19.2	138.1	583.8
2002	8.9	*	*	*	*	*	*	*	1.2	16.8	123.1	543.7
2001	8.4	*	*	*	*	*	*	*	1.1	15.2	124.7	506.7
2000	7.5	*	*	*	*	*	*	*	1.2	15.8	109.9	445.1
1999	6.5	*	*	*	*	*	*	*	1.2	15.1	98.2	372.4
Black, male												
2003	6.0	*	*	*	*	*	*	*	*	20.3	136.2	459.8
2002	5.3	*	*	*	*	*	*	*	*	18.0	122.5	412.8
2001	5.1	*	*	*	*	*	*	*	*	16.1	121.5	414.1
2000	4.5	*	*	*	*	*	*	*	*	16.5	97.5	371.3
1999	4.0	*	*	*	*	*	*	*	*	16.5	89.8	317.5
Black, female												
2003	13.2	*	*	*	*	*	*	*	*	18.4	139.2	631.9
2002	12.0	*	*	*	*	*	*	*	*	15.9	123.4	593.9
2001	11.3	*	*	*	*	*	*	*	*	14.6	126.4	541.0
2000	10.2	*	*	*	*	*	*	*	*	15.2	116.9	472.6
1999	8.7	*	*	*	*	*	*	*	*	14.2	102.9	392.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
All races 3/, both sexes												
2003	310.3	13.9	1.5	0.8	3.5	10.3	38.2	113.4	284.9	748.3	2,170.8	7,138.0
2002	318.6	15.9	1.4	0.9	3.2	10.0	37.8	114.3	294.8	788.4	2,257.0	7,389.6
2001	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1
All races, male												
2003	296.3	15.2	1.4	1.0	4.2	12.9	51.1	160.9	393.3	976.3	2,636.1	7,329.3
2002	303.3	16.6	1.5	0.9	4.1	12.7	50.8	162.3	407.0	1,027.8	2,731.1	7,638.8
2001	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2
All races, female												
2003	323.9	12.5	1.5	0.7	2.8	7.5	25.3	67.5	184.4	557.8	1,860.0	7,053.5
2002	333.3	15.1	1.3	0.8	2.2	7.2	24.9	68.0	191.0	589.1	1,942.2	7,281.5
2001	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
White, both sexes												
2003	330.0	11.0	1.2	0.7	2.9	8.5	32.9	99.0	259.8	712.2	2,151.4	7,233.1
2002	339.4	13.4	1.3	0.8	2.7	8.1	33.0	100.6	269.9	755.1	2,234.8	7,482.7
2001	344.8	12.2	1.5	0.8	2.7	8.5	31.7	99.5	275.3	774.1	2,298.9	7,770.6
2000	354.7	14.6	1.3	0.8	2.8	7.9	31.6	102.0	292.7	814.2	2,370.8	8,133.9
1999	362.7	14.6	1.3	0.8	2.9	8.1	32.1	102.7	300.0	850.9	2,449.9	8,299.2
White, male												
2003	312.6	11.5	1.1	0.9	3.5	10.9	45.6	144.6	364.8	938.3	2,623.0	7,466.8
2002	320.9	14.3	1.5	0.8	3.5	10.5	45.7	147.1	378.3	994.8	2,721.2	7,766.6
2001	323.5	13.2	1.6	0.8	3.3	10.9	44.2	145.1	385.3	1,020.8	2,809.7	8,060.7
2000	334.2	15.3	1.6	0.9	3.3	10.1	44.1	150.2	411.8	1,080.7	2,912.4	8,585.0
1999	341.2	15.3	1.5	0.9	3.4	10.4	45.1	150.5	420.7	1,133.1	3,012.8	8,717.5
White, female												
2003	347.0	10.4	1.4	0.6	2.3	5.9	20.0	53.9	160.6	519.6	1,832.5	7,129.6
2002	357.5	12.5	1.1	0.7	1.8	5.6	20.0	54.7	167.7	551.7	1,908.4	7,359.3
2001	365.5	11.1	1.5	0.8	2.0	6.0	19.0	54.6	171.9	565.8	1,959.8	7,649.2
2000	374.8	13.9	1.1	0.6	2.2	5.5	19.0	54.6	180.9	590.8	2,014.4	7,950.6
1999	383.6	13.8	1.2	0.7	2.4	5.7	19.0	55.7	186.8	615.7	2,081.0	8,131.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Major cardiovascular diseases (I00-I78)												
Black, both sexes												
2003	273.1	26.2	2.8	1.4	6.9	22.9	78.8	238.8	556.2	1,207.1	2,713.5	6,862.0
2002	276.1	27.6	2.0	1.4	5.9	23.3	76.0	237.4	568.3	1,226.2	2,824.0	7,081.7
2001	280.4	28.6	3.5	1.3	6.6	22.6	78.3	238.2	578.0	1,276.2	2,891.8	7,316.7
2000	285.0	27.8	2.5	1.4	6.8	21.5	77.4	241.3	600.3	1,316.2	2,968.8	7,518.9
1999	289.4	29.2	2.4	1.4	6.8	20.7	81.6	241.9	614.6	1,368.1	3,024.7	7,582.5
Black, male												
2003	268.6	31.2	3.3	1.5	8.5	28.0	98.1	317.1	742.4	1,542.1	3,265.7	6,547.7
2002	268.5	28.2	1.9	1.5	7.2	28.9	94.7	313.1	762.5	1,556.1	3,287.6	6,929.0
2001	272.4	28.1	3.6	1.2	8.0	27.9	97.4	312.5	761.1	1,620.3	3,435.3	7,363.9
2000	275.6	29.1	2.1	1.4	7.5	25.4	95.8	317.5	796.8	1,639.9	3,501.8	7,687.4
1999	282.5	30.6	2.5	1.5	8.4	25.4	102.7	323.2	822.6	1,723.6	3,526.5	7,921.1
Black, female												
2003	277.2	21.0	2.4	1.3	5.3	18.1	61.7	171.5	406.5	968.7	2,401.2	6,984.1
2002	283.0	26.9	2.1	1.3	4.5	18.1	59.5	172.5	412.2	992.0	2,561.6	7,140.2
2001	287.8	29.0	3.4	1.4	5.1	17.7	61.4	174.3	430.5	1,033.4	2,586.6	7,299.2
2000	293.4	26.5	2.9	1.4	6.1	17.9	61.1	175.6	441.9	1,088.3	2,668.7	7,456.2
1999	295.7	27.9	2.4	1.2	5.1	16.4	63.0	172.3	447.7	1,117.3	2,743.8	7,456.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
All races 3/, both sexes												
2003	235.6	11.0	1.2	0.6	2.7	8.2	30.7	92.5	233.2	585.0	1,611.1	5,278.4
2002	241.7	12.4	1.1	0.6	2.5	7.9	30.5	93.7	241.5	615.9	1,677.2	5,446.8
2001	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0
All races, male												
2003	235.0	12.1	1.1	0.7	3.4	10.5	42.8	136.2	331.7	785.3	2,030.3	5,621.5
2002	240.7	12.9	1.1	0.7	3.3	10.5	43.1	138.4	343.4	827.1	2,110.1	5,823.5
2001	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6
All races, female												
2003	236.2	9.8	1.3	0.5	2.1	5.7	18.6	50.2	141.9	417.5	1,331.1	5,126.7
2002	242.7	11.8	1.0	0.6	1.7	5.2	18.0	50.6	147.2	440.1	1,389.7	5,283.3
2001	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8
2000	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
White, both sexes												
2003	251.7	8.8	1.0	0.5	2.3	6.7	27.2	82.9	216.5	561.8	1,603.6	5,363.5
2002	258.5	10.6	0.9	0.6	2.1	6.5	27.3	84.8	224.5	595.1	1,666.9	5,529.1
2001	262.8	9.9	1.2	0.6	2.0	6.6	25.9	83.9	229.6	612.2	1,714.3	5,752.1
2000	270.2	11.5	1.1	0.6	2.1	6.1	25.8	85.4	243.2	642.5	1,767.8	6,020.8
1999	277.7	11.8	1.0	0.6	2.3	6.4	26.3	86.8	251.1	677.7	1,837.5	6,152.7
White, male												
2003	249.5	9.4	0.8	0.6	2.8	9.0	39.1	125.3	313.2	761.1	2,030.1	5,747.2
2002	256.0	11.3	1.0	0.6	2.8	8.8	39.7	128.6	324.0	807.8	2,112.0	5,939.8
2001	257.8	10.6	1.2	0.6	2.6	8.7	37.8	127.0	330.8	829.1	2,175.8	6,157.2
2000	265.8	11.9	1.3	0.7	2.6	8.1	37.8	130.7	351.8	877.8	2,247.0	6,560.8
1999	272.9	12.0	1.2	0.7	2.7	8.4	38.6	132.1	361.4	928.4	2,338.8	6,669.0
White, female												
2003	253.8	8.1	1.1	0.4	1.7	4.4	15.1	41.1	125.2	392.0	1,315.2	5,193.6
2002	261.0	9.9	0.8	0.5	1.4	4.0	14.8	41.7	130.6	414.7	1,368.2	5,350.6
2001	267.7	9.2	1.1	0.6	1.5	4.4	13.9	41.5	134.3	429.0	1,407.9	5,582.5
2000	274.5	11.1	0.8	0.4	1.6	4.0	13.7	40.9	141.3	445.2	1,452.4	5,801.4
1999	282.4	11.6	0.9	0.5	1.8	4.3	13.8	42.2	147.8	468.7	1,509.0	5,945.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of heart (I00-I09,I11,I13,I20-I51)												
Black, both sexes												
2003	202.8	20.2	2.2	1.1	5.6	18.1	59.5	181.4	427.8	913.0	1,962.1	4,941.2
2002	205.6	21.3	1.6	1.0	4.9	18.4	57.8	180.9	442.9	924.5	2,048.8	5,115.2
2001	208.8	21.6	3.0	1.1	5.4	17.9	59.5	179.9	447.6	965.7	2,108.4	5,275.7
2000	211.8	19.8	2.0	1.1	5.6	17.2	57.4	181.5	466.0	1,002.6	2,159.7	5,389.5
1999	217.2	23.8	2.0	1.1	5.8	16.7	62.1	185.4	481.3	1,048.6	2,211.1	5,487.4
Black, male												
2003	206.0	24.1	2.5	1.2	7.3	22.2	77.1	248.1	580.9	1,195.5	2,426.6	4,850.3
2002	206.3	22.1	*	1.0	6.2	23.9	75.6	246.0	605.3	1,192.7	2,449.6	5,125.7
2001	209.0	19.5	3.2	1.0	6.9	22.8	76.7	242.6	602.2	1,245.8	2,569.3	5,459.9
2000	211.1	19.8	1.9	1.2	6.6	21.2	74.3	247.2	631.2	1,268.8	2,597.6	5,633.5
1999	218.5	23.4	2.1	1.3	7.4	21.0	81.9	255.0	660.0	1,346.4	2,623.5	5,858.8
Black, female												
2003	200.0	16.1	2.0	1.0	3.9	14.4	43.9	124.1	304.7	712.0	1,699.6	4,976.5
2002	205.0	20.6	1.8	1.0	3.7	13.5	42.1	124.9	312.3	734.0	1,821.9	5,111.2
2001	208.6	23.9	2.7	1.1	3.8	13.5	44.2	125.9	323.1	768.0	1,849.6	5,207.3
2000	212.6	19.8	2.0	1.0	4.7	13.6	42.4	125.0	332.8	815.2	1,913.1	5,298.7
1999	216.0	24.2	1.9	1.0	4.3	12.8	44.6	125.8	337.9	838.6	1,980.3	5,348.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
All races 3/, both sexes												
2003	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
2002	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0
All races, male												
2003	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
2002	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3
All races, female												
2003	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
2002	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
White, both sexes												
2003	1.4	*	*	*	*	*	0.2	0.4	1.2	3.8	10.2	21.8
2002	1.4	*	*	*	*	0.1	0.2	0.5	1.5	4.0	10.1	18.9
2001	1.3	*	*	*	*	*	0.2	0.5	1.4	4.1	10.0	18.7
2000	1.4	*	*	*	*	0.1	0.3	0.6	1.7	4.6	10.1	17.6
1999	1.4	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.5	18.9
White, male												
2003	0.9	*	*	*	*	*	0.1	0.3	0.9	3.0	8.1	17.3
2002	0.8	*	*	*	*	*	0.2	0.4	1.1	3.0	7.1	12.6
2001	0.8	*	*	*	*	*	0.2	0.4	1.1	2.8	6.9	13.3
2000	0.9	*	*	*	*	*	0.2	0.6	1.5	2.9	7.0	12.7
1999	0.8	*	*	*	*	*	0.2	0.5	1.1	3.7	7.0	11.1
White, female												
2003	1.8	*	*	*	*	*	0.2	0.5	1.4	4.6	11.6	23.8
2002	1.9	*	*	*	*	*	0.2	0.6	1.9	4.8	12.1	21.6
2001	1.8	*	*	*	*	*	0.2	0.6	1.6	5.2	12.0	21.0
2000	1.9	*	*	*	*	*	0.3	0.7	1.9	6.0	12.2	19.6
1999	2.0	*	*	*	*	*	0.3	0.7	2.2	5.7	12.8	22.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Black, both sexes												
2003	0.8	*	*	*	*	*	0.5	1.3	2.6	3.9	4.1	7.6
2002	0.7	*	*	*	*	*	0.3	0.9	2.4	2.9	4.4	*
2001	0.8	*	*	*	*	*	0.7	1.3	2.1	4.1	4.2	6.8
2000	0.7	*	*	*	*	*	0.6	1.0	2.0	3.5	3.9	8.4
1999	0.8	*	*	*	*	*	0.6	1.2	2.2	3.4	4.4	8.3
Black, male												
2003	0.6	*	*	*	*	*	*	0.9	2.7	3.0	*	*
2002	0.5	*	*	*	*	*	*	*	1.9	3.0	*	*
2001	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000	0.5	*	*	*	*	*	*	*	*	*	*	*
1999	0.5	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	1.0	*	*	*	*	*	*	1.7	2.5	4.5	5.1	8.1
2002	0.8	*	*	*	*	*	*	1.1	2.8	2.8	5.0	*
2001	1.1	*	*	*	*	*	0.9	1.7	2.6	5.3	5.3	*
2000	1.0	*	*	*	*	*	0.8	1.3	2.7	4.3	4.8	*
1999	1.0	*	*	*	*	*	0.7	1.4	2.6	4.1	4.5	11.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
All races 3/, both sexes												
2003	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
2002	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9
All races, male												
2003	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
2002	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6
All races, female												
2003	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
2002	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
White, both sexes												
2003	8.8	*	*	*	0.1	0.5	2.0	5.4	9.8	17.3	45.1	185.3
2002	8.3	*	*	*	*	0.4	1.9	5.1	9.1	16.7	42.9	177.3
2001	7.8	*	*	*	*	0.4	1.6	4.2	7.9	15.2	42.6	179.9
2000	7.6	*	*	*	*	0.3	1.5	3.7	7.6	15.0	41.5	181.8
1999	7.3	*	*	*	*	0.3	1.3	3.4	7.5	14.4	40.9	175.3
White, male												
2003	7.7	*	*	*	0.1	0.7	3.1	7.9	13.4	21.3	44.8	131.9
2002	7.1	*	*	*	*	0.7	2.8	7.4	12.7	19.9	42.6	124.8
2001	6.4	*	*	*	*	0.6	2.4	6.1	10.9	18.8	41.9	123.7
2000	6.1	*	*	*	*	0.4	2.1	5.5	10.4	18.6	39.6	129.3
1999	5.8	*	*	*	*	0.6	2.0	4.9	10.0	16.6	39.7	123.7
White, female												
2003	10.0	*	*	*	*	0.2	1.0	2.9	6.4	13.9	45.2	209.0
2002	9.5	*	*	*	*	0.2	0.9	2.8	5.8	14.1	43.1	200.1
2001	9.2	*	*	*	*	0.2	0.8	2.4	5.1	12.3	43.0	203.5
2000	9.0	*	*	*	*	0.2	0.9	1.9	5.0	11.9	42.9	203.1
1999	8.7	*	*	*	*	*	0.7	2.0	5.1	12.5	41.7	196.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart disease (I11)												
Black, both sexes												
2003	17.9	*	*	*	0.4	2.5	9.4	27.9	44.7	69.3	126.3	332.3
2002	17.2	*	*	*	*	2.8	8.6	24.3	44.4	68.0	133.0	320.4
2001	16.0	*	*	*	*	2.1	8.3	23.0	39.2	65.6	124.4	326.1
2000	16.1	*	*	*	*	2.1	7.4	21.2	42.2	66.8	131.9	340.0
1999	15.5	*	*	*	0.3	1.8	7.9	20.9	39.6	67.2	131.9	305.1
Black, male												
2003	18.1	*	*	*	*	3.0	11.2	35.0	57.1	92.3	137.1	287.8
2002	17.5	*	*	*	*	3.9	11.6	31.3	60.8	83.6	141.1	267.0
2001	15.9	*	*	*	*	2.6	10.3	30.2	49.8	79.1	137.1	284.8
2000	15.6	*	*	*	*	2.6	9.2	27.8	54.9	81.4	139.3	273.3
1999	15.2	*	*	*	*	2.1	10.1	26.3	50.6	80.4	142.3	263.6
Black, female												
2003	17.8	*	*	*	*	2.0	7.9	21.8	34.7	52.9	120.2	349.6
2002	16.9	*	*	*	*	1.7	6.0	18.3	31.2	56.8	128.5	340.9
2001	16.2	*	*	*	*	1.7	6.5	16.8	30.6	56.1	117.3	341.4
2000	16.5	*	*	*	*	1.7	5.8	15.5	31.9	56.6	127.7	364.9
1999	15.9	*	*	*	*	1.5	5.9	16.2	30.8	57.9	126.0	320.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
All races 3/, both sexes												
2003	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
2002	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1
All races, male												
2003	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
2002	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3
All races, female												
2003	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
2002	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
White, both sexes												
2003	0.9	*	*	*	*	*	0.1	0.2	0.5	1.4	6.0	23.4
2002	0.8	*	*	*	*	*	0.1	0.1	0.5	1.4	5.7	21.0
2001	0.8	*	*	*	*	*	0.1	0.2	0.5	1.5	5.7	21.0
2000	0.8	*	*	*	*	*	0.1	0.2	0.4	1.5	5.6	22.0
1999	1.0	*	*	*	*	*	0.1	0.2	0.5	1.5	7.3	26.1
White, male												
2003	0.7	*	*	*	*	*	*	0.2	0.6	1.6	6.6	24.3
2002	0.7	*	*	*	*	*	*	0.1	0.6	1.6	6.0	20.6
2001	0.7	*	*	*	*	*	*	0.2	0.5	1.6	5.9	22.7
2000	0.7	*	*	*	*	*	0.1	0.2	0.5	1.7	6.2	24.4
1999	0.8	*	*	*	*	*	*	0.2	0.6	1.7	8.2	28.8
White, female												
2003	1.0	*	*	*	*	*	*	*	0.4	1.3	5.5	22.9
2002	1.0	*	*	*	*	*	*	*	0.4	1.2	5.6	21.1
2001	1.0	*	*	*	*	*	*	0.1	0.5	1.4	5.5	20.3
2000	1.0	*	*	*	*	*	*	0.1	0.4	1.4	5.2	21.1
1999	1.1	*	*	*	*	*	*	0.1	0.5	1.4	6.7	25.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hypertensive heart and renal disease (I13)												
Black, both sexes												
2003	2.4	*	*	*	*	0.4	1.2	3.0	5.4	9.0	19.6	53.1
2002	2.4	*	*	*	*	*	0.9	2.7	6.1	9.9	19.6	56.1
2001	2.2	*	*	*	*	0.4	1.2	2.1	5.0	9.9	20.1	50.1
2000	2.2	*	*	*	*	*	0.8	2.2	4.6	9.9	22.8	50.4
1999	2.8	*	*	*	*	*	1.5	2.4	6.0	12.5	26.9	65.4
Black, male												
2003	2.3	*	*	*	*	*	1.7	4.0	7.1	8.9	18.9	50.0
2002	2.3	*	*	*	*	*	1.0	3.8	7.5	10.1	19.2	61.1
2001	2.1	*	*	*	*	*	1.6	3.2	5.9	11.9	18.5	44.3
2000	2.0	*	*	*	*	*	1.0	2.8	5.3	11.3	20.9	63.4
1999	2.5	*	*	*	*	*	1.8	3.5	7.6	13.5	25.8	59.8
Black, female												
2003	2.5	*	*	*	*	*	0.8	2.1	4.0	9.1	19.9	54.3
2002	2.5	*	*	*	*	*	0.9	1.8	4.9	9.7	19.8	54.2
2001	2.3	*	*	*	*	*	0.8	1.1	4.2	8.6	21.0	52.2
2000	2.4	*	*	*	*	*	*	1.6	4.1	9.0	23.9	45.5
1999	3.0	*	*	*	*	*	1.2	1.5	4.8	11.7	27.5	67.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
All races 3/, both sexes												
2003	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
2002	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1
All races, male												
2003	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
2002	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5
All races, female												
2003	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
2002	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
White, both sexes												
2003	178.3	*	*	*	0.3	2.5	16.4	58.8	162.0	418.9	1,159.0	3,671.7
2002	185.4	*	*	*	0.3	2.3	16.5	60.7	169.5	448.8	1,217.8	3,836.4
2001	190.5	*	*	*	0.3	2.6	15.9	60.7	175.1	465.8	1,268.5	4,022.7
2000	197.9	0.8	*	*	0.3	2.1	15.8	62.8	187.3	493.4	1,320.9	4,257.7
1999	205.1	*	*	*	0.3	2.3	16.2	64.4	194.3	524.8	1,383.7	4,389.2
White, male												
2003	185.5	*	*	*	0.5	3.8	25.1	93.5	243.5	587.9	1,529.5	4,106.1
2002	192.5	*	*	*	0.4	3.5	25.6	96.3	254.3	632.9	1,608.0	4,299.7
2001	195.8	*	*	*	0.5	3.9	24.7	96.3	262.3	653.8	1,672.4	4,494.0
2000	203.6	*	*	*	0.4	3.3	25.1	100.5	280.3	697.3	1,744.8	4,813.9
1999	210.5	*	*	*	0.4	3.4	25.5	102.8	289.0	741.3	1,827.8	4,933.9
White, female												
2003	171.4	*	*	*	0.2	1.2	7.5	24.6	85.0	274.9	908.5	3,479.3
2002	178.4	*	*	*	0.2	1.1	7.3	25.5	89.5	292.7	956.0	3,635.2
2001	185.3	*	*	*	0.2	1.3	7.0	25.6	93.1	307.2	1,000.3	3,825.5
2000	192.4	*	*	*	0.2	0.9	6.5	25.6	100.0	322.5	1,042.0	4,031.8
1999	200.0	*	*	*	0.1	1.3	6.8	26.7	105.5	344.3	1,092.6	4,170.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Ischemic heart diseases (I20-I25)												
Black, both sexes												
2003	129.2	*	*	*	0.7	4.3	24.7	99.3	273.1	615.2	1,363.0	3,310.1
2002	132.9	*	*	*	0.5	3.9	25.1	101.6	288.0	636.1	1,422.7	3,462.9
2001	136.9	*	*	*	0.9	4.6	25.6	102.3	297.9	673.0	1,479.0	3,615.1
2000	140.1	*	*	*	0.4	4.3	25.6	105.6	314.6	702.2	1,523.2	3,684.7
1999	143.7	*	*	*	0.8	3.9	27.5	108.5	320.1	735.6	1,552.7	3,805.8
Black, male												
2003	133.4	*	*	*	0.9	6.2	35.0	144.5	381.8	827.3	1,744.8	3,285.3
2002	135.3	*	*	*	*	5.1	35.2	144.3	407.3	840.8	1,748.8	3,556.2
2001	138.9	*	*	*	1.2	6.6	36.0	145.7	412.8	894.0	1,825.4	3,775.0
2000	141.5	*	*	*	*	5.9	35.6	149.4	439.2	907.9	1,877.0	3,885.2
1999	146.0	*	*	*	0.9	5.7	39.5	156.9	452.9	962.3	1,870.8	4,080.1
Black, female												
2003	125.3	*	*	*	*	2.5	15.6	60.5	185.6	464.3	1,147.2	3,319.7
2002	130.8	*	*	*	*	2.8	16.3	64.9	192.2	490.8	1,238.1	3,427.2
2001	135.2	*	*	*	*	2.8	16.4	65.0	205.3	517.1	1,284.5	3,555.7
2000	138.8	*	*	*	*	2.8	16.7	68.0	214.1	557.4	1,324.0	3,610.0
1999	141.7	*	*	*	0.7	2.3	16.9	67.0	213.6	575.8	1,374.6	3,703.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
All races 3/, both sexes												
2003	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
2002	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3
All races, male												
2003	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
2002	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1
All races, female												
2003	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
2002	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
White, both sexes												
2003	63.4	*	*	*	0.2	1.0	6.8	24.7	69.5	166.8	412.7	1,132.6
2002	67.3	*	*	*	0.1	1.0	6.9	25.9	73.5	184.2	443.3	1,201.2
2001	70.1	*	*	*	0.1	1.1	6.8	26.9	76.8	190.8	471.1	1,277.8
2000	74.1	*	*	*	0.1	0.9	6.8	28.2	84.9	206.7	502.6	1,353.5
1999	77.4	*	*	*	0.1	1.0	7.0	29.1	88.6	223.3	530.2	1,393.2
White, male												
2003	67.7	*	*	*	0.2	1.4	10.3	39.2	103.6	230.7	532.6	1,289.3
2002	71.8	*	*	*	0.2	1.4	10.5	40.4	108.9	257.3	572.0	1,386.1
2001	74.4	*	*	*	0.2	1.6	10.6	42.4	113.7	264.7	609.1	1,481.1
2000	78.8	*	*	*	0.2	1.3	10.6	44.7	125.0	287.2	652.5	1,592.9
1999	82.7	*	*	*	0.2	1.4	10.9	46.4	131.3	311.1	692.3	1,644.0
White, female												
2003	59.2	*	*	*	*	0.5	3.2	10.4	37.2	112.3	331.5	1,063.2
2002	62.8	*	*	*	*	0.5	3.3	11.5	40.1	122.2	357.0	1,120.8
2001	66.0	*	*	*	*	0.7	3.1	11.7	42.0	128.4	379.6	1,192.7
2000	69.6	*	*	*	*	0.4	2.9	11.9	47.3	139.3	404.1	1,256.2
1999	72.3	*	*	*	*	0.6	3.1	12.1	48.6	150.1	423.9	1,292.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute myocardial infarction (I21-I22)												
Black, both sexes												
2003	45.7	*	*	*	*	1.6	10.1	37.5	98.6	220.0	479.9	1,094.9
2002	48.5	*	*	*	*	1.6	9.7	37.9	106.9	238.8	517.3	1,201.2
2001	50.0	*	*	*	0.3	1.8	10.0	39.2	109.8	252.6	537.8	1,239.9
2000	52.2	*	*	*	*	1.7	9.9	41.1	120.1	268.7	571.1	1,275.7
1999	53.3	*	*	*	*	1.6	10.3	40.9	120.9	279.5	584.6	1,318.8
Black, male												
2003	46.6	*	*	*	*	2.3	13.8	53.0	133.3	283.0	600.2	1,113.5
2002	48.3	*	*	*	*	2.1	13.3	52.0	141.6	300.1	623.4	1,280.1
2001	49.2	*	*	*	*	2.5	13.2	51.9	143.8	317.1	638.0	1,382.0
2000	52.0	*	*	*	*	2.2	13.4	54.9	160.3	335.8	687.3	1,418.5
1999	53.4	*	*	*	*	2.3	14.2	56.6	165.5	353.6	685.6	1,498.7
Black, female												
2003	44.9	*	*	*	*	1.0	6.7	24.1	70.7	175.2	411.9	1,087.7
2002	48.8	*	*	*	*	1.2	6.5	25.9	79.1	195.3	457.2	1,171.0
2001	50.6	*	*	*	*	1.2	7.1	28.2	82.4	207.0	481.5	1,187.2
2000	52.5	*	*	*	*	1.3	6.8	29.1	87.7	221.6	505.6	1,222.5
1999	53.2	*	*	*	*	0.9	6.9	27.4	85.1	227.4	528.1	1,251.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
All races 3/, both sexes												
2003	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
2002	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9
All races, male												
2003	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
2002	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3
All races, female												
2003	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
2002	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
White, both sexes												
2003	1.1	*	*	*	*	*	0.2	0.6	1.3	3.0	6.3	20.2
2002	1.2	*	*	*	*	*	0.2	0.8	1.6	3.3	7.0	21.4
2001	1.2	*	*	*	*	*	0.2	0.7	1.6	3.3	6.8	21.5
2000	1.2	*	*	*	*	*	0.2	0.7	1.8	3.3	7.1	22.1
1999	1.2	*	*	*	*	*	0.1	0.6	1.7	3.5	6.8	22.4
White, male												
2003	1.1	*	*	*	*	*	0.3	0.9	1.8	4.1	7.6	19.7
2002	1.3	*	*	*	*	*	0.2	1.1	2.3	4.4	8.6	21.9
2001	1.2	*	*	*	*	*	0.2	1.1	2.4	4.4	8.5	20.7
2000	1.3	*	*	*	*	*	0.3	1.1	2.6	4.4	8.9	21.0
1999	1.2	*	*	*	*	*	0.2	0.8	2.5	4.7	8.6	22.9
White, female												
2003	1.1	*	*	*	*	*	*	0.3	0.8	2.0	5.5	20.5
2002	1.2	*	*	*	*	*	*	0.4	0.9	2.3	5.9	21.1
2001	1.2	*	*	*	*	*	*	0.3	0.9	2.4	5.7	21.7
2000	1.2	*	*	*	*	*	*	0.3	0.9	2.3	5.9	22.6
1999	1.2	*	*	*	*	*	*	0.3	1.1	2.5	5.6	22.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute ischemic heart diseases (I24)												
Black, both sexes												
2003	1.1	*	*	*	*	*	0.5	1.3	3.0	5.5	9.9	19.8
2002	1.2	*	*	*	*	*	0.3	1.7	3.0	5.9	9.3	21.1
2001	1.2	*	*	*	*	*	0.5	1.5	3.9	5.7	8.5	24.0
2000	1.2	*	*	*	*	*	0.4	1.5	4.2	5.6	10.8	21.9
1999	1.2	*	*	*	*	*	0.5	1.8	3.6	5.7	10.7	17.2
Black, male												
2003	1.4	*	*	*	*	*	*	1.9	4.8	7.6	16.1	24.0
2002	1.4	*	*	*	*	*	*	2.7	4.2	9.1	11.8	21.4
2001	1.4	*	*	*	*	*	*	2.4	5.5	8.5	10.3	23.8
2000	1.5	*	*	*	*	*	*	2.4	6.1	8.6	13.9	*
1999	1.4	*	*	*	*	*	0.7	2.6	6.1	8.2	12.6	*
Black, female												
2003	0.9	*	*	*	*	*	*	*	1.5	4.0	6.4	18.2
2002	1.0	*	*	*	*	*	*	0.8	2.0	3.6	8.0	20.9
2001	1.1	*	*	*	*	*	*	*	2.6	3.7	7.4	24.0
2000	1.0	*	*	*	*	*	*	*	2.7	3.5	9.0	22.3
1999	1.0	*	*	*	*	*	*	1.1	1.5	3.8	9.6	16.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
All races 3/, both sexes												
2003	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
2002	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0
All races, male												
2003	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
2002	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1
All races, female												
2003	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
2002	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
White, both sexes												
2003	113.8	*	*	*	0.2	1.5	9.4	33.5	91.3	249.1	740.0	2,518.8
2002	116.9	*	*	*	0.2	1.4	9.4	34.0	94.4	261.3	767.5	2,613.9
2001	119.1	*	*	*	0.2	1.4	8.9	33.1	96.7	271.7	790.5	2,723.5
2000	122.6	*	*	*	0.2	1.2	8.9	33.9	100.6	283.4	811.2	2,882.1
1999	126.5	*	*	*	0.1	1.3	9.0	34.7	104.0	298.0	846.7	2,973.6
White, male												
2003	116.6	*	*	*	0.2	2.3	14.4	53.4	138.1	353.1	989.3	2,797.1
2002	119.4	*	*	*	0.2	2.1	14.9	54.8	143.1	371.2	1,027.4	2,891.6
2001	120.1	*	*	*	0.3	2.2	13.9	52.9	146.2	384.6	1,054.9	2,992.2
2000	123.5	*	*	*	0.2	1.9	14.2	54.8	152.7	405.6	1,083.4	3,200.0
1999	126.5	*	*	*	0.2	2.0	14.4	55.5	155.2	425.5	1,126.9	3,267.1
White, female												
2003	111.0	*	*	*	*	0.7	4.2	13.9	47.0	160.6	571.4	2,395.5
2002	114.4	*	*	*	*	0.6	3.9	13.6	48.5	168.1	593.1	2,493.2
2001	118.1	*	*	*	*	0.6	3.9	13.6	50.2	176.4	615.0	2,611.0
2000	121.6	*	*	*	*	0.5	3.5	13.4	51.8	180.9	632.0	2,753.0
1999	126.5	*	*	*	*	0.6	3.6	14.3	55.9	191.7	663.1	2,855.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)												
Black, both sexes												
2003	82.3	*	*	*	0.4	2.6	14.2	60.6	171.5	389.7	873.2	2,195.3
2002	83.2	*	*	*	*	2.2	15.1	61.9	178.1	391.4	896.1	2,240.6
2001	85.8	*	*	*	0.5	2.7	15.1	61.6	184.2	414.8	932.7	2,351.2
2000	86.6	*	*	*	*	2.5	15.3	63.1	190.2	427.8	941.3	2,387.1
1999	89.3	*	*	*	0.5	2.3	16.8	65.8	195.7	450.4	957.4	2,469.8
Black, male												
2003	85.5	*	*	*	*	3.9	20.6	89.6	243.7	536.7	1,128.5	2,147.8
2002	85.7	*	*	*	*	2.9	21.7	89.5	261.4	531.5	1,113.6	2,254.6
2001	88.2	*	*	*	0.7	4.2	22.2	91.3	263.6	568.3	1,177.1	2,369.1
2000	88.0	*	*	*	*	3.6	21.6	92.1	272.7	563.5	1,175.7	2,446.0
1999	91.1	*	*	*	*	3.3	24.6	97.7	281.3	600.5	1,172.6	2,561.5
Black, female												
2003	79.4	*	*	*	*	1.4	8.5	35.7	113.4	285.0	728.9	2,213.8
2002	81.0	*	*	*	*	1.5	9.4	38.2	111.1	291.9	772.9	2,235.2
2001	83.5	*	*	*	*	1.5	8.9	36.1	120.3	306.4	795.5	2,344.5
2000	85.3	*	*	*	*	1.5	9.7	38.1	123.7	332.3	809.4	2,365.2
1999	87.6	*	*	*	*	1.3	9.9	38.5	127.0	344.6	836.9	2,435.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
All races 3/, both sexes												
2003	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
2002	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4
All races, male												
2003	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
2002	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6
All races, female												
2003	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
2002	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
White, both sexes												
2003	23.6	*	*	*	*	0.5	3.2	12.6	29.0	56.5	135.5	439.3
2002	24.1	*	*	*	*	0.4	3.1	12.1	29.4	58.1	138.4	470.6
2001	24.3	*	*	*	*	0.4	3.0	11.6	28.8	58.3	144.5	489.7
2000	25.1	*	*	*	*	0.3	2.8	11.8	29.4	61.2	149.2	529.5
1999	26.1	*	*	*	*	0.3	3.0	11.3	30.4	64.8	157.0	557.5
White, male												
2003	25.3	*	*	*	*	0.8	5.0	20.8	44.8	80.0	171.9	458.7
2002	25.5	*	*	*	*	0.7	5.1	19.9	45.9	82.9	174.7	488.0
2001	25.3	*	*	*	*	0.6	4.7	19.2	45.1	82.7	184.2	498.6
2000	26.0	*	*	*	*	0.5	4.6	19.6	46.3	87.8	187.5	545.6
1999	26.6	*	*	*	*	0.5	4.9	18.6	46.9	92.5	195.2	571.2
White, female												
2003	21.9	*	*	*	*	0.2	1.4	4.5	14.1	36.5	110.8	430.7
2002	22.7	*	*	*	*	0.1	1.1	4.5	13.8	37.0	114.0	463.0
2001	23.3	*	*	*	*	0.2	1.2	4.1	13.4	37.8	118.2	486.0
2000	24.3	*	*	*	*	*	1.0	4.2	13.5	39.0	123.9	522.9
1999	25.6	*	*	*	*	0.2	1.0	4.2	15.0	41.6	131.9	552.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)												
Black, both sexes												
2003	26.0	*	*	*	*	0.9	4.9	25.7	69.7	128.1	241.6	543.2
2002	27.0	*	*	*	*	0.8	5.8	25.5	71.4	136.3	259.0	575.5
2001	27.7	*	*	*	*	0.7	5.6	25.3	77.5	143.4	266.5	594.5
2000	28.5	*	*	*	*	0.7	5.1	26.9	78.2	145.5	278.4	646.6
1999	29.9	*	*	*	*	0.7	6.3	27.5	81.4	158.6	291.2	668.6
Black, male												
2003	30.5	*	*	*	*	1.5	7.7	40.4	106.0	190.3	331.4	583.9
2002	31.1	*	*	*	*	1.1	8.6	38.7	111.4	197.2	357.2	599.3
2001	31.5	*	*	*	*	0.9	8.5	39.2	117.4	208.2	351.7	609.3
2000	32.3	*	*	*	*	1.0	7.8	42.0	120.0	207.3	369.7	701.2
1999	33.8	*	*	*	*	1.0	9.6	42.8	123.3	229.3	380.0	715.9
Black, female												
2003	21.9	*	*	*	*	*	2.6	13.1	40.5	83.9	190.9	527.4
2002	23.3	*	*	*	*	*	3.3	14.2	39.2	93.0	203.5	566.4
2001	24.3	*	*	*	*	*	3.1	13.3	45.4	97.7	218.6	589.0
2000	25.0	*	*	*	*	*	2.8	14.0	44.6	102.0	227.1	626.3
1999	26.4	*	*	*	*	*	3.4	14.4	47.8	108.8	241.5	650.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
All races 3/, both sexes												
2003	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
2002	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6
All races, male												
2003	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
2002	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4
All races, female												
2003	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
2002	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
White, both sexes												
2003	90.2	*	*	*	0.1	1.0	6.2	20.9	62.2	192.6	604.6	2,079.5
2002	92.8	*	*	*	0.1	0.9	6.3	21.9	65.1	203.2	629.1	2,143.3
2001	94.8	*	*	*	0.1	1.0	5.9	21.5	67.9	213.4	646.0	2,233.8
2000	97.4	*	*	*	0.1	0.9	6.1	22.1	71.2	222.2	662.0	2,352.7
1999	100.4	*	*	*	0.1	1.0	6.1	23.4	73.5	233.2	689.7	2,416.1
White, male												
2003	91.3	*	*	*	0.2	1.5	9.4	32.5	93.3	273.1	817.4	2,338.4
2002	93.9	*	*	*	0.1	1.4	9.8	34.9	97.3	288.3	852.7	2,403.6
2001	94.8	*	*	*	0.2	1.6	9.2	33.7	101.1	301.9	870.7	2,493.6
2000	97.5	*	*	*	0.1	1.4	9.6	35.2	106.4	317.8	895.9	2,654.4
1999	99.9	*	*	*	0.2	1.4	9.5	36.9	108.3	333.0	931.7	2,695.9
White, female												
2003	89.1	*	*	*	*	0.5	2.9	9.4	32.9	124.0	460.7	1,964.9
2002	91.7	*	*	*	*	0.4	2.8	9.1	34.7	131.1	479.1	2,030.2
2001	94.9	*	*	*	*	0.4	2.6	9.6	36.7	138.6	496.8	2,125.0
2000	97.3	*	*	*	*	0.4	2.5	9.3	38.3	141.9	508.1	2,230.1
1999	100.8	*	*	*	*	0.5	2.6	10.1	40.9	150.1	531.1	2,303.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Black, both sexes												
2003	56.3	*	*	*	0.3	1.8	9.3	34.9	101.8	261.5	631.6	1,652.1
2002	56.2	*	*	*	*	1.4	9.4	36.4	106.7	255.1	637.0	1,665.1
2001	58.0	*	*	*	0.4	2.0	9.5	36.3	106.7	271.4	666.3	1,756.7
2000	58.1	*	*	*	*	1.8	10.2	36.1	112.0	282.3	662.9	1,740.5
1999	59.3	*	*	*	0.3	1.5	10.4	38.3	114.2	291.8	666.2	1,801.2
Black, male												
2003	55.0	*	*	*	*	2.4	13.0	49.1	137.6	346.5	797.1	1,563.9
2002	54.5	*	*	*	*	1.8	13.1	50.8	150.1	334.3	756.5	1,655.3
2001	56.7	*	*	*	*	3.2	13.7	52.1	146.2	360.1	825.4	1,759.8
2000	55.8	*	*	*	*	2.6	13.9	50.2	152.7	356.1	806.0	1,744.8
1999	57.3	*	*	*	*	2.3	15.0	54.9	157.9	371.2	792.6	1,845.5
Black, female												
2003	57.5	*	*	*	*	1.2	6.0	22.6	72.9	201.1	538.0	1,686.4
2002	57.7	*	*	*	*	1.1	6.1	24.0	71.9	198.9	569.4	1,668.8
2001	59.2	*	*	*	*	1.0	5.9	22.8	75.0	208.7	576.9	1,755.5
2000	60.3	*	*	*	*	1.1	6.9	24.1	79.2	230.3	582.3	1,738.9
1999	61.2	*	*	*	*	0.8	6.4	24.1	79.1	235.8	595.4	1,784.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
All races 3/, both sexes												
2003	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
2002	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9
All races, male												
2003	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
2002	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9
All races, female												
2003	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
2002	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
White, both sexes												
2003	62.2	8.2	0.9	0.5	1.8	3.6	8.6	18.1	43.1	120.3	383.4	1,461.4
2002	62.6	10.0	0.9	0.5	1.7	3.6	8.7	18.4	43.9	124.2	390.3	1,475.5
2001	62.3	9.3	1.1	0.5	1.6	3.6	8.1	18.4	44.6	125.6	387.6	1,509.6
2000	62.5	10.6	1.0	0.5	1.7	3.6	8.1	18.2	46.1	128.1	389.6	1,541.6
1999	62.9	11.0	1.0	0.5	1.9	3.6	8.4	18.1	47.2	132.2	395.1	1,543.3
White, male												
2003	54.7	8.8	0.8	0.6	2.2	4.4	10.8	23.3	54.7	147.2	441.1	1,467.6
2002	54.9	10.5	1.0	0.5	2.3	4.5	11.1	24.3	55.3	150.4	448.3	1,482.1
2001	54.1	9.6	1.2	0.5	2.0	4.2	10.4	24.0	55.9	152.2	448.7	1,503.4
2000	54.6	10.8	1.2	0.6	2.1	4.3	10.3	23.9	59.2	157.4	449.5	1,580.5
1999	55.1	11.4	1.1	0.6	2.2	4.3	10.9	23.7	60.6	165.1	456.1	1,571.5
White, female												
2003	69.6	7.6	1.0	0.4	1.5	2.8	6.3	13.0	32.1	97.3	344.4	1,458.6
2002	70.2	9.6	0.8	0.5	1.1	2.6	6.4	12.6	33.1	101.9	351.4	1,472.6
2001	70.4	8.9	1.1	0.5	1.3	2.9	5.7	12.8	34.0	103.1	347.1	1,512.2
2000	70.2	10.4	0.7	0.4	1.4	2.8	6.0	12.5	33.9	103.4	350.1	1,525.8
1999	70.5	10.7	0.8	0.5	1.6	2.8	6.0	12.6	34.5	104.7	355.1	1,531.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other heart diseases (I26-I51)												
Black, both sexes												
2003	52.5	19.6	2.0	1.0	4.4	10.9	23.6	49.9	102.1	215.7	449.2	1,238.1
2002	52.5	20.2	1.5	0.8	4.1	11.3	22.8	51.4	102.0	207.6	469.1	1,270.4
2001	52.8	20.1	2.9	1.0	4.2	10.5	23.7	51.1	103.5	213.0	480.6	1,277.7
2000	52.7	19.0	1.9	1.1	4.7	10.3	23.0	51.6	102.6	220.2	477.9	1,306.0
1999	54.4	22.3	2.0	1.0	4.5	10.5	24.7	52.4	113.3	229.9	495.2	1,302.8
Black, male												
2003	51.5	23.2	2.2	1.1	5.7	12.4	28.6	63.8	132.3	264.0	523.5	1,220.9
2002	50.8	20.9	*	0.9	5.1	14.2	27.4	65.9	127.9	255.2	537.3	1,238.3
2001	51.7	17.7	3.2	0.9	5.3	12.8	28.3	62.5	132.1	258.5	585.9	1,349.1
2000	51.5	18.9	1.9	1.2	5.4	12.2	28.1	66.5	130.5	266.0	558.0	1,402.3
1999	54.3	21.8	2.1	1.1	5.8	12.7	30.1	67.3	147.3	287.8	580.2	1,454.1
Black, female												
2003	53.4	15.8	1.8	0.8	3.2	9.4	19.1	38.0	77.8	181.3	407.1	1,244.8
2002	54.0	19.4	1.8	0.8	3.1	8.7	18.7	39.0	81.1	173.8	430.6	1,282.7
2001	53.8	22.7	2.5	1.0	3.0	8.5	19.6	41.3	80.5	180.9	421.5	1,251.2
2000	53.9	19.2	2.0	1.0	4.1	8.7	18.6	38.7	80.1	187.9	432.8	1,270.2
1999	54.5	22.8	1.8	0.8	3.3	8.5	19.9	39.7	86.1	189.1	447.7	1,246.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
All races 3/, both sexes												
2003	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
2002	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9
All races, male												
2003	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
2002	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7
All races, female												
2003	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
2002	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
White, both sexes												
2003	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.1
2002	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001	0.4	*	*	*	*	0.1	0.2	0.4	0.5	1.2	2.4	2.6
2000	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.2	2.4	2.7
1999	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.4	3.0
White, male												
2003	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.3	4.4
2002	0.5	*	*	*	*	0.2	0.2	0.5	0.6	1.5	3.2	3.6
2001	0.5	*	*	*	*	*	0.2	0.5	0.7	1.3	3.5	3.9
2000	0.5	*	*	*	*	*	0.3	0.5	0.8	1.4	3.5	4.2
1999	0.5	*	*	*	*	*	0.2	0.4	0.7	1.4	3.4	5.1
White, female												
2003	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.6
2002	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001	0.3	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000	0.3	*	*	*	*	*	0.2	0.2	0.4	1.0	1.7	2.1
1999	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and subacute endocarditis (I33)												
Black, both sexes												
2003	0.6	*	*	*	*	*	0.6	1.3	1.9	2.2	3.0	*
2002	0.5	*	*	*	*	*	0.4	1.2	1.6	2.3	*	*
2001	0.6	*	*	*	*	*	0.7	1.3	1.6	2.6	4.1	*
2000	0.6	*	*	*	*	*	0.5	1.3	1.6	1.9	2.8	*
1999	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*
Black, male												
2003	0.7	*	*	*	*	*	*	2.0	2.2	*	*	*
2002	0.6	*	*	*	*	*	*	1.5	1.9	2.9	*	*
2001	0.8	*	*	*	*	*	0.7	1.9	2.1	3.8	*	*
2000	0.6	*	*	*	*	*	0.8	1.8	*	*	*	*
1999	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*
Black, female												
2003	0.5	*	*	*	*	*	*	*	1.7	2.0	*	*
2002	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001	0.5	*	*	*	*	*	0.7	*	*	*	3.3	*
2000	0.5	*	*	*	*	*	*	1.0	1.6	*	*	*
1999	0.5	*	*	*	*	*	*	*	1.6	2.8	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
All races 3/, both sexes												
2003	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
2002	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0
All races, male												
2003	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
2002	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5
All races, female												
2003	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
2002	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
White, both sexes												
2003	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	1.7
2002	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.1	2.2
2001	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.9
2000	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.2	1.8
1999	0.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.3	2.0
White, male												
2003	0.3	*	*	*	0.1	*	0.2	0.3	0.4	1.1	1.3	1.7
2002	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.1	2.5
2001	0.3	*	*	*	*	0.2	0.2	0.3	0.5	0.7	1.2	2.0
2000	0.3	*	*	*	*	*	0.2	0.2	0.4	0.7	1.5	2.1
1999	0.3	*	*	*	*	0.1	0.3	0.3	0.6	0.8	1.4	2.5
White, female												
2003	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.1	1.7
2002	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.0
2001	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.9
2000	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.0	1.6
1999	0.3	*	*	*	*	*	0.2	0.3	0.4	0.7	1.2	1.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Black, both sexes												
2003	0.4	*	*	*	*	0.4	0.5	0.5	*	*	*	*
2002	0.4	*	*	*	*	*	*	0.6	0.8	*	2.1	*
2001	0.4	*	*	*	*	*	0.4	0.7	*	*	*	*
2000	0.4	*	*	*	*	*	0.4	0.5	*	*	*	*
1999	0.4	*	*	*	*	*	0.5	0.5	*	1.2	*	*
Black, male												
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	*	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
All races 3/, both sexes												
2003	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
2002	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9
All races, male												
2003	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
2002	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0
All races, female												
2003	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
2002	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
White, both sexes												
2003	21.8	0.6	*	*	0.1	0.2	0.5	1.9	7.8	32.2	134.3	660.4
2002	21.6	*	*	*	*	0.2	0.5	1.9	8.0	32.5	133.3	664.5
2001	22.1	*	*	*	*	0.2	0.5	2.0	8.5	33.0	137.0	695.1
2000	21.8	*	*	*	*	0.1	0.4	1.8	8.2	33.4	136.7	703.7
1999	21.6	*	*	*	*	0.1	0.5	1.7	8.0	34.1	136.7	699.2
White, male												
2003	17.1	*	*	*	*	0.2	0.6	2.3	9.6	38.6	150.4	650.8
2002	16.7	*	*	*	*	0.2	0.6	2.5	9.5	37.7	150.3	641.9
2001	16.9	*	*	*	*	0.2	0.6	2.6	10.3	39.0	154.6	666.5
2000	16.7	*	*	*	*	0.1	0.6	2.2	10.0	40.0	154.6	690.0
1999	16.4	*	*	*	*	*	0.6	2.1	9.8	41.3	152.1	680.0
White, female												
2003	26.3	*	*	*	*	0.1	0.3	1.4	6.1	26.8	123.4	664.7
2002	26.4	*	*	*	*	0.1	0.3	1.4	6.5	28.1	121.8	674.3
2001	27.1	*	*	*	*	0.2	0.4	1.4	6.9	28.0	125.3	707.1
2000	26.7	*	*	*	*	*	0.3	1.3	6.6	27.9	124.9	709.3
1999	26.6	*	*	*	*	*	0.3	1.2	6.3	28.0	126.6	707.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Heart failure (I50)												
Black, both sexes												
2003	14.0	*	*	*	*	0.7	2.3	6.8	20.6	54.3	150.4	531.8
2002	13.7	*	*	*	*	0.8	1.8	7.2	18.6	53.2	156.9	528.0
2001	13.8	*	*	*	*	0.6	1.9	6.4	19.6	54.5	157.3	543.8
2000	13.6	*	*	*	*	0.4	1.6	5.7	18.6	55.4	154.4	558.4
1999	13.8	*	*	*	*	0.5	1.6	6.4	18.9	56.4	163.4	552.9
Black, male												
2003	11.8	*	*	*	*	0.8	2.7	8.3	26.3	63.2	163.9	513.0
2002	11.5	*	*	*	*	1.0	2.2	9.0	22.9	64.7	174.4	479.2
2001	11.6	*	*	*	*	0.9	2.2	7.4	22.2	65.3	183.3	541.2
2000	11.2	*	*	*	*	*	1.7	7.1	21.4	61.9	173.5	581.2
1999	11.8	*	*	*	*	*	1.9	8.4	22.9	65.2	179.2	603.5
Black, female												
2003	15.9	*	*	*	*	*	1.9	5.6	16.0	48.0	142.7	539.2
2002	15.8	*	*	*	*	*	1.5	5.6	15.1	45.1	147.0	546.7
2001	15.8	*	*	*	*	*	1.7	5.5	17.6	46.9	142.6	544.8
2000	15.7	*	*	*	*	*	1.5	4.5	16.4	50.9	143.7	549.9
1999	15.7	*	*	*	*	*	1.4	4.7	15.7	50.2	154.6	534.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
All races 3/, both sexes												
2003	38.0	8.6	0.9	0.5	1.9	4.1	9.1	18.2	38.1	91.2	246.0	781.5
2002	38.5	10.5	0.8	0.5	1.9	4.0	9.3	18.6	38.8	93.7	254.4	793.7
2001	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0
1999	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1
All races, male												
2003	35.8	9.2	0.9	0.6	2.3	4.8	11.3	23.3	48.5	112.1	287.0	795.9
2002	36.3	10.9	0.8	0.5	2.5	5.0	11.5	24.1	49.0	114.8	294.9	821.3
2001	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7
All races, female												
2003	40.1	7.9	1.0	0.4	1.5	3.4	7.0	13.3	28.4	73.7	218.6	775.1
2002	40.6	10.0	0.8	0.4	1.3	3.1	7.1	13.2	29.3	76.2	227.6	781.7
2001	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
White, both sexes												
2003	39.8	7.1	0.7	0.4	1.6	3.3	7.7	15.6	34.3	86.0	245.4	796.1
2002	40.4	9.1	0.7	0.4	1.6	3.2	7.9	15.9	34.9	89.7	253.4	806.1
2001	39.6	8.2	0.9	0.4	1.5	3.2	7.2	15.7	35.2	90.7	246.9	809.9
2000	40.0	9.6	0.8	0.4	1.6	3.3	7.3	15.8	37.0	92.8	249.3	833.4
1999	40.6	10.3	0.8	0.5	1.8	3.3	7.6	15.9	38.1	96.1	254.7	839.0
White, male												
2003	36.9	7.4	0.7	0.5	1.9	4.0	9.7	20.2	43.9	106.3	286.1	810.7
2002	37.5	9.5	0.8	0.5	2.1	4.0	10.0	21.0	44.6	110.3	293.7	834.2
2001	36.5	8.2	0.9	0.5	1.8	3.8	9.4	20.6	44.6	111.2	289.4	831.0
2000	37.1	9.4	0.9	0.5	1.9	4.0	9.2	20.9	48.0	115.3	290.0	884.2
1999	37.9	10.7	1.0	0.5	2.0	4.0	9.8	20.9	49.4	121.6	299.2	883.9
White, female												
2003	42.6	6.8	0.8	0.3	1.3	2.6	5.7	11.1	25.2	68.8	217.9	789.7
2002	43.2	8.7	0.6	0.4	1.0	2.3	5.8	10.9	25.8	72.2	226.4	793.9
2001	42.6	8.2	0.8	0.4	1.1	2.6	5.0	11.0	26.4	73.3	218.8	801.1
2000	42.8	9.7	0.7	0.3	1.2	2.5	5.4	10.8	26.6	73.9	222.6	812.7
1999	43.2	10.0	0.7	0.4	1.5	2.6	5.4	11.0	27.4	74.9	225.5	821.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)												
Black, both sexes												
2003	37.5	15.0	1.6	0.8	3.9	9.7	20.3	41.3	79.2	158.2	294.5	703.7
2002	37.8	17.2	1.1	0.7	3.8	10.1	20.3	42.4	81.0	150.9	308.2	738.5
2001	38.0	17.2	2.3	0.9	3.7	9.6	20.7	42.8	81.5	155.1	318.1	730.2
2000	38.2	15.6	1.4	0.9	4.3	9.4	20.5	44.0	81.7	161.8	319.6	742.0
1999	39.6	20.2	1.4	0.8	4.2	9.6	21.8	44.3	91.7	169.8	328.2	746.1
Black, male												
2003	38.6	17.8	1.8	0.9	5.0	10.9	24.8	52.9	103.3	197.6	355.1	707.9
2002	38.3	18.9	*	0.8	4.7	12.7	24.3	54.5	102.0	186.7	359.2	753.7
2001	38.9	15.0	2.6	0.8	4.8	11.4	25.1	52.4	106.8	188.2	395.5	805.6
2000	39.2	14.7	1.6	1.0	4.9	11.2	25.0	57.2	106.9	200.2	379.7	816.5
1999	41.5	19.8	*	1.1	5.4	11.5	26.6	56.4	121.0	218.8	398.1	849.5
Black, female												
2003	36.5	12.1	*	0.7	2.7	8.5	16.2	31.4	59.7	130.2	260.2	702.0
2002	37.4	15.4	*	*	2.9	7.7	16.7	32.0	64.1	125.5	279.3	732.8
2001	37.2	19.5	2.0	0.9	2.6	7.9	16.7	34.5	61.1	131.7	274.6	702.2
2000	37.3	16.5	*	0.8	3.7	7.8	16.5	32.7	61.4	134.8	285.7	714.3
1999	37.9	20.5	*	*	3.0	7.9	17.5	34.0	68.2	135.1	289.0	707.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
All races 3/, both sexes												
2003	7.5	*	*	*	0.1	0.2	0.8	2.5	6.3	16.9	51.7	188.9
2002	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	16.0	48.2	180.4
2001	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1
All races, male												
2003	5.7	*	*	*	*	0.3	1.0	3.3	7.4	18.4	48.9	153.5
2002	5.4	*	*	*	*	0.2	1.0	2.8	6.6	17.1	47.3	153.4
2001	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4
All races, female												
2003	9.3	*	*	*	*	0.2	0.7	1.8	5.3	15.7	53.5	204.6
2002	8.6	*	*	*	*	0.2	0.6	1.8	4.9	15.1	48.8	192.1
2001	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
White, both sexes												
2003	7.2	*	*	*	*	0.1	0.5	1.7	4.4	13.0	46.2	181.0
2002	6.6	*	*	*	*	0.1	0.4	1.5	4.1	12.8	42.6	170.4
2001	6.3	*	*	*	*	0.1	0.4	1.4	4.0	12.1	42.3	163.3
2000	6.0	*	*	*	*	0.1	0.4	1.4	4.1	11.8	40.4	153.2
1999	5.6	*	*	*	*	0.1	0.4	1.3	3.7	12.1	37.9	142.0
White, male												
2003	5.2	*	*	*	*	0.1	0.7	2.2	5.2	14.5	43.8	145.3
2002	4.9	*	*	*	*	0.1	0.5	1.9	4.9	13.8	41.5	145.2
2001	4.7	*	*	*	*	0.2	0.5	1.8	4.5	13.2	40.4	142.2
2000	4.5	*	*	*	*	0.1	0.6	1.9	5.0	12.6	40.3	132.5
1999	4.2	*	*	*	*	0.2	0.5	1.7	4.5	13.4	37.3	121.0
White, female												
2003	9.1	*	*	*	*	0.1	0.4	1.2	3.6	11.7	47.8	196.8
2002	8.3	*	*	*	*	*	0.3	1.0	3.3	11.9	43.4	181.3
2001	8.0	*	*	*	*	*	0.3	1.0	3.5	11.3	43.6	172.2
2000	7.4	*	*	*	*	*	0.3	0.9	3.2	11.2	40.4	161.6
1999	7.0	*	*	*	*	*	0.2	1.0	3.0	11.0	38.3	150.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Essential (primary) hypertension and hypertensive renal disease (I10,I12)												
Black, both sexes												
2003	11.8	*	*	*	*	1.0	3.1	9.3	24.1	54.4	119.5	301.1
2002	11.1	*	*	*	*	0.8	3.3	9.0	21.0	47.6	118.0	308.8
2001	10.9	*	*	*	*	1.0	3.1	9.9	22.0	47.4	112.6	288.0
2000	10.6	*	*	*	*	0.6	3.1	9.7	22.9	48.7	107.1	283.4
1999	10.3	*	*	*	*	0.7	3.1	9.1	21.5	46.7	111.7	269.4
Black, male												
2003	10.4	*	*	*	*	1.3	3.4	12.7	29.3	60.5	119.6	274.2
2002	9.6	*	*	*	*	0.9	4.1	11.0	23.5	52.0	127.4	274.5
2001	9.7	*	*	*	*	1.1	3.8	12.9	25.5	54.4	118.3	275.7
2000	9.1	*	*	*	*	*	3.7	11.9	27.8	50.5	109.0	269.9
1999	9.1	*	*	*	*	*	3.7	11.6	24.6	51.3	118.7	266.0
Black, female												
2003	13.0	*	*	*	*	*	2.7	6.4	20.0	50.1	119.4	311.5
2002	12.5	*	*	*	*	0.7	2.5	7.2	19.0	44.5	112.7	322.0
2001	11.9	*	*	*	*	0.9	2.4	7.4	19.2	42.5	109.4	292.6
2000	12.0	*	*	*	*	*	2.6	7.7	18.9	47.5	106.0	288.5
1999	11.4	*	*	*	*	0.7	2.6	7.0	19.0	43.4	107.7	270.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
All races 3/, both sexes												
2003	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
2002	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8
All races, male												
2003	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
2002	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8
All races, female												
2003	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
2002	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
White, both sexes												
2003	57.0	2.0	0.2	0.2	0.5	1.2	4.2	11.4	29.4	103.7	403.0	1,382.2
2002	59.5	2.3	0.3	0.2	0.4	1.2	4.2	11.6	31.4	110.7	423.5	1,459.5
2001	60.5	1.9	0.3	0.2	0.4	1.3	4.4	11.4	31.7	113.0	436.4	1,511.2
2000	62.8	2.5	0.3	0.2	0.5	1.3	4.5	12.4	34.8	119.4	452.4	1,601.6
1999	63.3	2.3	0.3	0.2	0.4	1.3	4.4	11.7	34.0	121.1	461.6	1,631.6
White, male												
2003	44.2	2.0	*	0.2	0.5	1.3	4.4	12.9	33.3	117.3	422.4	1,247.0
2002	45.7	2.6	0.4	0.2	0.5	1.2	4.2	12.9	35.6	123.8	437.5	1,327.4
2001	46.6	2.2	*	0.2	0.5	1.4	4.5	12.7	36.1	128.5	458.8	1,386.2
2000	48.4	3.0	*	0.2	0.5	1.4	4.5	13.6	39.7	133.8	480.0	1,490.7
1999	48.7	2.8	*	0.2	0.5	1.3	4.6	12.8	38.9	136.0	490.3	1,522.7
White, female												
2003	69.5	2.0	*	0.1	0.4	1.2	4.0	10.0	25.8	92.1	389.9	1,442.1
2002	73.0	2.0	*	0.2	0.3	1.2	4.3	10.4	27.4	99.5	414.1	1,516.9
2001	74.0	1.6	0.4	0.2	0.4	1.3	4.2	10.2	27.6	99.9	421.6	1,563.5
2000	76.9	2.0	*	0.2	0.4	1.3	4.4	11.2	30.2	107.3	434.2	1,646.7
1999	77.6	1.9	*	0.1	0.4	1.2	4.3	10.7	29.4	108.6	442.8	1,675.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cerebrovascular diseases (I60-I69)												
Black, both sexes												
2003	49.3	4.9	*	*	0.9	3.1	13.8	41.0	89.8	201.1	531.1	1,348.7
2002	50.0	5.5	*	0.4	0.6	3.1	12.8	40.7	88.0	215.1	552.3	1,383.9
2001	51.1	6.6	*	*	0.8	2.9	13.2	42.6	91.1	222.5	560.3	1,452.7
2000	52.5	7.5	*	*	0.7	2.9	14.5	43.4	93.8	222.9	588.8	1,530.0
1999	52.2	4.6	*	*	0.7	2.7	14.3	41.5	96.3	231.9	590.5	1,492.5
Black, male												
2003	43.2	5.9	*	*	0.8	3.3	14.3	46.9	112.1	237.4	588.9	1,180.3
2002	43.5	5.8	*	*	0.7	2.9	12.2	46.5	110.3	262.9	587.8	1,252.2
2001	44.6	8.1	*	*	0.7	3.1	13.2	48.8	111.9	269.2	613.9	1,349.1
2000	46.1	8.6	*	*	*	2.7	14.7	49.5	115.4	268.5	659.2	1,458.8
1999	45.9	*	*	*	*	2.8	14.7	48.6	117.9	274.7	650.6	1,448.3
Black, female												
2003	54.8	*	*	*	1.1	2.8	13.3	36.0	71.8	175.3	498.3	1,414.2
2002	55.8	*	*	*	*	3.3	13.4	35.7	70.1	181.2	532.2	1,434.3
2001	56.9	*	*	*	1.0	2.8	13.1	37.3	74.4	189.5	530.3	1,491.2
2000	58.3	*	*	*	0.8	3.1	14.4	38.1	76.4	190.9	549.2	1,556.5
1999	57.9	*	*	*	0.8	2.7	13.9	35.3	78.9	201.7	556.8	1,509.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
All races 3/, both sexes												
2003	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
2002	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0
All races, male												
2003	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
2002	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0
All races, female												
2003	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
2002	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
White, both sexes												
2003	5.0	*	*	*	*	*	0.1	0.4	1.6	6.5	30.3	159.5
2002	5.4	*	*	*	*	*	0.1	0.4	1.7	7.1	32.1	174.7
2001	5.6	*	*	*	*	*	0.1	0.5	1.8	7.3	33.8	185.9
2000	5.7	*	*	*	*	*	0.1	0.4	1.8	7.7	35.1	195.2
1999	6.0	*	*	*	*	*	0.1	0.5	2.0	8.0	36.4	207.7
White, male												
2003	3.7	*	*	*	*	*	0.2	0.6	2.1	7.5	33.7	143.3
2002	4.0	*	*	*	*	*	0.1	0.5	2.3	8.8	35.6	163.2
2001	4.1	*	*	*	*	*	0.1	0.6	2.3	9.0	36.4	173.9
2000	4.3	*	*	*	*	*	0.2	0.5	2.5	9.7	40.3	181.4
1999	4.4	*	*	*	*	*	0.1	0.7	2.6	9.5	40.2	193.7
White, female												
2003	6.3	*	*	*	*	*	*	0.3	1.1	5.6	28.0	166.7
2002	6.7	*	*	*	*	*	*	0.3	1.3	5.7	29.8	179.7
2001	7.0	*	*	*	*	*	*	0.3	1.3	5.9	32.1	190.9
2000	7.1	*	*	*	*	*	*	0.2	1.1	6.1	31.7	200.8
1999	7.6	*	*	*	*	*	*	0.3	1.4	6.8	34.0	213.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Atherosclerosis (I70)												
Black, both sexes												
2003	2.6	*	*	*	*	*	*	0.8	2.6	9.2	30.7	121.1
2002	2.7	*	*	*	*	*	*	1.0	2.9	8.0	36.5	117.0
2001	2.6	*	*	*	*	*	*	0.7	3.3	8.9	30.7	125.6
2000	2.9	*	*	*	*	*	*	1.1	3.2	8.9	35.2	138.3
1999	3.0	*	*	*	*	*	*	0.7	3.2	9.2	34.9	154.3
Black, male												
2003	2.3	*	*	*	*	*	*	1.2	3.6	12.4	40.3	104.3
2002	2.3	*	*	*	*	*	*	1.4	4.2	10.1	44.2	110.4
2001	2.1	*	*	*	*	*	*	1.0	3.8	10.9	37.1	113.5
2000	2.4	*	*	*	*	*	*	1.4	3.9	11.0	40.3	147.6
1999	2.5	*	*	*	*	*	*	*	4.4	11.8	44.0	164.0
Black, female												
2003	2.9	*	*	*	*	*	*	*	1.8	7.0	25.2	127.6
2002	3.1	*	*	*	*	*	*	*	1.9	6.5	32.2	119.5
2001	3.1	*	*	*	*	*	*	*	2.9	7.5	27.1	130.1
2000	3.3	*	*	*	*	*	*	*	2.7	7.5	32.4	134.8
1999	3.4	*	*	*	*	*	*	*	2.3	7.4	29.8	150.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
All races 3/, both sexes												
2003	8.5	*	*	*	0.2	0.4	1.0	2.9	8.1	26.9	67.6	146.0
2002	8.6	*	*	*	0.2	0.4	1.0	2.7	8.6	29.1	68.7	148.4
2001	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2
All races, male												
2003	9.2	*	*	*	0.2	0.5	1.4	4.1	11.3	37.0	91.8	179.7
2002	9.4	*	*	*	0.2	0.5	1.3	3.7	11.9	39.8	92.3	187.4
2001	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5
All races, female												
2003	7.8	*	*	*	0.1	0.2	0.7	1.8	5.2	18.4	51.4	131.1
2002	7.9	*	*	*	0.1	0.3	0.7	1.7	5.5	20.1	53.0	131.5
2001	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
White, both sexes												
2003	9.2	*	*	*	0.2	0.3	0.9	2.5	7.9	27.3	68.3	146.9
2002	9.3	*	*	*	0.2	0.3	0.9	2.3	8.2	29.4	69.7	148.9
2001	9.6	*	*	*	0.1	0.4	0.9	2.3	8.3	29.5	72.1	158.2
2000	10.0	*	*	*	0.1	0.3	0.8	2.4	8.8	32.7	75.1	163.0
1999	10.0	*	*	*	0.1	0.3	1.0	2.4	9.1	32.1	76.4	165.1
White, male												
2003	10.0	*	*	*	0.2	0.4	1.2	3.6	11.0	37.9	93.1	184.1
2002	10.2	*	*	*	0.2	0.4	1.2	3.2	11.4	40.6	94.6	190.9
2001	10.4	*	*	*	0.2	0.6	1.2	3.1	11.6	41.1	98.3	201.3
2000	11.1	*	*	*	0.2	0.5	1.0	3.4	12.8	46.7	104.7	219.6
1999	11.0	*	*	*	0.2	0.5	1.2	3.3	13.4	45.8	106.2	211.1
White, female												
2003	8.3	*	*	*	*	0.2	0.5	1.5	4.9	18.3	51.5	130.5
2002	8.5	*	*	*	*	0.3	0.6	1.4	5.2	19.9	52.9	130.7
2001	8.8	*	*	*	*	0.3	0.6	1.5	5.2	19.7	54.7	140.1
2000	8.9	*	*	*	*	0.2	0.6	1.4	5.1	21.0	55.6	140.1
1999	9.1	*	*	*	*	0.2	0.7	1.5	5.2	20.6	56.9	146.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of circulatory system (I71-I78)												
Black, both sexes												
2003	6.6	*	*	*	*	0.7	2.3	6.2	11.8	29.3	70.1	150.0
2002	6.7	*	*	*	*	0.9	2.0	5.9	13.5	31.1	68.3	156.8
2001	7.1	*	*	*	*	0.7	2.4	5.1	13.9	31.7	79.8	174.7
2000	7.1	*	*	*	*	0.7	2.2	5.5	14.4	33.0	78.0	177.7
1999	6.7	*	*	*	*	0.5	2.0	5.2	12.3	31.7	76.6	178.9
Black, male												
2003	6.7	*	*	*	*	1.1	3.0	8.2	16.4	36.3	90.3	138.7
2002	6.7	*	*	*	*	1.2	2.6	8.1	19.3	38.4	78.6	166.2
2001	6.9	*	*	*	*	0.9	3.4	7.3	17.8	40.0	96.8	165.7
2000	6.9	*	*	*	*	0.9	2.8	7.5	18.4	41.1	95.7	177.6
1999	6.5	*	*	*	*	0.9	2.3	7.2	15.7	39.4	89.8	184.0
Black, female												
2003	6.5	*	*	*	*	*	1.7	4.5	8.2	24.2	58.7	154.3
2002	6.6	*	*	*	*	*	1.4	4.0	8.9	25.8	62.5	153.2
2001	7.2	*	*	*	*	*	1.6	3.3	10.8	26.0	70.2	178.1
2000	7.2	*	*	*	*	*	1.5	3.9	11.1	27.3	68.0	177.7
1999	7.0	*	*	*	*	*	1.7	3.5	9.6	26.2	69.2	177.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
All races 3/, both sexes												
2003	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
2002	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3
All races, male												
2003	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
2002	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3
All races, female												
2003	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
2002	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
White, both sexes												
2003	5.6	*	*	*	0.1	0.2	0.6	1.7	5.2	18.6	44.0	72.4
2002	5.7	*	*	*	0.1	0.2	0.6	1.5	5.4	19.9	45.0	71.2
2001	5.9	*	*	*	0.1	0.3	0.6	1.5	5.4	19.9	46.9	79.7
2000	6.2	*	*	*	0.1	0.2	0.6	1.6	6.0	22.7	48.5	80.8
1999	6.3	*	*	*	0.1	0.2	0.6	1.5	6.1	22.6	50.1	82.2
White, male												
2003	6.9	*	*	*	0.1	0.3	1.0	2.7	8.2	27.3	64.3	110.0
2002	7.0	*	*	*	0.1	0.3	0.9	2.3	8.4	29.8	65.5	111.8
2001	7.2	*	*	*	0.1	0.5	1.0	2.3	8.4	29.9	68.5	120.3
2000	7.8	*	*	*	*	0.3	0.8	2.5	9.5	34.5	73.0	132.8
1999	7.8	*	*	*	*	0.4	0.9	2.4	9.8	34.3	75.8	129.8
White, female												
2003	4.3	*	*	*	*	0.2	0.2	0.8	2.4	11.1	30.3	55.7
2002	4.3	*	*	*	*	*	0.3	0.7	2.6	11.4	31.2	53.5
2001	4.6	*	*	*	*	*	0.2	0.6	2.6	11.5	32.5	62.6
2000	4.6	*	*	*	*	0.1	0.3	0.7	2.8	12.8	32.3	59.6
1999	4.8	*	*	*	*	*	0.4	0.7	2.7	12.8	33.3	63.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Aortic aneurysm and dissection (I71)												
Black, both sexes												
2003	3.2	*	*	*	*	0.5	1.6	4.2	7.1	14.5	30.0	46.4
2002	3.2	*	*	*	*	0.6	1.4	4.2	7.5	14.6	29.9	50.5
2001	3.4	*	*	*	*	0.4	1.6	3.5	7.6	14.6	37.6	55.3
2000	3.5	*	*	*	*	0.5	1.6	3.2	8.2	17.0	36.3	62.3
1999	3.3	*	*	*	*	0.4	1.4	3.7	6.5	16.7	33.8	60.3
Black, male												
2003	3.8	*	*	*	*	1.0	2.4	6.4	10.8	18.0	39.2	51.1
2002	3.8	*	*	*	*	1.0	2.2	6.6	12.1	19.0	35.0	49.3
2001	3.8	*	*	*	*	*	2.6	5.3	11.2	20.0	47.4	48.8
2000	3.9	*	*	*	*	*	2.4	4.7	11.5	22.1	46.9	69.2
1999	3.6	*	*	*	*	*	1.9	5.8	9.1	21.9	38.4	76.2
Black, female												
2003	2.7	*	*	*	*	*	1.0	2.4	4.2	11.9	24.7	44.6
2002	2.7	*	*	*	*	*	*	2.2	3.8	11.5	27.0	50.9
2001	3.0	*	*	*	*	*	0.7	1.9	4.6	10.8	32.1	57.7
2000	3.2	*	*	*	*	*	0.9	2.0	5.5	13.4	30.3	59.7
1999	3.0	*	*	*	*	*	0.9	2.0	4.4	13.0	31.2	54.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
All races 3/, both sexes												
2003	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
2002	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9
All races, male												
2003	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
2002	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2
All races, female												
2003	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
2002	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
White, both sexes												
2003	3.6	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.2	74.6
2002	3.7	*	*	*	*	0.1	0.3	0.8	2.8	9.6	24.7	77.8
2001	3.7	*	*	*	*	0.1	0.3	0.9	2.9	9.5	25.2	78.5
2000	3.8	*	*	*	*	0.1	0.3	0.8	2.8	10.0	26.6	82.3
1999	3.8	*	*	*	*	0.1	0.3	0.9	3.0	9.5	26.3	82.9
White, male												
2003	3.1	*	*	*	*	*	0.2	0.9	2.9	10.6	28.8	74.1
2002	3.2	*	*	*	*	*	0.3	0.9	3.1	10.8	29.1	79.1
2001	3.2	*	*	*	*	*	0.2	0.8	3.2	11.2	29.8	80.9
2000	3.4	*	*	*	*	0.1	0.2	0.9	3.3	12.2	31.6	86.9
1999	3.2	*	*	*	*	*	0.3	0.9	3.6	11.4	30.4	81.3
White, female												
2003	4.0	*	*	*	*	*	0.3	0.7	2.5	7.2	21.2	74.8
2002	4.2	*	*	*	*	0.1	0.3	0.7	2.6	8.5	21.7	77.2
2001	4.2	*	*	*	*	0.1	0.4	0.9	2.6	8.2	22.1	77.5
2000	4.2	*	*	*	*	*	0.3	0.7	2.3	8.2	23.3	80.4
1999	4.3	*	*	*	*	*	0.3	0.8	2.5	7.9	23.6	83.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Black, both sexes												
2003	3.4	*	*	*	*	*	0.7	2.0	4.7	14.8	40.1	103.6
2002	3.5	*	*	*	*	*	0.6	1.7	6.1	16.5	38.5	106.3
2001	3.7	*	*	*	*	*	0.9	1.7	6.4	17.1	42.2	119.4
2000	3.6	*	*	*	*	*	0.6	2.3	6.2	16.0	41.7	115.4
1999	3.5	*	*	*	*	*	0.6	1.5	5.8	15.0	42.8	118.6
Black, male												
2003	2.9	*	*	*	*	*	*	1.8	5.6	18.3	51.1	87.6
2002	3.0	*	*	*	*	*	*	1.5	7.2	19.4	43.6	116.9
2001	3.2	*	*	*	*	*	0.9	2.0	6.5	20.0	49.4	116.9
2000	3.1	*	*	*	*	*	*	2.8	7.0	19.0	48.7	108.4
1999	2.9	*	*	*	*	*	*	1.4	6.6	17.5	51.3	107.8
Black, female												
2003	3.8	*	*	*	*	*	0.7	2.1	4.0	12.3	34.0	109.8
2002	3.9	*	*	*	*	*	0.8	1.9	5.1	14.4	35.6	102.3
2001	4.2	*	*	*	*	*	0.8	1.4	6.2	15.1	38.2	120.4
2000	4.1	*	*	*	*	*	0.7	1.8	5.5	13.9	37.7	118.0
1999	4.0	*	*	*	*	*	0.8	1.6	5.2	13.3	38.0	122.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
All races 3/, both sexes												
2003	1.6	0.9	*	*	0.1	0.3	0.8	1.4	2.0	4.1	9.0	24.0
2002	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4
All races, male												
2003	1.4	1.0	*	*	0.1	0.3	0.8	1.6	2.4	4.2	8.5	19.1
2002	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8
All races, female												
2003	1.9	*	*	*	0.1	0.4	0.8	1.2	1.7	4.0	9.4	26.2
2002	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
White, both sexes												
2003	1.6	0.9	*	*	0.1	0.3	0.6	1.2	1.9	3.8	8.6	23.6
2002	1.6	0.6	*	*	0.1	0.3	0.6	1.2	2.0	3.6	9.5	24.3
2001	1.7	*	*	*	0.1	0.3	0.5	1.0	2.0	4.3	9.1	26.0
2000	1.6	0.7	*	*	0.1	0.3	0.6	1.2	1.9	4.0	9.1	25.5
1999	1.5	*	*	*	*	0.2	0.5	0.9	1.8	3.5	8.9	23.2
White, male												
2003	1.3	*	*	*	*	0.2	0.7	1.4	2.2	3.9	7.9	18.5
2002	1.3	*	*	*	*	0.3	0.6	1.3	2.2	3.6	9.0	19.9
2001	1.4	*	*	*	*	0.2	0.5	1.3	2.2	4.6	9.5	21.7
2000	1.4	*	*	*	*	0.2	0.8	1.5	2.0	4.4	9.5	22.0
1999	1.2	*	*	*	*	0.2	0.6	0.9	1.9	3.9	9.2	20.7
White, female												
2003	1.9	*	*	*	*	0.3	0.5	1.1	1.5	3.8	9.1	25.8
2002	1.9	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.8	26.3
2001	1.9	*	*	*	*	0.3	0.5	0.8	1.8	4.1	8.9	27.9
2000	1.9	*	*	*	*	0.3	0.5	1.0	1.8	3.6	8.9	26.9
1999	1.7	*	*	*	*	0.2	0.4	0.9	1.7	3.2	8.7	24.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of circulatory system (I80-I99)												
Black, both sexes												
2003	2.2	*	*	*	*	0.9	1.9	2.6	4.3	7.5	15.4	33.6
2002	2.1	*	*	*	*	1.0	1.6	2.6	4.9	6.8	16.6	30.0
2001	2.0	*	*	*	*	0.8	1.5	2.2	4.9	7.8	14.4	30.1
2000	2.1	*	*	*	*	0.9	1.5	2.8	4.5	8.9	14.4	30.0
1999	1.7	*	*	*	*	0.5	1.4	2.4	2.9	6.6	13.8	28.4
Black, male												
2003	2.0	*	*	*	*	0.9	1.3	3.4	5.2	8.7	17.5	32.3
2002	2.0	*	*	*	*	1.0	1.8	3.0	5.6	7.7	18.4	32.2
2001	1.9	*	*	*	*	*	1.7	2.4	6.4	9.0	15.9	29.5
2000	1.9	*	*	*	*	0.9	1.8	3.4	5.1	9.1	13.3	30.0
1999	1.6	*	*	*	*	*	1.8	2.8	2.6	8.2	16.9	*
Black, female												
2003	2.3	*	*	*	*	0.9	2.5	1.9	3.6	6.7	14.2	34.0
2002	2.2	*	*	*	*	0.9	1.5	2.3	4.4	6.1	15.6	29.2
2001	2.1	*	*	*	*	0.9	1.4	2.1	3.6	7.1	13.6	30.3
2000	2.2	*	*	*	*	0.9	1.3	2.3	3.9	8.8	15.0	30.0
1999	1.8	*	*	*	*	*	1.1	2.0	3.1	5.5	12.0	30.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
All races 3/, both sexes												
2003	22.4	8.0	1.0	0.4	0.5	0.9	2.2	5.2	11.2	37.3	151.1	666.1
2002	22.8	6.5	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.5	156.9	696.6
2001	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8
All races, male												
2003	20.1	9.4	1.1	0.3	0.6	1.1	2.5	6.4	13.2	46.5	186.8	757.8
2002	20.4	7.8	0.8	0.2	0.5	0.9	2.4	5.8	13.6	46.4	194.7	805.8
2001	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7
All races, female												
2003	24.6	6.6	0.9	0.4	0.5	0.7	1.9	4.1	9.4	29.5	127.2	625.6
2002	25.1	5.1	0.6	0.3	0.4	0.8	1.9	3.8	9.0	30.1	131.8	649.2
2001	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
White, both sexes												
2003	24.4	6.7	0.9	0.3	0.5	0.8	2.0	4.7	10.3	35.9	151.5	679.3
2002	24.9	5.2	0.5	0.2	0.4	0.8	1.9	4.4	10.4	36.7	156.9	709.1
2001	23.6	5.3	0.7	0.2	0.4	0.7	1.8	4.0	9.9	35.3	147.2	695.0
2000	25.2	5.8	0.6	0.2	0.4	0.8	2.1	4.1	10.9	37.1	160.0	758.3
1999	24.8	6.2	0.6	0.2	0.4	0.7	1.9	4.0	10.1	36.1	157.8	764.3
White, male												
2003	21.4	8.0	1.1	0.3	0.6	1.0	2.2	5.6	11.6	44.0	184.9	767.1
2002	21.9	6.6	0.6	0.2	0.4	0.8	2.1	5.2	12.3	45.2	193.1	815.9
2001	20.7	6.2	0.7	0.2	0.5	0.8	2.1	4.9	11.8	43.0	182.6	814.4
2000	22.0	6.4	0.5	0.2	0.5	0.9	2.5	4.8	12.9	45.1	197.8	906.1
1999	21.5	6.7	0.5	0.3	0.4	0.8	2.1	4.9	11.5	44.7	196.1	905.7
White, female												
2003	27.3	5.4	0.8	0.4	0.5	0.6	1.8	3.7	9.0	29.0	129.0	640.4
2002	27.8	3.7	0.5	0.3	0.3	0.7	1.7	3.6	8.5	29.6	132.7	662.6
2001	26.4	4.4	0.6	0.2	0.3	0.7	1.6	3.2	8.1	28.9	123.8	645.0
2000	28.2	5.1	0.7	0.2	0.4	0.8	1.7	3.4	8.9	30.3	135.0	698.3
1999	27.9	5.6	0.7	0.2	0.4	0.6	1.8	3.0	8.7	28.9	132.6	707.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza and pneumonia (J10-J18)												
Black, both sexes												
2003	15.4	14.1	1.1	0.4	0.7	1.9	4.4	10.4	20.6	54.2	155.9	525.4
2002	15.6	12.9	1.6	*	0.6	1.5	4.3	8.7	21.2	50.3	166.2	575.2
2001	15.5	16.3	1.0	*	0.7	1.6	4.8	9.7	18.9	48.3	169.0	581.3
2000	16.4	16.7	1.2	*	0.7	1.7	4.5	10.0	22.0	60.3	171.1	600.9
1999	16.2	19.0	1.7	*	0.8	2.0	5.5	10.6	20.9	54.1	157.4	634.8
Black, male												
2003	16.0	16.1	*	*	0.8	2.3	5.3	13.2	29.1	78.2	218.6	641.2
2002	15.4	13.9	1.7	*	0.6	1.7	5.1	11.2	29.0	65.9	228.9	689.4
2001	15.9	15.9	*	*	0.8	2.1	5.7	13.3	26.3	69.4	242.2	714.8
2000	16.7	17.9	1.7	*	0.7	2.3	5.6	13.9	30.0	83.5	234.1	755.4
1999	16.4	18.9	2.0	*	0.8	2.2	6.6	15.1	27.2	76.9	213.7	803.8
Black, female												
2003	14.9	12.1	*	*	*	1.5	3.6	7.9	13.8	37.2	120.4	480.4
2002	15.7	11.8	*	*	*	1.3	3.7	6.5	15.0	39.1	130.7	531.5
2001	15.2	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.9	531.7
2000	16.0	15.5	*	*	0.8	1.2	3.4	6.7	15.6	43.9	135.7	543.4
1999	16.1	19.1	*	*	0.9	1.7	4.4	6.8	15.8	38.0	125.8	571.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
All races 3/, both sexes												
2003	0.6	0.8	0.4	0.1	0.1	*	0.1	0.1	0.3	1.0	3.6	17.0
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2
All races, male												
2003	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.0
2002	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9
All races, female												
2003	0.7	*	0.3	0.1	*	*	*	0.1	0.3	0.9	3.5	17.1
2002	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
White, both sexes												
2003	0.7	0.7	0.3	0.1	0.1	*	0.1	0.1	0.3	1.1	3.8	18.2
2002	0.3	*	*	*	*	*	*	*	0.1	0.3	1.5	9.4
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000	0.7	*	*	*	*	*	*	0.1	0.4	1.1	4.2	21.4
1999	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.9	22.8
White, male												
2003	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.9	18.3
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.7	9.7
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
2000	0.5	*	*	*	*	*	*	*	0.4	1.3	4.2	21.6
1999	0.5	*	*	*	*	*	*	*	0.4	0.9	4.4	21.7
White, female												
2003	0.8	*	*	*	*	*	*	0.1	0.3	1.0	3.7	18.1
2002	0.4	*	*	*	*	*	*	*	*	*	1.4	9.3
2001	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000	0.9	*	*	*	*	*	*	0.1	0.4	0.9	4.2	21.3
1999	0.9	*	*	*	*	*	*	*	0.3	0.9	3.5	23.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Influenza (J10-J11)												
Black, both sexes												
2003	0.3	*	*	*	*	*	*	*	*	*	*	5.8
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	9.4
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
All races 3/, both sexes												
2003	21.8	7.3	0.7	0.2	0.5	0.9	2.2	5.1	10.9	36.3	147.5	649.1
2002	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6
All races, male												
2003	19.6	8.5	0.7	0.2	0.5	1.1	2.5	6.2	12.9	45.4	183.2	740.9
2002	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7
All races, female												
2003	23.9	6.0	0.6	0.3	0.4	0.7	1.9	4.0	9.1	28.6	123.7	608.5
2002	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
White, both sexes												
2003	23.7	6.0	0.6	0.2	0.5	0.8	1.9	4.5	10.0	34.8	147.7	661.1
2002	24.6	5.0	0.5	0.2	0.4	0.8	1.9	4.3	10.2	36.4	155.4	699.6
2001	23.5	5.1	0.6	0.2	0.4	0.7	1.8	4.0	9.8	35.2	146.8	692.7
2000	24.4	5.6	0.5	0.2	0.4	0.8	2.0	4.0	10.4	36.0	155.8	737.0
1999	24.1	5.9	0.6	0.2	0.4	0.7	1.9	3.9	9.8	35.2	153.9	741.6
White, male												
2003	20.9	7.1	0.7	0.2	0.5	1.0	2.2	5.5	11.3	42.8	181.0	748.8
2002	21.7	6.4	0.5	0.1	0.4	0.8	2.1	5.1	12.1	44.8	191.4	806.2
2001	20.6	5.8	0.6	0.2	0.5	0.8	2.1	4.8	11.7	42.9	182.2	812.1
2000	21.5	6.1	0.4	0.2	0.5	0.8	2.4	4.7	12.5	43.8	193.6	884.4
1999	21.0	6.7	0.4	0.2	0.4	0.8	2.1	4.8	11.1	43.8	191.7	884.0
White, female												
2003	26.5	4.8	0.5	0.2	0.4	0.6	1.7	3.6	8.7	28.0	125.2	622.3
2002	27.4	3.6	0.5	0.2	0.3	0.7	1.7	3.5	8.5	29.3	131.3	653.3
2001	26.2	4.4	0.6	0.2	0.3	0.7	1.6	3.1	8.0	28.7	123.3	642.8
2000	27.3	5.0	0.6	0.2	0.3	0.7	1.7	3.2	8.5	29.4	130.8	677.1
1999	27.1	5.2	0.7	0.2	0.3	0.5	1.7	3.0	8.5	27.9	129.1	684.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonia (J12-J18)												
Black, both sexes												
2003	15.1	13.0	0.9	*	0.6	1.8	4.4	10.1	20.3	53.7	154.5	519.6
2002	15.5	12.7	1.5	*	0.6	1.5	4.3	8.7	21.1	49.9	165.8	572.8
2001	15.5	16.3	1.0	*	0.7	1.6	4.7	9.7	18.8	48.2	168.6	579.7
2000	16.1	16.4	1.2	*	0.7	1.7	4.4	9.9	21.8	59.3	169.7	591.5
1999	16.1	18.3	1.7	*	0.8	1.9	5.4	10.5	20.7	53.3	155.7	629.1
Black, male												
2003	15.8	15.2	*	*	0.7	2.2	5.2	12.9	28.7	77.7	217.5	638.1
2002	15.3	13.7	1.6	*	*	1.7	5.1	11.2	28.7	65.5	228.4	687.2
2001	15.8	15.9	*	*	0.7	2.0	5.7	13.3	26.1	69.1	242.2	712.6
2000	16.5	17.9	1.7	*	0.7	2.2	5.5	13.6	29.7	82.8	232.5	742.7
1999	16.3	17.9	1.9	*	0.8	2.2	6.6	15.0	27.1	76.0	211.8	800.3
Black, female												
2003	14.6	10.8	*	*	*	1.4	3.6	7.7	13.6	36.6	119.0	473.5
2002	15.6	11.8	*	*	*	1.3	3.7	6.5	14.9	38.8	130.4	529.0
2001	15.1	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.2	530.5
2000	15.8	14.9	*	*	0.8	1.2	3.4	6.6	15.4	42.7	134.3	535.3
1999	15.9	18.8	*	*	0.9	1.7	4.3	6.8	15.5	37.2	124.3	565.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
All races 3/, both sexes												
2003	0.1	1.3	0.2	*	*	*	*	*	0.1	0.2	0.6	3.5
2002	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0
All races, male												
2003	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.4
2002	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0
All races, female												
2003	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
2002	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
White, both sexes												
2003	0.2	1.3	*	*	*	*	*	*	*	0.2	0.7	3.7
2002	0.1	1.1	0.2	*	*	*	*	*	*	0.2	0.5	3.5
2001	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000	0.2	0.8	*	*	*	*	*	*	*	0.3	0.7	4.7
1999	0.2	1.2	*	*	*	*	*	*	*	0.2	0.9	6.3
White, male												
2003	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.7
2002	0.1	1.6	*	*	*	*	*	*	*	*	0.5	2.9
2001	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.8
1999	0.1	*	*	*	*	*	*	*	*	*	1.1	4.0
White, female												
2003	0.2	*	*	*	*	*	*	*	*	*	0.6	4.2
2002	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000	0.2	*	*	*	*	*	*	*	*	0.2	0.7	5.1
1999	0.3	*	*	*	*	*	*	*	*	*	0.8	7.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other acute lower respiratory infections (J20-J22)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.0	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
All races 3/, both sexes												
2003	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2002	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6
All races, male												
2003	0.1	1.7	*	*	*	*	*	*	*	*	*	*
2002	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6
All races, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2002	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
White, both sexes												
2003	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2002	0.1	1.0	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2001	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.1
2000	0.1	0.8	*	*	*	*	*	*	*	0.2	0.4	2.7
1999	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.8
White, male												
2003	0.1	1.6	*	*	*	*	*	*	*	*	*	*
2002	0.1	1.5	*	*	*	*	*	*	*	*	*	1.7
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.6	2.4
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	2.6
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2002	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.9
1999	0.2	*	*	*	*	*	*	*	*	*	0.5	4.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	4.0	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
All races 3/, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	1.3
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4
All races, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
White, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	0.3	1.4
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5
White, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2002	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999	0.1	*	*	*	*	*	*	*	*	*	0.3	2.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Unspecified acute lower respiratory infection (J22)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
All races 3/, both sexes												
2003	43.5	0.8	0.3	0.3	0.5	0.7	2.1	8.7	43.3	163.2	383.0	635.1
2002	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0
All races, male												
2003	42.4	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.5	183.6	464.9	865.9
2002	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7
All races, female												
2003	44.4	*	*	0.2	0.4	0.6	2.3	8.2	40.3	146.0	328.3	533.0
2002	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
White, both sexes												
2003	49.5	0.7	0.2	0.2	0.3	0.5	1.9	8.5	45.2	173.8	403.0	664.5
2002	49.2	0.8	0.2	0.2	0.3	0.6	1.9	8.4	44.2	173.2	405.6	666.2
2001	49.0	0.7	0.2	0.2	0.3	0.5	1.8	8.3	46.0	177.7	398.6	671.8
2000	49.0	0.8	0.2	0.2	0.3	0.5	1.7	8.2	45.8	178.7	404.4	675.6
1999	50.2	*	0.3	0.2	0.4	0.5	1.6	8.1	49.3	186.2	414.3	672.9
White, male												
2003	47.4	1.3	*	0.2	0.4	0.5	1.7	8.9	47.6	191.6	478.5	894.4
2002	47.8	*	0.3	0.2	0.4	0.6	1.8	8.8	46.0	192.3	495.2	923.4
2001	47.6	*	*	0.2	0.3	0.5	1.7	8.6	48.0	198.3	489.4	943.6
2000	48.3	*	*	0.2	0.4	0.5	1.6	8.4	48.6	201.4	503.6	997.4
1999	50.6	*	*	0.3	0.4	0.5	1.5	8.1	53.1	217.3	525.4	1,029.4
White, female												
2003	51.5	*	*	0.1	0.3	0.6	2.1	8.1	42.9	158.6	352.0	562.8
2002	50.5	*	*	0.2	0.3	0.6	2.0	8.1	42.4	157.0	345.4	554.5
2001	50.3	*	*	0.1	0.3	0.5	1.9	8.0	44.1	160.4	338.3	557.9
2000	49.7	*	*	0.2	0.3	0.5	1.8	7.9	43.2	159.6	339.1	544.8
1999	49.8	*	*	0.2	0.4	0.6	1.8	8.1	45.6	160.3	341.5	529.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)												
Black, both sexes												
2003	20.2	*	1.0	0.9	1.3	1.9	4.3	12.5	40.5	113.2	235.9	353.0
2002	20.7	*	1.2	1.1	1.3	1.9	5.1	12.9	40.7	113.5	248.7	368.2
2001	20.4	*	0.9	0.8	1.3	2.4	4.8	12.5	40.7	118.9	234.1	375.8
2000	20.8	*	0.8	1.0	1.4	2.1	4.8	14.2	43.9	121.7	234.7	373.8
1999	21.9	*	*	1.0	1.3	2.6	4.6	13.8	45.8	128.5	262.6	376.2
Black, male												
2003	23.3	*	1.6	1.2	1.5	2.8	4.0	13.3	50.5	155.1	382.2	601.6
2002	24.1	*	1.6	1.2	1.9	2.2	5.7	14.4	52.3	158.0	392.2	645.4
2001	23.6	*	*	0.9	1.6	2.7	4.7	13.3	49.8	168.0	380.8	671.7
2000	24.3	*	*	1.2	1.7	2.8	4.8	15.0	54.6	176.9	370.3	693.1
1999	26.2	*	*	1.4	1.8	2.7	4.7	15.3	59.3	184.6	434.4	701.9
Black, female												
2003	17.3	*	*	0.6	1.1	1.1	4.6	11.9	32.4	83.3	153.2	256.4
2002	17.7	*	*	1.0	0.6	1.6	4.6	11.6	31.5	82.0	167.4	262.0
2001	17.5	*	*	0.7	1.0	2.1	4.9	11.7	33.3	84.3	151.7	266.1
2000	17.6	*	*	0.8	1.0	1.5	4.7	13.4	35.3	82.9	158.4	255.0
1999	18.0	*	*	0.6	0.9	2.6	4.6	12.6	34.9	88.9	166.4	254.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
All races 3/, both sexes												
2003	0.3	0.5	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002	0.3	0.6	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1
2001	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7
All races, male												
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.6	2.2	7.0
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9
All races, female												
2003	0.3	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2001	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
White, both sexes												
2003	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.5
2002	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.3	8.6
2001	0.4	*	*	*	*	*	*	0.1	0.3	0.7	2.1	9.6
2000	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.7
1999	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	11.2
White, male												
2003	0.3	*	*	*	*	*	*	*	0.2	0.5	2.2	7.3
2002	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.6	9.3
2001	0.3	*	*	*	*	*	*	*	0.3	0.9	2.3	9.4
2000	0.4	*	*	*	*	*	*	*	0.3	1.0	3.7	13.8
1999	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4
White, female												
2003	0.4	*	*	*	*	*	*	*	*	0.6	1.8	7.6
2002	0.4	*	*	*	*	*	*	*	0.2	0.7	2.2	8.3
2001	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.7
2000	0.5	*	*	*	*	*	*	*	0.3	0.8	2.2	12.2
1999	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Bronchitis, chronic and unspecified (J40-J42)												
Black, both sexes												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	2.2	*
1999	0.2	*	*	*	*	*	*	*	*	*	2.9	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
All races 3/, both sexes												
2003	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
2002	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1
All races, male												
2003	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
2002	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4
All races, female												
2003	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
2002	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
White, both sexes												
2003	5.9	*	*	*	*	*	0.2	1.1	6.7	23.9	48.5	57.0
2002	6.2	*	*	*	*	*	0.2	1.2	6.9	25.7	50.7	62.3
2001	6.6	*	*	*	*	*	0.2	1.2	7.7	28.2	53.5	65.1
2000	6.8	*	*	*	*	*	0.2	1.2	7.9	29.7	55.6	65.8
1999	7.3	*	*	*	*	*	0.2	1.3	9.1	32.9	59.8	64.7
White, male												
2003	5.9	*	*	*	*	*	0.2	1.4	7.2	27.1	59.8	80.4
2002	6.4	*	*	*	*	*	0.3	1.3	7.5	29.3	65.8	93.0
2001	6.8	*	*	*	*	*	0.2	1.4	8.2	32.3	68.1	105.1
2000	7.0	*	*	*	*	*	0.2	1.4	8.4	34.3	72.1	105.5
1999	7.8	*	*	*	*	*	0.2	1.4	9.9	39.4	79.0	109.2
White, female												
2003	5.8	*	*	*	*	*	0.2	0.9	6.2	21.2	40.8	46.6
2002	5.9	*	*	*	*	*	*	1.0	6.2	22.7	40.5	48.9
2001	6.3	*	*	*	*	*	0.1	0.9	7.2	24.7	43.7	48.4
2000	6.5	*	*	*	*	*	0.2	1.0	7.4	25.9	44.7	49.7
1999	6.8	*	*	*	*	*	0.2	1.1	8.3	27.4	47.2	46.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Emphysema (J43)												
Black, both sexes												
2003	2.1	*	*	*	*	*	*	1.2	5.1	12.9	27.3	30.9
2002	2.3	*	*	*	*	*	0.4	1.2	4.4	15.9	28.0	32.1
2001	2.2	*	*	*	*	*	*	1.3	5.6	15.3	23.6	35.3
2000	2.5	*	*	*	*	*	*	2.0	6.0	17.0	26.9	37.2
1999	2.6	*	*	*	*	*	0.4	1.8	6.1	17.1	31.3	37.6
Black, male												
2003	2.7	*	*	*	*	*	*	1.7	7.3	18.2	46.0	70.9
2002	3.0	*	*	*	*	*	*	1.8	6.5	24.8	48.8	64.3
2001	2.7	*	*	*	*	*	*	1.6	8.0	22.6	39.7	68.1
2000	3.2	*	*	*	*	*	*	2.9	8.1	27.4	41.5	87.6
1999	3.5	*	*	*	*	*	*	2.9	8.9	29.0	54.7	72.6
Black, female												
2003	1.6	*	*	*	*	*	*	*	3.4	9.2	16.7	15.4
2002	1.6	*	*	*	*	*	*	*	2.7	9.5	16.2	19.7
2001	1.7	*	*	*	*	*	*	1.1	3.6	10.2	14.5	23.2
2000	1.8	*	*	*	*	*	*	1.3	4.4	9.7	18.8	18.5
1999	1.7	*	*	*	*	*	*	1.0	3.9	8.7	18.2	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
All races 3/, both sexes												
2003	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
2002	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3
All races, male												
2003	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
2002	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7
All races, female												
2003	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
2002	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
White, both sexes												
2003	1.2	*	*	0.2	0.2	0.4	0.7	1.1	1.6	2.4	5.4	14.3
2002	1.3	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.7	6.1	14.7
2001	1.3	*	*	0.1	0.2	0.4	0.7	1.1	1.6	3.0	6.0	15.7
2000	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.7	3.4	6.5	15.9
1999	1.5	*	*	0.2	0.3	0.4	0.7	1.2	1.9	3.5	7.2	16.6
White, male												
2003	0.8	*	*	0.2	0.3	0.4	0.5	0.8	1.2	1.7	4.0	10.6
2002	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.2	1.9	4.6	10.7
2001	0.8	*	*	0.2	0.2	0.3	0.6	0.8	1.1	2.2	4.0	11.3
2000	0.9	*	*	0.2	0.3	0.4	0.6	1.0	1.2	2.5	4.7	11.4
1999	0.9	*	*	0.2	0.4	0.4	0.5	0.8	1.4	2.5	5.3	10.8
White, female												
2003	1.6	*	*	0.1	0.2	0.5	0.8	1.4	2.0	3.1	6.3	15.9
2002	1.7	*	*	*	0.2	0.5	0.9	1.3	1.9	3.3	7.2	16.4
2001	1.7	*	*	*	0.2	0.4	0.8	1.5	2.1	3.7	7.3	17.6
2000	1.8	*	*	0.2	0.2	0.4	0.8	1.3	2.1	4.1	7.7	17.7
1999	2.0	*	*	0.1	0.3	0.4	0.9	1.7	2.4	4.4	8.5	18.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Asthma (J45-J46)												
Black, both sexes												
2003	2.7	*	0.9	0.9	1.2	1.6	2.7	4.8	5.1	7.0	9.1	12.0
2002	2.9	*	0.9	1.0	1.2	1.6	3.4	4.8	6.4	7.8	8.0	11.9
2001	3.0	*	0.8	0.8	1.1	2.1	3.3	4.6	6.8	7.7	10.6	16.3
2000	3.2	*	*	1.0	1.3	1.9	3.1	5.4	6.9	9.2	10.4	22.2
1999	3.2	*	*	1.0	1.2	2.4	2.9	5.2	6.8	9.6	10.4	20.4
Black, male												
2003	2.4	*	1.5	1.2	1.4	2.4	2.3	3.9	4.2	5.3	8.7	*
2002	2.8	*	*	1.1	1.8	2.0	3.6	3.9	5.7	7.3	6.9	*
2001	2.6	*	*	0.9	1.4	2.4	3.0	4.1	4.6	6.8	8.8	*
2000	2.8	*	*	1.2	1.5	2.5	2.9	3.8	6.1	7.9	6.4	25.4
1999	2.8	*	*	1.4	1.6	2.6	2.8	3.8	5.4	9.1	8.3	*
Black, female												
2003	2.9	*	*	*	1.0	0.9	3.1	5.6	5.8	8.2	9.2	13.8
2002	3.0	*	*	1.0	*	1.2	3.3	5.6	7.0	8.2	8.6	12.3
2001	3.3	*	*	0.7	0.9	1.9	3.5	5.0	8.5	8.3	11.6	16.8
2000	3.5	*	*	0.8	1.0	1.4	3.3	6.8	7.5	10.2	12.6	21.0
1999	3.5	*	*	0.6	0.9	2.1	3.1	6.4	7.9	10.0	11.5	21.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
All races 3/, both sexes												
2003	36.6	*	*	*	0.1	0.1	1.0	5.9	34.8	137.5	329.2	558.9
2002	36.1	*	*	*	*	0.1	0.9	5.9	33.8	135.1	330.1	555.5
2001	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8
All races, male												
2003	35.9	*	*	*	*	0.1	1.0	6.4	37.8	155.4	400.1	769.0
2002	35.7	*	*	*	*	0.1	1.0	6.4	36.2	153.4	409.8	785.1
2001	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7
All races, female												
2003	37.4	*	*	*	*	*	1.0	5.5	32.1	122.6	281.9	466.0
2002	36.5	*	*	*	*	0.1	0.9	5.4	31.5	119.8	277.2	455.8
2001	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2
2000	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
White, both sexes												
2003	42.1	*	*	*	*	0.1	1.0	6.2	36.7	146.9	347.2	585.8
2002	41.3	*	*	*	*	0.1	1.0	6.1	35.6	144.0	346.4	580.7
2001	40.8	*	*	*	*	0.1	0.9	5.8	36.5	145.9	337.0	581.3
2000	40.4	*	*	*	*	*	0.8	5.7	36.0	144.7	339.5	581.2
1999	41.0	*	*	*	*	0.1	0.7	5.6	37.8	148.8	344.5	580.3
White, male												
2003	40.4	*	*	*	*	*	0.9	6.6	39.0	162.2	412.5	796.0
2002	40.2	*	*	*	*	*	1.0	6.5	37.1	160.2	422.1	810.4
2001	39.7	*	*	*	*	0.1	0.9	6.3	38.5	162.9	415.0	817.8
2000	39.9	*	*	*	*	*	0.8	6.0	38.7	163.6	423.0	866.7
1999	41.5	*	*	*	*	*	0.7	5.9	41.4	174.2	438.0	897.0
White, female												
2003	43.7	*	*	*	*	*	1.0	5.8	34.5	133.8	303.0	492.7
2002	42.5	*	*	*	*	*	0.9	5.7	34.1	130.3	295.6	480.9
2001	41.8	*	*	*	*	*	1.0	5.4	34.6	131.5	285.3	482.3
2000	40.9	*	*	*	*	*	0.8	5.5	33.4	128.9	284.4	465.2
1999	40.4	*	*	*	*	0.1	0.7	5.2	34.5	127.7	283.2	453.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic lower respiratory diseases (J44,J47)												
Black, both sexes												
2003	15.2	*	*	*	*	*	1.2	6.3	29.9	92.9	198.3	304.6
2002	15.4	*	*	*	*	*	1.2	6.7	29.7	89.5	211.3	320.1
2001	15.1	*	*	*	*	*	1.3	6.4	28.1	95.0	198.7	320.9
2000	14.9	*	*	*	*	*	1.3	6.6	30.9	94.5	195.2	310.6
1999	15.9	*	*	*	*	*	1.2	6.4	32.5	100.8	218.0	313.7
Black, male												
2003	18.0	*	*	*	*	*	1.3	7.4	38.7	131.3	325.5	520.3
2002	18.2	*	*	*	*	*	1.4	8.3	39.9	125.3	335.1	565.0
2001	18.1	*	*	*	*	*	1.4	7.4	37.0	137.4	330.5	583.2
2000	18.2	*	*	*	*	*	1.3	8.3	40.1	140.8	320.4	576.6
1999	19.6	*	*	*	*	*	1.5	8.2	44.6	145.4	366.5	607.0
Black, female												
2003	12.6	*	*	*	*	*	1.2	5.3	22.9	65.5	126.5	220.8
2002	12.9	*	*	*	*	*	1.1	5.3	21.5	64.0	141.3	226.3
2001	12.3	*	*	*	*	*	1.2	5.6	21.0	65.1	124.8	223.5
2000	12.0	*	*	*	*	*	1.3	5.1	23.4	62.0	124.8	211.6
1999	12.5	*	*	*	*	*	1.0	4.9	22.8	69.4	134.9	204.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
All races 3/, both sexes												
2003	0.4	*	*	*	*	*	*	*	0.3	1.3	3.8	6.4
2002	0.4	*	*	*	*	*	*	*	0.3	1.2	3.8	6.8
2001	0.4	*	*	*	*	*	*	*	0.3	1.5	4.1	6.2
2000	0.4	*	*	*	*	*	*	*	0.2	1.5	3.9	7.5
1999	0.4	*	*	*	*	*	*	*	0.2	1.5	4.1	6.1
All races, male												
2003	0.7	*	*	*	*	*	*	*	0.5	2.7	9.1	19.8
2002	0.7	*	*	*	*	*	*	*	0.5	2.6	9.0	21.6
2001	0.8	*	*	*	*	*	*	*	0.5	3.1	9.9	20.2
2000	0.8	*	*	*	*	*	*	*	0.5	3.2	9.6	24.9
1999	0.8	*	*	*	*	*	*	*	0.5	3.2	10.1	20.4
All races, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
White, both sexes												
2003	0.4	*	*	*	*	*	*	*	0.3	1.4	4.0	6.9
2002	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	7.2
2001	0.5	*	*	*	*	*	*	*	0.3	1.6	4.4	6.5
2000	0.5	*	*	*	*	*	*	*	0.3	1.7	4.1	7.9
1999	0.5	*	*	*	*	*	*	*	0.2	1.6	4.4	6.3
White, male												
2003	0.9	*	*	*	*	*	*	*	0.6	2.9	9.6	21.3
2002	0.9	*	*	*	*	*	*	*	0.6	2.8	9.4	22.8
2001	0.9	*	*	*	*	*	*	*	0.5	3.4	10.5	21.2
2000	0.9	*	*	*	*	*	*	*	0.5	3.4	10.0	26.4
1999	0.9	*	*	*	*	*	*	*	0.5	3.4	10.7	21.0
White, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2001	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumoconioses and chemical effects (J60-J66,J68)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	2.3	*
2002	0.2	*	*	*	*	*	*	*	*	*	2.4	*
2001	0.2	*	*	*	*	*	*	*	*	*	2.2	*
2000	0.2	*	*	*	*	*	*	*	*	*	2.5	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	6.2	*
2002	0.3	*	*	*	*	*	*	*	*	*	6.0	*
2001	0.3	*	*	*	*	*	*	*	*	*	6.2	*
2000	0.3	*	*	*	*	*	*	*	*	*	6.4	*
1999	0.4	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
All races 3/, both sexes												
2003	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6
All races, male												
2003	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4
All races, female												
2003	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
White, both sexes												
2003	6.5	*	*	*	0.1	0.2	0.3	1.0	2.6	9.3	44.7	178.7
2002	6.7	*	*	*	0.1	0.2	0.4	0.8	2.4	9.5	46.5	189.5
2001	6.6	*	*	*	0.1	0.2	0.4	0.9	2.5	9.6	45.8	190.8
2000	6.5	*	*	*	0.1	0.1	0.4	0.9	2.3	9.8	44.4	191.0
1999	5.9	*	*	*	0.1	0.2	0.3	0.7	2.3	9.0	40.7	178.0
White, male												
2003	6.6	*	*	*	0.1	0.2	0.4	1.2	3.2	12.4	61.1	250.8
2002	6.9	*	*	*	0.1	0.2	0.5	1.1	3.0	12.5	64.0	274.6
2001	6.7	*	*	*	*	0.2	0.5	1.2	3.1	12.9	63.3	275.4
2000	6.6	*	*	*	*	0.1	0.5	1.1	2.9	13.4	62.1	284.0
1999	6.1	*	*	*	*	0.2	0.4	0.8	2.9	12.2	58.7	263.6
White, female												
2003	6.4	*	*	*	*	0.1	0.3	0.7	2.0	6.7	33.6	146.7
2002	6.6	*	*	*	*	0.2	0.3	0.6	1.9	6.9	34.8	152.5
2001	6.6	*	*	*	*	0.2	0.3	0.7	2.1	6.9	34.2	155.3
2000	6.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	32.8	153.2
1999	5.8	*	*	*	*	0.2	0.3	0.6	1.7	6.4	29.0	143.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)												
Black, both sexes												
2003	4.2	*	*	*	*	*	0.7	1.3	5.1	13.7	54.4	163.4
2002	4.1	*	*	*	*	*	0.7	1.4	3.7	14.8	52.2	165.1
2001	4.3	*	*	*	*	*	0.6	1.6	4.0	15.1	53.2	185.5
2000	4.2	*	*	*	*	*	0.7	1.9	4.2	17.5	51.5	159.8
1999	4.2	*	*	*	*	*	0.7	1.5	4.4	16.2	52.2	167.1
Black, male												
2003	4.3	*	*	*	*	*	*	1.6	7.1	19.6	77.8	212.7
2002	4.2	*	*	*	*	*	0.8	2.0	4.8	21.7	73.4	233.7
2001	4.2	*	*	*	*	*	0.7	1.9	5.3	20.4	75.3	243.9
2000	4.4	*	*	*	*	*	0.8	2.7	5.0	27.8	74.5	209.9
1999	4.5	*	*	*	*	*	0.9	1.8	6.4	25.0	80.9	228.5
Black, female												
2003	4.1	*	*	*	*	*	0.8	1.0	3.6	9.6	41.1	144.2
2002	3.9	*	*	*	*	*	*	0.9	2.9	9.9	40.1	138.8
2001	4.4	*	*	*	*	*	*	1.4	2.9	11.3	40.8	163.8
2000	3.9	*	*	*	*	*	*	1.2	3.5	10.3	38.5	141.2
1999	3.8	*	*	*	*	*	*	1.2	2.8	10.0	36.1	144.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
All races 3/, both sexes												
2003	8.8	7.0	0.5	0.1	0.3	0.5	1.1	3.3	8.8	28.1	68.5	138.1
2002	8.7	6.9	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.7	68.0	133.5
2001	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6
All races, male												
2003	8.7	8.0	0.6	0.2	0.4	0.6	1.1	3.5	10.2	34.1	85.3	165.0
2002	8.5	7.9	0.7	0.2	0.3	0.4	1.3	3.3	10.5	35.3	83.9	156.6
2001	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3
All races, female												
2003	8.9	5.9	0.5	0.1	0.3	0.5	1.1	3.0	7.4	23.1	57.2	126.2
2002	8.8	5.8	0.4	0.2	0.3	0.5	1.1	3.1	8.0	23.1	57.5	123.4
2001	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
White, both sexes												
2003	9.6	5.4	0.4	0.1	0.3	0.5	0.9	2.9	8.6	29.0	70.8	143.2
2002	9.5	4.8	0.5	0.2	0.3	0.4	1.0	2.9	8.9	29.3	69.9	137.9
2001	10.0	5.0	0.5	0.2	0.3	0.4	1.1	3.0	9.3	30.6	75.4	146.0
2000	9.9	5.6	0.5	0.2	0.3	0.6	1.2	3.2	9.7	30.0	73.7	145.9
1999	9.7	4.9	0.5	0.2	0.3	0.6	1.1	3.0	9.2	29.6	72.2	150.7
White, male												
2003	9.4	5.8	0.4	0.2	0.3	0.5	1.0	3.3	9.9	35.1	87.6	170.5
2002	9.2	5.4	0.6	0.2	0.3	0.4	1.1	3.0	10.1	36.1	86.1	162.0
2001	9.7	5.9	0.5	0.2	0.3	0.4	1.2	3.4	10.6	37.4	92.0	173.5
2000	9.7	6.6	0.5	0.2	0.3	0.6	1.3	3.4	11.1	36.1	92.5	184.4
1999	9.6	5.6	0.6	0.2	0.4	0.6	1.2	3.2	10.3	36.8	92.0	185.7
White, female												
2003	9.8	5.0	0.4	*	0.3	0.4	0.9	2.6	7.2	23.8	59.4	131.0
2002	9.7	4.2	*	0.2	0.2	0.4	1.0	2.8	7.7	23.4	59.1	127.4
2001	10.2	4.0	0.6	0.2	0.3	0.5	0.9	2.6	8.2	24.9	64.3	134.4
2000	10.0	4.6	0.6	0.2	0.3	0.6	1.1	2.9	8.4	25.0	61.4	130.3
1999	9.8	4.3	0.5	0.2	0.2	0.5	0.9	2.8	8.2	23.6	59.3	136.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)												
Black, both sexes												
2003	5.9	15.0	1.1	*	0.6	1.0	2.4	6.4	12.2	24.2	50.8	89.6
2002	6.0	16.6	1.1	0.4	0.7	0.9	2.3	6.1	13.6	26.5	51.5	90.3
2001	6.7	18.0	1.0	*	0.6	1.3	2.9	7.3	13.6	29.5	56.7	109.6
2000	6.5	16.7	0.8	*	0.7	1.1	2.7	6.8	13.4	30.5	58.0	98.2
1999	6.5	15.4	1.0	*	0.4	1.3	2.9	6.4	13.4	30.8	57.9	111.6
Black, male												
2003	6.0	18.4	*	*	0.8	1.1	2.2	6.3	15.1	30.8	68.5	113.6
2002	6.0	19.5	*	*	*	0.9	2.7	6.5	16.1	31.7	65.7	100.8
2001	6.4	23.1	*	*	*	1.3	2.6	7.6	15.3	32.9	69.4	129.3
2000	6.5	16.3	*	*	0.8	1.2	3.1	7.8	15.2	36.0	73.0	132.6
1999	6.4	17.6	*	*	*	1.0	2.9	7.1	15.5	35.1	72.6	145.3
Black, female												
2003	5.7	11.4	*	*	*	1.0	2.7	6.5	9.8	19.4	40.8	80.2
2002	6.0	13.6	*	*	0.7	0.8	2.0	5.8	11.5	22.9	43.5	86.3
2001	7.0	12.6	*	*	*	1.2	3.1	7.0	12.2	27.1	49.6	102.3
2000	6.4	17.2	*	*	*	1.0	2.2	5.9	11.9	26.6	49.6	85.4
1999	6.7	13.1	*	*	*	1.6	2.8	5.8	11.6	27.8	49.7	99.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
All races 3/, both sexes												
2003	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
2002	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6
All races, male												
2003	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
2002	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5
All races, female												
2003	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
2002	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
White, both sexes												
2003	1.4	*	*	*	*	0.1	0.2	0.6	1.3	3.2	9.8	26.9
2002	1.5	*	*	*	*	*	0.2	0.6	1.4	3.6	10.3	29.1
2001	1.7	*	*	*	*	*	0.2	0.7	1.5	3.9	11.5	34.4
2000	1.7	*	*	*	*	*	0.2	0.7	1.8	4.0	11.5	36.3
1999	1.8	*	*	*	*	*	0.2	0.6	1.7	4.4	12.3	35.3
White, male												
2003	1.4	*	*	*	*	*	0.2	0.8	1.6	4.1	11.4	29.0
2002	1.5	*	*	*	*	*	0.3	0.8	1.8	4.4	12.3	32.5
2001	1.5	*	*	*	*	*	0.3	0.8	2.0	4.5	12.9	35.7
2000	1.7	*	*	*	*	*	0.3	1.0	2.4	5.2	13.5	40.7
1999	1.7	*	*	*	*	*	0.2	0.8	2.2	5.9	14.9	38.4
White, female												
2003	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.8	26.0
2002	1.6	*	*	*	*	*	0.1	0.5	1.1	2.9	9.0	27.6
2001	1.9	*	*	*	*	*	0.1	0.5	1.1	3.3	10.7	33.9
2000	1.8	*	*	*	*	*	0.1	0.4	1.2	3.0	10.2	34.5
1999	1.9	*	*	*	*	*	0.2	0.4	1.3	3.2	10.7	34.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Peptic ulcer (K25-K28)												
Black, both sexes												
2003	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.0	16.9
2002	1.0	*	*	*	*	*	0.6	1.2	2.6	4.1	8.0	19.0
2001	1.1	*	*	*	*	*	0.6	1.3	2.7	4.4	9.1	20.9
2000	1.2	*	*	*	*	*	0.5	1.7	2.7	4.8	10.0	26.6
1999	1.2	*	*	*	*	*	0.8	1.5	2.8	5.7	10.9	25.5
Black, male												
2003	1.2	*	*	*	*	*	*	2.3	3.4	5.2	16.1	*
2002	1.2	*	*	*	*	*	0.8	1.9	3.8	5.9	11.5	*
2001	1.4	*	*	*	*	*	1.0	2.2	4.2	7.0	12.4	23.8
2000	1.3	*	*	*	*	*	0.8	2.3	3.7	7.0	13.9	31.1
1999	1.4	*	*	*	*	*	1.2	2.3	4.4	8.1	12.3	28.1
Black, female												
2003	0.8	*	*	*	*	*	*	0.8	*	*	8.1	15.8
2002	0.8	*	*	*	*	*	*	*	1.6	2.8	6.0	18.5
2001	0.9	*	*	*	*	*	*	*	1.5	2.7	7.3	19.8
2000	1.0	*	*	*	*	*	*	1.1	1.8	3.3	7.8	24.9
1999	1.1	*	*	*	*	*	*	*	*	3.9	10.2	24.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
All races 3/, both sexes												
2003	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5
All races, male												
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
2002	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9
All races, female												
2003	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
2002	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
White, both sexes												
2003	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.1
2001	0.1	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
1999	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.5
White, male												
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.5
2002	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.4	2.0
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.5
2000	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.0
1999	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.7	1.3
2002	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.9
2000	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Diseases of appendix (K35-K38)												
Black, both sexes												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
All races 3/, both sexes												
2003	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
2002	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7
All races, male												
2003	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
2002	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2
All races, female												
2003	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
2002	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
White, both sexes												
2003	0.6	0.9	*	*	*	*	0.1	0.3	0.5	1.3	3.6	12.3
2002	0.6	0.8	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.4
2001	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.8
2000	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.3
1999	0.6	1.1	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0
White, male												
2003	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
2002	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.9	11.9
2001	0.4	*	*	*	*	*	*	0.2	0.5	1.2	3.4	11.8
2000	0.5	1.5	*	*	*	*	*	0.2	0.4	1.1	4.1	12.7
1999	0.5	1.3	*	*	*	*	0.1	0.2	0.6	1.2	4.1	12.9
White, female												
2003	0.7	*	*	*	*	*	*	0.2	0.6	1.3	3.4	12.2
2002	0.8	*	*	*	*	*	*	0.2	0.5	1.2	4.1	14.0
2001	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.2
2000	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.6
1999	0.7	*	*	*	*	*	*	*	0.4	1.1	3.5	13.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hernia (K40-K46)												
Black, both sexes												
2003	0.4	*	*	*	*	*	*	*	1.0	1.4	4.0	9.0
2002	0.4	*	*	*	*	*	*	*	0.8	1.6	3.1	7.7
2001	0.4	*	*	*	*	*	*	*	0.9	1.5	3.6	11.1
2000	0.4	*	*	*	*	*	*	*	*	1.3	4.1	12.5
1999	0.4	*	*	*	*	*	*	*	*	1.5	4.2	9.9
Black, male												
2003	0.4	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.5	*	*	*	*	*	*	*	*	*	3.8	8.1
2002	0.4	*	*	*	*	*	*	*	*	*	*	*
2001	0.4	*	*	*	*	*	*	*	*	*	*	8.8
2000	0.5	*	*	*	*	*	*	*	*	*	4.4	12.4
1999	0.5	*	*	*	*	*	*	*	*	*	4.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
All races 3/, both sexes												
2003	9.5	*	*	*	*	0.9	6.8	18.3	23.0	29.5	30.0	20.1
2002	9.5	*	*	*	0.1	0.9	7.0	18.0	22.9	29.4	31.4	21.4
2001	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2
All races, male												
2003	12.5	*	*	*	*	1.2	9.2	27.5	33.8	39.1	36.9	27.5
2002	12.3	*	*	*	*	1.2	9.3	26.9	33.1	38.5	38.4	28.4
2001	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1
All races, female												
2003	6.5	*	*	*	*	0.5	4.4	9.4	13.1	21.4	25.4	16.8
2002	6.7	*	*	*	*	0.7	4.8	9.5	13.5	21.9	26.7	18.4
2001	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
White, both sexes												
2003	10.2	*	*	*	*	0.9	7.1	18.4	23.1	30.7	31.6	20.9
2002	10.1	*	*	*	0.1	0.9	7.3	18.1	23.1	30.7	32.8	22.3
2001	10.1	*	*	*	*	1.0	7.5	18.3	22.7	30.9	31.5	23.2
2000	10.0	*	*	*	0.1	1.0	7.4	17.5	23.8	30.6	32.3	23.9
1999	9.9	*	*	*	*	1.0	7.2	17.0	23.3	31.4	33.0	23.8
White, male												
2003	13.4	*	*	*	*	1.3	9.6	27.8	33.8	40.5	38.5	28.4
2002	13.2	*	*	*	*	1.2	9.7	27.1	33.2	39.8	40.2	29.6
2001	13.1	*	*	*	*	1.3	9.9	27.3	32.8	41.1	38.4	33.5
2000	13.2	*	*	*	*	1.4	10.3	26.8	34.5	40.9	39.9	35.5
1999	13.1	*	*	*	*	1.3	10.3	26.4	34.1	42.4	40.8	34.2
White, female												
2003	7.0	*	*	*	*	0.5	4.4	9.2	13.1	22.4	27.0	17.5
2002	7.2	*	*	*	*	0.6	4.8	9.1	13.5	23.0	27.9	19.1
2001	7.1	*	*	*	*	0.6	5.0	9.4	13.3	22.4	27.0	18.9
2000	6.8	*	*	*	*	0.6	4.5	8.2	13.7	21.9	27.2	19.2
1999	6.8	*	*	*	*	0.7	4.2	7.8	13.2	22.2	27.9	19.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic liver disease and cirrhosis (K70,K73-K74)												
Black, both sexes												
2003	6.9	*	*	*	*	0.7	5.7	19.8	25.8	23.8	14.4	12.3
2002	6.9	*	*	*	*	0.9	5.5	20.8	25.0	21.6	18.5	12.2
2001	7.5	*	*	*	*	0.9	7.3	22.6	25.9	25.0	18.5	10.4
2000	7.5	*	*	*	*	1.0	7.8	22.1	25.9	26.2	18.0	13.8
1999	7.8	*	*	*	*	1.0	8.0	22.7	29.3	26.6	21.5	17.2
Black, male												
2003	9.5	*	*	*	*	1.0	7.1	30.4	40.8	33.4	20.6	21.9
2002	9.0	*	*	*	*	1.0	6.7	30.0	37.8	32.4	22.4	*
2001	10.4	*	*	*	*	1.0	10.0	34.5	41.5	39.1	22.4	*
2000	9.9	*	*	*	*	1.0	10.5	32.7	38.0	37.7	24.5	*
1999	10.7	*	*	*	*	1.1	11.1	35.4	43.3	40.7	29.5	*
Black, female												
2003	4.5	*	*	*	*	*	4.4	10.6	13.7	16.9	10.8	8.5
2002	4.9	*	*	*	*	0.8	4.4	12.9	14.7	14.0	16.2	10.7
2001	4.8	*	*	*	*	0.7	4.8	12.4	13.4	15.0	16.4	9.7
2000	5.2	*	*	*	*	1.0	5.5	13.1	16.2	18.1	14.3	11.6
1999	5.2	*	*	*	*	0.9	5.3	11.8	18.1	16.7	17.0	15.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
All races 3/, both sexes												
2003	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
2002	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8
All races, male												
2003	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
2002	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1
All races, female												
2003	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
2002	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
White, both sexes												
2003	4.5	*	*	*	*	0.6	4.6	10.7	11.7	10.0	5.9	1.9
2002	4.4	*	*	*	*	0.6	4.6	10.5	11.7	10.4	5.8	2.1
2001	4.5	*	*	*	*	0.7	4.8	10.8	11.5	10.2	5.8	2.3
2000	4.5	*	*	*	*	0.7	4.9	10.3	12.2	10.5	6.1	2.3
1999	4.4	*	*	*	*	0.6	4.8	10.2	11.8	10.5	6.4	2.7
White, male												
2003	6.8	*	*	*	*	0.9	6.2	16.4	18.5	16.4	10.4	3.7
2002	6.6	*	*	*	*	0.8	6.1	16.0	18.5	16.8	9.9	4.5
2001	6.7	*	*	*	*	0.9	6.5	16.3	18.4	16.6	10.4	5.4
2000	6.8	*	*	*	*	1.0	6.8	16.2	19.2	17.5	11.7	5.4
1999	6.7	*	*	*	*	0.8	6.7	16.2	19.0	17.9	11.1	5.8
White, female												
2003	2.3	*	*	*	*	0.3	3.0	5.1	5.3	4.5	2.9	1.1
2002	2.3	*	*	*	*	0.3	3.0	5.1	5.2	5.0	3.1	1.1
2001	2.3	*	*	*	*	0.4	3.1	5.3	5.1	4.8	2.8	1.0
2000	2.1	*	*	*	*	0.3	3.0	4.4	5.6	4.6	2.4	1.0
1999	2.1	*	*	*	*	0.4	2.8	4.4	5.0	4.3	3.3	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcoholic liver disease (K70)												
Black, both sexes												
2003	3.2	*	*	*	*	*	3.3	9.9	12.5	9.3	3.4	*
2002	3.2	*	*	*	*	0.5	2.9	10.0	12.3	8.6	5.3	*
2001	3.6	*	*	*	*	0.5	4.2	11.5	12.9	10.7	4.7	*
2000	3.8	*	*	*	*	0.6	5.0	12.0	13.3	10.3	5.5	*
1999	4.1	*	*	*	*	0.5	5.0	12.8	15.1	11.8	7.1	*
Black, male												
2003	4.8	*	*	*	*	*	4.1	15.8	20.1	16.0	6.8	*
2002	4.4	*	*	*	*	*	3.8	14.7	18.7	15.0	9.2	*
2001	5.3	*	*	*	*	*	5.9	17.1	21.4	19.5	7.7	*
2000	5.3	*	*	*	*	*	6.9	17.4	19.4	18.1	10.0	*
1999	5.9	*	*	*	*	*	6.7	19.7	24.2	21.8	12.6	*
Black, female												
2003	1.8	*	*	*	*	*	2.5	4.7	6.4	4.5	*	*
2002	2.0	*	*	*	*	*	2.2	5.9	7.1	4.1	*	*
2001	2.1	*	*	*	*	*	2.7	6.7	6.1	4.4	*	*
2000	2.4	*	*	*	*	*	3.3	7.3	8.4	4.7	*	*
1999	2.4	*	*	*	*	*	3.4	6.9	7.9	4.8	4.0	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
All races 3/, both sexes												
2003	5.2	*	*	*	*	0.3	2.4	7.8	11.4	19.7	24.3	18.2
2002	5.2	*	*	*	0.1	0.3	2.6	7.7	11.3	19.3	25.7	19.3
2001	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4
All races, male												
2003	6.2	*	*	*	*	0.3	3.2	11.4	15.5	23.0	26.8	23.9
2002	6.1	*	*	*	*	0.4	3.5	11.2	14.8	22.0	28.7	23.7
2001	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0
All races, female												
2003	4.3	*	*	*	*	0.2	1.5	4.3	7.6	17.0	22.6	15.8
2002	4.5	*	*	*	*	0.3	1.8	4.3	8.1	17.1	23.7	17.4
2001	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1
2000	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
White, both sexes												
2003	5.7	*	*	*	*	0.3	2.5	7.7	11.4	20.8	25.7	19.0
2002	5.7	*	*	*	0.1	0.3	2.7	7.5	11.4	20.3	27.0	20.2
2001	5.6	*	*	*	*	0.3	2.7	7.5	11.2	20.7	25.7	20.9
2000	5.5	*	*	*	*	0.3	2.5	7.2	11.6	20.1	26.1	21.6
1999	5.5	*	*	*	*	0.4	2.5	6.8	11.5	20.9	26.6	21.1
White, male												
2003	6.6	*	*	*	*	0.3	3.4	11.4	15.3	24.1	28.1	24.7
2002	6.5	*	*	*	*	0.4	3.7	11.1	14.7	23.0	30.3	25.1
2001	6.4	*	*	*	*	0.4	3.4	11.0	14.4	24.4	28.0	28.1
2000	6.4	*	*	*	*	0.3	3.5	10.6	15.3	23.4	28.2	30.1
1999	6.4	*	*	*	*	0.5	3.5	10.2	15.1	24.4	29.7	28.4
White, female												
2003	4.7	*	*	*	*	0.2	1.5	4.1	7.8	17.9	24.0	16.5
2002	4.9	*	*	*	*	0.3	1.8	4.0	8.3	18.0	24.8	18.1
2001	4.8	*	*	*	*	0.2	1.9	4.2	8.2	17.6	24.2	17.9
2000	4.7	*	*	*	*	0.2	1.5	3.8	8.1	17.3	24.8	18.2
1999	4.6	*	*	*	*	0.3	1.4	3.4	8.2	17.9	24.6	18.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other chronic liver disease and cirrhosis (K73-K74)												
Black, both sexes												
2003	3.7	*	*	*	*	0.4	2.4	9.9	13.3	14.5	11.0	10.8
2002	3.7	*	*	*	*	0.5	2.5	10.8	12.7	13.0	13.2	9.8
2001	3.8	*	*	*	*	0.4	3.1	11.1	13.0	14.3	13.9	7.7
2000	3.7	*	*	*	*	0.4	2.8	10.1	12.6	16.0	12.5	12.2
1999	3.8	*	*	*	*	0.5	3.0	9.9	14.2	14.8	14.5	12.8
Black, male												
2003	4.7	*	*	*	*	*	3.0	14.6	20.8	17.3	13.8	*
2002	4.6	*	*	*	*	*	2.9	15.3	19.1	17.4	13.2	*
2001	5.1	*	*	*	*	*	4.1	17.4	20.1	19.5	14.7	*
2000	4.7	*	*	*	*	*	3.6	15.2	18.6	19.6	14.5	*
1999	4.8	*	*	*	*	*	4.4	15.8	19.1	19.0	16.9	*
Black, female												
2003	2.7	*	*	*	*	*	1.9	5.9	7.3	12.4	9.4	8.1
2002	2.9	*	*	*	*	*	2.2	7.0	7.6	9.9	13.2	9.9
2001	2.7	*	*	*	*	*	2.1	5.7	7.3	10.6	13.4	*
2000	2.8	*	*	*	*	*	2.2	5.8	7.8	13.4	11.4	11.2
1999	2.9	*	*	*	*	*	1.9	4.9	10.2	11.9	13.1	13.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
All races 3/, both sexes												
2003	1.0	*	*	*	*	0.1	0.1	0.3	0.7	2.7	8.2	21.8
2002	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9
All races, male												
2003	0.9	*	*	*	*	*	0.1	0.3	0.8	3.2	10.1	22.7
2002	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7
All races, female												
2003	1.1	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
White, both sexes												
2003	1.1	*	*	*	*	*	0.1	0.3	0.6	2.7	8.2	21.9
2002	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.6	23.5
2001	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	7.4	23.9
2000	1.1	*	*	*	*	*	0.1	0.2	0.7	2.3	7.4	25.6
1999	1.1	*	*	*	*	*	0.1	0.3	0.7	2.5	7.8	24.4
White, male												
2003	1.0	*	*	*	*	*	*	0.3	0.7	3.2	10.0	23.0
2002	1.0	*	*	*	*	*	0.1	0.4	0.9	3.1	9.8	24.7
2001	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.3	26.7
2000	1.0	*	*	*	*	*	0.1	0.2	0.9	2.8	9.6	28.9
1999	1.0	*	*	*	*	*	0.1	0.3	0.9	3.0	9.6	28.1
White, female												
2003	1.2	*	*	*	*	*	*	0.2	0.5	2.2	6.9	21.4
2002	1.2	*	*	*	*	*	*	0.2	0.6	2.4	6.2	23.0
2001	1.2	*	*	*	*	*	*	0.3	0.7	2.4	6.1	22.8
2000	1.2	*	*	*	*	*	*	0.2	0.5	1.8	6.0	24.3
1999	1.2	*	*	*	*	*	*	0.3	0.6	2.1	6.7	22.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)												
Black, both sexes												
2003	0.7	*	*	*	*	*	*	0.7	1.2	2.9	7.9	20.7
2002	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.7	14.8
2001	0.8	*	*	*	*	*	*	0.7	1.7	4.7	8.3	19.7
2000	0.8	*	*	*	*	*	*	0.7	1.2	3.9	8.3	17.8
1999	0.7	*	*	*	*	*	0.4	0.5	1.5	3.6	7.2	15.3
Black, male												
2003	0.7	*	*	*	*	*	*	*	*	3.2	10.4	20.9
2002	0.6	*	*	*	*	*	*	*	*	*	10.9	*
2001	0.7	*	*	*	*	*	*	*	2.1	4.9	6.2	27.2
2000	0.6	*	*	*	*	*	*	*	*	3.7	8.2	*
1999	0.6	*	*	*	*	*	*	*	*	3.7	7.1	*
Black, female												
2003	0.8	*	*	*	*	*	*	0.8	*	2.7	6.5	20.7
2002	0.8	*	*	*	*	*	*	*	*	3.7	9.1	16.0
2001	0.9	*	*	*	*	*	*	*	*	4.5	9.4	16.8
2000	1.0	*	*	*	*	*	*	*	*	4.1	8.4	19.3
1999	0.8	*	*	*	*	*	*	*	*	3.5	7.2	15.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
All races 3/, both sexes												
2003	14.6	4.5	*	0.1	0.2	0.7	1.8	4.9	13.6	40.1	109.5	293.1
2002	14.2	4.3	*	0.1	0.2	0.7	1.7	4.7	13.0	39.2	109.1	288.6
2001	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9
All races, male												
2003	14.3	5.2	*	*	0.3	0.7	2.1	6.0	15.4	45.3	136.8	387.6
2002	13.9	5.3	*	*	0.2	0.6	1.8	5.5	14.3	45.8	136.2	385.5
2001	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1
All races, female												
2003	14.9	3.8	*	*	0.2	0.7	1.5	3.9	12.0	35.6	91.3	251.4
2002	14.5	3.2	*	*	0.2	0.7	1.5	4.0	11.8	33.7	91.1	246.6
2001	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
White, both sexes												
2003	14.3	3.8	*	*	0.2	0.5	1.2	3.4	10.6	33.9	102.4	284.8
2002	13.9	3.6	*	*	0.1	0.5	1.1	3.2	9.9	33.4	102.2	278.8
2001	13.5	2.4	*	*	0.2	0.4	1.2	3.2	9.7	34.2	96.9	278.2
2000	12.9	2.9	*	0.1	0.2	0.4	1.1	3.0	9.7	31.9	94.0	268.7
1999	12.3	3.4	*	0.1	0.1	0.4	1.0	2.7	9.1	31.5	90.1	255.7
White, male												
2003	14.0	4.5	*	*	0.2	0.5	1.4	4.1	11.9	39.0	130.1	383.6
2002	13.7	4.4	*	*	0.1	0.5	1.2	3.8	10.7	40.0	130.4	377.4
2001	13.3	2.8	*	*	0.2	0.3	1.4	3.8	10.9	40.4	123.5	388.9
2000	12.7	3.8	*	*	0.2	0.4	1.3	3.6	11.0	37.7	121.1	376.2
1999	12.1	4.1	*	*	*	0.4	1.2	3.1	10.6	36.9	117.2	360.8
White, female												
2003	14.5	3.0	*	*	0.1	0.5	0.9	2.7	9.3	29.5	83.6	241.0
2002	14.1	2.7	*	*	0.1	0.5	1.0	2.7	9.1	27.8	83.3	235.9
2001	13.7	2.1	*	*	0.2	0.5	1.0	2.6	8.6	29.0	79.2	231.9
2000	13.0	2.0	*	*	0.2	0.4	0.9	2.4	8.4	27.1	76.2	225.0
1999	12.4	2.8	*	*	0.2	0.4	0.9	2.3	7.8	26.9	72.3	213.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)												
Black, both sexes												
2003	20.6	9.1	*	*	0.5	2.2	6.2	16.4	42.8	103.3	207.0	440.5
2002	19.8	8.2	*	*	0.6	2.0	5.3	16.2	41.0	100.1	203.2	441.6
2001	19.6	8.1	*	*	0.4	1.8	5.3	15.8	41.6	102.5	202.1	429.6
2000	18.9	11.5	*	*	0.4	1.8	5.3	15.8	40.6	99.9	195.1	406.4
1999	18.6	9.6	*	*	0.5	1.9	5.4	15.2	37.2	95.4	195.2	433.3
Black, male												
2003	20.0	9.5	*	*	0.6	2.4	7.1	21.2	51.5	120.5	246.0	518.2
2002	19.1	10.2	*	*	*	1.9	6.0	20.1	49.1	115.0	235.2	561.8
2001	18.0	8.7	*	*	*	2.2	5.2	19.3	45.4	114.0	233.4	511.7
2000	17.7	13.4	*	*	*	1.9	6.5	20.3	44.9	104.8	231.6	490.1
1999	17.5	8.1	*	*	*	1.8	6.5	18.9	40.4	102.9	250.0	526.1
Black, female												
2003	21.1	8.7	*	*	*	1.9	5.4	12.2	35.8	91.0	185.0	410.4
2002	20.5	6.1	*	*	0.7	2.1	4.7	12.8	34.4	89.6	185.0	395.5
2001	21.0	7.6	*	*	*	1.4	5.5	12.7	38.5	94.4	184.6	399.1
2000	20.0	9.6	*	*	*	1.7	4.3	11.9	37.1	96.5	174.6	375.2
1999	19.5	11.1	*	*	*	1.9	4.5	12.0	34.6	90.1	164.5	398.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
All races 3/, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
All races, male												
2003	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999	0.1	*	*	*	*	*	*	*	*	*	0.7	*
All races, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	0.6
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
White, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.5
2002	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2001	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2000	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.1
1999	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8
White, male												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.5	*
1999	0.1	*	*	*	*	*	*	*	*	*	0.6	*
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	0.8
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	*
2000	0.1	*	*	*	*	*	*	*	*	*	0.3	1.0
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
All races 3/, both sexes												
2003	0.2	*	*	*	*	*	0.0	0.1	0.2	0.6	1.3	2.8
2002	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.8
2001	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2
All races, male												
2003	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.6	4.1
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0
All races, female												
2003	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.2
2002	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
White, both sexes												
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.6
2002	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
2001	0.2	*	*	*	*	*	*	0.1	0.1	0.6	1.1	2.9
2000	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.4
1999	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9
White, male												
2003	0.2	*	*	*	*	*	*	0.1	*	0.5	1.4	3.8
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.8
2001	0.2	*	*	*	*	*	*	*	*	0.7	1.7	4.0
2000	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	4.0
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	2.2	4.6
White, female												
2003	0.2	*	*	*	*	*	*	*	0.2	0.5	1.0	2.1
2002	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.0
2001	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	1.8
1999	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)												
Black, both sexes												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	2.2	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	2.2	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
All races 3/, both sexes												
2003	14.4	4.2	*	0.1	0.2	0.7	1.7	4.8	13.4	39.3	107.8	289.7
2002	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8
All races, male												
2003	14.1	4.8	*	*	0.2	0.7	2.0	5.7	15.2	44.5	134.8	383.1
2002	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6
All races, female												
2003	14.6	3.7	*	*	0.2	0.6	1.5	3.8	11.8	34.9	89.7	248.4
2002	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
White, both sexes												
2003	14.0	3.5	*	*	0.2	0.5	1.1	3.3	10.3	33.2	100.7	281.5
2002	13.6	3.4	*	*	0.1	0.4	1.1	3.1	9.6	32.7	100.5	275.3
2001	13.2	2.2	*	*	0.2	0.4	1.1	3.1	9.5	33.4	95.3	274.6
2000	12.6	2.7	*	*	0.2	0.4	1.1	2.9	9.4	31.2	92.2	265.0
1999	12.0	3.3	*	*	0.1	0.4	1.0	2.5	8.9	30.8	87.9	251.9
White, male												
2003	13.8	4.1	*	*	0.2	0.5	1.3	3.9	11.7	38.3	128.3	379.3
2002	13.4	4.2	*	*	*	0.4	1.2	3.6	10.4	39.2	128.3	372.7
2001	13.0	2.6	*	*	0.2	0.3	1.3	3.7	10.7	39.5	121.5	383.9
2000	12.4	3.6	*	*	0.2	0.4	1.2	3.4	10.6	36.9	118.9	370.4
1999	11.8	3.9	*	*	*	0.4	1.1	3.0	10.3	36.2	114.4	354.9
White, female												
2003	14.2	2.8	*	*	*	0.5	0.9	2.7	9.0	28.8	82.1	238.2
2002	13.8	2.6	*	*	*	0.5	1.0	2.6	8.9	27.2	81.8	232.9
2001	13.4	1.8	*	*	0.2	0.4	0.9	2.5	8.4	28.3	78.0	228.8
2000	12.8	1.7	*	*	0.2	0.4	0.9	2.3	8.2	26.5	74.6	222.1
1999	12.2	2.8	*	*	0.1	0.4	0.8	2.2	7.6	26.2	70.6	210.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Renal failure (N17-N19)												
Black, both sexes												
2003	20.3	8.9	*	*	0.5	2.1	6.1	16.1	42.5	102.0	204.4	436.4
2002	19.5	7.7	*	*	0.5	1.9	5.3	15.7	40.6	99.6	200.2	434.2
2001	19.3	8.1	*	*	0.4	1.7	5.2	15.4	41.1	101.3	200.0	425.6
2000	18.5	11.1	*	*	0.4	1.7	5.1	15.2	40.1	98.6	192.4	400.4
1999	18.2	9.6	*	*	0.4	1.8	5.2	14.8	36.4	94.1	191.8	425.3
Black, male												
2003	19.7	9.2	*	*	*	2.4	7.0	20.8	51.2	118.8	241.7	513.0
2002	18.8	9.6	*	*	*	1.7	5.9	19.6	48.8	114.2	231.5	555.3
2001	17.7	8.7	*	*	*	2.1	5.0	18.8	44.7	112.4	231.6	508.3
2000	17.3	13.4	*	*	*	1.8	6.3	19.5	44.2	103.3	229.2	480.9
1999	17.1	8.1	*	*	*	1.7	6.2	18.5	39.7	101.5	245.7	513.2
Black, female												
2003	20.9	8.7	*	*	*	1.8	5.3	12.1	35.5	90.0	183.2	406.7
2002	20.2	*	*	*	*	2.0	4.7	12.3	34.1	89.2	182.6	387.7
2001	20.7	7.6	*	*	*	1.3	5.4	12.6	38.2	93.5	182.3	394.9
2000	19.6	8.6	*	*	*	1.6	4.0	11.5	36.7	95.3	171.7	370.4
1999	19.1	11.1	*	*	*	1.9	4.3	11.7	33.8	88.9	161.6	392.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
All races 3/, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
White, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other disorders of kidney (N25,N27)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
All races 3/, both sexes												
2003	0.3	*	*	*	*	*	0.1	0.2	0.2	0.6	1.9	5.9
2002	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.3
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5
All races, male												
2003	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8
All races, female												
2003	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.1
2002	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.7
2001	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
White, both sexes												
2003	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.0
2002	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.4
2001	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.9	6.1
2000	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999	0.3	*	*	*	*	*	0.1	0.1	0.2	0.7	2.0	7.7
White, male												
2003	0.2	*	*	*	*	*	*	0.1	*	0.4	1.3	5.8
2002	0.2	*	*	*	*	*	*	*	*	0.5	1.4	5.4
2001	0.2	*	*	*	*	*	*	*	*	0.4	1.4	4.3
2000	0.2	*	*	*	*	*	*	*	*	*	1.5	5.9
1999	0.2	*	*	*	*	*	*	*	0.2	0.5	1.8	7.2
White, female												
2003	0.4	*	*	*	*	*	*	0.2	0.3	0.8	2.2	6.0
2002	0.4	*	*	*	*	*	*	*	0.2	0.7	2.3	6.8
2001	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	6.9
2000	0.4	*	*	*	*	*	*	*	0.3	0.7	2.2	8.3
1999	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.2	7.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Infections of kidney (N10-N12,N13.6,N15.1)												
Black, both sexes												
2003	0.2	*	*	*	*	*	*	*	*	*	2.4	5.8
2002	0.2	*	*	*	*	*	*	*	*	*	*	6.2
2001	0.2	*	*	*	*	*	*	*	*	*	2.3	*
2000	0.3	*	*	*	*	*	*	*	*	*	3.1	7.8
1999	0.2	*	*	*	*	*	*	*	*	1.2	*	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	0.3	*	*	*	*	*	*	*	*	*	*	*
2002	0.3	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.4	*	*	*	*	*	*	*	*	*	3.6	9.0
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
All races 3/, both sexes												
2003	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.3
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.8
2001	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.5
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.0
All races, male												
2003	0.3	*	*	*	*	*	*	*	*	0.6	3.2	17.2
2002	0.3	*	*	*	*	*	*	*	*	0.5	3.2	15.8
2001	0.3	*	*	*	*	*	*	*	*	0.5	2.8	18.6
2000	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999	0.3	*	*	*	*	*	*	*	*	0.6	3.3	17.5
All races, female												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
White, both sexes												
2003	0.2	*	*	*	*	*	*	*	*	0.3	1.3	5.5
2002	0.2	*	*	*	*	*	*	*	*	0.2	1.3	4.9
2001	0.2	*	*	*	*	*	*	*	*	0.3	1.1	5.7
2000	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.2
1999	0.2	*	*	*	*	*	*	*	*	0.2	1.3	5.1
White, male												
2003	0.4	*	*	*	*	*	*	*	*	0.6	3.2	18.0
2002	0.3	*	*	*	*	*	*	*	*	0.5	3.2	16.1
2001	0.4	*	*	*	*	*	*	*	*	0.5	2.9	19.4
2000	0.3	*	*	*	*	*	*	*	*	0.5	3.3	18.1
1999	0.3	*	*	*	*	*	*	*	*	0.5	3.4	17.9
White, female												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Hyperplasia of prostate (N40)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.2	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
All races 3/, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2000	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.3	*
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
White, both sexes												
2003	0.0	*	*	*	*	*	*	*	*	*	0.2	0.7
2002	0.0	*	*	*	*	*	*	*	*	*	0.2	0.8
2001	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2000	0.0	*	*	*	*	*	*	*	*	*	0.2	0.6
1999	0.0	*	*	*	*	*	*	*	*	*	0.2	*
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	0.4	1.0
2002	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2001	0.1	*	*	*	*	*	*	*	*	*	0.4	1.2
2000	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
1999	0.1	*	*	*	*	*	*	*	*	*	0.4	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Inflammatory diseases of female pelvic organs (N70-N76)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
All races 3/, both sexes												
2003	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2002	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*
1999	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	0.4	...	...	*	0.6	1.2	0.7	0.1	*	*	*	*
2002	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2001	0.3	...	...	*	0.6	1.0	0.5	*	*	*	*	*
2000	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
1999	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
White, both sexes												
2003	0.1	...	...	*	0.2	0.4	0.3	0.1	*	*	*	*
2002	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.2	0.4	0.1	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.3	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	0.3	...	...	*	0.4	0.9	0.5	0.1	*	*	*	*
2002	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
2001	0.2	...	...	*	0.4	0.7	0.3	*	*	*	*	*
2000	0.2	...	...	*	0.4	0.7	0.4	*	*	*	*	*
1999	0.2	...	...	*	0.3	0.6	0.4	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy, childbirth and the puerperium (O00-O99)												
Black, both sexes												
2003	0.5	...	...	*	0.7	1.5	1.0	*	*	*	*	*
2002	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2001	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2000	0.4	...	...	*	0.6	1.2	0.7	*	*	*	*	*
1999	0.4	...	...	*	0.8	1.3	0.7	*	*	*	*	*
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	1.0	...	...	*	1.4	2.9	1.9	*	*	*	*	*
2002	0.8	...	...	*	1.5	2.3	1.4	*	*	*	*	*
2001	0.8	...	...	*	1.5	2.3	1.4	*	*	*	*	*
2000	0.7	...	...	*	1.2	2.3	1.2	*	*	*	*	*
1999	0.9	...	...	*	1.6	2.5	1.4	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
All races 3/, both sexes												
2003	0.0	...	...	*	*	*	*	*	*	*	*	*
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	0.0	...	...	*	*	*	*	*	*	*	*	*
2002	0.0	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	0.0	...	...	*	*	*	*	*	*	*	*	*
1999	0.0	...	...	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
White, both sexes												
2003	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	0.0	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Pregnancy with abortive outcome (O00-O07)												
Black, both sexes												
2003	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	*	...	...	*	*	*	*	*	*	*	*	*
2002	*	...	...	*	*	*	*	*	*	*	*	*
2001	*	...	...	*	*	*	*	*	*	*	*	*
2000	*	...	...	*	*	*	*	*	*	*	*	*
1999	*	...	...	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
All races 3/, both sexes												
2003	0.2	...	...	*	0.3	0.5	0.4	0.1	*	*	*	*
2002	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.2	0.4	0.3	*	*	*	*	*
All races, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
All races, female												
2003	0.3	...	...	*	0.5	1.1	0.7	0.1	*	*	*	*
2002	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
1999	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
White, both sexes												
2003	0.1	...	...	*	0.2	0.4	0.3	0.1	*	*	*	*
2002	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
2001	0.1	...	...	*	0.2	0.3	0.1	*	*	*	*	*
2000	0.1	...	...	*	0.2	0.3	0.2	*	*	*	*	*
1999	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
White, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
White, female												
2003	0.3	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2002	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
2001	0.2	...	...	*	0.4	0.7	0.3	*	*	*	*	*
2000	0.2	...	...	*	0.4	0.6	0.4	*	*	*	*	*
1999	0.2	...	...	*	0.3	0.6	0.4	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other complications of pregnancy, childbirth and the puerperium (O10-O99)												
Black, both sexes												
2003	0.5	...	...	*	0.6	1.4	0.9	*	*	*	*	*
2002	0.4	...	...	*	0.7	1.1	0.6	*	*	*	*	*
2001	0.4	...	...	*	0.6	1.1	0.7	*	*	*	*	*
2000	0.3	...	...	*	0.5	1.0	0.6	*	*	*	*	*
1999	0.4	...	...	*	0.7	1.1	0.7	*	*	*	*	*
Black, male												
2003	...	...	...	...	...	...	...	...	...	...	...	...
2002	...	...	...	...	...	...	...	...	...	...	...	...
2001	...	...	...	...	...	...	...	...	...	...	...	...
2000	...	...	...	...	...	...	...	...	...	...	...	...
1999	...	...	...	...	...	...	...	...	...	...	...	...
Black, female												
2003	0.9	...	...	*	1.1	2.7	1.8	*	*	*	*	*
2002	0.7	...	...	*	1.4	2.2	1.2	*	*	*	*	*
2001	0.7	...	...	*	1.3	2.1	1.3	*	*	*	*	*
2000	0.6	...	...	*	1.0	2.0	1.1	*	*	*	*	*
1999	0.8	...	...	*	1.5	2.2	1.3	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
All races 3/, both sexes												
2003	4.9	356.0	0.5	*	*	*	*	*	*	*	*	*
2002	4.9	349.7	0.4	0.1	0.1	*	*	*	*	*	*	*
2001	4.9	340.5	0.5	0.1	*	*	*	*	*	*	*	*
2000	5.0	365.7	0.5	0.1	*	*	*	*	*	*	*	*
1999	5.1	371.0	0.6	0.1	*	*	*	*	*	*	*	*
All races, male												
2003	5.7	397.4	0.5	*	*	*	*	*	*	*	*	*
2002	5.7	385.7	0.5	*	*	*	*	*	*	*	*	*
2001	5.6	376.6	0.6	0.1	*	*	*	*	*	*	*	*
2000	5.8	403.9	0.7	0.1	*	*	*	*	*	*	*	*
1999	5.9	408.6	0.7	0.1	*	*	*	*	*	*	*	*
All races, female												
2003	4.2	312.8	0.5	*	*	*	*	*	*	*	*	*
2002	4.2	312.0	0.3	*	*	*	*	*	*	*	*	*
2001	4.2	302.6	0.4	*	*	*	*	*	*	*	*	*
2000	4.3	325.6	0.4	*	*	*	*	*	*	*	*	*
1999	4.4	331.7	0.5	0.1	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
White, both sexes												
2003	3.8	285.1	0.4	*	*	*	*	*	*	*	*	*
2002	3.7	277.2	0.3	0.1	*	*	*	*	*	*	*	*
2001	3.7	268.5	0.4	0.1	*	*	*	*	*	*	*	*
2000	3.7	283.7	0.4	0.1	*	*	*	*	*	*	*	*
1999	3.8	285.9	0.5	0.1	*	*	*	*	*	*	*	*
White, male												
2003	4.4	319.1	0.4	*	*	*	*	*	*	*	*	*
2002	4.3	309.4	*	*	*	*	*	*	*	*	*	*
2001	4.2	297.1	0.5	*	*	*	*	*	*	*	*	*
2000	4.3	312.4	0.5	0.1	*	*	*	*	*	*	*	*
1999	4.3	315.0	0.5	*	*	*	*	*	*	*	*	*
White, female												
2003	3.2	249.5	0.3	*	*	*	*	*	*	*	*	*
2002	3.2	243.5	*	*	*	*	*	*	*	*	*	*
2001	3.1	238.6	*	*	*	*	*	*	*	*	*	*
2000	3.2	253.5	*	*	*	*	*	*	*	*	*	*
1999	3.2	255.3	0.4	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Certain conditions originating in the perinatal period (P00-P96)												
Black, both sexes												
2003	12.8	734.4	1.2	*	*	*	*	*	*	*	*	*
2002	13.1	724.8	1.0	*	*	*	*	*	*	*	*	*
2001	13.0	738.1	0.9	*	*	*	*	*	*	*	*	*
2000	13.7	809.5	1.1	*	*	*	*	*	*	*	*	*
1999	14.2	844.6	1.2	*	*	*	*	*	*	*	*	*
Black, male												
2003	15.3	819.8	*	*	*	*	*	*	*	*	*	*
2002	15.2	783.2	*	*	*	*	*	*	*	*	*	*
2001	15.5	815.9	*	*	*	*	*	*	*	*	*	*
2000	16.3	901.3	*	*	*	*	*	*	*	*	*	*
1999	16.9	937.7	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	10.5	645.5	*	*	*	*	*	*	*	*	*	*
2002	11.2	663.8	*	*	*	*	*	*	*	*	*	*
2001	10.8	656.2	*	*	*	*	*	*	*	*	*	*
2000	11.3	714.6	*	*	*	*	*	*	*	*	*	*
1999	11.8	748.5	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
All races 3/, both sexes												
2003	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
2002	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8
All races, male												
2003	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
2002	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5
All races, female												
2003	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
2002	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
White, both sexes												
2003	3.5	137.6	3.2	0.9	1.1	1.1	1.3	1.7	2.5	2.3	3.7	6.9
2002	3.6	138.3	3.3	1.0	1.2	1.2	1.3	1.8	2.4	2.5	3.9	6.9
2001	3.6	135.0	3.2	0.9	1.2	1.2	1.3	1.7	2.4	2.5	3.8	7.4
2000	3.6	147.1	2.9	0.9	1.1	1.2	1.2	1.7	2.4	2.5	4.1	7.8
1999	3.6	141.3	3.2	0.9	1.1	1.2	1.2	1.8	2.4	2.6	4.0	8.0
White, male												
2003	3.7	143.2	3.1	0.9	1.2	1.2	1.4	1.7	2.6	2.3	3.4	6.9
2002	3.8	143.3	3.4	1.0	1.3	1.4	1.3	1.9	2.5	2.6	3.9	7.3
2001	3.8	143.0	3.2	1.0	1.4	1.3	1.4	1.7	2.4	2.9	4.2	7.5
2000	3.9	151.9	2.9	0.9	1.3	1.4	1.4	1.9	2.4	2.6	4.3	11.1
1999	3.8	146.5	3.5	0.9	1.3	1.2	1.3	1.9	2.6	2.5	4.1	8.6
White, female												
2003	3.4	131.8	3.2	0.9	0.9	1.0	1.2	1.6	2.4	2.2	3.9	6.9
2002	3.5	133.1	3.1	1.0	1.1	1.0	1.2	1.7	2.3	2.4	3.9	6.7
2001	3.4	126.7	3.3	0.7	1.0	1.1	1.2	1.6	2.5	2.2	3.6	7.3
2000	3.5	142.0	2.9	0.9	0.9	1.0	1.1	1.5	2.4	2.5	4.0	6.5
1999	3.5	135.7	2.9	1.0	0.9	1.2	1.2	1.7	2.3	2.6	4.0	7.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)												
Black, both sexes												
2003	4.5	157.5	4.7	1.1	1.5	1.4	1.2	1.6	2.2	2.3	3.1	5.8
2002	4.7	155.2	4.5	1.2	1.6	1.4	1.6	1.9	2.6	2.4	3.3	*
2001	4.6	158.4	5.3	1.1	1.9	1.3	1.6	1.7	1.9	2.2	3.0	6.1
2000	4.9	179.2	5.0	1.3	1.5	1.4	1.6	1.6	2.4	2.6	2.7	*
1999	4.8	169.3	5.5	1.6	1.4	1.1	1.3	1.9	1.9	2.8	4.2	7.7
Black, male												
2003	5.1	170.7	4.8	1.1	1.7	1.5	1.2	1.9	2.7	3.5	*	*
2002	5.1	151.7	4.6	1.4	1.9	1.7	1.7	2.3	3.4	3.0	*	*
2001	5.0	160.8	5.5	1.0	2.4	1.2	1.7	1.7	2.0	*	*	*
2000	5.4	181.4	5.5	1.5	2.0	1.8	1.9	2.1	*	3.8	*	*
1999	5.2	172.7	6.2	1.6	1.8	1.2	1.6	2.3	2.1	*	*	*
Black, female												
2003	3.9	143.8	4.6	1.0	1.2	1.3	1.2	1.4	1.8	*	*	*
2002	4.3	158.9	4.4	1.0	1.3	1.1	1.5	1.5	2.0	*	*	*
2001	4.3	155.9	5.1	1.2	1.4	1.3	1.4	1.7	1.8	2.5	*	*
2000	4.4	176.9	4.6	1.2	1.1	1.1	1.4	1.1	3.0	*	*	*
1999	4.3	165.8	4.8	1.6	1.1	1.0	1.2	1.5	1.8	3.1	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
All races 3/, both sexes												
2003	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
2002	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7
All races, male												
2003	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
2002	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3
All races, female												
2003	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
2002	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
White, both sexes												
2003	11.1	68.1	1.4	0.4	2.0	3.2	5.1	6.4	6.9	13.2	39.0	228.3
2002	10.5	71.0	1.3	0.3	1.8	3.2	5.1	5.9	7.0	12.7	38.6	211.6
2001	11.6	70.9	1.5	0.4	2.4	3.8	6.8	8.6	8.5	14.4	39.6	210.0
2000	11.4	77.2	1.5	0.4	2.1	3.7	6.5	7.8	8.6	15.6	40.2	212.2
1999	9.4	74.7	1.3	0.3	1.4	2.5	4.1	5.1	7.1	13.9	35.1	190.2
White, male												
2003	10.1	82.4	1.4	0.4	2.7	4.4	6.6	8.4	8.8	16.3	41.7	180.0
2002	9.8	82.5	1.4	0.3	2.4	4.3	6.5	7.7	9.2	15.3	42.9	178.2
2001	11.2	82.0	1.7	0.4	3.2	5.2	8.9	11.4	11.0	17.2	43.9	177.0
2000	11.3	89.4	1.7	0.4	2.9	5.1	8.9	10.7	11.2	19.2	45.5	186.8
1999	9.0	86.4	1.6	0.3	2.0	3.2	5.4	6.9	9.3	17.3	38.7	167.3
White, female												
2003	12.0	53.3	1.3	0.4	1.3	2.0	3.7	4.4	5.1	10.5	37.2	249.7
2002	11.2	59.0	1.2	0.2	1.1	2.0	3.6	4.2	5.0	10.5	35.7	226.1
2001	11.9	59.3	1.4	0.4	1.5	2.4	4.8	5.8	6.2	12.1	36.7	223.8
2000	11.5	64.3	1.3	0.3	1.3	2.3	4.1	4.9	6.1	12.7	36.7	222.5
1999	9.9	62.4	1.1	0.3	0.8	1.7	2.7	3.4	5.0	11.2	32.8	199.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)												
Black, both sexes												
2003	12.3	159.3	3.1	0.8	2.4	5.0	8.5	13.4	14.2	23.4	45.3	210.9
2002	12.3	162.3	2.6	0.7	2.5	5.1	8.5	13.7	13.7	20.1	49.6	196.0
2001	12.6	169.5	2.8	0.6	2.7	5.0	9.0	13.9	14.8	24.4	49.5	191.6
2000	13.7	197.1	2.3	0.7	3.2	5.8	9.2	14.7	17.4	28.9	55.5	186.4
1999	12.7	192.8	2.7	0.5	2.6	4.8	9.0	13.2	15.9	24.5	54.1	182.4
Black, male												
2003	13.5	170.7	3.4	0.8	3.0	6.7	11.1	17.6	20.3	31.4	52.5	193.9
2002	14.1	183.0	3.5	0.8	3.0	6.8	11.3	19.4	19.7	26.4	68.3	181.2
2001	13.9	187.4	3.5	0.8	3.7	6.0	11.1	18.0	23.3	30.4	51.8	184.9
2000	15.5	227.1	2.3	0.8	3.7	7.7	10.8	20.6	25.2	39.9	65.1	161.5
1999	14.6	217.6	2.8	0.6	3.5	6.8	11.5	19.4	23.1	32.5	66.1	184.0
Black, female												
2003	11.3	147.5	2.9	0.7	1.9	3.3	6.1	9.8	9.2	17.7	41.3	217.5
2002	10.6	140.8	1.8	0.7	2.0	3.6	6.0	8.8	8.9	15.7	39.0	201.7
2001	11.4	150.6	2.0	*	1.7	4.1	7.2	10.3	7.9	20.2	48.3	194.1
2000	12.0	166.0	2.3	*	2.6	4.1	7.7	9.7	11.0	21.2	50.1	195.7
1999	10.9	167.1	2.7	*	1.8	3.1	6.8	7.8	10.1	18.9	47.3	181.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
All races 3/, both sexes												
2003	69.3	29.7	4.0	2.3	4.9	8.4	18.9	38.9	67.1	144.4	411.5	1,508.4
2002	67.5	28.4	4.2	2.1	4.6	7.9	18.9	37.8	65.3	140.6	405.3	1,493.0
2001	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7
All races, male												
2003	57.3	34.8	4.2	2.6	5.6	9.3	22.3	47.2	76.5	155.6	415.1	1,258.1
2002	55.8	33.4	4.6	2.3	5.1	9.1	22.4	46.3	74.5	152.6	406.7	1,262.9
2001	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4
All races, female												
2003	81.0	24.5	3.8	1.9	4.2	7.4	15.6	31.0	58.4	135.0	409.1	1,619.1
2002	78.8	23.0	3.9	1.9	4.0	6.6	15.5	29.6	56.8	130.6	404.4	1,592.8
2001	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
White, both sexes												
2003	73.9	25.5	3.7	2.2	4.7	7.5	17.4	35.9	63.8	142.0	412.1	1,534.2
2002	71.8	25.2	3.9	2.1	4.2	7.1	17.3	34.8	62.0	137.8	405.2	1,517.6
2001	69.4	23.7	3.9	2.1	4.2	6.9	16.0	32.8	59.7	138.1	395.1	1,498.0
2000	67.0	24.9	3.5	2.0	4.3	6.9	16.0	32.0	58.6	132.7	386.9	1,464.6
1999	65.0	25.2	4.0	2.1	4.3	7.0	15.5	30.3	56.9	132.6	371.6	1,434.0
White, male												
2003	59.9	31.5	3.8	2.5	5.7	8.8	21.0	44.0	72.4	151.4	411.4	1,279.1
2002	58.2	30.8	4.1	2.2	5.0	8.4	21.3	43.2	70.5	148.1	403.4	1,274.7
2001	55.8	25.9	4.4	2.1	5.1	7.8	19.5	39.9	66.8	149.4	394.8	1,282.9
2000	54.1	26.8	3.9	2.2	5.2	8.3	19.5	39.3	66.0	143.4	387.1	1,274.8
1999	52.7	29.2	4.3	2.1	5.1	8.4	19.6	36.7	64.9	143.7	379.3	1,231.6
White, female												
2003	87.6	19.3	3.7	1.8	3.7	6.2	13.6	27.9	55.5	133.9	412.6	1,647.2
2002	85.2	19.3	3.6	1.9	3.4	5.6	13.3	26.6	54.0	129.0	406.3	1,623.1
2001	82.7	21.4	3.4	2.0	3.4	5.8	12.6	25.8	53.0	128.6	395.2	1,588.0
2000	79.6	22.9	3.2	1.8	3.4	5.4	12.5	24.7	51.6	123.6	386.7	1,541.7
1999	77.0	20.9	3.7	2.1	3.4	5.5	11.4	24.0	49.5	123.3	366.5	1,515.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14, N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)												
Black, both sexes												
2003	60.9	51.8	5.3	2.7	6.5	15.1	33.0	69.6	111.4	196.9	480.2	1,435.7
2002	59.7	45.2	5.9	2.6	6.8	14.5	33.7	68.4	108.1	194.2	477.7	1,425.4
2001	59.1	51.3	5.4	2.4	6.5	14.1	33.9	70.4	107.2	196.2	470.9	1,418.0
2000	58.1	49.6	5.9	2.8	6.5	15.0	34.2	69.3	107.4	191.9	467.7	1,388.9
1999	56.2	56.7	5.3	3.0	6.7	12.9	35.3	67.8	103.4	199.8	440.9	1,286.2
Black, male												
2003	55.7	55.3	6.1	2.9	5.9	15.0	35.4	82.2	132.3	233.6	544.4	1,224.1
2002	54.8	48.8	6.4	2.8	6.7	15.4	35.3	81.5	129.1	229.4	522.1	1,319.8
2001	54.6	58.4	5.1	2.4	6.6	14.9	38.2	84.2	126.4	229.8	524.1	1,313.9
2000	54.3	59.8	6.1	2.5	7.0	15.6	38.7	81.1	130.2	228.4	523.5	1,325.1
1999	52.8	64.7	5.4	3.5	7.1	13.4	39.3	80.3	121.7	226.2	505.1	1,291.3
Black, female												
2003	65.6	48.2	4.5	2.6	7.1	15.2	30.9	58.8	94.6	170.7	444.0	1,517.9
2002	64.1	41.5	5.4	2.3	6.9	13.7	32.2	57.2	91.3	169.2	452.5	1,465.9
2001	63.1	43.8	5.6	2.5	6.3	13.3	30.0	58.5	91.8	172.6	441.0	1,456.6
2000	61.6	39.0	5.7	3.0	6.0	14.4	30.2	59.1	89.1	166.3	436.2	1,412.7
1999	59.3	48.3	5.1	2.5	6.2	12.5	31.9	57.0	88.7	181.2	405.0	1,284.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
All races 3/, both sexes												
2003	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
2002	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4
All races, male												
2003	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
2002	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4
All races, female												
2003	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
2002	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
White, both sexes												
2003	39.5	20.0	10.6	6.1	40.1	33.0	39.0	38.5	32.3	43.8	104.0	291.1
2002	38.7	18.6	9.9	6.2	40.8	32.8	38.1	36.2	30.7	43.9	103.6	286.1
2001	37.0	20.6	11.0	6.4	38.2	30.9	35.8	33.1	29.4	42.4	102.7	285.5
2000	35.9	19.1	11.1	6.9	38.3	29.9	34.3	31.3	30.2	41.1	96.5	282.8
1999	36.0	18.9	11.5	7.1	37.2	30.0	33.8	30.5	29.2	43.5	100.9	290.4
White, male												
2003	51.3	22.5	12.8	7.6	57.4	49.3	55.6	54.9	45.8	58.3	134.4	365.7
2002	50.4	22.3	11.8	7.6	58.7	49.3	54.7	52.3	43.5	59.2	133.0	356.6
2001	48.4	22.1	13.3	7.8	55.8	46.7	52.3	48.6	41.1	56.3	134.3	368.1
2000	47.0	21.7	13.2	8.4	55.0	45.4	51.0	46.7	42.1	54.2	124.3	369.8
1999	46.9	21.8	13.2	8.8	52.8	45.4	50.0	45.4	41.5	58.0	131.8	383.6
White, female												
2003	28.0	17.4	8.3	4.7	21.7	15.9	22.1	22.3	19.5	31.4	83.5	258.0
2002	27.3	14.7	7.8	4.7	21.8	15.5	21.3	20.3	18.6	30.9	83.9	255.4
2001	25.9	18.9	8.5	5.0	19.5	14.4	19.2	17.8	18.3	30.7	81.6	251.0
2000	25.1	16.4	9.0	5.4	20.4	13.8	17.4	16.2	18.9	30.1	78.1	247.4
1999	25.3	15.8	9.6	5.4	20.5	14.1	17.4	15.8	17.7	31.3	80.6	253.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)												
Black, both sexes												
2003	32.4	43.5	13.4	7.7	25.6	29.9	38.6	48.4	42.3	50.4	82.7	160.2
2002	33.1	48.9	14.1	9.1	27.8	31.4	39.1	46.8	41.4	49.3	83.6	173.4
2001	33.5	43.1	13.6	9.4	28.7	30.2	40.0	47.7	42.0	51.0	88.5	183.6
2000	33.5	44.5	16.2	9.3	28.7	32.5	39.7	47.0	40.8	52.5	84.9	181.1
1999	35.2	41.6	17.7	10.5	29.5	32.2	40.0	47.8	44.7	57.8	103.3	207.6
Black, male												
2003	46.1	47.6	15.7	9.7	38.6	46.7	57.8	74.5	67.9	74.3	123.3	225.2
2002	47.9	51.1	18.5	11.6	41.8	50.6	58.4	72.2	68.7	76.9	129.7	260.5
2001	48.2	48.5	15.4	12.8	43.2	47.4	59.9	77.2	67.6	74.9	136.0	255.3
2000	49.0	47.7	18.8	12.0	43.0	51.8	62.0	75.4	68.8	79.4	132.6	275.6
1999	51.0	45.9	21.0	13.8	44.9	51.4	62.0	78.6	71.1	88.2	150.0	316.4
Black, female												
2003	19.9	39.3	11.0	5.7	12.5	14.5	21.6	26.1	21.7	33.4	59.8	134.9
2002	19.7	46.6	9.6	6.6	13.8	14.1	22.0	25.1	19.5	29.7	57.5	140.1
2001	20.1	37.5	11.9	6.0	14.2	14.7	22.4	22.3	21.4	34.2	61.8	157.0
2000	19.5	41.3	13.5	6.5	14.6	15.0	19.9	22.6	18.3	33.5	58.0	145.9
1999	20.8	37.3	14.3	7.1	14.3	14.9	20.6	21.4	23.5	36.4	77.1	166.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
All races 3/, both sexes												
2003	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
2002	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0
All races, male												
2003	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
2002	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5
All races, female												
2003	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
2002	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
White, both sexes												
2003	17.0	3.6	4.0	4.4	29.8	18.7	17.4	16.6	15.7	17.5	26.5	30.5
2002	17.3	2.7	3.9	4.2	31.3	19.1	17.3	16.4	15.6	18.4	27.5	29.6
2001	17.0	3.7	4.3	4.3	29.5	19.2	17.6	15.8	15.1	17.9	28.5	29.9
2000	17.0	4.1	4.3	4.7	30.0	18.5	17.1	15.6	16.0	17.9	27.5	32.2
1999	17.0	4.6	4.1	4.8	29.3	18.7	16.8	15.2	15.7	19.4	28.6	32.3
White, male												
2003	23.9	4.0	4.5	5.2	41.1	27.8	25.0	24.1	22.7	23.7	36.7	52.8
2002	24.2	3.1	4.5	5.0	43.4	28.3	24.7	24.0	22.3	24.9	37.7	53.8
2001	23.9	3.2	4.9	5.1	41.7	28.7	25.1	22.9	21.2	24.0	40.3	54.1
2000	23.6	4.5	4.7	5.5	41.6	27.4	24.7	22.6	22.0	23.7	38.1	61.6
1999	23.4	5.2	4.4	5.7	40.1	27.3	23.8	21.8	21.9	25.8	40.6	63.7
White, female												
2003	10.3	3.1	3.4	3.5	17.7	9.1	9.6	9.1	9.1	12.2	19.7	20.6
2002	10.5	2.2	3.4	3.4	18.3	9.4	9.8	8.9	9.4	12.8	20.6	19.1
2001	10.3	4.1	3.6	3.5	16.5	9.3	10.0	8.8	9.4	12.7	20.7	19.7
2000	10.5	3.7	3.8	4.0	17.7	9.3	9.4	8.7	10.3	12.9	20.5	20.3
1999	10.7	3.9	3.8	3.9	17.8	9.8	9.6	8.6	10.0	14.0	20.7	19.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Transport accidents (V01-V99,Y85)												
Black, both sexes												
2003	15.1	4.2	5.0	4.1	19.7	19.3	18.3	18.1	18.5	18.2	22.4	21.0
2002	15.4	5.3	5.0	4.6	21.2	20.4	18.2	17.8	17.1	17.1	21.7	18.4
2001	15.9	*	5.2	5.3	22.2	19.9	18.7	18.8	18.1	18.8	21.2	17.5
2000	16.0	6.5	5.6	4.7	21.7	21.4	18.6	19.3	17.4	19.6	20.0	21.6
1999	16.3	7.3	7.1	5.4	22.0	19.9	19.0	19.0	18.3	19.3	26.0	27.7
Black, male												
2003	22.6	*	5.1	5.1	29.3	30.6	28.4	29.0	29.1	26.6	37.5	42.7
2002	23.1	*	6.4	5.3	30.5	33.1	27.5	27.8	28.3	28.5	40.4	46.1
2001	23.8	*	5.7	6.9	32.5	31.6	28.9	30.2	29.6	27.4	36.2	35.2
2000	24.1	6.7	6.2	5.7	31.6	34.4	29.7	29.6	29.4	32.2	36.3	45.0
1999	24.2	*	8.0	6.8	32.0	31.8	29.0	30.3	28.9	29.1	43.4	68.0
Black, female												
2003	8.3	*	5.0	3.1	9.8	9.0	9.4	8.8	9.9	12.2	13.9	12.6
2002	8.3	*	3.4	3.9	11.8	9.0	9.9	9.2	8.1	9.0	11.0	*
2001	8.7	*	4.7	3.7	11.9	9.4	9.7	9.1	8.9	12.7	12.7	10.9
2000	8.5	*	5.0	3.6	11.9	9.7	8.7	10.5	7.6	10.8	10.7	12.9
1999	9.2	8.7	6.2	4.0	12.2	9.0	10.1	9.4	9.7	12.4	16.4	12.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
All races 3/, both sexes												
2003	15.4	3.6	3.9	4.0	26.6	17.1	15.7	14.9	14.2	16.2	24.9	28.8
2002	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4
All races, male												
2003	21.4	3.9	4.4	4.8	36.9	25.5	22.5	21.5	20.0	21.3	34.3	50.0
2002	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8
All races, female												
2003	9.5	3.3	3.5	3.2	15.8	8.5	8.9	8.5	8.8	11.9	18.7	19.5
2002	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
White, both sexes												
2003	15.8	3.5	3.8	4.1	28.6	17.5	15.9	14.9	13.9	16.1	25.2	29.5
2002	16.2	2.6	3.7	3.9	30.3	18.0	15.9	14.8	14.1	17.1	26.3	28.8
2001	15.7	3.5	3.9	4.0	28.4	17.8	15.9	14.1	13.6	16.3	27.0	28.8
2000	15.7	3.9	4.0	4.3	28.7	17.1	15.4	14.0	14.3	16.4	26.1	31.1
1999	15.5	4.5	3.8	4.3	27.8	17.0	14.9	13.3	13.9	17.7	26.8	30.7
White, male												
2003	22.0	3.9	4.3	4.9	39.2	25.9	22.6	21.3	19.6	21.3	34.4	50.7
2002	22.4	2.9	4.2	4.6	41.9	26.6	22.3	21.2	19.6	22.4	35.5	52.4
2001	21.9	3.1	4.4	4.7	40.0	26.4	22.4	20.1	18.6	21.3	37.8	52.6
2000	21.6	4.2	4.5	5.0	39.6	25.1	21.8	20.0	19.2	21.1	35.6	59.2
1999	21.1	5.0	4.1	5.0	37.8	24.6	20.9	18.8	18.7	23.0	37.9	60.3
White, female												
2003	9.8	3.0	3.2	3.2	17.2	8.6	9.0	8.5	8.6	11.7	19.0	20.1
2002	10.1	2.2	3.2	3.2	17.9	9.0	9.4	8.5	8.9	12.6	20.1	18.5
2001	9.8	3.8	3.4	3.2	16.0	8.8	9.3	8.1	8.8	12.0	19.8	18.9
2000	10.0	3.5	3.6	3.7	17.1	8.9	8.9	8.0	9.8	12.4	19.8	19.7
1999	10.0	3.9	3.5	3.6	17.1	9.1	8.8	7.9	9.3	13.3	19.6	18.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)												
Black, both sexes												
2003	14.2	4.2	4.9	3.9	18.8	18.4	17.1	16.5	17.0	17.0	21.0	20.7
2002	14.5	5.2	4.8	4.2	20.6	19.7	16.9	16.1	15.6	15.9	20.0	18.1
2001	14.8	*	5.0	4.9	21.5	18.9	17.2	16.9	16.4	17.3	19.7	16.6
2000	15.0	6.5	5.4	4.4	20.9	20.4	17.1	17.9	16.1	18.2	18.9	20.3
1999	15.1	7.1	6.4	5.0	21.1	18.6	17.4	17.1	16.5	17.9	24.1	25.5
Black, male												
2003	21.1	*	5.0	4.7	27.9	29.0	26.4	25.9	26.3	24.2	35.0	41.7
2002	21.5	*	6.3	4.9	29.6	31.7	25.3	24.7	25.2	25.5	36.1	45.0
2001	22.0	*	5.3	6.3	31.4	29.9	26.1	26.9	26.3	24.2	34.1	35.2
2000	22.5	6.7	5.8	5.4	30.2	32.6	27.2	27.1	27.1	29.4	34.5	43.8
1999	22.2	*	7.0	6.0	30.5	29.7	26.3	27.0	25.5	26.5	41.2	60.9
Black, female												
2003	8.0	*	4.8	3.1	9.6	8.7	8.9	8.4	9.6	11.9	13.1	12.6
2002	8.0	*	3.4	3.5	11.6	8.8	9.4	8.8	7.9	9.0	10.9	*
2001	8.2	*	4.6	3.3	11.7	8.9	9.2	8.3	8.3	12.5	11.6	9.7
2000	8.2	*	4.9	3.5	11.7	9.4	8.2	10.0	7.3	10.3	10.1	11.6
1999	8.7	8.7	5.8	3.9	11.8	8.6	9.6	8.7	9.2	11.9	14.5	12.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)												
All races 3/, both sexes												
2003	0.5	*	*	0.1	0.6	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002	0.4	*	*	0.2	0.4	0.3	0.5	0.5	0.4	0.4	0.5	*
2001	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3
All races, male												
2003	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.8	1.0	*
2002	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.7	1.0	*
2001	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9
All races, female												
2003	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	*	0.4	*
2002	0.1	*	*	0.1	0.1	0.2	0.2	0.1	*	*	*	*
2001	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9, V89.1,V89.3,V89.9)												
White, both sexes												
2003	0.5	*	*	0.2	0.7	0.5	0.5	0.5	0.5	0.4	0.6	0.5
2002	0.4	*	*	0.2	0.4	0.4	0.5	0.5	0.4	0.4	0.5	*
2001	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.6
2000	0.5	*	0.2	0.3	0.7	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999	0.8	*	0.3	0.3	1.0	0.8	0.9	0.8	0.8	0.8	1.3	1.3
White, male												
2003	0.7	*	*	0.2	1.1	0.8	0.8	0.9	0.8	0.7	1.0	*
2002	0.6	*	*	0.2	0.7	0.5	0.8	0.9	0.6	0.6	0.9	*
2001	0.7	*	*	0.2	0.8	0.8	0.9	0.9	0.8	0.9	1.1	*
2000	0.8	*	*	0.3	1.0	0.9	1.1	0.8	0.8	0.8	1.2	*
1999	1.1	*	*	0.4	1.4	1.2	1.4	1.2	1.2	1.1	1.8	2.8
White, female												
2003	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.3	*	0.4	*
2002	0.1	*	*	*	*	0.2	0.1	0.1	*	*	*	*
2001	0.2	*	*	0.1	0.2	*	0.2	0.3	*	0.3	0.5	*
2000	0.3	*	*	0.2	0.3	0.2	0.2	0.3	*	0.3	0.4	*
1999	0.4	*	*	0.2	0.5	0.4	0.4	0.4	0.4	0.4	0.9	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)												
Black, both sexes												
2003	0.5	*	*	*	0.4	0.5	0.7	1.0	*	*	*	*
2002	0.5	*	*	*	*	*	0.7	0.9	*	*	*	*
2001	0.6	*	*	0.3	0.4	0.5	1.0	0.9	0.9	*	*	*
2000	0.6	*	*	*	0.5	0.6	0.9	0.8	*	*	*	*
1999	0.7	*	*	0.3	0.6	0.8	0.9	1.0	*	*	*	*
Black, male												
2003	0.9	*	*	*	0.7	1.0	1.0	1.9	*	*	*	*
2002	0.8	*	*	*	*	*	1.2	1.9	*	*	*	*
2001	0.9	*	*	*	*	*	1.8	1.5	*	*	*	*
2000	0.9	*	*	*	0.9	1.0	1.6	1.4	*	*	*	*
1999	1.1	*	*	0.6	1.0	1.3	1.5	1.8	*	*	*	*
Black, female												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.3	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
All races 3/, both sexes												
2003	0.7	*	*	0.2	0.5	0.6	0.9	1.1	1.2	0.9	0.7	*
2002	0.7	*	*	0.1	0.5	0.6	0.9	1.0	1.1	0.9	0.8	0.4
2001	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*
1999	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*
All races, male												
2003	1.1	*	*	0.2	0.8	0.9	1.5	1.8	2.2	1.6	1.3	*
2002	1.1	*	*	0.1	0.8	1.1	1.5	1.8	1.9	1.9	1.5	*
2001	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*
All races, female												
2003	0.2	*	*	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	*	0.3	*
2001	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
White, both sexes												
2003	0.7	*	*	0.2	0.5	0.7	1.0	1.2	1.3	0.9	0.7	*
2002	0.7	*	*	0.1	0.5	0.7	0.9	1.1	1.2	0.9	0.7	*
2001	0.8	*	0.2	0.1	0.6	0.9	1.1	1.2	1.1	1.0	0.8	*
2000	0.7	*	*	0.1	0.6	0.9	1.0	1.0	1.1	1.0	0.7	*
1999	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.5	*
White, male												
2003	1.2	*	*	0.2	0.8	1.0	1.6	2.0	2.3	1.7	1.3	*
2002	1.2	*	*	0.1	0.8	1.2	1.6	1.9	2.0	1.9	1.4	*
2001	1.3	*	*	0.2	0.9	1.5	1.8	1.9	1.8	1.8	1.4	*
2000	1.3	*	*	0.2	1.0	1.4	1.8	1.8	1.9	1.8	1.3	*
1999	1.2	*	*	0.3	0.9	1.5	1.5	1.8	2.0	1.6	0.9	*
White, female												
2003	0.3	*	*	0.2	0.2	0.3	0.4	0.4	0.3	0.3	*	*
2002	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	*	0.3	*
2001	0.3	*	*	*	0.3	0.4	0.5	0.4	0.4	0.3	0.4	*
2000	0.2	*	*	*	0.3	0.2	0.3	0.3	0.4	0.3	*	*
1999	0.3	*	*	0.1	0.3	0.4	0.4	0.3	0.3	0.3	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)												
Black, both sexes												
2003	0.4	*	*	*	0.4	*	0.5	0.7	0.7	*	*	*
2002	0.4	*	*	*	*	0.4	0.6	0.8	0.8	*	*	*
2001	0.5	*	*	*	*	0.6	0.5	1.0	0.9	*	*	*
2000	0.4	*	*	*	*	0.5	0.5	0.6	*	*	*	*
1999	0.5	*	*	*	0.3	0.5	0.7	0.8	1.1	*	*	*
Black, male												
2003	0.7	*	*	*	0.8	*	1.0	1.2	*	*	*	*
2002	0.8	*	*	*	*	*	1.0	1.3	1.8	*	*	*
2001	0.8	*	*	*	*	0.9	1.0	1.8	2.0	*	*	*
2000	0.7	*	*	*	*	0.9	1.0	1.1	*	*	*	*
1999	0.8	*	*	*	*	0.9	1.2	1.4	2.3	*	*	*
Black, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.2	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
All races 3/, both sexes												
2003	21.0	19.9	6.8	2.1	9.3	13.2	20.7	22.3	16.9	26.6	75.6	249.2
2002	20.2	20.3	6.4	2.4	8.9	12.7	20.1	20.2	15.7	25.9	74.4	246.7
2001	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3
All races, male												
2003	26.0	22.4	8.5	2.8	14.7	19.9	29.4	31.6	24.1	35.3	96.3	300.5
2002	25.2	23.3	8.1	3.2	14.2	19.6	29.1	29.3	22.7	35.3	93.9	293.6
2001	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9
All races, female												
2003	16.2	17.3	5.0	1.4	3.6	6.3	12.0	13.3	10.3	19.3	61.8	226.5
2002	15.4	17.2	4.7	1.5	3.2	5.7	11.1	11.5	9.2	18.1	61.4	226.3
2001	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
White, both sexes												
2003	22.5	16.4	6.6	1.8	10.3	14.3	21.6	21.9	16.6	26.3	77.5	260.6
2002	21.4	15.9	5.9	1.9	9.6	13.7	20.8	19.8	15.1	25.5	76.1	256.5
2001	20.0	16.9	6.7	2.1	8.7	11.7	18.2	17.3	14.3	24.6	74.2	255.7
2000	18.9	15.1	6.9	2.2	8.2	11.4	17.2	15.8	14.2	23.2	69.0	250.6
1999	19.0	14.3	7.3	2.3	7.9	11.3	17.0	15.3	13.5	24.1	72.3	258.1
White, male												
2003	27.4	18.6	8.3	2.3	16.3	21.5	30.6	30.7	23.1	34.6	97.8	312.9
2002	26.2	19.2	7.4	2.6	15.3	20.9	29.9	28.3	21.3	34.3	95.2	302.8
2001	24.5	18.9	8.4	2.7	14.1	18.0	27.2	25.7	19.9	32.3	94.0	314.0
2000	23.4	17.3	8.4	2.9	13.4	18.0	26.2	24.1	20.2	30.4	86.2	308.2
1999	23.5	16.5	8.8	3.0	12.8	18.1	26.1	23.5	19.6	32.3	91.3	319.8
White, female												
2003	17.7	14.2	4.9	1.1	3.9	6.8	12.4	13.2	10.4	19.2	63.8	237.4
2002	16.8	12.5	4.4	1.2	3.5	6.1	11.5	11.3	9.2	18.0	63.3	236.3
2001	15.6	14.8	4.8	1.5	3.0	5.0	9.2	9.0	9.0	18.0	60.9	231.2
2000	14.6	12.8	5.2	1.4	2.7	4.5	8.0	7.5	8.6	17.2	57.7	227.2
1999	14.6	11.8	5.8	1.5	2.7	4.3	7.8	7.2	7.7	17.3	59.9	233.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nontransport accidents (W00-X59,Y86)												
Black, both sexes												
2003	17.2	39.3	8.4	3.6	5.9	10.6	20.3	30.3	23.9	32.2	60.3	139.2
2002	17.8	43.6	9.2	4.5	6.6	11.0	20.9	29.0	24.3	32.2	61.9	155.0
2001	17.6	40.2	8.4	4.1	6.5	10.3	21.3	28.9	23.8	32.3	67.3	166.1
2000	17.6	38.0	10.6	4.6	7.0	11.0	21.1	27.7	23.5	32.8	64.9	159.5
1999	18.9	34.4	10.6	5.1	7.5	12.3	21.0	28.7	26.4	38.5	77.2	179.8
Black, male												
2003	23.5	44.0	10.6	4.6	9.2	16.2	29.5	45.5	38.7	47.7	85.7	182.5
2002	24.8	45.9	12.1	6.3	11.2	17.5	30.9	44.4	40.4	48.4	89.2	214.4
2001	24.4	46.4	9.7	5.8	10.7	15.8	31.0	47.1	37.9	47.5	99.8	220.1
2000	24.9	40.9	12.6	6.3	11.4	17.4	32.3	45.8	39.3	47.3	96.3	230.6
1999	26.8	40.0	13.0	7.0	12.9	19.5	32.9	48.3	42.2	59.1	106.7	248.4
Black, female												
2003	11.6	34.3	6.0	2.6	2.6	5.5	12.2	17.3	11.9	21.2	45.9	122.3
2002	11.4	41.2	6.2	2.7	2.0	5.1	12.0	15.9	11.4	20.7	46.4	132.3
2001	11.5	33.7	7.1	2.4	2.4	5.3	12.6	13.2	12.5	21.6	49.1	146.1
2000	11.0	35.1	8.5	2.9	2.7	5.3	11.3	12.1	10.6	22.7	47.2	133.1
1999	11.7	28.5	8.1	3.1	2.0	5.9	10.5	12.0	13.8	24.0	60.8	154.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
All races 3/, both sexes												
2003	5.9	*	0.3	0.1	0.6	0.7	1.4	2.6	4.4	11.2	40.8	135.9
2002	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2
All races, male												
2003	6.2	*	0.5	0.1	0.9	1.2	2.4	4.0	6.2	14.5	51.6	165.1
2002	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5
All races, female												
2003	5.6	*	*	*	0.2	0.2	0.5	1.2	2.7	8.4	33.5	122.9
2002	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
White, both sexes												
2003	6.7	*	0.3	0.1	0.6	0.8	1.5	2.5	4.4	11.5	42.8	144.4
2002	6.4	*	0.2	0.1	0.7	0.9	1.5	2.6	4.1	11.1	40.4	138.0
2001	6.0	*	0.2	0.2	0.8	0.9	1.5	2.6	4.0	10.4	37.0	128.0
2000	5.3	*	0.2	0.1	0.7	0.8	1.4	2.3	3.9	9.2	32.6	119.1
1999	5.2	*	0.4	0.1	0.7	0.9	1.5	2.2	3.5	9.1	32.9	115.9
White, male												
2003	6.9	*	0.5	0.2	1.0	1.3	2.4	3.9	6.2	14.7	53.8	175.1
2002	6.6	*	*	0.1	1.1	1.5	2.4	3.9	6.0	14.8	50.2	166.4
2001	6.4	*	0.3	0.2	1.3	1.6	2.5	4.2	5.8	13.5	47.4	162.3
2000	5.7	*	*	0.2	1.2	1.5	2.2	3.7	5.7	12.1	41.4	152.8
1999	5.6	*	0.5	0.2	1.2	1.5	2.4	3.5	5.4	12.3	41.9	149.1
White, female												
2003	6.5	*	*	*	0.2	0.3	0.5	1.1	2.8	8.8	35.4	130.8
2002	6.2	*	*	*	0.2	0.2	0.7	1.2	2.4	8.0	33.9	125.6
2001	5.5	*	*	*	0.1	0.2	0.4	1.1	2.2	7.7	30.2	113.6
2000	5.0	*	*	*	0.2	0.2	0.5	0.9	2.1	6.7	26.8	105.4
1999	4.9	*	*	*	0.2	0.2	0.6	0.8	1.8	6.4	27.1	102.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Falls (W00-W19)												
Black, both sexes												
2003	2.4	*	*	*	*	0.5	1.3	3.3	4.8	8.9	19.4	44.1
2002	2.2	*	*	*	*	0.4	1.3	2.9	4.2	8.2	16.1	49.3
2001	2.3	*	*	*	*	0.5	1.6	2.9	4.4	8.7	18.5	44.5
2000	2.1	*	*	*	*	0.4	1.3	2.7	4.7	8.3	14.9	38.8
1999	2.4	*	*	*	0.4	0.8	1.3	3.4	5.5	9.2	17.1	45.3
Black, male												
2003	3.0	*	*	*	*	0.8	2.2	5.2	7.1	12.5	24.8	56.3
2002	3.0	*	*	*	*	0.8	2.2	4.8	6.7	11.8	24.4	77.2
2001	3.1	*	*	*	*	0.8	2.6	5.4	7.0	12.6	28.5	68.1
2000	2.8	*	*	*	*	*	2.1	4.7	8.2	11.4	21.2	61.1
1999	3.3	*	*	*	*	1.3	2.5	5.8	8.8	14.7	24.3	58.6
Black, female												
2003	1.9	*	*	*	*	*	*	1.6	2.9	6.3	16.3	39.3
2002	1.5	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.6
2001	1.6	*	*	*	*	*	*	*	2.3	5.9	12.9	35.8
2000	1.4	*	*	*	*	*	*	1.0	2.0	6.1	11.4	30.5
1999	1.7	*	*	*	*	*	*	1.3	2.9	5.3	13.1	40.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
All races 3/, both sexes												
2003	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
All races, male												
2003	0.5	*	*	0.2	0.9	0.5	0.5	0.4	0.4	0.4	0.6	*
2002	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*
All races, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
White, both sexes												
2003	0.2	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002	0.3	*	*	0.1	0.4	0.3	0.3	0.2	0.2	0.2	0.2	*
2001	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*
White, male												
2003	0.4	*	*	0.2	0.8	0.5	0.5	0.4	0.5	0.4	0.6	*
2002	0.4	*	*	0.2	0.8	0.5	0.4	0.4	0.4	0.4	0.5	*
2001	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999	0.5	*	*	0.3	1.0	0.6	0.5	0.4	0.4	0.4	*	*
White, female												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000	0.1	*	*	*	*	*	0.1	*	*	*	*	*
1999	0.1	*	*	*	*	*	0.1	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental discharge of firearms (W32-W34)												
Black, both sexes												
2003	0.3	*	*	*	0.8	0.6	*	*	*	*	*	*
2002	0.4	*	*	*	1.0	0.7	0.3	*	*	*	*	*
2001	0.3	*	*	*	0.8	*	*	*	*	*	*	*
2000	0.4	*	*	*	0.9	0.6	*	*	*	*	*	*
1999	0.4	*	*	*	1.1	0.4	*	*	*	*	*	*
Black, male												
2003	0.6	*	*	*	1.5	1.3	*	*	*	*	*	*
2002	0.8	*	*	*	1.9	1.5	*	*	*	*	*	*
2001	0.6	*	*	*	1.5	*	*	*	*	*	*	*
2000	0.7	*	*	*	1.6	1.1	*	*	*	*	*	*
1999	0.8	*	*	*	2.1	0.8	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	*	*	*	*	*	*	*	*
1999	0.1	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
All races 3/, both sexes												
2003	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
2002	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6
All races, male												
2003	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
2002	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3
All races, female												
2003	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
2002	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
White, both sexes												
2003	1.1	1.4	3.1	0.5	1.3	0.9	1.0	1.0	0.9	1.1	1.3	1.6
2002	1.1	1.6	3.0	0.5	1.4	1.0	1.1	0.9	0.8	0.9	1.3	1.6
2001	1.1	1.7	3.3	0.5	1.3	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000	1.2	1.6	3.5	0.7	1.5	1.0	1.0	0.9	0.9	1.0	1.2	1.5
1999	1.2	1.5	3.5	0.6	1.5	1.0	1.1	1.0	0.9	1.0	1.3	1.6
White, male												
2003	1.8	1.5	4.0	0.7	2.3	1.5	1.7	1.5	1.5	1.8	2.1	3.3
2002	1.8	1.9	4.0	0.7	2.5	1.8	1.9	1.4	1.2	1.5	2.1	3.7
2001	1.7	1.6	4.2	0.7	2.4	1.4	1.6	1.5	1.2	1.5	2.0	3.5
2000	1.9	1.8	4.4	0.9	2.6	1.7	1.7	1.5	1.6	1.6	1.8	2.7
1999	1.9	1.7	4.3	0.8	2.7	1.8	1.8	1.7	1.4	1.4	2.2	2.4
White, female												
2003	0.5	1.3	2.2	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.7	0.9
2002	0.4	*	2.0	0.3	0.2	0.3	0.4	0.3	0.4	0.3	0.7	0.7
2001	0.5	1.8	2.4	0.4	0.3	0.3	0.4	0.3	0.4	0.3	0.7	1.0
2000	0.5	*	2.5	0.4	0.3	0.2	0.3	0.3	0.3	0.5	0.8	1.0
1999	0.5	1.4	2.6	0.3	0.3	0.2	0.4	0.3	0.4	0.5	0.6	1.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental drowning and submersion (W65-W74)												
Black, both sexes												
2003	1.3	*	2.1	1.3	1.6	1.1	0.9	1.0	1.2	*	*	*
2002	1.6	*	2.5	2.0	2.1	1.4	1.3	1.2	0.8	*	*	*
2001	1.6	*	1.9	2.0	2.2	1.2	1.2	1.1	1.3	*	*	*
2000	1.6	4.2	2.2	2.0	2.3	1.1	1.4	1.2	*	*	*	*
1999	1.8	3.5	2.4	2.4	2.4	1.6	1.1	0.9	1.2	1.3	*	*
Black, male												
2003	2.2	*	3.0	1.9	3.0	2.1	1.8	2.0	2.4	*	*	*
2002	2.8	*	3.4	3.1	4.0	2.5	2.4	2.2	*	*	*	*
2001	2.7	*	2.2	3.0	4.2	2.4	2.2	2.0	2.3	*	*	*
2000	2.7	*	2.8	3.0	4.3	2.0	2.6	2.3	*	*	*	*
1999	3.1	*	3.2	3.7	4.5	3.1	2.1	1.9	2.2	*	*	*
Black, female												
2003	0.4	*	*	0.7	*	*	*	*	*	*	*	*
2002	0.5	*	*	0.8	*	*	*	*	*	*	*	*
2001	0.5	*	1.6	0.9	*	*	*	*	*	*	*	*
2000	0.6	*	1.7	1.0	*	*	*	*	*	*	*	*
1999	0.5	*	1.6	1.0	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
All races 3/, both sexes												
2003	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
2002	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9
All races, male												
2003	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
2002	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3
All races, female												
2003	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
2002	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
White, both sexes												
2003	1.1	0.7	1.2	0.4	0.5	0.6	0.9	1.2	1.3	2.0	3.1	4.7
2002	1.0	*	1.0	0.4	0.4	0.6	0.8	1.0	1.1	1.8	3.0	4.0
2001	1.0	1.1	1.2	0.5	0.5	0.6	0.9	0.9	1.3	1.9	2.8	4.8
2000	1.0	0.8	1.3	0.5	0.5	0.5	0.8	1.0	1.3	1.8	3.1	5.1
1999	1.0	0.9	1.5	0.5	0.5	0.7	0.8	0.9	1.2	1.8	3.3	4.7
White, male												
2003	1.3	*	1.4	0.4	0.6	0.7	1.2	1.5	1.7	2.6	3.6	6.7
2002	1.2	*	1.1	0.6	0.6	0.8	1.1	1.3	1.5	2.5	4.1	5.8
2001	1.3	*	1.6	0.6	0.7	0.7	1.2	1.2	1.6	2.2	3.4	7.0
2000	1.3	*	1.6	0.5	0.6	0.7	1.1	1.5	1.7	2.3	4.1	8.2
1999	1.3	*	1.7	0.5	0.7	1.0	1.2	1.2	1.5	2.3	4.2	8.0
White, female												
2003	0.9	*	1.0	0.4	0.4	0.5	0.6	0.8	0.9	1.5	2.7	3.8
2002	0.7	*	1.0	0.3	0.3	0.4	0.5	0.7	0.7	1.3	2.2	3.2
2001	0.8	*	0.8	0.4	0.3	0.4	0.5	0.6	0.9	1.6	2.5	3.8
2000	0.8	*	1.1	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.4	3.9
1999	0.8	*	1.3	0.5	0.3	0.4	0.5	0.5	0.8	1.4	2.6	3.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)												
Black, both sexes												
2003	2.0	*	3.0	1.4	0.6	0.9	1.2	2.6	2.9	5.3	10.2	14.9
2002	2.1	*	3.5	1.4	0.8	1.1	1.0	2.1	3.3	5.0	8.6	15.7
2001	2.3	*	3.3	1.3	0.7	1.2	1.8	2.4	3.5	6.1	9.7	17.8
2000	2.5	*	4.8	1.5	0.7	1.2	1.7	2.4	4.2	5.9	13.2	20.3
1999	2.5	*	4.5	1.5	0.7	1.0	1.8	2.1	3.4	6.9	13.2	21.7
Black, male												
2003	2.4	*	3.2	1.5	*	1.1	1.4	3.3	4.3	7.5	16.1	*
2002	2.6	*	4.1	1.6	1.0	1.4	1.7	3.0	4.7	7.8	10.9	25.7
2001	2.9	*	3.6	1.6	0.8	1.4	2.7	3.9	5.0	9.0	12.9	27.2
2000	3.2	*	5.6	1.7	0.9	1.5	2.4	3.5	6.3	8.8	18.5	27.7
1999	3.0	*	5.0	1.7	0.8	1.2	2.7	3.2	4.1	10.1	18.1	29.3
Black, female												
2003	1.7	*	2.7	1.3	0.6	*	1.0	2.0	1.7	3.7	6.9	13.4
2002	1.6	*	2.8	1.3	*	0.9	*	1.4	2.2	3.0	7.3	11.9
2001	1.7	*	2.9	1.0	*	1.1	1.0	1.1	2.4	4.1	7.9	14.3
2000	2.0	*	4.1	1.3	*	0.9	1.1	1.4	2.4	3.9	10.2	17.6
1999	2.0	*	3.9	1.3	*	0.8	1.0	1.3	2.7	4.6	10.5	18.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
All races 3/, both sexes												
2003	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
2002	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4
All races, male												
2003	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
2002	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4
All races, female												
2003	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
2002	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
White, both sexes												
2003	7.1	*	0.3	0.1	5.8	9.7	14.8	13.3	4.8	2.0	2.3	3.5
2002	6.3	*	*	0.1	4.9	8.7	14.0	11.3	3.8	2.1	2.3	3.9
2001	5.0	*	0.2	0.1	3.9	6.9	11.4	8.4	2.8	1.5	2.1	3.3
2000	4.6	*	*	0.1	3.4	6.4	10.4	7.5	2.6	1.4	2.1	4.1
1999	4.3	*	0.2	0.1	2.8	6.1	10.0	7.1	2.5	1.5	2.2	3.3
White, male												
2003	9.7	*	*	0.1	8.8	13.9	19.8	17.6	5.8	2.2	2.8	3.7
2002	8.8	*	*	0.2	7.4	12.7	19.2	15.2	4.5	2.3	2.6	5.2
2001	7.2	*	*	*	6.0	10.2	16.2	11.9	3.4	1.9	2.5	3.7
2000	6.7	*	*	*	5.3	9.7	15.2	10.9	2.9	1.6	2.1	5.9
1999	6.4	*	*	*	4.2	9.5	14.9	10.5	3.2	1.9	2.5	4.5
White, female												
2003	4.5	*	*	*	2.6	5.2	9.8	9.0	3.8	1.8	1.9	3.4
2002	3.9	*	*	*	2.2	4.5	8.7	7.5	3.0	1.8	2.1	3.4
2001	2.9	*	*	0.1	1.6	3.5	6.6	5.0	2.3	1.2	1.8	3.1
2000	2.5	*	*	*	1.3	3.0	5.5	4.1	2.3	1.3	2.0	3.4
1999	2.3	*	*	*	1.2	2.6	5.1	3.8	1.8	1.2	2.1	2.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)												
Black, both sexes												
2003	6.3	*	*	*	1.4	5.7	13.5	17.9	7.6	4.1	2.0	*
2002	6.3	*	*	*	1.4	5.6	13.8	17.9	8.4	2.7	2.8	*
2001	5.8	*	*	*	1.6	5.2	13.1	16.7	6.5	2.5	3.6	6.4
2000	5.5	*	*	*	1.3	5.6	13.0	15.5	5.5	2.6	*	8.1
1999	5.7	*	*	*	1.6	6.1	13.1	15.5	6.0	3.0	*	*
Black, male												
2003	8.8	*	*	*	2.0	8.0	18.6	26.2	12.8	7.6	*	*
2002	9.1	*	*	*	2.2	8.5	19.3	26.5	15.0	5.0	*	*
2001	8.4	*	*	*	2.3	7.5	17.9	26.6	11.3	3.9	5.9	*
2000	8.2	*	*	*	1.7	8.4	18.7	24.9	9.5	4.5	*	*
1999	8.7	*	*	*	2.7	9.1	19.4	25.9	11.0	4.4	*	*
Black, female												
2003	3.9	*	*	*	0.7	3.6	8.9	10.8	3.3	*	*	*
2002	3.7	*	*	*	0.7	3.0	8.9	10.5	3.2	*	*	*
2001	3.5	*	*	*	0.9	3.1	9.0	8.2	2.6	*	*	*
2000	3.1	*	*	*	1.0	3.0	7.8	7.4	2.3	*	*	*
1999	3.0	*	*	*	*	3.4	7.6	6.5	2.0	2.1	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
All races 3/, both sexes												
2003	5.9	16.9	1.8	0.5	1.6	2.0	3.0	3.9	5.1	9.6	27.5	102.5
2002	6.0	16.8	1.6	0.7	1.6	2.1	3.0	3.7	5.2	9.8	28.8	105.6
2001	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9
All races, male												
2003	6.9	19.5	2.2	0.8	2.6	3.4	4.8	6.0	7.8	13.1	34.9	120.2
2002	6.9	18.8	2.0	0.9	2.6	3.4	4.8	6.0	7.9	13.4	36.0	118.6
2001	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7
All races, female												
2003	4.9	14.1	1.3	0.3	0.5	0.7	1.2	1.9	2.6	6.7	22.6	94.7
2002	5.0	14.6	1.2	0.4	0.5	0.6	1.2	1.5	2.7	6.8	24.0	100.0
2001	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
White, both sexes												
2003	6.3	13.6	1.7	0.5	1.7	2.2	3.1	3.8	4.9	9.4	27.8	106.3
2002	6.3	12.9	1.4	0.7	1.7	2.2	3.1	3.7	5.1	9.4	28.8	108.8
2001	6.6	13.4	1.7	0.6	1.8	2.1	3.1	4.1	5.1	9.7	30.6	117.7
2000	6.6	12.2	1.6	0.7	1.7	2.4	3.2	3.8	5.4	9.5	29.8	120.5
1999	6.9	11.2	1.8	0.8	1.9	2.3	3.3	3.9	5.2	10.6	32.5	132.4
White, male												
2003	7.3	15.7	2.2	0.8	2.8	3.6	4.9	5.8	7.4	12.9	34.9	123.9
2002	7.3	15.3	1.8	0.9	2.9	3.7	5.0	6.0	7.6	12.7	35.7	121.2
2001	7.5	15.2	2.0	0.9	2.9	3.5	5.1	6.4	7.4	12.9	38.0	137.0
2000	7.4	14.0	2.0	0.9	2.9	4.0	5.4	6.2	7.9	12.5	36.2	137.6
1999	7.8	12.9	2.1	1.1	3.1	3.8	5.4	6.2	7.7	14.0	40.1	155.2
White, female												
2003	5.3	11.4	1.2	0.3	0.5	0.7	1.2	1.8	2.5	6.5	23.0	98.5
2002	5.4	10.5	1.1	0.4	0.5	0.7	1.1	1.5	2.6	6.5	24.2	103.4
2001	5.8	11.5	1.4	0.4	0.6	0.6	1.1	1.9	3.0	7.1	25.7	109.7
2000	5.7	10.3	1.2	0.4	0.5	0.6	1.0	1.5	3.0	7.1	25.6	113.6
1999	6.1	9.5	1.4	0.5	0.6	0.7	1.2	1.6	2.8	7.7	27.4	123.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)												
Black, both sexes												
2003	4.9	34.9	2.0	0.5	1.3	1.8	3.2	5.4	7.3	12.8	27.6	73.2
2002	5.2	37.5	2.4	0.8	1.1	1.7	3.1	4.7	7.3	15.2	32.3	83.1
2001	5.4	34.2	2.5	0.6	1.0	1.9	3.2	5.6	7.9	14.1	33.8	96.1
2000	5.4	30.6	2.3	0.7	1.5	2.2	3.5	5.6	8.2	14.9	33.7	90.1
1999	6.1	27.9	2.5	0.7	1.3	2.4	3.4	6.6	10.0	17.9	42.9	105.5
Black, male												
2003	6.4	40.5	2.8	0.7	1.9	2.9	5.1	8.5	11.9	18.3	39.8	91.8
2002	6.6	38.9	3.3	1.1	1.7	2.9	4.7	7.4	12.1	21.6	45.6	106.1
2001	6.8	40.4	2.8	0.9	1.5	3.2	5.1	9.0	11.9	19.8	48.8	114.6
2000	7.3	32.0	2.5	1.0	2.5	3.7	6.0	10.1	13.8	20.3	53.0	124.5
1999	8.0	33.2	3.4	0.9	2.3	4.0	5.7	11.1	15.7	26.9	57.2	153.5
Black, female												
2003	3.6	29.1	*	*	0.7	0.7	1.5	2.8	3.6	8.8	20.7	66.0
2002	4.0	36.0	*	*	*	*	1.7	2.3	3.5	10.7	24.7	74.3
2001	4.1	27.7	2.2	*	*	0.7	1.6	2.7	4.6	10.0	25.3	89.3
2000	3.8	29.1	2.0	*	*	0.9	1.3	1.8	3.6	11.1	22.8	77.3
1999	4.4	22.5	1.6	*	*	1.0	1.4	2.7	5.4	11.6	34.9	87.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
All races 3/, both sexes												
2003	10.8	...	...	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
2002	11.0	...	...	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 4/	10.8	...	...	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3
All races, male												
2003	17.6	...	...	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
2002	17.9	...	...	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 4/	17.6	...	...	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000	17.1	...	...	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999	17.1	...	...	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1
All races, female												
2003	4.3	...	...	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
2002	4.3	...	...	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 4/	4.1	...	...	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000	4.0	...	...	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999	4.0	...	...	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
White, both sexes												
2003	12.1	...	...	0.6	10.2	13.7	16.6	17.9	15.2	13.8	17.6	18.1
2002	12.2	...	...	0.7	10.7	13.6	17.0	17.6	15.0	14.7	19.0	19.1
2001 4/	11.9	...	...	0.7	10.6	13.8	16.4	16.9	14.3	14.3	18.6	18.6
2000	11.5	...	...	0.8	10.7	12.8	16.1	15.9	13.3	13.5	18.8	21.0
1999	11.5	...	...	0.6	10.5	13.5	15.8	15.3	13.2	14.4	19.3	20.7
White, male												
2003	19.5	...	...	0.9	16.9	21.8	25.7	27.3	24.3	25.2	37.5	51.4
2002	19.9	...	...	1.0	17.7	21.6	26.2	27.0	24.3	26.8	40.6	53.9
2001 4/	19.5	...	...	1.0	17.6	22.4	25.4	26.0	22.8	26.3	40.2	55.0
2000	18.8	...	...	1.2	17.9	20.7	24.9	24.5	21.2	24.3	41.1	61.6
1999	18.7	...	...	1.0	17.5	21.8	24.4	23.4	21.2	26.4	41.2	61.5
White, female												
2003	4.7	...	...	0.3	3.1	5.1	7.5	8.7	6.5	4.1	4.2	3.3
2002	4.8	...	...	0.4	3.1	5.2	7.8	8.4	6.2	4.4	4.4	4.0
2001 4/	4.6	...	...	0.3	3.1	4.9	7.3	8.0	6.4	4.2	4.3	3.4
2000	4.4	...	...	0.3	3.1	4.6	7.2	7.5	6.0	4.4	4.1	4.5
1999	4.5	...	...	0.3	3.2	4.9	7.2	7.4	5.7	4.4	4.9	4.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)												
Black, both sexes												
2003	5.1	...	...	0.6	7.1	9.3	7.2	5.6	5.0	4.5	4.5	*
2002	5.1	...	...	0.5	6.5	9.3	7.6	5.9	4.8	4.9	5.5	5.9
2001 4/	5.3	...	...	0.6	7.1	9.1	7.3	6.2	5.2	5.5	5.7	*
2000	5.4	...	...	0.6	8.1	8.7	7.6	6.5	4.2	5.1	5.8	*
1999	5.4	...	...	0.4	8.1	9.2	7.6	5.7	4.9	5.8	5.7	*
Black, male												
2003	8.8	...	...	1.0	12.1	16.5	12.2	9.1	8.7	8.3	11.3	*
2002	9.1	...	...	0.8	11.3	17.3	13.0	10.2	8.4	9.7	13.8	*
2001 4/	9.2	...	...	0.9	13.0	16.5	12.3	9.9	9.4	10.7	13.5	*
2000	9.4	...	...	1.0	14.2	15.5	13.2	11.2	7.5	11.1	12.1	*
1999	9.6	...	...	0.6	14.3	16.8	13.3	10.0	9.1	11.8	13.8	*
Black, female												
2003	1.8	...	...	*	2.0	2.7	2.9	2.6	2.0	*	*	*
2002	1.5	...	...	*	1.7	2.0	2.9	2.2	1.9	*	*	*
2001 4/	1.7	...	...	*	1.3	2.4	2.9	3.0	1.8	*	*	*
2000	1.7	...	...	*	2.2	2.6	2.7	2.5	1.6	*	*	*
1999	1.6	...	...	*	1.9	2.3	2.6	2.0	1.6	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
All races 3/, both sexes												
2003	5.8	...	...	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
2002	5.9	...	...	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001	5.9	...	...	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000	5.9	...	...	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999	5.9	...	...	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6
All races, male												
2003	10.4	...	...	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
2002	10.6	...	...	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001	10.6	...	...	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000	10.5	...	...	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999	10.6	...	...	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6
All races, female												
2003	1.4	...	...	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
2002	1.4	...	...	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001	1.5	...	...	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000	1.5	...	...	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999	1.5	...	...	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
White, both sexes												
2003	6.6	...	...	0.2	5.3	6.4	7.6	9.2	9.3	10.3	13.4	12.9
2002	6.7	...	...	0.2	5.5	6.5	7.9	8.9	9.4	10.8	14.2	12.9
2001	6.7	...	...	0.2	5.6	7.1	7.5	8.7	9.2	10.5	14.1	13.2
2000	6.6	...	...	0.3	6.1	6.6	7.9	8.5	8.3	10.1	14.1	14.3
1999	6.6	...	...	0.3	6.2	7.0	7.7	8.2	8.6	10.6	14.1	13.6
White, male												
2003	11.7	...	...	0.3	9.4	10.9	12.6	15.6	16.5	20.2	30.9	39.9
2002	11.9	...	...	0.4	9.6	10.8	13.3	15.2	16.7	21.4	33.3	40.5
2001	11.8	...	...	0.4	9.7	12.0	12.6	14.9	16.1	20.9	32.9	42.1
2000	11.6	...	...	0.5	10.6	11.2	13.1	14.4	14.7	19.8	33.3	46.5
1999	11.7	...	...	0.4	10.8	11.8	12.6	13.9	15.2	21.0	33.3	44.9
White, female												
2003	1.6	...	...	*	1.0	1.7	2.4	2.9	2.5	1.8	1.6	0.9
2002	1.6	...	...	*	1.0	1.9	2.5	2.6	2.5	1.7	1.5	0.9
2001	1.7	...	...	*	1.1	1.8	2.4	2.7	2.7	1.8	1.6	1.1
2000	1.7	...	...	*	1.2	1.8	2.7	2.6	2.4	1.9	1.5	1.2
1999	1.7	...	...	0.1	1.3	2.0	2.6	2.6	2.4	1.9	1.5	1.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by discharge of firearms (X72-X74)												
Black, both sexes												
2003	2.6	...	...	*	4.3	5.1	2.6	2.5	2.7	3.3	4.0	*
2002	2.8	...	...	*	4.1	5.0	3.5	2.9	2.8	3.3	4.4	*
2001	2.9	...	...	*	4.9	4.9	3.4	2.8	2.8	3.7	5.2	*
2000	3.0	...	...	*	5.2	5.0	3.4	3.3	2.9	3.7	3.5	*
1999	3.1	...	...	*	5.6	5.0	3.5	3.0	3.4	4.0	4.7	*
Black, male												
2003	5.0	...	...	*	7.8	10.0	4.8	4.7	5.1	6.6	9.9	*
2002	5.4	...	...	*	7.5	10.0	6.3	5.5	5.5	7.1	11.5	*
2001	5.5	...	...	*	9.2	9.4	6.3	4.8	5.7	7.7	12.4	*
2000	5.5	...	...	*	9.6	9.3	6.2	5.8	5.7	8.8	8.2	*
1999	5.9	...	...	*	10.3	9.6	6.7	5.6	7.0	8.7	12.0	*
Black, female												
2003	0.5	...	...	*	0.7	*	0.7	*	*	*	*	*
2002	0.5	...	...	*	0.7	*	0.9	*	*	*	*	*
2001	0.6	...	...	*	*	0.9	0.9	1.0	*	*	*	*
2000	0.7	...	...	*	0.9	1.1	0.8	1.1	*	*	*	*
1999	0.6	...	...	*	1.0	0.8	0.7	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
All races 3/, both sexes												
2003	5.0	...	...	0.4	4.6	6.7	8.3	7.8	5.5	3.5	4.0	5.1
2002	5.0	...	...	0.4	4.7	6.6	8.2	7.9	5.2	3.8	4.5	6.0
2001 4/	4.8	...	...	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000	4.5	...	...	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999	4.5	...	...	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7
All races, male												
2003	7.3	...	...	0.6	7.1	10.2	12.1	10.6	7.3	5.0	6.4	11.1
2002	7.3	...	...	0.6	7.4	10.2	11.8	10.8	7.1	5.2	7.1	12.9
2001 4/	7.1	...	...	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000	6.6	...	...	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999	6.6	...	...	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5
All races, female												
2003	2.8	...	...	0.2	2.1	3.2	4.5	5.2	3.7	2.2	2.5	2.4
2002	2.9	...	...	0.2	2.0	3.0	4.7	5.2	3.5	2.5	2.8	3.0
2001 4/	2.6	...	...	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000	2.5	...	...	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999	2.5	...	...	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
White, both sexes												
2003	5.5	...	...	0.4	4.9	7.3	9.1	8.7	5.9	3.6	4.2	5.2
2002	5.5	...	...	0.4	5.2	7.1	9.1	8.8	5.6	3.9	4.7	6.2
2001 4/	5.3	...	...	0.5	5.0	6.8	8.8	8.2	5.2	3.8	4.5	5.5
2000	4.9	...	...	0.5	4.6	6.2	8.1	7.5	5.0	3.4	4.6	6.7
1999	4.9	...	...	0.4	4.3	6.5	8.2	7.2	4.6	3.8	5.2	7.1
White, male												
2003	7.9	...	...	0.6	7.5	10.9	13.1	11.7	7.7	5.0	6.6	11.5
2002	8.0	...	...	0.6	8.1	10.7	12.9	11.8	7.6	5.4	7.4	13.4
2001 4/	7.7	...	...	0.7	7.9	10.4	12.8	11.0	6.8	5.4	7.3	12.9
2000	7.1	...	...	0.7	7.3	9.5	11.8	10.0	6.5	4.5	7.8	15.2
1999	7.0	...	...	0.6	6.6	9.9	11.8	9.6	6.1	5.4	7.9	16.6
White, female												
2003	3.1	...	...	0.2	2.1	3.4	5.1	5.8	4.1	2.3	2.6	2.4
2002	3.2	...	...	0.3	2.1	3.3	5.3	5.8	3.7	2.6	2.9	3.1
2001 4/	2.9	...	...	0.2	2.0	3.0	4.9	5.3	3.7	2.4	2.7	2.3
2000	2.8	...	...	0.2	1.9	2.8	4.5	4.9	3.6	2.5	2.6	3.3
1999	2.8	...	...	0.1	1.9	2.9	4.6	4.8	3.3	2.5	3.4	3.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)												
Black, both sexes												
2003	2.5	...	...	0.5	2.8	4.2	4.6	3.1	2.2	1.2	*	*
2002	2.3	...	...	0.4	2.4	4.2	4.2	3.0	2.0	1.5	*	*
2001 4/	2.3	...	...	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000	2.4	...	...	0.5	2.9	3.7	4.2	3.2	1.3	1.4	2.3	*
1999	2.3	...	...	0.4	2.5	4.2	4.1	2.7	1.5	1.9	*	*
Black, male												
2003	3.8	...	...	0.9	4.3	6.5	7.3	4.4	3.6	*	*	*
2002	3.7	...	...	0.6	3.8	7.3	6.7	4.7	2.9	*	*	*
2001 4/	3.7	...	...	0.6	3.8	7.1	6.1	5.0	3.7	3.0	*	*
2000	3.9	...	...	0.9	4.6	6.3	6.9	5.4	1.8	*	*	*
1999	3.7	...	...	*	4.0	7.2	6.6	4.4	2.2	3.1	*	*
Black, female												
2003	1.3	...	...	*	1.3	2.0	2.2	2.1	*	*	*	*
2002	1.1	...	...	*	1.0	1.5	1.9	1.6	1.4	*	*	*
2001 4/	1.1	...	...	*	0.7	1.5	2.0	2.0	*	*	*	*
2000	1.0	...	...	*	1.3	1.5	1.9	1.4	*	*	*	*
1999	1.0	...	...	*	0.9	1.6	1.9	1.2	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
All races 3/, both sexes												
2003	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
2002	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 4/	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4
All races, male												
2003	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
2002	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 4/	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4
All races, female												
2003	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
2002	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 4/	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
White, both sexes												
2003	3.7	6.9	1.7	0.5	6.7	6.1	4.6	3.5	2.2	1.9	2.1	2.0
2002	3.7	5.4	2.0	0.6	6.7	5.9	4.8	3.3	2.6	1.9	1.9	1.8
2001 4/	4.9	6.2	1.9	0.7	7.2	8.3	7.3	5.0	3.4	2.5	2.1	2.2
2000	3.6	6.6	1.7	0.7	6.4	5.7	4.7	3.3	2.4	1.9	2.0	2.2
1999	3.8	6.7	1.7	0.8	6.8	6.0	4.8	3.3	2.4	2.2	2.0	2.1
White, male												
2003	5.4	8.1	1.5	0.6	10.6	9.2	6.3	5.0	3.1	2.7	2.8	3.0
2002	5.4	6.2	2.1	0.6	10.6	8.9	6.8	4.6	3.7	2.6	2.4	2.7
2001 4/	7.2	7.3	2.2	0.7	11.2	12.3	10.7	7.4	4.9	3.2	2.6	2.8
2000	5.2	8.2	1.9	0.9	9.9	8.4	6.5	4.6	3.5	2.4	2.5	2.8
1999	5.5	7.9	1.8	0.9	10.3	8.8	6.7	4.8	3.6	2.9	2.7	2.7
White, female												
2003	2.0	5.7	1.8	0.3	2.5	2.8	2.8	2.1	1.4	1.3	1.7	1.6
2002	2.0	4.6	1.8	0.6	2.5	2.8	2.8	2.1	1.5	1.3	1.6	1.5
2001 4/	2.6	5.1	1.6	0.6	3.0	4.1	3.8	2.7	2.0	1.8	1.8	2.0
2000	2.1	5.0	1.5	0.6	2.7	2.9	2.8	2.1	1.4	1.5	1.8	1.9
1999	2.1	5.4	1.7	0.7	3.0	3.1	2.9	1.8	1.3	1.6	1.6	1.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)												
Black, both sexes												
2003	22.0	16.1	6.1	2.2	47.6	45.2	23.4	15.4	8.1	6.2	6.6	*
2002	22.0	17.3	6.3	2.0	46.8	45.3	23.7	15.5	8.3	5.7	7.4	*
2001 4/	22.1	18.9	6.8	1.4	47.3	43.7	25.0	16.1	8.9	7.1	6.7	*
2000	21.5	22.8	5.8	1.8	47.8	40.8	23.9	14.8	8.3	7.6	7.0	*
1999	21.1	19.5	6.1	2.2	47.7	39.3	23.2	14.0	8.2	6.3	7.4	*
Black, male												
2003	38.9	17.8	7.4	2.9	84.6	82.5	40.2	26.3	14.8	11.1	11.6	*
2002	38.4	16.3	7.1	2.5	83.1	82.2	38.8	27.1	15.4	10.6	13.8	*
2001 4/	38.3	21.0	8.0	2.0	85.7	78.8	38.5	26.9	15.0	12.6	10.0	*
2000	37.2	23.3	5.9	2.1	85.3	73.9	38.5	26.0	14.7	13.9	11.8	*
1999	36.1	19.5	5.9	2.7	84.7	69.2	37.2	23.4	15.1	11.8	12.0	*
Black, female												
2003	6.6	14.2	4.8	1.5	10.1	11.1	8.5	6.0	2.7	2.7	3.8	*
2002	7.0	18.5	5.5	1.5	10.3	11.9	10.4	5.6	2.5	2.3	3.7	*
2001 4/	7.4	16.7	5.6	0.8	8.9	12.1	13.0	6.8	3.9	3.3	4.8	*
2000	7.2	22.2	5.7	1.6	10.7	10.9	11.0	5.3	3.3	3.1	4.3	*
1999	7.6	19.5	6.3	1.7	11.2	12.4	10.8	5.9	2.6	2.5	4.8	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
All races 3/, both sexes												
2003	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
2002	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6
All races, male												
2003	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
2002	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*
All races, female												
2003	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
2002	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
White, both sexes												
2003	2.2	*	0.2	0.3	5.0	4.2	2.7	1.9	1.1	0.7	0.6	0.5
2002	2.2	*	0.3	0.3	5.0	4.1	2.8	1.8	1.2	0.7	0.8	*
2001	2.2	*	0.3	0.4	5.0	4.2	2.9	1.7	1.2	0.9	0.7	*
2000	2.1	*	*	0.4	4.6	3.7	2.7	1.8	1.3	0.8	0.6	0.6
1999	2.2	*	0.2	0.4	4.9	3.8	2.7	1.7	1.2	0.9	0.7	0.6
White, male												
2003	3.5	*	*	0.4	8.5	6.9	3.9	2.8	1.6	1.1	0.5	*
2002	3.5	*	0.3	0.4	8.4	6.6	4.2	2.5	1.8	0.9	1.0	*
2001	3.5	*	*	0.5	8.3	6.7	4.3	2.6	1.6	1.1	0.8	*
2000	3.2	*	*	0.5	7.7	5.9	3.9	2.6	1.8	1.0	0.6	*
1999	3.4	*	*	0.5	8.0	6.1	4.0	2.6	1.9	1.2	0.7	*
White, female												
2003	0.9	*	*	0.1	1.3	1.4	1.5	1.0	0.7	0.4	0.6	*
2002	1.0	*	*	0.3	1.4	1.5	1.4	1.1	0.7	0.6	0.6	*
2001	1.0	*	*	0.3	1.5	1.6	1.5	0.9	0.8	0.7	0.6	*
2000	1.0	*	*	0.2	1.4	1.5	1.4	1.0	0.8	0.7	0.6	*
1999	0.9	*	*	0.3	1.5	1.5	1.5	0.9	0.6	0.7	0.6	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)												
Black, both sexes												
2003	16.8	*	*	1.4	41.9	38.8	15.9	8.7	3.9	2.5	*	*
2002	16.7	*	*	1.2	41.8	38.2	16.3	8.5	4.1	2.4	2.2	*
2001	15.8	*	*	0.8	41.8	35.1	15.0	7.5	3.7	2.7	*	*
2000	15.6	*	*	1.0	41.2	33.2	15.0	8.4	3.5	2.7	*	*
1999	15.4	*	*	1.2	42.0	31.4	14.1	8.0	4.4	2.4	2.3	*
Black, male												
2003	31.6	*	*	1.9	76.8	74.1	29.1	15.9	7.5	4.6	*	*
2002	31.0	*	*	1.6	76.1	72.8	29.2	15.1	7.6	4.7	*	*
2001	29.8	*	*	1.3	78.2	66.9	26.0	13.7	7.1	5.6	*	*
2000	29.2	*	*	1.5	76.5	63.3	26.7	15.1	6.3	5.4	*	*
1999	28.4	*	*	1.8	76.9	59.1	24.0	13.9	8.9	4.6	*	*
Black, female												
2003	3.3	*	*	0.8	6.5	6.6	4.2	2.5	*	*	*	*
2002	3.6	*	*	0.8	7.3	6.9	4.8	2.8	1.4	*	*	*
2001	3.1	*	*	*	5.4	6.3	5.4	2.2	*	*	*	*
2000	3.2	*	*	*	6.4	6.1	4.6	2.6	*	*	*	*
1999	3.7	*	*	0.6	7.6	6.5	5.3	3.0	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
All races 3/, both sexes												
2003	2.0	8.3	2.1	0.3	2.3	2.4	2.6	2.2	1.4	1.5	1.8	1.7
2002	2.0	7.3	2.4	0.4	2.2	2.6	2.7	2.2	1.6	1.4	1.4	1.7
2001 4/	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8
All races, male												
2003	2.6	9.7	2.3	0.4	3.1	3.0	3.5	3.1	2.1	2.0	2.6	2.5
2002	2.6	7.8	2.6	0.3	3.0	3.2	3.4	3.2	2.4	2.1	1.9	2.3
2001 4/	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6
All races, female												
2003	1.4	6.9	2.0	0.3	1.6	1.8	1.8	1.3	0.8	1.0	1.2	1.4
2002	1.4	6.8	2.2	0.4	1.4	1.9	1.9	1.3	0.9	0.8	1.1	1.4
2001 4/	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7
2000	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
White, both sexes												
2003	1.5	6.9	1.5	0.2	1.7	1.8	1.9	1.6	1.1	1.2	1.5	1.6
2002	1.5	5.3	1.7	0.3	1.7	1.8	2.0	1.6	1.3	1.2	1.1	1.4
2001 4/	2.6	6.0	1.6	0.3	2.2	4.1	4.4	3.3	2.3	1.6	1.4	1.8
2000	1.5	6.5	1.5	0.4	1.8	2.0	2.0	1.5	1.1	1.0	1.5	1.5
1999	1.6	6.6	1.5	0.4	1.9	2.1	2.0	1.6	1.2	1.3	1.4	1.5
White, male												
2003	1.9	8.0	1.4	0.3	2.1	2.3	2.4	2.2	1.5	1.5	2.2	2.3
2002	1.9	6.2	1.8	0.2	2.2	2.3	2.5	2.1	1.9	1.8	1.3	2.0
2001 4/	3.7	7.0	1.9	0.3	2.8	5.6	6.4	4.8	3.3	2.1	1.8	2.3
2000	2.0	8.1	1.7	0.4	2.3	2.5	2.6	2.0	1.7	1.4	1.9	2.1
1999	2.1	7.7	1.5	0.4	2.3	2.7	2.7	2.3	1.7	1.7	2.0	1.9
White, female												
2003	1.1	5.6	1.6	0.2	1.2	1.4	1.4	1.0	0.7	0.9	1.1	1.2
2002	1.1	4.5	1.6	0.4	1.1	1.3	1.4	1.0	0.8	0.7	1.0	1.2
2001 4/	1.6	4.9	1.4	0.4	1.5	2.5	2.3	1.8	1.3	1.1	1.2	1.6
2000	1.1	4.8	1.3	0.4	1.3	1.4	1.4	1.1	0.6	0.7	1.2	1.3
1999	1.2	5.4	1.5	0.5	1.4	1.6	1.4	0.9	0.7	0.9	1.0	1.4

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)												
Black, both sexes												
2003	5.2	15.2	5.3	0.8	5.7	6.4	7.5	6.8	4.1	3.7	4.9	*
2002	5.3	16.5	5.8	0.8	5.0	7.1	7.4	7.1	4.1	3.4	5.2	*
2001 4/	6.3	18.6	6.1	0.5	5.5	8.7	9.9	8.5	5.2	4.4	4.8	*
2000	5.9	21.6	5.5	0.8	6.5	7.6	8.9	6.5	4.9	4.8	5.1	*
1999	5.7	19.2	5.4	1.0	5.7	7.9	9.1	6.0	3.8	4.0	5.1	*
Black, male												
2003	7.4	16.1	6.5	0.9	7.7	8.4	11.1	10.5	7.3	6.5	8.7	*
2002	7.3	15.7	6.4	0.9	7.0	9.4	9.6	12.0	7.8	5.9	9.2	*
2001 4/	8.5	20.7	7.3	0.7	7.5	11.8	12.6	13.2	8.0	7.0	*	*
2000	8.0	21.1	5.5	*	8.8	10.6	11.7	10.8	8.3	8.5	8.2	*
1999	7.7	19.2	5.4	0.9	7.8	10.1	13.2	9.6	6.2	7.2	7.1	*
Black, female												
2003	3.3	14.2	4.1	0.7	3.6	4.6	4.2	3.6	1.6	*	*	*
2002	3.4	17.3	5.1	0.7	3.0	5.0	5.6	2.8	*	*	*	*
2001 4/	4.3	16.4	4.8	*	3.5	5.8	7.6	4.6	2.9	2.7	4.3	*
2000	4.0	22.2	5.5	1.1	4.3	4.8	6.4	2.7	2.1	2.3	3.4	*
1999	3.9	19.1	5.4	1.1	3.6	5.9	5.5	2.9	1.9	*	4.0	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
All races 3/, both sexes												
2003	0.1	*	*	*	0.2	0.3	0.3	0.1	*	*	*	*
2002	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
2000	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.2	0.3	0.3	0.1	0.1	*	*	*
All races, male												
2003	0.3	*	*	*	0.4	0.6	0.5	0.2	*	*	*	*
2002	0.3	*	*	*	0.4	0.5	0.5	0.3	*	*	*	*
2001	0.3	*	*	*	0.4	0.5	0.5	0.3	0.2	*	*	*
2000	0.3	*	*	*	0.4	0.5	0.4	0.3	*	*	*	*
1999	0.3	*	*	*	0.3	0.7	0.5	0.3	0.2	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
White, both sexes												
2003	0.1	*	*	*	0.2	0.2	0.3	0.1	*	*	*	*
2002	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	0.1	*	*	*	*
2000	0.1	*	*	*	0.1	0.2	0.2	0.1	*	*	*	*
1999	0.1	*	*	*	0.1	0.3	0.2	0.1	*	*	*	*
White, male												
2003	0.2	*	*	*	0.3	0.5	0.5	0.2	*	*	*	*
2002	0.2	*	*	*	0.3	0.4	0.4	0.2	*	*	*	*
2001	0.2	*	*	*	0.4	0.5	0.4	0.2	*	*	*	*
2000	0.2	*	*	*	0.3	0.4	0.3	0.3	*	*	*	*
1999	0.2	*	*	*	0.2	0.5	0.4	0.3	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Legal intervention (Y35,Y89.0)												
Black, both sexes												
2003	0.3	*	*	*	0.6	1.0	0.4	*	*	*	*	*
2002	0.3	*	*	*	0.6	0.5	0.5	*	*	*	*	*
2001	0.2	*	*	*	0.4	0.4	0.4	*	*	*	*	*
2000	0.3	*	*	*	0.5	0.6	0.6	*	*	*	*	*
1999	0.3	*	*	*	0.5	0.8	0.6	*	*	*	*	*
Black, male												
2003	0.6	*	*	*	1.0	2.0	0.8	*	*	*	*	*
2002	0.6	*	*	*	1.2	1.1	1.0	*	*	*	*	*
2001	0.4	*	*	*	0.8	0.8	0.8	*	*	*	*	*
2000	0.6	*	*	*	1.0	1.3	1.2	*	*	*	*	*
1999	0.7	*	*	*	1.1	1.6	1.3	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
All races 3/, both sexes												
2003	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
2002	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4
All races, male												
2003	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
2002	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1
All races, female												
2003	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
2002	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
White, both sexes												
2003	1.8	1.7	0.2	0.1	1.5	2.3	3.7	3.1	1.3	0.6	0.7	1.2
2002	1.7	1.8	0.4	0.2	1.3	2.1	3.4	2.8	1.2	0.7	0.7	1.4
2001	1.5	1.4	0.3	0.1	1.1	1.9	3.1	2.5	0.9	0.5	0.9	1.0
2000	1.3	1.4	0.2	0.1	1.0	1.7	2.8	2.1	0.9	0.5	0.6	1.3
1999	1.4	1.2	0.2	0.1	1.1	1.9	2.8	2.0	0.9	0.6	0.8	1.4
White, male												
2003	2.3	1.9	*	0.2	2.1	3.0	4.7	3.9	1.5	0.7	1.0	1.5
2002	2.2	1.8	0.4	0.2	1.9	2.9	4.4	3.3	1.5	0.9	0.9	2.3
2001	2.0	1.7	*	0.2	1.6	2.7	4.0	3.2	1.1	0.7	1.2	*
2000	1.8	*	*	0.2	1.4	2.4	3.8	2.8	1.1	0.7	1.0	*
1999	1.9	1.4	*	0.2	1.8	2.5	3.8	2.6	1.1	0.8	1.1	2.1
White, female												
2003	1.3	1.4	*	*	0.9	1.5	2.7	2.4	1.0	0.6	0.6	1.0
2002	1.2	1.8	0.4	0.1	0.7	1.3	2.5	2.2	0.9	0.6	0.6	1.1
2001	1.0	*	*	*	0.6	1.2	2.2	1.7	0.6	0.4	0.6	0.8
2000	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.4	0.4	1.1
1999	0.9	*	*	*	0.4	1.2	1.8	1.4	0.7	0.4	0.6	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)												
Black, both sexes												
2003	1.8	3.2	*	*	0.8	1.8	3.8	4.0	1.9	1.2	*	*
2002	2.0	5.5	*	*	0.9	2.3	4.2	4.2	1.5	*	*	*
2001	1.7	5.5	*	*	0.7	1.7	3.9	3.8	1.3	*	*	*
2000	1.8	*	*	0.3	0.9	1.9	4.1	3.7	1.8	*	*	*
1999	1.9	*	*	*	1.2	2.2	4.2	4.1	1.2	*	*	*
Black, male												
2003	2.7	*	*	*	1.4	2.9	5.3	6.4	3.2	*	*	*
2002	2.9	*	*	*	1.4	3.5	6.0	6.4	2.5	*	*	*
2001	2.7	6.0	*	*	1.3	2.6	6.2	6.3	2.4	*	*	*
2000	2.8	*	*	*	1.6	3.1	6.0	5.9	3.0	*	*	*
1999	3.0	*	*	*	1.9	3.6	6.4	7.2	2.5	*	*	*
Black, female												
2003	0.9	*	*	*	*	*	2.4	1.9	*	*	*	*
2002	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001	0.9	*	*	*	*	0.9	1.9	1.6	*	*	*	*
2000	0.9	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999	0.9	*	*	*	*	1.0	2.3	1.6	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
All races 3/, both sexes												
2003	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
All races, male												
2003	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.3	0.2	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.6	0.2	0.2	0.1	*	*	*	*
All races, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
White, both sexes												
2003	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2002	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2001	0.1	*	*	*	0.1	0.1	0.1	0.1	*	*	*	*
2000	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
1999	0.1	*	*	*	0.3	0.1	0.1	0.1	*	*	*	*
White, male												
2003	0.1	*	*	*	0.2	0.1	0.1	0.1	*	*	*	*
2002	0.1	*	*	*	0.3	0.2	0.2	0.1	*	*	*	*
2001	0.1	*	*	*	0.2	0.2	0.2	*	*	*	*	*
2000	0.1	*	*	*	0.3	0.2	0.1	*	*	*	*	*
1999	0.2	*	*	*	0.5	0.2	0.2	*	*	*	*	*
White, female												
2003	0.0	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	0.0	*	*	*	*	*	*	*	*	*	*	*
2000	0.0	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Discharge of firearms, undetermined intent (Y22-Y24)												
Black, both sexes												
2003	0.1	*	*	*	*	*	*	*	*	*	*	*
2002	0.1	*	*	*	*	*	*	*	*	*	*	*
2001	0.1	*	*	*	*	*	*	*	*	*	*	*
2000	0.1	*	*	*	0.3	*	*	*	*	*	*	*
1999	0.2	*	*	*	0.6	*	*	*	*	*	*	*
Black, male												
2003	0.2	*	*	*	*	*	*	*	*	*	*	*
2002	0.2	*	*	*	*	*	*	*	*	*	*	*
2001	0.2	*	*	*	*	*	*	*	*	*	*	*
2000	0.3	*	*	*	*	*	*	*	*	*	*	*
1999	0.3	*	*	*	1.0	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
All races 3/, both sexes												
2003	1.7	1.9	0.2	0.1	1.2	2.0	3.5	3.0	1.2	0.7	0.7	1.1
2002	1.6	2.5	0.4	0.2	1.0	2.0	3.3	2.8	1.2	0.7	0.7	1.4
2001	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3
All races, male												
2003	2.2	2.4	*	0.1	1.7	2.7	4.5	3.9	1.5	0.7	0.8	1.4
2002	2.1	2.5	0.3	0.2	1.4	2.7	4.2	3.4	1.5	0.9	0.8	2.2
2001	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8
All races, female												
2003	1.2	1.3	*	*	0.7	1.3	2.5	2.2	1.0	0.7	0.6	0.9
2002	1.1	2.5	0.4	0.1	0.6	1.2	2.3	2.1	0.8	0.5	0.6	1.1
2001	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
White, both sexes												
2003	1.7	1.7	0.2	0.1	1.4	2.2	3.6	3.0	1.2	0.6	0.7	1.2
2002	1.6	1.8	0.4	0.1	1.1	2.0	3.3	2.7	1.2	0.6	0.7	1.4
2001	1.4	1.4	0.3	0.1	1.0	1.8	3.0	2.4	0.8	0.5	0.8	1.0
2000	1.3	1.4	0.2	0.1	0.8	1.6	2.7	2.1	0.8	0.5	0.6	1.2
1999	1.3	1.2	0.2	0.1	0.8	1.8	2.6	1.9	0.9	0.5	0.7	1.3
White, male												
2003	2.2	1.9	*	0.1	1.9	2.9	4.6	3.8	1.3	0.6	0.8	*
2002	2.0	1.8	0.4	0.1	1.5	2.8	4.2	3.2	1.5	0.7	0.8	2.3
2001	1.9	1.7	*	0.2	1.4	2.5	3.9	3.1	1.0	0.6	1.0	*
2000	1.7	*	*	0.2	1.2	2.2	3.7	2.8	1.0	0.6	0.9	*
1999	1.7	1.4	*	0.1	1.3	2.3	3.5	2.5	1.1	0.7	0.9	*
White, female												
2003	1.3	1.4	*	*	0.8	1.5	2.6	2.3	1.0	0.6	0.6	1.0
2002	1.2	1.8	0.4	0.1	0.7	1.3	2.4	2.2	0.9	0.6	0.6	1.1
2001	1.0	*	*	*	0.6	1.1	2.1	1.7	0.6	0.3	0.6	0.8
2000	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.3	0.4	1.1
1999	0.8	*	*	*	0.4	1.2	1.7	1.4	0.7	0.4	0.6	1.1

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)												
Black, both sexes												
2003	1.7	3.2	*	*	0.5	1.6	3.7	3.9	1.9	1.2	*	*
2002	1.9	5.5	*	*	0.6	2.1	4.1	4.2	1.4	*	*	*
2001	1.7	5.5	*	*	0.5	1.6	3.9	3.7	1.1	*	*	*
2000	1.7	*	*	*	0.5	1.7	3.9	3.6	1.6	*	*	*
1999	1.7	*	*	*	0.6	2.0	4.1	4.0	1.1	*	*	*
Black, male												
2003	2.6	*	*	*	0.8	2.7	5.2	6.3	3.2	*	*	*
2002	2.7	*	*	*	0.8	3.1	5.8	6.3	2.5	*	*	*
2001	2.5	6.0	*	*	0.8	2.4	6.2	6.1	2.1	*	*	*
2000	2.5	*	*	*	1.0	2.6	5.8	5.8	2.8	*	*	*
1999	2.7	*	*	*	0.9	3.1	6.2	6.9	2.5	*	*	*
Black, female												
2003	0.9	*	*	*	*	*	2.3	1.9	*	*	*	*
2002	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001	0.8	*	*	*	*	0.9	1.8	1.6	*	*	*	*
2000	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999	0.9	*	*	*	*	0.9	2.2	1.6	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
All races 3/, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	0.0	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
All races, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
White, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	0.0	*	*	*	*	*	*	*	*	*	*	*
White, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Operations of war and their sequelae (Y36,Y89.1)												
Black, both sexes												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, male												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*
Black, female												
2003	*	*	*	*	*	*	*	*	*	*	*	*
2002	*	*	*	*	*	*	*	*	*	*	*	*
2001	*	*	*	*	*	*	*	*	*	*	*	*
2000	*	*	*	*	*	*	*	*	*	*	*	*
1999	*	*	*	*	*	*	*	*	*	*	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
All races 3/, both sexes												
2003	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
2002	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7
All races, male												
2003	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
2002	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5
All races, female												
2003	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
2002	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
White, both sexes												
2003	1.0	*	*	*	0.1	0.2	0.4	0.6	1.2	2.8	6.1	11.0
2002	1.0	*	*	0.1	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.2
2001	1.1	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.1
2000	1.1	*	*	0.1	0.1	0.2	0.3	0.7	1.4	3.3	6.4	13.1
1999	1.0	1.2	0.2	*	0.1	0.2	0.3	0.5	1.2	3.0	6.3	12.1
White, male												
2003	1.0	*	*	*	0.1	0.2	0.4	0.6	1.4	3.1	7.0	12.8
2002	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.9	6.3	11.8
2001	0.9	*	*	*	*	0.2	0.3	0.6	1.4	2.8	7.7	13.6
2000	1.1	*	*	*	*	0.2	0.3	0.7	1.4	3.7	8.2	15.8
1999	1.0	1.4	*	*	*	0.2	0.3	0.5	1.3	3.2	7.5	15.6
White, female												
2003	1.0	*	*	*	*	0.2	0.3	0.5	1.1	2.5	5.4	10.2
2002	1.1	*	*	*	*	0.1	0.3	0.5	1.2	3.0	6.0	10.9
2001	1.2	*	*	*	*	0.2	0.3	0.6	1.3	2.7	6.2	11.5
2000	1.1	*	*	*	*	0.2	0.3	0.7	1.3	3.0	5.3	12.0
1999	1.1	*	*	*	*	0.1	0.3	0.6	1.1	2.9	5.6	10.7

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)												
Black, both sexes												
2003	1.1	*	*	*	*	*	0.6	1.4	2.7	4.6	8.8	15.2
2002	1.2	*	*	*	*	*	0.8	1.6	2.7	4.6	8.3	14.0
2001	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.3	9.2	23.3
2000	1.3	*	*	*	*	0.4	0.5	1.4	3.5	6.1	10.5	21.0
1999	1.2	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.1
Black, male												
2003	1.0	*	*	*	*	*	*	1.8	2.8	3.9	7.6	*
2002	1.0	*	*	*	*	*	*	1.3	2.5	5.3	7.2	*
2001	1.4	*	*	*	*	*	1.0	1.8	4.2	7.8	9.1	29.5
2000	1.1	*	*	*	*	*	*	1.3	3.1	7.6	10.3	*
1999	1.1	*	*	*	*	*	*	1.4	2.8	5.9	12.3	23.4
Black, female												
2003	1.3	*	*	*	*	*	0.7	1.1	2.5	5.0	9.4	15.4
2002	1.3	*	*	*	*	*	1.1	1.8	2.7	4.1	8.9	11.5
2001	1.4	*	*	*	*	*	0.7	1.3	2.7	5.3	9.3	21.0
2000	1.5	*	*	*	*	*	*	1.5	3.8	5.0	10.6	21.9
1999	1.3	*	*	*	*	*	*	1.2	2.4	4.8	9.6	18.8

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/												
All races 3/, both sexes												
2003	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.1	9.2	4.1
2002 6/	7.0	*	*	*	0.3	1.5	8.1	16.8	18.4	15.5	9.3	4.4
2001 6/	7.1	*	*	*	0.3	1.6	8.3	17.2	18.2	15.5	9.6	5.0
2000 6/	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4
1999 6/	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5
All races, male												
2003	10.9	*	*	*	0.5	2.3	11.4	27.0	29.8	25.4	16.8	7.8
2002 6/	10.8	*	*	*	0.4	2.3	11.5	26.3	30.0	26.0	16.6	9.7
2001 6/	10.8	*	*	*	0.4	2.3	11.9	26.6	29.7	26.3	17.2	11.6
2000 6/	10.9	*	*	*	0.4	2.5	12.5	25.9	30.0	27.1	18.8	11.9
1999 6/	10.9	*	*	*	0.4	2.3	12.6	26.2	30.5	27.4	19.4	11.8
All races, female												
2003	3.4	*	*	*	0.1	0.6	4.7	7.9	8.1	6.5	4.1	2.5
2002 6/	3.4	*	*	*	*	0.7	4.7	7.7	7.6	6.8	4.5	2.1
2001 6/	3.4	*	*	*	*	0.9	4.8	8.1	7.5	6.6	4.6	2.3
2000 6/	3.2	*	*	*	*	0.7	4.5	7.0	8.3	6.6	4.1	2.7
1999 6/	3.2	*	*	*	0.1	0.9	4.5	7.0	7.7	6.5	4.8	3.0

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/												
White, both sexes												
2003	7.4	*	*	*	0.3	1.5	8.2	17.4	18.2	15.2	9.3	4.2
2002 6/	7.2	*	*	*	0.3	1.5	8.3	16.9	17.8	15.4	9.3	4.3
2001 6/	7.2	*	*	*	0.2	1.6	8.3	16.8	17.6	15.3	9.6	4.8
2000 6/	7.1	*	*	*	0.2	1.6	8.4	15.8	18.0	15.6	9.8	5.3
1999 6/	7.0	*	*	*	0.3	1.6	8.2	15.7	17.7	15.3	10.3	5.3
White, male												
2003	11.3	*	*	*	0.5	2.4	11.8	26.8	29.0	25.4	16.8	7.8
2002 6/	11.1	*	*	*	0.4	2.3	11.8	26.4	29.0	25.4	16.5	9.3
2001 6/	11.0	*	*	*	0.4	2.3	11.7	26.0	28.4	25.4	17.2	11.1
2000 6/	11.0	*	*	*	0.4	2.5	12.3	25.3	28.8	26.4	18.6	11.6
1999 6/	10.9	*	*	*	0.4	2.3	12.2	25.2	28.9	26.0	18.9	11.4
White, female												
2003	3.5	*	*	*	*	0.7	4.7	8.0	8.0	6.5	4.3	2.5
2002 6/	3.4	*	*	*	*	0.7	4.7	7.6	7.3	7.0	4.5	2.2
2001 6/	3.4	*	*	*	*	0.8	4.8	7.6	7.4	6.7	4.6	2.2
2000 6/	3.2	*	*	*	*	0.7	4.4	6.5	7.8	6.5	4.1	2.7
1999 6/	3.1	*	*	*	*	0.9	4.1	6.4	7.2	6.5	4.6	2.9

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 5/												
Black, both sexes												
2003	6.3	*	*	*	*	1.1	6.8	18.4	24.2	16.9	8.9	*
2002 6/	6.4	*	*	*	*	1.3	6.7	18.5	25.3	18.3	11.2	*
2001 6/	7.3	*	*	*	*	1.3	8.7	22.5	25.9	20.5	11.1	*
2000 6/	7.4	*	*	*	*	1.7	9.2	22.7	26.4	21.0	11.5	6.3
1999 6/	7.8	*	*	*	*	1.3	9.9	23.4	29.1	23.2	14.6	7.7
Black, male												
2003	10.0	*	*	*	*	1.8	9.6	31.2	42.2	30.7	18.9	*
2002 6/	10.0	*	*	*	*	2.0	9.4	29.3	43.5	35.8	21.8	*
2001 6/	11.6	*	*	*	*	1.9	13.2	35.6	46.3	40.0	21.8	*
2000 6/	11.4	*	*	*	*	2.5	13.6	36.0	43.8	39.9	24.2	*
1999 6/	12.2	*	*	*	*	1.8	14.4	37.7	51.0	45.3	28.3	*
Black, female												
2003	2.9	*	*	*	*	*	4.3	7.4	9.8	7.1	3.2	*
2002 6/	3.2	*	*	*	*	*	4.4	9.3	10.7	5.8	5.2	*
2001 6/	3.5	*	*	*	*	0.8	4.6	11.1	9.4	6.8	5.1	*
2000 6/	3.7	*	*	*	*	1.0	5.2	11.3	12.3	7.7	4.3	*
1999 6/	3.9	*	*	*	*	0.9	5.9	11.2	11.6	7.7	6.9	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/												
All races 3/, both sexes												
2003	9.9	0.6	0.2	0.1	6.4	12.2	20.6	20.0	8.0	4.1	4.2	6.0
2002 6/	9.0	0.7	0.2	0.1	5.5	11.1	19.7	17.9	6.9	3.6	3.8	5.7
2001 6/	7.6	0.5	0.2	0.1	4.5	9.4	17.0	14.8	5.4	3.0	3.5	5.1
2000 6/	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7
1999 6/	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8
All races, male												
2003	12.9	*	0.3	0.1	9.2	16.6	26.5	25.7	9.4	4.6	4.9	7.8
2002 6/	11.8	1.0	*	0.1	7.8	15.5	25.6	23.0	8.1	3.7	4.2	7.3
2001 6/	10.2	*	0.3	0.1	6.6	13.2	22.7	19.7	6.3	3.5	3.8	5.7
2000 6/	9.5	*	*	0.1	5.8	12.5	21.8	18.2	5.4	2.8	3.8	7.5
1999 6/	9.4	*	*	*	5.0	12.7	21.9	17.5	6.0	3.4	3.8	6.2
All races, female												
2003	7.0	*	*	0.1	3.3	7.6	14.8	14.6	6.7	3.7	3.7	5.2
2002 6/	6.3	*	0.3	*	3.0	6.6	13.9	13.1	5.7	3.4	3.5	4.9
2001 6/	5.1	*	*	0.1	2.4	5.6	11.5	10.0	4.6	2.6	3.3	4.9
2000 6/	4.6	*	*	*	2.1	5.1	10.2	8.5	4.4	2.5	3.2	4.9
1999 6/	4.4	*	*	*	2.0	5.0	9.7	8.0	3.8	2.7	3.8	4.3

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/												
White, both sexes												
2003	10.4	*	0.2	0.1	7.6	13.6	21.7	20.0	7.8	4.1	4.5	6.3
2002 6/	9.4	*	*	0.1	6.4	12.3	20.6	17.5	6.6	3.7	3.9	5.8
2001 6/	7.8	*	*	0.1	5.2	10.3	17.5	14.0	5.1	3.0	3.6	5.1
2000 6/	7.1	*	*	0.1	4.6	9.5	16.1	12.4	4.8	2.7	3.7	5.6
1999 6/	6.9	*	*	0.1	3.9	9.4	15.6	11.7	4.6	3.0	4.0	4.8
White, male												
2003	13.5	*	*	*	11.0	18.6	27.8	25.1	8.7	4.3	5.2	7.8
2002 6/	12.2	*	*	0.1	9.1	17.1	26.6	21.9	7.2	3.6	4.2	7.6
2001 6/	10.4	*	*	*	7.6	14.4	23.2	18.1	5.5	3.4	3.8	5.6
2000 6/	9.6	*	*	0.1	6.7	13.3	21.9	16.5	4.9	2.7	4.0	7.2
1999 6/	9.3	*	*	*	5.5	13.3	21.7	15.6	5.2	3.3	3.9	6.4
White, female												
2003	7.4	*	*	*	3.9	8.3	15.6	15.0	7.0	3.9	4.0	5.6
2002 6/	6.7	*	*	*	3.5	7.3	14.5	13.2	6.0	3.7	3.8	4.9
2001 6/	5.4	*	*	*	2.7	6.0	11.7	10.1	4.8	2.7	3.5	4.9
2000 6/	4.7	*	*	*	2.3	5.5	10.2	8.4	4.7	2.7	3.4	4.9
1999 6/	4.5	*	*	*	2.2	5.2	9.5	7.9	4.1	2.8	4.1	4.2

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7- F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/												
Black, both sexes												
2003	9.2	*	*	*	1.8	8.1	20.8	26.9	11.7	5.0	2.2	*
2002 6/	9.2	*	*	*	2.1	8.5	20.7	26.8	11.5	3.1	2.7	*
2001 6/	8.5	*	*	*	1.9	7.7	20.0	25.2	8.8	3.2	2.6	6.1
2000 6/	8.3	*	*	*	2.0	8.3	20.1	23.8	7.5	3.0	2.4	7.5
1999 6/	8.6	*	*	*	2.1	9.1	20.9	23.7	8.3	3.5	2.2	*
Black, male												
2003	12.7	*	*	*	2.5	10.7	27.7	38.9	19.0	8.6	*	*
2002 6/	12.8	*	*	*	2.9	12.0	27.7	39.2	19.9	5.4	*	*
2001 6/	12.2	*	*	*	2.9	10.8	27.0	39.3	15.1	5.4	*	*
2000 6/	12.0	*	*	*	2.7	12.0	28.2	37.6	12.4	5.3	*	*
1999 6/	12.7	*	*	*	3.2	13.3	30.1	38.2	15.2	5.4	*	*
Black, female												
2003	6.1	*	*	*	1.1	5.8	14.6	16.6	5.8	2.4	*	*
2002 6/	5.8	*	*	*	1.3	5.3	14.5	16.1	4.8	*	*	*
2001 6/	5.1	*	*	*	0.9	4.8	13.8	13.0	3.8	*	*	*
2000 6/	4.9	*	*	*	1.3	4.9	13.0	11.9	3.5	*	*	*
1999 6/	4.8	*	*	*	1.0	5.2	12.9	11.3	2.8	2.1	*	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/												
All races 3/, both sexes												
2003	10.4	*	0.3	0.8	16.6	15.5	11.5	11.2	10.1	10.4	13.5	12.5
2002	10.5	*	0.4	0.8	16.7	15.4	12.1	10.8	10.2	10.9	14.4	12.5
2001	10.4	*	0.5	0.8	16.7	15.5	11.7	10.5	10.1	10.9	14.3	12.8
2000	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2
1999	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5
All races, male												
2003	18.3	*	0.3	1.2	29.2	27.1	19.1	18.8	17.7	20.3	30.2	37.8
2002	18.4	*	0.5	1.2	29.3	26.5	20.1	18.2	18.0	21.3	32.9	38.9
2001	18.2	*	0.5	1.2	29.6	26.7	19.2	17.8	17.4	21.1	32.8	40.2
2000	17.8	*	0.4	1.4	29.4	24.9	19.4	17.6	16.3	20.3	32.2	44.7
1999	18.1	*	0.5	1.5	30.6	25.5	18.9	17.0	17.1	21.5	32.8	43.3
All races, female												
2003	2.7	*	0.3	0.3	3.3	3.6	4.0	3.8	2.9	2.2	2.2	1.3
2002	2.8	*	0.3	0.5	3.5	4.0	4.2	3.6	3.1	2.3	2.1	1.1
2001	2.8	*	0.4	0.4	3.2	4.0	4.3	3.5	3.3	2.4	2.2	1.3
2000	2.8	*	*	0.4	3.5	4.0	4.4	3.6	3.0	2.5	2.0	1.7
1999	2.9	*	0.4	0.5	3.9	4.1	4.4	3.6	2.9	2.4	2.1	1.5

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/												
White, both sexes												
2003	9.2	*	0.2	0.6	11.1	11.2	10.8	11.5	10.8	11.3	14.4	13.5
2002	9.3	*	0.3	0.7	11.3	11.2	11.3	11.1	10.9	11.8	15.3	13.4
2001	9.4	*	0.4	0.8	11.4	11.9	11.1	10.9	10.7	11.7	15.2	13.8
2000	9.1	*	0.2	0.9	11.5	10.9	11.2	10.6	9.9	11.2	15.0	15.2
1999	9.2	*	0.3	0.9	12.0	11.5	11.0	10.3	10.1	11.8	15.1	14.5
White, male												
2003	16.0	*	*	1.0	19.2	18.8	17.5	19.2	18.9	21.9	32.3	41.0
2002	16.1	*	0.4	0.9	19.4	18.5	18.5	18.4	19.0	22.9	34.8	41.7
2001	16.2	*	0.4	1.1	19.5	19.9	18.0	18.3	18.3	22.4	34.7	43.3
2000	15.6	*	*	1.3	19.6	18.1	17.9	17.7	17.0	21.4	34.5	48.4
1999	15.9	*	*	1.3	20.5	19.2	17.6	17.2	17.6	22.8	34.7	46.8
White, female												
2003	2.6	*	*	0.2	2.5	3.1	4.0	4.0	3.2	2.3	2.3	1.3
2002	2.7	*	*	0.4	2.6	3.5	4.1	3.8	3.3	2.4	2.2	1.2
2001	2.7	*	*	0.4	2.7	3.6	4.1	3.7	3.6	2.6	2.3	1.4
2000	2.7	*	*	0.3	2.8	3.5	4.3	3.7	3.2	2.8	2.1	1.8
1999	2.8	*	*	0.5	3.0	3.6	4.3	3.6	3.1	2.6	2.2	1.6

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/												
Black, both sexes												
2003	20.1	*	0.9	1.7	47.7	45.4	19.0	11.4	6.8	6.0	6.1	*
2002	20.2	*	*	1.5	47.7	44.7	20.5	11.8	7.2	5.8	7.1	*
2001	19.3	*	0.9	1.2	48.1	40.7	19.0	10.6	6.9	6.6	7.5	*
2000	19.3	*	*	1.4	48.1	39.4	19.2	12.1	6.7	6.9	5.5	*
1999	19.4	*	1.1	1.6	49.9	37.7	18.3	11.5	8.2	6.6	7.5	*
Black, male												
2003	37.8	*	*	2.5	87.6	87.2	34.8	21.1	12.9	11.7	13.8	*
2002	37.8	*	*	2.1	87.1	85.6	36.9	21.4	13.5	12.1	17.5	*
2001	36.4	*	*	1.8	90.3	77.7	33.2	19.1	13.7	13.8	18.0	*
2000	36.1	*	*	2.2	89.3	74.8	34.3	21.7	12.6	15.1	12.1	*
1999	36.0	*	*	2.3	91.4	71.3	32.2	20.4	16.5	14.0	18.1	*

Worktable 290R. Death rates from 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2003

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin reported multiple-race data in 2003. The multiple-race data for these States were bridged to the single race categories of the 1977 Office of Management and Budget standards for comparability with other States; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003." For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/												
Black, female												
2003	3.9	*	*	0.9	7.4	7.3	5.0	3.1	2.0	2.0	*	*
2002	4.2	*	*	0.9	8.1	7.5	5.9	3.5	2.1	*	*	*
2001	3.8	*	*	*	6.1	7.3	6.5	3.2	1.5	*	*	*
2000	4.0	*	*	*	7.6	7.4	5.7	3.8	1.9	*	*	*
1999	4.4	*	*	0.8	9.0	7.4	6.1	3.8	1.6	*	*	*

* Figure does not meet standards of reliability or precision.
0.0 Quantity more than zero but less than 0.05.
… Category not applicable.
1/ Figures for age not stated included in "All ages" but not distributed among age groups.
2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).
3/ Includes races other than white and black.
4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."
5/ Included in selected categories.
6/ Figures may differ from those previously shown as they are based on a modified list of causes; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2003."

Source: CDC/NCHS, National Vital Statistics System, Mortality.